# An improved convergence theorem for the Newton method under relaxed continuity assumptions

Andrei Dubin ITEP, 117218, B.Cheremushkinskaya 25, Moscow, Russia

## **Abstract**

In the framework of the majorization technique, an improved condition is proposed for the semilocal convergence of the Newton method under the mild assumption that the derivative  $F'(x)$  of the involved operator  $F(x)$  is continuous. Our starting point is the Argyros' representation of the optimal upper bound for the distance  $||x_{k+1} - x_k||$  between the adjacent members of the Newton sequence  $\{x_k\}$ . The major novel element of our proposal is the optimally reconstructed «first integral» approximation  $v_{k+1} = \psi(v_k)$  to the recurrence relation defining the scalar majorizing sequence  $\{v_k\}$ . Compared to the previous results of Argyros, it enables one to obtain a weaker convergence condition that leads to a better bound on the location of the solution of the equation  $F(x) = 0$  and allows for a wider choice of initial guesses  $x_0$ . In the simplest case of the Lipschitz continuous operator  $F'(x)$ , the new convergence condition improves the famous Kantorovich condition provided that  $l_0/l < (6-4\sqrt{2})$ , where  $l_0$  and l stand for the center-Lipschitz and Lipschitz constants respectively.

**Keywords:** Newton method, Banach space, relaxed continuity assumptions, majorization technique, convergence conditions, Kantorovich theorem.

## **1 Introduction**

Resolution of many different problems of the fundamental and applied mathematics involves the Newton method generating the corresponding sequence  $\{x_k\}$  through the recurrence relation

<span id="page-0-0"></span>
$$
x_{k+1} = x_k - (F'(x_k))^{-1} F(x_k) = T(x_k) \qquad , \qquad k \ge 0, \tag{1-1}
$$

where  $F(x) \in Y$  is a continuously differentiable operator which defines a mapping  $D_F \subseteq X \to Y$ between an open subset  $D_F$  of a Banach space X into a given Banach space Y.

The application of this method is built on the so-called Newton-Kantorovich (NK) theorem due to Kantorovich, see [\[ 1](#page-5-0) ] and [\[ 2](#page-5-1) ] for the original proof and further references. The major limitation of this famous theorem is that one has to impose rather stringy smoothness restriction that the operator  $F(x)$  is twice differentiable or at least that the Frechet derivative  $F'(x)$  of  $F(x)$  is Lipschitz continuous. The latter restriction cannot be satisfied in many interesting applied problems. Therefore, the practically important objective is to obtain Kantorovich-type theorems for the Newton method relaxing as much as possible the assumptions about the degree of the continuity of the operator  $F'(y)$ .

Recently, an interesting new proposal  $\lceil 3 \rceil$ ,  $\lceil 4 \rceil$  in this direction is made presuming only the continuity of  $F'(x)$ . The employed majorization technique is built on the novel upper bound for the distance  $||x_{k+1} - x_k||$ . The bound is expressed in terms of certain continuity measure depending *only* on the relative distances  $||x_q - x_0||$  of  $x_q$  ( $q = k, k - 1$ ) from the initial point  $x_0$ . The subtle feature which calls for an optimization is that the resulting convergence conditions introduce a larger number of elementary restrictions compared to the NK theorem.

To accomplish such optimization of  $\lceil 3 \rceil$ ,  $\lceil 4 \rceil$ , we put forward a new implementation of the majorization technique. It optimally introduces the approximate «first integral» representation [\[ 2](#page-5-1) ] of the recurrence relation defining the majorizing sequence. Under the same computational cost, the proposed technique enables one to improve the convergence condition of [4] formulating the results under the relaxed continuity assumptions in accordance with the pattern of the NK theorem.

# **2 Improved convergence theorem**

<span id="page-1-0"></span> $\overline{a}$ 

## **2.1 «First integral» form of the recurrence relations**

To formulate and prove the improved convergence theorem, we derive such implementation of the general majorization condition

<span id="page-1-3"></span>
$$
||x_{k+1} - x_k|| \leq v_{k+1} - v_k \quad , \quad \forall k \geq 0 ,
$$
\n(2-1)

that the scalar majorizing sequence  $\{v_k\}$  is generated by the recurrence relation in the so-called «first integral» form  $\lceil 2 \rceil$ ,  $\lceil 1 \rceil$ :

<span id="page-1-2"></span>
$$
v_{k+1} = \psi(v_k) \qquad , \qquad \forall k \ge 0 \,, \tag{2-2}
$$

where the continuous non-decreasing function  $\psi(\nu) \ge 0$  assumes the form

<span id="page-1-1"></span>
$$
\psi(\nu) = \eta + 2(1 - \omega_0(\nu))^{-1} \cdot \int_0^{\nu} \omega_0(l) dl. \tag{2-3}
$$

Here, the continuous non-decreasing function  $\omega_0(r)$  facilitates the affine-invariant upper bound<sup>1</sup>

$$
|| (F'(x_0))^{-1} \cdot (F'(x) - F'(x_0)) || \le \omega_0(r) \qquad , \qquad || x - x_0 || \le r \quad , \quad r \le R \,, \tag{2-4}
$$

where  $\overline{B}(x_0, R) \subset D_F$  while  $\overline{B}(x_0, R)$  denotes the closed ball of the center  $x_0$  and the radius R. The (uniform) continuity of  $F'(x)$  on  $\overline{B}(x_0, R)$  is implemented through the restriction  $\omega_0(0) = 0$ .

Note that the characteristic feature of the bound [\( 2-4](#page-1-0) ) is that this continuity measure is *centered* at  $x_0$ , i.e. depends on the relative distance  $||x-x_0||$  of x from the *stationary* point identified with  $x_0$ . Being easier to compute, the continuity measure  $\omega_0(\cdot)$  is generically smaller than its conventional non-centered counterpart  $\omega(r)$  and, furthermore,  $\omega_0(r) \ll \omega(r)$  in a variety of practically interesting cases [\[ 4](#page-5-3) ].

 $1$  The matrix norm is presumed to be submultiplicative and consistent with the corresponding vector norm.

#### **2.2 Statement of the theorem**

**The set** Y. Consider a continuously differentiable operator  $F(x)$ :  $D_F \subseteq X \rightarrow Y$  such that 1  $(F'(x_0))^{-1} \in L(X,Y)$  for some  $x_0 \in D_F$ . Assume that the condition (2-4) and the restriction 1  $(\Gamma(x_0)) \in L(X, Y)$  for some  $x_0 \in D_F$ . Assume that the condition (2-4) and the restriction  $\eta \ge ||(F'(x_0))^{-1}F(x_0)|| = ||x_1 - x_0||$  are valid for some constants  $\eta > 0$ ,  $R > 0$  and a continuous non-decreasing function  $\omega_0(r) \ge 0$  with  $\omega_0(0) = 0$ . Presume also that (2-3) defines such continuous non-decreasing function  $\psi(\nu)$  that the (associated to (2-2)) fixed point equation

<span id="page-2-0"></span>
$$
U = \psi(U) \tag{2-5}
$$

has a minimal solution  $v_*$  satisfying the restrictions  $\eta < v_* \le R$ ,  $\overline{B}(x_0, R) \subset D_F$  where  $\overline{B}(x_0, v_*)$ denotes the closed ball of the center  $x_0$  and the radius  $v_*$ . Finally, let the scalar sequence  $\{v_k\}$ be generated by the relation (2-2) supplemented by the initial conditions  $v_0 = 0$ ,  $v_1 = \psi(0) = \eta$ .

**Theorem.** Under the set  $\Upsilon$  of the conditions, the Newton sequence  $\{x_k\} \equiv \{T^k x_0\}$  is welldefined, remains in  $\overline{B}(x_0, v_*)$  and converges to a solution  $x_*$  of the equation  $F(x) = 0$ . Moreover, the majorization estimates [\( 2-1](#page-1-3) ) and

<span id="page-2-1"></span>
$$
||x_* - x_k|| \leq v_* - v_k \qquad , \qquad \forall k \geq 0,
$$
\n
$$
(2-6)
$$

are valid where the scalar sequence  $\{v_k\}$ , being non-decreasing, converges to the minimal solution  $v_*$  of (2-5).

In the above stated theorem, the central role is played by the condition that there exists a solution  $v_*$  of the equation [\( 2-5](#page-2-0)). To make contact with the standard formulation of the NK theorem, it is sufficient to observe that the NK theorem may be reformulated in terms of the similar requirement. If  $2l\eta \leq 1$ , the NK fixed point equation  $\eta - \nu + l\nu^2 / 2 = 0$  has a solution  $\nu_* > \eta$  (for  $\eta$  > 0) where *l* denotes the Lipschitz constant in the affine-invariant framework (e.g., see [4]).

Let us also note without proof that a minor modification of the arguments in [4] enables one to verify that the considered in the theorem solution  $x_*$  is unique in the open ball  $B(x_0, v_*)$ .

## **2.3 Proof of the theorem**

Assume that there is such a majorizing scalar sequence  $\{v_k\}$  that converges to a limiting point  $\nu_* = \lim_{k \to \infty} \nu_k \le R$  and facilitates the upper bound (2-1) once  $\{x_k\} \in \overline{B}(x_0, R) \subset D_F$ . Then, the standard arguments [2] verify that the Newton sequence  $\{x_k\}$  complies with the Cauchy criterion and, therefore, converges to a limiting point  $x_*$  so that the condition (2-6) is valid and  $x_*, x_k \in \overline{B}(x_0, v_*)$  for  $\forall k \ge 0$ . In turn, the continuity of  $F(x)$  implies that  $F(x_*) = 0$ .

Once the sequence  $\{v_k\}$  is generated by the recurrence relation (2-2), the convenient sufficient conditions for the existence of the limiting point  $v_*$  are proposed by Kantorovich [1]. The proof of Theorem 1 of Section 2 in Chapter XVIII of  $\lceil 1 \rceil$  includes the verification of the following useful Lemma. Presume that the fixed point equation [\( 2-5](#page-2-0) ) has a minimal solution  $\nu_* \in [0, R]$  and the continuous function  $\psi(\nu) \ge 0$  is non-decreasing when  $\nu \in [0, R]$ . As long as  $\{v_k\}$  is generated by (2-2) with the considered initial conditions, this sequence is nondecreasing and converges to  $v_*$  so that  $\{v_k\} \in [0, R]$ .

Next, let us prove that, under the set of the restrictions  $\Upsilon$ , the majorization condition  $(2-1)$  is valid where the continuous non-decreasing function  $\psi(v) \ge 0$  is defined on [0, R] by [\( 2-3](#page-1-1) ). As a starting point of the proof, we use the Rheinboldt majorization technique [\[ 2](#page-5-1) ]. It introduces such a function  $\Omega(t, s, r) \ge 0$  that, being continuous and non-decreasing in  $t, s, r \ge 0$ , *optimally* implements the upper bound  $d(T^2)$ that, being continuous and non-decreasing in  $t, s, r \ge 0$ <br> $d(T^2x, Tx) \le \Omega(d(Tx, x), d(Tx, x_0), d(x, x_0))$  presuming that  $\forall x, Tx \in D \subset D_F$ . Here,  $d(x, y) = ||x - y||$ ,  $T(x) = Tx$ ,  $T(T(x)) = T^2x$ ,  $x_0$  denotes the starting point of the Newton sequence  $\{x_k\} \in D \subset D_F$  defined by (1-1) and one identifies  $D = \overline{B}(x_0, R)$ . In turn, the latter upper bound is converted into the recurrence relation [2]:

$$
u_{k+1} - u_k = \Omega(u_k - u_{k-1}, u_k, u_{k-1}) \qquad , \qquad \forall k \ge 1.
$$
 (2-7)

<span id="page-3-0"></span>Given the initial conditions  $u_0 = v_0 = 0$  and  $u_1 = v_1 = \eta > 0$ , it defines such scalar nondecreasing sequence  $\{u_k\}$  that facilitates the estimate  $||x_{k+1} - x_k|| \le u_{k+1} - u_k$  for  $\forall k \ge 0$ .

As [\( 2-7](#page-3-0) ) is generically too complex to be analyzed exactly, we propose the following general approach. Given the upper bound  $\Omega(\cdot)$ , the idea is to *optimally* replace  $\Omega(\cdot)$  by such its majorant  $\overline{\Omega}(\cdot)$  that admits the so-called «first-integral» representation [2] (Section 12.5):

<span id="page-3-1"></span>
$$
\Omega(t, t+r, r) \le \overline{\Omega}(t, t+r, r) = \psi(t+r) - \psi(r) \qquad , \qquad \forall r \in [0, R] \quad , \quad \forall t \in [0, R-r], \qquad (2-8)
$$

where  $\psi(r)$  is restricted to be a continuous and non-decreasing function. It is to be compared with the more ambitious suggestion [2] to try to derive such representation directly for  $\Omega(\cdot)$ that is *not* implemented so far for the Newton method under relaxed continuity assumptions.

Given (2-8) and (2-2) together with the monotonicity of  $\Omega(\cdot)$ , it is straightforward to justify by induction that  $u_k - u_{k-1} \le v_k - v_{k-1}$  for  $\forall k \ge 1$  if  $u_0 = v_0 = 0$  and  $u_1 = v_1 = \psi(0) = \eta \ge 0$ . In turn, combining it with the estimate  $||x_{k+1} - x_k|| \le u_{k+1} - u_k$ , one reproduces (2-1). As for the justification of the inequality  $u_k - u_{k-1} \le v_k - v_{k-1}$ , note first that it is obviously valid for  $k = 1$ . Assume that this equality holds true for  $1 \le k \le q$  which implies that  $u_k \le v_k$  when  $1 \le k \le q$ . Assume that this equality holds true for  $1 \le k \le q$  which implies that  $u_k \le v_k$  when  $1 \le k \le q$ .<br>Then,  $u_{q+1} - u_q \le \Omega(v_q - v_{q-1}, v_q, v_{q-1}) \le \psi(v_q) - \psi(v_{q-1}) = v_{q+1} - v_q$  that completes the justification.

It remains to optimally reconstruct the explicit form of the relevant pattern of  $\Omega(\cdot)$  and then derive such its majorant  $\overline{\Omega}(\cdot)$  that, complying with (2-8), verifies the expression (2-3) for  $\psi(\nu)$ . The standard transformations lead to

<span id="page-3-2"></span>
$$
\Omega(t,s,r) = \gamma(s) \cdot \left( \int_{r}^{r+t} \omega_0(l) dl + \omega_0(r) t \right) , \qquad \gamma(s) = (1 - \omega_0(s))^{-1} , \qquad (2-9)
$$

that, in fact, is the particular case of the general expression obtained in [\[ 3](#page-5-2) ] for a generic Newton-like method (see the equation  $(34)$  where one is to identify  $a = 0$  and  $w_1(r) = w_0(r) = \omega_0(r)$ ). Thus defined  $\Omega(t, s, r)$  is continuous and non-decreasing in t, s, r due to the continuity and monotonicity of  $\omega_0(r)$  and the verified below relation  $\omega_0(\nu_*) \in ]0,1[$ . In turn, Newton-like method (see the equation (34) where one is to identify  $a = 0$  and  $w_1(r) = w_0(r) = \omega_0(r)$ ). Thus defined  $\Omega(t, s, r)$  is continuous and non-decreasing the continuity and monotonicity of  $\omega_0(r)$  and the verified be

$$
w_1(r) = w_0(r) = \omega_0(r)
$$
. Thus defined  $\Omega(t, s, r)$  is continuous and non-decreasing in  $t, s, r$  due to the continuity and monotonicity of  $\omega_0(r)$  and the verified below relation  $\omega_0(\nu_*) \in ]0,1[$ . In turn,  $\Omega(t, s, r) \le 2\gamma(s) \int_{r}^{r+t} \omega_0(l)dl \le 2\gamma(s) \int_{0}^{r+t} \omega_0(l)dl - 2\gamma(r) \int_{0}^{r} \omega_0(l)dl = \overline{\Omega}(t, s, r)$ , (2-10)

<span id="page-4-0"></span>where it is presumed that  $r \leq s$  (with  $\gamma(r) \leq \gamma(s)$ ) in accordance with the subsequent identification  $r + t = s$ . In sum, (2-10) and (2-8) lead to (2-3).

Finally, to prove that thus defined functions  $\psi(v)$  and  $\Omega(t, s, r)$  are non-negative, it is sufficient to demonstrate that  $\omega_0(\nu_*) \in ]0,1[$  which also implies that  $\| (F'(x_k))^{-1}$  $\| (F'(x_k))^{-1} F'(x_0) \| < \infty$  for  $\forall k \ge 1$  (i.e., the sequence  $\{x_k\}$  is well-defined). In turn, this property of  $\omega_0(\nu_*)$  directly follows from the pattern (2-3) of (2-5) and the imposed restriction  $v_* \in ]\eta, R]$ .

## **3 Application to the case of the Lipschitz continuous operator**  $F'(x)$

When  $F'(x)$  is Lipschitz continuous, the upper bound (2-4) is introduced with  $\omega_0(r) = l_0 r$ where  $l_0$  is the center-Lipschitz constant [4]. In this case, the implementation (2-3) of (2-5 ) is reduced to the quadratic in  $\upsilon$  equation  $2l_0\upsilon^2 - (1 + l_0\eta)\upsilon + \eta = 0$  with the discriminant  $D(l_0\eta) = (1 + l_0\eta)^2 - 8l_0\eta = (l_0\eta)^2 - 6l_0\eta + 1 \ge 0$ . For the latter equation to possess a solution, the  $2^2-8l_0\eta = (l_0\eta)^2-6l_0\eta+1 \ge 0$ . For the latter equation to possess a solution, the secondary quadratic function  $D(\lambda)$  should be non-negative which constrains that  $l_0 \eta \leq 3 - 2\sqrt{2}$ (i.e.,  $\eta \le \eta_{\text{max}} = (3 - 2\sqrt{2})/l_0$ ) where  $\gamma_* = 3 - 2\sqrt{2} \approx 0.171$  is the minimal root of the equation  $D(\gamma) = 0$ . It is straightforward to demonstrate that the corresponding minimal solution  $\nu_*$ satisfies the required restriction  $v_* > \eta$  when  $\eta > 0$ .

The Kantorovich condition [1] reads  $l\eta \le 0.5$  ( $\eta \le \tilde{\eta}_{max} = (2l)^{-1}$ ) where *l* stands for the standard Lipschitz constant [\[ 4](#page-5-3) ] formulated in the affine-invariant way. The above condition standard Lipschitz constant [4] formulated in the affine-invariant way. The above condition  $l_0 \eta \leq 3 - 2\sqrt{2}$  is weaker if  $l_0 / l < 2(3 - 2\sqrt{2})$  when  $\eta_{\text{max}} = (l_0)^{-1}(3 - 2\sqrt{2}) > \tilde{\eta}_{\text{max}} = (2l)^{-1}$ . It is noteworthy that the critical value  $6-4\sqrt{2} \approx 0,343$  of the ratio  $l_0/l$  is fairly moderate because it may often be that  $l_0 / l \ll 1$  (see [4]). The price to pay for this advantage is the linear (rather than quadratic as in the Kantorovich case) convergence of the majorizing sequence  $\{v_k\}$ .

# **4 Comparison with the Argyros convergence theorem**

The proposal of [\[ 4](#page-5-3) ] (Section 2.7) derives certain approximate estimates starting directly from the bound  $\Omega(\cdot)$  of (2-9). The single condition of the existence of the minimal solution  $\nu_* \in ]\eta, R]$  of the fixed point equation (2-5) is replaced in [4] by the <u>two</u> different restrictions. The first one requires the existence of a solution  $r_0$  of equation  $f(r) = r$  and, in our terms,

<span id="page-5-4"></span>
$$
f(v) = (1 - \omega_0(v))^{-1} \cdot \left( \int_0^v \omega_0(l) dl + \omega_0(v) v + \eta \right) \ge \psi(v) \qquad , \qquad v \in [0, R].
$$
 (4-1)

In the derivation of the inequality  $f(v) \ge \psi(v)$ , the monotonicity of  $\omega_0(v)$  is taken into where  $\psi(\nu)$  is introduced by (2-3). In particular,  $f(\nu) > \psi(\nu)$  for  $\forall \nu \in ]0, R]$  when  $\eta > 0$  and  $\omega_0(\nu)$ is strictly increasing (with  $\omega_0(v) > 0$  for  $v > 0$ ). As a result, compared to the first restriction of [4], our condition is weaker leading to a finer localization of the limiting point  $x_* \in \overline{B}(x_0, v_*)$ and a larger maximal admissible value  $\eta_{\text{max}}$  of the upper bound  $\eta \geq ||(F'(x_0))^{-1}F(x_0)||$ restricting the choice of the initial point  $x_0$ . As for the second restriction of  $[4]$ , in our terms it reads  $q(r_0) = 2\omega_0(r_0)/(1-\omega_0(r_0))$  <1 that is *not* generically necessitated by the first restriction.

In particular, given (4-1), the counterpart of our condition  $l_0 \eta \leq 3 - 2\sqrt{2} \approx 0.171$  may be obtained in the form  $l_0 \eta \le 0, 1$  while in [3] it is argued that  $l_0 \eta \le (2 - \sqrt{3})/2 \approx 0,134$  using a slightly different technique. Both these upper bounds are less favorable than our bound.

# **5 References**

- <span id="page-5-0"></span>**[ 1 ]** L.V. Kantorovich, G.P. Akilov, Functional Analysis, Pergamon Press, Oxford, 1982.
- <span id="page-5-1"></span>**[ 2 ]** J.M. Ortega, W.C. Rheinboldt, Iterative Solution of Nonlinear Equations in Several Variables, Academic Press, New York, 2000.
- <span id="page-5-2"></span>**[ 3 ]** I. K. Argyros, Relaxing the convergence conditions for Newton-like methods, J. Appl. Math. and Computing, 21 (2006), No. 1 - 2, 119 – 126.
- <span id="page-5-3"></span>**[ 4 ]** I. K. Argyros, Convergence and Applications of Newton-Type Iterations, Springer, New York, 2008.