Efficient Evaluation of Natural Stochastic Policies in Offline Reinforcement Learning

Nathan Kallus
Department of Operations Research and Information Engineering
and Cornell Tech Cornell University

Masatoshi Uehara *
Department of Computer Science and Cornell Tech
Cornell University

Abstract

We study the efficient off-policy evaluation of natural stochastic policies, which are defined in terms of deviations from the behavior policy. This is a departure from the literature on off-policy evaluation where most work consider the evaluation of explicitly specified policies. Crucially, offline reinforcement learning with natural stochastic policies can help alleviate issues of weak overlap, lead to policies that build upon current practice, and improve policies' implementability in practice. Compared with the classic case of a pre-specified evaluation policy, when evaluating natural stochastic policies, the efficiency bound, which measures the best-achievable estimation error, is inflated since the evaluation policy itself is unknown. In this paper we derive the efficiency bounds of two major types of natural stochastic policies: tilting policies and modified treatment policies. We then propose efficient nonparametric estimators that attain the efficiency bounds under very lax conditions. These also enjoy a (partial) double robustness property.

Keywords: Off-policy evaluation, Dynamic treatment regime, Reinforcement Learning, Semiparametric Inference, Double Robustness,

^{*}Corresponding author mu223@cornell.edu

1 Introduction

In many emerging application domains for reinforcement learning (RL), exploration is highly limited and simulation unreliable, such as in healthcare (Kosorok and Moodie, 2015; Gottesman et al., 2019). In these domains, we must use offline RL, where we evaluate and learn new sequential decision policies from existing observational data (Murphy, 2003; Robins, 2004; Zhang et al., 2013; Bibaut et al., 2019; Kallus and Uehara, 2019b; Nachum et al., 2019). A key task in offline RL is that of off-policy evaluation (OPE), in which we evaluate a new policy from data logged by another behavior policy. In many applications such as mobile health, the horizon is often long and possibly infinite (Boruvka et al., 2018; Luckett et al., 2018; Liao et al., 2020; Shi et al., 2020). In such settings, the naïve sequential importance sampling estimator suffers from the curse of horizon (Liu et al., 2018) in the sense that its mean squared error (MSE) grows exponentially in the horizon. Recent work in OPE (Kallus and Uehara, 2020a, 2019a) has shown how efficiently leveraging problem structure, such as Markovianness and time-homogeneity, can significantly improve OPE and address issues such as the curse of horizon.

In most of the literature on OPE, including the above, the policy to be evaluated is *pre-specified*, that is, it is a given and known function from states to a distribution over actions. In a departure from this, in this paper we consider the evaluation of *natural stochastic policies*, which may depend on the *natural* value of the action, that is, the treatment that is observed in the data without intervention (Muñoz and Van Der Laan, 2012; Shpitser and Pearl, 2012; Haneuse and Rotnitzky, 2013; Young et al., 2014, 2019; Richardson and Robins, 2013; Díaz and van der Laan, 2018). Specifically, we consider policies defined as *deviations* from the behavior policy that generated the observed data.

There are two primary advantages to natural stochastic policies. A first advantage is *imple*mentability. Subjects are often unable or reluctant to undertake an assigned treatment if the deviation from the treatment they would have naturally undertaken is large. For example, consider intervening on leisure-time physical activity to reduce mortality among the elderly (as in Díaz and van der Laan, 2018). An evaluation policy assigning $a + \delta$ minutes of weekly activity to an individual whose current physical activity level is a (i.e., the natural value) would be a realistic intervention for small to moderate δ . On the other hand, evaluation policies assigning any arbitrary level of physical activity level ignoring the current level of physical activity is unrealistic and rarely implementable. Another example is intervening on air pollution levels to improve the health of children (as in Díaz and van Der Laan, 2013). A possible evaluation policy is enforcing the pollution levels below a certain cutoff point if the observed pollution level (i.e., the natural value) exceeds the threshold. A second advantage is that we can relax or more easily satisfy the positivity assumption, which requires some overlap between the evaluation and behavior policies and is fundamentally necessary for OPE. Often, we cannot know a priori whether the positivity assumption is satisfied for a given intervention in an observational study or how good is the overlap. We can, however, easily consider policies that only deviate slightly from the behavior policy, ensuring a good overlap and reliable evaluation by design.

In this paper, we derive efficiency bounds and develop efficient estimators for two major types of natural stochastic policies: tilting policies and modified treatment policies. The efficiency bounds quantify the statistical limits to evaluation by showing what is the best-achievable MSE asymptotically. We study how much the efficiency bounds inflate in comparison with the case of a pre-specified evaluation policy (see Table 1). Our central message is that the order of the efficiency bound is surprisingly the same as in the case of a pre-specified policy. Importantly, this implies the curse of the horizon is avoidable since the horizon dependence is polynomial, while good overlap can be achieved by design. We then develop efficient estimation methods achieving these efficiency bounds under lax conditions. We also demonstrate how methods that are efficient for pre-specified policies (Kallus and Uehara, 2020a, 2019a) break. Unlike the pre-specified case, the estimator has an interesting partial double robustness property, which is a different from the usual double robustness (Robins et al., 1999; Jiang and Li, 2016; Kallus and Uehara, 2020a), due to a special bias structure.

	Form of $\pi_t^{\mathrm{e}}(a_t \mid s_t)$	Inflation of the bound compared with (a)	Order of the bound
(a) Pre-specified	Explicit function	0	$CC'R_{\max}^2H^2$
(b) Tilting	$\frac{u_t(a_t)\pi_t^{\mathrm{b}}(a_t s_t)}{\int u_t(\tilde{a}_t)\pi_t^{\mathrm{b}}(\tilde{a}_t s_t)\mathrm{d}\tilde{a}_t}$	$\sum_{t=0}^{H} \mathbb{E}[\operatorname{var}[\mu_t q_t \mid s_t]]$	$CC'R_{\max}^2H^2$
(c) Modified treatment	$\pi_t^{\mathrm{b}}(\tilde{\tau}_t(s_t, a_t) \mid s_t) \tilde{\tau}_t'(s_t, a_t)$	$\sum_{t=0}^{H} \mathbb{E}[\mu_t^2 \operatorname{var}[q_{t+1}^{\tau} \mid s_{t+1}]]$	$CC'R_{\max}^2H^2$

The article is organized as follows. Section 2 sets up the problem and definitions and discusses related literature. Section 3 establishes the efficiency bounds for OPE of natural stochastic policies in time-varying Markov decision processes (TMDPs). Section 4 develops efficient estimators. Section 5 provides the efficiency bound and efficient estimators when we remove the Markovian assumption. Section 6 provides the efficiency bound and efficient estimators when we additionally impose time invariance on the decision process. Section 7 studies the performance of our approach empirically. Section 8 provides concluding remarks on our findings.

2 Setup and Background

We setup the problem and notation and summarize the literature on OPE and natural stochastic policies.

2.1 Problem Setup and Definitions

Consider an H-long time-varying Markov decision process (TMDP), with states $s_t \in \mathcal{S}_t$, actions $a_t \in \mathcal{A}_t$, rewards $r_t \in \mathbb{R}$, initial state distribution $p_1(s_1)$, transition distributions $p_{t+1}(s_{t+1} \mid s_t, a_t)$, and reward distributions $p_t(r_t \mid s_t, a_t)$, for $t = 1, \ldots, H$. A policy $(\pi_t(a_t \mid s_t))_{t \leq H}$ induces a distribution over trajectories $\mathcal{T} = (s_1, a_1, r_1, \ldots, s_T, a_H, r_H, s_{H+1})$:

$$p_{\pi}(\mathcal{T}) = p_1(s_1) \prod_{t=1}^{H} \pi_t(a_t \mid s_t) p_t(r_t \mid s_t, a_t) p_{t+1}(s_{t+1} \mid s_t, a_t). \tag{1}$$

Given an evaluation policy π^e , which we consider as unknown in this paper, we are interested in its value, $J = \mathbb{E}_{p_{\pi^e}} \left[\sum_{t=1}^{H} r_t \right]$, where the expectation is taken with respect to (w.r.t.) the density induced by the evaluation policy, p_{π^e} . In the *off-policy* setting, our data consists of trajectory observations from some fixed policy, π^b , known as the *behavior policy*:

$$\mathcal{T}^{(1)}, \dots, \mathcal{T}^{(n)} \sim p_{\pi^b}, \quad \mathcal{T}^{(i)} = (S_1^{(i)}, A_1^{(i)}, R_1^{(i)}, \dots, S_H^{(i)}, A_H^{(i)}, R_H^{(i)}).$$
 (Off-policy data)

In observational studies, as we consider herein, $\pi^{\rm b}$ is *unknown*, and the observed action $A_j^{(i)}$ is considered the natural value of the action in the sense that it is the one naturally observed in the absence of our intervention. Our goal is to estimate J from the observed data $\{\mathcal{T}^{(i)}\}_{i=1}^n$.

absence of our intervention. Our goal is to estimate J from the observed data $\{\mathcal{T}^{(i)}\}_{i=1}^n$. We define the following variables. Let $q_t = \mathbb{E}_{p_{\pi^e}} \left[\sum_{k=t}^H r_t \mid s_t, a_t \right], \ v_t = \mathbb{E}_{p_{\pi^e}} \left[\sum_{k=t}^H r_t \mid s_t \right]$ be the q- and v-functions for π^e . Further, let the instantaneous, cumulative, marginal state, and marginal state-action density ratios be, respectively, $\eta_t = \frac{\pi_t^e(a_t|s_t)}{\pi_t^b(a_t|s_t)}, \ \lambda_t = \prod_{k=1}^t \eta_k, \ w_t = \frac{p_{\pi^e}(s_t)}{p_{\pi^b}(s_t)}, \ \mu_t = \eta_t w_t,$

¹Although we do not explicitly use a counterfactual notation this is the same as the counterfactual value of following π^e instead of π^b if we had used potential outcomes and assumed the usual sequential ignorability and consistency assumptions (Ertefaie and Strawderman, 2018; Luckett et al., 2018).

where $p_{\pi}(s_t)$ is a marginal density at s_t under p_{π} . We assume throughout the paper that $0 \le r_t \le R_{\max}$, $\eta_t \le C$, $w_t \le C'$, $\forall t \le H$.

Given trajectory data, $\mathcal{T}^{(1)}, \ldots, \mathcal{T}^{(n)}$, we define the *empirical expectation* as $\mathbb{P}_n f = \frac{1}{n} \sum_{i=1}^n f(\mathcal{T}^{(i)})$. Unless otherwise noted, all expectations, variances, and probabilities are w.r.t. p_{π^b} . Define the L_2 norm by $||f||_2 = {\mathbb{E}[f^2(\mathcal{T})]}^{1/2}$.

2.2 Natural Stochastic Policies

In OPE, π^e is often pre-specified. Our focus is instead the case where π^e depends on the natural value of the treatment in an observational study. Importantly, in this setting, both π^e and π^b are unknown. Natural stochastic policies are widely studied in the non-sequential (bandit) setting where H=1 (Muñoz and Van Der Laan, 2012; Haneuse and Rotnitzky, 2013). However, it has not been extensively studied in the longitudinal (RL) setting but for a few exceptions. Kennedy (2019) considers OPE with binary actions under a tilting policy in an NMDP (non-Markov decision process). In comparison, we allow actions to be arbitrary and focus on the Markovian setting that is central to RL. Young et al. (2014) considers OPE under a modified treatment policy in an NMDP using a parametric approach. In comparison, our methods are nonparametric and globally efficient, and we focus on the Markovian setting common in RL.

In this paper, we consider two types of natural stochastic policies: modified treatment policies and tilting polices. These constructions are inspired by previous work focusing on the bandit and NMDP settings (Díaz and Hejazi, 2020; Muñoz and Van Der Laan, 2012; Haneuse and Rotnitzky, 2013).

Definition 1 (Tilting policy). A tilting policy is specified by $u_t: \mathcal{A}_t \to \mathbb{R}$ and defined as

$$\pi_t^{\mathrm{e}}(a_t \mid s_t) = u_t(a_t)\pi_t^{\mathrm{b}}(a_t \mid s_t) / \int u_t(\tilde{a}_t)\pi_t^{\mathrm{b}}(\tilde{a}_t \mid s_t) \mathrm{d}\tilde{a}_t. \tag{2}$$

Tilting policies tilt the behavior policy slightly toward actions with higher values of u_t . For example, for binary action, letting $u_t(1) = \delta$, $u_t(0) = 1$ yields

$$\pi_t^{\mathrm{e}}(a_t \mid s_t) = \mathrm{I}(a_t = 1) \frac{\delta \pi_t^{\mathrm{b}}(1 \mid s_t)}{1 + (\delta - 1)\pi_t^{\mathrm{b}}(1 \mid s_t)} + \mathrm{I}(a_t = 0) \frac{\delta^{-1} \pi_t^{\mathrm{b}}(0 \mid s_t)}{1 + (\delta^{-1} - 1)\pi_t^{\mathrm{b}}(0 \mid s_t)},\tag{3}$$

as considered by Kennedy (2019) in the binary-action NMDP setting. For $\delta=1$ we get $\pi^{\rm e}=\pi^{\rm b}$; as δ shrinks, we tilt toward action 0; and, as δ grows, we tilt toward action 1. The parameter δ directly controls the amount of overlap; specifically $\pi_t^{\rm e}(a_t\mid s_t)/\pi_t^{\rm b}(a_t\mid s_t) \leq \max(\delta,\delta^{-1})$. For the general case in Definition 1, we have that $\pi_t^{\rm e}(a_t\mid s_t)/\pi_t^{\rm b}(a_t\mid s_t) \leq \max_{\tilde{a}_t} u_t(\tilde{a}_t)/\min_{\tilde{a}_t} u_t(\tilde{a}_t)$ so that the variation in u_t can directly control the overlap. Tilting policies ensure that $\pi_t^{\rm e}(\cdot\mid s_t)$ is absolutely continuous w.r.t. $\pi_t^{\rm b}(\cdot\mid s_t)$ so that the density ratio always exists. In contrast, if $\pi_t^{\rm e}$ is pre-specified and $\pi_t^{\rm b}$ is unknown, we cannot always ensure that the density ratio exists, let alone is bounded.

Definition 2 (Modified treatment policy). A modified treatment policy is specified by the maps $\tau_t : \mathcal{S}_t \times \mathcal{A}_t \to \mathcal{A}_t$ and assigns the action $\tau_t(s_t, a_t)$ in state s_t when the natural action value is a_t .

Notice that the modified treatment policy is the same value J as the evaluation policy defined by letting $\pi_t^{\rm e}(\cdot\mid s_t)$ be the distribution of $a_t=\tau_t(s_t,\tilde{a}_t)$ under $\tilde{a}_t\sim\pi_t^{\rm b}(\cdot\mid s_t)$. For the purpose of OPE, we can therefore equivalently define the modified treatment policy as this transformation of $\pi^{\rm b}$. For example, if for each s_t , $\tau_t(s_t,\cdot)$ has a differentiable inverse $\tilde{\tau}_t(s_t,\cdot)$, then $\pi_t^{\rm e}(a_t\mid s_t)=\pi_t^{\rm b}(\tilde{\tau}_t(s_t,a_t)\mid s_t)\tilde{\tau}_t'(s_t,a_t)$, where ' denotes a differentiation w.r.t a_t . The simplest example of a modified treatment policy is $\tau_t(s_t,a_t)=a_t+b_t(s_t)$ for some function $b_t(s_t)$, for which $\pi_t^{\rm e}(a_t|s_t)=\pi_t^{\rm b}(a_t-b_t(s_t)\mid s_t)$. The function $b_t(s_t)$ quantifies the deviation from the natural value. Keeping $b_t(s_t)$ small ensures implementability.

2.3 Off-Policy Evaluation

Step-wise importance sampling (IS; Precup et al., 2000) and direct estimation of q-functions (DM; Munos and Szepesvári, 2008; Ernst et al., 2005) are two common approaches for OPE. However, the former is known to suffer from the high variance and the latter from model misspecification. To alleviate this, the doubly robust estimate combines the two (Murphy et al., 2001; Jiang and Li, 2016; Thomas and Brunskill, 2016). However, the asymptotic MSE of these can still grow exponentially in the horizon. Kallus and Uehara (2020a) show that the efficiency bound in the TMDP case is actually polynomial in H, $\mathfrak{G}(CC'R_{\max}^2H^2/n)$, and give an estimator achieving it by combining marginalized IS (Xie et al., 2019) and q-modeling using cross-fold estimation. When we additionally assume time invariance on TMDPs, Kallus and Uehara (2019a) show an orders-smaller efficiency bound and develop an efficient estimator leveraging time invariance.

All of the above methods focus on the case where $\pi^{\rm e}$ is given explicitly. If the behavior policy is known, then natural stochastic policies can also be regarded as given explicitly and these still apply. When $\pi^{\rm b}$ is unknown, as in observational studies, we can still operationalize these methods for evaluating natural stochastic policies by first estimating $\pi^{\rm b}$ from the data, plugging this into $\pi^{\rm e}$, and then treating $\pi^{\rm e}$ as specified by this estimate. However, this will fail to be efficient, as we discuss in Section 4. In fact, the efficiency bounds for evaluating natural stochastic policies are different than the pre-specified case.

3 Efficiency Bounds

In this section we calculate the efficiency bounds for evaluating natural stochastic policies in RL. We first briefly explain what the efficiency bound is (see van Der Laan and Robins, 2003 for more detail).

A fundamental question is what is the smallest-possible error we can hope to achieve in estimating J. In parametric models, the Cramér-Rao bound lower bounds the asymptotic MSE of all (regular) estimators. Our model, however, is nonparametric as it consists of all TMDP distributions, i.e., any choice for $p_t(r_t \mid s_t, a_t)$, $p_{t+1}(s_{t+1} \mid s_t, a_t)$, and $\pi_t(a_t \mid s_t)$ in Eq. (1). Semiparametric theory gives an answer to this question by extending the notion of a Cramér-Rao lower bound to nonparametric models. We first informally state the key property of the efficient influence function (EIF) from semiparametric theory in our setting, i.e., the estimand is J and the model is all TMDP distributions denoted by $\mathcal{M}_{\text{TMDP}}$.

Theorem 1 (Informal description of Theorem 25.20 of van der Vaart (1998)). The EIF $\phi(\mathcal{T})$ is the gradient of J regarding $\mathcal{M}_{\text{TMDP}}$ that has the smallest L_2 norm, and it satisfies that, for any estimator \hat{J} of J that is regular w.r.t. $\mathcal{M}_{\text{TMDP}}$, $\text{AMSE}[\hat{J}] \geq \text{var}[\phi(\mathcal{T})]$, where $\text{AMSE}[\hat{J}]$ is the second moment of the limiting distribution of $\sqrt{n}(\hat{J}-J)$.

A regular estimator is any whose limiting distribution is insensitive to perturbations of order $\mathfrak{G}(1/\sqrt{n})$ to the data-generating process p_{π^b} that remain in $\mathcal{M}_{\text{TMDP}}$ (i.e., keep it a TMDP distribution). That is, a regular estimator is one that "works" for all problem instances in the model, and in this sense the result crucially depends on the model being considered. Here, $\text{var}[\phi(\mathcal{T})]$ is called the *efficiency bound* as it is a lower bound on the asymptotic MSE of all regular estimators, which is a very general class. This class is so general that in fact this also implies local minimax bound for *all* estimators (see van der Vaart, 1998, Theorem 25.21).

3.1 Tilting Policies

In the next result we calculate the EIF and efficiency bound for OPE of tiling policies.

Theorem 2. Let π^e be as in Definition 1. Then the EIF and efficiency bound of J w.r.t. the model \mathcal{M}_{TMDP} are, respectively,

$$-J + \sum_{t=1}^{H} (\mu_t(r_t - v_t) + \mu_{t-1}v_t), \quad \Upsilon_{\text{TI1}} = \sum_{t=0}^{H} \mathbb{E}[\text{var}[\mu_t(r_t + v_{t+1}) \mid s_t]],$$

where $\mu_0 = 1, v_0 = r_0 = 0$. Moreover, Υ_{TI1} is upper bounded by $CC'R_{\text{max}}^2H^2$.

Note that the function u_t that specifies the tilting policy is implicit in the variables μ_t, v_t above, which depend on π_t^e . While the efficiency bound is larger than in the case of a pre-specified evaluation policy (Kallus and Uehara, 2020a), the overall order, $CC'R_{\text{max}}^2H^2$, is the same. This implies we can circumvent the curse of horizon by using and efficient estimator.

Remark 1 (Comparison to pre-specified evaluation policy). In a pre-specified evaluation policy case, Kallus and Uehara (2020a) show that the EIF and efficiency bound are, respectively,

$$-J + \sum_{t=1}^{H} (\mu_t(r_t - q_t) + \mu_{t-1}v_t), \quad \sum_{t=0}^{H} \mathbb{E}[\mu_t^2 \operatorname{var}[r_t + v_{t+1} \mid s_t, a_t]]. \tag{4}$$

Specifically, if we let $\pi^{\rm e}$ be as in Definition 1 and assume that $\pi^{\rm b}$ is *known* then this is the efficiency bound. Compared with this quantity, $\Upsilon_{\rm TI1}$ is larger by $\sum_{t=1}^{H} \mathbb{E}[{\rm var}[\mu_t q_t \mid s_t]]$.

Remark 2 (Non-Markovian Decision Processes). Kennedy (2019) provides the EIF for the binary-action tilting policy Eq. (3) under an NMDP. In comparison, our Theorem 2 handles the Markovian case relevant to RL as well as a general action space. In particular, as we will see in Section 5, OPE under the NMDP model necessarily suffers from the curse of horizon and therefore cannot handle long horizons. An NMDP model can actually be embedded in a TMDP model (but not vice versa) by including the whole state-action history up to time t in the state variable s_t . Using this transformation, Theorem 2 recovers the result of Kennedy (2019) as a special case. For further discussion of Kennedy (2019), refer to Section 5.

Remark 3 (Bandit case). When H = 1, the EIF is $\eta_1(r_1 - v_1(s_1)) + v_1(s_1)$.

3.2 Modified Treatment Policies

We next handle the case of modified treatment policies. Again, we will see that we can potentially circumvent the curse of horizon with an efficient estimator.

Theorem 3. Let π^e be as in Definition 2. Then the EIF and efficiency bound of J w.r.t. the model \mathcal{M}_{TMDP} are, respectively,

$$-J + \sum_{t=1}^{H} \mu_t(r_t - q_t) + \mu_{t-1}q_t^{\tau}, \quad \Upsilon_{\text{MO1}} = \sum_{t=0}^{H} \mathbb{E}[\mu_t^2 \text{var}[r_t + q_{t+1}^{\tau} \mid s_t, a_t]]$$
 (5)

where $q_t^{\tau}(s_t, a_t) = q_t(s_t, \tau_t(s_t, a_t))$. Moreover, Υ_{MO1} is upper bounded by $CC'R_{\text{max}}^2H^2$.

Remark 4 (Comparison to pre-specified evaluation policy). Compared with the efficiency bound for a pre-specified evaluation policy, Υ_{MO1} is larger by $\sum_{t=0}^{H} \mathbb{E}[\mu_t^2 \text{var}[q_{t+1}^{\tau} \mid s_{t+1}]]$.

Remark 5 (Bandit case). When H = 1, the EIF is $\eta_1(r_1 - q_1(s_1, a_1)) + q_1^{\tau}(s_1, a_1)$. This matches the results in Díaz and van Der Laan (2013); Díaz and van der Laan (2018).

Algorithm 1 Efficient Off-Policy Evaluation for Natural Stochastic Policies

Take a K-fold random partition of $\{1,\ldots,n\}=I_1\cup\cdots\cup I_K$ such that the size of each fold, $|I_k|$, is within 1 of n/K; set $\mathcal{U}_k=\{\mathcal{T}^{(i)}:i\in I_k\},\,\mathcal{L}_k=\{\mathcal{T}^{(i)}:i\notin I_k\}$ for $k\in\{1,\cdots,K\}$ do

Using only \mathcal{L}_{k} as data, construct nuisance estimators $\hat{w}_{t}^{(k)}$, $\hat{\pi}_{t}^{b,(k)}$, $\hat{q}_{t}^{(k)}$ for $t \leq H$ Set $\hat{\pi}_{t}^{e,(k)}(a_{t} \mid s_{t}) = u_{t}(a_{t})\hat{\pi}_{t}^{b,(k)}(a_{t} \mid s_{t})/\int u_{t}(\tilde{a}_{t})\hat{\pi}_{t}^{b,(k)}(\tilde{a}_{t} \mid s_{t})d\tilde{a}_{t}$ $\hat{\eta}_{t}^{(k)}(s_{t}, a_{t}) = \hat{\pi}_{t}^{e,(k)}(a_{t} \mid s_{t})/\hat{\pi}_{t}^{b,(k)}(a_{t} \mid s_{t}), \hat{v}_{t}^{(k)}(s_{t}) = \int \hat{q}_{t}^{(k)}(s_{t}, a_{t})\hat{\pi}_{t}^{e,(k)}(a_{t} \mid s_{t})da_{t}$ Set $\hat{J}_{k} = \frac{1}{|I_{k}|}\sum_{\mathcal{T} \in \mathcal{U}_{k}}\hat{\phi}^{(k)}(\mathcal{T})$, where

$$\hat{\phi}^{(k)}(\mathcal{T}) = \sum_{t=1}^{H} \hat{w}_{t}^{(k)}(s_{t}) \hat{\eta}_{t}^{(k)}(s_{t}, a_{t}) (r_{t} - \hat{v}_{t}^{(k)}(s_{t})) + \hat{w}_{t-1}^{(k)}(s_{t-1}) \hat{\eta}_{t-1}^{(k)}(s_{t-1}, a_{t-1}) \hat{v}_{t}^{(k)}(s_{t})$$
(6)

end for

Return $\hat{J}_{TI1} = \frac{1}{n} \sum_{k=1}^{K} |I_k| \hat{J}_k$

4 Efficient and (Partially) Doubly Robust Estimation

We next propose efficient estimators for evaluating natural stochastic policies based on the obtained EIFs. Since both EIFs have second-order bias w.r.t. nuisances, we can obtain efficient estimators by estimating nuisances under nonparametric rate conditions and plugging these into the EIFs with a sample-splitting cross-fitting technique (Zheng and van Der Laan, 2011; Chernozhukov et al., 2018).

4.1 Tilting Policies

We propose an estimator \hat{J}_{TI1} for tilting polices in Algorithm 1. This is a meta-algorithm given estimation procedures for the nuisances w_t, π_t^b, q_t , which we discuss how to estimate in Section 4.3. We next prove \hat{J}_{TI1} is efficient under nonparametric rate conditions on nuisance estimators, which crucially can be *slower* than $\mathfrak{G}_p(n^{-1/2})$ and do not require metric entropy conditions.

Theorem 4 (Efficiency). Suppose $\forall k \leq K, \forall j \leq H, \|\hat{w}_{j}^{(k)}(s_{j}) - w_{j}(s_{j})\|_{2} \leq \alpha_{1}, \|\hat{\pi}_{j}^{b,(k)}(a_{j} \mid s_{j}) - \pi_{j}^{b}(a_{j} \mid s_{j})\|_{2} \leq \alpha_{2}, \|\hat{q}_{j}^{(k)}(s_{j}, a_{j}) - q_{j}(s_{j}, a_{j})\|_{2} \leq \beta, \text{ where } \alpha_{1} = \mathbb{O}_{p}(n^{-1/4}), \ \alpha_{2} = \mathrm{o}_{p}(n^{-1/4}), \ \beta = \mathbb{O}_{p}(n^{-1/4}), \ \alpha_{1}\beta = \mathrm{o}_{p}(n^{-1/2}). \ Then, \sqrt{n}(\hat{J}_{\text{TI1}} - J) \xrightarrow{d} \mathcal{N}(0, \Upsilon_{\text{TI1}}).$

The result essentially follows by showing that $|\hat{J}_{\text{TII}} - J - \mathbb{P}_n[\phi(\mathcal{T})]| \leq \alpha_1 \alpha_2 + \alpha_1 \beta + \alpha_2 \beta + \alpha_2^2 + o_p(n^{-1/2})$, where $\phi(\mathcal{T})$ is the EIF. Under the above rate assumptions, the right-hand side is $o_p(n^{-1/2})$ and the result is immediately concluded from CLT. Notice that if we knew the behavior policy so that $\alpha_2 = 0$, this becomes simply $\alpha_1 \beta + o_p(n^{-1/2})$ and we recover the doubly robust structure of the pre-specified case (Kallus and Uehara, 2020a): the estimator is consistent if $either\ w_t$ or q_t is consistently estimated. In our setting, because of the term α_2^2 , the consistent estimation of π^b is always required to estimate J consistently. So, we have a partial double robustness in the sense that the estimator is consistent as long as π^b and either w or q are consistently estimated.

Theorem 5 (Partial double robustness). Suppose $\forall k \leq K, \forall j \leq H$, for some $w_j^{\dagger}, q_j^{\dagger}, \|\hat{w}_j^{(k)}(s_j) - w_j^{\dagger}(s_j)\|_2 = o_p(1), \|\hat{q}_j^{(k)}(s_j, a_j) - q_j^{\dagger}(s_j, a_j)\|_2 = o_p(1), \text{ and } \|\hat{\pi}_j^{b,(k)}(a_j|s_j) - \pi_j^b(a_j|s_j)\|_2 = o_p(1). \text{ As long as either } q_j^{\dagger} = q_j \text{ or } w_j^{\dagger} = w_j, \text{ we have } \hat{J}_{\text{TII}} \xrightarrow{p} J.$

Remark 6 (Comparison to Kallus and Uehara, 2020a). Since we have to estimate $\pi^{\rm b}$ (and hence $\pi^{\rm e}$) consistently anyway for our estimator to work, a careful reader might wonder whether we might

as well plug in the estimated π^e into estimators that are efficient for the pre-specified case such as Kallus and Uehara (2020a). Specifically, we could replace Eq. (6) in Algorithm 1 with

$$\phi^{(k)}(\mathcal{T}) = \sum_{t=1}^{H} \hat{w}_{t}^{(k)}(s_{t}) \hat{\eta}_{t}^{(k)}(s_{t}, a_{t}) (r_{t} - \hat{q}_{t}^{(k)}(s_{t}, a_{t})) + \hat{w}_{t-1}^{(k)}(s_{t-1}) \hat{\eta}_{t-1}^{(k)}(s_{t-1}, a_{t-1}) \hat{v}_{t}^{(k)}(s_{t}),$$

which corresponds to plugging our estimated nuisances into the EIF derived in Kallus and Uehara (2020a). However, this can fail to achieve a \sqrt{n} -convergence rate, let alone efficiency. Specifically, in Theorem 4, we used the fact that Eq. (6) has a second-order bias structure w.r.t. w_t, π_t^b, q_t to ensure the \sqrt{n} -consistency and efficiency. In contrast, the above does not have this structure. It only has such a structure when $\hat{v}_t^{(k)}$ is the integral of $\hat{q}_t^{(k)}$ with respect to the *true* π_t^e , which Kallus and Uehara (2020a) use to achieve efficiency, but that is not the case here.

Remark 7 (Estimation of v-functions). Although $\hat{q}_t^{(k)}$ does not explicitly appear in Eq. (6), we do need to estimate $\hat{q}_t^{(k)}$ first and then compute $\hat{v}_t^{(k)}$ based on it as in Algorithm 1, instead of directly estimating v_t . The reason is that we cannot generally say that Eq. (6) has a second-order bias structure w.r.t. $w_t, \pi_t^{\rm b}, v_t$. Therefore, the efficiency result would not be guaranteed. To achieve the efficiency, it is crucial to use the specific construction of $\hat{v}_t^{(k)}$ in Algorithm 1, which ensures a certain compatibility between the nuisance estimators, as they all use the same estimated behavior policy.

4.2 Modified Treatment Policies

We similarly define the estimator \hat{J}_{MO1} for the case of modified treatment policies by taking Algorithm 1 and (a) replacing $\hat{\pi}_t^{e,(k)}(a_t \mid s_t)$ by $\hat{\pi}_t^{e,(k)}(a_t \mid s_t) = \hat{\pi}_t^{b,(k)}(\tilde{\tau}_t(s_t, a_t) \mid s_t)\tilde{\tau}_t'(s_t, a_t)$ and (b) replacing Eq. (6) by

$$\hat{\phi}^{(k)}(\mathcal{T}) = \sum_{t=1}^{H} \hat{w}_{t}^{(k)}(s_{t}) \hat{\eta}_{t}^{(k)}(s_{t}, a_{t}) (r_{t} - \hat{q}_{t}^{(k)}(s_{t}, a_{t})) + \hat{w}_{t-1}^{(k)}(s_{t-1}) \hat{\eta}_{t-1}^{(k)}(s_{t-1}, a_{t-1}) \hat{q}_{t}^{(k)}(s_{t}, \tau_{t}(s_{t}, a_{t})).$$

We then have the following efficiency and (full) double robustness results.

Theorem 6 (Efficiency). Suppose $\forall k \leq K, \forall j \leq H, \|\hat{w}_{j}^{(k)}(s_{j}) - w_{j}(s_{j})\|_{2} \leq \alpha_{1}, \|\hat{\pi}_{j}^{b,(k)}(a_{j}|s_{j}) - \pi_{j}^{b,(k)}(a_{j}|s_{j})\|_{2} \leq \alpha_{2}, \|\hat{q}_{j}^{(k)}(s_{j}, a_{j}) - q_{j}(s_{j}, a_{j})\|_{2} \leq \beta \text{ where } (\alpha_{1} + \alpha_{2})\beta = o_{p}(n^{-1/2}), \max\{\alpha_{1}, \alpha_{2}, \beta\} = o_{p}(1). \text{ Then, } \sqrt{n}(\hat{J}_{\text{MO1}} - J) \xrightarrow{d} \mathcal{N}(0, \Upsilon_{\text{MO1}}).$

Theorem 7 (Double robustness). Assume $\forall k \leq K, \forall j \leq H$, for some $\pi_j^{b\dagger}, q_j^{\dagger}, w_j^{\dagger}, \|w_j^{(k)}(s_j) - w_j^{\dagger}(s_j)\|_2 = o_p(1), \|\hat{\pi}_j^b(a_j|s_j) - \pi_j^{b\dagger}(a_j|s_j)\|_2 = o_p(1), \|\hat{q}_j^{(k)}(s_j, a_j) - q_j^{\dagger}(s_j, a_j)\|_2 = o_p(1).$ Then as long as either $q_j^{\dagger} = q_j$ or $\pi_j^{b\dagger} = \pi_j^b, w_j^{\dagger} = w_j$, we have $\hat{J}_{\text{MO1}} \stackrel{p}{\longrightarrow} J$.

These theorems arise from the bias structure $|\hat{J}_{\text{MO1}} - J - \mathbb{P}_n[\phi(\mathcal{T})]| \leq (\alpha_1 + \alpha_2)\beta + o_p(n^{-1/2})$. The conditions on nuisance estimates in these theorems are weaker than the ones for tilting policies. Comparing Theorems 4 and 6, the condition in Theorem 6 is satisfied even if some of $\alpha_1, \alpha_2, \beta$ are slower than $o_p(n^{-1/4})$. Comparing Theorems 5 and 7, the condition in Theorem 7 is satisfied even if the behavior policy model is misspecified. The intuitive reason is that for a modified treatment policy, J can be specified in a form not depending on π^b , while this is not true for tilting policies.

4.3 Nuisance Estimation

Our estimators for both types of stochastic policies require that we estimate π_t^b, w_t, q_t at some slow rate. Here we discuss some standard ways to estimate these nuisance functions. We focus on the case of tilting policies for brevity.

First of all, estimating π_t^b amounts to fitting a probabilistic classification model in the case of finitely many actions and to conditional density estimation in the case of continuous actions. Once we fit π_t^b , we also immediately have an estimate of π_t^e . We can then use standard methods for estimating w_t and q_t that assume π_t^e is given by plugging in our estimate for it as follows. Generally speaking, if the estimate for q_t or w_t would have had some convergence rate r_n if π_t^e were given exactly, then this rate does not deteriorate as long as the plugged-in estimate for π_t^e also has rate r_n . For all of our nuisance estimators, we do not require any metric entropy conditions.

Estimation of q-functions In the tabular case, a model-based approach is the most common way to estimate q-functions from off-policy data. In the non-tabular case, we have to rely on some function approximation. The key equation to derive these methods is the Bellman equation:

$$q_t(s_t, a_t) = \mathbb{E}[r_t + q_{t+1}(s_{t+1}, \pi^e) \mid s_t, a_t].$$

where $q_t(s_t, \pi) = \int q_t(s_t, a_t)\pi(a_t \mid s_t)da_t$. One of the most common ways to operationalize this is using fitted q-iteration (Antos et al., 2008; Le et al., 2019; Duan and Wang, 2020; expressed here using an estimated evaluation policy, $\hat{\pi}^e$): set $\hat{q}_{H+1} \equiv 0$, and for t = H, ..., 1 estimate \hat{q}_t by regressing $r_t + \hat{q}_{t+1}(s_{t+1}, \hat{\pi}^e)$ onto s_t, a_t using any given regression method. Note that the regressions can also be replaced with Z-estimation (Ueno et al., 2011; Luckett et al., 2018) based on the moment equations:

$$\mathbb{E}[g_t(s_t, a_t) (r_t + q_{t+1}(s_{t+1}, \pi^e) - q_t(s_t, a_t))] = 0, \quad \forall g_t, t \le H.$$

When we let q_t and g_t be the class of linear functions, this leads to the LSTD method (Lagoudakis and Parr, 2004).

Estimation of Marginal Density Ratios In the tabular case, a model-based approach (Yin and Wang, 2020) would be a competitive way to estimate marginal density ratios:

$$\hat{w}_t(s_t) = \frac{1}{\hat{p}_{\pi_t^{\rm b}}(s_t)} \int \hat{p}_t(s_t|s_{t-1}, a_{t-1}) \prod_{k=0}^{t-1} (\pi_k^{\rm e}(a_k|s_k) \hat{p}_k(s_k|s_{k-1}, a_{k-1})) \, \mathrm{d}(h_{a_{t-1}}),$$

where $\hat{p}_t, \hat{p}_{\pi_t^b}$ are each empirical frequency (histogram) estimators, and $h_{a_{t-1}} = (s_1, a_1, \dots, a_{t-1})$. When the behavior policy is known, Xie et al. (2019) also proposed a similar method to estimate w_t . In the non-tabular case, we have to rely on some function approximation methods. In this case, we can use the equation $w_t = \mathbb{E}[\lambda_{t-1} \mid s_t]$. Thus, for example, w_t is estimated by regressing λ_{t-1} onto s_t . Alternatively, we can use the recurrence $w_t = \mathbb{E}[\eta_{t-1}w_{t-1} \mid s_t]$ and recursively construct regression estimates similarly to fitted q-iteration but going forward in time rather than backward.

5 Application to Non-Markov Decision Processes

So far, we have so focused on the TMDP setting where the trajectory distribution p_{π} is given by Eq. (1). For the sake of completeness and comparison, we next also consider the *non-Markov* decision process (NMDP) model. Our results for TMDP may in fact be applied directly by recognoizing that an NMDP can be embedded in a TMDP by including the whole history in the state variable. (The opposite cannot be done and the results for TMDP are novel.) We use this to derive the efficiency bound for the NMDP case and compare to Kennedy (2019); Díaz et al. (2020).

Relaxing the Markov assumption of the TMDP model, yields the NMDP model $\mathcal{M}_{\text{NMDP}}$, where the trajectory distribution $p_{\pi}(\mathcal{T})$ is

$$p_1(s_1) \prod_{t=1}^{H} \pi_t(a_t \mid j_{s_t}) p_t(r_t \mid j_{a_t}) p_{t+1}(s_{t+1} \mid j_{r_t}),$$

where j_{a_t} is $(s_1, a_1, r_1, \dots, a_t)$, j_{s_t} is $(s_1, a_1, r_1, \dots, s_t)$ and j_{r_t} is $(s_1, a_1, r_1, \dots, r_t)$.

Notice that in the embedding of an NMDP as a TMDP, the marginal density ratio μ_t in the TMDP is *exactly* the cumulative importance ratio λ_t in the original NMDP. We therefore have the following for tilting policies as a corollary of Theorem 2.

Theorem 8. Let π^e be as in Definition 1. Then the EIF and the efficiency of J bound w.r.t. the model \mathcal{M}_{NMDP} are, respectively,

$$\sum_{t=1}^{H} \lambda_{t} \{ r_{t} - v_{t}(j_{s_{t}}) \} + \lambda_{t-1} v_{t}(j_{s_{t}}), \sum_{t=0}^{H} \mathbb{E}[\lambda_{t-1}^{2}(\mathcal{J}_{s_{t-1}}) \text{var}[\eta_{t}(\mathcal{J}_{a_{t}}) \{ R_{t} + v_{t+1}(\mathcal{J}_{s_{t+1}}) \} \mid \mathcal{J}_{s_{t}}]].$$

When actions are binary $(\mathcal{A}_t = \{0,1\})$ and $r_1 = \cdots = r_{H-1} = 0$ so rewards only occur at the end (r_H) , this coincides with Theorem 2 of Kennedy (2019).

The main issue with the NMDP model is that OPE estimators that are regular w.r.t. the NMDP model necessarily suffer from the curse of horizon, in the sense that their MSE is exponentially growing in horizon.

Theorem 9. Suppose $\exp(\mathbb{E}_{p_{\pi^c}}[\log(\eta_t)]) \geq C_{\min}$ and $\exp(\mathbb{E}[\log(\operatorname{var}[\eta_t(j_{a_t})\{r_t + v_{t+1}(j_{s_{t+1}})\} \mid j_{s_t}])]) \geq V_{\min}^2$. Then, the efficiency bound in Theorem 8 is lower bounded by $C_{\min}^{H-1}V_{\min}^2$.

Notice $\mathbb{E}_{p_{\pi^e}}[\log(\eta_t)]$ is exactly the Kullback-Leibler divergence between $\pi^e(\cdot \mid s_t)$ and $\pi^b(\cdot \mid s_t)$ averaged over $p_{\pi^e}(s_t)$.

We can similarly extend Theorem 3 (modified treatment policies) to the NMDP case as follows, and again see that it suffers from the curse of horizon.

Theorem 10. Let π^{e} be as in Definition 2. Then the EIF and efficiency bound of J w.r.t. the model \mathcal{M}_{NMDP} are, respectively,

$$\sum_{t=1}^{H} \lambda_t \{ r_t - q_t(j_{a_t}) \} + \lambda_{t-1} q_t(j_{s_t}, \tau_t(j_{a_t})), \sum_{t=0}^{H} \mathbb{E}[\lambda_t^2 \operatorname{var}[r_t + q_{t+1}(j_{s_{t+1}}, \tau_{t+1}(j_{a_{t+1}})) \mid j_{a_t}]].$$

Again, when $r_1 = \cdots = r_{H-1} = 0$ so rewards only occur at the end (r_H) , this coincides with Theorem 2 of Díaz et al. (2020). Again, this still suffers from the curse of horizon.

Theorem 11. Suppose $\exp(\mathbb{E}_{p_{\pi^c}}[\log(\eta_t)]) \geq C_{\min}$ and $\exp(\mathbb{E}[\log(\operatorname{var}[r_t + q_{t+1}(j_{s_{t+1}}, \tau_{t+1}(j_{a_{t+1}})) \mid j_{a_t}])]) \geq V_{\min}^2$. Then, the efficiency bound in Theorem 10 is lower bounded by $C_{\min}^H V_{\min}^2$.

6 Extension to Infinite-Horizon Time-Homogeneous Markov Decision Processes

We next extend the results to time-homogeneous MDPs (MDPs) where transitions, rewards, and policies do not depend on t, i.e., $p_t(r|s,a) = p(r|s,a), p_t(s'|s,a) = p(s'|s,a), \pi_t^{\rm b} = \pi^{\rm b}, \tau_t = \tau$. Here, the estimand we consider is an average discounted reward, $J(\gamma) = (1-\gamma) \lim_{T\to\infty} \mathbb{E}_{\pi^{\rm c}}[\sum_{t=1}^T \gamma^{t-1} r_t]$ when the initial state distribution is $p_e^{(1)}(s)$. In this section, We derive the EIF and efficiency bound of $J(\gamma)$ w.r.t. the MDP model. Then, we discuss the relationship to the efficiency bound and estimators for pre-specified evaluation policies.

Although we can still apply methods for TMDP in the MDP case, if we correctly leverage the time-homogeneity of MDP we should do much better in that the rate of MSE should be $\mathfrak{G}(1/NH)$, not $\mathfrak{G}(1/N)$, when we observe N trajectories of length H. Per Liu et al. (2018),

$$J(\gamma) = \mathbb{E}_{s \sim p_b(s), a \sim \pi^{\mathrm{b}}(a|s), r \sim p(r|s, a)} [r \ p_{e, \gamma}^{(\infty)}(s) \pi^{\mathrm{e}}(a \mid s) / (p_b(s) \pi^{\mathrm{b}}(a \mid s))], \tag{7}$$

where $p_b(s)$ is any distribution² and $p_{e,\gamma}^{(\infty)}$ is the γ -discounted average visitation distribution associated with the MDP, π^e , and the initial state distribution $p_e^{(1)}(s)$.

When $\pi^{\rm e}$ is pre-specified, Kallus and Uehara (2019a) derived the efficiency bound of $J(\gamma)$ under a nonparametric model and proposed an efficient estimator. Here, we present corresponding results for the efficient evaluation of natural stochastic policies in time-homogeneous infinite-horizon RL. We consider the observed data to be n i.i.d. draws from the stationary behavior distribution: for $i=1,\ldots,n$, n

$$(s^{(i)}, a^{(i)}, r^{(i)}, s'^{(i)}, a'^{(i)}) \sim p_b(s, a, r, s', a') = p_b(s)\pi^{\mathsf{b}}(a \mid s)p(s' \mid s, a)p(r \mid s, a)\pi^{\mathsf{b}}(a' \mid s').$$

We consider a fully nonparametric model \mathcal{M}_{MDP} in that we make no restrictions on the above distributions. In this section, we define $q(s,a) = \mathbb{E}_{p_{\pi_e}}[\sum_{t=1}^{\infty} \gamma^{t-1} r_t | s_1 = s, a_1 = a], \ q^{\tau}(s,a) = q(s,\tau(s,a)), \ v(s) = \mathbb{E}_{\pi^e(a|s)}[q(s,a)|s], \ w^*(s) = p_{e,\gamma}^{(\infty)}(s)/p_b(s), \ \text{and} \ \mu^*(s,a) = w^*(s)\eta(s,a).$

6.1 Tilting Policies

We consider first the case of tilting policies.

Theorem 12. Let π^e be as in Definition 1. Then the EIF and efficiency bound of $J(\gamma)$ w.r.t. \mathcal{M}_{MDP} are, respectively,

$$\mu^*(s, a)(r + \gamma v(s') - v(s)), \quad \Upsilon_{\text{TI2}} = \mathbb{E}[\text{var}[\mu^*(s, a)(r + \gamma v(s')) \mid s]].$$

We can construct an efficient estimator by following a similar but slightly different cross-fitting strategy as before. With additional data $s_1^{(j)} \sim p_e^{(1)}(s)$, $j=1,\ldots,m$ where $m=\Omega(n)$ (or, if $p_e^{(1)}$ is known), and given nuisance estimators $\hat{\pi}^{b,(k)}$, $\hat{q}^{(k)}$, $\hat{w}^{*(k)}$, we propose the estimator \hat{J}_{TI2} for $J(\gamma)$ by taking Algorithm 1 and replacing \hat{J}_k with

$$\hat{J}_{k} = \frac{1 - \gamma}{m} \sum_{j=1}^{m} \hat{v}^{(k)}(s_{1}^{(j)}) + \frac{1}{|I_{k}|} \sum_{i \in I_{k}} \hat{w}^{*(k)}(s^{(i)}) \hat{\eta}^{(k)}(s^{(i)}, a^{(i)}) (r^{(i)} + \gamma \hat{v}^{(k)}(s^{'(i)}) - \hat{v}^{(k)}(s^{(i)})), \quad (8)$$

$$\hat{\pi}^{e,(k)}(a \mid s) = u(a)\hat{\pi}^{b,(k)}(a \mid s) / \int u(\tilde{a})\hat{\pi}^{b,(k)}(\tilde{a} \mid s)d\tilde{a},$$

$$\hat{\eta}^{(k)}(s,a) = \hat{\pi}^{e,(k)}(a \mid s)/\hat{\pi}^{b,(k)}(a \mid s), \ \hat{v}^{(k)}(s) = \int \hat{q}^{(k)}(s,a)\hat{\pi}^{e,(k)}(a \mid s)da. \tag{9}$$

To estimate $\pi^{\rm b}$, we can follow Section 4.3. To estimate w^*, q , we can solve the following conditional moment equations (cf. Liu et al., 2018; Kallus and Uehara, 2019a) either by instantiating with some set test functions and solving the empirical moments or using a method such as Bennett et al. (2019):

$$0 = (1 - \gamma) \mathbb{E}_{s_1 \sim p_1^{(1)}}[f(s_1)] + \mathbb{E}_{(s,a,s') \sim p_b}[\gamma w^*(s)(\eta(s,a)f(s') - f(s))] \qquad \forall f(s), \tag{10}$$

$$0 = \mathbb{E}_{(s,a,r,s') \sim p_b}[g(s,a)(r + \gamma v(s') - q(s,a))] \qquad \forall g(s,a). \tag{11}$$

Given any nuisance estimators that satisfy certain weak rate assumptions (and no other assumptions), we prove \hat{J}_{TI2} is efficient and partially doubly robust.

²In practice, it is often take as the stationary distribution associated with the MDP and the behavior policy $\pi^{\rm b}$. Our theory holds not only for this case but also for any $p_b(s)$.

³Considering the data to instead be N=n/H observations of H-long trajectories can be also handled if we impose some additional mixing assumptions following Kallus and Uehara (2019a), which ensures the dependent trajectories look approximately like independent observations. This only changes the analysis. The estimator itself is unchanged.

⁴The observation of the ensuing action $a'^{(i)}$ is actually only needed for the case of modified treatment policies.

Theorem 13 (Efficiency). Suppose $\forall k \leq K$, $\|\hat{w}^{*(k)}(s) - w^{*}(s)\|_{2} \leq \alpha_{1}$, $\|\hat{\pi}^{b}(a|s) - \pi^{b}(a|s)\|_{2} \leq \alpha_{2}$, $\|\hat{q}^{(k)}(s,a) - q(s,a)\|_{2} \leq \beta$, where $\alpha_{1} = \mathbb{O}_{p}(n^{-1/4})$, $\alpha_{2} = \mathrm{o}_{p}(n^{-1/4})$, $\beta = \mathbb{O}_{p}(n^{-1/4})$, $\alpha_{1}\beta = \mathrm{o}_{p}(n^{-1/2})$. Then, $\sqrt{n}(\hat{J}_{\text{TI2}} - J) \stackrel{d}{\to} \mathcal{N}(0, \Upsilon_{\text{TI2}})$.

Theorem 14 (Partial Double Robustness). Assume $\forall k \leq K$, for some $w^{*\dagger}(s), q^{\dagger}(s, a), \|\hat{w}^{*(k)}(s) - w^{*\dagger}(s)\|_2 = o_p(1), \|\hat{\pi}^b(a|s) - \pi^b(a|s)\|_2 = o_p(1), \|\hat{q}^{(k)}(s, a) - q^{\dagger}(s, a)\|_2 = o_p(1)$. Then, as long as $w^{*\dagger}(s) = w^*(s)$ or $q^{\dagger}(s, a) = q(s, a), \hat{J}_{\text{TI2}} \stackrel{p}{\rightarrow} J$.

The result again essentially follows by showing that $|\hat{J}_{\text{TI1}} - J - \mathbb{P}_n[\phi(s, a, r, s')]| \leq \alpha_1 \alpha_2 + \alpha_1 \beta + \alpha_2 \beta + \alpha_2^2 + o_p(n^{-1/2})$, where $\phi(s, a, r, s')$ is the EIF. Under the above rate assumptions, the right-hand side is $o_p(n^{-1/2})$ and the result is immediately concluded from CLT. Partially doubly robustness similarly holds as in Section 4.

6.2 Modified Treatment Policies

We next consider the case of modified treatment policies.

Theorem 15. Let π^e be as in Definition 2. Then the EIF and efficiency bound of $J(\gamma)$ w.r.t. \mathcal{M}_{MDP} are, respectively,

$$\mu^*(s, a)(r + \gamma q^{\tau}(s', a') - q(s, a)), \quad \Upsilon_{MO2} = \mathbb{E}[\mu^{*2}(s, a)var[r + \gamma q^{\tau}(s', a') \mid s, a]].$$

We can construct an efficient estimator as follows. With additional data $(s^{(j)}, a^{(j)}) \sim p_e^{(1)}(s)\pi^b(a \mid s)$, $j = 1, \ldots, m$ where $m = \Omega(n)$, and given nuisance estimators $\hat{w}^{*(k)}, \hat{\pi}^{b,(k)}, \hat{q}^{(k)}$, we propose the estimator \hat{J}_{MO2} by taking Algorithm 1 and replacing \hat{J}_k with

$$\hat{J}_{k} = \frac{1}{|I_{k}|} \sum_{i \in I_{k}} \hat{w}^{*(k)}(s) \hat{\eta}^{(k)}(s^{(i)}, a^{(i)}) (r^{(i)} + \gamma \hat{q}^{(k)\tau}(s^{'(i)}, a^{'(i)}) - \hat{q}^{(k)}(s^{(i)}, a^{(i)}))$$

$$+ \frac{(1 - \gamma)}{m} \sum_{j=1}^{m} \hat{q}^{(k)\tau}(s^{(j)}, a^{(j)}), \quad \hat{\pi}^{e,(k)}(a_{t} \mid s_{t}) = \hat{\pi}^{b,(k)}(\tilde{\tau}(s_{t}, a_{t}) \mid s_{t})\tilde{\tau}'(s_{t}, a_{t}),$$

and $\hat{\eta}^{(k)}$, $\hat{v}^{(k)}$ are as in Eq. (9). To estimate w^* we can use Eq. (10) and to estimate q we can use:

$$0 = \mathbb{E}_{(s,a,r,s',a') \sim p_b} [g(s,a) (r - q(s,a) + \gamma q^{\tau}(s',a'))] \qquad \forall g(s,a).$$
 (12)

We next prove \hat{J}_{MO2} is efficient and doubly robust.

Theorem 16 (Efficiency). Assume $\forall k \leq K$, $\|\hat{w}^{*(k)}(s) - w^*(s)\|_2 \leq \alpha_1$, $\|\hat{\pi}^{b,(k)}(a|s) - \pi^b(a|s)\|_2 \leq \alpha_2$, $\|\hat{q}^{(k)}(s,a) - q(s,a)\|_2 \leq \beta$, where $(\alpha_1 + \alpha_2)\beta = o_p(n^{-1/4})$, $\max\{\alpha_1, \alpha_2, \beta\} = o_p(1)$. Then, $\sqrt{n}(\hat{J}_{MO2} - J) \stackrel{d}{\to} \mathcal{N}(0, \Upsilon_{MO2})$.

Theorem 17 (Double Robustness). Assume $\forall k \leq K$, for some $w^{*\dagger}, \pi^{b\dagger}, q^{\dagger}, \|\hat{w}^{*(k)}(s) - w^{*\dagger}(s)\|_2 = o_p(1), \|\hat{\pi}^{b,(k)}(a|s) - \pi^{b\dagger}(a|s)\|_2 = o_p(1), \|\hat{q}^{(k)}(s,a) - q^{\dagger}(s,a)\|_2 = o_p(1)$. Then, as long as $w^{*\dagger} = w^*, \pi^{b\dagger} = \pi^b$ or $q^{\dagger} = q$, $\hat{J}_{\text{MO2}} \stackrel{p}{\to} J$.

6.3 Comparison with Kallus and Uehara (2019a); Tang et al. (2020)

We next compare our results with some related work for the case of pre-specified policies.

Comparison with Kallus and Uehara (2019a) When the evaluation policy is pre-specified, Kallus and Uehara (2019a) proposed an estimator that is similar but uses $\hat{q}^{(k)}$ in place of the last $\hat{v}^{(k)}$ in Eq. (8). Then, under similar rate conditions to Theorem 13, they prove it achieves the asymptotic MSE

$$\Upsilon_{PR} = \mathbb{E}[\mu^{*2}(s, a) \text{var}[r + \gamma v(s') \mid s, a]],$$

which is the efficiency bound when the evaluation policy is pre-specified so the estimator is efficient. Notice that Υ_{PR} is smaller than Υ_{TI2} by $\mathbb{E}[w^{*2}(s)\text{var}[\eta(s,a)(r+\gamma v(s'))\mid s]]$. As in Remark 6, naïvely plugging in an estimated π^e into the EIF for the pre-specified case can fail to be efficient or even \sqrt{n} -consistent.

Comparison with Tang et al. (2020) In the case of a pre-specified evaluation policy and known behavior policy, Tang et al. (2020) propose an estimator with a form similar to Eq. (8) without sample splitting and where \hat{v} is directly estimated (rather than computed as a function of other nuisance estimates). The similarity to Eq. (8) appears to be coincidental and superficial. In the case of pre-specified evaluation policy, even if we used oracle values for all nuisances, the estimator of Tang et al. (2020) is inefficient since its variance would be equal to Υ_{TI2} , which is larger than Υ_{PR} . Tang et al. (2020) do not study the asymptotic properties of their estimator, but we can show that using cross-fitting of nuisance esimates, the asymptotic MSE of the estimator is also equal to Υ_{TI2} (see Appendix B). Again, this is inefficient in the case of a pre-specified evaluation policy. And, in the case of a natural stochastic policy, \hat{v} must be computed in a fashion compatible with \hat{q} and $\hat{\pi^b}$ in order to ensure the second-order bias structure and hence efficiency.

7 Empirical Study

In this section we examine the performance of different OPE estimators in a time-invariant infinitehorizon setting. We use the Taxi environment, which is a commonly used tabular environment for OPE, which has $\mathcal{S} = \{1, \dots, 2000\}, \mathcal{A} = \{1, \dots, 6\}$ (Dietterich, 2000; we also refer the reader to Liu et al., 2018, Section 5 for more details), and we consider separate experiments for the case of tilting and modified treatment policies. We consider observing of a single (N=1) trajectory of varying length $H(\{1, 2.5, 5, 10\} \times 10^4)$. For each H we run 60 replications of the experiment. We compare the stationary marginal IS estimator \hat{J}_{MIS} (Liu et al., 2018), the direct method \hat{J}_{DM} , and one of our proposed estimators $\hat{J}_{\text{TI2}}, \hat{J}_{\text{MO2}}$, depending on whether we are considering tilting or modified treatment policies. We do not compare to step-wise IS (Precup et al., 2000) and DR (Jiang and Li, 2016) as these estimators do not converge when given single-trajectory data (as shown in Kallus and Uehara, 2019a, Section 7). Behavior and evaluation policies are set as follows. We run 150 iterations of q-learning to learn a near-optimal policy for the MDP and define this to be π^b . We consider evaluating either a tilting policy with $u(a) = \lceil a/2 \rceil$ or a modified treatment policy with $\tau(s,a) = (s+a) \mod 6$. We set $\gamma = 0.98$. We estimate π^b as $\hat{\pi}^b(a \mid s) = \sum_{i=1}^n \mathbb{I}\left[a^{(i)} = a, s^{(i)} = s\right] / \sum_{i=1}^n \mathbb{I}\left[s^{(i)} = s\right]$ and w^* - and q-functions by solving Eqs. (10) to (12) using $\{\mathbb{I}[s=i]: i=1,\ldots,2000\}$ and $\{\mathbb{I}[s=i,a=j]:i=1,\ldots,2000,j=1,\ldots,6\}$ as test functions, respectively. We use these nuisance estimates to construct all estimators. To validate double robustness, we also add Gaussian noise $\mathcal{N}(3.0, 1.0)$ to either the q- or w^* -function estimates to simulate misspecification. In Figs. 1 to 6, we report the MSE of each estimator over the 60 replications with 95% confidence intervals.

We find the performance of \hat{J}_{TI2} or of \hat{J}_{MO2} is consistently good, with or without of model specification due to double robustness. While MIS and DM fail when their respective model is misspecified, they do well when well-specified. Since either parametric misspecification or nonparametric rates for w^* and q is unavoidable in practice for large or continuous state-action spaces, \hat{J}_{TI2} and \hat{J}_{MO2} are seen to be superior to \hat{J}_{DM} and \hat{J}_{MIS} .




Figure 4: Well-specified q, w^*

Figure 5: Misspecified q

Figure 6: Misspecified w^*

8 Conclusions

We considered the evaluation of natural stochastic policies in RL, both in finite and infinite horizons. We derived the efficiency bounds and proposed estimators that achieved them under lax conditions on nuisance estimators that permit flexible machine learning methods. An important next question is *learning* natural stochastic policies. This can perhaps be done using an off-policy policy gradient approach by extending Kallus and Uehara (2020b) to natural stochastic policies, where we take gradients in u or τ that specify how and where we deviate from π^b .

References

Antos, A., C. Szepesvári, and R. Munos (2008). Learning near-optimal policies with bellman-residual minimization based fitted policy iteration and a single sample path. *Machine Learning* 71, 89–129.

Bennett, A., N. Kallus, and T. Schnabel (2019). Deep generalized method of moments for instrumental variable analysis. In *Advances in Neural Information Processing Systems*, pp. 3564–3574.

Bibaut, A., I. Malenica, N. Vlassis, and M. Van Der Laan (2019). More efficient off-policy evaluation through regularized targeted learning. In *Proceedings of the 36th International Conference on Machine Learning*, Volume 97, pp. 654–663.

Boruvka, A., D. Almirall, K. Witkiewitz, and S. A. Murphy (2018). Assessing time-varying causal effect moderation in mobile health. *Journal of the American Statistical Association* 113 (523), 1112 – 1121.

Chernozhukov, V., D. Chetverikov, M. Demirer, E. Duflo, C. Hansen, W. Newey, and J. Robins (2018). Double/debiased machine learning for treatment and structural parameters. *Econometrics Journal* 21, C1–C68.

Díaz, I. and M. J. van Der Laan (2013). Assessing the causal effect of policies: an example using stochastic interventions. *The international journal of biostatistics 9*, 161–174.

- Díaz, I. and M. J. van der Laan (2018). Stochastic Treatment Regimes, pp. 219–232. Springer International Publishing.
- Díaz, I., N. Williams, K. L. Hoffman, and E. J. Schenck (2020). Non-parametric causal effects based on longitudinal modified treatment policies. arXiv preprint arXiv:2006.01366.
- Dietterich, T. G. (2000). Hierarchical reinforcement learning with the maxq value function decomposition. *Journal of Artificial Intelligence Research* 13, 227–303.
- Duan, Y. and M. Wang (2020). Minimax-optimal off-policy evaluation with linear function approximation. *ICML 2020 (To appear)*.
- Díaz, I. and N. S. Hejazi (2020). Causal mediation analysis for stochastic interventions. Journal of the Royal Statistical Society: Series B (Statistical Methodology).
- Ernst, D., P. Geurts, and L. Wehenkel (2005). Tree-based batch mode reinforcement learning. *Journal of Machine Learning Research* 6, 503–556.
- Ertefaie, A. and R. L. Strawderman (2018). Constructing dynamic treatment regimes over indefinite time horizons. *Biometrika* 105, 963–977.
- Gottesman, O., F. Johansson, M. Komorowski, A. Faisal, D. Sontag, F. Doshi-Velez, and L. A. Celi (2019). Guidelines for reinforcement learning in healthcare. *Nat Med* 25, 16–18.
- Haneuse, S. and A. Rotnitzky (2013). Estimation of the effect of interventions that modify the received treatment. *Statistics in Medicine 32*, 5260–5277.
- Jiang, N. and L. Li (2016). Doubly robust off-policy value evaluation for reinforcement learning. In Proceedings of the 33rd International Conference on International Conference on Machine Learning-Volume, 652–661.
- Kallus, N. and M. Uehara (2019a). Efficiently breaking the curse of horizon: Double reinforcement learning in infinite-horizon processes. arXiv preprint arXiv:1909.05850.
- Kallus, N. and M. Uehara (2019b). Intrinsically efficient, stable, and bounded off-policy evaluation for reinforcement learning. In *Advances in Neural Information Processing Systems* 32, pp. 3320–3329.
- Kallus, N. and M. Uehara (2020a). Double reinforcement learning for efficient off-policy evaluation in markov decision processes. *Journal of Machine Learning Research* 21, 1–63.
- Kallus, N. and M. Uehara (2020b). Statistically efficient off-policy policy gradients. *ICML 2020 (To appear)*.
- Kennedy, E. H. (2019). Nonparametric causal effects based on incremental propensity score interventions. Journal of the American Statistical Association 114, 645–656.
- Kosorok, M. R. and E. E. Moodie (2015). Adaptive Treatment Strategies in Practice: Planning Trials and Analyzing Data for Personalized Medicine. USA: Society for Industrial and Applied Mathematics.
- Lagoudakis, M. and R. Parr (2004). Least-squares policy iteration. *Journal of Machine Learning Research* 4(6), 1107–1149.
- Le, H., C. Voloshin, and Y. Yue (2019). Batch policy learning under constraints. In *Proceedings of the 36th International Conference on Machine Learning*, pp. 3703–3712.
- Liao, P., P. Klasnja, and S. Murphy (2020). Off-policy estimation of long-term average outcomes with applications to mobile health. *Journal of the American Statistical Association (To appear)*.

- Liu, Q., L. Li, Z. Tang, and D. Zhou (2018). Breaking the curse of horizon: Infinite-horizon off-policy estimation. In *Advances in Neural Information Processing Systems* 31, pp. 5356–5366.
- Luckett, D. J., E. B. Laber, A. R. Kahkoska, D. M. Maahs, E. Mayer-Davis, and M. R. Kosorok (2018). Estimating dynamic treatment regimes in mobile health using v-learning. *Journal of the American Statistical Association*, 1–34.
- Munos, R. and C. Szepesvári (2008). Finite-time bounds for fitted value iteration. Journal of Machine Learning Research 9, 815–857.
- Murphy, S. A. (2003). Optimal dynamic treatment regimes. *Journal of the Royal Statistical Society:* Series B (Statistical Methodology) 65, 331–355.
- Murphy, S. A., M. J. Van Der Laan, and J. M. Robins (2001). Marginal mean models for dynamic regimes. *Journal of the American Statistical Association* 96, 1410–1423.
- Muñoz, I. D. and M. Van Der Laan (2012). Population intervention causal effects based on stochastic interventions. *Biometrics* 68, 541–549.
- Nachum, O., Y. Chow, B. Dai, and L. Li (2019). Dualdice: Behavior-agnostic estimation of discounted stationary distribution corrections. In *Advances in Neural Information Processing Systems* 32.
- Precup, D., R. S. Sutton, and S. P. Singh (2000). Eligibility Traces for Off-Policy Policy Evaluation. In *Proceedings of the 17th International Conference on Machine Learning*, pp. 759–766.
- Richardson, T. S. and J. M. Robins (2013). Single world intervention graphs (swigs): a unification of the counterfactual and graphical approaches to causality. CSSS Working Paper 128.
- Robins, J. (2004). Optimal structural nested models for optimal sequential decisions. In Proceedings of the Second Seattle Symposium in Biostatistics: Analysis of Correlated Data.
- Robins, J. M., A. Rotnitzky, and D. O. Scharfstein (1999). Sensitivity Analysis for Selection Bias and Unmeasured Confounding in Missing Data and Causal Inference Models, Volume 116 of Statistical Models in Epidemiology: The Environment and Clinical Trials. NY: Springer-Verlag.
- Shi, C., S. Zhang, W. Lu, and R. Song (2020). Statistical inference of the value function for reinforcement learning in infinite horizon settings. arXiv preprint arXiv:2001.04515.
- Shpitser, I. and J. Pearl (2012). Effects of treatment on the treated: Identification and generalization.

 Proceedings of the 25th Conference On Uncertainty in Artificial Intelligence.
- Tang, Z., Y. Feng, L. Li, D. Zhou, and Q. Liu (2020). Harnessing infinite-horizon off-policy evaluation: Double robustness via duality. *ICLR 2020 (To appear)*.
- Thomas, P. and E. Brunskill (2016). Data-efficient off-policy policy evaluation for reinforcement learning. In Proceedings of the 33rd International Conference on Machine Learning, 2139–2148.
- Ueno, T., M. Kawanabe, T. Mori, S.-I. Maeda, and S. Ishii (2011). Generalized td learning. *Journal of Machine Learning Research* 12, 1977–2020.
- van Der Laan, M. J. and J. M. Robins (2003). *Unified Methods for Censored Longitudinal Data and Causality*. Springer Series in Statistics, New York, NY: Springer New York.
- van der Vaart, A. W. (1998). Asymptotic statistics. Cambridge, UK: Cambridge University Press.
- Xie, T., Y. Ma, and Y.-X. Wang (2019). Towards optimal off-policy evaluation for reinforcement learning with marginalized importance sampling. In Advances in Neural Information Processing Systems 32, pp. 9665–9675.

- Yin, M. and Y.-X. Wang (2020). Asymptotically efficient off-policy evaluation for tabular reinforcement learning. In Proceedings of the 23nd International Workshop on Artificial Intelligence and Statistics (To appear).
- Young, J. G., M. A. Herńan, and J. M. Robins (2014). Identification, estimation and approximation of risk under interventions that depend on the natural value of treatment using observational data. *Epidemiologic methods* 3, 1–19.
- Young, J. G., R. W. Logan, J. M. Robins, and M. A. Hernán (2019). Inverse probability weighted estimation of risk under representative interventions in observational studies. *Journal of the American Statistical Association* 114, 938–947.
- Zhang, B., A. Tsiatis, E. Laber, and M. Davidian (2013). Robust estimation of optimal dynamic treatment regimes for sequential treatment decisions. *Biometrika*, 681.
- Zheng, W. and M. J. van Der Laan (2011). Cross-validated targeted minimum-loss-based estimation. In Targeted Learning: Causal Inference for Observational and Experimental Data, Springer Series in Statistics, pp. 459–474. New York, NY: Springer New York.

SUPPLEMENTARY MATERIAL

A Proofs

Throughout this paper, in an MDP, we define

$$c(s) = 1/\int u(a)\pi^{\mathbf{b}}(a|s)\mathrm{d}a.$$

Then, $\pi^{e}(a|s) = u(a)\pi^{b}(a|s)c(s)$. Similarly, in a TMDP, we define

$$c_t(s_t) = 1/\int u(a_t)\pi^{\mathrm{b}}(a_t|s_t)\mathrm{d}a_t.$$

Then, $\pi_t^e(a_t|s_t) = u_t(a_t)\pi_t^b(a_t|s_t)c_t(s_t)$. In addition, note that $\hat{\pi}^e(a|s)$ and $\hat{\pi}_t^e(a_t|s_t)$ are expressed as

$$\hat{\pi}^e(a|s) = u(a)\hat{\pi}^b(a|s)\hat{c}(s), \, \hat{\pi}^e_t(a_t|s_t) = u(a_t)\hat{\pi}^b(a_t|s_t)\hat{c}(s_t).$$

A.1 Warm-up

As a warm-up, we prove the results for the bandit case and NMDP case. In a bandit setting, we drop an index t.

Proof of Remark 3. The entire regular parametric sumbodel is

$$\{p_{\theta}(s)p_{\theta}(a \mid s)p_{\theta}(r|s,a)\},\$$

where it matches with a true pdf at $\theta = 0$. The score functions of the nonparametric model is decomposed as

$$g(\mathcal{J}) = \log p_{\theta}(s) + \log p_{\theta}(a \mid s) + \log p_{\theta}(r \mid s, a) = g_S + g_{A\mid S} + g_{R\mid S, A}.$$

We calculate the gradient of the target functional $J(\pi^{e})$ w.r.t. the nonparametric model. Since

$$J(\pi^{\mathrm{e}}) = \int r p_{\theta}(r \mid a, s) \pi^{\mathrm{e}}(a \mid s; \pi^{\mathrm{b}}) p_{\theta}(s) \mathrm{d}(a, s, r),$$

we have

$$\nabla J(\pi^{\mathrm{e}}) = \mathbb{E}\left[\frac{\pi^{\mathrm{e}}(A\mid S)}{\pi^{\mathrm{b}}(A\mid S)} \{R - q(S,A)\}g(\mathcal{J}) + v(S)g(\mathcal{J}) + R\frac{\nabla_{\theta}\pi^{\mathrm{e}}(A\mid S; \pi^{\mathrm{b}})}{\pi^{\mathrm{b}}(A\mid S)}\right].$$

Especially, the third term is

$$\begin{split} & \mathbb{E}\left[\frac{R\nabla_{\theta}\pi^{\mathrm{e}}(A\mid S; \pi^{\mathrm{b}})}{\pi^{\mathrm{b}}(A\mid S)}\right] \\ & = \mathbb{E}\left[\left\{\mathbb{E}[R|S, A]\frac{u(A)}{c(S)} - \mathbb{E}\left[R\frac{u(A)}{c(S)}\mid S\right]\frac{u(A)}{c(S)}\right\}g_{A\mid S}\right] \\ & = \mathbb{E}\left[\left\{\mathbb{E}[R\mid S, A]\frac{u(A)}{c(S)} - \mathbb{E}\left[R\frac{u(A)}{c(S)}\mid S\right]\frac{u(A)}{c(S)}\right\}g(\mathcal{J})\right] \\ & = \mathbb{E}\left[\frac{\pi^{\mathrm{e}}(A\mid S)}{\pi^{\mathrm{b}}(A\mid S)}\{q(S, A) - v(S)\}g(\mathcal{J})\right]. \end{split}$$

Then, we have

$$\nabla J(\pi^{\mathrm{e}}) = \mathbb{E}\left[\left\{\frac{\pi^{\mathrm{e}}(A\mid S)}{\pi^{\mathrm{b}}(A\mid S)}(R - v(S)) + v(S)\right\}g(\mathfrak{F})\right].$$

Since the gradient is unique for the current case, this concludes that the following is the EIF:

$$\frac{\pi^{e}(a|s)}{\pi^{b}(a|s)} \{r - v(s)\} + v(s).$$

Proof of Remark 5. The proof is also mentioned in Muñoz and Van Der Laan (2012). We add the proof here since this would improve the reader's understanding in the RL case. The entire regular parametric sumbodel is

$$\{p_{\theta}(s)p_{\theta}(a \mid s)p_{\theta}(r|s,a)\}$$

where it matches with the true pdf at $\theta = 0$. The score functions of the nonparametric model is decomposed as

$$g(\mathcal{J}) = \log p_{\theta}(s) + \log p_{\theta}(a \mid s) + \log p_{\theta}(r \mid s, a)$$
$$= g_S + g_{A\mid S} + g_{R\mid S, A}.$$

We calculate the gradient of the target functional $J(\pi^{\rm e})$ w.r.t. the nonparametric model. Since

$$J(\pi^{e}) = \int rp_{\theta}(r \mid \tau(s, a), s) \pi^{b}(a \mid s) p_{\theta}(s) d(a, s, r)$$
$$= \int rp_{\theta}(r \mid a, s) \pi^{b}(\tilde{\tau}(s, a) \mid s) \tilde{\tau}'(s, a) p_{\theta}(s) d(a, s, r),$$

we have

$$\nabla_{\theta} J(\pi^{\mathbf{e}}) = \mathbb{E}\left[\left\{\frac{\pi^{\mathbf{b}}(\tilde{\tau}(S,A)\mid S)\tilde{\tau}'(S,A)}{\pi^{\mathbf{b}}(A\mid S)}\{R - q(S,A)\} + \mathbb{E}[R\mid S, \tau(S,A)]\right\}g(\mathcal{J})\right].$$

Since the gradient is unique for the current case, this concludes that the following is the EIF:

$$\frac{\pi^{e}(a \mid s)}{\pi^{b}(a \mid s)} \{r - q(s, a)\} + q(s, \tau(s, a)).$$

A.2 Proofs for Section 3

Proof of Theorem 2.

Calculation of derivatives under a nonpametric TMDP The entire regular parametric submodel is

$$\{p_{\theta}(s_1)p_{\theta}(a_1|s_1)p_{\theta}(r_1|s_1,a_1)p_{\theta}(s_1|s_1,a_1)p_{\theta}(a_1|s_1)p_{\theta}(r_1|s_1,a_1)\cdots p_{\theta}(r_H|s_H,a_H)\}.$$

The score function of the parametric submodel is

$$g(j_{r_H}) = \sum_{k=1}^{H} \nabla_{\theta} \log p_{\theta}(s_k \mid s_{k-1}, a_{k-1}) + \nabla_{\theta} \log p_{\theta}(a_k \mid s_k) + \nabla_{\theta} \log p_{\theta}(r_k \mid s_k, a_k)$$

$$= \sum_{k=1}^{H} g_{S_k|S_{k-1},A_{k-1}} + \sum_{k=1}^{H} g_{A_k|S_k} + \sum_{k=1}^{H} g_{R_k|S_k,A_k}.$$

We have

$$\begin{split} &\nabla_{\theta} \mathbb{E}_{\pi^{e}}[\sum_{t=1}^{H} r_{t}] = \nabla_{\theta} \int \sum_{t=1}^{H} r_{t} \left\{ \prod_{k=1}^{t} p_{\theta}(s_{k}|a_{k-1}, s_{k-1}) p_{\pi_{k}^{e}}(a_{k}|s_{k}; \theta) p_{\theta}(r_{k}|a_{k}, s_{k}) \right\} \mathrm{d}\mu(j_{r_{H}}) \\ &= \sum_{c=1}^{H} \{ \mathbb{E}_{\pi^{e}}\left[(\mathbb{E}_{\pi^{e}}[r_{c}|s_{1}] - \mathbb{E}_{\pi^{e}}[r_{c}]) g_{s_{1}} \right] + \mathbb{E}_{\pi^{e}}\left[(r_{c} - \mathbb{E}_{\pi^{e}}[r_{c}|s_{c}, a_{c}]) g_{R_{c}|S_{c}, A_{c}} \right] \\ &+ \mathbb{E}_{\pi^{e}} \left[\left(\mathbb{E}_{\pi^{e}}\left[\sum_{c=t+1}^{H} r_{t}|s_{c+1} \right] - \mathbb{E}_{\pi^{e}}\left[\sum_{c=t+1}^{H} r_{t}|s_{c}, a_{c} \right] \right) g_{S_{c+1}|S_{c}, A_{c}} \right] \\ &+ \mathbb{E}_{\pi^{b}} \left[\mu_{c}(s_{c}, a_{c}) \left(\mathbb{E}_{\pi^{e}}\left[\sum_{c=t}^{H} r_{t}|s_{c}, a_{c} \right] - \mathbb{E}_{\pi^{e}}\left[\sum_{c=t}^{H} r_{t}|s_{c} \right] \right) g_{A_{c}|S_{c}} \right] \right\}. \end{split}$$

Except for the third term, the calculation is the same as Theorem 2 (Kallus and Uehara, 2020a). The third term has been calculated as the proof of Theorem 8. Then, $\nabla_{\theta} E_{\pi^c} [\sum_{t=1}^{H} r_t]$ is equal to

$$\begin{split} &\sum_{c=1}^{H} \{ \mathbb{E}_{\pi^{b}} [(\mathbb{E}[r_{c}|s_{1}] - \mathbb{E}_{\pi^{e}}[r_{c}])g] + \mathbb{E}_{\pi^{b}} \left[\mu_{c}(s_{c}, a_{c})(r_{c} - \mathbb{E}[r_{c}|s_{c}, a_{c}])g \right] \\ &+ \mathbb{E}_{\pi^{b}} \left[\mu_{c}(s_{c}, a_{c})(\mathbb{E}[\sum_{t=c+1}^{H} r_{t}|s_{c+1}] - \mathbb{E}[\sum_{t=c+1}^{H} r_{t}|s_{c}, a_{c}])g \right] \\ &+ \mathbb{E}_{\pi^{b}} \left[\mu_{c}(s_{c}, a_{c}) \left(\mathbb{E}_{\pi^{e}}[\sum_{c=t}^{H} r_{t}|s_{c}, a_{c}] - \mathbb{E}_{\pi^{e}}[\sum_{c=t}^{H} r_{t}|s_{c}] \right)g \right] \} \\ &= \mathbb{E}_{\pi^{b}} \left[\left[-J + \sum_{c=1}^{H} \left\{ \mu_{c}(s_{c}, a_{c}) \left\{ r_{c} - \mathbb{E}_{\pi_{e}}[\sum_{t=c}^{H} r_{t}|s_{c}] \right\} + \mu_{c-1}(s_{c-1}, a_{c-1}) \sum_{t=c}^{H} \mathbb{E}_{\pi_{e}}[r_{t}|s_{c}] \right\} \right] g(\mathcal{J}_{r_{H}}) \right] \end{split}$$

In the end, we can conclude that the following is a derivative:

$$-J + \sum_{c=1}^{H} \mu_c \{ r_c - v_c(s_c) \} + \mu_{c-1} v_c(s_c). \tag{13}$$

Projection onto the tangent space Then, based on the argument in Appendix B (Kallus and Uehara, 2020b), we need to project it onto the tangent space spanned by the nonparametric model deduced by an MDP. Writing the gradient Eq. (13) as ϕ , the projection of ϕ onto the tangent space is calculated as follows:

$$\begin{split} &\sum_{j=1}^{H} \mathbb{E}[\phi|R_{j}, S_{j}, A_{j}] - \mathbb{E}[\phi|S_{j}, A_{j}] + \mathbb{E}[\phi|A_{j}, S_{j}] - \mathbb{E}[\phi|S_{j}] + \mathbb{E}[\phi|S_{j}, A_{j-1}, S_{j-1}] - \mathbb{E}[\phi|A_{j-1}, S_{j-1}] \\ &= \{\sum_{j=1}^{H} \mu_{j}(R_{j} - \mathbb{E}[R_{j} \mid S_{j}, A_{j}])\} \\ &+ \mathbb{E}[\sum_{c=j}^{H} \mu_{c}\{R_{c} - v_{c}(S_{c}) + v_{c+1}(S_{c+1})\} \mid A_{j}, S_{j}] - \mathbb{E}[\sum_{c=j}^{H} \mu_{c}\{R_{c} - v_{c}(S_{c}) + v_{c+1}(S_{c+1})\} \mid S_{j}] \end{split}$$

$$\begin{split} &+ \mathbb{E}[\sum_{c=j}^{H} \mu_{c}\{R_{c} - v_{c}(S_{c}) + v_{c+1}(S_{c+1})\} \mid S_{j}] - \mathbb{E}[\sum_{c=j}^{H} \mu_{c}\{R_{c} - v_{c}(S_{c}) + v_{c+1}(S_{c+1}) \mid S_{j-1}, A_{j-1}] \\ &+ \mu_{j-1}v_{j}(S_{j}) - \mathbb{E}[\mu_{j-1}v_{j}(S_{j}) \mid S_{j-1}, A_{j-1}] \\ &= \{\sum_{j=1}^{H} \mu_{j}(R_{j} - \mathbb{E}[R_{j} \mid S_{j}, A_{j}])\} + \mu_{j}\{\mathbb{E}[R_{j} \mid S_{j}, A_{j}] - v_{j}(S_{j}) + \mathbb{E}[v_{j+1}(S_{j+1}) \mid S_{j}, A_{j}]\} \\ &+ \mu_{j-1}v_{j}(S_{j}) - \mathbb{E}[\mu_{j-1}v_{j}(S_{j}) \mid S_{j-1}, A_{j-1}] \\ &= -J + v_{1}(S_{1}) + \sum_{j=1}^{H} \mu_{j}(S_{j}, A_{j})\{R_{j} - v_{j}(S_{j}) + v_{j+1}(S_{j+1})\} = \phi. \end{split}$$

This concludes that ϕ is actually the EIF.

Efficiency bound From the law of the total variance, the efficiency bound is

$$\operatorname{var}\left[\sum_{c=1}^{H} \mu_{c} \{R_{c} - v_{c}(S_{c})\} + \mu_{c-1} v_{c}(S_{c})\right]$$

$$= \sum_{t=0}^{H} \mathbb{E}[\operatorname{var}\left[\mathbb{E}\left[\sum_{c=1}^{H} \mu_{c} \{R_{c} + v_{c+1}(S_{c+1}) - v_{c}(S_{c})\} | \mathcal{J}_{S_{t+1}}\right] | \mathcal{J}_{S_{t}}\right]\right]$$

$$= \sum_{t=0}^{H} \mathbb{E}[\operatorname{var}\left[\mu_{t} \{R_{t} + v_{t+1}(S_{t+1})\} | \mathcal{J}_{S_{t}}\right]\right] = \sum_{t=0}^{H} \mathbb{E}[w_{t}^{2}(S_{t}) \operatorname{var}\left[\eta_{t}(S_{t}, A_{t}) \{R_{t} + v_{t+1}(S_{t+1})\} | S_{t}]\right].$$

Order of the efficiency bounds We use importance sampling as follows,

$$\begin{split} &\sum_{t=0}^{H} \mathbb{E}_{\pi^{\mathbf{b}}}[w_{t}^{2} \text{var}_{\pi^{\mathbf{b}}}[\lambda_{t}\{r_{t} + v_{t}(s_{t+1})\} \mid s_{t}]] = \sum_{t=0}^{H} \mathbb{E}_{\pi^{\mathbf{b}}}[w_{t}^{2} \mathbb{E}_{\pi^{\mathbf{b}}}[\lambda_{t}^{2}\{r_{t} + v_{t}(s_{t+1}) - v_{t}(s_{t})\}^{2} \mid s_{t}]] \\ &\leq CC' \sum_{t=0}^{H} \mathbb{E}_{\pi^{\mathbf{c}}}[\mathbb{E}_{\pi^{\mathbf{c}}}[\{r_{t} + v_{t}(s_{t+1}) - v_{t}(s_{t})\}^{2} \mid s_{t}]] = CC' \sum_{t=0}^{H} \mathbb{E}_{\pi^{\mathbf{c}}}[\{r_{t} + v_{t}(s_{t+1}) - v_{t}(s_{t})\}^{2}] \\ &= CC' \mathbb{E}_{\pi^{\mathbf{c}}}[\{\sum_{t=0}^{H} r_{t} + v_{t}(s_{t+1}) - v_{t}(s_{t})\}^{2}] = CC' \mathbb{E}_{\pi^{\mathbf{c}}}[\{[\sum_{t=1}^{H} r_{t}] - J\}^{2}] \leq CC' (R_{\max}H)^{2}. \end{split}$$

Remark 8. Noting

$$\mathbb{E}[\operatorname{var}[f(Z)|X]] = \operatorname{var}[\mathbb{E}[f(Z)|X,Y]|X] + \mathbb{E}[\operatorname{var}[f(Z)|X,Y]|X].$$

for random variables X, Y, Z, the difference of this efficiency bound and the efficiency bound of the pre-specified evaluation policies is

$$\sum_{t=0}^{H} \mathbb{E}[w_t^2(S_t) \text{var}[\eta_t(S_t, A_t) q_t(S_t, A_t) \mid S_t]].$$

Proof of Theorem 3.

EIF and efficiency bound under a TMDP The entire regular parametric submodel is

$$\{p_{\theta}(s_1)p_{\theta}(a_1|s_1)p_{\theta}(r_1|s_1,a_1)p_{\theta}(s_1|s_1,a_1)p_{\theta}(a_1|s_1)p_{\theta}(r_1|s_1,a_1)\cdots p_{\theta}(r_H|s_H,a_H)\},$$

where it matches with the true pdf at $\theta = 0$. The score function of the parametric submodel is

$$g(j_{r_H}) = \sum_{k=1}^{H} \nabla_{\theta} \log p_{\theta}(s_k \mid s_{k-1}, a_{k-1}) + \nabla_{\theta} \log p_{\theta}(a_k \mid s_{k-1}) + \nabla_{\theta} \log p_{\theta}(r_k \mid s_k, a_k)$$

$$= \sum_{k=1}^{H} g_{S_k \mid S_{k-1}, A_{k-1}} + \sum_{k=1}^{H} g_{A_k \mid S_{k-1}} + \sum_{k=1}^{H} g_{R_k \mid S_k, A_k}.$$

The target functional is

$$\int \sum_{t=1}^{H} r_t \left\{ \prod_{k=1}^{t} p_{\theta}(s_k | \tau(a_{k-1}, s_{k-1}), s_{k-1}) \pi_k^{b}(a_k | s_k; \theta) p_{\theta}(r_k | \tau(a_k, s_k), s_k) \right\} d\mu(j_{r_H})
= \int \sum_{t=1}^{H} r_t \left\{ \prod_{k=1}^{t} p_{\theta}(s_k | a_{k-1}, s_{k-1}) \pi_k^{e}(a_k | s_k; \theta) p_{\theta}(r_k | a_k, s_k) \right\} d\mu(j_{r_H}),$$

where $\pi_k^{\mathrm{e}}(a_k \mid s_k)$ is $\pi_k^{\mathrm{b}}(\tilde{\tau}(s, a) \mid s)\tilde{\tau}'(s, a)$, $\tilde{\tau}_k(\cdot, s)$ is the inverse function of $\tau_k(\cdot, s)$. Here, we use a change of variables: $\tau(a_{k-1}, s_{k-1}) = u_{k-1}$, and write u_k as a_k . We have

$$\begin{split} &\nabla_{\theta} \mathbb{E}_{\pi^{e}} [\sum_{t=1}^{H} r_{t}] = \nabla_{\theta} \int \sum_{t=1}^{H} r_{t} \left\{ \prod_{k=1}^{t} p_{\theta}(s_{k} | \tau(s_{k-1}, a_{k-1}), s_{k-1}) \pi_{k}^{e}(a_{k} | s_{k}; \theta) p_{\theta}(r_{k} | a_{k}, s_{k}) \right\} d\mu(j_{r_{H}}) \\ &= \sum_{c=1}^{H} \{ \mathbb{E}_{\pi^{e}} \left[(\mathbb{E}_{\pi^{e}} [r_{c} | s_{1}] - \mathbb{E}_{\pi^{e}} [r_{c}]) g_{S_{1}} \right] + \mathbb{E}_{\pi^{e}} \left[(r_{c} - \mathbb{E}_{\pi^{e}} [r_{c} | s_{c}, a_{c}]) g_{R_{c} | S_{c}, A_{c}} \right] \\ &+ \mathbb{E}_{\pi^{e}} \left[\left(\mathbb{E}_{\pi^{e}} [\sum_{c=t+1}^{H} r_{t} | s_{c+1}] - \mathbb{E}_{\pi^{e}} [\sum_{c=t+1}^{H} r_{t} | s_{c}, a_{c}] \right) g_{S_{c+1} | S_{c}, A_{c}} \right] \\ &+ \mathbb{E}_{\pi^{b}} \left[\prod_{k=1}^{t-1} \eta_{k} \left(\mathbb{E}_{\pi^{e}} [\sum_{c=t}^{H} r_{t} | s_{c}, \tau(s_{c}, a_{c})] - \mathbb{E}_{\pi^{e}} [\sum_{c=t}^{H} r_{t} | s_{c}] \right) g_{A_{c} | S_{c}} \right]. \end{split}$$

Except for the third line, the proof is almost the same as that of Kallus and Uehara (2020a, Theorem 2). The third line is proved by

$$\begin{split} &\sum_{c=1}^{H} \int \sum_{t=1}^{H} r_{t} \left\{ \prod_{k \neq c}^{t} p(s_{k} | a_{k-1}, s_{k-1}) p_{\pi_{k}^{e}}(a_{k} | s_{k}) p(r_{k} | a_{k}, s_{k}) \right\} p(s_{c} | \tau(a_{c-1}, s_{c-1}), s_{c-1}) \\ &\times \nabla \pi^{b}(a_{c} | s_{c}; \theta) p(r_{c} | \tau(a_{c}, s_{c}), s_{c}) \mathrm{d}\mu(j_{r_{H}}) \\ &= \sum_{c=1}^{H} \int \sum_{t=c}^{H} r_{t} \left\{ \prod_{k \neq c}^{t} p(s_{k} | a_{k-1}, s_{k-1}) p_{\pi_{k}^{e}}(a_{k} | s_{k}) p(r_{k} | a_{k}, s_{k}) \right\} p(s_{c} | \tau(a_{c-1}, s_{c-1}), s_{c-1}) \\ &\times \pi^{b}(a_{c} | s_{c}) p(r_{c} | \tau(a_{c}, s_{c}), s_{c}) g_{A_{c} | S_{c}} \mathrm{d}\mu(j_{r_{H}}) \\ &= \mathbb{E}_{\pi^{b}} \left[\prod_{k=1}^{t-1} \eta_{k} \mathbb{E}_{\pi^{e}} \left[\sum_{c=t}^{H} r_{t} | s_{c}, \tau(s_{c}, a_{c}) \right] q_{A_{c} | S_{c}} \right] \\ &= \mathbb{E}_{\pi^{b}} \left[\prod_{k=1}^{t-1} \eta_{k} \left(\mathbb{E}_{\pi^{e}} \left[\sum_{c=t}^{H} r_{t} | s_{c}, \tau(s_{c}, a_{c}) \right] - \mathbb{E}_{\pi^{e}} \left[\sum_{c=t}^{H} r_{t} | s_{c} \right] \right) g_{A_{c} | S_{c}} \right]. \end{split}$$

Then,

$$\nabla_{\theta} \mathbb{E}_{\pi^{e}} \left[\sum_{t=1}^{H} r_{t} \right] = \sum_{c=1}^{H} \left\{ \mathbb{E}_{\pi^{b}} \left[\left(\mathbb{E}_{\pi^{c}} [r_{c} | s_{1}] - \mathbb{E}_{\pi^{c}} [r_{c}] \right) g \right] + \mathbb{E}_{\pi^{b}} \left[\mu_{c}(s_{c}, a_{c}) (r_{c} - \mathbb{E}[r_{c} | s_{c}, a_{c}]) g \right] \right. \\ + \mathbb{E}_{\pi^{b}} \left[\mu_{c} \left(\left[\mathbb{E}_{\pi^{c}} \left[\sum_{t=c+1}^{H} r_{t} | s_{c+1} \right] - \mathbb{E}_{\pi^{c}} \left[\sum_{t=c+1}^{H} r_{t} | s_{c}, a_{c} \right] \right) g \right] \\ + \mathbb{E}_{\pi^{b}} \left[\mu_{c-1} \left(\mathbb{E}_{\pi^{c}} \left[\sum_{c=t}^{H} r_{t} | s_{c}, \tau(s_{c}, a_{c}) \right] - \mathbb{E}_{\pi^{c}} \left[\sum_{c=t}^{H} r_{t} | s_{c} \right] \right) g \right] \right\}.$$

Here, note that we define $\mathbb{E}_{\pi^b}[\prod_{k=1}^t \eta_k \mid s_t, a_t] = \mu_t$. Then, $\nabla_{\theta} \mathbf{E}_{\pi^e}[\sum_{t=1}^H r_t]$ is equal to

$$\begin{split} &= \mathbb{E}\left[\left[-J + \sum_{c=1}^{H} \left\{\mu_{c} \left\{r_{c} - \mathbb{E}_{\pi_{e}}[\sum_{t=c}^{H} r_{t} | s_{c}, a_{c}]\right\} + \mu_{c-1} \sum_{t=c}^{H} \mathbb{E}_{\pi_{e}}[r_{t} | s_{c}]\right\}\right] g(\mathcal{J}_{r_{H}})\right] \\ &+ \mathbb{E}\left[\mu_{c-1} \left(\mathbb{E}_{\pi^{e}}[\sum_{c=t}^{H} r_{t} | s_{c}, \tau(s_{c}, a_{c})] - \mathbb{E}_{\pi^{e}}[\sum_{c=t}^{H} r_{t} | s_{c}]\right) g\right] \\ &= \mathbb{E}\left[\left[-J + \sum_{c=1}^{H} \left\{\mu_{c} \left\{r_{c} - \mathbb{E}_{\pi_{e}}[\sum_{t=c}^{H} r_{t} | s_{c}, s_{c}]\right\} + \mu_{c-1} \sum_{t=c}^{H} \mathbb{E}_{\pi_{e}}[r_{t} | s_{c}, \tau(a_{c}, s_{c})]\right\}\right] g(\mathcal{J}_{r_{H}})\right]. \end{split}$$

Then, as in the proof of Theorem 2 via the projection onto the tangent space, the EIF becomes

$$-J(\pi^{e}) + \sum_{c=1}^{H} \mu_{c} \{r_{c} - q_{c}(s_{c}, a_{c})\} + \mu_{c-1} q_{c}(s_{c}, \tau(a_{c}, s_{c})).$$

In addition, as in the proof of Theorem 2, the efficiency bound is

$$\sum_{c=0}^{H} \mathbb{E}[\mu_c^2(S_c, A_c) \text{var}[R_c + q_{c+1}(S_c, \tau(A_c, S_c)) \mid S_c, A_c]].$$

Order of the efficiency bound First, we observe

$$\operatorname{var}[R_c + q_{c+1}(S_{c+1}, \tau(A_{c+1}, S_{c+1})) \mid S_c = s_c, A_c = a_c] = \operatorname{var}_{\pi^e}[r_c + q_{c+1}(s_{c+1}, a_{c+1}) \mid s_c, a_c].$$
 (14)

This is proved by

$$\begin{aligned} & \operatorname{var}[R_c + q_{c+1}(S_{c+1}, \tau(A_{c+1}, S_{c+1})) \mid S_c = s_c, A_c = a_c] \\ & = \int \{r_c + q_{c+1}(s_{c+1}, \tau(a_{c+1}, s_{c+1})) - q_c(s_c, a_c)\}^2 p(r_c \mid s_c, a_c) p(s_{c+1} \mid s_c, a_c) \pi^{\mathsf{b}}(a_{c+1} \mid s_{c+1}) \operatorname{d}(r_c, s_{c+1}, a_{c+1}) \\ & = \int \{r_c + q_{c+1}(s_{c+1}, u_{c+1})) - q_c(s_c, a_c)\}^2 p(r_c \mid s_c, a_c) p(s_{c+1} \mid s_c, a_c) \times \\ & \pi^{\mathsf{b}}(\tilde{\tau}(u_{c+1}, s_{c+1}) \mid s_{c+1}) \tilde{\tau}'(u_{c+1}, s_{c+1}) \operatorname{d}(r_c, s_{c+1}, u_{c+1}) \\ & = \int \{r_c + q_{c+1}(s_{c+1}, u_{c+1})) - q_c(s_c, a_c)\}^2 p(r_c \mid s_c, a_c) p(s_{c+1} \mid s_c, a_c) \pi^{\mathsf{e}}(u_{c+1} \mid s_{c+1}) \operatorname{d}(r_c, s_{c+1}, u_{c+1}) \\ & = \operatorname{var}_{\pi^{\mathsf{e}}}[r_c + q_{c+1}(s_{c+1}, a_{c+1})) \mid s_c, a_c]. \end{aligned}$$

Then, we have

$$\sum_{c=0}^{H} \mathbb{E}[\mu_{c}^{2}(S_{c}, A_{c}) \operatorname{var}[R_{c} + q_{c+1}(S_{c}, \tau(A_{c+1}, S_{c+1})) \mid S_{c}, A_{c}]]$$

$$\leq CC' \sum_{c=0}^{H} \mathbb{E}_{\pi^{e}}[\operatorname{var}_{\pi^{b}}[r_{c} + q_{c+1}(s_{c+1}, \tau(a_{c+1}, s_{c+1})) \mid s_{c}, a_{c}]]$$

$$= CC' \sum_{c=0}^{H} \mathbb{E}_{\pi^{e}}[\operatorname{var}_{\pi^{e}}[r_{c} + q_{c+1}(a_{c+1}, a_{c+1})) \mid s_{c}, a_{c}]]$$

$$= CC' \sum_{c=0}^{H} \mathbb{E}_{\pi^{e}}[\{r_{c} + q_{c+1}(s_{c+1}, a_{c+1}) - q_{c}(s_{c}, a_{c})\}^{2}]$$

$$= CC' \mathbb{E}_{\pi^{e}}[\{\sum_{c=0}^{H} r_{c} + q_{c+1}(s_{c+1}, a_{c+1}) - q_{c}(s_{c}, a_{c})\}^{2}] = CC' \mathbb{E}_{\pi^{e}}[\{\sum_{c=1}^{H} r_{c}\} - J\}^{2}]$$

$$\leq CC' R_{\max}^{2} H^{2}.$$

From the first line to the second line, we use $\mu_c \leq CC'$. From the second line to the third line, we use Eq. (14). From the fourth line to the fifth line, we use a fact that the cross-term is equal to 0. \Box

Remark 9. Noting

$$\mathbb{E}[\operatorname{var}[f(Z)|X]] = \operatorname{var}[\mathbb{E}[f(Z)|X,Y]|X] + \mathbb{E}[\operatorname{var}[f(Z)|X,Y]|X],$$

for random variables X, Y, Z, the difference regarding the efficiency bound between the above and that of the pre-specified evaluation policy is

$$\sum_{t=1}^{H} \mathbb{E}[\mu_t^2(S_t, A_t) \operatorname{var}[R_t + q_{t+1}(S_{t+1}, \tau(S_{t+1}, A_{t+1})) \mid S_{t+1}, R_t, S_t, A_t]]$$

$$= \sum_{t=1}^{H} \mathbb{E}[\mu_t^2(S_t, A_t) \operatorname{var}[q_{t+1}(S_{t+1}, \tau(S_{t+1}, A_{t+1})) \mid S_{t+1}]].$$

Proof of Theorem 4. For simplicitly, we consider the case K=2. First, we prove

$$\mathbb{P}_{u_{1}}[\phi(\hat{w}^{(1)}, \hat{\pi}^{b(1)}, \hat{q}^{(1)})|\mathcal{L}_{1}] - \mathbb{P}_{u_{1}}[\phi(w, \pi^{b}, q)|\mathcal{L}_{1}]
= \mathbb{G}_{u_{1}}[\phi(\hat{w}^{(1)}, \hat{\pi}^{b(1)}, \hat{q}^{(1)}) - \phi(w, \pi^{b}, q)]
+ \mathbb{E}[\phi(\hat{w}^{(1)}, \hat{\pi}^{b(1)}, \hat{q}^{(1)})|\mathcal{L}_{1}] - \mathbb{E}[\phi(w, \pi^{b}, q)|\mathcal{L}_{1}]
= o_{p}(n^{-1/2}).$$
(15)

From now on, we drop an index (1) for simplicity. In addition, we use a notation of $c_t(s_t)$, $\hat{c}_t(s_t)$.

Part1: Eq. (16) is $o_p(1/\sqrt{n})$ First, we consider controlling the following term:

$$\mathbb{E}\left[\sum_{t=1}^{H} \hat{w}_{t}(S_{t}) \hat{\eta}_{t}(S_{t}, A_{t}) \{R_{t} - \hat{q}_{t}(S_{t}, A_{t})\} + \hat{w}_{t-1}(S_{t-1}) \hat{\eta}_{t-1}(S_{t-1}, A_{t-1}) \hat{v}_{t}(S_{t}) | \mathcal{L}_{1}\right] - J.$$

By some algebra, we have

$$\mathbb{E}\left[\sum_{t=1}^{H} \hat{w}_{t}(S_{t}) \hat{\eta}_{t}(S_{t}, A_{t}) \{R_{t} - \hat{q}_{t}(S_{t}, A_{t})\} + \hat{w}_{t-1}(S_{t-1}) \hat{\eta}_{t-1}(S_{t-1}, A_{t-1}) \hat{v}_{t}(S_{t}) | \mathcal{L}_{1}\right] - J$$

$$= \mathbb{E}\left[\sum_{t=1}^{H} \hat{w}_{t}(S_{t})\hat{\eta}_{t}(S_{t}, A_{t})\{R_{t} - \hat{q}_{t}(S_{t}, A_{t})\}\right] \\
+ \mathbb{E}_{\pi^{e}}\left[\hat{q}_{t}(s_{t})|A_{t}\right]|\mathcal{L}_{1}\right] - J + \mathbb{E}\left[\sum_{t=1}^{H} \hat{w}_{t-1}(S_{t-1})\hat{\eta}_{t-1}(S_{t-1}, A_{t-1})\{\hat{v}_{t}(S_{t}) - \mathbb{E}_{\pi^{e}}[\hat{q}_{t}(s_{t}, A_{t})|A_{t}]\}|\mathcal{L}_{1}\right] \\
= \mathbb{E}\left[\sum_{t=1}^{H} \{\hat{w}_{t}(S_{t})\hat{\eta}_{t}(S_{t}, A_{t}) - w_{t}(S_{t})\eta_{t}(S_{t}, A_{t})\}\{\hat{q}_{t}(S_{t}, A_{t}) - q_{t}(S_{t}, A_{t})\}|\mathcal{L}_{1}\right] \\
+ \mathbb{E}\left[\sum_{t=1}^{H} \{\hat{w}_{t-1}(S_{t-1})\hat{\eta}_{t-1}(S_{t-1}, A_{t-1}) - w_{t}(S_{t-1})\eta_{t-1}(S_{t-1}, A_{t-1})\}\{-\hat{v}_{t}(S_{t}) + v_{t}(S_{t})\}|\mathcal{L}_{1}\right] \\
+ \mathbb{E}\left[\sum_{t=1}^{H} \mu_{t}(S_{t}, A_{t})\{-\hat{q}_{t}(S_{t}, A_{t}) + q_{t}(S_{t}, A_{t})\} + \mu_{t-1}(S_{t-1}, A_{t-1})\{\hat{v}_{t}(S_{t}) - v_{t}(S_{t})\}|\mathcal{L}_{1}\right] \\
+ \mathbb{E}\left[\sum_{t=1}^{H} \{\hat{\mu}_{t}(S_{t}, A_{t}) - \mu_{t}(S_{t}, A_{t})\}\{R_{t} - q_{t}(S_{t}, A_{t}) + v_{t+1}(S_{t+1})\}\}|\mathcal{L}_{1}\right] \\
+ \mathbb{E}\left[\sum_{t=1}^{H} \hat{w}_{t-1}(S_{t-1})\hat{\eta}_{t-1}(S_{t-1}, A_{t-1})\int \{\hat{\pi}_{t}^{e}(a_{t}|S_{t}) - \pi_{t}^{e}(a_{t}|S_{t})\}\hat{q}_{t}(a_{t}, S_{t})da_{t}|\mathcal{L}_{1}\right]. \tag{19}$$

Here, the terms from Eq. (17) to Eq. (19) are the same as the decomposition in the same as Theorem 2 (Kallus and Uehara, 2020a). Since Eq. (19) and Eq. (18) are equal to 0, the above is equal to

$$\mathbb{E}\left[\sum_{t=1}^{H} \{\hat{w}_{t}(S_{t})\hat{\eta}_{t}(S_{t}, A_{t}) - w_{t}(S_{t})\eta_{t}(S_{t}, A_{t})\} \{\hat{q}_{t}(S_{t}, A_{t}) - q_{t}(S_{t}, A_{t})\} | \mathcal{L}_{1}\right] \\
+ \mathbb{E}\left[\sum_{t=1}^{H} \{\hat{w}_{t-1}(S_{t-1})\hat{\eta}_{t-1}(S_{t-1}, A_{t-1}) - w_{t}(S_{t-1})\eta_{t-1}(S_{t-1}, A_{t-1})\} \{-\hat{v}_{t}(S_{t}) + v_{t}(S_{t})\} | \mathcal{L}_{1}\right] \\
+ \mathbb{E}\left[\sum_{t=1}^{H} \{\hat{w}_{t-1}(S_{t-1})\hat{\eta}_{t-1}(S_{t-1}, A_{t-1}) - w_{t-1}(S_{t-1})\eta_{t-1}(S_{t-1}, A_{t-1})\} \int \{\hat{\pi}_{t}^{e}(a_{t}|S_{t}) - \pi_{t}^{e}(a_{t}|S_{t})\} \hat{q}_{t}(a_{t}, S_{t}) da_{t} | \mathcal{L}_{1}\right] \\
+ \mathbb{E}\left[\sum_{t=1}^{H} w_{t-1}(S_{t-1})\eta_{t-1}(S_{t-1}, A_{t-1}) \int \{\hat{\pi}_{t}^{e}(a_{t}|S_{t}) - \pi_{t}^{e}(a_{t}|S_{t})\} \hat{q}_{t}(a_{t}, S_{t}) da_{t} | \mathcal{L}_{1}\right] \\
= \alpha_{1}\beta + \alpha_{2}\beta + (\alpha_{1} + \alpha_{2})\alpha_{2} + \mathbb{E}\left[\sum_{t=1}^{H} w_{t}(S_{t}) \int \{\hat{\pi}_{t}^{e}(a_{t}|S_{t}) - \pi_{t}^{e}(a_{t}|S_{t})\} \hat{q}_{t}(a_{t}, S_{t}) da_{t} | \mathcal{L}_{1}\right].$$

Here, we use $\|\hat{v}_t - v_t\|_2 \le C \|\hat{q}_t - q_t\|_2 \lesssim \beta$. Next, we also use a fact $\|\hat{c}_t(S_t) - c_t(S_t)\|_2^2 = \alpha_2^2$ since

$$\|\hat{c}_t(S_t) - c_t(S_t)\|_2^2 = \int \left\{ u(a_t)\hat{\pi}_t^b(a_t|s_t) - \pi_t^b(a_t|s_t)u(a_t) \right\}^2 p_{\pi^b}(s_t) d\mu(a_t, s_t)$$

$$\lessapprox \|\hat{\pi}_t^b(A_t|S_t) - \pi_t^b(A_t|S_t)\|_2^2.$$

Therefore, noting what we want to control

$$\mathbb{E}\left[\sum_{t=1}^{H} \hat{w}_{t}(S_{t}) \hat{\eta}_{t}(S_{t}, A_{t}) \{R_{t} - \hat{v}_{t}(S_{t})\} + \hat{w}_{t-1}(S_{t-1}) \hat{\eta}_{t-1}(S_{t-1}, A_{t-1}) \hat{v}_{t}(S_{t}) | \mathcal{L}_{1}\right] - J,$$

the following holds:

$$\mathbb{E}\left[\sum_{t=1}^{H} \hat{w}_{t}(S_{t}) \hat{\eta}_{t}(S_{t}, A_{t}) \{R_{t} - \hat{v}_{t}(S_{t})\} + \hat{w}_{t-1}(S_{t-1}) \hat{\eta}_{t-1}(S_{t-1}, A_{t-1}) \hat{v}_{t}(S_{t}) | \mathcal{L}_{1}\right] - J$$

$$= \mathbb{E}\left[\sum_{t=1}^{H} \hat{w}_{t}(S_{t}) \hat{\eta}_{t}(S_{t}, A_{t}) \{-\hat{v}_{t}(S_{t}) + \hat{q}_{t}(S_{t}, A_{t})\} + w_{t}(S_{t}) \int \{\hat{\pi}_{t}^{e}(a_{t}|S_{t}) - \pi_{t}^{e}(a_{t}|S_{t})\} \hat{q}_{t}(a_{t}, S_{t}) da_{t} | \mathcal{L}_{1}\right] + \alpha_{1}\beta + \alpha_{2}\beta + \alpha_{1}\alpha_{2} + \alpha_{2}^{2}.$$

Then, the main term in the above is further expanded as follows.

$$\mathbb{E}\left[\sum_{t=1}^{H} w_{t}(S_{t})\hat{\eta}_{t}(S_{t}, A_{t})\{-\hat{v}_{t}(S_{t}) + \hat{q}_{t}(S_{t}, A_{t})\}|\mathcal{L}_{1}\right] + \mathbb{E}\left[\sum_{t=1}^{H} w_{t}(S_{t}) \int \{\hat{\pi}_{t}^{e}(a_{t}|S_{t}) - \pi_{t}^{e}(a_{t}|S_{t})\}\hat{q}_{t}(a_{t}, S_{t})da_{t}|\mathcal{L}_{1}\right] \\
= \mathbb{E}\left[\sum_{t=1}^{H} w_{t}(S_{t})\hat{c}_{t}(S_{t})u_{t}(A_{t})\{-\hat{c}_{t}(S_{t}) \int u_{t}(a_{t})\hat{q}_{t}(a_{t}, S_{t})\hat{\pi}_{t}^{b}(a_{t}|S_{t})da_{t} + \hat{q}_{t}(S_{t}, A_{t})\}|\mathcal{L}_{1}\right] \\
+ \mathbb{E}\left[\sum_{t=1}^{H} w_{t}(S_{t}) \int u_{t}(a_{t})\{\hat{c}_{t}(S_{t})\hat{\pi}_{t}^{b}(a_{t}|S_{t}) - c_{t}(S_{t})\pi_{t}^{b}(a_{t}|S_{t})\}\hat{q}_{t}(a_{t}, S_{t})da_{t}|\mathcal{L}_{1}\right] \\
= \mathbb{E}\left[\sum_{t=1}^{H} w_{t}(S_{t}) \int \hat{c}_{t}(S_{t})u_{t}(g_{t})\pi_{t}^{b}(g_{t}|S_{t})\{-\hat{c}_{t}(S_{t}) \int u_{t}(a_{t})\hat{q}_{t}(a_{t}, S_{t})\hat{\pi}_{t}^{b}(a_{t}|S_{t})da_{t} + \hat{q}_{t}(g_{t}, S_{t})\}dg_{t}|\mathcal{L}_{1}\right] \\
+ \mathbb{E}\left[\sum_{t=1}^{H} w_{t}(S_{t}) \int u_{t}(a_{t})\{\hat{c}_{t}(S_{t})\hat{\pi}_{t}^{b}(a_{t}|S_{t}) - c_{t}(S_{t})\pi_{t}^{b}(a_{t}|S_{t})\}\hat{q}_{t}(a_{t}, S_{t})da_{t}\right]|\mathcal{L}_{1}\right].$$

Noting we have $1/c_t(S_t) = \int u_t(g_t)\pi_t^b(g_t|S_t)dg_t$, the following holds:

$$\begin{split} &= \mathbb{E}[\sum_{t=1}^{H} w_{t}(S_{t}) \int \{\hat{c}_{t}(S_{t}) - c_{t}(S_{t})\} u_{t}(g_{t}) \pi_{t}^{b}(g_{t}|S_{t}) \hat{q}_{t}(g_{t}, S_{t}) dg_{t}|\mathcal{L}_{1}] + \\ &+ \mathbb{E}[\sum_{t=1}^{H} w_{t}(S_{t}) \{-\hat{c}_{t}^{2}(S_{t})/c_{t}(S_{t}) + \hat{c}_{t}(S_{t})\} \int u_{t}(a_{t}) \hat{\pi}_{t}^{b}(a_{t}|S_{t}) \hat{q}_{t}(a_{t}, S_{t}) da_{t}|\mathcal{L}_{1}] \\ &= \mathbb{E}[\sum_{t=1}^{H} w_{t}(S_{t}) \int \{\hat{c}_{t}(S_{t}) - c_{t}(S_{t})\} u_{t}(g_{t}) \{\pi_{t}^{b}(g_{t}|S_{t}) - \hat{\pi}_{t}^{b}(g_{t}|S_{t})\} \hat{q}_{t}(g_{t}, S_{t}) dg_{t}|\mathcal{L}_{1}] + \\ &+ \mathbb{E}[\sum_{t=1}^{H} w_{t}(S_{t}) \{-\hat{c}_{t}^{2}(S_{t})/c_{t}(S_{t}) + \hat{c}_{t}(S_{t}) + \hat{c}_{t}(S_{t}) - c_{t}(S_{t})\} \int u_{t}(a_{t}) \hat{\pi}_{t}^{b}(a_{t}|S_{t}) \hat{q}_{t}(a_{t}, S_{t}) da_{t}|\mathcal{L}_{1}] \\ &= \mathbb{E}[\sum_{t=1}^{H} w_{t}(S_{t}) \int \{\hat{c}_{t}(S_{t}) - c_{t}(S_{t})\} u_{t}(g_{t}) \{\pi_{t}^{b}(g_{t}|S_{t}) - \hat{\pi}_{t}^{b}(g_{t}|S_{t})\} \hat{q}_{t}(g_{t}, S_{t}) dg_{t}|\mathcal{L}_{1}] + \\ &+ \mathbb{E}[\sum_{t=1}^{H} w_{t}(S_{t}) \{-\hat{c}_{t}(S_{t}) + c_{t}(S_{t})\}^{2}/c_{t}(S_{t}) \int u_{t}(a_{t}) \hat{\pi}_{t}^{b}(a_{t}|S_{t}) \hat{q}_{t}(a_{t}, S_{t}) da_{t}|\mathcal{L}_{1}] \\ &\lesssim \|\pi_{t}^{b}(A_{t}|S_{t}) - \hat{\pi}_{t}^{b}(A_{t}|S_{t})\|_{2} \lesssim \alpha_{2}^{2}. \end{split}$$

This concludes that Eq. (16) is $\alpha_1\beta + \alpha_2\beta + \alpha_1\alpha_2 + \alpha_2^2$. Under the assumption for the convergence rates, this is equal to $o_p(n^{-1/2})$.

Part 2:Eq. (15) is $o_p(1/\sqrt{n})$ Following Kallus and Uehara (2020a), this is proved if the following is proved:

$$\mathbb{E}[\{\phi(\hat{w}^{(1)}, \hat{\pi}^{b(1)}, \hat{q}^{(1)}) - \phi(w, \pi^b, q)\}^2 | \mathcal{L}_1] = o_p(1).$$

This is proved as in the Part 1 using the same decomposition.

Final part

$$\begin{split} & \mathbb{P}_{u_1}[\phi(\hat{w}^{(1)}, \hat{\pi}^{b(1)}, \hat{q}^{(1)}) | \mathcal{L}_1] + \mathbb{E}_{u_2}[\phi(\hat{w}^{(2)}, \hat{\pi}^{b(2)}, \hat{q}^{(2)}) | \mathcal{L}_2] \\ & = \mathbb{P}_n[\phi(w, \pi^b, q)] + o_n(n^{-1/2}). \end{split}$$

Then, CLT concludes the proof.

Proof of Theorem 5. As we have ever seen,

$$\hat{J}_{\text{TI}} - \mathbb{P}_n[\phi(w^{\dagger}, \pi^b, q^{\dagger})] = \alpha_1 \beta + \alpha_2 \beta + \alpha_1 \alpha_2 + \alpha_2^2 + o_p(n^{-1/2}).$$

Under the assumption, the above is equal to $o_p(1)$. In addition, the mean of $\mathbb{P}_n[\phi(w^{\dagger}, \pi^b, q^{\dagger})]$ is J. Therefore, the statement holds from the law of large numbers.

Proof of Theorem 6. For simplicity, we consider the case K=2. First, we prove

$$\mathbb{P}_{u_{1}}[\phi(\hat{w}^{(1)}, \hat{\pi}^{b(1)}, \hat{q}^{(1)})|\mathcal{L}_{1}] - \mathbb{P}_{u_{1}}[\phi(w, \pi^{b}, q)|\mathcal{L}_{1}]
= \mathbb{G}_{u_{1}}[\phi(\hat{w}^{(1)}, \hat{\pi}^{b(1)}, \hat{q}^{(1)}) - \phi(w, \pi^{b}, q)]
+ \mathbb{E}[\phi(\hat{w}^{(1)}, \hat{\pi}^{b(1)}, \hat{q}^{(1)})|\mathcal{L}_{1}] - \mathbb{E}[\phi(w, \pi^{b}, q)|\mathcal{L}_{1}]
= o_{p}(n^{-1/2}).$$
(20)

Part1: Eq. (21) is $o_p(n^{-1/2})$

$$\begin{split} & \operatorname{Eq.} \left(21\right) = \mathbb{E}[\phi(\hat{w}^{(1)}, \hat{\pi}^{b(1)}, \hat{q}^{(1)}) | \mathcal{L}_{1}] - \mathbb{E}[\phi(w, \pi^{b}, q) | \mathcal{L}_{1}] \\ & = \mathbb{E}[\sum_{t=1}^{H} \{\hat{\mu}_{t}(S_{t}, A_{t}) - \mu_{t}(S_{t}, A_{t})\} \{ -\hat{q}_{t}(S_{t}, A_{t}) + q_{t}(S_{t}, A_{t}) \} \\ & + \mathbb{E}[\sum_{t=1}^{H} \{\hat{\mu}_{t-1}(S_{t-1}, A_{t-1}) - \mu_{t-1}(S_{t-1}, A_{t-1})\} \{ -\hat{q}_{t}(S_{t}, \tau(S_{t}, A_{t})) + q_{t}(S_{t}, \tau(S_{t}, A_{t})) \} | \mathcal{L}_{1}] \\ & + \mathbb{E}[\sum_{t=1}^{H} \mu_{t}(S_{t}, A_{t}) \{ -\hat{q}_{t}(S_{t}, A_{t}) + q_{t}(S_{t}, A_{t}) \} + \mu_{t-1}(S_{t-1}, A_{t-1}) \{ \hat{q}_{t}(S_{t}, \tau(S_{t}, A_{t})) - q_{t}(S_{t}, \tau(S_{t}, A_{t})) \} | \mathcal{L}_{1}] \\ & + \mathbb{E}[\sum_{t=1}^{H} \{ \hat{\mu}_{t}(S_{t}, A_{t}) - \mu_{t}(S_{t}, A_{t}) \} \{ R_{t} - q_{t}(S_{t}, A_{t}) + q_{t+1}(S_{t+1}, \tau(S_{t+1}, A_{t+1})) | \mathcal{L}_{1}] \\ & = \mathbb{E}[\sum_{t=1}^{H} \{ \hat{\mu}_{t}(S_{t}, A_{t}) - \mu_{t}(S_{t}, A_{t}) \} \{ -\hat{q}_{t}(S_{t}, A_{t}) + q_{t}(S_{t}, A_{t}) \} | \mathcal{L}_{1}] \\ & + \mathbb{E}[\sum_{t=1}^{H} \{ \hat{\mu}_{t-1}(S_{t-1}, A_{t-1}) - \mu_{t-1}(S_{t-1}, A_{t-1}) \} \{ -\hat{q}_{t}(S_{t}, \tau(S_{t}, A_{t})) + q_{t}(S_{t}, \tau(S_{t}, A_{t})) \} | \mathcal{L}_{1}] \\ & = \sum_{t=1}^{H} \|\hat{\mu}_{t}(S_{t}, A_{t}) - \mu_{t}(S_{t}, A_{t}) \|_{2} \| -\hat{q}_{t}(S_{t}, A_{t}) + q_{t}(S_{t}, T(S_{t}, A_{t})) + q_{t}(S_{t}, \tau(S_{t}, A_{t})) \|_{2} \\ & + \|\hat{\mu}_{t-1}(S_{t-1}, A_{t-1}) - \mu_{t-1}(S_{t-1}, A_{t-1}) \|_{2} \| -\hat{q}_{t}(S_{t}, \tau(S_{t}, A_{t})) + q_{t}(S_{t}, \tau(S_{t}, A_{t})) \|_{2} \\ & = (\alpha_{1} + \alpha_{2})\beta = o_{n}(n^{-1/2}). \end{split}$$

Here, we use

$$\mathbb{E}[\mu_k(S_k, A_k) f(S_k, A_k)] = \mathbb{E}[\prod_{t=1}^k \eta_t(S_t, A_t) f(S_k, A_k)] = \mathbb{E}[(\prod_{t=1}^{k-1} \eta_t(S_t, A_t)) \eta_k(S_k, A_k) f(S_k, A_k)]$$

$$= \mathbb{E}[(\prod_{t=1}^{k-1} \eta_t(S_t, A_t)) f(S_k, \tau_k(A_k, S_k))] = \mathbb{E}[\mu_{k-1}(S_{k-1}, A_{k-1}) f(S_k, \tau_k(A_k, S_k))],$$

and

$$\begin{aligned} \|\hat{q}(S,\tau(S,A)) - q(S,\tau(S,A))\|_{2}^{2} &= \int \{\hat{q}(s,\tau(s,a)) - q(s,\tau(s,a))\}^{2} \pi^{b}(a|s) p(s) d(s,a) \\ &= \int \{\hat{q}(s,a) - q(s,a)\}^{2} \pi^{e}(a|s) p(s) d(s,a) \\ &\leq C' \|\hat{q}(S,A) - q(S,A)\|_{2}^{2}. \end{aligned}$$

Part 2:Eq. (20) is $o_p(n^{-1/2})$ Following Kallus and Uehara (2020a), this is proved if the following is proved:

$$\mathbb{E}[\{\phi(\hat{w}^{(1)}, \hat{\pi}^{b(1)}, \hat{q}^{(1)}) - \phi(w, \pi^b, q)\}^2 | \mathcal{L}_1] = o_p(1).$$

This is proved as in the Part 1 using the same decomposition.

Final part

$$\mathbb{P}_{u_1}[\phi(\hat{w}^{(1)}, \hat{\pi}^{b(1)}, \hat{q}^{(1)}) | \mathcal{L}_1] + \mathbb{E}_{u_2}[\phi(\hat{w}, \hat{\pi}^{b(2)}, \hat{q}^{(2)}) | \mathcal{L}_2] \\
= \mathbb{P}_n[\phi(w, \pi^b, q)] + o_p(n^{-1/2}).$$

Then, CLT concludes the proof.

Proof of Theorem 7. From the proof of Theorem 6, we have

$$\hat{J}_{\text{TI1}} = \mathbb{P}_n[\phi(w^{\dagger}, \pi^{b\dagger}, q^{\dagger})] + (\alpha_1 + \alpha_2)\beta + o_p(n^{-1/2}).$$

From the law of large numbers, noting the expectation of $\mathbb{P}_n[\phi(w^{\dagger}, \pi^{b\dagger}, q^{\dagger})]$ is J under the assumption, the statement is concluded.

A.3 Proofs for Section 5

Proof of Theorem 8.

Calculation of derivatives under the nonparemtric NMDP. The entire regular parametric submodel corresponding an NMDP is

$$\{p_{\theta}(s_1)p_{\theta}(a_1|s_1)p_{\theta}(r_1|j_{a_1})p_{\theta}(s_2|j_{r_1})p_{\theta}(a_2|j_{s_2})p_{\theta}(r_2|j_{a_2})\cdots p_{\theta}(r_H|j_{a_H})\},$$

where it matches with the true pdf at $\theta = 0$. The score function of the model is decomposed as

$$g(j_{r_H}) = \sum_{k=1}^{H} \nabla \log p_{\theta}(s_k|j_{r_{k-1}}) + \sum_{k=1}^{H} \nabla \log p_{\theta}(a_k|j_{s_k}) + \sum_{k=1}^{H} \nabla \log p_{\theta}(r_k|j_{a_k})$$

$$= \sum_{k=1}^{H} g_{S_{k}|\mathcal{I}_{r_{k-1}}} + \sum_{k=1}^{H} g_{A_{k}|\mathcal{I}_{s_{k}}} + \sum_{k=1}^{H} g_{R_{k}|\mathcal{I}_{a_{k}}}.$$

Then, we have

$$\nabla_{\theta} \mathbf{E}_{\pi^e} \left[\sum_{t=1}^{H} r_t \right] = \mathbb{E}[\{-J + \sum_{c=1}^{H} \lambda_c \{r_c - v_c(\mathcal{J}_{s_c})\} + \lambda_{c-1} v_c(\mathcal{J}_{s_c})\} g(\mathcal{J}_{r_T})].$$

This is proved by

$$\begin{split} &\nabla_{\theta} \mathbb{E}_{\pi^{e}} \left[\sum_{t=1}^{H} r_{t} \right] = \nabla_{\theta} \left[\int \sum_{t=1}^{H} r_{t} \left\{ \prod_{k=1}^{H} p_{\theta}(s_{k} | j_{r_{k-1}}) \pi^{e}(a_{k} | j_{s_{k}}; \theta) p_{\theta}(r_{k} | j_{a_{k}}) \right\} d\mu(j_{r_{H}}) \right] \\ &= \sum_{c=1}^{H} \left\{ \mathbb{E}_{\pi^{e}} \left[\left\{ \mathbb{E}_{\pi^{e}}(r_{c} | s_{1}) - \mathbb{E}_{\pi^{e}}(r_{c}) \right\} g_{s_{1}} \right] + \mathbb{E}_{\pi^{e}} \left[\left\{ r_{c} - \mathbb{E}_{\pi^{e}}(R_{c} | \mathcal{J}_{a_{c}}) \right\} g_{r_{c} | \mathcal{J}_{a_{c}}} \right] \right. \\ &+ \mathbb{E}_{\pi^{e}} \left[\left(\mathbb{E}_{\pi^{e}} \left[\sum_{t=c+1}^{H} r_{t} | \mathcal{J}_{s_{c+1}} \right] - \mathbb{E}_{\pi^{e}} \left[\sum_{t=c+1}^{H} r_{t} | \mathcal{J}_{a_{c}} \right] \right) g_{S_{c+1} | \mathcal{J}_{r_{c}}} \right] \\ &+ \mathbb{E} \left[\lambda_{c} \left\{ \mathbb{E}_{\pi^{e}} \left[\sum_{t=c}^{H} r_{t} | \mathcal{J}_{a_{c}} \right] - \mathbb{E}_{\pi^{e}} \left[\sum_{t=c}^{H} r_{t} | \mathcal{J}_{s_{c}} \right] \right\} g_{A_{c} | \mathcal{J}_{s_{c}}} \right] \right\}. \end{split}$$

Note that except for the third term, the proof is the same as Theorem 2. The third term is calculated by

$$\begin{split} &\sum_{m=1}^{H} \int \sum_{t=1}^{H} \left\{ r_{t} \prod_{k=1, k \neq m}^{t} p_{\theta}(s_{k}|j_{r_{k-1}}) \pi^{e}(a_{k}|j_{s_{k}}; \theta) p_{\theta}(r_{k}|j_{a_{k}}) \right\} \times \\ &\pi_{m}^{e} \left\{ g_{A_{m}|\mathcal{I}_{s_{m}}} - \left[\int g_{A_{m}|H_{S_{m}}} \pi^{e}(a_{m} \mid j_{s_{m}}) \mathrm{d}(a_{m}) \right] \right\} \mathrm{d}\mu(j_{r_{H}}) \\ &= \sum_{m=1}^{H} \mathbb{E}_{\pi^{e}} \left[\sum_{t=m}^{H} r_{t} g_{A_{m}|\mathcal{I}_{s_{m}}} - \sum_{t=m}^{H} r_{t} \mathbb{E}_{\pi^{e}} \left[g_{A_{m}|H_{s_{m}}} \mid j_{s_{m}} \right] \right] \\ &= \sum_{m=1}^{H} \mathbb{E}_{\pi^{e}} \left[\mathbb{E}_{\pi^{e}} \left[\sum_{t=m}^{H} r_{t} \mid j_{A_{m}} \right] g_{A_{m}|\mathcal{I}_{s_{m}}} - \mathbb{E}_{\pi^{e}} \left[\sum_{t=m}^{H} r_{t} \mid j_{s_{m}} \right] \mathbb{E}_{\pi^{e}} \left[g_{A_{m}|H_{s_{m}}} \mid j_{s_{m}} \right] \right] \\ &= \mathbb{E}[\lambda_{c} \left\{ \mathbb{E}_{\pi^{e}} \left[\sum_{t=n}^{H} r_{t} \mid \mathcal{I}_{a_{c}} \right] - \mathbb{E}_{\pi^{e}} \left[\sum_{t=n}^{H} r_{t} \mid \mathcal{I}_{s_{c}} \right] \right\} g_{A_{c}|\mathcal{I}_{s_{c}}} \right]. \end{split}$$

Then, $\nabla_{\theta} \mathbf{E}_{\pi^e} \left[\sum_{t=1}^{H} r_t \right]$ is equal to

$$\begin{split} &\sum_{c=1}^{H} \{\mathbb{E}_{\pi^{e}} [\{\mathbb{E}_{\pi^{e}} (r_{c} | s_{1}) - \mathbb{E}_{\pi^{e}} (r_{c})\} g(\mathcal{J}_{r_{H}})] + \mathbb{E}_{\pi^{e}} [\{r_{c} - \mathbb{E}_{\pi^{e}} (r_{c} | \mathcal{J}_{a_{c}})\} g(\mathcal{J}_{r_{H}})] \\ &+ \mathbb{E}_{\pi^{e}} \left[\left(\mathbb{E}_{\pi^{e}} \left[\sum_{t=c+1}^{H} r_{t} | \mathcal{J}_{s_{c}+1} \right] - \mathbb{E}_{\pi^{e}} \left[\sum_{t=c+1}^{H} r_{t} | \mathcal{J}_{a_{c}} \right] \right) g(\mathcal{J}_{r_{H}}) \right] \} \\ &+ \mathbb{E} [\lambda_{c} \{\mathbb{E} [\sum_{t=c}^{H} r_{t} | \mathcal{J}_{a_{c}}] - \mathbb{E} [\sum_{t=c}^{H} r_{t} | \mathcal{J}_{s_{c}}] \} g(\mathcal{J}_{r_{H}})] \} \\ &= \mathbb{E} \left(\left[-J + \sum_{c=1}^{H} \left\{ \lambda_{c} r_{c} - \lambda_{c} \sum_{t=c}^{H} \mathbb{E}_{\pi_{e}} (r_{t} | \mathcal{J}_{a_{c}}) + \lambda_{c-1} \sum_{t=c}^{H} \mathbb{E}_{\pi_{e}} (r_{t} | \mathcal{J}_{s_{c}}) \right\} \right] g(\mathcal{J}_{r_{H}}) \right) + \end{split}$$

$$+ \mathbb{E}\left[\sum_{c=1}^{H} \left\{\lambda_{c} \sum_{t=c}^{H} \mathbb{E}_{\pi_{e}}(r_{t} | \mathcal{J}_{a_{c}}) - \lambda_{c} \mathbb{E}_{\pi_{e}}\left[\sum_{t=c}^{H} r_{t} | \mathcal{J}_{s_{c}}\right]\right\} g(\mathcal{J}_{r_{H}})\right]$$

$$= \mathbb{E}\left[\left\{-J + \sum_{c=1}^{H} \lambda_{c} \left\{r_{c} - v_{c}(\mathcal{J}_{s_{c}})\right\} + \lambda_{c-1} v_{c}(\mathcal{J}_{s_{c}})\right\} g(\mathcal{J}_{r_{H}})\right].$$

This concludes that the following is a derivative:

$$-J + \sum_{c=1}^{H} \lambda_c \{ r_c - v_c(j_{s_c}) \} + \lambda_{c-1} v_c(j_{s_c}).$$
 (22)

This concludes that ϕ is actually the EIF since the derivative is unique.

Calculation of the efficiency bound From the law of total variance, the efficiency bound is

$$\begin{aligned} & \operatorname{var}\left[\sum_{c=1}^{H} \lambda_c \{R_c - v_c(\mathcal{J}_{S_c})\} + \lambda_{c-1} v_c(\mathcal{J}_{S_c})\right] \\ & = \sum_{t=0}^{H} \mathbb{E}[\operatorname{var}[\mathbb{E}[\sum_{c=1}^{H} \lambda_c \{R_c + v_{c+1}(\mathcal{J}_{S_{c+1}}) - v_c(\mathcal{J}_{S_c})\} | \mathcal{J}_{S_{t+1}}] | \mathcal{J}_{S_t}]] \\ & = \sum_{t=0}^{H} \mathbb{E}[\operatorname{var}[\lambda_t \{R_t + v_{t+1}(\mathcal{J}_{S_{t+1}})\} \mid \mathcal{J}_{S_t}]] = \sum_{t=0}^{H} \mathbb{E}[\lambda_t^2 \operatorname{var}[R_t + v_{t+1}(\mathcal{J}_{S_{t+1}}) \mid \mathcal{J}_{S_t}]]. \end{aligned}$$

Proof of Theorem 9.

$$\begin{split} &\sum_{t=0}^{H} \mathcal{E}_{\pi_{b}} \left[\lambda_{t-1}^{2} \text{var} \left\{ \eta_{t} \{ r_{t} + v_{t+1}(j_{s_{t+1}}) \} \mid j_{s_{t}} \right\} \right] \\ &= \sum_{t=0}^{H} \mathcal{E}_{\pi_{e}} \left[\lambda_{t-1} \text{var} \left\{ \eta_{t} \{ r_{t} + v_{t+1}(j_{s_{t+1}}) \} \mid j_{s_{t}} \right\} \right] \\ &\geq \sum_{t=0}^{H} \exp \mathcal{E}_{\pi^{e}} \left[\log(\lambda_{t-1} \text{var} \left\{ \eta_{t}(r_{t} + v_{t+1}(j_{s_{t+1}}) \mid j_{s_{t}} \right\}) \right] \\ &\geq \sum_{t=0}^{H} \exp(\mathcal{E}_{\pi^{e}} \left[\log(\lambda_{t-1}) \right] + \mathcal{E}_{\pi^{e}} \left[\log(\text{var} \left\{ (\eta_{t} \{ r_{t} + v_{t+1}(j_{s_{t+1}}) \} \mid j_{s_{t}} \right\}) \right]) \\ &\geq \sum_{t=0}^{H} \exp((t-1) \log C_{\min} + \log V_{\min}^{2}) \geq V_{\min}^{2} C_{\min}^{H-1}. \end{split}$$

 $Proof\ of\ Theorem\ 10.$ Almost the same as the proof of Theorem 3

Proof of Theorem 11.

$$\sum_{t=0}^{H} \mathbb{E}[\lambda_t^2 \text{var}[r_t + q_{t+1}(j_{s_{t+1}}, \tau_{t+1}(j_{a_{t+1}})) \mid j_{a_t}]]$$

$$\begin{split} &= \sum_{t=0}^{H} \mathbb{E}_{\pi^{e}}[\lambda_{t} \text{var}[r_{t} + q_{t+1}(j_{s_{t+1}}, \tau_{t+1}(j_{a_{t+1}})) \mid j_{a_{t}}]] \\ &\geq \sum_{t=0}^{H} \exp \mathbb{E}_{\pi^{e}}[\log(\lambda_{t} \text{var}[r_{t} + q_{t+1}(j_{s_{t+1}}, \tau_{t+1}(j_{a_{t+1}})) \mid j_{a_{t}}])] \\ &\geq \sum_{t=0}^{H} \exp(\mathbb{E}_{\pi^{e}}[\log(\lambda_{t})] + \mathbb{E}_{\pi^{e}}[\log(\text{var}[r_{t} + q_{t+1}(j_{s_{t+1}}, \tau_{t+1}(j_{a_{t+1}})) \mid j_{a_{t}}])]) \\ &\geq \sum_{t=0}^{H} \exp((t) \log C_{\min} + \log V_{\min}^{2}) \geq V_{\min}^{2} C_{\min}^{H}. \end{split}$$

A.4 Proofs for Section 6

Proof of Theorem 12. In this section, we define $c_T(\gamma) = \sum_{t=1}^T \gamma^{t-1}$. Here, the entire regular parametric submodel is

$$\{p_{\theta}(a \mid s)p_{\theta}(r|a,s)p_{\theta}(s'|s,a)p_{\theta}(r|s,a)\},\$$

where it matches with the true pdf at $\theta = 0$. The score function of the model \mathcal{M}_{MDP} is decomposed as

$$g(o) = \log p_{\theta}(s) + \log p_{\theta}(a|s) + \log p_{\theta}(s'|s,a) + \log p_{\theta}(r|s,a) = g_S + g_{A|S} + g_{R|S,A} + g_{S'|S,A}.$$

Here, the corresponding tangent space is

$$\begin{split} & \mathcal{T}_{1} \times \mathcal{T}_{2} \times \mathcal{T}_{3} \times \mathcal{T}_{4}, \\ & \mathcal{T}_{1} = \{f(s); \mathbb{E}[f] = 0, \text{var}[f] < \infty\} \;, \\ & \mathcal{T}_{2} = \{f(s,a); \mathbb{E}[f(s,a)|s] = 0, \text{var}[f(s,a)|s] < \infty\} \;, \\ & \mathcal{T}_{3} = \{f(s,a,s'); \mathbb{E}[f(s,a,s')|s,a] = 0, \text{var}[f(s,a,s')|s,a] < \infty\} \;, \\ & \mathcal{T}_{4} = \{f(s,a,r); \mathbb{E}[f(s,a,r)|s,a] = 0, \text{var}[f(s,a,r)|s,a] < \infty\} \;. \end{split}$$

We calculate the gradient of the target functional $J(\pi^{e})$ w.r.t. the model \mathcal{M}_{MDP} . Since

$$J(\pi^{e}) = \int r p_{e,\gamma}^{(\infty)}(s) \pi^{e}(a \mid s) p(r \mid s, a) d(s, a, r)$$

$$= \lim_{T \to \infty} \int c_{T}(\gamma) \sum_{t=1}^{H} \int \gamma^{t-1} r_{t} p_{\theta}(r_{t} | s_{t}, a_{t}) \left\{ \prod_{k=1}^{t} \pi^{e}(a_{k} | s_{k}; \theta) p_{\theta}(s_{k+1} | s_{k}, a_{k}) \right\} p_{e}^{(1)}(s_{1}) d(h_{s_{t+1}})$$

$$= \lim_{T \to \infty} \int c_{T}(\gamma) \sum_{t=1}^{H} \mathbb{E}_{\pi^{e}}[\gamma^{t-1} r_{t}],$$

what we need is deriving the gradient ϕ satisfying

$$\nabla J(\pi^{\mathbf{e}}) = \mathbb{E}_{p_b}[\phi(s, a, r, s')g(s, a, r, s')].$$

This is done as follows: $\nabla J(\pi^{\rm e}) =$

$$\lim_{T \to \infty} \int c_{T}(\gamma) \sum_{t=1}^{H} \int \gamma^{t-1} r_{t} \nabla p_{\theta}(r_{t}|s_{t}, a_{t}) \left\{ \prod_{k=1}^{t} \pi^{e}(a_{k}|s_{k}; \theta) p_{\theta}(s_{k+1}|s_{k}, a_{k}) \right\} p_{e}^{(1)}(s_{1}) d(j_{s_{t+1}})$$

$$+ \lim_{T \to \infty} \int c_{T}(\gamma) \sum_{t=1}^{H} \int \gamma^{t-1} r_{t} p_{\theta}(r_{t}|s_{t}, a_{t}) \left\{ \prod_{k=1}^{t} \pi^{e}(a_{k}|s_{k}; \theta) \nabla p_{\theta}(s_{k+1}|s_{k}, a_{k}) \right\} p_{e}^{(1)}(s_{1}) d(j_{s_{t+1}})$$

$$+ \lim_{T \to \infty} \int c_{T}(\gamma) \sum_{t=1}^{H} \int \gamma^{t-1} r_{t} p_{\theta}(r_{t}|s_{t}, a_{t}) \left\{ \prod_{k=1}^{t} \nabla \pi^{e}(a_{k}|s_{k}; \theta) p_{\theta}(s_{k+1}|s_{k}, a_{k}) \right\} p_{e}^{(1)}(s_{1}) d(j_{s_{t+1}}).$$

First term This is equal to

$$\lim_{T \to \infty} \int c_T(\gamma) \sum_{t=1}^H \mathbb{E}_{\pi^e} [\gamma^{t-1} r_t g_{R|S,A}(r_t|s_t, a_t)] = \mathbb{E}_{p_{e,\gamma}^{(\infty)}} [r g_{R|S,A}(r|s, a)]$$

$$= \mathbb{E}_{p_{e,\gamma}^{(\infty)}} [\{r - \mathbb{E}[r|s, a]\} g_{R|S,A}(r|s, a)] = \mathbb{E}_{p_{e,\gamma}^{(\infty)}} [\{r - \mathbb{E}_{p_{e,\gamma}^{(\infty)}}[r|s, a]\} g(s, a, r, s')].$$

Second term This is equal to

$$\begin{split} &\lim_{T \to \infty} \gamma \int c_T(\gamma) \sum_{c=1}^H \gamma^c \sum_{t=c+1}^H \mathbb{E}_{\pi^e} [\gamma^{t-c-2} r_t g_{S'|S,A}(s_{t+1}|s_t, a_t)] = \gamma \mathbb{E}_{p_{e,\gamma}^{(\infty)}} [v(s') g_{S'|S,A}(s'|s, a)] \\ &= \gamma \mathbb{E}_{p_{e,\gamma}^{(\infty)}} [v(s') g_{S'|S,A}(s'|s, a)] = \gamma \mathbb{E}_{p_{e,\gamma}^{(\infty)}} [\{v(s') - \mathbb{E}_{p_{e,\gamma}^{(\infty)}} [v(s')|s, a]\} g_{S'|S,A}(s'|s, a)] \\ &= \gamma \mathbb{E}_{p_{e,\gamma}^{(\infty)}} [\{v(s') - \mathbb{E}_{p_{e,\gamma}^{(\infty)}} [v(s')|s, a]\} g(s, a, r, s')]. \end{split}$$

Third term This is equal to

$$\lim_{T \to \infty} \int c_T(\gamma) \sum_{c=1}^H \gamma^c \sum_{t=c+1}^H \mathbb{E}_{\pi^c} [\gamma^{t-c-1} r_t \{ g_{A|S}(a_t|s_t) - \mathbb{E}_{\pi^c} [g_{A|S}(a_t|s_t)|s_t] \}]$$

$$= \mathbb{E}_{p_{e,\gamma}^{(\infty)}} [\{ q(s,a) - v(s) \} g_{S'|S,A}(s'|s,a)] = \mathbb{E}_{p_{e,\gamma}^{(\infty)}} [\{ q(s,a) - v(s) \} g(s,a,r,s')].$$

In summary,

$$\begin{split} \nabla J(\pi^{\mathrm{e}}) &= \mathbb{E}_{p_{e,\gamma}^{(\infty)}} \left[\left\{ r - \mathbb{E}_{p_{e,\gamma}^{(\infty)}}[r|s,a] + \gamma v(s') - \gamma \mathbb{E}[v(s')|s,a] + q(s,a) - v(s) \right\} g(s,a,r,s') \right] \\ &= \mathbb{E}_{p_{e,\gamma}^{(\infty)}} [\{ r + \gamma v(s') - v(s) \} g(s,a,r,s')] \\ &= \mathbb{E}_{p_{b}} [\mu^{*}(s,a) \{ r + \gamma v(s') - v(s) \} g(s,a,r,s')]. \end{split}$$

Thus, the following function:

$$\mu^*(s, a)\{r + \gamma v(s') - v(s)\}.$$

is a gradient. Besides, since it belongs to a tangent space, this is the EIF. The efficiency bound is

$$\mathbb{E}[w^{*2}(S)\operatorname{var}[\eta(S,A)\{R+\gamma v(S')\}|S]].$$

Proof of Theorem 13. For simplicity, we consider the case K=2. Here, we have

$$\mathbb{P}_{u_{1}}[\phi(\hat{w}^{*(1)}, \hat{\pi}^{b(1)}, \hat{q}^{(1)})|\mathcal{L}_{1}] - \mathbb{P}_{u_{1}}[\phi(w^{*}, \pi^{b}, q)|\mathcal{L}_{1}]
= \mathbb{G}_{u_{1}}[\phi(\hat{w}^{*(1)}, \hat{\pi}^{b(1)}, \hat{q}^{(1)}) - \phi(w^{*}, \pi^{b}, q)]
+ \mathbb{E}[\phi(\hat{w}^{*(1)}, \hat{\pi}^{b(1)}, \hat{q}^{(1)})|\mathcal{L}_{1}] - \mathbb{E}[\phi(w^{*}, \pi^{b}, q)|\mathcal{L}_{1}]
= o_{p}(n^{-1/2}).$$
(23)

Then, the proof is immediately concluded from CLT. For the rest of the proof, we prove Eq. (24) is $o_p(n^{-1/2})$. The part Eq. (23) is $o_p(n^{-1/2})$ is similarly proved from the same decomposition. We have

$$\mathbb{E}[\hat{w}^*(S)\hat{\eta}(S,A)\{R - \hat{q}(S,A) + \gamma \hat{v}(S')\}] + \mathbb{E}_{n^{(1)}}[\hat{v}(S)] - J$$

$$\begin{split} &= \mathbb{E}[\{\hat{w}^*(S)\hat{\eta}(S,A) - w^*(S)\eta(S,A)\}\{R - q(S,A) + \gamma v(S')\}] \\ &+ \mathbb{E}[w^*(s)\eta(S,A)\{q(S,A) - \hat{q}(S,A)\} + w^*(s)\{\gamma \hat{v}(S') - \gamma v(S')\}] + (1-\gamma)\mathbb{E}_{p_e^{(1)}}[\hat{v}(s_1) - v(s_1)] \\ &+ \mathbb{E}[\{\hat{w}^*(S)\hat{\eta}(S,A) - w^*(S)\eta(S,A)\}\{q(S,A) - \hat{q}(S,A) + \gamma \hat{v}(S') - \gamma v(S')\}] \\ &\lessapprox \mathbb{E}[w^*(S)\eta(S,A)\{-\hat{q}(S,A) + \gamma \hat{v}(S')\}] + (1-\gamma)\mathbb{E}_{p_e^{(1)}}[\hat{v}(s_1)] \\ &+ \|\hat{q}(S,A) - \hat{q}(S,A)\|_2 \|\hat{\mu}(S,A) - \hat{\mu}(S,A)\|_2 \\ &\lessapprox \mathbb{E}[w^*(S)\eta(S,A)\{-\hat{q}(S,A) + \gamma \hat{v}(S')\}] + (1-\gamma)\mathbb{E}_{p_e^{(1)}}[\hat{v}(s_1)] + \alpha_1\alpha_2 + \alpha_1\beta + \alpha_2\beta + \alpha_2^2. \end{split}$$

Then, noting what we want to control is $\mathbb{E}[\hat{w}^*(S)\hat{\eta}(S,A)\{R-\hat{v}(S)+\gamma\hat{v}(S')\}]+\mathbb{E}_{p^{(1)}}[\hat{v}(S)]$, we have

$$\begin{split} & \mathbb{E}[\hat{w}^*(S)\hat{\eta}(S,A)\{R-\hat{v}(S)+\gamma\hat{v}(S')\}] + \mathbb{E}_{p_e^{(1)}}[\hat{v}(S)] - J \\ & = \mathbb{E}[w^*(S)\eta(S,A)\{-\hat{q}(S,A)+\gamma\hat{v}(S')\}] + (1-\gamma)\mathbb{E}_{p_e^{(1)}}[\hat{v}(S)] \\ & + \mathbb{E}[\hat{w}^*(S)\hat{\eta}(S,A)\{\hat{q}(S,A)-\hat{v}(S)\}] + \alpha_1\alpha_2 + \alpha_1\beta + \alpha_2\beta + \alpha_2^2 \\ & = \mathbb{E}[w^*(S)\eta(S,A)\{-\hat{q}(S,A)+\gamma v(S')\}] + \mathbb{E}[w^*(S)\hat{\eta}(S,A)\{\hat{q}(S,A)-\hat{v}(S)\}] \\ & + \mathbb{E}[-\gamma w^*(S)\eta(S,A)\hat{v}(S') + w^*(S)\hat{v}(S)] + \alpha_1\alpha_2 + \alpha_1\beta + \alpha_2\beta + \alpha_2^2 \\ & = \mathbb{E}[w^*(S)\hat{\eta}(S,A)\{\hat{q}(S,A)-\hat{v}(S)\}] + \mathbb{E}[w^*(S)\eta(S,A)\{-\hat{q}(S,A)+\hat{v}(S)\}] + \alpha_1\alpha_2 + \alpha_1\beta + \alpha_2\beta + \alpha_2^2. \end{split}$$

The last term is equal to α_2^2 as follows:

$$\begin{split} &\mathbb{E}[w^*(S)\hat{\eta}(S,A)\{\hat{q}(S,A) - \hat{v}(S)\}] + \mathbb{E}[w^*(S)\eta(S,A)\{-\hat{q}(S,A) + \hat{v}(S)\}] \\ &= \mathbb{E}\left[w^*(S)\hat{c}(S)u(A)\hat{\pi}^b(A|S)\{-\hat{v}(S) + \hat{q}(S,A)\}\right] + \mathbb{E}[w^*(S)\int\{\hat{\pi}^e(a|S) - \pi^e(a|S)\}\hat{q}(a,S)\mathrm{d}a] \\ &= \mathbb{E}\left[w^*(S)\int\hat{c}(S)u(g)\pi^b(g|s)\{-\hat{c}(S)\int u(a)\hat{q}(a,S)\hat{\pi}^b(a|S)\mathrm{d}a + \hat{q}(g,S)\}\mathrm{d}g\right] \\ &+ \mathbb{E}[w^*(S)\int u(a)\{\hat{c}(S)\hat{\pi}^b(a|S) - c(S)\pi^b(a|S)\}\hat{q}(a,S)\mathrm{d}a] \\ &= \mathbb{E}[w^*(S)\int\{\hat{c}(S) - c(S)\}u(g)\pi^b(g|s)\hat{q}(g,S)\mathrm{d}g] + \\ &+ \mathbb{E}[w^*(S)\{-\hat{c}^2(S)/c(S) + \hat{c}(S)\}\int u(a)\hat{\pi}^b(a|S)\hat{q}(a,S)\mathrm{d}a] \\ &= \mathbb{E}[w^*(S)\int\{\hat{c}(S) - c(S)\}u(g)\{\pi^b(g|s) - \hat{\pi}^b(g|S)\}\hat{q}(g,S)\mathrm{d}g] + \\ &+ \mathbb{E}[w^*(S)\{-\hat{c}^2(S)/c(S) + \hat{c}(S) + \hat{c}(S) - c(S)\}\int u(a)\hat{\pi}^b(a|S)\hat{q}(a,S)\mathrm{d}a] \\ &= \mathbb{E}[w^*(S)\int\{\hat{c}(S) - c(S)\}u(g)\{\pi^b(g|S) - \hat{\pi}^b(g|S)\}\hat{q}(g,S)\mathrm{d}g] + \\ &+ \mathbb{E}[w^*(S)\{-\hat{c}(S) + c(S)\}^2/c(S)\int u(a)\hat{\pi}^b(a|S)\hat{q}(a,S)\mathrm{d}a] \\ &\lesssim \|\hat{\pi}^b(A|S) - \pi^b(A|S)\|_2^2 = \alpha_2^2. \end{split}$$

In summary,

$$\mathbb{E}[\phi(\hat{w}^{*(1)}, \hat{\pi}^{b(1)}, \hat{q}^{(1)}) | \mathcal{L}_1] - \mathbb{E}[\phi(w^*, \pi^b, q) | \mathcal{L}_1] = \alpha_1 \alpha_2 + \alpha_1 \beta + \alpha_2 \beta + \alpha_2^2.$$

Under the assumption regarding convergence rates, this is equal to $o_p(n^{-1/2})$.

Proof of Theorem 14. This is immediately concluded from

$$\mathbb{E}[\phi(\hat{w}^{*(1)}, \hat{\pi}^{b(1)}, \hat{q}^{(1)})|\mathcal{L}_1] - \mathbb{E}[\phi(w^{*\dagger}, \pi^{b\dagger}, q^{\dagger})|\mathcal{L}_1] = \alpha_1 \alpha_2 + \alpha_1 \beta + \alpha_2 \beta + \alpha_2^2,$$

which is obtained in the proof of Theorem 13.

Proof of Theorem 15. This is similarly proved as in Theorem 12 by some calculation. \Box

Proof of Theorem 16. Here, we have

$$\mathbb{P}_{u_{1}}[\phi(\hat{w}^{*(1)}, \hat{\pi}^{b(1)}, \hat{q}^{(1)})|\mathcal{L}_{1}] - \mathbb{P}_{u_{1}}[\phi(w^{*}, \pi^{b}, q)|\mathcal{L}_{1}]
= \mathbb{G}_{u_{1}}[\phi(\hat{w}^{*(1)}, \hat{\pi}^{b(1)}, \hat{q}^{(1)}) - \phi(w^{*}, \pi^{b}, q)]
+ \mathbb{E}[\phi(\hat{w}^{*(1)}, \hat{\pi}^{b(1)}, \hat{q}^{(1)})|\mathcal{L}_{1}] - \mathbb{E}[\phi(w^{*}, \pi^{b}, q)|\mathcal{L}_{1}]
= o_{p}(n^{-1/2}).$$
(25)

Then, the proof is immediately concluded from CLT. It remains to bound Eq. (26) is $o_p(n^{-1/2})$. The part that Eq. (25) is $o_p(n^{-1/2})$ is similarly proved from the same decomposition. We have

$$\mathbb{E}[\hat{w}^*(S)\hat{\eta}(S,A)\{R - \hat{q}(S,A) + \gamma \hat{q}(S',\tau(A',S'))\}] + (1-\gamma)\mathbb{E}_{p_e^{(1)}}[\hat{v}(s_1)] - J$$

$$= \mathbb{E}[\{\hat{w}^*(S)\hat{\eta}(S,A) - w^*(S)\eta(S,A)\}\{R - q(S,A) + \gamma q(S',\tau(A',S'))\}]$$

$$+ \mathbb{E}[w^*(S)\eta(S,A)\{q(S,A) - \hat{q}(S,A)\} + w^*(S)\{\gamma \hat{q}(S,\tau(S,A)) - \gamma q(S,\tau(S,A))\}]$$

$$+ (1-\gamma)\mathbb{E}_{p_e^{(1)}}[\hat{v}(s_1) - v(s_1)]$$

$$+ \mathbb{E}[\{\hat{w}^*(S)\hat{\eta}(S,A) - w^*(S)\eta(S,A)\}\{q(S,A) - \hat{q}(S,A) + \gamma \hat{q}(S',\tau(A',S')) - \gamma q(S',\tau(A',S'))\}]$$

$$\lessapprox \|\hat{w}^*(S)\hat{\eta}(S,A) - w^*(S)\eta(S,A)\|_2 \|\hat{q}(S,A) - q(S,A)\|_2 = (\alpha_1 + \alpha_2)\beta.$$
(27)

Here, we use Eq. (27) and Eq. (28) are 0.

Proof of Theorem 17. This is immediately concluded from the proof of Theorem 16 as in Corollary 16 (Kallus and Uehara, 2020a). \Box

Proof of Theorem 18. Immediately, concluded from Theorem 19. \Box

Proof of Theorem 19. Here, we have

$$\mathbb{P}_{u_{1}}[\phi(\hat{w}^{*(1)}, \hat{v}^{(1)})|\mathcal{L}_{1}] - \mathbb{P}_{u_{1}}[\phi(w^{*}, v)|\mathcal{L}_{1}]
= \mathbb{G}_{u_{1}}[\phi(\hat{w}^{*(1)}, \hat{v}^{(1)}) - \phi(w^{*}, v)]
+ \mathbb{E}[\phi(\hat{w}^{*(1)}, \hat{v}^{(1)})|\mathcal{L}_{1}] - \mathbb{E}[\phi(w^{*}, v)|\mathcal{L}_{1}]
= o_{p}(n^{-1/2}).$$
(29)

Then, the proof is immediately concluded from CLT. For the rest of the proof, we prove Eq. (30) is $o_p(n^{-1/2})$. The part Eq. (29) is $o_p(n^{-1/2})$ is similarly proved from the same decomposition. We have

$$\begin{split} &\mathbb{E}[\hat{w}^*(S)\eta(S,A)\{R-\hat{v}(S)+\gamma\hat{v}(S')\}] + (1-\gamma)\mathbb{E}_{p_e^{(1)}}[\hat{v}(s_1)] - J \\ &= \mathbb{E}[\{\hat{w}^*(S)\eta(S,A)-w^*(S)\eta(S,A)\}\{R-v(S)+\gamma v(S')\}] \\ &+ \mathbb{E}[w^*(S)\eta(S,A)\{v(S)-\hat{v}(S)\}+w^*(S)\{\gamma\hat{v}(S)-\gamma v(S')\}] + (1-\gamma)\mathbb{E}_{p_e^{(1)}}[\hat{v}(s_1)-v(s_1)] \\ &+ \mathbb{E}[\{\hat{w}^*(S)\eta(S,A)-w^*(S)\eta(S,A)\}\{v(S)-\hat{v}(S)+\gamma\hat{v}(S')-\gamma v(S')\}] \\ &\lesssim \|\hat{w}^*(S)\eta(S,A)-w^*(S)\eta(S,A)\|_2\|\hat{v}(S)-v(S)\|_2 = \alpha_1\beta = o_p(n^{-1/2}). \end{split}$$

Here, we use the following fact:

$$\mathbb{E}[\eta(S, A)\{R - v(S) + \gamma v(S')\}] = \mathbb{E}[\mathbb{E}[\eta(S, A)\{R + \gamma v(S')\}|S] - v(S)] = 0,$$

and we derived that Eq. (32) and Eq. (31) are 0.

Remark 10. In the finite horizon case, we also have a double robustness and efficiency. The argument is as follows. Here, we have

$$\mathbb{P}_{u_{1}}[\phi(\hat{w}^{(1)}, \hat{v}^{(1)})|\mathcal{L}_{1}] - \mathbb{P}_{u_{1}}[\phi(w, v)|\mathcal{L}_{1}]
= \mathbb{G}_{u_{1}}[\phi(\hat{w}^{(1)}, \hat{v}^{(1)}) - \phi(w, v)]
+ \mathbb{E}[\phi(\hat{w}^{(1)}, \hat{v}^{(1)})|\mathcal{L}_{1}] - \mathbb{E}[\phi(w, v)|\mathcal{L}_{1}]
= o_{p}(n^{-1/2}).$$
(33)

We prove Eq. (34) is $o_p(n^{-1/2})$. The fact Eq. (33) is $o_p(n^{-1/2})$ is similarly proved as follows. Then, the final statement is concluded from CLT. The proof is as follows:

$$\mathbb{E}\left[\sum_{t=1}^{H} \hat{w}_{t}(S_{t})\eta(S_{t}, A_{t})\{R_{t} - \hat{v}_{t}(S_{t})\} + \hat{w}_{t-1}(S_{t-1})\eta(S_{t-1}, A_{t-1})\hat{v}_{t}(S_{t})\right]$$

$$= \mathbb{E}\left[\sum_{t=1}^{H} \{\hat{w}_{t}(S_{t}) - w_{t}(S_{t})\}\eta_{t}(S_{t}, A_{t})\{-\hat{v}_{t}(S_{t}) + v_{t}(S_{t})\}\right]$$

$$+ \mathbb{E}\left[\sum_{t=1}^{H} \{\hat{w}_{t-1}(S_{t-1}) - w_{t-1}(S_{t-1})\}\eta_{t-1}(S_{t-1}, A_{t-1})\{-\hat{v}_{t}(S_{t}) + v_{t}(S_{t})\}\right]$$

$$+ \mathbb{E}\left[\sum_{t=1}^{H} w_{t}(S_{t})\{-\hat{v}_{t}(S_{t}) + v_{t}(S_{t})\} + w_{t-1}(S_{t-1})\{\hat{v}_{t}(S_{t}) - v_{t}(S_{t})\}\right]$$

$$+ \mathbb{E}\left[\sum_{t=1}^{H} \{\hat{w}_{t}(S_{t}) - w_{t}(S_{t})\}\eta_{t}(S_{t}, A_{t})\{R_{t} - v_{t}(S_{t}) + v_{t+1}(S_{t+1})\}\right]$$

$$\lessapprox \sum_{t=1}^{H} \|\hat{v}_{t}(S_{t}) - v_{t}(S_{t})\|_{2} \|\hat{w}_{t}(S_{t}) - w_{t}(S_{t})\|_{2} = \alpha_{1}\beta.$$

B Properties of the estimator of Tang et al. (2020)

In this section we consider a sample-splitting version of the estimator of Tang et al. (2020) in the case of a pre-specified evaluation policy and establish its (inefficient) asymptotic behavior and double robustness. Given nuisance estimators $\hat{v}^{(k)}(s)$, $\hat{w}^{*(k)}(s)$, the estimator we consider is given by replacing Eq. (6) in Algorithm 1 with

$$\hat{J}_k = \frac{1}{|I_k|} \sum_{i \in I_k} [\hat{w}^{*(k)}(S^{(i)}) \eta(S^{(i)}, A^{(i)}) \{ R^{(i)} + \gamma \hat{v}^{(k)}(S^{'(i)}) - \hat{v}^{(k)}(S^{(i)}) \}]$$

$$+ (1 - \gamma) \mathbb{E}_{p_e^{(1)}(s_1)} [\hat{v}^{(k)}(s_1)].$$

Here, we assume the behavior policy is known. We denote this estimator as \hat{J}_{V2} . We can similarly consider an analogous estimator, \hat{J}_{V1} , for the finite horizon case, where, given nuisance estimators $\hat{v}_t^{(k)}(s)$, $\hat{w}_t^{(k)}(s)$, we use

$$\hat{J}_k = \frac{1}{|I_k|} \sum_{i \in I_k} \left[\hat{v}_1^{(k)}(S_1^{(i)}) + \sum_{t=1}^H \hat{w}_t^{(k)}(S_t^{(i)}) \eta_t(S_t^{(i)}, A_t^{(i)}) \{ R_t^{(i)} + \hat{v}_{t+1}^{(k)}(S_{t+1}^{(i)}) - \hat{v}_t^{(k)}(S_t^{(i)}) \} \right].$$

Importantly, these estimators allow direct estimation of the value functions. In this sense, they are different from \hat{J}_{TI1} , \hat{J}_{TI2} , where the construction of $\hat{v}(s)$ is restricted to $\hat{c}(s) \int u(a)\hat{\pi}^b(a|s)\hat{q}(s,a)\mathrm{d}a$, $\hat{c}(s) = 1/\int u(a)\hat{\pi}^b(a|s)\mathrm{d}a$.

The estimator \hat{J}_{V2} is doubly robust if the behavior policy is known in the sense that it is consistent as long as the either model of $w^*(s)$ and v(s) is correct.

Theorem 18 (Double Robustness). Suppose
$$\forall k \leq K$$
, for some $w^{*\dagger}, v^{\dagger}$, $\|\hat{w}^{*(k)}(s) - w^{*\dagger}(s)\|_2 = o_p(1)$, $\|\hat{v}^{(k)}(s) - v^{\dagger}(s)\|_2 = o_p(1)$. Then, as long as $w^{*\dagger} = w^*$ or $v^{\dagger} = v$, $\hat{J}_{V2} \stackrel{p}{\to} J$.

In addition, the asymptotic MSE can be calculated under nonparametric rate conditions if the behavior policy is known.

Theorem 19 (Asymptotic Results). Suppose
$$\forall k \leq K$$
, $\|\hat{w}^{*(k)}(s) - w^*(s)\|_2 = \alpha_1$, $\|\hat{v}^{(k)}(s) - v(s)\|_2 = \beta$, where $\alpha_1 \beta = o_p(n^{-1/2})$, $\alpha_1 = o_p(1)$, $\beta = o_p(1)$. Then, $\sqrt{n}(\hat{J}_{V2} - J) \stackrel{d}{\to} \mathcal{N}(0, \Upsilon_{TI2})$.

Again note that this asymptotic MSE is larger than the efficiency bound in the pre-specified evaluation policy case (Kallus and Uehara, 2019a).