Transformers are Universal In-context Learners

Takashi Furuya Shimane Univ. takashi.furuya01010gmail.com Maarten V. de Hoop Rice Univ. mvd2@rice.edu Gabriel Peyré CNRS, ENS, PSL Univ. gabriel.peyre@ens.fr

Abstract

Transformers are deep architectures that define "in-context mappings" which enable predicting new tokens based on a given set of tokens (such as a prompt in NLP applications or a set of patches for vision transformers). This work studies in particular the ability of these architectures to handle an arbitrarily large number of context tokens. To mathematically and uniformly address the expressivity of these architectures, we consider the case that the mappings are conditioned on a context represented by a probability distribution of tokens (discrete for a finite number of tokens). The related notion of smoothness corresponds to continuity in terms of the Wasserstein distance between these contexts. We demonstrate that deep transformers are universal and can approximate continuous in-context mappings to arbitrary precision, uniformly over compact token domains. A key aspect of our results, compared to existing findings, is that for a fixed precision, a single transformer can operate on an arbitrary (even infinite) number of tokens. Additionally, it operates with a fixed embedding dimension of tokens (this dimension). The use of MLP layers between multi-head attention layers is also explicitly controlled.

1 Introduction

Transformers have revolutionized the field of machine learning with their powerful attention mechanisms, as introduced by Vaswani et al. [35]. The exceptional performance and expressivity of large-scale transformers have been empirically well established for both NLP [4] and vision applications [10]. One key property of these architectures is their ability to leverage contexts of arbitrary length, which enables the parameterization of "in context" mappings with an arbitrarily large complexity. In this paper, we present a rigorous formalism to model inputs and the associated context with an arbitrarily large number of tokens, defining a notion of continuity that enables the analysis of their expressivity.

1.1 Previous work

Universality, from neural networks to neural operators. Multilayer Perceptrons (MLP) with two layers are universal approximators, as shown decades ago in [8, 18], with a comprehensive review in [29]. The significance of depth in enhancing expressivity is explored in [17, 39]. These results have been extended to cover a variety of architectural constraints on the networks, for instance, using weight sharing in Convolutional Neural Networks (CNN) [42] and skip connexions in ResNets [7, 34]. It is also possible to design equivariant architectures, in particular for graph neural networks [23, 19, 38] and neural networks operating on sets of points [30, 9]. The connection between transformers and graph neural networks is exposed in [27]. We focus on this article from a related but different point of view, viewing transformers as operating on probability distribution rather than sets of points. Related to this setup are extensions of neural networks to infinite dimensional input functional spaces using the concept of neural operator [20], which universality is studied in [14]. They can be generalized to cope with data in metric spaces, addressing topological obstructions, in [22].

Mathematical modeling of transformers. It is now customary to describe transformers as performing "in context" prediction, which means that it maps token to token, but that this map depends on a set of previously seen tokens. The size of this context might be very long, possibly arbitrarily long, which is the focus of this article. The ability of trained transformers to effectively perform in-context computation has been supported by both empirical studies [36] and theoretical results [2, 25, 32, 41] on simplified architectures (typically linear attention) and specific data generation processes.

To make a rigorous analysis of arbitrarily long token lengths, and also describe a "mean field" limit of an infinite number of tokens, it is convenient to view attention as operating over probability distributions of tokens [37, 31]. The smoothness (Lipschitz continuity) of these attention layers is analyzed in [5]. Deep transformers (with the residual connection) can be described by a coupled system of particles evolving across the layers. The analysis of the clustering properties of such an evolution is studied in [15, 16].

Universality of transformers. [40] provides, to the best of our knowledge, the most detailed account of the universality of transformers. The authors rely on shallow transformers with only 2 heads but require that the transformers operate over an embedding dimension which grows with the number of tokens. This result is refined in [28] which highlights the difficulty of attention mechanisms to capture smooth functions. Our focus is different, since we consider deep transformers with a fixed embedding dimension, but which are universal for an arbitrary number of tokens.

We note that there exist variations over the initial transformers architecture which enjoys universality results, for instance, the Sumformer [3] and stochastic deep netorks [9], which also requires an embedding dimension that grows with the number of tokens. We also mention probabilistic transformers [21] which can approximate embeddings of metric spaces. The work of [1] provides an abstract universal interpolation result for equivariant architectures under genericity conditions, but it is not known whether there exist generic attention maps.

While this is not directly related to our results, a line of works studies the expressivity of transformers as operating on a discrete set of tokens as formal systems [6, 26, 33, 12]. Another line of work studies the impact of positional encoding on the expressivity [24]

1.2 Our contributions

Our work provides a rigorous formalization of transformer expressivity and continuity as operating over the space of probability distributions. The main mathematical result is the universality presented in Theorem 1. Our approach effectively handles an arbitrary number of tokens and leverages deep architectures without requiring arbitrary width. The embedding dimension and the number of heads are proportional to the dimension of the input tokens and are independent of precision.

A limitation of our approach is that it does not consider masked attention, which would require going beyond the current framework based on Wasserstein distances. Despite this, our formalism provides a significant step forward in the theoretical modeling of transformers, offering new insights into their underlying mechanics and potential applications.

1.3 Notation

For natural number $N \in \mathbb{N}$, we denote by $[N] := \{1, ..., N\}$. For vector $x \in \mathbb{R}^d$, the Euclidean norm of x is denoted by |x|. For two vectors $x, y \in \mathbb{R}^d$, the Euclidean inner product of x and y is denoted by $\langle x, y \rangle$ and the component-wise multiplication of x and y is denoted by $x \odot y$. The vector $\mathbf{1}_d$ is the vector of dimension d with all coordinates equal to 1, that is, $\mathbf{1}_d := (1, ..., 1) \in \mathbb{R}^d$.

In the following, we denote by $\mu \in \mathcal{P}(\Omega)$ a probability measure on some compact domain $\Omega \subset \mathbb{R}^d$ of tokens' embeddings. We denote by $\mathcal{C}(\Omega)$ the space of continuous functions from Ω to \mathbb{R} . Our construction makes use of the push-forward operator T_{\sharp} . For $T : \Omega \subset \mathbb{R}^d \to \Omega' \subset \mathbb{R}^{d'}$ a measurable map, a measure $\mu \in \mathcal{P}(\Omega)$ is mapped to $T_{\sharp}\mu \in \mathcal{P}(\Omega')$. It operates over discrete measures by simply displacing the support

$$T_{\sharp}\left(\frac{1}{n}\sum_{i=1}^{n}\delta_{x_{i}}\right) \coloneqq \frac{1}{n}\sum_{i=1}^{n}\delta_{T(x_{i})}$$

For a general measure, $\nu = T_{\sharp}\mu$ is defined by a change of variables in integration:

$$\forall g \in \mathcal{C}(\Omega'), \quad \int_{\Omega'} g(y) \, d\nu(y) \coloneqq \int_{\Omega} g(T(x)) \, d\mu(x). \tag{1}$$

We emply the weak^{*} topology on $\mathcal{P}(\Omega)$, which is associated with the following notion of convergence of sequences:

$$\mu_k \rightharpoonup^* \mu \quad \Leftrightarrow \quad \left(\forall f \in \mathcal{C}(\Omega), \ \int f(x) \, d\mu_k(x) \to \int f(x) \, d\mu(x) \right).$$

Intuitively, this corresponds to a "soft" notion of convergence where the support of μ_k approaches that of μ .

In the special case of discrete measures with a fixed number n of points, this corresponds, up to relabeling of the points, to the usual convergence of points in finite dimensions:

$$\left(\frac{1}{n}\sum_{i=1}^{n}\delta_{x_{i}^{k}} \stackrel{\rightharpoonup^{*}}{\longrightarrow} \frac{1}{n}\sum_{i=1}^{n}\delta_{x_{i}}\right) \quad \Leftrightarrow \quad \left(X_{k} = (x_{i,k})_{i} \in \mathbb{R}^{d \times n} \to X = (x_{i})_{i} \in \mathbb{R}^{d \times n}\right).$$

While we do not make use of this in this paper (since our claims are not quantitative), it is possible to metrize this weak^{*} topology using the Wasserstein Optimal Transport distance, which is defined, for $1 \le p < +\infty$, as

$$W_p(\mu,\nu)^p \coloneqq \min_{\pi \in \mathcal{P}(\Omega^2)} \left\{ \int \|x - y\|^p \, d\pi(x,y) : \pi_1 = \mu, \pi_2 = \nu \right\},$$

where $\pi_i = (P_i)_{\sharp}\pi$ are the marginals of π with $P_1(x, y) = x$ and $P_2(x, y) = y$. One has

$$\mu_k \rightharpoonup^* \mu \quad \Leftrightarrow \quad W_p(\mu, \nu) \to 0.$$

An avenue for future work is to obtain quantitative approximation results for in-context mappings that are, for instance, Lipschitz continuous according to the Wasserstein distance.

2 Measure-theoretic in-context mappings

Transformers are defined by alternating multi-head attention layers (which compute interactions between tokens), MLP and normalization layers (which operate independently over each token). For the sake of simplicity, we omit normalization in the following analysis. We first recall their definition and then explain how they can be equivalently re-written using in-context mappings. This definition provides new insights and can also be generalized to an "infinite" number of tokens encoded in a probability measure.

2.1 Attention as in context mappings on token ensembles

Classical definition. A set of *n* tokens $x_i \in \mathbb{R}^{d_{\text{in}}}$ is denoted by $X = (x_i)_{i=1}^n \in \mathbb{R}^{d_{\text{in}} \times n}$. An attention head maps these *n* tokens to the same number *n* of tokens in $\mathbb{R}^{d_{\text{head}}}$ using

$$\forall X \in \mathbb{R}^{d_{\text{in}} \times n}, \quad \text{Att}_{\theta}(X) \coloneqq VX \text{ SoftMax}(X^{\top}Q^{\top}KX/\sqrt{k}) \in \mathbb{R}^{d_{\text{head}} \times n},$$

where the parameters are the (Key, Query, Value) matrices $\theta := (Q, K, V) \in \mathbb{R}^{k \times d_{\text{in}}} \times \mathbb{R}^{d_{\text{head}} \times d_{\text{in}}}$. Here, the SoftMax function operates in a row-wise manner:

$$\forall Z \in \mathbb{R}^{n \times n}, \quad \text{SoftMax}(Z) \coloneqq \left(\frac{e^{Z_{i,j}}}{\sum_{\ell} e^{Z_{i,\ell}}}\right)_{i,j=1}^n \in \mathbb{R}^{n \times n}_+.$$

Multiple heads with different parameters $\theta := (W^h, \theta^h)_{h=1}^H$ are combined in a linear way in a multi-head attention:

$$\operatorname{MAtt}_{\theta}(X) \coloneqq \sum_{h=1}^{H} W^{h} \operatorname{Att}_{\theta^{h}}(X),$$

where $W^h \in \mathbb{R}^{d_{\text{out}} \times d_{\text{head}}}$ and $\theta^h \coloneqq (Q^h, K^h, V^h)$.

In the following, we denote the various dimensions of a multi-head attention layer by: $d_{in}(\theta)$, $d_{out}(\theta)$, $d_{head}(\theta)$ for the input, output, and head dimensions, respectively, and $k(\theta)$ for the key/query dimensions, and $H(\theta)$ for the number of heads.

In-context mappings form. The mapping $X \mapsto MAtt_{\theta}(X)$ can be re-written as the application of an "in context" function $G_{\theta}(X, \cdot)$ to each token,

$$x_i \mapsto G_{\theta}(X, x_i)$$
 i.e. $\operatorname{MAtt}_{\theta}(X) = (G_{\theta}(X, x_i))_{i=1}^n$

where the in-context mapping is

$$\forall (X,x) \in \mathbb{R}^{d_{\mathrm{in}} \times n} \times \mathbb{R}^{d_{\mathrm{in}}}, \quad G_{\theta}(X,x) \coloneqq \sum_{h=1}^{H} W^{h} \sum_{j=1}^{n} \frac{\exp\left(\frac{1}{\sqrt{k}} \langle Q^{h}x, K^{h}x_{j} \rangle\right)}{\sum_{\ell=1}^{n} \exp\left(\frac{1}{\sqrt{k}} \langle Q^{h}x, K^{h}x_{\ell} \rangle\right)} V^{h}x_{j}.$$
(2)

Here, the terminology "in context" refers to the fact that $G_{\theta}(X, \cdot)$ depends on the tokens X themselves, and can thus be seen as a parametric map that is modified for each token depending on its interactions with the other tokens. While this re-writing is equivalent to the original one, it highlights the fact that transformers define spatial mappings. This also allows us to clearly state the associated mathematical question at the core of this paper, which is the approximation of arbitrary in-context mappings by (compositions of) such parametric maps. Another interest in this reformulation is that it enables the definition of generalized attention operating over a possibly infinite number of tokens, as explained in Section 2.2.

Composition of in-context mappings. A transformer (ignoring normalization layers at this moment) is a composition of L attention layers and Multi-Layer Perceptrons (MLP):

$$\mathrm{MLP}_{\xi_L} \circ \mathrm{MAtt}_{\theta_L} \circ \ldots \circ \mathrm{MLP}_{\xi_1} \circ \mathrm{MAtt}_{\theta_1} \,. \tag{3}$$

Here, the MLP_{ξ} functions process each token independently from one another:

$$\mathrm{MLP}_{\xi}(X) = (F_{\xi}(x_i))_{i=1}^n,$$

i.e., they are "context-free" mappings (in the above notation, $F_{\xi}(X, x) = F_{\xi}(x)$), while the attention maps, $G_{\theta}(X, \cdot)$, depend on the context X.

On the level of in-context mappings, the composition of layers in (3) induces a new "in-context" composition rule, which we denote by \diamond ,

$$\forall X = (x_i)_{i=1}^n, \ \forall x : \quad (G_2 \diamond G_1)(X, x) \coloneqq G_2(X_1, G_1(X, x)) \quad \text{where} \quad X_1 = (G_1(X, x_i))_i.$$
(4)

This rule can be applied whether $G_1(X, \cdot)$ or $G_2(X, \cdot)$ depends on the context X or not (such as for the F_{ξ} mappings above, which are independent of the context). Using this rule, the transformer's definition (3) translates into a composition of in-context and context-free maps:

$$F_{\xi_L} \diamond G_{\theta_L} \diamond \dots \diamond F_{\xi_1} \diamond G_{\theta_1}. \tag{5}$$

The core question this paper addresses is the uniform approximation of a continuous (in a suitable topology) in-context map $(X, x) \mapsto G(X, x)$ by transformers' in-context mappings of the form (3), with clear control of the dimensions and the number of heads involved in the different layers. The main originality of our approach is that we aim to do so for an arbitrary number n of tokens, as we now explain.

2.2 Measure-theoretic in-context mappings

A key observation is that the definition in (2) makes sense irrespective of the number, n, of tokens. To make this more explicit, and also handle the limit of an infinite number of tokens, we represent a set X of tokens using a probability distribution $\mu \in \mathcal{P}(\mathbb{R}^{d_{\text{in}}})$ over $\mathbb{R}^{d}_{\text{in}}$. A finite number of tokens is encoded using a discrete empirical measure,

$$\mu = \frac{1}{n} \sum_{i=1}^{n} \delta_{x_i} \in \mathcal{P}(\mathbb{R}^{d_{\text{in}}}).$$
(6)

This encoding is not only for notional convenience, it also allows us to define clearly a correct notion of smoothness for the in-context mappings. This smoothness corresponds to the displacement of the tokens and is quantified through the optimal transport distance as presented in Section 1.3. This enables us to compare context with different sizes and, for instance, to compare a set of tokens with a large (but finite) n to a continuous distribution.

Using probability distributions, the in-context mapping (2) is now defined as

$$\forall (\mu, x) \in \mathcal{P}(\mathbb{R}^{d_{\mathrm{in}}}) \times \mathbb{R}^{d_{\mathrm{in}}}, \quad \Gamma_{\theta}(\mu, x) \coloneqq \sum_{h=1}^{H} W^{h} \int \frac{\exp\left(\frac{1}{\sqrt{k}} \langle Q^{h}x, K^{h}y \rangle\right)}{\int \exp\left(\frac{1}{\sqrt{k}} \langle Q^{h}x, K^{h}z \rangle\right) \mathrm{d}\mu(z)} V^{h}y \,\mathrm{d}\mu(y). \tag{7}$$

We can include a skip connection, that is, Γ_{θ} can be redefined as

$$\forall (\mu, x) \in \mathcal{P}(\mathbb{R}^{d_{\mathrm{in}}}) \times \mathbb{R}^{d_{\mathrm{in}}}, \quad \Gamma_{\theta}(\mu, x) \coloneqq x + \sum_{h=1}^{H} W^{h} \int \frac{\exp\left(\frac{1}{\sqrt{k}} \langle Q^{h}x, K^{h}y \rangle\right)}{\int \exp\left(\frac{1}{\sqrt{k}} \langle Q^{h}x, K^{h}z \rangle\right) \mathrm{d}\mu(z)} V^{h}y \,\mathrm{d}\mu(y). \tag{8}$$

In this case, we assume that $d_{in} = d_{out}$.

The discrete case is contained in this more general definition in the sense that

$$\forall X = (x_i)_{i=1}^n, \quad G_\theta(X, x) = \Gamma_\theta \left(\frac{1}{n} \sum_{i=1}^n \delta_{x_i}, x\right).$$

In the following, we will invoke, whenever convenient, the following slight abuse of notation,

$$\Gamma_{\theta}(\mu, x) = \Gamma_{\theta}(\mu)(x),$$

so that $\Gamma_{\theta}(\mu) : \mathbb{R}^{d_{\text{in}}} \to \mathbb{R}^{d_{\text{out}}}$ defines a map between Euclidean spaces. Using this general definition, the attention map $X \mapsto \text{MAtt}_{\theta}(X)$ can be rewritten as displacing the tokens' positions, which corresponds to applying a push-forward to the measure as defined in (1),

$$\mu \in \mathcal{P}(\mathbb{R}^{d_{\mathrm{in}}}) \longmapsto \Gamma_{\theta}(\mu)_{\sharp} \mu \in \mathcal{P}(\mathbb{R}^{d_{\mathrm{out}}}).$$

This formulation of transformers as a mapping between probability measures was introduced in [31] and also used in [5] to prove a convergence result of deep transformers. We re-use it here but put emphasis on the in-context mapping itself, which is the object of interest of this paper (rather than on studying the mapping between measures).

Composition of in-context measure-theoretic mappings. The definition of composition in (4) generalizes to the measure-theoretic setting as

$$(\Gamma_2 \diamond \Gamma_1)(\mu, x) \coloneqq \Gamma_2(\mu_1, \Gamma_1(\mu, x)), \quad \text{where} \quad \mu_1 \coloneqq \Gamma_1(\mu)_{\sharp} \mu, \tag{9}$$

i.e., $(\Gamma_2 \diamond \Gamma_1)(\mu) = \Gamma_2(\mu_1) \circ \Gamma_1(\mu)$. Transformers operating over an arbitrary (possibly infinite) number of tokens are then obtained by replacing the original definition (5) by

$$F_{\xi_L} \diamond \Gamma_{\theta_L} \diamond \dots \diamond F_{\xi_1} \diamond \Gamma_{\theta_1}. \tag{10}$$

Here, the F_{ξ} are "context-free" MLP mappings, i.e., $F_{\xi}(\mu, x) = F_{\xi}(x)$ is independent of μ . It is important to keep in mind that when restricted to finite discrete empirical measures of the form (6), definitions (5) and (10) coincide. Our theory encompasses classical transformers as well as their "mean field" limits operating over arbitrary measures.

2.3 Training transformers and applications

Transformers are trained in a supervised or unsupervised way to approximate some (unknown) incontext map $\Lambda^*(\mu, x)$ to perform in-context prediction by minimizing some loss function ℓ :

$$\min_{(\theta_{\ell},\xi_{\ell})_{\ell=1}^{L}} \sum_{p} \ell(F_{\xi_{L}} \diamond \cdots \diamond \Gamma_{\theta_{1}}(\mu_{p}, x_{p}), y_{p}),$$

where (μ_p, x_p) are pairs of contexts and tokens, and y_p are the labels to predict (typically another token). In most situations of practical interest, x_p belongs to the support of the measure μ_p .

In computer vision applications, in particular for generative image modeling (diffusion models) using Vision Transformers, μ_p is a set of patches (with positional encoding added) extracted from a noisy image, and $\Lambda^*(\mu_p, x_p)$ maps a noisy patch x_p to approximate its denoised version y_p . In NLP applications, the leading paradigm is next-token prediction through self-supervised learning. Tokens extracted from a sentence are of the form $\{(x(t),t)\}_{t=1}^n$, and the measure representation is $\mu_p = \frac{1}{n} \sum_{t=1}^T \delta_{(x(t),t)}$. Note that here, following previous work, we have appended the token index t in the sentence (since for NLP, transformers are not permutation equivariant). In this case, for a token $x_p = x(t)$ at some position t in the sentence, the token to predict is $y_p = x(t+1)$. For these NLP applications, the self-attention maps should be made causal to ensure that the problem is not trivial and to enable use in a generative model through auto-regressive evaluation of the map. This is made explicit in the definition,

$$\Gamma_{\theta}(\mu, (x, t)) \coloneqq \sum_{h=1}^{H} W^{h} \int \frac{\exp\left(\frac{1}{\sqrt{k}} \langle Q^{h}x, K^{h}y \rangle\right) \mathbf{1}_{t \leq t'}}{\int \exp\left(\frac{1}{\sqrt{k}} \langle Q^{h}x, K^{h}z \rangle\right) \mathbf{1}_{t \leq s} \, \mathrm{d}\mu(z, s)} V^{h}y \, \mathrm{d}\mu(y, t'), \tag{11}$$

where $1_{t \leq t'}$ is a masking function that is 1 if $t \leq t'$ and 0 otherwise. We leave the extension of our results to such a causal in-context mapping for future work.

2.4 Main result and discussion

Our main result is the following uniform approximation theorem.

Theorem 1. Let $\Omega \subset \mathbb{R}^d$ be a compact set and $\Lambda^* : \mathcal{P}(\Omega) \times \Omega \to \mathbb{R}^{d'}$ be continuous, where $\mathcal{P}(\Omega)$ is endowed with the weak^{*} topology. Then for all $\varepsilon > 0$, there exist L and parameters $(\theta_{\ell}, \xi_{\ell})_{\ell=1}^{L}$, such that

 $\forall (\mu, x) \in \mathcal{P}(\Omega) \times \Omega, \quad |F_{\xi_L} \diamond \Gamma_{\theta_L} \diamond \ldots \diamond F_{\xi_1} \diamond \Gamma_{\theta_1}(\mu, x) - \Lambda^{\star}(\mu, x)| \le \varepsilon,$

with $d_{\rm in}(\theta_\ell), d_{\rm out}(\theta_\ell) \le d + 3d', d_{\rm head}(\theta_\ell) = k(\theta_\ell) = 1, \ H(\theta_\ell) = d'.$

The two main strengths of this result are (i) the approximating architecture performs the approximation independently of n (it even works for an infinite number of tokens), and (ii) the number of heads and the embedding dimension do not depend on ε .

A weakness is that we have no explicit control over the dependency of the number of MLP parameters ξ_{ℓ} on ε . Another limitation of our proof technique is that the number of heads grows proportionally to the output dimension while each head only outputs a scalar $d_{\text{head}}(\theta_{\ell}) = 1$. Obtaining a better balance between these two parameters is an interesting problem. As explained in the proof, these MLPs approximate a real-valued squaring operator $a \in \mathbb{R} \mapsto a^2 \in \mathbb{R}$, so we expect this dependency to be well-behaved in common situations, but our construction does not provide any a priori bound on how the magnitude of the tokens grows through the layers. The main hypothesis of Theorem 1 is that the underlying map, Λ^* , is a smooth (at least continuous) map for the weak* topology over measures (see Section 1.3 for some background). Since our results are not quantitative, this is not a strong restriction, and it enables a unifying study of transformers for any number, n, of tokens. It is, however, not clear how much this is a good model to conduct further quantitative studies, and we leave this exploration for future work.

3 Proof of Theorem 1

The proof of Theorem 1 is divided into three steps. First, we approximate the map Λ^* by "cylindrical maps", which are spanned by pointwise multiplication of self-attention maps, including also affine transformations. This approximation leverages the Stone-Weierstrass theorem. Second, we approximate the multiplication obtained in the first step using MLPs. We note that these MLPs are "context-free", that is, independent of the measures. Finally, we show how the obtained approximation can be represented using a deep transformer operating over a higher dimensional embedding space.

3.1 Step 1: Approximation by cylindrical mapping

In this section, we approximate the map Λ^* by a class of functions that are pointwise products of elementary attention and affine transforms. We coin these functions "cylindrical mappings" because similar functions have been used in [13] to define Sobolev regularity over the Wasserstein space. In contrast to deep transformer maps, which operate by composition, we consider here simple multiplication. Section 3.3 details how to switch from these multiplications to compositions. To this end, we apply the Stone-Weierstrass theorem to the *h*-th component map, $(\mu, x) \mapsto \Lambda^*(\mu, x)_h$, for each $h \in [d']$, where $\Lambda^*(\mu, x)_h \in \mathbb{R}$ is *h*-th component of vector $\Lambda^*(\mu, x) \in \mathbb{R}^d$. The key is to prove that the subalgebra of cylindrical mapping, constructed by the span of compositions of self-attentions and affine transforms separates points. This is proved by using the injectivity of the Radon transform (see Appendix A).

For $t \in [T]$ and $n \in [N]$, we define

$$\Gamma_{\tilde{\theta}_{t,n}}(\mu, x) := x + \sum_{h=1}^{d'} \tilde{W}_{t,n}^h \int \frac{\exp\left(\langle \tilde{Q}_{t,n}^h x, \tilde{K}_{t,n}^h y \rangle\right)}{\int \exp\left(\langle \tilde{Q}_{t,n}^h x, \tilde{K}_{t,n}^h y \rangle\right) \mathrm{d}\mu(y)} \tilde{V}_{t,n}^h y \,\mathrm{d}\mu(y), \ x \in \mathbb{R}^{d'}, \tag{12}$$

where $d_{\text{in}}(\tilde{\theta}_{t,n}) = d_{\text{out}}(\tilde{\theta}_{t,n}) = d', d_{\text{head}}(\tilde{\theta}_{t,n}) = k(\tilde{\theta}_{t,n}) = 1$ and

$$\tilde{\theta}_{t,n} := \{ \tilde{W}_{t,n}^h, \tilde{V}_{t,n}^h, \tilde{Q}_{t,n}^h, \tilde{K}_{t,n}^h \}_{h=1,\dots,d'} \subset \mathbb{R}^{d' \times 1} \times \mathbb{R}^{1 \times d'} \times \mathbb{R}^{1 \times d'} \times \mathbb{R}^{1 \times d'}.$$

We define affine transforms, $\mathcal{A}_{t,n} : \mathbb{R}^d \to \mathbb{R}^{d'}$, by

$$\mathcal{A}_{t,n}(x) := A_{t,n}x + b_{t,n},\tag{13}$$

where $A_{t,n} \in \mathbb{R}^{d' \times d}$, $b_{t,n} \in \mathbb{R}^{d'}$. Then we have the composition,

$$\Gamma_{\tilde{\theta}_{t,n}} \diamond \mathcal{A}_{t,n}(\mu, x) = \Gamma_{\tilde{\theta}_{t,n}}((\mathcal{A}_{t,n})_{\sharp} \mu, \mathcal{A}_{t,n}(x)) = A_{t,n} x + b_{t,n}$$

$$+ \sum_{h=1}^{d'} \tilde{W}_{t,n}^{h} \int \frac{\exp\left(\langle \tilde{Q}_{t,n}^{h}(A_{t,n} x + b_{t,n}), \tilde{K}_{t,n}^{h}(A_{t,n} y + b_{t,n})\rangle\right)}{\int \exp\left(\langle \tilde{Q}_{t,n}^{h}(A_{t,n} x + b_{t,n}), \tilde{K}_{t,n}^{h}(A_{t,n} y + b_{t,n})\rangle\right) d\mu(y)} \tilde{V}_{t,n}^{h}(A_{t,n} y + b_{t,n}) d\mu(y).$$

In what follows, we write

$$\tilde{\Gamma}_{t,n}(\mu, x) := \Gamma_{\tilde{\theta}_{t,n}} \diamond \mathcal{A}_{t,n}(\mu, x).$$
(14)

Lemma 1. For any $\varepsilon > 0$, there exist $T, N \in \mathbb{N}$, $\{\tilde{\theta}_{t,n}\}_{t \in [T], n \in [N]}$, $\{A_{t,n}, b_{t,n}\}_{t \in [T], n \in [N]}$ such that

$$\forall (\mu, x) \in \mathcal{P}(\Omega) \times \Omega, \quad \left| \left(\sum_{n=1}^{N} \tilde{\Gamma}_{T,n}(\mu, x) \odot \cdots \odot \tilde{\Gamma}_{1,n}(\mu, x) \right) - \Lambda^{\star}(\mu, x) \right| \leq \varepsilon.$$

Proof. We write

$$A_{t,n} = (a_{t,n}^1, \dots, a_{t,n}^{d'}), \quad b_{t,n} = (b_{t,n}^1, \dots, b_{t,n}^{d'}),$$

where $a_{t,n}^h \in \mathbb{R}^{d'}$ and $b_{t,n}^h \in \mathbb{R}$. We choose

$$\tilde{W}_{t,n}^h = (0, ..., 0, \underbrace{1}_{h-\text{th}}, 0, ..., 0) = e_h,$$

independently of t, n, where $\{e_h\}_{h \in [d']}$ is the standard basis in $\mathbb{R}^{d'}$, and

$$\tilde{V}_{t,n}^{h} = (0, ..., 0, \underbrace{v_{t,n}^{h}}_{h-\text{th}}, 0..., 0), \ \tilde{Q}_{t,n}^{h} = (0, ..., 0, \underbrace{c_{t,n}^{h}}_{h-\text{th}}, 0..., 0), \ \tilde{K}_{t,n}^{h} = (0, ..., 0, \underbrace{1}_{h-\text{th}}, 0..., 0),$$

where $v_{t,n}^h \in \mathbb{R}, c_{t,n}^h \in \mathbb{R}$. Then, as

$$\begin{split} \tilde{Q}_{t,n}^{h}(A_{t,n}x+b_{t,n}) &= c_{t,n}^{h}(\langle a_{t,n}^{h}, x \rangle + b_{t,n}^{h}), \\ \tilde{K}_{t,n}^{h}(A_{t,n}y+b_{t,n}) &= \langle a_{t,n}^{h}, x \rangle + b_{t,n}^{h}, \ \tilde{V}_{t,n}^{h}(A_{t,n}y+b_{t,n}) = v_{t,n}^{h}(\langle a_{t,n}^{h}, x \rangle + b_{t,n}^{h}), \end{split}$$

we see that

$$\begin{split} \tilde{\Gamma}_{t,n}(\mu, x) &= \Gamma_{\tilde{\theta}_{t,n}} \diamond \mathcal{A}_{t,n}(\mu, x) = \sum_{h=1}^{d'} e_h \bigg[\langle a_{t,n}^h, x \rangle + b_{t,n}^h \\ &+ \int \frac{\exp\left((\langle a_{t,n}^h, x \rangle + b_{t,n}^h) c_{t,n}^h (\langle a_{t,n}^h, y \rangle + b_{t,n}^h) \right)}{\int \exp\left((\langle a_{t,n}^h, x \rangle + b_{t,n}^h) c_{t,n}^h (\langle a_{t,n}^h, z \rangle + b_{t,n}^h) \bigg] \, \mathrm{d}\mu(z)} v_{t,n}^h (\langle a_{t,n}^h, y \rangle + b_{t,n}^h) \, \mathrm{d}\mu(y) \bigg]. \end{split}$$

Thus, we only need to show that the set,

$$\mathcal{A} := \operatorname{span} \left\{ \mathcal{P}(\Omega) \times \Omega \ni (\mu, x) \mapsto \sum_{n=1}^{N} \tilde{m}_{\lambda_{T,n}}(\mu, x) \cdots \tilde{m}_{\lambda_{1,n}}(\mu, x) \in \mathbb{R} \right\},\$$

is dense in $\mathcal{C}(\mathcal{P}(\Omega) \times \Omega; \mathbb{R})$; here, $\tilde{m}_{\lambda_{t,n}}(\mu, x)$ is defined by

$$\begin{split} \tilde{m}_{\lambda_{t,n}}(\mu, x) &\coloneqq \langle a_{t,n}, x \rangle + b_{t,n} \\ &+ \int \frac{\exp\left((\langle a_{t,n}, x \rangle + b_{t,n})c_{t,n}(\langle a_{t,n}, y \rangle + b_{t,n})\right)}{\int \exp\left((\langle a_{t,n}, x \rangle + b_{t,n})c_{t,n}(\langle a_{t,n}, z \rangle + b_{t,n})\right) \mathrm{d}\mu(z)} v_{t,n}(\langle a_{t,n}, y \rangle + b_{t,n}) \,\mathrm{d}\mu(y), \end{split}$$

where

 $\lambda_{t,n} := \{a_{t,n}, b_{t,n}, c_{t,n}, v_{t,n}\} \subset \mathbb{R}^d \times \mathbb{R} \times \mathbb{R} \times \mathbb{R}.$

If we can show that, by applying the denseness result to the *h*-th component map, $(\mu, x) \mapsto \Lambda^*(\mu, x)_h \in \mathbb{R}$ for each $h \in [d']$, where $\Lambda^*(\mu, x)_h \in \mathbb{R}$ is the *h*-th component of the vector $\Lambda^*(\mu, x) \in \mathbb{R}^{d'}$, then we can approximate the *h*-th component map with an element in \mathcal{A} . The concatenation of obtained elements takes the form $\sum_{n=1}^{N} \tilde{\Gamma}_{T,n}(\mu, x) \odot \cdots \odot \tilde{\Gamma}_{1,n}(\mu, x)$, thereby proving Lemma 1. In what follows, we verify the application of the Stone-Weierstrass theorem to the set \mathcal{A} .

It is straightforward to establish the stability of the sum and scalar product. The set \mathcal{A} contains the constant function, namely, by choosing T = N = 1, $c_{1,1} = v_{1,1} = 0$, $a_{1,1} = 0$ and $b_{1,1} = 1$.

For the separation, we first assume that $x \neq x'$ (that is, assume that $\exists i \in [d]$ such that $x_i \neq x'_i$). Specifically, by choosing T = N = 1, $a_{1,1} = (0, ..., 0, \underbrace{1}_{i-th}, 0, ..., 0)$, $c_{1,1} = v_{1,1} = 0$ and $b_{1,1} = 0$, we see

that

$$a_{\lambda_{1,1}}(\mu, x) = \langle a_{1,1}, x \rangle = x_i \neq x'_i = \langle a_{1,1}, x' \rangle = \tilde{m}_{\lambda_{1,1}}(\mu', x').$$

Next, we assume that x = x'. Again, we choose T = N = 1. It is enough to show that

$$\forall \lambda_{1,1}, \ \tilde{m}_{\lambda_{1,1}}(\mu, x) = \tilde{m}_{\lambda_{1,1}}(\mu', x) \text{ implies that } \mu = \mu'.$$

We note that the left-hand side implies that

 \tilde{m}

$$\frac{\int \exp\left(\left(\langle a_{1,1}, x \rangle + b_{t,n} \right) c_{1,1} \langle a_{1,1}, y \rangle\right) \langle a_{1,1}, y \rangle \,\mathrm{d}\mu(y)}{\int \exp\left(\langle a_{1,1}, x \rangle + b_{1,1} \rangle c_{1,1} \langle a_{1,1}, z \rangle\right) \,\mathrm{d}\mu(z)} = \frac{\int \exp\left(\left(\langle a_{1,1}, x \rangle + b_{1,1} \rangle c_{1,1} \langle a_{1,1}, y \rangle\right) \langle a_{1,1}, y \rangle \,\mathrm{d}\mu'(y)}{\int \exp\left(\left(\langle a_{1,1}, x \rangle + b_{1,1} \rangle c_{1,1} \langle a_{1,1}, z \rangle\right) \,\mathrm{d}\mu'(z)}.$$

By choosing $b_{1,1} \in \mathbb{R}$ so that $\langle a_{1,1}, x \rangle + b_{1,1} \neq 0$ (while x is frozen), letting $a_{1,1} = a$, and choosing $c_{1,1} = c/(\langle a_{1,1}, x \rangle + b_{1,1})$ with $a \in \mathbb{R}^d$ and $c \in \mathbb{R}$ arbitrary, we obtain

$$L(\mu)(a,c) = L(\mu')(a,c), \ \forall a \in \mathbb{R}^d, c \in \mathbb{R},$$

where

$$L(\mu)(a,c) := \frac{\int \exp(c\langle a, y \rangle) \langle a, y \rangle \, \mathrm{d}\mu(y)}{\int \exp(c\langle a, z \rangle) \, \mathrm{d}\mu(z)}$$

Then, Lemma 5 in Appendix A implies that $\mu = \mu'$ and the proof is complete.

3.2 Step 2: Approximation of the multiplication map by MLPs

In Section 3.1 (Step 1), we showed that the map $(\mu, x) \mapsto \Lambda^*(\mu, x)$ can be approximated by a cylindrical map

$$(\mu, x) \mapsto \sum_{n=1}^{N} \tilde{\Gamma}_{T,n}(\mu, x) \odot \cdots \odot \tilde{\Gamma}_{1,n}(\mu, x),$$

where $\Gamma_{t,n}$ was defined in (14). To obtain an approximator that can be represented by deep transformers, we will further approximate the multiplication map in the above using MLPs. We note that approximation of multiplication by MLP has been studied in detail, see for instance [11, Lemma 6.2]. We now prove the following result.

Lemma 2. Let $\tilde{\Gamma}_{t,n} : \mathcal{P}(\mathbb{R}^{d'}) \times \mathbb{R}^{d'} \to \mathbb{R}^{d'}$ be defined in (14). For any $\varepsilon > 0$, there exists an MLP $\Phi : \mathbb{R}^{2d'} \to \mathbb{R}^{d'}$ such that

$$\begin{aligned} \forall (\mu, x) \in \mathcal{P}(\Omega) \times \Omega, \quad \left| \sum_{n=1}^{N} \tilde{\Gamma}_{T,n}(\mu, x) \odot \cdots \odot \tilde{\Gamma}_{1,n}(\mu, x) \right. \\ \left. \left. - \sum_{n=1}^{N} \Phi\left(\tilde{\Gamma}_{T,n}(\mu, x), \Phi\left(\tilde{\Gamma}_{T-1,n}(\mu, x), \cdots \Phi\left(\tilde{\Gamma}_{2,n}(\mu, x), \Phi\left(\tilde{\Gamma}_{1,n}(\mu, x), \mathbf{1}_{d'} \right) \right) \right) \right) \right| &\leq \varepsilon. \end{aligned}$$

Proof. We note that

$$\tilde{\Gamma}_{T,n}(\mu)(x)\odot\cdots\odot\tilde{\Gamma}_{1,n}(\mu)(x) = \tilde{\Gamma}_{T,n}(\mu)(x)\odot\left(\tilde{\Gamma}_{T-1,n}(\mu)(x)\odot\cdots\odot\left(\tilde{\Gamma}_{2,n}(\mu)(x)\odot\left(\tilde{\Gamma}_{1,n}(\mu)(x)\odot\mathbf{1}_{d'}\right)\right)\right)$$

Because the component-wise multiplication map $(x, y) \in \mathbb{R}^{2d'} \mapsto x \odot y \in \mathbb{R}^{d'}$ is continuous, by the universality of MLPs, for any $\varepsilon > 0$ and R > 0, there exists an MLP $\Phi : \mathbb{R}^{2d'} \to \mathbb{R}^{d'}$ such that

 $\forall (x,y) \in B_{\mathbb{R}^{2d'}}(0,R), \quad |x \odot y - \Phi(x,y)| \le \varepsilon.$ (15)

Since $\Omega \subset \mathbb{R}^d$ is compact then $0 \leq C_\Omega := \sup_{x \in \Omega} \|x\|_2$ is finite. Thus we estimate that

$$\begin{split} \left| \tilde{\Gamma}_{t,n}(\mu, x) \right| &\leq \|A_{t,n}\|_2 C_{\Omega} + \|b_{t,n}\|_2 + \sum_{h=1}^{d'} (\|A_{t,n}\|_2 C_{\Omega} + \|b_{t,n}\|_2) \|\tilde{W}_{t,n}^h \tilde{V}_{t,n}^h\|_2 \\ &\leq \max_{t \in [T], n \in [N]} \left((\|A_{t,n}\|_2 C_{\Omega} + \|b_{t,n}\|_2)(1 + \sum_{h=1}^{d'} \|\tilde{W}_{t,n}^h \tilde{V}_{t,n}^h\|_2) \right) =: C_{\tilde{\Gamma}} \text{ for all } (\mu, x) \in \mathcal{P}(\Omega) \times \Omega, \quad (16) \end{split}$$

where the constant, $C_{\tilde{\Gamma}} > 0$, depends on Ω , $\tilde{W}_{t,n}^h, \tilde{V}_{t,n}^h, A_{t,n}, b_{t,n}$, but is independent of t, n, μ and x. Thus, using the universality in (15), choosing a large radius R > 0 depending on the constant $C_{\tilde{\Gamma}} > 0$, we can show that there exists an MLP $\Phi : \mathbb{R}^{2d'} \to \mathbb{R}^{d'}$ such that

$$\left| \tilde{\Gamma}_{T,n}(\mu, x) \odot \cdots \odot \tilde{\Gamma}_{1,n}(\mu, x) - \Phi \left(\tilde{\Gamma}_{T,n}(\mu, x), \Phi \left(\tilde{\Gamma}_{T-1,n}(\mu, x), \cdots \Phi \left(\tilde{\Gamma}_{2,n}(\mu, x), \Phi \left(\tilde{\Gamma}_{1,n}(\mu, x), \mathbf{1}_{d'} \right) \right) \right) \right) \right| \leq \frac{\varepsilon}{N},$$

which completes the proof.

Remark 1. (The challenge to derive quantitative estimates.) The key is to approximate and capture the mentioned multiplicity by MLPs, for which quantitative estimates have been studied, e.g., [11, Lemma 6.2], which is a variant of [39, Proposition 2]. However, the depth and width of MLPs depend on the bound of input variables. Specifically, an existential Φ in the above depends on the bound $C_{\tilde{\Gamma}}$ (see (16)), which in turn depends on parameters in $\tilde{\Gamma}_{t,n}$ that are chosen to approximate Γ^* within ε through the application of the Stone-Weierstrass theorem (see Lemma 1). Thus, providing the quantitative estimate for the MLP Φ is challenging.

3.3 Step 3: Realization of cylindrical mappings by deep transformers

In (Step 1) and (Step 2), we have shown that the map $(\mu, x) \mapsto \Lambda^*(\mu, x)$ can be approximated by the map

$$\sum_{n=1}^{N} \Phi\left(\tilde{\Gamma}_{T,n}(\mu, x), \Phi\left(\tilde{\Gamma}_{T-1,n}(\mu, x), \cdots \Phi\left(\tilde{\Gamma}_{2,n}(\mu, x), \Phi\left(\tilde{\Gamma}_{1,n}(\mu, x), \mathbf{1}_{d'}\right)\right)\right)\right),$$

where $\tilde{\Gamma}_{t,n}$, defined by (14), is the composition of self-attention and an affine transform, and Φ is some "context-free" MLP. To obtain the main result (Theorem 1), it is sufficient to show that this map can be represented by a deep transformer in the form of (10). Here, $\Gamma_{\theta_{t,n}} : \mathcal{P}(\mathbb{R}^{d+3d'}) \times \mathbb{R}^{d+3d'} \to \mathbb{R}^{d+3d'}$ is given by

$$\Gamma_{\theta_{t,n}}(\mu_{t,n}, (x, u, p, w)) = (x, u, p, w)$$

$$+ \sum_{h=1}^{d'} W_{t,n}^{h} \int \frac{\exp\left(\langle Q_{t,n}^{h}(x, u, p, w), K_{t,n}^{h}(y', v', q', z')\rangle\right)}{\int \exp\left(\langle Q_{t,n}^{h}(x, u, p, w), K_{t,n}^{h}(y, v, q, z)\rangle\right) d\mu_{t,n}(y, v, q, z)} V_{t,n}^{h}(y', v', q', z') d\mu_{t,n}(y', v', q', z')$$

for $\mu_{t,n} \in \mathcal{P}(\mathbb{R}^{d+3d'}), x \in \mathbb{R}^d, u, p, w \in \mathbb{R}^{d'}$, where

$$\theta_{t,n} = \{ W_{t,n}^h, V_{t,n}^h, Q_{t,n}^h, K_{t,n}^h \}_{h=1,\dots,d'} \subset \mathbb{R}^{(d+3d')\times 1} \times \mathbb{R}^{1\times (d+3d')} \times \mathbb{R}^{1\times (d+3d')} \times \mathbb{R}^{1\times (d+3d')};$$

thus, $\Gamma_{\theta_{t,n}}$ has the following size,

$$d_{\mathrm{in}}(\theta_{t,n}) = d_{\mathrm{out}}(\theta_{t,n}) = d + 3d', \quad d_{\mathrm{head}}(\theta_{t,n}) = k(\theta_{t,n}) = 1, \quad H(\theta_{t,n}) = d'.$$

We denote the MLPs by $F_{\xi_{t,n}} : \mathbb{R}^{d+3d'} \to \mathbb{R}^{d+3d'}$, with weight and bias parameters $\xi_{t,n}$.

The aim of this section is to prove, by construction, the following result.

Lemma 3. Let $\tilde{\Gamma}_{t,n} : \mathcal{P}(\mathbb{R}^{d'}) \times \mathbb{R}^{d'} \to \mathbb{R}^{d'}$ be defined in (14). Let $\Phi : \mathbb{R}^{2d'} \to \mathbb{R}^{d'}$ be an MLP. There exist $\xi_0, \xi_{t,n}, \xi_*$, and $\theta_0, \theta_{t,n}, \theta_*$ such that

$$\begin{aligned} \forall (\mu, x) \in \mathcal{P}(\Omega) \times \Omega, \quad \sum_{n=1}^{N} \Phi\left(\tilde{\Gamma}_{T,n}(\mu, x), \Phi\left(\tilde{\Gamma}_{T-1,n}(\mu, x), \cdots \Phi\left(\tilde{\Gamma}_{2,n}(\mu, x), \Phi\left(\tilde{\Gamma}_{1,n}(\mu, x), \mathbf{1}_{d'}\right)\right)\right)\right) \\ &= F_{\xi_*} \diamond \Gamma_{\theta_*} \diamond \left(\diamond_{n=1}^N \diamond_{t=1}^T F_{\xi_{t,n}} \diamond \Gamma_{\theta_{t,n}}\right) \diamond F_{\xi_0} \diamond \Gamma_{\theta_0}(\mu, x). \end{aligned}$$

Proof. The proof is based on the following scheme:

$$\begin{aligned} x \xrightarrow{F_{\xi_{0}} \diamond \Gamma_{\theta_{0}}} \begin{pmatrix} x \\ \mathcal{A}_{1,1}(x) \\ \varphi_{1,1}(x) \\ f_{1}(x) \end{pmatrix} \\ \xrightarrow{F_{\xi_{1,1}} \diamond \Gamma_{\theta_{1,1}}} \begin{bmatrix} x \\ \mathcal{A}_{2,1}(x) \\ \varphi_{2,1}(x) \\ f_{1}(x) \end{pmatrix} \xrightarrow{F_{\xi_{2,1}} \diamond \Gamma_{\theta_{2,1}}} \cdots \xrightarrow{F_{\xi_{T-1,1}} \diamond \Gamma_{\theta_{T-1,1}}} \begin{bmatrix} x \\ \mathcal{A}_{T,1}(x) \\ \varphi_{T,1}(x) \\ f_{1}(x) \end{pmatrix} \xrightarrow{F_{\xi_{T,1}} \diamond \Gamma_{\theta_{T,1}}} \begin{pmatrix} x \\ \mathcal{A}_{1,2}(x) \\ \varphi_{1,2}(x) \\ f_{2}(x) \end{pmatrix} \\ \xrightarrow{F_{\xi_{1,2}} \diamond \Gamma_{\theta_{1,2}}} \begin{bmatrix} x \\ \mathcal{A}_{2,2}(x) \\ \varphi_{2,2}(x) \\ f_{2}(x) \end{pmatrix} \xrightarrow{F_{\xi_{2,2}} \diamond \Gamma_{\theta_{2,2}}} \cdots \xrightarrow{F_{\xi_{T-1,2}} \diamond \Gamma_{\theta_{T-1,2}}} \begin{bmatrix} x \\ \mathcal{A}_{T,2}(x) \\ \varphi_{T,2}(x) \\ f_{2}(x) \end{pmatrix} \xrightarrow{F_{\xi_{T,2}} \diamond \Gamma_{\theta_{T,2}}} \begin{pmatrix} x \\ \mathcal{A}_{1,3}(x) \\ \varphi_{1,3}(x) \\ f_{3}(x) \end{pmatrix} \\ \vdots \end{aligned}$$

$$\frac{F_{\xi_{1,N}} \circ \Gamma_{\theta_{1,N}}}{[\mathbf{Step B}]} \begin{pmatrix} x \\ \mathcal{A}_{2,N}(x) \\ \varphi_{2,N}(x) \\ f_{N}(x) \end{pmatrix} \xrightarrow{F_{\xi_{2,N}} \circ \Gamma_{\theta_{2,N}}}{[\mathbf{Step B}]} \cdots \xrightarrow{F_{\xi_{T-1,N}} \circ \Gamma_{\theta_{T-1,N}}}{[\mathbf{Step B}]} \begin{pmatrix} x \\ \mathcal{A}_{T,N}(x) \\ \varphi_{T,N}(x) \\ f_{N}(x) \end{pmatrix} \xrightarrow{F_{\xi_{T,N}} \circ \Gamma_{\theta_{T,N}}}{[\mathbf{Step C}]} \begin{pmatrix} x \\ \mathcal{A}_{1,N+1}(x) \\ \varphi_{1,N+1}(x) \\ f_{N+1}(x) \end{pmatrix} \xrightarrow{F_{\xi_{*}} \circ \Gamma_{\theta_{*}}}{[\mathbf{Step D}]} f_{N+1}(x) = \sum_{n=1}^{N} \Phi\left(\tilde{\Gamma}_{T,n}(\mu, x), \Phi\left(\tilde{\Gamma}_{T-1,n}(\mu, x), \cdots \Phi\left(\tilde{\Gamma}_{2,n}(\mu, x), \Phi\left(\tilde{\Gamma}_{1,n}(\mu, x), \mathbf{1}_{d'}\right)\right)\right)\right),$$

where $\varphi_{t,n}: \mathbb{R}^d \to \mathbb{R}^{d'}$ is given by

$$\varphi_{t,n}(x) := \begin{cases} \Phi\left(\tilde{\Gamma}_{t-1,n}(\mu, x), \Phi\left(\tilde{\Gamma}_{t-2,n}(\mu, x), \cdots \Phi\left(\tilde{\Gamma}_{2,n}(\mu, x), \Phi\left(\tilde{\Gamma}_{1,n}(\mu, x), \mathbf{1}_{d'}\right)\right)\right)\right), & t \ge 2 \\ \mathbf{1}_{d'}, & t = 1 \end{cases},$$

and $f_n : \mathbb{R}^d \to \mathbb{R}^{d'}$ by

$$f_n(x) := \begin{cases} \sum_{i=1}^{n-1} \varphi_{T,i}(x), & n \ge 2\\ 0 & n = 1 \end{cases}$$

Furthermore, the $\mathcal{A}_{t,n} : \mathbb{R}^d \to \mathbb{R}^{d'}$ are the affine transforms chosen in Lemma 1. Here, Γ_{θ_0} , $\Gamma_{\theta_{t,n}}$, Γ_{θ_*} , F_{ξ_0} , $F_{\xi_{t,n}}$ and F_{ξ_*} will be specified below, in the following steps:

[Step A] Let $\Gamma_{\theta_0}(\mu) : \mathbb{R}^d \to \mathbb{R}^d$ be

 $\Gamma_{\theta_0}(\mu, x) = x,$

and let $F_{\xi_0}: \mathbb{R}^d \to \mathbb{R}^{d+3d'}$ be the affine transform defined by

$$F_{\xi_0}(x) := (x, A_{1,1}x + b_{1,1}, \mathbf{1}_{d'}, 0) = (x, \mathcal{A}_{1,1}(x), \varphi_{1,1}(x), f_1(x)).$$

Then we see that

$$F_{\xi_0} \diamond \Gamma_{\theta_0}(\mu, x) = (x, \mathcal{A}_{1,1}(x), \varphi_{1,1}(x), f_1(x)),$$

and

$$\mu_{1,1} := (F_{\xi_0} \diamond \Gamma_{\theta_0}(\mu))_{\sharp} \mu = (\mu, (\mathcal{A}_{1,1})_{\sharp} \mu, (\varphi_{1,1})_{\sharp} \mu, (f_1)_{\sharp} \mu).$$

We proceed with [Step B] in which we handle the case when n = t = 1.

[Step B] Let t = 1, ..., T - 1 and n = 1, ..., N. We already have that

$$\left(\diamond_{j=1}^{t-1}F_{\xi_{j,n}}\diamond\Gamma_{\theta_{j,n}}\right)\diamond\left(\diamond_{i=1}^{n-1}\diamond_{s=1}^{T}F_{\xi_{s,i}}\diamond\Gamma_{\theta_{s,i}}\right)\diamond F_{\xi_{0}}\diamond\Gamma_{\theta_{0}}(\mu,x)=(x,\mathcal{A}_{t,n}(x),\varphi_{t,n}(x),f_{n}(x))$$

and

$$\mu_{t,n} := \left(\left(\diamond_{j=1}^{t-1} F_{\xi_{j,n}} \diamond \Gamma_{\theta_{j,n}} \right) \diamond \left(\diamond_{i=1}^{n-1} \diamond_{s=1}^{T} F_{\xi_{s,i}} \diamond \Gamma_{\theta_{s,i}} \right) \diamond F_{\xi_{0}} \diamond \Gamma_{\theta_{0}}(\mu) \right)_{\sharp} \mu \\ = \left(\mu, (\mathcal{A}_{t,n})_{\sharp} \mu, (\varphi_{t,n})_{\sharp} \mu, (f_{n})_{\sharp} \mu \right).$$

When n = 1 or t = 1, the above reduces to $\diamond_{i=1}^{n-1} \diamond_{s=1}^T F_{\xi_{s,i}} \diamond \Gamma_{\theta_{s,i}} = I_{d+3d'}$ or $\diamond_{j=1}^{t-1} F_{\xi_{j,n}} \diamond \Gamma_{\theta_{j,n}} = I_{d+3d'}$. Let $\Gamma_{\theta_{t,n}}(\mu_{t,n}) : \mathbb{R}^{d+3d'} \to \mathbb{R}^{d+3d'}$ be given by

$$\begin{split} \Gamma_{\theta_{t,n}}(\mu_{t,n},(x,u,p,w)) \\ &= \left(x,u + \sum_{h=1}^{d'} \tilde{W}^{h}_{t,n} \int \frac{\exp\left(\langle \tilde{Q}^{h}_{t,n}u, \tilde{K}^{h}_{t,n}v'\rangle\right)}{\int \exp\left(\langle \tilde{Q}^{h}_{t,n}u, \tilde{K}^{h}_{t,n}v\rangle\right) \mathrm{d}\mu_{t,n}(y,v,q,z)} \tilde{V}^{h}_{t,n}v' \,\mathrm{d}\mu_{t,n}(y',v',q',z'), p, w\right) \\ &= \left(x, \Gamma_{\tilde{\theta}_{t,n}}((\mathcal{A}_{t,n})_{\sharp}\mu,u), p, w\right), \end{split}$$

where

$$\{\tilde{W}_{t,n}^h, \tilde{V}_{t,n}^h, \tilde{Q}_{t,n}^h, \tilde{K}_{t,n}^h\}_{h=1,\dots,d'} \subset \mathbb{R}^{d' \times 1} \times \mathbb{R}^{1 \times d'} \times \mathbb{R}^{1 \times d'} \times \mathbb{R}^{1 \times d'},$$

which were specified in Lemma 1. Here, we choose

$$W_{t,n}^{h} = (O, \tilde{W}_{t,n}^{h}, O, O), \ V_{t,n}^{h} = (O, \tilde{V}_{t,n}^{h}, O, O), \ Q_{t,n}^{h} = (O, \tilde{Q}_{t,n}^{h}, O, O), \ K_{t,n}^{h} = (O, \tilde{K}_{t,n}^{h}, O, O).$$

Let $F_{\xi_{t,n}}: \mathbb{R}^{d+3d'} \rightarrow \mathbb{R}^{d+3d'}$ be an MLP defined by

 $F_{\xi_{t,n}}(x, u, p, w) = (x, A_{t+1,n}x + b_{t+1,n}, \Phi(u, p), w) = (x, \mathcal{A}_{t+1,n}(x), \Phi(u, p), w).$

Then we have

$$\begin{split} (\diamond_{j=1}^{t} F_{\xi_{j,n}} \diamond \Gamma_{\theta_{j,n}}) &\diamond (\diamond_{i=1}^{n-1} \diamond_{s=1}^{T} F_{\xi_{s,i}} \diamond \Gamma_{\theta_{s,i}}) \diamond F_{\xi_{0}} \diamond \Gamma_{\theta_{0}}(\mu, x) \\ &= F_{\xi_{t,n}} \diamond \Gamma_{\theta_{t,n}} \diamond (\diamond_{j=1}^{t-1} F_{\xi_{j,n}} \diamond \Gamma_{\theta_{j,n}}) \diamond (\diamond_{i=1}^{n-1} \diamond_{s=1}^{T} F_{\xi_{s,i}} \diamond \Gamma_{\theta_{s,i}}) \diamond F_{\xi_{0}} \diamond \Gamma_{\theta_{0}}(\mu, x) \\ &= F_{\xi_{t,n}} \diamond \Gamma_{\theta_{t,n}}(\mu_{t,n}, (x, \mathcal{A}_{t,n}(x), \varphi_{t,n}(x), f_{n}(x))) \\ &= F_{\xi_{t,n}}(x, \Gamma_{\tilde{\theta}_{t,n}}((\mathcal{A}_{t,n})_{\sharp}\mu, \mathcal{A}_{t,n}(x)), \varphi_{t,n}(x), f_{n}(x)) \\ &= F_{\xi_{t,n}}(x, \tilde{\Gamma}_{t,n}(\mu, x), \varphi_{t,n}(x), f_{n}(x)) \\ &= (x, \mathcal{A}_{t+1,n}(x), \varphi_{t+1,n}(x), f_{n}(x))) \end{split}$$

and

$$\mu_{t+1,n} := \left(\left(\diamond_{j=1}^t F_{\xi_{j,n}} \diamond \Gamma_{\theta_{j,n}} \right) \diamond \left(\diamond_{i=1}^{n-1} \diamond_{s=1}^T F_{\xi_{s,i}} \diamond \Gamma_{\theta_{s,i}} \right) \diamond F_{\xi_0} \diamond \Gamma_{\theta_0}(\mu) \right)_{\sharp} \mu$$
$$= \left(\mu, (\mathcal{A}_{t+1,n})_{\sharp} \mu, (\varphi_{t+1,n})_{\sharp} \mu, (f_n)_{\sharp} \mu \right).$$

We repeat [Step B] until obtaining $\mu_{T,n}$. Once $\mu_{T,n}$ is obtained, we proceed with [Step C].

[Step C] Let $\Gamma_{\theta_{T,n}}(\mu_{T,n}) : \mathbb{R}^{d+3d'} \to \mathbb{R}^{d+3d'}$ be given by

$$\begin{split} \Gamma_{\theta_{T,n}}(\mu_{T,n},(x,u,p,w)) \\ &= \left(x,u + \sum_{h=1}^{d'} \tilde{W}^{h}_{T,n} \int \frac{\exp\left(\langle \tilde{Q}^{h}_{T,n}u, \tilde{K}^{h}_{T,n}v'\rangle\right)}{\int \exp\left(\langle \tilde{Q}^{h}_{T,n}u, \tilde{K}^{h}_{T,n}v\rangle\right) \mathrm{d}\mu_{T,n}(y,v,q,z)} \tilde{V}^{h}_{T,n}v' \,\mathrm{d}\mu_{T,n}(y',v',q',z'), p, w\right) \\ &= \left(x, \Gamma_{\tilde{\theta}_{T,n}}((\mathcal{A}_{T,n})_{\sharp}\mu, u), p, w\right) \end{split}$$

Let $F_{\xi_{T,n}} : \mathbb{R}^{d+3d'} \to \mathbb{R}^{d+3d'}$ be an MLP defined by

 $F_{\xi_{T,n}}(x, u, p, w) = (x, W_{1,n+1}x + b_{1,n+1}, \mathbf{1}_{d'}, w + \Phi(u, p)) = (x, \mathcal{A}_{1,n+1}(x), \varphi_{1,n+1}(x), w + \Phi(u, p)).$ When n = N, we define by $\mathcal{A}_{1,N+1}(x) := 0$ and $\varphi_{1,N+1} := 0$ in the above. We find that

$$(\diamond_{j=1}^T F_{\xi_{j,n}} \diamond \Gamma_{\theta_{j,n}}) \diamond (\diamond_{i=1}^{n-1} \diamond_{s=1}^T F_{\xi_{s,i}} \diamond \Gamma_{\theta_{s,i}}) \diamond F_{\xi_0} \diamond \Gamma_{\theta_0}(\mu, x)$$
$$= F_{\xi_{T,n}}(x, \tilde{\Gamma}_{T,n}(\mu, x), \varphi_{T,n}(x), f_n(x))$$
$$= (x, \mathcal{A}_{1,n+1}(x), \varphi_{1,n+1}(x), f_{n+1}(x)))$$

and

$$\mu_{T+1,n} := \left(\left(\diamond_{j=1}^T F_{\xi_{j,n}} \diamond \Gamma_{\theta_{j,n}} \right) \diamond \left(\diamond_{i=1}^{n-1} \diamond_{s=1}^T F_{\xi_{s,i}} \diamond \Gamma_{\theta_{s,i}} \right) \diamond F_{\xi_0} \diamond \Gamma_{\theta_0}(\mu) \right)_{\sharp} \mu$$
$$= \left(\mu, (\mathcal{A}_{1,n+1})_{\sharp} \mu, (\varphi_{1,n+1})_{\sharp} \mu, (f_{n+1})_{\sharp} \mu \right).$$

Denoting

$$\mu_{1,n+1} := \mu_{T+1,n},$$

we return to [Step B], and repeat [Step B] and [Step C] until obtaining $\mu_{T+1,N}$. Once $\mu_{T+1,N}$ is obtained, we proceed with [Step D].

[Step D] Let $\Gamma_{\theta_*}(\mu_{T+1,N}) : \mathbb{R}^{d+3d'} \to \mathbb{R}^{d+3d'}$ be given by

$$\Gamma_{\theta_*}(\mu_{T+1,N}, (x, u, p, w)) = (x, u, p, w)$$

and let $F_{\xi_*}:\mathbb{R}^{d+3d'}\to\mathbb{R}^{d'}$ be the affine transform defined by

$$F_{\xi_*}(x, u, p, w) := w$$

Then we conclude that

$$F_{\xi_*} \diamond \Gamma_{\theta_*} \diamond \left(\diamond_{n=1}^N \diamond_{s=1}^T F_{\xi_{s,n}} \diamond \Gamma_{\theta_{s,n}} \right) \diamond F_{\xi_0} \diamond \Gamma_{\theta_0}(\mu, x)$$

$$= F_{\xi_*} \diamond \Gamma_{\theta_*} \left(\mu_{T+1,N}, (x, \mathcal{A}_{1,N+1}(x), \varphi_{1,n+1}(x), f_{N+1}(x)) \right) \right) = f_{N+1}(x)$$

$$= \sum_{n=1}^N \Phi \left(\tilde{\Gamma}_{T,n}(\mu, x), \Phi \left(\tilde{\Gamma}_{T-1,n}(\mu, x), \cdots \Phi \left(\tilde{\Gamma}_{2,n}(\mu, x), \Phi \left(\tilde{\Gamma}_{1,n}(\mu, x), \mathbf{1}_{d'} \right) \right) \right) \right).$$

Discussion

A limitation of our method is that it is not quantitative. Using, for instance, the Wasserstein distance between token distributions could be a way to impose smoothness on the map to obtain quantitative bounds. Our proof relies on the approximation of the map along each dimension and the use of a commuting architecture (the transformer layers are multiplied together to obtain the output). This results in a growth of the number of heads proportional to the dimension. Lowering this dependency would require the development of new proof techniques beyond the use of the Stone-Weierstrass theorem. Additionally, it does not handle masked attention, which means that the architectures considered are not causal and are permutation invariant.

Acknowledgements

T. Furuya was partially supported by JSPS KAKENHI Grant Number JP24K16949. M.V. de Hoop carried out the work while he was an invited professor at the Centre Sciences des Données at Ecole Normale Supérieure, Paris. He acknowledges the support of the Simons Foundation under the MATH + X Program, the Department of Energy under grant DE-SC0020345, and the corporate members of the Geo-Mathematical Imaging Group at Rice University. The work of G. Peyré was supported by the European Research Council (ERC project WOLF) and the French government under management of Agence Nationale de la Recherche as part of the "Investissements d'avenir" program, reference ANR-19-P3IA-0001 (PRAIRIE 3IA Institute).

References

- [1] Andrei Agrachev and Cyril Letrouit. Generic controllability of equivariant systems and applications to particle systems and neural networks. *arXiv preprint arXiv:2404.08289*, 2024.
- [2] Kwangjun Ahn, Xiang Cheng, Hadi Daneshmand, and Suvrit Sra. Transformers learn to implement preconditioned gradient descent for in-context learning. Advances in Neural Information Processing Systems, 36, 2024.
- [3] Silas Alberti, Niclas Dern, Laura Thesing, and Gitta Kutyniok. Sumformer: Universal approximation for efficient transformers. In *Topological, Algebraic and Geometric Learning Workshops* 2023, pages 72–86. PMLR, 2023.
- [4] Tom Brown, Benjamin Mann, Nick Ryder, Melanie Subbiah, Jared D Kaplan, Prafulla Dhariwal, Arvind Neelakantan, Pranav Shyam, Girish Sastry, Amanda Askell, et al. Language models are few-shot learners. Advances in neural information processing systems, 33:1877–1901, 2020.
- [5] Valérie Castin, Pierre Ablin, and Gabriel Peyré. How smooth is attention? In ICML 2024, 2024.
- [6] David Chiang, Peter Cholak, and Anand Pillay. Tighter bounds on the expressivity of transformer encoders. In *International Conference on Machine Learning*, pages 5544–5562. PMLR, 2023.
- [7] Christa Cuchiero, Martin Larsson, and Josef Teichmann. Deep neural networks, generic universal interpolation, and controlled odes. SIAM Journal on Mathematics of Data Science, 2(3):901–919, 2020.
- [8] George Cybenko. Approximation by superpositions of a sigmoidal function. Mathematics of control, signals and systems, 2(4):303-314, 1989.
- [9] Gwendoline De Bie, Gabriel Peyré, and Marco Cuturi. Stochastic deep networks. In *International Conference on Machine Learning*, pages 1556–1565. PMLR, 2019.
- [10] Alexey Dosovitskiy, Lucas Beyer, Alexander Kolesnikov, Dirk Weissenborn, Xiaohua Zhai, Thomas Unterthiner, Mostafa Dehghani, Matthias Minderer, Georg Heigold, Sylvain Gelly, et al. An image is worth 16x16 words: Transformers for image recognition at scale. arXiv preprint arXiv:2010.11929, 2020.

- [11] Dennis Elbrächter, Philipp Grohs, Arnulf Jentzen, and Christoph Schwab. Dnn expression rate analysis of high-dimensional pdes: Application to option pricing. *Constructive Approximation*, 55(1):3–71, 2022.
- [12] Nelson Elhage, Neel Nanda, Catherine Olsson, Tom Henighan, Nicholas Joseph, Ben Mann, Amanda Askell, Yuntao Bai, Anna Chen, Tom Conerly, et al. A mathematical framework for transformer circuits. *Transformer Circuits Thread*, 1(1):12, 2021.
- [13] Massimo Fornasier, Giacomo E Sodini, and Giuseppe Savaré. Density of subalgebras of lipschitz functions in metric sobolev spaces and applications to sobolev-wasserstein spaces. *Journal of Functional Analysis*, 285(11), 2023.
- [14] Takashi Furuya, Michael Puthawala, Matti Lassas, and Maarten V de Hoop. Globally injective and bijective neural operators. arXiv preprint arXiv:2306.03982, 2023.
- [15] Borjan Geshkovski, Cyril Letrouit, Yury Polyanskiy, and Philippe Rigollet. The emergence of clusters in self-attention dynamics. arXiv preprint arXiv:2305.05465, 2023.
- [16] Borjan Geshkovski, Cyril Letrouit, Yury Polyanskiy, and Philippe Rigollet. A mathematical perspective on transformers. arXiv preprint arXiv:2312.10794, 2023.
- [17] Boris Hanin and Mark Sellke. Approximating continuous functions by relu nets of minimal width. arXiv preprint arXiv:1710.11278, 2017.
- [18] Kurt Hornik, Maxwell Stinchcombe, and Halbert White. Multilayer feedforward networks are universal approximators. *Neural networks*, 2(5):359–366, 1989.
- [19] Nicolas Keriven and Gabriel Peyré. Universal invariant and equivariant graph neural networks. Advances in Neural Information Processing Systems, 32, 2019.
- [20] Nikola Kovachki, Zongyi Li, Burigede Liu, Kamyar Azizzadenesheli, Kaushik Bhattacharya, Andrew Stuart, and Anima Anandkumar. Neural operator: Learning maps between function spaces with applications to pdes. *Journal of Machine Learning Research*, 24(89):1–97, 2023.
- [21] Anastasis Kratsios, Valentin Debarnot, and Ivan Dokmanić. Small transformers compute universal metric embeddings. Journal of Machine Learning Research, 24(170):1–48, 2023.
- [22] Anastasis Kratsios, Chong Liu, Matti Lassas, Maarten V de Hoop, and Ivan Dokmanić. An approximation theory for metric space-valued functions with a view towards deep learning. arXiv preprint arXiv:2304.12231, 2023.
- [23] Anastasis Kratsios and Léonie Papon. Universal approximation theorems for differentiable geometric deep learning. Journal of Machine Learning Research, 23(196):1–73, 2022.
- [24] Shengjie Luo, Shanda Li, Shuxin Zheng, Tie-Yan Liu, Liwei Wang, and Di He. Your transformer may not be as powerful as you expect. Advances in Neural Information Processing Systems, 35:4301–4315, 2022.
- [25] Arvind Mahankali, Tatsunori B Hashimoto, and Tengyu Ma. One step of gradient descent is provably the optimal in-context learner with one layer of linear self-attention. arXiv preprint arXiv:2307.03576, 2023.
- [26] William Merrill and Ashish Sabharwal. The expressive power of transformers with chain of thought. arXiv preprint arXiv:2310.07923, 2023.
- [27] Luis Müller, Mikhail Galkin, Christopher Morris, and Ladislav Rampášek. Attending to graph transformers. arXiv preprint arXiv:2302.04181, 2023.
- [28] Swaroop Nath, Harshad Khadilkar, and Pushpak Bhattacharyya. Transformers are expressive, but are they expressive enough for regression? *arXiv preprint arXiv:2402.15478*, 2024.
- [29] Allan Pinkus. Approximation theory of the mlp model in neural networks. Acta numerica, 8:143– 195, 1999.

- [30] Charles R Qi, Hao Su, Kaichun Mo, and Leonidas J Guibas. Pointnet: Deep learning on point sets for 3d classification and segmentation. In *Proceedings of the IEEE conference on computer* vision and pattern recognition, pages 652–660, 2017.
- [31] Michael E Sander, Pierre Ablin, Mathieu Blondel, and Gabriel Peyré. Sinkformers: Transformers with doubly stochastic attention. In *International Conference on Artificial Intelligence and Statistics*, pages 3515–3530. PMLR, 2022.
- [32] Michael E Sander, Raja Giryes, Taiji Suzuki, Mathieu Blondel, and Gabriel Peyré. How do transformers perform in-context autoregressive learning? arXiv preprint arXiv:2402.05787, 2024.
- [33] Lena Strobl, William Merrill, Gail Weiss, David Chiang, and Dana Angluin. What formal languages can transformers express? a survey. Transactions of the Association for Computational Linguistics, 12:543–561, 2024.
- [34] Paulo Tabuada and Bahman Gharesifard. Universal approximation power of deep residual neural networks through the lens of control. *IEEE Transactions on Automatic Control*, 2022.
- [35] Ashish Vaswani, Noam Shazeer, Niki Parmar, Jakob Uszkoreit, Llion Jones, Aidan N Gomez, Lukasz Kaiser, and Illia Polosukhin. Attention is all you need. Advances in neural information processing systems, 30, 2017.
- [36] Johannes von Oswald, Eyvind Niklasson, Maximilian Schlegel, Seijin Kobayashi, Nicolas Zucchet, Nino Scherrer, Nolan Miller, Mark Sandler, Max Vladymyrov, Razvan Pascanu, et al. Uncovering mesa-optimization algorithms in transformers. arXiv preprint arXiv:2309.05858, 2023.
- [37] James Vuckovic, Aristide Baratin, and Remi Tachet des Combes. A mathematical theory of attention. arXiv preprint arXiv:2007.02876, 2020.
- [38] Keyulu Xu, Weihua Hu, Jure Leskovec, and Stefanie Jegelka. How powerful are graph neural networks? arXiv preprint arXiv:1810.00826, 2018.
- [39] Dmitry Yarotsky. Error bounds for approximations with deep relu networks. *Neural Networks*, 94:103–114, 2017.
- [40] Chulhee Yun, Srinadh Bhojanapalli, Ankit Singh Rawat, Sashank J Reddi, and Sanjiv Kumar. Are transformers universal approximators of sequence-to-sequence functions? arXiv preprint arXiv:1912.10077, 2019.
- [41] Ruiqi Zhang, Spencer Frei, and Peter L Bartlett. Trained transformers learn linear models incontext. arXiv preprint arXiv:2306.09927, 2023.
- [42] Ding-Xuan Zhou. Universality of deep convolutional neural networks. Applied and computational harmonic analysis, 48(2):787–794, 2020.

A Radon transform and injectivity

Lemma 4. Let $\Omega \subset \mathbb{R}$ be a compact set, and let $\mu, \nu \in \mathcal{P}(\Omega)$. Then,

$$L_1(\mu)(c) = L_1(\nu)(c), \ \forall c \in \mathbb{R}, \ \Rightarrow \ \mu = \nu.$$

where, for $k \in \mathbb{N}$,

$$L_k(\mu)(c) := \frac{\int e^{cy} y^k \,\mathrm{d}\mu(y)}{\int e^{cy} \,\mathrm{d}\mu(y)}$$

Proof. One has

$$L_k(\mu)'(c) = L_{k+1}(\mu)(c) - L_k(\mu)(c)L_1(\mu)(c).$$

So by recursion, we have that

$$L_1(\mu)(c) = L_1(\nu)(c), \ \forall c \in \mathbb{R}, \quad \Rightarrow \quad L_k(\mu)(c) = L_k(\nu)(c), \ \forall c \in \mathbb{R}, \ \forall k \ge 1,$$

So evaluating this at c = 0, we obtain that

$$L_k(\mu)(0) = L_k(\nu)(0), \ \forall k \ge 1 \quad \Leftrightarrow \quad \forall k, \int y^k d\mu(y) = \int y^k d\nu(y)$$

which is equivalent to $\mu = \nu$.

Lemma 5. Let $\Omega \subset \mathbb{R}^d$ be a compact set, and let $\mu, \nu \in \mathcal{P}(\Omega)$. Then,

$$L(\mu)(a,c) = L(\nu)(a,c), \ \forall a \in \mathbb{R}^d, \forall c \in \mathbb{R}, \ \Rightarrow \ \mu = \nu$$

where

$$L(\mu)(a,c) := \frac{\int \exp(c\langle a, y \rangle) \langle a, y \rangle \, \mathrm{d}\mu(y)}{\int \exp(c\langle a, y \rangle) \, \mathrm{d}\mu(y)}.$$

Proof. We define

$$\forall e \in \mathbb{S}^d, \quad \mu^e := (P_e)_{\sharp} \mu$$

where \mathbb{S}^d is the *d*-dimensional sphere, and $P_e(x) = \langle x, e \rangle$ is the projection on *e*. We see that

$$L(\mu)(e,c) = \frac{\int \exp(c\langle e, y \rangle) \langle e, y \rangle \,\mathrm{d}\mu(y)}{\int \exp(c\langle e, y \rangle) \,\mathrm{d}\mu(y)} = \frac{\int e^{cs} s \,\mathrm{d}\mu^e(s)}{\int e^{cs} \,\mathrm{d}\mu^e(s)}.$$

By Lemma 4, we can show that

 $\forall e, (P_e)_{\sharp} \mu = (P_e)_{\sharp} \nu$

which implies that by the injectivity of the Radon transform

 $\mu = \nu.$

		-
		н