

The Generating Idempotent Is a Minimum-Weight Codeword for Some Binary BCH Codes

Yaron Shany¹ and Amit Berman¹

¹Samsung Semiconductor Israel R&D Center, 146 Derech Menachem Begin St., Tel Aviv 6492103, Israel. Emails: {yaron.shany, amit.berman}@samsung.com

August 23, 2024

Abstract

In a paper from 2015, Ding *et al.* (IEEE Trans. IT, May 2015) conjectured that for odd m , the minimum distance of the binary BCH code of length $2^m - 1$ and designed distance $2^{m-2} + 1$ is equal to the Bose distance calculated in the same paper.

In this paper, we prove the conjecture. In fact, we prove a stronger result: the weight of the generating idempotent is equal to the Bose distance for both odd and even m . Our main tools are some new properties of the so-called fibbinary integers, in particular, the splitting field of related polynomials, and the relation of these polynomials to the idempotent of the BCH code.

1 Introduction

BCH codes are widely used in storage systems and various communication systems, including optical communications, digital video broadcasting, and more. Despite being introduced more than sixty years ago [Hoc59], [BR60], the exact minimum distance of BCH codes is known only for a small number of cases; we refer the reader to the recent survey [DL24] for an account of the current known results. The main objective of the current paper is to resolve the minimum distance for a case that was left as a conjecture in [DDZ15].

More specifically, for positive integers m, d with d odd, let $\text{BCH}(m, d)$ be the (primitive, narrow-sense) binary BCH code of length $2^m - 1$ and designed distance d , that is, the binary cyclic code of length $2^m - 1$ having zeros exactly $\alpha, \alpha^2, \dots, \alpha^{d-1}$ and their conjugates, where α is a primitive element of \mathbb{F}_{2^m} .¹ Equivalently, $\text{BCH}(m, d)$ is the binary cyclic code with generator polynomial $\text{lcm} \{M_{\alpha^i}(X) \mid i \in \{1, \dots, d-1\}\}$, where for

¹For a prime power q , we let \mathbb{F}_q be the finite field of q elements.

$\beta \in \mathbb{F}_{2^m}$, $M_\beta(X)$ is the minimal polynomial of β over \mathbb{F}_2 . It is well known by the *BCH bound* that the minimum distance² of $\text{BCH}(m, d)$ is at least d .

It may happen that $\text{BCH}(m, d_1) = \text{BCH}(m, d_2)$ for $d_2 > d_1$: starting with d_1 and including all conjugates of $\alpha, \alpha^2, \dots, \alpha^{d_1-1}$ as zeros, it may turn out that the longer progression $\alpha, \alpha^2, \dots, \alpha^{d_2-1}$ appears in the set of zeros. Starting with a *designed distance* of δ , the largest odd d for which $\text{BCH}(m, \delta) = \text{BCH}(m, d)$ is called the *Bose distance*. So, the minimum distance of $\text{BCH}(m, \delta)$ is *at least* the Bose distance, but it is possible that the minimum distance of $\text{BCH}(m, \delta)$ is larger than the Bose distance. In general, determining the minimum distance of BCH codes is a notoriously hard problem [DL24].

In a paper from 2015 [DDZ15], Ding *et al.* found the Bose distance d_B of $\text{BCH}(m, \delta)$ for $\delta := 2^{m-2} + 1$, proved that for even m it is equal to the minimum distance, and posed the following conjecture:

Conjecture 1.1 (Ding *et al.*, Conjecture 1 of [DDZ15]). *Let m be odd. Then the minimum distance d of $\text{BCH}(m, 2^{m-2} + 1)$ equals the Bose distance, that is,*

$$d = d_B = \frac{2^m + 1}{3}.$$

It is also mentioned in [DDZ15] that to prove the conjecture, it is sufficient to find some codeword whose weight equals d_B , and that checking some small values of m , it seems that the generating idempotent of the code has this property. However, the authors of [DDZ15] state that they were not able to prove this in general, and invite the reader to attack this open problem [DDZ15, p. 2355].³

The main result of the current paper is the following theorem.

Theorem 1.2. *For all integer $m \geq 4$ (both odd and even), the weight w of the generating idempotent of $\text{BCH}(m, 2^{m-2} + 1)$ is equal to the Bose distance found in [DDZ15], that is,*

$$w = d_B = \begin{cases} \frac{1}{3}(2^m + 1) & \text{if } m \text{ is odd} \\ \frac{1}{3}(2^m - 1) & \text{if } m \text{ is even.} \end{cases}$$

In particular, Theorem 1.2 proves Conjecture 1.1. Moreover, although for the case of even m the minimum distance was settled in [DDZ15], it was not previously known that for even m , the generating idempotent attains the minimum weight.

1.1 Proof technique

Our main tools are the relation of the relevant idempotents to “fibbinary polynomials” (see Definition 3.3 ahead), as well as some new properties of these polynomials. In particular, we specify the splitting fields of the fibbinary polynomials (Proposition 3.8), which seems interesting for its own sake.

The proof proceeds along the following steps:

²Throughout, “distance” and “weight” stand for Hamming distance and Hamming weight, respectively.

³It is also stated in [DDZ15] that [AS94] may be useful for this purpose, but eventually we have used a different approach.

- Identify a polynomial $g(X) \in \mathbb{F}_2[X]$ whose number of roots in $\mathbb{F}_{2^m}^*$ equals the weight of the generating idempotent. Hence, the number of roots is at least d_B .
- Show that g factors as $g = Xg_1g_2$, where $\deg(g_1) = d_B$ and g_2 is a power of a fibbinary polynomial that splits in $\mathbb{F}_{2^{m-1}}$.
- As $\mathbb{F}_{2^{m-1}}^* \cap \mathbb{F}_{2^m}^* = \mathbb{F}_2^*$, g_2 may contribute at most the root 1 on top of the roots of g_1 in $\mathbb{F}_{2^m}^*$. However, we show that when $g_2(1) = 0$, also $g_1(1) = 0$, so that g_2 cannot contribute any additional root in $\mathbb{F}_{2^m}^*$. Hence, the number of roots of g in $\mathbb{F}_{2^m}^*$ is at most $\deg(g_1) = d_B$.
- So, the weight of the generating idempotent must equal d_B .

1.2 Paper outline

Section 2 includes some required preliminaries. The proof of Theorem 1.2 then appears in Section 3 following the above steps: In Section 3.1, it is shown that the weight of the generating idempotent is equal to the number of roots of a “cyclic fibbinary polynomial” (Definition 3.3) in $\mathbb{F}_{2^m}^*$. Then, in Section 3.2, it is shown that the cyclic fibbinary polynomial factors as the product Xg_1g_2 described above, and in Section 3.3, the splitting field of g_2 is identified. Finally, all the intermediate results are wrapped up in Section 3.4 to complete the proof. We conclude the paper in Section 4 with some concluding remarks.

2 Preliminaries

This section includes some notation and definitions that will be used throughout the paper.

For a positive integer m and for an integer $i \in \{0, 1, \dots, 2^m - 1\}$, the m -bit **binary representation** of i is the vector $(i_{m-1}, i_{m-2}, \dots, i_0) \in \{0, 1\}^m$ such that $i = \sum_{j=0}^{m-1} i_j 2^j$. To simplify notation, we write the binary representation in a string form $i_{m-1}i_{m-2} \cdots i_0$.

2.1 Cyclic codes

Let us first recall some basic definitions related to cyclic codes. For more details, see, e.g., [MS78, Chs. 7,8] or [Rot06, Ch. 8]. A cyclic code over \mathbb{F}_q (for q a prime power) is an \mathbb{F}_q -linear code that is invariant under cyclic shifts. Equivalently, a cyclic code of length n is an ideal in $\mathbb{F}_q[X]/(X^n - 1)$. By correspondence of ideals of $\mathbb{F}_q[X]$ and those of $\mathbb{F}_q[X]/(X^n - 1)$, any ideal of the latter ring (that is, any cyclic code) is the image of an ideal of the former ring that includes $(X^n - 1)$, that is, the image of $(g(X))$ for some $g(X)$ with $g(X)|(X^n - 1)$. The **generator polynomial** of a cyclic code $C \subseteq \mathbb{F}_q[X]/(X^n - 1)$ is the unique monic $g(X)$ whose image in $\mathbb{F}_q[X]/(X^n - 1)$ generates C as an ideal.

We will assume that n is coprime to q , so that $X^n - 1$ is separable. The **zeros** of a cyclic code $C \subseteq \mathbb{F}_q[X]/(X^n - 1)$ are the roots of its generator polynomial, which all lie in the group of n -th roots of unity in the splitting field \mathbb{F}_{q^m} of $X^n - 1$. The set of zeros defines g , and hence determines the code uniquely. We note also that as $g \in \mathbb{F}_q[X]$, its set of roots (i.e., the zeros of the generated cyclic code) must be invariant under the Galois group of $\mathbb{F}_{q^m}/\mathbb{F}_q$. Hence if β is a root, so are all its conjugates $\beta, \beta^q, \beta^{q^2}, \dots$

From this point on, we will only consider the binary case, where $q = 2$, and, furthermore, $n := 2^m - 1$ for some positive integer m . Hence the zeros of the relevant codes are all in $\mathbb{F}_{2^m}^*$. As already mentioned in the introduction, we are concerned with primitive, narrow-sense, binary BCH codes of length $2^m - 1$, which are defined as the binary cyclic codes whose zero set is comprised of $\alpha, \alpha^2, \dots, \alpha^{d-1}$ and their conjugates, where $\alpha \in \mathbb{F}_{2^m}^*$ is a primitive element, and d is an odd integer ≥ 3 .

2.2 Cyclotomic cosets

The orbit of an element $\beta \in \mathbb{F}_{2^m}^*$ under the action of the Galois group of $\mathbb{F}_{2^m}/\mathbb{F}_2$ is $\{\beta, \beta^2, \dots, \beta^{2^{m'-1}}\}$, where $m'|m$ is the degree of the minimal polynomial of β over \mathbb{F}_2 (equivalently, the extension degree $[\mathbb{F}_2(\beta) : \mathbb{F}_2]$). Fixing a primitive element α and writing $\beta = \alpha^i$ for some $i \in \{0, \dots, n-1\}$, a convenient way to represent this orbit is to record the “base- α logarithms” $\{i, 2 \cdot i \bmod n, 2^2 \cdot i \bmod n, \dots, 2^{m'-1} \cdot i \bmod n\}$, where $2^{m'} \cdot i \bmod n = i$, and m' is the smallest positive integer with this property. Such a set is called the **cyclotomic coset** of i modulo n .

2.3 Fourier transforms on $\mathbb{F}_{2^m}^*$

Fix $m \in \mathbb{N}^*$ and a primitive element $\alpha \in \mathbb{F}_{2^m}^*$, and recall that $n = 2^m - 1$. For a polynomial $f \in \mathbb{F}_{2^m}[X]$ with $\deg(f) \leq n-1$, we let the **Fourier transform** of f be the polynomial⁴

$$\hat{f} := \sum_{i=0}^{n-1} f(\alpha^i) X^i.$$

It is well-known that f is determined from \hat{f} as

$$f = \sum_{i=0}^{n-1} \hat{f}(\alpha^{-i}) X^i \quad (1)$$

(this can be shown, e.g., by verifying that the inverse of the Vandermonde matrix $\{\alpha^{ij}\}_{0 \leq i, j \leq n-1}$ is $\{\alpha^{-ij}\}_{0 \leq i, j \leq n-1}$). In fact, the Fourier transform, as well as the inversion formula (1), are well-defined on $\mathbb{F}_{2^m}[X]/(X^n + 1)$, since $\alpha^n = 1$.

2.4 The generating idempotent of a binary cyclic code

We continue to restrict attention to the case $n = 2^m - 1$ of interest to the paper. It is well known [MS78, Ch. 8] that any binary cyclic code $C \subseteq \mathbb{F}_2[X]/(X^n + 1)$ has a unique codeword⁵ $e(X)$ such that $e^2 = e$ in $\mathbb{F}_2[X]/(X^n + 1)$ and also e generates C as an ideal in $\mathbb{F}_2[X]/(X^n + 1)$. Polynomials $e(X)$ with $e^2 = e$ in $\mathbb{F}_2[X]/(X^n + 1)$ are called **idempotents**, and an idempotent e that generates a binary cyclic code C , as above, is called the **generating idempotent** of C .

It is easily verified that any idempotent evaluates to either 0 or 1 on $\mathbb{F}_{2^m}^*$, and that the generating idempotent of a cyclic code C evaluates to 1 exactly on the set of non-zeros

⁴More precisely, the Fourier transform is the mapping $f \mapsto \hat{f}$.

⁵For simplicity, we identify polynomials of degree $< n$ with their image in $\mathbb{F}_2[X]/(X^n + 1)$.

of C in $\mathbb{F}_{2^m}^*$. Hence, if we fix a primitive element $\alpha \in \mathbb{F}_{2^m}$ and let $N \subseteq \{0, \dots, n-1\}$ be such that the set of non-zeros of C is $\{\alpha^i | i \in N\}$, then the Fourier transform \hat{e} of the generating idempotent e of C is given by $\hat{e} = \sum_{i \in N} X^i$.

3 Proof of Theorem 1.2

Throughout this section, m is a positive integer, $n := 2^m - 1$, and α is a primitive element of \mathbb{F}_{2^m} .

3.1 Finding a polynomial whose number of roots equals the weight of the idempotent

By definition, the generating idempotent of a cyclic code takes the value 0 for any zero of the code, and the value 1 for any non-zero of the code. In what follows, for a polynomial $a(X) \in \mathbb{F}_{2^m}[X]$ of degree up to $n-1$, we write $a(X) = \sum_{i=0}^{n-1} a_i X^i$, and let $\mathbf{a} := (a_0, \dots, a_{n-1})$.

To continue, it will be useful to define the following sets of integers. We note that the integers defined in the first part of the definition are the so-called *fibbinary integers* [Slo].

Definition 3.1. 1. Let F_m be the set of integers $i \in \{0, \dots, n-1\}$ such that the binary representation of i has no two consecutive ones.

2. Let $F_m^{(\text{cyc})}$ be the set of integers $i \in \{1, \dots, n-1\}$ such that the m -bit binary representation of i has no two **cyclically** consecutive ones.

For example, $F_3 = \{0, 1, 2, 4, 5\}$, while $F_3^{(\text{cyc})} = \{1, 2, 4\}$.

Observation 3.2. *It holds that*

$$\max F_m = \begin{cases} \frac{2}{3}(2^m + 1) - 1 & \text{if } m \text{ is odd} \\ \frac{2}{3}(2^m - 1) & \text{if } m \text{ is even} \end{cases} \quad (2)$$

and

$$\max F_m^{(\text{cyc})} = \begin{cases} \frac{2}{3}(2^m + 1) - 2 & \text{if } m \text{ is odd} \\ \frac{2}{3}(2^m - 1) & \text{if } m \text{ is even.} \end{cases} \quad (3)$$

Proof. When m is odd, the m -bit binary representation of the largest integer in F_m is $1010 \cdots 101$, while that of the largest integer in $F_m^{(\text{cyc})}$ is $1010 \cdots 100$. When m is even, the m -bit binary representation of the largest integer in both F_m and $F_m^{(\text{cyc})}$ is $1010 \cdots 10$. \square

It will also be useful to define the following polynomials related to F_m and $F_m^{(\text{cyc})}$.

Definition 3.3. Let $f_m := \sum_{i \in F_m} X^i \in \mathbb{F}_2[X]$ and $f_m^{(\text{cyc})}(X) := \sum_{i \in F_m^{(\text{cyc})}} X^i \in \mathbb{F}_2[X]$. We refer to f_m as the m -th **fibbinary polynomial**, and to $f_m^{(\text{cyc})}$ as the m -th **cyclic fibbinary polynomial**.

Note that $\deg(f_m) = \max F_m$ and $\deg(f_m^{(\text{cyc})}) = \max F_m^{(\text{cyc})}$ are given by (2) and (3), respectively.

We will later explore the relations between the fibbinary polynomials and the cyclic fibbinary polynomials, as well as useful properties of the fibbinary polynomials. In particular, we will pin down the splitting field of the f_m . For now, we only relate the cyclic fibbinary polynomials to the idempotent of $\text{BCH}(m, 2^{m-2} + 1)$.

From this point on, “a BCH code” means a primitive, narrow-sense, binary BCH code.

Proposition 3.4. *Let $e(X)$ be the generating idempotent of $\text{BCH}(m, 2^{m-2} + 1)$. Then the weight of e equals the number of roots of $f_m^{(\text{cyc})}(X)$ in $\mathbb{F}_{2^m}^*$.*

Proof. It is well-known that for a BCH code of designed distance δ (for odd δ), all representatives of the conjugacy classes of the zeros of the code are contained in $\alpha, \alpha^2, \dots, \alpha^{\delta-2}$ (as $\alpha^{\delta-1} = (\alpha^{(\delta-1)/2})^2$). So, the zeros of $\text{BCH}(m, 2^{m-2} + 1)$ are α^i , where i runs on the cyclotomic cosets of integers ≥ 1 that are at most $2^{m-2} - 1$.

Note that the m -bit binary representation of $2^{m-2} - 1$ is $0011 \dots 1$. This implies that the set of zeros of $\text{BCH}(m, 2^{m-2} + 1)$ is exactly $\{\alpha^i \mid i \in J_m\}$, where J_m is the set of integers j in $\{1, \dots, n-1\}$ such that the binary representation of j has two cyclically consecutive zeros.⁶ In other words, the set of **non**-zeros of the code consists of α^i for i in $\{0\} \cup \bar{J}_m$, where \bar{J}_m consists of the integers j in $\{1, \dots, n-1\}$ whose binary representation does not have two cyclically consecutive zeros. Hence

$$\hat{e}(X) = 1 + \sum_{i \in \bar{J}_m} X^i.$$

The second summand is invariant under squaring modulo $X^n + 1$ (as \bar{J}_m is clearly the union of complete cyclotomic cosets), and therefore evaluates to either 0 or 1 on \mathbb{F}_{2^m} .

By (1), the weight of e is the number of non-zeros of $\hat{e}(X)$ in $\mathbb{F}_{2^m}^*$, which, by the above comment, is the number of zeros of $v(X) := \sum_{i \in \bar{J}_m} X^i$ in $\mathbb{F}_{2^m}^*$. This number does not change if we replace the exponent set \bar{J}_m by $\{2^m - 1 - i \mid i \in \bar{J}_m\}$, as this change has the effect of getting $v(\beta^{-1})$ when substituting $\beta \in \mathbb{F}_{2^m}^*$. But moving from i to $2^m - 1 - i$ just flips $0 \leftrightarrow 1$ in the m -bit binary representation, and therefore the last set is exactly $F_m^{(\text{cyc})}$, completing the proof. \square

3.2 Recursions for $f_m(X)$, $f_m^{(\text{cyc})}(X)$, and a factorization of $f_m^{(\text{cyc})}(X)$

By Proposition 3.4, it is of interest to find the number of roots of $f_m^{(\text{cyc})}$ in $\mathbb{F}_{2^m}^*$. Toward this end, we will study in this section some recurrence relations for the $f_m(X)$, $f_m^{(\text{cyc})}(X)$, as well as their consequence for the required number of roots.

In what follows, we will use freely the relation $g(X^{2^i}) = g(X)^{2^i}$ for $g(X) \in \mathbb{F}_2[X]$ and $i \in \mathbb{N}$.

Proposition 3.5. *For $m \geq 4$, it holds that*

$$f_m^{(\text{cyc})}(X) = X f_{m-3}(X^4) + 1 + f_{m-1}(X^2). \quad (4)$$

⁶If there are two cyclically consecutive zeros, they can be made the two MSBs by a cyclic shift, resulting in a number $\leq 2^{m-2} - 1$. On the other hand, if there are no two cyclically consecutive zeros, then for any cyclic shift, at least one of the two MSBs is 1, so that the number is $> 2^{m-2} - 1$.

Proof. Write

$$F_m^{(\text{cyc})} = A_m \cup B_m, \quad (5)$$

where A_m, B_m are the subsets consisting of integers whose binary representation has LSB 0, 1, respectively. Then we claim that

$$A_m = \{2j | j \in F_{m-1}\} \setminus \{0\} \quad (6)$$

and

$$B_m = \{4j + 1 | j \in F_{m-3}\}. \quad (7)$$

To see this, note that the binary representation of the numbers in B_m must have 2nd LSB and MSB of 0 (because the LSB is 1), and then the part of the binary representation obtained by omitting the two LSBs and the MSB can be chosen freely from F_{m-3} . A similar argument justifies also (6). Now the assertion follows at once from (5)–(7). \square

Let us now turn to some recursions for the f_m .

Proposition 3.6. *It holds that $f_1(X) = 1 + X$, $f_2(X) = 1 + X + X^2$. Also, for $m \geq 3$, it holds that*

$$f_m = f_{m-1}(X^2) + X f_{m-2}(X^4), \quad (8)$$

alternatively,

$$f_m = f_{m-1}(X) + X^{2^{m-1}} f_{m-2}(X). \quad (9)$$

In addition,

$$1 + f_m = X f_{m-1} f_{m-2}^2. \quad (10)$$

Finally, let $N_m := |F_m|$ be the the number of monomials appearing in f_m . Then $N_1 = 2$, $N_2 = 3$, and for $m \geq 3$, $N_m = N_{m-1} + N_{m-2}$. Hence

$$f_m(1) = 0 \quad \iff \quad 3 | (m - 1). \quad (11)$$

Proof. First, (8) follows from decomposing F_m as the union of the subset of integers having LSB 0, and the subset having LSB 1, and hence two LSBs 01 (alternatively, it follows from the power series decomposition in [Arn10, p. 756]). Also, (9) follows similarly by considering MSBs instead of LSBs.

To prove (10), we use induction. For $m = 3$, the equation can be verified directly. Assume, therefore, that (10) holds for some $m \geq 3$. Using (8), we get

$$\begin{aligned} 1 + f_{m+1} &= 1 + f_m^2 + X f_{m-1}^4 \\ &= X^2 f_{m-1}^2 f_{m-2}^4 + X f_{m-1}^4 \quad (\text{induction hypothesis}) \\ &= X f_{m-1}^2 (f_{m-1}^2 + X f_{m-2}^4) \\ &= X f_m f_{m-1}^2, \end{aligned}$$

as required.

Finally, the last assertion follows from either of (8) or (9). \square

Remark. The above Fibonacci recursion for the N_m is a well known property of the sets F_m of fibbinary integers; it is, in fact, the reason for the name [Slo].

Corollary 3.7. For $m \geq 4$, it holds that $f_m^{(\text{cyc})} = Xu_m f_{m-3}^4$ for some polynomial $u_m(X) \in \mathbb{F}_2[X]$ with

$$\deg(u_m) = \begin{cases} \frac{1}{3}(2^m + 1) & \text{if } m \text{ is odd} \\ \frac{1}{3}(2^m - 1) & \text{if } m \text{ is even.} \end{cases} \quad (12)$$

Hence, for both even and odd m , $\deg(u_m) = d_B$, where d_B is the Bose distance found in [DDZ15]. Moreover, if $f_{m-3}(1) = 0$, then $u_m(1) = 0$.

Proof. By (4), it holds that $f_m^{(\text{cyc})} = Xf_{m-3}^4 + (1 + f_{m-1})^2$, while by (10), $1 + f_{m-1} = Xf_{m-2}f_{m-3}^2$. It follows that $f_m^{(\text{cyc})}$ is divisible by Xf_{m-3}^4 . Now (12) follows from subsisting the degrees of $f_m^{(\text{cyc})}$ and f_{m-3} from (2) and (3).

For the last assertion, by substituting (10) in (4) as mentioned above, it can be verified that $u_m = 1 + Xf_{m-2}^2$, so that $u_m(1) = 0$ iff $f_{m-2}(1) = 1$. But by (11), this must hold if $f_{m-3}(1) = 0$. \square

3.3 The splitting field of the fibbinary polynomials

Proposition 3.8. For all integer $m \geq 3$, f_m divides $X^{2^{m+2}-1} + 1$. Specifically, it holds that

$$X^{2^{m+2}-1} + 1 = (1 + X)f_m^2 + Xf_{m-1}^2f_m + X^3f_{m-1}^4f_m^4. \quad (13)$$

Hence for all $m \geq 3$, the splitting field of f_m is $\mathbb{F}_{2^{m+2}}$.

Proof. We will prove (13) by induction. For the basis, one can verify (13) directly.

To continue, let us write t_m for the right-hand side of (13). Assume now that $m \geq 4$, and (13) holds for $m - 1$, that is, $t_{m-1} = X^{2^{m+1}-1} + 1$. Since

$$X^{2^{m+2}-1} + 1 = X((X^{2^{m+1}-1} + 1) + 1)^2 + 1,$$

it is sufficient to prove that

$$X(t_{m-1} + 1)^2 + 1 = t_m. \quad (14)$$

Now,

$$\begin{aligned} X(t_{m-1} + 1)^2 + 1 &= X(1 + X)^2 f_{m-1}^4 + \underbrace{X^3 f_{m-2}^4 f_{m-1}^2}_{X(Xf_{m-1}f_{m-2}^2)^2} + \underbrace{X^7 f_{m-2}^8 f_{m-1}^8}_{X^3 f_{m-1}^4 (Xf_{m-1}f_{m-2}^2)^4} + X + 1 \\ &= X(1 + X)^2 f_{m-1}^4 + X(1 + f_m^2) + X^3 f_{m-1}^4 (1 + f_m^4) + X + 1 \quad (\text{by 10}) \\ &= Xf_{m-1}^4 + X^3 f_{m-1}^4 + X + Xf_m^2 + X^3 f_{m-1}^4 + X^3 f_{m-1}^4 f_m^4 + X + 1 \\ &= 1 + Xf_{m-1}^4 + Xf_m^2 + X^3 f_{m-1}^4 f_m^4. \end{aligned}$$

The last expression is equal to t_m iff

$$1 + Xf_{m-1}^4 + Xf_m^2 = (1 + X)f_m^2 + Xf_{m-1}^2 f_m,$$

that is, iff $f_m^2 = 1 + Xf_{m-1}^2(f_m + f_{m-1}^2)$. However, by (8), the right-hand side of the last expression equals $1 + X^2 f_{m-1}^2 f_{m-2}^4$, which, in turn, equals f_m^2 , by (10). This completes the proof of (13).

Hence the splitting field of f_m is a subfield of $\mathbb{F}_{2^{m+2}}$, and f_m is separable. Considering the degree of f_m from (2), the only possible subfield of $\mathbb{F}_{2^{m+2}}$ that can contain all roots is $\mathbb{F}_{2^{m+2}}$ itself. \square

3.4 Completing the proof of Theorem 1.2

Proof of Theorem 1.2. By Proposition 3.4, the weight w of the generating idempotent is equal to the number of roots of $f_m^{(\text{cyc})}$ in $\mathbb{F}_{2^m}^*$. By Corollary 3.7, $f_m^{(\text{cyc})} = Xu_m f_{m-3}^4$, and by Proposition 3.8, all the roots of f_{m-3} are in $\mathbb{F}_{2^{m-1}}$, whose intersection with \mathbb{F}_{2^m} is \mathbb{F}_2 . So, the only root possibly contributed by f_{m-3} in $\mathbb{F}_{2^m}^*$ is 1, but by Corollary 3.7, if 1 is a root of f_{m-3} , then it is already a root of u_m .

Hence the total number of roots is at most $\deg(u_m) = d_B$, but also the number of roots is at least d_B as it is the weight of the idempotent. \square

4 Concluding remarks

It follows from the proof of Theorem 1.2 that the polynomial $u_m(X)$ from the proof of Corollary 3.7 splits in \mathbb{F}_{2^m} , and its roots are α^i exactly for those i for which X^i appears in $e(X)$. Hence $u_m(X)$ is nothing but the *locator polynomial* of the generating idempotent. Note, however, that in most references, including [AS94], the term “locator polynomial” refers to the “reversed” polynomial $X^{\deg(u_m)}u_m(X^{-1})$, having $1/\alpha^i$ for i as above as its roots.

From the proof of Corollary 3.7 it follows that $u_m = 1 + Xf_{m-2}^2$, so that u_m has only odd powers of X , apart from the free coefficient. Considering that u_m is reversed with respect to the definition of the locator polynomial in [AS94], this agrees with [AS94, Theorem 2]. However, without the above analysis, it is not automatically clear that $u_m = 1 + Xf_{m-2}^2$ splits in \mathbb{F}_{2^m} , and that it is the locator of the *generating* idempotent. It would be interesting to see if the methods of [AS94] can provide an alternative proof for Theorem 1.2.

Finally, we remark that the above analysis also recovers the dimension of $\text{BCH}(m, 2^{m-2} + 1)$ from [DDZ15, Corollary 10]. The dimension is the number of non-zeros of $e(X)$, which equals $1 + |\bar{J}_m| = 1 + |F_m^{(\text{cyc})}|$, as shown in the proof of Proposition 3.4. From Proposition 3.5, it can be verified that $1 + |F_m^{(\text{cyc})}| = |F_{m-3}| + |F_{m-1}|$, and as $|F_j|$ is the $(j + 2)$ -th Fibonacci number (Proposition 3.6), Binet’s formula for the Fibonacci numbers can be used to recover [DDZ15, Corollary 10]; we omit the details.

Acknowledgment

The authors are grateful to Ariel Doubchak for many helpful comments on an earlier draft.

References

- [Arn10] Jörg Arndt. *Matters Computational - Ideas, Algorithms, Source Code [The fxtbook]*. www.jjj.de, 2010.
- [AS94] D. Augot and N. Sendrier. Idempotents and the BCH bound. *IEEE Trans. Inf. Theory*, 40(1):204–207, 1994.

- [BR60] R. C. Bose and Dwijendra K. Ray-Chaudhuri. On a class of error correcting binary group codes. *Inf. Control.*, 3(1):68–79, 1960.
- [DDZ15] Cunsheng Ding, Xiaoni Du, and Zhengchun Zhou. The Bose and minimum distance of a class of BCH codes. *IEEE Trans. Inf. Theory*, 61(5):2351–2356, 2015.
- [DL24] Cunsheng Ding and Chengju Li. BCH cyclic codes. *Discret. Math.*, 347(5), 2024.
- [Hoc59] A. Hocquenghem. Codes correcteurs d’erreurs. *Chiffres*, 2:147–156, 1959.
- [MS78] F.J. MacWilliams and N.J.A. Sloane. *The Theory of Error-Correcting Codes*. North-holland Publishing Company, 2nd edition, 1978.
- [Rot06] Ron Roth. *Introduction to Coding Theory*. Cambridge University Press, USA, 2006.
- [Slo] N. J. A. Sloane. Sequence A003714 (fibbinary numbers). *The On-Line Encyclopedia of Integer Sequences, OEIS*.