

Random Walk Integrals

Jonathan M. Borwein¹ and Dirk Nuyens^{2†} and Armin Straub³ and James Wan⁴

¹University of Newcastle, Australia. Email: jonathan.borwein@newcastle.edu.au

²K.U.Leuven, Belgium. Email: dirk.nuyens@cs.kuleuven.be

³Tulane University, New Orleans, USA. Email: astraub@tulane.edu

⁴University of Newcastle, Australia. Email: james.wan@newcastle.edu.au

Abstract. We study the expected distance of a two-dimensional walk in the plane with unit steps in random directions. A series evaluation and recursions are obtained making it possible to explicitly formulate this distance for small number of steps. Formulae for all the moments of a 2-step and a 3-step walk are given, and an expression is conjectured for the 4-step walk. The paper makes use of the combinatorial features exhibited by the even moments which, for instance, lead to analytic continuations of the underlying integral.

Résumé.

Resumen.

Keywords: some well classifying words, **mandatory!**

1 Introduction and Preliminaries

This is an extended abstract of (BNSW09) which contains the exposition given here complemented with much more details. In particular, we often refer to (BNSW09) for full proofs of statements that we present.

Throughout, we consider the n -dimensional integral

$$W_n(s) := \int_{[0,1]^n} \left| \sum_{k=1}^n e^{2\pi x_k i} \right|^s d\mathbf{x} \quad (1)$$

which occurs in the theory of uniform random walk integrals in the plane, where at each step a unit-step is taken in a random direction, see Figure 1. As such, the integral (1) expresses the s th moment of the expected distance to the origin after n steps. Particularly interesting is the special case of the expected distance $W_n(1)$ after n steps.

A lot is known about the one-dimensional random walk. E.g., its expected distance after n unit-steps is $(n-1)!!/(n-2)!!$ when n is even and $n!!/(n-1)!!$ when n is odd (and asymptotically this distance is

[†]part of this work done while a research associate at the Department of Mathematics and Statistics, University of New South Wales, Australia

$\sqrt{2n/\pi}$. For the two-dimensional walk no such explicit expressions were known, although the term random walk first appears in a (related) question by Karl Pearson in *Nature* in 1905 (Pea1905) for explicitly this two-dimensional walk under consideration. Pearson triggered answers by Lord Rayleigh (Ray1905) on the asymptotic behaviour of the probability for n very large and by Benett (referred to in (Pea1905b)) for the case $n = 2$, after which he concluded that there still was a large interest for the unresolved case of small n which is dramatically different from the case of large n . Note that the expected value for the root-mean-square distance is well known to be just \sqrt{n} (in that case the implicit square root in (1) disappears which greatly simplifies the problem).




Fig. 1: Random walks in the plane.

We picked up the special case $s = 1$ of (1) from the whiteboard in the common room at UNSW where it was written as a generalization of a discrete problem in a cryptographic context by Peter Donovan, discussed in (Don09). However, the problem in itself appears in numerous applications, e.g., in problems involving Brownian motion in physics. Numerical values of W_n evaluated at integers can be seen in Tables 1 and 2. One immediately notices the apparent integer sequences for the even moments—which are the moments of the squared expected distance (thus the square root for $s = 2$ gives the root-mean-square distance \sqrt{n}). By experimentation and some sketchy arguments we quickly conjectured and believed that, for k a nonnegative integer,

$$W_3(k) = \operatorname{Re} {}_3F_2 \left(\begin{matrix} \frac{1}{2}, -\frac{k}{2}, -\frac{k}{2} \\ 1, 1 \end{matrix} \middle| 4 \right). \quad (2)$$

(In fact, (2) also holds for negative odd integers.) This was for long a mystery, but it will be proven in the final section of the paper.

In Section 2 we develop an infinite series expression for $W_n(s)$ which holds for all real $s > 0$, see Theorem 2.1. From this it then follows in Corollary 2.2 that the even moments of $W_n(s)$ are given by integer sequences. The combinatorial features of $f_n(k) := W_n(2k)$, k a nonnegative integer, are studied in Section 3. We show that there is a recurrence relation for the numbers $f_n(k)$ and confirm that indeed, an observation from Table 1, the last digit in the column for $s = 10$ is always $n \bmod 10$.

n	$s = 2$	$s = 4$	$s = 6$	$s = 8$	$s = 10$	(Slo09)
2	2	6	20	70	252	A000984
3	3	15	93	639	4653	A002893
4	4	28	256	2716	31504	A002895
5	5	45	545	7885	127905	
6	6	66	996	18306	384156	

Tab. 1: $W_n(s)$ at even integers.

n	$s = 1$	$s = 3$	$s = 5$	$s = 7$	$s = 9$
2	1.27324	3.39531	10.86498	37.25135	132.44925
3	1.57460	6.45168	36.70519	241.54389	1714.61960
4	1.79909	10.12068	82.65146	822.27337	9169.61964
5	2.00816	14.28958	152.31577	2037.14185	31392.80994
6	2.19386	18.91327	248.75895	4186.18720	82718.49864

Tab. 2: $W_n(s)$ at odd integers.

In Section 4 some analytic and numerical results for $n = 1, 2, 3$ are given and we lift the recursion for $f_n(k)$ to $W_n(s)$ by the use of Carlson’s theorem. The recursions for $n = 2, 3, 4$ are given explicitly as an example. These recursions then give further information on the poles of the analytic continuations of W_n (graphs of W_n for $n = 3, 4, 5, 6$ and their analytic continuations are shown in Figure 2). From here we conjecture the recursion

$$W_{2n}(s) \stackrel{?[1]}{=} \sum_{j \geq 0} \binom{s/2}{j}^2 W_{2n-1}(s - 2j),$$

based on analytic continuations, and the explicit form, related to (2),

$$W_4(k) \stackrel{?[2]}{=} \operatorname{Re} \sum_{j \geq 0} \binom{s/2}{j}^2 {}_3F_2 \left(\begin{matrix} \frac{1}{2}, -\frac{k}{2} + j, -\frac{k}{2} + j \\ 1, 1 \end{matrix} \middle| 4 \right)$$

for k a positive integer. High precision numerical evaluations for $n = 3$ and $n = 4$ are given.

In the final section we explore the underlying probability model more closely, starting with another answer to Pearson, this time by Kluver (Klu1906). Finally, considering conditional densities, we are able to give an alternative form for $W_3(s)$ which eventually leads to a proof of (2).

2 A Series Evaluation of $W_n(s)$

Theorem 2.1 For real $s > 0$,

$$W_n(s) = n^s \sum_{m \geq 0} (-1)^m \binom{s/2}{m} \sum_{k=0}^m (-1)^k \binom{m}{k} \left\{ n^{-2k} \sum_{a_1 + \dots + a_n = k} \binom{k}{a_1, \dots, a_n}^2 \right\}. \quad (3)$$

Proof:




Fig. 2: Various W_n and their analytic continuations.

We first exploit the binomial theorem to show that for integer $n > 1$ and real $s \geq 0$,

$$W_n(s) = n^s \sum_{m \geq 0} (-1)^m \binom{s/2}{m} n^{-2m} \int_{[0,1]^n} \left(4 \sum_{1 \leq i < j \leq n} \sin^2(\pi(x_j - x_i)) \right)^m d\mathbf{x}. \quad (4)$$

Next we evaluate the trigonometric integral in (4). To this end, we show that it is the constant term of

$$(n^2 - (x_1 + \cdots + x_n)(1/x_1 + \cdots + 1/x_n))^m.$$

The details appear in (BNSW09). Alternatively, one may start with the observation that $W_n(s)$ is the constant term of

$$((x_1 + \cdots + x_n)(1/x_1 + \cdots + 1/x_n))^{s/2} \quad (5)$$

which follows directly from the integral definition. \square

From Theorem 2.1 and the fact that the binomial transform is an involution we additionally learn that the even moments are integer sequences as detailed by the following corollary.

Corollary 2.2 For nonnegative integers k ,

$$W_n(2k) = \sum_{a_1 + \cdots + a_n = k} \binom{k}{a_1, \dots, a_n}^2. \quad (6)$$

An outline of the genesis of these evaluations is also given in (BNSW09).

3 Further Combinatorial Features

In light of Corollary 2.2, we consider the combinatorial sums $f_n(k) := W_n(2k)$ of multinomial coefficients squared. These numbers also appear in (RS09) in the following way: $f_n(k)$ counts the number of *abelian squares* of length $2k$ over an alphabet with n letters (that is strings xx' of length $2k$ from an alphabet with n letters such that x' is a permutation of x). It is not hard to see that, (Bar64),

$$f_{n_1+n_2}(k) = \sum_{j=0}^k \binom{k}{j}^2 f_{n_1}(j) f_{n_2}(k-j), \quad (7)$$

for two non-overlapping alphabets with n_1 and n_2 letters. In particular, we may use (7) to obtain $f_1(k) = 1$, $f_2(k) = \binom{2k}{k}$, as well as

$$f_3(k) = \sum_{j=0}^k \binom{k}{j}^2 \binom{2j}{j} = {}_3F_2 \left(\begin{matrix} \frac{1}{2}, -k, -k \\ 1, 1 \end{matrix} \middle| 4 \right) = \binom{2k}{k} {}_3F_2 \left(\begin{matrix} -k, -k, -k \\ 1, -k + \frac{1}{2} \end{matrix} \middle| \frac{1}{4} \right), \quad (8)$$

$$f_4(k) = \sum_{j=0}^k \binom{k}{j}^2 \binom{2j}{j} \binom{2(k-j)}{k-j} = \binom{2k}{k} {}_4F_3 \left(\begin{matrix} \frac{1}{2}, -k, -k, -k \\ 1, 1, -k + \frac{1}{2} \end{matrix} \middle| 1 \right). \quad (9)$$

Here and below ${}_pF_q$ denotes the hypergeometric function.

The following result is established in (Bar64) with the recursions for $n \leq 6$ given explicitly.

Theorem 3.1 *For fixed $n \geq 2$, the sequence $f_n(k)$ satisfies a recurrence of order $\lambda := \lceil n/2 \rceil$ with polynomial coefficients of degree $n-1$:*

$$c_{n,0}(k)f_n(k) + \dots + c_{n,\lambda}(k)f_n(k+\lambda) = 0.$$

Remark 3.2 For fixed k , the map $n \mapsto f_n(k)$ is a polynomial of degree k . This follows from

$$f_n(k) = \sum_{j=0}^k \binom{n}{j} \sum_{\substack{a_1+\dots+a_j=k \\ a_i>0}} \binom{k}{a_1, \dots, a_j}^2, \quad (10)$$

because the right-hand side is a linear combination (with positive coefficients only depending on k) of the polynomials $\binom{n}{0}, \binom{n}{1}, \dots, \binom{n}{k}$ of respective degrees $0, 1, \dots, k$. From (10) the coefficient of $\binom{n}{k}$ is seen to be $(k!)^2$. We therefore obtain the first-order approximation $W_n(s) \approx_n n^{s/2} \Gamma(s/2 + 1)$ for n approaching infinity, see also (Klu1906). In particular, $W_n(1) \approx_n \sqrt{n\pi}/2$. Similarly, the coefficient of $\binom{n}{k-1}$ is $\frac{k-1}{4}(k!)^2$ which gives rise to the second-order approximation

$$(k!)^2 \binom{n}{k} + \frac{k-1}{4} (k!)^2 \binom{n}{k-1} = k!n^k - \frac{k(k-1)}{4} k!n^{k-1} + O(n^{k-2}).$$

of $f_n(k)$. We therefore obtain

$$W_n(s) \approx_n n^{s/2-1} \left\{ \left(n - \frac{1}{2} \right) \Gamma \left(\frac{s}{2} + 1 \right) + \Gamma \left(\frac{s}{2} + 2 \right) - \frac{1}{4} \Gamma \left(\frac{s}{2} + 3 \right) \right\},$$

which is exact for $s = 0, 2, 4$. In particular, $W_n(1) \approx_n \sqrt{n\pi}/2 + \sqrt{\pi/n}/32$. More general approximations are given in (Cra09). \diamond

Remark 3.3 It follows straight from (6) that, for primes p , $f_n(p) \equiv n$ modulo p . Further, for $k \geq 1$, $f_n(k) \equiv n$ modulo 2. This may be derived inductively from the recurrence (7) since, assuming that $f_n(k) \equiv n$ modulo 2 for some n and all $k \geq 1$,

$$f_{n+1}(k) = \sum_{j=0}^k \binom{k}{j}^2 f_n(j) \equiv k + \sum_{j=1}^k \binom{k}{j} j \equiv k + \sum_{\substack{j=1 \\ j \text{ odd}}}^k \binom{k}{j} = k + 2^{k-1} \equiv k \pmod{2}.$$

Hence for odd primes p ,

$$f_n(p) \equiv n \pmod{2p}. \quad (11)$$

The congruence (11) also holds for $p = 2$ since $f_n(2) = (2n - 1)n$, compare (10). \diamond

Remark 3.4 The integers $f_3(k)$ (respectively $f_4(k)$), the first of which are given in Table 1, also arise in physics, see for instance (BBBG08), and are referred to as *hexagonal* (respectively *diamond*) *lattice integers*. The following formulae (BBBG08, (186)–(188)) relate these sequences in non-obvious ways:

$$\begin{aligned} \left(\sum_{k \geq 0} f_3(k) (-x)^k \right)^2 &= \sum_{k \geq 0} f_2(k)^3 \frac{x^{3k}}{((1+x)^3(1+9x))^{k+\frac{1}{2}}} \\ &= \sum_{k \geq 0} f_2(k) f_3(k) \frac{(-x(1+x)(1+9x))^k}{((1-3x)(1+3x))^{2k+1}} \\ &= \sum_{k \geq 0} f_4(k) \frac{x^k}{((1+x)(1+9x))^{k+1}}. \end{aligned}$$

It would be instructive to similarly engage $f_5(k)$. \diamond

4 Analytic and Numerical Results

We start with investigating the analyticity of $W_n(s)$ for a given n . In (BNSW09)[Proposition 1], we show that $W_n(s)$, as defined in (1), is analytic at least for $\operatorname{Re} s > 0$. Furthermore, it is shown (based on the results of Section 4.2) that (1) is indeed finite and analytic for $\operatorname{Re} s > -2$, for each integer $n > 2$ (compare the graphs of the W_n shown in Figure 2).

4.1 $n = 1$, $n = 2$, and $n = 3$

The case $n = 1$ is trivial: it follows straight from the integral definition (1) that $W_1(s) = 1$.

In the case $n = 2$, direct integration of (18) with $n = 2$ yields

$$W_2(s) = 2^{s+1} \int_0^{1/2} \cos(\pi t)^s dt = \binom{s}{s/2}, \quad (12)$$

which may also be obtained using (3).

For $n = 3$, based on (8) we define

$$V_3(s) := {}_3F_2 \left(\begin{matrix} \frac{1}{2}, -\frac{s}{2}, -\frac{s}{2} \\ 1, 1 \end{matrix} \middle| 4 \right), \quad (13)$$

so that by Corollary 2.2 and (8), $W_3(2k) = V_3(2k)$ for nonnegative integers k . This led us to explore $V_3(s)$ more generally numerically and so to conjecture the following which we prove in the penultimate section:

Theorem 4.1 *For nonnegative even integers and all odd integers k :*

$$W_3(k) = \operatorname{Re} V_3(k). \quad (14)$$

From here, we derive the following equivalent expressions for $W_3(1)$:

$$\begin{aligned} W_3(1) &= \frac{4\sqrt{3}}{3} \left({}_3F_2 \left(\begin{matrix} -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2} \\ 1, 1 \end{matrix} \middle| \frac{1}{4} \right) - \frac{1}{\pi} \right) + \frac{\sqrt{3}}{24} {}_3F_2 \left(\begin{matrix} \frac{1}{2}, \frac{1}{2}, \frac{1}{2} \\ 2, 2 \end{matrix} \middle| \frac{1}{4} \right) \\ &= 2\sqrt{3} \frac{K^2(k_3)}{\pi^2} + \sqrt{3} \frac{1}{K^2(k_3)} \\ &= \frac{3}{16} \frac{2^{1/3}}{\pi^4} \Gamma^6 \left(\frac{1}{3} \right) + \frac{27}{4} \frac{2^{2/3}}{\pi^4} \Gamma^6 \left(\frac{2}{3} \right). \end{aligned}$$

These rely on using Legendre's identity and several Clausen-like product formulae, plus Legendre's evaluation of $K(k_3)$ where $k_3 := \frac{\sqrt{3}-1}{2\sqrt{2}}$ is the *third singular value* as in (BB87). Similar expressions can be given for W_3 evaluated at odd integers.

4.2 Carlson's Theorem

We may lift the recursive structure of f_n , defined in Section 3, to W_n to a fair degree on appealing to Carlson's theorem (Tit39, 5.81):

Theorem 4.2 (Carlson) *Let f be analytic in the right half-plane $\operatorname{Re} z \geq 0$ and of exponential type (meaning that $|f(z)| \leq M e^{c|z|}$ for some M and c), with the additional requirement that*

$$|f(z)| \leq M e^{d|z|}$$

for some $d < \pi$ on the imaginary axis $\operatorname{Re} z = 0$. If $f(k) = 0$ for $k = 0, 1, 2, \dots$ then $f(z) = 0$ identically.

By verifying that Carlson's theorem applies, we get:

Theorem 4.3 *Given that $f_n(k)$ satisfies a recurrence*

$$c_{n,0}(k)f_n(k) + \dots + c_{n,\lambda}(k)f_n(k + \lambda) = 0$$

with polynomial coefficients $c_{n,j}(k)$ (see Theorem 3.1) then $W_n(s)$ satisfies the corresponding functional equation

$$c_{n,0}(s/2)W_n(s) + \dots + c_{n,\lambda}(s/2)W_n(s + 2\lambda) = 0,$$

for $\operatorname{Re} s \geq 0$.

Example 4.4 For $n = 2, 3, 4$ we find

$$\begin{aligned}(s+2)W_2(s+2) - 4(s+1)W_2(s) &= 0, \\ (s+4)^2W_3(s+4) - 2(5s^2 + 30s + 46)W_3(s+2) + 9(s+2)^2W_3(s) &= 0, \\ (s+4)^3W_4(s+4) - 4(s+3)(5s^2 + 30s + 48)W_4(s+2) + 64(s+2)^3W_4(s) &= 0.\end{aligned}$$

Note that for all complex s , the function $V_3(s)$ defined in (13) also satisfies the recursion given above for $W_3(s)$ —as is routine to prove symbolically. \diamond

We note that in each case the recursion lets us determine significant information about the nature and position of any poles of W_n . Details appear in (BNSW09). In particular, for $n \geq 3$, the recursion guaranteed by Theorem 4.3 provides an analytic continuation of W_n to all of the complex plane with poles at certain negative integers. Here, we confine ourselves to show the continuations of W_3, W_4, W_5 , and W_6 on the negative real axis in Figure 2. These illustrate the fact that, e.g., W_3 and W_5 have simple poles at $-2, -4, -6, \dots$ whereas W_4 has double poles at these integers. It is further shown in (BNSW09) that, for instance, $\text{Res}_{-2}(W_3) = \frac{2}{\sqrt{3}\pi}$.

Below we use $\stackrel{?[1]}{=}$ and the like to indicate equivalent conjectural equalities. Our next somewhat audacious conjecture is:

Conjecture 4.5 For positive integer s and n one has

$$W_{2n}(s) \stackrel{?[1]}{=} \sum_{j \geq 0} \binom{s/2}{j}^2 W_{2n-1}(s-2j). \quad (15)$$

In (15) we use the recursion/continuation of W_n on the righthand side as given above for $n = 2, 3, 4$. By (7) Conjecture 4.5 clearly holds for s an even positive integer. Further, it follows from (12) that the conjecture holds for $n = 1$.

Recall that the real part of $V_3(k)$ as defined in (13) gives $W_3(k)$ for nonnegative integers k . Define

$$V_4(s) := \sum_{j \geq 0} \binom{s/2}{j}^2 V_3(s-2j) = \sum_{j \geq 0} \binom{s/2}{j}^2 {}_3F_2 \left(\begin{matrix} \frac{1}{2}, -\frac{s}{2} + j, -\frac{s}{2} + j \\ 1, 1 \end{matrix} \middle| 4 \right). \quad (16)$$

This combines with the much better substantiated special case $n = 2$ of Conjecture 4.5 to provide:

Conjecture 4.6 For all integers k ,

$$W_4(k) \stackrel{?[2]}{=} \text{Re } V_4(k). \quad (17)$$

4.3 Numerical Evaluations

Note that the following one-dimensional reduction of the integral may be achieved by taking periodicity into account.

$$W_n(s) = \int_{[0,1]^{n-1}} \left| 1 + \sum_{k=1}^{n-1} e^{2\pi x_k i} \right|^s d(x_1, \dots, x_{n-1}). \quad (18)$$

$n = 3$ Using this reduction, David Bailey (running tanh-sinh integration on a 256-core LBNL system for roughly 15 minutes) has confirmed that the first 175 digits of $W_3(1)$ are given by

$$\begin{aligned} W_3(1) \approx & 1.5745972375518936574946921830765196902216661807585191701936930983 \\ & 018311805944543821310853133622419530649842236115540882056173012611 \\ & 081031331499438143442975115786527521008424458. \end{aligned}$$

This agreed with the evaluation $W_3(1) = \operatorname{Re} V_3(1)$ originally conjectured in (14). He has also confirmed 175 digits for $W_3(s) = \operatorname{Re} V_3(s)$ for $s = 2, \dots, 7$.

$n = 4$ Using Conjecture 4.6 we provide the approximation

$$\begin{aligned} W_4(1) \approx & 1.7990924798428510335326028458461089100662820032916204566266417735 \\ & 988542669321205752411619305734748280560170144445179836872885. \end{aligned}$$

It is worthwhile observing that this level of approximation is made possible by the fact that, roughly, one correct digit is added by each term of the sum.

5 More Probability

As noted such problems have a long lineage. For example, in response to the question posed by Pearson in *Nature*, Kluyver (Klu1906) makes a lovely analysis of the cumulative distribution function of the distance traveled by a “rambler” in the plane for various step lengths. In particular, for our uniform walk Kluyver provides the Bessel representation

$$P_n(t) = t \int_0^\infty J_1(xt) J_0^n(x) dx. \quad (19)$$

Thus, $W_n(s) = \int_0^n t^s p_n(t) dt$, where $p_n = P'_n$. From here, David Broadhurst (Bro09) obtains

$$W_n(s) = 2^{s+1-k} \frac{\Gamma(1 + \frac{s}{2})}{\Gamma(k - \frac{s}{2})} \int_0^\infty x^{2k-s-1} \left(-\frac{1}{x} \frac{d}{dx}\right)^k J_0^n(x) dx \quad (20)$$

for real s with $2k > s > \max(-2, -\frac{n}{2})$. (20) enables Broadhurst (Bro09) to verify Conjecture 4.5 for $n = 2, 3, 4, 5$ and odd $s < 50$ to a precision of 50 digits.

Remark 5.1 For $n = 3, 4$, symbolic integration in *Mathematica* of (20) leads to interesting analytic continuations (Cra09) such as

$$W_3(s) = \frac{1}{2^{2s+1}} \tan\left(\frac{\pi s}{2}\right) \binom{s}{\frac{s-1}{2}}^2 {}_3F_2\left(\begin{matrix} \frac{1}{2}, \frac{1}{2}, \frac{1}{2} \\ \frac{s+1}{2}, \frac{s+3}{2} \end{matrix} \middle| \frac{1}{4}\right) + \binom{s}{\frac{s}{2}} {}_3F_2\left(\begin{matrix} -\frac{s}{2}, -\frac{s}{2}, -\frac{s}{2} \\ 1, -\frac{s-1}{2} \end{matrix} \middle| \frac{1}{4}\right), \quad (21)$$

and

$$W_4(s) = \frac{1}{2^{2s}} \tan\left(\frac{\pi s}{2}\right) \binom{s}{\frac{s-1}{2}}^3 {}_4F_3\left(\begin{matrix} \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{s}{2} + 1 \\ \frac{s+3}{2}, \frac{s+3}{2}, \frac{s+3}{2} \end{matrix} \middle| 1\right) + \binom{s}{\frac{s}{2}} {}_4F_3\left(\begin{matrix} \frac{1}{2}, -\frac{s}{2}, -\frac{s}{2}, -\frac{s}{2} \\ 1, 1, -\frac{s-1}{2} \end{matrix} \middle| 1\right). \quad (22)$$

We note that for $s = 2k = 0, 2, 4, \dots$ the first term in (21) (resp. (22)) is zero and the second is a formula given in (8) (resp. (9)). Thence, one can in principle prove (21) and (22) by applying Carlson’s theorem—after showing the singularities at $1, 3, 5, \dots$ are removable. \diamond

Herein, we will take a related probabilistic approach so as to be able to express our quantities of interest in terms of special functions which allows us to explicitly resolve $W_3(2k+1)$ for all odd values.

It is elementary to express the distance y of an $(n+1)$ -step walk conditioned on a given distance x of an n -step walk. Since, by a simple application of the cosine rule we find

$$y^2 = x^2 + 1 + 2x \cos(\theta), \quad (23)$$

where θ is the outside angle of the triangle with sides $x, 1, y$. It follows, for details see (BNSW09), that the conditional density for the distance y of an $(n+1)$ -step walk as an extension of an n -step walk with distance x is

$$h_x(y) = \frac{2y}{\pi \sqrt{4x^2 - (y^2 - x^2 - 1)^2}} \quad (24)$$

which, of course, is independent of n .

We therefore have the following trivial evaluation

$$W_{n+1}(s) = \mathbb{E}(y^s) = \mathbb{E}(\mathbb{E}(y^s | x)) = \int_0^n \left(\int_{|x-1|}^{x+1} y^s h_x(y) dy \right) p_n(x) dx, \quad (25)$$

under the assumption that the probability density p_n for the n -step walk is known. Clearly, for the 1-step walk we have $p_1(x) = \delta_1(x)$, a Dirac delta at $x = 1$. It then follows immediately that the probability density for a 2-step walk is given by $p_2(x) = \frac{2}{\pi \sqrt{4-x^2}}$ for $0 \leq x \leq 1$ and 0 otherwise.

5.1 Applications to W_3

The explicit form of $p_2(x)$ leads to some alternative probabilistically inspired formulae for $W_3(s)$. The inner integral in (25) is in fact expressible in terms of the hypergeometric function with details appearing in (BNSW09). For instance, in the case $s = 1$ we find

$$\int_{|x-1|}^{x+1} y h_x(y) dy = \frac{2(x+1)}{\pi} E\left(\frac{2\sqrt{x}}{x+1}\right), \quad (26)$$

(for $x > 0$ and $x \neq 1$) where $E(k) = \frac{\pi}{2} {}_2F_1\left(\frac{1}{2}, -\frac{1}{2}; 1; k^2\right)$ denotes the complete elliptic integral of the second kind with parameter k . This leads to the following expression for the 3-step walk:

$$W_3(1) = \int_0^2 \frac{4(x+1)}{\pi^2 \sqrt{4-x^2}} E\left(\frac{2\sqrt{x}}{x+1}\right) dx. \quad (27)$$

We are now in a position to prove Theorem 4.1.

Proof of Theorem 4.1: It remains to prove the result for odd integers. Since, as noted, for all complex s , the function $V_3(s)$ defined in (13) also satisfies the recursion given for $W_3(s)$ in Example 4.4, it suffices to show that the values given for $s = 1$ and $s = -1$ are correct. First, (BB87, Exercise 1c), p. 16) allows us to write

$$(x+1) E\left(2 \frac{\sqrt{x}}{x+1}\right) = \operatorname{Re} (2 E(x) - (1-x^2) K(x))$$

for $0 < x < \infty$ where we have used Jacobi's imaginary transformations (BB87, Exercises 7a) & 8b), p. 73) to introduce the real part for $x > 1$. Thus, from (27),

$$\begin{aligned} W_3(1) &= \frac{4}{\pi^2} \operatorname{Re} \int_0^{\pi/2} \left(2 E(2 \sin(t)) - (1 - 4 \sin^2(t)) K(2 \sin(t)) \right) dt \\ &= \frac{4}{\pi^2} \operatorname{Re} \int_0^{\pi/2} \int_0^{\pi/2} 2 \sqrt{1 - 4 \sin^2(t) \sin^2(r)} dt dr \\ &\quad - \frac{4}{\pi^2} \operatorname{Re} \int_0^{\pi/2} \int_0^{\pi/2} \frac{1 - 4 \sin^2(t)}{\sqrt{1 - 4 \sin^2(t) \sin^2(r)}} dt dr. \end{aligned}$$

Joining up the two last integrals and parameterizing, we consider

$$Q(a) := \frac{4}{\pi^2} \int_0^{\pi/2} \int_0^{\pi/2} \frac{1 + a^2 \sin^2(t) - 2 a^2 \sin^2(t) \sin^2(r)}{\sqrt{1 - a^2 \sin^2(t) \sin^2(r)}} dt dr. \quad (28)$$

We now use the binomial theorem to integrate (28) term-by-term for $|a| < 1$ and substitute

$$\frac{2}{\pi} \int_0^{\pi/2} \sin^{2m}(t) dt = (-1)^m \binom{-1/2}{m}$$

throughout. Moreover, $(-1)^m \binom{-\alpha}{m} = (\alpha)_m / m!$ where the latter denoted the *Pochhammer* symbol. Evaluation of the consequent infinite sum produces:

$$\begin{aligned} Q(a) &= \sum_{k \geq 0} (-1)^k \binom{-1/2}{k} \left(a^{2k} \binom{-1/2}{k}^2 - a^{2k+2} \binom{-1/2}{k} \binom{-1/2}{k+1} - 2 a^{2k+2} \binom{-1/2}{k+1}^2 \right) \\ &= \sum_{k \geq 0} (-1)^k a^{2k} \binom{-1/2}{k}^3 \frac{1}{(1 - 2k)^2} \\ &= {}_3F_2 \left(\begin{matrix} -\frac{1}{2}, -\frac{1}{2}, \frac{1}{2} \\ 1, 1 \end{matrix} \middle| a^2 \right). \end{aligned}$$

Analytic continuation to $a = 2$ yields the claimed result as per formula (13) for $s = 1$. The case $s = -1$ is similar, see (BNSW09). \square

6 Conclusion

The behaviour of these two-dimensional walks provides a fascinating blend of probabilistic, analytic, algebraic and combinatorial challenges. Conjecture 4.6 remains mysterious to us as does does its less compelling parent Conjecture 4.5.

Acknowledgements We are grateful to David Bailey for his substantial computational help, to Richard Crandall for helpful suggestions, to Bruno Salvy for reminding us of the existence of (Bar64), Michael Mossinghoff for showing us (Klu1906), and to Peter Donovan for stimulating this research and for much subsequent useful discussion.

References

- [BBBG08] David H. Bailey, Jonathan M. Borwein, David J. Broadhurst, and M. L. Glasser. “Elliptic integral evaluations of Bessel moments and applications.” *J. Phys. A: Math. Theor.*, **41** (2008), 5203–5231.
- [Bar64] P. Barrucand. “Sur la somme des puissances des coefficients multinomiaux et les puissances successives d’une fonction de Bessel.” *Comptes rendus hebdomadaires des séances de l’Académie des sciences*, **258** (1964), 5318–5320.
- [BBC07] J. Borwein, D. Bailey, N. Calkin, R. Girgensohn, R. Luke, and V. Moll, *Experimental Mathematics in Action*, A.K. Peters, 2007.
- [BB87] Jonathan M. Borwein and Peter B. Borwein. *Pi and the AGM: A Study in Analytic Number Theory and Computational Complexity*. Wiley, 1987.
- [BNSW09] J. Borwein, D. Nuyens, A. Straub, and J. Wan. “Random walk integrals.” Preprint, October 2009.
- [Bro09] David Broadhurst. “Bessel moments, random walks and Calabi-Yau equations.” Preprint, November 2009.
- [BS08] Jonathan M. Borwein and Bruno Salvy. “A proof of a recursion for Bessel moments.” *Experimental Mathematics*, **17** (2008), 223–230. [D-drive Preprint 346].
- [CKN06] Ronald Cools, Frances Y. Kuo, and Dirk Nuyens. “Constructing embedded lattice rules for multivariate integration.” *SIAM J. Sci. Comput.*, **28**(2006), 2162–2188.
- [Cra09] R. E. Crandall, “Analytic representations for random walk moments,” preprint, September 2009.
- [CG96] John H. Conway and Richard K. Guy. *The Book of Numbers*. Springer-Verlag, 1996.
- [Don09] Peter Donovan. “The flaw in the JN-25 series of ciphers, II.” *Cryptologia*, (2010), in press.
- [Klu1906] J. C. Kluyver. A local probability problem. *Nederl. Acad. Wetensch. Proc.*, **8** (1906), 341–350.
- [Pea1905] Karl Pearson. “The random walk.” *Nature*, **72** (1905), 294.
- [Pea1905b] Karl Pearson. “The problem of the random walk.” *Nature*, **72** (1905), 342.
- [Ray1905] Lord Rayleigh. “The problem of the random walk.” *Nature*, **72** (1905), 318.
- [RS09] L. B. Richmond and J. Shallit. “Counting abelian squares.” *The Electronic Journal of Combinatorics*, **16**, 2009.
- [Slo09] N. J. A. Sloane. The On-Line Encyclopedia of Integer Sequences, 2009. Published electronically at <http://www.research.att.com/sequences>.
- [Tit39] E. Titchmarsh. *The Theory of Functions*. Oxford University Press, 2nd edition, 1939.