



- (51) **International Patent Classification:**  
G01N 33/564 (2006.01) G01N 33/68 (2006.01)
- (21) **International Application Number:**  
PCT/IB2015/059303
- (22) **International Filing Date:**  
2 December 2015 (02.12.2015)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**  
62/086,549 2 December 2014 (02.12.2014) US  
62/157,903 6 May 2015 (06.05.2015) US
- (71) **Applicant:** NESTEC S.A. [CH/CH]; Avenue Nestlé 55, CH-1800 Vevey (CH).
- (72) **Inventors:** SINGH, Sharat; 9410 Carroll Park Drive, San Diego, California 92121 (US). JAIN, Anjali; 9410 Carroll Park Drive, San Diego, California 92121 (US). KONDRA-GUNTA, Venkateswarlu; 9410 Carroll Park Drive, San Diego, California 92121 (US).
- (81) **Designated States** (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM,

AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.


- (84) **Designated States** (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

**Published:**  
— with international search report (Art. 21(3))

WO 2016/088068 A1

(54) **Title:** METHODS FOR ESTABLISHING A VEDOLIZUMAB DOSING REGIMEN TO TREAT PATIENTS WITH IRRITABLE BOWEL DISEASE

FIG. 1A



Higher TNFa levels at baseline result in lower vedolizumab at week 6

(57) **Abstract:** The present invention provides methods for predicting whether an individual having inflammatory bowel disease (IBD) is likely to respond to vedolizumab treatment. Also provided are methods for predicting whether an individual with IBD such as Crohn's disease or ulcerative colitis will develop autoantibodies against vedolizumab. The present invention also provides a treatment regimen for an IBD patient which includes measuring the level of one or more predictive markers of response to vedolizumab prior to administering the anti- $\alpha 4\beta 7$  integrin drug.

**METHODS FOR ESTABLISHING A VEDOLIZUMAB DOSING  
5 REGIMEN TO TREAT PATIENTS WITH IRRITABLE BOWEL  
DISEASE**

**CROSS-REFERENCE TO RELATED APPLICATIONS**

[0001] This application claims priority to U.S. Provisional Application Nos. 62/086,549,  
filed December 2, 2014 and 62/157,903, filed May 6, 2015, the contents of which are hereby  
10 incorporated by reference in their entirety for all purposes.

**BACKGROUND OF THE INVENTION**

[0002] Inflammatory bowel disease (IBD), which occurs world-wide and afflicts millions  
of people, is the collective term used to describe three gastrointestinal disorders of unknown  
etiology: Crohn's disease (CD), ulcerative colitis (UC), and indeterminate colitis (IC). IBD,  
15 together with irritable bowel syndrome (IBS), will affect one-half of all Americans during  
their lifetime, at a cost of greater than \$2.6 billion dollars for IBD and greater than \$8 billion  
dollars for IBS. A primary determinant of these high medical costs is the difficulty of  
diagnosing digestive diseases and how these diseases will progress. The cost of IBD and IBS  
is compounded by lost productivity, with people suffering from these disorders missing at  
20 least 8 more days of work annually than the national average.

[0003] Despite the successes of anti-TNF $\alpha$  therapies in the treatment of IBD, a  
subpopulation of patients are refractory to treatment, highlighting an unmet medical need for  
new therapies. Vedolizumab is a gut-specific,  $\alpha 4\beta 7$  integrin-neutralizing monoclonal Ab,  
which does not affect peripheral blood cell counts and appears to lack systemic effects.

25 Vedolizumab is a new anti-inflammatory treatment option for the management of therapy-  
refractory patients.

[0004] There is a need in the art for methods of therapeutic management of diseases such  
as ulcerative colitis and Crohn's Disease using an individualized approach to monitor drug  
efficacy and optimize therapy accordingly. The methods need to include assessing disease  
30 course and clinical parameters such as pharmacodynamics, disease activity indices, disease  
burden, and inflammatory biomarkers. There is a need to predict whether a patient will  
respond to vedolizumab therapy. The present invention satisfies this and other needs.

**BRIEF SUMMARY OF THE INVENTION**

**[0005]** The present invention provides methods for predicting whether a subject having inflammatory bowel disease (IBD) will respond to anti- $\alpha 4\beta 7$  integrin drug treatment. The method comprises: (a) detecting the presence or level of at least one predictive marker  
5 selected from the group consisting of TNF $\alpha$ , human serum albumin (HSA), VEGF, angiopoietin-1 (ANG-1), angiopoietin-2 (ANG-2), adenosine deaminase (ADA), serum  $\alpha 4\beta 7$  integrin, IL-12p40, C-reactive protein (CRP), matrix metalloproteinase 9 (MMP9), MAdCAM-1, VCAM-1, ICAM-1, or a combination thereof in a sample from the subject; and  
10 (b) classifying the subject as a responder or a non-responder to the anti- $\alpha 4\beta 7$  integrin drug treatment according to a predictive marker profile based on a higher or lower level of the at least one predictive marker compared to a corresponding reference value. In some  
embodiments, classifying comprises applying a statistical analysis to the predictive marker  
profile to determine if the subject is a responder or a non-responder to the anti- $\alpha 4\beta 7$  integrin  
drug treatment. In some embodiments, the at least one predictive marker comprises at least 1,  
15 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13 or more predictive markers. In some embodiments, the inflammatory bowel disease is ulcerative colitis (UC) or Crohn's Disease (CD).

**[0006]** In some embodiments, the subject is classified as a responder if the level of ANG-1, ADA, HSA, IL-12p40, MMP9, ICAM-1, and/or serum  $\alpha 4\beta 7$  integrin is higher than the  
20 corresponding reference value. In some instances, the subject is classified as a responder if the level of MadCAM-1, VCAM-1 and/or TNF $\alpha$  is lower than the corresponding reference value. In other embodiments, the subject is classified as a non-responder if the level of TNF $\alpha$ , VEGF, ANG-2, CRP, and/or VCAM-1 is higher than the corresponding reference value.

**[0007]** In some embodiments, detecting the presence or level of the at least one predictive  
25 marker comprises performing a proximity dual detection assay or an immunoassay. In some cases, the proximity dual detection assay is a Collaborative Enzyme Enhanced Reactive Immunoassay (CEER<sup>TM</sup>).

**[0008]** In some embodiments, the anti- $\alpha 4\beta 7$  integrin drug is ENTYVIO<sup>®</sup> (vedolizumab).

**[0009]** In some embodiments, the sample is selected from the group consisting of a whole  
30 blood, serum or plasma sample.

**[0010]** In some embodiments, the subject has had an inadequate response with, has lost response to, or was intolerant to an anti-TNF $\alpha$  drug. The anti-TNF $\alpha$  drug can be a member

selected from the group consisting of REMICADE<sup>®</sup> (infliximab), ENBREL<sup>®</sup> (etanercept), HUMIRA<sup>®</sup> (adalimumab), CIMZIA<sup>®</sup> (certolizumab pegol), SIMPONI<sup>®</sup> (golimumab), STELARA<sup>®</sup> (ustekinumab), and combinations thereof. In other embodiments, the subject has not previously been administered an anti- $\alpha 4\beta 7$  integrin drug.

5 [0011] In some instances, the method further comprises applying a statistical analysis to the predictive marker profile to predict whether the subject will develop autoantibodies against an anti- $\alpha 4\beta 7$  integrin drug.

[0012] Also provided herein is a method for predicting whether a subject having inflammatory bowel disease (IBD) is likely to develop autoantibodies against an anti- $\alpha 4\beta 7$  integrin drug. The method comprises (a) detecting the presence or level of at least one  
10 predictive marker selected from the group consisting of TNF $\alpha$ , human serum albumin (HSA), VEGF, angiopoietin-1 (ANG-1), angiopoietin-2 (ANG-2), adenosine deaminase (ADA), serum  $\alpha 4\beta 7$  integrin, IL-12p40, C-reactive protein (CRP), MMP9, MAdCAM-1, VCAM-1, ICAM-1, and a combination thereof in a sample from the subject; and (b) applying a  
15 statistical analysis to the presence or level of the at least one predictive marker of step (a) to generate a predictive marker profile to determine whether the subject is likely to develop or not to develop autoantibodies against the anti- $\alpha 4\beta 7$  integrin drug. In one embodiment, the method further comprises determining the presence or level of anti-drug antibodies in the sample from the subject. In some embodiments, the anti- $\alpha 4\beta 7$  integrin drug is ENTYVIO<sup>®</sup>  
20 (vedolizumab).

[0013] In some embodiments, the inflammatory bowel disease is ulcerative colitis (UC) or Crohn's Disease (CD).

[0014] The at least one predictive marker can include at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13 or more predictive markers. In some embodiments, detecting the presence or level of  
25 at least one predictive marker comprises performing a proximity dual detection assay or an immunoassay. In one embodiment, the proximity dual detection assay is a Collaborative Enzyme Enhanced Reactive Immunoassay (CEER<sup>™</sup>).

[0015] In some embodiments, the subject is likely not to develop autoantibodies if the level of ANG-1, ADA, HSA, IL-12p40, MMP9, ICAM-1, and/or serum  $\alpha 4\beta 7$  integrin is higher  
30 than the corresponding reference value. In other embodiments, the subject is likely not to develop autoantibodies if the level of MAdCAM-1, VCAM-1 and/or TNF $\alpha$  is lower than the corresponding reference value. In other embodiments, the subject is likely to develop

autoantibodies if the level of TNF $\alpha$ , VEGF, ANG-2, CRP, and/or VCAM-1 is higher than the corresponding reference value.

5 [0016] In some embodiments, the subject has been previously administered the anti- $\alpha$ 4 $\beta$ 7 integrin drug. In some cases, the subject has not been administered the anti- $\alpha$ 4 $\beta$ 7 integrin drug.

[0017] In some embodiments, the subject is maintained or is recommended to be maintained on the anti- $\alpha$ 4 $\beta$ 7 integrin drug if the subject is likely not to develop autoantibodies. The subject may be administered the anti- $\alpha$ 4 $\beta$ 7 integrin drug.

10 [0018] In other embodiments, the subject is not maintained or is recommended not to be maintained on the anti- $\alpha$ 4 $\beta$ 7 integrin drug if the subject is likely to develop autoantibodies. The subject may not be administered the anti- $\alpha$ 4 $\beta$ 7 integrin drug.

[0019] In some embodiments, the sample is selected from the group consisting of a whole blood, serum or plasma sample.

[0020] In some embodiments, the anti- $\alpha$ 4 $\beta$ 7 integrin drug is ENTYVIO<sup>®</sup> (vedolizumab).

15 [0021] In some embodiments, the subject has had an inadequate response with, has lost response to, or is intolerant to an anti-TNF $\alpha$  drug. The anti-TNF $\alpha$  drug may be a member selected from the group consisting of REMICADE<sup>®</sup> (infliximab), ENBREL<sup>®</sup> (etanercept), HUMIRA<sup>®</sup> (adalimumab), CIMZIA<sup>®</sup> (certolizumab pegol), SIMPONI<sup>®</sup> (golimumab), STELARA<sup>®</sup> (ustekinumab), and combinations thereof.

20 [0022] The present invention also provides a method for predicting whether a subject having inflammatory bowel disease (IBD) will develop autoantibodies against an anti- $\alpha$ 4 $\beta$ 7 integrin drug at a later time point during a course of therapy with the anti- $\alpha$ 4 $\beta$ 7 integrin drug. The method includes (a) measuring the presence or level of at least one predictive marker at a first time point in a sample from the subject to determine a first predictive marker profile,  
25 wherein the at least one predictive marker is selected from the group consisting of TNF $\alpha$ , human serum albumin (HSA), VEGF, angiopoietin-1 (ANG-1), angiopoietin-2 (ANG-2), adenosine deaminase (ADA), serum  $\alpha$ 4 $\beta$ 7 integrin, IL-12p40, C-reactive protein (CRP), MMP9, MAdCAM-1, VCAM-1, ICAM-1, and a combination thereof; (b) measuring the presence of level of the same at least one predictive marker at a later time point in a second  
30 sample from the subject to determine a second predictive marker profile; and (c) applying a statistical analysis to the first and second predictive marker profiles to determine that the

subject will develop autoantibodies to the anti- $\alpha 4\beta 7$  integrin drug during the course of therapy.

[0023] In some embodiments, the at least one predictive marker comprises at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, or more predictive markers.

5 [0024] In some embodiments, detecting the presence or level of at least one predictive marker comprises performing a proximity dual detection assay or an immunoassay. In some instances, the proximity dual detection assay is a Collaborative Enzyme Enhanced Reactive Immunoassay (CEER™).

[0025] In some embodiments, the anti- $\alpha 4\beta 7$  integrin drug is ENTYVIO® (vedolizumab).

10 [0026] In some embodiments, the later time point is at weeks 2, 4, 6, 8, 10, 12, 16, 24, 32, 40, 48, or 52 during the course of therapy.

[0027] Provided herein is a method for inducing clinical remission in a subject having inflammatory bowel disease (IBD) and after receiving an induction therapy of an anti- $\alpha 4\beta 7$  integrin drug. The method comprises administering a maintenance therapy of an anti- $\alpha 4\beta 7$  integrin drug to a subject with IBD and having a higher level of serum  $\alpha 4\beta 7$  integrin and a lower level of MadCAM-1 compared to corresponding levels at baseline. In some  
15 embodiments, clinical remission corresponds to a score on the Crohn's Disease Activity Index (CDAI) of less than or equal to 150. The maintenance therapy can include administering a dose of about 300 mg of the anti- $\alpha 4\beta 7$  integrin drug, *e.g.*, vedolizumab every  
20 8 weeks during the course of therapy. Alternatively, the maintenance therapy includes administering a dose of about 300 mg of ENTYVIO® (vedolizumab) every 4 weeks of the regimen.

[0028] Also provided herein is a method for inducing a clinical response in a subject having inflammatory bowel disease (IBD), comprising administering an induction therapy of  
25 an anti- $\alpha 4\beta 7$  integrin drug to a subject with IBD and having a higher level of ANG-1, ADA, HSA, IL-12p40, MMP9, ICAM-1, and/or serum  $\alpha 4\beta 7$  integrin, and/or a lower level of MadCAM-1, VCAM-1, and/or TNF $\alpha$  compared to a corresponding reference value. In some embodiments, the induction therapy comprises administering a dose of about 300 mg of the anti- $\alpha 4\beta 7$  integrin drug, *e.g.*, vedolizumab at weeks 0, 2 and 6 during the course of therapy.  
30 In some embodiments, clinical response corresponds to a 70-point or greater decrease in CDAI score. In other embodiments, clinical response in a subject with CD corresponds to a

100-point or more decrease in CDAI score. In some embodiments, clinical response in a subject with UC corresponds to a reduction in the Mayo Clinical score of at least 3 points and a decrease of at least 30% from the baseline score, with a decrease of at least 1 point on the rectal bleeding score of 0 or 1 for UC.

5 [0029] The present invention provides a therapy regimen for an anti- $\alpha 4\beta 7$  integrin drug, the therapy regimen comprising measuring the presence or level of at least one predictive marker selected from the group consisting of TNF $\alpha$ , human serum albumin (HSA), VEGF, angiopoietin-1 (ANG-1), angiopoietin-2 (ANG-2), adenosine deaminase (ADA), serum  $\alpha 4\beta 7$  integrin, IL-12p40, C-reactive protein (CRP), MMP9, MAdCAM-1, VCAM-1, ICAM-1, and  
10 a combination thereof in a sample from the subject; and administering the therapy regimen of the anti- $\alpha 4\beta 7$  integrin drug according to a predictive marker profile based on a higher or lower level of the at least one predictive marker compared to a corresponding reference value.

[0030] In some embodiments, the therapy regimen of the anti- $\alpha 4\beta 7$  integrin drug is  
15 administered if the level of MadCAM-1 and/or TNF $\alpha$  is lower than the corresponding reference value. In other embodiments, the therapy regimen is not administered if the level of TNF $\alpha$ , VEGF, ANG-2, CRP, and/or VCAM-1 is higher than the corresponding reference value. In some instances, the therapy regimen is an induction therapy regimen. For instance, an induction therapy regimen for ENTYVIO<sup>®</sup> (vedolizumab) comprises administering a dose  
20 of about 300 mg of the anti- $\alpha 4\beta 7$  integrin drug at weeks 0, 2 and 6 of the regimen.

[0031] In some embodiments, the therapy regimen of the anti- $\alpha 4\beta 7$  integrin drug is administered if the level of MadCAM-1 is lower and/or the level of serum  $\alpha 4\beta 7$  integrin is higher than the corresponding reference value. In some instances, the therapy regimen is a  
25 maintenance therapy regimen. For instance, a maintenance therapy regimen for ENTYVIO<sup>®</sup> (vedolizumab) comprises administering a dose of about 300 mg of the anti- $\alpha 4\beta 7$  integrin drug every 8 weeks of the regimen. In other cases, the maintenance therapy regimen for ENTYVIO<sup>®</sup> (vedolizumab) includes administering a dose of about 300 mg of the anti- $\alpha 4\beta 7$  integrin drug every 4 weeks of the regimen.

[0032] These and other aspects, objects, embodiments will become more apparent when  
30 read with the detailed description and figures that follow.

## BRIEF DESCRIPTION OF THE DRAWINGS

[0033] FIGS. 1A and 1B depict the associated between TNF $\alpha$  levels and vedolizumab in patient with CD or UC. FIG. 1A shows a diagram of an exemplary embodiment of the present invention that shows higher TNF $\alpha$  levels at baseline result in lower vedolizumab at week 6. FIG. 1B shows a diagram of an exemplary embodiment of the present invention that shows higher TNF $\alpha$  levels at week 2 result in lower vedolizumab at week 6.

[0034] FIG. 2 shows a diagram of an exemplary embodiment of the present invention that shows higher VEGF levels at baseline result in lower vedolizumab at week 6.

[0035] FIGS. 3A and 3B show the association between two angiogenesis markers ANG-1 and ANG-2 and vedolizumab levels. FIG. 3A shows a diagram of an exemplary embodiment of the present invention that shows higher ANG-2 levels at baseline result in lower vedolizumab at week 6. FIG. 3B shows a diagram of an exemplary embodiment of the present invention that shows higher higher ANG-1 levels at baseline result in higher vedolizumab at week 6.

[0036] FIGS. 4A and 4B show the relationship between level of adenosine deaminase (ADA) and vedolizumab during therapy. FIG. 4A shows a diagram of an exemplary embodiment of the present invention that shows higher ADA levels at baseline result in higher vedolizumab at week 6. FIG. 4B shows a diagram of an exemplary embodiment of the present invention that shows higher ADA levels at week 2 result in higher vedolizumab at week 6.

[0037] FIGS. 5A and 5B provide bivariate fit log graphs used to determine whether a specific biomarker is associated with the level of vedolizumab. FIG. 5A shows higher albumin levels at baseline result in higher vedolizumab at week 6. FIG. 5B shows a lack of association between vedolizumab at week 6 and hemoglobin (HGB) at baseline.

[0038] FIGS. 6A and 6B illustrate the relationship between either IL-12p40 or MMP9 and vedolizumab. FIG. 6A shows a positive association between vedolizumab at week 6 and IL-12p40 levels at baseline. FIG. 6B shows a positive association between vedolizumab at week 6 and MMP9 levels at week 2.

[0039] FIG. 7A shows a positive association between vedolizumab at week 6 and MAdCAM-1 at week 2. FIG. 7B shows an association between vedolizumab at week 6 and CRP at baseline.



[0040] FIGS. 8A and 8B shows a diagram of an exemplary embodiment of the present invention that shows that treatment with vedolizumab result in increase in serum  $\alpha 4\beta 7$  integrin (FIG. 8A) and decrease in MAdCAM-1 (FIG. 8B) levels.

[0041] FIG. 9 shows the changes in serum  $\alpha 4\beta 7$  integrin levels during the course of vedolizumab therapy (*e.g.*, at baseline, post week 0, week 2 and week 6) in samples tested.

[0042] FIG. 10 shows a decrease in serum MAdCAM-1 levels during the course of vedolizumab therapy (*e.g.*, at baseline, post week 0, week 2 and week 6) in samples tested.

[0043] FIGS. 11A and 11B provide graphs of multiple regression models used to evaluate the correlations between two predictive biomarkers at baseline of vedolizumab treatment and levels of vedolizumab at week 6. FIG. 11A represent VCAM-1 and  $\alpha 4\beta 7$  integrin at baseline to predict vedolizumab levels at week 6. FIG. 11B shows multiple regression model with interaction between VCAM-1 and MAdCAM-1 (baseline) to predict vedolizumab at week 6.

[0044] FIGS. 12A and 12B provide bivariate fit log graphs of MadCAM-1 levels versus ICAM-1 or VCAM-1. FIG. 12A shows that baseline MAdCAM-1 levels are negatively correlated with soluble ICAM-1 levels. FIG. 12B shows a negative correlation between MAdCAM-1 at baseline and soluble VCAM-1 at baseline.

[0045] FIG. 13 shows the longitudinal change of MadCAM-1 levels during vedolizumab induction therapy and pairwise p-values.

[0046] FIG. 14 shows no change of VCAM-1 levels during vedolizumab induction therapy.

[0047] FIG. 15 shows the longitudinal change of serum  $\alpha 4\beta 7$  integrin levels during vedolizumab induction therapy.

[0048] FIG. 16 shows stable levels of vedolizumab across induction therapy.

[0049] FIG. 17 shows the longitudinal change of MAdCAM-1 across visits in patients responding to vedolizumab. P-values are calculated for change from baseline for each visit from induction therapy through maintenance therapy. The differences are statistically significant.

[0050] FIG. 18 shows the longitudinal change of serum  $\alpha 4\beta 7$  integrin across visits in patients responding to vedolizumab. P-values are calculated for change from baseline for

each visit from induction therapy through maintenance therapy. The differences are statistically significant.

[0051] FIGS. 19A and 19B provide bivariate fit log graphs that compare serum albumin levels and vedolizumab levels. FIG. 19A shows that higher levels of serum albumin at baseline are associated with higher levels of vedolizumab at week 6. FIG. 19B shows that higher levels of serum albumin at baseline are associated with higher levels of vedolizumab at week 14.

[0052] FIGS. 20A and 20B provide bivariate fit log graphs that compare TNF $\alpha$  levels and vedolizumab levels. FIG. 20A shows that lower levels of TNF $\alpha$  (at baseline) are associated with higher levels of vedolizumab at week 2. FIG. 20B shows that higher levels of TNF $\alpha$  at baseline are associated with lower levels of vedolizumab at week 6.

[0053] FIG. 21 provides a graph of the changes in s- $\alpha$ 4 $\beta$ 7 integrin levels between clinical responders of vedolizumab and non-responders, as determined by the Physician's Global Assessment (PGA). The graph shows changes in the levels from week 0 (baseline) to week 2 ("1"), from week 0 to week 6 ("2") and from week 0 to week 14 ("3") of vedolizumab treatment in clinical responder or non-responders. The change in s- $\alpha$ 4 $\beta$ 7 integrin levels was determined by subtracting the level at the earlier time point from the level at the later time point. The patients in this study were administered standard induction therapy and maintenance therapy of vedolizumab.

[0054] FIG. 22 provides a graph of the changes in s- $\alpha$ 4 $\beta$ 7 integrin levels between clinical responders of vedolizumab and non-responders, as determined by the withdrawal of steroid therapy. The graph shows changes in the levels from week 0 (baseline) to week 2 ("1"), from week 0 to week 6 ("2") and from week 0 to week 14 ("3") of vedolizumab treatment in clinical responder or non-responders. The change in s- $\alpha$ 4 $\beta$ 7 integrin levels was determined by subtracting the level at the earlier time point from the level at the later time point.

## DETAILED DESCRIPTION OF THE INVENTION

### I. DEFINITIONS

[0055] The term "inflammatory bowel disease" or "IBD" includes gastrointestinal disorders such as, *e.g.*, Crohn's disease (CD), ulcerative colitis (UC), and indeterminate colitis (IC). Inflammatory bowel diseases (*e.g.*, CD, UC, and IC) are distinguished from all

other disorders, syndromes, and abnormalities of the gastroenterological tract, including irritable bowel syndrome (IBS).

**[0056]** The term “sample” as used herein includes any biological specimen obtained from a patient. Samples include, without limitation, whole blood, plasma, serum, red blood cells, white blood cells (*e.g.*, peripheral blood mononuclear cells (PBMC), polymorphonuclear (PMN) cells), ductal lavage fluid, nipple aspirate, lymph (*e.g.*, disseminated tumor cells of the lymph node), bone marrow aspirate, saliva, urine, stool (*i.e.*, feces), sputum, bronchial lavage fluid, tears, fine needle aspirate (*e.g.*, harvested by random periareolar fine needle aspiration), any other bodily fluid, a tissue sample such as a biopsy of a site of inflammation (*e.g.*, needle biopsy), and cellular extracts thereof. In some embodiments, the sample is whole blood or a fractional component thereof such as plasma, serum, or a cell pellet. In other embodiments, the sample is obtained by isolating PBMCs and/or PMN cells using any technique known in the art. In yet other embodiments, the sample is a tissue biopsy, *e.g.*, from a site of inflammation such as a portion of the gastrointestinal tract or synovial tissue.

**[0057]** The term “marker” or “biomarker” includes any biochemical marker, serological marker, genetic marker, or other clinical or echographic characteristic that can be used predicting whether a subject having inflammatory bowel disease (IBD) will respond to vedolizumab treatment. The marker can be used to classify a sample from the subject is a responder or a non-responder to vedolizumab therapy. In some embodiments, the markers are utilized in combination with a statistical analysis to provide a prognosis of IBD in an individual.

**[0058]** The term “classifying” includes “to associate” or “to categorize” a sample with a disease state. In certain instances, “classifying” is based on statistical evidence, empirical evidence, or both. In certain embodiments, the methods and systems of classifying use a so-called training set of samples having known disease states. Once established, the training data set serves as a basis, model, or template against which the features of an unknown sample are compared, in order to classify the unknown disease state of the sample. In certain instances, classifying the sample is akin to diagnosing the disease state of the sample. In certain other instances, classifying the sample is akin to differentiating the disease state of the sample from another disease state.

**[0059]** The present invention relies, in part, on determining the presence (or absence) or level (*e.g.*, concentration) of at least one marker in a sample obtained from an individual. As

used herein, the term “detecting the presence of at least one marker” includes determining the presence of each marker of interest by using any quantitative or qualitative assay known to one of skill in the art. In certain instances, qualitative assays that determine the presence or absence of a particular trait, variable, genotype, and/or biochemical or serological substance (e.g., protein or antibody) are suitable for detecting each marker of interest. In certain other instances, quantitative assays that determine the presence or absence of DNA, RNA, protein, antibody, or activity are suitable for detecting each marker of interest. As used herein, the term “detecting the level of at least one marker” includes determining the level of each marker of interest by using any direct or indirect quantitative assay known to one of skill in the art. In certain instances, quantitative assays that determine, for example, the relative or absolute amount of DNA, RNA, protein, antibody, or activity are suitable for detecting the level of each marker of interest. One skilled in the art will appreciate that any assay useful for detecting the level of a marker is also useful for detecting the presence or absence of the marker.

**[0060]** The term “predictive profile” includes 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 30, 35, 40, 45, 50, or more marker(s) of an individual, wherein the marker(s) can be a serological marker, a protein marker, a genetic marker, and the like. In some embodiments, the marker profile together with a statistical analysis can provide physicians and caregivers valuable diagnostic insight. In other embodiments, the marker profile with optionally a statistical analysis provides a projected disease condition (e.g., IBD or CD). By using multiple markers (e.g., serological, inflammation, protein, etc.) in conjunction with statistical analyses, the assays described herein provide diagnostic, prognostic and therapeutic value by identifying patients with IBD or a clinical subtype thereof, predicting risk of developing complicated disease, assisting in assessing the rate of disease progression (e.g., rate of progression to complicated disease or surgery), and assisting in the selection of therapy.

**[0061]** The term “individual,” “subject,” or “patient” typically refers to humans, but also to other animals including, e.g., other primates, rodents, canines, felines, equines, ovines, porcines, and the like.

**[0062]** The term “prognosis” includes a prediction of the probable course and outcome of UC or CD or the likelihood of recovery from the disease.

**[0063]** The term “monitoring the progression or regression of UC or CD” includes the use of the methods of the present invention to determine the disease state (*e.g.*, severity of UC) of an individual. In some aspects, the methods of the present invention can also be used to predict the progression of UC or CD, *e.g.*, by determining a likelihood for UC to progress either rapidly or slowly in an individual based on the presence or level of at least one marker in a sample. In other aspects, the methods of the present invention can also be used to predict the regression of UC, *e.g.*, by determining a likelihood for UC to regress either rapidly or slowly in an individual based on the presence or level of at least one marker in a sample.

**[0064]** The term “course of therapy” includes any therapeutic approach taken to relieve or prevent one or more symptoms associated with UC or CD. The term encompasses administering any compound, drug, procedure, or regimen useful for improving the health of an individual with UC and includes any of the therapeutic agents as well as surgery. One skilled in the art will appreciate that either the course of therapy or the dose of the current course of therapy can be changed, *e.g.*, based upon the methods of the present invention.

**[0065]** The phrase “determining the course of therapy” and the like includes the use of an empirically derived index, score or analysis to select for example, selecting a dose of drug, selecting an appropriate drug, or a course or length of therapy, a therapy regimen, or maintenance of an existing drug or dose. In certain aspects, a derived or measured index can be used to determine the course of therapy.

**[0066]** As used herein, the phrase “at a later time point” includes phrases such as “by a later time point” and “within the later time point.” For example, a method for predicting whether a subject will develop autoantibodies to an anti- $\alpha 4\beta 7$  integrin drug (*e.g.*, vedolizumab) at a later time point during a course of therapy includes a method for predicting whether a subject will develop autoantibodies to an anti- $\alpha 4\beta 7$  integrin drug by the later time point during the course of therapy as well as a method for predicting whether a subject will develop autoantibodies to an anti- $\alpha 4\beta 7$  integrin drug within the later time point during the course of therapy.

**[0067]** In “quartile analysis”, there are three numbers (values) that divide a range of data into four equal parts. The first quartile (also called the ‘lower quartile’) is the number below which lies the bottom 25 percent of the data. The second quartile (the ‘median’) divides the range in the middle and has 50 percent of the data below it. The third quartile (also called the ‘upper quartile’) has 75 percent of the data below it and the top 25 percent of the data above

it. As a non-limiting example, quartile analysis can be applied to the concentration level of a marker such as an antibody or other protein marker described herein, such that a marker level in the first quartile (<25%) is assigned a value of 1, a marker level in the second quartile (25-50%) is assigned a value of 2, a marker level in the third quartile (51%-<75%) is assigned a value of 3, and a marker level in the fourth quartile (75%-100%) is assigned a value of 4.

## II. Detailed Description of the Embodiments

[0068] The present invention provides methods for predicting whether a subject having inflammatory bowel disease (IBD) will respond to vedolizumab treatment, the method comprises:

- 10 (a) determining a predictive marker profile by detecting the presence or level of at least one predictive marker in a sample from a subject; and
- (b) classifying the sample as either a responder or a non-responder to vedolizumab therapy.

[0069] The present invention also provides methods for predicting whether a subject having inflammatory bowel disease (IBD) will develop anti-drug antibodies (autoantibodies) against an anti- $\alpha 4\beta 7$  integrin drug (*e.g.*, vedolizumab), the method comprises:

- 15 (a) determining a predictive marker profile by detecting the presence or level of at least one predictive marker in a sample from a subject; and
- (b) applying a statistical analysis to the predictive marker profile to determine that the subject is likely to develop or not develop anti-drug antibodies.

[0070] Additionally, the present invention also provides methods for predicting whether a subject having inflammatory bowel disease (IBD) will develop anti-drug antibodies (autoantibodies) against an anti- $\alpha 4\beta 7$  integrin drug (*e.g.*, vedolizumab) at a later time point during a course of therapy with the anti- $\alpha 4\beta 7$  integrin drug, the method comprises:

- 25 (a) measuring the presence or level of at least one predictive marker at a first time point in a sample from the subject to determine a first predictive marker profile;
- (b) measuring the presence or level of the same at least one predictive marker at a later time point in a sample from the subject to determine a second predictive marker profile; and

(c) applying a statistical analysis to the first and second predictive marker profiles to determine that the subject will develop autoantibodies to the anti- $\alpha 4\beta 7$  integrin drug, during the course of therapy.

[0071] In certain aspects, vedolizumab (ENTYVIO<sup>®</sup>) is indicated in adult patients with moderately to severely active ulcerative colitis (UC) or Crohn's Disease (CD) who have had an inadequate response with, lost response to, or were intolerant to a tumor necrosis factor (TNF) therapy, or had an inadequate response with, were intolerant to, or demonstrated dependence on corticosteroids for inducing and maintaining clinical response, inducing and maintaining clinical remission, improving endoscopic appearance of the mucosa, and achieving corticosteroid-free remission.

[0072] Vedolizumab can be administered as an intravenous infusion over 30 minutes. The recommended dosage of in adults with ulcerative colitis or Crohn's disease is 300 mg administered by intravenous infusion at zero, two and six weeks (0, 2, and 6 as induction therapy) and then every eight weeks thereafter (14 weeks, 22 weeks, *etc.* as maintenance therapy). Alternatively, the maintenance therapy may include administering vedolizumab every 4 weeks. Therapy may be discontinued in patients who show no evidence of therapeutic benefit by week 14.

#### A. Inflammatory bowel disease (IBD)

[0073] Inflammatory bowel disease (IBD) is a group of inflammatory conditions of the large intestine and small intestine. The main forms of IBD are Crohn's disease (CD) and ulcerative colitis (UC). Other less common forms of IBD include, *e.g.*, indeterminate colitis (IC), collagenous colitis, lymphocytic colitis, ischemic colitis, diversion colitis, Behçet's syndrome, infective colitis, and the like. U.S. Patent Publication 2008/0131439, entitled "Methods of Diagnosing Inflammatory Bowel Disease" is incorporated herein by reference for all purposes.

##### 1. Crohn's Disease

[0074] Crohn's disease (CD) is a disease of chronic inflammation that can involve any part of the gastrointestinal tract. Commonly, the distal portion of the small intestine, *i.e.*, the ileum, and the cecum are affected. In other cases, the disease is confined to the small intestine, colon, or anorectal region. CD occasionally involves the duodenum and stomach, and more rarely the esophagus and oral cavity.

[0075] The variable clinical manifestations of CD are, in part, a result of the varying anatomic localization of the disease. The most frequent symptoms of CD are abdominal pain, diarrhea, and recurrent fever. CD is commonly associated with intestinal obstruction or fistula, an abnormal passage between diseased loops of bowel. CD also includes  
5 complications such as inflammation of the eye, joints, and skin, liver disease, kidney stones, and amyloidosis. In addition, CD is associated with an increased risk of intestinal cancer.

[0076] Several features are characteristic of the pathology of CD. The inflammation associated with CD, known as transmural inflammation, involves all layers of the bowel wall. Thickening and edema, for example, typically also appear throughout the bowel wall, with  
10 fibrosis present in long-standing forms of the disease. The inflammation characteristic of CD is discontinuous in that segments of inflamed tissue, known as “skip lesions,” are separated by apparently normal intestine. Furthermore, linear ulcerations, edema, and inflammation of the intervening tissue lead to a “cobblestone” appearance of the intestinal mucosa, which is distinctive of CD.

[0077] A hallmark of CD is the presence of discrete aggregations of inflammatory cells, known as granulomas, which are generally found in the submucosa. Some CD cases display typical discrete granulomas, while others show a diffuse granulomatous reaction or a nonspecific transmural inflammation. As a result, the presence of discrete granulomas is  
15 indicative of CD, although the absence of granulomas is also consistent with the disease.

Thus, transmural or discontinuous inflammation, rather than the presence of granulomas, is a preferred diagnostic indicator of CD (Rubin and Farber, Pathology (Second Edition), Philadelphia, J.B. Lippincott Company (1994)).

[0078] Crohn’s disease may be categorized by the behavior of disease as it progresses. This was formalized in the Vienna classification of Crohn’s disease. See, Gasche *et al.*,  
25 *Inflamm. Bowel Dis.*, 6:8-15 (2000). There are three categories of disease presentation in Crohn’s disease: (1) stricturing, (2) penetrating, and (3) inflammatory. Stricturing disease causes narrowing of the bowel which may lead to bowel obstruction or changes in the caliber of the feces. Penetrating disease creates abnormal passageways (fistulae) between the bowel and other structures such as the skin. Inflammatory disease (also known as non-stricturing,  
30 non-penetrating disease) causes inflammation without causing strictures or fistulae.

[0079] As such, Crohn’s disease represents a number of heterogeneous disease subtypes that affect the gastrointestinal tract and may produce similar symptoms. As used herein in



reference to CD, the term “clinical subtype” includes a classification of CD defined by a set of clinical criteria that distinguish one classification of CD from another. As non-limiting examples, subjects with CD can be classified as having stricturing (*e.g.*, internal stricturing), penetrating (*e.g.*, internal penetrating), or inflammatory disease as described herein, or these  
5 subjects can additionally or alternatively be classified as having fibrostenotic disease, small bowel disease, internal perforating disease, perianal fistulizing disease, UC-like disease, the need for small bowel surgery, the absence of features of UC, or combinations thereof.

**[0080]** In certain instances, subjects with CD can be classified as having complicated CD, which is a clinical subtype characterized by stricturing or penetrating phenotypes. In certain  
10 other instances, subjects with CD can be classified as having a form of CD characterized by one or more of the following complications: fibrostenosis, internal perforating disease, and the need for small bowel surgery. In further instances, subjects with CD can be classified as having an aggressive form of fibrostenotic disease requiring small bowel surgery. Criteria relating to these subtypes have been described, for example, in Gasche *et al.*, *Inflamm. Bowel*  
15 *Dis.*, 6:8-15 (2000); Abreu *et al.*, *Gastroenterology*, 123:679-688 (2002); Vasiliauskas *et al.*, *Gut*, 47:487-496 (2000); Vasiliauskas *et al.*, *Gastroenterology*, 110:1810-1819 (1996); and Greenstein *et al.*, *Gut*, 29:588-592 (1988).

**[0081]** The “fibrostenotic subtype” of CD is a classification of CD characterized by one or more accepted characteristics of fibrostenosing disease. Such characteristics of  
20 fibrostenosing disease include, but are not limited to, documented persistent intestinal obstruction or an intestinal resection for an intestinal obstruction. The fibrostenotic subtype of CD can be accompanied by other symptoms such as perforations, abscesses, or fistulae, and can further be characterized by persistent symptoms of intestinal blockage such as nausea, vomiting, abdominal distention, and inability to eat solid food. Intestinal X-rays of  
25 patients with the fibrostenotic subtype of CD can show, for example, distention of the bowel before the point of blockage.

**[0082]** The requirement for small bowel surgery in a subject with the fibrostenotic subtype of CD can indicate a more aggressive form of this subtype. Additional subtypes of CD are also known in the art and can be identified using defined clinical criteria. For example,  
30 internal perforating disease is a clinical subtype of CD defined by current or previous evidence of entero-enteric or entero-vesicular fistulae, intra-abdominal abscesses, or small bowel perforation. Perianal perforating disease is a clinical subtype of CD defined by current

or previous evidence of either perianal fistulae or abscesses or rectovaginal fistula. The UC-like clinical subtype of CD can be defined by current or previous evidence of left-sided colonic involvement, symptoms of bleeding or urgency, and crypt abscesses on colonic biopsies. Disease location can be classified based on one or more endoscopic, radiologic, or pathologic studies.

[0083] One skilled in the art understands that overlap can exist between clinical subtypes of CD and that a subject having CD can have more than one clinical subtype of CD. For example, a subject having CD can have the fibrostenotic subtype of CD and can also meet clinical criteria for a clinical subtype characterized by the need for small bowel surgery or the internal perforating disease subtype. Similarly, the markers described herein can be associated with more than one clinical subtype of CD.

## 2. Ulcerative Colitis

[0084] Ulcerative colitis (UC) is a disease of the large intestine characterized by chronic diarrhea with cramping, abdominal pain, rectal bleeding, loose discharges of blood, pus, and mucus. The manifestations of UC vary widely. A pattern of exacerbations and remissions typifies the clinical course for about 70% of UC patients, although continuous symptoms without remission are present in some patients with UC. Local and systemic complications of UC include arthritis, eye inflammation such as uveitis, skin ulcers, and liver disease. In addition, UC, and especially the long-standing, extensive form of the disease is associated with an increased risk of colon carcinoma.

[0085] UC is a diffuse disease that usually extends from the most distal part of the rectum for a variable distance proximally. The term “left-sided colitis” describes an inflammation that involves the distal portion of the colon, extending as far as the splenic flexure. Sparing of the rectum or involvement of the right side (proximal portion) of the colon alone is unusual in UC. The inflammatory process of UC is limited to the colon and does not involve, for example, the small intestine, stomach, or esophagus. In addition, UC is distinguished by a superficial inflammation of the mucosa that generally spares the deeper layers of the bowel wall. Crypt abscesses, in which degenerated intestinal crypts are filled with neutrophils, are also typical of UC (Rubin and Farber, *supra*).

[0086] In certain instances, with respect to UC, the variability of symptoms reflect differences in the extent of disease (*i.e.*, the amount of the colon and rectum that are inflamed) and the intensity of inflammation. Disease starts at the rectum and moves “up” the

colon to involve more of the organ. UC can be categorized by the amount of colon involved. Typically, patients with inflammation confined to the rectum and a short segment of the colon adjacent to the rectum have milder symptoms and a better prognosis than patients with more widespread inflammation of the colon.

5 [0087] In comparison with CD, which is a patchy disease with frequent sparing of the rectum, UC is characterized by a continuous inflammation of the colon that usually is more severe distally than proximally. The inflammation in UC is superficial in that it is usually limited to the mucosal layer and is characterized by an acute inflammatory infiltrate with neutrophils and crypt abscesses. In contrast, CD affects the entire thickness of the bowel wall  
10 with granulomas often, although not always, present. Disease that terminates at the ileocecal valve, or in the colon distal to it, is indicative of UC, while involvement of the terminal ileum, a cobblestone-like appearance, discrete ulcers, or fistulas suggests CD.

[0088] The different types of ulcerative colitis are classified according to the location and the extent of inflammation. As used herein in reference to UC, the term “clinical subtype”  
15 includes a classification of UC defined by a set of clinical criteria that distinguish one classification of UC from another. As non-limiting examples, subjects with UC can be classified as having ulcerative proctitis, proctosigmoiditis, left-sided colitis, pancolitis, fulminant colitis, and combinations thereof. Criteria relating to these subtypes have been described, for example, in Kornbluth *et al.*, *Am. J. Gastroenterol.*, 99: 1371-85 (2004).

20 [0089] Ulcerative proctitis is a clinical subtype of UC defined by inflammation that is limited to the rectum. Proctosigmoiditis is a clinical subtype of UC which affects the rectum and the sigmoid colon. Left-sided colitis is a clinical subtype of UC which affects the entire left side of the colon, from the rectum to the place where the colon bends near the spleen and begins to run across the upper abdomen (the splenic flexure). Pancolitis is a clinical subtype  
25 of UC which affects the entire colon. Fulminant colitis is a rare, but severe form of pancolitis. Patients with fulminant colitis are extremely ill with dehydration, severe abdominal pain, protracted diarrhea with bleeding, and even shock.

[0090] In some embodiments, classification of the clinical subtype of UC is important in planning an effective course of treatment. While ulcerative proctitis, proctosigmoiditis, and  
30 left-sided colitis can be treated with local agents introduced through the anus, including steroid-based or other enemas and foams, pancolitis must be treated with oral medication so that active ingredients can reach all of the affected portions of the colon.

[0091] One skilled in the art understands that overlap can exist between clinical subtypes of UC and that a subject having UC can have more than one clinical subtype of UC. Similarly, the prognostic markers described herein can be associated with more than one clinical subtype of UC.

### 5           3.       **Patients with CD or UC**

[0092] In some embodiments, the subjects of methods disclosed herein are patients with moderate to severe CD or a score of about 220 to 450 on the Crohn's Disease Activity Index (CDAI ranges from 0 to about 600, with higher scores indicating greater disease activity. In other embodiments, the subjects have moderate to severe UC or a Mayo Clinic score ranging  
10 from about 6 to 12 (Mayo Clinic scores range from 0 to 12 with higher scores indicating active disease), with a sigmoidoscopy subscore of at least 2, and disease that extends 15 cm or more from the anal verge.

[0093] In some embodiments, the subject has not received an anti- $\alpha 4\beta 7$  integrin drug (*e.g.*, vedolizumab). In some embodiments, the subject has not received an anti-TNF $\alpha$  therapy.  
15 The subject may be predicted to be nonresponsive to an anti-TNF $\alpha$  drug. In other embodiments, the subject has developed an intolerance to the anti-TNF $\alpha$  drug. In some instances, the subject has had an inadequate response to the anti-TNF $\alpha$  drug. In other instances, the subject has lost response to the anti-TNF $\alpha$  drug.

[0094] In some aspects of the present invention, the method is performed at baseline (*e.g.*,  
20 prior to receiving an anti- $\alpha 4\beta 7$  integrin drug). The presence or level of one or more predictive markers described herein may be detected or quantitated at a single time point. In other aspects, the method is performed during induction therapy (*e.g.*, at week 0 to week 6 of anti- $\alpha 4\beta 7$  integrin drug treatment). In some embodiments, the presence or level of one or more predictive markers are measured at one or more time points during induction therapy. In  
25 yet other aspects, the method is performed during maintenance therapy (*e.g.*, at week 8 or later of anti- $\alpha 4\beta 7$  integrin drug treatment). In some instances, the presence or level of one or more predictive markers are measured at one or more time points during maintenance therapy.

### **B.       Markers for Predicting Response to Vedolizumab**

30 [0095] A variety of IBD markers, including biochemical markers, serological markers, protein markers, genetic markers, and other clinical or echographic characteristics, are

suitable for use in the methods of the present invention for predicting response to vedolizumab therapy. In certain aspects, prognostic methods described herein utilize the application of an algorithm (*e.g.*, statistical analysis) to the presence or concentration level determined for one or more of the markers to aid or assist in a prognosis regarding whether a UC or CD patient will respond to vedolizumab therapy.

**[0096]** The following predictive markers are suitable for use in the present invention. The markers can make up a marker profile. Suitable markers include, but are not limited to, TNF $\alpha$ , human serum albumin (HSA), VEGF, angiopoietin-1 (ANG-1), angiopoietin-2 (ANG-2), adenosine deaminase (ADA), serum  $\alpha 4\beta 7$  integrin, IL-12p40, C-reactive protein (CRP), MMP9, MAdCAM-1, VCAM-1, and ICAM-1.

**[0097]** In some embodiments, the methods provided herein include measuring/detecting the presence or level one or more, *e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 or more predictive markers. In some embodiments, the method includes detecting the presence or level of serum  $\alpha 4\beta 7$  integrin and MAdCAM-1. In some embodiments, the method includes detecting the presence or level of serum  $\alpha 4\beta 7$  integrin and TNF $\alpha$ . In some embodiments, the method includes detecting the presence or level of serum  $\alpha 4\beta 7$  integrin and HSA. In some embodiments, the method includes detecting the presence or level of serum  $\alpha 4\beta 7$  integrin and ANG-1. In some embodiments, the method includes detecting the presence or level of serum  $\alpha 4\beta 7$  integrin and ANG-2. In some embodiments, the method includes detecting the presence or level of serum  $\alpha 4\beta 7$  integrin and IL-12p40. In some embodiments, the method includes detecting the presence or level of serum  $\alpha 4\beta 7$  integrin and CRP. In some embodiments, the method includes detecting the presence or level of serum  $\alpha 4\beta 7$  integrin and MMP9. In other embodiments, the method includes detecting the presence or level of serum  $\alpha 4\beta 7$  integrin and ICAM-1. In other embodiments, the method includes detecting the presence or level of serum  $\alpha 4\beta 7$  integrin and VCAM-1. In some aspects, the method includes detecting the presence or level of serum  $\alpha 4\beta 7$  integrin, MAdCAM-1, and one or more additional markers provided herein.

**[0098]** In some embodiments, the method includes detecting the presence or level of MAdCAM-1 and VCAM-1. In other embodiments, the method includes detecting the presence or level of MAdCAM-1 and ICAM-1. In some embodiments, the method includes detecting the presence or level of MAdCAM-1 and TNF $\alpha$ .

[0099] In some embodiments, the method includes detecting the presence or level of ANG-1, ADA, HSA, IL-12p40, MMP9, ICAM-1, VCAM-1, and serum  $\alpha 4\beta 7$  integrin. In some embodiments, the method includes detecting the presence or level of ANG-1, ADA, HSA, IL-12p40, MMP9, ICAM-1, and VCAM-1. In some embodiments, the method includes detecting the presence or level of ANG-1, ADA, HAS, IL-12p40, MMP9, ICAM-1, and VCAM-1.

[0100] In some embodiments, the methods include measuring/detecting the presence or level one or more, *e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 or more predictive markers at one or more, *e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more time points.

In some embodiments, the method includes detecting the presence or level of one or more, *e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 or more predictive markers at, for example, baseline (prior to the administration of vedolizumab), post week 0 (immediately after the initial administration of the drug), week 2, week 4, week 6, week 8, week 10, week 12, week 14, week 16, week 18, week 20, week 22, week 24, week 26, week 28, week 30, week 32, week 34, week 36, week 38, week 40, week 42, week 44, week 46, week 48, week 50, or week 52 of drug treatment, or any combination thereof. In other instances, the presence or level of one or more predictive markers, *e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 or more predictive markers, is measured once a week or less often during the course of treatment.

[0101] In some embodiments, the presence or level of one or more, *e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 or more predictive markers are detected at baseline. In some embodiments, the presence or level of one or more, *e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 or more predictive markers are detected at week 2. In some embodiments, the presence or level of one or more, *e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 or more predictive markers are detected at week 4. In some embodiments, the presence or level of one or more, *e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 or more predictive markers are detected at week 6. In some embodiments, the presence or level of one or more, *e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 or more predictive markers are detected prior to the development of anti-drug antibodies. In other embodiments, the presence or level of one or more, *e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 or more predictive markers are detected prior to the development of anti-drug antibodies at any time point during maintenance therapy.

[0102] In certain instances, the presence or level of a particular biomarker is detected at the level of mRNA expression with an assay such as, for example, a hybridization assay or an amplification-based assay. In certain other instances, the presence or level of a particular biomarker is detected at the level of protein expression using, for example, an immunoassay (e.g., ELISA), an immunohistochemical assay or a proximity dual detection assay. Suitable ELISA kits for determining the presence or level of a biomarker in a sample such as a serum, plasma, saliva, or urine sample are available from, e.g., R&D Systems, Inc. (Minneapolis, MN), Neogen Corp. (Lexington, KY), Alpco Diagnostics (Salem, NH), Assay Designs, Inc. (Ann Arbor, MI), BD Biosciences Pharmingen (San Diego, CA), Invitrogen (Camarillo, CA), Calbiochem (San Diego, CA), CHEMICON International, Inc. (Temecula, CA), Antigenix America Inc. (Huntington Station, NY), QIAGEN Inc. (Valencia, CA), Bio-Rad Laboratories, Inc. (Hercules, CA), and/or Bender MedSystems Inc. (Burlingame, CA). In some embodiments, the proximity dual detection assay is a CEER™ (Collaborative Enzyme Enhanced Reactive immunoassay) assay, an antibody-microarray based platform is utilized to form a unique “triple-antibody-enzyme-channeling” immuno-complex capable of measuring analytes of limited availability in a sample. A detailed description of CEER™ is found in, e.g., U.S. Patent No. 8,163,499, which is hereby incorporated by reference in its entirety for all purposes.

### 1. Growth Factors

[0103] The determination of the presence or level of one or more growth factors in a sample is also useful in the present invention. As used herein, the term “growth factor” includes any of a variety of peptides, polypeptides, or proteins that are capable of stimulating cellular proliferation and/or cellular differentiation. In certain aspects, the presence or level of at least one growth factor including, but not limited to, vascular endothelial growth factor (VEGF).

[0104] The term “VEGF” or “vascular endothelial growth factor” refers to isolated nucleic acids, polypeptides and polymorphic variants, alleles, mutants, and interspecies homologues thereof and as further described herein, that: (1) have an amino acid sequence that has greater than about 60% amino acid sequence identity, 65%, 70%, 75%, 80%, 85%, 90%, preferably 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% or greater amino acid sequence identity, preferably over a region of at least about 50, 75, 100, 150, 200, 250, 300, 400, 410, or more amino acids, to a human VEGF sequence shown below; (2) bind to antibodies, e.g.,

polyclonal antibodies, raised against an immunogen comprising an amino acid sequence shown below, or conservatively modified variants thereof; (3) bind to a VEGF binding protein; (4) compete with a naturally occurring VEGF ligand binding to a VEGF ligand binding protein; (5) induce vasculogenesis and/or angiogenesis in cells having a membrane-bound VEGF binding protein; (6) specifically hybridize under stringent hybridization conditions to a nucleic acid sequence shown below, or conservatively modified variants thereof; (7) have a nucleic acid sequence that has greater than about 90%, preferably greater than about 96%, 97%, 98%, 99%, or higher nucleotide sequence identity, preferably over a region of at least about 100, 200, 300, 400 or more nucleotides, to a human VEGF mRNA sequence; and/or (8) have at least 25, often 50, 75, 100, 125 or 143 contiguous amino acid residues of a human VEGF polypeptide sequence.

**[0105]** The human VEGF polypeptide sequence is set forth in, *e.g.*, Genbank Accession No. NP\_001020537, NP\_001020538, NP\_001020539, NP\_001020540, NP\_001020541, NP\_001028928, NP\_001165093, NP\_001165094, NP\_001165095, NP\_001165096, NP\_001165097, NP\_001165098, NP\_001165099, NP\_001165100, NP\_001165101, NP\_001191313, and NP\_001191314. The human VEGF mRNA (coding) sequence is set forth in, *e.g.*, Genbank Accession No. NM\_001025366, NM\_001025367, NM\_001025368, NM\_001025370, NM\_001025356, NM\_001033756, NM\_001171622, NM\_001171623, NM\_00117624, NM\_001171625, NM\_001171626, NM\_001171627, NM\_001171628, NM\_001171629, NM\_001171630, NM\_001204384, NM\_001204385, and NM\_003376. One skilled in the art will appreciate that VEGF is also known as vascular endothelial growth factor, VEGF1, VEGF-A, VEGFA, VPF, vascular permeability factor, and MVCD1. One skilled in the art will appreciate that variants, isoforms, alternative sequences of VEGF are also useful in the present invention.

## 25                    2.        **Intercellular Adhesion Molecule-1 (ICAM-1)**

**[0106]** The term “ICAM-1” or “intercellular adhesion molecule 1” refers to isolated nucleic acids, polypeptides and polymorphic variants, alleles, mutants, and interspecies homologues thereof and as further described herein, that: (1) have an amino acid sequence that has greater than about 60% amino acid sequence identity, 65%, 70%, 75%, 80%, 85%, 90%, preferably 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% or greater amino acid sequence identity, preferably over a region of at least about 50, 75, 100, 150, 200, 250, 300, 400, 500, 525, or more amino acids, to a human ICAM-1 sequence shown below; (2) bind to antibodies, *e.g.*,



polyclonal antibodies, raised against an immunogen comprising an amino acid sequence shown below, or conservatively modified variants thereof; (3) bind to a ICAM-1 binding protein; (4) specifically hybridize under stringent hybridization conditions to a nucleic acid sequence shown below, or conservatively modified variants thereof; (5) have a nucleic acid sequence that has greater than about 90%, preferably greater than about 96%, 97%, 98%, 99%, or higher nucleotide sequence identity, preferably over a region of at least about 100, 200, 300, 400 or more nucleotides, to a human ICAM-1 mRNA sequence; and/or (6) have at least 25, often 50, 75, 100, 125 or 143 contiguous amino acid residues of a human ICAM-1 polypeptide sequence. The human ICAM-1 polypeptide sequence is set forth in, *e.g.*, Genbank Accession No. NP\_000192. The human ICAM-1 mRNA (coding) sequence is set forth in, *e.g.*, Genbank Accession No. NM\_000201. One skilled in the art will appreciate that variants, isoforms, alternative sequences of ICAM-1 are also useful in the present invention.

**[0107]** ICAM-1 is a transmembrane cellular adhesion protein that is continuously present in low concentrations in the membranes of leukocytes and endothelial cells. Upon cytokine stimulation, the concentrations greatly increase. ICAM-1 can be induced by IL-1 and TNF $\alpha$  and is expressed by the vascular endothelium, macrophages, and lymphocytes. In IBD, proinflammatory cytokines cause inflammation by upregulating expression of adhesion molecules such as ICAM-1. The increased expression of adhesion molecules recruit more lymphocytes to the infected tissue, resulting in tissue inflammation (*see, Goke et al., J., Gastroenterol.*, 32:480 (1997); and Rijcken *et al., Gut*, 51:529 (2002)). ICAM-1 is encoded by the intercellular adhesion molecule 1 gene (ICAM1; Entrez GeneID: 3383; Genbank Accession No. NM\_000201) and is produced after processing of the intercellular adhesion molecule 1 precursor polypeptide (Genbank Accession No. NP\_000192).

### 3. Vascular Cell Adhesion Molecule-1 (VCAM-1)

**[0108]** The term “VCAM-1” or “vascular cell adhesion molecule 1” refers to isolated nucleic acids, polypeptides and polymorphic variants, alleles, mutants, and interspecies homologues thereof and as further described herein, that: (1) have an amino acid sequence that has greater than about 60% amino acid sequence identity, 65%, 70%, 75%, 80%, 85%, 90%, preferably 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% or greater amino acid sequence identity, preferably over a region of at least about 50, 75, 100, 150, 200, 250, 300, 400, 500, 600, 700, 720, 730, or more amino acids, to a human VCAM-1 sequence shown below; (2) bind to antibodies, *e.g.*, polyclonal antibodies, raised against an immunogen

comprising an amino acid sequence shown below, or conservatively modified variants thereof; (3) bind to a VCAM-1 binding protein; (4) specifically hybridize under stringent hybridization conditions to a nucleic acid sequence shown below, or conservatively modified variants thereof; (5) have a nucleic acid sequence that has greater than about 90%, preferably greater than about 96%, 97%, 98%, 99%, or higher nucleotide sequence identity, preferably over a region of at least about 100, 200, 300, 400 or more nucleotides, to a human VCAM-1 mRNA sequence; and/or (6) have at least 25, often 50, 75, 100, 125 or 143 contiguous amino acid residues of a human VCAM-1 polypeptide sequence.

[0109] VCAM-1 is a transmembrane cellular adhesion protein that mediates the adhesion of lymphocytes, monocytes, eosinophils, and basophils to vascular endothelium. Upregulation of VCAM-1 in endothelial cells by cytokines occurs as a result of increased gene transcription (*e.g.*, in response to Tumor necrosis factor- $\alpha$  (TNF $\alpha$ ) and Interleukin-1 (IL-1)). VCAM-1 is encoded by the vascular cell adhesion molecule 1 gene (VCAM1; Entrez GeneID: 7412) and is produced after differential splicing of the transcript (Genbank Accession No. NM\_001078 (variant 1) or NM\_080682 (variant 2)), and processing of the precursor polypeptide splice isoform (Genbank Accession No. NP\_001069 (isoform a) or NP\_542413 (isoform b)).

[0110] In certain instances, the presence or level of an IgSF CAM is detected at the level of mRNA expression with an assay such as, *e.g.*, a hybridization assay or an amplification-based assay. In certain other instances, the presence or level of an IgSF CAM such as ICAM-1 or VCAM-1 is detected at the level of protein expression using, for example, an immunoassay (*e.g.*, ELISA or an immuno electrochemiluminescence assay) or an immunohistochemical assay. Suitable antibodies and/or ELISA kits for determining the presence or level of ICAM-1 and/or VCAM-1 in a sample such as a tissue sample, biopsy, serum, plasma, saliva, urine, or stool are available from, *e.g.*, Invitrogen (Camarillo, CA), Santa Cruz Biotechnology, Inc. (Santa Cruz, CA), and/or Abcam Inc. (Cambridge, MA).

#### 4. Acute Phase Proteins

[0111] The determination of the presence or level of one or more acute-phase proteins in a sample is also useful in the present invention. Acute-phase proteins are a class of proteins whose plasma concentrations increase (positive acute-phase proteins) or decrease (negative acute-phase proteins) in response to inflammation. This response is called the acute-phase

reaction (also called acute-phase response). Examples of positive acute-phase proteins include, but are not limited to, C-reactive protein (CRP).

**[0112]** The term “CRP” or “C-reactive protein” refers to isolated nucleic acids, polypeptides and polymorphic variants, alleles, mutants, and interspecies homologues thereof and as further described herein, that: (1) have an amino acid sequence that has greater than about 60% amino acid sequence identity, 65%, 70%, 75%, 80%, 85%, 90%, preferably 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% or greater amino acid sequence identity, preferably over a region of at least about 50, 75, 100, 150, 200, 210, 220 or more amino acids, to a human CRP sequence shown below; (2) bind to antibodies, *e.g.*, polyclonal antibodies, raised against an immunogen comprising an amino acid sequence shown below, or conservatively modified variants thereof; (3) bind to a CRP binding protein; (4) specifically hybridize under stringent hybridization conditions to a nucleic acid sequence shown below, or conservatively modified variants thereof; (5) have a nucleic acid sequence that has greater than about 90%, preferably greater than about 96%, 97%, 98%, 99%, or higher nucleotide sequence identity, preferably over a region of at least about 100, 200, 300, 400 or more nucleotides, to a human CRP mRNA sequence; and/or (6) have at least 25, often 50, 75, 100, 125 or 143 contiguous amino acid residues of a human CRP polypeptide sequence.

**[0113]** CRP is a protein found in the blood in response to inflammation (an acute-phase protein). CRP is typically produced by the liver and by fat cells (adipocytes). It is a member of the pentraxin family of proteins. The human CRP polypeptide sequence is set forth in, *e.g.*, Genbank Accession No. NP\_000558. The human CRP mRNA (coding) sequence is set forth in, *e.g.*, Genbank Accession No. NM\_000567. One skilled in the art will appreciate that CRP is also known as PTX1, MGC88244, and MGC149895.

**[0114]** In certain instances, the presence or level of a particular acute-phase protein is detected at the level of mRNA expression with an assay such as, for example, a hybridization assay or an amplification-based assay. In certain other instances, the presence or level of a particular acute-phase protein is detected at the level of protein expression using, for example, an immunoassay (*e.g.*, ELISA) or an immunohistochemical assay. For example, a sandwich colorimetric ELISA assay available from Alpco Diagnostics (Salem, NH) can be used to determine the level of CRP in a serum, plasma, urine, or stool sample. Similarly, an ELISA kit available from Biomeda Corporation (Foster City, CA) can be used to detect CRP

levels in a sample. Other methods for determining CRP levels in a sample are described in, *e.g.*, U.S. Patent Nos. 6,838,250; 6,406,862; and 7,439,019; and U.S. Patent Publication No. 2006/0019410. Additional methods for determining CRP levels include, *e.g.*,

immunoturbidimetry assays, rapid immunodiffusion assays, and visual agglutination assays.

- 5 Suitable ELISA kits for determining the presence or level of SAA in a sample such as serum, plasma, saliva, urine, or stool are available from, *e.g.*, Antigenix America Inc. (Huntington Station, NY), Abazyme (Needham, MA), USCN Life (Missouri City, TX), and/or U.S. Biological (Swampscott, MA).

### 5. Mucosal Addressin Cell Adhesion Molecule (MAdCAM-1)

- 10 **[0115]** The term “mucosal addressin cell adhesion molecule” or “MAdCAM-1” refers to isolated nucleic acids, polypeptides and polymorphic variants, alleles, mutants, and interspecies homologues thereof and as further described herein, that: (1) have an amino acid sequence that has greater than about 60% amino acid sequence identity, 65%, 70%, 75%, 80%, 85%, 90%, preferably 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% or greater
- 15 amino acid sequence identity, preferably over a region of at least about 50, 75, 100, 150, 200, 250, 281, 290 or more amino acids, to a human MAdCAM-1 sequence shown below; (2) bind to antibodies, *e.g.*, polyclonal antibodies, raised against an immunogen comprising an amino acid sequence shown below, or conservatively modified variants thereof; (3) bind to a MAdCAM-1 binding protein; (4) specifically hybridize under stringent hybridization
- 20 conditions to a nucleic acid sequence shown below, or conservatively modified variants thereof; (5) have a nucleic acid sequence that has greater than about 90%, preferably greater than about 96%, 97%, 98%, 99%, or higher nucleotide sequence identity, preferably over a region of at least about 100, 200, 300, 400 or more nucleotides, to a human MAdCAM-1 mRNA sequence; and/or (6) have at least 25, often 50, 75, 100, 125 or 143 contiguous amino
- 25 acid residues of a human MAdCAM-1 polypeptide sequence.

- [0116]** MAdCAM-1 is a predictive marker which is essential in mediating the infiltration of leucocytes into chronically inflamed tissues and plays a pivotal role in T-lymphocyte homing to the gut. The human MAdCAM-1 polypeptide sequence is set forth in, *e.g.*, Genbank Accession No. NP\_570118. The human MAdCAM-1 mRNA (coding) sequence is set forth
- 30 in, *e.g.*, Genbank Accession No. NM\_130762. One skilled in the art will appreciate that variants, isoforms, alternative sequences of MAdCAM-1 are also useful in the present invention.

[0117] MAdCAM-1 is expressed by intestinal endothelium and its expression is increased under conditions of inflammation, including in the setting of inflammatory bowel disease (IBD). This molecule has been detected in body fluids, such as urine and serum, using a sandwich ELISA assay; however, the mechanism by which it is cleaved from the endothelial surface and released into circulation as soluble (s)-MAdCAM-1 is not well defined. s-MAdCAM-1 was detected during treatment with vedolizumab, a novel alpha-4 beta-7 ( $\alpha 4\beta 7$ ) antagonist approved for treatment of ulcerative colitis (UC) and Crohn's disease (CD).

## 6. TNF $\alpha$

[0118] The term "TNF $\alpha$ " or "tumor necrosis factor  $\alpha$ " refers to isolated nucleic acids, polypeptides and polymorphic variants, alleles, mutants, and interspecies homologues thereof and as further described herein, that: (1) have an amino acid sequence that has greater than about 60% amino acid sequence identity, 65%, 70%, 75%, 80%, 85%, 90%, preferably 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% or greater amino acid sequence identity, preferably over a region of at least about 50, 75, 100, 150, 200, 225, 230 or more amino acids, to a human TNF $\alpha$  sequence shown below; (2) bind to antibodies, *e.g.*, polyclonal antibodies, raised against an immunogen comprising an amino acid sequence shown below, or conservatively modified variants thereof; (3) bind to a TNF $\alpha$  binding protein; (4) compete with a naturally occurring TNF $\alpha$  ligand binding to a TNF $\alpha$  ligand binding protein; (5) induce apoptosis in cells having a membrane-bound TNF $\alpha$  binding protein; (6) specifically hybridize under stringent hybridization conditions to a nucleic acid sequence shown below, or conservatively modified variants thereof; (7) have a nucleic acid sequence that has greater than about 90%, preferably greater than about 96%, 97%, 98%, 99%, or higher nucleotide sequence identity, preferably over a region of at least about 100, 200, 300, 400 or more nucleotides, to a human TNF $\alpha$  mRNA sequence; and/or (8) have at least 25, often 50, 75, 100, 125 or 143 contiguous amino acid residues of a human TNF $\alpha$  polypeptide sequence. The human TNF $\alpha$  polypeptide sequence is set forth in, *e.g.*, Genbank Accession No. NP\_000585. The human TNF $\alpha$  mRNA (coding) sequence is set forth in, *e.g.*, Genbank Accession No. NM\_000594. One skilled in the art will appreciate that variants, isoforms, alternative sequences of TNF $\alpha$  are also useful in the present invention.

[0119] In other embodiments, an immunoassay such as a sandwich assay or ELISA can be used to measure TNF $\alpha$ . Non-limiting examples include Human TNF $\alpha$  High Sensitivity ELISA (Cat. No. BMS223HS, eBioscience, San Diego, CA), Erenna Human TNF $\alpha$

immunoassay (Cat. No. 03-0022-xx, Singulex, Alameda, CA), Human TNF $\alpha$  cytokine assay (Cat. No. K151BHA-5, Meso Scale Diagnostics (MSD), Rockville, MD)) and a multi-marker immunoassay (*e.g.*, as described in U.S. Patent No. 8,450,069; Singulex).

### 7. Human Serum Albumin (HSA)

5 [0120] The term “human serum albumin” or “HSA” refers to isolated nucleic acids, polypeptides and polymorphic variants, alleles, mutants, and interspecies homologues thereof and as further described herein, that: (1) have an amino acid sequence that has greater than about 60% amino acid sequence identity, 65%, 70%, 75%, 80%, 85%, 90%, preferably 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% or greater amino acid sequence identity,  
10 preferably over a region of at least about 50, 75, 100, 150, 200, 250, 281, 300, 400, 500, 600, or more amino acids, to a human HSA sequence shown below; (2) bind to antibodies, *e.g.*, polyclonal antibodies, raised against an immunogen comprising an amino acid sequence shown below, or conservatively modified variants thereof; (3) bind to an HSA binding protein; (4) specifically hybridize under stringent hybridization conditions to a nucleic acid  
15 sequence shown below, or conservatively modified variants thereof; (5) have a nucleic acid sequence that has greater than about 90%, preferably greater than about 96%, 97%, 98%, 99%, or higher nucleotide sequence identity, preferably over a region of at least about 100, 200, 300, 400 or more nucleotides, to a human HSA mRNA sequence; and/or (6) have at least 25, often 50, 75, 100, 125 or 143 contiguous amino acid residues of a human HSA  
20 polypeptide sequence. The human HSA polypeptide sequence is set forth in, *e.g.*, Genbank Accession No. NP\_000468. The human HSA mRNA (coding) sequence is set forth in, *e.g.*, Genbank Accession No. NM\_000477. One skilled in the art will appreciate that variants, isoforms, alternative sequences of HSA are also useful in the present invention.

### 8. Angiopoietin-1 and -2 (ANG-1 and ANG-2)

25 [0121] The term “ANG-1” or “angiopoietin-1” refers to isolated nucleic acids, polypeptides and polymorphic variants, alleles, mutants, and interspecies homologues thereof and as further described herein, that: (1) have an amino acid sequence that has greater than about 60% amino acid sequence identity, 65%, 70%, 75%, 80%, 85%, 90%, preferably 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% or greater amino acid sequence identity, preferably  
30 over a region of at least about 50, 75, 100, 150, 200, 250, 281, 300, 400, 450, 475, or more amino acids, to a human ANG-1 sequence shown below; (2) bind to antibodies, *e.g.*, polyclonal antibodies, raised against an immunogen comprising an amino acid sequence

shown below, or conservatively modified variants thereof; (3) bind to a ANG-1 binding protein; (4) compete with a naturally occurring ANG-1 ligand binding to a ANG-1 ligand binding protein; (5) induce angiogenesis in cells having a membrane-bound ANG-1 binding protein; (6) specifically hybridize under stringent hybridization conditions to a nucleic acid sequence shown below, or conservatively modified variants thereof; (7) have a nucleic acid sequence that has greater than about 90%, preferably greater than about 96%, 97%, 98%, 99%, or higher nucleotide sequence identity, preferably over a region of at least about 100, 200, 300, 400 or more nucleotides, to a human ANG-1 mRNA sequence; and/or (8) have at least 25, often 50, 75, 100, 125 or 143 contiguous amino acid residues of a human ANG-1 polypeptide sequence. The human ANG-1 polypeptide sequence is set forth in, *e.g.*, Genbank Accession No. NP\_001137 and NP\_001146. The human ANG-1 mRNA (coding) sequence is set forth in, *e.g.*, Genbank Accession No. NM\_001146 and NM\_001199859. One skilled in the art will appreciate that variants, isoforms, alternative sequences of ANG-1 are also useful in the present invention.

15 **[0122]** The term “ANG-2” or “angiopoietin-2” refers to isolated nucleic acids, polypeptides and polymorphic variants, alleles, mutants, and interspecies homologues thereof and as further described herein, that: (1) have an amino acid sequence that has greater than about 60% amino acid sequence identity, 65%, 70%, 75%, 80%, 85%, 90%, preferably 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% or greater amino acid sequence identity, preferably over a region of at least about 50, 75, 100, 150, 200, 250, or 281 amino acids, to a human ANG-2 sequence shown below; (2) bind to antibodies, *e.g.*, polyclonal antibodies, raised against an immunogen comprising an amino acid sequence shown below, or conservatively modified variants thereof; (3) bind to a ANG-2 binding protein; (4) compete with a naturally occurring ANG-2 ligand binding to a ANG-2 ligand binding protein; (5) induce angiogenesis in cells having a membrane-bound ANG-2 binding protein; (6) specifically hybridize under stringent hybridization conditions to a nucleic acid sequence shown below, or conservatively modified variants thereof; (7) have a nucleic acid sequence that has greater than about 90%, preferably greater than about 96%, 97%, 98%, 99%, or higher nucleotide sequence identity, preferably over a region of at least about 100, 200, 300, 400 or more nucleotides, to a human ANG-2 mRNA sequence; and/or (8) have at least 25, often 50, 75, 100, 125 or 143 contiguous amino acid residues of a human ANG-2 polypeptide sequence. The human ANG-2 polypeptide sequence is set forth in, *e.g.*, Genbank Accession Nos. NP\_001112359, NP\_001112360 and NP\_001138. The human ANG-2 mRNA (coding) sequence is set forth

in, *e.g.*, Genbank Accession Nos. NM\_001118887, NM\_001118888 and NM\_001147. One skilled in the art will appreciate that variants, isoforms, alternative sequences of ANG-2 are also useful in the present invention.

### 9. Adenosine Deaminase (ADA)

5 [0123] The term “adenosine deaminase” or “ADA” refers to isolated nucleic acids, polypeptides and polymorphic variants, alleles, mutants, and interspecies homologues thereof and as further described herein, that: (1) have an amino acid sequence that has greater than about 60% amino acid sequence identity, 65%, 70%, 75%, 80%, 85%, 90%, preferably 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% or greater amino acid sequence identity,  
10 preferably over a region of at least about 50, 75, 100, 150, 200, 250, or 281 amino acids, to a human ADA sequence shown below; (2) bind to antibodies, *e.g.*, polyclonal antibodies, raised against an immunogen comprising an amino acid sequence shown below, or conservatively modified variants thereof; (3) bind to an ADA binding protein; (4) specifically hybridize under stringent hybridization conditions to a nucleic acid sequence shown below,  
15 or conservatively modified variants thereof; (5) have a nucleic acid sequence that has greater than about 90%, preferably greater than about 96%, 97%, 98%, 99%, or higher nucleotide sequence identity, preferably over a region of at least about 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, or more nucleotides, to a human ADA mRNA sequence; and/or (6) have  
20 at least 25, often 50, 75, 100, 125 or 143 contiguous amino acid residues of a human ADA polypeptide sequence. The human ADA polypeptide sequence is set forth in, *e.g.*, Genbank Accession No. NP\_000013. The human ADA mRNA (coding) sequence is set forth in, *e.g.*, Genbank Accession No. NM\_000022. One skilled in the art will appreciate that variants, isoforms, alternative sequences of ADA are also useful in the present invention.

### 10. $\alpha 4\beta 7$ Integrin

25 [0124] The term “ $\alpha 4\beta 7$  integrin” refers to isolated nucleic acids, polypeptides and polymorphic variants, alleles, mutants, and interspecies homologues thereof and as further described herein, that: (1) have an amino acid sequence that has greater than about 60% amino acid sequence identity, 65%, 70%, 75%, 80%, 85%, 90%, preferably 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% or greater amino acid sequence identity, preferably over  
30 a region of at least about 50, 75, 100, 150, 200, 250, or 281 amino acids, to a human  $\alpha 4\beta 7$  integrin sequence shown below; (2) bind to antibodies, *e.g.*, polyclonal antibodies, raised against an immunogen comprising an amino acid sequence shown below, or conservatively



modified variants thereof; (3) bind to an  $\alpha 4\beta 7$  integrin binding protein; (4) specifically hybridize under stringent hybridization conditions to a nucleic acid sequence shown below, or conservatively modified variants thereof; (5) have a nucleic acid sequence that has greater than about 90%, preferably greater than about 96%, 97%, 98%, 99%, or higher nucleotide sequence identity, preferably over a region of at least about 100, 200, 300, 400 or more nucleotides, to a human  $\alpha 4\beta 7$  integrin mRNA sequence; and/or (6) have at least 25, often 50, 75, 100, 125 or 143 contiguous amino acid residues of a human  $\alpha 4\beta 7$  integrin polypeptide sequence. The human  $\alpha 4\beta 7$  integrin polypeptide sequence is set forth in, *e.g.*, Genbank Accession Nos. NP\_000876 and NP\_000880. The human  $\alpha 4\beta 7$  integrin mRNA (coding) sequence is set forth in, *e.g.*, Genbank Accession Nos. NM\_000885 and NM\_000889. One skilled in the art will appreciate that variants, isoforms, alternative sequences of  $\alpha 4\beta 7$  integrin are also useful in the present invention.

### 11. Hemoglobin (Hgb)

[0125] The term “hemoglobin” or “Hb” or “Hgb” refers to isolated nucleic acids, polypeptides and polymorphic variants, alleles, mutants, and interspecies homologues thereof and as further described herein, that: (1) have an amino acid sequence that has greater than about 60% amino acid sequence identity, 65%, 70%, 75%, 80%, 85%, 90%, preferably 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% or greater amino acid sequence identity, preferably over a region of at least about 50, 75, 100, 150, 200, 250, or 281 amino acids, to a human Hb sequence shown below; (2) bind to antibodies, *e.g.*, polyclonal antibodies, raised against an Hb immunogen comprising an amino acid sequence shown below, or conservatively modified variants thereof; (3) bind to an Hb binding protein; (4) specifically hybridize under stringent hybridization conditions to a nucleic acid sequence shown below, or conservatively modified variants thereof; (5) have a nucleic acid sequence that has greater than about 90%, preferably greater than about 96%, 97%, 98%, 99%, or higher nucleotide sequence identity, preferably over a region of at least about 100, 200, 300, 400 or more nucleotides, to human Hb subunits alpha, beta, delta and/or gamma RNA sequences; and/or (6) have at least 25, often 50, 75, 100, 125 or 143 contiguous amino acid residues of human Hb subunits alpha, beta, delta and/or gamma polypeptide sequences.

[0126] The human hemoglobin alpha polypeptide sequence is set forth in, *e.g.*, Genbank Accession No. NP\_000558. The human hemoglobin alpha mRNA (coding) sequence is set forth in, *e.g.*, Genbank Accession No. NM\_000549. The human hemoglobin beta

polypeptide sequence is set forth in, *e.g.*, Genbank Accession No. NP\_000509. The human hemoglobin beta mRNA (coding) sequence is set forth in, *e.g.*, Genbank Accession No. NM\_000518. The human hemoglobin delta polypeptide sequence is set forth in, *e.g.*, Genbank Accession No. NP\_000510. The human hemoglobin delta mRNA (coding) sequence is set forth in, *e.g.*, Genbank Accession No. NM\_000519. The human hemoglobin gamma polypeptide sequence is set forth in, *e.g.*, Genbank Accession No. NP\_000550 and NP\_000175. The human hemoglobin gamma mRNA (coding) sequence is set forth in, *e.g.*, Genbank Accession No. NM\_000559 and NM\_000184. One skilled in the art will appreciate that variants, isoforms, alternative sequences of the hemoglobin alpha, beta, delta and gamma subunits are also useful in the present invention.

## 12. Matrix Metalloproteinase 9 (MMP9)

[0127] The term “MMP9” or “matrix metalloproteinase 9” refers to isolated nucleic acids, polypeptides and polymorphic variants, alleles, mutants, and interspecies homologues thereof and as further described herein, that: (1) have an amino acid sequence that has greater than about 60% amino acid sequence identity, 65%, 70%, 75%, 80%, 85%, 90%, preferably 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% or greater amino acid sequence identity, preferably over a region of at least about 50, 75, 100, 150, 200, 250, 300, 400, 500, 600, 700, or 707 amino acids to a human MMP9 sequence shown below; (2) bind to antibodies, *e.g.*, polyclonal antibodies, raised against an MMP9 immunogen comprising an amino acid sequence shown below, or conservatively modified variants thereof; (3) bind to an MMP9 binding protein; (4) specifically hybridize under stringent hybridization conditions to a nucleic acid sequence shown below, or conservatively modified variants thereof; (5) have a nucleic acid sequence that has greater than about 90%, preferably greater than about 96%, 97%, 98%, 99%, or higher nucleotide sequence identity, preferably over a region of at least about 100, 200, 300, 400, 500, 1000, 1500, 2000, 2300, or more nucleotides, to a human MMP9 mRNA sequence; and/or (6) have at least 25, often 50, 75, 100, 125, 143 or more contiguous amino acid residues of a human MMP9 polypeptide sequences.

[0128] The human MMP9 polypeptide sequence is set forth in, *e.g.*, Genbank Accession No. NP\_004985. The human MMP9 mRNA (coding) sequence is set forth in, *e.g.*, Genbank Accession No. NM\_004994. One skilled in the art will appreciate that variants, isoforms, alternative sequences of MMP9 are also useful in the present invention.

### 13. IL-12p40

[0129] The term “IL-12p40” or “IL-12 subunit p40” refers to isolated nucleic acids, polypeptides and polymorphic variants, alleles, mutants, and interspecies homologues thereof and as further described herein, that: (1) have an amino acid sequence that has greater than  
5 about 60% amino acid sequence identity, 65%, 70%, 75%, 80%, 85%, 90%, preferably 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% or greater amino acid sequence identity, preferably over a region of at least about 50, 75, 100, 150, 200, 250, 300, or 328 amino acids to a human IL-12p40 sequence shown below; (2) bind to antibodies, *e.g.*, polyclonal antibodies, raised against an IL-12p40 immunogen comprising an amino acid sequence  
10 shown below, or conservatively modified variants thereof; (3) bind to an IL-12p40 binding protein; (4) specifically hybridize under stringent hybridization conditions to a nucleic acid sequence shown below, or conservatively modified variants thereof; (5) have a nucleic acid sequence that has greater than about 90%, preferably greater than about 96%, 97%, 98%, 99%, or higher nucleotide sequence identity, preferably over a region of at least about 100,  
15 200, 300, 400, 500, 600, 1000, 2000, 2300, or more nucleotides, to a human IL-12p40 mRNA sequence; and/or (6) have at least 25, often 50, 75, 100, 125 or 143 contiguous amino acid residues of a human IL-12p40 polypeptide sequence.

[0130] The human IL-12p40 polypeptide sequence is set forth in, *e.g.*, Genbank Accession No. NP\_002178. The human IL-12p40 mRNA (coding) sequence is set forth in, *e.g.*,  
20 Genbank Accession No. NM\_002187. One skilled in the art will appreciate that variants, isoforms, alternative sequences of IL-12p40 are also useful in the present invention.

### 14. Predicting Vedolizumab Response or Non-Response

[0131] In some embodiments, the presence of a higher level of TNF $\alpha$  in patient serum at baseline or at week 2 of vedolizumab treatment, compared to a cut-off value indicates that the  
25 patient is likely to have lower vedolizumab levels at week 6 compared to a cut-off value. In other embodiments, if a patient has a higher level of VEGF at baseline or at week 2 compared to a cut-off value, then the patient is likely to have a lower vedolizumab level at week 6 compared to a cut-off value. In yet other embodiments, if a patient has a higher level of ANG-2 at baseline or at week 2 compared to a cut-off value, then the patient is likely to have  
30 a lower vedolizumab level at week 6 compared to a cut-off value. In still other embodiments, if a patient has a higher level of CRP at baseline or at week 2 compared to a cut-off value, then the patient is likely to have a lower vedolizumab level at week 6 compared to a cut-off

value. In some embodiments, if a patient has a higher level of VCAM-1 at baseline or at week 2 compared to a cut-off value, then the patient is likely to have a lower vedolizumab level at week 6 compared to a cut-off value. In some instances, the presence of a lower vedolizumab level at week 6 indicates that the patient has autoantibodies against

5 vedolizumab. The cut-off value for a specific predictive marker can be established from a reference population of subjects who are likely to have higher vedolizumab levels at week 6. In some cases, the reference population includes subject who had a clinical response (*e.g.*, a greater than or equal to 70-point decrease in CDAI score for CD; a reduction in the Mayo Clinical score of at least 3 points and a decrease of at least 30% from the baseline score, with  
10 a decrease of at least 1 point on the rectal bleeding score of 0 or 1 for UC) to vedolizumab. The lower level of vedolizumab may be relative or in comparison to the level of vedolizumab in a sample from a patient who has had a clinical response, is in clinical remission or does not have autoantibodies against vedolizumab. In some embodiments, the lower level of serum vedolizumab is less than about 33-34 µg/mL for a subject with CD or UC.

15 **[0132]** In some embodiments, the presence of a higher level of ANG-1 in patient serum at baseline compared to a cut-off value indicates that the patient is likely to have a higher vedolizumab level at week 6 compared to a cut-off value. In some embodiments, if a patient has higher ADA levels at baseline or at week 2 of vedolizumab treatment compared to a cut-off value, the patient has higher levels of vedolizumab compared to a cut-off value. In other  
20 embodiments, if the patient has a higher serum albumin level at week 2 compared to a cut-off value, the patient is likely to have a higher level of vedolizumab compared to a cut-off value. In another embodiment, if a patient has higher IL-12p40 levels at baseline or at week 2 of vedolizumab treatment compared to a cut-off value, the patient has higher levels of vedolizumab compared to a cut-off value. In yet other embodiments, if the patient has a  
25 higher MMP9 level at baseline or week 2 compared to a cut-off value, the patient is likely to have a higher level of vedolizumab compared to a cut-off value. In some embodiments, if the patient has a higher ICAM-1 level at baseline or week 2 compared to a cut-off value, the patient is likely to have a higher level of vedolizumab compared to a cut-off value. In one embodiment, if the patient has a higher serum  $\alpha$ 4 $\beta$ 7 level at baseline or week 2 compared to a  
30 cut-off value, the patient is likely to have a higher level of vedolizumab compared to a cut-off value. In some instances, the presence of a higher vedolizumab level at week 6 indicates that the patient does not have autoantibodies against vedolizumab. In some cases, the cut-off value for a specific predictive marker is determined using a reference population of subjects.

In some cases, the reference population includes subjects who have not had a clinical response (*e.g.*, a greater than or equal to 70–point decrease in CDAI score; a reduction in the Mayo Clinical score of at least 3 points and a decrease of at least 30% from the baseline score, with a decrease of at least 1 point on the rectal bleeding score of 0 or 1 for UC) to vedolizumab. The higher level of vedolizumab may be relative to the level of vedolizumab in a sample from a patient who has autoantibodies against vedolizumab. In some embodiments, the higher level of serum vedolizumab is about 33-34  $\mu\text{g/mL}$  or greater for a subject with CD or UC.

**[0133]** In some embodiments, if the patient has a lower MadCAM-1 level at baseline or week 2 compared to a cut-off value, the patient is likely to have a higher level of vedolizumab compared to a cut-off value. In other embodiments, if the patient has a lower  $\text{TNF}\alpha$  level at baseline or week 2 compared to a cut-off value, the patient is likely to have a higher level of vedolizumab compared to a cut-off value. In some cases, the reference population includes subjects who have not had a clinical response (*e.g.*, a greater than or equal to 70–point decrease in CDAI score for CD; a reduction in the Mayo Clinical score of at least 3 points and a decrease of at least 30% from the baseline score, with a decrease of at least 1 point on the rectal bleeding score of 0 or 1 for UC) to vedolizumab. The higher level of vedolizumab may be relative to the level of vedolizumab in a sample from a patient who has autoantibodies against vedolizumab. In some embodiments, the higher level of serum vedolizumab is about 33-34  $\mu\text{g/mL}$  or greater for a subject with CD or UC.

**[0134]** In some embodiments, there is a positive association (correlation) between the level of vedolizumab at week 6 and the level of IL-12p40 at baseline. In some embodiments, there is a positive association (correlation) between the level of vedolizumab at week 6 and the level of MMP9 at week 2. In other embodiments, CRP levels at baseline are not associated with vedolizumab levels at week 6 of drug therapy.

**[0135]** In some embodiments for patients who are likely to have a clinical response to vedolizumab, for example, at week 6 of treatment, the level of serum  $\alpha 4\beta 7$  integrin increases and the level of MadCAM-1 decreases during the course of vedolizumab treatment. For instance, a CD or UC patient can undergo clinical response at week 6 after initiating a vedolizumab induction treatment regimen if the level of serum  $\alpha 4\beta 7$  integrin increases and the level of MadCAM-1 decreases compared to baseline.

[0136] In some embodiments for patients (*e.g.*, CD or UC patients) who are likely to undergo clinical remission, for example, at week 6 or week 52 of treatment, the level of serum  $\alpha 4\beta 7$  integrin increases and the level of MadCAM-1 decreases during therapy such as from a first time point to a later time point. In some cases, a CD or UC patient can undergo clinical remission upon receiving a vedolizumab maintenance treatment regimen if the patient has an increase in serum  $\alpha 4\beta 7$  integrin levels and a decrease in MadCAM-1 levels relative to baseline levels.

[0137] In some embodiments, if a patient has a high TNF $\alpha$ , VEGF, ANG-2, CRP, and/or VCAM-1 level(s) at baseline, the patient is likely to develop autoantibodies against vedolizumab. Those with a high TNF $\alpha$  level during induction therapy are predicted to develop autoantibodies. For instance, a patient with a high baseline level of TNF $\alpha$  is likely to have autoantibodies against vedolizumab at week 2 and week 6.

[0138] Patients with low level(s) of MadCAM-1, VCAM-1, and/or TNF $\alpha$  are not predicted to develop such autoantibodies. In other embodiments, patients with a high level(s) of ANG-1, ADA, HSA, IL-12p40, MMP9, ICAM-1 and/or serum  $\alpha 4\beta 7$  integrin at baseline are not likely to develop autoantibodies to an anti- $\alpha 4\beta 7$  integrin drug. For instance, patients with a higher HSA level are predicted to not develop autoantibodies against vedolizumab at week 6 or week 14.

[0139] A patient with autoantibodies against an anti- $\alpha 4\beta 7$  integrin drug may have had higher levels of TNF $\alpha$ , VEGF, ANG-2, CRP, VCAM-1 or any combination thereof at baseline or week 2 of therapy, compared to a patient who does not have the autoantibodies. A patient who does not have the autoantibodies may have had a lower level of MadCAM-1, VCAM-1, and TNF $\alpha$ , and higher levels ANG-1, ADA, HSA, IL-12p40, MMP9, ICAM-1, serum  $\alpha 4\beta 7$  integrin, or any combination thereof at baseline or week 2, compared to a patient with the autoantibodies.

[0140] Provided herein are methods for treating a subject with CD or UC with an anti- $\alpha 4\beta 7$  integrin drug such as vedolizumab. The method includes administering the drug to a subject having CD or UC and having a level of ANG-1, ADA, HSA, IL-12p40, MMP9, ICAM-1 and/or serum  $\alpha 4\beta 7$  integrin that is higher than a corresponding reference value.

Alternatively, the method may include administering the drug to a subject having CD or UC and having a level of MadCAM-1, VCAM-1 or TNF $\alpha$  that is lower than a corresponding reference value. In some cases, the UC subject administered vedolizumab may undergo

mucosal healing. In other cases, the CD or UC subject receiving the drug may be withdrawn from steroid treatment or undergo glucocorticoid-free remission (*e.g.*, clinical remission without glucocorticoid therapy).

### C. Setting Cut-off Values

5 [0141] Once the sample(s) from the human subject have been assayed for the criteria listed above, a value is generated for predicting likelihood of clinical response (*e.g.*, as defined by the Physician's Global Assessment for CD or UC, the Crohn's Disease Activity Index, the Mayo Clinic Score, or any other standard assessment criteria or scale for IBS) to vedolizumab or likelihood of having clinical remission (*e.g.*, as defined by the Physician's  
10 Global Assessment for CD or UC, the Crohn's Disease Activity Index, the Mayo Clinic Score, or any other standard assessment criteria or scale for IBS). When two or more predictive markers or other criteria are used in the method described herein, the level of each marker can be weighted and combined. Thus, a test value may be provided by (a) weighting the determined level of each marker with a predefined coefficient, and (b) combining the  
15 weighted level to provide a test value. The combining step can be either by straight addition or averaging (*i.e.*, weighted equally) or by a different predefined coefficient.

[0142] Once generated, the value from a sample can be compared to one or more cut-off or threshold value(s) to provide a likelihood of clinical response or clinical remission. In order to establish a cut-off value for practicing the method, a reference population of subjects can  
20 be used. In some embodiments, a population of patients with CD or UC can be used. In some instances, the patients have had a clinical response to vedolizumab. In other instances, the patients have not had a clinical response to vedolizumab or have autoantibodies against vedolizumab. Alternatively, the patients may have a clinical response to the anti- $\alpha 4\beta 7$  integrin drug. In some embodiments, the patients are in clinical remission from CD or in  
25 clinical remission from UC.

[0143] In some embodiments, these patients are within the appropriate parameters, if applicable, for the purpose of screening for and/or monitoring CD or UC using the methods of the present disclosure. Optionally, the patients are of similar age or similar ethnic background. The status of the selected patients can be confirmed by well established,  
30 routinely employed methods including but not limited to general physical examination of the individuals and general review of their medical history. Furthermore, the selected group of patients will generally be of sufficient size, such that the average value in the sample obtained

from the group can be reasonably regarded as representative of a particular indication, for example indicative of reoccurrence of CD or UC or not after a set period of time (*e.g.*, 2 years) after treatment.

[0144] Once an average value is established based on the individual values found in each subject of the selected group, this average or median or representative value or profile can be used as a cut-off value. For example, a sample value over the cut-off value can indicate a more than average likelihood of clinical response or clinical remission depending on the predictive marker used. In some embodiments, a standard deviation is also determined during the same process. In some cases, separate cut-off values may be established for separately defined groups having distinct characteristics such as age, gender, or ethnic background.

[0145] According to the methods described herein, the sample is compared to one or more reference or threshold values. In some embodiments, the sample value is deemed “high” if it is at least 1, 2, 3, 4, 5, 10, 15, 20 or more standard deviations greater than the reference value subjects. In other embodiments, the sample value is below the threshold if the sample value is at least 1, 2, 3, 4, 5, 10, 15, 20 or more standard deviations lower than the reference or threshold value.

[0146] In some embodiments, a computer-based analysis program is used to translate the raw data generated by the detection methods described herein (*e.g.*, the presence, absence, or amount of a given marker or markers) into a score of predictive value to a clinician.

[0147] The predictive marker profile or score, as determined according to the methods above, can predict that the patient has an above-average likelihood of clinical response or remission. In some cases, the patient has a high likelihood of clinical response or remission. The score can also predict that the patient has an average or below-average likelihood of clinical response or remission. In such instances, the patient can have a low or intermediate likelihood of clinical response or remission.

#### **D. Markers for Measuring Anti- $\alpha 4\beta 7$ Integrin drug and Anti-Drug Antibody (ADA) Levels**

[0148] In some embodiments, the method comprises determining the presence and/or level of anti- $\alpha 4\beta 7$  integrin drug (*e.g.*, level of free anti- $\alpha 4\beta 7$  integrin therapeutic antibody such as vedolizumab) and/or anti-drug antibody (ADA) (*e.g.*, level of autoantibody to the anti- $\alpha 4\beta 7$



integrin drug such as HAHA) in a patient sample (*e.g.*, a serum sample from a patient on anti- $\alpha 4\beta 7$  integrin drug therapy) at multiple time points, *e.g.*, before, during, and/or after the course of therapy.

5 [0149] In some embodiments, the presence and/or level of anti- $\alpha 4\beta 7$  integrin drug and/or ADA is determined with a homogeneous mobility shift assay (HMSA) using size exclusion chromatography. These methods are described in U.S. Patent No. 8,574,855, and 8,865,417 and U.S. Patent Publication Nos. 2014/0051184 and 2014/0141983, the disclosures of which are hereby incorporated by reference in their entirety for all purposes. The methods are particularly useful for measuring the presence or level of  $\alpha 4\beta 7$  integrin inhibitors as well as  
10 autoantibodies (*e.g.*, HACA, HAHA, *etc.*) that are generated against them.

[0150] In other embodiments, the presence and/or level of anti- $\alpha 4\beta 7$  integrin drug and/or ATV is determined with an immunoassay, such as an enzyme-linked immunosorbent assay (ELISA). In yet other embodiments, the presence and/or level of anti- $\alpha 4\beta 7$  integrin drug and/or ATV is determined with a flow cytometry assay such as FACS.

#### 15 E. Statistical Analysis

[0151] In some aspects, the present invention provides methods for selecting anti- $\alpha 4\beta 7$  integrin drug therapy, optimizing anti- $\alpha 4\beta 7$  integrin drug therapy, reducing toxicity associated with anti- $\alpha 4\beta 7$  integrin drug therapy, and/or monitoring the efficacy of anti- $\alpha 4\beta 7$  integrin drug treatment by applying one or more statistical algorithm to one or more (*e.g.*, a  
20 combination of two, three, four, five, six, seven, eight, nine, ten, eleven, twelve or more) pharmacodynamic and/or predictive markers. In particular embodiments, quantile analysis is applied to the presence and/or level of one or more markers to guide treatment decisions for patients receiving anti- $\alpha 4\beta 7$  integrin drug therapy. In other embodiments, one or a  
25 combination of two of more learning statistical classifier systems are applied to the presence and/or level of one or more markers to guide treatment decisions for patients receiving anti- $\alpha 4\beta 7$  integrin drug therapy. The statistical analyses of the methods of the present invention advantageously assist in determining when or how to adjust or modify (*e.g.*, increase or decrease) the subsequent dose of an anti- $\alpha 4\beta 7$  integrin drug, to combine an anti- $\alpha 4\beta 7$  integrin drug (*e.g.*, at an increased, decreased, or same dose) with one or more immunosuppressive  
30 agents such as methotrexate (MTX) or azathioprine (AZA), and/or to change the current course of therapy (*e.g.*, switch to a different anti- $\alpha 4\beta 7$  integrin drug).

**[0152]** The term “statistical analysis” or “statistical algorithm” or “statistical process” includes any of a variety of statistical methods and models used to determine relationships between variables. In the present invention, the variables are the presence or level of at least one marker of interest. Any number of markers can be analyzed using a statistical analysis described herein. For example, the presence or level of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 35, 40, 45, 50, 55, 60, or more markers can be included in a statistical analysis. In one embodiment, logistic regression is used. In another embodiment, linear regression is used. In yet another embodiment, ordinary least squares regression or unconditional logistic regression is used. In certain preferred embodiments, the statistical analyses of the present invention comprise a quantile measurement of one or more markers, *e.g.*, within a given population, as a variable. Quantiles are a set of “cut points” that divide a sample of data into groups containing (as far as possible) equal numbers of observations. For example, quartiles are values that divide a sample of data into four groups containing (as far as possible) equal numbers of observations. The lower quartile is the data value a quarter way up through the ordered data set; the upper quartile is the data value a quarter way down through the ordered data set. Quintiles are values that divide a sample of data into five groups containing (as far as possible) equal numbers of observations. The present invention can also include the use of percentile ranges of marker levels (*e.g.*, tertiles, quartile, quintiles, *etc.*), or their cumulative indices (*e.g.*, quartile sums of marker levels to obtain quartile sum scores (QSS), *etc.*) as variables in the statistical analyses (just as with continuous variables).

**[0153]** In certain embodiments, the present invention involves detecting or determining the presence and/or level (*e.g.*, magnitude) of one or more markers of interest using quartile analysis. In this type of statistical analysis, the level of a marker of interest is defined as being in the first quartile (<25%), second quartile (25-50%), third quartile (51%-<75%), or fourth quartile (75-100%) in relation to a reference database of samples. These quartiles may be assigned a quartile score of 1, 2, 3, and 4, respectively. In certain instances, a marker that is not detected in a sample is assigned a quartile score of 0 or 1, while a marker that is detected (*e.g.*, present) in a sample (*e.g.*, sample is positive for the marker) is assigned a quartile score of 4. In some embodiments, quartile 1 represents samples with the lowest marker levels, while quartile 4 represent samples with the highest marker levels. The reference database of samples can include a large spectrum of patients with a TNF $\alpha$ -mediated disease or disorder such as, *e.g.*, IBD. From such a database, quartile cut-offs can be

established. A non-limiting example of quartile analysis suitable for use in the present invention is described in, *e.g.*, Mow *et al.*, *Gastroenterology*, 126:414-24 (2004).

**[0154]** In some embodiments, the statistical analyses of the present invention comprise one or more learning statistical classifier systems. As used herein, the term “learning statistical classifier system” includes a machine learning algorithmic technique capable of adapting to complex data sets (*e.g.*, panel of markers of interest) and making decisions based upon such data sets. In some embodiments, a single learning statistical classifier system such as a decision/classification tree (*e.g.*, random forest (RF) or classification and regression tree (C&RT)) is used. In other embodiments, a combination of 2, 3, 4, 5, 6, 7, 8, 9, 10, or more learning statistical classifier systems are used, preferably in tandem. Examples of learning statistical classifier systems include, but are not limited to, those using inductive learning (*e.g.*, decision/classification trees such as random forests, classification and regression trees (C&RT), boosted trees, *etc.*), Probably Approximately Correct (PAC) learning, connectionist learning (*e.g.*, neural networks (NN), artificial neural networks (ANN), neuro fuzzy networks (NFN), network structures, the Cox Proportional-Hazards Model (CPHM), perceptrons such as multi-layer perceptrons, multi-layer feed-forward networks, applications of neural networks, Bayesian learning in belief networks, *etc.*), reinforcement learning (*e.g.*, passive learning in a known environment such as naïve learning, adaptive dynamic learning, and temporal difference learning, passive learning in an unknown environment, active learning in an unknown environment, learning action-value functions, applications of reinforcement learning, *etc.*), and genetic algorithms and evolutionary programming. Other learning statistical classifier systems include support vector machines (*e.g.*, Kernel methods), multivariate adaptive regression splines (MARS), Levenberg-Marquardt algorithms, Gauss-Newton algorithms, mixtures of Gaussians, gradient descent algorithms, and learning vector quantization (LVQ).

**[0155]** Random forests are learning statistical classifier systems that are constructed using an algorithm developed by Leo Breiman and Adele Cutler. Random forests use a large number of individual decision trees and decide the class by choosing the mode (*i.e.*, most frequently occurring) of the classes as determined by the individual trees. Random forest analysis can be performed, *e.g.*, using the RandomForest software available from Salford Systems (San Diego, CA). *See, e.g.*, Breiman, *Machine Learning*, 45:5-32 (2001); and [http://stat-www.berkeley.edu/users/breiman/RandomForests/cc\\_home.htm](http://stat-www.berkeley.edu/users/breiman/RandomForests/cc_home.htm), for a description of random forests.

[0156] Classification and regression trees represent a computer intensive alternative to fitting classical regression models and are typically used to determine the best possible model for a categorical or continuous response of interest based upon one or more predictors.

Classification and regression tree analysis can be performed, *e.g.*, using the C&RT software available from Salford Systems or the Statistica data analysis software available from StatSoft, Inc. (Tulsa, OK). A description of classification and regression trees is found, *e.g.*, in Breiman *et al.* "Classification and Regression Trees," Chapman and Hall, New York (1984); and Steinberg *et al.*, "CART: Tree-Structured Non-Parametric Data Analysis," Salford Systems, San Diego, (1995).

[0157] Neural networks are interconnected groups of artificial neurons that use a mathematical or computational model for information processing based on a connectionist approach to computation. Typically, neural networks are adaptive systems that change their structure based on external or internal information that flows through the network. Specific examples of neural networks include feed-forward neural networks such as perceptrons, single-layer perceptrons, multi-layer perceptrons, backpropagation networks, ADALINE networks, MADALINE networks, Learnmatrix networks, radial basis function (RBF) networks, and self-organizing maps or Kohonen self-organizing networks; recurrent neural networks such as simple recurrent networks and Hopfield networks; stochastic neural networks such as Boltzmann machines; modular neural networks such as committee of machines and associative neural networks; and other types of networks such as instantaneously trained neural networks, spiking neural networks, dynamic neural networks, and cascading neural networks. Neural network analysis can be performed, *e.g.*, using the Statistica data analysis software available from StatSoft, Inc. See, *e.g.*, Freeman *et al.*, In "Neural Networks: Algorithms, Applications and Programming Techniques," Addison-Wesley Publishing Company (1991); Zadeh, Information and Control, 8:338-353 (1965); Zadeh, "IEEE Trans. on Systems, Man and Cybernetics," 3:28-44 (1973); Gersho *et al.*, In "Vector Quantization and Signal Compression," Kluwer Academic Publishers, Boston, Dordrecht, London (1992); and Hassoun, "Fundamentals of Artificial Neural Networks," MIT Press, Cambridge, Massachusetts, London (1995), for a description of neural networks.

[0158] Support vector machines are a set of related supervised learning techniques used for classification and regression and are described, *e.g.*, in Cristianini *et al.*, "An Introduction to Support Vector Machines and Other Kernel-Based Learning Methods," Cambridge University Press (2000). Support vector machine analysis can be performed, *e.g.*, using the

SVM<sup>light</sup> software developed by Thorsten Joachims (Cornell University) or using the LIBSVM software developed by Chih-Chung Chang and Chih-Jen Lin (National Taiwan University).

[0159] The various statistical methods and models described herein can be trained and tested using a cohort of samples (*e.g.*, serological and/or genomic samples) from healthy individuals and patients with a TNF $\alpha$ -mediated disease or disorder such as, *e.g.*, IBD (*e.g.*, CD and/or UC) or rheumatoid arthritis. For example, samples from patients diagnosed by a physician, preferably by a gastroenterologist, as having IBD or a clinical subtype thereof using a biopsy, colonoscopy, or an immunoassay as described in, *e.g.*, U.S. Patent No. 6,218,129, are suitable for use in training and testing the statistical methods and models of the present invention. Samples from patients diagnosed with IBD can also be stratified into Crohn's disease or ulcerative colitis using an immunoassay as described in, *e.g.*, U.S. Patent Nos. 5,750,355 and 5,830,675. Samples from healthy individuals can include those that were not identified as IBD samples. One skilled in the art will know of additional techniques and diagnostic criteria for obtaining a cohort of patient samples that can be used in training and testing the statistical methods and models of the present invention.

#### F. Predictive Modeling

[0160] In certain aspects, the present invention provides pharmacokinetic models to predict the likelihood of developing anti-drug antibodies.

[0161] Pharmacokinetic models are ways to mathematically understand the fate of drugs in vivo. In a one compartment model, the drug-concentration time profile shows a monophasic response, and is described by a single exponential. In addition, the body is assumed to be a homogeneous unit with instantaneous distribution of the drug. A one-compartment model shows a linear relationship between log concentrations in plasma ( $C_p$ ) versus time.

[0162] A two-compartment model resolves the body into two units, a central unit and a peripheral unit. In the two-compartment model, the log concentration in plasma ( $C_p$ ) versus time profile is biphasic. In the biphasic model, there is a rapid decline in drug concentration followed by a slower decline. D. Ternant *et al.*, *Ther Drug Monit*, 30(4), 523-529 (2008), showed that infliximab pharmacokinetics followed a two compartment model, with an elimination half-life of close to 3 weeks.

[0163] In other aspects, the present invention provides an algorithmic model to predict patient response to anti- $\alpha 4\beta 7$  integrin drug. The model uses one or more markers such as an inflammatory marker which include cytokines and chemokines and the like, a signaling molecule, an acute phase protein, a cellular adhesion molecule and a combination thereof.

5 The markers also include the presence or absence of ADA, the levels of  $\alpha 4\beta 7$  integrin, the levels of MAdCAM-1, the concentration or levels of anti- $\alpha 4\beta 7$  integrin drugs and the like.

[0164] An algorithmic model includes any of a variety of statistical methods and models used to determine relationships between variables. In the present invention, the variables are the presence or level of at least one marker of interest. Any number of markers can be

10 analyzed using a statistical analysis described herein (*see*, “Statistical Analysis” section). For example, the presence or level of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, or more markers can be included in a statistical analysis.

[0165] In particular embodiments, quantile analysis is applied to the presence and/or level of one or more markers to guide treatment decisions for patients receiving anti- $\alpha 4\beta 7$  integrin drug therapy. In other embodiments, one or a combination of two or more learning statistical classifier systems are applied to the presence and/or level of one or more markers to guide treatment decisions for patients receiving anti- $\alpha 4\beta 7$  integrin drug therapy. The statistical analyses of the methods of the present invention advantageously assist in determining when or how to adjust or modify (*e.g.*, increase or decrease) the subsequent dose of an anti- $\alpha 4\beta 7$  integrin drug, to combine an anti- $\alpha 4\beta 7$  integrin drug (*e.g.*, at an increased, decreased, or same dose) with one or more immunosuppressive agents such as methotrexate (MTX) or azathioprine (AZA), and/or to change the current course of therapy (*e.g.*, switch to a different anti- $\alpha 4\beta 7$  integrin drug).

25 [0166] The algorithmic model includes the level or concentration of the one or more markers along with a statistic algorithm such as a learning statistical algorithm. In certain instances, the model has been trained with known outcomes using a training set cohort of samples. The algorithm is then validated using a validation cohort. Patient unknown samples can then be predicted based on the trained algorithms.

[0167] In some aspects, the present invention provides a system for predicting the level of an anti- $\alpha 4\beta 7$  integrin drug in a subject at a later time point during a course of therapy with the anti- $\alpha 4\beta 7$  integrin drug. In other aspects, the present invention provides a system for predicting whether a subject will develop autoantibodies to an anti- $\alpha 4\beta 7$  integrin drug at a

later time point during a course of therapy with the anti- $\alpha 4\beta 7$  integrin drug. In yet other aspects, the present invention provides a system for predicting a clinical outcome of a subject at a later time point during a course of therapy with the anti- $\alpha 4\beta 7$  integrin drug.

**[0168]** In certain embodiments, the system comprises: a data acquisition module

5 configured to produce a data set comprising one or more predictor variables for the subject determined at an earlier time point during the course of therapy and/or prior to the initiation of the course of therapy; a data processing module configured to process the data set by applying a statistical analysis to the data set to produce a statistically derived decision predicting the level of the anti- $\alpha 4\beta 7$  integrin drug or predicting whether the subject will  
10 develop autoantibodies to the anti- $\alpha 4\beta 7$  integrin drug or predicting a clinical outcome of the subject receiving the anti- $\alpha 4\beta 7$  integrin drug based upon the one or more predictor variables; and a display module configured to display the statistically derived decision.

**[0169]** In some embodiments, the system includes an intelligence module, such as a computer, having a processor and memory module. The intelligence module may also

15 include communication modules for transmitting and receiving information over one or more direct connections (*e.g.*, USB, Firewire, or other interface) and one or more network connections (*e.g.*, including a modem or other network interface device). The memory module may include internal memory devices and one or more external memory devices.

The intelligence module also includes a display module, such as a monitor, screen, or printer.

20 In one aspect, the intelligence module receives data such as patient test results from a data acquisition module such as a test system, either through a direct connection or over a network. For example, the test system may be configured to run multianalyte tests on one or more patient samples and automatically provide the test results to the intelligence module.

The data may also be provided to the intelligence module via direct input by a user or it may  
25 be downloaded from a portable medium such as a compact disk (CD) or a digital versatile disk (DVD). The test system may be integrated with the intelligence module, directly coupled to the intelligence module, or it may be remotely coupled with the intelligence module over the network. The intelligence module may also communicate data to and from one or more client systems over the network as is well known. For example, a requesting  
30 physician or healthcare provider may obtain and view a report from the intelligence module, which may be resident in a laboratory or hospital, using a client system.

[0170] The network can be a LAN (local area network), WAN (wide area network), wireless network, point-to-point network, star network, token ring network, hub network, or other configuration. As the most common type of network in current use is a TCP/IP (Transfer Control Protocol and Internet Protocol) network such as the global internetwork of networks often referred to as the "Internet" with a capital "I," that will be used in many of the examples herein, but it should be understood that the networks that the present invention might use are not so limited, although TCP/IP is the currently preferred protocol.

[0171] Several elements in the system may include conventional, well-known elements that need not be explained in detail here. For example, the intelligence module could be

implemented as a desktop personal computer, workstation, mainframe, laptop, *etc.* Each client system could include a desktop personal computer, workstation, laptop, cell phone, tablet, PDA, or any WAP-enabled device or any other computing device capable of interfacing directly or indirectly to the Internet or other network connection. A client system typically runs an HTTP client, *e.g.*, a browsing program, such as Microsoft's Internet Explorer™ browser, Google's Chrome™ browser, or a WAP-enabled browser or mobile application in the case of a cell phone, tablet, PDA, or other wireless device, or the like, allowing a user of the client system to access, process, and view information and pages available to it from the intelligence module over the network. Each client system also typically includes one or more user interface devices, such as a keyboard, a mouse, touch screen, pen, or the like, for interacting with a graphical user interface (GUI) provided by the browser on a display (*e.g.*, monitor screen, cell phone or tablet screen, LCD display, *etc.*) in conjunction with pages, forms, and other information provided by the intelligence module.

As discussed above, the present invention is suitable for use with the Internet, which refers to a specific global internetwork of networks. However, it should be understood that other networks can be used instead of the Internet, such as an intranet, an extranet, a virtual private network (VPN), a non-TCP/IP based network, any LAN or WAN, or the like.

[0172] According to one embodiment, each client system and all of its components are operator configurable using applications, such as a browser, including computer code run using a central processing unit such as an Intel® Pentium® processor or the like. Similarly, the intelligence module and all of its components might be operator configurable using application(s) including computer code run using a central processing unit such as an Intel® Pentium® processor or the like, or multiple processor units. Computer code for operating and configuring the intelligence module to process data and test results as described herein is



preferably downloaded and stored on a hard disk, but the entire program code, or portions thereof, may also be stored in any other volatile or non-volatile memory medium or device as is well known, such as a ROM or RAM, or provided on any other computer readable medium capable of storing program code, such as a compact disk (CD) medium, digital versatile disk (DVD) medium, a floppy disk, ROM, RAM, and the like.

[0173] The computer code for implementing various aspects and embodiments of the present invention can be implemented in any programming language that can be executed on a computer system such as, for example, in C, C++, C#, HTML, Java, JavaScript, or any other scripting language, such as VBScript. Additionally, the entire program code, or portions thereof, may be embodied as a carrier signal, which may be transmitted and downloaded from a software source (*e.g.*, server) over the Internet, or over any other conventional network connection as is well known (*e.g.*, extranet, VPN, LAN, *etc.*) using any communication medium and protocols (*e.g.*, TCP/IP, HTTP, HTTPS, Ethernet, *etc.*) as are well known.

### III. EXAMPLES

[0174] The present invention will be described in greater detail by way of specific examples. The following examples are offered for illustrative purposes, and are not intended to limit the present invention in any manner. Those of skill in the art will readily recognize a variety of noncritical parameters which can be changed or modified to yield essentially the same results.

#### Example 1: Methods for Predicting Response to Vedolizumab Based on the Presence or Level of One or More Biomarkers in a Patient Sample.

[0175] The aim of this example was to determine whether one or more markers can predict a response to therapy in IBD patients treated with vedolizumab. The markers tested included vedolizumab, antibodies against vedolizumab, TNF $\alpha$ , human serum albumin (HSA), VEGF, angiopoietin-1 (ANG-1), angiopoietin-2 (ANG-2), adenosine deaminase (ADA), serum  $\alpha 4\beta 7$  integrin ( $\alpha 4\beta 7$ ), matrix metalloproteinase 9 (MMP9), MAdCAM-1, hemoglobin (Hb), C-reactive protein (CRP), VCAM-1, and ICAM-1.

[0176] Sera were obtained from patients with IBD prior to the onset of vedolizumab therapy with (5 patients with ulcerative colitis and 7 patients with Crohn's disease) and after initiation of vedolizumab therapy (week 2 and 6). Concentrations of s-MAdCAM-1, TNF $\alpha$ , serum  $\alpha 4\beta 7$  integrin, vedolizumab, and antibodies to vedolizumab were measured by

Collaborative Enzyme Enhanced Reactive assay (CEER™), an ultrasensitive enzyme linked immunoassay, and by homogenous mobility shift assay (HMSA). *See, e.g., Wang et al., J Immunol Methods, 382:177-188 (2012), Kim et al., Proteome Sci; 9:75, 2011; Tao et al., Sci Signal, 7:rs29, 2014; and Elkabets et al., Sci Transl Med, 5:196ra99 (2013).* In addition, the markers human serum albumin (HSA), VEGF, angiopoietin-1 (ANG-1), antiopoietin-2 (ANG-2), adenosine deaminase (ADA), MMP9, hemoglobin (Hb), C-reactive protein (CRP), VCAM-1, ICAM-1, TWEAK, and IL-20p40 were also measured by ELISA or CEER™.

**[0177]** Results: The mean age of patients with IBD was 42.8 years old with a mean baseline albumin of 3.9 g/dL, hemoglobin of 12.2 g/dL, and CRP of 67 mg/L. The mean concentration of s-MAdCAM-1 was 25,843 pg/mL in patients with IBD prior to initiation of vedolizumab and decreased to 3288 pg/mL at week 6 after two induction doses of vedolizumab. Soluble  $\alpha 4\beta 7$  concentration increased from a mean of 1360 pg/mL at baseline to 3880 pg/mL at week 6. Mean vedolizumab concentrations were 38.3  $\mu$ g/mL at week 2 and 29.0  $\mu$ g/mL at week 6. There was no significant change in concentrations of CRP or TNF $\alpha$  during induction therapy with vedolizumab. Higher baseline albumin concentrations correlated with higher vedolizumab concentrations at week 6 (adjusted R-squared = 0.34); however, TNF $\alpha$  concentrations at baseline correlated with lower vedolizumab concentrations at week 6 (Adjusted R-square = 0.87). The correlation between specific marker pairs at baseline and at week 2 are provided in Table 1.

Table 1. Subject Characteristics

	N	Mean	Q1	Median	Q3
age	12	42.83333333333333	34.5	40	56.75
baseline alb	12	3.875	3.275	4	4.375
baseline hgb	12	12.23333333333333	11.225	11.7	13.175
CRP ng/mL baseline	9	6684.2222222222	1682	4024	12117
SAA ng/mL baseline	9	39822	2307.5	13891	80165.5
sICAM-1 ng/mL baseline	9	394.1111111111111	265	301	443
sVCAM-1 ng/mL baseline	9	733.888888888889	475	576	941
a4b7 pg/ml baseline	9	1360.1111111111111	176.5	346	999
Madcam pg/ml baseline	9	25843.33333333333	17765.5	19783	36125
ADA pg/ml baseline	9	1937.88888888889	784.5	1625	2257
TNF $\alpha$ pg/ml baseline	9	8.35555555555556	3.95	6.1	13.25
Ang1 baseline	9	44265.5555555556	34005	43250	53440
MMP9 baseline	9	1487488.8888889	829250	1311000	1972500
P40 baseline	9	1020.7055555556	328.375	437.6	1837.75

Tweak baseline	9	304.9	213.2	328	404.75
Ang2 baseline	9	7234.64444444444	902.3	1671	6907
VEGF pg/mL	9	294.782444444	96.57	147.284	509.87
Vedo ug/ml wk6	7	28.971428571429	11.31	26.33	51.92
CRP ng/mL wk6	7	5960.5714285714	482	4213	11962
SAA ng/mL wk6	7	19099.714285714	1635	12908	42020
sICAM-1 ng/mL wk6	7	350.14285714286	270	291	507
sVCAM-1 ng/mL wk6	7	592.85714285714	460	520	855
a4b7 pg/ml wk6	7	3880.4285714286	2696	3782	4195
Madcam pg/ml wk6	7	3288	2409	2890	3249
ADA pg/ml wk6	7	1109.7142857143	801	905	1174
TNFa pg/ml wk6	7	4.6142857142857	2.1	4.3	6.6
Ang1 wk6	7	38805.714285714	30520	37580	52100
MMP9 wk6	7	879657.14285714	589600	924400	1159000
P40 wk6	7	342.30714285714	188.3	215.35	480.3
Tweak wk6	7	288.12857142857	175.7	342.2	374.3
Ang2 wk6	7	3110.8571428571	638.7	3184	6416

[0178] Univariate regression analysis was performed to identify markers that can predict vedolizumab levels at week 6 of induction therapy. Baseline TNF $\alpha$  levels were higher in samples having lower levels of vedolizumab at week 6 (FIG. 1A). Also, TNF $\alpha$  levels at week 2 were higher in samples having lower vedolizumab levels at week 6 (FIG. 1B). VEGF levels at baseline were higher in samples having lower drug levels at week 6 (FIG. 2). Similarly, ANG-1 levels at baseline were higher in samples having higher vedolizumab levels at week 6 (FIG. 3B). Higher CRP levels at baseline were associated with lower drug levels at week 6 (FIG. 7B). ANG-2 levels at baseline were higher in samples having lower vedolizumab levels at week 6 (FIG. 3A). High ADA levels at baseline and at week 2 also resulted in higher drug levels at week 6 (FIGS. 4A and 4B). Higher human serum albumin levels at baseline were detected when vedolizumab levels were higher at week 6 (FIG. 5A). The data also showed a positive association between the anti- $\alpha$ 4 $\beta$ 7 integrin drug at week 6 and either IL-12p40 levels at baseline (FIG. 6A), MMP9 levels at week 2 (FIG. 6B) and MAdCAM-1 at week 2 (FIG. 7A). In other words, high levels of Il-12p40 at baseline correlated with high levels of vedolizumab at week 6. High levels of MMP9 at week 9 also correlated with high levels of drug at week 6. No such association was calculated for hemoglobin levels at baseline (FIG. 5B). Further analysis was performed to investigate the levels of  $\alpha$ 4 $\beta$ 7 integrin or MAdCAM-1 relative drug levels during the course of treatment. Vedolizumab therapy increased serum  $\alpha$ 4 $\beta$ 7 integrin (FIGS. 8A and 9) and decrease

MAdCAM-1 levels (FIGS. 8B and 10) over time. A summary of the correlations between the markers tested is provided in Table 2.

Table 2. Summary of Correlations

Marker 1	Marker 2	Correlation	P-value	Time point
ANG-1	TNF $\alpha$	-0.7183	0.0293	Baseline
IL-12p40	$\alpha$ 4 $\beta$ 7 integrin	0.7262	0.0267	Baseline
Tweak	CRP	0.6847	0.0418	Baseline
Tweak	TNF $\alpha$	0.7249	0.0271	Baseline
ANG-2	MMP9	-0.7528	0.0192	Baseline
CRP	Vedolizumab	-0.6849	0.0418	Wk 2
TNF $\alpha$	CRP	0.77	0.0152	Wk 2
MMP9	ANG-1	0.806	0.0087	Wk 2
MMP9	ICAM-1	-0.7039	0.03	Wk 2
MMP9	VCAM-1	-0.7926	0.01	Wk 2
ANG-1	ICAM-1	-0.7022	0.0349	Wk 2
ANG-1	VCAM-1	-0.6453	0.06 ( n.s.)	Wk 2
ANG-1	TNF $\alpha$	-0.6727	0.047	Wk 2

5 [0179] Using a multiple regression model it was determined that there is an interaction (correlation) between VCAM-1 and  $\alpha$ 4 $\beta$ 7 integrin levels at baseline and a prediction of vedolizumab levels at week 6 (FIG. 11A). Further, an interaction between VCAM-1 and MadCAM-1 at baseline and a vedolizumab levels at week 6 was also established (FIG. 11B). The data suggested that the presence or level of one or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) markers at baseline (a first time point) can be used to predict drug levels at a future time point (a second time point) during the course of therapy. The results showed that VCAM-1 and  $\alpha$ 4 $\beta$ 7 integrin levels at baseline can predict the presence of vedolizumab at week 6 and the absence of autoantibodies to the drug.

15 [0180] The results showed that MAdCAM-1 levels negatively correlated to ICAM-1 levels (FIG. 12A) and VCAM-1 levels (FIG. 12B).

[0181] Longitudinal analyses (*e.g.*, analysis at baseline, post- week 0, week 2 and week 6 of therapy) of marker levels and vedolizumab levels were performed. MAdCAM-1 levels decreased during the course of therapy at the initiation phase and induction phase (FIG. 13), and in the maintenance phase (FIG. 17). VCAM-1 levels did not change statistically significantly with time (FIG. 14). Serum  $\alpha$ 4 $\beta$ 7 integrin levels were low at baseline and

20

increased during induction therapy (FIG. 15) and remained high during maintenance therapy (FIG. 18). In these samples, vedolizumab levels were about the same during induction therapy (FIG. 16). The results of the analyses show that the patients did not develop autoantibodies against vedolizumab. The data shows that a decrease in MadCAM-1 levels and an increase in serum 4β7 integrin levels across induction therapy predicts that patient with CD or UC will likely have a clinical response to vedolizumab at week 6 of treatment. The patient is also likely to have clinical remission during maintenance therapy.

[0182] Further analysis revealed that there is a positive correlation between the levels of human serum albumin at baseline and levels of vedolizumab at week 6 (induction phase) and week 14 (maintenance phase) of treatment (FIGS. 19A and 19B). There was a negative correlation between the levels of TNFα at baseline and levels of vedolizumab at week 2 and week 6 of treatment (FIGS. 20A and 20B).

[0183] Table 3 provides data of the bivariate fit graphs that analyze vedolizumab vs. TNFα (FIGS. 1A and 1B), vedolizumab vs. VEGF (FIG. 2), vedolizumab vs. ANG-2 (FIG. 3A), and vedolizumab vs. ANG-1 (FIG. 3B).

Table 3. Summary of Bivariate Fit Graphs of Vedolizumab with TNFα, VEGF, ANG-2 and ANG-1

	log Vedo (ug/ml at wk 6) v. log TNFa (pg/ml at baseline)	log Vedo (ug/ml at wk 6) v. log TNFa (pg/ml at wk 2)	log Vedo (ug/ml at wk 6) v. log VEGF (pg/ml at baseline)	log Vedo (ug/ml at wk 6) v. log ANG-2 (pg/ml at baseline)	log Vedo (ug/ml at wk 6) v. log ANG-1 (pg/ml at baseline)
linear fit	log Vedo (ug/ml at wk 6) - 5.5242934 - 1.7970711*[ log TNFa (pg/ml at baseline)]	log Vedo (ug/ml at wk 6) - 4.3374543 - 1.8306673*[ log TNFa (pg/ml at wk 2)]	log Vedo (ug/ml at wk 6) = 10.277741 - 1.424572*[ log VEGF (pg/ml at baseline)]	log Vedo (ug/ml at wk 6) = 8.731798 - 0.9235765 * [log ANG-2 (pg/ml at baseline)]	log Vedo (ug/ml at wk 6) = 5.45025 + 5.179299 * [log ANG-1 (pg/ml at baseline)]
Rsquare	0.901884	0.882787	0.511011	0.501428	0.841118
Rsquare Adj	0.869178	0.843715	0.348015	0.252148	0.788158
Root Mean Square Error	0.600528	0.656373	1.340638	1.379359	0.764186
Mean of Response	2.398325	2.398325	2.398325	1.986356	2.398325
Observations (or Sum Weights)	5	5	5	4	5
Analysis of Variance (Model DF; Sum of Squares; Mean Square, F Ratio)	1; 9.944800; 9.94480; 27.5759	1; 9.734222; 9.73422; 22.5943	1; 5.634768; 5.63477; 3.1351	1; 3.8270572; 3.82706; 2.0115	1; 9.274757; 9.27476; 15.8820
Analysis of Variance (Error;	3; 1.081900; 0.36063; --	3; 1.292477; 0.43083; --	3; 5.391932; 1.79731; --	2; 3.8052615; 1.90263; --; --	3; 1.751943; 0.58398 ;--

DF; Sum of Squares; Mean Square, F Ratio)					
Analysis of Variance (C Total; DF; Sum of Squares; Mean Square, F Ratio)	4; 11.026700; --; --	4; 11.026700; --; --	4; 11.026700; --; --	3; 7.6323188; -; -	4; 11.026700; --; --
Prob > F	0.0134	0.0177	0.1748	0.2919	0.0283
Parameter Estimate (Intercept; estimate std error; t ratio; Prob>[t])	5.5242934; 0.653056; 8.46; 0.0035	4.3374543; 0.502582; 8.63; 0.0033	10.27741; 4.490284; 2.29; 0.1061	8.7831798; 4.841747; 1.81; 0.2113	-53.45025; 14.0181; -3.81; 0.0317
Parameter Estimate (X-value; estimate std error; t ratio; Prob>[t])	-1.797071, 0.342216, -5.25, 0.0134	-1830667; 0.385132; -4.75; 0.0177	-1.424572; 0.804559; -1.77; 0.1748	-0.923576; 0.6581205; -1.42; 0.299	5.179299; 1.299628; 3.99; 0.0283
Robust Fit (Sigma; chi square; P-value; LogWorth)	NA	NA	1.41317; 6.55162; 0.01018; 1.97968	1.45399; 423246; 0.3966; 1.40167	0.80553; 53.843; 2.2e-13; 12.6632
Parameter (Intercept; robust estimate; Std error)	NA	NA	10.2777; 2.62271	8.78318; 2.87003	-53.45; 7.42013
Parameter (X-value; robust estimate; Std error)			-1.4246; 0.55656	-0.9236; 0.44893	5.1793; 0.70584

[0184] Table 4 provides data of the bivariate fit graphs that analyze vedolizumab vs. ADA (FIGS. 4A and 4B), vedolizumab vs. albumin (FIG. 5A), and vedolizumab vs. hemoglobin (FIG. 5B).

5 Table 4. Summary of Bivariate Fit Graphs of Vedolizumab with ADA, Albumin and Hemoglobin

	log Vedo (ug/ml at wk 6) v. log ADA(pg/ml at baseline)	log Vedo (ug/ml at wk 6) v. log ADA(pg/ml at week 2)	log Vedo (ug/ml at wk 6) v. log albumin (pg/ml at baseline)	log Vedo (ug/ml at wk 6) v. log hemoglobin (pg/ml at baseline)
linear fit	log Vedo (ug/ml at wk 6) = -18.26969+2.8022482 log ADA(pg/ml at baseline)	log Vedo (ug/ml at wk 6) = -19.71982+3.0702953 * [log ADA(pg/ml at week 2)]	log Vedo (ug/ml at wk 6) = -3.112497+4.5289224 * [log albumin (pg/ml at baseline)]	log Vedo (ug/ml at wk 6) = 8.5923412-2.3740808 * [log hemoglobin (pg/ml at baseline)]
Rsquare	0.456554	0.799121	0.405153	0.12482
Rsquare Adj	0.275405	0.732161	0.286184	-0.18502
Root Mean Square	1.41332	0.85927	1.292116	1.664837

Error				
Mean of Response	2.398325	2.398325	2.811286	2.811286
Observations (or Sum Weights)	5	5	7	7
Analysis of Variance (Model DF; Sum of Squares; Mean Square, F Ratio)	1; 5.034280; 5.03428; 2.5203	1; 8.811665; 8.1166; 11.9343	1; 5.685742; 5.68574; 3.4055	1; 0.175160; 0.17516; 0.0632
Analysis of Variance (Error; DF; Sum of Squares; Mean Square, F Ratio)	3; 5.992419; 1.99747; --; --	3; 2.215035; 0.73835; --	5; 8.347823; 1.66956; --	5; 13.858404; 2.77168; --
Analysis of Variance (C Total; DF; Sum of Squares; Mean Square, F Ratio)	4; 11.026700; --; --	4; 11.026700; --; --	6; 14.033565; --; --	6; 14.033565; --; --
Prob > F	0.2106	0.0408	0.1243	0.8115
Parameter Estimate (Intercept; estimate std error; t ratio; Prob>[t])	-18.26969; 1303412; -1.40; 0.2556	-19.71982; 6.414018; -3.07; 0.0544	-3.112497; 3.246954; -0.96; 0.3818	8.5923412; 23.00505; 0.37; 0.7241
Parameter Estimate (X-value; estimate std error; t ratio; Prob>[t])	2.8022482; 1765137; 1.59; 0.106	3.0702953; 0.88753; 3.45; 0.0408	4.5289224; 2.45416; 1.85; 0.1243	-2.374081; 9.443849; -0.25; 0.8115
Robust Fit (Sigma; chi square; P-value; LogWorth)	1.48979; 508451; 0.02414; 1.61725	0.90576; 15.7548; 7.21e-5; 4.14204	1.36203; 5.51481; 0.01886; 1.72455	1.75491; 0.80918; 0.36836; 0.43372
Parameter (Intercept; robust estimate; Std error)	-18.27; 9.44979	-19.72; 5.58927	-3.1152; 2.88796	8.59234; 6.7798
Parameter (X-value; robust estimate; Std error)	2.80225; 1.24274	3.0703; 0.77352	4.52892; 1.92854	-2.3741; 2.63921

[0185] Table 5 provides data of the bivariate fit graphs that analyze vedolizumab vs. IL-12p40 (FIG. 6A), vedolizumab vs. MadCAM-1 at week 2 (FIG. 7A), vedolizumab vs. MMP9 at week 2 (FIG. 6B), MadCAM-1 vs. soluble ICAM-1 (FIG. 12A), and MadCAM-1 vs.

5 soluble VCAM-1 (FIG. 12B).

Table 5. Summary of Bivariate Fit Graphs of Vedolizumab with IL-12p40, MadCAM-1, and MMP9, and MadCAM-1 with ICAM-1 and VCAM-1

	log Vedo (ug/ml at wk 6) v. log IL-12p40 (pg/ml at baseline)	log Vedo (ug/ml at wk 6) v. log MMP9 (pg/ml at wk 2)	log Vedo (ug/ml at wk 6) v. log MadCAM-1 (pg/ml at wk 2)	log MadCAM-1 (pg/ml at baseline) v. log sICAM-1 (ng/ml at baseline)	log MadCAM-1 (pg/ml at baseline) v. log sVCAM-1 (ng/ml at baseline)
linear fit	log Vedo (ug/ml at wk 6) = -11.88877	log Vedo (ug/ml at wk 6)=-30.96904+2.3590	log Vedo (ug/ml at wk 6)=-8.529237+1.2524	log MadCAM-1 (pg/ml at baseline)=12.4669	log MadCAM-1 (pg/ml at baseline)=13.1332

	+2.3841137 *[log IL-12p40 (pg/ml at baseline)]	076*[log MMP9 (pg/ml at week2)]	517*[log MadCAM-1 (pg/ml at week 2)]	26-0.40506*[log sICAM-1 (ng/ml at baseline)]	67-0.4690828*[log sVCAM-1 (ng/ml at baseline)]
Rsquare	0.538091	0.637475	0.140385	0.208307	0.29805
Rsquare Adj	0.384122	0.516633	-0.14615	0.095208	0.197771
Root Mean Square Error	1.302988	1.154333	1.777519	0.376394	0.35442
Mean of Response	2.398325	2.398325	2.398325	10.08734	10.08734
Observations (or Sum Weights)	5	5	5	9	9
Analysis of Variance (Model DF; Sum of Squares; Mean Square, F Ratio)	1; 5.933370; 5.93337; 3.4948	1; 7.029242; 7.02924; 5.2753	1; 1.547979; 1.5498; 0.4899	1; 0.2609350; 0.260935; 1.8418	1; 0.3733504; 0.373350; 2.9722
Analysis of Variance (Error; DF; Sum of Squares; Mean Square, F Ratio)	3; 5.093330; 1.69778; --	3; 3.997457; 1.33249; --	; 9.478721; 3.15957; --	7; 0.9917089; 0.141673; --	7; 0.8792935; 0.125613; --
Analysis of Variance (C Total; DF; Sum of Squares; Mean Square, F Ratio)	4; 11.026700; --; --	4; 11.026700; --; --	4; 11.026700; --; --	8; 1.2526439; --; --	8; 1.2526439; --; --
Prob > F	0.1583	0.1053	0.5344	0.2169	0.1284
Parameter Estimate (Intercept; estimate std error; t ratio; Prob>[t])	-11.88877; 7.66465; -1.55; 0.2187	-30.96904; 14.53694; -2.13; 0.1230	-8.529237; 15.6321; -0.55; 0.6233	12.466926; 1.75787; 7.09; 0.0002	13.133267; 1.77071; 7.42; 0.0001
Parameter Estimate (X-value; estimate std error; t ratio; Prob>[t])	2.3841137 ;1.275312; 1.87; 0.1583	2.3590076; 1.027085; 2.30; 0.1503	1.2524517; 1.78934; 0.70; 0.5344	-0.40506; 0.298467; -1.36; 0.2169	-0.469083; 0.272088; -1.72; 0.1284
Robust Fit (Sigma; chi square; P-value; LogWorth)	1.37349; 7.84119; 0.00511; 2.29184	1.21679; 16.0954; 0.00006; 4.22018	1.87369; 2.94639; 0.08607; 1.06515	0.39676; 4.70911; 0.03; 1.52283	0.3736; 6.5995; 0.0102; 1.99137
Parameter (Intercept; robust estimate; Std error)	-11.889; 5.44922	-30.969; 8.31154	-8.5292; 6.89747	12.4669; 1.17294	13.1333; 1.27585
Parameter (X-value; robust estimate; Std error)	2.38411; 0.8514	2.35901; 0.588	1.25245; 0.72965	-0.4051; 0.18666	-0.4691; 0.1826



[0186] Table 6 provides data of the bivariate fit graphs that analyze vedolizumab vs. CRP (FIG. 7B), vedolizumab vs. albumin (FIGS. 19A and 19B), and vedolizumab vs. TNF $\alpha$  (FIGS. 20A and 20B).

Table 6. Summary of Bivariate Fit Graphs of Vedolizumab with CRP, Albumin and TNF $\alpha$ .

	log Vedo (ug/ml at wk 6) v. log CRP (ng/ml at baseline)	log Vedo (ug/ml at wk 6) v. log albumin (ng/ml at baseline)	log Vedo (ug/ml at wk 14) v. log albumin (ng/ml at baseline)	log Vedo (ug/ml at wk 2) v. log TNF $\alpha$ (pg/ml at baseline)	log Vedo (ug/ml at wk 6) v. log TNF $\alpha$ (pg/ml at baseline)
linear fit	log Vedo (ug/ml at week 6)=7.4766583-0.6358414*[log CRP (ng/ml at baseline)]	log Vedo (ug/ml at week 6)=-0.652687+0.9211269*[log albumin (ng/ml at baseline)]	log Vedo (ug/ml at week 14)=-1.769356+1.0637125*[log albumin (ng/ml at baseline)]	log Vedo (ug/ml at week 2)=4.3372739-0.6198702*[log TNF $\alpha$ (np/ml at baseline)]	log Vedo (ug/ml at week 6)=5.3935304-1.4873362*[ log TNF $\alpha$ (np/ml at baseline)]
Rsquare	0.404505	0.596686	0.40339	0.668319	0.636842
Rsquare Adj	0.206007	0.551873	0.3371	0.626859	0.596491
Root Mean Square Error	1.479453	0.397241	0.714957	0.325704	0.767368
Mean of Response	2.398325	3.140682	2.582196	3.176041	2.831056
Observations (or Sum Weights)	5	11	11	10	11
Analysis of Variance (Model DF; Sum of Squares; Mean Square, F Ratio)	1; 4.460357; 4.46036; 2.0378	1; 2.1011322; 2.10113; 13.3151	1; 3.1105533; 3.11055; 6.0852	1; 1.7100103; 1.71001; 16.1196	1; 9.293624; 9.29362; 15.7826
Analysis of Variance (Error; DF; Sum of Squares; Mean Square, F Ratio)	3; 6.566343; 2.18878; --	9; 1.4202036; 0.15780; --	9 ;4.6004780; 0.51116; --	8; 0.8486634; 0.10608; --	9; 5.299677; 0.58885; --
Analysis of Variance (C Total; DF; Sum of Squares; Mean Square, F Ratio)	4; 11.026700; --; --	10; 3.5213359; --; --	10; 7.7110313; --; --	9; 2.5586737; --; --	10 ;14.593301; --; --
Prob > F	0.2487	0.0053	0.0358	0.0039	0.0032
Parameter Estimate (Intercept; estimate std error; t ratio; Prob>[t])	7.466583; 3.618443; 2.07; 0.1307	-0.652687; 1.046444; -0.62; 0.5483	-1.769356; 1.777151; -1.00; 0.3455	4.3372739; 0.307021; 14.13; <0.0001	5.3935304; 0.685257; 7.87; <0.0001
Parameter Estimate (X-value; estimate std error; t ratio; Prob>[t])	-0.635841; 0.445415; -1.43; 0.2487	0.9211269; 0.252433; 3.65; 0.0053	1.0637125; 0.431207; 2.47; 0.358	-0.61987; 0.154392; -4.01; m0.0039	-1.487336; 0.374386; -3.97; 0.00076; 3.11798
Robust Fit (Sigma; chi square; P-value;	1.5595; 3.74565; 0.05295; 1.27617	0.41873; 14.5684; 0.00014; 3.86915	0.75364; 8.92485; 0.00281; 2.55081	0.33085; 18.2345; 0.00002; 4.70927	0.80889; 11.3313; 0.00076; 3.11798

LogWorth)					
Parameter (Intercept; robust estimate; Std error)	7.47555; 2.23707	-0.6527; 1.04094	-1.7694; 1.35522	4.32507; 0.39341	5.39353; 0.64127
Parameter (X-value; robust estimate; Std error)	-0.6358; 0.3254	0.92113; 0.24133	1.06371; 0.35606	-0.6157; 0.14419	-1.4873; 0.44184

[0187] Table 7 provides data of the multivariate regression analysis of vedolizumab levels at week 6 in association with either (a)  $\alpha\beta$  integrin at baseline and soluble VCAM-1 at baseline (FIG. 11A), or (b) soluble VCAM-1 at baseline and MadCAM-1 at baseline (FIG. 11B).

Table 7. Summary of Multivariate Regression Models with Interactions between VCAM-1 and  $\alpha\beta$  Integrin or VCAM-1 and MadCAM-1 for Predicting Vedolizumab Levels at Week 6

Log [Vedo ug/ml at week 6] vs. Log [Vedo ug/ml at week 6]	
Predicted P	0.0724
R Square	0.996763
R Square Adj	0.987054
Root Mean Square Error	0.188915
Mean of Response	2.398325
Observations (Sum Wgts)	5
Model (DF; Sum of Squares; Mean Square; F Ratio)	3; 10.991011; 3.66367; 102.6559
Error (DF; Sum of Squares; Mean Square; F Ratio)	1; 0.035689; 0.3569; --
C. Total (DF; Sum of Squares; Mean Square; F Ratio)	4; 11.026700; --; --
Prob > F	0.0724
Intercept (Estimate; Std Error; t ratio; Prob > [t])	17.211202; 1.145423; 15.03; 0.423
Log [sVCAM-1 ng/ml at baseline) (Estimate; Std Error; t ratio; Prob > [t])	-4.392047; 0.277274; -15.84; 0.0401
Log [a4b7 integrin pg/ml at baseline) (Estimate; Std Error; t ratio; Prob > [t])	0.2952687; 0.261342; 8.78; 0.722
Log [sVCAM-1 ng/ml at baseline]-6.51781)*(Log[a4b7 pg/ml at baseline]-5.55963) (Estimate; Std Error; t ratio; Prob > [t])	5.0931735; 0.354773; 14.36; 0.0443
Predicted P	0.1528
Rsquare	0.985532
Rsquare Adj	0.942127
Root Mean Square Error	0.399423

Mean of Response	2.398325
Observations (Sum Wgts)	5
Model (DF; Sum of Squares; Mean Square; F Ratio)	3; 10.867161; 3.62239; 22.7054
Error (DF; Sum of Squares; Mean Square; F Ratio)	1 ;0.159538; 0.15954; --
C. Total (DF; Sum of Squares; Mean Square; F Ratio)	4; 11.026700; --; --
Prob > F	0.1528
Intercept (Estimate; Std Error; t ratio; Prob > [t])	125.73107; 18.19193; 6.91; 0.0915
Log [sVCAM-1 ng/ml at baseline) (Estimate; Std Error; t ratio; Prob > [t])	-6.301264; 0.798176; -7.89; 0.0802
Log [MadCAM-1 pg/ml at baseline) (Estimate; Std Error; t ratio; Prob > [t])	-8.489875; 1.37403; -6.18; 0.1021
Log [sVCAM-1 ng/ml at baseline]-6.51781)*(Log[MadCAM-1 pg/ml at baseline]-9.92376) (Estimate; Std Error; t ratio; Prob > [t])	-19.3602; 2.753055; -7.03; 0.0899

**[0188]** Additional analysis of responders to vedolizumab and levels of  $\alpha 4\beta 7$  integrin showed that patients who had a clinical response as based on Physician’s Global Assessment (PGA) had a greater change or increase in  $\alpha 4\beta 7$  integrin levels compared to patient who were non-responders. FIG. 21 shows increases between levels at week 0 and week 2 (bars labeled “1”) at week 0 and week 6 (bars labeled “2”) and at week 0 and week 14 (bars labeled “3”). The differences in the increased levels between responders and non-responders were statistically significant when comparing those at week 0 to week 6 (“2”) and at week 0 to week 14 (“3”).

**[0189]** The correlation between clinical responders and increases in  $\alpha 4\beta 7$  integrin levels was also found when clinical response was defined as withdrawal from steroid therapy. FIG. 22 shows that clinical responders had a greater or larger increase in  $\alpha 4\beta 7$  integrin levels when changes between week 0 and week 14 were calculated. The difference between the responders and non-responders was statistically significant between the increased level at week 0 and the increased level at week 14 (p-value=0.0107). As such, s- $\alpha 4\beta 7$  integrin level increases in the presence of vedolizumab treatment appear to be high in clinical responders than non-responders, based on PGA and withdrawal from steroid therapy. Changes in baseline of s- $\alpha 4\beta 7$  integrin can serve as an effective biomarker for responsiveness to vedolizumab.

[0190] In summary, one or more markers, such as TNF $\alpha$ , human serum albumin (HSA), VEGF, angiopoietin-1 (ANG-1), angiopoietin-2 (ANG-2), adenosine deaminase (ADA),  $\alpha$ 4 $\beta$ 7 integrin ( $\alpha$ 4 $\beta$ 7), MAdCAM-1, hemoglobin (Hgb), C-reactive protein (CRP), matrix metalloproteinase 9 (MMP9), VCAM-1, and ICAM-1, can be used to predict response to therapy in patients treated with vedolizumab.

[0191] All publications and patent applications cited in this specification are herein incorporated by reference as if each individual publication or patent application were specifically and individually indicated to be incorporated by reference. Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be readily apparent to those of ordinary skill in the art in light of the teachings of this invention that certain changes and modifications may be made thereto without departing from the spirit or scope of the appended claims.

WHAT IS CLAIMED IS:

1                   **1.**       A method for predicting that a subject having inflammatory bowel  
2 disease (IBD) will have a clinical response to an anti- $\alpha 4\beta 7$  integrin drug treatment, the  
3 method comprising:

4                   (a) detecting the presence or level of at least one predictive marker selected  
5 from the group consisting of TNF $\alpha$ , human serum albumin (HSA), VEGF, angiopoietin-1  
6 (ANG-1), angiopoietin-2 (ANG-2), adenosine deaminase (ADA), serum  $\alpha 4\beta 7$  integrin, IL-  
7 12p40, C-reactive protein (CRP), matrix metalloproteinase 9 (MMP9), MAdCAM-1, VCAM-  
8 1, ICAM-1, and a combination thereof in a sample from the subject; and

9                   (b) classifying the subject as a responder or a non-responder to the anti- $\alpha 4\beta 7$   
10 integrin drug treatment according to a predictive marker profile based on a higher or lower  
11 level of the at least one predictive marker compared to a corresponding reference value.

1                   **2.**       The method of claim **1**, wherein the inflammatory bowel disease is  
2 ulcerative colitis (UC) or Crohn's Disease (CD).

1                   **3.**       The method of claim **1** or **2**, wherein classifying comprises applying a  
2 statistical analysis to said predictive marker profile to determine if the subject is a responder  
3 or a non-responder to an anti- $\alpha 4\beta 7$  integrin drug treatment.

1                   **4.**       The method of any one of claims **1** to **3**, wherein the at least one  
2 predictive marker comprises at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13 or more predictive  
3 markers.

1                   **5.**       The method of any one of claims **1** to **4**, wherein the subject is  
2 classified as a responder if the level of ANG-1, ADA, HSA, IL-12p40, MMP9, ICAM-1,  
3 and/or serum  $\alpha 4\beta 7$  integrin is higher than the corresponding reference value.

1                   **6.**       The method of any one of claims **1** to **4**, wherein the subject is  
2 classified as a responder if the level of MadCAM-1, VCAM-1, and/or TNF $\alpha$  is lower than the  
3 corresponding reference value.

1                   **7.**       The method of any one of claims **1** to **4**, wherein the subject is  
2 classified as a non-responder if the level of TNF $\alpha$ , VEGF, ANG-2, CRP, and/or VCAM-1 is  
3 higher than the corresponding reference value.

1           **8.**       The method of any one of claims **1** to **7**, wherein detecting the presence  
2 or level of the at least one predictive marker comprises performing a proximity dual detection  
3 assay or an immunoassay.

1           **9.**       The method of claim **8**, wherein the proximity dual detection assay is a  
2 Collaborative Enzyme Enhanced Reactive Immunoassay (CEER™).

1           **10.**     The method of any one of claims **1** to **9**, wherein the anti- $\alpha 4\beta 7$  integrin  
2 drug is ENTYVIO® (vedolizumab).

1           **11.**     The method of any one of claims **1** to **10**, wherein the sample is  
2 selected from the group consisting of a whole blood, serum or plasma sample.

1           **12.**     The method of any one of claims **1** to **11**, wherein the subject has had  
2 an inadequate response with, has lost response to, or was intolerant to an anti-TNF $\alpha$  drug.

1           **13.**     The method of claims **12**, wherein the anti-TNF $\alpha$  drug is a member  
2 selected from the group consisting of REMICADE® (infliximab), ENBREL® (etanercept),  
3 HUMIRA® (adalimumab), CIMZIA® (certolizumab pegol), SIMPONI® (golimumab),  
4 STELARA® (ustekinumab), and combinations thereof.

1           **14.**     The method of any one of claims **1** to **13**, wherein the subject has not  
2 previously been administered the anti- $\alpha 4\beta 7$  integrin drug.

1           **15.**     The method of any one of claims **1** to **14**, wherein the method further  
2 comprising applying a statistical analysis to the predictive marker profile to predict whether  
3 the subject will develop autoantibodies against the anti- $\alpha 4\beta 7$  integrin drug.

1           **16.**     A method for predicting whether a subject having inflammatory bowel  
2 disease (IBD) is likely to develop autoantibodies to an anti- $\alpha 4\beta 7$  integrin drug, the method  
3 comprising:

4           (a) detecting the presence or level of at least one predictive marker selected  
5 from the group consisting of TNF $\alpha$ , human serum albumin (HSA), VEGF, angiopoietin-1  
6 (ANG-1), angiopoietin-2 (ANG-2), adenosine deaminase (ADA), serum  $\alpha 4\beta 7$  integrin, IL-  
7 12p40, C-reactive protein (CRP), matrix metalloproteinase 9 (MMP9), MAdCAM-1, VCAM-  
8 1, ICAM-1, and a combination thereof in a sample from the subject; and

9 (b) applying a statistical analysis to the presence or level of the at least one  
10 predictive marker of step (a) to generate a predictive marker profile to determine whether the  
11 subject is likely to develop or not to develop the autoantibodies.

1 17. The method of claim 16, wherein the inflammatory bowel disease is  
2 ulcerative colitis (UC) or Crohn's Disease (CD).

1 18. The method of any one of claims 16 to 17, wherein the at least one  
2 predictive marker comprises at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13 or more predictive  
3 markers.

1 19. The method of any one of claims 16 to 18, wherein detecting the  
2 presence or level of at least one predictive marker comprises performing a proximity dual  
3 detection assay or an immunoassay.

1 20. The method of claim 19, wherein the proximity dual detection assay is  
2 a Collaborative Enzyme Enhanced Reactive Immunoassay (CEER™).

1 21. The method of any one of claims 16 to 20, wherein the subject is likely  
2 not to develop the autoantibodies if the level of ANG-1, ADA, HSA, IL-12p40, MMP9,  
3 ICAM-1, and/or serum  $\alpha 4\beta 7$  integrin is higher than the corresponding reference value.

1 22. The method of any one of claims, 16 to 20, wherein the subject is  
2 likely not to develop autoantibodies if the level of MadCAM-1, VCAM-1, and/or TNF $\alpha$  is  
3 lower than the corresponding reference value.

1 23. The method of any one of claims 16 to 20, wherein the subject is likely  
2 to develop autoantibodies if the level of TNF $\alpha$ , VEGF, ANG-2, CRP, and/or VCAM-1 is  
3 higher than the corresponding reference value.

1 24. The method of any one of claims 16 to 23, wherein the subject has  
2 been previously administered the anti- $\alpha 4\beta 7$  integrin drug.

1 25. The method of any one of claims 16 to 24, further comprising  
2 determining the presence or level of autoantibodies in the sample from the subject.

1           **26.**     The method of any one of claims **16** to **25**, wherein the subject is  
2 maintained or is recommended to be maintained on the anti- $\alpha 4\beta 7$  integrin drug if the subject  
3 is likely to not develop autoantibodies.

1           **27.**     The method of any one of claims **16** to **26** , wherein the subject is not  
2 maintained or is recommended not to be maintained on the anti- $\alpha 4\beta 7$  integrin drug if the  
3 subject is likely to develop autoantibodies.

1           **28.**     The method of any one of claims **16** to **27**, wherein the sample is  
2 selected from the group consisting of a whole blood, serum or plasma sample.

1           **29.**     The method of any one of claims **16** to **28**, wherein the subject has had  
2 an inadequate response with, has lost response to, or was intolerant to an anti-TNF $\alpha$  drug.

1           **30.**     The method of claim **29**, wherein the anti-TNF $\alpha$  drug is a member  
2 selected from the group consisting of REMICADE<sup>®</sup> (infliximab), ENBREL<sup>®</sup> (etanercept),  
3 HUMIRA<sup>®</sup> (adalimumab), CIMZIA<sup>®</sup> (certolizumab pegol), SIMPONI<sup>®</sup> (golimumab),  
4 STELARA<sup>®</sup> (ustekinumab), and combinations thereof.

1           **31.**     The method of any one of claims **16** to **30**, wherein the anti- $\alpha 4\beta 7$   
2 integrin drug is ENTYVIO<sup>®</sup> (vedolizumab).

1           **32.**     A method for predicting whether a subject having inflammatory bowel  
2 disease (IBD) will develop autoantibodies against an anti- $\alpha 4\beta 7$  integrin drug at a later time  
3 point during a course of therapy with the anti- $\alpha 4\beta 7$  integrin drug, the method comprising:

4           (a) measuring the presence or level of at least one predictive marker at a first  
5 time point in a sample from the subject to determine a first predictive marker profile, wherein  
6 the at least one predictive marker is selected from the group consisting of TNF $\alpha$ , human  
7 serum albumin (HSA), VEGF, angiopoietin-1 (ANG-1), angiopoietin-2 (ANG-2), adenosine  
8 deaminase (ADA), serum  $\alpha 4\beta 7$  integrin, IL-12p40, C-reactive protein (CRP), matrix  
9 metalloproteinase 9 (MMP9), MAdCAM-1, VCAM-1, ICAM-1, and a combination thereof;

10           (b) measuring the presence of level of the same at least one predictive marker  
11 at a later time point in a second sample from the subject to determine a second predictive  
12 marker profile; and



13 (c) applying a statistical analysis to the first and second predictive marker  
14 profiles to determine that the subject will develop autoantibodies against the anti- $\alpha 4\beta 7$   
15 integrin drug during the course of therapy.

1 **33.** The method of claim **32**, wherein the at least one predictive marker  
2 comprises at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13 or more predictive markers.

1 **34.** The method of claim **32** or **33**, wherein detecting the presence or level  
2 of at least one predictive marker comprises performing a proximity dual detection assay or an  
3 immunoassay.

1 **35.** The method of claim **34**, wherein the proximity dual detection assay is  
2 a Collaborative Enzyme Enhanced Reactive Immunoassay (CEER™).

1 **36.** The method of any one of claims **32** to **35**, wherein the anti- $\alpha 4\beta 7$   
2 integrin drug is ENTYVIO® (vedolizumab).

1 **37.** The method of any one of claims **32** to **36**, wherein the later time point  
2 is at weeks 2, 4, 6, 8, 10, 12, 16, 24, 32, 40, 48, or 52 during the course of therapy.

1 **38.** A method for inducing clinical remission in a subject having  
2 inflammatory bowel disease (IBD) and after receiving an induction therapy of an anti- $\alpha 4\beta 7$   
3 integrin drug, comprising administering a maintenance therapy of an anti- $\alpha 4\beta 7$  integrin drug  
4 to a subject with IBD and having a higher level of serum  $\alpha 4\beta 7$  integrin and/or a lower level  
5 of MadCAM-1 compared to corresponding levels at baseline.

1 **39.** The method of claim **38**, wherein the maintenance therapy comprises  
2 administering a dose of the anti- $\alpha 4\beta 7$  integrin drug every 8 weeks during the course of  
3 therapy.

1 **40.** A method for inducing a clinical response in a subject having  
2 inflammatory bowel disease (IBD), comprising administering an induction therapy of an anti-  
3  $\alpha 4\beta 7$  integrin drug to a subject with IBD and having a higher level of ANG-1, ADA, HSA,  
4 IL-12p40, MMP9, ICAM-1, and/or serum  $\alpha 4\beta 7$  integrin, and/or a lower level of MadCAM-1,  
5 VCAM-1, and/or TNF $\alpha$  compared to a corresponding reference level.

6                   **41.**     The method of claim **40**, wherein the induction therapy comprises  
7 administering a dose of about 300 mg of the anti- $\alpha 4\beta 7$  integrin drug at weeks 0, 2 and 6  
8 during the course of therapy.

1                   **42.**     A method for inducing clinical remission in a subject having  
2 inflammatory bowel disease (IBD) and after receiving an induction therapy of an anti- $\alpha 4\beta 7$   
3 integrin drug, comprising administering a maintenance therapy of an anti- $\alpha 4\beta 7$  integrin drug  
4 to a subject with IBD and having a higher level of serum  $\alpha 4\beta 7$  integrin and/or a lower level  
5 of MadCAM-1 compared to corresponding levels at baseline.

1                   **43.**     The method of claim **42**, wherein the maintenance therapy comprises  
2 administering a dose of about 300 mg of the anti- $\alpha 4\beta 7$  integrin drug every 8 weeks during the  
3 course of therapy.

1                   **44.**     A therapy regimen for an anti- $\alpha 4\beta 7$  integrin drug, said therapy regimen  
2 comprising:  
3                   measuring the presence or level of at least one predictive marker selected from  
4 the group consisting of TNF $\alpha$ , human serum albumin (HSA), VEGF, angiopoietin-1 (ANG-  
5 1), angiopoietin-2 (ANG-2), adenosine deaminase (ADA), serum  $\alpha 4\beta 7$  integrin, IL-12p40, C-  
6 reactive protein (CRP), MMP9, MAdCAM-1, VCAM-1, ICAM-1, and a combination thereof  
7 in a sample from the subject; and  
8                   administering the therapy regimen of the anti- $\alpha 4\beta 7$  integrin drug according to  
9 a predictive marker profile based on a higher or lower level of the at least one predictive  
10 marker compared to a corresponding reference value.

1                   **45.**     The method of claim **44**, wherein the therapy regimen of the anti- $\alpha 4\beta 7$   
2 integrin drug is administered if the level of ANG-1, ADA, HSA, IL-12p40, MMP9, ICAM-1,  
3 and/or serum  $\alpha 4\beta 7$  integrin level is higher and/or the level of MadCAM-1, VCAM-1, and/or  
4 TNF $\alpha$  is lower than the corresponding reference value.

1                   **46.**     The method of claim **44**, wherein the therapy regimen is not  
2 administered if the level of TNF $\alpha$ , VEGF, ANG-2, CRP, and/or VCAM-1 is higher than the  
3 corresponding reference value.


1                   **47.**     The method of claim **45** or **46**, wherein the therapy regimen comprises  
2 an induction therapy regimen.

1                   **48.**     The method of claim **44**, wherein the therapy regimen of the anti- $\alpha 4\beta 7$   
2 integrin drug is administered if the level of MadCAM-1 is lower and/or the level serum  $\alpha 4\beta 7$   
3 integrin is higher than the corresponding reference value.

1                   **49.**     The method of claim **48**, wherein the therapy regimen comprises a  
2 maintenance therapy regimen.


1                   **50.**     The method of any one of claims **38** to **49**, wherein the anti- $\alpha 4\beta 7$   
2 integrin drug is ENTYVIO<sup>®</sup> (vedolizumab).

**FIG. 1A**




Higher TNFa levels at baseline result in lower vedolizumab at week 6

**FIG. 1B**



2/22




Higher VEGF levels at baseline result in lower vedolizumab at week 6

FIG. 2


**FIG. 3A**

Higher ANG-2 levels at baseline result in lower vedolizumab at week 6



**FIG. 3B**


Bivariate Fit of Log[Vedo ug/ml\_wk6] By Log[Ang1\_baseline]



Higher ANG-1 levels at baseline result in higher vedolizumab at week 6


4/22

**FIG. 4A**



Higher ADA levels at baseline result in Higher vedolizumab at week 6


**FIG. 4B**



Higher ADA levels at week 2 result in Higher vedolizumab at week 6

5/22


**FIG. 5A**



Higher human serum albumin levels at baseline result in higher vedolizumab at week 6

**FIG. 5B**

Association between levels of vedolizumab at week 6 and baseline HGB (Hemoglobin): No association






6/22

**FIG. 6A**


**Bivariate Fit of Log[Vedo ug/ml\_wk6] By Log[P40\_baseline]**



Association between level of vedolizumab at week 6 and level of IL-12p40 at baseline (increased levels)

**FIG. 6B**


**Bivariate Fit of Log[Vedo ug/ml\_wk6] By Log[MMP9\_wk2]**



Association between levels of vedolizumab at week 6 and levels of MMP9 at week 2 (increased levels)


7/22

**FIG. 7A**



**Association between vedolizumab at wk 6 and MadCAM-1 at week 2**

**FIG. 7B**



**Association between Vedolizumab at week 6 and CRP at baseline**

8/22

FIG. 8A





FIG. 8B



9/22

Longitudinal Analysis of  $\alpha 4\beta 7$  Integrin Levels in Serum




FIG. 9

10/22

Longitudinal Analysis of MAdCAM-1 Levels in Serum





FIG. 10

11/22


**FIG. 11A**

Multiple regression model with interaction between VCAM and  $\alpha 4\beta 7$  integrin (baseline) to predict vedolizumab at week 6.




**FIG. 11B**

Multiple regression model with interaction between VCAM and MAdCAM-1 (baseline) to predict vedolizumab at week 6.




12/22

**FIG. 12A**



**Baseline MadCAM-1 negatively correlated with ICAM-1 (-0.4564: p-value=0.21)**

**FIG. 12B**



**MadCAM-1 at baseline is negatively correlated to VCAM-1 at baseline (-0.5459, p-value=0.1284)**

MadCAM-1 longitudinal change & pairwise p-values

- Log[Madcam pg/ml\_baseline]
- Log[Madcam pg/ml\_post\_wk0]
- Log[Madcam pg/ml\_wk2]
- Log[Madcam pg/ml\_wk6]




FIG. 13



14/22

### VCAM-1 longitudinal change




FIG. 14

### $\alpha 4\beta 7$ integrin longitudinal change




FIG. 15

Vedolizumab longitudinal change






FIG. 16



MadCam expression decreases with Vedo treatment


FIG. 17

**α4β7 integrin longitudinal change**




**FIG. 18**

**FIG. 19A**




**FIG. 19B**




20/22

**FIG. 20A**



**FIG. 20B**






FIG. 21






FIG. 22

**INTERNATIONAL SEARCH REPORT**

International application No  
PCT/IB2015/059303

**A. CLASSIFICATION OF SUBJECT MATTER**  
 INV. G01N33/564 G01N33/68  
 ADD.  
 According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**  
 Minimum documentation searched (classification system followed by classification symbols)  
 G01N  
 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)  
 EPO-Internal, WPI Data, BIOSIS, EMBASE

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2014/055824 A1 (GENENTECH INC [US]; HOFFMANN LA ROCHE [CH]) 10 April 2014 (2014-04-10) p. 7, paragraph [0019] - p. 20, paragraph [0046], p. 23, paragraph [0062], p. 67, paragraph [0220] - p. 74, paragraph [0251], claims 1-42	1-14, 38-50
X	WO 2009/140684 A2 (GENENTECH INC [US]; GELZLEICHTER THOMAS RICHARD [US]; HIRARAGI HAJIME) 19 November 2009 (2009-11-19) p. 2, paragraph [0009] - p. 3, paragraph [0045], p. 16, paragraph [0232] - p. 31, paragraph [0394], claims 1-25 ----- -/--	1-14, 38-50

Further documents are listed in the continuation of Box C.

See patent family annex.

\* Special categories of cited documents :

<p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier application or patent but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p>	<p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</p> <p>"&amp;" document member of the same patent family</p>
---	---

Date of the actual completion of the international search <b>29 January 2016</b>	Date of mailing of the international search report <b>05/02/2016</b>
---	---

Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer <b>R. von Eggelkraut-G.</b>
--	---

## INTERNATIONAL SEARCH REPORT

International application No  
PCT/IB2015/059303

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2012/061074 A1 (ELAN PHARM INC [US]; CHACKERIAN ALISSA A [US]) 10 May 2012 (2012-05-10) p. 2, line 32 - p. 5, line 17, p. 9, line 5 - p. 23, line 21, p. 28, line 21 - p. 37, line 17, claims 1-21 -----	1-14
X	WILLIAM J. SANDBORN ET AL: "Vedolizumab as Induction and Maintenance Therapy for Crohn's Disease", NEW ENGLAND JOURNAL OF MEDICINE, vol. 369, no. 8, 22 August 2013 (2013-08-22), pages 711-721, XP055245598, US ISSN: 0028-4793, DOI: 10.1056/NEJMoa1215739 the whole document -----	1-50
X	ROSARIO M ET AL: "P489 Exposure-response relationship during vedolizumab induction therapy in adults with ulcerative colitis", JOURNAL OF CROHN'S AND COLITIS, vol. 8, 1 February 2014 (2014-02-01), XP028618082, ISSN: 1873-9946, DOI: 10.1016/S1873-9946(14)60609-7 Posters P488 and P489 -----	1-14, 40-44, 46-50
X	SMITH MICHAEL A ET AL: "Vedolizumab: An a alpha(4)beta(7) Integrin Inhibitor for Inflammatory Bowel Diseases", ANNALS OF PHARMACOTHERAPY, vol. 48, no. 12, 1 December 2014 (2014-12-01), pages 1629-1635, XP009188218, the whole document -----	38-50

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/IB2015/059303

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2014055824 A1	10-04-2014	AU 2013326921 A1	02-04-2015
		CA 2884368 A1	10-04-2014
		CN 104870056 A	26-08-2015
		EP 2903691 A1	12-08-2015
		KR 20150058270 A	28-05-2015
		US 2015307611 A1	29-10-2015
		WO 2014055824 A1	10-04-2014
		-----	
WO 2009140684 A2	19-11-2009	AU 2009246071 A1	19-11-2009
		CA 2723614 A1	19-11-2009
		CN 102124344 A	13-07-2011
		DK 2279004 T3	02-02-2015
		EP 2279004 A2	02-02-2011
		ES 2533480 T3	10-04-2015
		JP 2011521236 A	21-07-2011
		JP 2013253987 A	19-12-2013
		KR 20110018365 A	23-02-2011
		KR 20130038946 A	18-04-2013
		SI 2279004 T1	29-05-2015
		US 2010255508 A1	07-10-2010
		US 2013109032 A1	02-05-2013
		WO 2009140684 A2	19-11-2009
-----			
WO 2012061074 A1	10-05-2012	AU 2011323820 A1	23-05-2013
		CA 2816016 A1	10-05-2012
		EP 2632492 A1	04-09-2013
		JP 2014500950 A	16-01-2014
		JP 2015215363 A	03-12-2015
		US 2013337470 A1	19-12-2013
		WO 2012061074 A1	10-05-2012
-----			