



(86) Date de dépôt PCT/PCT Filing Date: 2008/12/22
 (87) Date publication PCT/PCT Publication Date: 2009/07/16
 (85) Entrée phase nationale/National Entry: 2010/06/28
 (86) N° demande PCT/PCT Application No.: US 2008/088066
 (87) N° publication PCT/PCT Publication No.: 2009/088753
 (30) Priorité/Priority: 2008/01/03 (US61/018,880)

(51) Cl.Int./Int.Cl. *C12Q 1/533* (2006.01),
A61K 38/52 (2006.01), *C07K 16/40* (2006.01),
C12N 15/61 (2006.01), *C12N 9/90* (2006.01)

(71) Demandeur/Applicant:
 VERENIUM CORPORATION, US

(72) Inventeurs/Inventors:
 WEINER, DAVID, US;
 BURKE, ELLEN, US;
 LUGINBUHL, PETER, US;
 BUENO, ANALIA, US;
 CUENCA, JOSLIN, US;
 DE SOUZA, MERVYN L., US;
 KOLLMANN, SHERRY, US

(74) Agent: OGILVY RENAULT LLP/S.E.N.C.R.L.,S.R.L.

(54) Titre : ISOMERASES, ACIDES NUCLEIQUES CODANT POUR CELLES-CI ET LEURS PROCEDES DE FABRICATION D'UTILISATION

(54) Title: ISOMERASES, NUCLEIC ACIDS ENCODING THEM AND METHODS FOR MAKING AND USING THEM

(57) **Abrégé/Abstract:**

This invention relates generally to enzymes, polynucleotides encoding the enzymes, the use of such polynucleotides and polypeptides and more specifically to enzymes having isomerase activity, e.g., racemase activity, e.g., amino acid racemase activity, alanine racemase activity, and/or epimerase activity, and/or catalyze the re[arrangement of atoms within a molecule, catalyze the conversion of one isomer into another, catalyze the conversion of an optically active substrate into a raceme, which is optically inactive, catalyze the interconversion of substrate enantiomers, catalyze the stereochemical inversion around the asymmetric carbon atom in a substrate having only one center of asymmetry, catalyze the stereochemical inversion of the configuration around an asymmetric carbon atom in a substrate having more than one asymmetric center, and/or catalyze the racemization of amino acids. Thus, the invention provides enzymes, compositions, methods for production of pharmaceutical compositions, pharmaceutical intermediates, antibiotics, sweeteners, peptide enzymes, peptide hormones, fuel and fuel additive compositions, foods and food additives, beverage and beverage additives, feeds and feed additives, drugs and drug additives, dietary supplements, textiles, wood, paper, pulp, and detergents comprising the polypeptides or polynucleotides in accordance with the invention.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
16 July 2009 (16.07.2009)

PCT

(10) International Publication Number
WO 2009/088753 A1

(51) International Patent Classification:

C12Q 1/533 (2006.01) *A61K 38/52* (2006.01)
C12N 9/90 (2006.01) *C07K 16/40* (2006.01)
C12N 15/61 (2006.01)

92126 (US). **DE SOUZA, Mervyn, L.** [US/US]; 10935
38th Avenue N., Plymouth, MN 55441 (US). **KOLL-**
MANN, Sherry [US/US]; 12031 99th Avenue N., Maple
Grove, MN 55249 (US).

(21) International Application Number:

PCT/US2008/088066

(74) **Agents: EINHORN, Gregory, P.** et al.; Morrison & Fo-
erster LLP, 12531 High Bluff Drive, Suite 100, San Diego,
CA 92130-2040 (US).

(22) International Filing Date:

22 December 2008 (22.12.2008)

(81) **Designated States** (*unless otherwise indicated, for every
kind of national protection available*): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA,
CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE,
EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID,
IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK,
LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW,
MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT,
RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ,
TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM,
ZW.

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/018,880 3 January 2008 (03.01.2008) US

(71) **Applicant** (*for all designated States except US*): **VERE-**
NIUM CORPORATION [US/US]; 4955 Directors Place,
San Diego, CA 92121 (US).

(72) Inventors; and

(75) **Inventors/Applicants** (*for US only*): **WEINER, David**
[GB/US]; 13416 Portofino Drive, Del Mar, CA 92014
(US). **BURKE, Ellen** [US/US]; 3428 Talbot Street, San
Diego, CA 92106 (US). **LUGINBUHL, Peter** [CH/US];
10802 Calle De La Siena, San Diego, CA 92130 (US).
BUENO, Analia [US/US]; 4498 Ocean View Boulevard,
San Diego, CA 92113 (US). **CUENCA, Joslin** [US/US];
8847 Spectrum Center Boulevard, #6116, San Diego, CA

(84) **Designated States** (*unless otherwise indicated, for every
kind of regional protection available*): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL,
NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG,
CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:— *with international search report*

(54) **Title:** ISOMERASES, NUCLEIC ACIDS ENCODING THEM AND METHODS FOR MAKING AND USING THEM

(57) **Abstract:** This invention relates generally to enzymes, polynucleotides encoding the enzymes, the use of such polynucleotides and polypeptides and more specifically to enzymes having isomerase activity, e.g., racemase activity, e.g., amino acid racemase activity, alanine racemase activity, and/or epimerase activity, and/or catalyze the re-arrangement of atoms within a molecule, catalyze the conversion of one isomer into another, catalyze the conversion of an optically active substrate into a raceme, which is optically inactive, catalyze the interconversion of substrate enantiomers, catalyze the stereochemical inversion around the asymmetric carbon atom in a substrate having only one center of asymmetry, catalyze the stereochemical inversion of the configuration around an asymmetric carbon atom in a substrate having more than one asymmetric center, and/or catalyze the racemization of amino acids. Thus, the invention provides enzymes, compositions, methods for production of pharmaceutical compositions, pharmaceutical intermediates, antibiotics, sweeteners, peptide enzymes, peptide hormones, fuel and fuel additive compositions, foods and food additives, beverage and beverage additives, feeds and feed additives, drugs and drug additives, dietary supplements, textiles, wood, paper, pulp, and detergents comprising the polypeptides or polynucleotides in accordance with the invention.

WO 2009/088753 A1

DEMANDE OU BREVET VOLUMINEUX

LA PRÉSENTE PARTIE DE CETTE DEMANDE OU CE BREVET COMPREND PLUS D'UN TOME.

CECI EST LE TOME 1 DE 2
CONTENANT LES PAGES 1 À 311

NOTE : Pour les tomes additionels, veuillez contacter le Bureau canadien des brevets

JUMBO APPLICATIONS/PATENTS

THIS SECTION OF THE APPLICATION/PATENT CONTAINS MORE THAN ONE VOLUME

THIS IS VOLUME 1 OF 2
CONTAINING PAGES 1 TO 311

NOTE: For additional volumes, please contact the Canadian Patent Office

NOM DU FICHER / FILE NAME :

NOTE POUR LE TOME / VOLUME NOTE:

ISOMERASES, NUCLEIC ACIDS ENCODING THEM AND METHODS FOR MAKING AND USING THEM

5 REFERENCE TO SEQUENCE LISTING SUBMITTED VIA EFS-WEB

The entire content of the following electronic submission of the sequence listing via the USPTO EFS-WEB server, as authorized and set forth in MPEP §1730 II.B.2.(a)(C), is incorporated herein by reference in its entirety for all purposes. The sequence listing is identified on the electronically filed .txt file as follows:

File Name	Date of Creation	Size
564462016240Seqlist.txt	December 18, 2008	1,294,466

10 FIELD OF THE INVENTION

This invention relates generally to enzymes, polynucleotides encoding the enzymes, the use of such polynucleotides and polypeptides and more specifically to enzymes having isomerase activity, e.g., racemase activity, e.g., amino acid racemase activity, alanine racemase activity, and/or epimerase activity, and/or catalyze the re-
15 arrangement of atoms within a molecule, catalyze the conversion of one isomer into another, catalyze the conversion of an optically active substrate into a raceme, which is optically inactive, catalyze the interconversion of substrate enantiomers, catalyze the stereochemical inversion around the asymmetric carbon atom in a substrate having only one center of asymmetry, catalyze the stereochemical inversion of the configuration
20 around an asymmetric carbon atom in a substrate having more than one asymmetric center, and/or catalyze the racemization of amino acids. Thus, the invention provides enzymes, compositions, methods for production of pharmaceutical (drug) compositions, pharmaceutical (drug) precursors and intermediates, antibiotics, sweeteners, peptide enzymes, peptide hormones, fuel and fuel additive compositions, foods and food additives,
25 beverage and beverage additives, feeds and feed additives, drugs and drug additives, dietary supplements, textiles, wood, paper, pulp, and detergents comprising the polypeptides or polynucleotides in accordance with the invention.

BACKGROUND

Isomerases, e.g., racemases, e.g., amino acid racemases, alanine racemases, and/or
30 epimerases catalyze the re-arrangement of atoms within a molecule, catalyze the conversion of one isomer into another, catalyze the conversion of an optically active substrate into a raceme, which is optically inactive, catalyze the interconversion of

substrate enantiomers, catalyze the stereochemical inversion around the asymmetric carbon atom in a substrate having only one center of asymmetry, catalyze the stereochemical inversion of the configuration around an asymmetric carbon atom in a substrate having more than one asymmetric center, and/or catalyze the racemization of amino acids. Isomerases, e.g., racemases, e.g., amino acid racemases, alanine racemases, and/or epimerases are of considerable commercial value, being used in the pharmaceutical industry, in the food, feed and beverage industries, e.g. for the production of sweeteners, in the wood/paper industry and in the fuel industry.

10

SUMMARY OF THE INVENTION

This invention provides enzymes having isomerase activity, e.g., racemase activity, e.g., amino acid racemase activity, alanine racemase activity, and/or epimerase activity, and/or catalyze the re-arrangement of atoms within a molecule, catalyze the conversion of one isomer into another, catalyze the conversion of an optically active substrate into a raceme, which is optically inactive, catalyze the interconversion of substrate enantiomers, catalyze the stereochemical inversion around the asymmetric carbon atom in a substrate having only one center of asymmetry, catalyze the stereochemical inversion of the configuration around an asymmetric carbon atom in a substrate having more than one asymmetric center, and/or catalyze the racemization of amino acids. The invention further provides enzymes having isomerase activity, e.g., racemase activity, e.g., amino acid racemase activity, alanine racemase activity, and/or epimerase activity and nucleic acids encoding them, vectors and cells comprising them, probes for amplifying and identifying these an isomerase-, e.g., a racemase-, e.g., an amino acid racemase-, an alanine racemase-, and/or an epimerase-isomerase-, e.g., racemase-, e.g., amino acid racemase-, alanine racemase-, and/or epimerase-encoding nucleic acids, and methods for making and using these polypeptides and peptides.

The invention provides enzymes, compositions, methods for production of pharmaceutical (drug) compositions, pharmaceutical (drug) precursors and intermediates, antibiotics, sweeteners, peptide enzymes, peptide hormones, fuel and fuel additive compositions, foods and food additives, beverage and beverage additives, feeds and feed additives, drugs and drug additives, dietary supplements, textiles, wood, paper, pulp, and detergents comprising the polypeptides or polynucleotides in accordance with the invention. These compositions can be formulated in a variety of forms, such as tablets,

gels, pills, implants, liquids, sprays, films, micelles, powders, food, feed pellets or as any type of encapsulated form.

In some embodiments, the isomerases, e.g., racemases, e.g., amino acid racemases, alanine racemases, and/or epimerases and/or compositions thereof may be useful in
5 pharmaceutical, industrial, and/or agricultural contexts.

In some embodiments, the isomerases, e.g., racemases, e.g., amino acid racemases, alanine racemases, and/or epimerases and/or compositions thereof may be useful for catalyzing the inversion of stereochemistry in biological molecules. In some
10 embodiments, the isomerases, e.g., racemases, e.g., amino acid racemases, alanine racemases, and/or epimerases and/or compositions thereof may be useful for catalyzing the interconversion of substrate enantiomers. In some embodiments, isomerases, e.g., racemases, e.g., amino acid racemases, and/or alanine racemases catalyze the stereochemical inversion around the asymmetric carbon atom in a substrate having only one center of asymmetry. In some embodiments, isomerases, e.g., epimerases catalyze the
15 stereochemical inversion of the configuration around an asymmetric carbon atom in a substrate having more than one asymmetric center. In some embodiments, isomerases, e.g., racemases, e.g., amino acid racemases, and/or alanine racemases are provided that catalyze the racemization of amino acids. In some embodiments, racemases are provided that catalyze the racemization of a specific amino acid. In some embodiments,
20 isomerases, e.g., racemases, e.g., amino acid racemases, and/or alanine racemases are provided that catalyze the racemization of several amino acids.

In some embodiments, the racemases, e.g., amino acid racemases, alanine racemases, and/or epimerases and/or compositions thereof may be useful in D-amino acid metabolism. D-amino acids are necessary for bacterial growth and for peptidoglycan
25 assembly and cross linking. D-amino acids are also present in the brains of newborn humans. Bacterial serine racemase plays an important role in vancomycin resistance. Some amino acid racemases are PLP dependent; other amino acid racemases are PLP independent. (see, e.g., Yoshimura, T., N. Esaki, 2003, Journal of Bioscience and Bioengineering. 96:103-109). In alternative embodiments, the racemases, e.g., amino acid
30 racemases, alanine racemases, and/or epimerases and/or combinations thereof are components in pharmaceutical (drug) compositions, pharmaceutical (drug) precursors and/or intermediates, e.g. as antibiotics or for treatment of amino acid deficiencies.

In alternative embodiments, the isomerases, e.g., racemases, e.g., amino acid racemases, alanine racemases, and/or epimerases of the invention and/or compositions thereof of may be useful as an antibiotic or in the preparation of antibiotics (see, e.g., Strych, U., M.J. Benedik. 2002, Journal of Bacteriology. 184:4321-4325).

5 In alternative embodiments, the isomerases, e.g., racemases, e.g., amino acid racemases, alanine racemases, and/or epimerases of the invention and/or compositions thereof of may be useful as in mediation of mammal nervous transmission and maintenance of bacterial cell wall rigidity and strength (see, e.g. Liu, L., K. Iwata, M. Yohda, K. Miki. 2002, FEBS. 528:114-118).

10 In alternative embodiments, the invention provides enzymes and processes for the bioconversion of any biomass into fuel, e.g. biofuel, e.g., ethanol, propanol, butanol, methanol, and/or biodiesel or biofuels such as synthetic liquids or gases, such as syngas, and the production of other fermentation products, e.g. succinic acid, lactic acid, or acetic acid.

15 In alternative embodiments, the invention provides polypeptides (and the nucleic acids that encode them) having at least one conservative amino acid substitution and retaining its isomerase activity, e.g., racemase activity, e.g., amino acid racemase activity, alanine racemase activity, and/or epimerase activity; or, wherein the at least one conservative amino acid substitution comprises substituting an amino acid with another
20 amino acid of like characteristics; or, a conservative substitution comprises: replacement of an aliphatic amino acid with another aliphatic amino acid; replacement of a Serine with a Threonine or vice versa; replacement of an acidic residue with another acidic residue; replacement of a residue bearing an amide group with another residue bearing an amide group; exchange of a basic residue with another basic residue; or replacement of an
25 aromatic residue with another aromatic residue;

In alternative embodiments, the invention provides polypeptides (and the nucleic acids that encode them) having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity but lacking a signal sequence, a prepro domain and/or other domain.

30 In alternative embodiments, the invention provides polypeptides (and the nucleic acids that encode them) having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity further comprising a heterologous sequence; and in one aspect, the heterologous sequence comprises, or consists of a sequence encoding: (i) a heterologous signal sequence, a

heterologous domain, a heterologous dockerin domain, a heterologous catalytic domain (CD), or a combination thereof; (ii) the sequence of (i), wherein the heterologous signal sequence, domain or catalytic domain (CD) is derived from a heterologous enzyme; or, (iii) a tag, an epitope, a targeting peptide, a cleavable sequence, a detectable moiety or an enzyme; and in one aspect, the heterologous signal sequence targets the encoded protein to a vacuole, the endoplasmic reticulum, a chloroplast or a starch granule.

In alternative embodiments, the invention provides polypeptides (and the nucleic acids that encode them) having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity, wherein the polypeptides are cofactor dependent or cofactor independent. In one embodiment, a cofactor dependent polypeptide requires the presence of a non-protein component to be functional. In one embodiment, the cofactor comprises a metal ion, a coenzyme, a pyridoxal-phosphate and or a phosphopantetheine.

The invention provides isolated, synthetic or recombinant nucleic acids comprising(a) a nucleic acid (polynucleotide) encoding at least one polypeptide, wherein the nucleic acid comprises a sequence having at least about 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more or complete (100%) sequence identity to the nucleic acid (polynucleotide) sequence of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO:31, SEQ ID NO:33, SEQ ID NO:35, SEQ ID NO:37, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:43, SEQ ID NO:45, SEQ ID NO:47, SEQ ID NO:49, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:57, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:89, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95, SEQ ID NO:97, SEQ ID NO:99, SEQ ID NO:101, SEQ ID NO:103, SEQ ID NO:105, SEQ ID NO:107, SEQ ID NO:109, SEQ ID NO:111, SEQ ID NO:113, SEQ ID NO:115, SEQ ID NO:117, SEQ ID NO:119, SEQ ID NO:121, SEQ ID NO:123, SEQ ID NO:125, SEQ ID NO:127, SEQ ID NO:129, SEQ ID NO:131, SEQ ID NO:133, SEQ ID NO:135, SEQ ID NO:137, SEQ ID NO:139, SEQ ID NO:141, SEQ ID NO:143, SEQ ID NO:145, SEQ ID NO:147,

SEQ ID NO:149, SEQ ID NO:151, SEQ ID NO:153, SEQ ID NO:155, SEQ ID NO:157,
SEQ ID NO:159, SEQ ID NO:161, SEQ ID NO:163, SEQ ID NO:165, SEQ ID NO:167,
SEQ ID NO:169, SEQ ID NO:171, SEQ ID NO:173, SEQ ID NO:175, SEQ ID NO:177,
SEQ ID NO:179, SEQ ID NO:181, SEQ ID NO:183, SEQ ID NO:185, SEQ ID NO:187,
5 SEQ ID NO:189, SEQ ID NO:191, SEQ ID NO:193, SEQ ID NO:195, SEQ ID NO:197,
SEQ ID NO:199, SEQ ID NO:201, SEQ ID NO:203, SEQ ID NO:205, SEQ ID NO:207,
SEQ ID NO:209, SEQ ID NO:211, SEQ ID NO:213, SEQ ID NO:215, SEQ ID NO:217,
SEQ ID NO:219, SEQ ID NO:221, SEQ ID NO:223, SEQ ID NO:225, SEQ ID NO:227,
SEQ ID NO:229, SEQ ID NO:231, SEQ ID NO:233, SEQ ID NO:235, SEQ ID NO:237,
10 SEQ ID NO:239, SEQ ID NO:241, SEQ ID NO:243, SEQ ID NO:245, SEQ ID NO:247,
SEQ ID NO:249, SEQ ID NO:251, SEQ ID NO:253, SEQ ID NO:255, SEQ ID NO:257,
SEQ ID NO:259, SEQ ID NO:261, SEQ ID NO:263, SEQ ID NO:265, SEQ ID NO:267,
SEQ ID NO:269, SEQ ID NO:271, SEQ ID NO:273, SEQ ID NO:275, SEQ ID NO:277,
SEQ ID NO:279, SEQ ID NO:281, SEQ ID NO:283, SEQ ID NO:285, SEQ ID NO:287,
15 SEQ ID NO:289, SEQ ID NO:291, SEQ ID NO:293, SEQ ID NO:295, SEQ ID NO:297,
SEQ ID NO:299, SEQ ID NO:301, SEQ ID NO:303, SEQ ID NO:305, SEQ ID NO:307,
SEQ ID NO:309, SEQ ID NO:311, SEQ ID NO:313, SEQ ID NO:315, SEQ ID NO:317,
SEQ ID NO:319, SEQ ID NO:321, SEQ ID NO:323, SEQ ID NO:325, SEQ ID NO:327,
SEQ ID NO:329, SEQ ID NO:331, SEQ ID NO:333, SEQ ID NO:335, SEQ ID NO:337,
20 SEQ ID NO:339, SEQ ID NO:341, SEQ ID NO:343, SEQ ID NO:345, SEQ ID NO:347,
SEQ ID NO:349, SEQ ID NO:351, SEQ ID NO:353, SEQ ID NO:355, SEQ ID NO:357,
SEQ ID NO:359, SEQ ID NO:361, SEQ ID NO:363, SEQ ID NO:365, SEQ ID NO:367,
SEQ ID NO:369, SEQ ID NO:371, SEQ ID NO:373, SEQ ID NO:375, SEQ ID NO:377,
SEQ ID NO:379, SEQ ID NO:381, SEQ ID NO:383, SEQ ID NO:385, SEQ ID NO:387,
25 SEQ ID NO:389, SEQ ID NO:391, SEQ ID NO:393, SEQ ID NO:395, SEQ ID NO:397,
SEQ ID NO:399, SEQ ID NO:401, SEQ ID NO:403, SEQ ID NO:405, SEQ ID NO:407,
SEQ ID NO:409, SEQ ID NO:411, SEQ ID NO:413, SEQ ID NO:415, SEQ ID NO:417,
SEQ ID NO:419, SEQ ID NO:421, SEQ ID NO:423, SEQ ID NO:425, SEQ ID NO:427,
SEQ ID NO:429, SEQ ID NO:431, SEQ ID NO:433, SEQ ID NO:435, SEQ ID NO:437,
30 SEQ ID NO:439, SEQ ID NO:441, SEQ ID NO:443, SEQ ID NO:445, SEQ ID NO:447,
SEQ ID NO:449, SEQ ID NO:451, SEQ ID NO:453, SEQ ID NO:455, SEQ ID NO:457,
SEQ ID NO:459, SEQ ID NO:461, SEQ ID NO:463, SEQ ID NO:465, SEQ ID NO:467,
SEQ ID NO:469, SEQ ID NO:471, SEQ ID NO:473, SEQ ID NO:475, SEQ ID NO:477,
SEQ ID NO:479, SEQ ID NO:481, SEQ ID NO:483, SEQ ID NO:485, SEQ ID NO:487,

SEQ ID NO:489, SEQ ID NO:491, SEQ ID NO:493, SEQ ID NO:495 or SEQ ID
NO:497; wherein the nucleic acid encodes at least one polypeptide having an isomerase
activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine
racemase activity, and/or an epimerase activity, or encodes a polypeptide or peptide
5 capable of generating an isomerase specific antibody, e.g., a racemase specific antibody,
e.g., an amino acid racemase, an alanine racemase, and/or an epimerase specific antibody
(a polypeptide or peptide that acts as an epitope or immunogen),

(b) the nucleic acid (polynucleotide) of (a), wherein the sequence identities are
determined: (A) by analysis with a sequence comparison algorithm or by a visual
10 inspection, or (B) over a region of at least about 20, 30, 40, 50, 75, 100, 150, 200, 250, 300,
350, 400, 450, 500, 550, 600, 650, 700, 750, 800, 850, 900, 950, 1000, 1050, 1100, 1150 or
more residues, or over the full length of a cDNA, transcript (mRNA) or gene;

(c) the nucleic acid (polynucleotide) of (a) or (b), wherein the sequence
comparison algorithm is a BLAST version 2.2.2 algorithm where a filtering setting is set
15 to blastall -p blastp -d "nr pataa" -F F, and all other options are set to default;

(d) a nucleic acid (polynucleotide) encoding at least one polypeptide or peptide,
wherein the nucleic acid comprises a sequence that hybridizes under stringent conditions
to a nucleic acid comprising the sequence of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5,
SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID
20 NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27,
SEQ ID NO:29, SEQ ID NO:31, SEQ ID NO:33, SEQ ID NO:35, SEQ ID NO:37, SEQ
ID NO:39, SEQ ID NO:41, SEQ ID NO:43, SEQ ID NO:45, SEQ ID NO:47, SEQ ID
NO:49, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:57, SEQ ID NO:59,
SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ
25 ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID
NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:89, SEQ ID NO:91,
SEQ ID NO:93, SEQ ID NO:95, SEQ ID NO:97, SEQ ID NO:99, SEQ ID NO:101, SEQ
ID NO:103, SEQ ID NO:105, SEQ ID NO:107, SEQ ID NO:109, SEQ ID NO:111, SEQ
ID NO:113, SEQ ID NO:115, SEQ ID NO:117, SEQ ID NO:119, SEQ ID NO:121, SEQ
30 ID NO:123, SEQ ID NO:125, SEQ ID NO:127, SEQ ID NO:129, SEQ ID NO:131, SEQ
ID NO:133, SEQ ID NO:135, SEQ ID NO:137, SEQ ID NO:139, SEQ ID NO:141, SEQ
ID NO:143, SEQ ID NO:145, SEQ ID NO:147, SEQ ID NO:149, SEQ ID NO:151, SEQ
ID NO:153, SEQ ID NO:155, SEQ ID NO:157, SEQ ID NO:159, SEQ ID NO:161, SEQ
ID NO:163, SEQ ID NO:165, SEQ ID NO:167, SEQ ID NO:169, SEQ ID NO:171, SEQ

ID NO:173, SEQ ID NO:175, SEQ ID NO:177, SEQ ID NO:179, SEQ ID NO:181, SEQ
ID NO:183, SEQ ID NO:185, SEQ ID NO:187, SEQ ID NO:189, SEQ ID NO:191, SEQ
ID NO:193, SEQ ID NO:195, SEQ ID NO:197, SEQ ID NO:199, SEQ ID NO:201, SEQ
ID NO:203, SEQ ID NO:205, SEQ ID NO:207, SEQ ID NO:209, SEQ ID NO:211, SEQ
5 ID NO:213, SEQ ID NO:215, SEQ ID NO:217, SEQ ID NO:219, SEQ ID NO:221, SEQ
ID NO:223, SEQ ID NO:225, SEQ ID NO:227, SEQ ID NO:229, SEQ ID NO:231, SEQ
ID NO:233, SEQ ID NO:235, SEQ ID NO:237, SEQ ID NO:239, SEQ ID NO:241, SEQ
ID NO:243, SEQ ID NO:245, SEQ ID NO:247, SEQ ID NO:249, SEQ ID NO:251, SEQ
ID NO:253, SEQ ID NO:255, SEQ ID NO:257, SEQ ID NO:259, SEQ ID NO:261, SEQ
10 ID NO:263, SEQ ID NO:265, SEQ ID NO:267, SEQ ID NO:269, SEQ ID NO:271, SEQ
ID NO:273, SEQ ID NO:275, SEQ ID NO:277, SEQ ID NO:279, SEQ ID NO:281, SEQ
ID NO:283, SEQ ID NO:285, SEQ ID NO:287, SEQ ID NO:289, SEQ ID NO:291, SEQ
ID NO:293, SEQ ID NO:295, SEQ ID NO:297, SEQ ID NO:299, SEQ ID NO:301, SEQ
ID NO:303, SEQ ID NO:305, SEQ ID NO:307, SEQ ID NO:309, SEQ ID NO:311, SEQ
15 ID NO:313, SEQ ID NO:315, SEQ ID NO:317, SEQ ID NO:319, SEQ ID NO:321, SEQ
ID NO:323, SEQ ID NO:325, SEQ ID NO:327, SEQ ID NO:329, SEQ ID NO:331, SEQ
ID NO:333, SEQ ID NO:335, SEQ ID NO:337, SEQ ID NO:339, SEQ ID NO:341, SEQ
ID NO:343, SEQ ID NO:345, SEQ ID NO:347, SEQ ID NO:349, SEQ ID NO:351, SEQ
ID NO:353, SEQ ID NO:355, SEQ ID NO:357, SEQ ID NO:359, SEQ ID NO:361, SEQ
20 ID NO:363, SEQ ID NO:365, SEQ ID NO:367, SEQ ID NO:369, SEQ ID NO:371, SEQ
ID NO:373, SEQ ID NO:375, SEQ ID NO:377, SEQ ID NO:379, SEQ ID NO:381, SEQ
ID NO:383, SEQ ID NO:385, SEQ ID NO:387, SEQ ID NO:389, SEQ ID NO:391, SEQ
ID NO:393, SEQ ID NO:395, SEQ ID NO:397, SEQ ID NO:399, SEQ ID NO:401, SEQ
ID NO:403, SEQ ID NO:405, SEQ ID NO:407, SEQ ID NO:409, SEQ ID NO:411, SEQ
25 ID NO:413, SEQ ID NO:415, SEQ ID NO:417, SEQ ID NO:419, SEQ ID NO:421, SEQ
ID NO:423, SEQ ID NO:425, SEQ ID NO:427, SEQ ID NO:429, SEQ ID NO:431, SEQ
ID NO:433, SEQ ID NO:435, SEQ ID NO:437, SEQ ID NO:439, SEQ ID NO:441, SEQ
ID NO:443, SEQ ID NO:445, SEQ ID NO:447, SEQ ID NO:449, SEQ ID NO:451, SEQ
ID NO:453, SEQ ID NO:455, SEQ ID NO:457, SEQ ID NO:459, SEQ ID NO:461, SEQ
30 ID NO:463, SEQ ID NO:465, SEQ ID NO:467, SEQ ID NO:469, SEQ ID NO:471, SEQ
ID NO:473, SEQ ID NO:475, SEQ ID NO:477, SEQ ID NO:479, SEQ ID NO:481, SEQ
ID NO:483, SEQ ID NO:485, SEQ ID NO:487, SEQ ID NO:489, SEQ ID NO:491, SEQ
ID NO:493, SEQ ID NO:495 or SEQ ID NO:497,

and the stringent conditions comprise a wash step comprising a wash in 0.2X SSC at a temperature of about 65°C for about 15 minutes;

(e) the nucleic acid (polynucleotide) of any of (a) to (d) having a length of at least about 20, 25, 30, 50, 75, 100, 125, 150, 175, 200, 225, 300, 350, 400, 450, 500, 550, 600, 5 650, 700, 750, 800, 850, 900, 950, 1000, 1050, 1100, 1150 or more nucleotide residues, or the full length of a gene or a transcript;

(f) a nucleic acid (polynucleotide) encoding at least one polypeptide having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity, wherein the polypeptide comprises
10 the sequence of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:22, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:28, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:42, SEQ ID NO:44, SEQ ID NO:46, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:52,
15 SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:58, SEQ ID NO:60, SEQ ID NO:62, SEQ ID NO:64, SEQ ID NO:66, SEQ ID NO:68, SEQ ID NO:70, SEQ ID NO:72, SEQ ID NO:74, SEQ ID NO:76, SEQ ID NO:78, SEQ ID NO:80, SEQ ID NO:82, SEQ ID NO:84, SEQ ID NO:86, SEQ ID NO:88, SEQ ID NO:90, SEQ ID NO:92, SEQ ID NO:94, SEQ ID NO:96, SEQ ID NO:98, SEQ ID NO:100, SEQ ID NO:102, SEQ ID NO:104, SEQ ID NO:106, SEQ ID NO:108, SEQ ID NO:110, SEQ ID NO:112, SEQ ID NO:114, SEQ ID NO:116, SEQ ID NO:118, SEQ ID NO:120, SEQ ID NO:122, SEQ ID NO:124, SEQ ID NO:126, SEQ ID NO:128, SEQ ID NO:130, SEQ ID NO:132, SEQ ID NO:134, SEQ ID NO:136, SEQ ID NO:138, SEQ ID NO:140, SEQ ID NO:142, SEQ ID NO:143, SEQ ID NO:146, SEQ ID NO:148, SEQ ID NO:150, SEQ ID NO:152, SEQ ID NO:154, SEQ ID NO:156, SEQ ID NO:158, SEQ ID NO:160, SEQ ID NO:162, SEQ ID NO:164, SEQ ID NO:166, SEQ ID NO:168, SEQ ID NO:170, SEQ ID NO:172, SEQ ID NO:174, SEQ ID NO:176, SEQ ID NO:178, SEQ ID NO:180, SEQ ID NO:182, SEQ ID NO:184, SEQ ID NO:186, SEQ ID NO:188, SEQ ID NO:190, SEQ ID NO:192, SEQ ID NO:194, SEQ ID NO:196, SEQ ID NO:198, SEQ ID NO:200, SEQ ID NO:202, SEQ ID NO:204, SEQ ID NO:206, SEQ ID NO:208, SEQ ID NO:210, SEQ ID NO:212, SEQ ID NO:214, SEQ ID NO:216, SEQ ID NO:218, SEQ ID NO:220, SEQ ID NO:222, SEQ ID NO:224, SEQ ID NO:226, SEQ ID NO:228, SEQ ID NO:230, SEQ ID NO:232, SEQ ID NO:234, SEQ ID NO:236, SEQ ID NO:238, SEQ ID NO:240, SEQ ID NO:242, SEQ ID NO:244, SEQ ID NO:246, SEQ ID NO:248, SEQ ID NO:250, SEQ ID NO:252, SEQ ID NO:254, SEQ ID

NO:256, SEQ ID NO:258, SEQ ID NO:260, SEQ ID NO:262, SEQ ID NO:264, SEQ ID
NO:266, SEQ ID NO:268, SEQ ID NO:270, SEQ ID NO:272, SEQ ID NO:274, SEQ ID
NO:276, SEQ ID NO:278, SEQ ID NO:280, SEQ ID NO:282, SEQ ID NO:284, SEQ ID
NO:286, SEQ ID NO:288, SEQ ID NO:290, SEQ ID NO:292, SEQ ID NO:294, SEQ ID
5 NO:296, SEQ ID NO:298, SEQ ID NO:300, SEQ ID NO:302, SEQ ID NO:304, SEQ ID
NO:306, SEQ ID NO:308, SEQ ID NO:310, SEQ ID NO:312, SEQ ID NO:314, SEQ ID
NO:316, SEQ ID NO:318, SEQ ID NO:320, SEQ ID NO:322, SEQ ID NO:324, SEQ ID
NO:326, SEQ ID NO:328, SEQ ID NO:330, SEQ ID NO:332, SEQ ID NO:334, SEQ ID
NO:336, SEQ ID NO:338, SEQ ID NO:340, SEQ ID NO:342, SEQ ID NO:344, SEQ ID
10 NO:346, SEQ ID NO:348, SEQ ID NO:350, SEQ ID NO:352, SEQ ID NO:354, SEQ ID
NO:356, SEQ ID NO:358, SEQ ID NO:360, SEQ ID NO:362, SEQ ID NO:364, SEQ ID
NO:366, SEQ ID NO:368, SEQ ID NO:370, SEQ ID NO:372, SEQ ID NO:374, SEQ ID
NO:376, SEQ ID NO:378, SEQ ID NO:380, SEQ ID NO:382, SEQ ID NO:384, SEQ ID
NO:386, SEQ ID NO:388, SEQ ID NO:390, SEQ ID NO:392, SEQ ID NO:394, SEQ ID
15 NO:396, SEQ ID NO:398, SEQ ID NO:400, SEQ ID NO:402, SEQ ID NO:404, SEQ ID
NO:406, SEQ ID NO:408, SEQ ID NO:410, SEQ ID NO:412, SEQ ID NO:414, SEQ ID
NO:416, SEQ ID NO:418, SEQ ID NO:420, SEQ ID NO:422, SEQ ID NO:424, SEQ ID
NO:426, SEQ ID NO:428, SEQ ID NO:430, SEQ ID NO:432, SEQ ID NO:434, SEQ ID
NO:436, SEQ ID NO:438, SEQ ID NO:440, SEQ ID NO:442, SEQ ID NO:444, SEQ ID
20 NO:446, SEQ ID NO:448, SEQ ID NO:450, SEQ ID NO:452, SEQ ID NO:454, SEQ ID
NO:456, SEQ ID NO:458, SEQ ID NO:460, SEQ ID NO:462, SEQ ID NO:464, SEQ ID
NO:466, SEQ ID NO:468, SEQ ID NO:470, SEQ ID NO:472, SEQ ID NO:474, SEQ ID
NO:476, SEQ ID NO:478, SEQ ID NO:480, SEQ ID NO:482, SEQ ID NO:484, SEQ ID
NO:486, SEQ ID NO:488, SEQ ID NO:490, SEQ ID NO:492, SEQ ID NO:494, SEQ ID
25 NO:496 or SEQ ID NO:498, or enzymatically active fragments thereof;

(g) the nucleic acid (polynucleotide) of any of (a) to (f) and encoding a polypeptide
having at least one conservative amino acid substitution and retaining its isomerase activity,
e.g., racemase activity, e.g., amino acid racemase activity, alanine racemase activity, and/or
epimerase activity, wherein the at least one conservative amino acid substitution comprises
30 substituting an amino acid with another amino acid of like characteristics; or, a conservative
substitution comprises: replacement of an aliphatic amino acid with another aliphatic amino
acid; replacement of a Serine with a Threonine or vice versa; replacement of an acidic residue
with another acidic residue; replacement of a residue bearing an amide group with another

residue bearing an amide group; exchange of a basic residue with another basic residue; or replacement of an aromatic residue with another aromatic residue;

(h) the nucleic acid (polynucleotide) of any of (a) to (g) encoding a polypeptide having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity but lacking a signal sequence, a prepro domain, and/or other domain;

(i) the nucleic acid (polynucleotide) of any of (a) to (h) encoding a polypeptide having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity further comprising a heterologous sequence;

(j) the nucleic acid (polynucleotide) of (i), wherein the heterologous sequence comprises, or consists of a sequence encoding: (A) a heterologous signal sequence, a heterologous domain, a heterologous dockerin domain, a heterologous catalytic domain (CD), or a combination thereof; (B) the sequence of (i), wherein the heterologous signal sequence, domain or catalytic domain (CD) is derived from a heterologous enzyme; or, (C) a tag, an epitope, a targeting peptide, a cleavable sequence, a detectable moiety or an enzyme;

(k) the nucleic acid (polynucleotide) of (j), wherein the heterologous signal sequence targets the encoded protein to a vacuole, the endoplasmic reticulum, a chloroplast or a starch granule; or

(l) a nucleic acid sequence (polynucleotide) fully (completely) complementary to the sequence of any of (a) to (k).

The invention provides isolated, synthetic or recombinant nucleic acids comprising a nucleic acid encoding at least one polypeptide having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity, wherein the polypeptide has a sequence as set forth in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:22, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:28, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:42, SEQ ID NO:44, SEQ ID NO:46, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:58, SEQ ID NO:60, SEQ ID NO:62, SEQ ID NO:64, SEQ ID NO:66, SEQ ID NO:68, SEQ ID NO:70, SEQ ID NO:72, SEQ ID NO:74, SEQ ID NO:76, SEQ ID NO:78, SEQ ID NO:80, SEQ ID NO:82, SEQ ID NO:84, SEQ ID NO:86, SEQ ID NO:88, SEQ ID NO:90, SEQ ID NO:92, SEQ ID NO:94, SEQ ID NO:96, SEQ ID

NO:98, SEQ ID NO:100, SEQ ID NO:102, SEQ ID NO:104, SEQ ID NO:106, SEQ ID
NO:108, SEQ ID NO:110, SEQ ID NO:112, SEQ ID NO:114, SEQ ID NO:116, SEQ ID
NO:118, SEQ ID NO:120, SEQ ID NO:122, SEQ ID NO:124, SEQ ID NO:126, SEQ ID
NO:128, SEQ ID NO:130, SEQ ID NO:132, SEQ ID NO:134, SEQ ID NO:136, SEQ ID
5 NO:138, SEQ ID NO:140, SEQ ID NO:142, SEQ ID NO:143, SEQ ID NO:146, SEQ ID
NO:148, SEQ ID NO:150, SEQ ID NO:152, SEQ ID NO:154, SEQ ID NO:156, SEQ ID
NO:158, SEQ ID NO:160, SEQ ID NO:162, SEQ ID NO:164, SEQ ID NO:166, SEQ ID
NO:168, SEQ ID NO:170, SEQ ID NO:172, SEQ ID NO:174, SEQ ID NO:176, SEQ ID
NO:178, SEQ ID NO:180, SEQ ID NO:182, SEQ ID NO:184, SEQ ID NO:186, SEQ ID
10 NO:188, SEQ ID NO:190, SEQ ID NO:192, SEQ ID NO:194, SEQ ID NO:196, SEQ ID
NO:198, SEQ ID NO:200, SEQ ID NO:202, SEQ ID NO:204, SEQ ID NO:206, SEQ ID
NO:208, SEQ ID NO:210, SEQ ID NO:212, SEQ ID NO:214, SEQ ID NO:216, SEQ ID
NO:218, SEQ ID NO:220, SEQ ID NO:222, SEQ ID NO:224, SEQ ID NO:226, SEQ ID
NO:228, SEQ ID NO:230, SEQ ID NO:232, SEQ ID NO:234, SEQ ID NO:236, SEQ ID
15 NO:238, SEQ ID NO:240, SEQ ID NO:242, SEQ ID NO:244, SEQ ID NO:246, SEQ ID
NO:248, SEQ ID NO:250, SEQ ID NO:252, SEQ ID NO:254, SEQ ID NO:256, SEQ ID
NO:258, SEQ ID NO:260, SEQ ID NO:262, SEQ ID NO:264, SEQ ID NO:266, SEQ ID
NO:268, SEQ ID NO:270, SEQ ID NO:272, SEQ ID NO:274, SEQ ID NO:276, SEQ ID
NO:278, SEQ ID NO:280, SEQ ID NO:282, SEQ ID NO:284, SEQ ID NO:286, SEQ ID
20 NO:288, SEQ ID NO:290, SEQ ID NO:292, SEQ ID NO:294, SEQ ID NO:296, SEQ ID
NO:298, SEQ ID NO:300, SEQ ID NO:302, SEQ ID NO:304, SEQ ID NO:306, SEQ ID
NO:308, SEQ ID NO:310, SEQ ID NO:312, SEQ ID NO:314, SEQ ID NO:316, SEQ ID
NO:318, SEQ ID NO:320, SEQ ID NO:322, SEQ ID NO:324, SEQ ID NO:326, SEQ ID
NO:328, SEQ ID NO:330, SEQ ID NO:332, SEQ ID NO:334, SEQ ID NO:336, SEQ ID
25 NO:338, SEQ ID NO:340, SEQ ID NO:342, SEQ ID NO:344, SEQ ID NO:346, SEQ ID
NO:348, SEQ ID NO:350, SEQ ID NO:352, SEQ ID NO:354, SEQ ID NO:356, SEQ ID
NO:358, SEQ ID NO:360, SEQ ID NO:362, SEQ ID NO:364, SEQ ID NO:366, SEQ ID
NO:368, SEQ ID NO:370, SEQ ID NO:372, SEQ ID NO:374, SEQ ID NO:376, SEQ ID
NO:378, SEQ ID NO:380, SEQ ID NO:382, SEQ ID NO:384, SEQ ID NO:386, SEQ ID
30 NO:388, SEQ ID NO:390, SEQ ID NO:392, SEQ ID NO:394, SEQ ID NO:396, SEQ ID
NO:398, SEQ ID NO:400, SEQ ID NO:402, SEQ ID NO:404, SEQ ID NO:406, SEQ ID
NO:408, SEQ ID NO:410, SEQ ID NO:412, SEQ ID NO:414, SEQ ID NO:416, SEQ ID
NO:418, SEQ ID NO:420, SEQ ID NO:422, SEQ ID NO:424, SEQ ID NO:426, SEQ ID
NO:428, SEQ ID NO:430, SEQ ID NO:432, SEQ ID NO:434, SEQ ID NO:436, SEQ ID

NO:438, SEQ ID NO:440, SEQ ID NO:442, SEQ ID NO:444, SEQ ID NO:446, SEQ ID NO:448, SEQ ID NO:450, SEQ ID NO:452, SEQ ID NO:454, SEQ ID NO:456, SEQ ID NO:458, SEQ ID NO:460, SEQ ID NO:462, SEQ ID NO:464, SEQ ID NO:466, SEQ ID NO:468, SEQ ID NO:470, SEQ ID NO:472, SEQ ID NO:474, SEQ ID NO:476, SEQ ID NO:478, SEQ ID NO:480, SEQ ID NO:482, SEQ ID NO:484, SEQ ID NO:486, SEQ ID NO:488, SEQ ID NO:490, SEQ ID NO:492, SEQ ID NO:494, SEQ ID NO:496 or SEQ ID NO:498, or enzymatically active fragments thereof, including the sequences described herein and in Tables 1, 2 and 3, and the Sequence Listing (all of these sequences are “exemplary enzymes/ polypeptides of the invention”), and enzymatically active subsequences (fragments) thereof and/or immunologically active subsequences thereof (such as epitopes or immunogens) (all “peptides of the invention”) and variants thereof (all of these sequences encompassing polypeptide and peptide sequences of the invention) (or, hereinafter referred to as the exemplary polypeptide sequences of the inventions).

The invention provides isolated, synthetic or recombinant nucleic acids comprising sequences completely complementary to all of these nucleic acid sequences of the invention (complementary (non-coding) and coding sequences also hereinafter collectively referred to as nucleic acid sequences of the invention).

In one aspect, the sequence identity is at least about 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% (complete) sequence identity (homology). In one aspect, the sequence identity is over a region of at least about 150, 175, 200, 225, 250, 275, 300, 350, 400, 450, 500, 550, 600, 650, 700, 750, 800, 850, 900, 950, 1000, 1050, 1100, 1150 or more residues, or the full length of a gene or a transcript. For example, the invention provides isolated, synthetic or recombinant nucleic acids comprising a nucleic acid sequence of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO:31, SEQ ID NO:33, SEQ ID NO:35, SEQ ID NO:37, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:43, SEQ ID NO:45, SEQ ID NO:47, SEQ ID NO:49, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:57, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ

ID NO:89, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95, SEQ ID NO:97, SEQ ID
NO:99, SEQ ID NO:101, SEQ ID NO:103, SEQ ID NO:105, SEQ ID NO:107, SEQ ID
NO:109, SEQ ID NO:111, SEQ ID NO:113, SEQ ID NO:115, SEQ ID NO:117, SEQ ID
NO:119, SEQ ID NO:121, SEQ ID NO:123, SEQ ID NO:125, SEQ ID NO:127, SEQ ID
5 NO:129, SEQ ID NO:131, SEQ ID NO:133, SEQ ID NO:135, SEQ ID NO:137, SEQ ID
NO:139, SEQ ID NO:141, SEQ ID NO:143, SEQ ID NO:145, SEQ ID NO:147, SEQ ID
NO:149, SEQ ID NO:151, SEQ ID NO:153, SEQ ID NO:155, SEQ ID NO:157, SEQ ID
NO:159, SEQ ID NO:161, SEQ ID NO:163, SEQ ID NO:165, SEQ ID NO:167, SEQ ID
NO:169, SEQ ID NO:171, SEQ ID NO:173, SEQ ID NO:175, SEQ ID NO:177, SEQ ID
10 NO:179, SEQ ID NO:181, SEQ ID NO:183, SEQ ID NO:185, SEQ ID NO:187, SEQ ID
NO:189, SEQ ID NO:191, SEQ ID NO:193, SEQ ID NO:195, SEQ ID NO:197, SEQ ID
NO:199, SEQ ID NO:201, SEQ ID NO:203, SEQ ID NO:205, SEQ ID NO:207, SEQ ID
NO:209, SEQ ID NO:211, SEQ ID NO:213, SEQ ID NO:215, SEQ ID NO:217, SEQ ID
NO:219, SEQ ID NO:221, SEQ ID NO:223, SEQ ID NO:225, SEQ ID NO:227, SEQ ID
15 NO:229, SEQ ID NO:231, SEQ ID NO:233, SEQ ID NO:235, SEQ ID NO:237, SEQ ID
NO:239, SEQ ID NO:241, SEQ ID NO:243, SEQ ID NO:245, SEQ ID NO:247, SEQ ID
NO:249, SEQ ID NO:251, SEQ ID NO:253, SEQ ID NO:255, SEQ ID NO:257, SEQ ID
NO:259, SEQ ID NO:261, SEQ ID NO:263, SEQ ID NO:265, SEQ ID NO:267, SEQ ID
NO:269, SEQ ID NO:271, SEQ ID NO:273, SEQ ID NO:275, SEQ ID NO:277, SEQ ID
20 NO:279, SEQ ID NO:281, SEQ ID NO:283, SEQ ID NO:285, SEQ ID NO:287, SEQ ID
NO:289, SEQ ID NO:291, SEQ ID NO:293, SEQ ID NO:295, SEQ ID NO:297, SEQ ID
NO:299, SEQ ID NO:301, SEQ ID NO:303, SEQ ID NO:305, SEQ ID NO:307, SEQ ID
NO:309, SEQ ID NO:311, SEQ ID NO:313, SEQ ID NO:315, SEQ ID NO:317, SEQ ID
NO:319, SEQ ID NO:321, SEQ ID NO:323, SEQ ID NO:325, SEQ ID NO:327, SEQ ID
25 NO:329, SEQ ID NO:331, SEQ ID NO:333, SEQ ID NO:335, SEQ ID NO:337, SEQ ID
NO:339, SEQ ID NO:341, SEQ ID NO:343, SEQ ID NO:345, SEQ ID NO:347, SEQ ID
NO:349, SEQ ID NO:351, SEQ ID NO:353, SEQ ID NO:355, SEQ ID NO:357, SEQ ID
NO:359, SEQ ID NO:361, SEQ ID NO:363, SEQ ID NO:365, SEQ ID NO:367, SEQ ID
NO:369, SEQ ID NO:371, SEQ ID NO:373, SEQ ID NO:375, SEQ ID NO:377, SEQ ID
30 NO:379, SEQ ID NO:381, SEQ ID NO:383, SEQ ID NO:385, SEQ ID NO:387, SEQ ID
NO:389, SEQ ID NO:391, SEQ ID NO:393, SEQ ID NO:395, SEQ ID NO:397, SEQ ID
NO:399, SEQ ID NO:401, SEQ ID NO:403, SEQ ID NO:405, SEQ ID NO:407, SEQ ID
NO:409, SEQ ID NO:411, SEQ ID NO:413, SEQ ID NO:415, SEQ ID NO:417, SEQ ID
NO:419, SEQ ID NO:421, SEQ ID NO:423, SEQ ID NO:425, SEQ ID NO:427, SEQ ID

NO:429, SEQ ID NO:431, SEQ ID NO:433, SEQ ID NO:435, SEQ ID NO:437, SEQ ID NO:439, SEQ ID NO:441, SEQ ID NO:443, SEQ ID NO:445, SEQ ID NO:447, SEQ ID NO:449, SEQ ID NO:451, SEQ ID NO:453, SEQ ID NO:455, SEQ ID NO:457, SEQ ID NO:459, SEQ ID NO:461, SEQ ID NO:463, SEQ ID NO:465, SEQ ID NO:467, SEQ ID NO:469, SEQ ID NO:471, SEQ ID NO:473, SEQ ID NO:475, SEQ ID NO:477, SEQ ID NO:479, SEQ ID NO:481, SEQ ID NO:483, SEQ ID NO:485, SEQ ID NO:487, SEQ ID NO:489, SEQ ID NO:491, SEQ ID NO:493, SEQ ID NO:495 or SEQ ID NO:497, e.g., as described in Tables 1, 2 and 3 and in the Sequence Listing (all of these sequences are “exemplary p of the invention”), and enzymatically active subsequences (fragments) thereof.

10 The invention provides isolated, synthetic or recombinant nucleic acids encoding a polypeptide having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity, wherein the nucleic acid has at least one sequence modification of an exemplary sequence of the invention, or, any sequence of the invention.

15 In one aspect (optionally), the isolated, synthetic or recombinant nucleic acids of the invention have an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity, e.g., wherein the activity comprises catalyzing the re-arrangement of atoms within a molecule, catalyzing the conversion of one isomer into another, catalyzing the conversion of an
20 optically active substrate into a raceme, which is optically inactive, catalyzing the interconversion of substrate enantiomers, catalyzing the stereochemical inversion around the asymmetric carbon atom in a substrate having only one center of asymmetry, catalyzing the stereochemical inversion of the configuration around an asymmetric carbon atom in a substrate having more than one asymmetric center, and/or catalyzing the
25 racemization of amino acids.

In one aspect, the isomerase activity, e.g., racemase activity, e.g., amino acid racemase activity, alanine racemase activity, and/or epimerase activity is thermostable, e.g., wherein the polypeptide retains an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase
30 activity under conditions comprising a temperature range from about -100°C to about -80°C, about -80°C to about -40°C, about -40°C to about -20°C, about -20°C to about 0°C, about 0°C to about 5°C, about 5°C to about 15°C, about 15°C to about 25°C, about 25°C to about 37°C, about 37°C to about 45°C, about 45°C to about 55°C, about 55°C to about 70°C, about 70°C to about 75°C, about 75°C to about 85°C, about 85°C to about 90°C,

about 90°C to about 95°C, about 95°C to about 100°C, about 100°C to about 105°C, about 105°C to about 110°C, about 110°C to about 120°C, or 95°C, 96°C, 97°C, 98°C, 99°C, 100°C, 101°C, 102°C, 103°C, 104°C, 105°C, 106°C, 107°C, 108°C, 109°C, 110°C, 111°C, 112°C, 113°C, 114°C, 115°C or more. In some embodiments, the thermostable
5 polypeptides according to the invention retains activity, e.g., an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity, at a temperature in the ranges described above, at about pH 3.0, about pH 3.5, about pH 4.0, about pH 4.5, about pH 5.0, about pH 5.5, about pH 6.0, about pH 6.5, about pH 7.0, about pH 7.5, about pH 8.0, about pH 8.5, about pH 9.0, about
10 pH 9.5, about pH 10.0, about pH 10.5, about pH 11.0, about pH 11.5, about pH 12.0 or more.

In one aspect, the isomerase activity, e.g., racemase activity, e.g., amino acid racemase activity, alanine racemase activity, and/or epimerase activity is thermotolerant, e.g., wherein the polypeptide retains an isomerase activity, e.g., a racemase activity, e.g.,
15 an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity after exposure to a temperature in the range from about -100°C to about -80°C, about -80°C to about -40°C, about -40°C to about -20°C, about -20°C to about 0°C, about 0°C to about 5°C, about 5°C to about 15°C, about 15°C to about 25°C, about 25°C to about 37°C, about 37°C to about 45°C, about 45°C to about 55°C, about 55°C to about 70°C,
20 about 70°C to about 75°C, about 75°C to about 85°C, about 85°C to about 90°C, about 90°C to about 95°C, about 95°C to about 100°C, about 100°C to about 105°C, about 105°C to about 110°C, about 110°C to about 120°C, or 95°C, 96°C, 97°C, 98°C, 99°C, 100°C, 101°C, 102°C, 103°C, 104°C, 105°C, 106°C, 107°C, 108°C, 109°C, 110°C, 111°C, 112°C, 113°C, 114°C, 115°C or more. The thermotolerant polypeptides according to the
25 invention can retain activity, e.g. an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity, after exposure to a temperature in the range from about -100°C to about -80°C, about -80°C to about -40°C, about -40°C to about -20°C, about -20°C to about 0°C, about 0°C to about 5°C, about 5°C to about 15°C, about 15°C to about 25°C, about 25°C to about 37°C,
30 about 37°C to about 45°C, about 45°C to about 55°C, about 55°C to about 70°C, about 70°C to about 75°C, about 75°C to about 85°C, about 85°C to about 90°C, about 90°C to about 95°C, about 95°C to about 100°C, about 100°C to about 105°C, about 105°C to about 110°C, about 110°C to about 120°C, or 95°C, 96°C, 97°C, 98°C, 99°C, 100°C, 101°C, 102°C, 103°C, 104°C, 105°C, 106°C, 107°C, 108°C, 109°C, 110°C, 111°C, 112°C,

113°C, 114°C, 115°C or more. In some embodiments, the thermotolerant polypeptides according to the invention retains activity, e.g. an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity, after exposure to a temperature in the ranges described above, at about
5 pH 3.0, about pH 3.5, about pH 4.0, about pH 4.5, about pH 5.0, about pH 5.5, about pH 6.0, about pH 6.5, about pH 7.0, about pH 7.5, about pH 8.0, about pH 8.5, about pH 9.0, about pH 9.5, about pH 10.0, about pH 10.5, about pH 11.0, about pH 11.5, about pH 12.0 or more.

In one aspect, the isomerase activity, e.g., racemase activity, e.g., amino acid
10 racemase activity, alanine racemase activity, and/or epimerase activity of polypeptides encoded by nucleic acids of the invention retain activity under acidic conditions comprising about pH 6.5, pH 6, pH 5.5, pH 5, pH 4.5, pH 4.0, pH 3.5, pH 3.0 or less (more acidic) pH, or, retain an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity after
15 exposure to acidic conditions comprising about pH 6.5, pH 6, pH 5.5, pH 5, pH 4.5, pH 4.0, pH 3.5, pH 3.0 or less (more acidic) pH; or, retain activity under basic conditions comprising about pH 7, pH 7.5 pH 8.0, pH 8.5, pH 9, pH 9.5, pH 10, pH 10.5, pH 11, pH 11.5, pH 12, pH 12.5 or more (more basic) or, retain an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity,
20 and/or an epimerase activity after exposure to basic conditions comprising about pH 7, pH 7.5 pH 8.0, pH 8.5, pH 9, pH 9.5, pH 10, pH 10.5, pH 11, pH 11.5, pH 12, pH 12.5 or more (more basic). In one aspect, isomerase activity, e.g., racemase activity, e.g., amino acid racemase activity, alanine racemase activity, and/or epimerase activity of polypeptides encoded by nucleic acids of the invention retain activity at a temperature of
25 at least about 80°C, 81°C, 82°C, 83°C, 84°C, 85°C, 86°C, 87°C, 88°C, 89°C, 90°C, 91°C, 92°C, 93°C, 94°C, 95°C, 96°C, 97°C, 98°C, 99°C, 100°C, 101°C, 102°C, 103°C, 103.5°C, 104°C, 105°C, 107°C, 108°C, 109°C or 110°C, or more, and a basic pH of at least about pH 7.5 pH 8.0, pH 8.5, pH 9, pH 9.5, pH 10, pH 10.5, pH 11, pH 11.5, pH 12, pH 12.5 or more (more basic).

30 The invention provides expression cassettes, cloning vehicles, or a vector (e.g., expression vectors) comprising a nucleic acid comprising a sequence of the invention. The cloning vehicle can comprise a viral vector, a plasmid, a phage, a phagemid, a cosmid, a fosmid, a bacteriophage or an artificial chromosome. The viral vector can comprise an adenovirus vector, a retroviral vector or an adeno-associated viral vector. The

cloning vehicle can comprise an artificial chromosome comprising a bacterial artificial chromosome (BAC), a bacteriophage P1-derived vector (PAC), a yeast artificial chromosome (YAC), or a mammalian artificial chromosome (MAC).

5 The invention provides nucleic acid probes for identifying a nucleic acid encoding a polypeptide with an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity, wherein the probe comprises at least about 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 100, 125, 150, 175, 200, 225, 250, 275, 300 or more consecutive bases of a nucleic acid comprising an exemplary sequence of the invention, or, any sequence of the invention (as
10 defined herein), wherein in one aspect (optionally) the probe comprises an oligonucleotide comprising between at least about 10 to 300, about 25 to 250, about 10 to 50, about 20 to 60, about 30 to 70, about 40 to 80, about 60 to 100, or about 50 to 150 or more consecutive bases.

The invention provides amplification primer pairs for amplifying a nucleic acid
15 encoding a polypeptide having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity, wherein the primer pair is capable of amplifying a nucleic acid comprising an exemplary sequence of the invention, or, any sequence of the invention (as defined herein), or a subsequence thereof, wherein optionally a member of the amplification primer sequence
20 pair comprises an oligonucleotide comprising at least about 10 to 50 consecutive bases of the sequence, or, about 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 or more consecutive bases of the sequence. The invention provides amplification primer pairs wherein the primer pair comprises a first member having a sequence as set forth by about the first (the 5') 10, 11, 12, 13, 14, 15, 16, 17, 18,
25 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 or more residues of an exemplary sequence of the invention, or, any sequence of the invention (as defined herein), and a second member having a sequence as set forth by about the first (the 5') 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 or more residues of the complementary strand of the first member.

30 The invention provides an isomerase-, e.g., a racemase-, e.g., an amino acid racemase-, an alanine racemase-, and/or an epimerase- encoding nucleic acids generated by amplification of a polynucleotide using an amplification primer pair of the invention, wherein optionally the amplification is by polymerase chain reaction (PCR). In one aspect, the nucleic acid is generated by amplification of a gene library, wherein in one

aspect (optionally) the gene library is an environmental library. The invention provides isolated, synthetic or recombinant isomerases, e.g., racemases, e.g., amino acid racemases, alanine racemases, and/or epimerases encoded by an isomerase-, e.g., a racemase-, e.g., an amino acid racemase-, an alanine racemase-, and/or an epimerase- encoding nucleic acid
5 generated by amplification of a polynucleotide using an amplification primer pair of the invention. The invention provides methods of amplifying a nucleic acid encoding a polypeptide having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity, the methods comprising the step of amplification of a template nucleic acid with an amplification
10 primer sequence pair capable of amplifying an exemplary sequence of the invention, or, any sequence of the invention (as defined herein), or a subsequence thereof.

The invention provides expression cassette, a vector or a cloning vehicle comprising a nucleic acid comprising a sequence of the invention, wherein optionally the cloning vehicle comprises a viral vector, a plasmid, a phage, a phagemid, a cosmid, a
15 fosmid, a bacteriophage or an artificial chromosome. The viral vector can comprise an adenovirus vector, a retroviral vector or an adeno-associated viral vector, or, the artificial chromosome comprises a bacterial artificial chromosome (BAC), a bacteriophage P1-derived vector (PAC), a yeast artificial chromosome (YAC), or a mammalian artificial chromosome (MAC).

20 The invention provides transformed cells comprising a nucleic acid or vector of the invention, or an expression cassette or cloning vehicle of the invention. The transformed cell can be a bacterial cell, a mammalian cell, a fungal cell, a yeast cell, an insect cell or a plant cell.

The invention provides transgenic non-human animals comprising a sequence of
25 the invention. The transgenic non-human animal can be a mouse, a rat, a rabbit, a sheep, a pig, a chicken, a goat, a fish, a dog, or a cow. The invention provides transgenic plants comprising a sequence of the invention, e.g., wherein the plant is a corn plant, a sorghum plant, a potato plant, a tomato plant, a wheat plant, an oilseed plant, a rapeseed plant, a soybean plant, a rice plant, a barley plant, a grass, or a tobacco plant. The invention
30 provides transgenic seeds comprising a sequence of the invention, e.g., wherein the seed is a corn seed, a wheat kernel, an oilseed, a rapeseed, a soybean seed, a palm kernel, a sunflower seed, a sesame seed, a rice, a barley, a peanut or a tobacco plant seed.

The invention provides antisense oligonucleotides comprising a nucleic acid sequence complementary to or capable of hybridizing under stringent conditions to a sequence of the invention (including, e.g., exemplary sequences of the invention), or a subsequence thereof, wherein optionally the antisense oligonucleotide is between about 10 to 50, about 20 to 60, about 30 to 70, about 40 to 80, or about 60 to 100 bases in length, and in one aspect (optionally) the stringent conditions comprise a wash step comprising a wash in 0.2X SSC at a temperature of about 65°C for about 15 minutes.

The invention provides methods of inhibiting the translation of an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase message in a cell comprising administering to the cell or expressing in the cell an antisense oligonucleotide comprising a nucleic acid sequence complementary to or capable of hybridizing under stringent conditions to a sequence of the invention (including, e.g., exemplary sequences of the invention).

The invention provides double-stranded inhibitory RNA (RNAi) molecules comprising a subsequence of a sequence of the invention (including, e.g., exemplary sequences of the invention). The double-stranded inhibitory RNA (RNAi) molecule can be about 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, or 30 or more duplex nucleotides in length. The invention provides methods of inhibiting the expression of an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase in a cell comprising administering to the cell or expressing in the cell a double-stranded inhibitory RNA (iRNA), wherein the RNA comprises a subsequence of a sequence of the invention (including, e.g., exemplary sequences of the invention).

The invention provides isolated, synthetic or recombinant polypeptides having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity, or polypeptides capable of generating an immune response specific for an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase (e.g., an epitope); and in alternative aspects peptide and polypeptide of the invention comprise a sequence:

(a) comprising an amino acid sequence having at least about 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more, or has 100% (complete) sequence identity to SEQ ID NO:2, SEQ

ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID
NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:22, SEQ ID
NO:24, SEQ ID NO:26, SEQ ID NO:28, SEQ ID NO:30, SEQ ID NO:32, SEQ ID
NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:42, SEQ ID
5 NO:44, SEQ ID NO:46, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:52, SEQ ID
NO:54, SEQ ID NO:56, SEQ ID NO:58, SEQ ID NO:60, SEQ ID NO:62, SEQ ID
NO:64, SEQ ID NO:66, SEQ ID NO:68, SEQ ID NO:70, SEQ ID NO:72, SEQ ID
NO:74, SEQ ID NO:76, SEQ ID NO:78, SEQ ID NO:80, SEQ ID NO:82, SEQ ID
NO:84, SEQ ID NO:86, SEQ ID NO:88, SEQ ID NO:90, SEQ ID NO:92, SEQ ID
10 NO:94, SEQ ID NO:96, SEQ ID NO:98, SEQ ID NO:100, SEQ ID NO:102, SEQ ID
NO:104, SEQ ID NO:106, SEQ ID NO:108, SEQ ID NO:110, SEQ ID NO:112, SEQ
ID NO:114, SEQ ID NO:116, SEQ ID NO:118, SEQ ID NO:120, SEQ ID NO:122,
SEQ ID NO:124, SEQ ID NO:126, SEQ ID NO:128, SEQ ID NO:130, SEQ ID
NO:132, SEQ ID NO:134, SEQ ID NO:136, SEQ ID NO:138, SEQ ID NO:140, SEQ
15 ID NO:142, SEQ ID NO:143, SEQ ID NO:146, SEQ ID NO:148, SEQ ID NO:150,
SEQ ID NO:152, SEQ ID NO:154, SEQ ID NO:156, SEQ ID NO:158, SEQ ID
NO:160, SEQ ID NO:162, SEQ ID NO:164, SEQ ID NO:166, SEQ ID NO:168, SEQ
ID NO:170, SEQ ID NO:172, SEQ ID NO:174, SEQ ID NO:176, SEQ ID NO:178,
SEQ ID NO:180, SEQ ID NO:182, SEQ ID NO:184, SEQ ID NO:186, SEQ ID
20 NO:188, SEQ ID NO:190, SEQ ID NO:192, SEQ ID NO:194, SEQ ID NO:196, SEQ
ID NO:198, SEQ ID NO:200, SEQ ID NO:202, SEQ ID NO:204, SEQ ID NO:206,
SEQ ID NO:208, SEQ ID NO:210, SEQ ID NO:212, SEQ ID NO:214, SEQ ID
NO:216, SEQ ID NO:218, SEQ ID NO:220, SEQ ID NO:222, SEQ ID NO:224, SEQ
ID NO:226, SEQ ID NO:228, SEQ ID NO:230, SEQ ID NO:232, SEQ ID NO:234,
25 SEQ ID NO:236, SEQ ID NO:238, SEQ ID NO:240, SEQ ID NO:242, SEQ ID
NO:244, SEQ ID NO:246, SEQ ID NO:248, SEQ ID NO:250, SEQ ID NO:252, SEQ
ID NO:254, SEQ ID NO:256, SEQ ID NO:258, SEQ ID NO:260, SEQ ID NO:262,
SEQ ID NO:264, SEQ ID NO:266, SEQ ID NO:268, SEQ ID NO:270, SEQ ID
NO:272, SEQ ID NO:274, SEQ ID NO:276, SEQ ID NO:278, SEQ ID NO:280, SEQ
30 ID NO:282, SEQ ID NO:284, SEQ ID NO:286, SEQ ID NO:288, SEQ ID NO:290,
SEQ ID NO:292, SEQ ID NO:294, SEQ ID NO:296, SEQ ID NO:298, SEQ ID
NO:300, SEQ ID NO:302, SEQ ID NO:304, SEQ ID NO:306, SEQ ID NO:308, SEQ
ID NO:310, SEQ ID NO:312, SEQ ID NO:314, SEQ ID NO:316, SEQ ID NO:318,
SEQ ID NO:320, SEQ ID NO:322, SEQ ID NO:324, SEQ ID NO:326, SEQ ID

NO:328, SEQ ID NO:330, SEQ ID NO:332, SEQ ID NO:334, SEQ ID NO:336, SEQ ID NO:338, SEQ ID NO:340, SEQ ID NO:342, SEQ ID NO:344, SEQ ID NO:346, SEQ ID NO:348, SEQ ID NO:350, SEQ ID NO:352, SEQ ID NO:354, SEQ ID NO:356, SEQ ID NO:358, SEQ ID NO:360, SEQ ID NO:362, SEQ ID NO:364, SEQ ID NO:366, SEQ ID NO:368, SEQ ID NO:370, SEQ ID NO:372, SEQ ID NO:374, SEQ ID NO:376, SEQ ID NO:378, SEQ ID NO:380, SEQ ID NO:382, SEQ ID NO:384, SEQ ID NO:386, SEQ ID NO:388, SEQ ID NO:390, SEQ ID NO:392, SEQ ID NO:394, SEQ ID NO:396, SEQ ID NO:398, SEQ ID NO:400, SEQ ID NO:402, SEQ ID NO:404, SEQ ID NO:406, SEQ ID NO:408, SEQ ID NO:410, SEQ ID NO:412, SEQ ID NO:414, SEQ ID NO:416, SEQ ID NO:418, SEQ ID NO:420, SEQ ID NO:422, SEQ ID NO:424, SEQ ID NO:426, SEQ ID NO:428, SEQ ID NO:430, SEQ ID NO:432, SEQ ID NO:434, SEQ ID NO:436, SEQ ID NO:438, SEQ ID NO:440, SEQ ID NO:442, SEQ ID NO:444, SEQ ID NO:446, SEQ ID NO:448, SEQ ID NO:450, SEQ ID NO:452, SEQ ID NO:454, SEQ ID NO:456, SEQ ID NO:458, SEQ ID NO:460, SEQ ID NO:462, SEQ ID NO:464, SEQ ID NO:466, SEQ ID NO:468, SEQ ID NO:470, SEQ ID NO:472, SEQ ID NO:474, SEQ ID NO:476, SEQ ID NO:478, SEQ ID NO:480, SEQ ID NO:482, SEQ ID NO:484, SEQ ID NO:486, SEQ ID NO:488, SEQ ID NO:490, SEQ ID NO:492, SEQ ID NO:494, SEQ ID NO:496 or SEQ ID NO:498, or enzymatically active fragments thereof,

wherein the polypeptide or peptide of (i) or (ii) has an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity, or the polypeptide or peptide is capable of generating an isomerase specific antibody, e.g., a racemase specific antibody, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase specific antibody (a polypeptide or peptide that acts as an epitope or immunogen),

(b) the polypeptide or peptide of (a), wherein the sequence identities are determined: (A) by analysis with a sequence comparison algorithm or by a visual inspection, or (B) over a region of at least about 20, 25, 30, 35, 40, 45, 50, 55, 60, 75, 100, 150, 200, 250, 300 or more amino acid residues, or over the full length of the polypeptide or peptide or enzyme, and/or enzymatically active subsequences (fragments) thereof,

(c) the polypeptide or peptide of (a) of (b), wherein the sequence identities are determined by analysis with a sequence comparison algorithm or by a visual inspection, and optionally the sequence comparison algorithm is a BLAST version 2.2.2 algorithm where

a filtering setting is set to blastall -p blastp -d "nr pataa" -F F, and all other options are set to default;

(d) an amino acid sequence encoded by the nucleic acid of claim 1, wherein the polypeptide has (i) an isomerase activity, e.g., a racemase activity, e.g., an amino acid
5 racemase activity, an alanine racemase activity, and/or an epimerase activity, or, (ii) has immunogenic activity in that it is capable of generating an antibody that specifically binds to a polypeptide having a sequence of (a), and/or enzymatically active subsequences (fragments) thereof;

(e) the amino acid sequence of any of (a) to (d), and comprising at least one amino
10 acid residue conservative substitution, and the polypeptide or peptide retains isomerase activity, e.g., racemase activity, e.g., amino acid racemase activity, alanine racemase activity, and/or epimerase activity;

(e) the amino acid sequence of (d), wherein the conservative substitution comprises replacement of an aliphatic amino acid with another aliphatic amino acid; replacement of a
15 serine with a threonine or vice versa; replacement of an acidic residue with another acidic residue; replacement of a residue bearing an amide group with another residue bearing an amide group; exchange of a basic residue with another basic residue; or, replacement of an aromatic residue with another aromatic residue, or a combination thereof,

(f) the amino acid sequence of (e), wherein the aliphatic residue comprises Alanine,
20 Valine, Leucine, Isoleucine or a synthetic equivalent thereof; the acidic residue comprises Aspartic acid, Glutamic acid or a synthetic equivalent thereof; the residue comprising an amide group comprises Aspartic acid, Glutamic acid or a synthetic equivalent thereof; the basic residue comprises Lysine, Arginine or a synthetic equivalent thereof; or, the aromatic residue comprises Phenylalanine, Tyrosine or a synthetic equivalent thereof;

(g) the polypeptide of any of (a) to (f) having an isomerase activity, e.g., a racemase
25 activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity but lacking a signal sequence, a prepro domain, and/or other domain,

(h) the polypeptide of any of (a) to (g) having an isomerase activity, e.g., a racemase
30 activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity further comprising a heterologous sequence;

(i) the polypeptide of (h), wherein the heterologous sequence comprises, or consists of:
(A) a heterologous signal sequence, a heterologous domain, a heterologous dockerin domain, a heterologous catalytic domain (CD), or a combination thereof; (B) the sequence of (A), wherein the heterologous signal sequence, domain or catalytic domain (CD) is derived from a

heterologous enzyme; and/or, (C) a tag, an epitope, a targeting peptide, a cleavable sequence, a detectable moiety or an enzyme;

(j) polypeptide of (i), wherein the heterologous signal sequence targets the encoded protein to a vacuole, the endoplasmic reticulum, a chloroplast or a starch granule; or

5 (k) comprising an amino acid sequence encoded any nucleic acid sequence of this invention.

In one aspect, the isomerase activity, e.g., racemase activity, e.g., amino acid racemase activity, alanine racemase activity, and/or epimerase activity comprises catalyzing the re-arrangement of atoms within a molecule, catalyzing the conversion of
10 one isomer into another, catalyzing the conversion of an optically active substrate into a raceme, which is optically inactive, catalyzing the interconversion of substrate enantiomers, catalyzing the stereochemical inversion around the asymmetric carbon atom in a substrate having only one center of asymmetry, catalyzing the stereochemical
15 inversion of the configuration around an asymmetric carbon atom in a substrate having more than one asymmetric center, and/or catalyzing the racemization of amino acids.

The invention provides isolated, synthetic or recombinant polypeptides comprising a polypeptide of the invention and lacking a signal sequence or a prepro sequence. The invention provides isolated, synthetic or recombinant polypeptides comprising a
20 polypeptide of the invention and having a heterologous signal sequence or a heterologous prepro sequence.

In one aspect, a polypeptide of the invention has isomerase activity, e.g., racemase activity, e.g., amino acid racemase activity, alanine racemase activity, and/or epimerase activity comprising a specific activity at about 37°C in the range from about 100 to about 1000 units per milligram of protein, from about 500 to about 750 units per milligram of
25 protein, from about 500 to about 1200 units per milligram of protein, or from about 750 to about 1000 units per milligram of protein. In alternative aspects, polypeptides of the invention have isomerase activity, e.g., racemase activity, e.g., amino acid racemase activity, alanine racemase activity, and/or epimerase activity in the range of between about 0.05 to 20 units per gram, or 0.05, 0.10, 0.20, 0.30, 0.40, 0.50, 0.60, 0.70, 0.80, 0.90, 1.0,
30 1.5, 2.0, 2.5, 3.0, 3.5, 4.0, 4.5, 5.0, 5.5, 6.0, 6.5, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20 or more units per gram, where a unit equals one μmol of product released per minute per mg of enzyme. In one embodiment, for racemases, one unit of activity equals one μmol of an isomer with inverted configuration (from the starting isomer) produced per minute per mg of enzyme (formed from the respective alpha-amino acid or amine). In an

alternative embodiment, for amino acid racemases, one unit of activity equals one umol of R-amino acid produced per minute per mg of enzyme formed from the corresponding S-amino acid. In an alternative embodiment, for amino acid racemases, one unit of activity equals one umol of S-amino acid produced per minute per mg of enzyme formed from the
5 corresponding R-amino acid.

In one aspect, the polypeptides of the invention comprise at least one glycosylation site or further comprises a polysaccharide. The glycosylation can be an N-linked glycosylation, e.g., wherein the polypeptide is glycosylated after being expressed in a *P. pastoris* or a *S. pombe*.

10 The invention provides protein preparation comprising a polypeptide of the invention, wherein the protein preparation comprises a liquid, a slurry, a solid or a gel. The invention provides heterodimers comprising a polypeptide of the invention and a second domain. The second domain can be a polypeptide and the heterodimer is a fusion protein. the second domain can be an epitope or a tag.

15 The invention provides homodimers or heterodimers comprising a polypeptide of the invention. The invention provides immobilized polypeptides, wherein the polypeptide comprises a sequence of the invention, or a subsequence thereof, or a polypeptide encoded by a nucleic acid of the invention, or a polypeptide comprising a polypeptide of the invention and a second domain, e.g., wherein the polypeptide is
20 immobilized on or inside a cell, a vesicle, a liposome, a film, a membrane, a metal, a resin, a polymer, a ceramic, a glass, a microelectrode, a graphitic particle, a bead, a gel, a plate, an array, a capillary tube, a crystal, a tablet, a pill, a capsule, a powder, an agglomerate, a surface, a porous structure, or materials such as wood chips, brownstock, pulp, paper, and materials deriving therefrom.

25 The isomerases, e.g., racemases, e.g., amino acid racemases, alanine racemases, and/or epimerases of the invention can be used or formulated alone or as mixture (a “cocktail”) of isomerases, e.g., racemases, e.g., amino acid racemases, alanine racemases, and/or epimerases, and other hydrolytic enzymes such as cellulases, mannanases, proteases, lipases, amylases, or redox enzymes such as laccases, peroxidases, catalases,
30 oxidases, or reductases. They can be used formulated in a solid form such as a powder, a lyophilized preparation, a granule, a tablet, a bar, a crystal, a capsule, a pill, a pellet, or in a liquid form such as in an aqueous solution, an aerosol, a gel, a paste, a slurry, an aqueous/oil emulsion, a cream, a capsule, or in a vesicular or micellar suspension. The formulations of the invention can comprise any or a combination of the following

ingredients: polyols such as a polyethylene glycol, a polyvinylalcohol, a glycerol, a sugar such as a sucrose, a sorbitol, a trehalose, a glucose, a fructose, a maltose, a mannose, a gelling agent such as a guar gum, a carageenan, an alginate, a dextrans, a cellulosic derivative, a pectin, a salt such as a sodium chloride, a sodium sulfate, an ammonium sulfate, a calcium chloride, a magnesium chloride, a zinc chloride, a zinc sulfate, a salt of a fatty acid and a fatty acid derivative, a metal chelator such as an EDTA, an EGTA, a sodium citrate, an antimicrobial agent such as a fatty acid or a fatty acid derivative, a paraben, a sorbate, a benzoate, an additional modulating compound to block the impact of an enzyme such as a protease, a bulk proteins such as a BSA, a wheat hydrolysate, a borate compound, an amino acid or a peptide, an appropriate pH or temperature modulating compound, an emulsifier such as a non-ionic and/or an ionic detergent, a redox agent such as a cystine/cysteine, a glutathione, an oxidized glutathione, a reduced or an antioxidant compound such as an ascorbic acid, or a dispersant. Cross-linking and protein modification such as pegylation, fatty acid modification, glycosylation can also be used to improve enzyme stability.

The invention provides arrays comprising immobilized polypeptide(s) and/or nucleic acids of the invention, and arrays comprising an immobilized oligonucleotide of the invention. The enzymes, fragments thereof and nucleic acids which encode the enzymes, or probes of the invention, and fragments thereof, can be affixed to a solid support; and these embodiments can be economical and efficient in the use of enzymes and nucleic acids of the invention in industrial, medical, research, pharmaceutical, food and feed and food and feed supplement processing and other applications and processes. For example, a consortium or cocktail of enzymes (or active fragments thereof), which are used in a specific chemical reaction, can be attached to a solid support and dunked into a process vat. The enzymatic reaction can occur. Then, the solid support can be taken out of the vat, along with the enzymes affixed thereto, for repeated use. In one embodiment of the invention, the isolated nucleic acid is affixed to a solid support. In another embodiment of the invention, the solid support is selected from the group of a gel, a resin, a polymer, a ceramic, a glass, a microelectrode and any combination thereof.

For example, solid supports useful in this invention include gels. Some examples of gels include sepharose, gelatin, glutaraldehyde, chitosan-treated glutaraldehyde, albumin-glutaraldehyde, chitosan-Xanthan, toyopearl gel (polymer gel), alginate, alginate-polylysine, carrageenan, agarose, glyoxyl agarose, magnetic agarose, dextran-agarose, poly(Carbamoyl Sulfonate) hydrogel, BSA-PEG hydrogel, phosphorylated polyvinyl

alcohol (PVA), monoaminoethyl-N-aminoethyl (MANA), amino, or any combination thereof. Another solid support useful in the present invention are resins or polymers. Some examples of resins or polymers include cellulose, acrylamide, nylon, rayon, polyester, anion-exchange resin, AMBERLITE™ XAD-7, AMBERLITE™ XAD-8, 5 AMBERLITE™ IRA-94, AMBERLITE™ IRC-50, polyvinyl, polyacrylic, polymethacrylate, or any combination thereof. Another type of solid support useful in the present invention is ceramic. Some examples include non-porous ceramic, porous ceramic, SiO₂, Al₂O₃. Another type of solid support useful in the present invention is glass. Some examples include non-porous glass, porous glass, aminopropyl glass or any 10 combination thereof. Another type of solid support which can be used is a microelectrode. An example is a polyethyleneimine-coated magnetite. Graphitic particles can be used as a solid support. Another example of a solid support is a cell, such as a red blood cell.

There are many methods which would be known to one of skill in the art for immobilizing enzymes or fragments thereof, or nucleic acids, onto a solid support. Some 15 examples of such methods include electrostatic droplet generation, electrochemical means, via adsorption, via covalent binding, via cross-linking, via chemical reaction or process, via encapsulation, via entrapment, via calcium alginate, or via poly (2-hydroxyethyl methacrylate). Like methods are described in *Methods in Enzymology, Immobilized Enzymes and Cells*, Part C. 1987. Academic Press. Edited by S. P. Colowick and N. O. 20 Kaplan. Volume 136; and *Immobilization of Enzymes and Cells*. 1997. Humana Press. Edited by G. F. Bickerstaff. Series: Methods in Biotechnology, Edited by J. M. Walker.

The invention provides isolated, synthetic or recombinant antibodies that specifically binds to a polypeptide of the invention. The antibody can be a monoclonal or a polyclonal antibody, or is a single chained antibody. The invention provides hybridomas 25 comprising an antibody that specifically binds to a polypeptide of the invention.

The invention provides methods of isolating or identifying a polypeptide with an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity comprising the steps of: (a) providing an antibody of the invention; (b) providing a sample comprising polypeptides; 30 and (c) contacting the sample of step (b) with the antibody of step (a) under conditions wherein the antibody can specifically bind to the polypeptide, thereby isolating or identifying a polypeptide having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity. The invention provides methods of making an anti-isomerase, e.g., anti-racemase, e.g.,

anti-amino acid racemase, anti-alanine racemase, and/or anti-epimerase antibody comprising administering to a non-human animal a nucleic acid of the invention or a subsequence thereof in an amount sufficient to generate a humoral immune response, thereby making an anti-isomerase, e.g., anti-racemase, e.g., anti-amino acid racemase, anti-alanine racemase, and/or anti-epimerase antibody. The invention provides methods of making an anti-isomerase, e.g., anti-racemase, e.g., anti-amino acid racemase, anti-alanine racemase, and/or anti-epimerase antibody comprising administering to a non-human animal a polypeptide of the invention or a subsequence thereof in an amount sufficient to generate a humoral immune response, thereby making an anti-isomerase, e.g., anti-racemase, e.g., anti-amino acid racemase, anti-alanine racemase, and/or anti-epimerase antibody.

The invention provides methods of producing a recombinant polypeptide comprising the steps of: (a) providing a nucleic acid operably linked to a promoter, wherein the nucleic acid comprises a sequence of the invention; and (b) expressing the nucleic acid of step (a) under conditions that allow expression of the polypeptide, thereby producing a recombinant polypeptide. The method can further comprise transforming a host cell with the nucleic acid of step (a) followed by expressing the nucleic acid of step (a), thereby producing a recombinant polypeptide in a transformed cell.

The invention provides methods for identifying a polypeptide having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity comprising: (a) providing a polypeptide of the invention; (b) providing an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase substrate; and (c) contacting the polypeptide with the substrate of step (b) and detecting a decrease in the amount of substrate or an increase in the amount of a reaction product, wherein a decrease in the amount of the substrate or an increase in the amount of the reaction product detects a polypeptide having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity.

The invention provides methods for identifying an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase substrate comprising: (a) providing a polypeptide of the invention; (b) providing a test substrate; and (c) contacting the polypeptide of step (a) with the test substrate of step (b) and detecting a decrease in the amount of substrate or an increase in the amount of reaction product, wherein a decrease in the amount of the substrate or an increase in the amount of

a reaction product identifies the test substrate as an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase substrate.

The invention provides methods of determining whether a test compound specifically binds to a polypeptide comprising: (a) expressing a nucleic acid or a vector
5 comprising the nucleic acid under conditions permissive for translation of the nucleic acid to a polypeptide, wherein the nucleic acid has a sequence of the invention; (b) providing a test compound; (c) contacting the polypeptide with the test compound; and (d) determining whether the test compound of step (b) specifically binds to the polypeptide.

The invention provides methods of determining whether a test compound
10 specifically binds to a polypeptide comprising: (a) providing a polypeptide of the invention; (b) providing a test compound; (c) contacting the polypeptide with the test compound; and (d) determining whether the test compound of step (b) specifically binds to the polypeptide.

The invention provides methods for identifying a modulator of an isomerase
15 activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity comprising: (a) providing a polypeptide of the invention; (b) providing a test compound; (c) contacting the polypeptide of step (a) with the test compound of step (b) and measuring an activity of the isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase, wherein a
20 change in the isomerase activity, e.g., racemase activity, e.g., amino acid racemase activity, alanine racemase activity, and/or epimerase activity measured in the presence of the test compound compared to the activity in the absence of the test compound provides a determination that the test compound modulates the isomerase activity, e.g., racemase activity, e.g., amino acid racemase activity, alanine racemase activity, and/or epimerase
25 activity. The isomerase activity, e.g., racemase activity, e.g., amino acid racemase activity, alanine racemase activity, and/or epimerase activity can be measured by providing an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase substrate and detecting a decrease in the amount of the substrate or an increase in the amount of a reaction product, or, an increase in the amount
30 of the substrate or a decrease in the amount of a reaction product. In one aspect, a decrease in the amount of the substrate or an increase in the amount of the reaction product with the test compound as compared to the amount of substrate or reaction product without the test compound identifies the test compound as an activator of an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an

alanine racemase activity, and/or an epimerase activity. In one aspect, an increase in the amount of the substrate or a decrease in the amount of the reaction product with the test compound as compared to the amount of substrate or reaction product without the test compound identifies the test compound as an inhibitor of an isomerase activity, e.g., a
5 racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity.

The invention provides computer systems comprising a processor and a data storage device wherein said data storage device has stored thereon a polypeptide sequence or a nucleic acid sequence, wherein the polypeptide sequence comprises sequence of the
10 invention, a polypeptide encoded by a nucleic acid of the invention. The computer systems can further comprise a sequence comparison algorithm and a data storage device having at least one reference sequence stored thereon. In another aspect, the sequence comparison algorithm comprises a computer program that indicates polymorphisms. In one aspect, the computer system can further comprise an identifier that identifies one or
15 more features in said sequence. The invention provides computer readable media having stored thereon a polypeptide sequence or a nucleic acid sequence of the invention. The invention provides methods for identifying a feature in a sequence comprising the steps of: (a) reading the sequence using a computer program which identifies one or more features in a sequence, wherein the sequence comprises a polypeptide sequence or a nucleic acid
20 sequence of the invention; and (b) identifying one or more features in the sequence with the computer program. The invention provides methods for comparing a first sequence to a second sequence comprising the steps of: (a) reading the first sequence and the second sequence through use of a computer program which compares sequences, wherein the first sequence comprises a polypeptide sequence or a nucleic acid sequence of the invention;
25 and (b) determining differences between the first sequence and the second sequence with the computer program. The step of determining differences between the first sequence and the second sequence can further comprise the step of identifying polymorphisms. In one aspect, the method can further comprise an identifier that identifies one or more features in a sequence. In another aspect, the method can comprise reading the first
30 sequence using a computer program and identifying one or more features in the sequence.

The invention provides methods for isolating or recovering a nucleic acid encoding a polypeptide having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity from an environmental sample comprising the steps of: (a) providing an amplification primer

sequence pair for amplifying a nucleic acid encoding a polypeptide having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity, wherein the primer pair is capable of amplifying a nucleic acid of the invention; (b) isolating a nucleic acid from the environmental sample or treating the environmental sample such that nucleic acid in the sample is accessible for hybridization to the amplification primer pair; and, (c) combining the nucleic acid of step (b) with the amplification primer pair of step (a) and amplifying nucleic acid from the environmental sample, thereby isolating or recovering a nucleic acid encoding a polypeptide having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity from an environmental sample. One or each member of the amplification primer sequence pair can comprise an oligonucleotide comprising at least about 10 to 50 consecutive bases of a sequence of the invention. In one aspect, the amplification primer sequence pair is an amplification pair of the invention.

The invention provides methods for isolating or recovering a nucleic acid encoding a polypeptide having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity from an environmental sample comprising the steps of: (a) providing a polynucleotide probe comprising a nucleic acid of the invention or a subsequence thereof; (b) isolating a nucleic acid from the environmental sample or treating the environmental sample such that nucleic acid in the sample is accessible for hybridization to a polynucleotide probe of step (a); (c) combining the isolated nucleic acid or the treated environmental sample of step (b) with the polynucleotide probe of step (a); and (d) isolating a nucleic acid that specifically hybridizes with the polynucleotide probe of step (a), thereby isolating or recovering a nucleic acid encoding a polypeptide having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity from an environmental sample. The environmental sample can comprise a water sample, a liquid sample, a soil sample, an air sample or a biological sample. In one aspect, the biological sample can be derived from a bacterial cell, a protozoan cell, an insect cell, a yeast cell, a plant cell, a fungal cell or a mammalian cell.

The invention provides methods of generating a variant of a nucleic acid encoding a polypeptide having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity comprising the steps of: (a) providing a template nucleic acid comprising a nucleic acid of the

invention; and (b) modifying, deleting or adding one or more nucleotides in the template sequence, or a combination thereof, to generate a variant of the template nucleic acid. In one aspect, the method can further comprise expressing the variant nucleic acid to generate a variant isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase polypeptide. The modifications, additions or deletions can be introduced by a method comprising error-prone PCR, shuffling, oligonucleotide-directed mutagenesis, assembly PCR, sexual PCR mutagenesis, *in vivo* mutagenesis, cassette mutagenesis, recursive ensemble mutagenesis, exponential ensemble mutagenesis, site-specific mutagenesis, gene reassembly (e.g., GeneReassembly, see, e.g., U.S. Patent No. 6,537,776), Gene Site Saturation Mutagenesis (GSSM), synthetic ligation reassembly (SLR) or a combination thereof. In another aspect, the modifications, additions or deletions are introduced by a method comprising recombination, recursive sequence recombination, phosphothioate-modified DNA mutagenesis, uracil-containing template mutagenesis, gapped duplex mutagenesis, point mismatch repair mutagenesis, repair-deficient host strain mutagenesis, chemical mutagenesis, radiogenic mutagenesis, deletion mutagenesis, restriction-selection mutagenesis, restriction-purification mutagenesis, artificial gene synthesis, ensemble mutagenesis, chimeric nucleic acid multimer creation and a combination thereof.

In one aspect, the method can be iteratively repeated until an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase having an altered or different activity or an altered or different stability from that of a polypeptide encoded by the template nucleic acid is produced. In one aspect, the variant isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase polypeptide is thermotolerant, and retains some activity after being exposed to an elevated temperature. In another aspect, the variant isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase polypeptide has increased glycosylation as compared to the isomerase, e.g., the racemase, e.g., the amino acid racemase, the alanine racemase, and/or the epimerase encoded by a template nucleic acid. Alternatively, the variant isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase polypeptide has an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity under a high temperature, wherein the isomerase, e.g., the racemase, e.g., the amino acid racemase, the alanine racemase, and/or the epimerase encoded by the template nucleic acid is not active under the high temperature. In one aspect, the method can be iteratively

repeated until an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase coding sequence having an altered codon usage from that of the template nucleic acid is produced. In another aspect, the method can be iteratively repeated until an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine
5 racemase, and/or an epimerase gene having higher or lower level of message expression or stability from that of the template nucleic acid is produced. In another aspect, formulation of the final isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase product enables an increase or modulation of the performance of the isomerase, e.g., the racemase, e.g., the amino acid racemase, the alanine racemase, and/or the
10 epimerase in the product.

The invention provides methods for modifying codons in a nucleic acid encoding a polypeptide having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity to increase its expression in a host cell, the method comprising: (a) providing a nucleic acid of the
15 invention encoding a polypeptide having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity; and, (b) identifying a non-preferred or a less preferred codon in the nucleic acid of step (a) and replacing it with a preferred or neutrally used codon encoding the same amino acid as the replaced codon, wherein a preferred codon is a codon over-represented
20 in coding sequences in genes in the host cell and a non-preferred or less preferred codon is a codon under-represented in coding sequences in genes in the host cell, thereby modifying the nucleic acid to increase its expression in a host cell.

The invention provides methods for modifying codons in a nucleic acid encoding a polypeptide having an isomerase activity, e.g., a racemase activity, e.g., an amino acid
25 racemase activity, an alanine racemase activity, and/or an epimerase activity; the method comprising: (a) providing a nucleic acid of the invention; and, (b) identifying a codon in the nucleic acid of step (a) and replacing it with a different codon encoding the same amino acid as the replaced codon, thereby modifying codons in a nucleic acid encoding an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an
30 epimerase.

The invention provides methods for modifying codons in a nucleic acid encoding a polypeptide having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity to increase its expression in a host cell, the method comprising: (a) providing a nucleic acid of the

invention encoding an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase polypeptide; and, (b) identifying a non-preferred or a less preferred codon in the nucleic acid of step (a) and replacing it with a preferred or neutrally used codon encoding the same amino acid as the replaced codon, wherein a preferred codon is a codon over-represented in coding sequences in genes in the host cell and a non-preferred or less preferred codon is a codon under-represented in coding sequences in genes in the host cell, thereby modifying the nucleic acid to increase its expression in a host cell.

The invention provides methods for modifying a codon in a nucleic acid encoding a polypeptide having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity to decrease its expression in a host cell, the method comprising: (a) providing a nucleic acid of the invention; and (b) identifying at least one preferred codon in the nucleic acid of step (a) and replacing it with a non-preferred or less preferred codon encoding the same amino acid as the replaced codon, wherein a preferred codon is a codon over-represented in coding sequences in genes in a host cell and a non-preferred or less preferred codon is a codon under-represented in coding sequences in genes in the host cell, thereby modifying the nucleic acid to decrease its expression in a host cell. In one aspect, the host cell can be a bacterial cell, a fungal cell, an insect cell, a yeast cell, a plant cell or a mammalian cell.

The invention provides methods for producing a library of nucleic acids encoding a plurality of modified isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase active sites or substrate binding sites, wherein the modified active sites or substrate binding sites are derived from a first nucleic acid comprising a sequence encoding a first active site or a first substrate binding site the method comprising: (a) providing a first nucleic acid encoding a first active site or first substrate binding site, wherein the first nucleic acid sequence comprises a sequence that hybridizes under stringent conditions to a sequence of the invention, or a subsequence thereof, and the nucleic acid encodes an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase active site or an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase substrate binding site; (b) providing a set of mutagenic oligonucleotides that encode naturally-occurring amino acid variants at a plurality of targeted codons in the first nucleic acid; and, (c) using the set of mutagenic oligonucleotides to generate a set of active site-encoding or substrate binding site-encoding variant nucleic acids encoding a range of amino acid variations at

each amino acid codon that was mutagenized, thereby producing a library of nucleic acids encoding a plurality of modified isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase active sites or substrate binding sites. In one aspect, the method comprises mutagenizing the first nucleic acid of step (a) by a method
5 comprising an optimized directed evolution system, Gene Site Saturation Mutagenesis (GSSM), or a synthetic ligation reassembly (SLR). In one aspect, the method comprises mutagenizing the first nucleic acid of step (a) or variants by a method comprising error-prone PCR, shuffling, oligonucleotide-directed mutagenesis, assembly PCR, sexual PCR mutagenesis, in vivo mutagenesis, cassette mutagenesis, recursive ensemble mutagenesis,
10 exponential ensemble mutagenesis, site-specific mutagenesis, gene reassembly (GeneReassembly, U.S. Patent No. 6,537,776), Gene Site Saturation Mutagenesis (GSSM), synthetic ligation reassembly (SLR) and a combination thereof. In one aspect, the method comprises mutagenizing the first nucleic acid of step (a) or variants by a method comprising recombination, recursive sequence recombination, phosphothioate-
15 modified DNA mutagenesis, uracil-containing template mutagenesis, gapped duplex mutagenesis, point mismatch repair mutagenesis, repair-deficient host strain mutagenesis, chemical mutagenesis, radiogenic mutagenesis, deletion mutagenesis, restriction-selection mutagenesis, restriction-purification mutagenesis, artificial gene synthesis, ensemble mutagenesis, chimeric nucleic acid multimer creation and a combination thereof.

20 The invention provides methods for making a small molecule comprising: (a) providing a plurality of biosynthetic enzymes capable of synthesizing or modifying a small molecule, wherein one of the enzymes comprises an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase enzyme encoded by a nucleic acid of the invention; (b) providing a substrate for at least one of the enzymes
25 of step (a); and (c) reacting the substrate of step (b) with the enzymes under conditions that facilitate a plurality of biocatalytic reactions to generate a small molecule by a series of biocatalytic reactions. The invention provides methods for modifying a small molecule comprising: (a) providing an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase enzyme, wherein the enzyme comprises a
30 polypeptide of the invention, or, a polypeptide encoded by a nucleic acid of the invention, or a subsequence thereof; (b) providing a small molecule; and (c) reacting the enzyme of step (a) with the small molecule of step (b) under conditions that facilitate an enzymatic reaction catalyzed by the isomerase, e.g., the racemase, e.g., the amino acid racemase, the alanine racemase, and/or the epimerase enzyme, thereby modifying a small molecule by

an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase enzymatic reaction. In one aspect, the method can comprise a plurality of small molecule substrates for the enzyme of step (a), thereby generating a library of modified small molecules produced by at least one enzymatic reaction catalyzed by the isomerase, e.g., the racemase, e.g., the amino acid racemase, the alanine racemase, and/or the epimerase enzyme. In one aspect, the method can comprise a plurality of additional enzymes under conditions that facilitate a plurality of biocatalytic reactions by the enzymes to form a library of modified small molecules produced by the plurality of enzymatic reactions. In another aspect, the method can further comprise the step of testing the library to determine if a particular modified small molecule that exhibits a desired activity is present within the library. The step of testing the library can further comprise the steps of systematically eliminating all but one of the biocatalytic reactions used to produce a portion of the plurality of the modified small molecules within the library by testing the portion of the modified small molecule for the presence or absence of the particular modified small molecule with a desired activity, and identifying at least one specific biocatalytic reaction that produces the particular modified small molecule of desired activity.

The invention provides methods for determining a functional fragment of an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase enzyme comprising the steps of: (a) providing an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase enzyme, wherein the enzyme comprises a polypeptide of the invention, or a polypeptide encoded by a nucleic acid of the invention, or a subsequence thereof; and (b) deleting a plurality of amino acid residues from the sequence of step (a) and testing the remaining subsequence for an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity, thereby determining a functional fragment of an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase enzyme. In one aspect, the isomerase activity, e.g., racemase activity, e.g., amino acid racemase activity, alanine racemase activity, and/or epimerase activity is measured by providing an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase substrate and detecting a decrease in the amount of the substrate or an increase in the amount of a reaction product.

The invention provides methods for whole cell engineering of new or modified phenotypes by using real-time metabolic flux analysis, the method comprising: (a) making a modified cell by modifying the genetic composition of a cell, wherein the genetic composition is modified by addition to the cell of a nucleic acid of the invention; (b) 5 culturing the modified cell to generate a plurality of modified cells; (c) measuring at least one metabolic parameter of the cell by monitoring the cell culture of step (b) in real time; and, (d) analyzing the data of step (c) to determine if the measured parameter differs from a comparable measurement in an unmodified cell under similar conditions, thereby identifying an engineered phenotype in the cell using real-time metabolic flux analysis. In 10 one aspect, the genetic composition of the cell can be modified by a method comprising deletion of a sequence or modification of a sequence in the cell, or, knocking out the expression of a gene. In one aspect, the method can further comprise selecting a cell comprising a newly engineered phenotype. In another aspect, the method can comprise culturing the selected cell, thereby generating a new cell strain comprising a newly 15 engineered phenotype.

The invention provides isolated, synthetic or recombinant signal sequences consisting of, or comprising, a sequence as set forth in residues 1 to 12, 1 to 13, 1 to 14, 1 to 15, 1 to 16, 1 to 17, 1 to 18, 1 to 19, 1 to 20, 1 to 21, 1 to 22, 1 to 23, 1 to 24, 1 to 25, 1 to 26, 1 to 27, 1 to 28, 1 to 28, 1 to 30, 1 to 31, 1 to 32, 1 to 33, 1 to 34, 1 to 35, 1 to 36, 1 to 20 to 37, 1 to 38, 1 to 40, 1 to 41, 1 to 42, 1 to 43 or 1 to 44, of a polypeptide of the invention, including exemplary polypeptide sequences of the invention.

The invention provides chimeric polypeptides comprising at least a first domain comprising a signal peptide (SP) and at least a second domain comprising a heterologous polypeptide or peptide comprising a sequence of the invention, or a subsequence thereof, 25 wherein the heterologous polypeptide or peptide is not naturally associated with the signal peptide (SP). In one aspect, the signal peptide (SP) is not derived from an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase. The heterologous polypeptide or peptide can be amino terminal to, carboxy terminal to or on both ends of the signal peptide (SP) or an isomerase, e.g., a racemase, e.g., an amino acid 30 racemase, an alanine racemase, and/or an epimerase catalytic domain (CD). The invention provides isolated, synthetic or recombinant nucleic acids encoding a chimeric polypeptide, wherein the chimeric polypeptide comprises at least a first domain comprising signal peptide (SP) and at least a second domain comprising a heterologous polypeptide or peptide comprising a sequence of the invention, or a subsequence thereof, wherein the

heterologous polypeptide or peptide is not naturally associated with the signal peptide (SP).

The invention provides methods of increasing thermotolerance or thermostability of an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase polypeptide, the method comprising glycosylating an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase polypeptide, wherein the polypeptide comprises at least thirty contiguous amino acids of a polypeptide of the invention; or a polypeptide encoded by a nucleic acid sequence of the invention, thereby increasing the thermotolerance or thermostability of the isomerase, e.g., the racemase, e.g., the amino acid racemase, the alanine racemase, and/or the epimerase polypeptide. In one aspect, the isomerase, e.g., the racemase, e.g., the amino acid racemase, the alanine racemase, and/or the epimerase-specific activity can be thermostable or thermotolerant at a temperature in the range from greater than about 0°C to about 20°C, about 20°C to about 37°C, about 37°C to about 50°C, about 50°C to about 70°C, about 70°C to about 75°C, about 75°C to about 80°C, about 80°C to about 85°C, about 85°C to about 90°C, about 90°C to about 95°C, about 95°C to about 100°C, about 100°C to about 110°C, or higher.

The invention provides methods for overexpressing a recombinant isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase polypeptide in a cell comprising expressing a vector comprising a nucleic acid comprising a nucleic acid of the invention or a nucleic acid sequence of the invention, wherein the sequence identities are determined by analysis with a sequence comparison algorithm or by visual inspection, wherein overexpression is effected by use of a high activity promoter, a dicistronic vector or by gene amplification of the vector.

The invention provides methods of making a transgenic plant and seeds comprising: (a) introducing a heterologous nucleic acid sequence into the cell, wherein the heterologous nucleic sequence comprises a nucleic acid sequence of the invention, thereby producing a transformed plant or seed cell; and (b) producing a transgenic plant from the transformed cell or seed. In one aspect, the step (a) can further comprise introducing the heterologous nucleic acid sequence by electroporation or microinjection of plant cell protoplasts. In another aspect, the step (a) can further comprise introducing the heterologous nucleic acid sequence directly to plant tissue by DNA particle bombardment. Alternatively, the step (a) can further comprise introducing the heterologous nucleic acid

sequence into the plant cell DNA using an *Agrobacterium tumefaciens* host. In one aspect, the plant cell can be a potato, corn, rice, wheat, tobacco, or barley cell.

The invention provides methods of expressing a heterologous nucleic acid sequence in a plant cell comprising: (a) transforming the plant cell with a heterologous nucleic acid sequence operably linked to a promoter, wherein the heterologous nucleic sequence comprises a nucleic acid of the invention; (b) growing the plant under conditions wherein the heterologous nucleic acids sequence is expressed in the plant cell. The invention provides methods of expressing a heterologous nucleic acid sequence in a plant cell comprising: (a) transforming the plant cell with a heterologous nucleic acid sequence operably linked to a promoter, wherein the heterologous nucleic sequence comprises a sequence of the invention; (b) growing the plant under conditions wherein the heterologous nucleic acids sequence is expressed in the plant cell.

The invention provides detergent compositions comprising a polypeptide of the invention, or a polypeptide encoded by a nucleic acid of the invention, wherein the polypeptide has an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity. The isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase can be nonsurface-active or surface-active. The isomerase, e.g., the racemase, e.g., the amino acid racemase, the alanine racemase, and/or the epimerase can be formulated in a non-aqueous liquid composition, a cast solid, a granular form, a particulate form, a compressed tablet, a gel form, a paste or a slurry form. The invention provides methods for washing an object comprising: (a) providing a composition comprising a polypeptide of the invention having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity, or a polypeptide encoded by a nucleic acid of the invention; (b) providing an object; and (c) contacting the polypeptide of step (a) and the object of step (b) under conditions wherein the composition can wash the object.

The invention provides textiles or fabrics, including, e.g., threads, comprising a polypeptide of the invention, or a polypeptide encoded by a nucleic acid of the invention. The invention provides methods for treating a textile or fabric (e.g., removing a stain from a composition) comprising: (a) providing a composition comprising a polypeptide of the invention having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity, or a polypeptide encoded by a nucleic acid of the invention; (b) providing a textile or fabric;

and (c) contacting the polypeptide of step (a) and the composition of step (b) under conditions wherein the isomerase, e.g., the racemase, e.g., the amino acid racemase, the alanine racemase, and/or the epimerase can treat the textile or fabric (e.g., remove the stain). The invention provides methods for improving the finish of a fabric comprising:

5 (a) providing a composition comprising a polypeptide of the invention having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity, or a polypeptide encoded by a nucleic acid of the invention; (b) providing a fabric; and (c) contacting the polypeptide of step (a) and the fabric of step (b) under conditions wherein the polypeptide can treat the

10 fabric thereby improving the finish of the fabric. In one aspect, the fabric is a wool or a silk. In another aspect, the fabric is a cellulosic fiber or a blend of a natural fiber and a synthetic fiber.

The invention provides feeds, foods, feed supplements, food supplements, dietary compositions or dietary aids comprising a polypeptide of the invention, or a polypeptide

15 encoded by a nucleic acid of the invention. The food or the feed can be, e.g., a cereal, a grain, a corn and the like.

The invention provides dough, bread or baked products and/or dough, bread or baked product precursors comprising a polypeptide having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity,

20 and/or an epimerase activity, wherein the polypeptide comprises a sequence of the invention, or the polypeptide is encoded by a nucleic acid comprising a sequence of the invention, or an enzymatically active fragment thereof.

The invention provides beverages and beverage precursors comprising a polypeptide, or an enzymatically active fragment thereof, having an isomerase activity,

25 e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity, wherein the polypeptide comprises a sequence of the invention, or the polypeptide is encoded by a nucleic acid comprising a sequence of the invention. The invention provides methods of beverage production comprising administration of at least one polypeptide having an isomerase activity, e.g., a racemase

30 activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity, wherein the polypeptide comprises a sequence of the invention, or the polypeptide is encoded by a nucleic acid comprising a sequence of the invention, or an enzymatically active fragment thereof, to a beverage or a beverage precursor, wherein in one aspect (optionally) the beverage or beverage precursor is a wort or a beer.

The invention provides food, feed or nutritional supplements, e.g. for a human or an animal comprising a polypeptide of the invention, e.g., a polypeptide encoded by the nucleic acid of the invention. In one aspect, the polypeptide in the food or nutritional supplement can be glycosylated. The invention provides edible enzyme delivery matrices comprising a polypeptide of the invention, e.g., a polypeptide encoded by the nucleic acid of the invention. In one aspect, the delivery matrix comprises a pellet. In one aspect, the polypeptide can be glycosylated. In one aspect, the isomerase activity, e.g., racemase activity, e.g., amino acid racemase activity, alanine racemase activity, and/or epimerase activity is thermotolerant. In another aspect, the isomerase activity, e.g., racemase activity, e.g., amino acid racemase activity, alanine racemase activity, and/or epimerase activity is thermostable.

In one aspect, the isomerase, e.g., the racemase, e.g., the amino acid racemase, the alanine racemase, and/or the epimerase enzyme can be prepared by expression of a polynucleotide encoding the isomerase, e.g., the racemase, e.g., the amino acid racemase, the alanine racemase, and/or the epimerase in an organism selected from the group consisting of a bacterium, a yeast, a plant, an insect, a fungus and an animal. The organism can be selected from the group consisting of an *S. pombe*, *S. cerevisiae*, *Pichia pastoris*, *Pseudomonas* sp., *E. coli*, *Streptomyces* sp., *Bacillus* sp. and *Lactobacillus* sp.

The invention provides edible enzyme delivery matrix comprising a thermostable recombinant isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase enzyme, e.g., a polypeptide of the invention. The invention provides methods for delivering an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase supplement to an animal, the method comprising: preparing an edible enzyme delivery matrix in the form of pellets comprising a granulate edible carrier and a thermostable recombinant isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase enzyme, wherein the pellets readily disperse the isomerase, e.g., the racemase, e.g., the amino acid racemase, the alanine racemase, and/or the epimerase enzyme contained therein into aqueous media, and administering the edible enzyme delivery matrix to the animal. The recombinant isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase enzyme can comprise a polypeptide of the invention. The granulate edible carrier can comprise a carrier selected from the group consisting of a grain germ, a grain germ that is spent of oil, a hay, an alfalfa, a timothy, a soy hull, a sunflower seed meal and a wheat midd. The edible carrier can comprise grain germ that is spent of oil. The isomerase, e.g., the racemase, e.g., the

amino acid racemase, the alanine racemase, and/or the epimerase enzyme can be glycosylated to provide thermostability at pelletizing conditions. The delivery matrix can be formed by pelletizing a mixture comprising a grain germ and an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase. The pelletizing conditions can include application of steam. The pelletizing conditions can comprise application of a temperature in excess of about 80°C for about 5 minutes and the enzyme retains a specific activity of at least 350 to about 900 units per milligram of enzyme.

The invention provides methods for treating, e.g. improving texture and flavor of a dairy product comprising: (a) providing a polypeptide of the invention having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity, or an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase encoded by a nucleic acid of the invention; (b) providing a dairy product; and (c) contacting the polypeptide of step (a) and the dairy product of step (b) under conditions wherein the isomerase, e.g., the racemase, e.g., the amino acid racemase, the alanine racemase, and/or the epimerase can treat, e.g. improve the texture or flavor of the dairy product. In one aspect, the dairy product comprises a cheese or a yogurt. The invention provides dairy products comprising an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase of the invention, or is encoded by a nucleic acid of the invention.

The invention provides methods for improving the extraction of oil from an oil-rich plant material comprising: (a) providing a polypeptide of the invention having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity, or an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase encoded by a nucleic acid of the invention; (b) providing an oil-rich plant material; and (c) contacting the polypeptide of step (a) and the oil-rich plant material. In one aspect, the oil-rich plant material comprises an oil-rich seed. The oil can be a soybean oil, an olive oil, a rapeseed (canola) oil or a sunflower oil.

The invention provides methods for preparing a fruit or vegetable juice, syrup, puree or extract comprising: (a) providing a polypeptide of the invention having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity, or an isomerase, e.g., a racemase,

e.g., an amino acid racemase, an alanine racemase, and/or an epimerase encoded by a nucleic acid of the invention; (b) providing a composition or a liquid comprising a fruit or vegetable material; and (c) contacting the polypeptide of step (a) and the composition, thereby preparing the fruit or vegetable juice, syrup, puree or extract.

5 The invention provides methods for treating a wood, a wood product, a paper, a paper product, a pulp, a pulp product, a paper waste or a paper recycling composition comprising: (a) providing a polypeptide of the invention having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity, or an isomerase, e.g., a racemase, e.g., an amino acid
10 racemase, an alanine racemase, and/or an epimerase encoded by a nucleic acid of the invention; (b) providing a composition comprising a wood, a wood product, a paper, a paper product, a pulp, a pulp product, a paper waste or a paper recycling composition; and (c) contacting the polypeptide of step (a) and the composition, thereby treating the wood, wood product, paper, paper product, pulp, pulp product, paper waste or paper recycling
15 composition. In one aspect of the invention, the treatment comprises reducing or solubilizing lignin (delignification), bleaching or decoloring, and/or deinking.

 The invention provides papers or paper products or paper pulp comprising an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase of the invention, or a polypeptide encoded by a nucleic acid of the invention.

20 The invention provides methods for treating a paper or a paper or wood pulp comprising: (a) providing a polypeptide of the invention having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity, or an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase encoded by a nucleic acid of the invention; (b)
25 providing a composition comprising a paper or a paper or wood pulp; and (c) contacting the polypeptide of step (a) and the composition of step (b) under conditions wherein the isomerase, e.g., the racemase, e.g., the amino acid racemase, the alanine racemase, and/or the epimerase can treat the paper or paper or wood pulp.

 The invention provides methods for bleaching a thread, fabric, yarn, cloth or textile
30 comprising contacting the fabric, yarn, cloth or textile with an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase under conditions suitable to produce a whitening of the textile, wherein the isomerase, e.g., the racemase, e.g., the amino acid racemase, the alanine racemase, and/or the epimerase comprises a polypeptide of the invention, or an enzymatically active fragment thereof. The thread,

fabric, yarn, cloth or textile can comprise a non-cotton cellulosic thread, fabric, yarn, cloth or textile. The invention provides fabrics, yarns, cloths or textiles comprising a polypeptide having a sequence of the invention, or a polypeptide encoded by a nucleic acid comprising a sequence of the invention, or an enzymatically active fragment thereof, 5 wherein in one aspect (optionally) the fabric, yarn, cloth or textile comprises a non-cotton cellulosic fabric, yarn, cloth or textile.

The invention provides wood, wood chips, wood pulp, wood products, paper pulps, paper products, newspapers or paper waste comprising a polypeptide of the invention, or an enzymatically active fragment thereof. The invention provides thread, fabric, yarn, 10 cloth or textile comprising a polypeptide of the invention, or an enzymatically active fragment thereof.

The invention provides methods for making ethanol comprising contacting an organic material, e.g. a biomass with a polypeptide having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, 15 and/or an epimerase activity, wherein the polypeptide has a sequence of the invention, or the polypeptide is encoded by a nucleic acid comprising a sequence of the invention, or an enzymatically active fragment thereof. The invention provides compositions comprising an ethanol and a polypeptide having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase 20 activity, wherein the polypeptide has a sequence of the invention, or the polypeptide is encoded by a nucleic acid comprising a sequence of the invention, or an enzymatically active fragment thereof. The invention provides methods for making ethanol comprising: (a) providing at least one polypeptide having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity, or an enzymatically active fragment thereof; (b) providing an organic 25 composition; and (c) contacting the composition of step (b) with the polypeptide of step (a).

The invention provides methods of making a pharmaceutical (drug) composition, a pharmaceutical (drug) precursor, or a drug intermediate, comprising using a polypeptide of 30 this invention having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity.

The invention provides pharmaceutical (drug) compositions and pharmaceutical (drug) precursors and intermediates comprising a polypeptide having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine

racemase activity, and/or an epimerase activity, wherein the polypeptide comprises a sequence of the invention, or the polypeptide is encoded by a nucleic acid comprising a sequence of the invention, or an enzymatically active fragment thereof. In one aspect, the pharmaceutical composition acts as a digestive aid, is an antibiotic or is useful for
5 treatment of amino acid deficiencies. In one aspect, the treatment is prophylactic.

In one aspect, the invention provides oral care products comprising a polypeptide of the invention having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity, or an
10 isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase encoded by a nucleic acid of the invention. The oral care product can comprise a toothpaste, a dental cream, a gel or a tooth powder, an odontic, a mouth wash, a pre- or post brushing rinse formulation, a chewing gum, a lozenge or a candy. The invention provides contact lens cleaning compositions comprising a polypeptide of the invention having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase
15 activity, an alanine racemase activity, and/or an epimerase activity, or an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase encoded by a nucleic acid of the invention.

The invention provides chimeric isomerases, e.g., racemases, e.g., amino acid racemases, alanine racemases, and/or epimerases comprising a polypeptide sequence of
20 the invention and at least one heterologous domain, e.g. a binding domain or a dockerin domain. The invention provides methods for designing a chimeric isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase having a new specificity or an enhanced specificity, comprising inserting a heterologous or an additional endogenous domain, e.g. a binding domain or a dockerin domain, into an isomerase, e.g., a
25 racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase, wherein the domain is inserted into an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase sequence of the invention.

The invention provides enzyme mixtures, or “cocktails” comprising at least one enzyme of the invention and one or more other enzyme(s), which can be another
30 isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase, or any other enzyme; for example, the “cocktails” of the invention, in addition to at least one enzyme of this invention, can comprise any other enzyme, such as xylanase, cellulases, lipases, esterases, proteases, or endoglycosidases, endo-beta.-1,4-glucanases, beta-glucanases, endo-beta-1,3(4)-glucanases, cutinases, peroxidases, catalases, laccases,

amylases, glucoamylases, pectinases, transferases, transaminases, amino transferases, dehydrogenases, oxidoreductases, reductases, oxidases, phenoloxidases, ligninases, pullulanases, arabinanases, hemicellulases, mannanases, xyloglucanases, pectin acetyl esterases, rhamnogalacturonan acetyl esterases, polygalacturonases, 5 rhamnogalacturonases, galactanases, pectin lyases, pectin methylesterases, cellobiohydrolases and/or transglutaminases, to name just a few examples. In alternative embodiments, these enzyme mixtures, or “cocktails” comprising at least one enzyme of the invention can be used in any process or method of the invention, or composition of the invention, e.g., in foods or feeds, food or feed supplements, textiles, papers, processed 10 woods, etc. and methods for making them, and in compositions and methods for treating paper, pulp, wood, paper, pulp or wood waste or by-products, and the like, and in the final products thereof.

The details of one or more embodiments of the invention are set forth in the accompanying drawings and the description below. Other features, objects, and 15 advantages of the invention will be apparent from the description and drawings, and from the claims.

All publications, patents, patent applications, GenBank sequences and ATCC deposits, cited herein are hereby expressly incorporated by reference for all purposes.

BRIEF DESCRIPTION OF THE DRAWINGS

20 The following drawings are illustrative of aspects of the invention and are not meant to limit the scope of the invention as encompassed by the claims.

The patent or application file contains at least one drawing executed in color. Copies of this patent or patent application publication with color drawing(s) will be provided by the Office upon request and payment of the necessary fee.

25 Figure 1 is a block diagram of a computer system.

Figure 2 is a flow diagram illustrating one aspect of a process for comparing a new nucleotide or protein sequence with a database of sequences in order to determine the homology levels between the new sequence and the sequences in the database.

30 Figure 3 is a flow diagram illustrating one aspect of a process in a computer for determining whether two sequences are homologous.

Figure 4 is a flow diagram illustrating one aspect of an identifier process 300 for detecting the presence of a feature in a sequence.

Like reference symbols in the various drawings indicate like elements.

DETAILED DESCRIPTION OF THE INVENTION

The invention provides isomerases, e.g., racemases, e.g., amino acid racemases, alanine racemases, and/or epimerases, and polynucleotides encoding them and methods of making and using them. Isomerases, e.g., racemases, e.g., amino acid racemases, alanine racemases, and/or epimerases, of the polypeptides of the invention encompasses enzymes
5 having isomerase activity, e.g., racemase activity, e.g., amino acid racemase activity, alanine racemase activity, and/or epimerase activity, and/or catalyze the re-arrangement of atoms within a molecule, catalyze the conversion of one isomer into another, catalyze the conversion of an optically active substrate into a raceme, which is optically inactive,
10 catalyze the interconversion of substrate enantiomers, catalyze the stereochemical inversion around the asymmetric carbon atom in a substrate having only one center of asymmetry, catalyze the stereochemical inversion of the configuration around an asymmetric carbon atom in a substrate having more than one asymmetric center, and/or catalyze the racemization of amino acids. The enzymes, e.g., racemases of the invention
15 can be used to make and/or process pharmaceutical (drug) compositions, pharmaceutical (drug) precursors and intermediates, such as molecules comprising unnatural amino acids or antibiotics, sweeteners, peptide enzymes, peptide hormones, fuel and fuel additive compositions, foods and food additives, beverage and beverage additives, feeds and feed additives, drugs and drug additives, dietary supplements, textiles, wood, paper, pulp,
20 detergents and the like.

In one aspect, an enzyme of the invention is thermotolerant and/or tolerant of high and/or low pH conditions. For example, in one aspect, an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase of the invention retains activity under conditions comprising a temperature of at least about 80°C, 85°C,
25 86°C, 87°C, 88°C, 89°C, 90°C, 91°C, 92°C, 93°C, 94°C, 95°C, 96°C, 97°C, 98°C, 99°C, 100°C, 101°C, 102°C, 103°C, 103.5°C, 104°C, 105°C, 107°C, 108°C, 109°C or 110°C, or more, and a basic pH of at least about pH 11, or more.

The invention provides isolated, synthetic or recombinant nucleic acids comprising a nucleic acid encoding at least one polypeptide having an isomerase activity, e.g., a
30 racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity, or other activity as described herein, wherein the nucleic acid comprises a sequence having at least about 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%,

88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more, or complete (100%) sequence identity (homology) to SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, 5 SEQ ID NO:29, SEQ ID NO:31, SEQ ID NO:33, SEQ ID NO:35, SEQ ID NO:37, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:43, SEQ ID NO:45, SEQ ID NO:47, SEQ ID NO:49, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:57, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:89, SEQ ID NO:91, 10 SEQ ID NO:93, SEQ ID NO:95, SEQ ID NO:97, SEQ ID NO:99, SEQ ID NO:101, SEQ ID NO:103, SEQ ID NO:105, SEQ ID NO:107, SEQ ID NO:109, SEQ ID NO:111, SEQ ID NO:113, SEQ ID NO:115, SEQ ID NO:117, SEQ ID NO:119, SEQ ID NO:121, SEQ ID NO:123, SEQ ID NO:125, SEQ ID NO:127, SEQ ID NO:129, SEQ ID NO:131, SEQ ID NO:133, SEQ ID NO:135, SEQ ID NO:137, SEQ ID NO:139, SEQ ID NO:141, SEQ ID NO:143, SEQ ID NO:145, SEQ ID NO:147, SEQ ID NO:149, SEQ ID NO:151, SEQ ID NO:153, SEQ ID NO:155, SEQ ID NO:157, SEQ ID NO:159, SEQ ID NO:161, SEQ ID NO:163, SEQ ID NO:165, SEQ ID NO:167, SEQ ID NO:169, SEQ ID NO:171, SEQ ID NO:173, SEQ ID NO:175, SEQ ID NO:177, SEQ ID NO:179, SEQ ID NO:181, SEQ ID NO:183, SEQ ID NO:185, SEQ ID NO:187, SEQ ID NO:189, SEQ ID NO:191, SEQ ID NO:193, SEQ ID NO:195, SEQ ID NO:197, SEQ ID NO:199, SEQ ID NO:201, SEQ ID NO:203, SEQ ID NO:205, SEQ ID NO:207, SEQ ID NO:209, SEQ ID NO:211, SEQ ID NO:213, SEQ ID NO:215, SEQ ID NO:217, SEQ ID NO:219, SEQ ID NO:221, SEQ ID NO:223, SEQ ID NO:225, SEQ ID NO:227, SEQ ID NO:229, SEQ ID NO:231, SEQ ID NO:233, SEQ ID NO:235, SEQ ID NO:237, SEQ ID NO:239, SEQ ID NO:241, SEQ ID NO:243, SEQ ID NO:245, SEQ ID NO:247, SEQ ID NO:249, SEQ ID NO:251, SEQ ID NO:253, SEQ ID NO:255, SEQ ID NO:257, SEQ ID NO:259, SEQ ID NO:261, SEQ ID NO:263, SEQ ID NO:265, SEQ ID NO:267, SEQ ID NO:269, SEQ ID NO:271, SEQ ID NO:273, SEQ ID NO:275, SEQ ID NO:277, SEQ ID NO:279, SEQ ID NO:281, SEQ ID NO:283, SEQ ID NO:285, SEQ ID NO:287, SEQ ID NO:289, SEQ ID NO:291, SEQ ID NO:293, SEQ ID NO:295, SEQ ID NO:297, SEQ ID NO:299, SEQ ID NO:301, SEQ ID NO:303, SEQ ID NO:305, SEQ ID NO:307, SEQ ID NO:309, SEQ ID NO:311, SEQ ID NO:313, SEQ ID NO:315, SEQ ID NO:317, SEQ ID NO:319, SEQ ID NO:321, SEQ ID NO:323, SEQ ID NO:325, SEQ ID NO:327, SEQ ID NO:329, SEQ ID NO:331, SEQ

ID NO:333, SEQ ID NO:335, SEQ ID NO:337, SEQ ID NO:339, SEQ ID NO:341, SEQ ID NO:343, SEQ ID NO:345, SEQ ID NO:347, SEQ ID NO:349, SEQ ID NO:351, SEQ ID NO:353, SEQ ID NO:355, SEQ ID NO:357, SEQ ID NO:359, SEQ ID NO:361, SEQ ID NO:363, SEQ ID NO:365, SEQ ID NO:367, SEQ ID NO:369, SEQ ID NO:371, SEQ ID NO:373, SEQ ID NO:375, SEQ ID NO:377, SEQ ID NO:379, SEQ ID NO:381, SEQ ID NO:383, SEQ ID NO:385, SEQ ID NO:387, SEQ ID NO:389, SEQ ID NO:391, SEQ ID NO:393, SEQ ID NO:395, SEQ ID NO:397, SEQ ID NO:399, SEQ ID NO:401, SEQ ID NO:403, SEQ ID NO:405, SEQ ID NO:407, SEQ ID NO:409, SEQ ID NO:411, SEQ ID NO:413, SEQ ID NO:415, SEQ ID NO:417, SEQ ID NO:419, SEQ ID NO:421, SEQ ID NO:423, SEQ ID NO:425, SEQ ID NO:427, SEQ ID NO:429, SEQ ID NO:431, SEQ ID NO:433, SEQ ID NO:435, SEQ ID NO:437, SEQ ID NO:439, SEQ ID NO:441, SEQ ID NO:443, SEQ ID NO:445, SEQ ID NO:447, SEQ ID NO:449, SEQ ID NO:451, SEQ ID NO:453, SEQ ID NO:455, SEQ ID NO:457, SEQ ID NO:459, SEQ ID NO:461, SEQ ID NO:463, SEQ ID NO:465, SEQ ID NO:467, SEQ ID NO:469, SEQ ID NO:471, SEQ ID NO:473, SEQ ID NO:475, SEQ ID NO:477, SEQ ID NO:479, SEQ ID NO:481, SEQ ID NO:483, SEQ ID NO:485, SEQ ID NO:487, SEQ ID NO:489, SEQ ID NO:491, SEQ ID NO:493, SEQ ID NO:495 or SEQ ID NO:497, and as described herein and in Tables 1, 2 and 3, and the Sequence Listing (all of these sequences are “exemplary polynucleotides of the invention”), and enzymatically active subsequences (fragments) thereof, over a region of between about 10 to 2500, or more residues, or the full length of a cDNA, transcript (mRNA) or gene. Nucleic acids of the invention includes those encoding a polypeptide of this invention, having at least 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more, or 100% (complete) sequence identity to an exemplary polypeptide of the invention, which includes, e.g., SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:22, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:28, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:42, SEQ ID NO:44, SEQ ID NO:46, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:58, SEQ ID NO:60, SEQ ID NO:62, SEQ ID NO:64, SEQ ID NO:66, SEQ ID NO:68, SEQ ID NO:70, SEQ ID NO:72, SEQ ID NO:74, SEQ ID NO:76, SEQ ID NO:78, SEQ ID NO:80, SEQ ID NO:82, SEQ ID NO:84,

SEQ ID NO:86, SEQ ID NO:88, SEQ ID NO:90, SEQ ID NO:92, SEQ ID NO:94, SEQ
ID NO:96, SEQ ID NO:98, SEQ ID NO:100, SEQ ID NO:102, SEQ ID NO:104, SEQ ID
NO:106, SEQ ID NO:108, SEQ ID NO:110, SEQ ID NO:112, SEQ ID NO:114, SEQ ID
NO:116, SEQ ID NO:118, SEQ ID NO:120, SEQ ID NO:122, SEQ ID NO:124, SEQ ID
5 NO:126, SEQ ID NO:128, SEQ ID NO:130, SEQ ID NO:132, SEQ ID NO:134, SEQ ID
NO:136, SEQ ID NO:138, SEQ ID NO:140, SEQ ID NO:142, SEQ ID NO:143, SEQ ID
NO:146, SEQ ID NO:148, SEQ ID NO:150, SEQ ID NO:152, SEQ ID NO:154, SEQ ID
NO:156, SEQ ID NO:158, SEQ ID NO:160, SEQ ID NO:162, SEQ ID NO:164, SEQ ID
NO:166, SEQ ID NO:168, SEQ ID NO:170, SEQ ID NO:172, SEQ ID NO:174, SEQ ID
10 NO:176, SEQ ID NO:178, SEQ ID NO:180, SEQ ID NO:182, SEQ ID NO:184, SEQ ID
NO:186, SEQ ID NO:188, SEQ ID NO:190, SEQ ID NO:192, SEQ ID NO:194, SEQ ID
NO:196, SEQ ID NO:198, SEQ ID NO:200, SEQ ID NO:202, SEQ ID NO:204, SEQ ID
NO:206, SEQ ID NO:208, SEQ ID NO:210, SEQ ID NO:212, SEQ ID NO:214, SEQ ID
NO:216, SEQ ID NO:218, SEQ ID NO:220, SEQ ID NO:222, SEQ ID NO:224, SEQ ID
15 NO:226, SEQ ID NO:228, SEQ ID NO:230, SEQ ID NO:232, SEQ ID NO:234, SEQ ID
NO:236, SEQ ID NO:238, SEQ ID NO:240, SEQ ID NO:242, SEQ ID NO:244, SEQ ID
NO:246, SEQ ID NO:248, SEQ ID NO:250, SEQ ID NO:252, SEQ ID NO:254, SEQ ID
NO:256, SEQ ID NO:258, SEQ ID NO:260, SEQ ID NO:262, SEQ ID NO:264, SEQ ID
NO:266, SEQ ID NO:268, SEQ ID NO:270, SEQ ID NO:272, SEQ ID NO:274, SEQ ID
20 NO:276, SEQ ID NO:278, SEQ ID NO:280, SEQ ID NO:282, SEQ ID NO:284, SEQ ID
NO:286, SEQ ID NO:288, SEQ ID NO:290, SEQ ID NO:292, SEQ ID NO:294, SEQ ID
NO:296, SEQ ID NO:298, SEQ ID NO:300, SEQ ID NO:302, SEQ ID NO:304, SEQ ID
NO:306, SEQ ID NO:308, SEQ ID NO:310, SEQ ID NO:312, SEQ ID NO:314, SEQ ID
NO:316, SEQ ID NO:318, SEQ ID NO:320, SEQ ID NO:322, SEQ ID NO:324, SEQ ID
25 NO:326, SEQ ID NO:328, SEQ ID NO:330, SEQ ID NO:332, SEQ ID NO:334, SEQ ID
NO:336, SEQ ID NO:338, SEQ ID NO:340, SEQ ID NO:342, SEQ ID NO:344, SEQ ID
NO:346, SEQ ID NO:348, SEQ ID NO:350, SEQ ID NO:352, SEQ ID NO:354, SEQ ID
NO:356, SEQ ID NO:358, SEQ ID NO:360, SEQ ID NO:362, SEQ ID NO:364, SEQ ID
NO:366, SEQ ID NO:368, SEQ ID NO:370, SEQ ID NO:372, SEQ ID NO:374, SEQ ID
30 NO:376, SEQ ID NO:378, SEQ ID NO:380, SEQ ID NO:382, SEQ ID NO:384, SEQ ID
NO:386, SEQ ID NO:388, SEQ ID NO:390, SEQ ID NO:392, SEQ ID NO:394, SEQ ID
NO:396, SEQ ID NO:398, SEQ ID NO:400, SEQ ID NO:402, SEQ ID NO:404, SEQ ID
NO:406, SEQ ID NO:408, SEQ ID NO:410, SEQ ID NO:412, SEQ ID NO:414, SEQ ID
NO:416, SEQ ID NO:418, SEQ ID NO:420, SEQ ID NO:422, SEQ ID NO:424, SEQ ID

NO:426, SEQ ID NO:428, SEQ ID NO:430, SEQ ID NO:432, SEQ ID NO:434, SEQ ID NO:436, SEQ ID NO:438, SEQ ID NO:440, SEQ ID NO:442, SEQ ID NO:444, SEQ ID NO:446, SEQ ID NO:448, SEQ ID NO:450, SEQ ID NO:452, SEQ ID NO:454, SEQ ID NO:456, SEQ ID NO:458, SEQ ID NO:460, SEQ ID NO:462, SEQ ID NO:464, SEQ ID NO:466, SEQ ID NO:468, SEQ ID NO:470, SEQ ID NO:472, SEQ ID NO:474, SEQ ID NO:476, SEQ ID NO:478, SEQ ID NO:480, SEQ ID NO:482, SEQ ID NO:484, SEQ ID NO:486, SEQ ID NO:488, SEQ ID NO:490, SEQ ID NO:492, SEQ ID NO:494, SEQ ID NO:496 or SEQ ID NO:498, including the sequences described herein and in Tables 1, 2 and 3, below, and in the Sequence Listing (all of these sequences are “exemplary enzymes/ polypeptides of the invention”), and enzymatically active subsequences (fragments) thereof.

Tables 1, 2 and 3, below, are charts describing selected characteristics of exemplary nucleic acids and polypeptides of the invention, including sequence identity comparison of the exemplary sequences to public databases.

Table 1, below, describes the assigned activity (as determined by experimental data, see Examples 1-19, below) of the exemplary polypeptides (encoded by the exemplary polynucleotides) of the invention. Table 1 further indicates whether the polynucleotide (encoding a polypeptide) of the invention is a clone (a genomic sequence isolated from the original source, as described in Table 2) or is a subclone (where the clone is manipulated by, e.g. removal of a native signal sequence, addition of a start Methionine, addition of a tag, etc). Table 1 also indicates the clone and subclone relationship, e.g. which subclone was derived from which clone. For aid in reading Table 1, for example, Columns 1 and 4, rows 1 and 2, indicate that SEQ ID NO:34 (encoded by SEQ ID NO:33) is a clone with the corresponding subclone being SEQ ID NO:464 (encoded by SEQ ID NO:463), denoted as “Clone/subclone pair 1”.

Table 2, below, indicates the source from which the exemplary nucleic acids and polypeptides of the invention were first derived. Table 2, below, also indicates the “Signalp Cleavage Site” for the exemplary enzyme’s signal sequence (or “signal peptide”, or SP), as determined by the paradigm Signalp, as discussed below (see Nielsen (1997), *infra*); the “Predicted Signal Sequence” is listed from the amino terminal to the carboxy terminal, for example, for the polypeptide SEQ ID NO:42, the signal peptide is “MPFCRTLLAVSLGLLITGQAPLYA” (amino acids 1-24 of SEQ ID NO:42).

Table 3, below describes selected characteristics of exemplary nucleic acids and polypeptides of the invention, including sequence identity comparison of the exemplary sequences to public databases. To further aid in reading Table 3, for example, the first

row, labeled "SEQ ID NO:", the numbers "1, 2" represent the exemplary polypeptide of the invention having a sequence as set forth in SEQ ID NO:2, encoded by, e.g., SEQ ID NO:1. All sequences described in Table 2 (all the exemplary sequences of the invention) have been subject to a BLAST search (as described in detail, below) against two sets of
5 databases. The first database set is available through NCBI (National Center for Biotechnology Information). All results from searches against these databases are found in the columns entitled "NR Description", "NR Accession Code", "NR Evaluate" or "NR Organism". "NR" refers to the Non-Redundant nucleotide database maintained by NCBI. This database is a composite of GenBank, GenBank updates, and EMBL updates. The
10 entries in the column "NR Description" refer to the definition line in any given NCBI record, which includes a description of the sequence, such as the source organism, gene name/protein name, or some description of the function of the sequence. The entries in the column "NR Accession Code" refer to the unique identifier given to a sequence record. The entries in the column "NR Evaluate" refer to the Expect value (Evaluate), which
15 represents the probability that an alignment score as good as the one found between the query sequence (the sequences of the invention) and a database sequence would be found in the same number of comparisons between random sequences as was done in the present BLAST search. The entries in the column "NR Organism" refer to the source organism of the sequence identified as the closest BLAST hit. The second set of databases is
20 collectively known as the GENESEQTM database, which is available through Thomson Derwent (Philadelphia, PA). All results from searches against this database are found in the columns entitled "GENESEQTM Protein Description", "GENESEQTM Protein Accession Code", "Evaluate", "GENESEQTM DNA Description", "GENESEQTM DNA Accession Code" or "Evaluate". The information found in these columns is comparable to
25 the information found in the NR columns described above, except that it was derived from BLAST searches against the GENESEQTM database instead of the NCBI databases. In addition, this table includes the column "Predicted EC No.". An EC number is the number assigned to a type of enzyme according to a scheme of standardized enzyme nomenclature developed by the Enzyme Commission of the Nomenclature Committee of the
30 International Union of Biochemistry and Molecular Biology (IUBMB). The results in the "Predicted EC No." column are determined by a BLAST search against the Kegg (Kyoto Encyclopedia of Genes and Genomes) database. If the top BLAST match has an Evaluate equal to or less than e-6, the EC number assigned to the top match is entered into the table. The EC number of the top hit is used as a guide to what the EC number of the sequence of

the invention might be. The columns “Query DNA Length” and “Query Protein Length” refer to the number of nucleotides or the number amino acids, respectively, in the sequence of the invention that was searched or queried against either the NCBI or GENESEQ™ databases. The columns “Subject DNA Length” and “Subject Protein Length” refer to the number of nucleotides or the number amino acids, respectively, in the sequence of the top match from the BLAST searches. The results provided in these columns are from the search that returned the lower Evalue, either from the NCBI databases or the Geneseq database. The columns “%ID Protein” and “%ID DNA” refer to the percent sequence identity between the sequence of the invention and the sequence of the top BLAST match. The results provided in these columns are from the search that returned the lower Evalue, either from the NCBI databases or the GENESEQ™ database.

Table 1

<u>Clone/subclone pair</u>	<u>SEQ ID NO:</u>	<u>Activity</u>	<u>Sequence type (Clone or subclone)</u>
1	33, 34	Racemase	Clone
1	463, 464	Racemase	Subclone
2	131, 132	Racemase	Clone
2	457, 458	Racemase	Subclone
3	13, 14	Racemase	Clone
3	387, 388	Racemase	Subclone
4	25, 26	Racemase	Clone
4	389, 390	Racemase	Subclone
5	23, 24	Racemase	Clone
5	391, 392	Racemase	Subclone
6	61, 62	Racemase	Clone
6	411, 412	Racemase	Subclone (leaderless)
6	489, 490	Racemase	Subclone
7	297, 298	Racemase	Clone
7	467, 468	Racemase	Subclone
8	11, 12	Racemase	Clone
8	397, 398	Racemase	Subclone
9	311, 312	Racemase	Clone
9	469, 470	Racemase	Subclone
10	3, 4	Racemase	Clone
10	393, 394	Racemase	Subclone
11	49, 50	Racemase	Clone
11	423, 424	Racemase	Subclone
12	367, 368	Racemase	Clone
12	471, 472	Racemase	Subclone
13	41, 42	Racemase	Clone
13	399, 400	Racemase	Subclone (leaderless)
13	491, 492	Racemase	Subclone
14	43, 44	Racemase	Clone
14	431, 432	Racemase	Subclone

<u>Clone/subclone pair</u>	<u>SEQ ID NO:</u>	<u>Activity</u>	<u>Sequence type (Clone or subclone)</u>
15	45, 46	Racemase	Clone
15	409, 410	Racemase	Subclone (leaderless)
15	495, 496	Racemase	Subclone
16	47, 48	Racemase	Clone
16	407, 408	Racemase	Subclone (leaderless)
16	493, 494	Racemase	Subclone
17	287, 288	Racemase	Clone
17	453, 454	Racemase	Subclone
18	35, 36	Racemase	Clone
18	385, 386	Racemase	Subclone
19	51, 52	Racemase	Clone
19	401, 402	Racemase	Subclone (leaderless)
19	497, 498	Racemase	Subclone
20	53, 54	Racemase	Clone
20	403, 404	Racemase	Subclone (leaderless)
20	427, 428	Racemase	Subclone
21	55, 56	Racemase	Clone
21	433, 434	Racemase	Subclone
22	107, 108	Racemase	Clone
22	473, 474	Racemase	Subclone
23	57, 58	Racemase	Clone
23	405, 406	Racemase	Subclone (leaderless)
23	429, 430	Racemase	Subclone
24	109, 110	Racemase	Clone
24	415, 416	Racemase	Subclone
25	121, 122	Racemase	Clone
25	425, 426	Racemase	Subclone
26	301, 302	Racemase	Clone
26	477, 478	Racemase	Subclone
27	123, 124	Racemase	Clone
27	439, 440	Racemase	Subclone
28	125, 126	Racemase	Clone
28	461, 462	Racemase	Subclone
29	299, 300	Racemase	Clone
29	475, 476	Racemase	Subclone
30	111, 112	Racemase	Clone
30	417, 418	Racemase	Subclone
31	119, 120	Racemase	Clone
31	459, 460	Racemase	Subclone
32	113, 114	Racemase	Clone
32	435, 436	Racemase	Subclone
33	115, 116	Racemase	Clone
33	419, 420	Racemase	Subclone
34	117, 118	Racemase	Clone
34	421, 422	Racemase	Subclone
35	223, 224	Racemase	Clone
35	441, 442	Racemase	Subclone
36	217, 218	Racemase	Clone
36	443, 444	Racemase	Subclone
37	233, 234	Racemase	Clone

Clone/subclone pair	SEQ ID NO:	Activity	Sequence type (Clone or subclone)
37	445, 446	Racemase	Subclone
38	243, 244	Racemase	Clone
38	447, 448	Racemase	Subclone
39	247, 248	Racemase	Clone
39	449, 450	Racemase	Subclone
40	273, 274	Racemase	Clone
40	451, 452	Racemase	Subclone
41	105, 106	Racemase	Clone
41	465, 466	Racemase	Subclone
42	103, 104	Racemase	Clone
42	437, 438	Racemase	Subclone
43	7, 8	Racemase	Clone
43	413, 414	Racemase	Subclone
44	9, 10	Racemase	Clone
44	395, 396	Racemase	Subclone
45	129, 130	Racemase	Clone
45	455, 456	Racemase	Subclone
	379, 380	Epimerase	Clone
	381, 382	Epimerase	Clone
	369, 370	Epimerase	Clone
	375, 376	Epimerase	Clone
	383, 384	Isomerase	Clone
	373, 374	Epimerase	Clone
	371, 372	Epimerase	Clone
	377, 378	Epimerase	Clone
	17, 18	Racemase	Clone
	19, 20	Racemase	Clone
	15, 16	Racemase	Clone
	29, 30	Racemase	Clone
	27, 28	Racemase	Clone
	255, 256	Racemase	Clone
	321, 322	Racemase	Clone
	323, 324	Racemase	Clone
	327, 328	Racemase	Clone
	307, 308	Racemase	Clone
	303, 304	Racemase	Clone
	309, 310	Racemase	Clone
	305, 306	Racemase	Clone
	21, 22	Racemase	Clone
	479, 480	Racemase	Clone
	313, 314	Racemase	Clone
	315, 316	Racemase	Clone
	1, 2	Racemase	Clone
	85, 86	Racemase	Clone
	87, 88	Racemase	Clone
	89, 90	Racemase	Clone
	91, 92	Racemase	Clone
	93, 94	Racemase	Clone
	99, 100	Racemase	Clone
	77, 78	Racemase	Clone
	331, 332	Racemase	Clone

<u>Clone/subclone pair</u>	<u>SEQ ID NO:</u>	<u>Activity</u>	<u>Sequence type</u> (Clone or subclone)
	345, 346	Racemase	Clone
	347, 348	Racemase	Clone
	333, 334	Racemase	Clone
	325, 326	Racemase	Clone
	319, 320	Racemase	Clone
	335, 336	Racemase	Clone
	349, 350	Racemase	Clone
	339, 340	Racemase	Clone
	341, 342	Racemase	Clone
	343, 344	Racemase	Clone
	355, 356	Racemase	Clone
	353, 354	Racemase	Clone
	351, 352	Racemase	Clone
	317, 318	Racemase	Clone
	329, 330	Racemase	Clone
	167, 168	Racemase	Clone
	213, 214	Racemase	Clone
	285, 286	Racemase	Clone
	289, 290	Racemase	Clone
	37, 38	Racemase	Clone
	39, 40	Racemase	Clone
	483, 484	Racemase	Clone
	485, 486	Racemase	Clone
	487, 488	Racemase	Clone
	31, 32	Racemase	Clone
	101, 102	Racemase	Clone
	169, 170	Racemase	Clone
	171, 172	Racemase	Clone
	59, 60	Racemase	Clone
	135, 136	Racemase	Clone
	173, 174	Racemase	Clone
	137, 138	Racemase	Clone
	337, 338	Racemase	Clone
	357, 358	Racemase	Clone
	359, 360	Racemase	Clone
	361, 362	Racemase	Clone
	363, 364	Racemase	Clone
	365, 366	Racemase	Clone
	175, 176	Racemase	Clone
	177, 178	Racemase	Clone
	179, 180	Racemase	Clone
	181, 182	Racemase	Clone
	143, 144	Racemase	Clone
	187, 188	Racemase	Clone
	189, 190	Racemase	Clone
	133, 134	Racemase	Clone
	145, 146	Racemase	Clone
	481, 482	Racemase	Clone
	63, 64	Racemase	Clone
	193, 194	Racemase	Clone
	153, 154	Racemase	Clone

<u>Clone/subclone pair</u>	<u>SEQ ID NO:</u>	<u>Activity</u>	<u>Sequence type</u> (Clone or subclone)
	155, 156	Racemase	Clone
	195, 196	Racemase	Clone
	157, 158	Racemase	Clone
	159, 160	Racemase	Clone
	161, 162	Racemase	Clone
	163, 164	Racemase	Clone
	165, 166	Racemase	Clone
	65, 66	Racemase	Clone
	147, 148	Racemase	Clone
	149, 150	Racemase	Clone
	191, 192	Racemase	Clone
	151, 152	Racemase	Clone
	71, 72	Racemase	Clone
	69, 70	Racemase	Clone
	73, 74	Racemase	Clone
	75, 76	Racemase	Clone
	95, 96	Racemase	Clone
	97, 98	Racemase	Clone
	79, 80	Racemase	Clone
	81, 82	Racemase	Clone
	83, 84	Racemase	Clone
	67, 68	Racemase	Clone
	183, 184	Racemase	Clone
	139, 140	Racemase	Clone
	141, 142	Racemase	Clone
	185, 186	Racemase	Clone
	197, 198	Racemase	Clone
	199, 200	Racemase	Clone
	201, 202	Racemase	Clone
	203, 204	Racemase	Clone
	205, 206	Racemase	Clone
	207, 208	Racemase	Clone
	209, 210	Racemase	Clone
	211, 212	Racemase	Clone
	219, 220	Racemase	Clone
	221, 222	Racemase	Clone
	225, 226	Racemase	Clone
	227, 228	Racemase	Clone
	229, 230	Racemase	Clone
	231, 232	Racemase	Clone
	235, 236	Racemase	Clone
	237, 238	Racemase	Clone
	239, 240	Racemase	Clone
	241, 242	Racemase	Clone
	293, 294	Racemase	Clone
	245, 246	Racemase	Clone
	295, 296	Racemase	Clone
	249, 250	Racemase	Clone
	251, 252	Racemase	Clone
	253, 254	Racemase	Clone
	257, 258	Racemase	Clone

Clone/subclone pair	SEQ ID NO:	Activity	Sequence type (Clone or subclone)
	259, 260	Racemase	Clone
	261, 262	Racemase	Clone
	263, 264	Racemase	Clone
	265, 266	Racemase	Clone
	267, 268	Racemase	Clone
	269, 270	Racemase	Clone
	271, 272	Racemase	Clone
	275, 276	Racemase	Clone
	277, 278	Racemase	Clone
	279, 280	Racemase	Clone
	281, 282	Racemase	Clone
	283, 284	Racemase	Clone
	291, 292	Racemase	Clone
	215, 216	Racemase	Clone
	5, 6	Racemase	Clone
	127, 128	Racemase	Clone

Table 2

SEQ ID NO:	Source	Signalp Cleavage Site	Predicted Signal Sequence
1, 2	Aquifex aeolicus		
3, 4	Aquifex aeolicus		
5, 6	Bacteria		
7, 8	Bacteria		
9, 10	Bacteria		
11, 12	Unknown		
13, 14	Unknown		
15, 16	Unknown		
17, 18	Unknown		
19, 20	Unknown		
21, 22	Unknown		
23, 24	Unknown		
25, 26	Unknown		
27, 28	Pelagibacter ubique		
29, 30	Unknown		
31, 32	Unknown		
33, 34	Unknown		
35, 36	Unknown		
37, 38	Unknown		
39, 40	Unknown		
41, 42	Unknown	Probability: 0.999 AA1: 24 AA2: 25	MPFCRTLLAVSLGLLITGQAPLYA
43, 44	Unknown	Probability: 0.999 AA1: 24 AA2: 25	MPFCRTLLAVSLGLLITGQAPLYA
45, 46	Unknown	Probability: 1.000 AA1: 24 AA2: 25	MPFSRTLLAVSLGLLITGQAPLYA
47, 48	Unknown	Probability: 1.000 AA1: 24 AA2: 25	MPFSRTLLAVSLGLLITGQAPLYA

SEQ ID NO:	Source	Signalp Cleavage Site	Predicted Signal Sequence
49, 50	Unknown	Probability: 1.000 AA1: 24 AA2: 25	MPFCRTLLAASLALLITGQAPLYA
51, 52	Unknown	Probability: 1.000 AA1: 23 AA2: 24	MPFRRTLLALSGLVWQGVHA
53, 54	Unknown	Probability: 1.000 AA1: 23 AA2: 24	MPFCRTLLALSGLVWQQAHA
55, 56	Unknown	Probability: 1.000 AA1: 23 AA2: 24	MPFCRTLLALSGLVWQGVHA
57, 58	Unknown	Probability: 1.000 AA1: 24 AA2: 25	MPFSRTLLAASLALLITGQAPLYA
59, 60	Unknown	Probability: 1.000 AA1: 24 AA2: 25	MPFCRTLLAASLALLITGQAPLYA
61, 62	Unknown	Probability: 0.999 AA1: 33 AA2: 34	MFTMIFMKKKFCLLFATIILFITCLC FLLKSVS
63, 64	Unknown	Probability: 1.000 AA1: 24 AA2: 25	MPFRRTLLAASLALLITGLAPLYA
65, 66	Unknown	Probability: 1.000 AA1: 24 AA2: 25	MPFPRTLLAASLALLITGQAPLYA
67, 68	Unknown	Probability: 1.000 AA1: 24 AA2: 25	MPFRRTLLAASLALLVTAQAPLYA
69, 70	Unknown	Probability: 1.000 AA1: 24 AA2: 25	MPFCRTLLAASLALLITGQAPLYA
71, 72	Unknown	Probability: 1.000 AA1: 24 AA2: 25	MPFRRTLLAASLALLITGQAPLYA
73, 74	Unknown	Probability: 1.000 AA1: 24 AA2: 25	MPFCRTLLAASLALLITGQAPLYA
75, 76	Unknown	Probability: 1.000 AA1: 24 AA2: 25	MPFPRTLLAASLALLITGQAPLYA
77, 78	Unknown	Probability: 1.000 AA1: 24 AA2: 25	MPFRRTLLAASLALLITGQAPLYA
79, 80	Unknown	Probability: 1.000 AA1: 24 AA2: 25	MPFCRTLLAASLALLITGQAPLYA
81, 82	Unknown	Probability: 1.000 AA1: 24 AA2: 25	MPFCRTLLAASLALLITGQAPLYA
83, 84	Unknown	Probability: 1.000 AA1: 24 AA2: 25	MPFCRTLLAASLALLITGQAPLYA
85, 86	Unknown	Probability: 1.000 AA1: 24 AA2: 25	MPFSRTLLAASLALLITGQAPLFA
87, 88	Unknown	Probability: 1.000 AA1: 24 AA2: 25	MPFCRTLLAASLALLITGQAPLYA
89, 90	Unknown	Probability: 1.000 AA1: 24 AA2: 25	MPFRRTLLAASLALLITGQAPLFA
91, 92	Unknown	Probability: 1.000 AA1: 24 AA2: 25	MPFCRTLLAASLALLITGQAPLYA
93, 94	Unknown	Probability: 1.000 AA1: 24 AA2: 25	MPFRRTLLAASLALLITGQAPLYA
95, 96	Unknown	Probability: 1.000 AA1: 24 AA2: 25	MPFPRTLLAASLALLITGQAPLYA
97, 98	Unknown	Probability: 1.000 AA1: 24 AA2: 25	MPFRRTLLAASLALLITGQAPLYA
99, 100	Unknown	Probability: 1.000 AA1: 24 AA2: 25	MPFCRTLLAASLALLITGQAPLYA
101, 102	Unknown		
103, 104	Unknown	Probability: 0.826 AA1: 19 AA2: 20	MKNNKCIAILGGMGQASS
105, 106	Unknown		

SEQ ID NO:	Source	Signalp Cleavage Site	Predicted Signal Sequence
107, 108	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLILGLLAGQAVA
109, 110	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLIFGLLAGQAVA
111, 112	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLILGLLAGQAVA
113, 114	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLVFGLLAGQAVA
115, 116	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLVFGLLAGQAVA
117, 118	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLVFGLLAGQAVA
119, 120	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLVFGLLAGQAVA
121, 122	Unknown		
123, 124	Unknown		
125, 126	Unknown		
127, 128	Unknown		
129, 130	Unknown		
131, 132	Unknown		
133, 134	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLILGLLAGQAVA
135, 136	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLIFGLLAGQAVA
137, 138	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLIFGLLAGQAVA
139, 140	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLILGLLAGQAVA
141, 142	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLIFGLLAGQAVA
143, 144	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLILGLLAGQAVA
145, 146	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLILGLLAGQAVA
147, 148	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLILGLLAGQAVA
149, 150	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLVFGLLAGQAVA
151, 152	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLVFGLLAGQAVA
153, 154	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLILGLLAGQAVA
155, 156	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLILGLLAGQAVA
157, 158	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLVFGLLAGQAVA
159, 160	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLVFGLLAGQAVA
161, 162	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLILGLLAGQAVA
163, 164	Unknown		
165, 166	Unknown		
167, 168	Unknown		
169, 170	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLILGLLAGQAVA

SEQ ID NO:	Source	Signalp Cleavage Site	Predicted Signal Sequence
171, 172	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLILGLLAGQAVA
173, 174	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLIFGLLAGQAVA
175, 176	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLVFGLLAGQAVA
177, 178	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLILGLLAGQAVA
179, 180	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLILGLLAGQAVA
181, 182	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLILGLLAGQAVA
183, 184	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLILGLLAGQAVA
185, 186	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLIFGLLAGQAVA
187, 188	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLILGLLAGQAVA
189, 190	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLILGLLAGQAVA
191, 192	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLVFGLLAGQAVA
193, 194	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLVFGLLAGQAVA
195, 196	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLVFGLLAGQAVA
197, 198	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLVFGLLAGQAVA
199, 200	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLIFGLLAGQAVA
201, 202	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLIFGLLAGQAVA
203, 204	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLIFGLLAGQAVA
205, 206	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLIFGLLAGQAVA
207, 208	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLVFGLLAGQAVA
209, 210	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLVFGLLAGQAVA
211, 212	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLVFGLLAGQAVA
213, 214	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLVFGLLAGQAVA
215, 216	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLILGLLAGQAVA
217, 218	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLVFGLLAGQAVA
219, 220	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLILGLLAGQAVA
221, 222	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLILGLLAGQAVA
223, 224	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLILGLLAGQAVA
225, 226	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLILGLLAGQAVA

SEQ ID NO:	Source	Signalp Cleavage Site	Predicted Signal Sequence
227, 228	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLVFGLLAGQAVA
229, 230	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLVFGLLAGQAVA
231, 232	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLILGLLAGQAVA
233, 234	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLILGLLAGQAVA
235, 236	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLIFGLLAGQAVA
237, 238	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLILGLLAGQAVA
239, 240	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLILGLLAGQAVA
241, 242	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLILGLLAGQAVA
243, 244	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLIFGLLAGQAVA
245, 246	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLIFGLLAGQAVA
247, 248	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLIFGLLAGQAVA
249, 250	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLIFGLLAGQAVA
251, 252	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLIFGLLAGQAVA
253, 254	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLIFGLLAGQAVA
255, 256	Unknown		
257, 258	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLILGLLAGQAVA
259, 260	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLILGLLAGQAVA
261, 262	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLILGLLAGQAVA
263, 264	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLILGLLAGQAVA
265, 266	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLILGLLAGQAVA
267, 268	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLILGLLAGQAVA
269, 270	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLVGLLAGQAVA
271, 272	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLVFGLLAGQAVA
273, 274	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLILGLLAGQAVA
275, 276	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLILGLLAGQAVA
277, 278	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLVFGLLAGQAVA
279, 280	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLVFGLLAGQAVA
281, 282	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLVFGLLAGQAVA
283, 284	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLVFGLLAGQAVA

SEQ ID NO:	Source	Signalp Cleavage Site	Predicted Signal Sequence
285, 286	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLILGLLAGQAVA
287, 288	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLILGLLAGQAVA
289, 290	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLILGLLAGQAVA
291, 292	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLVFGLLAGQAVA
293, 294	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLILGLLAGQAVA
295, 296	Unknown	Probability: 1.000 AA1: 21 AA2: 22	MHKKTLLATLIFGLLAGQAVA
297, 298	Unknown		
299, 300	Unknown	Probability: 1.000 AA1: 23 AA2: 24	MPFTRTVLALSLGLVLLQSQVHA
301, 302	Unknown	Probability: 1.000 AA1: 22 AA2: 23	MKFTPTLLAVALAGCLSTQVQA
303, 304	Unknown		
305, 306	Unknown		
307, 308	Unknown		
309, 310	Unknown		
311, 312	Unknown		
313, 314	Unknown		
315, 316	Unknown		
317, 318	Unknown		
319, 320	Unknown		
321, 322	Unknown		
323, 324	Unknown		
325, 326	Unknown		
327, 328	Unknown		
329, 330	Unknown		
331, 332	Unknown		
333, 334	Unknown		
335, 336	Unknown		
337, 338	Unknown		
339, 340	Unknown		
341, 342	Unknown		
343, 344	Unknown		
345, 346	Unknown		
347, 348	Unknown		
349, 350	Unknown		
351, 352	Unknown		
353, 354	Unknown		
355, 356	Unknown		
357, 358	Unknown		
359, 360	Unknown		
361, 362	Unknown		
363, 364	Unknown		
365, 366	Unknown		
367, 368	Pseudomonas straminea ATCC 33636		

SEQ ID NO:	Source	Signalp Cleavage Site	Predicted Signal Sequence
369, 370	Unknown		
371, 372	Unknown		
373, 374	Unknown		
375, 376	Unknown		
377, 378	Unknown		
379, 380	Unknown		
381, 382	Unknown		
383, 384	Unknown		
385, 386	Unknown		
387, 388	Unknown		
389, 390	Unknown		
391, 392	Unknown		
393, 394	Unknown		
395, 396	Unknown		
397, 398	Unknown		
399, 400	Unknown		
401, 402	Unknown		
403, 404	Unknown		
405, 406	Unknown		
407, 408	Unknown		
409, 410	Unknown		
411, 412	Unknown		
413, 414	Unknown		
415, 416	Unknown		
417, 418	Unknown		
419, 420	Unknown		
421, 422	Unknown		
423, 424	Unknown		
425, 426	Unknown		
427, 428	Unknown	Probability: 1.000 AA1: 23 AA2: 24	MPFCRTLALSLGLVLWQQQAHA
429, 430	Unknown	Probability: 1.000 AA1: 24 AA2: 25	MPFSRTLLAASLALLITGQAPLYA
431, 432	Unknown		
433, 434	Unknown		
435, 436	Unknown		
437, 438	Unknown	Probability: 0.826 AA1: 19 AA2: 20	MKNNKCIAILGGMGPQASS
439, 440	Unknown		
441, 442	Unknown		
443, 444	Unknown		
445, 446	Unknown		
447, 448	Unknown		
449, 450	Unknown		
451, 452	Unknown		
453, 454	Unknown		
455, 456	Unknown		
457, 458	Unknown	Probability: 0.742 AA1: 30 AA2: 31	MARVLRWARSAYIRITTGSHALF ADATLA
459, 460	Unknown		
461, 462	Unknown		

SEQ ID NO:	Source	Signalp Cleavage Site	Predicted Signal Sequence
463, 464	Unknown		
465, 466	Unknown		
467, 468	Unknown		
469, 470	Unknown		
471, 472	Unknown		
473, 474	Unknown		
475, 476	Unknown		
477, 478	Unknown		
479, 480	Unknown		
481, 482	Unknown		
483, 484	Unknown		
485, 486	Unknown		
487, 488	Unknown		

5

[REST OF PAGE INTENTIONALLY LEFT BLANK]

Table 3

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Geneseq DNA Description	Geneseq DNA Accession Code	Genes eq DNA Accession Code	Eval ue	Predi cted EC Number	Qu ery DN A Le ngt h	Qu ery Pro tein Le ngt h	Subj ect DN A Le ngt h	Subj ect Prot ein Len gth	% ID Pro tein	% ID DN A
1, 2	hypothetical protein [Aquifex aeolicus].	1560 5814	1.00 E-128	Aquifex aeolicus	Prokaryotic essential gene #34740.	ABU25 646	Drosophila melanogaster polypeptide SEQ ID NO: 24465.	ABLO 3829	0.48	2.7.3.	68 7	22 8	228	687	10 0	10 0	
3, 4	alanine racemase [Aquifex aeolicus].	1560 6873	0	Aquifex aeolicus	Aquifex pyrophilus heat resistant alanine racemase encoding DNA.	ABB06 296	Aquifex pyrophilus heat resistant alanine racemase encoding DNA.	ABL4 9607	2.00 E-07	5.1.1.1	10 17	33 8	0 338	0	10 0		
5, 6	hypothetical protein SAV6126 [Streptomyces avermitilis MA-4680]	2983 2668	1.00 E-101	Streptomyces avermitilis MA-4680	Prokaryotic essential gene #34740.	ABU19 499	Human immune/haematopoietic antigen genomic sequence SEQ ID NO:41436.	AAK8 0650	0.03 2	2.7.3.	72 0	23 9	0 239	0	80		

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Genes eq Protein Accession Code	Geneseq DNA Description	Genes eq DNA Accession Code	Genes Evaluate	Predicted EC Number	Query DN A Length	Query Protein Length	Subject DN A Length	Subject Protein Length	% ID Protein	% ID DN A
7, 8	hypothetical protein SAV4292 [Streptomyces avermitilis MA-4680]	29830835	0	Streptomyces avermitilis MA-4680	Propionibacterium acnes predicted ORF-encoded polypeptide #300.	ABM54358	EST clone EP219.	AAV88076	2.00 E-53	0.048	1032	343	0	343	89	
9, 10	alanine racemase [Streptomyces coelicolor A3(2)]	21223124	1.00 E-166	Streptomyces coelicolor A3(2)	Prokaryotic essential gene #34740.	ABU34223	Pseudomonas aeruginosa polypeptide #3.	ABD08732	6.00 E-90	5.1.1.1	1149	382	0	391	78	
11, 12	alanine racemase [Rhodospseudomonas palustris HaA2]	86748627	1.00 E-106	Rhodospseudomonas palustris HaA2	Pseudomonas aeruginosa polypeptide #3.	ABO84274	Pseudomonas aeruginosa polypeptide #3.	ABD06378	4.00 E-50	5.1.1.1	1122	373	0	388	58	
13, 14	alanine racemase [Azoarcus sp. EbN1]	56477426	1.00 E-128	Azoarcus sp. EbN1	Prokaryotic essential gene #34740.	ABU41398	Pseudomonas aeruginosa polypeptide #3.	ABD08732	1.00 E-110	5.1.1.1	1065	354	0	353	64	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Genes eq Protein Accession Code	Evaluate	Geneseq DNA Description	Genes eq DNA Accession Code	Evaluate	Predicted EC Number	Query DNA Length	Query Protein Length	Subject DNA Length	Subject Protein Length	% ID Protein	% ID DNA
15, 16	Aspartate racemase [Marinobacter algicola DG893]	1.49 E+08	4.00 E-93	Marinobacter algicola DG893	Klebsiella pneumoniae polypeptide seqid 7178.	ABO65 503	1.00 E-71	Klebsiella pneumoniae polypeptide seqid 7178.	ACH9 9054	0.00 E-2	5.1.1.13	702	233	0	231	74	
17, 18	glutamate racemase [Cytophaga hutchinsonii ATCC 33406]	1.11 E+08	1.00 E-106	Cytophaga hutchinsonii ATCC 33406	Prokaryotic essential gene #34740.	ABU25 174	2.00 E-44	Human phosphodiesterase 4D amino acid sequence N1 SEQ ID NO:7.	AEB8 5185	0.00 E-9	5.1.1.3	807	268	0	269	72	
19, 20	glutamate racemase [Geobacter sulfurreducens PCA]	3999 8014	1.00 E-74	Geobacter sulfurreducens PCA	M. xanthus protein sequence, seq id 9726.	ABM90 755	2.00 E-67	M. xanthus protein sequence, seq id 9726.	ACL6 4794	0.03 E-7	5.1.1.3	822	273	0	272	53	
21, 22	Putative decarboxylase [Bordetella parapertussis 12822]	3359 6748	5.00 E-40	Bordetella parapertussis 12822	Thermococcus kodakaraensis KOD1 protein sequence SeqID4.	ADN46 910	5.00 E-27	Bacterial polypeptide #10001.	ADS6 1806	1.00 E-04	5.2.1.1	768	255	0	265	41	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Geneseq DNA Description	Geneseq DNA Accession Code	Eval	Predicted EC Number	Query DNALength	Query ProteinLength	Subject DNALength	Subject ProteinLength	% ID Protein	% ID DNA
23, 24	alanine racemase [Enterobacter sp. 638]	1.46 E+08	0	Enterobacter sp. 638	Enterobacter cloacae protein amino acid sequence - SEQ ID 5666.	AEH63094	Enterobacter cloacae protein amino acid sequence - SEQ ID 5666.	AEH55699	0	5.1.1.1	1080	359	0	359	95	
25, 26	alanine racemase [Enterobacter sp. 638]	1.46 E+08	0	Enterobacter sp. 638	Enterobacter cloacae protein amino acid sequence - SEQ ID 5666.	AEH63094	Enterobacter cloacae protein amino acid sequence - SEQ ID 5666.	AEH55699	0	5.1.1.1	1080	359	0	359	95	
27, 28	hypothetical protein SAR11_0361 [Candidatus Pelagibacter ubique HTCC1062]	7108 3067	1.00 E-121	Candidatus Pelagibacter ubique HTCC1062	Prokaryotic essential gene #34740.	ABU24290	Prokaryotic essential gene #34740.	ACA27845	0.002	2.7.3.	651	216	0	216	100	
29, 30	putative proline racemase [Brucella ovis ATCC 25840]	1.49 E+08	1.00 E-130	Brucella ovis ATCC 25840	Pseudomonas aeruginosa polypeptide #3.	ABO82134	Pseudomonas aeruginosa polypeptide #3.	ABD15676	1.00 E-08	5.1.1.4	1032	343	0	342	64	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Genes eq Protein Accession Code	Eval ue	Geneseq DNA Description	Genes eq DNA Accession Code	Eval ue	Predicted EC Number	Qu ery DN A Le ngt h	Qu ery Protein Le ngt h	Subj ect DN A Le ngt h	Subj ect Protein Len gth	% ID Protein	% ID DN A
31, 32	proline racemase, putative [Stappia aggregata IAM 12614] gij118435940 gb E AV42584.1 proline racemase, putative [Stappia aggregata IAM 12614]	1.19 E+08	1.00 E-109	Stappia aggregata IAM 12614	Acinetobacter baumannii protein #19.	ADA35 228	1.00 E-105	Pseudomonas aeruginosa polypeptide #3.	ABD1 5666	9.00 E-16	5.1.1.4	10 11	33 6	0	310	58	
33, 34	alanine racemase [Mesorhizobium sp. BNC1] putative alanine racemase [Aurantimonas sp. SI85-9A1] gij9033811 gb E AS51762.1 putative alanine racemase [Aurantimonas sp. SI85-9A1]	1.11 E+08	1.00 E-124	Mesorhizobium sp. BNC1	Prokaryotic essential gene #34740.	ABU38 829	3.00 E-46	Human cancer associated cDNA SEQ ID NO 9.	AEH9 8530	0.01 E-3	5.1.1.1	11 16	37 1	0	385	61	
35, 36	putative alanine racemase [Aurantimonas sp. SI85-9A1] gij9033811 gb E AS51762.1 putative alanine racemase [Aurantimonas sp. SI85-9A1]	9041 8439	6.00 E-49	Aurantimonas sp. SI85-9A1	Achromobacter xylosoxidans DTA SEQ ID NO 6.	AEH19 277	5.00 E-47	N. meningitidis partial DNA sequence gnm_640 SEQ ID NO:640.	AAA8 1486	3.2	5.1.1.1	11 01	36 6	0	368	33	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Genes eq Protein Accession Code	Geneseq DNA Description	Genes eq DNA Accession Code	Eval ue	Predicted EC Number	Qu ery DN A Le ngth	Qu ery Protein Le ngth	Subj ect DN A Le ngth	Subj ect Protein Le ngth	% ID Protein	% ID DNA
37, 38	Alanine racemase [Magnetospirillum gryphiswaldense MSR-1]	1.45 E+08	1.00 E-38	Magnetospirillum gryphiswaldense MSR-1	Aquifex pyrophilus heat resistant alanine racemase encoding DNA.	ABB06 296	Pseudomonas aeruginosa polypeptide #3.	ABD1 2586	2.00 E-20	5.1.1.1	489	162	0	369	51	
39, 40	putative proline racemase [Burkholderia xenovorans LB400]	9177 9222	2.00 E-36	Burkholderia xenovorans LB400	Prokaryotic essential gene #34740.	ABU21 813	Prokaryotic essential gene #34740.	ACA2 6332	5.00 E-37	5.1.1.4	372	123	960	318		
41, 42	alanine racemase [Pseudomonas putida F1]	1.49 E+08	0	Pseudomonas putida F1	T. maritima D-alanine-D-alanine ligase.	AED11 803	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	0	5.1.1.1	1230	409	0	409	97	
43, 44	alanine racemase [Pseudomonas putida F1]	1.49 E+08	0	Pseudomonas putida F1	T. maritima D-alanine-D-alanine ligase.	AED11 803	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	0	5.1.1.1	1230	409	0	409	96	
45, 46	alanine racemase [Pseudomonas putida F1]	1.49 E+08	0	Pseudomonas putida F1	T. maritima D-alanine-D-alanine ligase.	AED11 803	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	0	5.1.1.1	1230	409	0	409	97	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Genes eq Protein Accession Code	Geneseq DNA Description	Genes eq DNA Accession Code	Eval ue	Predicted EC Number	Query DN A Length	Query Protein Length	Subj ect DN A Length	Subj ect Protein Length	% ID Protein	% ID DN A
47, 48	alanine racemase [Pseudomonas putida F1]	1.49 E+08	0	Pseudomonas putida F1	T. maritima D-alanine-D-alanine ligase.	AED11 803	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	0	5.1.1.1	1230	409	0	409	97	
49, 50	alanine racemase [Pseudomonas putida F1]	1.49 E+08	0	Pseudomonas putida F1	T. maritima D-alanine-D-alanine ligase.	AED11 803	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	0	5.1.1.1	1230	409	0	409	96	
51, 52	alanine racemase [Pseudomonas putida GB-1] gij126314851 gb E AZ66019.1 alanine racemase [Pseudomonas putida GB-1]	1.26 E+08	0	Pseudomonas putida GB-1	T. maritima D-alanine-D-alanine ligase.	AED11 803	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9537	1.00 E-120	5.1.1.1	1230	409	0	409	77	
53, 54	alanine racemase [Pseudomonas putida GB-1] gij126314851 gb E AZ66019.1 alanine racemase [Pseudomonas putida GB-1]	1.26 E+08	1.00 E-180	Pseudomonas putida GB-1	T. maritima D-alanine-D-alanine ligase.	AED11 803	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9537	1.00 E-124	5.1.1.1	1230	409	0	409	77	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Geneseq DNA Description	Geneseq DNA Accession Code	Genes Evaluate	Predicted EC Number	Query DNA Length	Query Protein Length	Subject DNA Length	Subject Protein Length	% ID Protein	% ID DNA
55, 56	alanine racemase [Pseudomonas putida GB-1] gij126314851 gb E AZ66019.1 alanine racemase [Pseudomonas putida GB-1]	1.26 E+08	1.00 E-180	Pseudomonas putida GB-1	T. maritima D-alanine-D-alanine ligase.	AED11803	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99537	1.00 E-124	5.1.1.1	1230	409	0	409	77	
57, 58	alanine racemase [Pseudomonas putida F1]	1.49 E+08	0	Pseudomonas putida F1	T. maritima D-alanine-D-alanine ligase.	AED11804	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	0	5.1.1.1	1230	409	0	409	96	
59, 60	alanine racemase [Pseudomonas putida F1]	1.49 E+08	0	Pseudomonas putida F1	T. maritima D-alanine-D-alanine ligase.	AED11804	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	0	5.1.1.1	1230	409	0	409	95	
61, 62	alanine racemase [Fusobacterium nucleatum subsp. polymorphum ATCC 10953]	1.48 E+08	7.00 E-19	Fusobacterium nucleatum subsp. polymorphum ATCC 10953	Aquifex pyrophilus heat resistant alanine racemase encoding DNA.	ABB06296	Human prostate expression marker cDNA 6604.	ABV52203	1.00 E-12	5.1.1.1	1614	537	0	354	19	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Genes eq Protein Accession Code	Geneseq DNA Description	Genes eq DNA Accession Code	Eval ue	Predicted EC Number	Query DN A Length	Query Protein Length	Subject DN A Length	Subject Protein Length	% ID Protein	% ID DN A
63, 64	alanine racemase [Pseudomonas putida F1]	1.49 E+08	0	Pseudomonas putida F1	T. maritima D-alanine-D-alanine ligase.	AED11 803	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9537	0	5.1.1.1	12 30	40 9	0	409	98	
65, 66	alanine racemase [Pseudomonas putida F1]	1.49 E+08	0	Pseudomonas putida F1	T. maritima D-alanine-D-alanine ligase.	AED11 803	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9537	0	5.1.1.1	12 30	40 9	0	409	97	
67, 68	alanine racemase [Pseudomonas putida GB-1] gi 126314851 gb E AZ66019.1 alanine racemase [Pseudomonas putida GB-1]	1.26 E+08	0	Pseudomonas putida GB-1	T. maritima D-alanine-D-alanine ligase.	AED11 803	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	0	5.1.1.1	12 30	40 9	0	409	90	
69, 70	alanine racemase [Pseudomonas putida GB-1] gi 126314851 gb E AZ66019.1 alanine racemase [Pseudomonas putida GB-1]	1.26 E+08	0	Pseudomonas putida GB-1	T. maritima D-alanine-D-alanine ligase.	AED11 803	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9537	0	5.1.1.1	12 30	40 9	0	409	95	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Genes eq Protein Accession Code	Eval ue	Geneseq DNA Description	Genes eq DNA Accession Code	Eval ue	Predicted EC Number	Qu ery DN A Length	Qu ery Protein Length	Subj ect DN A Length	Subj ect Protein Length	% ID Protein	% ID DN A
71, 72	alanine racemase [Pseudomonas putida GB-1] gi 126314851 gb E AZ66019.1 alanine racemase [Pseudomonas putida GB-1]	1.26 E+08	0	Pseudomonas putida GB-1	T. maritima D-alanine-D-alanine ligase.	AED11 803	0	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9537	0	5.1.1.1	12 30	40 9	0	409	96	
73, 74	alanine racemase [Pseudomonas putida GB-1] gi 126314851 gb E AZ66019.1 alanine racemase [Pseudomonas putida GB-1]	1.26 E+08	0	Pseudomonas putida GB-1	T. maritima D-alanine-D-alanine ligase.	AED11 803	0	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9537	0	5.1.1.1	12 30	40 9	0	409	95	
75, 76	alanine racemase [Pseudomonas putida GB-1] gi 126314851 gb E AZ66019.1 alanine racemase [Pseudomonas putida GB-1]	1.26 E+08	0	Pseudomonas putida GB-1	T. maritima D-alanine-D-alanine ligase.	AED11 803	0	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9537	0	5.1.1.1	12 30	40 9	0	409	95	
77, 78	alanine racemase [Pseudomonas putida F1]	1.49 E+08	0	Pseudomonas putida F1	T. maritima D-alanine-D-alanine ligase.	AED11 803	0	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	0	5.1.1.1	12 30	40 9	0	409	95	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Genes eq Protein Accession Code	Geneseq DNA Description	Genes eq DNA Accession Code	Eval ue	Predicted EC Number	Qu ery DN A Length	Qu ery Protein Length	Subj ect DN A Length	Subj ect Protein Length	% ID Protein	% ID DN A
79, 80	alanine racemase [Pseudomonas putida F1]	1.49 E+08	0	Pseudomonas putida F1	T. maritima D-alanine-D-alanine ligase.	AED11 804	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	0	5.1.1.1	12 30	40 9	0	409	95	
81, 82	alanine racemase [Pseudomonas putida F1]	1.49 E+08	0	Pseudomonas putida F1	T. maritima D-alanine-D-alanine ligase.	AED11 804	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	0	5.1.1.1	12 30	40 9	0	409	96	
83, 84	alanine racemase [Pseudomonas putida F1]	1.49 E+08	0	Pseudomonas putida F1	T. maritima D-alanine-D-alanine ligase.	AED11 804	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	0	5.1.1.1	12 30	40 9	0	409	94	
85, 86	alanine racemase [Pseudomonas putida F1]	1.49 E+08	0	Pseudomonas putida F1	T. maritima D-alanine-D-alanine ligase.	AED11 803	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9537	0	5.1.1.1	12 30	40 9	0	409	96	
87, 88	alanine racemase [Pseudomonas putida F1]	1.49 E+08	0	Pseudomonas putida F1	T. maritima D-alanine-D-alanine ligase.	AED11 803	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	0	5.1.1.1	12 30	40 9	0	409	96	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Genes eq Protein Accession Code	Geneseq DNA Description	Genes eq DNA Accession Code	Eval ue	Predicted EC Number	Query DN A Length	Query Protein Length	Subject DN A Length	Subject Protein Length	% ID Protein	% ID DN A
89, 90	alanine racemase [Pseudomonas putida KT2440]	2699 0430	0	Pseudomonas putida KT2440	T. maritima D-alanine-D-alanine ligase.	AED11 803	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9537	0	5.1.1.1	12 30	40 9	0	409	96	96
91, 92	alanine racemase [Pseudomonas putida F1]	1.49 E+08	0	Pseudomonas putida F1	T. maritima D-alanine-D-alanine ligase.	AED11 803	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	0	5.1.1.1	12 30	40 9	0	409	96	96
93, 94	alanine racemase [Pseudomonas putida F1] alanine racemase [Pseudomonas putida GB-1] gi 126314851 gb E AZ66019.1	1.49 E+08	0	Pseudomonas putida F1	T. maritima D-alanine-D-alanine ligase.	AED11 804	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	0	5.1.1.1	12 30	40 9	0	409	96	96
95, 96	alanine racemase [Pseudomonas putida GB-1]	1.26 E+08	0	Pseudomonas putida GB-1	T. maritima D-alanine-D-alanine ligase.	AED11 803	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9537	0	5.1.1.1	12 30	40 9	0	409	95	95

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Genes eq Protein Accession Code	Eval ue	Geneseq DNA Description	Genes eq DNA Accession Code	Eval ue	Predicted EC Number	Qu ery DN A Le ngth	Qu ery Protein Le ngth	Subj ect DN A Len gth	Subj ect Protein Len gth	% ID Protein	% ID DN A
97, 98	alanine racemase [Pseudomonas putida GB-1] gij126314851 gb E AZ66019.1 alanine racemase [Pseudomonas putida GB-1]	1.26 E+08	0	Pseudomonas putida GB-1	T. maritima D-alanine-D-alanine ligase.	AED11803	0	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99537	0	5.1.1.1	1230	409	0	409	96	
99, 100	alanine racemase [Pseudomonas putida F1]	1.49 E+08	0	Pseudomonas putida F1	T. maritima D-alanine-D-alanine ligase.	AED11803	0	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99537	0	5.1.1.1	1230	409	0	409	97	
101, 102	alanine racemase 2, catabolic [Escherichia coli O157:H7 EDL933].	15801412	2.00 E-58	Escherichia coli O157:H7 EDL933	Prokaryotic essential gene #34740.	ABU45089	2.00 E-58	Enterobacter cloacae protein amino acid sequence - SEQ ID 5666.	AEH53103	5.00 E-27	5.1.1.1	384	107	356	86	76	
103, 104	aspartate racemase [Pyrococcus abyssii].	14521575	8.00 E-29	Pyrococcus abyssii	Thermococcus kodakaraensis KOD1 protein sequence SeqID4.	ADN46207	2.00 E-25	Prokaryotic essential gene #34740.	ACA27539	0.13	5.1.1.13	714	228	687	32	48	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Geneseq DNA Description	Geneseq DNA Accession Code	Eval ue	Predic ted EC Number	Qu ery DN A Length	Qu ery Protein Length	Subj ect DN A Length	Subj ect Protein Length	% ID Protein	% ID DN A
105, 106	proline racemase [Microscilla marina ATCC 23134] gij 123984081 gb EAY24454.1]	1.24 E+08	1.00 E-150	Microscilla marina ATCC 23134	Bacterial polypeptide #10001.	ADS22 995	Bovine ABCG2 related PCR primer #31.	AEN6 9487	1.00 E-128	5.1.1.4	1002	333	0	333	75	
107, 108	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter brevis mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	1.00 E-131	5.1.1.1	1227	408	0	408	96	
109, 110	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter brevis mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	1.00 E-129	5.1.1.1	1227	408	0	408	95	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Genes eq Protein Accession Code	Evaluate	Geneseq DNA Description	Genes eq DNA Accession Code	Evaluate	Predicted EC Number	Query DN A Length	Query Protein Length	Subject DN A Length	Subject Protein Length	% ID Protein	% ID DNA
111, 112	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	1.00 E-131	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	2.00 E-111	5.1.1.1	12 27	40 8	0	408	99	
113, 114	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	1.00 E-131	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	2.00 E-141	5.1.1.1	12 24	40 7	0	408	92	
115, 116	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	1.00 E-132	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	2.00 E-08	5.1.1.1	12 27	40 8	0	408	92	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Geneseq DNA Description	Geneseq DNA Accession Code	Eval	Predicted EC Number	Query DN A Length	Query Protein Length	Subject DN A Length	Subject Protein Length	% ID Protein	% ID DNA
117, 118	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter brevis mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	6.00 E-11	5.1.1.1	12 27	40 8	0	408	92	
119, 120	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter brevis mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	1.00 E-12	5.1.1.1	12 27	40 8	0	408	95	
121, 122	alanine racemase [Pseudomonas putida F1]	1.49 E+08	1.00 E-136	Pseudomonas putida F1	T. maritima D-alanine-D-alanine ligase.	AED11 803	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	2.00 E-26	5.1.1.1	10 86	36 1	122 7	409		
123, 124	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter brevis mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	6.00 E-11	5.1.1.1	10 86	36 1	0	408	88	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Genes eq Protein Accession Code	Evaluate	Geneseq DNA Description	Genes eq DNA Accession Code	Evaluate	Predicted EC Number	Query DN A Length	Query Protein Length	Subject DN A Length	Subject Protein Length	% ID Protein	% ID DNA
125, 126	alanine racemase [Pseudomonas putida GB-1] gij126314851 gb E AZ66019.1 alanine racemase [Pseudomonas putida GB-1]	1.26 E+08	1.00 E-155	Pseudomonas putida GB-1	Empedobacter brevis mature peptide synthetase enzyme SEQ ID NO:3.	AED10581	1.00 E-159	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	4.00 E-95	5.1.1.1	1086	361	1227	386		
127, 128	glutamate racemase [Bradyrhizobium japonicum USDA 110] proline racemase, putative [Stappia aggregata IAM 12614] gij118435940 gb E AV42584.1 proline racemase, putative [Stappia aggregata IAM 12614]	27380813	3.00 E-26	Bradyrhizobium japonicum USDA 110	Photorhabdus luminescens protein sequence #59.	ABM68637	7.00 E-18	Prokaryotic essential gene #34740.	ACA37866	2.2	5.1.1.3	261	86	0	265		63
129, 130	putative proline racemase [Myxococcus xanthus DK 1622]	1.19 E+08	1.00 E-103	Stappia aggregata IAM 12614	Prokaryotic essential gene #34740.	ABU21813	1.00 E-101	Prokaryotic essential gene #34740.	ACA26332	5.00 E-11	5.1.1.4	957	318	0	310		56
131, 132	putative proline racemase [Myxococcus xanthus DK 1622]	1.09 E+08	5.00 E-75	Myxococcus xanthus DK 1622	M. xanthus protein sequence, seq id 9726.	ABM96637	1.00 E-75	Prokaryotic essential gene #34740.	ACA26332	1.00 E-18	5.1.1.4	1074	357	960	311		

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Evaluate	Geneseq DNA Description	Geneseq DNA Accession Code	Evaluate	Predicted EC Number	Query DNA Length	Query Protein Length	Subject DNA Length	Subject Protein Length	% ID Protein	% ID DNA
133, 134	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	1.00 E-130	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	6.00 E-11	5.1.1.1	12 27	40 8	0	408	99	
135, 136	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	1.00 E-129	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	6.00 E-05	5.1.1.1	12 27	40 8	0	408	95	
137, 138	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	1.00 E-128	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	0.00 E-4	5.1.1.1	12 21	40 6	0	408	95	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Evaluate	Geneseq DNA Description	Geneseq DNA Accession Code	Evaluate	Predicted EC Number	Query DN Length	Query Protein Length	Subject DN Length	Subject Protein Length	% ID Protein	% ID DNA
139, 140	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	1.00 E-131	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	6.00 E-11	5.1.1.1	12 24	40 7	0	408	99	
141, 142	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	1.00 E-130	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	1.00 E-15	5.1.1.1	12 27	40 8	0	408	89	
143, 144	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	1.00 E-132	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	6.00 E-11	5.1.1.1	12 27	40 8	0	408	100	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Gene Eval	Geneseq DNA Description	Geneseq DNA Accession Code	Gene Eval	Predicted EC Number	Query DNA Length	Query Protein Length	Subject DNA Length	Subject Protein Length	% ID Protein	% ID DNA
145, 146	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-131	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	2.00 E-11	5.1.1.1	1224	407	0	408	99	
147, 148	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-131	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	3.00 E-19	5.1.1.1	1224	407	0	408	97	
149, 150	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-132	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	6.00 E-11	5.1.1.1	1227	408	0	408	93	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Geneseq DNA Description	Geneseq DNA Accession Code	Gene Eval	Predicted EC Number	Query DNA Length	Query Protein Length	Subject DNA Length	Subject Protein Length	% ID Protein	% ID DNA
151, 152	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter brevis mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	1.00 E-131	5.1.1.1	1227	408	0	408	92	
153, 154	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter brevis mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	1.00 E-131	5.1.1.1	1224	407	0	408	97	
155, 156	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter brevis mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	2.00 E-14	5.1.1.1	1227	408	0	408	96	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Genes eq Protein Accession Code	Evaluate	Geneseq DNA Description	Genes eq DNA Accession Code	Evaluate	Predicted EC Number	Query DNA Length	Query Protein Length	Subject DNA Length	Subject Protein Length	% ID Protein	% ID DNA
157, 158	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-132	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	6.00 E-11	5.1.1.1	1227	408	0	408	92	
159, 160	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-131	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	6.00 E-11	5.1.1.1	1227	408	0	408	92	
161, 162	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-131	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	3.00 E-19	5.1.1.1	1227	408	0	408	97	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Geneseq DNA Description	Geneseq DNA Accession Code	Eval	Predicted EC Number	Query DN Length	Query Protein Length	Subject DN Length	Subject Protein Length	% ID Protein	% ID DNA
163, 164	alanine racemase [Pseudomonas putida F1]	1.49 E+08	1.00 E-153	Pseudomonas putida F1	Empedobacter brevis mature peptide synthetase enzyme SEQ ID NO:3.	AED10581	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	1.00 E-64	5.1.1.1	1086	361	1227	386		
165, 166	alanine racemase [Pseudomonas putida F1] proline racemase [Flavobacteriales bacterium HTCC2170] gi 88708932 gb E AR01166.1 proline racemase [Flavobacteriales bacterium HTCC2170]	1.49 E+08	1.00 E-153	Pseudomonas putida F1 Flavobacteriales bacterium HTCC2170	Empedobacter brevis mature peptide synthetase enzyme SEQ ID NO:3. Bacterial polypeptide #10001.	AED10581	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	2.00 E-91	5.1.1.1	1083	360	1227	386		
167, 168		8871 2394	1.00 E-69	HTCC2170	Bacterial polypeptide #10001.	ADS22995	Bacterial polypeptide #10001.	ADS60041	0.00 E-7	5.1.1.1	657	218	0	335		58

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Evaluate	Geneseq DNA Description	Geneseq DNA Accession Code	Evaluate	Predicted EC Number	Query DN A Length	Query Protein Length	Subject DN A Length	Subject Protein Length	% ID Protein	% ID DNA
169, 170	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter brevis mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-131	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	6.00 E-11	5.1.1.1	1227	408	0	408	97	
171, 172	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter brevis mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-131	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	6.00 E-11	5.1.1.1	1227	408	0	408	96	
173, 174	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter brevis mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-129	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	6.00 E-05	5.1.1.1	1224	407	0	408	95	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Gene Eval	Geneseq DNA Description	Geneseq DNA Accession Code	Eval	Predicted EC Number	Query DNA Length	Query Protein Length	Subject DNA Length	Subject Protein Length	% ID Protein	% ID DNA
175, 176	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter brevis mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-130	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	2.00 E-14	5.1.1.1	1227	408	0	408	89	
177, 178	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter brevis mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-132	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	6.00 E-11	5.1.1.1	1227	408	0	408	99	
179, 180	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter brevis mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-132	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	3.00 E-10	5.1.1.1	1227	408	0	408	99	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Evaluate	Geneseq DNA Description	Geneseq DNA Accession Code	Evaluate	Predicted EC Number	Query DN A Length	Query Protein Length	Subject DN A Length	Subject Protein Length	% ID Protein	% ID DN A
181, 182	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	1.00 E-130	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	6.00 E-11	5.1.1.1	12 27	40 8	0	408	99	
183, 184	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	1.00 E-132	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	6.00 E-11	5.1.1.1	12 27	40 8	0	408	100	
185, 186	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	1.00 E-129	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	1.00 E-15	5.1.1.1	12 27	40 8	0	408	89	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Evaluate	Geneseq DNA Description	Geneseq DNA Accession Code	Evaluate	Predicted EC Number	Query DNA Length	Query Protein Length	Subject DNA Length	Subject Protein Length	% ID Protein	% ID DNA
187, 188	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	1.00 E-131	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	6.00 E-11	5.1.1.1	12 27	40 8	0	408	99	
189, 190	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	1.00 E-131	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	6.00 E-11	5.1.1.1	12 27	40 8	0	408	99	
191, 192	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	1.00 E-131	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	6.00 E-11	5.1.1.1	12 27	40 8	0	408	91	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Genes eq Protein Accession Code	Evaluate	Geneseq DNA Description	Genes eq DNA Accession Code	Evaluate	Predicted EC Number	Query DNA Length	Query Protein Length	Subject DNA Length	Subject Protein Length	% ID Protein	% ID DNA
193, 194	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter brevis mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	1.00 E-131	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	6.00 E-11	5.1.1.1	12 27	40 8	0	408	92	
195, 196	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter brevis mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	1.00 E-131	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9537	6.00 E-11	5.1.1.1	12 27	40 8	0	408	92	
197, 198	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter brevis mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	1.00 E-131	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	2.00 E-14	5.1.1.1	12 27	40 8	0	408	91	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Geneseq DNA Description	Geneseq DNA Accession Code	Eval ue	Pred icted EC Num ber	Qu ery DN A Len gth	Qu ery Pro tein Len gth	Subj ect DN A Len gth	Subj ect Prot ein Len gth	% ID Pro tein	% ID DN A
199, 200	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	1.00 E-130	5.1.1.1	12 27	40 8	0	408	90	
201, 202	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	1.00 E-129	5.1.1.1	12 27	40 8	0	408	89	
203, 204	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	4.00 E-18	5.1.1.1	12 27	40 8	0	408	92	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Genes Protein Description	Genes Protein Accession Code	GeneSeq Protein Description	GeneSeq Protein Accession Code	GeneSeq Description	Genes eq DNA Accession Code	GeneSeq Description	GeneSeq Accession Code	GeneSeq Description	GeneSeq Accession Code	Eval ue	Predi cted EC Number	Qu ery DN A Len gth	Qu ery Pro tein Len gth	Subj ect DN A Len gth	Subj ect Prot ein Len gth	% ID Pro tein	% ID DN A	
205, 206	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	1.00 E-130	1.00 E-15	5.1.1.1	12 27	40 8	0	408	89							
207, 208	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	1.00 E-133	3.00 E-16	5.1.1.1	12 24	40 7	0	408	92							
209, 210	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	1.00 E-131	1.00 E-12	5.1.1.1	12 27	40 8	0	408	91							

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Gene Eval	Geneseq DNA Description	Geneseq DNA Accession Code	Gene Eval	Predicted EC Number	Query DN Length	Query Protein Length	Subject DN Length	Subject Protein Length	% ID Protein	% ID DNA
211, 212	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-131	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	2.00 E-14	5.1.1.1	1224	407	0	408	89	
213, 214	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-133	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	1.00 E-15	5.1.1.1	1224	407	0	408	91	
215, 216	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-130	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	6.00 E-11	5.1.1.1	1227	408	0	408	96	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Geneseq DNA Description	Geneseq DNA Accession Code	Eval	Predicted EC Number	Query DNA Length	Query Protein Length	Subject DNA Length	Subject Protein Length	% ID Protein	% ID DNA
217, 218	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	1.00 E-132	7.00 E-141	1224	407	0	408	92	
219, 220	alanine racemase [Aeromonas salmonicida subsp. salmonicida A449]	1.45 E+08	0	Aeromonas salmonicida subsp. salmonicida A449	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	1.00 E-129	6.00 E-111	1227	408	0	408	95	
221, 222	alanine racemase [Aeromonas salmonicida subsp. salmonicida A449]	1.45 E+08	0	Aeromonas salmonicida subsp. salmonicida A449	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	1.00 E-128	6.00 E-111	1224	407	0	408	95	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Geneseq DNA Description	Geneseq DNA Accession Code	Eval E-value	Predicted EC Number	Query DN A Length	Query Protein Length	Subject DN A Length	Subject Protein Length	% ID Protein	% ID DNA
223, 224	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter brevis mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	1.00 E-133	5.1.1.1	1224	407	0	408	93	
225, 226	alanine racemase [Aeromonas salmonicida subsp. salmonicida A449]	1.45 E+08	0	Aeromonas salmonicida subsp. salmonicida A449	Empedobacter brevis mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	1.00 E-130	5.1.1.1	1227	408	0	408	94	
227, 228	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter brevis mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	1.00 E-132	5.1.1.1	1227	408	0	408	91	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Evaluate	Geneseq DNA Description	Geneseq DNA Accession Code	Evaluate	Predicted EC Number	Query DNA Length	Query Protein Length	Subject DNA Length	Subject Protein Length	% ID Protein	% ID DNA
229, 230	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	1.00 E-130	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	3.00 E-13	5.1.1.1	12 27	40 8	0	408	91	
231, 232	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	1.00 E-130	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	6.00 E-11	5.1.1.1	12 27	40 8	0	408	95	
233, 234	alanine racemase [Aeromonas salmonicida subsp. salmonicida A449]	1.45 E+08	0	Aeromonas salmonicida subsp. salmonicida A449	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	1.00 E-128	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	6.00 E-11	5.1.1.1	12 27	40 8	0	408	95	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Geneseq DNA Description	Geneseq DNA Accession Code	Eval ue	Predi cted EC Number	Qu ery DN A Length	Qu ery Protein Length	Subj ect DN A Length	Subj ect Protein Length	% ID Protein	% ID DN A
235, 236	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	1.00 E-128	5.1.1.1	12 24	40 7	0	408	95	
237, 238	alanine racemase [Aeromonas salmonicida subsp. salmonicida A449]	1.45 E+08	0	Aeromonas salmonicida subsp. salmonicida A449	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	1.00 E-128	5.1.1.1	12 27	40 8	0	408	96	
239, 240	alanine racemase [Aeromonas salmonicida subsp. salmonicida A449]	1.45 E+08	0	Aeromonas salmonicida subsp. salmonicida A449	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	1.00 E-128	5.1.1.1	12 24	40 7	0	408	94	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Evaluate	Geneseq DNA Description	Geneseq DNA Accession Code	Evaluate	Predicted EC Number	Query DNA Length	Query Protein Length	Subject DNA Length	Subject Protein Length	% ID Protein	% ID DNA
241, 242	alanine racemase [Aeromonas salmonicida subsp. salmonicida A449]	1.45 E+08	0	Aeromonas salmonicida subsp. salmonicida A449	Empedobacter brevis mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-130	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	6.00 E-11	5.1.1.1	127	408	0	408	95	95
243, 244	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter brevis mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-129	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	4.00 E-06	5.1.1.1	127	408	0	408	95	95
245, 246	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter brevis mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-129	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	4.00 E-06	5.1.1.1	127	408	0	408	95	95

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Evaluate	Geneseq DNA Description	Geneseq DNA Accession Code	Evaluate	Predicted EC Number	Query DN A Length	Query Protein Length	Query DN A Length	Subject DN A Length	Subject Protein Length	% ID Protein	% ID DN A
247, 248	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter brevis mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-130	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	4.00 E-06	5.1.1.1	1227	408	0	408	96		
249, 250	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter brevis mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-129	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	4.00 E-06	5.1.1.1	1227	408	0	408	95		
251, 252	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter brevis mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-130	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	4.00 E-06	5.1.1.1	1227	408	0	408	95		

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Evaluate	Geneseq DNA Description	Geneseq DNA Accession Code	Evaluate	Predicted EC Number	Query DNA Length	Query Protein Length	Subject DNA Length	Subject Protein Length	% ID Protein	% ID DNA
253, 254	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter brevis mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-129	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	4.00 E-06	5.1.1.1	1224	407	0	408	95	
255, 256	hypothetical protein [Mesorhizobium loti].	13474743	1.00 E-170	Mesorhizobium loti	Propionibacterium acnes predicted ORF-encoded polypeptide #300.	ABM37068	1.00 E-77	Plant polypeptide, SEQ ID 5546.	ADT17374	0.2	5.1.1.1	1086	361	1827	608	83	80
257, 258	alanine racemase [Aeromonas salmonicida subsp. salmonicida A449]	1.45 E+08	0	Aeromonas salmonicida subsp. salmonicida A449	Empedobacter brevis mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-129	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	2.00 E-08	5.1.1.1	1227	408	0	408	95	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Evalue	Geneseq DNA Description	Geneseq DNA Accession Code	Evalue	Predicted EC Number	Query DNALength	Query ProteinLength	Subject DNALength	Subject ProteinLength	% ID Protein	% ID DNA
259, 260	alanine racemase [Aeromonas salmonicida subsp. salmonicida A449]	1.45 E+08	0	Aeromonas salmonicida subsp. salmonicida A449	Empedobacter mature peptide synthetase enzyme SEQ ID NO:3.	AED10581	1.00 E-130	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	6.00 E-11	5.1.1.1	127	408	0	408	95	
261, 262	alanine racemase [Aeromonas salmonicida subsp. salmonicida A449]	1.45 E+08	0	Aeromonas salmonicida subsp. salmonicida A449	Empedobacter mature peptide synthetase enzyme SEQ ID NO:3.	AED10581	1.00 E-130	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	2.00 E-08	5.1.1.1	127	408	0	408	94	
263, 264	alanine racemase [Aeromonas salmonicida subsp. salmonicida A449]	1.45 E+08	0	Aeromonas salmonicida subsp. salmonicida A449	Empedobacter mature peptide synthetase enzyme SEQ ID NO:3.	AED10581	1.00 E-128	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	6.00 E-11	5.1.1.1	127	408	0	408	95	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Geneseq Protein Evaluation	Geneseq DNA Description	Geneseq DNA Accession Code	Geneseq Evaluation	Predicted EC Number	Query DNA Length	Query Protein Length	Subject DNA Length	Subject Protein Length	% ID Protein	% ID DNA
265, 266	alanine racemase [Aeromonas salmonicida subsp. salmonicida A449]	1.45 E+08	0	Aeromonas salmonicida subsp. salmonicida A449	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-129	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	6.00 E-11	5.1.1.1	1227	408	0	408	95	
267, 268	alanine racemase [Aeromonas salmonicida subsp. salmonicida A449]	1.45 E+08	0	Aeromonas salmonicida subsp. salmonicida A449	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-129	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	6.00 E-11	5.1.1.1	1227	408	0	408	94	
269, 270	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-132	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	1.00 E-12	5.1.1.1	1227	408	0	408	92	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Genes eq Protein Accession Code	Geneseq DNA Description	Genes eq DNA Accession Code	Eval ue	Predi cted EC Number	Qu ery DN A Length	Qu ery Protein Length	Subj ect DN A Length	Subj ect Protein Length	% ID Protein	% ID DN A
271, 272	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	3.00 E-13	5.1.1.1	12 30	40 9	0	408	92	
273, 274	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	6.00 E-11	5.1.1.1	12 27	40 8	0	408	96	
275, 276	alanine racemase [Aeromonas salmonicida subsp. salmonicida A449]	1.45 E+08	0	Aeromonas salmonicida subsp. salmonicida A449	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	6.00 E-11	5.1.1.1	12 27	40 8	0	408	97	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Genes eq Protein Accession Code	Evaluate	Geneseq DNA Description	Genes eq DNA Accession Code	Evaluate	Predicted EC Number	Query DN A Length	Query Protein Length	Subject DN A Length	Subject Protein Length	% ID Protein	% ID DNA
277, 278	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	1.00 E-133	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	2.00 E-14	5.1.1.1	12 27	40 8	0	408	92	
279, 280	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	1.00 E-132	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	1.00 E-15	5.1.1.1	12 27	40 8	0	408	92	
281, 282	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	1.00 E-132	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	1.00 E-18	5.1.1.1	12 27	40 8	0	408	92	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Evaluate	Geneseq DNA Description	Geneseq DNA Accession Code	Evaluate	Predicted EC Number	Query DNA Length	Query Protein Length	Subject DNA Length	Subject Protein Length	% ID Protein	% ID DNA
283, 284	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter brevis mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-132	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	1.00 E-18	5.1.1.1	1227	408	0	408	92	
285, 286	alanine racemase [Aeromonas salmonicida subsp. salmonicida A449]	1.45 E+08	0	Aeromonas salmonicida subsp. salmonicida A449	Empedobacter brevis mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-130	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	6.00 E-11	5.1.1.1	1227	408	0	408	95	
287, 288	alanine racemase [Aeromonas salmonicida subsp. salmonicida A449]	1.45 E+08	0	Aeromonas salmonicida subsp. salmonicida A449	Empedobacter brevis mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-129	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	6.00 E-11	5.1.1.1	1227	408	0	408	95	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Geneseq DNA Description	Geneseq DNA Accession Code	Eval ue	Predi cted EC Number	Qu ery DNA Length	Qu ery Protein Length	Subj ect DNA Length	Subj ect Protein Length	% ID Protein	% ID DNA
289, 290	alanine racemase [Aeromonas salmonicida subsp. salmonicida A449]	1.45 E+08	0	Aeromonas salmonicida subsp. salmonicida A449	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	1.00 E-129	5.1.1.1	12 27	40 8	0	408	95	
291, 292	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	1.00 E-131	5.1.1.1	12 27	40 8	0	408	92	
293, 294	alanine racemase [Aeromonas salmonicida subsp. salmonicida A449]	1.45 E+08	0	Aeromonas salmonicida subsp. salmonicida A449	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	1.00 E-127	5.1.1.1	12 27	40 8	0	408	94	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Gene Eval	Geneseq DNA Description	Geneseq DNA Accession Code	Gene Eval	Predicted EC Number	Query DNA Length	Query Protein Length	Subject DNA Length	Subject Protein Length	% ID Protein	% ID DNA
295, 296	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empeobacter brevis mature peptide synthetase enzyme SEQ ID NO:3.	AED10581	1.00 E-129	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	4.00 E-06	5.1.1.1	127	408	0	408	95	
297, 298	proline racemase [Burkholderia phymatum STM815] gi 117982269 gb EAU96656.1 proline racemase [Burkholderia phymatum STM815]	1.18 E+08	104	Burkholderia phymatum STM815	Acinetobacter baumannii protein #19.	ADA35228	1.00 E-105	Prokaryotic essential gene #34740.	ACA19650	9.00 E-22	5.1.1.4	960	319	945	331		
299, 300	alanine racemase [Pseudomonas putida GB-1] gi 126314851 gb EAZ66019.1 alanine racemase [Pseudomonas putida GB-1]	1.26 E+08	170	Pseudomonas putida GB-1	T. maritima D-alanine-D-alanine ligase.	AED11803	1.00 E-169	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	5.00 E-95	5.1.1.1	1230	409	0	409	73	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Geneseq Protein Evaluate	Geneseq DNA Description	Geneseq DNA Accession Code	Geneseq Evaluate	Predicted EC Number	Query DN A Length	Query Protein Length	Subject DN A Length	Subject Protein Length	% ID Protein	% ID DN A
301, 302	alanine racemase [Pseudomonas putida KT2440]	2699 0430	1.00 E-143	Pseudomonas putida KT2440	T. maritima D-alanine-D-alanine ligase.	AED11 803	1.00 E-144	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	5.00 E-27	5.1.1.1	122 7	41 2	122 7	409		
303, 304	hypothetical protein PaerP_01003954 [Pseudomonas aeruginosa PA7]	9441 4235	4.00 E-33	Pseudomonas aeruginosa PA7	Pseudomonas aeruginosa polypeptide #3.	ABO82 155	4.00 E-33	Prokaryotic gene #34740.	ACA4 4058	7.00 E-14	5.1.1.4	37 5	12 4	0	314	60	
305, 306	AGR_L_3051p [Agrobacterium tumefaciens].	1589 1641	1.00 E-128	Agrobacterium tumefaciens	L. pneumophila protein SEQ ID NO 3367.	AEB41 596	3.00 E-39	Prokaryotic gene #34740.	ACA2 7267	0.21	5.1.1.1	11 25	37 4	116 7	388	64	66
307, 308	proline racemase [Microscilla marina ATCC 23134] gi 123984081 gb EAY24454.1] proline racemase [Microscilla marina ATCC 23134]	1.24 E+08	3.00 E-59	Microscilla marina ATCC 23134	Bacterial polypeptide #10001.	ADS22 995	3.00 E-47	Bacterial polypeptide #10001.	ADS6 0041	2.00 E-05	5.1.1.4	42 9	14 2	0	333	73	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Genes eq Protein Accession Code	Evaluate	Geneseq DNA Description	Genes eq DNA Accession Code	Evaluate	Predicted EC Number	Query DN A Length	Query Protein Length	Subject DN A Length	Subject Protein Length	% ID Protein	% ID DNA
309, 310	alanine racemase [Stenotrophomonas maltophilia R551-3] g EAX22642.1 alanine racemase [Stenotrophomonas maltophilia R551-3]	1.2E+08	2.00 E-57	Stenotrophomonas maltophilia R551-3	Prokaryotic essential gene #34740.	ABU41398	8.00 E-52	Prokaryotic essential gene #34740.	ACA43665	3.00 E-13	5.1.1.1	408	135	0	355	80	
311, 312	mandelate racemase/muconate lactonizing enzyme family protein [Sulfitobacter sp. NAS-14.1] g EAP81462.1 mandelate racemase/muconate lactonizing enzyme family protein [Sulfitobacter sp. NAS-14.1]	83953326	1.00 E-143	Sulfitobacter sp. NAS-14.1	Klebsiella pneumoniae polypeptide seqid 7178.	ABO61307	2.00 E-71	Novel mar regulated protein (NIMR) #29.	AAS46252	0.67	5.1.2.2	939	312	0	321	81	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Geneseq Evaluate	Geneseq DNA Description	Geneseq DNA Accession Code	Genes Evaluate	Predicted EC Number	Query DNA Length	Query Protein Length	Subject DNA Length	Subject Protein Length	% ID Protein	% ID DNA
313, 314	proline racemase [Mesorhizobium loti].	13473394	1.00 E-120	Mesorhizobium loti	Bacterial polypeptide #10001.	ADS22995	1.00 E-120	C. botulinum active BoNT/A modified open reading frame, SEQ ID No.7.	AEF99607	0.65	5.1.1.4	909	302	1002	333	70	62
315, 316	proline racemase [Flavobacteriales bacterium HTCC2170] gij88708932 gb E AR01166.1 proline racemase [Flavobacteriales bacterium HTCC2170]	88712394	1.00 E-106	Flavobacteriales bacterium HTCC2170	Bacterial polypeptide #10001.	ADS22995	1.00 E-72	Haemophilus influenzae (NTHi) protein - SEQ ID 618.	ADT05504	0.03	5.1.1.4	684	227	0	335	79	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Genes eq Protein Accession Code	Geneseq DNA Description	Genes eq DNA Accession Code	Eval ue	Predicted EC Number	Qu ery DN A Le ngt h	Qu ery Protein Le ngt h	Subj ect DN A Le ngt h	Subj ect Protein Le ngt h	% ID Protein	% ID DN A
317, 318	Mandelate racemase/muconate lactonizing enzyme [Flavobacteriales bacterium HTCC2170] gij88709134 gb E AR01368.1] Mandelate racemase/muconate lactonizing enzyme [Flavobacteriales bacterium HTCC2170]	8871 2596	1.00 E-110	Flavobacteriales bacterium HTCC2170	Bacteroides fragilis strain 14062 protein, SEQ:5227.	AEX28 600	Prokaryotic gene #34740.	ACA2 3465	5.00 E-34	5.5.1.1	10 32	34 3	0	348	57	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Genes eq Protein Accession Code	Geneseq DNA Description	Genes eq DNA Accession Code	Genes Evaluate	Predicted EC Number	Query DN Length	Query Protein Length	Subject DN Length	Subject Protein Length	% ID Protein	% ID DNA
319, 320	COG3938: Proline racemase [Pseudomonas aeruginosa C3719] gi 84328204 ref ZP_00976211.1 COG3938: Proline racemase [Pseudomonas aeruginosa 2192] gi 107100719 ref ZP_01364637.1 hypothetical protein PaerPA_0100174 6 [Pseudomonas aeruginosa PACS2] gi 12616632	8432 1952	1.00 E-60	Pseudomonas aeruginosa C3719	Pseudomonas aeruginosa polypeptide #3.	ABO82 155	Pseudomonas aeruginosa polypeptide #3.	ABD1 5666	4.00 E-61	5.1.1.4	63 6	21 1	486	339		
321, 322	aspartate racemase [Pyrococcus abyssi].	1452 1575	2.00 E-28	Pyrococcus abyssi	Thermococcus kodakaraensis KOD1 protein sequence SeqID4.	ADN46 207	Breast cancer related marker, seq id 2.	ACN8 9360	7.00 E-25	5.1.1.13	69 9	23 2	687	228	31	46

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Genes eq Protein Accession Code	Geneseq DNA Description	Genes eq DNA Accession Code	Genes Evaluate	Predicted EC Number	Query DNA Length	Query Protein Length	Subject DNA Length	Subject Protein Length	% ID Protein	% ID DNA
323, 324	proline racemase, putative [Psychroflexus torquis ATCC 700755] gi 91184469 gb EAS70852.1 proline racemase, putative [Psychroflexus torquis ATCC 700755]	91217361	1.00 E-152	Psychroflexus torquis ATCC 700755	Bacterial polypeptide #10001.	ADS22995	Bacterial polypeptide #10001.	ADS60041	1.00 E-110	5.1.1.4	885	294	0	336	85	
325, 326	proline racemase [Microscilla marina ATCC 23134] gi 123984081 gb EAY24454.1 proline racemase [Microscilla marina ATCC 23134]	1.24 E+08	1.00 E-145	Microscilla marina ATCC 23134	Bacterial polypeptide #10001.	ADS22995	Bacterial polypeptide #10001.	ADS60041	1.00 E-126	5.1.1.4	1005	334	0	333	73	
327, 328	proline racemase [Microscilla marina ATCC 23134] gi 123984081 gb EAY24454.1 proline racemase [Microscilla marina ATCC 23134]	1.24 E+08	1.00 E-147	Microscilla marina ATCC 23134	Bacterial polypeptide #10001.	ADS22995	Bacterial polypeptide #10001.	ADS60041	1.00 E-132	5.1.1.4	1002	333	0	333	74	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Geneseq Evaluate	Geneseq DNA Description	Geneseq DNA Accession Code	Geneseq Evaluate	Predicted EC Number	Query DNA Length	Query Protein Length	Query Protein Le ngth	Subject DNA Length	Subject Protein Length	% ID Protein	% ID DNA
329, 330	putative proline racemase [Myxococcus xanthus DK 1622]	1.09 E+08	1.00 E-101	Myxococcus xanthus DK 1622	M. xanthus protein sequence, seq id 9726.	ABM96 637	1.00 E-101	Prokaryotic essential gene #34740.	ACA2 3259	1.00 E-17	5.1.1.4	957	318	311	0	311	58	58
331, 332	AGR_L_3051p [Agrobacterium tumefaciens].	1589 1641	7.00 E-44	Agrobacterium tumefaciens	Bacterial polypeptide #19.	ADF07 948	5.00 E-21	Mycobacterium tuberculosis strain H37Rv genome SEQ ID NO 2.	AAI99 682	1.1	5.1.1.1	420	139	388	1167	388	61	68
333, 334	putative alanine racemase [Rhizobium leguminosarum bv. viciae 3841]	1.16 E+08	9.00 E-97	Rhizobium leguminosarum bv. viciae 3841	Bacterial polypeptide #19.	ADF07 948	6.00 E-29	Plant full length insert polynucleotide seqid 4980.	ADX5 1636	0.67	5.1.1.1	939	312	377	0	377	58	58
335, 336	alanine racemase [Moorella thermoacetica ATCC 39073]	8359 0990	5.00 E-78	Moorella thermoacetica ATCC 39073	Prokaryotic essential gene #34740.	ABU24 721	1.00 E-63	Enterobacter cloacae protein amino acid sequence - SEQ ID 5666.	AEH5 4260	0.05 3	5.1.1.1	1143	380	373	0	373	41	41

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Geneseq Protein Evaluate	Geneseq DNA Description	Geneseq DNA Accession Code	Geneseq Evaluate	Predicted EC Number	Query DNA Length	Query Protein Length	Subject DNA Length	Subject Protein Length	% ID Protein	% ID DNA
337, 338	alanine racemase [Moorella thermoacetica ATCC 39073]	8359 0990	3.00 E-77	Moorella thermoacetica ATCC 39073	Prokaryotic essential gene #34740.	ABU24 721	1.00 E-60	Bacillus licheniformis araA gene fragment amplifying PCR primer #1.	AAD2 9866	8.00 E-04	5.1.1.1	11 07	36 8	0	373	44	
339, 340	Alanine racemase [Azotobacter vinelandii AvOP] gij67087270 gb EAM06737.1 Alanine racemase [Azotobacter vinelandii AvOP]	6715 5469	1.00 E-115	Azotobacter vinelandii AvOP	Prokaryotic essential gene #34740.	ABU39 795	1.00 E-115	Prokaryotic essential gene #34740.	ACA2 3293	5.00 E-05	5.1.1.1	10 68	35 5	0	418	60	
341, 342	mandelate racemase/muconate lactonizing enzyme [Sphingopyxis alaskensis RB2256]	1.03 E+08	2.00 E-99	Sphingopyxis alaskensis RB2256	Klebsiella pneumoniae polypeptide seqid 7178.	ABO61 307	1.00 E-45	Geranylgeranyl pyrophosphate synthase polypeptide #7.	ADM9 8687	0.04 E-6	5.5.1.1	99 0	32 9	0	335	57	
343, 344	putative alanine racemase [Rhizobium leguminosarum bv. viciae 3841]	1.16 E+08	1.00 E-127	Rhizobium leguminosarum bv. viciae 3841	Pseudomonas aeruginosa polypeptide #3.	ABO84 274	8.00 E-42	Streptomyces cattleya NRRL 8057 orfY protein.	ADO5 1695	0.83 E-0	5.1.1.1	11 34	37 7	0	377	62	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Evaluate	Geneseq DNA Description	Geneseq DNA Accession Code	Evaluate	Predicted EC Number	Query DNA Length	Query Protein Length	Subject DNA Length	Subject Protein Length	% ID Protein	% ID DNA
345, 346	AGR_L_3051p [Agrobacterium tumefaciens].	1589 1641	2.00 E-79	Agrobacterium tumefaciens	Photorehabilitation luminescence protein sequence #59.	ABM69 114	1.00 E-27	Prokaryotic essential gene #34740.	ACA2 5617	2.2	5.1.1.1	77 7	25 8	116 7	388	57	65
347, 348	putative alanine racemase [Rhizobium leguminosarum bv. viciae 3841]	1.16 E+08	1.00 E-114	Rhizobium leguminosarum bv. viciae 3841	Prokaryotic essential gene #34740.	ABU41 398	1.00 E-36	Plant full length insert polynucleotide seqid 4980.	ADX5 1636	0.81	5.1.1.1	11 13	37 0	0 0	377	58	
349, 350	Mandelate racemase/muconate lactonizing enzyme; C-terminal domain protein [Methylobacterium sp. 4-46]	1.49 E+08	2.00 E-87	Methylobacterium sp. 4-46	Klebsiella pneumoniae polypeptide seqid 7178.	ABO61 307	1.00 E-63	Prokaryotic essential gene #34740.	ACA4 2641	0.72	5.1.2.2	99 3	33 0	0 0	326	51	
351, 352	Mandelate racemase/muconate lactonizing enzyme; C-terminal domain protein [Sphingomonas wittichii RW1]	1.49 E+08	3.00 E-93	Sphingomonas wittichii RW1	Klebsiella pneumoniae polypeptide seqid 7178.	ABO61 307	3.00 E-48	Klebsiella pneumoniae polypeptide seqid 7178.	ACH9 4858	5.00 E-05	5.5.1.1	10 05	33 4	0 0	354	52	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Evaluate	Geneseq DNA Description	Geneseq DNA Accession Code	Evaluate	Predicted EC Number	Query DNALength	Query ProteinLength	Subject DNALength	Subject ProteinLength	% ID Protein	% ID DNA
353, 354	putative proline racemase [Burkholderia xenovorans LB400]	9177 9222	1.00 E-38	Burkholderia xenovorans LB400	Prokaryotic essential gene #34740.	ABU21 813	3.00 E-39	Prokaryotic essential gene #34740.	ACA1 9650	3.00 E-22	5.1.1.4	38 1	12 6	945	318		
355, 356	alanine racemase [Lentisphaera araneosa HTCC2155]	1.49 E+08	2.00 E-67	Lentisphaera araneosa HTCC2155	Prokaryotic essential gene #34740.	ABU23 921	1.00 E-56	Prokaryotic essential gene #34740.	ACA3 8093	9.00 E-04	5.1.1.1	11 55	38 4	0	360	39	
357, 358	alanine racemase [Roseiflexus castenholzii DSM 13941] gi 118010932 gb E AV24951.1 alanine racemase [Roseiflexus castenholzii DSM 13941]	1.18 E+08	1.00 E-107	Roseiflexus castenholzii DSM 13941	Prokaryotic essential gene #34740.	ABU23 921	5.00 E-75	Prokaryotic essential gene #34740.	ACA4 0224	0.88	5.1.1.1	12 03	40 0	0	849	52	
359, 360	alanine racemase [Roseiflexus castenholzii DSM 13941] gi 118010932 gb E AV24951.1 alanine racemase [Roseiflexus castenholzii DSM 13941]	1.18 E+08	1.00 E-107	Roseiflexus castenholzii DSM 13941	Prokaryotic essential gene #34740.	ABU23 921	1.00 E-75	T. versicolor pyrF PCR primer SEQ ID 5.	AAF2 6441	0.88	5.1.1.1	12 03	40 0	0	849	52	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Genes eq Protein Accession Code	Geneseq DNA Description	Genes eq DNA Accession Code	Eval ue	Predi cted EC Number	Qu ery DN A Length	Qu ery Protein Length	Subj ect DN A Length	Subj ect Protein Length	% ID Protein	% ID DN A
361, 362	alanine racemase [Roseiflexus castenholzii DSM 13941] gij118010932 gb E AV24951.1 alanine racemase [Roseiflexus castenholzii DSM 13941]	1.18 E+08	1.00 E-107	Roseiflexus castenholzii DSM 13941	Prokaryotic essential gene #34740.	ABU23 921	T. versicolor pyrF PCR primer SEQ ID 5.	AAF2 6441	5.00 E-75	5.1.1.1	12 03	40 0	0 0	849	52	
363, 364	alanine racemase [Roseiflexus castenholzii DSM 13941] gij118010932 gb E AV24951.1 alanine racemase [Roseiflexus castenholzii DSM 13941]	1.18 E+08	1.00 E-107	Roseiflexus castenholzii DSM 13941	Prokaryotic essential gene #34740.	ABU23 921	T. versicolor pyrF PCR primer SEQ ID 5.	AAF2 6441	3.00 E-75	5.1.1.1	12 03	40 0	0 0	849	52	
365, 366	alanine racemase [Roseiflexus castenholzii DSM 13941] gij118010932 gb E AV24951.1 alanine racemase [Roseiflexus castenholzii DSM 13941]	1.18 E+08	1.00 E-107	Roseiflexus castenholzii DSM 13941	Prokaryotic essential gene #34740.	ABU23 921	T. versicolor pyrF PCR primer SEQ ID 5.	AAF2 6441	3.00 E-76	5.1.1.1	12 03	40 0	0 0	849	52	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Geneseq DNA Description	Geneseq DNA Accession Code	Genes Eval	Predicted EC Number	Query DNA Length	Query Protein Length	Subject DNA Length	Subject Protein Length	% ID Protein	% ID DNA
367, 368	alanine racemase [Pseudomonas mendocina ymp]	1.46 E+08	1.00 E-168	Pseudomonas mendocina ymp	Prokaryotic essential gene #34740.	ABU41 398	Prokaryotic essential gene #34740.	ACA4 3665	7.00 E-32	5.1.1.1	10 74	35 7	0	362	82	
369, 370	diaminopimelate epimerase [Nostoc sp. PCC 7120].	1723 2333	3.00 E-78	Nostoc sp. PCC 7120	Bacterial polypeptide #10001.	ADS29 848	S. lavendulae gene mutagenic PCR primer #2.	ADE1 0236	9.00 E-79	5.1.1.7	86 1	28 6	141 3	285		
371, 372	diaminopimelate epimerase [Syntrophus aciditrophicus SB]	8585 7916	2.00 E-76	Syntrophus aciditrophicus SB	Bacterial polypeptide #10001.	ADN17 489	Bacterial polypeptide #10001.	ADT4 6187	5.00 E-60	5.1.1.7	81 3	27 0	0	277	57	
373, 374	diaminopimelate epimerase [Nostoc sp. PCC 7120].	1723 2333	2.00 E-71	Nostoc sp. PCC 7120	Bacterial polypeptide #10001.	ADS29 848	Environmental isolate hydrolyase, SEQ ID NO:44.	AEH4 7413	7.00 E-72	5.1.1.7	87 3	29 0	900	285		
375, 376	Diaminopimelate epimerase [Clostridium thermocellum ATCC 27405]	1.26 E+08	5.00 E-33	Clostridium thermocellum ATCC 27405	Bacterial polypeptide #10001.	ADN27 169	Bacterial polypeptide #10001.	ADT4 2245	2.00 E-31	5.1.1.7	82 8	27 5	0	280	29	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Genes eq Protein Accession Code	Geneseq DNA Description	Genes eq DNA Accession Code	Genes Evaluate	Predicted EC Number	Query DNA Length	Query Protein Length	Subject DNA Length	Subject Protein Length	% ID Protein	% ID DNA
377, 378	UDP-N-acetylglucosamine 2-epimerase [Chloroflexus aggregans DSM 9485] gij117997290 gb EAV11478.1 UDP-N-	1.18 E+08	2.00 E-97	Chloroflexus aggregans DSM 9485	Prokaryotic essential gene #34740.	ABU24 769	Prokaryotic essential gene #34740.	ACA2 7041	6.00 E-80	5.1.3.14	1200	399	0	386	50	
379, 380	NmrA family protein [Burkholderia vietnamiensis G4]	1.34 E+08	2.00 E-88	Burkholderia vietnamiensis G4	Bacterial polypeptide #10001.	ADS22 779	Bacterial polypeptide #10001.	ADS5 9825	2.00 E-84	1.6.5.3	879	292	0	287	55	
381, 382	hypothetical protein RL1205 [Rhizobium leguminosarum bv. viciae 3841]	1.16 E+08	2.00 E-87	Rhizobium leguminosarum bv. viciae 3841	Bacterial polypeptide #10001.	ADS22 779	Bacterial polypeptide #10001.	ADS5 9825	1.00 E-08	1.6.5.3	879	292	0	289	55	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Genes eq Protein Accession Code	Evaluate	Geneseq DNA Description	Genes eq DNA Accession Code	Evaluate	Predicted EC Number	Query DN A Length	Query Protein Length	Subj ect DN A Length	Subj ect Protein Length	% ID Protein	% ID DN A
383, 384	similar to chloromuconate cycloisomerase [Gloeobacter violaceus PCC 7421]	3752 2668	1.00 E-93	Gloeobacter violaceus PCC 7421	L. pneumophila protein SEQ ID NO 3367.	AEB37 282	4.00 E-48	Novel canine microarray-related DNA sequence SeqID100 21.	ADQ5 3782	3.1	5.1.2.2	10 71	35 6	0	356	50	
1, 2	hypothetical protein [Aquifex aeolicus].	1560 5814	1.00 E-128	Aquifex aeolicus	Prokaryotic essential gene #34740.	ABU25 646	3.00 E-47	Geneseq DNA Description Drosophila melanogaster polypeptide SEQ ID NO 24465.	ABLO 3829	0.48	2.7.3.	68 7	22 8	687	228	10 0	10 0

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Genes eq Protein Accession Code	Evaluate	Geneseq DNA Description	Genes eq DNA Accession Code	Evaluate	Predicted EC Number	Query DNA Length	Query Protein Length	Subject DNA Length	Subject Protein Length	% ID Protein	% ID DNA
3, 4	alanine racemase [Aquifex aeolicus].	1560 6873	0	Aquifex aeolicus	Aquifex pyrophilus heat resistant alanine racemase encoding DNA.	ABB06 296	1.00 E- 147	Aquifex pyrophilus heat resistant alanine racemase encoding DNA.	ABL4 9607	2.00 E- 07	5.1.1. 1	10 17	33 8	0 0	338 0	10 0	
5, 6	hypothetical protein SAV6126 [Streptomyces avermitilis MA- 4680]	2983 2668	1.00 E- 101	Streptom yces avermitili s MA- 4680	Prokaryotic essential gene #34740.	ABU19 499	5.00 E- 37	Human immune/h ematopoi etic antigen genomic sequence SEQ ID NO:41436	AAK8 0650	0.03 2	2.7.3.	72 0	23 9	0 0	239 0	80 0	
7, 8	hypothetical protein SAV4292 [Streptomyces avermitilis MA- 4680]	2983 0835	0	Streptom yces avermitili s MA- 4680	Propioniba cterium acnes predicted ORF- encoded polypeptide #300.	ABM54 358	2.00 E- 53	EST clone EP219.	AAV8 8076	0.04 8		10 32	34 3	0 0	343 0	89 0	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Genes eq Protein Accession Code	Geneseq DNA Description	Genes eq DNA Accession Code	Eval ue	Predicted EC Number	Qu ery DN A Le ngt h	Qu ery Protein Le ngt h	Subj ect DN A Le ngt h	Subj ect Protein Le ngt h	% ID Protein	% ID DN A
9, 10	alanine racemase [Streptomyces coelicolor A3(2)]	2122 3124	1.00 E-166	Streptomyces coelicolor A3(2)	Prokaryotic essential gene #34740.	ABU34 223	Pseudomonas aeruginosa polypeptide #3.	ABD0 8732	6.00 E-90	5.1.1.1	11 49	38 2	0	391	78	
11, 12	alanine racemase [Rhodospseudomonas palustris HaA2]	8674 8627	1.00 E-106	Rhodospseudomonas palustris HaA2	Pseudomonas aeruginosa polypeptide #3.	ABO84 274	Pseudomonas aeruginosa polypeptide #3.	ABD0 6378	4.00 E-50	5.1.1.1	11 22	37 3	0	388	58	
13, 14	alanine racemase [Azoarcus sp. EbN1]	5647 7426	1.00 E-128	Azoarcus sp. EbN1	Prokaryotic essential gene #34740.	ABU41 398	Pseudomonas aeruginosa polypeptide #3.	ABD0 8732	1.00 E-110	5.1.1.1	10 65	35 4	0	353	64	
15, 16	Aspartate racemase [Marinobacter algicola DG893]	1.49 E+08	4.00 E-93	Marinobacter algicola DG893	Klebsiella pneumoniae polypeptide seqid 7178.	ABO65 503	Klebsiella pneumoniae polypeptide seqid 7178.	ACH9 9054	1.00 E-71	5.1.1.1	70 2	23 3	0	231	74	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Genes eq Protein Accession Code	Eval ue	Geneseq DNA Description	Genes eq DNA Accession Code	Eval ue	Predi cted EC Number	Qu ery DN A Le ngt h	Qu ery Pro tein Le ngt h	Subj ect DN A Le ngt h	Subj ect Prot ein Len gth	% ID Pro tein	% ID DN A
17, 18	glutamate racemase [Cytophaga hutchinsonii ATCC 33406]	1.11 E+08	1.00 E-106	Cytophaga hutchinsonii ATCC 33406	Prokaryotic essential gene #34740.	ABU25 174	2.00 E-44	Human phosphodiesterase 4D amino acid sequence N1 SEQ ID NO:7.	AEB8 5185	0.00 9	5.1.1.3	80 7	26 8	0	269	72	
19, 20	glutamate racemase [Geobacter sulfurreducens PCA]	3999 8014	1.00 E-74	Geobacter sulfurreducens PCA	M. xanthus protein sequence, seq id 9726.	ABM90 755	2.00 E-67	M. xanthus protein sequence, seq id 9726.	ACL6 4794	0.03 7	5.1.1.3	82 2	27 3	0	272	53	
21, 22	Putative decarboxylase [Bordetella parapertussis 12822]	3359 6748	5.00 E-40	Bordetella parapertussis 12822	Thermococcus kodakaraensis KOD1 protein sequence SeqID4.	ADN46 910	5.00 E-27	Bacterial polypeptide #10001.	ADS6 1806	1.00 E-04	5.2.1.1	76 8	25 5	0	265	41	
23, 24	alanine racemase [Enterobacter sp. 638]	1.46 E+08	0	Enterobacter sp. 638	Enterobacter cloacae protein amino acid sequence - SEQ ID 5666.	AEH63 094	0	Enterobacter cloacae protein amino acid sequence - SEQ ID 5666.	AEH5 5699	0 1	5.1.1.1	10 80	35 9	0	359	95	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Geneseq DNA Description	Geneseq DNA Accession Code	Eval	Predicted EC Number	Query DNALength	Query ProteinLength	Query DNALength	Subject DNALength	Subject ProteinLength	% ID Protein	% ID DNA
25, 26	alanine racemase [Enterobacter sp. 638]	1.46 E+08	0	Enterobacter sp. 638	Enterobacter cloacae protein amino acid sequence - SEQ ID 5666.	AEH63094	Enterobacter cloacae protein amino acid sequence - SEQ ID 5666.	AEH55699	0	5.1.1.1	1080	359	0	0	359	95	
27, 28	hypothetical protein SAR11_0361 [Candidatus Pelagibacter ubique HTCC1062]	71083067	1.00 E-121	Candidatus Pelagibacter ubique HTCC1062	Prokaryotic essential gene #34740.	ABU24290	Prokaryotic essential gene #34740.	ACA27845	0.002	2.7.3.	651	216	0	0	216	100	
29, 30	putative proline racemase [Brucella ovis ATCC 25840]	1.49 E+08	1.00 E-130	Brucella ovis ATCC 25840	Pseudomonas aeruginosa polypeptide #3.	ABO82134	Pseudomonas aeruginosa polypeptide #3.	ABD15676	1.00 E-08	5.1.1.4	1032	343	0	0	342	64	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Genes eq Protein Accession Code	Geneseq DNA Description	Genes eq DNA Accession Code	Eval ue	Predicted EC Number	Qu ery DN A Le ngth	Qu ery Protein Le ngth	Subj ect DN A Le ngth	Subj ect Protein Len gth	% ID Pro tein	% ID DN A
31, 32	proline racemase, putative [Stappia aggregata IAM 12614] gij118435940 gb E AV42584.1 proline racemase, putative [Stappia aggregata IAM 12614]	1.19 E+08	1.00 E-109	Stappia aggregata IAM 12614	Acinetobacter baumannii protein #19.	ADA35 228	Pseudomonas aeruginosa polypeptide #3.	ABD1 5666	1.00 E-105	9.00 E-164	10 11	33 6	0	310	58	
33, 34	alanine racemase [Mesorhizobium sp. BNC1] putative alanine racemase [Aurantimonas sp. SI85-9A1] gij9033811 gb E AS51762.1 putative alanine racemase [Aurantimonas sp. SI85-9A1]	1.11 E+08	1.00 E-124	Mesorhizobium sp. BNC1	Prokaryotic essential gene #34740.	ABU38 829	Human cancer associated cDNA SEQ ID NO 9.	AEH9 8530	3.00 E-46	0.01 E-31	11 16	37 1	0	385	61	
35, 36	putative alanine racemase [Aurantimonas sp. SI85-9A1] gij9033811 gb E AS51762.1 putative alanine racemase [Aurantimonas sp. SI85-9A1]	9041 8439	6.00 E-49	Aurantimonas sp. SI85-9A1	Achromobacter xylosoxidans DTA SEQ ID NO 6.	AEH19 277	N. meningitidis partial DNA sequence gnm_640 SEQ ID NO:640.	AAA8 1486	5.00 E-47	3.2 E-5.1.1.	11 01	36 6	0	368	33	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Geneseq DNA Description	Geneseq DNA Accession Code	Eval	Predicted EC Number	Query DN A Length	Query Protein Length	Subject DN A Length	Subject Protein Length	% ID Protein	% ID DN A
37, 38	Alanine racemase [Magnetospirillum gryphiswaldense MSR-1]	1.45 E+08	1.00 E-38	Magnetospirillum gryphiswaldense MSR-1	Aquifex pyrophilus heat resistant alanine racemase encoding DNA.	ABB06 296	Pseudomonas aeruginosa polypeptide #3.	ABD1 2586	2.00 E-20	5.1.1.1	489	162	0	369	51	
39, 40	putative proline racemase [Burkholderia xenovorans LB400]	9177 9222	2.00 E-36	Burkholderia xenovorans LB400	Prokaryotic essential gene #34740.	ABU21 813	Prokaryotic essential gene #34740.	ACA2 6332	5.00 E-37	5.1.1.4	372	123	960	318		
41, 42	alanine racemase [Pseudomonas putida F1]	1.49 E+08	0	Pseudomonas putida F1	T. maritima D-alanine-D-alanine ligase.	AED11 803	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	0	5.1.1.1	1230	409	0	409	97	
43, 44	alanine racemase [Pseudomonas putida F1]	1.49 E+08	0	Pseudomonas putida F1	T. maritima D-alanine-D-alanine ligase.	AED11 803	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	0	5.1.1.1	1230	409	0	409	96	
45, 46	alanine racemase [Pseudomonas putida F1]	1.49 E+08	0	Pseudomonas putida F1	T. maritima D-alanine-D-alanine ligase.	AED11 803	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	0	5.1.1.1	1230	409	0	409	97	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Genes eq Protein Accession Code	Geneseq DNA Description	Genes eq DNA Accession Code	Eval ue	Pred icted EC Num ber	Qu ery DN A Le ngth	Qu ery Pro tein Le ngth	Subj ect DN A Le ngth	Subj ect Prot ein Len gth	% ID Pro tein	% ID DN A
47, 48	alanine racemase [Pseudomonas putida F1]	1.49 E+08	0	Pseudomonas putida F1	T. maritima D-alanine-D-alanine ligase.	AED11 803	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	0	5.1.1.1	12 30	40 9	0	409	97	
49, 50	alanine racemase [Pseudomonas putida F1]	1.49 E+08	0	Pseudomonas putida F1	T. maritima D-alanine-D-alanine ligase.	AED11 803	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	0	5.1.1.1	12 30	40 9	0	409	96	
51, 52	alanine racemase [Pseudomonas putida GB-1] gij126314851 gb E AZ66019.1] alanine racemase [Pseudomonas putida GB-1]	1.26 E+08	0	Pseudomonas putida GB-1	T. maritima D-alanine-D-alanine ligase.	AED11 803	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9537	1.00 E-120	5.1.1.1	12 30	40 9	0	409	77	
53, 54	alanine racemase [Pseudomonas putida GB-1] gij126314851 gb E AZ66019.1] alanine racemase [Pseudomonas putida GB-1]	1.26 E+08	1.00 E-180	Pseudomonas putida GB-1	T. maritima D-alanine-D-alanine ligase.	AED11 803	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9537	1.00 E-124	5.1.1.1	12 30	40 9	0	409	77	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Genes eq Protein Accession Code	Geneseq DNA Description	Genes eq DNA Accession Code	Genes Evaluate	Predicted EC Number	Query DNA Length	Query Protein Length	Subject DNA Length	Subject Protein Length	% ID Protein	% ID DNA
55, 56	alanine racemase [Pseudomonas putida GB-1] gi 126314851 gb E AZ66019.1	1.26 E+08	1.00 E-180	Pseudomonas putida GB-1	T. maritima D-alanine-D-alanine ligase.	AED11803	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99537	1.00 E-124	5.1.1.1	1230	409	0	409	77	
57, 58	alanine racemase [Pseudomonas putida F1]	1.49 E+08	0	Pseudomonas putida F1	T. maritima D-alanine-D-alanine ligase.	AED11804	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	0	5.1.1.1	1230	409	0	409	96	
59, 60	alanine racemase [Pseudomonas putida F1]	1.49 E+08	0	Pseudomonas putida F1	T. maritima D-alanine-D-alanine ligase.	AED11804	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	0	5.1.1.1	1230	409	0	409	95	
61, 62	alanine racemase [Fusobacterium nucleatum subsp. polymorphum ATCC 10953]	1.48 E+08	7.00 E-19	Fusobacterium nucleatum subsp. polymorphum ATCC 10953	Aquifex pyrophilus heat resistant alanine racemase encoding DNA.	ABB06296	Human prostate expression marker cDNA 6604.	ABV52203	1.00 E-12	5.1.1.1	1614	0	354	19		

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Geneseq DNA Description	Geneseq DNA Accession Code	Eval	Predicted EC Number	Query DNA Length	Query Protein Length	Subject DNA Length	Subject Protein Length	% ID Protein	% ID DNA
63, 64	alanine racemase [Pseudomonas putida F1]	1.49 E+08	0	Pseudomonas putida F1	T. maritima D-alanine-D-alanine ligase.	AED11803	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99537	0	5.1.1.1	1230	409	0	409	98	
65, 66	alanine racemase [Pseudomonas putida F1]	1.49 E+08	0	Pseudomonas putida F1	T. maritima D-alanine-D-alanine ligase.	AED11803	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99537	0	5.1.1.1	1230	409	0	409	97	
67, 68	alanine racemase [Pseudomonas putida GB-1]	1.26 E+08	0	Pseudomonas putida GB-1	T. maritima D-alanine-D-alanine ligase.	AED11803	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	0	5.1.1.1	1230	409	0	409	90	
69, 70	alanine racemase [Pseudomonas putida GB-1]	1.26 E+08	0	Pseudomonas putida GB-1	T. maritima D-alanine-D-alanine ligase.	AED11803	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99537	0	5.1.1.1	1230	409	0	409	95	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Genes eq Protein Accession Code	Geneseq DNA Description	Genes eq DNA Accession Code	Eval ue	Predicted EC Number	Qu ery DN A Le ngth	Qu ery Protein Le ngth	Subj ect DN A Le ngth	Subj ect Protein Le ngth	% ID Protein	% ID DN A
71, 72	alanine racemase [Pseudomonas putida GB-1] gi 126314851 gb E AZ66019.1] alanine racemase [Pseudomonas putida GB-1]	1.26 E+08	0	Pseudomonas putida GB-1	T. maritima D-alanine-D-alanine ligase.	AED11 803	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9537	0	5.1.1.1	12 30	40 9	0	409	96	
73, 74	alanine racemase [Pseudomonas putida GB-1] gi 126314851 gb E AZ66019.1] alanine racemase [Pseudomonas putida GB-1]	1.26 E+08	0	Pseudomonas putida GB-1	T. maritima D-alanine-D-alanine ligase.	AED11 803	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9537	0	5.1.1.1	12 30	40 9	0	409	95	
75, 76	alanine racemase [Pseudomonas putida GB-1] gi 126314851 gb E AZ66019.1] alanine racemase [Pseudomonas putida GB-1]	1.26 E+08	0	Pseudomonas putida GB-1	T. maritima D-alanine-D-alanine ligase.	AED11 803	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9537	0	5.1.1.1	12 30	40 9	0	409	95	
77, 78	alanine racemase [Pseudomonas putida F1]	1.49 E+08	0	Pseudomonas putida F1	T. maritima D-alanine-D-alanine ligase.	AED11 803	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	0	5.1.1.1	12 30	40 9	0	409	95	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Geneseq DNA Description	Geneseq DNA Accession Code	Eval	Predicted EC Number	Query DNA Length	Query Protein Length	Subject DNA Length	Subject Protein Length	% ID Protein	% ID DNA
79, 80	alanine racemase [Pseudomonas putida F1]	1.49 E+08	0	Pseudomonas putida F1	T. maritima D-alanine-D-alanine ligase.	AED11 804	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	0	5.1.1.1	1230	409	0	409	95	
81, 82	alanine racemase [Pseudomonas putida F1]	1.49 E+08	0	Pseudomonas putida F1	T. maritima D-alanine-D-alanine ligase.	AED11 804	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	0	5.1.1.1	1230	409	0	409	96	
83, 84	alanine racemase [Pseudomonas putida F1]	1.49 E+08	0	Pseudomonas putida F1	T. maritima D-alanine-D-alanine ligase.	AED11 804	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	0	5.1.1.1	1230	409	0	409	94	
85, 86	alanine racemase [Pseudomonas putida F1]	1.49 E+08	0	Pseudomonas putida F1	T. maritima D-alanine-D-alanine ligase.	AED11 803	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9537	0	5.1.1.1	1230	409	0	409	96	
87, 88	alanine racemase [Pseudomonas putida F1]	1.49 E+08	0	Pseudomonas putida F1	T. maritima D-alanine-D-alanine ligase.	AED11 803	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	0	5.1.1.1	1230	409	0	409	96	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Genes eq Protein Accession Code	Geneseq DNA Description	Genes eq DNA Accession Code	Eval ue	Predicted EC Number	Query DN A Length	Query Protein Length	Subject DN A Length	Subject Protein Length	% ID Protein	% ID DN A
89, 90	alanine racemase [Pseudomonas putida KT2440]	2699 0430	0	Pseudomonas putida KT2440	T. maritima D-alanine-D-alanine ligase.	AED11 803	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9537	0	5.1.1.1	12 30	40 9	0	409	96	96
91, 92	alanine racemase [Pseudomonas putida F1]	1.49 E+08	0	Pseudomonas putida F1	T. maritima D-alanine-D-alanine ligase.	AED11 803	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	0	5.1.1.1	12 30	40 9	0	409	96	96
93, 94	alanine racemase [Pseudomonas putida F1] alanine racemase [Pseudomonas putida GB-1] gi 126314851 gb E AZ66019.1	1.49 E+08	0	Pseudomonas putida F1	T. maritima D-alanine-D-alanine ligase.	AED11 804	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	0	5.1.1.1	12 30	40 9	0	409	96	96
95, 96	alanine racemase [Pseudomonas putida GB-1]	1.26 E+08	0	Pseudomonas putida GB-1	T. maritima D-alanine-D-alanine ligase.	AED11 803	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9537	0	5.1.1.1	12 30	40 9	0	409	95	95

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Genes eq Protein Accession Code	Eval ue	Geneseq DNA Description	Genes eq DNA Accession Code	Eval ue	Predicted EC Number	Qu ery DN A Le ngth	Qu ery Protein Le ngth	Subj ect DN A Len gth	Subj ect Protein Len gth	% ID Protein	% ID DN A
97, 98	alanine racemase [Pseudomonas putida GB-1] gij126314851 gb E AZ66019.1 alanine racemase [Pseudomonas putida GB-1]	1.26 E+08	0	Pseudomonas putida GB-1	T. maritima D-alanine-D-alanine ligase.	AED11803	0	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99537	0	5.1.1.1	1230	409	0	409	96	
99, 100	alanine racemase [Pseudomonas putida F1]	1.49 E+08	0	Pseudomonas putida F1	T. maritima D-alanine-D-alanine ligase.	AED11803	0	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99537	0	5.1.1.1	1230	409	0	409	97	
101, 102	alanine racemase 2, catabolic [Escherichia coli O157:H7 EDL933].	15801412	2.00 E-58	Escherichia coli O157:H7 EDL933	Prokaryotic essential gene #34740.	ABU45089	2.00 E-58	Enterobacter cloacae protein amino acid sequence - SEQ ID 5666.	AEH53103	5.00 E-27	5.1.1.1	384	127	1071	356	86	76
103, 104	aspartate racemase [Pyrococcus abyssii].	14521575	8.00 E-29	Pyrococcus abyssii	Thermococcus kodakaraensis KOD1 protein sequence SeqID4.	ADN46207	2.00 E-25	Prokaryotic essential gene #34740.	ACA27539	0.13	5.1.1.13	714	237	687	228	32	48

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Evaluate	Geneseq DNA Description	Geneseq DNA Accession Code	Evaluate	Predicted EC Number	Query DN A Length	Query Protein Length	Subject DN A Length	Subject Protein Length	% ID Protein	% ID DNA
105, 106	proline racemase [Microscilla marina ATCC 23134] gij123984081 gb EAY24454.1 proline racemase [Microscilla marina ATCC 23134]	1.24 E+08	1.00 E-150	Microscilla marina ATCC 23134	Bacterial polypeptide #10001.	ADS22 995	1.00 E-128	Bovine ABCG2 related PCR primer #31.	AEN6 9487	2.9	5.1.1.4	10023	333	0	333	75	
107, 108	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter brevis mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	1.00 E-131	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	6.00 E-11	5.1.1.1	1227	408	0	408	96	
109, 110	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter brevis mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	1.00 E-129	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	6.00 E-05	5.1.1.1	1227	408	0	408	95	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Evaluate	Geneseq DNA Description	Geneseq DNA Accession Code	Evaluate	Predicted EC Number	Query DN A Length	Query Protein Length	Subject DN A Length	Subject Protein Length	% ID Protein	% ID DN A
111, 112	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-131	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	2.00 E-11	5.1.1.1	1227	408	0	408	99	
113, 114	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-131	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	2.00 E-14	5.1.1.1	1224	407	0	408	92	
115, 116	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-132	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	2.00 E-08	5.1.1.1	1227	408	0	408	92	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Geneseq DNA Description	Geneseq DNA Accession Code	Eval	Predicted EC Number	Query DN A Length	Query Protein Length	Subject DN A Length	Subject Protein Length	% ID Protein	% ID DN A
117, 118	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	6.00 E-11	5.1.1.1	12 27	40 8	0	408	92	
119, 120	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	1.00 E-12	5.1.1.1	12 27	40 8	0	408	95	
121, 122	alanine racemase [Pseudomonas putida F1]	1.49 E+08	1.00 E-136	Pseudomonas putida F1	T. maritima D-alanine-D-alanine ligase.	AED11 803	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	2.00 E-26	5.1.1.1	10 86	36 1	122 7	409		
123, 124	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	6.00 E-11	5.1.1.1	10 86	36 1	0	408	88	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Evaluate	Geneseq DNA Description	Geneseq DNA Accession Code	Evaluate	Predicted EC Number	Query DN A Length	Query Protein Length	Subject DN A Length	Subject Protein Length	% ID Protein	% ID DN A
125, 126	alanine racemase [Pseudomonas putida GB-1] gi 126314851 gb E AZ66019.1 alanine racemase [Pseudomonas putida GB-1]	1.26 E+08	1.00 E-155	Pseudomonas putida GB-1	Empedobacter brevis mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-159	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	4.00 E-95	5.1.1.1	1086	361	1227	386		
127, 128	glutamate racemase [Bradyrhizobium japonicum USDA 110]	27380813	3.00 E-26	Bradyrhizobium japonicum USDA 110	Photorhabdus luminescens protein sequence #59.	ABM68637	7.00 E-18	Prokaryotic essential gene #34740.	ACA37866	2.2	5.1.1.3	261	86	0	265	63	
129, 130	proline racemase, putative [Stappia aggregata IAM 12614] gi 118435940 gb E AV42584.1 proline racemase, putative [Stappia aggregata IAM 12614]	1.19 E+08	1.00 E-103	Stappia aggregata IAM 12614	Prokaryotic essential gene #34740.	ABU21813	1.00 E-101	Prokaryotic essential gene #34740.	ACA26332	5.00 E-11	5.1.1.4	957	318	0	310	56	
131, 132	putative proline racemase [Myxococcus xanthus DK 1622]	1.09 E+08	5.00 E-75	Myxococcus xanthus DK 1622	M. xanthus protein sequence, seq id 9726.	ABM96637	1.00 E-75	Prokaryotic essential gene #34740.	ACA26332	1.00 E-18	5.1.1.4	1074	357	960	311		

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Genes eq Protein Accession Code	Evaluate	Geneseq DNA Description	Genes eq DNA Accession Code	Evaluate	Predicted EC Number	Query DN A Length	Query Protein Length	Subject DN A Length	Subject Protein Length	% ID Protein	% ID DNA
133, 134	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthetase enzyme SEQ ID NO:3.	AED10 581	1.00 E-130	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	6.00 E-11	5.1.1.1	12 27	40 8	0	408	99	
135, 136	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthetase enzyme SEQ ID NO:3.	AED10 581	1.00 E-129	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	6.00 E-05	5.1.1.1	12 27	40 8	0	408	95	
137, 138	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthetase enzyme SEQ ID NO:3.	AED10 581	1.00 E-128	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	0.00 E-4	5.1.1.1	12 21	40 6	0	408	95	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Evaluate	Geneseq DNA Description	Geneseq DNA Accession Code	Evaluate	Predicted EC Number	Query DNA Length	Query Protein Length	Subject DNA Length	Subject Protein Length	% ID Protein	% ID DNA
139, 140	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-131	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	6.00 E-11	5.1.1.1	1224	407	0	408	99	
141, 142	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-130	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	1.00 E-15	5.1.1.1	1227	408	0	408	89	
143, 144	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-132	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	6.00 E-11	5.1.1.1	1227	408	0	408	100	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Gene Eval	Geneseq DNA Description	Geneseq DNA Accession Code	Eval	Predicted EC Number	Query DNA Length	Query Protein Length	Subject DNA Length	Subject Protein Length	% ID Protein	% ID DNA
145, 146	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter brevis mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-131	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	2.00 E-11	5.1.1.1	1224	407	0	408	99	
147, 148	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter brevis mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-131	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	3.00 E-19	5.1.1.1	1224	407	0	408	97	
149, 150	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter brevis mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-132	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	6.00 E-11	5.1.1.1	1227	408	0	408	93	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Evaluate	Geneseq DNA Description	Geneseq DNA Accession Code	Evaluate	Predicted EC Number	Query DNA Length	Query Protein Length	Subject DNA Length	Subject Protein Length	% ID Protein	% ID DNA
151, 152	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	1.00 E-131	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	3.00 E-19	5.1.1.1	12 27	40 8	0	408	92	
153, 154	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	1.00 E-131	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	3.00 E-19	5.1.1.1	12 24	40 7	0	408	97	
155, 156	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	1.00 E-133	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	2.00 E-14	5.1.1.1	12 27	40 8	0	408	96	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Genes eq Protein Accession Code	Evaluate	Geneseq DNA Description	Genes eq DNA Accession Code	Evaluate	Predicted EC Number	Query DN A Length	Query Protein Length	Subject DN A Length	Subject Protein Length	% ID Protein	% ID DN A
157, 158	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	1.00 E-132	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	6.00 E-11	5.1.1.1	12 27	40 8	0	408	92	
159, 160	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	1.00 E-131	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	6.00 E-11	5.1.1.1	12 27	40 8	0	408	92	
161, 162	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	1.00 E-131	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	3.00 E-19	5.1.1.1	12 27	40 8	0	408	97	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Eval ue	Geneseq DNA Description	Geneseq DNA Accession Code	Eval ue	Predicted EC Number	Qu ery DN A Le ngt h	Qu ery Protein Le ngt h	Subj ect DN A Le ngt h	Subj ect Protein Len gth	% ID Protein	% ID DN A
163, 164	alanine racemase [Pseudomonas putida F1]	1.49 E+08	1.00 E-153	Pseudomonas putida F1	Empedobacter brevis mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	0	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	1.00 E-64	5.1.1.1	1086	361	1227	386		
165, 166	alanine racemase [Pseudomonas putida F1]	1.49 E+08	1.00 E-153	Pseudomonas putida F1	Empedobacter brevis mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	0	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	2.00 E-91	5.1.1.1	1083	360	1227	386		
167, 168	proline racemase [Flavobacteriales bacterium HTCC2170] gij88708932 gb E AR01166.1 proline racemase [Flavobacteriales bacterium HTCC2170]	88712394	1.00 E-69	Flavobacteriales bacterium HTCC2170	Bacterial polypeptide #10001.	ADS22995	2.00 E-57	Bacterial polypeptide #10001.	ADS60041	0.00 E-7	5.1.1.4	657	218	0	335		58

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Eval ue	Geneseq DNA Description	Geneseq DNA Accession Code	Eval ue	Predicted EC Number	Qu ery DN A Length	Qu ery Protein Length	Subj ect DN A Length	Subj ect Protein Length	% ID Protein	% ID DN A
169, 170	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthetase enzyme SEQ ID NO:3.	AED10 581	1.00 E-131	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	6.00 E-11	5.1.1.1	12 27	40 8	0	408	97	
171, 172	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthetase enzyme SEQ ID NO:3.	AED10 581	1.00 E-131	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	6.00 E-11	5.1.1.1	12 27	40 8	0	408	96	
173, 174	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthetase enzyme SEQ ID NO:3.	AED10 581	1.00 E-129	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	6.00 E-05	5.1.1.1	12 24	40 7	0	408	95	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Gene Eval	Geneseq DNA Description	Geneseq DNA Accession Code	Gene Eval	Predicted EC Number	Query DN Length	Query Protein Length	Subject DN Length	Subject Protein Length	% ID Protein	% ID DNA
175, 176	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-130	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	2.00 E-14	5.1.1.1	1227	408	0	408	89	
177, 178	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-132	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	6.00 E-11	5.1.1.1	1227	408	0	408	99	
179, 180	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-132	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	3.00 E-10	5.1.1.1	1227	408	0	408	99	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Genes eq Protein Accession Code	Geneseq DNA Description	Genes eq DNA Accession Code	Eval ue	Predi cted EC Number	Qu ery DN A Length	Qu ery Protein Length	Subj ect DN A Length	Subj ect Protein Length	% ID Protein	% ID DN A
181, 182	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	1.00 E-130	5.1.1.1	12 27	40 8	0	408	99	
183, 184	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	1.00 E-132	5.1.1.1	12 27	40 8	0	408	10 0	
185, 186	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	1.00 E-129	5.1.1.1	12 27	40 8	0	408	89	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Genes eq Protein Accession Code	Evaluate	Geneseq DNA Description	Genes eq DNA Accession Code	Evaluate	Predicted EC Number	Query DN A Length	Query Protein Length	Subject DN A Length	Subject Protein Length	% ID Protein	% ID DNA
187, 188	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	1.00 E-131	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	6.00 E-11	5.1.1.1	12 27	40 8	0	408	99	
189, 190	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	1.00 E-131	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	6.00 E-11	5.1.1.1	12 27	40 8	0	408	99	
191, 192	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	1.00 E-131	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	6.00 E-11	5.1.1.1	12 27	40 8	0	408	91	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Gene Eval	Geneseq DNA Description	Gene Accession Code	Gene Eval	Predicted EC Number	Query DNA Length	Query Protein Length	Subject DNA Length	Subject Protein Length	% ID Protein	% ID DNA
193, 194	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	1.00 E-131	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	6.00 E-11	5.1.1.1	12 27	40 8	0	408	92	
195, 196	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	1.00 E-131	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9537	6.00 E-11	5.1.1.1	12 27	40 8	0	408	92	
197, 198	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	1.00 E-131	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	2.00 E-14	5.1.1.1	12 27	40 8	0	408	91	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Genes eq Protein Accession Code	Geneseq DNA Description	Genes eq DNA Accession Code	Eval ue	Predi cted EC Number	Qu ery DN A Length	Qu ery Protein Length	Subj ect DN A Length	Subj ect Protein Length	% ID Protein	% ID DN A
199, 200	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	1.00 E-15	5.1.1.1	12 27	40 8	0	408	90	
201, 202	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	1.00 E-129	5.1.1.1	12 27	40 8	0	408	89	
203, 204	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	4.00 E-18	5.1.1.1	12 27	40 8	0	408	92	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Evaluate	Geneseq DNA Description	Geneseq DNA Accession Code	Evaluate	Predicted EC Number	Query DN A Length	Query Protein Length	Subject DN A Length	Subject Protein Length	% ID Protein	% ID DN A
205, 206	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	1.00 E-130	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	1.00 E-15	5.1.1.1	12 27	40 8	0	408	89	
207, 208	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	1.00 E-133	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	3.00 E-16	5.1.1.1	12 24	40 7	0	408	92	
209, 210	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	1.00 E-131	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	1.00 E-12	5.1.1.1	12 27	40 8	0	408	91	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Evaluate	Geneseq DNA Description	Geneseq DNA Accession Code	Evaluate	Predicted EC Number	Query DNA Length	Query Protein Length	Subject DNA Length	Subject Protein Length	% ID Protein	% ID DNA
211, 212	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	1.00 E-131	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	2.00 E-14	5.1.1.1	12 24	40 7	0	408	89	
213, 214	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	1.00 E-133	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	1.00 E-15	5.1.1.1	12 24	40 7	0	408	91	
215, 216	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	1.00 E-130	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	6.00 E-11	5.1.1.1	12 27	40 8	0	408	96	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Evaluate	Geneseq DNA Description	Geneseq DNA Accession Code	Evaluate	Predicted EC Number	Query DN A Length	Query Protein Length	Subject DN A Length	Subject Protein Length	% ID Protein	% ID DN A
217, 218	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthetase enzyme SEQ ID NO:3.	AED10 581	1.00 E-132	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	7.00 E-14	5.1.1.1	12 24	40 7	0	408	92	
219, 220	alanine racemase [Aeromonas salmonicida subsp. salmonicida A449]	1.45 E+08	0	Aeromonas salmonicida subsp. salmonicida A449	Empedobacter mature peptide synthetase enzyme SEQ ID NO:3.	AED10 581	1.00 E-129	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	6.00 E-11	5.1.1.1	12 27	40 8	0	408	95	
221, 222	alanine racemase [Aeromonas salmonicida subsp. salmonicida A449]	1.45 E+08	0	Aeromonas salmonicida subsp. salmonicida A449	Empedobacter mature peptide synthetase enzyme SEQ ID NO:3.	AED10 581	1.00 E-128	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	6.00 E-11	5.1.1.1	12 24	40 7	0	408	95	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Eval ue	Geneseq DNA Description	Geneseq DNA Accession Code	Eval ue	Predicted EC Number	Qu ery DN A Le ngt h	Qu ery Protein Le ngt h	Subj ect DN A Le ngt h	Subj ect Protein Len gth	% ID Protein	% ID DNA
223, 224	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	1.00 E-133	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	2.00 E-14	5.1.1.1	12 24	40 7	0	408	93	
225, 226	alanine racemase [Aeromonas salmonicida subsp. salmonicida A449]	1.45 E+08	0	Aeromonas salmonicida subsp. salmonicida A449	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	1.00 E-130	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	6.00 E-11	5.1.1.1	12 27	40 8	0	408	94	
227, 228	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	1.00 E-132	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	2.00 E-14	5.1.1.1	12 27	40 8	0	408	91	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Gene Eval	Geneseq DNA Description	Geneseq DNA Accession Code	Gene Eval	Predicted EC Number	Query DN A Length	Query Protein Length	Subject DN A Length	Subject Protein Length	% ID Protein	% ID DNA
229, 230	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	1.00 E-130	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	3.00 E-13	5.1.1.1	12 27	40 8	0	408	91	
231, 232	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	1.00 E-130	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	6.00 E-11	5.1.1.1	12 27	40 8	0	408	95	
233, 234	alanine racemase [Aeromonas salmonicida subsp. salmonicida A449]	1.45 E+08	0	Aeromonas salmonicida subsp. salmonicida A449	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	1.00 E-128	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	6.00 E-11	5.1.1.1	12 27	40 8	0	408	95	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Genes eq Protein Accession Code	Geneseq DNA Description	Genes eq DNA Accession Code	Genes Evaluate	Predicted EC Number	Query DNA Length	Query Protein Length	Subject DNA Length	Subject Protein Length	% ID Protein	% ID DNA
235, 236	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter brevis mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	6.00 E-05	5.1.1.1	1224	407	0	408	95	
237, 238	alanine racemase [Aeromonas salmonicida subsp. salmonicida A449]	1.45 E+08	0	Aeromonas salmonicida subsp. salmonicida A449	Empedobacter brevis mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	6.00 E-05	5.1.1.1	1227	408	0	408	96	
239, 240	alanine racemase [Aeromonas salmonicida subsp. salmonicida A449]	1.45 E+08	0	Aeromonas salmonicida subsp. salmonicida A449	Empedobacter brevis mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	6.00 E-11	5.1.1.1	1224	407	0	408	94	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Evaluate	Geneseq DNA Description	Geneseq DNA Accession Code	Evaluate	Predicted EC Number	Query DNA Length	Query Protein Length	Subject DNA Length	Subject Protein Length	% ID Protein	% ID DNA
241, 242	alanine racemase [Aeromonas salmonicida subsp. salmonicida A449]	1.45 E+08	0	Aeromonas salmonicida subsp. salmonicida A449	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-130	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	6.00 E-11	5.1.1.1	127	408	0	408	95	
243, 244	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-129	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	4.00 E-06	5.1.1.1	127	408	0	408	95	
245, 246	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-129	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	4.00 E-06	5.1.1.1	127	408	0	408	95	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Evaluate	Geneseq DNA Description	Geneseq DNA Accession Code	Evaluate	Predicted EC Number	Query DNA Length	Query Protein Length	Subject DNA Length	Subject Protein Length	% ID Protein	% ID DNA
247, 248	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter brevis mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-130	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	4.00 E-06	5.1.1.1	1227	408	0	408	96	
249, 250	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter brevis mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-129	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	4.00 E-06	5.1.1.1	1227	408	0	408	95	
251, 252	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter brevis mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-130	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	4.00 E-06	5.1.1.1	1227	408	0	408	95	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Evaluate	Geneseq DNA Description	Geneseq DNA Accession Code	Evaluate	Predicted EC Number	Query DNALength	Query ProteinLength	Query DNALength	Subject DNALength	Subject ProteinLength	% ID Protein	% ID DNA
253, 254	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter brevis mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-129	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	4.00 E-06	5.1.1.1	1224	407	0	408	95	95	
255, 256	hypothetical protein [Mesorhizobium loti].	13474743	1.00 E-170	Mesorhizobium loti	Propionibacterium acnes predicted ORF-encoded polypeptide #300.	ABM37068	1.00 E-77	Plant polypeptide, SEQ ID 5546.	ADT17374	0.2	5.1.1.1	1086	361	1827	608	83	80	
257, 258	alanine racemase [Aeromonas salmonicida subsp. salmonicida A449]	1.45 E+08	0	Aeromonas salmonicida subsp. salmonicida A449	Empedobacter brevis mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-129	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	2.00 E-08	5.1.1.1	1227	408	0	408	95	95	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Gene Eval	Geneseq DNA Description	Geneseq DNA Accession Code	Gene Eval	Predicted EC Number	Query DNALength	Query ProteinLength	Subject DNALength	Subject ProteinLength	% ID Protein	% ID DNA
259, 260	alanine racemase [Aeromonas salmonicida subsp. salmonicida A449]	1.45 E+08	0	Aeromonas salmonicida subsp. salmonicida A449	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-130	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	6.00 E-11	5.1.1.1	127	408	0	408	95	95
261, 262	alanine racemase [Aeromonas salmonicida subsp. salmonicida A449]	1.45 E+08	0	Aeromonas salmonicida subsp. salmonicida A449	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-130	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	2.00 E-08	5.1.1.1	127	408	0	408	94	94
263, 264	alanine racemase [Aeromonas salmonicida subsp. salmonicida A449]	1.45 E+08	0	Aeromonas salmonicida subsp. salmonicida A449	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-128	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	6.00 E-11	5.1.1.1	127	408	0	408	95	95

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Gene Eval	Geneseq DNA Description	Geneseq DNA Accession Code	Gene Eval	Predicted EC Number	Query DN Length	Query Protein Length	Subject DN Length	Subject Protein Length	% ID Protein	% ID DNA
265, 266	alanine racemase [Aeromonas salmonicida subsp. salmonicida A449]	1.45 E+08	0	Aeromonas salmonicida subsp. salmonicida A449	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-129	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	6.00 E-11	5.1.1.1	127	408	0	408	95	
267, 268	alanine racemase [Aeromonas salmonicida subsp. salmonicida A449]	1.45 E+08	0	Aeromonas salmonicida subsp. salmonicida A449	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-129	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	6.00 E-11	5.1.1.1	127	408	0	408	94	
269, 270	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-132	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	1.00 E-12	5.1.1.1	127	408	0	408	92	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Gene Eval	Geneseq DNA Description	Geneseq DNA Accession Code	Gene Eval	Predicted EC Number	Query DN A Length	Query Protein Length	Subject DN A Length	Subject Protein Length	% ID Protein	% ID DNA
271, 272	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthetase enzyme SEQ ID NO:3.	AED10 581	1.00 E-132	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	3.00 E-13	5.1.1.1	12 30	40 9	0	408	92	
273, 274	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthetase enzyme SEQ ID NO:3.	AED10 581	1.00 E-130	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	6.00 E-11	5.1.1.1	12 27	40 8	0	408	96	
275, 276	alanine racemase [Aeromonas salmonicida subsp. salmonicida A449]	1.45 E+08	0	Aeromonas salmonicida subsp. salmonicida A449	Empedobacter mature peptide synthetase enzyme SEQ ID NO:3.	AED10 581	1.00 E-128	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	6.00 E-11	5.1.1.1	12 27	40 8	0	408	97	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Eval ue	Geneseq DNA Description	Geneseq DNA Accession Code	Eval ue	Predicted EC Number	Qu ery DN A Le ngt h	Qu ery Protein Le ngt h	Subj ect DN A Le ngt h	Subj ect Protein Len gth	% ID Protein	% ID DN A
277, 278	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthetase enzyme SEQ ID NO:3.	AED10 581	1.00 E-133	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	2.00 E-14	5.1.1.1	12 27	40 8	0	408	92	
279, 280	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthetase enzyme SEQ ID NO:3.	AED10 581	1.00 E-132	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	1.00 E-15	5.1.1.1	12 27	40 8	0	408	92	
281, 282	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthetase enzyme SEQ ID NO:3.	AED10 581	1.00 E-132	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	1.00 E-18	5.1.1.1	12 27	40 8	0	408	92	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Eval ue	Geneseq DNA Description	Geneseq DNA Accession Code	Eval ue	Predicted EC Number	Qu ery DN A Le ngt h	Qu ery Protein Le ngt h	Subj ect DN A Le ngt h	Subj ect Protein Len gth	% ID Protein	% ID DN A
283, 284	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthetase enzyme SEQ ID NO:3.	AED10 581	1.00 E-132	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	1.00 E-18	5.1.1.1	12 27	40 8	0	408	92	
285, 286	alanine racemase [Aeromonas salmonicida subsp. salmonicida A449]	1.45 E+08	0	Aeromonas salmonicida subsp. salmonicida A449	Empedobacter mature peptide synthetase enzyme SEQ ID NO:3.	AED10 581	1.00 E-130	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	6.00 E-11	5.1.1.1	12 27	40 8	0	408	95	
287, 288	alanine racemase [Aeromonas salmonicida subsp. salmonicida A449]	1.45 E+08	0	Aeromonas salmonicida subsp. salmonicida A449	Empedobacter mature peptide synthetase enzyme SEQ ID NO:3.	AED10 581	1.00 E-129	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	6.00 E-11	5.1.1.1	12 27	40 8	0	408	95	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Eval ue	Geneseq DNA Description	Geneseq DNA Accession Code	Eval ue	Predicted EC Number	Qu ery DN A Le ngth	Qu ery Protein Le ngth	Subj ect DN A Le ngth	Subj ect Protein Le ngth	% ID Protein	% ID DN A
289, 290	alanine racemase [Aeromonas salmonicida subsp. salmonicida A449]	1.45 E+08	0	Aeromonas salmonicida subsp. salmonicida A449	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	1.00 E-129	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	6.00 E-11	5.1.1.1	12 27	40 8	0	408	95	
291, 292	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	1.00 E-131	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	2.00 E-14	5.1.1.1	12 27	40 8	0	408	92	
293, 294	alanine racemase [Aeromonas salmonicida subsp. salmonicida A449]	1.45 E+08	0	Aeromonas salmonicida subsp. salmonicida A449	Empedobacter mature peptide synthesizing enzyme SEQ ID NO:3.	AED10 581	1.00 E-127	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	6.00 E-11	5.1.1.1	12 27	40 8	0	408	94	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Eval ue	Geneseq DNA Description	Geneseq DNA Accession Code	Eval ue	Predi cted EC Number	Qu ery DN A Le ngt h	Qu ery Pro tein Le ngt h	Subj ect DN A Le ngt h	Subj ect Prot ein Len gth	% ID Pro tein	% ID DN A
295, 296	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	0	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter brevis mature peptide synthetizing enzyme SEQ ID NO:3.	AED10581	1.00 E-129	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	4.00 E-06	5.1.1.1	127	408	0	408	95	
297, 298	proline racemase [Burkholderia phymatum STM815] gi 117982269 gb EAU96656.1 proline racemase [Burkholderia phymatum STM815]	1.18 E+08	104	Burkholderia phymatum STM815	Acinetobacter baumannii protein #19.	ADA35228	1.00 E-105	Prokaryotic essential gene #34740.	ACA19650	9.00 E-22	5.1.1.4	960	319	945	331		
299, 300	alanine racemase [Pseudomonas putida GB-1] gi 126314851 gb EAZ66019.1 alanine racemase [Pseudomonas putida GB-1]	1.26 E+08	170	Pseudomonas putida GB-1	T. maritima D-alanine-D-alanine ligase.	AED11803	1.00 E-169	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	5.00 E-95	5.1.1.1	1230	409	0	409	73	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Evaluate	Geneseq DNA Description	Geneseq DNA Accession Code	Evaluate	Predicted EC Number	Query DNALength	Query ProteinLength	Subject DNALength	Subject ProteinLength	% ID Protein	% ID DNA
301, 302	alanine racemase [Pseudomonas putida KT2440]	2699 0430	1.00 E-143	Pseudomonas putida KT2440	T. maritima D-alanine-D-alanine ligase.	AED11 803	1.00 E-144	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB9 9538	5.00 E-27	5.1.1.1	12 39	41 2	122 7	409		
303, 304	hypothetical protein PaerP_01003954 [Pseudomonas aeruginosa PA7]	9441 4235	4.00 E-33	Pseudomonas aeruginosa PA7	Pseudomonas aeruginosa polypeptide #3.	ABO82 155	4.00 E-33	Prokaryotic gene #34740.	ACA4 4058	7.00 E-14	5.1.1.4	37 5	12 4	0	314	60	
305, 306	AGR_L_3051p [Agrobacterium tumefaciens]. proline racemase [Microscilla marina ATCC 23134] gi 123984081 gb EAY24454.1	1589 1641	1.00 E-128	Agrobacterium tumefaciens	L. pneumophila protein SEQ ID NO 3367.	AEB41 596	3.00 E-39	Prokaryotic gene #34740.	ACA2 7267	0.21	5.1.1.1	11 25	37 4	116 7	388	64	66
307, 308	proline racemase [Microscilla marina ATCC 23134]	1.24 E+08	3.00 E-59	Microscilla marina ATCC 23134	Bacterial polypeptide #10001.	ADS22 995	3.00 E-47	Bacterial polypeptide #10001.	ADS6 0041	2.00 E-05	5.1.1.4	42 9	14 2	0	333	73	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Geneseq DNA Description	Geneseq DNA Accession Code	Eval ue	Predi cted EC Number	Qu ery DN A Le ngth	Qu ery Protein Le ngth	Subj ect DN A Le ngth	Subj ect Protein Len gth	% ID Protein	% ID DN A
309, 310	alanine racemase [Stenotrophomonas maltophilia R551-3] gij119820021 gb EAX22642.1 alanine racemase [Stenotrophomonas maltophilia R551-3]	1.2E+08	2.00 E-57	Stenotrophomonas maltophilia R551-3	Prokaryotic essential gene #34740.	ABU41398	Prokaryotic essential gene #34740.	ACA43665	8.00 E-52	5.1.1.1	408	135	0	355	80	
311, 312	mandelate racemase/muconate lactonizing enzyme family protein [Sulfitobacter sp. NAS-14.1] gij83842294 gb EAP81462.1 mandelate racemase/muconate lactonizing enzyme family protein [Sulfitobacter sp. NAS-14.1]	83953326	1.00 E-143	Sulfitobacter sp. NAS-14.1	Klebsiella pneumoniae polypeptide seqid 7178.	ABO61307	Novel mar regulated protein (NIMR) #29.	AAS46252	2.00 E-71	5.1.2.2	939	312	0	321	81	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Evaluate	Geneseq DNA Description	Geneseq DNA Accession Code	Evaluate	Predicted EC Number	Query DNA Length	Query Protein Length	Subject DNA Length	Subject Protein Length	% ID Protein	% ID DNA
313, 314	proline racemase [Mesorhizobium loti].	13473394	1.00 E-120	Mesorhizobium loti	Bacterial polypeptide #10001.	ADS22995	1.00 E-120	C. botulinum active BoNT/A modified open reading frame, SEQ ID No:7.	AEF99607	0.65	5.1.1.4	909	302	1002	333	70	62
315, 316	proline racemase [Flavobacteriales bacterium HTCC2170] gij8708932 gb E AR01166.1 proline racemase [Flavobacteriales bacterium HTCC2170]	88712394	1.00 E-106	Flavobacteriales bacterium HTCC2170	Bacterial polypeptide #10001.	ADS22995	1.00 E-72	Haemophilus influenzae (NTHi) protein - SEQ ID 618.	ADT05504	0.03	5.1.1.4	684	227	0	335	79	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Genes eq Protein Accession Code	Evaluate	Geneseq DNA Description	Genes eq DNA Accession Code	Evaluate	Predicted EC Number	Query DN A Length	Query Protein Length	Subject DN A Length	Subject Protein Length	% ID Protein	% ID DN A
317, 318	Mandelate racemase/muconate lactonizing enzyme [Flavobacteriales bacterium HTCC2170] gi 88709134 gb E AR01368.1	8871 2596	1.00 E-110	Flavobacteriales bacterium HTCC2170	Bacteroides fragilis strain 14062 protein, SEQ:5227.	AEX28 600	5.00 E-34	Prokaryotic essential gene #34740.	ACA2 3465	2.9	5.5.1.1	10 32	34 3	0	348	57	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Geneseq DNA Description	Geneseq DNA Accession Code	Eval ue	Predicted EC Number	Qu ery DN A Le ngt h	Qu ery Protein Le ngt h	Subj ect DN A Le ngt h	Subj ect Protein Le ngt h	% ID Protein	% ID DN A
319, 320	COG3938: Proline racemase [Pseudomonas aeruginosa C3719] gij84328204 ref Z P_00976211.1 COG3938: Proline racemase [Pseudomonas aeruginosa 2192] gij107100719 ref Z P_01364637.1 hypothetical protein PaerPA_0100174 6 [Pseudomonas aeruginosa PACS2] gij12616632	8432 1952	1.00 E-60	Pseudomonas aeruginosa C3719	Pseudomonas aeruginosa polypeptide #3.	ABO82 155	Pseudomonas aeruginosa polypeptide #3.	ABD1 5666	4.00 E-61	5.1.1.4	63 6	21 1	486	339		
321, 322	aspartate racemase [Pyrococcus abyssii].	1452 1575	2.00 E-28	Pyrococcus abyssii	Thermococcus kodakaraensis KOD1 protein sequence SeqID4.	ADN46 207	Breast cancer related marker, seq id 2.	ACN8 9360	7.00 E-25	5.1.1.13	69 9	23 2	687	228	31	46

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Genes eq Protein Accession Code	Geneseq DNA Description	Genes eq DNA Accession Code	Eval ue	Pred icted EC Num ber	Qu ery DN A Le ngt h	Qu ery Pro tein Le ngt h	Subj ect DN A Le ngt h	Subj ect Prot ein Len gth	% ID Pro tein	% ID DN A
323, 324	proline racemase, putative [Psychroflexus torquis ATCC 700755] gi 91184469 gb EAS70852.1 proline racemase, putative [Psychroflexus torquis ATCC 700755]	9121 7361	1.00 E-152	Psychrofl exus torquis ATCC 700755	Bacterial polypeptide #10001.	ADS22 995	Bacterial polypeptide #10001.	ADS6 0041	1.00 E-110	5.1.1.4	88 5	29 4	0	336	85	
325, 326	proline racemase [Microscilla marina ATCC 23134] gi 123984081 gb EAY24454.1 proline racemase [Microscilla marina ATCC 23134]	1.24 E+08	1.00 E-145	Microscill a marina ATCC 23134	Bacterial polypeptide #10001.	ADS22 995	Bacterial polypeptide #10001.	ADS6 0041	1.00 E-126	5.1.1.4	10 05	33 4	0	333	73	
327, 328	proline racemase [Microscilla marina ATCC 23134] gi 123984081 gb EAY24454.1 proline racemase [Microscilla marina ATCC 23134]	1.24 E+08	1.00 E-147	Microscill a marina ATCC 23134	Bacterial polypeptide #10001.	ADS22 995	Bacterial polypeptide #10001.	ADS6 0041	1.00 E-132	5.1.1.4	10 02	33 3	0	333	74	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Evaluate	Geneseq DNA Description	Geneseq DNA Accession Code	Evaluate	Predicted EC Number	Query DNA Length	Query Protein Length	Subject DNA Length	Subject Protein Length	% ID Protein	% ID DNA
329, 330	putative proline racemase [Myxococcus xanthus DK 1622]	1.09 E+08	1.00 E-101	Myxococcus xanthus DK 1622	M. xanthus protein sequence, seq id 9726.	ABM96 637	1.00 E-101	Prokaryotic essential gene #34740.	ACA2 3259	1.00 E-17	5.1.1.4	957	318	0	311	58	58
331, 332	AGR_L_3051p [Agrobacterium tumefaciens].	1589 1641	7.00 E-44	Agrobacterium tumefaciens	Bacterial polypeptide #19.	ADF07 948	5.00 E-21	Mycobacterium tuberculosis strain H37Rv genome SEQ ID NO 2.	AAI99 682	1.1	5.1.1.1	420	139	1167	388	61	68
333, 334	putative alanine racemase [Rhizobium leguminosarum bv. viciae 3841]	1.16 E+08	9.00 E-97	Rhizobium leguminosarum bv. viciae 3841	Bacterial polypeptide #19.	ADF07 948	6.00 E-29	Plant full length insert polynucleotide seqid 4980.	ADX5 1636	0.67	5.1.1.1	939	312	0	377	58	58
335, 336	alanine racemase [Moorella thermoacetica ATCC 39073]	8359 0990	5.00 E-78	Moorella thermoacetica ATCC 39073	Prokaryotic essential gene #34740.	ABU24 721	1.00 E-63	Enterobacter cloacae protein amino acid sequence - SEQ ID 5666.	AEH5 4260	0.053	5.1.1.1	1143	380	0	373	41	41

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Geneseq Evaluate	Geneseq DNA Description	Geneseq DNA Accession Code	Geneseq Evaluate	Predicted EC Number	Query DNALength	Query ProteinLength	Subject DNALength	Subject ProteinLength	% ID Protein	% ID DNA
337, 338	alanine racemase [Moorella thermoacetica ATCC 39073]	8359 0990	3.00 E-77	Moorella thermoacetica ATCC 39073	Prokaryotic essential gene #34740.	ABU24 721	1.00 E-60	Bacillus licheniformis araA gene fragment amplifying PCR primer #1.	AAD2 9866	8.00 E-04	5.1.1.1	368	373	0	44		
339, 340	Alanine racemase [Azotobacter vinelandii AvOP] gij67087270 gb E AM06737.1 Alanine racemase [Azotobacter vinelandii AvOP]	6715 5469	1.00 E-115	Azotobacter vinelandii AvOP	Prokaryotic essential gene #34740.	ABU39 795	1.00 E-115	Prokaryotic essential gene #34740.	ACA2 3293	5.00 E-05	5.1.1.1	355	418	0	60		
341, 342	mandelate racemase/muconate lactonizing enzyme [Sphingopyxis alaskensis RB2256]	1.03 E+08	2.00 E-99	Sphingopyxis alaskensis RB2256	Klebsiella pneumoniae polypeptide seqid 7178.	ABO61 307	1.00 E-45	Geranylgeranyl pyrophosphate synthase polypeptide #7.	ADM9 8687	0.04 E-6	5.5.1.1	329	335	0	57		
343, 344	putative alanine racemase [Rhizobium leguminosarum bv. viciae 3841]	1.16 E+08	1.00 E-127	Rhizobium leguminosarum bv. viciae 3841	Pseudomonas aeruginosa polypeptide #3.	ABO84 274	8.00 E-42	Streptomyces cattleya NRRL 8057 orfY protein.	ADO5 1695	0.83	5.1.1.1	377	377	0	62		

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Evaluate	Geneseq DNA Description	Geneseq DNA Accession Code	Evaluate	Predicted EC Number	Query DN A Length	Query Protein Length	Query DN A Length	Query Protein Length	Subject DN A Length	Subject Protein Length	% ID DN A	% ID Protein
345, 346	AGR_L_3051p [Agrobacterium tumefaciens].	1589 1641	2.00 E-79	Agrobacterium tumefaciens	Photophosphorylation protein sequence #59.	ABM69 114	1.00 E-27	Prokaryotic essential gene #34740.	ACA2 5617	2.2	5.1.1.1	77 7	25 8	116 7	388	57	65		
347, 348	putative alanine racemase [Rhizobium leguminosarum bv. viciae 3841]	1.16 E+08	1.00 E-114	Rhizobium leguminosarum bv. viciae 3841	Prokaryotic essential gene #34740.	ABU41 398	1.00 E-36	Plant full length insert polynucleotide seqid 4980.	ADX5 1636	0.81	5.1.1.1	11 13	37 0	0 0	377	58			
349, 350	Mandelate racemase/muconate lactonizing enzyme; C-terminal domain protein [Methylobacterium sp. 4-46]	1.49 E+08	2.00 E-87	Methylobacterium sp. 4-46	Klebsiella pneumoniae polypeptide seqid 7178.	ABO61 307	1.00 E-63	Prokaryotic essential gene #34740.	ACA4 2641	0.72	5.1.2.2	99 3	33 0	0 0	326	51			
351, 352	Mandelate racemase/muconate lactonizing enzyme; C-terminal domain protein [Sphingomonas wittichii RW1]	1.49 E+08	3.00 E-93	Sphingomonas wittichii RW1	Klebsiella pneumoniae polypeptide seqid 7178.	ABO61 307	3.00 E-48	Klebsiella pneumoniae polypeptide seqid 7178.	ACH9 4858	5.00 E-05	5.5.1.1	10 05	33 4	0 0	354	52			

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Evaluate	Geneseq DNA Description	Geneseq DNA Accession Code	Evaluate	Predicted EC Number	Query DNALength	Query ProteinLength	Subject DNALength	Subject ProteinLength	% ID Protein	% ID DNA
353, 354	putative proline racemase [Burkholderia xenovorans LB400]	9177 9222	1.00 E-38	Burkholderia xenovorans LB400	Prokaryotic essential gene #34740.	ABU21 813	3.00 E-39	Prokaryotic essential gene #34740.	ACA1 9650	3.00 E-22	5.1.1.4	381	126	945	318		
355, 356	alanine racemase [Lentisphaera araneosa HTCC2155]	1.49 E+08	2.00 E-67	Lentisphaera araneosa HTCC2155	Prokaryotic essential gene #34740.	ABU23 921	1.00 E-56	Prokaryotic essential gene #34740.	ACA3 8093	9.00 E-04	5.1.1.1	1155	384	0	360	39	
357, 358	alanine racemase [Roseiflexus castenholzii DSM 13941] gij118010932 gb E AV24951.1 alanine racemase [Roseiflexus castenholzii DSM 13941]	1.18 E+08	1.00 E-107	Roseiflexus castenholzii DSM 13941	Prokaryotic essential gene #34740.	ABU23 921	5.00 E-75	Prokaryotic essential gene #34740.	ACA4 0224	0.88	5.1.1.1	1203	400	0	849	52	
359, 360	alanine racemase [Roseiflexus castenholzii DSM 13941] alanine racemase [Roseiflexus castenholzii DSM 13941] gij118010932 gb E AV24951.1	1.18 E+08	1.00 E-107	Roseiflexus castenholzii DSM 13941	Prokaryotic essential gene #34740.	ABU23 921	1.00 E-75	T. versicolor pyrF PCR primer SEQ ID 5.	AAF2 6441	0.88	5.1.1.1	1203	400	0	849	52	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Genes eq Protein Accession Code	Evaluate	Geneseq DNA Description	Genes eq DNA Accession Code	Evaluate	Predicted EC Number	Query DN A Length	Query Protein Length	Subject DN A Length	Subject Protein Length	% ID Protein	% ID DN A
361, 362	alanine racemase [Roseiflexus castenholzii DSM 13941] gi 118010932 gb E AV24951.1 alanine racemase [Roseiflexus castenholzii DSM 13941]	1.18 E+08	1.00 E-107	Roseiflexus castenholzii DSM 13941	Prokaryotic essential gene #34740.	ABU23 921	5.00 E-75	T. versicolor pyrF PCR primer SEQ ID 5.	AAF2 6441	0.88	5.1.1.1	1203	400	0	849	52	
363, 364	alanine racemase [Roseiflexus castenholzii DSM 13941] gi 118010932 gb E AV24951.1 alanine racemase [Roseiflexus castenholzii DSM 13941]	1.18 E+08	1.00 E-107	Roseiflexus castenholzii DSM 13941	Prokaryotic essential gene #34740.	ABU23 921	3.00 E-75	T. versicolor pyrF PCR primer SEQ ID 5.	AAF2 6441	0.88	5.1.1.1	1203	400	0	849	52	
365, 366	alanine racemase [Roseiflexus castenholzii DSM 13941] gi 118010932 gb E AV24951.1 alanine racemase [Roseiflexus castenholzii DSM 13941]	1.18 E+08	1.00 E-107	Roseiflexus castenholzii DSM 13941	Prokaryotic essential gene #34740.	ABU23 921	3.00 E-76	T. versicolor pyrF PCR primer SEQ ID 5.	AAF2 6441	0.88	5.1.1.1	1203	400	0	849	52	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Evaluate	Geneseq DNA Description	Geneseq DNA Accession Code	Evaluate	Predicted EC Number	Query DNALength	Query ProteinLength	Subject DNALength	Subject ProteinLength	% ID Protein	% ID DNA
367, 368	alanine racemase [Pseudomonas mendocina ymp]	1.46 E+08	1.00 E-168	Pseudomonas mendocina ymp	Prokaryotic essential gene #34740.	ABU41398	1.00 E-160	Prokaryotic essential gene #34740.	ACA43665	7.00 E-32	5.1.1.1	1074	357	0	362	82	
369, 370	diaminopimelate epimerase [Nostoc sp. PCC 7120].	1723 2333	3.00 E-78	Nostoc sp. PCC 7120	Bacterial polypeptide #10001.	ADS29848	9.00 E-79	S. lavendulae gene mutagenic PCR primer #2.	ADE10236	0.16	5.1.1.7	861	286	1413	285		
371, 372	diaminopimelate epimerase [Syntrophus aciditrophicus SB]	8585 7916	2.00 E-76	Syntrophus aciditrophicus SB	Bacterial polypeptide #10001.	ADN17489	5.00 E-60	Bacterial polypeptide #10001.	ADT46187	2.00 E-06	5.1.1.7	813	270	0	277	57	
373, 374	diaminopimelate epimerase [Nostoc sp. PCC 7120].	1723 2333	2.00 E-71	Nostoc sp. PCC 7120	Bacterial polypeptide #10001.	ADS29848	7.00 E-72	Environmental isolate hydrolyase, SEQ ID NO:44.	AEH47413	0.16	5.1.1.7	873	290	900	285		
375, 376	Diaminopimelate epimerase [Clostridium thermocellum ATCC 27405]	1.26 E+08	5.00 E-33	Clostridium thermocellum ATCC 27405	Bacterial polypeptide #10001.	ADN27169	2.00 E-31	Bacterial polypeptide #10001.	ADT42245	9.2	5.1.1.7	828	275	0	280	29	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Genes eq Protein Accession Code	Evaluate	Geneseq DNA Description	Genes eq DNA Accession Code	Evaluate	Predicted EC Number	Query DN A Length	Query Protein Length	Subject DN A Length	Subject Protein Length	% ID Protein	% ID DN A
377, 378	UDP-N-acetylglucosamine 2-epimerase [Chloroflexus aggregans DSM 9485] gi 117997290 gb EAV11478.1 UDP-N-acetylglucosamine 2-epimerase [Chloroflexus aggregans DSM 9485]	1.18 E+08	2.00 E-97	Chloroflexus aggregans DSM 9485	Prokaryotic essential gene #34740.	ABU24 769	6.00 E-80	Prokaryotic essential gene #34740.	ACA2 7041	6.00 E-08	5.1.3.14	1200	399	0	386	50	
379, 380	NmrA family protein [Burkholderia vietnamiensis G4]	1.34 E+08	2.00 E-88	Burkholderia vietnamiensis G4	Bacterial polypeptide #10001.	ADS22 779	2.00 E-84	Bacterial polypeptide #10001.	ADS5 9825	3.00 E-06	1.6.5.3	879	292	0	287	55	
381, 382	hypothetical protein RL1205 [Rhizobium leguminosarum bv. viciae 3841]	1.16 E+08	2.00 E-87	Rhizobium leguminosarum bv. viciae 3841	Bacterial polypeptide #10001.	ADS22 779	1.00 E-83	Bacterial polypeptide #10001.	ADS5 9825	1.00 E-08	1.6.5.3	879	292	0	289	55	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Geneseq Evaluate	Geneseq DNA Description	Geneseq DNA Accession Code	Geneseq Evaluate	Predicted EC Number	Query DNA Length	Query Protein Length	Subject DNA Length	Subject Protein Length	% ID Protein	% ID DNA
383, 384	similar to chloromuconate cycloisomerase [Gloeobacter violaceus PCC 7421]	3752 2668	1.00 E-93	Gloeobacter violaceus PCC 7421	L. pneumophila protein SEQ ID NO 3367.	AEB37 282	4.00 E-48	Novel canine microarray-related DNA sequence SeqID10021.	ADQ5 3782	3.1	5.1.2.2	10 71	35 6	0	356	50	
385, 386				Aurantimonas sp. SI85-9A1		AEH19 277	5.00 E-47			3.4	5.1.1.-	11 01	36 6	287 30	379	34	95
387, 388				Azoarcus sp. EbN1		ABU41 398	1.00 E-110			1.00 E-08	5.1.1.1	10 65	35 4	146 4	357	57	88
389, 390				Enterobacter sp. 638		AEH63 094	0			0	5.1.1.1	10 80	35 9	108 9	362	96	87
391, 392				Enterobacter sp. 638		AEH63 094	0			0	5.1.1.1	10 80	35 9	108 9	362	96	87
393, 394				Aquifex aeolicus VF5		ABB06 296	1.00 E-147			2.00 E-07	5.1.1.1	10 17	33 8	146 8	341	72	95
395, 396				Streptomyces coelicolor A3(2)		ABU34 223	6.00 E-90			2.00 E-05	5.1.1.1	11 49	38 2	146 4	391	50	92

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Evaluate	Geneseq DNA Description	Geneseq DNA Accession Code	Evaluate	Predicted EC Number	Query DN A Length	Query Protein Length	Subject DN A Length	Subject Protein Length	% ID Protein	% ID DN A
397, 398				Rhodospirillum rubrum HaA2		ABO84274	4.00 E-50			0.014	5.1.1.1	1122	373	2046	368	38	100
399, 400				Pseudomonas putida F1		AED11803	0			0	5.1.1.1	1161	386	1227	409	96	92
401, 402				Pseudomonas putida GB-1		AED11803	1.00 E-174			1.00	5.1.1.1	1164	387	1227	409	77	85
403, 404				Pseudomonas putida GB-1		AED11804	1.00 E-175			1.00	5.1.1.1	1164	387	1227	409	78	85
405, 406				Pseudomonas putida F1		AED11804	0			0	5.1.1.1	1161	386	1227	409	97	96
407, 408				Pseudomonas putida F1		AED11803	0			0	5.1.1.1	1161	386	1227	409	96	92
409, 410				Pseudomonas putida F1		AED11803	0			0	5.1.1.1	1161	386	1227	409	96	92
411, 412				alpha proteobacterium HTCC2255		ABB06296	9.00 E-13			0.077	5.1.1.1	1521	506	633	341	28	100

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Evaluate	Geneseq DNA Description	Geneseq DNA Accession Code	Evaluate	Predicted EC Number	Query DN A Length	Query Protein Length	Query DN A Length	Query Protein Length	Subject DN A Length	Subject Protein Length	% ID Protein	% ID DN A
413, 414				Streptomyces avermitilis MA-4680		ABM54358	2.00E-53			0.052		1032	343	306	346			37	100
415, 416				Aeromonas hydrophila subsp. hydrophila ATCC 7966		AED10581	1.00E-129			6.00E-05	5.1.1.1	1167	388	1227	386			61	83
417, 418				Aeromonas hydrophila subsp. hydrophila ATCC 7966		AED10581	1.00E-131			2.00E-11	5.1.1.1	1167	388	1227	386			60	83
419, 420				Aeromonas hydrophila subsp. hydrophila ATCC 7966		AED10581	1.00E-132			2.00E-08	5.1.1.1	1167	388	1227	386			62	88

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Geneseq DNA Description	Geneseq DNA Accession Code	Gene Eval	Predicted EC Number	Query DN A Length	Query Protein Length	Subject DN A Length	Subject Protein Length	% ID Protein	% ID DN A
421, 422				Aeromonas hydrophila subsp. hydrophila ATCC 7966		AED10581			7.00 E-11	5.1.1.1	1167	388	1227	386	61	90
423, 424				Pseudomonas putida F1		AED11803			0	5.1.1.1	1161	386	1227	409	95	92
425, 426				Pseudomonas putida F1		AED11803			2.00 E-26	5.1.1.1	1092	363	1227	409	64	81
427, 428				Pseudomonas putida GB-1		AED11803			1.00 E-124	5.1.1.1	1230	409	1227	409	76	85
429, 430				Pseudomonas putida F1		AED11804			0	5.1.1.1	1230	409	1227	409	97	96
431, 432				Pseudomonas putida F1		AED11803			0	5.1.1.1	1161	386	1227	409	95	91
433, 434				Pseudomonas putida GB-1		AED11804			1.00 E-124	5.1.1.1	1164	387	1227	409	78	85

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Genes eq Protein Accession Code	Evaluate	Geneseq DNA Description	Genes eq DNA Accession Code	Evaluate	Predicted EC Number	Query DN A Length	Query Protein Length	Query DN A Length	Query Protein Length	Subject DN A Length	Subject Protein Length	% ID Protein	% ID DN A
435, 436				Aeromonas hydrophila subsp. hydrophila ATCC 7966		AED10581	1.00 E-131			2.00 E-14	5.1.1.1	1167	388	1227	386	1227	386	61	91
437, 438				Pyrococcus abyssi GE5		ADN46207	2.00 E-25			0.14	5.1.1.13	714	237	1056	232	1056	232	31	100
439, 440				Aeromonas hydrophila subsp. hydrophila ATCC 7966		AED10581	1.00 E-127			6.00 E-11	5.1.1.1	1095	364	1227	386	1227	386	61	86
441, 442				Aeromonas hydrophila subsp. hydrophila ATCC 7966		AED10581	1.00 E-133			2.00 E-14	5.1.1.1	1167	388	1227	386	1227	386	60	91
443, 444				Aeromonas hydrophila subsp. hydrophila ATCC 7966		AED10581	1.00 E-132			7.00 E-14	5.1.1.1	1167	388	1227	386	1227	386	61	81

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Geneseq DNA Description	Geneseq DNA Accession Code	Gene	Eval	Predicted EC Number	Query DNA Length	Query Protein Length	Subject DNA Length	Subject Protein Length	% ID Protein	% ID DNA
445, 446				Aeromonas salmonicida subsp. salmonicida A449		AED10581				1.00 E-128	5.1.1.1	1167	388	1227	386	59	86
447, 448				Aeromonas hydrophila subsp. hydrophila ATCC 7966		AED10581				1.00 E-129	5.1.1.1	1167	388	1227	386	61	83
449, 450				Aeromonas hydrophila subsp. hydrophila ATCC 7966		AED10581				1.00 E-130	5.1.1.1	1167	388	1227	386	61	83
451, 452				Aeromonas hydrophila subsp. hydrophila ATCC 7966		AED10581				1.00 E-130	5.1.1.1	1167	388	1227	386	62	86

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Genes eq Protein Accession Code	Evaluate	Geneseq DNA Description	Genes eq DNA Accession Code	Evaluate	Predicted EC Number	Query DN A Length	Query Protein Length	Query Protein Le ngth	Subject DN A Length	Subject Protein Length	% ID Protein	% ID DN A
453, 454				Aeromonas salmonicida subsp. salmonicida A449		AED10581	1.00 E-129			7.00 E-11	5.1.1.1	1167	388	1227	386	61	86	
455, 456				Stappia aggregata IAM 12614		ABU21813	1.00 E-101			5.00 E-11	5.1.1.4	957	318	960	318	60	90	
457, 458				Myxococcus xanthus DK 1622		ABM96637	1.00 E-75			1.00 E-18	5.1.1.4	1074	357	960	311	51	88	
459, 460				Aeromonas hydrophila subsp. hydrophila ATCC 7966		AED10581	1.00 E-130			1.00 E-12	5.1.1.1	1167	388	1227	386	61	90	
461, 462				Pseudomonas putida GB-1		AED10581	1.00 E-159			4.00 E-95	5.1.1.1	1092	363	1227	386	77	85	
463, 464				Mesorhizobium sp. BNC1		ABU38829	3.00 E-46			0.01 E-5	5.1.1.1	1143	380	801	358	38	100	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Genes Protein Accession Code	Evaluate	Geneseq DNA Description	Genes eq DNA Accession Code	Evaluate	Predicted EC Number	Query DN A Length	Query Protein Length	Subject DN A Length	Subject Protein Length	% ID Protein	% ID DNA
465, 466				Microscilla marina ATCC 23134		ADS22995	1.00 E-128			3.1	5.1.1.4	1002	333	84480	333	65	95
467, 468				Burkholderia phymatum STM815		ADA35228	1.00 E-105			1.00 E-21	5.1.1.4	960	319	945	331	60	91
469, 470				Sulfitobacter sp. NAS-14.1		ABO61307	2.00 E-71			0.74	5.1.2.2	939	312	10118	347	50	93
471, 472				Pseudomonas mendocinaymp		ABU41398	1.00 E-160			6.00 E-45	5.1.1.1	1074	357	1071	357	78	82
473, 474				Aeromonas hydrophila subsp. hydrophila ATCC 7966		AED10581	1.00 E-131			7.00 E-11	5.1.1.1	1167	388	1227	386	62	86
475, 476				Pseudomonas putida GB-1		AED10581	1.00 E-167			5.00 E-95	5.1.1.1	1164	387	1227	386	76	85

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Geneseq Protein Accession Code	Evaluate	Geneseq DNA Description	Geneseq DNA Accession Code	Evaluate	Predicted EC Number	Query DNALength	Query ProteinLength	Query DNALength	Subject DNALength	Subject ProteinLength	% ID Protein	% ID DNA
477, 478				Pseudomonas putida KT2440		AED11803	1.00 E-141			5.00 E-27	5.1.1.1	1176	391	1227	409	63	81	
479, 480	protein of unknown function DUF453 [Novosphingobium aromaticivorans DSM 12444]	87200831	1.00 E-137	Novosphingobium aromaticivorans DSM 12444	Klebsiella pneumoniae polypeptide seqid 7178.	ABO64182	1.00 E-113	Bacterial polypeptide #23667.	ADS50054	0.79		1056	351	0	352	68		
481, 482	alanine racemase [Aeromonas hydrophila subsp. hydrophila ATCC 7966]	1.18 E+08	1.00 E-177	Aeromonas hydrophila subsp. hydrophila ATCC 7966	Empedobacter brevis mature peptide synthesizing enzyme SEQ ID NO:3.	AED10581	1.00 E-129	Pseudomonas putida racemase peptide, SEQ ID 5.	ADB99538	6.00 E-14	5.1.1.1	1089	362	0	408	83		
483, 484	threonine aldolase family protein [Caulobacter crescentus CB15] gij13424764 gb AK25068.1 threonine aldolase family protein [Caulobacter crescentus CB15]	16127336	0	Caulobacter crescentus CB15					19172958	0	5.1.1.-	1110	369	4E+06	369	100	100	

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Genes eq Protein Accession Code	Geneseq DNA Description	Genes eq DNA Accession Code	Eval ue	Predic ted EC Number	Qu ery DN A Le ngt h	Qu ery Protein Le ngt h	Subj ect DN A Le ngt h	Subj ect Protein Len gth	% ID Protein	% ID DN A
485, 486	hypothetical protein AGR_L_1837 [Agrobacterium tumefaciens str. C58] gij17937630 ref NP_534419.1 hypothetical protein Atu3927 [Agrobacterium tumefaciens str. C58] gij15159365 gb AK89493.1 AGR_L_1837p [Agrobacterium tumefaciens str. C58] gij17742368 gb AL44735.1 conserved hypothetical protein [Agrobacterium tumefaciens str. C58]	1589 1036	0	Agrobacterium tumefaciens str. C58				16445 346	0	5.1.1.-	1089	362	2E+06	362	100	100

SEQ ID NO:	NR Description	NR Accession Code	NR Evaluate	Organism	Geneseq Protein Description	Genes eq Protein Accession Code	Geneseq DNA Description	Genes eq DNA Accession Code	Eval ue	Predi cted EC Number	Qu ery DN A Le ngt h	Qu ery Protein Le ngt h	Subj ect DN A Le ngt h	Subj ect Protein Len gth	% ID Protein	% ID DN A
487, 488	hypothetical protein AGR_L_1808 [Agrobacterium tumefaciens str. C58] gj 15159347 gb AK89478.1 AGR_L_1808p [Agrobacterium tumefaciens str. C58]	1589 1021	0	Agrobacterium tumefaciens str. C58				16445 346	0	5.1.1. -	11 46	38 1	2E+ 06	397	10 0	10 0

The invention provides variants of polynucleotides or polypeptides of the invention, which comprise sequences modified at one or more base pairs, codons, introns, exons, or amino acid residues (respectively) yet still retain the biological activity of an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase of the invention. Variants can be produced by any number of means included methods such as, for example, error-prone PCR, shuffling, oligonucleotide-directed mutagenesis, assembly PCR, sexual PCR mutagenesis, *in vivo* mutagenesis, cassette mutagenesis, recursive ensemble mutagenesis, exponential ensemble mutagenesis, site-specific mutagenesis, gene reassembly (e.g., GeneReassembly, see, e.g., U.S. Patent No. 6,537,776), GSSM and any combination thereof.

The term “saturation mutagenesis”, “gene site saturation mutagenesis” or “GSSM” includes a method that uses degenerate oligonucleotide primers to introduce point mutations into a polynucleotide, as described in detail, below.

The term “optimized directed evolution system” or “optimized directed evolution” includes a method for reassembling fragments of related nucleic acid sequences, e.g., related genes, and explained in detail, below.

The term “synthetic ligation reassembly” or “SLR” includes a method of ligating oligonucleotide fragments in a non-stochastic fashion, and explained in detail, below.

Generating and Manipulating Nucleic Acids

The invention provides nucleic acids (e.g., nucleic acids encoding polypeptides having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity; including enzymes having at least one sequence modification of an exemplary nucleic acid sequence of the invention (as defined above), wherein the sequence modification comprises one or more nucleotide residue changes (or the equivalent thereof), including expression cassettes such as expression vectors, encoding the polypeptides of the invention.

The invention also includes methods for discovering new isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase sequences using the nucleic acids of the invention. The invention also includes methods for inhibiting the expression of isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase genes, transcripts and polypeptides using the nucleic acids of the invention. Also provided are methods for modifying the nucleic acids of the invention by,

e.g., synthetic ligation reassembly, optimized directed evolution system and/or saturation mutagenesis.

The nucleic acids of the invention can be made, isolated and/or manipulated by, e.g., cloning and expression of cDNA libraries, amplification of message or genomic DNA
5 by PCR, and the like.

In one aspect, the invention also provides an isomerase-, e.g., a racemase-, e.g., an amino acid racemase-, an alanine racemase-, and/or an epimerase-isomerase-, e.g.,
10 racemase-, e.g., amino acid racemase-, alanine racemase-, and/or epimerase-encoding nucleic acids with a common novelty in that they are derived from an environmental source, or a bacterial source, or an archaeal source.

In practicing the methods of the invention, homologous genes can be modified by manipulating a template nucleic acid, as described herein. The invention can be practiced in conjunction with any method or protocol or device known in the art, which are well described in the scientific and patent literature.

15 One aspect of the invention is an isolated nucleic acid comprising one of the sequences of The invention and sequences substantially identical thereto, the sequences complementary thereto, or a fragment comprising at least 10, 15, 20, 25, 30, 35, 40, 50, 75, 100, 150, 200, 300, 400, or 500 consecutive bases of one of the sequences of a Sequence of the invention (or the sequences complementary thereto). The isolated, nucleic acids may
20 comprise DNA, including cDNA, genomic DNA and synthetic DNA. The DNA may be double-stranded or single-stranded and if single stranded may be the coding strand or non-coding (anti-sense) strand. Alternatively, the isolated nucleic acids may comprise RNA.

Accordingly, another aspect of the invention is an isolated nucleic acid which encodes one of the polypeptides of the invention, or fragments comprising at least 5, 10,
25 15, 20, 25, 30, 35, 40, 50, 75, 100, or 150 consecutive amino acids of one of the polypeptides of the invention. The coding sequences of these nucleic acids may be identical to one of the coding sequences of one of the nucleic acids of the invention, or a fragment thereof or may be different coding sequences which encode one of the polypeptides of the invention, sequences substantially identical thereto and fragments
30 having at least 5, 10, 15, 20, 25, 30, 35, 40, 50, 75, 100, or 150 consecutive amino acids of one of the polypeptides of the invention, as a result of the redundancy or degeneracy of the genetic code. The genetic code is well known to those of skill in the art and can be obtained, for example, on page 214 of B. Lewin, Genes VI, Oxford University Press, 1997.

The isolated nucleic acid which encodes one of the polypeptides of the invention and sequences substantially identical thereto, may include, but is not limited to: only the coding sequence of a nucleic acid of the invention and sequences substantially identical thereto and additional coding sequences, such as leader sequences or proprotein sequences and non-coding sequences, such as introns or non-coding sequences 5' and/or 3' of the coding sequence. Thus, as used herein, the term "polynucleotide encoding a polypeptide" encompasses a polynucleotide which includes only the coding sequence for the polypeptide as well as a polynucleotide which includes additional coding and/or non-coding sequence.

Alternatively, the nucleic acid sequences of the invention and sequences substantially identical thereto, may be mutagenized using conventional techniques, such as site directed mutagenesis, or other techniques familiar to those skilled in the art, to introduce silent changes into the polynucleotides of the invention and sequences substantially identical thereto. As used herein, "silent changes" include, for example, changes which do not alter the amino acid sequence encoded by the polynucleotide. Such changes may be desirable in order to increase the level of the polypeptide produced by host cells containing a vector encoding the polypeptide by introducing codons or codon pairs which occur frequently in the host organism.

The invention also relates to polynucleotides which have nucleotide changes which result in amino acid substitutions, additions, deletions, fusions and truncations in the polypeptides of the invention and sequences substantially identical thereto. Such nucleotide changes may be introduced using techniques such as site directed mutagenesis, random chemical mutagenesis, exonuclease III deletion and other recombinant DNA techniques. Alternatively, such nucleotide changes may be naturally occurring allelic variants which are isolated by identifying nucleic acids which specifically hybridize to probes comprising at least 10, 15, 20, 25, 30, 35, 40, 50, 75, 100, 150, 200, 300, 400, or 500 consecutive bases of one of the sequences of The invention and sequences substantially identical thereto (or the sequences complementary thereto) under conditions of high, moderate, or low stringency as provided herein.

General Techniques

The nucleic acids used to practice this invention, whether RNA, iRNA, antisense nucleic acid, cDNA, genomic DNA, vectors, viruses or hybrids thereof, may be isolated from a variety of sources, genetically engineered, amplified, and/or expressed/ generated

recombinantly. Recombinant polypeptides (e.g., isomerases, e.g., racemases, e.g., amino acid racemases, alanine racemases, and/or epimerases of the invention) generated from these nucleic acids can be individually isolated or cloned and tested for a desired activity. Any recombinant expression system can be used, including bacterial, mammalian, yeast, 5 insect or plant cell expression systems.

Alternatively, these nucleic acids can be synthesized *in vitro* by well-known chemical synthesis techniques, as described in, e.g., Adams (1983) J. Am. Chem. Soc. 105:661; Belousov (1997) Nucleic Acids Res. 25:3440-3444; Frenkel (1995) Free Radic. Biol. Med. 19:373-380; Blommers (1994) Biochemistry 33:7886-7896; Narang (1979) 10 Meth. Enzymol. 68:90; Brown (1979) Meth. Enzymol. 68:109; Beaucage (1981) Tetra. Lett. 22:1859; U.S. Patent No. 4,458,066.

Techniques for the manipulation of nucleic acids, such as, e.g., subcloning, labeling probes (e.g., random-primer labeling using Klenow polymerase, nick translation, amplification), sequencing, hybridization and the like are well described in the scientific 15 and patent literature, see, e.g., Sambrook, ed., MOLECULAR CLONING: A LABORATORY MANUAL (2ND ED.), Vols. 1-3, Cold Spring Harbor Laboratory, (1989); CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, Ausubel, ed. John Wiley & Sons, Inc., New York (1997); LABORATORY TECHNIQUES IN BIOCHEMISTRY AND MOLECULAR BIOLOGY: HYBRIDIZATION WITH 20 NUCLEIC ACID PROBES, Part I. Theory and Nucleic Acid Preparation, Tijssen, ed. Elsevier, N.Y. (1993).

Another useful means of obtaining and manipulating nucleic acids used to practice the methods of the invention is to clone from genomic samples, and, if desired, screen and re-clone inserts isolated or amplified from, e.g., genomic clones or cDNA clones. Sources 25 of nucleic acid used in the methods of the invention include genomic or cDNA libraries contained in, e.g., mammalian artificial chromosomes (MACs), see, e.g., U.S. Patent Nos. 5,721,118; 6,025,155; human artificial chromosomes, see, e.g., Rosenfeld (1997) Nat. Genet. 15:333-335; yeast artificial chromosomes (YAC); bacterial artificial chromosomes (BAC); P1 artificial chromosomes, see, e.g., Woon (1998) Genomics 50:306-316; P1- 30 derived vectors (PACs), see, e.g., Kern (1997) Biotechniques 23:120-124; cosmids, recombinant viruses, phages or plasmids.

In one aspect, a nucleic acid encoding a polypeptide of the invention is assembled in appropriate phase with a leader sequence capable of directing secretion of the translated polypeptide or fragment thereof.

The invention provides fusion proteins and nucleic acids encoding them. A polypeptide of the invention can be fused to a heterologous peptide or polypeptide, such as N-terminal identification peptides which impart desired characteristics, such as increased stability or simplified purification. Peptides and polypeptides of the invention can also be synthesized and expressed as fusion proteins with one or more additional domains linked thereto for, e.g., producing a more immunogenic peptide, to more readily isolate a recombinantly synthesized peptide, to identify and isolate antibodies and antibody-expressing B cells, and the like. Detection and purification facilitating domains include, e.g., metal chelating peptides such as polyhistidine tracts and histidine-tryptophan modules that allow purification on immobilized metals, protein A domains that allow purification on immobilized immunoglobulin, and the domain utilized in the FLAGS extension/affinity purification system (Immunex Corp, Seattle WA). The inclusion of a cleavable linker sequences such as Factor Xa or enterokinase (Invitrogen, San Diego CA) between a purification domain and the motif-comprising peptide or polypeptide to facilitate purification. For example, an expression vector can include an epitope-encoding nucleic acid sequence linked to six histidine residues followed by a thioredoxin and an enterokinase cleavage site (see e.g., Williams (1995) *Biochemistry* 34:1787-1797; Dobeli (1998) *Protein Expr. Purif.* 12:404-414). The histidine residues facilitate detection and purification while the enterokinase cleavage site provides a means for purifying the epitope from the remainder of the fusion protein. Technology pertaining to vectors encoding fusion proteins and application of fusion proteins are well described in the scientific and patent literature, see e.g., Kroll (1993) *DNA Cell. Biol.*, 12:441-53.

The phrases “nucleic acid” or “nucleic acid sequence” as used herein refer to an oligonucleotide, nucleotide, polynucleotide, or to a fragment of any of these, to DNA or RNA of genomic or synthetic origin which may be single-stranded or double-stranded and may represent a sense or antisense strand, to peptide nucleic acid (PNA), or to any DNA-like or RNA-like material, natural or synthetic in origin. The phrases “nucleic acid” or “nucleic acid sequence” includes oligonucleotide, nucleotide, polynucleotide, or to a fragment of any of these, to DNA or RNA (e.g., mRNA, rRNA, tRNA, iRNA) of genomic or synthetic origin which may be single-stranded or double-stranded and may represent a sense or antisense strand, to peptide nucleic acid (PNA), or to any DNA-like or RNA-like material, natural or synthetic in origin, including, e.g., iRNA, ribonucleoproteins (e.g., e.g., double stranded iRNAs, e.g., iRNPs). The term encompasses nucleic acids, i.e., oligonucleotides, containing known analogues of natural nucleotides. The term also

encompasses nucleic-acid-like structures with synthetic backbones, see e.g., Mata (1997) Toxicol. Appl. Pharmacol. 144:189-197; Strauss-Soukup (1997) Biochemistry 36:8692-8698; Samstag (1996) Antisense Nucleic Acid Drug Dev 6:153-156. "Oligonucleotide" includes either a single stranded polydeoxynucleotide or two complementary
5 polydeoxynucleotide strands that may be chemically synthesized. Such synthetic oligonucleotides have no 5' phosphate and thus will not ligate to another oligonucleotide without adding a phosphate with an ATP in the presence of a kinase. A synthetic oligonucleotide can ligate to a fragment that has not been dephosphorylated.

A "coding sequence of" or a "nucleotide sequence encoding" a particular
10 polypeptide or protein, is a nucleic acid sequence which is transcribed and translated into a polypeptide or protein when placed under the control of appropriate regulatory sequences.

The term "gene" means the segment of DNA involved in producing a polypeptide chain; it includes regions preceding and following the coding region (leader and trailer) as well as, where applicable, intervening sequences (introns) between individual coding
15 segments (exons). "Operably linked" as used herein refers to a functional relationship between two or more nucleic acid (e.g., DNA) segments. Typically, it refers to the functional relationship of transcriptional regulatory sequence to a transcribed sequence. For example, a promoter is operably linked to a coding sequence, such as a nucleic acid of the invention, if it stimulates or modulates the transcription of the coding sequence in an
20 appropriate host cell or other expression system. Generally, promoter transcriptional regulatory sequences that are operably linked to a transcribed sequence are physically contiguous to the transcribed sequence, i.e., they are cis-acting. However, some transcriptional regulatory sequences, such as enhancers, need not be physically contiguous or located in close proximity to the coding sequences whose transcription they enhance.

25 The term "expression cassette" as used herein refers to a nucleotide sequence which is capable of affecting expression of a structural gene (i.e., a protein coding sequence, such as an isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase of the invention) in a host compatible with such sequences. Expression cassettes include at least a promoter operably linked with the polypeptide
30 coding sequence; and, in one aspect, with other sequences, e.g., transcription termination signals. Additional factors necessary or helpful in effecting expression may also be used, e.g., enhancers. Thus, expression cassettes also include plasmids, expression vectors, recombinant viruses, any form of recombinant "naked DNA" vector, and the like. A "vector" comprises a nucleic acid that can infect, transfect, transiently or permanently

transduce a cell. It will be recognized that a vector can be a naked nucleic acid, or a nucleic acid complexed with protein or lipid. The vector in one aspect comprises viral or bacterial nucleic acids and/or proteins, and/or membranes (e.g., a cell membrane, a viral lipid envelope, etc.). Vectors include, but are not limited to replicons (e.g., RNA
5 replicons, bacteriophages) to which fragments of DNA may be attached and become replicated. Vectors thus include, but are not limited to RNA, autonomous self-replicating circular or linear DNA or RNA (e.g., plasmids, viruses, and the like, see, e.g., U.S. Patent No. 5,217,879), and include both the expression and non-expression plasmids. Where a recombinant microorganism or cell culture is described as hosting an "expression vector"
10 this includes both extra-chromosomal circular and linear DNA and DNA that has been incorporated into the host chromosome(s). Where a vector is being maintained by a host cell, the vector may either be stably replicated by the cells during mitosis as an autonomous structure, or is incorporated within the host's genome.

As used herein, the term "promoter" includes all sequences capable of driving
15 transcription of a coding sequence in a cell, e.g., a plant cell. Thus, promoters used in the constructs of the invention include *cis*-acting transcriptional control elements and regulatory sequences that are involved in regulating or modulating the timing and/or rate of transcription of a gene. For example, a promoter can be a *cis*-acting transcriptional control element, including an enhancer, a promoter, a transcription terminator, an origin of
20 replication, a chromosomal integration sequence, 5' and 3' untranslated regions, or an intronic sequence, which are involved in transcriptional regulation. These *cis*-acting sequences typically interact with proteins or other biomolecules to carry out (turn on/off, regulate, modulate, etc.) transcription. "Constitutive" promoters are those that drive expression continuously under most environmental conditions and states of development
25 or cell differentiation. "Inducible" or "regulatable" promoters direct expression of the nucleic acid of the invention under the influence of environmental conditions or developmental conditions. Examples of environmental conditions that may affect transcription by inducible promoters include anaerobic conditions, elevated temperature, drought, or the presence of light.

30 "Tissue-specific" promoters are transcriptional control elements that are only active in particular cells or tissues or organs, e.g., in plants or animals. Tissue-specific regulation may be achieved by certain intrinsic factors that ensure that genes encoding proteins specific to a given tissue are expressed. Such factors are known to exist in mammals and plants so as to allow for specific tissues to develop.

As used herein, the term “isolated” means that the material (e.g., a nucleic acid, a polypeptide, a cell) is removed from its original environment (e.g., the natural environment if it is naturally occurring). For example, a naturally-occurring polynucleotide or polypeptide present in a living animal is not isolated, but the same
5 polynucleotide or polypeptide, separated from some or all of the coexisting materials in the natural system, is isolated. Such polynucleotides could be part of a vector and/or such polynucleotides or polypeptides could be part of a composition and still be isolated in that such vector or composition is not part of its natural environment. As used herein, the term “purified” does not require absolute purity; rather, it is intended as a relative definition.

10 Individual nucleic acids obtained from a library have been conventionally purified to electrophoretic homogeneity. The sequences obtained from these clones could not be obtained directly either from the library or from total human DNA. The purified nucleic acids of the invention have been purified from the remainder of the genomic DNA in the organism by at least 10^4 - 10^6 fold. However, the term “purified” also includes nucleic acids
15 that have been purified from the remainder of the genomic DNA or from other sequences in a library or other environment by at least one order of magnitude, typically two or three orders and more typically four or five orders of magnitude.

As used herein, the term “recombinant” means that the nucleic acid is adjacent to a “backbone” nucleic acid to which it is not adjacent in its natural environment. Additionally,
20 to be “enriched” the nucleic acids will represent 5% or more of the number of nucleic acid inserts in a population of nucleic acid backbone molecules. Backbone molecules according to the invention include nucleic acids such as expression vectors, self-replicating nucleic acids, viruses, integrating nucleic acids and other vectors or nucleic acids used to maintain or manipulate a nucleic acid insert of interest. Typically, the enriched nucleic acids represent
25 15% or more of the number of nucleic acid inserts in the population of recombinant backbone molecules. More typically, the enriched nucleic acids represent 50% or more of the number of nucleic acid inserts in the population of recombinant backbone molecules. In a one aspect, the enriched nucleic acids represent 90% or more of the number of nucleic acid inserts in the population of recombinant backbone molecules.

30 “Plasmids” are designated by a lower case “p” preceded and/or followed by capital letters and/or numbers. The starting plasmids herein are either commercially available, publicly available on an unrestricted basis, or can be constructed from available plasmids in accord with published procedures. In addition, equivalent plasmids to those described herein are known in the art and will be apparent to the ordinarily skilled artisan.

“Plasmids” can be commercially available, publicly available on an unrestricted basis, or can be constructed from available plasmids in accord with published procedures.

Equivalent plasmids to those described herein are known in the art and will be apparent to the ordinarily skilled artisan.

5 “Digestion” of DNA refers to catalytic cleavage of the DNA with a restriction enzyme that acts only at certain sequences in the DNA. The various restriction enzymes used herein are commercially available and their reaction conditions, cofactors and other requirements were used as would be known to the ordinarily skilled artisan. For analytical purposes, typically 1 μg of plasmid or DNA fragment is used with about 2 units of enzyme
10 in about 20 μl of buffer solution. For the purpose of isolating DNA fragments for plasmid construction, typically 5 to 50 μg of DNA are digested with 20 to 250 units of enzyme in a larger volume. Appropriate buffers and substrate amounts for particular restriction enzymes are specified by the manufacturer. Incubation times of about 1 hour at 37°C are ordinarily used, but may vary in accordance with the supplier's instructions. After
15 digestion, gel electrophoresis may be performed to isolate the desired fragment.

“Hybridization” refers to the process by which a nucleic acid strand joins with a complementary strand through base pairing. Hybridization reactions can be sensitive and selective so that a particular sequence of interest can be identified even in samples in which it is present at low concentrations. Suitably stringent conditions can be defined by,
20 for example, the concentrations of salt or formamide in the prehybridization and hybridization solutions, or by the hybridization temperature and are well known in the art. In particular, stringency can be increased by reducing the concentration of salt, increasing the concentration of formamide, or raising the hybridization temperature. In alternative aspects, nucleic acids of the invention are defined by their ability to hybridize under
25 various stringency conditions (e.g., high, medium, and low), as set forth herein.

For example, hybridization under high stringency conditions could occur in about 50% formamide at about 37°C to 42°C. Hybridization could occur under reduced stringency conditions in about 35% to 25% formamide at about 30°C to 35°C. In particular, hybridization could occur under high stringency conditions at 42°C in 50%
30 formamide, 5X SSPE, 0.3% SDS and 200 $\mu\text{g}/\text{ml}$ sheared and denatured salmon sperm DNA. Hybridization could occur under reduced stringency conditions as described above, but in 35% formamide at a reduced temperature of 35°C. The temperature range corresponding to a particular level of stringency can be further narrowed by calculating the

purine to pyrimidine ratio of the nucleic acid of interest and adjusting the temperature accordingly. Variations on the above ranges and conditions are well known in the art.

Transcriptional and translational control sequences

The invention provides nucleic acid (e.g., DNA) sequences of the invention
5 operatively linked to expression (e.g., transcriptional or translational) control sequence(s),
e.g., promoters or enhancers, to direct or modulate RNA synthesis/ expression. The
expression control sequence can be in an expression vector. Exemplary bacterial
promoters include *lacI*, *lacZ*, T3, T7, *gpt*, lambda PR, PL and *trp*. Exemplary eukaryotic
promoters include CMV immediate early, HSV thymidine kinase, early and late SV40,
10 LTRs from retrovirus, and mouse metallothionein I. A promoter sequence is “operably
linked to” a coding sequence when RNA polymerase which initiates transcription at the
promoter will transcribe the coding sequence into mRNA.

Promoters suitable for expressing a polypeptide in bacteria include the *E.*
coli lac or *trp* promoters, the *lacI* promoter, the *lacZ* promoter, the T3 promoter, the T7
15 promoter, the *gpt* promoter, the lambda PR promoter, the lambda PL promoter, promoters
from operons encoding glycolytic enzymes such as 3-phosphoglycerate kinase (PGK), and
the acid phosphatase promoter. Eukaryotic promoters include the CMV immediate early
promoter, the HSV thymidine kinase promoter, heat shock promoters, the early and late
SV40 promoter, LTRs from retroviruses, and the mouse metallothionein-I promoter.
20 Other promoters known to control expression of genes in prokaryotic or eukaryotic cells or
their viruses may also be used. Promoters suitable for expressing the polypeptide or
fragment thereof in bacteria include the *E. coli lac* or *trp* promoters, the *lacI* promoter, the
lacZ promoter, the T3 promoter, the T7 promoter, the *gpt* promoter, the *lambda P_R*
promoter, the *lambda P_L* promoter, promoters from operons encoding glycolytic enzymes
25 such as 3-phosphoglycerate kinase (PGK) and the acid phosphatase promoter. Fungal
promoters include the ∇ factor promoter. Eukaryotic promoters include the CMV
immediate early promoter, the HSV thymidine kinase promoter, heat shock promoters, the
early and late SV40 promoter, LTRs from retroviruses and the mouse metallothionein-I
promoter. Other promoters known to control expression of genes in prokaryotic or
30 eukaryotic cells or their viruses may also be used.

Tissue-Specific Plant Promoters

The invention provides expression cassettes that can be expressed in a tissue-specific manner, e.g., that can express an isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase of the invention in a tissue-specific manner.

5 The invention also provides plants or seeds that express an isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase of the invention in a tissue-specific manner. The tissue-specificity can be seed specific, stem specific, leaf specific, root specific, fruit specific and the like.

In one aspect, a constitutive promoter such as the CaMV 35S promoter can be used
10 for expression in specific parts of the plant or seed or throughout the plant. For example, for overexpression, a plant promoter fragment can be employed which will direct expression of a nucleic acid in some or all tissues of a plant, e.g., a regenerated plant. Such promoters are referred to herein as "constitutive" promoters and are active under most environmental conditions and states of development or cell differentiation.

15 Examples of constitutive promoters include the cauliflower mosaic virus (CaMV) 35S transcription initiation region, the 1'- or 2'- promoter derived from T-DNA of *Agrobacterium tumefaciens*, and other transcription initiation regions from various plant genes known to those of skill. Such genes include, e.g., *ACT11* from *Arabidopsis* (Huang (1996) *Plant Mol. Biol.* 33:125-139); *Cat3* from *Arabidopsis* (GenBank No. U43147, Zhong (1996) *Mol. Gen. Genet.* 251:196-203); the gene encoding stearyl-acyl carrier protein desaturase from *Brassica napus* (Genbank No. X74782, Solocombe (1994) *Plant Physiol.* 104:1167-1176); *GPc1* from maize (GenBank No. X15596; Martinez (1989) *J. Mol. Biol.* 208:551-565); the *Gpc2* from maize (GenBank No. U45855, Manjunath (1997) *Plant Mol. Biol.* 33:97-112); plant promoters described in U.S. Patent Nos. 4,962,028;
20 5,633,440.
25

The invention uses tissue-specific or constitutive promoters derived from viruses which can include, e.g., the tobamovirus subgenomic promoter (Kumagai (1995) *Proc. Natl. Acad. Sci. USA* 92:1679-1683; the rice tungro bacilliform virus (RTBV), which replicates only in phloem cells in infected rice plants, with its promoter which drives
30 strong phloem-specific reporter gene expression; the cassava vein mosaic virus (CVMV) promoter, with highest activity in vascular elements, in leaf mesophyll cells, and in root tips (Verdaguer (1996) *Plant Mol. Biol.* 31:1129-1139).

Alternatively, the plant promoter may direct expression of an isomerase-, e.g., a racemase-, e.g., an amino acid racemase-, an alanine racemase-, and/or an epimerase-isomerase-, e.g., racemase-, e.g., amino acid racemase-, alanine racemase-, and/or epimerase-expressing nucleic acid in a specific tissue, organ or cell type (*i.e.* tissue-specific promoters) or may be otherwise under more precise environmental or developmental control or under the control of an inducible promoter. Examples of environmental conditions that may affect transcription include anaerobic conditions, elevated temperature, the presence of light, or sprayed with chemicals/hormones. For example, the invention incorporates the drought-inducible promoter of maize (Busk (1997) supra); the cold, drought, and high salt inducible promoter from potato (Kirch (1997) *Plant Mol. Biol.* 33:897-909).

Tissue-specific promoters can promote transcription only within a certain time frame of developmental stage within that tissue. See, e.g., Blazquez (1998) *Plant Cell* 10:791-800, characterizing the *Arabidopsis* LEAFY gene promoter. See also Cardon (1997) *Plant J* 12:367-77, describing the transcription factor SPL3, which recognizes a conserved sequence motif in the promoter region of the *A. thaliana* floral meristem identity gene AP1; and Mandel (1995) *Plant Molecular Biology*, Vol. 29, pp 995-1004, describing the meristem promoter eIF4. Tissue specific promoters which are active throughout the life cycle of a particular tissue can be used. In one aspect, the nucleic acids of the invention are operably linked to a promoter active primarily only in cotton fiber cells. In one aspect, the nucleic acids of the invention are operably linked to a promoter active primarily during the stages of cotton fiber cell elongation, e.g., as described by Rinehart (1996) supra. The nucleic acids can be operably linked to the Fbl2A gene promoter to be preferentially expressed in cotton fiber cells (Ibid). See also, John (1997) *Proc. Natl. Acad. Sci. USA* 89:5769-5773; John, et al., U.S. Patent Nos. 5,608,148 and 5,602,321, describing cotton fiber-specific promoters and methods for the construction of transgenic cotton plants. Root-specific promoters may also be used to express the nucleic acids of the invention. Examples of root-specific promoters include the promoter from the alcohol dehydrogenase gene (DeLisle (1990) *Int. Rev. Cytol.* 123:39-60). Other promoters that can be used to express the nucleic acids of the invention include, e.g., ovule-specific, embryo-specific, endosperm-specific, integument-specific, seed coat-specific promoters, or some combination thereof; a leaf-specific promoter (see, e.g., Busk (1997) *Plant J.* 11:1285-1295, describing a leaf-specific promoter in maize); the ORF13 promoter from *Agrobacterium rhizogenes* (which exhibits high activity in roots, see, e.g.,

Hansen (1997) supra); a maize pollen specific promoter (see, e.g., Guerrero (1990) Mol. Gen. Genet. 224:161-168); a tomato promoter active during fruit ripening, senescence and abscission of leaves and, to a lesser extent, of flowers can be used (see, e.g., Blume (1997) Plant J. 12:731-746); a pistil-specific promoter from the potato SK2 gene (see, e.g., Ficker
5 (1997) Plant Mol. Biol. 35:425-431); the Blec4 gene from pea, which is active in epidermal tissue of vegetative and floral shoot apices of transgenic alfalfa making it a useful tool to target the expression of foreign genes to the epidermal layer of actively growing shoots or fibers; the ovule-specific BEL1 gene (see, e.g., Reiser (1995) Cell 83:735-742, GenBank No. U39944); and/or, the promoter in Klee, U.S. Patent No.
10 5,589,583, describing a plant promoter region is capable of conferring high levels of transcription in meristematic tissue and/or rapidly dividing cells.

Alternatively, plant promoters which are inducible upon exposure to plant hormones, such as auxins, are used to express the nucleic acids of the invention. For example, the invention can use the auxin-response elements E1 promoter fragment
15 (AuxREs) in the soybean (*Glycine max* L.) (Liu (1997) Plant Physiol. 115:397-407); the auxin-responsive *Arabidopsis* GST6 promoter (also responsive to salicylic acid and hydrogen peroxide) (Chen (1996) Plant J. 10: 955-966); the auxin-inducible parC promoter from tobacco (Sakai (1996) 37:906-913); a plant biotin response element (Streit (1997) Mol. Plant Microbe Interact. 10:933-937); and, the promoter responsive to the
20 stress hormone abscisic acid (Sheen (1996) Science 274:1900-1902).

The nucleic acids of the invention can also be operably linked to plant promoters which are inducible upon exposure to chemicals reagents which can be applied to the plant, such as herbicides or antibiotics. For example, the maize In2-2 promoter, activated by benzenesulfonamide herbicide safeners, can be used (De Veylder (1997) Plant Cell
25 Physiol. 38:568-577); application of different herbicide safeners induces distinct gene expression patterns, including expression in the root, hydathodes, and the shoot apical meristem. Coding sequence can be under the control of, e.g., a tetracycline-inducible promoter, e.g., as described with transgenic tobacco plants containing the *Avena sativa* L. (oat) arginine decarboxylase gene (Masgrau (1997) Plant J. 11:465-473); or, a salicylic
30 acid-responsive element (Stange (1997) Plant J. 11:1315-1324). Using chemically- (e.g., hormone- or pesticide-) induced promoters, i.e., promoter responsive to a chemical which can be applied to the transgenic plant in the field, expression of a polypeptide of the invention can be induced at a particular stage of development of the plant. Thus, the invention also provides for transgenic plants containing an inducible gene encoding for

polypeptides of the invention whose host range is limited to target plant species, such as corn, rice, barley, wheat, potato or other crops, inducible at any stage of development of the crop.

5 One of skill will recognize that a tissue-specific plant promoter may drive expression of operably linked sequences in tissues other than the target tissue. Thus, a tissue-specific promoter is one that drives expression preferentially in the target tissue or cell type, but may also lead to some expression in other tissues as well.

10 The nucleic acids of the invention can also be operably linked to plant promoters which are inducible upon exposure to chemicals reagents. These reagents include, e.g., herbicides, synthetic auxins, or antibiotics which can be applied, e.g., sprayed, onto transgenic plants. Inducible expression of the an isomerase-, e.g., a racemase-, e.g., an amino acid racemase-, an alanine racemase-, and/or an epimerase-isomerase-, e.g., racemase-, e.g., amino acid racemase-, alanine racemase-, and/or epimerase-producing nucleic acids of the invention will allow the grower to select plants with the optimal
15 isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase expression and/or activity. The development of plant parts can thus controlled. In this way the invention provides the means to facilitate the harvesting of plants and plant parts. For example, in various embodiments, the maize In2-2 promoter, activated by benzenesulfonamide herbicide safeners, is used (De Veylder (1997) Plant Cell Physiol.
20 38:568-577); application of different herbicide safeners induces distinct gene expression patterns, including expression in the root, hydathodes, and the shoot apical meristem. Coding sequences of the invention are also under the control of a tetracycline-inducible promoter, e.g., as described with transgenic tobacco plants containing the *Avena sativa* L. (oat) arginine decarboxylase gene (Masgrau (1997) Plant J. 11:465-473); or, a salicylic
25 acid-responsive element (Stange (1997) Plant J. 11:1315-1324).

In some aspects, proper polypeptide expression may require polyadenylation region at the 3'-end of the coding region. The polyadenylation region can be derived from the natural gene, from a variety of other plant (or animal or other) genes, or from genes in the *Agrobacterial* T-DNA.

30 The term "plant" (e.g., as in a transgenic plant or plant seed of this invention, or plant promoter used in a vector of the invention) includes whole plants, plant parts (e.g., leaves, stems, flowers, roots, etc.), plant protoplasts, seeds and plant cells and progeny of same; the classes of plants that can be used to practice this invention (including compositions and methods) can be as broad as the class of higher plants, including plants

amenable to transformation techniques, including angiosperms (monocotyledonous and dicotyledonous plants), as well as gymnosperms; also including plants of a variety of ploidy levels, including polyploid, diploid, haploid and hemizygous states. As used herein, the term "transgenic plant" includes plants or plant cells into which a heterologous nucleic acid sequence has been inserted, e.g., the nucleic acids and various recombinant constructs (e.g., expression cassettes, such a vectors) of the invention. Transgenic plants of the invention are also discussed, below.

Expression vectors and cloning vehicles

The invention provides expression vectors and cloning vehicles comprising nucleic acids of the invention, e.g., sequences encoding the isomerases, e.g., racemases, e.g., amino acid racemases, alanine racemases, and/or epimerases of the invention. Expression vectors and cloning vehicles of the invention can comprise viral particles, baculovirus, phage, plasmids, phagemids, cosmids, fosmids, bacterial artificial chromosomes, viral DNA (e.g., vaccinia, adenovirus, fowl pox virus, pseudorabies and derivatives of SV40), P1-based artificial chromosomes, yeast plasmids, yeast artificial chromosomes, and any other vectors specific for specific hosts of interest (such as bacillus, Aspergillus and yeast). Vectors of the invention can include chromosomal, non-chromosomal and synthetic DNA sequences. Large numbers of suitable vectors are known to those of skill in the art, and are commercially available. Exemplary vectors are include: bacterial: pQE vectors (Qiagen), pBluescript plasmids, pNH vectors, (lambda-ZAP vectors (Stratagene); ptrc99a, pKK223-3, pDR540, pRIT2T (Pharmacia); Eukaryotic: pXT1, pSG5 (Stratagene), pSVK3, pBPV, pMSG, pSVLSV40 (Pharmacia). However, any other plasmid or other vector may be used so long as they are replicable and viable in the host. Low copy number or high copy number vectors may be employed with the present invention.

The expression vector can comprise a promoter, a ribosome binding site for translation initiation and a transcription terminator. The vector may also include appropriate sequences for amplifying expression. Mammalian expression vectors can comprise an origin of replication, any necessary ribosome binding sites, a polyadenylation site, splice donor and acceptor sites, transcriptional termination sequences, and 5' flanking non-transcribed sequences. In some aspects, DNA sequences derived from the SV40 splice and polyadenylation sites may be used to provide the required non-transcribed genetic elements.

In one aspect, the expression vectors contain one or more selectable marker genes to permit selection of host cells containing the vector. Such selectable markers include genes encoding dihydrofolate reductase or genes conferring neomycin resistance for eukaryotic cell culture, genes conferring tetracycline or ampicillin resistance in *E. coli*,
5 and the *S. cerevisiae* TRP1 gene. Promoter regions can be selected from any desired gene using chloramphenicol transferase (CAT) vectors or other vectors with selectable markers.

Vectors for expressing the polypeptide or fragment thereof in eukaryotic cells can also contain enhancers to increase expression levels. Enhancers are cis-acting elements of DNA, usually from about 10 to about 300 bp in length that act on a promoter to increase
10 its transcription. Examples include the SV40 enhancer on the late side of the replication origin bp 100 to 270, the cytomegalovirus early promoter enhancer, the polyoma enhancer on the late side of the replication origin, and the adenovirus enhancers.

A nucleic acid sequence can be inserted into a vector by a variety of procedures. In general, the sequence is ligated to the desired position in the vector following digestion
15 of the insert and the vector with appropriate restriction endonucleases. Alternatively, blunt ends in both the insert and the vector may be ligated. A variety of cloning techniques are known in the art, e.g., as described in Ausubel and Sambrook. Such procedures and others are deemed to be within the scope of those skilled in the art.

The vector can be in the form of a plasmid, a viral particle, or a phage. Other
20 vectors include chromosomal, non-chromosomal and synthetic DNA sequences, derivatives of SV40; bacterial plasmids, phage DNA, baculovirus, yeast plasmids, vectors derived from combinations of plasmids and phage DNA, viral DNA such as vaccinia, adenovirus, fowl pox virus, and pseudorabies. A variety of cloning and expression vectors for use with prokaryotic and eukaryotic hosts are described by, e.g., Sambrook.

25 Particular bacterial vectors which can be used include the commercially available plasmids comprising genetic elements of the well known cloning vector pBR322 (ATCC 37017), pKK223-3 (Pharmacia Fine Chemicals, Uppsala, Sweden), GEM1 (Promega Biotec, Madison, WI, USA) pQE70, pQE60, pQE-9 (Qiagen), pD10, psiX174 pBluescript II KS, pNH8A, pNH16a, pNH18A, pNH46A (Stratagene), ptrc99a, pKK223-3, pKK233-
30 3, DR540, pRIT5 (Pharmacia), pKK232-8 and pCM7. Particular eukaryotic vectors include pSV2CAT, pOG44, pXT1, pSG (Stratagene) pSVK3, pBPV, pMSG, and pSVL (Pharmacia). However, any other vector may be used as long as it is replicable and viable in the host cell.

The nucleic acids of the invention can be expressed in expression cassettes, vectors or viruses and transiently or stably expressed in plant cells and seeds. One exemplary transient expression system uses episomal expression systems, e.g., cauliflower mosaic virus (CaMV) viral RNA generated in the nucleus by transcription of an episomal mini-chromosome containing supercoiled DNA, see, e.g., Covey (1990) Proc. Natl. Acad. Sci. USA 87:1633-1637. Alternatively, coding sequences, i.e., all or sub-fragments of sequences of the invention can be inserted into a plant host cell genome becoming an integral part of the host chromosomal DNA. Sense or antisense transcripts can be expressed in this manner. A vector comprising the sequences (e.g., promoters or coding regions) from nucleic acids of the invention can comprise a marker gene that confers a selectable phenotype on a plant cell or a seed. For example, the marker may encode biocide resistance, particularly antibiotic resistance, such as resistance to kanamycin, G418, bleomycin, hygromycin, or herbicide resistance, such as resistance to chlorosulfuron or Basta.

Expression vectors capable of expressing nucleic acids and proteins in plants are well known in the art, and can include, e.g., vectors from *Agrobacterium* spp., potato virus X (see, e.g., Angell (1997) EMBO J. 16:3675-3684), tobacco mosaic virus (see, e.g., Casper (1996) Gene 173:69-73), tomato bushy stunt virus (see, e.g., Hillman (1989) Virology 169:42-50), tobacco etch virus (see, e.g., Dolja (1997) Virology 234:243-252), bean golden mosaic virus (see, e.g., Morinaga (1993) Microbiol Immunol. 37:471-476), cauliflower mosaic virus (see, e.g., Cecchini (1997) Mol. Plant Microbe Interact. 10:1094-1101), maize Ac/Ds transposable element (see, e.g., Rubin (1997) Mol. Cell. Biol. 17:6294-6302; Kunze (1996) Curr. Top. Microbiol. Immunol. 204:161-194), and the maize suppressor-mutator (Spm) transposable element (see, e.g., Schlappi (1996) Plant Mol. Biol. 32:717-725); and derivatives thereof.

In one aspect, the expression vector can have two replication systems to allow it to be maintained in two organisms, for example in mammalian or insect cells for expression and in a prokaryotic host for cloning and amplification. Furthermore, for integrating expression vectors, the expression vector can contain at least one sequence homologous to the host cell genome. It can contain two homologous sequences which flank the expression construct. The integrating vector can be directed to a specific locus in the host cell by selecting the appropriate homologous sequence for inclusion in the vector. Constructs for integrating vectors are well known in the art.

Expression vectors of the invention may also include a selectable marker gene to allow for the selection of bacterial strains that have been transformed, e.g., genes which render the bacteria resistant to drugs such as ampicillin, chloramphenicol, erythromycin, kanamycin, neomycin and tetracycline. Selectable markers can also include biosynthetic
5 genes, such as those in the histidine, tryptophan and leucine biosynthetic pathways.

The DNA sequence in the expression vector is operatively linked to an appropriate expression control sequence(s) (promoter) to direct RNA synthesis. Particular named bacterial promoters include *lacI*, *lacZ*, *T3*, *T7*, *gpt*, *lambda P_R*, *P_L* and *trp*. Eukaryotic promoters include CMV immediate early, HSV thymidine kinase, early and late SV40,
10 LTRs from retrovirus and mouse metallothionein-I. Selection of the appropriate vector and promoter is well within the level of ordinary skill in the art. The expression vector also contains a ribosome binding site for translation initiation and a transcription terminator. The vector may also include appropriate sequences for amplifying expression. Promoter regions can be selected from any desired gene using chloramphenicol transferase
15 (CAT) vectors or other vectors with selectable markers. In addition, the expression vectors preferably contain one or more selectable marker genes to provide a phenotypic trait for selection of transformed host cells such as dihydrofolate reductase or neomycin resistance for eukaryotic cell culture, or such as tetracycline or ampicillin resistance in *E. coli*.

20 Mammalian expression vectors may also comprise an origin of replication, any necessary ribosome binding sites, a polyadenylation site, splice donor and acceptor sites, transcriptional termination sequences and 5' flanking nontranscribed sequences. In some aspects, DNA sequences derived from the SV40 splice and polyadenylation sites may be used to provide the required nontranscribed genetic elements.

25 Vectors for expressing the polypeptide or fragment thereof in eukaryotic cells may also contain enhancers to increase expression levels. Enhancers are cis-acting elements of DNA, usually from about 10 to about 300 bp in length that act on a promoter to increase its transcription. Examples include the SV40 enhancer on the late side of the replication origin bp 100 to 270, the cytomegalovirus early promoter enhancer, the polyoma enhancer
30 on the late side of the replication origin and the adenovirus enhancers.

In addition, the expression vectors typically contain one or more selectable marker genes to permit selection of host cells containing the vector. Such selectable markers include genes encoding dihydrofolate reductase or genes conferring neomycin resistance

for eukaryotic cell culture, genes conferring tetracycline or ampicillin resistance in *E. coli* and the *S. cerevisiae TRP1* gene.

In some aspects, the nucleic acid encoding one of the polypeptides of the invention and sequences substantially identical thereto, or fragments comprising at least about 5, 10, 5 15, 20, 25, 30, 35, 40, 50, 75, 100, or 150 consecutive amino acids thereof is assembled in appropriate phase with a leader sequence capable of directing secretion of the translated polypeptide or fragment thereof. The nucleic acid can encode a fusion polypeptide in which one of the polypeptides of the invention and sequences substantially identical thereto, or fragments comprising at least 5, 10, 15, 20, 25, 30, 35, 40, 50, 75, 100, or 150 10 consecutive amino acids thereof is fused to heterologous peptides or polypeptides, such as N-terminal identification peptides which impart desired characteristics, such as increased stability or simplified purification.

The appropriate DNA sequence may be inserted into the vector by a variety of procedures. In general, the DNA sequence is ligated to the desired position in the vector 15 following digestion of the insert and the vector with appropriate restriction endonucleases. Alternatively, blunt ends in both the insert and the vector may be ligated. A variety of cloning techniques are disclosed in Ausubel *et al.* *Current Protocols in Molecular Biology*, John Wiley 503 Sons, Inc. 1997 and Sambrook *et al.*, Molecular Cloning: A Laboratory Manual 2nd Ed., Cold Spring Harbor Laboratory Press (1989). Such procedures and others 20 are deemed to be within the scope of those skilled in the art.

The vector may be, for example, in the form of a plasmid, a viral particle, or a phage. Other vectors include chromosomal, nonchromosomal and synthetic DNA sequences, derivatives of SV40; bacterial plasmids, phage DNA, baculovirus, yeast plasmids, vectors derived from combinations of plasmids and phage DNA, viral DNA 25 such as vaccinia, adenovirus, fowl pox virus and pseudorabies. A variety of cloning and expression vectors for use with prokaryotic and eukaryotic hosts are described by Sambrook, *et al.*, Molecular Cloning: A Laboratory Manual, 2nd Ed., Cold Spring Harbor, N.Y., (1989).

Host cells and transformed cells

30 The invention also provides transformed cells comprising a nucleic acid sequence of the invention, e.g., a sequence encoding an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase of the invention, or a vector of the invention. The host cell may be any of the host cells familiar to those skilled in the art,

including prokaryotic cells, eukaryotic cells, such as bacterial cells, fungal cells, yeast cells, mammalian cells, insect cells, or plant cells. Exemplary bacterial cells include any species within the genera *Escherichia*, *Bacillus*, *Streptomyces*, *Salmonella*, *Pseudomonas* and *Staphylococcus*, including, e.g., *Escherichia coli*, *Lactococcus lactis*, *Bacillus subtilis*,
5 *Bacillus cereus*, *Salmonella typhimurium*, *Pseudomonas fluorescens*. Exemplary fungal cells include any species of *Aspergillus*. Exemplary yeast cells include any species of *Pichia*, *Saccharomyces*, *Schizosaccharomyces*, or *Schwanniomyces*, including *Pichia pastoris*, *Saccharomyces cerevisiae*, or *Schizosaccharomyces pombe*. Exemplary insect cells include any species of *Spodoptera* or *Drosophila*, including *Drosophila S2* and
10 *Spodoptera Sf9*. Exemplary animal cells include CHO, COS or Bowes melanoma or any mouse or human cell line. The selection of an appropriate host is within the abilities of those skilled in the art. Techniques for transforming a wide variety of higher plant species are well known and described in the technical and scientific literature. See, e.g., Weising (1988) Ann. Rev. Genet. 22:421-477; U.S. Patent No. 5,750,870.

15 The vector can be introduced into the host cells using any of a variety of techniques, including transformation, transfection, transduction, viral infection, gene guns, or Ti-mediated gene transfer. Particular methods include calcium phosphate transfection, DEAE-Dextran mediated transfection, lipofection, or electroporation (Davis, L., Dibner, M., Battey, I., Basic Methods in Molecular Biology, (1986)).

20 In one aspect, the nucleic acids or vectors of the invention are introduced into the cells for screening, thus, the nucleic acids enter the cells in a manner suitable for subsequent expression of the nucleic acid. The method of introduction is largely dictated by the targeted cell type. Exemplary methods include CaPO₄ precipitation, liposome fusion, lipofection (e.g., LIPOFECTIN™), electroporation, viral infection, etc. The
25 candidate nucleic acids may stably integrate into the genome of the host cell (for example, with retroviral introduction) or may exist either transiently or stably in the cytoplasm (i.e. through the use of traditional plasmids, utilizing standard regulatory sequences, selection markers, etc.). As many pharmaceutically important screens require human or model mammalian cell targets, retroviral vectors capable of transfecting such targets are can be
30 used.

Where appropriate, the engineered host cells can be cultured in conventional nutrient media modified as appropriate for activating promoters, selecting transformants or amplifying the genes of the invention. Following transformation of a suitable host strain and growth of the host strain to an appropriate cell density, the selected promoter may be

induced by appropriate means (e.g., temperature shift or chemical induction) and the cells may be cultured for an additional period to allow them to produce the desired polypeptide or fragment thereof.

Cells can be harvested by centrifugation, disrupted by physical or chemical means, and the resulting crude extract is retained for further purification. Microbial cells employed for expression of proteins can be disrupted by any convenient method, including freeze-thaw cycling, sonication, mechanical disruption, or use of cell lysing agents. Such methods are well known to those skilled in the art. The expressed polypeptide or fragment thereof can be recovered and purified from recombinant cell cultures by methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Protein refolding steps can be used, as necessary, in completing configuration of the polypeptide. If desired, high performance liquid chromatography (HPLC) can be employed for final purification steps.

The constructs in host cells can be used in a conventional manner to produce the gene product encoded by the recombinant sequence. Depending upon the host employed in a recombinant production procedure, the polypeptides produced by host cells containing the vector may be glycosylated or may be non-glycosylated. Polypeptides of the invention may or may not also include an initial methionine amino acid residue.

Cell-free translation systems can also be employed to produce a polypeptide of the invention. Cell-free translation systems can use mRNAs transcribed from a DNA construct comprising a promoter operably linked to a nucleic acid encoding the polypeptide or fragment thereof. In some aspects, the DNA construct may be linearized prior to conducting an in vitro transcription reaction. The transcribed mRNA is then incubated with an appropriate cell-free translation extract, such as a rabbit reticulocyte extract, to produce the desired polypeptide or fragment thereof.

The expression vectors can contain one or more selectable marker genes to provide a phenotypic trait for selection of transformed host cells such as dihydrofolate reductase or neomycin resistance for eukaryotic cell culture, or such as tetracycline or ampicillin resistance in *E. coli*.

Host cells containing the polynucleotides of interest, e.g., nucleic acids of the invention, can be cultured in conventional nutrient media modified as appropriate for activating promoters, selecting transformants or amplifying genes. The culture conditions,

such as temperature, pH and the like, are those previously used with the host cell selected for expression and will be apparent to the ordinarily skilled artisan. The clones which are identified as having the specified enzyme activity may then be sequenced to identify the polynucleotide sequence encoding an enzyme having the enhanced activity.

5 The invention provides a method for overexpressing a recombinant isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase in a cell comprising expressing a vector comprising a nucleic acid of the invention, e.g., a nucleic acid comprising a nucleic acid sequence with at least about 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%,
10 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more sequence identity to a sequence of the invention over a region of at least about 100 residues, wherein the sequence identities are determined by analysis with a sequence comparison algorithm or by visual inspection, or, a nucleic acid that hybridizes under
15 stringent conditions to a nucleic acid sequence of the invention, or a subsequence thereof. The overexpression can be effected by any means, e.g., use of a high activity promoter, a dicistronic vector or by gene amplification of the vector.

 The nucleic acids of the invention can be expressed, or overexpressed, in any in vitro or in vivo expression system. Any cell culture systems can be employed to express,
20 or over-express, recombinant protein, including bacterial, insect, yeast, fungal or mammalian cultures. Over-expression can be effected by appropriate choice of promoters, enhancers, vectors (e.g., use of replicon vectors, dicistronic vectors (see, e.g., Gurtu (1996) *Biochem. Biophys. Res. Commun.* 229:295-8), media, culture systems and the like. In one aspect, gene amplification using selection markers, e.g., glutamine synthetase (see,
25 e.g., Sanders (1987) *Dev. Biol. Stand.* 66:55-63), in cell systems are used to overexpress the polypeptides of the invention.

 Additional details regarding this approach are in the public literature and/or are known to the skilled artisan. In a particular non-limiting exemplification, such publicly available literature includes, for example: *Appl Environ Microbiol.* 2004 Jun;70(6):3298-
30 304; *Biotechnol Bioeng.* 2007 Nov 1;98(4):812-24 and *FEMS Microbiol Lett.* 2001 Mar 15;196(2):93-8, although these references do not teach the inventive enzymes of the instant application.

The host cell may be any of the host cells familiar to those skilled in the art, including prokaryotic cells, eukaryotic cells, mammalian cells, insect cells, or plant cells. As representative examples of appropriate hosts, there may be mentioned: bacterial cells, such as *E. coli*, *Streptomyces*, *Bacillus subtilis*, *Bacillus cereus*, *Salmonella typhimurium*
5 and various species within the genera *Pseudomonas*, *Streptomyces* and *Staphylococcus*, fungal cells, such as *Aspergillus*, yeast such as any species of *Pichia*, *Saccharomyces*, *Schizosaccharomyces*, *Schwanniomyces*, including *Pichia pastoris*, *Saccharomyces cerevisiae*, or *Schizosaccharomyces pombe*, insect cells such as *Drosophila S2* and *Spodoptera Sf9*, animal cells such as CHO, COS or Bowes melanoma and adenoviruses.

10 The selection of an appropriate host is within the abilities of those skilled in the art.

The vector may be introduced into the host cells using any of a variety of techniques, including transformation, transfection, transduction, viral infection, gene guns, or Ti-mediated gene transfer. Particular methods include calcium phosphate transfection, DEAE-Dextran mediated transfection, lipofection, or electroporation (Davis, L., Dibner,
15 M., Battey, I., Basic Methods in Molecular Biology, (1986)).

Where appropriate, the engineered host cells can be cultured in conventional nutrient media modified as appropriate for activating promoters, selecting transformants or amplifying the genes of the invention. Following transformation of a suitable host strain and growth of the host strain to an appropriate cell density, the selected promoter may be
20 induced by appropriate means (e.g., temperature shift or chemical induction) and the cells may be cultured for an additional period to allow them to produce the desired polypeptide or fragment thereof.

Cells are typically harvested by centrifugation, disrupted by physical or chemical means and the resulting crude extract is retained for further purification. Microbial cells
25 employed for expression of proteins can be disrupted by any convenient method, including freeze-thaw cycling, sonication, mechanical disruption, or use of cell lysing agents. Such methods are well known to those skilled in the art. The expressed polypeptide or fragment thereof can be recovered and purified from recombinant cell cultures by methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation
30 exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Protein refolding steps can be used, as necessary, in completing configuration of the polypeptide. If desired, high performance liquid chromatography (HPLC) can be employed for final purification steps.

Various mammalian cell culture systems can also be employed to express recombinant protein. Examples of mammalian expression systems include the COS-7 lines of monkey kidney fibroblasts (described by Gluzman, *Cell*, 23:175, 1981) and other cell lines capable of expressing proteins from a compatible vector, such as the C127, 3T3, CHO, HeLa and BHK cell lines.

The constructs in host cells can be used in a conventional manner to produce the gene product encoded by the recombinant sequence. Depending upon the host employed in a recombinant production procedure, the polypeptides produced by host cells containing the vector may be glycosylated or may be non-glycosylated. Polypeptides of the invention may or may not also include an initial methionine amino acid residue.

Alternatively, the polypeptides of amino acid sequences of the invention, or fragments comprising at least 5, 10, 15, 20, 25, 30, 35, 40, 50, 75, 100, or 150 consecutive amino acids thereof can be synthetically produced by conventional peptide synthesizers. In other aspects, fragments or portions of the polypeptides may be employed for producing the corresponding full-length polypeptide by peptide synthesis; therefore, the fragments may be employed as intermediates for producing the full-length polypeptides.

Cell-free translation systems can also be employed to produce one of the polypeptides of amino acid sequences of the invention, or fragments comprising at least 5, 10, 15, 20, 25, 30, 35, 40, 50, 75, 100, or 150 consecutive amino acids thereof using mRNAs transcribed from a DNA construct comprising a promoter operably linked to a nucleic acid encoding the polypeptide or fragment thereof. In some aspects, the DNA construct may be linearized prior to conducting an *in vitro* transcription reaction. The transcribed mRNA is then incubated with an appropriate cell-free translation extract, such as a rabbit reticulocyte extract, to produce the desired polypeptide or fragment thereof.

Amplification of Nucleic Acids

In practicing the invention, nucleic acids of the invention and nucleic acids encoding the isomerases, e.g., racemases, e.g., amino acid racemases, alanine racemases, and/or epimerases of the invention, or modified nucleic acids of the invention, can be reproduced by amplification. Amplification can also be used to clone or modify the nucleic acids of the invention. Thus, the invention provides amplification primer sequence pairs for amplifying nucleic acids of the invention. One of skill in the art can design amplification primer sequence pairs for any part of or the full length of these sequences.

In one aspect, the invention provides a nucleic acid amplified by a primer pair of the invention, e.g., a primer pair as set forth by about the first (the 5') 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25 residues of a nucleic acid of the invention, and about the first (the 5') 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25 residues of the complementary
5 strand.

The invention provides an amplification primer sequence pair for amplifying a nucleic acid encoding a polypeptide having an isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase activity, wherein the primer pair is capable of amplifying a nucleic acid comprising a sequence of the invention, or fragments or
10 subsequences thereof. One or each member of the amplification primer sequence pair can comprise an oligonucleotide comprising at least about 10 to 50 consecutive bases of the sequence, or about 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25 consecutive bases of the sequence. The invention provides amplification primer pairs, wherein the primer pair comprises a first member having a sequence as set forth by about the first (the
15 5') 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25 residues of a nucleic acid of the invention, and a second member having a sequence as set forth by about the first (the 5') 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25 residues of the complementary strand of the first member. The invention provides isomerases, e.g., racemases, e.g., amino acid racemases, alanine racemases, and/or epimerases generated by amplification,
20 e.g., polymerase chain reaction (PCR), using an amplification primer pair of the invention. The invention provides methods of making an isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase by amplification, e.g., polymerase chain reaction (PCR), using an amplification primer pair of the invention. In one aspect, the amplification primer pair amplifies a nucleic acid from a library, e.g., a gene library, such
25 as an environmental library.

Amplification reactions can also be used to quantify the amount of nucleic acid in a sample (such as the amount of message in a cell sample), label the nucleic acid (e.g., to apply it to an array or a blot), detect the nucleic acid, or quantify the amount of a specific nucleic acid in a sample. In one aspect of the invention, message isolated from a cell or a
30 cDNA library are amplified.

The skilled artisan can select and design suitable oligonucleotide amplification primers. Amplification methods are also well known in the art, and include, e.g., polymerase chain reaction, PCR (see, e.g., PCR PROTOCOLS, A GUIDE TO METHODS AND APPLICATIONS, ed. Innis, Academic Press, N.Y. (1990) and PCR STRATEGIES

(1995), ed. Innis, Academic Press, Inc., N.Y., ligase chain reaction (LCR) (see, e.g., Wu (1989) Genomics 4:560; Landegren (1988) Science 241:1077; Barringer (1990) Gene 89:117); transcription amplification (see, e.g., Kwoh (1989) Proc. Natl. Acad. Sci. USA 86:1173); and, self-sustained sequence replication (see, e.g., Guatelli (1990) Proc. Natl. Acad. Sci. USA 87:1874); Q Beta replicase amplification (see, e.g., Smith (1997) J. Clin. Microbiol. 35:1477-1491), automated Q-beta replicase amplification assay (see, e.g., Burg (1996) Mol. Cell. Probes 10:257-271) and other RNA polymerase mediated techniques (e.g., NASBA, Cangene, Mississauga, Ontario); see also Berger (1987) Methods Enzymol. 152:307-316; Sambrook; Ausubel; U.S. Patent Nos. 4,683,195 and 4,683,202; Sooknanan (1995) Biotechnology 13:563-564.

Determining the degree of sequence identity

The invention provides nucleic acids comprising sequences having at least about 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more, or complete (100%) sequence identity to an exemplary nucleic acid of the invention (as defined above) over a region of at least about 50, 75, 100, 150, 200, 250, 300, 350, 400, 450, 500, 550, 600, 650, 700, 750, 800, 850, 900, 950, 1000, 1050, 1100, 1150, 1200, 1250, 1300, 1350, 1400, 1450, 1500, 1550 or more, residues. The invention provides polypeptides comprising sequences having at least about 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more, or complete (100%) sequence identity to an exemplary polypeptide of the invention. The extent of sequence identity (homology) may be determined using any computer program and associated parameters, including those described herein, such as BLAST 2.2.2. or FASTA version 3.0t78, with the default parameters.

As used herein, the terms “computer,” “computer program” and “processor” are used in their broadest general contexts and incorporate all such devices, as described in detail, below. A “coding sequence of” or a “sequence encodes” a particular polypeptide or protein, is a nucleic acid sequence which is transcribed and translated into a polypeptide or protein when placed under the control of appropriate regulatory sequences.

The phrase “substantially identical” in the context of two nucleic acids or polypeptides, refers to two or more sequences that have, e.g., at least about 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 5 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more nucleotide or amino acid residue (sequence) identity, when compared and aligned for maximum correspondence, as measured using one of the known sequence comparison algorithms or by visual inspection. Typically, the substantial identity exists over a region of at least about 100 residues and most commonly the 10 sequences are substantially identical over at least about 150-200 residues. In some aspects, the sequences are substantially identical over the entire length of the coding regions.

Additionally a “substantially identical” amino acid sequence is a sequence that differs from a reference sequence by one or more conservative or non-conservative 15 amino acid substitutions, deletions, or insertions, particularly when such a substitution occurs at a site that is not the active site of the molecule and provided that the polypeptide essentially retains its functional properties. A conservative amino acid substitution, for example, substitutes one amino acid for another of the same class (*e.g.*, substitution of one hydrophobic amino acid, such as isoleucine, valine, leucine, or methionine, for another, or 20 substitution of one polar amino acid for another, such as substitution of arginine for lysine, glutamic acid for aspartic acid or glutamine for asparagine). One or more amino acids can be deleted, for example, from an isomerase, *e.g.*, racemase, *e.g.*, amino acid racemase, alanine racemase, and/or epimerase polypeptide, resulting in modification of the structure of the polypeptide, without significantly altering its biological activity. For example, 25 amino- or carboxyl-terminal amino acids that are not required for isomerase, *e.g.*, racemase, *e.g.*, amino acid racemase, alanine racemase, and/or epimerase biological activity can be removed. Modified polypeptide sequences of the invention can be assayed for isomerase, *e.g.*, racemase, *e.g.*, amino acid racemase, alanine racemase, and/or epimerase biological activity by any number of methods, including contacting the 30 modified polypeptide sequence with an isomerase, *e.g.*, racemase, *e.g.*, amino acid racemase, alanine racemase, and/or epimerase substrate and determining whether the modified polypeptide decreases the amount of specific substrate in the assay or increases the bioproducts of the enzymatic reaction of a functional isomerase, *e.g.*, racemase, *e.g.*, amino acid racemase, alanine racemase, and/or epimerase polypeptide with the substrate.

Nucleic acid sequences of the invention can comprise at least 10, 15, 20, 25, 30, 35, 40, 50, 75, 100, 150, 200, 300, 400, or 500 consecutive nucleotides of an exemplary sequence of the invention and sequences substantially identical thereto.

Nucleic acid sequences of the invention can comprise homologous sequences and
5 fragments of nucleic acid sequences and sequences substantially identical thereto, refer to a sequence having at least 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more sequence identity (homology)
10 to these sequences. Homology may be determined using any of the computer programs and parameters described herein, including FASTA version 3.0t78 with the default parameters. Homologous sequences also include RNA sequences in which uridines replace the thymines in the nucleic acid sequences of the invention. The homologous sequences may be obtained using any of the procedures described herein or may result
15 from the correction of a sequencing error. It will be appreciated that the nucleic acid sequences of the invention and sequences substantially identical thereto, can be represented in the traditional single character format (See the inside back cover of Stryer, Lubert. Biochemistry, 3rd Ed., W. H Freeman & Co., New York.) or in any other format which records the identity of the nucleotides in a sequence.

20 Various sequence comparison programs identified elsewhere in this patent specification are particularly contemplated for use in this aspect of the invention. Protein and/or nucleic acid sequence homologies may be evaluated using any of the variety of sequence comparison algorithms and programs known in the art. Such algorithms and programs include, but are by no means limited to, TBLASTN, BLASTP, FASTA,
25 TFASTA and CLUSTALW (Pearson and Lipman, Proc. Natl. Acad. Sci. USA 85(8):2444-2448, 1988; Altschul *et al.*, J. Mol. Biol. 215(3):403-410, 1990; Thompson *et al.*, Nucleic Acids Res. 22(2):4673-4680, 1994; Higgins *et al.*, Methods Enzymol. 266:383-402, 1996; Altschul *et al.*, J. Mol. Biol. 215(3):403-410, 1990; Altschul *et al.*, Nature Genetics 3:266-272, 1993).

30 Homology or identity is often measured using sequence analysis software (*e.g.*, Sequence Analysis Software Package of the Genetics Computer Group, University of Wisconsin Biotechnology Center, 1710 University Avenue, Madison, WI 53705). Such software matches similar sequences by assigning degrees of homology to various deletions, substitutions and other modifications. The terms "homology" and "identity" in the context of

two or more nucleic acids or polypeptide sequences, refer to two or more sequences or subsequences that are the same or have a specified percentage of amino acid residues or nucleotides that are the same when compared and aligned for maximum correspondence over a comparison window or designated region as measured using any number of sequence
5 comparison algorithms or by manual alignment and visual inspection.

For sequence comparison, typically one sequence acts as a reference sequence, to which test sequences are compared. When using a sequence comparison algorithm, test and reference sequences are entered into a computer, subsequence coordinates are designated, if necessary and sequence algorithm program parameters are designated.

10 Default program parameters can be used, or alternative parameters can be designated. The sequence comparison algorithm then calculates the percent sequence identities for the test sequences relative to the reference sequence, based on the program parameters.

A “comparison window”, as used herein, includes reference to a segment of any one of the number of contiguous positions selected from the group consisting of from 20
15 to 600, usually about 50 to about 200, more usually about 100 to about 150 in which a sequence may be compared to a reference sequence of the same number of contiguous positions after the two sequences are optimally aligned. Methods of alignment of sequence for comparison are well-known in the art. Optimal alignment of sequences for comparison can be conducted, *e.g.*, by the local homology algorithm of Smith & Waterman, *Adv. Appl. Math.* 2:482, 1981, by the homology alignment algorithm of Needleman & Wunsch, *J. Mol. Biol.* 48:443, 1970, by the search for similarity method of person & Lipman, *Proc. Nat'l. Acad. Sci. USA* 85:2444, 1988, by computerized implementations of these algorithms (GAP, BESTFIT, FASTA and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, WI), or by manual alignment and visual
25 inspection. Other algorithms for determining homology or identity include, for example, in addition to a BLAST program (Basic Local Alignment Search Tool at the National Center for Biological Information), ALIGN, AMAS (Analysis of Multiply Aligned Sequences), AMPS (Protein Multiple Sequence Alignment), ASSET (Aligned Segment Statistical Evaluation Tool), BANDS, BESTSCOR, BIOSCAN (Biological Sequence Comparative
30 Analysis Node), BLIMPS (BLOCKS IMPROVED Searcher), FASTA, Intervals & Points, BMB, CLUSTAL V, CLUSTAL W, CONSENSUS, LCONSENSUS, WCONSENSUS, Smith-Waterman algorithm, DARWIN, Las Vegas algorithm, FNAT (Forced Nucleotide Alignment Tool), Framealign, Framesearch, DYNAMIC, FILTER, FSAP (Fristensky Sequence Analysis Package), GAP (Global Alignment Program), GENAL, GIBBS,

GenQuest, ISSC (Sensitive Sequence Comparison), LALIGN (Local Sequence Alignment), LCP (Local Content Program), MACAW (Multiple Alignment Construction & Analysis Workbench), MAP (Multiple Alignment Program), MBLKP, MBLKN, PIMA (Pattern-Induced Multi-sequence Alignment), SAGA (Sequence Alignment by Genetic Algorithm) and WHAT-IF. Such alignment programs can also be used to screen genome databases to identify polynucleotide sequences having substantially identical sequences. A number of genome databases are available, for example, a substantial portion of the human genome is available as part of the Human Genome Sequencing Project. At least twenty-one other genomes have already been sequenced, including, for example, *M. genitalium* (Fraser *et al.*, 1995), *M. jannaschii* (Bult *et al.*, 1996), *H. influenzae* (Fleischmann *et al.*, 1995), *E. coli* (Blattner *et al.*, 1997) and yeast (*S. cerevisiae*) (Mewes *et al.*, 1997) and *D. melanogaster* (Adams *et al.*, 2000). Significant progress has also been made in sequencing the genomes of model organism, such as mouse, *C. elegans* and *Arabidopsis sp.* Several databases containing genomic information annotated with some functional information are maintained by different organization and are accessible via the internet

One example of a useful algorithm is BLAST and BLAST 2.0 algorithms, which are described in Altschul *et al.*, Nuc. Acids Res. 25:3389-3402, 1997 and Altschul *et al.*, J. Mol. Biol. 215:403-410, 1990, respectively. Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information. This algorithm involves first identifying high scoring sequence pairs (HSPs) by identifying short words of length W in the query sequence, which either match or satisfy some positive-valued threshold score T when aligned with a word of the same length in a database sequence. T is referred to as the neighborhood word score threshold (Altschul *et al.*, *supra*). These initial neighborhood word hits act as seeds for initiating searches to find longer HSPs containing them. The word hits are extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Cumulative scores are calculated using, for nucleotide sequences, the parameters M (reward score for a pair of matching residues; always >0). For amino acid sequences, a scoring matrix is used to calculate the cumulative score. Extension of the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T and X determine the sensitivity and speed of the alignment. The BLASTN program (for nucleotide sequences) uses as defaults a

wordlength (W) of 11, an expectation (E) of 10, M=5, N=-4 and a comparison of both strands. For amino acid sequences, the BLASTP program uses as defaults a wordlength of 3 and expectations (E) of 10 and the BLOSUM62 scoring matrix (see Henikoff & Henikoff, Proc. Natl. Acad. Sci. USA 89:10915, 1989) alignments (B) of 50, expectation
5 (E) of 10, M=5, N= -4 and a comparison of both strands.

The BLAST algorithm also performs a statistical analysis of the similarity between two sequences (see, *e.g.*, Karlin & Altschul, Proc. Natl. Acad. Sci. USA 90:5873, 1993). One measure of similarity provided by BLAST algorithm is the smallest sum probability (P(N)), which provides an indication of the probability by which a match
10 between two nucleotide or amino acid sequences would occur by chance. For example, a nucleic acid is considered similar to a references sequence if the smallest sum probability in a comparison of the test nucleic acid to the reference nucleic acid is less than about 0.2, more preferably less than about 0.01 and most preferably less than about 0.001.

In one aspect, protein and nucleic acid sequence homologies are evaluated
15 using the Basic Local Alignment Search Tool ("BLAST") In particular, five specific BLAST programs are used to perform the following task:

- (1) BLASTP and BLAST3 compare an amino acid query sequence against a protein sequence database;
- (2) BLASTN compares a nucleotide query sequence against a
20 nucleotide sequence database;
- (3) BLASTX compares the six-frame conceptual translation products of a query nucleotide sequence (both strands) against a protein sequence database;
- (4) TBLASTN compares a query protein sequence against a nucleotide sequence database translated in all six reading frames (both strands); and
- (5) TBLASTX compares the six-frame translations of a nucleotide
25 query sequence against the six-frame translations of a nucleotide sequence database.

The BLAST programs identify homologous sequences by identifying similar segments, which are referred to herein as "high-scoring segment pairs," between a
30 query amino or nucleic acid sequence and a test sequence which is preferably obtained from a protein or nucleic acid sequence database. High-scoring segment pairs are preferably identified (*i.e.*, aligned) by means of a scoring matrix, many of which are known in the art. Preferably, the scoring matrix used is the BLOSUM62 matrix (Gonnet *et al.*, Science 256:1443-1445, 1992; Henikoff and Henikoff, Proteins 17:49-61, 1993).

Less preferably, the PAM or PAM250 matrices may also be used (see, *e.g.*, Schwartz and Dayhoff, eds., 1978, *Matrices for Detecting Distance Relationships: Atlas of Protein Sequence and Structure*, Washington: National Biomedical Research Foundation).

BLAST programs are accessible through the U.S. National Library of Medicine.

5 The parameters used with the above algorithms may be adapted depending on the sequence length and degree of homology studied. In some aspects, the parameters may be the default parameters used by the algorithms in the absence of instructions from the user.

Computer systems and computer program products

10 To determine and identify sequence identities, structural homologies, motifs and the like in silico, a nucleic acid or polypeptide sequence of the invention can be stored, recorded, and manipulated on any medium which can be read and accessed by a computer.

 Accordingly, the invention provides computers, computer systems, computer readable mediums, computer programs products and the like recorded or stored thereon the nucleic acid and polypeptide sequences of the invention. As used herein, the words “recorded” and “stored” refer to a process for storing information on a computer medium. A skilled artisan can readily adopt any known methods for recording information on a computer readable medium to generate manufactures comprising one or more of the nucleic acid and/or polypeptide sequences of the invention.

20 The polypeptides of the invention include the exemplary sequences of the invention, and sequences substantially identical thereto, and fragments of any of the preceding sequences. Substantially identical, or homologous, polypeptide sequences refer to a polypeptide sequence having at least 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%, 72%, 25 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more, or complete (100%) sequence identity to an exemplary sequence of the invention, *e.g.*, a polypeptide sequences of the invention.

30 Homology may be determined using any of the computer programs and parameters described herein, including FASTA version 3.0t78 with the default parameters or with any modified parameters. The homologous sequences may be obtained using any of the procedures described herein or may result from the correction of a sequencing error. The polypeptide fragments comprise at least about 10, 15, 20, 25, 30, 35, 40, 45, 50, 75, 100, 150,

200, 250, 300, 350, 400, 450, 500 or more consecutive amino acids of the polypeptides of the invention and sequences substantially identical thereto. It will be appreciated that the polypeptide codes of amino acid sequences of the invention and sequences substantially identical thereto, can be represented in the traditional single character format or three letter
5 format (See Stryer, Lubert. Biochemistry, 3rd Ed., supra) or in any other format which relates the identity of the polypeptides in a sequence.

A nucleic acid or polypeptide sequence of the invention can be stored, recorded and manipulated on any medium which can be read and accessed by a computer. As used herein, the words "recorded" and "stored" refer to a process for storing information
10 on a computer medium. A skilled artisan can readily adopt any of the presently known methods for recording information on a computer readable medium to generate manufactures comprising one or more of the nucleic acid sequences of the invention and sequences substantially identical thereto, one or more of the polypeptide sequences of the invention and sequences substantially identical thereto. Another aspect of the invention is a computer
15 readable medium having recorded thereon at least 2, 5, 10, 15, or 20 or more nucleic acid sequences of the invention and sequences substantially identical thereto.

Another aspect of the invention is a computer readable medium having recorded thereon one or more of the nucleic acid sequences of the invention and sequences substantially identical thereto. Another aspect of the invention is a computer readable
20 medium having recorded thereon one or more of the polypeptide sequences of the invention and sequences substantially identical thereto. Another aspect of the invention is a computer readable medium having recorded thereon at least 2, 5, 10, 15, or 20 or more of the sequences as set forth above.

Computer readable media include magnetically readable media, optically
25 readable media, electronically readable media and magnetic/optical media. For example, the computer readable media may be a hard disk, a floppy disk, a magnetic tape, CD-ROM, Digital Versatile Disk (DVD), Random Access Memory (RAM), or Read Only Memory (ROM) as well as other types of other media known to those skilled in the art.

Aspects of the invention include systems (*e.g.*, internet based systems),
30 particularly computer systems which store and manipulate the sequence information described herein. One example of a computer system 100 is illustrated in block diagram form in Figure 1. As used herein, "a computer system" refers to the hardware components, software components and data storage components used to analyze a nucleotide sequence of a nucleic acid sequence of the invention and sequences substantially identical thereto, or a

polypeptide sequence as set forth in the amino acid sequences of the invention. The computer system 100 typically includes a processor for processing, accessing and manipulating the sequence data. The processor 105 can be any well-known type of central processing unit, such as, for example, the Pentium III from Intel Corporation, or similar
5 processor from Sun, Motorola, Compaq, AMD or International Business Machines.

Typically the computer system 100 is a general purpose system that comprises the processor 105 and one or more internal data storage components 110 for storing data and one or more data retrieving devices for retrieving the data stored on the data storage components. A skilled artisan can readily appreciate that any one of the currently
10 available computer systems are suitable.

In one particular aspect, the computer system 100 includes a processor 105 connected to a bus which is connected to a main memory 115 (preferably implemented as RAM) and one or more internal data storage devices 110, such as a hard drive and/or other computer readable media having data recorded thereon. In some aspects, the computer
15 system 100 further includes one or more data retrieving device 118 for reading the data stored on the internal data storage devices 110.

The data retrieving device 118 may represent, for example, a floppy disk drive, a compact disk drive, a magnetic tape drive, or a modem capable of connection to a remote data storage system (*e.g.*, via the internet) etc. In some aspects, the internal data
20 storage device 110 is a removable computer readable medium such as a floppy disk, a compact disk, a magnetic tape, etc. containing control logic and/or data recorded thereon. The computer system 100 may advantageously include or be programmed by appropriate software for reading the control logic and/or the data from the data storage component once inserted in the data retrieving device.

25 The computer system 100 includes a display 120 which is used to display output to a computer user. It should also be noted that the computer system 100 can be linked to other computer systems 125a-c in a network or wide area network to provide centralized access to the computer system 100.

Software for accessing and processing the nucleotide sequences of a nucleic
30 acid sequence of the invention and sequences substantially identical thereto, or a polypeptide sequence of the invention and sequences substantially identical thereto, (such as search tools, compare tools and modeling tools etc.) may reside in main memory 115 during execution.

In some aspects, the computer system 100 may further comprise a sequence comparison algorithm for comparing a nucleic acid sequence of the invention and sequences substantially identical thereto, or a polypeptide sequence of the invention and sequences substantially identical thereto, stored on a computer readable medium to a reference nucleotide or polypeptide sequence(s) stored on a computer readable medium. A “sequence comparison algorithm” refers to one or more programs which are implemented (locally or remotely) on the computer system 100 to compare a nucleotide sequence with other nucleotide sequences and/or compounds stored within a data storage means. For example, the sequence comparison algorithm may compare the nucleotide sequences of a nucleic acid sequence of the invention and sequences substantially identical thereto, or a polypeptide sequence of the invention and sequences substantially identical thereto, stored on a computer readable medium to reference sequences stored on a computer readable medium to identify homologies or structural motifs.

Figure 2 is a flow diagram illustrating one aspect of a process 200 for comparing a new nucleotide or protein sequence with a database of sequences in order to determine the homology levels between the new sequence and the sequences in the database. The database of sequences can be a private database stored within the computer system 100, or a public database such as GENBANK that is available through the Internet.

The process 200 begins at a start state 201 and then moves to a state 202 wherein the new sequence to be compared is stored to a memory in a computer system 100. As discussed above, the memory could be any type of memory, including RAM or an internal storage device.

The process 200 then moves to a state 204 wherein a database of sequences is opened for analysis and comparison. The process 200 then moves to a state 206 wherein the first sequence stored in the database is read into a memory on the computer. A comparison is then performed at a state 210 to determine if the first sequence is the same as the second sequence. It is important to note that this step is not limited to performing an exact comparison between the new sequence and the first sequence in the database. Well-known methods are known to those of skill in the art for comparing two nucleotide or protein sequences, even if they are not identical. For example, gaps can be introduced into one sequence in order to raise the homology level between the two tested sequences. The parameters that control whether gaps or other features are introduced into a sequence during comparison are normally entered by the user of the computer system.

Once a comparison of the two sequences has been performed at the state 210, a determination is made at a decision state 210 whether the two sequences are the same. Of course, the term "same" is not limited to sequences that are absolutely identical. Sequences that are within the homology parameters entered by the user will be marked as "same" in the
5 process 200.

If a determination is made that the two sequences are the same, the process 200 moves to a state 214 wherein the name of the sequence from the database is displayed to the user. This state notifies the user that the sequence with the displayed name fulfills the homology constraints that were entered. Once the name of the stored sequence is displayed
10 to the user, the process 200 moves to a decision state 218 wherein a determination is made whether more sequences exist in the database. If no more sequences exist in the database, then the process 200 terminates at an end state 220. However, if more sequences do exist in the database, then the process 200 moves to a state 224 wherein a pointer is moved to the next sequence in the database so that it can be compared to the new sequence. In this
15 manner, the new sequence is aligned and compared with every sequence in the database.

It should be noted that if a determination had been made at the decision state 212 that the sequences were not homologous, then the process 200 would move immediately to the decision state 218 in order to determine if any other sequences were available in the database for comparison.

Accordingly, one aspect of the invention is a computer system comprising a processor, a data storage device having stored thereon a nucleic acid sequence of the invention and sequences substantially identical thereto, or a polypeptide sequence of the invention and sequences substantially identical thereto, a data storage device having
20 retrievably stored thereon reference nucleotide sequences or polypeptide sequences to be compared to a nucleic acid sequence of the invention and sequences substantially identical thereto, or a polypeptide sequence of the invention and sequences substantially identical thereto and a sequence comparer for conducting the comparison. The sequence comparer may indicate a homology level between the sequences compared or identify structural motifs in the above described nucleic acid code of nucleic acid sequences of the invention
25 and sequences substantially identical thereto, or a polypeptide sequence of the invention and sequences substantially identical thereto, or it may identify structural motifs in sequences which are compared to these nucleic acid codes and polypeptide codes. In some aspects, the data storage device may have stored thereon the sequences of at least 2, 5, 10, 15, 20, 25, 30 or 40 or more of the nucleic acid sequences of the invention and sequences
30

substantially identical thereto, or the polypeptide sequences of the invention and sequences substantially identical thereto.

Another aspect of the invention is a method for determining the level of homology between a nucleic acid sequence of the invention and sequences substantially identical thereto, or a polypeptide sequence of the invention and sequences substantially identical thereto and a reference nucleotide sequence. The method including reading the nucleic acid code or the polypeptide code and the reference nucleotide or polypeptide sequence through the use of a computer program which determines homology levels and determining homology between the nucleic acid code or polypeptide code and the reference nucleotide or polypeptide sequence with the computer program. The computer program may be any of a number of computer programs for determining homology levels, including those specifically enumerated herein, (*e.g.*, BLAST2N with the default parameters or with any modified parameters). The method may be implemented using the computer systems described above. The method may also be performed by reading at least 2, 5, 10, 15, 20, 25, 30 or 40 or more of the above described nucleic acid sequences of the invention, or the polypeptide sequences of the invention through use of the computer program and determining homology between the nucleic acid codes or polypeptide codes and reference nucleotide sequences or polypeptide sequences.

Figure 3 is a flow diagram illustrating one aspect of a process 250 in a computer for determining whether two sequences are homologous. The process 250 begins at a start state 252 and then moves to a state 254 wherein a first sequence to be compared is stored to a memory. The second sequence to be compared is then stored to a memory at a state 256. The process 250 then moves to a state 260 wherein the first character in the first sequence is read and then to a state 262 wherein the first character of the second sequence is read. It should be understood that if the sequence is a nucleotide sequence, then the character would normally be either A, T, C, G or U. If the sequence is a protein sequence, then it is preferably in the single letter amino acid code so that the first and sequence sequences can be easily compared.

A determination is then made at a decision state 264 whether the two characters are the same. If they are the same, then the process 250 moves to a state 268 wherein the next characters in the first and second sequences are read. A determination is then made whether the next characters are the same. If they are, then the process 250 continues this loop until two characters are not the same. If a determination is made that

the next two characters are not the same, the process 250 moves to a decision state 274 to determine whether there are any more characters either sequence to read.

If there are not any more characters to read, then the process 250 moves to a state 276 wherein the level of homology between the first and second sequences is displayed to the user. The level of homology is determined by calculating the proportion of characters between the sequences that were the same out of the total number of sequences in the first sequence. Thus, if every character in a first 100 nucleotide sequence aligned with a every character in a second sequence, the homology level would be 100%.

Alternatively, the computer program may be a computer program which compares the nucleotide sequences of a nucleic acid sequence as set forth in the invention, to one or more reference nucleotide sequences in order to determine whether the nucleic acid code of a nucleic acid sequence of the invention and sequences substantially identical thereto, differs from a reference nucleic acid sequence at one or more positions. In one aspect such a program records the length and identity of inserted, deleted or substituted nucleotides with respect to the sequence of either the reference polynucleotide or a nucleic acid sequence of the invention and sequences substantially identical thereto. In one aspect, the computer program may be a program which determines whether a nucleic acid sequence of the invention and sequences substantially identical thereto, contains a single nucleotide polymorphism (SNP) with respect to a reference nucleotide sequence.

Another aspect of the invention is a method for determining whether a nucleic acid sequence of the invention and sequences substantially identical thereto, differs at one or more nucleotides from a reference nucleotide sequence comprising the steps of reading the nucleic acid code and the reference nucleotide sequence through use of a computer program which identifies differences between nucleic acid sequences and identifying differences between the nucleic acid code and the reference nucleotide sequence with the computer program. In some aspects, the computer program is a program which identifies single nucleotide polymorphisms. The method may be implemented by the computer systems described above and the method illustrated in Figure 3. The method may also be performed by reading at least 2, 5, 10, 15, 20, 25, 30, or 40 or more of the nucleic acid sequences of the invention and sequences substantially identical thereto and the reference nucleotide sequences through the use of the computer program and identifying differences between the nucleic acid codes and the reference nucleotide sequences with the computer program.

In other aspects the computer based system may further comprise an identifier for identifying features within a nucleic acid sequence of the invention or a polypeptide sequence of the invention and sequences substantially identical thereto.

An “identifier” refers to one or more programs which identifies certain features within a nucleic acid sequence of the invention and sequences substantially identical thereto, or a polypeptide sequence of the invention and sequences substantially identical thereto. In one aspect, the identifier may comprise a program which identifies an open reading frame in a nucleic acid sequence of the invention and sequences substantially identical thereto.

Figure 4 is a flow diagram illustrating one aspect of an identifier process 300 for detecting the presence of a feature in a sequence. The process 300 begins at a start state 302 and then moves to a state 304 wherein a first sequence that is to be checked for features is stored to a memory 115 in the computer system 100. The process 300 then moves to a state 306 wherein a database of sequence features is opened. Such a database would include a list of each feature’s attributes along with the name of the feature. For example, a feature name could be “Initiation Codon” and the attribute would be “ATG”. Another example would be the feature name “TAATAA Box” and the feature attribute would be “TAATAA”. An example of such a database is produced by the University of Wisconsin Genetics Computer Group. Alternatively, the features may be structural polypeptide motifs such as alpha helices, beta sheets, or functional polypeptide motifs such as enzymatic active sites, helix-turn-helix motifs or other motifs known to those skilled in the art.

Once the database of features is opened at the state 306, the process 300 moves to a state 308 wherein the first feature is read from the database. A comparison of the attribute of the first feature with the first sequence is then made at a state 310. A determination is then made at a decision state 316 whether the attribute of the feature was found in the first sequence. If the attribute was found, then the process 300 moves to a state 318 wherein the name of the found feature is displayed to the user.

The process 300 then moves to a decision state 320 wherein a determination is made whether more features exist in the database. If no more features do exist, then the process 300 terminates at an end state 324. However, if more features do exist in the database, then the process 300 reads the next sequence feature at a state 326 and loops back to the state 310 wherein the attribute of the next feature is compared against the first sequence. It should be noted, that if the feature attribute is not found in

the first sequence at the decision state 316, the process 300 moves directly to the decision state 320 in order to determine if any more features exist in the database.

Accordingly, another aspect of the invention is a method of identifying a feature within a nucleic acid sequence of the invention and sequences substantially identical thereto, or a polypeptide sequence of the invention and sequences substantially identical thereto, comprising reading the nucleic acid code(s) or polypeptide code(s) through the use of a computer program which identifies features therein and identifying features within the nucleic acid code(s) with the computer program. In one aspect, computer program comprises a computer program which identifies open reading frames. The method may be performed by reading a single sequence or at least 2, 5, 10, 15, 20, 25, 30, or 40 of the nucleic acid sequences of the invention and sequences substantially identical thereto, or the polypeptide sequences of the invention and sequences substantially identical thereto, through the use of the computer program and identifying features within the nucleic acid codes or polypeptide codes with the computer program.

A nucleic acid sequence of the invention and sequences substantially identical thereto, or a polypeptide sequence of the invention and sequences substantially identical thereto, may be stored and manipulated in a variety of data processor programs in a variety of formats. For example, a nucleic acid sequence of the invention and sequences substantially identical thereto, or a polypeptide sequence of the invention and sequences substantially identical thereto, may be stored as text in a word processing file, such as Microsoft WORD™ or WORDPERFECT™ or as an ASCII file in a variety of database programs familiar to those of skill in the art, such as DB2™, SYBASE™, or ORACLE™. In addition, many computer programs and databases may be used as sequence comparison algorithms, identifiers, or sources of reference nucleotide sequences or polypeptide sequences to be compared to a nucleic acid sequence of the invention and sequences substantially identical thereto, or a polypeptide sequence of the invention and sequences substantially identical thereto. The following list is intended not to limit the invention but to provide guidance to programs and databases which are useful with the nucleic acid sequences of the invention and sequences substantially identical thereto, or the polypeptide sequences of the invention and sequences substantially identical thereto.

The programs and databases which may be used include, but are not limited to: MacPattern (EMBL), DiscoveryBase (Molecular Applications Group), GeneMine (Molecular Applications Group), Look (Molecular Applications Group), MacLook (Molecular Applications Group), BLAST and BLAST2 (NCBI), BLASTN and BLASTX (Altschul et al,

J. Mol. Biol. 215: 403, 1990), FASTA (Pearson and Lipman, Proc. Natl. Acad. Sci. USA, 85: 2444, 1988), FASTDB (Brutlag *et al.* Comp. App. Biosci. 6:237-245, 1990), Catalyst (Molecular Simulations Inc.), Catalyst/SHAPE (Molecular Simulations Inc.), Cerius².DBAccess (Molecular Simulations Inc.), HypoGen (Molecular Simulations Inc.),
5 Insight II, (Molecular Simulations Inc.), Discover (Molecular Simulations Inc.), CHARMM (Molecular Simulations Inc.), Felix (Molecular Simulations Inc.), DelPhi, (Molecular Simulations Inc.), QuanteMM, (Molecular Simulations Inc.), Homology (Molecular Simulations Inc.), Modeler (Molecular Simulations Inc.), ISIS (Molecular Simulations Inc.), Quanta/Protein Design (Molecular Simulations Inc.), WebLab (Molecular Simulations Inc.),
10 WebLab Diversity Explorer (Molecular Simulations Inc.), Gene Explorer (Molecular Simulations Inc.), SeqFold (Molecular Simulations Inc.), the MDL Available Chemicals Directory database, the MDL Drug Data Report data base, the Comprehensive Medicinal Chemistry database, Derwents's World Drug Index database, the BioByteMasterFile database, the Genbank database and the Genseqn database. Many other programs and data
15 bases would be apparent to one of skill in the art given the present disclosure.

Motifs which may be detected using the above programs include sequences encoding leucine zippers, helix-turn-helix motifs, glycosylation sites, ubiquitination sites, alpha helices and beta sheets, signal sequences encoding signal peptides which direct the secretion of the encoded proteins, sequences implicated in transcription regulation such as
20 homeoboxes, acidic stretches, enzymatic active sites, substrate binding sites and enzymatic cleavage sites.

Hybridization of nucleic acids

The invention provides isolated, synthetic or recombinant nucleic acids that hybridize under stringent conditions to an exemplary sequence of the invention. The
25 stringent conditions can be highly stringent conditions, medium stringent conditions and/or low stringent conditions, including the high and reduced stringency conditions described herein. In one aspect, it is the stringency of the wash conditions that set forth the conditions which determine whether a nucleic acid is within the scope of the invention, as discussed below.

30 In alternative aspects, nucleic acids of the invention as defined by their ability to hybridize under stringent conditions can be between about five residues and the full length of nucleic acid of the invention; e.g., they can be at least 5, 10, 15, 20, 25, 30, 35, 40, 50, 55, 60, 65, 70, 75, 80, 90, 100, 150, 200, 250, 300, 350, 400, 450, 500, 550, 600, 650, 700,

750, 800, 850, 900, 950, 1000, or more, residues in length. Nucleic acids shorter than full length are also included. These nucleic acids can be useful as, e.g., hybridization probes, labeling probes, PCR oligonucleotide probes, iRNA (single or double stranded), antisense or sequences encoding antibody binding peptides (epitopes), motifs, active sites and the like.

In one aspect, nucleic acids of the invention are defined by their ability to hybridize under high stringency comprising conditions of about 50% formamide at about 37°C to 42°C. In one aspect, nucleic acids of the invention are defined by their ability to hybridize under reduced stringency comprising conditions in about 35% to 25% formamide at about 30°C to 35°C.

Alternatively, nucleic acids of the invention are defined by their ability to hybridize under high stringency comprising conditions at 42°C in 50% formamide, 5X SSPE, 0.3% SDS, and a repetitive sequence blocking nucleic acid, such as cot-1 or salmon sperm DNA (e.g., 200 ug/ml sheared and denatured salmon sperm DNA). In one aspect, nucleic acids of the invention are defined by their ability to hybridize under reduced stringency conditions comprising 35% formamide at a reduced temperature of 35°C.

In nucleic acid hybridization reactions, the conditions used to achieve a particular level of stringency will vary, depending on the nature of the nucleic acids being hybridized. For example, the length, degree of complementarity, nucleotide sequence composition (e.g., GC v. AT content) and nucleic acid type (e.g., RNA v. DNA) of the hybridizing regions of the nucleic acids can be considered in selecting hybridization conditions. An additional consideration is whether one of the nucleic acids is immobilized, for example, on a filter.

Hybridization may be carried out under conditions of low stringency, moderate stringency or high stringency. As an example of nucleic acid hybridization, a polymer membrane containing immobilized denatured nucleic acids is first prehybridized for 30 minutes at 45°C in a solution consisting of 0.9 M NaCl, 50 mM NaH₂PO₄, pH 7.0, 5.0 mM Na₂EDTA, 0.5% SDS, 10X Denhardt's and 0.5 mg/ml polyriboadenylic acid. Approximately 2×10^7 cpm (specific activity $4-9 \times 10^8$ cpm/ug) of ³²P end-labeled oligonucleotide probe are then added to the solution. After 12-16 hours of incubation, the membrane is washed for 30 minutes at room temperature in 1X SET (150 mM NaCl, 20 mM Tris hydrochloride, pH 7.8, 1 mM Na₂EDTA) containing 0.5% SDS, followed by a 30

minute wash in fresh 1X SET at $T_m-10^\circ\text{C}$ for the oligonucleotide probe. The membrane is then exposed to auto-radiographic film for detection of hybridization signals.

All of the foregoing hybridizations would be considered to be under conditions of high stringency.

5 Following hybridization, a filter can be washed to remove any non-specifically bound detectable probe. The stringency used to wash the filters can also be varied depending on the nature of the nucleic acids being hybridized, the length of the nucleic acids being hybridized, the degree of complementarity, the nucleotide sequence composition (*e.g.*, GC v. AT content) and the nucleic acid type (*e.g.*, RNA v. DNA). Examples of progressively
10 higher stringency condition washes are as follows: 2X SSC, 0.1% SDS at room temperature for 15 minutes (low stringency); 0.1X SSC, 0.5% SDS at room temperature for 30 minutes to 1 hour (moderate stringency); 0.1X SSC, 0.5% SDS for 15 to 30 minutes at between the hybridization temperature and 68°C (high stringency); and 0.15M NaCl for 15 minutes at 72°C (very high stringency). A final low stringency wash can be conducted in
15 0.1X SSC at room temperature. The examples above are merely illustrative of one set of conditions that can be used to wash filters. One of skill in the art would know that there are numerous recipes for different stringency washes. Some other examples are given below.

Nucleic acids which have hybridized to the probe are identified by autoradiography or other conventional techniques.

20 The above procedure may be modified to identify nucleic acids having decreasing levels of homology to the probe sequence. For example, to obtain nucleic acids of decreasing homology to the detectable probe, less stringent conditions may be used. For example, the hybridization temperature may be decreased in increments of 5°C from 68°C to 42°C in a hybridization buffer having a Na^+ concentration of approximately 1M. Following
25 hybridization, the filter may be washed with 2X SSC, 0.5% SDS at the temperature of hybridization. These conditions are considered to be "moderate" conditions above 50°C and "low" conditions below 50°C . A specific example of "moderate" hybridization conditions is when the above hybridization is conducted at 55°C . A specific example of "low stringency" hybridization conditions is when the above hybridization is conducted at 45°C .

30 Alternatively, the hybridization may be carried out in buffers, such as 6X SSC, containing formamide at a temperature of 42°C . In this case, the concentration of formamide in the hybridization buffer may be reduced in 5% increments from 50% to 0% to identify clones having decreasing levels of homology to the probe. Following hybridization, the filter

may be washed with 6X SSC, 0.5% SDS at 50°C. These conditions are considered to be “moderate” conditions above 25% formamide and “low” conditions below 25% formamide. A specific example of “moderate” hybridization conditions is when the above hybridization is conducted at 30% formamide. A specific example of “low stringency” hybridization
5 conditions is when the above hybridization is conducted at 10% formamide.

However, the selection of a hybridization format is not critical - it is the stringency of the wash conditions that set forth the conditions which determine whether a nucleic acid is within the scope of the invention. Wash conditions used to identify nucleic acids within the scope of the invention include, e.g.: a salt concentration of about 0.02 molar at pH 7
10 and a temperature of at least about 50°C or about 55°C to about 60°C; or, a salt concentration of about 0.15 M NaCl at 72°C for about 15 minutes; or, a salt concentration of about 0.2X SSC at a temperature of at least about 50°C or about 55°C to about 60°C for about 15 to about 20 minutes; or, the hybridization complex is washed twice with a solution with a salt concentration of about 2X SSC containing 0.1% SDS at room
15 temperature for 15 minutes and then washed twice by 0.1X SSC containing 0.1% SDS at 68°C for 15 minutes; or, equivalent conditions. See Sambrook, Tijssen and Ausubel for a description of SSC buffer and equivalent conditions.

These methods may be used to isolate nucleic acids of the invention. For example, the preceding methods may be used to isolate nucleic acids having a sequence with at least
20 about 97%, at least 95%, at least 90%, at least 85%, at least 80%, at least 75%, at least 70%, at least 65%, at least 60%, at least 55%, or at least 50% homology to a nucleic acid sequence selected from the group consisting of one of the sequences of The invention and sequences substantially identical thereto, or fragments comprising at least about 10, 15, 20, 25, 30, 35, 40, 50, 75, 100, 150, 200, 300, 400, or 500 consecutive bases thereof and the
25 sequences complementary thereto. Homology may be measured using the alignment algorithm. For example, the homologous polynucleotides may have a coding sequence which is a naturally occurring allelic variant of one of the coding sequences described herein. Such allelic variants may have a substitution, deletion or addition of one or more nucleotides when compared to the nucleic acids of The invention or the sequences
30 complementary thereto.

Additionally, the above procedures may be used to isolate nucleic acids which encode polypeptides having at least about 99%, 95%, at least 90%, at least 85%, at least 80%, at least 75%, at least 70%, at least 65%, at least 60%, at least 55%, or at least 50%

homology to a polypeptide having the sequence of one of amino acid sequences of the invention, or fragments comprising at least 5, 10, 15, 20, 25, 30, 35, 40, 50, 75, 100, or 150 consecutive amino acids thereof as determined using a sequence alignment algorithm (e.g., such as the FASTA version 3.0t78 algorithm with the default parameters).

5 Oligonucleotides probes and methods for using them

The invention also provides nucleic acid probes that can be used, e.g., for identifying nucleic acids encoding a polypeptide with an isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase activity or fragments thereof or for identifying isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase genes. In one aspect, the probe comprises at least 10 consecutive bases of a nucleic acid of the invention. Alternatively, a probe of the invention can be at least about 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, 100, 110, 120, 130, 150 or about 10 to 50, about 20 to 60 about 30 to 70, consecutive bases of a sequence as set forth in a nucleic acid of the invention. The probes identify a nucleic acid by binding and/or hybridization. The probes can be used in arrays of the invention, see discussion below, including, e.g., capillary arrays. The probes of the invention can also be used to isolate other nucleic acids or polypeptides.

The isolated nucleic acids of the invention and sequences substantially identical thereto, the sequences complementary thereto, or a fragment comprising at least 10, 15, 20, 25, 30, 35, 40, 50, 75, 100, 150, 200, 300, 400, or 500 consecutive bases of one of the sequences of The invention and sequences substantially identical thereto, or the sequences complementary thereto may also be used as probes to determine whether a biological sample, such as a soil sample, contains an organism having a nucleic acid sequence of the invention or an organism from which the nucleic acid was obtained. In such procedures, a biological sample potentially harboring the organism from which the nucleic acid was isolated is obtained and nucleic acids are obtained from the sample. The nucleic acids are contacted with the probe under conditions which permit the probe to specifically hybridize to any complementary sequences from which are present therein.

Where necessary, conditions which permit the probe to specifically hybridize to complementary sequences may be determined by placing the probe in contact with complementary sequences from samples known to contain the complementary sequence as well as control sequences which do not contain the complementary sequence.

Hybridization conditions, such as the salt concentration of the hybridization buffer, the

formamide concentration of the hybridization buffer, or the hybridization temperature, may be varied to identify conditions which allow the probe to hybridize specifically to complementary nucleic acids.

5 If the sample contains the organism from which the nucleic acid was isolated, specific hybridization of the probe is then detected. Hybridization may be detected by labeling the probe with a detectable agent such as a radioactive isotope, a fluorescent dye or an enzyme capable of catalyzing the formation of a detectable product.

10 Many methods for using the labeled probes to detect the presence of complementary nucleic acids in a sample are familiar to those skilled in the art. These include Southern Blots, Northern Blots, colony hybridization procedures and dot blots. Protocols for each of these procedures are provided in Ausubel *et al.* Current Protocols in Molecular Biology, John Wiley 503 Sons, Inc. (1997) and Sambrook *et al.*, Molecular Cloning: A Laboratory Manual 2nd Ed., Cold Spring Harbor Laboratory Press (1989).

15 Alternatively, more than one probe (at least one of which is capable of specifically hybridizing to any complementary sequences which are present in the nucleic acid sample), may be used in an amplification reaction to determine whether the sample contains an organism containing a nucleic acid sequence of the invention (*e.g.*, an organism from which the nucleic acid was isolated). Typically, the probes comprise oligonucleotides. In one aspect, the amplification reaction may comprise a PCR reaction. 20 PCR protocols are described in Ausubel and Sambrook, *supra*. Alternatively, the amplification may comprise a ligase chain reaction, 3SR, or strand displacement reaction. (See Barany, F., "The Ligase Chain Reaction in a PCR World", *PCR Methods and Applications* 1:5-16, 1991; E. Fahy *et al.*, "Self-sustained Sequence Replication (3SR): An Isothermal Transcription-based Amplification System Alternative to PCR", *PCR Methods and Applications* 1:25-33, 1991; and Walker G.T. *et al.*, "Strand Displacement 25 Amplification-an Isothermal *in vitro* DNA Amplification Technique", *Nucleic Acid Research* 20:1691-1696, 1992). In such procedures, the nucleic acids in the sample are contacted with the probes, the amplification reaction is performed and any resulting amplification product is detected. The amplification product may be detected by performing gel electrophoresis on 30 the reaction products and staining the gel with an intercalator such as ethidium bromide. Alternatively, one or more of the probes may be labeled with a radioactive isotope and the presence of a radioactive amplification product may be detected by autoradiography after gel electrophoresis.

Probes derived from sequences near the ends of the sequences of The invention and sequences substantially identical thereto, may also be used in chromosome walking procedures to identify clones containing genomic sequences located adjacent to the sequences of The invention and sequences substantially identical thereto. Such methods
5 allow the isolation of genes which encode additional proteins from the host organism.

The isolated nucleic acids of the invention and sequences substantially identical thereto, the sequences complementary thereto, or a fragment comprising at least 10, 15, 20, 25, 30, 35, 40, 50, 75, 100, 150, 200, 300, 400, or 500 consecutive bases of one of the sequences of the invention and sequences substantially identical thereto, or the sequences
10 complementary thereto may be used as probes to identify and isolate related nucleic acids. In some aspects, the related nucleic acids may be cDNAs or genomic DNAs from organisms other than the one from which the nucleic acid was isolated. For example, the other organisms may be related organisms. In such procedures, a nucleic acid sample is contacted with the probe under conditions which permit the probe to specifically hybridize
15 to related sequences. Hybridization of the probe to nucleic acids from the related organism is then detected using any of the methods described above.

By varying the stringency of the hybridization conditions used to identify nucleic acids, such as cDNAs or genomic DNAs, which hybridize to the detectable probe, nucleic acids having different levels of homology to the probe can be identified and isolated.
20 Stringency may be varied by conducting the hybridization at varying temperatures below the melting temperatures of the probes. The melting temperature, T_m , is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly complementary probe. Very stringent conditions are selected to be equal to or about 5°C lower than the T_m for a particular probe. The melting temperature of the probe may be
25 calculated using the following formulas:

For probes between 14 and 70 nucleotides in length the melting temperature (T_m) is calculated using the formula: $T_m = 81.5 + 16.6(\log [Na^+]) + 0.41(\text{fraction G+C}) - (600/N)$ where N is the length of the probe.

If the hybridization is carried out in a solution containing formamide, the melting
30 temperature may be calculated using the equation: $T_m = 81.5 + 16.6(\log [Na^+]) + 0.41(\text{fraction G+C}) - (0.63\% \text{ formamide}) - (600/N)$ where N is the length of the probe.

Prehybridization may be carried out in 6X SSC, 5X Denhardt's reagent, 0.5% SDS, 100 µg/ml denatured fragmented salmon sperm DNA or 6X SSC, 5X Denhardt's reagent, 0.5% SDS, 100 µg/ml denatured fragmented salmon sperm DNA, 50% formamide. The formulas for SSC and Denhardt's solutions are listed in Sambrook *et al.*, *supra*.

5 Hybridization is conducted by adding the detectable probe to the prehybridization solutions listed above. Where the probe comprises double stranded DNA, it is denatured before addition to the hybridization solution. The filter is contacted with the hybridization solution for a sufficient period of time to allow the probe to hybridize to cDNAs or genomic DNAs containing sequences complementary thereto or homologous thereto. For probes over
10 200 nucleotides in length, the hybridization may be carried out at 15-25°C below the T_m . For shorter probes, such as oligonucleotide probes, the hybridization may be conducted at 5-10°C below the T_m . Typically, for hybridizations in 6X SSC, the hybridization is conducted at approximately 68°C. Usually, for hybridizations in 50% formamide containing solutions, the hybridization is conducted at approximately 42°C.

15 Inhibiting Expression of Isomerases

The invention provides nucleic acids complementary to (e.g., antisense sequences to) the nucleic acids of the invention, e.g., an isomerase-, e.g., a racemase-, e.g., an amino acid racemase-, an alanine racemase-, and/or an epimerase-isomerase-, e.g., racemase-, e.g., amino acid racemase-, alanine racemase-, and/or epimerase-encoding nucleic acids.
20 Antisense sequences are capable of inhibiting the transport, splicing or transcription of an isomerase-, e.g., a racemase-, e.g., an amino acid racemase-, an alanine racemase-, and/or an epimerase-isomerase-, e.g., racemase-, e.g., amino acid racemase-, alanine racemase-, and/or epimerase- encoding genes. The inhibition can be effected through the targeting of genomic DNA or messenger RNA. The transcription or function of targeted nucleic acid
25 can be inhibited, for example, by hybridization and/or cleavage. One particularly useful set of inhibitors provided by the present invention includes oligonucleotides which are able to either bind an isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase gene or message, in either case preventing or inhibiting the production or function of an isomerase, e.g., racemase, e.g., amino acid racemase, alanine
30 racemase, and/or epimerase. The association can be through sequence specific hybridization. Another useful class of inhibitors includes oligonucleotides which cause inactivation or cleavage of an isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase message. The oligonucleotide can have enzyme

activity which causes such cleavage, such as ribozymes. The oligonucleotide can be chemically modified or conjugated to an enzyme or composition capable of cleaving the complementary nucleic acid. A pool of many different such oligonucleotides can be screened for those with the desired activity. Thus, the invention provides various
5 compositions for the inhibition of isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase expression on a nucleic acid and/or protein level, e.g., antisense, iRNA and ribozymes comprising isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase sequences of the invention and the anti-isomerase, e.g., anti-racemase, e.g., anti-amino acid racemase, anti-alanine racemase,
10 and/or anti-epimerase antibodies of the invention.

The compositions of the invention for the inhibition of isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase expression (e.g., antisense, iRNA, ribozymes, antibodies) can be used as pharmaceutical compositions.

Antisense Oligonucleotides

15 The invention provides antisense oligonucleotides capable of binding an isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase message which can inhibit, for example, isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase activity by targeting mRNA. Strategies for designing antisense oligonucleotides are well described in the scientific and patent
20 literature, and the skilled artisan can design such isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase oligonucleotides using the novel reagents of the invention. For example, gene walking/ RNA mapping protocols to screen for effective antisense oligonucleotides are well known in the art, see, e.g., Ho (2000) *Methods Enzymol.* 314:168-183, describing an RNA mapping assay, which is based on
25 standard molecular techniques to provide an easy and reliable method for potent antisense sequence selection. See also Smith (2000) *Eur. J. Pharm. Sci.* 11:191-198.

Naturally occurring nucleic acids are used as antisense oligonucleotides. The antisense oligonucleotides can be of any length; for example, in alternative aspects, the antisense oligonucleotides are between about 5 to 100, about 10 to 80, about 15 to 60,
30 about 18 to 40. The optimal length can be determined by routine screening. The antisense oligonucleotides can be present at any concentration. The optimal concentration can be determined by routine screening. A wide variety of synthetic, non-naturally occurring nucleotide and nucleic acid analogues are known which can address this potential

problem. For example, peptide nucleic acids (PNAs) containing non-ionic backbones, such as N-(2-aminoethyl) glycine units can be used. Antisense oligonucleotides having phosphorothioate linkages can also be used, as described in WO 97/03211; WO 96/39154; Mata (1997) Toxicol Appl Pharmacol 144:189-197; Antisense Therapeutics, ed. Agrawal
5 (Humana Press, Totowa, N.J., 1996). Antisense oligonucleotides having synthetic DNA backbone analogues provided by the invention can also include phosphoro-dithioate, methylphosphonate, phosphoramidate, alkyl phosphotriester, sulfamate, 3'-thioacetal, methylene(methylimino), 3'-N-carbamate, and morpholino carbamate nucleic acids, as described above.

10 Combinatorial chemistry methodology can be used to create vast numbers of oligonucleotides that can be rapidly screened for specific oligonucleotides that have appropriate binding affinities and specificities toward any target, such as the sense and antisense isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase sequences of the invention (see, e.g., Gold (1995) J. of Biol. Chem. 270:13581-
15 13584).

Inhibitory Ribozymes

The invention provides ribozymes capable of binding an isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase message. These ribozymes can inhibit isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or
20 epimerase activity by, e.g., targeting mRNA. Strategies for designing ribozymes and selecting the an isomerase-, e.g., a racemase-, e.g., an amino acid racemase-, an alanine racemase-, and/or an epimerase-isomerase-, e.g., racemase-, e.g., amino acid racemase-, alanine racemase-, and/or epimerase-specific antisense sequence for targeting are well described in the scientific and patent literature, and the skilled artisan can design such
25 ribozymes using the novel reagents (e.g. nucleic acids) of the invention. Ribozymes act by binding to a target RNA through the target RNA binding portion of a ribozyme which is held in close proximity to an enzymatic portion of the RNA that cleaves the target RNA. Thus, the ribozyme recognizes and binds a target RNA through complementary base-pairing, and once bound to the correct site, acts enzymatically to cleave and inactivate the
30 target RNA. Cleavage of a target RNA in such a manner will destroy its ability to direct synthesis of an encoded protein if the cleavage occurs in the coding sequence. After a ribozyme has bound and cleaved its RNA target, it can be released from that RNA to bind and cleave new targets repeatedly.

In some circumstances, the enzymatic nature of a ribozyme can be advantageous over other technologies, such as antisense technology (where a nucleic acid molecule simply binds to a nucleic acid target to block its transcription, translation or association with another molecule) as the effective concentration of ribozyme necessary to effect a therapeutic treatment can be lower than that of an antisense oligonucleotide. This potential advantage reflects the ability of the ribozyme to act enzymatically. Thus, a single ribozyme molecule is able to cleave many molecules of target RNA. In addition, a ribozyme is typically a highly specific inhibitor, with the specificity of inhibition depending not only on the base pairing mechanism of binding, but also on the mechanism by which the molecule inhibits the expression of the RNA to which it binds. That is, the inhibition is caused by cleavage of the RNA target and so specificity is defined as the ratio of the rate of cleavage of the targeted RNA over the rate of cleavage of non-targeted RNA. This cleavage mechanism is dependent upon factors additional to those involved in base pairing. Thus, the specificity of action of a ribozyme can be greater than that of antisense oligonucleotide binding the same RNA site.

The ribozyme of the invention, e.g., an enzymatic ribozyme RNA molecule, can be formed in a hammerhead motif, a hairpin motif, as a hepatitis delta virus motif, a group I intron motif and/or an RNaseP-like RNA in association with an RNA guide sequence. Examples of hammerhead motifs are described by, e.g., Rossi (1992) *Aids Research and Human Retroviruses* 8:183; hairpin motifs by Hampel (1989) *Biochemistry* 28:4929, and Hampel (1990) *Nuc. Acids Res.* 18:299; the hepatitis delta virus motif by Perrotta (1992) *Biochemistry* 31:16; the RNaseP motif by Guerrier-Takada (1983) *Cell* 35:849; and the group I intron by Cech U.S. Pat. No. 4,987,071. The recitation of these specific motifs is not intended to be limiting. Those skilled in the art will recognize that a ribozyme of the invention, e.g., an enzymatic RNA molecule of this invention, can have a specific substrate binding site complementary to one or more of the target gene RNA regions. A ribozyme of the invention can have a nucleotide sequence within or surrounding that substrate binding site which imparts an RNA cleaving activity to the molecule.

RNA interference (RNAi)

In one aspect, the invention provides an RNA inhibitory molecule, a so-called "RNAi" molecule, comprising an isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase enzyme sequence of the invention. The RNAi molecule can comprise a double-stranded RNA (dsRNA) molecule, e.g., siRNA, miRNA

and/or short hairpin RNA (shRNA) molecules. The RNAi molecule, e.g., siRNA (small inhibitory RNA) can inhibit expression of an isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase enzyme gene, and/or miRNA (micro RNA) to inhibit translation of an isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase message. In one aspect, the RNAi molecule, e.g., siRNA and/or miRNA, is about 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29 or more duplex nucleotides in length. While the invention is not limited by any particular mechanism of action, the RNAi can enter a cell and cause the degradation of a single-stranded RNA (ssRNA) of similar or identical sequences, including endogenous mRNAs. When a cell is exposed to double-stranded RNA (dsRNA), mRNA from the homologous gene is selectively degraded by a process called RNA interference (RNAi). A possible basic mechanism behind RNAi is the breaking of a double-stranded RNA (dsRNA) matching a specific gene sequence into short pieces called short interfering RNA, which trigger the degradation of mRNA that matches its sequence. In one aspect, the RNAi's of the invention are used in gene-silencing therapeutics, see, e.g., Shuey (2002) Drug Discov. Today 7:1040-1046. In one aspect, the invention provides methods to selectively degrade RNA using the RNAi's molecules, e.g., siRNA and/or miRNA, of the invention. The process may be practiced *in vitro*, *ex vivo* or *in vivo*. In one aspect, the RNAi molecules of the invention can be used to generate a loss-of-function mutation in a cell, an organ or an animal.

In one aspect, intracellular introduction of the RNAi is by internalization of a target cell specific ligand bonded to an RNA binding protein comprising an RNAi (e.g., microRNA) is adsorbed. The ligand is specific to a unique target cell surface antigen. The ligand can be spontaneously internalized after binding to the cell surface antigen. If the unique cell surface antigen is not naturally internalized after binding to its ligand, internalization can be promoted by the incorporation of an arginine-rich peptide, or other membrane permeable peptide, into the structure of the ligand or RNA binding protein or attachment of such a peptide to the ligand or RNA binding protein. See, e.g., U.S. Patent App. Pub. Nos. 20060030003; 20060025361; 20060019286; 20060019258. In one aspect, the invention provides lipid-based formulations for delivering, e.g., introducing nucleic acids of the invention as nucleic acid-lipid particles comprising an RNAi molecule to a cell, see .g., U.S. Patent App. Pub. No. 20060008910.

Methods for making and using RNAi molecules, e.g., siRNA and/or miRNA, for selectively degrade RNA are well known in the art, see, e.g., U.S. Patent No. 6,506,559; 6,511,824; 6,515,109; 6,489,127.

Modification of Nucleic Acids

5 The invention provides methods of generating variants of the nucleic acids of the invention, e.g., those encoding an isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase. These methods can be repeated or used in various combinations to generate isomerases, e.g., racemases, e.g., amino acid racemases, alanine racemases, and/or epimerases having an altered or different activity or an altered or
10 different stability from that of an isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase encoded by the template nucleic acid. These methods also can be repeated or used in various combinations, e.g., to generate variations in gene/
message expression, message translation or message stability. In another aspect, the genetic composition of a cell is altered by, e.g., modification of a homologous gene ex
15 vivo, followed by its reinsertion into the cell.

A nucleic acid of the invention can be altered by any means. For example, random or stochastic methods, or, non-stochastic, or “directed evolution,” methods, see, e.g., U.S. Patent No. 6,361,974. Methods for random mutation of genes are well known in the art, see, e.g., U.S. Patent No. 5,830,696. For example, mutagens can be used to randomly
20 mutate a gene. Mutagens include, e.g., ultraviolet light or gamma irradiation, or a chemical mutagen, e.g., mitomycin, nitrous acid, photoactivated psoralens, alone or in combination, to induce DNA breaks amenable to repair by recombination. Other chemical mutagens include, for example, sodium bisulfite, nitrous acid, hydroxylamine, hydrazine or formic acid. Other mutagens are analogues of nucleotide precursors, e.g.,
25 nitrosoguanidine, 5-bromouracil, 2-aminopurine, or acridine. These agents can be added to a PCR reaction in place of the nucleotide precursor thereby mutating the sequence. Intercalating agents such as proflavine, acriflavine, quinacrine and the like can also be used.

Any technique in molecular biology can be used, e.g., random PCR mutagenesis,
30 see, e.g., Rice (1992) Proc. Natl. Acad. Sci. USA 89:5467-5471; or, combinatorial multiple cassette mutagenesis, see, e.g., Cramer (1995) Biotechniques 18:194-196. Alternatively, nucleic acids, e.g., genes, can be reassembled after random, or “stochastic,” fragmentation, see, e.g., U.S. Patent Nos. 6,291,242; 6,287,862; 6,287,861; 5,955,358;

5,830,721; 5,824,514; 5,811,238; 5,605,793. In alternative aspects, modifications, additions or deletions are introduced by error-prone PCR, shuffling, oligonucleotide-directed mutagenesis, assembly PCR, sexual PCR mutagenesis, in vivo mutagenesis, cassette mutagenesis, recursive ensemble mutagenesis, exponential ensemble mutagenesis, site-specific mutagenesis, gene reassembly (e.g., GeneReassembly, see, e.g., U.S. Patent No. 6,537,776), gene site saturation mutagenesis (GSSM), synthetic ligation reassembly (SLR), recombination, recursive sequence recombination, phosphothioate-modified DNA mutagenesis, uracil-containing template mutagenesis, gapped duplex mutagenesis, point mismatch repair mutagenesis, repair-deficient host strain mutagenesis, chemical mutagenesis, radiogenic mutagenesis, deletion mutagenesis, restriction-selection mutagenesis, restriction-purification mutagenesis, artificial gene synthesis, ensemble mutagenesis, chimeric nucleic acid multimer creation, and/or a combination of these and other methods.

The following publications describe a variety of recursive recombination procedures and/or methods which can be incorporated into the methods of the invention:

Stemmer (1999) "Molecular breeding of viruses for targeting and other clinical properties" *Tumor Targeting* 4:1-4; Ness (1999) *Nature Biotechnology* 17:893-896; Chang (1999) "Evolution of a cytokine using DNA family shuffling" *Nature Biotechnology* 17:793-797; Minshull (1999) "Protein evolution by molecular breeding" *Current Opinion in Chemical Biology* 3:284-290; Christians (1999) "Directed evolution of thymidine kinase for AZT phosphorylation using DNA family shuffling" *Nature Biotechnology* 17:259-264; Crameri (1998) "DNA shuffling of a family of genes from diverse species accelerates directed evolution" *Nature* 391:288-291; Crameri (1997) "Molecular evolution of an arsenate detoxification pathway by DNA shuffling," *Nature Biotechnology* 15:436-438; Zhang (1997) "Directed evolution of an effective fucosidase from a galactosidase by DNA shuffling and screening" *Proc. Natl. Acad. Sci. USA* 94:4504-4509; Patten et al. (1997) "Applications of DNA Shuffling to Pharmaceuticals and Vaccines" *Current Opinion in Biotechnology* 8:724-733; Crameri et al. (1996) "Construction and evolution of antibody-phage libraries by DNA shuffling" *Nature Medicine* 2:100-103; Gates et al. (1996) "Affinity selective isolation of ligands from peptide libraries through display on a lac repressor 'headpiece dimer'" *Journal of Molecular Biology* 255:373-386; Stemmer (1996) "Sexual PCR and Assembly PCR" In: *The Encyclopedia of Molecular Biology*. VCH Publishers, New York. pp.447-457; Crameri and Stemmer (1995) "Combinatorial multiple cassette mutagenesis creates all the permutations of mutant and wildtype cassettes"

BioTechniques 18:194-195; Stemmer et al. (1995) "Single-step assembly of a gene and entire plasmid from large numbers of oligodeoxyribonucleotides" *Gene*, 164:49-53; Stemmer (1995) "The Evolution of Molecular Computation" *Science* 270: 1510; Stemmer (1995) "Searching Sequence Space" *Bio/Technology* 13:549-553; Stemmer (1994) "Rapid evolution of a protein in vitro by DNA shuffling" *Nature* 370:389-391; and Stemmer (1994) "DNA shuffling by random fragmentation and reassembly: In vitro recombination for molecular evolution." *Proc. Natl. Acad. Sci. USA* 91:10747-10751.

Mutational methods of generating diversity include, for example, site-directed mutagenesis (Ling et al. (1997) "Approaches to DNA mutagenesis: an overview" *Anal Biochem.* 254(2): 157-178; Dale et al. (1996) "Oligonucleotide-directed random mutagenesis using the phosphorothioate method" *Methods Mol. Biol.* 57:369-374; Smith (1985) "In vitro mutagenesis" *Ann. Rev. Genet.* 19:423-462; Botstein & Shortle (1985) "Strategies and applications of in vitro mutagenesis" *Science* 229:1193-1201; Carter (1986) "Site-directed mutagenesis" *Biochem. J.* 237:1-7; and Kunkel (1987) "The efficiency of oligonucleotide directed mutagenesis" in *Nucleic Acids & Molecular Biology* (Eckstein, F. and Lilley, D. M. J. eds., Springer Verlag, Berlin)); mutagenesis using uracil containing templates (Kunkel (1985) "Rapid and efficient site-specific mutagenesis without phenotypic selection" *Proc. Natl. Acad. Sci. USA* 82:488-492; Kunkel et al. (1987) "Rapid and efficient site-specific mutagenesis without phenotypic selection" *Methods in Enzymol.* 154, 367-382; and Bass et al. (1988) "Mutant Trp repressors with new DNA-binding specificities" *Science* 242:240-245); oligonucleotide-directed mutagenesis (*Methods in Enzymol.* 100: 468-500 (1983); *Methods in Enzymol.* 154: 329-350 (1987); Zoller (1982) "Oligonucleotide-directed mutagenesis using M13-derived vectors: an efficient and general procedure for the production of point mutations in any DNA fragment" *Nucleic Acids Res.* 10:6487-6500; Zoller & Smith (1983) "Oligonucleotide-directed mutagenesis of DNA fragments cloned into M13 vectors" *Methods in Enzymol.* 100:468-500; and Zoller (1987) Oligonucleotide-directed mutagenesis: a simple method using two oligonucleotide primers and a single-stranded DNA template" *Methods in Enzymol.* 154:329-350); phosphorothioate-modified DNA mutagenesis (Taylor (1985) "The use of phosphorothioate-modified DNA in restriction enzyme reactions to prepare nicked DNA" *Nucl. Acids Res.* 13: 8749-8764; Taylor (1985) "The rapid generation of oligonucleotide-directed mutations at high frequency using phosphorothioate-modified DNA" *Nucl. Acids Res.* 13: 8765-8787 (1985); Nakamaye (1986) "Inhibition of restriction endonuclease Nci I cleavage by phosphorothioate groups

and its application to oligonucleotide-directed mutagenesis" Nucl. Acids Res. 14: 9679-9698; Sayers (1988) "Y-T Exonucleases in phosphorothioate-based oligonucleotide-directed mutagenesis" Nucl. Acids Res. 16:791-802; and Sayers et al. (1988) "Strand specific cleavage of phosphorothioate-containing DNA by reaction with restriction
5 endonucleases in the presence of ethidium bromide" Nucl. Acids Res. 16: 803-814); mutagenesis using gapped duplex DNA (Kramer et al. (1984) "The gapped duplex DNA approach to oligonucleotide-directed mutation construction" Nucl. Acids Res. 12: 9441-9456; Kramer & Fritz (1987) Methods in Enzymol. "Oligonucleotide-directed construction of mutations via gapped duplex DNA" 154:350-367; Kramer (1988) "Improved enzymatic
10 in vitro reactions in the gapped duplex DNA approach to oligonucleotide-directed construction of mutations" Nucl. Acids Res. 16: 7207; and Fritz (1988) "Oligonucleotide-directed construction of mutations: a gapped duplex DNA procedure without enzymatic reactions *in vitro*" Nucl. Acids Res. 16: 6987-6999).

Additional protocols that can be used to practice the invention include point
15 mismatch repair (Kramer (1984) "Point Mismatch Repair" Cell 38:879-887), mutagenesis using repair-deficient host strains (Carter et al. (1985) "Improved oligonucleotide site-directed mutagenesis using M13 vectors" Nucl. Acids Res. 13: 4431-4443; and Carter (1987) "Improved oligonucleotide-directed mutagenesis using M13 vectors" Methods in Enzymol. 154: 382-403), deletion mutagenesis (Eghtedarzadeh (1986) "Use of
20 oligonucleotides to generate large deletions" Nucl. Acids Res. 14: 5115), restriction-selection and restriction-selection and restriction-purification (Wells et al. (1986) "Importance of hydrogen-bond formation in stabilizing the transition state of subtilisin" Phil. Trans. R. Soc. Lond. A 317: 415-423), mutagenesis by total gene synthesis (Nambiar et al. (1984) "Total synthesis and cloning of a gene coding for the ribonuclease S protein"
25 Science 223: 1299-1301; Sakamar and Khorana (1988) "Total synthesis and expression of a gene for the α -subunit of bovine rod outer segment guanine nucleotide-binding protein (transducin)" Nucl. Acids Res. 14: 6361-6372; Wells et al. (1985) "Cassette mutagenesis: an efficient method for generation of multiple mutations at defined sites" Gene 34:315-323; and Grundstrom et al. (1985) "Oligonucleotide-directed mutagenesis by microscale
30 `shot-gun` gene synthesis" Nucl. Acids Res. 13: 3305-3316), double-strand break repair (Mandecki (1986); Arnold (1993) "Protein engineering for unusual environments" Current Opinion in Biotechnology 4:450-455. "Oligonucleotide-directed double-strand break repair in plasmids of *Escherichia coli*: a method for site-specific mutagenesis" Proc. Natl. Acad. Sci. USA, 83:7177-7181). Additional details on many of the above methods can be

found in Methods in Enzymology Volume 154, which also describes useful controls for trouble-shooting problems with various mutagenesis methods.

Protocols that can be used to practice the invention are described, e.g., in U.S. Patent Nos. 5,605,793 to Stemmer (Feb. 25, 1997), "Methods for In Vitro
 5 Recombination;" U.S. Pat. No. 5,811,238 to Stemmer et al. (Sep. 22, 1998) "Methods for Generating Polynucleotides having Desired Characteristics by Iterative Selection and Recombination;" U.S. Pat. No. 5,830,721 to Stemmer et al. (Nov. 3, 1998), "DNA Mutagenesis by Random Fragmentation and Reassembly;" U.S. Pat. No. 5,834,252 to Stemmer, et al. (Nov. 10, 1998) "End-Complementary Polymerase Reaction;" U.S. Pat.
 10 No. 5,837,458 to Minshull, et al. (Nov. 17, 1998), "Methods and Compositions for Cellular and Metabolic Engineering;" WO 95/22625, Stemmer and Cramer, "Mutagenesis by Random Fragmentation and Reassembly;" WO 96/33207 by Stemmer and Lipschutz "End Complementary Polymerase Chain Reaction;" WO 97/20078 by Stemmer and Cramer "Methods for Generating Polynucleotides having Desired Characteristics by
 15 Iterative Selection and Recombination;" WO 97/35966 by Minshull and Stemmer, "Methods and Compositions for Cellular and Metabolic Engineering;" WO 99/41402 by Punnonen et al. "Targeting of Genetic Vaccine Vectors;" WO 99/41383 by Punnonen et al. "Antigen Library Immunization;" WO 99/41369 by Punnonen et al. "Genetic Vaccine Vector Engineering;" WO 99/41368 by Punnonen et al. "Optimization of
 20 Immunomodulatory Properties of Genetic Vaccines;" EP 752008 by Stemmer and Cramer, "DNA Mutagenesis by Random Fragmentation and Reassembly;" EP 0932670 by Stemmer "Evolving Cellular DNA Uptake by Recursive Sequence Recombination;" WO 99/23107 by Stemmer et al., "Modification of Virus Tropism and Host Range by Viral Genome Shuffling;" WO 99/21979 by Apt et al., "Human Papillomavirus Vectors;"
 25 WO 98/31837 by del Cardayre et al. "Evolution of Whole Cells and Organisms by Recursive Sequence Recombination;" WO 98/27230 by Patten and Stemmer, "Methods and Compositions for Polypeptide Engineering;" WO 98/27230 by Stemmer et al., "Methods for Optimization of Gene Therapy by Recursive Sequence Shuffling and Selection," WO 00/00632, "Methods for Generating Highly Diverse Libraries," WO
 30 00/09679, "Methods for Obtaining in Vitro Recombined Polynucleotide Sequence Banks and Resulting Sequences," WO 98/42832 by Arnold et al., "Recombination of Polynucleotide Sequences Using Random or Defined Primers," WO 99/29902 by Arnold et al., "Method for Creating Polynucleotide and Polypeptide Sequences," WO 98/41653 by Vind, "An in Vitro Method for Construction of a DNA Library," WO 98/41622 by

Borchert et al., "Method for Constructing a Library Using DNA Shuffling," and WO 98/42727 by Pati and Zaring, "Sequence Alterations using Homologous Recombination."

Protocols that can be used to practice the invention (providing details regarding various diversity generating methods) are described, e.g., in U.S. Patent application serial
5 no. (USSN) 09/407,800, "SHUFFLING OF CODON ALTERED GENES" by Patten et al. filed Sep. 28, 1999; "EVOLUTION OF WHOLE CELLS AND ORGANISMS BY RECURSIVE SEQUENCE RECOMBINATION" by del Cardayre et al., United States Patent No. 6,379,964; "OLIGONUCLEOTIDE MEDIATED NUCLEIC ACID
10 RECOMBINATION" by Crameri et al., United States Patent Nos. 6,319,714; 6,368,861; 6,376,246; 6,423,542; 6,426,224 and PCT/US00/01203; "USE OF CODON-VARIED OLIGONUCLEOTIDE SYNTHESIS FOR SYNTHETIC SHUFFLING" by Welch et al., United States Patent No. 6,436,675; "METHODS FOR MAKING CHARACTER STRINGS, POLYNUCLEOTIDES & POLYPEPTIDES HAVING DESIRED CHARACTERISTICS" by Selifonov et al., filed Jan. 18, 2000, (PCT/US00/01202) and,
15 e.g. "METHODS FOR MAKING CHARACTER STRINGS, POLYNUCLEOTIDES & POLYPEPTIDES HAVING DESIRED CHARACTERISTICS" by Selifonov et al., filed Jul. 18, 2000 (U.S. Ser. No. 09/618,579); "METHODS OF POPULATING DATA STRUCTURES FOR USE IN EVOLUTIONARY SIMULATIONS" by Selifonov and Stemmer, filed Jan. 18, 2000 (PCT/US00/01138); and "SINGLE-STRANDED NUCLEIC
20 ACID TEMPLATE-MEDIATED RECOMBINATION AND NUCLEIC ACID FRAGMENT ISOLATION" by Affholter, filed Sep. 6, 2000 (U.S. Ser. No. 09/656,549); and United States Patent Nos. 6,177,263; 6,153,410.

Non-stochastic, or "directed evolution," methods include, e.g., saturation mutagenesis (GSSM), synthetic ligation reassembly (SLR), or a combination thereof are
25 used to modify the nucleic acids of the invention to generate isomerases, e.g., racemases, e.g., amino acid racemases, alanine racemases, and/or epimerases with new or altered properties (e.g., activity under highly acidic or alkaline conditions, high or low temperatures, and the like). Polypeptides encoded by the modified nucleic acids can be screened for an activity. Any testing modality or protocol can be used, e.g., using a
30 capillary array platform. See, e.g., U.S. Patent Nos. 6,361,974; 6,280,926; 5,939,250.

Gene Site Saturation Mutagenesis, or, GSSM

The invention also provides methods for making enzyme using Gene Site Saturation mutagenesis, or, GSSM, as described herein, and also in U.S. Patent Nos.

6,171,820 and 6,579,258. In one aspect, codon primers containing a degenerate N,N,G/T sequence are used to introduce point mutations into a polynucleotide, e.g., an isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase or an antibody of the invention, so as to generate a set of progeny polypeptides in which a full range of single amino acid substitutions is represented at each amino acid position, e.g., an amino acid residue in an enzyme active site or ligand binding site targeted to be modified. These oligonucleotides can comprise a contiguous first homologous sequence, a degenerate N,N,G/T sequence, and, in one aspect, a second homologous sequence. The downstream progeny translational products from the use of such oligonucleotides include all possible amino acid changes at each amino acid site along the polypeptide, because the degeneracy of the N,N,G/T sequence includes codons for all 20 amino acids. In one aspect, one such degenerate oligonucleotide (comprised of, e.g., one degenerate N,N,G/T cassette) is used for subjecting each original codon in a parental polynucleotide template to a full range of codon substitutions. In another aspect, at least two degenerate cassettes are used – either in the same oligonucleotide or not, for subjecting at least two original codons in a parental polynucleotide template to a full range of codon substitutions. For example, more than one N,N,G/T sequence can be contained in one oligonucleotide to introduce amino acid mutations at more than one site. This plurality of N,N,G/T sequences can be directly contiguous, or separated by one or more additional nucleotide sequence(s). In another aspect, oligonucleotides serviceable for introducing additions and deletions can be used either alone or in combination with the codons containing an N,N,G/T sequence, to introduce any combination or permutation of amino acid additions, deletions, and/or substitutions.

In one aspect, simultaneous mutagenesis of two or more contiguous amino acid positions is done using an oligonucleotide that contains contiguous N,N,G/T triplets, i.e. a degenerate (N,N,G/T)_n sequence. In another aspect, degenerate cassettes having less degeneracy than the N,N,G/T sequence are used. For example, it may be desirable in some instances to use (e.g. in an oligonucleotide) a degenerate triplet sequence comprised of only one N, where said N can be in the first second or third position of the triplet. Any other bases including any combinations and permutations thereof can be used in the remaining two positions of the triplet. Alternatively, it may be desirable in some instances to use (e.g. in an oligo) a degenerate N,N,N triplet sequence.

In one aspect, use of degenerate triplets (e.g., N,N,G/T triplets) allows for systematic and easy generation of a full range of possible natural amino acids (for a total of 20 amino acids) into each and every amino acid position in a polypeptide (in alternative aspects, the methods also include generation of less than all possible substitutions per amino acid residue, or codon, position). For example, for a 100 amino acid polypeptide, 2000 distinct species (i.e. 20 possible amino acids per position X 100 amino acid positions) can be generated. Through the use of an oligonucleotide or set of oligonucleotides containing a degenerate N,N,G/T triplet, 32 individual sequences can code for all 20 possible natural amino acids. Thus, in a reaction vessel in which a parental polynucleotide sequence is subjected to saturation mutagenesis using at least one such oligonucleotide, there are generated 32 distinct progeny polynucleotides encoding 20 distinct polypeptides. In contrast, the use of a non-degenerate oligonucleotide in site-directed mutagenesis leads to only one progeny polypeptide product per reaction vessel. Nondegenerate oligonucleotides can in one aspect be used in combination with degenerate primers disclosed; for example, nondegenerate oligonucleotides can be used to generate specific point mutations in a working polynucleotide. This provides one means to generate specific silent point mutations, point mutations leading to corresponding amino acid changes, and point mutations that cause the generation of stop codons and the corresponding expression of polypeptide fragments.

In one aspect, each saturation mutagenesis reaction vessel contains polynucleotides encoding at least 20 progeny polypeptide (e.g., isomerases, e.g., racemases, e.g., amino acid racemases, alanine racemases, and/or epimerases) molecules such that all 20 natural amino acids are represented at the one specific amino acid position corresponding to the codon position mutagenized in the parental polynucleotide (other aspects use less than all 20 natural combinations). The 32-fold degenerate progeny polypeptides generated from each saturation mutagenesis reaction vessel can be subjected to clonal amplification (e.g. cloned into a suitable host, e.g., *E. coli* host, using, e.g., an expression vector) and subjected to expression screening. When an individual progeny polypeptide is identified by screening to display a favorable change in property (when compared to the parental polypeptide, such as increased isomerase activity, e.g., racemase activity, e.g., amino acid racemase activity, alanine racemase activity, and/or epimerase activity under alkaline or acidic conditions), it can be sequenced to identify the correspondingly favorable amino acid substitution contained therein.

In one aspect, upon mutagenizing each and every amino acid position in a parental polypeptide using saturation mutagenesis as disclosed herein, favorable amino acid changes may be identified at more than one amino acid position. One or more new progeny molecules can be generated that contain a combination of all or part of these favorable amino acid substitutions. For example, if 2 specific favorable amino acid changes are identified in each of 3 amino acid positions in a polypeptide, the permutations include 3 possibilities at each position (no change from the original amino acid, and each of two favorable changes) and 3 positions. Thus, there are $3 \times 3 \times 3$ or 27 total possibilities, including 7 that were previously examined - 6 single point mutations (i.e. 2 at each of three positions) and no change at any position.

In yet another aspect, site-saturation mutagenesis can be used together with shuffling, chimerization, recombination and other mutagenizing processes, along with screening. This invention provides for the use of any mutagenizing process(es), including saturation mutagenesis, in an iterative manner. In one exemplification, the iterative use of any mutagenizing process(es) is used in combination with screening.

The invention also provides for the use of proprietary codon primers (containing a degenerate N,N,N sequence) to introduce point mutations into a polynucleotide, so as to generate a set of progeny polypeptides in which a full range of single amino acid substitutions is represented at each amino acid position (gene site saturation mutagenesis (GSSM)). The oligos used are comprised contiguously of a first homologous sequence, a degenerate N,N,N sequence and preferably but not necessarily a second homologous sequence. The downstream progeny translational products from the use of such oligos include all possible amino acid changes at each amino acid site along the polypeptide, because the degeneracy of the N,N,N sequence includes codons for all 20 amino acids.

In one aspect, one such degenerate oligo (comprised of one degenerate N,N,N cassette) is used for subjecting each original codon in a parental polynucleotide template to a full range of codon substitutions. In another aspect, at least two degenerate N,N,N cassettes are used – either in the same oligo or not, for subjecting at least two original codons in a parental polynucleotide template to a full range of codon substitutions. Thus, more than one N,N,N sequence can be contained in one oligo to introduce amino acid mutations at more than one site. This plurality of N,N,N sequences can be directly contiguous, or separated by one or more additional nucleotide sequence(s). In another aspect, oligos serviceable for introducing additions and deletions can be used either alone

or in combination with the codons containing an N,N,N sequence, to introduce any combination or permutation of amino acid additions, deletions and/or substitutions.

In a particular exemplification, it is possible to simultaneously mutagenize two or more contiguous amino acid positions using an oligo that contains contiguous N,N,N triplets, *i.e.* a degenerate (N,N,N)_n sequence.

In another aspect, the present invention provides for the use of degenerate cassettes having less degeneracy than the N,N,N sequence. For example, it may be desirable in some instances to use (*e.g.* in an oligo) a degenerate triplet sequence comprised of only one N, where the N can be in the first second or third position of the triplet. Any other bases including any combinations and permutations thereof can be used in the remaining two positions of the triplet. Alternatively, it may be desirable in some instances to use (*e.g.*, in an oligo) a degenerate N,N,N triplet sequence, N,N,G/T, or an N,N, G/C triplet sequence.

It is appreciated, however, that the use of a degenerate triplet (such as N,N,G/T or an N,N, G/C triplet sequence) as disclosed in the instant invention is advantageous for several reasons. In one aspect, this invention provides a means to systematically and fairly easily generate the substitution of the full range of possible amino acids (for a total of 20 amino acids) into each and every amino acid position in a polypeptide. Thus, for a 100 amino acid polypeptide, the invention provides a way to systematically and fairly easily generate 2000 distinct species (*i.e.*, 20 possible amino acids per position times 100 amino acid positions). It is appreciated that there is provided, through the use of an oligo containing a degenerate N,N,G/T or an N,N, G/C triplet sequence, 32 individual sequences that code for 20 possible amino acids. Thus, in a reaction vessel in which a parental polynucleotide sequence is subjected to saturation mutagenesis using one such oligo, there are generated 32 distinct progeny polynucleotides encoding 20 distinct polypeptides. In contrast, the use of a non-degenerate oligo in site-directed mutagenesis leads to only one progeny polypeptide product per reaction vessel.

This invention also provides for the use of nondegenerate oligos, which can in one aspect be used in combination with degenerate primers disclosed. It is appreciated that in some situations, it is advantageous to use nondegenerate oligos to generate specific point mutations in a working polynucleotide. This provides a means to generate specific silent point mutations, point mutations leading to corresponding amino acid changes and point mutations that cause the generation of stop codons and the corresponding expression of polypeptide fragments.

Thus, in one aspect of this invention, each saturation mutagenesis reaction vessel contains polynucleotides encoding at least 20 progeny polypeptide molecules such that all 20 amino acids are represented at the one specific amino acid position corresponding to the codon position mutagenized in the parental polynucleotide. The 32-fold degenerate
5 progeny polypeptides generated from each saturation mutagenesis reaction vessel can be subjected to clonal amplification (*e.g.*, cloned into a suitable *E. coli* host using an expression vector) and subjected to expression screening. When an individual progeny polypeptide is identified by screening to display a favorable change in property (when compared to the parental polypeptide), it can be sequenced to identify the correspondingly
10 favorable amino acid substitution contained therein.

It is appreciated that upon mutagenizing each and every amino acid position in a parental polypeptide using saturation mutagenesis as disclosed herein, favorable amino acid changes may be identified at more than one amino acid position. One or more new progeny molecules can be generated that contain a combination of all or part of these
15 favorable amino acid substitutions. For example, if 2 specific favorable amino acid changes are identified in each of 3 amino acid positions in a polypeptide, the permutations include 3 possibilities at each position (no change from the original amino acid and each of two favorable changes) and 3 positions. Thus, there are $3 \times 3 \times 3$ or 27 total possibilities, including 7 that were previously examined - 6 single point mutations (*i.e.*, 2
20 at each of three positions) and no change at any position.

Thus, in a non-limiting exemplification, this invention provides for the use of saturation mutagenesis in combination with additional mutagenization processes, such as process where two or more related polynucleotides are introduced into a suitable host cell such that a hybrid polynucleotide is generated by recombination and reductive
25 reassortment.

In addition to performing mutagenesis along the entire sequence of a gene, the instant invention provides that mutagenesis can be used to replace each of any number of bases in a polynucleotide sequence, wherein the number of bases to be mutagenized is preferably every integer from 15 to 100,000. Thus, instead of mutagenizing every position
30 along a molecule, one can subject every or a discrete number of bases (preferably a subset totaling from 15 to 100,000) to mutagenesis. Preferably, a separate nucleotide is used for mutagenizing each position or group of positions along a polynucleotide sequence. A group of 3 positions to be mutagenized may be a codon. The mutations are preferably introduced using a mutagenic primer, containing a heterologous cassette, also referred to

as a mutagenic cassette. Exemplary cassettes can have from 1 to 500 bases. Each nucleotide position in such heterologous cassettes be N, A, C, G, T, A/C, A/G, A/T, C/G, C/T, G/T, C/G/T, A/G/T, A/C/T, A/C/G, or E, where E is any base that is not A, C, G, or T (E can be referred to as a designer oligo).

5 In a general sense, saturation mutagenesis is comprised of mutagenizing a complete set of mutagenic cassettes (wherein each cassette is preferably about 1-500 bases in length) in defined polynucleotide sequence to be mutagenized (wherein the sequence to be mutagenized is preferably from about 15 to 100,000 bases in length). Thus, a group of mutations (ranging from 1 to 100 mutations) is introduced into each cassette to be
10 mutagenized. A grouping of mutations to be introduced into one cassette can be different or the same from a second grouping of mutations to be introduced into a second cassette during the application of one round of saturation mutagenesis. Such groupings are exemplified by deletions, additions, groupings of particular codons and groupings of particular nucleotide cassettes.

15 Defined sequences to be mutagenized include a whole gene, pathway, cDNA, an entire open reading frame (ORF) and entire promoter, enhancer, repressor/transactivator, origin of replication, intron, operator, or any polynucleotide functional group. Generally, a “defined sequences” for this purpose may be any polynucleotide that a 15 base-
20 polynucleotide sequence and polynucleotide sequences of lengths between 15 bases and 15,000 bases (this invention specifically names every integer in between). Considerations in choosing groupings of codons include types of amino acids encoded by a degenerate mutagenic cassette.

In one exemplification a grouping of mutations that can be introduced into a mutagenic cassette, this invention specifically provides for degenerate codon substitutions
25 (using degenerate oligos) that code for 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 and 20 amino acids at each position and a library of polypeptides encoded thereby.

Synthetic Ligation Reassembly (SLR)

The invention provides a non-stochastic gene modification system termed
30 “synthetic ligation reassembly,” or simply “SLR,” a “directed evolution process,” to generate polypeptides, e.g., isomerases, e.g., racemases, e.g., amino acid racemases, alanine racemases, and/or epimerases, or antibodies of the invention, with new or altered properties.

SLR is a method of ligating oligonucleotide fragments together non-stochastically. This method differs from stochastic oligonucleotide shuffling in that the nucleic acid building blocks are not shuffled, concatenated or chimerized randomly, but rather are assembled non-stochastically. See, e.g., U.S. Patent Nos. 6,773,900; 6,740,506; 5 6,713,282; 6,635,449; 6,605,449; 6,537,776. In one aspect, SLR comprises: (a) providing a template polynucleotide, wherein the template polynucleotide comprises sequence encoding a homologous gene; (b) providing a plurality of building block polynucleotides, wherein the building block polynucleotides are designed to cross-over reassemble with the template polynucleotide at a predetermined sequence, and a building block polynucleotide 10 comprises a sequence that is a variant of the homologous gene and a sequence homologous to the template polynucleotide flanking the variant sequence; (c) combining a building block polynucleotide with a template polynucleotide such that the building block polynucleotide cross-over reassembles with the template polynucleotide to generate polynucleotides comprising homologous gene sequence variations.

15 SLR does not depend on the presence of high levels of homology between polynucleotides to be rearranged. Thus, this method can be used to non-stochastically generate libraries (or sets) of progeny molecules comprised of over 10^{100} different chimeras. SLR can be used to generate libraries comprised of over 10^{1000} different progeny chimeras. Thus, aspects of the present invention include non-stochastic methods 20 of producing a set of finalized chimeric nucleic acid molecule shaving an overall assembly order that is chosen by design. This method includes the steps of generating by design a plurality of specific nucleic acid building blocks having serviceable mutually compatible ligatable ends, and assembling these nucleic acid building blocks, such that a designed overall assembly order is achieved.

25 The mutually compatible ligatable ends of the nucleic acid building blocks to be assembled are considered to be "serviceable" for this type of ordered assembly if they enable the building blocks to be coupled in predetermined orders. Thus, the overall assembly order in which the nucleic acid building blocks can be coupled is specified by the design of the ligatable ends. If more than one assembly step is to be used, then the 30 overall assembly order in which the nucleic acid building blocks can be coupled is also specified by the sequential order of the assembly step(s). In one aspect, the annealed building pieces are treated with an enzyme, such as a ligase (e.g. T4 DNA ligase), to achieve covalent bonding of the building pieces.

In one aspect, the design of the oligonucleotide building blocks is obtained by analyzing a set of progenitor nucleic acid sequence templates that serve as a basis for producing a progeny set of finalized chimeric polynucleotides. These parental oligonucleotide templates thus serve as a source of sequence information that aids in the design of the nucleic acid building blocks that are to be mutagenized, e.g., chimerized or shuffled. In one aspect of this method, the sequences of a plurality of parental nucleic acid templates are aligned in order to select one or more demarcation points. The demarcation points can be located at an area of homology, and are comprised of one or more nucleotides. These demarcation points are preferably shared by at least two of the progenitor templates. The demarcation points can thereby be used to delineate the boundaries of oligonucleotide building blocks to be generated in order to rearrange the parental polynucleotides. The demarcation points identified and selected in the progenitor molecules serve as potential chimerization points in the assembly of the final chimeric progeny molecules. A demarcation point can be an area of homology (comprised of at least one homologous nucleotide base) shared by at least two parental polynucleotide sequences. Alternatively, a demarcation point can be an area of homology that is shared by at least half of the parental polynucleotide sequences, or, it can be an area of homology that is shared by at least two thirds of the parental polynucleotide sequences. Even more preferably a serviceable demarcation points is an area of homology that is shared by at least three fourths of the parental polynucleotide sequences, or, it can be shared by at almost all of the parental polynucleotide sequences. In one aspect, a demarcation point is an area of homology that is shared by all of the parental polynucleotide sequences.

In one aspect, a ligation reassembly process is performed exhaustively in order to generate an exhaustive library of progeny chimeric polynucleotides. In other words, all possible ordered combinations of the nucleic acid building blocks are represented in the set of finalized chimeric nucleic acid molecules. At the same time, in another aspect, the assembly order (i.e. the order of assembly of each building block in the 5' to 3' sequence of each finalized chimeric nucleic acid) in each combination is by design (or non-stochastic) as described above. Because of the non-stochastic nature of this invention, the possibility of unwanted side products is greatly reduced.

In another aspect, the ligation reassembly method is performed systematically. For example, the method is performed in order to generate a systematically compartmentalized library of progeny molecules, with compartments that can be screened systematically, e.g. one by one. In other words this invention provides that, through the selective and

judicious use of specific nucleic acid building blocks, coupled with the selective and judicious use of sequentially stepped assembly reactions, a design can be achieved where specific sets of progeny products are made in each of several reaction vessels. This allows a systematic examination and screening procedure to be performed. Thus, these methods allow a potentially very large number of progeny molecules to be examined systematically in smaller groups. Because of its ability to perform chimerizations in a manner that is highly flexible yet exhaustive and systematic as well, particularly when there is a low level of homology among the progenitor molecules, these methods provide for the generation of a library (or set) comprised of a large number of progeny molecules. Because of the non-stochastic nature of the instant ligation reassembly invention, the progeny molecules generated preferably comprise a library of finalized chimeric nucleic acid molecules having an overall assembly order that is chosen by design. The saturation mutagenesis and optimized directed evolution methods also can be used to generate different progeny molecular species. It is appreciated that the invention provides freedom of choice and control regarding the selection of demarcation points, the size and number of the nucleic acid building blocks, and the size and design of the couplings. It is appreciated, furthermore, that the requirement for intermolecular homology is highly relaxed for the operability of this invention. In fact, demarcation points can even be chosen in areas of little or no intermolecular homology. For example, because of codon wobble, i.e. the degeneracy of codons, nucleotide substitutions can be introduced into nucleic acid building blocks without altering the amino acid originally encoded in the corresponding progenitor template. Alternatively, a codon can be altered such that the coding for an originally amino acid is altered. This invention provides that such substitutions can be introduced into the nucleic acid building block in order to increase the incidence of intermolecular homologous demarcation points and thus to allow an increased number of couplings to be achieved among the building blocks, which in turn allows a greater number of progeny chimeric molecules to be generated.

Synthetic gene reassembly

In one aspect, the present invention provides a non-stochastic method termed synthetic gene reassembly (e.g., GeneReassembly, see, e.g., U.S. Patent No. 6,537,776), which differs from stochastic shuffling in that the nucleic acid building blocks are not shuffled or concatenated or chimerized randomly, but rather are assembled non-stochastically.

The synthetic gene reassembly method does not depend on the presence of a high level of homology between polynucleotides to be shuffled. The invention can be used to non-stochastically generate libraries (or sets) of progeny molecules comprised of over 10^{100} different chimeras. Conceivably, synthetic gene reassembly can even be used to generate libraries comprised of over 10^{1000} different progeny chimeras.

Thus, in one aspect, the invention provides a non-stochastic method of producing a set of finalized chimeric nucleic acid molecules having an overall assembly order that is chosen by design, which method is comprised of the steps of generating by design a plurality of specific nucleic acid building blocks having serviceable mutually compatible ligatable ends and assembling these nucleic acid building blocks, such that a designed overall assembly order is achieved.

In one aspect, synthetic gene reassembly comprises a method of: 1) preparing a progeny generation of molecule(s) (including a molecule comprising a polynucleotide sequence, e.g., a molecule comprising a polypeptide coding sequence), that is mutagenized to achieve at least one point mutation, addition, deletion, &/or chimerization, from one or more ancestral or parental generation template(s); 2) screening the progeny generation molecule(s), e.g., using a high throughput method, for at least one property of interest (such as an improvement in an enzyme activity); 3) in one aspect obtaining &/or cataloguing structural &/or and functional information regarding the parental &/or progeny generation molecules; and 4) in one aspect repeating any of steps 1) to 3). In one aspect, there is generated (e.g., from a parent polynucleotide template), in what is termed "codon site-saturation mutagenesis," a progeny generation of polynucleotides, each having at least one set of up to three contiguous point mutations (i.e. different bases comprising a new codon), such that every codon (or every family of degenerate codons encoding the same amino acid) is represented at each codon position. Corresponding to, and encoded by, this progeny generation of polynucleotides, there is also generated a set of progeny polypeptides, each having at least one single amino acid point mutation. In a one aspect, there is generated, in what is termed "amino acid site-saturation mutagenesis", one such mutant polypeptide for each of the 19 naturally encoded polypeptide-forming alpha-amino acid substitutions at each and every amino acid position along the polypeptide. This yields, for each and every amino acid position along the parental polypeptide, a total of 20 distinct progeny polypeptides including the original amino acid, or potentially more than 21 distinct progeny polypeptides if additional amino acids are used either instead of or in addition to the 20 naturally encoded amino acids

Thus, in another aspect, this approach is also serviceable for generating mutants containing, in addition to &/or in combination with the 20 naturally encoded polypeptide-forming alpha-amino acids, other rare &/or not naturally-encoded amino acids and amino acid derivatives. In yet another aspect, this approach is also serviceable for generating
5 mutants by the use of, in addition to &/or in combination with natural or unaltered codon recognition systems of suitable hosts, altered, mutagenized, &/or designer codon recognition systems (such as in a host cell with one or more altered tRNA molecules.

In yet another aspect, this invention relates to recombination and more specifically to a method for preparing polynucleotides encoding a polypeptide by a method of *in vivo*
10 re-assortment of polynucleotide sequences containing regions of partial homology, assembling the polynucleotides to form at least one polynucleotide and screening the polynucleotides for the production of polypeptide(s) having a useful property.

In yet another aspect, this invention is serviceable for analyzing and cataloguing, with respect to any molecular property (e.g. an enzymatic activity) or combination of
15 properties allowed by current technology, the effects of any mutational change achieved (including particularly saturation mutagenesis). Thus, a comprehensive method is provided for determining the effect of changing each amino acid in a parental polypeptide into each of at least 19 possible substitutions. This allows each amino acid in a parental polypeptide to be characterized and catalogued according to its spectrum of potential
20 effects on a measurable property of the polypeptide.

In one aspect, an intron may be introduced into a chimeric progeny molecule by way of a nucleic acid building block. Introns often have consensus sequences at both termini in order to render them operational. In addition to enabling gene splicing, introns may serve an additional purpose by providing sites of homology to other nucleic acids to
25 enable homologous recombination. For this purpose, and potentially others, it may be sometimes desirable to generate a large nucleic acid building block for introducing an intron. If the size is overly large easily generating by direct chemical synthesis of two single stranded oligos, such a specialized nucleic acid building block may also be generated by direct chemical synthesis of more than two single stranded oligos or by using
30 a polymerase-based amplification reaction

The mutually compatible ligatable ends of the nucleic acid building blocks to be assembled are considered to be "serviceable" for this type of ordered assembly if they enable the building blocks to be coupled in predetermined orders. Thus, in one aspect, the overall assembly order in which the nucleic acid building blocks can be coupled is

specified by the design of the ligatable ends and, if more than one assembly step is to be used, then the overall assembly order in which the nucleic acid building blocks can be coupled is also specified by the sequential order of the assembly step(s). In a one aspect of the invention, the annealed building pieces are treated with an enzyme, such as a ligase
5 (e.g., T4 DNA ligase) to achieve covalent bonding of the building pieces.

Coupling can occur in a manner that does not make use of every nucleotide in a participating overhang. The coupling is particularly lively to survive (e.g. in a transformed host) if the coupling reinforced by treatment with a ligase enzyme to form what may be referred to as a “gap ligation” or a “gapped ligation”. This type of coupling
10 can contribute to generation of unwanted background product(s), but it can also be used advantageously increase the diversity of the progeny library generated by the designed ligation reassembly. Certain overhangs are able to undergo self-coupling to form a palindromic coupling. A coupling is strengthened substantially if it is reinforced by treatment with a ligase enzyme. Lack of 5’ phosphates on these overhangs can be used
15 advantageously to prevent this type of palindromic self-ligation. Accordingly, this invention provides that nucleic acid building blocks can be chemically made (or ordered) that lack a 5’ phosphate group. Alternatively, they can be removed, e.g. by treatment with a phosphatase enzyme, such as a calf intestinal alkaline phosphatase (CIAP), in order to prevent palindromic self-ligations in ligation reassembly processes.

20 In a another aspect, the design of nucleic acid building blocks is obtained upon analysis of the sequences of a set of progenitor nucleic acid templates that serve as a basis for producing a progeny set of finalized chimeric nucleic acid molecules. These progenitor nucleic acid templates thus serve as a source of sequence information that aids in the design of the nucleic acid building blocks that are to be mutagenized, *i.e.* chimerized
25 or shuffled.

In one exemplification, the invention provides for the chimerization of a family of related genes and their encoded family of related products. In a particular exemplification, the encoded products are enzymes. The isomerases, e.g., racemases, e.g., amino acid racemases, alanine racemases, and/or epimerases of the present invention can be
30 mutagenized in accordance with the methods described herein.

Thus according to one aspect of the invention, the sequences of a plurality of progenitor nucleic acid templates (e.g., polynucleotides of The invention) are aligned in order to select one or more demarcation points, which demarcation points can be located at an area of homology. The demarcation points can be used to delineate the boundaries of

nucleic acid building blocks to be generated. Thus, the demarcation points identified and selected in the progenitor molecules serve as potential chimerization points in the assembly of the progeny molecules.

Typically a serviceable demarcation point is an area of homology (comprised of at least one homologous nucleotide base) shared by at least two progenitor templates, but the demarcation point can be an area of homology that is shared by at least half of the progenitor templates, at least two thirds of the progenitor templates, at least three fourths of the progenitor templates and preferably at almost all of the progenitor templates. Even more preferably still a serviceable demarcation point is an area of homology that is shared by all of the progenitor templates.

In a one aspect, the gene reassembly process is performed exhaustively in order to generate an exhaustive library. In other words, all possible ordered combinations of the nucleic acid building blocks are represented in the set of finalized chimeric nucleic acid molecules. At the same time, the assembly order (*i.e.* the order of assembly of each building block in the 5' to 3' sequence of each finalized chimeric nucleic acid) in each combination is by design (or non-stochastic). Because of the non-stochastic nature of the method, the possibility of unwanted side products is greatly reduced.

In another aspect, the method provides that the gene reassembly process is performed systematically, for example to generate a systematically compartmentalized library, with compartments that can be screened systematically, *e.g.*, one by one. In other words the invention provides that, through the selective and judicious use of specific nucleic acid building blocks, coupled with the selective and judicious use of sequentially stepped assembly reactions, an experimental design can be achieved where specific sets of progeny products are made in each of several reaction vessels. This allows a systematic examination and screening procedure to be performed. Thus, it allows a potentially very large number of progeny molecules to be examined systematically in smaller groups.

Because of its ability to perform chimerizations in a manner that is highly flexible yet exhaustive and systematic as well, particularly when there is a low level of homology among the progenitor molecules, the instant invention provides for the generation of a library (or set) comprised of a large number of progeny molecules. Because of the non-stochastic nature of the instant gene reassembly invention, the progeny molecules generated preferably comprise a library of finalized chimeric nucleic acid molecules having an overall assembly order that is chosen by design. In a particularly aspect, such a

generated library is comprised of greater than 10^3 to greater than 10^{1000} different progeny molecular species.

In one aspect, a set of finalized chimeric nucleic acid molecules, produced as described is comprised of a polynucleotide encoding a polypeptide. According to one aspect, this polynucleotide is a gene, which may be a man-made gene. According to another aspect, this polynucleotide is a gene pathway, which may be a man-made gene pathway. The invention provides that one or more man-made genes generated by the invention may be incorporated into a man-made gene pathway, such as pathway operable in a eukaryotic organism (including a plant).

In another exemplification, the synthetic nature of the step in which the building blocks are generated allows the design and introduction of nucleotides (*e.g.*, one or more nucleotides, which may be, for example, codons or introns or regulatory sequences) that can later be in one aspect removed in an *in vitro* process (*e.g.*, by mutagenesis) or in an *in vivo* process (*e.g.*, by utilizing the gene splicing ability of a host organism). It is appreciated that in many instances the introduction of these nucleotides may also be desirable for many other reasons in addition to the potential benefit of creating a serviceable demarcation point.

Thus, according to another aspect, the invention provides that a nucleic acid building block can be used to introduce an intron. Thus, the invention provides that functional introns may be introduced into a man-made gene of the invention. The invention also provides that functional introns may be introduced into a man-made gene pathway of the invention. Accordingly, the invention provides for the generation of a chimeric polynucleotide that is a man-made gene containing one (or more) artificially introduced intron(s).

Accordingly, the invention also provides for the generation of a chimeric polynucleotide that is a man-made gene pathway containing one (or more) artificially introduced intron(s). Preferably, the artificially introduced intron(s) are functional in one or more host cells for gene splicing much in the way that naturally-occurring introns serve functionally in gene splicing. The invention provides a process of producing man-made intron-containing polynucleotides to be introduced into host organisms for recombination and/or splicing.

A man-made gene produced using the invention can also serve as a substrate for recombination with another nucleic acid. Likewise, a man-made gene pathway produced using the invention can also serve as a substrate for recombination with another nucleic

acid. In a one aspect, the recombination is facilitated by, or occurs at, areas of homology between the man-made, intron-containing gene and a nucleic acid, which serves as a recombination partner. In one aspect, the recombination partner may also be a nucleic acid generated by the invention, including a man-made gene or a man-made gene pathway.

5 Recombination may be facilitated by or may occur at areas of homology that exist at the one (or more) artificially introduced intron(s) in the man-made gene.

The synthetic gene reassembly method of the invention utilizes a plurality of nucleic acid building blocks, each of which preferably has two ligatable ends. The two ligatable ends on each nucleic acid building block may be two blunt ends (*i.e.* each having an overhang of zero nucleotides), or preferably one blunt end and one overhang, or more preferably still two overhangs.

A useful overhang for this purpose may be a 3' overhang or a 5' overhang. Thus, a nucleic acid building block may have a 3' overhang or alternatively a 5' overhang or alternatively two 3' overhangs or alternatively two 5' overhangs. The overall order in which the nucleic acid building blocks are assembled to form a finalized chimeric nucleic acid molecule is determined by purposeful experimental design and is not random.

In one aspect, a nucleic acid building block is generated by chemical synthesis of two single-stranded nucleic acids (also referred to as single-stranded oligos) and contacting them so as to allow them to anneal to form a double-stranded nucleic acid building block.

A double-stranded nucleic acid building block can be of variable size. The sizes of these building blocks can be small or large. Exemplary sizes for building block range from 1 base pair (not including any overhangs) to 100,000 base pairs (not including any overhangs). Other exemplary size ranges are also provided, which have lower limits of from 1 bp to 10,000 bp (including every integer value in between) and upper limits of from 2 bp to 100,000 bp (including every integer value in between).

Many methods exist by which a double-stranded nucleic acid building block can be generated that is serviceable for the invention; and these are known in the art and can be readily performed by the skilled artisan.

30 According to one aspect, a double-stranded nucleic acid building block is generated by first generating two single stranded nucleic acids and allowing them to anneal to form a double-stranded nucleic acid building block. The two strands of a double-stranded nucleic acid building block may be complementary at every nucleotide apart from any that form an overhang; thus containing no mismatches, apart from any

overhang(s). According to another aspect, the two strands of a double-stranded nucleic acid building block are complementary at fewer than every nucleotide apart from any that form an overhang. Thus, according to this aspect, a double-stranded nucleic acid building block can be used to introduce codon degeneracy. The codon degeneracy can be
5 introduced using the site-saturation mutagenesis described herein, using one or more N,N,G/T cassettes or alternatively using one or more N,N,N cassettes.

The *in vivo* recombination method of the invention can be performed blindly on a pool of unknown hybrids or alleles of a specific polynucleotide or sequence. However, it is not necessary to know the actual DNA or RNA sequence of the specific polynucleotide.

10 The approach of using recombination within a mixed population of genes can be useful for the generation of any useful proteins, for example, interleukin I, antibodies, tPA and growth hormone. This approach may be used to generate proteins having altered specificity or activity. The approach may also be useful for the generation of hybrid nucleic acid sequences, for example, promoter regions, introns, exons, enhancer
15 sequences, 3' untranslated regions or 5' untranslated regions of genes. Thus this approach may be used to generate genes having increased rates of expression. This approach may also be useful in the study of repetitive DNA sequences. Finally, this approach may be useful to mutate ribozymes or aptamers.

In one aspect the invention described herein is directed to the use of repeated
20 cycles of reductive reassortment, recombination and selection which allow for the directed molecular evolution of highly complex linear sequences, such as DNA, RNA or proteins thorough recombination.

Optimized Directed Evolution System

The invention provides a non-stochastic gene modification system termed
25 "optimized directed evolution system" to generate polypeptides, e.g., isomerases, e.g., racemases, e.g., amino acid racemases, alanine racemases, and/or epimerases, or antibodies of the invention, with new or altered properties. Optimized directed evolution is directed to the use of repeated cycles of reductive reassortment, recombination and selection that allow for the directed molecular evolution of nucleic acids through
30 recombination. Optimized directed evolution allows generation of a large population of evolved chimeric sequences, wherein the generated population is significantly enriched for sequences that have a predetermined number of crossover events.

A crossover event is a point in a chimeric sequence where a shift in sequence occurs from one parental variant to another parental variant. Such a point is normally at the juncture of where oligonucleotides from two parents are ligated together to form a single sequence. This method allows calculation of the correct concentrations of oligonucleotide sequences so that the final chimeric population of sequences is enriched for the chosen number of crossover events. This provides more control over choosing chimeric variants having a predetermined number of crossover events.

In addition, this method provides a convenient means for exploring a tremendous amount of the possible protein variant space in comparison to other systems. Previously, if one generated, for example, 10^{13} chimeric molecules during a reaction, it would be extremely difficult to test such a high number of chimeric variants for a particular activity. Moreover, a significant portion of the progeny population would have a very high number of crossover events which resulted in proteins that were less likely to have increased levels of a particular activity. By using these methods, the population of chimeric molecules can be enriched for those variants that have a particular number of crossover events. Thus, although one can still generate 10^{13} chimeric molecules during a reaction, each of the molecules chosen for further analysis most likely has, for example, only three crossover events. Because the resulting progeny population can be skewed to have a predetermined number of crossover events, the boundaries on the functional variety between the chimeric molecules is reduced. This provides a more manageable number of variables when calculating which oligonucleotide from the original parental polynucleotides might be responsible for affecting a particular trait.

One method for creating a chimeric progeny polynucleotide sequence is to create oligonucleotides corresponding to fragments or portions of each parental sequence. Each oligonucleotide preferably includes a unique region of overlap so that mixing the oligonucleotides together results in a new variant that has each oligonucleotide fragment assembled in the correct order. Additional information can also be found, e.g., in USSN 09/332,835; U.S. Patent No. 6,361,974.

The number of oligonucleotides generated for each parental variant bears a relationship to the total number of resulting crossovers in the chimeric molecule that is ultimately created. For example, three parental nucleotide sequence variants might be provided to undergo a ligation reaction in order to find a chimeric variant having, for example, greater activity at high temperature. As one example, a set of 50 oligonucleotide sequences can be generated corresponding to each portions of each parental variant.

Accordingly, during the ligation reassembly process there could be up to 50 crossover events within each of the chimeric sequences. The probability that each of the generated chimeric polynucleotides will contain oligonucleotides from each parental variant in alternating order is very low. If each oligonucleotide fragment is present in the ligation
5 reaction in the same molar quantity it is likely that in some positions oligonucleotides from the same parental polynucleotide will ligate next to one another and thus not result in a crossover event. If the concentration of each oligonucleotide from each parent is kept constant during any ligation step in this example, there is a 1/3 chance (assuming 3 parents) that an oligonucleotide from the same parental variant will ligate within the
10 chimeric sequence and produce no crossover.

Accordingly, a probability density function (PDF) can be determined to predict the population of crossover events that are likely to occur during each step in a ligation reaction given a set number of parental variants, a number of oligonucleotides corresponding to each variant, and the concentrations of each variant during each step in
15 the ligation reaction. The statistics and mathematics behind determining the PDF is described below. By utilizing these methods, one can calculate such a probability density function, and thus enrich the chimeric progeny population for a predetermined number of crossover events resulting from a particular ligation reaction. Moreover, a target number of crossover events can be predetermined, and the system then programmed to calculate
20 the starting quantities of each parental oligonucleotide during each step in the ligation reaction to result in a probability density function that centers on the predetermined number of crossover events. These methods are directed to the use of repeated cycles of reductive reassortment, recombination and selection that allow for the directed molecular evolution of a nucleic acid encoding a polypeptide through recombination. This system
25 allows generation of a large population of evolved chimeric sequences, wherein the generated population is significantly enriched for sequences that have a predetermined number of crossover events. A crossover event is a point in a chimeric sequence where a shift in sequence occurs from one parental variant to another parental variant. Such a point is normally at the juncture of where oligonucleotides from two parents are ligated
30 together to form a single sequence. The method allows calculation of the correct concentrations of oligonucleotide sequences so that the final chimeric population of sequences is enriched for the chosen number of crossover events. This provides more control over choosing chimeric variants having a predetermined number of crossover events.

In addition, these methods provide a convenient means for exploring a tremendous amount of the possible protein variant space in comparison to other systems. By using the methods described herein, the population of chimeric molecules can be enriched for those variants that have a particular number of crossover events. Thus, although one can still
5 generate 10^{13} chimeric molecules during a reaction, each of the molecules chosen for further analysis most likely has, for example, only three crossover events. Because the resulting progeny population can be skewed to have a predetermined number of crossover events, the boundaries on the functional variety between the chimeric molecules is reduced. This provides a more manageable number of variables when calculating which
10 oligonucleotide from the original parental polynucleotides might be responsible for affecting a particular trait.

In one aspect, the method creates a chimeric progeny polynucleotide sequence by creating oligonucleotides corresponding to fragments or portions of each parental sequence. Each oligonucleotide preferably includes a unique region of overlap so that
15 mixing the oligonucleotides together results in a new variant that has each oligonucleotide fragment assembled in the correct order. See also USSN 09/332,835.

Determining Crossover Events

Aspects of the invention include a system and software that receive a desired crossover probability density function (PDF), the number of parent genes to be
20 reassembled, and the number of fragments in the reassembly as inputs. The output of this program is a "fragment PDF" that can be used to determine a recipe for producing reassembled genes, and the estimated crossover PDF of those genes. The processing described herein is preferably performed in MATLAB™ (The Mathworks, Natick, Massachusetts) a programming language and development environment for technical
25 computing.

Iterative Processes

In practicing the invention, these processes can be iteratively repeated. For example, a nucleic acid (or, the nucleic acid) responsible for an altered or new isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase phenotype
30 is identified, re-isolated, again modified, re-tested for activity. This process can be iteratively repeated until a desired phenotype is engineered. For example, an entire biochemical anabolic or catabolic pathway can be engineered into a cell, including, e.g.,

isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase activity.

Similarly, if it is determined that a particular oligonucleotide has no affect at all on the desired trait (e.g., a new isomerase, e.g., racemase, e.g., amino acid racemase, alanine
5 racemase, and/or epimerase phenotype), it can be removed as a variable by synthesizing larger parental oligonucleotides that include the sequence to be removed. Since incorporating the sequence within a larger sequence prevents any crossover events, there will no longer be any variation of this sequence in the progeny polynucleotides. This iterative practice of determining which oligonucleotides are most related to the desired
10 trait, and which are unrelated, allows more efficient exploration all of the possible protein variants that might be provide a particular trait or activity.

In vivo shuffling

In vivo shuffling of molecules is use in methods of the invention that provide variants of polypeptides of the invention, e.g., antibodies, isomerases, e.g.,
15 racemases, e.g., amino acid racemases, alanine racemases, and/or epimerases, and the like. *In vivo* shuffling can be performed utilizing the natural property of cells to recombine multimers. While recombination *in vivo* has provided the major natural route to molecular diversity, genetic recombination remains a relatively complex process that involves 1) the recognition of homologies; 2) strand cleavage, strand invasion, and metabolic steps
20 leading to the production of recombinant chiasma; and finally 3) the resolution of chiasma into discrete recombined molecules. The formation of the chiasma requires the recognition of homologous sequences.

In another aspect, the invention includes a method for producing a hybrid polynucleotide from at least a first polynucleotide and a second polynucleotide. The
25 invention can be used to produce a hybrid polynucleotide by introducing at least a first polynucleotide and a second polynucleotide which share at least one region of partial sequence homology into a suitable host cell. The regions of partial sequence homology promote processes which result in sequence reorganization producing a hybrid polynucleotide. The term “hybrid polynucleotide”, as used herein, is any nucleotide
30 sequence which results from the method of the present invention and contains sequence from at least two original polynucleotide sequences. Such hybrid polynucleotides can result from intermolecular recombination events which promote sequence integration between DNA molecules. In addition, such hybrid polynucleotides can result from

intramolecular reductive reassortment processes which utilize repeated sequences to alter a nucleotide sequence within a DNA molecule.

In *vivo* reassortment is focused on “inter-molecular” processes collectively referred to as “recombination” which in bacteria, is generally viewed as a “RecA-
5 dependent” phenomenon. The invention can rely on recombination processes of a host cell to recombine and re-assort sequences, or the cells’ ability to mediate reductive processes to decrease the complexity of quasi-repeated sequences in the cell by deletion. This process of “reductive reassortment” occurs by an “intra-molecular”, RecA-independent process.

10 Therefore, in another aspect of the invention, novel polynucleotides can be generated by the process of reductive reassortment. The method involves the generation of constructs containing consecutive sequences (original encoding sequences), their insertion into an appropriate vector and their subsequent introduction into an appropriate host cell. The reassortment of the individual molecular identities occurs by combinatorial
15 processes between the consecutive sequences in the construct possessing regions of homology, or between quasi-repeated units. The reassortment process recombines and/or reduces the complexity and extent of the repeated sequences and results in the production of novel molecular species. Various treatments may be applied to enhance the rate of reassortment. These could include treatment with ultra-violet light, or DNA damaging
20 chemicals and/or the use of host cell lines displaying enhanced levels of “genetic instability”. Thus the reassortment process may involve homologous recombination or the natural property of quasi-repeated sequences to direct their own evolution.

Repeated or “quasi-repeated” sequences play a role in genetic instability. In the present invention, “quasi-repeats” are repeats that are not restricted to their original unit
25 structure. Quasi-repeated units can be presented as an array of sequences in a construct; consecutive units of similar sequences. Once ligated, the junctions between the consecutive sequences become essentially invisible and the quasi-repetitive nature of the resulting construct is now continuous at the molecular level. The deletion process the cell performs to reduce the complexity of the resulting construct operates between the quasi-
30 repeated sequences. The quasi-repeated units provide a practically limitless repertoire of templates upon which slippage events can occur. The constructs containing the quasi-repeats thus effectively provide sufficient molecular elasticity that deletion (and potentially insertion) events can occur virtually anywhere within the quasi-repetitive units.

When the quasi-repeated sequences are all ligated in the same orientation, for instance head to tail or vice versa, the cell cannot distinguish individual units. Consequently, the reductive process can occur throughout the sequences. In contrast, when for example, the units are presented head to head, rather than head to tail, the inversion delineates the endpoints of the adjacent unit so that deletion formation will favor the loss of discrete units. Thus, it is preferable with the present method that the sequences are in the same orientation. Random orientation of quasi-repeated sequences will result in the loss of reassortment efficiency, while consistent orientation of the sequences will offer the highest efficiency. However, while having fewer of the contiguous sequences in the same orientation decreases the efficiency, it may still provide sufficient elasticity for the effective recovery of novel molecules. Constructs can be made with the quasi-repeated sequences in the same orientation to allow higher efficiency.

Sequences can be assembled in a head to tail orientation using any of a variety of methods, including the following:

- 15 a) Primers that include a poly-A head and poly-T tail which when made single-stranded would provide orientation can be utilized. This is accomplished by having the first few bases of the primers made from RNA and hence easily removed RNaseH.
- b) Primers that include unique restriction cleavage sites can be utilized.
20 Multiple sites, a battery of unique sequences and repeated synthesis and ligation steps would be required.
- c) The inner few bases of the primer could be thiolated and an exonuclease used to produce properly tailed molecules.

The recovery of the re-assorted sequences relies on the identification of cloning vectors with a reduced repetitive index (RI). The re-assorted encoding sequences can then be recovered by amplification. The products are re-cloned and expressed. The recovery of cloning vectors with reduced RI can be affected by:

- 1) The use of vectors only stably maintained when the construct is reduced in complexity.
- 30 2) The physical recovery of shortened vectors by physical procedures. In this case, the cloning vector would be recovered using standard plasmid isolation procedures and size fractionated on either an agarose gel, or column with a low molecular weight cut off utilizing standard procedures.

- 3) The recovery of vectors containing interrupted genes which can be selected when insert size decreases.
- 4) The use of direct selection techniques with an expression vector and the appropriate selection.

5 Encoding sequences (for example, genes) from related organisms may demonstrate a high degree of homology and encode quite diverse protein products. These types of sequences are particularly useful in the present invention as quasi-repeats. However, while the examples illustrated below demonstrate the reassortment of nearly identical original encoding sequences (quasi-repeats), this process is not limited to such nearly
10 identical repeats.

The following example demonstrates a method of the invention. Encoding nucleic acid sequences (quasi-repeats) derived from three (3) unique species are described. Each sequence encodes a protein with a distinct set of properties. Each of the sequences differs by a single or a few base pairs at a unique position in the sequence. The quasi-repeated
15 sequences are separately or collectively amplified and ligated into random assemblies such that all possible permutations and combinations are available in the population of ligated molecules. The number of quasi-repeat units can be controlled by the assembly conditions. The average number of quasi-repeated units in a construct is defined as the repetitive index (RI).

20 Once formed, the constructs may, or may not be size fractionated on an agarose gel according to published protocols, inserted into a cloning vector and transfected into an appropriate host cell. The cells are then propagated and “reductive reassortment” is effected. The rate of the reductive reassortment process may be stimulated by the introduction of DNA damage if desired. Whether the reduction in RI is mediated by
25 deletion formation between repeated sequences by an “intra-molecular” mechanism, or mediated by recombination-like events through “inter-molecular” mechanisms is immaterial. The end result is a reassortment of the molecules into all possible combinations.

In one aspect (optionally), the method comprises the additional step of screening
30 the library members of the shuffled pool to identify individual shuffled library members having the ability to bind or otherwise interact, or catalyze a particular reaction (*e.g.*, such as catalytic domain of an enzyme) with a predetermined macromolecule, such as for

example a proteinaceous receptor, an oligosaccharide, virion, or other predetermined compound or structure.

The polypeptides that are identified from such libraries can be used for therapeutic, diagnostic, research and related purposes (*e.g.*, catalysts, solutes for increasing osmolarity of an aqueous solution and the like) and/or can be subjected to one or more additional
5 cycles of shuffling and/or selection.

In another aspect, it is envisioned that prior to or during recombination or reassortment, polynucleotides generated by the method of the invention can be subjected to agents or processes which promote the introduction of mutations into the original
10 polynucleotides. The introduction of such mutations would increase the diversity of resulting hybrid polynucleotides and polypeptides encoded therefrom. The agents or processes which promote mutagenesis can include, but are not limited to: (+)-CC-1065, or a synthetic analog such as (+)-CC-1065-(N3-Adenine (*See* Sun and Hurley, (1992); an N-acetylated or deacetylated 4'-fluro-4-aminobiphenyl adduct capable of inhibiting DNA
15 synthesis (*See* , for example, van de Poll *et al.* (1992)); or a N-acetylated or deacetylated 4-aminobiphenyl adduct capable of inhibiting DNA synthesis (*See* also, van de Poll *et al.* (1992), pp. 751-758); trivalent chromium, a trivalent chromium salt, a polycyclic aromatic hydrocarbon (PAH) DNA adduct capable of inhibiting DNA replication, such as 7-bromomethyl-benz[*a*]anthracene ("BMA"), tris(2,3-dibromopropyl)phosphate ("Tris-BP"),
20 BP"), 1,2-dibromo-3-chloropropane ("DBCP"), 2-bromoacrolein (2BA), benzo[*a*]pyrene-7,8-dihydrodiol-9-10-epoxide ("BPDE"), a platinum(II) halogen salt, N-hydroxy-2-amino-3-methylimidazo[4,5-*f*]-quinoline ("N-hydroxy-IQ") and N-hydroxy-2-amino-1-methyl-6-phenylimidazo[4,5-*f*]-pyridine ("N-hydroxy-PhIP"). Exemplary means for slowing or halting PCR amplification consist of UV light (+)-CC-1065 and (+)-CC-1065-(N3-
25 Adenine). Particularly encompassed means are DNA adducts or polynucleotides comprising the DNA adducts from the polynucleotides or polynucleotides pool, which can be released or removed by a process including heating the solution comprising the polynucleotides prior to further processing.

In another aspect the invention is directed to a method of producing recombinant
30 proteins having biological activity by treating a sample comprising double-stranded template polynucleotides encoding a wild-type protein under conditions according to the invention which provide for the production of hybrid or re-assorted polynucleotides.

Producing sequence variants

The invention also provides additional methods for making sequence variants of the nucleic acid (e.g., isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase) sequences of the invention. The invention also provides
5 additional methods for isolating isomerases, e.g., racemases, e.g., amino acid racemases, alanine racemases, and/or epimerases using the nucleic acids and polypeptides of the invention. In one aspect, the invention provides for variants of an isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase coding sequence (e.g., a gene, cDNA or message) of the invention, which can be altered by any means,
10 including, e.g., random or stochastic methods, or, non-stochastic, or “directed evolution,” methods, as described above.

The isolated variants may be naturally occurring. Variant can also be created *in vitro*. Variants may be created using genetic engineering techniques such as site directed mutagenesis, random chemical mutagenesis, Exonuclease III deletion procedures, and
15 standard cloning techniques. Alternatively, such variants, fragments, analogs, or derivatives may be created using chemical synthesis or modification procedures. Other methods of making variants are also familiar to those skilled in the art. These include procedures in which nucleic acid sequences obtained from natural isolates are modified to generate new nucleic acids which encode polypeptides having characteristics which
20 enhance their value in industrial, medical, laboratory (research), pharmaceutical, food and feed and food and feed supplement processing and other applications and processes. In such procedures, a large number of variant sequences having one or more nucleotide differences with respect to the sequence obtained from the natural isolate are generated and characterized. These nucleotide differences can result in amino acid changes with
25 respect to the polypeptides encoded by the nucleic acids from the natural isolates.

For example, variants may be created using error prone PCR. In error prone PCR, PCR is performed under conditions where the copying fidelity of the DNA polymerase is low, such that a high rate of point mutations is obtained along the entire length of the PCR product. Error prone PCR is described, e.g., in Leung, D.W., et al., *Technique*, 1:11-15,
30 1989) and Caldwell, R. C. & Joyce G.F., *PCR Methods Applic.*, 2:28-33, 1992. Briefly, in such procedures, nucleic acids to be mutagenized are mixed with PCR primers, reaction buffer, MgCl₂, MnCl₂, Taq polymerase and an appropriate concentration of dNTPs for achieving a high rate of point mutation along the entire length of the PCR product. For example, the reaction may be performed using 20 fmoles of nucleic acid to be

mutagenized, 30 pmole of each PCR primer, a reaction buffer comprising 50mM KCl, 10mM Tris HCl (pH 8.3) and 0.01% gelatin, 7mM MgCl₂, 0.5mM MnCl₂, 5 units of Taq polymerase, 0.2mM dGTP, 0.2mM dATP, 1mM dCTP, and 1mM dTTP. PCR may be performed for 30 cycles of 94°C for 1 min, 45°C for 1 min, and 72°C for 1 min. However, it will be appreciated that these parameters may be varied as appropriate. The mutagenized nucleic acids are cloned into an appropriate vector and the activities of the polypeptides encoded by the mutagenized nucleic acids are evaluated.

Variants may also be created using oligonucleotide directed mutagenesis to generate site-specific mutations in any cloned DNA of interest. Oligonucleotide mutagenesis is described, e.g., in Reidhaar-Olson (1988) Science 241:53-57. Briefly, in such procedures a plurality of double stranded oligonucleotides bearing one or more mutations to be introduced into the cloned DNA are synthesized and inserted into the cloned DNA to be mutagenized. Clones containing the mutagenized DNA are recovered and the activities of the polypeptides they encode are assessed.

Another method for generating variants is assembly PCR. Assembly PCR involves the assembly of a PCR product from a mixture of small DNA fragments. A large number of different PCR reactions occur in parallel in the same vial, with the products of one reaction priming the products of another reaction. Assembly PCR is described in, e.g., U.S. Patent No. 5,965,408.

Still another method of generating variants is sexual PCR mutagenesis. In sexual PCR mutagenesis, forced homologous recombination occurs between DNA molecules of different but highly related DNA sequence *in vitro*, as a result of random fragmentation of the DNA molecule based on sequence homology, followed by fixation of the crossover by primer extension in a PCR reaction. Sexual PCR mutagenesis is described, e.g., in Stemmer (1994) Proc. Natl. Acad. Sci. USA 91:10747-10751. Briefly, in such procedures a plurality of nucleic acids to be recombined are digested with DNase to generate fragments having an average size of 50-200 nucleotides. Fragments of the desired average size are purified and resuspended in a PCR mixture. PCR is conducted under conditions which facilitate recombination between the nucleic acid fragments. For example, PCR may be performed by resuspending the purified fragments at a concentration of 10-30 ng/μl in a solution of 0.2mM of each dNTP, 2.2mM MgCl₂, 50mM KCL, 10mM Tris HCl, pH 9.0, and 0.1% Triton X-100. 2.5 units of Taq polymerase per 100:1 of reaction mixture is added and PCR is performed using the following regime: 94°C for 60 seconds, 94°C for 30 seconds, 50-55°C for 30 seconds, 72°C for 30 seconds (30-45

times) and 72°C for 5 minutes. However, it will be appreciated that these parameters may be varied as appropriate. In some aspects, oligonucleotides may be included in the PCR reactions. In other aspects, the Klenow fragment of DNA polymerase I may be used in a first set of PCR reactions and Taq polymerase may be used in a subsequent set of PCR
5 reactions. Recombinant sequences are isolated and the activities of the polypeptides they encode are assessed.

Variants may also be created by *in vivo* mutagenesis. In some aspects, random mutations in a sequence of interest are generated by propagating the sequence of interest in a bacterial strain, such as an E. coli strain, which carries mutations in one or
10 more of the DNA repair pathways. Such “mutator” strains have a higher random mutation rate than that of a wild-type parent. Propagating the DNA in one of these strains will eventually generate random mutations within the DNA. Mutator strains suitable for use for *in vivo* mutagenesis are described in PCT Publication No. WO 91/16427, published October 31, 1991, entitled “Methods for Phenotype Creation from Multiple Gene
15 Populations”.

Variants may also be generated using cassette mutagenesis. In cassette mutagenesis a small region of a double stranded DNA molecule is replaced with a synthetic oligonucleotide “cassette” that differs from the native sequence. The oligonucleotide often contains completely and/or partially randomized native sequence.

20 Recursive ensemble mutagenesis may also be used to generate variants. Recursive ensemble mutagenesis is an algorithm for protein engineering (protein mutagenesis) developed to produce diverse populations of phenotypically related mutants whose members differ in amino acid sequence. This method uses a feedback mechanism to control successive rounds of combinatorial cassette mutagenesis. Recursive ensemble
25 mutagenesis is described in Arkin, A.P. and Youvan, D.C., PNAS, USA, 89:7811-7815, 1992.

In some aspects, variants are created using exponential ensemble mutagenesis. Exponential ensemble mutagenesis is a process for generating combinatorial libraries with a high percentage of unique and functional mutants, wherein small groups of
30 residues are randomized in parallel to identify, at each altered position, amino acids which lead to functional proteins. Exponential ensemble mutagenesis is described in Delegrave, S. and Youvan, D.C., Biotechnology Research, 11:1548-1552, 1993. Random and site-directed mutagenesis are described in Arnold, F.H., Current Opinion in Biotechnology, 4:450-455, 1993.

In some aspects, the variants are created using shuffling procedures wherein portions of a plurality of nucleic acids which encode distinct polypeptides are fused together to create chimeric nucleic acid sequences which encode chimeric polypeptides as described in U.S. Patent No. 5,965,408, filed July 9, 1996, entitled,
5 “Method of DNA Reassembly by Interrupting Synthesis” and U.S. Patent No. 5,939,250, filed May 22, 1996, entitled, “Production of Enzymes Having Desired Activities by Mutagenesis.

The variants of the polypeptides of the invention may be variants in which one or more of the amino acid residues of the polypeptides of the invention are substituted
10 with a conserved or non-conserved amino acid residue (preferably a conserved amino acid residue) and such substituted amino acid residue may or may not be one encoded by the genetic code.

Conservative substitutions are those that substitute a given amino acid in a polypeptide by another amino acid of like characteristics. Typically seen as conservative
15 substitutions are the following replacements: replacements of an aliphatic amino acid such as Alanine, Valine, Leucine and Isoleucine with another aliphatic amino acid; replacement of a Serine with a Threonine or vice versa; replacement of an acidic residue such as Aspartic acid and Glutamic acid with another acidic residue; replacement of a residue bearing an amide group, such as Asparagine and Glutamine, with another residue bearing
20 an amide group; exchange of a basic residue such as Lysine and Arginine with another basic residue; and replacement of an aromatic residue such as Phenylalanine, Tyrosine with another aromatic residue.

Other variants are those in which one or more of the amino acid residues of the polypeptides of the invention includes a substituent group.

25 Still other variants are those in which the polypeptide is associated with another compound, such as a compound to increase the half-life of the polypeptide (for example, polyethylene glycol).

Additional variants are those in which additional amino acids are fused to the polypeptide, such as a leader sequence, a secretory sequence, a proprotein sequence or
30 a sequence which facilitates purification, enrichment, or stabilization of the polypeptide.

In some aspects, the fragments, derivatives and analogs retain the same biological function or activity as the polypeptides of the invention and sequences substantially identical thereto. In other aspects, the fragment, derivative, or analog

includes a proprotein, such that the fragment, derivative, or analog can be activated by cleavage of the proprotein portion to produce an active polypeptide.

Optimizing codons to achieve high levels of protein expression in host cells

The invention provides methods for modifying an isomerase-, e.g., a racemase-,
5 e.g., an amino acid racemase-, an alanine racemase-, and/or an epimerase-isomerase-, e.g.,
racemase-, e.g., amino acid racemase-, alanine racemase-, and/or epimerase-encoding
nucleic acids to modify codon usage. In one aspect, the invention provides methods for
modifying codons in a nucleic acid encoding an isomerase, e.g., racemase, e.g., amino acid
racemase, alanine racemase, and/or epimerase to increase or decrease its expression in a
10 host cell. The invention also provides nucleic acids encoding an isomerase, e.g.,
racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase modified to
increase its expression in a host cell, enzymes so modified, and methods of making the
modified enzymes. The method comprises identifying a “non-preferred” or a “less
preferred” codon in an isomerase-, e.g., racemase-, e.g., amino acid racemase-, alanine
15 racemase-, and/or epimerase-encoding nucleic acid and replacing one or more of these
non-preferred or less preferred codons with a “preferred codon” encoding the same amino
acid as the replaced codon and at least one non-preferred or less preferred codon in the
nucleic acid has been replaced by a preferred codon encoding the same amino acid. A
preferred codon is a codon over-represented in coding sequences in genes in the host cell
20 and a non-preferred or less preferred codon is a codon under-represented in coding
sequences in genes in the host cell.

Host cells for expressing the nucleic acids, expression cassettes and vectors of the
invention include bacteria, yeast, fungi, plant cells, insect cells and mammalian cells.
Thus, the invention provides methods for optimizing codon usage in all of these cells,
25 codon-altered nucleic acids and polypeptides made by the codon-altered nucleic acids.
Exemplary host cells include gram negative bacteria, such as *Escherichia coli* and
Pseudomonas fluorescens; gram positive bacteria, such as *Lactobacillus gasseri*,
Lactococcus lactis, *Lactococcus cremoris*, *Bacillus subtilis*. Exemplary host cells also
include eukaryotic organisms, e.g., various yeast, such as *Saccharomyces* sp., including
30 *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, *Pichia pastoris*, and
Kluyveromyces lactis, *Hansenula polymorpha*, *Aspergillus niger*, and mammalian cells
and cell lines and insect cells and cell lines. Other exemplary host cells include bacterial
cells, such as *E. coli*, *Streptomyces*, *Bacillus subtilis*, *Bacillus cereus*, *Salmonella*

typhimurium and various species within the genera *Pseudomonas*, *Streptomyces* and *Staphylococcus*, fungal cells, such as *Aspergillus*, yeast such as any species of *Pichia*, *Saccharomyces*, *Schizosaccharomyces*, *Schwanniomyces*, including *Pichia pastoris*, *Saccharomyces cerevisiae*, or *Schizosaccharomyces pombe*, insect cells such as
5 *Drosophila S2* and *Spodoptera Sf9*, animal cells such as CHO, COS or Bowes melanoma and adenoviruses. The selection of an appropriate host is within the abilities of those skilled in the art. Thus, the invention also includes nucleic acids and polypeptides optimized for expression in these organisms and species.

For example, the codons of a nucleic acid encoding an isomerase, e.g.,
10 racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase isolated from a bacterial cell are modified such that the nucleic acid is optimally expressed in a bacterial cell different from the bacteria from which the isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase was derived, a yeast, a fungi, a plant cell, an insect cell or a mammalian cell. Methods for optimizing codons are well known in the art,
15 see, e.g., U.S. Patent No. 5,795,737; Baca (2000) Int. J. Parasitol. 30:113-118; Hale (1998) Protein Expr. Purif. 12:185-188; Narum (2001) Infect. Immun. 69:7250-7253. See also Narum (2001) Infect. Immun. 69:7250-7253, describing optimizing codons in mouse systems; Outchkourov (2002) Protein Expr. Purif. 24:18-24, describing optimizing codons in yeast; Feng (2000) Biochemistry 39:15399-15409, describing optimizing codons in *E.*
20 *coli*; Humphreys (2000) Protein Expr. Purif. 20:252-264, describing optimizing codon usage that affects secretion in *E. coli*.

Transgenic non-human animals

The invention provides transgenic non-human animals comprising a nucleic acid, a polypeptide (e.g., an isomerase, e.g., racemase, e.g., amino acid racemase, alanine
25 racemase, and/or epimerase), an expression cassette or vector or a transfected or transformed cell of the invention. The invention also provides methods of making and using these transgenic non-human animals.

The transgenic non-human animals can be, e.g., goats, rabbits, sheep, pigs, cows, rats, horses, dogs, fish and mice, comprising the nucleic acids of the invention. These
30 animals can be used, e.g., as *in vivo* models to study isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase activity, or, as models to screen for agents that change the isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase activity *in vivo*. The coding sequences for the polypeptides to

be expressed in the transgenic non-human animals can be designed to be constitutive, or, under the control of tissue-specific, developmental-specific or inducible transcriptional regulatory factors. Transgenic non-human animals can be designed and generated using any method known in the art; see, e.g., U.S. Patent Nos. 6,211,428; 6,187,992; 6,156,952; 5 6,118,044; 6,111,166; 6,107,541; 5,959,171; 5,922,854; 5,892,070; 5,880,327; 5,891,698; 5,639,940; 5,573,933; 5,387,742; 5,087,571, describing making and using transformed cells and eggs and transgenic mice, rats, rabbits, sheep, pigs, chickens, goats, fish and cows. See also, e.g., Pollock (1999) J. Immunol. Methods 231:147-157, describing the production of recombinant proteins in the milk of transgenic dairy animals; Baguisi (1999) 10 Nat. Biotechnol. 17:456-461, demonstrating the production of transgenic goats. U.S. Patent No. 6,211,428, describes making and using transgenic non-human mammals which express in their brains a nucleic acid construct comprising a DNA sequence. U.S. Patent No. 5,387,742, describes injecting cloned recombinant or synthetic DNA sequences into fertilized mouse eggs, implanting the injected eggs in pseudo-pregnant females, and 15 growing to term transgenic mice whose cells express proteins related to the pathology of Alzheimer's disease. U.S. Patent No. 6,187,992, describes making and using a transgenic mouse whose genome comprises a disruption of the gene encoding amyloid precursor protein (APP).

“Knockout animals” can also be used to practice the methods of the invention. For 20 example, in one aspect, the transgenic or modified animals of the invention comprise a “knockout animal,” e.g., a “knockout mouse,” engineered not to express an endogenous gene, which is replaced with a gene expressing an isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase of the invention, or, a fusion protein comprising an isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, 25 and/or epimerase of the invention.

Transgenic Plants and Seeds

The invention provides transgenic plants and seeds comprising a nucleic acid, a polypeptide (e.g., an isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase), an expression cassette or vector or a transfected or 30 transformed cell of the invention. The invention also provides plant products or byproducts, e.g., fruits, oils, seeds, leaves, extracts and the like, including any plant part, comprising a nucleic acid and/or a polypeptide (e.g., an isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase) of the invention, e.g., wherein

the nucleic acid or polypeptide of the invention is heterologous to the plant, plant part, seed etc. The transgenic plant (which includes plant parts, fruits, seeds etc.) can be dicotyledonous (a dicot) or monocotyledonous (a monocot). The invention also provides methods of making and using these transgenic plants and seeds. The transgenic plant or
5 plant cell expressing a polypeptide of the present invention may be constructed in accordance with any method known in the art. See, for example, U.S. Patent No. 6,309,872.

Nucleic acids and expression constructs of the invention can be introduced into a plant cell by any means. For example, nucleic acids or expression constructs can be
10 introduced into the genome of a desired plant host, or, the nucleic acids or expression constructs can be episomes. Introduction into the genome of a desired plant can be such that the host's isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase production is regulated by endogenous transcriptional or translational control elements. The invention also provides "knockout plants" where insertion of gene
15 sequence by, e.g., homologous recombination, has disrupted the expression of the endogenous gene. Means to generate "knockout" plants are well-known in the art, see, e.g., Strepp (1998) Proc Natl. Acad. Sci. USA 95:4368-4373; Miao (1995) Plant J 7:359-365. See discussion on transgenic plants, below.

The nucleic acids of the invention can be used to confer desired traits on
20 essentially any plant, e.g., on starch-producing plants, such as potato, wheat, rice, barley, and the like. Nucleic acids of the invention can be used to manipulate metabolic pathways of a plant in order to optimize or alter host's expression of an isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase. The can change
25 isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase activity in a plant. Alternatively, an isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase of the invention can be used in production of a transgenic plant to produce a compound not naturally produced by that plant. This can lower production costs or create a novel product.

In one aspect, the first step in production of a transgenic plant involves making an
30 expression construct for expression in a plant cell. These techniques are well known in the art. They can include selecting and cloning a promoter, a coding sequence for facilitating efficient binding of ribosomes to mRNA and selecting the appropriate gene terminator sequences. One exemplary constitutive promoter is CaMV35S, from the cauliflower mosaic virus, which generally results in a high degree of expression in plants. Other

promoters are more specific and respond to cues in the plant's internal or external environment. An exemplary light-inducible promoter is the promoter from the *cab* gene, encoding the major chlorophyll a/b binding protein.

In one aspect, the nucleic acid is modified to achieve greater expression in a plant cell. For example, a sequence of the invention is likely to have a higher percentage of A-T nucleotide pairs compared to that seen in a plant, some of which prefer G-C nucleotide pairs. Therefore, A-T nucleotides in the coding sequence can be substituted with G-C nucleotides without significantly changing the amino acid sequence to enhance production of the gene product in plant cells.

Selectable marker gene can be added to the gene construct in order to identify plant cells or tissues that have successfully integrated the transgene. This may be necessary because achieving incorporation and expression of genes in plant cells is a rare event, occurring in just a few percent of the targeted tissues or cells. Selectable marker genes encode proteins that provide resistance to agents that are normally toxic to plants, such as antibiotics or herbicides. Only plant cells that have integrated the selectable marker gene will survive when grown on a medium containing the appropriate antibiotic or herbicide. As for other inserted genes, marker genes also require promoter and termination sequences for proper function.

In one aspect, making transgenic plants or seeds comprises incorporating sequences of the invention and, in one aspect (optionally), marker genes into a target expression construct (e.g., a plasmid), along with positioning of the promoter and the terminator sequences. This can involve transferring the modified gene into the plant through a suitable method. For example, a construct may be introduced directly into the genomic DNA of the plant cell using techniques such as electroporation and microinjection of plant cell protoplasts, or the constructs can be introduced directly to plant tissue using ballistic methods, such as DNA particle bombardment. For example, see, e.g., Christou (1997) *Plant Mol. Biol.* 35:197-203; Pawlowski (1996) *Mol. Biotechnol.* 6:17-30; Klein (1987) *Nature* 327:70-73; Takumi (1997) *Genes Genet. Syst.* 72:63-69, discussing use of particle bombardment to introduce transgenes into wheat; and Adam (1997) *supra*, for use of particle bombardment to introduce YACs into plant cells. For example, Rinehart (1997) *supra*, used particle bombardment to generate transgenic cotton plants. Apparatus for accelerating particles is described U.S. Pat. No. 5,015,580; and, the commercially available BioRad (Biolistics) PDS-2000 particle acceleration

instrument; see also, John, U.S. Patent No. 5,608,148; and Ellis, U.S. Patent No. 5, 681,730, describing particle-mediated transformation of gymnosperms.

In one aspect, protoplasts can be immobilized and injected with a nucleic acids, e.g., an expression construct. Although plant regeneration from protoplasts is not easy
5 with cereals, plant regeneration is possible in legumes using somatic embryogenesis from protoplast derived callus. Organized tissues can be transformed with naked DNA using gene gun technique, where DNA is coated on tungsten microprojectiles, shot 1/100th the size of cells, which carry the DNA deep into cells and organelles. Transformed tissue is then induced to regenerate, usually by somatic embryogenesis. This technique has been
10 successful in several cereal species including maize and rice.

Nucleic acids, e.g., expression constructs, can also be introduced in to plant cells using recombinant viruses. Plant cells can be transformed using viral vectors, such as, e.g., tobacco mosaic virus derived vectors (Rouwental (1997) *Plant Mol. Biol.* 33:989-999), see Porta (1996) "Use of viral replicons for the expression of genes in plants," *Mol.*
15 *Biotechnol.* 5:209-221.

Alternatively, nucleic acids, e.g., an expression construct, can be combined with suitable T-DNA flanking regions and introduced into a conventional *Agrobacterium tumefaciens* host vector. The virulence functions of the *Agrobacterium tumefaciens* host will direct the insertion of the construct and adjacent marker into the plant cell DNA when
20 the cell is infected by the bacteria. *Agrobacterium tumefaciens*-mediated transformation techniques, including disarming and use of binary vectors, are well described in the scientific literature. See, e.g., Horsch (1984) *Science* 233:496-498; Fraley (1983) *Proc. Natl. Acad. Sci. USA* 80:4803 (1983); *Gene Transfer to Plants*, Potrykus, ed. (Springer-Verlag, Berlin 1995). The DNA in an *A. tumefaciens* cell is contained in the bacterial
25 chromosome as well as in another structure known as a Ti (tumor-inducing) plasmid. The Ti plasmid contains a stretch of DNA termed T-DNA (~20 kb long) that is transferred to the plant cell in the infection process and a series of vir (virulence) genes that direct the infection process. *A. tumefaciens* can only infect a plant through wounds: when a plant root or stem is wounded it gives off certain chemical signals, in response to which, the vir
30 genes of *A. tumefaciens* become activated and direct a series of events necessary for the transfer of the T-DNA from the Ti plasmid to the plant's chromosome. The T-DNA then enters the plant cell through the wound. One speculation is that the T-DNA waits until the plant DNA is being replicated or transcribed, then inserts itself into the exposed plant DNA. In order to use *A. tumefaciens* as a transgene vector, the tumor-inducing section of

T-DNA have to be removed, while retaining the T-DNA border regions and the vir genes. The transgene is then inserted between the T-DNA border regions, where it is transferred to the plant cell and becomes integrated into the plant's chromosomes.

The invention provides for the transformation of monocotyledonous plants using
5 the nucleic acids of the invention, including important cereals, see Hiei (1997) *Plant Mol. Biol.* 35:205-218. See also, *e.g.*, Horsch, *Science* (1984) 233:496; Fraley (1983) *Proc. Natl. Acad. Sci USA* 80:4803; Thykjaer (1997) *supra*; Park (1996) *Plant Mol. Biol.* 32:1135-1148, discussing T-DNA integration into genomic DNA. See also D'Halluin, U.S. Patent No. 5,712,135, describing a process for the stable integration of a DNA
10 comprising a gene that is functional in a cell of a cereal, or other monocotyledonous plant.

In one aspect, the third step can involve selection and regeneration of whole plants capable of transmitting the incorporated target gene to the next generation. Such regeneration techniques rely on manipulation of certain phytohormones in a tissue culture growth medium, typically relying on a biocide and/or herbicide marker that has been
15 introduced together with the desired nucleotide sequences. Plant regeneration from cultured protoplasts is described in Evans et al., *Protoplasts Isolation and Culture, Handbook of Plant Cell Culture*, pp. 124-176, MacMillan Publishing Company, New York, 1983; and Binding, *Regeneration of Plants, Plant Protoplasts*, pp. 21-73, CRC Press, Boca Raton, 1985. Regeneration can also be obtained from plant callus, explants,
20 organs, or parts thereof. Such regeneration techniques are described generally in Klee (1987) *Ann. Rev. of Plant Phys.* 38:467-486. To obtain whole plants from transgenic tissues such as immature embryos, they can be grown under controlled environmental conditions in a series of media containing nutrients and hormones, a process known as tissue culture. Once whole plants are generated and produce seed, evaluation of the
25 progeny begins.

After the expression cassette is stably incorporated in transgenic plants, it can be introduced into other plants by sexual crossing. Any of a number of standard breeding techniques can be used, depending upon the species to be crossed. Since transgenic expression of the nucleic acids of the invention leads to phenotypic changes, plants
30 comprising the recombinant nucleic acids of the invention can be sexually crossed with a second plant to obtain a final product. Thus, the seed of the invention can be derived from a cross between two transgenic plants of the invention, or a cross between a plant of the invention and another plant. The desired effects (*e.g.*, expression of the polypeptides of the invention to produce a plant in which flowering behavior is altered) can be enhanced

when both parental plants express the polypeptides (e.g., an isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase) of the invention. The desired effects can be passed to future plant generations by standard propagation means.

The nucleic acids and polypeptides of the invention are expressed in or inserted in
5 any plant or seed. Transgenic plants of the invention can be dicotyledonous or monocotyledonous. Examples of monocot transgenic plants of the invention are grasses, such as meadow grass (blue grass, *Poa*), forage grass such as festuca, lolium, temperate grass, such as *Agrostis*, and cereals, e.g., wheat, oats, rye, barley, rice, sorghum, and maize (corn). Examples of dicot transgenic plants of the invention are tobacco, legumes, such as
10 lupins, potato, sugar beet, pea, bean and soybean, and cruciferous plants (family *Brassicaceae*), such as cauliflower, rape seed, and the closely related model organism *Arabidopsis thaliana*. Thus, the transgenic plants and seeds of the invention include a broad range of plants, including, but not limited to, species from the genera *Anacardium*, *Arachis*, *Asparagus*, *Atropa*, *Avena*, *Brassica*, *Citrus*, *Citrullus*, *Capsicum*, *Carthamus*,
15 *Cocos*, *Coffea*, *Cucumis*, *Cucurbita*, *Daucus*, *Elaeis*, *Fragaria*, *Glycine*, *Gossypium*, *Helianthus*, *Heterocallis*, *Hordeum*, *Hyoscyamus*, *Lactuca*, *Linum*, *Lolium*, *Lupinus*, *Lycopersicon*, *Malus*, *Manihot*, *Majorana*, *Medicago*, *Nicotiana*, *Olea*, *Oryza*, *Panicum*, *Pannisetum*, *Persea*, *Phaseolus*, *Pistachia*, *Pisum*, *Pyrus*, *Prunus*, *Raphanus*, *Ricinus*,
20 *Secale*, *Senecio*, *Sinapis*, *Solanum*, *Sorghum*, *Theobromus*, *Trigonella*, *Triticum*, *Vicia*, *Vitis*, *Vigna*, and *Zea*. Transgenic plants and seeds of the invention can be any monocot or dicot, e.g., a monocot corn, sugarcane, rice, wheat, barley, switchgrass or *Miscanthus*; or a dicot oilseed crop, soy, canola, rapeseed, flax, cotton, palm oil, sugar beet, peanut, tree, poplar or lupine.

In alternative embodiments, the nucleic acids of the invention are expressed in
25 plants (and/or their seeds) which contain fiber cells, including, e.g., cotton, silk cotton tree (Kapok, *Ceiba pentandra*), desert willow, creosote bush, winterfat, balsa, ramie, kenaf, hemp, roselle, jute, sisal abaca and flax. In alternative embodiments, the transgenic plants of the invention can be members of the genus *Gossypium*, including members of any *Gossypium* species, such as *G. arboreum*; *G. herbaceum*, *G. barbadense*, and *G.*
30 *hirsutum*.

The invention also provides for transgenic plants (and/or their seeds) to be used for producing large amounts of the polypeptides (e.g., an isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase or antibody) of the invention. For example, see Palmgren (1997) Trends Genet. 13:348; Chong (1997) Transgenic Res.

6:289-296 (producing human milk protein beta-casein in transgenic potato plants using an auxin-inducible, bidirectional mannopine synthase (mas1',2') promoter with *Agrobacterium tumefaciens*-mediated leaf disc transformation methods).

Using known procedures, one of skill can screen for plants (and/or their seeds) of the invention by detecting the increase or decrease of transgene mRNA or protein in transgenic plants. Means for detecting and quantitation of mRNAs or proteins are well known in the art.

Polypeptides and peptides

In one aspect, the invention provides isolated, synthetic or recombinant polypeptides and peptides having isomerase activity, e.g., racemase activity, e.g., amino acid racemase activity (e.g., resolving a D- and/or an L- amino acid from a racemic mixture), alanine racemase activity, and/or epimerase activity, or polypeptides and peptides capable of generating an antibody that specifically binds to an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase, including an enzyme of this invention, including the amino acid sequences of the invention, which include those having at least 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more, or 100% (complete) sequence identity to an exemplary polypeptide of the invention (as defined above, including SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:22, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:28, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:42, SEQ ID NO:44, SEQ ID NO:46, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:58, SEQ ID NO:60, SEQ ID NO:62, SEQ ID NO:64, SEQ ID NO:66, SEQ ID NO:68, SEQ ID NO:70, SEQ ID NO:72, SEQ ID NO:74, SEQ ID NO:76, SEQ ID NO:78, SEQ ID NO:80, SEQ ID NO:82, SEQ ID NO:84, SEQ ID NO:86, SEQ ID NO:88, SEQ ID NO:90, SEQ ID NO:92, SEQ ID NO:94, SEQ ID NO:96, SEQ ID NO:98, SEQ ID NO:100, SEQ ID NO:102, SEQ ID NO:104, SEQ ID NO:106, SEQ ID NO:108, SEQ ID NO:110, SEQ ID NO:112, SEQ ID NO:114, SEQ ID NO:116, SEQ ID NO:118, SEQ ID NO:120, SEQ ID NO:122, SEQ ID NO:124, SEQ ID NO:126, SEQ ID NO:128, SEQ ID NO:130, SEQ ID NO:132, SEQ ID

NO:134, SEQ ID NO:136, SEQ ID NO:138, SEQ ID NO:140, SEQ ID NO:142, SEQ ID
NO:143, SEQ ID NO:146, SEQ ID NO:148, SEQ ID NO:150, SEQ ID NO:152, SEQ ID
NO:154, SEQ ID NO:156, SEQ ID NO:158, SEQ ID NO:160, SEQ ID NO:162, SEQ ID
NO:164, SEQ ID NO:166, SEQ ID NO:168, SEQ ID NO:170, SEQ ID NO:172, SEQ ID
5 NO:174, SEQ ID NO:176, SEQ ID NO:178, SEQ ID NO:180, SEQ ID NO:182, SEQ ID
NO:184, SEQ ID NO:186, SEQ ID NO:188, SEQ ID NO:190, SEQ ID NO:192, SEQ ID
NO:194, SEQ ID NO:196, SEQ ID NO:198, SEQ ID NO:200, SEQ ID NO:202, SEQ ID
NO:204, SEQ ID NO:206, SEQ ID NO:208, SEQ ID NO:210, SEQ ID NO:212, SEQ ID
NO:214, SEQ ID NO:216, SEQ ID NO:218, SEQ ID NO:220, SEQ ID NO:222, SEQ ID
10 NO:224, SEQ ID NO:226, SEQ ID NO:228, SEQ ID NO:230, SEQ ID NO:232, SEQ ID
NO:234, SEQ ID NO:236, SEQ ID NO:238, SEQ ID NO:240, SEQ ID NO:242, SEQ ID
NO:244, SEQ ID NO:246, SEQ ID NO:248, SEQ ID NO:250, SEQ ID NO:252, SEQ ID
NO:254, SEQ ID NO:256, SEQ ID NO:258, SEQ ID NO:260, SEQ ID NO:262, SEQ ID
NO:264, SEQ ID NO:266, SEQ ID NO:268, SEQ ID NO:270, SEQ ID NO:272, SEQ ID
15 NO:274, SEQ ID NO:276, SEQ ID NO:278, SEQ ID NO:280, SEQ ID NO:282, SEQ ID
NO:284, SEQ ID NO:286, SEQ ID NO:288, SEQ ID NO:290, SEQ ID NO:292, SEQ ID
NO:294, SEQ ID NO:296, SEQ ID NO:298, SEQ ID NO:300, SEQ ID NO:302, SEQ ID
NO:304, SEQ ID NO:306, SEQ ID NO:308, SEQ ID NO:310, SEQ ID NO:312, SEQ ID
NO:314, SEQ ID NO:316, SEQ ID NO:318, SEQ ID NO:320, SEQ ID NO:322, SEQ ID
20 NO:324, SEQ ID NO:326, SEQ ID NO:328, SEQ ID NO:330, SEQ ID NO:332, SEQ ID
NO:334, SEQ ID NO:336, SEQ ID NO:338, SEQ ID NO:340, SEQ ID NO:342, SEQ ID
NO:344, SEQ ID NO:346, SEQ ID NO:348, SEQ ID NO:350, SEQ ID NO:352, SEQ ID
NO:354, SEQ ID NO:356, SEQ ID NO:358, SEQ ID NO:360, SEQ ID NO:362, SEQ ID
NO:364, SEQ ID NO:366, SEQ ID NO:368, SEQ ID NO:370, SEQ ID NO:372, SEQ ID
25 NO:374, SEQ ID NO:376, SEQ ID NO:378, SEQ ID NO:380, SEQ ID NO:382, SEQ ID
NO:384, SEQ ID NO:386, SEQ ID NO:388, SEQ ID NO:390, SEQ ID NO:392, SEQ ID
NO:394, SEQ ID NO:396, SEQ ID NO:398, SEQ ID NO:400, SEQ ID NO:402, SEQ ID
NO:404, SEQ ID NO:406, SEQ ID NO:408, SEQ ID NO:410, SEQ ID NO:412, SEQ ID
NO:414, SEQ ID NO:416, SEQ ID NO:418, SEQ ID NO:420, SEQ ID NO:422, SEQ ID
30 NO:424, SEQ ID NO:426, SEQ ID NO:428, SEQ ID NO:430, SEQ ID NO:432, SEQ ID
NO:434, SEQ ID NO:436, SEQ ID NO:438, SEQ ID NO:440, SEQ ID NO:442, SEQ ID
NO:444, SEQ ID NO:446, SEQ ID NO:448, SEQ ID NO:450, SEQ ID NO:452, SEQ ID
NO:454, SEQ ID NO:456, SEQ ID NO:458, SEQ ID NO:460, SEQ ID NO:462, SEQ ID
NO:464, SEQ ID NO:466, SEQ ID NO:468, SEQ ID NO:470, SEQ ID NO:472, SEQ ID

NO:474, SEQ ID NO:476, SEQ ID NO:478, SEQ ID NO:480, SEQ ID NO:482, SEQ ID NO:484, SEQ ID NO:486, SEQ ID NO:488, SEQ ID NO:490, SEQ ID NO:492, SEQ ID NO:494, SEQ ID NO:496 or SEQ ID NO:498), including the sequences described herein and in Tables 1, 2 and 3, and the Sequence Listing (all of these sequences are “exemplary enzymes/ polypeptides of the invention”), and enzymatically active subsequences (fragments) thereof.

In one aspect, the invention provides chimeric enzymes, including an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase, having heterologous domains, e.g., a binding domain or a dockerin domain, e.g., for use in the processes of the invention and in various industrial, medical, pharmaceutical, research, food and feed and food and feed supplement processing and other applications. For example, in one aspect the invention provides enzymes, e.g., isomerases, e.g., racemases, e.g., amino acid racemases, alanine racemases, and/or epimerases comprising one or more domain of an enzyme of the invention. In another aspect, domains between different enzymes of the invention can be swapped; or, alternatively, one or more domains of one or more enzymes of the invention can be spliced into an enzyme, e.g., an isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase. In one aspect of the invention, the domains are selected from a binding domain or a dockerin domain.

The invention further provides chimeric enzymes having heterologous, non-natural substrates; including chimeric enzymes having multiple substrates by nature of their “spliced-in” heterologous domains – thus giving the chimeric enzyme new specificity or enhanced binding. The heterologous domains of the chimeric enzymes of the invention can be designed to be modular, i.e., to be appended to a catalytic module or catalytic domain (e.g., an active site), which also can be heterologous or can be homologous to the enzyme.

Utilization of just the catalytic module of an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase (e.g., an enzyme of the invention) has been shown to be effective. Thus, the invention provides peptides and polypeptides consisting of, or comprising, modular domains/active site modules, which can be homologously paired or joined as chimeric (heterologous) active site module pairs. Thus, these chimeric polypeptides/ peptides of the invention can be used to improve or alter the performance of an individual enzyme, e.g., an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase enzyme. A chimeric catalytic module of the invention (comprising, e.g., at least one domain of the invention)

can be designed to target the enzyme to particular regions of a substrate. For example, in one aspect, this is achieved by making fusions of the isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase and various domains (either an isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase of the invention with a heterologous domain, or, a domain of the invention with another enzyme, e.g., an isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase.

Thus, the invention provides chimeric isomerases, e.g., racemases, e.g., amino acid racemases, alanine racemases, and/or epimerases, e.g., a fusion of an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase with different (e.g., heterologous) domains. In one aspect, the chimeric isomerases, e.g., racemases, e.g., amino acid racemases, alanine racemases, and/or epimerases comprise an enzyme of the invention. In one aspect, the chimeric enzyme comprises fusions of different domains. The invention also provides methods comprising recombining different domains with different isomerases, e.g., racemases, e.g., amino acid racemases, alanine racemases, and/or epimerases (e.g., domains of the invention and/or isomerases, e.g., racemases, e.g., amino acid racemases, alanine racemases, and/or epimerases of the invention) and screening the resultant chimerics to find the best combination for a particular application or substrate.

Other variations also are within the scope of this invention, e.g., where one, two, three, four or five or more residues are removed from the carboxy- or amino- terminal ends of any polypeptide of the invention. Another variation includes modifying any residue to increase or decrease pI of a polypeptide, e.g., removing or modifying (e.g., to another amino acid) a glutamate. This method was used as a general scheme for improving the enzyme's properties without creating regulatory issues since no amino acids are mutated; and this general scheme can be used with any polypeptide of the invention.

The invention provides isolated, synthetic or recombinant polypeptides having isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase activity, wherein the polypeptide has a sequence modification of any polypeptide of the invention, including any exemplary amino acid sequence of the invention, including SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:22, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:28, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:42, SEQ ID NO:44,

SEQ ID NO:46, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:52, SEQ ID NO:54, SEQ
ID NO:56, SEQ ID NO:58, SEQ ID NO:60, SEQ ID NO:62, SEQ ID NO:64, SEQ ID
NO:66, SEQ ID NO:68, SEQ ID NO:70, SEQ ID NO:72, SEQ ID NO:74, SEQ ID NO:76,
SEQ ID NO:78, SEQ ID NO:80, SEQ ID NO:82, SEQ ID NO:84, SEQ ID NO:86, SEQ
5 ID NO:88, SEQ ID NO:90, SEQ ID NO:92, SEQ ID NO:94, SEQ ID NO:96, SEQ ID
NO:98, SEQ ID NO:100, SEQ ID NO:102, SEQ ID NO:104, SEQ ID NO:106, SEQ ID
NO:108, SEQ ID NO:110, SEQ ID NO:112, SEQ ID NO:114, SEQ ID NO:116, SEQ ID
NO:118, SEQ ID NO:120, SEQ ID NO:122, SEQ ID NO:124, SEQ ID NO:126, SEQ ID
NO:128, SEQ ID NO:130, SEQ ID NO:132, SEQ ID NO:134, SEQ ID NO:136, SEQ ID
10 NO:138, SEQ ID NO:140, SEQ ID NO:142, SEQ ID NO:143, SEQ ID NO:146, SEQ ID
NO:148, SEQ ID NO:150, SEQ ID NO:152, SEQ ID NO:154, SEQ ID NO:156, SEQ ID
NO:158, SEQ ID NO:160, SEQ ID NO:162, SEQ ID NO:164, SEQ ID NO:166, SEQ ID
NO:168, SEQ ID NO:170, SEQ ID NO:172, SEQ ID NO:174, SEQ ID NO:176, SEQ ID
NO:178, SEQ ID NO:180, SEQ ID NO:182, SEQ ID NO:184, SEQ ID NO:186, SEQ ID
15 NO:188, SEQ ID NO:190, SEQ ID NO:192, SEQ ID NO:194, SEQ ID NO:196, SEQ ID
NO:198, SEQ ID NO:200, SEQ ID NO:202, SEQ ID NO:204, SEQ ID NO:206, SEQ ID
NO:208, SEQ ID NO:210, SEQ ID NO:212, SEQ ID NO:214, SEQ ID NO:216, SEQ ID
NO:218, SEQ ID NO:220, SEQ ID NO:222, SEQ ID NO:224, SEQ ID NO:226, SEQ ID
NO:228, SEQ ID NO:230, SEQ ID NO:232, SEQ ID NO:234, SEQ ID NO:236, SEQ ID
20 NO:238, SEQ ID NO:240, SEQ ID NO:242, SEQ ID NO:244, SEQ ID NO:246, SEQ ID
NO:248, SEQ ID NO:250, SEQ ID NO:252, SEQ ID NO:254, SEQ ID NO:256, SEQ ID
NO:258, SEQ ID NO:260, SEQ ID NO:262, SEQ ID NO:264, SEQ ID NO:266, SEQ ID
NO:268, SEQ ID NO:270, SEQ ID NO:272, SEQ ID NO:274, SEQ ID NO:276, SEQ ID
NO:278, SEQ ID NO:280, SEQ ID NO:282, SEQ ID NO:284, SEQ ID NO:286, SEQ ID
25 NO:288, SEQ ID NO:290, SEQ ID NO:292, SEQ ID NO:294, SEQ ID NO:296, SEQ ID
NO:298, SEQ ID NO:300, SEQ ID NO:302, SEQ ID NO:304, SEQ ID NO:306, SEQ ID
NO:308, SEQ ID NO:310, SEQ ID NO:312, SEQ ID NO:314, SEQ ID NO:316, SEQ ID
NO:318, SEQ ID NO:320, SEQ ID NO:322, SEQ ID NO:324, SEQ ID NO:326, SEQ ID
NO:328, SEQ ID NO:330, SEQ ID NO:332, SEQ ID NO:334, SEQ ID NO:336, SEQ ID
30 NO:338, SEQ ID NO:340, SEQ ID NO:342, SEQ ID NO:344, SEQ ID NO:346, SEQ ID
NO:348, SEQ ID NO:350, SEQ ID NO:352, SEQ ID NO:354, SEQ ID NO:356, SEQ ID
NO:358, SEQ ID NO:360, SEQ ID NO:362, SEQ ID NO:364, SEQ ID NO:366, SEQ ID
NO:368, SEQ ID NO:370, SEQ ID NO:372, SEQ ID NO:374, SEQ ID NO:376, SEQ ID
NO:378, SEQ ID NO:380, SEQ ID NO:382, SEQ ID NO:384, SEQ ID NO:386, SEQ ID

NO:388, SEQ ID NO:390, SEQ ID NO:392, SEQ ID NO:394, SEQ ID NO:396, SEQ ID NO:398, SEQ ID NO:400, SEQ ID NO:402, SEQ ID NO:404, SEQ ID NO:406, SEQ ID NO:408, SEQ ID NO:410, SEQ ID NO:412, SEQ ID NO:414, SEQ ID NO:416, SEQ ID NO:418, SEQ ID NO:420, SEQ ID NO:422, SEQ ID NO:424, SEQ ID NO:426, SEQ ID NO:428, SEQ ID NO:430, SEQ ID NO:432, SEQ ID NO:434, SEQ ID NO:436, SEQ ID NO:438, SEQ ID NO:440, SEQ ID NO:442, SEQ ID NO:444, SEQ ID NO:446, SEQ ID NO:448, SEQ ID NO:450, SEQ ID NO:452, SEQ ID NO:454, SEQ ID NO:456, SEQ ID NO:458, SEQ ID NO:460, SEQ ID NO:462, SEQ ID NO:464, SEQ ID NO:466, SEQ ID NO:468, SEQ ID NO:470, SEQ ID NO:472, SEQ ID NO:474, SEQ ID NO:476, SEQ ID NO:478, SEQ ID NO:480, SEQ ID NO:482, SEQ ID NO:484, SEQ ID NO:486, SEQ ID NO:488, SEQ ID NO:490, SEQ ID NO:492, SEQ ID NO:494, SEQ ID NO:496 or SEQ ID NO:498, including the sequences described herein and in Tables 1, 2 and 3, and the Sequence Listing (all of these sequences are “exemplary enzymes/ polypeptides of the invention”), and enzymatically active subsequences (fragments) thereof. The sequence change(s) can also comprise any amino acid modification to change the pI of a polypeptide, e.g., deletion or modification of a glutamate, or changing from a glutamate to another residue.

The invention further provides isolated, synthetic or recombinant polypeptides having a sequence identity (e.g., at least about 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more, or complete (100%) sequence identity) to an exemplary sequence of the invention.

In one aspect, the polypeptide of the invention has an isomerase activity, e.g., racemase activity, e.g., amino acid racemase activity, alanine racemase activity, and/or epimerase activity, and/or catalyze the re-arrangement of atoms within a molecule, catalyze the conversion of one isomer into another, catalyze the conversion of an optically active substrate into a raceme, which is optically inactive, catalyze the interconversion of substrate enantiomers, catalyze the stereochemical inversion around the asymmetric carbon atom in a substrate having only one center of asymmetry, catalyze the stereochemical inversion of the configuration around an asymmetric carbon atom in a substrate having more than one asymmetric center, and/or catalyze the racemization of amino acids.

In one aspect, a polypeptide of the invention has a 5-substituted hydantoin racemase activity, e.g., it can catalyze racemization reaction of an optically active 5-substituted hydantoin compound, such as a D- or L-5-substituted hydantoin compound.

5 In one aspect, a polypeptide of the invention has peptidyl-prolyl isomerase activity, and can be part of a signaling pathway that leads to T-cell activation, and/or correct protein folding and/or protein trafficking, and also can be involved in assembly/disassembly of protein complexes and regulation of protein activity.

10 In one aspect, a polypeptide of the invention has a racemase, or isomerase, activity that catalyzes inversion of a molecule's configuration around an asymmetric carbon atom in a substrate having a single center of asymmetry, thereby interconverting two racemers.

In one aspect, a polypeptide of the invention has a racemase, or an epimerase activity, that catalyzes inversion of configuration around an asymmetric carbon atom in a substrate with more than one center of symmetry, thereby interconverting two epimers. Racemases and epimerases of this invention can act on amino acids and their derivatives, hydroxy acids and their derivatives, and carbohydrates and their derivatives. For example, the interconversion of UDP-galactose and UDP-glucose can be catalyzed by an enzyme of this invention having a UDP-galactose-4'-epimerase activity; proper regulation and function of this epimerase is essential to the synthesis of glycoproteins and glycolipids. Elevated blood galactose levels have been correlated with UDP-galactose-4'-epimerase deficiency in screening programs of infants.

15
20

In one aspect, a polypeptide of the invention has a serine racemase enzyme activity, and this enzyme can be used to increase or decrease D-serine formation, which can be used as a pharmaceutical (drug) in an individual, e.g., to increase or decrease NMDA receptor activation; a decrease in D-serine formation (by using a serine racemase enzyme of this invention) can aid in the prevention of neuron damage following an ischemic event, such as stroke; and regulation of D-serine formation by a serine racemase enzyme of this invention also can be effective for treating a neurodegenerative condition caused by the over- or under-activation of the glutamate NMDA receptor.

25

Polypeptides of the invention having a serine racemase enzyme activity can be used to regulate D-serine levels, e.g., to prevent or minimize neuron damage caused, for example, by primary and/or secondary disorders after brain injury, motor unit-like neurogenic and myopathic disorders, neurodegenerative disorders like Alzheimer's and Parkinson's disease, disorders leading to peripheral and chronic pain. See, e.g., U.S. Pat. App. Pub. No. 20030175941.

30

Any isomerase activity, e.g., racemase activity, e.g., amino acid racemase activity, alanine racemase activity, and/or epimerase activity assay known in the art can be used to determine if a polypeptide has isomerase activity, e.g., racemase activity, e.g., amino acid racemase activity, alanine racemase activity, and/or epimerase activity and is within scope
5 of the invention. For example, Schonfeld and Bornscheuer (Anal Chem. 2004 Feb 15;76(4):1184-8) describe a polarimetric assay which identifies alpha-amino acid racemase activity using a glutamate racemase from *Lactobacillus fermentii*, expressed in *E. coli*, and measuring the time-dependent change of the optical rotation using the l-glutamate as the substrate.

10 Another exemplary method for detecting racemase activity to determine if a polypeptide is within the scope of this invention is described in U.S. Pat. App. Pub. No. 20070128601, and comprises providing a reaction medium containing a D-amino acid specific to the racemase to be detected; reacting the D-amino acid with a D-amino oxidase with a prosthetic group to form a reduced prosthetic group by oxidation of the D-amino
15 acid; reacting the reduced prosthetic group with oxygen to form hydrogen peroxide; and detecting the hydrogen peroxide thus formed; wherein the detection of hydrogen peroxide indicates racemase activity in the reaction medium. An exemplary method for detecting a D-amino acid in a sample comprises oxidatively deaminating a D-amino acid by reaction with a D-amino acid oxidase in a prosthetic group; and, detecting the hydrogen peroxide
20 generated by the oxidative deamination; wherein the presence of hydrogen peroxide is indicative of the presence of a D-amino acid in the sample.

Another exemplary method for detecting racemase activity to determine if a polypeptide is within the scope of this invention is described in U.S. Pat. App. Pub. No. 20060014162, and comprises detecting a D-amino acid by providing a reaction medium
25 containing a D-amino acid; reacting the D-amino acid with a D-amino oxidase with a prosthetic group to form a reduced prosthetic group by oxidative deamination of the D-amino acid with a primary amine or oxidation of the D-amino acid with a secondary amine; reacting the reduced prosthetic group with oxygen to form hydrogen peroxide; and detecting the hydrogen peroxide thus formed.

30 The polypeptides of the invention include isomerases, e.g., racemases, e.g., amino acid racemases, alanine racemases, and/or epimerases in an active or inactive form. For example, the polypeptides of the invention include proproteins before “maturation” or processing of prepro sequences, e.g., by a proprotein-processing enzyme, such as a proprotein convertase to generate an “active” mature protein. The polypeptides of the

invention include isomerases, e.g., racemases, e.g., amino acid racemases, alanine racemases, and/or epimerases inactive for other reasons, e.g., before “activation” by a post-translational processing event, e.g., an endo- or exo-peptidase or proteinase action, a phosphorylation event, an amidation, a glycosylation or a sulfation, a dimerization event, and the like. The polypeptides of the invention include all active forms, including active subsequences, e.g., catalytic domains or active sites, of the isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase.

Methods for identifying "prepro" domain sequences and signal sequences are well known in the art, see, e.g., Van de Ven (1993) Crit. Rev. Oncog. 4(2):115-136. For example, to identify a prepro sequence, the protein is purified from the extracellular space and the N-terminal protein sequence is determined and compared to the unprocessed form.

The invention includes polypeptides with or without a signal sequence and/or a prepro sequence. The invention includes polypeptides with heterologous signal sequences and/or prepro sequences. The prepro sequence (including a sequence of the invention used as a heterologous prepro domain) can be located on the amino terminal or the carboxy terminal end of the protein. The invention also includes isolated, synthetic or recombinant signal sequences, prepro sequences and catalytic domains (e.g., “active sites”) comprising sequences of the invention.

The percent sequence identity can be over the full length of the polypeptide, or, the identity can be over a region of at least about 50, 60, 70, 80, 90, 100, 150, 200, 250, 300, 350, 400, 450, 500, 550, 600, 650, 700 or more residues. Polypeptides of the invention can also be shorter than the full length of exemplary polypeptides. In alternative aspects, the invention provides polypeptides (peptides, fragments) ranging in size between about 5 and the full length of a polypeptide, e.g., an enzyme, such as an isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase; exemplary sizes being of about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 100, 125, 150, 175, 200, 250, 300, 350, 400, 450, 500, 550, 600, 650, 700, or more residues, e.g., contiguous residues of an exemplary isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase of the invention.

Peptides of the invention (e.g., a subsequence of an exemplary polypeptide of the invention) can be useful as, e.g., labeling probes, antigens, toleragens, motifs, active sites (e.g., “catalytic domains”), signal sequences and/or prepro domains.

Polypeptides and peptides of the invention can be isolated from natural sources, be synthetic, or be recombinantly generated polypeptides. Peptides and proteins can be recombinantly expressed *in vitro* or *in vivo*. The peptides and polypeptides of the invention can be made and isolated using any method known in the art. Polypeptides and peptides of the invention can also be synthesized, whole or in part, using chemical methods well known in the art. See e.g., Caruthers (1980) Nucleic Acids Res. Symp. Ser. 215-223; Horn (1980) Nucleic Acids Res. Symp. Ser. 225-232; Banga, A.K., Therapeutic Peptides and Proteins, Formulation, Processing and Delivery Systems (1995) Technomic Publishing Co., Lancaster, PA. For example, peptide synthesis can be performed using various solid-phase techniques (see e.g., Roberge (1995) Science 269:202; Merrifield (1997) Methods Enzymol. 289:3-13) and automated synthesis may be achieved, e.g., using the ABI 431A Peptide Synthesizer (Perkin Elmer) in accordance with the instructions provided by the manufacturer.

The peptides and polypeptides of the invention can also be glycosylated. The glycosylation can be added post-translationally either chemically or by cellular biosynthetic mechanisms, wherein the later incorporates the use of known glycosylation motifs, which can be native to the sequence or can be added as a peptide or added in the nucleic acid coding sequence. The glycosylation can be O-linked or N-linked.

“Amino acid” or “amino acid sequence” as used herein refer to an oligopeptide, peptide, polypeptide, or protein sequence, or to a fragment, portion, or subunit of any of these and to naturally occurring or synthetic molecules. “Amino acid” or “amino acid sequence” include an oligopeptide, peptide, polypeptide, or protein sequence, or to a fragment, portion, or subunit of any of these, and to naturally occurring or synthetic molecules. The term “polypeptide” as used herein, refers to amino acids joined to each other by peptide bonds or modified peptide bonds, *i.e.*, peptide isosteres and may contain modified amino acids other than the 20 gene-encoded amino acids. The polypeptides may be modified by either natural processes, such as post-translational processing, or by chemical modification techniques that are well known in the art. Modifications can occur anywhere in the polypeptide, including the peptide backbone, the amino acid side-chains and the amino or carboxyl termini. It will be appreciated that the same type of modification may be present in the same or varying degrees at several sites in a given polypeptide. Also a given polypeptide may have many types of modifications. Modifications include acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a

nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of a phosphatidylinositol, cross-linking cyclization, disulfide bond formation, demethylation, formation of covalent cross-links, formation of cysteine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, pegylation, phosphorylation, prenylation, racemization, selenoylation, sulfation and transfer-RNA mediated addition of amino acids to protein such as arginylation. (See Creighton, T.E., Proteins – Structure and Molecular Properties 2nd Ed., W.H. Freeman and Company, New York (1993); Posttranslational Covalent Modification of Proteins, B.C. Johnson, Ed., Academic Press, New York, pp. 1-12 (1983)). The peptides and polypeptides of the invention also include all “mimetic” and “peptidomimetic” forms, as described in further detail, below.

“Recombinant” polypeptides or proteins refer to polypeptides or proteins produced by recombinant DNA techniques; *i.e.*, produced from cells transformed by an exogenous DNA construct encoding the desired polypeptide or protein. “Synthetic” nucleic acids (including oligonucleotides), polypeptides or proteins of the invention include those prepared by any chemical synthesis, *e.g.*, as described, below. Solid-phase chemical peptide synthesis methods can also be used to synthesize the polypeptide or fragments of the invention. Such method have been known in the art since the early 1960's (Merrifield, R. B., *J. Am. Chem. Soc.*, 85:2149-2154, 1963) (See also Stewart, J. M. and Young, J. D., Solid Phase Peptide Synthesis, 2nd Ed., Pierce Chemical Co., Rockford, Ill., pp. 11-12)) and have recently been employed in commercially available laboratory peptide design and synthesis kits (Cambridge Research Biochemicals). Such commercially available laboratory kits have generally utilized the teachings of H. M. Geysen *et al*, *Proc. Natl. Acad. Sci., USA*, 81:3998 (1984) and provide for synthesizing peptides upon the tips of a multitude of “rods” or “pins” all of which are connected to a single plate. When such a system is utilized, a plate of rods or pins is inverted and inserted into a second plate of corresponding wells or reservoirs, which contain solutions for attaching or anchoring an appropriate amino acid to the pin's or rod's tips. By repeating such a process step, *i.e.*, inverting and inserting the rod's and pin's tips into appropriate solutions, amino acids are built into desired peptides. In addition, a number of available Fmoc peptide synthesis systems are available. For example, assembly of a polypeptide or fragment can be carried out on a solid support using an Applied Biosystems, Inc. Model 431A automated peptide synthesizer. Such equipment provides ready access to

the peptides of the invention, either by direct synthesis or by synthesis of a series of fragments that can be coupled using other known techniques.

“Fragments” as used herein are a portion of a naturally occurring protein which can exist in at least two different conformations. Fragments can have the same or substantially the same amino acid sequence as the naturally occurring protein. “Substantially the same” means that an amino acid sequence is largely, but not entirely, the same, but retains at least one functional activity of the sequence to which it is related. In general two amino acid sequences are “substantially the same” or “substantially homologous” if they are at least about 85% identical. Fragments which have different three dimensional structures as the naturally occurring protein are also included. An example of this, is a “pro-form” molecule, such as a low activity proprotein that can be modified by cleavage to produce a mature enzyme with significantly higher activity.

The peptides and polypeptides of the invention, as defined above, include all “mimetic” and “peptidomimetic” forms. The terms “mimetic” and “peptidomimetic” refer to a synthetic chemical compound which has substantially the same structural and/or functional characteristics of the polypeptides of the invention. The mimetic can be either entirely composed of synthetic, non-natural analogues of amino acids, or, is a chimeric molecule of partly natural peptide amino acids and partly non-natural analogs of amino acids. The mimetic can also incorporate any amount of natural amino acid conservative substitutions as long as such substitutions also do not substantially alter the mimetic’s structure and/or activity. As with polypeptides of the invention which are conservative variants, routine experimentation will determine whether a mimetic is within the scope of the invention, i.e., that its structure and/or function is not substantially altered. Thus, in one aspect, a mimetic composition is within the scope of the invention if it has an isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase activity.

Polypeptide mimetic compositions of the invention can contain any combination of non-natural structural components. In alternative aspect, mimetic compositions of the invention include one or all of the following three structural groups: a) residue linkage groups other than the natural amide bond (“peptide bond”) linkages; b) non-natural residues in place of naturally occurring amino acid residues; or c) residues which induce secondary structural mimicry, i.e., to induce or stabilize a secondary structure, e.g., a beta turn, gamma turn, beta sheet, alpha helix conformation, and the like. For example, a polypeptide of the invention can be characterized as a mimetic when all or some of its

residues are joined by chemical means other than natural peptide bonds. Individual peptidomimetic residues can be joined by peptide bonds, other chemical bonds or coupling means, such as, e.g., glutaraldehyde, N-hydroxysuccinimide esters, bifunctional maleimides, N,N'-dicyclohexylcarbodiimide (DCC) or N,N'-diisopropylcarbodiimide (DIC). Linking groups that can be an alternative to the traditional amide bond ("peptide bond") linkages include, e.g., ketomethylene (e.g., -C(=O)-CH₂- for -C(=O)-NH-), aminomethylene (CH₂-NH), ethylene, olefin (CH=CH), ether (CH₂-O), thioether (CH₂-S), tetrazole (CN₄-), thiazole, retroamide, thioamide, or ester (see, e.g., Spatola (1983) in Chemistry and Biochemistry of Amino Acids, Peptides and Proteins, Vol. 7, pp 267-357, "Peptide Backbone Modifications," Marcell Dekker, NY).

A polypeptide of the invention can also be characterized as a mimetic by containing all or some non-natural residues in place of naturally occurring amino acid residues. Non-natural residues are well described in the scientific and patent literature; a few exemplary non-natural compositions useful as mimetics of natural amino acid residues and guidelines are described below. Mimetics of aromatic amino acids can be generated by replacing by, e.g., D- or L- naphylalanine; D- or L- phenylglycine; D- or L-2 thiencylalanine; D- or L-1, -2, 3-, or 4- pyrenylalanine; D- or L-3 thiencylalanine; D- or L-(2-pyridinyl)-alanine; D- or L-(3-pyridinyl)-alanine; D- or L-(2-pyrazinyl)-alanine; D- or L-(4-isopropyl)-phenylglycine; D-(trifluoromethyl)-phenylglycine; D-(trifluoromethyl)-phenylalanine; D-p-fluoro-phenylalanine; D- or L-p-biphenylphenylalanine; D- or L-p-methoxy-biphenylphenylalanine; D- or L-2-indole(alkyl)alanines; and, D- or L-alkylainines, where alkyl can be substituted or unsubstituted methyl, ethyl, propyl, hexyl, butyl, pentyl, isopropyl, iso-butyl, sec-isotyl, iso-pentyl, or a non-acidic amino acids. Aromatic rings of a non-natural amino acid include, e.g., thiazolyl, thiophenyl, pyrazolyl, benzimidazolyl, naphthyl, furanyl, pyrrolyl, and pyridyl aromatic rings.

Mimetics of acidic amino acids can be generated by substitution by, e.g., non-carboxylate amino acids while maintaining a negative charge; (phosphono)alanine; sulfated threonine. Carboxyl side groups (e.g., aspartyl or glutamyl) can also be selectively modified by reaction with carbodiimides (R'-N-C-N-R') such as, e.g., 1-cyclohexyl-3(2-morpholinyl-(4-ethyl) carbodiimide or 1-ethyl-3(4-azonia- 4,4-dimetholpentyl) carbodiimide. Aspartyl or glutamyl can also be converted to asparaginyll and glutaminyll residues by reaction with ammonium ions. Mimetics of basic amino acids can be generated by substitution with, e.g., (in addition to lysine and arginine) the amino

acids ornithine, citrulline, or (guanidino)-acetic acid, or (guanidino)alkyl-acetic acid, where alkyl is defined above. Nitrile derivative (e.g., containing the CN-moiety in place of COOH) can be substituted for asparagine or glutamine. Asparaginyl and glutaminyl residues can be deaminated to the corresponding aspartyl or glutamyl residues. Arginine residue mimetics can be generated by reacting arginyl with, e.g., one or more conventional reagents, including, e.g., phenylglyoxal, 2,3-butanedione, 1,2-cyclo-hexanedione, or ninhydrin, preferably under alkaline conditions. Tyrosine residue mimetics can be generated by reacting tyrosyl with, e.g., aromatic diazonium compounds or tetranitromethane. N-acetylimidizol and tetranitromethane can be used to form O-acetyl tyrosyl species and 3-nitro derivatives, respectively. Cysteine residue mimetics can be generated by reacting cysteinyl residues with, e.g., alpha-haloacetates such as 2-chloroacetic acid or chloroacetamide and corresponding amines; to give carboxymethyl or carboxyamidomethyl derivatives. Cysteine residue mimetics can also be generated by reacting cysteinyl residues with, e.g., bromo-trifluoroacetone, alpha-bromo-beta-(5-imidozoyl) propionic acid; chloroacetyl phosphate, N-alkylmaleimides, 3-nitro-2-pyridyl disulfide; methyl 2-pyridyl disulfide; p-chloromercuribenzoate; 2-chloromercuri-4-nitrophenol; or, chloro-7-nitrobenzo-oxa-1,3-diazole. Lysine mimetics can be generated (and amino terminal residues can be altered) by reacting lysinyl with, e.g., succinic or other carboxylic acid anhydrides. Lysine and other alpha-amino-containing residue mimetics can also be generated by reaction with imidoesters, such as methyl picolinimate, pyridoxal phosphate, pyridoxal, chloroborohydride, trinitrobenzenesulfonic acid, O-methylisourea, 2,4, pentanedione, and transamidase-catalyzed reactions with glyoxylate. Mimetics of methionine can be generated by reaction with, e.g., methionine sulfoxide. Mimetics of proline include, e.g., pipercolic acid, thiazolidine carboxylic acid, 3- or 4- hydroxy proline, dehydroproline, 3- or 4-methylproline, or 3,3-dimethylproline. Histidine residue mimetics can be generated by reacting histidyl with, e.g., diethylprocarbonate or para-bromophenacyl bromide. Other mimetics include, e.g., those generated by hydroxylation of proline and lysine; phosphorylation of the hydroxyl groups of seryl or threonyl residues; methylation of the alpha-amino groups of lysine, arginine and histidine; acetylation of the N-terminal amine; methylation of main chain amide residues or substitution with N-methyl amino acids; or amidation of C-terminal carboxyl groups.

A residue, e.g., an amino acid, of a polypeptide of the invention can also be replaced by an amino acid (or peptidomimetic residue) of the opposite chirality. Thus, any amino acid naturally occurring in the L-configuration (which can also be referred to as the R or S, depending upon the structure of the chemical entity) can be replaced with the
5 amino acid of the same chemical structural type or a peptidomimetic, but of the opposite chirality, referred to as the D- amino acid, but also can be referred to as the R- or S- form.

The invention also provides methods for modifying the polypeptides of the invention by either natural processes, such as post-translational processing (e.g., phosphorylation, acylation, etc), or by chemical modification techniques, and the resulting
10 modified polypeptides. Modifications can occur anywhere in the polypeptide, including the peptide backbone, the amino acid side-chains and the amino or carboxyl termini. It will be appreciated that the same type of modification may be present in the same or varying degrees at several sites in a given polypeptide. Also a given polypeptide may have many types of modifications. Modifications include acetylation, acylation, ADP-
15 ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of a phosphatidylinositol, cross-linking cyclization, disulfide bond formation, demethylation, formation of covalent cross-links, formation of cysteine, formation of pyroglutamate, formylation, gamma-carboxylation,
20 glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, pegylation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, and transfer-RNA mediated addition of amino acids to protein such as arginylation. See, e.g., Creighton, T.E., *Proteins – Structure and Molecular Properties* 2nd Ed., W.H. Freeman and Company, New York
25 (1993); *Posttranslational Covalent Modification of Proteins*, B.C. Johnson, Ed., Academic Press, New York, pp. 1-12 (1983).

Solid-phase chemical peptide synthesis methods can also be used to synthesize the polypeptide or fragments of the invention. Such method have been known in the art since the early 1960's (Merrifield, R. B., *J. Am. Chem. Soc.*, 85:2149-2154, 1963) (See also
30 Stewart, J. M. and Young, J. D., *Solid Phase Peptide Synthesis*, 2nd Ed., Pierce Chemical Co., Rockford, Ill., pp. 11-12)) and have recently been employed in commercially available laboratory peptide design and synthesis kits (Cambridge Research Biochemicals). Such commercially available laboratory kits have generally utilized the teachings of H. M. Geysen et al, *Proc. Natl. Acad. Sci., USA*, 81:3998 (1984) and provide

for synthesizing peptides upon the tips of a multitude of “rods” or “pins” all of which are connected to a single plate. When such a system is utilized, a plate of rods or pins is inverted and inserted into a second plate of corresponding wells or reservoirs, which contain solutions for attaching or anchoring an appropriate amino acid to the pin's or rod's tips. By repeating such a process step, i.e., inverting and inserting the rod's and pin's tips into appropriate solutions, amino acids are built into desired peptides. In addition, a number of available Fmoc peptide synthesis systems are available. For example, assembly of a polypeptide or fragment can be carried out on a solid support using an Applied Biosystems, Inc. Model 431A™ automated peptide synthesizer. Such equipment provides ready access to the peptides of the invention, either by direct synthesis or by synthesis of a series of fragments that can be coupled using other known techniques.

The invention includes isomerases, e.g., racemases, e.g., amino acid racemases, alanine racemases, and/or epimerases of the invention with and without signal. The polypeptide comprising a signal sequence of the invention can be an isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase of the invention or another isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase or another enzyme or other polypeptide.

The invention includes immobilized isomerases, e.g., racemases, e.g., amino acid racemases, alanine racemases, and/or epimerases and anti-isomerase, e.g., anti-racemase, e.g., anti-amino acid racemase, anti-alanine racemase, and/or anti-epimerase antibodies and fragments thereof. The invention provides methods for inhibiting isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase activity, e.g., using dominant negative mutants or anti-isomerase, e.g., anti-racemase, e.g., anti-amino acid racemase, anti-alanine racemase, and/or anti-epimerase antibodies of the invention. The invention includes heterocomplexes, e.g., fusion proteins, heterodimers, etc., comprising the isomerases, e.g., racemases, e.g., amino acid racemases, alanine racemases, and/or epimerases of the invention.

Polypeptides of the invention can have an isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase activity under various conditions, e.g., extremes in pH and/or temperature, oxidizing agents, and the like. The invention provides methods leading to alternative isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase preparations with different catalytic efficiencies and stabilities, e.g., towards temperature, oxidizing agents and changing wash conditions. In one aspect, isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or

epimerase variants can be produced using techniques of site-directed mutagenesis and/or random mutagenesis. In one aspect, directed evolution can be used to produce a great variety of isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase variants with alternative specificities and stability.

5 The proteins of the invention are also useful as research reagents to identify isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase modulators, e.g., activators or inhibitors of isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase activity. Briefly, test samples (compounds, broths, extracts, and the like) are added to isomerase, e.g., racemase, e.g., amino acid
10 racemase, alanine racemase, and/or epimerase assays to determine their ability to inhibit substrate cleavage. Inhibitors identified in this way can be used in industry and research to reduce or prevent undesired proteolysis. Inhibitors can be combined to increase the spectrum of activity.

The enzymes of the invention are also useful as research reagents to digest proteins
15 or in protein sequencing. For example, the isomerases, e.g., racemases, e.g., amino acid racemases, alanine racemases, and/or epimerases may be used to break polypeptides into smaller fragments for sequencing using, e.g. an automated sequencer.

The invention also provides methods of discovering new isomerases, e.g., racemases, e.g., amino acid racemases, alanine racemases, and/or epimerases using the
20 nucleic acids, polypeptides and antibodies of the invention. In one aspect, phagemid libraries are screened for expression-based discovery of isomerases, e.g., racemases, e.g., amino acid racemases, alanine racemases, and/or epimerases. In another aspect, lambda phage libraries are screened for expression-based discovery of isomerases, e.g., racemases, e.g., amino acid racemases, alanine racemases, and/or epimerases. Screening of the phage
25 or phagemid libraries can allow the detection of toxic clones; improved access to substrate; reduced need for engineering a host, by-passing the potential for any bias resulting from mass excision of the library; and, faster growth at low clone densities. Screening of phage or phagemid libraries can be in liquid phase or in solid phase. In one aspect, the invention provides screening in liquid phase. This gives a greater flexibility in
30 assay conditions; additional substrate flexibility; higher sensitivity for weak clones; and ease of automation over solid phase screening.

The invention provides screening methods using the proteins and nucleic acids of the invention and robotic automation to enable the execution of many thousands of biocatalytic reactions and screening assays in a short period of time, e.g., per day, as well

as ensuring a high level of accuracy and reproducibility (see discussion of arrays, below). As a result, a library of derivative compounds can be produced in a matter of weeks. For further teachings on modification of molecules, including small molecules, see PCT/US94/09174.

5 Another aspect of the invention is an isolated or purified polypeptide comprising the sequence of one of the invention and sequences substantially identical thereto, or fragments comprising at least about 5, 10, 15, 20, 25, 30, 35, 40, 50, 75, 100, or 150 consecutive amino acids thereof. As discussed above, such polypeptides may be obtained by inserting a nucleic acid encoding the polypeptide into a vector such that the coding
10 sequence is operably linked to a sequence capable of driving the expression of the encoded polypeptide in a suitable host cell. For example, the expression vector may comprise a promoter, a ribosome binding site for translation initiation and a transcription terminator. The vector may also include appropriate sequences for amplifying expression.

Another aspect of the invention is polypeptides or fragments thereof which have at
15 least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or more than about 95% homology to one of the polypeptides of the invention and sequences substantially identical thereto, or a fragment comprising at least 5, 10, 15, 20, 25, 30, 35, 40, 50, 75, 100, or 150 consecutive amino acids thereof.

20 Homology may be determined using any of the programs described above which aligns the polypeptides or fragments being compared and determines the extent of amino acid identity or similarity between them. It will be appreciated that amino acid "homology" includes conservative amino acid substitutions such as those described above.

The polypeptides or fragments having homology to one of the polypeptides of the
25 invention, or a fragment comprising at least about 5, 10, 15, 20, 25, 30, 35, 40, 50, 75, 100, or 150 consecutive amino acids thereof may be obtained by isolating the nucleic acids encoding them using the techniques described above.

Alternatively, the homologous polypeptides or fragments may be obtained through biochemical enrichment or purification procedures. The sequence of potentially
30 homologous polypeptides or fragments may be determined by gel electrophoresis and/or microsequencing. The sequence of the prospective homologous polypeptide or fragment can be compared to one of the polypeptides of the invention, or a fragment comprising at least about 5, 10, 15, 20, 25, 30, 35, 40, 50, 75, 100, or 150 consecutive amino acids thereof using any of the programs described above.

Another aspect of the invention is an assay for identifying fragments or variants of the invention, which retain the enzymatic function of the polypeptides of the invention. For example the fragments or variants of said polypeptides, may be used to catalyze biochemical reactions, which indicate that the fragment or variant retains the enzymatic activity of the polypeptides of the invention.

The assay for determining if fragments or variants retain the enzymatic activity of the polypeptides of the invention includes the steps of: contacting the polypeptide fragment or variant with a substrate molecule under conditions which allow the polypeptide fragment or variant to function and detecting either a decrease in the level of substrate or an increase in the level of the specific reaction product of the reaction between the polypeptide and substrate.

The present invention exploits the unique catalytic properties of enzymes. Whereas the use of biocatalysts (i.e., purified or crude enzymes, non-living or living cells) in chemical transformations normally requires the identification of a particular biocatalyst that reacts with a specific starting compound, the present invention uses selected biocatalysts and reaction conditions that are specific for functional groups that are present in many starting compounds, such as small molecules. Each biocatalyst is specific for one functional group, or several related functional groups and can react with many starting compounds containing this functional group.

The biocatalytic reactions produce a population of derivatives from a single starting compound. These derivatives can be subjected to another round of biocatalytic reactions to produce a second population of derivative compounds. Thousands of variations of the original small molecule or compound can be produced with each iteration of biocatalytic derivatization.

Enzymes react at specific sites of a starting compound without affecting the rest of the molecule, a process which is very difficult to achieve using traditional chemical methods. This high degree of biocatalytic specificity provides the means to identify a single active compound within the library. The library is characterized by the series of biocatalytic reactions used to produce it, a so called "biosynthetic history". Screening the library for biological activities and tracing the biosynthetic history identifies the specific reaction sequence producing the active compound. The reaction sequence is repeated and the structure of the synthesized compound determined. This mode of identification, unlike other synthesis and screening approaches, does not require immobilization technologies and compounds can be synthesized and tested free in solution using virtually any type of

screening assay. It is important to note, that the high degree of specificity of enzyme reactions on functional groups allows for the "tracking" of specific enzymatic reactions that make up the biocatalytically produced library.

5 Many of the procedural steps are performed using robotic automation enabling the execution of many thousands of biocatalytic reactions and screening assays per day as well as ensuring a high level of accuracy and reproducibility. As a result, a library of derivative compounds can be produced in a matter of weeks which would take years to produce using current chemical methods.

10 In a particular aspect, the invention provides a method for modifying small molecules, comprising contacting a polypeptide encoded by a polynucleotide described herein or enzymatically active fragments thereof with a small molecule to produce a modified small molecule. A library of modified small molecules is tested to determine if a modified small molecule is present within the library which exhibits a desired activity. A specific biocatalytic reaction which produces the modified small molecule of desired activity is identified by systematically eliminating each of the biocatalytic reactions used to produce a portion of the library and then testing the small molecules produced in the portion of the library for the presence or absence of the modified small molecule with the desired activity. The specific biocatalytic reactions which produce the modified small molecule of desired activity is in one aspect (optionally) repeated. The biocatalytic reactions are conducted with a group of biocatalysts that react with distinct structural moieties found within the structure of a small molecule, each biocatalyst is specific for one structural moiety or a group of related structural moieties; and each biocatalyst reacts with many different small molecules which contain the distinct structural moiety.

Isomerase signal sequences, prepro and catalytic domains

25 The invention provides isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase signal sequences (e.g., signal peptides (SPs)), prepro domains and catalytic domains (CDs). The SPs, prepro domains and/or CDs of the invention can be isolated, synthetic or recombinant peptides or can be part of a fusion protein, e.g., as a heterologous domain in a chimeric protein. The invention provides nucleic acids encoding these catalytic domains (CDs), prepro domains and signal sequences (SPs, e.g., a peptide having a sequence comprising/ consisting of amino terminal residues of a polypeptide of the invention). In one aspect, the invention provides a signal sequence comprising a peptide comprising/ consisting of a sequence as set forth in

30

residues 1 to 12, 1 to 13, 1 to 14, 1 to 15, 1 to 16, 1 to 17, 1 to 18, 1 to 19, 1 to 20, 1 to 21, 1 to 22, 1 to 23, 1 to 24, 1 to 25, 1 to 26, 1 to 27, 1 to 28, 1 to 28, 1 to 30, 1 to 31, 1 to 32, 1 to 33, 1 to 34, 1 to 35, 1 to 36, 1 to 37, 1 to 38, 1 to 39, 1 to 40, 1 to 41, 1 to 42, 1 to 43, 1 to 44, 1 to 45, 1 to 46, 1 to 47, 1 to 48, 1 to 49 or 1 to 50, of a polypeptide of the
5 invention.

The isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase signal sequences (SPs) and/or prepro sequences of the invention can be isolated peptides, or, sequences joined to another isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase or a non-isomerase, e.g., non-racemase,
10 e.g., non-amino acid racemase, non-alanine racemase, and/or non-epimerase polypeptide, e.g., as a fusion (chimeric) protein. In one aspect, the invention provides polypeptides comprising isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase signal sequences of the invention. In one aspect, polypeptides comprising isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase
15 signal sequences SPs and/or prepro of the invention comprise sequences heterologous to an isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase of the invention (e.g., a fusion protein comprising an SP and/or prepro of the invention and sequences from another isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase or a non-isomerase, e.g., non-racemase,
20 e.g., non-amino acid racemase, non-alanine racemase, and/or non-epimerase polypeptide protein). In one aspect, the invention provides isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase of the invention with heterologous SPs and/or prepro sequences, e.g., sequences with a yeast signal sequence. An isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase of the invention
25 can comprise a heterologous SP and/or prepro in a vector, e.g., a pPIC series vector (Invitrogen, Carlsbad, CA).

In one aspect, SPs and/or prepro sequences of the invention are identified following identification of novel isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase polypeptides. The pathways by which proteins are
30 sorted and transported to their proper cellular location are often referred to as protein targeting pathways. One of the most important elements in all of these targeting systems is a short amino acid sequence at the amino terminus of a newly synthesized polypeptide called the signal sequence. This signal sequence directs a protein to its appropriate location in the cell and is removed during transport or when the protein reaches its final

destination. Most lysosomal, membrane, or secreted proteins have an amino-terminal signal sequence that marks them for translocation into the lumen of the endoplasmic reticulum. More than 100 signal sequences for proteins in this group have been determined. The signal sequences can vary in length from between about 10 to 50 amino acid residues. Various methods of recognition of signal sequences are known to those of skill in the art. For example, in one aspect, novel isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase signal peptides are identified by a method referred to as SignalP. SignalP uses a combined neural network which recognizes both signal peptides and their cleavage sites; see, e.g., Nielsen (1997) "Identification of prokaryotic and eukaryotic signal peptides and prediction of their cleavage sites." Protein Engineering 10:1-6.

It should be understood that in some aspects isomerases, e.g., racemases, e.g., amino acid racemases, alanine racemases, and/or epimerases of the invention may not have SPs and/or prepro sequences, or "domains." In one aspect, the invention provides the isomerases, e.g., racemases, e.g., amino acid racemases, alanine racemases, and/or epimerases of the invention lacking all or part of an SP and/or a prepro domain. In one aspect, the invention provides a nucleic acid sequence encoding a signal sequence (SP) and/or prepro from one isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase operably linked to a nucleic acid sequence of a different isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase or, in one aspect (optionally), a signal sequence (SPs) and/or prepro domain from a non-isomerase, e.g., non-racemase, e.g., non-amino acid racemase, non-alanine racemase, and/or non-epimerase protein may be desired.

The invention also provides isolated, synthetic or recombinant polypeptides comprising signal sequences (SPs), prepro domain and/or catalytic domains (CDs) of the invention and heterologous sequences. The heterologous sequences are sequences not naturally associated (e.g., to an isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase) with an SP, prepro domain and/or CD. The sequence to which the SP, prepro domain and/or CD are not naturally associated can be on the SP's, prepro domain and/or CD's amino terminal end, carboxy terminal end, and/or on both ends of the SP and/or CD. In one aspect, the invention provides an isolated, synthetic or recombinant polypeptide comprising (or consisting of) a polypeptide comprising a signal sequence (SP), prepro domain and/or catalytic domain (CD) of the invention with the proviso that it is not associated with any sequence to which it is naturally associated (e.g.,

an isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase sequence). Similarly in one aspect, the invention provides isolated, synthetic or recombinant nucleic acids encoding these polypeptides. Thus, in one aspect, the isolated, synthetic or recombinant nucleic acid of the invention comprises coding sequence
5 for a signal sequence (SP), prepro domain and/or catalytic domain (CD) of the invention and a heterologous sequence (i.e., a sequence not naturally associated with the a signal sequence (SP), prepro domain and/or catalytic domain (CD) of the invention). The heterologous sequence can be on the 3' terminal end, 5' terminal end, and/or on both ends of the SP, prepro domain and/or CD coding sequence.

10 Hybrid (chimeric) isomerases and peptide libraries

In one aspect, the invention provides hybrid isomerases, e.g., racemases, e.g., amino acid racemases, alanine racemases, and/or epimerases and fusion proteins, including peptide libraries, comprising sequences of the invention. The peptide libraries of the invention can be used to isolate peptide modulators (e.g., activators or inhibitors) of
15 targets, such as isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase substrates, receptors, enzymes. The peptide libraries of the invention can be used to identify formal binding partners of targets, such as ligands, e.g., cytokines, hormones and the like. In one aspect, the invention provides chimeric proteins comprising a signal sequence (SP), prepro domain and/or catalytic domain (CD) of the invention or a
20 combination thereof and a heterologous sequence (see above).

In one aspect, the fusion proteins of the invention (e.g., the peptide moiety) are conformationally stabilized (relative to linear peptides) to allow a higher binding affinity for targets. The invention provides fusions of isomerases, e.g., racemases, e.g., amino acid racemases, alanine racemases, and/or epimerases of the invention and other peptides,
25 including known and random peptides. They can be fused in such a manner that the structure of the isomerases, e.g., racemases, e.g., amino acid racemases, alanine racemases, and/or epimerases is not significantly perturbed and the peptide is metabolically or structurally conformationally stabilized. This allows the creation of a peptide library that is easily monitored both for its presence within cells and its quantity.

30 Amino acid sequence variants of the invention can be characterized by a predetermined nature of the variation, a feature that sets them apart from a naturally occurring form, e.g., an allelic or interspecies variation of an isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase sequence. In one aspect,

the variants of the invention exhibit the same qualitative biological activity as the naturally occurring analogue. Alternatively, the variants can be selected for having modified characteristics. In one aspect, while the site or region for introducing an amino acid sequence variation is predetermined, the mutation per se need not be predetermined. For example, in order to optimize the performance of a mutation at a given site, random mutagenesis may be conducted at the target codon or region and the expressed isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase variants screened for the optimal combination of desired activity. Techniques for making substitution mutations at predetermined sites in DNA having a known sequence are well known, as discussed herein for example, M13 primer mutagenesis and PCR mutagenesis. Screening of the mutants can be done using, e.g., assays of isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase activity. In alternative aspects, amino acid substitutions can be single residues; insertions can be on the order of from about 1 to 20 amino acids, although considerably larger insertions can be done. Deletions can range from about 1 to about 20, 30, 40, 50, 60, 70 residues or more. To obtain a final derivative with the optimal properties, substitutions, deletions, insertions or any combination thereof may be used. Generally, these changes are done on a few amino acids to minimize the alteration of the molecule. However, larger changes may be tolerated in certain circumstances.

The invention provides isomerases, e.g., racemases, e.g., amino acid racemases, alanine racemases, and/or epimerases where the structure of the polypeptide backbone, the secondary or the tertiary structure, e.g., an alpha-helical or beta-sheet structure, has been modified. In one aspect, the charge or hydrophobicity has been modified. In one aspect, the bulk of a side chain has been modified. Substantial changes in function or immunological identity are made by selecting substitutions that are less conservative. For example, substitutions can be made which more significantly affect: the structure of the polypeptide backbone in the area of the alteration, for example a alpha-helical or a beta-sheet structure; a charge or a hydrophobic site of the molecule, which can be at an active site; or a side chain. The invention provides substitutions in polypeptide of the invention where (a) a hydrophilic residues, e.g. seryl or threonyl, is substituted for (or by) a hydrophobic residue, e.g. leucyl, isoleucyl, phenylalanyl, valyl or alanyl; (b) a cysteine or proline is substituted for (or by) any other residue; (c) a residue having an electropositive side chain, e.g. lysyl, arginyl, or histidyl, is substituted for (or by) an electronegative residue, e.g. glutamyl or aspartyl; or (d) a residue having a bulky side chain, e.g.

DEMANDE OU BREVET VOLUMINEUX

LA PRÉSENTE PARTIE DE CETTE DEMANDE OU CE BREVET COMPREND PLUS D'UN TOME.

CECI EST LE TOME 1 DE 2
CONTENANT LES PAGES 1 À 311

NOTE : Pour les tomes additionels, veuillez contacter le Bureau canadien des brevets

JUMBO APPLICATIONS/PATENTS

THIS SECTION OF THE APPLICATION/PATENT CONTAINS MORE THAN ONE VOLUME

THIS IS VOLUME 1 OF 2
CONTAINING PAGES 1 TO 311

NOTE: For additional volumes, please contact the Canadian Patent Office

NOM DU FICHER / FILE NAME :

NOTE POUR LE TOME / VOLUME NOTE:

WHAT IS CLAIMED IS:

1. An isolated, synthetic or recombinant nucleic acid (polynucleotide) comprising

(a) a nucleic acid (polynucleotide) encoding at least one polypeptide, wherein the
5 nucleic acid comprises a sequence having at least about 50%, 51%, 52%, 53%, 54%, 55%,
56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%,
71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%,
86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more
or complete (100%) sequence identity to the nucleic acid (polynucleotide) sequence of
10 SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID
NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21,
SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO:31, SEQ
ID NO:33, SEQ ID NO:35, SEQ ID NO:37, SEQ ID NO:39, SEQ ID NO:41, SEQ ID
NO:43, SEQ ID NO:45, SEQ ID NO:47, SEQ ID NO:49, SEQ ID NO:51, SEQ ID NO:53,
15 SEQ ID NO:55, SEQ ID NO:57, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ
ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID
NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85,
SEQ ID NO:87, SEQ ID NO:89, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95, SEQ
ID NO:97, SEQ ID NO:99, SEQ ID NO:101, SEQ ID NO:103, SEQ ID NO:105, SEQ ID
20 NO:107, SEQ ID NO:109, SEQ ID NO:111, SEQ ID NO:113, SEQ ID NO:115, SEQ ID
NO:117, SEQ ID NO:119, SEQ ID NO:121, SEQ ID NO:123, SEQ ID NO:125, SEQ ID
NO:127, SEQ ID NO:129, SEQ ID NO:131, SEQ ID NO:133, SEQ ID NO:135, SEQ ID
NO:137, SEQ ID NO:139, SEQ ID NO:141, SEQ ID NO:143, SEQ ID NO:145, SEQ ID
NO:147, SEQ ID NO:149, SEQ ID NO:151, SEQ ID NO:153, SEQ ID NO:155, SEQ ID
25 NO:157, SEQ ID NO:159, SEQ ID NO:161, SEQ ID NO:163, SEQ ID NO:165, SEQ ID
NO:167, SEQ ID NO:169, SEQ ID NO:171, SEQ ID NO:173, SEQ ID NO:175, SEQ ID
NO:177, SEQ ID NO:179, SEQ ID NO:181, SEQ ID NO:183, SEQ ID NO:185, SEQ ID
NO:187, SEQ ID NO:189, SEQ ID NO:191, SEQ ID NO:193, SEQ ID NO:195, SEQ ID
NO:197, SEQ ID NO:199, SEQ ID NO:201, SEQ ID NO:203, SEQ ID NO:205, SEQ ID
30 NO:207, SEQ ID NO:209, SEQ ID NO:211, SEQ ID NO:213, SEQ ID NO:215, SEQ ID
NO:217, SEQ ID NO:219, SEQ ID NO:221, SEQ ID NO:223, SEQ ID NO:225, SEQ ID
NO:227, SEQ ID NO:229, SEQ ID NO:231, SEQ ID NO:233, SEQ ID NO:235, SEQ ID
NO:237, SEQ ID NO:239, SEQ ID NO:241, SEQ ID NO:243, SEQ ID NO:245, SEQ ID
NO:247, SEQ ID NO:249, SEQ ID NO:251, SEQ ID NO:253, SEQ ID NO:255, SEQ ID

NO:257, SEQ ID NO:259, SEQ ID NO:261, SEQ ID NO:263, SEQ ID NO:265, SEQ ID
NO:267, SEQ ID NO:269, SEQ ID NO:271, SEQ ID NO:273, SEQ ID NO:275, SEQ ID
NO:277, SEQ ID NO:279, SEQ ID NO:281, SEQ ID NO:283, SEQ ID NO:285, SEQ ID
NO:287, SEQ ID NO:289, SEQ ID NO:291, SEQ ID NO:293, SEQ ID NO:295, SEQ ID
5 NO:297, SEQ ID NO:299, SEQ ID NO:301, SEQ ID NO:303, SEQ ID NO:305, SEQ ID
NO:307, SEQ ID NO:309, SEQ ID NO:311, SEQ ID NO:313, SEQ ID NO:315, SEQ ID
NO:317, SEQ ID NO:319, SEQ ID NO:321, SEQ ID NO:323, SEQ ID NO:325, SEQ ID
NO:327, SEQ ID NO:329, SEQ ID NO:331, SEQ ID NO:333, SEQ ID NO:335, SEQ ID
NO:337, SEQ ID NO:339, SEQ ID NO:341, SEQ ID NO:343, SEQ ID NO:345, SEQ ID
10 NO:347, SEQ ID NO:349, SEQ ID NO:351, SEQ ID NO:353, SEQ ID NO:355, SEQ ID
NO:357, SEQ ID NO:359, SEQ ID NO:361, SEQ ID NO:363, SEQ ID NO:365, SEQ ID
NO:367, SEQ ID NO:369, SEQ ID NO:371, SEQ ID NO:373, SEQ ID NO:375, SEQ ID
NO:377, SEQ ID NO:379, SEQ ID NO:381, SEQ ID NO:383, SEQ ID NO:385, SEQ ID
NO:387, SEQ ID NO:389, SEQ ID NO:391, SEQ ID NO:393, SEQ ID NO:395, SEQ ID
15 NO:397, SEQ ID NO:399, SEQ ID NO:401, SEQ ID NO:403, SEQ ID NO:405, SEQ ID
NO:407, SEQ ID NO:409, SEQ ID NO:411, SEQ ID NO:413, SEQ ID NO:415, SEQ ID
NO:417, SEQ ID NO:419, SEQ ID NO:421, SEQ ID NO:423, SEQ ID NO:425, SEQ ID
NO:427, SEQ ID NO:429, SEQ ID NO:431, SEQ ID NO:433, SEQ ID NO:435, SEQ ID
NO:437, SEQ ID NO:439, SEQ ID NO:441, SEQ ID NO:443, SEQ ID NO:445, SEQ ID
20 NO:447, SEQ ID NO:449, SEQ ID NO:451, SEQ ID NO:453, SEQ ID NO:455, SEQ ID
NO:457, SEQ ID NO:459, SEQ ID NO:461, SEQ ID NO:463, SEQ ID NO:465, SEQ ID
NO:467, SEQ ID NO:469, SEQ ID NO:471, SEQ ID NO:473, SEQ ID NO:475, SEQ ID
NO:477, SEQ ID NO:479, SEQ ID NO:481, SEQ ID NO:483, SEQ ID NO:485, SEQ ID
NO:487, SEQ ID NO:489, SEQ ID NO:491, SEQ ID NO:493, SEQ ID NO:495 or SEQ ID
25 NO:497, wherein the nucleic acid encodes at least one polypeptide having an isomerase
activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine
racemase activity, and/or an epimerase activity, or encodes a polypeptide or peptide
capable of generating an isomerase specific antibody, e.g., a racemase specific antibody,
e.g., an amino acid racemase, an alanine racemase, and/or an epimerase specific antibody
30 (a polypeptide or peptide that acts as an epitope or immunogen),

(b) the nucleic acid (polynucleotide) of (a), wherein the sequence identities are
determined: (A) by analysis with a sequence comparison algorithm or by a visual
inspection, or (B) over a region of at least about 20, 30, 40, 50, 75, 100, 150, 200, 250,

300, 350, 400, 450, 500, 550, 600, 650, 700, 750, 800, 850, 900, 950, 1000, 1050, 1100, 1150 or more residues, or over the full length of a cDNA, transcript (mRNA) or gene;

(c) the nucleic acid (polynucleotide) of (a) or (b), wherein the sequence comparison algorithm is a BLAST version 2.2.2 algorithm where a filtering setting is set to blastall -p blastp -d "nr pataa" -F F, and all other options are set to default;

(d) a nucleic acid (polynucleotide) encoding at least one polypeptide or peptide, wherein the nucleic acid comprises a sequence that hybridizes under stringent conditions to a nucleic acid comprising the nucleic acid (polynucleotide) sequence of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO:31, SEQ ID NO:33, SEQ ID NO:35, SEQ ID NO:37, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:43, SEQ ID NO:45, SEQ ID NO:47, SEQ ID NO:49, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:57, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:89, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95, SEQ ID NO:97, SEQ ID NO:99, SEQ ID NO:101, SEQ ID NO:103, SEQ ID NO:105, SEQ ID NO:107, SEQ ID NO:109, SEQ ID NO:111, SEQ ID NO:113, SEQ ID NO:115, SEQ ID NO:117, SEQ ID NO:119, SEQ ID NO:121, SEQ ID NO:123, SEQ ID NO:125, SEQ ID NO:127, SEQ ID NO:129, SEQ ID NO:131, SEQ ID NO:133, SEQ ID NO:135, SEQ ID NO:137, SEQ ID NO:139, SEQ ID NO:141, SEQ ID NO:143, SEQ ID NO:145, SEQ ID NO:147, SEQ ID NO:149, SEQ ID NO:151, SEQ ID NO:153, SEQ ID NO:155, SEQ ID NO:157, SEQ ID NO:159, SEQ ID NO:161, SEQ ID NO:163, SEQ ID NO:165, SEQ ID NO:167, SEQ ID NO:169, SEQ ID NO:171, SEQ ID NO:173, SEQ ID NO:175, SEQ ID NO:177, SEQ ID NO:179, SEQ ID NO:181, SEQ ID NO:183, SEQ ID NO:185, SEQ ID NO:187, SEQ ID NO:189, SEQ ID NO:191, SEQ ID NO:193, SEQ ID NO:195, SEQ ID NO:197, SEQ ID NO:199, SEQ ID NO:201, SEQ ID NO:203, SEQ ID NO:205, SEQ ID NO:207, SEQ ID NO:209, SEQ ID NO:211, SEQ ID NO:213, SEQ ID NO:215, SEQ ID NO:217, SEQ ID NO:219, SEQ ID NO:221, SEQ ID NO:223, SEQ ID NO:225, SEQ ID NO:227, SEQ ID NO:229, SEQ ID NO:231, SEQ ID NO:233, SEQ ID NO:235, SEQ ID NO:237, SEQ ID NO:239, SEQ ID NO:241, SEQ ID NO:243, SEQ ID NO:245, SEQ ID NO:247, SEQ ID NO:249, SEQ ID NO:251, SEQ ID NO:253, SEQ ID NO:255, SEQ ID NO:257, SEQ ID NO:259, SEQ ID NO:261, SEQ ID NO:263, SEQ ID NO:265, SEQ ID NO:267, SEQ ID

NO:269, SEQ ID NO:271, SEQ ID NO:273, SEQ ID NO:275, SEQ ID NO:277, SEQ ID
NO:279, SEQ ID NO:281, SEQ ID NO:283, SEQ ID NO:285, SEQ ID NO:287, SEQ ID
NO:289, SEQ ID NO:291, SEQ ID NO:293, SEQ ID NO:295, SEQ ID NO:297, SEQ ID
NO:299, SEQ ID NO:301, SEQ ID NO:303, SEQ ID NO:305, SEQ ID NO:307, SEQ ID
5 NO:309, SEQ ID NO:311, SEQ ID NO:313, SEQ ID NO:315, SEQ ID NO:317, SEQ ID
NO:319, SEQ ID NO:321, SEQ ID NO:323, SEQ ID NO:325, SEQ ID NO:327, SEQ ID
NO:329, SEQ ID NO:331, SEQ ID NO:333, SEQ ID NO:335, SEQ ID NO:337, SEQ ID
NO:339, SEQ ID NO:341, SEQ ID NO:343, SEQ ID NO:345, SEQ ID NO:347, SEQ ID
NO:349, SEQ ID NO:351, SEQ ID NO:353, SEQ ID NO:355, SEQ ID NO:357, SEQ ID
10 NO:359, SEQ ID NO:361, SEQ ID NO:363, SEQ ID NO:365, SEQ ID NO:367, SEQ ID
NO:369, SEQ ID NO:371, SEQ ID NO:373, SEQ ID NO:375, SEQ ID NO:377, SEQ ID
NO:379, SEQ ID NO:381, SEQ ID NO:383, SEQ ID NO:385, SEQ ID NO:387, SEQ ID
NO:389, SEQ ID NO:391, SEQ ID NO:393, SEQ ID NO:395, SEQ ID NO:397, SEQ ID
NO:399, SEQ ID NO:401, SEQ ID NO:403, SEQ ID NO:405, SEQ ID NO:407, SEQ ID
15 NO:409, SEQ ID NO:411, SEQ ID NO:413, SEQ ID NO:415, SEQ ID NO:417, SEQ ID
NO:419, SEQ ID NO:421, SEQ ID NO:423, SEQ ID NO:425, SEQ ID NO:427, SEQ ID
NO:429, SEQ ID NO:431, SEQ ID NO:433, SEQ ID NO:435, SEQ ID NO:437, SEQ ID
NO:439, SEQ ID NO:441, SEQ ID NO:443, SEQ ID NO:445, SEQ ID NO:447, SEQ ID
NO:449, SEQ ID NO:451, SEQ ID NO:453, SEQ ID NO:455, SEQ ID NO:457, SEQ ID
20 NO:459, SEQ ID NO:461, SEQ ID NO:463, SEQ ID NO:465, SEQ ID NO:467, SEQ ID
NO:469, SEQ ID NO:471, SEQ ID NO:473, SEQ ID NO:475, SEQ ID NO:477, SEQ ID
NO:479, SEQ ID NO:481, SEQ ID NO:483, SEQ ID NO:485, SEQ ID NO:487, SEQ ID
NO:489, SEQ ID NO:491, SEQ ID NO:493, SEQ ID NO:495 or SEQ ID NO:497,

wherein the polypeptide or peptide has an isomerase activity, e.g., a racemase
25 activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an
epimerase activity or is capable of generating an isomerase-specific antibody, e.g., a
racemase-specific antibody, e.g., an amino acid racemase-specific antibody, an alanine
racemase-specific antibody, and/or an epimerase-specific antibody (a polypeptide or
peptide that acts as an epitope or immunogen),

30 and the stringent conditions comprise a wash step comprising a wash in 0.2X SSC
at a temperature of about 65°C for about 15 minutes;

(e) the nucleic acid (polynucleotide) of any of (a) to (d) having a length of at least about 20, 25, 30, 50, 75, 100, 125, 150, 175, 200, 225, 300, 350, 400, 450, 500, 550, 600, 650, 700, 750, 800, 850, 900, 950, 1000, 1050, 1100, 1150 or more nucleotide residues, or the full length of a gene or a transcript;

5 (f) a nucleic acid (polynucleotide) encoding at least one polypeptide having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity, wherein the polypeptide comprises the sequence of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, 10 SEQ ID NO:22, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:28, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:42, SEQ ID NO:44, SEQ ID NO:46, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:58, SEQ ID NO:60, SEQ ID NO:62, SEQ ID NO:64, SEQ ID NO:66, SEQ ID NO:68, SEQ ID NO:70, SEQ ID NO:72, SEQ ID NO:74, SEQ ID NO:76, SEQ ID NO:78, SEQ ID NO:80, SEQ ID NO:82, SEQ ID NO:84, 15 SEQ ID NO:86, SEQ ID NO:88, SEQ ID NO:90, SEQ ID NO:92, SEQ ID NO:94, SEQ ID NO:96, SEQ ID NO:98, SEQ ID NO:100, SEQ ID NO:102, SEQ ID NO:104, SEQ ID NO:106, SEQ ID NO:108, SEQ ID NO:110, SEQ ID NO:112, SEQ ID NO:114, SEQ ID NO:116, SEQ ID NO:118, SEQ ID NO:120, SEQ ID NO:122, SEQ ID NO:124, SEQ ID NO:126, SEQ ID NO:128, SEQ ID NO:130, SEQ ID NO:132, SEQ ID NO:134, SEQ ID NO:136, SEQ ID NO:138, SEQ ID NO:140, SEQ ID NO:142, SEQ ID NO:143, SEQ ID NO:146, SEQ ID NO:148, SEQ ID NO:150, SEQ ID NO:152, SEQ ID NO:154, SEQ ID NO:156, SEQ ID NO:158, SEQ ID NO:160, SEQ ID NO:162, SEQ ID NO:164, SEQ ID NO:166, SEQ ID NO:168, SEQ ID NO:170, SEQ ID NO:172, SEQ ID NO:174, SEQ ID NO:176, SEQ ID NO:178, SEQ ID NO:180, SEQ ID NO:182, SEQ ID NO:184, SEQ ID NO:186, SEQ ID NO:188, SEQ ID NO:190, SEQ ID NO:192, SEQ ID NO:194, SEQ ID NO:196, SEQ ID NO:198, SEQ ID NO:200, SEQ ID NO:202, SEQ ID NO:204, SEQ ID NO:206, SEQ ID NO:208, SEQ ID NO:210, SEQ ID NO:212, SEQ ID NO:214, SEQ ID NO:216, SEQ ID NO:218, SEQ ID NO:220, SEQ ID NO:222, SEQ ID NO:224, SEQ ID NO:226, SEQ ID NO:228, SEQ ID NO:230, SEQ ID NO:232, SEQ ID NO:234, SEQ ID NO:236, SEQ ID NO:238, SEQ ID NO:240, SEQ ID NO:242, SEQ ID NO:244, SEQ ID NO:246, SEQ ID NO:248, SEQ ID NO:250, SEQ ID NO:252, SEQ ID NO:254, SEQ ID NO:256, SEQ ID NO:258, SEQ ID NO:260, SEQ ID NO:262, SEQ ID NO:264, SEQ ID NO:266, SEQ ID NO:268, SEQ ID NO:270, SEQ ID NO:272, SEQ ID NO:274, SEQ ID

NO:276, SEQ ID NO:278, SEQ ID NO:280, SEQ ID NO:282, SEQ ID NO:284, SEQ ID
NO:286, SEQ ID NO:288, SEQ ID NO:290, SEQ ID NO:292, SEQ ID NO:294, SEQ ID
NO:296, SEQ ID NO:298, SEQ ID NO:300, SEQ ID NO:302, SEQ ID NO:304, SEQ ID
NO:306, SEQ ID NO:308, SEQ ID NO:310, SEQ ID NO:312, SEQ ID NO:314, SEQ ID
5 NO:316, SEQ ID NO:318, SEQ ID NO:320, SEQ ID NO:322, SEQ ID NO:324, SEQ ID
NO:326, SEQ ID NO:328, SEQ ID NO:330, SEQ ID NO:332, SEQ ID NO:334, SEQ ID
NO:336, SEQ ID NO:338, SEQ ID NO:340, SEQ ID NO:342, SEQ ID NO:344, SEQ ID
NO:346, SEQ ID NO:348, SEQ ID NO:350, SEQ ID NO:352, SEQ ID NO:354, SEQ ID
NO:356, SEQ ID NO:358, SEQ ID NO:360, SEQ ID NO:362, SEQ ID NO:364, SEQ ID
10 NO:366, SEQ ID NO:368, SEQ ID NO:370, SEQ ID NO:372, SEQ ID NO:374, SEQ ID
NO:376, SEQ ID NO:378, SEQ ID NO:380, SEQ ID NO:382, SEQ ID NO:384, SEQ ID
NO:386, SEQ ID NO:388, SEQ ID NO:390, SEQ ID NO:392, SEQ ID NO:394, SEQ ID
NO:396, SEQ ID NO:398, SEQ ID NO:400, SEQ ID NO:402, SEQ ID NO:404, SEQ ID
NO:406, SEQ ID NO:408, SEQ ID NO:410, SEQ ID NO:412, SEQ ID NO:414, SEQ ID
15 NO:416, SEQ ID NO:418, SEQ ID NO:420, SEQ ID NO:422, SEQ ID NO:424, SEQ ID
NO:426, SEQ ID NO:428, SEQ ID NO:430, SEQ ID NO:432, SEQ ID NO:434, SEQ ID
NO:436, SEQ ID NO:438, SEQ ID NO:440, SEQ ID NO:442, SEQ ID NO:444, SEQ ID
NO:446, SEQ ID NO:448, SEQ ID NO:450, SEQ ID NO:452, SEQ ID NO:454, SEQ ID
NO:456, SEQ ID NO:458, SEQ ID NO:460, SEQ ID NO:462, SEQ ID NO:464, SEQ ID
20 NO:466, SEQ ID NO:468, SEQ ID NO:470, SEQ ID NO:472, SEQ ID NO:474, SEQ ID
NO:476, SEQ ID NO:478, SEQ ID NO:480, SEQ ID NO:482, SEQ ID NO:484, SEQ ID
NO:486, SEQ ID NO:488, SEQ ID NO:490, SEQ ID NO:492, SEQ ID NO:494, SEQ ID
NO:496 or SEQ ID NO:498, or enzymatically active fragments thereof;

(g) (A) the nucleic acid (polynucleotide) of any of (a) to (f) and encoding a
25 polypeptide having at least one conservative amino acid substitution and retaining its
isomerase activity, e.g., racemase activity, e.g., amino acid racemase activity, alanine
racemase activity, and/or epimerase activity; or, (B) the nucleic acid of (i)(A), wherein the
at least one conservative amino acid substitution comprises substituting an amino acid
with another amino acid of like characteristics; or, a conservative substitution comprises:
30 replacement of an aliphatic amino acid with another aliphatic amino acid; replacement of a
Serine with a Threonine or vice versa; replacement of an acidic residue with another acidic
residue; replacement of a residue bearing an amide group with another residue bearing an
amide group; exchange of a basic residue with another basic residue; or replacement of an
aromatic residue with another aromatic residue;

(h) the nucleic acid (polynucleotide) of any of (a) to (g) encoding a polypeptide having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity but lacking a signal sequence, a prepro domain, and/or other domain;

5 (i) the nucleic acid (polynucleotide) of any of (a) to (h) encoding a polypeptide having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity further comprising a heterologous sequence;

(j) the nucleic acid (polynucleotide) of (i), wherein the heterologous sequence
10 comprises, or consists of a sequence encoding: (A) a heterologous signal sequence, a heterologous domain, a heterologous dockerin domain, a heterologous catalytic domain (CD), or a combination thereof; (B) the sequence of (l), wherein the heterologous signal sequence, domain or catalytic domain (CD) is derived from a heterologous enzyme; or,
15 (C) a tag, an epitope, a targeting peptide, a cleavable sequence, a detectable moiety or an enzyme;

(k) the nucleic acid (polynucleotide) of (j), wherein the heterologous signal sequence targets the encoded protein to a vacuole, the endoplasmic reticulum, a chloroplast or a starch granule; or

(l) a nucleic acid sequence (polynucleotide) fully (completely) complementary to
20 the sequence of any of (a) to (k).

2. The isolated, synthetic or recombinant nucleic acid of claim 1, wherein the isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase activity comprises catalyzing the re-arrangement of atoms within a molecule, catalyzing
25 the conversion of one isomer into another, catalyzing the conversion of an optically active substrate into a raceme, which is optically inactive, catalyzing the interconversion of substrate enantiomers, catalyzing the stereochemical inversion around the asymmetric carbon atom in a substrate having only one center of asymmetry, catalyzing the stereochemical inversion of the configuration around an asymmetric carbon atom in a
30 substrate having more than one asymmetric center, and/or catalyzing the racemization of amino acids.

3. The isolated, synthetic or recombinant nucleic acid of claim 1, wherein (a) the isomerase activity, e.g., racemase activity, e.g., amino acid racemase activity, alanine racemase activity, and/or epimerase activity is thermostable; or (b) the polypeptide retains an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity under conditions comprising a temperature range of between 0°C to about 20°C, about 20°C to about 37°C, about 37°C to about 50°C, about 50°C to about 70°C, about 70°C to about 75°C, about 75°C to about 80°C, about 80°C to about 85°C, about 85°C to about 90°C, about 90°C to about 95°C, about 95°C to about 100°C, about 100°C to about 110°C, or higher.

10

4. The isolated, synthetic or recombinant nucleic acid of claim 1, wherein (a) the isomerase activity, e.g., racemase activity, e.g., amino acid racemase activity, alanine racemase activity, and/or epimerase activity is thermotolerant; or (b) the polypeptide retains an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity after exposure to a temperature in the range from about 0°C to about 20°C, about 20°C to about 37°C, about 37°C to about 50°C, about 50°C to about 70°C, about 70°C to about 75°C, about 75°C to about 80°C, about 80°C to about 85°C, about 85°C to about 90°C, about 90°C to about 95°C, about 95°C to about 100°C, about 100°C to about 110°C, or higher.

15

5. The isolated, synthetic or recombinant nucleic acid of claim 1, wherein the isomerase activity, e.g., racemase activity, e.g., amino acid racemase activity, alanine racemase activity, and/or epimerase activity retains activity under acidic conditions comprising about pH 6.5, pH 6, pH 5.5, pH 5, pH 4.5, pH 4.0, pH 3.5, pH 3.0 or less (more acidic) pH, or, retains an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity after exposure to acidic conditions comprising about pH 6.5, pH 6, pH 5.5, pH 5, pH 4.5, pH 4.0, pH 3.5, pH 3.0 or less (more acidic) pH.

20

6. The isolated, synthetic or recombinant nucleic acid of claim 1, wherein the isomerase activity, e.g., racemase activity, e.g., amino acid racemase activity, alanine racemase activity, and/or epimerase activity retains activity under basic conditions comprising about pH 7, pH 7.5, pH 8.0, pH 8.5, pH 9, pH 9.5, pH 10, pH 10.5, pH 11, pH 11.5, pH 12, pH 12.5 or more (more basic) or, retains an isomerase activity, e.g., a

25

30

racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity after exposure to basic conditions comprising about pH 7, pH 7.5, pH 8.0, pH 8.5, pH 9, pH 9.5, pH 10, pH 10.5, pH 11, pH 11.5, pH 12, pH 12.5 or more (more basic).

5

7. A nucleic acid probe for identifying a nucleic acid encoding a polypeptide with an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity, wherein the probe comprises at least about 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 100, 125, 150, 175, 200,
10 225 or more consecutive bases of the nucleic acid of any of claims 1 to 6.

8. An amplification primer pair for amplifying a nucleic acid encoding a polypeptide having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity, wherein (a)
15 the primer pair is capable of amplifying a nucleic acid comprising the sequence of any of claims 1 to 6; or (b) the primer pair of (a), wherein a member of the amplification primer pair comprises an oligonucleotide comprising at least about 10 to 50 consecutive bases of the sequence, or, about 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27,
20 28, 29, 30, 31, 32, 33, 34, 35 or more consecutive bases of the sequence.

20

9. An amplification primer pair, wherein the primer pair comprises a first member having a sequence as set forth by about the first (the 5') 10, 11, 12, 13, 14, 15, 16,
17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 or more residues of the sequence of any of claims 1 to 6, and a second member having a sequence as set forth
25 by about the first (the 5') 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 or more residues of the complementary strand of the first member.

10. An isomerase-, e.g., a racemase-, e.g., an amino acid racemase-, an alanine
30 racemase-, and/or an epimerase- encoding nucleic acid generated by amplification of a polynucleotide using the amplification primer pair of claim 8 or claim 9, wherein optionally the amplification is by polymerase chain reaction (PCR).

11. The isomerase-, e.g., racemase-, e.g., amino acid racemase-, alanine racemase-, and/or epimerase- encoding nucleic acid of claim 10, wherein the nucleic acid generated by amplification of a gene library, wherein optionally the gene library is an environmental library.

5

12. A method of amplifying a nucleic acid encoding a polypeptide having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity comprising amplification of a template nucleic acid with an amplification primer sequence pair of claim 8 or claim 9.

10

13. An expression cassette, a vector or a cloning vehicle comprising a nucleic acid comprising the sequence of any of claims 1 to 6, wherein optionally the cloning vehicle comprises a viral vector, a plasmid, a phage, a phagemid, a cosmid, a fosmid, a bacteriophage or an artificial chromosome.

15

14. The cloning vehicle of claim 13, wherein the viral vector comprises an adenovirus vector, a retroviral vector or an adeno-associated viral vector, or, the artificial chromosome comprises a bacterial artificial chromosome (BAC), a bacteriophage P1-derived vector (PAC), a yeast artificial chromosome (YAC), or a mammalian artificial chromosome (MAC).

20

15. A transformed cell comprising a nucleic acid having the sequence of claim 1, or comprising the expression cassette, vector or cloning vehicle of claim 13 or claim 14, wherein optionally the cell is a bacterial cell, a mammalian cell, a fungal cell, a yeast cell, an insect cell or a plant cell.

25

16. A transgenic non-human animal comprising a nucleic acid having the sequence of any of claims 1 to 6, or comprising the expression cassette, vector or cloning vehicle of claim 13 or claim 14, or the transformed cell of claim 15, wherein optionally the animal is a mouse, a rat, a rabbit, a sheep, a pig, a chicken, a goat, a fish, or a cow.

30

17. A transgenic plant, plant part or plant seed comprising a nucleic acid having the sequence of claim 1, wherein optionally the plant is a corn plant, a sorghum plant, a potato plant, a tomato plant, a wheat plant, an oilseed plant, a rapeseed plant, a

soybean plant, a rice plant, a barley plant, a grass, a cotton plant, a cottonseed plant, a palm, a sesame plant, a peanut plant, a sunflower plant or a tobacco plant.

18. An antisense oligonucleotide comprising a nucleic acid sequence
5 complementary to or capable of hybridizing under stringent conditions to the sequence of any of claim 1, wherein optionally the antisense oligonucleotide is between about 10 to 50, about 20 to 60, about 30 to 70, about 40 to 80, or about 60 to 100 bases in length,

and optionally the stringent conditions comprise a wash step comprising a wash in 0.2X SSC at a temperature of about 65°C for about 15 minutes.

10

19. A double-stranded inhibitory RNA (RNAi) molecule comprising a subsequence of the sequence of any of any of claims 1 to 6, wherein optionally the RNAi is about 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, or 30 or more duplex nucleotides in length.

15

20. A method of inhibiting the translation of an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase message in a cell, or inhibiting the expression of a an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase in a cell, comprising administering to
20 the cell or expressing in the cell the antisense oligonucleotide of claim 18 or the double-stranded inhibitory RNA (RNAi) molecule of claim 19.

21. An isolated, synthetic or recombinant polypeptide or peptide having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an
25 alanine racemase activity, and/or an epimerase activity

(a) comprising an amino acid sequence having at least about 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%,
30 98%, 99%, or more, or has 100% (complete) sequence identity to the amino acid sequence of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:22, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:28, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:42, SEQ ID

NO:44, SEQ ID NO:46, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:52, SEQ ID NO:54,
SEQ ID NO:56, SEQ ID NO:58, SEQ ID NO:60, SEQ ID NO:62, SEQ ID NO:64, SEQ
ID NO:66, SEQ ID NO:68, SEQ ID NO:70, SEQ ID NO:72, SEQ ID NO:74, SEQ ID
NO:76, SEQ ID NO:78, SEQ ID NO:80, SEQ ID NO:82, SEQ ID NO:84, SEQ ID NO:86,
5 SEQ ID NO:88, SEQ ID NO:90, SEQ ID NO:92, SEQ ID NO:94, SEQ ID NO:96, SEQ
ID NO:98, SEQ ID NO:100, SEQ ID NO:102, SEQ ID NO:104, SEQ ID NO:106, SEQ ID
NO:108, SEQ ID NO:110, SEQ ID NO:112, SEQ ID NO:114, SEQ ID NO:116, SEQ ID
NO:118, SEQ ID NO:120, SEQ ID NO:122, SEQ ID NO:124, SEQ ID NO:126, SEQ ID
NO:128, SEQ ID NO:130, SEQ ID NO:132, SEQ ID NO:134, SEQ ID NO:136, SEQ ID
10 NO:138, SEQ ID NO:140, SEQ ID NO:142, SEQ ID NO:143, SEQ ID NO:146, SEQ ID
NO:148, SEQ ID NO:150, SEQ ID NO:152, SEQ ID NO:154, SEQ ID NO:156, SEQ ID
NO:158, SEQ ID NO:160, SEQ ID NO:162, SEQ ID NO:164, SEQ ID NO:166, SEQ ID
NO:168, SEQ ID NO:170, SEQ ID NO:172, SEQ ID NO:174, SEQ ID NO:176, SEQ ID
NO:178, SEQ ID NO:180, SEQ ID NO:182, SEQ ID NO:184, SEQ ID NO:186, SEQ ID
15 NO:188, SEQ ID NO:190, SEQ ID NO:192, SEQ ID NO:194, SEQ ID NO:196, SEQ ID
NO:198, SEQ ID NO:200, SEQ ID NO:202, SEQ ID NO:204, SEQ ID NO:206, SEQ ID
NO:208, SEQ ID NO:210, SEQ ID NO:212, SEQ ID NO:214, SEQ ID NO:216, SEQ ID
NO:218, SEQ ID NO:220, SEQ ID NO:222, SEQ ID NO:224, SEQ ID NO:226, SEQ ID
NO:228, SEQ ID NO:230, SEQ ID NO:232, SEQ ID NO:234, SEQ ID NO:236, SEQ ID
20 NO:238, SEQ ID NO:240, SEQ ID NO:242, SEQ ID NO:244, SEQ ID NO:246, SEQ ID
NO:248, SEQ ID NO:250, SEQ ID NO:252, SEQ ID NO:254, SEQ ID NO:256, SEQ ID
NO:258, SEQ ID NO:260, SEQ ID NO:262, SEQ ID NO:264, SEQ ID NO:266, SEQ ID
NO:268, SEQ ID NO:270, SEQ ID NO:272, SEQ ID NO:274, SEQ ID NO:276, SEQ ID
NO:278, SEQ ID NO:280, SEQ ID NO:282, SEQ ID NO:284, SEQ ID NO:286, SEQ ID
25 NO:288, SEQ ID NO:290, SEQ ID NO:292, SEQ ID NO:294, SEQ ID NO:296, SEQ ID
NO:298, SEQ ID NO:300, SEQ ID NO:302, SEQ ID NO:304, SEQ ID NO:306, SEQ ID
NO:308, SEQ ID NO:310, SEQ ID NO:312, SEQ ID NO:314, SEQ ID NO:316, SEQ ID
NO:318, SEQ ID NO:320, SEQ ID NO:322, SEQ ID NO:324, SEQ ID NO:326, SEQ ID
NO:328, SEQ ID NO:330, SEQ ID NO:332, SEQ ID NO:334, SEQ ID NO:336, SEQ ID
30 NO:338, SEQ ID NO:340, SEQ ID NO:342, SEQ ID NO:344, SEQ ID NO:346, SEQ ID
NO:348, SEQ ID NO:350, SEQ ID NO:352, SEQ ID NO:354, SEQ ID NO:356, SEQ ID
NO:358, SEQ ID NO:360, SEQ ID NO:362, SEQ ID NO:364, SEQ ID NO:366, SEQ ID
NO:368, SEQ ID NO:370, SEQ ID NO:372, SEQ ID NO:374, SEQ ID NO:376, SEQ ID
NO:378, SEQ ID NO:380, SEQ ID NO:382, SEQ ID NO:384, SEQ ID NO:386, SEQ ID

NO:388, SEQ ID NO:390, SEQ ID NO:392, SEQ ID NO:394, SEQ ID NO:396, SEQ ID
NO:398, SEQ ID NO:400, SEQ ID NO:402, SEQ ID NO:404, SEQ ID NO:406, SEQ ID
NO:408, SEQ ID NO:410, SEQ ID NO:412, SEQ ID NO:414, SEQ ID NO:416, SEQ ID
NO:418, SEQ ID NO:420, SEQ ID NO:422, SEQ ID NO:424, SEQ ID NO:426, SEQ ID
5 NO:428, SEQ ID NO:430, SEQ ID NO:432, SEQ ID NO:434, SEQ ID NO:436, SEQ ID
NO:438, SEQ ID NO:440, SEQ ID NO:442, SEQ ID NO:444, SEQ ID NO:446, SEQ ID
NO:448, SEQ ID NO:450, SEQ ID NO:452, SEQ ID NO:454, SEQ ID NO:456, SEQ ID
NO:458, SEQ ID NO:460, SEQ ID NO:462, SEQ ID NO:464, SEQ ID NO:466, SEQ ID
NO:468, SEQ ID NO:470, SEQ ID NO:472, SEQ ID NO:474, SEQ ID NO:476, SEQ ID
10 NO:478, SEQ ID NO:480, SEQ ID NO:482, SEQ ID NO:484, SEQ ID NO:486, SEQ ID
NO:488, SEQ ID NO:490, SEQ ID NO:492, SEQ ID NO:494, SEQ ID NO:496 or SEQ ID
NO:498, or enzymatically active fragments thereof, wherein the polypeptide or peptide of
(i) or (ii) has an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase
activity, an alanine racemase activity, and/or an epimerase activity, or the polypeptide or
15 peptide is capable of generating an isomerase specific antibody, e.g., a racemase specific
antibody, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase specific
antibody (a polypeptide or peptide that acts as an epitope or immunogen),

(b) the polypeptide or peptide of (a), wherein the sequence identities are
determined: (A) by analysis with a sequence comparison algorithm or by a visual
20 inspection, or (B) over a region of at least about 20, 25, 30, 35, 40, 45, 50, 55, 60, 75, 100,
150, 200, 250, 300 or more amino acid residues, or over the full length of the polypeptide
or peptide or enzyme, and/or enzymatically active subsequences (fragments) thereof,

(c) the polypeptide or peptide of (a) of (b), wherein the sequence identities are
determined by analysis with a sequence comparison algorithm or by a visual inspection,
25 and optionally the sequence comparison algorithm is a BLAST version 2.2.2 algorithm
where a filtering setting is set to blastall -p blastp -d "nr pataa" -F F, and all other options
are set to default;

(d) an amino acid sequence encoded by the nucleic acid of any of claims 1 to 6,
wherein the polypeptide has (i) an isomerase activity, e.g., a racemase activity, e.g., an
30 amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity,
or, (ii) has immunogenic activity in that it is capable of generating an antibody that
specifically binds to a polypeptide having a sequence of (a), and/or enzymatically active
subsequences (fragments) thereof;

(e) the amino acid sequence of any of (a) to (d), and comprising at least one amino acid residue conservative substitution, and the polypeptide or peptide retains isomerase activity, e.g., racemase activity, e.g., amino acid racemase activity, alanine racemase activity, and/or epimerase activity;

5 (e) the amino acid sequence of (d), wherein the conservative substitution comprises replacement of an aliphatic amino acid with another aliphatic amino acid; replacement of a serine with a threonine or vice versa; replacement of an acidic residue with another acidic residue; replacement of a residue bearing an amide group with another residue bearing an amide group; exchange of a basic residue with another basic residue; or, replacement of an
10 aromatic residue with another aromatic residue, or a combination thereof,

(f) the amino acid sequence of (e), wherein the aliphatic residue comprises Alanine, Valine, Leucine, Isoleucine or a synthetic equivalent thereof; the acidic residue comprises Aspartic acid, Glutamic acid or a synthetic equivalent thereof; the residue comprising an amide group comprises Aspartic acid, Glutamic acid or a synthetic
15 equivalent thereof; the basic residue comprises Lysine, Arginine or a synthetic equivalent thereof; or, the aromatic residue comprises Phenylalanine, Tyrosine or a synthetic equivalent thereof;

(g) the polypeptide of any of (a) to (f) having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity,
20 and/or an epimerase activity but lacking a signal sequence, a prepro domain, and/or other domain,

(h) the polypeptide of any of (a) to (g) having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity further comprising a heterologous sequence;

25 (i) the polypeptide of (h), wherein the heterologous sequence comprises, or consists of: (A) a heterologous signal sequence, a heterologous domain, a heterologous dockerin domain, a heterologous catalytic domain (CD), or a combination thereof; (B) the sequence of (A), wherein the heterologous signal sequence, domain or catalytic domain (CD) is derived from a heterologous enzyme; and/or, (C) a tag, an epitope, a targeting
30 peptide, a cleavable sequence, a detectable moiety or an enzyme;

(j) polypeptide of (i), wherein the heterologous signal sequence targets the encoded protein to a vacuole, the endoplasmic reticulum, a chloroplast or a starch granule;
or

(k) comprising an amino acid sequence encoded by the nucleic acid sequence of any of claims 1 to 6.

22. The isolated, synthetic or recombinant polypeptide of claim 21, wherein the
5 isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase
activity comprises catalyzing the conversion of an optically active substrate into a raceme,
which is optically inactive, catalyzing the interconversion of substrate enantiomers,
catalyzing the stereochemical inversion around the asymmetric carbon atom in a substrate
10 configuration around an asymmetric carbon atom in a substrate having more than one
asymmetric center, and/or catalyzing the racemization of amino acids.

23. The isolated, synthetic or recombinant polypeptide of any of claim 21,
wherein the isomerase activity, e.g., racemase activity, e.g., amino acid racemase activity,
15 alanine racemase activity, and/or epimerase activity is thermostable, and optionally the
polypeptide retains an isomerase activity, e.g., a racemase activity, e.g., an amino acid
racemase activity, an alanine racemase activity, and/or an epimerase activity under
conditions comprising a temperature range of between about 0°C to about 20°C, about
20°C to about 37°C, about 37°C to about 50°C, about 50°C to about 70°C, about 70°C to
20 about 75°C, about 75°C to about 80°C, about 80°C to about 85°C, about 85°C to about
90°C, about 90°C to about 95°C, about 95°C to about 100°C, about 100°C to about 110°C,
or higher.

24. The isolated, synthetic or recombinant polypeptide of any of claim 21,
25 wherein the isomerase activity, e.g., racemase activity, e.g., amino acid racemase activity,
alanine racemase activity, and/or epimerase activity is thermotolerant, and optionally the
polypeptide retains an isomerase activity, e.g., a racemase activity, e.g., an amino acid
racemase activity, an alanine racemase activity, and/or an epimerase activity after
exposure to a temperature in the range from about 0°C to about 20°C, about 20°C to about
30 37°C, about 37°C to about 50°C, about 50°C to about 70°C, about 70°C to about 75°C,
about 75°C to about 80°C, about 80°C to about 85°C, about 85°C to about 90°C, about
90°C to about 95°C, about 95°C to about 100°C, about 100°C to about 110°C, or higher,
wherein optionally the thermotolerance comprises retention of at least half of the
specific activity of the isomerase, e.g., the racemase, e.g., the amino acid racemase, the

alanine racemase, and/or the epimerase at 37°C after being heated to an elevated temperature, or, optionally the thermotolerance comprises retention of specific activity at 37°C in the range from about 500 to about 1200 units per milligram of protein after being heated to an elevated temperature, and optionally the elevated temperature is at least about
5 0°C to about 20°C, about 20°C to about 37°C, about 37°C to about 50°C, about 50°C to about 70°C, about 70°C to about 75°C, about 75°C to about 80°C, about 80°C to about 85°C, about 85°C to about 90°C, about 90°C to about 95°C, about 95°C to about 100°C, about 100°C to about 110°C, or higher.

10 25. The isolated, synthetic or recombinant polypeptide of any of claim 21, wherein the isomerase activity, e.g., racemase activity, e.g., amino acid racemase activity, alanine racemase activity, and/or epimerase activity retains activity under acidic conditions comprising about pH 6.5, pH 6, pH 5.5, pH 5, pH 4.5, pH 4.0, pH 3.5, pH 3.0 or less (more acidic) pH, or, retains an isomerase activity, e.g., a racemase activity, e.g., an
15 amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity after exposure to acidic conditions comprising about pH 6.5, pH 6, pH 5.5, pH 5, pH 4.5, pH 4.0, pH 3.5, pH 3.0 or less (more acidic) pH.

 26. The isolated, synthetic or recombinant polypeptide of any of claim 21,
20 wherein the isomerase activity, e.g., racemase activity, e.g., amino acid racemase activity, alanine racemase activity, and/or epimerase activity retains activity under basic conditions comprising about pH 7, pH 7.5 pH 8.0, pH 8.5, pH 9, pH 9.5, pH 10, pH 10.5, pH 11, pH 11.5, pH 12, pH 12.5 or more (more basic) or, retains an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity,
25 and/or an epimerase activity after exposure to basic conditions comprising about pH 7, pH 7.5 pH 8.0, pH 8.5, pH 9, pH 9.5, pH 10, pH 10.5, pH 11, pH 11.5, pH 12 pH 12.5 or more (more basic).

 27. An isolated, synthetic or recombinant polypeptide comprising a polypeptide
30 of any of claims 21 to 26, and lacking a signal sequence or a prepro sequence.

 28. An isolated, synthetic or recombinant polypeptide comprising a polypeptide of any of claims 21 to 27, and having a heterologous signal sequence or a heterologous prepro sequence.

29. The isolated, synthetic or recombinant polypeptide of any of claims 21 to 28, wherein the isomerase activity, e.g., racemase activity, e.g., amino acid racemase activity, alanine racemase activity, and/or epimerase activity comprises a specific activity
5 at about 37°C in the range from about 100 to about 1000 units per milligram of protein, from about 500 to about 750 units per milligram of protein, from about 500 to about 1200 units per milligram of protein, or from about 750 to about 1000 units per milligram of protein.

10 30. The isolated, synthetic or recombinant polypeptide of any of claims 21 to 29, wherein the polypeptide comprises at least one glycosylation site or further comprises a polysaccharide, wherein optionally the glycosylation is an N-linked glycosylation, and optionally the polypeptide is glycosylated after being expressed in a *P. pastoris* or a *S. pombe*.

15 31. A protein preparation comprising the polypeptide of any of claims 21 to 30, wherein the protein preparation comprises a liquid, a solid or a gel.

20 32. A heterodimer comprising the polypeptide of any of claims 21 to 30, and a second domain, wherein optionally the second domain is a polypeptide and the heterodimer is a fusion protein, or optionally the second domain is an epitope or a tag.

25 33. A homodimer comprising the polypeptide of any of claims 21 to 30, and optionally the homodimer is a fusion protein.

30 34. An immobilized polypeptide, wherein the polypeptide comprises the polypeptide of any of claims 21 to 30, wherein optionally the polypeptide is immobilized on a wood chip, a paper, a cell, a metal, a resin, a polymer, a ceramic, a glass, a microelectrode, a graphitic particle, a bead, a gel, a plate, an array or a capillary tube.

35. An isolated, synthetic or recombinant antibody that specifically binds to the polypeptide of any of claims 21 to 30, wherein optionally the antibody is a monoclonal or a polyclonal antibody, or is a single chained antibody.

36. A hybridoma comprising an antibody as set forth in claim 35.

37. An array comprising: an immobilized polypeptide, wherein the polypeptide comprises the polypeptide of any of claims 21 to 30; an immobilized nucleic acid, wherein
5 the nucleic acid comprises the nucleic acid of any of claims 1 to 6; the antibody of claim 35; or, a combination thereof.

38. A method of isolating or identifying a polypeptide with an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine
10 racemase activity, and/or an epimerase activity comprising the steps of:
(a) providing the antibody of claim 35;
(b) providing a sample comprising polypeptides; and
(c) contacting the sample of step (b) with the antibody of step (a) under conditions wherein the antibody can specifically bind to the polypeptide, thereby isolating or identifying
15 a polypeptide having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity.

39. A method of making an anti-isomerase, e.g., anti-racemase, e.g., anti-amino acid racemase, anti-alanine racemase, and/or anti-epimerase antibody comprising
20 administering to a non-human animal the nucleic acid of any of claims 1 to 6, in an amount sufficient to generate a humoral immune response, thereby making an anti-isomerase, e.g., anti-racemase, e.g., anti-amino acid racemase, anti-alanine racemase, and/or anti-epimerase antibody.

25 40. A method of making an anti-isomerase, e.g., anti-racemase, e.g., anti-amino acid racemase, anti-alanine racemase, and/or anti-epimerase antibody comprising administering to a non-human animal the polypeptide of any of claims 21 to 30, in an amount sufficient to generate a humoral immune response, thereby making an anti-isomerase, e.g., anti-racemase, e.g., anti-amino acid racemase, anti-alanine racemase,
30 and/or anti-epimerase antibody.

41. A method of producing a recombinant polypeptide comprising the steps of:
(a) providing a nucleic acid operably linked to a promoter, wherein the nucleic acid comprises the sequence of any of claims 1 to 6; and (b) expressing the nucleic acid of step

(a) under conditions that allow expression of the polypeptide, thereby producing a recombinant polypeptide, and optionally the method further comprises transforming a host cell with the nucleic acid of step (a) followed by expressing the nucleic acid of step (a), thereby producing a recombinant polypeptide in a transformed cell.

5

42. A method for identifying a polypeptide having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity comprising:

(a) providing the polypeptide of any of claims 21 to 30;

10

(b) providing an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase substrate; and

15

(c) contacting the polypeptide with the substrate of step (b) and detecting a decrease in the amount of substrate or an increase in the amount of a reaction product, wherein a decrease in the amount of the substrate or an increase in the amount of the reaction product detects a polypeptide having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity.

20

43. A method for identifying an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase substrate comprising:

(a) providing the polypeptide of any of claims 21 to 30;

(b) providing a test substrate; and

25

(c) contacting the polypeptide of step (a) with the test substrate of step (b) and detecting a decrease in the amount of substrate or an increase in the amount of reaction product, wherein a decrease in the amount of the substrate or an increase in the amount of a reaction product identifies the test substrate as an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase substrate.

30

44. A method of determining whether a test compound specifically binds to a polypeptide comprising:

(a) expressing a nucleic acid or a vector comprising the nucleic acid under conditions permissive for translation of the nucleic acid to a polypeptide, wherein the nucleic acid has the sequence of any of claims 1 to 6;

(b) providing a test compound;

(c) contacting the polypeptide with the test compound; and
(d) determining whether the test compound of step (b) specifically binds to the polypeptide.

5 45. A method of determining whether a test compound specifically binds to a polypeptide comprising:

(a) providing the polypeptide of any of claims 21 to 30;
(b) providing a test compound;
(c) contacting the polypeptide with the test compound; and
10 (d) determining whether the test compound of step (b) specifically binds to the polypeptide.

 46. A method for identifying a modulator of an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity,
15 and/or an epimerase activity comprising:

(a) providing the polypeptide of any of claims 21 to 30;
(b) providing a test compound;
(c) contacting the polypeptide of step (a) with the test compound of step (b) and measuring an activity of the isomerase, e.g., racemase, e.g., amino acid racemase, alanine
20 racemase, and/or epimerase, wherein a change in the isomerase activity, e.g., racemase activity, e.g., amino acid racemase activity, alanine racemase activity, and/or epimerase activity measured in the presence of the test compound compared to the activity in the absence of the test compound provides a determination that the test compound modulates the isomerase activity, e.g., racemase activity, e.g., amino acid racemase activity, alanine
25 racemase activity, and/or epimerase activity,

 wherein optionally the isomerase activity, e.g., racemase activity, e.g., amino acid racemase activity, alanine racemase activity, and/or epimerase activity is measured by providing an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase substrate and detecting a decrease in the amount of the
30 substrate or an increase in the amount of a reaction product, or, an increase in the amount of the substrate or a decrease in the amount of a reaction product, wherein optionally a decrease in the amount of the substrate or an increase in the amount of the reaction product with the test compound as compared to the amount of substrate or reaction product without the test compound identifies the test compound as an activator of an

isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity,

wherein optionally, an increase in the amount of the substrate or a decrease in the amount of the reaction product with the test compound as compared to the amount of
5 substrate or reaction product without the test compound identifies the test compound as an inhibitor of an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity.

47. A computer system comprising a processor and a data storage device, or,
10 computer readable medium having stored thereon a polypeptide sequence or a nucleic acid sequence, wherein the data storage device or computer readable medium has stored thereon a polypeptide sequence or a nucleic acid sequence, wherein the polypeptide sequence comprises the amino acid sequence of any of claims 21 to 30, a polypeptide encoded the nucleic acid sequence of any of claims 1 to 6,

15 wherein optionally the system further comprises a sequence comparison algorithm and a data storage device having at least one reference sequence stored thereon, or, optionally further comprises an identifier that identifies one or more features in the sequence,

and optionally the sequence comparison algorithm comprises a computer program
20 that indicates polymorphisms.

48. A method for identifying a feature in a sequence comprising the steps of:
(a) reading the sequence using a computer program which identifies one or more features in a sequence, wherein the sequence comprises a polypeptide sequence or a nucleic acid
25 sequence, wherein the polypeptide sequence comprises the amino acid sequence of any of claims 21 to 30, a polypeptide encoded the nucleic acid sequence of any of claims 1 to 6; and (b) identifying one or more features in the sequence with the computer program.

49. A method for comparing a first sequence to a second sequence comprising
30 the steps of: (a) reading the first sequence and the second sequence through use of a computer program which compares sequences, wherein the first sequence comprises a polypeptide sequence or a nucleic acid sequence, wherein the polypeptide sequence comprises the amino acid sequence of any of claims 21 to 30, a polypeptide encoded the

nucleic acid sequence of any of claims 1 to 6; and (b) determining differences between the first sequence and the second sequence with the computer program,

wherein optionally the step of determining differences between the first sequence and the second sequence further comprises the step of identifying polymorphisms,

5 and optionally the method further comprises an identifier that identifies one or more features in a sequence, and optionally the method further comprises reading the first sequence using a computer program and identifying one or more features in the sequence.

10 50. A method for isolating or recovering a nucleic acid encoding a polypeptide with an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity from an environmental sample comprising the steps of:

(a) providing the polynucleotide probe of claim 7;

15 (b) isolating a nucleic acid from the environmental sample or treating the environmental sample such that nucleic acid in the sample is accessible for hybridization to a polynucleotide probe of step (a);

(c) combining the isolated nucleic acid or the treated environmental sample of step (b) with the polynucleotide probe of step (a); and

20 (d) isolating a nucleic acid that specifically hybridizes with the polynucleotide probe of step (a), thereby isolating or recovering a nucleic acid encoding a polypeptide with an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity from an environmental sample,

wherein optionally the environmental sample comprises a water sample, a liquid sample, a soil sample, an air sample or a biological sample;

25 and optionally the biological sample is derived from a bacterial cell, a protozoan cell, an insect cell, a yeast cell, a plant cell, a fungal cell or a mammalian cell.

30 51. A method for isolating or recovering a nucleic acid encoding a polypeptide with an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity from an environmental sample comprising the steps of:

(a) providing the amplification primer pair of claim 8 or claim 9;

(b) isolating a nucleic acid from the environmental sample or treating the environmental sample such that nucleic acid in the sample is accessible for hybridization to the amplification primer pair; and,

(c) combining the nucleic acid of step (b) with the amplification primer pair of step (a) and amplifying nucleic acid from the environmental sample, thereby isolating or recovering a nucleic acid encoding a polypeptide with an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity from an environmental sample.

52. A method of generating a variant of a nucleic acid encoding a polypeptide with an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity comprising the steps of:

(a) providing a template nucleic acid comprising the sequence of any of claims 1 to 6; and

(b) modifying, deleting or adding one or more nucleotides in the template sequence, or a combination thereof, to generate a variant of the template nucleic acid.

wherein optionally the method further comprises expressing the variant nucleic acid to generate a variant isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase polypeptide,

and optionally the modifications, additions or deletions are introduced by a method comprising error-prone PCR, shuffling, oligonucleotide-directed mutagenesis, assembly PCR, sexual PCR mutagenesis, *in vivo* mutagenesis, cassette mutagenesis, recursive ensemble mutagenesis, exponential ensemble mutagenesis, site-specific mutagenesis, gene reassembly, Gene Site Saturation Mutagenesis (GSSM), synthetic ligation reassembly (SLR) and a combination thereof, or, the modifications, additions or deletions are introduced by a method comprising recombination, recursive sequence recombination, phosphothioate-modified DNA mutagenesis, uracil-containing template mutagenesis, gapped duplex mutagenesis, point mismatch repair mutagenesis, repair-deficient host strain mutagenesis, chemical mutagenesis, radiogenic mutagenesis, deletion mutagenesis, restriction-selection mutagenesis, restriction-purification mutagenesis, artificial gene synthesis, ensemble mutagenesis, chimeric nucleic acid multimer creation and a combination thereof,

wherein optionally, the method is iteratively repeated until an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase having an altered or different activity or an altered or different stability from that of a polypeptide encoded by the template nucleic acid is produced,

5 wherein optionally the variant isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase polypeptide is thermotolerant, and retains some activity after being exposed to an elevated temperature, or, optionally the variant isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase polypeptide has increased glycosylation as compared to the isomerase, e.g., the racemase, 10 e.g., the amino acid racemase, the alanine racemase, and/or the epimerase encoded by a template nucleic acid, or optionally the variant isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase polypeptide has an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity under a high temperature, wherein the isomerase, e.g., the 15 racemase, e.g., the amino acid racemase, the alanine racemase, and/or the epimerase encoded by the template nucleic acid is not active under the high temperature,

wherein optionally, the method is iteratively repeated until an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase coding sequence having an altered codon usage from that of the template nucleic acid is produced,

20 wherein optionally the method is iteratively repeated until an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase gene having higher or lower level of message expression or stability from that of the template nucleic acid is produced.

25 53. A method for modifying codons in a nucleic acid encoding a polypeptide with an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity to increase its expression in a host cell, the method comprising:

(a) providing a nucleic acid encoding a polypeptide with an isomerase activity, 30 e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity comprising the sequence of any of claims 1 to 6; and,

(b) identifying a non-preferred or a less preferred codon in the nucleic acid of step (a) and replacing it with a preferred or neutrally used codon encoding the same amino acid as the replaced codon, wherein a preferred codon is a codon over-represented in coding

sequences in genes in the host cell and a non-preferred or less preferred codon is a codon under-represented in coding sequences in genes in the host cell, thereby modifying the nucleic acid to increase its expression in a host cell.

5 54. A method for modifying codons in a nucleic acid encoding an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase polypeptide, the method comprising:

 (a) providing a nucleic acid encoding a polypeptide with an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase
10 activity, and/or an epimerase activity comprising the sequence of any of claims 1 to 6; and,

 (b) identifying a codon in the nucleic acid of step (a) and replacing it with a different codon encoding the same amino acid as the replaced codon, thereby modifying codons in a nucleic acid encoding an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase.

15

 55. A method for modifying codons in a nucleic acid encoding an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase polypeptide to increase its expression in a host cell, the method comprising:

 (a) providing a nucleic acid encoding an isomerase, e.g., a racemase, e.g., an amino
20 acid racemase, an alanine racemase, and/or an epimerase polypeptide comprising the sequence of any of claims 1 to 6; and,

 (b) identifying a non-preferred or a less preferred codon in the nucleic acid of step (a) and replacing it with a preferred or neutrally used codon encoding the same amino acid as the replaced codon, wherein a preferred codon is a codon over-represented in coding
25 sequences in genes in the host cell and a non-preferred or less preferred codon is a codon under-represented in coding sequences in genes in the host cell, thereby modifying the nucleic acid to increase its expression in a host cell.

 56. A method for modifying a codon in a nucleic acid encoding a polypeptide
30 having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity to decrease its expression in a host cell, the method comprising:

(a) providing a nucleic acid encoding an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase polypeptide comprising the sequence of any of claims 1 to 6; and

(b) identifying at least one preferred codon in the nucleic acid of step (a) and replacing it with a non-preferred or less preferred codon encoding the same amino acid as the replaced codon, wherein a preferred codon is a codon over-represented in coding sequences in genes in a host cell and a non-preferred or less preferred codon is a codon under-represented in coding sequences in genes in the host cell, thereby modifying the nucleic acid to decrease its expression in a host cell,

wherein optionally the host cell is a bacterial cell, a fungal cell, an insect cell, a yeast cell, a plant cell or a mammalian cell.

57. A method for producing a library of nucleic acids encoding a plurality of modified isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase active sites or substrate binding sites, wherein the modified active sites or substrate binding sites are derived from a first nucleic acid comprising a sequence encoding a first active site or a first substrate binding site the method comprising:

(a) providing a first nucleic acid encoding a first active site or first substrate binding site, wherein the first nucleic acid sequence comprises a sequence that hybridizes under stringent conditions to the sequence of any of claims 1 to 6, or a subsequence thereof, and the nucleic acid encodes an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase active site or an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase substrate binding site;

(b) providing a set of mutagenic oligonucleotides that encode naturally-occurring amino acid variants at a plurality of targeted codons in the first nucleic acid; and,

(c) using the set of mutagenic oligonucleotides to generate a set of active site-encoding or substrate binding site-encoding variant nucleic acids encoding a range of amino acid variations at each amino acid codon that was mutagenized, thereby producing a library of nucleic acids encoding a plurality of modified isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase active sites or substrate binding sites.

58. The method of claim 57, comprising mutagenizing the first nucleic acid of step (a) or variants by a method comprising an optimized directed evolution system, Gene Site Saturation Mutagenesis (GSSM), a synthetic ligation reassembly (SLR), error-prone PCR, shuffling, oligonucleotide-directed mutagenesis, assembly PCR, sexual PCR
5 mutagenesis, in vivo mutagenesis, cassette mutagenesis, recursive ensemble mutagenesis, exponential ensemble mutagenesis, site-specific mutagenesis, gene reassembly, recombination, recursive sequence recombination, phosphothioate-modified DNA mutagenesis, uracil-containing template mutagenesis, gapped duplex mutagenesis, point mismatch repair mutagenesis, repair-deficient host strain mutagenesis, chemical
10 mutagenesis, radiogenic mutagenesis, deletion mutagenesis, restriction-selection mutagenesis, restriction-purification mutagenesis, artificial gene synthesis, ensemble mutagenesis, chimeric nucleic acid multimer creation or a combination thereof.

59. A method for making a small molecule comprising:

- 15 (a) providing a plurality of biosynthetic enzymes capable of synthesizing or modifying a small molecule, wherein one of the enzymes comprises an isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase encoded by a nucleic acid comprising the sequence any of claims 1 to 6;
- (b) providing a substrate for at least one of the enzymes of step (a); and
- 20 (c) reacting the substrate of step (b) with the enzymes under conditions that facilitate a plurality of biocatalytic reactions to generate a small molecule by a series of biocatalytic reactions.

60. A method for modifying a small molecule comprising:

- 25 (a) providing an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase enzyme, wherein the enzyme comprises the polypeptide of any of claims 21 to 30, or a polypeptide encoded by a nucleic acid comprising the sequence of any of claims 1 to 6;
- (b) providing a small molecule; and
- 30 (c) reacting the enzyme of step (a) with the small molecule of step (b) under conditions that facilitate an enzymatic reaction catalyzed by the isomerase, e.g., the racemase, e.g., the amino acid racemase, the alanine racemase, and/or the epimerase enzyme, thereby modifying a small molecule by an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase enzymatic reaction

wherein optionally the method comprises providing a plurality of small molecule substrates for the enzyme of step (a), thereby generating a library of modified small molecules produced by at least one enzymatic reaction catalyzed by the isomerase, e.g., the racemase, e.g., the amino acid racemase, the alanine racemase, and/or the epimerase enzyme.

61. The method of claim 60, further comprising a plurality of additional enzymes under conditions that facilitate a plurality of biocatalytic reactions by the enzymes to form a library of modified small molecules produced by the plurality of enzymatic reactions, wherein optionally the method further comprises the step of testing the library to determine if a particular modified small molecule which exhibits a desired activity is present within the library, and optionally the step of testing the library further comprises the steps of systematically eliminating all but one of the biocatalytic reactions used to produce a portion of the plurality of the modified small molecules within the library by testing the portion of the modified small molecule for the presence or absence of the particular modified small molecule with a desired activity, and identifying at least one specific biocatalytic reaction that produces the particular modified small molecule of desired activity.

62. A method for determining a functional fragment of an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase enzyme comprising the steps of:

(a) providing an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase enzyme, wherein the enzyme comprises the polypeptide of any of claims 21 to 30, or a polypeptide encoded by a nucleic acid comprising the sequence of any of claims 1 to 6; and

(b) deleting a plurality of amino acid residues from the sequence of step (a) and testing the remaining subsequence for an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity, thereby determining a functional fragment of an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase enzyme,

wherein optionally the isomerase activity, e.g., racemase activity, e.g., amino acid racemase activity, alanine racemase activity, and/or epimerase activity is measured by providing an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine

racemase, and/or an epimerase substrate and detecting a decrease in the amount of the substrate or an increase in the amount of a reaction product.

5 63. A method for whole cell engineering of new or modified phenotypes by using real-time metabolic flux analysis, the method comprising:

(a) making a modified cell by modifying the genetic composition of a cell, wherein the genetic composition is modified by addition to the cell of a nucleic acid comprising a sequence as set forth in any one of claims 1 to 6;

(b) culturing the modified cell to generate a plurality of modified cells;

10 (c) measuring at least one metabolic parameter of the cell by monitoring the cell culture of step (b) in real time; and,

(d) analyzing the data of step (c) to determine if the measured parameter differs from a comparable measurement in an unmodified cell under similar conditions, thereby identifying an engineered phenotype in the cell using real-time metabolic flux analysis,

15 wherein optionally the genetic composition of the cell is modified by a method comprising deletion of a sequence or modification of a sequence in the cell, or, knocking out the expression of a gene,

and optionally the method further comprises selecting a cell comprising a newly engineered phenotype, and optionally further comprising culturing the selected cell,

20 thereby generating a new cell strain comprising a newly engineered phenotype.

64. A chimeric polypeptide comprising at least a first domain comprising a signal peptide (SP) and at least a second domain comprising a heterologous polypeptide or peptide comprising a the amino acid sequence of any of claims 21 to 30, or a subsequence
25 thereof, wherein the heterologous polypeptide or peptide is not naturally associated with the signal peptide (SP), wherein optionally the signal peptide (SP) is not derived from an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase, and optionally the heterologous polypeptide or peptide is amino terminal to, carboxy terminal to or on both ends of the signal peptide (SP) or an isomerase, e.g., a
30 racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase catalytic domain (CD).

65. An isolated, synthetic or recombinant nucleic acid encoding a chimeric polypeptide, wherein the chimeric polypeptide comprises at least a first domain comprising signal peptide (SP) and at least a second domain comprising a heterologous polypeptide or peptide, wherein the signal peptide (SP) comprises a signal sequence as set
5 forth in claim 64.

66. A method of increasing thermotolerance or thermostability of an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase polypeptide, the method comprising glycosylating an isomerase, e.g., a racemase, e.g., an
10 amino acid racemase, an alanine racemase, and/or an epimerase, wherein the polypeptide comprises the polypeptide of any of claims 21 to 30, or a polypeptide encoded by a nucleic acid comprising the sequence of any of claims 1 to 6, or enzymatically active fragments thereof, thereby increasing the thermotolerance or thermostability of the isomerase, e.g.,
15 racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase.

67. A method for overexpressing a recombinant isomerase, e.g., racemase, e.g., amino acid racemase, alanine racemase, and/or epimerase in a cell comprising expressing a vector comprising a nucleic acid sequence as set forth in any of claims 1 to 6, or a
20 nucleic acid encoding the polypeptide of any of claims 21 to 30, or enzymatically active fragments thereof, wherein overexpression is effected by use of a high activity promoter, a dicistronic vector or by gene amplification of the vector.

68. A method of making a transgenic plant comprising:
(a) introducing a heterologous nucleic acid sequence into the cell, wherein the
25 heterologous nucleic sequence comprises the sequence of any of claims 1 to 6, or a nucleic acid encoding the polypeptide of any of claims 21 to 30, or enzymatically active fragments thereof, thereby producing a transformed plant cell;
(b) producing a transgenic plant from the transformed cell,
wherein optionally step (a) further comprises introducing the heterologous nucleic
30 acid sequence by electroporation or microinjection of plant cell protoplasts,
and optionally step (a) comprises introducing the heterologous nucleic acid sequence directly to plant tissue by DNA particle bombardment or by using an *Agrobacterium tumefaciens* host.

69. A method of expressing a heterologous nucleic acid sequence in a plant cell comprising:

(a) transforming the plant cell with a heterologous nucleic acid sequence operably linked to a promoter, wherein the heterologous nucleic acid comprising the sequence of any of claims 1 to 6, or a nucleic acid encoding the polypeptide of any of claims 21 to 30, or enzymatically active fragments thereof;

(b) growing the plant under conditions wherein the heterologous nucleic acid sequence is expressed in the plant cell.

70. A dough, bread or baked products and/or dough, bread or baked product precursors comprising a polypeptide having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity, wherein the polypeptide comprises a the amino acid sequence of any of claims 21 to 30, or the polypeptide is encoded by a nucleic acid comprising the sequence of any of claims 1 to 6, or an enzymatically active fragment thereof, or the enzyme mixture or cocktail of claim 88.

71. A beverage or beverage precursor comprising a polypeptide having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity, wherein the polypeptide comprises a the amino acid sequence of any of claims 21 to 30, or the polypeptide is encoded by a nucleic acid comprising the sequence of any of claims 1 to 6, or an enzymatically active fragment thereof, or the enzyme mixture or cocktail of claim 88.

72. A method of beverage production comprising administration of at least one polypeptide having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity, wherein the polypeptide comprises a the amino acid sequence of any of claims 21 to 30, or the polypeptide is encoded by a nucleic acid comprising the sequence of any of claims 1 to 6, or an enzymatically active fragment thereof, or the enzyme mixture or cocktail of claim 88, to a beverage or a beverage precursor,

wherein optionally the beverage or beverage precursor is a wort or a beer.

73. A feed, food, food or feed additive, food or feed supplement, or dietary aid comprising a polypeptide having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity, wherein the polypeptide comprises a the amino acid sequence of any of claims 21 to 30, or
5 the polypeptide is encoded by a nucleic acid comprising the sequence of any of claims 1 to 6, or an enzymatically active fragment thereof, or the enzyme mixture or cocktail of claim 88,

wherein optionally, wherein optionally the feed, food, food or feed additive, food or feed supplement, or dietary aid further comprises a carrier selected from the group
10 consisting of a grain germ, a grain germ that is spent of oil, a hay, an alfalfa, a timothy, a soy hull, a sunflower seed meal and a wheat midd,

and optionally the carrier comprises grain germ that is spent of oil, or optionally the isomerase, e.g., the racemase, e.g., the amino acid racemase, the alanine racemase, and/or the epimerase enzyme is glycosylated to provide thermostability at pelletizing
15 conditions, and optionally the delivery matrix is formed by pelletizing a mixture comprising a grain germ and an isomerase, e.g., a racemase, e.g., an amino acid racemase, an alanine racemase, and/or an epimerase, and optionally the pelletizing conditions include application of steam, and optionally the pelletizing conditions comprise application of a temperature in excess of about 80°C for about 5 minutes and the enzyme retains a specific
20 activity of at least 350 to about 900 units per milligram of enzyme.

74. A granule, pellet or particle comprising a polypeptide having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity, wherein the polypeptide comprises a the
25 amino acid sequence of any of claims 21 to 30, or the polypeptide is encoded by a nucleic acid comprising the sequence of any of claims 1 to 6, or an enzymatically active fragment thereof, or the enzyme mixture or cocktail of claim 88,

wherein optionally the isomerase, e.g., the racemase, e.g., the amino acid racemase, the alanine racemase, and/or the epimerase is thermostable.
30

75. A cleaning composition comprising a polypeptide having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity, wherein the polypeptide comprises a the amino acid sequence of any of claims 21 to 30, or the polypeptide is encoded by a nucleic

acid comprising the sequence of any of claims 1 to 6, or an enzymatically active fragment thereof, or the enzyme mixture or cocktail of claim 88,

wherein optionally, the cleaning composition is a detergent, disinfectant or cleanser, or wherein optionally, the cleaning composition is for cleaning fabrics,
5 dishwashing, laundry, oral cleaning, denture cleaning, or contact lenses.

76. A pharmaceutical composition comprising a polypeptide having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity, wherein the polypeptide comprises
10 a the amino acid sequence of any of claims 21 to 30, or the polypeptide is encoded by a nucleic acid comprising the sequence of any of claims 1 to 6, or an enzymatically active fragment thereof, or the enzyme mixture or cocktail of claim 88.

77. A method for treating a wood, a wood product, a paper, a paper product, a
15 pulp, a pulp product, a paper waste or a paper recycling composition, comprising contacting the wood, wood product, paper, paper product, pulp, pulp product, paper waste or paper recycling composition with a polypeptide having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity,
20 and/or an epimerase activity, wherein the polypeptide comprises a the amino acid sequence of any of claims 21 to 30, or the polypeptide is encoded by a nucleic acid comprising the sequence of any of claims 1 to 6, or an enzymatically active fragment thereof, or the enzyme mixture or cocktail of claim 88,

wherein optionally, the treatment comprises reducing or solubilizing lignin (delignification), bleaching or decoloring, and/or deinking.
25

78. A biomass, wood, wood pulp, wood product, paper pulp, paper product, newspaper or paper waste comprising a polypeptide having a the amino acid sequence of any of claims 21 to 30, or a polypeptide encoded by a nucleic acid comprising the sequence of any of claims 1 to 6, or an enzymatically active fragment thereof, or the
30 enzyme mixture or cocktail of claim 88.

79. A fabric, yarn, cloth or textile comprising a polypeptide having a the amino acid sequence of any of claims 21 to 30, or a polypeptide encoded by a nucleic acid comprising the sequence of any of claims 1 to 6, or an enzymatically active fragment thereof, or the enzyme mixture or cocktail of claim 88,

5 wherein optionally the fabric, yarn, cloth or textile comprises a non-cotton cellulosic fabric, yarn, cloth or textile.

80. A method for treating an organic material comprising:

10 (a) providing at least one polypeptide having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity having the amino acid sequence of any of claims 21 to 30, or the isomerase, e.g., the racemase, e.g., the amino acid racemase, the alanine racemase, and/or the epimerase is encoded by a nucleic acid comprising the sequence of any of claims 1 to 6, or an enzymatically active fragment thereof, or the enzyme mixture or
15 cocktail of claim 88;

(b) providing an organic material; and

(c) contacting the organic material of (b) with the polypeptide of step (a),

wherein optionally the organic material is a biomass, wood, a wood pulp, a Kraft pulp, a paper, a paper product or a paper pulp,

20 and wherein optionally, the biomass, wood, wood pulp, Kraft pulp, paper, paper product or paper pulp comprises a softwood and hardwood, or the wood, wood pulp, Kraft pulp, paper or paper pulp is derived from a softwood and hardwood.

81. A composition comprising an organic material comprising a polypeptide
25 having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity, wherein the polypeptide has the amino acid sequence of any of claims 21 to 30, or the polypeptide is encoded by a nucleic acid comprising the sequence of any of claims 1 to 6, or an enzymatically active fragment thereof, or the enzyme mixture or cocktail of claim 88,

30 wherein optionally the organic material is a biomass, wood, a wood pulp, a Kraft pulp, a paper, a paper product or a paper pulp,

and wherein optionally, the biomass, wood, wood pulp, Kraft pulp, paper, paper product or paper pulp comprises a softwood and hardwood, or the wood, wood pulp, Kraft pulp, paper or paper pulp is derived from a softwood and hardwood.

82. A method for making an alcohol comprising contacting an organic material with a polypeptide having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity, wherein
5 the polypeptide has the amino acid sequence of any of claims 21 to 30, or the polypeptide is encoded by a nucleic acid comprising the sequence of any of claims 1 to 6, or an enzymatically active fragment thereof, or the enzyme mixture or cocktail of claim 88,

wherein optionally the organic material is a biomass, wood, a wood pulp, a Kraft pulp, a paper, a paper product or a paper pulp,

10 and wherein optionally, the biomass, wood, wood pulp, Kraft pulp, paper, paper product or paper pulp comprises a softwood and hardwood, or the wood, wood pulp, Kraft pulp, paper or paper pulp is derived from a softwood and hardwood,

and optionally the method further comprises fermentation,

and optionally the alcohol is or comprises an ethanol.

15 83. A composition comprising an alcohol and a polypeptide having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity, wherein the polypeptide has the amino acid sequence of any of claims 21 to 30, or the polypeptide is encoded by a nucleic
20 acid comprising the sequence of any of claims 1 to 6, or an enzymatically active fragment thereof, or the enzyme mixture or cocktail of claim 88,

and optionally the alcohol is or comprises an ethanol.

25 84. A waste treatment solution comprising a polypeptide having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity, wherein the polypeptide has the amino acid sequence of any of claims 21 to 30, or the polypeptide is encoded by a nucleic acid comprising the sequence of any of claims 1 to 6, or an enzymatically active fragment
30 thereof, or the enzyme mixture or cocktail of claim 88.

85. A chewing gum, lozenge or candy comprising a polypeptide having an isomerase activity, e.g., a racemase activity, e.g., an amino acid racemase activity, an alanine racemase activity, and/or an epimerase activity, wherein the polypeptide has the amino acid sequence of any of claims 21 to 30, or the polypeptide is encoded by a nucleic

acid comprising the sequence of any of claims 1 to 6, or an enzymatically active fragment thereof, or the enzyme mixture or cocktail of claim 88.

86. A chimeric isomerase, e.g., racemase, e.g., amino acid racemase, alanine
5 racemase, and/or epimerase comprising (a) a the amino acid sequence of any of claims 21
to 30 and at least one heterologous domain; or (b) the chimeric isomerase, e.g., racemase,
e.g., amino acid racemase, alanine racemase, and/or epimerase of (a), wherein optionally
the domain comprises a binding domain or a dockerin domain.

10 87. A method for designing a chimeric glycosidase, isomerase, e.g., racemase,
e.g., amino acid racemase, alanine racemase, and/or epimerase having a new specificity or
an enhanced specificity, comprising inserting a heterologous or an additional endogenous
domain into a glycosidase, wherein the domain comprises a binding domain or a dockerin
domain.

15 88. An enzyme mixture or cocktail comprising (a) at least one enzyme of any
of any of claims 21 to 30, and one or more other enzyme(s); (b) the mixture or cocktail of
(a), wherein the one or more other enzyme(s) is another isomerase, e.g., racemase, e.g.,
amino acid racemase, alanine racemase, and/or epimerase, a mannanase and/or a
20 glucanase, cellulases, lipases, esterases, proteases, or endoglycosidases, endo-beta.-1,4-
glucanases, beta-glucanases, endo-beta-1,3(4)-glucanases, cutinases, peroxidases,
catalases, laccases, amylases, glucoamylases, pectinases, transferases, transaminases,
amino transferases, dehydrogenases, oxidoreductases, reductases, oxidases,
phenoloxidases, ligninases, pullulanases, arabinanases, hemicellulases, mannanases,
25 xyloglucanases, xylanase, a mannanase and/or a glucanases, pectin acetyl esterases,
rhamnogalacturonan acetyl esterases, polygalacturonases, rhamnogalacturonases,
galactanases, pectin lyases, pectin methylesterases, cellobiohydrolases and/or
transglutaminases.

30 89. A liquid composition comprising (a) an alcohol and a polypeptide of any of
any of claims 21 to 30, or the enzyme mixture or cocktail of claim 88; (b) the liquid
composition of (a), wherein the alcohol is or comprises ethanol, propanol, butanol and/or
methanol; (c) the liquid composition of (a) or (b) comprising or contained in a fuels, a
biofuel, a synthetic liquid or gas or a syngas.

DocCode – SCORE

SCORE Placeholder Sheet for IFW Content

Application Number: PCT/US08/88066 Document Date: 12/22/2008

The presence of this form in the IFW record indicates that the following document type was received in electronic format on the date identified above. This content is stored in the SCORE database.

- Design Drawing

Since this was an electronic submission, there is no physical artifact folder, no artifact folder is recorded in PALM, and no paper documents or physical media exist. The TIFF images in the IFW record were created from the original documents that are stored in SCORE.

To access the documents in the SCORE database, refer to instructions developed by SIRA.

At the time of document entry (noted above):

- Examiners may access SCORE content via the eDAN interface.
- Other USPTO employees can bookmark the current SCORE URL (<http://es/ScoreAccessWeb/>).
- External customers may access SCORE content via the Public and Private PAIR interfaces.

Form Revision Date: February 8, 2006

Figure 1




Figure 2




FIGURE 3




Figure 4

