




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

| | | |
|---|---|---|
| (51) International Patent Classification⁵ : A61K 37/547 // (A61K 37/547 A61K 37:02) | A1 | (11) International Publication Number: WO 93/23074 (43) International Publication Date: 25 November 1993 (25.11.93) |
| (21) International Application Number: PCT/US93/04493 (22) International Filing Date: 12 May 1993 (12.05.93) (30) Priority data: 07/882,202 13 May 1992 (13.05.92) US 08/021,615 19 February 1993 (19.02.93) US (60) Parent Application or Grant (63) Related by Continuation US 021,615 (CON) Filed on 19 February 1993 (19.02.93) (71) Applicant (for all designated States except US): OKLAHOMA MEDICAL RESEARCH FOUNDATION [US/US]; 825 Northeast 13th Street, Oklahoma City, OK 73104 (US). | (72) Inventors; and (75) Inventors/Applicants (for US only): MORRISSEY, James, H. [US/US]; 228 W. Eubanks, Oklahoma City, OK 73118 (US). COMP, Philip, C. [US/US]; 9805 Silver Lake Drive, Oklahoma City, OK 73118 (US). (74) Agents: HANSEN, Eugenia, S. et al.; Richards, Medlock & Andrews, 4500 Renaissance Tower, 1201 Elm Street, Dallas, TX 75270-2197 (US). (81) Designated States: AU, CA, JP, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i> | |

(54) Title: TRUNCATED TISSUE FACTOR AND FVIIa OR FVII ACTIVATOR FOR BLOOD COAGULATION

**(57) Abstract**

It has been discovered that it is possible to administer truncated tissue factor, not having the transmembrane region, (tTF) in combination with factor VIIa (FVIIa) or an activator of endogenous factor VII to treat bleeding disorders such as those resulting from hemophilia or cirrhosis of the liver. The tTF is administered to produce up to 10 µg tTF/ml of plasma. The FVIIa or FVII activator is administered to produce levels of between 40 ng VIIa/ml and 700 ng FVIIa/ml of plasma. The effective dosages of both tTF and VIIa/factor VII activator are significantly and surprisingly less than the administration of either alone to stop bleeding. Examples demonstrate safety and efficacy in normal and hemophilic dogs.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

| | | | | | |
|----|--------------------------|----|--|----|--------------------------|
| AT | Austria | FR | France | MR | Mauritania |
| AU | Australia | GA | Gabon | MW | Malawi |
| BB | Barbados | GB | United Kingdom | NL | Netherlands |
| BE | Belgium | GN | Guinea | NO | Norway |
| BF | Burkina Faso | GR | Greece | NZ | New Zealand |
| BG | Bulgaria | HU | Hungary | PL | Poland |
| BJ | Benin | IE | Ireland | PT | Portugal |
| BR | Brazil | IT | Italy | RO | Romania |
| CA | Canada | JP | Japan | RU | Russian Federation |
| CF | Central African Republic | KP | Democratic People's Republic of Korea | SD | Sudan |
| CG | Congo | KR | Republic of Korea | SE | Sweden |
| CH | Switzerland | KZ | Kazakhstan | SK | Slovak Republic |
| CI | Côte d'Ivoire | LI | Liechtenstein | SN | Senegal |
| CM | Cameroon | LK | Sri Lanka | SU | Soviet Union |
| CS | Czechoslovakia | LU | Luxembourg | TD | Chad |
| CZ | Czech Republic | MC | Monaco | TG | Togo |
| DE | Germany | MG | Madagascar | UA | Ukraine |
| DK | Denmark | ML | Mali | US | United States of America |
| ES | Spain | MN | Mongolia | VN | Viet Nam |
| FI | Finland | | | | |

**TRUNCATED TISSUE FACTOR AND FVIIa OR FVII ACTIVATOR
FOR BLOOD COAGULATION**

Technical Field

The present invention relates to a composition and method for treatment of bleeding.

5 **Background of the Invention**

The United States government has certain rights in this invention by virtue of grant R01 HL 44225 awarded by the National Institutes of Health to James H. Morrissey.

10 This is a continuation of U.S. Serial No. 08/021615
filed February 19, 1993, entitled, "Treatment of Bleeding
with Modified Tissue Factor In Combination with An
Activator of FVII" which is a continuation-in-part of U.S.
Serial No. 07/882,202 entitled "Treatment of Bleeding with
Modified Tissue Factor in Combination with FVIIa" filed
15 May 13, 1992, both by James H. Morrissey and Philip C.
Comp.

Blood coagulation results from the production of thrombin, a proteolytic enzyme inducing platelet
20 aggregation and cleaving fibrinogen to fibrin, which
stabilizes the platelet plug. A number of proenzymes and
procofactors circulating in the blood interact in this
process through several stages during which they are
sequentially or simultaneously converted to the activated
form, ultimately resulting in the activation of
25 prothrombin to thrombin by activated factor X (fXa) in the
presence of factor Va, ionic calcium, and platelets.

Factor X can be activated by either of two pathways,
termed the extrinsic and intrinsic pathways. The
intrinsic pathway, or surface-mediated activation pathway,
30 consists of a series of reactions where a protein
precursor is cleaved to form an active protease, beginning

with activation of factor XII to factor XIIa, which converts factor XI to factor XIa, which, in the presence of calcium, converts factor IX to factor IXa. Factors IX and X can also be activated via the extrinsic pathway by
5 tissue factor (TF) in combination with activated factor VII (factor VIIa; fVIIa). Activated factor IX, in the presence of calcium, phospholipid (platelets), and factor VIIIa, activates factor X to factor Xa.

Physiologically, the major pathway involved in
10 coagulation is believed to be the extrinsic pathway, an essential step of which is tissue factor-mediated activation of factor VII to factor VIIa. Tissue factor is an integral membrane glycoprotein having a protein and a phospholipid component. It has been isolated from a
15 variety of tissues and species and reported to have a molecular mass of between 42,000 and 53,000. DNA encoding tissue factor and methods for expression of the protein have now been reported, for example, in European Patent Application 0 278 776 by Genentech, Inc. and by J. H.
20 Morrissey, et al. Cell 50, 129-135 (1987).

The complex of factor VIIa and its essential cofactor, TF, is the most potent known trigger of the clotting cascade. Factor VII is present in plasma at a concentration of 0.5 $\mu\text{g/ml}$ plasma. In contrast, factor
25 VIIa is present in plasma at trace levels of roughly 1 ng/ml. Accordingly, factor VII is normally in considerable excess over factor VIIa. Factor VIIa circulates with a relatively long half-life of about two hours in plasma. This is an unusual property among
30 activated coagulation enzymes, which typically are inactivated very rapidly by various protease inhibitors in plasma.

Hemophilia A is characterized by the absence of active coagulation factor VIII or the presence of

inhibitors to factor VIII. Hemophilia B is characterized by the absence of active factor IX. Both types of hemophilia are associated with bleeding diatheses that can lead to crippling injuries or death. Traditionally, patients with either type of hemophilia were treated with infusion of plasma concentrates to stop bleeding episodes. The problems with the concentrates are many, especially infection, most recently with HIV. Highly purified factor VIII or factor IX have been introduced to obviate these problems, as have methods of treating factor concentrates to inactivate viruses. However, some patients develop high-titer, inhibitory antibodies to factor VIII. Therefore, such patients can no longer be treated with conventional factor VIII replacement therapy.

As described by Hedner and Kisiel, J. Clin. Invest. 71, 1836-1841 (1983), purified naturally produced factor VIIa can be administered to hemophilia A patients with high titers of antibodies against factor VIII:C and restore hemostasis in these patients. As reported by Brinkhous, et al., Proc. Natl. Acad. Sci. USA 86, 1382-1386 (1989), recombinant factor VIIa (rFVIIa) can be administered to hemophilic and von Willebrand disease dogs and stop bleeding in both hemophilic A and B dogs, but not the von Willebrand disease dogs. Telgt, et al., Thrombosis Res. 56, 603-609 (1989), reported that, at high levels, rFVIIa was believed to act by direct activation of factor X, in the presence of calcium and phospholipid but in the absence of TF. Teitel, Thrombosis and Haemostasis 66(5), 559-564 (1991), reported that the important ingredient in prothrombin complex concentrates for efficacy in treating hemophilia is factor VIIa.

Hedner, "Experiences with Recombinant Factor VIIa in Haemophiliacs" in Biotechnology of Plasma Proteins Lenfant, Mannucci, Sixma, eds., Curr. Stud. Hematol. Blood

Transf. No. 58, 63-68 (1991), review the use of prothrombin complex concentrates (effective in only 50 to 60% of the bleeds), as well as the use of plasma-derived (pFVIIa) and recombinant factor VIIa (rFVIIa). Dosages of 10 to 15 $\mu\text{g}/\text{kg}$ of pFVIIa were effective in some hemophilia A patients. Safety studies in dogs and rabbits indicated that the recombinant factor VIIa was safe and efficacious at dosages of up to 150 $\mu\text{g}/\text{kg}$. A number of patients were also successfully treated, using dosages of between 54 $\mu\text{g}/\text{kg}$ and 90 $\mu\text{g}/\text{kg}$ during surgery and bleeding complications. Gringeri, et al., reported that treatment of hemophiliacs with rFVIIa is not always effective, even at dosages of 75 $\mu\text{g}/\text{kg}$ at intervals of every two to three hours. The authors noted that perhaps larger dosages, more frequent infusions, and/or the concomitant use of antifibrinolytic medication might be required in such cases. rFVIIa is currently in clinical trials in the United States for treatment of hemophilia, particularly hemophilia in patients with inhibitors who do not benefit from conventional factor VIII or factor IX replacement therapy. Doses of rFVIIa currently employed are typically 45 to 90 μg rFVIIa/kg body weight, and are repeated every two to four hours. These doses are designed to achieve a level of circulating rFVIIa of approximately 4 $\mu\text{g}/\text{ml}$, extremely high compared to the normal plasma concentrations of FVII of approximately 0.5 $\mu\text{g}/\text{ml}$ or FVIIa of approximately 1 ng/ml.

O'Brien, et al., J. Clin. Invest. 82, 206-211 (1988), reported that apo-TF, a delipidated preparation of the normally lipid-associated TF glycoprotein could be used to normalize bleeding in animals having antibodies to factor VIII. Since purified apo-TF is inactive unless incorporated into a phospholipid membrane, the theoretical basis for infusing apo-TF is the hypothesis that it would

spontaneously and preferentially incorporate into exposed membrane surfaces, particularly into damaged cells at the sites of injury. Subsequent studies have indicated there are dangers associated with the use of purified apo-TF in the treatment of hemophilia. The apo-TF can spontaneously incorporate into many types of lipid membranes and become active at sites where clotting is not desired, resulting in thrombosis or disseminated intravascularization (DIC). Indeed, O'Brien, et al., reported evidence of DIC in some animals receiving apo-TF, and Sakai and Kisiel, Thromb. Res. 60, 213-222 (1990), recently demonstrated that apo-TF will spontaneously combine with plasma lipoproteins to form active TF. This latter phenomenon is cause for concern because of a number of studies which have demonstrated that intravenous administration of active TF is a potent inducer of DIC.

Recently, a soluble, truncated form of TF (tTF) has been reported which retains some cofactor function towards factor VIIa as measured *in vitro* using purified proteins. However, this form of TF has been dismissed as an alternative to TF because it has almost no procoagulant activity when tested with normal plasma, as reported by Paborsky, et al., J. Biol. Chem. 266:21911-21916 (1991).

As described in U.S. Serial No. 07/683,682 entitled "Quantitative Clotting Assay for Activated Factor VII" filed April 10, 1991 by James H. Morrissey, the reason tTF was reported to lack procoagulant activity in the prior art is because, although tTF retains cofactor function toward factor VIIa, tTF had lost the ability to promote conversion of factor VII to factor VIIa. As a consequence, tTF can clot plasma only in conjunction with significantly elevated levels of factor VIIa, as compared to normal plasma, which contains only trace levels of factor VIIa.

It is therefore an object of the present invention to provide a method and composition for treatment of significant bleeding disorders, such as hemophilia, including those hemophiliacs with high titers of anti-factor VIII antibodies.

It is a further object of the present invention to provide a method and compositions for treatment of patients with bleeding disorders that are relatively safe and can be produced in commercial quantities.

10

Summary of the Invention

It has been discovered that it is possible to administer truncated tissue factor, the isolated extracellular domain of tissue factor, (tTF) in combination with factor VIIa (FVIIa), or truncated tissue factor in combination with an activator of factor VIIa such as the combination of factor Xa/phospholipid, factor IXa/phospholipid, thrombin, factor XIIa, or the factor VII activator from the venom of *Oxyuranus scutellatus*/phospholipid, to treat bleeding disorders such as, for example, those associated with hemophilia or cirrhosis of the liver. The minimal effective dosages of both tTF and FVIIa are significantly and surprisingly less than the administration of either alone to stop bleeding.

Examples demonstrate safety and efficacy in normal and hemophilic dogs.

Brief Description of the Drawings

Figure 1a is a graph of dilute thromboplastin clotting time (seconds) for factor VIII-deficient plasma when tTF and FVIIa (ng/ml) are added (dark triangles) or when FVIIa alone (ng/ml) are added (dark circles).

Figure 1b is a graph of dilute thromboplastin clotting time (seconds) for normal plasma when tTF and

FVIIa (ng/ml) are added (dark triangles) or when FVIIa alone (ng/ml) are added (dark circles).

Detailed Description of the Invention

5 Truncated tissue factor (tTF) is used as an adjuvant for factor VIIa (FVIIa) therapy of excessive bleeding in patients such as hemophiliacs. The tTF is administered in combination with the FVIIa, or a composition which
10 activates endogenous FVII to form FVIIa, such as the combination of FXa and phospholipid, or immediately before or after the FVIIa. As used herein, "FVIIa" refers to administration of either FVIIa or an activator of FVII to FVIIa. The minimal effective dosage of FVIIa is
15 significantly decreased by the tTF, while the thrombogenicity of the mixture is reduced as compared to a mixture of VIIa and TF in its native form.

The advantage, and necessity, of using a combination of FVIIa plus tTF over tTF alone is that tTF is active in promoting blood clotting only in the presence of pre-
20 formed FVIIa. The advantage of using the combination of FVIIa plus tTF over FVIIa alone is that FVIIa is approximately 22-fold more potent at clotting plasma in the presence of tTF than in the absence of tTF. Furthermore, as with FVIIa alone, the complex of tTF plus
25 FVIIa activates factor X at markedly enhanced rates in the presence of negatively charged phospholipids such as phosphatidyl serine, even though tTF is not itself incorporated into lipid membranes. Normally, negatively
30 charged phospholipids are sequestered into the inner leaflet of the plasma membrane of cells, and thus are not exposed to the cell surface unless the cells are damaged, or, in the case of platelets, the platelets have undergone surface activation. Thus, negatively charged phospholipids should be preferentially exposed on cell

surfaces at anatomic sites that have experienced trauma sufficient to cause cell lysis and/or platelet activation. This means that the complex of tTF and FVIIa is predicted from *in vitro* studies to be active in promoting blood clotting preferentially at sites of injury, where it would be needed to confer hemostasis. Furthermore, tTF, by itself or in complex with FVIIa, remains a soluble entity and does not become incorporated into lipid membranes. This means that it should clear relatively rapidly from the circulation, diminishing the risk of thrombosis. Because of these properties, the complex of tTF and FVIIa should not cause systemic activation of the blood clotting system leading to DIC, as can injection of wild-type TF. These *in vitro* results are consistent with results from *in vivo* studies in normal and hemophiliac dogs described below.

The combination of tTF plus FVIIa should be useful for treatment of hemophilia A, hemophilia B, congenital or acquired deficiencies in any other blood coagulation factor, or platelet defects. Other patients that can be treated with the combination include patients suffering from severe trauma, postoperative bleeding or those with cirrhosis.

Truncated Tissue Factor

As used herein, "truncated tissue factor" is a soluble tissue factor having only the extracellular domains, which is not bound to a phospholipid membrane surface, and therefore does not support conversion of fVII to fVIIa, as described in U.S. Serial No. 07/683,682 filed April 10, 1991, corresponding to International Application PCT/US92/02898 filed April 9, 1992, the teachings of which are incorporated herein. In the preferred embodiment, truncated tissue factor is a recombinant protein produced *in vitro* in cell culture using a mammalian cell line such

as CHO-K1 cells, American Type Culture Collection CCL 61. These cells are stably transfected with a mutant form of the human tissue factor cDNA carried in a commercially available expression vector, and secrete a form of tissue factor consisting only of amino acids 1-219 (numbered according to Morrissey, et al., Cell 50:129-135 (1987), the teachings of which are incorporated herein).

The recombinant, truncated tissue factor is purified from the culture medium using an immobilized monoclonal antibody to human tissue factor, such as TF9-5B7, described in Morrissey, et al. Thromb. Res. 52:247-261 (1988). Hybridomas for production of monoclonal antibodies can be propagated by ascites growth and the monoclonal antibodies (MAbs) purified from ascites fluid using the BioRad MAPS II system for mAb purification, as described by Morrissey, et al., Thromb. Res. 52:247-261 (1988).

The TF9-5B7 is coupled to AffigelTM beads. Detergent is not used during the purification of TF₂₁₉. After removal of cellular debris by centrifugation, the culture medium is made 25 mM in Tris.HCl (pH 7.4), 10 mM in sodium EDTA (pH 7.4), and 0.1% in sodium azide by the addition of concentrated stock solutions. In order to remove proteins that bind to agarose beads non-specifically, the culture medium is gently agitated for 4 hr at 4°C with AffiGelTM-10 beads that had previously been blocked chemically with glycine ethyl ester (GEE-AffiGel). The GEE-AffiGel beads are removed by filtration through a sintered glass funnel, and the supernatant is agitated overnight at 4°C with the MAb TF9-5B7 coupled to AffiGel beads (typically 2 ml of beads). The TF9-5B7-AffiGel beads are collected on a sintered glass funnel, and the beads are washed on the funnel with 100 ml of TBS-EDTA (TBS = 100 mM NaCl, 50 mM Tris.HCl pH 7.4, 0.02% sodium azide; TBS-EDTA = TBS with

10 mM EDTA included). The beads are then transferred to a chromatography column and washed with 40 ml TBS followed by 40 ml of a solution consisting of 1 M NaCl, 10 mM Tris.HCl pH 7.4, 0.02% sodium azide. Truncated TF is
5 eluted from the beads using 100 mM glycine.HCl pH 2.5, with 1 ml fractions being collected into tubes containing 57 μ l 1 M Tris base (to immediately neutralize the acidic glycine buffer). Fractions containing protein are detected using the BCA protein assay (Pierce), pooled,
10 dialyzed against TBS, and then stored at -70°C . Protein concentrations are determined using standard methods such as the BCA assay (Pierce Chemical Co.) based on a bovine serum albumin standard of known concentration.

For production of recombinant truncated TF, cells are
15 typically grown to confluence in roller bottles in Dulbecco's Modified Eagle's Medium (DMEM) supplemented with 10% iron-supplemented calf serum (HyClone Laboratories, Logan, Utah). Upon reaching confluence, the cultures are shifted to DMEM containing 2% serum, and
20 culture medium is collected every four days thereafter.

The nucleotide and amino acid sequence of truncated tissue factor (tTF) is shown below as SEQ. ID. NO: 1 and 2, respectively. The truncated tissue factor protein lacks the predicted transmembrane and cytoplasmic domains
25 of tissue factor. This version of the protein retains cofactor activity, as reported by Ruf, et al., Thromb. Haemost. 62, 347 (abstract) (1989) and Ruf, et al., J. Biol. Chem. 266: 2158-2166 (1991). This truncated form of tissue factor fails to support conversion of factor VII to
30 VIIa, allowing it to be used in a specific clotting assay for factor VIIa free from interference by factor VII.

Since the vast majority of FVII in plasma is in the inactive, zymogen form, and since tTF is selectively deficient in promoting conversion of factor VII to VIIa,

tTF has extremely little procoagulant activity towards normal plasma. Since tTF lacks the membrane-anchoring domain of the protein, it lacks the ability to become incorporated into exposed phospholipid surfaces as wild-type apo-TF can. Although the deficiency of tTF in the conversion of FVII to FVIIa was not previously appreciated, its extremely low procoagulant activity when tested with normal plasma has made it appear to be unacceptable as a therapeutic agent to control bleeding in hemophilia.

The essential difference between truncated tissue factor and wild-type tissue factor is that truncated tissue factor is no longer tethered to the phospholipid membrane surface. It is therefore expected that other methods for preparing truncated tissue factor can be used to generate an essentially equivalent soluble form of tissue factor that retains FVIIa cofactor activity while no longer stimulating conversion of factor VII to factor VIIa. Methods include chemical and/or enzymatic cleavage of wild-type tissue factor to separate the predicted extracellular domain from the transmembrane region. Recombinant human TF is available from Calbiochem Corporation. Precise positioning of the stop codon following amino acid 219 is believed to not be essential to make functional truncated TF, and other placements of a stop codon near amino acid 219 are predicted to yield an essentially equivalent product with respect to its ability in conjugation with FVIIa to serve as a treatment for bleeding disorders.

Activated Factor VII

Factor VII can be prepared as described by Fair, Blood 62, 784-791 (1983). The coding portion of the human factor VII cDNA sequence reported by Hagen et al., Proc. Natl. Acad. Sci. USA 83:2412-2416 (1986) is shown below as

SEQ. ID. NO: 3, along with the translated amino acid sequence, SEQ. ID. NO: 4. The amino acid sequence from 1 to 60 corresponds to the pre-pro/leader sequence that is removed by the cell prior to secretion. The mature FVII
5 polypeptide chain consists of amino acids 61 to 466. FVII is converted to FVIIa by cleavage of a single peptide bond between arginine-212 and isoleucine-213.

FVII can be converted *in vitro* to fVIIa by incubation of the purified protein with factor Xa immobilized on
10 Affi-GelTM 15 beads (Bio-Rad). Conversion can be monitored by SDS-polyacrylamide gel electrophoresis of reduced samples. Free factor Xa in the fVIIa preparation can be detected with the chromogenic substrate
15 methoxycarbonyl-D-cyclohexylglycyl-glycyl-arginine-p-nitroanilide acetate (SpectrozymeTM FXa, American Diagnostica, Greenwich, CT) at 0.2 mM final concentration in the presence of 50 mM EDTA.

Recombinant fVIIa can also be purchased from Novo Biolabs (Danbury, CT).

20 **Formation of FVIIa *in vivo***

Alternatively, the FVIIa can be formed *in vivo*, at the time of, or shortly before, administration of the truncated tissue factor. In a preferred embodiment, endogenous FVII is converted into FVIIa by infusion of an
25 activator of VIIa, such as factor Xa (FXa) in combination with phospholipid (PCPS).

Activators of factor VII *in vivo* include FXa/PCPS, FIXa/PCPS, thrombin, FXIIa, and the FVII activator from the venom of *Oxyuranus scutellatus* in combination with
30 PCPS. These have been shown to activate FVII to FVIIa *in vitro*, although never in combination with truncated tissue factor:

FXa and Thrombin: Radcliffe, R. and Nemerson, Y. (1975). Activation and control of factor VII by activated

factor X and thrombin: Isolation and characterization of a single chain form of factor VII. J. Biol. Chem. 250:388-395. This paper points out that activation of FVII by FXa is accelerated when phospholipids are present.

5 FIXa and FXa: Masys, D.R., Bjaaj, S.P. and Rapaport, S.I. (1982). Activation of human factor VII by activated factors IX and X. Blood 60:1143-1150. This paper describes activation of factor VII by both factors IXa and Xa, and the fact that phospholipids accelerate both
10 reactions.

 FXa plus phospholipids in vivo: Giles, A.R., Mann, K.G. and Nesheim, M.E. (1988). A combination of factor Xa and phosphatidylcholine-phosphatidylserine vesicles bypasses factor VIII in vivo. Br. J. Haematol. 69:491-
15 497. This paper describes the factor VIII bypassing activity of FXa/PCPS in hemophilic dogs. It does not deal with activation of factor VII.

 FXIIa: Kisiel, W., Fujikawa, K. and Davie, E.W. (1977). Activation of bovine factor VII (proconvertin) by
20 factor XIIa (activated Hageman factor). Biochemistry 16:4189-4194. Describes activation of bovine FVII by FXIIa (no phospholipids).

 Snake venom activator of FVII: Nakagaki, T., Lin, P. and Kisiel, W. (1992). Activation of human factor VII by
25 the prothrombin activator from the venom of Oxyuranus scutellatus (Taipan snake). Thromb. Res. 65:105-116. This paper describes the isolation of an enzyme from this snake venom which, in combination with phospholipids, will activate factor VII to VIIa in vitro.

30 Activation of FVII to FVIIa for Xa/PCPS in vivo has also been measured directly.

 In general, the factor VII activator is administered in a dosage between 1 and 10 $\mu\text{g/ml}$ of carrier.

The phospholipid can be provided in a number of forms but the preferred form is as phosphatidyl choline/phosphatidyl serine vesicles (PCPS). The PCPS vesicle preparations and the method of administration of Xa/PCPS is described in Giles, et al., (1988), the teachings of which are specifically incorporated herein. Other phospholipid preparations can be substituted for PCPS, so long as they accelerate the activation of FVII by FXa. Effectiveness, and therefore determination of optimal composition and dose, can be monitored as described below.

A highly effective dose of Xa/PCPS, which elevates FVIIa levels in vivo in the chimpanzee, was 26 pmoles FXa + 40 pmoles PCPS per kg body weight. That dose yielded an eighteen fold increase in endogenous levels of FVIIa (to 146 ng/ml). A marginally detectable effect was observed using a smaller dose in dogs, where the infusion of 12 pmoles FXa + 19 pmoles PCPS per kg body weight yielded a three fold increase in endogenous FVIIa levels. Accordingly, doses of FXa that are at least 12 pmoles FXa per kg body weight, and preferably 26 pmoles FXa per kg body weight, should be useful. Doses of PCPS that are at least 19 pmoles PCPS per kg body weight, and preferably 40 pmoles PCPS per kg body weight, are similarly useful.

The effectiveness of any infusible FVII activator can be monitored, following intravenous administration, by drawing citrated blood samples at varying times (at 2, 5, 10, 20, 30, 60, 90 and 120 min) following a bolus infusion of the activator, and preparing platelet-poor plasma from the blood samples. The amount of endogenous FVIIa can then be measured in the citrated plasma samples by performing our tTF-based FVIIa clotting assay. Desired levels of endogenous FVIIa would be the same as the target levels of plasma FVIIa indicated for co-infusion of

purified FVIIa and tTF. Therefore, other activators of FVII could be tested in vivo for generation of FVIIa, without undue experimentation, and the dose adjusted to generate the desirable levels of FVIIa, using the tTF-
5 based FVIIa assay of plasma samples. The proper dose of the FVII activator (yielding the desired level of endogenous FVIIa) can then be used in combination with the recommended amounts of tTF.

Doses can be timed to provide prolong elevation in
10 FVIIa levels. Preferably doses would be administered every two hours until the desired hemostatic effect is achieved, and then repeated as needed to control bleeding. The half-life of FVIIa in vivo has been reported to be approximately two hours, although this could vary with
15 different therapeutic modalities and individual patients. Therefore, the half-life of FVIIa in the plasma in a given treatment modality should be determined with the tTF-based clotting assay.

20 Administration of the combination of the tTF and FVIIa

The tTF and VIIa are administered by infusion in the preferred embodiment, using a pharmaceutically acceptable carrier such as saline or buffered saline. The tTF and VIIa can also be administered topically either by direct
25 application using a conventional topical base such as petrolatum or a water based gel, or as an aerosol spray.

The tTF is administered in a dosage effective to produce in the plasma an effective level of between 100 ng/ml and 50 μ g/ml, or a preferred level of between 1
30 μ g/ml and 10 μ g/ml or 60 to 600 μ g/kg body weight, when administered systemically; or an effective level of between 10 μ g/ml and 50 μ g/ml, or a preferred level of between 10 μ g/ml and 50 μ g/ml, when administered topically.

The FVIIa is administered in a dosage effective to produce in the plasma an effective level of between 20 ng/ml and 10 μ g/ml (1.2 to 600 μ g/kg), or a preferred level of between 40 ng/ml and 4 μ g/ml (2.4 to 240 μ g/kg),
5 or a level of between 1 to 10 μ g FVIIa/ml when administered topically.

The FVIIa activator is administered in a dosage effective to produce in the plasma an effective level of FVIIa, as defined above. However, the maximum amount of
10 FVIIa converted from endogenous FVII is about 700 ng per ml.

In general, one would administer tTF and FVIIa to produce levels of up to 10 μ g tTF/ml plasma and between 40 ng and 4 μ g VIIa/ml plasma. For hemophilic patients, one
15 would administer tTF and FVIIa to produce levels of up to 10 μ g tTF and between 100 and 300 ng FVIIa/ml. For patients with cirrhosis, one would administer the same amount of tTF but up to 1 μ g FVIIa/ml plasma.

Since tTF cannot stimulate conversion of FVII to
20 FVIIa, it should only be functional in conjunction with elevated factor VIIa levels. Therefore, tTF is expected to be an effective hemostatic agent only in association with FVIIa therapy or in individuals who have elevated FVIIa levels for some other reason.

25 Other conditions that can be treated with this combination include surgical bleeding from the microvasculature, bleeding at skin graft harvest sites, postoperative bleeding, including following orthopedic surgery, brain surgery or brain trauma, bleeding secondary
30 to thrombocytopenia, and platelet dysfunction.

Example 1: Safety of tTF and FVIIa in normal dogs.

Three beagle dogs were treated with either tTF or tTF plus FVIIa.

Dog #5533 was treated with tTF alone, in a dosage of 60 μg tTF/kg body weight by intravenous bolus infusion. ELISA studies showed 1.23 μg tTF/ml in plasma at time zero. Half-life of tTF was 2.2 hr in this dog.

5 Dog #5534 was also treated with tTF alone, in a dosage of 41 μg tTF/kg body weight by intravenous bolus infusion. Assuming a blood volume of 60 ml/kg, this should have produced a level of 0.67 μg tTF/ml in the blood. ELISA studies showed 0.60 μg tTF/ml in plasma at
10 time zero. Half-life of tTF was 1.3 hr in this dog.

Dog #5734 was treated with tTF in combination with FVIIa, in a dosage of 41 μg tTF/kg body weight and 6.37 μg Novo recombinant factor VIIa/kg body weight by intravenous bolus infusion. ELISA studies showed 0.51 μg tTF/ml in
15 plasma at time zero. Half-life of tTF was biphasic (1 hr followed by 3.0 hr) in this dog.

The dogs all remained healthy and active. Blood tests of the clotting system indicated no significant decrease in fibrinogen levels or platelet counts and no
20 measurable increase in fibrin degradation products. Prothrombin times and APTT times were normal, as were white blood cell counts, and red blood cell counts. Therefore, administration of either tTF alone, or tTF in conjunction with FVIIa, caused no measurable disseminated
25 intravascular coagulation or other detectable coagulopathy.

A complete necropsy was performed on the animal receiving truncated tissue factor and factor VIIa. No evidence of thrombosis was found in veins, arteries or in
30 the capillaries. There was no evidence on gross and microscopic examination of myocardial infarction or of cerebral infarction (stroke).

Example 2: *In vitro* correction of clotting time of hemophilic plasma with tTF in combination with FVIIa.

A modified prothrombin test (PT) was performed with
5 diluted thromboplastin (since hemophiliacs have normal
PT's unless the thromboplastin is diluted) using either
normal or congenital factor VIII deficient (Hemophilia A)
plasma. Sigma Chemical Co. rabbit brain thromboplastin
was diluted 1:500 with TBS/0.1% BSA/rabbit brain cephalin
10 (Sigma Chemical Co.).

12 x 75 mm glass test tubes were pre-warmed in a 37°C
water bath.

Diluted thromboplastin was added (0.1 ml) and allowed
to warm to 37°C for more than two minutes.

15 Plasma sample (0.1 ml) was added and allowed to warm
to exactly 30 sec.

Pre-warmed 25 mM CaCl₂ (at 37°C) was added and the
clotting time was determined by the manual tilt-tube
method.

20 The results are shown in Figure 1a and b. Figure 1a
is a graph of dilute thromboplastin clotting time
(seconds) for factor VIII-deficient plasma when tTF (1
µg/ml) and varying concentrations of FVIIa (ng/ml) are
added (dark triangles) or when varying concentrations of
25 FVIIa alone (ng/ml) are added (dark circles). Figure 1b
is a graph of dilute thromboplastin clotting time
(seconds) for normal plasma when tTF (1 µg/ml) and varying
concentrations FVIIa (ng/ml) are added (dark triangles) or
when varying concentration of FVIIa alone (ng/ml) are
30 added (dark circles). The clotting time of the hemophilic
plasma without any added FVIIa or tTF was 88.5 sec which
is indicated by the upper dotted horizontal line. The
clotting time of normal plasma without added FVIIa or tTF
was 53.0 sec which is indicated by the lower dotted

horizontal line. Without added tTF, the 88.5 sec clotting time of the hemophilic plasma was reduced to that of normal plasma (53 sec) at 808 ng FVIIa/ml of plasma. With added tTF (at 1 μ g/ml), the clotting time of the
5 hemophilic plasma was reduced to that of normal plasma at 36.8 ng FVIIa/ml plasma.

Therefore, in the presence of 1 μ g/ml tTF in plasma, correction of the prolonged clotting time of hemophilic plasma was achieved at a level of added FVIIa that was 22-
10 fold lower than in the absence of added tTF. In addition, tTF was not able to correct the prolonged clotting time of hemophilic plasma in the absence of added FVIIa.

Example 3: Efficacy of the combination of tTF and FVIIa in the treatment of Hemophilic Dogs.

15 As described by Brinkhous, et al., Proc. Natl. Acad. Sci. USA 82, 8752-8756 (1985), Graham, et al., J. Exp. Med. 90, 97-111 (1949), and Brinkhous, et al., Ann. N.Y. Acad. Sci. 370, 191-204 (1981), a colony of hemophilic dogs has been developed at the University of North
20 Carolina at Chapel Hill. Dogs have hemophilia A. A test (secondary cuticle bleeding time) is used to measure bleeding tendency.

A modified toenail bleeding time (BT) is used to test the hemostatic effectiveness of infused preparations. The
25 paw of the front leg is warmed by placing it in isotonic saline at 37°C and a toenail is trimmed to expose only the distal matrix. The bleeding nail is placed in isotonic saline at 37°C and the time until cessation of bleeding recorded as the primary BT. At 2 to 4 hours, the site is
30 shaved to remove the clot and as little nail matrix as possible. The paw is again placed in saline at 37°C. A discrete stream of extruding blood is visible. With hemophilic dogs in the absence of treatment, bleeding typically continues for 30 min or longer. This time to

cessation of bleeding is recorded as a secondary BT. For normal dogs, the primary BT is two to five minutes and the secondary BT is less than five minutes. For hemophilic dogs the primary BT is similar to that of normal dogs while secondary BT is greater than 15 minutes.

Dog #V02 was administered a low dose of FVIIa, 6 μ g FVIIa/kg body weight, which was designed to provide only a weak hemostatic effect. After 15 min equilibrium, secondary cuticle bleeding time was measured at 2 min 30 sec, which spontaneously rebled for 5 min 15 sec. Rebleeding challenge (wipe off clot with gauze): 9 min 20 sec bleeding time, which again spontaneously rebled for 15 min, at which time the nail was cauterized to prevent further bleeding. It was difficult to stop bleeding and oozing from the phlebotomy sites in this dog, which continued after FVIIa administration.

The results indicate that this dosage of FVIIa had a very weak hemostatic effect.

The dog was then administered 42 μ g tTF/kg body weight. After 15 min equilibration, secondary cuticle bleeding time was measured at 50 sec, which did not spontaneously rebleed. This was dramatically shorter than when the animal had received FVIIa alone. Wiping the clot off with gauze to provoke rebleeding caused rebleeding for only 35 sec, and no subsequent rebleeding was observed. Oozing from two phlebotomy sites in forepaw veins also stopped following infusion of tTF.

The results indicated that tTF in combination with low dose FVIIa has an excellent hemostatic effect.

The same dog was administered 42 μ g tTF/kg body weight, without FVIIa, two days later (since the half-lives of both tTF and FVIIa are about 2 hr, this was more than sufficient to ensure clearance of either substance from the plasma). After 15 minutes equilibration, the

secondary cuticle bleeding time was greater than 15 minutes and had to be stopped by cauterization.

The results indicated that tTF alone had no measurable hemostatic effect.

5 The dog was then administered 6 μ g FVIIa/kg body weight. After 15 minutes equilibration, secondary cuticle bleeding time was measured at 3 minutes 15 seconds, which did not spontaneously rebleed. Wiping the clot off with gauze caused rebleeding for 3 minutes, and no subsequent
10 rebleeding was observed.

The results again demonstrate that tTF in combination with low dose FVIIa has a very good hemostatic effect.

**Example 4: Infusion of Factor Xa in combination with
 phospholipid vesicles (PCPS) to generate
15 Factor VIIa to inhibit bleeding.**

Three chimpanzees were infused with a bolus injection of FXa/PCPS as described by Giles, et al., (1988), the teachings of which are incorporated herein by reference. Briefly, factor Xa/PCPS was infused intravenously. The
20 chimpanzee receiving the highest dose received 26 pmoles FXa + 40 pmoles PCPS per kg body weight. Plasma samples were drawn at the following times after infusion, giving the indicated plasma FVIIa levels:

| | | |
|----|---------------------|--------------------------------|
| 25 | | Plasma <u>FVIIa (ng/ml)</u> |
| | Pre-infusion | 8.0 |
| | 2 min post-infusion | 24.6 |
| 30 | 5 " " " | 65.4 |
| | 10 " " " | 125.5 |
| | 15 " " " | 146.5 |
| | 20 " " " | 132.0 |
| | 20 " " " | 101.8 |
| 35 | 60 " " " | 43.6 |
| | 90 " " " | 4.4 |

The results indicate that infusion of the tTF in combination with factor Xa/PCPS greatly increased factor VIIa levels.

Modifications and variations of the present
5 invention, a method and compositions for the treatment of excessive bleeding, will be obvious from the foregoing detailed description and are intended to come within the scope of the appended claims.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT: Morrissey, James H.
Comp, Philip C.
- (ii) TITLE OF INVENTION: Truncated Tissue Factor and FVIIa or
FVII Activator for Blood Coagulation
- (iii) NUMBER OF SEQUENCES: 4
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: Richards, Medlock & Andrews
 - (B) STREET: 1201 Elm Street, Suite 4500
 - (C) CITY: Dallas
 - (D) STATE: Texas
 - (E) COUNTRY: US
 - (F) ZIP: 75270-2197
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Floppy disk
 - (B) COMPUTER: IBM PC compatible
 - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 - (D) SOFTWARE: PatentIn Release #1.0, Version #1.25
- (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER:
 - (B) FILING DATE:
 - (C) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA:
 - (A) APPLICATION NUMBER: US 07/882202
 - (B) FILING DATE: 13-MAY-1992
- (vii) PRIOR APPLICATION DATA:
 - (A) APPLICATION NUMBER: US 08/021615
 - (B) FILING DATE: 19-FEB-1993
- (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: Trujillo, Doreen Y.
 - (B) REGISTRATION NUMBER: 35,719
 - (C) REFERENCE/DOCKET NUMBER: OMRF B34290CIPC/PCT
- (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: 214-939-4500
 - (B) TELEFAX: 214-939-4600

(2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 795 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: double
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- (iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(F) TISSUE TYPE: Fibroblast

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 34..786

(D) OTHER INFORMATION: /product= "Truncated Tissue Factor"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

```

CGTTCGGCTC GATCTCGCCG CCAACTGGTA GAC ATG GAG ACC CCT GCC TGG CCC      54
                               Met Glu Thr Pro Ala Trp Pro
                               1                               5

CGG GTC CCG CGC CCC GAG ACC GCC GTC GCT CGG ACG CTC CTG CTC GGC      102
Arg Val Pro Arg Pro Glu Thr Ala Val Ala Arg Thr Leu Leu Leu Gly
                               10                               15                               20

TGG GTC TTC GCC CAG GTG GCC GGC GCT TCA GGC ACT ACA AAT ACT GTG      150
Trp Val Phe Ala Gln Val Ala Gly Ala Ser Gly Thr Thr Asn Thr Val
                               25                               30                               35

GCA GCA TAT AAT TTA ACT TGG AAA TCA ACT AAT TTC AAG ACA ATT TTG      198
Ala Ala Tyr Asn Leu Thr Trp Lys Ser Thr Asn Phe Lys Thr Ile Leu
                               40                               45                               50                               55

GAG TGG GAA CCC AAA CCC GTC AAT CAA GTC TAC ACT GTT CAA ATA AGC      246
Glu Trp Glu Pro Lys Pro Val Asn Gln Val Tyr Thr Val Gln Ile Ser
                               60                               65                               70

ACT AAG TCA GGA GAT TGG AAA AGC AAA TGC TTT TAC ACA ACA GAC ACA      294
Thr Lys Ser Gly Asp Trp Lys Ser Lys Cys Phe Tyr Thr Thr Asp Thr
                               75                               80                               85

GAG TGT GAC CTC ACC GAC GAG ATT GTG AAG GAT GTG AAG CAG ACG TAC      342
Glu Cys Asp Leu Thr Asp Glu Ile Val Lys Asp Val Lys Gln Thr Tyr
                               90                               95                               100

TTG GCA CGG GTC TTC TCC TAC CCG GCA GGG AAT GTG GAG AGC ACC GGT      390
Leu Ala Arg Val Phe Ser Tyr Pro Ala Gly Asn Val Glu Ser Thr Gly
                               105                               110                               115

TCT GCT GGG GAG CCT CTG TAT GAG AAC TCC CCA GAG TTC ACA CCT TAC      438
Ser Ala Gly Glu Pro Leu Tyr Glu Asn Ser Pro Glu Phe Thr Pro Tyr
                               120                               125                               130                               135

CTG GAG ACA AAC CTC GGA CAG CCA ACA ATT CAG AGT TTT GAA CAG GTG      486
Leu Glu Thr Asn Leu Gly Gln Pro Thr Ile Gln Ser Phe Glu Gln Val
                               140                               145                               150

GGA ACA AAA GTG AAT GTG ACC GTA GAA GAT GAA CGG ACT TTA GTC AGA      534
Gly Thr Lys Val Asn Val Thr Val Glu Asp Glu Arg Thr Leu Val Arg
                               155                               160                               165

AGG AAC AAC ACT TTC CTA AGC CTC CGG GAT GTT TTT GGC AAG GAC TTA      582
Arg Asn Asn Thr Phe Leu Ser Leu Arg Asp Val Phe Gly Lys Asp Leu
                               170                               175                               180

```

| | |
|---|-----|
| ATT TAT ACA CTT TAT TAT TGG AAA TCT TCA AGT TCA GGA AAG AAA ACA | 630 |
| Ile Tyr Thr Leu Tyr Tyr Trp Lys Ser Ser Ser Ser Gly Lys Lys Thr | |
| 185 190 195 | |
| GCC AAA ACA AAC ACT AAT GAG TTT TTG ATT GAT GTG GAT AAA GGA GAA | 678 |
| Ala Lys Thr Asn Thr Asn Glu Phe Leu Ile Asp Val Asp Lys Gly Glu | |
| 200 205 210 215 | |
| AAC TAC TGT TTC AGT GTT CAA GCA GTG ATT CCC TCC CGA ACA GTT AAC | 726 |
| Asn Tyr Cys Phe Ser Val Gln Ala Val Ile Pro Ser Arg Thr Val Asn | |
| 220 225 230 | |
| CGG AAG AGT ACA GAC AGC CCG GTA GAG TGT ATG GGC CAG GAG AAA GGG | 774 |
| Arg Lys Ser Thr Asp Ser Pro Val Glu Cys Met Gly Gln Glu Lys Gly | |
| 235 240 245 | |
| GAA TTT AGA GAA TAAGAATTC | 795 |
| Glu Phe Arg Glu | |
| 250 | |

(2) INFORMATION FOR SEQ ID NO:2:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 251 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

| | |
|---|--|
| Met Glu Thr Pro Ala Trp Pro Arg Val Pro Arg Pro Glu Thr Ala Val | |
| 1 5 10 15 | |
| Ala Arg Thr Leu Leu Leu Gly Trp Val Phe Ala Gln Val Ala Gly Ala | |
| 20 25 30 | |
| Ser Gly Thr Thr Asn Thr Val Ala Ala Tyr Asn Leu Thr Trp Lys Ser | |
| 35 40 45 | |
| Thr Asn Phe Lys Thr Ile Leu Glu Trp Glu Pro Lys Pro Val Asn Gln | |
| 50 55 60 | |
| Val Tyr Thr Val Gln Ile Ser Thr Lys Ser Gly Asp Trp Lys Ser Lys | |
| 65 70 75 80 | |
| Cys Phe Tyr Thr Thr Asp Thr Glu Cys Asp Leu Thr Asp Glu Ile Val | |
| 85 90 95 | |
| Lys Asp Val Lys Gln Thr Tyr Leu Ala Arg Val Phe Ser Tyr Pro Ala | |
| 100 105 110 | |
| Gly Asn Val Glu Ser Thr Gly Ser Ala Gly Glu Pro Leu Tyr Glu Asn | |
| 115 120 125 | |
| Ser Pro Glu Phe Thr Pro Tyr Leu Glu Thr Asn Leu Gly Gln Pro Thr | |
| 130 135 140 | |
| Ile Gln Ser Phe Glu Gln Val Gly Thr Lys Val Asn Val Thr Val Glu | |
| 145 150 155 160 | |

Asp Glu Arg Thr Leu Val Arg Arg Asn Asn Thr Phe Leu Ser Leu Arg
 165 170 175

Asp Val Phe Gly Lys Asp Leu Ile Tyr Thr Leu Tyr Tyr Trp Lys Ser
 180 185 190

Ser Ser Ser Gly Lys Lys Thr Ala Lys Thr Asn Thr Asn Glu Phe Leu
 195 200 205

Ile Asp Val Asp Lys Gly Glu Asn Tyr Cys Phe Ser Val Gln Ala Val
 210 215 220

Ile Pro Ser Arg Thr Val Asn Arg Lys Ser Thr Asp Ser Pro Val Glu
 225 230 235 240

Cys Met Gly Gln Glu Lys Gly Glu Phe Arg Glu
 245 250

(2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 1440 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: double
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
 - (A) ORGANISM: Homo sapiens
 - (F) TISSUE TYPE: Blood
- (ix) FEATURE:
 - (A) NAME/KEY: CDS
 - (B) LOCATION: 36..1433
 - (D) OTHER INFORMATION: /product= "Tissue Factor"
 /note= "Coding portion of human factor VIII cDNA"
 /citation= ([1])
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

| | |
|---|-----|
| TCAACAGGCA GGGGCAGCAC TGCAGAGATT TCATC ATG GTC TCC CAG GCC CTC | 53 |
| Met Val Ser Gln Ala Leu | |
| 1 5 | |
| AGG CTC CTC TGC CTT CTG CTT GGG CTT CAG GGC TGC CTG GCT GCA GGC | 101 |
| Arg Leu Leu Cys Leu Leu Leu Gly Leu Gln Gly Cys Leu Ala Ala Gly | |
| 10 15 20 | |
| GGG GTC GCT AAG GCC TCA GGA GGA GAA ACA CGG GAC ATG CCG TGG AAG | 149 |
| Gly Val Ala Lys Ala Ser Gly Gly Glu Thr Arg Asp Met Pro Trp Lys | |
| 25 30 35 | |
| CCG GGG CCT CAC AGA GTC TTC GTA ACC CAG GAG GAA GCC CAC GGC GTC | 197 |
| Pro Gly Pro His Arg Val Phe Val Thr Gln Glu Glu Ala His Gly Val | |
| 40 45 50 | |

| | | | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| CTG | CAC | CGG | CGC | CGG | CGC | GCC | AAC | GCG | TTC | CTG | GAG | GAG | CTG | CGG | CCG | 245 |
| Leu | His | Arg | Arg | Arg | Arg | Ala | Asn | Ala | Phe | Leu | Glu | Glu | Leu | Arg | Pro | |
| 55 | | | | | 60 | | | | | 65 | | | | | 70 | |
| GGC | TCC | CTG | GAG | AGG | GAG | TGC | AAG | GAG | GAG | CAG | TGC | TCC | TTC | GAG | GAG | 293 |
| Gly | Ser | Leu | Glu | Arg | Glu | Cys | Lys | Glu | Glu | Gln | Cys | Ser | Phe | Glu | Glu | |
| | | | | 75 | | | | | 80 | | | | | | 85 | |
| GCC | CGG | GAG | ATC | TTC | AAG | GAC | GCG | GAG | AGG | ACG | AAG | CTG | TTC | TGG | ATT | 341 |
| Ala | Arg | Glu | Ile | Phe | Lys | Asp | Ala | Glu | Arg | Thr | Lys | Leu | Phe | Trp | Ile | |
| | | | 90 | | | | | 95 | | | | | | 100 | | |
| TCT | TAC | AGT | GAT | GGG | GAC | CAG | TGT | GCC | TCA | AGT | CCA | TGC | CAG | AAT | GGG | 389 |
| Ser | Tyr | Ser | Asp | Gly | Asp | Gln | Cys | Ala | Ser | Ser | Pro | Cys | Gln | Asn | Gly | |
| | | 105 | | | | | 110 | | | | | | 115 | | | |
| GGC | TCC | TGC | AAG | GAC | CAG | CTC | CAG | TCC | TAT | ATC | TGC | TTC | TGC | CTC | CCT | 437 |
| Gly | Ser | Cys | Lys | Asp | Gln | Leu | Gln | Ser | Tyr | Ile | Cys | Phe | Cys | Leu | Pro | |
| | 120 | | | | | 125 | | | | | 130 | | | | | |
| GCC | TTC | GAG | GGC | CGG | AAC | TGT | GAG | ACG | CAC | AAG | GAT | GAC | CAG | CTG | ATC | 485 |
| Ala | Phe | Glu | Gly | Arg | Asn | Cys | Glu | Thr | His | Lys | Asp | Asp | Gln | Leu | Ile | |
| 135 | | | | | 140 | | | | | 145 | | | | | 150 | |
| TGT | GTG | AAC | GAG | AAC | GGC | GGC | TGT | GAG | CAG | TAC | TGC | AGT | GAC | CAC | ACG | 533 |
| Cys | Val | Asn | Glu | Asn | Gly | Gly | Cys | Glu | Gln | Tyr | Cys | Ser | Asp | His | Thr | |
| | | | | 155 | | | | | 160 | | | | | | 165 | |
| GGC | ACC | AAG | CGC | TCC | TGT | CGG | TGC | CAC | GAG | GGG | TAC | TCT | CTG | CTG | GCA | 581 |
| Gly | Thr | Lys | Arg | Ser | Cys | Arg | Cys | His | Glu | Gly | Tyr | Ser | Leu | Leu | Ala | |
| | | | 170 | | | | | 175 | | | | | | 180 | | |
| GAC | GGG | GTG | TCC | TGC | ACA | CCC | ACA | GTT | GAA | TAT | CCA | TGT | GGA | AAA | ATA | 629 |
| Asp | Gly | Val | Ser | Cys | Thr | Pro | Thr | Val | Glu | Tyr | Pro | Cys | Gly | Lys | Ile | |
| | | 185 | | | | | 190 | | | | | | 195 | | | |
| CCT | ATT | CTA | GAA | AAA | AGA | AAT | GCC | AGC | AAA | CCC | CAA | GGC | CGA | ATT | GTG | 677 |
| Pro | Ile | Leu | Glu | Lys | Arg | Asn | Ala | Ser | Lys | Pro | Gln | Gly | Arg | Ile | Val | |
| | 200 | | | | | 205 | | | | | 210 | | | | | |
| GGG | GGC | AAG | GTG | TGC | CCC | AAA | GGG | GAG | TGT | CCA | TGG | CAG | GTC | CTG | TTG | 725 |
| Gly | Gly | Lys | Val | Cys | Pro | Lys | Gly | Glu | Cys | Pro | Trp | Gln | Val | Leu | Leu | |
| 215 | | | | | 220 | | | | | 225 | | | | | 230 | |
| TTG | GTG | AAT | GGA | GCT | CAG | TTG | TGT | GGG | GGG | ACC | CTG | ATC | AAC | ACC | ATC | 773 |
| Leu | Val | Asn | Gly | Ala | Gln | Leu | Cys | Gly | Gly | Thr | Leu | Ile | Asn | Thr | Ile | |
| | | | | 235 | | | | | 240 | | | | | 245 | | |
| TGG | GTG | GTC | TCC | GCG | GCC | CAC | TGT | TTC | GAC | AAA | ATC | AAG | AAC | TGG | AGG | 821 |
| Trp | Val | Val | Ser | Ala | Ala | His | Cys | Phe | Asp | Lys | Ile | Lys | Asn | Trp | Arg | |
| | | | 250 | | | | | 255 | | | | | | 260 | | |
| AAC | CTG | ATC | GCG | GTG | CTG | GGC | GAG | CAC | GAC | CTC | AGC | GAG | CAC | GAC | GGG | 869 |
| Asn | Leu | Ile | Ala | Val | Leu | Gly | Glu | His | Asp | Leu | Ser | Glu | His | Asp | Gly | |
| | | 265 | | | | | 270 | | | | | | | 275 | | |
| GAT | GAG | CAG | AGC | CGG | CGG | GTG | GCG | CAG | GTC | ATC | ATC | CCC | AGC | ACG | TAC | 917 |
| Asp | Glu | Gln | Ser | Arg | Arg | Val | Ala | Gln | Val | Ile | Ile | Pro | Ser | Thr | Tyr | |
| | 280 | | | | | 285 | | | | | 290 | | | | | |

| | |
|---|------|
| GTC CCG GGC ACC ACC AAC CAC GAC ATC GCG CTG CTC CGC CTG CAC CAG Val Pro Gly Thr Thr Asn His Asp Ile Ala Leu Leu Arg Leu His Gln 295 300 305 310 | 965 |
| CCC GTG GTC CTC ACT GAC CAT GTG GTG CCC CTC TGC CTG CCC GAA CGG Pro Val Val Leu Thr Asp His Val Val Pro Leu Cys Leu Pro Glu Arg 315 320 325 | 1013 |
| ACG TTC TCT GAG AGG ACG CTG GCC TTC GTG CGC TTC TCA TTG GTC AGC Thr Phe Ser Glu Arg Thr Leu Ala Phe Val Arg Phe Ser Leu Val Ser 330 335 340 | 1061 |
| GGC TGG GGC CAG CTG CTG GAC CGT GGC GCC ACG GCC CTG GAG CTC ATG Gly Trp Gly Gln Leu Leu Asp Arg Gly Ala Thr Ala Leu Glu Leu Met 345 350 355 | 1109 |
| GTG CTC AAC GTG CCC CGG CTG ATG ACC CAG GAC TGC CTG CAG CAG TCA Val Leu Asn Val Pro Arg Leu Met Thr Gln Asp Cys Leu Gln Gln Ser 360 365 370 | 1157 |
| CGG AAG GTG GGA GAC TCC CCA AAT ATC ACG GAG TAC ATG TTC TGT GCC Arg Lys Val Gly Asp Ser Pro Asn Ile Thr Glu Tyr Met Phe Cys Ala 375 380 385 390 | 1205 |
| GGC TAC TCG GAT GGC AGC AAG GAC TCC TGC AAG GGG GAC AGT GGA GGC Gly Tyr Ser Asp Gly Ser Lys Asp Ser Cys Lys Gly Asp Ser Gly Gly 395 400 405 | 1253 |
| CCA CAT GCC ACC CAC TAC CGG GGC ACG TGG TAC CTG ACG GGC ATC GTC Pro His Ala Thr His Tyr Arg Gly Thr Trp Tyr Leu Thr Gly Ile Val 410 415 420 | 1301 |
| AGC TGG GGC CAG GGC TGC GCA ACC GTG GGC CAC TTT GGG GTG TAC ACC Ser Trp Gly Gln Gly Cys Ala Thr Val Gly His Phe Gly Val Tyr Thr 425 430 435 | 1349 |
| AGG GTC TCC CAG TAC ATC GAG TGG CTG CAA AAG CTC ATG CGC TCA GAG Arg Val Ser Gln Tyr Ile Glu Trp Leu Gln Lys Leu Met Arg Ser Glu 440 445 450 | 1397 |
| CCA CGC CCA GGA GTC CTC CTG CGA GCC CCA TTT CCC TAGCCCA Pro Arg Pro Gly Val Leu Arg Ala Pro Phe Pro 455 460 465 | 1440 |

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 466 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Met Val Ser Gln Ala Leu Arg Leu Leu Cys Leu Leu Leu Gly Leu Gln
1 5 10 15
Gly Cys Leu Ala Ala Gly Gly Val Ala Lys Ala Ser Gly Gly Glu Thr
20 25 30

Arg Asp Met Pro Trp Lys Pro Gly Pro His Arg Val Phe Val Thr Gln
 35 40 45
 Glu Glu Ala His Gly Val Leu His Arg Arg Arg Ala Asn Ala Phe
 50 55 60
 Leu Glu Glu Leu Arg Pro Gly Ser Leu Glu Arg Glu Cys Lys Glu Glu
 65 70 75 80
 Gln Cys Ser Phe Glu Glu Ala Arg Glu Ile Phe Lys Asp Ala Glu Arg
 85 90 95
 Thr Lys Leu Phe Trp Ile Ser Tyr Ser Asp Gly Asp Gln Cys Ala Ser
 100 105 110
 Ser Pro Cys Gln Asn Gly Gly Ser Cys Lys Asp Gln Leu Gln Ser Tyr
 115 120 125
 Ile Cys Phe Cys Leu Pro Ala Phe Glu Gly Arg Asn Cys Glu Thr His
 130 135 140
 Lys Asp Asp Gln Leu Ile Cys Val Asn Glu Asn Gly Gly Cys Glu Gln
 145 150 155 160
 Tyr Cys Ser Asp His Thr Gly Thr Lys Arg Ser Cys Arg Cys His Glu
 165 170 175
 Gly Tyr Ser Leu Leu Ala Asp Gly Val Ser Cys Thr Pro Thr Val Glu
 180 185 190
 Tyr Pro Cys Gly Lys Ile Pro Ile Leu Glu Lys Arg Asn Ala Ser Lys
 195 200 205
 Pro Gln Gly Arg Ile Val Gly Gly Lys Val Cys Pro Lys Gly Glu Cys
 210 215 220
 Pro Trp Gln Val Leu Leu Leu Val Asn Gly Ala Gln Leu Cys Gly Gly
 225 230 235 240
 Thr Leu Ile Asn Thr Ile Trp Val Val Ser Ala Ala His Cys Phe Asp
 245 250 255
 Lys Ile Lys Asn Trp Arg Asn Leu Ile Ala Val Leu Gly Glu His Asp
 260 265 270
 Leu Ser Glu His Asp Gly Asp Glu Gln Ser Arg Arg Val Ala Gln Val
 275 280 285
 Ile Ile Pro Ser Thr Tyr Val Pro Gly Thr Thr Asn His Asp Ile Ala
 290 295 300
 Leu Leu Arg Leu His Gln Pro Val Val Leu Thr Asp His Val Val Pro
 305 310 315 320
 Leu Cys Leu Pro Glu Arg Thr Phe Ser Glu Arg Thr Leu Ala Phe Val
 325 330 335
 Arg Phe Ser Leu Val Ser Gly Trp Gly Gln Leu Leu Asp Arg Gly Ala
 340 345 350

Thr Ala Leu Glu Leu Met Val Leu Asn Val Pro Arg Leu Met Thr Gln
355 360 365

Asp Cys Leu Gln Gln Ser Arg Lys Val Gly Asp Ser Pro Asn Ile Thr
370 375 380

Glu Tyr Met Phe Cys Ala Gly Tyr Ser Asp Gly Ser Lys Asp Ser Cys
385 390 395 400

Lys Gly Asp Ser Gly Gly Pro His Ala Thr His Tyr Arg Gly Thr Trp
405 410 415

Tyr Leu Thr Gly Ile Val Ser Trp Gly Gln Gly Cys Ala Thr Val Gly
420 425 430

His Phe Gly Val Tyr Thr Arg Val Ser Gln Tyr Ile Glu Trp Leu Gln
435 440 445

Lys Leu Met Arg Ser Glu Pro Arg Pro Gly Val Leu Leu Arg Ala Pro
450 455 460

Phe Pro
465

We claim:

1. A composition for treatment of patients with prolonged or excessive bleeding, comprising truncated tissue factor and a substance effective to produce a plasma level of factor VIIa which in combination with said truncated tissue factor will control or stop said excessive bleeding.

2. The composition of claim 1, wherein said substance is selected from factor VIIa and an activator which promotes the conversion of said patient's endogenous factor VII to factor VIIa.

3. The composition of claim 1 wherein said truncated tissue factor is in a dosage for administration to a patient in an amount effective to produce a concentration of between 100 ng and 50 μ g truncated tissue factor per milliliter of plasma and said substance is in a dosage in an amount effective to produce levels of between 20 ng and 700 μ g factor VIIa per milliliter of plasma.

4. The composition of claim 1, further comprising a pharmaceutically acceptable carrier and wherein said truncated tissue factor is in a concentration of between 100 ng and 50 μ g/ml of carrier and wherein the factor VIIa activator which activates factor VII to yield factor VIIa is in a concentration of between 1 and 10 μ g/ml of carrier.

5. The composition of claim 2 wherein said substance is said activator which is selected from the group consisting of factor Xa in combination with phospholipid, factor IXa in combination with phospholipid, thrombin, factor XIIa, and the FVII activator from the venom of *Oxyuranus scutellatus* in combination with phospholipid.

6. The composition according to claim 1, wherein the truncated tissue factor and factor VIIa are in a pharmaceutically acceptable carrier for topical administration to a patient, and the truncated tissue factor is in a concentration of between 100 ng and 50 $\mu\text{g/ml}$ of carrier and the factor VIIa is in a concentration of between 1 and 10 $\mu\text{g/ml}$ of carrier.

7. The composition according to claim 6, wherein said combination is formulated so that a dose of said combination provides a plasma level of between about 100 ng and 50 μg truncated tissue factor per milliliter of plasma and between about 20 ng and 10 μg factor VIIa per milliliter of plasma.

8. The composition of claim 6, wherein said truncated tissue factor is in a concentration of between 10 and 50 $\mu\text{g/ml}$.

9. A method for treating patients with excessive bleeding or at risk of excessive bleeding comprising administering to the patients truncated tissue factor, wherein the tissue factor is administered in an amount
5 effective to produce between 100 ng/ml and 50 μ g truncated tissue factor/ml of plasma and an activator which promotes the conversion of endogenous factor VII to factor VIIa in an amount effective to produce levels of between about 20 ng FVIIa/ml and 10 μ g FVIIa/ml of plasma.

10. The method of claim 9 wherein the truncated tissue factor and FVIIa activator are administered in combination.

11. The method of claim 9 wherein the truncated tissue factor and FVIIa activator are administered sequentially.

12. The method of claim 9 wherein the truncated tissue factor and FVIIa activator are administered systemically to a patient in need of such therapy.

13. The method of claim 12 comprising administering truncated tissue factor and FVIIa activator systemically to produce levels of between 1 and 10 μ g tTF/ml plasma and between 40 ng and 700 ng VIIa/ml plasma.

14. The method of claim 9 wherein the truncated tissue factor and FVIIa activator are administered topically to a patient in an effective amount to decrease the bleeding time.

15. The method of claim 14 comprising administering truncated tissue factor and FVIIa activator topically to produce levels of between 10 and 50 μ g tTF and between 40 and 700 ng FVIIa/ml.

16. The method of claim 9 wherein the activator of endogenous FVII is selected from the group consisting of factor Xa in combination with phospholipid, factor IXa in combination with phospholipid, thrombin, factor XIIa, and
5 the FVII activator from the venom of *Oxyuranus scutellatus* in combination with phospholipid.

17. The method of claim 16 wherein the activator is factor Xa in a range of 12 to greater than 26 pmoles per kg body weight in combination with phospholipid in a range of 19 pmoles to greater than 40 pmoles per kg body weight.

18. A method for treating patients with excessive bleeding or at risk of excessive bleeding comprising administering to the patients truncated tissue factor to produce between 100 ng/ml and 50 μ g truncated tissue factor/ml of plasma and factor VIIa to produce levels of
5 between 20 ng FVIIa/ml and 10 μ g FVIIa/ml of plasma.

19. The method of claim 18 wherein the truncated tissue factor and FVIIa are administered in combination.

20. The method of claim 18 wherein the truncated tissue factor and FVIIa are administered sequentially.

21. The method of claim 18 wherein the truncated tissue factor and FVIIa are administered systemically to a patient in need of such therapy.

22. The method of claim 21 comprising administering truncated tissue factor and FVIIa systemically to produce levels of between 1 and 10 μ g tTF/ml plasma and between 40 ng and 4 μ g VIIa/ml plasma.

23. The method of claim 18 wherein the truncated tissue factor and FVIIa are administered topically to a patient in an effective amount to decrease the bleeding time.

24. The method of claim 23 comprising administering truncated tissue factor and FVIIa topically to produce levels of between 10 and 50 μ g tTF and between 1 and 10 μ g FVIIa/ml.




FIG. 1A




FIG. 1B

INTERNATIONAL SEARCH REPORT

PCT/US 93/04493

International Application No

| | | |
|--|--|-------------------------------------|
| I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) ⁶ | | |
| According to International Patent Classification (IPC) or to both National Classification and IPC Int.C1. 5 A61K37/547; //(A61K37/547, 37:02) | | |
| II. FIELDS SEARCHED | | |
| Minimum Documentation Searched ⁷ | | |
| Classification System | Classification Symbols | |
| Int.C1. 5 | A61K | |
| Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched ⁸ | | |
| III. DOCUMENTS CONSIDERED TO BE RELEVANT⁹ | | |
| Category ¹⁰ | Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹² | Relevant to Claim No. ¹³ |
| A | JOURNAL OF BIOLOGICAL CHEMISTRY. vol. 266, no. 4, 5 February 1991, BALTIMORE US pages 2158 - 2166 W. RUF ET AL. 'PHOSPHOLIPID-INDEPENDENT AND -DEPENDENT INTERACTIONS REQUIRED FOR TISSUE FACTOR RECEPTOR AND COFACTOR FUNCTION.' cited in the application see page 2158, right column, line 29 - line 46 --- -/-- | 1-24 |
| <p>¹⁰ Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&" document member of the same patent family</p> | | |
| IV. CERTIFICATION | | |
| Date of the Actual Completion of the International Search | Date of Mailing of this International Search Report | |
| 02 SEPTEMBER 1993 | 23. 09. 93 | |
| International Searching Authority | Signature of Authorized Officer | |
| EUROPEAN PATENT OFFICE | RYCKEBOSCH A.O. | |

| III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET) | | |
|--|---|-----------------------|
| Category ° | Citation of Document, with indication, where appropriate, of the relevant passages | Relevant to Claim No. |
| A | <p>FASEB JOURNAL vol. 6, no. 1, 1 January 1992, BETHESDA, MD US page A330 R.C. HAPAK ET AL. 'THE LOCATION OF THE ACTIVE SITE OF FACTOR VIIa ABOVE THE MEMBRANE SURFACE IS ALTERED BY EITHER FULL-LENGTH OR TRUNCATED TISSUE FACTOR.' see abstract nr. 1900 ----</p> | 1-24 |
| A | <p>EP,A,0 225 160 (NOVO INDUSTRI A/S) 10 June 1987 see page 2, line 30 - line 38; claims ----</p> | 1-24 |
| A | <p>CHEMICAL ABSTRACTS, vol. 109, no. 17, 24 October 1988, Columbus, Ohio, US; abstract no. 142300r, A.R. GILES ET AL. 'A COMBINATION OF FACTOR Xa AND PHOSPHATIDYLCHOLINE-PHOSPHATIDYLSERINE VESICLES BYPASSES FACTOR VIII IN VIVO.' page 42 ; see abstract & BR. J. HAEMATOL. vol. 69, no. 4, 1988, pages 491 - 497 cited in the application -----</p> | 1-24 |

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 93/04493

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Remark: Although claims 9-24 are directed to a method of treatment of the human/animal body the research has been carried out and based on the alleged effects of the compound/composition.

2. Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

 The additional search fees were accompanied by the applicant's protest. No protest accompanied the payment of additional search fees.

**ANNEX TO THE INTERNATIONAL SEARCH REPORT
ON INTERNATIONAL PATENT APPLICATION NO.**

US 9304493
SA 74681

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information. 02/09/93

| Patent document cited in search report | Publication date | Patent family member(s) | Publication date |
|--|------------------|-------------------------|------------------|
| EP-A-0225160 | 10-06-87 | AU-B- 593042 | 01-02-90 |
| | | AU-A- 6567086 | 28-05-87 |
| | | DE-A- 3680994 | 26-09-91 |
| | | US-A- 5180583 | 19-01-93 |
| | | JP-A- 62195335 | 28-08-87 |
| ----- | | | |

EPO FORM P0479

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82