

(19)



(11) Publication number:

SG 193447 A1

(43) Publication date:

30.10.2013

(51) Int. Cl:

A61K 39/395, A61K 39/00;

(12)

Patent Application

(21) Application number: **2013069026**

(71) Applicant:

**IMMUNOGEN, INC. 830 WINTER
STREET WALTHAM, MA 02451 MC US**

(22) Date of filing: **30.03.2012**

(30) Priority: **US 61/470,863 01.04.2011**

(72) Inventor:

**DECKERT, JUTTA 16 EATON ROAD
LEXINGTON, MA 02420 US
SETIADY, JULIANTO 132
CLOCKTOWER DRIVE UNIT 4110
WALTHAM, MA 02452 US
PARK, PETER U. 25 WEBSTER AVENUE
#301 SOMERVILLE, MA 02143 US**

(54) **Title:**

**CD37-BINDING MOLECULES AND IMMUNOCONJUGATES
THEREOF**

(57) **Abstract:**

Methods of using CD37 agents, including, but not limited to, antibodies and immunoconjugates, that bind to CD37 to deplete B-cells (e.g., non-cancerous B-cells) and methods of treating autoimmune and inflammatory diseases are further provided.



- (51) International Patent Classification:
A61K 39/395 (2006.01) *A61P 29/00* (2006.01)
C12N 5/0781 (2010.01) *C07K 16/28* (2006.01)
- (21) International Application Number:
PCT/US2012/031648
- (22) International Filing Date:
30 March 2012 (30.03.2012)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
61/470,863 1 April 2011 (01.04.2011) US
- (71) Applicant (for all designated States except US): **IMMUNOGEN, INC.** [US/US]; 830 Winter Street, Waltham, MA 02451 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **DECKERT, Jutta** [US/US]; 16 Eaton Road, Lexington, MA 02420 (US). **SETIADY, Julianto** [ID/US]; 132 Clocktower Drive, Unit 4110, Waltham, MA 02452 (US). **PARK, Peter U.**

[US/US]; 25 Webster Avenue, #301, Somerville, MA 02143 (US).

- (74) Agents: **STEFFE, Eric K.** et al.; Sterne Kessler Goldstein Fox P.L.L.C., 1100 New York Avenue N.W., Washington, DC 20005 (US).


- (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU,

[Continued on next page]

(54) Title: CD37-BINDING MOLECULES AND IMMUNOCONJUGATES THEREOF

Figure 4



(57) Abstract: Methods of using CD37 agents, including, but not limited to, antibodies and immunoconjugates, that bind to CD37 to deplete B-cells (e.g., non-cancerous B-cells) and methods of treating autoimmune and inflammatory diseases are further provided.



LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK,
SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).

— with (an) indication(s) in relation to deposited biological material furnished under Rule 13bis separately from the description (Rules 13bis.4(d)(i) and 48.2(a)(viii))

Published:

— without international search report and to be republished upon receipt of that report (Rule 48.2(g))

CD37-BINDING MOLECULES AND IMMUNOCONJUGATES THEREOF

Field of the Invention

[0001] The field of the invention generally relates to antibodies, antigen-binding fragments thereof, polypeptides, and immunoconjugates that bind to CD37, as well as to methods of using such CD37-binding molecules for the treatment of diseases, such as autoimmune diseases and inflammatory diseases.

Background of the Invention

[0002] Leukocyte antigen CD37 ("CD37"), also known as GP52-40, tetraspanin-26, or TSPAN26, is a transmembrane protein of the tetraspanin superfamily (Maecker et al., 1997 FASEB J. 11:428-442). It is a heavily glycosylated protein with four transmembrane domains that is expressed on B cells during the pre-B to peripheral mature B-cell stages, but is reportedly absent on terminal differentiation to plasma cells. (Link et al., 1987, J Pathol. 152:12-21). The CD37 antigen is only weakly expressed on T-cells, myeloid cells, and granulocytes (Schwartz-Albiez et al. 1988, J. Immunol., 140(3)905-914). However, CD37 is also expressed on malignant B-cells such as those founding non-Hodgkin's lymphoma (NHL) and chronic lymphoid leukemia (CLL) (Moore et al. 1986, J Immunol. 137(9):3013-8).

[0003] While the exact physiological role of CD37 is unclear, studies in CD37-deficient mice suggest an immunoregulatory function. Although mice deficient in CD37 expression have normal development (Knobeloch et al. 2000, Mol Cell Biol., 20(15):5363-9), in the C57/Bl6 background, CD37^{-/-} T cells are hyper-proliferative (van Spriël et al., J Immunol. 172, 2953 (2004)), CD37^{-/-} dendritic cells (DC) exhibit an increased antigen presentation (Sheng et al., Eur J Immunol. 39, 50 (2009)), and CD37^{-/-} macrophages show increased dectin-1-induced IL-6 production (Meyer-Wentrup et al., J Immunol. 178, 154 (2007)). CD37-deficient C57/Bl6 mice also contain significantly higher level of IgA than the wild-type mice (van Spriël et al., PLoS Pathol. 5, e1000338 (2009) and Rops et al., Am J Pathol. 176, 2188 (2010)). All of these results suggest a general regulatory role of CD37 in the immune system. Interestingly, crosslinking of CD37 antigen by antibody on human T cells inhibits T cell proliferation induced by CD3 stimulation (van Spriël et al., J Immunol. 172, 2953 (2004)).

[0004] Antibodies are emerging as a promising method to treat human diseases including autoimmune diseases. Currently, an anti-CD20 antibody called rituximab has been approved for rheumatoid arthritis (RA) treatment (Edwards JC et al. 2006, Nat Rev Immunol. 6: 119). Rituximab is used in the United States in combination with methotrexate (MTX) to reduce signs and symptoms in adult patients with moderately- to severely-active RA who have had an inadequate response to at least one TNF antagonist. Many studies address the use of rituximab in a variety of non-malignant autoimmune or inflammatory disorders, including RA, in which B-cells and autoantibodies appear to play a role in disease pathophysiology. Edwards et al., Biochem Soc. Trans. 30:824-828 (2002). Targeting of CD20

using anti-CD20 antibody has been reported to potentially relieve signs and symptoms of a number of autoimmune or inflammatory diseases including, for example, RA (Leandro et al., *Ann. Rheum. Dis.* 61:883-888 (2002); Edwards et al., *Arthritis Rheum.*, 46 (Suppl. 9): S46 (2002); Stahl et al., *Ann. Rheum. Dis.*, 62 (Suppl. 1): OP004 (2003); Emery et al., *Arthritis Rheum.* 48(9): S439 (2003)), lupus (Eisenberg, *Arthritis. Res. Ther.* 5:157-159 (2003); Leandro et al. *Arthritis Rheum.* 46: 2673-2677 (2002); Gorman et al., *Lupus*, 13: 312-316 (2004)), immune thrombocytopenic purpura (D'Arena et al., *Leuk. Lymphoma* 44:561-562 (2003); Stasi et al., *Blood*, 98: 952-957 (2001); Saleh et al., *Semin. Oncol.*, 27 (Supp 12):99-103 (2000); Zaja et al., *Haematologica*, 87:189-195 (2002); Ratanatharathorn et al., *Ann. Int. Med.*, 133:275-279 (2000)), pure red cell aplasia (Auner et al., *Br. J. Haematol.*, 116:725-728 (2002)); autoimmune anemia (Zaja et al., *supra* (erratum appears in *Haematologica* 87:336 (2002)), cold agglutinin disease (Layios et al., *Leukemia*, 15:187-8 (2001); Berentsen et al., *Blood*, 103: 2925-2928 (2004); Berentsen et al., *Br. J. Haematol.*, 115:79-83 (2001); Bauduer, *Br. J. Haematol.*, 112:1083-1090 (2001); Zaja et al., *Br. J. Haematol.*, 115:232-233 (2001)), type B syndrome of severe insulin resistance (Coll et al., *N. Engl. J. Med.*, 350:310-311 (2004), mixed cryoglobulinemia (DeVita et al., *Arthritis Rheum.* 46 Suppl. 9:S206/S469 (2002)), myasthenia gravis (Zaja et al., *Neurology*, 55:1062-1063 (2000); Wylam et al., *J. Pediatr.*, 143:674-677 (2003)), Wegener's granulomatosis (Specks et al., *Arthritis & Rheumatism* 44:2836-2840 (2001)), microscopic polyangiitis (MPA), refractory pemphigus vulgaris (Dupuy et al., *Arch Dermatol.*, 140:91-96 (2004)), dermatomyositis (Levine, *Arthritis Rheum.*, 46 (Suppl. 9):S1299 (2002)), Sjogren's syndrome (Sommer et al., *Arthritis & Rheumatism*, 49:394-398 (2003)), active type-II mixed cryoglobulinemia (Zaja et al., *Blood*, 101:3827-3834 (2003)), pemphigus vulgaris (Dupay et al., *Arch. Dermatol.*, 140:91-95 (2004)), autoimmune neuropathy (Pestronk et al., *J. Neurol. Neurosurg. Psychiatry* 74:485-489 (2003)), paraneoplastic opsoclonus-myoclonus syndrome (Pranzatelli et al. *Neurology* 60 (Suppl. 1) PO5.128:A395 (2003)), and relapsing-remitting multiple sclerosis (RRMS). Cross et al. (abstract) "Preliminary Results from a Phase II Trial of Rituximab in MS" Eighth Annual Meeting of the Americas Committees for Research and Treatment in Multiple Sclerosis, 20-21 (2003).

[0005] In animal models, B-cell depletion using antibodies against B-cell antigens such as CD20 has been shown to inhibit or ameliorate several autoimmune diseases including systemic lupus erythematosus (SLE), experimental autoimmune encephalomyelitis (EAE; mouse model of multiple sclerosis), type-1 diabetes (T1D) and rheumatoid arthritis (RA). Rituximab has been shown to deplete both malignant and normal B cells *in vivo* in animal models as well as patients (Maloney DG et al, *Blood*. 1994;84(8):2457-66; Reff ME, et al. *Blood*. 1994;83(2):435-45; Schröder C, et al. *Transpl Immunol*. 2003;12(1):19-28). It can also deplete normal B-cells from human peripheral blood mononuclear cells (PBMCs) in *in vitro* experiments (Vugmeyster Y, et al, *Cytometry A*. 2003;52(2):101-9; Vugmeyster Y and Howell K. *Int Immunopharmacol*. 2004;4(8):1117-24).

[0006] Campath-1H (alumtuzumab), an anti-CD52 chimeric IgG1, binds to the CD52 antigen, which is highly expressed on all lymphocytes (Ginaldi L, et al, Leuk Res. 1998 Feb;22(2):185-91; Hale G, et al, Tissue Antigens. 1990 Mar;35(3):118-27). It is used in patients to deplete malignant lymphocytes and is approved for treating chronic lymphocytic leukemia. It has also shown efficacy in treating multiple sclerosis and is currently in Phase III clinical testing (N Engl J Med 2008; 359:1786-1801; ClinicalTrials.gov NCT00530348 & NCT00548405). It has been shown to deplete normal lymphocytes *in vitro* as well (Hale G, et al. Blood. 1983 Oct;62(4):873-82; Waldmann H and Hale G Philos Trans R Soc Lond B Biol Sci. 2005 Sep 29;360(1461):1707-11).

[0007] CD37-binding agents are also being tested as potential therapeutics for B-cell malignancies. Emergent Biosolutions (formerly Trubion Pharmaceuticals) developed the CD37-binding agents SMIP-016 and TRU-016 (Zhao et al., 2007, Blood, 110:2569-2577). SMIP-016 is a single chain polypeptide that includes variable regions from a hybridoma and engineered human constant regions. TRU-016 is a humanized version of the anti-CD37 SMIP protein. See e.g. U.S. Published Application No. 2007/0059306. TRU-016 is being tested clinically for the treatment of chronic lymphocytic leukemia (CLL). Boehringer Ingelheim has also disclosed a CD37 binding agent in International Published Application No. WO 2009/019312. However, no CDC activity has been described for any of these binding agents and no *in vitro* pro-apoptotic activity has been described in the absence of cross-linking agents.

[0008] Radio-immunotherapy (RIT) has been attempted using a radio-labeled anti-CD37 antibody MB-1 in two separate trials. Therapeutic doses of ¹³¹I-MB-1 were administered to six relapsed NHL patients (Press et al. 1989 J Clin Oncol. 7(8):1027-38; Press et al. 1993, N Engl J Med. 329(17):1219-24). All six patients achieved a complete remission (CR) with a duration of four to thirty-one months. In another trial, ¹³¹I-MB-1 was administered to ten relapsed NHL patients (Kaminski et al. 1992 J Clin Oncol. 10(11):1696-711). A total of four patients had a response ranging in duration from two to six months, although only one CR was reported. However, not all patients could be treated due to an unfavorable biodistribution of the radio-label which raised concern for radiation exposure of vital non-target organs. Indeed, RIT related toxicities were observed in these trials including severe myelosuppression and cardiopulmonary toxicity. While these clinical data suggest that anti-CD37 radio-immunoconjugates may be effective, these therapies are cumbersome to administer, and at relapse post-RIT patients cannot be retreated with RIT due to the risks associated with high doses of radiation.

[0009] To overcome the limitations of RIT, antibody-cytotoxic agent conjugates (ACC), also called antibody-drug conjugates (ADC), have been developed. These are immunoconjugates that include a cytotoxic agent covalently linked to an antibody through a chemical linker which can allow for specific delivery of cytotoxic drugs to cells expressing a protein recognized by the antibody. However, proteins that are poorly internalized are not considered to be favorable targets for such therapeutics. CD37 is

structurally similar to CD20 as both antigens contain four transmembrane domains, although CD20 is not part of the tetraspanin family (Tedder et al. 1989, J. Immun. 142: 2560-2568). Antibodies against several B-cell antigens including CD37 and CD20 have been studied for their ability to undergo endocytosis and degradation (Press et al. 1989, Cancer Res. 49(17):4906-12, and Press et al. 1994, Blood. 83(5):1390-7). The anti-CD37 antibody MB-1 was retained on the cell surface and internalized slowly in Daudi lymphoma cells *in vitro*. The MB-1 antibody also had a low rate of endocytosis and intracellular metabolism in NHL patient cells *in vitro*. Similar results were obtained with the anti-CD20 antibody 1F5, which was also retained mainly on the lymphoma cell surface and internalized poorly. ADCs of CD20 antibodies have been studied previously but have not demonstrated significantly strong potency, especially when non-disulfide or acid stable linkers are used (see for example Polson et al., 2009, Cancer Res., 69(6):2358-2364). In light of these observations, CD37 has not been considered a favorable target for antibody-drug conjugates.

[0010] While their role in cancer treatment has been studied, the potential effect of CD37-directed therapies such as antibodies, antibody derivatives or radio-immunoconjugates on cells involved in autoimmune diseases, inflammatory diseases or other disorders of the immune system is not well understood. Furthermore, none of the compounds described above have been demonstrated to induce depletion of target cells involved in manifestation or progression of these types of diseases.

[0011] Therefore, there exists a need for CD37 binding agents including antibodies, antigen-binding fragments thereof, and antibody-drug conjugates (immunoconjugates) as a means to treat autoimmune diseases, inflammatory diseases, or other disorders of the immune system. The present invention addresses that need.

BRIEF SUMMARY OF THE INVENTION

[0012] In one aspect, the present disclosure provides a method for depleting B-cells or treating a disease associated with aberrant B-cell activity, comprising administering to a patient an effective amount of a humanized CD37 targeting antibody or immunoconjugate provided herein. In some embodiments, the B-cells are non-cancerous B-cells. In some embodiments, the B-cells do not overexpress CD37.

[0013] In certain embodiments, the disease associated with aberrant B-cell activity is a disease associated with B-cell autoantibody production, and/or a disease associated with inappropriate T-cell stimulation in connection with a B-cell pathway.

[0014] In certain embodiments, the disease characterized by autoantibody production is rheumatoid arthritis, multiple sclerosis, type I diabetes mellitus, idiopathic inflammatory myopathy, systemic lupus erythematosus (SLE), myasthenia gravis, Grave's disease, dermatomyositis, polymyositis, or other autoimmune diseases.

[0015] In certain embodiments, the present disclosure provides a method for depleting a B-cell comprising contacting a B-cell (e.g., in a population of cells comprising a non-cancerous B-cell) with an antibody or antigen binding fragment thereof that specifically binds to CD37, wherein the antibody or fragment thereof is capable of inducing apoptosis *in vitro* in the absence of a cross-linking agent. In certain embodiments, the present disclosure provides a method for treating a patient having an autoimmune or inflammatory disease comprising administering to the patient a therapeutically effective amount of an antibody or antigen-binding fragment thereof that specifically binds to CD37, wherein the antibody or fragment thereof is capable of inducing apoptosis *in vitro* in the absence of a cross-linking agent. In some embodiments, the antibody or antigen-binding fragment thereof is also capable of inducing complement dependent cytotoxicity (CDC). In some embodiments, the antibody or antigen-binding fragment thereof is also capable of inducing antibody dependent cell mediated cytotoxicity (ADCC). In some embodiments, the antibody or antigen-binding fragment thereof has a long serum half-life.

In certain embodiments, the present disclosure provides a method for depleting a B-cell comprising contacting a B-cell (e.g., in a population of cells comprising a non-cancerous B-cell) with an antibody or antigen binding fragment thereof that specifically binds to the same CD37 epitope as an antibody selected from the group consisting of: (a) an antibody comprising the polypeptide of SEQ ID NO:55 and the polypeptide of SEQ ID NO:72; (b) an antibody comprising the polypeptide of SEQ ID NO:56 and the polypeptide of SEQ ID NO:73; (c) an antibody comprising the polypeptide of SEQ ID NO:57 and the polypeptide of SEQ ID NO:74; (d) an antibody comprising the polypeptide of SEQ ID NO:58 and the polypeptide of SEQ ID NO:74; (e) an antibody comprising the polypeptide of SEQ ID NO:59 and the polypeptide of SEQ ID NO:75; (f) an antibody comprising the polypeptide of SEQ ID NO:60 and the polypeptide of SEQ ID NO:76; (g) an antibody comprising the polypeptide of SEQ ID NO:61 and the polypeptide of SEQ ID NO:77; (h) an antibody comprising the polypeptide of SEQ ID NO:62 and the polypeptide of SEQ ID NO:78; (i) an antibody comprising the polypeptide of SEQ ID NO:63 and the polypeptide of SEQ ID NO:79; (j) an antibody comprising the polypeptide of SEQ ID NO:64 and the polypeptide of SEQ ID NO:80; (k) an antibody comprising the polypeptide of SEQ ID NO:65 and the polypeptide of SEQ ID NO:81; (l) an antibody comprising the polypeptide of SEQ ID NO:66 and the polypeptide of SEQ ID NO:82; (m) an antibody comprising the polypeptide of SEQ ID NO:67 and the polypeptide of SEQ ID NO:83; (n) an antibody comprising the polypeptide of SEQ ID NO:68 and the polypeptide of SEQ ID NO:84; (o) an antibody comprising the polypeptide of SEQ ID NO:69 and the polypeptide of SEQ ID NO:85; (p) an antibody comprising the polypeptide of SEQ ID NO:70 and the polypeptide of SEQ ID NO:86; (q) an antibody comprising the polypeptide of SEQ ID NO:71 and the polypeptide of SEQ ID NO:87; and (r) an antibody comprising the polypeptide of SEQ ID NO:177 and the polypeptide of SEQ ID NO:178.

[0016] In certain embodiments, the present disclosure provides a method for treating a patient having an autoimmune or inflammatory disease comprising administering to the patient a therapeutically effective amount of an antibody or antigen-binding fragment thereof that specifically binds to the same CD37 epitope as an antibody selected from the group described above. In some embodiments, the antibody or antigen-binding fragment thereof competitively inhibits an antibody selected from the group described above.

[0017] In certain embodiments, the present disclosure provides a method for depleting a B-cell comprising contacting a B-cell (e.g., in a population of cells comprising a non-cancerous B-cell) with an antibody or antigen-binding fragment thereof that specifically binds to CD37 and specifically binds to the polypeptide of SEQ ID NO: 184. In certain embodiments, the present disclosure provides a method for treating a patient having an autoimmune or inflammatory disease comprising administering to the patient a therapeutically effective amount of an antibody or antigen-binding fragment thereof that specifically binds to CD37 and specifically binds to the polypeptide of SEQ ID NO: 184. In some embodiments, the antibody or antigen-binding fragment thereof does not bind to the polypeptide of SEQ ID NO: 185.

[0018] In certain embodiments, the present disclosure provides a method for depleting a B-cell comprising contacting a B-cell (e.g., in a population of cells comprising a non-cancerous B-cell) with an antibody or antigen-binding fragment thereof that specifically binds to CD37 and does not specifically bind to the polypeptide of SEQ ID NO: 185. In certain embodiments, the present disclosure provides a method for treating a patient having an autoimmune or inflammatory disease comprising administering to the patient a therapeutically effective amount of an antibody or antigen-binding fragment thereof that specifically binds to CD37 and does not specifically bind to the polypeptide of SEQ ID NO: 185.

[0019] In certain embodiments, the present disclosure provides a method for depleting a B-cell comprising contacting a B-cell (e.g., in a population of cells comprising a non-cancerous B-cell) with an antibody or antigen-binding fragment thereof produced by a hybridoma selected from the group consisting of ATCC Deposit Designation PTA-10664, deposited with the ATCC on February 18, 2010, ATCC Deposit Designation PTA-10665, deposited with the ATCC on February 18, 2010, ATCC Deposit Designation PTA-10666, deposited with the ATCC on February 18, 2010, ATCC Deposit Designation PTA-10667, deposited with the ATCC on February 18, 2010, ATCC Deposit Designation PTA-10668, deposited with the ATCC on February 18, 2010, ATCC Deposit Designation PTA-10669, deposited with the ATCC on February 18, 2010, and ATCC Deposit Designation PTA-10670, deposited with the ATCC on February 18, 2010. In certain embodiments, the present disclosure provides a method for treating a patient having an autoimmune or inflammatory disease comprising administering to the patient a therapeutically effective amount of an antibody or antigen-binding fragment thereof produced by a hybridoma described above.

[0020] In certain embodiments, the present disclosure provides a method for depleting a B-cell comprising contacting a B-cell (e.g., in a population of cells comprising a non-cancerous B-cell) with an antibody or antigen-binding fragment thereof that specifically binds to CD37, wherein the antibody comprises polypeptide sequences selected from the group consisting of: (a) SEQ ID NOs: 4, 5, and 6 and SEQ ID NOs: 28, 29, and 30; (b) SEQ ID NOs: 7, 8, and 9 and SEQ ID NOs: 31, 32, and 33; (c) SEQ ID NOs: 10, 11, and 12 and SEQ ID NOs: 34, 35, and 36; (d) SEQ ID NOs: 13, 14, and 15 and SEQ ID NOs: 37, 38, and 39; (e) SEQ ID NOs: 13, 14, and 15 and SEQ ID NOs: 37, 40, and 39; (f) SEQ ID NOs: 16, 17, and 18 and SEQ ID NOs: 41, 42, and 43; (g) SEQ ID NOs: 19, 20, and 21 and SEQ ID NOs: 44, 45, and 46; (h) SEQ ID NOs: 19, 20, and 21 and SEQ ID NOs: 44, 47, and 46; (i) SEQ ID NOs: 22, 23, and 24 and SEQ ID NOs: 48, 49, and 50; (j) SEQ ID NOs: 22, 23, and 24 and SEQ ID NOs: 48, 51, and 50; (k) SEQ ID NOs: 25, 26, and 27 and SEQ ID NOs: 52, 53, and 54; (l) SEQ ID NOs: 171, 172 or 181, and 173 and SEQ ID NOs: 174, 175, and 176; (m) variants of (a) to (l) comprising 1, 2, 3, or 4 conservative amino acid substitutions. In certain embodiments, the present disclosure provides a method for treating a patient having an autoimmune or inflammatory disease comprising administering to the patient a therapeutically effective amount of an antibody or antigen-binding fragment thereof with an antibody or antigen-binding fragment thereof that specifically binds to CD37, wherein the antibody comprises polypeptide sequences selected from the group described above. In some embodiments, the antibody or antigen-binding fragment thereof comprises polypeptide sequences that are at least 90% identical to polypeptide sequences described above. In some embodiments, the polypeptide sequences are at least 95% identical to the polypeptide sequences. In some embodiments, the polypeptide sequences are at least 99% identical to the polypeptide sequences. In some embodiments, the antibody or antigen-binding fragment thereof comprises polypeptide sequences that are at least 90% identical, at least 95% identical, at least 99% identical, or identical to the polypeptide sequences of SEQ ID NO: 57 and SEQ ID NO:74. In some embodiments, the antibody or antigen-binding fragment thereof comprises polypeptide sequences that are at least 90% identical, at least 95% identical, at least 99% identical, or identical to the polypeptide sequences of SEQ ID NO: 58 and SEQ ID NO:74. In some embodiments, the antibody or antigen-binding fragment thereof comprises polypeptide sequences that are at least 90% identical, at least 95% identical, at least 99% identical, or identical to the polypeptide sequences of SEQ ID NO: 63 and SEQ ID NO:79. In some embodiments, the antibody or antigen-binding fragment thereof comprises polypeptide sequences that are at least 90% identical, at least 95% identical, at least 99% identical, or identical to the polypeptide sequences of SEQ ID NO: 65 and SEQ ID NO:81.

[0021] In some embodiments, the antibody or antigen binding fragment thereof is murine, non-human, humanized, chimeric, resurfaced, or human.

[0022] In some embodiments, the antibody or antibody fragment is capable of inducing apoptosis of a cell expressing CD37 *in vitro* in the absence of cross-linking agents. In some embodiments, the antibody

or antigen binding fragment is capable of inducing complement dependent cytotoxicity (CDC). In some embodiments, the antibody is capable of inducing antibody dependent cell mediated cytotoxicity (ADCC).

[0023] In certain embodiments, the present disclosure provides a method for depleting a B-cell comprising contacting a B-cell (e.g., in a population of cells comprising a non-cancerous B-cell) with a human or humanized antibody or antigen binding fragment thereof that specifically binds to CD37, wherein the antibody or fragment thereof is capable of inducing apoptosis of a cell expressing CD37 *in vitro* in the absence of cross-linking agents. In certain embodiments, the present disclosure provides a method for treating a patient having an autoimmune or inflammatory disease comprising administering to the patient a therapeutically effective amount of a human or humanized antibody or antigen binding fragment thereof that specifically binds to CD37, wherein the antibody or fragment thereof is capable of inducing apoptosis of a cell expressing CD37 *in vitro* in the absence of cross-linking agents. In some embodiments, the human or humanized antibody or antigen binding fragment thereof is also capable of inducing complement dependent cytotoxicity (CDC). In some embodiments, the human or humanized antibody or antigen binding fragment thereof is also capable of inducing antibody dependent cell mediated cytotoxicity (ADCC).

[0024] In some embodiments, the antibody or antigen-binding fragment binds to human CD37 and macaque CD37.

[0025] In some embodiments, the antibody is a full length antibody. In some embodiments, an antigen-binding fragment is used. In some embodiments, the antibody or antigen-binding fragment thereof comprises a Fab, Fab', F(ab')₂, Fd, single chain Fv or scFv, disulfide linked Fv, V-NAR domain, IgNar, intrabody, IgGACH2, minibody, F(ab')₃, tetrabody, triabody, diabody, single-domain antibody, DVD-Ig, Fcab, mAb2, (scFv)₂, or scFv-Fc.

[0026] In some embodiments, the antibody or antigen-binding fragment thereof is linked via a linker (L) to a cytotoxic agent (C) to form an immunoconjugate.

[0027] In certain embodiments, the present disclosure provides a method for depleting a B-cell comprising contacting a B-cell (e.g., in a population of cells comprising a non-cancerous B-cell) with a composition comprising an immunoconjugate having the formula (A) - (L) - (C), wherein: (A) is an antibody or antigen binding fragment that specifically binds to CD37; (L) is a non-cleavable linker; and (C) is a cytotoxic agent; and wherein the linker (L) links (A) to (C). In certain embodiments, the present disclosure provides a method for treating a patient having an autoimmune or inflammatory disease comprising administering to the patient a therapeutically effective amount of a composition comprising an immunoconjugate having the formula (A) - (L) - (C), wherein: (A) is an antibody or antigen binding fragment that specifically binds to CD37; (L) is a non-cleavable linker; and (C) is a cytotoxic agent; and wherein the linker (L) links (A) to (C). In some embodiments, the immunoconjugate has a serum half-life that is comparable to that of the naked antibody.

[0028] In certain embodiments, the present disclosure provides a method for depleting a B-cell comprising contacting a B-cell (e.g., in a population of cells comprising a non-cancerous B-cell) with a composition comprising an immunoconjugate having the formula (A) - (L) - (C), wherein: (A) is an antibody or antigen binding fragment that specifically binds to CD37; (L) is a linker; and (C) is a maytansinoid; and wherein the linker (L) links (A) to (C). In certain embodiments, the present disclosure provides a method for treating a patient having an autoimmune or inflammatory disease comprising administering to the patient a therapeutically effective amount of a composition comprising an immunoconjugate having the formula (A) - (L) - (C), wherein: (A) is an antibody or antigen binding fragment that specifically binds to CD37; (L) is a linker; and (C) is a maytansinoid; and wherein the linker (L) links (A) to (C).

[0029] In some embodiments, the linker is a non-cleavable linker. In some embodiments, the immunoconjugate further comprises a second (C). In some embodiments, the immunoconjugate further comprises a third (C). In some embodiments, the immunoconjugate further comprises a fourth (C). In some embodiments, the immunoconjugate comprises 2-6 (C). In some embodiments, the immunoconjugate comprises 3-4 (C).

[0030] In some embodiments, the linker is selected from the group consisting of a cleavable linker, a non-cleavable linker, a hydrophilic linker, and a dicarboxylic acid based linker. In some embodiments, the linker is selected from the group consisting of: N-succinimidyl 4-(2-pyridyldithio)pentanoate (SPP); N-succinimidyl 4-(2-pyridyldithio)butanoate (SPDB) or N-succinimidyl 4-(2-pyridyldithio)-2-sulfobutanoate (sulfo-SPDB); N-succinimidyl 4-(maleimidomethyl) cyclohexanecarboxylate (SMCC); N-sulfosuccinimidyl 4-(maleimidomethyl) cyclohexanecarboxylate (sulfoSMCC); N-succinimidyl-4-(iodoacetyl)-aminobenzoate (SIAB); and N-succinimidyl-[(N-maleimidopropionamido)-tetraethyleneglycol] ester (NHS-PEG4-maleimide). In some embodiments, the linker is N-succinimidyl-[(N-maleimidopropionamido)-tetraethyleneglycol] ester (NHS-PEG4-maleimide).

[0031] In some embodiments, the cytotoxic agent is selected from the group consisting of a maytansinoid, maytansinoid analog, doxorubicin, a modified doxorubicin, benzodiazepine, taxoid, CC-1065, CC-1065 analog, duocarmycin, duocarmycin analog, calicheamicin, dolastatin, dolastatin analog, aristatin, tomaymycin derivative, and leptomyacin derivative or a prodrug of the agent. In some embodiments, the cytotoxic agent is a maytansinoid. In some embodiments, the cytotoxic agent is N(2')-deacetyl-N(2')-(3-mercapto-1-oxopropyl)-maytansine (DM1) or N(2')-deacetyl-N2-(4-mercapto-4-methyl-1-oxopentyl)-maytansine (DM4).

[0032] In some embodiments, the composition comprising an immunoconjugate comprises multiple cytotoxic agents (C) with an average of about 3 to about 4 (C) per (A). In some embodiments, the immunoconjugates have an average of about 3.5 (C) per (A). In some embodiments, the immunoconjugates have an average of about 3.5 ± 0.5 (C) per (A).

[0033] In some embodiments, the composition comprising an immunoconjugate comprises an antibody comprising SEQ ID NO:57 and SEQ ID NO:74 or SEQ ID NO:58 and SEQ ID NO:74, an SMCC linker, and DM1. In some embodiments, the composition comprising an immunoconjugate comprises an antibody comprising SEQ ID NO:63 and SEQ ID NO:79, an SMCC linker, and DM1. In some embodiments, the composition comprising an immunoconjugate comprises an antibody comprising SEQ ID NO:65 and SEQ ID NO:81, an SMCC linker, and DM1.

[0034] In some embodiments, the antibody or antigen-binding fragment is capable of depleting B-cells. In some embodiments, the antibody or antigen-binding fragment is capable of inhibiting T-cell responses.

[0035] In some embodiments, the B-cell is in a composition further comprising a T-cell. In some embodiments, the B-cell is in a composition comprising peripheral blood mononuclear cells. In some embodiments, the peripheral blood mononuclear cells were obtained from a human. In some embodiments, the B-cell is in whole blood. In some embodiments, the whole blood was obtained from a human. In some embodiments, the B-cell is in an organism. In some embodiments, the B-cell is in a patient having an autoimmune or inflammatory disease.

[0036] In some embodiments, the B-cell is an autoreactive B-cell.

[0037] In some embodiments, at least about 30% of B-cells are depleted. In some embodiments, less than about 5% of T-cells are depleted.

[0038] In some embodiments, a second therapeutic agent is administered. In some embodiments, the second therapeutic is selected from the group consisting of methotrexate, an anti-CD20 therapeutic, an anti-IL-6 receptor therapeutic, an anti-IL-12/23p40 therapeutic, a chemotherapeutic, an immunosuppressant, an anti-Interferon beta-1a therapeutic, glatiramer acetate, an anti- α 4-integrin therapeutic, fingolimod, an anti-BLys therapeutic, CTLA-Fc, or an anti-TNF therapeutic. In some embodiments, the second therapeutic is an antibody directed against an antigen selected from a group consisting of CD3, CD14, CD19, CD20, CD22, CD25, CD28, CD30, CD33, CD36, CD38, CD40, CD44, CD52, CD55, CD59, CD56, CD70, CD79, CD80, CD103, CD134, CD137, CD138, and CD152. In some embodiments, the second therapeutic is an antibody directed against an antigen selected from the group consisting of IL-2, IL-6, IL-12, IL-23, IL-12/23 p40, IL-17, IFN γ , TNF α , IFN α , IL-15, IL-21, IL-1a, IL-1b, IL-18, IL-8, IL-4, GM-CSF, IL-3, and IL-5.

[0039] In some embodiments, the autoimmune or inflammatory disease is selected from the group consisting of rheumatoid arthritis, multiple sclerosis, type I diabetes mellitus, idiopathic inflammatory myopathy, systemic lupus erythematosus (SLE), myasthenia gravis, Grave's disease, dermatomyositis, polymyositis, Crohn's disease, ulcerative colitis, gastritis, Hashimoto's thyroiditis, asthma, psoriasis, psoriatic arthritis, dermatitis, systemic scleroderma and sclerosis, inflammatory bowel disease (IBD), respiratory distress syndrome, meningitis, encephalitis, uveitis, glomerulonephritis, eczema,

atherosclerosis, leukocyte adhesion deficiency, Raynaud's syndrome, Sjögren's syndrome, Reiter's disease, Behcet's disease, immune complex nephritis, IgA nephropathy, IgM polyneuropathies, immune-mediated thrombocytopenias, acute idiopathic thrombocytopenic purpura, chronic idiopathic thrombocytopenic purpura, hemolytic anemia, myasthenia gravis, lupus nephritis, atopic dermatitis, pemphigus vulgaris, opsoclonus-myoclonus syndrome, pure red cell aplasia, mixed cryoglobulinemia, ankylosing spondylitis, hepatitis C-associated cryoglobulinemic vasculitis, chronic focal encephalitis, bullous pemphigoid, hemophilia A, membranoproliferative glomerulonephritis, adult and juvenile dermatomyositis, adult polymyositis, chronic urticaria, primary biliary cirrhosis, neuromyelitis optica, Graves' dysthyroid disease, bullous pemphigoid, membranoproliferative glomerulonephritis, Churg-Strauss syndrome, juvenile onset diabetes, hemolytic anemia, atopic dermatitis, systemic sclerosis, Sjögren's syndrome and glomerulonephritis, dermatomyositis, ANCA, aplastic anemia, autoimmune hemolytic anemia (AIHA), factor VIII deficiency, hemophilia A, autoimmune neutropenia, Castleman's syndrome, Goodpasture's syndrome, solid organ transplant rejection, graft versus host disease (GVHD), autoimmune hepatitis, lymphoid interstitial pneumonitis, HIV, bronchiolitis obliterans (non-transplant), Guillain-Barre Syndrome, large vessel vasculitis, giant cell (Takayasu's) arteritis, medium vessel vasculitis, Kawasaki's Disease, polyarteritis nodosa, Wegener's granulomatosis, microscopic polyangiitis (MPA), Omenn's syndrome, chronic renal failure, acute infectious mononucleosis, HIV and herpes virus associated diseases.

BRIEF DESCRIPTION OF THE DRAWINGS/FIGURES

[0040] Figure 1 depicts an FL2-H (PE) histogram overlay for a flow cytometry experiment with human B-cells. The following conditions are shown: antibody control (dark filled), isotype control stain (light filled), anti-CD37 stain (thick black line), and anti-CD20 stain (dashed line) for CD19+ B-cells.

[0041] Figure 2 depicts the results of *in vitro* depletion experiments using purified human PBMC samples treated with 10 µg/mL of huCD37-3, huCD37-3-SMCC-DM1, huCD37-50, huCD37-50-SMCC-DM1, rituximab, TRU-016, or alemtuzumab. Results from two different donors are shown in panel A and B.

[0042] Figure 3 depicts the results of *in vitro* depletion experiments using purified human PBMC samples treated with varying concentrations of huCD37-3-SMCC-DM1. Results from two different donors are shown in panels A and B. Figure 3 (C) shows the results using huCD37-3, huCD37-38, huCD37-50 and huCD37-56.

[0043] Figure 4 depicts the results of *in vitro* depletion experiments using unpurified whole human blood samples treated with 10 µg/mL of huCD37-3, huCD37-3-SMCC-DM1, huCD37-50, huCD37-50-SMCC-DM1, rituximab, TRU-016, or alemtuzumab.

[0044] Figure 5 depicts the results of *in vitro* depletion experiments using unpurified whole human blood samples treated with varying concentrations of (A) huCD37-3, huCD37-3-SMCC-DM1, and rituximab and (B) huCD37-3, huCD37-3-SMCC-DM1, huCD37-50, and rituximab.

[0045] Figure 6 depicts release of IFN- γ (Interferon), TNF- α (Tumor Necrosis Factor) and IL-6 (Interleukin-6) measured by ELISpot as number of spots per 5×10^5 peripheral blood mononuclear cells (PBMCs) from one healthy human donor incubated for 18-20 hours with compounds at a concentration of 2.5 ng/mL to 250 μ g/mL.

[0046] Figure 7 depicts release of IFN- γ (Interferon), TNF- α (Tumor Necrosis Factor) and IL-6 (Interleukin-6) measured by ELISpot as number of spots per 5×10^5 peripheral blood mononuclear cells (PBMCs) from a second healthy human donor incubated for 18-20 hours with compounds at a concentration of 2.5 ng/mL to 250 μ g/mL.

[0047] Figure 8 depicts the binding curve of anti-muCD37 monoclonal antibody clone 252-3.

[0048] Figure 9 shows the activity of the 252-3 antibody in depleting peripheral blood B cells (A) and in inhibiting EAE (B) in C57Bl/6 mice. In (A), each symbol represent one mouse; to compare the B cell level in control vs. experimental mice, B cell level was normalized with T cell level and ratio of B/T cell in control mice was considered 100%. In (B), open and closed symbols represent mean of EAE score in control group (n=10) and 252-3 antibody treated group (n=10), respectively; arrow indicates day of antibody injection.

[0049] Figure 10 shows the activity of the 252-3 antibody in depleting peripheral blood B cells (A) and in inhibiting T1D (B) in NOD mice. In (A), each symbol represent one mouse; to compare the B cell level in control vs. experimental mice, B cell level was normalized with T cell level and ratio of B/T cell in control mice was considered 100%. In (B), open and closed symbols represent the diabetes incidence in control group (n=6) and 252-3 antibody treated group (n=6), respectively.

[0050] Figure 11 shows the activity of the 252-3 antibody in depleting peripheral blood B cells (A) and in inhibiting CIA (B) in DBA/1 mice. In (A), each symbol represent one mouse; to compare the B cell level in control vs. experimental mice, B cell level was normalized with T cell level and ratio of B/T cell in control mice was considered 100%. In (B), open and closed symbols represents mean of CIA score in control group (n=12) and 252-3 antibody treated group (n=12), respectively; arrow indicates day of antibody injection.

DETAILED DESCRIPTION OF THE INVENTION

[0051] The present invention provides methods of depleting B-cells and of treating diseases associated with aberrant B-cell activity and/or aberrant T-cell stimulation in connection with a B-cell pathway using CD37 binding molecules.

I. Definitions

[0052] To facilitate an understanding of the present invention, a number of terms and phrases are defined below.

[0053] The term CD37 as used herein, refers to any native CD37, unless otherwise indicated. CD37 is also referred to as GP52-40, leukocyte antigen CD37, and Tetraspanin-26. The term "CD37" encompasses "full-length," unprocessed CD37 as well as any form of CD37 that results from processing in the cell. The term also encompasses naturally occurring variants of CD37, e.g., splice variants, allelic variants, and isoforms. The CD37 polypeptides described herein can be isolated from a variety of sources, such as from human tissue types or from another source, or prepared by recombinant or synthetic methods.

[0054] The term "antibody" means an immunoglobulin molecule that recognizes and specifically binds to a target, such as a protein, polypeptide, peptide, carbohydrate, polynucleotide, lipid, or combinations of the foregoing through at least one antigen recognition site within the variable region of the immunoglobulin molecule. As used herein, the term "antibody" encompasses intact polyclonal antibodies, intact monoclonal antibodies, antibody fragments (such as Fab, Fab', F(ab')₂, and Fv fragments), single chain Fv (scFv) mutants, multispecific antibodies such as bispecific antibodies generated from at least two intact antibodies, chimeric antibodies, humanized antibodies, human antibodies, fusion proteins comprising an antigen determination portion of an antibody, and any other modified immunoglobulin molecule comprising an antigen recognition site so long as the antibodies exhibit the desired biological activity. An antibody can be of any the five major classes of immunoglobulins: IgA, IgD, IgE, IgG, and IgM, or subclasses (isotypes) thereof (e.g. IgG1, IgG2, IgG3, IgG4, IgA1 and IgA2), based on the identity of their heavy-chain constant domains referred to as alpha, delta, epsilon, gamma, and mu, respectively. The different classes of immunoglobulins have different and well known subunit structures and three-dimensional configurations. Antibodies can be naked or conjugated to other molecules such as toxins, radioisotopes, etc.

[0055] A "blocking" antibody or an "antagonist" antibody is one which inhibits or reduces biological activity of the antigen it binds, such as CD37. In some embodiments, blocking antibodies or antagonist antibodies substantially or completely inhibit the biological activity of the antigen. The biological activity can be reduced by 10%, 20%, 30%, 50%, 70%, 80%, 90%, 95%, or even 100%.

[0056] The term "anti-CD37 antibody" or "an antibody that binds to CD37" refers to an antibody that is capable of binding CD37 with sufficient affinity such that the antibody is useful as a diagnostic and/or therapeutic agent in targeting CD37. The extent of binding of an anti-CD37 antibody to an unrelated, non-CD37 protein can be less than about 10% of the binding of the antibody to CD37 as measured, e.g., by a radioimmunoassay (RIA). In certain embodiments, an antibody that binds to CD37 has a dissociation constant (K_d) of $\leq 1 \mu\text{M}$, $\leq 100 \text{ nM}$, $\leq 10 \text{ nM}$, $\leq 1 \text{ nM}$, or $\leq 0.1 \text{ nM}$.

[0057] The term "antibody fragment" refers to a portion of an intact antibody and refers to the antigenic determining variable regions of an intact antibody. Examples of antibody fragments include, but are not limited to Fab, Fab', F(ab')₂, and Fv fragments, linear antibodies, single chain antibodies, and multispecific antibodies formed from antibody fragments.

[0058] A "monoclonal antibody" refers to a homogeneous antibody population involved in the highly specific recognition and binding of a single antigenic determinant, or epitope. This is in contrast to polyclonal antibodies that typically include different antibodies directed against different antigenic determinants. The term "monoclonal antibody" encompasses both intact and full-length monoclonal antibodies as well as antibody fragments (such as Fab, Fab', F(ab')₂, Fv), single chain (scFv) mutants, fusion proteins comprising an antibody portion, and any other modified immunoglobulin molecule comprising an antigen recognition site. Furthermore, "monoclonal antibody" refers to such antibodies made in any number of manners including but not limited to by hybridoma, phage selection, recombinant expression, and transgenic animals.

[0059] The term "humanized antibody" refers to forms of non-human (e.g. murine) antibodies that are specific immunoglobulin chains, chimeric immunoglobulins, or fragments thereof that contain minimal non-human (e.g., murine) sequences. Typically, humanized antibodies are human immunoglobulins in which residues from the complementary determining region (CDR) are replaced by residues from the CDR of a non-human species (e.g. mouse, rat, rabbit, hamster) that have the desired specificity, affinity, and capability (Jones et al., 1986, *Nature*, 321:522-525; Riechmann et al., 1988, *Nature*, 332:323-327; Verhoeyen et al., 1988, *Science*, 239:1534-1536). In some instances, the Fv framework region (FR) residues of a human immunoglobulin are replaced with the corresponding residues in an antibody from a non-human species that has the desired specificity, affinity, and capability. The humanized antibody can be further modified by the substitution of additional residues either in the Fv framework region and/or within the replaced non-human residues to refine and optimize antibody specificity, affinity, and/or capability. In general, the humanized antibody will comprise substantially all of at least one, and typically two or three, variable domains containing all or substantially all of the CDR regions that correspond to the non-human immunoglobulin whereas all or substantially all of the FR regions are those of a human immunoglobulin consensus sequence. The humanized antibody can also comprise at least a portion of an immunoglobulin constant region or domain (Fc), typically that of a human immunoglobulin. Examples of methods used to generate humanized antibodies are described in U.S. Pat. 5,225,539.

[0060] A "variable region" of an antibody refers to the variable region of the antibody light chain or the variable region of the antibody heavy chain, either alone or in combination. The variable regions of the heavy and light chain each consist of four framework regions (FR) connected by three complementarity determining regions (CDRs) also known as hypervariable regions. The CDRs in each

chain are held together in close proximity by the FRs and, with the CDRs from the other chain, contribute to the formation of the antigen-binding site of antibodies. There are at least two techniques for determining CDRs: (1) an approach based on cross-species sequence variability (i.e., Kabat et al. Sequences of Proteins of Immunological Interest, (5th ed., 1991, National Institutes of Health, Bethesda Md.)); and (2) an approach based on crystallographic studies of antigen-antibody complexes (Al-lazikani et al (1997) *J. Molec. Biol.* 273:927-948)). In addition, combinations of these two approaches are sometimes used in the art to determine CDRs.

[0061] The Kabat numbering system is generally used when referring to a residue in the variable domain (approximately residues 1-107 of the light chain and residues 1-113 of the heavy chain) (e.g., Kabat et al., Sequences of Immunological Interest. 5th Ed. Public Health Service, National Institutes of Health, Bethesda, Md. (1991)).

[0062] The amino acid position numbering as in Kabat, refers to the numbering system used for heavy chain variable domains or light chain variable domains of the compilation of antibodies in Kabat et al., Sequences of Proteins of Immunological Interest, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, Md. (1991). Using this numbering system, the actual linear amino acid sequence can contain fewer or additional amino acids corresponding to a shortening of, or insertion into, a FR or CDR of the variable domain. For example, a heavy chain variable domain can include a single amino acid insert (residue 52a according to Kabat) after residue 52 of H2 and inserted residues (e.g. residues 82a, 82b, and 82c, etc according to Kabat) after heavy chain FR residue 82. The Kabat numbering of residues can be determined for a given antibody by alignment at regions of homology of the sequence of the antibody with a "standard" Kabat numbered sequence. Chothia refers instead to the location of the structural loops (Chothia and Lesk *J. Mol. Biol.* 196:901-917 (1987)). The end of the Chothia CDR-H1 loop when numbered using the Kabat numbering convention varies between H32 and H34 depending on the length of the loop (this is because the Kabat numbering scheme places the insertions at H35A and H35B; if neither 35A nor 35B is present, the loop ends at 32; if only 35A is present, the loop ends at 33; if both 35A and 35B are present, the loop ends at 34). The AbM hypervariable regions represent a compromise between the Kabat CDRs and Chothia structural loops, and are used by Oxford Molecular's AbM antibody modeling software.

Loop	Kabat	AbM	Chothia
L1	L24-L34	L24-L34	L24-L34
L2	L50-L56	L50-L56	L50-L56
L3	L89-L97	L89-L97	L89-L97
H1	H31-H35B	H26-H35B (Kabat Numbering)	H26-H32..34
H1	H31-H35	H26-H35 (Chothia Numbering)	H26-H32
H2	H50-H65	H50-H58	H52-H56
H3	H95-H102	H95-H102	H95-H102

[0063] The term "human antibody" means an antibody produced by a human or an antibody having an amino acid sequence corresponding to an antibody produced by a human made using any technique known in the art. This definition of a human antibody includes intact or full-length antibodies, fragments thereof, and/or antibodies comprising at least one human heavy and/or light chain polypeptide such as, for example, an antibody comprising murine light chain and human heavy chain polypeptides.

[0064] The term "chimeric antibodies" refers to antibodies wherein the amino acid sequence of the immunoglobulin molecule is derived from two or more species. Typically, the variable region of both light and heavy chains corresponds to the variable region of antibodies derived from one species of mammals (e.g. mouse, rat, rabbit, etc) with the desired specificity, affinity, and capability while the constant regions are homologous to the sequences in antibodies derived from another (usually human) to avoid eliciting an immune response in that species.

[0065] The term "epitope" or "antigenic determinant" are used interchangeably herein and refer to that portion of an antigen capable of being recognized and specifically bound by a particular antibody. When the antigen is a polypeptide, epitopes can be formed both from contiguous amino acids and noncontiguous amino acids juxtaposed by tertiary folding of a protein. Epitopes formed from contiguous amino acids are typically retained upon protein denaturing, whereas epitopes formed by tertiary folding are typically lost upon protein denaturing. An epitope typically includes at least 3, and more usually, at least 5 or 8-10 amino acids in a unique spatial conformation.

[0066] "Binding affinity" generally refers to the strength of the sum total of noncovalent interactions between a single binding site of a molecule (e.g., an antibody) and its binding partner (e.g., an antigen). Unless indicated otherwise, as used herein, "binding affinity" refers to intrinsic binding affinity which reflects a 1:1 interaction between members of a binding pair (e.g., antibody and antigen). The affinity of a molecule X for its partner Y can generally be represented by the dissociation constant (K_d). Affinity can be measured by common methods known in the art, including those described herein. Low-affinity antibodies generally bind antigen slowly and tend to dissociate readily, whereas high-affinity antibodies

generally bind antigen faster and tend to remain bound longer. A variety of methods of measuring binding affinity are known in the art, any of which can be used for purposes of the present invention. Specific illustrative embodiments are described in the following.

[0067] "Or better" when used herein to refer to binding affinity refers to a stronger binding between a molecule and its binding partner. "Or better" when used herein refers to a stronger binding, represented by a smaller numerical Kd value. For example, an antibody which has an affinity for an antigen of "0.6 nM or better", the antibody's affinity for the antigen is <0.6 nM, i.e. 0.59 nM, 0.58 nM, 0.57 nM etc. or any value less than 0.6 nM.

[0068] By "specifically binds," it is generally meant that an antibody binds to an epitope via its antigen binding domain, and that the binding entails some complementarity between the antigen binding domain and the epitope. According to this definition, an antibody is said to "specifically bind" to an epitope when it binds to that epitope, via its antigen binding domain more readily than it would bind to a random, unrelated epitope. The term "specificity" is used herein to qualify the relative affinity by which a certain antibody binds to a certain epitope. For example, antibody "A" may be deemed to have a higher specificity for a given epitope than antibody "B," or antibody "A" may be said to bind to epitope "C" with a higher specificity than it has for related epitope "D."

[0069] By "preferentially binds," it is meant that the antibody specifically binds to an epitope more readily than it would bind to a related, similar, homologous, or analogous epitope. Thus, an antibody which "preferentially binds" to a given epitope would more likely bind to that epitope than to a related epitope, even though such an antibody may cross-react with the related epitope.

[0070] An antibody is said to "competitively inhibit" binding of a reference antibody to a given epitope if it preferentially binds to that epitope to the extent that it blocks, to some degree, binding of the reference antibody to the epitope. Competitive inhibition may be determined by any method known in the art, for example, competition ELISA assays. An antibody may be said to competitively inhibit binding of the reference antibody to a given epitope by at least 90%, at least 80%, at least 70%, at least 60%, or at least 50%.

[0071] The phrase "substantially similar," or "substantially the same", as used herein, denotes a sufficiently high degree of similarity between two numeric values (generally one associated with an antibody of the invention and the other associated with a reference/comparator antibody) such that one of skill in the art would consider the difference between the two values to be of little or no biological and/or statistical significance within the context of the biological characteristic measured by said values (e.g., Kd values). The difference between said two values can be less than about 50%, less than about 40%, less than about 30%, less than about 20%, or less than about 10% as a function of the value for the reference/comparator antibody.

[0072] A polypeptide, antibody, polynucleotide, vector, cell, or composition which is "isolated" is a polypeptide, antibody, polynucleotide, vector, cell, or composition which is in a form not found in nature. Isolated polypeptides, antibodies, polynucleotides, vectors, cell or compositions include those which have been purified to a degree that they are no longer in a form in which they are found in nature. In some embodiments, an antibody, polynucleotide, vector, cell, or composition which is isolated is substantially pure.

[0073] As used herein, "substantially pure" refers to material which is at least 50% pure (i.e., free from contaminants), at least 90% pure, at least 95% pure, at least 98% pure, or at least 99% pure.

[0074] The term "immunoconjugate" or "conjugate" as used herein refers to a compound or a derivative thereof that is linked to a cell binding agent (i.e., an anti-CD37 antibody or fragment thereof) and is defined by a generic formula: C-L-A, wherein C = cytotoxin, L = linker, and A = cell binding agent or anti-CD37 antibody or antibody fragment. Immunoconjugates can also be defined by the generic formula in reverse order: A-L-C.

[0075] A "linker" is any chemical moiety that is capable of linking a compound, usually a drug, such as a maytansinoid, to a cell-binding agent such as an anti CD37 antibody or a fragment thereof in a stable, covalent manner. Linkers can be susceptible to or be substantially resistant to acid-induced cleavage, light-induced cleavage, peptidase-induced cleavage, esterase-induced cleavage, and disulfide bond cleavage, at conditions under which the compound or the antibody remains active. Suitable linkers are well known in the art and include, for example, disulfide groups, thioether groups, acid labile groups, photolabile groups, peptidase labile groups and esterase labile groups. Linkers also include charged linkers, and hydrophilic forms thereof as described herein and known in the art.

[0076] The terms "cancer" and "cancerous" refer to or describe the physiological condition in mammals in which a population of cells are characterized by unregulated cell growth. Examples of cancer include, but are not limited to, carcinoma, lymphoma, blastoma, sarcoma, and leukemia. "Tumor" and "neoplasm" refer to one or more cells that result from excessive cell growth or proliferation, either benign (noncancerous) or malignant (cancerous) including pre-cancerous lesions. Examples of "cancer" or "tumorigenic" diseases which can be treated and/or prevented include B-cell lymphomas including NHL, precursor B-cell lymphoblastic leukemia/lymphoma and mature B-cell neoplasms, such as B-cell chronic lymphocytic leukemia (CLL)/small lymphocytic lymphoma (SLL), B-cell prolymphocytic leukemia, lymphoplasmacytic lymphoma, mantle cell lymphoma (MCL), follicular lymphoma (FL), including low-grade, intermediate-grade and high-grade FL, cutaneous follicle center lymphoma, marginal zone B-cell lymphoma (MALT type, nodal and splenic type), hairy cell leukemia, diffuse large B-cell lymphoma, Burkitt's lymphoma, plasmacytoma, plasma cell myeloma, post-transplant lymphoproliferative disorder, and anaplastic large-cell lymphoma (ALCL). Non-cancerous cells are cells that do not result in the

formation of tumors or neoplasms or the development of cancer. However, non-cancerous cells can contribute to disease, e.g., autoimmune diseases, and include, for example auto-reactive B-cells.

[0077] The terms "cancer cell," "tumor cell," and grammatical equivalents refer to the total population of cells derived from a tumor or a pre-cancerous lesion, including both non-tumorigenic cells, which comprise the bulk of the tumor cell population, and tumorigenic stem cells (cancer stem cells). As used herein, the term "tumor cell" will be modified by the term "non-tumorigenic" when referring solely to those tumor cells lacking the capacity to renew and differentiate to distinguish those tumor cells from cancer stem cells.

[0078] The term "autoreactive" refers to a cell, tissue, protein, antibody or other substance that produces an immune response directed against an organism's own cells, tissues, proteins, antibodies, or other substances.

[0079] The term "subject" refers to any animal (e.g., a mammal), including, but not limited to humans, non-human primates, rodents, and the like, which is to be the recipient of a particular treatment. Typically, the terms "subject" and "patient" are used interchangeably herein in reference to a human subject.

[0080] Administration "in combination with" one or more further therapeutic agents includes simultaneous (concurrent) and consecutive administration in any order.

[0081] The term "pharmaceutical formulation" refers to a preparation which is in such form as to permit the biological activity of the active ingredient to be effective, and which contains no additional components which are unacceptably toxic to a subject to which the formulation would be administered. The formulation can be sterile.

[0082] An "effective amount" of an antibody as disclosed herein is an amount sufficient to carry out a specifically stated purpose. An "effective amount" can be determined empirically and in a routine manner, in relation to the stated purpose.

[0083] The term "therapeutically effective amount" refers to an amount of an antibody or other drug effective to "treat" a disease or disorder in a subject or mammal. In some embodiments, the therapeutically effective amount of the drug can reduce the number of B-cells; reduce the number of autoreactive B-cells; decrease the symptoms of disease; or slow the progression of disease. See the definition herein of "treating". A "prophylactically effective amount" refers to an amount effective, at dosages and for periods of time necessary, to achieve the desired prophylactic result. Typically but not necessarily, since a prophylactic dose is used in subjects prior to or at an earlier stage of disease, the prophylactically effective amount will be less than the therapeutically effective amount.

[0084] The word "label" when used herein refers to a detectable compound or composition which is conjugated directly or indirectly to the antibody so as to generate a "labeled" antibody. The label can be

detectable by itself (e.g., radioisotope labels or fluorescent labels) or, in the case of an enzymatic label, can catalyze chemical alteration of a substrate compound or composition which is detectable.

[0085] Terms such as "treating" or "treatment" or "to treat" or "alleviating" or "to alleviate" refer to therapeutic measures that cure, slow down, lessen symptoms of, and/or halt progression of a diagnosed pathologic condition or disorder. Thus, those in need of treatment include those already diagnosed with or suspected of having the disorder. Prophylactic or preventative measures refer to therapeutic measures that prevent and/or slow the development of a targeted pathologic condition or disorder. Thus, those in need of prophylactic or preventative measures include those prone to have the disorder and those in whom the disorder is to be prevented. In certain embodiments, a subject is successfully "treated" if the patient shows one or more of the following: decreased B-cells; decreased autoreactive B-cells; decreased B-cell activity; decreased aberrant B-cell activity; decreased non-malignant B-cells, decreased non-cancerous B-cells, reduced immunoglobulin level; reduced morbidity and mortality; improvement in quality of life; or some combination of effects.

[0086] "Polynucleotide," or "nucleic acid," as used interchangeably herein, refer to polymers of nucleotides of any length, and include DNA and RNA. The nucleotides can be deoxyribonucleotides, ribonucleotides, modified nucleotides or bases, and/or their analogs, or any substrate that can be incorporated into a polymer by DNA or RNA polymerase. A polynucleotide can comprise modified nucleotides, such as methylated nucleotides and their analogs. If present, modification to the nucleotide structure can be imparted before or after assembly of the polymer. The sequence of nucleotides can be interrupted by non-nucleotide components. A polynucleotide can be further modified after polymerization, such as by conjugation with a labeling component. Other types of modifications include, for example, "caps", substitution of one or more of the naturally occurring nucleotides with an analog, internucleotide modifications such as, for example, those with uncharged linkages (e.g., methyl phosphonates, phosphotriesters, phosphoamidates, cabamates, etc.) and with charged linkages (e.g., phosphorothioates, phosphorodithioates, etc.), those containing pendant moieties, such as, for example, proteins (e.g., nucleases, toxins, antibodies, signal peptides, poly-L-lysine, etc.), those with intercalators (e.g., acridine, psoralen, etc.), those containing chelators (e.g., metals, radioactive metals, boron, oxidative metals, etc.), those containing alkylators, those with modified linkages (e.g., alpha anomeric nucleic acids, etc.), as well as unmodified forms of the polynucleotide(s). Further, any of the hydroxyl groups ordinarily present in the sugars can be replaced, for example, by phosphonate groups, phosphate groups, protected by standard protecting groups, or activated to prepare additional linkages to additional nucleotides, or can be conjugated to solid supports. The 5' and 3' terminal OH can be phosphorylated or substituted with amines or organic capping group moieties of from 1 to 20 carbon atoms. Other hydroxyls can also be derivatized to standard protecting groups. Polynucleotides can also contain analogous forms of ribose or deoxyribose sugars that are generally known in the art, including, for example, 2'-O-methyl-, 2'-O-allyl,

2'-fluoro- or 2'-azido-ribose, carbocyclic sugar analogs, .alpha.-anomeric sugars, epimeric sugars such as arabinose, xyloses or lyxoses, pyranose sugars, furanose sugars, sedoheptuloses, acyclic analogs and abasic nucleoside analogs such as methyl riboside. One or more phosphodiester linkages can be replaced by alternative linking groups. These alternative linking groups include, but are not limited to, embodiments wherein phosphate is replaced by P(O)S ("thioate"), P(S)S ("dithioate"), "(O)NR₂" ("amidate"), P(O)R, P(O)OR', CO or CH₂ ("formacetal"), in which each R or R' is independently H or substituted or unsubstituted alkyl (1-20 C) optionally containing an ether (--O--) linkage, aryl, alkenyl, cycloalkyl, cycloalkenyl or araldyl. Not all linkages in a polynucleotide need be identical. The preceding description applies to all polynucleotides referred to herein, including RNA and DNA.

[0087] The term "vector" means a construct, which is capable of delivering, and optionally expressing, one or more gene(s) or sequence(s) of interest in a host cell. Examples of vectors include, but are not limited to, viral vectors, naked DNA or RNA expression vectors, plasmid, cosmid or phage vectors, DNA or RNA expression vectors associated with cationic condensing agents, DNA or RNA expression vectors encapsulated in liposomes, and certain eukaryotic cells, such as producer cells..

[0088] The terms "polypeptide," "peptide," and "protein" are used interchangeably herein to refer to polymers of amino acids of any length. The polymer can be linear or branched, it can comprise modified amino acids, and it can be interrupted by non-amino acids. The terms also encompass an amino acid polymer that has been modified naturally or by intervention; for example, disulfide bond formation, glycosylation, lipidation, acetylation, phosphorylation, or any other manipulation or modification, such as conjugation with a labeling component. Also included within the definition are, for example, polypeptides containing one or more analogs of an amino acid (including, for example, unnatural amino acids, etc.), as well as other modifications known in the art. It is understood that, because the polypeptides of this invention are based upon antibodies, in certain embodiments, the polypeptides can occur as single chains or associated chains.

[0089] The terms "identical" or percent "identity" in the context of two or more nucleic acids or polypeptides, refer to two or more sequences or subsequences that are the same or have a specified percentage of nucleotides or amino acid residues that are the same, when compared and aligned (introducing gaps, if necessary) for maximum correspondence, not considering any conservative amino acid substitutions as part of the sequence identity. The percent identity can be measured using sequence comparison software or algorithms or by visual inspection. Various algorithms and software are known in the art that can be used to obtain alignments of amino acid or nucleotide sequences. One such non-limiting example of a sequence alignment algorithm is the algorithm described in Karlin et al, 1990, *Proc. Natl. Acad. Sci.*, 87:2264-2268, as modified in Karlin et al., 1993, *Proc. Natl. Acad. Sci.*, 90:5873-5877, and incorporated into the NBLAST and XBLAST programs (Altschul et al., 1991, *Nucleic Acids Res.*, 25:3389-3402). In certain embodiments, Gapped BLAST can be used as described in Altschul et al.,

1997, *Nucleic Acids Res.* 25:3389-3402. BLAST-2, WU-BLAST-2 (Altschul et al., 1996, *Methods in Enzymology*, 266:460-480), ALIGN, ALIGN-2 (Genentech, South San Francisco, California) or Megalign (DNASTAR) are additional publicly available software programs that can be used to align sequences. In certain embodiments, the percent identity between two nucleotide sequences is determined using the GAP program in GCG software (e.g., using a NWSgapdna.CMP matrix and a gap weight of 40, 50, 60, 70, or 90 and a length weight of 1, 2, 3, 4, 5, or 6). In certain alternative embodiments, the GAP program in the GCG software package, which incorporates the algorithm of Needleman and Wunsch (*J. Mol. Biol.* (48):444-453 (1970)) can be used to determine the percent identity between two amino acid sequences (e.g., using either a Blossum 62 matrix or a PAM250 matrix, and a gap weight of 16, 14, 12, 10, 8, 6, or 4 and a length weight of 1, 2, 3, 4, 5). Alternatively, in certain embodiments, the percent identity between nucleotide or amino acid sequences is determined using the algorithm of Myers and Miller (*CABIOS*, 4:11-17 (1989)). For example, the percent identity can be determined using the ALIGN program (version 2.0) and using a PAM120 with residue table, a gap length penalty of 12 and a gap penalty of 4. Appropriate parameters for maximal alignment by particular alignment software can be determined by one skilled in the art. In certain embodiments, the default parameters of the alignment software are used. In certain embodiments, the percentage identity "X" of a first amino acid sequence to a second sequence amino acid is calculated as $100 \times (Y/Z)$, where Y is the number of amino acid residues scored as identical matches in the alignment of the first and second sequences (as aligned by visual inspection or a particular sequence alignment program) and Z is the total number of residues in the second sequence. If the length of a first sequence is longer than the second sequence, the percent identity of the first sequence to the second sequence will be longer than the percent identity of the second sequence to the first sequence.

[0090] As a non-limiting example, whether any particular polynucleotide has a certain percentage sequence identity (e.g., is at least 80% identical, at least 85% identical, at least 90% identical, and in some embodiments, at least 95%, 96%, 97%, 98%, or 99% identical) to a reference sequence can, in certain embodiments, be determined using the Bestfit program (Wisconsin Sequence Analysis Package, Version 8 for Unix, Genetics Computer Group, University Research Park, 575 Science Drive, Madison, WI 53711). Bestfit uses the local homology algorithm of Smith and Waterman, *Advances in Applied Mathematics* 2: 482-489 (1981), to find the best segment of homology between two sequences. When using Bestfit or any other sequence alignment program to determine whether a particular sequence is, for instance, 95% identical to a reference sequence according to the present invention, the parameters are set such that the percentage of identity is calculated over the full length of the reference nucleotide sequence and that gaps in homology of up to 5% of the total number of nucleotides in the reference sequence are allowed.

[0091] In some embodiments, two nucleic acids or polypeptides of the invention are substantially identical, meaning they have at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, and in some embodiments at least 95%, 96%, 97%, 98%, 99% nucleotide or amino acid residue identity, when

compared and aligned for maximum correspondence, as measured using a sequence comparison algorithm or by visual inspection. Identity can exist over a region of the sequences that is at least about 10, about 20, about 40-60 residues in length or any integral value therebetween, and can be over a longer region than 60-80 residues, for example, at least about 90-100 residues, and in some embodiments, the sequences are substantially identical over the full length of the sequences being compared, such as the coding region of a nucleotide sequence for example.

[0092] A "conservative amino acid substitution" is one in which one amino acid residue is replaced with another amino acid residue having a similar side chain. Families of amino acid residues having similar side chains have been defined in the art, including basic side chains (e.g., lysine, arginine, histidine), acidic side chains (e.g., aspartic acid, glutamic acid), uncharged polar side chains (e.g., glycine, asparagine, glutamine, serine, threonine, tyrosine, cysteine), nonpolar side chains (e.g., alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan), beta-branched side chains (e.g., threonine, valine, isoleucine) and aromatic side chains (e.g., tyrosine, phenylalanine, tryptophan, histidine). For example, substitution of a phenylalanine for a tyrosine is a conservative substitution. In some embodiments, conservative substitutions in the sequences of the polypeptides and antibodies of the invention do not abrogate the binding of the polypeptide or antibody containing the amino acid sequence, to the antigen(s), i.e., the CD37 to which the polypeptide or antibody binds. Methods of identifying nucleotide and amino acid conservative substitutions which do not eliminate antigen binding are well-known in the art (see, e.g., Brummell et al., *Biochem.* 32: 1180-1 187 (1993); Kobayashi et al. *Protein Eng.* 12(10):879-884 (1999); and Burks et al. *Proc. Natl. Acad. Sci. USA* 94:412-417 (1997)).

[0093] As used in the present disclosure and claims, the singular forms "a," "an," and "the" include plural forms unless the context clearly dictates otherwise.

[0094] It is understood that wherever embodiments are described herein with the language "comprising," otherwise analogous embodiments described in terms of "consisting of" and/or "consisting essentially of" are also provided.

[0095] The term "and/or" as used in a phrase such as "A and/or B" herein is intended to include both "A and B," "A or B," "A," and "B." Likewise, the term "and/or" as used in a phrase such as "A, B, and/or C" is intended to encompass each of the following embodiments: A, B, and C; A, B, or C; A or C; A or B; B or C; A and C; A and B; B and C; A (alone); B (alone); and C (alone).

II. CD37 binding agents

[0096] The present invention provides agents that specifically bind CD37. These agents are referred to herein as "CD37 binding agents." Exemplary CD37-binding agents have been described in U.S. Published Application No. 2011/0256153, which is herein incorporated by reference in its entirety.

[0097] The full-length amino acid sequences for human, macaca, and murine CD37 are known in the art and also provided herein as represented by SEQ ID NOs:1-3, respectively.

[0098] Human CD37:

[0099] MSAQESCLSLIKYFLFVFNLFVFFVLGSLIFCFGIWILIDKTSFVSVFVGLAFVPLQIWSKVL
AISGIFTMGIALLGCVGALKELRCLLGLYFGMLLLLFATQITLGILISTQRAQLERSLRDVEKTIQ
KYGTNPEETAAEESWDYVQFQLRCCGWHYPQDWFQVLILRNGNSEAHRVPCSCYNLSATNDSTI
LDKVILPQLSRLGHLARSRHADICAVPAESHIYREGCAQGLQKWLHNNLISIVGICLGVGLLELG
FMFLSIFLCRNLDHVYNRLAYR (SEQ ID NO:1)

[00100] Macaca mulatta CD37:

[00101] MSAQESCLSLIKYFLFVFNLFVFFVILGSLIFCFGIWILIDKTSFVSVFVGLAFVPLQIWSKV
LAISGVFTMGLALLGCVGALKELRCLLGLYFGMLLLLFATQITLGILISTQRAQLERSLQDIVEKTI
QRYHTNPEETAAEESWDYVQFQLRCCGWHSPQDWFQVLTLRGNGNSEAHRVPCSCYNLSATNDS
TILDKVILPQLSRLGQLARSRHSTDICAVPANSIYREGCARSLQKWLHNNLISIVGICLGVGLLEL
GFMTLSIFLCRNLDHVYNRLRYR (SEQ ID NO:2)

[00102] Murine CD37 (NP_031671):

[00103] MSAQESCLSLIKYFLFVFNLFVFFVLGGLIFCFGTWILIDKTSFVSVFVGLSFVPLQTWSKV
LAVSGVLTMALALLGCVGALKELRCLLGLYFGMLLLLFATQITLGILISTQVRRLERRVQELVLR
TIQSYRTNPDETAAEESWDYAQFQLRCCGWQSPRDWNKAQMLKANESEEPFVPCSCYNSTATN
DSTVFDKLFQSRLGPRAKLRQTADICALPAKAHIYREGCAQSLQKWLHNNIISIVGICLGVGL
LELGFMTLSIFLCRNLDHVYDRLARYR (SEQ ID NO:3)

[00104] In certain embodiments, the CD37 binding agents are antibodies, immunoconjugates or polypeptides. In some embodiments, the CD37 binding agents are humanized antibodies.

[00105] In certain embodiments, the CD37-binding agents are capable of inducing complement dependent cytotoxicity. Examples of CD37-binding agents that are capable of inducing complement dependent cytotoxicity are disclosed, for example, in U.S. Published Application No. 2011/0256153, which is herein incorporated by reference in its entirety. For example, treatment of cells with the CD37-binding agents can result in CDC activity that reduces cell viability to less than about 80%, less than about 70%, less than about 60%, less than about 50%, less than about 40% or less than about 35% of the cell viability of untreated cells. Treatment of cells with the CD37-binding agents can also result in CDC activity that reduces cell viability to about 70-80%, about 60-70%, about 50-60%, about 40-50%, or about 30-40% of the cell viability of untreated cells. In some particular embodiments, the CD37-binding agents are capable of inducing complement dependent cytotoxicity in Ramos cells.

[00106] In certain embodiments, the CD37-binding agents are capable of inducing antibody dependent cell mediated cytotoxicity (ADCC). Examples of CD-37 binding agents that are capable of inducing antibody dependent cell mediated cytotoxicity (ADCC) are disclosed, for example, in U.S. Published Application No. 2011/0256153, which is herein incorporated by reference in its entirety. For example, treatment of cells with the CD37-binding agents can result in ADCC activity that produces at least about

15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, or at least about 60% cell lysis. Treatment of cells with the CD37-binding agents can result in ADCC activity that produces about 10-20%, about 20-30%, about 30-40%, or about 40-50% cell lysis. Treatment of cells with the CD37-binding agents can also result in ADCC activity that produces about 10-50%, about 20-50%, about 30-50%, or about 40-50% cell lysis. In some particular embodiments, the CD37-binding agents are capable of inducing ADCC in Daudi, Ramos, and/or Granata-519 cells.

[00107] In some embodiments, the CD37-binding agents are capable of inducing apoptosis. In some embodiment, the CD37-binding agents are capable of inducing apoptosis in the absence of cross-linking agents. Examples of CD37-binding agents that are capable of inducing apoptosis *in vitro* in the absence of a cross-linking agent are disclosed, for example, in U.S. Published Application No. 2011/0256153, which is herein incorporated by reference in its entirety. For example, treatment of cells with the CD37-binding agents can induce apoptosis in at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, or at least about 55% of cells. In some particular embodiments, the CD37-binding agents are capable of inducing apoptosis in Ramos cells and/or Raji cells.

[00108] In some embodiments, the CD37-binding agents are capable of depleting B-cells. In some embodiments, the B-cells are autoreactive B-cells. In some embodiments, the B-cells are not cancer cells. In some embodiments, the B-cells are not tumor cells. In some embodiments, the B-cells are not cancerous cells. In some embodiments, the B-cells overexpress CD37. In some embodiments, the B-cells do not overexpress CD37.

[00109] Treatment of cells with CD37-binding agents can result in depletion of at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, or least about 75% of B-cells.

[00110] In some embodiments, the CD37-binding agents do not deplete T-cells under the same conditions in which B-cells are depleted. For example, treatment of cells with CD37-binding agents can result in depletion of less than about 20%, less than about 15%, less than about 10%, or less than about 5% of T-cells. In certain embodiments, the CD37-binding agents deplete at least about 25% of B-cells and deplete less than about 10% of T-cells. In certain embodiments, the CD37-binding agents deplete at least about 30% of B-cells and deplete less than about 5% of T-cells.

[00111] In some embodiments, the CD37-binding agents do not deplete monocytes under the same conditions in which B-cells are depleted. For example, treatment of cells with CD37-binding agents can result in depletion of less than about 20%, less than about 15%, less than about 10%, or less than about 5% of monocytes. In certain embodiments, the CD37-binding agents deplete at least about 25% of B-cells

and deplete less than about 10% of monocytes. In certain embodiments, the CD37-binding agents deplete at least about 30% of B-cells and deplete less than about 5% of monocytes.

[00112] In certain embodiments, immunoconjugates or other agents that specifically bind human CD37 trigger cell death via a cytotoxic agent. For example, in certain embodiments, an antibody to human CD37 is conjugated to a maytansinoid that is activated in cells expressing the CD37 by protein internalization. In certain alternative embodiments, the agent or antibody is not conjugated to a maytansinoid or other cytotoxic molecule.

[00113] The CD37-binding agents include CD37 antibodies such as CD37-3, CD37-12, CD37-38, CD37-50, CD37-51, CD37-56 and CD37-57 and fragments, variants and derivatives thereof. The CD37-binding agents also include CD37-binding agents that specifically bind to the same CD37 epitope as an antibody selected from the group consisting of CD37-3, CD37-12, CD37-38, CD37-50, CD37-51, CD37-56 and CD37-57. The CD37-binding agents also include CD37-binding agents that competitively inhibit an antibody selected from the group consisting of CD37-3, CD37-12, CD37-38, CD37-50, CD37-51, CD37-56 and CD37-57.

[00114] In some particular embodiments, CD37-binding agents can be characterized by their ability to bind chimeric CD37 polypeptides, including murine/human and macaca/human chimeric polypeptides described in U.S. Published Application No. 2011/0256153, which is herein incorporated by reference in its entirety, and provided in the table below.

Chimeric Poly-peptide	Sequence
hCD37-M1	MSAQESCLSLIKYFLFVFNLFVFLGSLIFCFGIWILIDKTSFVSFVGLAFVPLQIWSKV LAISGIFTMGIALLGCVGALKELRCLLGLYFGMLLLLFATQITLGILISTQRVRLERRV QELVLRTIQSYRTNPDETA AEESWDYVQFQLRCCGWHYPQDWFQVLILRGNGSEAH RVPCSCYNLSATNDSTILDKVILPQLSRLGHLARSRHADICAVPAESHYREGCAQGL QKWLHNNLISIVGICLGVGELLELGFMTLSIFLCRNLDHVYNRLARYR (SEQ ID NO:184)
muCD37-R176	ISTQRVRLERRVQELVLRTIQSYRTNPDETA AEESWDYAQFQLRCCGWQSPRDWNK AQMLKANEEPRVPCSCYNSTATNDSTVFDKLFSSQLSRLGPRAKLRQTADICALPA KAHIYREGCAQSLQ (SEQ ID NO:185)
hCD37-M45	MSAQESCLSLIKYFLFVFNLFVFLGSLIFCFGIWILIDKTSFVSFVGLAFVPLQIWSKV LAISGIFTMGIALLGCVGALKELRCLLGLYFGMLLLLFATQITLGILISTQRAQLERSLR DVVEKTIQKYGTNPEETA AEESWDYVQFQLRCCGWHYPQDWFQVLILRGNGSEAH RVPCSCYNLSATNDSTILDKVILPQLSRLGPRAKLRQTADICALPAKAHIYREGCAQS LQKWLHNNLISIVGICLGVGELLELGFMTLSIFLCRNLDHVYNRLARYR (SEQ ID NO:186)
hCD37m ECD-H45	MSAQESCLSLIKYFLFVFNLFVFLGSLIFCFGIWILIDKTSFVSFVGLAFVPLQIWSKV LAISGIFTMGIALLGCVGALKELRCLLGLYFGMLLLLFATQITLGILISTQRVRLERRV QELVLRTIQSYRTNPDETA AEESWDYAQFQLRCCGWQSPRDWNKAQMLKANEEEP RVPCSCYNSTATNDSTVFDKLFSSQLSRLGHLARSRHADICAVPAESHYREGCAQG LQKWLHNNLISIVGICLGVGELLELGFMTLSIFLCRNLDHVYNRLARYR (SEQ ID NO: 187)
hCD37m ECD-H5	MSAQESCLSLIKYFLFVFNLFVFLGSLIFCFGIWILIDKTSFVSFVGLAFVPLQIWSKV LAISGIFTMGIALLGCVGALKELRCLLGLYFGMLLLLFATQITLGILISTQRVRLERRV

	QELVLRTIQSYRTNPDETAEEESWDYAQFQLRCCGWQSPRDWNKAQMLKANEESEP RVPCSCYNSTATNDSTVFDKLFSSQLSRLGPRAKLRQTADICAVPAESHIYREGCAQG LQKWLHNNLISIVGICLGVGLLELGFMTLSIFLCRNLDHVYNRLARYR (SEQ ID NO: 188)
hCD37m ECD-H4	MSAQESCLSLIKYFLFVFNLFVGLSLIFCFGIWILIDKTSFVSFVGLAFVPLQIWSKV LAISGIFTMGIALLGCVGALKELRCLLGLYFGMLLLLFATQITLGIISTQRVRLERRV QELVLRTIQSYRTNPDETAEEESWDYAQFQLRCCGWQSPRDWNKAQMLKANEESEP RVPCSCYNSTATNDSTVFDKLFSSQLSRLGHLARSRHSDICALPAKAHIYREGCAQS LQKWLHNNLISIVGICLGVGLLELGFMTLSIFLCRNLDHVYNRLARYR (SEQ ID NO: 189)
hCD37- Mac4	MSAQESCLSLIKYFLFVFNLFVGLSLIFCFGIWILIDKTSFVSFVGLAFVPLQIWSKV LAISGIFTMGIALLGCVGALKELRCLLGLYFGMLLLLFATQITLGIISTQRAQLERSLR DVVEKTIQKYGTNPEETAEEESWDYVQFQLRCCGWHYPQDWFQVLILRGNGSEAH RVPCSCYNLSATNDSTILDKVILPQLSRLGQLARSRHSTDICAVPAESHIYREGCAQGL QKWLHNNLISIVGICLGVGLLELGFMTLSIFLCRNLDHVYNRLARYR (SEQ ID NO: 190)
hCD37- Mac45	MSAQESCLSLIKYFLFVFNLFVGLSLIFCFGIWILIDKTSFVSFVGLAFVPLQIWSKV LAISGIFTMGIALLGCVGALKELRCLLGLYFGMLLLLFATQITLGIISTQRAQLERSLR DVVEKTIQKYGTNPEETAEEESWDYVQFQLRCCGWHYPQDWFQVLILRGNGSEAH RVPCSCYNLSATNDSTILDKVILPQLSRLGQLARSRHSTDICAVPANSIHYREGCARSL QKWLHNNLISIVGICLGVGLLELGFMTLSIFLCRNLDHVYNRLARYR (SEQ ID NO: 191)
hCD37- Mac5	MSAQESCLSLIKYFLFVFNLFVGLSLIFCFGIWILIDKTSFVSFVGLAFVPLQIWSKV LAISGIFTMGIALLGCVGALKELRCLLGLYFGMLLLLFATQITLGIISTQRAQLERSLR DVVEKTIQKYGTNPEETAEEESWDYVQFQLRCCGWHYPQDWFQVLILRGNGSEAH RVPCSCYNLSATNDSTILDKVILPQLSRLGHLARSRHSDICAVPANSIHYREGCARSL QKWLHNNLISIVGICLGVGLLELGFMTLSIFLCRNLDHVYNRLARYR (SEQ ID NO: 192)

[00115] In some particular embodiments, the binding of the CD37-binding agents to CD37 does not require human CD37 amino acids 109-138. Thus, some CD37-binding agents bind to a polypeptide comprising the amino acid sequence of SEQ ID NO:184. In other embodiments, the binding of the CD37-binding agents to CD37 is disrupted by mutation of human CD37 amino acids 202-243. Thus, some CD37-binding agents do not bind to a polypeptide comprising the amino acid sequence of SEQ ID NO:185.

[00116] In some embodiments, the CD37-binding agents bind to a polypeptide of SEQ ID NO:184 and to a polypeptide of SEQ ID NO:186, but do not bind to a polypeptide of SEQ ID NO:185.

[00117] In some embodiments, the CD37-binding agents bind to a polypeptide of SEQ ID NO:187. In some embodiments, the CD37-binding agents bind to a polypeptide of SEQ ID NO:187 and a polypeptide of SEQ ID NO:188. In some embodiments, the CD37-binding agents bind to a polypeptide of SEQ ID NO:187 and a polypeptide of SEQ ID NO:189.

[00118] In some embodiments, the CD37-binding agent binds to a polypeptide of SEQ ID NO:190, but does not bind to a polypeptide of SEQ ID NO:191. In some embodiments, the CD37-binding agent binds to a polypeptide of SEQ ID NO:192, but does not bind to a polypeptide of SEQ ID NO:191.

[00119] CD37 peptide fragments to which certain CD37-binding agents bind to include, but are not limited to, CD37 fragments comprising, consisting essentially of, or consisting of amino acids 200-243 of SEQ ID NO: 1, amino acids 202-220 or SEQ ID NO:1, or amino acids 221-243 of SEQ ID NO:1. In some embodiments, the CD37-binding agent is specifically binds to a human CD37 epitope comprising amino acids 202-243 of SEQ ID NO:1. In some embodiments, the binding of the CD37-binding agent to CD37 requires amino acids 202-243 of SEQ ID NO:1. In some embodiments, the binding of the CD37-binding agent to CD37 requires amino acids 200-220 of SEQ ID NO:1. In some embodiments, the binding of the CD37-binding agent to CD37 requires amino acids 221-243 of SEQ ID NO:1.

[00120] Examples of CD37-binding agents with the aforementioned binding properties are described in U.S. Published Application No. 2011/0256153, which is herein incorporated by reference in its entirety.

[00121] The CD37-binding agents also include CD37-binding agents that comprise the heavy and light chain CDR sequences of CD37-3, CD37-12, CD37-38, CD37-50, CD37-51, CD37-56 or CD37-57. The heavy and light chain CDRs of CD37-38, CD37-50, CD37-51, CD37-56 and CD37-57 contain related sequences. Therefore, the CD37-binding agents can also comprise heavy and light chain CDR sequences that comprise a consensus sequence obtained by the alignment of CD37-38, CD37-50, CD37-51, CD37-56 and CD37-57. The CDR sequences of CD37-3, CD37-12, CD37-38, CD37-50, CD37-51, CD37-56 and CD37-57, as well as the consensus sequence of CD37-38, CD37-50, CD37-51, CD37-56 and CD37-57 are described in Tables 1 and 2 below.

Table 1: Variable heavy chain CDR amino acid sequences

Antibody	VH-CDR1	VH-CDR2	VH-CDR3
CD37-3	TSGVS (SEQ ID NO:4)	VIWGDGSTN (SEQ ID NO:5)	GGYSLAH (SEQ ID NO:6)
CD37-12	KYGMN (SEQ ID NO:7)	WINTNTGESR (SEQ ID NO:8)	GTVVAD (SEQ ID NO:9)
CD37-38	SGFGWH (SEQ ID NO:10)	YILYSGGTD (SEQ ID NO:11)	GYGYGAWFVY (SEQ ID NO:12)
CD37-50	SGFAWH (SEQ ID NO:13)	YILYSGSTV (SEQ ID NO:14)	GYGYGAWFAY (SEQ ID NO:15)
CD37-51	SGFAWH (SEQ ID NO:16)	YIHYSGSTN (SEQ ID NO:17)	GYGFGAWFVY (SEQ ID NO:18)
CD37-56	SGFAWH (SEQ ID NO:19)	YIHYSGGTN (SEQ ID NO:20)	GYGFGAWFAY (SEQ ID NO:21)
CD37-57	SGFAWH (SEQ ID NO:22)	YILYSGSTV (SEQ ID NO:23)	GYGYGAWFAY (SEQ ID NO:24)
CONSENSUS	SGF[A or G]WH (SEQ ID NO:25)	YI[L or H]YSG[G or S]T[D,V,or N] (SEQ ID NO:26)	GYYG[Y or F]GAWF[V or A]Y (SEQ ID NO:27)
252-3	SYGMS (SEQ ID NO:171)	TISSGGSYTYSPDSVKG (SEQ ID NO:172)	HSYYDTSVDY (SEQ ID NO:173)
252-3	SYGMS (SEQ ID NO:171)	TISSGGSYTY (SEQ ID NO:181)	HSYYDTSVDY (SEQ ID NO:173)

Table 2: Variable light chain CDR amino acid sequences

Antibody	VL-CDR1	VL-CDR2	VL-CDR3
CD37-3	RASENIRSNLA (SEQ ID NO:28)	VATNLAD (SEQ ID NO:29)	QHYWGTTWT (SEQ ID NO:30)
CD37-12	RASQSVSTSSYSYLY (SEQ ID NO:31)	YASNLAS (SEQ ID NO:32)	QHSWEIPYT (SEQ ID NO:33)
CD37-38	SASSSVTYMH (SEQ ID NO:34)	DTSKLAS (SEQ ID NO:35)	QQWISNPPT (SEQ ID NO:36)
CD37-50	SATSSSVTYMH (SEQ ID NO:37)	DTSKLPY (SEQ ID NO:38)	QQWSDNPPT (SEQ ID NO:39)
		Humanized DTSNLPY (SEQ ID NO:40)	
CD37-51	SATSSSVTYMH (SEQ ID NO:41)	DTSKLAS (SEQ ID NO:42)	QQWSSNPPT (SEQ ID NO:43)
CD37-56	SASSSVTYMH (SEQ ID NO:44)	DTSKLAS (SEQ ID NO:45)	QQWISDPPT (SEQ ID NO:46)
		Humanized DTSNLAS (SEQ ID NO:47)	
CD37-57	SATSSSVTYMH (SEQ ID NO:48)	DTSKLAS (SEQ ID NO:49)	QQWSDNPPT (SEQ ID NO:50)
		Humanized DTSNLAS (SEQ ID NO:51)	
CONSENSUS	SA[T or S]SSSVTYMH (SEQ ID NO:52)	DTS[K or N]L[A or P][S or Y] (SEQ ID NO:53)	QQW[I or S][S or D][N or D]PPT (SEQ ID NO:54)
252-3	RASQDISNYLN (SEQ ID NO:174)	YTSKLHS (SEQ ID NO:175)	QQGNALPWT (SEQ ID NO:176)

[00122] The CD37 binding molecules can be antibodies or antigen binding fragments that specifically bind to CD37 that comprise the CDRs of CD37-3, CD37-12, CD37-50, CD37-51, CD37-56, or CD37-57 with up to four (i.e., 0, 1, 2, 3, or 4) conservative amino acid substitutions per CDR.

[00123] The CD37 binding molecules can comprise one of the individual variable light chains or variable heavy chains described herein. Antibodies and polypeptides can also comprise both a variable light chain and a variable heavy chain. The variable light chain and variable heavy chain sequences of murine, chimeric, and humanized CD37-3, CD37-12, CD37-50, CD37-51, CD37-56, and CD37-57 antibodies are provided in Tables 3 and 4 below.

Table 3: Variable heavy chain amino acid sequences

Antibody	VH Amino Acid Sequence (SEQ ID NO)
muCD37-3	QVQVKESGPGLVAPSQSLSICTVSGFSLTTSQVSWVRQPPGKGLEWLGVIWGDGSTNYHSALKSRLSIKKDHSKSQVFLKLNLSLQTDATYYCAKGGYSLA

	HWGOGTLVTVSA (SEQ ID NO:55)
chCD37-3	QVQVKESGPGGLVAPSQLSITCTVSGFSLTTSQVSWVRQPPGKGLEWLGVIW GDGSTNYHSALKSRLSIKKDHKSQVFLKLNLSLQITDDTATYYCAKGGYSLA HWGOGTLVTVSA (SEQ ID NO:56)
huCD37-3v1.0	QVQVQESGPGGLVAPSQLSITCTVSGFSLTTSQVSWVRQPPGKGLEWLGVIW GDGSTNYHPSLKSRLSIKKDHKSQVFLKLNLSLTAADTATYYCAKGGYSLA HWGOGTLVTVSS (SEQ ID NO:57)
huCD37-3v1.1	QVQVQESGPGGLVAPSQLSITCTVSGFSLTTSQVSWVRQPPGKGLEWLGVIW GDGSTNYHSSLKSRLSIKKDHKSQVFLKLNLSLTAADTATYYCAKGGYSLA HWGOGTLVTVSS (SEQ ID NO:58)
muCD37-12	QIQLVQSGPELKKPGETVKISCKASGYTFTKYGMNWVKQAQGKGLKWMG WINTNTGESRNAEEFKGRFAFSLETSASTAYLQINNLYEDTATYFCGRGTV VADWGQGTTLTVSS (SEQ ID NO:59)
chCD37-12	QIQLVQSGPELKKPGETVKISCKASGYTFTKYGMNWVKQAQGKGLKWMG WINTNTGESRNAEEFKGRFAFSLETSASTAYLQINNLYEDTATYFCGRGTV VADWGQGTTLTVSS (SEQ ID NO:60)
muCD37-38	DVQLQESGPDLVKPSQSLSLTCTVTGYSITSGFGWHWIRQFPGNKLEWMAY ILYSGGTDYNPSLKSRLSITRDTSKNQFFLRLSSVTTEDTATYYCARGYYGYG AWFVYWGQGTTLTVSA (SEQ ID NO:61)
chCD37-38	QVQLQESGPDLVKPSQSLSLTCTVTGYSITSGFGWHWIRQFPGNKLEWMAY ILYSGGTDYNPSLKSRLSITRDTSKNQFFLRLSSVTTEDTATYYCARGYYGYG AWFVYWGQGTTLTVSA (SEQ ID NO:62)
huCD37-38	QVQLQESGPGGLVKPSQSLSLTCTVSGYSITSGFGWHWIRQFPGKGLEWMAYI LYSGGTDYNPSLKSRLSITRDTSKNQFFLRLSSVTAADTATYYCARGYYGYG AWFVYWGQGTTLTVSS (SEQ ID NO:63)
muCD37-50	DVQLQESGPDLLKPSQSLSLTCTVTGYSITSGFAWHWIRQFPGNKLEWMGYI LYSGSTVYSPSLKSRLSITRDTSKNHFFLQLNSVTTEDTATYYCARGYYGYG AWFAYWGQGTTLTVSA (SEQ ID NO:64)
huCD37-50	QVQLQESGPGLLKPSQSLSLTCTVSGYSITSGFAWHWIRQHPGNKLEWMGYI ILYSGSTVYSPSLKSRLSITRDTSKNHFFLQLNSVTAADTATYYCARGYYGYG AWFAYWGQGTTLTVSA (SEQ ID NO:65)
muCD37-51	DVQLQESGPDLLKPSQSLSLTCTVTGYSISSGFAWHWIRQFPGNKLEWMGYI HYSGSTNYSPLKSRISITRDTSSKNQFFLQLNSVTTEDTATYYCARGYYGFGA WFVYWGQGTTLTVSA (SEQ ID NO:66)
huCD37-51	EVQLVESGPEVLKPGESLSLCTVSGYSISSGFAWHWIRQFPGKGLEWMGYI HYSGSTNYSPLQGRISITRDTSSINQFFLQLNSVTASDTATYYCARGYYGFGA WFVYWGQGTTLTVSA (SEQ ID NO:67)
muCD37-56	DVQLQESGPDLVKPSQSLSLTCTVTGYSITSGFAWHWIRQFPGNKLEWMGYI IHYSGGTNYNPSLKSRLSITRDTSKNQFFLQLNSVTTEDTATYYCARGYYGF GAWFAYWGQGTLPVSA (SEQ ID NO:68)
huCD37-56	QVQLQESGPGGLVKPSQSLSLTCTVSGYSITSGFAWHWIRQFPGKGLEWMGYI HYSGGTNYNPSLKSRLSITRDTSKNQFFLQLNSVTAADTATYYCARGYYGF GAWFAYWGQGTLPVSA (SEQ ID NO:69)
muCD37-57	DVQLQESGPDLLKPSQSLSLTCTVTGYSITSGFAWHWIRQFPGNKLEWMGYI LYSGSTVYSPSLKSRISITRDTSKNQFFLQLNSVTTEDTATYYCARGYYGYG AWFAYWGQGTTLTVSA (SEQ ID NO:70)
huCD37-57	QVQLQESGPGLLKPSQSLSLTCTVSGYSITSGFAWHWIRQFPGKGLEWMGYI LYSGSTVYSPSLKSRISITRDTSKNQFFLQLNSVTAADTATYYCARGYYGYG AWFAYWGQGTTLTVSA (SEQ ID NO:71)
252-3	EVQVVESGGDLVKPGGSLKLSAASGFTFSSYGMSWVRQTPDKRLEWVATI SSGGSYTYSVSPDKRFTISRDNACKTLYLQMSLKSSEDAMYYCARHSYY DTSVDYWGQGTSTVTVSS (SEQ ID NO:177)

Table 4: Variable light chain amino acid sequences

Antibody	VL Amino Acid Sequence (SEQ ID NO)
muCD37-3	DIQMTQSPASLSVSVGETVTITCRASENIRSNLAWYQQKQKGKSPQLLVNVAT NLADGVPSRFSGSGSGTQYSLKINSLQSEDFGTYYCQHYWGTTWTFGGGK LEIKR (SEQ ID NO:72)
chCD37-3	DIQMTQSPASLSVSVGETVTITCRASENIRSNLAWYQQKQKGKSPQLLVNVAT NLADGVPSRFSGSGSGTQYSLKINSLQSEDFGTYYCQHYWGTTWTFGGGK LEIKR (SEQ ID NO:73)
huCD37-3 (1.0 and 1.1)	DIQMTQSPSSLSVSVGERVTITCRASENIRSNLAWYQQKPGKSPKLLNVVAT NLADGVPSRFSGSGSGTDYSLKINSLQPEDFGTYYCQHYWGTTWTFGQGTK LEIKR (SEQ ID NO:74)
muCD37-12	DIVLTQSPASLAVSLGQRATISCRASQSVSTSSYSYLYWFQOKPGQPPKLLIK YASNLASGVPARFSGSGSGTDFTLNIHPVEEEDTATYYCQHSWEIPYTFGGG TKLEIKR (SEQ ID NO:75)
chCD37-12	DIVLTQSPASLAVSLGQRATISCRASQSVSTSSYSYLYWFQOKPGQPPKLLIK YASNLASGVPARFSGSGSGTDFTLNIHPVEEEDTATYYCQHSWEIPYTFGGG TKLEIKR (SEQ ID NO:76)
muCD37-38	QIVLTQSPAIMASAPGEKVTMTCSASSSVTYMHWYQQKSGTSPKRWIYDTS KLASGVPARFSGGGSGTYSYSLTISSMEAEDAATYYCQQWISNPPTFGGGTKL EIKR (SEQ ID NO:77)
chCD37-38	QIVLTQSPAIMASAPGEKVTMTCSASSSVTYMHWYQQKSGTSPKRWIYDTS KLASGVPARFSGGGSGTYSYSLTISSMEAEDAATYYCQQWISNPPTFGGGTKL EIKR (SEQ ID NO:78)
huCD37-38	DIVLTQSPASMSASPGERVMTMTCSASSSVTYMHWYQQKPGTSPKRWIYDTS KLASGVPARFSGSGSGTYSYSLTISSMEAEDAATYYCQQWISNPPTFGGGTKL EIKR (SEQ ID NO:79)
muCD37-50	QIVLTQSPAIMASAPGEKVTMTCSATSSSVTYMHWYQQKSGTSPKRWIYDTS KLPYGVPRFSGSGSGTYSYSLTISSMEAEDAATYYCQQWSDNPPTFGSGTKL EIKR (SEQ ID NO:80)
huCD37-50	EIVLTQSPATMSASPGERVMTMTCSATSSSVTYMHWYQQKPGQSPKRWIYDTS NLPYGVPARFSGSGSGTYSYSLTISSMEAEDAATYYCQQWSDNPPTFGQGTKL EIKR (SEQ ID NO:81)
muCD37-51	QIVLTQSPAIMASAPGEKVTMTCSATSSSVTYMHWYQQKSGTSPKRWIYDTS KLASGVPARFSGSGSGTYSYSLTISNMEAEDAATYYCQQWSSNPPTFGSGTKL EIKR (SEQ ID NO:82)
huCD37-51	EIVLTQSPATMSASPGERVMTMTCSATSSSVTYMHWYQQKPGQSPKRWIYDTS KLASGVPARFSGSGSGTYSYSLTISSMEAEDAATYYCQQWSSNPPTFGQGTKL EIKR (SEQ ID NO:83)
muCD37-56	QIVLTQSPAFMSASPGDKVTMTCSASSSVTYMHWYQQKSGTSPKRWIYDTS KLASGVPARFSGGGSGTYSYSLTISTMEAEDAATYYCQQWISDPPTFGGGTKL EIKR (SEQ ID NO:84)
huCD37-56	DIVLTQSPAFMSASPGEKVTMTCSASSSVTYMHWYQQKPDQSPKRWIYDTS NLASGVPSRFSGSGSGTDYSLTISSMEAEDAATYYCQQWISDPPTFGQGTKL EIKR (SEQ ID NO:85)
muCD37-57	QIVLTQSPAIMASAPGEKVTMTCSATSSSVTYMHWYQQKSGTSPKRWIYDTS KLASGVPARFSGSGSGTYSYSLTISSMEAEDAATYYCQQWSDNPPTFGSGTKL EIKR (SEQ ID NO:86)
huCD37-57	EIVLTQSPATMSASPGERVMTMTCSATSSSVTYMHWYQQKPGQSPRRWIYDTS

	NLASGVPARFSGSGSGTSYSLTISSMEAEDAATYYCQQWSDNPPTFGQGTKL EIKR (SEQ ID NO:87)
252-3	DIQMTQTTSSLSASLGDRVTISCRASQDISNYLNWYQQKPDGTVKLLIYYTS KLHSGVPSRFSGSGSGTDYSLTISNLEQEDIATYFCQQGNALPWTFGGGTKL ELKR (SEQ ID NO:178)

[00124] Also provided are polypeptides that comprise: (a) a polypeptide having at least about 90% sequence identity to SEQ ID NOs:55-71 or 177; and/or (b) a polypeptide having at least about 90% sequence identity to SEQ ID NOs:72-87 or 178. In certain embodiments, the polypeptide comprises a polypeptide having at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% sequence identity to SEQ ID NOs:55-87, 177, or 178. Thus, in certain embodiments, the polypeptide comprises (a) a polypeptide having at least about 95% sequence identity to SEQ ID NOs:55-71 or 177, and/or (b) a polypeptide having at least about 95% sequence identity to SEQ ID NOs:72-87 or 178. In certain embodiments, the polypeptide comprises (a) a polypeptide having the amino acid sequence of SEQ ID NOs:55-71 or 177; and/or (b) a polypeptide having the amino acid sequence of SEQ ID NOs:72-87 or 178. In certain embodiments, the polypeptide is an antibody and/or the polypeptide specifically binds CD37. In certain embodiments, the polypeptide is a murine, chimeric, or humanized antibody that specifically binds CD37. In certain embodiments, the polypeptide having a certain percentage of sequence identity to SEQ ID NOs:55-87, 177, or 178 differs from SEQ ID NOs:55-87 by conservative amino acid substitutions only.

[00125] Polypeptides can comprise one of the individual light chains or heavy chains described herein. Antibodies and polypeptides can also comprise both a light chain and a heavy chain. The light chain and variable chain sequences of murine, chimeric, and humanized CD37-3, CD37-12, CD37-50, CD37-51, CD37-56, and CD37-57 antibodies are provided in Tables 5 and 6 below.

Table 5: Full-length heavy chain amino acid sequences

Antibody	Full-Length Heavy Chain Amino Acid Sequence (SEQ ID NO)
muCD37-3	QVQVKESGPGLVAPSQLSITCTVSGFSLTTSQVSWVRQPPGKGLEWLGVIW GDGSTNYHSALKSRLSIKKDHKSQVFLKLNLSLQTDATYYCAKGGYSLA HWGQGLTVTVSAAKTTAPSVYPLAPVCGDITGSSVTLGCLVKGYFPEPVTL TWNSGSLSSGVHTFPAVLQSDLYTLSSSVTVTSSTWPSQSITCNVAHPASSTK VDKKIEPRGPTIKPCPPCKCPAPNLLGGPSVFIFPPKIKDVLMIKSLPIVTCVVV DVSEDDPDVQISWVFNVEVHTAQTQTHREDYNSTLRVVSALPIQHQDWM SGKEFKCKVNNKDLPAPIERTISKPKGVRAPQVYVLPPEEEMTKKQVTLT CMVTDMPEDIYVEWTNNGKTELNYKNTEPVLDSGYSYFMYSKLRVEKKN WVERNSYSCSVVHEGLHNHHTTKSFSRTPGK (SEQ ID NO:88)
chCD37-3	QVQVKESGPGLVAPSQLSITCTVSGFSLTTSQVSWVRQPPGKGLEWLGVIW GDGSTNYHSALKSRLSIKKDHKSQVFLKLNLSLQTDATYYCAKGGYSLA HWGQGLTVTVSAASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVS WNSGALTSQVHTFPAVLQSSGLYSLSSVTVTPSSSLGTQTYICNVNHKPSNT KVDKKVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQ

	DWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQ VSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDK SRWQOGNVFSCSV MHEALHNHYTQKSLSLSPGK (SEQ ID NO:89)
huCD37-3v1.0	QVQVQESGPGVLVAPSQTLSITCTVSGFSLTTSVSGVSWVRQPPGKGLEWLGVIW GDGSTNYHPSLKSRLSIKKDHSKSQVFLKLNLSLTAADTATYYCAKGGYSLA HWGQGTLLTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVS WNSGALTSKVHTFPFAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNT KVDKKVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQ DWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQ VSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDK SRWQOGNVFSCSV MHEALHNHYTQKSLSLSPG (SEQ ID NO:90)
huCD37-3v1.1	QVQVQESGPGVLVAPSQTLSITCTVSGFSLTTSVSGVSWVRQPPGKGLEWLGVIW GDGSTNYHSSLKSRLSIKKDHSKSQVFLKLNLSLTAADTATYYCAKGGYSLA HWGQGTLLTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVS WNSGALTSKVHTFPFAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNT KVDKKVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQ DWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQ VSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDK SRWQOGNVFSCSV MHEALHNHYTQKSLSLSPG (SEQ ID NO:91)
muCD37-12	QIQLVQSGPELKKPGETVKISCKASGYTFTKYGMNWWKQAQGGKGLKWMG WINTNTGESRNAEEFKGRFAFSLETSASTAYLQINNLKYEDTATYFCGRGTV VADWGQGTLLTVSSAKTTAPSVYPLAPVCGDITGSSVTLGCLVKGYFPEPV TLTWNSSGLSSGVHTFPFAVLQSDLYTLSSSVTVTSSWPSQSITCNVAHPASS TKVDKKIEPRGPTIKPCPPCKCPAPNLLGGPSVFIFPPKIKDVLMSLSPVTCV VVDVSEDDPDVQISWVFNNEVHTAQTQTHREDYNSTLRVVSALPIQHQQD WMSGKEFKCKVNNKDLPAPIERTISKPKGSVRAPQVYVLPPEEEMTKKQV TLTCMVTDFMPEDIYVEWTNNGKTELNYKNTEPVLDSDGSYFMYSKLRVE KKNWVERNSYSCSVVHEGLHNHHTTKSFSRTPGK (SEQ ID NO:92)
chCD37-12	QIQLVQSGPELKKPGETVKISCKASGYTFTKYGMNWWKQAQGGKGLKWMG WINTNTGESRNAEEFKGRFAFSLETSASTAYLQINNLKYEDTATYFCGRGTV VADWGQGTLLTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVT VSWNSGALTSKVHTFPFAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPS NTKVDKKVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVT CVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVL HQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTV DKSRWQOGNVFSCSV MHEALHNHYTQKSLSLSPGK (SEQ ID NO:93)
muCD37-38	DVQLQESGPDLVKPSQSLSLTCTVTGYSITSGFGWHWIRQFPGNKLEWMAY ILYSGGTDYNPSLKSRLSITRDTSKNQFFLRLSSVTTEDTATYYCARGYYGYG AWFVYWGQGTLLTVSAAKTTPPSVYPLAPGSAAQINSMVTLGCLVKGYFP EPVTVTWNSSGLSSGVHTFPFAVLES DLYTLSSSVTVPSSMRPSETVTCNVAH PASSTKVDKKIVPRDCGCKPCICTVPEVSSVFIFPPKPKDVLITLTPKVTVCV VDISKDDPEVQFSWFVDDVEVHTAQTQPREEQFNSTFRSVSELPIMHQDWL NGKEFKCRVNSAAFPAPIEKTISKTKGRPKAPQVYTI PPPKEQMAKDKVSLT CMITDFFPEDITVEWQWNGQPAENYKNTQPIMN TNGSYFVYSKLNQKSN WEAGNTFTCSVLHEGLHNHHTTEKSLSHSPGK (SEQ ID NO:94)
chCD37-38	QVQLQESGPDLVKPSQSLSLTCTVTGYSITSGFGWHWIRQFPGNKLEWMAY ILYSGGTDYNPSLKSRLSITRDTSKNQFFLRLSSVTTEDTATYYCARGYYGYG AWFVYWGQGTLLTVSAASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPE PVTVSWNSGALTSKVHTFPFAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNH KPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTP

	<p>EVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLT VLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDEL KNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLT VDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPG (SEQ ID NO:95)</p>
huCD37-38	<p>QVQLQESGPGLVKPSQSLSTCTVSGYSITSGFGWHWIRQFPKGLWMAI LYSGGTDYNPSLKSRIITRDTSKNQFFLRLSSVTAADTATYYCARGYYGYG AWFVYWGQGTILVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPE PVTVSWNSGALTSKVHTFPAVLQSSGLYSLSSVTVTPSSSLGTQTYICNVNH KPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRT EVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLT VLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDEL KNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLT VDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPG (SEQ ID NO:96)</p>
muCD37-50	<p>DVQLQESGPDLLKPSQSLSTCTVTGYISITSGFAWHWIRQFPGNKLEWMGYI LYSGSTVYSPSLKSRISITRDTSKNHFFLQLNSVTTEDTATYYCARGYYGYG AWFAYWGQGTILVTVSAAKTTPSVYPLAPVCGDTTGSSTVTLGCLVKGYFP EPVTLTWNSGSLSSGVHTFPAVLQSDLYTLSSSVTVTSSTWPSQSITCNVAHP ASSTKVDKKIEPRGPTIKPCPPCKCPAPNLLGGPSVFIFPPKIKDVLMSLSPIV TCVVVDVSEDDPDVQISWVFNVEVHTAQTQTHREDYNSTLRVVSALPIQH QDWMGKEFKCKVNNKDLPAPIERTISKPKGSRAPQVYVLPPEEEMTKK QVTLTCMVTDMPEDIYVEWTNNGKTELNYKNTEPVLDSGYSYFMYSKLR VEKKNWVERNSYSCSVVHEGLHNHHTTKSFSRTPGK (SEQ ID NO:97)</p>
huCD37-50	<p>QVQLQESGPGLLKPSQSLSTCTVSGYSITSGFAWHWIRQHPGNKLEWMGYI ILYSGSTVYSPSLKSRISITRDTSKNHFFLQLNSVTAADTATYYCARGYYGYG AWFAYWGQGTILVTVSAASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPE PVTVSWNSGALTSKVHTFPAVLQSSGLYSLSSVTVTPSSSLGTQTYICNVNH KPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRT EVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLT VLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDEL KNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLT VDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPG (SEQ ID NO:98)</p>
muCD37-51	<p>DVQLQESGPDLLKPSQSLSTCTVTGYSISSGFAWHWIRQFPGNKLEWMGYI HYSGSTNYSPSLKSRISITRDSSKNQFFLQLNSVTTEDTATYYCARGYYGFGA WVVYWGQGTILVTVSAAKTTPSVYPLAPVCGDTTGSSTVTLGCLVKGYFPEP VTLTWNSGSLSSGVHTFPAVLQSDLYTLSSSVTVTSSTWPSQSITCNVAHPAS STKVDKKIEPRGPTIKPCPPCKCPAPNLLGGPSVFIFPPKIKDVLMSLSPIVTC VVVVDVSEDDPDVQISWVFNVEVHTAQTQTHREDYNSTLRVVSALPIQH QDWMGKEFKCKVNNKDLPAPIERTISKPKGSRAPQVYVLPPEEEMTKKQV TLTCMVTDMPEDIYVEWTNNGKTELNYKNTEPVLDSGYSYFMYSKLRVE KKNWVERNSYSCSVVHEGLHNHHTTKSFSRTPGK (SEQ ID NO:99)</p>
huCD37-51	<p>EVQLVESGPEVLKPGESLSLTCTVSGYSISSGFAWHWIRQFPKGLWMAI HYSGSTNYSPSLQGRISITRDSSINQFFLQLNSVTAASDTATYYCARGYYGFGA WVVYWGQGTILVTVSAASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEP VTVSWNSGALTSKVHTFPAVLQSSGLYSLSSVTVTPSSSLGTQTYICNVNHK PSNTKVDKKVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPE VTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLT VLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDEL KNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLT VDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPG (SEQ ID NO:100)</p>
muCD37-56	<p>DVQLQESGPDLVKPSQSLSTCTVTGYISITSGFAWHWIRQFPGNKLEWMGYI IHYSGGTNYNPSLKSRIITRDTSKNQFFLQLNSVTTEDTATYYCARGYYGF GAWFAYWGQGTILVTVSAAKTTPSVYPLAPGSAQAQTNMVTGCLVKGYF PEPVTVTWNSGSLSSGVHTFPAVLESGLYTLSSSVTVTPSSMRPSETVTCNVA</p>

	HPASSTKVDKKIVPRDCGCKPCICTVPEVSSVFIFPPKPKDVLTIITLTPKVTCV VVDISKDDPEVQFSWFVDDVEVHTAQTQPREEQFNSTFRSVSELPIMHQDW LNGKEFKCRVNSAAFPAPIEKTISKTKGRPKAPQVYTIPPPKEQMAKDKVSL TCMITDFFPEDITVEWQWNGQPAENYKNTQPMNINGSYFVYSKLNQKSN WEAGNTFTCSVLHEGLHNHHTTEKSLSHSPGK (SEQ ID NO:101)
huCD37-56	QVQLQESGPGGLVKPSQSLSLTCTVSGYSITSGFAWHWIRQFPGKGLEWMGYI HYSGGTNYNPSLKSRSVITRDTSKNQFFLQLNSVTAADTATYYCARGYYGF GAWFAYWGQGLVPVSAASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFP EPVTVSWNSGALTSQVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVN HKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRT PEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVL TVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDEL TKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKL TVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPG (SEQ ID NO:102)
muCD37-57	DVQLQESGPDLLKPSQSLSLTCTVTGYSITSGFAWHWIRQFPGNKLEWMGYI LYSGSTVYSPSLKSRISITRDTSKNQFFLQLNSVTTEDTATYYCARGYYGYG AWFAYWGQGLVTVSAAKTTAPSVYPLAPVCGDTTGSSVTLGCLVKGYFP EPVTLTWNSSGLSSGVHTFPAVLQSDLYTLSSSVTVTSSTWPSQSITCNVAHP ASSTKVDKKIEPRGPTIKPCPPCKCPAPNLLGGPSVFIFPPKIKDVLMISSPIV TCVVVDVSEDDPDVQISWFVNNVEVHTAQTQTHREDYNSTLRVVSALPIQH QDWMSGKEFKCKVNNKDLPAPIERTISKPKGSRAPQVYVLPPEEEMTKK QVTLTCMVTDMPEDIYVEWTNNGKTELNYKNTEPVLDSGDSYFMYSKLR VEKKNWVERNSYSCSVVHEGLHNHHTTKSFSRTPGK (SEQ ID NO:103)
huCD37-57	QVQLQESGPGGLKPSQSLSLTCTVSGYSITSGFAWHWIRQFPGKGLEWMGYI LYSGSTVYSPSLKSRISITRDTSKNQFFLQLNSVTAADTATYYCARGYYGYG AWFAYWGQGLVTVSAAASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPE PVTVSWNSGALTSQVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNH KPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRT EVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLT VLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDEL KNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLT VDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPG (SEQ ID NO:104)
252-3	EVQVVESGGDLVKPFGGSLKLSCAASGFTFSSYGMSWVRQTPDKRLEWVATI SSGGSYTYSPDSVKGRFTISRDNAAKTLYLQMSSLKSEDTAMYYCARHSYY DTSVDYWGQGTSTVSSAKTTAPSVYPLAPVCGDTTGSSVTLGCLVKGYFP EPVILTWNSSGLSSGVHTFPAVLQSDLYTLSSSVTVTSSTWPSQSITCNVAHP ASSTKVDKKIEPRGPTIKPCPPCKCPAPNLLGGPSVFIFPPKIKDVLMISSPIV TCVVVDVSEDDPDVQISWFVNNVEVHTAQTQTHREDYNSTLRVVSALPIQH QDWMSGKEFKCKVNNKDLPAPIERTISKPKGSRAPQVYVLPPEEEMTKK QVTLTCMVTDMPEDIYVEWTNNGKTELNYKNTEPVLDSGDSYFMYSKLR VEKKNWVERNSYSCSVVHEGLHNHHTTKSFSRTPGK (SEQ ID NO:179)

Table 6: Full-length light chain amino acid sequences

Antibody	Full-length Light Chain Amino Acid Sequence (SEQ ID NO)
muCD37-3	DIQMTQSPASLSVSVGETVTITCRASENIRSNLAWYQQKQKSPQLLVNVAI NLADGVPSRFSGSGSGTQYSLKINSLQSEDFGTYYCQHYWGTTWTFGGGK LEIKRADAAPTIVSIFPPSSEQLTSGGASVVCFLNFPKIDINVKWKIDGSERQ NGVLNSWTDQSDKSTYSMSSTLTTLTKDEYERHNSYCEATHKTSSTPIVKS FNRNEC (SEQ ID NO:105)
chCD37-3	DIQMTQSPASLSVSVGETVTITCRASENIRSNLAWYQQKQKSPQLLVNVAI

	NLADGVPSRFSGSGSGTQYSLKINSLQSEDFGTYYCQHYWGTTWTFGGGK LEIKRTVAAPS VFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQ SGNSQESVTEQDSKDSTYLSSTLTLKADYEKHKVYACEVTHQGLSSPVTK SFNRGEC (SEQ ID NO:106)
huCD37-3 (1.0 and 1.1)	DIQMTQSPSSLSVSVGERVTITCRASENIRS NLAWYQQKPGKSPKLLVNVAT NLADGVPSRFSGSGSGT DYSLKINSLQPEDFGTYYCQHYWGTTWTFGGGK LEIKRTVAAPS VFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQ SGNSQESVTEQDSKDSTYLSSTLTLKADYEKHKVYACEVTHQGLSSPVTK SFNRGEC (SEQ ID NO:107)
muCD37-12	DIVLTQSPASLAVSLGQRATISCRASQSVSTSSYSYLYWFQQKPGQPPELLIK YASNLAGVVPARFSGSGSGTDFTLNIHPVEEEDTATYYCQHSWEIPYTFGGG TKLEIKRADAAPT VSIFFPSSEQLTSGGASVVCFLNNFYPKDINVKWKIDGSE RQNGVLNSWTDQDSKDSTYSMSSTLTLTKDEYERHNSYTCEATHKTSTSPI VKSFNRENC (SEQ ID NO:108)
chCD37-12	DIVLTQSPASLAVSLGQRATISCRASQSVSTSSYSYLYWFQQKPGQPPELLIK YASNLAGVVPARFSGSGSGTDFTLNIHPVEEEDTATYYCQHSWEIPYTFGGG TKLEIKRTVAAPS VFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNA LQSGNSQESVTEQDSKDSTYLSSTLTLKADYEKHKVYACEVTHQGLSSPV TKSFNRGEC (SEQ ID NO:109)
muCD37-38	QIVLTQSPAIMASAPGKVTMTCSASSSVTYMHWYQQKSGTSPKRWIYDTS KLASGVPARFSGGGSGTSYSLTISSMEAEDAATYYCQQWISNPPTFGGGTKL EIKRADAAPT VSIFFPSSEQLTSGGASVVCFLNNFYPKDINVKWKIDGSERQN GVLNSWTDQDSKDSTYSMSSTLTLTKDEYERHNSYTCEATHKTSTSPIVKSF NRNEC (SEQ ID NO:110)
chCD37-38	QIVLTQSPAIMASAPGKVTMTCSASSSVTYMHWYQQKSGTSPKRWIYDTS KLASGVPARFSGGGSGTSYSLTISSMEAEDAATYYCQQWISNPPTFGGGTKL EIKRTVAAPS VFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQS GNSQESVTEQDSKDSTYLSSTLTLKADYEKHKVYACEVTHQGLSSPVTKS FNRGEC (SEQ ID NO:111)
huCD37-38	DIVLTQSPASMSAPGERVTMTCSASSSVTYMHWYQQKPGTSPKRWIYDTS KLASGVPARFSGSGSGTSYSLTISSMEAEDAATYYCQQWISNPPTFGGGTKL EIKRTVAAPS VFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQS GNSQESVTEQDSKDSTYLSSTLTLKADYEKHKVYACEVTHQGLSSPVTKS FNRGEC (SEQ ID NO:112)
muCD37-50	QIVLTQSPAIMASAPGKVTMTCSATSSSVTYMHWYQQKSGTSPKRWIYDTS KLPGVPGRFSGSGSGTSYSLTISSMEAEDAATYYCQQWSDNPPTFGSGTKL EIKRADAAPT VSIFFPSSEQLTSGGASVVCFLNNFYPKDINVKWKIDGSERQN GVLNSWTDQDSKDSTYSMSSTLTLTKDEYERHNSYTCEATHKTSTSPIVKSF NRNEC (SEQ ID NO:113)
huCD37-50	EIVLTQSPATMSAPGERVTMTCSATSSSVTYMHWYQQKPGQSPKRWIYDTS NLPYGVPARFSGSGSGTSYSLTISSMEAEDAATYYCQQWSDNPPTFGGQTKL EIKRTVAAPS VFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQS GNSQESVTEQDSKDSTYLSSTLTLKADYEKHKVYACEVTHQGLSSPVTKS FNRGEC (SEQ ID NO:114)
muCD37-51	QIVLTQSPAIMASAPGKVTMTCSATSSSVTYMHWYQQKSGTSPKRWIYDTS KLASGVPARFSGSGSGTSYSLTISNMEAEDAATYYCQQWSSNPPTFGSGTKL EIKRADAAPT VSIFFPSSEQLTSGGASVVCFLNNFYPKDINVKWKIDGSERQN GVLNSWTDQDSKDSTYSMSSTLTLTKDEYERHNSYTCEATHKTSTSPIVKSF NRNEC (SEQ ID NO:115)
huCD37-51	EIVLTQSPATMSAPGERVTMTCSATSSSVTYMHWYQQKPGQSPKRWIYDTS KLASGVPARFSGSGSGTSYSLTISSMEAEDAATYYCQQWSSNPPTFGGQTKL EIKRTVAAPS VFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQS

	GNSQESVTEQDSKDYSLSSSTLTLTKADYKHKVYACEVTHQGLSSPVTKS FNRGEC (SEQ ID NO:116)
muCD37-56	QIVLTQSPAFMSASPGDKVTMTCSASSSVTYMHWYQQKSGTSPKRWIYDTS KLAGVVPARFSGGGSGTYSYSLTISTMEAEDAATYYCQQWISDPPTFGGGTKL EIKRADAAPTVSIFPPSSEQLTSGGASVVCFLNMFYPKDINVKWKIDGSRQN GVLNSWTDQDSKDYSLSSSTLTLTKDEYERHNSYTCEATHKTSTSPIVKSF NRNEC (SEQ ID NO:117)
huCD37-56	DIVLTQSPAFMSASPGDKVTMTCSASSSVTYMHWYQQKPDQSPKRWIYDTS NLAGVPSRFSGGGSGTDYSLTISSMEAEDAATYYCQQWISDPPTFGQGTKL EIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNMFYPREKAVQWKVDNALQS GNSQESVTEQDSKDYSLSSSTLTLTKADYKHKVYACEVTHQGLSSPVTKS FNRGEC (SEQ ID NO:118)
muCD37-57	QIVLTQSPAIMASASPGDKVTMTCSATSSSVTYMHWYQQKSGTSPKRWIYDTS KLAGVVPARFSGSGSGTYSYSLTISSMEAEDAATYYCQQWSDNPPTFGSGTKL EIKRADAAPTVSIFPPSSEQLTSGGASVVCFLNMFYPKDINVKWKIDGSRQN GVLNSWTDQDSKDYSLSSSTLTLTKDEYERHNSYTCEATHKTSTSPIVKSF NRNEC (SEQ ID NO:119)
huCD37-57	EIVLTQSPATMSASPGERVTMTCSATSSSVTYMHWYQQKPGQSPRRWIYDTS NLAGVVPARFSGSGSGTYSYSLTISSMEAEDAATYYCQQWSDNPPTFGQGTKL EIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNMFYPREKAVQWKVDNALQS GNSQESVTEQDSKDYSLSSSTLTLTKADYKHKVYACEVTHQGLSSPVTKS FNRGEC (SEQ ID NO:120)
252-3	DIQMTQTSSLSASLGDRVTISCRASQDISNYLNWYQQKPDGTVKLLIYYTS KLHSGVPSRFSGSGSGTDYSLTISNLEQEDIATYFCQQGNALPWTFGGGTKL ELKRADAAPTVSIFPPSSEQLTSGGASVVCFLNMFYPKDINVKWKIDGSRQ NGVLNSWTDQDSKDYSLSSSTLTLTKDEYERHNSYTCEATHKTSTSPIVKS FNRNEC (SEQ ID NO:180)

[00126] Also provided are polypeptides that comprise: (a) a polypeptide having at least about 90% sequence identity to SEQ ID NOs:88-104 or 179; and/or (b) a polypeptide having at least about 90% sequence identity to SEQ ID NOs:105-120 or 180. In certain embodiments, the polypeptide comprises a polypeptide having at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% sequence identity to SEQ ID NOs:88-120, 179, or 180. Thus, in certain embodiments, the polypeptide comprises (a) a polypeptide having at least about 95% sequence identity to SEQ ID NOs:88-104 or 179, and/or (b) a polypeptide having at least about 95% sequence identity to SEQ ID NOs:105-120 or 180. In certain embodiments, the polypeptide comprises (a) a polypeptide having the amino acid sequence of SEQ ID NOs:88-104 or 179; and/or (b) a polypeptide having the amino acid sequence of SEQ ID NOs:105-120 or 180. In certain embodiments, the polypeptide is an antibody and/or the polypeptide specifically binds CD37. In certain embodiments, the polypeptide is a murine, chimeric, or humanized antibody that specifically binds CD37. In certain embodiments, the polypeptide having a certain percentage of sequence identity to SEQ ID NOs:88-120, 179, or 180 differs from SEQ ID NOs:88-120, 179, or 180 by conservative amino acid substitutions only.

[00127] In certain embodiments, the CD37 antibody can be the antibody produced from a hybridoma selected from the group consisting of consisting of ATCC Deposit Designation PTA-10664, deposited with the ATCC on February 18, 2010, ATCC Deposit Designation PTA-10665, deposited with the ATCC

on February 18, 2010, ATCC Deposit Designation PTA-10666, deposited with the ATCC on February 18, 2010, ATCC Deposit Designation PTA-10667 deposited with the ATCC on February 18, 2010, ATCC Deposit Designation PTA-10668, deposited with the ATCC on February 18, 2010, ATCC Deposit Designation PTA-10669, deposited with the ATCC on February 18, 2010, and ATCC Deposit Designation PTA-10670, deposited with the ATCC on February 18, 2010 (American Type Culture Collection (ATCC) at 10801 University Boulevard, Manassas, Virginia 20110). In certain embodiments, the antibody comprises the VH-CDRs and the VL-CDRS of the antibody produced from a hybridoma selected from the group consisting of PTA-10665, PTA-10666, PTA-10667, PTA-10668, PTA-10669, and PTA-10670.

[00128] In certain embodiments, the CD37 antibody can comprise a light chain encoded by the recombinant plasmid DNA phuCD37-3LC (ATCC Deposit Designation PTA-10722, deposited with the ATCC on March 18, 2010). In certain embodiments, the CD37 antibody can comprise a heavy chain encoded by the recombinant plasmid DNA phuCD37-3HCv.1.0 (ATCC Deposit Designation PTA-10723, deposited with the ATCC on March 18, 2010). In certain embodiments, the CD37 antibody can comprise a light chain encoded by the recombinant plasmid DNA phuCD37-3LC (PTA-10722) and a heavy chain encoded by the recombinant plasmid DNA phuCD37-3HCv.1.0 (PTA-10723). In certain embodiments, the CD37 antibody can comprise the VL-CDRs encoded by the recombinant plasmid DNA phuCD37-3LC (PTA-10722) and the VH-CDRs encoded by the recombinant plasmid DNA phuCD37-3HCv.1.0 (PTA-10723).

[00129] Monoclonal antibodies can be prepared using hybridoma methods, such as those described by Kohler and Milstein (1975) Nature 256:495. Using the hybridoma method, a mouse, hamster, or other appropriate host animal, is immunized as described above to elicit the production by lymphocytes of antibodies that will specifically bind to an immunizing antigen. Lymphocytes can also be immunized *in vitro*. Following immunization, the lymphocytes are isolated and fused with a suitable myeloma cell line using, for example, polyethylene glycol, to form hybridoma cells that can then be selected away from unfused lymphocytes and myeloma cells. Hybridomas that produce monoclonal antibodies directed specifically against a chosen antigen as determined by immunoprecipitation, immunoblotting, or by an *in vitro* binding assay (e.g. radioimmunoassay (RIA); enzyme-linked immunosorbent assay (ELISA)) can then be propagated either *in vitro* culture using standard methods (Goding, Monoclonal Antibodies: Principles and Practice, Academic Press, 1986) or *in vivo* as ascites tumors in an animal. The monoclonal antibodies can then be purified from the culture medium or ascites fluid as described for polyclonal antibodies above.

[00130] Alternatively monoclonal antibodies can also be made using recombinant DNA methods as described in U.S. Patent 4,816,567. The polynucleotides encoding a monoclonal antibody are isolated from mature B-cells or hybridoma cell, such as by RT-PCR using oligonucleotide primers that specifically

amplify the genes encoding the heavy and light chains of the antibody, and their sequence is determined using conventional procedures. The isolated polynucleotides encoding the heavy and light chains are then cloned into suitable expression vectors, which when transfected into host cells such as *E. coli* cells, simian COS cells, Chinese hamster ovary (CHO) cells, or myeloma cells that do not otherwise produce immunoglobulin protein, monoclonal antibodies are generated by the host cells. Also, recombinant monoclonal antibodies or fragments thereof of the desired species can be isolated from phage display libraries expressing CDRs of the desired species as described (McCafferty et al., 1990, *Nature*, 348:552-554; Clackson et al., 1991, *Nature*, 352:624-628; and Marks et al., 1991, *J. Mol. Biol.*, 222:581-597).

[00131] The polynucleotide(s) encoding a monoclonal antibody can further be modified in a number of different manners using recombinant DNA technology to generate alternative antibodies. In some embodiments, the constant domains of the light and heavy chains of, for example, a mouse monoclonal antibody can be substituted 1) for those regions of, for example, a human antibody to generate a chimeric antibody or 2) for a non-immunoglobulin polypeptide to generate a fusion antibody. In some embodiments, the constant regions are truncated or removed to generate the desired antibody fragment of a monoclonal antibody. Site-directed or high-density mutagenesis of the variable region can be used to optimize specificity, affinity, etc. of a monoclonal antibody.

[00132] In some embodiments, the monoclonal antibody against the human CD37 is a humanized antibody. In certain embodiments, such antibodies are used therapeutically to reduce antigenicity and HAMA (human anti-mouse antibody) responses when administered to a human subject. Humanized antibodies can be produced using various techniques known in the art. In certain alternative embodiments, the antibody to CD37 is a human antibody.

[00133] Human antibodies can be directly prepared using various techniques known in the art. Immortalized human B lymphocytes immunized *in vitro* or isolated from an immunized individual that produce an antibody directed against a target antigen can be generated (See, e.g., Cole et al., *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, p. 77 (1985); Boemer et al., 1991, *J. Immunol.*, 147 (1):86-95; and U.S. Patent 5,750,373). Also, the human antibody can be selected from a phage library, where that phage library expresses human antibodies, as described, for example, in Vaughan et al., 1996, *Nat. Biotech.*, 14:309-314, Sheets et al., 1998, *Proc. Nat'l. Acad. Sci.*, 95:6157-6162, Hoogenboom and Winter, 1991, *J. Mol. Biol.*, 227:381, and Marks et al., 1991, *J. Mol. Biol.*, 222:581). Techniques for the generation and use of antibody phage libraries are also described in U.S. Patent Nos. 5,969,108, 6,172,197, 5,885,793, 6,521,404; 6,544,731; 6,555,313; 6,582,915; 6,593,081; 6,300,064; 6,653,068; 6,706,484; and 7,264,963; and Rothe et al., 2007, *J. Mol. Bio.*, 376:1182 (each of which is incorporated by reference in its entirety). Affinity maturation strategies and chain shuffling strategies (Marks et al., 1992, *Bio/Technology* 10:779-783, incorporated by reference in its entirety) are known in the art and can be employed to generate high affinity human antibodies.

[00134] Humanized antibodies can also be made in transgenic mice containing human immunoglobulin loci that are capable upon immunization of producing the full repertoire of human antibodies in the absence of endogenous immunoglobulin production. This approach is described in U.S. Patents 5,545,807; 5,545,806; 5,569,825; 5,625,126; 5,633,425; and 5,661,016.

[00135] This invention also encompasses bispecific antibodies that specifically recognize a CD37. Bispecific antibodies are antibodies that are capable of specifically recognizing and binding at least two different epitopes. The different epitopes can either be within the same molecule (e.g. the same CD37) or on different molecules such that both, for example, the antibodies can specifically recognize and bind a CD37 as well as, for example, 1) an effector molecule on a leukocyte such as a T-cell receptor (e.g. CD3) or Fc receptor (e.g. CD64, CD32, or CD16) or 2) a cytotoxic agent as described in detail below.

[00136] Exemplary bispecific antibodies can bind to two different epitopes, at least one of which originates in a polypeptide of the invention. Alternatively, an anti-antigenic arm of an immunoglobulin molecule can be combined with an arm which binds to a triggering molecule on a leukocyte such as a T cell receptor molecule (e.g. CD2, CD3, CD28, or B7), or Fc receptors for IgG so as to focus cellular defense mechanisms to the cell expressing the particular antigen. Bispecific antibodies can also be used to direct cytotoxic agents to cells which express a particular antigen. These antibodies possess an antigen-binding arm and an arm which binds a cytotoxic agent or a radionuclide chelator, such as EOTUBE, DPTA, DOTA, or TETA. Techniques for making bispecific antibodies are common in the art (Millstein et al., 1983, *Nature* 305:537-539; Brennan et al., 1985, *Science* 229:81; Suresh et al, 1986, *Methods in Enzymol.* 121:120; Traunecker et al., 1991, *EMBO J.* 10:3655-3659; Shalaby et al., 1992, *J. Exp. Med.* 175:217-225; Kostelny et al., 1992, *J. Immunol.* 148:1547-1553; Gruber et al., 1994, *J. Immunol.* 152:5368; and U.S. Patent 5,731,168). Antibodies with more than two valencies are also contemplated. For example, trispecific antibodies can be prepared (Tutt et al., *J. Immunol.* 147:60 (1991)). Thus, in certain embodiments the antibodies to CD37 are multispecific.

[00137] In certain embodiments are provided an antibody fragment to, for example, increase tissue penetration. Various techniques are known for the production of antibody fragments. Traditionally, these fragments are derived via proteolytic digestion of intact antibodies (for example Morimoto et al., 1993, *Journal of Biochemical and Biophysical Methods* 24:107-117; Brennan et al., 1985, *Science*, 229:81). In certain embodiments, antibody fragments are produced recombinantly. Fab, Fv, and scFv antibody fragments can all be expressed in and secreted from *E. coli* or other host cells, thus allowing the production of large amounts of these fragments. Such antibody fragments can also be isolated from the antibody phage libraries discussed above. The antibody fragment can also be linear antibodies as described in U.S. Patent 5,641,870, for example, and can be monospecific or bispecific. Other techniques for the production of antibody fragments will be apparent to the skilled practitioner.

[00138] According to the present invention, techniques can be adapted for the production of single-chain antibodies specific to CD37 (see U.S. Pat. No. 4,946,778). In addition, methods can be adapted for the construction of Fab expression libraries (Huse, et al., Science 246:1275-1281 (1989)) to allow rapid and effective identification of monoclonal Fab fragments with the desired specificity for CD37, or derivatives, fragments, analogs or homologs thereof. Antibody fragments can be produced by techniques in the art including, but not limited to: (a) a F(ab')₂ fragment produced by pepsin digestion of an antibody molecule; (b) a Fab fragment generated by reducing the disulfide bridges of an F(ab')₂ fragment, (c) a Fab fragment generated by the treatment of the antibody molecule with papain and a reducing agent, and (d) Fv fragments.

[00139] It can further be desirable, especially in the case of antibody fragments, to modify an antibody in order to increase its serum half-life. This can be achieved, for example, by incorporation of a salvage receptor binding epitope into the antibody fragment by mutation of the appropriate region in the antibody fragment or by incorporating the epitope into a peptide tag that is then fused to the antibody fragment at either end or in the middle (e.g., by DNA or peptide synthesis).

[00140] Heteroconjugate antibodies are also within the scope of the present invention. Heteroconjugate antibodies are composed of two covalently joined antibodies. Such antibodies have, for example, been proposed to target immune cells to unwanted cells (U.S. Pat. No. 4,676,980). It is contemplated that the antibodies can be prepared *in vitro* using known methods in synthetic protein chemistry, including those involving crosslinking agents. For example, immunotoxins can be constructed using a disulfide exchange reaction or by forming a thioether bond. Examples of suitable reagents for this purpose include iminothiolate and methyl-4-mercaptobutyrimidate.

[00141] For the purposes of the present invention, it should be appreciated that modified antibodies can comprise any type of variable region that provides for the association of the antibody with the polypeptides of a human CD37. In this regard, the variable region can comprise or be derived from any type of mammal that can be induced to mount a humoral response and generate immunoglobulins against the desired antigen. As such, the variable region of the modified antibodies can be, for example, of human, murine, non-human primate (e.g. cynomolgus monkeys, macaques, etc.) or lupine origin. In some embodiments both the variable and constant regions of the modified immunoglobulins are human. In other embodiments the variable regions of compatible antibodies (usually derived from a non-human source) can be engineered or specifically tailored to improve the binding properties or reduce the immunogenicity of the molecule. In this respect, variable regions useful in the present invention can be humanized or otherwise altered through the inclusion of imported amino acid sequences.

[00142] In certain embodiments, the variable domains in both the heavy and light chains are altered by at least partial replacement of one or more CDRs and, if necessary, by partial framework region replacement and sequence changing. Although the CDRs can be derived from an antibody of the same

class or even subclass as the antibody from which the framework regions are derived, it is envisaged that the CDRs will be derived from an antibody of different class and possibly from an antibody from a different species. It is not always necessary to replace all of the CDRs with the complete CDRs from the donor variable region to transfer the antigen binding capacity of one variable domain to another. Rather, in some cases it is only necessary to transfer those residues that are necessary to maintain the activity of the antigen binding site. Given the explanations set forth in U.S. Pat. Nos. 5,585,089, 5,693,761 and 5,693,762, it will be well within the competence of those skilled in the art, either by carrying out routine experimentation or by trial and error testing to obtain a functional antibody with reduced immunogenicity.

[00143] Alterations to the variable region notwithstanding, those skilled in the art will appreciate that the modified antibodies of this invention will comprise antibodies (e.g., full-length antibodies or immunoreactive fragments thereof) in which at least a fraction of one or more of the constant region domains has been deleted or otherwise altered so as to provide desired biochemical characteristics such as reduced serum half-life when compared with an antibody of approximately the same immunogenicity comprising a native or unaltered constant region. In some embodiments, the constant region of the modified antibodies will comprise a human constant region. Modifications to the constant region compatible with this invention comprise additions, deletions or substitutions of one or more amino acids in one or more domains. That is, the modified antibodies disclosed herein can comprise alterations or modifications to one or more of the three heavy chain constant domains (CH1, CH2 or CH3) and/or to the light chain constant domain (CL). In some embodiments, modified constant regions wherein one or more domains are partially or entirely deleted are contemplated. In some embodiments, the modified antibodies will comprise domain deleted constructs or variants wherein the entire CH2 domain has been removed (Δ CH2 constructs). In some embodiments, the omitted constant region domain will be replaced by a short amino acid spacer (e.g. 10 residues) that provides some of the molecular flexibility typically imparted by the absent constant region.

[00144] Besides their configuration, it is known in the art that the constant region mediates several effector functions. For example, binding of the C1 component of complement to antibodies activates the complement system. Activation of complement is important in the opsonisation and lysis of cell pathogens. The activation of complement also stimulates the inflammatory response and can also be involved in autoimmune hypersensitivity. Further, antibodies bind to cells via the Fc region, with a Fc receptor site on the antibody Fc region binding to a Fc receptor (FcR) on a cell. There are a number of Fc receptors which are specific for different classes of antibody, including IgG (gamma receptors), IgE (eta receptors), IgA (alpha receptors) and IgM (mu receptors). Binding of antibody to Fc receptors on cell surfaces triggers a number of important and diverse biological responses including engulfment and destruction of antibody-coated particles, clearance of immune complexes, lysis of antibody-coated target

cells by killer cells (called antibody-dependent cell-mediated cytotoxicity, or ADCC), release of inflammatory mediators, placental transfer and control of immunoglobulin production.

[00145] In certain embodiments, the CD37-binding antibodies provide for altered effector functions that, in turn, affect the biological profile of the administered antibody. For example, the deletion or inactivation (through point mutations or other means) of a constant region domain can reduce Fc receptor binding of the circulating modified antibody. In other cases, it can be that constant region modifications, consistent with this invention, moderate complement binding and thus reduce the serum half life and nonspecific association of a conjugated cytotoxin. Yet other modifications of the constant region can be used to eliminate disulfide linkages or oligosaccharide moieties that allow for enhanced localization due to increased antigen specificity or antibody flexibility. Similarly, modifications to the constant region in accordance with this invention can easily be made using well known biochemical or molecular engineering techniques well within the purview of the skilled artisan.

[00146] In certain embodiments, a CD37-binding agent that is an antibody does not have one or more effector functions. For instance, in some embodiments, the antibody has no antibody-dependent cellular cytotoxicity (ADCC) activity and/or no complement-dependent cytotoxicity (CDC) activity. In certain embodiments, the antibody does not bind to an Fc receptor and/or complement factors. In certain embodiments, the antibody has no effector function.

[00147] It will be noted that in certain embodiments, the modified antibodies can be engineered to fuse the CH3 domain directly to the hinge region of the respective modified antibodies. In other constructs it can be desirable to provide a peptide spacer between the hinge region and the modified CH2 and/or CH3 domains. For example, compatible constructs could be expressed wherein the CH2 domain has been deleted and the remaining CH3 domain (modified or unmodified) is joined to the hinge region with a 5-20 amino acid spacer. Such a spacer can be added, for instance, to ensure that the regulatory elements of the constant domain remain free and accessible or that the hinge region remains flexible. However, it should be noted that amino acid spacers can, in some cases, prove to be immunogenic and elicit an unwanted immune response against the construct. Accordingly, in certain embodiments, any spacer added to the construct will be relatively non-immunogenic, or even omitted altogether, so as to maintain the desired biochemical qualities of the modified antibodies.

[00148] Besides the deletion of whole constant region domains, it will be appreciated that the antibodies of the present invention can be provided by the partial deletion or substitution of a few or even a single amino acid. For example, the mutation of a single amino acid in selected areas of the CH2 domain can be enough to substantially reduce Fc binding. Similarly, it can be desirable to simply delete that part of one or more constant region domains that control the effector function (e.g. complement CLQ binding) to be modulated. Such partial deletions of the constant regions can improve selected characteristics of the antibody (serum half-life) while leaving other desirable functions associated with the

subject constant region domain intact. Moreover, as alluded to above, the constant regions of the disclosed antibodies can be modified through the mutation or substitution of one or more amino acids that enhances the profile of the resulting construct. In this respect it can be possible to disrupt the activity provided by a conserved binding site (e.g. Fc binding) while substantially maintaining the configuration and immunogenic profile of the modified antibody. Certain embodiments can comprise the addition of one or more amino acids to the constant region to enhance desirable characteristics such as decreasing or increasing effector function or provide for more cytotoxin or carbohydrate attachment. In such embodiments it can be desirable to insert or replicate specific sequences derived from selected constant region domains.

[00149] The present invention further embraces variants and equivalents which are substantially homologous to the chimeric, humanized and human antibodies, or antibody fragments thereof, set forth herein. These can contain, for example, conservative substitution mutations, i.e. the substitution of one or more amino acids by similar amino acids. For example, conservative substitution refers to the substitution of an amino acid with another within the same general class such as, for example, one acidic amino acid with another acidic amino acid, one basic amino acid with another basic amino acid or one neutral amino acid by another neutral amino acid. What is intended by a conservative amino acid substitution is well known in the art.

[00150] The polypeptides of the present invention can be recombinant polypeptides, natural polypeptides, or synthetic polypeptides comprising an antibody, or fragment thereof, against a human CD37. It will be recognized in the art that some amino acid sequences of the invention can be varied without significant effect of the structure or function of the protein. Thus, the invention further includes variations of the polypeptides which show substantial activity or which include regions of an antibody, or fragment thereof, against CD37 protein. Such mutants include deletions, insertions, inversions, repeats, and type substitutions.

[00151] The polypeptides and analogs can be further modified to contain additional chemical moieties not normally part of the protein. Those derivatized moieties can improve the solubility, the biological half life or absorption of the protein. The moieties can also reduce or eliminate any desirable side effects of the proteins and the like. An overview for those moieties can be found in REMINGTON'S PHARMACEUTICAL SCIENCES, 20th ed., Mack Publishing Co., Easton, PA (2000).

[00152] The isolated polypeptides described herein can be produced by any suitable method known in the art. Such methods range from direct protein synthetic methods to constructing a DNA sequence encoding isolated polypeptide sequences and expressing those sequences in a suitable transformed host. In some embodiments, a DNA sequence is constructed using recombinant technology by isolating or synthesizing a DNA sequence encoding a wild-type protein of interest. Optionally, the sequence can be

mutagenized by site-specific mutagenesis to provide functional analogs thereof. See, e.g. Zoeller et al., Proc. Nat'l. Acad. Sci. USA 81:5662-5066 (1984) and U.S. Pat. No. 4,588,585.

[00153] In some embodiments a DNA sequence encoding a polypeptide of interest would be constructed by chemical synthesis using an oligonucleotide synthesizer. Such oligonucleotides can be designed based on the amino acid sequence of the desired polypeptide and selecting those codons that are favored in the host cell in which the recombinant polypeptide of interest will be produced. Standard methods can be applied to synthesize an isolated polynucleotide sequence encoding an isolated polypeptide of interest. For example, a complete amino acid sequence can be used to construct a back-translated gene. Further, a DNA oligomer containing a nucleotide sequence coding for the particular isolated polypeptide can be synthesized. For example, several small oligonucleotides coding for portions of the desired polypeptide can be synthesized and then ligated. The individual oligonucleotides typically contain 5' or 3' overhangs for complementary assembly.

[00154] Once assembled (by synthesis, site-directed mutagenesis or another method), the polynucleotide sequences encoding a particular isolated polypeptide of interest will be inserted into an expression vector and operatively linked to an expression control sequence appropriate for expression of the protein in a desired host. Proper assembly can be confirmed by nucleotide sequencing, restriction mapping, and expression of a biologically active polypeptide in a suitable host. As is well known in the art, in order to obtain high expression levels of a transfected gene in a host, the gene must be operatively linked to transcriptional and translational expression control sequences that are functional in the chosen expression host.

[00155] In certain embodiments, recombinant expression vectors are used to amplify and express DNA encoding antibodies, or fragments thereof, against human CD37. Recombinant expression vectors are replicable DNA constructs which have synthetic or cDNA-derived DNA fragments encoding a polypeptide chain of an anti-CD37 antibody, or fragment thereof, operatively linked to suitable transcriptional or translational regulatory elements derived from mammalian, microbial, viral or insect genes. A transcriptional unit generally comprises an assembly of (1) a genetic element or elements having a regulatory role in gene expression, for example, transcriptional promoters or enhancers, (2) a structural or coding sequence which is transcribed into mRNA and translated into protein, and (3) appropriate transcription and translation initiation and termination sequences, as described in detail below. Such regulatory elements can include an operator sequence to control transcription. The ability to replicate in a host, usually conferred by an origin of replication, and a selection gene to facilitate recognition of transformants can additionally be incorporated. DNA regions are operatively linked when they are functionally related to each other. For example, DNA for a signal peptide (secretory leader) is operatively linked to DNA for a polypeptide if it is expressed as a precursor which participates in the secretion of the polypeptide; a promoter is operatively linked to a coding sequence if it controls the transcription of the

sequence; or a ribosome binding site is operatively linked to a coding sequence if it is positioned so as to permit translation. Structural elements intended for use in yeast expression systems include a leader sequence enabling extracellular secretion of translated protein by a host cell. Alternatively, where recombinant protein is expressed without a leader or transport sequence, it can include an N-terminal methionine residue. This residue can optionally be subsequently cleaved from the expressed recombinant protein to provide a final product.

[00156] The choice of expression control sequence and expression vector will depend upon the choice of host. A wide variety of expression host/vector combinations can be employed. Useful expression vectors for eukaryotic hosts, include, for example, vectors comprising expression control sequences from SV40, bovine papilloma virus, adenovirus and cytomegalovirus. Useful expression vectors for bacterial hosts include known bacterial plasmids, such as plasmids from *E. coli*, including pCR 1, pBR322, pMB9 and their derivatives, wider host range plasmids, such as M13 and filamentous single-stranded DNA phages.

[00157] Suitable host cells for expression of a CD37-binding polypeptide or antibody (or a CD37 protein to use as an antigen) include prokaryotes, yeast, insect or higher eukaryotic cells under the control of appropriate promoters. Prokaryotes include gram negative or gram positive organisms, for example *E. coli* or bacilli. Higher eukaryotic cells include established cell lines of mammalian origin as described below. Cell-free translation systems could also be employed. Appropriate cloning and expression vectors for use with bacterial, fungal, yeast, and mammalian cellular hosts are described by Pouwels et al. (*Cloning Vectors: A Laboratory Manual*, Elsevier, N.Y., 1985), the relevant disclosure of which is hereby incorporated by reference. Additional information regarding methods of protein production, including antibody production, can be found, e.g., in U.S. Patent Publication No. 2008/0187954, U.S. Patent Nos. 6,413,746 and 6,660,501, and International Patent Publication No. WO 04009823, each of which is hereby incorporated by reference herein in its entirety.

[00158] Various mammalian or insect cell culture systems are also advantageously employed to express recombinant protein. Expression of recombinant proteins in mammalian cells can be performed because such proteins are generally correctly folded, appropriately modified and completely functional. Examples of suitable mammalian host cell lines include the COS-7 lines of monkey kidney cells, described by Gluzman (*Cell* 23:175, 1981), and other cell lines capable of expressing an appropriate vector including, for example, L cells, C127, 3T3, Chinese hamster ovary (CHO), HeLa and BHK cell lines. Mammalian expression vectors can comprise nontranscribed elements such as an origin of replication, a suitable promoter and enhancer linked to the gene to be expressed, and other 5' or 3' flanking nontranscribed sequences, and 5' or 3' nontranslated sequences, such as necessary ribosome binding sites, a polyadenylation site, splice donor and acceptor sites, and transcriptional termination sequences.

Baculovirus systems for production of heterologous proteins in insect cells are reviewed by Luckow and Summers, *Bio/Technology* 6:47 (1988).

[00159] The proteins produced by a transformed host can be purified according to any suitable method. Such standard methods include chromatography (e.g., ion exchange, affinity and sizing column chromatography), centrifugation, differential solubility, or by any other standard technique for protein purification. Affinity tags such as hexahistidine, maltose binding domain, influenza coat sequence and glutathione-S-transferase can be attached to the protein to allow easy purification by passage over an appropriate affinity column. Isolated proteins can also be physically characterized using such techniques as proteolysis, nuclear magnetic resonance and x-ray crystallography.

[00160] For example, supernatants from systems which secrete recombinant protein into culture media can be first concentrated using a commercially available protein concentration filter, for example, an Amicon or Millipore Pellicon ultrafiltration unit. Following the concentration step, the concentrate can be applied to a suitable purification matrix. Alternatively, an anion exchange resin can be employed, for example, a matrix or substrate having pendant diethylaminoethyl (DEAE) groups. The matrices can be acrylamide, agarose, dextran, cellulose or other types commonly employed in protein purification. Alternatively, a cation exchange step can be employed. Suitable cation exchangers include various insoluble matrices comprising sulfopropyl or carboxymethyl groups. Finally, one or more reversed-phase high performance liquid chromatography (RP-HPLC) steps employing hydrophobic RP-HPLC media, e.g., silica gel having pendant methyl or other aliphatic groups, can be employed to further purify a CD37-binding agent. Some or all of the foregoing purification steps, in various combinations, can also be employed to provide a homogeneous recombinant protein.

[00161] Recombinant protein produced in bacterial culture can be isolated, for example, by initial extraction from cell pellets, followed by one or more concentration, salting-out, aqueous ion exchange or size exclusion chromatography steps. High performance liquid chromatography (HPLC) can be employed for final purification steps. Microbial cells employed in expression of a recombinant protein can be disrupted by any convenient method, including freeze-thaw cycling, sonication, mechanical disruption, or use of cell lysing agents.

[00162] Methods known in the art for purifying antibodies and other proteins also include, for example, those described in U.S. Patent Publication No. 2008/0312425, 2008/0177048, and 2009/0187005, each of which is hereby incorporated by reference herein in its entirety.

[00163] In certain embodiments, the CD37-binding agent is a polypeptide that is not an antibody. A variety of methods for identifying and producing non-antibody polypeptides that bind with high affinity to a protein target are known in the art. See, e.g., Skerra, *Curr. Opin. Biotechnol.*, 18:295-304 (2007), Hosse et al., *Protein Science*, 15:14-27 (2006), Gill et al., *Curr. Opin. Biotechnol.*, 17:653-658 (2006), Nygren, *FEBS J.*, 275:2668-76 (2008), and Skerra, *FEBS J.*, 275:2677-83 (2008), each of which is incorporated by

reference herein in its entirety. In certain embodiments, phage display technology has been used to identify/produce the CD37-binding polypeptide. In certain embodiments, the polypeptide comprises a protein scaffold of a type selected from the group consisting of protein A, a lipocalin, a fibronectin domain, an ankyrin consensus repeat domain, and thioredoxin.

[00164] In some embodiments, the agent is a non-protein molecule. In certain embodiments, the agent is a small molecule. Combinatorial chemistry libraries and techniques useful in the identification of non-protein CD37-binding agents are known to those skilled in the art. See, e.g., Kennedy et al., *J. Comb. Chem.*, 10:345-354 (2008), Dolle et al., *J. Comb. Chem.*, 9:855-902 (2007), and Bhattacharyya, *Curr. Med. Chem.*, 8:1383-404 (2001), each of which is incorporated by reference herein in its entirety. In certain further embodiments, the agent is a carbohydrate, a glycosaminoglycan, a glycoprotein, or a proteoglycan.

[00165] In certain embodiments, the agent is a nucleic acid aptamer. Aptamers are polynucleotide molecules that have been selected (e.g., from random or mutagenized pools) on the basis of their ability to bind to another molecule. In some embodiments, the aptamer comprises a DNA polynucleotide. In certain alternative embodiments, the aptamer comprises an RNA polynucleotide. In certain embodiments, the aptamer comprises one or more modified nucleic acid residues. Methods of generating and screening nucleic acid aptamers for binding to proteins are well known in the art. See, e.g., U.S. Patent No. 5,270,163, U.S. Patent No. 5,683,867, U.S. Patent No. 5,763,595, U.S. Patent No. 6,344,321, U.S. Patent No. 7,368,236, U.S. Patent No. 5,582,981, U.S. Patent No. 5,756,291, U.S. Patent No. 5,840,867, U.S. Patent No. 7,312,325, U.S. Patent No. 7,329,742, International Patent Publication No. WO 02/077262, International Patent Publication No. WO 03/070984, U.S. Patent Application Publication No. 2005/0239134, U.S. Patent Application Publication No. 2005/0124565, and U.S. Patent Application Publication No. 2008/0227735, each of which is incorporated by reference herein in its entirety.

III. Immunoconjugates



[00166] The present invention is also directed to conjugates (also referred to herein as immunoconjugates), comprising the anti-CD37 antibodies, antibody fragments, and their functional equivalents as disclosed herein, linked or conjugated to a drug or prodrug. Suitable drugs or prodrugs are known in the art. The drugs or prodrugs can be cytotoxic agents. The cytotoxic agent used in the cytotoxic conjugate of the present invention can be any compound that results in the death of a cell, or induces cell death, or in some manner decreases cell viability, and includes, for example, maytansinoids and maytansinoid analogs. Other suitable cytotoxic agents are for example benzodiazepines, taxoids, CC-1065 and CC-1065 analogs, duocarmycins and duocarmycin analogs, enediynes, such as calicheamicins, dolastatin and dolastatin analogs including auristatins, tomaymycin derivatives, leptomyacin derivatives, methotrexate, cisplatin, carboplatin, daunorubicin, doxorubicin, vincristine, vinblastine, melphalan, mitomycin C, chlorambucil and morpholino doxorubicin.

[00167] Such conjugates can be prepared by using a linking group in order to link a drug or prodrug to the antibody or functional equivalent. Suitable linking groups are well known in the art and include, for example, disulfide groups, thioether groups, acid labile groups, photolabile groups, peptidase labile groups and esterase labile groups.

[00168] The drug or prodrug can, for example, be linked to the anti-CD37 antibody or fragment thereof through a disulfide bond. The linker molecule or crosslinking agent comprises a reactive chemical group that can react with the anti-CD37 antibody or fragment thereof. The reactive chemical groups for reaction with the cell-binding agent can be *N*-succinimidyl esters and *N*-sulfosuccinimidyl esters. Additionally the linker molecule comprises a reactive chemical group, which can be a dithiopyridyl group that can react with the drug to form a disulfide bond. Linker molecules include, for example, *N*-succinimidyl 3-(2-pyridyldithio) propionate (SPDP) (see, e.g., Carlsson et al., *Biochem. J.*, 173: 723-737 (1978)), *N*-succinimidyl 4-(2-pyridyldithio)butanoate (SPDB) (see, e.g., U.S. Patent No. 4,563,304), *N*-succinimidyl 4-(2-pyridyldithio)2-sulfobutanoate (sulfo-SPDB) (see US Publication No. 20090274713), *N*-succinimidyl 4-(2-pyridyldithio) pentanoate (SPP) (see, e.g., CAS Registry number 341498-08-6), 2-iminothiolane, or acetylsuccinic anhydride. For example, the antibody or cell binding agent can be modified with crosslinking reagents and the antibody or cell binding agent containing free or protected thiol groups thus derived is then reacted with a disulfide- or thiol-containing maytansinoid to produce conjugates. The conjugates can be purified by chromatography, including but not limited to HPLC, size-exclusion, adsorption, ion exchange and affinity capture, dialysis or tangential flow filtration.

[00169] In another aspect of the present invention, the anti-CD37 antibody is linked to cytotoxic drugs via disulfide bonds and a polyethylene glycol spacer in enhancing the potency, solubility or the efficacy of the immunoconjugate. Such cleavable hydrophilic linkers are described in WO2009/0134976. The additional benefit of this linker design is the desired high monomer ratio and the minimal aggregation of the antibody-drug conjugate. Specifically contemplated in this aspect are conjugates of cell-binding agents and drugs linked via disulfide group (-S-S-) bearing polyethylene glycol spacers ((CH₂CH₂O)_{n=1-14}) with a narrow range of drug load of 2-8 are described that show relatively high potent biological activity toward cells and have the desired biochemical properties of high conjugation yield and high monomer ratio with minimal protein aggregation.

[00170] Specifically contemplated in this aspect is an anti-CD37 antibody drug conjugate of formula (I) or a conjugate of formula (I')



wherein:

[00171] CB represents an anti-CD37 antibody or fragment;

[00172] D represents a drug;

[00173] X represents an aliphatic, an aromatic or a heterocyclic unit attached to the cell-binding agent via a thioether bond, an amide bond, a carbamate bond, or an ether bond;

[00174] Y represents an aliphatic, an aromatic or a heterocyclic unit attached to the drug via a disulfide bond;

[00175] l is 0 or 1;

[00176] m is an integer from 2 to 8; and

[00177] n is an integer from 1 to 24.

[00178] In some embodiments, m is an integer from 2 to 6.

[00179] In some embodiments, m is an integer from 3 to 5.



[00180] In some embodiments, n is an integer from 2 to 8. Alternatively, as disclosed in, for example, U.S. Patent No. 6,441,163 and 7,368,565, the drug can be first modified to introduce a reactive ester suitable to react with a cell-binding agent. Reaction of these drugs containing an activated linker moiety with a cell-binding agent provides another method of producing a cell-binding agent drug conjugate. Maytansinoids can also be linked to anti-CD37 antibody or fragment using PEG linking groups, as set forth for example in U.S. Patent 6,716,821. These PEG non-cleavable linking groups are soluble both in water and in non-aqueous solvents, and can be used to join one or more cytotoxic agents to a cell binding agent. Exemplary PEG linking groups include heterobifunctional PEG linkers that react with cytotoxic agents and cell binding agents at opposite ends of the linkers through a functional sulfhydryl or disulfide group at one end, and an active ester at the other end. As a general example of the synthesis of a cytotoxic conjugate using a PEG linking group, reference is again made to U.S. Patent 6,716,821 which is incorporated entirely by reference herein. Synthesis begins with the reaction of one or more cytotoxic agents bearing a reactive PEG moiety with a cell-binding agent, resulting in displacement of the terminal active ester of each reactive PEG moiety by an amino acid residue of the cell binding agent, to yield a cytotoxic conjugate comprising one or more cytotoxic agents covalently bonded to a cell binding agent through a PEG linking group. Alternatively, the cell binding can be modified with the bifunctional PEG crosslinker to introduce a reactive disulfide moiety (such as a pyridyldisulfide), which can then be treated with a thiol-containing maytansinoid to provide a conjugate. In another method, the cell binding can be modified with the bifunctional PEG crosslinker to introduce a thiol moiety which can then be treated with a reactive disulfide-containing maytansinoid (such as a pyridyldisulfide), to provide a conjugate.

[00181] Antibody-maytansinoid conjugates with non-cleavable links can also be prepared. Such crosslinkers are described in the art (see US Publication No. 20050169933) and include but are not limited to, *N*-succinimidyl 4-(maleimidomethyl) cyclohexanecarboxylate (SMCC). In some embodiments, the antibody is modified with crosslinking reagents such as succinimidyl 4-(*N*-maleimidomethyl)-cyclohexane-1-carboxylate (SMCC), sulfo-SMCC, maleimidobenzoyl-*N*-hydroxysuccinimide ester

(MBS), sulfo-MBS or succinimidyl-iodoacetate, as described in the literature, to introduce 1-10 reactive groups (Yoshitake et al, Eur. J. Biochem., 101:395-399 (1979); Hashida et al, J. Applied Biochem., 56-63 (1984); and Liu et al, Biochem., 18:690-697 (1979)). The modified antibody is then reacted with the thiol-containing maytansinoid derivative to produce a conjugate. The conjugate can be purified by gel filtration through a Sephadex G25 column or by dialysis or tangential flow filtration. The modified antibodies are treated with the thiol-containing maytansinoid (1 to 2 molar equivalent/maleimido group) and antibody-maytansinoid conjugates are purified by gel filtration through a Sephadex G-25 column, chromatography on a ceramic hydroxyapatite column, dialysis or tangential flow filtration or a combination of methods thereof. Typically, an average of 1-10 maytansinoids per antibody are linked. One method is to modify antibodies with succinimidyl 4-(N-maleimidomethyl)-cyclohexane-1-carboxylate (SMCC) to introduce maleimido groups followed by reaction of the modified antibody with a thiol-containing maytansinoid to give a thioether-linked conjugate. Again conjugates with 1 to 10 drug molecules per antibody molecule result. Maytansinoid conjugates of antibodies, antibody fragments, and other proteins are made in the same way.

[00182] In another aspect of the invention, the CD37 antibody is linked to the drug via a non-cleavable bond through the intermediacy of a PEG spacer. Suitable crosslinking reagents comprising hydrophilic PEG chains that form linkers between a drug and the anti-CD37 antibody or fragment are also well known in the art, or are commercially available (for example from Quanta Biodesign, Powell, Ohio). Suitable PEG-containing crosslinkers can also be synthesized from commercially available PEGs themselves using standard synthetic chemistry techniques known to one skilled in the art. The drugs can be reacted with bifunctional PEG-containing cross linkers to give compounds of the following formula, $Z-X_1-(CH_2-CH_2-O)_n-Y_p-D$, by methods described in detail in US Patent Publication 20090274713 and in WO2009/0134976, which can then react with the cell binding agent to provide a conjugate. Alternatively, the cell binding can be modified with the bifunctional PEG crosslinker to introduce a thiol-reactive group (such as a maleimide or haloacetamide) which can then be treated with a thiol-containing maytansinoid to provide a conjugate. In another method, the cell binding can be modified with the bifunctional PEG crosslinker to introduce a thiol moiety which can then be treated with a thiol-reactive maytansinoid (such as a maytansinoid bearing a maleimide or haloacetamide), to provide a conjugate.

[00183] Accordingly, another aspect of the present invention is an anti-CD37 antibody drug conjugate of formula (II) or of formula (II'):



wherein, CB represents an anti-CD37 antibody or fragment;

[00184] D represents a drug;

[00185] X represents an aliphatic, an aromatic or a heterocyclic unit bonded to the cell-binding agent via a thioether bond, an amide bond, a carbamate bond, or an ether bond;

[00186] Y represents an aliphatic, an aromatic, or a heterocyclic unit bonded to the drug via a covalent bond selected from the group consisting of a thioether bond, an amide bond, a carbamate bond, an ether bond, an amine bond, a carbon-carbon bond and a hydrazone bond;

[00187] l is 0 or 1;

[00188] p is 0 or 1;

[00189] m is an integer from 2 to 15; and

[00190] n is an integer from 1 to 2000.

[00191] In some embodiments, m is an integer from 2 to 8; and

[00192] In some embodiments, n is an integer from 1 to 24.

[00193] In some embodiments, m is an integer from 2 to 6.

[00194] In some embodiments, m is an integer from 3 to 5.

[00195] In some embodiments, n is an integer from 2 to 8. Examples of suitable PEG-containing linkers include linkers having an *N*-succinimidyl ester or *N*-sulfosuccinimidyl ester moiety for reaction with the anti-CD37 antibody or fragment thereof, as well as a maleimido- or haloacetyl-based moiety for reaction with the compound. A PEG spacer can be incorporated into any crosslinker known in the art by the methods described herein.

[00196] Many of the linkers disclosed herein are described in detail in U.S. Patent Publication Nos. 20050169933 and 20090274713, and in WO2009/0134976; the contents of which are entirely incorporated herein by reference.

[00197] The present invention includes aspects wherein about 2 to about 8 drug molecules ("drug load"), for example, maytansinoid, are linked to an anti-CD37 antibody or fragment thereof. "Drug load", as used herein, refers to the number of drug molecules (e.g., a maytansinoid) that can be attached to a cell binding agent (e.g., an anti-CD37 antibody or fragment thereof). In one aspect, the number of drug molecules that can be attached to a cell binding agent can average from about 2 to about 8 (e.g., 1.9, 2.0, 2.1, 2.2, 2.3, 2.4, 2.5, 2.6, 2.7, 2.8, 2.9, 3.0, 3.1, 3.2, 3.3, 3.4, 3.5, 3.6, 3.7, 3.8, 3.9, 4.0, 4.1, 4.2, 4.3, 4.4, 4.5, 4.6, 4.7, 4.8, 4.9, 5.0, 5.1, 5.2, 5.3, 5.4, 5.5, 5.6, 5.7, 5.8, 5.9, 6.0, 6.1, 6.2, 6.3, 6.4, 6.5, 6.6, 6.7, 6.8, 6.9, 7.0, 7.1, 7.2, 7.3, 7.4, 7.5, 7.6, 7.7, 7.8, 7.9, 8.0, 8.1). $N^{2'}$ -deacetyl- $N^{2''}$ -(3-mercapto-1-oxopropyl)-maytansine (DM1) and $N^{2'}$ -deacetyl- $N^{2''}$ -(4-mercapto-4-methyl-1-oxopentyl) maytansine (DM4) can be used.

[00198] Thus, in one aspect, an immunoconjugate comprises 1 maytansinoid per antibody. In another aspect, an immunoconjugate comprises 2 maytansinoids per antibody. In another aspect, an immunoconjugate comprises 3 maytansinoids per antibody. In another aspect, an immunoconjugate comprises 4 maytansinoids per antibody. In another aspect, an immunoconjugate comprises 5

maytansinoids per antibody. In another aspect, an immunoconjugate comprises 6 maytansinoids per antibody. In another aspect, an immunoconjugate comprises 7 maytansinoids per antibody. In another aspect, an immunoconjugate comprises 8 maytansinoids per antibody.

[00199] In one aspect, an immunoconjugate comprises about 1 to about 8 maytansinoids per antibody. In another aspect, an immunoconjugate comprises about 2 to about 7 maytansinoids per antibody. In another aspect, an immunoconjugate comprises about 2 to about 6 maytansinoids per antibody. In another aspect, an immunoconjugate comprises about 2 to about 5 maytansinoids per antibody. In another aspect, an immunoconjugate comprises about 3 to about 5 maytansinoids per antibody. In another aspect, an immunoconjugate comprises about 3 to about 4 maytansinoids per antibody.

[00200] In one aspect, a composition comprising immunoconjugates has an average of about 2 to about 8 (e.g., 1.9, 2.0, 2.1, 2.2, 2.3, 2.4, 2.5, 2.6, 2.7, 2.8, 2.9, 3.0, 3.1, 3.2, 3.3, 3.4, 3.5, 3.6, 3.7, 3.8, 3.9, 4.0, 4.1, 4.2, 4.3, 4.4, 4.5, 4.6, 4.7, 4.8, 4.9, 5.0, 5.1, 5.2, 5.3, 5.4, 5.5, 5.6, 5.7, 5.8, 5.9, 6.0, 6.1, 6.2, 6.3, 6.4, 6.5, 6.6, 6.7, 6.8, 6.9, 7.0, 7.1, 7.2, 7.3, 7.4, 7.5, 7.6, 7.7, 7.8, 7.9, 8.0, 8.1) drug molecules (e.g., maytansinoids) attached per antibody. In one aspect, a composition comprising immunoconjugates has an average of about 1 to about 8 drug molecules (e.g., maytansinoids) per antibody. In one aspect, a composition comprising immunoconjugates has an average of about 2 to about 7 drug molecules (e.g., maytansinoids) per antibody. In one aspect, a composition comprising immunoconjugates has an average of about 2 to about 6 drug molecules (e.g., maytansinoids) per antibody. In one aspect, a composition comprising immunoconjugates has an average of about 2 to about 5 drug molecules (e.g., maytansinoids) per antibody. In one aspect, a composition comprising immunoconjugates has an average of about 3 to about 5 drug molecules (e.g., maytansinoids) per antibody. In one aspect, a composition comprising immunoconjugates has an average of about 3 to about 4 drug molecules (e.g., maytansinoids) per antibody.

[00201] In one aspect, a composition comprising immunoconjugates has an average of about 2 ± 0.5 , about 3 ± 0.5 , about 4 ± 0.5 , about 5 ± 0.5 , about 6 ± 0.5 , about 7 ± 0.5 , or about 8 ± 0.5 drug molecules (e.g., maytansinoids) attached per antibody. In one aspect, a composition comprising immunoconjugates has an average of about 3.5 ± 0.5 drug molecules (e.g., maytansinoids) per antibody.

[00202] The anti-CD37 antibody or fragment thereof can be modified by reacting a bifunctional crosslinking reagent with the anti-CD37 antibody or fragment thereof, thereby resulting in the covalent attachment of a linker molecule to the anti-CD37 antibody or fragment thereof. As used herein, a "bifunctional crosslinking reagent" is any chemical moiety that covalently links a cell-binding agent to a drug, such as the drugs described herein. In another method, a portion of the linking moiety is provided by the drug. In this respect, the drug comprises a linking moiety that is part of a larger linker molecule that is used to join the cell-binding agent to the drug. For example, to form the maytansinoid DM1, the side chain at the C-3 hydroxyl group of maytansine is modified to have a free sulfhydryl group (SH). This

thiolated form of maytansine can react with a modified cell-binding agent to form a conjugate. Therefore, the final linker is assembled from two components, one of which is provided by the crosslinking reagent, while the other is provided by the side chain from DM1.


[00203] The drug molecules can also be linked to the antibody molecules through an intermediary carrier molecule such as serum albumin.

[00204] As used herein, the expression "linked to a cell-binding agent" or "linked to an anti-CD37 antibody or fragment" refers to the conjugate molecule comprising at least one drug derivative bound to a cell-binding agent anti-CD37 antibody or fragment via a suitable linking group, or a precursor thereof. One linking group is SMCC.


[00205] In certain embodiments, cytotoxic agents useful in the present invention are maytansinoids and maytansinoid analogs. Examples of suitable maytansinoids include esters of maytansinol and maytansinol analogs. Included are any drugs that inhibit microtubule formation and that are highly toxic to mammalian cells, as are maytansinol and maytansinol analogs.

[00206] Examples of suitable maytansinol esters include those having a modified aromatic ring and those having modifications at other positions. Such suitable maytansinoids are disclosed in U.S. Patent Nos. 4,424,219; 4,256,746; 4,294,757; 4,307,016; 4,313,946; 4,315,929; 4,331,598; 4,361,650; 4,362,663; 4,364,866; 4,450,254; 4,322,348; 4,371,533; 5,208,020; 5,416,064; 5,475,092; 5,585,499; 5,846,545; 6,333,410; 7,276,497 and 7,473,796.


[00207] In a certain embodiment, the immunoconjugates of the invention utilize the thiol-containing maytansinoid (DM1), formally termed $N^{2'}$ -deacetyl- $N^{2'}$ -(3-mercapto-1-oxopropyl)-maytansine, as the cytotoxic agent. DM1 is represented by the following structural formula (III):



[00208] In another embodiment, the conjugates of the present invention utilize the thiol-containing maytansinoid $N^{2'}$ -deacetyl- $N^{2'}$ -(4-methyl-4-mercapto-1-oxopentyl)-maytansine (e.g., DM4) as the cytotoxic agent. DM4 is represented by the following structural formula (IV):




[00209] Another maytansinoid comprising a side chain that contains a sterically hindered thiol bond is *N*²-deacetyl-*N*²-(4-mercapto-1-oxopentyl)-maytansine (termed DM3), represented by the following structural formula (V):



[00210] Each of the maytansinoids taught in US Patent No. 5,208,020 and 7,276,497, can also be used in the conjugate of the present invention. In this regard, the entire disclosure of 5,208,020 and 7,276,697 is incorporated herein by reference.


[00211] Many positions on maytansinoids can serve as the position to chemically link the linking moiety. For example, the C-3 position having a hydroxyl group, the C-14 position modified with hydroxymethyl, the C-15 position modified with hydroxy and the C-20 position having a hydroxy group are all expected to be useful. In some embodiments, the C-3 position serves as the position to chemically link the linking moiety, and in some particular embodiments, the C-3 position of maytansinol serves as the position to chemically link the linking moiety.

[00212] Structural representations of some conjugates are shown below:




Ab-PEG-Mal-DM1/DM4

(VI)






Ab-PEG4-Mal-DM1



(VII)



Ab-PEG-SIA-DM1/DM4

(VIII)






[00213] Several descriptions for producing such antibody-maytansinoid conjugates are provided in U.S. Patent Nos. 6,333,410, 6,441,163, 6,716,821, and 7,368,565, each of which is incorporated herein in its entirety.

[00214] In general, a solution of an antibody in aqueous buffer can be incubated with a molar excess of maytansinoids having a disulfide moiety that bears a reactive group. The reaction mixture can be quenched by addition of excess amine (such as ethanolamine, taurine, etc.). The maytansinoid-antibody conjugate can then be purified by gel filtration.

[00215] The number of maytansinoid molecules bound per antibody molecule can be determined by measuring spectrophotometrically the ratio of the absorbance at 252 nm and 280 nm. The average number of maytansinoid molecules/antibody can be, for example, about 1-10, 2-5, 3-4, or about 3.5. In one aspect, the average number of maytansinoid molecules/antibody is about 3.5 ± 0.5 .

[00216] Anthracycline compounds, as well as derivatives, intermediates and modified versions thereof, can also be used to prepare anti-CD37 immunoconjugates. For example, doxorubicin, doxorubicin derivatives, doxorubicin intermediates, and modified doxorubicins can be used in anti-CD37 conjugates. Exemplary compounds are described in WO 2010/009124, which is herein incorporated by reference in its entirety. Such compounds include, for example, compounds of the following formula:



wherein R_1 is a hydrogen atom, hydroxy or methoxy group and R_2 is a C_1 - C_5 alkoxy group, or a pharmaceutically acceptable salt thereof.

[00217] Conjugates of antibodies with maytansinoid or other drugs can be evaluated for their ability to suppress proliferation of various unwanted cell lines *in vitro*. For example, cell lines such as the human lymphoma cell line Daudi and the human lymphoma cell line Ramos, can easily be used for the assessment of cytotoxicity of these compounds. Cells to be evaluated can be exposed to the compounds for 4 to 5 days and the surviving fractions of cells measured in direct assays by known methods. IC_{50} values can then be calculated from the results of the assays.

[00218] The immunoconjugates can, according to some embodiments described herein, be internalized into cells. The immunoconjugate, therefore, can exert a therapeutic effect when it is taken up by, or internalized, by a CD37-expressing cell. In some particular embodiments, the immunoconjugate comprises an antibody, antibody fragment, or polypeptide, linked to a cytotoxic agent by a cleavable linker, and the cytotoxic agent is cleaved from the antibody, antibody fragment, or polypeptide, wherein it is internalized by a CD37-expressing cell.

[00219] In some embodiments, the immunoconjugates are capable of depleting B-cells, e.g. autoreactive B-cells. For example, in some embodiments, treatment with an immunoconjugate results in a depletion of at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, or at least about 75% of B-cells.

[00220] In another aspect of the invention siRNA molecules can be linked to the antibodies of the present invention instead of a drug. siRNAs can be linked to the antibodies of the present invention by methods commonly used for the modification of oligonucleotides (see, for example, US Patent Publications 20050107325 and 20070213292). Thus the siRNA in its 3' or 5'-phosphorimidite form can be reacted with one end of the crosslinker bearing a hydroxyl functionality to give an ester bond between the siRNA and the crosslinker. Similarly reaction of the siRNA phosphoramidite with a crosslinker bearing a terminal amino group results in linkage of the crosslinker to the siRNA through an amine. Alternatively, the siRNA can be derivatized by standard chemical methods to introduce a thiol group.

This thiol-containing siRNA can be reacted with an antibody, that has been modified to introduce an active disulfide or maleimide moiety, to produce a cleavable or non cleavable conjugate. Between 1 - 20 siRNA molecules can be linked to an antibody by this method.

III. Polynucleotides

[00221] In certain embodiments, the invention encompasses polynucleotides comprising polynucleotides that encode a polypeptide that specifically binds CD37 or a fragment of such a polypeptide. For example, the invention provides a polynucleotide comprising a nucleic acid sequence that encodes an antibody to a human CD37 or encodes a fragment of such an antibody. The polynucleotides of the invention can be in the form of RNA or in the form of DNA. DNA includes cDNA, genomic DNA, and synthetic DNA; and can be double-stranded or single-stranded, and if single stranded can be the coding strand or non-coding (anti-sense) strand.

[00222] In certain embodiments, the polynucleotides are isolated. In certain embodiments, the polynucleotides are substantially pure.

[00223] The invention provides a polynucleotide comprising a polynucleotide encoding a polypeptide comprising a sequence selected from the group consisting of SEQ ID NOs:4-120.

[00224] The invention further provides a polynucleotide comprising a sequence selected from those shown in Tables 7-10 below.

Table 7: Variable heavy chain polynucleotide sequences

Antibody	VH Polynucleotide Sequence (SEQ ID NO)
muCD37-3	cagggtgcagggtgaaggagtcaggacctggcctggtggcgccctcacagagcctgtccattacatgcactg tctcagggftctcaftaaccacctctggtgtaagctgggttcgccagcctccaggaaagggtctggagtg gctgggagtaatatggggtgacgggagcacaactatcattcagctctcaatccagactgagcatcaag aaggatcactccaagagccaagtttcttaaaactgaacagtctgcaactgatgacacagccacgtact actgtgccaaaggaggctactcgttggtcactggggccaagggactctggtcacagtctctgca (SEQ ID NO:121)
chCD37-3	aagcttgccaccatggctgctcctggcactgctcctctgctggtgacataccaagctgtgctcctacacaggtgcaggtg aaggagtcaggacctggcctggtggcgccctcacagagcctgtccattacatgcactgtctcagggftctcaftaaccac ctctggtgtaagctgggttcgccagcctccaggaaagggtctggagtggtgggagtaatatggggtgacgggagcac aaactatcattcagctctcaatccagactgagcatcaagaaggatcactccaagagccaagtttcttaaaactgaacagt ctgcaactgatgacacagccacgtactactgtgccaaggaggctactcgttggtcactggggccaagggactctgg tcactctctcagcctctacgaaggccc (SEQ ID NO:122)
huCD37-3v1.0	aagcttgccaccatgggtggagctgattattctgtttctggtggccaccgccaccgggtgctcactcacaagtccaagtc caagaatctggtccaggtctggtggccccctccaaactctgagcatcactgtaccgtttctggttttagccttaccacctc tggtgtgagttgggtacgccaaccaccggtaagggtctcgaatggctgggtgtaatctggggtgatggtccacaatt accatcctccctcaagtccegccttagcatcaaaaaggatcacagcaaaagtcaagtttctgaaactgaatagctgac agcagccgatacagccactactattgcccagggtggttatagcttgcacactggggccaaggtaccctcgttaccgt ctcctcaactagtagccaaggccc (SEQ ID NO:123)
huCD37-3v1.1	aagcttgccaccatgggctggagctgtatcattctgtttctggtggcgacagctactggggccactcccaagtcaggta caagagtcgggctggtggtgcaccaagccagacctctctatcactgtaccgttagcgggttctctctgacaacc agtcaactgagttgggtgagcagccaccaggaaagggactcagtggtgggggtgattggggcagggcagca

	caaacatcattccagctttaaactcgggttgccatfaaaaaagaccatagtaaactcaagtttctgaaactcaatagcctgacagccgcagacactgctacgtattactgocgcaaaaggaggatagcagctctggctcactggggacaggggacctggtagaccgtgctatccgcatcaacaaggccc (SEQ ID NO:124)
muCD37-12	cagatccagttggtgcagctcggacctgagctgaagaagcctggagagacagtcagatcctcctgcaaggcttctgggtataccttcacaaglatggaatgaactgggtgaagcaggctcaaggaaagggtttaaagtgatgggctggataaacaccaactggagagtcagaaatgctgaagaatcaaggagcgggttgcctctctttggaaacctctccagcactgctatttgcagatcaacaacctcaaatatgaggacacggctacatatttctgtggaaggggcacggtagtagggactggggccaaggcaccactctcacagtctctca (SEQ ID NO:125)
chCD37-12	aagcttgcaccatgggggtgctcagcataatcctcttctggctcgtactgctaccgggtgctcactcacagatcagctggttcaaaagtggcccagagctgaaaaagccaggggaaacagtgaaaaataagttgcaaggcatccgggtfacacttcacaaagtacggcatgaactgggtcaagcaggcccaggggcaaggggctcaaatggatgggttggatcaatacacaactggcgagtctaggaatgctgaggagtttaaggccgggttgccttcagcctggagacaagtgccagcacagctacctgcaaatcaaatctgaagatgaggatagcaacciatctctgcccggcggcactgctgctgagactggggacaaggataccaactgactgataccagtgccagcactaaggccc (SEQ ID NO:126)
muCD37-38	gatgtgcagctcaggagtcaggacctgacctgggtgaaacctctcagtcacttccactcactgcactgtcactggctactccatcaccagttggttggctggcactggatccggcagttccaggaaacaagctggaatggatggcctacatactacagtggtggcactgactacaacceatctcfaaaagtcgaatctctactcagagacactccaagaaccagttctctcctgcccgttgagttctgtgactactgaggacacagccacatattactgtgcaagaggctactatggttacgggcccgtggttggactggggccaagggactctgtgctactgtctctgca (SEQ ID NO:127)
chCD37-38	aagcttgcaccatgggctggagttgatcattctgttttgggtggccaccgccactggagtcattccaagtgcactccaggaatctggccctgacctggttaagccatcagagocctcctgacctgactgttacagatactcaatcacatcaggctttggctggcactggatcagacaattcccgggaacaagttggaatggatggcttacattctgtatagcgggggtaccgattacaatccttccctcaagagccgaatctctataccaggggatacaagaagaaccaatttttctccgctcagctctgtgactaccgaagataccgctacttactattgtgccaggggctactatggatattggctcagttgctatggggccagggaacctggtgactgtgagcgtctcctctaccaaggccc (SEQ ID NO:128)
huCD37-38	aagcttgcaccatggggtggagctgcatcattcttctctggtcgtactgcaactggagtcactcacaggtccagctgc aagagtcoggtcctgggcttggtaaaaccagccagtcctcagctcactgactgtctctggtactctattaccagtggttccgctggcaatggattagggacttcccggtaaggggctggagtgatggcatatatectgtacagcggaggaaaccgattacaaccaagctgaagagcaggatcagcattaccgggacacaagcaaaaaccagtttctctcctggctctctagtggtacagctgcagacaccgctacttactattgtctcggggtactatggctatgggctgggttgggttattggggacaaggcactctgtgaccgtgagcagcctcaacaaggccc (SEQ ID NO:129)
muCD37-50	gatgtgcagctcaggagtcaggacctgacctgfgaaacctctcagtcacttccactcactgcactgtcactggctactccatcaccagttggttgcctggcactggatccggcagttccaggaaacaactggaatggatgggctacatactacagtggtgacactgtctacageccatctcfaaaagtcgaatctctactcagagacacatccaagaaccacttctcctgagttgaatctgtgactactgaggacacagccacatattactgtgcaagaggatctatggttacggcctggttgcctactggggccaagggactctgtgctactgtctctgca (SEQ ID NO:130)
huCD37-50	aagcttgcaccatgggggtgctcagcataatccttctctggtgctactgctaccggagtcactcacaggtgcagctgcaggagtcogggcccggcctgcicaagcctctcagagctgagctgactgtactgttctggctacagcataaccagcggtttcgctggcactggafagacagcagatcccggcaacaactggagtgatgggafacatactgtactcaggctcaactgctattcccctcctgaaatcccggatcagttatcccggtagacttcaagaaccatttttctcagctgaacagcgttaccgagctgacactgcaacctactctgtcccgggatafialggafacggagctgggtctgctactggggccaaggcaccctcgtactgtgagtgctctccaccaaggccc (SEQ ID NO:193)
muCD37-51	gatgtgcagctcaggagtcaggacctgacctgfgaaacctctcagtcacttccactcactgcactgtcactggctactccatcaccagttggttgcctggcactggafccggcagttccaggaaacaactggaatggatgggctacatactacagtggtgacactaacacagccatctctcaaaagtogaatctctactcagagacacatccaagaaccagttctcctgagttgaatctgtgactactgaggacacagccacatattactgtgcaagaggatctatggttccggcctggttgggtactggggccaagggactctgtgctactgtctctgca (SEQ ID NO:131)
huCD37-51	Aagcttgcaccatggggtggctcagcactcctgctctggtggcactgcccactggcgtgcaatcagaagttcagttgtgagtcogggcccagaagtgctgaaccggcgaatcactctcctgactgtgactgtcaggttatagcatcagcagc

	ggctttgcttggcactggattcggcagttccaggcaagggactggaatggatgggctacatccattacagtggctcaac caattacagccclagcctgcagggccgaatctctattaccagggatggtctattaaccagttttctcgcagcttaattccgt gactgcctctgacacagcaactactattgcgcccgtggctactacgggtcggagcctggtttgatactggggcaggg caccctgctactgctcagccgctctaccaaggggccc (SEQ ID NO:194)
muCD37-56	gatgtgcagcttcaggagtcaggacctgacctggtgaaacctctcagtcactttcactcaactgcactg tactggctactccatcaccagtggtttgctggcactggatccggcagttccaggaaacaaactgga atggatgggctacatacactacagtggtggcactaactacaaccatctcctcaaaagtcagctctctac actcgagacacatccaagaaccagttctcctgcagttgaattctgtgactactgaggacacagccacatattactgtgcaa gaggctactatggttcgggctggtttgcttactggggccaagggactctggtccc tctctgca (SEQ ID NO:132)
huCD37-56	aagcttgccaccatggggtggagctgcaatctctgtctcgtcggaccgcaaccggcgtccactcccaggtgcagct gcaagaaagcgggcccaggattggtaaaacctccagctctctgagcttactgtaccgtatctggatacagtatcacatct ggcttcgctggcattggattcggcagttcccggcaaggggcttgagtgatgggtatattcattattctggaggtacca actacaaccttccctgaagagtcgagctcaatfaccagggacacttcaagaaccaattcttttgcagcttaattcagtg accgctgcccagaccgctacttactgcccgggctactatgggttggctgctggtcgcctactggggccaggg accctgctcccctctctgctgctccacaaggggccc (SEQ ID NO:133)
muCD37-57	gatgtgcagcttcaggagtcaggacctgacctggtgaaacctctcagtcactttcactcaactgcactg tactggctactccatcaccagtggtttgctggcactggatccggcagttccaggaaacaaactgga atggatgggctacatacactacagtggtgactgtctacagccatctcctcaaaagtcgaatctctac actcgagacacatccaagaaccagttctcctgcagttgaattctgtgactactgaggacacagccacatattactgtgcaa gagggtactatggttacggcgcctggttgcctactggggccaagggactctggtcactgtctgca (SEQ ID NO:134)
huCD37-57	aagcttgccaccatggggtggagctgcaatctctgttctggtggccacagcaactggcgttcacagtcagtccaactg caggagagcggccccggactcctgaaacctctcagtcactcagctgacatgactgtgagcggctacagcattacctc aggcttcgctggcattggatcaggcagttcccggaaaaggtctggagtgatgggttacattctgtacagcggcagta cagtgattaccctcctgaaatctaggatatcaatcacagtgatacaagcaaaaatcagttcttctccagctgaactcc gtcaccgcccagacacagcaacctattattgtctcgggatactacggalatggcgcagttctgcctattggggcca ggggacactctgaccgttccggcctccacaaggggccc (SEQ ID NO:135)
252-3	gaggtgcaggtggtggagctctgggggagacttagtgaagcctggagggtccctgaaactctcctgtgcagccttgat tcaattcagtagctatggcatgcttgggtcgcagactccagacaagaggctggagtggtcgaaccattagtagtg gtggtggttacactactctccagacagtggaagggcgcattaccatctccagagacaatgcaagaaaacctgtac ctgcaaatgagcagctgaagctgaggacacagccatgtattactgtgcaagacatagttactacgatactagcgtgcac tactgggtcaaggaaacctcagtcaccgtctctca (SEQ ID NO:182)

Table 8: Variable light chain polynucleotide sequences

Antibody	VL Polynucleotide Sequence (SEQ ID NO)
muCD37-3	gacatccagatgactcagtcctcagcctcccttctgtatctgtgggagaaactgcaccatcacatgctc gagcaagtgagaatattcgcagtaatttagcatggtatcagcagaacagggaaaatctcctcagctcct ggtcaatgttgcaacaaacttagcagatggtgtgccatcaaggtcagtgagcagtgatcaggcacacag tattccctcaagatcaacagcctgcagctctgaagatttgggacttattactgtcaacattattgggta ctacgtggacttccggtggaggcaccacaaactgaaatcaaacct (SEQ ID NO:136)
chCD37-3	gaattcgccaccatgagtggtgcccactcaggctcctggggtgctgctgctggtgcttacagatgccagatgacatccag atgactcagctccagcctcccttctgtatctgtgggagaaactgtaccatcacatgctgagcaagtgagaatattcga gtaatttagcatggtatcagcagaacagggaaaatctcctcagctcctggtcaatgtgcaacaaacttagcagatggtg gcatcaaggtcagtggtcagtgatcaggcacacagttccctcaagatcaacagcctgcagctctgaagatttggga cttattactgtcaacattattgggtactacgtggagctcgtggaggccaagctggaatcaaacgtacg (SEQ ID NO:137)
huCD37-3 (1.0 and 1.1)	gaattcgccaccatgggttggctcctgcatctctgttctcgtggccacagccaccggttctactctgatatacaaatgac tcaagccctccagttgagcgtgaagtgtgggtgaacgcgtaacaatcactgtagagctagtgaaaacatccgcagta atctccatggtaccaacaaaaggcaagtaagtcacctcctctgaaatgtctaccaacctcctgctgaggtgctc

	cttcacgattctctggtcaggttccggtaaccgattatcacttaagatcaactcactccaaccagaagattfeggtacatatta ctgtcaacactactggggtagcaccctggacattcggtaaggtactaagctggaaatcaagcgtacg (SEQ ID NO:138)
muCD37-12	gacattgtgctaacacagtcctctgcttcttagctgtatctctggggcagagggccaccatctcatgca gggcccagecaaaagtgacgtacatctagctatagttatgtactggtccagcagaaaccaggacagcc acccaaactcctcatcaagatgatccaacctagcactctggggtccctgccaggttcagtggtcagtggtg tctgggacagactcaccctcaacatccatctctgiggaggaggaggatactgcaacatattactgtcaac acagttgggagattccgtacacgtfegggagggggaccaaaactggaaataaaacgg (SEQ ID NO:139)
chCD37-12	gaattcgcaccatggggttggctctgtataatcctgttcttggggccaccgctactggcgttcagtgatattgtactact cagtcaccagccagctctggcaggtgctccctggggccagcgtgcccaccatctctgccgggctcacagtcctgtgagcacta gtcttattcctatctctactggttcaacagaagccaggacagccccctaaagctgctgatcaagtaoecctcaacctgc cagcggcgttcccgctagattctctgggtccggtagcggaaactgattcacttgaacatccaccccggtgaggaagagga taccgcacttacattgcaaacactcttgggagattcctacaccttggaggaggaaacaaagctcgaattaaagcgtacg (SEQ ID NO:140)
muCD37-38	caaatgttctcaccagctctccagcaatcatgtctgcactcaccaggggagaaggtcaccatgacctgca gtgccagctcaagtgttaactatgcactggtaccagcagaagtcaggcacctccccaaaagatggat ttatgacacatccaaactggctctggagtcctctgctcagtggtgggtgggtctgggacctttac tctctcaaatcagcagcatggagctgaagatgctgccacttattactgccagcagtggtatagtaacc caccacggtcggggggggaccaagctggaattaaacgg (SEQ ID NO:141)
chCD37-38	gaattcgcaccatgggctggtcctgtatcactctgttctctggtggccacagctacaggtgtcaattcagattgtgctgac ccaateaccagctattatgtccgctagccccggcgagaaagtgacaatgacatgtagcgtactcttctgtgacttacat gcattggatcaacagaagtcaggfaccagtcccaagcgttggatciacgacacatccaaactggcctccggagtcctg ccagggtcagcggaggtgggtccggcaccagttatcactgaccatctctctatggaagctgaagatgctgctacttatta ttgtcaacaatggatttcaacccccccaccttgggtggcggaacaaagctggagatcaagcgtacg (SEQ ID NO:142)
huCD37-38	gaattcgcaccatgggatggtcctgctalttctgttcttggctgccactgctactggcgttactctgacattgtgctcaca cagtcctccagcctcaatgtctgtctccccgggtgagcgggtgaccatgacatgctctgccagttccctcgatataatg attggtatcagcaaaaaccggtaacctctccaaaagatggatctacgacacttcaaaagctgcatcagcgttctctgcca gatttccgggtctgggtctggcacttcatacagctgaccattagttccatggaagctgaagatgcagccacttactgt cagcagtggaattcaaatcctctactctcggcggcggaaacaaactggagataaaagcgtacg (SEQ ID NO:143)
muCD37-50	caaatgttctcaccagctctccagcaatcatgtctgcactcaccaggggagaaggtcaccatgacctgca gtgccacctcaagtgtgacttacatgcactggtaccagcagaagtcaggcacctccccaaaagatggattatgacaca tccaaactgccttattggagtcctgtgctggtcagtggttagtgggtctgggaccttactctctcaaatcagcagcatgg aggctgaagatgctgccacttattactgccagcagtggaagtataaccaccaccgctcggctcggggacaaaagttgga aataaagcgg (SEQ ID NO:144)
huCD37-50	gaattcgcaccatggggttggctcatctattctgttcttgggtgctaccgcaacaggagtacatagtgatagtcctcac ccaaagtcctgtactatgtctgccagcccaggagagcgtgtgaccatgactgtctgcaacctcaagtgtgacatacat gcattggatcagcaaaagcctggccaatccctaaaaggtggatctacgatacttcaatctgccatacgggtgcccgc aagggtctccgggagtggtgagtcaccagttatagctgaccatcagttcaatggaagcagaggatgcagcaacctatt attgtcagcagtggtccgataatccccctacttttgggtcagggtacaaagctggagattaagcgtacg (SEQ ID NO:145)
muCD37-51	caaatgttctcaccagctctccagcaatcatgtctgcactcaccaggggagaaggtcaccatgacctgca gtgccacctcaagtgtgacttacatgcactggtaccagcagaagtcaggcacctccccaaaagatggattatgacaca tccaaactggcttctggagtcctgtctgctcagtggtcagtggtctgggaccttactctctcaaatcagcaacatgg aggctgaagatgctgccacttattactgccagcagtggaagtataaccaccaccgctcggctcggggacaaaagttgga aataaagcgg (SEQ ID NO:146)
huCD37-51	gaattcgcaccatgggatggagctgtatttctgttcttgggtgctactgctactggcgtcattccgagatagtcctcac ccagagcccccaacctagtgctcctccctggggagcagtgactatgactgttccgaccttctcagttacctat gcattggatcagcagaacctggacagctcctcaaacggttgattacgacacctcaacctggctcaggagttctctg taggttcagcggatctgggtctggcacaagtattcactcaccattagttccatggaggccgaagatgccgctacttact tgtcagcagtggtgagcagcaacccccctacattcgggcagggaactaagctggagatcaaacgtacg (SEQ ID NO:147)

muCD37-56	caaattgttctcaccagtcctccagcattcatgtctgcatctccaggggataaggtcaccatgacctgca gtgccagtcaagtgaacttacatgactggtatcagcagaagtcaggcacctccccaaaagatggattatgacacat ccaaactggctctggagtcctgctcgtcagtgccgggtgggtctgggacctctac tctctacaatcagccatggaggctgaagatgctgccacttattactgccagcagtgattagtgacc caccacattcgaaggggaccaagctggaataaaacg (SEQ ID NO:148)
huCD37-56	gaattcgccaccatgggctggtctgtatcctctgttctggtggcaaccgactggtggtcactctgatattgtctgac acagagtcagcctcatgagtgcttctccggagaaaaggtcacaatgactgttcagcttctcctcctcgtacatacatg cattggtaccagcagaagcctgaccagagtcctaagaggtggatctatgatacaagcaatctggctccgggtgccccctc ccgctttcaggcggcgaagcggaaactgactatagcctaccatctcctcaatggaagccgaggacgctgctacatatt actgccagcaatggatcagcgcacctctactttcggacagggaaacaaattgaaattaagcgtacg (SEQ ID NO:149)
muCD37-57	caaattgttctcaccagtcctccagcaatcatgtctgcatctccaggggagaaggtcaccatgacctgca gtgccacctcaagtgtgacttacatgactggtaccagcagaagtcaggcacctccccaaaagatggattatgacaca tccaaactggctctggagtcctgctcgttcagtgccagtggtgctgggacctctactctcacaatcagcagcatgg aggctgaagatgctgccacttattactgccagcagtgagtgataaccaccacgctcggctcggggacaaaagttgga aataaacggg (SEQ ID NO:150)
huCD37-57	gaattcgccaccatggggtggtcctgtattatcctgttctggtcgaaccgccacaggcgtcactccgagatcgtgtga ctcagagcccagccaccatgtccgctccccggggagagagtgacaatgactgttccgccacaagttctgtaacctac atgcattggtaccagcaaaaaccaggacagagtcctcctgctggtgattatgataacctaaactggttcaggcgttctg cccgttttctgtagtgatctgggacttctatagccttaccataagctctatggaagccgaggacccgctacatacta ctgccagcagtgagtgataacccccaccttcgggcagggaaacaaattggagatcaaacgtacg (SEQ ID NO:151)
252-3	gatatccagatgacacagactacatcctcctgtctgctctctgggagacagagtcaccatcagttgcagggc aagtcaggacattagcaattattaaactggatcagcagaaccgatggaactgttaaactcctgatctactac acatcaaaattactcagagtcctcatcaaggttcagtggcagtggtctggaacagattattctcaccatt agcaacctggagcaagaagatattgccacttctttccaacagggtaatgcgcttccgtggacgttcggtgg aggcaccagctggaactcaaacgg (SEQ ID NO:183)

Table 9: Full-length heavy chain polynucleotide sequences

Antibody	Full-Length Heavy Chain Polynucleotide Sequence (SEQ ID NO)
chCD37-3	aagcttgcaccatggctgtcctggcactgctcctctgctggtgacataccaagctgtgctctacacaggtgcaggtg aaggagtccagacctggcctggtggcgcctcagagcctgtccattacatgactgtctcagggttctcattaaccac ctctggtgtaagctgggttcgccagcctccaggaaagggtctggagtggtgggagtaatatgggtgacgggagcac aaactatcattcagctcctcaatccagactgagcatcaagaaggtcactccaagagccaagtttctaaactgaacagt ctgcaaaactgatgacacagccagctactactgtgccaagagggtactcgttggctcactggggccaagggactctgg tcacagctctgcaagcctctacgaaggcccatcagtttcccccttggctccaagttctaaatccacaagcggtggaacag ctgcaactgtgctcgtgtaagattatctcctgagcctgtgacagtgagctggaatagcggagcattgactcaggtgt gcacactttccgctgtgtgctcctcggctgtactcactgtccagtgctgtaaccgtccctctagcagcttgggaa cccagacctacatctgtaacgtcaaccataaacatccaacacaaaggtggataagaaggtgaaaccaagagctgtga taagacacafacatgcccctctgtcctgaccagagctcctcggagggtccatctgtgtcctgtttcccccaaaccaag gacactcttatgatctctgactccagaggtcactgtgtgtgtgacgtgagccatgaagatcccgaggttaaafca ctggtactggtgagtgaggttcacaatgccaagaccaagcccaggaggagcaatataattctacatacgggta gtgagcgttctgacctgtctccaccaagattggctcaatggaaaagagtacaagtgcaaggtgtccaacaaggctctcc cgctcccattgagaaaactatctcaaaagccaaggggcagccacgggaaccccagggtgatacattgccccatctaga gacagctgaccaagaaccagtgagctcactgtctggtcaaggggtttacccttctgacattgctgtagagtgaggag tctaacggacagccagaaaactacaagacaactccccagtgctggacagcagggagcttctcctactcca agttgactgtagacaagtctagatggcagcaaggaaacgtttctcctgctcagtaatgcatgaggtctgcacaactacta tacccaagaatcactgtcccttaaccagggtgactca (SEQ ID NO:152)
huCD37-3v1.0	aagcttgcaccatggctgtcctggcactgctcctctgctggtgacataccaagctgtgctctacacaggtgcaggtg

	<p>caagaatctgggtccaggtctgggtggcccccctccaaactctgagcatcacctgtaccgtttctgggtttgacctaccacctc tgggtgagttgggtacgccaaccacccggtaagggtctcgaatggctgggtgtaatctgggtgatggtccacaaatt accatcctccctcaagtccecgcttagcaiaaaaaggatcacagcaaaagcaagtttctgaaactgaatagtctgac agcagccgatacagccacctactattgcgccaagggtgggtatagtctgacactggggcaaggtaccctcgtaccgt ctctcagctagtaccaagggccatcagtttccccctggctccaagttcfaaafccacaagcgggtggaacagctgcaact gggatgcccgttaagattattccctgagcctgtgacagtgagctggaaatagcgggacattgactcagggtgtgcacac tttcccgctgtgtgagctctccggctgtactactgtccagtgctgtaaccgtccctctagcagcttgggaaccaga cctacatctgtaacgtaaccataaacatccaacacaaaagggtgataagaagggtgaaccaaagagctgtgataagac acatacatgccctcctgtcctgcaccagagctcctcggagggtccatctgtgttccgtttcccccaaaccaaggacact ctatgatctctcgtactccagaggtcacctgtgtgtgtgctgacgtgagccatgaagatcccagggttaaaftcaactggta cgtggatggagctgaggttcacaatgccaaagcaagccagggaggagcaatataaattctacafatcgggttagtgagc gttctgaccgtgtccaccaagattggctcaatggaaaagagtacaagtgcaaggtgtccaacaaggctctccccgtcc cattgagaaaactatctccaaagccaaaggccagccacgggaaccccagggtgtatacattgccccatctagagacga gctgaccaagaaccagggtgagctcactgtctgtcgaaggggtttaccctctgacattgctgtagagtgggagctaac ggacagccagaaaacaactacaagacaactccccagtgctggacagcgaacgggagcttctctactccaaggtga ctgtagacaagctagatggcagcaaggaaacgttttctcctcctcagtaatgatgaggtctgcacaatcaatatacc agaaatcactgtcccttagccagggtgactcgag (SEQ ID NO: 153)</p>
huCD37-3v1.1	<p>aagcttgcaccatgggctggagctgtatcattctgtttctgtgtggcgacagctactgggtccactccaagtgcaggtg caagagtccgggctggattggctgcaccaagccagacccctctctatcactgtaccgttagcgggtctctctgacaacc agtggagtgagttgggtgagcgaccaccaggaaaggactggagtggtcgggggtgattggggcgacggcgagca caactatcattccagcttaaatctcgggtgtccafiaaaaaagaccatagtaaatctcaagtttctgaaactcaatagcct gacagccgacagactgctactgtactgctgcccnaaggaggatacagctgtgctcactggggacaggggacccctgg gaccgtgtcatccgcatcaaaaaggccatcagtttccccctggctccaagtttaaatccacaagcgggtggaacag ctgcaactgggatgctctgtaagattattccctgagcctgtgacagtgagctggaaiaagcgggacattgactcaggtgt gcacactttcccgctgtgtgagctctccggctgtactactgtccagtgctgtaaccgtccctctagcagcttgggaa cccagacctacatctgtaacgtaaccataaacatccaacacaaaagggtgataagaagggtgaaccaaagagctgtga taagacacatacatgccctcctgtcctgcaccagagctcctcggagggtccatctgtgttctgttcccccaaaccaag gacactctatgatctctcgtactccagaggtcacctgtgtgtgtgacgtgagccatgaagatcccagggttaaaftca ctggtagctggatggagctgaggttcacaatgccaaagccagccagggaggagcaatataattctacatctcgggtg gtgagcgttctgaccgtctcccaagattggctcaatggaaaagagtacaagtgcaaggtgtccaacaaggctctcc cgtctccattgagaaaactatctccaaagccaaaggccagccacgggaaccccagggtgtatacattgccccatctaga gacgagctgaccaagaaccagggtgagctcactgtctgtcgaaggggtttaccctctgacattgctgtagagtgggag tctaacggacagccagaaaacaactacaagacaactccccagtgctggacagcgaacgggagcttctctctactcca agttgactgtagacaagctagatggcagcaaggaaacgttttctcctcctcagtaatgatgaggtctgcacaatcaat taccagaaatcactgtcccttagccagggtgactcgag (SEQ ID NO: 154)</p>
chCD37-12	<p>aagcttgcaccatggggtgtcatgtataatcctcttctgtgtcgtactgctaccgggtgtgcactcagagattcagctgg ttcaaaagtggccagagctgaaaaagccaggggaacagtgaaaataaggtgcaaggcattccggttacactttcacaaa gtaccgcatgaaactgggtcaagcagggccagggcaaggggctcaaatggatgggttggatcaatccaactggcg agtctaggaaatgctgaggagttaaggccgggttgccttcagccctggagacaagtgccagcagcacttaccgcaaatc aacaatctgaagfatgaggatacagcaacctatttctgcccggcggcactgtcgtgacagctggggacaaggtacca ccttgactgtatccagtccagcactaaggggccatcagtttccccctggctccaagttcfaaatccacaagcgggtggaa cagctgcaactgggatgctcgttaagattattccctgagcctgtgacagtgagctggaaatagcgggacattgactcag gtgtgcaacttttcccgctgtgtgagctctccggctgtactactgtccagtgctgtaaccgtccctctagcagcttgg gaaccagacctacatctgtaacgtaaccataaacatccaacacaaaagggtgataagaagggtgaaccaaagagctg tgataagacacatacatgccctcctgtcctgcaccagagctcctcggagggtccatctgtgttctgttcccccaaaccc aaggacactctatgatctctcgtactccagaggtcacctgtgtgtgtgacgtgagccatgaagatcccagggttaaaft caactgtgactggatggagctgaggttcacaatgccaaagccagccagggaggagcaatataattctacatctcgg gtagtgagcgttctgaccgtctcccaagattggctcaatggaaaagagtacaagtgcaaggtgtccaacaaggctct tcccgctccattgagaaaactatctccaaagccaaaggccagccacgggaaccccagggtgtatacattgccccatct agagacgagctgaccaagaaccagggtgagctcactgtctgtcgaaggggtttaccctctgacattgctgtagagtgg gagctaacggacagccagaaaacaactacaagacaactccccagtgctggacagcgaacgggagcttctctact ccaagttgactgtagacaagctagatggcagcaaggaaacgttttctcctcctcagtaatgatgaggtctgcacaatc actataccagaaatcactgtcccttagccagggtgactcgag (SEQ ID NO: 155)</p>

chCD37-38	<p>aagcttgccaccatgggctggagttgtatcattctgttttgggtggccaccgccactggagtcattcccaagtgaactcc aggaaatctggccctgacctggtaagccaatcagagcctctccctgacctgactgttacaggatacacaatcacatcag gctttggctggcactggatcagacaattcccgggaacaagttggaatggatggcttacattctgtatagcgggggtaccg attacaatccttcccctaagagccgaatctctatcaccagggatacaagcaagaaccaatftttctccgectcagctctgtg actaccgaagataccgctactactattgtgccaggggctactatggatatgtgcatggttcgtctattggggccaggga accctgtgactgtgagcgtgcctctaccaagggcccatcagtttcccctggctccaagttctaaatccacaagcgggt ggaacagctgactgggatgctcgttaaagattattccctgagcctgtgacagtgagctggaatagcggagcattgact tcaggtgtgcacactttcccgtgtgtgagcctccggctgtactcactgtccagtgtcgaaccgtccctctagcag cttgggaaccagactacatctgaacgtcaaccataaacatccaacacaaggtggataagaaggtgaaccaaag agctgtgataagacacatacatgcccctctgtctgcaccagagctcctcggaggctccatctgtgttctgtttccccc aaccgaaggacactctatgatctctgtactccagaggtcactgtgtgtgtgacgtgagccatgaagatcccagg ftaaattcaactggtacgtggatggagtcgaggttcacaatgccaagaccaagcccaggaggagcaatataattctaca tatcgggtagtgagcgttctgaccgtgtccaccaagattggctcaatggaaaagagtacaagtgaaggtgtccaaca ggctcttcccgtcccattgagaaaactatctccaagccaaggggagccacgggaaccccagggtgtatacattgcc ccatctagagacgagctgaccaagaaccagggtgagctcactgtctgtgcaaggggtttaccctctgacattgctgtg agtgaggagttaacggacagccagaaaactacaagacaactccccagtgctggacagcagcgggagcttctcc tctactccaagttgactgtagacaagctagatggcagcaaggaaacgttttctcctgctcagtaatgcatgaggctctgca caatcactatacccaaaatcactgtcccctagcccagggtgactcag (SEQ ID NO:156)</p>
huCD37-38	<p>aagcttgccaccatgggtggagctgcatcatttctggtcgtactgcaactggagtcactcagaggtccagctgc aagagtccggctcctggctgtgaaaccagccagtcctcagctcactgtactgtctgtgactacttaccagtg gttcggctggcattggattagcagtttcccggtaaggggctggagtggatggcatatctgtacagcggaggaacc gattacaaccaagctgaagagcaggatcagcattaccgggacacaagcaaaaaccagtttcttccgctgtctagt gttacagctgcagacaccgctactactattgtgctcgggttactatggctatggggctgtgtgtattggggacaag gcactctgtgaccgtgagcagcgcctcaacaagggcccatcagtttcccctggctccaagttctaaatccacaagc gtggaacagctgactgggatgctcgttaaagattattccctgagcctgtgacagtgagctggaatagcggagcattg acttcaggtgtgcacactttcccgtgtgtgagcctccggctgtactcactgtccagtgtcgaaccgtccctctagc agcttgggaaccagactacatctgaacgtcaaccataaacatccaacacaaggtggataagaaggtgaaccaa agagctgtgataagacacatacatgcccctctgtctgcaccagagctcctcggaggctccatctgtgttctgtttcccc caaaccgaagacactctatgatctctgtactccagaggtcactgtgtgtgtgacgtgagccatgaagatcccga ggftaaattcaactggtacgtggatggagtcgaggttcacaatgccaagaccaagcccaggaggagcaatataattcta catatcgggtagtgagcgttctgaccgtgtccaccaagattggctcaatggaaaagagtacaagtgaaggtgtccaac aaggtcttcccgtcccattgagaaaactatctccaagccaaggggagccacgggaaccccagggtgtatacattgc ccccatctagagacgagctgaccaagaaccagggtgagctcactgtctgtgcaaggggtttaccctctgacattgctgt agagtgggagttaacggacagccagaaaactacaagacaactccccagtgctggacagcagcgggagcttct cctctactccaagttgactgtagacaagctagatggcagcaaggaaacgttttctcctgctcagtaatgcatgaggctctg caaatcactatacccaaaatcactgtcccctagcccagggtgactcag (SEQ ID NO:157)</p>
huCD37-50	<p>aagcttgccaccatgggtggctcgtcataatccttctggttctactgctaccggagtcactcagaggtgcagctgc aggagtccggccccgctgctcaagcctctcagagctgagctgactgtactgtttctggctacagcataaccagcg gtttcgttggcactggatcagacagatcccggcaacaactggagtgatgggatacactgtactcaggtcaact gtctattccccctcctgaaatcccggatcagattaccctgacacttctaagaaccatfttttctgcagctgaacagcgtt accgcagctgacactgcaacctactactgtgccgggatattatggatacggagcttgggtctactgtgggccaagg caccctcgtactgtgagtgctgttccaccaagggcccatcagtttcccctggctccaagttctaaatccacaagcgggt ggaacagctgactgggatgctcgttaaagattattccctgagcctgtgacagtgagctggaatagcggagcattgact tcaggtgtgcacactttcccgtgtgtgagcctccggctgtactcactgtccagtgtcgaaccgtccctctagcag cttgggaaccagactacatctgaacgtcaaccataaacatccaacacaaggtggataagaaggtgaaccaaag agctgtgataagacacatacatgcccctctgtctgcaccagagctcctcggaggctccatctgtgttctgtttccccc aaccgaaggacactctatgatctctgtactccagaggtcactgtgtgtgtgacgtgagccatgaagatcccagg ftaaattcaactggtacgtggatggagtcgaggttcacaatgccaagaccaagcccaggaggagcaatataattctaca tatcgggtagtgagcgttctgaccgtgtccaccaagattggctcaatggaaaagagtacaagtgaaggtgtccaaca ggctcttcccgtcccattgagaaaactatctccaagccaaggggagccacgggaaccccagggtgtatacattgcc ccatctagagacgagctgaccaagaaccagggtgagctcactgtctgtgcaaggggtttaccctctgacattgctgtg agtgaggagttaacggacagccagaaaactacaagacaactccccagtgctggacagcagcgggagcttctcc tctactccaagttgactgtagacaagctagatggcagcaaggaaacgttttctcctgctcagtaatgcatgaggctctgca</p>

<p>huCD37-51</p>	<p>caatcactatacccagaaatcactgicccttagcccagggtagctcgag (SEQ ID NO:158)</p> <p>aagcttgcaccatgggttggcttgcatacctgttctgtggccactgccactggogtgcattcagaaggtcagttggtt ggagtcggccagaagtgctgaaacccggogaatcactgtccctgacttctacaggtgcaggttatagatcagcagc ggcttctctggcactggatcggcagttccaggcaaggactggaatggatgggtacatcattacagtggtcacaac caattacagccctagcctgcagggccgaatcttattaccaggatagttcttaataaccagtttctctgcagcttaattccgt gactgccctgacacagcaacttactatggcccgtggctactacgggttcggagccctggtttatactgggtcaggg caccctggicactgtctcagccgctctaccaagggccatcagtttcccttggctcgaagttctaaatccacaagcgggt ggaacagctgcactgggatgctcgttaagatttccctgagcctgtgacagtgagctggaatagcggagcattgact tcagggtgtgcacactttcccgctgtgtgacagctctccggctgtactcactgtccagtgctgtaaccgtccctctagcag cttgggaaccagacctacatctgtaacgtcaaccataaaccatccaacacaaagggtggataagaaggtgaaccaaag agctgtgataagacacatacatgcccctctgtcctgcaaccagagctcctggagggtccatctgtgttctgttccccca aaccacagacactcttatgatctctctgactccagaggtcactgtgtgtgtcgcactgtgagccatgaagatcccagg ttaaattcaactgggtacgtggatggagtcgaggttcacaatgccaaagaccaagcccaggggaggagcaatataaattcaca tatcgggtagttagcgttctgaccgtgtccaccaagattggctcaatggaaaagagtacaagtgcaagggtgtccacaa ggctcttcccgtcccattgagaaaactatctcaaaagccaaggggcagccacgggaaccccagggtatatacattgccc ccatctagagacgagctgaccagaaccagggtgagctcactgtctggcgaaggggtttaccctctgacattgtctgtag agtgaggagctaacggacagccagaaaacaactacaagacaactccccagtgctggacagcgcagggagctctcc ctactccaagttgactgtgacaaagtctagatggcagcaaggaaaacgtttctcctgctcagtaatgcatgaggctctgca caatcactatacccagaaatcactgicccttagcccagggtagctcgag (SEQ ID NO:159)</p>
<p>huCD37-56</p>	<p>aagcttgcaccatgggttggagctgcattatcctgttctcgtcgcaccgcaccggogtccactcccagggtgcagct gcaagaaagcgggcccaggatgglaaaaacctcccagctctctgagttactgtacagctatctggatcagatfcaatct ggcttctcctggcattggatcggcagttcccggcaaggggcttggatgggtatattcatttctggagggtacca actacaaccttcccgaagagtgagctcaattaccagggacactccaagaaccaattcttttgcagcttaattcagtg accgctgccgacaccgciacttactactgcgcceggggctactatgggttgggtcctggctgcctactggggccaggg gaacctgggtcccgtgtctgctgcctccacaaggcccacatcagtttccccttggctccaagttctaaatccacaagcgg tggaacagctgcactgggatgctcgttaagatttccctgagcctgtgacagtgagctggaatagcggagcattgac ttcagggtgtgcacactttcccgtgtgtgacagctcctggctctgactcactgtccagtgctgtaaccgtccctctagcag cttgggaaccagacctacatctgtaacgtcaaccataaaccatccaacacaaagggtggataagaaggtgaaccaaag agctgtgataagacacatacatgcccctctgtcctgcaaccagagctcctggagggtccatctgtgttctgttccccca aaccacagacactcttatgatctctctgactccagaggtcactgtgtgtgtcgcactgtgagccatgaagatcccagg ttaaattcaactgggtacgtggatggagtcgaggttcacaatgccaaagaccaagcccaggggaggagcaatataaattcaca tatcgggtagttagcgttctgaccgtgtccaccaagattggctcaatggaaaagagtacaagtgcaagggtgtccacaa ggctcttcccgtcccattgagaaaactatctcaaaagccaaggggcagccacgggaaccccagggtatatacattgccc ccatctagagacgagctgaccagaaccagggtgagctcactgtctggcgaaggggtttaccctctgacattgtctgtag agtgaggagctaacggacagccagaaaacaactacaagacaactccccagtgctggacagcgcagggagctctcc ctactccaagttgactgtgacaaagtctagatggcagcaaggaaaacgtttctcctgctcagtaatgcatgaggctctgca caatcactatacccagaaatcactgicccttagcccagggtagctcgag (SEQ ID NO:160)</p>
<p>huCD37-57</p>	<p>aagcttgcaccatgggttggagctgcataatctgttctcgttggccacagcaactggcgttcacagtcaggtccaactg caggagagcggccccggactcctgaaacctctcagtcactcagctgacatgactgtgagcggctacagcattacct aggcttctcgttggcattggatcaggcagttcccggaaaaggcttggagtgatgggttacattctgtacagcggcagta cagtgattcaccctctctgaaatctaggatcaatcacacgtgatacaagcaaaaatcagttctctccagctgaactcc gtcaccggcgcagacacagcaacctatattgtctcgggatactacggatagggcagctgttctcctattggggcca ggggacactctgaccgttccgcgcctccacaaggcccacatcagtttccccttggctccaagttctaaatccacaag cgggtggaacagctgcactgggatgctcgttaagatttccctgagcctgtgacagtgagctggaatagcggagcatt tgactcaggtgtgcacactttcccgtgtgtgacagctctccggctgtactcactgtccagtgctgtaaccgtccctctc gcagcttgggaaccagacctacatctgtaacgtcaaccataaaccatccaacacaaagggtggataagaaggtgaacc aaagagctgtgataagacacatacatgcccctctgtcctgcaaccagagctcctggagggtccatctgtgttctgttccc cccaaacccaagacactcttatgatctctctgactccagaggtcacctgtgtgtgtcgcactgtgagccatgaagatccc gaggttaaattcaactgggtacgtggatggagtcgaggttcacaatgccaaagaccaagcccaggggaggagcaatataa ctacatacgggtagttagcgttctgaccgtgtccaccaagattggctcaatggaaaagagtacaagtgcaagggtgtcc aacaaggtcttcccgtcccattgagaaaactatctcaaaagccaaggggcagccacgggaaccccagggtatatac tgcccccatctagagacgagctgaccagaaccagggtgagctcactgtctggcgaaggggtttaccctctgacattg ctgtagagtgaggctcaacggacagccagaaaacaactacaagacaactccccagtgctggacagcgcagggagc</p>

ttcttctctactccaagttgactgtagacaagctagatggcagcaaggaaacgtttctcctgctcagtaatgcatgagge ctcgcacaatcactataccagaaatcactgtcccttagccccagggtgactcgaag (SEQ ID NO:161)
--

Table 10: Full-length light chain polynucleotide sequences

Antibody	Full-length Light Chain Polynucleotide Sequence (SEQ ID NO)
chCD37-3	gaatfcgccaccatgagtggtgcccactcaggctctgggggtgctgctgctggtgcttacagatgccagatgtagacatccag atgactcagctccagcctcccctttctgtatctgtgggagaaaactgtcaccatcacatgtcagcaagtgagaatattcga gtaatttagcatggtatcagcagaaaacagggaatctcctcagcicctggcctcaatgtgcaacaacttagcagatgggt goccatcaaggtcagtggtcagtgatcaggcacacagtafctccctcaagaicaacagccigcagctcgaagatttggga cttatactgtcaacattatggggactacgtggacgtfcggtggaggcaccagctggaatcaaacgtacgggtggcig caccatctgtctcctctccgccatctgatgagcagttgaaatctggaaactgctctgtgtgctgctgaataacttcta tcccagagaggccaaagtacagtggaaggtggataacgccctccaatcgggtaactcccaggagagtgacacagagc aggacagcaaggacagcacctacagcctcagcagcaccctgacgtgagcaaaagcagactacgagaacacaaagt ctacgcctgcgaagtcacccatcagggtcctgagctcggccgtcacaagagctcaacaggggagagtgtag (SEQ ID NO:162)
huCD37-3 (1.0 and 1.1)	gaatfcgccaccatgggttggtectgcatcatctgtttctcgtggccacagccaccgggtgtcactctgatatacaaatgac tcaaaagcccctccagttfagcgtgaaggtgggtaacgccgtaacaatcactgtagagctagtgaaaacatccgcagta atctcgcagtggtaccaacaaaagccaggtaagteacctaagctctcgtgaatgtgctaccaacctcgcctgatggtg cttcaagattctctggtcaggttccggtaccgattatcacttaagatcaactcactccaaccagaagattfcggtacatatta ctgtcaaacactactggggtagcaccctggacatfcggtcaaggtactaagctggaaatcaagcgtacgggtgctgacccat ctgtcttcatctcccgccatctgatgagcagttgaaatctggaactgcctctgtgtgtgctcgtgaataacttctatcca gagaggccaaagtacagtggaaggtggataacgccctccaatcgggtaactcccaggagagtgctcacagagcaggac agcaaggacagcacctacagcctcagcagcaccctgacgtgagcaaaagcagactacgagaacacaaagtctacgc ctgcgaagtcacccatcagggtcctgagctcggccgtcacaagagctcaacaggggagagtgtag (SEQ ID NO:163)
chCD37-12	gaatfcgccaccatgggttggtectgtataatctgttcttgggtggccaccgctactggcgttcatagtgatattgtaetact cagtcaccagccagctctggcagtgctcctggccagcgtgcccaccatctctgcccgggctcagctccctgagcacta gctcttattcctatctctactgtttcaacagaagccaggacagccccctaaagctgctgatcaagtagcctccaacctcgc cagcggcgttcccctagatctctggttccggtagcggaaactgattcacttgaacatccaccccgttggaggaaggga taccgccactactattgtcaaacactctgggagattcctacaccttggaggaggaacaaagctcgaatlaagcgtacg gtggctgcaccatctgtcttctctccgccatctgatgagcagttgaaatctggaactgcctctgtgtgctcctgctgat aaactctatcccagagaggccaaagtacagtggaaggtggataacgccctccaatcgggtaactcccaggagagtgta cagagcaggacagcaaggacagcactacagcctcagcagcaccctgacgtgagcaaaagcagactacgagaacaca caaagtctacgcctgcgaagtcacccatcagggtcctgagctcggccgtcacaagagctcaacaggggagagtgtag (SEQ ID NO:164)
chCD37-38	gaatfcgccaccatgggttggtectgtataatctgttctcgtggccacagctacaggtgttcaattctcagattgtgctgac ccaatcaccagctattatgtccgctagccccggcgagaaagtgacaatgacatgtagcgtactcttctgtgactfacat gcattggatcaacagaagtcaggtaccagtcaccaagcgttggatctacgacacatccaaactggcctccggagtcctg ccaggtcagcggagggtgggtccggcaccagttatcactgaccatctctatggaagctgaagatgctgctactaita ttgtcaacaatggatttctaacccccacccttgggtggcgaacaaagctggagatcaagcgtacgggtgctgacccat ctgtcttcatctcccgccatctgatgagcagttgaaatctggaactgcctctgtgtgctcctgctgaataacttctatcca gagaggccaaagtacagtggaaggtggataacgccctccaatcgggtaactcccaggagagtgtagacagagcaggac agcaaggacagcacctacagcctcagcagcaccctgacgtgagcaaaagcagactacgagaacacaaagtctacgc ctgcgaagtcacccatcagggtcctgagctcggccgtcacaagagctcaacaggggagagtgtag (SEQ ID NO:165)
huCD37-38	gaatfcgccaccatgggttggtectgtataatctgttctcgtggccacagctacaggtgttcaattctcagattgtgctgac cagctccagcctcaatgtctcttccccgggtgagcgggtgaccatgacatgctctgcccagttcctcctgacatataatgc attggatcagcaaaaacccggtagctctccaaaagatggatctacgacactcaaaagcttgcacagggcgttctcctcca gatttccgggtctgggtctggcactcctacagctgaccattagttccatggaagctgaagatgagcaccctattactgt cagcagtggaattcaaaactctctacctcggcggcgaaccaaactggagataaaagcgtacgtggtgctgacccatctgt

	<p>cttcacatcccgccatctgatgagcagttgaaatctggaactgcctctgtgtgtgctctgaataactctatcccagaga ggccaaagtacagtggaaggtggataacgccctccaatcgggtaactcccaggagagtgctacagagcaggacagca aggacagcacctacagcctcagcagcaccctgacgctgagcaaaagcagactacgagaaacacaaaagtctacgctge gaagtcacccatcagggcctgagctcggccgacacaagagctcaacaggggagagtgtag (SEQ ID NO:166)</p>
<p>huCD37-50</p>	<p>gaattcgcaccatggggtggtatgcaattatctgttctggttgcctaccgcaacaggagtagatagtgatagtcctcac ccaaagtctctgctactatgtctgccagcccaggagagcgtgtgacctgactgctctgcaacctcaagtgtagacatacat gcattgggtatcagcaaaaagccctggccaatcccctaaaagggtggatctacgatacttaactctccatacgggtgtgcccgc aaggttctccgggagtggtgagtgccaccagttatagctgacctcagttcaatggaagcagaggatgcageaacctatt attgtcagcagtggtccgataatcccctacttttgggtcagggtacaaaagctggagattaaagctacgggtgctgaccat ctgtcttcacatctcccgccatctgatgagcagttgaaatctggaactgcctctgtgtgtgctctgtaataactctatcca gagaggccaaagtacagtggaaggtggataacgccctccaatcgggtaactcccaggagagtgctacagagcaggac agcaaggacagcacctacagcctcagcagcaccctgacgctgagcaaaagcagactacgagaaacacaaagtctacgc ctgcgaagtcacccatcagggcctgagctcggccgacaaaagagctcaacaggggagagtgtag (SEQ ID NO:167)</p>
<p>huCD37-51</p>	<p>gaattcgcaccatgggatggagctgtatattctgttctggttgcctactgctactggcgtccattccgagatagtcctcac ccagagcccgcacaacctgagtgctcctcctggggagcggagtactatgactgttccgccacttctcagttacatata gcattgggtatcagcagaaaacctggacagctcctcaaaagcgttgatttacgacacctccaacctggctcaggagttctge taggttcagcggatctgggtctggcacaagttatcaccatgattccatggaggccgaagatgcccgtacttactac tgtcagcagtgaggcagcaacccccctacattcgggcagggaaactaagctggagatcaaacgtacgggtgctgacca tctgtcttcacatctcccgccatctgatgagcagttgaaatctggaactgcctctgtgtgtgctctgtaataactctatcca gagaggccaaagtacagtggaaggtggataacgccctccaatcgggtaactcccaggagagtgctacagagcaggac agcaaggacagcacctacagcctcagcagcaccctgacgctgagcaaaagcagactacgagaaacacaaagtctacgc ctgcgaagtcacccatcagggcctgagctcggccgacaaaagagctcaacaggggagagtgtag (SEQ ID NO:168)</p>
<p>huCD37-56</p>	<p>gaattcgcaccatgggctggtctgtatcctctgttctggtggcaaccgctactgggggtcactctgatattgtctgac acagagtcagccctcatgagtgcttctccggagaaaaggctacaatgactgttcagcttctctcctcctcctcacatacatg cattgggtaccagcagaagcctgaccagagtcctaagaggtggatctatgatacaagcaatcggcttccgggtgctccctc ccgcttttcaggcggcggaaagcggaaactgactatagccttaccatctcaatggaagccgaggacgctgctacatatt actgcccagcaatggatcagcgaacctcctacttctggacagggaaacaaaattggaaitaagcgtacgggtgctgacc atctgtcttcacatctcccgccatctgatgagcagttgaaatctggaactgcctctgtgtgtgctctgtaataactctatcca agagaggccaaagtacagtggaaggtggataacgccctccaatcgggtaactcccaggagagtgctacagagcaggga cagcaaggacagcacctacagcctcagcagcaccctgacgctgagcaaaagcagactacgagaaacacaaagtctacg cctgcgaagtcacccatcagggcctgagctcggccgacaaaagagctcaacaggggagagtgtag (SEQ ID NO:169)</p>
<p>huCD37-57</p>	<p>gaattcgcaccatggggtggtctgtatattctgttctggttgcctaccgcaacaggcgttactccgagatcgtgtgta ctcagagcccagccaccatgtcgccttccccggggagagagtgacaatgactgttccgccacaagttctgtaacctac atgcattgggtaccagcaaaaaccaggacagagtcctccgctggtggattatgataccttaacctggctcaggcgttctg cccgttttctggtagtggatctgggacttctatagccttaccataagctctatggaagccgaggaccccgtacatacta ctgccagcagtggtgagtgataacccccacccttgggcagggaaacaaattggagatcaaacgtacgggtgctgacc atctgtcttcacatctcccgccatctgatgagcagttgaaatctggaactgcctctgtgtgtgctctgtaataactctatcca agagaggccaaagtacagtggaaggtggataacgccctccaatcgggtaactcccaggagagtgctacagagcaggga cagcaaggacagcacctacagcctcagcagcaccctgacgctgagcaaaagcagactacgagaaacacaaagtctacg cctgcgaagtcacccatcagggcctgagctcggccgacaaaagagctcaacaggggagagtgtag (SEQ ID NO:170)</p>

[00225] Also provided is a polynucleotide having at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% sequence identity to SEQ ID NOs:121-170, 182, or 183. Thus, in certain embodiments, the polynucleotide comprises (a) a polynucleotide having at least about 95% sequence identity to SEQ ID NOs:121-135, 152-161, or 182, and/or (b) a polynucleotide having at least about 95% sequence identity to SEQ ID NOs:136-151, 162-170, or 183. In certain embodiments, the

polynucleotide comprises (a) a polynucleotide having the nucleic acid sequence of SEQ ID NOs: 121-135, 152-161 or 182; and/or (b) a polynucleotide having the nucleic acid sequence of SEQ ID NOs: 136-151, 162-170, or 183.

[00226] In some embodiments, the polynucleotide encodes the light chain encoded by the recombinant plasmid DNA phuCD37-3LC (ATCC Deposit Designation PTA-10722, deposited with the ATCC on March 18, 2010) or a light chain that is at least about 85%, at least about 90%, at least about 95%, or at least about 99% to the light chain encoded by phuCD37-3LC (PTA-10722). In some embodiments, the polynucleotide encodes the heavy chain encoded by the recombinant plasmid DNA phuCD37-3HCv.1.0 (ATCC Deposit Designation PTA-10723, deposited with the ATCC on March 18, 2010) or a heavy chain that is at least about 85%, at least about 90%, at least about 95%, or at least about 99% identical to the heavy chain encoded by phuCD37-3HCv.1.0 (PTA-10723). In certain embodiments the polynucleotide is the recombinant plasmid DNA phuCD37-3LC (PTA-10722) or the recombinant plasmid phuCD37-3HCv.1.0 (PTA-10723).

[00227] In certain embodiments the polynucleotides comprise the coding sequence for the mature polypeptide fused in the same reading frame to a polynucleotide which aids, for example, in expression and secretion of a polypeptide from a host cell (e.g. a leader sequence which functions as a secretory sequence for controlling transport of a polypeptide from the cell). The polypeptide having a leader sequence is a preprotein and can have the leader sequence cleaved by the host cell to form the mature form of the polypeptide. The polynucleotides can also encode for a proprotein which is the mature protein plus additional 5' amino acid residues. A mature protein having a prosequence is a proprotein and is an inactive form of the protein. Once the prosequence is cleaved an active mature protein remains.

[00228] In certain embodiments the polynucleotides comprise the coding sequence for the mature polypeptide fused in the same reading frame to a marker sequence that allows, for example, for purification of the encoded polypeptide. For example, the marker sequence can be a hexa-histidine tag supplied by a pQE-9 vector to provide for purification of the mature polypeptide fused to the marker in the case of a bacterial host, or the marker sequence can be a hemagglutinin (HA) tag derived from the influenza hemagglutinin protein when a mammalian host (e.g. COS-7 cells) is used.

[00229] The present invention further relates to variants of the hereinabove described polynucleotides encoding, for example, fragments, analogs, and derivatives.

[00230] The polynucleotide variants can contain alterations in the coding regions, non-coding regions, or both. In some embodiments the polynucleotide variants contain alterations which produce silent substitutions, additions, or deletions, but do not alter the properties or activities of the encoded polypeptide. In some embodiments, nucleotide variants are produced by silent substitutions due to the degeneracy of the genetic code. Polynucleotide variants can be produced for a variety of reasons, e.g., to

optimize codon expression for a particular host (change codons in the human mRNA to those preferred by a bacterial host such as *E. coli*).

[00231] Vectors and cells comprising the polynucleotides described herein are also provided.

IV. Methods of use and pharmaceutical compositions

[00232] The CD37-binding agents (including antibodies, immunoconjugates, and polypeptides) of the invention are useful in a variety of applications including, but not limited to, therapeutic treatment methods, such as the treatment of cancer, such as B-cell malignancies, autoimmune diseases, and inflammatory diseases. In certain embodiments, the agents are useful for depleting B-cells. In certain embodiments, the agents are useful for depleting autoreactive B-cells. In certain embodiments, the agents are useful for depleting peripheral B-cells. In certain embodiments, the agents are useful for preventing inappropriate T-cell stimulation. The T-cell stimulation can be in connection with a B-cell pathway. The methods of use can be *in vitro*, *ex vivo*, or *in vivo* methods. In certain embodiments, the CD37-binding agent or antibody or immunoconjugate, or polypeptide is an antagonist of the human CD37 to which it binds.

[00233] In one aspect, anti-CD37 antibodies and immunoconjugates of the invention are useful for detecting the presence of CD37 in a biological sample. The term "detecting" as used herein encompasses quantitative or qualitative detection. In certain embodiments, a biological sample comprises a cell or tissue. In certain embodiments, such tissues include tissues that express CD37 at higher levels relative to other tissues, for example, B-cells and/or B-cell associated tissues.

[00234] In one aspect, the invention provides a method of detecting the presence of CD37 in a biological sample. In certain embodiments, the method comprises contacting the biological sample with an anti-CD37 antibody under conditions permissive for binding of the anti-CD37 antibody to CD37, and detecting whether a complex is formed between the anti-CD37 antibody and CD37.

[00235] In one aspect, the invention provides a method of diagnosing a disorder associated with increased expression of CD37. In certain embodiments, the method comprises contacting a test cell with an anti-CD37 antibody; determining the level of expression (either quantitatively or qualitatively) of CD37 by the test cell by detecting binding of the anti-CD37 antibody to CD37; and comparing the level of expression of CD37 by the test cell with the level of expression of CD37 by a control cell (e.g., a normal cell of the same tissue origin as the test cell or a cell that expresses CD37 at levels comparable to such a normal cell), wherein a higher level of expression of CD37 by the test cell as compared to the control cell indicates the presence of a disorder associated with increased expression of CD37. In certain embodiments, the test cell is obtained from an individual suspected of having an autoimmune disorder or inflammatory disorder. In some embodiments, the disorder is associated with increased expression of CD37. In some embodiments, the disorder is associated with increased number of B-cells. In some embodiments, the disorder is associated with increased activity of B-cells.

[00236] In certain embodiments, a method of diagnosis or detection, such as those described above, comprises detecting binding of an anti-CD37 antibody to CD37 expressed on the surface of a cell or in a membrane preparation obtained from a cell expressing CD37 on its surface. In certain embodiments, the method comprises contacting a cell with an anti-CD37 antibody under conditions permissive for binding of the anti-CD37 antibody to CD37, and detecting whether a complex is formed between the anti-CD37 antibody and CD37 on the cell surface. An exemplary assay for detecting binding of an anti-CD37 antibody to CD37 expressed on the surface of a cell is a "FACS" assay.

[00237] Certain other methods can be used to detect binding of anti-CD37 antibodies to CD37. Such methods include, but are not limited to, antigen-binding assays that are well known in the art, such as western blots, radioimmunoassays, ELISA (enzyme linked immunosorbent assay), "sandwich" immunoassays, immunoprecipitation assays, fluorescent immunoassays, protein A immunoassays, and immunohistochemistry (IHC).

[00238] In certain embodiments, anti-CD37 antibodies are labeled. Labels include, but are not limited to, labels or moieties that are detected directly (such as fluorescent, chromophoric, electron-dense, chemiluminescent, and radioactive labels), as well as moieties, such as enzymes or ligands, that are detected indirectly, e.g., through an enzymatic reaction or molecular interaction.

[00239] In certain embodiments, anti-CD37 antibodies are immobilized on an insoluble matrix. Immobilization entails separating the anti-CD37 antibody from any CD37 that remains free in solution. This conventionally is accomplished by either insolubilizing the anti-CD37 antibody before the assay procedure, as by adsorption to a water-insoluble matrix or surface (Bennich et al., U.S. Pat. No. 3,720,760), or by covalent coupling (for example, using glutaraldehyde cross-linking), or by insolubilizing the anti-CD37 antibody after formation of a complex between the anti-CD37 antibody and CD37, e.g., by immunoprecipitation.

[00240] Any of the above embodiments of diagnosis or detection can be carried out using an immunoconjugate of the invention in place of or in addition to an anti-CD37 antibody.

[00241] In certain embodiments, the disease treated with the CD37-binding agent is an autoimmune or inflammatory disease. In certain embodiments, the autoimmune or inflammatory disease is selected from the group consisting of psoriasis, dermatitis, systemic scleroderma and sclerosis, responses associated with inflammatory bowel disease, Crohn's disease, ulcerative colitis, respiratory distress syndrome, adult respiratory distress syndrome (ARDS), dermatitis, meningitis, encephalitis, uveitis, colitis, glomerulonephritis, allergic conditions, eczema, asthma, conditions involving infiltration of T cells and chronic inflammatory responses, atherosclerosis, leukocyte adhesion deficiency, rheumatoid arthritis, systemic lupus erythematosus (SLE), diabetes mellitus, multiple sclerosis, Reynaud's syndrome, autoimmune thyroiditis, allergic encephalomyelitis, Sjorgen's syndrome, juvenile onset diabetes, immune responses associated with acute and delayed hypersensitivity mediated by cytokines and T-lymphocytes,

tuberculosis, sarcoidosis, polymyositis, granulomatosis, vasculitis, pernicious anemia (Addison's disease), diseases involving leukocyte diapedesis, central nervous system (CNS) inflammatory disorder, multiple organ injury syndrome, hemolytic anemia, myasthenia gravis, antigen-antibody complex mediated diseases, anti-glomerular basement membrane disease, antiphospholipid syndrome, allergic neuritis, Graves' disease, Lambert-Eaton myasthenic syndrome, pemphigoid bullous, pemphigus, autoimmune polyendocrinopathies, Reiter's disease, stiff-man syndrome, Behcet disease, giant cell arteritis, immune complex nephritis, IgA nephropathy, IgM polyneuropathies, idiopathic thrombocytopenic purpura (ITP) and autoimmune thrombocytopenia.

[00242] In some embodiments, the autoimmune or inflammatory disease is selected from the group consisting of: RA, lupus, immune thrombocytopenic purpura, pure red cell aplasia, autoimmune anemia, cold agglutinin disease, type B syndrome of severe insulin resistance, mixed cryoglobulinemia, myasthenia gravis, Wegener's granulomatosis, microscopic polyangiitis (MPA), refractory pemphigus vulgaris, dermatomyositis, Sjogren's syndrome, active type-II mixed cryoglobulinemia, pemphigus vulgaris, autoimmune neuropathy, paraneoplastic opsoclonus-myoclonus syndrome, and relapsing-remitting multiple sclerosis (RRMS).

[00243] In certain embodiments, the autoimmune disease or inflammatory disease is characterized by CD37 expressing cells to which the CD37-binding agent (e.g., antibody) binds.

[00244] The present invention provides for methods of treating autoimmune and inflammatory diseases comprising administering a therapeutically effective amount of a CD37-binding agent to a subject (e.g., a subject in need of treatment). In certain embodiments, the subject is a human.

[00245] The present invention further provides methods for depleting B-cells, e.g., autoreactive B-cells, using the antibodies or other agents described herein. In certain embodiments, the method of depleting B-cells comprises contacting a B-cell with a CD37-binding agent (e.g., antibody) *in vitro*. For example, a cell line that expresses CD37 is cultured in medium to which is added the antibody or other agent to deplete the cells. In some embodiments, the cells are isolated from a patient sample such as, for example, a tissue biopsy, pleural effusion, or blood sample and cultured in medium to which is added an CD37-binding agent to deplete the cells.

[00246] In some embodiments, the method of depleting B-cells, e.g. autoreactive B-cells, comprises contacting the cells with the CD37-binding agent (e.g., antibody) *in vivo*. In certain embodiments, contacting a cell with a CD37-binding agent is undertaken in an animal model. For example, CD37-binding agents can be administered to xenografts expressing one or more CD37s that have been grown in immunocompromised mice (e.g. NOD/SCID mice). In some embodiments, cells are isolated from a patient sample such as, for example, a tissue biopsy, pleural effusion, or blood sample and injected into immunocompromised mice that are then administered a CD37-binding agent to deplete B-cells. In some embodiments, the CD37-binding agent is administered at the same time or shortly after introduction of

cells into the animal. In further examples, CD37 binding agents can be administered *in vivo* to mice expressing one or more CD37 antigens. In some embodiments, these mice can be engineered to express human CD37 in addition to, or instead of, murine CD37. In some embodiments, these mice are disease models, e.g. models for autoimmune disease. In some embodiments, administering a CD37 binding agent depletes B-cells *in vivo*. In some embodiments, a CD37 binding agent prevents T-cell stimulation. In some embodiments, administering a CD37 binding agent prevents or alleviates an autoimmune disease.

[00247] In certain embodiments, the B-cells overexpress CD37. In other embodiments, the B-cells do not overexpress CD37. In some embodiments, the B-cells are not cancer cells. In some embodiments, the B-cells are not tumor cells. In some embodiments, the B-cells are not cancerous cells.

[00248] The present invention further provides pharmaceutical compositions comprising one or more of the CD37-binding agents described herein. In certain embodiments, the pharmaceutical compositions further comprise a pharmaceutically acceptable vehicle. These pharmaceutical compositions find use in treating autoimmune and inflammatory disease in human patients.

[00249] In certain embodiments, formulations are prepared for storage and use by combining a purified antibody or agent of the present invention with a pharmaceutically acceptable vehicle (e.g. carrier, excipient) (Remington, The Science and Practice of Pharmacy 20th Edition Mack Publishing, 2000). Suitable pharmaceutically acceptable vehicles include, but are not limited to, nontoxic buffers such as phosphate, citrate, and other organic acids; salts such as sodium chloride; antioxidants including ascorbic acid and methionine; preservatives (e.g. octadecyldimethylbenzyl ammonium chloride; hexamethonium chloride; benzalkonium chloride; benzethonium chloride; phenol, butyl or benzyl alcohol; alkyl parabens, such as methyl or propyl paraben; catechol; resorcinol; cyclohexanol; 3-pentanol; and m-cresol); low molecular weight polypeptides (e.g. less than about 10 amino acid residues); proteins such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, histidine, arginine, or lysine; carbohydrates such as monosaccharides, disaccharides, glucose, mannose, or dextrans; chelating agents such as EDTA; sugars such as sucrose, mannitol, trehalose or sorbitol; salt-forming counter-ions such as sodium; metal complexes (e.g. Zn-protein complexes); and non-ionic surfactants such as TWEEN or polyethylene glycol (PEG).

[00250] The pharmaceutical compositions of the present invention can be administered in any number of ways for either local or systemic treatment. Administration can be topical (such as to mucous membranes including vaginal and rectal delivery) such as transdermal patches, ointments, lotions, creams, gels, drops, suppositories, sprays, liquids and powders; pulmonary (e.g., by inhalation or insufflation of powders or aerosols, including by nebulizer; intratracheal, intranasal, epidermal and transdermal); oral; or parenteral including intravenous, intraarterial, subcutaneous, intraperitoneal or intramuscular injection or infusion; or intracranial (e.g., intrathecal or intraventricular) administration.

[00251] An antibody or immunoconjugate of the invention can be combined in a pharmaceutical combination formulation, or dosing regimen as combination therapy, with a second compound having anti-autoimmune or inflammatory properties. The second compound of the pharmaceutical combination formulation or dosing regimen can have complementary activities to CD37-binding agent of the combination such that they do not adversely affect each other. Pharmaceutical compositions comprising the CD37-binding agent and the second agent are also provided. For example, CD37-binding agents can be administered in combination with CD20-binding agents, such as Rituximab. In other embodiments, CD37-binding agents can be administered in combination with salicylate; nonsteroidal anti-inflammatory drugs such as indomethacin, phenylbutazone, phenylacetic acid derivatives (e.g., ibuprofen and fenoprofen), naphthalene acetic acids (naproxen), pyrrolealkanoic acid (tometin), indoleacetic acids (sulindac), halogenated anthranilic acid (meclofenamate sodium), piroxicam, zomepirac and diflunisal; antimalarials such as chloroquine; gold salts; penicillamine; or immunosuppressive agents such as methotrexate or corticosteroids. In some embodiments, the CD37-binding agent is administered in combination with a second therapeutic selected from the group consisting of methotrexate, an anti-CD20 therapeutic, an anti-IL-6 receptor therapeutic, an anti-IL-12/23p40 therapeutic, a chemotherapeutic, an immunosuppressant, an anti-interferon beta-1a therapeutic, glatiramer acetate, an anti- α 4-integrin therapeutic, fingolimod, an anti-BLys therapeutic, CTLA-Fc, or an anti-TNF therapeutic. In some embodiments, the CD37-binding agent is administered in combination with a second therapeutic that is an antibody directed against an antigen selected from a group consisting of CD3, CD14, CD19, CD20, CD22, CD25, CD28, CD30, CD33, CD36, CD38, CD40, CD44, CD52, CD55, CD59, CD56, CD70, CD79, CD80, CD103, CD134, CD137, CD138, and CD152. In some embodiments, the CD37-binding agent is administered in combination with a second therapeutic that is an antibody directed against a target selected from the group consisting of IL-2, IL-6, IL-12, IL-23, IL-12/23 p40, IL-17, IFN γ , TNF α , IFN α , IL-15, IL-21, IL-1a, IL-1b, IL-18, IL-8, IL-4, GM-CSF, IL-3, and IL-5. In some embodiments, the CD37-binding agents are administered in combination with methotrexate.

[00252] For the treatment of the disease, the appropriate dosage of an antibody or agent of the present invention depends on the type of disease to be treated, the severity and course of the disease, the responsiveness of the disease, whether the antibody or agent is administered for therapeutic or preventative purposes, previous therapy, patient's clinical history, and so on all at the discretion of the treating physician. The antibody or agent can be administered one time or over a series of treatments lasting from several days to several months, or until a cure is affected or a diminution of the disease state is achieved. Optimal dosing schedules can be calculated from measurements of drug accumulation in the body of the patient and will vary depending on the relative potency of an individual antibody or agent. The administering physician can easily determine optimum dosages, dosing methodologies and repetition rates. In certain embodiments, dosage is from 0.01 μ g to 100 mg per kg of body weight, and can be given

once or more daily, weekly, monthly or yearly. In certain embodiments, the antibody or other CD37-binding agent is given once every two weeks or once every three weeks. In certain embodiments, the dosage of the antibody or other CD37-binding agent is from about 0.1 mg to about 20 mg per kg of body weight. The treating physician can estimate repetition rates for dosing based on measured residence times and concentrations of the drug in bodily fluids or tissues.

[00253] The combination therapy can provide "synergy" and prove "synergistic", i.e. the effect achieved when the active ingredients used together is greater than the sum of the effects that results from using the compounds separately. A synergistic effect can be attained when the active ingredients are: (1) co-formulated and administered or delivered simultaneously in a combined, unit dosage formulation; (2) delivered by alternation or in parallel as separate formulations; or (3) by some other regimen. When delivered in alternation therapy, a synergistic effect can be attained when the compounds are administered or delivered sequentially, e.g. by different injections in separate syringes. In general, during alternation therapy, an effective dosage of each active ingredient is administered sequentially, i.e. serially, whereas in combination therapy, effective dosages of two or more active ingredients are administered together.

VI. Kits comprising CD37-binding agents

[00254] The present invention provides kits that comprise the antibodies, immunoconjugates or other agents described herein and that can be used to perform the methods described herein. In certain embodiments, a kit comprises at least one purified antibody against CD37 in one or more containers. In some embodiments, the kits contain all of the components necessary and/or sufficient to perform a detection assay, including all controls, directions for performing assays, and any necessary software for analysis and presentation of results. A label or indicator describing, or a set of instructions for use of, kit components in a ligand detection method of the present invention, can also be included. The instructions may be associated with a package insert and/or the packaging of the kit or the components thereof. One skilled in the art will readily recognize that the disclosed antibodies, immunoconjugates or other agents of the present invention can be readily incorporated into one of the established kit formats which are well known in the art. Such kits can also include, for example, other compounds and/or compositions, a device(s) for administering the compounds and/or compositions, and written instructions in a form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products.

[00255] Further provided are kits comprising a CD37-binding agent (e.g., a CD37-binding antibody), as well as a second agent. In certain embodiments, the second agent is rituximab. In certain embodiments, the second agent is methotrexate.

* * *

[00256] Embodiments of the present disclosure can be further defined by reference to the following non-limiting examples, which describe in detail preparation of certain antibodies of the present disclosure

and methods for using antibodies of the present disclosure. It will be apparent to those skilled in the art that many modifications, both to materials and methods, can be practiced without departing from the scope of the present disclosure.

Examples

[00257] It is understood that the examples and embodiments described herein are for illustrative purposes only and that various modifications or changes in light thereof will be suggested to persons skilled in the art and are to be included within the spirit and purview of this application.

[00258] All publications, patents, patent applications, internet sites, and accession numbers/database sequences (including both polynucleotide and polypeptide sequences) cited herein are hereby incorporated by reference in their entirety for all purposes to the same extent as if each individual publication, patent, patent application, internet site, or accession number/database sequence were specifically and individually indicated to be so incorporated by reference.

Example 1

CD37 expression in normal human PBMCs

[00259] The CD37 antigen was reported to be expressed on B-cells from the pre-B stage to the peripheral mature B-cell stage, while being absent on B-cell progenitors and terminally differentiated plasma cells. (Link et al., 1987, J Pathol. 152:12-21). In addition, the CD37 antigen is only weakly expressed on T-cells, myeloid cells and granulocytes (Schwartz-Albiez et al. 1988, J. Immunol., 140(3)905-914).

[00260] The ability of antibodies (including certain CD37 antibodies and immunoconjugates previously described in U.S. Published Application No. 2011/0256153, which is herein incorporated by reference in its entirety) to bind to normal human B-cells was measured using flow cytometry assays with fluorescently labeled antibodies. In addition, the commercially available QuantiBRITE system from BD Biosciences was used to estimate antigen density based on the number of antibodies bound to the cells (ABC). The QuantiBRITE system from BD Biosciences utilizes the following reagents: anti-CD20-PE supplied at 100 µg/mL and QuantiBRITE PE supplied as lyophilized PE-labeled beads. In addition, the huCD37-3 antibody was labeled with PE to obtain an antibody-PE conjugate with an Ab:PE ratio of approximately 1:1.

[00261] Fresh buffy coats from healthy donors were obtained from Research Blood Components (Brighton, MA, US) as a source of normal blood cells. Buffy coats were prepared by centrifugation of a unit of whole blood and collecting the interface between the plasma and the red blood cells. This unpurified buffy coat contains PBMCs, neutrophils, platelets, red blood cells, and plasma and was used

for experiments on the same day it was drawn. Peripheral blood mononuclear cells (PBMCs) were prepared from buffy coats by standard density gradient centrifugation using Ficoll-Paque as follows. Blood was diluted 1:3 with 1x HBSS containing 5mM EDTA and up to 30 mL were added to a 50 mL conical tube. Ten mL of Ficoll-Paque (GE Healthcare) were slowly added to the bottom of each tube. Samples were centrifuged at 500 x g with no brake at RT for 30 minutes to obtain a layer of PBMCs below the plasma and to remove red blood cells and most granulocytes. The PBMCs were transferred to new tubes and washed twice with 1x HBSS containing 5mM EDTA by centrifugation at 400 x g for 10 minutes at RT. Staining buffer (1x HBSS, 1% BSA, 0.1% sodium azide) was then used to resuspend the PBMC pellets at 6.25×10^6 cells/mL. Eighty μ L of cells were transferred to a round-bottom 96-well plate to achieve 5×10^5 cells/assay and 20 μ L of human serum (Sigma H4522) were added to block Fc receptor-mediated binding and incubated with cells on ice for 20 min in the dark. Fluorescently labeled antibodies obtained from Miltenyi were used to identify PBMC populations: anti-CD3-allophycocyanin (APC) was used to identify T-cells, anti-CD19-APC for B-cells, anti-CD56-APC for natural killer (NK) cells and anti-CD14-APC for monocytes.

[00262] Cells were co-stained for CD37 expression using 20 μ L of huCD37-3-PE for a final concentration of approximately 10 μ g/mL. Likewise, cells were co-stained for CD20 expression using 20 μ L of anti-CD20-PE. As a control a non-binding PE-labeled huIgG1 isotype control antibody was used at 10 μ g/mL. Staining was carried out for 1 hour on ice in the dark. Samples were washed twice with staining buffer and fixed in 200 μ L of 1% formaldehyde in 1x PBS. Samples were stored at 4°C in the dark until acquisition, which was performed within 4 days of sample preparation.

[00263] A fresh tube of QuantiBRITE beads was reconstituted in the supplied tube with 0.5 mL of staining buffer just prior to sample acquisition. Samples were acquired on a FACSCalibur flow cytometer (BD Biosciences). Compensation controls were run with each assay to select appropriate instrument settings and at least 10,000 events were collected for each sample. Instrument settings for fluorescence and compensation were kept the same for both cell sample and bead sample acquisition to allow for an accurate comparison. CellQuest (version 5.2.1, BD Biosciences) was used for acquisition control and analysis.

[00264] The QuantiBRITE analysis utilizes on a bead standard with 4 bead populations conjugated with a known number of PE molecules. For data analysis, a G1 gate was drawn around the bead singlets on an FSC-H/SSC-H scatter plot. This gated bead population was subsequently analyzed using a histogram plot of FL2-H to evaluate the level of PE staining. Separate markers were drawn around the peaks of the four bead populations (M1-M4) and the geometric mean for FL2 of each bead population was determined. The FL2 geometric mean of each bead was plotted against the lot specific PE/bead values in a log-log plot. Linear regression was performed to obtain a standard curve using the following equation:

$y = mx + c$, with “m” equal to the slope and “c” equal to the y-intercept.

[00265] For PBMC sample analysis, a G1 gate was drawn around the positive fluorescent cell population of interest on an SSC-H/FL4-H dot plot. This gated cell population was subsequently analyzed using a histogram plot of FL2-H to evaluate the level of PE-labeled antibody staining. The FL2 geometric mean was determined for each blood cell sample stained with anti-CD37-PE or anti-CD20-PE, as well as unstained control samples. All geometric mean values for FL2 were plotted against the bead standard curve and values for PE per cell were extrapolated. Since both antibody-PE conjugates were at a PE:Ab ratio of approximately 1:1, the values for PE per cell correspond to the number of antibodies bound per cell (ABC) value. Experiments were performed with duplicate samples for each assay. The mean and standard deviation was determined from several assays for each blood cell population.

[00266] CD37 expression was evaluated in normal blood cells from 4 independent donors. Results were compared to CD20 staining, unstained cells and a non-binding huIgG-PE conjugate as controls. An example of a typical staining profile of normal B-cells is given in a histograms in Figure 1. The average ABC values of 4 different experiments for CD37 and CD20 were calculated and listed in Table 1.

[00267] Table 1: ABC values for CD37 and CD20 expression on human PBMC samples

	CD37 ABC	CD20 ABC	No Ab control	huIgG-PE control
CD19+ B-cells	77,440	94,598	80	76
CD3+ T cells	2,016	336	74	68
CD56+ NK cells	3,090	264	85	88
CD14+ monocytes	5,244	794	180	215

[00268] The highest overall CD37 staining level was found in CD19+ B-cells at approximately 77,000 ABC. In addition, CD37 staining was seen at low levels in other PBMC populations examined, with CD14+ monocytes showing CD37 staining at approximately 5,000 ABC, CD56+ NK cells at 3,000 ABC, and CD3+ T cells at 2,000 ABC. Staining with the non-binding huIgG-PE control resulted in ABC values of approximately 70 – 90 for B, T and NK cells and approximately 200 for monocytes. In the same 4 donors CD20 expression was evaluated in comparison to CD37. In accordance with published findings, the CD20 staining was restricted mainly to CD19+ B-cells with an ABC value of approximately 95,000 ABC. The CD20 expression level was just slightly higher than the CD37 expression level. Only minimal CD20 staining was observed in other PBMC populations examined, with CD14+ monocytes showing CD20 staining at 794 ABC, CD56+ NK cells at 264 ABC and CD3+ T cells at 336 ABC.

[00269] This result demonstrates that high CD37 expression is mainly restricted to B-cells in peripheral blood samples with only minor expression on peripheral T cells, NK cells and monocytes. This is consistent with published findings ((Moore et al. 1986, J Immunol. 137(9):3013-8; Schwartz-Albiez et al. 1988, J. Immunol., 140(3)905-914). In addition, we found that the CD37 expression levels on

peripheral B-cells is similar to the level of CD20 expression. This expression pattern strongly suggest that CD37 directed therapies may be a suitable for targeting B-cells in diseases such as B-cell malignancies, autoimmune diseases, inflammatory diseases or other disorders of the immune system analogous to the use of CD20 directed therapies.

Example 2A

In vitro B-cell depletion using purified PBMCs

[00270] The ability of humanized antibodies to deplete B-cells was measured using *in vitro* assays with human PBMCs according to published studies performed with rituximab (Vugmeyster et al. Cytometry A. 2003;52(2):101-9 and Vugmeyster et al. Int Immunopharmacol. 2004;4(8):1117-24). Alemtuzumab (Campath) was used as apposite control, since it has been reported to efficiently deplete lymphocytes *in vivo* and *in vitro* (Hale, Blood. 1983 Oct;62(4):873-82 and Waldmann, Philos Trans R Soc Lond B Biol Sci. 2005 Sep 29;360(1461):1707-11).

[00271] Fresh buffy coats from healthy donors were obtained from Research Blood Components (Brighton, MA, US) as a source of normal blood cells for all experiments within this study. Buffy coats were prepared by centrifugation of a unit of whole blood and collecting the interface between the plasma and the red blood cells. This unpurified buffy coat contains PBMCs, neutrophils, platelets, red blood cells, and plasma and was used for experiments on the same day it was drawn. Peripheral blood mononuclear cells (PBMCs) were prepared from buffy coats by standard density gradient centrifugation using Ficoll-Paque as follows. Blood was diluted 1:3 with 1x HBSS containing 5mM EDTA and up to 30 mL were added to a 50 mL conical tube. Ten mL of Ficoll-Paque (GE Healthcare) were slowly added to the bottom of each tube. Samples were centrifuged at 500 x g with no brake at RT for 30 minutes to obtain a layer of PBMCs below the plasma and to remove red blood cells and most granulocytes. The PBMCs were transferred to new tubes and washed twice with 1x HBSS containing 5mM EDTA by centrifugation at 400 x g for 10 minutes at RT. Staining buffer (1x HBSS, 1% BSA, 0.1% sodium azide) was then used to resuspend the PBMC pellets in the initial blood volume to achieve the original cell density.

[00272] To assess the effect of huCD37-3, huCD37-3-SMCC-DM1, huCD37-50, huCD37-50-SMCC-DM1, rituximab, alemtuzumab (Campath), and TRU-016 on PBMC depletion, 90 μ L of purified cells were added to 12 x 75 mm polystyrene tubes and incubated with 10 μ L of a 100 μ g/mL solution of each sample or a huIgG isotype control antibody for 1 hr at 37°C in a humidified 5% CO₂ incubator. The final antibody (Ab) concentration was 10 μ g/mL in a final volume of 100 μ L in staining buffer. Three independent samples were prepared for each treatment.

[00273] To identify populations of PBMCs, all samples were co-stained immediately after Ab incubation with 10-20 μ L of fluorescently labeled Abs obtained from, for example, BD Biosciences or

Miltenyi. Anti-CD3-PerCP-Cy5.5 was used to identify T cells, anti-CD19-APC for B-cells, and anti-CD14-FITC for monocytes. Staining was carried out in a total of 150 μ L for 30 min in the dark at RT. CountBright Absolute Counting Beads (Invitrogen) were vortexed and added to each sample at 50 μ L per tube. For PBMC prep samples, cells were washed once with 1 mL staining buffer and centrifuged at 400 x g for 3-5 min. Supernatant was removed with a 1 mL pipette and cells were resuspended in 500 μ L of 1% formaldehyde in 1x PBS. Samples were stored at 4°C in the dark until acquisition, which was performed within 4 days of sample preparation.

[00274] TreeStar FlowJo software (version PC 7.5) was used for data analysis. A gate was drawn around the CountBright bead population on an FSC-H vs SSC-H dot plot to determine a total bead count for the sample. To determine the total count for each PBMC population of interest, a separate gate was drawn around the positive fluorescent population on an SSC-H vs FL(x)-H dot plot, where x is the channel of interest. Specifically, a total count for T cells in a sample was found by gating the positive population on an SSC-H vs FL3-H dot plot; for B-cells, the positive population was found on an SSC-H vs FL4-H dot plot; for NK cells, an SSC-H vs FL2-H dot plot was used; for monocytes, an SSC-H vs FL1-H dot plot was used. The ratio of CD19+ cells for B-cells (CD3+ cells for T cells, CD56+ cells for NK cells, or CD14+ cells for monocytes) relative to beads was determined and multiplied by 100. Percent depletion was then calculated by taking the ratio of the cell to bead ratio in treated samples relative to the cell to bead ratio in isotype control treated samples, subtracting this from 1 and multiplying by 100. This corresponds to the following formula: Percent Depletion = $100 \times (1 - \frac{\text{cell to bead ratio of treated sample}}{\text{cell to bead ratio of control sample}})$. Data for all cell types was analyzed in the same manner.

[00275] For two donors tested, treatment of purified PBMC samples with huCD37-3, huCD37-3-SMCC-DM1, huCD37-50 or huCD37-50-SMCC-DM1 resulted in approximately 55-70% depletion of B-cells (see Figure 2). There was less than 10% depletion of T cells or monocytes. The B-cell restricted depletion effect indicates that this activity is linked to the high CD37 expression on B-cells. In comparison, treatment with the anti-CD20 antibody rituximab resulted in approximately 30-40% depletion of B-cells. Treatment with the anti-CD37 SMIP™ TRU-016 resulted in only 20-30% depletion of B-cells. Alemtuzumab treatment resulted in depletion of 60-70% of B-cells, 55-65% of T cells and 40-65% of monocytes.

Example 2B

Dose response for *in vitro* B-cell depletion using purified PBMCs

[00276] To evaluate the dose-response of the antibodies and conjugates, purified PBMCs from 2 donors were incubated with a 5-fold sample dilution series. Each sample dilution was added at 10 μ L per tube to 90 μ L of purified cells in triplicate and incubated for 1 hour at 37°C in a humidified 5% CO₂ incubator. The final concentration ranged from 10 μ g/mL to 0.13 ng/mL. The same amount of a non-binding huIgG Ab was used as an isotype control.

[00277] For two donors tested, treatment of purified PBMC samples with huCD37-3-SMCC-DM1 resulted in a clear dose-response for the B-cell depletion activity (see Figure 3A and B). Incubation with huCD37-3-SMCC-DM1 caused *in vitro* depletion of approximately 60% of B-cells with an EC₅₀ of 40-75 ng/mL. For an additional donor tested, treatment of purified PBMC samples with huCD37-3, huCD37-38, huCD37-50, and huCD37-56 antibodies also resulted in a clear dose-response for the B-cell depletion activity (see Figure 3C). Incubation with these antibodies caused *in vitro* depletion of approximately 60-70% of B-cells with an EC₅₀ of 20-30 ng/mL.

Example 2C

In vitro B-cell depletion using whole blood

[00278] The ability of humanized antibodies to deplete B-cells was measured using *in vitro* assays with whole blood according to published studies performed with rituximab (Vugmeyster et al. Cytometry A. 2003;52(2):101-9 and Vugmeyster et al. Int Immunopharmacol. 2004;4(8):1117-24).

[00279] Fresh buffy coats from healthy donors were obtained from Research Blood Components (Brighton, MA, US) as a source of normal blood cells for all experiments within this study. To assess the effect of huCD37-3, huCD37-3-SMCC-DM1, rituximab, alemtuzumab (Campath), and TRU-016 on peripheral blood cells (PBCs) in a whole blood matrix, 90 μ L of whole blood from a buffy coat were incubated with Abs or isotype control as detailed above in a total volume of 100 μ L. Three independent samples were prepared for each Ab treatment.

[00280] To identify populations of blood cells, all samples were co-stained immediately after Ab incubation with 10 - 20 μ L of fluorescently labeled Abs obtained from, for example, BD Biosciences or Miltenyi. Anti-CD3-PerCP-Cy5.5 was used to identify T cells, anti-CD19-APC for B-cells, anti-CD56-PE for NK cells, and anti-CD14-FITC for monocytes. Staining was carried out in a total of 150 μ L for 30 min in the dark at RT. CountBright Absolute Counting Beads (Invitrogen #C36950) were vortexed and added to each sample at 50 μ L per tube to allow standardization of cell counts.

[00281] Following cell staining, 2 mL of BD FACS Lysing Solution (BD Biosciences, diluted 1:10 in dH₂O according to the manufacturer's instructions) were added to each sample in order to lyse the RBCs present. Samples were incubated at RT for 15-20 min in the dark, centrifuged at 400 x g for 3-5 min, and resuspended in 500 μ L of 1% formaldehyde in 1x PBS. Samples were stored at 4°C in the dark until acquisition, which was performed within 4 days of sample preparation. Samples were acquired on a BD FACSCalibur. Compensation controls were run with each assay to confirm instrument settings. A total of 160,000 ungated events were acquired for each sample using BD CellQuest software (version 5.2). TreeStar FlowJo software (version PC 7.5) was used for data analysis as described above.

[00282] For one donors tested, treatment of purified PBMC samples with huCD37-3, huCD37-3-SMCC-DM1, huCD37-50 or huCD37-50-SMCC-DM1 resulted in approximately 40% depletion of B-

cells (see Figure 4). There was less than 10% depletion of T cells, NK cells or monocytes. As seen for purified PBMCs, the *in vitro* depletion is restricted to B-cells indicating that the activity is linked to the high CD37 expression on B-cells. In comparison, treatment with the anti-CD20 antibody rituximab or the anti-CD37 SMIP™ TRU-016 resulted in a less than 10% depletion of B-cells. Alemtuzumab treatment resulted in depletion of 40% of B-cells, 80% of T cells, 15% of NK cells and 20% of monocytes.

Example 2D

Dose response for *in vitro* B-cell depletion using whole blood

[00283] To evaluate the dose-response of the antibodies and conjugates, whole blood from 2 donors was incubated with a 10-fold sample dilution series. Each sample dilution was added at 10 μ L per tube to 90 μ L of purified cells in triplicate and incubated for 1 hr at 37°C in a humidified 5% CO₂ incubator. The final concentration ranged from 10 μ g/mL to 0.1 ng/mL. The same amount of a non-binding huIgG Ab was used as an isotype control.

[00284] For two donors tested, treatment of whole blood samples with huCD37-3 or huCD37-3-SMCC-DM1 resulted in a clear dose response for the B-cell depletion activity (see Figure 5A and B). In addition, huCD37-50 was tested for one donor and also showed a similar dose response for the B-cell depletion activity (see Figure 5B). Incubation with huCD37-3, huCD37-3-SMCC-DM1 or huCD37-50 caused a maximum response of *in vitro* depletion of approximately 30-45% of B-cells with an EC₅₀ of 40-120 ng/mL.

[00285] In addition to the *in vitro* experiment described above, the capacity of CD37 antibodies to deplete B cells *in vivo* can be tested in huCD37 expressing mice (described in Example 3) and, for antibodies that crossreact with macaque CD37, in monkey.

Example 2E

In vitro cytokine release studies using human PBMCs

[00286] *In vitro* cytokine release was measured by ELISpot for IFN- γ (Interferon), TNF- α (Tumor Necrosis Factor) and IL-6 (Interleukin-6) using peripheral blood mononuclear cells (PBMCs) from healthy human donors incubated for 18-20 hours with compounds at a concentration of 2.5 ng/mL to 250 μ g/mL. The ELISpot method is designed to measure the number of cells secreting cytokine by capturing the cytokine onto the assay plate during the entire length of the incubation. In all assays the positive control anti-CD3 antibody CD3-2, as well as a negative non-binding isotype huIgG control antibody was included. Alemtuzumab (Campath®) and rituximab (Rituxan®) were used in comparison, since both have been reported to induce cytokine release in patients (Wing. *J Clin Invest.* 98:2819-26 (1996) and Winkler, *Blood* 94:2217-2224 (1999)). The assay conditions were chosen to reflect conditions that are relevant for antibody therapeutics. The highest concentration of 250 μ g/mL tested corresponds to the maximum

serum concentration of an antibody, such as for example the CD20-directed rituximab, in patient plasma after an infusion of 10 mg/kg of antibody.

[00287] As can be seen in Figures 6 and 7, the positive control anti-CD3 antibody induced release of very high levels of IFN- γ , TNF- α and IL-6 with PBMCs from two different donors. In the same assays, alemtuzumab caused intermediate cytokine release, while rituximab caused moderate cytokine release with PBMCs from two different donors. In contrast, huCD37-3, huCD37-50, huCD37-3-SMCC-DM1 or huCD37-50-SMCC-DM1 did not cause significant cytokine release in our assays.

[00288] This underscores the utility of the described CD37-targeting antibodies or conjugates as therapeutics as they combine potent activity, such as B-cell depletion, with a favorable safety profile with respect to cytokine release.

Example 3

In vivo models to evaluate the activity of CD37 directed antibodies or conjugates

[00289] B-cell depletion is known to ameliorate autoimmune diseases. In fact, rituximab has been approved for rheumatoid arthritis treatment (Edwards JC et al. Nat Rev Immunol. 6: 119 (2006)). In animal models, B-cell depletion using antibodies against B-cell antigens such as CD20, CD19 and CD79 has been shown to inhibit or ameliorate several autoimmune diseases including systemic lupus erythematosus (SLE), experimental autoimmune encephalomyelitis (EAE; mouse model of multiple sclerosis), type-1 diabetes (T1D) and rheumatoid arthritis (RA). The CD37 antigen is expressed at high levels in human B-cells. Therefore, antibodies or immunoconjugates directed against the CD37 antigen could potentially deplete B-cells and be therefore useful to treat multiple autoimmune diseases.

[00290] To test the utility of CD37 targeting antibodies and immunoconjugates to treat human autoimmune diseases, the activity of such CD37 targeting antibodies and immunoconjugates can be studied in mice using several murine autoimmune disease models.

[00291] For example, anti-murine CD37 antibodies can be generated using CD37-knock-out mice or other species such as rat and hamster, and antibodies that deplete B-cell *in vivo* effectively can be selected. The therapeutic potential of anti-CD37 antibodies can be tested in mouse models representing human autoimmune diseases, for example, a spontaneous T1D model in NOD mice, a myelin oligodendrocyte glycoprotein (MOG) peptide induced EAE model in wild type C57/Bl6 mice, a collagen induced rheumatoid arthritis model in DBA/1 mice or a spontaneous systemic lupus erythematosus (SLE) model in MRL/lpr mice. Examples of murine CD37 antibodies and their therapeutic efficacy in various animal models of autoimmune disease are provided below.

[00292] Alternatively, the therapeutic potential of anti-human CD37 antibodies and immunoconjugates can also be tested in murine autoimmune disease models that have been engineered to

express the human CD37 antigen. Such human CD37 (huCD37) expressing mice can be generated using standard knock in (KI) or transgenic (Tg) approaches. For example, to generate huCD37 KI mice, human CD37 cDNA can be inserted into the murine CD37 locus in the C57/Bl6 embryonic stem (ES) cells. The homozygous huCD37 KI mice will express human CD37 cDNA under the regulation of the endogenous murine CD37 promoter, thus the expression pattern of the huCD37 would mimic that of the endogenous muCD37. The different approach utilizes bacterial artificial chromosome (BAC) containing the human CD37 gene that can be randomly inserted into the mouse genome. This transgenic approach has been used successfully to generate huCD20 Tg mice resulting in B-cell specific high level expression of the antigen.

[00293] The resulting huCD37 expressing mice based on the C57/Bl6 background can be used to further develop several autoimmune disease model. For examples, MOG peptide immunization in the C57/Bl6 strain background can induces severe EAE in two weeks. In addition, introducing a FcγRIIB knock out phenotype by breeding huCD37 expressing mice with C57/Bl6 FcγRIIB knock out mice should yield a mouse model that spontaneously develop SLE and develop RA upon immunization with collagen II antigen. Alternatively, backcrossing of the huCD37 expressing C57/Bl6 mice into the NOD or MRL/lpr background for 10 generations can provide spontaneous T1D and SLE models, respectively.

Example 4A

Generation of anti-muCD37 monoclonal antibody clone 252-3

[00294] To develop proof of concept that CD37 targeting antibody and immunoconjugate can inhibit autoimmune disease, anti-murine CD37 (muCD37) monoclonal antibodies were generated by immunizing CD37-knock-out C57Bl/6 mice with 300-19, a murine pre-B cell line that endogenously expresses the muCD37 antigen. The immunogen was injected subcutaneously at the dose of 5×10^6 cells per mouse every 2 weeks for 5 times. Three days before being sacrificed for hybridoma generation, the immunized mice received intraperitoneal injection of another dose of antigen. The spleen cells were fused with murine myeloma P3X63Ag8.653 cells (P3 cells) (J. F. Kearney et al. 1979, *J Immunol*, 123: 1548-1550) at ratio of 1 P3 cells: 3 spleen cells according to standard procedure. The fused cells were cultured in RPMI-1640 selection medium containing hypoxanthine-aminopterin-thymidine (HAT) (Sigma Aldrich) in 5% CO₂ incubator at 37°C until hybridoma clones were ready for antibody screening.

[00295] Screening was done using flow cytometric binding assay with spleen cells from wild type mice and CD37-knock-out mice. The spleen cells were counterstained with anti-CD45R (B220) antibody to identify B cells that constitutively express CD37 antigen. The hybridomas producing antibody that bound the wild type, but not CD37-knock-out, B cells were subcloned by limiting dilution. One stable subclone (clone 252-3) was obtained. The 252-3 hybridoma was expanded in low IgG serum containing media and the antibody was purified using standard methods with protein A/G chromatography.

Example 4B

Characterization of anti-muCD37 monoclonal antibody clone 252-3

[00296] The purified 252-3 monoclonal antibody was identified as a mouse IgG2a with IsoStrip mouse monoclonal antibody isotyping kit (Roche Diagnostics Corporation, Indianapolis, IN). To determine the binding affinity to the muCD37 antigen, various concentrations of 252-3 antibody were incubated with 300-19 cells, a murine pre-B cell line that expresses the muCD37 antigen, for 30 minutes at 4°C. Cells were then washed and counterstained with anti muIgG-PE conjugate (Jackson Immunoresearch, West Grove, PA) for 30 minutes at 4°C. The cells were finally washed, fixed in formalin and analyzed by flow cytometry using a FACSarray (BD Bioscience, San Jose, CA). The flow cytometry data were analyzed using FlowJo (Tree Star Inc., Ashland, OR) and the geometric mean fluorescence intensity was plotted against the antibody concentration in a semi-log plot (Figure 8). A dose-response curve was generated by non-linear regression and the EC50 value of the curve, which corresponds to the apparent dissociation constant (Kd) of the antibody, was calculated using GraphPad Prism (GraphPad Software Inc., La Jolla, CA). It was found that the Kd of the 252-3 antibody was 14 nM. In contrast, the 252-3 antibody did not bind to human tumor cells expressing the human CD37 antigen. The 252-3 antibody was then used as a surrogate antibody in murine autoimmune disease models to demonstrate the therapeutic potential of a CD37-targeting antibody for the treatment of autoimmune diseases (Examples 5-7).

Example 5

Anti-muCD37 monoclonal antibody inhibits experimental autoimmune encephalomyelitis

[00297] Experimental autoimmune encephalomyelitis (EAE) is an animal model of inflammatory demyelinating disease of the central nervous system (CNS), including multiple sclerosis in human. Murine EAE is commonly induced by immunization of spinal cord homogenates, brain extracts, or CNS protein such as myelin protein or peptide, followed by injection of pertussis toxin to break down the blood-brain barrier and allow immune cells access to the CNS tissue. This immunization leads to multiple small disseminated lesions of demyelination in the brain and spinal cord, causing tail paralysis followed by limb paralysis.

[00298] To test the activity of anti-muCD37 antibody in the EAE model, we first studied the capacity of the 252-3 antibody to deplete B cells *in vivo*. C57Bl/6 mice were injected intraperitoneally with 25 mg/kg of 252-3 antibody or polyclonal murine IgG (Jackson Immunoresearch, West Grove, PA) as a control. Peripheral blood was collected at different time points and analyzed for B and T cell levels by flow cytometry. Allophycocyanin (APC)-conjugated anti-mouse CD45R (B220) antibody (ebioscience,

San Diego, CA) and fluorescein isothiocyanate (FITC)-conjugated anti CD3 ϵ antibody (ebioscience, San Diego, CA) were used to stain B and T cell populations, respectively. B cell depletion was assessed by calculating the ratio of B to T cells for each sample and the B/T ratio was normalized by setting the average B/T ratio of murine IgG-treated samples to 100%. The normalized B/T cell ratio was plotted for muIgG control mice and 252-3 antibody treated mice (Figure 9A). The result show that the B cell level of the mice treated with 252-3 antibody was rapidly reduced within a few hours after the antibody injection. The B cell depletion reached ~70% at 3h and peaked at day 3 (> 95%). After day 3, the B cell level slowly increased and reached ~60% of the normal level at day 14. This data suggests that the 252-3 antibody can rapidly and efficiently deplete peripheral blood B cells, and this effect was sustained for at least 7 days after the antibody injection.

[00299] The second study tested the capacity of 252-3 antibody to inhibit EAE. In this study, EAE was induced in C57Bl/6 mice by subcutaneous immunization of MOG₃₅₋₅₅ peptide emulsified in complete Freund's adjuvant (EAE kit from Hooke Laboratories, Lawrence, MA) into the upper and lower back at day 0 and two intraperitoneal injections of pertussis toxin at 2h and 24h after antigen immunization. Mice were checked for EAE signs daily starting on day 7 after immunization. The disease severity was scored on a scale of 0 to 5 using the following criteria:

Score	Clinical Observations
0	No obvious changes in motor functions of the mouse in comparison to non-immunized mice. When picked up by the tail, the tail has tension and is erect. Hind legs are usually spread apart. When the mouse is walking, there is no gait or head tilting.
1	Limp tail. When the mouse is picked up by tail, instead of being erect, the whole tail drapes over your finger.
2	Limp tail and weakness of hind legs. When the mouse is picked up by tail, legs are not spread apart, but held closer together. When the mouse is observed when walking, it has clearly apparent wobbly walk.
3	Limp tail and complete paralysis of hind legs (most common) OR, Limb tail with paralysis of one front and one hind leg. OR, ALL of: * Severe head tilting * Walking only along the edges of the cage * Pushing against the cage wall * Spinning when picked up by the tail

4	Limp tail, complete hind leg and partial front leg paralysis. Mouse is minimally moving around the cage but appears alert and feeding. Usually, euthanasia is recommended after the mouse scores level 4 for 2 days. When the mouse is euthanized because of severe paralysis, a score of 5 is entered for that mouse for the rest of the experiment.
5	Complete hind and front leg paralysis, no movement around the cage. OR, Mouse is spontaneously rolling in the cage. OR, Mouse is found dead due to paralysis. If mouse is alive, euthanize the mouse immediately if it scores 5. Once mouse scored 5, the same score is entered for all the days for the rest of the experiment.

[00300] All mice started to show signs of EAE between 12 to 18 days after antigen immunization. At the disease onset, mice were randomized and the 252-3 antibody or polyclonal muIgG was injected once intraperitoneally at a 25 mg/kg dose. A total of 10 mice were enrolled for each group. At the end of the study (18 days after the disease onset), the data were synchronized based on the day of disease onset for each mouse. The disease progression plot (Figure 9B) shows that mice from both groups had relapsing-remitting form of EAE. During the first wave of clinical symptoms, the control mice reached the mean of 3 while the mice treated with 252-3 antibody had a mean of 2. The difference in disease severity between these two groups was sustained for more than 2 weeks after the disease onset. Taken together, this data suggests that the 252-2 antibody treatment rapidly depletes the B cell population and alleviates EAE.

Example 6

Anti-muCD37 monoclonal antibody inhibits type-1 diabetes in NOD mice

[00301] Type-1 diabetes (T1D) or juvenile diabetes or insulin-dependent diabetes mellitus (IDDM) is caused by auto-immune reaction against insulin-producing pancreatic beta cells. Destruction of beta cells reduces insulin production and increases glucose level that produces various clinical symptoms. T1D incidence in Northern Europe and the US is between 8 and 17/100,000. Insulin supplement is the most common treatment of the disease.

[00302] Non-obese diabetic (NOD) mice spontaneously develop T1D and have been widely used to model the human disease. In NOD mice, the disease starts with leukocytic infiltration of the pancreatic islets (called insulinitis) as early as 4 weeks of age. The insulinitis progresses rapidly, leading to destruction of pancreatic islets and diabetes starting at 12-15 weeks of age. B cell depletion using anti-CD20 antibody in the early stage of insulinitis has been reported to delay the disease onset (Hu et al., J Clin Invest. 117, 3857 (2007)), suggesting that B cells play a critical role in the disease pathogenesis in NOD mice.

[00303] To test the activity of anti-muCD37 antibody, the 252-3 antibody was injected into six female NOD mice intraperitoneally at 25 mg/kg every 10 days for a total of 4 injections starting at 5 weeks of age (n=6). The control mice (n=6) were injected with polyclonal murine IgG (Jackson ImmunoResearch, West Grove, PA). Three days after the last injection, the B and T cell levels in peripheral blood were examined by flow cytometry. Allophycocyanin (APC)-conjugated anti-mouse CD45R (B220) antibody (ebioscience, San Diego, CA) and fluorescein isothiocyanate (FITC)-conjugated anti CD3 ϵ antibody (ebioscience, San Diego, CA) were used to stain B and T cell populations, respectively. The B/T cell ratio was normalized to murine IgG control treated samples as described above and the normalized B/T cell ratio was plotted for muIgG control mice and 252-3 antibody treated mice (Figure 10A). The results show that the B cell level of the mice treated with 252-3 antibody was significantly reduced as compared to the control mice, suggesting that the 252-3 antibody efficiently depletes peripheral blood B cells in NOD mice. To examine the effect of B cell depletion by anti-muCD37 antibody, blood glucose level was measured weekly starting at 12 weeks of age. Mice with blood glucose level \geq 250 mg/dL in two consecutive weeks are considered diabetic. The data in Figure 10B shows that the control mice started to develop diabetes on week 15 and 83% of the mice had diabetes on week 22. In contrast, the mice treated with 252-3 antibody started to develop diabetes on week 17 and only 50% of the mice were diabetic on week 27. This data shows that treatment of 252-3 antibody efficiently depletes B cells in NOD mice, delays the onset of diabetes and significantly reduces the disease incidence.

Example 7

Anti-muCD37 monoclonal antibody inhibits collagen-induced arthritis

[00304] Collagen-induced arthritis (CIA) is an animal model of rheumatoid arthritis (RA) that is widely used to investigate disease pathogenesis and to validate therapeutic targets. Arthritis is normally induced in mice or rats by immunization with autologous or heterologous type II collagen in adjuvant. This immunization elicits a robust T- and B- cell response to the antigen leading to proliferative synovitis with infiltration of polymorphonuclear and mononuclear cells, pannus formation, cartilage degradation, bone erosion and fibrosis.

[00305] Since different mouse strains have different susceptibility to antibody-mediated B cell depletion (Ahuja et al., *J. Immunol.*, 179: 3351-3361 (2007)), to test the activity of anti-muCD37 antibody in CIA model, we first studied the capacity of the 252-3 antibody to deplete B cells in DBA/1 mice. Mice were injected intraperitoneally with 25 mg/kg of 252-3 antibody or polyclonal murine IgG (Jackson ImmunoResearch, West Grove, PA) as control. Peripheral blood was collected at different time points and analyzed for B and T cell levels by flow cytometry. Allophycocyanin (APC)-conjugated anti-mouse CD45R (B220) antibody (ebioscience, San Diego, CA) and fluorescein isothiocyanate (FITC)-conjugated anti CD3 ϵ antibody (ebioscience, San Diego, CA) were used to stain B and T cell populations,

respectively. The normalized B/T cell ratio was calculated as described above and compared between the muIgG control mice and 252-3 antibody treated mice (Figure 11A). The result show that the 252-3 antibody significantly reduced the peripheral blood B cell level to ~20% and ~8% in 1 and 3 days after the antibody injection, and this low B cell level was maintained at 7 days after the antibody injection. This data suggests that the 252-3 antibody can rapidly and efficiently deplete peripheral blood B cells, and this effect was sustained for at least 7 days after the antibody injection.

[00306] The second study tests the capacity of 252-3 antibody to inhibit CIA. In this study, CIA was induced in DBA/1 mice by subcutaneous immunization of chicken collagen/CFA (complete Freund's adjuvant) on day 0 and chicken collagen/IFA (incomplete Freund's adjuvant) on day 21 (Hooke Laboratories, Lawrence, MA). Mice were checked for CIA signs daily starting on day 21 after immunization. The CIA severity was scored on a scale of 0 to 16 (based on a score of 0 to 4 for each paw) using the following criteria:

Paw Score	Clinical Observations
0	Normal paw.
1	One toe inflamed and swollen.
2	More than one toe, but not entire paw, inflamed and swollen, OR Mild swelling of entire paw.
3	Entire paw inflamed and swollen.
4	Very inflamed and swollen paw or ankylosed paw. If the paw is ankylosed, the mouse cannot grip the wire top of the cage.

[00307] At the onset of arthritis symptoms, mice were randomized into two groups and injected with the 252-3 antibody or polyclonal muIgG intraperitoneally at 10 mg/kg dose at three consecutive days. A total of 12 mice were enrolled for each group. At the end of the study (21 days after the disease onset), the data were synchronized based on the day of disease onset for each mouse. The disease progression plot (Figure 11B) shows that the disease severity in control mice increased rapidly from mean score of 2 at day 1 to 9.5 at day 7. In contrast, the disease in mice treated with the 252-3 antibody progressed significantly slower with mean score of 4.4 at day 7. Altogether, this data suggests that the 252-2 antibody treatment significantly depletes the B cell population and alleviates CIA.

[00308] In conclusion, the above experiments using a surrogate anti-muCD37 antibody provide evidence that a CD37-targeting antibody, or an immunoconjugate that includes a CD37 antibody, can inhibit autoimmune diseases in animal models.

[00309] It is to be appreciated that the Detailed Description section, and not the Abstract section, is intended to be used to interpret the claims. The Abstract may set forth one or more but not all

exemplary embodiments of the present invention as contemplated by the inventors, and thus, is not intended to limit the present invention and the appended claims in any way.

[00310] The present invention has been described above with the aid of functional building blocks illustrating the implementation of specified functions and relationships thereof. The boundaries of these functional building blocks have been arbitrarily defined herein for the convenience of the description. Alternate boundaries can be defined so long as the specified functions and relationships thereof are appropriately performed.

[00311] The foregoing description of the specific embodiments will so fully reveal the general nature of the invention that others can, by applying knowledge within the skill of the art, readily modify and/or adapt for various applications such specific embodiments, without undue experimentation, without departing from the general concept of the present invention. Therefore, such adaptations and modifications are intended to be within the meaning and range of equivalents of the disclosed embodiments, based on the teaching and guidance presented herein. It is to be understood that the phraseology or terminology herein is for the purpose of description and not of limitation, such that the terminology or phraseology of the present specification is to be interpreted by the skilled artisan in light of the teachings and guidance.

[00312] The breadth and scope of the present invention should not be limited by any of the above-described exemplary embodiments, but should be defined only in accordance with the following claims and their equivalents.

[00313] All publications, patents, and patent applications mentioned in this specification are herein incorporated by reference to the same extent as if each independent publication, patent, or patent application was specifically and individually indicated to be incorporated by reference.

WHAT IS CLAIMED IS:

1. A method for depleting a B-cell comprising contacting a population of cells comprising a non-cancerous B-cell with an antibody or antigen binding fragment thereof that specifically binds to CD37, wherein said antibody or fragment thereof is capable of inducing apoptosis *in vitro* in the absence of a cross-linking agent.
2. A method for treating a patient having an autoimmune or inflammatory disease comprising administering to said patient a therapeutically effective amount of an antibody or antigen-binding fragment thereof that specifically binds to CD37, wherein said antibody or fragment thereof is capable of inducing apoptosis *in vitro* in the absence of a cross-linking agent.
3. The method of claim 1 or 2, wherein said antibody or antigen-binding fragment thereof is also capable of inducing complement dependent cytotoxicity (CDC).
4. The method of any of claims 1-3, wherein said antibody or antigen-binding fragment thereof is also capable of inducing antibody dependent cell mediated cytotoxicity (ADCC).
5. A method for depleting a B-cell comprising contacting a population of cells comprising a non-cancerous B-cell with an antibody or antigen binding fragment thereof that specifically binds to the same CD37 epitope as an antibody selected from the group consisting of:
 - (a) an antibody comprising the polypeptide of SEQ ID NO:55 and the polypeptide of SEQ ID NO:72;
 - (b) an antibody comprising the polypeptide of SEQ ID NO:56 and the polypeptide of SEQ ID NO:73;
 - (c) an antibody comprising the polypeptide of SEQ ID NO:57 and the polypeptide of SEQ ID NO:74;
 - (d) an antibody comprising the polypeptide of SEQ ID NO:58 and the polypeptide of SEQ ID NO:74;
 - (e) an antibody comprising the polypeptide of SEQ ID NO:59 and the polypeptide of SEQ ID NO:75;
 - (f) an antibody comprising the polypeptide of SEQ ID NO:60 and the polypeptide of SEQ ID NO:76;
 - (g) an antibody comprising the polypeptide of SEQ ID NO:61 and the polypeptide of SEQ ID NO:77;
 - (h) an antibody comprising the polypeptide of SEQ ID NO:62 and the polypeptide of SEQ ID NO:78;
 - (i) an antibody comprising the polypeptide of SEQ ID NO:63 and the polypeptide of SEQ ID NO:79;
 - (j) an antibody comprising the polypeptide of SEQ ID NO:64 and the polypeptide of SEQ ID NO:80;
 - (k) an antibody comprising the polypeptide of SEQ ID NO:65 and the polypeptide of SEQ ID NO:81;
 - (l) an antibody comprising the polypeptide of SEQ ID NO:66 and the polypeptide of SEQ ID NO:82;
 - (m) an antibody comprising the polypeptide of SEQ ID NO:67 and the polypeptide of SEQ ID NO:83;
 - (n) an antibody comprising the polypeptide of SEQ ID NO:68 and the polypeptide of SEQ ID NO:84;
 - (o) an antibody comprising the polypeptide of SEQ ID NO:69 and the polypeptide of SEQ ID NO:85;

- (p) an antibody comprising the polypeptide of SEQ ID NO:70 and the polypeptide of SEQ ID NO:86;
- (q) an antibody comprising the polypeptide of SEQ ID NO:71 and the polypeptide of SEQ ID NO:87; and
- (r) an antibody comprising the polypeptide of SEQ ID NO:177 and the polypeptide of SEQ ID NO:178.

6. A method for treating a patient having an autoimmune or inflammatory disease comprising administering to said patient a therapeutically effective amount of an antibody or antigen-binding fragment thereof that specifically binds to the same CD37 epitope as an antibody selected from the group consisting of:

- (a) an antibody comprising the polypeptide of SEQ ID NO:55 and the polypeptide of SEQ ID NO:72;
- (b) an antibody comprising the polypeptide of SEQ ID NO:56 and the polypeptide of SEQ ID NO:73;
- (c) an antibody comprising the polypeptide of SEQ ID NO:57 and the polypeptide of SEQ ID NO:74;
- (d) an antibody comprising the polypeptide of SEQ ID NO:58 and the polypeptide of SEQ ID NO:74;
- (e) an antibody comprising the polypeptide of SEQ ID NO:59 and the polypeptide of SEQ ID NO:75;
- (f) an antibody comprising the polypeptide of SEQ ID NO:60 and the polypeptide of SEQ ID NO:76;
- (g) an antibody comprising the polypeptide of SEQ ID NO:61 and the polypeptide of SEQ ID NO:77;
- (h) an antibody comprising the polypeptide of SEQ ID NO:62 and the polypeptide of SEQ ID NO:78;
- (i) an antibody comprising the polypeptide of SEQ ID NO:63 and the polypeptide of SEQ ID NO:79;
- (j) an antibody comprising the polypeptide of SEQ ID NO:64 and the polypeptide of SEQ ID NO:80;
- (k) an antibody comprising the polypeptide of SEQ ID NO:65 and the polypeptide of SEQ ID NO:81;
- (l) an antibody comprising the polypeptide of SEQ ID NO:66 and the polypeptide of SEQ ID NO:82;
- (m) an antibody comprising the polypeptide of SEQ ID NO:67 and the polypeptide of SEQ ID NO:83;
- (n) an antibody comprising the polypeptide of SEQ ID NO:68 and the polypeptide of SEQ ID NO:84;
- (o) an antibody comprising the polypeptide of SEQ ID NO:69 and the polypeptide of SEQ ID NO:85;
- (p) an antibody comprising the polypeptide of SEQ ID NO:70 and the polypeptide of SEQ ID NO:86;
- (q) an antibody comprising the polypeptide of SEQ ID NO:71 and the polypeptide of SEQ ID NO:87; and
- (r) an antibody comprising the polypeptide of SEQ ID NO:177 and the polypeptide of SEQ ID NO:178.

7. The method of claim 5 or 6, wherein the antibody or antigen-binding fragment thereof competitively inhibits an antibody selected from the group consisting of:

- (a) an antibody comprising the polypeptide of SEQ ID NO:55 and the polypeptide of SEQ ID NO:72;
- (b) an antibody comprising the polypeptide of SEQ ID NO:56 and the polypeptide of SEQ ID NO:73;
- (c) an antibody comprising the polypeptide of SEQ ID NO:57 and the polypeptide of SEQ ID NO:74;
- (d) an antibody comprising the polypeptide of SEQ ID NO:58 and the polypeptide of SEQ ID NO:74;
- (e) an antibody comprising the polypeptide of SEQ ID NO:59 and the polypeptide of SEQ ID NO:75;
- (f) an antibody comprising the polypeptide of SEQ ID NO:60 and the polypeptide of SEQ ID NO:76;

- (g) an antibody comprising the polypeptide of SEQ ID NO:61 and the polypeptide of SEQ ID NO:77;
- (h) an antibody comprising the polypeptide of SEQ ID NO:62 and the polypeptide of SEQ ID NO:78;
- (i) an antibody comprising the polypeptide of SEQ ID NO:63 and the polypeptide of SEQ ID NO:79;
- (j) an antibody comprising the polypeptide of SEQ ID NO:64 and the polypeptide of SEQ ID NO:80;
- (k) an antibody comprising the polypeptide of SEQ ID NO:65 and the polypeptide of SEQ ID NO:81;
- (l) an antibody comprising the polypeptide of SEQ ID NO:66 and the polypeptide of SEQ ID NO:82;
- (m) an antibody comprising the polypeptide of SEQ ID NO:67 and the polypeptide of SEQ ID NO:83;
- (n) an antibody comprising the polypeptide of SEQ ID NO:68 and the polypeptide of SEQ ID NO:84;
- (o) an antibody comprising the polypeptide of SEQ ID NO:69 and the polypeptide of SEQ ID NO:85;
- (p) an antibody comprising the polypeptide of SEQ ID NO:70 and the polypeptide of SEQ ID NO:86;
- (q) an antibody comprising the polypeptide of SEQ ID NO:71 and the polypeptide of SEQ ID NO:87; and
- (r) an antibody comprising the polypeptide of SEQ ID NO:177 and the polypeptide of SEQ ID NO:178.

8. A method for depleting a B-cell comprising contacting a population of cells comprising non-cancerous B-cell with an antibody or antigen-binding fragment thereof produced by hybridoma selected from the group consisting of ATCC Deposit Designation PTA-10664, deposited with the ATCC on February 18, 2010, ATCC Deposit Designation PTA-10665, deposited with the ATCC on February 18, 2010, ATCC Deposit Designation PTA-10666, deposited with the ATCC on February 18, 2010, ATCC Deposit Designation PTA-10667 deposited with the ATCC on February 18, 2010, ATCC Deposit Designation PTA-10668, deposited with the ATCC on February 18, 2010, ATCC Deposit Designation PTA-10669, deposited with the ATCC on February 18, 2010, and ATCC Deposit Designation PTA-10670, deposited with the ATCC on February 18, 2010.
9. A method for treating a patient having an autoimmune or inflammatory disease comprising administering to said patient a therapeutically effective amount of an antibody or antigen-binding fragment thereof produced by hybridoma selected from the group consisting of ATCC Deposit Designation PTA-10664, deposited with the ATCC on February 18, 2010, ATCC Deposit Designation PTA-10665, deposited with the ATCC on February 18, 2010, ATCC Deposit Designation PTA-10666, deposited with the ATCC on February 18, 2010, ATCC Deposit Designation PTA-10667 deposited with the ATCC on February 18, 2010, ATCC Deposit Designation PTA-10668, deposited with the ATCC on February 18, 2010, ATCC Deposit Designation PTA-10669, deposited with the ATCC on February 18, 2010, and ATCC Deposit Designation PTA-10670, deposited with the ATCC on February 18, 2010.

10. A method for depleting a B-cell comprising contacting a population of cells comprising a non-cancerous B-cell with an antibody or antigen-binding fragment thereof that specifically binds to CD37, wherein said antibody comprises polypeptide sequences selected from the group consisting of:

- (a) SEQ ID NOs: 4, 5, and 6 and SEQ ID NOs: 28, 29, and 30;
- (b) SEQ ID NOs: 7, 8, and 9 and SEQ ID NOs: 31, 32, and 33;
- (c) SEQ ID NOs: 10, 11, and 12 and SEQ ID NOs: 34, 35, and 36;
- (d) SEQ ID NOs: 13, 14, and 15 and SEQ ID NOs: 37, 38, and 39;
- (e) SEQ ID NOs: 13, 14, and 15 and SEQ ID NOs: 37, 40, and 39;
- (f) SEQ ID NOs: 16, 17, and 18 and SEQ ID NOs: 41, 42, and 43;
- (g) SEQ ID NOs: 19, 20, and 21 and SEQ ID NOs: 44, 45, and 46;
- (h) SEQ ID NOs: 19, 20, and 21 and SEQ ID NOs: 44, 47, and 46;
- (i) SEQ ID NOs: 22, 23, and 24 and SEQ ID NOs: 48, 49, and 50;
- (j) SEQ ID NOs: 22, 23, and 24 and SEQ ID NOs: 48, 51, and 50;
- (k) SEQ ID NOs: 25, 26, and 27 and SEQ ID NOs: 52, 53, and 54;
- (l) SEQ ID NOs: 171, 172, and 173 and SEQ ID NOs: 174, 175, and 176;
- (m) SEQ ID NOs: 171, 181, and 173 and SEQ ID NOs: 174, 175, and 176; and
- (n) variants of (a) to (m) comprising 1, 2, 3, or 4 conservative amino acid substitutions.

11. A method for treating a patient having an autoimmune or inflammatory disease comprising administering to said patient a therapeutically effective amount of an antibody or antigen-binding fragment thereof with an antibody or antigen-binding fragment thereof that specifically binds to CD37, wherein said antibody comprises polypeptide sequences selected from the group consisting of:

- (a) SEQ ID NOs: 4, 5, and 6 and SEQ ID NOs: 28, 29, and 30;
- (b) SEQ ID NOs: 7, 8, and 9 and SEQ ID NOs: 31, 32, and 33;
- (c) SEQ ID NOs: 10, 11, and 12 and SEQ ID NOs: 34, 35, and 36;
- (d) SEQ ID NOs: 13, 14, and 15 and SEQ ID NOs: 37, 38, and 39;
- (e) SEQ ID NOs: 13, 14, and 15 and SEQ ID NOs: 37, 40, and 39;
- (f) SEQ ID NOs: 16, 17, and 18 and SEQ ID NOs: 41, 42, and 43;
- (g) SEQ ID NOs: 19, 20, and 21 and SEQ ID NOs: 44, 45, and 46;
- (h) SEQ ID NOs: 19, 20, and 21 and SEQ ID NOs: 44, 47, and 46;
- (i) SEQ ID NOs: 22, 23, and 24 and SEQ ID NOs: 48, 49, and 50;
- (j) SEQ ID NOs: 22, 23, and 24 and SEQ ID NOs: 48, 51, and 50;
- (k) SEQ ID NOs: 25, 26, and 27 and SEQ ID NOs: 52, 53, and 54; and

(l) variants of (a) to (k) comprising 1, 2, 3, or 4 conservative amino acid substitutions.

12. The method of claim 10 or 11, wherein the antibody or antigen-binding fragment thereof comprises polypeptide sequences that are at least 90% identical to polypeptide sequences selected from the group consisting of:

- (a) SEQ ID NO:55 and SEQ ID NO:72;
- (b) SEQ ID NO:56 and SEQ ID NO:73;
- (c) SEQ ID NO:57 and SEQ ID NO:74;
- (d) SEQ ID NO:58 and SEQ ID NO:74;
- (e) SEQ ID NO:59 and SEQ ID NO:75;
- (f) SEQ ID NO:60 and SEQ ID NO:76;
- (g) SEQ ID NO:61 and SEQ ID NO:77;
- (h) SEQ ID NO:62 and SEQ ID NO:78;
- (i) SEQ ID NO:63 and SEQ ID NO:79;
- (j) SEQ ID NO:64 and SEQ ID NO:80;
- (k) SEQ ID NO:65 and SEQ ID NO:81;
- (l) SEQ ID NO:66 and SEQ ID NO:82;
- (m) SEQ ID NO:67 and SEQ ID NO:83;
- (n) SEQ ID NO:68 and SEQ ID NO:84;
- (o) SEQ ID NO:69 and SEQ ID NO:85;
- (p) SEQ ID NO:70 and SEQ ID NO:86;
- (q) SEQ ID NO:71 and SEQ ID NO:87; and
- (r) SEQ ID NO:177 and SEQ ID NO:178.

13. The method of claim 12, wherein the polypeptide sequences are at least 95% identical to polypeptide sequences selected from the group consisting of:

- (a) SEQ ID NO:55 and SEQ ID NO:72;
- (b) SEQ ID NO:56 and SEQ ID NO:73;
- (c) SEQ ID NO:57 and SEQ ID NO:74;
- (d) SEQ ID NO:58 and SEQ ID NO:74;
- (e) SEQ ID NO:59 and SEQ ID NO:75;
- (f) SEQ ID NO:60 and SEQ ID NO:76;
- (g) SEQ ID NO:61 and SEQ ID NO:77;
- (h) SEQ ID NO:62 and SEQ ID NO:78;

- (i) SEQ ID NO:63 and SEQ ID NO:79;
- (j) SEQ ID NO:64 and SEQ ID NO:80;
- (k) SEQ ID NO:65 and SEQ ID NO:81;
- (l) SEQ ID NO:66 and SEQ ID NO:82;
- (m) SEQ ID NO:67 and SEQ ID NO:83;
- (n) SEQ ID NO:68 and SEQ ID NO:84;
- (o) SEQ ID NO:69 and SEQ ID NO:85;
- (p) SEQ ID NO:70 and SEQ ID NO:86;
- (q) SEQ ID NO:71 and SEQ ID NO:87; and
- (r) SEQ ID NO:177 and SEQ ID NO:178.

14. The method of claim 13, wherein the polypeptide sequences are at least 99% identical to polypeptide sequences selected from the group consisting of:

- (a) SEQ ID NO:55 and SEQ ID NO:72;
- (b) SEQ ID NO:56 and SEQ ID NO:73;
- (c) SEQ ID NO:57 and SEQ ID NO:74;
- (d) SEQ ID NO:58 and SEQ ID NO:74;
- (e) SEQ ID NO:59 and SEQ ID NO:75;
- (f) SEQ ID NO:60 and SEQ ID NO:76;
- (g) SEQ ID NO:61 and SEQ ID NO:77;
- (h) SEQ ID NO:62 and SEQ ID NO:78;
- (i) SEQ ID NO:63 and SEQ ID NO:79;
- (j) SEQ ID NO:64 and SEQ ID NO:80;
- (k) SEQ ID NO:65 and SEQ ID NO:81;
- (l) SEQ ID NO:66 and SEQ ID NO:82;
- (m) SEQ ID NO:67 and SEQ ID NO:83;
- (n) SEQ ID NO:68 and SEQ ID NO:84;
- (o) SEQ ID NO:69 and SEQ ID NO:85;
- (p) SEQ ID NO:70 and SEQ ID NO:86;
- (q) SEQ ID NO:71 and SEQ ID NO:87; and
- (r) SEQ ID NO:177 and SEQ ID NO:178.

15. The method of any one of claims 1-14, wherein said antibody or antigen binding fragment thereof is murine, non-human, humanized, chimeric, resurfaced, or human.

16. The method of any one of claims 5-15, wherein said antibody or antibody fragment is capable of inducing apoptosis of a cell expressing CD37 *in vitro* in the absence of cross-linking agents.
17. The method of any one of claims 5-15, wherein said antibody or antigen binding fragment is capable of inducing complement dependent cytotoxicity (CDC).
18. The method of any one of claims 5-15, wherein said antibody is capable of inducing antibody dependent cell mediated cytotoxicity (ADCC).
19. A method for depleting a B-cell comprising contacting a population of cells comprising a non-cancerous B-cell with a human or humanized antibody or antigen binding fragment thereof that specifically binds to CD37, wherein said antibody or fragment thereof is capable of inducing complement dependent cytotoxicity (CDC).
20. A method for treating a patient having an autoimmune or inflammatory disease comprising administering to said patient a therapeutically effective amount of a human or humanized antibody or antigen binding fragment thereof that specifically binds to CD37, wherein said antibody or fragment thereof is capable of inducing complement dependent cytotoxicity (CDC).
21. The method of claim 19 or 20, wherein said human or humanized antibody or antigen binding fragment thereof is also capable of inducing inducing apoptosis *in vitro* in the absence of a cross-linking agent.
22. The method of any one of claims 20-21, wherein said human or humanized antibody or antigen binding fragment thereof is also capable of inducing antibody dependent cell mediated cytotoxicity (ADCC).
23. The method of any one of claims 1-22 wherein said antibody binds to human CD37 and macaque CD37.
24. The method of any one of claims 1-22, which is a full length antibody.
25. The method of any one of claims 1-22, which is an antigen-binding fragment.
26. The method of any one of claims 1-22, wherein said antibody or antigen-binding fragment thereof comprises a Fab, Fab', F(ab')₂, Fd, single chain Fv or scFv, disulfide linked Fv, V-NAR domain, IgNar, intrabody, IgGΔCH2, minibody, F(ab')₃, tetrabody, triabody, diabody, single-domain antibody, DVD-Ig, Fcab, mAb², (scFv)₂, or scFv-Fc.

27. The method of any one of claims 1-26, wherein the antibody or antigen-binding fragment thereof is linked via a linker (L) to a cytotoxic agent (C) to form an immunoconjugate.
28. A method for depleting a B-cell comprising contacting a population of cells comprising a non-cancerous B-cell with a composition comprising an immunoconjugate having the formula (A) - (L) - (C), wherein:
- (A) is an antibody or antigen binding fragment that specifically binds to CD37;
(L) is a non-cleavable linker; and
(C) is a cytotoxic agent; and
wherein said linker (L) links (A) to (C).
29. A method for treating a patient having an autoimmune or inflammatory disease comprising administering to said patient a therapeutically effective amount of a composition comprising an immunoconjugate having the formula (A) - (L) - (C), wherein:
- (A) is an antibody or antigen binding fragment that specifically binds to CD37;
(L) is a non-cleavable linker; and
(C) is a cytotoxic agent; and
wherein said linker (L) links (A) to (C).
30. A method for depleting a B-cell comprising contacting a population of cells comprising a non-cancerous B-cell with a composition comprising an immunoconjugate having the formula (A) - (L) - (C), wherein:
- (A) is an antibody or antigen binding fragment that specifically binds to CD37;
(L) is a linker; and
(C) is a maytansinoid; and
wherein said linker (L) links (A) to (C).
31. A method for treating a patient having an autoimmune or inflammatory disease comprising administering to said patient a therapeutically effective amount of a composition comprising an immunoconjugate having the formula (A) - (L) - (C), wherein:
- (A) is an antibody or antigen binding fragment that specifically binds to CD37;
(L) is a linker; and
(C) is a maytansinoid; and
wherein said linker (L) links (A) to (C).

32. The method of claim 30 or 31, wherein the linker is a non-cleavable linker.
33. The method of any one of claims 27-32, wherein the immunoconjugate further comprises a second (C).
34. The method of claim 33, wherein the immunoconjugate further comprises a third (C).
35. The method of claim 34, wherein the immunoconjugate further comprises a fourth (C).
36. The method of one of claims 27-32, wherein the immunoconjugate comprises 2-6 (C).
37. The method of claim 36, wherein the immunoconjugate comprises 3-4 (C).
38. The method of anyone of claims 27-37, wherein said linker is selected from the group consisting of a cleavable linker, a non-cleavable linker, a hydrophilic linker, and a dicarboxylic acid based linker.
39. The method of any one of claims 27-38, wherein said linker is selected from the group consisting of: N-succinimidyl 4-(2-pyridyldithio)pentanoate (SPP); N-succinimidyl 4-(2-pyridyldithio)butanoate (SPDB) or N-succinimidyl 4-(2-pyridyldithio)-2-sulfobutanoate (sulfo-SPDB); N-succinimidyl 4-(maleimidomethyl) cyclohexanecarboxylate (SMCC); N-sulfosuccinimidyl 4-(maleimidomethyl) cyclohexanecarboxylate (sulfoSMCC); N-succinimidyl-4-(iodoacetyl)-aminobenzoate (SIAB); and N-succinimidyl-[(N-maleimidopropionamido)-tetraethyleneglycol] ester (NHS-PEG4-maleimide).
40. The method of claim 39, wherein said linker is N-succinimidyl-[(N-maleimidopropionamido)-tetraethyleneglycol] ester (NHS-PEG4-maleimide).
41. The method of any one of claims 27-29 and 33-40, wherein said cytotoxic agent is selected from the group consisting of a maytansinoid, maytansinoid analog, doxorubicin, a modified doxorubicin, benzodiazepine, taxoid, CC-1065, CC-1065 analog, duocarmycin, duocarmycin analog, calicheamicin, dolastatin, dolastatin analog, aristatin, tomaymycin derivative, and leptomyacin derivative or a prodrug of the agent.
42. The method of claim 41, wherein said cytotoxic agent is a maytansinoid.
43. The method of any one of claims 30-32 or 42, wherein said cytotoxic agent is N(2')-deacetyl-N(2')-(3-mercapto-1-oxopropyl)-maytansine (DM1) or N(2')-deacetyl-N2-(4-mercapto-4-methyl-1-oxopentyl)-maytansine (DM4).

44. The method of any one of claims 27-43, wherein the composition comprising an immunoconjugate comprises multiple cytotoxic agents (C) with an average of about 3 to about 4 (C) per (A).
45. The method of claim 44, wherein the immunoconjugates have an average of about 3.5 ± 0.5 (C) per (A).
46. The method of claim 45, wherein the immunoconjugates have an average of about 3.5 (C) per (A).
47. The method of any one of 1-46, wherein said antibody or antigen-binding fragment is capable of depleting B-cells.
48. The method of any one of claims 1-47, wherein said antibody or antigen-binding fragment is capable of inhibiting T-cell responses.
49. The method of any one of claims 1, 3-5, 7, 8, 10, 12-19, 21-28, 30, or 32-48, wherein, the population of cells comprises a T-cell.
50. The method of any one of claims 1, 3-5, 7, 8, 10, 12-19, 21-28, 30, or 32-49, wherein the population of cells comprises peripheral blood mononuclear cells.
51. The method of claim 50, wherein the peripheral blood mononuclear cells were obtained from a human.
52. The method of any one of claims 1, 3-5, 7, 8, 10, 12-19, 21-28, 30, or 32-49, wherein the population of cells is in whole blood.
53. The method of claim 52, wherein the whole blood was obtained from a human.
54. The method of any one of claims 1, 3-5, 7, 8, 10, 12-19, 21-28, 30, or 32-49, wherein the population of cells is in an organism.
55. The method of claim 54, wherein the population of cells is in a patient having an autoimmune or inflammatory disease.
56. The method of any one of claims 1, 3-5, 7, 8, 10, 12-19, 21-28, 30, or 32-55, wherein the B-cell is an autoreactive B-cell.
57. The method of any one of claims 1, 3-5, 7, 8, 10, 12-19, 21-28, 30, or 32-56, wherein at least about 30% of B-cells in the population of cells are depleted.

58. The method of any one of claims 49-57, wherein less than about 5% of T-cells are depleted.
59. The method of any one of claims 2-4, 6, 7, 9, 11-18, 20-27, 29, or 31-48, further comprising administering a second therapeutic agent.
60. The method of claim 59, wherein the second therapeutic is selected from the group consisting of methotrexate, an anti-CD20 therapeutic, an anti-IL-6 receptor therapeutic, an anti-IL-12/23p40 therapeutic, a chemotherapeutic, an immunosuppressant, an anti-Interferon beta-1a therapeutic, glatiramer acetate, an anti- α 4-integrin therapeutic, fingolimod, an anti-BLys therapeutic, CTLA-Fc, or an anti-TNF therapeutic.
61. The method of claim 59, wherein the second therapeutic is an antibody directed against an antigen selected from a group consisting of CD3, CD14, CD19, CD20, CD22, CD25, CD28, CD30, CD33, CD36, CD38, CD40, CD44, CD52, CD55, CD59, CD56, CD70, CD79, CD80, CD103, CD134, CD137, CD138, and CD152.
62. The method of claim 59, wherein the second therapeutic is an antibody directed against an antigen selected from the group consisting of IL-2, IL-6, IL-12, IL-23, IL-12/23 p40, IL-17, IFN γ , TNF α , IFN α , IL-15, IL-21, IL-1a, IL-1b, IL-18, IL-8, IL-4, GM-CSF, IL-3, and IL-5.
63. The method of any one of 2-4, 6, 7, 9, 11-18, 20-28, 30, 32-47, or 58-62, wherein said autoimmune or inflammatory disease is selected from the group consisting of rheumatoid arthritis, multiple sclerosis, type I diabetes mellitus, idiopathic inflammatory myopathy, systemic lupus erythematosus (SLE), myasthenia gravis, Grave's disease, dermatomyositis, polymyositis, Crohn's disease, ulcerative colitis, gastritis, Hashimoto's thyroiditis, asthma, psoriasis, psoriatic arthritis, dermatitis, systemic scleroderma and sclerosis, inflammatory bowel disease (IBD), respiratory distress syndrome, meningitis, encephalitis, uveitis, glomerulonephritis, eczema, atherosclerosis, leukocyte adhesion deficiency, Raynaud's syndrome, Sjögren's syndrome, Reiter's disease, Behcet's disease, immune complex nephritis, IgA nephropathy, IgM polyneuropathies, immune-mediated thrombocytopenias, acute idiopathic thrombocytopenic purpura, chronic idiopathic thrombocytopenic purpura, hemolytic anemia, myasthenia gravis, lupus nephritis, atopic dermatitis, pemphigus vulgaris, opsoclonus-myoclonus syndrome, pure red cell aplasia, mixed cryoglobulinemia, ankylosing spondylitis, hepatitis C-associated cryoglobulinemic vasculitis, chronic focal encephalitis, bullous pemphigoid, hemophilia A, membranoproliferative glomerulonephritis, adult and juvenile dermatomyositis, adult polymyositis, chronic urticaria, primary biliary cirrhosis, neuromyelitis optica, Graves' dysthyroid disease, bullous pemphigoid, membranoproliferative glomerulonephritis, Churg-Strauss syndrome, juvenile onset diabetes,

hemolytic anemia, atopic dermatitis, systemic sclerosis, Sjögen's syndrome and glomerulonephritis, dermatomyositis, ANCA, aplastic anemia, autoimmune hemolytic anemia (AIHA), factor VIII deficiency, hemophilia A, autoimmune neutropenia, Castleman's syndrome, Goodpasture's syndrome, solid organ transplant rejection, graft versus host disease (GVHD), autoimmune hepatitis, lymphoid interstitial pneumonitis, HIV, bronchiolitis obliterans (non-transplant), Guillain-Barre Syndrome, large vessel vasculitis, giant cell (Takayasu's) arteritis, medium vessel vasculitis, Kawasaki's Disease, polyarteritis nodosa. Wegener's granulomatosis, microscopic polyangiitis (MPA), Omenn's syndrome, chronic renal failure, acute infectious mononucleosis, HIV and herpes virus associated diseases.

64. An antibody or antigen binding fragment thereof that specifically binds to the same CD37 epitope as an antibody comprising the polypeptide of SEQ ID NO:177 and the polypeptide of SEQ ID NO:178.
65. An antibody or antigen binding fragment thereof that specifically binds to CD37, wherein said antibody or fragment comprises the sequences of SEQ ID NOS: 171, 172 or 181, and 173 and SEQ ID NOS: 174, 175, and 176.
66. The antibody or antigen binding fragment of claim 64 or 65, wherein said antibody or fragment comprises polypeptide sequences at least 90%, at least 95%, or at least 99% identical to the sequences of SEQ ID NO:177 and SEQ ID NO:178.
67. An immunoconjugate comprising the antibody or antigen binding fragment of any one of claims 64-66, a linker, and a cytotoxic agent.
68. A method for depleting a B-cell comprising contacting a population of cells comprising a B-cell with the antibody or antigen-binding fragment of any one of claims 64-66 or the immunoconjugate of claim 67.