

(19) **DANMARK**

(10) **DK/EP 3548061 T3**



(12) **Oversættelse af  
europæisk patentskrift**

Patent- og  
Varemærkestyrelsen

- 
- (51) Int.Cl.: **A 61 K 38/00 (2006.01)**      **A 61 P 21/00 (2006.01)**      **A 61 P 25/00 (2006.01)**  
**A 61 P 25/02 (2006.01)**      **A 61 P 25/08 (2006.01)**      **A 61 P 25/14 (2006.01)**  
**A 61 P 25/16 (2006.01)**      **A 61 P 25/24 (2006.01)**      **A 61 P 25/28 (2006.01)**  
**C 07 K 14/00 (2006.01)**      **C 07 K 14/605 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2020-12-14**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2020-09-09**
- (86) Europæisk ansøgning nr.: **17811693.5**
- (86) Europæisk indleveringsdag: **2017-12-04**
- (87) Den europæiske ansøgnings publiceringsdag: **2019-10-09**
- (86) International ansøgning nr.: **GB2017053655**
- (87) Internationalt publikationsnr.: **WO2018104718**
- (30) Prioritet: **2016-12-05 GB 201620611**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **Lancaster University Business Enterprises Limited, University House , Bailrigg, Lancaster LA1 4YW, Storbritannien**
- (72) Opfinder: **HÖLSCHER, Christian, Lancaster University, Furness College, Lancaster, Lancashire LA1 4YQ, Storbritannien**
- (74) Fuldmægtig i Danmark: **HØIBERG P/S, Adelgade 12, 1304 København K, Danmark**
- (54) Benævnelse: **Behandling af neurologiske sygdomme**
- (56) Fremdragne publikationer:  
**WO-A1-2013/164483**  
**WO-A1-2013/192129**  
**US-A1- 2016 015 788**  
**JI CHENHUI ET AL: "A novel dual GLP-1 and GIP receptor agonist is neuroprotective in the MPTP mouse model of Parkinson's disease by increasing expression of BDNF", BRAIN RESEARCH, ELSEVIER, AMSTERDAM, NL, vol. 1634, 11 October 2015 (2015-10-11), pages 1-11, XP029449878, ISSN: 0006-8993, DOI: 10.1016/J.BRAINRES.2015.09.035**  
**VERSPHOHL ET AL: "Novel therapeutics for type 2 diabetes: Incretin hormone mimetics (glucagon-like peptide-1 receptor agonists) and dipeptidyl peptidase-4 inhibitors", PHARMACOLOGY AND THERAPEUTICS, ELSEVIER, GB, vol. 124, no. 1, 1 October 2009 (2009-10-01), pages 113-138, XP026501107, ISSN: 0163-7258, DOI: 10.1016/J.PHARMTHERA.2009.06.002 [retrieved on 2009-06-21]**



# DESCRIPTION

## Technical Field of the Invention

[0001] Aspects and embodiments of the present invention relate to the treatment of neurological disorders such as for example, Alzheimer's disease and Parkinson's disease. Particularly, certain embodiments relate to GIP/GLP-1 co-agonist peptides for use in the treatment of these two neurological disorders. Also included in the present invention are *inter alia* pharmaceutical compositions comprising the GIP/GLP-1 co-agonist peptides, together with methods of treating such disorders as well as other subject matter.

## Background to the Invention

[0002] Alzheimer's disease (AD) is a chronic neurodegenerative disorder for which there is at present no cure. Currently prescribed medication only temporarily relieves some of the symptoms. The main hallmarks of the disease are disorientation, loss of memory, loss of neurons and synapses in the brain, the accumulation of beta-amyloid protein in the brain (amyloid plaques), and intracellular aggregation of hyperphosphorylated tau protein (tangles) (LaFerla and Oddo, 2005; Blennow et al., 2006).

[0003] Parkinson's disease (PD) is also a chronic neurodegenerative disease for which currently only delaying medication is available. The main hallmarks are tremor, rigor, and a loss of ability to move, the degeneration of neurons in the basal brain (substantia nigra) and the loss of release of the neurotransmitter, dopamine (Shen, 2010).

[0004] Type 2 diabetes (T2DM) has been identified as a risk factor for AD and PD (Holscher, 2014), indicating that insulin signalling impairment may be a factor in initiating or accelerating the development of AD. Epidemiological studies found a clear correlation between T2DM and the risk of developing AD or other neurodegenerative disorders at a later stage (Luchsinger et al., 2004; Ristow, 2004; Ohara et al., 2011). It was also shown that insulin signalling in the brain is desensitised in AD patients. Recent studies demonstrated that brains of AD patients had increased levels of inactivated phosphorylated insulin receptors and IRS-1 second messengers, which are both indicative of insulin desensitisation (Moloney et al., 2010; Bomfim et al., 2012; Talbot et al., 2012). In PD, insulin signalling was also found to be impaired and linked to disease progression (Morris et al., 2011; Cereda et al., 2012).

[0005] Glucagon-like peptide (GLP-1) is an endogenous 30-amino acid peptide incretin hormone (Baggio and Drucker, 2007). GLP-1 receptor stimulation enhances beta-cell proliferation in the pancreas by activating stem cell proliferation, facilitates glucose-dependent insulin secretion and lowers blood glucose in patients with T2DM (Lovshin and Drucker, 2009). Three GLP-1 analogues are currently on the market as a treatment for diabetes, exendin-4

(Byetta®, Bydureon®), lixisenatide (Lyxumia®) and liraglutide (Victoza®) (Campbell and Drucker, 2013; Elkinson and Keating, 2013).

**[0006]** Glucose-dependent insulintropic peptide (GIP), also known as gastric inhibitory polypeptide, is a 42-amino acid incretin hormone which activates pancreatic islets to enhance insulin secretion and to help reduce postprandial hyperglycaemia, similar to GLP-1 (Gault et al., 2003). GIP is a member of the secretin/glucagon family of neuroregulatory polypeptides which also include the growth hormone releasing factor. It is expressed in pancreatic alpha cells, endocrine cells, and also in neurons in the brain (Nyberg et al., 2007; Campbell and Drucker, 2013). GIP has also been shown to promote pancreatic beta-cell growth, differentiation, proliferation and cell survival, documenting its growth-hormone properties (Gault et al., 2003). Therefore, research is on-going to develop GIP as a therapeutic tool for T2DM treatment (Irwin et al., 2006). There are currently no GIP analogues authorised for the treatment of T2DM.

**[0007]** Dual agonist peptides which target more than one receptor are being considered for the treatment of T2DM. Several GIP/GLP-1 co-agonist peptides are currently in development for the treatment of T2D. However, there are currently no GIP/GLP-1 dual receptor agonists authorised for use to treat T2DM.

**[0008]** Recent investigations of the neuroproperties of GLP-1 and GIP have indicated that these peptides may play a role in preventing neurodegenerative hallmarks in several mouse models of Alzheimer's disease (AD) and also in animal models of Parkinson's disease (PD).

**[0009]** Insulin as well as the incretins not only have growth-factor like properties in the brain, but also modulate synaptic activity (Holscher, 2014). Synapses are the contacts between neurons, and they are important for memory formation and information processing in the brain. Direct injection of GLP-1 or long-lasting GLP-1 analogues into the brain markedly enhanced long-term potentiation of synaptic transmission (LTP) in the hippocampus, a brain area that is involved in memory formation. LTP is considered a cellular correlate of memory formation (Bliss and Collingridge, 1993). The GLP-1 analogue, liraglutide, has been shown to upregulate LTP in the rat brain (McClellan et al., 2010).

**[0010]** In addition, GLP-1 analogues were able to prevent the impairment of LTP that was induced by beta-amyloid fragments (Gault and Holscher, 2008a; McClellan et al., 2011; Gengler et al., 2012; Han et al., 2013). This impairment of LTP by amyloid protein may be the mechanism by which amyloid causes memory loss (Cleary et al., 2005). A study testing liraglutide in an APP/PS1 mouse model of AD showed that the drug can prevent the impairment in memory formation and synaptic plasticity, the reduction of total numbers of synapses, normalise stem cell proliferation and neurogenesis in the dentate gyrus, reduce the inflammation response, and furthermore reduce amyloid plaque load in the cortex and total amyloid levels in the brain (McClellan et al., 2011). In another study, liraglutide also had protective and regenerative effects in very old transgenic mice, demonstrating that even at an advanced stage of disease progression, memory can be improved and plaque load be reduced

to some degree (McClellan and Holscher, 2014).

**[0011]** Based on these findings in animal models, a pilot clinical trial in Alzheimer's patients has been conducted. The finding was that the GLP-1 analogue liraglutide protected the disease-induced degeneration of brain activity and energy utilization. This demonstrates that the drug enters the brain and protects neurons from the damaging effects of Alzheimer's while keeping neurons active and functional (Gejl et al., 2016). A phase II clinical trial of liraglutide in AD patients has started (NCT01843075).

**[0012]** Furthermore, one prior art study has investigated the effects of the GLP-1 receptor agonist exendin-4 in the 6-hydroxydopamine model of PD. After the lesion was induced, rats were treated with exendin-4 and a protection of motor activity was observed. Histological analysis showed that exendin-4 significantly increased the number of both tyrosine hydroxylase- and vesicular monoamine transporter 2-positive neurons in the substantia nigra (Bertilsson et al., 2008). In a second study, two rodent models of PD, 6-hydroxydopamine (6-OHDA) and lipopolysaccharide (LPS), were used to test the effects of exendin-4. Motor control was much improved in the drug group, and striatal tissue concentrations of dopamine were markedly higher. In addition, exendin-4 reversed the loss of extracellular DA in the striatum (Harkavyi et al., 2008). When testing the GLP-1 receptor agonists liraglutide and lixisenatide in the MPTP mouse model of Parkinson's disease, it was found that both drugs showed good effects in preventing the motor impairment, chronic inflammation in the brain, and loss of dopaminergic neurons induced by the toxin MPTP (Liu et al., 2015).

**[0013]** Based on these studies, a clinical trial of exendin-4 in PD patients had been initiated. This study reported that in several motor assessments and in a cognitive test, patients had improved, and the improvements were maintained even after the drug had been discontinued for 12 months (Aviles-Olmos et al., 2013; Aviles-Olmos et al., 2014). A phase II clinical trial in Parkinson's patients testing the drug liraglutide has started (NCT02953665).

**[0014]** Studies have also been carried out to determine whether GIP or GIP analogues have an effect in AD. It has been found that GIP analogues can prevent the LTP impairment that beta-amyloid fragments induce on synaptic transmission in the brain (Gault and Holscher, 2008b). In a GIP receptor- deletion mouse strain, LTP was also impaired, and paired-pulse facilitation was reduced, indicating that the release of synaptic vesicles is reduced (Faivre et al., 2011). The long-lasting GIP analogue D-Ala<sup>2</sup>-GIP also had neuroprotective effects in an APP/PS1 mouse model of AD. In 12 months old mice, synaptic plasticity in area CA1 of the hippocampus and spatial memory formation was impaired in APP/PS1 mice but was unimpaired in D-Ala<sup>2</sup>-GIP treated APP/PS1 mice. In addition, the amyloid plaque load was much reduced, showing impressive effects in reducing the main hallmarks of AD (Faivre and Holscher, 2013b).

**[0015]** In aged 19 month old AD mice, the drug was still able to reverse some of the AD symptoms such as synapse loss (Faivre and Holscher, 2013a). In a longitudinal study, oxidative stress and the inflammation response in the brain was much reduced in APP/PS1

mice (Duffy and Holscher, 2013). This suggests that these analogues have neuroprotective properties in AD and protect synapses from the detrimental effects of beta-amyloid.

**[0016]** In the MPTP mouse model of Parkinson's disease, the GIP analogue D-Ala<sup>2</sup>-GIP was protective and prevented the impairment of motor activity, the loss of dopaminergic neurons, loss of synapses and the chronic inflammation response induced by MPTP (Li et al., 2015). When testing a novel dual GLP-1/GIP dual agonist in this mouse model, it was found that it showed good protective effects (Ji et al., 2015). However, that particular molecule was not as effective as GLP-1 analogues.

**[0017]** There remains a need to identify treatments for neurological disorders such as for example the neurodegenerative diseases, Alzheimer's disease and Parkinson's disease.

**[0018]** It is an aim of certain embodiments of the present invention to at least partly mitigate the problems associated with the prior art.

**[0019]** It is an aim of certain embodiments of the present invention to provide a therapeutic peptide for use in the treatment and/or prevention of a neurodegenerative disorder such as for example Alzheimer's disease and/or Parkinson's disease.

**[0020]** It is an aim of certain embodiments of the present invention to provide novel GIP/GLP-1 dual agonist peptides which have superior efficacy as compared to the best in class GLP-1 mono agonist liraglutide, for use in the treatment and/or prevention of Alzheimer's or Parkinson's disease.

**[0021]** It is an aim of certain embodiments of the present invention to provide GIP/GLP-1 dual agonist peptides which have superior properties as compared to a GIP mono agonist. Examples of superior properties include for example a greater decrease in beta-amyloid plaque load and/or reduction of motor skill impairment.

**[0022]** It is an aim of certain embodiments of the present invention to provide novel GIP/GLP-1 dual agonist peptides for use in the treatment and/or prevention of a neurodegenerative disorder such as for example Alzheimer's disease and/or Parkinson's disease.

### Summary of the Invention

**[0023]** The invention is defined in the claims. In one aspect of the present disclosure there is provided a GIP/GLP-1 co-agonist peptide or a pharmaceutically acceptable salt or solvate thereof for use in the treatment and/or prophylaxis of a neurological disorder, wherein the co-agonist peptide is represented by the general formula 1:

Tyr-Aib-Glu-Gly-Thr-Phe-Thr-Ser-Asp-Tyr-Ser-Ile-Tyr-Leu-Asp-Lys-Gln-Ala-Ala-Aib-  
Glu-Phe-Val-Xaa<sup>24</sup>-Trp-Leu-Leu-Ala-Gly-Y1-Y2-R<sup>2</sup>

wherein Xaa<sup>24</sup> is selected from Asn and Cys;

Y1 is selected from absent or an extension comprising at least eight amino acid molecules;

Y2 is selected from:

Lys-Lys-Lys-Lys-Lys (SEQ. ID. No 2);

Lys-Lys-Lys-Lys-Lys-Lys (SEQ. ID. No. 3)

Arg-Arg-Gln-Arg-Arg-Lys-Lys-Arg-Gly-Tyr (SEQ. ID. No 4); and

Lys-Arg-Arg-Gln-Arg-Arg-Lys-Lys-Arg-Gly-Tyr (SEQ. ID. No 5); and

R<sup>2</sup> is selected from -NH<sub>2</sub> or -OH,

or a derivative or a pharmaceutically acceptable salt or solvate of the peptide or the derivative.

**[0024]** In certain aspects of the disclosure Y2 is Lys-Lys-Lys-Lys-Lys or Lys-Lys-Lys-Lys-Lys-Lys. In certain aspects of the disclosure

Y2 is Arg-Arg-Gln-Arg-Arg-Lys-Lys-Arg-Gly-Tyr or Lys-Arg-Arg-Gln-Arg-Arg-Lys-Lys-Arg-Gly-Tyr.

**[0025]** In certain embodiments, of the disclosure

Xaa<sup>24</sup> is Cys. In certain embodiments, Xaa<sup>24</sup> is Asn.

**[0026]** In certain embodiments, of the disclosure

Y1 is an extension comprising at least 10 amino acids e.g. at least 11 amino acids.

**[0027]** In certain embodiments, of the disclosure Y1 is selected from:

Gly-Pro-Ser-Ser-Gly-Ala-Pro-Pro-Pro-Ser (SEQ. ID. No. 6);

Gly-Pro-Ser-Ser-Gly-Ala-Pro-Pro-Pro-Ser-Lys (SEQ. ID No. 7);

Gly-Pro-Ser-Ser-Gly-Ala-Pro-Pro-Pro-Ser-Cys (SEQ. ID. No. 8);

Pro-Ser-Ser-Gly-Ala-Pro-Pro-Pro-Ser (SEQ. ID. No. 9);

Pro-Ser-Ser-Gly-Ala-Pro-Pro-Ser (SEQ. ID. No. 10); and

absent.

**[0028]** Aptly, Y1 is Gly-Pro-Ser-Ser-Gly-Ala-Pro-Pro-Pro-Ser (SEQ. ID. No. 6). In the disclosure

Y1 is Gly-Pro-Ser-Ser-Gly-Ala-Pro-Pro-Pro-Ser-Cys (SEQ. ID. No 8). In an alternative

embodiment, Y1 is Gly-Pro-Ser-Ser-Gly-Ala-Pro-Pro-Pro-Ser-Lys (SEQ. ID. No. 7).

**[0029]** In certain embodiments, the peptide comprises a hydrophilic moiety covalently linked to an amino acid. In certain embodiments, the peptide comprises a hydrophilic moiety covalently linked to an amino acid at position 24.

**[0030]** In certain embodiments, the peptide comprises a hydrophilic moiety covalently linked to an amino acid at position 39 or 40, when Y1 is an extension comprising at least 10 amino acids or at least 11 amino acids.

**[0031]** In certain embodiments, Xaa<sup>39</sup> or Xaa<sup>40</sup> is Cys and the peptide comprises a hydrophilic moiety covalently linked to Cys(39) or Cys(40).

**[0032]** In certain embodiments, Xaa<sup>24</sup> is Cys and the peptide comprises a hydrophilic moiety covalently linked to Cys(24).

**[0033]** In certain embodiments, the hydrophilic moiety is a water-soluble polymer.

**[0034]** In certain embodiments, the water-soluble polymer is a polyethylene glycol moiety and optionally is a polyethylene glycol moiety having a molecular weight of between about 20,000 Daltons and about 60,000 Daltons. In certain embodiments, the peptide is conjugated with a lipophilic substituent.

**[0035]** In certain embodiments, the lipophilic substituent comprises a hydrocarbon chain having 8 to 24 carbon (C) atoms. In certain embodiments, the lipophilic substituent comprises an acyl group.

**[0036]** In certain embodiments, the lipophilic substituent is a fatty acid molecule. Aptly, the fatty acid molecule is selected from a C-8 octanoyl group, a C-10 decanoyl group, a C-12 lauroyl group, a C-14 myristoyl group, a C-16 palmitoyl group, a C-18 stearoyl group and a C-20 acyl group. In certain embodiments, the lipophilic substituent is attached to an amino acid at the carboxyl-terminus of the peptide. In certain embodiments, the lipophilic substituent is attached to an amino acid at position 40. Aptly, the lipophilic substituent is covalently bonded to a side chain of an amino acid of the peptide.

**[0037]** In certain embodiments, the peptide comprises a spacer which conjugates the lipophilic substituent to an amino acid residue. Aptly, the spacer is a residue from a naturally occurring or unnatural amino acid, wherein the spacer comprises a residue of Gly, Pro, Ala, Val, Leu, Ile, Met, Cys, Phe, Tyr, Trp, His, Lys, Arg, Gln, Asn,  $\alpha$ -Glu,  $\gamma$ -Glu,  $\epsilon$ -Lys, Asp, Ser, Thr, Gaba, Aib,  $\beta$ -aminohexonyl, 7-aminoheptanoyl, 8-aminooctanoyl, 9-aminononanoyl, 10-aminodecanoyl or 8-amino-3,6- dioxaoctanoyl.

**[0038]** In certain embodiments, the spacer is  $\gamma$ -Glu. In certain embodiments, the spacer is a dipeptide, optionally wherein the spacer comprises two negatively charged amino acids and



further optionally wherein the spacer is  $\gamma$ -Glu-  $\gamma$ -Glu.

**[0039]** In certain embodiments, the peptide comprises the amino acid sequence of:  
YXEGTFTSDYSIYLDKQAAXEFVNWLLAGGPSSGAPPPSKKKKKK-NH<sub>2</sub> (SEQ. ID. No. 11)  
wherein X = aminoisobutyric acid.

**[0040]** In certain embodiments, the peptide consists essentially of the amino acid sequence of:  
YXEGTFTSDYSIYLDKQAAXEFVNWLLAGGPSSGAPPPSKKKKKK-NH<sub>2</sub> (SEQ. ID. No. 11)  
wherein X = aminoisobutyric acid.

**[0041]** In certain embodiments, the peptide comprises the amino acid sequence of:  
YXEGTFTSDYSIYLDKQAAXEFVNWLLAGGPSSGAPPPSKRRQRRKKRGY-NH<sub>2</sub> (SEQ. ID. No. 12)  
wherein X = aminoisobutyric acid.

**[0042]** Aptly, the peptide consists essentially of the amino acid sequence of:  
YXEGTFTSDYSIYLDKQAAXEFVNWLLAGGPSSGAPPPSKRRQRRKKRGY-NH<sub>2</sub> (SEQ. ID. No. 12)  
wherein X = aminoisobutyric acid.

**[0043]** In a further aspect of the present invention, there is provided a peptide as described herein for use in the treatment and/or prophylaxis of a neurological disorder.

**[0044]** In a further aspect of the present invention, there is provided a peptide as described herein for use in the treatment of a neurological disorder which is caused by or associated with beta-amyloid protein plaque deposition in an area of the patient. In certain embodiments, the beta-amyloid plaque deposition is in the brain of the patient.

**[0045]** In a further aspect of the present invention, there is provided a peptide as described herein for use in the treatment and/or prophylaxis of a neurological disorder caused by, or associated with, dysfunction of long-term potentiation of synaptic transmission.

**[0046]** In a further aspect of the present invention, there is provided a peptide as described herein is for use in the treatment and/or prophylaxis of a neurological disorder caused by, or associated with, inflammation.

**[0047]** In a further aspect of the present invention, there is provided a peptide as described herein for use in the treatment of a neurological disorder associated with motor impairment.

**[0048]** In a further aspect of the present invention, there is provided a peptide as described herein for use in the treatment and/or prophylaxis of a neurological disorder affecting cognitive function, e.g. dementia, stroke, schizophrenia and/or bipolar disorder.

**[0049]** In certain embodiments, the disorder is cerebral ischemia associated with stroke.

**[0050]** In certain embodiments, the peptide is for use in the treatment and/or prophylaxis of a disorder selected from post-traumatic stress disorder, epilepsy, Tourette's syndrome, and hallucinations; and dysfunctional cognitive processes, optionally selected from attention, calculation, memory, judgment, insight, learning and reasoning.

**[0051]** In certain embodiments, the peptide is for use in the treatment and/or prophylaxis of a neurodegenerative disorder e.g. Alzheimer's disease, Parkinson's disease, Amyotrophic Lateral Sclerosis, peripheral neuropathy, Huntington's disease and Creutzfeldt-Jacob disease. Aptly, the neurological disorder is multiple sclerosis.

**[0052]** In certain embodiments, the peptide is for use in the treatment and/or prophylaxis of a neurological disorder selected from clinical or pre-clinical Alzheimer's disease, prodromal Alzheimer's disease, and clinical or preclinical amyloid angiopathy (CAA).

**[0053]** Aptly, the peptide is for use in the treatment and/or prophylaxis of clinical Alzheimer's disease. In a further aspect of the present invention, there is provided a peptide as described herein for use in the treatment and/or prophylaxis of Parkinson's disease.

**[0054]** Also provided is a method of treating and/or preventing a neurological disorder as described herein, the method comprising administering a pharmaceutically effective amount of a GIP/GLP-1 co-agonist peptide as described herein to a subject in need thereof.

**[0055]** In a further aspect of the present disclosure there is provided a method of treating and/or preventing a neurological disorder comprising administering to a patient in need thereof a pharmaceutical composition comprising a GIP/ GLP-1 co-agonist peptide, or a derivative or a pharmaceutically acceptable salt or solvate of the peptide or the derivative, wherein said co-agonist peptide is represented by the general Formula I:  
Tyr-Aib-Glu-Gly-Thr-Phe-Thr-Ser-Asp-Tyr-Ser-Ile-Tyr-Leu-Asp-Lys-Gln-Ala-Ala-Aib-Glu-Phe-Val-Xaa<sup>24</sup>-Trp-Leu-Leu-Ala-Gly-Y1-Y2-R<sup>2</sup>

(I) (SEQ. ID. No. 1)

wherein Xaa<sup>24</sup> is selected from Asn and Cys;

Y1 is selected from absent or an extension comprising at least eight amino acid molecules;

Y2 is selected from:

Lys-Lys-Lys-Lys-Lys (SEQ. ID. No 2);

Lys-Lys-Lys-Lys-Lys-Lys (SEQ. ID. No. 3)

Arg-Arg-Gln-Arg-Arg-Lys-Lys-Arg-Gly-Tyr (SEQ. ID. No 4); and

Lys-Arg-Arg-Gln-Arg-Arg-Lys-Lys-Arg-Gly-Tyr (SEQ. ID. No 5); and

R<sup>2</sup> is selected from -NH<sub>2</sub> or -OH,

or a derivative or a pharmaceutically acceptable salt or solvate of the peptide or the derivative.

**[0056]** Aptly, the peptide is as described herein.

**[0057]** In certain embodiments, the neurological disorder is selected from post-traumatic stress disorder, epilepsy, Tourette's syndrome, and hallucinations; and dysfunctional cognitive processes, optionally selected from attention, calculation, memory, judgment, insight, learning and reasoning.

**[0058]** In certain embodiments, the neurological disorder is a neurodegenerative disorder. Aptly, the neurodegenerative disorder is selected from Alzheimer's Disease, Parkinson's Disease, Huntington's disease, Amyotrophic Lateral Sclerosis, peripheral neuropathy, and Creutzfeldt-Jacob disease.

**[0059]** In certain embodiments, the neurological disorder is a neurological disorder selected from clinical or pre-clinical Alzheimer's disease, prodromal Alzheimer's disease, and clinical or preclinical amyloid angiopathy (CAA). APTLY, the neurological disorder is clinical Alzheimer's Disease.

**[0060]** In a further aspect of the present invention, there is provided a peptide for use in the treatment and/ or prophylaxis of clinical Alzheimer's Disease, the peptide consisting of the following sequence:

YXEGTFTSDYSIYLDKQAAXEFVNWLLAGGPSSGAPPPSKKKKKK-NH<sub>2</sub> (SEQ. ID. No. 11)

wherein X = aminoisobutyric acid.

**[0061]** In a further aspect of the present invention, there is provided a peptide for use in the treatment and/ or prophylaxis of clinical Alzheimer's Disease, the peptide consisting of the following sequence:

YXEGTFTSDYSIYLDKQAAXEFVNWLLAGGPSSGAPPPSKRRQRRKKRGY-NH<sub>2</sub> (SEQ. ID. No. 12)

wherein X = aminoisobutyric acid.

**[0062]** In one embodiment, the peptide comprises or consists of the following amino acid sequence:

YXEGTFTSDYSIYLDKQAAXEFVNWLLAGGPSSGAPPPSKKKKKK-NH<sub>2</sub> (SEQ. ID. No. 11) X = aminoisobutyric acid

**[0063]** This peptide is referred to herein as "DA4" and has an amino acid sequence as set forth in SEQ. ID. No. 11.

**[0064]** In one embodiment, the peptide comprises or consists of the following amino acid sequence:

YXEGTFTSDYSIYLDKQAAXEFVNWLLAGGPSSGAPPPSKRRQRRKKRGY-NH<sub>2</sub> (SEQ. ID. No.

12)

X = aminoisobutyric acid

**[0065]** This peptide is referred to herein as "DA5" and has an amino acid sequence as set forth in SEQ. ID. No. 12.

**[0066]** The ability to cross the blood-brain barrier may have utility in the treatment of neurodegenerative disorders. Without being bound by theory, it is considered that the peptides of certain embodiments of the present invention are able to cross the blood-brain barrier. Furthermore, the peptides of certain embodiments cross the blood-brain barrier at an enhanced rate as compared to known GIP/GLP-1 co-agonist peptides.

**[0067]** It is noted that there can be many difficulties in transmission of a peptide across the blood brain barrier. A number of different mechanisms for enhancing access across the blood-brain barrier have been considered, one of which is the modification of peptide sequences. However, whilst a number of peptide modifications are known to potentially assist in crossing the blood-brain barrier, their *in vivo* behaviour is difficult to predict. Their addition to a peptide or protein molecule may be ineffective or lead to inactivation of the peptide or protein e.g. by protein degradation processes, incorrect peptide folding patterns and/or a change in receptor binding properties of the peptide.

**[0068]** Aptly, the peptide is for use in the treatment and/or prevention of Alzheimer's disease or Parkinson's disease.

**[0069]** Also provided herein is a pharmaceutical composition which comprises GIP/GLP-1 receptor co-agonist peptides as described herein and a pharmaceutically acceptable carrier for use in the treatment and/or prophylaxis of a neurological disorder as described herein. Further provided in the present disclosure is a kit including such a pharmaceutical composition.

### **Brief Description of Drawings**

**[0070]** Further details of embodiments of the invention, with reference to the accompanying drawings by way of example only, are provided below:

As used herein, the abbreviation "DA4" refers to a GIP/GLP-1 co-agonist consisting of the amino acid sequence shown in SEQ ID. No. 11.

As used herein, the abbreviation "DA5" refers to a GIP/GLP-1 co-agonist consisting of the amino acid sequence shown in SEQ ID. No. 12.

**Figure 1** illustrates an immunohistochemical measurement of beta-amyloid plaque load in the cortex of transgenic mice (Alzheimer's disease model). All peptides reduced the plaque load. Both co-agonist peptides (DA4 and DA5) of embodiments of the present invention were superior to the single GIP or GLP-1 analogues and administration of DA4 and DA5 reduces

beta-amyloid plaque load as quantified by beta amyloid immunohistochemistry and determination of the % area positive for beta amyloid in cross sections of the brain cortex. \*=  $p < 0.05$ , \*\*= $p < 0.01$ ; \*\*\*= $p < 0.001$ . N=6 per group.

**Figure 2** is a graph illustrating that the DA4 and DA5 peptides protected from synapse loss in the cortex of transgenic mice (Alzheimer's disease model) as quantified by immunohistochemical measurement of synaptic densities in the cortex of the transgenic mice. All peptides protected from synapse loss. Both co-agonist peptides of embodiments of the present invention were superior to the single GIP or GLP-1 analogues. \*=  $p < 0.05$ , \*\*= $p < 0.01$ ; \*\*\*= $p < 0.001$ . N=6 per group.

**Figure 3** is a graph illustrating that the DA4 and DA5 peptides protect mice from an impairment of motor skills induced by MPTP to reduce the levels of dopamine. A rotarod motor skill test was carried out as described below. Each animal's endurance time was recorded, and the average was calculated. Data were analysed using a one-way ANOVA with post-hoc Bonferroni tests. Both co-agonist peptides of embodiments of the present invention were superior to the single GIP or GLP-1 analogues. Control = wild type mouse without MPTP. \*=  $p < 0.05$ , \*\*= $p < 0.01$ ; \*\*\*= $p < 0.001$ . N=6 per group.

**Figure 4** illustrates MPTP-induced reduction of the enzyme TH that synthesises dopamine in the substantia nigra, pars compacta. Both DA4 and DA5 protected dopaminergic neurons at a greater rate than the single GIP and GLP-1 analogues. \*=  $p < 0.05$ , \*\*= $p < 0.01$ ; \*\*\*= $p < 0.001$ . N=6 per group.

**Figure 5** illustrates that the DA5 peptide does not affect the locomotor and exploratory performance of APP/PS1 and Wild Type mice. An open field test was carried out to measure the spontaneous behaviour of saline/DA5 treated mice. Statistical analysis ( $p > 0.05$ ) did not show any difference between saline/DA5 groups (a,b).

**Figure 6** illustrates improved spatial working memory of APP/PS1 mice in a Y-maze test upon DA5 peptide treatment. The percentage of right alternation in the mice of APP/PS1 mice was significantly reduced. \*=  $p < 0.05$ , \*\*= $p < 0.01$ ; \*\*\*= $p < 0.001$  N=8-14 per group.

**Figure 7** illustrates the disorder in spatial memory observed in APP/PS1 mice can be ameliorated with DA5 peptide treatment. A classical Morris water maze was carried out and the spatial memory of mice was assessed by a probe trial, as described below. (Figure 7a,7b) Each animal's escape latency was recorded over a training period and the average was calculated. (Figure 7c, 7d). Probe test indicated the DA5 treated APP/PS1 mice recalled the escape route to a greater extent than APP/PS1-saline mice. \*=  $p < 0.05$ , \*\*= $p < 0.01$ ; \*\*\*= $p < 0.001$ . N=8-14 per group.

**Figure 8** illustrates that DA5 peptide protects APP/PS1 mice from decreased cognitive flexibility and restores their ability to relearn. A reverse Morris water maze test was carried out as described below. (Figure 8a, Figure 8b). Each animal's escape latency was recorded over an extended training period and the average was calculated (Figure 8c, 8d). Probe test indicated the DA5 treated APP/PS1 mice recalled a change in escape route to a greater extent than

APP/PS1-saline mice. \*= p<0.05, \*\*=p<0.01; \*\*\*=p<0.001. N=8-14 per group.

**Figure 9** illustrates that the DA5 peptide maintains synaptic strength in APP/PS1 mice (Figure 9a) Pre-treatment of mice with DA5 was able to reverse the depression of L-LTP observed in APP/PS1 transgenic mice (Figure 9b) Furthermore, DA5 treated APP/PS1 mice maintained high levels of L-LTP post neuronal stimuli. \*= p<0.05, \*\*=p<0.01; \*\*\*=p<0,001. N=6 per group.

**Figure 10** is a graph illustrating DA5 reduces neurofibrillary tangles in APP/PS1 mice. Quantification of immunofluorescence staining of p-tau protein indicated reduced neurofibrillary tangles in APP/PS1 DA5 treated mice compared to APP/PS1-saline. \*= p<0.05, \*\*=p<0.01; \*\*\*=p<0.001. N=4 per group.

**Figure 11** shows a western blot and graphs illustrating DA5 can protect cognitive behaviours and improve pathological changes in AD mice. Treatment of APP/PS1 mice with DA5 prevents excessive activation of GSK $\beta$  by upregulating P13/AKT signalling (a-g). \*= p<0.05, N=6 per group.

**Figure 12** details amino acid sequences of peptides as described herein.

### **Detailed Description of Embodiments of the Invention**

**[0071]** The practice of embodiments of the present invention will employ, unless otherwise indicated, conventional techniques of molecular biology, microbiology, recombinant DNA technology and immunology, which are within the skill of those working in the art.

**[0072]** Most general molecular biology, microbiology recombinant DNA technology and immunological techniques can be found in Sambrook et al, Molecular Cloning, A Laboratory Manual (2001) Cold Harbor-Laboratory Press, Cold Spring Harbor, N.Y. or Ausubel et al., Current protocols in molecular biology (1990) John Wiley and Sons, N.Y. Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this disclosure belongs. For example, the Concise Dictionary of Biomedicine and Molecular Biology, Juo, Pei-Show, 2nd ed., 2002, CRC Press; The Dictionary of Cell and Molecular Biology, 3rd ed., Academic Press; and the Oxford University Press, provide a person skilled in the art with a general dictionary of many of the terms used in this disclosure.

**[0073]** The present disclosure relates to the use of GIP/GLP-1 co-agonist peptides. The native human GLP-1 peptide and GIP peptide sequences are known in the art. The term "GLP-1", or "hGLP-1" as used herein refers to the human Glucagon-Like Peptide-1 (GLP-1 (7-37)), the sequence of which is included herein as SEQ ID. No. 13.

**[0074]** The peptide having the sequence of SEQ ID. No 13 may also be designated "native"

GLP-1.

**[0075]** The Homo sapiens GLP-1(7-37) sequence is:

HAEGTFTSDV SSYLEGQAAK EFWLWKGR G-OH (SEQ ID No. 13).

**[0076]** The term "GIP ", or "hGIP" as used herein refers to the human Gastric Inhibitory Peptide (also known as glucose-dependent insulinotropic peptide) the sequence of which is included herein as SEQ. ID. No. 14. The peptide having the sequence of SEQ. ID. No. 14 may also be designated "native" GIP:

YAEGTFISDYSIAMDKIHQQDFVNWLLAQKGKNDWKHNITQ-OH (SEQ ID. No 14).

**[0077]** As used herein, a general reference to "GIP" or "GLP-1" in the absence of any further designation is intended to mean native GIP or native GLP-1, respectively.

**[0078]** As used herein, the term "peptide" encompasses a sequence of 3 or more amino acids and typically less than 50 amino acids, wherein the amino acids are naturally occurring or non-naturally occurring amino acids. Non-naturally occurring amino acids refer to amino acids that do not naturally occur in vivo but which, nevertheless, can be incorporated into the peptide structures described herein.

**[0079]** As used herein, the terms "polypeptide" and "protein" are terms that are used interchangeably to refer to a polymer of amino acids, without regard to the length of the polymer. Typically, polypeptides and proteins have a polymer length that is greater than that of "peptides".

**[0080]** As used herein an amino acid "modification" refers to a substitution, addition or deletion of an amino acid, and includes substitution with or addition of any of the 20 amino acids commonly found in human proteins, as well as atypical or non-naturally occurring amino acids. Throughout the application, all references to a particular amino acid position by number (e.g. position 28) refer to the amino acid at that position in the GIP/GLP-1 dual agonist of embodiments of the present invention.

**[0081]** Throughout this specification, the conventional one letter and three letter codes for naturally occurring amino acids are used, as well as generally accepted three letter codes for other amino acids, such as for example Aib ( $\alpha$ -aminoisobutyric acid).

**[0082]** Thus, in an aspect of the present invention, there is provided a GIP/GLP-1 co-agonist peptide, or a derivative or a pharmaceutically acceptable salt or solvate of the peptide or the derivative, for use in the treatment and/or prophylaxis of a neurological disorder. Aply, the co-agonist peptide has a GLP-1 percentage potency within about 10-fold of the GIP percentage potency.

**[0083]** As used herein, the term "agonist" refers to a substance (ligand) that activates the receptor type in question. The terms "dual agonist" and "co-agonist" are used herein are

interchangeable and refer to a substance (ligand) that activate two receptor types. Aptly, the GIP/GLP-1 co-agonist peptides described herein have balanced activity at both the GLP-1R and the GIPR. In one embodiment, the co-agonist peptide has an EC<sub>50</sub> at the human GLP-1 receptor within about 10-fold of the EC<sub>50</sub> at the human GIP receptor. Activity in *in vitro* assays may be used as a measure of the peptides' activity.

**[0084]** Aptly, the peptides described herein are synthesized by means of solid-phase or liquid-phase peptide synthesis, as described in for example WO98/11125. Aptly, the peptides may be synthesized as described in Finan et al. (Sci. Transl. Med. 5 209ra151 (2013) pp 1 to 16).

**[0085]** The peptides of the present disclosure may be formulated as pharmaceutical compositions prepared for storage or administration for use in the treatment and/or prevention of a neurological disorder as described herein. Such a composition typically comprises a therapeutically effective amount of a GIP/GLP-1 co-agonist peptide, in the appropriate form, in a pharmaceutically acceptable carrier.

**[0086]** The therapeutically effective amount of a GIP/GLP-1 co-agonist peptide as described herein will depend on the route of administration, the type of mammal being treated, and the physical characteristics of the specific mammal under consideration. These factors and their relationship to determining this amount are well known to skilled practitioners in the medical arts. This amount and the method of administration can be tailored to achieve optimal efficacy, and may depend on such factors as weight, diet, concurrent medication and other factors, well known to those skilled in the medical arts. The peptides of the present disclosure may be particularly useful for treatment of humans.

**[0087]** An effective dosage and treatment protocol may be determined by conventional means, starting with a low dose in laboratory animals and then increasing the dosage while monitoring the effects, and systematically varying the dosage regimen as well. Numerous factors may be taken into consideration by a clinician when determining an optimal dosage for a given subject. Such considerations are known to the person skilled in the art.

**[0088]** The term "pharmaceutically acceptable carrier" includes any of the standard pharmaceutical carriers. Pharmaceutically acceptable carriers for therapeutic use are well known in the pharmaceutical art, and are described, for example, in Remington's Pharmaceutical Sciences, Mack Publishing Co. (A.R.Gennaro edit. 1985). For example, sterile saline and phosphate-buffered saline at slightly acidic or physiological pH may be used. pH buffering agents may be phosphate, citrate, acetate, tris/hydroxymethyl)aminomethane (TRIS), N-Tris(hydroxymethyl)methyl-3-aminopropanesulphonic acid (TAPS), ammonium bicarbonate, diethanolamine, histidine, arginine, lysine, or acetate or mixtures thereof. The term further encompasses any agents listed in the US Pharmacopeia for use in animals, including humans.

**[0089]** The term "pharmaceutically acceptable salt" refers to a salt of any one of the GIP/GLP-1 co-agonist peptide of embodiments of the invention. Salts include pharmaceutically acceptable salts such as acid addition salts and basic salts. Examples of acid addition salts



include hydrochloride salts, citrate salts and acetate salts. Examples of basis salts include salts where the cation is selected from alkali metals, such as sodium and potassium, alkaline earth metals, such as calcium, and ammonium ions  $^+N(R^3)_3(R^4)$ , where  $R^3$  and  $R^4$  independently designates optionally substituted  $C_{1-6}$ -alkyl, optionally substituted  $C_{2-6}$ -alkenyl, optionally substituted aryl, or optionally substituted heteroaryl. Other examples of pharmaceutically acceptable salts are described in "Remington's Pharmaceutical Sciences", 17th edition. Ed. Alfonso R. Gennaro (Ed.), Mark Publishing Company, Easton, PA, U.S.A., 1985 and more recent editions, and in the Encyclopaedia of Pharmaceutical Technology.

**[0090]** The term "solvate" in the context of the present disclosure refers to a complex of defined stoichiometry formed between a solute (e.g., a peptide or pharmaceutically acceptable salt thereof according to the present disclosure) and a solvent. The solvent in this connection may, for example, be water, ethanol or another pharmaceutically acceptable, typically small-molecular organic species, such as, but not limited to, acetic acid or lactic acid. When the solvent in question is water, such a solvate is normally referred to as a hydrate.

**[0091]** "Treatment" is an approach for obtaining beneficial or desired clinical results. For the purposes of the present disclosure, beneficial or desired clinical results include, but are not limited to, alleviation of symptoms, diminishment of extent of disease, stabilized (i.e., not worsening) state of disease, delay or slowing of disease progression, amelioration or palliation of the disease state, and remission (whether partial or total), whether detectable or undetectable. "Treatment" can also mean prolonging survival as compared to expected survival if not receiving treatment. "Treatment" is an intervention performed with the intention of preventing the development or altering the pathology of a disorder. Accordingly, "treatment" refers to both therapeutic treatment and prophylactic or preventative measures in certain embodiments. Those in need of treatment include those already with the disorder as well as those in which the disorder is to be prevented. By treatment is meant inhibiting or reducing an increase in pathology or symptoms when compared to the absence of treatment, and is not necessarily meant to imply complete cessation of the relevant condition.

**[0092]** The pharmaceutical compositions for use in the treatment of a neurological disorder can be in unit dosage form. In such form, the composition is divided into unit doses containing appropriate quantities of the active component. The unit dosage form can be a packaged preparation, the package containing discrete quantities of the preparations, for example, packeted tablets, capsules, and powders in vials or ampoules. The unit dosage form can also be a capsule, cachet, or tablet itself, or it can be the appropriate number of any of these packaged forms. It may be provided in single dose injectable form, for example in the form of a pen. In certain embodiments, packaged forms include a label or insert with instructions for use. Compositions may be formulated for any suitable route and means of administration. Pharmaceutically acceptable carriers or diluents include those used in formulations suitable for oral, rectal, nasal, topical (including buccal and sublingual), vaginal or parenteral (including subcutaneous, intramuscular, intravenous, intradermal, and transdermal) administration. The formulations may conveniently be presented in unit dosage form and may be prepared by any of the methods well known in the art of pharmacy.

**[0093]** As used herein an "effective" amount or a "therapeutically effective amount" of a peptide refers to a nontoxic but sufficient amount of the peptide to provide the desired effect. The amount that is "effective" will vary from subject to subject, depending on the age and general condition of the individual, mode of administration, and the like. An appropriate "effective" amount in any individual case may be determined by one of ordinary skill in the art using routine experimentation.

**[0094]** The terms "patient", "subject" and "individual" may be used interchangeably and refer to either a human or non-human mammal. Aptly, the subject is a human.

**[0095]** The peptides described herein may be used to treat and/or prevent a neurological disorder e.g. a neurodegenerative disorder. In one embodiment, the peptide is for use in the treatment and/or prevention of Alzheimer's disease. Alzheimer's disease (AD) is a neurodegenerative disorder that results in the loss of cortical neurons, especially in the associative neocortex and hippocampus which in turn leads to slow and progressive loss of cognitive functions, ultimately leading to dementia and death. Major hallmarks of the disease are aggregation and deposition of misfolded proteins such as aggregated beta-amyloid peptide as extracellular senile or neuritic 'plaques', and hyperphosphorylated tau protein as intracellular neurofibrillary 'tangles' (NFTs).

**[0096]** Genetically, AD is divided into two forms: (1) early-onset familial AD (<60 years), and (2) late-onset sporadic AD (>60 years). Rare, disease causing mutations in Amyloid precursor protein (APP), Presenilin 1 (PSEN1), and Presenilin 2 (PSEN2) genes are known to result in early-onset familial AD while, APOE (allele 4) is the single most important risk factor for late-onset AD.

**[0097]** Although Alzheimer's disease develops differently for every individual, there are many common symptoms. Early symptoms are often mistakenly thought to be 'age-related' concerns, or manifestations of stress. In the early stages, the most common symptom is difficulty in remembering recent events. When AD is suspected, the diagnosis is usually confirmed with tests that evaluate behaviour and thinking abilities, often followed by a brain scan if available, however, examination of brain tissue is required for a definitive diagnosis.

**[0098]** As the disease advances, symptoms can include confusion, irritability, aggression, mood swings, trouble with language, and long-term memory loss. As the sufferer declines they often withdraw from family and society. Gradually, bodily functions are lost, ultimately leading to death. Since the disease is different for each individual, predicting how it will affect the person is difficult. AD develops for an unknown and variable amount of time before becoming fully apparent, and it can progress undiagnosed for years. On average, the life expectancy following diagnosis is approximately seven years. Fewer than three percent of individuals live more than fourteen years after diagnosis.

**[0099]** In one embodiment, the co-agonist peptide is for use in treating Alzheimer's disease.

The peptide may be used to slow down and/or halt the progression of Alzheimer's disease in a subject. The peptide may be for use to slow down and/or prevent the progression to clinical Alzheimer's disease of a subject suffering from pre-clinical Alzheimer's Disease.

**[0100]** In one embodiment, the co-agonist peptide is for use in the treatment and/or prevention of Parkinson's disease. Parkinson's disease is a neurodegenerative disorder characterized by progressive neuronal loss, in particular of dopaminergic neurons of the substantia nigra, leading to motor disturbances such as tremor, rigidity, slowness of movement, and postural instability. In one embodiment, the peptide is for use to reduce and/or prevent motor disturbances associated with Parkinson's disease.

**[0101]** In addition, there is an increasing number of atypical Parkinson syndromes and subclasses of Parkinson's disease, which are associated with learning and memory deficits. Typical examples are Parkinson's Disease Dementia (PDD), Lewy Body Dementia (LBD) and Multi-Systems Atrophy (MSA). The peptide may be for use in the retrieval or improvement of learning processes or learning deficits and/or the prevention, retrieval or improvement of memory loss or memory impairment in a subject suffering from Parkinson's disease.

**[0102]** Throughout the description and claims of this specification, the words "comprise" and "contain" and variations of them mean "including but not limited to" and they are not intended to (and do not) exclude other moieties, additives, components, integers or steps. Throughout the description and claims of this specification, the singular encompasses the plural unless the context otherwise requires. In particular, where the indefinite article is used, the specification is to be understood as contemplating plurality as well as singularity, unless the context requires otherwise.

**[0103]** The reader's attention is directed to all papers and documents which are filed concurrently with or previous to this specification in connection with this application and which are open to public inspection with this specification,

**[0104]** Units, prefixes and symbols are denoted in their Système international d'unités (SI) accepted form. Numeric ranges are inclusive of the numbers defining the range. Unless other indicated, amino acid sequences are written left to right in amino to carboxy orientation. All amino acid residues in peptides of embodiments of the invention are preferably of the L-configuration. However, D-configuration amino acids may also be present.

## **Examples**

## **Molecular structures**

## **Peptide Synthesis**

[0105] Peptides were synthesized by GL Biochem Ltd. (Shanghai). The purity of the peptide was analysed by reversed-phase HPLC and characterised using matrix assisted laser desorption/ionisation time of flight (MALDI-TOF) mass spectrometry, with a purity > 99%. Peptides were reconstituted in ultrapure® water to a concentration of 1 mg/ml in polypropylene tubes and frozen in aliquots to permit fresh preparation of doses required for injection.

[0106] The peptides tested were:

1. 1. Liraglutide (GLP-1 analogue) -

Liraglutide is a peptide drug which is typically administered parenterally. Liraglutide is authorised for the treatment of Type 2 diabetes. Liraglutide is disclosed in WO98/08871 A1, Example 37. The amino acid sequence of liraglutide is shown in SEQ ID. No. 15. The fatty acid is linked to amino acid Lys at position 26. The amino acid Lys at position 34 has been replaced by amino acid Arg.

2. 2. DAAla<sup>2</sup>GIP-Lys<sup>37</sup>-γ-Glu-PAL (GIP analogue)

The sequence of DAAla<sup>2</sup>GIP-Lys<sup>37</sup>-γ-Glu-PAL is:

**Yd-AEGTFISDYSIAMDKIHQQDFVNWLLAQKGKK(γ-E-C16)NDWKHNITQ**

Liraglutide is a GLP-1 receptor mono agonist, whilst DAAla<sup>2</sup>GIP--LyS<sup>37</sup>- γ-Glu-PAL is a GIP receptor mono agonist.

3. 3. A peptide referred to herein as DA4 which has the following amino acid sequence:

**YXEGTFTSDYSIYLDKQAAXEFVNWLLAGGPSSGAPPPSKKKKKK-NH<sub>2</sub>**

X = amino-isobutyric acid

The DA4 peptide is a GLP-1/GIP dual agonist.

4. 4. A peptide referred to herein as DA5 which has the following amino acid sequence:

**YXEGTFTSDYSIYLDKQAAXEFVNWLLAGGPSSGAPPPSKRRQRKRGY-NH<sub>2</sub>**

X = amino-isobutyric acid

The DA5 peptide is a GLP-1/GIP dual agonist.

### Example 1

#### Transgenic Alzheimer's disease mouse model

[0107] Animal models have been developed to test novel drugs. A standard model for Alzheimer's disease is transgenic mice that express human mutated genes for amyloid production that are known to induce Alzheimer's disease in humans. These APP/PS1 transgenic mice recapitulate some of the symptoms of Alzheimer's disease such as the aggregation of amyloid in the brain, memory loss and the loss of synapses (Radde et al., 2006).

**[0108]** The APP/PS1 mice express human Swedish mutated APP and human mutated presenilin-1 which induce Alzheimer's Disease in humans.

**[0109]** The dose of peptide tested was 10nmol/kg body weight. The peptides were administered by a once-daily intraperitoneal injection for 8 weeks. Saline injections were administered as a control. Six animals per group were injected with a peptide or a saline control.

#### **Brain tissue analysis for amyloid plaques and synapse numbers in AD mice**

**[0110]** Animals (6 per group) were perfused transcardially with 30 ml of ice-cold PBS and 30 ml of ice-cold 4% paraformaldehyde to postfix the brain. The brains were removed and placed in fresh 30% sucrose solution in PBS to cryoprotect tissue and cut at a thickness of 40 µm on a cryostat. Sections were chosen according to stereological rules, with the first section taken at random and every 5th section afterwards. Between 7 and 13 sections were analysed per brain.

**[0111]** Immunostaining techniques were used to assess the neuronal plaque load (anti beta-amyloid rabbit polyclonal antibody (1:200, rabbit polyclonal-Invitrogen 71-5800) and of synaptophysin (polyclonal rabbit anti-synaptophysin primary antibody, 1:2000, Abcam, Cambridge, UK) to measure the amount of synapses in the cortex. Brain samples were first exposed to 99% formic acid for 7 minutes and then washed 3 times for 10 minutes in TBS. Pre-treatment with 99% formic acid is known to drastically increase the detection of beta-Amyloid in formalin-fixed brain samples. Then samples were incubated in 0.3% H<sub>2</sub>O<sub>2</sub> (Sigma Aldrich; Cat.-No.: 516813) in TBS for 30 minutes on the shaker to deplete endogenous peroxides activity, and were 3 times washed in TBS for 10 minutes afterwards.

**[0112]** To increase permeability of membranes in the brain tissue, samples were exposed to 0.3% Triton X-100 (Sigma Aldrich; Cat.-No.: 516813) in TBS for 10 minutes on the shaker. Unspecific proteinophilic binding in the tissue was saturated by incubating the samples with 5% goat serum (Gibco; Cat.-No.: 16210-064) in TBS for 30 minutes on the shaker.

**[0113]** A primary antibody against beta-Amyloid or synaptophysin was added and incubated overnight on the shaker at 4°C. The antibody was polyclonal and raised against a synthetic beta-Amyloid 1-43 peptide in rabbit. The primary antibody was used in a final dilution of 1 to 250 in TBS containing 2% goat serum and 10% Triton X-100. Afterwards, the samples were washed with TBS 3 times for 10 minutes and a secondary antibody was added for 90 minutes on the shaker at 4°C. The secondary antibody was a biotinylated anti-rabbit IgG raised in goat (Vectastain ABC Kit, Rabbit IgG; Vector Laboratories; Cat.-No.: PK-6101), and was used in a final dilution of 1 to 60 in TBS containing 1% goat serum and 10% Triton-X-100.

**[0114]** After washing 3 times for 10 minutes in TBS, samples were incubated in TBS containing 3% avidin solution (Vectastain ABC Kit, Rabbit IgG; Vector Laboratories; Cat.-No.: PK-6101)

and 3% biotinylated horseradish peroxidase solution (Vectastain ABC Kit, Rabbit IgG; Vector Laboratories; Cat.-No.: PK-6101) for 90 minutes at 4°C on the shaker. The samples were then washed 3 times in TBS for 10 minutes and then stained by adding phosphate-buffered saline containing 3% SG blue solution (SG Blue Peroxidase Kit; Vector Laboratories; Cat.-No.: SK-4700) and 3% H<sub>2</sub>O<sub>2</sub> solution for 5 minutes and then again washed 3 times for 10 minutes with phosphate-buffered saline.

**[0115]** To enhance staining, samples were incubated with ddH<sub>2</sub>O containing 0.5% CuSO<sub>4</sub> (w/w) for 5 minutes. After washing 3 times for 10 minutes in ddH<sub>2</sub>O, the brain slices were put onto silane-coated glass slides with a fine brush where they could dry overnight. Finally the slides were cover slipped with an aquatic mounting medium (VectaMountAQ Mounting Medium; Vector Laboratories, Cat.-No.: H-5501). As control served a set of samples from a 17 month old non-transgenic littermate processed together with the experimental samples.

**[0116]** Sections photographed under a microscope (Zeiss, Germany), randomised unbiased dissectors were overlain on to the brain section images and analysed using a Multi threshold plug-in with the software Image J (NIH, USA). Data were analysed using a one-way ANOVA with post-hoc Bonferroni tests.

### **Parkinson mouse model**

**[0117]** A standard model to induce Parkinson like symptoms in mice is the injection of the chemical (MPTP) (Li et al., 2009). This chemical impairs or kills neurons in the brain that produce dopamine. The mice develop motor impairments, and the dopaminergic neurons in the brain are reduced in numbers and function. The enzyme tyrosine hydroxylase (TH) is required to synthesise dopamine. A loss of TH signifies a loss of dopamine production (Harkavyi et al., 2008).

**[0118]** Adult male C57BL/6 mice were given the dopaminergic toxin MPTP (20 mg/kg in 0.1 mL of PBS i.p. at 2-h intervals of 4 doses MPTP; Sigma) or vehicle (PBS) for one day. This treatment selectively affects dopaminergic neurons and induces Parkinson -like symptoms in mice. One group did not receive MPTP as a non-lesioned control.

**[0119]** The dose of peptide tested was 10nmol/kg body weight. The peptides were administered by a once-daily intraperitoneal injection for 2 weeks. Saline injections were administered as a control. Six animals per group were injected with a peptide or a saline control.

### **Rotarod motor control test**

**[0120]** The rotarod consists of a rotating pole that accelerates over time. Mice are placed on

the rod, and motor skills are tested by accelerating the rotation. As the rotation increases, animals lose grip and fall on a cushion located below the rod. Mice were given three trials with 45 min inter-trial intervals on each of 2 consecutive days for 3 weeks. Each animal's endurance time was recorded, and the average was calculated. Data were analysed using a one-way ANOVA with post-hoc Bonferroni tests.

### **Immunohistochemistry for TH in the substantia nigra pars compacta and the striatum**

**[0121]** Six animals per group were analysed for expression of tyrosine hydroxylase (TH), a marker for dopamine production. Coronal brain sections (20µm) from striatum (bregma 11 to 10.2) and SNpc (bregma 24.80 to 26.04) were analysed by immunohistochemistry using antibodies recognising TH.

**[0122]** Sections were cut on a cryostat and postfixed in 4% paraformaldehyde, washed in PBS, treated with 0.3% H<sub>2</sub>O<sub>2</sub> in methanol for 20 min, and washed again. Incubation with TH (1:800) antibody was at 48°C overnight in PBS with 0.1 % Tween and 10% goat serum. Sections were incubated for 1 hr at room temperature with biotinylated secondary antibody diluted in 0.1% PBS-Tween. DAB staining was performed according to the Vectastain ABC kit instructions (Vector Laboratories). For each animal, three tissue sections from one level of striatum were stained and analysed for TH-positive fibre innervation. To achieve a TH cell count representative of the whole SN, each animal was analysed at four and three rostrocaudal levels (bregma - 4.80 to - 6.04). Two tissue sections from each level were quantified by immunohistochemistry.

**[0123]** Data were analysed using a one-way ANOVA with post-hoc Bonferroni tests. The results are illustrated in Figures 1 to 4.

### **Conclusion**

**[0124]** The results demonstrate that both GLP-1/GIP dual agonist peptides (DA4 and DA5) showed superior neuroprotection over single GLP-1 and GIP analogues in animal models of Alzheimer's and Parkinson's disease. Thus, the data of the present examples demonstrate that both peptides DA4 and DA5 may be considered suitable for use in the treatment and/or prevention of neurodegenerative disorders such as, for example, Alzheimer's Disease and/or Parkinson's Disease.

### **Example 2**

**[0125]** Materials and methods in Example 2 are the same as Example 1 except when stated. Reagents used in Example 2 only include antibodies to p-PI3Kp85, PI3Kp85, p-Akt (ser473),

Akt, p-GSK3 $\beta$  (Y216), and GSK3 $\beta$  from Abcam, Inc. Cambridge, UK. The antibody to p-tau (AT8) from Thermo Scientific USA. Mouse Anti- $\beta$ -actin, Anti-Rabbit IgG, Anti-mouse IgG, CY3-Anti-mouse IgG, RIPA Lysate, BCA Protein Assay Kit, DAPI staining solution and Anti-fluorescence decay agent from Boster (Wuhan, China).

#### **Open field test for investigating locomotor activity and exploratory behaviour.**

[0126] Before the experiment, the animals had a 30 min of adaptation in the laboratory. The open field was divided into 16 equal size squares with defining 4 squares in the middle as central area and the rest as peripheral area. Each mouse was placed individually in the apparatus (40cm  $\times$  40cm  $\times$  40cm) from the center for 5 min and the total distance as well as the percentage of time in center were recorded by Smart 3.0 software system (Panlab, Spain) at the same time. The apparatus was cleaned with 70% ethanol before each test.

#### **Y-Maze test for measuring the working memory in AD mice**

[0127] Spatial working memory was assessed by recording spontaneous alternation behaviour in a Y-maze. The maze was made of three 30 cm long, 7 cm high, 15 cm wide arms and the arms converged in an equilateral triangular central area. Each mouse was placed at the central triangular area and allowed to move freely through the maze during an 8-min session. The total arm entries and arm order was recorded by Smart 3.0 software system. Alternation was defined as successive entries into the three arms, on overlapping triplet sets. The percent alternation was calculated as the ratio of actual to possible alternations (defined as the total number of arm entries minus 2) multiplied by 100.

#### **Classical Morris water maze and reversal Morris water maze tests for measuring spatial learning and memory.**

[0128] Classical Morris water maze (MWM) test was used to investigate spatial reference memory of mouse. The water maze mainly included a stainless steel pool and a platform in the pool for escape. There were several landmarks fixed to the walls of the water maze. The pool was filled with tap water at 22 $\pm$ 2 °C. The first phase was acquisition trial phase which consisted of five training days (days 1-5) and four trials each day. The pool (diameter, 120 cm; high, 50 cm) was divided into four quadrants and the escape platform (diameter, 12 cm) hidden 1 cm beneath the water surface was placed in the center of the first quadrant. Four points equally spaced along the circumference of the pool were served as the starting position, which was chosen randomly across the four trials per day. When the mouse climbed onto the platform or the time reached 1 minute, the records stopped. The escape latencies were recorded by Ethovision 3.0 software system (Noldus Information Technology, the Netherlands). The second phase was probe trial phase. On the day after finishing the acquisition task (day



6), a probe trial was performed to assess the spatial memory. The platform was removed from the pool, and animals were allowed to swim freely for 1 min. The percentage of swimming time in target quadrant (the first quadrant) and frequency of platform crossing were recorded. Reversal MWM test was used to observe the ability of mice to relearn and cognitive flexibility. After classical MWM test, the platform was moved to reversal quadrant (the third quadrant), and mice were placed into the water for similar four days' acquisition trial and one day' probe trial as mentioned above. The third phase was visible platform test. Raise the platform 1cm above the water level and the mice were placed in two randomly selected quadrants. Swimming time to target was recorded.

#### **In vivo hippocampal measurements of L-LTP in AD mice**

[0129] Mice were anesthetized with urethane (Sigma, UK, 1.5 g/kg, i.p.) and placed in a stereotaxic apparatus (RWD Life Science, China) for L-LTP recording. Cut off the scalp and expose the skull, small holes were drilled on one side of the skull (2.0 mm posterior to bregma and 1.5 mm lateral to the midline).

[0130] A pair of parallel stimulating/recording electrodes (Sequim, WA, USA) was inserted at the Schaffer collateral and stratum radiatum in the CA1 region of hippocampus. Baseline fEPSPs were elicited by test stimuli at an interval of 30 seconds and 30 minutes fEPSPs were recorded to observe whether the underlying synaptic transmission is affected. Then two paired test stimuli with an interval of 50 ms were given before HFS to induce paired-pulse facilitation (PPF). The change in PPF ratio, calculated by dividing the slope of the second fEPSP by the slope of the first fEPSP, is viewed to be related with the changes in neurotransmitter release from presynaptic terminals. L-LTP was induced by three sets of high-frequency stimulation (HFS) with an interval of 5 minutes. Each set of HFS includes three trains of 20 pulses with 5 ms of interstimulus interval (200 Hz). The fEPSPs were recorded for 180 minutes after HFS. All events were recorded by a biological signal processing system (Chengdu Instruments Ltd, China). The percentage change of fEPSPs in each group after HFS was compared.

#### **Immunostaining for phosphorylated tau protein in the hippocampus**

[0131] After *in vivo* hippocampal L-LTP recording, the mice were anesthetized with chloral hydrate (0.007ml/g) by intraperitoneal injection and then the brain tissue was fixed with paraformaldehyde for 24 hours, after that the tissue was transferred to 30% sucrose solution for dehydration. The brain tissue was harvested, frozen in -80°C, and then sliced at a thickness of 30 µm, put the frozen section in PBS, blocked with normal goat serum (Solarbio, Beijing, China), followed by adding the primary antibody, second antibody and DAPI. Phosphorylated tau protein was observed by confocal laser scanning microscopy.

#### **Western blotting for assessing hippocampal levels of PI3K/AKT/GSK3β**

**[0132]** The hippocampi of mouse were dissected and homogenized in tissue protein extraction reagent, and protein concentration was measured using BCA Protein Assay Kit. A total of 50 µg of protein was used from each sample. Sample proteins were separated on 12% SDS-polyacrylamide gels. After that, the proteins were transferred onto PVDF membranes and nonspecific binding was blocked with 5% BSA. A primary antibody was used to incubate the membranes overnight at 4°C, followed by a secondary antibody for 2 h. The optical density of the target strip (p-PI3K, t-PI3K, p-AKT, t-AKT, p-GSK3β and t-GSK3β) was analyzed by using a gel image processing system.

**[0133]** Data were analysed using a one-way ANOVA with post-hoc Bonferroni tests. The results are illustrated in Figures 5 to 11.

### **Conclusion**

**[0134]** The results present in this study confirmed the protective effects of the GLP-1/GIP dual agonist DA5 peptide on the cognitive function and pathological characteristics in APP/PS1 transgenic AD mice. The neuroprotection of the DA5 peptide may be associated with the improvement of hippocampal synaptic plasticity and PI3K/AKT/GSK3β signalling pathway. The above results suggest that the DA5 peptide may be useful in the treatment of neurodegenerative disease, and particularly may be beneficial for AD patients, especially those with T2DM or hyperglycemia.

### **References**

#### **[0135]**

Aviles-Olmos I, Dickson J, Kefalopoulou Z, Djamshidian A, Ell P, Soderlund T, Whitton P, Wyse R, Isaacs T, Lees A, Limousin P, Foltynie T (2013) Exenatide and the treatment of patients with Parkinson's disease. *The Journal of clinical investigation* 123:2730-6.

Aviles-Olmos I, Dickson J, Kefalopoulou Z, Djamshidian A, Kahan J, Fmedsci PE, Whitton P, Wyse R, Isaacs T, Lees A, Limousin P, Foltynie T (2014) Motor and Cognitive Advantages Persist 12 Months After Exenatide Exposure in Parkinson's Disease. *Journal of Parkinson's disease* 4:337-344.

Baggio LL, Drucker DJ (2007) Biology of incretins: GLP-1 and GIP. *Gastroenterology* 132:2131-2157.

Bertilsson G, Patrone C, Zachrisson O, Andersson A, Dannaeus K, Heidrich J, Kortessmaa J, Mercer A, Nielsen E, Ronnholm H, Wikstrom L (2008) Peptide hormone exendin-4 stimulates subventricular zone neurogenesis in the adult rodent brain and induces recovery in an animal

model of Parkinson's disease. *Journal of neuroscience research* 86:326-338.

Blennow K, de Leon MJ, Zetterberg H (2006) Alzheimer's disease. *Lancet* 368:387-403. Bliss TVP, Collingridge GL (1993) A synaptic model of memory: long-term potentiation in the hippocampus. *Nature* 361:31-39.

Bomfim TR, Forny-Germano L, Sathler LB, Brito-Moreira J, Houzel JC, Decker H, Silverman MA, Kazi H, Melo HM, McClean PL, Holscher C, Arnold SE, Talbot K, Klein WL, Munoz DP, Ferreira ST, De Felice FG (2012) An anti-diabetes agent protects the mouse brain from defective insulin signaling caused by Alzheimer's disease-associated Abeta oligomers. *The Journal of clinical investigation* 122:1339-53.

Campbell JE, Drucker DJ (2013) Pharmacology, physiology, and mechanisms of incretin hormone action. *Cell metabolism* 17:819-837.

Cereda E, Barichella M, Cassani E, Caccialanza R, Pezzoli G (2012) Clinical features of Parkinson disease when onset of diabetes came first: A case-control study. *Neurology* 78:1507-1511.

Christensen M, Knop FK, Holst JJ, Vilsboll T (2009) Lixisenatide, a novel GLP-1 receptor agonist for the treatment of type 2 diabetes mellitus. *IDrugs : the investigational drugs journal* 12:503-513.

Cleary JP, Walsh DM, Hofmeister JJ, Shankar GM, Kuskowski MA, Selkoe DJ, Ashe KH (2005) Natural oligomers of the amyloid-beta protein specifically disrupt cognitive function. *Nature neuroscience* 8:79-84.

Duffy AM, Holscher C (2013) The incretin analogue D-Ala(2)GIP reduces plaque load, astrogliosis and oxidative stress in an APP/PS1 mouse model of Alzheimer's disease. *Neuroscience* 228:294-300.

Elkinson S, Keating GM (2013) Lixisenatide: first global approval. *Drugs* 73:383-391.

Faivre E, Holscher C (2013a) D-Ala2GIP facilitated synaptic plasticity and reduces plaque load in aged wild type mice and in an Alzheimer's disease mouse model. *Journal of Alzheimer's disease : JAD* 35:267-283.

Faivre E, Holscher C (2013b) Neuroprotective effects of D-Ala2GIP on Alzheimer's disease biomarkers in an APP/PS1 mouse model. *Alzheimer's research & therapy* 5:20-28.

Faivre E, Gault VA, Thorens B, Holscher C (2011) Glucose-dependent insulintropic polypeptide receptor knockout mice are impaired in learning, synaptic plasticity, and neurogenesis. *J Neurophysiol* 105:1574-1580.

Gault V, Holscher C (2008a) GLP-1 agonists facilitate hippocampal LTP and reverse the impairment of LTP induced by beta-amyloid. *European journal of pharmacology* 587:112-117.

Gault VA, Holscher C (2008b) Protease-resistant glucose-dependent insulintropic polypeptide

agonists facilitate hippocampal LTP and reverse the impairment of LTP induced by beta-amyloid. *J Neurophysiol* 99:1590-1595.

Gault VA, Flatt PR, O'Harte FP (2003) Glucose-dependent insulintropic polypeptide analogues and their therapeutic potential for the treatment of obesity-diabetes. *Biochemical and biophysical research communications* 308:207-213.

Gejl M, Gjedde A, Egefjord L, Møller A, Hansen SB, Vang K, Rodell AB, Braendgaard H, Gottrup H, Schacht A, Møller N, Brock B, Rungby J (2016) In Alzheimer's Disease, Six-Month Treatment with GLP-1 Analogue Prevents Decline of Brain Glucose Metabolism: Randomized, Placebo-Controlled, Double-Blind Clinical Trial. *Frontiers in aging neuroscience* 8:1-10.

Gengler S, McClean P, McCurtin R, Gault V, Holscher C (2012) Val(8)GLP-1 rescues synaptic plasticity and reduces dense core plaques in APP/PS1 mice. *Neurobiology of aging* 33:265-276.

Han B, Hu J, Shen J, Gao Y, Lu Y, Wang T (2013) Neuroprotective effect of hydroxysafflor yellow A on 6-hydroxydopamine-induced Parkinson's disease in rats. *European journal of pharmacology* 714:83-88.

Harkavyi A, Abuirmeileh A, Lever R, Kingsbury AE, Biggs CS, Whitton PS (2008) Glucagon-like peptide 1 receptor stimulation reverses key deficits in distinct rodent models of Parkinson's disease. *Journal of neuroinflammation* 5:19(11-19).

Holscher C (2014) Insulin, incretins and other growth factors as potential novel treatments for Alzheimer's and Parkinson's diseases. *Biochemical Society transactions* 42:593-599.

Hunter K, Holscher C (2012) Drugs developed to treat diabetes, liraglutide and lixisenatide, cross the blood brain barrier and enhance neurogenesis. *BMC neuroscience* 13:33-38.

Irwin N, O'Harte F P, Gault VA, Green BD, Greer B, Harriott P, Bailey CJ, Flatt PR (2006) GIP(Lys(16)PAL) and GIP(Lys(37)PAL): Novel Long-Acting Acylated Analogues of Glucose-Dependent Insulintropic Polypeptide with Improved Antidiabetic Potential. *Journal of medicinal chemistry* 49:1047-1054.

Ji C, Xue GF, Lijun C, Feng P, Li D, Li L, Li G, Holscher C (2015) A novel dual GLP-1 and GIP receptor agonist is neuroprotective in the MPTP mouse model of Parkinson's disease by increasing expression of BDNF. *Brain research* 1634:1-11.

LaFerla FM, Oddo S (2005) Alzheimer's disease: Abeta, tau and synaptic dysfunction. *Trends Mol Med* 11:170-176.

Li Y, Liu W, Li L, Holscher C (2015) Neuroprotective effects of a GIP analogue in the MPTP Parkinson's disease mouse model. *Neuropharmacology* 101:255-263.

Li Y, Perry T, Kindy MS, Harvey BK, Tweedie D, Holloway HW, Powers K, Shen H, Egan JM, Sambamurti K, Brossi A, Lahiri DK, Mattson MP, Hoffer BJ, Wang Y, Greig NH (2009) GLP-1 receptor stimulation preserves primary cortical and dopaminergic neurons in cellular and

rodent models of stroke and Parkinsonism. *Proceedings of the National Academy of Sciences of the United States of America* 106:1285-1290.

Liu L, Venkatraman SS, Yang YY, Guo K, Lu J, He B, Mochhala S, Kan L (2008) Polymeric micelles anchored with TAT for delivery of antibiotics across the blood-brain barrier. *Biopolymers* 90:617-623.

Liu W, Jalewa J, Sharma M, Li G, Li L, Hölscher C (2015) Neuroprotective effects of lixisenatide and liraglutide in the MPTP mouse model of Parkinson's disease. *Neuroscience* 303:42-50.

Lovshin JA, Drucker DJ (2009) Incretin-based therapies for type 2 diabetes mellitus. *Nature reviews Endocrinology* 5:262-269.

Luchsinger JA, Tang MX, Shea S, Mayeux R (2004) Hyperinsulinemia and risk of Alzheimer disease. *Neurology* 63:1187-1192.

McClellan P, Holscher C (2014) Liraglutide can reverse memory impairment, synaptic loss and reduce plaque load in aged APP/PS1 mice, a model of Alzheimer's disease. *Neuropharmacology* 76:57-67.

McClellan P, Parthasarathy V, Faivre E, Hölscher C (2011) The diabetes drug Liraglutide prevents degenerative processes in a mouse model of Alzheimer's disease. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 31:6587-6594.

McClellan PL, Gault VA, Harriott P, Holscher C (2010) Glucagon-like peptide-1 analogues enhance synaptic plasticity in the brain: A link between diabetes and Alzheimer's disease. *European journal of pharmacology* 630:158-162.

Moloney AM, Griffin RJ, Timmons S, O'Connor R, Ravid R, O'Neill C (2010) Defects in IGF-1 receptor, insulin receptor and IRS-1/2 in Alzheimer's disease indicate possible resistance to IGF-1 and insulin signalling. *Neurobiology of aging* 31:224-243.

Morris JK, Bomhoff GL, Gorres BK, Davis VA, Kim J, Lee PP, Brooks WM, Gerhardt GA, Geiger PC, Stanford JA (2011) Insulin resistance impairs nigrostriatal dopamine function. *Experimental neurology* 231:171-180.

Nyberg J, Jacobsson C, Anderson MF, Eriksson PS (2007) Immunohistochemical distribution of glucose-dependent insulinotropic polypeptide in the adult rat brain. *Journal of neuroscience research* 85:2099-2119.

Ohara T, Doi Y, Ninomiya T, Hirakawa Y, Hata J, Iwaki T, Kanba S, Kiyohara Y (2011) Glucose tolerance status and risk of dementia in the community: The Hisayama Study. *Neurology* 77:1126-1134.

Radde R, Bolmont T, Kaeser SA, Coomaraswamy J, Lindau D, Stoltze L, Calhoun ME, Jäger F, Wolburg H, Gengler S, Haass C, Ghetti B, Czech C, Holscher C, Mathews PM, Jucker M (2006) Aβ<sub>42</sub>-driven cerebral amyloidosis in transgenic mice reveals early and robust

pathology. EMBO Rep 7:940-647.

Ristow M (2004) Neurodegenerative disorders associated with diabetes mellitus. J Mol Med 82:510-529.

Scherrmann JM (2002) Drug delivery to brain via the blood-brain barrier. Vascul Pharmacol 38:349-354.

Shen J (2010) Impaired neurotransmitter release in Alzheimer's and Parkinson's diseases. Neuro-degenerative diseases 7:80-83.

Talbot K, Wang HY, Kazi H, Han LY, Bakshi KP, Stucky A, Fuino RL, Kawaguchi KR, Samoyedny AJ, Wilson RS, Arvanitakis Z, Schneider JA, Wolf BA, Bennett DA, Trojanowski JQ, Arnold SE (2012) Demonstrated brain insulin resistance in Alzheimer's disease patients is associated with IGF-1 resistance, IRS-1 dysregulation, and cognitive decline. The Journal of clinical investigation 122:1316-38.

Windisch M, Gschanes A, Hutter-Paier B (1998) Neurotrophic activities and therapeutic experience with a brain derived peptide preparation. Journal of neural transmission Supplementum 53:289-298.

## **REFERENCES CITED IN THE DESCRIPTION**

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

### **Patent documents cited in the description**

- [WO9811125A \[0084\]](#)
- [WO9808871A1 \[0106\]](#)

### **Non-patent literature cited in the description**

- **SAMBROOK et al.** Molecular Cloning, A Laboratory Manual Cold Harbor-Laboratory Press 2001 0000 [\[0072\]](#)

- **AUSUBEL et al.** Current protocols in molecular biology John Wiley and Sons 1990 0000 [0072]
- **JUO, PEI-SHOW** the Concise Dictionary of Biomedicine and Molecular Biology CRC Press 2002 0000 [0072]
- The Dictionary of Cell and Molecular Biology Academic Press [0072]
- **FINAN et al.** Sci. Transl. Med., 2013, 1-16 [0084]
- Remington's Pharmaceutical Sciences Mack Publishing Co. 1985 0000 [0088]
- Remington's Pharmaceutical Sciences Mark Publishing Company 1985 0000 [0089]
- **AVILES-OLMOS IDICKSON JKEFALOPOULOU ZDJAMSHIDIAN AELL PSODERLUND TWHITTON PWYSE RISAACS TLEES A** Exenatide and the treatment of patients with Parkinson's disease The Journal of clinical investigation, 2013, vol. 123, 2730-6 [0135]
- **AVILES-OLMOS IDICKSON JKEFALOPOULOU ZDJAMSHIDIAN AKAHAN JFMEDSCI PEWHITTON PWYSE RISAACS TLEES A** Motor and Cognitive Advantages Persist 12 Months After Exenatide Exposure in Parkinson's Disease Journal of Parkinson's disease, 2014, vol. 4, 337-344 [0135]
- **BAGGIO LLDRUCKER DJ** Biology of incretins: GLP-1 and GIP Gastroenterology, 2007, vol. 132, 2131-2157 [0135]
- **BERTILSSON GPATRONE CZACHRISSON OANDERSSON ADANNAEUS KHEIDRICH JKORTESMAA JMERCER ANIELSEN ERONNHOLM H** Peptide hormone exendin-4 stimulates subventricular zone neurogenesis in the adult rodent brain and induces recovery in an animal model of Parkinson's disease Journal of neuroscience research, 2008, vol. 86, 326-338 [0135]
- **BLENNOW KDE LEON MJZETTERBERG H** Alzheimer's disease Lancet, 2006, vol. 368, 387-403 [0135]
- **BLISS TVPCOLLINGRIDGE GLA** synaptic model of memory: long-term potentiation in the hippocampus Nature, 1993, vol. 361, 31-39 [0135]
- **BOMFIM TRFORNY-GERMANO LSATHLER LBBRITO-MOREIRA JHOUZEL JCDECKER HSILVERMAN MAKAZI HMELO HMMCCLEAN PL** An anti-diabetes agent protects the mouse brain from defective insulin signaling caused by Alzheimer's disease-associated Abeta oligomers The Journal of clinical investigation, 2012, vol. 122, 1339-53 [0135]
- **CAMPBELL JEDRUCKER DJ** Pharmacology, physiology, and mechanisms of incretin hormone action Cell metabolism, 2013, vol. 17, 819-837 [0135]
- **CEREDA EBARICHELLA MCASSANI ECACCIALANZA RPEZZOLI G** Clinical features of Parkinson disease when onset of diabetes came first: A case-control study Neurology, 2012, vol. 78, 1507-1511 [0135]
- **CHRISTENSEN MKNOP FKHOLST JJVILSBOLL T** Lixisenatide, a novel GLP-1 receptor agonist for the treatment of type 2 diabetes mellitus I Drugs : the investigational drugs journal, 2009, vol. 12, 503-513 [0135]
- **CLEARY JPWALSH DMHOFMEISTER JJSHANKAR GMKUSKOWSKI MASELKOE DJASHE KH** Natural oligomers of the amyloid-beta protein specifically disrupt cognitive function Nature neuroscience, 2005, vol. 8, 79-84 [0135]
- **DUFFY AMHOLSCHER C** The incretin analogue D-Ala(2)GIP reduces plaque load, astrogliosis and oxidative stress in an APP/PS1 mouse model of Alzheimer's

- disease *Neuroscience*, 2013, vol. 228, 294-300 [0135]
- **ELKINSON SKEATING GM** Lixisenatide: first global approval *Drugs*, 2013, vol. 73, 383-391 [0135]
  - **FAIVRE EHOLSCHER CD** Ala2GIP facilitated synaptic plasticity and reduces plaque load in aged wild type mice and in an Alzheimer's disease mouse model *Journal of Alzheimer's disease : JAD*, 2013, vol. 35, 267-283 [0135]
  - **FAIVRE EHOLSCHER C** Neuroprotective effects of D-Ala2GIP on Alzheimer's disease biomarkers in an APP/PS1 mouse model *Alzheimer's research & therapy*, 2013, vol. 5, 20-28 [0135]
  - **FAIVRE EGAULT VATHORENS BHOLSCHER C** Glucose-dependent insulinotropic polypeptide receptor knockout mice are impaired in learning, synaptic plasticity, and neurogenesis *J Neurophysiol*, 2011, vol. 105, 1574-1580 [0135]
  - **GAULT VHOLSCHER C** GLP-1 agonists facilitate hippocampal LTP and reverse the impairment of LTP induced by beta-amyloid *European journal of pharmacology*, 2008, vol. 587, 112-117 [0135]
  - **GAULT VAHOLSCHER C** Protease-resistant glucose-dependent insulinotropic polypeptide agonists facilitate hippocampal LTP and reverse the impairment of LTP induced by beta-amyloid *J Neurophysiol*, 2008, vol. 99, 1590-1595 [0135]
  - **GAULT VAFLATT PRO'HARTE F** Glucose-dependent insulinotropic polypeptide analogues and their therapeutic potential for the treatment of obesity-diabetes *Biochemical and biophysical research communications*, 2003, vol. 308, 207-213 [0135]
  - **GEJL MGJEDDE AEGEFJORD LMØLLER AHANSEN SBVANG KRODELL ABBRAENDGAARD HGOTTRUP HSCHACHT A** In Alzheimer's Disease, Six-Month Treatment with GLP-1 Analogue Prevents Decline of Brain Glucose Metabolism: Randomized, Placebo-Controlled, Double-Blind Clinical Trial *Frontiers in aging neuroscience*, 2016, vol. 8, 1-10 [0135]
  - **GENGLER SMCCLEAN PMCCURTIN RGAULT VHOLSCHER C** Val(8)GLP-1 rescues synaptic plasticity and reduces dense core plaques in APP/PS1 mice *Neurobiology of aging*, 2012, vol. 33, 265-276 [0135]
  - **HAN BHU JSHEN JGAO YLU YWANG T** Neuroprotective effect of hydroxysafflor yellow A on 6-hydroxydopamine-induced Parkinson's disease in rats *European journal of pharmacology*, 2013, vol. 714, 83-88 [0135]
  - **HARKAVYI AABUIRMEILEH ALEVER RKINGSBURY AEBIGGS CSWHITTON P** Glucagon-like peptide 1 receptor stimulation reverses key deficits in distinct rodent models of Parkinson's disease *Journal of neuroinflammation*, 2008, vol. 5, 1911-19 [0135]
  - **HOLSCHER C** Insulin, incretins and other growth factors as potential novel treatments for Alzheimer's and Parkinson's diseases *Biochemical Society transactions*, 2014, vol. 42, 593-599 [0135]
  - **HUNTER KHOLSCHER C** Drugs developed to treat diabetes, liraglutide and lixisenatide, cross the blood brain barrier and enhance neurogenesis *BMC neuroscience*, 2012, vol. 13, 33-38 [0135]
  - **IRWIN NO'HARTE F PGAULT VAGREEN BDGREER BHARRIOTT PBAILEY CJFLATT**



- PRGIP(Lys(16)PAL) and GIP(Lys(37)PAL): Novel Long-Acting Acylated Analogues of Glucose-Dependent Insulinotropic Polypeptide with Improved Antidiabetic Potential *Journal of medicinal chemistry*, 2006, vol. 49, 1047-1054 [0135]
- **JI CXUE GFLIJUN CFENG PLI DLI LLI GHOLSCHER CA** novel dual GLP-1 and GIP receptor agonist is neuroprotective in the MPTP mouse model of Parkinson's disease by increasing expression of BDNF *Brain research*, 2015, vol. 1634, 1-11 [0135]
  - **LAFERLA FMODDO SA** Alzheimer's disease: Abeta, tau and synaptic dysfunction *Trends Mol Med*, 2005, vol. 11, 170-176 [0135]
  - **LI Y, LIU W, LI L, HOLSCHER C** Neuroprotective effects of a GIP analogue in the MPTP Parkinson's disease mouse model *Neuropharmacology*, 2015, vol. 101, 255-263 [0135]
  - **LI Y PERRY TKINDY MSHARVEY BKTWEEDIE DHOLLOWAY HWPOWERS KSHEN HEGAN JMSAMBAMURTI K** GLP-1 receptor stimulation preserves primary cortical and dopaminergic neurons in cellular and rodent models of stroke and Parkinsonism *Proceedings of the National Academy of Sciences of the United States of America*, 2009, vol. 106, 1285-1290 [0135]
  - **LIU L VENKATRAMAN S SYANG Y YGUO KLU JHE BMOOCHHALA SKAN L** Polymeric micelles anchored with TAT for delivery of antibiotics across the blood-brain barrier *Biopolymers*, 2008, vol. 90, 617-623 [0135]
  - **LIU W JALEWA J SHARMA M LI GLI LHÖLSCHER C** Neuroprotective effects of lixisenatide and liraglutide in the MPTP mouse model of Parkinson's disease *Neuroscience*, 2015, vol. 303, 42-50 [0135]
  - **LOVSHIN JADRUCKER DJ** Incretin-based therapies for type 2 diabetes mellitus *Nature reviews Endocrinology*, 2009, vol. 5, 262-269 [0135]
  - **LUCHSINGER J ATANG M XSHEA S MAYEUX R** Hyperinsulinemia and risk of Alzheimer disease *Neurology*, 2004, vol. 63, 1187-1192 [0135]
  - **MCCLEAN PHOLSCHER C** Liraglutide can reverse memory impairment, synaptic loss and reduce plaque load in aged APP/PS1 mice, a model of Alzheimer's disease *Neuropharmacol*, 2014, vol. 76, 57-67 [0135]
  - **MCCLEAN P PARTHSARATHY V FAIVRE E HÖLSCHER C** The diabetes drug Liraglutide prevents degenerative processes in a mouse model of Alzheimer's disease *The Journal of neuroscience : the official journal of the Society for Neuroscience*, 2011, vol. 31, 6587-6594 [0135]
  - **MCCLEAN P LGAULT V AHARRIOTT PHOLSCHER C** Glucagon-like peptide-1 analogues enhance synaptic plasticity in the brain: A link between diabetes and Alzheimer's disease *European journal of pharmacology*, 2010, vol. 630, 158-162 [0135]
  - **MOLONEY AM GRIFFIN R J TIMMONS SO'CONNOR R RAVID RO'NEILL C** Defects in IGF-1 receptor, insulin receptor and IRS-1/2 in Alzheimer's disease indicate possible resistance to IGF-1 and insulin signalling *Neurobiology of aging*, 2010, vol. 31, 224-243 [0135]
  - **MORRIS J KBOMHOFF G LGORRES B K DAVIS V AKIM J LEE P PBROOKS W M GERHARDT G AGEIGER P C STANFORD J A** Insulin resistance impairs nigrostriatal dopamine function *Experimental neurology*, 2011, vol. 231, 171-180 [0135]
  - **NYBERG J JACOBSSON C ANDERSON M FERIKSSON P S** Immunohistochemical distribution of glucose-dependent insulinotropic polypeptide in the adult rat brain *Journal*

- of neuroscience research, 2007, vol. 85, 2099-2119 [\[0135\]](#)
- **OHARA T DOI Y NINOMIYA THIRAKAWA YHATA JIWAKI TKANBA SKIYOHARA Y** Glucose tolerance status and risk of dementia in the community: The Hisayama Study *Neurology*, 2011, vol. 77, 1126-1134 [\[0135\]](#)
  - **RADDE R BOLMONT TKAESER SACOOMARASWAMY JLINDAU DSTOLTZE L CALHOUN MEJAGGI FWOLBURG HGENGLER S** beta42-driven cerebral amyloidosis in transgenic mice reveals early and robust pathology *EMBO Rep*, 2006, vol. 7, 940-647 [\[0135\]](#)
  - **RISTOW M** Neurodegenerative disorders associated with diabetes mellitus *J Mol Med*, 2004, vol. 82, 510-529 [\[0135\]](#)
  - **SCHERRMANN J M** Drug delivery to brain via the blood-brain barrier *Vascul Pharmacol*, 2002, vol. 38, 349-354 [\[0135\]](#)
  - **SHEN J** Impaired neurotransmitter release in Alzheimer's and Parkinson's diseases *Neuro-degenerative diseases*, 2010, vol. 7, 80-83 [\[0135\]](#)
  - **TALBOT K KWANG H YKAZI H HAN LYBAKSHI K PSTUCKY A FUINO R L KAWAGUCHI K R SAMOYEDNY A J WILSON R S** Demonstrated brain insulin resistance in Alzheimer's disease patients is associated with IGF-1 resistance, IRS-1 dysregulation, and cognitive decline *The Journal of clinical investigation*, 2012, vol. 122, 1316-38 [\[0135\]](#)
  - **WINDISCH M G SCHANES A HUTTER-PAIER B** Neurotrophic activities and therapeutic experience with a brain derived peptide preparation *Journal of neural transmission Supplementum*, 1998, vol. 53, 289-298 [\[0135\]](#)

**PATENTKRAV**

1. GIP/GLP-1 co-agonist-peptid vist med den generelle formel I:

Tyr-Aib-Glu-Gly-Thr-Phe-Thr-Ser-Asp-Tyr-Ser-Ile-Tyr-Leu-Asp-Lys-Gln-Ala-Ala-Aib-  
Glu-Phe-Val-Xaa<sup>24</sup>-Trp-Leu-Leu-Ala-Gly-Y1-Y2-R<sup>2</sup>

5

(I) (SEQ. ID. No. 1)

hvor Xaa<sup>24</sup> er Asn;

Y1 er Gly-Pro-Ser-Ser-Gly-Ala-Pro-Pro-Pro-Ser-Lys (SEQ. ID. Nr. 7);

Y2 er valgt blandt:

Lys-Lys-Lys-Lys-Lys-Lys (SEQ. ID. Nr. 3); og

10

Arg-Arg-Gln-Arg-Arg-Lys-Lys-Arg-Gly-Tyr (SEQ. ID. Nr. 4); og

R<sup>2</sup> er -NH<sub>2</sub>,

eller et farmaceutisk acceptabelt salt eller solvat af peptidet.

2. Peptid ifølge krav 1, som består af aminosyresekvensen:

15

YXEGTFTSDYSIYLDKQAAXEFVNWLLAGGPSSGAPPPSKKKKKK-NH<sub>2</sub> (SEQ. ID Nr.  
11)

hvor X = aminoisomørsyre.

3. Peptid ifølge krav 1, som består af aminosyresekvensen:

20

YXEGTFTSDYSIYLDKQAAXEFVNWLLAGGPSSGAPPPSKRRQRRKKRGY-N H<sub>2</sub>  
(SEQ ID ID. 12)

hvor X = aminoisomørsyre.

4. Peptid ifølge et hvilket som helst af de foregående krav, der omfatter en hydrofil del,  
25 der er kovalent bundet til en aminosyre.

5. Peptid ifølge krav 4, der omfatter en hydrofil del, der er kovalent bundet til (i) en aminosyre i position 24 eller (ii) en aminosyre i position 39 eller 40.
6. Peptid ifølge et hvilket som helst af kravene 4 til 5, hvor den hydrofile del er en vandopløselig polymer, hvor den vandopløselige polymer eventuelt er en polyethylenglykol-del og eventuelt er en polyethylenglykol-del med en molekylvægt på mellem omkring 20.000 Dalton og omkring 60.000 Dalton.
7. Peptid ifølge et hvilket som helst af de foregående krav, hvor peptidet er konjugeret med en lipofil substituent.
8. Peptid ifølge krav 7, hvor den lipofile substituent omfatter en acylgruppe og eventuelt er et fedtsyremolekyle, og yderligere hvor fedtsyremolekylet eventuelt er valgt fra en C-8-oktanoylgruppe, en C-10-decanoylgruppe, en C-12-lauroylgruppe, en C-14-myristoylgruppe, en C-16-palmitoylgruppe, en C-18-stearoylgruppe og en C-20-acylgruppe.
9. Peptid ifølge krav 7 eller krav 8, hvor den lipofile substituent er forbundet til en aminosyre ved peptidets carboxyl-terminal.
10. Peptid ifølge et hvilket som helst af kravene 7 til 9, der omfatter en spacer, der konjugerer den lipofile substituent til en aminosyrerest.
11. Peptid ifølge krav 10, hvor spacer er en rest fra en naturligt forekommende eller unaturlig aminosyre, hvor spaceren omfatter en rest af Gly, Pro, Ala, Val, Leu, Ile, Met, Cys, Phe, Tyr, Trp, His, Lys, Arg, Gln, Asn,  $\alpha$ -Glu,  $\gamma$ -Glu,  $\epsilon$ -Lys, Asp, Ser, Thr, Gaba, Aib,  $\beta$ -aminohexonyl, 7-aminoheptanoyl, 8-aminooctanoyl, 9-aminononanoyl, 10-aminodecanoyl eller 8-amino-3,6-dioxaoctanoyl.
12. Peptid ifølge krav 11, hvor spaceren er  $\gamma$ -Glu, eller hvor spaceren er et dipeptid, eventuelt hvor spaceren omfatter to negativt ladede aminosyrer, og yderligere eventuelt hvor spaceren er  $\gamma$ -Glu- $\gamma$ -Glu.
13. Peptid ifølge et hvilket som helst af de foregående krav, som er til anvendelse til behandling og/eller profylakse af en neurologisk lidelse, såsom en neurologisk lidelse, der er forårsaget af eller associeret med beta-amyloid-

proteinaflejring i en del af patienten,  
eller en neurologisk lidelse, der er forårsaget af eller forbundet med:

a) dysfunktion ved langvarig forstærkning af synaptisk transmission; eller

b) betændelse.

- 5 14. Peptid ifølge krav 13, som er til anvendelse til behandling og/eller profylakse af en neurologisk lidelse, der påvirker den kognitive funktion, fx demens, slagtilfælde, skizofreni og/eller bipolar lidelse, hvor lidelsen eventuelt er cerebral iskæmi associeret med slagtilfælde,  
eller til anvendelse til behandling og/eller profylakse af en lidelse valgt blandt posttraumatisk stresslidelse, epilepsi, Tourettes syndrom og hallucinationer, og dysfunktionelle kognitive processer, eventuelt valgt blandt opmærksomhed, beregning, hukommelse, vurdering, indsigt, læring og ræsonnement,  
10 eller til anvendelse til behandling og/eller profylakse af en neurodegenerativ lidelse, fx Alzheimers sygdom, Parkinsons sygdom, amyotrof lateral sklerose, perifer neuropati, Huntingtons sygdom og Creutzfeldt-Jacob sygdom,  
15 eller til anvendelse til behandling og/eller profylakse af en neurologisk lidelse valgt blandt klinisk eller præklinisk Alzheimers sygdom, prodromal Alzheimers sygdom og klinisk eller præklinisk amyloid angiopati (CAA).

- 20 15. Peptid til anvendelse til behandling og/eller profylakse af klinisk Alzheimers sygdom, hvor peptidet består af følgende sekvens:

(a) YXEGTFTSDYSIYLDKQAAXEFVNWLLAGGPSSGAPPPSKKKKKK-NH<sub>2</sub>


(SEQ ID Nr. 11)

hvor X = aminoismørsyre; eller

- 25 (b) YXEGTFTSDYSIYLDKQAAXEFVNWLLAGGPSSGAPPP-SKRRQRRRRKRGY-NH<sub>2</sub> (SEQ ID Nr. 12)

hvor X = aminoismørsyre.

# DRAWINGS



## Quantification of synapses in the cortex of APP/PS1 mice




Figure 2




Figure 3




Dopamine synthesis (TH density) in substantia nigra *pc*

Figure 4




Figure 5





Figure 6

Figure 7






Figure 8




Figure 9




Figure 10




Figure 11



Figure 12

Tyr-Aib-Glu-Gly-Thr-Phe-Thr-Ser-Asp-Tyr-Ser-Ile-Tyr-Leu-Asp-Lys-Gln-Ala-Ala-Aib-Glu-Phe-  
Val-Xaa<sup>24</sup>-Trp-Leu-Leu-Ala-Gly-Y1-Y2-R<sup>2</sup>  
SEQ. ID. No 1

Lys-Lys-Lys-Lys-Lys (SEQ. ID. No 2);

Lys-Lys-Lys-Lys-Lys-Lys (SEQ. ID. No. 3)

Arg-Arg-Gln-Arg-Arg-Lys-Lys-Arg-Gly-Tyr (SEQ. ID. No 4); and

Lys-Arg-Arg-Gln-Arg-Arg-Lys-Lys-Arg-Gly-Tyr (SEQ. ID. No 5);

Gly-Pro-Ser-Ser-Gly-Ala-Pro-Pro-Pro-Ser (SEQ. ID. No. 6);

Gly-Pro-Ser-Ser-Gly-Ala-Pro-Pro-Pro-Ser-Lys (SEQ. ID. No. 7);

Gly-Pro-Ser-Ser-Gly-Ala-Pro-Pro-Pro-Ser-Cys (SEQ. ID. No. 8);

Pro-Ser-Ser-Gly-Ala-Pro-Pro-Pro-Ser (SEQ. ID. No. 9);

Pro-Ser-Ser-Gly-Ala-Pro-Pro-Ser (SEQ. ID. No. 10);

YXEGTFTSDYSIYLDKQAAXEFVNWLLAGGPSSGAPPPSKKKKKK-NH<sub>2</sub>

SEQ ID No 11

YXEGTFTSDYSIYLDKQAAXEFVNWLLAGGPSSGAPPPSKRRQRRKRGY-NH<sub>2</sub>

X = aminoisobutyric acid

SEQ. ID. No 12

HAEGTFTSDV SSYLEGQAAK EFlAWLVKGR G-OH (SEQ ID No. 13)

YAEgTFISDYSIAMDKIHQQDFVNWLLAQKGKKNDWKHNITQ-OH (SEQ ID. No 14)

His Ala Glu Gly Thr Phe Thr Ser Asp Val Ser Ser Tyr Leu Glu Gly Gln Ala Ala Xaa Glu Phe Ile  
Ala Trp Leu Val Arg Gly Arg (SEQ ID No. 15)

Y $\alpha$ -AEGTFISDYSIAMDKIHQQDFVNWLLAQKGKK( $\gamma$ -E-C16)NDWKHNITQ (SEQ ID. No  
16)