

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property  
Organization

International Bureau

(43) International Publication Date  
04 April 2019 (04.04.2019)



(10) International Publication Number  
**WO 2019/063756 A1**

(51) International Patent Classification:

G01N 33/68 (2006.01) A61K 38/18 (2006.01)  
A61K 31/00 (2006.01)

TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,  
KM, ML, MR, NE, SN, TD, TG).

(21) International Application Number:

PCT/EP2018/076391

Published:

- with international search report (Art. 21(3))
- with sequence listing part of description (Rule 5.2(a))

(22) International Filing Date:

28 September 2018 (28.09.2018)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

17194169.3 29 September 2017 (29.09.2017) EP  
18169317.7 25 April 2018 (25.04.2018) EP

(71) Applicant: **MERCK PATENT GMBH** [DE/DE]; Frank-  
furter Strasse 250, 64293 Darmstadt (DE).

(72) Inventors: **LADEL, Christoph H.**; Meisenweg 3, 64291  
Darmstadt (DE). **GUEHRING, Hans**; An den Drei Weisen  
28, 65366 Geisenheim (DE). **BAY-JENSEN, Anne-Chris-  
tine**; Ahrenkildes alle 9, 2300 Copenhagen (DK). **KARS-  
DAL, Morten**; Eckersbergsgade 13, 2100 Copenhagen  
(DK). **QVIST, Per**; Norre alle 19V, 4, 2200 Copenhagen  
(DK).

(74) Agent: **MERCK SERONO S.A.**; Terre Bonne Business  
Park A2 Route de Crassier 15, 1262 Eysins (CH).

(81) Designated States (unless otherwise indicated, for every  
kind of national protection available): AE, AG, AL, AM,  
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ,  
CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO,  
DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN,  
HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP,  
KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME,  
MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ,  
OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA,  
SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN,  
TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every  
kind of regional protection available): ARIPO (BW, GH,  
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ,  
UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,  
TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,  
EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,  
MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,

(54) Title: INFLAMMATORY BIOMARKERS FOR PREDICTING RESPONSIVENESS TO FGF-18 COMPOUND

(57) Abstract: The present invention relates to biomarkers associated with the clinical response to an FGF-18 compound before or during treatment of a cartilage disorder. It relates more particularly to specific proteins present in the blood, serum, synovial fluid or in the urine, which can be used as biomarkers for the diagnosis, pre-treatment of patients and during therapy of cartilage disorders. The invention can be used in predicting the response to an FGF-18 compound treatment, before starting the treatment or during the treatment. It could be used for selecting/identifying subjects to be treated according to specific doses and/or dosing regimens by intra-articular administration of an FGF-18 compound. The use of these biomarkers in diagnostics could result in increased benefit and reduced risk-benefit ratio in subjects.



WO 2019/063756 A1

## Inflammatory biomarkers for predicting responsiveness to FGF-18 compound

### Field of Invention

The present invention relates to biomarkers associated with the clinical response to an FGF-18 compound before or during treatment of a cartilage disorder. It relates more particularly to specific proteins present in the blood, serum, synovial fluid or in the urine, which can be used as biomarkers for the diagnosis, pre-treatment of patients and during therapy of cartilage disorders. The invention further discloses specific proteins that are related to cartilage response to an FGF-18 compound treatment as well as diagnostic tools and kits based on their quantities or expression profile of proteins related to tissue inflammation. Thus, the invention can be used in predicting the response to an FGF-18 compound treatment, before starting the treatment or during the treatment. It could be used for selecting/identifying subjects to be treated according to specific doses and/or dosing regimens by intra-articular administration of an FGF-18 compound. The use of these biomarkers in diagnostics could result in increased benefit and reduced risk-benefit ratio in subjects.

15

### Background of the invention

Cartilage disorders broadly refer to diseases characterized by degeneration of metabolic abnormalities in the connective tissues which are manifested by pain, stiffness and limitation of motion of the affected body parts. These disorders can be due to pathology or can be the result of trauma or injury. Among others, cartilage disorders include osteoarthritis (OA), cartilage injury (inclusive sports injuries of cartilage and joint, and surgical injuries such as microfracture(s)). Mature cartilage has limited ability to repair itself, notably because mature chondrocytes have little potential for proliferation and due to the absence of blood vessels. In addition, cartilage is not well nurtured and has a low oxygen pressure. Replacement of damaged cartilage, in particular articular cartilage, caused either by injury or disease is a major challenge for physicians, and available surgical treatment procedures are considered not completely predictable and effective for only a limited time. Therefore, the majority of younger subjects either does not seek treatment or are counselled to postpone treatment for as long as possible. When treatment is required, the standard procedure is age dependent and varies between total joint replacement, transplantation of pieces of cartilage or marrow stimulating technique (such as microfracture). Microfracture is a common procedure that involves penetration of the subchondral bone to stimulate cartilage deposition by bone marrow derived stem cells. However, it has been shown that this technique does not repair sufficiently the chondral defect and the new cartilage formed is mainly fibrocartilage, resulting in inadequate or altered function and biomechanics. Indeed, fibrocartilage does not have the same durability and may not adhere correctly to the surrounding hyaline cartilage. For this reason, the newly synthesized fibrocartilage may breakdown more easily (expected time frame: 5-10 years).

35

For subjects with osteoarthritis, non-surgical treatment consists notably of physical therapy, lifestyle modification(e.g. increasing physical activity), supportive devices, oral and injected drugs (e.g. non-steroidal anti-inflammatory drugs), walking aids and medical symptom management. Once these

treatments fail, surgery, such as joint replacement, is the main option for the subjects. Tibial or femoral osteotomies (cutting the bone to rebalance joint wear) may reduce symptoms, help to maintain an active lifestyle, and delay the need for total joint replacement. Total joint replacement can provide relief for the symptom of advanced osteoarthritis, but generally requires a change in a subject's lifestyle and/or activity level.

At that time, drug treatments on the market are mainly directed to pain relief. There is not yet commercially available treatment that restores or postpones the cartilage damages (see Lotz, 2010; Karsdal, 2016).

Fibroblast Growth factor 18 (FGF-18) is a member of the FGF family of proteins, closely related to FGF-8 and FGF-17. It has been shown that FGF-18 is a proliferative agent for chondrocytes and osteoblasts (Ellsworth et al., 2002; Shimoaka et al., 2002; Gigout et al., 2017). FGF-18 has been proposed for the treatment of cartilage disorder such as osteoarthritis and cartilage injury either alone (WO2008/023063) or in combination with hyaluronic acid (WO2004/032849).

Sprifermin, which is a truncated form of human FGF-18, is being investigated in clinical trials for treatment of both osteoarthritis and cartilage injury (for more details see for instance NCT01033994, NCT00911469 and NCT01066871). The current dosing regimen for sprifermin is once weekly for 3 weeks (one treatment cycle), the drug being administered via intraarticular injections. This treatment cycle can be repeated. This dosing regimen has been described in WO2008/023063.

At that time, OA and cartilage injury treatments with FGF-18, during clinical trials, are provided to subjects without predictive information on the response (Lohmander et al., 2014; Dahlberg et al., 2016), *i.e.* without knowledge on whether the treatment will likely be highly effective, moderately effective or show only little or no effect. Currently, numerous treated subject population exhibit an intermediate/high response to treatment according to the WOMAC scores with sprifermin after at least one treatment cycle, however, some others either do not respond to said treatment or respond while presenting higher WOMAC score compared to control.

WO2014/023703 describes genetic markers (combination of SNPs IL-1RN rs9005 and IL-1RN rs315952) that are associated with the quality of the clinical response to treatment of cartilage disorder such as OA, cartilage injury or microfracture(s) with FGF-18. Such markers are useful for identifying, through genetic screening prior to the treatment, subgroups of subjects that are more likely to exhibit a particular response to treatment with FGF-18, such as a very good clinical response to treatment with FGF-18 or on the contrary those for whom the therapy may fail.

Knowledge on the type of clinical response of a subject to treatment can be used to optimize therapy or select therapy, such as selecting treatment with FGF-18 as a first line therapy or adapting the dose and/or dosing regimen. Such information will be clinically useful for the medical management of cartilage disorder, such as of OA and/or cartilage injury, in subjects. For example, if an individual with OA or cartilage injury is known to be at increased risk for not responding to the FGF-18 treatment at a given dose, the physician may exclude said subject from the FGF-18 treatment or change dose and/or dosing regimen. In addition, such predictive information may also be clinically useful to guide decisions on the dosing regimen.

There is a need to identify further biomarkers helping in optimizing therapy or in selecting a therapy, in order to provide a full range of solution for the subject to be treated or for the doctor looking for the best therapy for his patient.

## 5 Summary of the invention

As described herein, inflammatory biomarkers such as of C1M, C3M and/or CRPM can be used either alone or in combination in the detection, diagnosis and treatment of patients with osteoarthritis or cartilage disorders. The expression levels (or quantity) of at least one of these biomarkers (or combinations thereof) can be used, for instance, to detect patients to be included in specific dosing regimen for certain therapies, such as sprifermin or a fusion protein comprising an FGF-18 moiety. The present invention is directed to a method of predicting the sensitivity to treatment with an FGF-18 compound in a subject having a cartilage disorder, the method comprising the steps of:

- a) Determining, from a sample, the quantity of at least one of the biomarkers selected from the group consisting of C1M, C3M and/or CRPM;
- b) Predicting from the result of step a) good or low sensitivity of said subject to treatment with an FGF-18 compound.

According to said method, the presence of lower than  $30.0 \pm 2SD$  ng/mL of C1M and/or lower than  $12.0 \pm 2SD$  ng/mL of C3M and/or lower than  $10.0 \pm 2SD$  ng/mL of CRPM is predictive of good response (i.e. response or sensitivity) to treatment with an FGF-18 compound. FGF-18 compound could thus be administered to said subject at the following dosing regimen: at least two treatment cycles of 100  $\mu$ g of the FGF-18 compound given once per week for 3 consecutive weeks per treatment cycle. On the contrary, the presence of higher than  $30.0 \pm 2SD$  ng/mL of C1M and/or higher than  $12.0 \pm 2SD$  ng/mL of C3M and/or higher than  $10.0 \pm 2SD$  ng/mL of CRPM is predictive of low response (i.e. low sensitivity) to treatment with an FGF-18 compound. FGF-18 compound could thus be administered to said subject preferably at the following dosing regimen: four treatment cycles of 100  $\mu$ g of FGF-18 compound given once per week for 3 consecutive weeks per treatment cycle.

Also described herein is a method for selecting a subject having a cartilage disorder for inclusion in or exclusion from a specific dosing regimen when said subject is to be treated with an FGF-18 compound, based on the likelihood of their sensitivity to said treatment, comprising the steps of:

- a) Determining, from a sample, the quantity of at least one of the biomarkers selected from the group consisting of C1M, C3M and/or CRPM; wherein the quantity of at least one of these proteins is predictive about the subject's risk of having good-sensitivity or low-sensitivity to said treatment, and
- b) Selecting the appropriate dosing regimen according to the subject's risk of having good-sensitivity or low-sensitivity to said treatment.

According to said method, the subject presenting higher than  $30.0 \pm 2SD$  ng/mL of C1M and/or higher than  $12.0 \pm 2SD$  ng/mL of C3M and/or higher than  $10.0 \pm 2SD$  ng/mL of CRPM will preferably be excluded (i.e. they will not be selected) from the treatment with FGF-18 compound given at the

following dosing regimen: two treatment cycles of 100 µg of FGF-18 compound given once per week for 3 consecutive weeks per treatment cycle. To the contrary, the subject presenting lower than  $30.0 \pm 2SD$  ng/mL of C1M and/or lower than  $12.0 \pm 2SD$  ng/mL of C3M and/or lower than  $10.0 \pm 2SD$  ng/mL of will be included (i.e. they will be selected) for the treatment with FGF-18 compound given at the following dosing regimen: at least two treatment cycles of 100 µg of FGF-18 compound given once per week for 3 consecutive weeks per treatment cycle.

Also described herein is a kit comprising means for performing the method according to the invention as a whole.

Also encompassed is an FGF-18 compound for use in the treatment of a subject having a cartilage disorder, characterized in that:

- a) If the subject has lower than  $30.0 \pm 2SD$  ng/mL of C1M and/or lower than  $12.0 \pm 2SD$  ng/mL of C3M and/or lower than  $10.0 \pm 2SD$  10 ng/mL of CRPM he can receive a treatment with an FGF-18 compound given at the following dosing regimen at least two treatment cycles of 100 µg of FGF-18 compound given once per week for 3 consecutive weeks per treatment cycle and,
- b) If the subject has higher than  $30.0 \pm 2SD$  ng/mL of C1M and/or higher than  $12.0 \pm 2SD$  ng/mL of C3M and/or higher than  $10.0 \pm 2SD$  ng/mL of CRPM he can receive a treatment with an FGF-18 compound given at the following dosing regimen: four treatment cycles of 100 µg of FGF-18 compound given once per week for 3 consecutive weeks per treatment cycle.

Also described is a method for treating a subject having a cartilage disorder, comprising the following steps:

- a) Determining, from a sample, the quantity of at least one of the biomarkers selected from the group consisting of C1M, C3M and/or CRPM, wherein said quantities are predictive about the subject's risk of having good or low sensitivity to a treatment with said FGF-18 compound,
- b) Selecting the appropriate dosing regimen according to the subject's risk of having good-sensitivity or low-sensitivity to said treatment, and
- c) Administering intraarticularly an FGF-18 compound according to the dosing regimen selected in step b).

According to said method, the presence of lower than  $30.0 \pm 2SD$  ng/mL of C1M and/or lower than  $12.0 \pm 2SD$  ng/mL of C3M and/or lower than  $10.0 \pm 2SD$  ng/mL of CRPM is predictive of good response (i.e. response or sensitivity) to treatment with an FGF-18 compound in a dose-dependent way. FGF-18 compound could thus be administered to said subject at the following dosing regimen: at least two treatment cycles of 100 µg of FGF-18 compound given once per week for 3 consecutive weeks per treatment cycle. On the contrary, the presence of higher than  $30.0 \pm 2SD$  ng/mL of C1M and/or higher than  $12.0 \pm 2SD$  ng/mL of C3M and/or higher than  $10.0 \pm 2SD$  ng/mL of CRPM is predictive of low response (i.e. low sensitivity) to treatment with an FGF-18 compound. FGF-18 compound could thus be administered to said subject preferably at the following dosing regimen:

four treatment cycles of 100 µg of FGF-18 compound given once per week for 3 consecutive weeks per treatment cycle.

In particular embodiments of the present invention as a whole, *i.e.* in any of the methods or uses mentioned herein, the FGF-18 compound to be used as a treatment is sprifermin or a fusion protein comprising an FGF-18 moiety and the subject has a cartilage disorder selected from the group consisting of osteoarthritis, cartilage injury, fractures affecting joint cartilage or surgical procedures with impact on joint cartilage (e.g. Microfracture).

It is to be understood that in any of the methods or uses mentioned herein, before determining the quantity of at least one of the proteins, it is needed to obtain a sample (or a test sample) of said subject, via for instance blood, serum, synovial fluid or urine collecting. Further, it is also to be understood that any of the methods or uses mentioned herein are performed *in vitro*, and not on the animal or human body.

### Definitions

- The term "FGF-18 compound" or "FGF-18", as used herein, is intended to be a protein maintaining at least one biological activity of the human FGF-18 protein. FGF-18 may be native, in its mature form, or a truncated form thereof. Biological activities of the human FGF-18 protein include notably the increase in osteoblastic activity (see WO98/16644) or in cartilage formation (see WO2008/023063). Native, or wild-type, human FGF-18 is a protein mostly produced during skeletal development and is involved in bone and cartilage formation (See Haque et al., 2007). Human FGF-18 was first designated zFGF-5 and is fully described in WO98/16644. SEQ ID NO:1 corresponds to the amino acid sequence of the native human FGF-18, with a signal peptide consisting of amino acid residues 1(Met) to 27(Ala). The mature form of human FGF-18 corresponds to the amino acid sequence from residue 28(Glu) to residue 207(Ala) of SEQ ID NO: 1 (180 amino acids).

The FGF-18 compound, in the present invention, may be produced by recombinant methods, such as taught by the application WO2006/063362. Depending on the expression systems and conditions, FGF-18 in the present invention is expressed in a recombinant host cell with a starting Methionine (Met) residue or with a signal sequence for secretion. When expressed in prokaryotic host, such as in *E. coli*, FGF-18 contains an additional Met residue in N-terminal of its sequence. For instance, the amino acid sequence of human FGF-18, when expressed in *E.coli*, starts with a Met residue in N-term (position 1) followed by residues 28 (Glu) to residue 207 (Ala) of SEQ ID NO: 1.

The term "FGF-18 compound" also includes variants or mutants of the native, mature form, or truncated forms of FGF-18, as well as fusion proteins comprising a (biologically) active FGF-18 moiety coupled to a heterologous protein or a chemical compound (such as those disclosed in EP17192467.3 patent family). In such fusion proteins, the FGF-18 moiety can be the native, mature form, or truncated forms of the FGF-18 protein or variants or mutants thereof.

- The term "truncated form" of FGF-18, as used herein, refers to a protein which comprises or consists of residues 28(Glu) to 196(Lys) of SEQ ID NO: 1. Preferably, the truncated form of FGF-18 protein is the polypeptide designated "trFGF-18" (170 amino acids), which starts with a Met residue

(in N-terminal) followed by amino acid residues 28 (Glu) -196 (Lys) of the wild-type human FGF-18. The amino acid sequence of trFGF-18 is shown in SEQ ID NO:2 (amino acid residues 2 to 170 of SEQ ID NO:2 correspond to amino acid residues 28 to 196 of SEQ ID NO:1). trFGF-18 is a recombinant truncated form of human FGF-18, produced in E.coli (see WO2006/063362). The International Non-proprietary Name (INN) for this particular form of FGF-18 is sprifermin. Sprifermin has been shown to display similar activities as the mature human FGF-18, e.g. it increases chondrocyte proliferation and cartilage deposition leading to repair and reconstruction for a variety of cartilaginous tissues (see WO2008/023063).

- The term "marker" or "biomarker" are used interchangeably. In the context of the present invention they are proteins. A "prognostic biomarker" is informative about the subject condition, including and not limited to disease evolution, disease severity or disease outcome, regardless of any therapy. A "predictive biomarker" is informative about the effect of a received therapy, including, but not limited to efficacy and safety outcome. The prognostic and predictive definitions are not mutually exclusive thus a biomarker can be both prognostic and predictive. The quantity of biomarker or the expression level of biomarker is herein expressed as ng, µg, mg or g of a given protein. Said quantity or level can be expressed as absolute value (e.g. 10 ng or 2 µg) or as concentration (e.g. 10 ng/mL or 2 µg/mL). In relation to the biomarkers, the term "quantity" or "expression level" can be used interchangeably.

- The term "inflammatory biomarker" refers to biomarkers such as CRPM, C1M, and C3M.

- The term "CRPM" refer to C-reactive protein degraded by metalloproteinase (MMP) enzymatic activity generating a neo-epitope with sequence KAFVFP as specific epitope recognised by the assay: CRPM is a novel tissue-related measure of C-reactive protein (CRP). The difference between standard CRP and CRPM is that CRPM measure a degradation product of CRP; whereas CRP is an acute reactant released from the liver, CRPM is a product released after CRP has exerted its effect on the tissue. Thus CRPM is a measure of tissue inflammation. CRPM has been measured in different OA and RA studies and have shown to be associated disease activity and pain measures. More CRPM have shown to be prognostic for OA incidence (see e.g. Siebuhr et al. 2014 or Saberi Hosnijeh et al., 2016)

- The term "C1M" refers to Collagen type I degraded by MMP enzymatic activity generating a neo-epitope with sequence KDGVRG as specific epitope recognised by C1M assay. C1M assay measures type I collagen degradation mediated by MMP. This reflects connective tissue degradation. A significant association between baseline C1M and change to joint space narrowing was observed in RA. In addition, C1M is significantly associated with disease activity (e.g. DAS) modulated by anti-inflammatories such as tocilizumab and infliximab in RA and SpA (see e.g. Siebuhr et al., 2013 or Siebuhr et al., 2016).

- The term "C3M" refers to Collagen type 3 degraded by MMP, generating a neo-epitope with sequence KNGETG as specific epitope recognised by C3M assay: C3M measures type III collagen degradation mediated by MMP. This reflects connective tissue degradation and seems to especially associate with joint inflammation. It is highly associated with inflammation, tissue health, disease

activity and response to treatment. A significant association between baseline C3M to disease activity and it is predictive of treatment response in RA. Furthermore, there is a significant association of C3M with occurrence of flares in gout and are associated with an inflammatory phenotype in OA. In addition, C3M is modulated by anti-inflammatories such as tocilizumab and infliximab in RA and SpA. (See e.g. Valdes et al., 2016 or Maijer et al. 2016).

- "Cartilage disorder", as used herein, encompasses disorders resulting from damages due to injury, such as traumatic injury, chondropathy or arthritis. Examples of cartilage disorders that may be treated by the administration of the FGF-18 formulation described herein include but are not restricted to arthritis, such as osteoarthritis, cartilage injury, fractures affecting joint cartilage or surgical procedures with impact on joint cartilage (e.g. Microfracture). Degenerative diseases/disorders of the cartilage or of the joint, such as chondrocalcinosis, polychondritis, relapsing polychondritis, ankylosing spondylitis or costochondritis are also encompassed by this wording. The International Cartilage Repair Society has proposed an arthroscopic grading system to assess the severity of the cartilage defect: grade 0: (normal) healthy cartilage, grade 1: the cartilage has a soft spot or blisters, grade 2: minor tears visible in the cartilage, grade 3: lesions have deep crevices (more than 50% of cartilage layer) and grade 4: the cartilage tear exposes the underlying (subchondral) bone (see for instance page 13 of [http://www.cartilage.org/files/contentmanagement/ICRS\\_evaluation.pdf](http://www.cartilage.org/files/contentmanagement/ICRS_evaluation.pdf)).

- The term "SD" means standard deviation and is linked to the usual deviations of any validation assays/systems.

- The term "Osteoarthritis" is used to intend the most common form of arthritis. The term "osteoarthritis" encompasses both primary osteoarthritis and secondary osteoarthritis (see for instance The Merck Manual, 17<sup>th</sup> edition, page 449). The most common way of classifying/grading osteoarthritis is the use of the Kellgren-Lawrence radiographic grading scale (see table below). Osteoarthritis may be caused by the breakdown of cartilage. Bits of cartilage may break off and cause pain and swelling in the joint between bones. Over time, the cartilage may wear away entirely, and the bones will rub together. Osteoarthritis can affect any joint but usually concerns hands and weight-bearing joints such as hips, knees, feet, and spine. In a preferred example, the osteoarthritis may be knee osteoarthritis or hip osteoarthritis. Osteoarthritis is one of the preferred cartilage disorders that can be treated by administering the FGF-18 compounds according to the present invention.

Kellgren-Lawrence Radiographic Grading Scale of Osteoarthritis is described as follow:

<b>Grade of Osteoarthritis</b>	<b>Description</b>
0-None	No radiographic findings of osteoarthritis
1-Doubtful	Doubtful narrowing of joint space and possible osteophytic lipping
2-Minimal	Definite osteophytes, definite narrowing of joint space
3-Moderate	Moderate multiple osteophytes, definite narrowing of joints space, some sclerosis and possible deformity of bone contour



4-Severe	Large osteophytes, marked narrowing of joint space, severe sclerosis and definite deformity of bone contour
----------	---

Grades 1 and 2 can be considered as less severe forms of the disease, whereas grades 3 and 4 can be considered as more severe forms of the disease.

- 5 - The term “cartilage injury” as used herein is a cartilage disorder or cartilage damage resulting notably from a trauma. Cartilage injuries can occur notably after traumatic mechanical destruction, notably further to an accident or surgery (for instance microfracture surgery). This term “cartilage injury” also includes chondral or osteochondral fracture, damage to meniscus, and the term microfracture. Also considered within this definition is sport-related injury or sport-related wear of tissues of the joint.
- 10 - “WOMAC total scores” or “WOMAC scores” (“WOMAC” for “Western Ontario and McMaster Universities Osteoarthritis Index”) measure pain (WOMAC pain score), function (WOMAC function score) and stiffness (WOMAC stiffness score). When applied to assessing of pain and dysfunction associated with cartilage injury, it consists of a questionnaire containing 24 items divided into 3 subscales (5 items for Pain, 2 items for Stiffness and 17 items for Physical Function)(see Bellamy et al., 1988; Wolfe, 1999). It is a well-known instrument, widely used notably in assessment of the OA severity.
- 15 - In order to evaluate cartilage repair, cartilage volume measurements were performed through magnetic resonance imaging (MRI) measurements, including but not exclusively: Lateral volume of cartilage (also referred as LFTC), Medial volume of cartilage (also referred as MFTC), Total volume of cartilage (also referred as LFTC + MFTC), and new total average cartilage thickness.
- 20 - The term “baseline” means before treatment (*i.e.* at study entry). It refers notably to clinical variables, such as, but not limited to, the cartilage volume and WOMAC total score of one given subject at study entry (*i.e.* before treatment with FGF-18 compound or placebo).
- 25 - The term “subject” or “patient” refers to both human and non-human animals. The term non human comprises mammals such as rodents (including mice), rabbits, cats, dogs, horses, cows, sheep, or primates.
- 30 - “Sensitives” are subjects that exhibit a response to treatment of a cartilage disorder with an FGF-18 compound. Preferably, sensitive subjects (or subjects showing sensitivity to treatment) exhibit notably a higher increase in total cartilage volume than placebo treated subjects, *i.e.* they show cartilage repair. In addition, sensitive subjects exhibit at least similar improvement in WOMAC total scores than placebos. The terms “good-responders” (or good sensitive), and “low-responders” (including “Low-sensitive”) refer to the different groups of subjects depending notably on the increase of the cartilage volume following specific dosing regimen in the frame of a FGF-18 compound treatment. A good responder displays a positive response (*i.e.* cartilage repair) to treatment with an FGF-18 compound whatever the dosing regimen, whereas low responder display a response only to a specific dosing regimen when treated with an FGF-18 compound. Both groups of responders have similar improvement in WOMAC total score than placebos, should they be treated according to

specific dosing regimen. The good responders can also be simply called responders or sensitive (patients). A good response can also be called response.

The proposed criteria for sensitivity are the following:

1. Positive cartilage increase compared to baseline,
- 5 2. Cartilage increase change significantly higher than change in placebo (e.g. as tested with a linear model adjusting for BMI, KL grade, sex and age and with  $\alpha = 5\%$ ),
3. WOMAC score improvement, i.e. diminution, (e.g. more than 5 points reduction) compared to baseline,
- 10 4. WOMAC score change not significantly higher than change in placebo (e.g. as tested with a linear model adjusting for BMI, KL grade, sex and age and with  $\alpha = 5\%$ )

- The "response", or "sensitivity" to an FGF-18 compound treatment is to be understood as 1 year or 2 years after the first injection and measured as 1) increase of cartilage volume and/or cartilage thickness, measured owing to MRI or X-Ray for instance, 2) decrease of WOMAC total scores, or 3) changes in WOMAC total scores not significantly higher than those from placebos (refer also to the  
15 definition of "sensitive").

- As used in the present invention, the term "MAD" means Multiple Ascending Dose. When this acronym is followed by a figure, the figure corresponds to the dose at which FGF-18 compound has been injected during treatment. For instance MAD100 refers to a treatment during which a subject received 100 mcg of FGF-18 compound per injection. The abbreviation "PL" (and "MADPL") refers  
20 to placebo.

- The term "storage device", as used herein, is intended to include any suitable computing or processing apparatus or other device configured or adapted for storing data or information. Examples of electronic apparatus suitable for use with the present invention include stand-alone computing  
25 apparatus, data telecommunications networks, including local area networks (LAN), wide area networks (WAN), Internet, Intranet, and Extranet, and local and distributed computer processing systems. Storage devices also include, but are not limited to: magnetic storage media, such as floppy discs, hard disc storage media, magnetic tape, optical storage media such as CD-ROM, DVD, electronic storage media such as RAM, ROM, EPROM, EEPROM and the like, general hard disks and hybrids of these categories such as magnetic/optical storage media.

- As used herein, the term "stored" refers to a process for encoding information on the storage device. Those skilled in the art can readily adopt any of the presently known methods for recording  
30 information on known media to generate manufactures comprising expression level information.

### Detailed description of the invention

35 There is a need to predict the clinical efficacy (notably with regards to delaying cartilage thinning and/or cartilage repair) of an FGF-18 compound treatment for the treatment of subjects having a cartilage disorder, such as osteoarthritis, cartilage injury, fractures affecting joint cartilage or surgical procedures with impact on joint cartilage (e.g. Microfracture). To optimize the treatment of said subjects, it is important to identify biomarkers that could be used as predictors of the response of a

given subject to a specific dosage regimen in the frame of a FGF-18 compound treatment, notably with regard to cartilage repair. Such predictive biomarkers may be used to identify groups having risks of being low-sensitives to a given dose regimen. For instance, if one subject having osteoarthritis is known to be at risk for having a low-response (or for being low-sensitive) to the FGF-18 compound treatment, the physician may decide to propose a four cycle- treatment with a FGF-18 compound, such as sprifermin, to said subject. Such predictive information may be clinically useful to guide medical decisions, notably on the dosing regimen to be applied to one patient.

The surprising finding of the present invention is based on a study aimed at identifying potential biomarkers associated with FGF-18 compound administration. The biomarkers used in this study were composed of numerous protein markers. The association between protein markers and clinical response variables was assessed. The rationale behind this type of analysis was to identify protein(s) useful as biomarker(s) that could be predictive of the clinical outcome (notably with regard to cartilage repair), for a subject to be treated with an FGF-18 compound such as sprifermin or a fusion protein comprising an FGF-18 moiety. The proteins could be used to stratify and target specific subject populations.

The inventors have surprisingly found an association with certain proteins and outcome (e.g. cartilage repair). Of special interest are the proteins C1M, C3M and/or CRPM. It is noted that although only C1M, C3M and/or CRPM are specifically described herein, other inflammatory biomarker such as high-sensitivity C-reactive protein (hs-CRP) could be used. Based on the teaching of the present invention, it would then be routine matter for the skilled person to find the thresholds for each of these biomarkers.

These proteins have been described in the literature, as being possibly related to inflammation in arthritis. For instance, C1M, C3M and/or CRPM could be considered as inflammatory biomarkers, specific for tissue inflammation. The results presented herein show that high levels of tissue inflammation abolish the therapeutic effect of an FGF-18 compound (such as sprifermin), especially at lower doses/dosing regimens. This abolishing effect can be surprisingly counteract with an FGF-18 compound (such as sprifermin) administered at a particular dosing regimen.

It has also been surprisingly found by the present inventors that when the quantity of inflammatory biomarkers such as C1M, C3M and/or CRPM is/are decreased, this is associated with a better response to treatment with a FGF-18 compound, such as sprifermin or a fusion protein comprising an FGF-18 moiety, whatever the dosing regimen in subjects afflicted with cartilage injury. These subjects are called good-sensitives or good-responders. To the opposite, it has also surprisingly been found by the present inventors that when the quantity of C1M, C3M and/or CRPM is/are increased, this is associated with a lower response to treatment with a FGF-18 compound (i.e. low-sensitivity to treatment with a FGF-18 compound), such as sprifermin or a fusion protein comprising an FGF-18 moiety, in subjects afflicted with cartilage disorder, except at a specific dosing regimen comprising four cycles of treatment. These subjects are called low-sensitives or low-responders. It was further surprisingly been found that each one of these biomarkers can be used alone to provide an efficient prediction of the response to FGF-18.

Therefore, it is a finding of the present invention that the biomarkers C1M, C3M and/or CRPM can be used either alone or in combination as predictive biomarkers of responsiveness of one subject to specific dosing regimen in the frame of a FGF-18 compound treatment, such as sprifermin or a fusion protein comprising an FGF-18 moiety. Preferably, the subject has a cartilage disorder, such as osteoarthritis, cartilage injury, fractures affecting joint cartilage or surgical procedures with impact on joint cartilage (e.g. Microfracture). In a particular embodiment, the subject will be predicted to be low-sensitive to FGF-18 compound treatment if the quantity of C1M is higher than  $30.0 \pm 2SD$  ng/mL and/or if the quantity of C3M is higher than  $12.0 \pm 2SD$  ng/mL and/or if the quantity of CRPM is higher than  $10.0 \pm 2SD$  ng/mL. On the contrary, the subject will be predicted to be sensitive (or responder or good responder) to FGF-18 compound treatment if the quantity of C1M is lower than  $30.0 \pm 2SD$  ng/mL and/or the quantity of C3M is lower than  $12.0 \pm 2SD$  ng/mL and/or the quantity of CRPM is lower than  $10.0 \pm 2SD$  ng/mL.

The present invention is therefore directed to a method of predicting the sensitivity to treatment with an FGF-18 compound in a subject having a cartilage disorder, the method comprising the steps of:

- a) Determining, from a biological sample of said subject, the quantity of at least one of the biomarkers selected from the group consisting of C1M, C3M and/or CRPM; and
- b) Predicting from the result of step a) good or low sensitivity of said subject to treatment with an FGF-18 compound.

Before determining the quantity of at least one of the biomarkers, it is needed to obtain a sample (or biological sample or test sample) of said subject, for instance by blood, serum, synovial fluid or urine collecting. Thus, the present invention is directed to a method of predicting the sensitivity to treatment with an FGF-18 compound in a subject having a cartilage disorder, the method comprising the steps of:

- a) Obtaining a sample from said subject;
- b) Determining, from a sample, the quantity of at least one of the biomarkers selected from the group consisting of C1M, C3M and/or CRPM; and
- c) Predicting from the result of step a) good or low sensitivity of said subject to treatment with an FGF-18 compound

According to said method, the presence of lower than  $30.0 \pm 2SD$  ng/mL of C1M and/or lower than  $12.0 \pm 2SD$  ng/mL of C3M and/or lower than  $10.0 \pm 2SD$  ng/mL of CRPM is predictive of good response (or response or sensitivity) to treatment with an FGF-18 compound. FGF-18 compound could thus be administered to said subject at the following dosing regimen at least two treatment cycles of 100  $\mu$ g of FGF-18 compound given once per week for 3 consecutive weeks per treatment cycle. On the contrary, the presence of higher than  $30.0 \pm 2SD$  ng/mL of C1M and/or higher than  $12.0 \pm 2SD$  ng/mL of C3M and/or higher than  $10.0 \pm 2SD$  ng/mL of CRPM is predictive of low response (i.e. low sensitivity) to treatment with an FGF-18 compound. FGF-18 compound could thus be administered to said subject preferably at the following dosing regimen: four treatment cycles of 100  $\mu$ g of FGF-18 compound given once per week for 3 consecutive weeks per treatment cycle.

The present application also encompasses a method for selecting a subject having a cartilage disorder for inclusion in or exclusion from a dosing regimen when said subject is to be treated with an FGF-18 compound, based on the likelihood of their sensitivity to said treatment, comprising the steps of:

- 5           a. Determining, from a sample, the quantity of at least one of the biomarkers selected from the group consisting of C1M, C3M and/or CRPM; wherein the quantity of at least one of these proteins is predictive about the subject's risk of having good-sensitivity or low-sensitivity to said treatment, and
- 10           b. Selecting the appropriate dosing regimen according to the subject's risk of having good-sensitivity or low-sensitivity to said treatment.

Before determining the quantity of at least one of the proteins, in the above disclosed assays, it is needed to obtain a sample (or test sample or biological sample) of said subject, for instance by blood, serum, synovial fluid or urine collecting. Thus, the present invention is directed to a method for selecting a subject having a cartilage disorder for inclusion in or exclusion from dosing regimen when

15 said subject is to be treated with an FGF-18 compound, based on the likelihood of their sensitivity to said treatment, comprising the steps of:

- a. Obtaining a biological sample from said subject,
- b. Determining, from a sample, the quantity of at least one of the biomarkers selected from the group consisting of C1M, C3M and/or CRPM; wherein the quantity of at least one of
- 20 these proteins is predictive about the subject's risk of having good-sensitivity or low-sensitivity to said treatment, and
- c. Selecting the appropriate dosing regimen according to the subject's risk of having good-sensitivity or low-sensitivity to said treatment.

According to said method, the subject presenting higher than  $30.0 \pm 2SD$  ng/mL of C1M and/or higher

25 than  $12.0 \pm 2SD$  ng/mL of C3M and/or higher than  $10.0 \pm 2SD$  ng/mL of CRPM will preferably be excluded (i.e. they will not be selected) from the treatment with FGF-18 compound given at the following dosing regimen: two treatment cycles of 100  $\mu$ g of FGF-18 compound given once per week for 3 consecutive weeks per treatment cycle. Said subject would preferably be proposed the following dosing regimen: four treatment cycles of 100  $\mu$ g/mL of FGF-18 compound (such as sprifermin) given

30 once per week for 3 consecutive weeks per treatment cycle. To the contrary, the subject presenting lower than  $30.0 \pm 2SD$  ng/mL of C1M and/or lower than  $12.0 \pm 2SD$  ng/mL of C3M and/or lower than  $10.0 \pm 2SD$  ng/mL of will be included (i.e. they will be selected) for the treatment with FGF-18 compound given at the following dosing regimen: at least two treatment cycles of 100  $\mu$ g of FGF-18 compound given once per week for 3 consecutive weeks per treatment cycle.

35 The method for selecting a human subject for a clinical trial for testing FGF-18 compound, may alternatively comprises the steps of:

- a) assaying a biological sample from said human subject diagnosed with a cartilage disorder for the quantity of at least one of C1M, C3M and/or CRPM,
- b) determining the quantity of the at least one of C1M, C3M and/or CRPM; and

c) selecting:

- i. for the clinical trial at the following dosing regimens: at least two treatment cycles of 100 µg of FGF-18 compound given once per week for 3 consecutive weeks per treatment cycle the human subject who presents lower than  $30.0 \pm 2SD$  ng/mL of C1M and/or lower than  $12.0 \pm 2SD$  ng/mL of C3M and/or lower than  $10.0 \pm 2SD$  ng/mL of CRPM, or
- ii. for the clinical trial at the following dosing regimens: four treatment cycles of 100 µg of FGF-18 compound given once per week for 3 consecutive weeks per treatment cycle the human subject who presents higher than  $30.0 \pm 2SD$  ng/mL of C1M and/or higher than  $12.0 \pm 2SD$  ng/mL of C3M and/or higher than  $10.0 \pm 2SD$  ng/mL of CRPM.

10 The present invention also describes a method of excluding a human subject from a clinical trial testing FGF-18 compound at the following dosing regimens: two treatment cycles of 100 µg of FGF-18 compound given once per week for 3 consecutive weeks per treatment cycle, the method comprising the steps of: (a) assaying a biological sample from said human subject diagnosed with a cartilage disorder for the quantity of at least one of C1M, C3M and/or CRPM; (b) determining from  
15 said assay the likelihood of the subject of low sensitive to said treatment and (c) excluding for the clinical trial the human subject who presents higher than  $30.0 \pm 2SD$  ng/mL of C1M and/or higher than  $12.0 \pm 2SD$  ng/mL of C3M and/or higher than  $10.0 \pm 2SD$  ng/mL of CRPM.

The quantity (or the expression level) of one or more biomarkers in a sample can be compared, if needed, to a reference quantity (or reference expression level) from a reference sample. Said  
20 reference level can be obtained from a healthy subject, or from the very same patient to be diagnosed or treated prior to or during said treatment.

FGF-18 compound is to be usually administered intraarticularly at a dose of 100 mcg per injection, once weekly for 3 weeks per treatment cycle. FGF-18 compound is to be usually administered for at least one treatment cycle of treatment. Preferably, said treatment cycle is repeated at least once, for  
25 instance 6 months (or about 26 weeks) after the start of the first treatment cycle. Up to four treatment cycles within 2 years have shown promising results (see Figure 1).

The present invention further encompasses an FGF-18 compound for use in the treatment of a subject having a cartilage disorder, wherein the FGF-18 compound is administered at the following dosing regimen: at least two treatment cycles of 100 µg of FGF-18 compound given once per week  
30 for 3 consecutive weeks per treatment cycle, characterized in that the subject has lower than  $30.0 \pm 2SD$  ng/mL of C1M and/or lower than  $12.0 \pm 2SD$  ng/mL of C3M and/or lower than  $10.0 \pm 2SD$  ng/mL of CRPM. A subject who does not meet these criteria is preferably administered FGF-18 compound according to the following dose regime: four treatment cycles of 100 µg of FGF-18 compound given once per week for 3 consecutive weeks per treatment cycle.

35 The present invention also relates to an assay to determine sensitivity to an FGF-18 compound treatment or to determine a treatment regimen with an FGF-18 compound, the assay comprising: (a) subjecting a test sample from a human subject, diagnosed as having a cartilage disorder, to at least one assay that determines the quantity of at least one of C1M, C3M and/or CRPM, (b) determining the quantity of at least one of C1M, C3M and/or CRPM and (c) determining from the result of step b)

good-sensitivity or low-sensitivity of said subject to treatment with an FGF-18 compound. According to said assay, the presence of higher than  $30.0 \pm 2SD$  ng/mL of C1M and/or higher than  $12.0 \pm 2SD$  ng/mL of C3M and/or higher than  $10.0 \pm 2SD$  ng/mL of CRPM is predictive of low-sensitivity to treatment with an FGF-18 compound, requiring preferably a treatment comprising four-cycles of FGF-18 treatment. On the contrary, the presence of lower than  $30.0 \pm 2SD$  ng/mL of C1M and/or lower than  $12.0 \pm 2SD$  ng/mL of C3M and/or lower than  $10.0 \pm 2SD$  ng/mL of CRPM is predictive of good sensitivity (or of sensitivity).

Before determining the quantity of at least one of the biomarkers, in the above disclosed assays, it is needed to obtain a biomarker (or test) sample of said subject, for instance by blood, serum, synovial fluid or urine collecting.

The present invention is also directed to an assay for selecting a treatment regimen for a human subject with a cartilage disorder, the assay comprising: (a) subjecting a test sample from a human subject, diagnosed as having a cartilage disorder, to at least one assay that determines the quantity of at least one of C1M, C3M and/or CRPM, (b) determining the likelihood that said subject is intermediate-sensitive or a high-sensitive to FGF-18 treatment, and (c) determining from the result of step b) the appropriate treatment regimen for said subject. Once the assay is performed, said subject can be selected for and treated with an appropriate dosing regimen consisting of 1) at least two treatment cycles of 100  $\mu$ g of FGF-18 compound given once per week for 3 consecutive weeks per treatment cycle when the subject has lower than  $30.0 \pm 2SD$  ng/mL of C1M and/or lower than  $12.0 \pm 2SD$  ng/mL of C3M and/or lower than  $10.0 \pm 2SD$  ng/mL of CRPM based on the recognition that said quantities are associated with a response to said compound (i.e. the subject is intermediate-sensitive or a high-sensitive to FGF-18 treatment), and excluding the subject from treatment with an FGF-18 compound according to said dosing regimen when the subject has higher than  $30.0 \pm 2SD$  ng/mL of C1M and/or higher than 11.7 ng/mL of C3M and/or higher than 10 ng/mL of CRPM based on the recognition that said quantities are associated with inadequate response to treatment with said compound.

Also described is a method for treating a subject having a cartilage disorder, comprising the following steps:

- a. Determining, from a sample, the quantity of at least one of the biomarkers selected from the group consisting of C1M, C3M and/or CRPM, wherein said quantities are predictive about the subject's risk of having good or low sensitivity to a treatment with said FGF-18 compound,
- b. Selecting the appropriate dosing regimen according to the subject's risk of having good-sensitivity or low-sensitivity to said treatment, and
- c. Administering intraarticularly an FGF-18 compound according to the dosing regimen selected in step b).

Further discloses is a method for treating a human subject with a cartilage disorder, comprising the steps of: (a) assaying a biological sample of a subject, who is diagnosed as having the cartilage disorder for the quantity of at least one of the biomarkers selected from the group consisting of C1M,

C3M and/or CRPM; and (b) administering a treatment regimen consisting of at least two treatment cycles of 100 µg of FGF-18 compound given once per week for 3 consecutive weeks per treatment cycle when the subject has lower than  $30.0 \pm 2SD$  ng/mL of C1M and/or lower than  $12.0 \pm 2SD$  ng/mL of C3M and/or lower than  $10.0 \pm 2SD$  ng/mL of CRPM.

5 In the context of the present invention as a whole, C1M, C3M and/or CRPM have been shown to be useful predictive biomarkers of the response to an FGF-18 compound. As such they are considered as predictive biomarkers.

In the context of the present invention as a whole, the assays or other determination of the quantity of at least one of the biomarkers of the invention can be performed before treatment or during  
10 treatment. Indeed, also during treatment the dosing regimen may have to be adapted to the new biomarkers situation.

In the context of the invention as a whole, for the patients being in the range of  $30 \pm 2SD$  ng/mL of C1M,  $12.0 \pm 2SD$  ng/mL of C3M or  $10.0 \pm 2SD$  ng/mL of CRPM, it is advisable to complete the diagnostic or the biomarker testing with another inflammatory biomarker. For instance, should the  
15 level of C1M be in the range of  $30 \pm 2SD$  ng/mL, then the level(s) of C3M and/or CRPM could be considered. In the rare case where the patient will present ranges of  $30 \pm 2SD$  ng/mL of C1M and  $12.0 \pm 2SD$  ng/mL of C3M and  $10.0 \pm 2SD$  ng/mL of CRPM, then it will be advisable to complete the diagnostic or the biomarker testing with another kind of biomarkers such as a metabolic biomarker or SNP biomarkers such as those disclosed in WO2014023703.

20 In the context of the present invention as a whole, a dosing regimen consisting of at least two treatment cycles of 100 µg of FGF-18 compound given once per week for 3 consecutive weeks per treatment cycle means any one of: 1) two treatment cycles of 100 µg of FGF-18 compound given once per week for 3 consecutive weeks per treatment cycle, 2) three treatment cycles of 100 µg of FGF-18 compound given once per week for 3 consecutive weeks per treatment cycle, or 3) four  
25 treatment cycles of 100 µg of FGF-18 compound given once per week for 3 consecutive weeks per treatment cycle.

In another embodiments of the invention, also provided are systems (and computer readable media for causing computer systems) for obtaining data. Said data can be used notably for assessing suitability of a treatment with FGF- compound in a subject, or monitoring FGF-18 compound  
30 treatment efficacy for a given subject or simply monitor disease progression. Said systems can be used during clinical trials, when a treatment with FGF-18 compound has to be envisaged or when a treatment with said compound is already ongoing.

Therefore, in an embodiment of the present invention is included a computer system for obtaining data from at least one test sample obtained from at least one subject with a cartilage disorder, the  
35 system comprising: (a) at least one determination module configured to receive said at least one test sample and perform at least one analysis on said at least one test sample to determine the quantity of at least one of the biomarkers according to the invention; (b) at least one storage device configured to store data output from said determination module; and (c) at least one display module for displaying a content based in part on the data output from said determination module, wherein the



content comprises a signal indicative of the presence of at least one of these conditions, and optionally the absence of any one of these conditions.

The computer readable medium can have computer readable instructions recorded thereon to define software modules for implementing a method on a computer. In such a case, said computer readable storage medium may comprise: (a) instructions for comparing the data stored on a storage device with reference data to provide a comparison result, wherein the comparison is based on the quantity of at least one of the biomarkers according to the invention; and (b) instructions for displaying a content based in part on the data output from said determination module, wherein the content comprises a signal indicative of the presence of at least one of the conditions, and optionally the absence of one or more of the conditions.

The computer readable storage media can be any available tangible media that can be accessed by a computer. Computer readable storage media includes volatile and non-volatile, removable and non-removable tangible media implemented in any method or technology for storage of information such as computer readable instructions, data structures, program modules or other data. Computer readable storage media includes, but is not limited to, RAM (random access memory), ROM (read only memory), EPROM (erasable programmable read only memory), EEPROM (electrically erasable programmable read only memory), flash memory or other memory technology, CD-ROM (compact disc read only memory), DVDs (digital versatile disks) or other optical storage media, magnetic cassettes, magnetic tape, magnetic disk storage or other magnetic storage media, other types of volatile and non-volatile memory, and any other tangible medium which can be used to store the desired information and which can be accessed by a computer including any suitable combination of the foregoing.

Computer-readable data embodied on one or more computer-readable media may define instructions, for example, as part of one or more programs that, as a result of being executed by a computer, instruct the computer to perform one or more of the functions described herein, and/or various embodiments, variations and combinations thereof. Such instructions may be written in any of a plurality of programming languages, for example, Java, J#, Visual Basic, C, C#, C++, Fortran, Pascal, Eiffel, Basic, COBOL assembly language, and the like, or any of a variety of combinations thereof. The computer-readable media on which such instructions are embodied may reside on one or more of the components of either of a system, or a computer readable storage medium described herein, may be distributed across one or more of such components.

The computer-readable media may be transportable such that the instructions stored thereon can be loaded onto any computer resource to implement the aspects of the present invention discussed herein.

The information determined in the determination module can be read by the storage device. The storage device is adapted or configured for having recorded thereon expression level or protein level information. Such information may be provided in digital form that can be transmitted and read electronically, e.g., via the Internet, on diskette, via USB (universal serial bus) or via any other suitable mode of communication.

In the context of the present invention as a whole, e.g... in the context of any one of the methods, uses, assays or kits according to the present invention, the preferred FGF-18 compound is a truncated FGF-18, such as sprifermin, or a fusion protein comprising an FGF-18 moiety, and the preferred cartilage disorder is selected from the group consisting of osteoarthritis, cartilage injury, fractures affecting joint cartilage or surgical procedures with impact on joint cartilage, such as microfracture.

It is to be understood that in the context of the present invention as a whole, e.g. of any one of the methods, uses, assays, computer system or kits according to the present invention, before determining the quantity of at least one of the biomarkers of the invention (i.e. C1M, C3M and/or CRPM), it is needed to obtain a sample (or biological sample or a test sample) of one subject, for instance by blood, serum, synovial fluid or urine collecting. It can also be obtained, without any limitation, from a cell, tissue, cartilage or synovial fluid.

An individual afflicted with a cartilage disorder and to be tested, tested and/or treated according to any of the methods, uses, assays, kits and other computer systems described herein is a human subject that is a candidate for treatment with an FGF-18 compound, such as sprifermin or a fusion protein comprising an FGF-18 moiety. In a preferred embodiment, the individual has been diagnosed with cartilage disorder, or exhibits a symptom of cartilage disorder.

In a further embodiment, the present invention encompasses a kit comprising means for performing the methods described above and instructions for use. Preferably, the kit comprises means for detecting the presence of at least one of the biomarkers according to the invention (e.g. C1M, C3M and/or CRPM) and for quantifying them. The kit may comprises means for detecting the presence of at least two of the biomarkers according to the invention and for quantifying them.

The methods and kits according to the present invention are useful in clinical diagnostic applications. However, as used herein, the term "diagnostic" is not limited to clinical or medical uses, and the diagnostic methods and kits of the invention claimed herein are also useful in any research application, and during clinical trials, for which it is desirable to test a subject for the presence or absence of any markers described herein.

In the context of the invention, the presence of at least one of the biomarkers according to the invention (e.g. C1M, C3M and/or CRPM) and their quantitation may be detected by any technique known per se to the skilled artisan, including ELISA for instance.

Other embodiments of the invention within the scope of the claims herein will be apparent to one skilled in the art from consideration of the specification or practice of the invention as disclosed herein. It is intended that the specification, together with the examples, be considered exemplary only, with the scope and spirit of the invention being indicated by the claims that follow the examples.

#### **Description of the Figures:**

**Figure 1:** Scheme of the dosing regimens used for sprifermin in the FORWARD study.

**Figure 2:** Mean and 95% CI in Absolute Change from Baseline Over Weeks by Treatment, Cartilage Thickness (mm) in the Total Femorotibial Joint – mITT Analysis Set.

**Figure 3:** Mean and 95% CI in Absolute Change from Baseline Over Weeks by Treatment, Cartilage Volume ( $\mu\text{L}$ ) in the Total Femorotibial Joint – mITT Analysis Set.

**Figure 4:** Mean and 95% CI for Change from Baseline Over Weeks by Treatment and Biochemical Marker of Inflammation, Cartilage Thickness (mm) in the Total Femorotibial Joint – ITT Analysis Set – C1M biomarker.

**Figure 5:** Mean and 95% CI for Change from Baseline Over Weeks by Treatment and Biochemical Marker of Inflammation, Cartilage Thickness (mm) in the Total Femorotibial Joint – ITT Analysis Set – C3M biomarker.

**Figure 6:** Mean and 95% CI for Change from Baseline Over Weeks by Treatment and Biochemical Marker of Inflammation, Cartilage Thickness (mm) in the Total Femorotibial Joint – ITT Analysis Set – CRPM biomarker.

**Figure 7:** Mean and 95% CI for Change from Baseline Over Weeks by Treatment and Biochemical Marker of Inflammation, Cartilage Thickness (mm) in the Total Femorotibial Joint – ITT Analysis Set – C1M and C3M biomarker.

**Figure 8:** Mean and 95% CI in Absolute Change from Baseline over Weeks by Treatment and Biochemical Marker of Inflammation Subgroups, WOMAC Total Score in the Target Knee – ITT Analysis Set – C1M biomarker.

**Figure 9:** Mean and 95% CI in Absolute Change from Baseline over Weeks by Treatment and Biochemical Marker of Inflammation Subgroups, WOMAC Total Score in the Target Knee – ITT Analysis Set – C3M biomarker.

**Figure 10:** Mean and 95% CI in Absolute Change from Baseline over Weeks by Treatment and Biochemical Marker of Inflammation Subgroups, WOMAC Total Score in the Target Knee – ITT Analysis Set – C1M and C3M biomarker.

## Description of the sequences:

SEQ ID NO.1: Amino acid sequence of the native human FGF-18.

SEQ ID NO.2: Amino acid sequence of the recombinant truncated FGF-18 (trFGF-18).

SEQ ID NO.3: Amino acid sequence of the marker C1M.

SEQ ID NO.4: Amino acid sequence of the marker C3M.

SEQ ID NO.5: Amino acid sequence of the marker CRPM.

## Examples

### 1. Materials and methods

#### 1.1. FGF-18 compound

The FGF-18 compound used as a treatment in the present examples is sprifermin. It is a truncated form of FGF-18, as defined in the section “definitions”. Two strengths of sprifermin were supplied for the study: 30  $\mu\text{g}$  and 100  $\mu\text{g}$ . Sprifermin was supplied as a white, sterile, freeze-dried powder in 3-mL glass vials. Each vial contained either 31.5  $\mu\text{g}$  or 105  $\mu\text{g}$  of sprifermin active substance; these quantities included a 5% overage, permitting extraction of respectively 30  $\mu\text{g}$  or 100  $\mu\text{g}$  of sprifermin

active substance following reconstitution with 0.9% w/v Sodium Chloride Injection (referred to herein as “saline solution”). Excipients of the formulation were sodium phosphate buffer (pH 7.2), sodium hydroxide, O-phosphoric acid, sucrose, and poloxamer 188. Kits for 30 µg treatments contained 1 glass vial of sprifermin (30 µg strength) and 1 glass ampule of sterile saline solution for injection (2 mL per ampule). Kits for 100 µg treatments contained 1 glass vial of sprifermin (100 µg strength) and 1 glass ampule of sterile saline solution for injection (2 mL per ampule). For all treatment groups, the volume administered was 2 mL.

## 1.2. Methods

### 10 FORWARD study

The present study was based on the FORWARD study (EMR700692-006). Five groups of patients were studied:

- Group 1 (4 cycles placebo; hereafter referred to as placebo): 108 subjects.
- Group 2 (2 cycles sprifermin 30 µg/injection alternating with 2 cycles placebo; hereafter referred to as sprifermin/placebo 30 µg): 110 subjects.
- 15 • Group 3 (4 cycles sprifermin 30 µg/injection; hereafter referred to as sprifermin 30 µg): 111 subjects.
- Group 4 (2 cycles sprifermin 100 µg/injection alternating with 2 cycles of placebo; hereafter referred to as sprifermin/placebo 100 µg): 110 subjects.
- 20 • Group 5 (4 cycles sprifermin 100 µg/injection; hereafter referred to as sprifermin 100 µg): 110 subjects.

According to the FORWARD study, the patients received 4 cycles of treatment (each consisting of 3 once-weekly intra articular injections over 3 consecutive weeks) at intervals of 6 months (see Figure 1). Injections were all done intraarticular (i.art.).

25 The primary efficacy endpoint was the change from Baseline in cartilage thickness in the total femorotibial joint as evaluated by MRI at 2 years.

Exploratory endpoints included Baseline protein markers and/or genetic markers associated with response to treatment or disease progression (response assessed by MRI and/or questionnaire).

### 30 Inclusion/exclusion criteria

The study enrolled adult subjects of either sex with primary femorotibial OA according to American College of Rheumatology (ACR) clinical and radiographic criteria who had Kellgren-Lawrence grades (KLG) of 2 or 3 and a minimum joint space width (JSW) of  $\geq 2.5$  mm in the medial compartment. Subjects must have had pain in the target knee on most days and/or require symptomatic treatment of knee pain with paracetamol (acetaminophen), systemic non-steroidal anti-inflammatory drugs (NSAIDs) including COX inhibitors (COXibs), or tramadol on most days of the previous month, and must have had both: 1) A history of pain due to OA in the target knee for at least 6 months, and 2) Pain score for the target knee of 4 to 9 points in response to Question 1 of the Western Ontario and McMaster Universities Osteoarthritis Index (WOMAC) pain index (“how much pain have you had [in

the target knee, over the past 48 hours] when walking on a flat surface?") at screening and Baseline, after washout of at least 5 half-lives of analgesic medication(s): acetaminophen, topical or oral systemic NSAIDS, COXibs, opioids, and/or tramadol. Women of childbearing potential must have used a form of contraception with a failure rate of less than 1% per year throughout the study.

- 5 Main exclusion criteria included malalignment of > 5 degrees in the femorotibial axis of the target knee, clinical signs of inflammation (i.e., redness) in the target knee, i.art. administration of corticosteroids or hyaluronic acid into either knee within 6 months before screening, any plan for knee surgery (affecting either the target or the contralateral knee) within the next 2 years, concomitant conditions or treatments deemed to be incompatible with study participation, 10 contraindications to MRI scanning (including inability to fit in the scanner or knee coil), pregnancy or breastfeeding, participation in another clinical study within the past 30 days, and legal incapacity or limited legal capacity.

Written informed consent must have been obtained prior to any study-related activity.

15 Statistical methods

The treatment effect on the primary endpoint was assessed through dose-ranging using a repeated measurement analysis of variance (ANOVA, using PROC MIXED in SAS) on absolute change from Baseline, including the baseline value, the treatment group, the time, and the country as factors and treatment-by-time point as interaction. The primary efficacy analysis consisted of testing the linear 20 dose relationship and the overall treatment effect at 2 years. The significance level was set at 5% 2-sided for both tests. Pairwise comparisons (sprifermin versus placebo, and between sprifermin dose and regimen groups) were performed within the context of this modelling framework. For each pairwise comparison, the difference between treatments and the corresponding 95% confidence interval (CI) and p-value are presented. The same ANOVA model used for the primary endpoint was 25 used to assess the treatment effect on continuous secondary endpoints such as MRI endpoints, WOMAC endpoints (total, pain, function, and stiffness scores), and X-ray endpoints at each time point and over time. Logistic regression was used to assess the treatment effect on the binary efficacy endpoints such as the OMERACT-OARSI responder rate. Point estimates for each pairwise comparison and corresponding 95% CIs and p-values are provided.

30

Pain and function assessments

The WOMAC is a validated instrument used to assess symptom modification in clinical OA studies. This clinical score was developed in 1981 and is regarded as a valid instrument by both clinical researchers and regulatory authorities. The WOMAC is widely used in clinical studies in hip and knee 35 OA, and has been extensively validated.

Subjects had to answer all of the 24 questions themselves (i.e. 5 for pain, 2 for stiffness and 17 for physical function assessment), using the 11-box NRS assessment (with categories of 0 to 10) with reference to the past 48 hours. Different forms of the questionnaire exist for the right and the left knees: in order to reduce confounding of WOMAC responses by symptoms in the contralateral knee,

subjects used the WOMAC questionnaire specific to the target knee. For administration of the questionnaire, instructions for the WOMAC 3.1 Index were followed.

#### X-Ray Assessment of JSW

- 5 Change in JSW as measured by X-ray is a recognized endpoint accepted by the European Medicines Agency and the United States Food and Drug Administration for use in efficacy studies in OA. The JSW was measured using standardized technique.

#### qMRI Assessment

- 10 The primary endpoint for the DBPC treatment phase was the change from Baseline in cartilage thickness in the total femorotibial joint as evaluated by qMRI at 2 years in the mITT. Cartilage thickness of the total femorotibial joint were calculated in 2 ways:

1. Average Cartilage Thickness (Total Volume divided by Total Surface Area)
2. Total Cartilage Thickness (sum of cartilage thickness in medial and lateral compartment).

- 15 The treatment effect on the primary endpoint was assessed through dose-ranging using a repeated measurement analysis of variance (ANOVA) on absolute change from Baseline, including the treatment group, the time point, and the (pooled) country as fixed factors and the baseline value as covariate and treatment by time point as interaction. Repeated measures over time were accounted for using an “unstructured” covariance pattern.

- 20 Pairwise comparisons of absolute change from Baseline in cartilage thickness (sprifermin treatment groups versus placebo) were performed within the context of the modelling framework described above. For each pairwise comparison, the difference between treatments and the corresponding 95% confidence interval (CI) and p-value are presented. P-values (corresponding to Type 3 tests of fixed effects) are reported for all covariates in the original “Overall” model for all time points combined  
25 (i.e., baseline value, treatment, time point, treatment-by-time point interaction, country) and for all time points. Estimated coefficients, p-values, and 95% CIs are presented overall and at each time point for (i) the dose relationship (linear trend) and (ii) each pairwise comparison between dose level and placebo.

- 30 In order to assess the robustness of the primary results, the tests for linear dose-relationship and for the overall treatment effect were repeated using the PP Analysis Set. For the mITT Analysis Set, a non-parametric analysis was conducted for the ordered data of cartilage thickness in the total femorotibial joint as an alternative method for the primary analysis. Data were ordered by the magnitude of absolute change-from-Baseline over 2 years during DBPC treatment phase using rank transformation.

35

#### Biomarkers measurement

Serological and urine biochemical markers of bone and joint tissue turnover as well as synovial inflammation were evaluated. Potential biomarkers of synovial inflammation included, but were not limited to: neo-epitope of collagen type I degradation (C1M), neo-epitope of collagen type III

degradation (C3M), and neo-epitope of synovial C-reactive protein degradation (CRPM). Blood and urine samples for systemic biomarker assessment were collected at the following time points: week 0 (before first injection of sprifermin), week 26, week 54, week 80 and week 104. For time points where injections were also administered, samples were collected before injection. Synovial fluid samples were collected at the time points. These samples were taken just before injection, as part of the i.art. injection procedure and using the same needle that the one used for the injection. For urine collection, second morning void samples were obtained.

The following assessment were made as exploratory endpoints:

- Change from Baseline in serum and urine markers associated with administration of the compound.
- Baseline protein markers and/or genetic markers associated with response to treatment or disease progression (response assessed by MRI and/or questionnaire).

## 2. Results

### Primary Endpoint (allcomer)

Total femorotibial joint: A statistically significant treatment effect on the change from Baseline in cartilage thickness in the total femorotibial joint at 2 years was observed (See figure 2). Both the sprifermin/placebo 100 µg and sprifermin 100 µg groups showed greater improvements (mean change from Baseline: +0.02 mm and +0.03 mm, respectively) compared with the placebo group (mean change from Baseline: -0.02 mm) at Week 104 ( $p < 0.001$  for both comparisons). The treatment group difference started at Week 78 in the sprifermin/placebo 100 µg group and Week 52 in the sprifermin 100 µg group. Statistical significance was maintained through Week 104 in both groups. The placebo group showed no improvement from Baseline at any visit through Week 104. There was a statistically significant difference overall (all weeks) for increased cartilage thickness in the sprifermin/placebo 100 µg and sprifermin 100 µg groups, compared with the placebo group ( $p = 0.002$  and  $p < 0.001$ , respectively). The ANCOVA model was statistically significant for treatment ( $p < 0.001$ ), week ( $p < 0.001$ ), treatment\*week ( $p = 0.029$ ), and pooled country ( $p = 0.009$ ).

Cartilage volume in the total femorotibial joint: There was a statistically significant effect of treatment on the change from Baseline in cartilage volume in the total femorotibial joint (Figure 3;  $p < 0.001$ ). The sprifermin/placebo 100 µg and sprifermin 100 µg groups showed greater improvements compared with the placebo group. The statistical significance of sprifermin 100 µg compared with placebo began at Week 78 and persisted through Week 104; the statistical significance of sprifermin/placebo 100 µg compared with placebo appeared at Week 26 and persisted from Week 78 through Week 104. The mean changes from Baseline to Week 104 were -28.2 µL, +9.5 µL, +96.6 µL ( $p < 0.001$ ), and +116.5 µL ( $p < 0.001$ ) in the sprifermin/placebo 30 µg, sprifermin 30 µg, sprifermin/placebo 100 µg, and sprifermin 100 µg groups, respectively, and -55.5 µL in the placebo group. There was a statistically significant effect of pooled country ( $p = 0.011$ ).

Exploratory Endpoints (biomarkers stratification): The overall objectives of the pharmacodynamic (PD)/biomarker analysis were:

- 5 - To identify predictive biomarker(s) for identification of patients who retain a positive structural outcome (based on differences in MRI total cartilage thickness with sprifermin versus placebo) while improving symptom outcomes (WOMAC total score and WOMAC pain index score) versus placebo.
- To identify predictive biomarker(s) for safety parameters such as AIRs
- To characterize functionality biomarkers as potential predictive biomarkers and evaluate potential predictive cut-offs
- 10 - To identify potential prognostic biomarkers (in the placebo group only)
- To evaluate biochemical biomarkers as PD biomarkers (e.g., C1M, C3M, CPRM).

Stratification and cartilage thickness in the total femorotibial joint: Changes from Baseline in cartilage thickness in the total femorotibial joint showed the following notable differences between biomarker subgroups at Week 104 (see Figures 4-6). Subjects with high levels of inflammatory biomarkers (CRPM, C1M, and C3M) at Baseline did not exhibit a therapeutic effect of sprifermin on total cartilage thickness, except in the sprifermin 100 µg treatment group, four cycles of treatment. Subjects with low levels of inflammatory markers C1M and C3M at Baseline showed a similar improvement in total cartilage thickness compared with allcomers. However, there was a non negligible improvement among subjects with low CRPM levels at Baseline when treated with sprifermin 100 µg treatment groups, either two or four cycles of treatment. These results suggest that high levels of tissue inflammation abolish the therapeutic effect of sprifermin, especially at lower doses. This abolition can surprisingly be counteracted by a more intense treatment with sprifermin, i.e. 100 µg treatment group, four cycles of treatment. This counteraction effect is even stronger when both C1M and C3M biomarkers are used (i.e. a combination of the biomarkers; Figure 7). Same trend is shown using both C3M/CRPM or C1M/CRPM (i.e. a combination of these biomarkers; data not shown).

Stratification and WOMAC total scores: Changes from Baseline in WOMAC total scores showed the following notable differences between biomarker subgroups at Week 104. Subjects with lower levels of CRPM biomarker at Baseline globally showed improved outcomes after 104 weeks of treatment with sprifermin (whatever the dosing regimen) for WOMAC total score compared with subjects receiving placebo or with higher levels of biomarkers of cartilage inflammation (data not shown). The results surprisingly show that joints with lower inflammation (low CRPM) not only have a better response to anabolic therapy such as sprifermin with regard to cartilage thickness but also have a positive impact in the WOMAC total scores. In addition, the lower doses, i.e. 2 times 100µg of FGF-18, or 2 times 30 µg of FGF-18 or 4 times 30 µg of FGF-18 showed no therapeutic efficacy in subjects with high CRPM

To the contrary subjects with higher levels of C1M or C3M biomarkers at Baseline globally showed similar outcomes after 104 weeks of treatment with sprifermin (whatever the dosing regimen) for



WOMAC total score compared with subjects receiving placebo (Figures 8 and 9). The results surprisingly show that chondrocytes with high inflammation (high C1M or C3M) not only have a better response to anabolic therapy such as sprifermin when administered according to an intense regimen with regard to cartilage thickness but also have a positive impact in the WOMAC total scores. This effect is even stronger when both C1M and C3M biomarkers are used (i.e. a combination of the biomarkers; Figure 10). Same trend is shown using both C3M/CRPM or C1M/CRPM (i.e. a combination of these biomarkers; data not shown).

10

### References

- 1) WO2008/023063
- 2) WO2004/032849
- 15 3) WO2014/023703
- 4) WO2006/063362
- 5) Haque et al., 2007, *Histol. Histopathol.*, 22 :97-105
- 6) [http://www.cartilage.org/\\_files/contentmanagement/ICRS\\_evaluation.pdf](http://www.cartilage.org/_files/contentmanagement/ICRS_evaluation.pdf)
- 7) Lotz, 2010, *Arthritis research therapy*, 12:211
- 20 8) Karsdal, 2016, *Osteoarthritis and Cartilage*, 24(12): 2013-2021
- 9) Ellsworth et al., 2002, *Osteoarthritis and Cartilage*, 10: 308-320
- 10) Shimoaka et al., 2002, *J. Bio. Chem.* 277(9):7493-7500
- 11) Gigout et al., 2017, *Osteoarthritis and Cartilage*, published online the 18.08.2018  
(<https://doi.org/10.1016/j.joca.2017.08.004>)
- 25 12) The Merck Manual, 17<sup>th</sup> edition, page 449
- 13) Bellamy et al., 1988, *J.Rheumatology*, 15:1833-1840
- 14) Wolfe, 1999, *Rheumatology*, 38:355-361
- 15) Siebuhr et al., 2014, *Osteoarthritis Cartilage*, 22(1):44-50.
- 16) Saberi Hosnijeh et al., 2016, *Arthritis Res Ther.* 18:81.
- 30 17) Siebuhr et al., 2013, *Arthritis Res Ther.*, 15(4):R86.
- 18) Siebuhr et al., 2016, *Biomark Med.*, 10(2):197-208.
- 19) Valdes et al., 2016, *Rheumatology (Oxford)*. 2016 Jun 2.
- 20) Maijer et al., 2016, *PLoS One.*, 11(3):e0149329. eCollection 2016.

35

**Abbreviations**

- OA = Osteoarthritis
- 5 CI=confidence interval,  
DBPC=double-blind placebo-controlled  
ICOAP=Measure of Intermittent and Constant Osteoarthritis Pain  
ITT=intention-to-treat,  
KOOS Symptom Index=Knee Injury and Osteoarthritis Outcome Score symptom index subscale
- 10 KOOS QOL=Knee Injury and Osteoarthritis Outcome Score quality of life subscale  
LOCF=last observation carried forward,  
LFTC = lateral femoro-tibial compartment  
MFTC = medial femoro-tibial compartment  
mITT=modified intention-to-treat
- 15 MOS SF-36=Medical Outcomes Study Short Form-36 General Health Survey  
MRI=magnetic resonance imaging  
NRS pain score=numerical rating scale pain score  
PGA=Patient's Global Assessment  
PGIC=Patient's Global Impression of Change
- 20 PK=pharmacokinetic  
W=Week  
WOMAC=Western Ontario and McMaster Universities Osteoarthritis Index.

**Claims**

1. A method of predicting the sensitivity to treatment with an FGF-18 compound in a subject having a cartilage disorder, the method comprising the steps of:
- 5 a) Determining, from a sample from said subject, the quantity of at least one of the biomarkers selected from the group consisting of C1M, C3M and/or CRPM; and
- b) Predicting from the result of step a) good or low sensitivity of said subject to treatment with an FGF-18 compound.
- 10 2. The method according to claim 1, comprising the steps of predicting low sensitivity to treatment with an FGF-18 compound from the presence of:
- a) higher than  $30.0 \pm 2SD$  ng/mL of C1M, and/or
- b) higher than  $12.0 \pm 2SD$  ng/mL of C3M, and/or
- 15 c) higher than  $10.0 \pm 2SD$  ng/mL of CRPM.
3. The method according to claim 1, comprising the steps of predicting good sensitivity, or sensitivity, to treatment with an FGF-18 compound from the presence of:
- a) lower than  $30.0 \pm 2SD$  ng/mL of C1M, and/or
- b) lower than  $12.0 \pm 2SD$  ng/mL of C3M, and/or
- 20 c) lower than  $10.0 \pm 2SD$  ng/mL of CRPM.
4. A method for selecting a subject having a cartilage disorder for inclusion in or exclusion from a specific dosing regimen when said subject is to be treated with an FGF-18 compound, based on the likelihood of their sensitivity to said treatment, comprising the steps of:
- 25 a) Determining, from a sample from said subject, the quantity of at least one of the biomarkers selected from the group consisting of C1M, C3M and/or CRPM; wherein the quantity of at least one of these proteins is predictive about the subject's risk of having good-sensitivity (or sensitivity) or low-sensitivity to said treatment, and
- b) Selecting the appropriate dosing regimen according to the subject's risk of having good-
- 30 sensitivity or low-sensitivity to said treatment,
- wherein said appropriate dosing regimen is at least two treatment cycles of 100  $\mu$ g of FGF-18 compound given once per week for 3 consecutive weeks per treatment cycle.
5. The method according to claim 4, comprising the steps of excluding from the treatment with FGF-
- 35 18 compound the subject presenting:
- a) higher than  $30.0 \pm 2SD$  ng/mL of C1M, and/or
- b) higher than  $12.0 \pm 2SD$  ng/mL of C3M, and/or
- c) higher than  $10.0 \pm 2SD$  ng/mL of CRPM,

wherein the excluded treatment with FGF-18 compound consists of the following dosing regimen: two treatment cycles of 100 µg of FGF-18 compound given once per week for 3 consecutive weeks per treatment cycle.

- 5 6. The method according to claim 4, comprising the steps of including for the treatment with FGF-18 compound, whatever the dosing regimen, the subject presenting:
- a) lower than  $30.0 \pm 2SD$  ng/mL of C1M, and/or
  - b) lower than  $12.0 \pm 2SD$  ng/mL of C3M, and/or
  - c) lower than  $10.0 \pm 2SD$  ng/mL of CRPM.
- 10 7. A kit comprising means for performing the method according to any of the claims 1 to 6 and instructions for use.
8. An FGF-18 compound for use in the treatment of a subject having a cartilage disorder,
- 15 characterized in that:
- i) If the subject has lower than  $30.0 \pm 2SD$  ng/mL of C1M and/or lower than  $12.0 \pm 2SD$  ng/mL of C3M and/or lower than  $10.0 \pm 2SD$  ng/mL of CRPM he can receive a treatment with an FGF-18 compound given at the following dosing regimen: at least two treatment cycles of 100 µg of FGF-18 compound given once per week for 3 consecutive weeks per treatment cycle,
  - 20 ii) If the subject has higher than  $30.0 \pm 2SD$  ng/mL of C1M and/or higher than  $12.0 \pm 2SD$  ng/mL of C3M and/or higher than  $10.0 \pm 2SD$  ng/mL of CRPM he can receive a treatment with an FGF-18 compound given at the following dosing regimen: four treatment cycles of 100 µg of FGF-18 compound given once per week for 3 consecutive weeks per treatment cycle.
- 25 9. A method for treating a subject having a cartilage disorder with an FGF-18 compound, comprising the following steps:
- a) Determining, from a sample, the quantity of at least one of the biomarkers selected from the group consisting of C1M, C3M and/or CRPM, wherein said quantities are predictive about the subject's risk of having good or low sensitivity to a treatment with said FGF-18 compound
  - 30 b) administering intraarticularly the FGF-18 compound at:
    - i. at least two treatment cycles of 100 µg of FGF-18 compound given once per week for 3 consecutive weeks per treatment cycle if the subject has lower than  $30.0 \pm 2SD$  ng/mL of C1M and/or lower than  $12.0 \pm 2SD$  ng/mL of C3M and/or lower than  $10.0 \pm 2SD$  ng/mL of CRPM or,
    - 35 ii. four treatment cycles of 100 µg of FGF-18 compound given once per week for 3 consecutive weeks per treatment cycle if the subject has higher than  $30.0 \pm 2SD$  ng/mL of C1M and/or higher than  $12.0 \pm 2SD$  ng/mL of C3M and/or higher than  $10.0 \pm 2SD$  ng/mL of CRPM.

10. The method according to any one of claims 1 to 6 and 9, the kit according to claim 7 or the FGF-18 compound for use according to any one of claims 8-9, wherein the FGF-18 compound is sprifermin or a fusion protein comprising an FGF-18 moiety.

- 5 11. The method according to any one of claims 1 to 6 and 9-10, the kit according to any one of claims 7 or 10 or the FGF-18 compound for use according to any one of claims 8-10, wherein the cartilage disorder is selected from the group consisting of osteoarthritis, cartilage injury, fractures affecting joint cartilage or surgical procedures with impact on joint cartilage, such as microfracture.




Figure 1




Figure 2




Figure 3






Figure 4




Figure 5




Figure 6




Figure 7




Figure 8




Figure 9




Figure 10

INTERNATIONAL SEARCH REPORT

International application No  
PCT/EP2018/076391

A. CLASSIFICATION OF SUBJECT MATTER  
INV. G01N33/68 A61K31/00 A61K38/18  
ADD.  
According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED  
Minimum documentation searched (classification system followed by classification symbols)  
G01N A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)  
EPO-Internal

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2014/023703 A1 (ARES TRADING SA [CH]) 13 February 2014 (2014-02-13) cited in the application	7,10,11
A	page 5, line 35 abstract claim 1	1-6,8,9
X	----- SIEBUHR ET AL.: OSTEOARTHRITIS CARTILAGE, vol. 22, no. 1, 2014, pages 44-50, XP002774971, cited in the application	7,11
A	page 45, paragraph "Biomarkers" ----- -/--	1-6,8-10

Further documents are listed in the continuation of Box C.

See patent family annex.

\* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search  25 October 2018	Date of mailing of the international search report  06/11/2018
--	--

Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer  Jacques, Patrice
--	--



## INTERNATIONAL SEARCH REPORT

International application No

PCT/EP2018/076391

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	ANNE SOFIE SIEBUHR ET AL: "Serological identification of fast progressors of structural damage with rheumatoid arthritis", ARTHRITIS RESEARCH AND THERAPY, BIOMED CENTRAL, LONDON, GB, vol. 15, no. 4, 14 August 2013 (2013-08-14), page R86, XP021175254, ISSN: 1478-6354, DOI: 10.1186/AR4266 cited in the application	7,11
A	page 3, paragraph "Biochemical markers" -----	1-6,8-10
A	WO 2008/023063 A2 (ARES TRADING SA [CH]; GIMONA ALBERTO [CH]; LADEL CHRISTOPH H [IT]; VOM) 28 February 2008 (2008-02-28) cited in the application the whole document -----	1-11
A	BAY-JENSEN A-C ET AL: "Elevated levels of CRPM, an inflammatory biomarker correlating with disease activity in RA, are prognostic of radiographic knee OA", OSTEOARTHRITIS AND CARTILAGE, vol. 25, 1 April 2017 (2017-04-01), XP029981908, ISSN: 1063-4584, DOI: 10.1016/J.JOCA.2017.02.065 the whole document -----	1-11
A	Y. MORI ET AL: "Identification of Fibroblast Growth Factor-18 as a Molecule to Protect Adult Articular Cartilage by Gene Expression Profiling", JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 289, no. 14, 4 April 2014 (2014-04-04), pages 10192-10200, XP055190480, ISSN: 0021-9258, DOI: 10.1074/jbc.M113.524090 the whole document -----	1-11
A	FATEMEH SABERI HOSNIJEH ET AL: "Association between biomarkers of tissue inflammation and progression of osteoarthritis: evidence from the Rotterdam study cohort", ARTHRITIS RESEARCH & THERAPY, vol. 18, no. 1, 1 April 2016 (2016-04-01), XP055418352, DOI: 10.1186/s13075-016-0976-3 the whole document -----	1-11

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2018/076391

Patent document cited in search report	Publication date	Patent family member(s)	Publication date	
WO 2014023703	A1	13-02-2014	AU 2013301606 A1	05-03-2015
			CA 2881024 A1	13-02-2014
			CN 104736723 A	24-06-2015
			EP 2880179 A1	10-06-2015
			ES 2660818 T3	26-03-2018
			HK 1207401 A1	29-01-2016
			JP 6262731 B2	17-01-2018
			JP 2015533078 A	19-11-2015
			JP 2018075019 A	17-05-2018
			US 2015218637 A1	06-08-2015
			WO 2014023703 A1	13-02-2014
			-----	
WO 2008023063	A2	28-02-2008	AR 062522 A1	12-11-2008
			AU 2007287510 A1	28-02-2008
			CA 2658511 A1	28-02-2008
			CY 1113136 T1	13-04-2016
			EA 200970224 A1	28-08-2009
			EP 2054050 A2	06-05-2009
			HR P20120643 T1	31-08-2012
			IL 197169 A	29-08-2013
			JP 5815206 B2	17-11-2015
			JP 2010501531 A	21-01-2010
			JP 2014088429 A	15-05-2014
			KR 20090051100 A	20-05-2009
			US 2010016223 A1	21-01-2010
			WO 2008023063 A2	28-02-2008
-----				