



(51) International Patent Classification:

A01H 5/10 (2006.01)

(21) International Application Number:

PCT/EP2017/060419

(22) International Filing Date:

02 May 2017 (02.05.2017)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

16169747.9	15 May 2016 (15.05.2016)	EP
16170036.4	17 May 2016 (17.05.2016)	EP

(71) Applicant: BAYER CROPSCIENCE NV [BE/BE]; J.E. Mommaertsiaan 14, Diegem 1831 (BE).

(72) Inventor: DESOMER, Jan; Technologiepark 38, 9052 Gent (BE).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to the identity of the inventor (Rule 4.17(i))
- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))
- of inventorship (Rule 4.17(iv))

(54) Title: METHOD FOR INCREASING YIELD IN BRASSICACEAE

(57) Abstract: The invention relates to a method for increasing yield in *Brassicaceae* comprising at least one event or allele as described, wherein the *Brassicaceae* plants, plant parts thereof, plant propagation material or the soil in which *Brassicaceae* plants are grown or intended to be grown are treated with active ingredients or mixtures thereof.



WO 2017/198449 A1

Method for increasing yield in *Brassicaceae*

The invention relates to a method for increasing yield in *Brassicaceae* comprising at least one event or allele as described, wherein that the *Brassicaceae* plants, plant parts thereof, plant propagation material or the soil in which *Brassicaceae* plants are grown or intended to be grown are treated with the active ingredients or mixtures thereof.

Brassicaceae represent an economically important family of flowering plants also called Cruciferae due to the cross-like shape of the flower . Important economic species are found in vegetables, e.g. Brassica oleracea (broccoli, cabbage, cauliflower, etc.), Brassica rapa (turnip, Chinese cabbage, etc.), oil crops, e.g. Brassica napus (rapeseed, Canola, etc.), ornamentals (Wallflower, Alyssum) or as model crops like Arabidopsis thaliana (thale cress) which is intensively investigated. Rapeseed oil is currently one of the most used cooking oils and is produced at over 18 millions metric tons for food and industrial use after palm oil and soybean oil. Therefore there is an ongoing interest to further secure and increase yield in *Brassicaceae* crops.

It has now been found that, surprisingly, certain active ingredients or mixtures thereof are particularly suitable for increasing yield in *Brassicaceae*, in particular in winter and spring oilseed rape or Canola.

The following active ingredients active ingredients can be used to treat *Brassicaceae* plants or parts thereof, including seeds, to increase yield:

Herbicides:

Herbicidally active compounds useful in the context of the present invention, also in mixed formulations or in tank mix, are, for example, known active compounds as they are described in, for example, Weed Research 26, 441-445 (1986), or "The Pesticide Manual", 16th edition, November 2012, The British Crop Protection Council and the Royal Soc. of Chemistry, and the literature cited therein, and which for example act as inhibitor of acetolactate synthase, acetyl-CoA-carboxylase, cellulose-synthase, enolpyruvylshikimat-3-phosphat-synthase, glutamin-synthetase, p-hydroxyphenylpyruvat-dioxygenase, phytoendesaturase, photosystem I, photosystem II, and/or protoporphyrinogen-oxidase.

Examples of active compounds which may be mentioned as herbicides or plant growth regulators which are known from the literature and which can be combined with the compounds according to the invention are the following (compounds are either described by "common name" in accordance with the International Organization for Standardization (ISO) or by chemical name or by a customary code number), and always comprise all applicable forms such as acids, salts, ester, or modifications such as isomers, like stereoisomers and optical isomers. As an example at least one applicable form and/or modifications can be mentioned.

Examples for **herbicides** useful in the context of the present invention are disclosed in group H-1:

Acetochlor, acifluorfen, acifluorfen-sodium, aclonifen, alachlor, allidochlor, alloxydim, alloxydim-sodium, ametryn, amicarbazone, amidochlor, amidosulfuron, 4-amino-3-chloro-6-(4-chloro-2-fluoro-3-methylphenyl)-5-fluoropyridine-2-carboxylic acid, aminocyclopyrachlor, aminocyclopyrachlor-potassium, aminocyclopyrachlor-methyl, aminopyralid, amitrole, ammoniumsulfamate, anilofos, 5 asulam, atrazine, azafenidin, azimsulfuron, beflubutamid, benazolin, benazolin-ethyl, benfluralin, benfuresate, bensulfuron, bensulfuron-methyl, bensulide, bentazone, benzobicyclon, benzofenap, bicyclopyron, bifenox, bilanafos, bilanafos-sodium, bispyribac, bispyribac-sodium, bromacil, bromobutide, bromofenoxim, bromoxynil, bromoxynil-butyrate, -potassium, -heptanoate, and -
10 octanoate, busoxinone, butachlor, butafenacil, butamifos, butenachlor, butralin, butroxydim, butylate, cafenstrole, carbetamide, carfentrazone, carfentrazone-ethyl, chloramben, chlorbromuron, chlorfenac, chlorfenac-sodium, chlorfenprop, chlorflurenol, chlorflurenol-methyl, chloridazon, chlorimuron, chlorimuron-ethyl, chlorophthalim, chlorotoluron, chlorthal-dimethyl, chlorsulfuron, cinidon, cinidon-ethyl, cinmethylin, cinosulfuron, clacyfos, clethodim, clodinafop, clodinafop-propargyl, clomazone, 15 clomeprop, clopyralid, cloransulam, cloransulam-methyl, cumyluron, cyanamide, cyanazine, cycloate, cyclopyrimorate, cyclosulfamuron, cycloxydim, cyhalofop, cyhalofop-butyl, cyprazine, 2,4-D, 2,4-D-butotyl, -butyl, -dimethylammonium, -diolamin, -ethyl, -2-ethylhexyl, -isobutyl, -isooctyl, -isopropyl-ammonium, -potassium, -triisopropanolammonium, and -trolamine, 2,4-DB, 2,4-DB-butyl, -dimethyl-ammonium, -isooctyl, -potassium, and -sodium, daimuron (dymron), dalapon, dazomet, n-decanol, 20 desmedipham, detosyl-pyrazolate (DTP), dicamba, dichlobenil, 2-(2,4-dichlorobenzyl)-4,4-dimethyl-1,2-oxazolidin-3-one, 2-(2,5-dichlorobenzyl)-4,4-dimethyl-1,2-oxazolidin-3-one, dichlorprop, dichlorprop-P, diclofop, diclofop-methyl, diclofop-P-methyl, diclosulam, difenzoquat, diflufenican, diflufenzopyr, diflufenzopyr-sodium, dimefuron, dimepiperate, dimethachlor, dimethametryn, dimethenamid, dimethenamid-P, dimetrasulfuron, dinitramine, dinoterb, diphenamid, diquat, diquat-25 dibromid, dithiopyr, diuron, DNOC, endothal, EPTC, esprocarb, ethalfluralin, ethametsulfuron, ethametsulfuron-methyl, ethiozin, ethofumesate, ethoxyfen, ethoxyfen-ethyl, ethoxysulfuron, etobenzanid, F-9600, F-5231, i.e. N-{2-chloro-4-fluoro-5-[4-(3-fluoropropyl)-5-oxo-4,5-dihydro-1H-tetrazol-1-yl]phenyl}ethanesulfonamide, F-7967, i. e. 3-[7-chloro-5-fluoro-2-(trifluoromethyl)-1H-benzimidazol-4-yl]-1-methyl-6-(trifluoromethyl)pyrimidine-2,4(1H,3H)-dione, fenoxaprop, fenoxaprop-P, fenoxa-30 prop-ethyl, fenoxaprop-P-ethyl, fenoxasulfone, fenquino-trione, fentrazamide, flamprop, flamprop-M-isopropyl, flamprop-M-methyl, flazasulfuron, florasulam, fluazifop, fluazifop-P, fluazifop-butyl, fluazifop-P-butyl, flucarbazone, flucarbazone-sodium, flucetosulfuron, fluchloralin, flufenacet, flufenpyr, flufenpyr-ethyl, flumetsulam, flumiclorac, flumiclorac-pentyl, flumioxazin, fluometuron, flurenol, flurenol-butyl, -dimethylammonium and -methyl, fluoroglycofen, fluoroglycofen-ethyl, flu-35 propanate, flupyr-sulfuron, flupyr-sulfuron-methyl-sodium, fluridone, flurochloridone, fluroxypyr, fluroxypyr-meptyl, flurtamone, fluthiacet, fluthiacet-methyl, fomesafen, fomesafen-sodium, foramsulfuron, fosamine, glufosinate, glufosinate-ammonium, glufosinate-P-sodium, glufosinate-P-ammonium, glufosinate-P-sodium, glyphosate, glyphosate-ammonium, -isopropyl-

ammonium, -diammonium, -dimethylammonium, -potassium, -sodium, and -trimesium, H-9201, i.e. O-(2,4-dimethyl-6-nitrophenyl) O-ethyl isopropylphosphoramidothioate, halauxifen, halauxifen-methyl, halosafen, halosulfuron, halosulfuron-methyl, haloxyfop, haloxyfop-P, haloxyfop-ethoxyethyl, haloxyfop-P-ethoxyethyl, haloxyfop-methyl, haloxyfop-P-methyl, hexazinone, HW-02, i.e. 1-

5 (dimethoxyphosphoryl) ethyl-(2,4-dichlorophenoxy)acetate, imazamethabenz, imazamethabenz-methyl, imazamox, imazamox-ammonium, imazapic, imazapic-ammonium, imazapyr, imazapyr-isopropylammonium, imazaquin, imazaquin-ammonium, imazethapyr, imazethapyr-immonium, imazosulfuron, indanofan, indaziflam, iodosulfuron, iodosulfuron-methyl-sodium, ioxynil, ioxynil-octanoate, -potassium and -sodium, ipfencarbazone, isoproturon, isouron, isoxaben, isoxaflutole,

10 karbutilate, KUH-043, i.e. 3-([5-(difluoromethyl)-1-methyl-3-(trifluoromethyl)-1H-pyrazol-4-yl]methyl)sulfonyl)-5,5-dimethyl-4,5-dihydro-1,2-oxazole, ketospiradox, lactofen, lenacil, linuron, MCPA, MCPA-butotyl, -dimethylammonium, -2-ethylhexyl, -isopropylammonium, -potassium, and -sodium, MCPB, MCPB-methyl, -ethyl and -sodium, mecoprop, mecoprop-sodium, and -butotyl, mecoprop-P, mecoprop-P-butotyl, -dimethylammonium, -2-ethylhexyl, and -potassium, mefenacet,

15 mefluidide, mesosulfuron, mesosulfuron-methyl, mesotrione, methabenzthiazuron, metam, metamifop, metamitron, metazachlor, metazosulfuron, methabenzthiazuron, methiopyrsulfuron, methiozolin, methyl isothiocyanate, metobromuron, metolachlor, S-metolachlor, metosulam, metoxuron, metribuzin, metsulfuron, metsulfuron-methyl, molinat, monolinuron, monosulfuron, monosulfuron-ester, MT-5950, i.e. N-(3-chloro-4-isopropylphenyl)-2-methylpentan amide, NGGC-011, napropamide, NC-310, i.e. [5-


20 (benzyloxy)-1-methyl-1H-pyrazol-4-yl](2,4-dichlorophenyl)methanone, neburon, nicosulfuron, nonanoic acid (pelargonic acid), norflurazon, oleic acid (fatty acids), orbencarb, orthosulfamuron, oryzalin, oxadiargyl, oxadiazon, oxasulfuron, oxaziclomefon, oxyfluorfen, paraquat, paraquat dichloride, pebulate, pendimethalin, penoxsulam, pentachlorophenol, pentoxazone, pethoxamid, petroleum oils, phenmedipham, picloram, picolinafen, pinoxaden, piperophos, pretilachlor,

25 primisulfuron, primisulfuron-methyl, prodiamine, profoxydim, prometon, prometryn, propachlor, propanil, propaquizafop, propazine, propham, propisochlor, propoxycarbazone, propoxycarbazone-sodium, propyrisulfuron, propyzamide, prosulfocarb, prosulfuron, pyraclonil, pyraflufen, pyraflufen-ethyl, pyrasulfotole, pyrazolynate (pyrazolate), pyrazosulfuron, pyrazosulfuron-ethyl, pyrazoxyfen, pyribambenz, pyribambenz-isopropyl, pyribambenz-propyl, pyribenzoxim, pyributicarb, pyridafol,

30 pyridate, pyriftalid, pyriminobac, pyriminobac-methyl, pyrimisulfan, pyrithiobac, pyrithiobac-sodium, pyroxasulfone, pyroxsulam, quinclorac, quinmerac, quinochloramine, quizalofop, quizalofop-ethyl, quizalofop-P, quizalofop-P-ethyl, quizalofop-P-tefuryl, rimsulfuron, saflufenacil, sethoxydim, siduron, simazine, simetryn, SL-261, sulcotrione, sulfentrazone, sulfometuron, sulfometuron-methyl, sulfosulfuron, SYN-523, SYP-249, i.e. 1-ethoxy-3-methyl-1-oxobut-3-en-2-yl 5-[2-chloro-4-

35 (trifluoromethyl)phenoxy]-2-nitrobenzoate, SYP-300, i.e. 1-[7-fluoro-3-oxo-4-(prop-2-yn-1-yl)-3,4-dihydro-2H-1,4-benzoxazin-6-yl]-3-propyl-2-thioxoimidazolidine-4,5-dione, 2,3,6-TBA, TCA (trichloroacetic acid), TCA-sodium, tebuthiuron, tefuryltrione, tembotrione, tepraloxym, terbacil, terbucarb, terbumeton, terbuthylazin, terbutryn, thenylchlor, thiazopyr, thiencarbazone, thiencarbazone-

methyl, thifensulfuron, thifensulfuron-methyl, thiobencarb, tiafenacil, tolypyralate, topramezone, tralkoxydim, triafamone, tri-allate, triasulfuron, triaziflam, tribenuron, tribenuron-methyl, triclopyr, trietazine, trifloxysulfuron, trifloxysulfuron-sodium, trifludimoxazin, trifluralin, triflusulfuron, triflusulfuron-methyl, tritosulfuron, urea sulfate, vernolate, XDE-848, ZJ-0862, i.e. 3,4-dichloro-N-{2-
5 [(4,6-dimethoxypyrimidin-2-yl)oxy]benzyl}aniline, and the following compounds:



Group H-2 discloses herbicides especially used in the context of the present invention are selected from the group consisting of:

Acetochlor, Alachlor, Dicamba, Diflufenican, Flufenacet, Fluroxypyr, Foramsulfuron, Glufosinate, L-
10 Glufosinate, Glyphosate, Hexazinone, Imazamox, Imazethapyr, Indaziflam, Iodosulfuron, Isoxaflutole, Mesosulfuron, Mesotrione, Metolachlor, Metribuzin, Metsulfuron, Pendimethalin, Penoxsulam, Picloram, Pinoxaden, Pyroxsulam, Quinmerac, Rimsulfuron, Thiencarbazone, Tembotrione, Thifensulfuron, and/or Tribenuron, and the esters and/or agronomically acceptable salts of these herbicides.

15 Group H-3 discloses preferred herbicides are selected from the group consisting of:

Acetochlor, Dicamba, Diflufenican, Flufenacet, Foramsulfuron, Glufosinate, L-Glufosinate, Glyphosate, Iodosulfuron, Isoxaflutole, Mesosulfuron, Mesotrione, Metribuzin, Pinoxaden, Tembotrione, and /or Thiencarbazone, and the esters and/or agronomically acceptable salts of these herbicides.

If Glyphosate salts are used as herbicide, preference is given to Glyphosate isopropylamine salt,
20 Glyphosate potassium salt, Glyphosate sodium salt, Glyphosate trimethylsulfonium salt.

If (L-)Glufosinate salts are used as herbicide, preference is given to (L-)Glufosinate-ammonium salt, (L-)Glufosinate potassium salt and (L-)Glufosinate sodium salt.

If Foramsulfuron salts are used as herbicide, preference is given to Foramsulfuron-sodium.

If Iodosulfuron esters and/or salts thereof are used as herbicide, preference is given to Iodosulfuron-methyl and Iodosulfuron-methyl-sodium.

5 If Mesosulfuron esters and/or salts thereof are used as herbicide, preference is given to Mesosulfuron-methyl and Mesosulfuron-methyl-sodium.

If Thiencarbazone esters and/or salts thereof are used as herbicide, preference is given to Thiencarbazone-methyl and Thiencarbazone-methyl-sodium.

Preferred mixtures of herbicides used in the context of the present invention are disclosed in group H-4: mixtures of Acetochlor and other herbicides mentioned herein, mixtures of Dicamba, esters and/or salts
10 thereof and other herbicides mentioned herein, mixtures of Diflufenican and other herbicides mentioned herein, mixtures of Flufenacet and other herbicides mentioned herein, mixtures of Glufosinate and/or salts thereof (preferably Glufosinate-ammonium salt, Glufosinate potassium salt and Glufosinate sodium salt) and other herbicides mentioned herein, mixtures of L-Glufosinate and/or salts thereof (preferably L-Glufosinate-ammonium salt, L-Glufosinate potassium salt and L-Glufosinate sodium salt) and other
15 herbicides mentioned herein, mixtures of Indaziflam and other herbicides mentioned herein, mixtures of Isoxaflutole and other herbicides mentioned herein, mixtures of Mesosulfuron esters and/or salts thereof (preferably Mesosulfuron-methyl(-sodium)) and other herbicides mentioned herein, mixtures of Metribuzin and other herbicides mentioned herein, mixtures of Thiencarbazone esters and/or salts thereof (preferably Thiencarbazone-methyl(-sodium)) and other herbicides mentioned herein, mixtures
20 of Tembotrione and other herbicides mentioned herein.

Particularly preferred mixtures of herbicides used in the context of the present invention are disclosed in group H-5:

25 mixtures of Acetochlor and Isoxaflutole, mixtures of Acetochlor and Tembotrione mixtures of Diflufenican and Flufenacet, mixtures of Diflufenican and Metribuzin, mixtures of Glufosinate and/or salts thereof and Indaziflam, in particular mixtures of Glufosinate-ammonium and Indaziflam, mixtures of L-Glufosinate and/or salts thereof and Indaziflam, in particular mixtures of L-Glufosinate-ammonium and Indaziflam, mixtures of Mesosulfuron-methyl(-sodium) and Iodosulfuron-methyl(-sodium), mixtures of Mesosulfuron-methyl(-sodium), Iodosulfuron-methyl(-sodium) and Flufenacet, mixtures of Tembotrione and Isoxaflutole, mixtures of Tembotrione, Isoxaflutole, and Acetochlor.

30 Examples for **plant growth regulators** useful in the context of the present invention are disclosed in group P-1:

Acibenzolar, acibenzolar-S-methyl, 5-aminolevulinic acid, ancymidol, 6-benzylaminopurine, Brassinolid, catechine, chlormequat chloride, cloprop, cyclanilide, 3-(cycloprop-1-enyl) propionic acid,

daminozide, dazomet, n-decanol, dikegulac, dikegulac-sodium, endothal, endothal-dipotassium, -disodium, and -mono(N,N-dimethylalkylammonium), ethephon, flumetralin, flurenol, flurenol-butyl, flurprimidol, forchlorfenuron, gibberellic acid, inabenfide, indol-3-acetic acid (IAA), 4-indol-3-ylbutyric acid, isoprothiolane, probenazole, jasmonic acid, maleic hydrazide, mepiquat chloride, 5 1-methylcyclopropene, methyl jasmonate, 2-(1-naphthyl)acetamide, 1-naphthylacetic acid, 2-naphthyl-oxyacetic acid, nitrophenolate-mixture, paclobutrazol, N-(2-phenylethyl)-beta-alanine, N-phenyl-phthalamic acid, prohexadione, prohexadione-calcium, prohydrojasmon, salicylic acid, strigolactone, tecnazene, thidiazuron, triacontanol, trinexapac, trinexapac-ethyl, tsitodef, uniconazole, uniconazole-P.

Examples for **safeners** useful in the context of the present invention are disclosed in group SAF-1 in 16 subgroups:

- S1) compounds of the group of heterocyclic carboxylic acid derivatives:
- S1^a) compounds of the type of dichlorophenylpyrazoline-3-carboxylic acid (S1^a), preferably compounds such as 1-(2,4-dichlorophenyl)-5-(ethoxycarbonyl)-5-methyl-2-pyrazoline-3-carboxylic acid, 15 ethyl 1-(2,4-dichlorophenyl)-5-(ethoxycarbonyl)-5-methyl-2-pyrazoline-3-carboxylate (S1-1) ("mefenpyr(-diethyl)"), and related compounds, as described in WO-A-91/07874;
- S1^b) derivatives of dichlorophenylpyrazolecarboxylic acid (S1^b), preferably compounds such as ethyl 1-(2,4-dichlorophenyl)-5-methylpyrazole-3-carboxylate (S1-2), ethyl 1-(2,4-dichlorophenyl)-5-isopropylpyrazole-3-carboxylate (S1-3), 20 ethyl 1-(2,4-dichlorophenyl)-5-(1,1-dimethylethyl)pyrazole-3-carboxylate (S1-4) and related compounds, as described in EP-A-333 131 and EP-A-269 806;
- S1^c) derivatives of 1,5-diphenylpyrazole-3-carboxylic acid (S1^c), preferably compounds such as ethyl 1-(2,4-dichlorophenyl)-5-phenylpyrazole-3-carboxylate (S1-5), methyl 1-(2-chlorophenyl)-5-phenylpyrazole-3-carboxylate (S1-6) and related compounds, as 25 described, for example, in EP-A-268554;
- S1^d) compounds of the type of triazolecarboxylic acids (S1^d), preferably compounds such as fenclorazole(-ethyl), i.e. ethyl 1-(2,4-dichlorophenyl)-5-trichloromethyl-(1H)-1,2,4-triazole-3-carboxylate (S1-7), and related compounds, as described in EP-A-174 562 and EP-A-346 620;
- S1^e) compounds of the type of 5-benzyl- or 5-phenyl-2-isoxazoline-3-carboxylic acid or 5,5-diphenyl-2-isoxazoline-3-carboxylic acid (S1^e), preferably compounds such as ethyl 5-(2,4-dichlorobenzyl)-2-isoxazoline-3-carboxylate (S1-8) or ethyl 5-phenyl-2-isoxazoline-3-carboxylate (S1-9) and related compounds, as described in WO-A-91/08202, or 5,5-diphenyl-2-isoxazolinecarboxylic acid (S1-10) or ethyl 5,5-diphenyl-2-isoxazolinecarboxylate (S1-11) ("isoxadifen-ethyl") or n-propyl 5,5-diphenyl-2-isoxazolinecarboxylate (S1-12) or ethyl 30

5-(4-fluorophenyl)-5-phenyl-2-isoxazoline-3-carboxylate (S1-13), as described in the patent application WO-A-95/07897;

S2) Compounds of the group of 8-quinolinoxy derivatives (S2):

S2^a) compounds of the type of 8-quinolinoxyacetic acid (S2^a), preferably
 5 1-methylhexyl (5-chloro-8-quinolinoxy)acetate (common name "cloquintocet-mexyl" (S2-1),
 1,3-dimethyl-but-1-yl (5-chloro-8-quinolinoxy)acetate (S2-2),
 4-allyloxybutyl (5-chloro-8-quinolinoxy)acetate (S2-3),
 1-allyloxyprop-2-yl (5-chloro-8-quinolinoxy)acetate (S2-4),
 ethyl (5-chloro-8-quinolinoxy)acetate (S2-5),
 10 methyl (5-chloro-8-quinolinoxy)acetate (S2-6),
 allyl (5-chloro-8-quinolinoxy)acetate (S2-7),
 2-(2-propylideneiminoxy)-1-ethyl (5-chloro-8-quinolinoxy)acetate (S2-8),
 2-oxo-prop-1-yl (5-chloro-8-quinolinoxy)acetate (S2-9) and related compounds, as described in
 EP-A-86 750, EP-A-94 349 and EP-A-191 736 or EP-A-0 492 366, and also (5-chloro-8-
 15 quinolinoxy)acetic acid (S2-10), its hydrates and salts, for example its lithium, sodium,
 potassium, calcium, magnesium, aluminium, iron, ammonium, quaternary ammonium,
 sulphonium or phosphonium salts, as described in WO-A-2002/34048;


S2^b) compounds of the type of (5-chloro-8-quinolinoxy)malonic acid (S2^b), preferably compounds
 20 such as diethyl (5-chloro-8-quinolinoxy)malonate, diallyl (5-chloro-8-quinolinoxy)malonate,
 methyl ethyl (5-chloro-8-quinolinoxy)malonate and related compounds, as described in
 EP-A-0 582 198;

S3) Active compounds of the type of dichloroacetamides (S3) which are frequently used as pre-
 emergence safeners (soil-acting safeners), such as, for example,
 "dichlormid" (N,N-diallyl-2,2-dichloroacetamide) (S3-1),
 25 "R-29148" (3-dichloroacetyl-2,2,5-trimethyl-1,3-oxazolidine) from Stauffer (S3-2),
 "R-28725" (3-dichloroacetyl-2,2-dimethyl-1,3-oxazolidine) from Stauffer (S3-3),
 "benoxacor" (4-dichloroacetyl-3,4-dihydro-3-methyl-2H-1,4-benzoxazine) (S3-4),
 "PPG-1292" (N-allyl-N-[(1,3-dioxolan-2-yl)methyl]dichloroacetamide) from PPG Industries
 (S3 5),
 30 "DKA-24" (N-allyl-N-[(allylaminocarbonyl)methyl]dichloroacetamide) from Sagro-Chem (S3-
 6),
 "AD-67" or "MON 4660" (3-dichloroacetyl-1-oxa-3-aza-spiro[4,5]decane) from Nitrokemia or
 Monsanto (S3-7),
 "TI-35" (1-dichloroacetylazepane) from TRI-Chemical RT (S3-8)
 35 "diclonon" (dicyclonon) or "BAS145138" or "LAB145138" (S3-9)
 ((RS)-1-dichloroacetyl-3,3,8a-trimethylperhydropyrrolo[1,2-a]pyrimidin-6-one) from BASF,

furilazole" or "MON 13900" ((RS)-3-dichloroacetyl-5-(2-furyl)-2,2-dimethyloxazolidine) (S3-10), and also its (R)-isomer (S3-11);

S4) Compounds of the class of acylsulphonamides (S4):

S4^a) N-acylsulphonamides of the formula (S4^a) and salts thereof, as described in WO-A-97/45016



in which

R_A^1 is (C₁-C₆)-alkyl, (C₃-C₆)-cycloalkyl, where the 2 last-mentioned radicals are substituted by v_A substituents from the group consisting of halogen, (C₁-C₄)-alkoxy, halo-(C₁-C₆)-alkoxy and (C₁-C₄)-alkylthio and, in the case of cyclic radicals, also (C₁-C₄)-alkyl and (C₁-C₄)-haloalkyl;

10


R_A^2 is halogen, (C₁-C₄)-alkyl, (C₁-C₄)-alkoxy, CF₃;

m_A is 1 or 2;

v_D is 0, 1, 2 or 3;

S4^b) compounds of the type of 4-(benzoylsulphamoyl)benzamides of the formula (S4^b) and salts thereof, as described in WO-A-99/16744,

15



in which

R_B^1, R_B^2 independently of one another are hydrogen, (C₁-C₆)-alkyl, (C₃-C₆)-cycloalkyl, (C₃-C₆)-alkenyl, (C₃-C₆)-alkynyl,

R_B^3 is halogen, (C₁-C₄)-alkyl, (C₁-C₄)-haloalkyl or (C₁-C₄)-alkoxy,

20

m_B is 1 or 2;

for example those in which

$R_B^1 =$ cyclopropyl, $R_B^2 =$ hydrogen and $(R_B^3) =$ 2-OMe ("cyprosulfamide", S4-1),


R_B^1 = cyclopropyl, R_B^2 = hydrogen and $(R_B^3) = 5\text{-Cl-2-OMe}$ (S4-2),

R_B^1 = ethyl, R_B^2 = hydrogen and $(R_B^3) = 2\text{-OMe}$ (S4-3),

R_B^1 = isopropyl, R_B^2 = hydrogen and $(R_B^3) = 5\text{-Cl-2-OMe}$ (S4-4) and

R_B^1 = isopropyl, R_B^2 = hydrogen and $(R_B^3) = 2\text{-OMe}$ (S4-5);

- 5 S4^c) compounds of the class of benzoylsulphamoylphenylureas of the formula (S4^c) as described in EP-A-365484,



in which

- 10 R_C^1, R_C^2 independently of one another are hydrogen, (C₁-C₈)-alkyl, (C₃-C₈)-cycloalkyl, (C₃-C₆)-alkenyl, (C₃-C₆)-alkynyl,


R_C^3 is halogen, (C₁-C₄)-alkyl, (C₁-C₄)-alkoxy, CF₃,

m_C is 1 or 2;

for example

- 15 1-[4-(N-2-methoxybenzoylsulphamoyl)phenyl]-3-methylurea,
1-[4-(N-2-methoxybenzoylsulphamoyl)phenyl]-3,3-dimethylurea,
1-[4-(N-4,5-dimethylbenzoylsulphamoyl)phenyl]-3-methylurea;

- S4^d) compounds of the type of N-phenylsulphonylterephthalamides of the formula (S4^d) and salts thereof, which are known, for example, from CN 101838227,



- 20 in which

R_D^4 is halogen, (C₁-C₄)-alkyl, (C₁-C₄)-alkoxy, CF₃;

m_D is 1 or 2;


R_D^5 is hydrogen, (C₁-C₆)-alkyl, (C₃-C₆)-cycloalkyl, (C₂-C₆)-alkenyl, (C₂-C₆)-alkynyl, (C₅-C₆)-cycloalkenyl;

5 S5) Active compounds from the class of hydroxyaromatics and aromatic-aliphatic carboxylic acid derivatives (S5), for example ethyl 3,4,5-triacetoxybenzoate, 3,5-dimethoxy-4-hydroxybenzoic acid, 3,5-dihydroxybenzoic acid, 4-hydroxysalicylic acid, 4-fluorosalicylic acid, 2-hydroxycinnamic acid, 2,4-dichlorocinnamic acid, as described in WO-A-2004/084631, WO-A-2005/015994, WO-A-2005/016001;

10 S6) Active compounds from the class of 1,2-dihydroquinoxalin-2-ones (S6), for example 1-methyl-3-(2-thienyl)-1,2-dihydroquinoxalin-2-one, 1-methyl-3-(2-thienyl)-1,2-dihydroquinoxaline-2-thione, 1-(2-aminoethyl)-3-(2-thienyl)-1,2-dihydroquinoxalin-2-one hydrochloride, 1-(2-methylsulphonylaminoethyl)-3-(2-thienyl)-1,2-dihydroquinoxalin-2-one, as described in WO-A-2005/112630;

15 S7) Compounds from the class of diphenylmethoxyacetic acid derivatives (S7), for example methyl diphenylmethoxyacetate (CAS-Reg.Nr. 41858-19-9) (S7-1), ethyl diphenylmethoxyacetate, or diphenylmethoxyacetic acid, as described in WO-A-98/38856;

S8) Compounds of the formula (S8), as described in WO-A-98/27049,



where the symbols and indices have the following meanings:

R_D^1 is halogen, (C₁-C₄)-alkyl, (C₁-C₄)-haloalkyl, (C₁-C₄)-alkoxy, (C₁-C₄)-haloalkoxy,

20 R_D^2 is hydrogen or (C₁-C₄)-alkyl,


R_D^3 is hydrogen, (C₁-C₈)-alkyl, (C₂-C₄)-alkenyl, (C₂-C₄)-alkynyl or aryl, where each of the carbon-containing radicals mentioned above is unsubstituted or substituted by one or more, preferably by up to three, identical or different radicals from the group consisting of halogen and alkoxy; or salts thereof,

25 n_D is an integer from 0 to 2;

S9) Active compounds from the class of 3-(5-tetrazolylcarbonyl)-2-quinolones (S9), for example 1,2-dihydro-4-hydroxy-1-ethyl-3-(5-tetrazolylcarbonyl)-2-quinolone (CAS Reg. No.: 219479-

18-2), 1,2-dihydro-4-hydroxy-1-methyl-3-(5-tetrazolylcarbonyl)-2-quinolone (CAS Reg. No.: 95855-00-8), as described in WO-A-1999/000020;

S10) Compounds of the formula (S10^a) or (S10^b) as described in WO-A-2007/023719 and WO-A-2007/023764



5 in which

R_E^1 is halogen, (C₁-C₄)-alkyl, methoxy, nitro, cyano, CF₃, OCF₃

Y_E, Z_E independently of one another are O or S,

n_E is an integer from 0 to 4,

R_E^2 is (C₁-C₁₆)-alkyl, (C₂-C₆)-alkenyl, (C₃-C₆)-cycloalkyl, aryl; benzyl, halobenzyl,

10 R_E^3 is hydrogen or (C₁-C₆)-alkyl;

S11) Active compounds of the type of oxyimino compounds (S11), which are known as seed dressings, such as, for example, "oxabetrinil" ((Z)-1,3-dioxolan-2-ylmethoxyimino(phenyl)acetonitrile) (S11-1), which is known as seed dressing safener for millet against metolachlor damage,

15 "fluxofenim" (1-(4-chlorophenyl)-2,2,2-trifluoro-1-ethanone O-(1,3-dioxolan-2-ylmethyl)oxime) (S11-2), which is known as seed dressing safener for millet against metolachlor damage, and


"cyometrinil" or "CGA-43089" ((Z)-cyanomethoxyimino(phenyl)acetonitrile) (S11-3), which is known as seed dressing safener for millet against metolachlor damage;

20 S12) Active compounds from the class of isothiochromanones (S12), such as, for example, methyl [(3-oxo-1H-2-benzothiopyran-4(3H)-ylidene)methoxy]acetate (CAS Reg. No.: 205121-04-6) (S12-1) and related compounds from WO-A-1998/13361;

S13) One or more compounds from group (S13):

- "naphthalic anhydrid" (1,8-naphthalenedicarboxylic anhydride) (S13-1), which is known as seed dressing safener for corn against thiocarbamate herbicide damage,
- "fencloirim" (4,6-dichloro-2-phenylpyrimidine) (S13-2), which is known as safener for pretilachlor in sown rice,
- 5 "flurazole" (benzyl 2-chloro-4-trifluoromethyl-1,3-thiazole-5-carboxylate) (S13-3), which is known as seed dressing safener for millet against alachlor and metolachlor damage,
- "CL 304415" (CAS Reg. No.: 31541-57-8)
(4-carboxy-3,4-dihydro-2H-1-benzopyran-4-acetic acid) (S13-4) from American Cyanamid, which is known as safener for corn against imidazolinone damage,
- 10 "MG 191" (CAS Reg. No.: 96420-72-3) (2-dichloromethyl-2-methyl-1,3-dioxolane) (S13-5) from Nitrokemia, which is known as safener for corn,
- "MG 838" (CAS Reg. No.: 133993-74-5)
(2-propenyl 1-oxa-4-azaspiro[4.5]decane-4-carbodithioate) (S13-6) from Nitrokemia,
- "disulphoton" (O,O-diethyl S-2-ethylthioethyl phosphorodithioate) (S13-7),
- 15 "dietholate" (O,O-diethyl O-phenyl phosphorothioate) (S13-8),
- "mephenate" (4-chlorophenyl methylcarbamate) (S13-9);
- S14) Active compounds which, besides a herbicidal effect against harmful plants, also have a safener effect on crop plants such as rice, such as, for example,
- 20 "dimepiperate" or "MY 93" (*S*-1-methyl-1-phenylethyl piperidine-1-carbothioate), which is known as safener for rice against molinate herbicide damage,
- "daimuron" or "SK 23" (1-(1-methyl-1-phenylethyl)-3-p-tolylurea), which is known as safener for rice against imazosulphuron herbicide damage,
- "cumyluron" = "JC 940" (3-(2-chlorophenylmethyl)-1-(1-methyl-1-phenylethyl)urea, see JP-A-60087254), which is known as safener for rice against some herbicide damage,
- 25 "methoxyphenone" or "NK 049" (3,3'-dimethyl-4-methoxybenzophenone), which is known as safener for rice against some herbicide damage,
- "CSB" (1-bromo-4-(chloromethylsulphonyl)benzene) from Kumiai (CAS Reg. No. 54091-06-4), which is known as safener against some herbicide damage in rice;

S15) Compounds of the formula (S15) or its tautomers,



as described in WO-A-2008/131861 and WO-A-2008/131860,

in which

R_H^1 is (C₁-C₆)-haloalkyl,

5 R_H^2 is hydrogen or halogen,

R_H^3, R_H^4 independently of one another are hydrogen, (C₁-C₁₆)-alkyl, (C₂-C₁₆)-alkenyl or (C₂-C₁₆)-alkynyl,

where each of the 3 last-mentioned radicals is unsubstituted or substituted by one or more radicals from the group consisting of halogen, hydroxy, cyano, (C₁-C₄)-alkoxy, (C₁-C₄)-haloalkoxy, (C₁-C₄)-alkylthio, (C₁-C₄)-alkylamino, di-[(C₁-C₄)-alkyl]-amino, [(C₁-C₄)-alkoxy]-carbonyl, [(C₁-C₄)-haloalkoxy]-carbonyl, unsubstituted or substituted (C₃-C₆)-cycloalkyl, unsubstituted or substituted phenyl, and unsubstituted or substituted heterocyclyl;

or (C₃-C₆)-cycloalkyl, (C₄-C₆)-cycloalkenyl, (C₃-C₆)-cycloalkyl which is at one site of the ring condensed with a 4 to 6-membered saturated or unsaturated carbocyclic ring, or (C₄-C₆)-cycloalkenyl which is at one site of the ring condensed with a 4 to 6-membered saturated or unsaturated carbocyclic ring,

where each of the 4 last-mentioned radicals is unsubstituted or substituted by one or more radicals from the group consisting of halogen, hydroxy, cyano, (C₁-C₄)-alkyl, (C₁-C₄)-haloalkyl, (C₁-C₄)-alkoxy, (C₁-C₄)-haloalkoxy, (C₁-C₄)-alkylthio, (C₁-C₄)-alkylamino, di-(C₁-C₄)-alkyl]-amino, [(C₁-C₄)-alkoxy]-carbonyl, [(C₁-C₄)-haloalkoxy]-carbonyl, unsubstituted or substituted (C₃-C₆)-cycloalkyl, unsubstituted or substituted phenyl, and unsubstituted or substituted heterocyclyl; or

R_H^3 is (C₁-C₄)-alkoxy, (C₂-C₄)-alkenyloxy, (C₂-C₆)-alkynyloxy or (C₂-C₄)-haloalkoxy, and

25 R_H^4 is hydrogen or (C₁-C₄)-alkyl, or

R_H^3 and R_H^4 together with the directly bound N-atom are a 4 to 8-membered heterocyclic ring, which can contain further hetero ring atoms besides the N-atom, preferably up to two further hetero

ring atoms from the group consisting of N, O and S, and which is unsubstituted or substituted by one or more radicals from the group consisting of halogen, cyano, nitro, (C₁-C₄)-alkyl, (C₁-C₄)-haloalkyl, (C₁-C₄)-alkoxy, (C₁-C₄)-haloalkoxy, and (C₁-C₄)-alkylthio;

- 5 S16) Active compounds which are primarily used as herbicides, but also have a safener effect on crop plants, for example
- | | | | |
|----|--|------|-------------|
| | (2,4-dichlorophenoxy)acetic | acid | (2,4-D), |
| | (4-chlorophenoxy)acetic | | acid, |
| | (R,S)-2-(4-chloro-o-tolyloxy)propionic | acid | (mecoprop), |
| | 4-(2,4-dichlorophenoxy)butyric | acid | (2,4-DB), |
| 10 | (4-chloro-o-tolyloxy)acetic | acid | (MCPA), |
| | 4-(4-chloro-o-tolyloxy)butyric | | acid, |
| | 4-(4-chlorophenoxy)butyric | | acid, |
| | 3,6-dichloro-2-methoxybenzoic | acid | (dicamba), |
| | 1-(ethoxycarbonyl)ethyl 3,6-dichloro-2-methoxybenzoate (lactidichlor-ethyl). | | |

- 15 Preferred safeners in the context of the present invention are disclosed in group SAF-2: cloquintocet-mexyl, cyprosulfamid, fenclorazole-ethyl, isoxadifen-ethyl, mefenpyr-diethyl, fenclorim, cumyluron, S4-1, and S4-5, wherein particular preference is given to the safeners cloquintocet-mexyl, cyprosulfamid, isoxadifen-ethyl, and mefenpyr-diethyl.

- 20 Particularly preferred safeners used in the context of the present invention are disclosed in group SAF-3: Isoxadifen and its esters, in particular Isoxadifen-ethyl, and Cyprosulfamide.

The safeners are preferably used together with one or more of the herbicides mentioned above. Preferred in the context of the present invention are mixtures of safeners disclosed in group SAF-4: mixtures of Isoxadifen-ethyl and the herbicides mentioned herein and mixtures of Cyprosulfamide and the herbicides mentioned herein.

- 25 The herbicides and safeners mentioned above are known, see e.g. "The Pesticide Manual", 16th edition, November 2012, The British Crop Protection Council and the Royal Soc. of Chemistry.

The herbicides and the mixtures of herbicides (and optionally safeners) mentioned herein may be used in pre-emergence applications and/or in post-emergence applications.

- 30 In accordance with the present invention, the herbicides and the mixtures of herbicides (and optionally safeners) mentioned herein may be applied as a split application over time. Another possibility is the application of the individual active ingredients or the mixtures comprising the active ingredients in a plurality of portions (sequential application), for example after pre-emergence applications, followed by

post-emergence applications or after early post-emergence applications, followed by applications at medium or late post-emergence.

Biologics:

Biologics as used in the context of the present invention are selected from the group B-1 consisting of:

5 *Bacillus cereus* EIP-N2 (Accession No. CNCM I-1562), *Myrothecium verrucaria*, in particular strain AARC-0255, *Flavobacterium sp* strain H492 (Accession No. B-50584), *Bacillus subtilis* AQ713 (Accession No. B-21661), *Bacillus subtilis* AQ30002 (Accession No. B-50421), *Bacillus subtilis* AQ 30004 (Accession No. B-50455), *Bacillus firmus*, especially *Bacillus firmus* GB126 (Accession No. CNCM I-1582) and *Bacillus firmus* strain I-1582 (Accession No. CNCMI-1582), *Paecilomyces lilacinus*
 10 251 (Accession No. DSM 23289), *Coniothyrium minitans* (Accession No. DSM 9660), *Bacillus pumilus* QST 2808 (Accession No. NRRL B-30087), *Bacillus pumilus* GB34 (Accession No. ATCC 700814), *Metarhizium anisopliae* (Accession No. DSM 3884), *Metarhizium anisopliae* (Accession No. ATCC 90448), *Bacillus subtilis/amyloliquefaciens* MBI600 (Accession No. NRRL B-50595), MBI600 + pyraclostrobin), *Paenibacillus polymyxa* strains Lu16774 (Accession No. DSM 26969), Lu17007
 15 (Accession No. DSM 26970), and Lu17015 (Accession No. DSM 26971), *Bacillus subtilis* FZB24 (EPA Registration Number: 72098-5 and EPA Establishment Number: 73386-DEU-001, a strain that is commercially available from Earth Bioscience, Inc.), *Pasteuria usage* strains (e.g., used in ECONEM), *Trichoderma virens* GI-3 (Accession No. ATCC 58678), *Bacillus amyloliquefaciens* TJ1000 (Accession No. ATCC BAA-390), or a combination of GI-3 with TJ1000 or FZB24, *Bacillus amyloliquefaciens*
 20 D747 (Accession No. FERM BP-8234), *Bradyrhizobium japonicum*, *Pseudomonas fluorescens*, *Pseudomonas chloropsis*, *Rhizobium spp*, especially *Rhizobium tropici* SP25, a recombinant exosporium-producing *Bacillus thuringiensis* cell (especially *B. thuringiensis* BT013A) that expresses a fusion protein comprising:

- 25
- 1) an endoglucanase, (especially from *Bacillus subtilis*) having the amino acid sequence of Sequence 2 as herein defined and
 - 2) a targeting sequence having at least about 43% identity with amino acids 20–35 of Sequence 1 as herein defined, wherein the identity with amino acids 25–35 is at least about 54%

and a recombinant exosporium-producing *Bacillus thuringiensis* cell that expresses a fusion protein comprising:

- 30
- 1) a phospholipase having the amino acid sequence of Sequence 3 as herein defined and
 - 2) a targeting sequence having at least about 43% identity with amino acids 20–35 of Sequence 1 as herein defined, wherein the identity with amino acids 25–35 is at least about 54%.

The term “targeting sequence” as used herein refers to a polypeptide sequence that results in the localization of a longer polypeptide or the protein to the exosporium of a *Bacillus cereus* family
 35 member.

BclA, the major constituent of the surface nap, has been shown to be attached to the exosporium with its amino-terminus (N-terminus) positioned at the basal layer and its carboxy-terminus (C- terminus) extending outward from the spore. It was previously discovered that certain sequences from the N-terminal regions of BclA and BclB could be used to target a peptide or protein to the exosporium of a *Bacillus cereus* endospore (see, e.g., US Patent Publication Nos. 2010/0233124 and 2011/0281316, and Thompson, et al., "Targeting of the BclA and BclB Proteins to the Bacillus anthracis Spore Surface," *Molecular Microbiology*, 70(2):421–34 (2008), the entirety of each of which is hereby incorporated by reference). It was also found that the BetA/BAS3290 protein of *Bacillus anthracis* localized to the exosporium.

In particular, amino acids 20–35 of BclA from *Bacillus anthracis* Sterne strain have been found to be sufficient for targeting to the exosporium. A sequence alignment of amino acids 1–41 of BclA (Sequence 1 as herein defined) with the corresponding N-terminal regions of several other *Bacillus cereus* family exosporium proteins and *Bacillus cereus* family proteins having related sequences is shown in WO 2016/044529 as Fig. 1. There is a region of high-homology among all of the proteins in the region corresponding to amino acids 20–41 of BclA. However, in these sequences, the amino acids corresponding to amino acids 36–41 of BclA contain secondary structure and are not necessary for fusion protein localization to the exosporium.

Sequences:

Sequence 1: is identical with SEQ ID NO.: 1 of WO 2016/044529 referring to the amino acid sequence of AA 1-41 of BclA (herein incorporated by reference)

Sequence 2: is identical with SEQ ID NO.: 107 of WO 2016/044529 referring to the amino acid sequence of endonuclease (herein incorporated by reference)

Sequence 3: is identical with SEQ ID NO.: 108 of WO 2016/044529 referring to the amino acid sequence of *B. thuringiensis* BT013A phospholipase C (herein incorporated by reference)

In one preferred embodiment, Biologics are selected from the group B-2 consisting of:

Bacillus subtilis AQ713, *Bacillus firmus*, GB126, *Bacillus firmus* strain I-1582, *Paecilomyces lilacinus* 251, *Coniothyrium minitans* (Accession No. DSM 9660), *Bacillus pumilus* QST 2808, *Bacillus pumilus* GB34, *Metarhizium anisopliae* (Accession No. DSM 3884), *Metarhizium anisopliae* (Accession No. ATCC 90448), *Bacillus subtilis/amyloliquefaciens* MBI600, *Paenibacillus polymyxa* strains Lu16774, Lu17007, Lu17015, *Bacillus subtilis* FZB24, *Pasteuria usage* strains (e.g. used in ECONEM), *Trichoderma virens* GI-3, *Bacillus amyloliquefaciens* TJ1000, a combination of GI-3 with TJ1000 or FZB24, *Bacillus amyloliquefaciens* D747, a recombinant exosporium-producing *Bacillus thuringiensis* cell (especially *B. thuringiensis* BT013A) that expresses a fusion protein comprising:

- 1) an endoglucanase, (especially from *Bacillus subtilis*) having the amino acid sequence of Sequence 2 as herein defined and
 - 2) a targeting sequence having at least about 43% identity with amino acids 20–35 of Sequence 1 as herein defined, wherein the identity with amino acids 25–35 is at least about 54%
- 5 and a recombinant exosporium-producing *Bacillus thuringiensis* cell that expresses a fusion protein comprising:
- 1) a phospholipase having the amino acid sequence of Sequence 3 as herein defined and
 - 2) a targeting sequence having at least about 43% identity with amino acids 20–35 of Sequence 1 as herein defined, wherein the identity with amino acids 25–35 is at least about 54%.
- 10

In a more preferred embodiment, Biologics are selected from the group B-3 consisting of:

Bacillus subtilis AQ713, *Bacillus firmus* GB126, *Paecilomyces lilacinus* strain 251, *Coniothyrium minitans* (Accession No. DSM 9660), *Bacillus pumilus* QST 2808, *Bacillus pumilus* GB34, a
15 recombinant exosporium-producing *Bacillus thuringiensis* cell (especially *B. thuringiensis* BT013A) that expresses a fusion protein comprising:

- 1) an endoglucanase, (especially from *Bacillus subtilis*) having the amino acid sequence of Sequence 2 as herein defined and
 - 2) a targeting sequence having at least about 43% identity with amino acids 20–35 of Sequence 1 as herein defined, wherein the identity with amino acids 25–35 is at least about 54%
- 20

and a recombinant exosporium-producing *Bacillus thuringiensis* cell that expresses a fusion protein comprising:

- 1) a phospholipase having the amino acid sequence of Sequence 3 as herein defined and
 - 2) a targeting sequence having at least about 43% identity with amino acids 20–35 of Sequence 1 as herein defined, wherein the identity with amino acids 25–35 is at least about 54%.
- 25

In a further more preferred embodiment, Biologics are selected from a group B-4 consisting of:

Metarhizium anisopliae (Accession No. DSM 3884), *Metarhizium anisopliae* (Accession No. ATCC 90448), *Bacillus subtilis/amyloliquefaciens* MBI600 (Accession No. NRRL B-50595), *Paenibacillus polymyxa* strains Lu16774, Lu17007, Lu17015, *Bacillus subtilis* FZB24, *Pasteuria usage* strains (e.g. used in ECONEM), *Trichoderma virens* GI-3, *Bacillus amyloliquefaciens* TJ1000, a combination of GI-3 with TJ1000 or FZB24) and *Bacillus amyloliquefaciens* D747.

30

In an even more preferred embodiment, Biologics are selected from the group B-5 consisting of:

Bacillus subtilis AQ713, *Bacillus firmus* GB126, *Paecilomyces lilacinus* strain 251, *Coniothyrium minitans* (Accession No. DSM 9660), *Bacillus pumilus* GB34, and *Bacillus pumilus* QST 2808.

Insecticides

- 5 The Insecticides and Nematicides identified herein by their common names are known and are described, for example, in the pesticide handbook ("The Pesticide Manual" 16th Ed., British Crop Protection Council 2012) or can be found on the Internet (e.g. <http://www.alanwood.net/pesticides>). The classification is based on the current IRAC Mode of Action Classification Scheme at the time of filing of this patent application.
- 10 Insecticides used in the context of the present invention are disclosed in the group I-1 comprising 29 subgroups:
- (1) Acetylcholinesterase (AChE) inhibitors, such as, for example, carbamates, for example alanycarb, aldicarb, bendiocarb, benfuracarb, butocarboxim, butoxycarboxim, carbaryl, carbofuran, carbosulfan, ethiofencarb, fenobucarb, formetanate, furathiocarb, isoprocarb, methiocarb, methomyl, metolcarb, 15 oxamyl, pirimicarb, propoxur, thiodicarb, thiofanox, triazamate, trimethacarb, XMC and xylylcarb; or organophosphates, for example acephate, azamethiphos, azinphos-ethyl, azinphos-methyl, cadusafos, chlorethoxyfos, chlorfenvinphos, chlormephos, chlorpyrifos-methyl, coumaphos, cyanophos, demeton-S-methyl, diazinon, dichlorvos/DDVP, dicrotophos, dimethoate, dimethylvinphos, disulfoton, EPN, ethion, ethoprophos, famphur, fenamiphos, fenitrothion, fenthion, fosthiazate, heptenophos, imicyafos, 20 isofenphos, isopropyl O-(methoxyaminothiophosphoryl) salicylate, isoxathion, malathion, mecarbam, methamidophos, methidathion, mevinphos, monocrotophos, naled, omethoate, oxydemeton-methyl, parathion-methyl, phenthoate, phorate, phosalone, phosmet, phosphamidon, phoxim, pirimiphos-methyl, profenofos, propetamphos, prothiofos, pyraclofos, pyridaphenthion, quinalphos, sulfotep, tebupirimfos, temephos, terbufos, tetrachlorvinphos, thiometon, triazophos, triclofon and vamidothion;
- 25 (2) GABA-gated chloride channel blockers, such as, for example, cyclodiene-organochlorines, for example chlordane and endosulfan or phenylpyrazoles (fiproles), for example ethiprole and fipronil;
- (3) Sodium channel modulators, such as, for example, pyrethroids, e.g. acrinathrin, allethrin, d-cis-trans allethrin, d-trans allethrin, bifenthrin, bioallethrin, bioallethrin s-cyclopentenyl isomer, bioresmethrin, cycloprothrin, cyfluthrin, beta-cyfluthrin, cyhalothrin, lambda-cyhalothrin, gamma-cyhalothrin, 30 cypermethrin, alpha-cypermethrin, beta-cypermethrin, theta-cypermethrin, zeta-cypermethrin, cyphenothrin [(1R)-trans-isomer], deltamethrin, empenthrin [(EZ)-(1R)-isomer], esfenvalerate, etofenprox, fenpropathrin, fenvalerate, flucythrinate, flumethrin, tau-fluvalinate, halfenprox, imiprothrin, kadethrin, momfluorothrin, permethrin, phenothrin [(1R)-trans-isomer], prallethrin,

- pyrethrins (pyrethrum), resmethrin, silafluofen, tefluthrin, tetramethrin, tetramethrin [(1R)- isomer], tralomethrin and transfluthrin or DDT or methoxychlor;
- (4) Nicotinic acetylcholine receptor (nAChR) competitive modulators, such as, for example, neonicotinoids, e.g. acetamiprid, clothianidin, dinotefuran, imidacloprid, nitenpyram, thiacloprid and thiamethoxam or nicotine or sulfoxaflor or flupyradifurone;
- 5 (5) Nicotinic acetylcholine receptor (nAChR) allosteric modulators, such as, for example, spinosyns, e.g. spinetoram and spinosad;
- (6) Glutamate-gated chloride channel (GluCl) allosteric modulators, such as, for example, avermectins/milbemycins, for example abamectin, emamectin benzoate, lepimectin and milbemectin;
- 10 (7) Juvenile hormone mimics, such as, for example, juvenile hormone analogues, e.g. hydroprene, kinoprene and methoprene or fenoxycarb or pyriproxyfen;
- (8) Miscellaneous non-specific (multi-site) inhibitors, such as, for example, alkyl halides, e.g. methyl bromide and other alkyl halides; or chloropicrine or sulphuryl fluoride or borax or tartar emetic or methyl isocyanate generators, e.g. diazomet and metam;
- 15 (9) Modulators of Chordotonal Organs, such as, for example pymetrozine or flonicamid;
- (10) Mite growth inhibitors, such as, for example clofentezine, hexythiazox and diflovidazin or etoxazole;
- (11) Microbial disruptors of the insect gut membrane, such as, for example *Bacillus thuringiensis* subspecies *israelensis*, *Bacillus sphaericus*, *Bacillus thuringiensis* subspecies *aizawai*, *Bacillus thuringiensis* subspecies *kurstaki*, *Bacillus thuringiensis* subspecies *tenebrionis*, and B.t. plant proteins: Cry1Ab, Cry1Ac, Cry1Fa, Cry1A.105, Cry2Ab, Vip3A, mCry3A, Cry3Ab, Cry3Bb, Cry34Ab1/35Ab1;
- 20 (12) Inhibitors of mitochondrial ATP synthase, such as, ATP disruptors such as, for example, diafenthiuron or organotin compounds, for example azocyclotin, cyhexatin and fenbutatin oxide or propargite or tetradifon;
- 25 (13) Uncouplers of oxidative phosphorylation via disruption of the proton gradient, such as, for example, chlorfenapyr, DNOC and sulfluramid;
- (14) Nicotinic acetylcholine receptor channel blockers, such as, for example, bensultap, cartap hydrochloride, thiocyclam, and thiosultap-sodium;
- (15) Inhibitors of chitin biosynthesis, type 0, such as, for example, bistrifluron, chlorfluazuron, diflubenzuron, flucycloxuron, flufenoxuron, hexaflumuron, lufenuron, novaluron, noviflumuron, teflubenzuron and triflumuron;
- 30

- (16) Inhibitors of chitin biosynthesis, type 1, for example buprofezin;
- (17) Moulting disruptor (in particular for Diptera, i.e. dipterans), such as, for example, cyromazine;
- (18) Ecdysone receptor agonists, such as, for example, chromafenozide, halofenozide, methoxyfenozide and tebufenozide;
- 5 (19) Octopamine receptor agonists, such as, for example, amitraz;
- (20) Mitochondrial complex III electron transport inhibitors, such as, for example, hydramethylnone or acequinocyl or fluacrypyrim;
- (21) Mitochondrial complex I electron transport inhibitors, such as, for example from the group of the METI acaricides, e.g. fenazaquin, fenpyroximate, pyrimidifen, pyridaben, tebufenpyrad and tolfenpyrad
10 or rotenone (Derris);
- (22) Voltage-dependent sodium channel blockers, such as, for example indoxacarb or metaflumizone;
- (23) Inhibitors of acetyl CoA carboxylase, such as, for example, tetrionic and tetramic acid derivatives, e.g. spiroadiclofen, spiromesifen and spirotetramat;
- (24) Mitochondrial complex IV electron transport inhibitors, such as, for example, phosphines, e.g.
15 aluminium phosphide, calcium phosphide, phosphine and zinc phosphide or cyanides, e.g. calcium cyanide, potassium cyanide and sodium cyanide;
- (25) Mitochondrial complex II electron transport inhibitors, such as, for example, beta-ketonitrile derivatives, e.g. cyenopyrafen and cyflumetofen and carboxanilides, such as, for example, pyflubumide;
- (26) Ryanodine receptor modulators, such as, for example, diamides, e.g. chlorantraniliprole,
20 cyantraniliprole and flubendiamide,
- (27) further active compounds such as, for example, Afidopyropen, Afoxolaner, Azadirachtin, Benclothiaz, Benzoximate, Bifenazate, Broflanilide, Bromopropylate, Chinomethionat, Chloroprallethrin, Cryolite, Cyclaniliprole, Cycloxaprid, Cyhalodiamide, Dicloromezotiaz, Dicofol, epsilon-Metofluthrin, epsilon-Momfluthrin, Flometoquin, Fluazaindolizine, Fluensulfone, Flufenerim,
25 Flufenoxystrobin, Flufiprole, Fluhexafon, Fluopyram, Fluralaner, Fluxametamide, Fufenozide, Guadipyr, Heptafluthrin, Imidaclathiz, Iprodione, kappa-Bifenthrin, kappa-Tefluthrin, Lotilaner, Meperfluthrin, Paichongding, Pyridalyl, Pyrifluquinazon, Pyriminostrobin, Spirobudiclofen, Tetramethylfluthrin, Tetraniliprole, Tetrachlorantraniliprole, Tioxazafen, Thiofluoximate, Triflumezopyrim and iodomethane; furthermore preparations based on Bacillus firmus (I-1582,
30 BioNeem, Votivo), and also the following compounds: 1-{2-fluoro-4-methyl-5-[(2,2,2-trifluoroethyl)sulphinyl]phenyl}-3-(trifluoromethyl)-1H-1,2,4-triazole-5-amine (known from

WO2006/043635) (CAS 885026-50-6), {1'-[(2E)-3-(4-chlorophenyl)prop-2-en-1-yl]-5-fluorospiro[indol-3,4'-piperidin]-1(2H)-yl}(2-chloropyridin-4-yl)methanone (known from WO2003/106457) (CAS 637360-23-7), 2-chloro-N-[2-{1-[(2E)-3-(4-chlorophenyl)prop-2-en-1-yl]piperidin-4-yl}-4-(trifluoromethyl)phenyl]isonicotinamide (known from WO2006/003494) (CAS 872999-66-1), 3-(4-chloro-2,6-dimethylphenyl)-4-hydroxy-8-methoxy-1,8-diazaspiro[4.5]dec-3-en-2-one (known from WO 2010052161) (CAS 1225292-17-0), 3-(4-chloro-2,6-dimethylphenyl)-8-methoxy-2-oxo-1,8-diazaspiro[4.5]dec-3-en-4-yl ethyl carbonate (known from EP2647626) (CAS 1440516-42-6), 4-(but-2-yn-1-yloxy)-6-(3,5-dimethylpiperidin-1-yl)-5-fluoropyrimidine (known from WO2004/099160) (CAS 792914-58-0), PF1364 (known from JP2010/018586) (CAS 1204776-60-2), N-[(2E)-1-[(6-chloropyridin-3-yl)methyl]pyridin-2(1H)-ylidene]-2,2,2-trifluoroacetamide (known from WO2012/029672) (CAS 1363400-41-2), (3E)-3-[1-[(6-chloro-3-pyridyl)methyl]-2-pyridylidene]-1,1,1-trifluoro-propan-2-one (known from WO2013/144213) (CAS 1461743-15-6), N-[3-(benzylcarbamoyl)-4-chlorophenyl]-1-methyl-3-(pentafluoroethyl)-4-(trifluoromethyl)-1H-pyrazole-5-carboxamide (known from WO2010/051926) (CAS 1226889-14-0), 5-bromo-4-chloro-N-[4-chloro-2-methyl-6-(methylcarbamoyl)phenyl]-2-(3-chloro-2-pyridyl)pyrazole-3-carboxamide (known from CN103232431) (CAS 1449220-44-3), 4-[5-(3,5-dichlorophenyl)-4,5-dihydro-5-(trifluoromethyl)-3-isoxazolyl]-2-methyl-N-(cis-1-oxido-3-thietanyl)-benzamide, 4-[5-(3,5-dichlorophenyl)-4,5-dihydro-5-(trifluoromethyl)-3-isoxazolyl]-2-methyl-N-(trans-1-oxido-3-thietanyl)-benzamide and 4-[(5S)-5-(3,5-dichlorophenyl)-4,5-dihydro-5-(trifluoromethyl)-3-isoxazolyl]-2-methyl-N-(cis-1-oxido-3-thietanyl) benzamide (known from WO 2013/050317 A1) (CAS 1332628-83-7), N-[3-chloro-1-(3-pyridinyl)-1H-pyrazol-4-yl]-N-ethyl-3-[(3,3,3-trifluoropropyl)sulfinyl]-propanamide, (+)-N-[3-chloro-1-(3-pyridinyl)-1H-pyrazol-4-yl]-N-ethyl-3-[(3,3,3-trifluoropropyl)sulfinyl]-propanamide and (-)-N-[3-chloro-1-(3-pyridinyl)-1H-pyrazol-4-yl]-N-ethyl-3-[(3,3,3-trifluoropropyl)sulfinyl]-propanamide (known from WO 2013/162715 A2, WO 2013/162716 A2, US 2014/0213448 A1) (CAS 1477923-37-7), 5-[[[(2E)-3-chloro-2-propen-1-yl]amino]-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-4-[(trifluoromethyl)sulfinyl]-1H-pyrazole-3-carbonitrile (known from CN 101337937 A) (CAS 1105672-77-2), 3-bromo-N-[4-chloro-2-methyl-6-[(methylamino)thioxomethyl]phenyl]-1-(3-chloro-2-pyridinyl)-1H-pyrazole-5-carboxamide, (Liudaibenjiaxuanan, known from CN 103109816 A) (CAS 1232543-85-9); N-[4-chloro-2-[[[(1,1-dimethylethyl)amino]carbonyl]-6-methylphenyl]-1-(3-chloro-2-pyridinyl)-3-(fluoromethoxy)-1H-Pyrazole-5-carboxamide (known from WO 2012/034403 A1) (CAS 1268277-22-0), N-[2-(5-amino-1,3,4-thiadiazol-2-yl)-4-chloro-6-methylphenyl]-3-bromo-1-(3-chloro-2-pyridinyl)-1H-pyrazole-5-carboxamide (known from WO 2011/085575 A1) (CAS 1233882-22-8), 4-[3-[2,6-dichloro-4-[(3,3-dichloro-2-propen-1-yl)oxy]phenoxy]propoxy]-2-methoxy-6-(trifluoromethyl)-pyrimidine (known from CN 101337940 A) (CAS 1108184-52-6); (2E)- and 2(Z)-2-[2-(4-cyanophenyl)-1-[3-(trifluoromethyl)phenyl]ethylidene]-N-[4-(difluoromethoxy)phenyl]-hydrazinecarboxamide (known from CN 101715774 A) (CAS 1232543-85-9); 3-(2,2-dichloroethenyl)-2,2-dimethyl-4-(1H-benzimidazol-2-yl)phenyl-cyclopropanecarboxylic acid ester (known from CN 103524422 A) (CAS 1542271-46-4); (4aS)-7-chloro-2,5-dihydro-2-[[methoxycarbonyl]4-[(trifluoromethyl)thio]phenyl]amino]carbonyl]-

indeno[1,2-e][1,3,4]oxadiazine-4a(3H)-carboxylic acid methyl ester (known from CN 102391261 A) (CAS 1370358-69-2); 6-deoxy-3-O-ethyl-2,4-di-O-methyl-, 1-[N-[4-[1-[4-(1,1,2,2,2-pentafluoroethoxy) phenyl]-1H-1,2,4-triazol-3-yl]phenyl]carbamate]- α -L-mannopyranose (known from US 2014/0275503 A1) (CAS 1181213-14-8); 8-(2-cyclopropylmethoxy-4-trifluoromethyl-phenoxy)-3-(6-trifluoromethyl-pyridazin-3-yl)-3-aza-bicyclo[3.2.1]octane (CAS 1253850-56-4), (8-anti)-8-(2-cyclopropylmethoxy-4-trifluoromethyl-phenoxy)-3-(6-trifluoromethyl-pyridazin-3-yl)-3-aza-bicyclo[3.2.1]octane (CAS 933798-27-7), (8-syn)-8-(2-cyclopropylmethoxy-4-trifluoromethyl-phenoxy)-3-(6-trifluoromethyl-pyridazin-3-yl)-3-aza-bicyclo[3.2.1]octane (known from WO 2007040280 A1, WO 2007040282 A1) (CAS 934001-66-8) and N-[3-chloro-1-(3-pyridinyl)-1H-pyrazol-4-yl]-N-ethyl-3-[(3,3,3-trifluoropropyl)thio]-propanamide (known from WO 2015/058021 A1, WO 2015/058028 A1) (CAS 1477919-27-9).

The Insecticides and combinations thereof may be used in a curative or preventive manner.

Insecticides especially used in the context of the present invention are selected from the group I-2 consisting of:

15 Abamectin, Al Phosphide, Azadirachtin, *Bacillus firmus* strain I-1582, Cadusafos, Carbofuran, Chloropicrin, Dichloropropene, Dimethyl-sulfid, Clothianidin, Spirotetramat, Flupyradifurone, Tetraniliprole, *Bacillus firmus* GB126, Imidacloprid, Flubendiamide, Deltamethrin, Thiacloprid, Spiromesifen, Ethiprole, Ethoprophos, *Paecilomyces lilacinus* 251, Terpenoid blend (e.g., natural extracts or simulated blend of *Chenopodium ambrosioides* (product known as Requiem®, which
20 contains a mixture of three terpenes, i.e. α -terpinene (around 10%), p-cymene (around 3.75%) and limonene (around 3%), Fenamiphos, *Flavobacterium sp* strain H492, Fipronil, Fluazaindolizine, Fluensulfone, Fluopyram, Fosthiazate, Beta-Cyfluthrin, Thiodicarb, Triflumuron, Methiocarb, Cyfluthrin, Spirodiclofen, Silafluofen, Fenthion, Imicyafos, Iprodione, Metam-Sodium, Methylbromide, *Myrothecium verrucaria* in particular strain AARC-0255, Oxamyl, Sulphuryl fluoride, Tioxazafen,
25 Inclosamide, Tralomethrin, Triafamone, Carbendazim, Carboxin, Difenoconazole, Ethoxysulfuron, Fenamidon, Fenoxaprop-P-Ethyl, Fluopicolide, Fluoxastrobin, Fluquinconazole, Fosetyl-Al, Prochloraz, Propineb, Prothioconazole, Spirotetramat, Tebuconazole, Tembotrione, Thiacloprid, Thiodicarb, Thiram, Triadimenol, Trifloxystrobin, Triflumuron, Cyflumetofen, Afidopyropen, Broflanilide, Sulfoxaflor, Cyazapyr, Rynaxypyr, Fluazaindolizine, Triflumezopyrim and Tioxazafen.

30 In one preferred embodiment, Insecticides are selected from the group I-3 consisting of:

Clothianidin, Spirotetramat, Flupyradifurone, Tetraniliprole, *Bacillus firmus* GB126, Imidacloprid, Flubendiamide, Deltamethrin, Thiacloprid, Spiromesifen, Ethiprole, *Paecilomyces lilacinus* 251, Terpenoid blend (e.g. Requiem®), Fipronil, Beta-Cyfluthrin, Thiodicarb, Triflumuron, Methiocarb, Cyfluthrin, Cyflumetofen, Afidopyropen, Broflanilide, Sulfoxaflor, Cyazapyr, Rynaxypyr,
35 Fluazaindolizine, Triflumezopyrim and Tioxazafen.

In a more preferred embodiment, Insecticides are selected from the group I-4 consisting of:

Clothianidin, Spirotetramat, Flupyradifurone, Tetraniliprole, *Bacillus firmus* GB126, Imidacloprid, Flubendiamide, Deltamethrin, Thiacloprid, Spiromesifen, Ethiprole, *Paecilomyces lilacinus* 251, Terpenoid blend (e.g. Requiem®) and Fipronil.

5 In another more preferred embodiment, Insecticides are selected from the group I-5 consisting of:

Cyflumetofen, Afidopyropen, Broflanilide, Sulfoxaflor, Cyazypyr, Rynaxypyr, Fluazaindolizine, Triflumezopyrim and Tioxazafen.

In an even more preferred embodiment, Insecticides are selected from the group I-6 consisting of:

Clothianidin, Spirotetramat, Flupyradifurone, Tetraniliprole and *Bacillus firmus* GB126.

10 Fungicides

The active ingredients specified herein by their Common Name are known and described, for example, in The Pesticide Manual (16th Ed. British Crop Protection Council) or can be searched in the internet (e.g. www.alanwood.net/pesticides).

In one embodiment the fungicidal active ingredient are selected from the group F-1 consisting of fifteen
15 subgroups being

- 1) Inhibitors of the ergosterol biosynthesis, for example (1.001) cyproconazole, (1.002) difenoconazole, (1.003) epoxiconazole, (1.004) fenhexamid, (1.005) fenpropidin, (1.006) fenpropimorph, (1.007) fenpyrazamine, (1.008) fluquinconazole, (1.009) flutriafol, (1.010) imazalil, (1.011) imazalil sulfate, (1.012) ipconazole, (1.013) metconazole, (1.014) myclobutanil, (1.015) paclobutrazol, (1.016) prochloraz, (1.017) propiconazole, (1.018) prothioconazole, (1.019) Pyrisoxazole, (1.020) spiroxamine, (1.021) tebuconazole, (1.022) tetraconazole, (1.023) triadimenol, (1.024) tridemorph, (1.025) triticonazole, (1.026) (1R,2S,5S)-5-(4-chlorobenzyl)-2-(chloromethyl)-2-methyl-1-(1H-1,2,4-triazol-1-ylmethyl)cyclopentanol, (1.027) (1S,2R,5R)-5-(4-chlorobenzyl)-2-(chloromethyl)-2-methyl-1-(1H-1,2,4-triazol-1-ylmethyl)cyclopentanol, (1.028) (2R)-2-(1-chlorocyclopropyl)-4-[(1R)-2,2-dichlorocyclopropyl]-1-(1H-1,2,4-triazol-1-yl)butan-2-ol, (1.029) (2R)-2-(1-chlorocyclopropyl)-4-[(1S)-2,2-dichlorocyclopropyl]-1-(1H-1,2,4-triazol-1-yl)butan-2-ol, (1.030) (2R)-2-[4-(4-chlorophenoxy)-2-(trifluoromethyl)phenyl]-1-(1H-1,2,4-triazol-1-yl)propan-2-ol, (1.031) (2S)-2-(1-chlorocyclopropyl)-4-[(1R)-2,2-dichlorocyclopropyl]-1-(1H-1,2,4-triazol-1-yl)butan-2-ol, (1.032) (2S)-2-(1-chlorocyclopropyl)-4-[(1S)-2,2-dichlorocyclopropyl]-1-(1H-1,2,4-triazol-1-yl)butan-2-ol, (1.033) (2S)-2-[4-(4-chlorophenoxy)-2-(trifluoromethyl)phenyl]-1-(1H-1,2,4-triazol-1-yl)propan-2-ol, (1.034) (R)-[3-(4-chloro-2-fluorophenyl)-5-(2,4-difluorophenyl)-1,2-oxazol-4-yl](pyridin-3-yl)methanol, (1.035) (S)-[3-(4-chloro-2-fluorophenyl)-5-(2,4-difluorophenyl)-1,2-oxazol-4-yl](pyridin-3-yl)methanol, (1.036) [3-(4-chloro-2-fluorophenyl)-5-(2,4-difluorophenyl)-1,2-oxazol-4-yl](pyridin-3-yl)methanol, (1.037) 1-

5
 10
 15
 20
 25
 30
 35

([(2R,4S)-2-[2-chloro-4-(4-chlorophenoxy)phenyl]-4-methyl-1,3-dioxolan-2-yl]methyl)-1H-1,2,4-triazole, (1.038) 1-([(2S,4S)-2-[2-chloro-4-(4-chlorophenoxy)phenyl]-4-methyl-1,3-dioxolan-2-yl]methyl)-1H-1,2,4-triazole, (1.039) 1-[[3-(2-chlorophenyl)-2-(2,4-difluorophenyl)oxiran-2-yl]methyl]-1H-1,2,4-triazol-5-yl thiocyanate, (1.040) 1-[[rel(2R,3R)-3-(2-chlorophenyl)-2-(2,4-difluorophenyl)oxiran-2-yl]methyl]-1H-1,2,4-triazol-5-yl thiocyanate, (1.041) 1-[[rel(2R,3S)-3-(2-chlorophenyl)-2-(2,4-difluorophenyl)oxiran-2-yl]methyl]-1H-1,2,4-triazol-5-yl thiocyanate, (1.042) 2-[(2R,4R,5R)-1-(2,4-dichlorophenyl)-5-hydroxy-2,6,6-trimethylheptan-4-yl]-2,4-dihydro-3H-1,2,4-triazole-3-thione, (1.043) 2-[(2R,4R,5S)-1-(2,4-dichlorophenyl)-5-hydroxy-2,6,6-trimethylheptan-4-yl]-2,4-dihydro-3H-1,2,4-triazole-3-thione, (1.044) 2-[(2R,4S,5R)-1-(2,4-dichlorophenyl)-5-hydroxy-2,6,6-trimethylheptan-4-yl]-2,4-dihydro-3H-1,2,4-triazole-3-thione, (1.045) 2-[(2R,4S,5S)-1-(2,4-dichlorophenyl)-5-hydroxy-2,6,6-trimethylheptan-4-yl]-2,4-dihydro-3H-1,2,4-triazole-3-thione, (1.046) 2-[(2S,4R,5R)-1-(2,4-dichlorophenyl)-5-hydroxy-2,6,6-trimethylheptan-4-yl]-2,4-dihydro-3H-1,2,4-triazole-3-thione, (1.047) 2-[(2S,4R,5S)-1-(2,4-dichlorophenyl)-5-hydroxy-2,6,6-trimethylheptan-4-yl]-2,4-dihydro-3H-1,2,4-triazole-3-thione, (1.048) 2-[(2S,4S,5R)-1-(2,4-dichlorophenyl)-5-hydroxy-2,6,6-trimethylheptan-4-yl]-2,4-dihydro-3H-1,2,4-triazole-3-thione, (1.049) 2-[(2S,4S,5S)-1-(2,4-dichlorophenyl)-5-hydroxy-2,6,6-trimethylheptan-4-yl]-2,4-dihydro-3H-1,2,4-triazole-3-thione, (1.050) 2-[1-(2,4-dichlorophenyl)-5-hydroxy-2,6,6-trimethylheptan-4-yl]-2,4-dihydro-3H-1,2,4-triazole-3-thione, (1.051) 2-[2-chloro-4-(2,4-dichlorophenoxy)phenyl]-1-(1H-1,2,4-triazol-1-yl)propan-2-ol, (1.052) 2-[2-chloro-4-(4-chlorophenoxy)phenyl]-1-(1H-1,2,4-triazol-1-yl)butan-2-ol, (1.053) 2-[4-(4-chlorophenoxy)-2-(trifluoromethyl)phenyl]-1-(1H-1,2,4-triazol-1-yl)butan-2-ol, (1.054) 2-[4-(4-chlorophenoxy)-2-(trifluoromethyl)phenyl]-1-(1H-1,2,4-triazol-1-yl)pentan-2-ol, (1.055) 2-[4-(4-chlorophenoxy)-2-(trifluoromethyl)phenyl]-1-(1H-1,2,4-triazol-1-yl)propan-2-ol, (1.056) 2-[[3-(2-chlorophenyl)-2-(2,4-difluorophenyl)oxiran-2-yl]methyl]-2,4-dihydro-3H-1,2,4-triazole-3-thione, (1.057) 2-[[rel(2R,3R)-3-(2-chlorophenyl)-2-(2,4-difluorophenyl)oxiran-2-yl]methyl]-2,4-dihydro-3H-1,2,4-triazole-3-thione, (1.058) 2-[[rel(2R,3S)-3-(2-chlorophenyl)-2-(2,4-difluorophenyl)oxiran-2-yl]methyl]-2,4-dihydro-3H-1,2,4-triazole-3-thione, (1.059) 5-(4-chlorobenzyl)-2-(chloromethyl)-2-methyl-1-(1H-1,2,4-triazol-1-ylmethyl)cyclopentanol, (1.060) 5-(allylsulfanyl)-1-[[3-(2-chlorophenyl)-2-(2,4-difluorophenyl)oxiran-2-yl]methyl]-1H-1,2,4-triazole, (1.061) 5-(allylsulfanyl)-1-[[rel(2R,3R)-3-(2-chlorophenyl)-2-(2,4-difluorophenyl)oxiran-2-yl]methyl]-1H-1,2,4-triazole, (1.062) 5-(allylsulfanyl)-1-[[rel(2R,3S)-3-(2-chlorophenyl)-2-(2,4-difluorophenyl)oxiran-2-yl]methyl]-1H-1,2,4-triazole, (1.063) N'-(2,5-dimethyl-4-[[3-(1,1,2,2-tetrafluoroethoxy)phenyl]sulfanyl]phenyl)-N-ethyl-N-methylimidoforamide, (1.064) N'-(2,5-dimethyl-4-[[3-(2,2,2-trifluoroethoxy)phenyl]sulfanyl]phenyl)-N-ethyl-N-methylimidoforamide, (1.065) N'-(2,5-dimethyl-4-[[3-(2,2,3,3-tetrafluoropropoxy)phenyl]sulfanyl]phenyl)-N-ethyl-N-methylimidoforamide, (1.066) N'-(2,5-dimethyl-4-[[3-(pentafluoroethoxy)phenyl]sulfanyl]phenyl)-N-ethyl-N-methylimidoforamide, (1.067) N'-(2,5-dimethyl-4-[[3-[(1,1,2,2-tetrafluoroethyl)sulfanyl]phenoxy]phenyl)-N-ethyl-N-methylimidoforamide, (1.068) N'-(2,5-dimethyl-4-[[3-[(2,2,2-trifluoroethyl)sulfanyl]phenoxy]phenyl)-N-ethyl-N-methylimidoforamide, (1.069) N'-(2,5-dimethyl-4-[[3-[(2,2,3,3-

- tetrafluoropropyl)sulfanyl]phenoxy}phenyl)-N-ethyl-N-methylimidoforamamide, (1.070) N'-(2,5-dimethyl-4-{3-[(pentafluoroethyl)sulfanyl]phenoxy}phenyl)-N-ethyl-N-methylimidoforamamide, (1.071) N'-(2,5-dimethyl-4-phenoxyphenyl)-N-ethyl-N-methylimidoforamamide, (1.072) N'-(4-{3-(difluoromethoxy)phenyl}sulfanyl}-2,5-dimethylphenyl)-N-ethyl-N-methylimidoforamamide, (1.073) N'-(4-{3-[(difluoromethyl)sulfanyl]phenoxy}-2,5-dimethylphenyl)-N-ethyl-N-methylimidoforamamide, (1.074) N'-[5-bromo-6-(2,3-dihydro-1H-inden-2-yloxy)-2-methylpyridin-3-yl]-N-ethyl-N-methylimidoforamamide, (1.075) N'-{4-[(4,5-dichloro-1,3-thiazol-2-yl)oxy]-2,5-dimethylphenyl}-N-ethyl-N-methylimidoforamamide, (1.076) N'-{5-bromo-6-[(1R)-1-(3,5-difluorophenyl)ethoxy]-2-methylpyridin-3-yl}-N-ethyl-N-methylimidoforamamide, (1.077) N'-{5-bromo-6-[(1S)-1-(3,5-difluorophenyl)ethoxy]-2-methylpyridin-3-yl}-N-ethyl-N-methylimidoforamamide, (1.078) N'-{5-bromo-6-[(cis-4-isopropylcyclohexyl)oxy]-2-methylpyridin-3-yl}-N-ethyl-N-methylimidoforamamide, (1.079) N'-{5-bromo-6-[(trans-4-isopropylcyclohexyl)oxy]-2-methylpyridin-3-yl}-N-ethyl-N-methylimidoforamamide, (1.080) N'-{5-bromo-6-[1-(3,5-difluorophenyl)ethoxy]-2-methylpyridin-3-yl}-N-ethyl-N-methylimidoforamamide, (1.081) Mefentrifluconazole, (1.082) Ipfentrifluconazole;
- 2) Inhibitors of the respiratory chain at complex I or II, for example (2.001) benzovindiflupyr, (2.002) bixafen, (2.003) boscalid, (2.004) carboxin, (2.005) fluopyram, (2.006) flutolanil, (2.007) fluxapyroxad, (2.008) furametpyr, (2.009) Isofetamid, (2.010) isopyrazam (anti-epimeric enantiomer 1R,4S,9S), (2.011) isopyrazam (anti-epimeric enantiomer 1S,4R,9R), (2.012) isopyrazam (anti-epimeric racemate 1RS,4SR,9SR), (2.013) isopyrazam (mixture of syn-epimeric racemate 1RS,4SR,9RS and anti-epimeric racemate 1RS,4SR,9SR), (2.014) isopyrazam (syn-epimeric enantiomer 1R,4S,9R), (2.015) isopyrazam (syn-epimeric enantiomer 1S,4R,9S), (2.016) isopyrazam (syn-epimeric racemate 1RS,4SR,9RS), (2.017) penflufen, (2.018) penthiopyrad, (2.019) pydiflumetofen, (2.020) Pyraziflumid, (2.021) sedaxane, (2.022) 1,3-dimethyl-N-(1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl)-1H-pyrazole-4-carboxamide, (2.023) 1,3-dimethyl-N-[(3R)-1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl]-1H-pyrazole-4-carboxamide, (2.024) 1,3-dimethyl-N-[(3S)-1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl]-1H-pyrazole-4-carboxamide, (2.025) 1-methyl-3-(trifluoromethyl)-N-[2'-(trifluoromethyl)biphenyl-2-yl]-1H-pyrazole-4-carboxamide, (2.026) 2-fluoro-6-(trifluoromethyl)-N-(1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl)benzamide, (2.027) 3-(difluoromethyl)-1-methyl-N-(1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl)-1H-pyrazole-4-carboxamide, (2.028) 3-(difluoromethyl)-1-methyl-N-[(3R)-1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl]-1H-pyrazole-4-carboxamide, (2.029) 3-(difluoromethyl)-1-methyl-N-[(3S)-1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl]-1H-pyrazole-4-carboxamide, (2.030) 3-(difluoromethyl)-N-(7-fluoro-1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl)-1-methyl-1H-pyrazole-4-carboxamide, (2.031) 3-(difluoromethyl)-N-[(3R)-7-fluoro-1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl]-1-methyl-1H-pyrazole-4-carboxamide, (2.032) 3-(difluoromethyl)-N-[(3S)-7-fluoro-1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl]-1-methyl-1H-pyrazole-4-carboxamide, (2.033) 5,8-difluoro-N-[2-(2-fluoro-4-{[4-(trifluoromethyl)pyridin-2-yl]oxy}phenyl)ethyl]quinazolin-4-amine, (2.034) N-(2-cyclopentyl-5-fluorobenzyl)-N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide, (2.035) N-(2-tert-butyl-5-methylbenzyl)-N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1-methyl-1H-

pyrazole-4-carboxamide, (2.036) N-(2-tert-butylbenzyl)-N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide, (2.037) N-(5-chloro-2-ethylbenzyl)-N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide, (2.038) N-(5-chloro-2-isopropylbenzyl)-N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide, (2.039) N-[(1R,4S)-9-(dichloromethylene)-1,2,3,4-tetrahydro-1,4-methanonaphthalen-5-yl]-3-(difluoromethyl)-1-methyl-1H-pyrazole-4-carboxamide, (2.040) N-[(1S,4R)-9-(dichloromethylene)-1,2,3,4-tetrahydro-1,4-methanonaphthalen-5-yl]-3-(difluoromethyl)-1-methyl-1H-pyrazole-4-carboxamide, (2.041) N-[1-(2,4-dichlorophenyl)-1-methoxypropan-2-yl]-3-(difluoromethyl)-1-methyl-1H-pyrazole-4-carboxamide, (2.042) N-[2-chloro-6-(trifluoromethyl)benzyl]-N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide, (2.043) N-[3-chloro-2-fluoro-6-(trifluoromethyl)benzyl]-N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide, (2.044) N-[5-chloro-2-(trifluoromethyl)benzyl]-N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide, (2.045) N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1-methyl-N-[5-methyl-2-(trifluoromethyl)benzyl]-1H-pyrazole-4-carboxamide, (2.046) N-cyclopropyl-3-(difluoromethyl)-5-fluoro-N-(2-fluoro-6-isopropylbenzyl)-1-methyl-1H-pyrazole-4-carboxamide, (2.047) N-cyclopropyl-3-(difluoromethyl)-5-fluoro-N-(2-isopropyl-5-methylbenzyl)-1-methyl-1H-pyrazole-4-carboxamide, (2.048) N-cyclopropyl-3-(difluoromethyl)-5-fluoro-N-(2-isopropylbenzyl)-1-methyl-1H-pyrazole-4-carbothioamide, (2.049) N-cyclopropyl-3-(difluoromethyl)-5-fluoro-N-(2-isopropylbenzyl)-1-methyl-1H-pyrazole-4-carboxamide, (2.050) N-cyclopropyl-3-(difluoromethyl)-5-fluoro-N-(5-fluoro-2-isopropylbenzyl)-1-methyl-1H-pyrazole-4-carboxamide, (2.051) N-cyclopropyl-3-(difluoromethyl)-N-(2-ethyl-4,5-dimethylbenzyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide, (2.052) N-cyclopropyl-3-(difluoromethyl)-N-(2-ethyl-5-fluorobenzyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide, (2.053) N-cyclopropyl-3-(difluoromethyl)-N-(2-ethyl-5-methylbenzyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide, (2.054) N-cyclopropyl-N-(2-cyclopropyl-5-fluorobenzyl)-3-(difluoromethyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide, (2.055) N-cyclopropyl-N-(2-cyclopropyl-5-methylbenzyl)-3-(difluoromethyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide, (2.056) N-cyclopropyl-N-(2-cyclopropylbenzyl)-3-(difluoromethyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide;

3) Inhibitors of the respiratory chain at complex III, for example (3.001) ametocradin, (3.002) amisulbrom, (3.003) azoxystrobin, (3.004) coumethoxystrobin, (3.005) coumoxystrobin, (3.006) cyazofamid, (3.007) dimoxystrobin, (3.008) enoxastrobin, (3.009) famoxadone, (3.010) fenamidone, (3.011) flufenoxystrobin, (3.012) fluoxastrobin, (3.013) kresoxim-methyl, (3.014) metominostrobin, (3.015) orysastrobin, (3.016) picoxystrobin, (3.017) pyraclostrobin, (3.018) pyrametostrobin, (3.019) pyraoxystrobin, (3.020) trifloxystrobin, (3.021) (2E)-2-{2-[[[(1E)-1-(3-[(E)-1-fluoro-2-phenylvinyl]oxy)phenyl]ethylidene]amino]oxy)methyl]phenyl}-2-(methoxyimino)-N-methylacetamide, (3.022) (2E,3Z)-5-[[1-(4-chlorophenyl)-1H-pyrazol-3-yl]oxy]-2-(methoxyimino)-N,3-dimethylpent-3-enamide, (3.023) (2R)-2-{2-[(2,5-dimethylphenoxy)methyl]phenyl}-2-methoxy-N-methylacetamide, (3.024) (2S)-2-{2-[(2,5-dimethylphenoxy)methyl]phenyl}-2-methoxy-N-methylacetamide, (3.025)

(3S,6S,7R,8R)-8-benzyl-3-[(3-[(isobutyryloxy)methoxy]-4-methoxypyridin-2-yl]carbonyl)amino]-6-methyl-4,9-dioxo-1,5-dioxonan-7-yl 2-methylpropanoate, (3.026) 2-{2-[(2,5-dimethylphenoxy)methyl]phenyl}-2-methoxy-N-methylacetamide, (3.027) N-(3-ethyl-3,5,5-trimethylcyclohexyl)-3-formamido-2-hydroxybenzamide, (3.028) (2E,3Z)-5-[[1-(4-chloro-2-fluorophenyl)-1H-pyrazol-3-yl]oxy]-2-(methoxyimino)-N,3-dimethylpent-3-enamide, (3.029) methyl {5-[3-(2,4-dimethylphenyl)-1H-pyrazol-1-yl]-2-methylbenzyl}carbamate;

4) Inhibitors of the mitosis and cell division, for example (4.001) carbendazim, (4.002) diethofencarb, (4.003) ethaboxam, (4.004) fluopicolide, (4.005) pencycuron, (4.006) thiabendazole, (4.007) thiophanate-methyl, (4.008) zoxamide, (4.009) 3-chloro-4-(2,6-difluorophenyl)-6-methyl-5-phenylpyridazine, (4.010) 3-chloro-5-(4-chlorophenyl)-4-(2,6-difluorophenyl)-6-methylpyridazine, (4.011) 3-chloro-5-(6-chloropyridin-3-yl)-6-methyl-4-(2,4,6-trifluorophenyl)pyridazine, (4.012) 4-(2-bromo-4-fluorophenyl)-N-(2,6-difluorophenyl)-1,3-dimethyl-1H-pyrazol-5-amine, (4.013) 4-(2-bromo-4-fluorophenyl)-N-(2-bromo-6-fluorophenyl)-1,3-dimethyl-1H-pyrazol-5-amine, (4.014) 4-(2-bromo-4-fluorophenyl)-N-(2-bromophenyl)-1,3-dimethyl-1H-pyrazol-5-amine, (4.015) 4-(2-bromo-4-fluorophenyl)-N-(2-chloro-6-fluorophenyl)-1,3-dimethyl-1H-pyrazol-5-amine, (4.016) 4-(2-bromo-4-fluorophenyl)-N-(2-chlorophenyl)-1,3-dimethyl-1H-pyrazol-5-amine, (4.017) 4-(2-bromo-4-fluorophenyl)-N-(2-fluorophenyl)-1,3-dimethyl-1H-pyrazol-5-amine, (4.018) 4-(2-chloro-4-fluorophenyl)-N-(2,6-difluorophenyl)-1,3-dimethyl-1H-pyrazol-5-amine, (4.019) 4-(2-chloro-4-fluorophenyl)-N-(2-chloro-6-fluorophenyl)-1,3-dimethyl-1H-pyrazol-5-amine, (4.020) 4-(2-chloro-4-fluorophenyl)-N-(2-chlorophenyl)-1,3-dimethyl-1H-pyrazol-5-amine, (4.021) 4-(2-chloro-4-fluorophenyl)-N-(2-fluorophenyl)-1,3-dimethyl-1H-pyrazol-5-amine, (4.022) 4-(4-chlorophenyl)-5-(2,6-difluorophenyl)-3,6-dimethylpyridazine, (4.023) N-(2-bromo-6-fluorophenyl)-4-(2-chloro-4-fluorophenyl)-1,3-dimethyl-1H-pyrazol-5-amine, (4.024) N-(2-bromophenyl)-4-(2-chloro-4-fluorophenyl)-1,3-dimethyl-1H-pyrazol-5-amine, (4.025) N-(4-chloro-2,6-difluorophenyl)-4-(2-chloro-4-fluorophenyl)-1,3-dimethyl-1H-pyrazol-5-amine;

5) Compounds capable to have a multisite action, for example (5.001) bordeaux mixture, (5.002) captafol, (5.003) captan, (5.004) chlorothalonil, (5.005) copper hydroxide, (5.006) copper naphthenate, (5.007) copper oxide, (5.008) copper oxychloride, (5.009) copper(2+) sulfate, (5.010) dithianon, (5.011) dodine, (5.012) folpet, (5.013) mancozeb, (5.014) maneb, (5.015) metiram, (5.016) metiram zinc, (5.017) oxine-copper, (5.018) propineb, (5.019) sulfur and sulfur preparations including calcium polysulfide, (5.020) thiram, (5.021) zineb, (5.022) ziram, (5.023) 6-ethyl-5,7-dioxo-6,7-dihydro-5H-pyrrolo[3',4':5,6][1,4]dithiino[2,3-c][1,2]thiazole-3-carbonitrile;

6) Compounds capable to induce a host defence, for example (6.001) acibenzolar-S-methyl, (6.002) isotianil, (6.003) probenazole, (6.004) tiadinil;

- 7) Inhibitors of the amino acid and/or protein biosynthesis, for example (7.001) cyprodinil, (7.002) kasugamycin, (7.003) kasugamycin hydrochloride hydrate, (7.004) oxytetracycline, (7.005) pyrimethanil, (7.006) 3-(5-fluoro-3,3,4,4-tetramethyl-3,4-dihydroisoquinolin-1-yl)quinolone;
- 8) Inhibitors of the ATP production, for example (8.001) silthiofam;
- 5 9) Inhibitors of the cell wall synthesis, for example (9.001) bentiavalicarb, (9.002) dimethomorph, (9.003) flumorph, (9.004) iprovalicarb, (9.005) mandipropamid, (9.006) pyrimorph, (9.007) valifenalate, (9.008) (2E)-3-(4-tert-butylphenyl)-3-(2-chloropyridin-4-yl)-1-(morpholin-4-yl)prop-2-en-1-one, (9.009) (2Z)-3-(4-tert-butylphenyl)-3-(2-chloropyridin-4-yl)-1-(morpholin-4-yl)prop-2-en-1-one;
- 10) Inhibitors of the lipid and membrane synthesis, for example (10.001) propamocarb, (10.002) propamocarb hydrochloride, (10.003) tolclofos-methyl;
- 11) Inhibitors of the melanin biosynthesis, for example (11.001) tricyclazole, (11.002) 2,2,2-trifluoroethyl {3-methyl-1-[(4-methylbenzoyl)amino]butan-2-yl}carbamate;
- 12) Inhibitors of the nucleic acid synthesis, for example (12.001) benalaxyl, (12.002) benalaxyl-M (kiralaxyl), (12.003) metalaxyl, (12.004) metalaxyl-M (mefenoxam);
- 15 13) Inhibitors of the signal transduction, for example (13.001) fludioxonil, (13.002) iprodione, (13.003) procymidone, (13.004) proquinazid, (13.005) quinoxifen, (13.006) vinclozolin;
- 14) Compounds capable to act as an uncoupler, for example (14.001) fluazinam, (14.002) meptyldinocap;
- 15) Further compounds, for example (15.001) Abscisic acid, (15.002) benthiazole, (15.003) bethoxazin, 20 (15.004) capsimycin, (15.005) carvone, (15.006) chinomethionat, (15.007) cufraneb, (15.008) cyflufenamid, (15.009) cymoxanil, (15.010) cyprosulfamide, (15.011) flutianil, (15.012) fosetyl-aluminium, (15.013) fosetyl-calcium, (15.014) fosetyl-sodium, (15.015) methyl isothiocyanate, (15.016) metrafenone, (15.017) mildiomyacin, (15.018) natamycin, (15.019) nickel dimethyldithiocarbamate, (15.020) nitrothal-isopropyl, (15.021) oxamocarb, (15.022) Oxathiapiprolin, (15.023) oxyfentiin, 25 (15.024) pentachlorophenol and salts, (15.025) phosphorous acid and its salts, (15.026) propamocarb-fosetilate, (15.027) pyriofenone (chlazafenone), (15.028) tebufloquin, (15.029) tecloftalam, (15.030) tolmanifanide, (15.031) 1-(4-{4-[(5R)-5-(2,6-difluorophenyl)-4,5-dihydro-1,2-oxazol-3-yl]-1,3-thiazol-2-yl}piperidin-1-yl)-2-[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]ethanone, (15.032) 1-(4-{4-[(5S)-5-(2,6-difluorophenyl)-4,5-dihydro-1,2-oxazol-3-yl]-1,3-thiazol-2-yl}piperidin-1-yl)-2-[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]ethanone, (15.033) 2-(6-benzylpyridin-2-yl)quinazoline, (15.034) 2,6-dimethyl-1H,5H-[1,4]dithiino[2,3-c:5,6-c']dipyrrole-1,3,5,7(2H,6H)-tetrone, (15.035) 2-[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]-1-[4-(4-{5-[2-(prop-2-yn-1-yloxy)phenyl]-4,5-dihydro-1,2-oxazol-3-yl]-1,3-thiazol-2-yl}piperidin-1-yl)ethanone, (15.036) 2-[3,5-bis(difluoromethyl)-1H-pyrazol-

1-yl]-1-[4-(4-{5-[2-chloro-6-(prop-2-yn-1-yloxy)phenyl]-4,5-dihydro-1,2-oxazol-3-yl]-1,3-thiazol-2-yl)piperidin-1-yl]ethanone, (15.037) 2-[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]-1-[4-(4-{5-[2-fluoro-6-(prop-2-yn-1-yloxy)phenyl]-4,5-dihydro-1,2-oxazol-3-yl]-1,3-thiazol-2-yl)piperidin-1-yl]ethanone, (15.038) 2-[6-(3-fluoro-4-methoxyphenyl)-5-methylpyridin-2-yl]quinazoline, (15.039) 2-((5R)-3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl]-3-chlorophenyl methanesulfonate, (15.040) 2-((5S)-3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl]-3-chlorophenyl methanesulfonate, (15.041) 2-{2-[(7,8-difluoro-2-methylquinolin-3-yl)oxy]-6-fluorophenyl}propan-2-ol, (15.042) 2-{2-fluoro-6-[(8-fluoro-2-methylquinolin-3-yl)oxy]phenyl}propan-2-ol, (15.043) 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl]-3-chlorophenyl methanesulfonate, (15.044) 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl]phenyl methanesulfonate, (15.045) 2-phenylphenol and salts, (15.046) 3-(4,4,5-trifluoro-3,3-dimethyl-3,4-dihydroisoquinolin-1-yl)quinoline, (15.047) 3-(4,4-difluoro-3,3-dimethyl-3,4-dihydroisoquinolin-1-yl)quinoline, (15.048) 4-amino-5-fluoropyrimidin-2-ol (tautomeric form: 4-amino-5-fluoropyrimidin-2(1H)-one), (15.049) 4-oxo-4-[(2-phenylethyl)amino]butanoic acid, (15.050) 5-amino-1,3,4-thiadiazole-2-thiol, (15.051) 5-chloro-N'-phenyl-N'-(prop-2-yn-1-yl)thiophene-2-sulfonohydrazide, (15.052) 5-fluoro-2-[(4-fluorobenzyl)oxy]pyrimidin-4-amine, (15.053) 5-fluoro-2-[(4-methylbenzyl)oxy]pyrimidin-4-amine, (15.054) 9-fluoro-2,2-dimethyl-5-(quinolin-3-yl)-2,3-dihydro-1,4-benzoxazepine, (15.055) but-3-yn-1-yl {6-[(Z)-(1-methyl-1H-tetrazol-5-yl)(phenyl)methylene]amino}oxy)methyl}pyridin-2-yl}carbamate, (15.056) ethyl (2Z)-3-amino-2-cyano-3-phenylacrylate, (15.057) phenazine-1-carboxylic acid, (15.058) propyl 3,4,5-trihydroxybenzoate, (15.059) quinolin-8-ol, (15.060) quinolin-8-ol sulfate (2:1), (15.061) tert-butyl {6-[(Z)-(1-methyl-1H-tetrazol-5-yl)(phenyl)methylene]amino}oxy)methyl}pyridin-2-yl}carbamate, (15.062) 5-fluoro-4-imino-3-methyl-1-[(4-methylphenyl)sulfonyl]-3,4-dihydropyrimidin-2(1H)-one.

The fungicides and the mixtures of fungicides mentioned herein may be used in a curative or preventive manner.

Fungicides used in the context of the present invention are selected from the group F-2 consisting of:

(3S,6S,7R,8R)-8-benzyl-3-[(3-[(isobutyryloxy)methoxy]-4-methoxypyridin-2-yl)carbonyl]amino]-6-methyl-4,9-dioxo-1,5-dioxonan-7-yl 2-methylpropanoate, 1-(4-{4-[(5R)-5-(2,6-difluorophenyl)-4,5-dihydro-1,2-oxazol-3-yl]-1,3-thiazol-2-yl}piperidin-1-yl)-2-[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]ethanone, 2,6-dimethyl-1H,5H-[1,4]dithiino[2,3-c:5,6-c']dipyrrole-1,3,5,7(2H,6H)-tetrone, 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl]-3-chlorophenyl methanesulfonate, 2-fluoro-6-(trifluoromethyl)-N-(1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl)benzamide, 3-(difluoromethyl)-1-methyl-N-(1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl)-1H-pyrazole-4-carboxamide, 3-(difluoromethyl)-N-(7-fluoro-1,1,3-trimethyl-2,3-dihydro-

1H-inden-4-yl)-1-methyl-1H-pyrazole-4-carboxamide, 3-chloro-4-(2,6-difluorophenyl)-6-methyl-5-phenylpyridazine, 4-(2-bromo-4-fluorophenyl)-N-(2-chloro-6-fluorophenyl)-1,3-dimethyl-1H-pyrazol-5-amine, 5-fluoro-2-[(4-methylbenzyl)oxy]pyrimidin-4-amine, Ametocradin, Amisulbrom, Azoxystrobin, Benthiavalicarb, Benzovindiflupyr, Bitertanol, Bixafen, Boscalid, Captan, Carbendazim, Carboxin, Carpropamid, Chloromequat-chloride, Chlorothalonil, Copper fungicides, in particular copper oxychloride, copper sulfate, Cyazofamid, Cyflufenamid, Cymoxanil, Cyproconazole, Cyprodinil, Diclocymet, Difenoconazole, Dimethomorph, Dimoxystrobin, Diniconazole, Dithianon, Edifenphos, Epoxiconazole, Famoxadone, Fenamidone, Fenhexamid, Fenpropidin, Fenpropimorph, Fenpyrazamine, Ferimzone, Fluazinam, Fludioxonil, Fluindapyr, Fluopicolide, Fluopyram, Fluoxastrobin, Fluquinconazole, Flusilazole, Fluthianil, Flutriafol, Fluxapyroxad, Folpet, Fosetyl or salts and esters thereof, in particular Fosetyl-aluminium, Furametpyr, Gentamycin, Hexaconazole, Hymexazol, Imazalil, Iaconazole, Ipfentrifluconazole, Iprobenfos (IBP), Iprodione, Iprovalicarb, Isofetamid, Isoprothiolane, Isopyrazam, Isotianil, Kasugamycin, Kresoxim-methyl, Lyserphenvalpyr, Mancozeb, Mandestrobin, Mandipropamid, Maneb, Mefenoxam, Mefentrifluconazole, Mepiquat-chloride, Meptyldinocap, Metalaxyl, Mefenoxam, Metconazole, Metiram, Metominostrobin, Metrafenone, Myclobutanil, N-(5-chloro-2-isopropylbenzyl)-N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide, Orysastrobin, Oxathiapiprolin, Paclobutrazole, Penconazole, Pencycuron, Penflufen, Penthiopyrad, Phosphonic acid (H₃PO₃), Phthalide, Picarbutrazox, Picoxystrobin, Probenazole, Prochloraz, Propamocarb, Propiconazole, Propineb, Proquinazid, Prothioconazole, Pydiflumetofen, Pyraclostrobin, Pyraziflumid, Pyrimethanil, Pyriofenone, Pyroquilon, Quinofumelin, Quinoxifen, Quinoxifen, Quintozene, Saponin, Sedaxane, Silthiofam, Simconazole, Spiroxamine, Streptomycin, Sulphur, Tebuconazole, Tebufloquin, Tetraconazole, Thiabendazole, Thifluzamide, Thiophanate-methyl, Thiram, Tiadinil, Tolprocarb, Triadimefon, Triadimenol, Tricyclazole, Trifloxystrobin, Triticonazole, Validamycin, Ziram and the esters and/or agronomically acceptable salts of these fungicides.

Preferred Fungicides are selected from the group F-3 consisting of:

(3S,6S,7R,8R)-8-benzyl-3-[(3-[(isobutyryloxy)methoxy]-4-methoxypyridin-2-yl)carbonylamino]-6-methyl-4,9-dioxo-1,5-dioxonan-7-yl 2-methylpropanoate, 1-(4-{4-[(5R)-5-(2,6-difluorophenyl)-4,5-dihydro-1,2-oxazol-3-yl]-1,3-thiazol-2-yl}piperidin-1-yl)-2-[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]ethanone, 2,6-dimethyl-1H,5H-[1,4]dithiino[2,3-c:5,6-c']dipyrrole-1,3,5,7(2H,6H)-tetrone, 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-3-chlorophenyl methanesulfonate, 2-fluoro-6-(trifluoromethyl)-N-(1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl)benzamide, 3-(difluoromethyl)-1-methyl-N-(1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl)-1H-pyrazole-4-carboxamide, 3-(difluoromethyl)-N-(7-fluoro-1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl)-1-methyl-1H-pyrazole-4-carboxamide, 3-chloro-4-(2,6-difluorophenyl)-6-methyl-5-phenylpyridazine, 4-(2-bromo-4-fluorophenyl)-N-(2-chloro-6-fluorophenyl)-1,3-dimethyl-1H-pyrazol-5-amine, 5-fluoro-2-[(4-methylbenzyl)oxy]pyrimidin-4-amine, Ametocradin, Amisulbrom,

Azoxystrobin, Benzovindiflupyr, Bitertanol, Bixafen, Boscalid, Carpropamid, Chlorothalonil, Cyproconazole, Difenoconazole, Epoxiconazole, Famoxadone, Fenamidone, Fenhexamid, Fludioxonil, Fluindapyr, Fluopicolide, Fluopyram, Fluoxastrobin, Fluquinconazole, Fluxapyroxad, Fosetyl or salts and esters thereof, in particular Fosetyl-aluminium, Ipconazole, Ipfentrifluconazole, Iprodione, 5 Iprovalicarb, Isofetamid, Isoprothiolane, Isopyrazam, Isotianil, Kresoxim-methyl, Lyserphenvalpyr, Mandestrobin, Mandipropamid, Mefenoxam, Mefentrifluconazole, Metalaxyl, Mefenoxam, Metconazole, Metominostrobin, N-(5-chloro-2-isopropylbenzyl)-N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide, Oxathiapiprolin, Pencycuron, Picarbutrazox, Picoxystrobin, Propamocarb, Propiconazole, Propineb, Prothioconazole, Pydiflumetofen, 10 Pyraclostrobin, Pyraziflumid, Pyrimethanil, Pyriofenone, Quinofumelin, Quinoxifen, Sedaxane, Silthiofam, Spiroxamine, Tebuconazole, Tetraconazole, Tolprocarb, Triadimefon, Triadimenol, Trifloxystrobin, and the esters and/or agronomically acceptable salts of these fungicides.

Preferred mixtures of fungicides used in the context of the present invention are selected from the group F4 consisting of :

15 mixtures of (3S,6S,7R,8R)-8-benzyl-3-[(3-[(isobutyryloxy)methoxy]-4-methoxypyridin-2-yl)carbonyl]amino]-6-methyl-4,9-dioxo-1,5-dioxonan-7-yl 2-methylpropanoate and other fungicides mentioned herein, mixtures of 2,6-dimethyl-1H,5H-[1,4]dithiino[2,3-c:5,6-c']dipyrrole-1,3,5,7(2H,6H)-tetrone, esters and/or salts thereof and other fungicides mentioned herein, mixtures of 2-[3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl]-3-chlorophenyl methanesulfonate and other fungicides mentioned herein, mixtures of 1-(4-{4-[(5R)-5-(2,6-difluorophenyl)-4,5-dihydro-1,2-oxazol-3-yl]-1,3-thiazol-2-yl}piperidin-1-yl)-2-[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]ethanone and other fungicides mentioned herein, mixtures of Ametoctradin and other fungicides mentioned herein, mixtures of Bitertanol and other fungicides mentioned herein, mixtures of Bixafen and/or salts thereof and other fungicides mentioned herein, 25 mixtures of Carpropamid and other fungicides mentioned herein, mixtures of Cyproconazole and other fungicides mentioned herein, mixtures of Epoxiconazole and other fungicides mentioned herein, mixtures of Fenhexamid and other fungicides mentioned herein, mixtures of Fenamidone and other fungicides mentioned herein, mixtures of Fluopyram and other fungicides mentioned herein, mixtures of Fluopicolide and other fungicides mentioned herein, mixtures of Fluoxastrobin and other fungicides mentioned herein, mixtures of Fluxapyroxad and other fungicides mentioned herein, mixtures of Fluquinconazole and other fungicides mentioned herein, mixtures of Fosetyl or salts and esters thereof, in particular Fosetyl-aluminium and other fungicides mentioned herein, mixtures of Ipfentrifluconazole and other fungicides mentioned herein, mixtures of Iprovalicarb, and other fungicides mentioned herein, mixtures of Isotianil and other fungicides mentioned herein, mixtures of Lyserphenvalpyr and other 30 fungicides mentioned herein, mixtures of Mefentrifluconazole and other fungicides mentioned herein, mixtures of Metalaxyl and other fungicides mentioned herein, mixtures of Mefenoxam and other fungicides mentioned herein, mixtures of Metominostrobin and other fungicides mentioned herein,

mixtures of N-(5-chloro-2-isopropylbenzyl)-N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide and other fungicides mentioned herein, mixtures of Pencycuron and other fungicides mentioned herein, mixtures of Penflufen and other fungicides mentioned herein, mixtures of Propamocarb and other fungicides mentioned herein, mixtures of Propineb and other fungicides mentioned herein, mixtures of Pyrimethanil and other fungicides mentioned herein, mixtures of Pyraclostrobin and other fungicides mentioned herein, mixtures of Prothioconazole and other fungicides mentioned herein, mixtures of Spiroxamine and other fungicides mentioned herein, mixtures of Tebuconazole and other fungicides mentioned herein, mixtures of Triadimenol and other fungicides mentioned herein, mixtures of Triadimefon and other fungicides mentioned herein, mixtures of Trifloxystrobin and other fungicides mentioned herein.

Particularly preferred mixtures of fungicides used in the context of the present invention are selected from the group F-5 consisting of:

mixtures of Prothioconazole and Fluopyram, mixtures of Prothioconazole and Tebuconazole, mixtures of Prothioconazole, Bixafen, and Tebuconazole,

mixtures of Prothioconazole, Bixafen, and Trifloxystrobin, mixtures of Prothioconazole, Bixafen, and Spiroxamine, mixtures of Prothioconazole, Bixafen, and Chlorothalonil, mixtures of Prothioconazole and Tebuconazole and Metalaxyl, mixtures of Prothioconazole, Bixafen, and Fluopyram, mixtures of Prothioconazole and Fluoxastrobin, mixtures of Prothioconazole and Trifloxystrobin, mixtures of Prothioconazole and Pencycuron, mixtures of Prothioconazole and Spiroxamine, mixtures of Tebuconazole and Bixafen, mixtures of Tebuconazole and Metalaxyl, mixtures of Tebuconazole and Fluoxastrobin, mixtures of Tebuconazole and Trifloxystrobin, mixtures of Tebuconazole and Fenhexamid, mixtures of Tebuconazole and Fluopicolide, mixtures of Tebuconazole and Spiroxamine, mixtures of Tebuconazole and Pencycuron, mixtures of Fluopyram and Tebuconazole, mixtures of Fluopyram and Spiroxamine, mixtures of Fluopyram and Pyrimethanil, mixtures of Fluopyram and Trifloxystrobin, mixtures of Fluopyram and Triadimenol, mixtures of Fluopyram and Fosetyl or salts and esters thereof, in particular Fosetyl-aluminium, mixtures of Fluopyram and Imidacloprid, mixtures of Fluopyram and Spiroxamine, mixtures of Fluopyram and Fluoxastrobin, mixtures of Bixafen and Spiroxamine, mixtures of Bixafen and Trifloxystrobin, mixtures of Bixafen and Chlorothalonil, mixtures of Bixafen and Fluoxastrobin, mixtures of Trifloxystrobin and Isotianil, mixtures of Trifloxystrobin and Metalaxyl,

mixtures of Trifloxystrobin and Propineb, mixtures of Trifloxystrobin and Pyrimethanil, mixtures of Trifloxystrobin and Fosetyl or salts and esters thereof, in particular Fosetyl-aluminium, mixtures of Trifloxystrobin and Fluopicolide, mixtures of Propineb and Fluopicolide, mixtures of Fosetyl or salts and esters thereof, in particular Fosetyl-aluminium,

and Fluopicolide, mixtures of Bixafen and Ipflentrifluconazole, mixtures of Bixafen and Mefentrifluconazole,

mixtures of Metalaxyl and Ipfentrifluconazole, mixtures of Metalaxyl and Mefentrifluconazole, mixtures
 of Prothioconazole and Ipfentrifluconazole, mixtures of Prothioconazole and Mefentrifluconazole,
 mixtures of Tebuconazole and Ipfentrifluconazole, mixtures of Tebuconazole and Mefentrifluconazole,
 mixtures of Trifloxystrobin and Ipfentrifluconazole, mixtures of Trifloxystrobin and
 5 Mefentrifluconazole, mixtures of Fluoxastrobin and Ipfentrifluconazole, mixtures of Fluoxastrobin and
 Mefentrifluconazole, mixtures of Fluopyram and Ipfentrifluconazole, mixtures of Fluopyram and
 Mefentrifluconazole, mixtures of Spiroxamine and Ipfentrifluconazole, mixtures of Spiroxamine and
 Mefentrifluconazole, mixtures of Pydiflumetofen and Ipfentrifluconazole, mixtures of Pydiflumetofen
 and Mefentrifluconazole, mixtures of Pyraclostrobin and Ipfentrifluconazole, mixtures of Pyraclostrobin
 10 and Mefentrifluconazole, mixtures of Fluxapyroxad and Ipfentrifluconazole, mixtures of Fluxapyroxad
 and Mefentrifluconazole.

SeedGrowth

15 Seeds of plants comprising the events and alleles as disclosed below in table A, B and C can also be
 treated with seed treatment.

Seeds can be coated with a coating comprising a compound selected from the following fungicides,
 insecticides, safeners and/or biologics.

Preferred fungicides for seed treatment of seeds of plants comprising the events and alleles as disclosed
 below in table A, B and C are selected from the group consisting of fungicides, insecticides, biologics as
 20 disclosed below.

Preferred fungicides for seed treatment of seeds are selected from the group SG-1 consisting of

Benzovindiflupyr, Carbendazim, Carboxin, Difenoconazole, Ethaboxam, Fludioxonil, Fluquinconazole,
 Fluxapyroxad, Iaconazole, Ipfentrifluconazole, Isotianil, Mefenoxam, Mefentrifluconazole, Metalaxyl,
 Pencycuron, Penflufen, Penthiopyrad, Prothioconazole, Prochloraz, Pyraclostrobin, Sedaxane,
 25 Silthiofam, Tebuconazole, Trifloxystrobin, Triticonazole, Ethaboxam (SCC), Penthiopyrad (DPX
 pipeline), Benzovindiflupyr (SYN pipeline), Bixafen, (see biologicals), Dimethomorph, Fenamidone,
 Fluopicolide, Fluoxastrobin, Flutolanil, Tolclophos-methyl, Azoxystrobin, Chlorothalonil,
 Cyproconazole, Cyprodinil, Diniconazole, Fluopyram, Flutriafol, Fluxapyroxad, Imazalil, Isopyrazam,
 Isotianil, Iprodione, Metconazole, Myclobutanil, Picoxystrobin, Pyrimethanil, Tetraconazole,
 30 Thiabendazole, Thiophanate-methyl, Triadimenol, Thiram, Triflumizole, Ziram, 2,6-dimethyl-1H,5H-
 [1,4]dithiino[2,3-c:5,6-c']dipyrrole-1,3,5,7(2H,6H)-tetrone, 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-
 pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-3-chlorophenyl
 methanesulfonate (Thiazolylpiperidin, N-(5-chloro-2-isopropylbenzyl)-N-cyclopropyl-3-

(difluoromethyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide Cypropanide, Picarbutrazox and Oxathiapiprolin.

Preferred insecticides for seed treatment of seeds are selected from the group SG-2 consisting of

Abamectin, Afidopyropen, Bifenthrin, Carbofuran, Carbendazim, Carboxin, Clothianidin, Cyazapyr,
 5 Cypermethrin, Deltamethrin, Difenconazole, Ethoxysulfuron, Fenamidon, Fenoxaprop-P-Ethyl,
 Ethiprole, Fipronil, Fluazaindolizine, Flupyradifurone, Flubendiamide, Flupicolide, Fluopyram,
 Fluoxastrobin, Fluquinconazole, Fosetyl-Al, Imidacloprid, Prochloraz, Propineb, Prothioconazole,
 Lambda-Cyhalothrin, Methiocarb, Rynaxypyr, Spirotetramat, Spinosad, Tebuconazole, Tembotrione,
 Thiacloprid, Thiram, Thiametoxam, Thiodicarb, Thioxazafen, (MON pipeline Nematicide),
 10 Fluazaindolizine (DPX pipeline Nematicide), Lambda-Cyhalothrin (SYN), Bifenthrin (FMC),
 Cypermethrin (few companies), Acetamiprid, β -Cyfluthrin, Flubendiamide, Thiacloprid,
 Chlorpyrifos, Metamidophos, Phorate, Sulfoxaflor, Tefluthrin, Triflumuron, Tetraniliprole,
 Triadimenol, Trifloxystrobin, Triflumuron, Broflanilide (MCI 8007 – Mitsui/BASF), Cycloxaprid and
 Spinetoram.

15

Preferred safeners for seed treatment of seeds are selected from the group SG-3 consisting of

Cyprosulfamide, Isoxadifen, Isoxadifen-ethyl, Lipo-Chito-Oligosaccharide (LCO), Mefenpyr and Mefenpyr-diethyl.

Preferred Biologicals and biological active ingredients for seed treatment of seeds are selected from the
 20 group SG-4 consisting of

Bacillus firmus, especially *Bacillus firmus* GB126, *Bacillus subtilis*, *Bacillus pumilus* especially *Bacillus pumilus* QST 2808, *Bacillus pumilus* GB34, *Pasteuria usage* strains (e.g. as used in ECONEM),
Rhizobium spp, especially *Rhizobium tropici* SP25, *Pseudomonas fluorescens*, *Pseudomonas chloropsis*,
Penicillium bilaii, *Rhizobium japonicum*, *Purnate Varroacide*, *Chenopodium quinoa saponins*, a
 25 recombinant exosporium-producing *Bacillus thuringiensis* cell (especially *B. thuringiensis* BT013A)
 that expresses a fusion protein comprising:

- 1) an endoglucanase, (especially from *Bacillus subtilis*) having the amino acid sequence of Sequence 2 as herein defined and
 - 2) a targeting sequence having at least about 43% identity with amino acids 20–35 of Sequence 1 as herein defined, wherein the identity with amino acids 25–35 is at least about 54%,
- 30

and a recombinant exosporium-producing *Bacillus thuringiensis* cell that expresses a fusion protein comprising:

- 1) a phospholipase having the amino acid sequence of Sequence 3 as herein defined and
- 2) a targeting sequence having at least about 43% identity with amino acids 20–35 of Sequence 1 as herein defined, wherein the identity with amino acids 25–35 is at least about 54%.

5 More preferred Biologicals and biological active ingredients for seed treatment of seeds are selected from the group SG-5 consisting of:

Bacillus firmus, especially *Bacillus firmus* GB126, *Bacillus subtilis*, *Bacillus pumilus* especially *Bacillus pumilus* QST 2808, *Bacillus pumilus* GB34, a recombinant exosporium-producing *Bacillus thuringiensis* cell (especially *B. thuringiensis* BT013A) that expresses a fusion protein comprising:

- 10 1) an endoglucanase, (especially from *Bacillus subtilis*) having the amino acid sequence of Sequence 2 as herein defined and
- 2) a targeting sequence having at least about 43% identity with amino acids 20–35 of Sequence 1 as herein defined, wherein the identity with amino acids 25–35 is at least about 54%,

and a recombinant exosporium-producing *Bacillus thuringiensis* cell that expresses a fusion protein
15 comprising:

- 1) a phospholipase having the amino acid sequence of Sequence 3 as herein defined and
- 2) a targeting sequence having at least about 43% identity with amino acids 20–35 of Sequence 1 as herein defined, wherein the identity with amino acids 25–35 is at least about 54%.

20 In another more preferred embodiment, preferred active ingredients for seed treatment of seeds are selected from the group SG-6 consisting of:

β -Cyfluthrin, *Bradyrhizobium japonicum*, Carbendazim, Carboxin Clothianidin, Cypermethrin, Deltamethrin, Difenoconazole, Ethoxysulfuron, Fenamidon, Fenoxaprop-P-Ethyl, Flubendiamide, Fluopicolide, Fluopyram, Fluoxastrobin, Fluquinconazole, Fosetyl-Al, Fipronil, Prochloraz, Propineb,
25 Prothioconazole, *Pseudomonas Fluorescens*, Spirotetramat, Tebuconazole, Tembotrione, Thiacloprid, Thiodicarb, Thiram, Triadimenol, Trifloxystrobin and Triflumuron.

Particularly preferred combinations of active ingredients for seed treatment used in the context of the present invention are selected from the group consisting of SG-7:

- combination of Clothianidin and *B. firmus* GB126,
- 30 combination of Imidacloprid and Thiodicarb,
- combination of Imidacloprid and Prothioconazole,
- combination of Clothianidin and Carboxin and Metalaxyl and Trifloxystrobin,
- combination of Metalaxyl and Prothioconazole and Tebuconazole,
- combination of Clothianidin and beta-Cyfluthrin,
- 35 combination of Prothioconazole and Tebuconazole,

- combination of Clothianidin and Imidacloprid and Prothioconazole and Tebuconazole,
combination of Carbendazim and Thiram,
combination of Imidacloprid and Methiocarb,
combination of Metalaxyl and Penflufen and Prothioconazole,
- 5 combination of Imidacloprid Prothioconazole,
combination of Imidacloprid Tefluthrin,
combination of Imidacloprid Pencycuron,
combination of Imidacloprid Penflufen,
combination of Fluoxastrobin and Prothioconazole and Tebuconazole,
- 10 combination of Metalaxyl and Trifloxystrobin ,
combination of Penflufen and Trifloxystrobin,
combination of Prothioconazole and Tebuconazole,
combination of Fluoxastrobin and Prothioconazole and Tebuconazole and Triazoxide,
combination of Imidacloprid and Methiocarb and Thiram,
- 15 combination of Clothianidin and beta-Cyfluthrin,
combination of Clothianidin and Fluoxastrobin and Prothioconazole and Tebuconazole,
combination of Fluopyram and Fluoxastrobin and Triadimenol,
combination of Metalaxyl and Trifloxystrobin,
combination of Imidacloprid and Ipconacole,
- 20 combination of Difenconazol and Fludioxonil and Tebuconazole,
combination of Imidacloprid + Tebuconazole,
combination of Imidacloprid and Prothioconazole and Tebuconazole and
combination of Metalaxyl and Prothioconazole and Tebuconazole.
- 25 In one preferred embodiment, a combination of active ingredients for seed treatment is selected from the
group SG-8 consisting of:
- combination of Clothianidin and *B. firmus* GB126,
combination of Imidacloprid and Thiodicarb,
- 30 combination of Imidacloprid and Prothioconazole,
combination of Clothianidin Carboxin, Metalaxyl, Trifloxystrobin,
combination of Metalaxyl, Prothioconazole Tebuconazole and
combination of Clothianidin and beta-Cyfluthrin and
combination of Prothioconazole and Tebuconazole.

35

In the context of the present invention, "increasing yield" means a significant increase in yield by compared with the untreated plant, preferably a significant increase by at least 1% such as by 1% to 3% , compared with the untreated plant (100% yield), i.e. the yield of the treated plants is at least 101%

compared to the yield (100%) of the untreated plant; more preferably, the yield is even more increased
ba at least 2%, more preferably at least 5% such as by 2% to 5% (yield from 102% to 105%), by 2 to 10
% (yield from 102% to 110%) or by 5% to 20% (yield of from 105% to 120 %). The yield increase may
be achieved by curative treatment, i.e. for treatment of already infected plants, or by protective
5 treatment, for protection of plants which have not yet been infected.

In the context of the present invention, "to treat with" means to contact a plant or part of a plant with an
effective amount of an active ingredient (AI) or a combination thereof or to coat a seed with an active AI
or a combination thereof.

"Active ingredient" shall refer to compounds used in agriculture. Agriculture encompasses the
10 production of food and feed crops, forestry, the protection of stored products including food, feed but
also other materials of plant origin. Preferably agriculture shall encompass the production of food and
feed crops, forestry, the protection of stored products being food, feed, and materials of plant origin.
Active ingredients according to the invention are not applied to humans or animals as a medical or
therapeutic treatment.

15 "Insecticides" shall refer to active ingredients which are capable of controlling phytopathogenic pests
including but not limited to insects, mites, nematodes.

In the context of the present invention the active ingredients according to the invention or to be used
according to the invention may be a composition (i. e. a physical mixture) comprising at least one active
ingredient. It may also be a combination of active ingredients composed from separate formulations of a
20 single active ingredient component being active ingredient (tank-mix). Another example of a
combination of active ingredients according to the invention is that the active ingredients are not
present together in the same formulation, but packaged separately (combipack), i.e., not jointly
preformulated. As such, combipacks include one or more separate containers such as vials, cans, bottles,
pouches, bags or canisters, each container containing a separate component for an agrochemical
25 composition, here at least one active ingredient. One example is a two-component combipack.
Accordingly the present invention also relates to a two-component combipack, comprising a first
component which in turn comprises an active ingredient, a liquid or solid carrier and, if appropriate, at
least one surfactant and/or at least one customary auxiliary, and a second component which in turn
comprises another active ingredient, a liquid or solid carrier and, if appropriate, at least one surfactant
30 and/or at least one customary auxiliary. More details, e.g. as to suitable liquid and solid carriers,
surfactants and customary auxiliaries are described below. A mixture or combination according to the
invention shall mean/encompass a tank mix or a combipack.

In one embodiment an active ingredient or a combination of active ingredients as described above is
used for increasing yield in Brassicaceae.

35 Thus, one embodiment also refers to a method to increase yield of a plant as described herein
comprising the step of treating a plant or part of a plant as described herein with an effective amount of

an active ingredient or a combination thereof.

Especially, the following *Brassicaceae* comprising an event or allele as disclosed in tables A, B, and C below can be treated with an active ingredient or a combination of active ingredients as described in groups H-1, H-2, H-3, H-4, H-5, P-1, SAF-1, SAF-2, SAF-3, SAF-4, B-1, B-2, B-3, B-4, B-5, I-1, I-2, I-3, I-4, I-5, I-6, F-1, F-2, F-3, F-4, F-5, F-6, SG-1, SG-2, SG-3, SG-4, SG-5, SG-6, SG-7, SG-8.

In one embodiment, *Brassicaceae* plants suitable for treatments according to the invention are plants comprising the events or alleles listed in Table A. The events are characterized by reference to patent applications (column PA – incorporated by reference) or regulatory files (column RF –incorporated by reference) wherein they are described. Transgenic plants comprising the events can also be described by reference to the SEQ ID entries of the Sequence listings of the referenced patent applications (column CS- herein incorporated by reference). These event characterizing sequences correspond to 5' flanking regions and junction sequences, 3' flanking regions and junction sequences, inserted transgene sequences, and/or complete sequences of the transgenic insert in the event. Transgenic plants suitable for treatments according to the invention comprise at least one, preferably all of the reference sequences.

Thus, in one embodiment, *Brassicaceae* plants suitable for treatments according to the invention are transgenic plants comprising at least one or more events characterized by the following OECD numbers: CGN-89111-8 or MON-88302-9 or MON-89249-2 or MON-00073-7 or ACS-BN003-6 or *Brassicaceae* plants comprising at least one or more the following alleles: POSH122, OGURA, R113, R40, R2000.

In another embodiment, *Brassicaceae* plants suitable for treatments according to the invention are transgenic *Brassicaceae* plants having at least one event or allele as present in reference seeds being deposited as: ATCC PTA-10955 or ATCC PTA-730 or ATCC PTA-730 or NCIMB41574 or ATCC PTA-850 or PTA-2485.

Table A

TYPE	NAME	ALTER NATIVE NAME S (List)	UNIQUE IDENTIFIER NUMBER	GENES	TYPE TRAIT	DEPOSIT NUMBER	PA	RF	CS
EVENT1	23-18-17	EVENT 18, pCGN3 828-212 / 86-18	CGN-89111-8	te, nptIII	Modified acid oil/fatty			US94-090-01p	
EVENT2	MON8 8302	PV- BNHT2 672	MON-88302-9	cp4 epsps	Glyphosate tolerance	ATCC PTA-10955	WO2011/153186	US11-188-01p	SEQ ID Nos. 3, 4, 5 and particularly SEQ ID No 6
EVENT3	RT200	GT200	MON-89249-2	cp4 epsps, goxv247	Glyphosate tolerance			US01-324-01p	
EVENT4	RT73	GT73, PV- BNGT0 4	MON-00073-7	cp4 epsps, goxv247	Glyphosate tolerance		WO2002/036831	US98-216-01p	SEQ ID Nos 7 and 8
EVENT5	MS8		ACS-BN005-8	barnase, bar	Male Sterility, Glufosinate Tolerance	ATCC PTA-730	WO2001/041558	US98-278-01p	SEQ ID Nos 13 and 18

TYPE	ALTER NATIVE NAME S (List)	UNIQUE IDENTIFIER NUMBER	GENES	TYPE TRAIT	DEPOSIT NUMBER	PA	RF	CS
EVENT6	RF3	ACS-BN003-6	barstar, bar	Fertility Restoration, Glufosinate Tolerance	ATCC PTA-730	WO2001/041558	US98-278-01p	SEQ ID Nos 24 and 30
ALLELE 1	POSHI 22	not applicable	ind	Seed Resistance	NCIMB41 574	WO2010/006732		SEQ ID No 27
EVENT7	MS11		bar, barnase, barstar	Male Glufosinate Tolerance	ATCC PTA-850 or PTA- 2485			SEQ ID Nos 8 and 10
ALLELE 2	Ogura	not applicable	cms	Male Sterility				
ALLELE 3	R113	not applicable	Restorer	Fertility Restoration				
ALLELE 4	R40	not applicable	Restorer	Fertility Restoration				
ALLELE 5	R2000	not applicable	Restorer	Fertility Restoration				

In another embodiment, *Brassicaceae* plants suitable for treatments according to the invention are commercialized plants comprising the events or alleles listed in Table B. The events are characterized by reference to patent applications (column PA – incorporated by reference) or regulatory files (column RF –incorporated by reference) wherein they are described. Transgenic plants comprising the events can also be described by reference to the SEQ ID entries of the Sequence listings of the referenced patent applications (column CS- herein incorporated by reference). These event characterizing sequences correspond to 5'flanking regions and junction sequences, 3' flanking regions and junction sequences, inserted transgene sequences, and/or complete sequences of the transgenic insert in the event. Transgenic plants suitable for treatments according to the invention comprise at least one, preferably all of the reference sequences.

Thus, in another embodiment, *Brassicaceae* plants suitable for treatments according to the invention are transgenic plants comprising at least one or more events characterized by the following OECD numbers: MON-88302-9 or MON-89249-2 or MON-00073-7 or ACS-BN003-6 or *Brassicaceae* plants comprising at least one or more the following alleles: POSH122, OGURA, R113, R40, R2000.

In another embodiment, *Brassicaceae* plants suitable for treatments according to the invention are transgenic *Brassicaceae* plants having at least one event or allele as present in reference seeds being deposited as: ATCC PTA-10955 or ATCC PTA-730 or ATCC PTA-730 or NCIMB41574 or ATCC PTA-850 or PTA-2485.

Table B (commercialized plants)

ALTER NATIVE NAME	UNIQUE IDENTIFIER NUMBER	GENES	TYPE TRAIT	DEPOSIT NUMBER	PA	RF	CS
MON88302	PV-BNHIT2672 MON-88302-9	cp4 epsps	Glyphosate tolerance	ATCC PTA-10955	WO2011/153186	US11-188-01p	SEQ ID Nos. 3, 4, 5 and particularly SEQ ID No 6
RT200	MON-89249-2 GT200	cp4 epsps, goxv247	Glyphosate tolerance			US01-324-01p	
RT73	GT73, PV-BNGT04 MON-00073-7	cp4 epsps, goxv247	Glyphosate tolerance		WO2002/036831	US98-216-01p	SEQ ID Nos 7 and 8
MS8	ACS-BN005-8	barnase, bar	Male Sterility, Glufosinate Tolerance	ATCC PTA-730	WO2001/041558	US98-278-01p	SEQ ID Nos 13 and 18
RF3	ACS-BN003-6	barstar, bar	Fertility Restoration, Glufosinate Tolerance	ATCC PTA-730	WO2001/041558	US98-278-01p	SEQ ID Nos 24 and 30

POSH122	not applicable	not applicable	not applicable	Seed Shatter Resistance	NCIMB41574	WO2010/006732	SEQ ID No 27	
MS11	bar, barnase, barstar	Male Sterility, Glufosinate Tolerance	ATCC PTA-850 or PTA-2485	WO2001/031042	SEQ ID Nos 8 and 10			
Ogura	not applicable	Male Sterility						
R113	not applicable	Fertility Restoration						
R40	not applicable	Fertility Restoration						
R2000	not applicable	Fertility Restoration						

In yet another embodiment, *Brassicaceae* plants suitable for treatments according to the invention are plants comprising the recently developed events or alleles listed in Table C. The events are characterized by reference to patent applications (column PA – incorporated by reference) or regulatory files (column RF –incorporated by reference) wherein they are described. Transgenic plants comprising
5 the events can also be described by reference to the SEQ ID entries of the Sequence listings of the referenced patent applications (column CS- herein incorporated by reference). These event characterizing sequences correspond to 5'flanking regions and junction sequences, 3' flanking regions and junction sequences, inserted transgene sequences, and/or complete sequences of the transgenic insert in the event. Transgenic plants suitable for treatments according to the invention comprise at least one, preferably all
10 of the reference sequences.

Thus, in another embodiment, *Brassicaceae* plants suitable for treatments according to the invention are transgenic plants comprising at least the event characterized by the following OECD numbers: MON-88302-9 or *Brassicaceae* plants comprising at least the following allele: POSH122.

In another embodiment, *Brassicaceae* plants suitable for treatments according to the invention are
15 transgenic *Brassicaceae* plants having at least one event or allele as present in reference seeds being deposited as: ATCC PTA-10955 or NCIMB41574.

Table C (Recently developed plants)

TYPE	NAME	ALTERNATIVE NAMES (List)	UNIQUE IDENTIFIER NUMBER	GENES	TYPE TRAIT	DEPOSIT NUMBER	PA	RF	CS
EVENT	MON 88302	PV- BNHT2672	MON- 88302-9	cp4 epsps	Glyphosate tolerance	ATCC PTA- 10955	WO2011/ 153186	US11-188-01p	SEQ ID Nos. 3, 4, 5 and particularly SEQ ID No 6
ALLELE	POSH 122		not applicable	not applicable	Seed Shatter Resistance	NCIMB41574	WO2010/ 006732		SEQ ID No 27

- In one embodiment at least one active ingredient selected from group H-1 is used on a plant comprising EVENT1 or plant parts thereof, preferably for increasing yield.
- In one embodiment at least one active ingredient selected from group H-2 is used on a plant comprising
5 EVENT1 or plant parts thereof, preferably for increasing yield.
- In one embodiment at least one active ingredient selected from group H-3 is used on a plant comprising EVENT1 or plant parts thereof, preferably for increasing yield.
- In one embodiment at least one active ingredient selected from group H-4 is used on a plant comprising EVENT1 or plant parts thereof, preferably for increasing yield.
- 10 In one embodiment at least one active ingredient selected from group H-5 is used on a plant comprising EVENT1 or plant parts thereof, preferably for increasing yield.
- In one embodiment at least one active ingredient selected from group P-1 is used on a plant comprising EVENT1 or plant parts thereof, preferably for increasing yield.
- In one embodiment at least one active ingredient selected from group SAF-1 is used on a plant
15 comprising EVENT1 or plant parts thereof, preferably for increasing yield.
- In one embodiment at least one active ingredient selected from group SAF-2 is used on a plant comprising EVENT1 or plant parts thereof, preferably for increasing yield.
- In one embodiment at least one active ingredient selected from group SAF-3 is used on a plant comprising EVENT1 or plant parts thereof, preferably for increasing yield.
- 20 In one embodiment at least one active ingredient selected from group SAF-4 is used on a plant comprising EVENT1 or plant parts thereof, preferably for increasing yield.
- In one embodiment at least one active ingredient selected from group B-1 is used on a plant comprising EVENT1 or plant parts thereof, preferably for increasing yield.
- In one embodiment at least one active ingredient selected from group B-2 is used on a plant comprising
25 EVENT1 or plant parts thereof, preferably for increasing yield.
- In one embodiment at least one active ingredient selected from group B-3 is used on a plant comprising EVENT1 or plant parts thereof, preferably for increasing yield.
- In one embodiment at least one active ingredient selected from group B-4 is used on a plant comprising EVENT1 or plant parts thereof, preferably for increasing yield.
- 30 In one embodiment at least one active ingredient selected from group B-5 is used on a plant comprising EVENT1 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group I-1 is used on a plant comprising EVENT1 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group I-2 is used on a plant comprising EVENT1 or plant parts thereof, preferably for increasing yield.

5 In one embodiment at least one active ingredient selected from group I-3 is used on a plant comprising EVENT1 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group I-4 is used on a plant comprising EVENT1 or plant parts thereof, preferably for increasing yield.

10 In one embodiment at least one active ingredient selected from group I-5 is used on a plant comprising EVENT1 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group I-6 is used on a plant comprising EVENT1 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group F-1 is used on a plant comprising EVENT1 or plant parts thereof, preferably for increasing yield.

15 In one embodiment at least one active ingredient selected from group F-2 is used on a plant comprising EVENT1 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group F-3 is used on a plant comprising EVENT1 or plant parts thereof, preferably for increasing yield.

20 In one embodiment at least one active ingredient selected from group F-4 is used on a plant comprising EVENT1 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group F-5 is used on a plant comprising EVENT1 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group F-6 is used on a plant comprising EVENT1 or plant parts thereof, preferably for increasing yield.

25 In one embodiment at least one active ingredient selected from group SG-1 is used on a plant comprising EVENT1 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SG-2 is used on a plant comprising EVENT1 or plant parts thereof, preferably for increasing yield.

30 In one embodiment at least one active ingredient selected from group SG-3 is used on a plant comprising EVENT1 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SG-4 is used on a plant comprising EVENT1 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SG-5 is used on a plant comprising EVENT1 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SG-6 is used on a plant comprising EVENT1 or plant parts thereof, preferably for increasing yield.

5 In one embodiment at least one active ingredient selected from group SG-7 is used on a plant comprising EVENT1 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SG-8 is used on a plant comprising EVENT1 or plant parts thereof, preferably for increasing yield.

10 In one embodiment at least one active ingredient selected from group H-1 is used on a plant comprising EVENT2 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group H-2 is used on a plant comprising EVENT2 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group H-3 is used on a plant comprising EVENT2 or plant parts thereof, preferably for increasing yield.

15 In one embodiment at least one active ingredient selected from group H-4 is used on a plant comprising EVENT2 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group H-5 is used on a plant comprising EVENT2 or plant parts thereof, preferably for increasing yield.

20 In one embodiment at least one active ingredient selected from group P-1 is used on a plant comprising EVENT2 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SAF-1 is used on a plant comprising EVENT2 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SAF-2 is used on a plant comprising EVENT2 or plant parts thereof, preferably for increasing yield.

25 In one embodiment at least one active ingredient selected from group SAF-3 is used on a plant comprising EVENT2 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SAF-4 is used on a plant comprising EVENT2 or plant parts thereof, preferably for increasing yield.

30 In one embodiment at least one active ingredient selected from group B-1 is used on a plant comprising EVENT2 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group B-2 is used on a plant comprising EVENT2 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group B-3 is used on a plant comprising EVENT2 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group B-4 is used on a plant comprising EVENT2 or plant parts thereof, preferably for increasing yield.

5 In one embodiment at least one active ingredient selected from group B-5 is used on a plant comprising EVENT2 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group I-1 is used on a plant comprising EVENT2 or plant parts thereof, preferably for increasing yield.

10 In one embodiment at least one active ingredient selected from group I-2 is used on a plant comprising EVENT2 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group I-3 is used on a plant comprising EVENT2 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group I-4 is used on a plant comprising EVENT2 or plant parts thereof, preferably for increasing yield.

15 In one embodiment at least one active ingredient selected from group I-5 is used on a plant comprising EVENT2 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group I-6 is used on a plant comprising EVENT2 or plant parts thereof, preferably for increasing yield.

20 In one embodiment at least one active ingredient selected from group F-1 is used on a plant comprising EVENT2 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group F-2 is used on a plant comprising EVENT2 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group F-3 is used on a plant comprising EVENT2 or plant parts thereof, preferably for increasing yield.

25 In one embodiment at least one active ingredient selected from group F-4 is used on a plant comprising EVENT2 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group F-5 is used on a plant comprising EVENT2 or plant parts thereof, preferably for increasing yield.

30 In one embodiment at least one active ingredient selected from group F-6 is used on a plant comprising EVENT2 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SG-1 is used on a plant comprising EVENT2 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SG-2 is used on a plant comprising EVENT2 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SG-3 is used on a plant comprising EVENT2 or plant parts thereof, preferably for increasing yield.

5 In one embodiment at least one active ingredient selected from group SG-4 is used on a plant comprising EVENT2 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SG-5 is used on a plant comprising EVENT2 or plant parts thereof, preferably for increasing yield.

10 In one embodiment at least one active ingredient selected from group SG-6 is used on a plant comprising EVENT2 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SG-7 is used on a plant comprising EVENT2 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SG-8 is used on a plant comprising EVENT2 or plant parts thereof, preferably for increasing yield.

15 In one embodiment at least one active ingredient selected from group H-1 is used on a plant comprising EVENT3 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group H-2 is used on a plant comprising EVENT3 or plant parts thereof, preferably for increasing yield.

20 In one embodiment at least one active ingredient selected from group H-3 is used on a plant comprising EVENT3 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group H-4 is used on a plant comprising EVENT3 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group H-5 is used on a plant comprising EVENT3 or plant parts thereof, preferably for increasing yield.

25 In one embodiment at least one active ingredient selected from group P-1 is used on a plant comprising EVENT3 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SAF-1 is used on a plant comprising EVENT3 or plant parts thereof, preferably for increasing yield.

30 In one embodiment at least one active ingredient selected from group SAF-2 is used on a plant comprising EVENT3 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SAF-3 is used on a plant comprising EVENT3 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SAF-4 is used on a plant comprising EVENT3 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group B-1 is used on a plant comprising EVENT3 or plant parts thereof, preferably for increasing yield.

5 In one embodiment at least one active ingredient selected from group B-2 is used on a plant comprising EVENT3 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group B-3 is used on a plant comprising EVENT3 or plant parts thereof, preferably for increasing yield.

10 In one embodiment at least one active ingredient selected from group B-4 is used on a plant comprising EVENT3 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group B-5 is used on a plant comprising EVENT3 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group I-1 is used on a plant comprising EVENT3 or plant parts thereof, preferably for increasing yield.

15 In one embodiment at least one active ingredient selected from group I-2 is used on a plant comprising EVENT3 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group I-3 is used on a plant comprising EVENT3 or plant parts thereof, preferably for increasing yield.

20 In one embodiment at least one active ingredient selected from group I-4 is used on a plant comprising EVENT3 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group I-5 is used on a plant comprising EVENT3 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group I-6 is used on a plant comprising EVENT3 or plant parts thereof, preferably for increasing yield.

25 In one embodiment at least one active ingredient selected from group F-1 is used on a plant comprising EVENT3 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group F-2 is used on a plant comprising EVENT3 or plant parts thereof, preferably for increasing yield.

30 In one embodiment at least one active ingredient selected from group F-3 is used on a plant comprising EVENT3 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group F-4 is used on a plant comprising EVENT3 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group F-5 is used on a plant comprising EVENT3 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group F-6 is used on a plant comprising EVENT3 or plant parts thereof, preferably for increasing yield.

5 In one embodiment at least one active ingredient selected from group SG-1 is used on a plant comprising EVENT3 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SG-2 is used on a plant comprising EVENT3 or plant parts thereof, preferably for increasing yield.

10 In one embodiment at least one active ingredient selected from group SG-3 is used on a plant comprising EVENT3 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SG-4 is used on a plant comprising EVENT3 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SG-5 is used on a plant comprising EVENT3 or plant parts thereof, preferably for increasing yield.

15 In one embodiment at least one active ingredient selected from group SG-6 is used on a plant comprising EVENT3 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SG-7 is used on a plant comprising EVENT3 or plant parts thereof, preferably for increasing yield.

20 In one embodiment at least one active ingredient selected from group SG-8 is used on a plant comprising EVENT3 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group H-1 is used on a plant comprising EVENT4 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group H-2 is used on a plant comprising EVENT4 or plant parts thereof, preferably for increasing yield.

25 In one embodiment at least one active ingredient selected from group H-3 is used on a plant comprising EVENT4 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group H-4 is used on a plant comprising EVENT4 or plant parts thereof, preferably for increasing yield.

30 In one embodiment at least one active ingredient selected from group H-5 is used on a plant comprising EVENT4 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group P-1 is used on a plant comprising EVENT4 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SAF-1 is used on a plant comprising EVENT4 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SAF-2 is used on a plant comprising EVENT4 or plant parts thereof, preferably for increasing yield.

5 In one embodiment at least one active ingredient selected from group SAF-3 is used on a plant comprising EVENT4 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SAF-4 is used on a plant comprising EVENT4 or plant parts thereof, preferably for increasing yield.

10 In one embodiment at least one active ingredient selected from group B-1 is used on a plant comprising EVENT4 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group B-2 is used on a plant comprising EVENT4 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group B-3 is used on a plant comprising EVENT4 or plant parts thereof, preferably for increasing yield.

15 In one embodiment at least one active ingredient selected from group B-4 is used on a plant comprising EVENT4 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group B-5 is used on a plant comprising EVENT4 or plant parts thereof, preferably for increasing yield.

20 In one embodiment at least one active ingredient selected from group I-1 is used on a plant comprising EVENT4 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group I-2 is used on a plant comprising EVENT4 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group I-3 is used on a plant comprising EVENT4 or plant parts thereof, preferably for increasing yield.

25 In one embodiment at least one active ingredient selected from group I-4 is used on a plant comprising EVENT4 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group I-5 is used on a plant comprising EVENT4 or plant parts thereof, preferably for increasing yield.

30 In one embodiment at least one active ingredient selected from group I-6 is used on a plant comprising EVENT4 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group F-1 is used on a plant comprising EVENT4 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group F-2 is used on a plant comprising EVENT4 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group F-3 is used on a plant comprising EVENT4 or plant parts thereof, preferably for increasing yield.

5 In one embodiment at least one active ingredient selected from group F-4 is used on a plant comprising EVENT4 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group F-5 is used on a plant comprising EVENT4 or plant parts thereof, preferably for increasing yield.

10 In one embodiment at least one active ingredient selected from group F-6 is used on a plant comprising EVENT4 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SG-1 is used on a plant comprising EVENT4 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SG-2 is used on a plant comprising EVENT4 or plant parts thereof, preferably for increasing yield.

15 In one embodiment at least one active ingredient selected from group SG-3 is used on a plant comprising EVENT4 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SG-4 is used on a plant comprising EVENT4 or plant parts thereof, preferably for increasing yield.

20 In one embodiment at least one active ingredient selected from group SG-5 is used on a plant comprising EVENT4 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SG-6 is used on a plant comprising EVENT4 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SG-7 is used on a plant comprising EVENT4 or plant parts thereof, preferably for increasing yield.

25 In one embodiment at least one active ingredient selected from group SG-8 is used on a plant comprising EVENT4 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group H-1 is used on a plant comprising EVENT5 or plant parts thereof, preferably for increasing yield.

30 In one embodiment at least one active ingredient selected from group H-2 is used on a plant comprising EVENT5 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group H-3 is used on a plant comprising EVENT5 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group H-4 is used on a plant comprising EVENT5 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group H-5 is used on a plant comprising EVENT5 or plant parts thereof, preferably for increasing yield.

5 In one embodiment at least one active ingredient selected from group P-1 is used on a plant comprising EVENT5 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SAF-1 is used on a plant comprising EVENT5 or plant parts thereof, preferably for increasing yield.

10 In one embodiment at least one active ingredient selected from group SAF-2 is used on a plant comprising EVENT5 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SAF-3 is used on a plant comprising EVENT5 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SAF-4 is used on a plant comprising EVENT5 or plant parts thereof, preferably for increasing yield.

15 In one embodiment at least one active ingredient selected from group B-1 is used on a plant comprising EVENT5 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group B-2 is used on a plant comprising EVENT5 or plant parts thereof, preferably for increasing yield.

20 In one embodiment at least one active ingredient selected from group B-3 is used on a plant comprising EVENT5 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group B-4 is used on a plant comprising EVENT5 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group B-5 is used on a plant comprising EVENT5 or plant parts thereof, preferably for increasing yield.

25 In one embodiment at least one active ingredient selected from group I-1 is used on a plant comprising EVENT5 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group I-2 is used on a plant comprising EVENT5 or plant parts thereof, preferably for increasing yield.

30 In one embodiment at least one active ingredient selected from group I-3 is used on a plant comprising EVENT5 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group I-4 is used on a plant comprising EVENT5 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group I-5 is used on a plant comprising EVENT5 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group I-6 is used on a plant comprising EVENT5 or plant parts thereof, preferably for increasing yield.

5 In one embodiment at least one active ingredient selected from group F-1 is used on a plant comprising EVENT5 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group F-2 is used on a plant comprising EVENT5 or plant parts thereof, preferably for increasing yield.

10 In one embodiment at least one active ingredient selected from group F-3 is used on a plant comprising EVENT5 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group F-4 is used on a plant comprising EVENT5 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group F-5 is used on a plant comprising EVENT5 or plant parts thereof, preferably for increasing yield.

15 In one embodiment at least one active ingredient selected from group F-6 is used on a plant comprising EVENT5 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SG-1 is used on a plant comprising EVENT5 or plant parts thereof, preferably for increasing yield.

20 In one embodiment at least one active ingredient selected from group SG-2 is used on a plant comprising EVENT5 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SG-3 is used on a plant comprising EVENT5 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SG-4 is used on a plant comprising EVENT5 or plant parts thereof, preferably for increasing yield.

25 In one embodiment at least one active ingredient selected from group SG-5 is used on a plant comprising EVENT5 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SG-6 is used on a plant comprising EVENT5 or plant parts thereof, preferably for increasing yield.

30 In one embodiment at least one active ingredient selected from group SG-7 is used on a plant comprising EVENT5 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SG-8 is used on a plant comprising EVENT5 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group H-1 is used on a plant comprising EVENT6 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group H-2 is used on a plant comprising EVENT6 or plant parts thereof, preferably for increasing yield.

5 In one embodiment at least one active ingredient selected from group H-3 is used on a plant comprising EVENT6 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group H-4 is used on a plant comprising EVENT6 or plant parts thereof, preferably for increasing yield.

10 In one embodiment at least one active ingredient selected from group H-5 is used on a plant comprising EVENT6 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group P-1 is used on a plant comprising EVENT6 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SAF-1 is used on a plant comprising EVENT6 or plant parts thereof, preferably for increasing yield.

15 In one embodiment at least one active ingredient selected from group SAF-2 is used on a plant comprising EVENT6 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SAF-3 is used on a plant comprising EVENT6 or plant parts thereof, preferably for increasing yield.

20 In one embodiment at least one active ingredient selected from group SAF-4 is used on a plant comprising EVENT6 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group B-1 is used on a plant comprising EVENT6 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group B-2 is used on a plant comprising EVENT6 or plant parts thereof, preferably for increasing yield.

25 In one embodiment at least one active ingredient selected from group B-3 is used on a plant comprising EVENT6 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group B-4 is used on a plant comprising EVENT6 or plant parts thereof, preferably for increasing yield.

30 In one embodiment at least one active ingredient selected from group B-5 is used on a plant comprising EVENT6 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group I-1 is used on a plant comprising EVENT6 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group I-2 is used on a plant comprising EVENT6 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group I-3 is used on a plant comprising EVENT6 or plant parts thereof, preferably for increasing yield.

5 In one embodiment at least one active ingredient selected from group I-4 is used on a plant comprising EVENT6 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group I-5 is used on a plant comprising EVENT6 or plant parts thereof, preferably for increasing yield.

10 In one embodiment at least one active ingredient selected from group I-6 is used on a plant comprising EVENT6 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group F-1 is used on a plant comprising EVENT6 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group F-2 is used on a plant comprising EVENT6 or plant parts thereof, preferably for increasing yield.

15 In one embodiment at least one active ingredient selected from group F-3 is used on a plant comprising EVENT6 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group F-4 is used on a plant comprising EVENT6 or plant parts thereof, preferably for increasing yield.

20 In one embodiment at least one active ingredient selected from group F-5 is used on a plant comprising EVENT6 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group F-6 is used on a plant comprising EVENT6 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SG-1 is used on a plant comprising EVENT6 or plant parts thereof, preferably for increasing yield.

25 In one embodiment at least one active ingredient selected from group SG-2 is used on a plant comprising EVENT6 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SG-3 is used on a plant comprising EVENT6 or plant parts thereof, preferably for increasing yield.

30 In one embodiment at least one active ingredient selected from group SG-4 is used on a plant comprising EVENT6 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SG-5 is used on a plant comprising EVENT6 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SG-6 is used on a plant comprising EVENT6 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SG-7 is used on a plant comprising EVENT6 or plant parts thereof, preferably for increasing yield.

- 5 In one embodiment at least one active ingredient selected from group SG-8 is used on a plant comprising EVENT6 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group H-1 is used on a plant comprising EVENT7 or plant parts thereof, preferably for increasing yield.

- 10 In one embodiment at least one active ingredient selected from group H-2 is used on a plant comprising EVENT7 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group H-3 is used on a plant comprising EVENT7 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group H-4 is used on a plant comprising EVENT7 or plant parts thereof, preferably for increasing yield.

- 15 In one embodiment at least one active ingredient selected from group H-5 is used on a plant comprising EVENT7 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group P-1 is used on a plant comprising EVENT7 or plant parts thereof, preferably for increasing yield.

- 20 In one embodiment at least one active ingredient selected from group SAF-1 is used on a plant comprising EVENT7 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SAF-2 is used on a plant comprising EVENT7 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SAF-3 is used on a plant comprising EVENT7 or plant parts thereof, preferably for increasing yield.

- 25 In one embodiment at least one active ingredient selected from group SAF-4 is used on a plant comprising EVENT7 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group B-1 is used on a plant comprising EVENT7 or plant parts thereof, preferably for increasing yield.

- 30 In one embodiment at least one active ingredient selected from group B-2 is used on a plant comprising EVENT7 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group B-3 is used on a plant comprising EVENT7 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group B-4 is used on a plant comprising EVENT7 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group B-5 is used on a plant comprising EVENT7 or plant parts thereof, preferably for increasing yield.

5 In one embodiment at least one active ingredient selected from group I-1 is used on a plant comprising EVENT7 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group I-2 is used on a plant comprising EVENT7 or plant parts thereof, preferably for increasing yield.

10 In one embodiment at least one active ingredient selected from group I-3 is used on a plant comprising EVENT7 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group I-4 is used on a plant comprising EVENT7 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group I-5 is used on a plant comprising EVENT7 or plant parts thereof, preferably for increasing yield.

15 In one embodiment at least one active ingredient selected from group I-6 is used on a plant comprising EVENT7 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group F-1 is used on a plant comprising EVENT7 or plant parts thereof, preferably for increasing yield.

20 In one embodiment at least one active ingredient selected from group F-2 is used on a plant comprising EVENT7 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group F-3 is used on a plant comprising EVENT7 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group F-4 is used on a plant comprising EVENT7 or plant parts thereof, preferably for increasing yield.

25 In one embodiment at least one active ingredient selected from group F-5 is used on a plant comprising EVENT7 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group F-6 is used on a plant comprising EVENT7 or plant parts thereof, preferably for increasing yield.

30 In one embodiment at least one active ingredient selected from group SG-1 is used on a plant comprising EVENT7 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SG-2 is used on a plant comprising EVENT7 or plant parts thereof, preferably for increasing yield.

- In one embodiment at least one active ingredient selected from group SG-3 is used on a plant comprising EVENT7 or plant parts thereof, preferably for increasing yield.
- In one embodiment at least one active ingredient selected from group SG-4 is used on a plant comprising EVENT7 or plant parts thereof, preferably for increasing yield.
- 5 In one embodiment at least one active ingredient selected from group SG-5 is used on a plant comprising EVENT7 or plant parts thereof, preferably for increasing yield.
- In one embodiment at least one active ingredient selected from group SG-6 is used on a plant comprising EVENT7 or plant parts thereof, preferably for increasing yield.
- In one embodiment at least one active ingredient selected from group SG-7 is used on a plant
10 comprising EVENT7 or plant parts thereof, preferably for increasing yield.
- In one embodiment at least one active ingredient selected from group SG-8 is used on a plant comprising EVENT7 or plant parts thereof, preferably for increasing yield.
- In one embodiment at least one active ingredient selected from group H-1 is used on a plant comprising ALLELE1 or plant parts thereof, preferably for increasing yield.
- 15 In one embodiment at least one active ingredient selected from group H-2 is used on a plant comprising ALLELE1 or plant parts thereof, preferably for increasing yield.
- In one embodiment at least one active ingredient selected from group H-3 is used on a plant comprising ALLELE1 or plant parts thereof, preferably for increasing yield.
- In one embodiment at least one active ingredient selected from group H-4 is used on a plant comprising
20 ALLELE1 or plant parts thereof, preferably for increasing yield.
- In one embodiment at least one active ingredient selected from group H-5 is used on a plant comprising ALLELE1 or plant parts thereof, preferably for increasing yield.
- In one embodiment at least one active ingredient selected from group P-1 is used on a plant comprising ALLELE1 or plant parts thereof, preferably for increasing yield.
- 25 In one embodiment at least one active ingredient selected from group SAF-1 is used on a plant comprising ALLELE1 or plant parts thereof, preferably for increasing yield.
- In one embodiment at least one active ingredient selected from group SAF-2 is used on a plant comprising ALLELE1 or plant parts thereof, preferably for increasing yield.
- In one embodiment at least one active ingredient selected from group SAF-3 is used on a plant
30 comprising ALLELE1 or plant parts thereof, preferably for increasing yield.
- In one embodiment at least one active ingredient selected from group SAF-4 is used on a plant comprising ALLELE1 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group B-1 is used on a plant comprising ALLELE1 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group B-2 is used on a plant comprising ALLELE1 or plant parts thereof, preferably for increasing yield.

5 In one embodiment at least one active ingredient selected from group B-3 is used on a plant comprising ALLELE1 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group B-4 is used on a plant comprising ALLELE1 or plant parts thereof, preferably for increasing yield.

10 In one embodiment at least one active ingredient selected from group B-5 is used on a plant comprising ALLELE1 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group I-1 is used on a plant comprising ALLELE1 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group I-2 is used on a plant comprising ALLELE1 or plant parts thereof, preferably for increasing yield.

15 In one embodiment at least one active ingredient selected from group I-3 is used on a plant comprising ALLELE1 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group I-4 is used on a plant comprising ALLELE1 or plant parts thereof, preferably for increasing yield.

20 In one embodiment at least one active ingredient selected from group I-5 is used on a plant comprising ALLELE1 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group I-6 is used on a plant comprising ALLELE1 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group F-1 is used on a plant comprising ALLELE1 or plant parts thereof, preferably for increasing yield.

25 In one embodiment at least one active ingredient selected from group F-2 is used on a plant comprising ALLELE1 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group F-3 is used on a plant comprising ALLELE1 or plant parts thereof, preferably for increasing yield.

30 In one embodiment at least one active ingredient selected from group F-4 is used on a plant comprising ALLELE1 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group F-5 is used on a plant comprising ALLELE1 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group F-6 is used on a plant comprising ALLELE1 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SG-1 is used on a plant comprising ALLELE1 or plant parts thereof, preferably for increasing yield.

5 In one embodiment at least one active ingredient selected from group SG-2 is used on a plant comprising ALLELE1 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SG-3 is used on a plant comprising ALLELE1 or plant parts thereof, preferably for increasing yield.

10 In one embodiment at least one active ingredient selected from group SG-4 is used on a plant comprising ALLELE1 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SG-5 is used on a plant comprising ALLELE1 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SG-6 is used on a plant comprising ALLELE1 or plant parts thereof, preferably for increasing yield.

15 In one embodiment at least one active ingredient selected from group SG-7 is used on a plant comprising ALLELE1 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SG-8 is used on a plant comprising ALLELE1 or plant parts thereof, preferably for increasing yield.

20 In one embodiment at least one active ingredient selected from group H-1 is used on a plant comprising ALLELE2 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group H-2 is used on a plant comprising ALLELE2 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group H-3 is used on a plant comprising ALLELE2 or plant parts thereof, preferably for increasing yield.

25 In one embodiment at least one active ingredient selected from group H-4 is used on a plant comprising ALLELE2 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group H-5 is used on a plant comprising ALLELE2 or plant parts thereof, preferably for increasing yield.

30 In one embodiment at least one active ingredient selected from group P-1 is used on a plant comprising ALLELE2 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SAF-1 is used on a plant comprising ALLELE2 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SAF-2 is used on a plant comprising ALLELE2 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SAF-3 is used on a plant comprising ALLELE2 or plant parts thereof, preferably for increasing yield.

5 In one embodiment at least one active ingredient selected from group SAF-4 is used on a plant comprising ALLELE2 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group B-1 is used on a plant comprising ALLELE2 or plant parts thereof, preferably for increasing yield.

10 In one embodiment at least one active ingredient selected from group B-2 is used on a plant comprising ALLELE2 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group B-3 is used on a plant comprising ALLELE2 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group B-4 is used on a plant comprising ALLELE2 or plant parts thereof, preferably for increasing yield.

15 In one embodiment at least one active ingredient selected from group B-5 is used on a plant comprising ALLELE2 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group I-1 is used on a plant comprising ALLELE2 or plant parts thereof, preferably for increasing yield.

20 In one embodiment at least one active ingredient selected from group I-2 is used on a plant comprising ALLELE2 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group I-3 is used on a plant comprising ALLELE2 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group I-4 is used on a plant comprising ALLELE2 or plant parts thereof, preferably for increasing yield.

25 In one embodiment at least one active ingredient selected from group I-5 is used on a plant comprising ALLELE2 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group I-6 is used on a plant comprising ALLELE2 or plant parts thereof, preferably for increasing yield.

30 In one embodiment at least one active ingredient selected from group F-1 is used on a plant comprising ALLELE2 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group F-2 is used on a plant comprising ALLELE2 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group F-3 is used on a plant comprising ALLELE2 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group F-4 is used on a plant comprising ALLELE2 or plant parts thereof, preferably for increasing yield.

5 In one embodiment at least one active ingredient selected from group F-5 is used on a plant comprising ALLELE2 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group F-6 is used on a plant comprising ALLELE2 or plant parts thereof, preferably for increasing yield.

10 In one embodiment at least one active ingredient selected from group SG-1 is used on a plant comprising ALLELE2 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SG-2 is used on a plant comprising ALLELE2 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SG-3 is used on a plant comprising ALLELE2 or plant parts thereof, preferably for increasing yield.

15 In one embodiment at least one active ingredient selected from group SG-4 is used on a plant comprising ALLELE2 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SG-5 is used on a plant comprising ALLELE2 or plant parts thereof, preferably for increasing yield.

20 In one embodiment at least one active ingredient selected from group SG-6 is used on a plant comprising ALLELE2 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SG-7 is used on a plant comprising ALLELE2 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SG-8 is used on a plant comprising ALLELE2 or plant parts thereof, preferably for increasing yield.

25 In one embodiment at least one active ingredient selected from group H-1 is used on a plant comprising ALLELE3 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group H-2 is used on a plant comprising ALLELE3 or plant parts thereof, preferably for increasing yield.

30 In one embodiment at least one active ingredient selected from group H-3 is used on a plant comprising ALLELE3 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group H-4 is used on a plant comprising ALLELE3 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group H-5 is used on a plant comprising ALLELE3 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group P-1 is used on a plant comprising ALLELE3 or plant parts thereof, preferably for increasing yield.

5 In one embodiment at least one active ingredient selected from group SAF-1 is used on a plant comprising ALLELE3 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SAF-2 is used on a plant comprising ALLELE3 or plant parts thereof, preferably for increasing yield.

10 In one embodiment at least one active ingredient selected from group SAF-3 is used on a plant comprising ALLELE3 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SAF-4 is used on a plant comprising ALLELE3 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group B-1 is used on a plant comprising ALLELE3 or plant parts thereof, preferably for increasing yield.

15 In one embodiment at least one active ingredient selected from group B-2 is used on a plant comprising ALLELE3 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group B-3 is used on a plant comprising ALLELE3 or plant parts thereof, preferably for increasing yield.

20 In one embodiment at least one active ingredient selected from group B-4 is used on a plant comprising ALLELE3 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group B-5 is used on a plant comprising ALLELE3 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group I-1 is used on a plant comprising ALLELE3 or plant parts thereof, preferably for increasing yield.

25 In one embodiment at least one active ingredient selected from group I-2 is used on a plant comprising ALLELE3 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group I-3 is used on a plant comprising ALLELE3 or plant parts thereof, preferably for increasing yield.

30 In one embodiment at least one active ingredient selected from group I-4 is used on a plant comprising ALLELE3 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group I-5 is used on a plant comprising ALLELE3 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group I-6 is used on a plant comprising ALLELE3 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group F-1 is used on a plant comprising ALLELE3 or plant parts thereof, preferably for increasing yield.

5 In one embodiment at least one active ingredient selected from group F-2 is used on a plant comprising ALLELE3 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group F-3 is used on a plant comprising ALLELE3 or plant parts thereof, preferably for increasing yield.

10 In one embodiment at least one active ingredient selected from group F-4 is used on a plant comprising ALLELE3 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group F-5 is used on a plant comprising ALLELE3 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group F-6 is used on a plant comprising ALLELE3 or plant parts thereof, preferably for increasing yield.

15 In one embodiment at least one active ingredient selected from group SG-1 is used on a plant comprising ALLELE3 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SG-2 is used on a plant comprising ALLELE3 or plant parts thereof, preferably for increasing yield.

20 In one embodiment at least one active ingredient selected from group SG-3 is used on a plant comprising ALLELE3 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SG-4 is used on a plant comprising ALLELE3 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SG-5 is used on a plant comprising ALLELE3 or plant parts thereof, preferably for increasing yield.

25 In one embodiment at least one active ingredient selected from group SG-6 is used on a plant comprising ALLELE3 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SG-7 is used on a plant comprising ALLELE3 or plant parts thereof, preferably for increasing yield.

30 In one embodiment at least one active ingredient selected from group SG-8 is used on a plant comprising ALLELE3 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group H-1 is used on a plant comprising ALLELE4 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group H-2 is used on a plant comprising ALLELE4 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group H-3 is used on a plant comprising ALLELE4 or plant parts thereof, preferably for increasing yield.

5 In one embodiment at least one active ingredient selected from group H-4 is used on a plant comprising ALLELE4 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group H-5 is used on a plant comprising ALLELE4 or plant parts thereof, preferably for increasing yield.

10 In one embodiment at least one active ingredient selected from group P-1 is used on a plant comprising ALLELE4 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SAF-1 is used on a plant comprising ALLELE4 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SAF-2 is used on a plant comprising ALLELE4 or plant parts thereof, preferably for increasing yield.

15 In one embodiment at least one active ingredient selected from group SAF-3 is used on a plant comprising ALLELE4 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SAF-4 is used on a plant comprising ALLELE4 or plant parts thereof, preferably for increasing yield.

20 In one embodiment at least one active ingredient selected from group B-1 is used on a plant comprising ALLELE4 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group B-2 is used on a plant comprising ALLELE4 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group B-3 is used on a plant comprising ALLELE4 or plant parts thereof, preferably for increasing yield.

25 In one embodiment at least one active ingredient selected from group B-4 is used on a plant comprising ALLELE4 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group B-5 is used on a plant comprising ALLELE4 or plant parts thereof, preferably for increasing yield.

30 In one embodiment at least one active ingredient selected from group I-1 is used on a plant comprising ALLELE4 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group I-2 is used on a plant comprising ALLELE4 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group I-3 is used on a plant comprising ALLELE4 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group I-4 is used on a plant comprising ALLELE4 or plant parts thereof, preferably for increasing yield.

5 In one embodiment at least one active ingredient selected from group I-5 is used on a plant comprising ALLELE4 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group I-6 is used on a plant comprising ALLELE4 or plant parts thereof, preferably for increasing yield.

10 In one embodiment at least one active ingredient selected from group F-1 is used on a plant comprising ALLELE4 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group F-2 is used on a plant comprising ALLELE4 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group F-3 is used on a plant comprising ALLELE4 or plant parts thereof, preferably for increasing yield.

15 In one embodiment at least one active ingredient selected from group F-4 is used on a plant comprising ALLELE4 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group F-5 is used on a plant comprising ALLELE4 or plant parts thereof, preferably for increasing yield.

20 In one embodiment at least one active ingredient selected from group F-6 is used on a plant comprising ALLELE4 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SG-1 is used on a plant comprising ALLELE4 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SG-2 is used on a plant comprising ALLELE4 or plant parts thereof, preferably for increasing yield.

25 In one embodiment at least one active ingredient selected from group SG-3 is used on a plant comprising ALLELE4 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SG-4 is used on a plant comprising ALLELE4 or plant parts thereof, preferably for increasing yield.

30 In one embodiment at least one active ingredient selected from group SG-5 is used on a plant comprising ALLELE4 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SG-6 is used on a plant comprising ALLELE4 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SG-7 is used on a plant comprising ALLELE4 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SG-8 is used on a plant comprising ALLELE4 or plant parts thereof, preferably for increasing yield.

5 In one embodiment at least one active ingredient selected from group H-1 is used on a plant comprising ALLELE5 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group H-2 is used on a plant comprising ALLELE5 or plant parts thereof, preferably for increasing yield.

10 In one embodiment at least one active ingredient selected from group H-3 is used on a plant comprising ALLELE5 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group H-4 is used on a plant comprising ALLELE5 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group H-5 is used on a plant comprising ALLELE5 or plant parts thereof, preferably for increasing yield.

15 In one embodiment at least one active ingredient selected from group P-1 is used on a plant comprising ALLELE5 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SAF-1 is used on a plant comprising ALLELE5 or plant parts thereof, preferably for increasing yield.

20 In one embodiment at least one active ingredient selected from group SAF-2 is used on a plant comprising ALLELE5 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SAF-3 is used on a plant comprising ALLELE5 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SAF-4 is used on a plant comprising ALLELE5 or plant parts thereof, preferably for increasing yield.

25 In one embodiment at least one active ingredient selected from group B-1 is used on a plant comprising ALLELE5 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group B-2 is used on a plant comprising ALLELE5 or plant parts thereof, preferably for increasing yield.

30 In one embodiment at least one active ingredient selected from group B-3 is used on a plant comprising ALLELE5 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group B-4 is used on a plant comprising ALLELE5 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group B-5 is used on a plant comprising ALLELE5 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group I-1 is used on a plant comprising ALLELE5 or plant parts thereof, preferably for increasing yield.

5 In one embodiment at least one active ingredient selected from group I-2 is used on a plant comprising ALLELE5 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group I-3 is used on a plant comprising ALLELE5 or plant parts thereof, preferably for increasing yield.

10 In one embodiment at least one active ingredient selected from group I-4 is used on a plant comprising ALLELE5 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group I-5 is used on a plant comprising ALLELE5 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group I-6 is used on a plant comprising ALLELE5 or plant parts thereof, preferably for increasing yield.

15 In one embodiment at least one active ingredient selected from group F-1 is used on a plant comprising ALLELE5 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group F-2 is used on a plant comprising ALLELE5 or plant parts thereof, preferably for increasing yield.

20 In one embodiment at least one active ingredient selected from group F-3 is used on a plant comprising ALLELE5 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group F-4 is used on a plant comprising ALLELE5 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group F-5 is used on a plant comprising ALLELE5 or plant parts thereof, preferably for increasing yield.

25 In one embodiment at least one active ingredient selected from group F-6 is used on a plant comprising ALLELE5 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SG-1 is used on a plant comprising ALLELE5 or plant parts thereof, preferably for increasing yield.

30 In one embodiment at least one active ingredient selected from group SG-2 is used on a plant comprising ALLELE5 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SG-3 is used on a plant comprising ALLELE5 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SG-4 is used on a plant comprising ALLELE5 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SG-5 is used on a plant comprising ALLELE5 or plant parts thereof, preferably for increasing yield.

5 In one embodiment at least one active ingredient selected from group SG-6 is used on a plant comprising ALLELE5 or plant parts thereof, preferably for increasing yield.

In one embodiment at least one active ingredient selected from group SG-7 is used on a plant comprising ALLELE5 or plant parts thereof, preferably for increasing yield.

10 In one embodiment at least one active ingredient selected from group SG-8 is used on a plant comprising ALLELE5 or plant parts thereof, preferably for increasing yield.

Uses

The treatment of the plants and plant parts with active ingredients or mixtures thereof is carried out directly or by acting on the environment, habitat or storage space using customary treatment methods, for example by dipping, spraying, atomizing, misting, evaporating, dusting, fogging, scattering,
15 foaming, painting on, spreading, injecting, drenching, trickle irrigation and, in the case of propagation material, in particular in the case of seed, furthermore by the dry seed treatment method, the wet seed treatment method, the slurry treatment method, by encrusting, by coating with one or more coats and the like. It is furthermore possible to apply the active ingredients or mixtures thereof by the ultra-low volume method or to inject the active ingredient preparation or the active ingredient itself into the soil.

20 A preferred direct treatment of the plants is the leaf application treatment, i.e. active ingredients or mixtures thereof or compositions comprising active ingredients or mixtures thereof are applied to the foliage, it being possible for the treatment frequency and the application rate to be matched to achieve the yield increase in question.

In the case of systemically active compounds, active ingredients or mixtures thereof or compositions
25 comprising active ingredients or mixtures thereof reach the plants via the root system. In this case, the treatment of the plants is effected by allowing active ingredients or mixtures thereof or compositions comprising active ingredients or mixtures thereof to act on the environment of the plant. This can be done for example by drenching, incorporating in the soil or into the nutrient solution, i.e. the location of the plant (for example the soil or hydroponic systems) is impregnated with a liquid form of active
30 ingredients or mixtures thereof or compositions comprising active ingredients or mixtures thereof, or by soil application, i.e. the active ingredients or mixtures thereof or compositions comprising active ingredients or mixtures thereof are incorporated into the location of the plants in solid form (for example in the form of granules).

35 More particularly, the inventive use exhibits the advantages described on *Brassicaceae* plants, plant parts thereof, plant propagation material or the soil in which *Brassicaceae* plants are grown or intended

to be grown in spray application, in seed treatment, in drip and drench applications, in-furrow application, chemigation, i.e. by addition of active ingredients or mixtures thereof to the irrigation water, and in hydroponic/mineral systems.

The combined use of active ingredients or mixtures thereof or compositions comprising active ingredients or mixtures thereof, with genetically modified cultivars, especially of transgenic oilseed rape cultivars, is additionally likewise possible.

Application Rates

When using the active ingredients or mixtures thereof, the application rates can be varied within a relatively wide range, depending on the kind of application. The application rate of the inventive active ingredients is generally known to the skilled person. Generally, the following ranges can be applied

- in the case of treatment of plant parts, for example leaves: from 0.1 to 10 000 g/ha, preferably from 10 to 1000 g/ha, more preferably from 50 to 500 g/ha (in the case of application by watering or dripping, it is even possible to reduce the application rate, especially when inert substrates such as rockwool or perlite are used);
- in the case of seed treatment: from 0.1 to 200 g per 100 kg of seed, preferably from 1 to 150 g per 100 kg of seed, more preferably from 2.5 to 25 g per 100 kg of seed, even more preferably from 2.5 to 12.5 g per 100 kg of seed;
- in the case of soil treatment: from 0.1 to 10 000 g/ha, preferably from 1 to 5000 g/ha.

These application rates are merely by way of example and are not limiting for the purposes of the invention.

Formulations

The present invention further relates to a composition for increasing yield, comprising at least one active ingredient or mixtures thereof. These are preferably fungicidal, bactericidal, herbicidal, insecticidal, acaricidal, or nematocidal compositions which comprise agriculturally suitable auxiliaries, solvents, carriers, surfactants or extenders.

According to the invention, a carrier is a natural or synthetic, organic or inorganic substance with which the active ingredients are mixed or combined for better applicability, in particular for application to plants or plant parts or seed. The carrier, which may be solid or liquid, is generally inert and should be suitable for use in agriculture.

Useful solid carriers include: for example ammonium salts and natural rock flours, such as kaolins, clays, talc, chalk, quartz, attapulgite, montmorillonite or diatomaceous earth, and synthetic rock flours, such as finely divided silica, alumina and silicates; useful solid carriers for granules include: for example, crushed and fractionated natural rocks such as calcite, marble, pumice, sepiolite and dolomite, and also synthetic granules of inorganic and organic flours, and granules of organic material such as paper, sawdust, coconut shells, maize cobs and tobacco stalks; useful emulsifiers and/or foam-formers

include: for example nonionic and anionic emulsifiers, such as polyoxyethylene fatty acid esters, polyoxyethylene fatty alcohol ethers, for example alkylaryl polyglycol ethers, alkylsulphonates, alkyl sulphates, arylsulphonates and also protein hydrolysates; suitable dispersants are nonionic and/or ionic substances, for example from the classes of the alcohol-POE and/or -POP ethers, acid and/or POP POE esters, alkylaryl and/or POP POE ethers, fat and/or POP POE adducts, POE- and/or POP-polyol derivatives, POE- and/or POP-sorbitan or -sugar adducts, alkyl or aryl sulphates, alkyl- or arylsulphonates and alkyl or aryl phosphates or the corresponding PO-ether adducts. Additionally suitable are oligo- or polymers, for example those derived from vinylic monomers, from acrylic acid, from EO and/or PO alone or in combination with, for example, (poly)alcohols or (poly)amines. It is also possible to use lignin and its sulphonic acid derivatives, unmodified and modified celluloses, aromatic and/or aliphatic sulphonic acids and also their adducts with formaldehyde.

The active ingredients or mixtures thereof can be converted to the customary formulations, such as solutions, emulsions, wettable powders, water- and oil-based suspensions, powders, dusts, pastes, soluble powders, soluble granules, granules for broadcasting, suspoemulsion concentrates, natural products impregnated with active ingredient, synthetic substances impregnated with active ingredients or mixtures thereof, fertilizers and also microencapsulations in polymeric substances.

The active ingredients or mixtures thereof can be applied as such, in the form of their formulations or the use forms prepared therefrom, such as ready-to-use solutions, emulsions, water- or oil-based suspensions, powders, wettable powders, pastes, soluble powders, dusts, soluble granules, granules for broadcasting, suspoemulsion concentrates, natural products impregnated with active ingredient, synthetic substances impregnated with active ingredient, fertilizers and also microencapsulations in polymeric substances. Application is accomplished in a customary manner, for example by watering, spraying, atomizing, broadcasting, dusting, foaming, spreading-on and the like. It is also possible to deploy the active ingredients by the ultra-low volume method or to inject the active ingredient preparation/the active ingredient itself into the soil. It is also possible to treat the seed of the plants.

The formulations mentioned can be prepared in a manner known per se, for example by mixing the active ingredients or mixtures thereof with at least one customary extender, solvent or diluent, emulsifier, dispersant and/or binder or fixing agent, wetting agent, a water repellent, if appropriate siccatives and UV stabilizers and if appropriate dyes and pigments, antifoams, preservatives, secondary thickeners, stickers, gibberellins and also other processing auxiliaries.

The present invention includes not only formulations which are already ready for use and can be deployed with a suitable apparatus to the plant or the seed, but also commercial concentrates which have to be diluted with water prior to use.

active ingredients or mixtures thereof may be present as such or in their (commercial) formulations and in the use forms prepared from these formulations as a mixture with other (known) active ingredients, such as insecticides, attractants, sterilants, bactericides, acaricides, nematocides, fungicides, growth regulators, herbicides, fertilizers, safeners and/or semiochemicals.

The auxiliaries used may be those substances which are suitable for imparting particular properties to the composition itself or and/or to preparations derived therefrom (for example spray liquors, seed dressings), such as certain technical properties and/or also particular biological properties. Typical auxiliaries include: extenders, solvents and carriers.

- 5 Suitable extenders are, for example, water, polar and nonpolar organic chemical liquids, for example from the classes of the aromatic and nonaromatic hydrocarbons (such as paraffins, alkylbenzenes, alkylnaphthalenes, chlorobenzenes), the alcohols and polyols (which may optionally also be substituted, etherified and/or esterified), the ketones (such as acetone, cyclohexanone), esters (including fats and oils) and (poly)ethers, the unsubstituted and substituted amines, amides, lactams (such as N-
10 alkylpyrrolidones) and lactones, the sulphones and sulphoxides (such as dimethyl sulphoxide).

Liquefied gaseous extenders or carriers are understood to mean liquids which are gaseous at standard temperature and under standard pressure, for example aerosol propellants such as halohydrocarbons, or else butane, propane, nitrogen and carbon dioxide.

- 15 In the formulations it is possible to use tackifiers such as carboxymethylcellulose, natural and synthetic polymers in the form of powders, granules or latices, such as gum arabic, polyvinyl alcohol and polyvinyl acetate, or else natural phospholipids such as cephalins and lecithins and synthetic phospholipids. Further additives may be mineral and vegetable oils.

- If the extender used is water, it is also possible to use, for example, organic solvents as auxiliary solvents. Useful liquid solvents are essentially: aromatics such as xylene, toluene or alkylnaphthalenes, chlorinated aromatics or chlorinated aliphatic hydrocarbons such as chlorobenzenes, chloroethylenes or
20 methylene chloride, aliphatic hydrocarbons such as cyclohexane or paraffins, for example petroleum fractions, alcohols such as butanol or glycol and their ethers and esters, ketones such as acetone, methyl ethyl ketone, methyl isobutyl ketone or cyclohexanone, strongly polar solvents such as dimethylformamide and dimethyl sulphoxide, or else water.

- 25 Compositions comprising active ingredients or mixtures thereof may additionally comprise further components, for example surfactants. Suitable surfactants are emulsifiers and/or foam formers, dispersants or wetting agents having ionic or nonionic properties, or mixtures of these surfactants. Examples thereof are salts of polyacrylic acid, salts of lignosulphonic acid, salts of phenolsulphonic acid or naphthalenesulphonic acid, polycondensates of ethylene oxide with fatty alcohols or with fatty acids
30 or with fatty amines, substituted phenols (preferably alkylphenols or arylphenols), salts of sulphosuccinic esters, taurine derivatives (preferably alkyl taurates), phosphoric esters of polyethoxylated alcohols or phenols, fatty esters of polyols, and derivatives of the compounds containing sulphates, sulphonates and phosphates, for example alkylaryl polyglycol ethers, alkylsulphonates, alkyl sulphates, arylsulphonates, protein hydrolysates, lignosulphite waste liquors and
35 methylcellulose. The presence of a surfactant is necessary if one of the active ingredients and/or one of the inert carriers is insoluble in water and when application is effected in water. The proportion of surfactants is between 5 and 40 per cent by weight of the inventive composition.

It is possible to use dyes such as inorganic pigments, for example iron oxide, titanium oxide and Prussian Blue, and organic dyes such as alizarin dyes, azo dyes and metal phthalocyanine dyes, and trace nutrients such as salts of iron, manganese, boron, copper, cobalt, molybdenum and zinc.

Further additives may be perfumes, mineral or vegetable, optionally modified oils, waxes and nutrients
5 (including trace nutrients), such as salts of iron, manganese, boron, copper, cobalt, molybdenum and zinc.

Additional components may be stabilizers, such as cold stabilizers, preservatives, antioxidants, light stabilizers, or other agents which improve chemical and/or physical stability.

If appropriate, other additional components may also be present, for example protective colloids,
10 binders, adhesives, thickeners, thixotropic substances, penetrants, stabilizers, sequestering agents, complex formers. In general, the active ingredients or mixtures thereof can be combined with any solid or liquid additive commonly used for formulation purposes.

The formulations contain generally between 0.05 and 99% by weight, 0.01 and 98% by weight, preferably between 0.1 and 95% by weight, more preferably between 0.5 and 90% of active ingredients
15 or mixtures thereof, most preferably between 10 and 70 per cent by weight.

The formulations described above can be used for increasing yield, in which the compositions comprising compounds of the formula (I) are applied to the *Brassicaceae* and/or in their habitat.

Plants

According to the invention all plants and plant parts can be treated. By plants is meant all plants and
20 plant populations such as desirable and undesirable wild plants, cultivars and plant varieties (whether or not protectable by plant variety or plant breeder's rights). Cultivars and plant varieties can be plants obtained by conventional propagation and breeding methods which can be assisted or supplemented by one or more biotechnological methods such as by use of double haploids, protoplast fusion, random and directed mutagenesis, molecular or genetic markers or by bioengineering and genetic engineering
25 methods. By plant parts is meant all above ground and below ground parts and organs of plants such as shoot, leaf, blossom and root, whereby for example leaves, needles, stems, branches, blossoms, fruiting bodies, fruits and seed as well as roots, corms and rhizomes are listed. Crops and vegetative and generative propagating material, for example cuttings, corms, rhizomes, runners, slips and seeds also belong to plant parts. Seeds are preferred plant parts.

30 In one embodiment crop plants belonging to the plant family Brassicaceae are Brassica plants.

In a preferred embodiment crop species, cultivars and varieties belonging to the plant genus Brassica are

- *Brassica carinata*: Abyssinian mustard or Abyssinian cabbage
- *Brassica elongata*: elongated mustard
- *Brassica fruticulosa*: Mediterranean cabbage

- *Brassica juncea*: Indian mustard, brown and leaf mustards, Sarepta mustard
- *Brassica napus* comprising winter rapeseed, spring rapeseed, rutabaga (*Brassica napus* subsp *rapifera* swede/Swedish turnip/swede turnip)
- *Brassica narinosa*: broadbeaked mustard
- 5 • *Brassica nigra*: black mustard
- *Brassica oleracea* comprising cultivars like kale, cabbage, broccoli, cauliflower, kai-lan, Brussels sprouts, kohlrabi
- *Brassica perviridis*: tender green, mustard spinach
- *Brassica rapa* (syn *B. campestris*) comprising Chinese cabbage, turnip, rapini, komatsuna
- 10 • *Brassica rupestris*: brown mustard
- *Brassica septiceps*: seventop turnip
- *Brassica tournefortii*: Asian mustard
- *Brassica alba* (syn *Sinapis alba*, white mustard)
- Canola varieties

15 To use the name canola, an oilseed plant must meet the following internationally regulated standard:

"Seeds of the genus *Brassica* (*Brassica napus*, *Brassica rapa* or *Brassica juncea*) from which the oil shall contain less than 2% erucic acid in its fatty acid profile and the solid component shall contain less than 30 micromoles of any one or any mixture of 3-butenyl glucosinolate, 4-pentenyl glucosinolate, 2-hydroxy-3 butenyl glucosinolate, and 2-hydroxy- 4-pentenyl glucosinolate per gram of air-dry, oil-free
20 solid."

Further preferred crop plants belonging to the plant family *Brassicaceae* are horseradish (*Armoracia rusticana*), radish (e.g. *Raphanus sativus* var. *oleiformis*, *Raphanus sativus* L. var. *sativus*).

More preferred *Brassica* plants, plant parts including seeds according to the present invention are oilseed rape plants, plant parts including seeds (*Brassica napus*), Canola plants, plant parts including seeds or
25 *Brassica juncea* plants, plant parts including seeds; more preferred winter oilseed rape plants, plant parts including seeds (*Brassica napus*), spring oilseed rape plants, plant parts including seeds or Canola, plant parts including seeds.

In one aspect *Brassica napus* or *juncea* plants, plant parts or seeds are hybrid plants, plant parts including seeds. In another aspect *Brassica napus* or *juncea* hybrids are Ogura hybrids, Ms8/Rf3 hybrids
30 (marketed under the tradename Invigor) or Ms11/Rf3 hybrids.

In another embodiment the *Brassica napus* or *juncea* plants, plant parts including seeds are tolerant to one or more of the herbicides selected from the group of glufosinate, glyphosate (tradename RoundupReady), imazamethabenz, imazamethabenz-methyl, imazamox, imazamox-ammonium,

imazapic, imazapic-ammonium, imazapyr, imazapyr-isopropyl-ammonium, imazaquin, imazaquin-ammonium, imazethapyr, imazethapyr-ammonium, atrazine, simazine.

The term "growth stage" refers to the growth stages as defined by the BBCH Codes in "Growth stages of mono- and dicotyledonous plants", 2nd edition 2001, edited by Uwe Meier from the Federal Biological
 5 Research Centre for Agriculture and Forestry. The BBCH codes are a well-established system for a uniform coding of phenologically similar growth stages of all mono- and dicotyledonous plant species. The abbreviation BBCH derives from "Biologische Bundesanstalt, Bundessortenamt und Chemische
 10 Industrie".

Some of these BBCH growth stages and BBCH codes for oilseed rape plants are indicated in the
 10 following.

Growth stage 1: Leaf development¹

BBCH 10 - Cotyledons completely unfolded

BBCH 11 - First leaf unfolded

BBCH 12 – 2nd leaf unfolded

15 BBCH 13 – 3rd leaf unfolded

BBCH 14 - 18 Stages continuous till . . . (4 – 8th leaf unfolded)

BBCH 19 - 9 or more leaves unfolded

Growth stage 2: Formation of side shoots

BBCH 20 - No side shoots

20 BBCH 21 - Beginning of side shoot development: first side shoot detectable

BBCH 22 – 2nd side shoots detectable

BBCH 23 – 3rd side shoots detectable

BBCH 24 – 4th side shoots detectable

BBCH 25 – 5th side shoots detectable

25 BBCH 26 - 28 - Stages continuous till . . . (6 – 8 side shoots detectable)

BBCH 29 End of side shoot development: 9 or more side shoots detectable

Growth stage 3: Stem elongation²

BBCH 30 - Beginning of stem elongation: no internodes ("rosette")

BBCH 31 - 1 visibly extended internode

30 BBCH 32 – 2nd visibly extended internode

BBCH 33 – 3rd visibly extended internode

...

BBCH 39 - 9 or more visibly extended internodes

...

Growth stage 8: Ripening

5 BBCH 80 - Beginning of ripening: seed green, filling pod cavity

BBCH 81 - 10% of pods ripe, seeds dark and hard

BBCH 82 - 20% of pods ripe, seeds dark and hard

BBCH 83 - 30% of pods ripe, seeds dark and hard

BBCH 84 - 40% of pods ripe, seeds dark and hard

10 BBCH 85 - 50% of pods ripe, seeds dark and hard

BBCH 86 - 60% of pods ripe, seeds dark and hard

BBCH 87 - 70% of pods ripe, seeds dark and hard

BBCH 88 - 80% of pods ripe, seeds dark and hard

BBCH 89 - Fully ripe: nearly all pods ripe, seeds dark and hard

15 Growth stage 9: Senescence

BBCH 97- Plant dead and dry

BBCH 99 - Harvested product

Particular preference is given in accordance with the invention to treating plants of the plant cultivars which are each commercially available or in use. Plant cultivars are understood to mean plants which have new properties ("traits") and which have been obtained by conventional breeding, by mutagenesis or with the aid of recombinant DNA techniques. Crop plants may accordingly be plants which can be obtained by conventional breeding and optimization methods or by biotechnology and genetic engineering methods or combinations of these methods, including the transgenic plants and including the plant varieties which can and cannot be protected by plant variety rights.

25 The method according to the invention can thus also be used for the treatment of genetically modified organisms (GMOs), for example plants or seeds. Genetically modified plants (or transgenic plants) are plants in which a heterologous gene has been integrated stably into the genome. The term "heterologous gene" means essentially a gene which is provided or assembled outside the plant and which, on introduction into the cell nucleus genome, imparts new or improved agronomic or other properties to the chloroplast genome or the mitochondrial genome of the transformed plant by virtue of it expressing a protein or polypeptide of interest or by virtue of another gene which is present in the plant, or other genes which are present in the plant, being downregulated or silenced (for example by means of antisense technology, co-suppression technology or RNAi technology [RNA interference]). A

30

heterologous gene present in the genome is likewise referred to as a transgene. A transgene which is defined by its specific presence in the plant genome is referred to as a transformation or transgenic event.

Plants and plant cultivars which are preferably treated according to the invention include all plants
5 which have genetic material which imparts particularly advantageous, useful traits to these plants (whether obtained by breeding and/or biotechnological means).

Seed Treatment

The treatment of the seed of plants has been known for a long time and is the subject of constant improvements. Nevertheless, the treatment of seed gives rise to a series of problems which cannot
10 always be solved in a satisfactory manner. For instance, it is desirable to develop methods for protecting the seed, the germinating plant and the resulting plants or plant parts, which dispense with, or at least significantly reduce, the additional deployment of crop protection products after planting or after emergence of the plants. It is additionally desirable to optimize the amount of active ingredients or mixtures thereof used in such a way as to provide the best possible protection for the seed and the
15 germinating plant in order to achieve the yield increase, but without damaging the Brassicaceae plant itself by the active ingredient used.

The present invention therefore relates more particularly also to a method for treating seed to increase yield in Brassicaceae plants which grow from the seed or seedlings, by treating the Brassicaceae seed with active ingredients or mixtures thereof.

20 In another embodiment a method for treating seed to increase yield in Brassicaceae plants at BBCH stage 10 or later which grow from the seed or seedlings, by treating the Brassicaceae seed at BBCH stage 00 with active ingredients or mixtures thereof.

In another embodiment a method for treating seed to increase yield in Brassica napus plants at BBCH stage 10 or later which grow from the seed or seedlings, by treating the Brassica napus seed at BBCH
25 stage 00 with active ingredients or mixtures thereof.

In another embodiment a method for treating seed to increase yield in canola plants at BBCH stage 10 or later which grow from the seed or seedlings, by treating the canola seed at BBCH stage 00 with active ingredients or mixtures thereof.

30 In another embodiment a method for treating seed to increase yield in canola hybrid plants at BBCH stage 10 or later which grow from the seed or seedlings, by treating the canola hybrid seed at BBCH stage 00 with active ingredients or mixtures thereof.

In another embodiment a method for treating seed to increase yield in herbicide tolerant canola hybrid plants at BBCH stage 10 or later which grow from the seed or seedlings, by treating the herbicide tolerant canola hybrid seed at BBCH stage 00 with active ingredients or mixtures thereof.

The invention likewise relates to the use of active ingredients or mixtures for treatment of seed to increase yield in the the plants or those plant parts which grow from the seeds.

It is likewise considered to be advantageous that active ingredients or mixtures thereof, can especially also be used in transgenic seed.

5 The active ingredients or mixtures thereof are applied to the seed alone or in a suitable formulation. Preferably, the seed is treated in a state in which it is stable enough to avoid damage during treatment. In general, the seed may be treated at any time between harvest and sowing. The seed typically used has been separated from the plant and freed from cobs, shells, stalks, coats, hairs or the fruit flesh. For example, it is possible to use seed which has been harvested, cleaned and dried to a moisture content of
10 less than 15% by weight. Alternatively, it is also possible to use seed which, after drying, for example, has been treated with water and then dried again.

When treating the seed, it must generally be ensured that the amount of active ingredients or mixtures thereof applied to the seed and/or of further additives is selected such that the germination of the seed is not impaired, and that the resulting plant is not damaged. This should be noted in particular in the case
15 of active ingredients which can have phytotoxic effects at particular application rates.

The active ingredients or mixtures thereof can be applied directly, i.e. without containing any further components and without having been diluted. In general, it is preferable to apply active ingredients or mixtures thereof, to the seed in the form of a suitable formulation. Suitable formulations and methods for seed treatment are known to those skilled in the art and are described, for example, in the following
20 documents: US 4,272,417 A, US 4,245,432 A, US 4,808,430 A, US 5,876,739 A, US 2003/0176428 A1, WO 2002/080675 A1, WO 2002/028186 A2.

The active ingredients or mixtures thereof can be converted to the customary seed dressing formulations, such as solutions, emulsions, suspensions, powders, foams, slurries or other coating materials for seed, and also ULV formulations.

25 These formulations are produced in a known manner, by mixing the active ingredients or active ingredient combinations with customary additives, for example customary extenders and solvents or diluents, dyes, wetting agents, dispersants, emulsifiers, defoamers, preservatives, secondary thickeners, stickers, gibberellins and also water.

Useful dyes which may be present in the seed dressing formulations usable in accordance with the
30 invention are all dyes customary for such purposes. It is possible to use both sparingly water-soluble pigments and water-soluble dyes. Examples include the dyes known under the Rhodamine B, C.I. Pigment Red 112 and C.I. Solvent Red 1 names.

The wetting agents which may be present in the seed dressing formulations usable in accordance with the invention include all substances which promote wetting and are customary for formulation of active
35 agrochemical ingredients. Usable with preference are alkyl naphthalenesulphonates, such as diisopropyl or diisobutyl naphthalenesulphonate.

The dispersants and/or emulsifiers which may be present in the seed dressing formulations usable in accordance with the invention include all nonionic, anionic and cationic disperants which are customary for formulation of active agrochemical ingredients. Usable with preference are nonionic or anionic dispersants or mixtures of nonionic or anionic dispersants. Suitable nonionic dispersants include especially ethylene oxide-propylene oxide block polymers, alkylphenol polyglycol ethers and tristyrylphenol polyglycol ethers, and the phosphated or sulphated derivatives thereof. Suitable anionic dispersants are especially lignosulphonates, polyacrylic acid salts and arylsulphonate-formaldehyde condensates.

The defoamers which may be present in the seed dressing formulations usable in accordance with the invention include all foam-inhibiting substances customary for formulation of active agrochemical ingredients. Usable with preference are silicone defoamers and magnesium stearate.

The preservatives which may be present in the seed dressing formulations usable in accordance with the invention include all substances usable for such purposes in agrochemical formulations. Examples include dichlorophene and benzyl alcohol hemiformal.

Useful secondary thickeners which may be present in the seed dressing formulations usable in accordance with the invention include all substances usable for such purposes in agrochemical formulations. Preferred examples include cellulose derivatives, acrylic acid derivatives, xanthan, modified clays and finely divided silica.

Useful stickers which may be present in the seed dressing formulations usable in accordance with the invention are all customary binders usable in seed dressing compositions. Preferred examples include polyvinylpyrrolidone, polyvinyl acetate, polyvinyl alcohol and tylose.

The gibberellins which may be present in the seed dressing formulations usable in accordance with the invention are preferably gibberellins A1, A3 (= gibberellic acid), A4 and A7, particular preference being given to using gibberellic acid. The gibberellins are known (cf. R. Wegler "Chemie der Pflanzenschutz- und Schädlingsbekämpfungsmittel" [Chemistry of Crop Protection and Pest Control Compositions], vol. 2, Springer Verlag, 1970, p. 401-412).

The seed dressing formulations usable in accordance with the invention can be used to treat either directly or after preceding dilution with water. The seed dressing preparations usable in accordance with the invention or the dilute preparations thereof can also be used to dress seed of transgenic plants. In this case, it is also possible for additional synergistic effects to occur in interaction with substances formed by expression.

For treatment of seed with the seed dressing formulations usable in accordance with the invention, or the preparations prepared therefrom by adding water, all mixing units usable customarily for the seed dressing are useful. Specifically, the seed dressing procedure is to introduce the seed into a mixer, to add the particular desired amount of seed dressing formulations, either as such or after preceding dilution

with water, and to mix until the formulation is distributed homogeneously on the seed. This may be followed by a drying operation.

The application rate of seed dressing formulations usable in accordance with the invention may vary within a relatively wide range. It is guided by the particular content of the active ingredients in the formulations and by the seed. The application rates of seed treatment compositions comprising active ingredients or mixtures thereof are: from 0.1 to 200 g per 100 kg of seed, preferably from 1 to 150 g per 100 kg of seed, more preferably from 2.5 to 25 g per 100 kg of seed, even more preferably from 2.5 to 12.5 g per 100 kg of seed;

The active ingredients or mixtures thereof, may be present in their commercially available formulations and in the use forms, prepared from these formulations, as a mixture with other active ingredients, such as insecticides, attractants, sterilants, bactericides, acaricides, nematocides, fungicides, growth regulators, herbicides, safeners, fertilizers or semiochemicals.

In another embodiment the active ingredients may be present in their commercially available formulations and in the use forms, prepared from these formulations, as a mixture with one or more active ingredients.

LIST OF ABBREVIATIONS

2mepsps	Glyphosate oxidase from <i>Ochrobactrum anthropi</i> strain LBAA	decreases binding affinity for glyphosate, thereby increasing tolerance to glyphosate herbicide
accdPc	l-amino-cyclopropane-1-carboxylic acid deaminase enzyme from <i>Pseudomonas chloroaphis</i>	metabolizes the precursor of the fruit ripening hormone ethylene, resulting in delayed fruit ripening
antisense Pg	no functional polygalacturonase enzyme is produced (transcription of the endogenous enzyme is suppressed by a gene silencing mechanism)	inhibits the production of polygalacturonase enzyme responsible for the breakdown of pectin molecules in the cell wall, and thus causes delayed softening of the fruit
athb17	a protein from <i>Arabidopsis thaliana</i> of the class II family of the homeodomain-leucine zipper (HD-Zip) transcription factors.	Modulates plant growth and development and regulates gene expression
avhppd-03	p-hydroxyphenylpyruvate dioxygenase from <i>Avena sativa</i>	Tolerance to Mesotrione herbicide
bar	phosphinothricin N-acetyltransferase (PAT) enzyme from <i>Streptomyces hygroscopicus</i>	eliminates herbicidal activity of glufosinate (phosphinothricin) herbicides by acetylation
barnase	barnase ribonuclease enzyme from <i>Bacillus amyloliquefaciens</i>	causes male sterility by interfering with RNA production in the tapetum cells of the anther
barstar	barnase ribonuclease inhibitor from <i>Bacillus amyloliquefaciens</i>	restores fertility by repressing the inhibitory effect of barnase on tapetum cells of the anther
bbx32	protein from <i>Arabidopsis thaliana</i> that interacts with one or more endogenous transcription factors to regulate the plant's day/night physiological processes	to modulate plant's diurnal biology and to enhance growth and reproductive development
bxn	nitrilase enzyme from <i>Klebsiella pneumoniae subsp. Ozaenae</i>	eliminates herbicidal activity of oxynil herbicides (eg. bromoxynil)
ccomt (inverted repeat)	dsRNA that suppresses endogenous S-adenosyl-L-methionine: trans-caffeoyl CoA 3-O-methyltransferase (CCOMT gene) RNA transcript levels via the RNA interference (RNAi) pathway	reduces content of guaiacyl (G) lignin
CordapA	dihydrodipicolinate synthase enzyme from <i>Corynebacterium glutamicum</i>	increases the production of amino acid lysine
cp4 epsps	herbicide tolerant form of 5-enolpyruvulshikimate-3-phosphate synthase (EPSPS) enzyme from <i>Agrobacterium tumefaciens</i> strain CP4	decreases binding affinity for glyphosate, thereby conferring increased tolerance to glyphosate herbicide

cry1A.105	Cry1A.105 protein which comprises the Cry1Ab, Cry1F and Cry1Ac proteins from <i>Bacillus thuringiensis</i> subsp. <i>Kumamotoensis</i>	confers resistance to lepidopteran insects by selectively damaging their midgut lining
cry1Ab	Cry1Ab delta-endotoxin from <i>Bacillus thuringiensis</i> subsp. <i>Kurstaki</i>	confers resistance to lepidopteran insects by selectively damaging their midgut lining
cry1Ac	Cry1Ac delta-endotoxin from <i>Bacillus thuringiensis</i> subsp. <i>Kurstaki</i> strain <i>HD73</i>	confers resistance to lepidopteran insects by selectively damaging their midgut lining
Cry1F	Cry1F delta-endotoxin from <i>Bacillus thuringiensis</i> var. <i>Aizawai</i>	confers resistance to lepidopteran insects by selectively damaging their midgut lining
cry1Fa2	synthetic form of cry1F gene derived from <i>Bacillus thuringiensis</i> var. <i>Aizawai</i>	confers resistance to lepidopteran insects by selectively damaging their midgut lining
cry2Ab2	Cry2Ab delta-endotoxin from <i>Bacillus thuringiensis</i> subsp. <i>Kumamotoensis</i>	confers resistance to lepidopteran insects by selectively damaging their midgut lining
cry2Ae	Cry2Ae delta-endotoxin from <i>Bacillus thuringiensis</i> subsp. <i>Dakota</i>	confers resistance to lepidopteran insects by selectively damaging their midgut lining
cry34Ab1	Cry34Ab1 delta-endotoxin from <i>Bacillus thuringiensis</i> strain <i>PS149B1</i>	confers resistance to coleopteran insects particularly corn rootworm by selectively damaging their midgut lining
cry35Ab1	Cry35Ab1 delta-endotoxin from <i>Bacillus thuringiensis</i> strain <i>PS149B1</i>	confers resistance to coleopteran insects particularly corn rootworm by selectively damaging their midgut lining
cry3A	cry3A delta endotoxin from <i>Bacillus thuringiensis</i> subsp. <i>Tenebrionis</i>	confers resistance to coleopteran insects by selectively damaging their midgut lining
cry3Bb1	Cry3Bb1 delta endotoxin from <i>Bacillus thuringiensis</i> subsp. <i>Kumamotoensis</i>	confers resistance to coleopteran insects particularly corn rootworm by selectively damaging their midgut lining
cspB	cold shock protein B from <i>Bacillus subtilis</i>	maintains normal cellular functions under water stress conditions by preserving RNA stability and translation
Dgat2A	diacylglycerol acyltransferase type 2A from <i>Mortierella ramanniana</i>	modifies oil profile
dmo	dicamba mono-oxygenase enzyme from <i>Stenotrophomonas maltophilia</i> strain <i>DI-6</i>	confers tolerance to the herbicide dicamba (2-methoxy-3,6-dichlorobenzoic acid) by using dicamba as substrate in an enzymatic reaction
dvsnf7 (inverted repeat)	double-stranded RNA transcript containing a 240 bp fragment of the Western Corn Rootworm <i>Snf7</i> gene	RNAi interference resulting to down-regulation of the function of the targeted <i>Snf7</i> gene leading to Western Corn Rootworm mortality.
epsps grg23 ace 5	modified 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS) protein or EPSPS ACE5 protein from <i>Arthrobacter</i>	confers tolerance to glyphosate herbicides

	<i>globiformis</i>	
fad2-1A (sense antisense)	no functional enzyme is produced (production of delta-12 desaturase enzyme is suppressed by RNA interference)	reduces desaturation of 18:1 oleic acid to 18:2 linoleic acid; increases the levels of monounsaturated oleic acid and decreases the levels of saturated linoleic acid in the seed
fatb1-A (sense antisense)	no functional enzyme is produced (production of FATB enzymes or acyl-acyl carrier protein thioesterases is suppressed by RNA interference)	decreases the transport of saturated fatty acids out of the plastid, thereby increasing their availability to desaturation to 18:1 oleic acid; reduces the levels of saturated fatty acids and increases the levels of 18:1 oleic acid
goxv247	glyphosate oxidase from <i>Ochrobactrum anthropi strain LBAA</i>	confers tolerance to glyphosate herbicides by degrading glyphosate into aminomethylphosphonic acid (AMPA) and glyoxylate
hppdPF W336	modified p-hydroxyphenylpyruvate dioxygenase (hppd) enzyme from <i>Pseudomonas fluorescens strain A32</i>	confers tolerance to HPPD-inhibiting herbicides (such as isoxaflutole) by reducing the specificity for the herbicide's bioactive constituent
Nc.Fad3	delta 15 desaturase protein from <i>Neurospora crassa</i>	desaturates certain endogenous fatty acids resulting in the production of stearidonic acid (SDA), an omega-3 fatty acid
nptII	neomycin phosphotransferase II enzyme	allows transformed plants to metabolize neomycin and kanamycin antibiotics during selection
pat	phosphinothricin N-acetyltransferase (PAT) enzyme - synthetic form of pat gene derived from <i>Streptomyces viridochromogenes strain Tu 494</i>	eliminates herbicidal activity of glufosinate (phosphinothricin) herbicides by acetylation
pinII	protease inhibitor protein from <i>Solanum tuberosum</i>	enhances defense against insect predators by reducing the digestibility and nutritional quality of the leaves
Pj.D6D	delta 6 desaturase protein from <i>Primula juliae</i>	desaturates certain endogenous fatty acids resulting in the production of stearidonic acid (SDA), an omega-3 fatty acid
PLRV orf1	putative replicase domain of the potato leaf roll virus (PLRV)	confers resistance to potato leaf roll virus (PLRV) through gene silencing mechanism
PLRV orf2	putative helicase domain of the potato leaf roll virus (PLRV)	confers resistance to potato leaf roll virus (PLRV) through gene silencing mechanism
PVY coat protein	coat protein of the potato virus Y (PVY)	confers resistance to potato virus Y (PVY) through "pathogen-derived resistance" mechanism

te	12:0 ACP thioesterase enzyme from <i>Umbellularia californica</i> (bay leaf)	increases the level of triacylglycerides containing esterified lauric acid (12:0)
uidA	beta-D-glucuronidase (GUS) enzyme from <i>Escherichia coli</i>	produces blue stain on treated transformed tissue, which allows visual selection
vip3A	VIP3A vegetative insecticidal protein from <i>Bacillus thuringiensis strain AB88</i>	confers resistance to feeding damage caused by lepidopteran insects by selectively damaging their midgut lining

Claims:

1. Method for increasing yield in *Brassicaceae* plants comprising at least one event or allele according to table A, wherein the *Brassicaceae* plants, plant parts thereof or the soil in which *Brassicaceae* plants are grown or intended to be grown are treated with the active ingredients or mixtures thereof.
- 5 2. Method according to Claim 1, wherein the active ingredients or mixtures thereof are applied as a foliar treatment to *Brassicaceae* plants, as a seed treatment to *Brassicaceae* seeds or as a soil application to the habitat where *Brassicaceae* plants are grown or will be intended to grow.
3. Method according to Claim 1 or 2 wherein the active ingredients or mixtures thereof are selected from the group H-1, H-2, H-3, H-4, H-5, P-1, SAF-1, SAF-2, SAF-3, SAF-4, B-1, B-2, B-3, B-4, B-5, I-1, I-2, I-3, I-4, I-5, I-6, F-1, F-2, F-3, F-4, F-5, F-6, SG-1, SG-2, SG-3, SG-4, SG-5, SG-6, SG-7, SG-8.
10
4. Method according to any one of Claims 1 to 3, wherein the events or alleles are selected from the group consisting of 23-18-17, MON88302, RT200, RT73, MS8, RF3, POSH122, MS11, Ogura, R113, R40, R2000.
5. Method according to any one of Claims 1 to 4, wherein the *Brassicaceae* plants are hybrid winter
15 oilseed rape or hybrid spring oilseed rape plants.
6. Method according to any one of Claims 1 to 5, wherein the *Brassicaceae* plants are hybrid Canola plants.

INTERNATIONAL SEARCH REPORT

International application No PCT/EP2017/060419

A. CLASSIFICATION OF SUBJECT MATTER INV. A01H5/10 ADD.				
According to International Patent Classification (IPC) or to both national classification and IPC				
B. FIELDS SEARCHED				
Minimum documentation searched (classification system followed by classification symbols) A01H C12N				
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched				
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) EPO-Internal, BIOSIS, EMBL, FSTA, WPI Data				
C. DOCUMENTS CONSIDERED TO BE RELEVANT				
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.		
Y	WO 2011/153186 A1 (MONSANTO TECHNOLOGY LLC [US]; BROWN ANDREW J [US]; BYRNE JAMES F [US];) 8 December 2011 (2011-12-08) cited in the application figure 2 -----	1-6		
Y	WO 2010/051926 A2 (BAYER CROPSCIENCE AG [DE]; MAUE MICHAEL [DE]; ADALT ISABELLE [DE]; GIE) 14 May 2010 (2010-05-14) page 32, line 1 - page 37, line 17 page 42, lines 11-21 -----	1-6		
Y	WO 2011/114232 A2 (BASF AGROCHEMICAL PRODUCTS BV [NL]; VANTIEGHEM HERVE [DE]; PFENNING MA) 22 September 2011 (2011-09-22) paragraph [0011] ----- -/--	1-6		
<table style="width: 100%; border: none;"> <tr> <td style="width: 50%; border: none;"><input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C.</td> <td style="width: 50%; border: none;"><input checked="" type="checkbox"/> See patent family annex.</td> </tr> </table>			<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C.	<input checked="" type="checkbox"/> See patent family annex.
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C.	<input checked="" type="checkbox"/> See patent family annex.			
* Special categories of cited documents :				
"A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier application or patent but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family			
Date of the actual completion of the international search	Date of mailing of the international search report			
19 July 2017	13/09/2017			
Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer Obel, Nicolai			

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2017/060419

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 2005/223425 A1 (CLINTON WILLIAM P [US] ET AL) 6 October 2005 (2005-10-06) paragraph [0031]; claim 28 -----	1-6
Y	US 2014/296070 A1 (FABRY CARL J [US] ET AL) 2 October 2014 (2014-10-02) paragraphs [0014] - [0015] paragraph [0025] -----	1-6

INTERNATIONAL SEARCH REPORT

International application No.
PCT/EP2017/060419

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. As all searchable claims could be searched without effort justifying an additional fees, this Authority did not invite payment of additional fees.
3. As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

1-6(partially)

Remark on Protest

- The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. claims: 1-6(partially)
Directed to EVENT1 (table A).

2. claims: 1-6(partially)
Directed to EVENT2 (table A).

3. claims: 1-6(partially)
Directed to EVENT3 (table A).

4. claims: 1-6(partially)
Directed to EVENT4 (table A).

5. claims: 1-6(partially)
Directed to EVENT5 (table A).

6. claims: 1-6(partially)
Directed to EVENT6 (table A).

7. claims: 1-6(partially)
Directed to EVENT6 (table A).

8. claims: 1-6(partially)
Directed to ALLELE1 (table A).

9. claims: 1-6(partially)
Directed to EVENT7 (table A).

10. claims: 1-6(partially)
Directed to ALLELE2 (table A).

11. claims: 1-6(partially)

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Directed to ALLELE3 (table A).

12. claims: 1-6(partially)

Directed to ALLELE4 (table A).

13. claims: 1-6(partially)

Directed to ALLELE5 (table A).

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No PCT/EP2017/060419

Patent document cited in search report	Publication date	Patent family member(s)	Publication date	
WO 2011153186	A1	08-12-2011	AU 2011261580 A1	29-11-2012
			CA 2742005 A1	04-12-2011
			CA 2935943 A1	04-12-2011
			CL 2012003412 A1	22-03-2013
			CN 103096712 A	08-05-2013
			EA 201291344 A1	29-03-2013
			EP 2575431 A1	10-04-2013
			JP 5957447 B2	27-07-2016
			JP 2013532963 A	22-08-2013
			KR 20130086599 A	02-08-2013
			NZ 603506 A	29-11-2013
			US 2011302667 A1	08-12-2011
			US 2014373190 A1	18-12-2014
			US 2015119248 A1	30-04-2015
			WO 2011153186 A1	08-12-2011

WO 2010051926	A2	14-05-2010	AR 074023 A1	15-12-2010
			AU 2009313134 A1	14-05-2010
			CA 2742435 A1	14-05-2010
			CN 102317263 A	11-01-2012
			CR 20110233 A	12-09-2011
			DO P2011000123 A	15-10-2011
			EC SP11011024 A	30-06-2011
			EP 2184273 A1	12-05-2010
			EP 2352723 A2	10-08-2011
			ES 2616230 T3	12-06-2017
			JP 5911725 B2	27-04-2016
			JP 2012507482 A	29-03-2012
			KR 20110082181 A	18-07-2011
			NZ 592596 A	31-05-2013
			RU 2011122534 A	20-12-2012
			SV 2011003894 A	29-08-2011
			TW 201029573 A	16-08-2010
			US 2011301181 A1	08-12-2011
			US 2015099766 A1	09-04-2015
			UY 32223 A	31-05-2010
WO 2010051926 A2	14-05-2010			
ZA 201103225 B	25-07-2012			

WO 2011114232	A2	22-09-2011	AR 084109 A1	24-04-2013
			AU 2011228751 A1	18-10-2012
			BR 112012023280 A2	06-10-2015
			CA 2793030 A1	22-09-2011
			CL 2012002511 A1	21-12-2012
			CN 102947443 A	27-02-2013
			EA 201290921 A1	30-04-2013
			EP 2547766 A2	23-01-2013
			US 2014082772 A1	20-03-2014
			UY 33281 A	31-10-2011
			WO 2011114232 A2	22-09-2011

US 2005223425	A1	06-10-2005	AR 048411 A1	26-04-2006
			AU 2005235111 A1	03-11-2005
			BR PI0509541 A	18-09-2007
			CA 2561145 A1	03-11-2005
			EP 1729580 A2	13-12-2006
			NZ 550322 A	24-02-2012
			US 2005223425 A1	06-10-2005

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No PCT/EP2017/060419

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
		US 2016081345 A1	24-03-2016
		UY 28769 A1	30-09-2005
		WO 2005102057 A2	03-11-2005

US 2014296070 A1	02-10-2014	US 8828909 B1	09-09-2014
		US 2014296070 A1	02-10-2014
