

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
11 October 2007 (11.10.2007)

PCT

(10) International Publication Number  
WO 2007/113172 A2

(51) International Patent Classification:  
C07K 16/18 (2006.01) A61K 39/395 (2006.01)  
A61P 25/28 (2006.01)

(74) Agents: VALENTINE, Jill, Barbara et al.; Glaxo-SmithKline, Corporate Intellectual Property (CN925.1), 980 Great West Road, Brentford Middlesex TW8 9GS (GB).

(21) International Application Number:  
PCT/EP2007/052928

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(22) International Filing Date: 27 March 2007 (27.03.2007)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/787,588 30 March 2006 (30.03.2006) US

(71) Applicant (for all designated States except US): GLAXO GROUP LIMITED [GB/GB]; Glaxo Wellcome House, Berkeley Avenue, Greenford Middlesex UB6 0NN (GB).

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (for US only): BURBIDGE, Stephen, Anthony [GB/GB]; GlaxoSmithKline, New Frontiers Science Park South, Third Avenue, Harlow Essex CM19 5AW (GB). ELLIS, Jonathan, Henry [GB/GB]; GlaxoSmithKline, Gunnels Wood Road, Stevenage Hertfordshire SG1 2NY (GB). FORD, Susannah, K. [GB/GB]; GlaxoSmithKline, Gunnels Wood Road, Stevenage Hertfordshire SG1 2NY (GB). GERMASCHEWSKI, Volker [DE/GB]; GlaxoSmithKline, Gunnels Wood Road, Stevenage Hertfordshire SG1 2NY (GB). KUMAR, Umesh [GB/GB]; GlaxoSmithKline, New Frontiers Science Park South, Third Avenue, Harlow Essex CM19 5AW (GB). PHILPOTT, Karen, Louise [GB/GB]; GlaxoSmithKline, New Frontiers Science Park South, Third Avenue, Harlow Essex CM19 5AW (GB). SODEN, Peter, Ernest [GB/GB]; GlaxoSmithKline, New Frontiers Science Park South, Third Avenue, Harlow Essex CM19 5AW (GB).

**Declarations under Rule 4.17:**

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))
- of inventorship (Rule 4.17(iv))

**Published:**

- without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: ANTIBODIES

(57) Abstract: Antibodies that bind human  $\beta$ -amyloid peptide, methods of treating diseases or disorders characterised by elevated  $\beta$ -amyloid levels or  $\beta$ -amyloid deposits with said antibodies, pharmaceutical compositions comprising said antibodies and methods of manufacture.



WO 2007/113172 A2

## Antibodies

### Field of the invention

The present invention relates to antibodies that bind  $\beta$ -amyloid peptide and in particular human  $\beta$ -amyloid peptide. The present invention also concerns methods of  
5 treating diseases or disorders characterised by elevated  $\beta$ - amyloid levels or  $\beta$ - amyloid deposits, particularly Alzheimer's disease, with said antibodies, pharmaceutical compositions comprising said antibodies and methods of manufacture. Other aspects of the present invention will be apparent from the description below.

10

### Background of the invention

Alzheimer's disease (AD) is the most common cause of age-related cognitive decline, affecting greater than 12 million individuals worldwide (Citron M (2002) Nat. Neurosci 5, Suppl 1055-1057). The earliest stages of the disease are characterized  
15 by a progressive loss of memory with associated cognitive decline and language and behavioural deficits. In the later stages of the disease, patients develop global amnesia and have greatly reduced motor function. Death typically occurs 9 years following diagnosis and is often associated with other conditions, typically pneumonia (Davis K.L. and Samules S.C. (1998) in Pharmacological Management of  
20 Neurological and Psychiatric Disorders eds Enna S.J. and Coyle J.T. (McGraw-Hill, New York pp267-316)). Current therapies represent symptomatic approaches, focussing on alleviating the cognitive impairment and ameliorating the behavioural symptoms associated with the progressing disease aetiology. In practice these treatments provide only a short lived cognitive benefit with the level of cognitive  
25 impairment reported only to last up to 2 years. The potential for a disease-modifying therapy that slows and possibly halts the progression of the disease is enormous. Such approaches would provide radical and sustained improvements to the quality of life of patients and importantly their carers as well as reducing the huge overall healthcare costs of this disease.

30

Clinical diagnosis of Alzheimer's disease is based upon a combination of physical and mental tests which lead to a diagnosis of possible or probable Alzheimer's disease. At post mortem the disease is confirmed by well characterised neurological hallmarks in the brain, which include the deposition of  $A\beta$  in parenchymal plaques  
35 and cerebral vessels, intraneuronal formation of neurofibrillary tangles, synaptic loss and loss of neuronal subpopulations in specific brain regions (Terry, RD (1991) J Neural Trans Suppl 53 : 141-145).

40

A plethora of genetic, histological and functional evidence suggests that the  $\beta$ - amyloid peptide ( $A\beta$ ) is key to the progression of Alzheimer's disease (Selkoe, D. J. (2001) Physiological Reviews **81**: 741-766).

$A\beta$  is known to be produced through the cleavage of the beta amyloid precursor protein (also known as APP) by an aspartyl protease enzyme known as BACE1 (also

known as  $\beta$ -secretase, Asp2 or Memapsin-2) (De Strooper, B. and Konig, G. (1999) Nature **402**: 471-472). In addition to the parenchymal and vascular deposition, soluble oligomeric forms of  $A\beta$  have been postulated to contribute to the onset of AD and they may affect neuronal function initially by impairing synaptic function (Lambert et. al. (1998) Proceedings of the National Academy of Science, U.S.A. **95** : 6448-6453). Although insoluble amyloid plaques are found early in AD and in MCI, the levels of soluble  $A\beta$  aggregates (referred to as oligomers or  $A\beta$ -derived diffusible ligands (ADDLs) are also increased in these individuals, and soluble  $A\beta$  levels correlate better with neurofibrillary degeneration, and the loss of synaptic markers than do amyloid plaques (Naslund et. al. (2000) J Am Med Assoc **283**: 1571-1577, Younkin, S. (2001) Nat. Med. **1**: 8-19). The highly amyloidogenic  $A\beta_{42}$  and aminoterminally truncated forms  $A\beta_{x-42}$  are the predominant species of  $A\beta$  found in both diffuse and senile plaques (Iwatsubo, T (1994) *Neuron*. **13**:45-53, Gravina, SA (1995) *J. Biol. Chem.* **270**:7013-7016) The relative levels of  $A\beta_{42}$  appear to be the key regulator of  $A\beta$  aggregation into amyloid plaques, indeed  $A\beta_{42}$  has been shown to aggregate more readily than other  $A\beta$  forms in vitro (Jarrett, JT (1993) *Biochemistry*.**32**:4693-4697) and as such  $A\beta_{42}$  has been implicated as the initiating molecule in the pathogenesis of AD (Younkin SG, (1998) *J. Physiol. (Paris)*. **92**:289-292). Although  $A\beta_{42}$  is a minor product of APP metabolism, small shifts in its production are associated with large effects on  $A\beta$  deposition therefore it has been postulated that reduction of  $A\beta_{42}$  alone may be an effective way of treating AD (Younkin SG, (1998) *J. Physiol. (Paris)*. **92**:289-292) In support of this, mutations in the amyloid precursor protein (APP) and presenilin genes have been reported to predominantly increase the relative levels of  $A\beta_{42}$  and therefore shortening the time to onset of Alzheimer's disease (AD) (Selkoe D.J., Podlisny M.B. (2002) *Annu. Rev. Genomics Hum. Genet.* **3**:67-99). It should be noted however, that the rate of deposition is also dependant on catabolism and  $A\beta$  clearance.

Animal models of amyloid deposition have been generated by overexpressing mutant human transgenes in mice. Mice overexpressing single human APP transgenes typically develop cerebral plaque-like  $\beta$ -amyloid deposits from 12 months of age (Games D. et al., (1995) *Nature* **373**: 523-527; Hsiao K. et al., (1996) *Science* **274**: 99-102), while mice carrying both mutant human APP and presenilin-1 (PS-1) transgenes typically develop cerebral plaque-like  $\beta$ -amyloid deposits as early as 2 months of age (Kurt M.A. et al., (2001) *Exp.Neurol.* **171**: 59-71; McGowan E. et al., (1999) *Neurobiol. Dis.* **6**: 231-244.

It has become increasingly apparent that the transport of exogenous  $A\beta$  between the central nervous system (CNS) and plasma plays a role in the regulation of brain amyloid levels (Shibata, et al (2000) *J Clin Invest* **106** : 1489-1499), with CSF  $A\beta$  being rapidly transported from CSF to plasma. Therefore active vaccination with  $A\beta$  peptides or passive administration of specific  $A\beta$  antibodies rapidly binds peripheral  $A\beta$  altering the dynamic equilibrium between the plasma, CSF and ultimately the CNS. Indeed there are now a plethora of studies demonstrated both these

approaches can lower A $\beta$  levels, reduce A $\beta$  pathology and provide cognitive benefit in various transgenic models of amyloidosis. Limited studies have also been conducted in higher species. Caribbean vervet monkeys (16-10 years old) were immunised with A $\beta$  peptide over 10 months. A $\beta$ 40 levels were elevated 2-5 fold in the plasma which peaked at 251d while the CSF levels of A $\beta$ 40 and A $\beta$ 42 were significantly decreased by 100d and returned towards baseline thereafter. This reduction in CSF was accompanied by a significant reduction in plaque burden (Lemere, CA (2004) Am J Pathology 165 : 283-297). Similar increases in plasma A $\beta$  levels were also detected following immunisation of aged (15-20 year old) Rhesus Monkeys (Gandy, S (2004) Alzheimer Dis Assoc Disord 18 : 44:46).

The first immune therapy targeting brain amyloid was Elan/Wyeth's AN-1792, an active vaccine. This treatment was terminated following the development of clinical signs consistent with meningoencephalitis. Subgroup analyses suggested that treatment slowed the decline of cognitive function (Nature Clin Pract Neurol (2005) 1:84-85). Post-mortem analysis of patients also showed evidence of plaque-clearance (Gilman S. et al, (2005) Neurology 64 (9) 1553-1562). Bapineuzumab (AAB-001, Elan/Wyeth), a passive MAb therapy has been shown to significantly improve cognition scores in a small phase I safety study.

Other diseases or disorders characterised by elevated  $\beta$ - amyloid levels or  $\beta$ -amyloid deposits include mild cognitive impairment (MCI, Blasko I (2006) Neurobiology of aging "Conversion from cognitive health to mild cognitive impairment and Alzheimer's disease: Prediction by plasma amyloid beta 42, medial temporal lobe atrophy and homocysteine" in press, e-published 19 Oct 2006), hereditary cerebral haemorrhage with  $\beta$ -amyloidosis of the Dutch type, cerebral  $\beta$ -amyloid angiopathy and various types of degenerative dementias, such as those associated with Parkinson's disease, progressive supranuclear palsy, cortical basal degeneration and diffuse Lewis body type of Alzheimer's disease (Mollenhauer B (2007) J Neural Transm e-published 23 Feb 2007, van Oijen, M Lancet Neurol. 2006 5:655-60) and Down syndrome (Mehta, PD (2007) J Neurol Sci. 254:22-7).

### **Summary of the invention**

In an embodiment of the present invention there is provided a therapeutic antibody which is an antibody or antigen binding fragment and/or derivative thereof which binds  $\beta$ -amyloid peptide 1-12 (SEQ ID No:15) with equilibrium constant KD less than 100pM but does not bind to  $\beta$ -amyloid peptide 2-13 (SEQ ID No:44), both determinations being made in a surface plasmon resonance assay utilising peptide captured on streptavidin chip .

In another embodiment of the present invention there is provided a therapeutic antibody which is an antibody or antigen binding fragment and/or derivative thereof which binds  $\beta$ -amyloid peptide 1-12 (SEQ ID No:15) with equilibrium constant KD less than 100pM and has an equilibrium constant KD for binding to  $\beta$ -amyloid peptide

2-13 (SEQ ID No:44) which is 1000-fold greater than that for peptide 1-12 (SEQ ID No:15), both determinations being made in a surface plasmon resonance assay utilising peptide captured on streptavidin chip.

- 5 In another embodiment of the present invention there is provided a therapeutic antibody which is an antibody or antigen binding fragment and/or derivative thereof which binds  $\beta$ -amyloid peptide 1-12 (SEQ ID No:15) with equilibrium constant KD less than 100pM and has an equilibrium constant KD for binding to  $\beta$ -amyloid peptide 2-13 (SEQ ID No:44) which is 10,000-fold greater than that for peptide 1-12 (SEQ ID  
10 No:15), both determinations being made in a surface plasmon resonance assay utilising peptide captured on streptavidin chip.

In one aspect the surface plasmon resonance assay utilising peptide captured on streptavidin chip is the Surface Plasmon Resonance assay described in the Example  
15 below. In another aspect the surface plasmon resonance assay utilising peptide captured on streptavidin chip is the Method A(i) described under SPR Biacore™ Analysis below.

In an alternative embodiment of the present invention there is provided a therapeutic  
20 antibody which is an antibody or antigen binding fragment and/or derivative thereof which binds  $\beta$ -amyloid peptide 1-40 with equilibrium constant KD less than 10nM but does not bind to  $\beta$ -amyloid peptide 2-13 (SEQ ID No:44), both determinations being made in the surface plasmon resonance assay described in Method B of the Examples below.

25 In another alternative embodiment of the present invention there is provided a therapeutic antibody which is an antibody or antigen binding fragment and/or derivative thereof which binds  $\beta$ -amyloid peptide 1-40 with equilibrium constant KD less than 10nM and has an equilibrium constant KD for binding to  $\beta$ -amyloid peptide  
30 2-13 (SEQ ID No:44) which is 1000-fold greater than that for peptide 1-12 (SEQ ID No:15), both determinations being made in the surface plasmon resonance assay described in Method B of the Examples below.

In another alternative embodiment of the present invention there is provided a  
35 therapeutic antibody which is an antibody or antigen binding fragment and/or derivative thereof which binds  $\beta$ -amyloid peptide 1-40 with equilibrium constant KD less than 10nM and has an equilibrium constant KD for binding to  $\beta$ -amyloid peptide 2-13 (SEQ ID No:44) which is 10,000-fold greater than that for peptide 1-12 (SEQ ID No:15), both determinations being made in the surface plasmon resonance assay  
40 described in Method B of the Examples below.

In an embodiment of the present invention there is provided a therapeutic antibody which is an antibody or antigen binding fragment and/or derivative thereof which binds  $\beta$ -amyloid peptide and which comprises the following CDRs:

CDRH1: DNGMA (SEQ ID No:1)  
 CDRH2: FISNLAYSIDYADTVTG (SEQ ID No:2)  
 CDRH3: GTWFAY (SEQ ID No:3)

5 within a human heavy chain variable region originating from the VH3 gene family  
 and:

CDRL1: RVSQSLLHSNGYTYLH (SEQ ID No:4)  
 CDRL2: KVSNRFS (SEQ ID No:5)  
 CDRL3: SQTRHVPYT (SEQ ID No:6)

10 within a human light chain variable region originating from the amino acid sequence  
 disclosed in GenPept entry CAA51135 (SEQ ID No:24).

Throughout this specification, the terms "CDR", "CDRL1", "CDRL2", "CDRL3",  
 "CDRH1", "CDRH2", "CDRH3" follow the Kabat numbering system as set forth in  
 15 Kabat *et al*; *Sequences of proteins of Immunological Interest* NIH, 1987. Therefore  
 the following defines the CDRs according to the invention:

**CDR: Residues**

CDRH1: 31-35B  
 CDRH2: 50-65  
 20 CDRH3: 95-102  
 CDRL1: 24-34  
 CDRL2: 50-56  
 CDRL3: 89-97

25 The VH3 gene family and related immunoglobulin gene nomenclature is described in  
 Matsuda et al (Journal of Experimental Medicine, 188:2151-2162, 1998) and Lefranc  
 & Lefranc (The Immunoglobulin Factsbook. 2001. Academic Press: London).

In a particular embodiment, the human heavy chain variable region originates from:

- 30
- A V gene selected from the following subset of VH3 family members: VH3-48,  
 VH3-21, VH3-11, VH3-7, VH3-13, VH3-74, VH3-64, VH3-23, VH3-38, VH3-  
 53, VH3-66, VH3-20, VH3-9 & VH3-43
  - A V gene selected from the following subset of VH3 family members: VH3-48,  
 VH3-21 & VH3-11
  - 35 • The VH3-48 gene  
 or an allele thereof.

The sequence in Genbank entry M99675 is an allele of the VH3-48 gene. M99675 is  
 a Genbank nucleotide sequence of a genomic piece of DNA including the two exons  
 40 that constitute the human heavy chain gene VH3-48 (SEQ ID No:22) and encode the  
 variable region amino acid sequence given in SEQ ID No:21. In a particular aspect  
 the human acceptor heavy chain framework is derived from M99675.

In order to construct a complete V-region a framework 4 has to be added to the germline encoded V-gene M99675. Suitable framework 4 sequences include that encoded by the human JH4 minigene (Kabat):  
YFDYWGQGTLVTVSS (SEQ ID No:23)

5 of which the initial four residues fall within the CDR3 region which is replaced by the incoming CDR from the donor antibody.

The skilled person appreciates that a germline V gene and a J gene do not include coding sequence for the entirety of heavy chain CDR3. However, in the antibodies of  
10 the invention, the CDR3 sequence is provided by the donor immunoglobulin. Accordingly, the combination of a VH gene such as VH3-48, a JH minigene such as JH4, and a set of heavy chain CDRs, such as SEQ ID No:1, SEQ ID No:2 and SEQ ID No:3 (assembled in a manner so as to mimic a mature, fully rearranged heavy chain variable region) is sufficient to define a heavy chain variable region of the  
15 invention such as that represented in SEQ ID No:26, 28, 30.

The variable region encoded by Genpept ID CAA51134 has the amino acid sequence given in SEQ ID No:24.

20 The light chain variable region framework sequence known by the GenPept ID CAA51134 is the deduced amino acid sequence of a fully rearranged light chain variable region and is identical to another amino acid sequence with the same frameworks in the database: Genpept accession number S40356, and is described in Klein, R., et al., Eur. J. Immunol. 23 (12), 3248-3262 (1993). The DNA coding  
25 sequence for CAA51134, accessible as Genbank Accesion No X72467, is given as SEQ ID No: 25.

In a particular aspect of the invention the human acceptor heavy chain framework is derived from M99675 and the JH4 minigene and the human acceptor light chain  
30 framework is derived from CAA51135, optionally containing one or more, such as one to four, more particularly one to three, substitutions of amino acid residues based on the corresponding residues found in the donor V<sub>H</sub> domain having the sequence: SEQ ID No:17 and V<sub>L</sub> domain having the sequence: SEQ ID No: 19 that maintain all or substantially all of the binding affinity of the donor antibody for  $\beta$ -amyloid peptide.  
35

By 'substantially all of the binding affinity' is meant that the therapeutic antibody has at most a ten-fold reduction in binding affinity compared to the donor antibody.

In a more particular aspect the human acceptor heavy chain framework derived from  
40 M99675 and JH4 has one to four amino acid residue substitutions selected from positions 24, 48, 93 and/or 94 (Kabat numbering).

In a more particular aspect of the invention the human acceptor heavy chain framework derived from M99675 and JH4 comprises the following residues (or a conservative substitute thereof):

(i)

5	<b>Position</b>	<b>Residue</b>
	93	V
	94	S

or

10 (ii)

<b>Position</b>	<b>Residue</b>
24	V
93	V
94	S

15

or

(iii)

20	<b>Position</b>	<b>Residue</b>
	48	I
	93	V
	94	S

In one embodiment of the invention there is provided a therapeutic antibody comprising a V<sub>H</sub> chain having the sequence set forth in SEQ ID No:26, 28 or 30 and a V<sub>L</sub> domain having the sequence set forth in SEQ ID No:32.

In another embodiment of the invention there is provided a therapeutic antibody, which antibody comprises a heavy chain having the sequence set forth in SEQ ID No:34, 36 or 38 and a light chain having the sequence set forth in SEQ ID No:40.

In another embodiment of the invention there is provided a pharmaceutical composition comprising a therapeutic antibody according to the invention.

In a further embodiment of the invention there is provided a method of treating a human patient afflicted with a  $\beta$ -amyloid peptide-related disease which method comprises the step of administering to said patient a therapeutically effective amount of a therapeutic antibody according to the invention.

Use of a therapeutic antibody according to the invention in the manufacture of a medicament for the treatment of a  $\beta$ -amyloid peptide-related disease is also provided.



In another embodiment of the invention there is provided a process for the manufacture of a therapeutic antibody according to the invention, which process comprises expressing polynucleotide encoding the antibody in a host cell.

- 5 In another embodiment of the invention there is provided a polynucleotide encoding a therapeutic antibody heavy chain comprising a  $V_H$  chain having the sequence set forth in SEQ ID No:26, 28 or 30.

- 10 In another embodiment of the invention there is provided a polynucleotide encoding a therapeutic antibody light chain comprising a  $V_L$  domain having the sequence set forth in SEQ ID No:32.

- 15 In another embodiment of the invention there is provided a polynucleotide encoding a therapeutic antibody heavy chain having the sequence set forth in SEQ ID No:34, 36 or 38.

In another embodiment of the invention there is provided a polynucleotide encoding a therapeutic antibody light chain having the sequence set forth in SEQ ID No:40.

- 20 In a more particular embodiment of the invention there is provided a polynucleotide encoding a therapeutic antibody heavy chain, which polynucleotide comprises the sequence set forth in SEQ ID No:35, 37, 39 or 42.

- 25 In another more particular embodiment of the invention there is provided a polynucleotide encoding a therapeutic antibody light chain, which polynucleotide comprises the sequence set forth in SEQ ID No:41 or 43.

- 30 In a particular embodiment the therapeutic antibody which is an antibody or fragment and/or derivative thereof essentially lacks the functions of a) activation of complement by the classical pathway; and b) mediating antibody-dependent cellular cytotoxicity.

- 35 In another embodiment of the invention there is provided an antibody or a fragment thereof comprising a  $V_H$  domain having the sequence: SEQ ID No:17 and a  $V_L$  domain having the sequence: SEQ ID No: 19.

- 40 In another embodiment of the invention there is provided a polynucleotide encoding an antibody heavy chain or a fragment thereof comprising a  $V_H$  domain having the sequence SEQ ID No:17, in particular the polynucleotide of SEQ ID No:18.

In another embodiment of the invention there is provided a polynucleotide encoding an antibody light chain or a fragment thereof comprising a  $V_L$  domain having the sequence SEQ ID No: 19, in particular the polynucleotide of SEQ ID No:20.

## **Detailed Description of the Invention**

### **1. Antibody Structures**

#### **1.1 Intact Antibodies**

5 Intact antibodies are usually heteromultimeric glycoproteins comprising at least two heavy and two light chains. Aside from IgM, intact antibodies are heterotetrameric glycoproteins of approximately 150KDa, composed of two identical light (L) chains and two identical heavy (H) chains. Typically, each light chain is linked to a heavy chain by one covalent disulfide bond while the number of disulfide linkages between  
10 the heavy chains of different immunoglobulin isotypes varies. Each heavy and light chain also has intrachain disulfide bridges. Each heavy chain has at one end a variable domain ( $V_H$ ) followed by a number of constant regions. Each light chain has a variable domain ( $V_L$ ) and a constant region at its other end; the constant region of the light chain is aligned with the first constant region of the heavy chain and the light chain variable domain is aligned with the variable domain of the heavy chain. The  
15 light chains of antibodies from most vertebrate species can be assigned to one of two types called Kappa and Lambda based on the amino acid sequence of the constant region. Depending on the amino acid sequence of the constant region of their heavy chains, human antibodies can be assigned to five different classes, IgA, IgD, IgE, IgG and IgM. IgG and IgA can be further subdivided into subclasses, IgG1, IgG2, IgG3 and IgG4; and IgA1 and IgA2. Species variants exist with mouse and rat having at least IgG2a, IgG2b. The variable domain of the antibody confers binding specificity upon the antibody with certain regions displaying particular variability called  
20 complementarity determining regions (CDRs). The more conserved portions of the variable region are called framework regions (FR). The variable domains of intact heavy and light chains each comprise four FR connected by three CDRs. The CDRs in each chain are held together in close proximity by the FR regions and with the CDRs from the other chain contribute to the formation of the antigen binding site of antibodies. The constant regions are not directly involved in the binding of the  
30 antibody to the antigen but exhibit various effector functions such as participation in antibody dependent cell-mediated cytotoxicity (ADCC), phagocytosis via binding to Fc $\gamma$  receptor, half-life/clearance rate via neonatal Fc receptor (FcRn) and complement dependent cytotoxicity via the C1q component of the complement cascade. The human IgG2 constant region lacks the ability to activate complement by the classical pathway or to mediate antibody-dependent cellular cytotoxicity. The  
35 IgG4 constant region lacks the ability to activate complement by the classical pathway and mediates antibody-dependent cellular cytotoxicity only weakly. Antibodies essentially lacking these effector functions may be termed 'non-lytic' antibodies.

40

#### **1.1.1 Human antibodies**

Human antibodies may be produced by a number of methods known to those of skill in the art. Human antibodies can be made by the hybridoma method using human myeloma or mouse-human heteromyeloma cells lines see Kozbor J.Immunol 133,

3001, (1984) and Brodeur, *Monoclonal Antibody Production Techniques and Applications*, pp51-63 (Marcel Dekker Inc, 1987). Alternative methods include the use of phage libraries or transgenic mice both of which utilize human V region repertoires (see Winter G, (1994), *Annu.Rev.Immunol* 12,433-455, Green LL (1999),  
5 J.Immunol.methods 231, 11-23).

Several strains of transgenic mice are now available wherein their mouse immunoglobulin loci has been replaced with human immunoglobulin gene segments (see Tomizuka K, (2000) *PNAS* 97,722-727; Fishwild D.M (1996) *Nature Biotechnol.*  
10 14,845-851, Mendez MJ, 1997, *Nature Genetics*, 15,146-156). Upon antigen challenge such mice are capable of producing a repertoire of human antibodies from which antibodies of interest can be selected.

Of particular note is the Trimer<sup>TM</sup> system (see Eren R *et al*, (1998) *Immunology* 93:154-161) where human lymphocytes are transplanted into irradiated mice, the Selected Lymphocyte Antibody System (SLAM, see Babcock *et al*, *PNAS* (1996) 93:7843-7848) where human (or other species) lymphocytes are effectively put through a massive pooled *in vitro* antibody generation procedure followed by deconvoluted, limiting dilution and selection procedure and the Xenomouse II<sup>TM</sup>  
15 (Abgenix Inc). An alternative approach is available from Morphotek Inc using the Morphodoma<sup>TM</sup> technology.

Phage display technology can be used to produce human antibodies (and fragments thereof), see McCafferty; *Nature*, 348, 552-553 (1990) and Griffiths AD *et al* (1994)  
25 *EMBO* 13:3245-3260. According to this technique antibody V domain genes are cloned in frame into either a major or minor coat of protein gene of a filamentous bacteriophage such as M13 or fd and displayed (usually with the aid of a helper phage) as functional antibody fragments on the surface of the phage particle. Selections based on the functional properties of the antibody result in selection of the  
30 gene encoding the antibody exhibiting those properties. The phage display technique can be used to select antigen specific antibodies from libraries made from human B cells taken from individuals afflicted with a disease or disorder described above or alternatively from unimmunized human donors (see Marks; *J.Mol.Bio.* 222,581-597, 1991). Where an intact human antibody is desired comprising a Fc  
35 domain it is necessary to reclone the phage displayed derived fragment into a mammalian expression vectors comprising the desired constant regions and establishing stable expressing cell lines.

The technique of affinity maturation (Marks; *Bio/technol* 10,779-783 (1992)) may be  
40 used to improve binding affinity wherein the affinity of the primary human antibody is improved by sequentially replacing the H and L chain V regions with naturally occurring variants and selecting on the basis of improved binding affinities. Variants of this technique such as "epitope imprinting" are now also available see WO 93/06213. See also Waterhouse; *Nucl.Acids Res* 21, 2265-2266 (1993).

## **1.2 Chimaeric and Humanised Antibodies**

The use of intact non-human antibodies in the treatment of human diseases or disorders carries with it the now well established problems of potential immunogenicity especially upon repeated administration of the antibody that is the immune system of the patient may recognise the non-human intact antibody as non-self and mount a neutralising response. In addition to developing fully human antibodies (see above) various techniques have been developed over the years to overcome these problems and generally involve reducing the composition of non-human amino acid sequences in the intact therapeutic antibody whilst retaining the relative ease in obtaining non-human antibodies from an immunised animal e.g. mouse, rat or rabbit. Broadly two approaches have been used to achieve this. The first are chimaeric antibodies, which generally comprise a non-human (e.g. rodent such as mouse) variable domain fused to a human constant region. Because the antigen-binding site of an antibody is localised within the variable regions the chimaeric antibody retains its binding affinity for the antigen but acquires the effector functions of the human constant region and are therefore able to perform effector functions such as described *supra*. Chimaeric antibodies are typically produced using recombinant DNA methods. DNA encoding the antibodies (e.g. cDNA) is isolated and sequenced using conventional procedures (e.g. by using oligonucleotide probes that are capable of binding specifically to genes encoding the H and L chain variable regions of the antibody of the invention, e.g. DNA of SEQ.I.D.NO 18 and 20 described *supra*). Hybridoma cells serve as a typical source of such DNA. Once isolated, the DNA is placed into expression vectors which are then transfected into host cells such as *E.Coli*, COS cells, CHO cells, PerC6 cells or myeloma cells that do not otherwise produce immunoglobulin protein to obtain synthesis of the antibody. The DNA may be modified by substituting the coding sequence for human L and H chains for the corresponding non-human (e.g. murine) H and L constant regions see e.g. Morrison; PNAS 81, 6851 (1984). Thus another embodiment of the invention there is provided a chimaeric antibody comprising a  $V_H$  domain having the sequence: SEQ ID No:17 and a  $V_L$  domain having the sequence: SEQ ID No: 19 fused to a human constant region (which maybe of a IgG isotype e.g. IgG1) .

The second approach involves the generation of humanised antibodies wherein the non-human content of the antibody is reduced by humanizing the variable regions. Two techniques for humanisation have gained popularity. The first is humanisation by CDR grafting. CDRs build loops close to the antibody's N-terminus where they form a surface mounted in a scaffold provided by the framework regions. Antigen-binding specificity of the antibody is mainly defined by the topography and by the chemical characteristics of its CDR surface. These features are in turn determined by the conformation of the individual CDRs, by the relative disposition of the CDRs, and by the nature and disposition of the side chains of the residues comprising the CDRs. A large decrease in immunogenicity can be achieved by grafting only the CDRs of a non-human (e.g. murine) antibodies ("donor" antibodies) onto a suitable

human framework ("acceptor framework") and constant regions (see Jones *et al* (1986) Nature 321,522-525 and Verhoeyen M *et al* (1988) Science 239, 1534-1536). However, CDR grafting *per se* may not result in the complete retention of antigen-binding properties and it is frequently found that some framework residues of the donor antibody need to be preserved (sometimes referred to as "backmutations") in the humanised molecule if significant antigen-binding affinity is to be recovered (see Queen C *et al* (1989) PNAS 86, 10,029-10,033, Co, M *et al* (1991) Nature 351, 501-502). In this case, human V regions showing the greatest sequence homology (typically 60% or greater) to the non-human donor antibody maybe chosen from a database in order to provide the human framework (FR). The selection of human FRs can be made either from human consensus or individual human antibodies. Where necessary key residues from the donor antibody are substituted into the human acceptor framework to preserve CDR conformations. Computer modelling of the antibody maybe used to help identify such structurally important residues, see WO99/48523.

Alternatively, humanisation maybe achieved by a process of "veneering". A statistical analysis of unique human and murine immunoglobulin heavy and light chain variable regions revealed that the precise patterns of exposed residues are different in human and murine antibodies, and most individual surface positions have a strong preference for a small number of different residues (see Padlan E.A. *et al*; (1991) Mol.Immunol.28, 489-498 and Pedersen J.T. *et al* (1994) J.Mol.Biol. 235; 959-973). Therefore it is possible to reduce the immunogenicity of a non-human Fv by replacing exposed residues in its framework regions that differ from those usually found in human antibodies. Because protein antigenicity can be correlated with surface accessibility, replacement of the surface residues may be sufficient to render the mouse variable region "invisible" to the human immune system (see also Mark G.E. *et al* (1994) in *Handbook of Experimental Pharmacology vol.113: The pharmacology of monoclonal Antibodies*, Springer-Verlag, pp105-134). This procedure of humanisation is referred to as "veneering" because only the surface of the antibody is altered, the supporting residues remain undisturbed. Further alternative approaches include that set out in WO04/006955 and the procedure of Humaneering™ (Kalobios) which makes use of bacterial expression systems and produces antibodies that are close to human germline in sequence (Alfenito-M Advancing Protein Therapeutics January 2007, San Diego, California).

It will be apparent to those skilled in the art that the term "derived" is intended to define not only the source in the sense of it being the *physical* origin for the material but also to define material which is structually identical to the material but which does not originate from the reference source. Thus "residues found in the donor antibody" need not necessarily have been purified from the donor antibody.

It is well recognised in the art that certain amino acid substitutions are regarded as being "conservative". Amino acids are divided into groups based on common side-

chain properties and substitutions within groups that maintain all or substantially all of the binding affinity of the therapeutic antibody of the invention are regarded as conservative substitutions, see the following Table 1:

**Table 1**

<u>Side chain</u>	<u>Members</u>
Hydrophobic	met, ala, val, leu, ile
neutral hydrophilic	cys, ser, thr
Acidic	asp, glu
Basic	asn, gln, his, lys, arg
residues that influence chain orientation	gly, pro
aromatic	trp, tyr, phe

5

### **1.3 Bispecific antibodies**

A bispecific antibody is an antibody derivative having binding specificities for at least two different epitopes and also forms part of the invention. Methods of making such antibodies are known in the art. Traditionally, the recombinant production of bispecific antibodies is based on the coexpression of two immunoglobulin H chain-L chain pairs, where the two H chains have different binding specificities see Millstein *et al*, Nature 305 537-539 (1983), WO93/08829 and Traunecker *et al* EMBO, 10, 1991, 3655-3659. Because of the random assortment of H and L chains, a potential mixture of ten different antibody structures are produced of which only one has the desired binding specificity. An alternative approach involves fusing the variable domains with the desired binding specificities to heavy chain constant region comprising at least part of the hinge region, CH2 and CH3 regions. It is preferred to have the CH1 region containing the site necessary for light chain binding present in at least one of the fusions. DNA encoding these fusions, and if desired the L chain are inserted into separate expression vectors and are then cotransfected into a suitable host organism. It is possible though to insert the coding sequences for two or all three chains into one expression vector. In one preferred approach, the bispecific antibody is composed of a H chain with a first binding specificity in one arm and a H-L chain pair, providing a second binding specificity in the other arm, see WO94/04690. See also Suresh *et al* Methods in Enzymology 121, 210, 1986.

Delivery of therapeutic proteins to the brain has been hampered by the presence of the blood brain barrier (BBB). Where it is desired to deliver an antibody of the invention or antibody fragment of the invention across the BBB various strategies have been proposed to enhance such delivery where needed.

In order to obtain required nutrients and factors from the blood, the BBB possesses some specific receptors, which transport compounds from the circulating blood to the brain. Studies have indicated that some compounds like insulin (see Duffy KR *et al* (1989) Brain Res. 420:32-38), transferin (see Fishman JB *et al* (1987) J. Neurosci 18:299-304) and insulin like growth factors 1 and 2 (see Pardridge WM (1986)

Endocrine Rev.7:314-330 and Duffy KR *et al* (1986) Metabolism 37:136-140 traverse the BBB via receptor-mediated transcytosis. Receptors for these molecules thus provide a potential means for therapeutic antibodies of the invention to access the brain using so-called "vectored" antibodies (see Pardridge WM (1999) Advanced Drug Delivery Review 36:299-321). For example, an antibody to transferrin receptor has been shown to be dynamically transported into the brain parenchyma (see Friden PM *et al* (1991) PNAS 88:4771-4775 and Friden PM *et al* (1993) Science 259:373-377). Thus one potential approach is to produce a bispecific antibody or bispecific fragment such as described *supra* wherein a first specificity is towards and a second specificity towards a transport receptor located at the BBB e.g. a second specificity towards the transferrin transport receptor.

#### **1.4 Antibody Fragments**

In certain embodiments of the invention there is provided therapeutic antibody which is an antigen binding fragment. Such fragments may be functional antigen binding fragments of intact and/or humanised and/or chimaeric antibodies such as Fab, Fd, Fab', F(ab')<sub>2</sub>, Fv, ScFv fragments of the antibodies described *supra*. Fragments lacking the constant region lack the ability to activate complement by the classical pathway or to mediate antibody-dependent cellular cytotoxicity. Traditionally such fragments are produced by the proteolytic digestion of intact antibodies by e.g. papain digestion (see for example, WO 94/29348) but may be produced directly from recombinantly transformed host cells. For the production of ScFv, see Bird *et al*; (1988) Science, 242, 423-426. In addition, antibody fragments may be produced using a variety of engineering techniques as described below.

Fv fragments appear to have lower interaction energy of their two chains than Fab fragments. To stabilise the association of the V<sub>H</sub> and V<sub>L</sub> domains, they have been linked with peptides (Bird *et al*, (1988) Science 242, 423-426, Huston *et al*, PNAS, 85, 5879-5883), disulphide bridges (Glockshuber *et al*, (1990) Biochemistry, 29, 1362-1367) and "knob in hole" mutations (Zhu *et al* (1997), Protein Sci., 6, 781-788). ScFv fragments can be produced by methods well known to those skilled in the art see Whitlow *et al* (1991) Methods companion Methods Enzymol, 2, 97-105 and Huston *et al* (1993) Int.Rev.Immunol 10, 195-217. ScFv may be produced in bacterial cells such as *E.Coli* but are more typically produced in eukaryotic cells. One disadvantage of ScFv is the monovalency of the product, which precludes an increased avidity due to polyvalent binding, and their short half-life. Attempts to overcome these problems include bivalent (ScFv)<sub>2</sub> produced from ScFv containing an additional C terminal cysteine by chemical coupling (Adams *et al* (1993) Can.Res 53, 4026-4034 and McCartney *et al* (1995) Protein Eng. 8, 301-314) or by spontaneous site-specific dimerization of ScFv containing an unpaired C terminal cysteine residue (see Kipriyanov *et al* (1995) Cell. Biophys 26, 187-204). Alternatively, ScFv can be forced to form multimers by shortening the peptide linker to between 3 to 12 residues to form "diabodies", see Holliger *et al* PNAS (1993), 90, 6444-6448. Reducing the linker still further can result in ScFv trimers ("triabodies",

see Kortt *et al* (1997) Protein Eng, 10, 423-433) and tetramers ("tetrabodies", see Le Gall *et al* (1999) FEBS Lett, 453, 164-168). Construction of bivalent ScFv molecules can also be achieved by genetic fusion with protein dimerizing motifs to form "miniantibodies" (see Pack *et al* (1992) Biochemistry 31, 1579-1584) and "minibodies" (see Hu *et al* (1996), Cancer Res. 56, 3055-3061). ScFv-Sc-Fv tandems ((ScFv)<sub>2</sub>) may also be produced by linking two ScFv units by a third peptide linker, see Kurucz *et al* (1995) J.Immol.154, 4576-4582. Bispecific diabodies can be produced through the noncovalent association of two single chain fusion products consisting of V<sub>H</sub> domain from one antibody connected by a short linker to the V<sub>L</sub> domain of another antibody, see Kipriyanov *et al* (1998), Int.J.Can 77,763-772. The stability of such bispecific diabodies can be enhanced by the introduction of disulphide bridges or "knob in hole" mutations as described *supra* or by the formation of single chain diabodies (ScDb) wherein two hybrid ScFv fragments are connected through a peptide linker see Kontermann *et al* (1999) J.Immunol.Methods 226 179-188. Tetravalent bispecific molecules are available by e.g. fusing a ScFv fragment to the CH3 domain of an IgG molecule or to a Fab fragment through the hinge region see Coloma *et al* (1997) Nature Biotechnol. 15, 159-163. Alternatively, tetravalent bispecific molecules have been created by the fusion of bispecific single chain diabodies (see Alt *et al*, (1999) FEBS Lett 454, 90-94. Smaller tetravalent bispecific molecules can also be formed by the dimerization of either ScFv-ScFv tandems with a linker containing a helix-loop-helix motif (DiBi miniantibodies, see Muller *et al* (1998) FEBS Lett 432, 45-49) or a single chain molecule comprising four antibody variable domains (V<sub>H</sub> and V<sub>L</sub>) in an orientation preventing intramolecular pairing (tandem diabody, see Kipriyanov *et al*, (1999) J.Mol.Biol. 293, 41-56). Bispecific F(ab')<sub>2</sub> fragments can be created by chemical coupling of Fab' fragments or by heterodimerization through leucine zippers (see Shalaby *et al*, (1992) J.Exp.Med. 175, 217-225 and Kostelny *et al* (1992), J.Immunol. 148, 1547-1553). Also available are isolated V<sub>H</sub> and V<sub>L</sub> domains, see US 6, 248,516; US 6,291,158; US 6, 172,197.

### 30 **1.5 Heteroconjugate antibodies**

Heteroconjugate antibodies are derivatives which also form an embodiment of the present invention. Heteroconjugate antibodies are composed of two covalently joined antibodies formed using any convenient cross-linking methods. See US 4,676,980.

35

### **1.6 Other Modifications.**

The interaction between the Fc region of an antibody and various Fc receptors (Fc<sub>γ</sub>R) is believed to mediate the effector functions of the antibody which include antibody-dependent cellular cytotoxicity (ADCC), fixation of complement, phagocytosis and half-life/clearance of the antibody. Various modifications to the Fc region of antibodies of the invention may be carried out depending on the desired effector property. In particular, human constant regions which essentially lack the functions of a) activation of complement by the classical pathway; and b) mediating antibody-dependent cellular cytotoxicity include the IgG4 constant region, the IgG2



constant region and IgG1 constant regions containing specific mutations as for example mutations at positions 234, 235, 236, 237, 297, 318, 320 and/or 322 disclosed in EP0307434 (WO8807089), EP 0629 240 (WO9317105) and WO 2004/014953. Mutations at residues 235 or 237 within the CH2 domain of the heavy chain constant region (Kabat numbering; EU Index system) have separately been described to reduce binding to Fc $\gamma$ RI, Fc $\gamma$ RII and Fc $\gamma$ RIII binding and therefore reduce antibody-dependent cellular cytotoxicity (ADCC) (Duncan et al. Nature 1988, 332; 563-564; Lund et al. J. Immunol. 1991, 147; 2657-2662; Chappel et al. PNAS 1991, 88; 9036-9040; Burton and Woof, Adv. Immunol. 1992, 51;1-84; Morgan et al., Immunology 1995, 86; 319-324; Hezareh et al., J. Virol. 2001, 75 (24); 12161-12168). Further, some reports have also described involvement of some of these residues in recruiting or mediating complement dependent cytotoxicity (CDC) (Morgan et al., 1995; Xu et al., Cell. Immunol. 2000; 200:16-26; Hezareh et al., J. Virol. 2001, 75 (24); 12161-12168). Residues 235 and 237 have therefore both been mutated to alanine residues (Brett et al. Immunology 1997, 91; 346-353; Bartholomew et al. Immunology 1995, 85; 41-48; and WO9958679) to reduce both complement mediated and Fc $\gamma$ R-mediated effects. Antibodies comprising these constant regions may be termed 'non-lytic' antibodies.

One may incorporate a salvage receptor binding epitope into the antibody to increase serum half life see US 5,739,277.

There are five currently recognised human Fc $\gamma$  receptors, Fc $\gamma$ R (I), Fc $\gamma$ RIIa, Fc $\gamma$ RIIb, Fc $\gamma$ RIIIa and neonatal FcRn. Shields *et al*, (2001) J.Biol.Chem 276, 6591-6604 demonstrated that a common set of IgG1 residues is involved in binding all Fc $\gamma$ Rs, while Fc $\gamma$ RII and Fc $\gamma$ RIII utilize distinct sites outside of this common set. One group of IgG1 residues reduced binding to all Fc $\gamma$ Rs when altered to alanine: Pro-238, Asp-265, Asp-270, Asn-297 and Pro-239. All are in the IgG CH2 domain and clustered near the hinge joining CH1 and CH2. While Fc $\gamma$ RI utilizes only the common set of IgG1 residues for binding, Fc $\gamma$ RII and Fc $\gamma$ RIII interact with distinct residues in addition to the common set. Alteration of some residues reduced binding only to Fc $\gamma$ RII (e.g. Arg-292) or Fc $\gamma$ RIII (e.g. Glu-293). Some variants showed improved binding to Fc $\gamma$ RII or Fc $\gamma$ RIII but did not affect binding to the other receptor (e.g. Ser-267Ala improved binding to Fc $\gamma$ RII but binding to Fc $\gamma$ RIII was unaffected). Other variants exhibited improved binding to Fc $\gamma$ RII or Fc $\gamma$ RIII with reduction in binding to the other receptor (e.g. Ser-298Ala improved binding to Fc $\gamma$ RIII and reduced binding to Fc $\gamma$ RII). For Fc $\gamma$ RIIIa, the best binding IgG1 variants had combined alanine substitutions at Ser-298, Glu-333 and Lys-334. The neonatal FcRn receptor is believed to be involved in protecting IgG molecules from degradation and thus enhancing serum half life and the transcytosis across tissues (see Junghans R.P (1997) Immunol.Res 16. 29-57 and Ghetie *et al* (2000) Annu.Rev.Immunol. 18, 739-766). Human IgG1 residues determined to interact directly with human FcRn includes Ile253, Ser254, Lys288, Thr307, Gln311, Asn434 and His435.

The therapeutic antibody of the invention may incorporate any of the above constant region modifications.

5 In a particular embodiment, the therapeutic antibody essentially lacks the functions of a) activation of complement by the classical pathway; and b) mediating antibody-dependent cellular cytotoxicity. In a more particular embodiment the present invention provides therapeutic antibodies of the invention having any one (or more) of the residue changes detailed above to modify half-life/clearance and/or effector functions such as ADCC and/or complement dependent cytotoxicity and/or  
10 complement lysis.

In a further aspect of the present invention the therapeutic antibody has a constant region of isotype human IgG1 with alanine (or other disrupting) substitutions at positions 235 (e.g. L235A) and 237 (e.g. G237A) (numbering according to the EU  
15 scheme outlined in Kabat.

Other derivatives of the invention include glycosylation variants of the antibodies of the invention. Glycosylation of antibodies at conserved positions in their constant regions is known to have a profound effect on antibody function, particularly effector  
20 functioning such as those described above, see for example, Boyd *et al* (1996), Mol.Immunol. 32, 1311-1318. Glycosylation variants of the therapeutic antibodies of the present invention wherein one or more carbohydrate moiety is added, substituted, deleted or modified are contemplated. Introduction of an asparagine-X-serine or asparagine-X-threonine motif creates a potential site for enzymatic  
25 attachment of carbohydrate moieties and may therefore be used to manipulate the glycosylation of an antibody. In Raju *et al* (2001) Biochemistry 40, 8868-8876 the terminal sialylation of a TNFR-IgG immunoadhesin was increased through a process of regalactosylation and/or resialylation using beta-1, 4-galactosyltransferase and/or alpha, 2,3 sialyltransferase. Increasing the terminal sialylation is believed to increase  
30 the half-life of the immunoglobulin. Antibodies, in common with most glycoproteins, are typically produced in nature as a mixture of glycoforms. This mixture is particularly apparent when antibodies are produced in eukaryotic, particularly mammalian cells. A variety of methods have been developed to manufacture defined glycoforms, see Zhang *et al* Science (2004), 303, 371, Sears *et al*, Science, (2001)  
35 291, 2344, Wacker *et al* (2002) Science, 298 1790, Davis *et al* (2002) Chem.Rev. 102, 579, Hang *et al* (2001) Acc.Chem.Res 34, 727. Thus the invention concerns a plurality of therapeutic antibodies (which maybe of the IgG isotype, e.g. IgG1) as described herein comprising a defined number (e.g. 7 or less, for example 5 or less such as two or a single) glycoform(s) of said antibodies.

40 Derivatives according to the invention also include therapeutic antibodies of the invention coupled to a non-proteinaeous polymer such as polyethylene glycol (PEG), polypropylene glycol or polyoxyalkylene. Conjugation of proteins to PEG is an established technique for increasing half-life of proteins, as well as reducing

antigenicity and immunogenicity of proteins. The use of PEGylation with different molecular weights and styles (linear or branched) has been investigated with intact antibodies as well as Fab' fragments, see Koumenis I.L. *et al* (2000) Int.J.Pharmaceut. 198:83-95. A particular embodiment comprises an antigen-binding fragment of the invention without the effector functions of a) activation of complement by the classical pathway; and b) mediating antibody-dependent cellular cytotoxicity; (such as a Fab fragment or a scFv) coupled to PEG.

## 10 **2. Production Methods**

Antibodies of the present invention may be produced in transgenic organisms such as goats (see Pollock *et al* (1999), J.Immunol.Methods 231:147-157), chickens (see Morrow KJJ (2000) Genet.Eng.News 20:1-55), mice (see Pollock *et al ibid*) or plants (see Doran PM, (2000) Curr.Opinion Biotechnol. 11, 199-204, Ma JK-C (1998), 15 Nat.Med. 4; 601-606, Baez J *et al*, BioPharm (2000) 13: 50-54, Stoger E *et al*; (2000) Plant Mol.Biol. 42:583-590). Antibodies may also be produced by chemical synthesis. However, antibodies of the invention are typically produced using recombinant cell culturing technology well known to those skilled in the art. A polynucleotide encoding the antibody is isolated and inserted into a replicable vector such as a plasmid for 20 further propagation or expression in a host cell. One useful expression system is a glutamate synthetase system (such as sold by Lonza Biologics), particularly where the host cell is CHO or NS0 (see below). Polynucleotide encoding the antibody is readily isolated and sequenced using conventional procedures (e.g. oligonucleotide probes). Vectors that may be used include plasmid, virus, phage, transposons, 25 minichromosomes of which plasmids are a typical embodiment. Generally such vectors further include a signal sequence, origin of replication, one or more marker genes, an enhancer element, a promoter and transcription termination sequences operably linked to the light and/or heavy chain polynucleotide so as to facilitate expression. Polynucleotide encoding the light and heavy chains may be inserted into 30 separate vectors and introduced (e.g. by transformation, transfection, electroporation or transduction) into the same host cell concurrently or sequentially or, if desired both the heavy chain and light chain can be inserted into the same vector prior to such introduction.

35 It will be immediately apparent to those skilled in the art that due to the redundancy of the genetic code, alternative polynucleotides to those disclosed herein are also available that will encode the polypeptides of the invention.

### **2.1 Signal sequences**

40 Antibodies of the present invention maybe produced as a fusion protein with a heterologous signal sequence having a specific cleavage site at the N terminus of the mature protein. The signal sequence should be recognised and processed by the host cell. For prokaryotic host cells, the signal sequence may be an alkaline phosphatase, penicillinase, or heat stable enterotoxin II leaders. For yeast secretion

the signal sequences may be a yeast invertase leader,  $\alpha$  factor leader or acid phosphatase leaders see e.g. WO90/13646. In mammalian cell systems, viral secretory leaders such as herpes simplex gD signal and native immunoglobulin signal sequences (such as human Ig heavy chain) are available. Typically the signal sequence is ligated in reading frame to polynucleotide encoding the antibody of the invention.

### **2.2 Origin of replication**

Origin of replications are well known in the art with pBR322 suitable for most gram-negative bacteria, 2 $\mu$  plasmid for most yeast and various viral origins such as SV40, polyoma, adenovirus, VSV or BPV for most mammalian cells. Generally the SV40 origin of replication component is not needed for integrated mammalian expression vectors. However the SV40 ori may be included since it contains the early promoter.

### **2.3 Selection marker**

Typical selection genes encode proteins that (a) confer resistance to antibiotics or other toxins e.g. ampicillin, neomycin, methotrexate or tetracycline or (b) complement auxotrophic deficiencies or supply nutrients not available in the complex media or (c) combinations of both. The selection scheme may involve arresting growth of the host cells that contain no vector or vectors. Cells, which have been successfully transformed with the genes encoding the therapeutic antibody of the present invention, survive due to e.g. drug resistance conferred by the co-delivered selection marker. One example is the DHFR-selection system wherein transformants are generated in DHFR negative host strains (eg see Page and Sydenham 1991 Biotechnology **9**: 64-68). In this system the DHFR gene is co-delivered with antibody polynucleotide sequences of the invention and DHFR positive cells then selected by nucleoside withdrawal. If required, the DHFR inhibitor methotrexate is also employed to select for transformants with DHFR gene amplification. By operably linking DHFR gene to the antibody coding sequences of the invention or functional derivatives thereof, DHFR gene amplification results in concomitant amplification of the desired antibody sequences of interest. CHO cells are a particularly useful cell line for this DHFR/methotrexate selection and methods of amplifying and selecting host cells using the DHFR system are well established in the art see Kaufman R.J. et al J.Mol.Biol. (1982) 159, 601-621, for review, see Werner RG, Noe W, Kopp K, Schluter M, "Appropriate mammalian expression systems for biopharmaceuticals", Arzneimittel-Forschung. 48(8):870-80, 1998 Aug. A further example is the glutamate synthetase expression system (Bebbington et al Biotechnology 1992 Vol 10 p169). A suitable selection gene for use in yeast is the trp1 gene; see Stinchcomb *et al* Nature 282, 38, 1979.

### **2.4 Promoters**

Suitable promoters for expressing antibodies of the invention are operably linked to DNA/polynucleotide encoding the antibody. Promoters for prokaryotic hosts include phoA promoter, Beta-lactamase and lactose promoter systems, alkaline

phosphatase, tryptophan and hybrid promoters such as Tac. Promoters suitable for expression in yeast cells include 3-phosphoglycerate kinase or other glycolytic enzymes e.g. enolase, glyceraldehyde 3 phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose 6 phosphate isomerase, 3-  
5 phosphoglycerate mutase and glucokinase. Inducible yeast promoters include alcohol dehydrogenase 2, isocytochrome C, acid phosphatase, metallothionein and enzymes responsible for nitrogen metabolism or maltose/galactose utilization. Promoters for expression in mammalian cell systems include RNA polymerase II promoters including viral promoters such as polyoma, fowlpox and adenoviruses  
10 (e.g. adenovirus 2), bovine papilloma virus, avian sarcoma virus, cytomegalovirus (in particular the immediate early gene promoter), retrovirus, hepatitis B virus, actin, rous sarcoma virus (RSV) promoter and the early or late Simian virus 40 and non-viral promoters such as EF-1alpha (Mizushima and Nagata Nucleic Acids Res 1990  
18(17):5322. The choice of promoter may be based upon suitable compatibility with  
15 the host cell used for expression.

### **2.5 Enhancer element**

Where appropriate, e.g. for expression in higher eukaryotes, additional enhancer elements can be included instead of or as well as those found located in the promoters  
20 described above. Suitable mammalian enhancer sequences include enhancer elements from globin, elastase, albumin, fetoprotein, metallothionein and insulin. Alternatively, one may use an enhancer element from a eukaryotic cell virus such as SV40 enhancer, cytomegalovirus early promoter enhancer, polyoma enhancer, baculoviral enhancer or murine IgG2a locus (see WO04/009823). Whilst such  
25 enhancers are typically located on the vector at a site upstream to the promoter, they can also be located elsewhere e.g. within the untranslated region or downstream of the polydenylation signal. The choice and positioning of enhancer may be based upon suitable compatibility with the host cell used for expression..

### **2.6 Polyadenylation/Termination**

In eukaryotic systems, polyadenylation signals are operably linked to polynucleotide encoding the antibody of this invention. Such signals are typically placed 3' of the open reading frame. In mammalian systems, non-limiting example signals include those derived from growth hormones, elongation factor-1 alpha and viral (eg SV40)  
35 genes or retroviral long terminal repeats. In yeast systems non-limiting examples of polydenylation/termination signals include those derived from the phosphoglycerate kinase (PGK) and the alcohol dehydrogenase 1 (ADH) genes. In prokaryotic system polyadenylation signals are typically not required and it is instead usual to employ shorter and more defined terminator sequences. The choice of  
40 polyadenylation/termination sequences may be based upon suitable compatibility with the host cell used for expression.

### **2.7 Other methods/elements for enhanced yields**

In addition to the above, other features that can be employed to enhance yields include chromatin remodelling elements, introns and host-cell specific codon modification. The codon usage of the antibody of this invention thereof can be modified to accommodate codon bias of the host cell such to augment transcript and/or product yield (eg Hoekema A et al Mol Cell Biol 1987 7(8):2914-24). The choice of codons may be based upon suitable compatibility with the host cell used for expression.

### **2.8 Host cells**

Suitable host cells for cloning or expressing vectors encoding antibodies of the invention are prokaryotic, yeast or higher eukaryotic cells. Suitable prokaryotic cells include eubacteria e.g. enterobacteriaceae such as *Escherichia* e.g. *E.Coli* (for example ATCC 31,446; 31,537; 27,325), *Enterobacter*, *Erwinia*, *Klebsiella* *Proteus*, *Salmonella* e.g. *Salmonella typhimurium*, *Serratia* e.g. *Serratia marcescans* and *Shigella* as well as Bacilli such as *B.subtilis* and *B.licheniformis* (see DD 266 710), *Pseudomonas* such as *P.aeruginosa* and *Streptomyces*. Of the yeast host cells, *Saccharomyces cerevisiae*, *schizosaccharomyces pombe*, *Kluyveromyces* (e.g. ATCC 16,045; 12,424; 24178; 56,500), *yarrowia* (EP402, 226), *Pichia Pastoris* (EP183, 070, see also Peng *et al* J.Biotechnol. 108 (2004) 185-192), *Candida*, *Trichoderma reesia* (EP244, 234), *Penicillin*, *Tolypocladium* and *Aspergillus* hosts such as *A.nidulans* and *A.niger* are also contemplated.

Although Prokaryotic and yeast host cells are specifically contemplated by the invention, typically however, host cells of the present invention are vertebrate cells. Suitable vertebrate host cells include mammalian cells such as COS-1 (ATCC No.CRL 1650) COS-7 (ATCC CRL 1651), human embryonic kidney line 293, , PerC6 (Crucell), baby hamster kidney cells (BHK) (ATCC CRL.1632), BHK570 (ATCC NO: CRL 10314), 293 (ATCC NO.CRL 1573), Chinese hamster ovary cells CHO (e.g. CHO-K1, ATCC NO: CCL 61, DHFR minus CHO cell line such as DG44 (Urlaub *et al*, Somat Cell Mol Genet (1986) Vol 12 pp555-566), particularly those CHO cell lines adapted for suspension culture, mouse sertoli cells, monkey kidney cells, African green monkey kidney cells (ATCC CRL-1587), HELA cells, canine kidney cells (ATCC CCL 34), human lung cells (ATCC CCL 75), Hep G2 and myeloma or lymphoma cells e.g. NS0 (see US 5,807,715), Sp2/0, Y0.

Thus in one embodiment of the invention there is provided a stably transformed host cell comprising a vector encoding a heavy chain and/or light chain of the therapeutic antibody as described herein. Typically such host cells comprise a first vector encoding the light chain and a second vector encoding said heavy chain.

Such host cells may also be further engineered or adapted to modify quality, function and/or yield of the antibody of this invention . Non-limiting examples include expression of specific modifying (eg glycosylation) enzymes and protein folding chaperones.

### **2.9 Cell Culturing Methods.**

Host cells transformed with vectors encoding the therapeutic antibodies of the invention may be cultured by any method known to those skilled in the art. Host cells  
5 may be cultured in spinner flasks, shake flasks, roller bottles, wave reactors (eg System 1000 from wavebiotech.com) or hollow fibre systems but it is preferred for large scale production that stirred tank reactors or bag reactors (eg Wave Biotech, Somerset, New Jersey USA) are used particularly for suspension cultures. Typically the stirred tankers are adapted for aeration using e.g. spargers, baffles or low shear  
10 impellers. For bubble columns and airlift reactors direct aeration with air or oxygen bubbles maybe used. Where the host cells are cultured in a serum free culture media this can be supplemented with a cell protective agent such as pluronic F-68 to help prevent cell damage as a result of the aeration process. Depending on the host cell characteristics, either microcarriers maybe used as growth substrates for  
15 anchorage dependent cell lines or the cells maybe adapted to suspension culture (which is typical). The culturing of host cells, particularly vertebrate host cells may utilise a variety of operational modes such as batch, fed-batch, repeated batch processing (see Drapeau *et al* (1994) *cytotechnology* 15: 103-109), extended batch process or perfusion culture. Although recombinantly transformed mammalian host  
20 cells may be cultured in serum-containing media such media comprising fetal calf serum (FCS), it is preferred that such host cells are cultured in serum –free media such as disclosed in Keen *et al* (1995) *Cytotechnology* 17:153-163, or commercially available media such as ProCHO-CDM or UltraCHO™ (Cambrex NJ, USA), supplemented where necessary with an energy source such as glucose and  
25 synthetic growth factors such as recombinant insulin. The serum-free culturing of host cells may require that those cells are adapted to grow in serum free conditions. One adaptation approach is to culture such host cells in serum containing media and repeatedly exchange 80% of the culture medium for the serum-free media so that the host cells learn to adapt in serum free conditions (see e.g. Scharfenberg *K et al*  
30 (1995) in *Animal Cell technology: Developments towards the 21st century* (Beuvery E.C. *et al* eds), pp619-623, Kluwer Academic publishers).

Antibodies of the invention secreted into the media may be recovered and purified from the media using a variety of techniques to provide a degree of purification  
35 suitable for the intended use. For example the use of therapeutic antibodies of the invention for the treatment of human patients typically mandates at least 95% purity as determined by reducing SDS-PAGE, more typically 98% or 99% purity, when compared to the culture media comprising the therapeutic antibodies. In the first instance, cell debris from the culture media is typically removed using centrifugation  
40 followed by a clarification step of the supernatant using e.g. microfiltration, ultrafiltration and/or depth filtration. Alternatively, the antibody can be harvested by microfiltration, ultrafiltration or depth filtration without prior centrifugation. A variety of other techniques such as dialysis and gel electrophoresis and chromatographic techniques such as hydroxyapatite (HA), affinity chromatography (optionally involving

an affinity tagging system such as polyhistidine) and/or hydrophobic interaction chromatography (HIC, see US 5, 429,746) are available. In one embodiment, the antibodies of the invention, following various clarification steps, are captured using Protein A or G affinity chromatography followed by further chromatography steps  
5 such as ion exchange and/or HA chromatography, anion or cation exchange, size exclusion chromatography and ammonium sulphate precipitation. Typically, various virus removal steps are also employed (e.g. nanofiltration using e.g. a DV-20 filter). Following these various steps, a purified (typically monoclonal) preparation comprising at least 10mg/ml or greater e.g. 100mg/ml or greater of the antibody of  
10 the invention is provided and therefore forms an embodiment of the invention. Concentration to 100mg/ml or greater can be generated by ultracentrifugation. Suitably such preparations are substantially free of aggregated forms of antibodies of the invention.

15 Bacterial systems are particularly suited for the expression of antibody fragments. Such fragments are localised intracellularly or within the periplasma. Insoluble periplasmic proteins can be extracted and refolded to form active proteins according to methods known to those skilled in the art, see Sanchez *et al* (1999) J. Biotechnol. 72, 13-20 and Cupit PM *et al* (1999) Lett Appl Microbiol, 29, 273-277.

### **3. Pharmaceutical Compositions**

Purified preparations of antibodies of the invention (particularly monoclonal preparations) as described *supra*, may be incorporated into pharmaceutical compositions for use in the treatment of human diseases and disorders such as  
25 those outlined above. Typically such compositions further comprise a pharmaceutically acceptable (i.e. inert) carrier as known and called for by acceptable pharmaceutical practice, see e.g. Remington's Pharmaceutical Sciences, 16th ed, (1980), Mack Publishing Co. Examples of such carriers include sterilised carrier such as saline, Ringers solution or dextrose solution, buffered with suitable buffers  
30 such as sodium acetate trihydrate to a pharmaceutically acceptable pH, such as a pH within a range of 5 to 8. Pharmaceutical compositions for injection (e.g. by intravenous, intraperitoneal, intradermal, subcutaneous, intramuscular or intraportal) or continuous infusion are suitably free of visible particulate matter and may comprise from 1mg to 10g of therapeutic antibody, typically 5mg to 1g, more specifically 5mg  
35 to 25mg or 50mg of antibody. Methods for the preparation of such pharmaceutical compositions are well known to those skilled in the art. In one embodiment, pharmaceutical compositions comprise from 1mg to 10g of therapeutic antibodies of the invention in unit dosage form, optionally together with instructions for use. Pharmaceutical compositions of the invention may be lyophilised (freeze dried) for  
40 reconstitution prior to administration according to methods well known or apparent to those skilled in the art. Where embodiments of the invention comprise antibodies of the invention with an IgG1 isotype, a chelator of metal ions including copper, such as citrate (e.g. sodium citrate) or EDTA or histidine, may be added to the pharmaceutical composition to reduce the degree of metal-mediated degradation of



antibodies of this isotype, see EP0612251. Pharmaceutical compositions may also comprise a solubiliser such as arginine base, a detergent/anti-aggregation agent such as polysorbate 80, and an inert gas such as nitrogen to replace vial headspace oxygen.

5

Effective doses and treatment regimes for administering the antibody of the invention are generally determined empirically and are dependent on factors such as the age, weight and health status of the patient and disease or disorder to be treated. Such factors are within the purview of the attending physician. Guidance in selecting appropriate doses may be found in e.g. Smith *et al* (1977) *Antibodies in human diagnosis and therapy*, Raven Press, New York but will in general be 1mg to 10g. In one embodiment, the dosing regime for treating a human patient is 1mg to 1g of therapeutic antibody of the invention administered subcutaneously once per week or every two weeks, or by intravenous infusion every 1 or 2 months. Such a dosage corresponds to 0.014-140mg/kg, such as 0.014-14mg/kg. Compositions of the present invention may also be used prophylactically.

10

15

#### **4. Clinical uses.**

It will be appreciated that diseases characterised by elevated  $\beta$ -amyloid levels or  $\beta$ -amyloid deposits include Alzheimer's disease, mild cognitive impairment, Down's syndrome, hereditary cerebral haemorrhage with  $\beta$ -amyloidosis of the Dutch type, cerebral  $\beta$ -amyloid angiopathy and various types of degenerative dementias, such as those associated with Parkinson's disease, progressive supranuclear palsy, cortical basal degeneration and diffuse Lewis body type of Alzheimer's disease.

20

Most preferably, the disease characterised by elevated  $\beta$ -amyloid levels or  $\beta$ -amyloid deposits is Alzheimer's disease.

Although the present invention has been described principally in relation to the treatment of human diseases or disorders, the present invention may also have applications in the treatment of similar diseases or disorders in non-human mammals.

25

30

**Examples****Methods**

	Biacore™/Biacore 3000	a device that allows measurement of real time kinetics of molecular interactions using SPR
5	SPR	(surface plasmon resonance) – physical phenomenon employed by Biacore™ instruments for measurement of mass changes on sensor chip
	CM5	Biacore™ sensor chip with general purpose surface coated with a carboxymethylated dextran matrix
10	ELISA	enzyme linked immunosorbent assay
	SRU	SRU BIND™ Biosensor technology allowing to monitor label-free biochemical interactions
	Integra CL1000	Mini-bioreactors sold by IBS Integra Biosciences
	FMAT	fluorometric microvolume assay technology (Applied Biosystems)
15	ABi8200	Applied Biosystems 8200 fluorescence macro confocal cellular detection system for FMAT
	FPLC	Fast protein liquid chromatography
	ProSepA HiTrap	Protein A columns for FPLC sold by GE Healthcare

20

**Materials**

	DMSO	dimethylsulphoxide
	HEPES	N-(2-hydroxyethyl)piperazine-N'-(2-ethanesulfonic acid)
25	EDTA	ethylenediaminetetraacetic acid
	Tris HCl	- tris-(hydroxymethyl)aminomethane hydrochloride
	NaCl	- sodium chloride
	Tween-20	- polyoxyethylenesorbitan monolaurate
	BSA	- bovine serum albumin
30	PBS	- phosphate buffered saline
	PFA	- paraformaldehyde
	IMS	- industrial methylated spirit
	DAB	- 3,3'diaminobenzidine
	DMEM	dulbecco's modified eagle's medium
35	FCS	fetal calf serum
	Opti-MEM	modified eagle's medium based medium by Invitrogen/Gibco
	Lipofectamine	cationic lipid based cell transfection agent sold by Invitrogen/Gibco
	Transfast	liposomal transfection agent sold by Promega
40	Versene	metal ion chelating agent (ethylenediaminetetraacetic acid)
	Glutamax	stable form of glutamine added to culture medium (dipeptide L-Ananyl-L-Glutamine supplement)
	Histoclear	tissue clearing agent

HBS-EP buffer            General purpose Biacore™ buffer containing 0.01M HEPES  
pH7.4, 0.15M NaCl, 3mM EDTA, 0.005% Surfactant P20

5    **Generation of mouse monoclonal antibody 2E7**

Mouse monoclonal antibody 2E7 was generated from a conventional immunisation of mice. Mice were immunised with soluble or aggregated  $\beta$ -amyloid 1-40 and 1-42 formulated in Freund's adjuvant. Following final boost without adjuvant, splenocytes were fused with myeloma cells. Fused cells were grown in 96-well plates from which hybridoma supernatants were screened for potential leads. Selected antibody 2E7, which was obtained from the immunisation with soluble  $\beta$ -amyloid 1-40, was of murine IgG2a isotype and had beta-amyloid binding activity in the efflux assay described below and an affinity of 36.1 pM for beta-amyloid 1-40 when measured by Biacore™, Method A(i) (Table 10A).

15

**Epitope Mapping of 2E7**

In order to finely map the binding of antibody 2E7 to the  $\beta$ -amyloid peptide, a peptide set (A) was utilised. Peptide set (A) consisted of a set of 31 12-mer overlapping peptides covering the complete sequence of the  $\beta$ -amyloid 1-42 peptide. Each sequential peptide was initiated at the sequential amino acid within the  $\beta$ -amyloid peptide, thus shifting the sequence covered between sequential peptides by a single amino acid. All peptides in set (A) contained a 3 amino acid C-terminal linker (glycine-serine-glycine) and a terminal biotinylated lysine residue. In addition, all peptides except peptide A $\beta$ 1 DAEFRHDSGYEVGSGK-biotin (SEQ ID No:15) were N-terminally acetylated. A second set of peptides (set (B)) consisted of sequential one amino acid C-terminal deletions from a peptide containing amino acids 1 to 10 of the  $\beta$ -amyloid sequence. All peptides in set (B) contained a 3 amino acid C-terminal linker (glycine-serine-glycine) and a terminal biotinylated lysine residue, but with additional glycine and serine residues to replace for deleted  $\beta$ -amyloid amino acids (Table 2). Thus all peptides in set (B) are of the same length.

Table 2

Sequences of biotinylated peptides (set (B)) that contained truncated N-terminal fragments of  $\beta$ -amyloid

	DAEFRHDSGYGSGGSK-biotin	(SEQ ID No:7)
	DAEFRHDSG--GSGSGSK-biotin	(SEQ ID No:8)
	DAEFRHDS--GSGGSGGK-biotin	(SEQ ID No:9)
40	DAEFRHD--GSGGSGGSK-biotin	(SEQ ID No:10)
	DAEFRH--GSGGSGGSGK-biotin	(SEQ ID No:11)
	DAEFR--GSGGSGGSGSK-biotin	(SEQ ID No:12)
	DAEF--GSGGSGGSGGSK-biotin	(SEQ ID No:13)
	DAE--GSGGSGGSGGSGK-biotin	(SEQ ID No:14)

### Monitoring the binding of 2E7 to $\beta$ -amyloid derived peptides using Optical Biosensors

96-well SRU Bind™ streptavidin-coated plates (SRU Biosystems) were washed with PBS containing 1% DMSO to remove glycerol and preservative. A volume of 50ul/well was left to equilibrate to room temperature to provide a constant base line. Biotinylated peptides were diluted to approx. 0.3ug/ml in PBS containing 1% DMSO and 50ul of each added to wells and incubated for approximately 1h. Replicate wells were prepared using different sectors of the plate and at least one no-peptide control well was used in each sector to reference subtract the data. After peptide capture the plate was washed with PBS containing 1% DMSO, leaving 50ul of fresh buffer per well to provide a new base line on the reader. No decay of peptide from the surface was seen. The buffer was then replaced with 40ul/well buffer containing test antibody at 20-64nM. for 2 hours. It was found that antibody 2E7 only bound to the peptide encompassing amino acids 1-12 of the  $\beta$ -amyloid peptide in peptide set (A) (peptide A $\beta$ 1, SEQ ID No:15) This result implies that the aspartic acid at residue 1 is critical for binding to this peptide.

In order to further characterise the binding site of antibody 2E7, peptide set (B) was utilised. Using SRU BIND™ biosensor methodology antibody 2E7 showed negligible binding to the peptides encompassing amino acids 1-3 and 1-4 of the  $\beta$ -amyloid peptide (SEQ ID No:14 and 13). Binding to a peptide encompassing amino acids 1-7 of the  $\beta$ -amyloid peptide (SEQ ID No:10) was comparable to the peptide encompassing amino acids 1-12 of the  $\beta$ -amyloid peptide (from peptide set (A)). Binding to peptides encompassing amino acids 1-5 or 1-6 of the  $\beta$ -amyloid peptide (SEQ ID No:12 or 11) was observed, but was weaker (as measured by stability after an additional washing step) than the binding to the peptide encompassing amino acids 1-7 of the  $\beta$ -amyloid peptide (SEQ ID No:10). Thus it has been shown that only residues 1-7 of the  $\beta$ -amyloid peptide are required for full binding as measured using this methodology.

### Surface Plasmon Resonance assay

In addition to the experiments described above, the Biacore™ 3000 optical biosensor was used to monitor the binding of 2E7 antibody to selected  $\beta$ -amyloid sequence derived peptides. Binding was measured by injecting test antibodies at up to 64nM for 5 minute over peptides captured on separate streptavidin chip surfaces (130-230 RU (resonance units)). A running buffer (HBS-EP) containing 0.01M HEPES pH7.4, 0.15M NaCl, 3mM EDTA and 0.005% Surfactant P20™ at 25°C was used at a flow rate of 20ul/min. All runs were double referenced against a blank streptavidin surface and blank injections. Analysis was carried out using the Biacore™ analysis software BIAevaluation™ version 4.1. Results from selected peptides in set (A) further confirmed the SRU BIND™ derived data indicating that 2E7 bound only to the peptide encompassing amino acids 1-12 (SEQ ID No:15) of the  $\beta$ -amyloid peptide

with an apparent equilibrium constant  $K_D$  of approximately 50pM. Under the same conditions, 2E7 did not bind to the peptide encompassing amino acids 2-13 of the  $\beta$ -amyloid peptide.

Peptide A $\beta$ 2-13                      AEFRHDSGYEVHGSGK-biotin (SEQ ID No:44)

5

The experimental method and conditions used allowed the detection of high but also lower affinity molecules – in the same experimental setup, by contrast to 2E7, another antibody recognising an N-terminal epitope of the  $\beta$ -amyloid peptide was shown to bind the 2-13 peptide (SEQ ID No:44) with an apparent  $K_D$  of 7nM. 2E7 did not bind to a selection of peptides in set (A) from mid regions of the  $\beta$ -amyloid peptide. In a separate experiment the  $\beta$ -amyloid 1-40 peptide was captured via its N-terminal aspartic acid residue that had been biotinylated. This peptide was captured onto a Biacore™ streptavidin coated chip as previously described. Antibody 2E7 injected at 66nM for 1 minute could not bind this peptide. Therefore, it is concluded that the previously described N-terminal binding site was masked by the linker and capture method, thus further confirming the extreme N-terminus as containing the core binding site

10

15

20

#### **Binding to Cell Expressed Amyloid Precursor Protein (APP)**

$\beta$ -Amyloid is composed of peptides formed by proteolytic cleavage of a type I transmembrane precursor protein named amyloid precursor protein (APP). As APP has a large extracellular domain, binding to this protein could potentially initiate an antibody-dependent cellular cytotoxicity reaction (ADCC).

25

To characterise binding of antibody to cell-surface full length APP an FMAT™ ABI8200 based assay was utilised.

#### **Transfection of HEK293T cells with wild type APP DNA**

30

HEK293T cells are maintained in DMEM F12 medium containing 10 % (v/v) FCS and 1x Glutamax. Cells are seeded in 75cm<sup>2</sup> tissue culture flasks and grown to 60-80 % confluency (18-24 h) prior to transfection. For transfection, 9 ug of DNA, (either wild type APP DNA (in PCDNA3.1 (Invitrogen) vector), or vector only controls), is mixed with 0.3 ml of Opti-MEM™ media. 30ul Lipofectamine™ transfection agent is mixed with 0.3ml Opti-MEM™ media and the two mixtures pooled. The pooled mixtures are incubated at room temperature for 30 min prior to the addition of a further 4.8 ml of Opti-MEM™ media. The final mixture is added to the cells (post washing with Opti-MEM™ media) for 5 h and 6 ml of 10 % (v/v) newborn calf serum in DMEM is then added. 48hrs post transfection, supernatant is removed and the monolayer washed in versene, and then 3ml of Versene™ chelating agent is added to each flask, incubated for 5mins at 37C, and the detached cells pelleted at 200g for 5mins. The resultant cell pellet is gently resuspended in 1ml of assay buffer (2% heat treated serum, 0.5% BSA, 0.1% NaN<sub>3</sub> in PBS pH7.4, filtered through a 0.2um filter) to create a single cell suspension.

35

40

FMAT™ ABI8200 based assay

Test antibodies (2E7, LN27 (Zymed) mouse IgG to extracellular domain of APP as a positive control, and an antibody G210 which recognises the x-40 form of the  $\beta$ -amyloid peptide as a negative control) were diluted to 10 $\mu$ g/ml in sterile filtered assay buffer (2% heat treat serum, 0.5% BSA, 0.1% NaN<sub>3</sub> in PBS pH7.2) in a polypropylene plate, and then a further six serial 1:1 dilutions were performed down the plate. Assay buffer only was used as a blank. 50ul of a suspension of HEK293T cells transfected with wild type APP DNA (Experiment 1: 10,000 cells; Experiment 2: 20,000 cells) was added to each well of a 96 well plate, to which 5ul of each of the antibody solutions were added to duplicate wells. 50ul/well of F-MAT™ blue anti mouse IgG stock, (antibody is labelled using F-MAT™ blue monofunctional reactive dye kit from ABI, 4328408), diluted 1:500 (Experiment 1) and 1:1000 (Experiment 2) in assay buffer, was then added to each well and the plate briefly shaken and left to settle for 1hr. The plate was then read using the ABI 8200 fluorescence macro confocal cellular detection system (Applied Biosystems).

Derived counts data were then interpreted using Excel™ spreadsheet software. Briefly, mock transfected counts were subtracted from the full length APP transfected cell counts to obtain a specific signal for each antibody. Two antibody concentrations that were on the linear part of the curve were chosen (1.25 and 0.63ug/ml) and the background corrected derived counts at these concentrations expressed as the percentage of the LN27 antibody binding, and averaged over the two antibody concentrations. The resultant data is described in Table 3 (% of LN27 binding  $\pm$ SE)

Thus, within this assay system, the binding of 2E7 to cell surface APP is indistinguishable from that of the negative control antibody G210.

**Table 3**

antibody	Experiment 1	Experiment 2
<b>LN27</b>	100.0 $\pm$ 7.1	100.0 $\pm$ 4.7
<b>G210</b>	5.5 $\pm$ 1.3	2.0 $\pm$ 1.6
<b>2E7</b>	9.9 $\pm$ 3.7	2.2 $\pm$ 1.4

30

**Binding to Amyloid Precursor Protein Derived Peptide**

The previously described epitope mapping studies have shown that antibody 2E7 binds to the extreme N-terminus of the  $\beta$ -amyloid peptide, with the aspartic acid residue at position 1 being essential for binding. This suggests that the antibody recognises a 'neo' epitope formed by cleavage of APP at the  $\beta$ -secretase site. This observation would suggest that antibody 2E7 should not recognise adjacent APP

35

peptide sequence. To test this hypothesis an APP peptide (Peptide APP1, SEQ ID No:16) was synthesised which included residues 1-7 of the  $\beta$ -amyloid peptide and the five adjacent APP derived amino acids. Thus peptide APP1 contained contiguous amino acids from position 5 N-terminal to the BACE-1 cleavage site to position 7 C-terminal to the BACE-1 cleavage site and was N-terminally acetylated. The ability of antibody 2E7 to bind to the APP derived peptide APP1 and the  $\beta$ -amyloid 1-12 peptide (peptide A $\beta$ 1) was compared using Biacore™ methodology (as previously described for epitope mapping). Antibody 2E7 showed high affinity binding to the  $\beta$ -amyloid peptide A $\beta$ -1, which contains the basic epitope 1-7. However, no binding was observed to the APP1 peptide which also contains the basic  $\beta$ -amyloid derived sequence 1-7.

Peptide A $\beta$ 1                      DAEFRHDSGYEVGSGK-biotin    SEQ ID No:15  
 APP1                                AcNH-SEVKMDAEFRHDGSGK-biotin    SEQ ID No:16

A combination of FMAT™ based cellular binding and Biacore™ based peptide mapping has been utilised to show that, in these formats, 2E7 has no binding affinity for the full length APP protein. Given that the aspartic acid residue at position 1 of the  $\beta$ -amyloid peptide is required for binding, it is concluded that 2E7 only recognises the 'neo' N-terminus of  $\beta$ -amyloid and hence should not bind cell surface expressed APP.

### **In vivo Biological Activity**

#### **<sup>125</sup>I $\beta$ -Amyloid Efflux Model**

A number of published studies have shown that  $\beta$ -amyloid antibodies can form complexes with  $\beta$ -amyloid peptide in the bloodstream. It is argued that this sequestration of peripheral  $\beta$ -amyloid allows for further efflux of CNS amyloid into the bloodstream (DeMattos RB, PNAS (2001), 98(15); 8850-8855). An acute pharmacodynamic model was developed to screen antibodies for their ability to complex with brain derived  $\beta$ -amyloid peptide in the bloodstream.

Anaesthesia (4% isoflurane) was induced in male C57/BL6J mice and maintained (1.5% isoflurane) in 100% oxygen. Animals were then placed in a stereotaxic frame. Following midline incision along the sagittal suture a bore hole was drilled through the skull and a guide cannula was inserted into the lateral cerebral ventricle (coordinates anterioposterior (AP) -0.5mm, lateral (L) +0.7mm, ventral (V) -2.5mm). A further two bore holes were drilled through the skull into which cortical screws were placed. The cannula was anchored in place by cyanoacrylate gel and the incision was sutured around the cyanoacrylate gel headcap. Post-operatively the mice received 0.3ml saline subcutaneously and were placed in a warm environment to recover from anaesthesia. On recovery of the righting reflex, mice were housed singly and received 5 days standard post-op care. No procedures were permitted for a further 5 days or until pre-operative body weight was regained. Following recovery, cannula placement was verified by the angiotensin II drinking response. Each mouse received an intracerebroventricular (ICV) administration (5 $\mu$ l) of 100ng angiotensin II

(All) (made up in 0.9% saline). Following administration of All, water intake was observed for 15 minutes. Mice with a positive dipsogenic response to All (sustained drinking) were included in the studies, which commenced no sooner than five days post All injection.

5

On the day of study the mice were placed for 5-10 minutes in a warm environment to induce vasodilation, necessary for ease of injection into the tail vein. Test antibody (600 $\mu$ g) or PBS vehicle (dose volume no greater than 10ml per kg body weight) was injected via the tail vein and mice were returned to their individual cages post-injection. At exactly one hour post tail vein injection, mice were slowly ICV injected (2 $\mu$ l per minute) with 2ng (1 $\mu$ Ci) of I<sup>125</sup> beta-amyloid 1-40 (Amersham Biosciences, UK) in a dose volume of 5 $\mu$ l. At exactly four hours post ICV dose, 50 $\mu$ l of trunk blood was collected and the radioactivity level measured on a scintillation counter.

10

15

Mice that had been injected into the tail-vein with 2E7 (n=6 per treatment group) showed a statistically significant increase in the radioactive signal (counts per minute – CPM) in 50 $\mu$ l of trunk blood compared with the CPM signal detected in vehicle injected mice – (CPM – vehicle: 1339.7  $\pm$  496.2 vs. 2E7 4387.9  $\pm$  980.3; ANOVA:F(2,13) = 4.97, p<0.05. Post-hoc LSD: p=0.01 2E7 vs. vehicle [post-hoc Duncans: p=0.02 2E7 vs, vehicle]).

20

In two further studies with 2E7 conducted with the identical protocol, similar increases in amyloid efflux into blood when compared with vehicle injected controls were observed (CPM blood: Vehicle 352  $\pm$  113 versus 2E7 2397  $\pm$  353, and Vehicle 1281  $\pm$  312 versus 2E7 5291  $\pm$  885; ANOVA with post-hoc LSD test p<0.001 vs. vehicle).

25

### **Transgenic CNS $\beta$ -Amyloid Lowering Models**

30

#### **1. $\beta$ -Amyloid Load following 4 week dosing of 2 month old TASTPM mice**

Male and female TASTPM transgenic mice (double-mutant APP<sup>swe</sup> x PS1.M146V, Howlett DR (2004) Brain Research 1017 (1-2) 130-136) aged between 61 and 65 days at the start of the study and were singly housed. Equal numbers of mice were assigned to each treatment group (N=12 per group) and were randomized according to gender and age. Treatment groups comprised the following: A: MOPC21 (antibody with unknown specificity, Holton et al (1987) J.Immunol 139(9) 3041-3049, negative control), B: 2E7 (test antibody). All antibodies were dissolved in PBS and were dosed by the intraperitoneal route. Irrespective of animal weight, 300 $\mu$ g of antibody was administered. Animals were dosed twice weekly for four weeks. One day after the final dose, animals were euthanased by overdose with sodium pentobarbital. Brains were dissected and hemisected. Hemisected brain samples were collected into pre-weighed 2ml eppendorf<sup>TM</sup> tubes and snap frozen. Samples were subsequently thawed, reweighed and 1ml of 5M guanidine HCl containing Complete protease inhibitor<sup>TM</sup> tablets (Boehringer Mannheim) added, before the

35

40



samples were homogenized and incubated at 4°C for >90 min with constant agitation.

5 Samples were then diluted 1 in 10 into assay buffer (50mM Tris HCl, pH7.4, 150mM NaCl, 0.05% Tween-20+ 1% BSA), vortexed and spun at 20,000G for 20 mins at 4°C. The supernatant was removed and added as triplicate samples to the assay plate.

10 The levels of A $\beta$ 40 and A $\beta$ 42 were measured using a sensitive plate based electrochemiluminescent immunoassay (BioVeris™) employing C-terminal specific  $\beta$ -amyloid antibodies (to A $\beta$ 40 or A $\beta$ 42) labelled with Oritag™ specific label to facilitate detection (BioVeris™) used to capture either A $\beta$ 40 or A $\beta$ 42, along with a biotinylated N-terminal specific A $\beta$  antibody. Antibody-A $\beta$  complexes were captured with streptavidin coated beads that bind biotinylated antibodies (Dynabeads™, Dynal)

15 incubated overnight at room temperature with vigorous mixing and assayed in a BioVeris™ M8 photodetector. Standard curves were constructed using human A $\beta$ 40 and A $\beta$ 42 peptides in assay buffer containing the required concentration of Guanidine HCl. Data was analysed using Excel Robosage™ statistical analysis software and A $\beta$  levels expressed as pmole/g tissue.

20

In this paradigm, treatment with 2E7 antibody reduced CNS A $\beta$ 42 load by 37% (p<0.001). and CNS A $\beta$ 40 by 23% (p<0.001).

25 In subsequent studies under similar experimental conditions, 2E7 antibody reduced CNS A $\beta$ 42 load by 38% (Study 1, males only), 22% (Study 2, non-significant) and 39% (Study 3, males, p=0.001) and 13% (Study 3, females, non-significant) when compared to PBS treated animals. In these studies 2E7 also reduced CNS A $\beta$ 40 by 18% (Study 3, males, p=0.017) and offered a non-significant reduction in CNS A $\beta$ 40 by 25% (Study 1, males only), <1% (Study 2) and a non-significant increase of 3%

30 (Study 3, females) when compared to PBS treated animals.

## 2. $\beta$ -Amyloid Load following 4 month dosing of 4 month old TASTPM mice

Briefly, 4 month old TASTPM transgenic mice were dosed 300 $\mu$ g of antibody once or twice weekly via an intraperitoneal (i.p.) route. After 4 months of dosing CNS  $\beta$ -amyloid levels were measured by ELISA and plaque load was measured by immunohistochemistry. Between the ages of 4 and 8 months, the CNS  $\beta$ -amyloid load increases exponentially and consequently, plaque pathology rapidly develops (Howlett DR (2004) Brain Research 1017 (1-2) 130-136).

35

40 Mice were aged between 120 and 128 days at the start of the study and were singly housed. Similar numbers of mice were assigned to each treatment group (N=20 or 21 per group) and were randomized according to gender and age. Treatment groups comprised the following: A: PBS (vehicle) dosed twice weekly, B: 2E7 dosed once weekly, C: 2E7 dosed twice weekly, D: PBS dosed once weekly. A 300 microgram

dose (79 microlitres volume) of 2E7 was administered via the intraperitoneal route. Vehicle treated animals received the same volume of PBS. Animals were dosed for eighteen weeks. TASTPM mice are liable to suffer spontaneous seizures and as a result a number of animals died during the course of the study. Final numbers were as follows A: 4 females, 9 males; B: 5 females, 8 males; C: 4 females, 9 males; D: 2 females, 9 males. Two or four days after the final dose (equal numbers per group) animals were euthanased by overdose with sodium pentobarbital. A tail tip sample from each mouse was taken for confirmation of the genotype. Brains were dissected and hemisected. The right hemisphere was fixed by immersion in 4% paraformaldehyde and processed for histology. The left hemisphere was collected into pre-weighed 2ml eppendorf<sup>TM</sup> tubes, frozen on dry ice and stored at -80°C for subsequent analysis of amyloid content. Prior to analysis, samples were thawed, reweighed and 1ml of 5M guanidine HCl containing Complete protease inhibitor<sup>TM</sup> tablets (Boehringer Mannheim) added, before the samples were homogenized and incubated at 4°C for >90 min with constant agitation.

Samples were then diluted 1 in 10 into assay buffer (50mM Tris HCl, pH7.4, 150mM NaCl, 0.05% Tween-20+ 1% BSA), vortexed and spun at 20,000G for 20 mins at 4°C. The supernatants were diluted a further 1:1000 and added as triplicate samples to the assay plate.

The levels of A $\beta$ 40 and A $\beta$ 42 were measured as for the 4 week dosing study.

An analysis of variance was used with treatment, sex and dosing schedule included in the model as fixed effects. All of the interactions between the three factors were also included. There were no significant differences between the two dosing schedules (once or twice weekly). With this experimental design, firstly it could be assessed if there were any significant differences between the dosing schedules and secondly, as there were no such significant differences, data from the two dosing schedules could be combined, thus increasing the power of the experiment by doubling the number of mice in the analysis.

In this paradigm, treatment with 2E7 antibody reduced CNS A $\beta$ 42 load by 22.5% (p=0.0152). Levels of CNS A $\beta$ 40 were also lowered by 12.1%, but this figure did not reach statistical significance (p=0.118).

A complex immunohistochemical analysis of these samples was performed to define the area of brain tissue showing plaque pathology. Sections were taken from the cortex at the level of the caudate and from the cortex at the level of the hippocampus. Adjacent sections were stained with either an A $\beta$ 40 or A $\beta$ 42 specific antibody or alternatively with the amyloid stain Congo Red. Using image analysis software, the area of the section stained for plaque was expressed as a percentage of the total section area.

After fixation, the PFA-immersed half brains were coronally cut in a brain matrix into 6 x 2mm thick sections. These 2mm sections will be referred to as sections A to F, A being the most rostral and F the most caudal. Sections A, B & C and D, E & F were placed in separate embedding cassettes numbered for each animal. Cassettes were held in PFA until ready for processing and embedding.

Embedding was undertaken on a Citadel™ 1000 (Shandon) tissue processor. All tissues received the following processing regimen:-

70% IMS – 1 hr

100% IMS – 3 x 1hr

100% ethanol – 2hr

100% isobutyl alcohol; 1 x 2hr; 1 x 1.5hr

Histoclear™ – 2 x 1.5hr

Paraffin wax – 2 x 2hr

On completion of the processing cycle, the wax impregnated tissue sections were transferred to molten-paraffin wax filled base moulds and embedded utilising a Histocentre™ (Shandon) paraffin embedding system. Tissue was embedded such that sections A, B & C went into one mould; D, E & F into a second mould. This was carried out for all sets of sections ie. each hemisected brain resulted in two wax blocks of three sections each. Sections were placed in the moulds such that the caudal surface of each piece became the future cutting surface. Care was taken to ensure that each section was pushed well down in the mould so that microtoming of each would occur in parallel. The perforated processing cassette was then carefully placed onto each mould which was then topped up with molten wax. Embedded blocks were then cooled on the refrigerated plate until they could be removed from the moulds. Blocks were stored at room temperature until required for microtoming. Blocks were cut at random and 5 micron sections floated onto prelabelled gelatine coated slides (Superfrost™, Erie Scientific Company) slides. Two sections were floated onto each slide. Wherever possible, consecutive sections were mounted and slides were numbered consecutively from 1 to 25. Fifty sections (25 slides) were taken from each block. Slides were dried on a hot plate and then stored at room temperature until required.

Immunohistochemistry was undertaken on sets of 30 slides. On each slide, the top section was labelled with an A $\beta$ 40 antibody (G30, rabbit polyclonal recognising x-40  $\beta$ -amyloid), the lower section with the A $\beta$ 42 antibody, 20G10, monoclonal antibody recognising x-42  $\beta$ -amyloid. A minimum of 5 sections per antibody per block were labelled.

Labelling was carried out as follows. Following dewaxing through Histoclear and graded alcohols, sections were immersed in 85% formic acid for 8 minutes and then blocked in 0.3% hydrogen peroxide for 30 minutes to block endogenous peroxidases. Antibodies G30 and 20G10 were both applied overnight at 1:1000 dilutions, sections

being left at 4°C. Development of the sections was with the respective biotinylated anti rabbit and anti-mouse secondaries. Colour development was accomplished with a diaminobenzidine tetrahydrochloride staining kit (DAB<sup>TM</sup>, Vector Labs). Sections were briefly counterstained with Mayer's hematoxylin before being dehydrated, cleared and cover-slipped.

Sections were left to dry for at least 48 hours before microscopy. Images were captured on a Leica DMRB<sup>TM</sup> microscope equipped with digital camera. Images were analysed using Qwin<sup>TM</sup> software (Leica) and results presented as % of the section area that was labelled by the A $\beta$  antibody.

An analysis of variance was used with treatment, sex and dosing schedule included in the model as fixed effects. All of the interactions between the three factors were also included. There were no significant differences between the two dosing schedules (once or twice weekly). With this experimental design, firstly we could assess if there were any significant differences between the dosing schedules and secondly, as there were no such significant differences, data from the two dosing schedules could be combined, thus increasing the power of the experiment by doubling the number of mice in the analysis.

In this paradigm, treatment with 2E7 antibody reduced plaque pathology as measured with an antibody recognising A $\beta$ 42. Plaque pathology was reduced by 27.1% (p=0.0026) in the cortex at the level of the hippocampus and 43% (p<0.0001) in the cortex at the level of the caudate. Plaque pathology was also reduced when measured with an antibody recognising A $\beta$ 40. Plaque pathology was reduced by 16.6% (p=0.0421) in the cortex at the level of the hippocampus and 17.3% (p=0.0342) in the cortex at the level of the caudate.

No evidence of microhaemorrhage (as determined by Perls' Prussian Blue) was observed in any mice from this study treated with vehicle or 2E7. This method visualises ferric iron (iron is an essential constituent of the oxygen-carrying haemoglobin found in red cells) by producing an insoluble blue compound. All levels of brain from all animals were clear.

### 35 **Cognition models**

Following the 4 month dosing of 4 month old TASTPM mice as described above, these mice were tested in two models of cognition: the object recognition assay and the fear conditioning assay.

### 40 **Object recognition assay**

The object recognition assay exploits the animals' natural propensity to explore novel objects and relies on the animals' ability to recall an object which had been explored previously (familiar object). Eight month old TASTPM mice have been reported to demonstrate a deficit in the ability to distinguish between novel and familiar objects

(Howlett et al., 2004) indicating impaired cognitive performance in these animals. In this study, however, 8 month-old TASTPM mice treated with vehicle failed to demonstrate cognitive impairment i.e. they were able to distinguish between novel and familiar objects. There was therefore no window to investigate any potential therapeutic effect resulting from treatment with 2E7.

#### Fear conditioning assay

The fear conditioning model was designed to test the animals' ability to correlate a previous painful stimulus with a contextual or cued signal and recall this when presented with the same context or tone following Xh delay. In this study 8-month old TASTPM mice treated with vehicle (once or twice weekly) exhibited a deficit in contextual differentiation indicative of cognitive impairment in these animals. This deficit was unaffected by treatment with 2E7 when administered once or twice weekly

15

#### 4 Month dosing of 6 month old TASTPM mice

This study involved the administration of 2E7 (300ug i.p. twice weekly) to TASTPM mice for 4 months, starting at 3 months of age. Control animals received IgG2A in PBS. As described above, brains were dissected and hemisected. The right hemisphere was fixed by immersion in 4% paraformaldehyde and processed for histology. The left hemisphere was collected into pre-weighed 2ml eppendorf™ tubes, frozen on dry ice and stored at -80°C for subsequent analysis of amyloid content.

A preliminary analysis of a single section from each of a random selection of brain samples (n=6 vehicle, n=7 2E7 treated group) by IHC was undertaken using the same general protocol as above. Statistical analysis (Student's t-test) shows that there was a significant decrease in Aβ42 plaque load in thalamus (71.9%, p=0.007) and in thalamus + cortex + hippocampus (54.1%, p=0.022) in mice dosed with 2E7 but no significant change in Aβ40.

For biochemical measurement of brain Aβ40 and Aβ42, samples were processed and measured as above (dilution factor 1:10,000). Aβ42 was significantly decreased (p=0.01) by 29.9% in mice dosed with 2E7 (n=12 control, n=16 treated). Aβ40 concentrations were also decreased (22.6%) but this decrease failed to reach statistical significance (p=0.052).

#### Cloning of Hybridoma Variable Regions

##### Variable Region Sequences

Total RNA was extracted from 2E7 hybridoma cells and heavy and light variable domain cDNA sequences were then generated by reverse transcription and polymerase chain reaction (RT-PCR). The forward primer for RT-PCR was a mixture of degenerate primers specific for murine immunoglobulin gene leader-sequences

and the reverse primer was specific for the antibody constant regions, in this case murine isotype IgG2a for the heavy chain and murine kappa for the light chain. Primers were designed according to the strategy described by Jones and Bendig (Bio/Technology 9:88, 1991). RT-PCR was carried out in duplicate for both V-region sequences to enable subsequent verification of the correct V-region sequences. The V-region products generated by RT-PCR were cloned (Invitrogen TA Cloning Kit) and sequence data obtained.

2E7 V<sub>H</sub> Amino Acid Sequence (SEQ ID No:17)

10 EVKLVESGGGLVQPGGSLKLSCAVSGFTFSDNGMAWVRQAPRKGPEWIAFISNLA  
YSIDYADTVTGRFTISRDNAKNTLYLEMSSLRSEDTAMYYCVSGTWFAWGGQTLV  
 TVSA

2E7 V<sub>H</sub> DNA Sequence (SEQ ID No:18)

15 GAGGTGAAGCTGGTGGAGTCTGGGGGAGGCTTAGTGCAGCCTGGAGGGTCCC  
 TGAAACTCTCCTGTGCAGTCTCTGGATTCACTTTCAGTGACAACGGAATGGCGT  
 GGGTTCGACAGGCTCCAAGGAAGGGGCCTGAGTGGATAGCGTTCATTAGTAAT  
 TTGGCATATAGTATCGACTACGCAGACACTGTGACGGGCCGATTACCATCTCT  
 AGAGATAATGCCAAGAATACCCTGTACCTGGAAATGAGCAGTCTGAGGTCTGAG  
 20 GACACGGCCATGTACTATTGTGTAAGCGGGACCTGGTTTGCTTACTGGGGCCAA  
 GGGACTCTGGTCACTGTCTCTGCA

2E7 V<sub>L</sub> Amino Acid Sequence (SEQ ID No:19)

25 DVVLTQTPLSLPVSLGDQASISCRVSQSLLHSNGYTYLHWYLQKPGQSPKLLIYKVS  
NRFSGVPDRFSGSGSGTDFTLKISRVEAEDLGVYFCSQTRHVPYTFGGGTKLEIK

2E7 V<sub>L</sub> DNA Sequence ((SEQ ID No:20)

30 GATGTTGTGCTGACCCAACTCCACTCTCCCTGCCTGTCAGTCTTGGAGATCAA  
 GCCTCCATCTCTTGCAGAGTTAGTCAGAGCCTTTTACACAGTAATGGATACACCT  
 ATTTACATTGGTACCTGCAGAAGCCAGGCCAGTCTCCAAAGCTCCTGATCTACA  
 AAGTTTCCAACCGATTTTCTGGGGTCCCAGACAGGTTCAAGTGGCAGTGGATCAG  
 GGACAGATTTCACTCAAGATCAGCAGAGTGGAGGCTGAGGATCTGGGAGTT  
 TATTTCTGCTCTCAAACCTAGACATGTTCCGTACACGTTCCGAGGGGGGACCAAG  
 CTGGAAATAAAA

35

Complementarity Determining Regions (CDRs) are underlined in the amino acid sequences.

#### **Cloning and Expression of 2E7 Chimera**

40 A chimeric 2E7 antibody (2E7c) consisting of the parent murine V regions grafted on to human IgG1 (Fc mutated (L235A, G237A)) for the heavy chain or human C kappa regions for the light chain was generated in order to express recombinant antibody material that could be used to confirm the correct cloning of functional murine V regions. DNA encoding 2E7 murine heavy and light chain V regions and endogenous

murine signal sequences was cloned in frame into the mammalian expression vectors RLD-bshe (for the heavy chain) and RLN-bshe (for the light chain) already containing human constant regions (IgG1 Fc mutated (L235A, G237A) or human C kappa, respectively).

5

Elements of RLD-bshe expression vector for heavy chain expression:

Base Pairs	Description of DNA segment
0 - 1014	Promoter (SV40/RSV)
1015 - 2442	Antibody heavy chain
2443 - 2765	Poly A
2766 - 3142	BG Promoter
3239 - 3802	DHFR
3803 - 4162	Poly A
4163 - 6322	Total backbone
5077 - 5937 (complementary strand)	Beta lactamase

(position of elements and overall size of vector given above are for illustration purposes only and will depend upon the size of the antibody chain insert)

10 Elements of RLN-bshe expression vector for light chain expression:

Base Pairs	Description of DNA segment
0 - 1014	Promoter (SV40/RSV)
1015 - 1731	Antibody light chain
1732 - 2051	Poly A
2388 - 2764	BG Promoter
2774 - 3568	Neomycin
3569 - 3876	Poly A
3877 - 6063	Total backbone
5077 - 5937 (complementary strand)	Beta lactamase

(position of elements and overall size of vector given above are for illustration purposes only and will depend upon the size of the antibody chain insert)

15 Clones with correctly cloned  $V_H$  and  $V_L$  sequences were identified and plasmids prepared for expression in suspension culture CHO cells. Expressed 2E7c antibody was purified from cell culture supernatant by protein A chromatography on a FPLC system, and then tested for binding to A $\beta$  by ELISA and SPR using Biacore™ technology. The results indicated that the correct 2E7 mouse V regions were cloned and expressed, resulting in a functional antibody with similar characteristics to the parent murine antibody 2E7.

20

### Light Chain Humanisation

A human acceptor sequence with the Genpept ID CAA51135 (SEQ ID No:24) and Genbank Accesion No X72467, which had 77% identity on the amino acid level (including CDRs) was selected as the acceptor framework. Construct L1 is a graft of the murine CDRs from the 2E7 VL domain into this acceptor framework.

25

**Heavy Chain Humanisation**

Human sequence Genbank accession No M99675 (SEQ ID No:21) an allele of the VH3-48 gene with 74% identity on the amino acid level (including CDRs 1 and 2) to the 2E7 mouse variable heavy region was selected as the human heavy chain acceptor framework together with the human JH4 minigene. Three humanised variable heavy chain variants were designed based on the M99675 sequence and JH4. H1 is a graft of the murine CDRs using the Kabat definition with two additional framework back mutations at positions 93 and 94. H2 and H3 were both derived from H1, but incorporated one additional framework mutation which were different in each construct; (positions 24 and 48 respectively; see Table 4).

**Table 4**

<b>Construct</b>	<b>Template frameworks</b>	<b>Residue (Kabat#)</b>	<b>Human</b>	<b>Mouse</b>
<b>H1</b>	M99675 and JH4	93,94	A and R respectively	V and S respectively
<b>H2</b>	H1	24	A	V
<b>H3</b>	H1	48	V	I

**Construction of Humanised Heavy and Light Chain DNA**

Humanised V regions were synthesised *de novo* by build up of overlapping oligos and PCR amplification. Restriction sites for cloning into mammalian expression vectors RLD-bshe and RLN-bshe and human immunoglobulin signal sequences derived from the chosen human acceptor frameworks were included. The DNAs encoding the humanised V regions (H1 (SEQ ID NO:27), H2 (SEQ ID NO:29), H3 (SEQ ID NO:31), L1 (SEQ ID NO:33)) together with signal sequences and restriction sites were then cloned in frame into mammalian expression vectors: H1, H2 and H3 into RLD-bshe to generate DNA encoding three full length human IgG1 Fc mutated heavy chains each containing mutations L235A and G237A, full length H1 (SEQ ID NO:35), full length H2 (SEQ ID NO:37) and full length H3 (SEQ ID NO:39); L1 was cloned in frame into RLN-bshe containing the DNA encoding human kappa constant region to generate DNA encoding a full length human kappa light chain (SEQ ID NO:41).

**Representative Examples of Expression of Humanised Heavy and Light Chain****Antibody Combinations**

CHOK1 cells were transiently transfected at small scale with all combinations of humanised light and heavy chain DNA constructs: L1+H1, L1+H2, L1+H3 (SEQ ID Nos: 35 + 41, 37 + 41, 39 + 41) in 6-well plates. CHOK1 cells passaged in DMEM F12, with 5% ultra low IgG foetal bovine serum and 2mM glutamine were grown to confluency in 6-well plates. The confluent cells were transfected with a total of 7.5 µg DNA: 30 µg Transfast lipid (Promega) in Optimem Glutamax medium (Invitrogen). Transfected cells were incubated at 37°C with 5% CO<sub>2</sub>. At 72 hours supernatants were harvested and assayed for antibody concentration and then tested for binding



to human A $\beta$  by ELISA. Humanized L1 combined with the three humanized heavy chains all expressed complete antibody that bound to human A $\beta$ .

Humanized antibodies were also expressed in large scale transient CHOK1 cell transfections using liposomal delivery of DNA (eg TransFast (Promega)) and expression in culture bottles. For optimization of expression levels in transient transfections a heavy to light chain expression vector DNA ratio of 1:6 was used. Material from transient transfection was purified using ProSepA columns or FPLC with ProSepA HiTrap columns.

### **Assessment of 2E7 Humanised Variants H1L1, H2L1 and H3 L1 in $\beta$ -Amyloid Binding ELISA**

2E7 H1L1, H2L1 and H3L1 humanised variants were assessed for binding to human A $\beta$  peptide (1-40) biotinylated at the C terminus. The chimeric 2E7 was used as a reference. Tables 5-7 show results with various batches of purified material from large scale transient transfections.

**Table 5**

ELISA	MAb	EC <sub>50</sub> ( $\mu$ g/ml)	Standard Error
<b>A<math>\beta</math> binding</b>	2E7c Chimera	0.033	0.00144
	H1L1	0.035	0.00142
	H2L1	0.048	0.00202
	H3L1	0.044	0.00105

**Table 6**

ELISA	MAb	EC <sub>50</sub> ( $\mu$ g/ml)	Standard Error
<b>A<math>\beta</math> binding</b>	2E7c Chimera	0.043	0.00183
	H1L1	0.051	0.00164
	H2L1	0.044	0.00191
	H3L1	0.055	0.00094

**Table 7**

ELISA	MAb	EC <sub>50</sub> ( $\mu$ g/ml)	Standard Error
<b>A<math>\beta</math> binding</b>	2E7c Chimera	0.044	0.00169
	H1L1	0.047	0.00265
	H2L1	0.041	0.00174
	H3L1	0.040	0.00116

These results indicated very similar A $\beta$  binding profiles for each of the 2E7-derived humanised variants. Comparison of the EC<sub>50</sub> values to the 2E7c showed little loss of A $\beta$  binding activity had been incurred through the humanization process.

### **Comparison of 2E7 Humanised Variants by Competition ELISA**

2E7c chimeric and humanised antibodies H1L1, H2L1 and H3L1 were assessed for their ability to inhibit the binding between the human A $\beta$  peptide and the parental mouse 2E7 MAb in a competition ELISA.

- 5 Two types of competition ELISA were established in order to compare the A $\beta$  binding activity of the three humanised variants compared to the 2E7 chimeric antibody.
- 1) *Immobilised  $\beta$ -amyloid*; biotinylated human A $\beta$  peptide (1-40) was immobilized through Streptavidin on ELISA plates. Mouse 2E7 antibody was added at a constant concentration along with a dilution series of 2E7-derived humanised variant
- 10 antibodies. Bound mouse 2E7 MAb was then detected with anti-mouse IgG conjugate. Table 8 shows results of two assays.

Table 8

Competitor MAb	Experiment 1		Experiment 2	
	IC <sub>50</sub> ( $\mu$ g/ml)	Standard Error	IC <sub>50</sub> ( $\mu$ g/ml)	Standard Error
2E7c Chimera	1.31	0.20	1.29	0.13
H1L1	1.62	0.40	1.76	0.21
H2L1	1.28	0.26	1.66	0.28
H3L1	1.53	0.16	1.39	0.23

- 15 2)  *$\beta$ -amyloid in solution*; a constant concentration of  $\beta$ -amyloid was pre-incubated with a dilution series of humanised 2E7 antibody variants – the mixture including complexed and free amyloid was added for a short time to wells containing immobilised mouse 2E7 MAb. The amount of free  $\beta$ -amyloid that was still available for binding the immobilised parental 2E7 MAb was then detected. Table 9 shows
- 20 results of two assays.

Table 9

Competitor MAb	Experiment 1		Experiment 2	
	IC <sub>50</sub> ( $\mu$ g/ml)	Standard Error	IC <sub>50</sub> ( $\mu$ g/ml)	Standard Error
2E7c Chimera	0.052	0.006	-	-
H1L1	0.114	0.014	0.140	0.024
H2L1	0.075	0.009	0.119	0.014
H3L1	0.069	0.004	0.115	0.013

- 25 All humanised antibody variants inhibited the binding of mouse 2E7 MAb to  $\beta$ -amyloid with a very similar profile. IC<sub>50</sub> values generated for H2L1 and H3L1 variants were consistently close to that of the 2E7c chimera (where used), which had the highest inhibitory activity in both assays. However, variant H1L1 showed a somewhat reduced inhibitory activity in both assays, indicating a possible slightly lower affinity for  $\beta$ -amyloid.

30

### SPR Biacore™ Analysis of 2E7, 2E7c, H1L1, H2L1, H3L1

The kinetics parameters of recombinant mouse 2E7 MAb, chimeric 2E7c and humanized variants H1L1, H2L1 and H3L1 binding to human beta-amyloid peptide (1-40) and (1-42) were assessed using Biacore™ analysis on a Biacore™ 3000. Two different assay formats were used.

5

#### Method A

(i) Briefly, <20 resonance units of beta-amyloid 1-40 peptide (biotinylated at the C-terminus) were captured on a streptavidin biosensor chip (as used for Table 10A). The antibodies were diluted down in HBS-EP buffer and passed over the streptavidin/beta-amyloid surface at concentrations ranging from 0.001nM-8nM (for Table 10A). Two separate runs were carried out; each run was carried out on a new streptavidin/beta-amyloid surface. Runs 1 and 2 were essentially the same though there were some differences in the parameters used; Run 1 was carried out using a chip surface on which 16 RU's of beta-amyloid were captured, and antibody concentrations of 0.001nM-8nM were used, an association time of 4 minutes and a dissociation time of 20 minutes were used at a flow rate of 50µl per minute. For Run 2, less than 10 RU's of beta-amyloid were captured and antibody concentrations of 0.003125nM-8nM were used. The flow rate and association times were the same as Run 1, however the dissociation time was reduced to 15 minutes.

(ii) Beta amyloid (1-40) and (1-42) were amine-coupled on different surfaces of a CM5 biosensor chip to a level of <20 resonance units (as used for Table 10B). The antibodies were diluted down in HBS-EP buffer and passed over the biosensor/beta-amyloid surface at concentrations ranging from 1nM-64nM (as used for Table 10B).

25

#### Method B

In the second instance the assay was reversed, in that antibodies were first captured to a level of 1000-2500 resonance units on an anti-mouse IgG polyclonal antibody surface (for recombinant mouse 2E7 MAb) or a protein A surface (for humanized H2L1) of a CM5 biosensor chip. Freshly prepared beta-amyloid (1-40) or (1-42) was diluted down in HBS-EP buffer and passed over the captured-antibody surface at concentrations ranging from 4-500nM (Table 10C and 10D).

30

In both methods, regeneration was via a pulse of 100mM H<sub>3</sub>PO<sub>4</sub>, and for Table 10A data also followed by a pulse of 50mM NaOH. The surface was shown to be stable and unaffected by regeneration. All runs were double referenced against buffer blank injections. Analysis was carried out using the Biacore™ analysis software BIAevaluation version 4.1.

35

#### Results

Method A(i) was used to rank order the antibodies by beta-amyloid binding kinetic data. The data obtained is shown in Table 10A. This shows that the parental 2E7 Mab has a KD of 36.1 pM for streptavidin-captured beta-amyloid. The chimeric mouse-human antibody showed a slightly reduced KD of 45.8 pM and the humanised

40

constructs range from 54 (H2L1) to 93.6 pM (H1L1). In conclusion this demonstrates that the humanisation procedure had been very successful and very little affinity had been lost. The additional backmutations introduced for H2 and H3 had a small but beneficial effect, although the differences between H2 and H3 constructs are within the standard deviations for these experiments.

Table 10A

Antibody		<i>ka</i>	<i>kd</i>	KD(pM)
2E7	Run 1	1.61e6	6.17e-5	38.3
	Run 2	1.69e6	5.72e-5	33.8
	Average(SD)	1.65e6	5.97e-5	36.1(3.2)
c2E7	Run 1	1.34e6	6.44e-5	48.1
	Run 2	1.3e6	5.65e-5	43.5
	Average(SD)	1.32e6	6.10e-5	45.8(3.3)
H1L1	Run 1	5.60e5	5.32e-5	95.0
	Run 2	6.37e5	5.87e-5	92.2
	Average(SD)	5.99e5	5.60e-5	93.6(2.0)
H2L1	Run 1	9.91e5	5.76e-5	58.1
	Run 2	1.1e6	5.49e-5	49.8
	Average(SD)	1.05e6	5.63e-5	54.0(5.9)
H3L1	Run 1	8.24e5	6.26e-5	76.0
	Run 2	8.3e5	4.75e-5	57.2
	Average(SD)	8.27e5	5.47e-5	66.6(13.3)

Method A(ii) was used to confirm that the additional two amino-acid residues on the C-terminus of beta-amyloid (1-42) compared to beta-amyloid (1-40) did not significantly alter the binding properties of 2E7 and H2L1. The data obtained is shown in Table 10B and did confirm this.

Table 10B

Antibody	Beta-amyloid	<i>ka</i> (Ms <sup>-1</sup> )	<i>kd</i> (s <sup>-1</sup> )	KD (pM)
2E7	1-40	4.05e5	1.28e-4	317
	1-42	3.82e5	1.51e-4	394
H2L1	1-40	3.33e5	1.22e-4	366
	1-42	3.40e5	1.55e-4	456

Method B was used to negate avidity effects potentially seen in the first assay format. Avidity effects, caused by both Fab domains of a single antibody molecule binding at the same time to two adjacent beta-amyloid molecules on the biosensor surface (or

in multimeric forms of beta-amyloid), would increase the apparent affinity of binding. Affinity measurements obtained using Method B are shown in Table 10C.

Table 10C

Antibody	$k_a$ ( $\text{Ms}^{-1}$ )	$k_d$ ( $\text{s}^{-1}$ )	KD (nM) With Standard Deviation n=3
2E7	2.83e5 $\pm 0.54\text{e}5$	4.28e-4 $\pm 0.65\text{e-}4$	1.58 $\pm 0.55$
H2L1	1.06e5 $\pm 0.27\text{e}5$	7.50e-4 $\pm 0.50$	7.32 $\pm 1.64$

5

Evidence that this assay provided true 1:1 binding affinities was obtained when Fab fragments of H2L1, obtained by papain digestion, bound streptavidin-captured beta-amyloid (1-40) by a similar method to Method A(i) with an estimated KD of 2.4nM.

10 Method B was also used to confirm that the additional two amino-acid residues on the C-terminus of beta-amyloid (1-42) compared to beta-amyloid (1-40) did not significantly alter the binding properties of an identical sequence clone to mouse 2E7 MAb, named 2F11. The data obtained is shown in Table 10D.

15

Table 10D

Antibody	Beta-amyloid	$k_a$ ( $\text{Ms}^{-1}$ )	$k_d$ ( $\text{s}^{-1}$ )	KD (nM)
2F11	1-42	2.39e5	2.74e-4	1.14
2F11	1-40	2.99e5	3.92e-4	1.31

20 In a study similar to the epitope mapping study on 2E7 using the Surface Plasmon Resonance assay described above, H2L1 behaved similarly to 2E7 in binding to the peptide encompassing amino acids 1-12 ( $\text{A}\beta_1$ , SEQ ID No:15) of the  $\beta$ -amyloid peptide and not to the peptide encompassing amino acids 2-13 of the  $\beta$ -amyloid peptide ( $\text{A}\beta_{2-13}$ , SEQ ID No:44).

#### Activity of H2L1 in the $\text{I}^{125}$ $\beta$ -Amyloid Efflux Model

25 In order to functionally compare the humanised H2L1 with the parent mouse monoclonal 2E7, both were tested on the same day in the  $\text{I}^{125}$   $\beta$ -amyloid efflux model described above.

30 Both H2L1 and 2E7 significantly increased counts per minute (CPM) in blood compared with vehicle control. CPM of radioactivity in blood was as follows (Vehicle:  $1940 \pm 166$ ; 2E7:  $10065 \pm 1386$ ; H2L1:  $10913 \pm 1535$ ). Statistics used were ANOVA

with post-hoc LSD test. n=7 vehicle, n=6 2E7, n=6 H2L1, (p<0.001 for each test compound vs. vehicle).

5 This data provides further evidence that the humanised H2L1 antibody has retained the functional properties shown with the mouse 2E7 molecule.

**Investigation of the Pharmacokinetics of H2L1 and 2E7**

10 The terminal half life of test antibody in mice was investigated. Test antibody was administered by a 1 h intravenous infusion to 4 mice to achieve a target dose of 400ug per mouse. Serial blood samples were taken from each mouse up to 5 days after dosing (one mouse from the 2E7 group did not complete the study and one from the H2L1 group was removed from subsequent analysis because it became apparent the dose had not been administered i.v.). Antibody levels were measured using a  $\beta$ -amyloid capture ELISA.

15 Analysis of the data indicates that the humanised antibody H2L1 has a terminal half life of circa 82 hours in mice (Table 11), which is comparable to that of the parent mouse monoclonal antibody 2E7 (circa 75 hours).

20

**Table 11**

Parameter	Mean $\pm$ SD (n=3)
Cmax (ug/mL)	291 $\pm$ 43
Tmax (h) #	2.0 (1.1-2.1)
CLp (mL/h/kg)	0.9 $\pm$ 0.1
t $\frac{1}{2}$ (h)	82 $\pm$ 4
Vss (mL/kg)	94 $\pm$ 12

# median and range

Cmax Observed maximum plasma concentration.

Tmax Time of the observed maximum plasma concentration

CLp Total plasma clearance; Dose/AUC<sub>(0-inf)</sub>.

25

t $\frac{1}{2}$  Terminal phase half-life was determined as the ratio of ln2/z where z is the terminal phase rate constant; calculated using unweighted linear regression analysis (after log transformation) on those concentration-time pairs occurring after the visually assessed onset of the terminal log-linear phase.

Vss Volume of distribution at steady-state; CLp x MRT<sub>0-inf</sub>.

30

**Effect of H2L1 on peripheral amyloid load in aged non-human primates**

35 A study was conducted in aged Cynomolgus monkeys (approximately 15 years old) to investigate the exposure response relationship with respect to amyloid/H2L1 complex formation and clearance and the subsequent effects on CSF and CNS amyloid levels. Weekly lumbar CSF (taken under ketamine sedation) and blood samples were collected 3 weeks prior to 1<sup>st</sup> dose of H2L1. Immediately following sampling on week 3, animals received placebo (n=10), 0.1mg/kg (n=5), 1mg/kg (n=5) or 10mg/kg (n=10) H2L1 and then every 2 weeks for 12 weeks. Blood samples for

- plasma analysis of H2L1 and total A $\beta_{42}$  were taken weekly. CSF samples for quantification of A $\beta_{40/42}$  were collected every 2 weeks. Following completion of the dosing period, animals were euthanased for the purpose of brain quantification of beta-amyloid by biochemical analyses as described above and investigation of microhaemorrhage. In the lowest dose group (0.1 mg/kg), animals were euthanased in a staggered fashion to evaluate the potential time course effect in brain levels as a consequence of termination of dosing and hence saturation of the plasma amyloid pool.
- 5
- 10 This study was approved by the Institutional Animal Care and Use Committee (IACUC) of MACCINE Pte Ltd, or "Maccine" prior to start of the experimental phase. The IACUC protocol number was #08-2006. GSK has performed a site visit of Maccine and has reviewed their ethical review process and found it acceptable
- 15 Plasma samples were serially diluted 1:10 to 1:50000 and added to A $\beta_{40}$  coated ELISA plates. Standard curves were created ranging from 0–10 $\mu$ g/ml H2L1 in diluent. Following an overnight incubation at 4°C H2L1 was visualised using anti-human IgG horseradish peroxidase (Amersham – diluted 1:2000 in diluent) and tetramethylbenzidine detection system. Following single and repeat iv bolus
- 20 administration, plasma levels of H2L1 appeared to increase in a dose dependent fashion. There was no evidence of severe non-linearities in the pharmacokinetics, indicating that for the majority of the dosing interval, excess molar concentrations of H2L1 in the plasma compared with free amyloid levels were achieved.
- 25 Total A $\beta_{42}$  was measured in neat plasma using a commercially available A $\beta$  1-42 ELISA kit (Innogenetics) in accordance with the manufacturers instructions, with standard curves ranging from 500-7pg/ml created in kit diluent. Samples and standards are incubated overnight at 4°C before assaying in duplicate according to kit instructions. It should be noted that due to the interference of the detection
- 30 antibody supplied with the A $\beta_{42}$  assay, this kit cannot be used to measure free A $\beta_{42}$  levels but measures the apparent 'total' A $\beta_{42}$ . There was a dose and concentration dependent increase in A $\beta_{42}$  (with plateau levels of approximately 300, 125 and 25 pg/ml detected following 10, 1 or 0.1mg/kg H2L1 respectively). From the analysis, the increase in the "total A $\beta_{42}$ " is likely to be due to the result of a
- 35 significant efflux of amyloid from outside the plasma pool, that appeared dependent upon H2L1 concentrations >1 ug/mL, and did not appear to be a result of lack of clearance of complex. This was evident by the elimination rate of the total A $\beta_{42}$  as well as the fluctuation in the total levels over a dosing interval.
- 40 To date only the plasma analysis has been completed and fully analysed. However preliminary analysis indicates that there was a trend towards reduction in CSF and increase in the hippocampal level of 'total' A $\beta_{42}$  (measured as generally described above) following treatment with 10mg/kg H2L1.

In some brain sections, small areas of microhaemorrhage were detected as shown by the Perls' Prussian Blue staining method. This method visualises ferric iron (iron is an essential constituent of the oxygen-carrying haemoglobin found in red cells) by producing an insoluble blue compound. However there was no difference in the incidence between vehicle and drug treated animals.

#### **Analyses on Aged Cynomolgus macaque monkeys for beta amyloid plaques in the brain and total beta amyloid in the plasma**

Cerebral spinal fluid (CSF) and tissue parameters of human AD have been displayed in the cynomolgus monkey. The aged cynomolgus monkey has been shown to have evidence of amyloid deposition. (Covance, The cynomolgus monkey as a model for Alzheimer's disease. In: Buse E, Habermann G, Friderichs-Gromoll S, Kaspereit J, Nowak P and Niggemann K, editors. Poster Presentation at the 44th Annual Meeting of the Society of Toxicology, New Orleans, Louisiana, 6 to 10 March 2005). The potential for H2L1 to elicit an inappropriate response (such as encephalitis) in an aged brain was investigated in old, ca. 20 years, ex-breeding female monkeys. In addition, safety, treatment-related microhaemorrhage, neutralization/clearance of test material, hypersensitivity, and immune complex disease were also investigated following intravenous administration for 8 weeks in two-weekly intervals. In addition CNS and blood samples were analysed for levels of A $\beta_{40/42}$ .

#### **Study Design**

Groups of 5 (group 1), 9 (group 2) or 10 (group 3) geriatric female cynomolgus monkeys were given 0 (vehicle), 50 or 100 mg/kg/dosing day H2L1 in vehicle (4 ml/kg) every second week for 8 weeks intravenously by slow bolus administration. The vehicle consisted of sodium acetate trihydrate 6.81 mg/mL, disodium edetate dehydrate 0.0186 mg/mL, polysorbate 80 0.2 mg/mL, and L-Arginine base 10 mg/mL, the pH was 5.5. Dose levels were chosen to investigate dose levels that were 5 and 10 fold intended clinical dose levels.

The following evaluations were performed pre-dose, daily (clinical signs, body weight, food consumption), week 4 and the week before necropsy: in-life animal observations, body weight, body temperature, haematology, clinical chemistry (including cerebrospinal fluid [CSF] analysis), urinalysis, and cytokine determination in CSF. Following necropsy, organ weights, macroscopic observations, and microscopic observations of the brain, cervical spinal cord and gross lesions were conducted on all animals. Toxicokinetic evaluation was performed after each dosing.

#### **Results**

There were no unscheduled deaths, and there were no signs which indicated an influence of the test item on the general condition of the animals at the administered doses. The only remarkable observations in clinical pathology (hematology and serum chemistry) were concluded to be age- and not test article related.



Systemic exposure to H2L1 (as measured by  $AUC_{0-\tau}$  and  $C_{max}$ ) increased approximately in proportion to dose. For both dose groups, there was no marked change ( $\geq 2$ -fold) in systemic exposure between the 1st dose and 4th dose sampling periods.

5

There were no signs of inflammatory reactions in the brain detected by CSF-analysis, and there were no macroscopic or microscopic findings at necropsy that suggested a test item influence, specifically no microhaemorrhage or encephalitis.

- 10 This study was conducted in compliance with the Good Laboratory Practice Regulations as outlined in German Chemical Law, annex 1 and 2 to §19a Chemikalien Gesetz, June 2002, the OECD Principles of Good Laboratory Practice (revised 1997, issued January 1998) ENV/MC/CHEM (98) 17, the Consensus Document "The Application of the OECD Principles of GLP to the Organisation and Management of Multi-Site Studies" ENV/JM/MONO(2002)9. Studies conducted in compliance with the above regulations and standards were considered acceptable to US FDA regulatory authorities.

#### Analysis of Plaque load in the CNS

- 20 The left brain hemispheres of the vehicle treated cynomolgus macaque monkeys from the above study were analysed by immunohistochemistry. A coronal section, at the level of the middle temporal sulcus containing portions of the dentate gyrus and hippocampus, was processed into wax as described above. For immunohistochemistry, sections were labelled with a pan-A $\beta$  antibody (1E8, 25 monoclonal antibody raised to A $\beta$  13-27), or with the A $\beta$ 42 antibody, (20G10, monoclonal antibody recognising A $\beta$  x-42), and labelling was developed as above. A visual count of the number of plaques was taken for each section. Tissue from all five vehicle-treated cynomolgus monkeys showed evidence of parenchymal A $\beta$  plaques. There was also evidence of cerebrovascular labelled A $\beta$  and intraneuronal A $\beta$ .

30

#### Analysis of beta amyloid/antibody complexes in plasma

- Biochemical analysis was carried out on plasma samples from two time points (at the end of weeks 4 and 8 after start of dosing) from animals dosed with 50mg/kg (n=9) or 100mg/kg (n=10) H2L1, or vehicle dosed controls (n=5). 100ul duplicate samples were analysed using the commercially available Innogenetics A $\beta$  1-42 ELISA kit, incubated overnight at 4°C. Control samples were analysed both neat and at 1:10 dilution (using the supplied diluent), while samples from the dosed animals were tested neat and at 1:25. Subsequent absorbance values were analysed, with unknown absorbance values backcalculated to pg/ml values using standard curves, and then corrected for any assay dilution. Total plasma levels of A $\beta$ 42 derived from these samples are shown in Table 12 below (figures in pg/ml  $\pm$  SE); all samples from animals treated with H2L1 contained significantly higher levels of A $\beta$ 42 ( $p < 0.001$  by student t-test) than in control groups.

40

Table 12

	Week 4 (pg/ml)	Week 8 (pg/ml)
Control (1:10)	104.1 ± 30.4	29.8 ± 7.9
50mg/kg (1:25)	830.5 ± 79.1	615.8 ± 50.2
100mg/kg (1:25)	1020.5 ± 84.4	492.7 ± 46.3

Data reported were obtained from diluted samples. Results from the neat samples were not used as many data points were either greater than the top standard, or due to sample volume limitation, only assayed as a single point.

5

### **Production process**

Expression vectors encoding H2L1 and operatively linked to amplifiable selection markers such as the DHFR or glutamine synthetase may be used to transfect or transduce appropriate parental CHO cell lines (eg CHODG44 or CHOK 1) to produce engineered cell lines suitable for production of monoclonal antibody at large scale (for review see Bebbington and Hentschel DNA Cloning Volume III; A practical approach (edited by Glover DM) (Oxford IRL press, 1987). In order to increase expression levels the coding sequence maybe codon optimized in order to avoid cis-acting sequence motifs and extreme GC contents (high or low). SEQ. ID Nos:42 and No:43 exemplify such a coding sequence for H2 heavy chain and L1 light chain. Large scale production maybe in stirred tank bioreactors using animal-derived-component-free medium, followed by purification. This may comprise clarification of the harvest, followed by Protein-A affinity chromatography, and further purification using ion (e.g. cation) exchange and mixed mode (e.g. ceramic hydroxyapatite) chromatography unit operations. A virus removal nanofiltration is followed by a final ultrafiltration/diafiltration step that enables formulation suitable for the intended route of administration.

25

### **Example of Pharmaceutical formulation**

<u>Ingredient</u>	<u>Quantity (per mL)</u>
H2L1	50mg
Sodium acetate trihydrate	6.81mg
Polysorbate 80	0.20mg
Arginine base	10.00mg
Sodium chloride	3.00mg
Disodium edetate dihydrate	0.0186mg
Hydrochloric acid	qs to give pH 5.5
Water for Injections	To make 1.0mL
Nitrogen	To fill headspace

40

### **Amino Acid Sequences of V-regions of Acceptor Frameworks and Humanised Variants**

**M99675 heavy chain acceptor framework V region, amino acid sequence (SEQ ID No:21)**

EVQLVESGGGLVQPGGSLRLSCAASGFTFSSYSMNWVRQAPGKGLEWVSYISSSS  
 5 STIYYADSVKGRFTISRDNKNSLYLQMNSLRAEDTAVYYCAR

**M99675 heavy chain acceptor framework V region DNA (SEQ ID No:22)**

GAGGTGCAGCTGGTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCC  
 TGAGACTCTCCTGTGCAGCCTCTGGATTCACCTTCAGTAGCTATAGCATGAACT  
 10 GGGTCCGCCAGGCTCCAGGGAAGGGGCTGGAGTGGGTTTCATACATTAGTAGT  
 AGTAGTAGTACCATATACTACGCAGACTCTGTGAAGGGCCGATTACCATCTCC  
 AGAGACAATGCCAAGAACTCACTGTATCTGCAAATGAACAGCCTGAGAGCCGAG  
 GACACGGCTGTGTATTACTGTGCGAGAGA

**CAA51135 light chain acceptor framework V region amino acid sequence (SEQ ID No:24)**

DIVMTQSPLSLPVTPGEPASISCRSSQSLLSHNGYNYLDWYLQKPGQSPQLLIYLG  
 15 NRASGVPDFRFSGSGSGTDFTLKISRVEAEDVGVYYCMQALQTPWTFGQGTKVEIK

**CAA51135 light chain acceptor framework V region DNA (SEQ ID No:25)**

GATATTGTGATGACTCAGTCTCCACTCTCCCTGCCCGTCACCCCTGGAGAGCCG  
 GCCTCCATCTCCTGCAGGTCTAGTCAGAGCCTCCTGCATAGTAATGGATAACA  
 TATTTGGATTGGTACCTGCAGAAGCCAGGGCAGTCTCCACAGCTCCTGATCTAT  
 TTGGGTTCTAATCGGGCCTCCGGGGTCCCTGACAGGTTCACTGGCAGTGGATC  
 25 AGGCACAGATTTTACACTGAAAATCAGCAGAGTGGAGGCTGAGGATGTTGGGG  
 TTTATTACTGCATGCAAGCTCTACAACTCCGTGGACGTTCCGGCCAAGGGACCA  
 AGGTGGAAATCAAA

**Humanised heavy chain V region variant H1 , amino acid sequence (SEQ ID No:26)**

EVQLVESGGGLVQPGGSLRLSCAASGFTFSDNGMAWVRQAPGKGLEWVFSISNL  
 30 AYSIDYADTVTGRFTISRDNKNSLYLQMNSLRAEDTAVYYCVSGTWFAFWGQGL  
 VTVSS

**Humanised heavy chain V region variant H1 DNA coding sequence (SEQ ID No:27)**

GAGGTGCAGCTGGTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCC  
 TGAGACTCTCCTGTGCAGCCTCTGGATTCACCTTCAGTGACAACGGAATGGCGT  
 GGGTCCGCCAGGCTCCAGGGAAGGGGCTGGAGTGGGTTTCATTATTAGTAAT  
 40 TTGGCATATAGTATCGACTACGCAGACACTGTGACGGGCCGATTACCATCTCC  
 AGAGACAATGCCAAGAACTCACTGTATCTGCAAATGAACAGCCTGAGAGCCGAG  
 GACACGGCTGTGTATTACTGTGTCAGCGGGACCTGGTTTGCTTACTGGGGCCA  
 GGGCACACTAGTCACAGTCTCCTCA

**Humanised heavy chain V region variant H2 , amino acid sequence (SEQ ID No:28)**

EVQLVESGGGLVQPGGSLRLSCAIVSGFTFSDNGMAWVRQAPGKGLEWVSFISNL  
AYSIDYADTVTGRFTISRDNKNSLYLQMNSLRAEDTAVYYCVSGTWFAYWGQGL  
 5 VTVSS

**Humanised heavy chain V region variant H2 DNA (SEQ ID No:29)**

GAGGTGCAGCTGGTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCC  
 TGAGACTCTCCTGTGCAGTCTCTGGATTCACCTTCAGTGACAACCGAATGGCGT  
 10 GGGTCCGCCAGGCTCCAGGGAAGGGGCTGGAGTGGGTTTCATTCATTAGTAAT  
 TTGGCATATAGTATCGACTACGCAGACACTGTGACGGGCCGATTCACCATCTCC  
 AGAGACAATGCCAAGAAGTCACTGTATCTGCAAATGAACAGCCTGAGAGCCGAG  
 GACACGGCTGTGTATTACTGTGTCAGCGGGACCTGGTTTGCTTACTGGGGCCA  
 GGGCACACTAGTCACAGTCTCCTCA  
 15

**Humanised heavy chain V region variant H3 , amino acid sequence (SEQ ID No:30)**

EVQLVESGGGLVQPGGSLRLSCAASGFTFSDNGMAWVRQAPGKGLEWISFISNLA  
YSIDYADTVTGRFTISRDNKNSLYLQMNSLRAEDTAVYYCVSGTWFAYWGQGLV  
 20 TVSS

**Humanised heavy chain V region variant H3 DNA (SEQ ID No:31)**

GAGGTGCAGCTGGTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCC  
 TGAGACTCTCCTGTGCAGCCTCTGGATTCACCTTCAGTGACAACCGAATGGCGT  
 25 GGGTCCGCCAGGCTCCAGGGAAGGGGCTGGAGTGGATCTCATTTCATTAGTAAT  
 TTGGCATATAGTATCGACTACGCAGACACTGTGACGGGCCGATTCACCATCTCC  
 AGAGACAATGCCAAGAAGTCACTGTATCTGCAAATGAACAGCCTGAGAGCCGAG  
 GACACGGCTGTGTATTACTGTGTCAGCGGGACCTGGTTTGCTTACTGGGGCCA  
 GGGCACACTAGTCACAGTCTCCTCA  
 30

**Humanised light chain V region variant L1 amino acid sequence (SEQ ID No:32)**

DIVMTQSPLSLPVTPGEPASISCRVSQSLLSNGYTYLHWYLQKPGQSPQLLIYKVS  
NRFSQVPDRFSGSGSGTDFTLKISRVEAEDVGVYYCSQTRHVPYTFGGGTKVEIK

**Humanised light chain V region variant L1 DNA (SEQ ID No:33)**

GATATTGTGATGACTCAGTCTCCACTCTCCCTGCCCGTCACCCCTGGAGAGCCG  
 GCCTCCATCTCCTGCAGAGTTAGTCAGAGCCTTTTACACAGTAATGGATACACC  
 TATTTACATTGGTACCTGCAGAAGCCAGGGCAGTCTCCACAGCTCCTGATCTAT  
 AAAGTTTCCAACCGATTTTCTGGGGTCCCTGACAGGTTTCAGTGGCAGTGGATCA  
 40 GGCACAGATTTTACACTGAAAATCAGCAGAGTGGAGGCTGAGGATGTTGGGGTT  
 TATTACTGCTCTCAAAGTACATGTTCCGTACACGTTCCGGCGGAGGGACCAAG  
 GTGGAAATCAAA

**Mature H1 heavy chain amino acid sequence (Fc mutated double mutation bold) (SEQ ID No:34)**

EVQLVESGGGLVQPGGSLRLSCAASGFTFSDNGMAWVRQAPGKGLEWVSFISNL  
AYSIDYADTVTGRFTISRDNAKNSLYLQMNSLRAEDTAVYYCVSGTWFAYWGQGT  
 5 VTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVH  
 TFPAYLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKHT  
 CPPCPAPELAGAPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDG  
 VEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISK  
 AKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTT  
 10 PPVLDSDGSFFLYSKLTVDKSRWQQGNVFCSSVMHEALHNHYTQKSLSLSPGK

**H1 Full length DNA (SEQ ID No:35)**

GAGGTGCAGCTGGTGGAGTCTGG  
 GGGAGGCTTGGTACAGCCTGGGGGGTCCCTGAGACTCTCCTGTGCAGCCTCTG  
 15 GATTCACCTTCAGTGACAACGGAATGG  
 CGTGGGTCCGCCAGGCTCCAGGGAAGGGGCTGGAGTGGGTTTCATTCATTAGT  
 AATTTGGCATATAGTATCGACTACGCA  
 GACTACTGTGACGGGCCGATTACCATCTCCAGAGACAATGCCAAGAACTCACTG  
 TATCTGCAAATGAACAGCCTGAGAGC  
 20 CGAGGACACGGCTGTGTACTGTGTCAGCGGGACCTGGTTTGCTTACTGGG  
 GCCAGGGCACACTAGTCACAGTCTCCT  
 CAGCCTCCACCAAGGGCCCATCGGTCTTCCCCCTGGCACCCCTCCTCCAAGAGC  
 ACCTCTGGGGGCACAGCGGCCCTGGGC  
 TGCCTGGTCAAGGACTACTTCCCCGAACCGGTGACGGTGTCTGCGTGGAACTCAGG  
 25 CGCCCTGACCAGCGGCGTGCACACCTT  
 CCCGGCTGTCTACAGTCTCAGGACTCTACTCCCTCAGCAGCGTGGTGACCG  
 TGCCCTCCAGCAGCTTGGGCACCCAGA  
 CCTACATCTGCAACGTGAATCACAAGCCCAGCAACACCAAGGTGGACAAGAAA  
 GTTGAGCCCAAATCTTGTGACAAA  
 30 CACACATGCCACCGTGCCCAGCACCTGAACTCGCGGGGGCACCGTCAGTCTT  
 CCTCTTCCCCCAAACCAAGGACAC  
 CCTCATGATCTCCCGGACCCCTGAGGTCACATGCGTGGTGGTGGACGTGAGCC  
 ACGAAGACCCTGAGGTCAAGTTCAACT  
 GGTACGTGGACGGCGTGGAGGTGCATAATGCCAAGACAAAGCCGCGGGAGGA  
 35 GCAGTACAACAGCACGTACCGTGTGGTC  
 AGCGTCCTCACCGTCTGACCAGGACTGGCTGAATGGCAAGGAGTACAAGTG  
 CAAGGTCTCCAACAAAGCCCTCCAGC  
 CCCCATCGAGAAAACCATCTCAAAGCCAAAGGGCAGCCCCGAGAACCACAGG  
 TGACACCCTGCCCCCATCCCGGGATG  
 40 AGCTGACCAAGAACCAGGTCAGCCTGACCTGCCTGGTCAAAGGCTTCTATCCCA  
 GCGACATCGCCGTGGAGTGGGAGAGC  
 AATGGGCAGCCGGAGAACAACTACAAGACCACGCCTCCCGTGCTGGACTCCGA  
 CGGCTCCTTCTCCTCTACAGCAAGCT

CACCGTGGACAAGAGCAGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGA  
 TGCATGAGGCTCTGCACAACCACTACA  
 CGCAGAAGAGCCTCTCCCTGTCTCCGGGTAAA

5 **Mature H2 heavy chain amino acid sequence, (Fc mutated double mutation bold) (SEQ ID No:36)**

EVQLVESGGGLVQPGGSLRLSCAVSGFTFSDNGMAWVRQAPGKGLEWVSFISNL  
AYSIDYADTVTGRFTISRDNAKNSLYLQMNSLRAEDTAVYYCVSGTWFA~~YWGQ~~TL  
 VTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVH  
 10 TFPVAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVKDKVEPKSCDKTHT  
 CPPCPAPELAGAPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDG  
 VEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISK  
 AKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTT  
 PPVLDSDGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPGK

15 **H2 Full length DNA (SEQ ID No:37)**

GAGGTGCAGCTGGTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCC  
 TGAGACTCTCCTGTGCAGTCTCTGGATT  
 CACCTTCAGTGACAACGGAATGGCGTGGGTCCGCCAGGCTCCAGGGAAGGGG  
 20 CTGGAGTGGGTTTCATTCATTAGTAATT  
 TGGCATATAGTATCGACTACGCAGACACTGTGACGGGCCGATTCACCATCTCCA  
 GAGACAATGCCAAGAAGTCACTGTAT  
 CTGCAAATGAACAGCCTGAGAGCCGAGGACACGGCTGTGTATTACTGTGTCAG  
 CGGGACCTGGTTTGCTTACTGGGGCCA  
 25 GGGCACACTAGTCACAGTCTCCTCAGCCTCCACCAAGGGCCCATCGGTCTTCC  
 CCCTGGCACCTCCTCCAAGAGCACCT  
 CTGGGGGCACAGCGGCCCTGGGCTGCCTGGTCAAGGACTACTTCCCCGAACC  
 GGTGACGGTGTTCGTGGAAGTCAAGCGCC  
 CTGACCAGCGGCGTGCACACCTTCCCGGCTGTCCTACAGTCCTCAGGACTCTA  
 30 CTCCCTCAGCAGCGTGGTGACCGTGCC  
 CTCCAGCAGCTTGGGCACCCAGACCTACATCTGCAACGTGAATCACAAGCCCA  
 GCAACACCAAGGTGGACAAGAAAGTTG  
 AGCCCAAATCTTGTGACAAAACCTCACACATGCCACCGTGCCCAGCACCTGAAC  
 TCGCGGGGGCACCGTCAGTCTTCCTC  
 35 TTCCCCCAAACCCAAGGACACCCTCATGATCTCCCGGACCCCTGAGGTCACA  
 TGCGTGGTGGTGGACGTGAGCCACGA  
 AGACCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGGAGGTGCATAATG  
 CCAAGACAAAGCCGCGGGAGGAGCAGT  
 ACAACAGCACGTACCGTGTGGTCAGCGTCCCTACCGTCCCTGCACCAGGACTGG  
 40 CTGAATGGCAAGGAGTACAAGTGCAAG  
 GTCTCCAACAAAGCCCTCCCAGCCCCCATCGAGAAAACCATCTCCAAAGCCAAA  
 GGGCAGCCCCGAGAACCACAGGTGTA  
 CACCCTGCCCCCATCCCGGGATGAGCTGACCAAGAACCAGGTGAGCCTGACCT  
 GCCTGGTCAAAGGCTTCTATCCCAGCG

ACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAACAACACTACAAGAC  
 CACGCCTCCCGTGCTGGACTCCGACGGC  
 TCCTTCTTCTCTACAGCAAGCTCACCGTGGACAAGAGCAGGTGGCAGCAGGG  
 GAACGTCTTCTCATGCTCCGTGATGCA  
 5 TGAGGCTCTGCACAACCACTACACGCAGAAGAGCCTCTCCCTGTCTCCGGGTA  
 AA

**Mature H3 heavy chain amino acid sequence (Fc mutated double mutation bold) (SEQ ID No:38)**

10 EVQLVESGGGLVQPGGSLRLSCAASGFTFSDNGMAWVRQAPGKGLEWISFISNLA  
YSIDYADTVTGRFTISRDNAKNSLYLQMNSLRAEDTAVYYCVSGTWFAYWGQGLV  
 TVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHT  
 FPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHT  
 CPPCPAPELAGAPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDG  
 15 VEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISK  
 AKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTT  
 PPVLDSDGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPGK

**H3 full length DNA (SEQ ID No:39)**

20 GAGGTGCAGCTGGTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCC  
 TGAGACTCTCCTGTGCAGCCTCTGGATTCACCTTCAGTGACAACGGAATGGCGT  
 GGGTCCGCCAGGCTCCAGGGAAGGGGCTGGAGTGGATCTCATTATTAGTAAT  
 TTGGCATATAGTATCGACTACGCAGACACTGTGACGGGCCGATTACCATCTCC  
 AGAGACAATGCCAAGAACTCACTGTATCTGCAAATGAACAGCCTGAGAGCCGAG  
 25 GACACGGCTGTGTATTACTGTGTCAGCGGGACCTGGTTTGCTTACTGGGGCCA  
 GGGCACACTAGTCACAGTCTCCTCAGCCTCCACCAAGGGCCCATCGGTCTTCC  
 CCCTGGCACCCCTCCTCCAAGAGCACCTCTGGGGGCACAGCGGCCCTGGGCTG  
 CCTGGTCAAGGACTACTTCCCCGAACCGGTGACGGTGTCTGGAAGTCAAGGCG  
 CCCTGACCAGCGGCGTGACACCTTCCCGGCTGTCTACAGTCTCAGGACTC  
 30 TACTCCCTCAGCAGCGTGGTGACCGTGCCCTCCAGCAGCTTGGGCACCCAGAC  
 CTACATCTGCAACGTGAATCACAAGCCCAGCAACACCAAGGTGGACAAGAAAGT  
 TGAGCCCAAATCTTGTGACAAAACCTCACACATGCCACCGTGCCAGCACCTGA  
 ACTCGCGGGGGCACCGTCAGTCTTCTTCTTCCCCCAAACCAAGGACACCC  
 TCATGATCTCCCGGACCCCTGAGGTACATGCGTGGTGGTGGACGTGAGCCAC  
 35 GAAGACCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGGAGGTGCATAA  
 TGCCAAGACAAAGCCGCGGGAGGAGCAGTACAACAGCACGTACCGTGTGGTCA  
 GCGTCTCACCGTCTGACCAGGACTGGCTGAATGGCAAGGAGTACAAGTGC  
 AAGGTCTCCAACAAAGCCCTCCAGCCCCATCGAGAAAACCATCTCCAAGCC  
 AAAGGGCAGCCCCGAGAACCACAGGTGTACACCCTGCCCCCATCCCGGGATGA  
 40 GCTGACCAAGAACCAGGTCAGCCTGACCTGCCTGGTCAAAGGCTTCTATCCCA  
 GCGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAACAACACTACAA  
 GACCACGCCTCCCGTGCTGGACTCCGACGGCTCCTTCTTCTCTACAGCAAGC  
 TCACCGTGGACAAGAGCAGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTG

ATGCATGAGGCTCTGCACAACCACTACACGCAGAAGAGCCTCTCCCTGTCTCCG  
GGTAAA

**Mature Light chain amino acid sequence (SEQ ID No:40)**

5 DIVMTQSPLSLPVTPGEPASISCRVSQSLLHSNGYTYLHWYLQKPGQSPQLLIYKVS  
NRFSGVPDRFSGSGSGTDFTLKISRVEAEDVGVYYCSQTRHVPYTFGGGTKVEIKR  
TVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVT  
EQDSKDYSLSTLTLTKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC

10 **L1 Full length DNA (SEQ ID No:41)**

GATATTGTGATGACTCAGTCTCCACTCTCCCTGCCCGTCACCCCTGGAGAGCCG  
GCCTCCATCTCCTGCAGAGTTAGTCAGAGCCTTTTACACAGTAATGGATACACC  
TATTTACATTGGTACCTGCAGAAGCCAGGGCAGTCTCCACAGCTCCTGATCTAT  
AAAGTTTCCAACCGATTTTCTGGGGTCCCTGACAGGTTCAAGTGGCAGTGGATCA  
15 GGCACAGATTTTACACTGAAAATCAGCAGAGTGGAGGCTGAGGATGTTGGGGTT  
TATTACTGCTCTCAAAGTACATGTTCCGTACACGTTCCGGCGGAGGGACCAAG  
GTGGAATCAAACGTACGGTGGCTGCACCATCTGTCTTCATCTTCCC GCCATCT  
GATGAGCAGTTGAAATCTGGAAGTGCCTCTGTTGTGTGCCTGCTGAATAACTTC  
TATCCAGAGAGGCCAAAGTACAGTGGAAAGTGGACAACGCCCTCCAATCGGG  
20 TAACTCCCAGGAGAGTGTACAGAGCAGGACAGCAAGGACAGCACCTACAGCC  
TCAGCAGCACCCCTGACGCTGAGCAAAGCAGACTACGAGAAACACAAAGTCTAC  
GCCTGCGAAGTACCCATCAGGGCCTGAGCTCGCCCGTCACAAAGAGCTTCAA  
CAGGGGAGAGTGT

25 **Optimised H2 heavy chain DNA (SEQ ID No:42)**

GAGGTGCAGCTGGTGGAGTCTGGCGGCGGACTGGTGCAGCCTGGCGGCAGCC  
TGAGACTGAGCTGTGCCGTGTCCGGCTTACCTTCAGCGACAACGGCATGGCC  
TGGGTGAGGCAGGCCCTGGCAAGGGCCTGGAGTGGGTGTCTTCATCAGCA  
ACCTGGCCTACAGCATCGACTACGCCGACACCGTGACCGGCAGATTCACCATC  
30 AGCCGGGACAACGCCAAGAACAGCCTGTACCTGCAGATGAACAGCCTGAGAGC  
CGAGGACACCGCGTGTACTACTGTGTGAGCGGCACCTGGTTCGCTACTGGG  
GCCAGGGCACCCCTGGTACCCTGTCCAGCGCCAGCACCAAGGGCCCCAGCGT  
GTTCCCCCTGGCCCCCAGCAGCAAGAGCACCAAGCGGCGGCACAGCCGCCCTG  
GGCTGCCTGGTGAAGGACTACTTCCCCGAACCGGTGACCGTGTCTGGAACAG  
35 CGGAGCCCTGACCAGCGGCGTGCACACCTTCCCCGCCGTGCTGCAGAGCAGC  
GGCCTGTACAGCCTGAGCAGCGTGGTACCCTGCCAGCAGCAGCCTGGGCA  
CCCAGACCTACATCTGTAACGTGAACCACAAGCCCAGCAACACCAAGGTGGAC  
AAGAAGGTGGAGCCCAAGAGCTGTGACAAGACCCACACCTGCCCCCCTGCCC  
TGCCCCGAGCTGGCCGGAGCCCCAGCGTGTTCCTGTTCCCCCCCCAAGCCTA  
40 AGGACACCCTGATGATCAGCAGAACCCCCGAGGTGACCTGTGTGGTGGTGGAT  
GTGAGCCACGAGGACCCTGAGGTGAAGTTCAACTGGTACGTGGACGGCGTGGGA  
GGTGCACAATGCCAAGACCAAGCCCAGGGAGGAGCAGTACAACAGCACCTACC  
GGGTGGTGTCCGTGCTGACCGTGTGACCAAGGATTGGCTGAACGGCAAGGA  
GTACAAGTGTAAAGGTGTCCAACAAGGCCCTGCCTGCCCTATCGAGAAAACCAT



CAGCAAGGCCAAGGGCCAGCCCAGAGAGCCCCAGGTGTACACCCTGCCCCCT  
AGCAGAGATGAGCTGACCAAGAACCAGGTGTCCCTGACCTGCCTGGTGAAGGG  
CTTCTACCCCAGCGACATCGCCGTGGAGTGGGAGAGCAACGGCCAGCCCGAG  
AACAACTACAAGACCACCCCCCTGTGCTGGACAGCGATGGCAGCTTCTTCCTG  
5 TACAGCAAGCTGACCGTGGACAAGAGCAGATGGCAGCAGGGCAACGTGTTTCA  
CTGCTCCGTGATGCACGAGGCCCTGCACAATCACTACACCCAGAAGAGCCTGA  
GCCTGTCCCCTGGCAAG

**Optimised L1 light chain DNA (SEQ ID No:43)**

10 GACATCGTGATGACCCAGAGCCCCCTGAGCCTGCCCGTGACCCCTGGCGA  
GCCCGCCAGCATCAGCTGTAGAGTGAGCCAGAGCCTGCTGCACAGCAACG  
GCTACACCTACCTGCACTGGTATCTGCAGAAGCCTGGCCAGAGCCCTCAG  
CTGCTGATCTACAAGGTGTCCAACCGGTTTCAGCGGCGTGCCTGATAGATTC  
AGCGGCAGCGGCTCCGGCACCGACTTCACCCTGAAGATCAGCAGAGTGG  
15 GGCCGAGGATGTGGGCGTGTACTACTGCTCCCAGACCAGACACGTGCCTT  
ACACCTTTGGCGGCGGAACAAAGGTGGAGATCAAGCGTACGGTGGCCGCC  
CCCAGCGTGTTTCATCTTCCCCCCCAGCGATGAGCAGCTGAAGAGCGGCAC  
CGCCAGCGTGGTGTGTCTGCTGAACAATTCTACCCCCGGGAGGCCAAGG  
TGCAGTGGAAGGTGGACAATGCCCTGCAGAGCGGCAACAGCCAGGAGAG  
20 CGTGACCGAGCAGGACAGCAAGGACTCCACCTACAGCCTGAGCAGCACCC  
TGACCCTGAGCAAGGCCGACTACGAGAAGCACAAGGTGTACGCCTGTGAG  
GTGACCCACCAGGGCCTGTCCAGCCCCGTGACCAAGAGCTTCAACCGGGG  
CGAGTGC

**Claims**

1. A therapeutic antibody which is an antibody or antigen binding fragment and/or derivative thereof which binds  $\beta$ -amyloid peptide and which comprises the following CDRs:

- 5 CDRH1: DNGMA (SEQ ID No:1)  
 CDRH2: FISNLAYSIDYADTVTG (SEQ ID No:2)  
 CDRH3: GTWFAY (SEQ ID No:3)

within a human heavy chain variable region originating from the VH3 gene family and:

10

- CDRL1: RVSQSLLSNGYTYLH (SEQ ID No:4)  
 CDRL2: KVSNRFS (SEQ ID No:5)  
 CDRL3: SQTRHVPYT (SEQ ID No:6)

15 within a human light chain variable region originating from the amino acid sequence disclosed in GenPept entry CAA51135 (SEQ ID No:24).

2. A therapeutic antibody according to claim 1, in which the human heavy chain variable region originates from:

- 20 • A V gene selected from the following subset of VH3 family members: VH3-48, VH3-21, VH3-11, VH3-7, VH3-13, VH3-74, VH3-64, VH3-23, VH3-38, VH3-53, VH3-66, VH3-20, VH3-9 & VH3-43  
 • A V gene selected from the following subset of VH3 family members: VH3-48, VH3-21 & VH3-11  
 • The VH3-48 gene

25 or an allele thereof.

3. A therapeutic antibody according to claim 2, in which the human acceptor heavy chain framework is derived from M99675 (SEQ ID No:21) together with a framework 4.

30

4. A therapeutic antibody according to claim 3, in which the framework 4 sequence is that encoded by the human JH4 minigene (Kabat):  
 YFDYWGGTLVTVSS (SEQ ID No:23)

35 of which the initial four residues fall within the CDR3 region is replaced by the incoming CDR from the donor antibody.

5. A therapeutic antibody according to claim 4 which contains one or more substitutions of amino acid residues based on the corresponding residues found in the donor V<sub>H</sub> domain having the sequence: SEQ ID No:17 and V<sub>L</sub> domain having the sequence: SEQ ID No: 19 that maintain all or substantially all of the binding affinity of the donor antibody for  $\beta$ -amyloid peptide.

40

6. A therapeutic antibody according to claim 5 in which the human acceptor heavy chain framework derived from M99675 and JH4 contains one to four amino

acid residue substitutions selected from positions 24, 48, 93 and/or 94 (Kabat numbering).

7. A therapeutic antibody according to claim 6 in which the human acceptor heavy chain framework comprises the following residues (or a conservative substitute thereof):

(i)

Position	Residue
93	V
94	S

or

(ii)

Position	Residue
24	V
93	V
94	S

or

(iii)

Position	Residue
48	I
93	V
94	S

8. A therapeutic antibody comprising a V<sub>H</sub> chain having the sequence set forth in SEQ ID No:26 and a V<sub>L</sub> domain having the sequence set forth in SEQ ID No:32.

9. A therapeutic antibody comprising a V<sub>H</sub> chain having the sequence set forth in SEQ ID No: 28 and a V<sub>L</sub> domain having the sequence set forth in SEQ ID No:32.

10. A therapeutic antibody comprising a V<sub>H</sub> chain having the sequence set forth in SEQ ID No:30 and a V<sub>L</sub> domain having the sequence set forth in SEQ ID No:32.

11. A therapeutic antibody which is an antibody or antigen binding fragment and/or derivative thereof which binds  $\beta$ -amyloid peptide 1-12 (SEQ ID No:15) with equilibrium constant K<sub>D</sub> less than 100pM and has an equilibrium constant K<sub>D</sub> for binding to  $\beta$ -amyloid peptide 2-13 (SEQ ID No:44) which is 1000-fold greater than that for peptide 1-12 (SEQ ID No:15), both determinations being made in a surface plasmon resonance assay utilising peptide captured on streptavidin chip.

12. A therapeutic antibody which is an antibody or antigen binding fragment and/or derivative thereof which binds  $\beta$ -amyloid peptide 1-40 with equilibrium constant K<sub>D</sub> less than 10nM and has an equilibrium constant K<sub>D</sub> for binding to  $\beta$ -

amyloid peptide 2-13 (SEQ ID No:44) which is 1000-fold greater than that for peptide 1-12 (SEQ ID No:15), both determinations being made in the surface plasmon resonance assay described in Method B of the Examples.

- 5 13. A therapeutic antibody according to any preceding claim which is of IgG1 isotype.
14. A therapeutic antibody according to any preceding claim which essentially lacks the functions of a) activation of complement by the classical pathway; and b)
- 10 mediating antibody-dependent cellular cytotoxicity.
15. A therapeutic antibody according to claim 13 in which residues 235 and 237 have been mutated to alanine
- 15 16. A therapeutic antibody according to claim 1, which antibody comprises a heavy chain having the sequence set forth in SEQ ID No:34, 36 or 38 and a light chain having the sequence set forth in SEQ ID No:40.
17. A pharmaceutical composition comprising a therapeutic antibody according to
- 20 any preceding claim.
18. A method of treating a human patient afflicted with a  $\beta$ -amyloid peptide-related disease which method comprises the step of administering to said patient a therapeutically effective amount of a therapeutic antibody according to any of claims
- 25 1 to 16.
19. Use of a therapeutic antibody according to any of claims 1 to 16 in the manufacture of a medicament for the treatment of a  $\beta$ -amyloid peptide-related disease.
- 30 20. An antibody or a fragment thereof comprising a  $V_H$  domain having the sequence: SEQ ID No:17 and a  $V_L$  domain having the sequence: SEQ ID No: 19.