

UK Patent Application (12) GB (19) 2 286 395 (13) A

(43) Date of A Publication 16.08.1995

(21) Application No 9502874.2

(22) Date of Filing 14.02.1995

(30) Priority Data

(31) 9402807

(32) 14.02.1994 (33) GB

(71) Applicant(s)

Xenova Limited

(Incorporated in the United Kingdom)

240 Bath Road, SLOUGH, Berkshire, SL1 4EF,
United Kingdom

(72) Inventor(s)

Stephen James Brocchini

Justin Stephen Bryans

Adrian John Folkes

Christopher John Latham

Jule Elizabeth Brumwell

(74) Agent and/or Address for Service

J A Kemp & Co

14 South Square, Gray's Inn, LONDON, WC1R 5LX,
United Kingdom

(51) INT CL⁶

C07D 241/02, A61K 31/495

(52) UK CL (Edition N)

C2C CAA CKH CKS CLK CMM CQS CRM CTP CTR
C1340 C1341 C1343 C1344 C1410 C1470 C1494 C1510
C1530 C1532 C1535 C1562 C1626 C200 C213 C215
C22Y C220 C221 C222 C225 C226 C246 C247 C25Y
C250 C251 C252 C253 C254 C255 C28X C280 C281
C282 C29X C29Y C30Y C31Y C313 C314 C32Y C321
C322 C323 C332 C337 C338 C34Y C342 C351 C352
C355 C36Y C360 C361 C362 C364 C366 C368 C37Y
C373 C386 C387 C388 C389 C396 C43X C464 C601
C604 C612 C614 C615 C62X C620 C621 C623 C624
C628 C630 C633 C634 C635 C645 C65X C650 C652
C658 C66X C660 C661 C662 C678 C681 C699 C761
C762 C763 C80Y C802

U1S S2415

(56) Documents Cited

Chemical Abstracts 65:16969e

(58) Field of Search


UK CL (Edition N) C2C CTR

INT CL⁶ C07D

Online: CAS ONLINE.

(54) Pharmaceutical piperazines

(57) A diketopiperazine of formula (A):



wherein one or both of R₁ and R₂, which may be the same or different, is:

X, substituted phenyl, CH=C(W)V or cyclohexyl;

and where appropriate, the other of R₁ and R₂ is an optionally substituted phenyl group

X is a naphthyl group or an optionally substituted five- or six-membered saturated or unsaturated heterocyclic group

W is hydrogen or a phenyl group;

V is an optionally substituted phenyl group,

and the pharmaceutically acceptable salts and esters thereof have activity as inhibitors of plasminogen activator inhibitor.

GB 2 286 395 A


PHARMACEUTICAL COMPOUNDS

The present invention relates to compounds useful as inhibitors of plasminogen activator inhibitor (PAI), to their preparation and to pharmaceutical and veterinary compositions containing them.

Plasminogen activators (PAs) are serine proteases which control the activation of the zymogen, plasminogen, to the active enzyme plasmin. Plasmin is important in a number of physiological and pathological processes including fibrinolysis, tissue remodelling, tumour growth and metastasis. The glycoprotein plasminogen activator inhibitor (PAI) is an endogenous fast-acting inhibitor of PA activity. PAI is a member of the serpin family and is synthesised by a variety of cells including endothelial cells. An imbalance between PAs and PAI contributes to a number of pathological conditions including haemostasis, inflammation, tumour growth and metastasis.

The present invention provides a diketopiperazine of formula (A) :

20



25

wherein one or both of R₁ and R₂, which may be the same or different, is:

(I) X, or a phenyl group which is substituted by X,
C(O)X, OC(O)CH₂X, OCH₂CH₂X, CH₂X, CONH(CH₂)_nX,
O(CH₂)_nCH(OH)(CH₂)_nX or $-\text{C}(\text{O})\text{NH}-\left(\text{C}_6\text{H}_4\right)_n\text{CH}_2\text{X}$

5 or which is fused to a group X;

(II) a phenyl group substituted by CH₂NR₁₂R₁₃,

OC(O)(CH₂)_nZ, CH(OR₁₂)(OR₁₃), (CH₂)_nNR₁₄C(O)(CH₂)_mNR₁₂R₁₃, -

CH₂NR₁₂-(CH₂)_nNR₁₅R₁₆ or O(CH₂)_nCH(OH)(CH₂)_nN(R₁₂R₁₃);

(III) a group CH=C(W)V; or

10 (IV) a cyclohexyl group;

and where appropriate, the other of R₁ and R₂ is a phenyl group optionally substituted by one or more groups independently selected from halogen, nitro, methoxy,

NHC(O)R₁₂, CO₂H, O(CH₂)_nN(R₁₂R₁₃), CH₂Y(CH₂)_nN(R₁₂R₁₃),

15 C₁-C₄ alkyl and (CH₂)_nC(O)OR₁₂;

X is a naphthyl group or a five- or six-membered saturated or unsaturated heterocyclic group containing one or more heteroatoms, which heteroatoms may be the same or different and are independently selected from O, N and S; the

20 heteroatom(s) when nitrogen being optionally substituted by hydrogen, methyl, oxygen, tertiary-butyloxycarbonyl,

-(CH₂)_nCH₂OH or SO₂Me; the heterocyclic ring being optionally substituted by halogen, Me, MeS, phenyl,

O(CH₂)_nNR₁₂R₁₃, -N(R₁₂)(CH₂)_nN(R₁₂R₁₃), -(CH₂)_nN(R₁₂R₁₃) or

25 -O(CH₂)_nO(CH₂)_nN(R₁₂R₁₃), or the heterocyclic ring optionally containing one or more carbonyl groups and being optionally fused to a benzene ring, which benzene ring is optionally substituted by 1 or 2 C₁-C₆ alkoxy groups;

Y is O or S;

Z is a C₃-C₆ cycloalkyl group;

R₁₂, R₁₃ and R₁₄, which may be the same or different, are hydrogen or C₁-C₆ alkyl;

5 R₁₅ and R₁₆, which may be the same or different, are hydrogen or C₁-C₆ alkyl, or R₁₅ and R₁₆ form, together with the nitrogen atom to which they are attached, a 5- or 6-membered heterocyclic group;

W is hydrogen or a phenyl group;

10 V is a phenyl group optionally substituted by one or more groups independently selected from nitro, alkoxy, O(CH₂)_nNR₁₂R₁₃, and NR₁₂R₁₃; and

m and n are each, independently, 0 or an integer having the value 1, 2, 3 or 4;

15 or a pharmaceutically acceptable salt or ester thereof.

A C₁-C₆ alkyl group is, for example, a C₁-C₄ alkyl group, such as a methyl, ethyl, propyl, i-propyl, n-butyl, sec-butyl or tert-butyl group.

A halogen may be F, Cl, Br or I.

20 In compounds of formula A free rotation may occur at room temperature about the single bonds connecting substituents R₁ and R₂ to the double bonds at positions 3 and 6 of the piperazine-2,5-dione ring.

In one embodiment at least one of R₁ and R₂, which may 25 be the same or different, is chosen from a naphthyl group, X, a phenyl group substituted by X, C(O)X, OC(O)CH₂X, OCH₂CH₂X, or CH₂X and a phenyl group which is fused to a group X; wherein X is a five- or six-membered saturated or

unsaturated heterocyclic group containing one or two heteroatoms, which heteroatoms may be the same or different and are independently selected from O, N and S, the heteroatom(s) when nitrogen being optionally substituted by


5 hydrogen, methyl, oxygen, tertiary-butyloxycarbonyl, -(CH₂)_nCH₂OH or SO₂Me, the heterocyclic ring being optionally substituted by hydrogen, halogen, methyl, MeS, phenyl, O(CH₂)_nNR₁₂R₁₃, O(CH₂)_nN(R₁₂R₁₃) or -O(CH₂)_nO(CH₂)_nN(R₁₂R₁₃); the heterocyclic ring optionally

10 containing one or more carbonyl groups, and being optionally fused to a benzene ring; and the other of R₁ and R₂ is a phenyl group optionally substituted at the 2, 3 or 4-position by CH₂NR₁₂R₁₃, (CH₂)_nNR₁₄C(O)(CH₂)_mNR₁₂R₁₃, halogen, nitro, -NHC(O)R₁₂, -O(CH₂)_nN(R₁₂R₁₃) or -CH₂Y(CH₂)_nN(R₁₂R₁₃)

15 wherein Y is O or S. In a particularly preferred series of compounds the said other of R₁ and R₂ is a phenyl group substituted at the 4-position by -O(CH₂)_nN(R₁₂R₁₃), -CH₂Y(CH₂)_nN(R₁₂R₁₃) or -(CH₂)_nNR₁₄C(O)(CH₂)_mNR₁₂R₁₃.

In a further embodiment one of R₁ and R₂ is X, a

20 phenyl group substituted by X, -CH₂X, -OCH₂CH₂X,

O(CH₂)_nCH(OH)CH₂X or  ; wherein X is a 5

or 6-membered saturated or unsaturated heterocyclic group as defined above which is optionally substituted and

25 optionally fused to a benzene ring, for instance a pyridyl, imidazolyl, furyl, pyrrolyl, pyrrolidinyl, thieryl, piperazinyl, piperidinyl, morpholinyl, quinolyl, isoquinolyl or indolyl group; and the other of R₁ and R₂ is

a phenyl group optionally substituted at the 4-position by
-O(CH₂)_nN(R₁₂R₁₃), -CH₂Y(CH₂)_nN(R₁₂R₁₃) or
-(CH₂)_nNR₁₄C(O)(CH₂)_mNR₁₂R₁₃. In this embodiment it is
particularly preferred for X to be a furyl, imidazolyl,
5 pyrrolyl, thieryl, morpholinyl, piperidinyl or isoquinolyl
group.

In a further embodiment, R₁₂ and R₁₃, which may be the same or different, are hydrogen or C₁-C₃ alkyl and n is an integer of value 1 or 2.

10 In a yet further embodiment one of R₁ and R₂ is a phenyl group which is substituted by X, CO(X), OCO(O)CH₂X, OCH₂CH₂X, CH₂X or which is fused to a group X, wherein X is a five- or six-membered heterocyclic ring containing one or two heteroatoms which may be the same or different,
15 independently selected from O, N and S, the heteroatom(s) when nitrogen being optionally substituted by methyl, and the heterocyclic ring being optionally fused to a benzene ring.

In another embodiment one of R₁ and R₂ is a phenyl
20 group substituted by CH₂NR₁₂R₁₃, OC(O)(CH₂)_nZ, CH(OR₁₂)(OR₁₃), (CH₂)_nNR₁₄C(O)(CH₂)_mN(R₁₂R₁₃); wherein R₁₂, R₁₃ and R₁₄, which may be the same or different, are independently selected from hydrogen or C₁-C₃ alkyl; Z is a C₅ or C₆ cycloalkyl group; and m and n are, independently, integers having the
25 values 1, 2 or 3.

In a further embodiment R₁₂, R₁₃ and R₁₄, which may be the same or different, are independently selected from hydrogen and C₁-C₂ alkyl; Z is a cyclopentyl group; and

m and n are, independently, integers having the values of 1 or 2.

In a yet further embodiment one of R₁ and R₂ is a phenyl group optionally substituted by one or more groups 5 independently selected from chloro, nitro, methoxy, NHCO₂R₁₂, CO₂H and O(CH₂)_nNR₁₂R₁₃; R₁₂ and R₁₃, which may be the same or different, are independently selected from hydrogen or methyl and n is an integer having the value 1 or 2.

In another embodiment one of R₁ and R₂ is a group 10 CH=C(W)V, W is a phenyl group optionally substituted by one of more groups independently selected from nitro, methoxy and O(CH₂)_nNMe₂ and n is an integer having the value 1, 2, 3 or 4.

In a further embodiment n is 1 or 2. 15 In a yet further embodiment one of R₁ and R₂ is a phenyl group optionally substituted by NHAc or methoxy.

In another embodiment one of R₁ and R₂ is cyclohexyl and the other is a phenyl group optionally substituted by NHC(O)R₁₂.

20 In a further embodiment one of R₁ and R₂ is cyclohexyl and the other is a phenyl group optionally substituted by NHC(O)Me.

In a further embodiment R₃ is C₁-C₂ alkyl or (CH₂)_nC(O)OR₁₂; R₁₂ is hydrogen or C₁-C₂ alkyl and n is an 25 integer of value 1 or 2.

In a yet further embodiment R₃ is methyl or CH₂C(O)OR₁₂ and R₁₂ is hydrogen or methyl.

Certain diketopiperazines have been disclosed as

having utility as bioactive agents. Yokoi et al in J. Antibiotics vol XLI No. 4, pp 494-501 (1988) describe structure-cytotoxicity relationship studies on a series of diketopiperazines related to neihumycin, a compound

5 obtained from the micro-organism Micromonospora neihuensis.

Kamei et al in J. Antibiotics vol XLIII No. 8, 1018-1020 disclose that two diketopiperazines, designated piperafizines A and B, have utility as potentiators of the cytotoxicity of vincristine.

10 Examples of specific compounds of formula A are as follows. The compound numbering is adhered to in the rest of the specification:

1926 (3Z,6Z)-3-Benzylidene-6-(4-imidazolyl)methylene-2,5-piperazinedione.

15 1930 (3Z,6Z)-3-Benzylidene-6-(4-(1-imidazolyl)benzylidene)-2,5-piperazinedione.

1929 (3Z,6Z)-3-Benzylidene-6-(4-(1-imidazolylmethyl)benzylidene)-2,5-piperazinedione.

1959 (3Z,6Z)-3,Benzylidene-6-(4-(2-dimethylaminoethoxy)-3-methoxybenzylidene)-2,5-piperazinedione hydrochloride.

20 1927 (3Z,6Z)-3-Benzylidene-6-(4-(5-methylimidazolyl)methylene-2,5-piperazinedione.

1921 (3Z,6Z)-3-Benzylidene-6-(4-dimethylaminocinnamylidene)-2,5-piperazinedione.

25 1976 (3Z,6Z)-3-(4-(3-Dimethylaminopropoxy)benzylidene)-6-(4-(1-imidazolyl)benzylidene-2,5-piperazinedione.

1910 (3Z,6Z)-3-Benzylidene-6-(4-(2-imidazolylethoxy)benzylidene)-2,5-piperazinedione.

- 1923 (3Z,6Z)-3-Benzylidene-6-(4-nitrocinnamylidene-2,5-piperazinedione.
- 1657 (3Z,6Z)-3-(4-Aminomethylbenzylidene)-6-(4-methoxybenzylidene)-2,5-piperazinedione.
- 5 1693 (3Z,6Z)-3-(1-methanesulfonyl-3-indolyl)methylene-6-(4-methoxybenzylidene)-2,5-piperazinedione.
- 1886 (3Z,6Z)-3-(4-Methoxybenzylidene)-6-(4-phthalimidoacetoxybenzylidene)-2,5-piperazinedione.
- 1922 (3Z,6Z)-3-Benzylidene-6-(γ -phenylcinnamylidene)-2,5-piperazinedione.
- 10 1618 (3Z,6Z)-3-(1-tert-butoxycarbonyl-3-indolyl)methylene-6-(2-thenylidene)-2,5-piperazinedione.
- 1560 (3Z,6Z)-3-(2,6-Dichlorobenzylidene)-6-(1-tert-butoxycarbonyl-3-indolyl)methylene-2,5-piperazinedione.
- 15 1950 (3Z,6Z)-3-Benzylidene-6-(4-(2-dimethylaminoethoxy)-3-methoxycinnamylidene)-2,5-piperazinedione.
- 1975 (3Z,6Z)-3-(4-(3-Dimethylaminopropoxy)benzylidene)-6-(4-(1-imidazolylmethyl)benzylidene)-2,5-piperazinedione.
- 1983 (3Z,6Z)-3-Benzylidene-6-(4-N-methyl-N-(4-(N-methylpiperidinyl))aminomethylbenzylidene-2,5-piperazinedione.
- 20 1509 (3Z,6Z)-3-Benzylidene-6-(3-indolylmethylene)-2,5-piperazinedione.
- 1542 (3Z,6Z)-3-(2,6-Dichlorobenzylidene)-6-(3-furylmethylene)-2,5-piperazinedione.
- 25 1545 (3Z,6Z)-3-(3-Indolylmethylene)-6-(4-methoxybenzylidene)-2,5-piperazinedione.
- 1507 (3Z,6Z)-3-(4-Methoxybenzylidene)-6-(2-(1-

- tertbutoxycarbonyl)pyrrolylmethylene-2,5-piperazinedione.
- 1506 (3Z,6Z)-3-(4-Methoxybenzylidene)-6-(3-(1-tert-butoxycarbonyl)indolyl)methylene-2,5-piperazinedione.
- 1471 (3Z,6Z)-3-Benzylidene-6-(3-(1-tert-butoxycarbonyl)indolyl)methylene-2,5-piperazinedione.
- 5 1474 (3Z,6Z)-3-(4-Methoxybenzylidene)-6-(2-thienylmethylene)-2,5-piperazinedione.
- 1476 (3Z,6Z)-3-(4-Methoxybenzylidene)-6-(3-furylmethylene)-2,5-piperazinedione.
- 10 1672 (3Z,6Z)-3-(Acetamidobenzylidene)-6-cyclohexylmethylen-2,5-piperazinedione.
- 1676 (3Z,6Z)-3-(4-Acetamidobenzylidene)-6-cinnamylidene-2,5-piperazinedione.
- 1891 (3Z,6Z)-3-Benzylidene-6-(diethoxymethylbenzylidene)-2,5-piperazinedione.
- 15 1982 (3Z,6Z)-3-Benzylidene-6-(4-(N-methyl-N-(2-dimethylaminoethyl)aminomethylbenzylidene)-2,5-piperazinedione hydrochloride.
- 1884 (3Z,6Z)-3-Benzylidene-6-cyclohexylmethylen-2,5-piperazinedione.
- 20 1845 (3Z,6Z)-3-(4-Acetamidobenzylidene)-6-(3,4-methylenedioxybenzylidene)-2,5-piperazinedione.
- 1950 (3Z,6Z)-3-benzylidene-6-(4-(2-dimethylaminoethoxy)-3-methoxycinnamylidene)-2,5-piperazinedione.

1718 (3Z,6Z)-3-(2-Indolylmethylene)-6-(4-methoxybenzylidene)-2,5-piperazinedione.

1808 (3Z,6Z)-3-Benzylidene-6-(3,4-methylenedioxybenzylidene)-2,5-piperazinedione.

5 1809 (3Z,6Z)-3-(4-Methoxybenzylidene)-6-(3,4-methylenedioxybenzylidene)-2,5-piperazinedione.

1470 (3Z,6Z)-3-Benzylidene-6-(2-(1-tertbutoxycarbonyl)pyrrolyl)methylene-2,5-piperazinedione.

5023 (3Z,6Z)-3-(4-Dimethylaminomethylbenzylidene)-6-(4-(3-dimethylaminopropoxy)benzylidene-2,5-piperazinedione.

5026 (3Z,6Z)-3-(4-(3-Dimethylaminopropoxy)benzylidene)-6-(4-(1-imidazolyl)methylbenzylidene)-2,5-piperazinedione.

5030 (3Z,6Z)-3-(4-(3-Dimethylaminopropoxy)benzylidene)-6-(4-(1-imidazolyl)benzylidene)-2,5-piperazinedione.

15 5367 (2-(4-((3Z,6Z)-6-(4-(3-Dimethylaminopropoxy)benzylidene)-2,5-dioxo-3-piperazinylidene)methylbenzoyl)-1,2,3,4-tetrahydroisoquinoline.

5386 N-(2-(1,2,3,4-Tetrahydro-2-isoquinolyl)ethyl)-4-

20 ((3Z,6Z)-6-(4-(3-dimethylaminopropoxy)benzylidene)-2,5-dioxo-3-piperazinylidene)methylbenzamide.

5397 N-(4-(1,2,3,4-Tetrahydro-2-isoquinolyl)butyl)-4-((3Z,6Z)-6-(4-(3-dimethylaminopropoxy)benzylidene)-2,5-dioxo-3-piperazinylidene)methylbenzamide.

25 5027 (3Z,6Z)-6-(4-(3-Dimethylaminopropoxy)benzylidene-3-(4-pyridylmethylene)-2,5-piperazinedione.

5028 (3Z,6Z)-6-(4-(3-Dimethylaminopropoxy)benzylidene)-3-(3-pyridylmethylene)-2,5-piperazinedione.

5041 (3Z,6Z)-6-(4-(3-Dimethylaminopropoxy)benzylidene)-3-furfurylidene-2,5-piperazinedione.

5042 (3Z,6Z)-6-(4-(3-Dimethylaminopropoxy)benzylidene)-3-(3-Thenylidene)-2,5-piperazinedione.

5 5046 (3Z,6Z)-6-(4-(3-Dimethylaminopropoxy)benzylidene)-3-(2-Thenylidene)-2,5-piperazinedione.

5052 (3Z,6Z)-6-(4-(3-Dimethylaminopropoxy)benzylidene)-3-(3-Furylmethylene)-2,5-piperazinedione.

10 5188 (3Z,6Z)-6-(4-(3-Dimethylaminopropoxy)benzylidene)-3-(2-Naphthylmethylene)-2,5-piperazinedione.

5200 (3Z,6Z)-6-(4-(3-Dimethylaminopropoxy)benzylidene)-3-(1-Naphthylmethylene)-2,5-piperazinedione.

5032 (3Z,6Z)-6-Benzylidene-3-(4-(3-dimethylamino-2-hydroxypropoxy)benzylidene)-2,5-piperazinedione.

15 5040 (3Z,6Z)-6-Benzylidene-3-(4-(2-hydroxy-3-morpholinopropoxy)benzylidene)-2,5-piperazinedione.

5057 (3Z,6Z)-6-Benzylidene-3-(4-(2-hydroxy-3-(1-imidazolyl)propoxy)benzylidene)-2,5-piperazinedione.

5043 (3Z,6Z)-6-Benzylidene-3-(4-(2-hydroxy-3-(4-(2-hydroxyethyl)-1-piperazinyl)propoxy)benzylidene)-2,5-piperazinedione.

5062 (3Z,6Z)-6-(4-(2-Dimethylaminoethoxy)benzylidene)-3-(3-furylmethylene)-2,5-piperazinedione.

25 5071 (3Z,6Z)-6-(4-(2-Dimethylaminoethoxy)benzylidene)-3-(3-thenylidene)-2,5-piperazinedione.

5072 (3Z,6Z)-6-(4-(2-Dimethylaminoethoxy)benzylidene)-3-(5-methylthio-2-thenylidene)-2,5-piperazinedione.

5054 (3Z,6Z)-6-Benzylidene-3-(4-(2-

- morpholinoethoxy)benzylidene)-2,5-piperazinedione.
- 5055 (3Z,6Z)-6-Benzylidene-3-(4-(2-(1-imidazolyl)ethoxy)benzylidene)2,5-piperazinedione.
- 5053 (3Z,6Z)-6-Benzylidene-3-(4-(2-(1-pyrrolidinyl)ethoxy)benzylidene)2,5-piperazinedione.
- 5069 (3Z,6Z)-6-(4-(2-Dimethylaminoethoxymethyl)benzylidene)-3-(3-thenylidene)-2,5-piperazinedione.
- 5077 (3Z,6Z)-6-(4-(2-Dimethylaminoethoxymethyl)benzylidene)-3-(3-furylmethylene)-2,5-piperazinedione.
- 5074 (3Z,6Z)-6-(4-Dimethylaminoacetamidomethylbenzylidene)-3-(3-thenylidene)-2,5-piperazinedione.
- 5079 (3Z,6Z)-3-(2-Bromobenzylidene)-6-(4-dimethylaminoacetamidomethylbenzylidene)-2,5-piperazinedione.
- 15 dimethylaminoacetamidomethylbenzylidene)-2,5-piperazinedione.
- 5081 (3Z,6Z)-6-(4-Dimethylaminoacetamidomethylbenzylidene)-3-(3-furylmethylene)-2,5-piperazinedione.
- 5061 (3Z,6Z)-6-Benzylidene-3-(4-dimethylaminoacetamidomethylbenzylidene)-2,5-piperazinedione.
- 20 dimethylaminoacetamidomethylbenzylidene)-2,5-piperazinedione.
- 5073 (3Z,6Z)-6-(4-(2-Dimethylaminoethylthiomethyl)benzylidene)-3-(3-furylmethylene)-2,5-piperazinedione.
- 25 5078 (3Z,6Z)-6-(4-(2-Dimethylaminoethylthiomethyl)benzylidene)-3-(3-thenylidene)-2,5-piperazinedione.
- 1912 (3Z,6Z)-6-Benzylidene-3-(4-

dimethylaminoacetamidoaminomethylbenzylidene)-2,5-piperazinedione.

5324 (3Z,6Z)-6-Benzylidene-3-(5-(2-dimethylaminoethoxy)-2-thienylmethylene)-2,5-piperazinedione.

5 5327 (3Z,6Z)-6-Benzylidene-3-(4-(2-dimethylaminoethoxy)-2-thienylmethylene)-2,5-piperazinedione.

5335 (3Z,6Z)-6-Benzylidene-3-(5-(2-dimethylaminoethyl)-2-thienylmethylene)-2,5-piperazinedione.

5388 (3Z,6Z)-6-Benzylidene-3-(5-(2-(2-

10 dimethylaminoethoxy)ethoxy)-2-thienylmethylene)-2,5-piperazinedione.

5389 (3Z,6Z)-6-Benzylidene-3-(5-(6-dimethylaminohexyloxy)-2-thienylmethylene)-2,5-piperazinedione.

5299 (3Z,6Z)-6-Benzylidene-3-(5-(2-

15 dimethylaminoethyl)methylamino-2-thienylmethylene)-2,5-piperazinedione.

5075 (3Z,6Z)-3-(2,5-Dichloro-3-thenylidene)-6-benzylidene-2,5-piperazinedione.

5371 N-(4-(1,2,3,4-Tetrahydro-2-isoquinolyl)butyl)-4-((3Z,6Z)-6-benzylidene-2,5-dioxo-3-piperazinylidene)methylbenzamide.

5391 N-(2-(1,2,3,4-Tetrahydro-2-isoquinolyl)ethyl)-4-((3Z,6Z)-6-benzylidene-2,5-dioxo-3-piperazinylidene)methylbenzamide.

25 5394 N-(3-(1,2,3,4-Tetrahydro-2-isoquinolyl)propyl)-4-((3Z,6Z)-6-benzylidene-2,5-dioxo-3-piperazinylidene)methylbenzamide.

5393 N-(4-(2-(1,2,3,4-Tetrahydro-2-

isoquinolyl)ethyl)phenyl-4-((3Z,6Z)-6-benzylidene-2,5-dioxo-3-piperazinylidene)methylbenzamide.


5402 N-(4-(2-(1,2,3,4-Tetrahydro-2-

isoquinolyl)ethyl)phenyl)-4-((3Z,6Z)-2,5-dioxo-6-(4-

5 nitrobenzylidene)-3-piperazinylidene)methylbenzamide.

Compounds of formula A, may be prepared by a process which comprises either (i) condensing compound of formula

(I)




wherein R₂ is as defined above and is optionally protected, with a compound of formula (II):

15



wherein R₁ is as defined above and is optionally protected, in the presence of a base in an organic solvent; or (ii) condensing a compound of formula (I'):

20



wherein R₁ is as defined above and is optionally protected, 25 with a compound of formula (III):



wherein R₂ is as defined above and is optionally protected,

in the presence of a base in an organic solvent; and, in either case (i) or (ii), if required, removing optionally present protecting groups and/or, if desired, converting one compound of formula A into another compound of formula

5 A, and/or, if desired, converting a compound of formula A into a pharmaceutically acceptable salt or ester thereof, and/or, if desired, converting a salt or ester into a free compound, and/or, if desired, separating a mixture of isomers of compounds of formula A into the single isomers.

10 A compound of formula A produced directly by the condensation reaction between (I) and (II) or (I') and (III) may be modified, if desired, by converting R₁ into a different R₁ group. These optional conversions may be carried out by methods known in themselves. For example, a
15 compound of formula A in which R₁ comprises an ester group may be converted to a compound of formula A wherein the corresponding substituent is a free -COOH or OH group, by acid or alkaline hydrolysis at a suitable temperature, for example from ambient temperature to 100°C.

20 A compound of formula A in which either or both of R₁ and R₂ includes an -OH group may be converted into a compound of formula A wherein the corresponding substituent is esterified, for example by treating with a suitable carboxylic acid in the presence of an appropriate coupling
25 agent, acid anhydride or acid chloride in an inert solvent.

A compound of formula A in which either or both of R₁ and R₂ includes a -CO₂H group may be converted into a

compound of formula A wherein the corresponding substituent is esterified, for example by treating the carboxylic acid with a suitable C₁-C₆ alkyl alcohol in the presence of 1,3-dicyclohexylcarbodiimide in an inert solvent.

- 5 A compound of formula A in which either or both of R₁ and R₂ includes a free -CO₂H group may be converted into a compound of formula A in which the corresponding substituent is a group -CON(R₁₁R₁₂), wherein R₁₁ and R₁₂ are as defined above, for example by treatment with ammonia or
10 an amine in the presence of 1,3-dicyclohexylcarbodiimide in an inert solvent.

A compound of formula A in which either or both of R₁ and R₂ includes a free -CO₂H group may be converted into a compound of formula A wherein the corresponding substituent
15 is a -CH₂OH group by reduction, for example using borane in a suitable solvent such as tetrahydrofuran.

- A compound of formula A in which either or both of R₁ and R₂ is a nitro group may be converted into a compound of formula A in which the corresponding substituent is an
20 amino group by reduction under standard conditions, for example by catalytic hydrogenation.

Protecting groups for substituents on R₁ and/or R₂ in any of the compounds of formulae (I), (I'), (II) and (III) are optionally introduced prior to step (i) or step (ii)
25 when either or both R₁ and R₂ include one or more groups which are sensitive to the condensation reaction conditions or incompatible with the condensation reaction, for example a -COOH, -CH₂OH or amino group. The protecting groups are

then removed at the end of the process. Any conventional protecting group suitable for the group R₁ and/or R₂ in question may be employed, and may be introduced and subsequently removed by well-known standard methods.

5 The condensation reaction between compounds (I) and (II) or (I') and (III) is suitably performed in the presence of a base which is potassium t-butoxide, sodium hydride, potassium carbonate, sodium carbonate, caesium carbonate, sodium acetate, potassium fluoride on alumina, 10 or triethylamine in a solvent such as dimethylformamide, potassium t-butoxide in t-butanol, or a mixture of t-butanol and dimethylformamide (DMF). The reaction is typically performed at a temperature from 0°C to the reflux temperature of the solvent.

15 The compounds of formula (I) may be prepared by a process comprising reacting 1,4-diacetyl-2,5-piperazinedione with a compound of formula (III) as defined above, in the presence of a base in an organic solvent. Similarly, the compounds of formula (I') may be prepared by 20 a process which comprises reacting 1,4-diacetyl-2,5-piperazinedione with a compound of formula (II) as defined above, in the presence of a base in an organic solvent.

If necessary, the resulting compound of formula (I) or (I') can be separated from other reaction products by 25 chromatography.

The reaction of 1,4-diacetyl-2,5-piperazinedione with the compound of formula (III) or (II) is suitably performed under the same conditions as described above for the

condensation between compounds (I) and (II), or (I') and (III).

The substituted aldehydes of formulae (II) and (III) are known compounds or can be prepared from readily available starting materials by conventional methods. The 1,4-diacetyl-2,5-piperazinedione used as a starting material in the preparation of compounds of formula (I) may be prepared by treating 2,5-piperazinedione (glycine anhydride) with an acetylating agent. The acetylation may be performed using any conventional acetylating agent, for example acetic anhydride under reflux or, alternatively, acetic anhydride at a temperature below reflux in the presence of 4-dimethylaminopyridine.

Compounds of formula (I) may also be prepared by the microwave irradiation of a mixture comprising 1,4-diacetyl-2,5-piperazinedione, a compound of formula (III) and potassium fluoride on alumina (as base) in the absence of solvent.

Compounds of formula (I) may alternatively be prepared directly from 2,5-piperazinedione (glycine anhydride) by a process which comprises treating the 2,5-piperazinedione with a mixture comprising a compound of formula (III), sodium acetate and acetic anhydride at an elevated temperature, for example under reflux.

Compounds of formula (I') may be prepared by analogous processes, replacing compound (III) in each case by a compound of formula (II).


Compounds of formula A may also be prepared by a

process comprising the microwave irradiation of (i) a mixture comprising a compound of formula (I) as defined above, a compound of formula (II) and potassium fluoride on alumina, or (ii) a mixture comprising a compound of formula 5 (I') a compound of formula (III) and potassium fluoride on alumina, or (iii) a mixture comprising 1,4-diacetyl-2,5-piperazinedione, a compound of formula (II), a compound of formula (III) and potassium fluoride on alumina. The irradiation is performed in the absence of a solvent.

10 Compounds of formula (A) may also be obtained directly by a process which comprises condensing together 1,4-diacetyl-2,5-piperazinedione, a compound of formula (II) and a compound of formula (III) in the presence of a base in an organic solvent. Suitable bases, solvents and 15 reaction conditions are as described above for the condensation reaction between, for example, compounds (I) and (II).


An alternative direct process for the preparation of compounds of formula (A) comprises condensing together 2,5-20 piperazinedione, a compound of formula (II) and a compound of formula (III) in the presence of sodium acetate and acetic anhydride at elevated temperature, for example under reflux.

An alternative process for the preparation of 25 compounds of formula (I) comprises treating a compound of formula (V) :



wherein R₆ to R₁₀ are as defined above, X is a halogen and R' is a C₁-C₆ alkyl group, with ammonia followed by acetic anhydride.

Compounds of formula (I') may be prepared by an
10 analogous process which comprises treating a compound of
formula (V') :



wherein R₁ to R₅, X and R' are as defined above, with ammonia followed by acetic anhydride.

X in formula (V) or (V') is typically iodine. R' is, for example, a C₁-C₄ alkyl group such as a methyl, ethyl,
20 propyl, i-propyl, butyl, sec-butyl or tert-butyl group.

A review of synthetic approaches to unsaturated 3-monosubstituted and 3,6-disubstituted-2,5-piperazinediones is provided in Heterocycles, 1983, 20, 1407 (C.Shin).

Compounds of formula (A) may be optionally washed
25 after any of the above preparative procedures with one or more of the following: water, ethanol, ethyl acetate and diethyl ether.

Where appropriate compounds of formula (A) may be

optionally recrystallised from a suitable solvent such as methanol or acetic acid.

Compounds of formula (A) may be converted into pharmaceutically acceptable salts, and salts may be
5 converted into the free compound, by conventional methods. Suitable salts include salts with pharmaceutically acceptable, inorganic or organic, acids or bases. Examples of inorganic bases include ammonia and carbonates, hydroxides and hydrogen carbonates of group I and group II
10 metals such as sodium, potassium, magnesium and calcium. Examples of organic bases include aliphatic and aromatic amines such as methylamine, triethylamine, benzylamine, dibenzylamine or α - or β -phenylethylamine, and heterocyclic bases such as piperidine, 1-methylpiperidine and
15 morpholine. Examples of inorganic acids include hydrochloric acid, sulphuric acid and orthophosphoric acid. Examples of organic acids include p-toluenesulphonic acid, methansulphonic acid, mucic acid and succinic acid.

Compounds of formula (A) may also be converted into
20 pharmaceutically acceptable esters. Suitable esters include branched or unbranched, saturated or unsaturated C₁-C₆ alkyl esters, for example methyl, ethyl and vinyl esters.

The diketopiperazines of formula (A), both novel and
25 known and their pharmaceutically acceptable salts and esters (referred to hereinafter as the "present compounds") have utility as inhibitors of PAI. Elevated levels of PAI-1, by reducing the net endogenous fibrinolytic capacity,

can contribute to the pathogenesis of various thrombotic disorders including myocardial infarction, deep vein thrombosis and disseminated intravascular coagulation. The present compounds therefore can act as inhibitors of the
5 tPA/PAI-1 interaction. The present compounds can be used in the treatment of haemostatic disorders. A human or animal, e.g. a mammal, can therefore be treated by a method comprising administration of a therapeutically effective amount of a diketopiperazine of formula (A) or a
10 pharmaceutically or veterinarilly acceptable salt thereof.

Tissue plasminogen activator (tPA) is used as a fibrinolytic agent in the treatment of thrombotic disorders. The efficacy of the tPA in this role may be enhanced if it is administered together with a PAI
15 inhibitor. A human or animal, e.g. a mammal, can therefore be treated by a method comprising the combined administration of a therapeutically effective amount of tPA and a therapeutically effective amount of any one of the present compounds. The present invention also provides
20 products containing a diketopiperazine of formula (A) or a pharmaceutically acceptable salt or ester thereof and tPA as a combined preparation for simultaneous, separate or sequential use in the treatment of thrombotic disorders, for example where there is inappropriate PAI activity. In
25 such products the present compound is formulated for oral or parenteral (intravenous, intramuscular or subcutaneous) administration and the tPA is formulated for intravenous administration.

As one example, during acute myocardial infarction (MI) one of the present compounds may be administered to a patient together with tPA to enhance the efficacy of the tPA treatment. As a further example, early re-occlusion 5 following treatment of a patient with tPA may be prevented by the post-MI administration of one of the present compounds.

The compounds of formula (A) have been tested in a PAI functional assay. In this assay, a compound is incubated 10 with PAI-1 prior to addition to the tPA assay system. Inhibition of PAI-1 results in the production of plasmin from plasminogen. In turn, plasmin cleaves the chromogenic substrate S2251 (Kabi Vitrum) producing pNA (p-nitroaniline) which is detected spectrophotometrically at 15 405 nm (K.Nilsson et al, Fibrinolysis (1987) 1, 163-168). The results of the assay are reported below.

The present compounds can be administered in a variety of dosage forms, for example orally such as in the form of tablets, capsules, sugar- or film-coated tablets, liquid 20 solutions or suspensions or parenterally, for example intramuscularly, intravenously or subcutaneously. The present compounds may therefore be given by injection or infusion.

The dosage depends on a variety of factors including 25 the age, weight and condition of the patient and the route of administration. Typically, however, the dosage adopted for each route of administration when a compound of the invention is administered alone to adult humans is 0.001 to

10 mg/kg, most commonly in the range of 0.01 to 5 mg/kg, body weight. Such a dosage may be given, for example, from 1 to 5 times daily by bolus infusion, infusion over several hours and/or repeated administration.

5 When one of the present compounds is administered in combination with tPA to adult humans, the dosage adopted for each route of administration is typically from 0.001 to 10 mg, more typically 0.01 to 5 mg per kg body weight for a compound of the invention and from 5 to 500mg administered
10 intravenously for the tPA. A suitable dosage regimen for the tPA is 100 mg given intravenously over 3 hours as follows: 10% of the total dose as an i.v. bolus over 1-2 minutes, 50% of the total dose as an infusion over 1 hour, 40% of the total dose as an infusion over the subsequent 2
15 hours.

A diketopiperazine of formula (A) or a pharmaceutically acceptable salt or ester thereof is formulated for use as a pharmaceutical or veterinary composition also comprising a pharmaceutically or
20 veterinarianily acceptable carrier or diluent. The compositions are typically prepared following conventional methods and are administered in a pharmaceutically or veterinarianily suitable form. An agent for use as an inhibitor of PAI comprising any one of the present
25 compounds is therefore provided.

For example, the solid oral forms may contain, together with the active compound, diluents such as lactose, dextrose, saccharose, cellulose, corn starch or

potato starch; lubricants such as silica, talc, stearic acid, magnesium or calcium stearate and/or polyethylene glycols; binding agents such as starches, arabic gums, gelatin, methylcellulose, carboxymethylcellulose, or 5 polyvinyl pyrrolidone; disintegrating agents such as starch, alginic acid, alginates or sodium starch glycolate; effervesing mixtures; dyestuffs, sweeteners; wetting agents such as lecithin, polysorbates, lauryl sulphates. Such preparations may be manufactured in known manners, for 10 example by means of mixing, granulating, tabletting, sugar coating, or film-coating processes.

Liquid dispersions for oral administration may be syrups, emulsions and suspensions. The syrups may contain as carrier, for example, saccharose or saccharose with 15 glycerol and/or mannitol and/or sorbitol. In particular, a syrup for diabetic patients can contain as carriers only products, for example sorbitol, which do not metabolise to glucose or which only metabolise a very small amount to glucose. The suspensions and the emulsions may contain as 20 carrier, for example, a natural gum, agar, sodium alginate, pectin, methylcellulose, carboxymethylcellulose or polyvinyl alcohol.

Suspensions or solutions for intramuscular injections may contain, together with the active compound, a 25 pharmaceutically acceptable carrier such as sterile water, olive oil, ethyl oleate, glycols such as propylene glycol, and, if desired, a suitable amount of lidocaine hydrochloride. Some of the present compounds are insoluble

in water. A compound may be encapsulated within liposomes.

TESTING OF THE PRESENT
COMPOUNDS AS PAI INHIBITORS

5 Compounds of formula (A) were tested in a PAI chromogenic substrate assay. In the assay (K.Nilsson, Fibrinolysis (1987) 1, 163-168) each compound was incubated with PAI-1 prior to addition to the tPA assay system. Inhibition of PAI-1 by the compound of formula (A) resulted
10 in the production of plasmin from plasminogen. In turn, the plasmin cleaved the chromogenic substrate S2251 (Kabi-Vitrum) producing pNA (p-nitroaniline) which was detected spectrophotometrically at 405 nm.

The degrees of inhibition observed in the chromogenic
15 substrate assay at various concentrations, and/or IC₅₀ values, of compounds of formula (A) are presented in Table 1. IC₅₀ values for some compounds, not shown in Table 1, are listed in Table 2 which follows Table 1.

20

TABLE 1: INHIBITION OF PAI-1 IN THE S2251
CHROMOGENIC SUBSTRATE ASSAY

25

Compound No.	Concentration in μm				
	100	50	25	12.5	6.25
1470	70	20	2	0	0
1471	80	60	20	6	0
1474	64	52	28		
1476	68	48	18		
1506	75	58	26	4	2
1507	78	62	45	1	1

	1509	58	35	1	1	1
	1542	75	41	9	1	1
	1545	87	64	39	5	1
	1560	50	48	46	34	13
5	1618	51	32	3	1	
	1649	34	0	1	0	
	1657	53	60	46	2	
	1672	70	44	13	4	1
10	1676	29	51	52	12	1
	1693	89	2	1	0	
	1718	62	1	0	0	1
	1808	76	48	73	2	1
	1809	81	76	84	7	1
	1845	14	30	49	60	53
15	1884	40	14	0	0	0
	1886	42	40	18	6	0
	1891	28	36	17	3	3
	1910	27	36	50	61	63
	1912	30	55	29	22	17
20	1921	65	43	25	14	16
	1922	13	11	26	13	14
	1923	38	31	20	12	13
	1926	36	35	12	6	10
	1927	33	39	20	22	14
25	1928	67	60	47	24	19
	1929	27	45	59	48	16
	1930	54	61	79	38	30
	1959	5	1	2	2	1
	1975	7	0	0	0	0
30	1976	3	0	0	0	0
	1950	19	3	2	2	1
	1982	48	49	28	6	1
	1983	34	14	0	0	0

Compound No.	Concentration in μM			IC_{50}
	100 μM	50 μM	20 μM	
5	5023		1	
	5026	34	10	
	5027	12	8	
	5028	11	4	
	5030	20	7	6
	5032	65	62	63
10	5040	0	1	0
	5041	1	0	0
	5042	77	64	42
	5043	21	15	1
	5048	55	19	11
	5052	77	76	86
15	5053	68	64	56
	5054	5	57	48
	5055	69	69	70
	5057	44	29	37
	5061	43	48	60
	5062	78	81	87
20	5069	70	71	75
	5071	80	82	73
	5072	60	61	61
	5073	63	70	14
	5074	47	57	26
	5075	88	88	52
25	5077	34	46	42
	5078	60	67	11
	5079	44	58	14
	5081	25	34	50
	5188	90		94
	5200	10		10
30	5205	56		3.50
			33	100.0

	5206	72	78	3.0
	5299			7.00
	5324			9.00
5	5327		17	
	5335			22.0
	5367			18.00
	5371			12.00
	5376			12.00
10	5379		65	15.00
	5386			18.00
	5388		58	9.00
	5388.HCl		60	12.00
	5389		55	2.50
15	5389.HCl		57	2.50
	5391		64	6.50
	5391.HCl		100	3.50
	5393		76	14.00
	5393.HCl		58	20.00
20	5394		59	16.00
	5394.HCl		62	17.00
	5397		42	
	5397.HCl		21	
	5402		37	
25	5402.HCl		37	

TABLE 2

Compound No.	IC50 (μ m)
1470	50.0 - 100.0
1471	25.0 - 50.0
1474	25.0 - 50.0
1476	50.0 - 100.0
1506	25.0 - 50.0
1507	25.0 - 50.0

	1509	50.0 - 100.0
	1542	50.0 - 100.0
	1560	50.0 - 100.0
	1618	50.0 - 100.0
5	1652	25.0 - 50.0
	1657	25.0 - 50.0
	1672	50.0 - 100.0
	1676	12.0 - 25.0
	1693	50.0 - 100.0
10	1718	50.0 - 100.0
	1808	25.0 - 12.0
	1809	25.0 - 12.0
	1845	10.0 - 5.0
	1888	50.0 - 100.0
15	1910	5.0 - 10.0
	1912	25.0 - 50.0
	1921	100.0 - 50.0
	1928	25.0 - 50.0
	1929	25.0 - 12.0
20	1930	25.0 - 12.0
	1982	50.0 - 25.0

25


Reference Example 1: Preparation of (3Z)-1-acetyl-3-benzylidene-2,5-piperazinedione

1,4-Diacetyl-2,5-piperazinedione (25.0g, 126 mmol), which is compound (8) mentioned in Reference Example 3, was heated at 120-130°C in DMF (200 ml) with triethylamine (17.6 ml, 126 mmol) and benzaldehyde (13.0 ml, 126 mmol). After 4 h the mixture was cooled to room temperature and

poured into EtOAc (1000 ml), and washed three times with
 brine. Any solid formed at this stage was filtered off.
 The filtrate was dried ($MgSO_4$) and the solvent removed in
vacuo. The residue was recrystallised from EtOAc:Hexane to
 give 11.78 g (38%) of the title compound as a yellow solid.
¹H NMR ($CDCl_3$, 400 MHz) δ =2.69 (3H, s) 4.54 (2H,
 s) 7.20 (1H, s) 7.40 (3H, m), 7.48 (2H, m),
 7.93 (1H, br.s)
 MS (DCI, NH_3): 262 (MNH_4^+ , 20%), 245 (MH^+ , 53%),
 220 (52%), 204 (100%), 203 (100%)

Microanalysis	C	H	N
Calc	63.93	4.95	11.47
Found	64.11	5.02	11.41
Found	64.05	4.90	11.44

Alternatively (3Z)-1-acetyl-3-benzylidene-2,5-piperazinedione can be produced as follows:



Compound 16 is treated with ammonia and subsequently with acetic anhydride to yield the title compound.

Reference Example 2: Preparation of (3Z)-1-acetyl-3-(4-acetamidobenzylidene)-2,5-piperazinedione

1,4-Diacetyl-2,5-piperazinedione (10.0g, 50 mmol), prepared by the published procedure mentioned in Example 3, was stirred in DMF (40 ml) with 4-acetamidobenzaldehyde (8.24 g, 50 mmol) and triethylamine (7 ml, 50 mmol) and heated to 120°C. After 2½ h the mixture was cooled to room temperature, diluted with EtOAc (100 ml) and stirred overnight. The solid formed was collected, washed with EtOAc and dried to give 8.46 g (56%) of a yellow solid.


15

¹H NMR (CDCl₃+TFA, 400 MHz) δ=2.32 (3H, s) 2.72 (3H, s) 4.68 (2H, s) 7.36 (1H, s) 7.45 (2H, d, J=8Hz) 7.60 (2H, d, J=8Hz)

20


Microanalysis	C	H	N
Calc	59.80	5.02	13.95
Found	60.08	5.09	13.89
	60.11	5.07	13.86

25 Reference Example 3: Preparation of 1,4-Diacetyl-2,5-piperazinedione



1,4-Diacetyl-2,5-piperazine dione (8) was prepared by the published procedure (S.M. Marcuccio and J.A. Elix, Aust. J. Chem., 1984, 37, 1791).

5 Reference Example 4: (3Z)-1-Acetyl-3-(4-
methoxybenzylidene)-2,5-
piperazinedione



15 (3Z)-1-Acetyl-3-(4-methoxybenzylidene)-2,5-piperazinedione (9) was prepared by the published procedure (T. Yokoi, L-M. Yang, T. Yokoi, R-Y. Wu, and K-H. Lee, J. Antibiot., 1988, 41, 494).

20 Reference Example 5: Preparation of (3Z)-1-acetyl-3-(2,6-dichlorobenzylidene)-2,5-piperazinedione

1,4-Diacetyl-2,5-piperazinedione prepared by the published procedure mentioned in Reference Example 3, was stirred in DMF with 2,6-dichlorobenzaldehyde and triethylamine and heated to 120-130°C for 1-3h. The title compound was obtained with a yield of 40%.

Reference Example 6: Preparation of (3Z)-1-acetyl-3-(4-(3-dimethylamino)propoxybenzylidene)-2,5-piperazinedione

5 1,4-Diacetyl-2,5-piperazinedione, prepared by the published procedure mentioned in Reference Example 3, was stirred in DMF with 4-(3-dimethylamino)propoxybenzaldehyde and triethylamine and heated to 120-130°C for 2-4h to give the title compound.

10 By the same method, using 4-(2-dimethylamino)ethoxybenzaldehyde in place of the above-mentioned aldehyde, (3Z)-1-acetyl-3-(4-(2-dimethylamino)ethoxybenzylidene)-2,5-piperazinedione was prepared.

15

Reference Example 7: (3Z,6Z)-3-(4-Hydroxybenzylidene)-6-(4-methoxybenzylidene)-2,5-piperazinedione

20 (3Z,6Z)-3-(4-Acetoxybenzylidene)-6-(4-methoxybenzylidene)-2,5-piperazinedione was treated with aqueous sodium hydroxide in THF at room temperature for 8 hrs to give the title compound (1519) in 30% yield.

Example 1: Preparation of 1470

25 3(Z)-1-Acetyl-3-benzylidene-2,5-piperazinedione (one equivalent), which is compound 18 prepared according to Reference Example 1, was treated with 1-tert-butoxycarbonylpyrrole-2-carboxaldehyde in the presence of

Cs_2CO_3 (1-1.1 equivalents) in DMF at 80-100°C for 1-6 hours. The title compound was obtained in 24% yield.

The crude product was optionally, washed with water, methanol, ethyl acetate or diethylether and optionally 5 recrystallised from methanol as appropriate.

By the same method, but replacing 1-tert-butoxycarbonylpyrrole-2-carboxaldehyde by the appropriately substituted aldehyde or benzaldehyde, the following compounds were prepared:

	Compound	Yield (%)
10	1471	52
	1652	37
	1983	45
	1921	54
	1922	43
	1924	44
	1910	31
	1926	27
	1927	26
	1928	20
15	1929	-
	1930	-
	1912	33
	5032	50
	5040	45
	5043	24
	5053	44
	5054	22
	5057	43
	5058	16
20		
25		
30		

Example 2: Preparation of 1474

3 (Z)-1-Acetyl-3-(4-methoxybenzylidene)-2,5-piperazinedione prepared according to Reference Example 4, was treated with 2-thiophenecarboxaldehyde in the presence 5 of Cs₂CO₃ (1-1.1 equivalents) in DMF at 80-100°C for 1-6 hours. The title compound was obtained in 76% yield.

By the same method, but replacing 2-thiophenecarboxaldehyde by the appropriately substituted aldehyde, the following compounds were prepared:

10

Compound	Yield (%)
1476	54
1479	84
1506	67
1507	7

15

The crude product was optionally washed with water, methanol, ethyl acetate and diethylether and optionally recrystallised from acetic acid or methanol as appropriate.

20 Example 3: Preparation of 1884

3 (Z)-1-Acetyl-3-benzylidene-2,5-piperazinedione (1 equivalent), prepared according to Reference Example 1, was treated with cyclohexanecarboxaldehyde (4 equivalents) in the presence of 0.5M potassium tert-butoxide in tertiary 25 butanol (2 equivalents) in DMF at 0-100°C for 2 hours. The title compound was obtained with a yield of 58%. Purification was effected by recrystallisation from acetic acid.

1672 was prepared as above but replacing the 3 (Z)-1-

acetyl-3-benzylidene-2,5-piperazinedione with 3(Z)-1-acetyl-3-(4-acetamidobenzylidene)-2,5-piperazinedione. The reaction was maintained for 18 hours. A low yield was obtained.

5

Example 4: Preparation of 1676

1-Acetyl-3-(4-acetamidobenzylidene)-2,5-piperazinedione (one equivalent), prepared according to Reference Example 2, was treated with cinnamaldehyde in the presence of Cs_2CO_3 (1-1.1 equivalents) in DMF at 80-100°C for 1-6 hours. The title compound was obtained in 46% yield.

15 Example 5: Preparation of 1618

1,4-Diacetyl-2,5-piperazinedione, prepared by the published procedure mentioned in Reference Example 3, was stirred in DMF with 2-thiophenecarboxaldehyde (1 equivalent) and triethylamine (1 equivalent) at 120°C for 2-4h. (3Z)-1-Acetyl-3-(2-thenylidene)-2,5-piperazinedione was obtained with a yield of 36%.

(3Z)-1-Acetyl-3-(2-thenylidene)-2,5-piperazinedione (1 equivalent) was stirred in DMF with 3-1-tert-butoxycarbonylindole-3-carboxyaldehyde (1 equivalent) in the presence of Cs_2CO_3 (1-1.1 equivalents) at 80-100°C for 2-3h. The title compound was obtained with a yield of 14%.

Example 6: Preparation of 1542

3 (Z)-1-Acetyl-3-(2,6-dichlorobenzylidene)-2,5-piperazinedione (1 equivalent), prepared according to Reference Example 5 was treated with 3-furaldehyde (1 5 equivalent) in the presence of Cs₂CO₃ (1-1.1 equivalents) in DMF at 80-100°C for 2-5 hours. The title compound was obtained in 46% yield.

By the same method, but replacing 3-furaldehyde by the appropriately substituted aldehyde, 1560 was obtained 10 with a yield of 39%.

Example 7: Preparation of 1982

3 (Z)-1-Acetyl-3-benzylidene-2,5-piperazinedione (1 equivalent), as prepared in Reference Example 1, was 15 treated with 4-(N-(3-dimethylaminoethyl)-N-methyl)aminomethylbenzaldehyde in the presence of Cs₂CO₃ (1-1.1 equivalents) in DMF at 80-100°C for 1-6h to give (3Z,6Z)-3-Benzylidene-6-(4-(N-dimethylaminoethyl)-N-methyl)aminomethylbenzylidene)-2,5-piperazinedione in a 20 yield of 50%.

Compound 1982, the hydrochloride salt of (3Z,6Z)-3-Benzylidene-6-(4-(N-(3-dimethylaminoethyl)-N-methyl)aminomethylbenzylidene)-2,5-piperazinedione, was prepared by bubbling HCl gas through a solution of the 25 corresponding free base in THF, followed by evaporation to dryness. The yield was 45%.


Example 8: Preparation of 1976

3 (Z)-1-Acetyl-3-(4-(3-dimethylamino)propoxybenzylidene)-2,5-piperazinedione (1 equivalent), prepared according to Reference Example 6 was 5 treated with 3-(imidazol-1-yl)benzaldehyde (1 equivalent) in the presence of Cs_2CO_3 (1-1.1 equivalent) in DMF at 80-90°C for 2-4 hours. The title compound was obtained in 52% yield.

10 Example 9: Preparation of 1886

1519 (1 equivalent), prepared in Reference Example 7, was treated in DMF with sodium hydride (1 equivalent) and N-phthaloylglycyl chloride (1 equivalent) in DMF at room temperature for 4h. The title compound was obtained with a 15 yield of 30%.

Example 10: Preparation of 5026




(3Z)-1-acetyl-3-(4-(3-dimethylamino)propoxybenzylidene)-2,5-piperazinedione, prepared as in Reference Example 6, was treated with compound 10.1 in dimethylformamide (DMF) in the presence of Cs_2CO_3 at a 5 temperature of 80°C-90°C for 2-4 hours. Compound 5026 was obtained in 95% yield.

By an analogous process, using the appropriately substituted benzaldehyde in place of compound 10.1, the following compounds were prepared:

10	Compound No.	Yield %
	5030	30
	5048	72
	5188	70

15 Example 11: Preparation of 5027




(3Z)-1-acetyl-3-(4-(3-dimethylamino)propoxybenzylidene)-2,5-piperazinedione, prepared as in Reference Example 6, was treated with compound 11.1 in DMF in the presence of Cs_2CO_3 at 80°C-90°C for 2-4 hours. Compound 5027 was 5 produced in 33% yield.

By the same method, but replacing 11.1 by the appropriately substituted aldehyde, the following compounds were prepared:

	Compound No.	Yield (%)
10	5028	44
	5029	25
	5041	39
	5042	39
	5046	37
15	5052	58


Example 12: Preparation of 5023



Compound 12.1 was treated with 4-(3-dimethylamino)propoxybenzaldehyde in DMF in the presence of Cs₂CO₃ at a temperature of 80°C-90°C for 2-4 hours. Compound 5023 was obtained in 36% yield.

5

Example 13: Preparation of 5062




20 (3Z)-1-acetyl-3-(4-(2-dimethylamino)ethoxybenzylidene)-2,5-piperazinedione, prepared as in Reference Example 6, was treated with compound 13.1 in DMF in the presence of Cs₂CO₃ at a temperature of 80°C-90°C for 2-4 hours. Compound 5062 was obtained in 12% yield.

25 By the same method, but using the appropriately substituted aldehyde in place of compound 13.1, the following compounds were prepared:

Compound No.	Yield (%)
5071	41
5072	86

5

Example 14: Preparation of compounds of formula (I)



Compound (I)

- 20 The 2,5-piperazinedione derivative 14.1 was treated with the aldehyde 14.2, the groups Ar and Subst. being as specified below, in DMF in the presence of Cs_2CO_3 at 80°C-90°C for 2-4 hours. The compounds of formula (I) listed below were prepared:


25

Ar	Subst.	Compound of formula (I)	Yield (%)	
Phenyl	-CH ₂ S(CH ₂) ₂ NMe ₂	5058	16	
3-furyl	-CH ₂ S(CH ₂) ₂ NMe ₂	5073	33	
3-thienyl	-CH ₂ S(CH ₂) ₂ NMe ₂	5078	38	
5	3-thienyl	-CH ₂ NHC(O)CH ₂ NMe ₂	5074	83
2-bromophenyl	-CH ₂ NHC(O)CH ₂ NMe ₂	5079	28	
3-furyl	-CH ₂ NHC(O)CH ₂ NMe ₂	5081	68	
3-thienyl	-CH ₂ O(CH ₂) ₂ NMe ₂	5069	29	
3-furyl	-CH ₂ O(CH ₂) ₂ NMe ₂	5077	20	

10

Example 15: Preparation of compounds of formula (I)


15



15.1

15.2

20



Compound (I)


The 2,5-piperazinedione derivative 15.1 was treated with the aldehyde 15.2 in which R₂₀ and R₂₁ are both H or are both OMe, the substituent Ar and linking group A being as specified below, in DMF in the presence of Cs₂CO₃ at 80°C to 5 90°C for 2-4 hours. The compounds of formula (I) listed below were prepared. In 5391, 5394 and 5371 R₂₀ and R₂₁ are both H. In 5393 and 5402 R₂₀ and R₂₁ are OMe.

	Ar	A	Compound of Formula (I)	Yield (%)
10	Phenyl	- (CH ₂) ₂ -	5391	21
	Phenyl	- (CH ₂) ₃ -	5394	47
	Phenyl	- (CH ₂) ₄ -	5371	56
15	Phenyl		5393	44
	4-nitrophenyl		5402	62

15


Example 16: Preparation of compounds of formula (I)

20



25

25



(3Z)-1-acetyl-3-benzylidene-2,5-dione prepared as in Reference Example 1 (compound 18), was treated with the aldehyde 16.1 in which substituent Y was as indicated 5 below, in DMF in the presence of Cs_2CO_3 at 80°C-90°C for 2-4 hours. The compounds of formula (I) listed below were prepared:

Y	Compound of formula (I)	Yield %)
5-O(CH ₂) ₂ NMe ₂	5324	34
10 4-O(CH ₂) ₂ NMe ₂	5327	51
5-(CH ₂) ₂ NMe ₂	5335	45
5-O(CH ₂) ₂ O(CH ₂) ₂ NMe ₂	5388	12
5-O(CH ₂) ₆ NMe ₂	5389	35
15 5-N(Me)(CH ₂) ₂ NMe ₂	5299	2

15

By the same method, but using 2,5-dichlorothiophene-4-carboxaldehyde in place of compound 16.1, 5075 was prepared in 31% yield.

20 Example 17: Preparation of salts

1. Hydrochloride salts of the following compounds of formula (I) were prepared by bubbling HCl gas through a solution of the corresponding free base in tetrahydrofuran 25 (THF) at room temperature. The salt was recovered in the yield indicated.

	Compound of formula (I)	Hydrochloride salt	Yield (%)
	1975	5026	95
	1976	5030	30
5	5048	5048.HCl	72
	5188	5206	24
	5200	5205	31
	5367	5376	47
10	5397	5397.2HCl	36
	5041	5041.HCl	63
	5042	5042.HCl	51
	5046	5046.HCl	32
	5052	5052.HCl	58
	5023	1988	50
15	5062	5062.HCl	-
	5071	5071.HCl	-
	5072	5072.HCl	-
	1910	5055	57
	1912	5061	47
20	5032	5032.HCl	39
	5053	5053.HCl	90
	5054	5053.HCl	88
	5073	5073.HCl	76
	5078	5078.HCl	78
25	1912	5061	47
	5074	5074.HCl	51
	5079	5079.HCl	73
	5081	5081.HCl	76
	5069	5069.HCl	-
30	5077	5077.HCl	-
	5324	5324.HCl	68
	5336	5336.HCl	74
	5335	5335.HCl	-

	5388	5388.HCl	79
	5389	5389.HCl	75
	5391	5391.HCl	-
	5394	5394.HCl	75
5	5371	5379	65

2. Hydrochloride salts of the following compounds of formula (I) were prepared by bubbling HCl gas through a solution of the corresponding free base in hot DMF. The salt was recovered in the yield indicated.

	Compound of formula (I)	Hydrochloride salt	Yield
	5386	5386.2HCl	79
15	5393	5393.HCl	60
	5402	5402.HCl	52

3. Hydrochloride salts of the following compounds of formula (I) were prepared by treating the free base with 2M HCl:

	Compound of formula (I)	Hydrochloride salt	Yield (%)
	5027	5027.HCl	67
25	5028	5028.HCl	92
	5029	5029.HCl	76
	5040	5040.HCl	90

4. 5043.HCl, the hydrochloride salt of 5043, was prepared by bubbling HCl gas through a solution of 5043 in MeOH. 5057.HCl, the salt of 5057, was prepared by bubbling HCl gas through a solution of 5057 in THF following by 5 recrystallisation from MeOH.

Example 18: PHARMACEUTICAL COMPOSITION

Tablets, each weighing 0.15 g and containing 25 mg of 10 a compound of the invention can be manufactured as follows:

Composition for 10,000 tablets

compound of the invention (250 g)

lactose (800 g)

corn starch (415 g)

15 talc powder (30 g)

magnesium stearate (5 g)

The compound of the invention, lactose and half of the corn starch are mixed. The mixture is then forced through a sieve 0.5 mm mesh size. Corn starch (10 g) is 20 suspended in warm water (90 ml). The resulting paste is used to granulate the powder. The granulate is dried and broken up into small fragments on a sieve of 1.4 mm mesh size. The remaining quantity of starch, talc and magnesium stearate is added, carefully mixed and processed into 25 tablets.

Example 19: Characterisation of compounds of formula

A

The compounds prepared in the preceding Examples,
were characterised by mass spectroscopic, microanalytical,
5 proton nuclear magnetic resonance and, in some cases,
infra-red techniques. The results are set out in the
Tables which follow:

No.	Mol. Formula ^a	Mass spec. data			¹ H nmr data		
		mass (intensity)	mode	solvent (field)	δ		
1910	C ₂₃ H ₂₀ N ₄ O ₃	401(100)	CI	d ₆ -DMSO/400MHz	4.28-4.32 (2H,t), 6.75-7.70 (14H,m), 10.15 (2H,brs).		
5023	C ₂₆ H ₃₂ N ₄ O ₃	449(100)	EI	CDCl ₃ /400MHz	2.00 (2H,m), 3.45 (2H,s), 4.05 (2H,t), (10H,m), 8.15 (2H,brs).	2.25 (12H,s), 2.46 (2H,t), 6.95-7.42	
5026	C ₂₇ H ₂₉ N ₅ O ₃ .2HCl			d ₆ -DMSO/400MHz	2.12 (2H,m), 4.11 (2H,t), 7.00 (2H,d), 7.55 (2H,d), 9.21 (1H,s), (1H,brs).	2.73 (6H,s), 5.48 (2H,s), 7.47 (2H,d), 7.65 (1H,s), 7.77 (1H,s), 10.12 (2H,brs), 10.45	21 (2H,m); 6.76 (2H,s); 7.50 (2H,d); 7.77 (1H,s);
5027	C ₂₂ H ₂₄ N ₄ O ₃ .2HCl			CDCl ₃ +CF ₃ CO ₂ H/400 MHz	2.00 (2H,t), 3.90 (2H,t), 7.35 (1H,s), 8.95 (2H,d).	3.00 (6H,s), 7.00 (2H,d), 7.45 (2H,d), 8.00 (2H,d).	3.45 (2H,m); 7.15 (1H,s);
5028	C ₂₂ H ₂₄ N ₄ O ₃ .2HCl			CDCl ₃ +CF ₃ CO ₂ D/400MHz	2.35 (2H,m), 4.15 (2H,t), 7.30 (1H,s), 8.50 (1H,d),	3.00 (6H,s), 7.00 (2H,d), 7.45 (2H,d), 8.95 (1H,d), 9.15 (1H,s).	3.45 (2H,t), 7.15 (1H,s);
5030				d ₆ -DMSO/400MHz	2.18 (2H,m), 4.10 (2H,t), 7.00 (2H,d), 7.71 (1H,m), 8.29 (1H,s), (1H,brs),	2.77 (6H,s), 6.77 (1H,s), 7.51 (2H,d), 7.85 (1H,s), 9.60 (1H,s), 10.50 (1H,brs), (1H,brs).	3.20 (2H,m); 6.81 (1H,s); 7.65 (2H,m); 7.96 (1H,s); 10.21

No.	Mol. Formula	Mass spec. data		1H nmr data	
		mass (intensity)	mode	solvent (field)	δ
5032	C ₂₃ H ₂₅ N ₃ O ₄ .HCl	408(20), 306(30)	CI	d ₆ -DMSO/400MHz	2.83 (6H,s), 4.30 (1H,m), 6.78 (1H,s), 7.42 (2H,m); 10.12 (2H,br).
5040	C ₂₅ H ₂₇ N ₃ O ₅ .HCl	450(10)	CI	d ₆ -DMSO/400MHz	3.20-3.55 (6H,m), (2H,d), 4.39 (1H,m), (1H,s), 6.78 (1H,s), (1H,m), 7.45 (2H,m), (3H,br)
5041	C ₂₁ H ₂₃ N ₃ O ₄ .HCl	382(100)	EI	d ₆ -DMSO/400MHz	2.09 (2H,m), 4.09 (2H,t), 6.78 (1H,s), 7.54 (2H,d), 9.75 (1H,brs),
5042	C ₂₁ H ₂₃ N ₃ O ₃ S.HCl	398(35)	EI	d ₆ -DMSO/400MHz	2.09 (2H,m), 4.10 (2H,t), 7.00 (2H,d), 7.62 (1H,m), 9.92 (1H,brs),
5043	C ₂₂ H ₃₂ N ₄ O ₅ .HCl	493(100)	CI	d ₆ -DMSO/400MHz	3.10-3.85 (14H,m), (1H,brs), 6.77 (1H,s), (2H,d), 7.32 (1H,m), (4H,m), 10.20 (2H,s).
5046	C ₂₁ H ₂₃ N ₃ O ₃ S.HCl	398(23), 169(100)	EI	d ₆ -DMSO/400MHz	2.09 (2H,m), 4.10 (2H,t), 7.00 (2H,d), 7.58 (1H,m), 10.16 (1H,brs).

No.	Mol. Formula	Mass spec. data mass (intensity)	mode	solvent (field)	¹ H nmr data	
					δ	
5048	C ₂₅ H ₂₈ N ₄ O ₄ .HCl	485 (100)	EI	d ₆ -DMSO/400MHz	2.05 (2H,s), 3.20 (2H,m), 6.75 (1H,s), 7.51 (2H,d), 10.15 (1H,brs).	2.14 (2H,m), 4.13 (2H,t), 7.0 (2H,d), 7.62 (2H,d), 10.20 (1H,brs).
5052				d ₆ -DMSO/400MHz	2.15 (2H,m), 4.10 (2H,t), 6.94 (1H,s), 7.76 (1H,s).	2.28 (6H,s), 6.68 (1H,s), 7.00 (2H,d), 8.23 (1H,s).
5053	C ₂₄ H ₂₅ N ₃ O ₃ .HCl			CDCl ₃ +CF ₃ CO ₂ D/400MHz	2.20 (4H,m), 4.00 (2H,m), 7.23 (1H,s).	3.20 (2H,m), 4.45 (2H,m), 7.39 (1H,s).
5054	C ₂₄ H ₂₅ N ₃ O ₄ .HCl			CDCl ₃ +CF ₃ CO ₂ D/400MHz	3.25 (2H,m), 4.05-4.20 (4H,m), (2H,d) 7.20 (1H,s), 7.51 (7H,m).	3.67 (2H,m), 4.47 (2H,m), 7.26 (1H,s), 7.39-
5055	C ₂₃ H ₂₀ N ₄ O ₃ .HCl	401(100)	ESI	d ₆ -DMSO/400MHz	4.40 (2H,t), 6.75 (1H,s), (7H,m), (1H,s), 10.20 (1H,brs)	4.60 (2H,t), 6.99 (2H,d), 7.65 (1H,s), 10.10 (1H,s), 10.20 (1H,brs).
5057	C ₂₄ H ₂₂ N ₄ O ₄ .HCl			d ₆ -DMSO/400MHz	4.00-4.05 (2H,m), (1H,m), (2H,d), (4H,m), (1H,s),	4.20-4.32 (2H,m), 6.77 (1H,s), 7.32 (2H,m), 7.71 (1H,m), 10.20 (2H,brs).
						4.48 7.03 7.42 (2H,m), 7.55 9.12

No.	Mol. Formula	Mass spec. data		^1H nmr data	
		mass (intensity)	mode	solvent (field)	δ
5058.HCl	$\text{C}_{23}\text{H}_{25}\text{N}_3\text{O}_2\text{S} \cdot \text{HCl}$	409(15)	CI	d_6 -DMSO/400MHz	2.70-2.75 (8H, m), 3.20-3.25 (2H, m), 3.85 (2H, s); 6.78 (2H, s); 7.32-7.55 (9H, m), 9.68 (1H, brs), 10.22 (1H, s), 10.24 (1H, s)
5061	$\text{C}_{23}\text{H}_{24}\text{N}_4\text{O}_3 \cdot \text{HCl}$			d_6 -DMSO/400MHz	2.84 (6H, s), 3.95 (2H, s), 4.40 (2H, d), 6.75 (1H, s), 6.77 (1H, s), 7.33-7.55 (9H, m), 9.15 (1H, t), 9.85 (1H, brs), 10.20 (1H, brs), 10.25 (1H, brs).
5062	$\text{C}_{20}\text{H}_{21}\text{N}_3\text{O}_4 \cdot \text{HCl}$			d_6 -DMSO/400MHz	2.76 (6H, d), 3.51 (2H, m), 4.38 (2H, t), 6.66 (1H, s), 6.75 (1H, s), 6.91 (1H, s), 7.05 (2H, d), 7.55 (2H, d), 7.74 (1H, s), 8.22 (1H, s), 9.76 (1H, s).
5069	$\text{C}_{21}\text{H}_{23}\text{N}_3\text{O}_3\text{S} \cdot \text{HCl}$	397(10)	CI	d_6 -DMSO/400MHz	2.80 (6H, s), 3.30 (2H, t), 3.76 (2H, t), 4.58 (2H, s), 6.82 (1H, s), 6.87 (1H, s), 7.45 (2H, m), 7.58 (2H, d), 7.65 (1H, m), 8.00 (1H, s), 9.78 (1H, s), 10.02 (1H, s), 10.18 (1H, s).
5071	$\text{C}_{20}\text{H}_{21}\text{N}_3\text{O}_3\text{S} \cdot \text{HCl}$			d_6 -DMSO/400MHz	2.86 (6H, d), 3.53 (2H, m), 4.38 (2H, t), 6.78 (1H, s), 6.84 (1H, s), 7.07 (2H, d), 7.43 (1H, m), 7.58 (2H, d), 7.65 (1H, m), 7.96 (1H, m), 9.55 (1H, s), 10.05 (1H, brs), 10.13 (1H, brs).
5072	$\text{C}_{21}\text{H}_{23}\text{N}_3\text{O}_3\text{S}_2 \cdot \text{HCl}$			d_6 -DMSO/400MHz	2.58 (3H, s), 2.78 (6H, s), 3.44 (2H, m), 4.36 (2H, t), 6.77 (1H, s), 6.85 (1H, s), 7.05 (2H, d), 7.12 (1H, d), 7.52 (1H, d), 7.58 (2H, d), 10.20 (1H, s).

No.	Mol. Formula ^a	Mass spec. data		¹ H nmr data		
		mass (intensity)	mode	solvent (field)	δ	
5073	C ₂₁ H ₂₃ N ₃ O ₃ S	398(15); 293(100)	EI	CDCl ₃ +CF ₃ CO ₂ D/400MHz	2.75 (2H,t), 3.78 (2H,s), 7.40 (4H,s),	2.90 (6H,s), 6.70 (1H,s), 7.10 (1H,s), 7.85 (1H,s).
5073.HCl	C ₂₁ H ₂₃ N ₃ O ₃ S.HCl			d ₆ -DMSO/400MHz	2.75 (6H,s), (2H,m), (1H,s), (2H,d), (1H,brs),	2.75-2.80 (2H,m), 3.20 (2H,s), 6.70 (1H,s), 7.40 (2H,d), 8.20 (1H,s), 10.00 (1H,brs),
5074				d ₆ -DMSO/400MHz	2.82 (6H,s), 6.81 (1H,s), 9.15 (1H,brs),	4.00 (2H,s), 6.88 (1H,s), 9.90 (1H,brs),
5075	C ₁₆ H ₁₀ Cl ₂ N ₂ O ₂ S			d ₆ -DMSO/400MHz	7.39-7.45 (3H,m),	4.41 (2H,d), 7.98 (2H,m), 10.04 (1H,brs).
5077	C ₂₁ H ₂₃ N ₃ O ₄ .HCl			d ₆ -DMSO/400MHz	2.55 (2H,t), 4.55 (2H,s), 6.95 (1H,s), 7.85 (1H,s), 10.01 (1H,s),	2.80 (6H,s), 6.70 (1H,s), 7.45 (2H,d), 8.30 (1H,s), 9.90 (1H,s),
5078	C ₂₁ H ₂₃ N ₃ O ₂ S ₂	414(15); 309(100)	EI	CDCl ₃ +CF ₃ CO ₂ D/400MHz	2.75 (2H,t), 3.88 (2H,s), (4H,s),	2.88 (6H,s), 7.22-7.28 (3H,m), 7.50-7.54 (1H,m), (1H,s).
						3.25 (2H,t), 7.45 (-7.66 (1H,s).

No.	Mol. Formula	Mass spec. data		¹ H nmr data		
		mass (intensity)	mode	solvent (field)	δ	
5078.HCl	C ₂₁ H ₂₃ N ₃ O ₂ S ₂ .HCl		d ₆ -DMSO/400MHz	2.72-2.78 (2H,m), 2.75 (6H,s), 3.20-3.25 (2H,m), 3.84 (2H,s), 6.75 (1H,s), 6.85 (1H,s), 7.40-7.45 (3H,m), 7.55 (2H,d); 7.64-7.67 (1H,m), 7.96-7.99 (1H,m), 9.85 (1H,brs), 10.05 (1H,brs), 10.18 (1H,brs).		
5079	C ₂₃ H ₂₃ BrN ₄ O ₃ .HCl		d ₆ -DMSO/400MHz	2.82 (6H,s), 4.00 (2H,s), 4.41 (2H,d); 6.74 (1H,s), 6.80 (1H,s), 7.30 (1H,m); 7.36 (2H,d), 7.45 (1H,m), 7.54 (2H,d); 7.60 (1H,d), 7.68 (1H,d) 9.56 (1H,brt); 9.90 (1H,brs), 10.36 (1H,brs), 10.48 (1H,brs).		
5081	C ₂₁ H ₂₂ N ₄ O ₄ .HCl		d ₆ -DMSO/400MHz	2.83 (6H,s), 4.01 (2H,s), 4.39 (2H,d); 6.68 (1H,s), 6.79 (1H,s), 6.94 (1H,s); 7.35 (2H,d), 7.54 (2H,d), 7.76 (1H,s); 8.22 (1H,s), 9.12 (1H,brt), 9.82 (2H,brs) 10.12 (1H,brs).		
5188	C ₂₇ H ₂₇ N ₃ O ₃	442(100)	ESI	d ₆ -DMSO/400MHz	1.8-1.9 (2H,m), 2.15 (6H,s), 2.38 (2H,t), 4.05 (2H,t), 6.78 (1H,s), 6.90 (1H,s), 6.99 (2H,d), 7.50-7.58 (4H,m); 7.61-7.65 (1H,m), 7.39-7.98 (3H,m), 8.11 (1H,s), 10.28 (2H,brs).	
5200	C ₂₇ H ₂₇ N ₃ O ₃	442(100)	ESI	d ₆ -DMSO/400MHz	1.81-1.91 (2H,m), 2.15 (6H,s), 2.35 (2H,t), 4.09 (2H,t), 6.75 (1H,s), 6.96 (2H,d), 7.21 (1H,s), 7.5-7.65 (7H,m), 7.94 (2H,d), 10.15 (2H,brs).	

No.	Mol. Formula	Mass spec. data		solvent (field)	¹ H nmr data	
		mass (intensity)	mode		δ	
5205	C ₂₇ H ₂₇ N ₃ O ₃ .HCl	442(40)	CI	d ₆ -DMSO/400MHz	2.12-2.20 (2H,m), 2.80 (6H,s), 3.20-3.25 (2H,m), 4.10 (2H,t), 6.75 (1H,s), 7.01 (2H,d), 7.24 (1H,s), 7.51-7.67 (6H,m), 7.92 (2H,d), 7.98-8.01 (1H,m), 10.1 (2H,brs), 10.25 (1H,brs).	
5206	C ₂₇ H ₂₇ N ₃ O ₃ .HCl			d ₆ -DMSO/400MHz	2.11-2.21 (2H,m), 2.60 (6H,s), 2.85-2.98 (2H,m), 4.09 (2H,t), 6.78 (1H,s), 6.94 (1H,s), 7.0 (2H,d), 7.50-7.59 (4H,m), 7.64 (1H,d), 7.90-7.99 (3H,m), 8.12 (1H,m), 10.21 (1H,brs), 10.43 (1H,brs).	
5324	C ₂₀ H ₂₁ N ₃ O ₃ S.HCl	384(100)	CI	d ₆ -DMSO/400MHz	2.85 (6H,s), 3.52 (2H,t), 4.50 (2H,t), 6.52 (1H,d), 6.78 (1H,s), 6.81 (1H,s), 7.31 (1H,d), 7.32 (1H,m), 7.45 (2H,m), 7.57 (2H,d), 9.70 (1H,s), 10.15 (1H,s), 10.41 (1H,brs).	
5327	C ₂₀ H ₂₁ N ₃ O ₃ S	384(20)	CI	d ₆ -DMSO/400MHz	2.22 (6H,s), 2.63 (2H,t), 4.05 (2H,t), 6.76 (1H,s), 6.82 (2x1H,s), 7.30 (1H,s), 7.33 (1H,m), 7.42 (2H,m), 7.55 (2H,d).	
5335	C ₂₀ H ₂₁ N ₃ O ₂ S.HCl	368(20)	CI	d ₆ -DMSO/400MHz	2.78 (6H,s), 3.28 (4H,m), 6.78 (1H,s), 6.89 (1H,s), 7.02 (1H,d), 7.38-7.45 (4H,m), 7.55 (2H,d), 9.68 (1H,brs), 10.40 (1H,br).	
5336	C ₂₀ H ₂₁ N ₃ O ₃ S.HCl	384(10)	CI	d ₆ -DMSO/400MHz	2.82 (6H,s), 3.49 (2H,t), 4.38 (2H,t), 6.78 (1H,s), 6.80 (1H,s), 6.94 (1H,s), 7.31 (1H,s), 7.32 (1H,m), 7.42 (2H,m), 7.55 (2H,d), 9.78 (1H,s), 10.25 (1H,s), 10.45 (1H,brs).	

No.	Mol. Formula	Mass spec. data		^1H nmr data	
		mass (intensity)	mode	solvent (field)	δ
5367	$\text{C}_{33}\text{H}_{34}\text{N}_4\text{O}_4$	551(100)	CI	$\text{CDCl}_3+\text{CF}_3\text{CO}_2\text{D}/400\text{MHz}$	1.72 (2H,m), 1.95-2.01 (2H,m), 2.24 (6H,m), 2.48 (2H,t), 2.96 (2H,m), 3.70 (1H,m), 4.07 (2H,t), 4.89 (1H,m), 7.0 (2H,d), 7.01 ((2H,s), 7.15-7.25 (4H,m), 7.35 (2H,d), 7.48 (2H,d), 7.57 (2H,d), 8.17 (2H,brs).
5371	$\text{C}_{32}\text{H}_{32}\text{N}_4\text{O}_3$	521(100)	CI	$\text{CDCl}_3/400\text{MHz}$	1.75-1.80 (4H,m), 2.55-2.60 (2H,m), 2.75 (2H,t), 2.88 (2H,t), 3.50-3.55 (2H,m), 3.65 (2H,s), 6.95 (1H,s), 6.98-7.02 (1H,m), 7.05-7.10 (4H,m), 7.15-7.20 (2H,m), 7.38-7.50 (5H,m), 7.65 (2H,d), 7.85 (1H,brs), 8.00 (1H,brs), 8.15 (1H,brs).
5379	$\text{C}_{32}\text{H}_{32}\text{N}_4\text{O}_3 \cdot \text{HCl}$			$d_6\text{-DMSO}/400\text{MHz}$	1.60-1.68 (2H,m), 1.80-1.88 (2H,m), 3.00-3.06 (1H,m), 3.15-3.35 (6H,m), 3.65 -3.75 (1H,m), 4.25-4.55 (2H,m), 6.80 (2H,brs), 7.18-7.45 (7H,m), 7.55-7.65 (4H,m), 7.89 (2H,d), 8.57 (1H,brs), 10.29 (2H,brs), 10.36 (1H,brs).
5386	$\text{C}_{35}\text{H}_{39}\text{N}_5\text{O}_4$	594(100), 97(50)	ESI	$d_6\text{-DMSO}/400\text{MHz}$	1.81-1.90 (2H,m), 2.15 (6H,s), 2.35 (2H,t), 2.62-2.70 (2H,m), 2.79-2.83 (2H,m), 3.46-3.53 (2H,m), 4.02 (2H,t), 6.73 (1H,s), 6.75 (1H,s), 6.98 (2H,d), 7.02-7.11 (4H,m), 7.50 (2H,d), 7.60 (2H,d), 7.78 (2H,d), 8.41-8.48 (1H,m), 10.22 (1H,brs)

No.	Mol. Formula	Mass spec. data	¹ H nmr data				
			mass (intensity)	mode	solvent (field)	δ	
5386 . 2HCl	C ₃₅ H ₃₉ N ₅ O ₄ . 2HCl	594(100), 297(58)	ESI	d ₆ -DMSO/400MHz	2.12-2.21 (2H,m); 3.76-3.82 (2H,m); 4.41 (2H,brs); 7.02 (2H,d); (1H,brs),	2.72 (6H,s); 4.12 (2H,t); 6.78 (1H,s); 9.05 (1H,brs); 10.19 (1H,brs).	3.1-3.25
5388	C ₂₂ H ₂₅ N ₃ O ₄ S			d ₆ -DMSO/400MHz	2.16 (6H,s); 3.75 (2H,t); 6.72 (1H,s); 7.32 (1H,m);	2.42 (2H,t); 4.23 (2H,t); 6.78 (1H,s); 7.42 (2H,m);	3.55 (2H,t); 6.43 (1H,d); 7.22 (1H,d); 7.53 (2H,d).
5388 . HCl	C ₂₂ H ₂₅ N ₃ O ₄ S . HCl	428(5)	CI	d ₆ -DMSO/400MHz	2.72 (6H,s); 4.32 (2H,t); 6.81 (1H,s); 7.42 (2H,m);	3.25 (2H,t); 6.47 (1H,d); 7.27 (1H,d); 7.55 (2H,d); (1H,brs).	3.81 (4H,m); 6.76 (1H,s); 7.32 (1H,m); 10.15
5389	C ₂₄ H ₂₉ N ₃ O ₃ S	440(5)	CI	d ₆ -DMSO/400MHz	1.28-1.45 (6H,m); (6H,s); (1H,d); (1H,d); (2H,d).	1.57 (2H,m); 2.20 (2H,t); 6.75 (1H,s); 7.32 (1H,m); (2H,d).	2.12 6.41 7.23 7.55
5389 . HCl	C ₂₄ H ₂₉ N ₃ O ₃ S . HCl	440(5)	CI	d ₆ -DMSO/400MHz	1.36 (2H,m); 1.76 (2H,m); 4.13 (2H,t); 6.80 (1H,s); 7.41 (2H,m); (3H,brs).	1.45 (2H,m); 2.72 (6H,s); 6.42 (1H,d); 7.25 (1H,d); 7.55 (2H,d); 10.06	1.66 (2H,m); 3.0 (2H,t); 6.75 (1H,s); 7.32 (1H,m); 10.06

No.	Mol. Formula	Mass spec. data	1H nmr data			
			mass (intensity)	mode	solvent (field)	δ
5391	C ₃₀ H ₂₈ N ₄ O ₃	493(100), 489(50)	ESI	CDCl ₃ +CF ₃ CO ₂ D/400MHz	3.15-3.25 (1H,m); 3.48-3.57 (1H,m); 3.60-3.68 (2H,m); 3.92-4.02 (3H,m). (1H,d), 7.11 (1H,d), 7.22-7.56 (12H,m), 7.85 (2H,d).	3.15-3.25 (1H,m); 3.48-3.57 (1H,m); 3.60-3.68 (2H,m); 4.33 (2H,d); 4.77 (1H,d), 7.11 (1H,d), 7.22-7.56 (12H,m), 7.85 (2H,d).
5391.HCl	C ₃₀ H ₂₈ N ₄ O ₃ .HCl	493(100)	ESI	d ₆ -DMSO/400MHz	3.01-3.10 (1H,m); 3.80-3.85 (3H,m). 4.61-4.70 (1H,m); (5H,m), 7.41 (2H,t), 7.58 (2H,d), 7.67 (2H,d), 7.99 (2H,d), 9.02 (1H,t), 10.29 (1H,brs). 10.39 (1H,brs). 10.99 (1H,brs).	3.38-3.45 (4H,m); 4.32-4.41 (1H,m); 6.80 (2H,s); 7.18-7.36 (2H,d), 7.99 (2H,d), 9.02 (1H,t), 10.29 (1H,brs). 10.39 (1H,brs). 10.99 (1H,brs).
5393	C ₃₈ H ₃₆ N ₄ O ₅			d ₆ -DMSO/400MHz	2.70 (6H,m); 3.70 (6H,s); 6.80 (1H,s); 7.32 (1H,m); 7.68 (4H,d); 10.35 (2H,br).	2.80 (2H,m); 6.63 (1H,s); 6.83 (1H,s); 7.42 (2H,m); 7.55 (2H,d); 7.68 (4H,d); 7.99 (2H,d), 10.15 (1H,s).
5393.HCl	C ₃₈ H ₃₆ N ₄ O ₅ .HCl	629(100)	CI	d ₆ -DMSO/400MHz	2.95-3.45 (8H,m); 4.50 (2H,m); 6.82 (1H,s); 7.32 (1H,m); 7.68 (2H,d); 10.28 (2H,s). 10.40 (1H,s), 10.80 (1H,brs).	3.75 (2x3H,s); 4.25- 6.79 (1H,s); 6.80 (1H,s); 6.83 (1H,s); 7.30 (2H,d); 7.41 (2H,m); 7.55 (2H,d); 7.68 (2H,d); 7.77 (2H,d); 8.01 (2H,d); 10.28 (2H,s). 10.40 (1H,s), 10.80 (1H,brs).
5394	C ₃₁ H ₃₀ N ₄ O ₃	507(15)	CI	d ₆ -DMSO/400MHz	1.75-1.85 (2H,m); 2.52-2.57 (2H,m), 2.67 (2H,t), 2.84 (2H,t), 3.34-3.40 (2H,m); 3.57 (2H,s); 6.75 (1H,s); 6.80 (1H,s); 7.05-7.10 (4H,m); 7.30-7.55 (7H,m); 7.84 (2H,d), 8.57 (1H,brt), 10.25 (2H,brs).	1.75-1.85 (2H,m); 2.52-2.57 (2H,m), 2.67 (2H,t), 2.84 (2H,t), 3.34-3.40 (2H,m); 3.57 (2H,s); 6.75 (1H,s); 6.80 (1H,s); 7.05-7.10 (4H,m); 7.30-7.55 (7H,m); 7.84 (2H,d), 8.57 (1H,brt), 10.25 (2H,brs).

No.	Mol. Formula	Mass spec. data		¹ H nmr data		
		mass (intensity)	mode	solvent (field)	δ	
5394.HCl	C ₃₁ H ₃₀ N ₄ O ₃ .HCl		d ₆ -DMSO/400MHz	2.02-2.10 (2H,m); 3.18-3.43 (6H,m); 4.23-4.53 (2H,m); (1H,s); 7.20-7.45 (7H,m); 7.65 (2H,d); 10.25 (1H,s); 10.35 (1H,s); 10.60 (1H,brs).	2.95-3.01 (1H,m); 3.65-3.70 (1H,m); 6.79 (1H,s); 6.81 (2H,d); 8.70 (1H,t);	
5397	C ₃₇ H ₄₃ N ₅ O ₄	622(80)	C1	CDCl ₃ /400MHz	1.75-1.83 (4H,m); (6H,s); 2.45 (2H,t); 2.75 (2H,t); 2.85-2.90 (2H,m); (1H,s); 3.62 (2H,s); 6.95-7.20 (10H,m); 7.65 (1H,d); 7.83 (1H,brs); 8.15 (1H,brs).	1.95-2.00 (2H,m); 2.58-2.61 (2H,m); 3.47-3.52 (2H,t); 4.05 (2H,t); 6.90 (10H,m); 7.35 (2H,d);
5397.2HCl	C ₃₇ H ₄₃ N ₅ O ₄ .2HCl		d ₆ -DMSO/400MHz		1.60-1.65 (2H,m); 2.12-2.20 (2H,m); 3.25-3.35 (8H,m); 4.13 (2H,t); 6.75 (1H,s); 6.78 (1H,s); 8.60 (1H,brt); 10.20 (1H,brs); 10.30 (1H,brs).	1.82-1.90 (2H,m); 2.79 (6H,d); 3.65-3.75 (2H,t); 4.25-4.55 (2H,m); 7.00 (2H,d);
5402	C ₃₈ H ₃₅ N ₅ O ₇		d ₆ -DMSO/400MHz			2.70 (6H,m); 3.70 (6H,s); 6.80 (1H,s); 7.68 (4H,d); 8.22 (2H,d); 8.22 (2H,d); 10.15 (1H,s); 10.55 (1H,brs).
						2.80 (2H,m); 6.61 (1H,s); 6.82 (1H,s); 7.82 (2H,d); 7.98 (2H,d);
						3.55 (2H,s); 6.63 (1H,s); 7.22 (2H,d);

No.	Mol. Formula	Mass spec. data			¹ H nmr data		
		mass (intensity)	mode	solvent (field)	δ		
5402.HC1	C ₃₈ H ₃₅ N ₅ O ₇ .HC1	674(80)	ESI	d ₆ -DMSO/400MHz	3.00-3.50 (8H,m), 3.73 (2x3H,s), 4.25 (2H,m), 6.75 (1H,s), 6.79 (1H,s), 6.86 (1H,s), 6.88 (1H,s), 7.29 (2H,d), 7.69 (2H,d), 7.77 (4H,m), 8.00 (2H,d), 8.25 (2H,d), 10.25 (1H,s), 10.55 (1H,brs), 10.70 (1H,brs).		
5376	C ₃₃ H ₃₄ N ₄ O ₄ .HC1	551(100)	ESI	d ₆ -DMSO/400MHz	2.11-2.20 (2H,m), 2.78 (6H,s), 2.83-2.82 (2H,m), 3.20 (2H,m), 3.62 (2H,brs), 4.09 (2H,t), 4.75 (2H,brs), 6.77 (1H,s), 6.79 (1H,s), 7.00 (2H,d), 7.19 (4H,brs), 7.50 (2H,d), 7.55 (2H,d), 7.60 (2H,d), 10.19 (1H,brs), 10.32 (1H,brs), 10.55 (1H,brs).		
5299	C ₂₁ H ₂₄ N ₄ O ₂ S			d ₆ -DMSO/400MHz	2.18 (6H,s), 2.47 (2H,t), 3.01 (3H,s), 3.40 (2H,d), 5.98 (1H,d), 6.71 (1H,s), 6.85 (1H,s), 7.26 (1H,d), 7.31 (1H,m), 7.41 (2H,m), 7.52 (2H,d), 9.85 (1H,brs).		
1912	C ₂₃ H ₂₄ N ₄ O ₃	404(55)	EI	d ₆ -DMSO/400MHz	2.25 (6H,s), 2.93 (2H,s), 4.30 (2H,d), 6.74 (1H,s), 6.76 (1H,s), 7.28-7.55 (9H,m), 8.25 (1H,t), 10.20 (2H,brs).		

No.	Mol. Formula (M. Wt.)	Mass spec m/z, mass intensity (mode)	¹ H nmr			Microanalysis	
			Solvent	δ	all 400 MHz	Calc	Found
1927	C ₁₆ H ₁₄ N ₄ O ₂ 294	291, 30%; 295, MH ⁺ 100% (DCI, NH ₃)	CDCl ₃ + TFA	2.45 (3H, s), (1H, s)	6.85		
				7.38 (1H, s), (5H, m)	7.48		
				8.95 (1H, s).			
1926	C ₁₅ H ₁₂ N ₄ O ₂ 280	281 MH ₊ 100% (DCI, NH ₃)	CDCl ₃ + TFA	7.20 (1H, s), (8H, m).	7.45		
1545	C ₂₁ H ₁₇ N ₃ O ₃ 359	192, 20%; 292, 10%. MH ⁺ 360 (DCI NH ₃)	CDCl ₃ + CF ₃ CO ₂ D	7.82 (1H, d), 7.75 (1H, d), 7.65 (1H, t), 7.48 (3H, m), 7.35 (2H, m), 7.25 (1H, s), 7.06 (2H, d), 3.98 (3H, s).			
1542	C ₁₆ H ₁₀ N ₂ O ₃ Cl ₂ 348	349, 351, 353, 100%; 366, 368, 370, 50%; 313, 39%. (DCI NH ₃)	CDCl ₃ /TFA	6.72 (1H, s), 7.18 (2H, 2x s), 7.34 (1H, t), 7.43 (2H, d), 7.59 (1H, s).			

No.	Mol. Formula (M. Wt.)	Mass spec m/z, mass intensity (mode)	¹ H nmr		Microanalysis	
			Solvent δ all 400 MHz	Calc	Found	
1509	C ₂₀ H ₁₅ N ₃ O ₂ 407	347 MNH ₄ ⁺ , 1%; 330 MH ⁺ , 100% (DCI NH ₃)	CDCl ₃ /TFA 7.22-7.40 (3H, m), 7.40-7.52 (6H, m), 7.60 (1H, s), 7.78 (1H, d, J=7Hz), 7.81 (1H, s), 8.10 (1H, s).			
1507	C ₂₂ H ₂₃ N ₃ O ₅ 407	310, 100%; 336, 20%; 351, 20%; MH ⁺ , 410, 5%; MNH ₄ ⁺ , 427, 2% (DCI NH ₃)	CDCl ₃ + CF ₃ CO ₂ D 7.65 (1H, s), 7.48 (1H, brs), 7.42 (2H, d), 7.22 (1H, s); 7.00 (2H, d), 6.72 (1H, brd), 6.39 (1H, brd), 3.90 (3H, s), 1.65 (9H, s).	C H N	64.54 5.66 10.26	64.45 5.61 10.46
1506	C ₂₆ H ₂₅ N ₃ O ₅ 459	360, 100%; MH ⁺ , 460, MNH ₄ ⁺ , 477, 2% (DCI NH ₃)	CDCl ₃ + CF ₃ CO ₂ D 8.27 (1H, d) 8.05 (1H, s), 7.70 (1H, d), 7.47 (3H, m), 7.38 (2H, pt), 7.25 (1H, s), 7.05 (2H, d), 3.90 (3H, s), 1.65 (9H, s).	C H N	67.96 5.48 9.14	67.54 5.35 9.21

No.	Mol. Formula (M. Wt.)	Mass spec m/z, mass intensity (mode)	¹ H nmr Solvent δ all 400 MHz	Microanalysis		
				Calc	Found	
1476	C ₁₇ H ₁₄ N ₂ O ₄ 310	279, 15%; MH ⁺ , 311; MNH ₄ ⁺ , 328, 2%; (DCI NH ₃)	CDCl ₃ + CF ₃ CO ₂ D 7.85 (1H, s), 7.60 (1H, brs), 7.42 (2H, d), 7.21 (1H, s), 7.08 (1H, s), 7.02 (2H, d), 6.72 (1H, brs), 3.90 (3H, s).	C 65.80 H 4.55 N 9.03	65.87 4.44 9.03	65.68 4.54 8.98
1474	C ₁₇ H ₁₄ N ₂ O ₃ 326	279, 10%; MH ⁺ , 327 (DCI NH ₃)	CDCl ₃ + CF ₃ CO ₂ D 7.60 (1H, d), 7.45 (3H, m), 7.35 (1H, s), 7.23 (2H, m), 7.05 (2H, d), 3.90 (3H, s).	C 62.56 H 4.32 N 8.58	62.41 4.41 8.57	62.39 4.46 8.55
1950	C ₂₅ H ₂₇ N ₃ O ₄ 433	MH ⁺ (100%) 434 CI/NH ₃	CDCl ₃ CF ₃ CO ₂ D 400 MHz 7.50-7.42 (m, 5H), 7.25-7.15 (m, 4H), 7.00 (d, 1H), 6.96 (d, 1H), 6.90 (d, 1H), 4.41 (t, 2H), 3.90 (2, 3H), 3.67 (t, 2H), 3.12 (s, 6H).	C 69.57 H 6.28 N 9.69	68.98 6.25 9.59	69.06 6.25 9.60

No.	Mol. Formula (M. Wt.)	Mass spec m/z, mass intensity (mode)	¹ H nmr		Microanalysis	
			Solvent δ all 400 MHz	Calc	Found	
1718	C ₂₁ H ₁₇ N ₃ O ₃ 359	MH ⁺ 360, 100% (DCI NH ₃)	DMSO 11.4 (1H, s), 10.08 (1H, s), 9.82 (1H, s), 7.55 (3H, m), 7.39 (1H, d), 7.18 (1H, t), 7.01 (4H, m), 6.85 (1H, s), 6.78 (1H, s), 3.80 (3H, s).			
1693	C ₂₂ H ₁₉ N ₃ O ₅ S 437	360, 85%; 402, 25%; MH ⁺ 438 (DCI NH ₃)	7.98 (1H, d), 7.88 (1H, s), 7.75 (1H, d), 7.45 (5H, m), 7.35 (1H, s), 7.02 (2H, d), 3.90 (3H, s), 3.30 (2.33H, s).			
1618	C ₂₃ H ₂₁ N ₃ O ₄ S 435	436, 100%; 336, 82%	CDCl ₃ TFA 1.75 (9H, s), 7.22-7.28 (overlapping solvent & sample signals), 7.36-7.50 (6H, overlapping signals), 7.61 (2H, overlapping signals), 8.10 (1H, s).			

No.	Mol. Formula (M. Wt.)	Mass spec <i>m/z</i> , mass intensity (mode)	¹ H nmr		Microanalysis	
			Solvent δ at 400 MHz	Calc	Found	Calc
1560	C ₂₅ H ₂₁ N ₃ O ₄ C1 ₂ 497	498/500/502 (100/69/15)% 398/400/402 (49/31/7)%	DMSO-D ₆ 1.68 (9H, s), 6.66 (1H, s), 6.92 (1H, s), 7.30-7.44 (3H, c), 7.49 (2H, d), 7.68 (1H, d), 8.08 (1H, d), 8.17 (1H, s).			
1470	C ₂₁ H ₂₁ N ₃ O ₄	397, MNH ₄ ⁺ , 4%; 380, MH ⁺ , 13%, 280, 100% (DCI NH ₃)	CDCl ₃ 1.64 ³ (9H, s), 6.33 (1H, br. s), 6.57 (1H, br. s), 7.00 (1H, s), 7.35-7.50 (7H, m), 8.10 (1H, br. s), 8.18 (1H, br. s)			
1471	C ₂₅ H ₂₃ N ₃ O ₄	447, MNH ₄ ⁺ , 17%; 430, MH ⁺ , 100%; 330, 82%	CDCl ₃ 1.72 ³ (9H, s), 7.07 (1H, s), 7.14 (1H, s), 7.30-7.50 (7H, m), 7.66 (1H, d, J=7Hz), 7.84 (1H, s), 8.03 (1H, br. s), 8.18 (2H, m)			

N. 66998A

No.	Mol. Formula (M. Wt.)	Mass spec <i>m/z</i> , mass intensity (mode)	¹ H nmr		Microanalysis	
			Solvent δ all 400 MHz	Calc	Found	
1729	C ₂₃ H ₁₉ N ₃ O ₅	435, MNH ₄ ⁺ , 23%; 418, MH ⁺ , 100% (DCI NH ₃)	CDCl ₃ 3.09 (4H, s), 3.92 (3H, s), 7.07 (2H, d, J=7Hz), 7.28 (1H, s), 7.30 (1H, s), 7.39 (2H, d, J=6Hz), 7.45 (2H, d, J=7Hz), 7.60 (2H, d, J=6Hz)			
1647	C ₂₃ H ₂₁ N ₃ O ₃ 387	405, M ⁺ NH ₄ , 7%; 388, M ⁺ H, 100%; 317, 43%; 459, 29% (DCI NH ₃)	CDCl ₃ 1.84-2.00 (4H, m), 3.13 (2H, t), 3.64 (2H, t), 6.98 (1H, s), 7.03 (1H, s), 7.32-7.50 (9H, m), 8.10 (1H, brs), 8.32 (1H, brs)			
1845	C ₂₁ H ₁₇ N ₃ O ₅ 391	409, M ⁺ NH ₄ , 35%; 392, MH ⁺ , 100% (DCI NH ₃)	CDCl ₃ + TFA 2.35 (3H, s, AC), 6.05 (2H, s, OCH ₂ O), 6.85-7.60 (9H, m)	C 64.45	63.99	63.94
1809	C ₂₀ H ₁₆ N ₂ O ₅ 364	382, M ⁺ +NH ₄ , 5%; 365, MH ⁺ , 100% (DCI NH ₃)	CDCl ₃ + TFA 3.85 (3H, s, OMe), 6.05 (2H, s, OCH ₂ O), 6.90-7.45 (9H, m)	C 65.93	65.85	65.96
				H N	4.43 7.69	4.38 7.60
					4.37 7.65	4.37 7.65

N. 66998A

No.	Mol. Formula (M. Wt.)	Mass spec	¹ H nmr				Microanalysis	
			m/z, mass intensity (mode)	Solvent δ all 400 MHz	Calc	Found		
1808	C ₁₉ H ₁₄ N ₂ O ₄ 334	M ⁺ +1, 100%	CDCl ₃ + TFA 6.05 (2H, s, OCH ₂ O), 6.90-7.50 (10H, m)	C H N	68.26 4.22 8.38	68.07 4.15 8.35	68.00 4.17 8.35	
1929	C ₂₂ H ₁₈ N ₄ O ₂ 370	MH ⁺ , 371 (DCI NH ₃)	CDCl ₃ + TFA 5.45 (2H, s); 7.18 (1H, s); 7.26 (1H, s); 7.30 (1H, s); 7.45 (10H, m); 8.88 (1H, s)					
1930		MH ⁺ , 357, 100% (DCI NH ₃)	CDCl ₃ + TFA 7.27 (1H, s); 7.30 (1H, s); 7.50 (5H, m); 7.65 (5H, m); 7.75 (1H, t); 9.10 (1H, s).					

No.	Mol. Formula (M. Wt.)	Mass spec <i>m/z</i> , mass intensity (mode)	¹ H nmr		Microanalysis	
			Solvent δ all 400 MHz	Calc	Found	Calc
1975	C ₂₇ H ₂₉ N ₅ O ₃	236, 25%; 257, 100%; 376, 20%; NH ⁺ , 472, 20%.	CDCl ₃ + TFA 2.35 (2H, m), 3.01 (6H, s), 3.45 (2H, t), 4.18 (2H, t), 5.40 (2H, s), 6.95 (2H, d), 7.20 (1H, m), 7.25 (1H, s), 7.40 (3H, m), 7.50 (3H, m).			
1976	C ₂₆ H ₂₇ N ₅ O ₃ 457	230, 100%; 247, 60%; NH ⁺ , 458, 90%.	CDCl ₃ + TFA 2.30 (2H, m), 2.05 (6H, s), 3.45 (2H, t), 4.18 (2H, t), 6.98 (2H, d), 7.25 (2H, d), 7.45 (2H, d), 7.55 (3H, m), 7.75 (3H, m), 9.18 (1H, s).			

No.	Mol. Formula (M. Wt.)	Mass spec m/z, mass intensity (mode)	¹ H nmr		Microanalysis	
			Solvent δ all 400 MHz	Calc.	Found	Calc.
1982	C ₂₄ H ₂₈ N ₄ O ₃ .2HCl 404+73	405, 100%, MH ⁺ EI ⁺	D ₂ O 2.98 (3H, s), 3.09 (6H, s), 3.75 (4H, brs), 4.50 (2H, s), 7.09 (1H, s), 7.13 (1H, s), 7.52-7.68 (5H, c), 7.67-7.77 (4H, overlapping signals).			
1983	C ₂₆ H ₃₀ N ₄ O ₂	431, 25%; MH ⁺ 332, 30%; 303, 18%; 84, 92%; 118, 100%. EI ⁺	DMSO-D ₆ 1.53 (2H, m), 1.71 (2H, d), 1.83 (2H, t), 2.12 (3H, s), 2.14 (3H, s), 2.35 (1H, m), 2.80 (2H, d), 3.57 (2H, s), 6.78 (2H, overlapping signals), 7.34 (3H, overlapping signals), 7.43 (2H, t), 7.50 (2H, d), 7.57 (2H, d).			

No.	Mol. Formula (M. Wt.)	Mass spec m/z, mass intensity (mode)	¹ H nmr		Microanalysis	
			Solvent of all 400 MHz	Calc.	Found	
1886	C ₂₉ H ₂₁ N ₃ O ₇		CDCl ₃ / TFA 3.90 (3H, s), 4.79 (2H, s), 7.01 (2H, d, J=8Hz), 7.21 (1H, s), 7.24 (1H, s), 7.27 (2H, d), J=8Hz, 7.41 (2H, d), J=8Hz, 7.47 (2H, d), J=8Hz, 7.82 (2H, m), 7.97 (2H, m).			
1657	C ₂₀ H ₁₉ N ₃ O ₃ 349	M ⁺ , 350, 12%; M ⁺ , 349, 13%; 333, 100%. CI NH ₃	CDCl ₃ / TFA 3.92 (3H, s), 4.32 (2H, s), 7.05 (2H, d), 7.24 (2H, d), 7.45 (2H, d), 7.52 (4H, s).			
1891	C ₂₃ H ₂₄ N ₂ O ₄ 392	392, M ⁺ , 25%; 347, M ⁺ - OCH ₂ CH ₃ , 100% EI	DMSO 1.15 (6H, t, J=6Hz, CH ₃), 3.45-3.60 (4H, m, CH ₂ CH ₃) 5.50 (1H, s, O ₂ CH), 6.75 (2H, s), 7.28-7.55 (9H, m, Ar), 10.25 (2H, br.s, NH)	C 70.39 H 6.16 N 7.14	70.31 6.16 7.03	70.03 6.16 7.09

N. 66998A

No.	Mol. Formula (M. Wt.)	Mass Spec <i>m/z</i> , mass intensity (mode)	¹ H nmr		Microanalysis	
			Solvent δ all 400 MHz	Calc	Found	Calc
1912	C ₂₃ H ₂₄ N ₄ O ₃ 404	404, M ⁺ . 55%; 303, M ⁺ -. NHC(O)CH ₂ NMe ₂ , 30%; EI	DMSO 2.25 (6H, s; 2xMe). 2.95 (2H, s). 4.30 (2H, d, J=6Hz). 6.74 (1H, s). 6.76 (1H, s). 7.28-7.55 (9H, m). 8.24-8.27 (1H, br. m, NH). 10.20 (2H, br. s, 2xNH).			
1676	C ₂₂ H ₁₉ O ₃ N ₃ 373	MH ⁺ . 100%, 374 (DCI/NH ₃)	CDCl ₃ , CF ₃ CO ₂ D 7.65 (2H, d). 7.58 (2H, d) 7.48 (2H, d) 7.41-7.35 (4H, m). 7.24 (1H, s) 7.12-7.07 (2H, m). 2.36+2.23 (3H, s, rotamers).			

No.	Mol. Formula (M. Wt.)	Mass spec m/z, mass intensity (mode)	¹ H nmr			Microanalysis	
			Solvent	δ	all 400 MHz	Calc	Found
1959	C ₂₅ H ₂₈ N ₃ O ₄ C1 469/471	CI/NH ₃	d ₆ -DMSO	400 MHz			
			10.85 (1H, s), 10.10 (1H, brs), 10.02 (1H, s), 7.6-7.30 (7H, m), 7.10 (2H, m), 6.85 (1H, d), 6.80 (1H, s), 6.58 (1H, d), 4.36 (2H, t), 3.87 (3H, s), 3.50 (2H, t), 2.88 (6H, s).				
1921	C ₂₂ H ₂₁ N ₃ O ₂ 359	MH ⁺ . 100%. 360 CI/NH ₃	CDCl ₃ + CF ₃ CO ₂ D	C	73.52	73.24	73.11
			7.81 (2H, d), 7.52 (2H, d), 7.40-7.50 (6H, m), 7.24 (1H, s), 6.98 (1H, d), 6.96 (1H, d), 3.33 (6H, s).	H	5.89	5.82	5.77
				N	11.69	11.50	11.52
1922	C ₂₆ H ₂₀ N ₂ O ₂ 392	MH ⁺ . 393. 100%: MNH ⁺ . 410. CI/NH ₃	d ₆ -DMSO				
			11.15 (1H, brs), 10.00 (1H, brs), 7.66 (1H, d), 7.51-7.30 (13H, m), 7.20 (2H, m), 6.78 (1H, s), 6.83 (1H, d).				


No.	Mol. Formula (M. Wt.)	Mass spec m/z, mass intensity (mode)	^1H nmr Solvent δ at 400 MHz	Microanalysis		
				Calc	Found	
1923	$\text{C}_{20}\text{H}_{15}\text{N}_3\text{O}_4$ 361	MH^+ , 362, 100% (DCI NH_3)	$\text{CDCl}_3, \text{CF}_3\text{CO}_2\text{D}$ 8.25 (2H, d). 7.83 (2H, d). 7.63 (1H, dd). 7.55-7.45 (5H, m). 7.35 (1H, s). 7.12 (1H, d). 7.08 (1H, d).	C 66.48 H 4.18 N 11.63	66.61 4.23 11.40	66.54 4.26 11.48
1672	$\text{C}_{20}\text{H}_{23}\text{N}_3\text{O}_3$ 353	MH^+ , 354, 100%; MNH^+ , 371, 10%; 271, 10%; 260, 10% (DCI NH_3)	$\text{CDCl}_3, \text{CF}_3\text{CO}_2\text{D}$ 7.59 (2H, d). 7.45 (2H, d). 7.18 (1H, s). 6.29 (1H, d). 2.55-2.47 (1H, m). 2.36-2.22 (3H, s, rotamers). 1.82-1.70 (5H, s). 1.51-1.40 (2H, m). 1.32-1.20 (3H, m).			
1884	$\text{C}_{18}\text{H}_{20}\text{N}_2\text{O}_2$ 296	MH^+ , 297, 100%; MNH^+ , 315, 10% (DCI NH_3)	$\text{CDCl}_3, \text{CF}_3\text{CO}_2\text{D}$ 7.48-7.38 (5H, m). 7.21 (1H, s). 6.26 (1H, d). 2.48 (1H, m). 1.83-1.70 (1H, m). 1.35 (2H, m). 1.30-1.19 (3H, m).			

No.	Mol. Formula (M. Wt.)	Mass spec		^1H nmr		Microanalysis	
		m/z, mass	intensity (mode)	Solvent δ all 400 MHz		Calc	Found
1570	$\text{C}_{17}\text{H}_{14}\text{N}_2\text{O}_2\text{S}$ 310	311, M ⁺ , 100% DCI-NH ₃		CDCl ₃ 4.13 (3H, s), 6.59 (1H, s), 7.10 (1H, m), 7.30-7.60 (8H, m), 8.09 (1H, brs).	C H N	65.79 4.55 9.03	65.24 4.53 8.73

CLAIMS

1. A piperazine of general formula (A) :

5



wherein one or both of R₁ and R₂, which may be the same or
10 different, is:

(I) X, or a phenyl group which is substituted by X,

C(O)X, OC(O)CH₂X, OCH₂CH₂X, CH₂X, CONH(CH₂)_nX,

O(CH₂)_nCH(OH)(CH₂)_nX or $-\text{C}(\text{O})\text{NH}\left(-\text{C}_6\text{H}_4\right)_n\text{CH}_2\text{X}$

15 or which is fused to a group X;

(II) a phenyl group substituted by CH₂NR₁₂R₁₃,

OC(O)(CH₂)_nZ, CH(OR₁₂)(OR₁₃), (CH₂)_nNR₁₄C(O)(CH₂)_mNR₁₂R₁₃,

-CH₂NR₁₂-(CH₂)_nNR₁₅R₁₆, O(CH₂)_nCH(OH)(CH₂)_nN(R₁₂R₁₃);

(III) a group CH=C(W)V; or

20 (IV) a cyclohexyl group;

and where appropriate, the other of R₁ and R₂ is a phenyl group optionally substituted by one or more groups independently selected from halogen, nitro, methoxy,

NHC(O)R₁₂, CO₂H, O(CH₂)_nN(R₁₂R₁₃), CH₂Y(CH₂)_nN(R₁₂R₁₃),

25 C₁-C₄ alkyl and (CH₂)_nC(O)OR₁₂;

X is a naphthyl group or a five- or six-membered saturated or unsaturated heterocyclic group containing one or more heteroatoms, which heteroatoms may be the same or different

and are independently selected from O, N and S; the heteroatom(s) when nitrogen being optionally substituted by hydrogen, methyl, oxygen, tertiary-butyloxycarbonyl, -(CH₂)_nCH₂OH or SO₂Me; the heterocyclic ring being

5 optionally substituted by halogen, Me, MeS, phenyl, O(CH₂)_nNR₁₂R₁₃, -N(R₁₂)(CH₂)_nN(R₁₂R₁₃), -(CH₂)_nN(R₁₂R₁₃) or -O(CH₂)_nO(CH₂)_nN(R₁₂R₁₃), or the heterocyclic ring optionally containing one or more carbonyl groups and being optionally fused to a benzene ring, which benzene ring is optionally

10 substituted by 1 or 2 C₁-C₆ alkoxy groups;

Y is O or S;

Z is a C₃-C₆ cycloalkyl group;

R₁₂, R₁₃ and R₁₄, which may be the same or different, are hydrogen or C₁-C₆ alkyl;

15 R₁₅ and R₁₆, which may be the same or different, are hydrogen or C₁-C₆ alkyl, or R₁₅ and R₁₆ form, together with the atom to which they are attached, a 5- or 6-membered heterocyclic group;

W is hydrogen or a phenyl group;

20 V is a phenyl group optionally substituted by one or more groups independently selected from nitro, alkoxy and O(CH₂)_nNR₁₂R₁₃;

m and n are each, independently, 0 or an integer having the value 1, 2, 3 or 4;

25 O(CH₂)_nNR₁₂R₁₃ or containing one or more carbonyl groups and being optionally fused to a benzene ring;

Z is a C₃-C₆ cycloalkyl group;

R₁₂, R₁₃ and R₁₄, which may be the same or different, are

hydrogen or C₁-C₄ alkyl;

W is hydrogen or a phenyl group;

V is a phenyl group optionally substituted by one or more groups independently selected from nitro, alkoxy and

5 O(CH₂)_nNR₁₂R₁₃; and

m and n are, independently, integers having the values 1, 2, 3 or 4;

or a pharmaceutically acceptable salt or ester thereof.

2. A compound according to claim 1, wherein one or both of R₁ and R₂, which may be the same or different, is chosen from X and a phenyl group substituted by X, C(O)X, OC(O)CH₂X, OCH₂CH₂X, CH₂X or which is fused to a group X; X is a five- or six-membered heterocyclic ring containing one or two heteroatoms, which may be the same or different, independently selected from O, N and S, the heteroatoms(s) when nitrogen being optionally substituted by hydrogen, methyl, oxygen, tertiary-butyloxycarbonyl, or SO₂Me, the heterocyclic ring being optionally substituted by hydrogen, methyl, phenyl, O(CH₂)_nN(R₁₂R₁₃) or optically containing one or more carbonyl groups and being optionally fused to a benzene ring; Y, R₁₂, R₁₃ and n are as defined in claim 1.

3. A compound according to claim 1 or 2, wherein R₁₂ and R₁₃, which may be the same or different, are hydrogen or C₁-C₃ alkyl and n is an integer of value 1 or 2.

4. A compound according to claim 1, 2, or 3 wherein one of R₁ and R₂ is a phenyl group which is substituted by X, C(X), OCO(O)CH₂X, OCH₂CH₂X, CH₂X or which

is fused to a group X; wherein X is a five- or six-membered heterocyclic ring containing one or two heteroatoms which may be the same or different, independently selected from O, N and S, the heteroatoms(s) when nitrogen being
5 optionally substituted by methyl, the heterocyclic ring being optionally fused to a benzene ring.

5. A compound according to claim 1, wherein one of R₁ and R₂ is a phenyl group substituted by CH₂NR₁₂R₁₃, OC(O)(CH₂)_nZ, CH(OR₁₂)(OR₁₃), (CH₂)_nNR₁₄C(O)(CH₂)_mNR₁₂R₁₃;
10 wherein R₁₂, R₁₃ and R₁₄, which may be the same or different, are independently selected from hydrogen or C₁-C₃ alkyl; Z is a C₅ or C₆ cycloalkyl group; and m and n are, independently, integers having the values 1, 2 or 3.

15 6. A compound according to claim 1 or 5, wherein R₁₂, R₁₃ and R₁₄, which may be the same or different, are independently selected from hydrogen and C₁-C₂ alkyl; Z is a cyclopentyl group; and m and n are, independently, integers having the values of 1
20 or 2.

7. A compound selected from
1926 (3Z,6Z)-3-Benzylidene-6-(4-imidazolyl)methylene-2,5-piperazinedione.
1930 (3Z,6Z)-3-Benzylidene-6-(4-(1-imidazolyl)benzylidene)-
25 2,5-piperazinedione.

1929 (3Z,6Z)-3-Benzylidene-6-(4-(1-imidazolylmethyl)benzylidene)-2,5-piperazinedione.

1959 (3Z,6Z)-3-Benzylidene-6-(4-(2-dimethylaminoethoxy)-3-

methoxybenzylidene)-2,5-piperazinedione hydrochloride.

1927 (3Z,6Z)-3-Benzylidene-6-(4-(5-

methylimidazolyl)methylene-2,5-piperazinedione.

1921 (3Z,6Z)-3-Benzylidene-6-(4-

5 dimethylaminocinnamylidene)-2,5-piperazinedione.

1976 (3Z,6Z)-3-(4-(3-Dimethylaminopropoxy)benzylidene)-6-

(4-(1-imidazolyl)benzylidene-2,5-piperazinedione.

1910 (3Z,6Z)-3-Benzylidene-6-(4-(2-

imidazolylethoxy)benzylidene)-2,5-piperazinedione.

10 1923 (3Z,6Z)-3-Benzylidene-6-(4-nitrocinnamylidene-2,5-

piperazinedione.

1657 (3Z,6Z)-3-(4-Aminomethylbenzylidene)-6-(4-

methoxybenzylidene)-2,5-piperazinedione.

1491 Methyl (3Z,6Z)-3-benzylidene-6-(4-methoxybenzylidene)-

15 2-oxo-1,2,3,6-tetrahydro-5-pyrazonyloxyacetate.

1693 (3Z,6Z)-3-(1-methanesulfonyl-3-indolyl)methylene-6-(4-

methoxybenzylidene)-2,5-piperazinedione.

1886 (3Z,6Z)-3-(4-Methoxybenzylidene)-6-(4-

phthalimidoacetoxybenzylidene)-2,5-piperazinedione.

20 1922 (3Z,6Z)-3-Benzylidene-6-(γ -phenylcinnamylidene)-2,5-

piperazinedione.

1618 (3Z,6Z)-3-(1-tert-butoxycarbonyl-3-indolyl)methylene-

6-(2-thenylidene)-2,5-piperazinedione.

1560 (3Z,6Z)-3-(2,6-Dichlorobenzylidene)-6-(1-tert-

25 butoxycarbonyl-3-indolyl)methylene-2,5-piperazinedione.

1950 (3Z,6Z)-3-Benzylidene-6-(4-(2-dimethylaminoethoxy)-3-

methoxycinnamylidene)-2,5-piperazinedione.

1975 (3Z,6Z)-3-(4-(3-Dimethylaminopropoxy)benzylidene)-6-

(4-(1-imidazolylmethyl)benzylidene)-2,5-piperazinedione.

1983 (3Z,6Z)-3-Benzylidene-6-(4-N-methyl-N-(4-(N-methylpiperidinyl)aminomethylbenzylidene)-2,5-piperazinedione.

5 1509 ((3Z,6Z)-3-Benzylidene-6-(3-indolylmethylen)-2,5-piperazinedione.

1542 (3Z,6Z)-3-(2,6-Dichlorobenzylidene)-6-(3-furylmethylene)-2,5-piperazinedione.

1545 (3Z,6Z)-3-(3-Indoxylmethylen)-6-(4-

10 methoxybenzylidene)-2,5-piperazinedione.

1560 (3Z,6Z)-3-(2,6-Dichlorobenzylidene)-6-(3-(1-tertbutoxycarbonyl)indolyl)methylene-2,5-piperazinedione.

1507 (3Z,6Z)-3-(4-Methoxybenzylidene)-6-(2-(1-tertbutoxycarbonyl)pyrrolyl)methylene-2,5-piperazinedione.

15 1506 (3Z,6Z)-3-(4-Methoxybenzylidene)-6-(3-(1-tert-butoxycarbonyl)indolyl)methylene-2,5-piperazinedione.

1471 (3Z,6Z)-3-Benzylidene-6-(3-(1-tert-butoxycarbonyl)indolyl)methylene-2,5-piperazinedione.

1474 (3Z,6Z)-3-(4-Methoxybenzylidene)-6-(2-

20 thienylmethylene)-2,5-piperazinedione.

1476 (3Z,6Z)-3-(4-Methoxybenzylidene)-6-(3-furylmethylene)-2,5-piperazinedione.

1672 (3Z,6Z)-3-(Acetamidobenzylidene)-6-cyclohexylmethylen-2,5-piperazinedione.

25 1676 (3Z,6Z)-3-(4-Acetamidobenzylidene)-6-cinnamylidene-2,5-piperazinedione.

1891 (3Z,6Z)-3-Benzylidene-6-(diethoxymethylbenzylidene)-2,5-piperazinedione.

1982 (3Z,6Z)-3-Benzylidene-6-(4-(N-methyl-N-(2-dimethylaminoethyl)aminomethylbenzylidene)-2,5-piperazinedione hydrochloride.

1884 (3Z,6Z)-3-Benzylidene-6-cyclohexylmethylen-2,5-5 piperazinedione.

1845 (3Z,6Z)-3-(4-Acetamidobenzylidene)-6-(3,4-methylenedioxybenzylidene)-2,5-piperazinedione.

1718 (3Z,6Z)-3-(2-Indolylmethylene)-6-(4-methoxybenzylidene)-2,5-piperazinedione.

10 1808 (3Z,6Z)-3-Benzylidene-6-(3,4-methylenedioxybenzylidene)-2,5-piperazinedione.

1809 (3Z,6Z)-3-(4-Methoxybenzylidene)-6-(3,4-methylenedioxybenzylidene)-2,5-piperazinedione.

1470 (3Z,6Z)-3-Benzylidene-6-(2-(1-tertbutoxycarbonyl)pyrrolyl)methylene-2,5-piperazinedione.

15 5023 (3Z,6Z)-3-(4-Dimethylaminomethylbenzylidene)-6-(4-(3-dimethylaminopropoxy)benzylidene)-2,5-piperazinedione.

5026 (3Z,6Z)-3-(4-(3-Dimethylaminopropoxy)benzylidene)-6-(4-(1-imidazolyl)methylbenzylidene)-2,5-piperazinedione.

20 5030 (3Z,6Z)-3-(4-(3-Dimethylaminopropoxy)benzylidene)-6-(4-(1-imidazolyl)benzylidene).

5367 (2-(4-((3Z,6Z)-6-(4-(3-Dimethylaminopropoxy)benzylidene)-2,5-dioxo-3-piperazinylidene)methylbenzoyl)-1,2,3,4-

25 tetrahydroisoquinoline.

5386 N-(2-(1,2,3,4-Tetrahydro-2-isoquinolyl)ethyl)-4-((3Z,6Z)-6-(4-(3-dimethylaminopropoxy)benzylidene)-2,5-dioxo-3-piperazinylidene)methylbenzamide.

- 5397 N- (4- (1,2,3,4-Tetrahydro-2-isoquinolyl)butyl)-4-
((3Z,6Z)-6- (4- (3-dimethylaminopropoxy)benzylidene)-2,5-
dioxo-3-piperazinylidene)methylbenzamide.
- 5027 (3Z,6Z)-6- (4- (3-Dimethylaminopropoxy)benzylidene-3-
5 (4-pyridylmethylene)-2,5-piperazinedione.
- 5028 (3Z,6Z)-6- (4- (3-Dimethylaminopropoxy)benzylidene)-3-
(3-pyridylmethylene)-2,5-piperazinedione.
- 5041 (3Z,6Z)-6- (4- (3-Dimethylaminopropoxy)benzylidene)-3-
furfurylidene-2,5-piperazinedione.
- 10 5042 (3Z,6Z)-6- (4- (3-Dimethylaminopropoxy)benzylidene)-3-
(3-Thenylidene)-2,5-piperazinedione.
- 5046 (3Z,6Z)-6- (4- (3-Dimethylaminopropoxy)benzylidene)-3-
(2-Thenylidene)-2,5-piperazinedione.
- 5052 (3Z,6Z)-6- (4- (3-Dimethylaminopropoxy)benzylidene)-3-
15 (3-Furylmethylene)-2,5-piperazinedione.
- 5188 (3Z,6Z)-6- (4- (3-Dimethylaminopropoxy)benzylidene)-3-
(2-Naphthylmethylene)-2,5-piperazinedione.
- 5200 (3Z,6Z)-6- (4- (3-Dimethylaminopropoxy)benzylidene)-3-
(1-Naphthylmethylene)-2,5-piperazinedione.
- 20 5032 (3Z,6Z)-6-Benzylidene-3- (4- (3-dimethylamino-2-
hydroxypropoxy)benzylidene)-2,5-piperazinedione.
- 5040 (3Z,6Z)-6-Benzylidene-3- (4- (2-hydroxy-3-
morpholinopropoxy)benzylidene)-2,5-piperazinedione.
- 5057 (3Z,6Z)-6-Benzylidene-3- (4- (2-hydroxy-3- (1-
25 imidazolyl)propoxy)benzylidene)-2,5-piperazinedione.
- 5043 (3Z,6Z)-6-Benzylidene-3- (4- (2-hydroxy-3- (4- (2-
hydroxyethyl)-1-piperazinyl)propoxy)benzylidene)-2,5-
piperazinedione.

- 5062 (3Z,6Z)-6-(4-(2-Dimethylaminoethoxy)benzylidene)-3-(3-Furymethylene)-2,5-piperazinedione.
- 5071 (3Z,6Z)-6-(4-(2-Dimethylaminoethoxy)benzylidene)-3-(3-thenylidene)-2,5-piperazinedione.
- 5 5072 (3Z,6Z)-6-(4-(2-Dimethylaminoethoxy)benzylidene)-3-(5-methylthio-2-thenylidene)-2,5-piperazinedione.
- 5054 (3Z,6Z)-6-Benzylidene-3-(4-(2-morpholinoethoxy)benzylidene)-2,5-piperazinedione.
- 5055 (3Z,6Z)-6-Benzylidene-3-(4-(2-(1-imidazolyl)ethoxy)benzylidene)-2,5-piperazinedione.
- 10 5053 (3Z,6Z)-6-Benzylidene-3-(4-(2-(1-pyrrolidinyl)ethoxy)benzylidene)-2,5-piperazinedione.
- 5069 (3Z,6Z)-6-(4-(2-Dimethylaminoethoxymethyl)benzylidene)-3-(3-thenylidene)-2,5-piperazinedione.
- 15 5077 (3Z,6Z)-6-(4-(2-Dimethylaminoethoxymethyl)benzylidene)-3-(3-furymethylene)-2,5-piperazinedione.
- 5074 (3Z,6Z)-6-(4-Dimethylaminoacetamidomethylbenzylidene)-3-(3-thenylidene)-2,5-piperazinedione.
- 20 5079 (3Z,6Z)-3-(2-Bromobenzylidene)-6-(4-dimethylaminoacetamidomethylbenzylidene)-2,5-piperazinedione.
- 5081 (3Z,6Z)-6-(4-Dimethylaminoacetamidomethylbenzylidene)-3-(3-furymethylene)-2,5-piperazinedione.
- 25 5061 (3Z,6Z)-6-Benzylidene-3-(4-dimethylaminoacetamidomethylbenzylidene)-2,5-

piperazinedione.

5073 (3Z,6Z)-6-(4-(2-Dimethylaminoethylthiomethyl)benzylidene)-3-(3-furylmethylene)-2,5-piperazinedione.

5 5078 (3Z,6Z)-6-(4-(2-Dimethylaminoethylthiomethyl)benzylidene)-3-(3-thenylidene)-2,5-piperazinedione.

1912 (3Z,6Z)-6-Benzylidene-3-(4-dimethylaminoacetamidoaminomethylbenzylidene)-2,5-

10 piperazinedione.

5324 (3Z,6Z)-6-Benzylidene-3-(5-(2-dimethylaminoethoxy)-2-thienylmethylene)-2,5-piperazinedione.

5327 (3Z,6Z)-6-Benzylidene-3-(4-(2-dimethylaminoethoxy)-2-thienylmethylene)-2,5-piperazinedione.

15 5335 (3Z,6Z)-6-Benzylidene-3-(5-(2-dimethylaminoethyl)-2-thienylmethylene)-2,5-piperazinedione.

5388 (3Z,6Z)-6-Benzylidene-3-(5-(2-(2-dimethylaminoethoxy)ethoxy)-2-thienylmethylene)-2,5-piperazinedione.

20 5389 (3Z,6Z)-6-Benzylidene-3-(5-(6-dimethylaminohexyloxy)-2-thienylmethylene)-2,5-piperazinedione.

5299 (3Z,6Z)-6-Benzylidene-3-(5-(2-dimethylaminoethyl)methylamino-2-thienylmethylene)-2,5-piperazinedione.

25 5075 (3Z,6Z)-3-(2,5-Dichloro-3-thenylidene)-6-benzylidene-2,5-piperazinedione.

5371 N-(4-(1,2,3,4-Tetrahydro-2-isouinolyl)butyl)-4-((3Z,6Z)-6-benzylidene-2,5-dioxo-3-

piperazinylidene)methylbenzamide.

5391 N- (2- (1,2,3,4-Tetrahydro-2-isoquinolyl)ethyl)-4-
((3Z,6Z)-6-benzylidene-2,5-dioxo-3-
piperazinylidene)methylbenzamide.

5 5394 N- (3- (1,2,3,4-Tetrahydro-2-isoquinolyl)propoyl)-4-
((3Z,6Z)-6-benzylidene-2,5-dioxo-3-
piperazinylidene)methylbenzamide.


5393 N- (4- (2- (1,2,3,4-Tetrahydro-2-
isoquinolyl)ethyl)phenyl)-4- ((3Z,6Z)-6-benzylidene-2,5-
10 dioxo-3-piperazinylidene)methylbenzamide.

5402 N- (4- (2- (1,2,3,4-Tetrahydro-2-
isoquinolyl)ethyl)phenyl)-4- ((3Z,6Z)-2,5-dioxo-6- (4-
nitrobenzylidene)-3-piperazinylidene)methylbenzamide.

8. A pharmaceutical or veterinary composition
15 comprising a pharmaceutically or veterinarily acceptable
carrier or diluent and, as an active principle, a compound
as defined in claim 1.

9. A process for preparing a compound of formula
(A) as defined in claim 1, the process comprising:

20 (a) condensing a compound of formula (I):



25


wherein R₂ are as defined in claim 1 and is optionally
protected, with a compound of formula (II):



wherein R₁ is as defined in claim 1 and is optionally protected, in the presence of a base in an organic solvent; or

(b) condensing a compound of formula (I'):

5




wherein R₁ is as defined in claim 1 and are optionally 10 protected with a compound of formula (III):



wherein R₂ is as defined in claim 1 and is optionally 15 protected, in the presence of a base in an organic solvent; and

(c) if required, removing optionally present protecting groups, and/or, if desired, converting one compound of formula A into another compound of formula A, 20 and/or, if desired, converting a compound of formula A into a pharmaceutically acceptable salt or ester thereof, and/or, if desired, converting a salt or ester into a free compound, and/or, if desired, separating a mixture of isomers into the single isomers.

25 10. Use of a diketopiperazine of formula (A):



wherein one or both of R₁ and R₂, which may be the same or different, is:

- (I) X, or a phenyl group which is substituted by X,
C(O)X, OC(O)CH₂X, OCH₂CH₂X, CH₂X, CONH(CH₂)_nX,
 - 5 O(CH₂)_nCH(OH)(CH₂)_nX or $\text{--C(O)NH} \left(\text{--} \begin{array}{c} \text{C}_6\text{H}_4 \\ | \\ \text{C}_6\text{H}_4 \end{array} \right)_n \text{--} (\text{CH}_2)_m \text{X}$
 - or which is fused to a group X;
 - (II) a phenyl group substituted by CH₂NR₁₂R₁₃,
 - OC(O)(CH₂)_nZ, CH(OR₁₂)(OR₁₃), (CH₂)_nNR₁₄C(O)(CH₂)_mNR₁₂R₁₃ or
 - 10 O(CH₂)_nCH(OH)(CH₂)_nN(R₁₂R₁₃);
 - (III) a group CH=C(W)V; or
 - (IV) a cyclohexyl group;
- and where appropriate, the other of R₁ and R₂ is a phenyl group optionally substituted by one or more groups
- 15 independently selected from halogen, nitro, methoxy, NHC(O)R₁₂, CO₂H, O(CH₂)_nN(R₁₂R₁₃) and CH₂Y(CH₂)_nN(R₁₂R₁₃);
 - R₃ is C₁-C₄ alkyl or (CH₂)_nC(O)OR₁₂;
 - X is a naphthyl group or a five- or six-membered saturated or unsaturated heterocyclic group containing one or more
 - 20 heteroatoms, which heteroatoms may be the same or different and are independently selected from O, N and S; the heteroatom(s) when nitrogen being optionally substituted by hydrogen, methyl, oxygen, tertiary-butyloxycarbonyl, -(CH₂)_nCH₂OH or SO₂Me; the heterocyclic ring being
 - 25 optionally substituted by halogen, Me, MeS, phenyl, O(CH₂)_nNR₁₂R₁₃, -N(R₁₂)(CH₂)_nN(R₁₂R₁₃), -(CH₂)_nN(R₁₂R₁₃) or -O(CH₂)_nO(CH₂)_nN(R₁₂R₁₃), or the heterocyclic ring optionally containing one or more carbonyl groups and being optionally

fused to a benzene ring, which benzene ring is optionally substituted by 1 or 2 C₁-C₆ alkoxy groups;

Y is O or S;

Z is a C₃-C₆ cycloalkyl group;

5 R₁₂, R₁₃ and R₁₄, which may be the same or different, are hydrogen or C₁-C₆ alkyl;

W is hydrogen or a phenyl group;

V is a phenyl group optionally substituted by one or more groups independently selected from nitro, alkoxy and

10 O(CH₂)_nNR₁₂R₁₃;

m and n are each, independently, 0 or an integer having the value 1, 2, 3 or 4;

O(CH₂)_nNR₁₂R₁₃ or containing one or more carbonyl groups and being optionally fused to a benzene ring;

15 Z is a C₃-C₆ cycloalkyl group;

R₁₂, R₁₃ and R₁₄, which may be the same or different, are hydrogen or C₁-C₄ alkyl;

W is hydrogen or a phenyl group;

V is a phenyl group optionally substituted by one or more groups independently selected from nitro, alkoxy and

20 O(CH₂)_nNR₁₂R₁₃;

m and n are, independently, integers having the values 1, 2, 3 or 4;

or a pharmaceutically acceptable salt or ester thereof; in

25 the manufacture of a medicament for use as an inhibitor of plasminogen activator inhibitor.

Relevant Technical Fields		Search Examiner P N DAVEY
(i) UK Cl (Ed.N)	C2C (CTR)	Date of completion of Search 20 MARCH 1995
(ii) Int Cl (Ed.6)		Documents considered relevant following a search in respect of Claims :- 1-10
Databases (see below) (i) UK Patent Office collections of GB, EP, WO and US patent specifications. (ii) ONLINE: CAS ONLINE		

Categories of documents

- X: Document indicating lack of novelty or of inventive step.
- Y: Document indicating lack of inventive step if combined with one or more other documents of the same category.
- A: Document indicating technological background and/or state of the art.
- P: Document published on or after the declared priority date but before the filing date of the present application.
- E: Patent document published on or after, but with priority date earlier than, the filing date of the present application.
- &: Member of the same patent family; corresponding document.

Category	Identity of document and relevant passages	Relevant to claim(s)
X	Chemical Abstracts 65: 16969e see compound with CAS Registry No 7670-71-5	1 at least

Databases: The UK Patent Office database comprises classified collections of GB, EP, WO and US patent specifications as outlined periodically in the Official Journal (Patents). The on-line databases considered for search are also listed periodically in the Official Journal (Patents).