



US 20100196409A1

(19) **United States**

(12) **Patent Application Publication**
Thonnard

(10) **Pub. No.: US 2010/0196409 A1**

(43) **Pub. Date: Aug. 5, 2010**

(54) **BASB006 POLYPEPTIDES FROM NEISSERIA
MENINGITIDIS AND IMMUNOGENIC
COMPOSITIONS THEREOF**

division of application No. 09/673,896, filed on Dec.
18, 2000, now Pat. No. 6,696,062, filed as application
No. PCT/EP99/02766 on Apr. 20, 1999.

(75) Inventor: **Joelle Thonnard, Gembloux (BE)**

Correspondence Address:
**SEED INTELLECTUAL PROPERTY LAW
GROUP PLLC
701 FIFTH AVE, SUITE 5400
SEATTLE, WA 98104 (US)**

(30) **Foreign Application Priority Data**

Apr. 24, 1998 (GB) 9808866.9

(73) Assignee: **GLAXOSMITHKLINE
BIOLOGICALS S.A., Rixensart
(BE)**

Publication Classification

(51) **Int. Cl.**
A61K 39/095 (2006.01)
C07K 14/22 (2006.01)
A61P 31/04 (2006.01)
A61P 37/04 (2006.01)

(52) **U.S. Cl.** **424/190.1; 530/350**

(21) Appl. No.: **12/180,055**

(57) **ABSTRACT**

(22) Filed: **Jul. 25, 2008**

The invention provides BASB006 polypeptides and poly-
nucleotides encoding BASB006 polypeptides and methods
for producing such polypeptides by recombinant techniques.
Also provided are diagnostic, prophylactic and therapeutic
uses.

Related U.S. Application Data

(60) Continuation of application No. 10/742,345, filed on
Dec. 19, 2003, now Pat. No. 7,419,824, which is a

**Identity to SeqID No:1 is indicated by a dot, while a dash (-) indicates a nonexistent
nucleotide.**

```

                *           20           *           40           *
Seqid1 : ATGAAAAACAACCGACAAACGGACAACCGAAACACACCGCAAAGCCCCGAA : 50
Seqid3 : ..... : 50

                60           *           80           *           100
Seqid1 : AACCGGTGCGCATCCGCTTCTCGCCTGCTTACTTAGCCATATGCCTGTCGT : 100
Seqid3 : .....C..... : 100

                *           120           *           140           *
Seqid1 : TCGGCATTCTTCCCAAGCCTGGGCGGGACACACTTATTTGCGCATCAAC : 150
Seqid3 : ..... : 150

                160           *           180           *           200
Seqid1 : TACCAATACTATCGCGACTTTGCCGAAAAATAAAGGCAAGTTTGCAGTCGG : 200
Seqid3 : ..... : 200

                *           220           *           240           *
Seqid1 : GGCGAAAGATATTGAGGTTTACAACAAAAAGGGGAGTTGGTCGGCAAAT : 250
Seqid3 : ..... : 250

                260           *           280           *           300
Seqid1 : CAATGACAAAAGCCCCGATGATTGATTTTCTGTGGTGTGCGGTAACGGC : 300
Seqid3 : ..... : 300

                *           320           *           340           *
Seqid1 : GTGGCGGCATTGGTGGGCGATCAATATATTGTGAGCGTGGCACATAACGG : 350
Seqid3 : ..... : 350

```

Figure 1A

Identity to SeqID No:1 is indicated by a dot, while a dash (-) indicates a nonexistent nucleotide.

```

                *           20           *           40           *
Seqid1 : ATGAAAACAACCGACAACCGACAACCGAAACACACCGCAAAGCCCCGAA : 50
Seqid3 : ..... : 50

                60           *           80           *           100
Seqid1 : AACCGGTCGCATCCGCTTCTCGCCTGCTTACTTAGCCATATGCCTGTCGT : 100
Seqid3 : .....C..... : 100

                *           120           *           140           *
Seqid1 : TCGGCATTCTTCCCCAAGCCTGGGCGGGACACACTTATTTTCGGCATCAAC : 150
Seqid3 : ..... : 150

                160           *           180           *           200
Seqid1 : TACCAATACTATCGCGACTTTGCCGAAAATAAAGGCAAGTTTGCAGTCGG : 200
Seqid3 : ..... : 200

                *           220           *           240           *
Seqid1 : GCGAAAAGATATTGAGGTTTACAACAAAAAAGGGGAGTTGGTTCGGCAAAT : 250
Seqid3 : ..... : 250

                260           *           280           *           300
Seqid1 : CAATGACAAAAGCCCCGATGATTGATTTTCTGTGGTGTTCGCGTAACGGC : 300
Seqid3 : ..... : 300

                *           320           *           340           *
Seqid1 : GTGGCGGCATTGGTGGGCGATCAATATATTGTGAGCGTGGCACATAACGG : 350
Seqid3 : ..... : 350
    
```

Figure 1B

```

          360          *          380          *          400
Seqid1 : CGGCTATAACAACGTTGATTTTGGTGCGGAGGGAAGCAATCCCGATCAGC : 400
Seqid3 : .....A.....A.....A..... : 400

          *          420          *          440          *
Seqid1 : ACCGTTTTTCATTATCAAATTGTGAAAAGAAATAATTATAAAGCAGGGACT : 450
Seqid3 : .T.....A.....A.....C.G..... : 450

          460          *          480          *          500
Seqid1 : AACGGTCATCCTTATGGTGGCGATTATCATATGCCGCGTTTACATAAAATT : 500
Seqid3 : ..A..C..... : 500

          *          520          *          540          *
Seqid1 : TGTAACCGATGCAGAACCTGTTGAAATGACCAGTTATATGGATGGGCGGA : 550
Seqid3 : ...C..A..... : 550

          560          *          580          *          600
Seqid1 : AATATATCGATCAAATAATTACCCTGACCGTGTTCGTATTGGGGCAGGC : 600
Seqid3 : ..... : 600

          *          620          *          640          *
Seqid1 : AGGCAATATTGGCGATCTGATGAAGATGAGCCCAATAACCGCGAAAGTTC : 650
Seqid3 : ..... : 650

          660          *          680          *          700
Seqid1 : ATATCATATTGCAAGTGCGTATTCTTGGCTCGTTGGTGGCAATACCTTTG : 700
Seqid3 : ..... : 700

          *          720          *          740          *
Seqid1 : CACAAAATGGATCAGGTGGTGGCACAGTCAACTTAGGTAGTGAAAAAATT : 750
Seqid3 : ..... : 750
```

Figure 1C

Seqid1 : AAACATAGCCCATATGGT*****TACCAACAGGAGGCTCATTTGGCGACAG : 800
Seqid3 : : 800

Seqid1 : TGGCTCACCAATGTTTATCTATGATGCCCAAAAGCAAAAGTGGTTAATTA : 850
Seqid3 : : 850

Seqid1 : ATGGGGTATTGCAAACGGGCAACCCCTATATAGGAAAAAGCAATGGCTTC : 900
Seqid3 : : 900

Seqid1 : CAGCTGGTTCGTAAAGATTGGTTCTATGATGAAATCTTTGCTGGAGATAC : 950
Seqid3 : : 950

Seqid1 : CCATTCAGTATTCTACGAACCACATCAAAATGGGAAATACACTTTTCACG : 1000
Seqid3 :G.....T....A... : 1000

Seqid1 : ACAATAATAATGGCACAGGAAAAATCAATGCCAAACATGAACACAATTCT : 1050
Seqid3 : ..G..... : 1050

Seqid1 : CTGCCTAATAGATTAAAAACACGAACCGTTCAATTGTTTAAATGTTTCTTT : 1100
Seqid3 : : 1100

Seqid1 : ATCCGAGACAGCAAGAGAACCTGTTTATCATGCTGCAGGTGGTGTCAACA : 1150
Seqid3 : : 1150

Figure 1D

Seqid1 : 1160 * 1180 * 1200 : 1200
Seqid3 : : 1200

Seqid1 : * 1220 * 1240 * : 1250
Seqid3 : : 1250

Seqid1 : 1260 * 1280 * 1300 : 1300
Seqid3 : : 1300

Seqid1 : * 1320 * 1340 * : 1350
Seqid3 : : 1350

Seqid1 : 1360 * 1380 * 1400 : 1400
Seqid3 : : 1400

Seqid1 : * 1420 * 1440 * : 1450
Seqid3 : : 1450

Seqid1 : 1460 * 1480 * 1500 : 1500
Seqid3 : ...C...C...G...G...G...C...T...AGGC..... : 1500

Seqid1 : * 1520 * 1540 * : 1550
Seqid3 : : 1550

Figure 1E

Seqid1 : CGATAATCAGTTCAACCCCGACAAACTCTATTTTCGGCTTTCGGCGCGGAC : 1600
Seqid3 : : 1600

Seqid1 : GTTTGGATTGAAACGGGCATTTCGCTTTCGTTCCACCGTATTCAAAATACC : 1650
Seqid3 :A..... : 1650

Seqid1 : GATGAAGGGCGATGATTGTCAACCACAATCAAGACAAAGAATCCACCGT : 1700
Seqid3 : : 1700

Seqid1 : TACCATTACAGGCAATAAAGATATTGCTACAACCGGCAATAACAACAGCT : 1750
Seqid3 : : 1750

Seqid1 : TGGATAGCAAAAAAGAAATTGCCTACAACGGTTGGTTTGGCGAGAAAGAT : 1800
Seqid3 : : 1800

Seqid1 : ACGACCAAACGAACGGGCGGCTCAACCTTGTTTACCAGCCCGCCGAGA : 1850
Seqid3 : : 1850

Seqid1 : AGACCGCACCCCTGCTGCTTTCGGCGGAACAAATTTAAACGGTAACATCA : 1900
Seqid3 :C..... : 1900

Seqid1 : CGCAAACAAACGGCAAACGTGTTTTTCAGCGGCAGACCGCACCGCC : 1950
Seqid3 :A..... : 1950

Figure 1F

Seqid1 : TACAATCATTTAGGAAGCGGGTGGTCAAAAATGGAAGGTATCCCACAAGG : 2000
Seqid3 :AACGA.CAT...GC...AA..G..C..T...GC.. : 2000

Seqid1 : AGAAATCGTGTGGGACAACGACTGGATCAACCGCACGTTTAAAGCGGAAA : 2050
Seqid3 : G.....A..... : 2050

Seqid1 : ATTTCCATATTCAGGGCGGGCAGGCGGTGATTTCCCGCAATGTTGCCAAA : 2100
Seqid3 : .C....A..A.A....A.....G..... : 2100

Seqid1 : GTGGAAGGCGATTGGCATTGAGCAATCACGCCCAAGCAGTTTTTGGTGT : 2150
Seqid3 : ...A..... : 2150

Seqid1 : CGCACCGCATCAAAGCCACACAATCTGTACACGTTTCGGACTGGACGGGTC : 2200
Seqid3 : : 2200

Seqid1 : TGACAAATTGTGTCGAAAAAACCATTACCGACGATAAAGTGATTGCTTCA : 2250
Seqid3 : : 2250

Seqid1 : TTGACTAAGACCGACATCAGCGGCAATGTCAGCCTTGCCGATCAGCTCA : 2300
Seqid3 :GAT..... : 2300

Seqid1 : TTTAAATCTCACAGGGCTTGCCCACTCAACGGCAATCTTAGTGCAAATG : 2350
Seqid3 : : 2350

Figure 1G

```

                2360          *          2380          *          2400
Seqid1 : GCGATACACGTTATACAGTCAGCCACAACGCCACCCAAAACGGCGACCTT : 2400
Seqid3 : .....A..... : 2400

                *          2420          *          2440          *
Seqid1 : AGCCTCGTGGGCAATGCCCAAGCAACATTTAATCAAGCCACATTAAACGG : 2450
Seqid3 : ..... : 2450

                2460          *          2480          *          2500
Seqid1 : CAACACATCGGCTTCGGGCAATGCTTCATTTAATCTAAGCAACAACGCCG : 2500
Seqid3 : .....G..C..... : 2500

                *          2520          *          2540          *
Seqid1 : TACAAAACGGCAGTCTGACGCTTCCGGCAACGCTAAGGCAAACGTAAGC : 2550
Seqid3 : ..... : 2550

                2560          *          2580          *          2600
Seqid1 : CATTCCGCACTCAACGGTAATGTCTCCCTAGCCGATAAGGCAGTATTCCA : 2600
Seqid3 : ..... : 2600

                *          2620          *          2640          *
Seqid1 : TTTTGAAAGCAGCCGCTTTACCGGACAAATCAGCGGCAGCAAGGATACGG : 2650
Seqid3 : .....G..... : 2650

                2660          *          2680          *          2700
Seqid1 : CATTACACTTAAAAGACAGCGAATGGACGCTGCCGTCAGGCACGGAATTA : 2700
Seqid3 : ..... : 2700

                *          2720          *          2740          *
Seqid1 : GGCAATTTAAACCTTGACAACGCCACCATTACACTCAATTCCGCCTATCG : 2750
Seqid3 : ..... : 2750
    
```


Figure 1H

2760 * 2780 * 2800
 Seqid1 : CCACGATGCGGCAGGGGCGCAAACCGGCAGTGCACAGATGCGCCGCGC- : 2799
 Seqid3 :C : 2800

* 2820 * 2840 *
 Seqid1 : -----CGCCGTTGCGCGCGTTCCTATTATCCGTTACACCTCCGGCT : 2841
 Seqid3 : GCCGTTGCG.....G..AA.. : 2850

2860 * 2880 * 2900
 Seqid1 : TCGGCAGAATCCCATTTCAACACGCTGACGGTAAACGGCAAATGAACGG : 2891
 Seqid3 : ...T.....G..... : 2900

* 2920 * 2940 *
 Seqid1 : TCAGGGAACATTCCGCTTTATGTCGGAACCTTCGGCTACCGAAGCGACA : 2941
 Seqid3 :C..... : 2950

2960 * 2980 * 3000
 Seqid1 : AATTGAAGCTGGCGGAAAGTTCCGAAGGCACTTACACCTTGGCGGTCAAC : 2991
 Seqid3 : : 3000

* 3020 * 3040 *
 Seqid1 : AATACCGCAACGAACCCGTAAGCCTCGATCAATTGACGGTAGTGAAGG : 3041
 Seqid3 :T.C.....T..A..... : 3050

3060 * 3080 * 3100
 Seqid1 : GAAAGACAACAAACCGCTGTCCGAAAACCTTAATTTACCCCTGCAAACG : 3091
 Seqid3 : A.....T.....T..... : 3100

* 3120 * 3140 *
 Seqid1 : AACACGTCGATGCCGGCGCGTGGCGTTACCAACTCATCCGCAAAGACGGC : 3141
 Seqid3 : : 3150

Figure 1I

Seqid1 : GAGTTCCGCCTGCATAATCCGGTCAAAGAACAAGAGCTTTCGACAAACT : 3191
Seqid3 : : 3200

Seqid1 : CGGCAAGGCAGAAGCCAAAAACAGGCGGGAAAAGACAACGCGCAAAGCC : 3241
Seqid3 :A..... : 3250

Seqid1 : TTGACGCGCTGATTGCGGCCGGGCGCGATGCCGTCGAAAAGACAGAAAGC : 3291
Seqid3 : : 3300

Seqid1 : GTTGCCGAACCGGCCCGGCAGGCAGGCGGGGAAAATGTCGGCATTATGCA : 3341
Seqid3 : : 3350

Seqid1 : GCGCGAGGAAGAGAAAAACGGGTGCAGGCGGATAAAGACACCGCCTTGG : 3391
Seqid3 : : 3400

Seqid1 : CGAAACAGCGCGAAGGGAAAACCGGCCGGCTACCACCGCCTTCCCCCGC : 3441
Seqid3 :C.G..... : 3450

Seqid1 : GCCCGCCGCGCCCGCGGGATTGCGCAACCGCAGCCCCAACCGCAACC : 3491
Seqid3 :T..A.....G.. : 3500

Seqid1 : CCAACCGCAGCGCGACCTGATCAGCCGTTATGCCAATAGCGGTTTGAGTG : 3541
Seqid3 : : 3550

Figure 1J

Seqid1 : AATTTTCCGCCACGCTCAACAGCGTTTTCGCCGTACAGGACGAATTAGAC : 3591
Seqid3 : : 3600

Seqid1 : CGCGTATTTGCCGAAGACCCGCCAACGCCGTTTGGACAAGCGGCATCCG : 3641
Seqid3 :A..... : 3650

Seqid1 : GGACACCAAACACTACCGTTCGCAAGATTTCGCGCCTACCGCCAACAAA : 3691
Seqid3 : : 3700

Seqid1 : CCGACCTGCGCCAAATCGGTATGCAGAAAACCTCGGCAGCGGGCGCGTC : 3741
Seqid3 : : 3750

Seqid1 : GGCATCCTGTTTTCGCACAACCCGGACCGAAAACACCTTCGACGACGGCAT : 3791
Seqid3 : : 3800

Seqid1 : CGGCAACTCGGCACGGCTTGCCCACGGCGCCGTTTTCGGGCAATACGGCA : 3841
Seqid3 : : 3850

Seqid1 : TCGGCAGGTTTCGACATCGGCATCAGCACGGGCGCGGGTTTTAGCAGCGGC : 3891
Seqid3 : ...A.....T.....G..... : 3900

Seqid1 : AGTCTTTCAGACGACATCGGAAGCAAATCCGCCCGCGTGCTGCATTA : 3941
Seqid3 : ..C.....G.....G..... : 3950

Figure 1K

3960 * 3980 * 4000
Seqid1 : CGGCATTCAGGCACGATACCGCGCCGGTTTCGGCGGATTCGGCATCGAAC : 3991
Seqid3 : : 4000

* 4020 * 4040 *
Seqid1 : CGCACATCGGCGCAACGCGCTATTTTCGTCCAAAAAGCGGATTACCGCTAC : 4041
Seqid3 : : 4050

4060 * 4080 * 4100
Seqid1 : GAAAACGTCAATATCGCCACCCCGGCTTTCGTTCAACCGCTACCGCGC : 4091
Seqid3 :A..... : 4100

* 4120 * 4140 *
Seqid1 : GGGCATTAAAGGCAGATTATTCATTCAAACCGGCGCAACACATTTCCATCA : 4141
Seqid3 : : 4150

4160 * 4180 * 4200
Seqid1 : CGCCTTATTTGAGCCTGTCTTATACCGATGCCGCTTCGGGCAAAGTCCGA : 4191
Seqid3 : : 4200

* 4220 * 4240 *
Seqid1 : ACGCGCGTCAATACCGCCGTATTGGCTCAGGATTTTCGGCAAACCCGCAG : 4241
Seqid3 : ..A..... : 4250

4260 * 4280 * 4300
Seqid1 : TGCGGAATGGGGCGTAAACGCCGAAATCAAAGGTTTCACGCTGTCCCTCC : 4291
Seqid3 :C..... : 4300

* 4320 * 4340 *
Seqid1 : ACGCTGCCGCGCCCAAAGCCGCAACTGGAAGCGCAACACAGCGCGGGC : 4341
Seqid3 : : 4350

Figure 1L

4360 *
Seqid1 : ATCAAATTAGGCTACCGCTGGTAA : 4365
Seqid3 : : 4374

Figure 2A

Identity to SeqID No:2 is indicated by a dot, while a dash (-) indicates a nonexistent amino acid.

```

                *           20           *           40           *
Seqid2 : MKTTDKRTTETHRKAPKTGRIRFSPAYLAICLSFGILPQAWAGHTYFGIN : 50
Seqid4 : ..... : 50

                60           *           80           *           100
Seqid2 : YQYYRDFAEKNGKFAVGAKDIEVYNKKGELVGKSMTKAPMIDFSVVS RNG : 100
Seqid4 : ..... : 100

                *           120           *           140           *
Seqid2 : VAALVGDQYIVSVAHNGGYNNVDFGAEGSNPDQHRFSYQIVKRNNYKAGT : 150
Seqid4 : .....R.....T.K..... : 150

                160           *           180           *           200
Seqid2 : NGHPYGGDYHMPRLHKFVTD AEPVEMTSYMDGRKYIDQNNYPDRVRIGAG : 200
Seqid4 : K..... : 200

                *           220           *           240           *
Seqid2 : RQYWRSD EDEPNNRESSYHIASAYSWL VGGNTFAQNGSGGGTVNLGSEKI : 250
Seqid4 : ..... : 250

                260           *           280           *           300
Seqid2 : KHSPYGF LPTGG SFGDSGSPMFIYDAQKQKWLINGVLQ TGNPYIGKSNGF : 300
Seqid4 : ..... : 300

                *           320           *           340           *
Seqid2 : QLVRKDW FYDEIFAGDTHSVFYEPHQNGKYTFHDNNNGTGKINAKHEHNS : 350
Seqid4 : .....R.....S.N.D..... : 350
    
```

Figure 2B

Seqid2 : LPNRLKTRTVQLFNVSLSETAREPVYHAAGGVNSYRPRLNNGENISFIDE : 400
Seqid4 : : 400

Seqid2 : GKGEILILTSNINQGAGGLYFQGDFTVSPENNETWQGAGVHISEDSTVTWK : 450
Seqid4 : : 450

Seqid2 : VNGVANDRLSKIGKGTLHVQAKGENQGSISVGDGKVVILDQQADENKKQA : 500
Seqid4 :T.....DKG... : 500

Seqid2 : FSEIGLVSGRGTVQLNADNQFNPKLYFGFRGGRLDLNGHSLSFHRIQNT : 550
Seqid4 : : 550

Seqid2 : DEGAMIVNHNQDKESTVTITGNKDIAT'GNNNSLDSKKEIAYNGWFGEKD : 600
Seqid4 : : 600

Seqid2 : TTKTNGRLNLVYQPAEDRTLLLSGGTNLNGNITQTNGKLFFSGRPTPHA : 650
Seqid4 : : 650

Seqid2 : YNHLGSGWSKMEGIPQGEIVWDNDWINRTFKAENFHIQGGQAVISRNVAK : 700
Seqid4 : ...NDH..QK...R.....Q.K...V..... : 700

Seqid2 : VEGDWHLNSNHAQAVFGVAPHQSHTICTRSDWTGLTNCVEKTIITDDKVIAS : 750
Seqid4 : .K..... : 750

Figure 2C

Seqid2 : LTKTDISGNVSLADHAHLNLTGLATLNGNLSANGDTRYTVSHNATQNGDL : 800
 Seqid4 :D.....N. : 800

Seqid2 : SLVGNAQATFNQATLNGNTSASGNASFNLSNNAVQNGSLTSLGNAKANVS : 850
 Seqid4 :DH..... : 850

Seqid2 : HSALNGNVSLADKAVFHFESSRFTGQISGSKDTALHLKDSEWTLPSGTEL : 900
 Seqid4 :G..... : 900

Seqid2 : GNLNLDNATITLNSAYRHDAAGAQTGSATDAPR---RRSRRLLSVTPPA : 947
 Seqid4 :RRS.....T : 950

Seqid2 : SAESHFNLTIVNGKLNQGTFRFMSELFGYRSDKLKLAESSEGYTLAVN : 997
 Seqid4 : .V..R..... : 1000

Seqid2 : NTGNEPVS LDQLTVVEGKDNKPLSENLFNLQNEHVDAGAWRYQLIRKDG : 1047
 Seqid4 :A..E.....F..... : 1050

Seqid2 : EFRLHNPVKEQELSDKLGKAEAKQAGKDNAQSLDALIAAGRDAVEKTES : 1097
 Seqid4 :E..... : 1100

Seqid2 : VAEPARQAGGENVGMQAEKRVQADKDTALAKQREGKTRPATTAFPR : 1147
 Seqid4 :AE..... : 1150

Figure 2D

Seqid2 : 1160 * 1180 * 1200
Seqid2 : ARRARRDLPPQPQPQPQPQRDLISRYANSGLSEFSATLNSVFAVQDELD : 1197
Seqid4 :L..... : 1200

Seqid2 : * 1220 * 1240 *
Seqid2 : RVFAEDRRNAVWTSGIRDTKHYRSQDFRAYRQQTDLRQIGMQKNLGSGRV : 1247
Seqid4 :E..... : 1250

Seqid2 : 1260 * 1280 * 1300
Seqid2 : GILFSHNRTENTFDDGIGNSARLAHGAVFGQYGIGRFDIGISTGAGFSSG : 1297
Seqid4 :D..Y...A..... : 1300

Seqid2 : * 1320 * 1340 *
Seqid2 : SLSDDIGSKIRRRVLHYGIQARYRAGFGGFGIEPHIGATRYFVQKADYRY : 1347
Seqid4 :G..G..... : 1350

Seqid2 : 1360 * 1380 * 1400
Seqid2 : ENVNIATPGLAFNRYRAGIKADYSFKPAQHISITPYLSLSYTDAAASGKVR : 1397
Seqid4 : : 1400

Seqid2 : * 1420 * 1440 *
Seqid2 : TRVNTAVLAQDFGKTRSAEWGVNAEIKGFTLSLHAAAAGKGPQLEAQSAG : 1447
Seqid4 : : 1450

Seqid2 : IKLGYRW : 1454
Seqid4 : : 1457

Figure 3





Figure 4



**BASB006 POLYPEPTIDES FROM NEISSERIA
MENINGITIDIS AND IMMUNOGENIC
COMPOSITIONS THEREOF**

**CROSS REFERENCE TO RELATED
APPLICATIONS**

[0001] This application is a divisional application of U.S. patent application Ser. No. 09/673,896 filed Dec. 18, 2000, which is the National Stage Application of International Application No. PCT/EP99/02766, filed Apr. 20, 1999 which was published under PCT article 21(2) in English, which claims the benefit of priority of Great Britain Patent Application Serial No. 9808866.9, filed Apr. 24, 1998. The disclosure of these applications are herein incorporated by reference in their entirety.

FIELD OF THE INVENTION

[0002] This invention relates to polynucleotides, (herein referred to as "BASB006 polynucleotide(s)"), polypeptides encoded by them (referred to herein as "BASB006" or "BASB006 polypeptide(s)"), recombinant materials and methods for their production. In another aspect, the invention relates to methods for using such polypeptides and polynucleotides, including vaccines against bacterial infections. In a further aspect, the invention relates to diagnostic assays for detecting infection of certain pathogens.

BACKGROUND OF THE INVENTION

[0003] *Neisseria meningitidis* (meningococcus) is a Gram negative bacterium frequently isolated from the human upper respiratory tract. It occasionally causes invasive bacterial diseases such as bacteremia and meningitis. The incidence of meningococcal disease shows geographical seasonal and annual differences (Schwartz, B., Moore, P. S., Broome, C. V.; Clin. Microbiol. Rev. 2 (Supplement), S18-S24, 1989). Most disease in temperate countries is due to strains of serogroup B and varies in incidence from 1-10/100,000/year total population sometimes reaching higher values (Kaczmarek, E. B. (1997), Commun. Dis. Rep. Rev. 7: R55-9, 1995; Scholten, R. J. P. M., Bijlmer, H. A., Poolman, J. T. et al. Clin. Infect. Dis. 16: 237-246, 1993; Cruz, C., Pavez, G., Aguilar, E., et al. Epidemiol. Infect. 105: 119-126, 1990).

[0004] Epidemics dominated by serogroup A meningococci, mostly in central Africa, are encountered, sometimes reaching levels up to 1000/100,000/year (Schwartz, B., Moore, P. S., Broome, C. V. Clin. Microbiol. Rev. 2 (Supplement), S18-S24, 1989). Nearly all cases as a whole of meningococcal disease are caused by serogroup A, B, C, W-135 and Y meningococci and a tetravalent A, C, W-135, Y polysaccharide vaccine is available (Armand, J., Arminjon, F., Mynard, M. C., Lafaix, C., J. Biol. Stand. 10: 335-339, 1982).

[0005] The polysaccharide vaccines are currently being improved by way of chemical conjugating them to carrier proteins (Lieberman, J. M., Chiu, S. S., Wong, V. K., et al. JAMA 275:1499-1503, 1996).

[0006] A serogroup B vaccine is not available, since the B capsular polysaccharide was found to be nonimmunogenic, most likely because it shares structural similarity to host components (Wyle, F. A., Artenstein, M. S., Brandt, M. L. et al. J. Infect. Dis. 126: 514-522, 1972; Finne, J. M., Leinonen, M., Mäkelä, P. M. Lancet ii.: 355-357, 1983).

[0007] For many years efforts have been initiated and carried out to develop meningococcal outer membrane based vaccines (de Moraes, J. C., Perkins, B., Camargo, M. C. et al. Lancet 340: 1074-1078, 1992; Bjune, G., Høiby, E. A. Gronnesby, J. K. et al. 338: 1093-1096, 1991). Such vaccines have demonstrated efficacies from 57%-85% in older children (>4 years) and adolescents.

[0008] Many bacterial outer membrane components are present in these vaccines, such as PorA, PorB, Rmp, Opc, Opa, FrpB and the contribution of these components to the observed protection still needs further definition. Other bacterial outer membrane components have been defined by using animal or human antibodies to be potentially relevant to the induction of protective immunity, such as TbpB and NspA (Martin, D., Cadieux, N., Hamel, J., Brodeux, B. R., J. Exp. Med. 185: 1173-1183, 1997; Lissolo, L., Maître-Wilmotte, C., Dumas, p. et al., Inf. Immun. 63: 884-890, 1995). The mechanisms of protective immunity will involve antibody mediated bactericidal activity and opsonophagocytosis.

[0009] A bacteremia animal model has been used to combine all antibody mediated mechanisms (Saukkonen, K., Leinonen, M., Abdillahi, H. Poolman, J. T. Vaccine 7: 325-328, 1989). It is generally accepted that the late complement component mediated bactericidal mechanism is crucial for immunity against meningococcal disease (Ross, S. C., Rosenthal P. J., Berberic, H. M., Densen, P. J. Infect. Dis. 155: 1266-1275, 1987).

[0010] The frequency of *Neisseria meningitidis* infections has risen dramatically in the past few decades. This has been attributed to the emergence of multiply antibiotic resistant strains and an increasing population of people with weakened immune systems. It is no longer uncommon to isolate *Neisseria meningitidis* strains that are resistant to some or all of the standard antibiotics. This phenomenon has created an unmet medical need and demand for new anti-microbial agents, vaccines, drug screening methods, and diagnostic tests for this organism.

SUMMARY OF THE INVENTION

[0011] The present invention relates to BASB006, in particular BASB006 polypeptides and BASB006 polynucleotides, recombinant materials and methods for their production. In another aspect, the invention relates to methods for using such polypeptides and polynucleotides, including prevention and treatment of microbial diseases, amongst others. In a further aspect, the invention relates to diagnostic assays for detecting diseases associated with microbial infections and conditions associated with such infections, such as assays for detecting expression or activity of BASB006 polynucleotides or polypeptides.

[0012] Various changes and modifications within the spirit and scope of the disclosed invention will become readily apparent to those skilled in the art from reading the following descriptions and from reading the other parts of the present disclosure.

BRIEF DESCRIPTION OF THE DRAWINGS

[0013] FIGS. 1A-1L show consecutive segments of sequence alignment for two BASB006-encoding polynucleotides.

[0014] FIGS. 2A-2D show consecutive segments of sequence alignment for two BASB006 polypeptides.

[0015] FIG. 3 shows a Coomassie stain SDS-PAGE gel analysis of purified BASB006 protein.

[0016] FIG. 4 shows a Western-blot of partially purified recombinant BASB006 protein probed with mice sera containing anti-BASB006 antibodies.

[0017] FIG. 5 shows a Western-blot of partially purified recombinant BASB006 protein probed with human convalescent sera and mice sera.

DESCRIPTION OF THE INVENTION

[0018] The invention relates to BASB006 polypeptides and polynucleotides as described in greater detail below. In particular, the invention relates to polypeptides and polynucleotides of BASB006 of *Neisseria meningitidis*, which is related by amino acid sequence homology to *H. influenzae* Hap polypeptide. The invention relates especially to BASB006 having the nucleotide and amino acid sequences set out in SEQ ID NO:1,3 and SEQ ID NO:2,4 respectively. It is understood that sequences recited in the Sequence Listing below as "DNA" represent an exemplification of one embodiment of the invention, since those of ordinary skill will recognize that such sequences can be usefully employed in polynucleotides in general, including ribopolynucleotides.

Polypeptides

[0019] In one aspect of the invention there are provided polypeptides of *Neisseria meningitidis* referred to herein as "BASB006" and "BASB006 polypeptides" as well as biologically, diagnostically, prophylactically, clinically or therapeutically useful variants thereof, and compositions comprising the same.

[0020] The present invention further provides for:

(a) an isolated polypeptide which comprises an amino acid sequence which has at least 85% identity, more preferably at least 90% identity, yet more preferably at least 95% identity, most preferably at least 97-99% or exact identity, to that of SEQ ID NO:2, 4;

(b) a polypeptide encoded by an isolated polynucleotide comprising a polynucleotide sequence which has at least 85% identity, more preferably at least 90% identity, yet more preferably at least 95% identity, even more preferably at least 97-99% or exact identity to SEQ ID NO:1, 3 over the entire length of SEQ ID NO:1, 3 respectively; or

(c) a polypeptide encoded by an isolated polynucleotide comprising a polynucleotide sequence encoding a polypeptide which has at least 85% identity, more preferably at least 90% identity, yet more preferably at least 95% identity, even more preferably at least 97-99% or exact identity, to the amino acid sequence of SEQ ID NO:2, 4;

[0021] The BASB006 polypeptides provided in SEQ ID NO:2,4 are the BASB006 polypeptides from *Neisseria meningitidis* strains American Type Culture Collection 13090 (herein "ATCC13090") and H44/76.

[0022] The invention also provides an immunogenic fragment of a BASB006 polypeptide, that is, a contiguous portion of the BASB006 polypeptide which has the same or substantially the same immunogenic activity as the polypeptide comprising the amino acid sequence of SEQ ID NO:2,4. That is to say, the fragment (if necessary when coupled to a carrier) is capable of raising an immune response which recognises the BASB006 polypeptide. Such an immunogenic fragment may include, for example, the BASB006 polypeptide lacking an N-terminal leader sequence, and/or a transmembrane domain

and/or a C-terminal anchor domain. In a preferred aspect the immunogenic fragment of BASB006 according to the invention comprises substantially all of the extracellular domain of a polypeptide which has at least 85% identity, more preferably at least 90% identity, yet more preferably at least 95% identity, most preferably at least 97-99% identity, to that of SEQ ID NO:2,4 over the entire length of SEQ ID NO:2

[0023] A fragment is a polypeptide having an amino acid sequence that is entirely the same as part but not all of any amino acid sequence of any polypeptide of the invention. As with BASB006 polypeptides, fragments may be "free-standing," or comprised within a larger polypeptide of which they form a part or region, most preferably as a single continuous region in a single larger polypeptide.

[0024] Preferred fragments include, for example, truncation polypeptides having a portion of an amino acid sequence of SEQ ID NO:2,4 or of variants thereof, such as a continuous series of residues that includes an amino- and/or carboxyl-terminal amino acid sequence. Degradation forms of the polypeptides of the invention produced by or in a host cell, are also preferred. Further preferred are fragments characterized by structural or functional attributes such as fragments that comprise alpha-helix and alpha-helix forming regions, beta-sheet and beta-sheet-forming regions, turn and turn-forming regions, coil and coil-forming regions, hydrophilic regions, hydrophobic regions, alpha amphipathic regions, beta amphipathic regions, flexible regions, surface-forming regions, substrate binding region, and high antigenic index regions.

[0025] Further preferred fragments include an isolated polypeptide comprising an amino acid sequence having at least 15, 20, 30, 40, 50 or 100 contiguous amino acids from the amino acid sequence of SEQ ID NO:2,4, or an isolated polypeptide comprising an amino acid sequence having at least 15, 20, 30, 40, 50 or 100 contiguous amino acids truncated or deleted from the amino acid sequence of SEQ ID NO:2,4.

[0026] Fragments of the polypeptides of the invention may be employed for producing the corresponding full-length polypeptide by peptide synthesis; therefore, these fragments may be employed as intermediates for producing the full-length polypeptides of the invention.

[0027] Particularly preferred are variants in which several, 5-10, 1-5, 1-3, 1-2 or 1 amino acids are substituted, deleted, or added in any combination.

[0028] The polypeptides, or immunogenic fragments, of the invention may be in the form of the "mature" protein or may be a part of a larger protein such as a precursor or a fusion protein. It is often advantageous to include an additional amino acid sequence which contains secretory or leader sequences, pro-sequences, sequences which aid in purification such as multiple histidine residues, or an additional sequence for stability during recombinant production. Furthermore, addition of exogenous polypeptide or lipid tail or polynucleotide sequences to increase the immunogenic potential of the final molecule is also considered.

[0029] In one aspect, the invention relates to genetically engineered soluble fusion proteins comprising a polypeptide of the present invention, or a fragment thereof, and various portions of the constant regions of heavy or light chains of immunoglobulins of various subclasses (IgG, IgM, IgA, IgE). Preferred as an immunoglobulin is the constant part of the heavy chain of human IgG, particularly IgG1, where fusion takes place at the hinge region. In a particular embodiment,

the Fc part can be removed simply by incorporation of a cleavage sequence which can be cleaved with blood clotting factor Xa.

[0030] Furthermore, this invention relates to processes for the preparation of these fusion proteins by genetic engineering, and to the use thereof for drug screening, diagnosis and therapy. A further aspect of the invention also relates to polynucleotides encoding such fusion proteins. Examples of fusion protein technology can be found in International Patent Application Nos. WO94/29458 and WO94/22914.

[0031] The proteins may be chemically conjugated, or expressed as recombinant fusion proteins allowing increased levels to be produced in an expression system as compared to non-fused protein. The fusion partner may assist in providing T helper epitopes (immunological fusion partner), preferably T helper epitopes recognised by humans, or assist in expressing the protein (expression enhancer) at higher yields than the native recombinant protein. Preferably the fusion partner will be both an immunological fusion partner and expression enhancing partner.

[0032] Fusion partners include protein D from *Haemophilus influenzae* and the non-structural protein from influenzae virus, NS1 (hemagglutinin). Another fusion partner is the protein known as LytA. Preferably the C terminal portion of the molecule is used. LytA is derived from *Streptococcus pneumoniae* which synthesizes an N-acetyl-L-alanine amidase, amidase LytA, (coded by the *lytA* gene {Gene, 43 (1986) page 265-272}) an autolysin that specifically degrades certain bonds in the peptidoglycan backbone. The C-terminal domain of the LytA protein is responsible for the affinity to the choline or to some choline analogues such as DEAE. This property has been exploited for the development of *E. coli* C-LytA expressing plasmids useful for expression of fusion proteins. Purification of hybrid proteins containing the C-LytA fragment at its amino terminus has been described {Biotechnology: 10, (1992) page 795-798}. It is possible to use the repeat portion of the LytA molecule found in the C terminal end starting at residue 178, for example residues 188-305.

[0033] The present invention also includes variants of the aforementioned polypeptides, that is polypeptides that vary from the referents by conservative amino acid substitutions, whereby a residue is substituted by another with like characteristics. Typical such substitutions are among Ala, Val, Leu and Ile; among Ser and Thr; among the acidic residues Asp and Glu; among Asn and Gln; and among the basic residues Lys and Arg; or aromatic residues Phe and Tyr.

[0034] Polypeptides of the present invention can be prepared in any suitable manner. Such polypeptides include isolated naturally occurring polypeptides, recombinantly produced polypeptides, synthetically produced polypeptides, or polypeptides produced by a combination of these methods. Means for preparing such polypeptides are well understood in the art.

[0035] It is most preferred that a polypeptide of the invention is derived from *Neisseria meningitidis*, however, it may preferably be obtained from other organisms of the same taxonomic genus. A polypeptide of the invention may also be obtained, for example, from organisms of the same taxonomic family or order.

Polynucleotides

[0036] It is an object of the invention to provide polynucleotides that encode BASB006 polypeptides, particularly polynucleotides that encode the polypeptide herein designated BASB006.

[0037] In a particularly preferred embodiment of the invention the polynucleotide comprises a region encoding BASB006 polypeptides comprising a sequence set out in SEQ ID NO:1,3 which includes a full length gene, or a variant thereof.

[0038] The BASB006 polynucleotides provided in SEQ ID NO:1,3 are the BASB006 polynucleotides from *Neisseria meningitidis* strains ATCC13090 and H44/76.

[0039] As a further aspect of the invention there are provided isolated nucleic acid molecules encoding and/or expressing BASB006 polypeptides and polynucleotides, particularly *Neisseria meningitidis* BASB006 polypeptides and polynucleotides, including, for example, unprocessed RNAs, ribozyme RNAs, mRNAs, cDNAs, genomic DNAs, B- and Z-DNAs. Further embodiments of the invention include biologically, diagnostically, prophylactically, clinically or therapeutically useful polynucleotides and polypeptides, and variants thereof, and compositions comprising the same.

[0040] Another aspect of the invention relates to isolated polynucleotides, including at least one full length gene, that encodes a BASB006 polypeptide having a deduced amino acid sequence of SEQ ID NO:2,4 and polynucleotides closely related thereto and variants thereof.

[0041] In another particularly preferred embodiment of the invention there is a BASB006 polypeptide from *Neisseria meningitidis* comprising or consisting of an amino acid sequence of SEQ ID NO:2,4 or a variant thereof.

[0042] Using the information provided herein, such as a polynucleotide sequence set out in SEQ ID NO:1, 3 a polynucleotide of the invention encoding BASB006 polypeptide may be obtained using standard cloning and screening methods, such as those for cloning and sequencing chromosomal DNA fragments from bacteria using *Neisseria meningitidis* cells as starting material, followed by obtaining a full length clone. For example, to obtain a polynucleotide sequence of the invention, such as a polynucleotide sequence given in SEQ ID NO:1,3, typically a library of clones of chromosomal DNA of *Neisseria meningitidis* in *E. coli* or some other suitable host is probed with a radiolabeled oligonucleotide, preferably a 17-mer or longer, derived from a partial sequence. Clones carrying DNA identical to that of the probe can then be distinguished using stringent hybridization conditions. By sequencing the individual clones thus identified by hybridization with sequencing primers designed from the original polypeptide or polynucleotide sequence it is then possible to extend the polynucleotide sequence in both directions to determine a full length gene sequence. Conveniently, such sequencing is performed, for example, using denatured double stranded DNA prepared from a plasmid clone. Suitable techniques are described by Maniatis, T., Fritsch, E. F. and Sambrook et al., *MOLECULAR CLONING, A LABORATORY MANUAL*, 2nd Ed.; Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989). (see in particular Screening By Hybridization 1.90 and Sequencing Denatured Double-Stranded DNA Templates 13.70). Direct genomic DNA sequencing may also be performed to obtain a full length gene sequence. Illustrative of the invention, each polynucleotide set out in SEQ ID NO:1,3 was discovered in a DNA library derived from *Neisseria meningitidis*.

[0043] Moreover, each DNA sequence set out in SEQ ID NO:1,3 contains an open reading frame encoding a protein having about the number of amino acid residues set forth in SEQ ID NO:2, 4 with a deduced molecular weight that can be

calculated using amino acid residue molecular weight values well known to those skilled in the art.

[0044] The polynucleotide of SEQ ID NO:1, between the start codon at nucleotide number 1 and the stop codon which begins at nucleotide number 4363 of SEQ ID NO:1, encodes the polypeptide of SEQ ID NO:2.

The polynucleotide of SEQ ID NO:3, between the start codon at nucleotide number 1 and the stop codon which begins at nucleotide number 4372 of SEQ ID NO:3, encodes the polypeptide of SEQ ID NO:4.

[0045] In a further aspect, the present invention provides for an isolated polynucleotide comprising or consisting of:

(a) a polynucleotide sequence which has at least 85% identity, more preferably at least 90% identity, yet more preferably at least 95% identity, even more preferably at least 97-99% or exact identity to SEQ ID NO:1,3 over the entire length of SEQ ID NO:1,3 respectively; or

(b) a polynucleotide sequence encoding a polypeptide which has at least 85% identity, more preferably at least 90% identity, yet more preferably at least 95% identity, even more preferably at least 97-99% or 100% exact, to the amino acid sequence of SEQ ID NO:2, 4 over the entire length of SEQ ID NO:2, 4 respectively.

[0046] A polynucleotide encoding a polypeptide of the present invention, including homologs and orthologs from species other than *Neisseria meningitidis*, may be obtained by a process which comprises the steps of screening an appropriate library under stringent hybridization conditions (for example, using a temperature in the range of 45-65° C. and an SDS concentration from 0.1±1%) with a labeled or detectable probe consisting of or comprising the sequence of SEQ ID NO: 1, 3 or a fragment thereof; and isolating a full-length gene and/or genomic clones containing said polynucleotide sequence.

[0047] The invention provides a polynucleotide sequence identical over its entire length to a coding sequence (open reading frame) in SEQ ID NO: 1, 3. Also provided by the invention is a coding sequence for a mature polypeptide or a fragment thereof, by itself as well as a coding sequence for a mature polypeptide or a fragment in reading frame with another coding sequence, such as a sequence encoding a leader or secretory sequence, a pre-, or pro- or prepro-protein sequence. The polynucleotide of the invention may also contain at least one non-coding sequence, including for example, but not limited to at least one non-coding 5' and 3' sequence, such as the transcribed but non-translated sequences, termination signals (such as rho-dependent and rho-independent termination signals), ribosome binding sites, Kozak sequences, sequences that stabilize mRNA, introns, and polyadenylation signals. The polynucleotide sequence may also comprise additional coding sequence encoding additional amino acids. For example, a marker sequence that facilitates purification of the fused polypeptide can be encoded. In certain embodiments of the invention, the marker sequence is a hexa-histidine peptide, as provided in the pQE vector (Qiagen, Inc.) and described in Gentz et al., *Proc. Natl. Acad. Sci., USA* 86: 821-824 (1989), or an HA peptide tag (Wilson et al., *Cell* 37: 767 (1984), both of which may be useful in purifying polypeptide sequence fused to them. Polynucleotides of the invention also include, but are not limited to, polynucleotides comprising a structural gene and its naturally associated sequences that control gene expression.

[0048] The nucleotide sequence encoding BASB006 polypeptide of SEQ ID NO:2, 4 may be identical to the

polypeptide encoding sequence contained in nucleotides 1 to 4362 of SEQ ID NO:1, or the polypeptide encoding sequence contained in nucleotides 1 to 4371 of SEQ ID NO:3, respectively. Alternatively it may be a sequence, which as a result of the redundancy (degeneracy) of the genetic code, also encodes the polypeptide of SEQ ID NO:2, 4.

[0049] The term "polynucleotide encoding a polypeptide" as used herein encompasses polynucleotides that include a sequence encoding a polypeptide of the invention, particularly a bacterial polypeptide and more particularly a polypeptide of the *Neisseria meningitidis* BASB006 having an amino acid sequence set out in SEQ ID NO:2, 4. The term also encompasses polynucleotides that include a single continuous region or discontinuous regions encoding the polypeptide (for example, polynucleotides interrupted by integrated phage, an integrated insertion sequence, an integrated vector sequence, an integrated transposon sequence, or due to RNA editing or genomic DNA reorganization) together with additional regions, that also may contain coding and/or non-coding sequences.

[0050] The invention further relates to variants of the polynucleotides described herein that encode variants of a polypeptide having a deduced amino acid sequence of SEQ ID NO:2, 4. Fragments of polynucleotides of the invention may be used, for example, to synthesize full-length polynucleotides of the invention.

[0051] Further particularly preferred embodiments are polynucleotides encoding BASB006 variants, that have the amino acid sequence of BASB006 polypeptide of SEQ ID NO:2, 4 in which several, a few, 5 to 10, 1 to 5, 1 to 3, 2, 1 or no amino acid residues are substituted, modified, deleted and/or added, in any combination. Especially preferred among these are silent substitutions, additions and deletions, that do not alter the properties and activities of BASB006 polypeptide.

[0052] Further preferred embodiments of the invention are polynucleotides that are at least 85% identical over their entire length to a polynucleotide encoding BASB006 polypeptide having an amino acid sequence set out in SEQ ID NO:2, 4, and polynucleotides that are complementary to such polynucleotides. In this regard, polynucleotides at least 90% identical over their entire length to the same are particularly preferred, and among these particularly preferred polynucleotides, those with at least 95% are especially preferred. Furthermore, those with at least 97% are highly preferred among those with at least 95%, and among these those with at least 98% and at least 99% are particularly highly preferred, with at least 99% being the more preferred.

[0053] Preferred embodiments are polynucleotides encoding polypeptides that retain substantially the same biological function or activity as the mature polypeptide encoded by a DNA of SEQ ID NO:1, 3.

[0054] In accordance with certain preferred embodiments of this invention there are provided polynucleotides that hybridize, particularly under stringent conditions, to BASB006 polynucleotide sequences, such as those polynucleotides in SEQ ID NO:1, 3.

[0055] The invention further relates to polynucleotides that hybridize to the polynucleotide sequences provided herein. In this regard, the invention especially relates to polynucleotides that hybridize under stringent conditions to the polynucleotides described herein. As herein used, the terms "stringent conditions" and "stringent hybridization conditions" mean hybridization occurring only if there is at least 95% and

preferably at least 97% identity between the sequences. A specific example of stringent hybridization conditions is overnight incubation at 42° C. in a solution comprising: 50% formamide, 5×SSC (150 mM NaCl, 15 mM trisodium citrate), 50 mM sodium phosphate (pH7.6), 5×Denhardt's solution, 10% dextran sulfate, and 20 micrograms/ml of denatured, sheared salmon sperm DNA, followed by washing the hybridization support in 0.1×SSC at about 65° C. Hybridization and wash conditions are well known and exemplified in Sambrook, et al., *Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbor, N.Y., (1989), particularly Chapter 11 therein. Solution hybridization may also be used with the polynucleotide sequences provided by the invention.

[0056] The invention also provides a polynucleotide consisting of or comprising a polynucleotide sequence obtained by screening an appropriate library containing the complete gene for a polynucleotide sequence set forth in SEQ ID NO:1, 3 under stringent hybridization conditions with a probe having the sequence of said polynucleotide sequence set forth in SEQ ID NO:1, 3 or a fragment thereof; and isolating said polynucleotide sequence. Fragments useful for obtaining such a polynucleotide include, for example, probes and primers fully described elsewhere herein.

[0057] As discussed elsewhere herein regarding polynucleotide assays of the invention, for instance, the polynucleotides of the invention, may be used as a hybridization probe for RNA, cDNA and genomic DNA to isolate full-length cDNAs and genomic clones encoding BASB006 and to isolate cDNA and genomic clones of other genes that have a high identity, particularly high sequence identity, to the BASB006 gene. Such probes generally will comprise at least 15 nucleotide residues or base pairs. Preferably, such probes will have at least 30 nucleotide residues or base pairs and may have at least 50 nucleotide residues or base pairs. Particularly preferred probes will have at least 20 nucleotide residues or base pairs and will have less than 30 nucleotide residues or base pairs.

[0058] A coding region of a BASB006 gene may be isolated by screening using a DNA sequence provided in SEQ ID NO:1, 3 to synthesize an oligonucleotide probe. A labeled oligonucleotide having a sequence complementary to that of a gene of the invention is then used to screen a library of cDNA, genomic DNA or mRNA to determine which members of the library the probe hybridizes to.

[0059] There are several methods available and well known to those skilled in the art to obtain full-length DNAs, or extend short DNAs, for example those based on the method of Rapid Amplification of cDNA ends (RACE) (see, for example, Frohman, et al., *PNAS USA* 85: 8998-9002, 1988). Recent modifications of the technique, exemplified by the Marathon™ technology (Clontech Laboratories Inc.) for example, have significantly simplified the search for longer cDNAs. In the Marathon™ technology, cDNAs have been prepared from mRNA extracted from a chosen tissue and an 'adaptor' sequence ligated onto each end. Nucleic acid amplification (PCR) is then carried out to amplify the "missing" 5' end of the DNA using a combination of gene specific and adaptor specific oligonucleotide primers. The PCR reaction is then repeated using "nested" primers, that is, primers designed to anneal within the amplified product (typically an adaptor specific primer that anneals further 3' in the adaptor sequence and a gene specific primer that anneals further 5' in the selected gene sequence). The products of this reaction can

then be analyzed by DNA sequencing and a full-length DNA constructed either by joining the product directly to the existing DNA to give a complete sequence, or carrying out a separate full-length PCR using the new sequence information for the design of the 5' primer.

[0060] The polynucleotides and polypeptides of the invention may be employed, for example, as research reagents and materials for discovery of treatments of and diagnostics for diseases, particularly human diseases, as further discussed herein relating to polynucleotide assays.

[0061] The polynucleotides of the invention that are oligonucleotides derived from a sequence of SEQ ID NOS:1-4 may be used in the processes herein as described, but preferably for PCR, to determine whether or not the polynucleotides identified herein in whole or in part are transcribed in bacteria in infected tissue. It is recognized that such sequences will also have utility in diagnosis of the stage of infection and type of infection the pathogen has attained.

[0062] The invention also provides polynucleotides that encode a polypeptide that is the mature protein plus additional amino or carboxyl-terminal amino acids, or amino acids interior to the mature polypeptide (when the mature form has more than one polypeptide chain, for instance). Such sequences may play a role in processing of a protein from precursor to a mature form, may allow protein transport, may lengthen or shorten protein half-life or may facilitate manipulation of a protein for assay or production, among other things. As generally is the case in vivo, the additional amino acids may be processed away from the mature protein by cellular enzymes.

[0063] For each and every polynucleotide of the invention there is provided a polynucleotide complementary to it. It is preferred that these complementary polynucleotides are fully complementary to each polynucleotide with which they are complementary.

[0064] A precursor protein, having a mature form of the polypeptide fused to one or more prosequences may be an inactive form of the polypeptide. When prosequences are removed such inactive precursors generally are activated. Some or all of the prosequences may be removed before activation. Generally, such precursors are called proproteins.

[0065] In addition to the standard A, G, C, T/U representations for nucleotides, the term "N" may also be used in describing certain polynucleotides of the invention. "N" means that any of the four DNA or RNA nucleotides may appear at such a designated position in the DNA or RNA sequence, except it is preferred that N is not a nucleic acid that when taken in combination with adjacent nucleotide positions, when read in the correct reading frame, would have the effect of generating a premature termination codon in such reading frame.

[0066] In sum, a polynucleotide of the invention may encode a mature protein, a mature protein plus a leader sequence (which may be referred to as a preprotein), a precursor of a mature protein having one or more prosequences that are not the leader sequences of a preprotein, or a preprotein, which is a precursor to a proprotein, having a leader sequence and one or more prosequences, which generally are removed during processing steps that produce active and mature forms of the polypeptide.

[0067] In accordance with an aspect of the invention, there is provided the use of a polynucleotide of the invention for therapeutic or prophylactic purposes, in particular genetic immunization.

[0068] The use of a polynucleotide of the invention in genetic immunization will preferably employ a suitable delivery method such as direct injection of plasmid DNA into muscles (Wolff et al., *Hum Mol Genet* (1992) 1: 363, Manthorpe et al., *Hum. Gene Ther.* (1983) 4: 419), delivery of DNA complexed with specific protein carriers (Wu et al., *J Biol Chem.* (1989) 264: 16985), coprecipitation of DNA with calcium phosphate (Benvenisty & Reshef, *PNAS USA*, (1986) 83: 9551), encapsulation of DNA in various forms of liposomes (Kaneda et al., *Science* (1989) 243: 375), particle bombardment (Tang et al., *Nature* (1992) 356:152, Eisenbraun et al., *DNA Cell Biol* (1993) 12: 791) and in vivo infection using cloned retroviral vectors (Seeger et al., *PNAS USA* (1984) 81: 5849).

Vectors, Host Cells, Expression Systems

[0069] The invention also relates to vectors that comprise a polynucleotide or polynucleotides of the invention, host cells that are genetically engineered with vectors of the invention and the production of polypeptides of the invention by recombinant techniques. Cell-free translation systems can also be employed to produce such proteins using RNAs derived from the DNA constructs of the invention.

[0070] Recombinant polypeptides of the present invention may be prepared by processes well known in those skilled in the art from genetically engineered host cells comprising expression systems. Accordingly, in a further aspect, the present invention relates to expression systems that comprise a polynucleotide or polynucleotides of the present invention, to host cells which are genetically engineered with such expression systems, and to the production of polypeptides of the invention by recombinant techniques.

[0071] For recombinant production of the polypeptides of the invention, host cells can be genetically engineered to incorporate expression systems or portions thereof or polynucleotides of the invention. Introduction of a polynucleotide into the host cell can be effected by methods described in many standard laboratory manuals, such as Davis, et al., *BASIC METHODS IN MOLECULAR BIOLOGY*, (1986) and Sambrook, et al., *MOLECULAR CLONING: A LABORATORY MANUAL*, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989), such as, calcium phosphate transfection, DEAE-dextran mediated transfection, transvection, microinjection, cationic lipid-mediated transfection, electroporation, transduction, scrape loading, ballistic introduction and infection.

[0072] Representative examples of appropriate hosts include bacterial cells, such as cells of streptococci, staphylococci, enterococci, *E. coli*, *streptomyces*, cyanobacteria, *Bacillus subtilis*, *Moraxella catarrhalis*, *Haemophilus influenzae* and *Neisseria meningitidis*; fungal cells, such as cells of a yeast, *Kluveromyces*, *Saccharomyces*, a basidiomycete, *Candida albicans* and *Aspergillus*; insect cells such as cells of *Drosophila* S2 and *Spodoptera* Sf9; animal cells such as CHO, COS, HeLa, C127, 3T3, BHK, 293, CV-1 and Bowes melanoma cells; and plant cells, such as cells of a gymnosperm or angiosperm.

[0073] A great variety of expression systems can be used to produce the polypeptides of the invention. Such vectors include, among others, chromosomal-, episomal- and virus-derived vectors, for example, vectors derived from bacterial plasmids, from bacteriophage, from transposons, from yeast episomes, from insertion elements, from yeast chromosomal elements, from viruses such as baculoviruses, papova viruses,

such as SV40, vaccinia viruses, adenoviruses, fowl pox viruses, pseudorabies viruses, picornaviruses, retroviruses, and alphaviruses and vectors derived from combinations thereof, such as those derived from plasmid and bacteriophage genetic elements, such as cosmids and phagemids. The expression system constructs may contain control regions that regulate as well as engender expression. Generally, any system or vector suitable to maintain, propagate or express polynucleotides and/or to express a polypeptide in a host may be used for expression in this regard. The appropriate DNA sequence may be inserted into the expression system by any of a variety of well-known and routine techniques, such as, for example, those set forth in Sambrook et al., *MOLECULAR CLONING, A LABORATORY MANUAL*, (supra).

[0074] In recombinant expression systems in eukaryotes, for secretion of a translated protein into the lumen of the endoplasmic reticulum, into the periplasmic space or into the extracellular environment, appropriate secretion signals may be incorporated into the expressed polypeptide. These signals may be endogenous to the polypeptide or they may be heterologous signals.

[0075] Polypeptides of the present invention can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Most preferably, ion metal affinity chromatography (IMAC) is employed for purification. Well known techniques for refolding proteins may be employed to regenerate active conformation when the polypeptide is denatured during intracellular synthesis, isolation and or purification.

[0076] The expression system may also be a recombinant live microorganism, such as a virus or bacterium. The gene of interest can be inserted into the genome of a live recombinant virus or bacterium. Inoculation and in vivo infection with this live vector will lead to in vivo expression of the antigen and induction of immune responses. Viruses and bacteria used for this purpose are for instance: poxviruses (e.g.; vaccinia, fowlpox, canarypox), alphaviruses (Sindbis virus, Semliki Forest Virus, Venezuelan Equine Encephalitis Virus), adenoviruses, adeno-associated virus, picornaviruses (poliovirus, rhinovirus), herpesviruses (varicella zoster virus, etc), *Listeria*, *Salmonella*, *Shigella*, *Neisseria*, BCG. These viruses and bacteria can be virulent, or attenuated in various ways in order to obtain live vaccines. Such live vaccines also form part of the invention.

Diagnostic, Prognostic, Serotyping and Mutation Assays

[0077] This invention is also related to the use of BASB006 polynucleotides and polypeptides of the invention for use as diagnostic reagents. Detection of BASB006 polynucleotides and/or polypeptides in a eukaryote, particularly a mammal, and especially a human, will provide a diagnostic method for diagnosis of disease, staging of disease or response of an infectious organism to drugs. Eukaryotes, particularly mammals, and especially humans, particularly those infected or suspected to be infected with an organism comprising the BASB006 gene or protein, may be detected at the nucleic acid or amino acid level by a variety of well known techniques as well as by methods provided herein.

[0078] Polypeptides and polynucleotides for prognosis, diagnosis or other analysis may be obtained from a putatively

infected and/or infected individual's bodily materials. Polynucleotides from any of these sources, particularly DNA or RNA, may be used directly for detection or may be amplified enzymatically by using PCR or any other amplification technique prior to analysis. RNA, particularly mRNA, cDNA and genomic DNA may also be used in the same ways. Using amplification, characterization of the species and strain of infectious or resident organism present in an individual, may be made by an analysis of the genotype of a selected polynucleotide of the organism. Deletions and insertions can be detected by a change in size of the amplified product in comparison to a genotype of a reference sequence selected from a related organism, preferably a different species of the same genus or a different strain of the same species. Point mutations can be identified by hybridizing amplified DNA to labeled BASB006 polynucleotide sequences. Perfectly or significantly matched sequences can be distinguished from imperfectly or more significantly mismatched duplexes by DNase or RNase digestion, for DNA or RNA respectively, or by detecting differences in melting temperatures or renaturation kinetics. Polynucleotide sequence differences may also be detected by alterations in the electrophoretic mobility of polynucleotide fragments in gels as compared to a reference sequence. This may be carried out with or without denaturing agents. Polynucleotide differences may also be detected by direct DNA or RNA sequencing. See, for example, Myers et al., *Science*, 230: 1242 (1985). Sequence changes at specific locations also may be revealed by nuclease protection assays, such as RNase, V1 and S1 protection assay or a chemical cleavage method. See, for example, Cotton et al., *Proc. Natl. Acad. Sci., USA*, 85: 4397-4401 (1985).

[0079] In another embodiment, an array of oligonucleotides probes comprising BASB006 nucleotide sequence or fragments thereof can be constructed to conduct efficient screening of, for example, genetic mutations, serotype, taxonomic classification or identification. Array technology methods are well known and have general applicability and can be used to address a variety of questions in molecular genetics including gene expression, genetic linkage, and genetic variability (see, for example, Chee et al., *Science*, 274: 610 (1996)).

[0080] Thus in another aspect, the present invention relates to a diagnostic kit which comprises:

(a) a polynucleotide of the present invention, preferably the nucleotide sequence of SEQ ID NO:1, 3, or a fragment thereof;

(b) a nucleotide sequence complementary to that of (a);

(c) a polypeptide of the present invention, preferably the polypeptide of SEQ ID NO:2, 4 or a fragment thereof; or

(d) an antibody to a polypeptide of the present invention, preferably to the polypeptide of SEQ ID NO:2, 4.

[0081] It will be appreciated that in any such kit, (a), (b), (c) or (d) may comprise a substantial component. Such a kit will be of use in diagnosing a disease or susceptibility to a disease, among others.

[0082] This invention also relates to the use of polynucleotides of the present invention as diagnostic reagents. Detection of a mutated form of a polynucleotide of the invention, preferable, SEQ ID NO:1, 3, which is associated with a disease or pathogenicity will provide a diagnostic tool that can add to, or define, a diagnosis of a disease, a prognosis of a course of disease, a determination of a stage of disease, or a susceptibility to a disease, which results from under-expression, over-expression or altered expression of the polynucle-

otide. Organisms, particularly infectious organisms, carrying mutations in such polynucleotide may be detected at the polynucleotide level by a variety of techniques, such as those described elsewhere herein.

[0083] Cells from an organism carrying mutations or polymorphisms (allelic variations) in a polynucleotide and/or polypeptide of the invention may also be detected at the polynucleotide or polypeptide level by a variety of techniques, to allow for serotyping, for example. For example, RT-PCR can be used to detect mutations in the RNA. It is particularly preferred to use RT-PCR in conjunction with automated detection systems, such as, for example, GeneScan. RNA, cDNA or genomic DNA may also be used for the same purpose, PCR. As an example, PCR primers complementary to a polynucleotide encoding BASB006 polypeptide can be used to identify and analyze mutations.

[0084] The invention further provides primers with 1, 2, 3 or 4 nucleotides removed from the 5' and/or the 3' end. These primers may be used for, among other things, amplifying BASB006 DNA and/or RNA isolated from a sample derived from an individual, such as a bodily material. The primers may be used to amplify a polynucleotide isolated from an infected individual, such that the polynucleotide may then be subject to various techniques for elucidation of the polynucleotide sequence. In this way, mutations in the polynucleotide sequence may be detected and used to diagnose and/or prognose the infection or its stage or course, or to serotype and/or classify the infectious agent.

[0085] The invention further provides a process for diagnosing disease, preferably bacterial infections, more preferably infections caused by *Neisseria meningitidis*, comprising determining from a sample derived from an individual, such as a bodily material, an increased level of expression of polynucleotide having a sequence of SEQ ID NO:1, 3. Increased or decreased expression of a BASB006 polynucleotide can be measured using any one of the methods well known in the art for the quantitation of polynucleotides, such as, for example, amplification, PCR, RT-PCR, RNase protection, Northern blotting, spectrometry and other hybridization methods.

[0086] In addition, a diagnostic assay in accordance with the invention for detecting over-expression of BASB006 polypeptide compared to normal control tissue samples may be used to detect the presence of an infection, for example. Assay techniques that can be used to determine levels of a BASB006 polypeptide, in a sample derived from a host, such as a bodily material, are well-known to those of skill in the art. Such assay methods include radioimmunoassays, competitive-binding assays, Western Blot analysis, antibody sandwich assays, antibody detection and ELISA assays.

[0087] The polynucleotides of the invention may be used as components of polynucleotide arrays, preferably high density arrays or grids. These high density arrays are particularly useful for diagnostic and prognostic purposes. For example, a set of spots each comprising a different gene, and further comprising a polynucleotide or polynucleotides of the invention, may be used for probing, such as using hybridization or nucleic acid amplification, using a probe obtained or derived from a bodily sample, to determine the presence of a particular polynucleotide sequence or related sequence in an individual. Such a presence may indicate the presence of a pathogen, particularly *Neisseria meningitidis*, and may be useful in diagnosing and/or prognosing disease or a course of disease. A grid comprising a number of variants of the polynucleotide sequence of SEQ ID NO:1, 3 are preferred. Also preferred is

a grid comprising a number of variants of a polynucleotide sequence encoding the polypeptide sequence of SEQ ID NO:2, 4.

Antibodies

[0088] The polypeptides and polynucleotides of the invention or variants thereof, or cells expressing the same can be used as immunogens to produce antibodies immunospecific for such polypeptides or polynucleotides respectively.

[0089] In certain preferred embodiments of the invention there are provided antibodies against BASB006 polypeptides or polynucleotides.

[0090] Antibodies generated against the polypeptides or polynucleotides of the invention can be obtained by administering the polypeptides and/or polynucleotides of the invention, or epitope-bearing fragments of either or both, analogues of either or both, or cells expressing either or both, to an animal, preferably a nonhuman, using routine protocols. For preparation of monoclonal antibodies, any technique known in the art that provides antibodies produced by continuous cell line cultures can be used. Examples include various techniques, such as those in Kohler, G. and Milstein, C., *Nature* 256: 495-497 (1975); Kozbor et al., *Immunology Today* 4: 72 (1983); Cole et al., pg. 77-96 in *MONOCLONAL ANTIBODIES AND CANCER THERAPY*, Alan R. Liss, Inc. (1985).

[0091] Techniques for the production of single chain antibodies (U.S. Pat. No. 4,946,778) can be adapted to produce single chain antibodies to polypeptides or polynucleotides of this invention. Also, transgenic mice, or other organisms or animals, such as other mammals, may be used to express humanized antibodies immunospecific to the polypeptides or polynucleotides of the invention.

[0092] Alternatively, phage display technology may be utilized to select antibody genes with binding activities towards a polypeptide of the invention either from repertoires of PCR amplified v-genes of lymphocytes from humans screened for possessing anti-BASB006 or from naive libraries (McCafferty, et al., (1990), *Nature* 348, 552-554; Marks, et al., (1992) *Biotechnology* 10, 779-783). The affinity of these antibodies can also be improved by, for example, chain shuffling (Clackson et al., (1991) *Nature* 352: 628).

[0093] The above-described antibodies may be employed to isolate or to identify clones expressing the polypeptides or polynucleotides of the invention to purify the polypeptides or polynucleotides by, for example, affinity chromatography.

[0094] Thus, among others, antibodies against BASB006-polypeptide or BASB006-polynucleotide may be employed to treat infections, particularly bacterial infections.

[0095] Polypeptide variants include antigenically, epitopically or immunologically equivalent variants form a particular aspect of this invention.

[0096] Preferably, the antibody or variant thereof is modified to make it less immunogenic in the individual. For example, if the individual is human the antibody may most preferably be "humanized," where the complementarity determining region or regions of the hybridoma-derived antibody has been transplanted into a human monoclonal antibody, for example as described in Jones et al. (1986), *Nature* 321, 522-525 or Tempest et al., (1991) *Biotechnology* 9, 266-273.

Antagonists and Agonists—Assays and Molecules

[0097] Polypeptides and polynucleotides of the invention may also be used to assess the binding of small molecule

substrates and ligands in, for example, cells, cell-free preparations, chemical libraries, and natural product mixtures. These substrates and ligands may be natural substrates and ligands or may be structural or functional mimetics. See, e.g., Coligan et al., *Current Protocols in Immunology* 1(2): Chapter 5 (1991).

[0098] The screening methods may simply measure the binding of a candidate compound to the polypeptide or polynucleotide, or to cells or membranes bearing the polypeptide or polynucleotide, or a fusion protein of the polypeptide by means of a label directly or indirectly associated with the candidate compound. Alternatively, the screening method may involve competition with a labeled competitor. Further, these screening methods may test whether the candidate compound results in a signal generated by activation or inhibition of the polypeptide or polynucleotide, using detection systems appropriate to the cells comprising the polypeptide or polynucleotide. Inhibitors of activation are generally assayed in the presence of a known agonist and the effect on activation by the agonist by the presence of the candidate compound is observed. Constitutively active polypeptide and/or constitutively expressed polypeptides and polynucleotides may be employed in screening methods for inverse agonists or inhibitors, in the absence of an agonist or inhibitor, by testing whether the candidate compound results in inhibition of activation of the polypeptide or polynucleotide, as the case may be. Further, the screening methods may simply comprise the steps of mixing a candidate compound with a solution containing a polypeptide or polynucleotide of the present invention, to form a mixture, measuring BASB006 polypeptide and/or polynucleotide activity in the mixture, and comparing the BASB006 polypeptide and/or polynucleotide activity of the mixture to a standard. Fusion proteins, such as those made from Fc portion and BASB006 polypeptide, as hereinbefore described, can also be used for high-throughput screening assays to identify antagonists of the polypeptide of the present invention, as well as of phylogenetically and/or functionally related polypeptides (see D. Bennett et al., *J Mol Recognition*, 8:52-58 (1995); and K. Johanson et al., *J Biol Chem*, 270(16):9459-9471 (1995)).

[0099] The polynucleotides, polypeptides and antibodies that bind to and/or interact with a polypeptide of the present invention may also be used to configure screening methods for detecting the effect of added compounds on the production of mRNA and/or polypeptide in cells. For example, an ELISA assay may be constructed for measuring secreted or cell associated levels of polypeptide using monoclonal and polyclonal antibodies by standard methods known in the art. This can be used to discover agents which may inhibit or enhance the production of polypeptide (also called antagonist or agonist, respectively) from suitably manipulated cells or tissues.

[0100] The invention also provides a method of screening compounds to identify those which enhance (agonist) or block (antagonist) the action of BASB006 polypeptides or polynucleotides, particularly those compounds that are bacteriostatic and/or bactericidal. The method of screening may involve high-throughput techniques. For example, to screen for agonists or antagonists, a synthetic reaction mix, a cellular compartment, such as a membrane, cell envelope or cell wall, or a preparation of any thereof, comprising BASB006 polypeptide and a labeled substrate or ligand of such polypeptide is incubated in the absence or the presence of a candidate molecule that may be a BASB006 agonist or antagonist. The

ability of the candidate molecule to agonize or antagonize the BASB006 polypeptide is reflected in decreased binding of the labeled ligand or decreased production of product from such substrate. Molecules that bind gratuitously, i.e., without inducing the effects of BASB006 polypeptide are most likely to be good antagonists. Molecules that bind well and, as the case may be, increase the rate of product production from substrate, increase signal transduction, or increase chemical channel activity are agonists. Detection of the rate or level of, as the case may be, production of product from substrate, signal transduction, or chemical channel activity may be enhanced by using a reporter system. Reporter systems that may be useful in this regard include but are not limited to colorimetric, labeled substrate converted into product, a reporter gene that is responsive to changes in BASB006 polynucleotide or polypeptide activity, and binding assays known in the art.

[0101] Another example of an assay for BASB006 agonists is a competitive assay that combines BASB006 and a potential agonist with BASB006-binding molecules, recombinant BASB006 binding molecules, natural substrates or ligands, or substrate or ligand mimetics, under appropriate conditions for a competitive inhibition assay. BASB006 can be labeled, such as by radioactivity or a colorimetric compound, such that the number of BASB006 molecules bound to a binding molecule or converted to product can be determined accurately to assess the effectiveness of the potential antagonist.

[0102] Potential antagonists include, among others, small organic molecules, peptides, polypeptides and antibodies that bind to a polynucleotide and/or polypeptide of the invention and thereby inhibit or extinguish its activity or expression. Potential antagonists also may be small organic molecules, a peptide, a polypeptide such as a closely related protein or antibody that binds the same sites on a binding molecule, such as a binding molecule, without inducing BASB006-induced activities, thereby preventing the action or expression of BASB006 polypeptides and/or polynucleotides by excluding BASB006 polypeptides and/or polynucleotides from binding.

[0103] Potential antagonists include a small molecule that binds to and occupies the binding site of the polypeptide thereby preventing binding to cellular binding molecules, such that normal biological activity is prevented. Examples of small molecules include but are not limited to small organic molecules, peptides or peptide-like molecules. Other potential antagonists include antisense molecules (see Okano, *J. Neurochem.* 56: 560 (1991); *OLIGODEOXYNUCLEOTIDES AS ANTISENSE INHIBITORS OF GENE EXPRESSION*, CRC Press, Boca Raton, Fla. (1988), for a description of these molecules). Preferred potential antagonists include compounds related to and variants of BASB006.

[0104] In a further aspect, the present invention relates to genetically engineered soluble fusion proteins comprising a polypeptide of the present invention, or a fragment thereof, and various portions of the constant regions of heavy or light chains of immunoglobulins of various subclasses (IgG, IgM, IgA, IgE). Preferred as an immunoglobulin is the constant part of the heavy chain of human IgG, particularly IgG1, where fusion takes place at the hinge region. In a particular embodiment, the Fc part can be removed simply by incorporation of a cleavage sequence which can be cleaved with blood clotting factor Xa. Furthermore, this invention relates to processes for the preparation of these fusion proteins by genetic engineering, and to the use thereof for drug screening,

diagnosis and therapy. A further aspect of the invention also relates to polynucleotides encoding such fusion proteins. Examples of fusion protein technology can be found in International Patent Application Nos. WO94/29458 and WO94/22914.

[0105] Each of the polynucleotide sequences provided herein may be used in the discovery and development of antibacterial compounds. The encoded protein, upon expression, can be used as a target for the screening of antibacterial drugs. Additionally, the polynucleotide sequences encoding the amino terminal regions of the encoded protein or Shine-Delgarno or other translation facilitating sequences of the respective mRNA can be used to construct antisense sequences to control the expression of the coding sequence of interest.

[0106] The invention also provides the use of the polypeptide, polynucleotide, agonist or antagonist of the invention to interfere with the initial physical interaction between a pathogen or pathogens and a eukaryotic, preferably mammalian, host responsible for sequelae of infection. In particular, the molecules of the invention may be used: in the prevention of adhesion of bacteria, in particular gram positive and/or gram negative bacteria, to eukaryotic, preferably mammalian, extracellular matrix proteins on in-dwelling devices or to extracellular matrix proteins in wounds; to block bacterial adhesion between eukaryotic, preferably mammalian, extracellular matrix proteins and bacterial BASB006 proteins that mediate tissue damage and/or; to block the normal progression of pathogenesis in infections initiated other than by the implantation of in-dwelling devices or by other surgical techniques.

[0107] In accordance with yet another aspect of the invention, there are provided BASB006 agonists and antagonists, preferably bacteriostatic or bactericidal agonists and antagonists.

[0108] The antagonists and agonists of the invention may be employed, for instance, to prevent, inhibit and/or treat diseases.

[0109] In a further aspect, the present invention relates to mimotopes of the polypeptide of the invention. A mimotope is a peptide sequence, sufficiently similar to the native peptide (sequentially or structurally), which is capable of being recognised by antibodies which recognise the native peptide; or is capable of raising antibodies which recognise the native peptide when coupled to a suitable carrier.

[0110] Peptide mimotopes may be designed for a particular purpose by addition, deletion or substitution of elected amino acids. Thus, the peptides may be modified for the purposes of ease of conjugation to a protein carrier. For example, it may be desirable for some chemical conjugation methods to include a terminal cysteine. In addition it may be desirable for peptides conjugated to a protein carrier to include a hydrophobic terminus distal from the conjugated terminus of the peptide, such that the free unconjugated end of the peptide remains associated with the surface of the carrier protein. Thereby presenting the peptide in a conformation which most closely resembles that of the peptide as found in the context of the whole native molecule. For example, the peptides may be altered to have an N-terminal cysteine and a C-terminal hydrophobic amidated tail. Alternatively, the addition or substitution of a D-stereoisomer form of one or more of the amino acids may be performed to create a beneficial derivative, for example to enhance stability of the peptide.

[0111] Alternatively, peptide mimotopes may be identified using antibodies which are capable themselves of binding to the polypeptides of the present invention using techniques such as phage display technology (EP 0 552 267 B1). This technique, generates a large number of peptide sequences which mimic the structure of the native peptides and are, therefore, capable of binding to anti-native peptide antibodies, but may not necessarily themselves share significant sequence homology to the native polypeptide.

Vaccines

[0112] Another aspect of the invention relates to a method for inducing an immunological response in an individual, particularly a mammal, preferably humans, which comprises inoculating the individual with BASB006 polynucleotide and/or polypeptide, or a fragment or variant thereof, adequate to produce antibody and/or T cell immune response to protect said individual from infection, particularly bacterial infection and most particularly *Neisseria meningitidis* infection. Also provided are methods whereby such immunological response slows bacterial replication. Yet another aspect of the invention relates to a method of inducing immunological response in an individual which comprises delivering to such individual a nucleic acid vector, sequence or ribozyme to direct expression of BASB006 polynucleotide and/or polypeptide, or a fragment or a variant thereof, for expressing BASB006 polynucleotide and/or polypeptide, or a fragment or a variant thereof in vivo in order to induce an immunological response, such as, to produce antibody and/or T cell immune response, including, for example, cytokine-producing T cells or cytotoxic T cells, to protect said individual, preferably a human, from disease, whether that disease is already established within the individual or not. One example of administering the gene is by accelerating it into the desired cells as a coating on particles or otherwise. Such nucleic acid vector may comprise DNA, RNA, a ribozyme, a modified nucleic acid, a DNA/RNA hybrid, a DNA-protein complex or an RNA-protein complex.

[0113] A further aspect of the invention relates to an immunological composition that when introduced into an individual, preferably a human, capable of having induced within it an immunological response, induces an immunological response in such individual to a BASB006 polynucleotide and/or polypeptide encoded therefrom, wherein the composition comprises a recombinant BASB006 polynucleotide and/or polypeptide encoded therefrom and/or comprises DNA and/or RNA which encodes and expresses an antigen of said BASB006 polynucleotide, polypeptide encoded therefrom, or other polypeptide of the invention. The immunological response may be used therapeutically or prophylactically and may take the form of antibody immunity and/or cellular immunity, such as cellular immunity arising from CTL or CD4+ T cells.

[0114] A BASB006 polypeptide or a fragment thereof may be fused with co-protein or chemical moiety which may or may not by itself produce antibodies, but which is capable of stabilizing the first protein and producing a fused or modified protein which will have antigenic and/or immunogenic properties, and preferably protective properties. Thus fused recombinant protein, preferably further comprises an antigenic co-protein, such as lipoprotein D from *Haemophilus influenzae*, Glutathione-S-transferase (GST) or beta-galactosidase, or any other relatively large co-protein which solubilizes the protein and facilitates production and purification

thereof. Moreover, the co-protein may act as an adjuvant in the sense of providing a generalized stimulation of the immune system of the organism receiving the protein. The co-protein may be attached to either the amino- or carboxy-terminus of the first protein.

[0115] Provided by this invention are compositions, particularly vaccine compositions, and methods comprising the polypeptides and/or polynucleotides of the invention and immunostimulatory DNA sequences, such as those described in Sato, Y. et al. *Science* 273: 352 (1996).

[0116] Also, provided by this invention are methods using the described polynucleotide or particular fragments thereof, which have been shown to encode non-variable regions of bacterial cell surface proteins, in polynucleotide constructs used in such genetic immunization experiments in animal models of infection with *Neisseria meningitidis*. Such experiments will be particularly useful for identifying protein epitopes able to provoke a prophylactic or therapeutic immune response. It is believed that this approach will allow for the subsequent preparation of monoclonal antibodies of particular value, derived from the requisite organ of the animal successfully resisting or clearing infection, for the development of prophylactic agents or therapeutic treatments of bacterial infection, particularly *Neisseria meningitidis* infection, in mammals, particularly humans.

[0117] The invention also includes a vaccine formulation which comprises an immunogenic recombinant polypeptide and/or polynucleotide of the invention together with a suitable carrier, such as a pharmaceutically acceptable carrier. Since the polypeptides and polynucleotides may be broken down in the stomach, each is preferably administered parenterally, including, for example, administration that is subcutaneous, intramuscular, intravenous, or intradermal. Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteristatic compounds and solutes which render the formulation isotonic with the bodily fluid, preferably the blood, of the individual; and aqueous and non-aqueous sterile suspensions which may include suspending agents or thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example, sealed ampoules and vials and may be stored in a freeze-dried condition requiring only the addition of the sterile liquid carrier immediately prior to use.

[0118] The vaccine formulation of the invention may also include adjuvant systems for enhancing the immunogenicity of the formulation. Preferably the adjuvant system raises preferentially a TH1 type of response.

[0119] An immune response may be broadly distinguished into two extreme categories, being a humoral or cell mediated immune responses (traditionally characterised by antibody and cellular effector mechanisms of protection respectively). These categories of response have been termed TH1-type responses (cell-mediated response), and TH2-type immune responses (humoral response).

[0120] Extreme TH1-type immune responses may be characterised by the generation of antigen specific, haplotype restricted cytotoxic T lymphocytes, and natural killer cell responses. In mice TH1-type responses are often characterised by the generation of antibodies of the IgG2a subtype, whilst in the human these correspond to IgG1 type antibodies. TH2-type immune responses are characterised by the generation of a broad range of immunoglobulin isotypes including in mice IgG1, IgA, and IgM.

[0121] It can be considered that the driving force behind the development of these two types of immune responses are cytokines. High levels of TH1-type cytokines tend to favour the induction of cell mediated immune responses to the given antigen, whilst high levels of TH2-type cytokines tend to favour the induction of humoral immune responses to the antigen.

[0122] The distinction of TH1 and TH2-type immune responses is not absolute. In reality an individual will support an immune response which is described as being predominantly TH1 or predominantly TH2. However, it is often convenient to consider the families of cytokines in terms of that described in murine CD4 +ve T cell clones by Mosmann and Coffman (Mosmann, T. R. and Coffman, R. L. (1989) *TH1 and TH2 cells: different patterns of lymphokine secretion lead to different functional properties. Annual Review of Immunology*, 7, p 145-173). Traditionally, TH1-type responses are associated with the production of the INF- γ and IL-2 cytokines by T-lymphocytes. Other cytokines often directly associated with the induction of TH1-type immune responses are not produced by T-cells, such as IL-12. In contrast, TH2-type responses are associated with the secretion of IL-4, IL-6 and IL-13.

[0123] It is known that certain vaccine adjuvants are particularly suited to the stimulation of either TH1 or TH2-type cytokine responses. Traditionally the best indicators of the TH1:TH2 balance of the immune response after a vaccination or infection includes direct measurement of the production of TH1 or TH2 cytokines by T lymphocytes in vitro after restimulation with antigen, and/or the measurement of the IgG1:IgG2a ratio of antigen specific antibody responses.

[0124] Thus, a TH1-type adjuvant is one which preferentially stimulates isolated T-cell populations to produce high levels of TH1-type cytokines when re-stimulated with antigen in vitro, and promotes development of both CD8+ cytotoxic T lymphocytes and antigen specific immunoglobulin responses associated with TH1-type isotype.

[0125] Adjuvants which are capable of preferential stimulation of the TH1 cell response are described in International Patent Application No. WO 94/00153 and WO 95/17209.

[0126] 3 De-O-acylated monophosphoryl lipid A (3D-MPL) is one such adjuvant. This is known from GB 2220211 (Ribi). Chemically it is a mixture of 3 De-O-acylated monophosphoryl lipid A with 4, 5 or 6 acylated chains and is manufactured by Ribi Immunochem, Montana. A preferred form of 3 De-O-acylated monophosphoryl lipid A is disclosed in European Patent 0 689 454 B1 (SmithKline Beecham Biologicals SA).

[0127] Preferably, the particles of 3D-MPL are small enough to be sterile filtered through a 0.22 micron membrane (European Patent number 0 689 454). 3D-MPL will be present in the range of 10 μ g-100 μ g preferably 25-50 μ g per dose wherein the antigen will typically be present in a range 2-50 μ g per dose.

[0128] Another preferred adjuvant comprises QS21, an Hplc purified non-toxic fraction derived from the bark of *Quillaja Saponaria Molina*. Optionally this may be admixed with 3 De-O-acylated monophosphoryl lipid A (3D-MPL), optionally together with a carrier.

[0129] The method of production of QS21 is disclosed in U.S. Pat. No. 5,057,540.

[0130] Non-reactogenic adjuvant formulations containing QS21 have been described previously (WO 96/33739). Such formulations comprising QS21 and cholesterol have been

shown to be successful TH1 stimulating adjuvants when formulated together with an antigen.

[0131] Further adjuvants which are preferential stimulators of TH1 cell response include immunomodulatory oligonucleotides, for example unmethylated CpG sequences as disclosed in WO 96/02555.

[0132] Combinations of different TH1 stimulating adjuvants, such as those mentioned hereinabove, are also contemplated as providing an adjuvant which is a preferential stimulator of TH1 cell response. For example, QS21 can be formulated together with 3D-MPL. The ratio of QS21:3D-MPL will typically be in the order of 1:10 to 10:1; preferably 1:5 to 5:1 and often substantially 1:1. The preferred range for optimal synergy is 2.5:1 to 1:1 3D-MPL: QS21.

[0133] Preferably a carrier is also present in the vaccine composition according to the invention. The carrier may be an oil in water emulsion, or an aluminium salt, such as aluminium phosphate or aluminium hydroxide.

[0134] A preferred oil-in-water emulsion comprises a metabolisable oil, such as squalene, alpha tocopherol and Tween 80. In a particularly preferred aspect the antigens in the vaccine composition according to the invention are combined with QS21 and 3D-MPL in such an emulsion. Additionally the oil in water emulsion may contain span 85 and/or lecithin and/or tricaprylin.

[0135] Typically for human administration QS21 and 3D-MPL will be present in a vaccine in the range of 1-200 μ g, such as 10-100 μ g, preferably 10 μ g-50 μ g per dose. Typically the oil in water will comprise from 2 to 10% squalene, from 2 to 10% alpha tocopherol and from 0.3 to 3% tween 80. Preferably the ratio of squalene:alpha tocopherol is equal to or less than 1 as this provides a more stable emulsion. Span 85 may also be present at a level of 1%. In some cases it may be advantageous that the vaccines of the present invention will further contain a stabiliser.

[0136] Non-toxic oil in water emulsions preferably contain a non-toxic oil, e.g. squalene or squalene, an emulsifier, e.g. Tween 80, in an aqueous carrier. The aqueous carrier may be, for example, phosphate buffered saline.

[0137] A particularly potent adjuvant formulation involving QS21, 3D-MPL and tocopherol in an oil in water emulsion is described in WO 95/17210.

[0138] The present invention also provides a polyvalent vaccine composition comprising a vaccine formulation of the invention in combination with other antigens, in particular antigens useful for treating cancers, autoimmune diseases and related conditions. Such a polyvalent vaccine composition may include a TH-1 inducing adjuvant as hereinbefore described.

[0139] While the invention has been described with reference to certain BASB006 polypeptides and polynucleotides, it is to be understood that this covers fragments of the naturally occurring polypeptides and polynucleotides, and similar polypeptides and polynucleotides with additions, deletions or substitutions which do not substantially affect the immunogenic properties of the recombinant polypeptides or polynucleotides.

[0140] The antigen can also be delivered in the form of whole bacteria (dead or alive) or as subcellular fractions, these possibilities do include *N. meningitidis* itself.

Compositions, Kits and Administration

[0141] In a further aspect of the invention there are provided compositions comprising a BASB006 polynucleotide and/or a BASB006 polypeptide for administration to a cell or to a multicellular organism.

[0142] The invention also relates to compositions comprising a polynucleotide and/or a polypeptide discussed herein or their agonists or antagonists. The polypeptides and polynucleotides of the invention may be employed in combination with a non-sterile or sterile carrier or carriers for use with cells, tissues or organisms, such as a pharmaceutical carrier suitable for administration to an individual. Such compositions comprise, for instance, a media additive or a therapeutically effective amount of a polypeptide and/or polynucleotide of the invention and a pharmaceutically acceptable carrier or excipient. Such carriers may include, but are not limited to, saline, buffered saline, dextrose, water, glycerol, ethanol and combinations thereof. The formulation should suit the mode of administration. The invention further relates to diagnostic and pharmaceutical packs and kits comprising one or more containers filled with one or more of the ingredients of the aforementioned compositions of the invention.

[0143] Polypeptides, polynucleotides and other compounds of the invention may be employed alone or in conjunction with other compounds, such as therapeutic compounds.

[0144] The pharmaceutical compositions may be administered in any effective, convenient manner including, for instance, administration by topical, oral, anal, vaginal, intravenous, intraperitoneal, intramuscular, subcutaneous, intranasal or intradermal routes among others.

[0145] In therapy or as a prophylactic, the active agent may be administered to an individual as an injectable composition, for example as a sterile aqueous dispersion, preferably isotonic.

[0146] In a further aspect, the present invention provides for pharmaceutical compositions comprising a therapeutically effective amount of a polypeptide and/or polynucleotide, such as the soluble form of a polypeptide and/or polynucleotide of the present invention, agonist or antagonist peptide or small molecule compound, in combination with a pharmaceutically acceptable carrier or excipient. Such carriers include, but are not limited to, saline, buffered saline, dextrose, water, glycerol, ethanol, and combinations thereof. The invention further relates to pharmaceutical packs and kits comprising one or more containers filled with one or more of the ingredients of the aforementioned compositions of the invention. Polypeptides, polynucleotides and other compounds of the present invention may be employed alone or in conjunction with other compounds, such as therapeutic compounds.

[0147] The composition will be adapted to the route of administration, for instance by a systemic or an oral route. Preferred forms of systemic administration include injection, typically by intravenous injection. Other injection routes, such as subcutaneous, intramuscular, or intraperitoneal, can be used. Alternative means for systemic administration include transmucosal and transdermal administration using penetrants such as bile salts or fusidic acids or other detergents. In addition, if a polypeptide or other compounds of the present invention can be formulated in an enteric or an encapsulated formulation, oral administration may also be possible. Administration of these compounds may also be topical and/or localized, in the form of salves, pastes, gels, solutions, powders and the like.

[0148] For administration to mammals, and particularly humans, it is expected that the daily dosage level of the active agent will be from 0.01 mg/kg to 10 mg/kg, typically around 1 mg/kg. The physician in any event will determine the actual

dosage which will be most suitable for an individual and will vary with the age, weight and response of the particular individual. The above dosages are exemplary of the average case. There can, of course, be individual instances where higher or lower dosage ranges are merited, and such are within the scope of this invention.

[0149] The dosage range required depends on the choice of peptide, the route of administration, the nature of the formulation, the nature of the subject's condition, and the judgment of the attending practitioner. Suitable dosages, however, are in the range of 0.1-100 $\mu\text{g}/\text{kg}$ of subject.

[0150] A vaccine composition is conveniently in injectable form. Conventional adjuvants may be employed to enhance the immune response. A suitable unit dose for vaccination is 0.5-5 microgram/kg of antigen, and such dose is preferably administered 1-3 times and with an interval of 1-3 weeks. With the indicated dose range, no adverse toxicological effects will be observed with the compounds of the invention which would preclude their administration to suitable individuals.

[0151] Wide variations in the needed dosage, however, are to be expected in view of the variety of compounds available and the differing efficiencies of various routes of administration. For example, oral administration would be expected to require higher dosages than administration by intravenous injection. Variations in these dosage levels can be adjusted using standard empirical routines for optimization, as is well understood in the art.

Sequence Databases, Sequences in a Tangible Medium, and Algorithms

[0152] Polynucleotide and polypeptide sequences form a valuable information resource with which to determine their 2- and 3-dimensional structures as well as to identify further sequences of similar homology. These approaches are most easily facilitated by storing the sequence in a computer readable medium and then using the stored data in a known macromolecular structure program or to search a sequence database using well known searching tools, such as the GCG program package.

[0153] Also provided by the invention are methods for the analysis of character sequences or strings, particularly genetic sequences or encoded protein sequences. Preferred methods of sequence analysis include, for example, methods of sequence homology analysis, such as identity and similarity analysis, DNA, RNA and protein structure analysis, sequence assembly, cladistic analysis, sequence motif analysis, open reading frame determination, nucleic acid base calling, codon usage analysis, nucleic acid base trimming, and sequencing chromatogram peak analysis.

[0154] A computer based method is provided for performing homology identification. This method comprises the steps of: providing a first polynucleotide sequence comprising the sequence of a polynucleotide of the invention in a computer readable medium; and comparing said first polynucleotide sequence to at least one second polynucleotide or polypeptide sequence to identify homology.

[0155] A computer based method is also provided for performing homology identification, said method comprising the steps of: providing a first polypeptide sequence comprising the sequence of a polypeptide of the invention in a computer readable medium; and comparing said first polypeptide sequence to at least one second polynucleotide or polypeptide sequence to identify homology.

[0156] All publications and references, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference in their entirety as if each individual publication or reference were specifically and individually indicated to be incorporated by reference herein as being fully set forth. Any patent application to which this application claims priority is also incorporated by reference herein in its entirety in the manner described above for publications and references.

DEFINITIONS

[0157] “Identity,” as known in the art, is a relationship between two or more polypeptide sequences or two or more polynucleotide sequences, as the case may be, as determined by comparing the sequences. In the art, “identity” also means the degree of sequence relatedness between polypeptide or polynucleotide sequences, as the case may be, as determined by the match between strings of such sequences. “Identity” can be readily calculated by known methods, including but not limited to those described in (*Computational Molecular Biology*, Lesk, A. M., ed., Oxford University Press, New York, 1988; *Biocomputing: Informatics and Genome Projects*, Smith, D. W., ed., Academic Press, New York, 1993; *Computer Analysis of Sequence Data*, Part I, Griffin, A. M., and Griffin, H. G., eds., Humana Press, New Jersey, 1994; *Sequence Analysis in Molecular Biology*, von Heine, G., Academic Press, 1987; and *Sequence Analysis Primer*, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991; and Carillo, H., and Lipman, D., *SIAM J. Applied Math.*, 48: 1073 (1988). Methods to determine identity are designed to give the largest match between the sequences tested. Moreover, methods to determine identity are codified in publicly available computer programs. Computer program methods to determine identity between two sequences include, but are not limited to, the GAP program in the GCG program package (Devereux, J., et al., *Nucleic Acids Research* 12(1): 387 (1984)), BLASTP, BLASTN (Altschul, S. F. et al., *J. Molec. Biol.* 215: 403-410 (1990), and FASTA (Pearson and Lipman Proc. Natl. Acad. Sci. USA 85: 2444-2448 (1988)). The BLAST family of programs is publicly available from NCBI and other sources (*BLAST Manual*, Altschul, S., et al., NCBI NLM NIH Bethesda, Md. 20894; Altschul, S., et al., *J. Mol. Biol.* 215: 403-410 (1990)). The well known Smith Waterman algorithm may also be used to determine identity.

[0158] Parameters for polypeptide sequence comparison include the following:

Algorithm: Needleman and Wunsch, *J. Mol. Biol.* 48: 443-453 (1970)

Comparison matrix: BLOSSUM62 from Henikoff and Henikoff,

Proc. Natl. Acad. Sci. USA. 89:10915-10919 (1992)

Gap Penalty: 8

Gap Length Penalty: 2

[0159] A program useful with these parameters is publicly available as the “gap” program from Genetics Computer Group, Madison Wis. The aforementioned parameters are the default parameters for peptide comparisons (along with no penalty for end gaps).

[0160] Parameters for polynucleotide comparison include the following:

Algorithm: Needleman and Wunsch, *J. Mol Biol.* 48: 443-453 (1970)

[0161] Comparison matrix: matches=+10, mismatch=0

Gap Penalty: 50

Gap Length Penalty: 3

[0162] Available as: The “gap” program from Genetics Computer Group, Madison Wis. These are the default parameters for nucleic acid comparisons.

[0163] A preferred meaning for “identity” for polynucleotides and polypeptides, as the case may be, are provided in (1) and (2) below.

[0164] (1) Polynucleotide embodiments further include an isolated polynucleotide comprising a polynucleotide sequence having at least a 50, 60, 70, 80, 85, 90, 95, 97 or 100% identity to the reference sequence of SEQ ID NO:1, wherein said polynucleotide sequence may be identical to the reference sequence of SEQ ID NO:1 or may include up to a certain integer number of nucleotide alterations as compared to the reference sequence, wherein said alterations are selected from the group consisting of at least one nucleotide deletion, substitution, including transition and transversion, or insertion, and wherein said alterations may occur at the 5' or 3' terminal positions of the reference nucleotide sequence or anywhere between those terminal positions, interspersed either individually among the nucleotides in the reference sequence or in one or more contiguous groups within the reference sequence, and wherein said number of nucleotide alterations is determined by multiplying the total number of nucleotides in SEQ ID NO:1 by the integer defining the percent identity divided by 100 and then subtracting that product from said total number of nucleotides in SEQ ID NO:1, or:

$$n_n \leq x_n - (x_n \cdot y),$$

wherein n_n is the number of nucleotide alterations, x_n is the total number of nucleotides in SEQ ID NO:1, y is 0.50 for 50%, 0.60 for 60%, 0.70 for 70%, 0.80 for 80%, 0.85 for 85%, 0.90 for 90%, 0.95 for 95%, 0.97 for 97% or 1.00 for 100%, and \cdot is the symbol for the multiplication operator, and wherein any non-integer product of x_n and y is rounded down to the nearest integer prior to subtracting it from x_n . Alterations of a polynucleotide sequence encoding the polypeptide of SEQ ID NO:2 may create nonsense, missense or frameshift mutations in this coding sequence and thereby alter the polypeptide encoded by the polynucleotide following such alterations.

[0165] By way of example, a polynucleotide sequence of the present invention may be identical to the reference sequence of SEQ ID NO:1, that is it may be 100% identical, or it may include up to a certain integer number of nucleic acid alterations as compared to the reference sequence such that the percent identity is less than 100% identity. Such alterations are selected from the group consisting of at least one nucleic acid deletion, substitution, including transition and transversion, or insertion, and wherein said alterations may occur at the 5' or 3' terminal positions of the reference polynucleotide sequence or anywhere between those terminal positions, interspersed either individually among the nucleic acids in the reference sequence or in one or more contiguous

groups within the reference sequence. The number of nucleic acid alterations for a given percent identity is determined by multiplying the total number of nucleic acids in SEQ ID NO:1 by the integer defining the percent identity divided by 100 and then subtracting that product from said total number of nucleic acids in SEQ ID NO:1, or:

$$n_n \leq x_n - (x_n \cdot y),$$

wherein n_n is the number of nucleic acid alterations, x_n is the total number of nucleic acids in SEQ ID NO:1, y is, for instance 0.70 for 70%, 0.80 for 80%, 0.85 for 85% etc., \cdot is the symbol for the multiplication operator, and wherein any non-integer product of x_n and y is rounded down to the nearest integer prior to subtracting it from x_n .

[0166] (2) Polypeptide embodiments further include an isolated polypeptide comprising a polypeptide having at least a 50, 60, 70, 80, 85, 90, 95, 97 or 100% identity to a polypeptide reference sequence of SEQ ID NO:2, wherein said polypeptide sequence may be identical to the reference sequence of SEQ ID NO:2 or may include up to a certain integer number of amino acid alterations as compared to the reference sequence, wherein said alterations are selected from the group consisting of at least one amino acid deletion, substitution, including conservative and non-conservative substitution, or insertion, and wherein said alterations may occur at the amino- or carboxy-terminal positions of the reference polypeptide sequence or anywhere between those terminal positions, interspersed either individually among the amino acids in the reference sequence or in one or more contiguous groups within the reference sequence, and wherein said number of amino acid alterations is determined by multiplying the total number of amino acids in SEQ ID NO:2 by the integer defining the percent identity divided by 100 and then subtracting that product from said total number of amino acids in SEQ ID NO:2, or:

$$n_a \leq x_a - (x_a \cdot y),$$

wherein n_a is the number of amino acid alterations, x_a is the total number of amino acids in SEQ ID NO:2, y is 0.50 for 50%, 0.60 for 60%, 0.70 for 70%, 0.80 for 80%, 0.85 for 85%, 0.90 for 90%, 0.95 for 95%, 0.97 for 97% or 1.00 for 100%, and \cdot is the symbol for the multiplication operator, and wherein any non-integer product of x_a and y is rounded down to the nearest integer prior to subtracting it from x_a .

[0167] By way of example, a polypeptide sequence of the present invention may be identical to the reference sequence of SEQ ID NO:2, that is it may be 100% identical, or it may include up to a certain integer number of amino acid alterations as compared to the reference sequence such that the percent identity is less than 100% identity. Such alterations are selected from the group consisting of at least one amino acid deletion, substitution, including conservative and non-conservative substitution, or insertion, and wherein said alterations may occur at the amino- or carboxy-terminal positions of the reference polypeptide sequence or anywhere between those terminal positions, interspersed either individually among the amino acids in the reference sequence or in one or more contiguous groups within the reference sequence. The number of amino acid alterations for a given % identity is determined by multiplying the total number of amino acids in SEQ ID NO:2 by the integer defining the percent identity divided by 100 and then subtracting that product from said total number of amino acids in SEQ ID NO:2, or:

$$n_a \leq x_a - (x_a \cdot y),$$

wherein n_a is the number of amino acid alterations, x_a is the total number of amino acids in SEQ ID NO:2, y is, for instance 0.70 for 70%, 0.80 for 80%, 0.85 for 85% etc., and \cdot is the symbol for the multiplication operator, and wherein any non-integer product of x_a and y is rounded down to the nearest integer prior to subtracting it from x_a .

[0168] "Individual(s)," when used herein with reference to an organism, means a multicellular eukaryote, including, but not limited to a metazoan, a mammal, an ovid, a bovid, a simian, a primate, and a human.

[0169] "Isolated" means altered "by the hand of man" from its natural state, i.e., if it occurs in nature, it has been changed or removed from its original environment, or both. For example, a polynucleotide or a polypeptide naturally present in a living organism is not "isolated," but the same polynucleotide or polypeptide separated from the coexisting materials of its natural state is "isolated", as the term is employed herein. Moreover, a polynucleotide or polypeptide that is introduced into an organism by transformation, genetic manipulation or by any other recombinant method is "isolated" even if it is still present in said organism, which organism may be living or non-living.

[0170] "Polynucleotide(s)" generally refers to any polyribonucleotide or polydeoxyribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA including single and double-stranded regions.

[0171] "Variant" refers to a polynucleotide or polypeptide that differs from a reference polynucleotide or polypeptide, but retains essential properties. A typical variant of a polynucleotide differs in nucleotide sequence from another, reference polynucleotide. Changes in the nucleotide sequence of the variant may or may not alter the amino acid sequence of a polypeptide encoded by the reference polynucleotide. Nucleotide changes may result in amino acid substitutions, additions, deletions, fusions and truncations in the polypeptide encoded by the reference sequence, as discussed below. A typical variant of a polypeptide differs in amino acid sequence from another, reference polypeptide. Generally, differences are limited so that the sequences of the reference polypeptide and the variant are closely similar overall and, in many regions, identical. A variant and reference polypeptide may differ in amino acid sequence by one or more substitutions, additions, deletions in any combination. A substituted or inserted amino acid residue may or may not be one encoded by the genetic code. A variant of a polynucleotide or polypeptide may be a naturally occurring such as an allelic variant, or it may be a variant that is not known to occur naturally. Non-naturally occurring variants of polynucleotides and polypeptides may be made by mutagenesis techniques or by direct synthesis.

[0172] "Disease(s)" means any disease caused by or related to infection by a bacteria, including, for example, upper respiratory tract infection, invasive bacterial diseases, such as bacteremia and meningitis.

EXAMPLES

[0173] The examples below are carried out using standard techniques, which are well known and routine to those of skill in the art, except where otherwise described in detail. The examples are illustrative, but do not limit the invention.

Example 1

Discovery and Confirmatory DNA Sequencing of the BASB006 Gene from Two *N. meningitidis* Strains

[0174] A: BASB006 in *N. meningitidis* Serogroup B Strain ATCC13090.

[0175] The BASB006 gene disclosed in SEQ ID NO:1 was first discovered in the Incyte PathoSeq database containing unfinished genomic DNA sequences of the *N. meningitidis* strain ATCC13090. The translation of the BASB006 polynucleotide sequence, shown in SEQ ID NO:2, showed significant similarity (56% identity in a 1455 amino acids overlap) to the Hap protein of *Haemophilus influenzae*, a polypeptide reported previously to function as an adhesin. The sequence of the BASB006 gene was confirmed experimentally. For this purpose, genomic DNA was extracted from 10¹⁰ cells of the *N. meningitidis* cells (strain ATCC 13090) using the QIAGEN genomic DNA extraction kit (Qiagen GmbH), and 1 µg of this material was submitted to Polymerase Chain Reaction DNA amplification using primers Hap01 (5'-GGG GGC TAG CAA AAC AAC CGA CAA ACG GAC AAC C-3') [SEQ ID NO:5] and Hap02 (5'-GGG GAA GCT TCC AGC GGT AGC GGT AGC CTA ATT TGA TGC C-3') [SEQ ID NO:6]. This PCR product was gel-purified and subjected to DNA sequencing using the Big Dye Cycle Sequencing kit (Perkin-Elmer) and an ABI 373A/PRISM DNA sequencer. DNA sequencing was performed on both strands with a redundancy of 2 and the full-length sequence was assembled using the SeqMan program from the DNASTAR Lasergene software package. The resulting DNA sequence turned out to be 100% identical to SEQ ID NO:1.

B: BASB006 in *N. meningitidis* Serogroup B Strain H44/76. **[0176]** The sequence of the BASB006 gene was also determined in another *N. meningitidis* serogroup B strain, the strain H44/76. For this purpose, genomic DNA was extracted from the *N. meningitidis* strain H44/76 using the experimental conditions presented in Example 1. This material (1 µg) was then submitted to Polymerase Chain Reaction DNA amplification using primers Hap01 and Hap02 specific for the BASB006 gene. A 4389 bp DNA fragment was obtained, digested by the NheI/HindIII restriction endonucleases and inserted into the corresponding sites of the pET-24b cloning/expression vector (Novagen) using standard molecular biology techniques (Molecular Cloning, a Laboratory Manual, Second Edition, Eds: Sambrook, Fritsch & Maniatis, Cold Spring Harbor press 1989). Recombinant pET-24b/BASB006 was then submitted to DNA sequencing using the Big Dyes kit (Applied biosystems) and analyzed on a ABI 373A DNA sequencer in the conditions described by the supplier. As a result, the polynucleotide and deduced polypeptide sequences, referred to as SEQ ID NO:3 and SEQ ID NO:4 respectively, were obtained. Using the PILEUP program from the GCG package, an alignment of the polynucleotide sequences of SEQ ID NO:1 and 3 was performed, and is displayed in FIG. 1; their level of identity amounts to 97.8%, as determined by the GAP program. Using the same PILEUP program, an alignment of the polypeptide sequences of SEQ ID NO:2 and 4 was performed, and is displayed in FIG. 2; their level of identity amounts to 97.0%, as determined by the GAP program. Taken together, these data indicate strong sequence conservation of the BASB006 gene among the two *N. meningitidis* serogroup B strains.

Example 2

Expression and Purification of Recombinant BASB006 Protein in *Escherichia coli*

[0177] The construction of the pET-24b/BASB006 cloning/expression vector was described in Example 1B. This

vector harbours the BASB006 gene isolated from the strain H44/76 in fusion with a stretch of 6 histidine residues, placed under the control of the strong bacteriophage T7 gene 10 promoter. For expression study, this vector was introduced into the *Escherichia coli* strain BL21 DE3 (Novagen), in which, the gene for the T7 polymerase is placed under the control of the isopropyl-beta-D thiogalactoside (IPTG)-regulatable lac promoter. Liquid cultures (100 ml) of the BL21 DE-3 [pET-24b/BASB006] *E. coli* recombinant strain were grown at 37° C. under agitation until the optical density at 600 nm (OD600) reached 0.6. At that time-point, IPTG was added at a final concentration of 1 mM and the culture was grown for 4 additional hours. The culture was then centrifuged at 10,000 rpm and the pellet was frozen at -20° C. for at least 10 hours. After thawing, the pellet was resuspended during 30 min at 25° C. in buffer A (6M guanidine hydrochloride, 0.1M NaH2PO4, 0.01M Tris, pH 8.0), passed three-times through a needle and clarified by centrifugation (20000 rpm, 15 min). The sample was then loaded at a flow-rate of 1 ml/min on a Ni2+-loaded Hitrap column (Pharmacia Biotech). After passage of the flowthrough, the column was washed successively with 40 ml of buffer B (8M Urea, 0.1M NaH2PO4, 0.01M Tris, pH 8.0), 40 ml of buffer C (8M Urea, 0.1M NaH2PO4, 0.01M Tris, pH 6.3). The recombinant protein BASB006/His6 was then eluted from the column with 30 ml of buffer C (8M Urea, 0.1M NaH2PO4, 0.01M Tris, pH 6.3) containing 500 mM of imidazole and 3 ml-size fractions were collected. In FIG. 3, substantially purified proteins were separated on a 4-20% gradient polyacrylamide gel under PAGE-SDS conditions and stained with Coomassie Blue 8250. The sample loaded on the gel corresponded to protein fractions enriched (more than 80%) in BASB006 (lane 1 and 2) and a molecular weight marker (MW). As shown in FIG. 3, a highly enriched (Purity estimated to more than 90% pure in coomassie staining) BASB006/His6 protein, migrating at 170 kDa (estimated relative molecular mass), was eluted from the column. This polypeptide was reactive against a mouse monoclonal antibody raised against the 5-histidine motif. Taken together, these data indicate that the BASB006 gene can be expressed and purified under a recombinant form (BASB006/His6) in *E. coli*.

Example 3

Immunization of Mice with Recombinant BASB006 and Recognition of the BASB006 Polypeptide on Different *N. meningitidis* Serogroup B Strains by Western Blotting

[0178] Partially purified recombinant BASB006 expressed in *E. coli* has been injected three times in BALB/C mice on days 0, 14 and 28 (10 animals/group). Animals were injected by the subcutaneous route with 5 µg of antigen r formulated in SB62 emulsion containing 5 µg MPL and 1 µg QS21 per dose. Mice were bled on days 29 (15 days Post II) and 35 (6 days Post III) in order to detect specific anti-BASB006 antibodies. Specific anti-BASB006 antibodies were measured on pooled sera (from 10 mice/group) by western-blotting on six different *Neisseria meningitidis* serogroup B strains (FIG. 4).

[0179] The six different *Neisseria meningitidis* B strains are: H44/76 (B:15:P1.7, 16, lineage ET-5), M97 250987 (B:4: P1.15), BZ10 (B:2b:P1.2, lineage A4), BZ198 (B:NT*-, lineage 3), and EG328 (B:NT*, lineage ST-18), and on partially purified recombinant BASB006 protein (mixed with two other candidate antigens). (*:NT: Not Typed).

[0180] Briefly, 15 μ l ($>10^8$ cells/lane) of each sample treated with sample buffer (10 min at 95° C.) are put into a SDS-PAGE gradient gel (Tris-glycine 4-20%, Novex, code n° EC60252). Electrophoretic migration occurs at 125 volts for 90 min. Afterwards, proteins are transferred to a nitrocellulose sheet (0.45 μ m, Bio-rad code n° 162-0114) at 100 volts for 1 hour using a Bio-rad Trans-blot system (code n° 170-3930). The filter was blocked with PBS—0.05% Tween 20 overnight at room temperature, before incubation with the mice sera containing the anti-BASB006 antibodies. These sera are diluted 100 times in PBS—0.05% Tween 20, and incubated on the nitrocellulose sheet for two hours at room temperature with gentle shaking, using a mini-blotter system (Miniprotean, Bio-rad code n° 170-4017). After three repeated washing steps in PBS—0.05% Tween 20 for 5 min., the nitrocellulose sheet is incubated at room temperature for 1 hour under gentle shaking with the appropriate conjugate (biotinylated anti-mouse Ig antibodies from sheep, Amersham code n° RPN1001) diluted at 1/500 in the same washing buffer. The membrane is washed three times as previously, and incubated for 30 min. with agitation using the streptavidin-peroxidase complex (Amersham code n° 1051) diluted at 1/1000 in the washing buffer. After the last three repeated washing steps, the revelation occurs during the 20 min incubation time in a 50 ml solution containing 30 mg 4-chloro-1-naphthol (Sigma), 10 ml methanol, 40 ml PBS, and 30 μ l of H₂O₂. The staining is stopped while washing the membrane several times in distilled water.

[0181] FIG. 4 shows recognition of the native BASB006 protein from several *Neisseria meningitidis* serogroup B strains by sera from immunized mice. Results illustrated in FIG. 4 show that all strains tested present a band around 95-100 kD (see arrow), which is probably the extracellular part of the BASB006 protein (after cleavage of the intact molecule into two pieces, which is known to occur in the *H. influenzae* Hap protein). This means that the BASB006 protein is probably expressed in most of the *Neisseria meningitidis* serogroup B strains. All other bands could be antibodies directed against degradation products, or against, cross-reacting antigens between *E. coli* and *Neisseria meningitidis* B strains, as the preparation used for immunization still contained *E. coli* contaminants.

Example 4

Presence of Anti-BASB006 Antibodies in Sera from Human Convalescent Patients

[0182] In this test, human convalescent sera were tested by western-blotting for recognition of the purified recombinant BASB006 protein.

[0183] 5 μ g of partially purified BASB006 protein mixed with two other *Neisseria meningitidis* serogroup B proteins are put into a SDS-PAGE gradient gel (4-20%, Novex, code n° EC60252) for electrophoretic migration. Proteins are transferred to nitrocellulose sheet (0.45 μ m, Bio-rad code n° 162-0114) at 100 volts for 1 hour using a Bio-rad Trans-blot system (code n° 170-3930). Afterwards, the filter is blocked with PBS—0.05% Tween 20 overnight at room temperature, before incubation with the human sera. These sera are diluted 100 times in PBS—0.05% Tween 20, and incubated on the

nitrocellulose sheet for two hours at room temperature with gentle shaking, using a mini-blotter system (Miniprotean, Bio-rad code n° 170-4017). After three repeated washing steps in PBS—0.05% Tween 20 for 5 min., the nitrocellulose sheet is incubated at room temperature for 1 hour under gentle shaking with the appropriate conjugate (biotinylated anti-human Ig antibodies, from sheep, Amersham code n° RPN1003) diluted at 1/500 in the same washing buffer. The membrane is washed three times as previously, and incubated for 30 min with agitation using the streptavidin-peroxidase complex (Amersham code n° 1051) diluted at 1/1000 in the washing buffer. After the last three repeated washing steps, the revelation occurs during the 20 min incubation time in a 50 ml solution containing 30 mg 4-chloro-1-naphthol (Sigma), 10 ml methanol, 40 ml of ultra-pure water, and 30 μ l of H₂O₂. The staining is stopped while washing the membrane several times in distilled water.

[0184] Results illustrated in FIG. 5 (Part B) show that all convalescent sera react against the intact BASB006 protein at around 160 kD, while 3 out of 7 convalescent sera are reacting against the possible processed BASB006 protein (+/-95-100 kD). The BASB006 bands are clearly visible at these two molecular weights (95-100 and 160 kD). In part A of the western-blot, it can be seen that mice sera (mixture of specific antibodies against three different Ag candidates) recognize the intact recombinant BASB006 protein at the same molecular weight, while at the lower MW, it is more difficult to discriminate which of the two bands around 95 kD is related to the processed BASB006 protein

Deposited Materials

[0185] A deposit containing a *Neisseria meningitidis* Serogroup B strain has been deposited with the American Type Culture Collection (herein "ATCC") on Jun. 22, 1997 and assigned deposit number 13090. The deposit was described as *Neisseria meningitidis* (Albrecht and Ghon) and is a freeze-dried, 1.5-2.9 kb insert library constructed from *N. meningitidis* isolate. The deposit is described in Int. Bull. Bacteriol. Nomencl. Taxon. 8: 1-15 (1958).

[0186] The *Neisseria meningitidis* strain deposit is referred to herein as "the deposited strain" or as "the DNA of the deposited strain."

[0187] The deposited strain contains the full length BASB006 gene. The sequence of the polynucleotides contained in the deposited strain, as well as the amino acid sequence of any polypeptide encoded thereby, are controlling in the event of any conflict with any description of sequences herein.

[0188] The deposit of the deposited strain has been made under the terms of the Budapest Treaty on the International Recognition of the Deposit of Micro-organisms for Purposes of patent Procedure. The strain will be irrevocably and without restriction or condition released to the public upon the issuance of a patent. The deposited strain is provided merely as convenience to those of skill in the art and is not an admission that a deposit is required for enablement, such as that required under 35 U.S.C. §112.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 6

<210> SEQ ID NO 1

<211> LENGTH: 4365

<212> TYPE: DNA

<213> ORGANISM: *Neisseria meningitidis*

<400> SEQUENCE: 1

```
atgaaaaaca ccgacaaacg gacaaaccgaa acacaccgca aagccccgaa aaccgggtcgc   60
atccgcttct cgctgctta cttagccata tgcctgtcgt tcggcattct tccccaaagcc  120
tgggcgggac acacttattt cggcatcaac taccaatact atcgcgactt tgccgaaaat  180
aaaggcaagt ttgcagtcgg gccgaaagat attgaggttt acaacaaaaa aggggagttg  240
gtcggcaaat caatgacaaa agccccgatg attgatTTTT ctgtggtgtc gcgtaacggc  300
gtggcggcat tgggtggcga tcaatatatt gtgagcgtgg cacataacgg cggctataac  360
aacgttgatt ttggtgcgga gggaagcaat cccgatcagc accgtTTTTc ttatcaaatt  420
gtgaaaagaa ataattataa agcagggact aacggtcatc cttatggtgg cgattatcat  480
atgccgcggt tacataaatt tgtaaccgat gcagaacctg ttgaaatgac cagttatatg  540
gatgggcgga aatatatcga tcaaaataat taccctgacc gtgttcgtat tggggcaggc  600
aggcaatatt gccgatctga tgaagatgag cccaataacc gcgaaagtcc atatcatatt  660
gcaagtgcgt attcttggct cgttgggtgc aataccttg cacaaaatgg atcaggtggt  720
ggcacagtca acttaggtag tgaaaaaatt aaacatagcc catatggttt tttaccaaca  780
ggaggctcat ttggcgacag tggctcacca atgtttatct atgatgccc aaagcaaaag  840
tggttaatta atgggttatt gcaaacgggc aaccctata taggaaaaag caatggcttc  900
cagctgggtc gtaaagattg gttctatgat gaaatctttg ctggagatac ccattcagta  960
ttctacgaac cacatcaaaa tgggaaatc acttttcacg acaataataa tggcacagga 1020
aaaaatcaatg ccaaacatga acacaattct ctgcctaata gattaaaaac acgaaccggt 1080
caattgttta atgtttctt atccgagaca gcaagagaac ctgtttatca tgctgcaggt 1140
ggtgtcaaca gtatcagacc cagactgaat aatggagaaa atatttcctt tattgacgaa 1200
ggaaaaggcg aattgatact taccagcaac atcaatcaag gtgctggagg attatatctc 1260
caaggagatt ttacggtctc gcctgaaaat aacgaaacgt ggcaagggtc gggcgttcac 1320
atcagtgaag acagtaccgt tacttgaaaa gtaaacggcg tggcaaacga ccgctgtcc 1380
aaaaatcggca aaggcagcgt gcacgttcaa gccaaagggg aaaaccaagg ctcgatcagc 1440
gtggcgacg gtaaagttaa tttagatcaa caagcagatg aaaataataa aaaacaagcc 1500
tttagtgaaa tcggcttggt cagcggcagc ggtacggtgc aactgaatgc cgataatcag 1560
ttcaaccccc aaaaactcta ttctggcttt cgcggcggac gtttgattt gaacgggcat 1620
tcgctttcgt tccaccgat tcaaaatacc gatgaagggg cgatgattgt caaccacaat 1680
caagacaaag aatccaccgt taccattaca ggcaataaag atattgctac aaccggcaat 1740
aacaacagct tggatagcaa aaaagaaatt gcctacaacg gttggtttgg cgagaaagat 1800
acgacaaaaa cgaacgggcg gctcaacctt gtttaccagc ccgccgcaga agaccgcacc 1860
ctgctgcttt ccggcggaac aaatttaaac ggtaacatca cgcaacaaa cggcaaacctg 1920
```

-continued

tttttcagcg gcagaccgac accgcacgcc tacaatcatt taggaagcgg gtggtcaaaa	1980
atggaaggta tcccacaagg agaaatcgtg tgggacaacg actggatcaa ccgcacgttt	2040
aaagcggaaa atttccatat tcaggggcgg caggcgggtga tttcccgcaa tgttgccaaa	2100
gtggaaggcg attggcattt gagcaatcac gcccaagcag tttttggtgt cgcaccgcat	2160
caaagccaca caatctgtac acgttcggac tggacgggtc tgacaaaattg tgtcgaaaaa	2220
accattaccg acgataaagt gattgcttca ttgactaaga ccgacatcag cggcaatgtc	2280
agccttgccg atcacgctca tttaaatctc acagggttg ccacactcaa cggcaatctt	2340
agtgcaaatg gcgatacacg ttatacagtc agccacaacg ccacccaaaa cggcgacctt	2400
agcctcgtgg gcaatgccca agcaacattt aatcaagcca cattaaacgg caacacatcg	2460
gcttcgggca atgcttcatt taatctaagc aacaacgccg taaaaaacgg cagtctgacg	2520
ctttccggca acgctaaggc aaacgtaagc cattccgcac tcaacggtaa tgtctccta	2580
gccgataaag cagtattcca ttttgaaaag agccgcttta ccggacaaat cagcggcagc	2640
aaggatacgg cattacactt aaaagacagc gaatggacgc tggcgtcagg cacggaatta	2700
ggcaatttaa accttgacaa cgccaccatt aactcaatt ccgcctatcg ccacgatgcg	2760
gcagggggcg aaaccggcag tgcgacagat gcgcccggcc gccgttcgcg ccgttccta	2820
ttatccgtta cacctccggc ttccgagaaa tcccatttca acacgctgac ggtaaaacggc	2880
aaattgaacg gtcagggaac attccgctt atgtcggaac tcttcggcta ccgaagcgc	2940
aaattgaagc tggcgaaaag ttccgaaggc acttacact tggcggtaa caataccggc	3000
aacgaacccg taagcctcga tcaattgacg gtagtggaag gaaagacaa caaacccgtg	3060
tccgaaaacc ttaatttcac cctgcaaac gaacacgctc atgccggcgc gtggcgttac	3120
caactcatcc gcaaaagcgg cgagttccgc ctgcataatc cggtaaaaga acaagagctt	3180
tccgacaaac tcggcaaggc agaagccaaa aaacaggcgg gaaaagacaa cgcgcaaacg	3240
cttgacgcgc tgattcggc cgggcgcgat gccgtcgaag agacagaaag cgttgccgaa	3300
ccggcccggc aggcaggcgg gaaaaatgtc ggcatatgc aggcggagga agagaaaaaa	3360
cggtgacgag cgataaaga caccgccttg gcgaacacgc gcgaaggaa aaccggccg	3420
gctaccaccg ctttccccg cgcgcgcgc gcccgccggg atttgccgca accgcagccc	3480
caaccgcaac cccaaccgca gcgcgacctg atcagccgtt atgccaatag cggtttgagt	3540
gaattttccg ccacgctcaa cagcgttttc gccgtacagg acgaattaga ccgcgtattt	3600
gccgaagacc gccgcaacgc cgtttggaca agcggcatcc gggacaccaa aactaccgt	3660
tcgcaagatt tccgcgccta ccgccaacaa accgacctgc gccaaatcgg tatgcagaaa	3720
aacctcggca gggggcgcgt cggcatcctg ttttcgcaca accggaccga aaacaccttc	3780
gacgacggca tcggcaactc ggcacggctt gccacggcg ccgttttcgg gcaatacggc	3840
atcggcaggt tcgacatcgg catcagcacg ggcgggggtt ttagcagcgg cagtcttca	3900
gacgacatcg gaagcaaat ccgcccgcgc gtgctgcat acggcattca ggcacgatac	3960
cgcgcgggtt tcggcggtt cggcatcga cgcacatcg gcgcaacgcg ctatttcgtc	4020
caaaaagcgg attaccgcta cgaaaacgtc aatcgcgca cccccgcct tgcgttcaac	4080
cgctaccgcg cgggcattaa ggcagattat tcattcaaac cggcgcaaca catttccatc	4140
acgccttatt tgagcctgtc ctataccgat gccgcttcgg gcaaaagtcg aacgcgcgtc	4200

-continued

```

aataccgccc tattggctca ggatttcggc aaaaccgcca gtgcggaatg gggcgtaaac 4260
gccgaaatca aaggtttcac gctgtccctc cacgctgccg ccgccaagg cccgcaactg 4320
gaagcgcaac acagcgcggg catcaaatga ggctaccgct ggtaa 4365

```

```

<210> SEQ ID NO 2
<211> LENGTH: 1454
<212> TYPE: PRT
<213> ORGANISM: Neisseria meningitidis

```

```

<400> SEQUENCE: 2

```

```

Met Lys Thr Thr Asp Lys Arg Thr Thr Glu Thr His Arg Lys Ala Pro
1 5 10 15
Lys Thr Gly Arg Ile Arg Phe Ser Pro Ala Tyr Leu Ala Ile Cys Leu
20 25 30
Ser Phe Gly Ile Leu Pro Gln Ala Trp Ala Gly His Thr Tyr Phe Gly
35 40 45
Ile Asn Tyr Gln Tyr Tyr Arg Asp Phe Ala Glu Asn Lys Gly Lys Phe
50 55 60
Ala Val Gly Ala Lys Asp Ile Glu Val Tyr Asn Lys Lys Gly Glu Leu
65 70 75 80
Val Gly Lys Ser Met Thr Lys Ala Pro Met Ile Asp Phe Ser Val Val
85 90 95
Ser Arg Asn Gly Val Ala Ala Leu Val Gly Asp Gln Tyr Ile Val Ser
100 105 110
Val Ala His Asn Gly Gly Tyr Asn Asn Val Asp Phe Gly Ala Glu Gly
115 120 125
Ser Asn Pro Asp Gln His Arg Phe Ser Tyr Gln Ile Val Lys Arg Asn
130 135 140
Asn Tyr Lys Ala Gly Thr Asn Gly His Pro Tyr Gly Gly Asp Tyr His
145 150 155 160
Met Pro Arg Leu His Lys Phe Val Thr Asp Ala Glu Pro Val Glu Met
165 170 175
Thr Ser Tyr Met Asp Gly Arg Lys Tyr Ile Asp Gln Asn Asn Tyr Pro
180 185 190
Asp Arg Val Arg Ile Gly Ala Gly Arg Gln Tyr Trp Arg Ser Asp Glu
195 200 205
Asp Glu Pro Asn Asn Arg Glu Ser Ser Tyr His Ile Ala Ser Ala Tyr
210 215 220
Ser Trp Leu Val Gly Gly Asn Thr Phe Ala Gln Asn Gly Ser Gly Gly
225 230 235 240
Gly Thr Val Asn Leu Gly Ser Glu Lys Ile Lys His Ser Pro Tyr Gly
245 250 255
Phe Leu Pro Thr Gly Gly Ser Phe Gly Asp Ser Gly Ser Pro Met Phe
260 265 270
Ile Tyr Asp Ala Gln Lys Gln Lys Trp Leu Ile Asn Gly Val Leu Gln
275 280 285
Thr Gly Asn Pro Tyr Ile Gly Lys Ser Asn Gly Phe Gln Leu Val Arg
290 295 300
Lys Asp Trp Phe Tyr Asp Glu Ile Phe Ala Gly Asp Thr His Ser Val
305 310 315 320
Phe Tyr Glu Pro His Gln Asn Gly Lys Tyr Thr Phe His Asp Asn Asn
325 330 335

```

-continued

Asn Gly Thr Gly Lys Ile Asn Ala Lys His Glu His Asn Ser Leu Pro
 340 345 350

Asn Arg Leu Lys Thr Arg Thr Val Gln Leu Phe Asn Val Ser Leu Ser
 355 360 365

Glu Thr Ala Arg Glu Pro Val Tyr His Ala Ala Gly Gly Val Asn Ser
 370 375 380

Tyr Arg Pro Arg Leu Asn Asn Gly Glu Asn Ile Ser Phe Ile Asp Glu
 385 390 395 400

Gly Lys Gly Glu Leu Ile Leu Thr Ser Asn Ile Asn Gln Gly Ala Gly
 405 410 415

Gly Leu Tyr Phe Gln Gly Asp Phe Thr Val Ser Pro Glu Asn Asn Glu
 420 425 430

Thr Trp Gln Gly Ala Gly Val His Ile Ser Glu Asp Ser Thr Val Thr
 435 440 445

Trp Lys Val Asn Gly Val Ala Asn Asp Arg Leu Ser Lys Ile Gly Lys
 450 455 460

Gly Thr Leu His Val Gln Ala Lys Gly Glu Asn Gln Gly Ser Ile Ser
 465 470 475 480

Val Gly Asp Gly Lys Val Ile Leu Asp Gln Gln Ala Asp Glu Asn Asn
 485 490 495

Lys Lys Gln Ala Phe Ser Glu Ile Gly Leu Val Ser Gly Arg Gly Thr
 500 505 510

Val Gln Leu Asn Ala Asp Asn Gln Phe Asn Pro Asp Lys Leu Tyr Phe
 515 520 525

Gly Phe Arg Gly Gly Arg Leu Asp Leu Asn Gly His Ser Leu Ser Phe
 530 535 540

His Arg Ile Gln Asn Thr Asp Glu Gly Ala Met Ile Val Asn His Asn
 545 550 555 560

Gln Asp Lys Glu Ser Thr Val Thr Ile Thr Gly Asn Lys Asp Ile Ala
 565 570 575

Thr Thr Gly Asn Asn Ser Leu Asp Ser Lys Lys Glu Ile Ala Tyr
 580 585 590

Asn Gly Trp Phe Gly Glu Lys Asp Thr Thr Lys Thr Asn Gly Arg Leu
 595 600 605

Asn Leu Val Tyr Gln Pro Ala Ala Glu Asp Arg Thr Leu Leu Leu Ser
 610 615 620

Gly Gly Thr Asn Leu Asn Gly Asn Ile Thr Gln Thr Asn Gly Lys Leu
 625 630 635 640

Phe Phe Ser Gly Arg Pro Thr Pro His Ala Tyr Asn His Leu Gly Ser
 645 650 655

Gly Trp Ser Lys Met Glu Gly Ile Pro Gln Gly Glu Ile Val Trp Asp
 660 665 670

Asn Asp Trp Ile Asn Arg Thr Phe Lys Ala Glu Asn Phe His Ile Gln
 675 680 685

Gly Gly Gln Ala Val Ile Ser Arg Asn Val Ala Lys Val Glu Gly Asp
 690 695 700

Trp His Leu Ser Asn His Ala Gln Ala Val Phe Gly Val Ala Pro His
 705 710 715 720

Gln Ser His Thr Ile Cys Thr Arg Ser Asp Trp Thr Gly Leu Thr Asn
 725 730 735

-continued

Cys Val Glu Lys Thr Ile Thr Asp Asp Lys Val Ile Ala Ser Leu Thr
 740 745 750
 Lys Thr Asp Ile Ser Gly Asn Val Ser Leu Ala Asp His Ala His Leu
 755 760 765
 Asn Leu Thr Gly Leu Ala Thr Leu Asn Gly Asn Leu Ser Ala Asn Gly
 770 775 780
 Asp Thr Arg Tyr Thr Val Ser His Asn Ala Thr Gln Asn Gly Asp Leu
 785 790 795 800
 Ser Leu Val Gly Asn Ala Gln Ala Thr Phe Asn Gln Ala Thr Leu Asn
 805 810 815
 Gly Asn Thr Ser Ala Ser Gly Asn Ala Ser Phe Asn Leu Ser Asn Asn
 820 825 830
 Ala Val Gln Asn Gly Ser Leu Thr Leu Ser Gly Asn Ala Lys Ala Asn
 835 840 845
 Val Ser His Ser Ala Leu Asn Gly Asn Val Ser Leu Ala Asp Lys Ala
 850 855 860
 Val Phe His Phe Glu Ser Ser Arg Phe Thr Gly Gln Ile Ser Gly Ser
 865 870 875 880
 Lys Asp Thr Ala Leu His Leu Lys Asp Ser Glu Trp Thr Leu Pro Ser
 885 890 895
 Gly Thr Glu Leu Gly Asn Leu Asn Leu Asp Asn Ala Thr Ile Thr Leu
 900 905 910
 Asn Ser Ala Tyr Arg His Asp Ala Ala Gly Ala Gln Thr Gly Ser Ala
 915 920 925
 Thr Asp Ala Pro Arg Arg Arg Ser Arg Arg Ser Leu Leu Ser Val Thr
 930 935 940
 Pro Pro Ala Ser Ala Glu Ser His Phe Asn Thr Leu Thr Val Asn Gly
 945 950 955 960
 Lys Leu Asn Gly Gln Gly Thr Phe Arg Phe Met Ser Glu Leu Phe Gly
 965 970 975
 Tyr Arg Ser Asp Lys Leu Lys Leu Ala Glu Ser Ser Glu Gly Thr Tyr
 980 985 990
 Thr Leu Ala Val Asn Asn Thr Gly Asn Glu Pro Val Ser Leu Asp Gln
 995 1000 1005
 Leu Thr Val Val Glu Gly Lys Asp Asn Lys Pro Leu Ser Glu Asn Leu
 1010 1015 1020
 Asn Phe Thr Leu Gln Asn Glu His Val Asp Ala Gly Ala Trp Arg Tyr
 1025 1030 1035 1040
 Gln Leu Ile Arg Lys Asp Gly Glu Phe Arg Leu His Asn Pro Val Lys
 1045 1050 1055
 Glu Gln Glu Leu Ser Asp Lys Leu Gly Lys Ala Glu Ala Lys Lys Gln
 1060 1065 1070
 Ala Gly Lys Asp Asn Ala Gln Ser Leu Asp Ala Leu Ile Ala Ala Gly
 1075 1080 1085
 Arg Asp Ala Val Glu Lys Thr Glu Ser Val Ala Glu Pro Ala Arg Gln
 1090 1095 1100
 Ala Gly Gly Glu Asn Val Gly Ile Met Gln Ala Glu Glu Glu Lys Lys
 1105 1110 1115 1120
 Arg Val Gln Ala Asp Lys Asp Thr Ala Leu Ala Lys Gln Arg Glu Gly
 1125 1130 1135
 Lys Thr Arg Pro Ala Thr Thr Ala Phe Pro Arg Ala Arg Arg Ala Arg

-continued

	1140		1145		1150	
Arg Asp Leu Pro Gln Pro Gln Pro Gln Pro Gln Pro Gln Pro Gln Arg	1155		1160		1165	
Asp Leu Ile Ser Arg Tyr Ala Asn Ser Gly Leu Ser Glu Phe Ser Ala	1170		1175		1180	
Thr Leu Asn Ser Val Phe Ala Val Gln Asp Glu Leu Asp Arg Val Phe	1185		1190		1195	1200
Ala Glu Asp Arg Arg Asn Ala Val Trp Thr Ser Gly Ile Arg Asp Thr		1205		1210		1215
Lys His Tyr Arg Ser Gln Asp Phe Arg Ala Tyr Arg Gln Gln Thr Asp		1220		1225		1230
Leu Arg Gln Ile Gly Met Gln Lys Asn Leu Gly Ser Gly Arg Val Gly		1235		1240		1245
Ile Leu Phe Ser His Asn Arg Thr Glu Asn Thr Phe Asp Asp Gly Ile		1250		1255		1260
Gly Asn Ser Ala Arg Leu Ala His Gly Ala Val Phe Gly Gln Tyr Gly		1265		1270		1275
Ile Gly Arg Phe Asp Ile Gly Ile Ser Thr Gly Ala Gly Phe Ser Ser		1285		1290		1295
Gly Ser Leu Ser Asp Asp Ile Gly Ser Lys Ile Arg Arg Arg Val Leu		1300		1305		1310
His Tyr Gly Ile Gln Ala Arg Tyr Arg Ala Gly Phe Gly Gly Phe Gly		1315		1320		1325
Ile Glu Pro His Ile Gly Ala Thr Arg Tyr Phe Val Gln Lys Ala Asp		1330		1335		1340
Tyr Arg Tyr Glu Asn Val Asn Ile Ala Thr Pro Gly Leu Ala Phe Asn		1345		1350		1355
Arg Tyr Arg Ala Gly Ile Lys Ala Asp Tyr Ser Phe Lys Pro Ala Gln		1365		1370		1375
His Ile Ser Ile Thr Pro Tyr Leu Ser Leu Ser Tyr Thr Asp Ala Ala		1380		1385		1390
Ser Gly Lys Val Arg Thr Arg Val Asn Thr Ala Val Leu Ala Gln Asp		1395		1400		1405
Phe Gly Lys Thr Arg Ser Ala Glu Trp Gly Val Asn Ala Glu Ile Lys		1410		1415		1420
Gly Phe Thr Leu Ser Leu His Ala Ala Ala Ala Lys Gly Pro Gln Leu		1425		1430		1435
Glu Ala Gln His Ser Ala Gly Ile Lys Leu Gly Tyr Arg Trp		1445		1450		

<210> SEQ ID NO 3
 <211> LENGTH: 4374
 <212> TYPE: DNA
 <213> ORGANISM: Neisseria meningitidis

<400> SEQUENCE: 3

atgaaaaaca cgcacaaacg gacaaccgaa acacaccgca aagccccgaa aaccggccgc	60
atccgcttct cgctgctta cttagccata tgctgtcgt tcggcattct tcccgaagcc	120
tgggcgggac acacttattt cggcatcaac taccaatact atcgcgactt tgccgaaaat	180
aaaggcaagt ttgcagtcgg ggcgaaagat attgaggttt acaacaaaaa aggggagttg	240
gtcggcaaat caatgacaaa agccccgatg attgattttt ctgtggtgtc gcgtaacggc	300

-continued

gtggcgcat	tggtaggca	tcaatatatt	gtgagcgtgg	cacataacgg	cggtataac	360
aacgttgatt	tggtaggca	aggaagaaat	cccgatcaac	atcgttttac	ttataaaatt	420
gtgaaacgga	ataattataa	agcagggact	aaaggccatc	cttatggtgg	cgattatcat	480
atgccgcggt	tacataaatt	tgtcacagat	gcagaacctg	ttgaaatgac	cagttatatg	540
gatggggcga	aatatatacg	tcaaaaaat	taccctgacc	gtgttcgtat	tggggcaggc	600
aggcaatatt	ggcgatctga	tgaagatgag	cccaataacc	gcgaaagtcc	atatcatatt	660
gcaagtgcgt	attcttggct	cgttggtggc	aatacctttg	cacaaaatgg	atcagggtgg	720
ggcacagtca	acttaggtag	tgaaaaaatt	aaacatagcc	catatggttt	tttaccaaca	780
ggaggetcat	tggcgacag	tggctcacca	atgtttatct	atgatgcca	aaagcaaaag	840
tggttaatta	atgggttatt	gcaaacgggc	aaccctata	taggaaaaag	caatggcttc	900
cagctgggtc	gtaaagattg	gttctatgat	gaaatctttg	ctggagatac	ccattcagta	960
ttctacgaac	cacgtcaaaa	tgggaaatc	tcttttaacg	acgataataa	tggcacagga	1020
aaaaatcaatg	ccaaacatga	acacaattct	ctgcctaata	gattaaaaac	acgaaccggt	1080
caattgttta	atgtttcttt	atccgagaca	gcaagagaac	ctgtttatca	tgctgcaggt	1140
ggtgtcaaca	gttatcgacc	cagactgaat	aatggagaaa	atatttcctt	tattgacgaa	1200
ggaaaaggcg	aattgatact	taccagcaac	atcaatcaag	gtgctggagg	attatatttc	1260
caaggagatt	ttacggtctc	gcctgaaaat	aacgaaactg	ggcaagggtc	ggcggttcat	1320
atcagtgaag	acagtaccgt	tacttgaaaa	gtaaacggcg	tggcaaacga	ccgctgtcc	1380
aaaaatcgga	aaggcacgct	gcacgttcaa	gccaaagggg	aaaaccaagg	ctcgatcagc	1440
gtggcgacg	gtacagtcat	tttgatcag	caggcagacg	ataaaggcaa	aaaacaagcc	1500
tttagtgaaa	tcggcttgg	cagcggcag	ggtacggtgc	aactgaatgc	cgataatcag	1560
ttcaaccccg	acaaactcta	tttcggcttt	cgcgcgac	gtttgattt	aaacgggcat	1620
tcgcttctgt	tccaccgtat	tcaaaaatcc	gatgaagggg	cgatgattgt	caaccacaat	1680
caagacaaag	aatccaccgt	taccattaca	ggcaataaag	atattgctac	aaccggcaat	1740
aaacaacagct	tggatagcaa	aaaagaaatt	gcctacaacg	gttggtttgg	cgagaaagat	1800
acgacaaaa	cgaacggg	gctcaacctt	gtttaccagc	ccgcccaga	agaccgcacc	1860
ctgctgcttt	ccggcggaac	aaatttaaac	ggcaacatca	cgcaacaaa	cgcaaacctg	1920
ttttcagcg	gcagaccaac	accgcacgcc	tacaatcatt	taaacgacca	ttggtcgcaa	1980
aaagagggca	ttccacgcg	ggaaatcgtg	tgggacaacg	actggatcaa	ccgcacattt	2040
aaagcgaaa	acttccaaat	taaaggcgga	caggcgttgg	ttcccgcaa	tgttgccaaa	2100
gtgaaaggcg	attggcattt	gagcaatcac	gccaagcag	ttttggtgt	cgcaccgcat	2160
caaagccaca	caatctgtac	acgttcggac	tggacgggtc	tgacaaattg	tgtcgaaaaa	2220
accattaccg	acgataaagt	gattgcttca	ttgactaaga	ccgacatcag	cggaatgtc	2280
gatcttgccg	atcacgctca	tttaaatctc	acagggttgg	ccacactcaa	cggaatctt	2340
agtgcaaatg	gcgatacacg	ttatacagtc	agccacaacg	ccacccaaaa	cggaacctt	2400
agcctcgtgg	gcaatgcccc	agcaacattt	aatcaagcca	cattaaacgg	caacacatcg	2460
gcttcgggca	atgcttcatt	taactaagc	gaccacgcg	tacaaaacgg	cagtctgacg	2520
cttccggca	acgctaaggc	aaacgtaagc	cattccgcac	tcaacggtaa	tgtctccta	2580

-continued

```

gccgataagg cagtattcca ttttgaaagc agccgcttta ccgacaaat cagcggcggc 2640
aaggatacgg cattacactt aaaagacagc gaatggacgc tgccgtcagg cacggaatta 2700
ggcaatttaa accttgacaa cgccaccatt acaactcaatt ccgcctatcg ccacgatgcg 2760
gcagggggcg aaaccggcag tgcgacagat gcgccgccc gccgttcgcg ccgttcgcg 2820
cgttccctat tatccgttac accgccaact tcggtagaat cccgtttcaa cacgctgacg 2880
gtaaaccggc aattgaacgg tcaggaaca ttccgcttta tgcggaact cttcggtac 2940
cgcagcgaca aattgaagct ggcgaaaagt tccgaaggca cttacacctt ggcggtaaac 3000
aataccggca acgaactgc aagccttgaa caattgacgg tagtggaagg aaaagacaac 3060
aaaccgctgt ccgaaaactt taatttcacc ttgcaaacg aacacgtcga tgcggcgcg 3120
tggcgtaacc aactcatccg caaagacggc gagttccgcc tgcataatcc ggtcaaagaa 3180
caagagcttt ccgacaaact cggcaaggca gaagcaaaa aacaggcggg aaaagacaac 3240
gcgcaaagcc ttgacgcgct gattgcggcc gggcgcgatg ccgtcgaaa gacagaaagc 3300
gttgccgaac cggcccggca ggcaggcggg gaaaatgtcg gcattatgca ggcggaggaa 3360
gagaaaaaac ggggtgcaggc ggataaagac accgccttg cgaaaacagc cgaagcggaa 3420
accggccgg ctaccaccgc cttccccgc gcccgccgcg cccgcccggg tttgcccaa 3480
ctgcaacccc aaccgcagcc ccaaccgcag cgcgacctga tcagccgta tgccaatagc 3540
ggtttgagt aattttccg cacgctcaac agcgttttcg ccgtacagga cgaattagc 3600
cgcgtatttg ccgaagaacg ccgcaacgcc gtttgacaa gcgcatccg ggacaccaa 3660
cactaccggt cgcaagattt ccgcccctac cgccaacaaa ccgacctgcg ccaaatcggt 3720
atgcagaaaa acctcggcag cgggcgcgct ggcatcctgt tttcgacaa ccggaccgaa 3780
aacaccttcg acgacggcat cggcaactcg gcacggcttg ccacggcgc cgttttcggg 3840
caatacggca tcgacagggt ctacatcgcc atcagcggcg gcgcggttt tagcagcggc 3900
agcctttcag acggcatcgg aggcaaaatc cgccgcccgc tgctgcatta cggcattcag 3960
gcacgatacc gcgcccgttt cggcggatc ggcatcgaac cgcacatcgg cgcaacgcgc 4020
tatttcgtcc aaaaaggcga ttaccgctac gaaaacgtca atatcgccac ccccgccct 4080
gcattcaacc gctaccgcgc gggcattaag gcagattatt cattcaaacc ggcgcaaac 4140
atttccatca cgccttattt gagcctgtcc tataccgatg ccgcttcggg caaagtccga 4200
acacgcgtca ataccgccg attggctcag gatttcggca aaaccgcag tgcggaatgg 4260
ggcgtaaacg ccgaaatcaa agccttcag ctgtccctcc acgctgccc cgccaaaggc 4320
ccgcaactgg aagcgcaaca cagcgcgggc atcaaattag gctaccgctg gtaa 4374

```

<210> SEQ ID NO 4

<211> LENGTH: 1457

<212> TYPE: PRT

<213> ORGANISM: *Neisseria meningitidis*

<400> SEQUENCE: 4

```

Met Lys Thr Thr Asp Lys Arg Thr Thr Glu Thr His Arg Lys Ala Pro
1           5           10           15

```

```

Lys Thr Gly Arg Ile Arg Phe Ser Pro Ala Tyr Leu Ala Ile Cys Leu
20           25           30

```

```

Ser Phe Gly Ile Leu Pro Gln Ala Trp Ala Gly His Thr Tyr Phe Gly

```

-continued

35					40					45					
Ile	Asn	Tyr	Gln	Tyr	Tyr	Arg	Asp	Phe	Ala	Glu	Asn	Lys	Gly	Lys	Phe
50					55					60					
Ala	Val	Gly	Ala	Lys	Asp	Ile	Glu	Val	Tyr	Asn	Lys	Lys	Gly	Glu	Leu
65					70					75					80
Val	Gly	Lys	Ser	Met	Thr	Lys	Ala	Pro	Met	Ile	Asp	Phe	Ser	Val	Val
				85					90					95	
Ser	Arg	Asn	Gly	Val	Ala	Ala	Leu	Val	Gly	Asp	Gln	Tyr	Ile	Val	Ser
			100					105						110	
Val	Ala	His	Asn	Gly	Gly	Tyr	Asn	Asn	Val	Asp	Phe	Gly	Ala	Glu	Gly
		115					120					125			
Arg	Asn	Pro	Asp	Gln	His	Arg	Phe	Thr	Tyr	Lys	Ile	Val	Lys	Arg	Asn
		130					135					140			
Asn	Tyr	Lys	Ala	Gly	Thr	Lys	Gly	His	Pro	Tyr	Gly	Gly	Asp	Tyr	His
		145					150					155			160
Met	Pro	Arg	Leu	His	Lys	Phe	Val	Thr	Asp	Ala	Glu	Pro	Val	Glu	Met
				165					170					175	
Thr	Ser	Tyr	Met	Asp	Gly	Arg	Lys	Tyr	Ile	Asp	Gln	Asn	Asn	Tyr	Pro
			180					185						190	
Asp	Arg	Val	Arg	Ile	Gly	Ala	Gly	Arg	Gln	Tyr	Trp	Arg	Ser	Asp	Glu
		195					200						205		
Asp	Glu	Pro	Asn	Asn	Arg	Glu	Ser	Ser	Tyr	His	Ile	Ala	Ser	Ala	Tyr
		210					215					220			
Ser	Trp	Leu	Val	Gly	Gly	Asn	Thr	Phe	Ala	Gln	Asn	Gly	Ser	Gly	Gly
		225					230					235			240
Gly	Thr	Val	Asn	Leu	Gly	Ser	Glu	Lys	Ile	Lys	His	Ser	Pro	Tyr	Gly
				245					250					255	
Phe	Leu	Pro	Thr	Gly	Gly	Ser	Phe	Gly	Asp	Ser	Gly	Ser	Pro	Met	Phe
			260					265						270	
Ile	Tyr	Asp	Ala	Gln	Lys	Gln	Lys	Trp	Leu	Ile	Asn	Gly	Val	Leu	Gln
		275					280					285			
Thr	Gly	Asn	Pro	Tyr	Ile	Gly	Lys	Ser	Asn	Gly	Phe	Gln	Leu	Val	Arg
		290					295					300			
Lys	Asp	Trp	Phe	Tyr	Asp	Glu	Ile	Phe	Ala	Gly	Asp	Thr	His	Ser	Val
		305					310					315			320
Phe	Tyr	Glu	Pro	Arg	Gln	Asn	Gly	Lys	Tyr	Ser	Phe	Asn	Asp	Asp	Asn
				325					330					335	
Asn	Gly	Thr	Gly	Lys	Ile	Asn	Ala	Lys	His	Glu	His	Asn	Ser	Leu	Pro
			340					345						350	
Asn	Arg	Leu	Lys	Thr	Arg	Thr	Val	Gln	Leu	Phe	Asn	Val	Ser	Leu	Ser
		355					360					365			
Glu	Thr	Ala	Arg	Glu	Pro	Val	Tyr	His	Ala	Ala	Gly	Gly	Val	Asn	Ser
		370					375					380			
Tyr	Arg	Pro	Arg	Leu	Asn	Asn	Gly	Glu	Asn	Ile	Ser	Phe	Ile	Asp	Glu
		385					390					395			400
Gly	Lys	Gly	Glu	Leu	Ile	Leu	Thr	Ser	Asn	Ile	Asn	Gln	Gly	Ala	Gly
				405					410					415	
Gly	Leu	Tyr	Phe	Gln	Gly	Asp	Phe	Thr	Val	Ser	Pro	Glu	Asn	Asn	Glu
			420					425						430	
Thr	Trp	Gln	Gly	Ala	Gly	Val	His	Ile	Ser	Glu	Asp	Ser	Thr	Val	Thr
		435					440						445		

-continued

Trp Lys Val Asn Gly Val Ala Asn Asp Arg Leu Ser Lys Ile Gly Lys
 450 455 460

Gly Thr Leu His Val Gln Ala Lys Gly Glu Asn Gln Gly Ser Ile Ser
 465 470 475 480

Val Gly Asp Gly Thr Val Ile Leu Asp Gln Gln Ala Asp Asp Lys Gly
 485 490 495

Lys Lys Gln Ala Phe Ser Glu Ile Gly Leu Val Ser Gly Arg Gly Thr
 500 505 510

Val Gln Leu Asn Ala Asp Asn Gln Phe Asn Pro Asp Lys Leu Tyr Phe
 515 520 525

Gly Phe Arg Gly Gly Arg Leu Asp Leu Asn Gly His Ser Leu Ser Phe
 530 535 540

His Arg Ile Gln Asn Thr Asp Glu Gly Ala Met Ile Val Asn His Asn
 545 550 555 560

Gln Asp Lys Glu Ser Thr Val Thr Ile Thr Gly Asn Lys Asp Ile Ala
 565 570 575

Thr Thr Gly Asn Asn Ser Leu Asp Ser Lys Lys Glu Ile Ala Tyr
 580 585 590

Asn Gly Trp Phe Gly Glu Lys Asp Thr Thr Lys Thr Asn Gly Arg Leu
 595 600 605

Asn Leu Val Tyr Gln Pro Ala Ala Glu Asp Arg Thr Leu Leu Leu Ser
 610 615 620

Gly Gly Thr Asn Leu Asn Gly Asn Ile Thr Gln Thr Asn Gly Lys Leu
 625 630 635 640

Phe Phe Ser Gly Arg Pro Thr Pro His Ala Tyr Asn His Leu Asn Asp
 645 650 655

His Trp Ser Gln Lys Glu Gly Ile Pro Arg Gly Glu Ile Val Trp Asp
 660 665 670

Asn Asp Trp Ile Asn Arg Thr Phe Lys Ala Glu Asn Phe Gln Ile Lys
 675 680 685

Gly Gly Gln Ala Val Val Ser Arg Asn Val Ala Lys Val Lys Gly Asp
 690 695 700

Trp His Leu Ser Asn His Ala Gln Ala Val Phe Gly Val Ala Pro His
 705 710 715 720

Gln Ser His Thr Ile Cys Thr Arg Ser Asp Trp Thr Gly Leu Thr Asn
 725 730 735

Cys Val Glu Lys Thr Ile Thr Asp Asp Lys Val Ile Ala Ser Leu Thr
 740 745 750

Lys Thr Asp Ile Ser Gly Asn Val Asp Leu Ala Asp His Ala His Leu
 755 760 765

Asn Leu Thr Gly Leu Ala Thr Leu Asn Gly Asn Leu Ser Ala Asn Gly
 770 775 780

Asp Thr Arg Tyr Thr Val Ser His Asn Ala Thr Gln Asn Gly Asn Leu
 785 790 795 800

Ser Leu Val Gly Asn Ala Gln Ala Thr Phe Asn Gln Ala Thr Leu Asn
 805 810 815

Gly Asn Thr Ser Ala Ser Gly Asn Ala Ser Phe Asn Leu Ser Asp His
 820 825 830

Ala Val Gln Asn Gly Ser Leu Thr Leu Ser Gly Asn Ala Lys Ala Asn
 835 840 845

-continued

Val	Ser	His	Ser	Ala	Leu	Asn	Gly	Asn	Val	Ser	Leu	Ala	Asp	Lys	Ala
	850					855					860				
Val	Phe	His	Phe	Glu	Ser	Ser	Arg	Phe	Thr	Gly	Gln	Ile	Ser	Gly	Gly
	865				870					875				880	
Lys	Asp	Thr	Ala	Leu	His	Leu	Lys	Asp	Ser	Glu	Trp	Thr	Leu	Pro	Ser
				885					890					895	
Gly	Thr	Glu	Leu	Gly	Asn	Leu	Asn	Leu	Asp	Asn	Ala	Thr	Ile	Thr	Leu
			900					905					910		
Asn	Ser	Ala	Tyr	Arg	His	Asp	Ala	Ala	Gly	Ala	Gln	Thr	Gly	Ser	Ala
		915					920					925			
Thr	Asp	Ala	Pro	Arg	Arg	Arg	Ser	Arg	Arg	Ser	Arg	Arg	Ser	Leu	Leu
	930					935					940				
Ser	Val	Thr	Pro	Pro	Thr	Ser	Val	Glu	Ser	Arg	Phe	Asn	Thr	Leu	Thr
	945				950					955				960	
Val	Asn	Gly	Lys	Leu	Asn	Gly	Gln	Gly	Thr	Phe	Arg	Phe	Met	Ser	Glu
				965					970					975	
Leu	Phe	Gly	Tyr	Arg	Ser	Asp	Lys	Leu	Lys	Leu	Ala	Glu	Ser	Ser	Glu
			980				985						990		
Gly	Thr	Tyr	Thr	Leu	Ala	Val	Asn	Asn	Thr	Gly	Asn	Glu	Pro	Ala	Ser
		995					1000					1005			
Leu	Glu	Gln	Leu	Thr	Val	Val	Glu	Gly	Lys	Asp	Asn	Lys	Pro	Leu	Ser
	1010					1015					1020				
Glu	Asn	Phe	Asn	Phe	Thr	Leu	Gln	Asn	Glu	His	Val	Asp	Ala	Gly	Ala
	1025				1030					1035				1040	
Trp	Arg	Tyr	Gln	Leu	Ile	Arg	Lys	Asp	Gly	Glu	Phe	Arg	Leu	His	Asn
				1045					1050					1055	
Pro	Val	Lys	Glu	Gln	Glu	Leu	Ser	Asp	Lys	Leu	Gly	Lys	Ala	Glu	Ala
			1060					1065					1070		
Lys	Lys	Gln	Ala	Glu	Lys	Asp	Asn	Ala	Gln	Ser	Leu	Asp	Ala	Leu	Ile
		1075					1080						1085		
Ala	Ala	Gly	Arg	Asp	Ala	Val	Glu	Lys	Thr	Glu	Ser	Val	Ala	Glu	Pro
	1090					1095						1100			
Ala	Arg	Gln	Ala	Gly	Gly	Glu	Asn	Val	Gly	Ile	Met	Gln	Ala	Glu	Glu
	1105				1110					1115				1120	
Glu	Lys	Lys	Arg	Val	Gln	Ala	Asp	Lys	Asp	Thr	Ala	Leu	Ala	Lys	Gln
				1125					1130					1135	
Arg	Glu	Ala	Glu	Thr	Arg	Pro	Ala	Thr	Thr	Ala	Phe	Pro	Arg	Ala	Arg
			1140					1145					1150		
Arg	Ala	Arg	Arg	Asp	Leu	Pro	Gln	Leu	Gln	Pro	Gln	Pro	Gln	Pro	Gln
		1155					1160						1165		
Pro	Gln	Arg	Asp	Leu	Ile	Ser	Arg	Tyr	Ala	Asn	Ser	Gly	Leu	Ser	Glu
	1170					1175						1180			
Phe	Ser	Ala	Thr	Leu	Asn	Ser	Val	Phe	Ala	Val	Gln	Asp	Glu	Leu	Asp
	1185				1190					1195				1200	
Arg	Val	Phe	Ala	Glu	Glu	Arg	Arg	Asn	Ala	Val	Trp	Thr	Ser	Gly	Ile
				1205					1210					1215	
Arg	Asp	Thr	Lys	His	Tyr	Arg	Ser	Gln	Asp	Phe	Arg	Ala	Tyr	Arg	Gln
			1220					1225					1230		
Gln	Thr	Asp	Leu	Arg	Gln	Ile	Gly	Met	Gln	Lys	Asn	Leu	Gly	Ser	Gly
		1235					1240						1245		
Arg	Val	Gly	Ile	Leu	Phe	Ser	His	Asn	Arg	Thr	Glu	Asn	Thr	Phe	Asp

-continued

1250	1255	1260
Asp Gly Ile Gly Asn Ser Ala Arg Leu Ala His Gly Ala Val Phe Gly 1265	1270	1275 1280
Gln Tyr Gly Ile Asp Arg Phe Tyr Ile Gly Ile Ser Ala Gly Ala Gly 1285	1290	1295
Phe Ser Ser Gly Ser Leu Ser Asp Gly Ile Gly Gly Lys Ile Arg Arg 1300	1305	1310
Arg Val Leu His Tyr Gly Ile Gln Ala Arg Tyr Arg Ala Gly Phe Gly 1315	1320	1325
Gly Phe Gly Ile Glu Pro His Ile Gly Ala Thr Arg Tyr Phe Val Gln 1330	1335	1340
Lys Ala Asp Tyr Arg Tyr Glu Asn Val Asn Ile Ala Thr Pro Gly Leu 1345	1350	1355 1360
Ala Phe Asn Arg Tyr Arg Ala Gly Ile Lys Ala Asp Tyr Ser Phe Lys 1365	1370	1375
Pro Ala Gln His Ile Ser Ile Thr Pro Tyr Leu Ser Leu Ser Tyr Thr 1380	1385	1390
Asp Ala Ala Ser Gly Lys Val Arg Thr Arg Val Asn Thr Ala Val Leu 1395	1400	1405
Ala Gln Asp Phe Gly Lys Thr Arg Ser Ala Glu Trp Gly Val Asn Ala 1410	1415	1420
Glu Ile Lys Gly Phe Thr Leu Ser Leu His Ala Ala Ala Ala Lys Gly 1425	1430	1435 1440
Pro Gln Leu Glu Ala Gln His Ser Ala Gly Ile Lys Leu Gly Tyr Arg 1445	1450	1455
Trp		
<210> SEQ ID NO 5		
<211> LENGTH: 34		
<212> TYPE: DNA		
<213> ORGANISM: Artificial Sequence		
<220> FEATURE:		
<223> OTHER INFORMATION: primer		
<400> SEQUENCE: 5		
gggggctagc aaaacaaccg acaaacggac aacc		34
<210> SEQ ID NO 6		
<211> LENGTH: 40		
<212> TYPE: DNA		
<213> ORGANISM: Artificial Sequence		
<220> FEATURE:		
<223> OTHER INFORMATION: primer		
<400> SEQUENCE: 6		
ggggaagcctt ccagcggtag cggtagccta atttgatgcc		40

What is claimed is:

1. An isolated polypeptide comprising an immunogenic fragment of at least 175 contiguous amino acids of SEQ ID NO:2, wherein the immunogenic fragment, when administered to a subject induces an antibody that binds to said fragment within SEQ ID NO:2.

2. An isolated recombinant fusion protein comprising the polypeptide of claim 1.

3. An immunogenic composition comprising the polypeptide of claim 1 and a pharmaceutically acceptable carrier.

4. The immunogenic composition of claim 3, further comprising an adjuvant.

5. The immunogenic composition of claim 3, further comprising at least one other *Neisseria meningitidis* antigen other than a polypeptide comprising SEQ ID NO:2.

6. An immunogenic composition comprising the fusion protein of claim 2 and a pharmaceutically acceptable carrier.

7. A method for inducing an antibody in a mammal comprising administering the polypeptide of claim 1 to the mammal in an amount effective to induce an antibody.

8. An isolated polypeptide comprising an immunogenic fragment of at least 957 contiguous amino acids of SEQ ID NO:4, wherein the immunogenic fragment, when administered to a subject induces an antibody that binds to said fragment within SEQ ID NO:4.

9. An isolated recombinant fusion protein comprising the polypeptide of claim 8.

10. An immunogenic composition comprising the polypeptide of claim 8 and a pharmaceutically acceptable carrier.

11. The immunogenic composition of claim 10, further comprising at least one other *Neisseria meningitidis* antigen other than a polypeptide comprising SEQ ID NO:4.

12. The immunogenic composition of claim 10, further comprising an adjuvant.

13. An immunogenic composition comprising the fusion protein of claim 9 and a pharmaceutically acceptable carrier.

14. A method for inducing an antibody in a mammal comprising administering the polypeptide of claim 8 to the mammal in an amount effective to induce an antibody.

15. An isolated protein comprising a polypeptide having at least 97% identity with the amino acid sequence of SEQ ID NO:2, wherein the protein, when administered to a subject induces an antibody that binds to said polypeptide.

16. An immunogenic composition comprising the protein of claim 15 and a pharmaceutically acceptable carrier.

17. The immunogenic composition of claim 16, further comprising an adjuvant.

18. A method for inducing an antibody in a mammal comprising administering the protein of claim 15 to the mammal in an amount effective to induce an antibody.

19. An isolated protein comprising a polypeptide having at least 99.9% identity with the amino acid sequence of SEQ ID NO:4, wherein the protein, when administered to a subject induces an antibody that binds to said polypeptide.

20. An immunogenic composition comprising the protein of claim 19 and a pharmaceutically acceptable carrier.

21. The immunogenic composition of claim 19, further comprising an adjuvant.

22. A method for inducing an antibody in a mammal comprising administering the protein of claim 19 to the mammal in an amount effective to induce an antibody.

* * * * *