




(86) Date de dépôt PCT/PCT Filing Date: 2003/02/21
 (87) Date publication PCT/PCT Publication Date: 2003/09/04
 (85) Entrée phase nationale/National Entry: 2004/08/19
 (86) N° demande PCT/PCT Application No.: US 2003/005151
 (87) N° publication PCT/PCT Publication No.: 2003/072715
 (30) Priorités/Priorities: 2002/02/21 (60/357,845) US;
 2002/12/20 (60/434,644) US

(51) Cl.Int.⁷/Int.Cl.⁷ A61K 48/00, A61K 31/7088, A61K 38/17
 (71) Demandeur/Applicant:
 WYETH, US
 (72) Inventeurs/Inventors:
 HILL, JENNIFER J., US;
 WOLFMAN, NEIL M., US
 (74) Agent: SMART & BIGGAR

(54) Titre : PROTEINE CONTENANT UN DOMAINE DE FOLLISTATINE
 (54) Title: A FOLLISTATIN DOMAIN CONTAINING PROTEIN



(57) Abrégé/Abstract:

The present invention relates to the use of a protein, GASP1, comprising at least one follistatin domain to modulate the level or activity of growth and differentiation factor -8 (GDF-8). More particularly, the invention relates to the use of GASP1 for treating disorders that are related to modulation of the level or activity of GDF-8. The invention is useful for treating muscular diseases and disorders, particularly those in which an increase in muscle tissue would be therapeutically beneficial. The invention is also useful for treating diseases and disorders related to metabolism, adipose tissue, and bone degeneration.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
4 September 2003 (04.09.2003)


PCT

(10) International Publication Number
WO 03/072715 A2

- (51) International Patent Classification⁷: C12N (74) Agent: GARRETT, Arthur, S.; Finnegan, Henderson, Farabown, Garrett & Dunner, LLP, 1300 I Street, NW, Washington, DC 20005-3315 (US).
- (21) International Application Number: PCT/US03/05151
- (22) International Filing Date: 21 February 2003 (21.02.2003) (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/357,845 21 February 2002 (21.02.2002) US
60/434,644 20 December 2002 (20.12.2002) US
- (71) Applicant (*for all designated States except US*): WYETH [US/US]; Five Giralda Farms, Madison, NJ 07940 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): HILL, Jennifer, J. [US/US]; Apt. 16, 157R Summer Street, Somerville, MA 02143 (US). WOLFMAN, Neil, M. [US/US]; 5 Phillips Lane, Dover, MA 02030 (US).
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:
— without international search report and to be republished upon receipt of that report

[Continued on next page]

(54) Title: A FOLLISTATIN DOMAIN CONTAINING PROTEIN



(57) Abstract: The present invention relates to the use of a protein, GASP1, comprising at least one follistatin domain to modulate the level or activity of growth and differentiation factor -8 (GDF-8). More particularly, the invention relates to the use of GASP1 for treating disorders that are related to modulation of the level or activity of GDF-8. The invention is useful for treating muscular diseases and disorders, particularly those in which an increase in muscle tissue would be therapeutically beneficial. The invention is also useful for treating diseases and disorders related to metabolism, adipose tissue, and bone degeneration.

WO 03/072715 A2

WO 03/072715 A2



For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

GASP1: A FOLLISTATIN DOMAIN CONTAINING PROTEIN

[001] This application claims the benefit of U.S. Provisional Application No. 60/357,845, filed February 21, 2002, and U.S. Provisional Application No. 60/434,644, filed December 20, 2002.

Field of the Invention

[002] The present invention relates to the use of proteins comprising at least one follistatin domain to modulate the level or activity of growth and differentiation factor-8 (GDF-8). More particularly, the invention relates to the use of proteins comprising at least one follistatin domain, excluding follistatin itself, for treating disorders that are related to modulation of the level or activity of GDF-8. The invention is useful for treating muscular diseases and disorders, particularly those in which an increase in muscle tissue would be therapeutically beneficial. The invention is also useful for treating diseases and disorders related to metabolism, adipose tissue, and bone degeneration.

Background of the Invention

[003] Growth and differentiation factor-8 (GDF-8), also known as myostatin, is a member of the transforming growth factor-beta (TGF- β) superfamily of structurally related growth factors, all of which possess important physiological growth-regulatory and morphogenetic properties (Kingsley *et al.* (1994) *Genes Dev.*, 8: 133-46; Hoodless *et al.* (1998) *Curr. Topics Microbiol. Immunol.*, 228: 235-72). GDF-8 is a negative regulator of skeletal muscle mass, and there is considerable interest in identifying factors which regulate its biological activity. For example, GDF-8 is highly expressed in the developing and adult skeletal muscle. The GDF-8 null mutation in transgenic mice is characterized by a marked hypertrophy and hyperplasia of the skeletal muscle (McPherron *et al.* (1997) *Nature*, 387: 83-90). Similar increases in skeletal muscle mass are evident in naturally occurring mutations of GDF-8 in cattle (Ashmore *et al.* (1974) *Growth*, 38: 501-507; Swatland and Kieffer (1994) *J. Anim. Sci.*, 38: 752-757; McPherron and Lee (1997) *Proc. Nat. Acad. Sci. U.S.A.*, 94: 12457-12461; and Kambadur *et al.* (1997) *Genome Res.*, 7: 910-915). Recent studies have also shown that muscle

- 2 -

wasting associated with HIV-infection in humans is accompanied by increases in GDF-8 protein expression (Gonzalez-Cadavid *et al.* (1998) *Proc. Natl. Acad. Sci. U.S.A.*, 95: 14938-43). In addition, GDF-8 can modulate the production of muscle-specific enzymes (*e.g.*, creatine kinase) and modulate myoblast cell proliferation (WO 00/43781).

[004] A number of human and animal disorders are associated with loss of or functionally impaired muscle tissue. To date, very few reliable or effective therapies exist for these disorders. However, the terrible symptoms associated with these disorders may be substantially reduced by employing therapies that increase the amount of muscle tissue in patients suffering from the disorders. While not curing the conditions, such therapies would significantly improve the quality of life for these patients and could ameliorate some of the effects of these diseases. Thus, there is a need in the art to identify new therapies that may contribute to an overall increase in muscle tissue in patients suffering from these disorders.

[005] In addition to its growth-regulatory and morphogenetic properties in skeletal muscle, GDF-8 may also be involved in a number of other physiological processes (*e.g.*, glucose homeostasis), as well as abnormal conditions, such as in the development of type 2 diabetes and adipose tissue disorders, such as obesity. For example, GDF-8 modulates preadipocyte differentiation to adipocytes (Kim *et al.* (2001) *B.B.R.C.* 281: 902-906). Thus, modulation of GDF-8 may be useful for treating these diseases, as well.

[006] The GDF-8 protein is synthesized as a precursor protein consisting of an amino-terminal propeptide and a carboxy-terminal mature domain (McPherron and Lee, (1997) *Proc. Nat. Acad. Sci. U.S.A.*, 94: 12457-12461). Before cleavage, the precursor GDF-8 protein forms a homodimer. The amino-terminal propeptide is then cleaved from the mature domain. The cleaved propeptide may remain noncovalently bound to the mature domain dimer, inactivating its biological activity (Miyazono *et al.* (1988) *J. Biol. Chem.*, 263: 6407-6415; Wakefield *et al.* (1988) *J. Biol. Chem.*, 263: 7646-7654; and

- 3 -

Brown *et al.* (1990) *Growth Factors*, 3: 35-43). It is believed that two GDF-8 propeptides bind to the GDF-8 mature dimer (Thies *et al.* (2001) *Growth Factors*, 18: 251-259). Due to this inactivating property, the propeptide is known as the "latency-associated peptide" (LAP), and the complex of mature domain and propeptide is commonly referred to as the "small latent complex" (Gentry and Nash (1990) *Biochemistry*, 29:6851-6857; Derynck *et al.* (1995) *Nature*, 316:701-705; and Massague (1990) *Ann. Rev. Cell Biol.*, 12: 597-641). Other proteins are also known to bind to GDF-8 or structurally related proteins and inhibit their biological activity. Such inhibitory proteins include follistatin (Gamer *et al.* (1999) *Dev. Biol.*, 208: 222-232). The mature domain of GDF-8 is believed to be active as a homodimer when the propeptide is removed.

[007] Clearly, GDF-8 is involved in the regulation of many critical biological processes. Due to its key function in these processes, GDF-8 may be a desirable target for therapeutic intervention. In particular, therapeutic agents that inhibit the activity of GDF-8 may be used to treat human or animal disorders in which an increase in muscle tissue would be therapeutically beneficial.

[008] Known proteins comprising at least one follistatin domain play roles in many biological processes, particularly in the regulation of TGF- β superfamily signaling and the regulation of extracellular matrix-mediated processes such as cell adhesion. Follistatin, follistatin related gene (FLRG, FSRP), and follistatin-related protein (FRP) have all been linked to TGF- β signaling, either through transcriptional regulation by TGF- β (Bartholin *et al.* (2001) *Oncogene*, 20: 5409-5419; Shibamura *et al.* (1993) *Eur. J. Biochem.* 217: 13-19) or by their ability to antagonize TGF- β signaling pathways (Phillips and de Kretser (1998) *Front. Neuroendocrin.*, 19: 287-322; Tsuchida *et al.* (2000) *J. Biol. Chem.*, 275: 40788-40796; Patel *et al.* (1996) *Dev. Biol.*, 178: 327-342; Amthor *et al.* (1996) *Dev. Biol.*, 178: 343-362). Protein names in parentheses are alternative names.

- 4 -

[009] Insulin growth factor binding protein 7 (IGFBP7, mac25), which comprise at least one follistatin domain, binds to insulin and blocks subsequent interaction with the insulin receptor. In addition, IGFBP7 has been shown to bind to activin, a TGF- β family member (Kato (2000) *Mol. Med.*, 6: 126-135).

[010] Agrins and agrin related proteins contain upwards of nine follistatin domains and are secreted from nerve cells to promote the aggregation of acetylcholine receptors and other molecules involved in the formation of synapses. It has been suggested that the follistatin domains may serve to localize growth factors to the synapse (Patthy *et al.* (1993) *Trends Neurosci.*, 16: 76-81).

[011] Osteonectin (SPARC, BM40) and hevin (SC1, mast9, QR1) are closely related proteins that interact with extracellular matrix proteins and regulate cell growth and adhesion (Motamed (1999) *Int. J. Biochem. Cell. Biol.*, 31: 1363-1366; Girard and Springer (1996) *J. Biol. Chem.*, 271: 4511-4517). These proteins comprise at least one follistatin domain.

[012] Other follistatin domain proteins have been described or uncovered from the NCBI database (National Center for Biotechnology Information, Bethesda, Maryland, USA), however their functions are presently unknown. These proteins include U19878 (G01639, very similar to tomoregulin-1), T46914, human GASP1 (GDF-associated serum protein 1; described herein; Figure 7), human GASP2 (WFIKKN; Trexler *et al.* (2001) *Proc. Natl. Acad. Sci. U.S.A.*, 98: 3705-3709; Figure 9), and the proteoglycan family of testican (SPOCK) proteins (Alliel *et al.* (1993) *Eur. J. Biochem.*, 214: 347-350). Amino acid and nucleotide sequences for mouse GASP1 (Figure 6) and mouse GASP2 (Figure 8) were also determined from the Celera database (Rockville, MD). As described herein, the nucleotide sequence of cloned mouse GASP1 matched the predicted Celera sequence, with the exception of some base substitutions in wobble codons that did not change the predicted amino acid sequence (see Figure 13).

Summary of the Invention

[013] Accordingly, the invention relates to proteins, other than follistatin, comprising a unique structural feature, namely, the presence of at least one follistatin domain. Follistatin itself is not encompassed by the invention. The proteins comprising at least one follistatin domain are specifically reactive with a mature GDF-8 protein or a fragment thereof, whether the GDF-8 protein is in monomeric form, a dimeric active form, or complexed in the GDF-8 latent complex. Proteins comprising at least one follistatin domain may bind to an epitope on the mature GDF-8 protein that results in a reduction in one or more of the biological activities associated with GDF-8, relative to a mature GDF-8 protein that is not bound by the same protein.

[014] The present invention provides methods for modulating the effects of GDF-8 on cells. Such methods comprise administering an effective amount of a protein comprising at least one follistatin domain. The present invention also encompasses methods for expressing a protein in a cell by administering a DNA molecule encoding a protein comprising at least one follistatin domain.

[015] According to the invention, proteins comprising at least one follistatin domain may be administered to a patient, in a therapeutically effective dose, to treat or prevent medical conditions in which an increase in muscle tissue would be therapeutically beneficial. Embodiments include treatment of diseases, disorders, and injuries involving cells and tissue that are associated with the production, metabolism, or activity of GDF-8.

[016] Proteins comprising at least one follistatin domain may be prepared in a pharmaceutical preparation. The pharmaceutical preparation may contain other components that aid in the binding of the mature GDF-8 protein or fragments thereof, whether it is in monomeric form, dimeric active form, or complexed in the GDF-8 latent complex.

[017] In addition, proteins comprising at least one follistatin domain may be used as a diagnostic tool to quantitatively or qualitatively detect

- 6 -

mature GDF-8 protein or fragments thereof, whether it is in monomeric form, dimeric active form, or complexed in the GDF-8 latent complex. For example, proteins comprising at least one follistatin domain may be used to detect the presence, absence, or amount of GDF-8 protein in a cell, bodily fluid, tissue, or organism. The presence or amount of mature GDF-8 protein detected may be correlated with one or more of the medical conditions listed herein.

[018] Proteins comprising at least one follistatin domain may be provided in a diagnostic kit to detect mature GDF-8 protein or fragments thereof, whether it is in monomeric form, dimeric active form, or complexed in the GDF-8 latent complex, and help correlate the results with one or more of the medical conditions described herein. Such a kit may comprise at least one protein comprising at least one follistatin domain, whether it is labeled or unlabeled, and at least one agent that bind to this proteins, such as a labeled antibody. The kit may also include the appropriate biological standards and control samples to which one could compare the results of the experimental detection. It may also include buffers or washing solutions and instructions for using the kit. Structural components may be included on which one may carry out the experiment, such as sticks, beads, papers, columns, vials, or gels.

Brief Description of the Figures

[019] Figure 1 shows antibody purification of the GDF-8 complex from wild-type mouse serum. A silver stained reducing gel shows proteins purified from wild type mouse serum using the JA16 monoclonal antibody covalently coupled to agarose beads. A control purification (0) with mock-coupled beads was performed in parallel. Subsequent elutions with buffer (mock elute), a competing peptide, and SDS sample buffer revealed two visible protein bands which were specifically eluted with peptide from the JA16-conjugated beads (indicated by arrows).

[020] Figure 2 shows the identification of mature and unprocessed GDF-8 in affinity purified samples from normal mouse serum. Figure 2A shows a representative MS/MS spectrum of a GDF-8 derived peptide (SEQ ID NO:19) identified from the 12 kDa band visible in the affinity purified sample.

- 7 -

Both N-terminal fragment ions (b ions) and C-terminal fragment ions (y ions) are visible. Notably, the most intense y fragment ions result from fragmentation before the proline residue, a common characteristic of proline containing peptides. Figure 2B shows a western blot probed with a polyclonal antibody that recognizes the mature region of GDF-8, confirming the presence of GDF-8 in the affinity purified samples. Both the mature and unprocessed forms of GDF-8 are visible.

[021] Figure 3 shows the GDF-8 propeptide and follistatin-like related gene (FLRG) bind to circulating GDF-8 isolated from normal mouse serum. Representative MS/MS spectra from GDF-8 propeptide (SEQ ID NO:23) (Figure 3A) and FLRG (SEQ ID NO:30) (Figure 3C) derived peptides identified in the 36 kDa band are shown. Figure 3B shows a western blot of affinity purified GDF-8 complex probed with a polyclonal antibody that specifically recognizes the propeptide region of GDF-8, confirming the mass spectrometric identification of this protein in the GDF-8 complex. Both the clipped propeptide and unprocessed GDF-8 are visible -- at longer exposures, unprocessed GDF-8 can also be seen in the SDS eluted sample. Figure 3D shows a western blot of affinity purified GDF-8 complex probed with a monoclonal antibody to FLRG.

[022] Figure 4 shows results from a thorough analysis of a large scale GDF-8 purification that identified GDF-8 propeptide, FLRG, and a novel protein as the major GDF-8 binding proteins in serum. A silver stained gel was dissected into 13 slices from the peptide eluted sample of both negative control and JA16 immunoprecipitates. The proteins in each slice were digested with trypsin and identified using nanoflow-LC-MS/MS and database searching. Proteins unique to the JA16 sample included only unprocessed and mature GDF-8, GDF-8 propeptide, FLRG, and a novel multidomain protease inhibitor (GDF-associated serum protein 1, GASP1). These proteins were identified from the noted regions of the gel.

[023] Figure 5 shows that a novel multidomain protease inhibitor, GASP1, is bound to GDF-8 in serum. Figures 5A (peptide assigned SEQ ID

- 8 -

NO:31) and 5B (peptide assigned SEQ ID NO:33) show representative MS/MS spectra from two GASP1 peptides, identified in band 3 of the silver stained gel of Figure 4.

[024] Figure 6A shows the predicted nucleotide sequence to mouse GASP1. Figure 6B shows a predicted alternative nucleotide sequence to mouse GASP1. Figure 6C shows the predicted amino acid sequence encoded by the nucleotide sequences shown in Figures 6A and 6B. The protein sequences encoded by the two nucleotide sequences are identical because the nucleotide differences are all in wobble codon positions. The follistatin domain is shown in bold and underlined.

[025] Figure 7A shows the predicted nucleotide sequence of human GASP1. Figure 7B shows the corresponding predicted amino acid sequence. The follistatin domain is shown in bold and underlined. Figure 7C shows the predicted nucleotide sequence of human GASP1 using an alternative start site. Figure 7D shows the corresponding predicted amino acid sequence. The follistatin domain is shown in bold and underlined. The end of the sequence is denoted by the asterisk.

[026] Figure 8A shows the predicted nucleotide sequence to mouse GASP2, while Figure 8B shows the corresponding predicted amino acid sequence. The follistatin domain is shown in bold and underlined.

[027] Figure 9A shows the predicted nucleotide sequence to human GASP2, while Figure 9B shows the corresponding predicted amino acid sequence. The follistatin domain is shown in bold and underlined.

[028] Figure 10 shows that mouse GASP1 is expressed in many adult tissues and during development. The figure shows tissue expression profiles of mouse GASP1. A 551 bp fragment of GASP1 was amplified from normalized first-strand cDNA panels from Clontech (Palo Alto, CA). A portion of glyceraldehyde-3-phosphate dehydrogenase (G3PDH) was amplified as a control. G3PDH expression is known to be high in skeletal muscle and low in testis. The cDNA panels were normalized against β -actin, phospholipase A2, and ribosomal protein S29, in addition to G3PDH.

- 9 -

[029] Figure 11 shows proteins isolated from human serum. Proteins from a JA16 immunoprecipitate or a control sample (0) were eluted in a mock PBS elution, a competing peptide elution, or a SDS elution. The proteins in the indicated regions of the gel were digested with trypsin and analyzed by LS-MS/MS and database searching. The proteins present in the JA16 sample but not in the control sample were mature GDF-8 (band 16), GDF-8 propeptide and FLRG (band 11), and human GASP1 (band 4). Figure 11B shows a western blot of an identical JA16 immunoprecipitate probed with an antibody that recognizes mature GDF-8. Bands corresponding to mature and unprocessed GDF-8 isolated from human serum are visible.

[030] Figure 12 shows representative mass spectra of a peptide derived from GDF-8 and associated proteins isolated from bands 4, 11, and 16 (Figure 11). The peptide sequence and N-terminal (b ions) and C-terminal (y ions) are shown. A complete listing of identified peptides is provided in Table 1. Spectra are shown from a GASP1 peptide (SEQ ID NO:44) (Figure 12A), a FLRG peptide (SEQ ID NO:41) (Figure 12B), a GDF-8 propeptide peptide (SEQ ID NO:24) (Figure 12C), and a mature GDF-8 peptide (SEQ ID NO:13) (Figure 12D).

[031] Figure 13 shows the nucleotide (SEQ ID NO:48) and amino acid (SEQ ID NO:49) sequences of cloned mouse GASP1. The peptides identified by mass spectrometry in JA16 affinity-purified samples are underlined. The end of the sequence is denoted by the asterisk.

[032] Figure 14A shows the domain structure of GASP1. GASP1 has a signal sequence/cleavage site after amino acid 29. In addition, GASP1 contains two Kunitz/BPTI serine protease inhibitor domains, a follistatin domain (including a Kazal serine protease inhibitor motif) and a netrin domain, which may inhibit metalloproteases. Figure 14B shows the phylogenetic tree of GASP1 and GASP2 predicted from the mouse and human genomic sequences. Mouse and human GASP1 are 90% identical. GASP1 and GASP2 are 54% identical.

- 10 -

[033] Figure 15 shows that recombinantly-produced GASP1 binds separately to both GDF-8 and GDF-8 propeptide. (A) JA16 was used to immunoprecipitate GDF-8 from mock- or GASP1-V5-His transfected COS cell conditioned media supplemented with recombinant purified GDF-8 and/or propeptide. Western blots with anti-V5 (top panel), anti-GDF-8 (middle panel), or anti-propeptide polyclonal antibodies were used to determine whether these proteins were present in the immunoprecipitate. (B) Recombinantly-produced GASP1 protein was immunoprecipitated by anti-V5 tag antibodies from mock- or GASP1-V5-His conditioned media supplemented with recombinant purified GDF-8 and/or propeptide. The immunoprecipitate was analyzed by western blotting as in (A).

[034] Figure 16 shows that GASP1 inhibits the biological activity of GDF-8 and the highly related BMP-11, but not activin or TGF- β . Various dilutions of conditioned media from mock (open circles) or GASP1-V5-His (filled squares) transfectants were incubated with (A) 10 ng/ml GDF-8, (B) 10 ng/ml BMP-11, (C) 10 ng/ml activin, or (D) 0.5 ng/ml TGF- β . These samples were then subjected to a luciferase reporter activity assay in A204 (A-C) or RD (D) cells to determine the activity of the added growth factors. Luciferase activity is shown in relative luciferase units. The activity resulting from each of the growth factors alone is shown by the filled diamonds and short dashed line. Without addition of any growth factor, the background activity in the assay is low, as shown by the long dashed line with no symbols.

[035] Figure 17 shows the potency of GASP1 inhibition of GDF-8. Purified GASP1 was tested for its ability to inhibit 20 ng/ml of myostatin in the (CAGA)₁₂ (SEQ ID NO:53) luciferase reporter assay in RD cells (filled squares). The activity resulting from GDF-8 alone is shown by the filled diamonds and short dashed line. The activity present when no growth factors are added is shown by the long dashed line.

Definitions

[036] The term "**follicle-statin domain**" refers to an amino acid domain or a nucleotide domain encoding for an amino acid domain, characterized by

- 11 -

cysteine rich repeats. A follistatin domain typically encompasses a 65-90 amino acid span and contains 10 conserved cysteine residues and a region similar to Kazal serine protease inhibitor domains. In general, the loop regions between the cysteine residues exhibit sequence variability in follistatin domains, but some conservation is evident. The loop between the fourth and fifth cysteines is usually small, containing only 1 or 2 amino acids. The amino acids in the loop between the seventh and eighth cysteines are generally the most highly conserved containing a consensus sequence of (G,A)-(S,N)-(S,N,T)-(D,N)-(G,N) followed by a (T,S)-Y motif. The region between the ninth and tenth cysteines generally contains a motif containing two hydrophobic residues (specifically V, I, or L) separated by another amino acid.

[037] The term **“protein comprising at least one follistatin domain”** refers to proteins comprising at least one, but possibly more than one follistatin domain. The term also refers to any variants of such proteins (including fragments; proteins with substitution, addition or deletion mutations; and fusion proteins) that maintain the known biological activities associated with the native proteins, especially those pertaining to GDF-8 binding activity, including sequences that have been modified with conservative or non-conservative changes to the amino acid sequence. These proteins may be derived from any source, natural or synthetic. The protein may be human or derived from animal sources, including bovine, chicken, murine, rat, porcine, ovine, turkey, baboon, and fish. Follistatin itself is not encompassed by the invention.

[038] The terms **“GDF-8”** or **“GDF-8 protein”** refer to a specific growth and differentiation factor. The terms include the full length unprocessed precursor form of the protein, as well as the mature and propeptide forms resulting from post-translational cleavage. The terms also refer to any fragments of GDF-8 that maintain the known biological activities associated with the protein, including sequences that have been modified with conservative or non-conservative changes to the amino acid sequence. These GDF-8 molecules may be derived from any source, natural or

- 12 -

synthetic. The protein may be human or derived from animal sources, including bovine, chicken, murine, rat, porcine, ovine, turkey, baboon, and fish. Various GDF-8 molecules have been described in McPherron *et al.* (1997) *Proc. Natl. Acad. Sci. USA*, 94: 12457-12461.

[039] “**Mature GDF-8**” refers to the protein that is cleaved from the carboxy-terminal domain of the GDF-8 precursor protein. The mature GDF-8 may be present as a monomer, homodimer, or in a GDF-8 latent complex. Depending on the *in vivo* or *in vitro* conditions, mature GDF-8 may establish an equilibrium between any or all of these different forms. It is believed to be biologically active as homodimer. In its biologically active form, the mature GDF-8 is also referred to as “**active GDF-8**.”

[040] “**GDF-8 propeptide**” refers to the polypeptide that is cleaved from the amino-terminal domain of the GDF-8 precursor protein. The GDF-8 propeptide is capable of binding to the propeptide binding domain on the mature GDF-8.

[041] “**GDF-8 latent complex**” refers to the complex of proteins formed between the mature GDF-8 homodimer and the GDF-8 propeptide. It is believed that two GDF-8 propeptides associate with the two molecules of mature GDF-8 in the homodimer to form an inactive tetrameric complex. The latent complex may include other GDF inhibitors in place of or in addition to one or more of the GDF-8 propeptides.

[042] The phrase “**GDF-8 activity**” refers to one or more of physiologically growth-regulatory or morphogenetic activities associated with active GDF-8 protein. For example, active GDF-8 is a negative regulator of skeletal muscle. Active GDF-8 can also modulate the production of muscle-specific enzymes (*e.g.*, creatine kinase), stimulate myoblast cell proliferation, and modulate preadipocyte differentiation to adipocytes. GDF-8 is also believed to increase sensitivity to insulin and glucose uptake in peripheral tissues, particularly in skeletal muscle or adipocytes. Accordingly, GDF-8 biological activities include but are not limited to inhibition of muscle formation, inhibition of muscle cell growth, inhibition of muscle development, decrease in

- 13 -

muscle mass, regulation of muscle-specific enzymes, inhibition of myoblast cell proliferation, modulation of preadipocyte differentiation to adipocytes, increasing sensitivity to insulin, regulations of glucose uptake, glucose hemostasis, and modulate neuronal cell development and maintenance.

[043] The terms “**isolated**” or “**purified**” refer to a molecule that is substantially free of its natural environment. For instance, an isolated protein is substantially free of cellular material or other contaminating proteins from the cell or tissue source from which it is derived. The phrase “**substantially free of cellular material**” refers to preparations where the isolated protein is at least 70% to 80% (w/w) pure, at least 80%-89% (w/w) pure, at least 90-95% pure, or at least 96%, 97%, 98%, 99% or 100% (w/w) pure.

[044] The term “**LC-MS/MS**” refers to liquid chromatography in line with a mass spectrometer programmed to isolate a molecular ion of particular mass/charge ratio, fragment this ion, and record the mass/charge ratio of the fragment ions. When analyzing peptide samples this technique allows upstream separation of complex samples through liquid chromatography, followed by the recording of fragment ion masses and subsequent determination of the peptide sequence.

[045] The term “**MS/MS**” refers to the process of using a mass spectrometer to isolate a molecular ion of a particular mass/charge ratio, fragment this ion, and record the mass/charge ratio of the resulting fragment ions. The fragment ions provide information about the sequence of a peptide.

[046] The term “**treating**” or “**treatment**” refers to both therapeutic treatment and prophylactic or preventative measures. Those in need of treatment may include individuals already having a particular medical disorder as well as those who may ultimately acquire the disorder (*i.e.*, those needing preventative measures). The term treatment includes both measures that address the underlying cause of a disorder and measures that reduce symptoms of a medical disorder without necessarily affecting its cause. Thus, improvement of quality of life and amelioration of symptoms are considered treatment, as are measures that counteract the cause of a disorder.

- 14 -

[047] The term “**medical disorder**” refers to disorders of muscle, bone, or glucose homeostasis, and include disorders associated with GDF-8 and/or other members of the TGF- β superfamily (e.g., BMP-11). Examples of such disorders include, but are not limited to, metabolic diseases and disorders such as insulin-dependent (type 1) diabetes mellitus, noninsulin-dependent (type 2) diabetes mellitus, hyperglycemia, impaired glucose tolerance, metabolic syndrome (e.g., syndrome X), and insulin resistance induced by trauma (e.g., burns or nitrogen imbalance), and adipose tissue disorders (e.g., obesity); muscle and neuromuscular disorders such as muscular dystrophy (including but not limited to severe or benign X-linked muscular dystrophy, limb-girdle dystrophy, facioscapulohumeral dystrophy, myotonic dystrophy, distal muscular dystrophy, progressive dystrophic ophthalmoplegia, oculopharyngeal dystrophy, Duchenne’s muscular dystrophy, and Fakuyama-type congenital muscular dystrophy); amyotrophic lateral sclerosis (ALS); muscle atrophy; organ atrophy; frailty; carpal tunnel syndrome; congestive obstructive pulmonary disease; congenital myopathy; myotonia congenital; familial periodic paralysis; paroxysmal myoglobinuria; myasthenia gravis; Eaton-Lambert syndrome; secondary myasthenia; denervation atrophy; paroxymal muscle atrophy; and sarcopenia, cachexia and other muscle wasting syndromes. Other examples include osteoporosis, especially in the elderly and/or postmenopausal women; glucocorticoid-induced osteoporosis; osteopenia; osteoarthritis; osteoporosis-related fractures; and traumatic or chronic injury to muscle tissue. Yet further examples include low bone mass due to chronic glucocorticoid therapy, premature gonadal failure, androgen suppression, vitamin D deficiency, secondary hyperparathyroidism, nutritional deficiencies, and anorexia nervosa.

[048] The term “**increase in mass**” refers to the presence of a greater amount of muscle after treatment with proteins comprising at least one follistatin domain relative to the amount of muscle mass present before the treatment.

- 15 -

[049] The term “**therapeutic benefit**” refers to an improvement of symptoms of a disorder, a slowing of the progression of a disorder, or a cessation in the progression of a disorder. The therapeutic benefit is determined by comparing an aspect of a disorder, such as the amount of muscle mass, before and after at least one protein comprising at least one follistatin domain is administered.

[050] The term “**modulating**” refers to varying a property of a protein by increasing, decreasing, or inhibiting the activity, behavior, or amount of the protein. For example, proteins comprising at least one follistatin domain may modulate GDF-8 by inhibiting its activity.

[051] The term “**stabilizing modification**” is any modification known in the art or set forth herein capable of stabilizing a protein, enhancing the in vitro half life of a protein, enhancing circulatory half life of a protein and/or reducing proteolytic degradation of a protein. Such stabilizing modifications include but are not limited to fusion proteins (including, for example, fusion proteins comprising a protein comprising at least one follistatin domain and a second protein), modification of a glycosylation site (including, for example, addition of a glycosylation site to a protein comprising at least one follistatin domain), and modification of carbohydrate moiety (including, for example, removal of carbohydrate moieties from a protein comprising at least one follistatin domain). In the case of a stabilizing modification which comprises a fusion protein (e.g., such that a second protein is fused to a protein comprising at least one follistatin domain), the second protein may be referred to as a “**stabilizer portion**” or “**stabilizer protein.**” For example, a protein a human protein comprising at least one follistatin domain may be fused with an IgG molecule, wherein IgG acts as the stabilizer protein or stabilizer portion. As used herein, in addition to referring to a second protein of a fusion protein, a “**stabilizer portion**” also includes nonproteinaceous modifications such as a carbohydrate moiety, or nonproteinaceous polymer.

[052] The term “**Fc region of an IgG molecule**” refers to the Fc domain of an immunoglobulin of the isotype IgG, as is well known to those

- 16 -

skilled in the art. The Fc region of an IgG molecule is that portion of IgG molecule (IgG1, IgG2, IgG3, and IgG4) that is responsible for increasing the in vivo serum half-life of the IgG molecule.

[053] **"In vitro half life"** refers to the stability of a protein measured outside the context of a living organism. Assays to measure in vitro half life are well known in the art and include but are not limited to SDS-PAGE, ELISA, cell-based assays, pulse-chase, western blotting, northern blotting, etc. These and other useful assays are well known in the art.

[054] **"In vivo half life"** refers to the stability of a protein in an organism. In vivo half life may be measured by a number of methods known in the art including but not limited to in vivo serum half life, circulatory half life, and assays set forth in the examples herein.

[055] **"In vivo serum half life"** refers to the half-life of a protein circulating in the blood of an organism. Methods known in the art may be used to measure in vivo serum half life. For example, radioactive protein can be administered to an animal and the amount of labeled protein in the serum can be monitored over time.

[056] To assist in the identification of the sequences listed in the specification and figures, the following table is provided, which lists the SEQ ID NO, the figure location, and a description of the sequence.

SEQ ID NO:	REFERENCE	DESCRIPTION
1	Figure 6A	predicted mouse GASP1 nucleotide sequence
2	Figure 6B	predicted mouse GASP1 alternative nucleotide sequence
3	Figure 6C	predicted mouse GASP1 amino acid sequence encoded by SEQ ID NOS:1 and 2
4	Figure 7A	predicted human GASP1 nucleotide sequence
5	Figure 7B	predicted human GASP1 amino acid sequence encoded by SEQ ID NO:4
6	Figure 7C	predicted human GASP1 nucleotide sequence, alternative start site
7	Figure 7D	predicted human GASP1 amino acid sequence, alternative start site encoded by SEQ ID NO:6
8	Figure 8A	predicted mouse GASP2 nucleotide sequence
9	Figure 8B	predicted mouse GASP2 amino acid sequence encoded by SEQ ID NO:8
10	Figure 9A	predicted human GASP2 nucleotide sequence
11	Figure 9B	predicted human GASP2 amino acid sequence encoded by SEQ ID NO:10
12	Example 2	competing peptide
13-20	Table 1, Examples 5, 6	mouse GDF-8 peptides
21-27	Table 1, Examples 5, 6	mouse GDF-8 propeptide peptides
28-30	Table 1, Example 5	mouse FLRG peptides
31-35	Table 1, Examples 5, 7	mouse GASP1 peptides
36-37	Table 1, Example 8	human GDF-8 peptides
38-39	Table 1, Example 8	human GDF-8 propeptide peptides
40-42	Table 1, Example 8	human FLRG peptides
43-45	Table 1, Example 8	human GASP1 peptides
46	Example 7	forward primer
47	Example 7	reverse primer
48	Figure 13	cloned mouse GASP1 nucleotide sequence
49	Figure 13	cloned mouse GASP1 amino acid sequence encoded by SEQ ID NO:48
50	Example 9	forward primer
51	Example 9	reverse primer
52	Example 9	illustrative N-terminal peptide sequence
53	Example 11	synthetic oligonucleotide

Detailed Description of the Invention**Proteins Comprising At Least One Follistatin Domain**

[057] The present invention relates to proteins, other than follistatin, having a unique structural feature, namely, that they comprise at least one follistatin domain. Follistatin itself is not encompassed by the invention. It is believed that proteins containing at least one follistatin domain will bind and inhibit GDF-8. Examples of proteins having at least one follistatin domain include, but are not limited to follistatin-like related gene (FLRG), FRP (flik, tsc 36), agrins, osteonectin (SPARC, BM40), hevin (SC1, mast9, QR1), IGFBP7 (mac25), and U19878. GASP1, comprising the nucleotide and amino acid sequences provided in Figures 6 and 7, and GASP2, comprising the nucleotide and amino acid sequences provided in Figures 8 and 9, are other examples of proteins comprising at least one follistatin domain.

[058] A follistatin domain, as stated above, is defined as an amino acid domain or a nucleotide domain encoding for an amino acid domain, characterized by cysteine rich repeats. A follistatin domain typically encompasses a 65-90 amino acid span and contains 10 conserved cysteine residues and a region similar to Kazal serine protease inhibitor domains. In general, the loop regions between the cysteine residues exhibit sequence variability in follistatin domains, but some conservation is evident. The loop between the fourth and fifth cysteines is usually small, containing only 1 or 2 amino acids. The amino acids in the loop between the seventh and eighth cysteines are generally the most highly conserved containing a consensus sequence of (G,A)-(S,N)-(S,N,T)-(D,N)-(G,N) followed by a (T,S)-Y motif. The region between the ninth and tenth cysteines generally contains a motif containing two hydrophobic residues (specifically V, I, or L) separated by another amino acid.

[059] Proteins comprising at least one follistatin domain, which may bind GDF-8, may be isolated using a variety of methods. For example, one may use affinity purification using GDF-8, as exemplified in the present

- 19 -

invention. In addition, one may use a low stringency screening of a cDNA library, or use degenerate PCR techniques using a probe directed toward a follistatin domain. As more genomic data becomes available, similarity searching using a number of sequence profiling and analysis programs, such as MotifSearch (Genetics Computer Group, Madison, WI), ProfileSearch (GCG), and BLAST (NCBI) could be used to find novel proteins containing significant homology with known follistatin domains.

[060] One of skill in the art will recognize that both GDF-8 or proteins comprising at least one follistatin domain may contain any number of conservative changes to their respective amino acid sequences without altering their biological properties. Such conservative amino acid modifications are based on the relative similarity of the amino acid side-chain substituents, for example, their hydrophobicity, hydrophilicity, charge, size, and the like. Exemplary conservative substitutions which take various of the foregoing characteristics into consideration are well known to those of skill in the art and include: arginine and lysine; glutamate and aspartate; serine and threonine; glutamine and asparagine; and valine, leucine, and isoleucine. Furthermore, proteins comprising at least one follistatin domain may be used to generate functional fragments comprising at least one follistatin domain. It is expected that such fragments would bind and inhibit GDF-8. In an embodiment of the invention, proteins comprising at least one follistatin domain specifically bind to mature GDF-8 or a fragment thereof, whether it is in monomeric form, active dimer form, or complexed in a GDF-8 latent complex, with an affinity of between 0.001 and 100 nM, or between 0.01 and 10 nM, or between 0.1 and 1 nM.

Nucleotide and Protein Sequences

[061] While not always necessary, if desired, one of ordinary skill in the art may determine the amino acid or nucleic acid sequences of a novel proteins comprising at least one follistatin domain. For example, the present

- 20 -

invention provides the amino acid and nucleotide sequences for GASP1 and GASP2, as shown in Figures 6-9.

[062] The present invention also include variants, homologues, and fragments of such nucleic and amino acid sequences. For example, the nucleic or amino acid sequence may comprise a sequence at least 70% to 79% identical to the nucleic or amino acid sequence of the native protein, or at least 80% to 89% identical, or at least 90% to 95% identical, or at least 96% to 100% identical. One of skill in the art will recognize that the region that binds GDF-8 can tolerate less sequence variation than the other portions of the protein not involved in binding. Thus, these non-binding regions of the protein may contain substantial variations without significantly altering the binding properties of the protein. However, one of skill in the art will also recognize that many changes can be made to specifically increase the affinity of the protein for its target. Such affinity-increasing changes are typically determined empirically by altering the protein, which may be in the binding region, and testing the ability to bind GDF-8 or the strength of the binding. All such alterations, whether within or outside the binding region, are included in the scope of the present invention.

[063] Relative sequence similarity or identity may be determined using the "Best Fit" or "Gap" programs of the Sequence Analysis Software Package™ (Version 10; Genetics Computer Group, Inc., University of Wisconsin Biotechnology Center, Madison, WI). "Gap" utilizes the algorithm of Needleman and Wunsch (Needleman and Wunsch, 1970) to find the alignment of two sequences that maximizes the number of matches and minimizes the number of gaps. "BestFit" performs an optimal alignment of the best segment of similarity between two sequences. Optimal alignments are found by inserting gaps to maximize the number of matches using the local homology algorithm of Smith and Waterman (Smith and Waterman, 1981; Smith, *et al.*, 1983).

[064] The Sequence Analysis Software Package described above contains a number of other useful sequence analysis tools for identifying

- 21 -

homologues of the presently disclosed nucleotide and amino acid sequences. For example, the "BLAST" program (Altschul, *et al.*, 1990) searches for sequences similar to a query sequence (either peptide or nucleic acid) in a specified database (e.g., sequence databases maintained at the NCBI; "FastA" (Lipman and Pearson, 1985; see *also* Pearson and Lipman, 1988; Pearson, *et al.*, 1990) performs a Pearson and Lipman search for similarity between a query sequence and a group of sequences of the same type (nucleic acid or protein); "TfastA" performs a Pearson and Lipman search for similarity between a protein query sequence and any group of nucleotide sequences (it translates the nucleotide sequences in all six reading frames before performing the comparison); "FastX" performs a Pearson and Lipman search for similarity between a nucleotide query sequence and a group of protein sequences, taking frameshifts into account. "TfastX" performs a Pearson and Lipman search for similarity between a protein query sequence and any group of nucleotide sequences, taking frameshifts into account (it translates both strands of the nucleic sequence before performing the comparison).

Modified Proteins

[065] The invention encompasses fragments of proteins comprising at least one follistatin domain. Such fragments will likely include all or a part of the follistatin domain. Fragments may include all, a part, or none of the sequences between the follistatin domain and the N-terminus and/or between the follistatin domain and the C-terminus.

[066] It is understood by one of ordinary skill in the art that certain amino acids may be substituted for other amino acids in a protein structure without adversely affecting the activity of the protein, e.g., binding characteristics of a protein comprising at least one follistatin domain. It is thus contemplated by the inventors that various changes may be made in the amino acid sequences of proteins comprising at least one follistatin domain, or DNA sequences encoding the proteins, without appreciable loss of their

- 22 -

biological utility or activity. Such changes may include deletions, insertions, truncations, substitutions, fusions, shuffling of motif sequences, and the like.

[067] In making such changes, the hydrophobic index of amino acids may be considered. The importance of the hydrophobic amino acid index in conferring interactive biological function on a protein is generally understood in the art (Kyte and Doolittle (1982) *J. Mol. Biol.*, 157: 105-132). It is accepted that the relative hydrophobic character of the amino acid contributes to the secondary structure of the resultant protein, which in turn defines the interaction of the protein with other molecules, for example, enzymes, substrates, receptors, DNA, antibodies, antigens, and the like.

[068] Each amino acid has been assigned a hydrophobic index on the basis of its hydrophobicity and charge characteristics (Kyte and Doolittle, 1982); these are isoleucine (+4.5), valine (+4.2), leucine (+3.8), phenylalanine (+2.8), cysteine/cystine (+2.5), methionine (+1.9), alanine (+1.8), glycine (-0.4), threonine (-0.7), serine (-0.8), tryptophan (-0.9), tyrosine (-1.3), proline (-1.6), histidine (-3.2), glutamate (-3.5), glutamine (-3.5), aspartate (-3.5), asparagine (-3.5), lysine (-3.9), and arginine (-4.5). In making such changes, the substitution of amino acids whose hydrophobic indices may be within ± 2 , within ± 1 , and within ± 0.5 .

[069] It is also understood in the art that the substitution of like amino acids can be made effectively on the basis of hydrophilicity. U.S. Patent 4,554,101 states that the greatest local average hydrophilicity of a protein, as governed by the hydrophilicity of its adjacent amino acids, correlates with a biological property of the protein.

[070] As detailed in U.S. Patent 4,554,101, the following hydrophilicity values have been assigned to amino acid residues: arginine (+3.0), lysine (+3.0), aspartate (+3.0 \pm 1), glutamate (+3.0 \pm 1), serine (+0.3), asparagine (+0.2), glutamine (+0.2), glycine (0), threonine (-0.4), proline (-0.5 \pm 1), alanine (-0.5), histidine (-0.5), cysteine (-1.0), methionine (-1.3), valine (-1.5), leucine (-1.8), isoleucine (-1.8), tyrosine (-2.3), phenylalanine (-2.5), and tryptophan

- 23 -

(-3.4). In making such changes, the substitution of amino acids whose hydrophilicity values may be within ± 2 , within ± 1 , and within ± 0.5 .

[071] The modifications may be conservative such that the structure or biological function of the protein is not affected by the change. Such conservative amino acid modifications are based on the relative similarity of the amino acid side-chain substituents, for example, their hydrophobicity, hydrophilicity, charge, size, and the like. Exemplary conservative substitutions which take various of the foregoing characteristics into consideration are well known to those of skill in the art and include: arginine and lysine; glutamate and aspartate; serine and threonine; glutamine and asparagine; and valine, leucine, and isoleucine. The amino acid sequence of proteins comprising at least one follistatin domain may be modified to have any number of conservative changes, so long as the binding of the protein to its target antigen is not adversely affected. Such changes may be introduced inside or outside of the binding portion of the protein comprising at least one follistatin domain. For example, changes introduced inside of the antigen binding portion of the protein may be designed to increase the affinity of the protein for its target.

Stabilizing Modification

[072] Stabilizing modifications are capable of stabilizing a protein, enhancing the *in vitro* and/or *in vivo* half life of a protein, enhancing circulatory half life of a protein and/or reducing proteolytic degradation of a protein. Such stabilizing modifications include but are not limited to fusion proteins, modification of a glycosylation site, and modification of carbohydrate moiety. A stabilizer protein may be any protein which enhances the overall stability of the modified GDF propeptide. As will be recognized by one of ordinary skill in the art, such fusion protein may optionally comprise a linker peptide between the propeptide portion and the stabilizing portion. As is well known in the art, fusion proteins are prepared such that the second protein is fused in frame with the first protein such that the resulting translated protein comprises both

- 24 -

the first and second proteins. For example, in the present invention, a fusion protein may be prepared such that a protein comprising at least one follistatin domain is fused to a second protein (e.g. a stabilizer protein portion.) Such fusion protein is prepared such that the resulting translated protein contains both the propeptide portion and the stabilizer portion.

[073] Proteins comprising at least one follistatin domain can be glycosylated or linked to albumin or a nonproteineous polymer. For instance, proteins comprising at least one follistatin domain may be linked to one of a variety of nonproteinaceous polymers, e.g., polyethylene glycol, polypropylene glycol, or polyoxyalkylenes, in the manner set forth in U.S. Patent Numbers 4,640,835; 4,496,689; 4,301,144; 4,670,417; 4,791,192 or 4,179,337. Proteins are chemically modified by covalent conjugation to a polymer to increase their circulating half-life, for example. Polymers, and methods to attach them to peptides, are also shown in U.S. Pat. Nos. 4,766,106; 4,179,337; 4,495,285; and 4,609,546.

[074] Proteins comprising at least one follistatin domain may be pegylated. Pegylation is a process whereby polyethylene glycol (PEG) is attached to a protein in order to extend the half-life of the protein in the body. Pegylation of proteins comprising at least one follistatin domain may decrease the dose or frequency of administration of the proteins needed for an optimal inhibition of GDF-8. Reviews of the technique are provided in Bhadra *et al.* (2002) *Pharmazie*, 57: 5-29, and in Harris *et al.* (2001) *Clin. Pharmacokinet.*, 40: 539-551.

[075] Proteins comprising at least one follistatin domain can be linked to an Fc region of an IgG molecule. Proteins comprising at least one follistatin domain may be fused adjacent to the Fc region of the IgG molecule, or attached to the Fc region of the IgG molecule via a linker peptide. Use of such linker peptides is well known in the protein biochemistry art. The Fc region may be derived from IgG1 or IgG4, for example.

[076] Proteins comprising at least one follistatin domain may be modified to have an altered glycosylation pattern (*i.e.*, altered from the original

- 25 -

or native glycosylation pattern). As used herein, "altered" means having one or more carbohydrate moieties deleted, and/or having one or more glycosylation sites added to the original protein.

[077] Glycosylation of proteins is typically either N-linked or O-linked. N-linked refers to the attachment of the carbohydrate moiety to the side chain of an asparagine residue. The tripeptide sequences asparagine-X-serine and asparagine-X-threonine, where X is any amino acid except proline, are the recognition sequences for enzymatic attachment of the carbohydrate moiety to the asparagine side chain. Thus, the presence of either of these tripeptide sequences in a polypeptide creates a potential glycosylation site. O-linked glycosylation refers to the attachment of one of the sugars N-acylgalactosamine, galactose, or xylose to a hydroxyamino acid, most commonly serine or threonine, although 5-hydroxyproline or 5-hydroxylysine may also be used.

[078] Addition of glycosylation sites to proteins comprising at least one follistatin domain is conveniently accomplished by altering the amino acid sequence such that it contains one or more of the above-described tripeptide sequences (for N-linked glycosylation sites). The alteration may also be made by the addition of, or substitution by, one or more serine or threonine residues to the sequence of the original protein (for O-linked glycosylation sites). For ease, the protein amino acid sequence may be altered through changes at the DNA level.

[079] Another means of increasing the number of carbohydrate moieties on proteins is by chemical or enzymatic coupling of glycosides to the amino acid residues of the protein. These procedures are advantageous in that they do not require production of the GDF peptide inhibitor in a host cell that has glycosylation capabilities for N- or O-linked glycosylation. Depending on the coupling mode used, the sugars may be attached to (a) arginine and histidine, (b) free carboxyl groups, (c) free sulfhydryl groups such as those of cysteine, (d) free hydroxyl groups such as those of serine, threonine, or hydroxyproline, (e) aromatic residues such as those of phenylalanine,

- 26 -

tyrosine, or tryptophan, or (f) the amide group of glutamine. These methods are described in WO 87/05330, and in Aplin and Wriston (1981) *CRC Crit. Rev. Biochem.*, 22: 259-306.

[080] Removal of any carbohydrate moieties present on proteins comprising at least one follistatin domain may be accomplished chemically or enzymatically. Chemical deglycosylation requires exposure of the protein to trifluoromethanesulfonic acid, or an equivalent compound. This treatment results in the cleavage of most or all sugars except the linking sugar (N-acetylglucosamine or N-acetylgalactosamine), while leaving the amino acid sequence intact.

[081] Chemical deglycosylation is described by Hakimuddin *et al.* (1987) *Arch. Biochem. Biophys.*, 259: 52; and Edge *et al.* (1981) *Anal. Biochem.*, 118: 131. Enzymatic cleavage of carbohydrate moieties on GDF peptide inhibitors can be achieved by the use of a variety of endo- and exo-glycosidases as described by Thotakura *et al.* (1987) *Meth. Enzymol.*, 138: 350.

[082] Proteins comprising at least one follistatin domain may be linked to the protein albumin or a derivative of albumin. Methods for linking proteins and polypeptides to albumin or albumin derivatives are well known in the art. See, for example, U.S. Patent No. 5,116,944.

Pharmaceutical Compositions

[083] The present invention provides compositions containing proteins comprising at least one follistatin domain. Such compositions may be suitable for pharmaceutical use and administration to patients. The compositions typically contain one or more proteins comprising at least one follistatin domain and a pharmaceutically acceptable excipient. As used herein, the phrase "pharmaceutically acceptable excipient" includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like, that are compatible with pharmaceutical administration. The use of such media and agents for

- 27 -

pharmaceutically active substances is well known in the art. The compositions may also contain other active compounds providing supplemental, additional, or enhanced therapeutic functions. The pharmaceutical compositions may also be included in a container, pack, or dispenser together with instructions for administration.

[084] A pharmaceutical composition of the invention is formulated to be compatible with its intended route of administration. Methods to accomplish the administration are known to those of ordinary skill in the art. The administration may, for example, be intravenous, intramuscular, or subcutaneous.

[085] Solutions or suspensions used for subcutaneous application typically include one or more of the following components: a sterile diluent such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetra acetic acid; buffers such as acetates, citrates or phosphates; and agents for the adjustment of tonicity such as sodium chloride or dextrose. The pH can be adjusted with acids or bases, such as hydrochloric acid or sodium hydroxide. Such preparations may be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic.

[086] Pharmaceutical compositions suitable for injection include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. For intravenous administration, suitable carriers include physiological saline, bacteriostatic water, Cremophor EL™ (BASF, Parsippany, NJ) or phosphate buffered saline (PBS). In all cases, the composition must be sterile and should be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol

- 28 -

(for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Prevention of the action of microorganisms can be achieved by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic acid, thimerosal, and the like. In many cases, one may include isotonic agents, for example, sugars, polyalcohols such as manitol, sorbitol, sodium chloride in the composition. Prolonged absorption of the injectable compositions can be brought about by including in the composition an agent which delays absorption, for example, aluminum monostearate and gelatin.

[087] In one embodiment, proteins comprising at least one follistatin domain are prepared with carriers that will protect the compound against rapid elimination from the body, such as a controlled release formulation, including implants and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Methods for preparation of such formulations will be apparent to those skilled in the art. The materials can also be obtained commercially from Alza Corporation and Nova Pharmaceuticals, Inc. Liposomal suspensions containing proteins comprising at least one follistatin domain can also be used as pharmaceutically acceptable carriers. These can be prepared according to methods known to those skilled in the art, for example, as described in U.S. Patent No. 4,522,811.

[088] Therapeutically useful agents, such as growth factors (e.g., BMPs, TGF- β , FGF, IGF), cytokines (e.g., interleukins and CDFs), antibiotics, and any other therapeutic agent beneficial for the condition being treated may optionally be included in or administered simultaneously or sequentially with, proteins comprising at least one follistatin domain.

- 29 -

[089] It is especially advantageous to formulate compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the subject to be treated; each unit containing a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the invention are dictated by and directly dependent on the unique characteristics of the active compound and the particular therapeutic effect to be achieved, and the limitations inherent in the art of compounding such an active compound for the treatment of individuals.

Treatment Indications

[090] Proteins comprising at least one follistatin domain are useful to prevent, diagnose, or treat various medical disorders in humans or animals. Accordingly, the present invention provides a method for treating diseases and disorders related to muscle cells and tissue, by administering to a subject a composition comprising at least one protein comprising at least one follistatin domain in an amount sufficient to ameliorate the symptoms of the disease. Such disorders include muscular dystrophies, including, but not limited to severe or benign X-linked muscular dystrophy, limb-girdle dystrophy, facioscapulohumeral dystrophy, myotonic dystrophy, distal muscular dystrophy, progressive dystrophic ophthalmoplegia, oculopharyngeal dystrophy, Duchenne's muscular dystrophy, and Fukuyama-type congenital muscular dystrophy); amyotrophic lateral sclerosis (ALS); muscle atrophy; organ atrophy; frailty; carpal tunnel syndrome; congestive obstructive pulmonary disease; congenital myopathy; myotonia congenital; familial periodic paralysis; paroxysmal myoglobinuria; myasthenia gravis; Eaton-Lambert syndrome; secondary myasthenia; denervation atrophy; paroxymal muscle atrophy; and sarcopenia, cachexia and other muscle

- 30 -

wasting syndromes. The invention also relates to traumatic or chronic injury to muscle tissue.

[091] In addition to providing therapy for muscle diseases and disorders, the present invention also provides for methods for preventing or treating metabolic diseases or disorders resulting from abnormal glucose homeostasis. Such diseases or disorders include metabolic diseases and disorders (such as insulin-dependent (type 1) diabetes mellitus, noninsulin-dependent (type 2) diabetes mellitus), hyperglycemia, impaired glucose tolerance, metabolic syndrome (e.g., syndrome X), obesity and insulin resistance induced by trauma (e.g., burns or nitrogen imbalance), adipose tissue disorders (such as obesity), or bone degenerative diseases (such as osteoporosis, especially in the elderly and/or postmenopausal women; glucocorticoid-induced osteoporosis; osteopenia; osteoarthritis; and osteoporosis-related fractures). Yet further examples include low bone mass due to chronic glucocorticoid therapy, premature gonadal failure, androgen suppression, vitamin D deficiency, secondary hyperparathyroidism, nutritional deficiencies, and anorexia nervosa.

[092] Normal glucose homeostasis requires the finely tuned orchestration of insulin secretion by pancreatic beta cells in response to subtle changes in blood glucose levels. One of the fundamental actions of insulin is to stimulate uptake of glucose from the blood into tissues, especially muscle and fat.

[093] Accordingly, the present invention provides a method for treating diabetes mellitus and related disorders, such as obesity or hyperglycemia, by administering to a subject a composition comprising at least one protein comprising at least one follistatin domain in an amount sufficient to ameliorate the symptoms of the disease. Type 2 or noninsulin-dependent diabetes mellitus (NIDDM), in particular, is characterized by a triad of (1) resistance to insulin action on glucose uptake in peripheral tissues, especially skeletal muscle and adipocytes, (2) impaired insulin action to inhibit hepatic glucose production, and (3) dysregulated insulin secretion (DeFronzo (1997) *Diabetes*

Rev. 5: 177-269). Therefore, subjects suffering from type 2 diabetes can be treated according to the present invention by administration of protein comprising at least one follistatin domain, which increases sensitivity to insulin and glucose uptake by cells.

[094] Similarly, other diseases and metabolic disorders characterized by insulin dysfunction (e.g., resistance, inactivity, or deficiency) and/or insufficient glucose transport into cells also can be treated according to the present invention by administration of a protein comprising at least one follistatin domain, which increases sensitivity to insulin and glucose uptake by cells.

Methods of Treatment Using Proteins

[095] Proteins comprising at least one follistatin domain may be used to inhibit or reduce one or more activities associated with the GDF-8 protein (whether in monomeric form, dimeric active form, or complexed in a GDF-8 latent complex), relative to a GDF-8 protein not bound by the same protein. In an embodiment, the activity of the mature GDF-8 protein, when bound by a protein comprising at least one follistatin domain, is inhibited at least 50%, or at least 60, 62, 64, 66, 68, 70, 72, 72, 76, 78, 80, 82, 84, 86, or 88%, or at least 90, 91, 92, 93, or 94%, or at least 95% to 100% relative to a mature GDF-8 protein that is not bound by a protein having a follistatin domain.

[096] Pharmaceutical preparations comprising proteins comprising at least one follistatin domain are administered in therapeutically effective amounts. As used herein, an "effective amount" of the protein is a dosage which is sufficient to reduce the activity of GDF-8 to achieve a desired biological outcome. The desired biological outcome may be any therapeutic benefit including an increase in muscle mass, an increase in muscle strength, improved metabolism, decreased adiposity, or improved glucose homeostasis. Such improvements may be measured by a variety of methods including those that measure lean and fat body mass (such as dual x-ray scanning analysis), muscle strength, serum lipids, serum leptin, serum

- 32 -

glucose, glycated hemoglobin, glucose tolerance, and improvement in the secondary complication of diabetes.

[097] Generally, a therapeutically effective amount may vary with the subject's age, condition, and sex, as well as the severity of the medical condition in the subject. The dosage may be determined by a physician and adjusted, as necessary, to suit observed effects of the treatment. Appropriate dosages for administering at least one protein comprising at least one follistatin domain may range from 5 mg to 100 mg, from 15 mg to 85 mg, from 30 mg to 70 mg, or from 40 mg to 60 mg. Proteins can be administered in one dose, or at intervals such as once daily, once weekly, and once monthly. Dosage schedules can be adjusted depending on the affinity of the protein for GDF-8, the half life of the protein, or the severity of the patient's condition. Generally, the compositions are administered as a bolus dose, to maximize the circulating levels of proteins comprising at least one follistatin domain for the greatest length of time after the dose. Continuous infusion may also be used after the bolus dose.

[098] Toxicity and therapeutic efficacy of such compounds can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., for determining the LD₅₀ (the dose lethal to 50% of the population) and the ED₅₀ (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio LD₅₀/ED₅₀. Proteins comprising at least one follistatin domain which exhibit large therapeutic indices may be used.

[099] Data obtained from the cell culture assays and animal studies can be used in evaluating a range of dosage for use in humans. The dosage of such compounds may lie within a range of circulating concentrations that include the ED₅₀ with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized. For any protein comprising at least one follistatin domain used in the present invention, the therapeutically effective dose can

- 33 -

be estimated initially from cell culture assays. A dose may be formulated in animal models to achieve a circulating plasma concentration range that includes the IC₅₀ (i.e., the concentration of the test protein which achieves a half-maximal inhibition of symptoms) as determined in cell culture. Levels in plasma may be measured, for example, by high performance liquid chromatography. The effects of any particular dosage can be monitored by a suitable bioassay. Examples of suitable bioassays include GDF-8 protein/receptor binding assays, creatine kinase assays, assays based on glucose uptake in adipocytes, and immunological assays.

Methods of Administering DNA

[0100] The present invention also provides gene therapy for the in vivo production of proteins comprising at least one follistatin domain. Such therapy would achieve its therapeutic effect by introduction of the polynucleotide sequences into cells or tissues having the disorders as listed herein.

[0101] Delivery of polynucleotide sequences of proteins comprising at least one follistatin domain can be achieved using a recombinant expression vector such as a chimeric virus or a colloidal dispersion system. Target liposomes may be used for therapeutic delivery of the polynucleotide sequences. Various viral vectors which can be utilized for gene therapy include adenovirus, herpes virus, vaccinia, or an RNA virus such as a retrovirus. The retroviral vector may be a derivative of a murine or avian retrovirus. Examples of retroviral vectors in which a single foreign gene can be inserted include, but are not limited to: Moloney murine leukemia virus (MoMuLV), Harvey murine sarcoma virus (HaMuSV), murine mammary tumor virus (MuMTV), and Rous sarcoma virus (RSV). A number of additional retroviral vectors can incorporate multiple genes. All of these vectors can transfer or incorporate a gene for a selectable marker so that transduced cells can be identified and generated. By inserting a GDF propeptide polynucleotide sequence of interest into the viral vector, along with another

- 34 -

gene which encodes the ligand for a receptor on a specific target cell, for example, the vector is now target specific.

[0102] Retroviral vectors can be made target specific by attaching, for example, a sugar, a glycolipid, or a protein. Targeting may be accomplished by using an antibody. Those of skill in the art will recognize that specific polynucleotide sequences can be inserted into the retroviral genome or attached to a viral envelope to allow target specific delivery of the retroviral vector containing the polynucleotide of proteins comprising at least one follistatin domain. In one embodiment, the vector is targeted to muscle cells or muscle tissue.

[0103] Since recombinant retroviruses are defective, they require helper cell lines that contain plasmids encoding all of the structural genes of the retrovirus under the control of regulatory sequences within the LTR. These plasmids are missing a nucleotide sequence which enables the packaging mechanism to recognize an RNA transcript for encapsidation. Helper cell lines which have deletions of the packaging signal include, but are not limited to PSI.2, PA317 and PA12, for example. These cell lines produce empty virions, since no genome is packaged. If a retroviral vector is introduced into such cells in which the packaging signal is intact, but the structural genes are replaced by other genes of interest, the vector can be packaged and vector virion produced.

[0104] Alternatively, other tissue culture cells can be directly transfected with plasmids encoding the retroviral structural genes gag, pol and env, by conventional calcium phosphate transfection. These cells are then transfected with the vector plasmid containing the genes of interest. The resulting cells release the retroviral vector into the culture medium.

[0105] Another targeted delivery system for a polynucleotide of a protein comprising at least one follistatin domain is a colloidal dispersion system. Colloidal dispersion systems include macromolecule complexes, nanocapsules, microspheres, beads, and lipid-based systems including oil-in-water emulsions, micelles, mixed micelles, and liposomes. Liposomes are

- 35 -

artificial membrane vesicles which are useful as delivery vehicles in vitro and in vivo. RNA, DNA and intact virions can be encapsulated within the aqueous interior and be delivered to cells in a biologically active form (see, for example, Fraley, *et al.* (1981) *Trends Biochem. Sci.*, 6: 77). Methods for efficient gene transfer using a liposome vehicle, are known in the art (see, for example, Mannino, *et al.* (1988) *Biotechniques*, 6: 682. The composition of the liposome is usually a combination of phospholipids, usually in combination with steroids, especially cholesterol. Other phospholipids or other lipids may also be used. The physical characteristics of liposomes depend on pH, ionic strength, and the presence of divalent cations.

[0106] Examples of lipids useful in liposome production include phosphatidyl compounds, such as phosphatidylglycerol, phosphatidylcholine, phosphatidylserine, phosphatidylethanolamine, sphingolipids, cerebroside, and gangliosides. Illustrative phospholipids include egg phosphatidylcholine, dipalmitoylphosphatidylcholine and distearoylphosphatidylcholine. The targeting of liposomes is also possible based on, for example, organ-specificity, cell-specificity, and organelle-specificity and is known in the art.

[0107] There is a wide range of methods which can be used to deliver the cells expressing proteins comprising at least one follistatin domain to a site for use in modulating a GDF-8 response. In one embodiment of the invention, the cells expressing follistatin protein can be delivered by direct application, for example, direct injection of a sample of such cells into the site of tissue damage. These cells can be purified. The such cells can be delivered in a medium or matrix which partially impedes their mobility so as to localize the cells to a site of injury. Such a medium or matrix could be semi-solid, such as a paste or gel, including a gel-like polymer. Alternatively, the medium or matrix could be in the form of a solid, a porous solid which will allow the migration of cells into the solid matrix, and hold them there while allowing proliferation of the cells.

Methods of Detection and Isolation of GDF-8

[0108] Proteins comprising at least one follistatin domain may be used to detect the presence or level of GDF-8, *in vivo* or *in vitro*. By correlating the presence or level of these proteins with a medical condition, one of skill in the art can diagnose the associated medical condition. The medical conditions that may be diagnosed by the proteins comprising at least one follistatin domain are set forth herein.

[0109] Such detection methods are well known in the art and include ELISA, radioimmunoassay, immunoblot, western blot, immunofluorescence, immuno-precipitation, and other comparable techniques. Proteins comprising at least one follistatin domain may further be provided in a diagnostic kit that incorporates one or more of these techniques to detect GDF-8. Such a kit may contain other components, packaging, instructions, or other material to aid the detection of the protein and use of the kit.

[0110] Where proteins comprising at least one follistatin domain are intended for diagnostic purposes, it may be desirable to modify them, for example with a ligand group (such as biotin) or a detectable marker group (such as a fluorescent group, a radioisotope or an enzyme). If desired, the proteins may be labeled using conventional techniques. Suitable labels include fluorophores, chromophores, radioactive atoms, electron-dense reagents, enzymes, and ligands having specific binding partners. Enzymes are typically detected by their activity. For example, horseradish peroxidase is usually detected by its ability to convert 3,3',5,5'-tetramethylbenzidine (TMB) to a blue pigment, quantifiable with a spectrophotometer. Other suitable binding partners include biotin and avidin or streptavidin, IgG and protein A, and the numerous receptor-ligand couples known in the art. Other permutations and possibilities will be readily apparent to those of ordinary skill in the art, and are considered as equivalents within the scope of the instant invention.

[0111] Proteins comprising at least one follistatin domain or fragments thereof may also be useful for isolating GDF-8 in a purification process. In

- 37 -

one type of process, proteins may be immobilized, for example, through incorporation into a column or resin. The proteins are used to bind GDF-8, and then subject to conditions which result in the release of the bound GDF-8. Such processes may be used for the commercial production of GDF-8.

[0112] The following examples provide embodiments of the invention. One of ordinary skill in the art will recognize the numerous modifications and variations that may be performed without altering the spirit or scope of the present invention. Such modifications and variations are believed to be encompassed within the scope of the invention. The examples do not in any way limit the invention. It is understood that all of the numbers in the specification and claims are modified by the term about, as small changes in dosages, for example, would be considered to be within the scope of the invention.

EXAMPLES

Example 1: Purification of JA16-Conjugated Beads

[0113] N-hydroxysuccinimidyl-activated beads (4% beaded agarose, Sigma H-8635, St Louis MO) were washed in MilliQ-H₂O and incubated for 4 hours at 4°C with the anti-GDF-8 JA16 monoclonal antibody (3-4 µg/µl in 100 mM MOPS, pH 7.5) at a ratio to allow a final concentration of 10 mg JA16/ml resin. Beads were washed extensively with 100 mM MOPS pH 7.5 and phosphate-buffered saline (PBS) (Ausubel *et al*, (1999) *Current Protocols in Molecular Biology*, John Wiley & Sons) and stored at 4°C in PBS until use. Control beads were prepared identically without JA16 antibody.

Example 2: Affinity Purification

[0114] A total of 40 µl of packed JA16-conjugated or control beads were incubated with 15 ml normal Balb/C mouse serum (Golden West Biologicals, Temecula CA) or 30 ml pooled normal human serum (ICN Biomedical, Aurora OH) for 3 hours at 4°C. Beads were washed twice in ~10 ml of cold 1% Triton X-100/PBS, twice in ~10 ml of cold 0.1% Triton X-100/PBS, and twice in ~1 ml of cold PBS. Proteins were eluted from the

- 38 -

beads in three subsequent steps. First, the beads were treated to a 'mock elution', where 100 μ l of PBS was added to the beads and incubated at 4°C for 30 minutes. The supernatant was collected and combined with 30 μ l 4x LDS sample buffer (Invitrogen, Carlsbad CA). Second, the beads were subject to a 'peptide elution', 100 μ l of 1 μ g/ μ l competing peptide (sequence: DFGLDSDEHSTESRSSRYPLTVDFEAFGWD-COOH (SEQ ID NO:12)) in PBS was added to the beads and again incubated at 4°C for 30 minutes. The supernatant was collected as before. Third, the beads were treated with an 'SDS elution' technique, where 30 μ l of 4x LDS buffer (Invitrogen) and 100 μ l of PBS was added to the beads and heated to 80°C for 10 minutes before transferring the supernatant to a fresh tube.

[0115] A silver stained gel of the proteins released in each of the elution steps is shown in Figure 1. Two protein bands in the silver-stained gel shown in Figure 1 of approximately 12 and 36 kDa were specifically eluted from JA16-conjugated beads, but not from unconjugated control beads.

Example 3: Mass Spectrometry

[0116] Samples were reduced with NuPage 10x reducing agent (Invitrogen) for 10 minutes at 80°C and alkylated with 110 μ M iodoacetamide for 30 minutes at 22°C in the dark. Samples were run immediately on 10% NuPage Bis-Tris gels in an MES buffer system according to manufacturer's recommendations (Invitrogen) and silver stained using a gluteraldehyde-free system (Shevchenko, *et al.*, (1996) *Anal. Chem.*, 68: 850-858). Bands were excised and digested with Sequencing Grade Modified Trypsin (Promega, Madison WI) in an Abimed Digest Pro (Langenfeld, Germany) or ProGest Investigator (Genomics Solutions, Ann Arbor MI). The volume of digested samples was reduced by evaporation and supplemented with 1% acetic acid to a final volume of ~20 μ l. Samples (5-10 μ l) were loaded onto a 10 cm x 75 μ m inner diameter C₁₈ reverse phase column packed in a Picofrit needle (New Objectives, Woburn MA). MS/MS data was collected using an LCQ Deca or LCQ Deca XP (Finnigan, San Jose CA) mass spectrometer and searched

against the NCBI non-redundant database using the Sequest program (Finnigan). Unless otherwise noted, all peptide sequences listed in this paper corresponded to MS/MS spectra that were deemed high quality by manual inspection and produced X_{corr} scores > 2.5 in the Sequest scoring system.

Example 4: Western Blots

[0117] Proteins were transferred to a 0.45 μm nitrocellulose membrane (Invitrogen) and blocked with blocking buffer (5% non-fat dry milk in Tris-buffered saline (TBS: 10 mM Tris-Cl, pH 7.5, 150 mM NaCl)) at 4°C overnight. Blots were then probed with primary antibody diluted 1:1000 in blocking buffer for 1-3 hours at room temperature, washed 5x with TBS, probed with horseradish peroxidase-conjugated secondary antibody in blocking buffer for 1-3 hours at room temperature, and washed as before. Signals were detected by autoradiography using the West Pico Substrate (Pierce).

Example 5: Isolation of GDF-8

[0118] An experiment using the methods described in the previous Examples resulted in the isolation of GDF-8. Since GDF-8 in its reduced form is 12 kDa, we speculated that the protein in the lower band from the silver-stained gel shown in Figure 1 was mature GDF-8. To confirm this hypothesis, we excised this band, digested it with trypsin, and obtained MS/MS spectra of the resulting peptides by LC-MS/MS. MS/MS spectra corresponding to six tryptic peptides confirmed that mature GDF-8 was isolated from this region of the gel, as shown in Figure 2A and Table 1.

[0119] Table 1 lists peptides derived from GDF-8 (SEQ ID NO:13-20), GDF-8 propeptide (SEQ ID NO:21-27), FLRG (SEQ ID NO:28-30), and GASP1 (SEQ ID NO:31-35) that were found in JA16 immunoprecipitates from mouse and human serum. The immediately preceding amino acid in the protein sequence is shown in parentheses for each peptide and the charge state of the peptide (z) and the Sequest program correlation coefficient (X_{corr} , a measure of confidence) are listed. The sequence listing numbers in the table refer only to the isolated peptides and their sequences. The preceding

- 40 -

amino acids in parentheses are not included in the peptides, but are provided only for reference. All spectra were confirmed by manual inspection.

[0120] Interestingly, the western blot also contained a band corresponding to unprocessed full-length GDF-8 (43 kDa), implying that some portion of this molecule is secreted into serum without undergoing proteolytic processing (Figure 2B). The presence of unprocessed GDF-8 was confirmed by mass spectrometry (data not shown). Thus, the affinity purification method effectively isolated GDF-8 from normal mouse serum.

[0121] Although the JA16 antibody recognizes both GDF-8 and the highly related protein BMP/GDF-11, we saw no evidence of BMP-11 peptides in our affinity purified samples by mass spectrometry.

Table 1: Peptides Identified in JA16 Immunoprecipitates

	mouse serum	z	X _{corr}
GDF-8 (mature)	(K) ANYCSGEGEFVFLQK (SEQ ID NO:13)	3+	4.63
	(K) MSPINMLYFNGK (SEQ ID NO:14)	2+	3.81
	(R) DFGLDCDEHSTESR (SEQ ID NO:15)	2+	3.47
	(K) ANYCSGEGEFVFLQK (SEQ ID NO:16)	2+	3.31
	(K) M*SPINMLYFNGK (SEQ ID NO:17)	3+	2.95
	(R) YPLTVDFEAFGWDWIIAPK (SEQ ID NO:18)	2+	2.86
	(K) M*SPINM*LYFNGK (SEQ ID NO:19)	2+	2.51
	(R) GSAGPCCTPTK (SEQ ID NO:20)	2+	2.43
GDF-8 (propeptide)	(K) LDM*SPGTGIWQSIDVK (SEQ ID NO:21)	2+	3.82
	(K) ALDENGHDLAVTFPGPGEDGLNPFLEVK (SEQ ID NO: 22)	3+	3.17
	(K) LDMSPGTGIWQSIDVK (SEQ ID NO:23)	2+	2.98
	(R) ELIDQYDVQR (SEQ ID NO:24)	2+	2.97
	(K) TPTTVFVQILR (SEQ ID NO:25)	2+	2.91
	(K) AQLWIYLRPVK (SEQ ID NO:26)	2+	2.77
	(K) EGLCNACAWR (SEQ ID NO:27)	2+	2.75
follistatin-like related gene (FLRG)	(R) PQSCLVDQTGSAHCVVCR (SEQ ID NO:28)	3+	3.34
	(K) DSCDGVCEGPGK (SEQ ID NO:29)	2+	2.99
novel multidomain	(K) SCAQVVCPR (SEQ ID NO:30)	2+	2.59
	(R) ECETDQECETYEK (SEQ ID NO:31)	2+	2.98
protease inhibitor (GASP1)	(R) ADFPLSVVR (SEQ ID NO:32)	2+	2.56
	(R) EACEESCPFPR (SEQ ID NO:33)	2+	2.95
	(R) SDFVILGR (SEQ ID NO:34)	2+	2.73
	(R) VSELTEEQDSGR (SEQ ID NO:35)	2+	3.88

M* = oxidized methionine

	human serum	z	X _{corr}
GDF-8 (mature)	(K) ANYCSGEGEFVFLQK (SEQ ID NO:36)	2+	4.21
	(R) DFGLDCDEHSTESR (SEQ ID NO:37)	3+	2.08
GDF-8 (propeptide)	(K) ALDENGHDLAVTFPGPGEDGLNPFLEVK (SEQ ID NO:38)	3+	3.71
	(R) ELIDQYDVQR (SEQ ID NO:39)	2+	3.01
follistatin-like related gene (FLRG)	(R) PQSCVVDQTGSAHCVVCR (SEQ ID NO:40)	3+	3.37
	(R) CECAPDCSGLPAR (SEQ ID NO:41)	2+	3.21
multidomain protease inhibitor (GASP1)	(R) LQVCGSDGATYR (SEQ ID NO:42)	2+	3.06
	(R) VSELTEEPDSGR (SEQ ID NO:43)	2+	2.44
	(R) CYMDAEACSK (SEQ ID NO:44)	2+	2.69
	(K) GITLAVVTCR (SEQ ID NO:45)	2+	2.42

Example 6: Isolation of Proteins Bound to GDF-8

[0122] Once it was confirmed that the affinity purification technique could successfully isolate GDF-8 from normal mouse serum, we proceeded to identify proteins that bind to GDF-8 under native conditions. The 36 kDa band on the silver-stained gel shown in Figure 1 was analyzed as described above. Mass spectrometry identified two proteins in this region of the gel that were specific to the JA16-immunopurified sample. These were determined to be the GDF-8 propeptide and follistatin-like related gene (FLRG). The peptides identified from each of these proteins are shown in Table 1 (SEQ ID NO:13-27). High quality MS/MS spectra were found for six unique peptides from GDF-8 propeptide and three unique peptides from FLRG; representative peptides are shown in Figures 3A and 3C. Furthermore, the presence of both of these proteins was confirmed by western blotting with polyclonal antibodies specific to GDF-8 propeptide and FLRG respectively (Figures 3B and 3D). Thus, circulating GDF-8 appears to bind to the GDF-8 propeptide and to FLRG *in vivo*.

Example 7: Isolation of Novel Proteins that Bind GDF-8

[0123] To characterize the major components of the circulating GDF-8 complex *in vivo*, native GDF-8 and its associated proteins from wild-type mouse serum were isolated by affinity purification with an agarose-conjugated anti-GDF-8 monoclonal antibody, JA16. JA16-bound proteins were subjected to subsequent elution steps with PBS buffer alone (mock elution), a peptide that could compete with GDF-8 for JA16 binding, and SDS detergent. These samples were concentrated, run on a one-dimensional SDS-PAGE gel, and visualized by silver stain (Figure 4). Two bands unique to the JA16 purified samples are visible—a 12 kDa band identified as GDF-8, and a 36 kDa band containing both GDF-8 propeptide and FLRG.

[0124] In order to determine if one could identify other proteins that were bound to GDF-8 *in vivo*, we scaled up the purification approximately five-fold and used mass spectrometry to search for proteins that were present in

- 43 -

the JA16 immunocomplex, but not in the negative control. To achieve this goal, we excised regions of the silver stained gel corresponding to molecular weights between 10 and 200 kDa into 13 gel slices, as shown in Figure 4. Each of these slices was subjected to in-gel trypsin digestion and LC-MS/MS. Comparison of the resulting MS/MS spectra to the non-redundant NCBI database of known proteins did not reveal any additional proteins specific to the JA16 immunoprecipitate, although the proteins previously described (mature GDF-8, GDF-8 propeptide, unprocessed GDF-8, and FLRG) were all identified in these samples (Figure 4). Background proteins that were found both in the JA16 immunocomplex and in the negative control sample included abundant serum proteins, such as albumin, immunoglobulins, and complement proteins. There was no evidence of other TGF- β superfamily members, including the highly related protein BMP-11/GDF-11, in the JA16 samples. Thus, the JA16 antibody specifically purified GDF-8 in these experiments.

[0125] Interestingly, we found no evidence of follistatin in our GDF-8 immunocomplexes, despite the fact that JA16 is capable of immunoprecipitating a GDF-8/follistatin complex *in vitro* (data not shown). Follistatin has been shown to inhibit GDF-8 activity by antagonizing the association of GDF-8 with the ActRIIB receptor (Lee and McPherron (2001) *Proc. Natl. Acad. Sci. U.S.A.*, 98: 9306-9311). Our results suggest that follistatin does not play a major role in the regulation of the activity of the circulating GDF-8 complex under normal conditions.

[0126] Since the identification of proteins by this MS/MS procedure is dependent on the content of the database being searched, we further analyzed the data from Figure 4 by comparing the MS/MS spectra collected from the 13 samples to a database of proteins predicted from the Celera mouse genomic sequence. This analysis identified an additional protein specific to the JA16-purified sample, and is hereby referred to as GDF-associated serum protein 1 (GASP1). Since the initial identification of this

- 44 -

protein, this sequence has been added to the NCBI nr database by the public genome sequencing effort under the accession number gi|20914039.

[0127] Five peptides corresponding to the sequence of GASP1 were identified on the basis of high-quality MS/MS spectra (Table 1 (SEQ ID NO:31-35); Figure 5A and B). The spectra corresponding to GASP1 peptides were found in band 3, which contains 70-80 kDa proteins. However, a specific band corresponding to this protein was not visible, probably due to the abundance of background immunoglobulins and albumin at this area (see Figure 4). Sequest X_{corr} scores above 2.3 are generally considered significant for 2^+ ions. Fortuitously, one of the peptides identified in our experiments (sequence = ECETDQECETYEK (SEQ ID NO:31)) spans the junction between the two exons that code for this protein, verifying the accuracy of Celera's gene prediction algorithm in this instance.

[0128] The sequences of the GASP1 transcript and protein were predicted prior to the actual cloning of GASP1 (Figure 6). GASP1 was predicted to be a 571 amino acid protein with a predicted molecular mass of 63 kDa. It has a putative signal sequence/cleavage site at its N-terminus and two possible sites for N-glycosylation at amino acids 314 and 514. Analysis of the GASP1 protein sequence by Pfam and BLAST (according to the techniques in Altschul *et al.* (1990) *J. Mol. Biol.*, 215: 403-410; Bateman *et al.* (2002) *Nucleic Acids Res.*, 30: 276-280) revealed that GASP1 contains many conserved domains, including a WAP domain, a follistatin/Kazal domain, an immunoglobulin domain, two tandem Kunitz domains, and a netrin domain (Figure 14A). WAP domains, originally identified in whey acidic protein, contain 8 cysteines that form a four-disulfide core and are often found in proteins with anti-protease activity (Hennighausen and Sippel (1982) *Nucleic Acids Res.*, 10: 2677-2684; Seemuller *et al.* (1986) *FEBS Lett.*, 199: 43-48). It is believed that the follistatin domain mediates the interaction between GDF-8 and GASP1. The C-terminal region of follistatin domains contains a similarity to Kazal serine protease inhibitor domains. In the case of GASP1, this region is even more closely related to Kazal domains than in follistatin or

- 45 -

FLRG, suggesting the possibility that this region may have an additional protease inhibitor function. Kunitz domains, originally identified in bovine pancreatic trypsin inhibitor, also inhibit serine proteases, thus establishing a likely role for GASP1 in the regulation of this class of proteins. Furthermore, netrin domains have been implicated in the inhibition of metalloproteases (Banyai and Patthy, 1999; Mott *et al.*, 2000). Thus, based on the presence of these conserved regions, GASP1 is likely to inhibit the activity of proteases, perhaps regulating GDF-8 processing or activation of the latent GDF-8 complex.

[0129] BLAST searches against the mouse Celera transcript database revealed a protein that has >50% identity with GASP1, referred to here as GASP2. GASP2 contains the same domain structure as GASP1, suggesting that these proteins define a two member family of multivalent protease inhibitors (Figure 14B). Interestingly, only peptides corresponding to GASP1, not GASP2, were found in our JA16 purified samples. This result suggests that GASP1 and GASP2 likely have different biological specificity. Both GASP1 and GASP2 are conserved in humans (>90% identity with mouse). The sequence for human GASP1 is now available in the NCBI nr database under the accession number gi|18652308. Although, the concentration of GDF-8 in human serum is considerably lower than that found in mouse serum (Hill *et al.* (2002) *J. Biol. Chem.*, 277: 40735-40741), the sensitivity of mass spectrometric analysis of proteins allowed us to identify 3 peptides corresponding to the human homolog of GASP1 from JA16 immunoprecipitations from human serum (Table 1). None of these peptides were found in the corresponding negative control. Again, there was no evidence of human GASP2 in these experiments. Thus, the interaction between GASP1 and GDF-8 is conserved between mouse and human.

[0130] GDF-8 is produced nearly exclusively in skeletal muscle. In order to determine the tissue distribution of GASP1 mRNA, a 551 bp fragment of GASP1 was amplified from first-strand cDNA produced from a variety of mouse tissues and staged embryos (Figure 10). A mouse GASP1 fragment

- 46 -

was amplified from normalized mouse first-strand cDNA panels (Clontech, Palo Alto CA) using the Advantage cDNA PCR kit (Clontech) according to the manufacturer's recommendations (forward primer: 5' TTGGCCACTGCCACCACAATCTCAACCACTT 3' (SEQ ID NO:46); reverse primer: 5' TCTCAGCATGGCCATGCCGCCGTCGA 3' (SEQ ID NO:47)). GASP1 appears to be fairly widely-expressed, with particularly high expression in skeletal muscle and heart. Significant expression is also seen in brain, lung, and testis. In contrast, liver and kidney express relatively low levels of GASP1 mRNA. Developmentally, the level of GASP1 mRNA remains fairly constant, perhaps increasing slightly between day 7 and day 11 of mouse embryogenesis.

Example 8: GDF-8 in Human and Mouse Serum

[0131] The concentration of GDF-8 in human serum is considerably lower than that found in mouse serum. Since GDF-8 has potential as a therapeutic target, it was a goal to determine the composition of the circulating GDF-8 complex in humans. This knowledge would determine the validity of the mouse model and potentially identify alternative therapeutic targets. Thus, the JA16-based affinity purification of GDF-8 was repeated using human serum. Due to the lower level of GDF-8 in human serum compared with mouse, no bands corresponding to mature GDF-8 and GDF-8 propeptide/FLRG were visualized (Figure 11A). However, western blotting with a polyclonal antibody that recognizes the mature region of GDF-8 revealed the presence of mature and unprocessed GDF-8 in the JA16-purified samples (Figure 11B).

[0132] We took advantage of the high sensitivity of mass spectrometry to identify proteins that co-purified with mature GDF8. The lanes corresponding to peptide eluted samples from both negative control and JA16-conjugated beads were sliced into 16 pieces. These gel slices were subjected to in-gel trypsin digestion, nanoflow LC-MS/MS, and analysis with Sequest as before.

- 47 -

[0133] Interestingly, the only proteins that were identified specifically in the JA16 samples and not the negative control were mature GDF-8, GDF-8 propeptide, human FLRG, and the human homolog of GASP1. The peptides found from each of these proteins are listed in Table 1 (SEQ ID NO:36-45) and representative MS/MS spectra are shown in Figure 12. Thus the *in vivo* GDF-8 complex appears to be conserved between mouse and human.

Example 9: Cloning and Characterization of Mouse GASP1

[0134] After identifying the predicted GASP1 sequence, it was a goal to determine the actual sequence of mouse GASP1. Based on the Celera predicted sequence, the GASP1 coding sequence was amplified from mouse heart QUICKCLONE cDNA (Clontech) by PCR with PfuTurbo polymerase (Stratagene) using the following primers (fp: 5' CACCATGTGTGCCCCAGGGTATCATCGGTTCTGG 3' (SEQ ID NO:50); rp: 5' TTGCAAGCCCAGGAAGTCCTTGAGGAC 3' (SEQ ID NO:51)). The PCR product from this reaction ran as a single major band of approximately 1700 base pairs on a 1% agarose gel. The amplified DNA was then cloned into the TOPO sites of the pcDNA3.1D/V5-His-TOPO vector (Invitrogen) so as to include an in-frame C-terminal V5-His tag according to manufacturers' recommendations. The full-length cDNA insert was sequenced on both strands. The nucleotide sequence of the mouse GASP1 clone is shown in Figure 13. This clone matched the predicted Celera sequence, with the exception of some base substitutions in wobble codons that did not change the predicted amino acid sequence (i.e., 288C:G; 294G:A; 615G:A; 738A:G; 768C:T; 1407A:G; 1419A:G; and 1584C:G, where the first base at the indicated position is that reported by Celera and the second base is that obtained from sequencing of the clone; see Figure 6A and B).

[0135] To determine the N-terminal processing of the GASP1 protein, we transfected COS1 cells with a mammalian expression vector encoding mouse GASP1 cloned with a C-terminal V5-His tag (GASP1-V5-His). Serum-free conditioned media was harvested 48 hours later and analyzed by western

- 48 -

blot analysis with an anti-V5 polyclonal antibody (Sigma). More specifically, conditioned media was collected 48 hours after transfection of COS1 cells with GASP1-V5-His/pcDNA3.1D-V5-His-TOPO or empty vector using the FuGENE 6 reagent (Roche) in serum-free Dulbecco's modified Eagle's medium.

[0136] A single band, running at approximately 80 kDa was seen, confirming that GASP1 is secreted into the conditioned media (data not shown). Approximately 10 ml of this conditioned media was run over a His-affinity column and further purified by reverse phase chromatography. This purification scheme yielded a band the expected size of full-length GASP1 on a Coomassie stained SDS-PAGE gel. Edman sequencing of this band determined an N-terminal sequence of L-P-P-I-R-Y-S-H-A-G-I (SEQ ID NO:52). Thus, amino acids 1-29 of GASP1 constitute the signal sequence that is removed during processing and secretion.

Example 10: Binding of Recombinantly-Produced GASP1 to GDF-8 Propeptide and Mature GDF-8

[0137] Next, it was determined that recombinantly-produced GASP1 had the same binding pattern to GDF-8 as GASP1 isolated from mouse serum. For immunoprecipitations with recombinant proteins, 400 μ l conditioned media from vector- or GASP1-transfected cells was combined with 1.2 μ g of recombinant purified GDF-8 and/or GDF-8 propeptide protein (Thies *et al.*, 2001). JA16 (10 μ l packed volume) or anti-V5 (30 μ l) conjugated agarose beads were incubated with the supplemented conditioned media for two hours at 4°C and washed twice in cold 1% Triton in phosphate-buffered saline (PBS) and twice in PBS. Beads were resuspended in 50 μ l 1x LDS buffer with DTT. Western blots were performed as previously described (Hill *et al.*, 2002).

[0138] To confirm and further characterize the interaction between GDF-8 and GASP1, we incubated purified recombinant GDF-8 and purified recombinant GDF-8 propeptide with conditioned media from COS1 cells transfected with either a vector control or GASP1-V5-His. We then

- 49 -

immunoprecipitated GDF-8 with JA16-conjugated agarose beads and looked for co-purification of GASP1 and GDF-8 propeptide using western blots (Figure 15A). Both GASP1 (lane 3) and GDF-8 propeptide (lane 1) co-immunoprecipitated with GDF-8, proving that GDF-8 can interact with both of these proteins. Neither GASP1 nor propeptide were detected in JA16 immunoprecipitates in the absence of GDF-8 (lane 4), eliminating the possibility of non-specific binding in these experiments. When all three proteins were present, both GASP1 and GDF-8 propeptide were pulled down with GDF-8, suggesting the possibility that these proteins may form a tertiary complex (lane 5). However, this experiment does not eliminate the possibility that GASP1 and propeptide are bound to the same epitope on separate GDF-8 molecules.

[0139] To further confirm the interaction between GASP1 and GDF-8, we performed the reverse immunoprecipitation by pulling down GASP1 from conditioned media supplemented with GDF-8 and/or GDF-8 propeptide recombinant protein. To achieve this, we used an agarose-conjugated monoclonal antibody directed against the V5 epitope of the C-terminal V5-His tag on GASP1. As expected, GDF-8 co-immunoprecipitated with GASP1 (Figure 15B, lanes 3 and 5), further confirming a direct interaction between these proteins. Surprisingly, GDF-8 propeptide also co-purified with GASP1, even in the absence of GDF-8 (lane 4), suggesting that GDF-8 propeptide can bind directly to GASP1. Thus, GASP1 binds to both GDF-8 and GDF-8 propeptide independently. This is in contrast to FLRG, another follistatin-domain protein, that binds exclusively to mature GDF-8 (Hill et al. (2002) *J. Biol. Chem.*, 277: 40735-40741). Addition of both GDF-8 and propeptide consistently showed less propeptide binding to GASP1 than when propeptide was added alone. This observation suggests that GASP1 may not bind to the GDF-8 small latent complex.

- 50 -

Example 11: GASP1-Mediated Inhibition of GDF-8 and BMP-11, But Not Activin or TGF- β 1, Activity

[0140] A luciferase reporter construct, pGL3-(CAGA)₁₂ (SEQ ID NO:53) (Dennler et al. (1998) *EMBO J.*, 17: 3091-3100) was transiently transfected into A204 or RD rhabdomyosarcoma cells. Dilutions of conditioned media from vector or GASP1 transfected cells were incubated for 30 minutes at 37 °C with 10 ng/ml GDF-8, 10 ng/ml BMP-11, 10 ng/ml rh activin A (R&D Systems), or 0.5 ng/ml rh TGF- β 1 (R&D Systems). Luciferase activity was measured according to Thies et al. (2001) *Growth Factors*, 18: 251-259 and Zimmers et al. (2002) *Science*, 296: 1486-1488. In this assay, A204 cells respond to GDF-8, BMP-11, and activin, but do not respond well to TGF- β 1. RD cells respond to both GDF-8 and TGF- β 1. Thus, we used A204 cells to test for the ability of GASP1 to inhibit GDF-8, BMP-11, and activin and RD cells to monitor the activity of TGF- β and GDF-8. Results for GDF-8 are shown from A204 cells, but were similar in RD cells. A standard curve measuring the concentration dependence of the luciferase activity induced by each of these growth factors was generated for each experiment (data not shown). The growth factor concentrations used fall in the linear region of this curve such that small changes in concentration result in measurable changes in luciferase activity.

[0141] Two follistatin-domain proteins, follistatin and FLRG inhibit GDF-8 activity in a (CAGA)₁₂ (SEQ ID NO:53) luciferase transcriptional reporter assay, but also inhibit the biological activity of the related proteins, activin and BMP-11. The ability of GASP1 to inhibit GDF-8, BMP-11, activin, and TGF- β 1 activity in the (CAGA)₁₂ (SEQ ID NO:53) reporter assay was also tested.

[0142] Various dilutions of conditioned media from COS cells transfected with V5-His tagged GASP1 or a vector control were incubated with purified recombinant GDF-8 (10 ng/ml), BMP-11 (10 ng/ml), activin (10 ng/ml), or TGF- β 1 (0.5 ng/ml) and assayed for growth factor activity in rhabdomyosarcoma cells expressing the (CAGA)₁₂ (SEQ ID NO:53) reporter construct. GASP1 potently inhibited GDF-8 activity in a concentration

- 51 -

dependent manner (Figure 16A). GASP1 similarly inhibited the activity of BMP-11 in this assay (Figure 16B), as might be expected since mature GDF-8 and BMP-11 are highly conserved and differ by only 11 amino acids. Surprisingly, GASP1 did not inhibit the activity of activin or TGF- β 1 (Figure 16C and D), suggesting a very high level of specificity, which is not demonstrated by follistatin itself. Thus, GASP1 exhibits specificity in its inhibition of GDF-8 and BMP-11.

[0143] The affinity of GASP1 for GDF-8 was evaluated by determining the IC₅₀ for inhibition of GDF-8 in the reporter gene assay. GASP1-V5-His protein was purified from conditioned media on a cobalt affinity column and eluted as described above. Fractions containing GASP1 were further purified by size exclusion chromatography in PBS using a BioSepS3000 column (Phenomenex). As shown in Figure 17, GASP1 inhibited GDF-8 with an IC₅₀ of approximately 3 nM.

Example 12: Treatment of Muscle Disorders

[0144] GASP1 may be administered to patients suffering from a disease or disorder related to the functioning of GDF-8 according to Table 2. Patients take the composition one time or at intervals, such as once daily, and the symptoms of their disease or disorder improve. For example, symptoms related to a muscle disorder are improved, as measured by muscle mass, muscle activity, and or muscle tone. This shows that the composition of the invention is useful for the treatment of diseases or disorders related to the functioning of GDF-8, such as muscle disorders.

Table 2: Administration of GASP1

Patient	Disease	Route of Administration	Dosage	Dosage Frequency	Predicted Results
1	muscular dystrophy	subcutaneous	25 mg	once daily	increase in muscle mass and improvement in muscle activity
2	"	"	50 mg	"	"
3	"	"	50 mg	once weekly	"
4	"	"	50 mg	once monthly	"
5	"	intramuscular	25 mg	once daily	"
6	"	"	50 mg	"	"
7	"	"	50 mg	once weekly	"
8	"	"	50 mg	once monthly	"
9	"	intravenous	25 mg	once daily	"
10	"	"	50 mg	"	"
11	"	"	50 mg	once weekly	"
12	"	"	50 mg	once monthly	"
13	diabetes	subcutaneous	50 mg	once daily	improvement in the management of blood sugar levels
14	"	"	50 mg	once weekly	"
15	"	intramuscular	50 mg	"	"
16	"	intravenous	50 mg	"	"
17	obesity	subcutaneous	50 mg	once daily	weight loss and increase in muscle mass
18	"	intramuscular	50 mg	once weekly	"
19	"	intravenous	50 mg	"	"

[0145] The entire contents of all references, patents and published patent applications cited throughout this application are herein incorporated by reference. The foregoing detailed description has been given for illustration purposes only. A wide range of changes and modifications can be made to the embodiments described above. It should therefore be understood that it is the following claims, including all equivalents, are intended to define the scope of the invention.

1

SEQUENCE LISTING

<110> HILL, JENNIFER J.
WOLFMAN, NEIL M.

<120> FOLLISTATIN DOMAIN CONTAINING PROTEINS

<130> 08702.0015-00

<140>

<141>

<150> 60/357,846

<151> 2002-02-21

<150> 60/434,645

<151> 2002-12-20

<160> 53

<170> PatentIn Ver. 2.1

<210> 1

<211> 1716

<212> DNA

<213> Mus sp.

<400> 1

```

atgtgtgccc cagggtatca tcggttctgg tttcactggg ggctgctggt gctgctgctc 60
ctcgaggctc cccttcgagg cctagcactg ccacccatcc gatactccca tgcggggcatc 120
tgccccaacg acatgaaccc caacctctgg gtggatgccc agagcacctg caagecgagag 180
tgtgaaacag accaggaatg tgagacctat gagaaatgct gcccgaatgt gtgtgggacc 240
aagagctgtg tggcagcccc ctacatggat gtgaaagggg agaagggccc tgtgggcatg 300
cccaaggagg ccacatgtga ccatttcatg tgccctgcagc agggctctga gtgtgacatc 360
tgggacggcc agcccgtgtg taagtgcaaa gatcgtctgt agaaggagcc cagcttcacc 420
tgtgcctctg atggccttac ctactacaac cgttgcttca tggacgccga agcctgctcc 480
aagggcatca cactgtctgt ggtcacctgt cgttatcact tcacctggcc taacaccagc 540
cctccaccgc ctgagaccac ggtgcatccc accaccgect ctccggagac tctcgggctg 600
gacatggcag ccccggccct gctcaaccac cctgtccatc agtcagtcac cgtgggtgag 660
actgtgagtt tcctctgtga cgtggtaggc cggcctcggc cagagctcac ttgggagaaa 720
cagctggagg accgagaaaa tgttgctcatg aggcccaacc acgtgcgcgg taatgtggtg 780
gtcactaaca ttgcccagct ggtcatctac aacgtccagc cccaggatgc tggcatatac 840
acctgtacag ctcgaaatgt cgctgggtgtc ctgagggctg acttcccgtt gtcgggtggc 900
aggggtggtc aggccagggc cacttcagag agcagttctc atggcacagc ttttccagca 960
acagagtgcc tgaagcccc agacagtgag gactgtggag aggagcagac acgctggcac 1020
ttcgacgcc aggctaacia ctgcctcact ttcaccttg gccactgcca ccacaatctc 1080
aaccactttg agacctacga ggcctgtatg ctggcttgta tgagtgggccc attggccacc 1140
tgcagcctgc ctgcctgca agggccttgc aaagcttatg tcccacgctg ggcctacaac 1200
agccagacag gcctatgcca gtccttcgtc tatggcggct gtgagggcaa cggtaacaac 1260
tttgaagacc gtgaggcttg tgaggagtgc tgcccttcc cgaggggtaa ccagcactgc 1320
cgggcctgca agccccggca aaaacttgtt accagcttct gtcggagtga ctttgtcatc 1380
ctgggcaggg tctctgagct gaccgaagag caagactcag gccgtgccct ggtgaccgtg 1440
gatgaggtct taaaagatga gaagatgggc ctcaagtttc tgggccggga gcctctggaa 1500
gtcacccctgc ttcattgtaga ctggacctgt ccttgcccca acgtgacagt ggggtgagaca 1560
ccactcatca tcatggggga ggtcgacggc ggcatggcca tgctgagacc cgatagcttt 1620
gtgggggcat cgagcacacg gcggttcagg aagctccgtg aggtcatgta caagaaaacc 1680
tgtgacgtcc tcaaggactt cctgggcttg caatga 1716

```


2

<210> 2
 <211> 1716
 <212> DNA
 <213> Mus sp.

<400> 2
 atgtgtgccc cagggtatca tcggttctgg ttactactggg ggctgctggt gctgctgctc 60
 ctcgaggctc cccttcgagg cctagcactg ccacccatcc gatactccca tgcgggcatc 120
 tgccccaacg acatgaaccc caacctctgg gtggatgccc agagcacctg caagcgagag 180
 tgtgaaacag accaggaatg tgagacctat gagaaatgct gcccacatgt gtgtgggacc 240
 aagagctgtg tggcagcccg ctacatggat gtgaaagggg agaagggggc tgtaggcatg 300
 cccaaggagg ccacatgtga ccatttcatg tgcctgcagc agggctctga gtgtgacatc 360
 tgggacggcc agcccgtgtg taagtgcaaa gatcgctgtg agaaggagcc cagcttcacc 420
 tgtgcctctg atggccttac ctactacaac cgttgcttca tggacgcca gaagctgctc 480
 aagggcatca cactgtctgt ggtcacctgt cgttatcact tcacctggcc taacaccagc 540
 cctccaccgc ctgagaccac ggtgcatccc accaccgctt ctccggagac tctcgggctg 600
 gacatggcag ccccagccct gctcaaccac cctgtccatc agtcagtcac cgtgggtgag 660
 actgtgagtt tcctctgtga cgtggtaggc cggcctcggc cagagctcac ttgggagaaa 720
 cagctggagg accgagagaa tgttgtcatg aggcccaacc acgtgctgtg taatgtggtg 780
 gtcactaaca ttgcccagct ggtcatctac aacgtccagc cccaggatgc tggcatatac 840
 acctgtacag ctgaaatgt cgctgggtgc ctgagggctg acttcccgtt gtcgggtggtc 900
 aggggtggtc aggccagggc cacttcagag agcagtctca atggcacagc ttttccagca 960
 acagagtgcc tgaagcccc agacagtgag gactgtggag aggagcagac acgctggcac 1020
 ttcgacgccc aggtaaaaa ctgcctcact ttcacctttg gccactgcca ccacaatctc 1080
 aaccactttg agacctacga ggcctgtatg ctggcttgta tgagtgggccc attggccacc 1140
 tgcagcctgc ctgccctgca agggccttgc aaagcttatg tcccacgctg ggcctacaac 1200
 agccagacag gcctatgcca gtccttcgtc tatggcggct gtgagggcaa cggtaacaac 1260
 tttgaaagcc gtgaggcttg tgaggagtcg tgtcccttcc cgaggggtaa ccagcactgc 1320
 cgggcctgca agccccggca aaaacttggt accagcttct gtcggagtga ctttgtcatc 1380
 ctgggcaggg tctctgagct gaccgaggag caagactcgg gccgtgccct ggtgaccgtg 1440
 gatgaggtct taaaagatga gaagatgggc ctcaagtttc tgggcccggga gcctctggaa 1500
 gtcaccctgc ttcattgtaga ctggacctgt ccttgcccca acgtgacagt ggggtgagaca 1560
 ccactcatca tcatggggga ggtggacggc ggcatggcca tgctgagacc cgatagcttt 1620
 gtggggggcat cgagcacacg gcggttcagg aagctccgtg aggtcatgta caagaaaacc 1680
 tgtgacgtcc tcaaggactt cctgggcttg caatga 1716

<210> 3
 <211> 571
 <212> PRT
 <213> Mus sp.

<400> 3
 Met Cys Ala Pro Gly Tyr His Arg Phe Trp Phe His Trp Gly Leu Leu
 1 5 10 15
 Leu Leu Leu Leu Leu Glu Ala Pro Leu Arg Gly Leu Ala Leu Pro Pro
 20 25 30
 Ile Arg Tyr Ser His Ala Gly Ile Cys Pro Asn Asp Met Asn Pro Asn
 35 40 45
 Leu Trp Val Asp Ala Gln Ser Thr Cys Lys Arg Glu Cys Glu Thr Asp
 50 55 60
 Gln Glu Cys Glu Thr Tyr Glu Lys Cys Cys Pro Asn Val Cys Gly Thr
 65 70 75 80

Lys Ser Cys Val Ala Ala Arg Tyr Met Asp Val Lys Gly Lys Lys Gly
 85 90 95
 Pro Val Gly Met Pro Lys Glu Ala Thr Cys Asp His Phe Met Cys Leu
 100 105 110
 Gln Gln Gly Ser Glu Cys Asp Ile Trp Asp Gly Gln Pro Val Cys Lys
 115 120 125
 Cys Lys Asp Arg Cys Glu Lys Glu Pro Ser Phe Thr Cys Ala Ser Asp
 130 135 140
 Gly Leu Thr Tyr Tyr Asn Arg Cys Phe Met Asp Ala Glu Ala Cys Ser
 145 150 155 160
 Lys Gly Ile Thr Leu Ser Val Val Thr Cys Arg Tyr His Phe Thr Trp
 165 170 175
 Pro Asn Thr Ser Pro Pro Pro Pro Glu Thr Thr Val His Pro Thr Thr
 180 185 190
 Ala Ser Pro Glu Thr Leu Gly Leu Asp Met Ala Ala Pro Ala Leu Leu
 195 200 205
 Asn His Pro Val His Gln Ser Val Thr Val Gly Glu Thr Val Ser Phe
 210 215 220
 Leu Cys Asp Val Val Gly Arg Pro Arg Pro Glu Leu Thr Trp Glu Lys
 225 230 235 240
 Gln Leu Glu Asp Arg Glu Asn Val Val Met Arg Pro Asn His Val Arg
 245 250 255
 Gly Asn Val Val Val Thr Asn Ile Ala Gln Leu Val Ile Tyr Asn Val
 260 265 270
 Gln Pro Gln Asp Ala Gly Ile Tyr Thr Cys Thr Ala Arg Asn Val Ala
 275 280 285
 Gly Val Leu Arg Ala Asp Phe Pro Leu Ser Val Val Arg Gly Gly Gln
 290 295 300
 Ala Arg Ala Thr Ser Glu Ser Ser Leu Asn Gly Thr Ala Phe Pro Ala
 305 310 315 320
 Thr Glu Cys Leu Lys Pro Pro Asp Ser Glu Asp Cys Gly Glu Glu Gln
 325 330 335
 Thr Arg Trp His Phe Asp Ala Gln Ala Asn Asn Cys Leu Thr Phe Thr
 340 345 350
 Phe Gly His Cys His His Asn Leu Asn His Phe Glu Thr Tyr Glu Ala
 355 360 365
 Cys Met Leu Ala Cys Met Ser Gly Pro Leu Ala Thr Cys Ser Leu Pro
 370 375 380

4

Ala Leu Gln Gly Pro Cys Lys Ala Tyr Val Pro Arg Trp Ala Tyr Asn
 385 390 395 400

Ser Gln Thr Gly Leu Cys Gln Ser Phe Val Tyr Gly Gly Cys Glu Gly
 405 410 415

Asn Gly Asn Asn Phe Glu Ser Arg Glu Ala Cys Glu Glu Ser Cys Pro
 420 425 430

Phe Pro Arg Gly Asn Gln His Cys Arg Ala Cys Lys Pro Arg Gln Lys
 435 440 445

Leu Val Thr Ser Phe Cys Arg Ser Asp Phe Val Ile Leu Gly Arg Val
 450 455 460

Ser Glu Leu Thr Glu Glu Gln Asp Ser Gly Arg Ala Leu Val Thr Val
 465 470 475 480

Asp Glu Val Leu Lys Asp Glu Lys Met Gly Leu Lys Phe Leu Gly Arg
 485 490 495

Glu Pro Leu Glu Val Thr Leu Leu His Val Asp Trp Thr Cys Pro Cys
 500 505 510

Pro Asn Val Thr Val Gly Glu Thr Pro Leu Ile Ile Met Gly Glu Val
 515 520 525

Asp Gly Gly Met Ala Met Leu Arg Pro Asp Ser Phe Val Gly Ala Ser
 530 535 540

Ser Thr Arg Arg Val Arg Lys Leu Arg Glu Val Met Tyr Lys Lys Thr
 545 550 555 560

Cys Asp Val Leu Lys Asp Phe Leu Gly Leu Gln
 565 570

<210> 4
 <211> 1923
 <212> DNA
 <213> Homo sapiens

<220>
 <221> modified_base
 <222> (732)
 <223> a, t, c or g

<400> 4
 atgaatccca acctctgggt ggacgcacag agcacctgca ggcgggagtg tgagacggac 60
 caggagtgtg agatggacca ggtgagtggg atccagaagc cacagtgtga ggcagaccag 120
 gtgaatgggg tccagaagcc gcaatgtgag atggaccaga agtgggagtg tgaggttgac 180
 caggtgagtg ggggtccagaa gccggtgtgt gaggcggacc aggtgagtg ggtccagaag 240
 ccacagtgtg agatggacca ggtgagtggg atccagaagc tggagtgtga ggcggaccag 300
 aagtgggagt atgaggtgga ccaggtgagt ggggtccaga agccacagtg tgagatggac 360
 caggtgagtg ggatccagaa gctggagtgt gaggcggacc aggagtgtga gacctatgag 420
 aagtgctgcc ccaacgtatg tgggaccaag agctgcgtgg cggcccgcta catggacgtg 480
 aaaggaaga agggcccagt gggcatgcc aaggaggcca catgtgacca cttcatgtgt 540
 ctgcagcagg gctctgagtg tgacatctgg gatggccagc ccgtgtgtaa gtgcaaagac 600

5

cgctgtgaga aggagcccag ctttacctgc gcctcggacg gcctcaccta ctataaccgc 660
 tgctacatgg atgccgaggg ctgctccaaa ggcatcacac tggccgttgt aacctgccgc 720
 tatcacttca cntggcccaa caccagcccc ccaccacctg agaccaccat gcaccccacc 780
 acagcctccc cagagacccc tgagctggac atggcggccc ctgcgctgct caacaacctt 840
 gtgcaccagt cggtcaccat ggggtgagaca gtgagcttcc tctgtgatgt ggtgggcccg 900
 ccccggcctg agatcacctg ggagaagcag ttggaggatc gggagaatgt ggtcatgcgg 960
 cccaaccatg tgcgtggcaa cgtgggtggc accaacattg cccagctggg catctataac 1020
 gcccagctgc aggatgctgg gatctacacc tgcacggccc ggaacgtggc tggggtcctg 1080
 agggctgatt tcccgtgtc ggtggtcagg ggtcatcagg ctgcagccac ctgagagagc 1140
 agccccaatg gcacggcttt cccggcggcc gagtgctga agcccccaga cagtgaggac 1200
 tgtggcgaag agcagaccgg ctggcacttc gatgcccagg ccaacaactg cctgaccttc 1260
 accttcggcc actgccaccg taacctcaac cactttgaga cctatgaggc ctgcatgctg 1320
 gcctgcatga gggggccgct ggccgcgtgc agcctgcccg ccctgcaggg gccctgcaaa 1380
 gcctacgcgc ctcgctgggc ttacaacagc cagacgggccc agtgccagtc ctttgtctat 1440
 ggtggctgcg agggcaatgg caacaacttt gagagccgtg aggccctgtga ggagtcgtgc 1500
 cccttcccca gggggaacca gcgctgtcgg gcctgcaagc ctcggcagaa gctcgttacc 1560
 agcttctgtc gcagcgactt tgtcatcctg ggccgagtct ctgagctgac cgaggagcct 1620
 gactcgggccc gcgcctggg gactgtggat gaggtcctaa aggatgagaa aatgggcctc 1680
 aagttcctgg gccaggagcc attggaggtc actctgcttc acgtggactg ggcattgccc 1740
 tgccccaacg tgaccgtgag cgagatgccg ctcattcatca tgggggaggt ggacggcggc 1800
 atggccatgc tgcgccccga tagctttgtg ggcgcatcga gtgcccgcgc ggtcaggaag 1860
 cttcgtgagg tcatgcacaa gaagacctgt gacgtcctca aggagtttct tggcttgcac 1920
 tga 1923

<210> 5

<211> 640

<212> PRT

<213> Homo sapiens

<400> 5

Met Asn Pro Asn Leu Trp Val Asp Ala Gln Ser Thr Cys Arg Arg Glu
 1 5 10 15

Cys Glu Thr Asp Gln Glu Cys Glu Met Asp Gln Val Ser Gly Ile Gln
 20 25 30

Lys Pro Gln Cys Glu Ala Asp Gln Val Asn Gly Val Gln Lys Pro Gln
 35 40 45

Cys Glu Met Asp Gln Lys Trp Glu Cys Glu Val Asp Gln Val Ser Gly
 50 55 60

Val Gln Lys Pro Val Cys Glu Ala Asp Gln Val Ser Gly Val Gln Lys
 65 70 75 80

Pro Gln Cys Glu Met Asp Gln Val Ser Gly Ile Gln Lys Leu Glu Cys
 85 90 95

Glu Ala Asp Gln Lys Trp Glu Tyr Glu Val Asp Gln Val Ser Gly Val
 100 105 110

Gln Lys Pro Gln Cys Glu Met Asp Gln Val Ser Gly Ile Gln Lys Leu
 115 120 125

Glu Cys Glu Ala Asp Gln Glu Cys Glu Thr Tyr Glu Lys Cys Cys Pro
 130 135 140

6

Asn Val Cys Gly Thr Lys Ser Cys Val Ala Ala Arg Tyr Met Asp Val
 145 150 155 160
 Lys Gly Lys Lys Gly Pro Val Gly Met Pro Lys Glu Ala Thr Cys Asp
 165 170 175
 His Phe Met Cys Leu Gln Gln Gly Ser Glu Cys Asp Ile Trp Asp Gly
 180 185 190
 Gln Pro Val Cys Lys Cys Lys Asp Arg Cys Glu Lys Glu Pro Ser Phe
 195 200 205
 Thr Cys Ala Ser Asp Gly Leu Thr Tyr Tyr Asn Arg Cys Tyr Met Asp
 210 215 220
 Ala Glu Ala Cys Ser Lys Gly Ile Thr Leu Ala Val Val Thr Cys Arg
 225 230 235 240
 Tyr His Phe Thr Trp Pro Asn Thr Ser Pro Pro Pro Pro Glu Thr Thr
 245 250 255
 Met His Pro Thr Thr Ala Ser Pro Glu Thr Pro Glu Leu Asp Met Ala
 260 265 270
 Ala Pro Ala Leu Leu Asn Asn Pro Val His Gln Ser Val Thr Met Gly
 275 280 285
 Glu Thr Val Ser Phe Leu Cys Asp Val Val Gly Arg Pro Arg Pro Glu
 290 295 300
 Ile Thr Trp Glu Lys Gln Leu Glu Asp Arg Glu Asn Val Val Met Arg
 305 310 315 320
 Pro Asn His Val Arg Gly Asn Val Val Val Thr Asn Ile Ala Gln Leu
 325 330 335
 Val Ile Tyr Asn Ala Gln Leu Gln Asp Ala Gly Ile Tyr Thr Cys Thr
 340 345 350
 Ala Arg Asn Val Ala Gly Val Leu Arg Ala Asp Phe Pro Leu Ser Val
 355 360 365
 Val Arg Gly His Gln Ala Ala Ala Thr Ser Glu Ser Ser Pro Asn Gly
 370 375 380
 Thr Ala Phe Pro Ala Ala Glu Cys Leu Lys Pro Pro Asp Ser Glu Asp
 385 390 395 400
 Cys Gly Glu Glu Gln Thr Arg Trp His Phe Asp Ala Gln Ala Asn Asn
 405 410 415
 Cys Leu Thr Phe Thr Phe Gly His Cys His Arg Asn Leu Asn His Phe
 420 425 430
 Glu Thr Tyr Glu Ala Cys Met Leu Ala Cys Met Ser Gly Pro Leu Ala
 435 440 445

7

Ala Cys Ser Leu Pro Ala Leu Gln Gly Pro Cys Lys Ala Tyr Ala Pro
 450 455 460

Arg Trp Ala Tyr Asn Ser Gln Thr Gly Gln Cys Gln Ser Phe Val Tyr
 465 470 475 480

Gly Gly Cys Glu Gly Asn Gly Asn Asn Phe Glu Ser Arg Glu Ala Cys
 485 490 495

Glu Glu Ser Cys Pro Phe Pro Arg Gly Asn Gln Arg Cys Arg Ala Cys
 500 505 510

Lys Pro Arg Gln Lys Leu Val Thr Ser Phe Cys Arg Ser Asp Phe Val
 515 520 525

Ile Leu Gly Arg Val Ser Glu Leu Thr Glu Glu Pro Asp Ser Gly Arg
 530 535 540

Ala Leu Val Thr Val Asp Glu Val Leu Lys Asp Glu Lys Met Gly Leu
 545 550 555 560

Lys Phe Leu Gly Gln Glu Pro Leu Glu Val Thr Leu Leu His Val Asp
 565 570 575

Trp Ala Cys Pro Cys Pro Asn Val Thr Val Ser Glu Met Pro Leu Ile
 580 585 590

Ile Met Gly Glu Val Asp Gly Gly Met Ala Met Leu Arg Pro Asp Ser
 595 600 605

Phe Val Gly Ala Ser Ser Ala Arg Arg Val Arg Lys Leu Arg Glu Val
 610 615 620

Met His Lys Lys Thr Cys Asp Val Leu Lys Glu Phe Leu Gly Leu His
 625 630 635 640

<210> 6

<211> 1731

<212> DNA

<213> Homo sapiens

<400> 6

atgtgggccc caaggtgtcg ccggttctgg tctcgctggg agcaggtggc agcgctgctg 60
 ctgctgctgc tactgctcgg ggtgcccccg cgaagcctgg cgctgccgcc catccgctat 120
 tcccacgccg gcatctgccc caacgacatg aatcccaacc tctgggtgga cgcacagagc 180
 acctgcaggc gggagtgtga gacggaccag gagtgtgaga cctatgagaa gtgctgcccc 240
 aacgtatgtg ggaccaagag ctgcgtggcg gcccgctaca tggacgtgaa agggaagaag 300
 ggcccagtg gcatgccc aa ggaggccaca tgtgaccact tcatgtgtct gcagcagggc 360
 tctgagtgtg acatctggga tggccagccc gtgtgtaagt gcaaagaccg ctgtgagaag 420
 gagcccagct ttacctgcgc ctcggacggc ctcacctact ataaccgctg ctacatggat 480
 gccgaggcct gctccaaagg catcacactg gccgttgtaa cctgccgcta tcaattcacc 540
 tggcccaaca ccagcccccc accacctgag accaccatgc accccaccac agcctccccca 600
 gagaccctg agctggacat ggcggcccct gcgctgctca acaaccctgt gcaccagtcg 660
 gtcaccatgg gtgagacagt gagtttctc tgtgatgtgg tgggccggcc ccggcctgag 720
 atcacctggg agaagcagtt ggaggatcgg gagaatgtgg tcatgcccgc caaccatgtg 780
 cgtggcaacg tgggtggtcac caacattgcc cagctggcca tctataacgc ccagctgcag 840
 gatgctggga tctacacctg cacggccccg aacgtggctg ggtcctgag ggctgatttc 900

```

ccgctgtcgg tggtcagggg tcatcaggct gcagccacct cagagagcag ccccaatggc 960
acggctttcc cggcggccga gtgcctgaag cccccagaca gtgaggactg tggcgaagag 1020
cagacccgct ggcacttcga tgcccaggcc aacaactgcc tgaccttcac cttcggccac 1080
tgccaccgta acctcaacca ctttgagacc tatgaggcct gcatgctggc ctgcatgagc 1140
gggccgctgg ccgcgtgcag cctgcccgcc ctgcaggggc cctgcaaagc ctacgcgctt 1200
cgctgggctt acaacagcca gacggggccag tgccagtcct ttgtctatgg tggctgcgag 1260
ggcaatggca acaactttga gagccgtgag gcctgtgagg agtcgtgccc cttccccagg 1320
gggaaccagc gctgtcgggc ctgcaagcct cggcagaagc tcgttaccag cttctgtcgc 1380
agcgactttg tcatcctggg ccgagtctct gagctgaccg aggagcctga ctcggggccgc 1440
gccctggtga ctgtggatga ggtcctaaag gatgagaaaa tgggcctcaa gttcctgggc 1500
caggagccat tggaggtcac tctgcttcac gtggactggg catgcccctg ccccaacgtg 1560
accgtgagcg agatgccgct catcatcatg ggggaggtgg acggcggcat ggccatgctg 1620
cgccccgata gctttgtggg cgcacgcagt gcccgccggg tcaggaagct tcgtgaggtc 1680
atgcacaaga agacctgtga cgtcctcaag gagtttcttg gcttgcactg a 1731

```

<210> 7

<211> 576

<212> PRT

<213> Homo sapiens

<400> 7

```

Met Trp Ala Pro Arg Cys Arg Arg Phe Trp Ser Arg Trp Glu Gln Val
  1                5                10                15

```

```

Ala Ala Leu Leu Leu Leu Leu Leu Leu Leu Leu Gly Val Pro Pro Arg Ser
                20                25                30

```

```

Leu Ala Leu Pro Pro Ile Arg Tyr Ser His Ala Gly Ile Cys Pro Asn
  35                40                45

```

```

Asp Met Asn Pro Asn Leu Trp Val Asp Ala Gln Ser Thr Cys Arg Arg
  50                55                60

```

```

Glu Cys Glu Thr Asp Gln Glu Cys Glu Thr Tyr Glu Lys Cys Cys Pro
  65                70                75                80

```

```

Asn Val Cys Gly Thr Lys Ser Cys Val Ala Ala Arg Tyr Met Asp Val
                85                90                95

```

```

Lys Gly Lys Lys Gly Pro Val Gly Met Pro Lys Glu Ala Thr Cys Asp
  100                105                110

```

```

His Phe Met Cys Leu Gln Gln Gly Ser Glu Cys Asp Ile Trp Asp Gly
  115                120                125

```

```

Gln Pro Val Cys Lys Cys Lys Asp Arg Cys Glu Lys Glu Pro Ser Phe
  130                135                140

```

```

Thr Cys Ala Ser Asp Gly Leu Thr Tyr Tyr Asn Arg Cys Tyr Met Asp
  145                150                155                160

```

```

Ala Glu Ala Cys Ser Lys Gly Ile Thr Leu Ala Val Val Thr Cys Arg
                165                170                175

```

```

Tyr His Phe Thr Trp Pro Asn Thr Ser Pro Pro Pro Pro Glu Thr Thr
                180                185                190

```

Met His Pro Thr Thr Ala Ser Pro Glu Thr Pro Glu Leu Asp Met Ala
 195 200 205

Ala Pro Ala Leu Leu Asn Asn Pro Val His Gln Ser Val Thr Met Gly
 210 215 220

Glu Thr Val Ser Phe Leu Cys Asp Val Val Gly Arg Pro Arg Pro Glu
 225 230 235 240

Ile Thr Trp Glu Lys Gln Leu Glu Asp Arg Glu Asn Val Val Met Arg
 245 250 255

Pro Asn His Val Arg Gly Asn Val Val Val Thr Asn Ile Ala Gln Leu
 260 265 270

Val Ile Tyr Asn Ala Gln Leu Gln Asp Ala Gly Ile Tyr Thr Cys Thr
 275 280 285

Ala Arg Asn Val Ala Gly Val Leu Arg Ala Asp Phe Pro Leu Ser Val
 290 295 300

Val Arg Gly His Gln Ala Ala Ala Thr Ser Glu Ser Ser Pro Asn Gly
 305 310 315 320

Thr Ala Phe Pro Ala Ala Glu Cys Leu Lys Pro Pro Asp Ser Glu Asp
 325 330 335

Cys Gly Glu Glu Gln Thr Arg Trp His Phe Asp Ala Gln Ala Asn Asn
 340 345 350

Cys Leu Thr Phe Thr Phe Gly His Cys His Arg Asn Leu Asn His Phe
 355 360 365

Glu Thr Tyr Glu Ala Cys Met Leu Ala Cys Met Ser Gly Pro Leu Ala
 370 375 380

Ala Cys Ser Leu Pro Ala Leu Gln Gly Pro Cys Lys Ala Tyr Ala Pro
 385 390 395 400

Arg Trp Ala Tyr Asn Ser Gln Thr Gly Gln Cys Gln Ser Phe Val Tyr
 405 410 415

Gly Gly Cys Glu Gly Asn Gly Asn Asn Phe Glu Ser Arg Glu Ala Cys
 420 425 430

Glu Glu Ser Cys Pro Phe Pro Arg Gly Asn Gln Arg Cys Arg Ala Cys
 435 440 445

Lys Pro Arg Gln Lys Leu Val Thr Ser Phe Cys Arg Ser Asp Phe Val
 450 455 460

Ile Leu Gly Arg Val Ser Glu Leu Thr Glu Glu Pro Asp Ser Gly Arg
 465 470 475 480

Ala Leu Val Thr Val Asp Glu Val Leu Lys Asp Glu Lys Met Gly Leu
 485 490 495

11

<400> 9

Met Pro Ala Pro Gln Pro Phe Leu Pro Leu Leu Phe Val Phe Val Leu
 1 5 10 15
 Ile His Leu Thr Ser Glu Thr Asn Leu Leu Pro Asp Pro Gly Ser His
 20 25 30
 Pro Gly Met Cys Pro Asn Glu Leu Ser Pro His Leu Trp Val Asp Ala
 35 40 45
 Gln Ser Thr Cys Glu Arg Glu Cys Thr Gly Asp Gln Asp Cys Ala Ala
 50 55 60
 Ser Glu Lys Cys Cys Thr Asn Val Cys Gly Leu Gln Ser Cys Val Ala
 65 70 75 80
 Ala Arg Phe Pro Ser Gly Gly Pro Ala Val Pro Glu Thr Ala Ala Ser
 85 90 95
 Cys Glu Gly Phe Gln Cys Pro Gln Gln Gly Ser Asp Cys Asp Ile Trp
 100 105 110
 Asp Gly Gln Pro Val Cys Arg Cys Arg Asp Arg Cys Glu Lys Glu Pro
 115 120 125
 Ser Phe Thr Cys Ala Ser Asp Gly Leu Thr Tyr Tyr Asn Arg Cys Tyr
 130 135 140
 Met Asp Ala Glu Ala Cys Leu Arg Gly Leu His Leu His Val Val Pro
 145 150 155 160
 Cys Lys His Ile Leu Ser Trp Pro Pro Ser Ser Pro Gly Pro Pro Glu
 165 170 175
 Thr Thr Ala Arg Pro Thr Pro Gly Ala Ala Pro Met Pro Pro Ala Leu
 180 185 190
 Tyr Asn Ser Pro Ser Pro Gln Ala Val His Val Gly Gly Thr Ala Ser
 195 200 205
 Leu His Cys Asp Val Ser Gly Arg Pro Pro Pro Ala Val Thr Trp Glu
 210 215 220
 Lys Gln Ser His Gln Arg Glu Asn Leu Ile Met Arg Pro Asp Gln Met
 225 230 235 240
 Tyr Gly Asn Val Val Val Thr Ser Ile Gly Gln Leu Val Leu Tyr Asn
 245 250 255
 Ala Gln Leu Glu Asp Ala Gly Leu Tyr Thr Cys Thr Ala Arg Asn Ala
 260 265 270
 Ala Gly Leu Leu Arg Ala Asp Phe Pro Leu Ser Val Leu Gln Arg Ala
 275 280 285
 Thr Thr Gln Asp Arg Asp Pro Gly Ile Pro Ala Leu Ala Glu Cys Gln
 290 295 300

12

Ala Asp Thr Gln Ala Cys Val Gly Pro Pro Thr Pro His His Val Leu
 305 310 315 320

Trp Arg Phe Asp Pro Gln Arg Gly Ser Cys Met Thr Phe Pro Ala Leu
 325 330 335

Arg Cys Asp Gly Ala Ala Arg Gly Phe Glu Thr Tyr Glu Ala Cys Gln
 340 345 350

Gln Ala Cys Val Arg Gly Pro Gly Asp Val Cys Ala Leu Pro Ala Val
 355 360 365

Gln Gly Pro Cys Gln Gly Trp Glu Pro Arg Trp Ala Tyr Ser Pro Leu
 370 375 380

Leu Gln Gln Cys His Pro Phe Val Tyr Ser Gly Cys Glu Gly Asn Ser
 385 390 395 400

Asn Asn Phe Glu Thr Arg Glu Ser Cys Glu Asp Ala Cys Pro Val Pro
 405 410 415

Arg Thr Pro Pro Cys Arg Ala Cys Arg Leu Lys Ser Lys Leu Ala Leu
 420 425 430

Ser Leu Cys Arg Ser Asp Phe Ala Ile Val Gly Arg Leu Thr Glu Val
 435 440 445

Leu Glu Glu Pro Glu Ala Ala Gly Gly Ile Ala Arg Val Ala Leu Asp
 450 455 460

Asp Val Leu Lys Asp Asp Lys Met Gly Leu Lys Phe Leu Gly Thr Lys
 465 470 475 480

Tyr Leu Glu Val Thr Leu Ser Gly Met Asp Trp Ala Cys Pro Cys Pro
 485 490 495

Asn Val Thr Ala Val Asp Gly Pro Leu Val Ile Met Gly Glu Val Arg
 500 505 510

Glu Gly Val Ala Val Leu Asp Ala Asn Ser Tyr Val Arg Ala Ala Ser
 515 520 525

Glu Lys Arg Val Lys Lys Ile Val Glu Leu Leu Glu Lys Lys Ala Cys
 530 535 540

Glu Leu Leu Asn Arg Phe Gln Asp
 545 550

<210> 10

<211> 1695

<212> DNA

<213> Homo sapiens

<400> 10

atgcccgcc tacgtccact cctgccgctc ctgctcctcc tccggctgac ctcgggggct 60
 ggcttgctgc cagggctggg gagccaccgc ggcgtgtgcc ccaaccagct cagccccaac 120
 ctgtgggtgg acgccagag cacctgtgag cgcgagtgtg gcagggacca ggactgtgcg 180

```

gctgctgaga agtgctgcat caacgtgtgt ggactgcaca gctgcgtggc agcacgcttc 240
cccggcagcc cagctgcgcc gacgacagcg gcctcctgcg agggctttgt gtgcccacag 300
cagggctcgg actgcgacat ctgggacggg cagcccgtgt gccgctgccg cgaccgctgt 360
gagaaggagc ccagcttcac ctgcgcctcg gacggcctca cctactacaa ccgctgctat 420
atggacgccg aggcctgcct gcggggcctg cacctccaca tcgtgccctg caagcacgtg 480
ctcagctggc cgcccagcag cccggggccg ccggagacca ctgcccgcc caccactggg 540
gccgcgcccg tgctcctgc cctgtacagc agcccctccc cacaggcggt gcaggttggg 600
ggtacggcca gcctccactg cgacgtcagc ggccgcccgc cgcttgcctg gacctgggag 660
aagcagagtc accagcgaga gaacctgatc atgcgcctg atcagatgta tggcaacgtg 720
gtggtcacca gcatcgggca gctggtgctc tacaacgcgc ggcccgaaga cgccggcctg 780
tacacctgca ccgcgcgcaa cgtgctggg ctgctgcggg ctgacttccc actctctgtg 840
gtccagcgag agccggccag ggacgcagcc cccagcatcc cagccccggc cgagtgcctg 900
ccggatgtgc aggcctgcac gggccccact tccccacacc ttgtcctctg gcactacgac 960
ccgcagcggg gcggctgcat gaccttccc gcccggtggt gtgatggggc ggcccgcggc 1020
tttgagacct acgagcatg ccagcaggcc tgtgcccgcg gccccggcga cgcttgcctg 1080
ctgcctgccg tgcagggccc ctgccggggc tgggagccgc gctgggccta cagcccgtg 1140
ctgcagcagt gccatccctt cgtgtacggg ggctgcgagg gcaacggcaa caacttccac 1200
agccgcgaga gctgcgagga tgctgcccc gtgcccgcga caccgccctg ccgcgcctgc 1260
cgcttccgga gcaagctggc gctgagcctg tgccgcagcg acttcgcat cgtggggcgg 1320
ctcacggagg tgctggagga gcccagggc gccggcggca tcgcccgcgt ggcgctcgag 1380
gacgtgctca aggatgacaa gatgggcctc aagttcttgg gcaccaagta cctggagggtg 1440
acgctgagtg gcatggactg ggctgcccc tgcccacaaca tgacggcggg cgacgggccg 1500
ctggtcatca tgggtgaggt gcgcgatggc gtggccgtgc tggacgccgg cagctacgtc 1560
cgcgccgcca gcgagaagcg cgtcaagaag atcttggagc tgctggagaa gcaggcctgc 1620
gagctgctca accgcttcca ggactagccc ccgcaggggc ctgcgccacc ccgtcctggt 1680
gaataaacgc actcc 1695

```

<210> 11
<211> 548
<212> PRT
<213> Homo sapiens

<400> 11
Met Pro Ala Leu Arg Pro Leu Leu Pro Leu Leu Leu Leu Leu Arg Leu
1 5 10 15
Thr Ser Gly Ala Gly Leu Leu Pro Gly Leu Gly Ser His Pro Gly Val
20 25 30
Cys Pro Asn Gln Leu Ser Pro Asn Leu Trp Val Asp Ala Gln Ser Thr
35 40 45
Cys Glu Arg Glu Cys Ser Arg Asp Gln Asp Cys Ala Ala Ala Glu Lys
50 55 60
Cys Cys Ile Asn Val Cys Gly Leu His Ser Cys Val Ala Ala Arg Phe
65 70 75 80
Pro Gly Ser Pro Ala Ala Pro Thr Thr Ala Ala Ser Cys Glu Gly Phe
85 90 95
Val Cys Pro Gln Gln Gly Ser Asp Cys Asp Ile Trp Asp Gly Gln Pro
100 105 110
Val Cys Arg Cys Arg Asp Arg Cys Glu Lys Glu Pro Ser Phe Thr Cys
115 120 125

14

Ala Ser Asp Gly Leu Thr Tyr Tyr Asn Arg Cys Tyr Met Asp Ala Glu
 130 135 140

Ala Cys Leu Arg Gly Leu His Leu His Ile Val Pro Cys Lys His Val
 145 150 155 160

Leu Ser Trp Pro Pro Ser Ser Pro Gly Pro Pro Glu Thr Thr Ala Arg
 165 170 175

Pro Thr Pro Gly Ala Ala Pro Val Pro Pro Ala Leu Tyr Ser Ser Pro
 180 185 190

Ser Pro Gln Ala Val Gln Val Gly Gly Thr Ala Ser Leu His Cys Asp
 195 200 205

Val Ser Gly Arg Pro Pro Pro Ala Val Thr Trp Glu Lys Gln Ser His
 210 215 220

Gln Arg Glu Asn Leu Ile Met Arg Pro Asp Gln Met Tyr Gly Asn Val
 225 230 235 240

Val Val Thr Ser Ile Gly Gln Leu Val Leu Tyr Asn Ala Arg Pro Glu
 245 250 255

Asp Ala Gly Leu Tyr Thr Cys Thr Ala Arg Asn Ala Ala Gly Leu Leu
 260 265 270

Arg Ala Asp Phe Pro Leu Ser Val Val Gln Arg Glu Pro Ala Arg Asp
 275 280 285

Ala Ala Pro Ser Ile Pro Ala Pro Ala Glu Cys Leu Pro Asp Val Gln
 290 295 300

Ala Cys Thr Gly Pro Thr Ser Pro His Leu Val Leu Trp His Tyr Asp
 305 310 315 320

Pro Gln Arg Gly Gly Cys Met Thr Phe Pro Ala Arg Gly Cys Asp Gly
 325 330 335

Ala Ala Arg Gly Phe Glu Thr Tyr Glu Ala Cys Gln Gln Ala Cys Ala
 340 345 350

Arg Gly Pro Gly Asp Ala Cys Val Leu Pro Ala Val Gln Gly Pro Cys
 355 360 365

Arg Gly Trp Glu Pro Arg Trp Ala Tyr Ser Pro Leu Leu Gln Gln Cys
 370 375 380

His Pro Phe Val Tyr Gly Gly Cys Glu Gly Asn Gly Asn Asn Phe His
 385 390 395 400

Ser Arg Glu Ser Cys Glu Asp Ala Cys Pro Val Pro Arg Thr Pro Pro
 405 410 415

Cys Arg Ala Cys Arg Leu Arg Ser Lys Leu Ala Leu Ser Leu Cys Arg
 420 425 430

15

Ser Asp Phe Ala Ile Val Gly Arg Leu Thr Glu Val Leu Glu Glu Pro
 435 440 445

Glu Ala Ala Gly Gly Ile Ala Arg Val Ala Leu Glu Asp Val Leu Lys
 450 455 460

Asp Asp Lys Met Gly Leu Lys Phe Leu Gly Thr Lys Tyr Leu Glu Val
 465 470 475 480

Thr Leu Ser Gly Met Asp Trp Ala Cys Pro Cys Pro Asn Met Thr Ala
 485 490 495

Gly Asp Gly Pro Leu Val Ile Met Gly Glu Val Arg Asp Gly Val Ala
 500 505 510

Val Leu Asp Ala Gly Ser Tyr Val Arg Ala Ala Ser Glu Lys Arg Val
 515 520 525

Lys Lys Ile Leu Glu Leu Leu Glu Lys Gln Ala Cys Glu Leu Leu Asn
 530 535 540

Arg Phe Gln Asp
 545

<210> 12
 <211> 30
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Illustrative
 competing peptide

<400> 12
 Asp Phe Gly Leu Asp Ser Asp Glu His Ser Thr Glu Ser Arg Ser Ser
 1 5 10 15

Arg Tyr Pro Leu Thr Val Asp Phe Glu Ala Phe Gly Trp Asp
 20 25 30

<210> 13
 <211> 15
 <212> PRT
 <213> Mus sp.

<400> 13
 Ala Asn Tyr Cys Ser Gly Glu Cys Glu Phe Val Phe Leu Gln Lys
 1 5 10 15

<210> 14
 <211> 12
 <212> PRT
 <213> Mus sp.

17

<210> 20
 <211> 11
 <212> PRT
 <213> Mus sp.

<400> 20
 Gly Ser Ala Gly Pro Cys Cys Thr Pro Thr Lys
 1 5 10

<210> 21
 <211> 16
 <212> PRT
 <213> Mus sp.

<400> 21
 Leu Asp Met Ser Pro Gly Thr Gly Ile Trp Gln Ser Ile Asp Val Lys
 1 5 10 15

<210> 22
 <211> 28
 <212> PRT
 <213> Mus sp.

<400> 22
 Ala Leu Asp Glu Asn Gly His Asp Leu Ala Val Thr Phe Pro Gly Pro
 1 5 10 15

Gly Glu Asp Gly Leu Asn Pro Phe Leu Glu Val Lys
 20 25

<210> 23
 <211> 16
 <212> PRT
 <213> Mus sp.

<400> 23
 Leu Asp Met Ser Pro Gly Thr Gly Ile Trp Gln Ser Ile Asp Val Lys
 1 5 10 15

<210> 24
 <211> 10
 <212> PRT
 <213> Mus sp.

<400> 24
 Glu Leu Ile Asp Gln Tyr Asp Val Gln Arg
 1 5 10

<210> 25
 <211> 11
 <212> PRT
 <213> Mus sp.

18

<400> 25

Thr	Pro	Thr	Thr	Val	Phe	Val	Gln	Ile	Leu	Arg
1				5					10	

<210> 26

<211> 11

<212> PRT

<213> Mus sp.

<400> 26

Ala	Gln	Leu	Trp	Ile	Tyr	Leu	Arg	Pro	Val	Lys
1				5					10	

<210> 27

<211> 10

<212> PRT

<213> Mus sp.

<400> 27

Glu	Gly	Leu	Cys	Asn	Ala	Cys	Ala	Trp	Arg
1				5					10

<210> 28

<211> 18

<212> PRT

<213> Mus sp.

<400> 28

Pro	Gln	Ser	Cys	Leu	Val	Asp	Gln	Thr	Gly	Ser	Ala	His	Cys	Val	Val
1				5					10					15	

Cys Arg

<210> 29

<211> 12

<212> PRT

<213> Mus sp.

<400> 29

Asp	Ser	Cys	Asp	Gly	Val	Glu	Cys	Gly	Pro	Gly	Lys
1				5					10		

<210> 30

<211> 9

<212> PRT

<213> Mus sp.

<400> 30

Ser	Cys	Ala	Gln	Val	Val	Cys	Pro	Arg
1				5				

19

<210> 31
 <211> 13
 <212> PRT
 <213> Mus sp.

<400> 31
 Glu Cys Glu Thr Asp Gln Glu Cys Glu Thr Tyr Glu Lys
 1 5 10

<210> 32
 <211> 9
 <212> PRT
 <213> Mus sp.

<400> 32
 Ala Asp Phe Pro Leu Ser Val Val Arg
 1 5

<210> 33
 <211> 11
 <212> PRT
 <213> Mus sp.

<400> 33
 Glu Ala Cys Glu Glu Ser Cys Pro Phe Pro Arg
 1 5 10

<210> 34
 <211> 8
 <212> PRT
 <213> Mus sp.

<400> 34
 Ser Asp Phe Val Ile Leu Gly Arg
 1 5

<210> 35
 <211> 12
 <212> PRT
 <213> Mus sp.

<400> 35
 Val Ser Glu Leu Thr Glu Glu Gln Asp Ser Gly Arg
 1 5 10

<210> 36
 <211> 15
 <212> PRT
 <213> Homo sapiens

<400> 36
 Ala Asn Tyr Cys Ser Gly Glu Cys Glu Phe Val Phe Leu Gln Lys
 1 5 10 15

20

<210> 37
 <211> 14
 <212> PRT
 <213> Homo sapiens

<400> 37
 Asp Phe Gly Leu Asp Cys Asp Glu His Ser Thr Glu Ser Arg
 1 5 10

<210> 38
 <211> 28
 <212> PRT
 <213> Homo sapiens

<400> 38
 Ala Leu Asp Glu Asn Gly His Asp Leu Ala Val Thr Phe Pro Gly Pro
 1 5 10 15

Gly Glu Asp Gly Leu Asn Pro Phe Leu Glu Val Lys
 20 25

<210> 39
 <211> 10
 <212> PRT
 <213> Homo sapiens

<400> 39
 Glu Leu Ile Asp Gln Tyr Asp Val Gln Arg
 1 5 10

<210> 40
 <211> 18
 <212> PRT
 <213> Homo sapiens

<400> 40
 Pro Gln Ser Cys Val Val Asp Gln Thr Gly Ser Ala His Cys Val Val
 1 5 10 15

Cys Arg

<210> 41
 <211> 13
 <212> PRT
 <213> Homo sapiens

<400> 41
 Cys Glu Cys Ala Pro Asp Cys Ser Gly Leu Pro Ala Arg
 1 5 10

21

<210> 42
 <211> 12
 <212> PRT
 <213> Homo sapiens

<400> 42
 Leu Gln Val Cys Gly Ser Asp Gly Ala Thr Tyr Arg
 1 5 10

<210> 43
 <211> 12
 <212> PRT
 <213> Homo sapiens

<400> 43
 Val Ser Glu Leu Thr Glu Glu Pro Asp Ser Gly Arg
 1 5 10

<210> 44
 <211> 10
 <212> PRT
 <213> Homo sapiens

<400> 44
 Cys Tyr Met Asp Ala Glu Ala Cys Ser Lys
 1 5 10

<210> 45
 <211> 10
 <212> PRT
 <213> Homo sapiens

<400> 45
 Gly Ile Thr Leu Ala Val Val Thr Cys Arg
 1 5 10

<210> 46
 <211> 31
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Primer

<400> 46
 ttggccactg ccaccacaat ctcaaccact t

31

<210> 47
 <211> 26
 <212> DNA
 <213> Artificial Sequence

22

<220>

<223> Description of Artificial Sequence: Primer

<400> 47

tctcagcatg gccatgccgc cgtcga

26

<210> 48

<211> 1716

<212> DNA

<213> Mus sp.

<220>

<221> CDS

<222> (1)..(1713)

<400> 48

atg	tgt	gcc	cca	ggg	tat	cat	cgg	ttc	tgg	ttt	cac	tgg	ggg	ctg	ctg	48
Met	Cys	Ala	Pro	Gly	Tyr	His	Arg	Phe	Trp	Phe	His	Trp	Gly	Leu	Leu	
1				5				10						15		

ttg	ctg	ctg	ctc	ctc	gag	gct	ccc	ctt	cga	ggc	cta	gca	ctg	cca	ccc	96
Leu	Leu	Leu	Leu	Leu	Glu	Ala	Pro	Leu	Arg	Gly	Leu	Ala	Leu	Pro	Pro	
			20					25					30			

atc	cga	tac	tcc	cat	gcg	ggc	atc	tgc	ccc	aac	gac	atg	aac	ccc	aac	144
Ile	Arg	Tyr	Ser	His	Ala	Gly	Ile	Cys	Pro	Asn	Asp	Met	Asn	Pro	Asn	
		35					40					45				

ctc	tgg	gtg	gat	gcc	cag	agc	acc	tgc	aag	cga	gag	tgt	gaa	aca	gac	192
Leu	Trp	Val	Asp	Ala	Gln	Ser	Thr	Cys	Lys	Arg	Glu	Cys	Glu	Thr	Asp	
	50					55					60					

cag	gaa	tgt	gag	acc	tat	gag	aaa	tgc	tgc	ccc	aat	gtg	tgt	ggg	acc	240
Gln	Glu	Cys	Glu	Thr	Tyr	Glu	Lys	Cys	Cys	Pro	Asn	Val	Cys	Gly	Thr	
65					70					75					80	

aag	agc	tgt	gtg	gca	gcc	cgc	tac	atg	gat	gtg	aaa	ggg	aag	aag	ggg	288
Lys	Ser	Cys	Val	Ala	Ala	Arg	Tyr	Met	Asp	Val	Lys	Gly	Lys	Lys	Gly	
				85				90						95		

cct	gta	ggc	atg	ccc	aag	gag	gcc	aca	tgt	gac	cat	ttc	atg	tgc	ctg	336
Pro	Val	Gly	Met	Pro	Lys	Glu	Ala	Thr	Cys	Asp	His	Phe	Met	Cys	Leu	
			100					105						110		

cag	cag	ggc	tct	gag	tgt	gac	atc	tgg	gac	ggc	cag	ccc	gtg	tgt	aag	384
Gln	Gln	Gly	Ser	Glu	Cys	Asp	Ile	Trp	Asp	Gly	Gln	Pro	Val	Cys	Lys	
		115					120					125				

tgc	aaa	gat	cgc	tgt	gag	aag	gag	ccc	agc	ttc	acc	tgt	gcc	tct	gat	432
Cys	Lys	Asp	Arg	Cys	Glu	Lys	Glu	Pro	Ser	Phe	Thr	Cys	Ala	Ser	Asp	
	130					135					140					

ggc	ctt	acc	tac	tac	aac	cgt	tgc	ttc	atg	gac	gcc	gaa	gcc	tgc	tcc	480
Gly	Leu	Thr	Tyr	Tyr	Asn	Arg	Cys	Phe	Met	Asp	Ala	Glu	Ala	Cys	Ser	
145					150					155					160	

23

aag ggc atc aca ctg tct gtg gtc acc tgt cgt tat cac ttc acc tgg	528
Lys Gly Ile Thr Leu Ser Val Val Thr Cys Arg Tyr His Phe Thr Trp	
165 170 175	
cct aac acc agc cct cca ccg cct gag acc acg gtg cat ccc acc acc	576
Pro Asn Thr Ser Pro Pro Pro Pro Glu Thr Thr Val His Pro Thr Thr	
180 185 190	
gcc tct ccg gag act ctc ggg ctg gac atg gca gcc cca gcc ctg ctc	624
Ala Ser Pro Glu Thr Leu Gly Leu Asp Met Ala Ala Pro Ala Leu Leu	
195 200 205	
aac cac cct gtc cat cag tca gtc acc gtg ggt gag act gtg agt ttc	672
Asn His Pro Val His Gln Ser Val Thr Val Gly Glu Thr Val Ser Phe	
210 215 220	
ctc tgt gac gtg gta ggc cgg cct cgg cca gag ctc act tgg gag aaa	720
Leu Cys Asp Val Val Gly Arg Pro Arg Pro Glu Leu Thr Trp Glu Lys	
225 230 235 240	
cag ctg gag gac cga gag aat gtt gtc atg agg ccc aac cac gtg cgt	768
Gln Leu Glu Asp Arg Glu Asn Val Val Met Arg Pro Asn His Val Arg	
245 250 255	
ggg aat gtg gtg gtc act aac att gcc cag ctg gtc atc tac aac gtc	816
Gly Asn Val Val Val Thr Asn Ile Ala Gln Leu Val Ile Tyr Asn Val	
260 265 270	
cag ccc cag gat gct ggc ata tac acc tgt aca gct cga aat gtc gct	864
Gln Pro Gln Asp Ala Gly Ile Tyr Thr Cys Thr Ala Arg Asn Val Ala	
275 280 285	
ggg gtc ctg agg gct gac ttc ccg ttg tcg gtg gtc agg ggt ggt cag	912
Gly Val Leu Arg Ala Asp Phe Pro Leu Ser Val Val Arg Gly Gly Gln	
290 295 300	
gcc agg gcc act tca gag agc agt ctc aat ggc aca gct ttt cca gca	960
Ala Arg Ala Thr Ser Glu Ser Ser Leu Asn Gly Thr Ala Phe Pro Ala	
305 310 315 320	
aca gag tgc ctg aag ccc cca gac agt gag gac tgt gga gag gag cag	1008
Thr Glu Cys Leu Lys Pro Pro Asp Ser Glu Asp Cys Gly Glu Glu Gln	
325 330 335	
aca cgc tgg cac ttc gac gcc cag gct aac aac tgc ctc act ttc acc	1056
Thr Arg Trp His Phe Asp Ala Gln Ala Asn Asn Cys Leu Thr Phe Thr	
340 345 350	
ttt ggc cac tgc cac cac aat ctc aac cac ttt gag acc tac gag gcc	1104
Phe Gly His Cys His His Asn Leu Asn His Phe Glu Thr Tyr Glu Ala	
355 360 365	
tgt atg ctg gct tgt atg agt ggg cca ttg gcc acc tgc agc ctg cct	1152
Cys Met Leu Ala Cys Met Ser Gly Pro Leu Ala Thr Cys Ser Leu Pro	
370 375 380	

24

gcc ctg caa ggg cct tgc aaa gct tat gtc cca cgc tgg gcc tac aac 1200
 Ala Leu Gln Gly Pro Cys Lys Ala Tyr Val Pro Arg Trp Ala Tyr Asn
 385 390 395 400

agc cag aca ggc cta tgc cag tcc ttc gtc tat ggc ggc tgt gag ggc 1248
 Ser Gln Thr Gly Leu Cys Gln Ser Phe Val Tyr Gly Gly Cys Glu Gly
 405 410 415

aac ggt aac aac ttt gaa agc cgt gag gct tgt gag gag tcg tgt ccc 1296
 Asn Gly Asn Asn Phe Glu Ser Arg Glu Ala Cys Glu Glu Ser Cys Pro
 420 425 430

ttc ccg agg ggt aac cag cac tgc cgg gcc tgc aag ccc cgg caa aaa 1344
 Phe Pro Arg Gly Asn Gln His Cys Arg Ala Cys Lys Pro Arg Gln Lys
 435 440 445

ctt gtt acc agc ttc tgt cgg agt gac ttt gtc atc ctg ggc agg gtc 1392
 Leu Val Thr Ser Phe Cys Arg Ser Asp Phe Val Ile Leu Gly Arg Val
 450 455 460

tct gag ctg acc gag gag caa gac tcg ggc cgt gcc ctg gtg acc gtg 1440
 Ser Glu Leu Thr Glu Glu Gln Asp Ser Gly Arg Ala Leu Val Thr Val
 465 470 475 480

gat gag gtc tta aaa gat gag aag atg ggc ctc aag ttt ctg ggc cgg 1488
 Asp Glu Val Leu Lys Asp Glu Lys Met Gly Leu Lys Phe Leu Gly Arg
 485 490 495

gag cct ctg gaa gtc acc ctg ctt cat gta gac tgg acc tgt cct tgc 1536
 Glu Pro Leu Glu Val Thr Leu Leu His Val Asp Trp Thr Cys Pro Cys
 500 505 510

ccc aac gtg aca gtg ggt gag aca cca ctc atc atc atg ggg gag gtg 1584
 Pro Asn Val Thr Val Gly Glu Thr Pro Leu Ile Ile Met Gly Glu Val
 515 520 525

gac ggc ggc atg gcc atg ctg aga ccc gat agc ttt gtg ggg gca tcg 1632
 Asp Gly Gly Met Ala Met Leu Arg Pro Asp Ser Phe Val Gly Ala Ser
 530 535 540

agc aca cgg cgg gtc agg aag ctc cgt gag gtc atg tac aag aaa acc 1680
 Ser Thr Arg Arg Val Arg Lys Leu Arg Glu Val Met Tyr Lys Lys Thr
 545 550 555 560

tgt gac gtc ctc aag gac ttc ctg ggc ttg caa tga 1716
 Cys Asp Val Leu Lys Asp Phe Leu Gly Leu Gln
 565 570

<210> 49
 <211> 571
 <212> PRT
 <213> Mus sp.

<400> 49
 Met Cys Ala Pro Gly Tyr His Arg Phe Trp Phe His Trp Gly Leu Leu
 1 5 10 15

25

Leu Leu Leu Leu Leu Glu Ala Pro Leu Arg Gly Leu Ala Leu Pro Pro
 20 25 30
 Ile Arg Tyr Ser His Ala Gly Ile Cys Pro Asn Asp Met Asn Pro Asn
 35 40 45
 Leu Trp Val Asp Ala Gln Ser Thr Cys Lys Arg Glu Cys Glu Thr Asp
 50 55 60
 Gln Glu Cys Glu Thr Tyr Glu Lys Cys Cys Pro Asn Val Cys Gly Thr
 65 70 75 80
 Lys Ser Cys Val Ala Ala Arg Tyr Met Asp Val Lys Gly Lys Lys Gly
 85 90 95
 Pro Val Gly Met Pro Lys Glu Ala Thr Cys Asp His Phe Met Cys Leu
 100 105 110
 Gln Gln Gly Ser Glu Cys Asp Ile Trp Asp Gly Gln Pro Val Cys Lys
 115 120 125
 Cys Lys Asp Arg Cys Glu Lys Glu Pro Ser Phe Thr Cys Ala Ser Asp
 130 135 140
 Gly Leu Thr Tyr Tyr Asn Arg Cys Phe Met Asp Ala Glu Ala Cys Ser
 145 150 155 160
 Lys Gly Ile Thr Leu Ser Val Val Thr Cys Arg Tyr His Phe Thr Trp
 165 170 175
 Pro Asn Thr Ser Pro Pro Pro Pro Glu Thr Thr Val His Pro Thr Thr
 180 185 190
 Ala Ser Pro Glu Thr Leu Gly Leu Asp Met Ala Ala Pro Ala Leu Leu
 195 200 205
 Asn His Pro Val His Gln Ser Val Thr Val Gly Glu Thr Val Ser Phe
 210 215 220
 Leu Cys Asp Val Val Gly Arg Pro Arg Pro Glu Leu Thr Trp Glu Lys
 225 230 235 240
 Gln Leu Glu Asp Arg Glu Asn Val Val Met Arg Pro Asn His Val Arg
 245 250 255
 Gly Asn Val Val Val Thr Asn Ile Ala Gln Leu Val Ile Tyr Asn Val
 260 265 270
 Gln Pro Gln Asp Ala Gly Ile Tyr Thr Cys Thr Ala Arg Asn Val Ala
 275 280 285
 Gly Val Leu Arg Ala Asp Phe Pro Leu Ser Val Val Arg Gly Gly Gln
 290 295 300
 Ala Arg Ala Thr Ser Glu Ser Ser Leu Asn Gly Thr Ala Phe Pro Ala
 305 310 315 320

26

Thr Glu Cys Leu Lys Pro Pro Asp Ser Glu Asp Cys Gly Glu Glu Gln
 325 330 335
 Thr Arg Trp His Phe Asp Ala Gln Ala Asn Asn Cys Leu Thr Phe Thr
 340 345 350
 Phe Gly His Cys His His Asn Leu Asn His Phe Glu Thr Tyr Glu Ala
 355 360 365
 Cys Met Leu Ala Cys Met Ser Gly Pro Leu Ala Thr Cys Ser Leu Pro
 370 375 380
 Ala Leu Gln Gly Pro Cys Lys Ala Tyr Val Pro Arg Trp Ala Tyr Asn
 385 390 395 400
 Ser Gln Thr Gly Leu Cys Gln Ser Phe Val Tyr Gly Gly Cys Glu Gly
 405 410 415
 Asn Gly Asn Asn Phe Glu Ser Arg Glu Ala Cys Glu Glu Ser Cys Pro
 420 425 430
 Phe Pro Arg Gly Asn Gln His Cys Arg Ala Cys Lys Pro Arg Gln Lys
 435 440 445
 Leu Val Thr Ser Phe Cys Arg Ser Asp Phe Val Ile Leu Gly Arg Val
 450 455 460
 Ser Glu Leu Thr Glu Glu Gln Asp Ser Gly Arg Ala Leu Val Thr Val
 465 470 475 480
 Asp Glu Val Leu Lys Asp Glu Lys Met Gly Leu Lys Phe Leu Gly Arg
 485 490 495
 Glu Pro Leu Glu Val Thr Leu Leu His Val Asp Trp Thr Cys Pro Cys
 500 505 510
 Pro Asn Val Thr Val Gly Glu Thr Pro Leu Ile Ile Met Gly Glu Val
 515 520 525
 Asp Gly Gly Met Ala Met Leu Arg Pro Asp Ser Phe Val Gly Ala Ser
 530 535 540
 Ser Thr Arg Arg Val Arg Lys Leu Arg Glu Val Met Tyr Lys Lys Thr
 545 550 555 560
 Cys Asp Val Leu Lys Asp Phe Leu Gly Leu Gln
 565 570

<210> 50

<211> 34

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

27

<400> 50
caccatgtgt gccccagggt atcatcggtt ctgg 34

<210> 51
<211> 27
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 51
ttgcaagccc aggaagtcct tgaggac 27

<210> 52
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Illustrative
N-terminal peptide sequence

<400> 52
Leu Pro Pro Ile Arg Tyr Ser His Ala Gly Ile
1 5 10

<210> 53
<211> 48
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 53
cagacagaca gacagacaga cagacagaca gacagacaga cagacaga 48

We claim:

1. A pharmaceutical composition comprising:
 - i) GASP1, and
 - ii) at least one pharmaceutically acceptable carrier.
2. The composition of claim 1, wherein the GASP1 has a stabilizing modification.
3. The composition of claim 2, wherein the modification is a fusion to the Fc region of an IgG molecule.
4. The composition of claim 3, wherein the IgG molecule is IgG1 or IgG4, or derivatives thereof.
5. The composition of claim 4, wherein the IgG molecule is IgG1 or a derivative thereof.
6. The composition of claim 3, wherein the IgG molecule is fused to the GASP1 by a linker peptide.
7. The composition of claim 2, wherein the modification comprises an altered glycosylation site.
8. The composition of claim 2, wherein the modification comprises at least one carbohydrate moiety.
9. The composition of claim 2, wherein the modification comprises albumin or an albumin derivative.
10. The composition of claim 2, wherein the modification comprises a nonproteinaceous polymer.
11. The composition of claim 2, wherein the modification comprises pegylation.

- 54 -

12. A diagnostic kit comprising GASP1 and at least one other kit component chosen from:
 - i) at least one agent that binds GASP1;
 - ii) at least one buffer and/or solution; and
 - iii) at least one structural component.
13. A recombinant cell comprising a nucleic acid encoding GASP1.
14. The recombinant cell of claim 13, wherein the GASP1 has a stabilizing modification.
15. A method of modulating GDF-8 comprising administering GASP1 and allowing the GASP1 to interact with GDF-8.
16. A method of treating a patient suffering from a medical disorder, comprising administering a therapeutically effective dose of GASP1 and allowing the GASP1 to interact with GDF-8.
17. A method of treating a patient suffering from a medical disorder, comprising administering a nucleic acid encoding GASP1, allowing the nucleic acid to be translated into GASP1, and allowing the translated GASP1 to interact with GDF-8.
18. A method of expressing a nucleic acid encoding GASP1, comprising administering a nucleic acid encoding GASP1 to a cell, allowing the nucleic acid to enter the cell, and allowing the cell to express the GASP1.
19. The method of claim 16, 17, or 18, wherein the GASP1 has a stabilizing modification.

- 55 -

20. The method of claim 19, wherein the modification is a fusion to the Fc region of an IgG molecule.
21. The method of claim 20, wherein the IgG molecule is IgG1 or IgG4, or derivatives thereof.
22. The method of claim 21, wherein the IgG molecule is IgG1 or a derivative thereof.
23. The method of claim 20, wherein the IgG molecule is fused to the GASP1 by a linker peptide.
24. The method of claim 19, wherein the modification comprises an altered glycosylation site.
25. The method of claim 19, wherein the modification comprises at least one carbohydrate moiety.
26. The method of claim 19, wherein the modification comprises albumin or an albumin derivative.
27. The method of claim 19, wherein the modification comprises a nonproteinaceous polymer.
28. The method of claim 19, wherein the modification comprises pegylation.
29. The method of claim 16, wherein the patient would therapeutically benefit from an increase in mass or quantity of muscle tissue.
30. The method of claim 16, wherein the disorder is a muscular disorder.
31. The method of claim 30, wherein the muscular disorder is muscular dystrophy.

- 56 -

32. The method of claim 31, wherein the muscular dystrophy is chosen from severe or benign X-linked muscular dystrophy, limb-girdle dystrophy, facioscapulohumeral dystrophy, myotonic dystrophy, distal muscular dystrophy, progressive dystrophic ophthalmoplegia, oculopharyngeal dystrophy, Duchenne's muscular dystrophy, and Fakuyama-type congenital muscular dystrophy.
33. The method of claim 30, wherein the disorder is chosen from amyotrophic lateral sclerosis, congestive obstructive pulmonary disease, congenital myopathy, myotonia congenital, familial periodic paralysis, paroxysmal myoglobinuria, myasthenia gravis, Eaton-Lambert syndrome, secondary myasthenia, denervation atrophy, organ atrophy, frailty, carpal tunnel syndrome, muscle atrophy, paroxymal muscle atrophy, sarcopenia, cachexia, and other muscle wasting syndromes.
34. The method of claim 30, wherein the disorder is a muscular disorder chosen from a traumatic injury to muscle tissue and a chronic injury to muscle tissue.
35. The method of claim 16, wherein the disorder is a metabolic disease or disorder.
36. The method of claim 35, wherein the disorder is insulin-dependent (type 1) diabetes mellitus, noninsulin-dependent (type 2) diabetes mellitus, hyperglycemia, impaired glucose tolerance, metabolic

- 57 -

syndrome (e.g., syndrome X), insulin resistance induced by trauma (e.g., burns or nitrogen imbalance), or obesity.

37. The method of claim 16, wherein the disorder is an adipose tissue disorder such as obesity.
38. The method of claim 16, wherein the disorder is a bone degenerative disease such as osteoporosis, glucocorticoid-induced osteoporosis, osteopenia, osteoarthritis, or osteoporosis-related fractures, or other disorders including low bone mass due to chronic glucocorticoid therapy, premature gonadal failure, androgen suppression, vitamin D deficiency, secondary hyperparathyroidism, nutritional deficiencies, and anorexia nervosa.
39. The method of claim 16, wherein the GASP1 is administered at one time, or at daily, weekly, or monthly intervals.
40. The method of claim 16, wherein the GASP1 is administered at a dose of from 5 mg to 100 mg.
41. The method of claim 16, wherein the GASP1 is administered at a dose of from 15 mg to 85 mg.
42. The method of claim 16, wherein the GASP1 is administered at a dose of from 30 mg to 70 mg.
43. The method of claim 16, wherein the GASP1 is administered at a dose of from 40 mg to 60 mg.
44. The composition of claim 1, wherein the GASP1 is chosen from:
 - i) SEQ ID No. 5;

- 58 -

- ii) SEQ ID No. 7; and
 - iii) substitution, addition, and/or deletion mutants of i) or ii).
45. The composition of claim 1, wherein the GASP1 is encoded by a nucleotide sequence chosen from:
- i) SEQ ID No. 4;
 - ii) SEQ ID No. 6;
 - iii) nucleotide sequences encoding substitution, addition, and/or deletion mutants of the sequences encoded by i) or ii); and
 - iv) nucleotide sequences encoding the same amino acid sequences as are encoded by i) or ii).
46. A pharmaceutical composition comprising:
- i) a fragment of GASP-1 chosen from
 - a) amino acids 174-239 of SEQ ID No. 5 representing the follistatin domain;
 - b) amino acids 110-175 of SEQ ID No. 7 representing the follistatin domain; and
 - c) substitution, addition, and/or deletion mutants of a) or b);and
 - ii) at least one pharmaceutically acceptable carrier.
47. The composition of claim 46, wherein the GASP1 fragment is encoded by a nucleotide sequence chosen from:
- i) nucleotides 520-717 of SEQ ID No. 4 encoding the follistatin domain;

- 59 -

- ii) nucleotides 328-525 of SEQ ID No. 6 encoding the follistatin domain;
- iii) nucleotide sequences encoding substitution, addition, and/or deletion mutants of the sequences encoded by i) or ii); and
- iv) nucleotide sequences encoding the same amino acid sequences as are encoded by i) or ii).

1/27




FIG. 1




FIG. 2A




FIG. 2B

3/27




FIG. 3A




FIG. 3B

4/27




FIG. 3C




FIG. 3D

5/27




FIG. 4

6/27




FIG. 5A




FIG. 5B

7/27

PREDICTED MOUSE GASP1 NUCLEOTIDE SEQUENCE
SEQ ID NO:1

```
1 atgtgtgccc cagggatatca tcggttctgg tttcactggg ggctgctggt gctgctgctc
61 ctcgaggctc cccttcgagg cctagcactg ccacccatcc gatactccca tgcgggcatc
121 tgccccaacg acatgaaccc caacctctgg gtggatgccc agagcacctg caagcgagag
181 tgtgaaacag accaggaatg tgagacctat gagaaatgct gcccacatgt gtgtgggacc
241 aagagctgtg tggcagcccg ctacatggat gtgaaagggg agaagggccc tgtgggcatg
301 cccaaggagg ccacatgtga ccatttcatg tgctgcagc agggctctga gtgtgacatc
361 tgggacggcc agcccgtgtg taagtgcaaa gatcgctgtg agaaggagcc cagcttcacc
421 tgtgcctctg atggccttac ctactacaac cgttgcttca tggacgccga agcctgctcc
481 aagggcatca cactgtctgt ggtcacctgt cgttatcact tcacctggcc taacaccagc
541 cctccaccgc ctgagaccac ggtgcatccc accaccgctt ctccggagac tctcgggctg
601 gacatggcag ccccgccctt gctcaaccac cctgtccatc agtcagtcac cgtgggtgag
661 actgtgagtt tcctctgtga cgtggtaggc cggcctcggc cagagctcac ttgggagaaa
721 cagctggagg accgagaaaa tgttgtcatg aggccaacc acgtgcgcgg taatgtggtg
781 gtcactaaca ttgcccagct ggatcatctac aacgtccagc cccaggatgc tggcatatac
841 acctgtacag ctcgaaatgt cgctgggtgc ctgagggctg acttcccgtt gtcgggtggtc
901 aggggtggtc aggccagggc cacttcagag agcagtctca atggcacagc tttccagca
961 acagagtgcc tgaagcccc agacagtgag gactgtggag aggagcagac acgctggcac
1021 ttcgacgccc aggctaacaa ctgcctcact ttcacctttg gccactgcca ccacaatctc
1081 aaccactttg agacctacga ggctgtatg ctggcttgta tgagtgggcc attggccacc
1141 tgcagcctgc ctgccctgca agggccttgc aaagcttatg tcccacgctg ggctacaac
1201 agccagacag gcctatgcca gtccttcgtc tatggcggct gtgagggcaa cggtaacaac
1261 tttgaaagcc gtgaggcttg tgaggagtcg tgtcccttcc cgaggggtaa ccagcactgc
1321 cgggcctgca agccccggca aaaacttgtt accagcttct gtcggagtga ctttgtcatc
1381 ctgggcaggg tctctgagct gaccgaagag caagactcag gccgtgccct ggtgaccgtg
1441 gatgaggtct taaaagatga gaagatgggc ctcaagtttc tgggccggga gcctctggaa
1501 gtcaccctgc ttcattgtaga ctggacctgt ccttgcccca acgtgacagt ggggtgagaca
1561 ccactcatca tcatggggga ggtcgacggc ggcattggca tgctgagacc cgatagcttt
1621 gtgggggcat cgagcacacg gcgggtcagg aagctccgtg aggtcatgta caagaaaacc
1681 tgtgacgtcc tcaaggactt cctgggcttg caatga
```

FIG. 6A

8/27

PREDICTED MOUSE GASP1 ALTERNATIVE NUCLEOTIDE SEQUENCE
SEQ ID NO:2

```

1  atgtgtgccc  cagggatatca  tcggttctgg  tttcactggg  ggctgctggt
51  gctgctgctc  ctcgaggctc  cccttcgagg  cctagcactg  ccacccatcc
101  gatactccca  tgcgggcatc  tgccccaacg  acatgaacc  caacctctgg
151  gtggatgccc  agagcacctg  caagcgagag  tgtgaaacag  accaggaatg
201  tgagacctat  gagaaatgct  gcccgaatgt  gtgtgggacc  aagagctgtg
251  tggcagcccg  ctacatggat  gtgaaagggg  agaagggggc  tgtaggcatg
301  cccaaggagg  ccacatgtga  ccatttcatg  tgccctgcag  agggctctga
351  gtgtgacatc  tgggacggcc  agcccgtgtg  taagtgcaaa  gatcgctgtg
401  agaaggagcc  cagcttcacc  tgtgcctctg  atggccttac  ctactacaac
451  cgttgcttca  tggacgccga  agcctgctcc  aagggcatca  cactgtctgt
501  ggtcacctgt  cgttatcact  tcacctggcc  taacaccagc  cctccaccgc
551  ctgagaccac  ggtgcatccc  accaccgcct  ctccggagac  tctcgggctg
601  gacatggcag  ccccagccct  gctcaaccac  cctgtccatc  agtcagtcac
651  cgtgggtgag  actgtgagtt  tcctctgtga  cgtggtaggc  cggcctcggc
701  cagagctcac  ttgggagaaa  cagctggagg  accgagagaa  tgttgtcatg
751  aggcccaacc  acgtgcgtgg  taatgtgggt  gtcactaaca  ttgcccagct
801  ggatcatctac  aacgtccagc  cccaggatgc  tggcatatac  acctgtacag
851  ctcgaaatgt  cgctgggtgtc  ctgagggctg  acttcccgtt  gtcgggtggc
901  aggggtggtc  aggccagggc  cacttcagag  agcagtctca  atggcacagc
951  ttttccagca  acagagtgcc  tgaagcccc  agacagtgag  gactgtggag
1001  aggagcagac  acgctggcac  ttcgacgccc  aggctaacia  ctgcctcact
1051  ttcacctttg  gccactgcca  ccacaatctc  aaccactttg  agacctacga
1101  ggctgtatg  ctggcttgta  tgagtggggc  attggccacc  tgcagcctgc
1151  ctgccctgca  agggccttgc  aaagcttatg  tcccacgctg  ggcttacaac
1201  agccagacag  gcctatgcca  gtccttcgtc  tatggcggct  gtgagggcaa
1251  cggtaacaac  tttgaaagcc  gtgaggcttg  tgaggagtcg  tgtcccttcc
1301  cgaggggtaa  ccagcactgc  cgggcctgca  agccccggca  aaaacttgtt
1351  accagcttct  gtcggagtga  ctttgtcatc  ctgggcaggg  tctctgagct
1401  gaccgaggag  caagactcgg  gccgtgccct  ggtgaccgtg  gatgaggtct
1451  taaaagatga  gaagatgggc  ctcaagtctc  tgggccggga  gcctctggaa
1501  gtcaccctgc  ttcattgtag  ctggacctgt  ccttgcccc  acgtgacagt
1551  ggggtgagaca  ccactcatca  tcatggggga  ggtggacggc  ggcatggcca
1601  tgctgagacc  cgatagcttt  gtggggggcat  cgagcacacg  gcgggtcagg
1651  aagctccgtg  aggtcatgta  caagaaaacc  tgtgacgtcc  tcaaggactt
1701  cctgggcttg  caatga

```

FIG. 6B

9/27

PREDICTED MOUSE GASP1 AMINO ACID SEQUENCE
SEQ ID NO:3

1 MCAPGYHRFW FHWGLLLLLL LEAPLRGLAL PPIRYSHAGI CPNDMNPNLW VDAQSTCKRE
61 CETDQECETY EKCCPNVCGT KSCVAARYMD VKGKKGPVGM PKEATCDHFM **CLQOGSECDI**
121 **WDGQPVCKCK** **DRCEKEPSFT** **CASDGLTYYN** **RCFMDAEACS** **KGITLSVVTG** RYHFTWPNTS
181 PPPPETTVHP TTASPETLGL DMAAPALLNH PVHQSVTVGE TVSFLCDVVG RPRPELTWEK
241 QLEDRENVVM RPNHVVRGNVV VTNIAQLVIY NVQPQDAGIY TCTARNVAGV LRADFPLSVV
301 RGGQARATSE SSLNGTAFPA TECLKPPDSE DCGEEQTRWH FDAQANNCLT FTFGHCHHNL
361 NHFETYEACM LACMSGPLAT CSLPALQGPC KAYVPRWAYN SQTGLCQSFV YGGCEGNGNN
421 FESREACEES CPFPRGNQHC RACKPRQKLV TSFCRSDFVI LGRVSELTEE QDSGRALVTV
481 DEVLKDEKMG LKFLGREPLE VTLLHVDWTC PCPNVTVGET PLIIMGEVDG GMAMLRPDSF
541 VGASSTRVR KLREVMYKKT CDVLKDFLGL Q

FIG. 6C

10/27

PREDICTED HUMAN GASP1 NUCLEOTIDE SEQUENCE
SEQ ID NO:4

```

1  atgaatccca acctctgggt ggacgcacag agcacctgca ggcgggagtg tgagacggac
61  caggagtgtg agatggacca ggtgagtggg atccagaagc cacagtgtga ggcagaccag
121  gtgaatgggg tccagaagcc gcaatgtgag atggaccaga agtgggagtg tgaggttgac
181  caggtgagtg ggggccagaa gccggtgtgt gaggcggacc aggtgagtgg ggtccagaag
241  ccacagtgtg agatggacca ggtgagtggg atccagaagc tggagtgtga ggcggaccag
301  aagtgggagt atgaggtgga ccaggtgagt ggggtccaga agccacagtg tgagatggac
361  caggtgagtg ggatccagaa gctggagtgt gaggcggacc aggagtgtga gacctatgag
421  aagtgctgcc ccaacgtatg tgggaccaag agctgcgtgg cggcccgcta catggacgtg
481  aaaggaaga agggcccagt gggcatgccc aaggaggcca catgtgacca cttcatgtgt
541  ctgcagcagg gctctgagtg tgacatctgg gatggccagc ccgtgtgtaa gtgcaaagac
601  cgctgtgaga aggagcccag ctttacctgc gcctcggacg gcctcaccta ctataaccgc
661  tgctacatgg atgccgaggc ctgctccaaa ggcacacac tggccgttgt aacctgccgc
721  taccacttca cntggcccaa caccagcccc ccaccacctg agaccaccat gcaccccacc
781  acagcctccc cagagacccc tgagctggac atggcggccc ctgcgctgct caacaaccct
841  gtgcaccagt cggtcaccat gggtgagaca gtgagcttcc tctgtgatgt ggtgggcccg
901  cccggcctg agatcacctg ggagaagcag ttggaggatc gggagaatgt ggtcatgcgg
961  cccaaccatg tgcgtggcaa cgtggtggtc accaacattg ccagctggt catctataac
1021  gccagctgc aggatgctgg gatctacacc tgcacggccc ggaacgtggc tggggtcctg
1081  agggctgatt tcccgctgtc ggtggtcagg ggtcatcagg ctgcagccac ctcagagagc
1141  agccccaatg gcacggcttt cccggcggcc gagtgcctga agccccaga cagtgaggac
1201  tgtggcgaag agcagaccag ctggcacttc gatgccagc ccaacaactg cctgaccttc
1261  accttcggcc actgccaccg taacctcaac cactttgaga cctatgaggc ctgcatgctg
1321  gcctgcatga gcgggcccgt ggccgcgtgc agcctgcccg ccctgcaggg gccctgcaaa
1381  gcctacgcgc ctgcctgggc ttacaacagc cagacgggcc agtgccagtc ctttgtctat
1441  ggtggctgcg agggcaatgg caacaacttt gagagccgtg aggctgtga ggagtcgtgc
1501  cccttcccc a gggggaacca gcgctgtcgg gcctgcaagc ctgcggcagaa gctcgttacc
1561  agcttctgtc gcagcgactt tgtcatcctg ggccgagtct ctgagctgac cgaggagcct
1621  gactcgggcc gcgccctggt gactgtggat gaggtcctaa aggatgagaa aatgggcctc
1681  aagttcctgg gccaggagcc attggaggtc actctgcttc acgtggactg ggcattgccc
1741  tgccccaacg tgaccgtgag cgagatgccg ctcatcatca tgggggaggt ggacggcggc
1801  atggccatgc tgcgccccga tagctttgtg ggcgcatcga gtgcccggcc ggtcaggaag
1861  cttcgtgagg tcatgcacaa gaagacctgt gacgtcctca aggagtttct tggcttgcac
1921  tga

```

FIG. 7A

11/27

PREDICTED AMINO ACID SEQUENCE OF HUMAN GASP1

SEQ ID NO:5

1 MNPNLWVDAQ STCRRECETD QCEMDQVSG IQKPQCEADQ VNGVQKPQCE MDQKWECEVD
61 QVSGVQKPVC EADQVSGVQK PQCEMDQVSG IQKLECEADQ KWEYEVDQVS GVQKPQCEMD
121 QVSGIQKLEC EADQECETYE KCCPNVCGTK SCVAARYMDV KGKKG P V G M P **KEATCDHFMC**
181 **LQQGSECDIW DGQPVCKCKD RCEKEPSFTC ASDGLTYYNR CYMDAEACSK GITLAVVTCR**
241 YHFTWPNTSP PPPETTMHPT TASPETPELD MAAPALLNNP VHQSVMGET VSFLCDVVGR
301 PRPEITWEKQ LEDRENVVMR PNHVRGNVVV TNIAQLVIYN AQLQDAGIYT CTARNVAGVL
361 RADFPLSVVR GHQAAATSES SPNGTAFPAA ECLKPPDSED CGEEQTRWHF DAQANNCLTF
421 TFGHCHRNLN HFETYEACML ACMSGPLAAC SLPALQGPK AYAPRWAYNS QTGQCQSFVY
481 GGCEGNGNNF ESREACEESC PFPRGNQRRCR ACKPRQKLV T SFCRSDFVIL GRVSELTEEP
541 DSGRALVTVD EVLKDEKMGL KFLGQEPLEV TLLHVDWACP CPNVTVSEMP LIIMGEVDGG
601 MAMLRPDSFV GASSARRVRK LREVMHKKTC DVLKEFLGLH

FIG. 7B

12/27

PREDICTED NUCLEOTIDE SEQUENCE OF HUMAN GASP1 USING AN ALTERNATIVE START SITE

SEQ ID NO:6

1 atgtgggccc caaggtgtcg ccggttctgg tctcgtggg agcaggtggc agcgtgctg
61 ctgctgctgc tactgctcgg ggtgcccccg cgaagcctgg cgctgccgcc catccgctat
121 tcccacgccg gcatctgccc caacgacatg aatcccaacc tctgggtgga cgcacagagc
181 acctgcaggc gggagtgtga gacggaccag gagtgtgaga cctatgagaa gtgctgcccc
241 aacgtatgtg ggaccaagag ctgctgtggc gcccgctaca tggacgtgaa agggaagaag
301 ggcccagtgg gcatgccc aa ggaggccaca tgtgaccact tcatgtgtct gcagcagggc
361 tctgagtgtg acatctggga tggccagccc gtgtgtaagt gcaaagaccg ctgtgagaag
421 gagcccagct ttacctgcgc ctcggacggc ctcacctact ataaccgctg ctacatggat
481 gccgaggcct gctccaaagg catcacactg gccgttgtaa cctgccgcta tcacttcacc
541 tggcccaaca ccagcccc accacctgag accaccatgc accccaccac agcctcccca
601 gagaccctg agctggacat ggcggcccct gcgctgctca acaaccctgt gcaccagtgc
661 gtcaccatgg gtgagacagt gagtttctc tgtgatgtgg tgggccggcc ccggcctgag
721 atcacctggg agaagcagtt ggaggatcgg gagaatgtgg tcatgcggcc caaccatgtg
781 cgtggcaacg tgggtggcac caacattgcc cagctggcca tctataacgc ccagctgcag
841 gatgctggga tctacacctg cacggcccgg aacgtggctg gggctctgag ggctgatttc
901 ccgctgtcgg tggtcagggg tcatcaggct gcagccacct cagagagcag cccaatggc
961 acggctttcc cggcggccga gtgcctgaag ccccagaca gtgaggactg tggcgaagag
1021 cagaccgct ggcacttcga tggccaggcc aacaactgcc tgacctcac cttcggccac
1081 tgccaccgta acctcaacca ctttgagacc tatgaggcct gcctgctggc ctgcatgagc
1141 gggccgctgg ccgctgtcag cctgcccgcc ctgcaggggc cctgcaaagc ctacgcgcct
1201 cgctgggctt acaacagcca gacgggccag tgccagtcct ttgtctatgg tggctgcgag
1261 ggcaatggca acaactttga gagccgtgag gcctgtgagg agtcgtgccc cttccccagg
1321 gggaaccagc gctgtcgggc ctgcaagcct cggcagaagc tcgttaccag cttctgtcgc
1381 agcgactttg tcatcctggg ccgagtctct gagctgaccg aggagcctga ctcgggccgc
1441 gccctggtga ctgtggatga ggtcctaaag gatgagaaaa tgggcctcaa gttcctgggc
1501 caggagccat tggaggtcac tctgcttcac gtggactggg catgcccctg cccaacgtg
1561 accgtgagcg agatgccgct catcatcatg ggggaggtgg acggcggcat ggccatgctg
1621 cgccccgata gctttgtggg cgcacatcagt gcccgccggg tcaggaagct tcgtgaggtc
1681 atgcacaaga agacctgtga cgtcctcaag gagtttcttg gcttgactg a

FIG. 7C

13/27

PREDICTED AMINO ACID SEQUENCE OF HUMAN GASP1 USING AN ALTERNATIVE START SITE

SEQ ID NO:7

1 MWAPRCRRFW SRWEQVAALL LLLLLLVVPP RSLALPPIRY SHAGICPNDM NPNLWVDAQS
61 TCRRECETDQ ECETYKCCP NVCGTKSCVA ARYMDVKGKK GPVGMPKEAT **CDHFMCLQQG**
121 **SECIDIWDGQP** **VCKCKDRCEK** **EPSFTCASDG** **LYYNYRCYMD** **AEACSKGITL** **AVVTCRYHFT**
181 WPNTSPPPPE TTMHPTTASP ETPELDMAAP ALLNNPVHQS VTMGETVSFL CDVVGRPRPE
241 ITWEKQLEDR ENVVMRPNHV RGNVVVTNIA QLVIYNAQLQ DAGIYTCTAR NVAGVLRADF
301 PLSVVRGHQA AATSESSPNG TAFPAAECLK PPDSEDCGEE QTRWHFDAQA NNCLTFTFGH
361 CHRNLNHFET YEACMLACMS GPLAACSLPA LQGPKAYAP RWAYNSQTGQ CQSFVYGGCE
421 GNGNMFESRE ACEESCPFPR GNQRCRACKP RQKLVTSFCR SDFVILGRVS ELTEEPDSGR
481 ALVTVDEVLK DEKMGLKFLG QEPLEVTLLH VDWACPCPNV TVSEMPLIIM GEVDGGMAML
541 RPDSFVGASS ARRVRKLREV MHKKTCDVLK EFLGLH*

FIG. 7D

14/27

PREDICTED MOUSE GASP2 NUCLEOTIDE SEQUENCE
SEQ ID NO:8

```
1 atgcctgccc cacagccatt cctgcctctg ctctttgtct tcgtgctcat ccatctgacc
61 tcggagacca acctgctgcc agatcccgga agccatcctg gcatgtgccc caacgagctc
121 agccccacc tgtgggtcga cgcccagagc acctgtgagc gtgagtgtac cggggaccag
181 gactgtgcgg catccgagaa gtgctgcacc aatgtgtgtg ggctgcagag ctgcgtggct
241 gcccgtttc ccagtggtag cccagctgta cctgagacag cagcctcctg tgaaggcttc
301 caatgccac aacagggttc tgactgtgac atctgggatg ggcagccagt ttgtcgctgc
361 cgtgaccgct gtgaaaaga acccagcttc acatgtgctt ctgatggcct tacctattac
421 aaccgctgct acatggacgc agaagcctgc ctgcggggtc tccacctgca cgttgtacct
481 tgtaagcaca ttctcagttg gccgcccagc agcccgggac caccgagac cactgctcgc
541 ccaaccctg gggctgctcc catgccacct gccctgtaca acagcccctc accacaggca
601 gtgcatgttg gggggacagc cagcctccac tgtgatgtta gtggccgtcc accacctgct
661 gtgacctggg agaagcagag ccatcagcgg gagaacctga tcatgcgccc tgaccaaagt
721 tatggcaacg tggttgtcac cagtatcggg cagctagtcc tctacaatgc tcagttggag
781 gatgcgggcc tgtatactg cactgcacga aacgctgccg gcctgctgcg ggccgacttt
841 cccctttccg ttttacagcg ggcaactact caggacaggg acccaggtat cccagccttg
901 gctgagtgcc aggccgacac acaagcctgt gttgggccac ctactcccca tcatgtcctt
961 tggcgctttg acccacagag aggcagctgc atgacattcc cagccctcag atgtgatggg
1021 gctgcccggg gctttgagac ctatgaggca tgccagcagg cctgtgttcg tggccccggg
1081 gatgtctgtg cactgcctgc agttcagggg ccctgccagg gctgggagcc acgctgggcc
1141 tacagcccac tgctacagca gtgccacccc tttgtataca gtggctgtga aggaaacagc
1201 aataactttg agaccggga gagctgtgag gatgcttgcc ctgtaccacg cacaccaccc
1261 tgctcgtgct gccgcctcaa gagcaagctg gctctgagct tgtgccgcag tgactttgcc
1321 atcgtgggga gactcacaga ggtcctggag gagcccagg ctgcaggcgg catagctcgt
1381 gtggccttgg atgatgtgct aaaggacgac aagatgggcc tcaagttctt gggcaccaaa
1441 tacctggagg tgacattgag tggcatggac tgggctgcc catgcccaca cgtgacagct
1501 gtcgatgggc cactggctcat catgggtgag gttcgtgaag gtgtggctgt gttggacgcc
1561 aacagctatg tccgtgctgc cagcgagaag cgagtcaaga agattgtgga actgctcgag
1621 aagaaggctt gtgaactgct caaccgcttc caagactag
```

FIG. 8A

15/27

PREDICTED MOUSE GASP2 AMINO ACID SEQUENCE
SEQ ID NO:9

1 MPAPQPFLPL LFVFLIHLT SETNLLPDPG SHPGMCPNEL SPHLWVDAQS TCERECTGDQ
61 DCAASEKCCT NVCGLQSCVA ARFPSGGPAV PETAA**SCEGF** **QCPQOGSDCD** **IWDGQPVCRG**
121 **RDRCEKEPSF** **TCASDGLTYT** **NRCYMDAEAC** **LRGLHLHVVP** CKHILSWPPS SPGPPETTAR
181 PTPGAAPMPP ALYNPSPQA VHVGGTASLH CDVSGRPPPA VTWEKQSHQR ENLIMRPDQM
241 YGNVVVTSIG QLVLYNAQLE DAGLYTCTAR NAAGLLRADF PLSVLQRATT QDRDPGIPAL
301 AECQADTQAC VGPPTPHHVL WRFDPQRGSC MTFPALRCDG AARGFETYEA CQQACVRGPG
361 DVCALPAVQG PCQGWEPWA YSPLLQQCHP FVYSGCEGNS NNFETRESCE DACPVPRTPP
421 CRACRLKSKL ALSLCRSDFA IVGRLTEVLE EPEAAGGIAR VALDDVLKDD KMGLKFLGTK
481 YLEVTLGMD WACPCPNVTA VDGPLVIMGE VREGVAVLDA NSYVRAASEK RVKKIVELLE
541 KKACELLNRF QD

FIG. 8B

16/27

PREDICTED NUCLEOTIDE SEQUENCE OF HUMAN GASP2

SEQ ID NO:10

1 atgcccgcc tacgtccact cctgccgctc ctgctcctcc tccggctgac ctcgggggct
61 ggcttgctgc cagggctggg gagccacccg ggcgtgtgcc ccaaccagct cagccccaac
121 ctgtgggtgg acgcccagag cacctgtgag cgcgagtgta gcagggacca ggactgtgcg
181 gctgctgaga agtgctgcat caacgtgtgt ggactgcaca gctgcgtggc agcacgcttc
241 cccggcagcc cagctgcgcc gacgacagcg gcctcctgcg agggctttgt gtgcccacag
301 cagggctcgg actgcgacat ctgggacggg cagcccgtgt gccgctgccg cgaccgctgt
361 gagaaggagc ccagcttcac ctgcgcctcg gacggcctca cctactaaa ccgctgctat
421 atggacgccg aggctgcct gcggggcctg cacctccaca tcgtgccttg caagcacgtg
481 ctcagctggc cgcccagcag cccggggccg ccggagacca ctgcccgcc cacacctggg
541 gccgcgcccg tgctcctgc cctgtacagc agcccctccc cacaggcggg gcaggttggg
601 ggtacggcca gcctccactg cgacgtcagc ggccgcccgc cgcctgctgt gacctgggag
661 aagcagagtc accagcgaga gaacctgatc atgcgccctg atcagatgta tggcaacgtg
721 gtggtcacca gcatcgggca gctggtgctc tacaacgcgc ggcccgaaga cgccggcctg
781 tacacctgca ccgcgcgcaa cgctgctggg ctgctgcggg ctgacttccc actctctgtg
841 gtccagcgag agccggccag ggacgcagcc cccagcatcc cagccccggc cgagtgcctg
901 ccggatgtgc aggctgcac gggcccact tccccacacc ttgtcctctg gcactacgac
961 ccgcagcggg gcggctgcat gacctcccg gcccggtggct gtgatggggc ggcccgcggc
1021 tttgagacct acgaggcatg ccagcaggcc tgtgcccgcg gccccggcga cgctgcgtg
1081 ctgcctgccg tgcagggccc ctgccggggc tgggagccgc gctgggccta cagcccgctg
1141 ctgcagcagt gccatccctt cgtgtacggg ggctgcgagg gcaacggcaa caacttccac
1201 agccgcgaga gctgcgagga tgctgcccc gtgccgcgca caccgccttg ccgcgcctgc
1261 cgctccgga gcaagctggc gctgagcctg tgccgcagcg acttcgcat cgtggggcgg
1321 ctcacggagg tgctggagga gcccgaggcc gccggcggca tcgcccgcgt ggcgctcgag
1381 gacgtgctca aggatgaaa gatgggcctc aagttcttgg gcaccaagta cctggagggtg
1441 acgctgagtg gcatggactg ggctgcccc tgcccaaca tgacggcggg cgacgggccc
1501 ctggtcatca tgggtgaggt gcgcgatggc gtggccgtgc tggacgccgg cagctacgtc
1561 cgcgccgcca gcgagaagcg cgtcaagaag atcttggagc tgctggagaa gcaggcctgc
1621 gagctgctca accgcttcca ggactagccc ccgcaggggc ctgcgccacc ccgtcctggt
1681 gaataaacgc actcc

FIG. 9A

17/27

PREDICTED AMINO ACID SEQUENCE OF HUMAN GASP-2

SEQ ID NO:11

1 MPALRPLLPL LLLRLTSGA GLLPGLGSHV GVCNQLSPN LWVDAQSTCE RECSRQDCA
61 AAEKCCINVC GLHSCVAARF PGSPAAPTTA **ASCEGFVCPQ** **QSDCDIWDG** **QPVCRCRDRG**
121 **EKEPSFTCAS** **DGLTYYNRCY** **MDAEACLRGL** **HLHIVPCKHV** LSWPPSSPGP PETTARPTPG
181 AAPVPPALYS SPSPQAVQVG GTASLHCDVS GRPPPAVTWE KQSHQRENLI MRPDQMYGNV
241 VVTSIGQLVL YNARPEDAGL YTCTARNAAG LLRADFPLSV VQREPARDAA PSIPAPAECL
301 PDVQACTGPT SPHLVLWHYD PQRGGCMTFP ARGCDGAARG FETYEACQQA CARGPGDACV
361 LPAVQGPCRG WEPRWAYSPL LQQCHPFVYG GCEGNGNNFH SRESCEDACP VPRTPPCRAC
421 RLRSKLALSL CRSDFAIVGR LTEVLEEPEA AGGIARVALE DVLKDDKMGL KFLGTKYLEV
481 TLSGMDWACP CPNMTAGDGP LVIMGEVRDG VAVLDAGSYV RAASEKRVKK ILELLEKQAC
541 ELLNRFQD

FIG. 9B

18/27




FIG. 10

19/27




FIG. 11B

FIG. 11A

20/27




FIG. 12A






FIG. 12B

21/27

**FIG. 12C****FIG. 12D**

22/27

CLONED MOUSE GASP1 NUCLEOTIDE AND AMINO ACID SEQUENCES
SEQ ID NO: 48

```

M C A P G Y H R F W F H W G L L L L L L L E A P L
1 ATGTGTGCCCCAGGG TATCATCGGTTCTGG TTCACTGGGGGCTG CTGTTGCTGCTGCTC CTCGAGGCTCCCCTT
R G L A L P P I R Y S H A G I C P N D M N P N L W
76 CGAGGCCTAGCACTG CCACCCATCCGATAC TCCCATGCGGGCATC TGCCCCAACGACATG AACCCCAACCTCTGG
V D A Q S T C K R E C E T D Q E C E T Y E K C C P
151 GTGGATGCCAGAGC ACCTGCAAGCGAGAG TGTGAAACAGACCAG GAATGTGAGACCTAT GAGAAATGCTGCCCC
N V C G T K S C V A A R Y M D V K G K K G P V G M
226 AATGTGTGTGGGACC AAGAGCTGTGTGGCA GCCCGCTACATGGAT GTGAAAGGGAAGAAG GGGCCTGTAGGCATG
P K E A T C D H F M C L Q Q G S E C D I W D G Q P
301 CCCAAGGAGGCCACA TGTGACCATTTTCATG TGCCTGCAGCAGGGC TCTGAGTGTGACATC TGGGACGGCCAGCCC
V C K C K D R C E K E P S F T C A S D G L T Y Y N
376 GTGTGTAAGTGCAAA GATCGCTGTGAGAAG GAGCCCAGCTTCACC TGTGCCTCTGATGGC CTTACCTACTACAAC
R C F M D A E A C S K G I T L S V V T C R Y H F T
451 CGTTGCTTCATGGAC GCCGAAGCCTGCTCC AAGGGCATCACACTG TCTGTGGTCACTGT CGTTATCACTTCACC
W P N T S P P P P E T T V H P T T A S P E T L G L
526 TGGCCTAACACCAGC CCTCCACCGCCTGAG ACCACGGTGCATCCC ACCACCGCCTCTCCG GAGACTCTCGGGCTG
D M A A P A L L N H P V H Q S V T V G E T V S F L
601 GACATGGCAGCCCCA GCCCTGCTCAACCAC CCTGTCCATCAGTCA GTCACCGTGGGTGAG ACTGTGAGTTTCCTC
C D V V G R P R P E L T W E K Q L E D R E N V V M
676 TGTGACGTGGTAGGC CGGCCTCGGCCAGAG CTCACTTGGGAGAAA CAGCTGGAGGACCGA GAGAATGTTGTCATG
R P N H V R G N V V V T N I A Q L V I Y N V Q P Q
751 AGGCCCAACCACGTG CGTGGTAATGTGGTG GTCACTAACATTGCC CAGCTGGTCACTAC AACGTCCAGCCCCAG
D A G I Y T C T A R N V A G V L R A D F P L S V V
826 GATGCTGGCATATAC ACCTGTACAGCTCGA AATGTCGCTGGTGTG CTGAGGGCTGACTTC CCGTTGTCGGTGGTC
R G G Q A R A T S E S S L N G T A F P A T E C L K
901 AGGGGTGGTCAGGCC AGGGCCACTTCAGAG AGCAGTCTCAATGGC ACAGCTTTTCCAGCA ACAGAGTGCCTGAAG
P P D S E D C G E E Q T R W H F D A Q A N N C L T
976 CCCCAGACAGTGAG GACTGTGGAGAGGAG CAGACACGCTGGCAC TTCGACGCCCAGGCT AACAACTGCCTCACT
F T F G H C H H N L N H F E T Y E A C M L A C M S
1051 TTCACCTTTGGCCAC TGCCACCACAATCTC AACCCTTTGAGACC TACGAGGCCTGTATG CTGGCTTGTATGAGT
G P L A T C S L P A L Q G P C K A Y V P R W A Y N
1126 GGGCATTGGCCACC TGCAGCCTGCCTGCC CTGCAAGGGCCTTGC AAAGCTTATGTCCA CGCTGGGCCTACAAC
S Q T G L C Q S F V Y G G C E G N G N N F E S R E
1201 AGCCAGACAGGCCTA TGCCAGTCCTTCGTC TATGGCGGCTGTGAG GGCAACGGTAACAAC TTTGAAAGCCGTGAG
A C E E S C P F P R G N Q H C R A C K P R Q K L V
1276 GCTTGTGAGGAGTCG TGTCCCTTCCCGAGG GGTAACCAGCACTGC CGGGCCTGCAAGCCC CGGCAAAAACCTTGT
T S F C R S D F V I L G R V S E L T E E Q D S G R
1351 ACCAGCTTCTGTCGG AGTGACTTTGTCATC CTGGGCAGGGTCTCT GAGCTGACCGAGGAG CAAGACTCGGGCCGT
A L V T V D E V L K D E K M G L K F L G R E P L E
1426 GCCCTGGTGACCGTG GATGAGGTCTTAAAA GATGAGAAGATGGGC CTCAAGTTTCTGGGC CGGGAGCCTCTGGAA
V T L L H V D W T C P C P N V T V G E T P L I I M
1501 GTCACCCTGCTTCAT GTAGACTGGACCTGT CCTTGCCCCAACGTG ACAGTGGGTGAGACA CCACTCATCATCATG
G E V D G G M A M L R P D S F V G A S S T R R V R
1576 GGGGAGGTGGACGGC GGCATGGCCATGCTG AGACCCGATAGCTTT GTGGGGGCATCGAGC ACACGGCGGGTCAGG
K L R E V M Y K K T C D V L K D F L G L Q *
1651 AAGCTCCGTGAGGTC ATGTACAAGAAAACC TGTGACGTCTCAAG GACTTCCTGGGCTTG CAATGA

```

FIG. 13



FIG. 14A




FIG. 14B




FIG. 15A

FIG. 15B

25/27

GDF-8




FIG. 16A

BMP-11




FIG. 16B




FIG. 16C






FIG. 16D

27/27



GASP-1 (nM)

FIG. 17



#	unique proteins in JA16
3	GASP1
5	GDF8 (unprocessed)
6	GDF8 (nearly full length)
7-9	GDF8 (propeptide region)
	FLRG
12-13	GDF8 (mature)