

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(10) International Publication Number
WO 2014/032184 A1

(43) International Publication Date
6 March 2014 (06.03.2014)

(51) International Patent Classification:

C07D 211/60 (2006.01) *C07H 5/04* (2006.01)
A61K 31/445 (2006.01) *C07H 7/00* (2006.01)
A61P 25/28 (2006.01) *C12N 9/24* (2006.01)

(21) International Application Number:

PCT/CA2013/050668

(22) International Filing Date:

29 August 2013 (29.08.2013)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/695,939 31 August 2012 (31.08.2012) US

(71) Applicants: **ALECTOS THERAPEUTICS INC.**
[CA/CA]; 8999 Nelson Way, Burnaby, British Columbia
V5A 4B5 (CA). **MERCK SHARP & DOHME CORP.**
[US/US]; 126 East Lincoln Avenue, Rahway, New Jersey
07065 (US).

(72) Inventors: **MCEACHERN, Ernest J.**; 8999 Nelson Way,
Burnaby, British Columbia V5A 4B5 (CA). **VOCADLO,
David J.**; 8999 Nelson Way, Burnaby, British Columbia
V5A 4B5 (CA). **ZHOU, Yuanxi**; 8999 Nelson Way, Burn-
aby, British Columbia V5A 4B5 (CA). **SELNICK, Harold
G.**; 770 Summeytown Pike, West Point, Pennsylvania
19486 (US).

(74) Agents: **NAUMAN, David** et al.; Borden Ladner Gervais
LLP, 1200 Waterfront Centre, 200 Burrard Street, P.O.
Box 48600, Vancouver, British Columbia V7X 1T2 (CA).

(81) Designated States (*unless otherwise indicated, for every
kind of national protection available*): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KN, KP, KR,
KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME,
MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ,
OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA,
SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM,
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM,
ZW.

(84) Designated States (*unless otherwise indicated, for every
kind of regional protection available*): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ,
UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,
TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,
EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,
MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))



WO 2014/032184 A1

(54) Title: GLYCOSIDASE INHIBITORS AND USES THEREOF

(57) Abstract: The invention provides compounds for inhibiting glycosidases, prodrugs of the compounds, and pharmaceutical compositions including the compounds or prodrugs of the compounds. The invention also provides methods of treating diseases and disorders related to deficiency or overexpression of O-GlcNAcase, accumulation or deficiency of O-GlcNAc.

GLYCOSIDASE INHIBITORS AND USES THEREOF

FIELD OF THE INVENTION

[0001] This application relates to compounds which inhibit glycosidases and uses thereof.

5

BACKGROUND OF THE INVENTION

[0002] A wide range of cellular proteins, both nuclear and cytoplasmic, are post-translationally modified by the addition of the monosaccharide 2-acetamido-2-deoxy- β -D-glucopyranoside (β -N-acetylglucosamine) which is attached via an O-glycosidic linkage.¹

10 This modification is generally referred to as O-linked N-acetylglucosamine or O-GlcNAc. The enzyme responsible for post-translationally linking β -N-acetylglucosamine (GlcNAc) to specific serine and threonine residues of numerous nucleocytoplasmic proteins is O-GlcNAc transferase (OGT).²⁻⁵ A second enzyme, known as glycoprotein 2-acetamido-2-deoxy- β -D-glucopyranosidase (O-GlcNAcase)^{6,7} removes this post-translational modification to liberate
15 proteins making the O-GlcNAc-modification a dynamic cycle occurring several times during the lifetime of a protein.⁸

[0003] O-GlcNAc-modified proteins regulate a wide range of vital cellular functions including, for example, transcription,⁹⁻¹² proteasomal degradation,¹³ and cellular signaling.¹⁴ O-GlcNAc is also found on many structural proteins.¹⁵⁻¹⁷ For example, it has been found on a
20 number of cytoskeletal proteins, including neurofilament proteins,^{18,19} synapsins,^{6,20} synapsin-specific clathrin assembly protein AP-3,⁷ and ankyrinG.¹⁴ O-GlcNAc modification has been found to be abundant in the brain.^{21,22} It has also been found on proteins clearly implicated in the etiology of several diseases including Alzheimer's disease (AD) and cancer.

[0004] For example, it is well established that AD and a number of related tauopathies
25 including Down's syndrome, Pick's disease, Niemann-Pick Type C disease, and amyotrophic lateral sclerosis (ALS) are characterized, in part, by the development of neurofibrillary tangles (NFTs). These NFTs are aggregates of paired helical filaments (PHFs) and are composed of an abnormal form of the cytoskeletal protein "tau". Normally tau stabilizes a key cellular network of microtubules that is essential for distributing proteins and nutrients
30 within neurons. In AD patients, however, tau becomes hyperphosphorylated, disrupting its normal functions, forming PHFs and ultimately aggregating to form NFTs. Six isoforms of

tau are found in the human brain. In AD patients, all six isoforms of tau are found in NFTs, and all are markedly hyperphosphorylated.^{23,24} Tau in healthy brain tissue bears only 2 or 3 phosphate groups, whereas those found in the brains of AD patients bear, on average, 8 phosphate groups.^{25,26} A clear parallel between NFT levels in the brains of AD patients and the severity of dementia strongly supports a key role for tau dysfunction in AD.²⁷⁻²⁹ The precise causes of this hyperphosphorylation of tau remain elusive. Accordingly, considerable effort has been dedicated toward: a) elucidating the molecular physiological basis of tau hyperphosphorylation;³⁰ and b) identifying strategies that could limit tau hyperphosphorylation in the hope that these might halt, or even reverse, the progression of Alzheimer's disease³¹⁻³⁴ Thus far, several lines of evidence suggest that up-regulation of a number of kinases may be involved in hyperphosphorylation of tau,^{21,35,36} although very recently, an alternative basis for this hyperphosphorylation has been advanced.²¹

[0005] In particular, it has emerged that phosphate levels of tau are regulated by the levels of O-GlcNAc on tau. The presence of O-GlcNAc on tau has stimulated studies that correlate O-GlcNAc levels with tau phosphorylation levels. The interest in this field stems from the observation that O-GlcNAc modification has been found to occur on many proteins at amino acid residues that are also known to be phosphorylated.³⁷⁻³⁹ Consistent with this observation, it has been found that increases in phosphorylation levels result in decreased O-GlcNAc levels and conversely, increased O-GlcNAc levels correlate with decreased phosphorylation levels.⁴⁰ This reciprocal relationship between O-GlcNAc and phosphorylation has been termed the "Yin-Yang hypothesis"⁴¹ and has gained strong biochemical support by the discovery that the enzyme OGT⁴ forms a functional complex with phosphatases that act to remove phosphate groups from proteins.⁴² Like phosphorylation, O-GlcNAc is a dynamic modification that can be removed and reinstalled several times during the lifespan of a protein. Suggestively, the gene encoding O-GlcNAcase has been mapped to a chromosomal locus that is linked to AD.^{7,43} Hyperphosphorylated tau in human AD brains has markedly lower levels of O-GlcNAc than are found in healthy human brains.²¹ It has been shown that O-GlcNAc levels of soluble tau protein from human brains affected with AD are markedly lower than those from healthy brain.²¹ Furthermore, PHF from diseased brain was suggested to lack completely any O-GlcNAc modification whatsoever.²¹ The molecular basis of this hypoglycosylation of tau is not known, although it may stem from increased activity of kinases and/or dysfunction of one of the enzymes involved in processing O-GlcNAc. Supporting this latter view, in both PC-12 neuronal cells and in brain tissue sections from

mice, a nonselective N-acetylglucosaminidase inhibitor was used to increase tau O-GlcNAc levels, whereupon it was observed that phosphorylation levels decreased.²¹ The implication of these collective results is that by maintaining healthy O-GlcNAc levels in AD patients, such as by inhibiting the action of O-GlcNAcase, one should be able to block

5 hyperphosphorylation of tau and all of the associated effects of tau hyperphosphorylation, including the formation of NFTs and downstream effects. However, because the proper functioning of the β -hexosaminidases is critical, any potential therapeutic intervention for the treatment of AD that blocks the action of O-GlcNAcase would have to avoid the concomitant inhibition of both hexosaminidases A and B.

10 [0006] Neurons do not store glucose and therefore the brain relies on glucose supplied by blood to maintain its essential metabolic functions. Notably, it has been shown that within brain, glucose uptake and metabolism decreases with aging.⁴⁴ Within the brains of AD patients marked decreases in glucose utilization occur and are thought to be a potential cause of neurodegeneration.⁴⁵ The basis for this decreased glucose supply in AD brain⁴⁶⁻⁴⁸ is
15 thought to stem from any of decreased glucose transport,^{49,50} impaired insulin signaling,^{51,52} and decreased blood flow.⁵³

[0007] In light of this impaired glucose metabolism, it is worth noting that of all glucose entering into cells, 2-5% is shunted into the hexosamine biosynthetic pathway, thereby regulating cellular concentrations of the end product of this pathway, uridine diphosphate-N-
20 acetylglucosamine (UDP-GlcNAc).⁵⁴ UDP-GlcNAc is a substrate of the nucleocytoplasmic enzyme O-GlcNAc transferase (OGT),²⁻⁵ which acts to post-translationally add GlcNAc to specific serine and threonine residues of numerous nucleocytoplasmic proteins. OGT recognizes many of its substrates^{55,56} and binding partners^{42,57} through its tetratricopeptide repeat (TPR) domains.^{58,59} As described above, O-GlcNAcase^{6,7} removes this post-
25 translational modification to liberate proteins making the O-GlcNAc-modification a dynamic cycle occurring several times during the lifetime of a protein.⁸ O-GlcNAc has been found in several proteins on known phosphorylation sites,^{10,38,39,60} including tau and neurofilaments.⁶¹ Additionally, OGT shows unusual kinetic behaviour making it exquisitely sensitive to intracellular UDP-GlcNAc substrate concentrations and therefore glucose supply.⁴²

30 [0008] Consistent with the known properties of the hexosamine biosynthetic pathway, the enzymatic properties of OGT, and the reciprocal relationship between O-GlcNAc and phosphorylation, it has been shown that decreased glucose availability in brain leads to tau hyperphosphorylation.⁴⁵ Therefore the gradual impairment of glucose transport and

metabolism, whatever its causes, leads to decreased O-GlcNAc and hyperphosphorylation of tau (and other proteins). Accordingly, the inhibition of O-GlcNAcase should compensate for the age related impairment of glucose metabolism within the brains of health individuals as well as patients suffering from AD or related neurodegenerative diseases.

5 [0009] These results suggest that a malfunction in the mechanisms regulating tau O-GlcNAc levels may be vitally important in the formation of NFTs and associated neurodegeneration. Good support for blocking tau hyperphosphorylation as a therapeutically useful intervention⁶² comes from recent studies showing that when transgenic mice harbouring human tau are treated with kinase inhibitors, they do not develop typical motor defects³⁴ and, in another case,³³ show decreased levels of insoluble tau. These studies provide a clear link between
10 lowering tau phosphorylation levels and alleviating AD-like behavioural symptoms in a murine model of this disease. Indeed, pharmacological modulation of tau hyperphosphorylation is widely recognized as a valid therapeutic strategy for treating AD and other neurodegenerative disorders.⁶³

15 [0010] Small-molecule O-GlcNAcase inhibitors, to limit tau hyperphosphorylation, have been considered for treatment of AD and related tauopathies.⁶⁴ Specifically, the O-GlcNAcase inhibitor thiamet-G has been implicated in the reduction of tau phosphorylation in cultured PC-12 cells at pathologically relevant sites.⁶⁴ Moreover, oral administration of thiamet-G to healthy Sprague-Dawley rats has been implicated in reduced phosphorylation of
20 tau at Thr231, Ser396 and Ser422 in both rat cortex and hippocampus.⁶⁴

[0011] There is also a large body of evidence indicating that increased levels of O-GlcNAc protein modification provides protection against pathogenic effects of stress in cardiac tissue, including stress caused by ischemia, hemorrhage, hypervolemic shock, and calcium paradox. For example, activation of the hexosamine biosynthetic pathway (HBP) by administration of
25 glucosamine has been demonstrated to exert a protective effect in animals models of ischemia/reperfusion,⁶⁵⁻⁷¹ trauma hemorrhage,⁷²⁻⁷⁴ hypervolemic shock,⁷⁵ and calcium paradox.^{65,76} Moreover, strong evidence indicates that these cardioprotective effects are mediated by elevated levels of protein O-GlcNAc modification.^{65,66,68,71,73,76-79} There is also evidence that the O-GlcNAc modification plays a role in a variety of neurodegenerative
30 diseases, including Parkinson's disease and Huntington's disease.⁸⁰

[0012] Humans have three genes encoding enzymes that cleave terminal β -N-acetylglucosamine residues from glycoconjugates. The first of these encodes O-GlcNAcase. O-

GlcNAcase is a member of family 84 of glycoside hydrolases that includes enzymes from organisms as diverse as prokaryotic pathogens to humans (for the family classification of glycoside hydrolases see Coutinho, P.M. & Henrissat, B. (1999) Carbohydrate-Active Enzymes server at URL: <http://afmb.cnrs-mrs.fr/CAZY/>.^{81,82} O-GlcNAcase acts to hydrolyse O-GlcNAc off of serine and threonine residues of post-translationally modified proteins.^{1,6,7,83,84} Consistent with the presence of O-GlcNAc on many intracellular proteins, the enzyme O-GlcNAcase appears to have a role in the etiology of several diseases including type II diabetes,^{14,85} AD,^{16,21,86} and cancer.^{22,87} Although O-GlcNAcase was likely isolated earlier on,^{18,19} about 20 years elapsed before its biochemical role in acting to cleave O-GlcNAc from serine and threonine residues of proteins was understood.⁶ More recently O-GlcNAcase has been cloned,⁷ partially characterized,²⁰ and suggested to have additional activity as a histone acetyltransferase.²⁰ However, little was known about the catalytic mechanism of this enzyme.

[0013] The other two genes, HEXA and HEXB, encode enzymes catalyzing the hydrolytic cleavage of terminal β -N-acetylglucosamine residues from glycoconjugates. The gene products of HEXA and HEXB predominantly yield two dimeric isozymes, hexosaminidase A and hexosaminidase B, respectively. Hexosaminidase A ($\alpha\beta$), a heterodimeric isozyme, is composed of an α - and a β -subunit. Hexosaminidase B ($\beta\beta$), a homodimeric isozyme, is composed of two β -subunits. The two subunits, α - and β -, bear a high level of sequence identity. Both of these enzymes are classified as members of family 20 of glycoside hydrolases and are normally localized within lysosomes. The proper functioning of these lysosomal β -hexosaminidases is critical for human development, a fact that is underscored by the tragic genetic illnesses, Tay-Sach's and Sandhoff diseases which stem from a dysfunction in, respectively, hexosaminidase A and hexosaminidase B.⁸⁸ These enzymatic deficiencies cause an accumulation of glycolipids and glycoconjugates in the lysosomes resulting in neurological impairment and deformation. The deleterious effects of accumulation of gangliosides at the organismal level are still being uncovered.⁸⁹

[0014] As a result of the biological importance of these β -N-acetyl-glucosaminidases, small molecule inhibitors of glycosidases⁹⁰⁻⁹³ have received a great deal of attention,⁹⁴ both as tools for elucidating the role of these enzymes in biological processes and in developing potential therapeutic applications. The control of glycosidase function using small molecules offers several advantages over genetic knockout studies including the ability to rapidly vary doses or to entirely withdraw treatment.

[0015] However, a major challenge in developing inhibitors for blocking the function of mammalian glycosidases, including O-GlcNAcase, is the large number of functionally related enzymes present in tissues of higher eukaryotes. Accordingly, the use of non-selective inhibitors in studying the cellular and organismal physiological role of one particular enzyme is complicated because complex phenotypes arise from the concomitant inhibition of such functionally related enzymes. In the case of β -N-acetylglucosaminidases, many compounds that act to block O-GlcNAcase function are non-specific and act potently to inhibit the lysosomal β -hexosaminidases.

[0016] A few of the better characterized inhibitors of β -N-acetyl-glucosaminidases which have been used in studies of O-GlcNAc post-translational modification within both cells and tissues are streptozotocin (STZ), 2'-methyl- α -D-glucopyrano-[2,1-d]- Δ 2'-thiazoline (NAG-thiazoline) and O-(2-acetamido-2-deoxy-D-glucopyranosylidene)amino N-phenylcarbamate (PUGNAc).^{14,95-98}

[0017] STZ has long been used as a diabetogenic compound because it has a particularly detrimental effect on β -islet cells.⁹⁹ STZ exerts its cytotoxic effects through both the alkylation of cellular DNA^{99,100} as well as the generation of radical species including nitric oxide.¹⁰¹ The resulting DNA strand breakage promotes the activation of poly(ADP-ribose) polymerase (PARP)¹⁰² with the net effect of depleting cellular NAD⁺ levels and, ultimately, leading to cell death.^{103,104} Other investigators have proposed instead that STZ toxicity is a consequence of the irreversible inhibition of O-GlcNAcase, which is highly expressed within β -islet cells.^{95,105} This hypothesis has, however, been brought into question by two independent research groups.^{106,107} Because cellular O-GlcNAc levels on proteins increase in response to many forms of cellular stress¹⁰⁸ it seems possible that STZ results in increased O-GlcNAc-modification levels on proteins by inducing cellular stress rather than through any specific and direct action on O-GlcNAcase. Indeed, Hanover and coworkers have shown that STZ functions as a poor and somewhat selective inhibitor of O-GlcNAcase¹⁰⁹ and although it has been proposed by others that STZ acts to irreversibly inhibit O-GlcNAcase,¹¹⁰ there has been no clear demonstration of this mode of action. More recently, it has been shown that STZ does not irreversibly inhibit O-GlcNAcase.¹¹¹

[0018] NAG-thiazoline has been found to be a potent inhibitor of family 20 hexosaminidases,^{93,112} and more recently, the family 84 O-GlcNAcases.¹¹¹ Despite its potency, a downside to using NAG-thiazoline in a complex biological context is that it lacks selectivity and therefore perturbs multiple cellular processes.

[0019] PUGNAc is another compound that suffers from the same problem of lack of selectivity, yet has enjoyed use as an inhibitor of both human O-GlcNAcase^{6,113} and the family 20 human β -hexosaminidases.¹¹⁴ This molecule, developed by Vasella and coworkers, was found to be a potent competitive inhibitor of the β -*N*-acetyl-glucosaminidases from *Canavalia ensiformis*, *Mucor rouxii*, and the β -hexosaminidase from bovine kidney.⁹¹ It has been demonstrated that administration of PUGNAc in a rat model of trauma hemorrhage decreases circulating levels of the pro-inflammatory cytokines TNF- α and IL-6.¹¹⁵ It has also been shown that administration of PUGNAc in a cell-based model of lymphocyte activation decreases production of the cytokine IL-2.¹¹⁶ Subsequent studies have indicated that PUGNAc can be used in an animal model to reduce myocardial infarct size after left coronary artery occlusions.¹¹⁷ Of particular significance is the fact that elevation of O-GlcNAc levels by administration of PUGNAc, an inhibitor of O-GlcNAcase, in a rat model of trauma hemorrhage improves cardiac function.^{115,118} In addition, elevation of O-GlcNAc levels by treatment with PUGNAc in a cellular model of ischemia/reperfusion injury using neonatal rat ventricular myocytes improved cell viability and reduced necrosis and apoptosis compared to untreated cells.¹¹⁹


[0020] More recently, it has been suggested that the selective O-GlcNAcase inhibitor NButGT exhibits protective activity in cell-based models of ischemia/reperfusion and cellular stresses, including oxidative stress.¹²⁰ This study suggests the use of O-GlcNAcase inhibitors to elevate protein O-GlcNAc levels and thereby prevent the pathogenic effects of stress in cardiac tissue.

[0021] International patent applications PCT/CA2006/000300, filed 1 March 2006, published under No. WO 2006/092049 on 8 September 2006; PCT/CA2007/001554, filed 31 August 2007, published under No. WO 2008/025170 on 6 March 2008; PCT/CA2009/001087, filed 31 July 2009, published under No. WO 2010/012106 on 4 February 2010; PCT/CA2009/001088, filed 31 July 2009, published under WO 2010/012107 on 4 February 2010; PCT/CA2009/001302, filed 16 September 2009, published under WO 2010/037207 on 8 April 2010; PCT/CA2011/000548, filed 10 May 2011, published under No. WO 2011/140640 on 17 November 2011; PCT/CA/2011/001241, filed 8 November 2011, published under WO 2012/061927 on 18 May 2012; PCT/US2011/059668, filed 8 November 2011, published under WO 2012/064680 on 18 May 2012; and PCT/CA2011/001397, filed 21 December 2011, published under WO 2012/083435 on 28 June 2012, describe selective inhibitors of O-GlcNAcase.

SUMMARY OF THE INVENTION

[0022] The invention provides, in part, compounds for inhibiting glycosidases, prodrugs of the compounds, uses of the compounds and the prodrugs, pharmaceutical compositions including the compounds or prodrugs of the compounds, and methods of treating diseases and disorders related to deficiency or overexpression of O-GlcNAcase, and/or accumulation or deficiency of O-GlcNAc.

[0023] In one aspect, the invention provides a compound of Formula (I) or a pharmaceutically acceptable salt thereof.



(I)

where R^1 may be H and R^2 may be $C(O)NR^{11}_2$, or R^1 may be $C(O)NR^{11}_2$ and R^2 may be H, or R^1 may be H and R^2 may be selected from the group consisting of: H, CH_3 , CH_2F , CHF_2 , and CH_2OH , where each R^{11} may be independently H or C_{1-6} alkyl; R^3 may be H and R^4 may be OH, or R^3 may be H and R^4 may be H, or R^3 may be H and R^4 may be F, or R^3 may be F and R^4 may be H, or R^3 may be F and R^4 may be F, or R^3 may be OH and R^4 may be H; R^5 may be H and R^6 may be OH, or R^5 may be H and R^6 may be H, or R^5 may be H and R^6 may be F, or R^5 may be F and R^6 may be H, or R^5 may be F and R^6 may be F, or R^5 may be OH and R^6 may be H; R^7 may be H and R^8 may be OH, or R^7 may be H and R^8 may be H, or R^7 may be H and R^8 may be F, or R^7 may be F and R^8 may be H, or R^7 may be F and R^8 may be F, or R^7 may be OH and R^8 may be H; R^9 may be selected from the group consisting of: H, CH_3 , CH_2F , CHF_2 , and CH_2OH ; and R^{10} may be selected from the group consisting of: H, C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, C_{1-10} acyl, C_{8-20} arylalkylacyl, C_{3-20} heteroarylalkylacyl, C_{7-20} arylalkyl, C_{8-20} arylalkenyl, C_{8-20} arylalkynyl, C_{2-20} heteroarylalkyl, C_{3-20} heteroarylalkenyl, and C_{3-20} heteroarylalkynyl, each excluding H optionally substituted from one to four substituents with one of more of halo, OH, OCF_3 , CN, SO_2NH_2 , SO_2Me , $C(O)NH_2$, CH_2F , CHF_2 , CF_3 , CH_2CH_2F , CH_2CF_3 , $CH_2CH_2CH_2F$, C_{1-6} alkyl, C_{1-6} alkoxy, or C_{3-7} cycloalkyl.

[0024] In alternative embodiments, the invention provides a compound of Formula (I) in which at least one of R^3 , R^4 , R^5 , R^6 , R^7 , or R^8 may be OH; or R^2 may be CH_2OH ; or R^9 may be CH_2OH .

5 [0025] In alternative embodiments, the invention provides a compound of Formula (I) in which at least one of R^3 , R^4 , R^5 , R^6 , R^7 , or R^8 may be F; or R^2 may be CH_2F or CHF_2 ; or R^9 may be CH_2F or CHF_2 .

[0026] In alternative embodiments, the invention provides a compound of Formula (I) in which R^2 may be CH_3 ; or R^3 and R^4 may be H; or R^5 and R^6 may be H; or R^7 and R^8 may be H; or R^9 may be CH_3 .

10 [0027] In alternative embodiments, the invention provides a compound of Formula (I) in which R^1 and R^2 may be H, or R^9 may be H.

[0028] In alternative embodiments, the invention provides a compound of Formula (I) in which R^1 and R^2 may be other than H.

15 [0029] In alternative embodiments, the invention provides a compound of Formula (I) in which R^9 may be other than H.

[0030] In alternative embodiments, the invention provides a compound of Formula (I) in which at least one of R^1 , R^2 , or R^9 may be other than H.

[0031] In alternative embodiments, the invention provides a compound of Formula (I) in which at least one of R^3 , R^4 , R^5 , R^6 , R^7 , or R^8 may be other than H.

20 [0032] In alternative embodiments, the invention provides a compound of Formula (I) in which at least one of R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , or R^9 may be other than H.

[0033] In alternative embodiments, the invention provides a compound of Formula (I) in which at least one of R^2 or R^9 may be other than CH_3 .

25 [0034] In alternative embodiments, the invention provides a compound of Formula (I) in which when R^2 may be CH_2OH , then R^9 may be other than H.

[0035] In alternative embodiments, the invention provides a compound of Formula (I) in which when R^2 may be CH_3 , then R^9 may be other than H.


[0036] In alternative embodiments, the invention provides a compound of Formula (I) in which at least one of R^3 or R^4 may be other than H.

[0037] In alternative embodiments, the invention provides a compound of Formula (I) in which at least one of R⁵ or R⁶ may be other than H.

[0038] In alternative embodiments, the invention provides a compound of Formula (I) in which at least one of R⁷ or R⁸ may be other than H.

5 [0039] In alternative embodiments, the invention provides a compound of Formula (I) in which R¹⁰ may be other than H.

[0040] In alternative embodiments, the invention provides a compound of Formula (Ia) or a pharmaceutically acceptable salt thereof:




10

(Ia)

where each R¹¹ may be independently H or C₁₋₆ alkyl; R³ may be H and R⁴ may be OH, or R³ may be H and R⁴ may be H, or R³ may be H and R⁴ may be F, or R³ may be F and R⁴ may be H, or R³ may be F and R⁴ may be F, or R³ may be OH and R⁴ may be H; R⁵ may be H and R⁶ may be OH, or R⁵ may be H and R⁶ may be H, or R⁵ may be H and R⁶ may be F, or R⁵ may be F and R⁶ may be H, or R⁵ may be F and R⁶ may be F, or R⁵ may be OH and R⁶ may be H; R⁷ may be H and R⁸ may be OH, or R⁷ may be H and R⁸ may be H, or R⁷ may be H and R⁸ may be F, or R⁷ may be F and R⁸ may be H, or R⁷ may be F and R⁸ may be F, or R⁷ may be OH and R⁸ may be H; R⁹ may be selected from the group consisting of: H, CH₃, CH₂F, CHF₂, and CH₂OH; and R¹⁰ may be selected from the group consisting of: H, C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl, C₁₋₁₀ acyl, C₈₋₂₀ arylalkylacyl, C₃₋₂₀ heteroarylalkylacyl, C₇₋₂₀ arylalkyl, C₈₋₂₀ arylalkenyl, C₈₋₂₀ arylalkynyl, C₂₋₂₀ heteroarylalkyl, C₃₋₂₀ heteroarylalkenyl, and C₃₋₂₀ heteroarylalkynyl, each excluding H optionally substituted from one to four substituents with one of more of halo, OH, OCF₃, CN, SO₂NH₂, SO₂Me, C(O)NH₂, CH₂F, CHF₂, CF₃, CH₂CH₂F, CH₂CF₃, CH₂CH₂CH₂F, C₁₋₆ alkyl, C₁₋₆ alkoxy, or C₃₋₇ cycloalkyl.


25 [0041] In alternative embodiments, the invention provides a compound of Formula (Ib) or a pharmaceutically acceptable salt thereof:



(Ib)

where each R¹¹ may be independently H or C₁₋₆ alkyl; R³ may be H and R⁴ may be OH, or R³ may be H and R⁴ may be H, or R³ may be H and R⁴ may be F, or R³ may be F and R⁴ may be H, or R³ may be F and R⁴ may be F, or R³ may be OH and R⁴ may be H; R⁵ may be H and R⁶ may be OH, or R⁵ may be H and R⁶ may be H, or R⁵ may be H and R⁶ may be F, or R⁵ may be F and R⁶ may be H, or R⁵ may be F and R⁶ may be F, or R⁵ may be OH and R⁶ may be H; R⁷ may be H and R⁸ may be OH, or R⁷ may be H and R⁸ may be H, or R⁷ may be H and R⁸ may be F, or R⁷ may be F and R⁸ may be H, or R⁷ may be F and R⁸ may be F, or R⁷ may be OH and R⁸ may be H; R⁹ may be selected from the group consisting of: H, CH₃, CH₂F, CHF₂, and CH₂OH; and R¹⁰ may be selected from the group consisting of: H, C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl, C₁₋₁₀ acyl, C₈₋₂₀ arylalkylacyl, C₃₋₂₀ heteroarylalkylacyl, C₇₋₂₀ arylalkyl, C₈₋₂₀ arylalkenyl, C₈₋₂₀ arylalkynyl, C₂₋₂₀ heteroarylalkyl, C₃₋₂₀ heteroarylalkenyl, and C₃₋₂₀ heteroarylalkynyl, each excluding H optionally substituted from one to four substituents with one or more of halo, OH, OCF₃, CN, SO₂NH₂, SO₂Me, C(O)NH₂, CH₂F, CHF₂, CF₃, CH₂CH₂F, CH₂CF₃, CH₂CH₂CH₂F, C₁₋₆ alkyl, C₁₋₆ alkoxy, or C₃₋₇ cycloalkyl.

[0042] In alternative embodiments, the invention provides a compound of Formula (Ic) or a pharmaceutically acceptable salt thereof:




(Ic)

where R² may be selected from the group consisting of: H, CH₃, CH₂F, CHF₂, CH₂OH, and C(O)NR¹¹₂, where each R¹¹ may be independently H or C₁₋₆ alkyl; R³ may be H and R⁴ may be OH, or R³ may be H and R⁴ may be H, or R³ may be H and R⁴ may be F, or R³

may be F and R⁴ may be H, or R³ may be F and R⁴ may be F, or R³ may be OH and R⁴ may be H; R⁵ may be H and R⁶ may be OH, or R⁵ may be H and R⁶ may be H, or R⁵ may be H and R⁶ may be F, or R⁵ may be F and R⁶ may be H, or R⁵ may be F and R⁶ may be F, or R⁵ may be OH and R⁶ may be H; R⁷ may be H and R⁸ may be OH, or R⁷ may be H and R⁸ may be H, or R⁷ may be H and R⁸ may be F, or R⁷ may be F and R⁸ may be H, or R⁷ may be F and R⁸ may be F, or R⁷ may be OH and R⁸ may be H; R⁹ may be selected from the group consisting of: H, CH₃, CH₂F, CHF₂, and CH₂OH; and R¹⁰ may be selected from the group consisting of: H, C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl, C₁₋₁₀ acyl, C₈₋₂₀ arylalkylacyl, C₃₋₂₀ heteroarylalkylacyl, C₇₋₂₀ arylalkyl, C₈₋₂₀ arylalkenyl, C₈₋₂₀ arylalkynyl, C₂₋₂₀ heteroarylalkyl, C₃₋₂₀ heteroarylalkenyl, and C₃₋₂₀ heteroarylalkynyl, each excluding H optionally substituted from one to four substituents with one of more of halo, OH, OCF₃, CN, SO₂NH₂, SO₂Me, C(O)NH₂, CH₂F, CHF₂, CF₃, CH₂CH₂F, CH₂CF₃, CH₂CH₂CH₂F, C₁₋₆ alkyl, C₁₋₆ alkoxy, or C₃₋₇ cycloalkyl.

[0043] In alternative embodiments, the invention provides a compound of Formula (Id) or a pharmaceutically acceptable salt thereof:




(Id)

where R³ may be H and R⁴ may be OH, or R³ may be H and R⁴ may be H, or R³ may be H and R⁴ may be F, or R³ may be F and R⁴ may be H, or R³ may be F and R⁴ may be F, or R³ may be OH and R⁴ may be H; R⁵ may be H and R⁶ may be OH, or R⁵ may be H and R⁶ may be H, or R⁵ may be H and R⁶ may be F, or R⁵ may be F and R⁶ may be H, or R⁵ may be F and R⁶ may be F, or R⁵ may be OH and R⁶ may be H; R⁷ may be H and R⁸ may be OH, or R⁷ may be H and R⁸ may be H, or R⁷ may be H and R⁸ may be F, or R⁷ may be F and R⁸ may be H, or R⁷ may be F and R⁸ may be F, or R⁷ may be OH and R⁸ may be H; R⁹ may be selected from the group consisting of: H, CH₃, CH₂F, CHF₂, and CH₂OH; and R¹⁰ may be selected from the group consisting of: H, C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl, C₁₋₁₀ acyl, C₈₋₂₀ arylalkylacyl, C₃₋₂₀ heteroarylalkylacyl, C₇₋₂₀ arylalkyl, C₈₋₂₀ arylalkenyl, C₈₋₂₀ arylalkynyl, C₂₋₂₀ heteroarylalkyl, C₃₋₂₀ heteroarylalkenyl, and C₃₋₂₀ heteroarylalkynyl, each

excluding H optionally substituted from one to four substituents with one of more of halo, OH, OCF₃, CN, SO₂NH₂, SO₂Me, C(O)NH₂, CH₂F, CHF₂, CF₃, CH₂CH₂F, CH₂CF₃, CH₂CH₂CH₂F, C₁₋₆ alkyl, C₁₋₆ alkoxy, or C₃₋₇ cycloalkyl.


[0044] In alternative embodiments, the invention provides a compound of Formula (Ie) or a pharmaceutically acceptable salt thereof:



(Ie)

where R³ may be H and R⁴ may be OH, or R³ may be H and R⁴ may be H, or R³ may be H and R⁴ may be F, or R³ may be F and R⁴ may be H, or R³ may be F and R⁴ may be F, or R³ may be OH and R⁴ may be H; R⁵ may be H and R⁶ may be OH, or R⁵ may be H and R⁶ may be H, or R⁵ may be H and R⁶ may be F, or R⁵ may be F and R⁶ may be H, or R⁵ may be F and R⁶ may be F, or R⁵ may be OH and R⁶ may be H; R⁷ may be H and R⁸ may be OH, or R⁷ may be H and R⁸ may be H, or R⁷ may be H and R⁸ may be F, or R⁷ may be F and R⁸ may be H, or R⁷ may be F and R⁸ may be F, or R⁷ may be OH and R⁸ may be H; R⁹ may be selected from the group consisting of: H, CH₃, CH₂F, CHF₂, and CH₂OH; and R¹⁰ may be selected from the group consisting of: H, C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl, C₁₋₁₀ acyl, C₈₋₂₀ arylalkylacyl, C₃₋₂₀ heteroarylalkylacyl, C₇₋₂₀ arylalkyl, C₈₋₂₀ arylalkenyl, C₈₋₂₀ arylalkynyl, C₂₋₂₀ heteroarylalkyl, C₃₋₂₀ heteroarylalkenyl, and C₃₋₂₀ heteroarylalkynyl, each excluding H optionally substituted from one to four substituents with one of more of halo, OH, OCF₃, CN, SO₂NH₂, SO₂Me, C(O)NH₂, CH₂F, CHF₂, CF₃, CH₂CH₂F, CH₂CF₃, CH₂CH₂CH₂F, C₁₋₆ alkyl, C₁₋₆ alkoxy, or C₃₋₇ cycloalkyl.


[0045] In alternative embodiments, the invention provides a compound of Formula (If) or a pharmaceutically acceptable salt thereof:



(If)

where R^3 may be H and R^4 may be OH, or R^3 may be H and R^4 may be H, or R^3 may be H and R^4 may be F, or R^3 may be F and R^4 may be H, or R^3 may be F and R^4 may be F, or R^3 may be OH and R^4 may be H; R^5 may be H and R^6 may be OH, or R^5 may be H and R^6 may be H, or R^5 may be H and R^6 may be F, or R^5 may be F and R^6 may be H, or R^5 may be F and R^6 may be F, or R^5 may be OH and R^6 may be H; R^7 may be H and R^8 may be OH, or R^7 may be H and R^8 may be H, or R^7 may be H and R^8 may be F, or R^7 may be F and R^8 may be H, or R^7 may be F and R^8 may be F, or R^7 may be OH and R^8 may be H; R^9 may be selected from the group consisting of: H, CH₃, CH₂F, CHF₂, and CH₂OH; and R^{10} may be selected from the group consisting of: H, C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl, C₁₋₁₀ acyl, C₈₋₂₀ arylalkylacyl, C₃₋₂₀ heteroarylalkylacyl, C₇₋₂₀ arylalkyl, C₈₋₂₀ arylalkenyl, C₈₋₂₀ arylalkynyl, C₂₋₂₀ heteroarylalkyl, C₃₋₂₀ heteroarylalkenyl, and C₃₋₂₀ heteroarylalkynyl, each excluding H optionally substituted from one to four substituents with one of more of halo, OH, OCF₃, CN, SO₂NH₂, SO₂Me, C(O)NH₂, CH₂F, CHF₂, CF₃, CH₂CH₂F, CH₂CF₃, CH₂CH₂CH₂F, C₁₋₆ alkyl, C₁₋₆ alkoxy, or C₃₋₇ cycloalkyl.

[0046] In alternative embodiments, the invention provides a compound of Formula (Ig) or a pharmaceutically acceptable salt thereof:




(Ig)

where R^1 may be H and R^2 may be C(O)NR¹¹₂, or R^1 may be C(O)NR¹¹₂ and R^2 may be H, or R^1 may be H and R^2 may be selected from the group consisting of: H, CH₃, CH₂F, CHF₂, and CH₂OH, where each R^{11} may be independently H or C₁₋₆ alkyl; R^5 may be H and

R^6 may be OH, or R^5 may be H and R^6 may be H, or R^5 may be H and R^6 may be F, or R^5 may be F and R^6 may be H, or R^5 may be F and R^6 may be F, or R^5 may be OH and R^6 may be H; R^7 may be H and R^8 may be OH, or R^7 may be H and R^8 may be H, or R^7 may be H and R^8 may be F, or R^7 may be F and R^8 may be H, or R^7 may be F and R^8 may be F, or R^7 may be OH and R^8 may be H; R^9 may be selected from the group consisting of: H, CH₃, CH₂F, CHF₂, and CH₂OH; and R^{10} may be selected from the group consisting of: H, C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl, C₁₋₁₀ acyl, C₈₋₂₀ arylalkylacyl, C₃₋₂₀ heteroarylalkylacyl, C₇₋₂₀ arylalkyl, C₈₋₂₀ arylalkenyl, C₈₋₂₀ arylalkynyl, C₂₋₂₀ heteroarylalkyl, C₃₋₂₀ heteroarylalkenyl, and C₃₋₂₀ heteroarylalkynyl, each excluding H optionally substituted from one to four substituents with one or more of halo, OH, OCF₃, CN, SO₂NH₂, SO₂Me, C(O)NH₂, CH₂F, CHF₂, CF₃, CH₂CH₂F, CH₂CF₃, CH₂CH₂CH₂F, C₁₋₆ alkyl, C₁₋₆ alkoxy, or C₃₋₇ cycloalkyl.

[0047] In alternative embodiments, the invention provides a compound of Formula (Ih) or a pharmaceutically acceptable salt thereof:



15


(Ih)

where R^1 may be H and R^2 may be C(O)NR¹¹₂, or R^1 may be C(O)NR¹¹₂ and R^2 may be H, or R^1 may be H and R^2 may be selected from the group consisting of: H, CH₃, CH₂F, CHF₂, and CH₂OH, where each R¹¹ may be independently H or C₁₋₆ alkyl; R^5 may be H and R^6 may be OH, or R^5 may be H and R^6 may be H, or R^5 may be H and R^6 may be F, or R^5 may be F and R^6 may be H, or R^5 may be F and R^6 may be F, or R^5 may be OH and R^6 may be H; R^7 may be H and R^8 may be OH, or R^7 may be H and R^8 may be H, or R^7 may be H and R^8 may be F, or R^7 may be F and R^8 may be H, or R^7 may be F and R^8 may be F, or R^7 may be OH and R^8 may be H; R^9 may be selected from the group consisting of: H, CH₃, CH₂F, CHF₂, and CH₂OH; and R^{10} may be selected from the group consisting of: H, C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl, C₁₋₁₀ acyl, C₈₋₂₀ arylalkylacyl, C₃₋₂₀ heteroarylalkylacyl, C₇₋₂₀ arylalkyl, C₈₋₂₀ arylalkenyl, C₈₋₂₀ arylalkynyl, C₂₋₂₀ heteroarylalkyl, C₃₋₂₀ heteroarylalkenyl, and C₃₋₂₀ heteroarylalkynyl, each excluding H optionally substituted from one to four

25

substituents with one of more of halo, OH, OCF₃, CN, SO₂NH₂, SO₂Me, C(O)NH₂, CH₂F, CHF₂, CF₃, CH₂CH₂F, CH₂CF₃, CH₂CH₂CH₂F, C₁₋₆ alkyl, C₁₋₆ alkoxy, or C₃₋₇ cycloalkyl.

[0048] In alternative embodiments, the invention provides a compound of Formula (Ii) or a pharmaceutically acceptable salt thereof:



(Ii)

5


where R¹ may be H and R² may be C(O)NR¹¹₂, or R¹ may be C(O)NR¹¹₂ and R² may be H, or R¹ may be H and R² may be selected from the group consisting of: H, CH₃, CH₂F, CHF₂, and CH₂OH, where each R¹¹ may be independently H or C₁₋₆ alkyl; R⁵ may be H and R⁶ may be OH, or R⁵ may be H and R⁶ may be H, or R⁵ may be H and R⁶ may be F, or R⁵ may be F and R⁶ may be H, or R⁵ may be F and R⁶ may be F, or R⁵ may be OH and R⁶ may be H; R⁷ may be H and R⁸ may be OH, or R⁷ may be H and R⁸ may be H, or R⁷ may be H and R⁸ may be F, or R⁷ may be F and R⁸ may be H, or R⁷ may be F and R⁸ may be F, or R⁷ may be OH and R⁸ may be H; R⁹ may be selected from the group consisting of: H, CH₃, CH₂F, CHF₂, and CH₂OH; and R¹⁰ may be selected from the group consisting of: H, C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl, C₁₋₁₀ acyl, C₈₋₂₀ arylalkylacyl, C₃₋₂₀ heteroarylalkylacyl, C₇₋₂₀ arylalkyl, C₈₋₂₀ arylalkenyl, C₈₋₂₀ arylalkynyl, C₂₋₂₀ heteroarylalkyl, C₃₋₂₀ heteroarylalkenyl, and C₃₋₂₀ heteroarylalkynyl, each excluding H optionally substituted from one to four substituents with one of more of halo, OH, OCF₃, CN, SO₂NH₂, SO₂Me, C(O)NH₂, CH₂F, CHF₂, CF₃, CH₂CH₂F, CH₂CF₃, CH₂CH₂CH₂F, C₁₋₆ alkyl, C₁₋₆ alkoxy, or C₃₋₇ cycloalkyl.

10

15

20


[0049] In alternative embodiments, the invention provides a compound of Formula (Ij) or a pharmaceutically acceptable salt thereof:



(Ij)

where R^1 may be H and R^2 may be $C(O)NR^{11}_2$, or R^1 may be $C(O)NR^{11}_2$ and R^2 may be H, or R^1 may be H and R^2 may be selected from the group consisting of: H, CH_3 , CH_2F , CHF_2 , and CH_2OH , where each R^{11} may be independently H or C_{1-6} alkyl; R^3 may be H and R^4 may be OH, or R^3 may be H and R^4 may be H, or R^3 may be H and R^4 may be F, or R^3 may be F and R^4 may be H, or R^3 may be F and R^4 may be F, or R^3 may be OH and R^4 may be H; R^7 may be H and R^8 may be OH, or R^7 may be H and R^8 may be H, or R^7 may be H and R^8 may be F, or R^7 may be F and R^8 may be H, or R^7 may be F and R^8 may be F, or R^7 may be OH and R^8 may be H; R^9 may be selected from the group consisting of: H, CH_3 , CH_2F , CHF_2 , and CH_2OH ; and R^{10} may be selected from the group consisting of: H, C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, C_{1-10} acyl, C_{8-20} arylalkylacyl, C_{3-20} heteroarylalkylacyl, C_{7-20} arylalkyl, C_{8-20} arylalkenyl, C_{8-20} arylalkynyl, C_{2-20} heteroarylalkyl, C_{3-20} heteroarylalkenyl, and C_{3-20} heteroarylalkynyl, each excluding H optionally substituted from one to four substituents with one or more of halo, OH, OCF_3 , CN, SO_2NH_2 , SO_2Me , $C(O)NH_2$, CH_2F , CHF_2 , CF_3 , CH_2CH_2F , CH_2CF_3 , $CH_2CH_2CH_2F$, C_{1-6} alkyl, C_{1-6} alkoxy, or C_{3-7} cycloalkyl.

[0050] In alternative embodiments, the invention provides a compound of Formula (Ik) or a pharmaceutically acceptable salt thereof:




(Ik)

where R^1 may be H and R^2 may be $C(O)NR^{11}_2$, or R^1 may be $C(O)NR^{11}_2$ and R^2 may be H, or R^1 may be H and R^2 may be selected from the group consisting of: H, CH_3 , CH_2F , CHF_2 , and CH_2OH , where each R^{11} may be independently H or C_{1-6} alkyl; R^3 may be H and R^4 may be OH, or R^3 may be H and R^4 may be H, or R^3 may be H and R^4 may be F, or R^3 may be F and R^4 may be H, or R^3 may be F and R^4 may be F, or R^3 may be OH and R^4 may be H; R^7 may be H and R^8 may be OH, or R^7 may be H and R^8 may be H, or R^7 may be H and R^8 may be F, or R^7 may be F and R^8 may be H, or R^7 may be F and R^8 may be F, or R^7 may be OH and R^8 may be H; R^9 may be selected from the group consisting of: H, CH_3 , CH_2F , CHF_2 , and CH_2OH ; and R^{10} may be selected from the group consisting of: H, C_{1-10} alkyl, C_{2-10}

10 alkenyl, C₂₋₁₀ alkynyl, C₁₋₁₀ acyl, C₈₋₂₀ arylalkylacyl, C₃₋₂₀ heteroarylalkylacyl, C₇₋₂₀
 arylalkyl, C₈₋₂₀ arylalkenyl, C₈₋₂₀ arylalkynyl, C₂₋₂₀ heteroarylalkyl, C₃₋₂₀ heteroarylalkenyl,
 and C₃₋₂₀ heteroarylalkynyl, each excluding H optionally substituted from one to four
 5 substituents with one of more of halo, OH, OCF₃, CN, SO₂NH₂, SO₂Me, C(O)NH₂, CH₂F,
 CHF₂, CF₃, CH₂CH₂F, CH₂CF₃, CH₂CH₂CH₂F, C₁₋₆ alkyl, C₁₋₆ alkoxy, or C₃₋₇ cycloalkyl.


[0051] In alternative embodiments, the invention provides a compound of Formula (II) or a
 pharmaceutically acceptable salt thereof:



(II)

10 where R¹ may be H and R² may be C(O)NR¹¹₂, or R¹ may be C(O)NR¹¹₂ and R² may
 be H, or R¹ may be H and R² may be selected from the group consisting of: H, CH₃, CH₂F,
 CHF₂, and CH₂OH, where each R¹¹ may be independently H or C₁₋₆ alkyl; R³ may be H and
 R⁴ may be OH, or R³ may be H and R⁴ may be H, or R³ may be H and R⁴ may be F, or R³
 may be F and R⁴ may be H, or R³ may be F and R⁴ may be F, or R³ may be OH and R⁴ may
 15 be H; R⁷ may be H and R⁸ may be OH, or R⁷ may be H and R⁸ may be H, or R⁷ may be H and
 R⁸ may be F, or R⁷ may be F and R⁸ may be H, or R⁷ may be F and R⁸ may be F, or R⁷ may
 be OH and R⁸ may be H; R⁹ may be selected from the group consisting of: H, CH₃, CH₂F,
 CHF₂, and CH₂OH; and R¹⁰ may be selected from the group consisting of: H, C₁₋₁₀ alkyl, C<sub>2-
 10</sub> alkenyl, C₂₋₁₀ alkynyl, C₁₋₁₀ acyl, C₈₋₂₀ arylalkylacyl, C₃₋₂₀ heteroarylalkylacyl, C₇₋₂₀
 20 arylalkyl, C₈₋₂₀ arylalkenyl, C₈₋₂₀ arylalkynyl, C₂₋₂₀ heteroarylalkyl, C₃₋₂₀ heteroarylalkenyl,
 and C₃₋₂₀ heteroarylalkynyl, each excluding H optionally substituted from one to four
 substituents with one of more of halo, OH, OCF₃, CN, SO₂NH₂, SO₂Me, C(O)NH₂, CH₂F,
 CHF₂, CF₃, CH₂CH₂F, CH₂CF₃, CH₂CH₂CH₂F, C₁₋₆ alkyl, C₁₋₆ alkoxy, or C₃₋₇ cycloalkyl.


[0052] In alternative embodiments, the invention provides a compound of Formula (Im) or a
 25 pharmaceutically acceptable salt thereof:



(Im)

where R¹ may be H and R² may be C(O)NR¹¹₂, or R¹ may be C(O)NR¹¹₂ and R² may be H, or R¹ may be H and R² may be selected from the group consisting of: H, CH₃, CH₂F, CHF₂, and CH₂OH, where each R¹¹ may be independently H or C₁₋₆ alkyl; R³ may be H and R⁴ may be OH, or R³ may be H and R⁴ may be H, or R³ may be H and R⁴ may be F, or R³ may be F and R⁴ may be H, or R³ may be F and R⁴ may be F, or R³ may be OH and R⁴ may be H; R⁵ may be H and R⁶ may be OH, or R⁵ may be H and R⁶ may be H, or R⁵ may be H and R⁶ may be F, or R⁵ may be F and R⁶ may be H, or R⁵ may be F and R⁶ may be F, or R⁵ may be OH and R⁶ may be H; R⁹ may be selected from the group consisting of: H, CH₃, CH₂F, CHF₂, and CH₂OH; and R¹⁰ may be selected from the group consisting of: H, C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl, C₁₋₁₀ acyl, C₈₋₂₀ arylalkylacyl, C₃₋₂₀ heteroarylalkylacyl, C₇₋₂₀ arylalkyl, C₈₋₂₀ arylalkenyl, C₈₋₂₀ arylalkynyl, C₂₋₂₀ heteroarylalkyl, C₃₋₂₀ heteroarylalkenyl, and C₃₋₂₀ heteroarylalkynyl, each excluding H optionally substituted from one to four substituents with one or more of halo, OH, OCF₃, CN, SO₂NH₂, SO₂Me, C(O)NH₂, CH₂F, CHF₂, CF₃, CH₂CH₂F, CH₂CF₃, CH₂CH₂CH₂F, C₁₋₆ alkyl, C₁₋₆ alkoxy, or C₃₋₇ cycloalkyl.

[0053] In alternative embodiments, the invention provides a compound of Formula (In) or a pharmaceutically acceptable salt thereof:




(In)

where R¹ may be H and R² may be C(O)NR¹¹₂, or R¹ may be C(O)NR¹¹₂ and R² may be H, or R¹ may be H and R² may be selected from the group consisting of: H, CH₃, CH₂F, CHF₂, and CH₂OH, where each R¹¹ may be independently H or C₁₋₆ alkyl; R³ may be H and

R^4 may be OH, or R^3 may be H and R^4 may be H, or R^3 may be H and R^4 may be F, or R^3 may be F and R^4 may be H, or R^3 may be F and R^4 may be F, or R^3 may be OH and R^4 may be H; R^5 may be H and R^6 may be OH, or R^5 may be H and R^6 may be H, or R^5 may be H and R^6 may be F, or R^5 may be F and R^6 may be H, or R^5 may be F and R^6 may be F, or R^5 may be OH and R^6 may be H; R^9 may be selected from the group consisting of: H, CH₃, CH₂F, CHF₂, and CH₂OH; and R^{10} may be selected from the group consisting of: H, C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl, C₁₋₁₀ acyl, C₈₋₂₀ arylalkylacyl, C₃₋₂₀ heteroarylalkylacyl, C₇₋₂₀ arylalkyl, C₈₋₂₀ arylalkenyl, C₈₋₂₀ arylalkynyl, C₂₋₂₀ heteroarylalkyl, C₃₋₂₀ heteroarylalkenyl, and C₃₋₂₀ heteroarylalkynyl, each excluding H optionally substituted from one to four substituents with one or more of halo, OH, OCF₃, CN, SO₂NH₂, SO₂Me, C(O)NH₂, CH₂F, CHF₂, CF₃, CH₂CH₂F, CH₂CF₃, CH₂CH₂CH₂F, C₁₋₆ alkyl, C₁₋₆ alkoxy, or C₃₋₇ cycloalkyl.

[0054] In alternative embodiments, the invention provides a compound of Formula (Io) or a pharmaceutically acceptable salt thereof:



15


(Io)

where R^1 may be H and R^2 may be C(O)NR¹¹₂, or R^1 may be C(O)NR¹¹₂ and R^2 may be H, or R^1 may be H and R^2 may be selected from the group consisting of: H, CH₃, CH₂F, CHF₂, and CH₂OH, where each R^{11} may be independently H or C₁₋₆ alkyl; R^3 may be H and R^4 may be OH, or R^3 may be H and R^4 may be H, or R^3 may be H and R^4 may be F, or R^3 may be F and R^4 may be H, or R^3 may be F and R^4 may be F, or R^3 may be OH and R^4 may be H; R^5 may be H and R^6 may be OH, or R^5 may be H and R^6 may be H, or R^5 may be H and R^6 may be F, or R^5 may be F and R^6 may be H, or R^5 may be F and R^6 may be F, or R^5 may be OH and R^6 may be H; R^9 may be selected from the group consisting of: H, CH₃, CH₂F, CHF₂, and CH₂OH; and R^{10} may be selected from the group consisting of: H, C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl, C₁₋₁₀ acyl, C₈₋₂₀ arylalkylacyl, C₃₋₂₀ heteroarylalkylacyl, C₇₋₂₀ arylalkyl, C₈₋₂₀ arylalkenyl, C₈₋₂₀ arylalkynyl, C₂₋₂₀ heteroarylalkyl, C₃₋₂₀ heteroarylalkenyl, and C₃₋₂₀ heteroarylalkynyl, each excluding H optionally substituted from one to four

25

substituents with one of more of halo, OH, OCF₃, CN, SO₂NH₂, SO₂Me, C(O)NH₂, CH₂F, CHF₂, CF₃, CH₂CH₂F, CH₂CF₃, CH₂CH₂CH₂F, C₁₋₆ alkyl, C₁₋₆ alkoxy, or C₃₋₇ cycloalkyl.

[0055] In alternative embodiments, the invention provides a compound of Formula (Ip) or a pharmaceutically acceptable salt thereof:



(Ip)

5


where R¹ may be H and R² may be C(O)NR¹¹₂, or R¹ may be C(O)NR¹¹₂ and R² may be H, or R¹ may be H and R² may be selected from the group consisting of: H, CH₃, CH₂F, CHF₂, and CH₂OH, where each R¹¹ may be independently H or C₁₋₆ alkyl; R³ may be H and R⁴ may be OH, or R³ may be H and R⁴ may be H, or R³ may be H and R⁴ may be F, or R³ may be F and R⁴ may be H, or R³ may be F and R⁴ may be F, or R³ may be OH and R⁴ may be H; R⁵ may be H and R⁶ may be OH, or R⁵ may be H and R⁶ may be H, or R⁵ may be H and R⁶ may be F, or R⁵ may be F and R⁶ may be H, or R⁵ may be F and R⁶ may be F, or R⁵ may be OH and R⁶ may be H; R⁷ may be H and R⁸ may be OH, or R⁷ may be H and R⁸ may be H, or R⁷ may be H and R⁸ may be F, or R⁷ may be F and R⁸ may be H, or R⁷ may be F and R⁸ may be F, or R⁷ may be OH and R⁸ may be H; and R¹⁰ may be selected from the group consisting of: H, C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl, C₁₋₁₀ acyl, C₈₋₂₀ arylalkylacyl, C₃₋₂₀ heteroarylalkylacyl, C₇₋₂₀ arylalkyl, C₈₋₂₀ arylalkenyl, C₈₋₂₀ arylalkynyl, C₂₋₂₀ heteroarylalkyl, C₃₋₂₀ heteroarylalkenyl, and C₃₋₂₀ heteroarylalkynyl, each excluding H optionally substituted from one to four substituents with one of more of halo, OH, OCF₃, CN, SO₂NH₂, SO₂Me, C(O)NH₂, CH₂F, CHF₂, CF₃, CH₂CH₂F, CH₂CF₃, CH₂CH₂CH₂F, C₁₋₆ alkyl, C₁₋₆ alkoxy, or C₃₋₇ cycloalkyl.

10

15

20


[0056] In alternative embodiments, the invention provides a compound of Formula (Iq) or a pharmaceutically acceptable salt thereof:



(Iq)

where R¹ may be H and R² may be C(O)NR¹¹₂, or R¹ may be C(O)NR¹¹₂ and R² may be H, or R¹ may be H and R² may be selected from the group consisting of: H, CH₃, CH₂F, CHF₂, and CH₂OH, where each R¹¹ may be independently H or C₁₋₆ alkyl; R³ may be H and R⁴ may be OH, or R³ may be H and R⁴ may be H, or R³ may be H and R⁴ may be F, or R³ may be F and R⁴ may be H, or R³ may be F and R⁴ may be F, or R³ may be OH and R⁴ may be H; R⁵ may be H and R⁶ may be OH, or R⁵ may be H and R⁶ may be H, or R⁵ may be H and R⁶ may be F, or R⁵ may be F and R⁶ may be H, or R⁵ may be F and R⁶ may be F, or R⁵ may be OH and R⁶ may be H; R⁷ may be H and R⁸ may be OH, or R⁷ may be H and R⁸ may be H, or R⁷ may be H and R⁸ may be F, or R⁷ may be F and R⁸ may be H, or R⁷ may be F and R⁸ may be F, or R⁷ may be OH and R⁸ may be H; and R¹⁰ may be selected from the group consisting of: H, C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl, C₁₋₁₀ acyl, C₈₋₂₀ arylalkylacyl, C₃₋₂₀ heteroarylalkylacyl, C₇₋₂₀ arylalkyl, C₈₋₂₀ arylalkenyl, C₈₋₂₀ arylalkynyl, C₂₋₂₀ heteroarylalkyl, C₃₋₂₀ heteroarylalkenyl, and C₃₋₂₀ heteroarylalkynyl, each excluding H optionally substituted from one to four substituents with one of more of halo, OH, OCF₃, CN, SO₂NH₂, SO₂Me, C(O)NH₂, CH₂F, CHF₂, CF₃, CH₂CH₂F, CH₂CF₃, CH₂CH₂CH₂F, C₁₋₆ alkyl, C₁₋₆ alkoxy, or C₃₋₇ cycloalkyl.


[0057] In alternative embodiments, the invention provides a compound of Formula (Ir) or a pharmaceutically acceptable salt thereof.



(Ir)

where R^1 may be H and R^2 may be $C(O)NR^{11}_2$, or R^1 may be $C(O)NR^{11}_2$ and R^2 may be H, or R^1 may be H and R^2 may be selected from the group consisting of: H, CH_3 , CH_2F , CHF_2 , and CH_2OH , where each R^{11} may be independently H or C_{1-6} alkyl; R^3 may be H and R^4 may be OH, or R^3 may be H and R^4 may be H, or R^3 may be H and R^4 may be F, or R^3 may be F and R^4 may be H, or R^3 may be F and R^4 may be F, or R^3 may be OH and R^4 may be H; R^5 may be H and R^6 may be OH, or R^5 may be H and R^6 may be H, or R^5 may be H and R^6 may be F, or R^5 may be F and R^6 may be H, or R^5 may be F and R^6 may be F, or R^5 may be OH and R^6 may be H; R^7 may be H and R^8 may be OH, or R^7 may be H and R^8 may be H, or R^7 may be H and R^8 may be F, or R^7 may be F and R^8 may be H, or R^7 may be F and R^8 may be F, or R^7 may be OH and R^8 may be H; and R^{10} may be selected from the group consisting of: H, C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, C_{1-10} acyl, C_{8-20} arylalkylacyl, C_{3-20} heteroarylalkylacyl, C_{7-20} arylalkyl, C_{8-20} arylalkenyl, C_{8-20} arylalkynyl, C_{2-20} heteroarylalkyl, C_{3-20} heteroarylalkenyl, and C_{3-20} heteroarylalkynyl, each excluding H optionally substituted from one to four substituents with one of more of halo, OH, OCF_3 , CN, SO_2NH_2 , SO_2Me , $C(O)NH_2$, CH_2F , CHF_2 , CF_3 , CH_2CH_2F , CH_2CF_3 , $CH_2CH_2CH_2F$, C_{1-6} alkyl, C_{1-6} alkoxy, or C_{3-7} cycloalkyl.

[0058] In alternative embodiments, the invention provides a compound of Formula (Is) or a pharmaceutically acceptable salt thereof:




(Is)

where R^1 may be H and R^2 may be $C(O)NR^{11}_2$, or R^1 may be $C(O)NR^{11}_2$ and R^2 may be H, or R^1 may be H and R^2 may be selected from the group consisting of: H, CH_3 , CH_2F , CHF_2 , and CH_2OH , where each R^{11} may be independently H or C_{1-6} alkyl; R^3 may be H and R^4 may be OH, or R^3 may be H and R^4 may be H, or R^3 may be H and R^4 may be F, or R^3 may be F and R^4 may be H, or R^3 may be F and R^4 may be F, or R^3 may be OH and R^4 may be H; R^5 may be H and R^6 may be OH, or R^5 may be H and R^6 may be H, or R^5 may be H and R^6 may be F, or R^5 may be F and R^6 may be H, or R^5 may be F and R^6 may be F, or R^5 may be OH and R^6 may be H; R^7 may be H and R^8 may be OH, or R^7 may be H and R^8 may be H,

or R⁷ may be H and R⁸ may be F, or R⁷ may be F and R⁸ may be H, or R⁷ may be F and R⁸ may be F, or R⁷ may be OH and R⁸ may be H; and R¹⁰ may be selected from the group consisting of: H, C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl, C₁₋₁₀ acyl, C₈₋₂₀ arylalkylacyl, C₃₋₂₀ heteroarylalkylacyl, C₇₋₂₀ arylalkyl, C₈₋₂₀ arylalkenyl, C₈₋₂₀ arylalkynyl, C₂₋₂₀ heteroarylalkyl, C₃₋₂₀ heteroarylalkenyl, and C₃₋₂₀ heteroarylalkynyl, each excluding H optionally substituted from one to four substituents with one of more of halo, OH, OCF₃, CN, SO₂NH₂, SO₂Me, C(O)NH₂, CH₂F, CHF₂, CF₃, CH₂CH₂F, CH₂CF₃, CH₂CH₂CH₂F, C₁₋₆ alkyl, C₁₋₆ alkoxy, or C₃₋₇ cycloalkyl.

[0059] In alternative embodiments, the invention provides a compound of Formula (II) or a pharmaceutically acceptable salt thereof:



(II)

where R¹ may be H and R² may be C(O)NR¹¹₂, or R¹ may be C(O)NR¹¹₂ and R² may be H, or R¹ may be H and R² may be selected from the group consisting of: H, CH₃, CH₂F, CHF₂, and CH₂OH, where each R¹¹ may be independently H or C₁₋₆ alkyl; R³ may be H and R⁴ may be OH, or R³ may be H and R⁴ may be H, or R³ may be H and R⁴ may be F, or R³ may be F and R⁴ may be H, or R³ may be F and R⁴ may be F, or R³ may be OH and R⁴ may be H; R⁵ may be H and R⁶ may be OH, or R⁵ may be H and R⁶ may be H, or R⁵ may be H and R⁶ may be F, or R⁵ may be F and R⁶ may be H, or R⁵ may be F and R⁶ may be F, or R⁵ may be OH and R⁶ may be H; R⁷ may be H and R⁸ may be OH, or R⁷ may be H and R⁸ may be H, or R⁷ may be H and R⁸ may be F, or R⁷ may be F and R⁸ may be H, or R⁷ may be F and R⁸ may be F, or R⁷ may be OH and R⁸ may be H; R⁹ may be selected from the group consisting of: H, CH₃, CH₂F, CHF₂, and CH₂OH; and R¹⁰ may be selected from the group consisting of: C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl, C₁₋₁₀ acyl, C₈₋₂₀ arylalkylacyl, C₃₋₂₀ heteroarylalkylacyl, C₇₋₂₀ arylalkyl, C₈₋₂₀ arylalkenyl, C₈₋₂₀ arylalkynyl, C₂₋₂₀ heteroarylalkyl, C₃₋₂₀ heteroarylalkenyl, and C₃₋₂₀ heteroarylalkynyl, each optionally substituted from one to four substituents with one of more of halo, OH, OCF₃, CN, SO₂NH₂, SO₂Me, C(O)NH₂, CH₂F, CHF₂, CF₃,

CH₂CH₂F, CH₂CF₃, CH₂CH₂CH₂F, C₁₋₆ alkyl, C₁₋₆ alkoxy, or C₃₋₇ cycloalkyl; with the proviso that R¹⁰ excludes benzyl and (CH₂)₃phenyl.

[0060] In alternative embodiments, the compound may be a prodrug; the compound may selectively inhibit an O-glycoprotein 2-acetamido-2-deoxy-β-D-glucopyranosidase (O-GlcNAcase); the compound may selectively bind an O-GlcNAcase (e.g., a mammalian O-GlcNAcase); the compound may selectively inhibit the cleavage of a 2-acetamido-2-deoxy-β-D-glucopyranoside (O-GlcNAc); the compound may not substantially inhibit a mammalian β-hexosaminidase.

[0061] In alternative embodiments, a compound according to Formula (I), Formula (Ia), Formula (Ib), Formula (Ic), Formula (Id), Formula (Ie), Formula (If), Formula (Ig), Formula (Ih), Formula (Ii), Formula (Ij), Formula (Ik), Formula (Il), Formula (Im), Formula (In), Formula (Io), Formula (Ip), Formula (Iq), Formula (Ir), or Formula (Is) may have enhanced permeability.


[0062] In alternative embodiments, a compound according to Formula (Ia) may have enhanced selectivity.

[0063] In alternative embodiments, a compound according to Formula (Ie), Formula (If), Formula (Ig), Formula (Ih), Formula (Ij), Formula (Ik), Formula (Im), Formula (In), Formula (Ip), or Formula (Iq), may have enhanced permeability.

[0064] In alternative embodiments, a compound according to Formula (Id), Formula (Il), Formula (Io), Formula (Ir), or Formula (Is) may have enhanced permeability.

[0065] In alternative aspects, the invention provides a pharmaceutical composition including a compound according to the invention, in combination with a pharmaceutically acceptable carrier.

[0066] In alternative aspects, the invention provides methods of selectively inhibiting an O-GlcNAcase, or of inhibiting an O-GlcNAcase in a subject in need thereof, or of increasing the level of O-GlcNAc, or of treating a neurodegenerative disease, a tauopathy, cancer or stress, in a subject in need thereof, by administering to the subject an effective amount of a compound of Formula (I) or a pharmaceutically acceptable salt thereof:




(I)

where R^1 may be H and R^2 may be $C(O)NR^{11}_2$, or R^1 may be $C(O)NR^{11}_2$ and R^2 may be H, or R^1 may be H and R^2 may be selected from the group consisting of: H, CH_3 , CH_2F , CHF_2 , and CH_2OH , where each R^{11} may be independently H or C_{1-6} alkyl; R^3 may be H and R^4 may be OH, or R^3 may be H and R^4 may be H, or R^3 may be H and R^4 may be F, or R^3 may be F and R^4 may be H, or R^3 may be F and R^4 may be F, or R^3 may be OH and R^4 may be H; R^5 may be H and R^6 may be OH, or R^5 may be H and R^6 may be H, or R^5 may be H and R^6 may be F, or R^5 may be F and R^6 may be H, or R^5 may be F and R^6 may be F, or R^5 may be OH and R^6 may be H; R^7 may be H and R^8 may be OH, or R^7 may be H and R^8 may be H, or R^7 may be H and R^8 may be F, or R^7 may be F and R^8 may be H, or R^7 may be F and R^8 may be F, or R^7 may be OH and R^8 may be H; R^9 may be selected from the group consisting of: H, CH_3 , CH_2F , CHF_2 , and CH_2OH ; and R^{10} may be selected from the group consisting of: H, C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, C_{1-10} acyl, C_{8-20} arylalkylacyl, C_{3-20} heteroarylalkylacyl, C_{7-20} arylalkyl, C_{8-20} arylalkenyl, C_{8-20} arylalkynyl, C_{2-20} heteroarylalkyl, C_{3-20} heteroarylalkenyl, and C_{3-20} heteroarylalkynyl, each excluding H optionally substituted from one to four substituents with one of more of halo, OH, OCF_3 , CN, SO_2NH_2 , SO_2Me , $C(O)NH_2$, CH_2F , CHF_2 , CF_3 , CH_2CH_2F , CH_2CF_3 , $CH_2CH_2CH_2F$, C_{1-6} alkyl, C_{1-6} alkoxy, or C_{3-7} cycloalkyl. The condition may be Alzheimer's disease, Amyotrophic lateral sclerosis (ALS), Amyotrophic lateral sclerosis with cognitive impairment (ALSci), Argyrophilic grain dementia, Bluit disease, Corticobasal degeneration (CBD), Dementia pugilistica, Diffuse neurofibrillary tangles with calcification, Down's syndrome, Familial British dementia, Familial Danish dementia, Frontotemporal dementia with parkinsonism linked to chromosome 17 (FTDP-17), Gerstmann-Straussler-Scheinker disease, Guadeloupean parkinsonism, Hallevorden-Spatz disease (neurodegeneration with brain iron accumulation type 1), Multiple system atrophy, Myotonic dystrophy, Niemann-Pick disease (type C), Pallido-ponto-nigral degeneration, Parkinsonism-dementia complex of Guam, Pick's disease (PiD), Post-encephalitic parkinsonism (PEP), Prion diseases (including Creutzfeldt-Jakob Disease (CJD), Variant Creutzfeldt-Jakob Disease (vCJD), Fatal Familial Insomnia, and

Kuru), Progressive superecortical gliosis, Progressive supranuclear palsy (PSP), Richardson's syndrome, Subacute sclerosing panencephalitis, Tangle-only dementia, Huntington's disease, Parkinson's disease, Schizophrenia, Mild Cognitive Impairment (MCI), Neuropathy (including peripheral neuropathy, autonomic neuropathy, neuritis, and diabetic neuropathy),
 5 or Glaucoma. The stress may be a cardiac disorder, e.g., ischemia; hemorrhage; hypovolemic shock; myocardial infarction; an interventional cardiology procedure; cardiac bypass surgery; fibrinolytic therapy; angioplasty; or stent placement.

[0067] In alternative aspects, the invention provides a method of treating an O-GlcNAcase-mediated condition that excludes a neurodegenerative disease, a tauopathy, cancer or stress,
 10 in a subject in need thereof, by administering to the subject an effective amount of a compound of Formula (I) or a pharmaceutically acceptable salt thereof:




(I)

where R^1 may be H and R^2 may be $C(O)NR^{11}_2$, or R^1 may be $C(O)NR^{11}_2$ and R^2 may
 15 be H, or R^1 may be H and R^2 may be selected from the group consisting of: H, CH_3 , CH_2F , CHF_2 , and CH_2OH , where each R^{11} may be independently H or C_{1-6} alkyl; R^3 may be H and R^4 may be OH, or R^3 may be H and R^4 may be H, or R^3 may be H and R^4 may be F, or R^3 may be F and R^4 may be H, or R^3 may be F and R^4 may be F, or R^3 may be OH and R^4 may be H; R^5 may be H and R^6 may be OH, or R^5 may be H and R^6 may be H, or R^5 may be H and
 20 R^6 may be F, or R^5 may be F and R^6 may be H, or R^5 may be F and R^6 may be F, or R^5 may be OH and R^6 may be H; R^7 may be H and R^8 may be OH, or R^7 may be H and R^8 may be H, or R^7 may be H and R^8 may be F, or R^7 may be F and R^8 may be H, or R^7 may be F and R^8 may be F, or R^7 may be OH and R^8 may be H; R^9 may be selected from the group consisting of: H, CH_3 , CH_2F , CHF_2 , and CH_2OH ; and R^{10} may be selected from the group consisting of:
 25 H, C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, C_{1-10} acyl, C_{8-20} arylalkylacyl, C_{3-20} heteroarylalkylacyl, C_{7-20} arylalkyl, C_{8-20} arylalkenyl, C_{8-20} arylalkynyl, C_{2-20} heteroarylalkyl, C_{3-20} heteroarylalkenyl, and C_{3-20} heteroarylalkynyl, each excluding H optionally substituted from one to four substituents with one of more of halo, OH, OCF_3 , CN, SO_2NH_2 , SO_2Me ,

C(O)NH₂, CH₂F, CHF₂, CF₃, CH₂CH₂F, CH₂CF₃, CH₂CH₂CH₂F, C₁₋₆ alkyl, C₁₋₆ alkoxy, or C₃₋₇ cycloalkyl. In some embodiments, the condition may be inflammatory or allergic diseases such as asthma, allergic rhinitis, hypersensitivity lung diseases, hypersensitivity pneumonitis, eosinophilic pneumonias, delayed-type hypersensitivity, atherosclerosis, interstitial lung disease (ILD) (*e.g.*, idiopathic pulmonary fibrosis, or ILD associated with rheumatoid arthritis, systemic lupus erythematosus, ankylosing spondylitis, systemic sclerosis, Sjogren's syndrome, polymyositis or dermatomyositis); systemic anaphylaxis or hypersensitivity responses, drug allergies, insect sting allergies; autoimmune diseases, such as rheumatoid arthritis, psoriatic arthritis, multiple sclerosis, Guillain-Barré syndrome, systemic lupus erythematosus, myasthenia gravis, glomerulonephritis, autoimmune thyroiditis, graft rejection, including allograft rejection or graft-versus-host disease; inflammatory bowel diseases, such as Crohn's disease and ulcerative colitis; spondyloarthropathies; scleroderma; psoriasis (including T-cell mediated psoriasis) and inflammatory dermatoses such as dermatitis, eczema, atopic dermatitis, allergic contact dermatitis, urticaria; vasculitis (*e.g.*, necrotizing, cutaneous, and hypersensitivity vasculitis); eosinophilic myotitis, and eosinophilic fasciitis; graft rejection, in particular but not limited to solid organ transplants, such as heart, lung, liver, kidney, and pancreas transplants (*e.g.* kidney and lung allografts); epilepsy; pain; fibromyalgia; stroke, *e.g.*, neuroprotection following a stroke.

[0068] In alternative embodiments, the administering may increase the level of O-GlcNAc in the subject. The subject may be a human.

[0069] In alternative aspects, the invention provides use of a compound of an effective amount of a compound of Formula (I) or a pharmaceutically acceptable salt thereof:




(I)

where R¹ may be H and R² may be C(O)NR¹¹₂, or R¹ may be C(O)NR¹¹₂ and R² may be H, or R¹ may be H and R² may be selected from the group consisting of: H, CH₃, CH₂F, CHF₂, and CH₂OH, where each R¹¹ may be independently H or C₁₋₆ alkyl; R³ may be H and R⁴ may be OH, or R³ may be H and R⁴ may be H, or R³ may be H and R⁴ may be F, or R³

may be F and R⁴ may be H, or R³ may be F and R⁴ may be F, or R³ may be OH and R⁴ may be H; R⁵ may be H and R⁶ may be OH, or R⁵ may be H and R⁶ may be H, or R⁵ may be H and R⁶ may be F, or R⁵ may be F and R⁶ may be H, or R⁵ may be F and R⁶ may be F, or R⁵ may be OH and R⁶ may be H; R⁷ may be H and R⁸ may be OH, or R⁷ may be H and R⁸ may be H, or R⁷ may be H and R⁸ may be F, or R⁷ may be F and R⁸ may be H, or R⁷ may be F and R⁸ may be F, or R⁷ may be OH and R⁸ may be H; R⁹ may be selected from the group consisting of: H, CH₃, CH₂F, CHF₂, and CH₂OH; and R¹⁰ may be selected from the group consisting of: H, C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl, C₁₋₁₀ acyl, C₈₋₂₀ arylalkylacyl, C₃₋₂₀ heteroarylalkylacyl, C₇₋₂₀ arylalkyl, C₈₋₂₀ arylalkenyl, C₈₋₂₀ arylalkynyl, C₂₋₂₀ heteroarylalkyl, C₃₋₂₀ heteroarylalkenyl, and C₃₋₂₀ heteroarylalkynyl, each excluding H optionally substituted from one to four substituents with one or more of halo, OH, OCF₃, CN, SO₂NH₂, SO₂Me, C(O)NH₂, CH₂F, CHF₂, CF₃, CH₂CH₂F, CH₂CF₃, CH₂CH₂CH₂F, C₁₋₆ alkyl, C₁₋₆ alkoxy, or C₃₋₇ cycloalkyl, in the preparation of a medicament. The medicament may be for selectively inhibiting an O-GlcNAcase, for increasing the level of O-GlcNAc, for treating a condition modulated by an O-GlcNAcase, for treating a neurodegenerative disease, a tauopathy, a cancer, or stress.

[0070] In alternative aspects, the invention provides a method for screening for a selective inhibitor of an O-GlcNAcase, by a) contacting a first sample with a test compound; b) contacting a second sample with a compound of Formula (I)



(I)

where R¹ may be H and R² may be C(O)NR¹¹₂, or R¹ may be C(O)NR¹¹₂ and R² may be H, or R¹ may be H and R² may be selected from the group consisting of: H, CH₃, CH₂F, CHF₂, and CH₂OH, where each R¹¹ may be independently H or C₁₋₆ alkyl; R³ may be H and R⁴ may be OH, or R³ may be H and R⁴ may be H, or R³ may be H and R⁴ may be F, or R³ may be F and R⁴ may be H, or R³ may be F and R⁴ may be F, or R³ may be OH and R⁴ may be H; R⁵ may be H and R⁶ may be OH, or R⁵ may be H and R⁶ may be H, or R⁵ may be H and R⁶ may be F, or R⁵ may be F and R⁶ may be H, or R⁵ may be F and R⁶ may be F, or R⁵ may

be OH and R⁶ may be H; R⁷ may be H and R⁸ may be OH, or R⁷ may be H and R⁸ may be H, or R⁷ may be H and R⁸ may be F, or R⁷ may be F and R⁸ may be H, or R⁷ may be F and R⁸ may be F, or R⁷ may be OH and R⁸ may be H; R⁹ may be selected from the group consisting of: H, CH₃, CH₂F, CHF₂, and CH₂OH; and R¹⁰ may be selected from the group consisting of:
5 H, C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl, C₁₋₁₀ acyl, C₈₋₂₀ arylalkylacyl, C₃₋₂₀ heteroarylalkylacyl, C₇₋₂₀ arylalkyl, C₈₋₂₀ arylalkenyl, C₈₋₂₀ arylalkynyl, C₂₋₂₀ heteroarylalkyl, C₃₋₂₀ heteroarylalkenyl, and C₃₋₂₀ heteroarylalkynyl, each excluding H optionally substituted from one to four substituents with one of more of halo, OH, OCF₃, CN, SO₂NH₂, SO₂Me, C(O)NH₂, CH₂F, CHF₂, CF₃, CH₂CH₂F, CH₂CF₃, CH₂CH₂CH₂F, C₁₋₆ alkyl, C₁₋₆ alkoxy, or
10 C₃₋₇ cycloalkyl; c) determining the level of inhibition of the O-GlcNAcase in the first and second samples, where the test compound is a selective inhibitor of a O-GlcNAcase if the test compound exhibits the same or greater inhibition of the O-GlcNAcase when compared to the compound of Formula (I).

[0071] This summary of the invention does not necessarily describe all features of the
15 invention.

DETAILED DESCRIPTION

[0072] The invention provides, in part, novel compounds that are capable of inhibiting an O-glycoprotein 2-acetamido-2-deoxy-β-D-glucopyranosidase (O-GlcNAcase). In some
20 embodiments, the O-GlcNAcase may be a mammalian O-GlcNAcase, such as a rat, mouse or human O-GlcNAcase.

[0073] In some embodiments, one or more of the compounds according to the invention may exhibit enhanced permeability. Permeability can be assessed using a variety of standard experimental techniques, including without limitation in situ perfusion, ex vivo tissue
25 diffusion, in vitro cell monolayers (e.g. Caco-2 cells, MDCK cells, LLC-PK1 cells), and artificial cell membranes (e.g. PAMPA assay); suitable techniques for measuring effective permeability (P_{eff}) or apparent permeability (P_{app}) are reviewed for example by Volpe in *The AAPS Journal*, **2010**, 12(4), 670-678. In some embodiments, one or more of the compounds according to the invention may show enhanced permeability when tested in one or more of
30 these assays for determining P_{eff} or P_{app}. In some embodiments, a compound that exhibits enhanced permeability may exhibit greater oral absorption. In some embodiments, a compound that exhibits enhanced permeability may exhibit greater brain penetration when

administered in vivo. In some embodiments, a compound that exhibits enhanced permeability may achieve higher brain concentrations when administered in vivo. In some embodiments, a compound that exhibits enhanced permeability may exhibit a higher brain/plasma concentration ratio when administered in vivo. In some embodiments,

5 “enhanced permeability” means an increase in measured P_{eff} or P_{app} by any value between 10% and 100%, or of any integer value between 10% and 100%, for example, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100%, or over 100%, or an increase by 1-fold, 2-fold, or 3-fold, or more, as compared to a suitable reference compound disclosed in for example WO 2006/092049 or WO 2008/025170. A suitable reference compound may be, for
10 example, (3aR,5R,6S,7R,7aR)-5-(hydroxymethyl)-2-propyl-5,6,7,7a-tetrahydro-3aH-pyrano[3,2-d]thiazole-6,7-diol, or (3aR,5R,6S,7R,7aR)-2-(ethylamino)-5-(hydroxymethyl)-5,6,7,7a-tetrahydro-3aH-pyrano[3,2-d]thiazole-6,7-diol, or (3aR,5R,6S,7R,7aR)-2-(dimethylamino)-5-(hydroxymethyl)-5,6,7,7a-tetrahydro-3aH-pyrano[3,2-d]thiazole-6,7-diol. In some embodiments, “enhanced permeability” means a measurable P_{app} value (i.e. a value
15 greater than zero) in the assay described below for determination of P_{app} in LLC-PK1 cells. In some embodiments, “enhanced permeability” means a P_{app} value greater than 2×10^{-6} cm/s in the assay described below for determination of P_{app} in LLC-PK1 cells. In alternative embodiments, “enhanced permeability” means a P_{app} value in the range 2×10^{-6} cm/s to 35×10^{-6} cm/s in the assay described below for determination of P_{app} in LLC-PK1 cells.

20 [0074] In some embodiments, a compound according to the invention may exhibit superior selectivity in inhibiting an O-GlcNAcase. In some embodiments, one or more of the compounds according to the invention may be more selective for an O-GlcNAcase over a β -hexosaminidase. In some embodiments, one or more of the compounds may selectively inhibit the activity of a mammalian O-GlcNAcase over a mammalian β -hexosaminidase. In
25 some embodiments, a selective inhibitor of an O-GlcNAcase may not substantially inhibit a β -hexosaminidase. In some embodiments, the β -hexosaminidase may be a mammalian β -hexosaminidase, such as a rat, mouse or human β -hexosaminidase. A compound that “selectively” inhibits an O-GlcNAcase is a compound that may inhibit the activity or biological function of an O-GlcNAcase, but may not substantially inhibit the activity or
30 biological function of a β -hexosaminidase. For example, in some embodiments, a selective inhibitor of an O-GlcNAcase may selectively inhibit the cleavage of 2-acetamido-2-deoxy- β -D-glucopyranoside (O-GlcNAc) from polypeptides. In some embodiments, a selective inhibitor of an O-GlcNAcase may selectively bind to an O-GlcNAcase. In some

embodiments, a selective inhibitor of an O-GlcNAcase may inhibit hyperphosphorylation of a tau protein and/or inhibit formations of NFTs. By “inhibit,” “inhibition” or “inhibiting” means a decrease by any value between 10% and 90%, or of any integer value between 30% and 60%, or over 100%, or a decrease by 1-fold, 2-fold, 5-fold, 10-fold or more. It is to be understood that the inhibiting does not require full inhibition. In some embodiments, a selective inhibitor of an O-GlcNAcase may elevate or enhance O-GlcNAc levels e.g., O-GlcNAc-modified polypeptide or protein levels, in cells, tissues, or organs (e.g., in brain, muscle, or heart (cardiac) tissue) and in animals. By “elevating” or “enhancing” is meant an increase by any value between 10% and 90%, or of any integer value between 30% and 60%, or over 100%, or an increase by 1-fold, 2-fold, 5-fold, 10-fold, 15-fold, 25-fold, 50-fold, 100-fold or more. In some embodiments, a selective inhibitor of an O-GlcNAcase may exhibit a selectivity ratio, as described herein, in the range 10 to 100000, or in the range 100 to 100000, or in the range 1000 to 100000, or at least 10, 20, 50, 100, 200, 500, 1000, 1500, 2000, 2500, 3000, 3500, 4000, 4500, 5000, 6000, 7000, 10,000, 25,000, 50,000, 75,000, or any value within or about the described range.


[0075] One or more of the compounds of the present invention may elevate O-GlcNAc levels on O-GlcNAc-modified polypeptides or proteins *in vivo* specifically via interaction with an O-GlcNAcase enzyme, and may be effective in treating conditions which require or respond to inhibition of O-GlcNAcase activity.

[0076] In some embodiments, one or more of the compounds of the present invention may be useful as agents that produce a decrease in tau phosphorylation and NFT formation. In some embodiments, one or more of the compounds may therefore be useful to treat Alzheimer’s disease and related tauopathies. In some embodiments, one or more of the compounds may thus be capable of treating Alzheimer’s disease and related tauopathies by lowering tau phosphorylation and reducing NFT formation as a result of increasing tau O-GlcNAc levels. In some embodiments, one or more of the compounds may produce an increase in levels of O-GlcNAc modification on O-GlcNAc-modified polypeptides or proteins, and may therefore be useful for treatment of disorders responsive to such increases in O-GlcNAc modification; these disorders may include, without limitation, neurodegenerative, inflammatory, cardiovascular, and immunoregulatory diseases. In some embodiments, a compound may also be useful as a result of other biological activities related to its ability to inhibit the activity of glycosidase enzymes. In alternative embodiments, one or more of the compounds of the

invention may be valuable tools in studying the physiological role of O-GlcNAc at the cellular and organismal level.

[0077] In alternative embodiments, the invention provides methods of enhancing or elevating levels of protein O-GlcNAc modification in animal subjects, such as, veterinary and human subjects. In alternative embodiments, the invention provides methods of selectively inhibiting an O-GlcNAcase enzyme in animal subjects, such as, veterinary and human subjects. In alternative embodiments, the invention provides methods of inhibiting phosphorylation of tau polypeptides, or inhibiting formation of NFTs, in animal subjects, such as, veterinary and human subjects.

10 [0078] In specific embodiments, the invention provides compounds described generally by Formula (I) and the salts, prodrugs, and enantiomeric forms thereof:



(I)

[0079] As set forth in Formula (I): R¹ may be H and R² may be C(O)NR¹¹₂, or R¹ may be C(O)NR¹¹₂ and R² may be H, or R¹ may be H and R² may be selected from the group consisting of: H, CH₃, CH₂F, CHF₂, and CH₂OH, where each R¹¹ may be independently H or C₁₋₆ alkyl; R³ may be H and R⁴ may be OH, or R³ may be H and R⁴ may be H, or R³ may be H and R⁴ may be F, or R³ may be F and R⁴ may be H, or R³ may be F and R⁴ may be F, or R³ may be OH and R⁴ may be H; R⁵ may be H and R⁶ may be OH, or R⁵ may be H and R⁶ may be H, or R⁵ may be H and R⁶ may be F, or R⁵ may be F and R⁶ may be H, or R⁵ may be F and R⁶ may be F, or R⁵ may be OH and R⁶ may be H; R⁷ may be H and R⁸ may be OH, or R⁷ may be H and R⁸ may be H, or R⁷ may be H and R⁸ may be F, or R⁷ may be F and R⁸ may be H, or R⁷ may be F and R⁸ may be F, or R⁷ may be OH and R⁸ may be H; R⁹ may be selected from the group consisting of: H, CH₃, CH₂F, CHF₂, and CH₂OH; and R¹⁰ may be selected from the group consisting of: H, C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl, C₁₋₁₀ acyl, C₈₋₂₀ arylalkylacyl, C₃₋₂₀ heteroarylalkylacyl, C₇₋₂₀ arylalkyl, C₈₋₂₀ arylalkenyl, C₈₋₂₀ arylalkynyl, C₂₋₂₀ heteroarylalkyl, C₃₋₂₀ heteroarylalkenyl, and C₃₋₂₀ heteroarylalkynyl, each excluding H optionally substituted from one to four substituents with one of more of halo, OH, OCF₃, CN,

SO₂NH₂, SO₂Me, C(O)NH₂, CH₂F, CHF₂, CF₃, CH₂CH₂F, CH₂CF₃, CH₂CH₂CH₂F, C₁₋₆ alkyl, C₁₋₆ alkoxy, or C₃₋₇ cycloalkyl.

[0080] In some embodiments, R¹ as set forth in Formula (I) may be H or C(O)NR¹¹₂, where each R¹¹ may be independently H or C₁₋₆ alkyl. In some embodiments, R¹ may be C(O)NR¹¹₂, where each R¹¹ may be independently H or C₁₋₆ alkyl. In some embodiments, R¹ may be C(O)NHMe or C(O)NHCH₂CH₃. In some embodiments, R¹ may be H.

[0081] In some embodiments, R² as set forth in Formula (I) may be selected from the group consisting of: H, CH₃, CH₂F, CHF₂, CH₂OH, and C(O)NR¹¹₂, where each R¹¹ may be independently H or C₁₋₆ alkyl. In some embodiments, R² may be C(O)NR¹¹₂, where each R¹¹ may be independently H or C₁₋₆ alkyl. In some embodiments, R² may be C(O)NHR¹¹, where R¹¹ may be H or C₁₋₆ alkyl. In some embodiments, R² may be C(O)NH₂, C(O)NHCH₃ or C(O)NHCH₂CH₃. In some embodiments, R² may be CH₂OH. In some embodiments, R² may be CH₂F. In some embodiments, R² may be CHF₂. In some embodiments, R² may be CH₃. In some embodiments, R² may be H.

[0082] In some embodiments, R³ as set forth in Formula (I) may be H, F, or OH. In some embodiments, R³ may be H or F. In some embodiments, R³ may be F.

[0083] In some embodiments, R⁴ as set forth in Formula (I) may be H, F, or OH. In some embodiments, R⁴ may be H or F. In some embodiments, R⁴ may be F.

[0084] In some embodiments, R⁵ as set forth in Formula (I) may be H, F, or OH. In some embodiments, R⁵ may be H or F. In some embodiments, R⁵ may be F.

[0085] In some embodiments, R⁶ as set forth in Formula (I) may be H, F, or OH. In some embodiments, R⁶ may be H or F. In some embodiments, R⁶ may be F.

[0086] In some embodiments, R⁷ as set forth in Formula (I) may be H, F, or OH. In some embodiments, R⁷ may be H or F. In some embodiments, R⁷ may be F.



[0087] In some embodiments, R⁸ as set forth in Formula (I) may be H, F, or OH. In some embodiments, R⁸ may be H or F. In some embodiments, R⁸ may be F.








[0088] In some embodiments, R⁹ as set forth in Formula (I) may be selected from the group consisting of: H, CH₃, CH₂F, CHF₂, and CH₂OH. In some embodiments, R⁹ may be CH₂OH. In some embodiments, R⁹ may be CH₂F or CHF₂. In some embodiments, R⁹ may be CH₃. In some embodiments, R⁹ may be H.

[0089] In some embodiments, R¹⁰ as set forth in Formula (I) may be selected from the group consisting of: H, C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl, C₁₋₁₀ acyl, C₈₋₂₀ arylalkylacyl, C₃₋₂₀ heteroarylalkylacyl, C₇₋₂₀ arylalkyl, C₈₋₂₀ arylalkenyl, C₈₋₂₀ arylalkynyl, C₂₋₂₀ heteroarylalkyl, C₃₋₂₀ heteroarylalkenyl, and C₃₋₂₀ heteroarylalkynyl, each excluding H optionally substituted from one to four substituents with one of more of halo, OH, OCF₃, CN, SO₂NH₂, SO₂Me, C(O)NH₂, CH₂F, CHF₂, CF₃, CH₂CH₂F, CH₂CF₃, CH₂CH₂CH₂F, C₁₋₆ alkyl, C₁₋₆ alkoxy, or C₃₋₇ cycloalkyl. In some embodiments, R¹⁰ may be selected from the group consisting of: C₁₋₁₀ alkyl, C₇₋₂₀ arylalkyl, and C₂₋₂₀ heteroarylalkyl, each optionally substituted from one to four substituents with one of more of halo, OH, OCF₃, CN, SO₂NH₂, SO₂Me, C(O)NH₂, CH₂F, CHF₂, CF₃, CH₂CH₂F, CH₂CF₃, CH₂CH₂CH₂F, C₁₋₆ alkyl, C₁₋₆ alkoxy, or C₃₋₇ cycloalkyl. In some embodiments, R¹⁰ may be selected from the group consisting of: H, CH₃, hexyl, (CH₂)₂(phenyl), (CH₂)₃(phenyl), (CH₂)₄(phenyl), 5-methylhexyl, (E)-3-([1,1'-biphenyl]-4-yl)allyl, 3-([1,1'-biphenyl]-4-yl)propyl, 3-phenylpropanoyl, 3-(6-fluoropyridin-3-yl)propyl, 4,4,4-trifluorobutyl, 3-(5-(trifluoromethoxy)benzo[d]thiazol-2-yl)propyl, (Z)-5,5,5-trifluoropent-3-en-1-yl, pent-4-yn-1-yl, butyryl, 3-(6-fluoropyridin-3-yl)propanoyl, cinnamyl, 3-phenylprop-2-yn-1-yl, (E)-3-(6-fluoropyridin-3-yl)allyl, and 3-(6-fluoropyridin-3-yl)prop-2-yn-1-yl.








[0090] In specific embodiments of the invention, compounds according to Formula (I) include the compounds described in Table 1.







20 **Table 1**







Example	Name	Structure
1	(2S,3R,4S,5R,6R)-3,4,5-trihydroxy-6-(hydroxymethyl)-N,1-dimethylpiperidine-2-carboxamide hydrochloride	
2	(2S,3R,4S,5R,6R)-1-((E)-3-([1,1'-biphenyl]-4-yl)allyl)-3,4,5-trihydroxy-6-(hydroxymethyl)-N-methylpiperidine-2-carboxamide	








Example	Name	Structure
3	(2S,3R,4S,5R,6R)-1-(3-([1,1'-biphenyl]-4-yl)propyl)-3,4,5-trihydroxy-6-(hydroxymethyl)-N-methylpiperidine-2-carboxamide	
4	(2S,3R,4S,5R,6R)-N-ethyl-3,4,5-trihydroxy-6-(hydroxymethyl)piperidine-2-carboxamide hydrochloride	
5	(2S,3R,4S,5R,6R)-3,4,5-trihydroxy-6-(hydroxymethyl)-N-methyl-1-(3-phenylpropyl)piperidine-2-carboxamide	
6	(2S,3R,4S,5R,6R)-N-ethyl-3,4,5-trihydroxy-6-(hydroxymethyl)-1-(3-phenylpropyl)piperidine-2-carboxamide	
7	(2S,3R,4S,5R,6R)-3,4,5-trihydroxy-6-(hydroxymethyl)-N-methyl-1-(4-phenylbutyl)piperidine-2-carboxamide	
8	(2S,3R,4S,5R,6R)-3,4,5-trihydroxy-6-(hydroxymethyl)-N-methyl-1-(5-methylhexyl)piperidine-2-carboxamide	
9	(2S,3R,4S)-1-hexyl-3,4-dihydroxy-N-methylpiperidine-2-carboxamide	








Example	Name	Structure
10	(2S,3R,4S,5R)-3,4,5-trihydroxy-N-methylpiperidine-2-carboxamide	
11	(2S,3R,4S,5R)-3,4,5-trihydroxy-N,1-dimethylpiperidine-2-carboxamide	
12	(2S,3R,4S,5R)-3,4,5-trihydroxy-1-methylpiperidine-2-carboxamide	
13	(2S,3R,4S,5R)-N-ethyl-3,4,5-trihydroxypiperidine-2-carboxamide	
14	(2S,3R,4S,5R)-N-ethyl-3,4,5-trihydroxy-1-methylpiperidine-2-carboxamide	
15	(2S,3R,4S,5R,6R)-N-ethyl-3,4,5-trihydroxy-6-methylpiperidine-2-carboxamide	
16	(2S,3R,4S,5R,6R)-N-ethyl-3,4,5-trihydroxy-6-methyl-1-phenethylpiperidine-2-carboxamide	
17	(2S,3R,4S,5R)-N-ethyl-3,4,5-trihydroxy-1-(3-phenylpropyl)piperidine-2-carboxamide	

Example	Name	Structure
18	(2S,3S,4R,5S,6R)-N-ethyl-4-fluoro-3,5-dihydroxy-6-methyl-1-phenethylpiperidine-2-carboxamide	
19	(2R,3R,4S,5R,6R)-N-ethyl-3-fluoro-4,5-dihydroxy-6-(hydroxymethyl)-1-(3-phenylpropyl)piperidine-2-carboxamide	
20	(2S,3R,4S,5R,6R)-N-ethyl-4-fluoro-3,5-dihydroxy-6-(hydroxymethyl)-1-(3-phenylpropyl)piperidine-2-carboxamide	
21	(2S,3R,4R,5R,6R)-N-ethyl-5-fluoro-3,4-dihydroxy-6-(hydroxymethyl)-1-(3-phenylpropyl)piperidine-2-carboxamide	
22	(2S,3R,4R,5R,6S)-N-ethyl-6-(fluoromethyl)-3,4,5-trihydroxy-1-(3-phenylpropyl)piperidine-2-carboxamide	
23	(2R,3S,4S,5R,6R)-N-ethyl-3-fluoro-4,5-dihydroxy-6-(hydroxymethyl)-1-(3-phenylpropyl)piperidine-2-carboxamide	
24	(2S,3R,4R,5R,6R)-N-ethyl-4-fluoro-3,5-dihydroxy-6-(hydroxymethyl)-1-(3-phenylpropyl)piperidine-2-carboxamide	

Example	Name	Structure
25	(2S,3R,4R,5S,6R)-N-ethyl-5-fluoro-3,4-dihydroxy-6-(hydroxymethyl)-1-(3-phenylpropyl)piperidine-2-carboxamide	
26	(2S,3R,4R,5R,6S)-6-(difluoromethyl)-N-ethyl-3,4,5-trihydroxy-1-(3-phenylpropyl)piperidine-2-carboxamide	
27	(2S,4R,5R,6R)-N-ethyl-4,5-dihydroxy-6-(hydroxymethyl)-1-(3-phenylpropyl)piperidine-2-carboxamide	
28	(2S,3S,5S,6R)-N-ethyl-3,5-dihydroxy-6-(hydroxymethyl)-1-(3-phenylpropyl)piperidine-2-carboxamide	
29	(2S,3R,4S,6S)-N-ethyl-3,4-dihydroxy-6-(hydroxymethyl)-1-(3-phenylpropyl)piperidine-2-carboxamide	
30	(2S,3R,4S,5R,6R)-N-ethyl-3,4,5-trihydroxy-6-methyl-1-(3-phenylpropyl)piperidine-2-carboxamide	

Example	Name	Structure
31	(2S,3R,4S,5R)-N-ethyl-3,4,5-trihydroxy-1-(3-phenylpropyl)piperidine-2-carboxamide	
32	(2S,3R,4S,5R,6S)-6-(difluoromethyl)-N-ethyl-4-fluoro-3,5-dihydroxy-1-(3-phenylpropyl)piperidine-2-carboxamide	
33	(2R,3S,4S,5R,6S)-N-ethyl-3-fluoro-6-(fluoromethyl)-4,5-dihydroxy-1-(3-phenylpropyl)piperidine-2-carboxamide	
34	(2R,3R,4S,5R,6S)-6-(difluoromethyl)-N-ethyl-3-fluoro-4,5-dihydroxy-1-(3-phenylpropyl)piperidine-2-carboxamide	
35	(2R,3R,4S,5R)-N-ethyl-3-fluoro-4,5-dihydroxy-1-(3-phenylpropyl)piperidine-2-carboxamide	
36	(2S,3R,4S,5R)-N-ethyl-4-fluoro-3,5-dihydroxy-1-(3-phenylpropyl)piperidine-2-carboxamide	

Example	Name	Structure
37	(2S,3R,4R,5R)-N-ethyl-5-fluoro-3,4-dihydroxy-1-(3-phenylpropyl)piperidine-2-carboxamide	
38	(2R,3S,4S,5R)-N-ethyl-3-fluoro-4,5-dihydroxy-1-(3-phenylpropyl)piperidine-2-carboxamide	
39	(2S,3R,4R,5R)-N-ethyl-4-fluoro-3,5-dihydroxy-1-(3-phenylpropyl)piperidine-2-carboxamide	
40	(2S,3R,4R,5S)-N-ethyl-5-fluoro-3,4-dihydroxy-1-(3-phenylpropyl)piperidine-2-carboxamide	
41	(2S,4R,5R)-N-ethyl-4,5-dihydroxy-1-(3-phenylpropyl)piperidine-2-carboxamide	
42	(2S,3S,5S)-N-ethyl-3,5-dihydroxy-1-(3-phenylpropyl)piperidine-2-carboxamide	
43	(2S,3R,4S)-N-ethyl-3,4-dihydroxy-1-(3-phenylpropyl)piperidine-2-carboxamide	










Example	Name	Structure
44	(2R,3R,4S,5R)-N-ethyl-3,4-difluoro-5-hydroxy-1-(3-phenylpropyl)piperidine-2-carboxamide	
45	(2S,3R,4S,5R,6R)-N-ethyl-3,4,5-trihydroxy-6-(hydroxymethyl)-1-(3-phenylpropanoyl)piperidine-2-carboxamide	
46	(2S,3R,4S,5R,6R)-N-ethyl-1-(3-(6-fluoropyridin-3-yl)propyl)-3,4,5-trihydroxy-6-(hydroxymethyl)piperidine-2-carboxamide	
47	(2S,3R,4S,5R,6R)-N-ethyl-3,4,5-trihydroxy-6-(hydroxymethyl)-1-(4,4,4-trifluorobutyl)piperidine-2-carboxamide	
48	(2S,3R,4S,5R,6R)-N-ethyl-3,4,5-trihydroxy-6-(hydroxymethyl)-1-(3-(5-(trifluoromethoxy)benzo[d]thiazol-2-yl)propyl)piperidine-2-carboxamide	
49	(2S,3R,4S,5R,6R)-N-ethyl-3,4,5-trihydroxy-6-(hydroxymethyl)-1-((Z)-5,5,5-trifluoropent-3-en-1-yl)piperidine-2-carboxamide	
50	(2S,3R,4S,5R,6R)-N-ethyl-3,4,5-trihydroxy-6-(hydroxymethyl)-1-(pent-4-yn-1-yl)piperidine-2-carboxamide	











Example	Name	Structure
51	(2S,3R,4S,5R,6R)-1-butyryl-N-ethyl-3,4,5-trihydroxy-6-(hydroxymethyl)piperidine-2-carboxamide	
52	(2S,3R,4S,5R,6R)-N-ethyl-1-(3-(6-fluoropyridin-3-yl)propanoyl)-3,4,5-trihydroxy-6-(hydroxymethyl)piperidine-2-carboxamide	
53	(2S,3R,4S,5R,6R)-1-cinnamyl-N-ethyl-3,4,5-trihydroxy-6-(hydroxymethyl)piperidine-2-carboxamide	
54	(2S,3R,4S,5R,6R)-N-ethyl-3,4,5-trihydroxy-6-(hydroxymethyl)-1-(3-phenylprop-2-yn-1-yl)piperidine-2-carboxamide	
55	(2S,3R,4S,5R,6R)-N-ethyl-1-((E)-3-(6-fluoropyridin-3-yl)allyl)-3,4,5-trihydroxy-6-(hydroxymethyl)piperidine-2-carboxamide	
56	(2S,3R,4S,5R,6R)-N-ethyl-1-(3-(6-fluoropyridin-3-yl)prop-2-yn-1-yl)-3,4,5-trihydroxy-6-(hydroxymethyl)piperidine-2-carboxamide	










[0091] In alternative embodiments of the invention, compounds according to Formula (I) include one or more of the compounds described in Table 2.










[0092] In alternative embodiments of the invention, one or more of the compounds described in Table 2 are specifically excluded from the compounds described in Formula (I).









Table 2








Name	Structure
(2S,3R,4S,5R,6R)-3,4,5-trihydroxy-6-(hydroxymethyl)-N-methylpiperidine-2-carboxamide	
(2R,3R,4S,5R,6R)-3,4,5-trihydroxy-6-(hydroxymethyl)-N-methylpiperidine-2-carboxamide	
(2R,3S,4R,5S,6R)-3,4,5-trihydroxy-6-methylpiperidine-2-carboxamide	
(2R,3S,4R,5S,6R)-3,4,5-trihydroxy-N,6-dimethylpiperidine-2-carboxamide	
(2R,3S,4R,5S,6R)-N-butyl-3,4,5-trihydroxy-6-methylpiperidine-2-carboxamide	
(2S,3R,4R,5S)-3,4,5-trihoxypiperidine-2-carboxamide	
(2S,3R,4R,5S)-3,4,5-trihydroxy-1-methylpiperidine-2-carboxamide	
(2R,3R,4R,5S)-3,4,5-trihydroxy-N-methylpiperidine-2-carboxamide	
(2S,3R,4R,5S)-N-(tert-butyl)-3,4,5-trihydroxypiperidine-2-carboxamide	

Name	Structure
(2S,3S,4R,5S)-N-(tert-butyl)-3,4,5-trihydroxypiperidine-2-carboxamide	
(2S,3R,4R,5R)-N-(tert-butyl)-3,4,5-trihydroxypiperidine-2-carboxamide	
(2S,3S,4R,5R)-N-(tert-butyl)-3,4,5-trihydroxypiperidine-2-carboxamide	
(2R,3S,5S)-3,5-dihydroxypiperidine-2-carboxamide	
(2R,3S,4R,5R,6R)-2,6-bis(hydroxymethyl)piperidine-3,4,5-triol	
(2R,3R,5R,6R)-2,6-bis(hydroxymethyl)piperidine-3,4,5-triol	
(2R,3S,5S,6R)-2,6-bis(hydroxymethyl)piperidine-3,4,5-triol	
(2R,3R,4S,5S,6R)-2,6-bis(hydroxymethyl)piperidine-3,4,5-triol	
(2R,3R,4S,5S,6R)-2,6-bis(hydroxymethyl)-1-methylpiperidine-3,4,5-triol	
(2R,3S,4R,5R,6R)-2,6-bis(hydroxymethyl)-1-methylpiperidine-3,4,5-triol	

Name	Structure
(2R,3S,5S,6R)-1-(2-hydroxyethyl)-2,6-bis(hydroxymethyl)piperidine-3,4,5-triol	
(2R,3S,5S,6R)-1-butyl-2,6-bis(hydroxymethyl)piperidine-3,4,5-triol	
(2R,3R,4R,5S,6R)-1-butyl-2,6-bis(hydroxymethyl)piperidine-3,4,5-triol	
(2R,3S,5S,6R)-1-hexyl-2,6-bis(hydroxymethyl)piperidine-3,4,5-triol	
(2R,3S,5S,6R)-2,6-bis(hydroxymethyl)-1-nonylpiperidine-3,4,5-triol	
(2R,3R,4R,5S,6R)-2-(hydroxymethyl)-6-methylpiperidine-3,4,5-triol	
(2R,3S,4R,5S,6R)-2-(hydroxymethyl)-6-methylpiperidine-3,4,5-triol	
(2R,3S,4S,5S,6R)-2-(hydroxymethyl)-6-methylpiperidine-3,4,5-triol	
(2R,3R,4S,5R,6R)-2-(hydroxymethyl)-6-methylpiperidine-3,4,5-triol	

Name	Structure
(2R,3R,4R,5R,6R)-2-(hydroxymethyl)-6-methylpiperidine-3,4,5-triol	
(2R,3R,4S,5S,6R)-2-(hydroxymethyl)-6-methylpiperidine-3,4,5-triol	
(2S,3R,4R,5S)-2-(fluoromethyl)piperidine-3,4,5-triol	
(2S,3R,4R,5S)-2-(fluoromethyl)-1-(2-hydroxyethyl)piperidine-3,4,5-triol	
(3R,4S,6S)-6-(fluoromethyl)piperidine-3,4-diol	
(2S,3R,4R,5S)-4-fluoro-2-(fluoromethyl)piperidine-3,5-diol	
(3R,4r,5S)-piperidine-3,4,5-triol	
(3R,4r,5S)-1-methylpiperidine-3,4,5-triol	
(3R,4r,5S)-1-(2-hydroxyethyl)piperidine-3,4,5-triol	


Name	Structure
(3R,4r,5S)-1-propylpiperidine-3,4,5-triol	
(3S,4s,5R)-1-propylpiperidine-3,4,5-triol	
2-((3R,4r,5S)-3,4,5-trihydroxypiperidin-1-yl)acetonitrile	
(3S,4s,5R)-1-butylpiperidine-3,4,5-triol	
(3R,4r,5S)-1-butylpiperidine-3,4,5-triol	
(3R,4r,5S)-1-pentylpiperidine-3,4,5-triol	
(3R,4r,5S)-1-hexylpiperidine-3,4,5-triol	
(3R,4r,5S)-1-octylpiperidine-3,4,5-triol	

Name	Structure
(3R,4r,5S)-1-nonylpiperidine-3,4,5-triol	
(3S,4s,5R)-1-nonylpiperidine-3,4,5-triol	
(3R,4r,5S)-1-decylpiperidine-3,4,5-triol	
6-((3R,4r,5S)-3,4,5-trihydroxypiperidin-1-yl)hexanenitrile	
(3R,4r,5S)-1-benzylpiperidine-3,4,5-triol	
(3S,4s,5R)-1-benzylpiperidine-3,4,5-triol	
(3R,4r,5S)-1-(3-phenylpropyl)piperidine-3,4,5-triol	

[0093] In alternative embodiments of the invention, compounds according to Formula (I) include one or more of the compounds described in Table 2 in which one or more OH groups are replaced by F or H.

[0094] In alternative embodiments of the invention, one or more of the compounds described in Table 2 in which one or more OH groups is replaced by F or H are specifically excluded from the compounds described in Formula (I).

[0095] As will be appreciated by a person skilled in the art, Formula (I) above may also be represented alternatively as follows:



[0096] As used herein the singular forms “a”, “and”, and “the” include plural referents unless the context clearly dictates otherwise. For example, “a compound” refers to one or more of such compounds, while “the enzyme” includes a particular enzyme as well as other family members and equivalents thereof as known to those skilled in the art.

[0097] Throughout this application, it is contemplated that the term “compound” or “compounds” refers to the compounds discussed herein and includes precursors and derivatives of the compounds, including acyl-protected derivatives, and pharmaceutically acceptable salts of the compounds, precursors, and derivatives. The invention also includes prodrugs of the compounds, pharmaceutical compositions including the compounds and a pharmaceutically acceptable carrier, and pharmaceutical compositions including prodrugs of the compounds and a pharmaceutically acceptable carrier.

[0098] The compounds of the present invention may contain one or more asymmetric centers and can thus occur as racemates and racemic mixtures, single enantiomers, diastereomeric mixtures and individual diastereomers. Additional asymmetric centers may be present depending upon the nature of the various substituents on the molecule. Each such asymmetric center will independently produce two optical isomers and it is intended that all of the possible optical isomers and diastereomers in mixtures and as pure or partially purified compounds are included within the ambit of this invention. Any formulas, structures or names of compounds described in this specification that do not specify a particular

stereochemistry are meant to encompass any and all existing isomers as described above and mixtures thereof in any proportion. When stereochemistry is specified, the invention is meant to encompass that particular isomer in pure form or as part of a mixture with other isomers in any proportion.

- 5 [0099] “Alkyl” refers to a straight or branched hydrocarbon chain group consisting solely of carbon and hydrogen atoms, containing no unsaturation and including, for example, from one to ten carbon atoms, such as 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 carbon atoms, and which is attached to the rest of the molecule by a single bond. In alternative embodiments, the alkyl group may contain from one to eight carbon atoms, such as 1, 2, 3, 4, 5, 6, 7, or 8 carbon atoms. In
10 alternative embodiments, the alkyl group may contain from one to six carbon atoms, such as 1, 2, 3, 4, 5, or 6 carbon atoms. Unless stated otherwise specifically in the specification, the alkyl group may be optionally substituted by one or more substituents as described herein. Unless stated otherwise specifically herein, it is understood that the substitution can occur on any carbon of the alkyl group.
- 15 [00100] “Alkenyl” refers to a straight or branched hydrocarbon chain group consisting solely of carbon and hydrogen atoms, containing at least one double bond and including, for example, from two to ten carbon atoms, such as 2, 3, 4, 5, 6, 7, 8, 9, or 10 carbon atoms, and which is attached to the rest of the molecule by a single bond or a double bond. In alternative
20 embodiments, the alkenyl group may contain from two to eight carbon atoms, such as 2, 3, 4, 5, 6, 7, or 8 carbon atoms. In alternative embodiments, the alkenyl group may contain from three to six carbon atoms, such as 3, 4, 5, or 6 carbon atoms. Unless stated otherwise specifically in the specification, the alkenyl group may be optionally substituted by one or more substituents as described herein. Unless stated otherwise specifically herein, it is understood that the substitution can occur on any carbon of the alkenyl group.
- 25 [00101] “Alkynyl” refers to a straight or branched hydrocarbon chain group consisting solely of carbon and hydrogen atoms, containing at least one triple bond and including, for example, from two to ten carbon atoms. In alternative embodiments, the alkynyl group may contain from two to eight carbon atoms, such as 2, 3, 4, 5, 6, 7, or 8 carbon atoms. In alternative
30 embodiments, the alkynyl group may contain from three to six carbon atoms, such as 3, 4, 5, or 6 carbon atoms. Unless stated otherwise specifically in the specification, the alkynyl group may be optionally substituted by one or more substituents as described herein.

[00102] “Aryl” refers to a mono- or bicyclic aromatic ring containing only carbon atoms, including for example, 6-14 members, such as 6, 7, 8, 9, 10, 11, 12, 13, or 14 members.

Examples of aryl groups include phenyl, biphenyl, naphthyl, indanyl, indenyl, tetrahydronaphthyl, 2,3-dihydrobenzofuranyl, dihydrobenzopyranyl, 1,4-benzodioxanyl, and the like. Unless stated otherwise specifically herein, the term “aryl” is meant to include aryl groups optionally substituted by one or more substituents as described herein.

[00103] “Heteroaryl” refers to a single or fused aromatic ring group containing one or more heteroatoms in the ring, for example N, O, S, including for example, 5-14 members, such as 5, 6, 7, 8, 9, 10, 11, 12, 13, or 14 members. Examples of heteroaryl groups include furan, thiophene, pyrrole, oxazole, thiazole, imidazole, pyrazole, isoxazole, isothiazole, 1,2,3-oxadiazole, 1,2,3-triazole, 1,2,4-triazole, 1,3,4-thiadiazole, tetrazole, pyridine, pyridazine, pyrimidine, pyrazine, 1,3,5-triazine, imidazole, benzimidazole, benzoxazole, benzothiazole, indolizine, indole, isoindole, benzofuran, benzothiophene, 1H-indazole, purine, 4H-quinolizine, quinoline, isoquinoline, cinnoline, phthalazine, quinazoline, quinoxaline, 1,8-naphthyridine, pteridine, and the like. Unless stated otherwise specifically herein, the term “heteroaryl” is meant to include heteroaryl groups optionally substituted by one or more substituents as described herein.

[00104] “Arylalkyl” refers to a group of the formula $-R_aR_b$ where R_a is a C_{1-10} alkyl group as described herein and R_b is one or more aryl moieties as described herein. The aryl group(s) may be optionally substituted as described herein.

[00105] “Arylalkenyl” refers to a group of the formula $-R_cR_b$ where R_c is a C_{2-10} alkenyl group as described herein and R_b is one or more aryl moieties as described herein. The aryl group(s) may be optionally substituted as described herein.

[00106] “Arylalkynyl” refers to a group of the formula $-R_dR_b$ where R_d is a C_{2-10} alkynyl group as described herein and R_b is one or more aryl moieties as described herein. The aryl group(s) may be optionally substituted as described herein.

[00107] “Heteroarylalkyl” refers to a group of the formula $-R_aR_e$ where R_a is a C_{1-10} alkyl group as described herein and R_e is one or more heteroaryl moieties as described herein. The aryl group(s) may be optionally substituted as described herein.

[00108] “Heteroarylalkenyl” refers to a group of the formula $-R_cR_e$ where R_c is a C_{2-10} alkenyl group as described herein and R_e is one or more heteroaryl moieties as described herein. The heteroaryl group(s) may be optionally substituted as described herein.

[00109] “Heteroarylalkynyl” refers to a group of the formula $-R_dR_e$ where R_d is a C_{2-10} alkynyl group as described herein and R_e is one or more heteroaryl moieties as described herein. The heteroaryl group(s) may be optionally substituted as described herein.

5 [00110] “Acyl” refers to a group of the formula $-C(O)R_f$, where R_f is H or a C_{1-10} alkyl group or a C_{1-6} alkyl group or a C_{3-15} cycloalkyl group as described herein. The alkyl or cycloalkyl group(s) may be optionally substituted as described herein.

[00111] “Arylalkylacyl” refers to a group of the formula $-C(O)R_gR_b$, where R_g is a C_{1-10} alkyl or a C_{1-6} alkyl group as described herein and R_b is one or more aryl moieties as described herein. The alkyl or aryl group(s) may be optionally substituted as described herein.

10 [00112] “Heteroarylalkylacyl” refers to a group of the formula $-C(O)R_gR_e$, where R_g is a C_{1-10} alkyl or a C_{1-6} alkyl group as described herein and R_e is one or more heteroaryl moieties as described herein. The alkyl or heteroaryl group(s) may be optionally substituted as described herein.

15 [00113] “Alkoxy” refers to a group of the formula $-OR_g$, where R_g is a C_{1-10} alkyl or a C_{1-6} alkyl group as described herein. The alkyl group(s) may be optionally substituted as described herein.

[00114] “Cycloalkyl” refers to a stable monovalent monocyclic, bicyclic or tricyclic hydrocarbon group consisting solely of carbon and hydrogen atoms, having for example from 3 to 15 carbon atoms, and which is saturated and attached to the rest of the molecule by a single bond. In alternative embodiments, the cycloalkyl group may contain from three to six carbon atoms, such as 3, 4, 5, or 6 carbon atoms. Unless otherwise stated specifically herein, the term “cycloalkyl” is meant to include cycloalkyl groups which are optionally substituted as described herein.

25 [00115] “Halo” refers to bromo, chloro, fluoro, iodo, etc. In some embodiments, suitable halogens include fluorine or chlorine.

[00116] “Optional” or “optionally” means that the subsequently described event of circumstances may or may not occur, and that the description includes instances where said event or circumstance occurs one or more times and instances in which it does not. For example, “optionally substituted alkyl” means that the alkyl group may or may not be substituted and that the description includes both substituted alkyl groups and alkyl groups having no substitution, and that said alkyl groups may be substituted one or more times. Examples of optionally substituted alkyl groups include, without limitation, methyl, ethyl,

propyl, etc. and including cycloalkyls such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, etc.; examples of optionally substituted alkenyl groups include allyl, crotyl, 2-pentenyl, 3-hexenyl, 2-cyclopentenyl, 2-cyclohexenyl, 2-cyclopentenylmethyl, 2-cyclohexenylmethyl, etc. In some embodiments, optionally substituted alkyl and alkenyl groups include C₁₋₆ alkyls or alkenyls.

Therapeutic Indications

[00117] The invention provides methods of treating conditions that are modulated, directly or indirectly, by an O-GlcNAcase enzyme or by O-GlcNAc-modified protein levels, for example, a condition that is benefited by inhibition of an O-GlcNAcase enzyme or by an elevation of O-GlcNAc-modified protein levels. Such conditions may include, without limitation, Glaucoma, Schizophrenia, tauopathies, such as Alzheimer's disease, neurodegenerative diseases, cardiovascular diseases, diseases associated with inflammation, diseases associated with immunosuppression and cancers. One or more of the compounds of the invention may also be useful in the treatment of diseases or disorders related to deficiency or over-expression of O-GlcNAcase or accumulation or depletion of O-GlcNAc, or any disease or disorder responsive to glycosidase inhibition therapy. Such diseases and disorders may include, but are not limited to, Glaucoma, Schizophrenia, neurodegenerative disorders, such as Alzheimer's disease (AD), or cancer. Such diseases and disorders may also include diseases or disorders related to the accumulation or deficiency in the enzyme OGT. Also included is a method of protecting or treating target cells expressing proteins that are modified by O-GlcNAc residues, the dysregulation of which modification may result in disease or pathology. The term "treating" as used herein includes treatment, prevention, and amelioration.

[00118] In alternative embodiments, the invention provides methods of enhancing or elevating levels of protein O-GlcNAc modification in animal subjects, such as, veterinary and human subjects. This elevation of O-GlcNAc levels may be useful for the prevention or treatment of Alzheimer's disease; prevention or treatment of other neurodegenerative diseases (e.g. Parkinson's disease, Huntington's disease); providing neuroprotective effects; preventing damage to cardiac tissue; and treating diseases associated with inflammation or immunosuppression.

[00119] In alternative embodiments, the invention provides methods of selectively inhibiting an O-GlcNAcase enzyme in animal subjects, such as veterinary and human subjects.

[00120] In alternative embodiments, the invention provides methods of inhibiting phosphorylation of tau polypeptides, or inhibiting formation of NFTs, in animal subjects, such as, veterinary and human subjects. Accordingly, a compound of the invention may be used to study and treat AD and other tauopathies.

[00121] In general, the methods of the invention may be effected by administering a compound according to the invention to a subject in need thereof, or by contacting a cell or a sample with a compound according to the invention, for example, a pharmaceutical composition comprising a therapeutically effective amount of the compound according to Formula (I). More particularly, they may be useful in the treatment of a disorder in which the regulation of O-GlcNAc protein modification is implicated, or any condition as described herein. Disease states of interest may include Alzheimer's disease (AD) and related neurodegenerative tauopathies, in which abnormal hyperphosphorylation of the microtubule-associated protein tau is involved in disease pathogenesis. In some embodiments, a compound may be used to block hyperphosphorylation of tau by maintaining elevated levels of O-GlcNAc on tau, thereby providing therapeutic benefit.

[00122] The effectiveness of a compound in treating pathology associated with the accumulation of toxic tau species (for example, Alzheimer's disease and other tauopathies) may be confirmed by testing the ability of a compound to block the formation of toxic tau species in established cellular¹²¹⁻¹²³ and/or transgenic animal models of disease.^{33,34}

[00123] Tauopathies that may be treated with a compound of the invention may include, without limitation: Alzheimer's disease, Amyotrophic lateral sclerosis (ALS), Amyotrophic lateral sclerosis with cognitive impairment (ALSci), Argyrophilic grain dementia, Bluit disease, Corticobasal degeneration (CBD), Dementia pugilistica, Diffuse neurofibrillary tangles with calcification, Down's syndrome, Familial British dementia, Familial Danish dementia, Frontotemporal dementia with parkinsonism linked to chromosome 17 (FTDP-17), Gerstmann-Straussler-Scheinker disease, Guadeloupean parkinsonism, Hallevorden-Spatz disease (neurodegeneration with brain iron accumulation type 1), Multiple system atrophy, Myotonic dystrophy, Niemann-Pick disease (type C), Pallido-ponto-nigral degeneration, Parkinsonism-dementia complex of Guam, Pick's disease (PiD), Post-encephalitic parkinsonism (PEP), Prion diseases (including Creutzfeldt-Jakob Disease (CJD), Variant

Creutzfeldt-Jakob Disease (vCJD), Fatal Familial Insomnia, and Kuru), Progressive supercortical gliosis, Progressive supranuclear palsy (PSP), Richardson's syndrome, Subacute sclerosing panencephalitis, Tangle-only dementia, and Glaucoma.

5 [00124] One or more of the compounds of this invention may also be useful in the treatment of conditions associate with tissue damage or stress, stimulating cells, or promoting differentiation of cells. Accordingly, in some embodiments, a compound of this invention may be used to provide therapeutic benefit in a variety of conditions or medical procedures involving stress in cardiac tissue; such conditions may include, without limitation: ischemia; hemorrhage; hypovolemic shock; myocardial infarction; an interventional cardiology
10 procedure; cardiac bypass surgery; fibrinolytic therapy; angioplasty; and stent placement.

[00125] The effectiveness of a compound in treating pathology associated with cellular stress (including ischemia, hemorrhage, hypovolemic shock, myocardial infarction, and other cardiovascular disorders) may be confirmed by testing the ability of a compound to prevent cellular damage in established cellular stress assays,^{108,119,120} and to prevent tissue damage
15 and promote functional recovery in animal models of ischemia-reperfusion,^{71,117} and trauma-hemorrhage.^{73,115,118}

[00126] Compounds that selectively inhibit O-GlcNAcase activity may be used for the treatment of diseases that are associated with inflammation; such conditions may include, without limitation: inflammatory or allergic diseases such as asthma, allergic rhinitis,
20 hypersensitivity lung diseases, hypersensitivity pneumonitis, eosinophilic pneumonias, delayed-type hypersensitivity, atherosclerosis, interstitial lung disease (ILD) (*e.g.*, idiopathic pulmonary fibrosis, or ILD associated with rheumatoid arthritis, systemic lupus erythematosus, ankylosing spondylitis, systemic sclerosis, Sjogren's syndrome, polymyositis or dermatomyositis); systemic anaphylaxis or hypersensitivity responses, drug allergies,
25 insect sting allergies; autoimmune diseases, such as rheumatoid arthritis, psoriatic arthritis, multiple sclerosis, Guillain-Barré syndrome, systemic lupus erythematosus, myasthenia gravis, glomerulonephritis, autoimmune thyroiditis, graft rejection, including allograft rejection or graft-versus-host disease; inflammatory bowel diseases, such as Crohn's disease and ulcerative colitis; spondyloarthropathies; scleroderma; psoriasis (including T-cell mediated
30 psoriasis) and inflammatory dermatoses such as dermatitis, eczema, atopic dermatitis, allergic contact dermatitis, urticaria; vasculitis (*e.g.*, necrotizing, cutaneous, and hypersensitivity vasculitis); eosinphilic myotis, eosiniphilic fasciitis; and cancers.

[00127] In addition, compounds that affect levels of protein O-GlcNAc modification may be used for the treatment of diseases associated with immunosuppression, such as, for example, in individuals undergoing chemotherapy, radiation therapy, enhanced wound healing and burn treatment, therapy for autoimmune disease or other drug therapy (*e.g.*, corticosteroid therapy) or combination of conventional drugs used in the treatment of autoimmune diseases and graft/transplantation rejection, which causes immunosuppression; or immunosuppression due to congenital deficiency in receptor function or other causes.

[00128] One or more of the compounds of the invention may be useful for treatment of neurodegenerative diseases; such conditions may include, without limitation, Parkinson's disease and Huntington's disease. Other conditions that may be treated are those triggered, affected, or in any other way correlated with levels of O-GlcNAc post-translational protein modification. It is expected that one or more of the compounds of this invention may be useful for the treatment of such conditions and in particular, but not limited to, the following for which a association with O-GlcNAc levels on proteins has been established: graft rejection, in particular but not limited to solid organ transplants, such as heart, lung, liver, kidney, and pancreas transplants (*e.g.* kidney and lung allografts); cancer, in particular but not limited to cancer of the breast, lung, prostate, pancreas, colon, rectum, bladder, kidney, ovary; as well as non-Hodgkin's lymphoma and melanoma; epilepsy, pain, fibromyalgia, or stroke, *e.g.*, for neuroprotection following a stroke.

Pharmaceutical & Veterinary Compositions, Dosages, And Administration

[00129] Pharmaceutical compositions including compounds according to the invention, or for use according to the invention, are contemplated as being within the scope of the invention. In some embodiments, pharmaceutical compositions including an effective amount of a compound of Formula (I) are provided.

[00130] The compounds of Formula (I) and their pharmaceutically acceptable salts, enantiomers, solvates, and derivatives may be useful because they may have pharmacological activity in animals, including humans. In some embodiments, one or more of the compounds according to the invention may be stable in plasma, when administered to a subject.

[00131] In some embodiments, a compound according to the invention, or for use according to the invention, may be provided in combination with any other active agents or pharmaceutical compositions where such combined therapy may be useful to modulate O-

GlcNAcase activity, for example, to treat neurodegenerative, inflammatory, cardiovascular, or immunoregulatory diseases, or any condition described herein. In some embodiments, a compound according to the invention, or for use according to the invention, may be provided in combination with one or more agents useful in the prevention or treatment of Alzheimer's disease. Examples of such agents may include, without limitation,

- acetylcholine esterase inhibitors (AChEIs) such as Aricept® (Donepezil), Exelon® (Rivastigmine), Razadyne® (Razadyne ER®, Reminyl®, Nivalin®, Galantamine), Cognex® (Tacrine), Dimebon, Huperzine A, Phenserine, Debio-9902 SR (ZT-1 SR), Zanzepil (TAK0147), ganstigmine, NP7557, etc.;
- NMDA receptor antagonists such as Namenda® (Axura®, Akatinol®, Ebixa®, Memantine), Dimebon, SGS-742, Neramexane, Debio-9902 SR (ZT-1 SR), etc.;
- gamma-secretase inhibitors and/or modulators such as Flurizan™ (Tarenflurbil, MPC-7869, R-flurbiprofen), LY450139, MK 0752, E2101, BMS-289948, BMS-299897, BMS-433796, LY-411575, GSI-136, etc.;
- beta-secretase inhibitors such as ATG-Z1, CTS-21166, MK-8931, etc.;
- alpha-secretase activators, such as NGX267, etc.;
- amyloid- β aggregation and/or fibrillization inhibitors such as Alzhemed™ (3APS, Tramiprosate, 3-amino-1-propanesulfonic acid), AL-108, AL-208, AZD-103, PBT2, Cereact, ONO-2506PO, PPI-558, etc.;
- tau aggregation inhibitors such as methylene blue, etc.;
- microtubule stabilizers such as AL-108, AL-208, paclitaxel, etc.;
- RAGE inhibitors, such as TTP488, etc.;
- 5-HT_{1a} receptor antagonists, such as Xaliproden, Lecoizotan, etc.;
- 5-HT₄ receptor antagonists, such as PRX-03410, etc.;
- kinase inhibitors such as SRN-003-556, amfurindamide, LiCl, AZD1080, NP031112, SAR-502250, etc.;
- humanized monoclonal anti-A β antibodies such as Bapineuzumab (AAB-001), LY2062430, RN1219, ACU-5A5, etc.;
- amyloid vaccines such as AN-1792, ACC-001, etc.;
- neuroprotective agents such as Cerebrolysin, AL-108, AL-208, Huperzine A, etc.;
- L-type calcium channel antagonists such as MEM-1003, etc.;
- nicotinic receptor antagonists, such as AZD3480, GTS-21, etc.;

- nicotinic receptor agonists, such as MEM 3454, Nefiracetam, etc.;
- peroxisome proliferator-activated receptor (PPAR) gamma agonists such as Avandia® (Rosglitazone), etc.;
- phosphodiesterase IV (PDE4) inhibitors, such as MK-0952, etc.;
- 5 • hormone replacement therapy such as estrogen (Premarin), etc.;
- monoamine oxidase (MAO) inhibitors such as NS2330, Rasagiline (Azilect®), TVP-1012, etc.;
- AMPA receptor modulators such as Ampalex (CX 516), etc.;
- nerve growth factors or NGF potentiators, such as CERE-110 (AAV-NGF), T-588, T-10 817MA, etc.;
- agents that prevent the release of luteinizing hormone (LH) by the pituitary gland, such as leuprolide (VP-4896), etc.;
- GABA receptor modulators such as AC-3933, NGD 97-1, CP-457920, etc.;
- benzodiazepine receptor inverse agonists such as SB-737552 (S-8510), AC-3933, etc.;
- 15 • noradrenaline-releasing agents such as T-588, T-817MA, etc.

[00132] It is to be understood that combination of compounds according to the invention, or for use according to the invention, with Alzheimer's agents is not limited to the examples described herein, but may include combination with any agent useful for the treatment of Alzheimer's disease. Combination of compounds according to the invention, or for use
20 according to the invention, and other Alzheimer's agents may be administered separately or in conjunction. The administration of one agent may be prior to, concurrent to, or subsequent to the administration of other agent(s).

[00133] In alternative embodiments, a compound may be supplied as a "prodrug" or protected forms, which release the compound after administration to a subject. For example,
25 a compound may carry a protective group which is split off by hydrolysis in body fluids, *e.g.*, in the bloodstream, thus releasing the active compound or is oxidized or reduced in body fluids to release the compound. Accordingly, a "prodrug" is meant to indicate a compound that may be converted under physiological conditions or by solvolysis to a biologically active compound of the invention. Thus, the term "prodrug" refers to a metabolic precursor of a
30 compound of the invention that is pharmaceutically acceptable. A prodrug may be inactive when administered to a subject in need thereof, but may be converted *in vivo* to an active compound of the invention. Prodrugs are typically rapidly transformed *in vivo* to yield the

parent compound of the invention, for example, by hydrolysis in blood. The prodrug compound often offers advantages of solubility, tissue compatibility or delayed release in a subject.

[00134] The term "prodrug" is also meant to include any covalently bonded carriers which release the active compound of the invention *in vivo* when such prodrug is administered to a subject. Prodrugs of a compound of the invention may be prepared by modifying functional groups present in the compound of the invention in such a way that the modifications are cleaved, either in routine manipulation or *in vivo*, to the parent compound of the invention. Prodrugs include compounds of the invention where a hydroxy, amino or mercapto group is bonded to any group that, when the prodrug of the compound of the invention is administered to a mammalian subject, cleaves to form a free hydroxy, free amino or free mercapto group, respectively. Examples of prodrugs include, but are not limited to, acetate, formate and benzoate derivatives of alcohol and acetamide, formamide, and benzamide derivatives of amine functional groups in one or more of the compounds of the invention and the like.

[00135] A discussion of prodrugs may be found in "Smith and Williams' Introduction to the Principles of Drug Design," H.J. Smith, Wright, Second Edition, London (1988); Bundgaard, H., *Design of Prodrugs* (1985), pp. 7-9, 21-24 (Elsevier, Amsterdam); The Practice of Medicinal Chemistry, Camille G. Wermuth et al., Ch 31, (Academic Press, 1996); A Textbook of Drug Design and Development, P. Krosgaard-Larson and H. Bundgaard, eds. Ch 5, pgs 113 191 (Harwood Academic Publishers, 1991); Higuchi, T., *et al.*, "Pro-drugs as Novel Delivery Systems," A.C.S. Symposium Series, Vol. 14; or in *Bioreversible Carriers in Drug Design*, ed. Edward B. Roche, American Pharmaceutical Association and Pergamon Press, 1987.

[00136] Suitable prodrug forms of one or more of the compounds of the invention may include embodiments in which one or more OH groups as set forth in Formula (I) may be protected as OC(O)R, where R may be optionally substituted alkyl, alkenyl, alkynyl, aryl, or heteroaryl. In these cases the ester groups may be hydrolyzed *in vivo* (e.g. in bodily fluids), liberating the OH groups and releasing the active compounds. Preferred prodrug embodiments of the invention may include compounds of Formula (I) where one or more OH groups may be protected with acetate, for example as OC(O)CH₃.

[00137] Compounds according to the invention, or for use according to the invention, may be provided alone or in combination with other compounds in the presence of a liposome, an adjuvant, or any pharmaceutically acceptable carrier, diluent or excipient, in a form suitable

for administration to a subject such as a mammal, for example, humans, cattle, sheep, etc. If desired, treatment with a compound according to the invention may be combined with more traditional and existing therapies for the therapeutic indications described herein.

Compounds according to the invention may be provided chronically or intermittently.

- 5 “Chronic” administration refers to administration of the compound(s) in a continuous mode as opposed to an acute mode, so as to maintain the initial therapeutic effect (activity) for an extended period of time. “Intermittent” administration is treatment that is not consecutively done without interruption, but rather is cyclic in nature. The terms “administration,” “administrable,” or “administering” as used herein should be understood to mean providing a
10 compound of the invention to the subject in need of treatment.

- [00138] “Pharmaceutically acceptable carrier, diluent or excipient” may include, without limitation, any adjuvant, carrier, excipient, glidant, sweetening agent, diluent, preservative, dye/colorant, flavor enhancer, surfactant, wetting agent, dispersing agent, suspending agent, stabilizer, isotonic agent, solvent, or emulsifier that has been approved, for example, by the
15 United States Food and Drug Administration or other governmental agency as being acceptable for use in humans or domestic animals.

- [00139] A compound of the present invention may be administered in the form of a pharmaceutically acceptable salt. In such cases, pharmaceutical compositions in accordance with this invention may comprise a salt of such a compound, preferably a physiologically
20 acceptable salt, which are known in the art. In some embodiments, the term “pharmaceutically acceptable salt” as used herein means an active ingredient comprising compounds of Formula I used in the form of a salt thereof, particularly where the salt form confers on the active ingredient improved pharmacokinetic properties as compared to the free form of the active ingredient or other previously disclosed salt form.

- 25 [00140] A “pharmaceutically acceptable salt” may include both acid and base addition salts. A “pharmaceutically acceptable acid addition salt” refers to those salts which retain the biological effectiveness and properties of the free bases, which are not biologically or otherwise undesirable, and which may be formed with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid and the like, and organic
30 acids such as acetic acid, trifluoroacetic acid, propionic acid, glycolic acid, pyruvic acid, oxalic acid, maleic acid, malonic acid, succinic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, p-toluenesulfonic acid, salicylic acid, and the like.

[00141] A “pharmaceutically acceptable base addition salt” refers to those salts which may retain the biological effectiveness and properties of the free acids, which may not be biologically or otherwise undesirable. These salts may be prepared from addition of an inorganic base or an organic base to the free acid. Salts derived from inorganic bases may include, but are not limited to, the sodium, potassium, lithium, ammonium, calcium, magnesium, iron, zinc, copper, manganese, aluminum salts and the like. Preferred inorganic salts may be the ammonium, sodium, potassium, calcium, and magnesium salts. Salts derived from organic bases may include, but are not limited to, salts of primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines and basic ion exchange resins, such as isopropylamine, trimethylamine, diethylamine, triethylamine, tripropylamine, ethanolamine, 2-dimethylaminoethanol, 2-diethylaminoethanol, dicyclohexylamine, lysine, arginine, histidine, caffeine, procaine, hydrabamine, choline, betaine, ethylenediamine, glucosamine, methylglucamine, theobromine, purines, piperazine, piperidine, N-ethylpiperidine, polyamine resins and the like. Particularly preferred organic bases may be isopropylamine, diethylamine, ethanolamine, trimethylamine, dicyclohexylamine, choline and caffeine.

[00142] Thus, the term “pharmaceutically acceptable salt” encompasses all acceptable salts including but not limited to acetate, lactobionate, benzenesulfonate, laurate, benzoate, malate, bicarbonate, maleate, bisulfate, mandelate, bitartrate, mesylate, borate, methylbromide, bromide, methylnitrite, calcium edetate, methylsulfate, camsylate, mucate, carbonate, napsylate, chloride, nitrate, clavulanate, N-methylglucamine, citrate, ammonium salt, dihydrochloride, oleate, edetate, oxalate, edisylate, pamoate (embonate), estolate, palmitate, esylate, pantothenate, fumarate, phosphate/diphosphate, gluceptate, polygalacturonate, gluconate, salicylate, glutame, stearate, glycolylarsanilate, sulfate, hexylresorcinate, subacetate, hydradamine, succinate, hydrobromide, tannate, hydrochloride, tartrate, hydroxynaphthoate, teoate, iodide, tosylate, isothionate, triethiodide, lactate, panoate, valerate, and the like.

[00143] Pharmaceutically acceptable salts of a compound of the present invention may be used as a dosage for modifying solubility or hydrolysis characteristics, or may be used in sustained release or prodrug formulations. Also, pharmaceutically acceptable salts of a compound of this invention may include those formed from cations such as sodium, potassium, aluminum, calcium, lithium, magnesium, zinc, and from bases such as ammonia, ethylenediamine, N-methyl-glutamine, lysine, arginine, ornithine, choline,

N,N'-dibenzylethylene-diamine, chloroprocaine, diethanolamine, procaine, N-benzylphenethyl-amine, diethylamine, piperazine, tris(hydroxymethyl)aminomethane, and tetramethylammonium hydroxide.

5 [00144] Pharmaceutical formulations may typically include one or more carriers acceptable for the mode of administration of the preparation, be it by injection, inhalation, topical administration, lavage, or other modes suitable for the selected treatment. Suitable carriers may be those known in the art for use in such modes of administration.

[00145] Suitable pharmaceutical compositions may be formulated by means known in the art and their mode of administration and dose determined by the skilled practitioner. For 10 parenteral administration, a compound may be dissolved in sterile water or saline or a pharmaceutically acceptable vehicle used for administration of non-water soluble compounds such as those used for vitamin K. For enteral administration, the compound may be administered in a tablet, capsule or dissolved in liquid form. The table or capsule may be enteric coated, or in a formulation for sustained release. Many suitable formulations are 15 known, including, polymeric or protein microparticles encapsulating a compound to be released, ointments, gels, hydrogels, or solutions which can be used topically or locally to administer a compound. A sustained release patch or implant may be employed to provide release over a prolonged period of time. Many techniques known to skilled practitioners are described in *Remington: the Science & Practice of Pharmacy* by Alfonso Gennaro, 20th ed., 20 Williams & Wilkins, (2000). Formulations for parenteral administration may, for example, contain excipients, polyalkylene glycols such as polyethylene glycol, oils of vegetable origin, or hydrogenated naphthalenes. Biocompatible, biodegradable lactide polymer, lactide/glycolide copolymer, or polyoxyethylene-polyoxypropylene copolymers may be used to control the release of a compound. Other potentially useful parenteral delivery systems for 25 modulatory compounds may include ethylene-vinyl acetate copolymer particles, osmotic pumps, implantable infusion systems, and liposomes. Formulations for inhalation may contain excipients, for example, lactose, or may be aqueous solutions containing, for example, polyoxyethylene-9-lauryl ether, glycocholate and deoxycholate, or may be oily solutions for administration in the form of nasal drops, or as a gel.

30 [00146] A compound or a pharmaceutical composition according to the present invention may be administered by oral or non-oral, e.g., intramuscular, intraperitoneal, intravenous, intracisternal injection or infusion, subcutaneous injection, transdermal or transmucosal routes. In some embodiments, a compound or pharmaceutical composition in accordance

with this invention or for use in this invention may be administered by means of a medical device or appliance such as an implant, graft, prosthesis, stent, etc. Implants may be devised which are intended to contain and release such compounds or compositions. An example would be an implant made of a polymeric material adapted to release the compound over a period of time. A compound may be administered alone or as a mixture with a pharmaceutically acceptable carrier *e.g.*, as solid formulations such as tablets, capsules, granules, powders, etc.; liquid formulations such as syrups, injections, etc.; injections, drops, suppositories, pessaries. In some embodiments, compounds or pharmaceutical compositions in accordance with this invention or for use in this invention may be administered by inhalation spray, nasal, vaginal, rectal, sublingual, or topical routes and may be formulated, alone or together, in suitable dosage unit formulations containing conventional non-toxic pharmaceutically acceptable carriers, adjuvants and vehicles appropriate for each route of administration.

[00147] A compound of the invention may be used to treat animals, including mice, rats, horses, cattle, sheep, dogs, cats, and monkeys. However, a compound of the invention may also be used in other organisms, such as avian species (*e.g.*, chickens). One or more of the compounds of the invention may also be effective for use in humans. The term “subject” or alternatively referred to herein as “patient” is intended to be referred to an animal, preferably a mammal, most preferably a human, who has been the object of treatment, observation or experiment. However, one or more of the compounds, methods and pharmaceutical compositions of the present invention may be used in the treatment of animals. Accordingly, as used herein, a “subject” may be a human, non-human primate, rat, mouse, cow, horse, pig, sheep, goat, dog, cat, etc. The subject may be suspected of having or at risk for having a condition that may require modulation of O-GlcNAcase activity.

[00148] An “effective amount” of a compound according to the invention may include a therapeutically effective amount or a prophylactically effective amount. A “therapeutically effective amount” refers to an amount effective, at dosages and for periods of time necessary, to achieve the desired therapeutic result, such as inhibition of an O-GlcNAcase, elevation of O-GlcNAc levels, inhibition of tau phosphorylation, or any condition described herein. A therapeutically effective amount of a compound may vary according to factors such as the disease state, age, sex, and weight of the individual, and the ability of the compound to elicit a desired response in the individual. Dosage regimens may be adjusted to provide the optimum therapeutic response. A therapeutically effective amount may also be one in which

any toxic or detrimental effects of the compound are outweighed by the therapeutically beneficial effects. A “prophylactically effective amount” may refer to an amount effective, at dosages and for periods of time necessary, to achieve the desired prophylactic result, such as inhibition of an O-GlcNAcase, elevation of O-GlcNAc levels, inhibition of tau phosphorylation, or any condition described herein. Typically, a prophylactic dose may be used in subjects prior to or at an earlier stage of disease, so that a prophylactically effective amount may be less than a therapeutically effective amount. A suitable range for therapeutically or prophylactically effective amounts of a compound may be any integer from 0.1 nM - 0.1 M, 0.1 nM - 0.05 M, 0.05 nM - 15 μM or 0.01 nM - 10 μM.

5
10 [00149] In alternative embodiments, in the treatment or prevention of conditions which may require modulation of O-GlcNAcase activity, an appropriate dosage level may generally be about 0.01 to 500 mg per kg subject body weight per day, and may be administered in single or multiple doses. In some embodiments, the dosage level may be about 0.1 to about 250 mg/kg per day. It will be understood that the specific dose level and frequency of dosage for any particular patient may be varied and may depend upon a variety of factors including the activity of the specific compound used, the metabolic stability and length of action of that compound, the age, body weight, general health, sex, diet, mode and time of administration, rate of excretion, drug combination, the severity of the particular condition, and the patient undergoing therapy.

15
20 [00150] It is to be noted that dosage values may vary with the severity of the condition to be alleviated. For any particular subject, specific dosage regimens may be adjusted over time according to the individual need and the professional judgement of the person administering or supervising the administration of the compositions. Dosage ranges set forth herein are exemplary only and do not limit the dosage ranges that may be selected by medical practitioners. The amount of active compound(s) in the composition may vary according to factors such as the disease state, age, sex, and weight of the subject. Dosage regimens may be adjusted to provide the optimum therapeutic response. For example, a single bolus may be administered, several divided doses may be administered over time or the dose may be proportionally reduced or increased as indicated by the exigencies of the therapeutic situation. It may be advantageous to formulate parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. In general, compounds of the invention should be used without causing substantial toxicity, and as described herein, one or more of the compounds may exhibit a suitable safety profile for therapeutic use. Toxicity of a

25
30

compound of the invention may be determined using standard techniques, for example, by testing in cell cultures or experimental animals and determining the therapeutic index, i.e., the ratio between the LD50 (the dose lethal to 50% of the population) and the LD100 (the dose lethal to 100% of the population). In some circumstances however, such as in severe disease conditions, it may be necessary to administer substantial excesses of the compositions.

[00151] In the compounds of generic Formula (I), the atoms may exhibit their natural isotopic abundances, or one or more of the atoms may be artificially enriched in a particular isotope having the same atomic number, but an atomic mass or mass number different from the atomic mass or mass number predominantly found in nature. The present invention is meant to include all suitable isotopic variations of the compounds of generic Formula (I). For example, different isotopic forms of hydrogen (H) include protium (^1H), deuterium (^2H) and tritium (^3H). Protium is the predominant hydrogen isotope found in nature. Enriching for deuterium may afford certain therapeutic advantages, such as increasing *in vivo* half-life or reducing dosage requirements, or may provide a compound useful as a standard for characterization of biological samples. Isotopically-enriched compounds within generic Formula (I) may be prepared without undue experimentation by conventional techniques well known to those skilled in the art or by processes analogous to those described in the Schemes and Examples herein using appropriate isotopically-enriched reagents and/or intermediates.

Other Uses and Assays

[00152] A compound of Formula (I) may be used in screening assays for compounds which modulate the activity of glycosidase enzymes, preferably the O-GlcNAcase enzyme. The ability of a test compound to inhibit O-GlcNAcase-dependent cleavage of O-GlcNAc from a model substrate may be measured using any assays, as described herein or known to one of ordinary skill in the art. For example, a fluorescence or UV-based assay known in the art may be used. A “test compound” may be any naturally-occurring or artificially-derived chemical compound. Test compounds may include, without limitation, peptides, polypeptides, synthesised organic molecules, naturally occurring organic molecules, and nucleic acid molecules. A test compound may “compete” with a known compound such as a compound of Formula (I) by, for example, interfering with inhibition of O-GlcNAcase-dependent cleavage of O-GlcNAc or by interfering with any biological response induced by a compound of Formula (I).

[00153] Generally, a test compound may exhibit any value between 10% and 200%, or over 500%, modulation when compared to a compound of Formula (I) or other reference compound. For example, a test compound may exhibit at least any positive or negative integer from 10% to 200% modulation, or at least any positive or negative integer from 30% to 150% modulation, or at least any positive or negative integer from 60% to 100% modulation, or any positive or negative integer over 100% modulation. A compound that is a negative modulator may in general decrease modulation relative to a known compound, while a compound that is a positive modulator may in general increase modulation relative to a known compound.

[00154] In general, test compounds may be identified from large libraries of both natural products or synthetic (or semi-synthetic) extracts or chemical libraries according to methods known in the art. Those skilled in the field of drug discovery and development will understand that the precise source of test extracts or compounds is not critical to the method(s) of the invention. Accordingly, virtually any number of chemical extracts or compounds may be screened using the exemplary methods described herein. Examples of such extracts or compounds may include, but are not limited to, plant-, fungal-, prokaryotic- or animal-based extracts, fermentation broths, and synthetic compounds, as well as modification of existing compounds. Numerous methods are also available for generating random or directed synthesis (e.g., semi-synthesis or total synthesis) of any number of chemical compounds, that may include, without limitation, saccharide-, lipid-, peptide-, and nucleic acid-based compounds. Synthetic compound libraries are commercially available. Alternatively, libraries of natural compounds in the form of bacterial, fungal, plant, and animal extracts are commercially available from a number of sources, including Biotics (Sussex, UK), Xenova (Slough, UK), Harbor Branch Oceanographic Institute (Ft. Pierce, FL, USA), and PharmaMar, MA, USA. In addition, natural and synthetically produced libraries may be produced, if desired, according to methods known in the art, e.g., by standard extraction and fractionation methods. Furthermore, if desired, any library or compound may be readily modified using standard chemical, physical, or biochemical methods.

[00155] When a crude extract is found to modulate inhibition of O-GlcNAcase-dependent cleavage of O-GlcNAc, or any biological response induced by a compound of Formula (I), further fractionation of the positive lead extract may be necessary to isolate chemical constituents responsible for the observed effect. Thus, the goal of the extraction, fractionation, and purification process is the careful characterization and identification of a

chemical entity within the crude extract having O-GlcNAcase- inhibitory activities. The same assays described herein for the detection of activities in mixtures of compounds may be used to purify the active component and to test derivatives thereof. Methods of fractionation and purification of such heterogeneous extracts are known in the art. If desired, compounds shown to be useful agents for treatment may be chemically modified according to methods known in the art. Compounds identified as being of therapeutic, prophylactic, diagnostic, or other value may be subsequently analyzed using a suitable animal model, as described herein on known in the art.

5

10

15

[00156] In some embodiments, one or more of the compounds may be useful in the development of animal models for studying diseases or disorders that may be related to deficiencies in O-GlcNAcase, over-expression of O-GlcNAcase, accumulation of O-GlcNAc, depletion of O-GlcNAc, and for studying treatment of diseases and disorders that may be related to deficiency or over-expression of O-GlcNAcase, or accumulation or depletion of O-GlcNAc. Such diseases and disorders may include neurodegenerative diseases, including Alzheimer's disease, and cancer.

[00157] Various alternative embodiments and examples of the invention are described herein. These embodiments and examples are illustrative and should not be construed as limiting the scope of the invention.

20

EXAMPLES

[00158] The following examples are intended to illustrate embodiments of the invention and are not intended to be construed in a limiting manner.

Abbreviations


ABCN	= 1,1'-azobis(cyclohexane-carbonitrile)
25 AcCl	= acetyl chloride
AIBN	= azobisisobutyronitrile
BCl ₃	= boron trichloride
BnBr	= benzyl bromide
Bu ₄ NI	= tetra- <i>n</i> -butylammonium iodide
30 Boc ₂ O	= di- <i>tert</i> -butyl dicarbonate

	BzCl	= benzoyl chloride
	DAST	= diethylaminosulfur trifluoride
	DCM	= dichloromethane
	DIAD	= diisopropyl azodicarboxylate
5	DIPEA	= diisopropylethylamine
	DMAP	= 4-dimethylaminopyridine
	DMF	= <i>N,N</i> -dimethylformamide
	DMP	= Dess-Martin periodinane
	DMSO	= dimethyl sulfoxide
10	EDCI	= 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide
	Et ₃ N	= triethylamine
	Et ₂ O	= diethyl ether
	HATU	= (<i>O</i> -(7-azabenzotriazol-1-yl)- <i>N,N,N',N'</i> -tetramethyluronium hexafluorophosphate)
15	PMB	= pentamethylbenzene
	TBDMSCl	= <i>tert</i> -butyldimethylsilyl chloride
	TBAF	= tetra- <i>n</i> -butylammonium fluoride
	TMSCF ₃	= (trifluoromethyl)trimethylsilane
	TFA	= 2,2,2-trifluoroacetic acid
20	THF	= tetrahydrofuran
	thio-CDI	= 1,1'-thiocarbonyl diimidazole


General Procedures and Intermediates

[00159] The compounds of the invention are synthesized according to standard schemes and procedures, as indicated for example, in Schemes 1 to 23, as appropriate. Intermediate **A** (Scheme 1) may be prepared as described in, for example, *Tetrahedron: Asymmetry* **1998**, 3505. Intermediate **B** (Scheme 16) may be prepared as described in, for example, *Bioorg. Med. Chem. Lett.* **2008**, 8273. It is to be understood that any suitable scheme within the

knowledge of one skilled in the art may be used to synthesize one or more of the compounds described herein.




Scheme 2




5


Scheme 3




Scheme 4




Scheme 5




Scheme 6




Scheme 7




Scheme 8




Scheme 9




Scheme 10




Scheme 11




Scheme 12




Scheme 13



Scheme 14




Scheme 15




5


Scheme 16




Scheme 17




Scheme 18




Scheme 19




Scheme 20




Scheme 21




Scheme 22



Scheme 23



(2S,3R,4S,5R,6R)-3,4,5-trihydroxy-6-(hydroxymethyl)-N-methylpiperidine-2-carboxamide (hydrochloride salt)



5

[00160] (2S,3R,4S,5R,6R)-3,4,5-trihydroxy-6-(hydroxymethyl)-N-methylpiperidine-2-carboxamide was prepared following known procedures (Shilvock et al. *Tetrahedron: Asymmetry* **1998**, 9, 3505). To a solution of (3S,4S,5S,6R)-6-(hydroxymethyl)-tetrahydro-2H-pyran-2,3,4,5-tetraol (D-Mannose) (100 g, 0.56 mol) in acetone (2.2 L) was added iodine (28.2 g, 0.11 mmol). After stirring for 12 hours at room temperature, the reaction solution was poured into saturated aqueous sodium thiosulphate (2.2 L) and neutralized by sodium bicarbonate. The resulting mixture was extracted with DCM (3x1 L), and the combined organic layer was washed with brine (2x700 mL), dried over anhydrous sodium sulfate and concentrated under reduced pressure to afford (3aS,6R,6aS)-6-((S)-2,2-dimethyl-1,3-

10

dioxolan-4-yl)-2,2-dimethyl-tetrahydrofuro[3,4-d][1,3]dioxol-4-ol (190 g, 88%) as a yellow solid, which was used in next step without further purification; (ES, m/z) $[M+H]^+$ 261.1; 1H NMR (300 MHz, $CDCl_3$) δ 4.84 - 4.80 (m, 1H), 4.78 - 4.75 (m, 1H), 4.44 - 4.37 (m, 1H), 4.21 - 4.17 (m, 1H), 4.19 - 4.02 (m, 3H), 1.48 (s, 3H), 1.45 (s, 3H), 1.40 (s, 3H), 1.37 (s, 3H).

5 [00161] A solution of the above material (10 g, 38.4 mmol) and NaCN (1.7 g, 34.7 mmol) in water (300 mL) was stirred for 12 hours at 50 °C. Then the resulting solution was extracted with DCM (2x100 mL) (to remove the organic impurities). The pH value of the aqueous phase was adjusted to 2 - 3 with 2N sulfuric acid and extracted with EtOAc (5x200 mL). The combined organic layer was dried over anhydrous sodium sulfate and concentrated under
10 reduced pressure to give a residue, which was dissolved in DCM (100 mL) and followed by the addition of DCC (7.9 g, 38.4 mmol). After additional 12 hours at room temperature, the resulting mixture was filtrated and the filtrate was concentrated under reduced pressure. The crude residue was purified by a silica gel column, eluted with 10% ~ 35% EtOAc in petroleum ether to afford (3aR,4R,7S,7aR)-4-((S)-2,2-dimethyl-1,3-dioxolan-4-yl)-7-
15 hydroxy-2,2-dimethyl-tetrahydro-[1,3] dioxolo[4,5-c]pyran-6-one (2.6 g, 24%) (R_f 0.4, EtOAc: hexane, 2:1) as a yellow solid; (ES, m/z) $[M+H]^+$ 289.1; 1H NMR (300 MHz, $CDCl_3$) δ 4.85 - 4.83 (m, 1H), 4.71 - 4.69 (m, 1H), 4.44 - 4.37 (m, 2H), 4.17 - 4.13 (m, 1H), 4.09 - 4.04 (m, 2H), 1.46 (s, 3H), 1.45 (s, 3H), 1.36 (s, 3H), 1.34 (s, 3H).

[00162] A solution of the above material (2 g, 6.9 mmol) and pyridine (1.1 g, 13.8 mmol) in
20 DCM (30 mL) was treated with Tf_2O (3 g, 10.3 mmol) for 15 min at -30 °C. Then the reaction was quenched by water (20 mL), extracted with DCM (3x20 mL). The combined organic layer was washed with Brine (2x20 mL), dried over anhydrous sodium sulfate and concentrated under reduced pressure to give a residue, which was dissolved in DMF (20 mL) and followed by the addition of NaN_3 (1.2 g, 18.5 mmol). After additional 12 hours at room
25 temperature, the reaction was quenched by water (20 mL) and extracted with DCM (3x30 mL). The combined organic layer was washed with brine (3x30 mL), dried over anhydrous sodium sulfate and concentrated under reduced pressure give a residue, which was purified by a silica gel column, eluted with 10% ~ 30% EtOAc in petroleum ether to afford
(3aR,4R,7S,7aR)-7-azido-4-((S)-2,2-dimethyl-1,3-dioxolan-4-yl)-2,2-dimethyl-tetrahydro-
30 [1,3] dioxolo[4,5-c]pyran-6-one (1.6 g, 74%) as a white solid; (ES, m/z) $[M+H]^+$ 314.2; 1H NMR (300 MHz, $CDCl_3$) δ 4.61 - 4.59 (m, 1H), 4.54 - 4.51 (m, 1H), 4.49 - 4.46 (m, 1H), 4.40 - 4.31 (m, 2H), 4.18 - 4.09 (m, 2H), 1.46 (s, 3H), 1.43 (s, 3H), 1.39 (s, 3H), 1.36 (s, 3H).

[00163] A solution of the above material (4 g, 12.7 mmol) in HOAc (40 mL) and water (10 mL) was stirred for 3.5 hours at 50 °C. Volatiles were distilled out under reduced pressure to give a residue, which was purified by a silica gel column, eluted with 1% ~ 3% MeOH in DCM to afford (3aR,4R,7S,7aR)-7-azido-4-((S)-1,2-dihydroxyethyl)-2,2-dimethyl-

5 tetrahydro-[1,3]dioxolo[4,5-c]pyran-6-one (3.1 g, 89%) as a white solid; (ES, *m/z*) [M+H]⁺ 274.2; ¹H NMR (300 MHz, CDCl₃) δ 4.93 - 4.79 (m, 2H), 4.24 - 4.21 (m, 1H), 4.12 - 4.07 (m, 1H), 3.90 - 3.75 (m, 3H), 1.47 (s, 3H), 1.40 (s, 3H).

[00164] To a solution of the above material (2.9 g, 10.6 mmol), Et₃N (1.7 g, 16.8 mmol) and DMAP (134 mg, 1.1 mmol) in DCM (30 mL) was added TBSCl (2 g, 13.4 mmol) in portions.

10 After stirring for 12 hours at room temperature, the reaction was quenched by MeOH (30 mL). Volatiles were distilled out under reduced pressure to give a residue, which was purified by a silica gel column, eluted with 5% ~ 20% EtOAc in petroleum ether to afford (3aR,4R,7S,7aR)-7-azido-4-((S)-2-(tert-butyldimethylsilyloxy)-1-hydroxyethyl)-2,2-dimethyl-tetra hydro-[1,3]dioxolo[4,5-c]pyran-6-one (2.8 g, 68%) as a white solid; (ES, *m/z*)

15 [M+H]⁺ 388.2; ¹H NMR (300 MHz, CDCl₃) δ 4.91 - 4.84 (m, 2H), 4.08 - 4.03 (m, 2H), 3.92 - 3.80 (m, 3H), 1.50 (s, 3H), 1.43 (s, 3H), 0.92 (s, 9H), 0.11 (s, 6H).

[00165] To a solution of the above material (2.5 g, 6.5 mmol) in DCM (40 mL) was added DMP (5.4 g, 12.7 mmol) at 0 °C. After stirring for 3 hours at room temperature, the reaction was quenched by saturated aqueous sodium thiosulphate (15 mL) and sodium bicarbonate (15

20 mL), extracted with DCM (3x30 mL). The combined organic layer was washed with Brine (2x30 mL), dried over anhydrous sodium sulfate and concentrated under reduced pressure to give a residue, which was purified by a silica gel column, eluted with 5% ~ 15% EtOAc in petroleum ether to afford (3aR,4S,7S,7aR)-7-azido-4-(2-(tert-butyldimethylsilyloxy)acetyl)-2,2-dimethyl-tetrahydro-[1,3] dioxolo[4,5-c]pyran-6-one (1.9 g, 76%) as a white solid; (ES, *m/z*) [M+H]⁺ 386.2; ¹H NMR (300 MHz, CDCl₃) δ 4.94 - 4.89 (m, 2H), 4.78 - 4.63 (m, 3H), 3.81 (d, *J* = 2.1 Hz, 1H), 1.50 (s, 3H), 1.37 (s, 3H), 0.95 (s, 9H), 0.12 (s, 6H).

[00166] A solution of the above material (2 g, 5.2 mmol) in anhydrous THF (30 mL) was treated with triethyl phosphite (1.72 g, 10.4 mmol) for 16 hours at room temperature under nitrogen atmosphere. Volatiles were distilled out under reduced pressure to give a residue,


30 which was purified by a silica gel column, eluted with 2% ~ 20% EtOAc in petroleum ether to afford (3aR,4S,7R,7aR)-6-(((tert-butyldimethylsilyl)oxy)methyl)-2,2-dimethyl-3a,4,7,7a-tetrahydro-7,4-(epoxymethano)[1,3]dioxolo[4,5-c]pyridin-9-one (1.2 g, 68%) as a light

yellow syrup; (ES, m/z) $[M+H]^+$ 342.1; 1H NMR (400 MHz, $CDCl_3$) δ 5.51 (d, $J = 4.0$ Hz, 1H), 5.22 (d, $J = 2.8$ Hz, 1H), 4.75 - 4.65 (m, 2H), 4.57 - 4.44 (m, 2H), 1.51 (s, 3H), 1.37 (s, 3H), 0.97 (s, 9H), 0.12 (s, 6H).

[00167] A solution of the above material (500 mg, 1.47 mmol) in acetic acid (10 mL) was
5 treated with sodium cyanoborohydride (110 mg, 1.76 mmol) for 30 min at room temperature. Volatiles were distilled out under reduced pressure to give a residue, which was purified by a silica gel column, eluted with 5% ~ 20% EtOAc in petroleum ether to afford 7-O-tert-Butyldimethylsilyl-2,6-dideoxy-2,6-imino-3,4-O-isopropylidene-D-glycero-D-taloheptono-1,5-lactone (320 mg, 64%) as a light yellow syrup; (ES, m/z) $[M+H]^+$ 344.1; 1H NMR (300
10 MHz, $CDCl_3$) δ 4.73 (d, $J = 2.0$ Hz, 1H), 4.50 - 4.47 (m, 1H), 4.36 - 4.33 (m, 1H), 4.01 - 3.95 (m, 1H), 3.75 - 3.70 (m, 2H), 3.43 - 3.41 (m, 1H), 1.56 (s, 3H), 1.31 (s, 3H), 0.83 (s, 9H), 0.06 (s, 6H).

[00168] Methylamine (33% w/w solution in MeOH, 0.4 ml, 3.5 mmol) was added to a
15 solution of the above material (200 mg, 0.58 mmol) in dry THF (10 ml) and the mixture stirred at room temperature for 2 hours. Volatiles were distilled out under reduced pressure to give a residue, which was purified by a silica gel column with 10 ~ 40 % EtOAc in petroleum ether to afford (3aR,4S,6R,7R,7aS)-6-((tert-butyldimethylsilyloxy)methyl)-7-hydroxy-N,2,2-trimethyl-hexahydro-[1,3]dioxolo[4,5-c]pyridine-4-carboxamide (173 mg, 80%) as light yellow oil; (ES, m/z) $[M+H]^+$ 375.2; 1H NMR (400 MHz, $CDCl_3$) δ 7.37 (br,
20 1H), 4.98 - 4.96 (m, 1H), 4.03 - 3.95 (m, 2H), 3.85 - 3.72 (m, 1H), 3.69 - 3.67 (m, 2H), 2.87 (d, $J = 4.8$ Hz, 3H), 2.49 (br, 1H), 2.47 - 2.45 (m, 1H), 1.56 (s, 3H), 1.48 (s, 3H), 0.92 (s, 9H), 0.06 (s, 6H).

[00169] A solution of the above material (330 mg, 0.88 mmol) in MeOH (10 mL) was
25 treated with 4N solution of HCl (g) in MeOH (3 mL, 12 mmol) for 3 hours at room temperature. Volatiles were distilled out under reduced pressure to give a residue, which was dissolved in water (20 mL), extracted with DCM (2x10 mL) (to remove the organic impurities). The aqueous layer was concentrated and dried by lyophilization to afford (2S,3R,4S,5R,6R)-3,4,5-trihydroxy-6-(hydroxymethyl)-N-methylpiperidine-2-carboxamide hydrochloride (198 mg, 88%) as a white syrup; (ES, m/z) $[M+H]^+$ 221.2; 1H
30 NMR (300 MHz, D_2O) δ 4.31 - 4.29 (m, 1H), 4.20 - 4.18 (m, 1H), 3.90 - 3.81 (m, 3H), 3.69 - 3.54 (m, 2H), 2.68 (s, 3H).

Examples**Example 1****(2S,3R,4S,5R,6R)-3,4,5-trihydroxy-6-(hydroxymethyl)-N,1-dimethylpiperidine-2-carboxamide hydrochloride**

5

[00170] To a suspension of D-mannose (100.0 g, 0.56 mol) in dry acetone (2.2 L) was added iodine (28.6 g, 0.11 mol). The resulting mixture was stirred for 16 hours at ambient temperature to give a homogeneous solution. The reaction was then quenched by the addition of water (2 L) and neutralized by the addition of sodium bicarbonate (15 g, 0.18 mol). The resulting mixture was extracted with EtOAc (3 x 1 L). The combined organic layers was washed with brine (2 x 500 mL) and saturated aqueous solution of sodium sulfite (3 x 200 mL). After dried over anhydrous sodium sulfate, a filtration was performed and the filtrate was concentrated under reduced pressure to give (3*a*S,6*R*,6*a*S)-6-((*S*)-2,2-dimethyl-1,3-dioxolan-4-yl)-2,2-dimethyl-tetrahydrofuro[3,4-*d*][1,3]dioxol-4-ol as a colorless solid (130 g, 90%): ¹H NMR (300 MHz, CDCl₃) δ 5.38-5.01 (m, 0.2H), 4.83-4.80 (dd, *J* = 3.6 Hz, 6.0 Hz, 0.8H), 4.78-4.75 (dd, *J* = 3.3 Hz, 6.0 Hz, 0.2H), 4.65-4.61 (m, 1H), 4.55-4.52 (m, 0.2H), 4.47-4.35 (m, 1.2H), 4.11-4.08 (m, 0.8H), 4.07-4.03 (m, 2.4H), 3.49-3.45 (m, 0.2H), 1.48-1.25 (m, 12H); MS (ESI, *m/z*): 261.0 [M + 1]⁺.

[00171] (3*a*R,4*R*,7*S*,7*a*R)-4-((*S*)-2,2-dimethyl-1,3-dioxolan-4-yl)-7-hydroxy-2,2-dimethyl-tetrahydro-[1,3]dioxolo[4,5-*c*]pyran-6-one and (3*a*R,4*R*,7*R*,7*a*R)-4-((*S*)-2,2-dimethyl-1,3-dioxolan-4-yl)-7-hydroxy-2,2-dimethyl-tetrahydro-[1,3]dioxolo[4,5-*c*]pyran-6-one were prepared according to procedures described in *Tetrahedron: Asymmetry* **1991**, 883. To a suspension of (3*a*S,6*R*,6*a*S)-6-((*S*)-2,2-dimethyl-1,3-dioxolan-4-yl)-2,2-dimethyl-tetrahydrofuro[3,4-*d*][1,3]dioxol-4-ol (100 g, 0.38 mol) in water (800 mL) was added sodium cyanide (17 g, 0.35 mol). The resulting mixture was stirred for 16 hours at 50 °C. The solid gradually dissolved and the solution went brown. The resulting solution was cooled down to ambient temperature and extracted with DCM (3 x 500 mL). The DCM layers were discarded. Sodium chloride (58 g, 1 mol) was dissolved in the aqueous layer. EtOAc (300 mL) was added to it and the resulting mixture was stirred vigorously followed by the addition of dilute sulfuric acid (2 M) to adjust the pH value to 3 and the EtOAc layer was separated

30

out. The aqueous layer was extracted with EtOAc (3 x 100 mL). The combined organic layers was dried over anhydrous sodium sulfate and filtered. The filtrate was concentrated under reduced pressure to give the crude product (33 g), which was pure enough for next step. A sample (1 g) was further purified by silica gel column chromatography, eluted with 5% to 20% EtOAc in petroleum ether to give (3aR,4R,7R,7aR)-4-((S)-2,2-dimethyl-1,3-dioxolan-4-yl)-7-hydroxy-2,2-dimethyl-tetrahydro-[1,3]dioxolo[4,5-c]pyran-6-one as a colorless solid (285 mg, 29%): $^1\text{H NMR}$ (300 MHz, CDCl_3) δ 4.72 (dd, $J = 1.2$ Hz, 3.3 Hz, 1H), 4.67 (dd, $J = 1.5$ Hz, 5.7 Hz, 1H), 4.61 (dd, $J = 2.4$ Hz, 7.5 Hz, 1H), 4.41 (s, 1H), 4.40-4.36 (m, 1H), 4.16-4.08 (m, 2H), 1.46 (s, 3H), 1.41 (s, 3H), 1.39 (s, 3H), 1.37 (s, 3H); MS (ESI, m/z): 289.0 $[\text{M} + 1]^+$. And (3aR,4R,7S,7aR)-4-((S)-2,2-dimethyl-1,3-dioxolan-4-yl)-7-hydroxy-2,2-dimethyl-tetrahydro-[1,3]dioxolo[4,5-c]pyran-6-one as a colorless solid (660 mg, 66%): $^1\text{H NMR}$ (300 MHz, CDCl_3) δ 4.83 (dd, $J = 3.6$ Hz, 7.6 Hz, 1H), 4.68 (dd, $J = 1.6$ Hz, 7.6 Hz, 1H), 4.45-4.41 (m, 1H), 4.35 (d, $J = 3.6$ Hz, 1H), 4.16-4.01 (m, 3H), 1.45 (s, 3H), 1.44 (s, 3H), 1.38 (s, 3H), 1.34 (s, 3H); MS (ESI, m/z): 289.0 $[\text{M} + 1]^+$.

[00172] To a solution of the above mixture (30 g, 104 mmol) in DCM (500 mL) was added dry pyridine (25 g, 312 mmol) followed by the addition of trifluoromethanesulfonic anhydride (44 g, 156 mmol) over 10 min at -30 °C to -15 °C. After 30 min, the reaction was quenched by water (500 mL) and the organic layer was separated out. The aqueous layer was extracted with DCM (3 x 100 mL). The combined organic layer was dried over anhydrous sodium sulfate and filtered. The filtration was concentrated under reduced pressure to give crude triflate, which was dissolved into DMF (150 mL) followed by the addition of sodium azide (10 g, 156 mmol) at ambient temperature. After stirring for 4 hours, the solvent was removed under reduced pressure and the residue was partitioned between DCM (150 mL) and water (60 mL). The organic layer was separated out and washed with brine (3 x 50 mL). The organic layer was dried over anhydrous sodium sulfate and filtered. The filtrate was concentrated under reduced pressure and the residue was purified by silica gel column chromatography, eluted with 10% to 20% EtOAc in petroleum ether to give (3aR,4R,7S,7aR)-7-azido-4-((S)-2,2-dimethyl-1,3-dioxolan-4-yl)-2,2-dimethyl-tetrahydro-[1,3]dioxolo[4,5-c]pyran-6-one as a colorless solid (24.3 g, 76%): $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 4.87 (dd, $J = 3.2$ Hz, 7.6 Hz, 1H), 4.70 (dd, $J = 1.6$ Hz, 8.0 Hz, 1H), 4.43-4.38 (m, 1H), 4.17-4.13 (dd, $J = 6.0$ Hz, 9.2 Hz, 1H), 4.09-4.06 (dd, $J = 3.6$ Hz, 9.2 Hz, 1H), 3.98-3.95 (dd, $J = 1.2$ Hz, 8.4 Hz, 1H), 3.83 (d, $J = 3.2$ Hz, 1H), 1.47 (s, 3H), 1.44 (s, 3H), 1.40 (s, 3H), 1.38 (s, 3H); MS (ESI, m/z): 314.0 $[\text{M} + 1]^+$.

[00173] A solution of (3aR,4R,7S,7aR)-7-azido-4-((S)-2,2-dimethyl-1,3-dioxolan-4-yl)-2,2-dimethyl-tetrahydro-[1,3]dioxolo[4,5-c]pyran-6-one (24 g, 77 mmol) in 80% acetic acid (200 ml) was heated at 50 °C for 4 hours. The solvent was removed under reduced pressure and the residue was purified by silica gel column chromatography, eluted with 20% to 80% EtOAc in petroleum ether to give (3aR,4R,7S,7aR)-7-azido-4-((S)-1,2-dihydroxyethyl)-2,2-dimethyl-tetrahydro-[1,3]dioxolo[4,5-c]pyran-6-one as a colorless solid (16.5 g, 85%): ¹H NMR (400 MHz, CDCl₃) δ 4.89 (dd, *J* = 3.2 Hz, 8.0 Hz, 1H), 4.82 (dd, *J* = 1.6 Hz, 8.0 Hz, 1H), 4.22 (dd, *J* = 1.2 Hz, 8.4 Hz, 1H), 4.10-4.06 (m, 1H), 3.95-3.85 (m, 3H), 1.49 (s, 3H), 1.42 (s, 3H); MS (ESI, *m/z*): 274.0 [M + 1]⁺.

10 [00174] To a solution of (3aR,4R,7S,7aR)-7-azido-4-((S)-1,2-dihydroxyethyl)-2,2-dimethyl-tetrahydro-[1,3]dioxolo[4,5-c]pyran-6-one (15 g, 54.9 mmol) in DCM (150 mL) was added triethylamine (8.3 g, 82.5 mmol) and N,N-dimethylpyridin-4-amine (0.7 g, 5.7 mmol) followed by the addition of tert-butyldimethylsilyl chloride (9.9 g, 66 mmol) in several portions at ambient temperature under a nitrogen atmosphere. After additional 16 hours, the reaction was quenched by the addition of MeOH (5 mL) and the resulting mixture was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with 5% to 15% EtOAc in petroleum ether to give (3aR,4R,7S,7aR)-7-azido-4-((S)-2-(tert-butyldimethylsilyloxy)-1-hydroxyethyl)-2,2-dimethyl-tetrahydro-[1,3]dioxolo[4,5-c]pyran-6-one as a colorless solid (12.2 g, 58%): ¹H NMR (400 MHz, CDCl₃) δ 4.85 (m, 2H), 4.03 (m, 2H), 3.82 (m, 3H), 2.67 (brs, 1H), 1.46 (s, 3H), 1.39 (s, 3H), 0.89 (s, 9H), 0.08 (s, 6H); MS (ESI, *m/z*): 388.0 [M + 1]⁺.

25 [00175] A solution of (3aR,4R,7S,7aR)-7-azido-4-((S)-2-(tert-butyldimethylsilyloxy)-1-hydroxyethyl)-2,2-dimethyl-tetrahydro-[1,3]dioxolo[4,5-c]pyran-6-one (20 g, 51.6 mmol) in DCM (150 mL) was treated with Dess-Martin periodinane (43.7 g, 103.2 mmol) for 1 hour at ambient temperature. The reaction was quenched by the addition of saturated aqueous sodium dicarbonate (100 mL) and saturated aqueous sodium thiosulphate (100 mL). The organic layer was separated and the aqueous layer was extracted with DCM (3 x 50 mL). The combined organic layers was dried over anhydrous sodium sulfate and filtered. The filtrate was concentrated under reduced pressure and the residue was purified by silica gel column chromatography, eluted with 5% to 20% EtOAc in petroleum ether to give (3aR,4S,7S,7aR)-7-azido-4-(2-(tert-butyldimethylsilyloxy)acetyl)-2,2-dimethyl-tetrahydro-[1,3]dioxolo[4,5-c]pyran-6-one as a colorless solid (15 g, 75%): ¹H NMR (400 MHz, CDCl₃) δ 4.94-4.88 (m,

2H), 4.77-4.63 (m, 3H), 3.81 (d, $J = 2.8$ Hz, 1H), 1.49 (s, 3H), 1.36 (s, 3H), 0.95 (s, 9H), 0.13 (s, 3H), 0.11 (s, 3H); MS (ESI, m/z): 386.0 $[M + 1]^+$.

[00176] A solution of (3aR,4S,7S,7aR)-7-azido-4-(2-(tert-butyl dimethylsilyloxy)acetyl)-2,2-dimethyl-tetrahydro-[1,3]dioxolo[4,5-c]pyran-6-one (10 g, 25.9 mmol) in dry THF (120 mL) was treated with triethyl phosphite (8.6 g, 51.8 mmol) for 16 hours at ambient temperature under a nitrogen atmosphere. Solvent was removed under reduced pressure to give crude 11- $\{[(\text{tert-butyl dimethylsilyl})\text{oxy}]\text{methyl}\}$ -4,4-dimethyl-3,5,8-trioxa-10-azatricyclo[5.2.2.0 $\{2,6\}$]undec-10-en-9-one as a colorless oil (9.3 g): MS (ESI, m/z): 386.0 $[M + 1]^+$. This material was used in the next step without further purification

[00177] To a solution of the above crude 11- $\{[(\text{tert-butyl dimethylsilyl})\text{oxy}]\text{methyl}\}$ -4,4-dimethyl-3,5,8-trioxa-10-azatricyclo[5.2.2.0 $\{2,6\}$]undec-10-en-9-one (9.3 g) in acetic acid (80 mL) was added sodium cyanoborohydride (2.4 g, 38.8 mmol) at ambient temperature. After additional 2 hours, solvent was removed under reduced pressure and the residue was purified by silica gel column chromatography, eluted with 10% to 30% EtOAc in petroleum ether to give 11- $\{[(\text{tert-butyl dimethylsilyl})\text{oxy}]\text{methyl}\}$ -4,4-dimethyl-3,5,8-trioxa-10-azatricyclo[5.2.2.0 $\{2,6\}$]undecan-9-one as a colorless oil (5.4 g, 61%): ^1H NMR (300 MHz, CDCl_3) δ 4.74-4.73 (m, 1H), 4.49-4.47 (m, 1H), 4.36-4.33 (m, 1H), 4.00-3.95 (m, 1H), 3.75-3.70 (m, 2H), 3.43-3.41 (m, 1H), 2.01 (brs, 1H), 1.56 (s, 3H), 1.31 (s, 3H), 0.83 (s, 9H), 0.07 (s, 6 H); MS (ESI, m/z): 344.0 $[M + 1]^+$.

[00178] A solution of 11- $\{[(\text{tert-butyl dimethylsilyl})\text{oxy}]\text{methyl}\}$ -4,4-dimethyl-3,5,8-trioxa-10-azatricyclo[5.2.2.0 $\{2,6\}$]undecan-9-one (3.2 g, 9.3 mmol) in THF (20 mL) was treated with 2 *M* solution of methanamine in THF (10 mL, 20 mmol) for 1 hour at ambient temperature. Solvent was removed under reduced pressure and the residue was purified by silica gel column chromatography, eluted with 10% to 50% EtOAc to give

(3aR,4S,6R,7R,7aS)-6-((tert-butyl dimethylsilyloxy)methyl)-7-hydroxy-N,2,2-trimethyl-hexahydro-[1,3]dioxolo[4,5-c]pyridine-4-carboxamide as a colorless solid (2.7 g, 77%): ^1H NMR (400 MHz, CDCl_3) δ 7.37 (brs, 1H), 4.97-4.95 (m, 1H), 3.99-3.94 (m, 2H), 3.84 (d, $J = 1.6$ Hz, 1H), 3.74-3.66 (m, 2H), 2.86 (d, $J = 4.8$ Hz, 3H), 2.65 (brs, 1H), 2.50-2.47 (m, 1H), 1.53 (s, 3H), 1.43 (s, 3H), 0.92 (s, 9H), 0.12 (s, 6H); MS (ESI, m/z): 375.0 $[M + 1]^+$.

[00179] To a solution of (3aR,4S,6R,7R,7aS)-6-((tert-butyl dimethylsilyloxy)methyl)-7-hydroxy-N,2,2-trimethyl-hexahydro-[1,3]dioxolo[4,5-c]pyridine-4-carboxamide (0.37 g, 1 mmol) in DMF (20 mL) was added potassium carbonate (0.28 g, 2 mmol) and iodomethane


(0.21 g, 1.5 mmol) at ambient temperature. After stirring for 1 h, the reaction was quenched by water (100 mL) and extracted with DCM (3 x 50 mL). The combined organic layers was washed with brine (3 x 50 mL) and dried over anhydrous sodium sulfate. After filtration, the filtrate was concentrated under reduced pressure and the residue was purified by silica gel column chromatography, eluted with 1% to 2% MeOH in DCM to give (3aR,4S,6R,7R,7aS)-6-((tert-butyltrimethylsilyloxy)methyl)-7-hydroxy-N,2,2,5-tetramethyl-hexahydro-[1,3]dioxolo[4,5-c]pyridine-4-carboxamide as a colorless solid (0.31 g, 80%): ¹H NMR (400 MHz, CDCl₃) δ 7.35 (brs, 1H), 5.01-4.96 (m, 1H), 3.99-3.94 (m, 2H), 3.84 (d, *J* = 1.6 Hz, 1H), 3.74-3.66 (m, 2H), 2.87 (d, *J* = 4.8 Hz, 3H), 2.71 (s, 3H), 2.63 (brs, 1H), 2.52-2.47 (m, 1H), 1.54 (s, 3H), 1.45 (s, 3H), 0.90 (s, 9H), 0.10 (s, 6H); MS (ESI, *m/z*): 389.0 [M + 1]⁺.

[00180] A solution of (3aR,4S,6R,7R,7aS)-6-((tert-butyltrimethylsilyloxy)methyl)-7-hydroxy-N,2,2,5-tetramethyl-hexahydro-[1,3]dioxolo[4,5-c]pyridine-4-carboxamide (20 mg, 0.05 mmol) in MeOH (3 mL) was treated with 1 M hydrochloride solution in MeOH (1 mL) for 2 hours at ambient temperature. The reaction was diluted with water (20 mL) and the resulting solution was extracted with EtOAc (2 x 10 mL). The aqueous layer was treated by freeze drying to give (2S,3R,4S,5R,6R)-3,4,5-trihydroxy-6-(hydroxymethyl)-N,1-dimethylpiperidine-2-carboxamide hydrochloride as a colorless solid (9.6 mg, 69%): ¹H NMR (400 MHz, D₂O) δ 4.70-4.69 (m, 1H), 4.24-4.21 (m, 1H), 4.18-4.10 (m, 1H), 4.81-4.09 (m, 4H), 2.87 (s, 3H), 2.71 (s, 3H); MS (ESI, *m/z*): 335.0 [M + 1-HCl]⁺.

20

Example 2

(2S,3R,4S,5R,6R)-1-((E)-3-([1,1'-biphenyl]-4-yl)allyl)-3,4,5-trihydroxy-6-(hydroxymethyl)-N-methylpiperidine-2-carboxamide



[00181] A solution of (3aR,4S,6R,7R,7aS)-6-((tert-butyltrimethylsilyloxy)methyl)-7-hydroxy-N,2,2-trimethyl-hexahydro-[1,3]dioxolo[4,5-c]pyridine-4-carboxamide (150 mg, 0.40 mmol) in MeOH (3 mL) was treated with 1 M hydrochloride solution in MeOH (1 mL) for 2 hours at ambient temperature. The reaction was diluted with water (20 mL) and the

resulting solution was extracted with EtOAc (2 x 10 mL). The aqueous layer was treated by freeze drying to give (2S,3R,4S,5R,6R)-3,4,5-trihydroxy-6-(hydroxymethyl)-N-methylpiperidine-2-carboxamide hydrochloride as a colorless solid (100 mg, 97%): ¹H NMR (400 MHz, D₂O) δ 4.30-4.28 (m, 1H), 4.19 (d, *J* = 4.0 Hz, 1H), 3.90-3.80 (m, 3H), 3.69-3.60 (m, 2H), 2.68 (s, 3H); MS (ESI, *m/z*): 221.0 [M + 1-HCl]⁺.

[00182] To a solution of 1-bromo-4-phenylbenzene (10 g, 42.9 mmol) in DMF (150 mL) was added ethyl acrylate (8.6 g, 85.8 mmol) followed by the addition of potassium carbonate (8.9 g, 64.5 mmol), triphenylphosphine (5.7 g, 22.1 mmol), tetrabutylammoniumbromide (7.1 g, 22.1 mmol) and palladium acetate (II) (5.1 g, 22.1 mmol) consecutively. The resulting mixture was stirred at 90 °C for 16 hours under a nitrogen atmosphere. After cooling to ambient, the reaction was quenched by water (500 mL) and the resulting mixture was extracted with EtOAc (3 x 200 mL). The combined organic layer was washed with brine (5 x 100 mL) and dried over anhydrous sodium sulfate. After filtration, the filtrate was concentrated under reduced pressure and the residue was purified by silica gel column chromatography, eluted with 1% to 5% EtOAc in petroleum ether to give ethyl (2*E*)-3-(4-phenylphenyl)prop-2-enoate as a colorless solid (8.1 g, 74%): ¹H NMR (300 MHz, CDCl₃) δ 7.70 (d, *J* = 15.9 Hz, 1H), 7.64-7.57 (m, 6H), 7.48-7.43 (m, 2H), 7.39-7.34 (m, 1H), 6.54 (d, *J* = 15.9 Hz, 1H), 4.28 (q, *J* = 7.2 Hz, 2H), 1.32 (t, *J* = 7.2 Hz, 3H); MS (ESI, *m/z*): 253.0 [M + 1]⁺.

[00183] To a solution of (2*E*)-3-(4-phenylphenyl)prop-2-enoate (1 g, 4 mmol) in toluene (20 mL) was added 1 *M* solution of diisobutylaluminumhydride in toluene (8 mL, 8 mmol) at -78 °C. The resulting mixture was stirred for 2 hours at ambient temperature and quenched by water (50 mL). The organic layer was separated out and the aqueous layer was extracted with EtOAc (2 x 50 mL). The combined organic layers was washed with brine and dried over anhydrous sodium sulfate. After filtration, the filtrate was concentrated under reduced pressure and the residue was purified by silica gel column chromatography, eluted with 5% to 10% EtOAc in petroleum ether to give (2*E*)-3-(4-phenylphenyl)prop-2-en-1-ol as a colorless solid (0.6 g, 73%): ¹H NMR (400 MHz, CDCl₃) δ 7.67-7.61 (m, 4H), 7.51-7.43 (m, 4H), 7.39-7.33 (m, 1H), 6.67 (d, *J* = 12.0 Hz, 1H), 6.48-6.41 (m, 1H), 4.39-4.33 (m, 2H); MS (ESI, *m/z*): 211.0 [M + 1]⁺.


[00184] A solution of (2*E*)-3-(4-phenylphenyl)prop-2-en-1-ol (100 mg, 0.47 mmol) in DCM was treated with thionyl chloride (0.5 mL) for 1 hour at ambient temperature. Solvent was evaporated out under reduced pressure to give crude 1-[(1*E*)-3-chloroprop-1-en-1-yl]-4-

phenylbenzene as a light yellow solid (100 mg), which was used directly in the next step without further purification.

[00185] To a solution of (2S,3R,4S,5R,6R)-3,4,5-trihydroxy-6-(hydroxymethyl)-N-methylpiperidine-2-carboxamide hydrochloride (100 mg, 0.39 mmol) in DMF (15 mL) was added potassium carbonate (107 mg, 0.78 mmol) and potassium iodide (65 mg, 0.39 mmol) at ambient temperature. The resulting mixture was stirred for 10 min at 45 °C followed by the addition of the above crude 1-[(1E)-3-chloroprop-1-en-1-yl]-4-phenylbenzene in several portions. After 2 hours, the reaction was quenched by water (5 mL) and the solvents were concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with 2% to 10% MeOH in DCM to give (2S,3R,4S,5R,6R)-3,4,5-trihydroxy-6-(hydroxymethyl)-N-methyl-1-[(2E)-3-(4-phenylphenyl)prop-2-en-1-yl]piperidine-2-carboxamide as a colorless solid (89 mg, 55%): ¹H NMR (300 MHz, CD₃OD) δ 7.64-7.60 (m, 4H), 7.57-7.46 (m, 4H), 7.41-7.33 (m, 1H), 6.58 (d, *J* = 15.9 Hz, 1H), 6.41-6.34 (m, 1H), 4.27-4.25 (m, 1H), 4.07-4.03 (m, 1H), 3.93-3.89 (m, 1H), 3.77-3.75 (m, 3H), 3.57-3.51 (m, 2H), 3.27-3.19 (m, 1H), 2.74 (s, 3H); MS (ESI, *m/z*): 413.0 [M + 1]⁺.

Example 3

(2S,3R,4S,5R,6R)-1-(3-([1,1'-biphenyl]-4-yl)propyl)-3,4,5-trihydroxy-6-(hydroxymethyl)-N-methylpiperidine-2-carboxamide








[00186] To a stirred solution of (2S,3R,4S,5R,6R)-3,4,5-trihydroxy-6-(hydroxymethyl)-N-methyl-1-[(2E)-3-(4-phenylphenyl)prop-2-en-1-yl]piperidine-2-carboxamide (60 mg, 0.15 mmol) in MeOH (20 mL) was added palladium on charcoal (20 mg, 10% w/w). The resulting mixture was kept under a hydrogen atmosphere (1 atm.) for 4 hours at ambient temperature. Then, the mixture was filtered through Celite and the filtrate was concentrated under reduced pressure. The residue was purified by silica gel column chromatography, eluted with 2% to 10% MeOH in DCM to give (2S,3R,4S,5R,6R)-3,4,5-trihydroxy-6-(hydroxymethyl)-N-methyl-1-[3-(4-phenylphenyl)propyl]piperidine-2-carboxamide as a colorless solid (27.1 mg,

45%): $^1\text{H NMR}$ (300 MHz, CD_3OD) δ 7.54 (d, $J = 7.2$ Hz, 2H), 7.47 (d, $J = 8.1$ Hz, 2H), 7.37 (t, $J = 7.2$ Hz, 2H), 7.22-7.28 (m, 3H), 4.17 (t, $J = 4.5$ Hz, 1H), 3.90-3.87 (m, 1H), 3.81-3.69 (m, 2H), 3.63-3.56 (m, 2H), 3.21-2.95 (m, 1H), 2.91-2.85 (m, 1H), 2.81-2.75 (m, 1H), 2.69 (s, 3H), 2.67-2.65 (m, 2H), 1.88-1.78 (m, 2H); MS (ESI, m/z): 415.0 $[\text{M} + 1]^+$

- 5 [00187] The following examples and Examples 9 to 56 in Table 1 are synthesized according to procedures analogous to the schemes and examples outlined herein.

Table 3

Example	Name	Structure	MH^+
4	(2S,3R,4S,5R,6R)-N-ethyl-3,4,5-trihydroxy-6-(hydroxymethyl)piperidine-2-carboxamide hydrochloride		235.0
5	(2S,3R,4S,5R,6R)-3,4,5-trihydroxy-6-(hydroxymethyl)-N-methyl-1-(3-phenylpropyl)piperidine-2-carboxamide		339.0
6	(2S,3R,4S,5R,6R)-N-ethyl-3,4,5-trihydroxy-6-(hydroxymethyl)-1-(3-phenylpropyl)piperidine-2-carboxamide		353.0
7	(2S,3R,4S,5R,6R)-3,4,5-trihydroxy-6-(hydroxymethyl)-N-methyl-1-(4-phenylbutyl)piperidine-2-carboxamide		353.0
8	(2S,3R,4S,5R,6R)-3,4,5-trihydroxy-6-(hydroxymethyl)-N-methyl-1-(5-methylhexyl)piperidine-2-carboxamide		319.0

Biological ActivityAssay for determination of K_I values for inhibition of O-GlcNAcase activity

[00188] Experimental procedure for kinetic analyses: Enzymatic reactions were carried out in a reaction containing 50 mM NaH_2PO_4 , 100 mM NaCl and 0.1% BSA (pH 7.0) using 2 mM 4-Methylumbelliferyl N-acetyl- β -D-glucosaminide dihydrate (Sigma M2133) dissolved in ddH_2O , as a substrate. The amount of purified human O-GlcNAcase enzyme used in the reaction was 0.7 nM. Test compound of varying concentrations was added to the enzyme prior to initiation of the reaction. The reaction was performed at room temperature in a 96-well plate and was initiated with the addition of substrate. The production of fluorescent product was measured every 60 sec for 45 min with a Tecan Infinite M200 plate-reader with excitation at 355 nM and emission detected at 460 nM, with 4-Methylumbelliferone (Sigma M1381) used to produce a standard curve. The slope of product production was determined for each concentration of compound tested and plotted, using standard curve fitting algorithms for sigmoidal dose response curves. The values for a four parameter logistic curve fit of the data were determined.

[00189] K_I values were determined using the Cheng-Prusoff equation; the K_m of O-GlcNAcase for substrate was 0.2 mM.

[00190] Many compounds of the invention exhibit K_I values for inhibition of O-GlcNAcase in the range 0.1 nM - 50 μM .

Assay for determination of K_I values for inhibition of β -hexosaminidase activity

[00191] Experimental procedure for kinetic analyses: Enzymatic reactions were carried out in a reaction containing 50 mM NaH_2PO_4 , 100 mM NaCl and 0.1% BSA (pH 7.0) using 2 mM 4-Methylumbelliferyl N-acetyl- β -D-glucosaminide dihydrate (Sigma M2133) dissolved in ddH_2O , as a substrate. The amount of purified human β -hexosaminidase enzyme used in the reaction was 24 nM. Test compound of varying concentrations was added to the enzyme prior to initiation of the reaction. The reaction was performed at room temperature in a 96-well plate and was initiated with the addition of substrate. The production of fluorescent product was measured every 60 sec for 45 min with a Tecan Infinite M200 plate-reader with excitation at 355 nM and emission detected at 460 nM, with 4-Methylumbelliferone (Sigma

M1381) used to produce a standard curve. The slope of product production was determined for each concentration of compound tested and plotted, using standard curve fitting algorithms for sigmoidal dose response curves. The values for a four parameter logistic curve fit of the data were determined.

5 [00192] K_I values were determined using the Cheng-Prusoff equation.

[00193] When tested in this assay, many of the compounds described herein exhibit K_I values for inhibition of β -hexosaminidase in the range 10 nM to greater than 100 μ M.

[00194] The selectivity ratio for inhibition of O-GlcNAcase over β -hexosaminidase is defined here as:

$$10 \quad K_I (\beta\text{-hexosaminidase})/K_I (\text{O-GlcNAcase})$$

[00195] In general, many of the compounds described herein exhibit a selectivity ratio in the range of about 10 to 100000. Thus, many compounds of the invention exhibit high selectivity for inhibition of O-GlcNAcase over β -hexosaminidase.

15 Assay for determination of cellular activity for compounds that inhibit O-GlcNAcase activity

[00196] Inhibition of O-GlcNAcase, which removes O-GlcNAc from cellular proteins, results in an increase in the level of O-GlcNAcylated protein in cells. An increase in O-GlcNAcylated protein can be measured by an antibody, such as RL-2, that binds to O-GlcNAcylated protein. The amount of O-GlcNAcylated protein:RL2 antibody interaction
20 can be measured by enzyme linked immunosorbant assay (ELISA) procedures.

[00197] A variety of tissue culture cell lines, expressing endogenous levels of O-GlcNAcase, can be utilized; examples include rat PC-12, and human U-87, or SK-N-SH cells. In this assay, rat PC-12 cells are plated in 96-well plates with approximately 10,000 cells / well. Compounds to be tested are dissolved in DMSO, either 2 or 10 mM stock solution, and then
25 diluted with DMSO and water in a two-step process using a Tecan workstation. Cells are treated with diluted compounds for 24 h (5.4 μ L into 200 μ L l well volume) to reach a final concentration of inhibitor desired to measure a compound concentration dependent response, typically, ten 3 fold dilution steps, starting at 10 μ M are used to determine a concentration response curve. To prepare a cell lysate, the media from compound treated cells is removed,
30 the cells are washed once with phosphate buffered saline (PBS) and then lysed for 5 minutes at room temperature in 50 μ L of Phosphosafe reagent (Novagen Inc, Madison, WI) with

protease inhibitors and PMSF. The cell lysate is collected and transferred to a new plate, which is then either coated to assay plates directly or frozen -80°C until used in the ELISA procedure. If desired, the total protein concentration of samples is determined using 20 µL of the sample using the BCA method.

5 [00198] The ELISA portion of the assay is performed in a black Maxisorp 96-well plate that is coated overnight at 4°C with 100 µL /well of the cell lysate (1:10 dilution of the lysate with PBS containing protease inhibitors, phosphatase inhibitors, and PMSF). The following day the wells are washed 3 times with 300 µL /well of Wash buffer (Tris-buffered saline with 0.1% Tween 20). The wells are blocked with 100 µL /well Blocking buffer (Tris buffered
10 saline w/0.05% Tween 20 and 2.5% Bovine serum albumin). Each well is then washed two times with 300 µL/well of wash buffer. The anti O-GlcNAc antibody RL-2 (Abcam, Cambridge, MA), diluted 1:1000 in blocking buffer, is added at 100 µL/well. The plate is sealed and incubated at 37°C for 2 h with gentle shaking. The wells are then washed 3-times with 300 µL/well wash buffer. To detect the amount of RL-2 bound horse-radish peroxidase
15 (HRP) conjugated goat anti-mouse secondary antibody (diluted 1:3000 in blocking buffer) is added at 100 µL /well. The plate is incubated for 60 min at 37°C with gentle shaking. Each well is then washed 3-times with 300 µL/well wash buffer. The detection reagent is added, 100 µL /well of Amplex Ultra RED reagent (prepared by adding 30 µL of 10 mM Amplex Ultra Red stock solution to 10 mL PBS with 18 µL 3% hydrogen peroxide, H₂O₂). The
20 detection reaction is incubated for 15 minutes at room temperature and then read with excitation at 530 nm and emission at 590 nm.

[00199] The amount of O-GlcNAcylated protein, as detected by the ELISA assay, is plotted for each concentration of test compound using standard curve fitting algorithms for sigmoidal dose response curves. The values for a four parameter logistic curve fit of the data are
25 determined, with the inflection point of the curve being the potency value for the test compound.

Assay for determination of apparent permeability (P_{app})

[00200] Bi-directional transport is evaluated in LLC-PK1 cells in order to determine
30 apparent permeability (P_{app}). LLC-PK1 cells can form a tight monolayer and therefore can be used to assess vectorial transport of compounds from basolateral to apical (B→A) and from apical to basolateral (A → B).

[00201] To determine P_{app} , LLC-PK1 cells are cultured in 96-well transwell culture plates (Millipore). Solutions containing the test compounds (1 μM) are prepared in Hank's Balanced Salt Solution with 10 mM HEPES. Substrate solution (150 μL) is added to either the apical (A) or the basolateral (B) compartment of the culture plate, and buffer (150 μL) is added to the compartment opposite to that containing the compound. At $t = 3$ h, 50 μL samples are removed from both sides of monolayers dosed with test compound and placed in 96 well plates, scintillant (200 μL) or internal standard (100 μL labetolol 1 μM) is added to the samples and concentration is determined by liquid scintillation counting in a MicroBeta Wallac Trilux scintillation counter (Perkin Elmer Life Sciences, Boston, MA) or by LCMS/MS (Applied Biosystems SCIEX API 5000 triple quadruple mass spectrometer). $[^3\text{H}]$ Verapamil (1 μM) is used as the positive control. The experiment is performed in triplicate.

[00202] The apparent permeability, P_{app} , is calculated by the following formula for samples taken at $t = 3$ h:








$$P_{app} = \frac{\text{Volume of Receptor Chamber (mL)}}{[\text{Area of membrane (cm}^2\text{)}][\text{Initial Concentration } (\mu\text{M)}]} \times \frac{\Delta \text{ in Concentration } (\mu\text{M})}{\Delta \text{ in Time (s)}}$$



Where: Volume of Receptor Chamber is 0.15 mL; Area of membrane is 0.11 cm^2 ; the Initial Concentration is the sum of the concentration measured in the donor plus concentration measured in receiver compartments at $t = 3$ h; Δ in Concentration is concentration in the receiver compartment at 3 h; and Δ in Time is the incubation time ($3 \times 60 \times 60 = 10800$ s). P_{app} is expressed as 10^{-6} cm/s. The P_{app} (LLC-PK1 cells) are the average of the P_{app} for transport from A to B and P_{app} for transport from B to A at $t = 3$ h:

$$P_{app}(\text{LLC - PK1 Cells}) = \frac{P_{app}(A \rightarrow B) + P_{app}(B \rightarrow A)}{2}$$

[00203] Representative data from the binding, cell-based, and permeability assays described above are shown in the following table. Certain compounds of the invention exhibit superior potency or permeability in one or more of these assays.

Table 4

Example	Structure	Cell-based ELISA EC ₅₀ (nM)	Fluorescence-based hOGA Ki (nM)	Papp LLC-PK1 cells (10 ⁻⁶ cm/s)
-		ND	165	ND
1		ND	1474	ND
2		ND	1808	ND
3		ND	9.4	3.0
4		ND	3297	ND
5		ND	1699	ND
6		ND	2496	ND

Example	Structure	Cell-based ELISA EC ₅₀ (nM)	Fluorescence-based hOGA Ki (nM)	Papp LLC-PK1 cells (10 ⁻⁶ cm/s)
7		ND	252	ND
8		ND	620	ND

[00204] The present invention has been described with regard to one or more embodiments. However, it will be apparent to persons skilled in the art that a number of variations and modifications can be made without departing from the scope of the invention as defined in the claims.

5

REFERENCES


1. C.R. Torres, G.W. Hart, *J Biol Chem* **1984**, 259, 3308-17.
2. R.S. Haltiwanger, G.D. Holt, and G.W. Hart, *J Biol Chem* **1990**, 265, 2563-8.
- 5 3. L.K. Kreppel, M.A. Blomberg, and G.W. Hart, *J Biol Chem* **1997**, 272, 9308-15.
4. W.A. Lubas, et al., *J Biol Chem* **1997**, 272, 9316-24.
5. W.A. Lubas, J.A. Hanover, *J Biol Chem* **2000**, 275, 10983-8.
6. D.L. Dong, G.W. Hart, *J Biol Chem* **1994**, 269, 19321-30.
7. Y. Gao, et al., *J Biol Chem* **2001**, 276, 9838-45.
- 10 8. E.P. Roquemore, et al., *Biochemistry* **1996**, 35, 3578-86.
9. S.P. Jackson, R. Tjian, *Cell* **1988**, 55, 125-33.
10. W.G. Kelly, M.E. Dahmus, and G.W. Hart, *J Biol Chem* **1993**, 268, 10416-24.
11. M.D. Roos, et al., *Mol Cell Biol* **1997**, 17, 6472-80.
12. N. Lamarre-Vincent, L.C. Hsieh-Wilson, *J Am Chem Soc* **2003**, 125, 6612-3.
- 15 13. F. Zhang, et al., *Cell* **2003**, 115, 715-25.
14. K. Vosseller, et al., *Proc Natl Acad Sci U S A* **2002**, 99, 5313-8.
15. W.A. Lubas, et al., *Biochemistry* **1995**, 34, 1686-94.
16. L.S. Griffith, B. Schmitz, *Biochem Biophys Res Commun* **1995**, 213, 424-31.
17. R.N. Cole, G.W. Hart, *J Neurochem* **1999**, 73, 418-28.
- 20 18. I. Braidman, et al., *Biochem J* **1974**, 143, 295-301.
19. R. Ueno, C.S. Yuan, *Biochim Biophys Acta* **1991**, 1074, 79-84.
20. C. Toleman, et al., *J Biol Chem* **2004**, 279, 53665-73.
21. F. Liu, et al., *Proc Natl Acad Sci U S A* **2004**, 101, 10804-9.
22. T.Y. Chou, G.W. Hart, *Adv Exp Med Biol* **2001**, 491, 413-8.
- 25 23. M. Goedert, et al., *Neuron* **1992**, 8, 159-68.
24. M. Goedert, et al., *Neuron* **1989**, 3, 519-26.
25. E. Kopke, et al., *J Biol Chem* **1993**, 268, 24374-84.
26. H. Ksiezak-Reding, W.K. Liu, and S.H. Yen, *Brain Res* **1992**, 597, 209-19.
27. P.V. Arriagada, et al., *Neurology* **1992**, 42, 631-9.
- 30 28. K.P. Riley, D.A. Snowdon, and W.R. Markesbery, *Ann Neurol* **2002**, 51, 567-77.
29. I. Alafuzoff, et al., *Acta Neuropathol (Berl)* **1987**, 74, 209-25.
30. C.X. Gong, et al., *J Neural Transm* **2005**, 112, 813-38.
31. K. Iqbal, et al., *J Neural Transm Suppl* **2002**, 309-19.
32. K. Iqbal, et al., *J Mol Neurosci* **2003**, 20, 425-9.
- 35 33. W. Noble, et al., *Proc Natl Acad Sci U S A* **2005**, 102, 6990-5.
34. S. Le Corre, et al., *Proc Natl Acad Sci U S A* **2006**, 103, 9673-8.
35. S.J. Liu, et al., *J Biol Chem* **2004**, 279, 50078-88.
36. G. Li, H. Yin, and J. Kuret, *J Biol Chem* **2004**, 279, 15938-45.
37. T.Y. Chou, G.W. Hart, and C.V. Dang, *J Biol Chem* **1995**, 270, 18961-5.
- 40 38. X. Cheng, G.W. Hart, *J Biol Chem* **2001**, 276, 10570-5.
39. X. Cheng, et al., *Biochemistry* **2000**, 39, 11609-20.
40. L.S. Griffith, B. Schmitz, *Eur J Biochem* **1999**, 262, 824-31.
41. K. Kamemura, G.W. Hart, *Prog Nucleic Acid Res Mol Biol* **2003**, 73, 107-36.
42. L. Wells, et al., *J Biol Chem* **2004**, 279, 38466-70.
- 45 43. L. Bertram, et al., *Science* **2000**, 290, 2302-3.
44. S. Hoyer, et al., *Journal of Neural Transmission* **1998**, 105, 423-438.
45. C.X. Gong, et al., *Journal of Alzheimers Disease* **2006**, 9, 1-12.

46. W.J. Jagust, et al., *Journal of Cerebral Blood Flow and Metabolism* **1991**, *11*, 323-330.
47. S. Hoyer, *Experimental Gerontology* **2000**, *35*, 1363-1372.
48. S. Hoyer, in *Frontiers in Clinical Neuroscience: Neurodegeneration and Neuroprotection, Vol. 541*, **2004**, 135-152.
- 5 49. R.N. Kalaria, S.I. Harik, *Journal of Neurochemistry* **1989**, *53*, 1083-1088.
50. I.A. Simpson, et al., *Annals of Neurology* **1994**, *35*, 546-551.
51. S.M. de la Monte, J.R. Wands, *Journal of Alzheimers Disease* **2005**, *7*, 45-61.
52. X.W. Zhu, G. Perry, and M.A. Smith, *Journal of Alzheimers Disease* **2005**, *7*, 81-84.
- 10 53. J.C. de la Torre, *Neurological Research* **2004**, *26*, 517-524.
54. S. Marshall, W.T. Garvey, and R.R. Traxinger, *Faseb J* **1991**, *5*, 3031-6.
55. S.P. Iyer, Y. Akimoto, and G.W. Hart, *J Biol Chem* **2003**, *278*, 5399-409.
56. K. Brickley, et al., *J Biol Chem* **2005**, *280*, 14723-32.
57. S. Knapp, C.H. Yang, and T. Haimowitz, *Tetrahedron Letters* **2002**, *43*, 7101-7104.
- 15 58. S.P. Iyer, G.W. Hart, *J Biol Chem* **2003**, *278*, 24608-16.
59. M. Jinek, et al., *Nat Struct Mol Biol* **2004**, *11*, 1001-7.
60. K. Kamemura, et al., *J Biol Chem* **2002**, *277*, 19229-35.
61. Y. Deng, et al., *FASEB J.* **2007**, fj.07-8309com.
62. L.F. Lau, et al., *Curr Top Med Chem* **2002**, *2*, 395-415.
- 20 63. M.P. Mazanetz, P.M. Fischer, *Nature Reviews Drug Discovery* **2007**, *6*, 464-479.
64. S.A. Yuzwa, et al., *Nat Chem Biol* **2008**, *4*, 483-490.
65. P. Bounelis, et al., *Shock* **2004**, *21 170 Suppl. 2*, 58-58.
66. N. Fulop, et al., *Circulation Research* **2005**, *97*, E28-E28.
67. J. Liu, R.B. Marchase, and J.C. Chatham, *Faseb Journal* **2006**, *20*, A317-A317.
- 25 68. R. Marchase, et al., *PCT Int. Appl. WO 2006016904* **2006**.
69. N. Fulop, et al., *Journal of Molecular and Cellular Cardiology* **2004**, *37*, 286-287.
70. N. Fulop, et al., *Faseb Journal* **2005**, *19*, A689-A690.
71. J. Liu, R.B. Marchase, and J.C. Chatham, *Journal of Molecular and Cellular Cardiology* **2007**, *42*, 177-185.
- 30 72. L.G. Not, et al., *Faseb Journal* **2006**, *20*, A1471-A1471.
73. S.L. Yang, et al., *Shock* **2006**, *25*, 600-607.
74. L.Y. Zou, et al., *Faseb Journal* **2005**, *19*, A1224-A1224.
75. R.B. Marchase, et al., *Circulation* **2004**, *110*, 1099-1099.
76. J. Liu, et al., *Journal of Molecular and Cellular Cardiology* **2006**, *40*, 303-312.
- 35 77. J. Liu, J.C. Chatham, and R.B. Marchase, *Faseb Journal* **2005**, *19*, A691-A691.
78. T. Nagy, et al., *American Journal of Physiology-Cell Physiology* **2006**, *290*, C57-C65.
79. N. Fulop, R.B. Marchase, and J.C. Chatham, *Cardiovascular Research* **2007**, *73*, 288-297.
80. T. Lefebvre, et al., *Expert Review of Proteomics* **2005**, *2*, 265-275.
- 40 81. B. Henrissat, A. Bairoch, *Biochem J* **1993**, *293 (Pt 3)*, 781-8.
82. B. Henrissat, A. Bairoch, *Biochem J* **1996**, *316 (Pt 2)*, 695-6.
83. L. Wells, K. Vosseller, and G.W. Hart, *Science* **2001**, *291*, 2376-8.
84. J.A. Hanover, *FASEB J* **2001**, *15*, 1865-76.
85. D.A. McClain, et al., *Proc Natl Acad Sci U S A* **2002**, *99*, 10695-9.
- 45 86. P.J. Yao, P.D. Coleman, *J Neurosci* **1998**, *18*, 2399-411.
87. W.H. Yang, et al., *Nature Cell Biology* **2006**, *8*, 1074-U53.
88. B. Triggs-Raine, D.J. Mahuran, and R.A. Gravel, *Adv Genet* **2001**, *44*, 199-224.
89. D. Zhou, et al., *Science* **2004**, 1786-89.
90. G. Legler, et al., *Biochim Biophys Acta* **1991**, *1080*, 89-95.
- 50 91. M. Horsch, et al., *Eur J Biochem* **1991**, *197*, 815-8.

92. J. Liu, et al., *Chem Biol* **2001**, *8*, 701-11.
93. S. Knapp, et al., *J. Am. Chem. Soc.* **1996**, *118*, 6804-6805.
94. V.H. Lillielund, et al., *Chem Rev* **2002**, *102*, 515-53.
95. R.J. Konrad, et al., *Biochem J* **2001**, *356*, 31-41.
- 5 96. K. Liu, et al., *J Neurochem* **2004**, *89*, 1044-55.
97. G. Parker, et al., *J Biol Chem* **2004**, *279*, 20636-42.
98. E.B. Arias, J. Kim, and G.D. Cartee, *Diabetes* **2004**, *53*, 921-30.
99. A. Junod, et al., *Proc Soc Exp Biol Med* **1967**, *126*, 201-5.
100. R.A. Bennett, A.E. Pegg, *Cancer Res* **1981**, *41*, 2786-90.
- 10 101. K.D. Kroncke, et al., *Biol Chem Hoppe Seyler* **1995**, *376*, 179-85.
102. H. Yamamoto, Y. Uchigata, and H. Okamoto, *Nature* **1981**, *294*, 284-6.
103. K. Yamada, et al., *Diabetes* **1982**, *31*, 749-53.
104. V. Burkart, et al., *Nat Med* **1999**, *5*, 314-9.
105. M.D. Roos, et al., *Proc Assoc Am Physicians* **1998**, *110*, 422-32.
- 15 106. Y. Gao, G.J. Parker, and G.W. Hart, *Arch Biochem Biophys* **2000**, *383*, 296-302.
107. R. Okuyama, M. Yachi, *Biochem Biophys Res Commun* **2001**, *287*, 366-71.
108. N.E. Zachara, et al., *J Biol Chem* **2004**, *279*, 30133-42.
109. J.A. Hanover, et al., *Arch Biochem Biophys* **1999**, *362*, 38-45.
110. K. Liu, et al., *Mol Cell Endocrinol* **2002**, *194*, 135-46.
- 20 111. M.S. Macauley, et al., *J Biol Chem* **2005**, *280*, 25313-22.
112. B.L. Mark, et al., *J Biol Chem* **2001**, *276*, 10330-7.
113. R.S. Haltiwanger, K. Grove, and G.A. Philipsberg, *J Biol Chem* **1998**, *273*, 3611-7.
114. D.J. Miller, X. Gong, and B.D. Shur, *Development* **1993**, *118*, 1279-89.
115. L.Y. Zou, et al., *Shock* **2007**, *27*, 402-408.
- 25 116. J.B. Huang, A.J. Clark, and H.R. Petty, *Cellular Immunology* **2007**, *245*, 1-6.
117. N.E. Zachara, et al., *Abstract 418 in Joint Meeting of the Society for Glycobiology and the Japanese Society of Carbohydrate Research*. Honolulu, Hawaii, **2004**.
118. L.Y. Zou, et al., *Faseb Journal* **2006**, *20*, A1471-A1471.
119. V. Champattanachai, R.B. Marchase, and J.C. Chatham, *American Journal of Physiology-Cell Physiology* **2007**, *292*, C178-C187.
- 30 120. V. Champattanachai, R.B. Marchase, and J.C. Chatham, *American Journal of Physiology-Cell Physiology* **2008**, *294*, C1509-C1520.
121. I. Khlistunova, et al., *Current Alzheimer Research* **2007**, *4*, 544-546.
122. P. Friedhoff, et al., *Biochemistry* **1998**, *37*, 10223-10230.
- 35 123. M. Pickhardt, et al., *Journal of Biological Chemistry* **2005**, *280*, 3628-3635.

WHAT IS CLAIMED IS:

1. A compound of Formula (I) or a pharmaceutically acceptable salt thereof:



(I)

5 wherein

R^1 is H and R^2 is $C(O)NR^{11}_2$, or R^1 is $C(O)NR^{11}_2$ and R^2 is H, or R^1 is H and R^2 is selected from the group consisting of: H, CH_3 , CH_2F , CHF_2 , and CH_2OH , where each R^{11} is independently H or C_{1-6} alkyl;

10 R^3 is H and R^4 is OH, or R^3 is H and R^4 is H, or R^3 is H and R^4 is F, or R^3 is F and R^4 is H, or R^3 is F and R^4 is F, or R^3 is OH and R^4 is H;

R^5 is H and R^6 is OH, or R^5 is H and R^6 is H, or R^5 is H and R^6 is F, or R^5 is F and R^6 is H, or R^5 is F and R^6 is F, or R^5 is OH and R^6 is H;

R^7 is H and R^8 is OH, or R^7 is H and R^8 is H, or R^7 is H and R^8 is F, or R^7 is F and R^8 is H, or R^7 is F and R^8 is F, or R^7 is OH and R^8 is H;

15 R^9 is selected from the group consisting of: H, CH_3 , CH_2F , CHF_2 , and CH_2OH ; and

R^{10} is selected from the group consisting of: H, C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, C_{1-10} acyl, C_{8-20} arylalkylacyl, C_{3-20} heteroarylalkylacyl, C_{7-20} arylalkyl, C_{8-20} arylalkenyl, C_{8-20} arylalkynyl, C_{2-20} heteroarylalkyl, C_{3-20} heteroarylalkenyl, and C_{3-20} heteroarylalkynyl, each excluding H optionally substituted from one to four substituents with one of more of halo, OH, OCF_3 , CN, SO_2NH_2 , SO_2Me , $C(O)NH_2$, CH_2F , CHF_2 , CF_3 , CH_2CH_2F , CH_2CF_3 , $CH_2CH_2CH_2F$, C_{1-6} alkyl, C_{1-6} alkoxy, or C_{3-7} cycloalkyl.

2. The compound of claim 1 wherein the compound is a compound described in Table 1.

3. The compound of claim 1 wherein the compound is selected from the following group:

(2S,3R,4S,5R,6R)-N-ethyl-3,4,5-trihydroxy-6-(hydroxymethyl)piperidine-2-carboxamide;


25 (2S,3R,4S,5R,6R)-3,4,5-trihydroxy-6-(hydroxymethyl)-N,1-dimethylpiperidine-2-carboxamide;

- (2S,3R,4S,5R,6R)-3,4,5-trihydroxy-6-(hydroxymethyl)-N-methyl-1-(5-methylhexyl)piperidine-2-carboxamide;
- (2S,3R,4S,5R,6R)-3,4,5-trihydroxy-6-(hydroxymethyl)-N-methyl-1-(3-phenylpropyl)piperidine-2-carboxamide;
- 5 (2S,3R,4S,5R,6R)-N-ethyl-3,4,5-trihydroxy-6-(hydroxymethyl)-1-(3-phenylpropyl)piperidine-2-carboxamide;
- (2S,3R,4S,5R,6R)-3,4,5-trihydroxy-6-(hydroxymethyl)-N-methyl-1-(4-phenylbutyl)piperidine-2-carboxamide;
- (2S,3R,4S,5R,6R)-1-((E)-3-([1,1'-biphenyl]-4-yl)allyl)-3,4,5-trihydroxy-6-(hydroxymethyl)-
- 10 N-methylpiperidine-2-carboxamide;
- (2S,3R,4S,5R,6R)-1-(3-([1,1'-biphenyl]-4-yl)propyl)-3,4,5-trihydroxy-6-(hydroxymethyl)-N-methylpiperidine-2-carboxamide (2S,3R,4S)-1-hexyl-3,4-dihydroxy-N-methylpiperidine-2-carboxamide;
- (2S,3R,4S,5R)-3,4,5-trihydroxy-N-methylpiperidine-2-carboxamide;
- 15 (2S,3R,4S,5R)-3,4,5-trihydroxy-N,1-dimethylpiperidine-2-carboxamide;
- (2S,3R,4S,5R)-3,4,5-trihydroxy-1-methylpiperidine-2-carboxamide;
- (2S,3R,4S,5R)-N-ethyl-3,4,5-trihydroxypiperidine-2-carboxamide;
- (2S,3R,4S,5R)-N-ethyl-3,4,5-trihydroxy-1-methylpiperidine-2-carboxamide;
- (2S,3R,4S,5R,6R)-N-ethyl-3,4,5-trihydroxy-6-methylpiperidine-2-carboxamide;
- 20 (2S,3R,4S,5R,6R)-N-ethyl-3,4,5-trihydroxy-6-methyl-1-phenethylpiperidine-2-carboxamide;
- (2S,3R,4S,5R)-N-ethyl-3,4,5-trihydroxy-1-(3-phenylpropyl)piperidine-2-carboxamide;
- (2S,3S,4R,5S,6R)-N-ethyl-4-fluoro-3,5-dihydroxy-6-methyl-1-phenethylpiperidine-2-carboxamide;
- (2R,3R,4S,5R,6R)-N-ethyl-3-fluoro-4,5-dihydroxy-6-(hydroxymethyl)-1-(3-
- 25 phenylpropyl)piperidine-2-carboxamide;
- (2S,3R,4S,5R,6R)-N-ethyl-4-fluoro-3,5-dihydroxy-6-(hydroxymethyl)-1-(3-phenylpropyl)piperidine-2-carboxamide;
- (2S,3R,4R,5R,6R)-N-ethyl-5-fluoro-3,4-dihydroxy-6-(hydroxymethyl)-1-(3-phenylpropyl)piperidine-2-carboxamide;
- 30 (2S,3R,4R,5R,6S)-N-ethyl-6-(fluoromethyl)-3,4,5-trihydroxy-1-(3-phenylpropyl)piperidine-2-carboxamide;
- (2R,3S,4S,5R,6R)-N-ethyl-3-fluoro-4,5-dihydroxy-6-(hydroxymethyl)-1-(3-phenylpropyl)piperidine-2-carboxamide;

- (2S,3R,4R,5R,6R)-N-ethyl-4-fluoro-3,5-dihydroxy-6-(hydroxymethyl)-1-(3-phenylpropyl)piperidine-2-carboxamide;
- (2S,3R,4R,5S,6R)-N-ethyl-5-fluoro-3,4-dihydroxy-6-(hydroxymethyl)-1-(3-phenylpropyl)piperidine-2-carboxamide;
- 5 (2S,3R,4R,5R,6S)-6-(difluoromethyl)-N-ethyl-3,4,5-trihydroxy-1-(3-phenylpropyl)piperidine-2-carboxamide;
- (2S,4R,5R,6R)-N-ethyl-4,5-dihydroxy-6-(hydroxymethyl)-1-(3-phenylpropyl)piperidine-2-carboxamide;
- (2S,3S,5S,6R)-N-ethyl-3,5-dihydroxy-6-(hydroxymethyl)-1-(3-phenylpropyl)piperidine-2-
- 10 carboxamide;
- (2S,3R,4S,6S)-N-ethyl-3,4-dihydroxy-6-(hydroxymethyl)-1-(3-phenylpropyl)piperidine-2-carboxamide;
- (2S,3R,4S,5R,6R)-N-ethyl-3,4,5-trihydroxy-6-methyl-1-(3-phenylpropyl)piperidine-2-carboxamide;
- 15 (2S,3R,4S,5R)-N-ethyl-3,4,5-trihydroxy-1-(3-phenylpropyl)piperidine-2-carboxamide;
- (2S,3R,4S,5R,6S)-6-(difluoromethyl)-N-ethyl-4-fluoro-3,5-dihydroxy-1-(3-phenylpropyl)piperidine-2-carboxamide;
- (2R,3S,4S,5R,6S)-N-ethyl-3-fluoro-6-(fluoromethyl)-4,5-dihydroxy-1-(3-phenylpropyl)piperidine-2-carboxamide;
- 20 (2R,3R,4S,5R,6S)-6-(difluoromethyl)-N-ethyl-3-fluoro-4,5-dihydroxy-1-(3-phenylpropyl)piperidine-2-carboxamide;
- (2R,3R,4S,5R)-N-ethyl-3-fluoro-4,5-dihydroxy-1-(3-phenylpropyl)piperidine-2-carboxamide;
- (2S,3R,4S,5R)-N-ethyl-4-fluoro-3,5-dihydroxy-1-(3-phenylpropyl)piperidine-2-carboxamide;
- 25 (2S,3R,4R,5R)-N-ethyl-5-fluoro-3,4-dihydroxy-1-(3-phenylpropyl)piperidine-2-carboxamide;
- (2R,3S,4S,5R)-N-ethyl-3-fluoro-4,5-dihydroxy-1-(3-phenylpropyl)piperidine-2-carboxamide;
- (2S,3R,4R,5R)-N-ethyl-4-fluoro-3,5-dihydroxy-1-(3-phenylpropyl)piperidine-2-carboxamide;
- 30 (2S,3R,4R,5S)-N-ethyl-5-fluoro-3,4-dihydroxy-1-(3-phenylpropyl)piperidine-2-carboxamide;
- (2S,4R,5R)-N-ethyl-4,5-dihydroxy-1-(3-phenylpropyl)piperidine-2-carboxamide;
- (2S,3S,5S)-N-ethyl-3,5-dihydroxy-1-(3-phenylpropyl)piperidine-2-carboxamide;
- (2S,3R,4S)-N-ethyl-3,4-dihydroxy-1-(3-phenylpropyl)piperidine-2-carboxamide;

- (2R,3R,4S,5R)-N-ethyl-3,4-difluoro-5-hydroxy-1-(3-phenylpropyl)piperidine-2-carboxamide;
- (2S,3R,4S,5R,6R)-N-ethyl-3,4,5-trihydroxy-6-(hydroxymethyl)-1-(3-phenylpropanoyl)piperidine-2-carboxamide;
- 5 (2S,3R,4S,5R,6R)-N-ethyl-1-(3-(6-fluoropyridin-3-yl)propyl)-3,4,5-trihydroxy-6-(hydroxymethyl)piperidine-2-carboxamide;
- (2S,3R,4S,5R,6R)-N-ethyl-3,4,5-trihydroxy-6-(hydroxymethyl)-1-(4,4,4-trifluorobutyl)piperidine-2-carboxamide;
- (2S,3R,4S,5R,6R)-N-ethyl-3,4,5-trihydroxy-6-(hydroxymethyl)-1-(3-(5-
- 10 (trifluoromethoxy)benzo[d]thiazol-2-yl)propyl)piperidine-2-carboxamide;
- (2S,3R,4S,5R,6R)-N-ethyl-3,4,5-trihydroxy-6-(hydroxymethyl)-1-((Z)-5,5,5-trifluoropent-3-en-1-yl)piperidine-2-carboxamide;
- (2S,3R,4S,5R,6R)-N-ethyl-3,4,5-trihydroxy-6-(hydroxymethyl)-1-(pent-4-yn-1-yl)piperidine-2-carboxamide;
- 15 (2S,3R,4S,5R,6R)-1-butyryl-N-ethyl-3,4,5-trihydroxy-6-(hydroxymethyl)piperidine-2-carboxamide;
- (2S,3R,4S,5R,6R)-N-ethyl-1-(3-(6-fluoropyridin-3-yl)propanoyl)-3,4,5-trihydroxy-6-(hydroxymethyl)piperidine-2-carboxamide;
- (2S,3R,4S,5R,6R)-1-cinnamyl-N-ethyl-3,4,5-trihydroxy-6-(hydroxymethyl)piperidine-2-
- 20 carboxamide;
- (2S,3R,4S,5R,6R)-N-ethyl-3,4,5-trihydroxy-6-(hydroxymethyl)-1-(3-phenylprop-2-yn-1-yl)piperidine-2-carboxamide;
- (2S,3R,4S,5R,6R)-N-ethyl-1-((E)-3-(6-fluoropyridin-3-yl)allyl)-3,4,5-trihydroxy-6-(hydroxymethyl)piperidine-2-carboxamide;
- 25 (2S,3R,4S,5R,6R)-N-ethyl-1-(3-(6-fluoropyridin-3-yl)prop-2-yn-1-yl)-3,4,5-trihydroxy-6-(hydroxymethyl)piperidine-2-carboxamide;
- or a pharmaceutically acceptable salt of any of the foregoing compounds.
4. The compound of claim 1 wherein the compound is a prodrug.
5. The compound of any one of claims 1 to 4 wherein the compound selectively inhibits
- 30 an O-glycoprotein 2-acetamido-2-deoxy- β -D-glucopyranosidase (O-GlcNAcase).
6. The compound of any one of claims 1 to 5 wherein the compound selectively binds an O-GlcNAcase.

7. The compound of any one of claims 1 to 6 wherein the compound selectively inhibits the cleavage of 2-acetamido-2-deoxy- β -D-glucopyranoside (O-GlcNAc).
8. The compound of claim 6 wherein the O-GlcNAcase is a mammalian O-GlcNAcase.
9. The compound of any one of claims 1 to 8 wherein the compound does not substantially
5 inhibit a mammalian β -hexosaminidase.
10. A pharmaceutical composition comprising the compound of any one of claims 1 to 9 or a pharmaceutically acceptable salt thereof in combination with a pharmaceutically acceptable carrier.
11. A method of selectively inhibiting an O-GlcNAcase in a subject in need thereof, the
10 method comprising administering to the subject an effective amount of a compound of Formula (I) or a pharmaceutically acceptable salt thereof:




(I)

wherein

- 15 R^1 is H and R^2 is $C(O)NR^{11}_2$, or R^1 is $C(O)NR^{11}_2$ and R^2 is H, or R^1 is H and R^2 is selected from the group consisting of: H, CH_3 , CH_2F , CHF_2 , and CH_2OH , where each R^{11} is independently H or C_{1-6} alkyl;
- R^3 is H and R^4 is OH, or R^3 is H and R^4 is H, or R^3 is H and R^4 is F, or R^3 is F and R^4 is H, or R^3 is F and R^4 is F, or R^3 is OH and R^4 is H;
- 20 R^5 is H and R^6 is OH, or R^5 is H and R^6 is H, or R^5 is H and R^6 is F, or R^5 is F and R^6 is H, or R^5 is F and R^6 is F, or R^5 is OH and R^6 is H;
- R^7 is H and R^8 is OH, or R^7 is H and R^8 is H, or R^7 is H and R^8 is F, or R^7 is F and R^8 is H, or R^7 is F and R^8 is F, or R^7 is OH and R^8 is H;
- R^9 is selected from the group consisting of: H, CH_3 , CH_2F , CHF_2 , and CH_2OH ; and

R^{10} is selected from the group consisting of: H, C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, C_{1-10} acyl, C_{8-20} arylalkylacyl, C_{3-20} heteroarylalkylacyl, C_{7-20} arylalkyl, C_{8-20} arylalkenyl, C_{8-20} arylalkynyl, C_{2-20} heteroarylalkyl, C_{3-20} heteroarylalkenyl, and C_{3-20} heteroarylalkynyl, each excluding H optionally substituted from one to four substituents with one of more of halo, OH, OCF_3 , CN, SO_2NH_2 , SO_2Me , $C(O)NH_2$, CH_2F , CHF_2 , CF_3 , CH_2CH_2F , CH_2CF_3 , $CH_2CH_2CH_2F$, C_{1-6} alkyl, C_{1-6} alkoxy, or C_{3-7} cycloalkyl.

12. A method of elevating the level of O-GlcNAc in a subject in need thereof, the method comprising administering to the subject an effective amount of a compound of Formula (I) or a pharmaceutically acceptable salt thereof:



(I)

wherein

R^1 is H and R^2 is $C(O)NR^{11}_2$, or R^1 is $C(O)NR^{11}_2$ and R^2 is H, or R^1 is H and R^2 is selected from the group consisting of: H, CH_3 , CH_2F , CHF_2 , and CH_2OH , where each R^{11} is independently H or C_{1-6} alkyl;

R^3 is H and R^4 is OH, or R^3 is H and R^4 is H, or R^3 is H and R^4 is F, or R^3 is F and R^4 is H, or R^3 is F and R^4 is F, or R^3 is OH and R^4 is H;

R^5 is H and R^6 is OH, or R^5 is H and R^6 is H, or R^5 is H and R^6 is F, or R^5 is F and R^6 is H, or R^5 is F and R^6 is F, or R^5 is OH and R^6 is H;


R^7 is H and R^8 is OH, or R^7 is H and R^8 is H, or R^7 is H and R^8 is F, or R^7 is F and R^8 is H, or R^7 is F and R^8 is F, or R^7 is OH and R^8 is H;

R^9 is selected from the group consisting of: H, CH_3 , CH_2F , CHF_2 , and CH_2OH ; and

R^{10} is selected from the group consisting of: H, C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, C_{1-10} acyl, C_{8-20} arylalkylacyl, C_{3-20} heteroarylalkylacyl, C_{7-20} arylalkyl, C_{8-20} arylalkenyl, C_{8-20} arylalkynyl, C_{2-20} heteroarylalkyl, C_{3-20} heteroarylalkenyl, and C_{3-20} heteroarylalkynyl, each excluding H optionally substituted from one to four substituents with

one of more of halo, OH, OCF₃, CN, SO₂NH₂, SO₂Me, C(O)NH₂, CH₂F, CHF₂, CF₃, CH₂CH₂F, CH₂CF₃, CH₂CH₂CH₂F, C₁₋₆ alkyl, C₁₋₆ alkoxy, or C₃₋₇ cycloalkyl.

13. A method of treating a condition that is modulated by an O-GlcNAcase, in a subject in need thereof, the method comprising administering to the subject an effective amount of a compound of Formula (I) or a pharmaceutically acceptable salt thereof:



(I)

wherein

- R¹ is H and R² is C(O)NR¹¹₂, or R¹ is C(O)NR¹¹₂ and R² is H, or R¹ is H and R² is selected from the group consisting of: H, CH₃, CH₂F, CHF₂, and CH₂OH, where each R¹¹ is independently H or C₁₋₆ alkyl;

R³ is H and R⁴ is OH, or R³ is H and R⁴ is H, or R³ is H and R⁴ is F, or R³ is F and R⁴ is H, or R³ is F and R⁴ is F, or R³ is OH and R⁴ is H;

- R⁵ is H and R⁶ is OH, or R⁵ is H and R⁶ is H, or R⁵ is H and R⁶ is F, or R⁵ is F and R⁶ is H, or R⁵ is F and R⁶ is F, or R⁵ is OH and R⁶ is H;

R⁷ is H and R⁸ is OH, or R⁷ is H and R⁸ is H, or R⁷ is H and R⁸ is F, or R⁷ is F and R⁸ is H, or R⁷ is F and R⁸ is F, or R⁷ is OH and R⁸ is H;

R⁹ is selected from the group consisting of: H, CH₃, CH₂F, CHF₂, and CH₂OH; and

- R¹⁰ is selected from the group consisting of: H, C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl, C₁₋₁₀ acyl, C₈₋₂₀ arylalkylacyl, C₃₋₂₀ heteroarylalkylacyl, C₇₋₂₀ arylalkyl, C₈₋₂₀ arylalkenyl, C₈₋₂₀ arylalkynyl, C₂₋₂₀ heteroarylalkyl, C₃₋₂₀ heteroarylalkenyl, and C₃₋₂₀ heteroarylalkynyl, each excluding H optionally substituted from one to four substituents with one of more of halo, OH, OCF₃, CN, SO₂NH₂, SO₂Me, C(O)NH₂, CH₂F, CHF₂, CF₃, CH₂CH₂F, CH₂CF₃, CH₂CH₂CH₂F, C₁₋₆ alkyl, C₁₋₆ alkoxy, or C₃₋₇ cycloalkyl.

14. The method of claim 13 wherein the condition is selected from one or more of the group consisting of an inflammatory disease, an allergy, asthma, allergic rhinitis,

hypersensitivity lung diseases, hypersensitivity pneumonitis, eosinophilic pneumonias, delayed-type hypersensitivity, atherosclerosis, interstitial lung disease (ILD), idiopathic pulmonary fibrosis, ILD associated with rheumatoid arthritis, systemic lupus erythematosus, ankylosing spondylitis, systemic sclerosis, Sjogren's syndrome, polymyositis or


5 dermatomyositis, systemic anaphylaxis or hypersensitivity response, drug allergy, insect sting allergy, autoimmune disease, rheumatoid arthritis, psoriatic arthritis, multiple sclerosis, Guillain-Barré syndrome, systemic lupus erythematosus, myasthenia gravis, glomerulonephritis, autoimmune thyroiditis, graft rejection, allograft rejection, graft-versus-host disease, inflammatory bowel disease, Crohn's disease, ulcerative colitis,

10 spondyloarthropathy, scleroderma, psoriasis, T-cell mediated psoriasis, inflammatory dermatosis, dermatitis, eczema, atopic dermatitis, allergic contact dermatitis, urticaria, vasculitis, necrotizing, cutaneous, and hypersensitivity vasculitis, eosinophilic myositis, eosinophilic fasciitis, solid organ transplant rejection, heart transplant rejection, lung transplant rejection, liver transplant rejection, kidney transplant rejection, pancreas transplant

15 rejection, kidney allograft, lung allograft, epilepsy, pain, fibromyalgia, stroke, neuroprotection.

15. A method of treating a condition selected from the group consisting of a neurodegenerative disease, a tauopathy, cancer and stress, in a subject in need thereof, the method comprising administering to the subject an effective amount of a compound of

20 Formula (I) or a pharmaceutically acceptable salt thereof:



(I)

wherein

R¹ is H and R² is C(O)NR¹¹₂, or R¹ is C(O)NR¹¹₂ and R² is H, or R¹ is H and R² is

25 selected from the group consisting of: H, CH₃, CH₂F, CHF₂, and CH₂OH, where each R¹¹ is independently H or C₁₋₆ alkyl;

R³ is H and R⁴ is OH, or R³ is H and R⁴ is H, or R³ is H and R⁴ is F, or R³ is F and R⁴ is H, or R³ is F and R⁴ is F, or R³ is OH and R⁴ is H;

R^5 is H and R^6 is OH, or R^5 is H and R^6 is H, or R^5 is H and R^6 is F, or R^5 is F and R^6 is H, or R^5 is F and R^6 is F, or R^5 is OH and R^6 is H;

R^7 is H and R^8 is OH, or R^7 is H and R^8 is H, or R^7 is H and R^8 is F, or R^7 is F and R^8 is H, or R^7 is F and R^8 is F, or R^7 is OH and R^8 is H;

5 R^9 is selected from the group consisting of: H, CH₃, CH₂F, CHF₂, and CH₂OH; and

R^{10} is selected from the group consisting of: H, C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl, C₁₋₁₀ acyl, C₈₋₂₀ arylalkylacyl, C₃₋₂₀ heteroarylalkylacyl, C₇₋₂₀ arylalkyl, C₈₋₂₀ arylalkenyl, C₈₋₂₀ arylalkynyl, C₂₋₂₀ heteroarylalkyl, C₃₋₂₀ heteroarylalkenyl, and C₃₋₂₀ heteroarylalkynyl, each excluding H optionally substituted from one to four substituents with
 10 one of more of halo, OH, OCF₃, CN, SO₂NH₂, SO₂Me, C(O)NH₂, CH₂F, CHF₂, CF₃, CH₂CH₂F, CH₂CF₃, CH₂CH₂CH₂F, C₁₋₆ alkyl, C₁₋₆ alkoxy, or C₃₋₇ cycloalkyl.

16. The method of claim 15 wherein the condition is selected from one or more of the group consisting of Alzheimer's disease, Amyotrophic lateral sclerosis (ALS), Amyotrophic lateral sclerosis with cognitive impairment (ALSci), Argyrophilic grain dementia, Bluit
 15 disease, Corticobasal degeneration (CBD), Dementia pugilistica, Diffuse neurofibrillary tangles with calcification, Down's syndrome, Familial British dementia, Familial Danish dementia, Frontotemporal dementia with parkinsonism linked to chromosome 17 (FTDP-17), Gerstmann-Straussler-Scheinker disease, Guadeloupean parkinsonism, Hallevorden-Spatz disease (neurodegeneration with brain iron accumulation type 1), Multiple system atrophy,
 20 Myotonic dystrophy, Niemann-Pick disease (type C), Pallido-ponto-nigral degeneration, Parkinsonism-dementia complex of Guam, Pick's disease (PiD), Post-encephalitic parkinsonism (PEP), Prion diseases (including Creutzfeldt-Jakob Disease (CJD), Variant Creutzfeldt-Jakob Disease (vCJD), Fatal Familial Insomnia, and Kuru), Progressive supercortical gliosis, Progressive supranuclear palsy (PSP), Richardson's syndrome,
 25 Subacute sclerosing panencephalitis, Tangle-only dementia, Huntington's disease, Parkinson's disease, Schizophrenia, Mild Cognitive Impairment (MCI), Neuropathy (including peripheral neuropathy, autonomic neuropathy, neuritis, and diabetic neuropathy), or Glaucoma.

17. The method of claim 15 wherein the stress is a cardiac disorder.

30 18. The method of claim 17 wherein the cardiac disorder is selected from one or more of the group consisting of ischemia; hemorrhage; hypovolemic shock; myocardial infarction; an


interventional cardiology procedure; cardiac bypass surgery; fibrinolytic therapy; angioplasty; and stent placement.

19. The method of any one of claims 11 to 18 wherein the compound is selected from the group consisting of one or more of the compounds described in Table 1.

5 20. The method of any one of claims 11 to 19 wherein said administering increases the level of O-GlcNAc in the subject.

21. The method of any one of claims 11 to 20 wherein the subject is a human.

22. Use of a compound of an effective amount of a compound of Formula (I) or a pharmaceutically acceptable salt thereof:



(I)

wherein

R^1 is H and R^2 is $C(O)NR^{11}_2$, or R^1 is $C(O)NR^{11}_2$ and R^2 is H, or R^1 is H and R^2 is selected from the group consisting of: H, CH_3 , CH_2F , CHF_2 , and CH_2OH , where each R^{11} is independently H or C_{1-6} alkyl;

R^3 is H and R^4 is OH, or R^3 is H and R^4 is H, or R^3 is H and R^4 is F, or R^3 is F and R^4 is H, or R^3 is F and R^4 is F, or R^3 is OH and R^4 is H;

R^5 is H and R^6 is OH, or R^5 is H and R^6 is H, or R^5 is H and R^6 is F, or R^5 is F and R^6 is H, or R^5 is F and R^6 is F, or R^5 is OH and R^6 is H;

R^7 is H and R^8 is OH, or R^7 is H and R^8 is H, or R^7 is H and R^8 is F, or R^7 is F and R^8 is H, or R^7 is F and R^8 is F, or R^7 is OH and R^8 is H;

R^9 is selected from the group consisting of: H, CH_3 , CH_2F , CHF_2 , and CH_2OH ; and

R^{10} is selected from the group consisting of: H, C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, C_{1-10} acyl, C_{8-20} arylalkylacyl, C_{3-20} heteroarylalkylacyl, C_{7-20} arylalkyl, C_{8-20} arylalkenyl, C_{8-20} arylalkynyl, C_{2-20} heteroarylalkyl, C_{3-20} heteroarylalkenyl, and C_{3-20}

heteroarylalkynyl, each excluding H optionally substituted from one to four substituents with one of more of halo, OH, OCF₃, CN, SO₂NH₂, SO₂Me, C(O)NH₂, CH₂F, CHF₂, CF₃, CH₂CH₂F, CH₂CF₃, CH₂CH₂CH₂F, C₁₋₆ alkyl, C₁₋₆ alkoxy, or C₃₋₇ cycloalkyl, in the preparation of a medicament.

- 5 23. The use of claim 22 wherein said medicament is for selectively inhibiting an O-GlcNAcase, for increasing the level of O-GlcNAc, for treating a condition modulated by an O-GlcNAcase, or for treating a neurodegenerative disease, a tauopathy, a cancer, or stress.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/CA2013/050668

A. CLASSIFICATION OF SUBJECT MATTER IPC: C07D 211/60 (2006.01) , A61K 31/445 (2006.01) , A61P 25/28 (2006.01) , C07H 5/04 (2006.01) , C07H 7/00 (2006.01) , C12N 9/24 (2006.01) According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED		
Minimum documentation searched (classification system followed by classification symbols) IPC: C07D 211/60 (2006.01), A61K 31/445 (2006.01), A61P 25/28 (2006.01), C07H 5/04 (2006.01), C07H 7/00 (2006.01), C12N 9/24 (2006.01)		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic database(s) consulted during the international search (name of database(s) and, where practicable, search terms used) STN CAPlus (structure search), Canadian Patent Database (IPC + keywords = piperidine, glycosidase, overexpression, neurodegenerative, O-GlcNAc*, tauopathy, cancer, and stress)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2006/037069 A1 (SCHWEIGHOFFER et al.) 6 April 2006 (06-04-2006) *Abstract, Pages 16-17, & Claims*	1-10, 22-23
X	WO 2010/015816 A2 (WILSON et al.) 11 February 2010 (11-02-2010) *Table 1, compounds 216, 223, 246, 257, 264, 269, 752, 797, 806, 890, 891; Claims*	1-10
X	WO 2010/049678 A2 (WILSON et al.) 6 May 2010 (06-05-2010) *Table1 & Claims*	1-10
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input checked="" type="checkbox"/> See patent family annex.		
* Special categories of cited documents : "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier application or patent but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search 30 October 2013 (30-10-2013)		Date of mailing of the international search report 31 October 2013 (31-10-2013)
Name and mailing address of the ISA/CA Canadian Intellectual Property Office Place du Portage I, C114 - 1st Floor, Box PCT 50 Victoria Street Gatineau, Quebec K1A 0C9 Facsimile No.: 001-819-953-2476		Authorized officer Tung Siu (819) 934-6735

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of the first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons :

1. Claim Nos. : 11-21
because they relate to subject matter not required to be searched by this Authority, namely :

Claims 11-21 are directed to a method for treatment of the human or animal body by surgery or therapy which the International Search Authority is not required to search. However, this Authority has carried out a search based on the alleged effects or purposes/uses of the product defined in claims 11-21.
2. Claim Nos. :
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically :
3. Claim Nos. :
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows :

1. As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.
3. As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claim Nos. :
4. No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claim Nos. :

- Remark on Protest** The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT
Information on patent family members

International application No.
PCT/CA2013/050668

Patent Document Cited in Search Report	Publication Date	Patent Family Member(s)	Publication Date
WO2006037069A1	06 April 2006 (06-04-2006)	US2008085920A1 US7884115B2	10 April 2008 (10-04-2008) 08 February 2011 (08-02-2011)
WO2010015816A2	11 February 2010 (11-02-2010)	EP2323652A2 GB0814322D0 GB0817446D0 GB0817859D0 GB0819523D0 GB0819543D0 GB0906175D0 GB0906179D0 GB0908661D0 GB0908666D0 US2011237538A1 WO2010015816A3	25 May 2011 (25-05-2011) 10 September 2008 (10-09-2008) 29 October 2008 (29-10-2008) 05 November 2008 (05-11-2008) 03 December 2008 (03-12-2008) 03 December 2008 (03-12-2008) 20 May 2009 (20-05-2009) 20 May 2009 (20-05-2009) 01 July 2009 (01-07-2009) 01 July 2009 (01-07-2009) 29 September 2011 (29-09-2011) 26 August 2010 (26-08-2010)
WO2010049678A2	06 May 2010 (06-05-2010)	GB0819941D0 GB0906161D0 GB0908702D0 GB0914471D0 WO2010049678A3	10 December 2008 (10-12-2008) 20 May 2009 (20-05-2009) 01 July 2009 (01-07-2009) 30 September 2009 (30-09-2009) 26 August 2010 (26-08-2010)