

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 2016254958 B2**

(54) Title
Butyrolactones as herbicides

(51) International Patent Classification(s)
C07D 307/24 (2006.01) **C07D 307/33** (2006.01)
A01N 43/08 (2006.01)

(21) Application No: **2016254958** (22) Date of Filing: **2016.04.19**

(87) WIPO No: **WO16/176082**

(30) Priority Data

(31) Number	(32) Date	(33) Country
62/153,477	2015.04.27	US

(43) Publication Date: **2016.11.03**

(44) Accepted Journal Date: **2020.04.23**

(71) Applicant(s)
FMC Corporation

(72) Inventor(s)
Stevenson, Thomas Martin;Satterfield, Andrew Duncan


(74) Agent / Attorney
Houlihan² Pty Ltd, PO Box 611, BALWYN NORTH, VIC, 3104, AU

(56) Related Art
US 3741989 A



- (51) **International Patent Classification:**
C07D 307/24 (2006.01) *C07D 307/33* (2006.01)
A01N 43/08 (2006.01)
- (21) **International Application Number:** PCT/US2016/028260
- (22) **International Filing Date:** 19 April 2016 (19.04.2016)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:** 62/153,477 27 April 2015 (27.04.2015) US
- (71) **Applicant:** E I DU PONT DE NEMOURS AND COMPANY [US/US]; Chestnut Run Plaza, 974 Centre Road, P.O. Box 2915, Wilmington, Delaware 19805 (US).
- (72) **Inventors:** STEVENSON, Thomas Martin; 103 Iroquois Court, Newark, Delaware 19702 (US). SATTERFIELD, Andrew Duncan; 1524 Robert Burns Court, Bear, Delaware 19701 (US).
- (74) **Agent:** COATS, Reed A.; E.I. du Pont de Nemours and Company Legal Patent Records Center, Chestnut Run Plaza 721/2340, 974 Centre Road PO Box 2915, Wilmington, Delaware 19805 (US).
- (81) **Designated States** (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- (84) **Designated States** (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).
- Published:**
— with international search report (Art. 21(3))

(54) **Title:** BUTYROLACTONES AS HERBICIDES



(1)

(57) **Abstract:** Disclosed are compounds of Formula 1, including all stereoisomers, *N*-oxides, and salts thereof, wherein R¹, R², R³, R⁴, R⁵, Q¹, Q², Y¹, and Y² are as defined in the disclosure. Also disclosed are compositions containing the compounds of Formula 1 and methods for controlling undesired vegetation comprising contacting the undesired vegetation or its environment with an effective amount of a compound or a composition of the invention.

TITLE

BUTYROLACTONES AS HERBICIDES

FIELD OF THE INVENTION


This invention relates to certain butyrolactones, their *N*-oxides, salts and compositions,
5 and methods of their use for controlling undesirable vegetation.

BACKGROUND OF THE INVENTION

The control of undesired vegetation is extremely important in achieving high crop
efficiency. Achievement of selective control of the growth of weeds especially in such
useful crops as rice, soybean, sugar beet, maize, potato, wheat, barley, tomato and plantation
10 crops, among others, is very desirable. Unchecked weed growth in such useful crops can
cause significant reduction in productivity and thereby result in increased costs to the
consumer. The control of undesired vegetation in noncrop areas is also important. Many
products are commercially available for these purposes, but the need continues for new
compounds that are more effective, less costly, less toxic, environmentally safer or have
15 different sites of action.

SUMMARY OF THE INVENTION

This invention is directed to a compound of Formula 1 (including all stereoisomers),
including *N*-oxides and salts thereof, agricultural compositions containing them and their use
as herbicides:



1

20

wherein

Q^1 is a phenyl ring or a naphthalenyl ring system, each ring or ring system optionally
substituted with up to 5 substituents independently selected from R^7 ; or a 5- to
6-membered fully unsaturated heterocyclic ring or an 8- to 10-membered
25 heteroaromatic bicyclic ring system, each ring or ring system containing ring
members selected from carbon atoms and 1 to 4 heteroatoms independently
selected from up to 2 O, up to 2 S and up to 4 N atoms, wherein up to 3 carbon
ring members are independently selected from $C(=O)$ and $C(=S)$, and the sulfur
atom ring members are independently selected from $S(=O)_u(=NR^8)_v$, each ring
30 or ring system optionally substituted with up to 5 substituents independently

selected from R⁷ on carbon atom ring members and selected from R⁹ on nitrogen atom ring members;

Q² is a phenyl ring or a naphthalenyl ring system, each ring or ring system optionally substituted with up to 5 substituents independently selected from R¹⁰; or a 5- to 6-membered fully unsaturated heterocyclic ring or an 8- to 10-membered heteroaromatic bicyclic ring system, each ring or ring system containing ring members selected from carbon atoms and 1 to 4 heteroatoms independently selected from up to 2 O, up to 2 S and up to 4 N atoms, wherein up to 3 carbon ring members are independently selected from C(=O) and C(=S), and the sulfur atom ring members are independently selected from S(=O)_u(=NR⁸)_v, each ring or ring system optionally substituted with up to 8 substituents independently selected from R¹⁰ on carbon atom ring members and selected from R¹¹ on nitrogen atom ring members;

Y¹ and Y² are each independently O, S or NR⁶;

R¹ is H, hydroxy, amino, C₁-C₆ alkyl, C₁-C₆ haloalkyl, C₂-C₆ alkenyl, C₃-C₆ alkynyl, C₂-C₈ alkoxyalkyl, C₂-C₈ haloalkoxyalkyl, C₂-C₈ alkylthioalkyl, C₂-C₈ alkylsulfinylalkyl, C₂-C₈ alkylsulfonylalkyl, C₂-C₈ alkylcarbonyl, C₂-C₈ haloalkylcarbonyl, C₄-C₁₀ cycloalkylcarbonyl, C₂-C₈ alkoxycarbonyl, C₂-C₈ haloalkoxycarbonyl, C₄-C₁₀ cycloalkoxycarbonyl, C₂-C₈ alkylaminocarbonyl, C₃-C₁₀ dialkylaminocarbonyl, C₄-C₁₀ cycloalkylaminocarbonyl, C₁-C₆ alkoxy, C₁-C₆ alkylthio, C₁-C₆ haloalkylthio, C₃-C₈ cycloalkylthio, C₁-C₆ alkylsulfinyl, C₁-C₆ haloalkylsulfinyl, C₃-C₈ cycloalkylsulfinyl, C₁-C₆ alkylsulfonyl, C₁-C₆ haloalkylsulfonyl, C₃-C₈ cycloalkylsulfonyl, C₁-C₆ alkylaminosulfonyl, C₂-C₈ dialkylaminosulfonyl, C₃-C₁₀ trialkylsilyl or G¹;

R² and R³ are each independently H, halogen, hydroxy, C₁-C₄ alkyl, C₁-C₄ haloalkyl or C₁-C₄ alkoxy; or

R² and R³ are taken together with the carbon atom to which they are bonded to form a C₃-C₇ cycloalkyl ring;

R⁴ and R⁵ are each independently H, halogen, hydroxy, C₁-C₄ alkyl or C₁-C₄ alkoxy; each R⁶ is independently H, cyano, C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₁-C₄ alkoxy,

C₁-C₄ haloalkoxy, -(C=O)CH₃ or -(C=O)CF₃;

each R⁸ is independently H, cyano, C₂-C₃ alkylcarbonyl or C₂-C₃ haloalkylcarbonyl;

each R⁷ and R¹⁰ is independently halogen, cyano, nitro, C₁-C₈ alkyl, C₁-C₄

cyanoalkyl, C₁-C₄ cyanoalkoxy, C₁-C₈ haloalkyl, C₁-C₈ nitroalkyl, C₂-C₈ alkenyl, C₂-C₈ haloalkenyl, C₂-C₈ nitroalkenyl, C₂-C₈ alkynyl, C₂-C₈ haloalkynyl, C₄-C₁₀ cycloalkylalkyl, C₄-C₁₀ halocycloalkylalkyl, C₅-C₁₂ alkylcycloalkylalkyl, C₅-C₁₂ cycloalkylalkenyl, C₅-C₁₂ cycloalkylalkynyl,

- C₃-C₈ cycloalkyl, C₃-C₈ halocycloalkyl, C₄-C₁₀ alkylcycloalkyl, C₆-C₁₂
 cycloalkylcycloalkyl, C₃-C₈ cycloalkenyl, C₃-C₈ halocycloalkenyl, C₂-C₈
 alkoxyalkyl, C₂-C₈ haloalkoxyalkyl, C₄-C₁₀ cycloalkoxyalkyl, C₃-C₁₀
 alkoxyalkoxyalkyl, C₂-C₈ alkylthioalkyl, C₂-C₈ alkylsulfinylalkyl, C₂-C₈
 alkylsulfonylalkyl, C₂-C₈ alkylaminoalkyl, C₂-C₈ haloalkylaminoalkyl, C₄-C₁₀
 cycloalkylaminoalkyl, C₃-C₁₀ dialkylaminoalkyl, -CHO, C₂-C₈ alkylcarbonyl,
 C₂-C₈ haloalkylcarbonyl, C₄-C₁₀ cycloalkylcarbonyl, -C(=O)OH, C₂-C₈
 alkoxycarbonyl, C₂-C₈ haloalkoxycarbonyl, C₄-C₁₀ cycloalkoxycarbonyl,
 C₅-C₁₂ cycloalkylalkoxycarbonyl, -C(=O)NH₂, C₂-C₈ alkylaminocarbonyl,
 C₄-C₁₀ cycloalkylaminocarbonyl, C₃-C₁₀ dialkylaminocarbonyl, C₁-C₈ alkoxy,
 C₁-C₈ haloalkoxy, C₂-C₈ alkoxyalkoxy, C₂-C₈ haloalkoxyalkoxy, C₂-C₈
 alkenyloxy, C₂-C₈ haloalkenyloxy, C₃-C₈ alkynyloxy, C₃-C₈ haloalkynyloxy,
 C₃-C₈ cycloalkoxy, C₃-C₈ halocycloalkoxy, C₄-C₁₀ cycloalkylalkoxy, C₃-C₁₀
 alkylcarbonylalkoxy, C₂-C₈ alkylcarbonyloxy, C₂-C₈ haloalkylcarbonyloxy,
 C₄-C₁₀ cycloalkylcarbonyloxy, C₁-C₈ alkylsulfonyloxy, C₁-C₈
 haloalkylsulfonyloxy, C₁-C₈ alkylthio, C₁-C₈ haloalkylthio, C₃-C₈
 cycloalkylthio, C₁-C₈ alkylsulfinyl, C₁-C₈ haloalkylsulfinyl, C₁-C₈
 alkylsulfonyl, C₁-C₈ haloalkylsulfonyl, C₃-C₈ cycloalkylsulfonyl, formylamino,
 C₂-C₈ alkylcarbonylamino, C₂-C₈ haloalkylcarbonylamino, C₂-C₈
 alkoxycarbonylamino, C₁-C₆ alkylsulfonylamino, C₁-C₆
 haloalkylsulfonylamino, -SF₅, -SCN, SO₂NH₂, C₃-C₁₂ trialkylsilyl, C₄-C₁₂
 trialkylsilylalkyl, C₄-C₁₂ trialkylsilylalkoxy or G²; or
 two adjacent R⁷ are taken together along with the carbon atoms to which they are
 bonded to form a C₃-C₇ cycloalkyl ring; or
 two adjacent R¹⁰ are taken together along with the carbon atoms to which they are
 bonded to form a C₃-C₇ cycloalkyl ring;
 each R⁹ and R¹¹ is independently cyano, C₁-C₃ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl,
 C₃-C₆ cycloalkyl, C₂-C₃ alkoxyalkyl, C₁-C₃ alkoxy, C₂-C₃ alkylcarbonyl,
 C₂-C₃ alkoxycarbonyl, C₂-C₃ alkylaminoalkyl or C₃-C₄ dialkylaminoalkyl;
 each G¹ is independently phenyl, phenylmethyl (i.e. benzyl), pyridinylmethyl,
 phenylcarbonyl (i.e. benzoyl), phenylcarbonyl(C₁-C₄ alkyl), phenoxy,
 phenylethynyl, phenylsulfonyl, or a 5- or 6-membered heteroaromatic ring, each
 optionally substituted on ring members with up to 5 substituents independently
 selected from R¹²;
 each G² is independently phenyl, phenylmethyl (i.e. benzyl), pyridinylmethyl,
 phenylcarbonyl (i.e. benzoyl), phenylcarbonyl(C₁-C₄ alkyl), phenoxy,
 phenylethynyl, phenylsulfonyl, pyridinyloxy, or a 5- or 6-membered

heteroaromatic ring, each optionally substituted on ring members with up to 5 substituents independently selected from R¹³;

each R¹² and R¹³ is independently halogen, cyano, hydroxy, amino, nitro, -CHO, -C(=O)OH, -C(=O)NH₂, -SO₂NH₂, C₁-C₆ alkyl, C₁-C₆ haloalkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₂-C₈ alkylcarbonyl, C₂-C₈ haloalkylcarbonyl, C₂-C₈ alkoxy, C₄-C₁₀ cycloalkoxy, C₅-C₁₂ cycloalkylalkoxy, C₂-C₈ alkylaminocarbonyl, C₃-C₁₀ dialkylaminocarbonyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, C₂-C₈ alkylcarbonyloxy, C₁-C₆ alkylthio, C₁-C₆ haloalkylthio, C₁-C₆ alkylsulfinyl, C₁-C₆ haloalkylsulfinyl, C₁-C₆ alkylsulfonyl, C₁-C₆ haloalkylsulfonyl, C₁-C₆ alkylaminosulfonyl, C₂-C₈ dialkylaminosulfonyl, C₃-C₁₀ trialkylsilyl, C₁-C₆ alkylamino, C₂-C₈ dialkylamino, C₂-C₈ alkylcarbonylamino, C₁-C₆ alkylsulfonylamino, phenyl, pyridinyl or thienyl; and

each u and v are independently 0, 1 or 2 in each instance of S(=O)_u(=NR⁸)_v, provided that the sum of u and v is 0, 1 or 2.

More particularly, this invention pertains to a compound of Formula 1 (including all stereoisomers), an *N*-oxide or a salt thereof. This invention also relates to a herbicidal composition comprising a compound of the invention (i.e. in a herbicidally effective amount) and at least one component selected from the group consisting of surfactants, solid diluents and liquid diluents. This invention further relates to a method for controlling the growth of undesired vegetation comprising contacting the vegetation or its environment with a herbicidally effective amount of a compound of the invention (e.g., as a composition described herein).

This invention also includes a herbicidal mixture comprising (a) a compound selected from Formula 1, *N*-oxides, and salts thereof, and (b) at least one additional active ingredient selected from (b1) through (b16); and salts of compounds of (b1) through (b16), as described below.

DETAILS OF THE INVENTION

As used herein, the terms “comprises,” “comprising,” “includes,” “including,” “has,” “having,” “contains,” “containing,” “characterized by” or any other variation thereof, are intended to cover a non-exclusive inclusion, subject to any limitation explicitly indicated. For example, a composition, mixture, process or method that comprises a list of elements is not necessarily limited to only those elements but may include other elements not expressly listed or inherent to such composition, mixture, process or method.

The transitional phrase “consisting of” excludes any element, step, or ingredient not specified. If in the claim, such would close the claim to the inclusion of materials other than those recited except for impurities ordinarily associated therewith. When the phrase

“consisting of” appears in a clause of the body of a claim, rather than immediately following the preamble, it limits only the element set forth in that clause; other elements are not excluded from the claim as a whole.

The transitional phrase “consisting essentially of” is used to define a composition or method that includes materials, steps, features, components, or elements, in addition to those literally disclosed, provided that these additional materials, steps, features, components, or elements do not materially affect the basic and novel characteristic(s) of the claimed invention. The term “consisting essentially of” occupies a middle ground between “comprising” and “consisting of”.

Where applicants have defined an invention or a portion thereof with an open-ended term such as “comprising,” it should be readily understood that (unless otherwise stated) the description should be interpreted to also describe such an invention using the terms “consisting essentially of” or “consisting of.”

Further, unless expressly stated to the contrary, “or” refers to an inclusive or and not to an exclusive or. For example, a condition A or B is satisfied by any one of the following: A is true (or present) and B is false (or not present), A is false (or not present) and B is true (or present), and both A and B are true (or present).

Also, the indefinite articles “a” and “an” preceding an element or component of the invention are intended to be nonrestrictive regarding the number of instances (i.e. occurrences) of the element or component. Therefore “a” or “an” should be read to include one or at least one, and the singular word form of the element or component also includes the plural unless the number is obviously meant to be singular.

As referred to herein, the term “seedling”, used either alone or in a combination of words means a young plant developing from the embryo of a seed.

As referred to herein, the term “broadleaf” used either alone or in words such as “broadleaf weed” means dicot or dicotyledon, a term used to describe a group of angiosperms characterized by embryos having two cotyledons.

In the above recitations, the term “alkyl”, used either alone or in compound words such as “alkylthio” or “haloalkyl” includes straight-chain or branched alkyl, such as, methyl, ethyl, *n*-propyl, *i*-propyl, or the different butyl, pentyl or hexyl isomers. “Alkenyl” includes straight-chain or branched alkenes such as ethenyl, 1-propenyl, 2-propenyl, and the different butenyl, pentenyl and hexenyl isomers. “Alkenyl” also includes polyenes such as 1,2-propadienyl and 2,4-hexadienyl. “Alkynyl” includes straight-chain or branched alkynes such as ethynyl, 1-propynyl, 2-propynyl and the different butynyl, pentynyl and hexynyl isomers. “Alkynyl” can also include moieties comprised of multiple triple bonds such as 2,5-hexadiynyl.

“Alkoxy” includes, for example, methoxy, ethoxy, *n*-propyloxy, isopropyloxy and the different butoxy, pentyloxy and hexyloxy isomers. “Alkoxyalkyl” denotes alkoxy substitution

on alkyl. Examples of "alkoxyalkyl" include CH_3OCH_2 , $\text{CH}_3\text{OCH}_2\text{CH}_2$, $\text{CH}_3\text{CH}_2\text{OCH}_2$, $\text{CH}_3\text{CH}_2\text{CH}_2\text{OCH}_2$ and $\text{CH}_3\text{CH}_2\text{OCH}_2\text{CH}_2$. "Alkoxyalkoxy" denotes alkoxy substitution on alkoxy. "Alkenyloxy" includes straight-chain or branched alkenyloxy moieties. Examples of "alkenyloxy" include $\text{H}_2\text{C}=\text{CHCH}_2\text{O}$, $(\text{CH}_3)_2\text{C}=\text{CHCH}_2\text{O}$, $(\text{CH}_3)\text{CH}=\text{CHCH}_2\text{O}$, $(\text{CH}_3)\text{CH}=\text{C}(\text{CH}_3)\text{CH}_2\text{O}$ and $\text{CH}_2=\text{CHCH}_2\text{CH}_2\text{O}$. "Alkynyloxy" includes straight-chain or branched alkynyloxy moieties. Examples of "alkynyloxy" include $\text{HC}\equiv\text{CCH}_2\text{O}$, $\text{CH}_3\text{C}\equiv\text{CCH}_2\text{O}$ and $\text{CH}_3\text{C}\equiv\text{CCH}_2\text{CH}_2\text{O}$. "Alkylthio" includes branched or straight-chain alkylthio moieties such as methylthio, ethylthio, and the different propylthio, butylthio, pentylthio and hexylthio isomers. "Alkylsulfinyl" includes both enantiomers of an alkylsulfinyl group. Examples of "alkylsulfinyl" include $\text{CH}_3\text{S}(\text{O})-$, $\text{CH}_3\text{CH}_2\text{S}(\text{O})-$, $\text{CH}_3\text{CH}_2\text{CH}_2\text{S}(\text{O})-$, $(\text{CH}_3)_2\text{CHS}(\text{O})-$ and the different butylsulfinyl, pentylsulfinyl and hexylsulfinyl isomers. Examples of "alkylsulfonyl" include $\text{CH}_3\text{S}(\text{O})_2-$, $\text{CH}_3\text{CH}_2\text{S}(\text{O})_2-$, $\text{CH}_3\text{CH}_2\text{CH}_2\text{S}(\text{O})_2-$, $(\text{CH}_3)_2\text{CHS}(\text{O})_2-$, and the different butylsulfonyl, pentylsulfonyl and hexylsulfonyl isomers. "Alkylthioalkyl" denotes alkylthio substitution on alkyl. Examples of "alkylthioalkyl" include CH_3SCH_2 , $\text{CH}_3\text{SCH}_2\text{CH}_2$, $\text{CH}_3\text{CH}_2\text{SCH}_2$, $\text{CH}_3\text{CH}_2\text{CH}_2\text{SCH}_2$ and $\text{CH}_3\text{CH}_2\text{SCH}_2\text{CH}_2$. "Cyanoalkyl" denotes an alkyl group substituted with one cyano group. Examples of "cyanoalkyl" include NCCH_2 , NCCH_2CH_2 and $\text{CH}_3\text{CH}(\text{CN})\text{CH}_2$. "Alkylamino", "dialkylamino", and the like, are defined analogously to the above examples. The terms "alkylthioalkyl", "alkylsulfinylamino", "alkylsulfonylamino", "alkylaminosulfonyl", "alkylsulfonylamino", "alkylaminoalkyl", "alkylthioalkyl", "alkylsulfinylalkyl", "alkylsulfonylalkyl", "dialkylaminoalkyl" are defined likewise.

"Cycloalkyl" includes, for example, cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl. The term "alkylcycloalkyl" denotes alkyl substitution on a cycloalkyl moiety and includes, for example, ethylcyclopropyl, *i*-propylcyclobutyl, 3-methylcyclopentyl and 4-methylcyclohexyl. The term "cycloalkylalkyl" denotes cycloalkyl substitution on an alkyl moiety. The term "cycloalkylalkenyl" denotes cycloalkyl substitution on an alkenyl moiety. The term "cycloalkylalkynyl" denotes cycloalkyl substitution on an alkynyl moiety. Examples of "cycloalkylalkyl" include cyclopropylmethyl, cyclopentylethyl, and other cycloalkyl moieties bonded to straight-chain or branched alkyl groups. The term "cycloalkoxy" denotes cycloalkyl linked through an oxygen atom such as cyclopentyloxy and cyclohexyloxy. "Cycloalkylalkoxy" denotes cycloalkylalkyl linked through an oxygen atom attached to the alkyl chain. Examples of "cycloalkylalkoxy" include cyclopropylmethoxy, cyclopentylethoxy, and other cycloalkyl moieties bonded to straight-chain or branched alkoxy groups. "Cycloalkenyl" includes groups such as cyclopentenyl and cyclohexenyl as well as groups with more than one double bond such as 1,3- and 1,4-cyclohexadienyl. The term "cycloalkylcycloalkyl" means a cycloalkyl substitution on a cycloalkyl moiety". The terms "cycloalkoxyalkyl", "alkylcycloalkyl",

“cycloalkylaminoalkyl”, “cycloalkylthio”, “cycloalkylsulfinyl”, “cycloalkylsulfonyl” and the like are defined likewise.

The term “halogen”, either alone or in compound words such as “haloalkyl”, or when used in descriptions such as “alkyl substituted with halogen” includes fluorine, chlorine, bromine or iodine. Further, when used in compound words such as “haloalkyl”, or when used in descriptions such as “alkyl substituted with halogen” said alkyl may be partially or fully substituted with halogen atoms which may be the same or different. Examples of “haloalkyl” or “alkyl substituted with halogen” include F_3C , $ClCH_2$, CF_3CH_2 and CF_3CCl_2 . The terms “halocycloalkyl”, “haloalkoxy”, “haloalkylthio”, “haloalkenyl”, “haloalkynyl”, and the like, are defined analogously to the term “haloalkyl”. Examples of “haloalkoxy” include CF_3O- , CCl_3CH_2O- , $HCF_2CH_2CH_2O-$ and CF_3CH_2O- . Examples of “haloalkylthio” include CCl_3S- , CF_3S- , CCl_3CH_2S- and $ClCH_2CH_2CH_2S-$. Examples of “haloalkylsulfinyl” include $CF_3S(O)-$, $CCl_3S(O)-$, $CF_3CH_2S(O)-$ and $CF_3CF_2S(O)-$. Examples of “haloalkylsulfonyl” include $CF_3S(O)_2-$, $CCl_3S(O)_2-$, $CF_3CH_2S(O)_2-$ and $CF_3CF_2S(O)_2-$. Examples of “haloalkenyl” include $(Cl)_2C=CHCH_2-$ and $CF_3CH_2CH=CHCH_2-$. Examples of “haloalkynyl” include $HC\equiv CCHCl-$, $CF_3C\equiv C-$, $CCl_3C\equiv C-$ and $FCH_2C\equiv CCH_2-$. Examples of “haloalkoxyalkoxy” include CF_3OCH_2O- , $ClCH_2CH_2OCH_2CH_2O-$, $Cl_3CCH_2OCH_2O-$ as well as branched alkyl derivatives. Examples of the “haloalkenyloxy” include $(Cl)_2C=CHCH_2O-$ and $CF_3CH_2CH=CHCH_2O-$. Examples of “haloalkoxyalkyl” include CF_3OCH_2- , $CCl_3CH_2OCH_2-$, $HCF_2CH_2CH_2OCH_2-$ and $CF_3CH_2OCH_2-$.

“Alkylcarbonyl” denotes a straight-chain or branched alkyl moieties bonded to a $C(=O)$ moiety. Examples of “alkylcarbonyl” include $CH_3C(=O)-$, $CH_3CH_2CH_2C(=O)-$ and $(CH_3)_2CHC(=O)-$. Examples of “alkoxycarbonyl” include $CH_3OC(=O)-$, $CH_3CH_2OC(=O)-$, $CH_3CH_2CH_2OC(=O)-$, $(CH_3)_2CHOC(=O)-$ and the different butoxy- or pentoxycarbonyl isomers. The terms “alkylcarbonylalkyl”, “alkylcarbonyloxy”, “alkoxycarbonylamino”, “alkylaminocarbonyl”, “dialkylaminosulfonyl”, “cycloalkylcarbonyl”, “cycloalkoxycarbonyl”, “cycloalkylcarbonyloxy”, “dialkylaminocarbonyl”, “cycloalkylaminocarbonyl”, “haloalkylcarbonyl” and “haloalkoxycarbonyl” are defined likewise. The terms “cyanoalkyl” refers to a cyano group attached to an alkyl group. The terms “cyanoalkoxy” refers to a cyano group attached to an alkoxy group. The terms “nitroalkyl” refers to a nitro group attached to an alkyl group. The terms “nitroalkenyl” refers to a nitro group attached to an alkenyl group.

The term “trialkylsilyl” means silyl substituted with three alkyl groups. The term “trialkylsilylalkyl” means refers to a trialkylsilyl group bonded through an alkyl group (e.g. $-CH_2TMS$). The term “trialkylsilyloxy” means refers to a trialkylsilyl group bonded through oxygen (e.g. $-OTMS$).

The total number of carbon atoms in a substituent group is indicated by the “C_i-C_j” prefix where i and j are numbers from 1 to 12. For example, C₁-C₄ alkylsulfonyl designates methylsulfonyl through butylsulfonyl; C₂ alkoxyalkyl designates CH₃OCH₂-; C₃ alkoxyalkyl designates, for example, CH₃CH(OCH₃)-, CH₃OCH₂CH₂- or CH₃CH₂OCH₂-; and C₄ alkoxyalkyl designates the various isomers of an alkyl group substituted with an alkoxy group containing a total of four carbon atoms, examples including CH₃CH₂CH₂OCH₂- and CH₃CH₂OCH₂CH₂-.

When a compound is substituted with a substituent bearing a subscript that indicates the number of said substituents can exceed 1, said substituents (when they exceed 1) are independently selected from the group of defined substituents, e.g., (R⁷)_n, n is 1, 2, 3, 4 or 5. Further, when the subscript indicates a range, e.g. (R)_{i-j}, then the number of substituents may be selected from the integers between i and j inclusive. When a group contains a substituent which can be hydrogen, for example R¹ or R², then when this substituent is taken as hydrogen, it is recognized that this is equivalent to said group being unsubstituted. When a variable group is shown to be optionally attached to a position, for example (R⁷)_n wherein n may be 0, then hydrogen may be at the position even if not recited in the variable group definition. When one or more positions on a group are said to be “not substituted” or “unsubstituted”, then hydrogen atoms are attached to take up any free valency.

Unless otherwise indicated, a “ring” or “ring system” as a component of Formula 1 (e.g., substituent Q¹) is carbocyclic or heterocyclic. The term “ring system” denotes two or more fused rings. The terms “bicyclic ring system” and “fused bicyclic ring system” denote a ring system consisting of two fused rings, in which either ring can be saturated, partially unsaturated, or fully unsaturated unless otherwise indicated. The term “fused heterobicyclic ring system” denotes a fused bicyclic ring system in which at least one ring atom is not carbon. A “bridged bicyclic ring system” is formed by bonding a segment of one or more atoms to nonadjacent ring members of a ring. The term “ring member” refers to an atom or other moiety (e.g., C(=O), C(=S), S(O) or S(O)₂) forming the backbone of a ring or ring system.

The terms “heterocyclic ring”, “heterocycle” or “heterocyclic ring system” denote a ring or ring system in which at least one atom forming the ring backbone is not carbon, e.g., nitrogen, oxygen or sulfur. Typically a heterocyclic ring contains no more than 4 nitrogens, no more than 2 oxygens and no more than 2 sulfurs. Unless otherwise indicated, a heterocyclic ring can be a saturated, partially unsaturated, or fully unsaturated ring. When a fully unsaturated heterocyclic ring satisfies Hückel’s rule, then said ring is also called a “heteroaromatic ring” or “aromatic heterocyclic ring”. Unless otherwise indicated, heterocyclic rings and ring systems can be attached through any available carbon or nitrogen by replacement of a hydrogen on said carbon or nitrogen.

“Aromatic” indicates that each of the ring atoms is essentially in the same plane and has a *p*-orbital perpendicular to the ring plane, and that $(4n + 2) \pi$ electrons, where *n* is a positive integer, are associated with the ring to comply with Hückel’s rule. The term “aromatic ring system” denotes a carbocyclic or heterocyclic ring system in which at least one ring of the ring system is aromatic. The term “aromatic carbocyclic ring system” denotes a carbocyclic ring system in which at least one ring of the ring system is aromatic. The term “aromatic heterocyclic ring system” denotes a heterocyclic ring system in which at least one ring of the ring system is aromatic. The term “nonaromatic ring system” denotes a carbocyclic or heterocyclic ring system that may be fully saturated, as well as partially or fully unsaturated, provided that none of the rings in the ring system are aromatic. The term “nonaromatic carbocyclic ring system” in which no ring in the ring system is aromatic. The term “nonaromatic heterocyclic ring system” denotes a heterocyclic ring system in which no ring in the ring system is aromatic.


The term “optionally substituted” in connection with the heterocyclic rings refers to groups which are unsubstituted or have at least one non-hydrogen substituent that does not extinguish the biological activity possessed by the unsubstituted analog. As used herein, the following definitions shall apply unless otherwise indicated. The term “optionally substituted” is used interchangeably with the phrase “substituted or unsubstituted” or with the term “(un)substituted.” Unless otherwise indicated, an optionally substituted group may have a substituent at each substitutable position of the group, and each substitution is independent of the other.







When Q^1 or Q^2 is a 5- or 6-membered nitrogen-containing heterocyclic ring, it may be attached to the remainder of Formula 1 through any available carbon or nitrogen ring atom, unless otherwise described. As noted above, Q^1 and Q^2 can be (among others) phenyl optionally substituted with one or more substituents selected from a group of substituents as defined in the Summary of the Invention. An example of phenyl optionally substituted with one to five substituents is the ring illustrated as U-1 in Exhibit 1, wherein R^v is R^7 as defined in the Summary of the Invention for Q^1 , or R^v is R^{10} as defined in the Summary of the Invention for Q^2 , and *r* is an integer from 0 to 5.


As noted above, Q^1 and Q^2 can be (among others) a 5- or 6-membered heterocyclic ring, which may be saturated or unsaturated, optionally substituted with one or more substituents selected from a group of substituents as defined in the Summary of the Invention. Examples of a 5- or 6-membered unsaturated aromatic heterocyclic ring optionally substituted with from one or more substituents include the rings U-2 through U-61 illustrated in Exhibit 1 wherein R^v is any substituent as defined in the Summary of the Invention for Q^1 and Q^2 (e.g. R^7 for Q^1 or R^{10} for Q^2) and *r* is an integer from 0 to 4, limited by the number of available positions on each U group. As U-29, U-30, U-36, U-37, U-38, U-39, U-40, U-41, U-42 and U-43 have only one available position, for these U groups *r* is

limited to the integers 0 or 1, and r being 0 means that the U group is unsubstituted and a hydrogen is present at the position indicated by $(R^V)_r$.

Exhibit 1








U-61

As noted above, Q¹ and Q² can be (among others) an 8-, 9- or 10-membered fused bicyclic ring system optionally substituted with one or more substituents selected from a group of substituents as defined in the Summary of the Invention (e.g. R⁷ for Q¹ or R¹⁰ for Q²). Examples of an 8-, 9- or 10-membered fused bicyclic ring system optionally substituted with from one or more substituents include the rings U-81 through U-123 illustrated in Exhibit 2 wherein R^V is any substituent as defined in the Summary of the Invention for Q¹ or Q² (e.g. R⁷ for Q¹ or R¹⁰ for Q²), and r is typically an integer from 0 to 5.

Exhibit 2




U-62

U-63

U-64

U-65




U-66

U-67

U-68

U-69




U-70

U-71

U-72

U-73




U-74

U-75

U-76

U-77




U-78

U-79

U-80

U-81




U-82

U-83

U-84

U-85




U-86

U-87

U-88

U-89




U-90

U-91

U-92

U-93




U-94

U-95

U-96

U-97



U-98

U-99

U-100


Although R^V groups are shown in the structures U-1 through U-100, it is noted that they do not need to be present since they are optional substituents. Note that when R^V is H when attached to an atom, this is the same as if said atom is unsubstituted. The nitrogen atoms that require substitution to fill their valence are substituted with H or R^V . Note that when the attachment point between $(R^V)_r$ and the U group is illustrated as floating, $(R^V)_r$ can be attached to any available carbon atom or nitrogen atom of the U group. Note that when the attachment point on the U group is illustrated as floating, the U group can be attached to the remainder of Formula 1 through any available carbon or nitrogen of the U group by replacement of a hydrogen atom. Note that some U groups can only be substituted with less than 4 R^V groups (e.g., U-2 through U-5, U-7 through U-48, and U-52 through U-61).

A wide variety of synthetic methods are known in the art to enable preparation of aromatic and nonaromatic heterocyclic rings and ring systems; for extensive reviews see the


eight volume set of *Comprehensive Heterocyclic Chemistry*, A. R. Katritzky and C. W. Rees editors-in-chief, Pergamon Press, Oxford, 1984 and the twelve volume set of *Comprehensive Heterocyclic Chemistry II*, A. R. Katritzky, C. W. Rees and E. F. V. Scriven editors-in-chief, Pergamon Press, Oxford, 1996.

5 Compounds of this invention can exist as one or more stereoisomers. The various stereoisomers include enantiomers, diastereomers, atropisomers and geometric isomers. Stereoisomers are isomers of identical constitution but differing in the arrangement of their atoms in space and include enantiomers, diastereomers, cis-trans isomers (also known as geometric isomers) and atropisomers. Atropisomers result from restricted rotation about
10 single bonds where the rotational barrier is high enough to permit isolation of the isomeric species. One skilled in the art will appreciate that one stereoisomer may be more active and/or may exhibit beneficial effects when enriched relative to the other stereoisomer(s) or when separated from the other stereoisomer(s). Additionally, the skilled artisan knows how to separate, enrich, and/or to selectively prepare said stereoisomers. The compounds of the
15 invention may be present as a mixture of stereoisomers, individual stereoisomers or as an optically active form. Particularly when R⁴ and R⁵ are each H, the C(Y²)N(Q²)(R¹) and Q¹ substituents are typically mostly in the thermodynamically preferred trans configuration on the butyrolactone ring.

For example the C(O)N(Q²)(R¹) moiety (bonded to the carbon at the 3-position of the
20 butyrolactone ring) and Q¹ (bonded to the carbon at the 4-position of the pyrrolidinone ring) are generally found in the *trans* configuration. These two carbon atoms (i.e. at the 3- and 4-positions of the butyrolactone ring of Formula **1**) both possess a chiral center. The two most prevalent pairs of enantiomers are depicted as Formula **1'** and Formula **1''** where the chiral centers are identified (i.e. as 3*S*,4*R* or as 3*R*,4*S*). While this invention pertains to all
25 stereoisomers, the preferred enantiomeric pair for biological operability is identified as Formula **1'** (i.e. the 3*S*,4*R* configuration). The skilled artisan will understand that in some Embodiments of the invention, the R or S designation is determined relative to other substituents around the same carbon and therefore a compound of the invention could also be given the 3*S*,4*S* designation. For a comprehensive discussion of all aspects of
30 stereoisomerism, see Ernest L. Eliel and Samuel H. Wilen, *Stereochemistry of Organic Compounds*, John Wiley & Sons, 1994.




1'



1''

The skilled artisan will also recognize that the carbon atom at the 5-position of the butyrolactone ring (i.e. the carbon atom to which both R² and R³ are bonded) also contains a stereocenter indicated by a (*) as shown in Formula 1''' when R² and R³ are other than the same substituent. This invention pertains to all stereoisomers, and therefore, when R² and R³ are other than the same substituent, then a mixture of diastereomers is possible.



1'''

Molecular depictions drawn herein follow standard conventions for depicting stereochemistry. To indicate stereoconfiguration, bonds rising from the plane of the drawing and towards the viewer are denoted by solid wedges wherein the broad end of the wedge is attached to the atom rising from the plane of the drawing towards the viewer. Bonds going below the plane of the drawing and away from the viewer are denoted by dashed wedges wherein the narrow end of the wedge is attached to the atom further away from the viewer. Constant width lines indicate bonds with a direction opposite or neutral relative to bonds shown with solid or dashed wedges; constant width lines also depict bonds in molecules or parts of molecules in which no particular stereoconfiguration is intended to be specified.

This invention comprises racemic mixtures, for example, equal amounts of the enantiomers of Formulae 1' and 1''. In addition, this invention includes compounds that are enriched compared to the racemic mixture in an enantiomer of Formula 1. Also included are the essentially pure enantiomers of compounds of Formula 1, for example, Formula 1' and Formula 1''.

When enantiomerically enriched, one enantiomer is present in greater amounts than the other, and the extent of enrichment can be defined by an expression of enantiomeric excess

("ee"), which is defined as $(2x-1) \cdot 100\%$, where x is the mole fraction of the dominant enantiomer in the mixture (e.g., an ee of 20 % corresponds to a 60:40 ratio of enantiomers).

Preferably the compositions of this invention have at least a 50 % enantiomeric excess; more preferably at least a 75 % enantiomeric excess; still more preferably at least a 90 % enantiomeric excess; and the most preferably at least a 94 % enantiomeric excess of the more active isomer. Of particular note are enantiomerically pure embodiments of the more active isomer.

Compounds of Formula 1 can comprise additional chiral centers. For example, substituents and other molecular constituents such as R^2 and R^3 may themselves contain chiral centers. This invention comprises racemic mixtures as well as enriched and essentially pure stereoconfigurations at these additional chiral centers.

Compounds of this invention can exist as one or more conformational isomers due to restricted rotation about the amide bond $C(Y^2)N(Q^2)(R^1)$ in Formula 1. This invention comprises mixtures of conformational isomers. In addition, this invention includes compounds that are enriched in one conformer relative to others.

Compounds of Formula 1 typically exist in more than one form, and Formula 1 thus includes all crystalline and non-crystalline forms of the compounds they represent. Non-crystalline forms include embodiments which are solids such as waxes and gums as well as embodiments which are liquids such as solutions and melts. Crystalline forms include embodiments which represent essentially a single crystal type and embodiments which represent a mixture of polymorphs (i.e. different crystalline types). The term "polymorph" refers to a particular crystalline form of a chemical compound that can crystallize in different crystalline forms, these forms having different arrangements and/or conformations of the molecules in the crystal lattice. Although polymorphs can have the same chemical composition, they can also differ in composition due the presence or absence of co-crystallized water or other molecules, which can be weakly or strongly bound in the lattice. Polymorphs can differ in such chemical, physical and biological properties as crystal shape, density, hardness, color, chemical stability, melting point, hygroscopicity, suspensibility, dissolution rate and biological availability. One skilled in the art will appreciate that a polymorph of a compound of Formula 1 can exhibit beneficial effects (e.g., suitability for preparation of useful formulations, improved biological performance) relative to another polymorph or a mixture of polymorphs of the same compound of Formula 1. Preparation and isolation of a particular polymorph of a compound of Formula 1 can be achieved by methods known to those skilled in the art including, for example, crystallization using selected solvents and temperatures. For a comprehensive discussion of polymorphism see R. Hilfiker, Ed., *Polymorphism in the Pharmaceutical Industry*, Wiley-VCH, Weinheim, 2006.

One skilled in the art will appreciate that not all nitrogen-containing heterocycles can form *N*-oxides since the nitrogen requires an available lone pair for oxidation to the oxide;

one skilled in the art will recognize those nitrogen-containing heterocycles which can form *N*-oxides. One skilled in the art will also recognize that tertiary amines can form *N*-oxides. Synthetic methods for the preparation of *N*-oxides of heterocycles and tertiary amines are very well known by one skilled in the art including the oxidation of heterocycles and tertiary amines with peroxy acids such as peracetic and *m*-chloroperbenzoic acid (MCPBA), hydrogen peroxide, alkyl hydroperoxides such as *t*-butyl hydroperoxide, sodium perborate, and dioxiranes such as dimethyldioxirane. These methods for the preparation of *N*-oxides have been extensively described and reviewed in the literature, see for example: T. L. Gilchrist in *Comprehensive Organic Synthesis*, vol. 7, pp 748–750, S. V. Ley, Ed., Pergamon Press; M. Tisler and B. Stanovnik in *Comprehensive Heterocyclic Chemistry*, vol. 3, pp 18–20, A. J. Boulton and A. McKillop, Eds., Pergamon Press; M. R. Grimmett and B. R. T. Keene in *Advances in Heterocyclic Chemistry*, vol. 43, pp 149–161, A. R. Katritzky, Ed., Academic Press; M. Tisler and B. Stanovnik in *Advances in Heterocyclic Chemistry*, vol. 9, pp 285–291, A. R. Katritzky and A. J. Boulton, Eds., Academic Press; and G. W. H. Cheeseman and E. S. G. Werstiuk in *Advances in Heterocyclic Chemistry*, vol. 22, pp 390–392, A. R. Katritzky and A. J. Boulton, Eds., Academic Press.

One skilled in the art recognizes that because in the environment and under physiological conditions salts of chemical compounds are in equilibrium with their corresponding nonsalt forms, salts share the biological utility of the nonsalt forms. Thus a wide variety of salts of a compound of Formula 1 are useful for control of undesired vegetation (i.e. are agriculturally suitable). The salts of a compound of Formula 1 include acid-addition salts with inorganic or organic acids such as hydrobromic, hydrochloric, nitric, phosphoric, sulfuric, acetic, butyric, fumaric, lactic, maleic, malonic, oxalic, propionic, salicylic, tartaric, 4-toluenesulfonic or valeric acids. When a compound of Formula 1 contains an acidic moiety such as a carboxylic acid or phenol, salts also include those formed with organic or inorganic bases such as pyridine, triethylamine or ammonia, or amides, hydrides, hydroxides or carbonates of sodium, potassium, lithium, calcium, magnesium or barium. Accordingly, the present invention comprises compounds selected from Formula 1, *N*-oxides and agriculturally suitable salts thereof.

Embodiments of the present invention as described in the Summary of the Invention include (where Formula 1 as used in the following Embodiments includes *N*-oxides and salts thereof):

Embodiment 1. A compound of Formula 1 (including all stereoisomers), *N*-oxides, and salts thereof, agricultural compositions containing them and their use as herbicides as described in the Summary of the Invention.

Embodiment 2. A compound of Formula 1 wherein when Q¹ is an 8- to 10-membered heteroaromatic bicyclic ring system optionally substituted with R⁷ and R⁹, the

remainder of Formula 1 is bonded to a fully unsaturated ring of said bicyclic ring system.

Embodiment 3. A compound of Formula 1 wherein Q¹ is a phenyl ring optionally substituted with up to 5 substituents independently selected from R⁷.

5 Embodiment 4. A compound of Embodiment 3 wherein Q¹ is a phenyl ring substituted with 1 to 3 substituents independently selected from R⁷.

Embodiment 5. A compound of Embodiment 4 wherein Q¹ is a phenyl ring substituted with 1 to 2 substituents independently selected from R⁷.

10 Embodiment 6. A compound of Formula 1 or any one of Embodiments 3 through 5 wherein Q¹ is a phenyl ring having a substituent selected from R⁷ at the para (4-) position (and optionally other substituents).

15 Embodiment 7. A compound of Formula 1 or any one of Embodiments 3 through 6 wherein when Q¹ is a phenyl ring substituted with at least two substituents selected from R⁷, then one substituent is at the para (4-) position and at least one other substituent is at a meta position (of the phenyl ring).

Embodiment 8. A compound of Formula 1 or any one of Embodiments 2 through 7 wherein when Q² is an 8- to 10-membered heteroaromatic bicyclic ring system optionally substituted with R¹⁰ and R¹¹, the remainder of Formula 1 is bonded to a fully unsaturated ring of said bicyclic ring system.

20 Embodiment 9. A compound of Formula 1 or any one of Embodiments 2 through 7 wherein Q² is a phenyl ring substituted with up to 5 substituents independently selected from R¹⁰.

Embodiment 10. A compound of Embodiment 9 wherein Q² is a phenyl ring substituted with 1 to 3 substituents independently selected from R¹⁰.

25 Embodiment 11. A compound of Embodiment 10 wherein Q² is a phenyl ring substituted with 1 to 2 substituents independently selected from R¹⁰.

Embodiment 12. A compound of Embodiment 11 wherein Q² is a phenyl ring having at least one substituent selected from R¹⁰ at an ortho (e.g., 2-) position (and optionally other substituents).

30 Embodiment 13. A compound of Formula 1 or any one of Embodiments 2 through 7 or Embodiments 9 through 12 wherein when Q² is a phenyl ring substituted with at least two substituents selected from R¹⁰, then at least one substituent is at an ortho (e.g., 2-) position and at least one substituent is at an adjacent meta (e.g., 3-) position (of the phenyl ring).

35 Embodiment 14. A compound of Formula 1 or any one of Embodiments 2 through 13 wherein, independently, each R⁷ and R¹⁰ is independently halogen, cyano, nitro, C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₂-C₄ alkenyl, C₂-C₄ haloalkenyl, C₂-C₄ alkynyl, C₂-C₄ haloalkynyl, C₁-C₄ nitroalkyl, C₂-C₄ nitroalkenyl, C₂-C₄

alkoxyalkyl, C₂-C₄ haloalkoxyalkyl, C₃-C₄ cycloalkyl, C₃-C₄ halocycloalkyl, cyclopropylmethyl, methylcyclopropyl, C₁-C₄ alkoxy, C₁-C₄ haloalkoxy, C₂-C₄ alkenyloxy, C₂-C₄ haloalkenyloxy, C₃-C₄ alkynyloxy, C₃-C₄ haloalkynyloxy, C₃-C₄ cycloalkoxy, C₁-C₄ alkylthio, C₁-C₄ haloalkylthio, C₁-C₄ alkylsulfinyl, C₁-C₄ haloalkylsulfinyl, C₁-C₄ alkylsulfonyl, C₁-C₄ haloalkylsulfonyl, hydroxy, formyl, C₂-C₄ alkylcarbonyl, C₂-C₄ alkylcarbonyloxy, C₁-C₄ alkylsulfonyloxy, C₁-C₄ haloalkylsulfonyloxy, amino, C₁-C₄ alkylamino, C₂-C₄ dialkylamino, formylamino, C₂-C₄ alkylcarbonylamino, -SF₅, -SCN, C₃-C₄ trialkylsilyl, trimethylsilylmethyl or trimethylsilylmethoxy.

Embodiment 15. A compound of Embodiment 14 wherein each R⁷ is independently halogen, cyano, C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₂-C₄ alkenyl, C₂-C₄ haloalkenyl, C₂-C₄ alkynyl, C₂-C₄ haloalkynyl, C₃-C₄ cycloalkyl, C₃-C₄ halocycloalkyl, cyclopropylmethyl, methylcyclopropyl, C₁-C₄ alkoxy, C₁-C₄ haloalkoxy, C₁-C₄ alkylthio, C₁-C₄ haloalkylthio, C₁-C₄ alkylsulfonyl, C₁-C₄ alkylsulfonyloxy or C₁-C₄ haloalkylsulfonyloxy.

Embodiment 16. A compound of Embodiment 15 wherein each R⁷ is independently halogen, cyano, C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₁-C₄ alkoxy or C₁-C₄ haloalkoxy.

Embodiment 17. A compound of Embodiment 16 wherein each R⁷ is independently halogen, C₁-C₂ alkyl or C₁-C₃ haloalkyl.

Embodiment 18. A compound of Embodiment 17 wherein each R⁷ is independently halogen or C₁-C₂ haloalkyl.

Embodiment 19. A compound of Embodiment 18 wherein each R⁷ is independently halogen or C₁ haloalkyl.

Embodiment 20. A compound of Embodiment 19 wherein each R⁷ is independently halogen or C₁ fluoroalkyl.

Embodiment 21. A compound of Embodiment 20 wherein each R⁷ is independently halogen or CF₃.

Embodiment 22. A compound of Embodiment 21 wherein each R⁷ is independently F, Cl, Br or CF₃.

Embodiment 23. A compound of Embodiment 22 wherein each R⁷ is independently F or CF₃.

Embodiment 24. A compound of any one of Embodiments 21 through 23 wherein at most only one CF₃ substituent is present on the Q¹ phenyl ring and is at the meta position of said phenyl ring.

Embodiment 25. A compound any one of Embodiments 14 through 24 wherein each R¹⁰ is independently halogen, cyano, C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₂-C₄

alkenyl, C₂-C₄ haloalkenyl, C₂-C₄ alkynyl, C₂-C₄ haloalkynyl, C₃-C₄ cycloalkyl, C₃-C₄ halocycloalkyl, cyclopropylmethyl, methylcyclopropyl, C₁-C₄ alkoxy, C₁-C₄ haloalkoxy, C₁-C₄ alkylthio, C₁-C₄ haloalkylthio, C₁-C₄ alkylsulfonyl, C₁-C₄ alkylsulfonyloxy or C₁-C₄ haloalkylsulfonyloxy.

- 5 Embodiment 26. A compound of Embodiment 25 wherein each R¹⁰ is independently halogen, cyano, C₁-C₄ alkyl, C₁-C₄ haloalkyl or C₁-C₄ alkylsulfonyl.
- Embodiment 27. A compound of Embodiment 26 wherein each R¹⁰ is independently halogen, cyano, C₁-C₂ alkyl, C₁-C₃ haloalkyl or C₁-C₃ alkylsulfonyl.
- Embodiment 28. A compound of Embodiment 27 wherein each R¹⁰ is independently
10 halogen, C₁-C₂ haloalkyl or C₁-C₃ alkylsulfonyl.
- Embodiment 29. A compound of Embodiment 28 wherein each R¹⁰ is independently halogen, C₁ haloalkyl or C₁ alkylsulfonyl.
- Embodiment 30. A compound of Embodiment 29 wherein each R¹⁰ is independently halogen or C₁ fluoroalkyl.
- 15 Embodiment 31. A compound of Embodiment 30 wherein each R¹⁰ is independently halogen or CF₃.
- Embodiment 32. A compound of Embodiment 31 wherein each R¹⁰ is independently F, Cl, Br or CF₃.
- Embodiment 33. A compound of Embodiment 32 wherein each R¹⁰ is independently F
20 or CF₃.
- Embodiment 34. A compound of Embodiment 33 wherein each R¹⁰ is F.
- Embodiment 35. A compound of Formula 1 or any one of Embodiments 2 through 34 wherein, independently, each R⁹ and R¹¹ is independently C₁-C₂ alkyl or C₂-C₃ alkoxy carbonyl.
- 25 Embodiment 36. A compound of Embodiment 35 wherein, independently, each R⁹ and R¹¹ is independently C₁-C₂ alkyl.
- Embodiment 37. A compound of Embodiment 36 wherein, independently, each R⁹ and R¹¹ is CH₃.
- Embodiment 38. A compound of Formula 1 or any one of Embodiments 2 through 37
30 wherein Y¹ is O.
- Embodiment 39. A compound of Formula 1 or any one of Embodiments 2 through 38 wherein Y² is O.
- Embodiment 40. A compound of Formula 1 or any one of Embodiments 2 through 39 wherein R² is H or CH₃.
- 35 Embodiment 41. A compound of Embodiment 40 wherein R² is H.
- Embodiment 42. A compound of Formula 1 or any one of Embodiments 2 through 41 wherein R³ is H or CH₃.
- Embodiment 43. A compound of Embodiment 42 wherein R³ is H.

Embodiment 44. A compound of Formula 1 or any one of Embodiments 2 through 44 wherein R⁴ is H or CH₃.

Embodiment 45. A compound of Embodiment 44 wherein R⁴ is H.

Embodiment 46. A compound of Formula 1 or any one of Embodiments 2 through 45
5 wherein R⁵ is H or CH₃.

Embodiment 47. A compound of Embodiment 46 wherein R⁵ is H.

Embodiment 48. A compound of Formula 1 or any one of Embodiments 2 through 47
10 wherein R¹ is H, hydroxy, amino, C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₂-C₄ alkenyl, C₃-C₄ alkynyl, C₂-C₄ alkoxyalkyl, C₂-C₄ haloalkoxyalkyl, C₂-C₄ alkylthioalkyl, C₂-C₄ alkylsulfinylalkyl, C₂-C₄ alkylsulfonylalkyl, C₂-C₄ alkylcarbonyl, C₂-C₄ haloalkylcarbonyl, C₄-C₆ cycloalkylcarbonyl, C₂-C₄ alkoxycarbonyl, C₂-C₄ haloalkoxycarbonyl or C₄-C₆ cycloalkoxycarbonyl.

Embodiment 49. A compound of Embodiment 48 wherein R¹ is H, hydroxy, amino, C₁-C₃ alkyl, C₁-C₃ haloalkyl, C₂-C₄ alkenyl or C₃-C₄ alkynyl.

Embodiment 50. A compound of Embodiment 49 wherein R¹ is H or C₁-C₃ alkyl.

Embodiment 51. A compound of Embodiment 50 wherein R¹ is H or CH₃.

Embodiment 52. A compound of Embodiment 51 wherein R¹ is H.

Embodiment 53. A compound of Formula 1 or any one of Embodiments 2 through 52
20 wherein the stereochemistry is (3*R*,4*S*) or (3*S*,4*R*).

Embodiment 54. A compound of Embodiment 53 wherein the stereochemistry is (3*R*,4*S*).

Embodiment 55. A compound of Embodiment 53 wherein the stereochemistry is (3*S*,4*R*).

Embodiment 56. A compound of Formula 1 or any one of Embodiments 2 through 46
25 wherein R⁵ is CH₃.

Embodiment 57. A compound of Formula 1 wherein Q¹ is a phenyl ring optionally substituted with 1 to 4 substituents independently selected from R⁷; or a 5- to 6-membered heteroaromatic ring containing ring members selected from carbon atoms and 1 to 4 heteroatoms independently selected from up to 2 O, up to 2 S and up to 4 N atoms, optionally substituted with up to 4 substituents independently selected from R⁷ on carbon atom ring members and selected from R⁹ on nitrogen atom ring members.
30

Embodiment 58. A compound of Formula 1 wherein Q² is a phenyl ring optionally substituted with up to 5 substituents independently selected from R¹⁰; or a 5- to 6-membered heteroaromatic ring containing ring members selected from carbon atoms and 1 to 4 heteroatoms independently selected from up to 2 O, up to 2 S and up to 4 N atoms, optionally substituted with up to 5 substituents
35

independently selected from R¹⁰ on carbon atom ring members and selected from R¹¹ on nitrogen atom ring members.

Embodiment 59. A compound of Formula 1 wherein Q¹ is a 5- to 6-membered heteroaromatic ring or an 8- to 10-membered heteroaromatic bicyclic ring system, each ring or ring system optionally substituted with up to 4 substituents independently selected from R⁷ on carbon atom ring members and selected from R⁹ on nitrogen atom ring members.

Embodiments of this invention, including Embodiments 1–59 above as well as any other embodiments described herein, can be combined in any manner, and the descriptions of variables in the embodiments pertain not only to the compounds of Formula 1 but also to the starting compounds and intermediate compounds useful for preparing the compounds of Formula 1. In addition, embodiments of this invention, including Embodiments 1–59 above as well as any other embodiments described herein, and any combination thereof, pertain to the compositions and methods of the present invention.

Combinations of Embodiments 1–59 are illustrated by:

Embodiment A. A compound of Formula 1 wherein

each R⁷ and R¹⁰ is independently halogen, cyano, nitro, C₁–C₄ alkyl, C₁–C₄ haloalkyl, C₂–C₄ alkenyl, C₂–C₄ haloalkenyl, C₂–C₄ alkynyl, C₂–C₄ haloalkynyl, C₁–C₄ nitroalkyl, C₂–C₄ nitroalkenyl, C₂–C₄ alkoxyalkyl, C₂–C₄ haloalkoxyalkyl, C₃–C₄ cycloalkyl, C₃–C₄ halocycloalkyl, cyclopropylmethyl, methylcyclopropyl, C₁–C₄ alkoxy, C₁–C₄ haloalkoxy, C₂–C₄ alkenyloxy, C₂–C₄ haloalkenyloxy, C₃–C₄ alkynyloxy, C₃–C₄ haloalkynyloxy, C₃–C₄ cycloalkoxy, C₁–C₄ alkylthio, C₁–C₄ haloalkylthio, C₁–C₄ alkylsulfinyl, C₁–C₄ haloalkylsulfinyl, C₁–C₄ alkylsulfonyl, C₁–C₄ haloalkylsulfonyl, hydroxy, formyl, C₂–C₄ alkylcarbonyl, C₂–C₄ alkylcarbonyloxy, C₁–C₄ alkylsulfonyloxy, C₁–C₄ haloalkylsulfonyloxy, amino, C₁–C₄ alkylamino, C₂–C₄ dialkylamino, formylamino, C₂–C₄ alkylcarbonylamino, -SF₅, -SCN, C₃–C₄ trialkylsilyl, trimethylsilylmethyl or trimethylsilylmethoxy; and

each R⁹ and R¹¹ is independently C₁–C₂ alkyl or C₂–C₃ alkoxy carbonyl.

Embodiment B. A compound of Embodiment A wherein

Y¹ is O;

Y² is O;

R¹ is H;

R² is H;

R³ is H;

R⁴ is H; and

R⁵ is H.

Embodiment C. A compound of Embodiment B wherein

Q¹ is a phenyl ring substituted with 1 to 3 substituents independently selected from R⁷; and

Q² is a phenyl ring substituted with 1 to 3 substituents independently selected from R¹⁰.

Embodiment D. A compound of Embodiment C wherein

each R⁷ is independently halogen, cyano, C₁-C₄ alkyl, C₁-C₄ haloalkyl C₁-C₄ alkoxy or C₁-C₄ haloalkoxy; and

each R¹⁰ is independently halogen, cyano, C₁-C₄ alkyl, C₁-C₄ haloalkyl or C₁-C₄ alkylsulfonyl.

Embodiment E. A compound of Embodiment D wherein

Q¹ is a phenyl ring substituted with 1 substituent selected from R⁷ at the para position or substituted with 2 substituents independently selected from R⁷ wherein one substituent is at the para position and the other substituent is at a meta position; and

Q² is a phenyl ring substituted with 1 substituent selected from R¹⁰ at an ortho position or substituted with 2 substituents independently selected from R¹⁰ wherein one substituent is at an ortho position and the other substituent is at the adjacent meta position.

Embodiment F. A compound of Embodiment E wherein

each R⁷ is independently F or CF₃; and

each R¹⁰ is F.

Embodiment G. A compound of Embodiment A wherein

Y¹ is O;

Y² is O;

R¹ is H;

R² is H;

R³ is H;

R⁴ is H; and

R⁵ is H or CH₃.

Specific embodiments include a compound of Formula 1 selected from the group consisting of:

4-(4-fluorophenyl)tetrahydro-2-oxo-*N*-[2-(trifluoromethyl)phenyl]-3-furancarboxamide;

(3*R*,4*S*)-4-(4-fluorophenyl)tetrahydro-2-oxo-*N*-[2-(trifluoromethyl)phenyl]-3-furancarboxamide; and

(3*S*,4*R*)-4-(4-fluorophenyl)tetrahydro-2-oxo-*N*-[2-(trifluoromethyl)phenyl]-3-furancarboxamide.

This invention also relates to a method for controlling undesired vegetation comprising applying to the locus of the vegetation herbicidally effective amounts of the compounds of the invention (e.g., as a composition described herein). Of note as embodiments relating to methods of use are those involving the compounds of embodiments described above.

5 Compounds of the invention are particularly useful for selective control of weeds in crops such as wheat, barley, maize, soybean, sunflower, cotton, oilseed rape and rice, and specialty crops such as sugarcane, citrus, fruit and nut crops.

Also noteworthy as embodiments are herbicidal compositions of the present invention comprising the compounds of embodiments described above.

10 This invention also includes a herbicidal mixture comprising (a) a compound selected from Formula 1, *N*-oxides, and salts thereof, and (b) at least one additional active ingredient selected from (b1) photosystem II inhibitors, (b2) acetohydroxy acid synthase (AHAS) inhibitors, (b3) acetyl-CoA carboxylase (ACCase) inhibitors, (b4) auxin mimics, (b5) 5-enol-pyruvylshikimate-3-phosphate (EPSP) synthase inhibitors, (b6) photosystem I
15 electron diverters, (b7) protoporphyrinogen oxidase (PPO) inhibitors, (b8) glutamine synthetase (GS) inhibitors, (b9) very long chain fatty acid (VLCFA) elongase inhibitors, (b10) auxin transport inhibitors, (b11) phytoene desaturase (PDS) inhibitors, (b12) 4-hydroxyphenyl-pyruvate dioxygenase (HPPD) inhibitors, (b13) homogentisate solenesyltransferase (HST) inhibitors, (b14) cellulose biosynthesis inhibitors, (b15) other
20 herbicides including mitotic disruptors, organic arsenicals, asulam, bromobutide, cinmethylin, cumyluron, dazomet, difenzoquat, dymron, etobenzanid, flurenol, fosamine, fosamine-ammonium, hydantocidin, metam, methyldymron, oleic acid, oxaziclomefone, pelargonic acid and pyributicarb, and (b16) herbicide safeners; and salts of compounds of (b1) through (b16).

25 "Photosystem II inhibitors" (b1) are chemical compounds that bind to the D-1 protein at the Q_B -binding niche and thus block electron transport from Q_A to Q_B in the chloroplast thylakoid membranes. The electrons blocked from passing through photosystem II are transferred through a series of reactions to form toxic compounds that disrupt cell membranes and cause chloroplast swelling, membrane leakage, and ultimately cellular
30 destruction. The Q_B -binding niche has three different binding sites: binding site A binds the triazines such as atrazine, triazinones such as hexazinone, and uracils such as bromacil, binding site B binds the phenylureas such as diuron, and binding site C binds benzothiadiazoles such as bentazon, nitriles such as bromoxynil and phenyl-pyridazines such as pyridate. Examples of photosystem II inhibitors include ametryn, amicarbazone, atrazine,
35 bentazon, bromacil, bromofenoxim, bromoxynil, chlorbromuron, chloridazon, chlorotoluron, chloroxuron, cumyluron, cyanazine, daimuron, desmedipham, desmetryn, dimefuron, dimethametryn, diuron, ethidimuron, fenuron, fluometuron, hexazinone, ioxynil, isoproturon, isouron, lenacil, linuron, metamitron, methabenzthiazuron, metobromuron,

metoxuron, metribuzin, monolinuron, neburon, pentanochlor, phenmedipham, prometon, prometryn, propanil, propazine, pyridafol, pyridate, siduron, simazine, simetryn, tebuthiuron, terbacil, terbumeton, terbuthylazine, terbutryn and trietazine. Of note is a compound of the invention mixed with atrazine, bromoxynil or metribuzin.

5 “AHAS inhibitors” (b2) are chemical compounds that inhibit acetohydroxy acid synthase (AHAS), also known as acetolactate synthase (ALS), and thus kill plants by inhibiting the production of the branched-chain aliphatic amino acids such as valine, leucine and isoleucine, which are required for protein synthesis and cell growth. Examples of AHAS inhibitors include amidosulfuron, azimsulfuron, bensulfuron-methyl, 10 bispyribac-sodium, cloransulam-methyl, chlorimuron-ethyl, chlorsulfuron, cinosulfuron, cyclosulfamuron, diclosulam, ethametsulfuron-methyl, ethoxysulfuron, flazasulfuron, florasulam, flucarbazone-sodium, flumetsulam, flupyrsulfuron-methyl, flupyrsulfuron-sodium, foramsulfuron, halosulfuron-methyl, imazamethabenz-methyl, imazamox, imazapic, imazapyr, imazaquin, imazethapyr, imazosulfuron, iodosulfuron-methyl (including sodium 15 salt), iofensulfuron (2-iodo-*N*-[[[(4-methoxy-6-methyl-1,3,5-triazin-2-yl)amino]carbonyl]benzenesulfonamide), mesosulfuron-methyl, metazosulfuron (3-chloro-4-(5,6-dihydro-5-methyl-1,4,2-dioxazin-3-yl)-*N*-[[[(4,6-dimethoxy-2-pyrimidinyl)amino]carbonyl]-1-methyl-1*H*-pyrazole-5-sulfonamide), metosulam, metsulfuron-methyl, nicosulfuron, oxasulfuron, penoxsulam, primisulfuron-methyl, 20 propoxycarbazone-sodium, propyrisulfuron (2-chloro-*N*-[[[(4,6-dimethoxy-2-pyrimidinyl)amino]carbonyl]-6-propylimidazo[1,2-*b*]pyridazine-3-sulfonamide), prosulfuron, pyrazosulfuron-ethyl, pyribenzoxim, pyriftalid, pyriminobac-methyl, pyrithiobac-sodium, rimsulfuron, sulfometuron-methyl, sulfosulfuron, thiencarbazone, thifensulfuron-methyl, triafamone (*N*-[2-[(4,6-dimethoxy-1,3,5-triazin-2-yl)carbonyl]-6-fluorophenyl]-1,1-difluoro-*N*-methylmethanesulfonamide), triasulfuron, tribenuron-methyl, 25 trifloxysulfuron (including sodium salt), triflusulfuron-methyl and tritosulfuron. Of note is a compound of the invention mixed with nicosulfuron, flupyrsulfuron or chlorimuron.

“ACCCase inhibitors” (b3) are chemical compounds that inhibit the acetyl-CoA carboxylase enzyme, which is responsible for catalyzing an early step in lipid and fatty acid 30 synthesis in plants. Lipids are essential components of cell membranes, and without them, new cells cannot be produced. The inhibition of acetyl CoA carboxylase and the subsequent lack of lipid production leads to losses in cell membrane integrity, especially in regions of active growth such as meristems. Eventually shoot and rhizome growth ceases, and shoot meristems and rhizome buds begin to die back. Examples of ACCCase inhibitors include 35 alloxydim, butoxydim, clethodim, clodinafop, cycloxydim, cyhalofop, diclofop, fenoxaprop, fluazifop, haloxyfop, pinoxaden, profoxydim, propaquizafop, quizalofop, sethoxydim, tepraloxym and tralkoxydim, including resolved forms such as fenoxaprop-P, fluazifop-P, haloxyfop-P and quizalofop-P and ester forms such as clodinafop-propargyl,

cyhalofop-butyl, diclofop-methyl and fenoxaprop-P-ethyl. Of note is a compound of the invention mixed with pinoxaden or quizalofop.

Auxin is a plant hormone that regulates growth in many plant tissues. "Auxin mimics" (b4) are chemical compounds mimicking the plant growth hormone auxin, thus causing uncontrolled and disorganized growth leading to plant death in susceptible species. Examples of auxin mimics include aminocyclopyrachlor (6-amino-5-chloro-2-cyclopropyl-4-pyrimidinecarboxylic acid) and its methyl and ethyl esters and its sodium and potassium salts, aminopyralid, benazolin-ethyl, chloramben, clacyfos, clomeprop, clopyralid, dicamba, 2,4-D, 2,4-DB, dichlorprop, fluroxypyr, halauxifen (4-amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxyphenyl)-2-pyridinecarboxylic acid), halauxifen-methyl (methyl 4-amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxyphenyl)-2-pyridinecarboxylate), MCPA, MCPB, mecoprop, picloram, quinclorac, quinmerac, 2,3,6-TBA, triclopyr, and methyl 4-amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxyphenyl)-5-fluoro-2-pyridinecarboxylate. Of note is a compound of the invention mixed with dicamba.

"EPSP synthase inhibitors" (b5) are chemical compounds that inhibit the enzyme, 5-enol-pyruvylshikimate-3-phosphate synthase, which is involved in the synthesis of aromatic amino acids such as tyrosine, tryptophan and phenylalanine. EPSP inhibitor herbicides are readily absorbed through plant foliage and translocated in the phloem to the growing points. Glyphosate is a relatively nonselective postemergence herbicide that belongs to this group. Glyphosate includes esters and salts such as ammonium, isopropylammonium, potassium, sodium (including sesquisodium) and trimesium (alternatively named sulfosate).

"Photosystem I electron diverters" (b6) are chemical compounds that accept electrons from Photosystem I, and after several cycles, generate hydroxyl radicals. These radicals are extremely reactive and readily destroy unsaturated lipids, including membrane fatty acids and chlorophyll. This destroys cell membrane integrity, so that cells and organelles "leak", leading to rapid leaf wilting and desiccation, and eventually to plant death. Examples of this second type of photosynthesis inhibitor include diquat and paraquat.

"PPO inhibitors" (b7) are chemical compounds that inhibit the enzyme protoporphyrinogen oxidase, quickly resulting in formation of highly reactive compounds in plants that rupture cell membranes, causing cell fluids to leak out. Examples of PPO inhibitors include acifluorfen-sodium, azafenidin, benzfendizone, bifenox, butafenacil, carfentrazone, carfentrazone-ethyl, chlomethoxyfen, cinidon-ethyl, fluazolate, flufenpyr-ethyl, flumiclorac-pentyl, flumioxazin, fluoroglycofen-ethyl, fluthiacet-methyl, fomesafen, halosafen, lactofen, oxadiargyl, oxadiazon, oxyfluorfen, pentoxazone, profluzol, pyraclonil, pyraflufen-ethyl, saflufenacil, sulfentrazone, thidiazimin, trifludimoxazin (dihydro-1,5-dimethyl-6-thioxo-3-[2,2,7-trifluoro-3,4-dihydro-3-oxo-4-(2-propyn-1-yl)-2H-1,4-benzoxazin-6-yl]-1,3,5-triazine-2,4(1*H*,3*H*)-dione) and tiafenacil (methyl *N*-[2-[[2-

chloro-5-[3,6-dihydro-3-methyl-2,6-dioxo-4-(trifluoromethyl)-1(2*H*)-pyrimidinyl]-4-fluorophenyl]thio]-1-oxopropyl]- β -alaninate).

“GS inhibitors” (b8) are chemical compounds that inhibit the activity of the glutamine synthetase enzyme, which plants use to convert ammonia into glutamine. Consequently, ammonia accumulates and glutamine levels decrease. Plant damage probably occurs due to the combined effects of ammonia toxicity and deficiency of amino acids required for other metabolic processes. The GS inhibitors include glufosinate and its esters and salts such as glufosinate-ammonium and other phosphinothricin derivatives, glufosinate-P ((2*S*)-2-amino-4-(hydroxymethylphosphinyl)butanoic acid) and bilanaphos.

“VLCFA elongase inhibitors” (b9) are herbicides having a wide variety of chemical structures, which inhibit the elongase. Elongase is one of the enzymes located in or near chloroplasts which are involved in biosynthesis of VLCFAs. In plants, very-long-chain fatty acids are the main constituents of hydrophobic polymers that prevent desiccation at the leaf surface and provide stability to pollen grains. Such herbicides include acetochlor, alachlor, anilofos, butachlor, cafenstrole, dimethachlor, dimethenamid, diphenamid, fenoxasulfone (3-[[[(2,5-dichloro-4-ethoxyphenyl)methyl]sulfonyl]-4,5-dihydro-5,5-dimethylisoxazole), fentrazamide, flufenacet, indanofan, mefenacet, metazachlor, metolachlor, naproanilide, napropamide, napropamide-M ((2*R*)-*N,N*-diethyl-2-(1-naphthalenyloxy)propanamide), pethoxamid, piperophos, pretilachlor, propachlor, propisochlor, pyroxasulfone, and thenylchlor, including resolved forms such as *S*-metolachlor and chloroacetamides and oxyacetamides. Of note is a compound of the invention mixed with flufenacet.

“Auxin transport inhibitors” (b10) are chemical substances that inhibit auxin transport in plants, such as by binding with an auxin-carrier protein. Examples of auxin transport inhibitors include diflufenzopyr, naptalam (also known as *N*-(1-naphthyl)phthalamic acid and 2-[(1-naphthalenylamino)carbonyl]benzoic acid).


“PDS inhibitors” (b11) are chemical compounds that inhibit carotenoid biosynthesis pathway at the phytoene desaturase step. Examples of PDS inhibitors include beflubutamid, diflufenican, fluridone, flurochloridone, flurtamone norflurzon and picolinafen.

“HPPD inhibitors” (b12) are chemical substances that inhibit the biosynthesis of synthesis of 4-hydroxyphenyl-pyruvate dioxygenase. Examples of HPPD inhibitors include benzobicyclon, benzofenap, bicyclopyrone (4-hydroxy-3-[[2-[(2-methoxyethoxy)methyl]-6-(trifluoromethyl)-3-pyridinyl]carbonyl]bicyclo[3.2.1]oct-3-en-2-one), fenquinotriene (2-[[8-chloro-3,4-dihydro-4-(4-methoxyphenyl)-3-oxo-2-quinoxaliny]carbonyl]-1,3-cyclohexanedione), isoxachlortole, isoxaflutole, mesotriene, pyrasulfotole, pyrazolynate, pyrazoxyfen, sulcotriene, tefuryltrione, tembotriene, tolpyralate (1-[[1-ethyl-4-[3-(2-methoxyethoxy)-2-methyl-4-(methylsulfonyl)benzoyl]-1*H*-pyrazol-5-yl]oxy]ethyl methyl carbonate), topramezone, 5-chloro-3-[(2-hydroxy-6-oxo-1-cyclohexen-1-yl)carbonyl]-1-(4-methoxyphenyl)-2(1*H*)-quinoxalinone, 4-(2,6-diethyl-4-methylphenyl)-5-hydroxy-2,6-

dimethyl-3(2*H*)-pyridazinone, 4-(4-fluorophenyl)-6-[(2-hydroxy-6-oxo-1-cyclohexen-1-yl)carbonyl]-2-methyl-1,2,4-triazine-3,5(2*H*,4*H*)-dione, 5-[(2-hydroxy-6-oxo-1-cyclohexen-1-yl)carbonyl]-2-(3-methoxyphenyl)-3-(3-methoxypropyl)-4(3*H*)-pyrimidinone, 2-methyl-*N*-(4-methyl-1,2,5-oxadiazol-3-yl)-3-(methylsulfinyl)-4-(trifluoromethyl)benzamide and 2-methyl-3-(methylsulfonyl)-*N*-(1-methyl-1*H*-tetrazol-5-yl)-4-(trifluoromethyl)benzamide. Of note is a compound of the invention mixed with mesotrione or pyrasulfatole.

“HST inhibitors” (b13) disrupt a plant’s ability to convert homogentisate to 2-methyl-6-solanyl-1,4-benzoquinone, thereby disrupting carotenoid biosynthesis. Examples of HST inhibitors include cyclopyrimorate (6-chloro-3-(2-cyclopropyl-6-methylphenoxy)-4-pyridazinyl 4-morpholinecarboxylate), haloxydine, pyriclor, 3-(2-chloro-3,6-difluorophenyl)-4-hydroxy-1-methyl-1,5-naphthyridin-2(1*H*)-one, 7-(3,5-dichloro-4-pyridinyl)-5-(2,2-difluoroethyl)-8-hydroxypyrido[2,3-*b*]pyrazin-6(5*H*)-one and 4-(2,6-diethyl-4-methylphenyl)-5-hydroxy-2,6-dimethyl-3(2*H*)-pyridazinone.

HST inhibitors also include compounds of Formulae **A** and **B**.



15

wherein R^{d1} is H, Cl or CF₃; R^{d2} is H, Cl or Br; R^{d3} is H or Cl; R^{d4} is H, Cl or CF₃; R^{d5} is CH₃, CH₂CH₃ or CH₂CHF₂; and R^{d6} is OH, or -OC(=O)-*i*-Pr; and R^{e1} is H, F, Cl, CH₃ or CH₂CH₃; R^{e2} is H or CF₃; R^{e3} is H, CH₃ or CH₂CH₃; R^{e4} is H, F or Br; R^{e5} is Cl, CH₃, CF₃, OCF₃ or CH₂CH₃; R^{e6} is H, CH₃, CH₂CHF₂ or C≡CH; R^{e7} is OH, -OC(=O)Et, -OC(=O)-*i*-Pr or -OC(=O)-*t*-Bu; and A^{e8} is N or CH.

20

“Cellulose biosynthesis inhibitors” (b14) inhibit the biosynthesis of cellulose in certain plants. They are most effective when applied preemergence or early postemergence on young or rapidly growing plants. Examples of cellulose biosynthesis inhibitors include chlorthiamid, dichlobenil, flupoxam, indaziflam (*N*²-[(1*R*,2*S*)-2,3-dihydro-2,6-dimethyl-1*H*-inden-1-yl]-6-(1-fluoroethyl)-1,3,5-triazine-2,4-diamine), isoxaben and triaziflam.

25

“Other herbicides” (b15) include herbicides that act through a variety of different modes of action such as mitotic disruptors (e.g., flamprop-M-methyl and flamprop-M-isopropyl), organic arsenicals (e.g., DSMA, and MSMA), 7,8-dihydropteroate synthase inhibitors, chloroplast isoprenoid synthesis inhibitors and cell-wall biosynthesis inhibitors. Other herbicides include those herbicides having unknown modes of action or do

30

not fall into a specific category listed in (b1) through (b14) or act through a combination of modes of action listed above. Examples of other herbicides include acclonifen, asulam, amitrole, bromobutide, cinmethylin, clomazone, cumyluron, daimuron, difenzoquat, etobenzanid, fluometuron, flurenol, fosamine, fosamine-ammonium, dazomet, dymron, ipfencarbazone (1-(2,4-dichlorophenyl)-*N*-(2,4-difluorophenyl)-1,5-dihydro-*N*-(1-methylethyl)-5-oxo-4*H*-1,2,4-triazole-4-carboxamide), metam, methyl dymron, oleic acid, oxaziclomefone, pelargonic acid, pyributicarb and 5-[[[(2,6-difluorophenyl)methoxy]methyl]-4,5-dihydro-5-methyl-3-(3-methyl-2-thienyl)isoxazole].

“Herbicide safeners” (b16) are substances added to a herbicide formulation to eliminate or reduce phytotoxic effects of the herbicide to certain crops. These compounds protect crops from injury by herbicides but typically do not prevent the herbicide from controlling undesired vegetation. Examples of herbicide safeners include but are not limited to benoxacor, cloquintocet-mexyl, cumyluron, cyometrinil, cyprosulfamide, daimuron, dichlormid, dicyclonon, dietholate, dimepiperate, fenchlorazole-ethyl, fenclorim, flurazole, fluxofenim, furilazole, isoxadifen-ethyl, mefenpyr-diethyl, mephenate, methoxyphenone, naphthalic anhydride, oxabetrinil, *N*-(aminocarbonyl)-2-methylbenzenesulfonamide and *N*-(aminocarbonyl)-2-fluorobenzenesulfonamide, 1-bromo-4-[(chloromethyl)sulfonyl]benzene, 2-(dichloromethyl)-2-methyl-1,3-dioxolane (MG 191), 4-(dichloroacetyl)-1-oxa-4-azospiro[4.5]decane (MON 4660), 2,2-dichloro-1-(2,2,5-trimethyl-3-oxazolidinyl)-ethanone and 2-methoxy-*N*-[[4-[(methylamino)carbonyl]amino]phenyl]sulfonyl]benzamide.


The compounds of Formula 1 can be prepared by general methods known in the art of synthetic organic chemistry. One or more of the following methods and variations as described in Schemes 1–5 can be used to prepare the compounds of Formula 1. The definitions of R¹, R², R³, R⁴, R⁵, Q¹, Q², Y¹, and Y² in the compounds of Formulae 1–6 below are as defined above in the Summary of the Invention unless otherwise noted. Compounds of Formulae 1a–1e are various subsets of the compounds of Formula 1, and all substituents for Formulae 1a–1e are as defined above for Formula 1 unless otherwise noted.

As shown in Scheme 1 compounds of Formula 1a (i.e. Formula 1 wherein R⁴ is H, and Y¹ and Y² are O) can be prepared by reaction of acids of Formula 2 with amines of Formula 3 in the presence of a dehydrative coupling reagent such as propylphosphonic anhydride (T3P), dicyclohexylcarbodiimide (DCC), *N*-(3-dimethylaminopropyl)-*N*'-ethylcarbodiimide, *N,N*-carbonyldiimidazole (EDC), 2-chloro-1,3-dimethylimidazolium chloride or 2-chloro-1-methylpyridinium iodide (Mukaiyama's Reagent). Polymer-supported reagents, such as polymer-supported cyclohexylcarbodiimide, are also suitable. These reactions are typically run at temperatures ranging from 0–60 °C in a solvent such as, but not limited to, dichloromethane, acetonitrile, *N,N*-dimethylformamide or ethyl acetate in the presence of a base such as triethylamine, *N,N*-diisopropylamine, or 1,8-diazabicyclo[5.4.0]undec-7-

ene. See *Organic Process Research & Development* **2009**, *13*, 900–906 for coupling conditions employing propylphosphonic anhydride. Substituents in the 3- and 4-positions of the furanone ring of compounds of Formula **1a**, i.e. C(O)N(Q²)(R¹) and Q¹, respectively, are predominantly in the trans configuration.


5

Scheme 1



As shown in Scheme 2 compounds of Formula **2** can be prepared by hydrolysis of esters of Formula **4** by methods well known to those skilled in the art. Hydrolysis is carried out with aqueous base or aqueous acid, typically in the presence of a co-solvent. Suitable bases for the reaction include, but are not limited to, hydroxides such as lithium, sodium and potassium hydroxide and carbonates such as sodium and potassium carbonate. Suitable acids for the reaction include, but are not limited to, inorganic acids such as hydrochloric acid, hydrobromic acid and sulfuric acid, and organic acids such as acetic acid and trifluoroacetic acid. A wide variety of co-solvents are suitable for the reaction including, but not limited to, methanol, ethanol and tetrahydrofuran. The reaction is conducted at temperatures ranging from -20 °C to the boiling point of the solvent, and typically from 0 to 100 °C. A representative procedure can be found in Ollis and co-workers: *J. Chem. Soc Perkin 1* **1975**, 1480.

Scheme 2




20

Esters of Formula 4 can be prepared by the reaction of epoxides of Formula 5 with substituted malonates of Formula 6. This transformation requires the presence of an acid acceptor such as sodium hydride, sodium methoxide or sodium ethoxide. Other alkali alkoxides and hydrides also may be successfully employed. The reaction can be carried out in a variety of solvents including protic solvents such as methanol and ethanol as well as aprotic solvents such as *N,N*-dimethylformamide, dimethyl sulfoxide and tetrahydrofuran. Temperatures from 0 °C to the boiling point of the solvent can be employed. Typical reaction conditions can be found in *Indian Journal of Chemistry, Section B: Organic Chemistry Including Medicinal Chemistry* **1981**, 20B (9), 807-8. Epoxides may be replaced in this reaction by 1,2-cyclic sulfites as reported by Nymann and Svendsen in *Acta Chemica Scandinavica* **1998**, 52(3), 338-349. Another route to esters of Formula 4 has been reported by Yamada and coworkers in *Journal of Organic Chemistry* **2008**, 73(24), 9535-9538. Alternatively, esters of Formula 4 can also be made by a method reported by Tran and colleagues in *Bioorganic & Medicinal Chemistry Letters* **2008**, 18(3), 1124-1130. Epoxides of Formula 5 are well known in the literature and can be prepared by the well-known olefin epoxidation reaction of known or commercially available styrenes. Alternatively the reaction of sulfoxonium ylides with known or commercially available aldehydes also can be employed to synthesize epoxides of Formula 5 as reviewed by Gobolobov and coworkers in *Tetrahedron* **1987**, 43, 2609.

20


Scheme 3




25

30

As shown in Scheme 4, mixtures of compounds of Formula 1b (i.e. Formula 1 wherein R⁵ is H, R⁴ is halogen and Y¹ and Y² are O) and Formula 1c (i.e. Formula 1 wherein R⁴ is H, R⁵ is halogen and Y¹ and Y² are O) can be prepared by reacting compounds of Formula 1a with a halogen source in a solvent, in the presence or absence of an initiator. Suitable halogen sources for this reaction include bromine, chlorine, *N*-chlorosuccinimide, *N*-bromosuccinimide and *N*-iodosuccinimide. Suitable initiators for this reaction include 2,2'-azobisisobutyronitrile (AIBN) and benzoyl peroxide. Typically, the reaction is carried out in solvents such as dichloromethane in the range of from 0 °C to the boiling point of the solvent.

Scheme 4

As shown in Scheme 5, compounds of Formula **1e** (i.e. Formula **1** wherein Y^1 and Y^2 are S) can be prepared by reacting compounds of Formula **1d** with at least two equivalents of a thionation reagent such as Lawesson's reagent, tetraphosphorus decasulfide or diphosphorus pentasulfide in a solvent such as tetrahydrofuran or toluene. Typically, the reaction is carried out at temperatures ranging from 0 to 115 °C. One skilled in the art recognizes that using less than two equivalents of the thionating reagent can provide mixtures comprising Formula **1** products wherein Y^1 is O and Y^2 is S, or Y^1 is S and Y^2 is O, which can be separated by conventional methods such as chromatography and crystallization.

Scheme 5

It is recognized by one skilled in the art that various functional groups can be converted into others to provide different compounds of Formula **1**. For a valuable resource that illustrates the interconversion of functional groups in a simple and straightforward fashion, see Larock, R. C., *Comprehensive Organic Transformations: A Guide to Functional Group Preparations*, 2nd Ed., Wiley-VCH, New York, 1999. For example, intermediates for the preparation of compounds of Formula **1** may contain aromatic nitro groups, which can be reduced to amino groups, and then be converted via reactions well known in the art such as the Sandmeyer reaction, to various halides, providing compounds of Formula **1**. The above reactions can also in many cases be performed in alternate order.

It is recognized that some reagents and reaction conditions described above for preparing compounds of Formula **1** may not be compatible with certain functionalities

present in the intermediates. In these instances, the incorporation of protection/deprotection sequences or functional group interconversions into the synthesis will aid in obtaining the desired products. The use and choice of the protecting groups will be apparent to one skilled in chemical synthesis (see, for example, Greene, T. W.; Wuts, P. G. M. *Protective Groups in Organic Synthesis*, 2nd ed.; Wiley: New York, 1991). One skilled in the art will recognize that, in some cases, after the introduction of a given reagent as depicted in any individual scheme, it may be necessary to perform additional routine synthetic steps not described in detail to complete the synthesis of compounds of Formula 1. One skilled in the art will also recognize that it may be necessary to perform a combination of the steps illustrated in the above schemes in an order other than that implied by the particular presented to prepare the compounds of Formula 1.

One skilled in the art will also recognize that compounds of Formula 1 and the intermediates described herein can be subjected to various electrophilic, nucleophilic, radical, organometallic, oxidation, and reduction reactions to add substituents or modify existing substituents.

Without further elaboration, it is believed that one skilled in the art using the preceding description can utilize the present invention to its fullest extent. The following non-limiting Examples are illustrative of the invention. Steps in the following Examples illustrate a procedure for each step in an overall synthetic transformation, and the starting material for each step may not have necessarily been prepared by a particular preparative run whose procedure is described in other Examples or Steps. Percentages are by weight except for chromatographic solvent mixtures or where otherwise indicated. Parts and percentages for chromatographic solvent mixtures are by volume unless otherwise indicated. ^1H NMR spectra are reported at 400 MHz in ppm downfield from tetramethylsilane; "s" means singlet, "d" means doublet, "t" means triplet, "q" means quartet and "m" means multiplet.

SYNTHESIS EXAMPLE 1

Preparation of 4-(4-fluorophenyl)tetrahydro-2-oxo-*N*-[2-(trifluoromethyl)phenyl]-3-furancarboxamide (Compound 6)

Step A: Preparation of 4-(4-fluorophenyl)tetrahydro-2-oxo-3-furancarboxylic acid ethyl ester

To a 250 mL two neck round bottom flask was added sodium metal (0.99 g, 0.043 mol) in ethanol (60 mL) and then diethyl malonate (6.6 g, 0.041 mol) was added slowly at room temperature. The reaction mixture was stirred for 15 min at room temperature, the temperature was then raised to 40 °C and 2-(4-fluorophenyl)oxirane (5.0 g, 0.036 mmol) was added slowly to the reaction mixture over 2 h. The reaction mixture was then stirred for 18 h at room temperature. The progress of the reaction was monitored by TLC analysis. After completion of reaction, the reaction mixture was neutralized using 1M aqueous hydrochloric acid and then concentrated to a residue. The resulting residue was diluted with water and

then extracted with ethyl acetate (3 x 300 mL). The combined organic layers were washed with water, dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford crude compound. This was purified by column chromatography using 20% ethyl acetate: petroleum ether to afford the title compound of Step A (3.0 g) as a colorless liquid.

5 ¹H NMR (DMSO-d₆) δ 7.48–7.44 (m, 2H), 7.23–7.19 (t, *J* = 8.8 Hz, 2H), 4.68–4.66 (t, *J* = 7.6 Hz, 1H), 4.22–4.19 (m, 2H), 4.17 (m, 1H), 4.15–4.11 (q, 2H), 1.17–1.14 (t, 3H).

Step B: Preparation of 4-(4-fluorophenyl)tetrahydro-2-oxo-3-furancarboxylic acid

To a solution of the compound from Step A (1.3 g, 5.158 mmol) in water (13 mL) was added potassium hydroxide (1.73 g, 30.957 mmol) and the reaction mixture was stirred at room temperature for 2 h. The progress of the reaction was monitored by TLC analysis. After completion of the reaction, the reaction mixture was cooled to 0 °C and concentrated hydrochloric acid was added and the resulting mixture was stirred for 15 min. The precipitate was filtered to afford the title compound of Step B (750 mg) as white solid melting at 138–140 °C.

15 ¹H NMR (DMSO-d₆) δ 13.2 (s, 1H), 7.46–7.45 (m, 2H), 7.44–7.21 (t, 2H), 4.67–4.63 (t, *J* = 7.6 Hz, 1H), 4.19–4.01 (m, 3H).


Step C: Preparation of 4-(4-fluorophenyl)tetrahydro-2-oxo-*N*-[2-(trifluoromethyl)phenyl]-3-furancarboxamide

To a solution of the compound from Step B (200 mg, 0.892 mmol) in *N,N*-dimethylformamide (3 mL) was added *N,N*-diisopropylethylamine (138 mg, 1.071 mmol) followed by HATU [1-[Bis(dimethylamino)methylene]-1*H*-1,2,3-triazolo[4,5-*b*]pyridinium 3-oxide hexafluorophosphate] (408 mg, 1.071 mmol) and 2-(trifluoromethyl)aniline (99 mg, 0.892 mmol). The reaction mixture was then stirred at room temperature for 6 h. The progress of the reaction was monitored by thin layer chromatography analysis. After completion of the reaction, the reaction mixture was poured into water (15 mL) and extracted with ethyl acetate (3 x 10 mL). The combined organic layers were washed with water followed by brine solution, dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford crude compound. This was purified by column chromatography using 30% ethyl acetate in petroleum ether to afford the title compound of Example 1, Compound 6 in Index Table A (100 mg) as white solid melting at 144–146 °C.

30 ¹H NMR (DMSO-d₆) δ 10.08 (s, 1H), 7.73–7.66 (m, 2H), 7.47–7.43 (m, 4H), 7.24–7.20 (t, 2H), 4.74–4.70 (t, *J* = 7.2 Hz, 1H), 4.21–4.19 (m, 3H).

By the procedures described herein together with methods known in the art, the following compounds of Tables 1 to 40 can be prepared. The following abbreviations are used in the Tables which follow: Me means methyl, *i*-Pr means isopropyl and Bu means butyl.

Table 1



Q ² is Ph(2-F); and Q ¹ is	Q ² is Ph(2-F); and Q ¹ is
Ph(3-Cl)	Ph(4-OCF ₂ H)
Ph(3-F)	Ph(4-OMe)
Ph(3-Br)	Ph(4-CH ₂ CF ₃)
Ph(3-Me)	Ph(4-O- <i>i</i> -Pr)
Ph(3-CF ₃)	Ph(4-OCF ₂ CF ₂ H)
Ph(3-CH ₂ CF ₃)	Ph(2,3-di-F)
Ph(3-OCF ₃)	Ph(2,4-di-F)
Ph(3-OCF ₂ H)	Ph(2,5-di-F)
Ph(3-O- <i>i</i> -Pr)	Ph(2,6-di-F)
Ph(3-OMe)	Ph(3,4-di-F)
Ph(3-OCF ₂ CF ₂ H)	Ph(3,5-di-F)
Ph(2-Cl)	Ph(3-Me,4-F)
Ph(2-F)	Ph(3-F,4-Me)
Ph(2-Br)	Ph(3-CF ₃ ,4-F)
Ph(2-Me)	Ph(3-F,4-CF ₃)
Ph(2-CF ₃)	Ph(2,3,4-tri-F)
Ph(2-OCF ₃)	Ph(3,4,5-tri-F)
Ph(2-OCF ₂ H)	2-Pyridinyl
Ph(2-OMe)	2-Pyridinyl(6-F)
Ph(2-OCF ₂ CF ₂ H)	2-Pyridinyl(6-CF ₃)
Ph(2-CH ₂ CF ₃)	2-Pyridinyl(6-Me)
Ph(2-O- <i>i</i> -Pr)	2-Pyridinyl(5-F)
Ph(4-Cl)	2-Pyridinyl(5-CF ₃)
Ph(4-F)	2-Pyridinyl(5-Me)
Ph(4-Br)	2-Pyridinyl(4-F)
Ph(4-Me)	2-Pyridinyl(4-CF ₃)
Ph(4-CF ₃)	2-Pyridinyl(4-Me)
Ph(4-OCF ₃)	2-Pyridinyl(3-F)

Q ² is Ph(2-F); and Q ¹ is	Q ² is Ph(2-F); and Q ¹ is
2-Pyridinyl(3-CF ₃)	3-Furyl
2-Pyridinyl(3-Me)	3-Furyl(4-CF ₃)
3-Pyridinyl	3-Furyl(5-CF ₃)
3-Pyridinyl(6-F)	Pyrazol-1-yl
3-Pyridinyl(6-CF ₃)	Pyrazol-1-yl(4-CF ₃)
3-Pyridinyl(6-Me)	Imidazol-1-yl
3-Pyridinyl(5-F)	Imidazol-1-yl(4-CF ₃)
3-Pyridinyl(5-CF ₃)	Imidazol-1-yl(2-CF ₃)
3-Pyridinyl(5-Me)	Imidazol-2-yl(1-Me)
3-Pyridinyl(4-F)	Imidazol-4-yl(1-Me)
3-Pyridinyl(4-CF ₃)	Imidazol-4-yl(3-Me)
3-Pyridinyl(4-Me)	Pyrazol-4-yl(1-Me)
3-Pyridinyl(2-F)	Triazol-4-yl(1-Me)
3-Pyridinyl(2-CF ₃)	Triazol-4-yl(2-Me)
3-Pyridinyl(2-Me)	Triazol-2-yl(4-Me)
4-Pyridinyl	Triazol-1-yl(4-Me)
4-Pyridinyl(6-F)	Pyrazin-2-yl
4-Pyridinyl(6-CF ₃)	Pyrazin-2-yl(5-CF ₃)
4-Pyridinyl(6-Me)	Pyrimidin-2-yl
4-Pyridinyl(5-F)	Pyrimidin-2-yl(5-CF ₃)
4-Pyridinyl(5-CF ₃)	Pyrimidin-5-yl
4-Pyridinyl(5-Me)	Pyrimidin-5-yl(2-CF ₃)
4-Pyridinyl(3-F)	1,3,5-Triazin-2-yl
4-Pyridinyl(3-CF ₃)	Thiazol-2-yl
4-Pyridinyl(3-Me)	Thiazol-2-yl(5-CF ₃)
4-Pyridinyl(2-F)	Thiazol-5-yl
4-Pyridinyl(2-CF ₃)	Thiazol-5-yl(2-CF ₃)
4-Pyridinyl(2-Me)	Oxazol-2-yl
2-Thienyl	Oxazol-2-yl(5-CF ₃)
2-Thienyl(4-CF ₃)	Oxazol-5-yl
2-Thienyl(5-CF ₃)	Oxazol-5-yl(2-CF ₃)
3-Thienyl	Isothiazol-5-yl
3-Thienyl(4-CF ₃)	Isothiazol-5-yl(3-CF ₃)
3-Thienyl(5-CF ₃)	Isothiazol-3-yl
2-Furyl	Isothiazol-3-yl(5-CF ₃)
2-Furyl(4-CF ₃)	Isoxazol-5-yl
2-Furyl(5-CF ₃)	Isoxazol-5-yl(3-CF ₃)

Q ² is Ph(2-F); and Q ¹ is	Q ² is Ph(2-F); and Q ¹ is
Isoxazol-3-yl	Pyrrolo[2,3-c]pyridin-1-yl
Isoxazol-3-yl(5-CF ₃)	Ph(3-OCH ₂ - <i>c</i> -Pr)
Tetrazol-1-yl	Ph(2-OCH ₂ - <i>c</i> -Pr)
Tetrazol-1-yl(5-Me)	Ph(4-OCH ₂ CH ₂ CH ₂ CH ₂ - <i>c</i> -hex)
Tetrazol-5-yl(1-Me)	Ph(CH ₂ - <i>c</i> -Pr)
1,2,4-Triazol-1-yl	Ph(4-CH ₂ CH ₂ CH ₂ CH ₂ - <i>c</i> -hex)
1,3,4-Oxadiazol-2-yl	Ph(2-(3,3-dichloroallyloxy))
1,3,4-Thiadiazol-2-yl	Ph(2-methoxyethoxy)
1,2,4-Oxadiazol-3-yl	Ph(3-propoxypropoxy)
1,2,4-Thiadiazol-3-yl	Ph(2-CH ₂ CH ₂ SCH ₃)
Tetrahydropyran-2-yl	Ph(2-CH ₂ CH ₂ SOCH ₃)
Tetrahydropyran-3-yl	Ph(2-CH ₂ CH ₂ SO ₂ CH ₃)
Tetrahydrofuran-2-yl	Ph(3-SMe)
Tetrahydrofuran-3-yl	Ph(3-SCF ₃)
1,3-Dioxolan-4-yl	Ph(3-S- <i>c</i> -Pr)
2,2-di-Fluoro-1,3-Dioxolan-4-yl	Ph(3-SOMe)
1,3-Dithiolan-4-yl	Ph(3-SOCF ₃)
1,4-Dioxolan-2-yl	Ph(3-SO- <i>c</i> -Pr)
1,4-Dithiolan-2-yl	Ph(3-SO ₂ Me)
1-Naphthyl	Ph(3-SO ₂ CF ₃)
2-Naphthyl	Ph(3-SO ₂ - <i>c</i> -Pr)
Benzofuran-2-yl	Ph(3-propargyl)
Benzothiophen-2-yl	Ph(3-(2-Butynyl))
1,3-Benzoxazol-2-yl	Ph(2-CH ₂ CH ₂ OCH ₂ CH ₃)
1,3-Benzthiazol-2-yl	Ph(2-C(=O)CH ₃)
7-Quinoly	Ph(2-OC(=O)CH ₃)
Indazol-1-yl	Ph(3-OC(=O)CH ₃)
Benzimidazol-1-yl	Ph(2-OC(=O)CF ₃)
Indol-1-yl	Ph(3-OC(=O)CF ₃)

Table 2 is constructed in the same manner except that the Row Heading “Q² is Ph(2-F); and Q¹ is” is replaced with the Row Heading listed for Table 2 below (i.e. “Q² is Ph(2,3-di-F); and Q¹ is”). Therefore the first entry in Table 2 is a compound of Formula 1 wherein Q² is Ph(2,3-di-F); and Q¹ is; and Q¹ is Ph(3-Cl) (i.e. 3-chlorophenyl). Tables 3 through 10 are constructed similarly.


5

Table	Row Heading
2	Q ² is Ph(2,3-di-F); and Q ¹ is

3	Q ² is Ph(2,4-di-F); and Q ¹ is
4	Q ² is Ph(2,3,4-tri-F); and Q ¹ is
5	Q ² is Ph(2-CF ₃); and Q ¹ is
6	Q ² is Ph(2-Me); and Q ¹ is
7	Q ² is Ph(2-NO ₂); and Q ¹ is
8	Q ² is Ph(2-Cl); and Q ¹ is
9	Q ² is Ph(2-SO ₂ Me); and Q ¹ is
10	Q ² is Ph(2-F,3-Cl); and Q ¹ is

Table 11

Table 11 is constructed the same way as Table 1 above, except the structure is replaced with the following:



5


Tables 12 through 20

This disclosure also includes Tables 12 through 20, each Table is constructed in the same fashion as Tables 2 through 10 above, except that the structure is replaced with the structure in Table 11 above.

10

Table 21

Table 21 is constructed the same way as Table 1 above, except the structure is replaced with the following:


Tables 22 through 30

15

This disclosure also includes Tables 22 through 30, each Table is constructed in the same fashion as Tables 2 through 10 above, except that the structure is replaced with the structure in Table 21 above.

Table 31

Table 31 is constructed the same way as Table 1 above, except the structure is replaced with the following:



5

Tables 32 through 40

This disclosure also includes Tables 32 through 40, each Table is constructed in the same fashion as Tables 2 through 10 above, except that the structure is replaced with the structure in Table 31 above.

A compound of this invention will generally be used as a herbicidal active ingredient in a composition, i.e. formulation, with at least one additional component selected from the group consisting of surfactants, solid diluents and liquid diluents, which serves as a carrier. The formulation or composition ingredients are selected to be consistent with the physical properties of the active ingredient, mode of application and environmental factors such as soil type, moisture and temperature.

Useful formulations include both liquid and solid compositions. Liquid compositions include solutions (including emulsifiable concentrates), suspensions, emulsions (including microemulsions, oil-in-water emulsions, flowable concentrates and/or suspoemulsions) and the like, which optionally can be thickened into gels. The general types of aqueous liquid compositions are soluble concentrate, suspension concentrate, capsule suspension, concentrated emulsion, microemulsion, oil-in-water emulsion, flowable concentrate and suspo-emulsion. The general types of nonaqueous liquid compositions are emulsifiable concentrate, microemulsifiable concentrate, dispersible concentrate and oil dispersion.

The general types of solid compositions are dusts, powders, granules, pellets, prills, pastilles, tablets, filled films (including seed coatings) and the like, which can be water-dispersible (“wettable”) or water-soluble. Films and coatings formed from film-forming solutions or flowable suspensions are particularly useful for seed treatment. Active ingredient can be (micro)encapsulated and further formed into a suspension or solid formulation; alternatively the entire formulation of active ingredient can be encapsulated (or “overcoated”). Encapsulation can control or delay release of the active ingredient. An emulsifiable granule combines the advantages of both an emulsifiable concentrate formulation and a dry granular formulation. High-strength compositions are primarily used as intermediates for further formulation.

Sprayable formulations are typically extended in a suitable medium before spraying. Such liquid and solid formulations are formulated to be readily diluted in the spray medium, usually water, but occasionally another suitable medium like an aromatic or paraffinic hydrocarbon or vegetable oil. Spray volumes can range from about one to several thousand liters per hectare, but more typically are in the range from about ten to several hundred liters per hectare. Sprayable formulations can be tank mixed with water or another suitable medium for foliar treatment by aerial or ground application, or for application to the growing medium of the plant. Liquid and dry formulations can be metered directly into drip irrigation systems or metered into the furrow during planting.

The formulations will typically contain effective amounts of active ingredient, diluent and surfactant within the following approximate ranges which add up to 100 percent by weight.

	Weight Percent		
	<u>Active Ingredient</u>	<u>Diluent</u>	<u>Surfactant</u>
Water-Dispersible and Water-soluble Granules, Tablets and Powders	0.001–90	0–99.999	0–15
Oil Dispersions, Suspensions, Emulsions, Solutions (including Emulsifiable Concentrates)	1–50	40–99	0–50
Dusts	1–25	70–99	0–5
Granules and Pellets	0.001–99	5–99.999	0–15
High Strength Compositions	90–99	0–10	0–2

Solid diluents include, for example, clays such as bentonite, montmorillonite, attapulgite and kaolin, gypsum, cellulose, titanium dioxide, zinc oxide, starch, dextrin, sugars (e.g., lactose, sucrose), silica, talc, mica, diatomaceous earth, urea, calcium carbonate, sodium carbonate and bicarbonate, and sodium sulfate. Typical solid diluents are described in Watkins et al., *Handbook of Insecticide Dust Diluents and Carriers*, 2nd Ed., Dorland Books, Caldwell, New Jersey.

Liquid diluents include, for example, water, *N,N*-dimethylalkanamides (e.g., *N,N*-dimethylformamide), limonene, dimethyl sulfoxide, *N*-alkylpyrrolidones (e.g., *N*-methylpyrrolidinone), alkyl phosphates (e.g., triethyl phosphate), ethylene glycol, triethylene glycol, propylene glycol, dipropylene glycol, polypropylene glycol, propylene carbonate, butylene carbonate, paraffins (e.g., white mineral oils, normal paraffins, isoparaffins), alkylbenzenes, alkylnaphthalenes, glycerine, glycerol triacetate, sorbitol, aromatic hydrocarbons, dearomatized aliphatics, alkylbenzenes, alkylnaphthalenes, ketones such as cyclohexanone, 2-heptanone, isophorone and 4-hydroxy-4-methyl-2-pentanone,

acetates such as isoamyl acetate, hexyl acetate, heptyl acetate, octyl acetate, nonyl acetate, tridecyl acetate and isobornyl acetate, other esters such as alkylated lactate esters, dibasic esters, alkyl and aryl benzoates and γ -butyrolactone, and alcohols, which can be linear, branched, saturated or unsaturated, such as methanol, ethanol, *n*-propanol, isopropyl alcohol, 5 *n*-butanol, isobutyl alcohol, *n*-hexanol, 2-ethylhexanol, *n*-octanol, decanol, isodecyl alcohol, isooctadecanol, cetyl alcohol, lauryl alcohol, tridecyl alcohol, oleyl alcohol, cyclohexanol, tetrahydrofurfuryl alcohol, diacetone alcohol, cresol and benzyl alcohol. Liquid diluents also include glycerol esters of saturated and unsaturated fatty acids (typically C₆–C₂₂), such as plant seed and fruit oils (e.g., oils of olive, castor, linseed, sesame, corn 10 (maize), peanut, sunflower, grapeseed, safflower, cottonseed, soybean, rapeseed, coconut and palm kernel), animal-sourced fats (e.g., beef tallow, pork tallow, lard, cod liver oil, fish oil), and mixtures thereof. Liquid diluents also include alkylated fatty acids (e.g., methylated, ethylated, butylated) wherein the fatty acids may be obtained by hydrolysis of glycerol esters from plant and animal sources, and can be purified by distillation. Typical 15 liquid diluents are described in Marsden, *Solvents Guide*, 2nd Ed., Interscience, New York, 1950.

The solid and liquid compositions of the present invention often include one or more surfactants. When added to a liquid, surfactants (also known as “surface-active agents”) generally modify, most often reduce, the surface tension of the liquid. Depending on the 20 nature of the hydrophilic and lipophilic groups in a surfactant molecule, surfactants can be useful as wetting agents, dispersants, emulsifiers or defoaming agents.

Surfactants can be classified as nonionic, anionic or cationic. Nonionic surfactants useful for the present compositions include, but are not limited to: alcohol alkoxylates such as alcohol alkoxylates based on natural and synthetic alcohols (which may be branched or 25 linear) and prepared from the alcohols and ethylene oxide, propylene oxide, butylene oxide or mixtures thereof; amine ethoxylates, alkanolamides and ethoxylated alkanolamides; alkoxylated triglycerides such as ethoxylated soybean, castor and rapeseed oils; alkylphenol alkoxylates such as octylphenol ethoxylates, nonylphenol ethoxylates, dinonyl phenol ethoxylates and dodecyl phenol ethoxylates (prepared from the phenols and ethylene oxide, 30 propylene oxide, butylene oxide or mixtures thereof); block polymers prepared from ethylene oxide or propylene oxide and reverse block polymers where the terminal blocks are prepared from propylene oxide; ethoxylated fatty acids; ethoxylated fatty esters and oils; ethoxylated methyl esters; ethoxylated tristyrylphenol (including those prepared from ethylene oxide, propylene oxide, butylene oxide or mixtures thereof); fatty acid esters, 35 glycerol esters, lanolin-based derivatives, polyethoxylate esters such as polyethoxylated sorbitan fatty acid esters, polyethoxylated sorbitol fatty acid esters and polyethoxylated glycerol fatty acid esters; other sorbitan derivatives such as sorbitan esters; polymeric surfactants such as random copolymers, block copolymers, alkyd peg (polyethylene glycol)

resins, graft or comb polymers and star polymers; polyethylene glycols (pegs); polyethylene glycol fatty acid esters; silicone-based surfactants; and sugar-derivatives such as sucrose esters, alkyl polyglycosides and alkyl polysaccharides.

Useful anionic surfactants include, but are not limited to: alkylaryl sulfonic acids and their salts; carboxylated alcohol or alkylphenol ethoxylates; diphenyl sulfonate derivatives; lignin and lignin derivatives such as lignosulfonates; maleic or succinic acids or their anhydrides; olefin sulfonates; phosphate esters such as phosphate esters of alcohol alkoxyates, phosphate esters of alkylphenol alkoxyates and phosphate esters of styryl phenol ethoxylates; protein-based surfactants; sarcosine derivatives; styryl phenol ether sulfate; sulfates and sulfonates of oils and fatty acids; sulfates and sulfonates of ethoxylated alkylphenols; sulfates of alcohols; sulfates of ethoxylated alcohols; sulfonates of amines and amides such as *N,N*-alkyltaurates; sulfonates of benzene, cumene, toluene, xylene, and dodecyl and tridecylbenzenes; sulfonates of condensed naphthalenes; sulfonates of naphthalene and alkyl naphthalene; sulfonates of fractionated petroleum; sulfosuccinamates; and sulfosuccinates and their derivatives such as dialkyl sulfosuccinate salts.

Useful cationic surfactants include, but are not limited to: amides and ethoxylated amides; amines such as *N*-alkyl propanediamines, tripropylenetriamines and dipropylenetetramines, and ethoxylated amines, ethoxylated diamines and propoxylated amines (prepared from the amines and ethylene oxide, propylene oxide, butylene oxide or mixtures thereof); amine salts such as amine acetates and diamine salts; quaternary ammonium salts such as quaternary salts, ethoxylated quaternary salts and diquaternary salts; and amine oxides such as alkyldimethylamine oxides and bis-(2-hydroxyethyl)-alkylamine oxides.

Also useful for the present compositions are mixtures of nonionic and anionic surfactants or mixtures of nonionic and cationic surfactants. Nonionic, anionic and cationic surfactants and their recommended uses are disclosed in a variety of published references including *McCutcheon's Emulsifiers and Detergents*, annual American and International Editions published by McCutcheon's Division, The Manufacturing Confectioner Publishing Co.; Sisely and Wood, *Encyclopedia of Surface Active Agents*, Chemical Publ. Co., Inc., New York, 1964; and A. S. Davidson and B. Milwidsky, *Synthetic Detergents*, Seventh Edition, John Wiley and Sons, New York, 1987.

Compositions of this invention may also contain formulation auxiliaries and additives, known to those skilled in the art as formulation aids (some of which may be considered to also function as solid diluents, liquid diluents or surfactants). Such formulation auxiliaries and additives may control: pH (buffers), foaming during processing (antifoams such polyorganosiloxanes), sedimentation of active ingredients (suspending agents), viscosity (thixotropic thickeners), in-container microbial growth (antimicrobials), product freezing (antifreezes), color (dyes/pigment dispersions), wash-off (film formers or stickers),

evaporation (evaporation retardants), and other formulation attributes. Film formers include, for example, polyvinyl acetates, polyvinyl acetate copolymers, polyvinylpyrrolidone-vinyl acetate copolymer, polyvinyl alcohols, polyvinyl alcohol copolymers and waxes. Examples of formulation auxiliaries and additives include those listed in *McCutcheon's Volume 2: Functional Materials*, annual International and North American editions published by
5 McCutcheon's Division, The Manufacturing Confectioner Publishing Co.; and PCT Publication WO 03/024222.

The compound of Formula 1 and any other active ingredients are typically incorporated into the present compositions by dissolving the active ingredient in a solvent or
10 by grinding in a liquid or dry diluent. Solutions, including emulsifiable concentrates, can be prepared by simply mixing the ingredients. If the solvent of a liquid composition intended for use as an emulsifiable concentrate is water-immiscible, an emulsifier is typically added to emulsify the active-containing solvent upon dilution with water. Active ingredient slurries, with particle diameters of up to 2,000 μm can be wet milled using media mills to obtain
15 particles with average diameters below 3 μm . Aqueous slurries can be made into finished suspension concentrates (see, for example, U.S. 3,060,084) or further processed by spray drying to form water-dispersible granules. Dry formulations usually require dry milling processes, which produce average particle diameters in the 2 to 10 μm range. Dusts and powders can be prepared by blending and usually grinding (such as with a hammer mill or
20 fluid-energy mill). Granules and pellets can be prepared by spraying the active material upon preformed granular carriers or by agglomeration techniques. See Browning, "Agglomeration", *Chemical Engineering*, December 4, 1967, pp 147–48, *Perry's Chemical Engineer's Handbook*, 4th Ed., McGraw-Hill, New York, 1963, pages 8–57 and following, and WO 91/13546. Pellets can be prepared as described in U.S. 4,172,714.
25 Water-dispersible and water-soluble granules can be prepared as taught in U.S. 4,144,050, U.S. 3,920,442 and DE 3,246,493. Tablets can be prepared as taught in U.S. 5,180,587, U.S. 5,232,701 and U.S. 5,208,030. Films can be prepared as taught in GB 2,095,558 and U.S. 3,299,566.

For further information regarding the art of formulation, see T. S. Woods, "The
30 Formulator's Toolbox – Product Forms for Modern Agriculture" in *Pesticide Chemistry and Bioscience, The Food–Environment Challenge*, T. Brooks and T. R. Roberts, Eds., Proceedings of the 9th International Congress on Pesticide Chemistry, The Royal Society of Chemistry, Cambridge, 1999, pp. 120–133. See also U.S. 3,235,361, Col. 6, line 16 through Col. 7, line 19 and Examples 10–41; U.S. 3,309,192, Col. 5, line 43 through Col. 7, line 62
35 and Examples 8, 12, 15, 39, 41, 52, 53, 58, 132, 138–140, 162–164, 166, 167 and 169–182; U.S. 2,891,855, Col. 3, line 66 through Col. 5, line 17 and Examples 1–4; Klingman, *Weed Control as a Science*, John Wiley and Sons, Inc., New York, 1961, pp 81–96; Hance et al.,

Weed Control Handbook, 8th Ed., Blackwell Scientific Publications, Oxford, 1989; and *Developments in formulation technology*, PJB Publications, Richmond, UK, 2000.

In the following Examples, all percentages are by weight and all formulations are prepared in conventional ways. Compound numbers refer to compounds in Index Table A. Without further elaboration, it is believed that one skilled in the art using the preceding description can utilize the present invention to its fullest extent. The following Examples are, therefore, to be construed as merely illustrative, and not limiting of the disclosure in any way whatsoever. Percentages are by weight except where otherwise indicated.

Example A

High Strength Concentrate

Compound 6	98.5%
silica aerogel	0.5%
synthetic amorphous fine silica	1.0%

10

Example B

Wettable Powder

Compound 6	65.0%
dodecylphenol polyethylene glycol ether	2.0%
sodium ligninsulfonate	4.0%
sodium silicoaluminate	6.0%
montmorillonite (calcined)	23.0%

Example C

Granule

Compound 6	10.0%
attapulgite granules (low volatile matter, 0.71/0.30 mm; U.S.S. No. 25–50 sieves)	90.0%

Example D

Extruded Pellet

Compound 6	25.0%
anhydrous sodium sulfate	10.0%
crude calcium ligninsulfonate	5.0%
sodium alkylnaphthalenesulfonate	1.0%
calcium/magnesium bentonite	59.0%

Example E

Emulsifiable Concentrate

Compound 6	10.0%
polyoxyethylene sorbitol hexoleate	20.0%
C ₆ –C ₁₀ fatty acid methyl ester	70.0%

Example FMicroemulsion

Compound 6	5.0%
polyvinylpyrrolidone-vinyl acetate copolymer	30.0%
alkylpolyglycoside	30.0%
glyceryl monooleate	15.0%
water	20.0%

Example GSuspension Concentrate

Compound 6	35%
butyl polyoxyethylene/polypropylene block copolymer	4.0%
stearic acid/polyethylene glycol copolymer	1.0%
styrene acrylic polymer	1.0%
xanthan gum	0.1%
propylene glycol	5.0%
silicone based defoamer	0.1%
1,2-benzisothiazolin-3-one	0.1%
water	53.7%

Example HEmulsion in Water

Compound 6	10.0%
butyl polyoxyethylene/polypropylene block copolymer	4.0%
stearic acid/polyethylene glycol copolymer	1.0%
styrene acrylic polymer	1.0%
xanthan gum	0.1%
propylene glycol	5.0%
silicone based defoamer	0.1%
1,2-benzisothiazolin-3-one	0.1%
aromatic petroleum based hydrocarbon	20.0
water	58.7%

Example IOil Dispersion

Compound 6	25%
polyoxyethylene sorbitol hexaoleate	15%
organically modified bentonite clay	2.5%
fatty acid methyl ester	57.5%

The present disclosure also includes Formulation Examples A through I above except “Compound 6” in each of the above Examples A through I is replaced with “Compound 1”, “Compound 2”, “Compound 3”, “Compound 4”, “Compound 5”, “Compound 7”, “Compound 8”, “Compound 9”, “Compound 10”, “Compound 11”, “Compound 12”, “Compound 13”, “Compound 14”, “Compound 15”, “Compound 16”, “Compound 17”, “Compound 18”, “Compound 19”, “Compound 20”, “Compound 21 or “Compound 22”.

Test results indicate that the compounds of the present invention are highly active preemergent and/or postemergent herbicides and/or plant growth regulants. The compounds of the invention generally show highest activity for postemergence weed control (i.e. applied after weed seedlings emerge from the soil) and preemergence weed control (i.e. applied before weed seedlings emerge from the soil). Many of them have utility for broad-spectrum pre- and/or postemergence weed control in areas where complete control of all vegetation is desired such as around fuel storage tanks, industrial storage areas, parking lots, drive-in theaters, air fields, river banks, irrigation and other waterways, around billboards and highway and railroad structures. Many of the compounds of this invention, by virtue of selective metabolism in crops versus weeds, or by selective activity at the locus of physiological inhibition in crops and weeds, or by selective placement on or within the environment of a mixture of crops and weeds, are useful for the selective control of grass and broadleaf weeds within a crop/weed mixture. One skilled in the art will recognize that the preferred combination of these selectivity factors within a compound or group of compounds can readily be determined by performing routine biological and/or biochemical assays. Compounds of this invention may show tolerance to important agronomic crops including, but is not limited to, alfalfa, barley, cotton, wheat, rape, sugar beets, corn (maize), sorghum, soybeans, rice, oats, peanuts, vegetables, tomato, potato, perennial plantation crops including coffee, cocoa, oil palm, rubber, sugarcane, citrus, grapes, fruit trees, nut trees, banana, plantain, pineapple, hops, tea and forests such as eucalyptus and conifers (e.g., loblolly pine), and turf species (e.g., Kentucky bluegrass, St. Augustine grass, Kentucky fescue and Bermuda grass). Compounds of this invention can be used in crops genetically transformed or bred to incorporate resistance to herbicides, express proteins toxic to invertebrate pests (such as *Bacillus thuringiensis* toxin), and/or express other useful traits. Those skilled in the art will appreciate that not all compounds are equally effective against all weeds. Alternatively, the subject compounds are useful to modify plant growth.

As the compounds of the invention have both preemergent and postemergent herbicidal activity, to control undesired vegetation by killing or injuring the vegetation or reducing its growth, the compounds can be usefully applied by a variety of methods involving contacting a herbicidally effective amount of a compound of the invention, or a

composition comprising said compound and at least one of a surfactant, a solid diluent or a liquid diluent, to the foliage or other part of the undesired vegetation or to the environment of the undesired vegetation such as the soil or water in which the undesired vegetation is growing or which surrounds the seed or other propagule of the undesired vegetation.

5 A herbicidally effective amount of the compounds of this invention is determined by a number of factors. These factors include: formulation selected, method of application, amount and type of vegetation present, growing conditions, etc. In general, a herbicidally effective amount of compounds of this invention is about 0.001 to 20 kg/ha with a preferred range of about 0.004 to 1 kg/ha. One skilled in the art can easily determine the herbicidally
10 effective amount necessary for the desired level of weed control.

In one common embodiment, a compound of the invention is applied, typically in a formulated composition, to a locus comprising desired vegetation (e.g., crops) and undesired vegetation (i.e. weeds), both of which may be seeds, seedlings and/or larger plants, in contact with a growth medium (e.g., soil). In this locus, a composition comprising a
15 compound of the invention can be directly applied to a plant or a part thereof, particularly of the undesired vegetation, and/or to the growth medium in contact with the plant.

Plant varieties and cultivars of the desired vegetation in the locus treated with a compound of the invention can be obtained by conventional propagation and breeding methods or by genetic engineering methods. Genetically modified plants (transgenic plants)
20 are those in which a heterologous gene (transgene) has been stably integrated into the plant's genome. A transgene that is defined by its particular location in the plant genome is called a transformation or transgenic event.

Genetically modified plant cultivars in the locus which can be treated according to the invention include those that are resistant against one or more biotic stresses (pests such as
25 nematodes, insects, mites, fungi, etc.) or abiotic stresses (drought, cold temperature, soil salinity, etc.), or that contain other desirable characteristics. Plants can be genetically modified to exhibit traits of, for example, herbicide tolerance, insect-resistance, modified oil profiles or drought tolerance. Useful genetically modified plants containing single gene transformation events or combinations of transformation events are listed in Exhibit C.
30 Additional information for the genetic modifications listed in Exhibit C can be obtained from publicly available databases maintained, for example, by the U.S. Department of Agriculture.

The following abbreviations, T1 through T37, are used in Exhibit C for traits. A “-” means the entry is not available; “tol.” means “tolerance” and “res.” means resistance.

Trait	Description	Trait	Description	Trait	Description
T1	Glyphosate tol.	T15	Cold tol.	T27	High tryptophan
T2	High lauric acid oil	T16	Imidazolinone herb. tol.	T28	Erect leaves semidwarf
T3	Glufosinate tol.	T17	Modified alpha-amylase	T29	Semidwarf

T4	Phytate breakdown	T18	Pollination control	T30	Low iron tol.
T5	Oxynil tol.	T19	2,4-D tol.	T31	Modified oil/fatty acid
T6	Disease res.	T20	Increased lysine	T32	HPPD tol.
T7	Insect res.	T21	Drought tol.	T33	High oil
T9	Modified flower color	T22	Delayed ripening/senescence	T34	Aryloxyalkanoate tol.
T11	ALS Herbicide tol.	T23	Modified product quality	T35	Mesotrione tol.
T12	Dicamba tol.	T24	High cellulose	T36	Reduced nicotine
T13	Anti-allergy	T25	Modified starch/carbohydrate	T37	Modified product
T14	Salt tol.	T26	Insect & disease resist.		

Exhibit C

Crop	Event Name	Event Code	Trait(s)	Gene(s)
Alfalfa	J101	MON-00101-8	T1	cp4 epsps (aroA:CP4)
Alfalfa	J163	MON-00163-7	T1	cp4 epsps (aroA:CP4)
Canola*	23-18-17 (Event 18)	CGN-89465-2	T2	te
Canola*	23-198 (Event 23)	CGN-89465-2	T2	te
Canola*	61061	DP-061061-7	T1	gat4621
Canola*	73496	DP-073496-4	T1	gat4621
Canola*	GT200 (RT200)	MON-89249-2	T1	cp4 epsps (aroA:CP4); goxv247
Canola*	GT73 (RT73)	MON-00073-7	T1	cp4 epsps (aroA:CP4); goxv247
Canola*	HCN10 (Topas 19/2)	-	T3	bar
Canola*	HCN28 (T45)	ACS-BN008-2	T3	pat (syn)
Canola*	HCN92 (Topas 19/2)	ACS-BN007-1	T3	bar
Canola*	MON88302	MON-88302-9	T1	cp4 epsps (aroA:CP4)
Canola*	MPS961	-	T4	phyA
Canola*	MPS962	-	T4	phyA
Canola*	MPS963	-	T4	phyA
Canola*	MPS964	-	T4	phyA
Canola*	MPS965	-	T4	phyA
Canola*	MS1 (B91-4)	ACS-BN004-7	T3	bar
Canola*	MS8	ACS-BN005-8	T3	bar
Canola*	OXY-235	ACS-BN011-5	T5	bxn
Canola*	PHY14	-	T3	bar
Canola*	PHY23	-	T3	bar
Canola*	PHY35	-	T3	bar
Canola*	PHY36	-	T3	bar
Canola*	RF1 (B93-101)	ACS-BN001-4	T3	bar
Canola*	RF2 (B94-2)	ACS-BN002-5	T3	bar

Canola*	RF3	ACS-BNØØ3-6	T3	bar
Bean	EMBRAPA 5.1	EMB-PV051-1	T6	ac1 (sense and antisense)
Brinjal #	EE-1	-	T7	cry1Ac
Cotton	19-51a	DD-Ø1951A-7	T11	S4-HrA
Cotton	281-24-236	DAS-24236-5	T3,T7	pat (syn); cry1F
Cotton	3006-210-23	DAS-21Ø23-5	T3,T7	pat (syn); cry1Ac
Cotton	31707	-	T5,T7	bxn; cry1Ac
Cotton	31803	-	T5,T7	bxn; cry1Ac
Cotton	31807	-	T5,T7	bxn; cry1Ac
Cotton	31808	-	T5,T7	bxn; cry1Ac
Cotton	42317	-	T5,T7	bxn; cry1Ac
Cotton	BNLA-601	-	T7	cry1Ac
Cotton	BXN10211	BXN10211-9	T5	bxn; cry1Ac
Cotton	BXN10215	BXN10215-4	T5	bxn; cry1Ac
Cotton	BXN10222	BXN10222-2	T5	bxn; cry1Ac
Cotton	BXN10224	BXN10224-4	T5	bxn; cry1Ac
Cotton	COT102	SYN-IR102-7	T7	vip3A(a)
Cotton	COT67B	SYN-IR67B-1	T7	cry1Ab
Cotton	COT202	-	T7	vip3A
Cotton	Event 1	-	T7	cry1Ac
Cotton	GMF Cry1A	GTL-GMF311-7	T7	cry1Ab-Ac
Cotton	GHB119	BCS-GH005-8	T7	cry2Ae
Cotton	GHB614	BCS-GH002-5	T1	2mepsps
Cotton	GK12	-	T7	cry1Ab-Ac
Cotton	LLCotton25	ACS-GH001-3	T3	bar
Cotton	MLS 9124	-	T7	cry1C
Cotton	MON1076	MON-89924-2	T7	cry1Ac
Cotton	MON1445	MON-01445-2	T1	cp4 epsps (aroA:CP4)
Cotton	MON15985	MON-15985-7	T7	cry1Ac; cry2Ab2
Cotton	MON1698	MON-89383-1	T7	cp4 epsps (aroA:CP4)
Cotton	MON531	MON-00531-6	T7	cry1Ac
Cotton	MON757	MON-00757-7	T7	cry1Ac
Cotton	MON88913	MON-88913-8	T1	cp4 epsps (aroA:CP4)
Cotton	Nqwe Chi 6 Bt	-	T7	-
Cotton	SKG321	-	T7	cry1A; CpTI
Cotton	T303-3	BCS-GH003-6	T3,T7	cry1Ab; bar
Cotton	T304-40	BCS-GH004-7	T3,T7	cry1Ab; bar
Cotton	CE43-67B	-	T7	cry1Ab
Cotton	CE46-02A	-	T7	cry1Ab
Cotton	CE44-69D	-	T7	cry1Ab

Cotton	1143-14A	-	T7	cry1Ab
Cotton	1143-51B	-	T7	cry1Ab
Cotton	T342-142	-	T7	cry1Ab
Cotton	PV-GHGT07 (1445)	-	T1	cp4 epsps (aroA:CP4)
Cotton	EE-GH3	-	T1	mepsps
Cotton	EE-GH5	-	T7	cry1Ab
Cotton	MON88701	MON-88701-3	T3,T12	Modified dmo; bar
Cotton	OsCr11	-	T13	Modified Cry j
Flax	FP967	CDC-FL001-2	T11	als
Lentil	RH44	-	T16	als
Maize	3272	SYN-E3272-5	T17	amy797E
Maize	5307	SYN-05307-1	T7	ecry3.1Ab
Maize	59122	DAS-59122-7	T3,T7	cry34Ab1; cry35Ab1; pat
Maize	676	PH-000676-7	T3,T18	pat; dam
Maize	678	PH-000678-9	T3,T18	pat; dam
Maize	680	PH-000680-2	T3,T18	pat; dam
Maize	98140	DP-098140-6	T1,T11	gat4621; zm-hra
Maize	Bt10	-	T3,T7	cry1Ab; pat
Maize	Bt176 (176)	SYN-EV176-9	T3,T7	cry1Ab; bar
Maize	BVLA430101	-	T4	phyA2
Maize	CBH-351	ACS-ZM004-3	T3,T7	cry9C; bar
Maize	DAS40278-9	DAS40278-9	T19	aad-1
Maize	DBT418	DKB-89614-9	T3,T7	cry1Ac; pinII; bar
Maize	DLL25 (B16)	DKB-89790-5	T3	bar
Maize	GA21	MON-00021-9	T1	mepsps
Maize	GG25	-	T1	mepsps
Maize	GJ11	-	T1	mepsps
Maize	FI117	-	T1	mepsps
Maize	GAT-ZM1	-	T3	pat
Maize	LY038	REN-00038-3	T20	cordapA
Maize	MIR162	SYN-IR162-4	T7	vip3Aa20
Maize	MIR604	SYN-IR604-5	T7	mcry3A
Maize	MON801 (MON80100)	MON801	T1,T7	cry1Ab; cp4 epsps (aroA:CP4); goxv247
Maize	MON802	MON-80200-7	T1,T7	cry1Ab; cp4 epsps (aroA:CP4); goxv247
Maize	MON809	PH-MON-809- 2	T1,T7	cry1Ab; cp4 epsps (aroA:CP4); goxv247
Maize	MON810	MON-00810-6	T1,T7	cry1Ab; cp4 epsps (aroA:CP4); goxv247
Maize	MON832	-	T1	cp4 epsps (aroA:CP4); goxv247
Maize	MON863	MON-00863-5	T7	cry3Bb1
Maize	MON87427	MON-87427-7	T1	cp4 epsps (aroA:CP4)
Maize	MON87460	MON-87460-4	T21	cspB

Maize	MON88017	MON-88017-3	T1,T7	cry3Bb1; cp4 epsps (aroA:CP4)
Maize	MON89034	MON-89034-3	T7	cry2Ab2; cry1A.105
Maize	MS3	ACS-ZM001-9	T3,T18	bar; barnase
Maize	MS6	ACS-ZM005-4	T3,T18	bar; barnase
Maize	NK603	MON-00603-6	T1	cp4 epsps (aroA:CP4)
Maize	T14	ACS-ZM002-1	T3	pat (syn)
Maize	T25	ACS-ZM003-2	T3	pat (syn)
Maize	TC1507	DAS-01507-1	T3,T7	cry1Fa2; pat
Maize	TC6275	DAS-06275-8	T3,T7	mocry1F; bar
Maize	VIP1034	-	T3,T7	vip3A; pat
Maize	43A47	DP-043A47-3	T3,T7	cry1F; cry34Ab1; cry35Ab1; pat
Maize	40416	DP-040416-8	T3,T7	cry1F; cry34Ab1; cry35Ab1; pat
Maize	32316	DP-032316-8	T3,T7	cry1F; cry34Ab1; cry35Ab1; pat
Maize	4114	DP-004114-3	T3,T7	cry1F; cry34Ab1; cry35Ab1; pat
Melon	Melon A	-	T22	sam-k
Melon	Melon B	-	T22	sam-k
Papaya	55-1	CUH-CP551-8	T6	prsv cp
Papaya	63-1	CUH-CP631-7	T6	prsv cp
Papaya	Huanong No. 1	-	T6	prsv rep
Papaya	X17-2	UFL-X17CP-6	T6	prsv cp
Plum	C-5	ARS-PLMC5-6	T6	ppv cp
Canola**	ZSR500	-	T1	cp4 epsps (aroA:CP4); goxv247
Canola**	ZSR502	-	T1	cp4 epsps (aroA:CP4); goxv247
Canola**	ZSR503	-	T1	cp4 epsps (aroA:CP4); goxv247
Rice	7Crp#242-95-7	-	T13	7crp
Rice	7Crp#10	-	T13	7crp
Rice	GM Shanyou 63	-	T7	cry1Ab; cry1Ac
Rice	Huahui-1/TT51-1	-	T7	cry1Ab; cry1Ac
Rice	LLRICE06	ACS-OS001-4	T3	bar
Rice	LLRICE601	BCS-OS003-7	T3	bar
Rice	LLRICE62	ACS-OS002-5	T3	bar
Rice	Tarom molaii + cry1Ab	-	T7	cry1Ab (truncated)
Rice	GAT-OS2	-	T3	bar
Rice	GAT-OS3	-	T3	bar
Rice	PE-7	-	T7	Cry1Ac
Rice	7Crp#10	-	T13	7crp
Rice	KPD627-8	-	T27	OASA1D
Rice	KPD722-4	-	T27	OASA1D
Rice	KA317	-	T27	OASA1D
Rice	HW5	-	T27	OASA1D
Rice	HW1	-	T27	OASA1D

Rice	B-4-1-18	-	T28	Δ OsBRI1
Rice	G-3-3-22	-	T29	OSGA2ox1
Rice	AD77	-	T6	DEF
Rice	AD51	-	T6	DEF
Rice	AD48	-	T6	DEF
Rice	AD41	-	T6	DEF
Rice	13pNasNa800725atAprt1	-	T30	HvNAS1; HvNAAT-A; APRT
Rice	13pAprt1	-	T30	APRT
Rice	gHvNAS1-gHvNAAT-1	-	T30	HvNAS1; HvNAAT-A; HvNAAT-B
Rice	gHvIDS3-1	-	T30	HvIDS3
Rice	gHvNAAT1	-	T30	HvNAAT-A; HvNAAT-B
Rice	gHvNAS1-1	-	T30	HvNAS1
Rice	NIA-OS006-4	-	T6	WRKY45
Rice	NIA-OS005-3	-	T6	WRKY45
Rice	NIA-OS004-2	-	T6	WRKY45
Rice	NIA-OS003-1	-	T6	WRKY45
Rice	NIA-OS002-9	-	T6	WRKY45
Rice	NIA-OS001-8	-	T6	WRKY45
Rice	OsCr11	-	T13	Modified Cry j
Rice	17053	-	T1	cp4 epsps (aroA:CP4)
Rice	17314	-	T1	cp4 epsps (aroA:CP4)
Rose	WKS82 / 130-4-1	IFD-52401-4	T9	5AT; bp40 (f3'5'h)
Rose	WKS92 / 130-9-1	IFD-52901-9	T9	5AT; bp40 (f3'5'h)
Soybean	260-05 (G94-1, G94-19, G168)	-	T9	gm-fad2-1 (silencing locus)
Soybean	A2704-12	ACS-GM005-3	T3	pat
Soybean	A2704-21	ACS-GM004-2	T3	pat
Soybean	A5547-127	ACS-GM006-4	T3	pat
Soybean	A5547-35	ACS-GM008-6	T3	pat
Soybean	CV127	BPS-CV127-9	T16	csr1-2
Soybean	DAS68416-4	DAS68416-4	T3	pat
Soybean	DP305423	DP-305423-1	T11,T31	gm-fad2-1 (silencing locus); gm-hra
Soybean	DP356043	DP-356043-5	T1,T31	gm-fad2-1 (silencing locus); gat4601
Soybean	FG72	MST-FG072-3	T32,T1	2mepsps; hppdPF W336
Soybean	GTS 40-3-2 (40-3-2)	MON-04032-6	T1	cp4 epsps (aroA:CP4)
Soybean	GU262	ACS-GM003-1	T3	pat
Soybean	MON87701	MON-87701-2	T7	cry IAc
Soybean	MON87705	MON-87705-6	T1,T31	fatb1-A (sense & antisense); fad2-1A (sense & antisense); cp4 epsps (aroA:CP4)

Soybean	MON87708	MON-87708-9	T1,T12	dmo; cp4 epsps (aroA:CP4)
Soybean	MON87769	MON-87769-7	T1,T31	Pj.D6D; Nc.Fad3; cp4 epsps (aroA:CP4)
Soybean	MON89788	MON-89788-1	T1	cp4 epsps (aroA:CP4)
Soybean	W62	ACS-GM002-9	T3	bar
Soybean	W98	ACS-GM001-8	T3	bar
Soybean	MON87754	MON-87754-1	T33	dgat2A
Soybean	DAS21606	DAS-21606	T34,T3	Modified aad-12; pat
Soybean	DAS44406	DAS-44406-6	T1,T3,T34	Modified aad-12; 2mepsps; pat
Soybean	SYHT04R	SYN-0004R-8	T35	Modified avhppd
Soybean	9582.814.19.1	-	T3,T7	cry1Ac, cry1F, PAT
Squash	CZW3	SEM-ØCZW3-2	T6	cmv cp, zymv cp, wmv cp
Squash	ZW20	SEM-ØZW20-7	T6	zymv cp, wmv cp
Sugar Beet	GTSB77 (T9100152)	SY-GTSB77-8	T1	cp4 epsps (aroA:CP4); goxv247
Sugar Beet	H7-1	KM-000H71-4	T1	cp4 epsps (aroA:CP4)
Sugar Beet	T120-7	ACS-BV001-3	T3	pat
Sugar Beet	T227-1	-	T1	cp4 epsps (aroA:CP4)
Sugarcane	NXI-1T	-	T21	EcbetA
Sunflower	X81359	-	T16	als
Pepper	PK-SP01	-	T6	cmv cp
Tobacco	C/F/93/08-02	-	T5	bxn
Tobacco	Vector 21-41	-	T36	NtQPT1 (antisense)
Sunflower	X81359	-	T16	als
Wheat	MON71800	MON-71800-3	T1	cp4 epsps (aroA:CP4)

* Argentine (*Brassica napus*), ** Polish (*B. rapa*), # Eggplant

Although most typically, compounds of the invention are used to control undesired vegetation, contact of desired vegetation in the treated locus with compounds of the invention may result in super-additive or synergistic effects with genetic traits in the desired vegetation, including traits incorporated through genetic modification. For example, resistance to phytophagous insect pests or plant diseases, tolerance to biotic/abiotic stresses or storage stability may be greater than expected from the genetic traits in the desired vegetation.

Compounds of this invention can also be mixed with one or more other biologically active compounds or agents including herbicides, herbicide safeners, fungicides, insecticides, nematocides, bactericides, acaricides, growth regulators such as insect molting inhibitors and rooting stimulants, chemosterilants, semiochemicals, repellents, attractants, pheromones, feeding stimulants, plant nutrients, other biologically active compounds or entomopathogenic bacteria, virus or fungi to form a multi-component pesticide giving an

even broader spectrum of agricultural protection. Mixtures of the compounds of the invention with other herbicides can broaden the spectrum of activity against additional weed species, and suppress the proliferation of any resistant biotypes. Thus the present invention also pertains to a composition comprising a compound of Formula 1 (in a herbicidally effective amount) and at least one additional biologically active compound or agent (in a biologically effective amount) and can further comprise at least one of a surfactant, a solid diluent or a liquid diluent. The other biologically active compounds or agents can be formulated in compositions comprising at least one of a surfactant, solid or liquid diluent. For mixtures of the present invention, one or more other biologically active compounds or agents can be formulated together with a compound of Formula 1, to form a premix, or one or more other biologically active compounds or agents can be formulated separately from the compound of Formula 1, and the formulations combined together before application (e.g., in a spray tank) or, alternatively, applied in succession.

A mixture of one or more of the following herbicides with a compound of this invention may be particularly useful for weed control: acetochlor, acifluorfen and its sodium salt, aclonifen, acrolein (2-propenal), alachlor, alloxydim, ametryn, amicarbazone, amidosulfuron, aminocyclopyrachlor and its esters (e.g., methyl, ethyl) and salts (e.g., sodium, potassium), aminopyralid, amitrole, ammonium sulfamate, anilofos, asulam, atrazine, azimsulfuron, beflubutamid, benazolin, benazolin-ethyl, bencarbazone, benfluralin, benfuresate, bensulfuron-methyl, bensulide, bentazone, benzobicyclon, benzofenap, bicyclopyrone, bifenox, bilanafos, bispyribac and its sodium salt, bromacil, bromobutide, bromofenoxim, bromoxynil, bromoxynil octanoate, butachlor, butafenacil, butamifos, butralin, butroxydim, butylate, cafenstrole, carbetamide, carfentrazone-ethyl, catechin, chlomethoxyfen, chloramben, chlorbromuron, chlorflurenol-methyl, chloridazon, chlorimuron-ethyl, chlorotoluron, chlorpropham, chlorsulfuron, chlorthal-dimethyl, chlorthiamid, cinidon-ethyl, cinmethylin, cinosulfuron, clacyfos, clefoxydim, clethodim, clodinafop-propargyl, clomazone, clomeprop, clopyralid, clopyralid-olamine, cloransulam-methyl, cumyluron, cyanazine, cycloate, cyclopyrimorate, cyclosulfamuron, cycloxydim, cyhalofop-butyl, 2,4-D and its butotyl, butyl, isooctyl and isopropyl esters and its dimethylammonium, diolamine and trolamine salts, daimuron, dalapon, dalapon-sodium, dazomet, 2,4-DB and its dimethylammonium, potassium and sodium salts, desmedipham, desmetryn, dicamba and its diglycolammonium, dimethylammonium, potassium and sodium salts, dichlobenil, dichlorprop, diclofop-methyl, diclosulam, difenzoquat metilsulfate, diflufenican, diflufenzopyr, dimefuron, dimepiperate, dimethachlor, dimethametryn, dimethenamid, dimethenamid-P, dimethipin, dimethylarsinic acid and its sodium salt, dinitramine, dinoterb, diphenamid, diquat dibromide, dithiopyr, diuron, DNOC, endothal, EPTC, esprocarb, ethalfluralin, ethametsulfuron-methyl, ethiozin, ethofumesate, ethoxyfen, ethoxysulfuron, etobenzanid, fenoxaprop-ethyl, fenoxaprop-P-ethyl, fenoxasulfone,

fenquino-trione, fentrazamide, fenuron, fenuron-TCA, flamprop-methyl, flamprop-M-isopropyl, flamprop-M-methyl, flazasulfuron, florasulam, fluazifop-butyl, fluazifop-P-butyl, fluazolate, flucarbazone, flucetosulfuron, fluchloralin, flufenacet, flufenpyr, flufenpyr-ethyl, flumetsulam, flumiclorac-pentyl, flumioxazin, fluometuron, 5 fluoroglycofen-ethyl, flupoxam, flupyr-sulfuron-methyl and its sodium salt, flurenol, flurenol-butyl, fluridone, flurochloridone, fluroxypyr, flurtamone, fluthiacet-methyl, fomesafen, foramsulfuron, fosamine-ammonium, glufosinate, glufosinate-ammonium, glufosinate-P, glyphosate and its salts such as ammonium, isopropylammonium, potassium, sodium (including sesquisodium) and trimesium (alternatively named sulfosate), halauxifen, 10 halauxifen-methyl, halosulfuron-methyl, haloxyfop-etotyl, haloxyfop-methyl, hexazinone, hydantocidin, imazamethabenz-methyl, imazamox, imazapic, imazapyr, imazaquin, imazaquin-ammonium, imazethapyr, imazethapyr-ammonium, imazosulfuron, indanofan, indaziflam, iofensulfuron, iodosulfuron-methyl, ioxynil, ioxynil octanoate, ioxynil-sodium, ipfencarbazone, isoproturon, isouron, isoxaben, isoxaflutole, isoxachlortole, lactofen, lenacil, 15 linuron, maleic hydrazide, MCPA and its salts (e.g., MCPA-dimethylammonium, MCPA-potassium and MCPA-sodium, esters (e.g., MCPA-2-ethylhexyl, MCPA-butotyl) and thioesters (e.g., MCPA-thioethyl), MCPB and its salts (e.g., MCPB-sodium) and esters (e.g., MCPB-ethyl), mecoprop, mecoprop-P, mefenacet, mefluidide, mesosulfuron-methyl, mesotrione, metam-sodium, metamifop, metamitron, metazachlor, metazosulfuron, 20 methabenzthiazuron, methylarsonic acid and its calcium, monoammonium, monosodium and disodium salts, methyldymron, metobenzuron, metobromuron, metolachlor, S-metolachlor, metosulam, metoxuron, metribuzin, metsulfuron-methyl, molinate, monolinuron, naproanilide, napropamide, napropamide-M, naptalam, neburon, nicosulfuron, norflurazon, orbencarb, orthosulfamuron, oryzalin, oxadiargyl, oxadiazon, oxasulfuron, oxaziclomefone, 25 oxyfluorfen, paraquat dichloride, pebulate, pelargonic acid, pendimethalin, penoxsulam, pentanochlor, pentoxazone, perfluidone, pethoxamid, pethoxyamid, phenmedipham, picloram, picloram-potassium, picolinafen, pinoxaden, piperophos, pretilachlor, primisulfuron-methyl, prodiamine, profoxydim, prometon, prometryn, propachlor, propanil, propaquiza-fop, propazine, propham, propisochlor, propoxycarbazone, propyrisulfuron, 30 propyzamide, prosulfocarb, prosulfuron, pyraclonil, pyraflufen-ethyl, pyrasulfotole, pyrazogyl, pyrazolynate, pyrazoxyfen, pyrazosulfuron-ethyl, pyribenzoxim, pyributicarb, pyridate, pyriftalid, pyriminobac-methyl, pyrimisulfan, pyrithiobac, pyrithiobac-sodium, pyroxasulfone, pyroxsulam, quinclorac, quinmerac, quinclamine, quizalofop-ethyl, quizalofop-P-ethyl, quizalofop-P-tefuryl, rimsulfuron, saflufenacil, sethoxydim, siduron, 35 simazine, simetryn, sulcotrione, sulfentrazone, sulfometuron-methyl, sulfosulfuron, 2,3,6-TBA, TCA, TCA-sodium, tebutam, tebuthiuron, tefuryltrione, tembotrione, tepraloxydim, terbacil, terbumeton, terbuthylazine, terbutryn, thenylchlor, thiazopyr, thiencarbazone, thifensulfuron-methyl, thiobencarb, tiafenacil, tiocarbazil, tolpyralate, topramezone,

tralkoxydim, tri-allate, triafamone, triasulfuron, triaziflam, tribenuron-methyl, triclopyr, triclopyr-butotyl, triclopyr-triethylammonium, tridiphane, trietazine, trifloxysulfuron, trifludimoxazin, trifluralin, triflusulfuron-methyl, tritosulfuron, vernolate, 3-(2-chloro-3,6-difluorophenyl)-4-hydroxy-1-methyl-1,5-naphthyridin-2(1H)-one, 5-chloro-3-[(2-hydroxy-6-oxo-1-cyclohexen-1-yl)carbonyl]-1-(4-methoxyphenyl)-2(1H)-quinoxalinone, 2-chloro-N-(1-methyl-1H-tetrazol-5-yl)-6-(trifluoromethyl)-3-pyridinecarboxamide, 7-(3,5-dichloro-4-pyridinyl)-5-(2,2-difluoroethyl)-8-hydroxypyrido[2,3-b]pyrazin-6(5H)-one, 4-(2,6-diethyl-4-methylphenyl)-5-hydroxy-2,6-dimethyl-3(2H)-pyridazinone), 5-[[[(2,6-difluorophenyl)methoxy]methyl]-4,5-dihydro-5-methyl-3-(3-methyl-2-thienyl)isoxazole (previously methioxolin), 4-(4-fluorophenyl)-6-[(2-hydroxy-6-oxo-1-cyclohexen-1-yl)carbonyl]-2-methyl-1,2,4-triazine-3,5(2H,4H)-dione, methyl 4-amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxyphenyl)-5-fluoro-2-pyridinecarboxylate, 2-methyl-3-(methylsulfonyl)-N-(1-methyl-1H-tetrazol-5-yl)-4-(trifluoromethyl)benzamide and 2-methyl-N-(4-methyl-1,2,5-oxadiazol-3-yl)-3-(methylsulfinyl)-4-(trifluoromethyl)benzamide. Other herbicides also include bioherbicides such as *Alternaria destruens* Simmons, *Colletotrichum gloeosporioides* (Penz.) Penz. & Sacc., *Drechslera monoceras* (MTB-951), *Myrothecium verrucaria* (Albertini & Schweinitz) Ditmar: Fries, *Phytophthora palmivora* (Butl.) Butl. and *Puccinia thlaspeos* Schub.

Compounds of this invention can also be used in combination with plant growth regulators such as aviglycine, N-(phenylmethyl)-1H-purin-6-amine, epocholeone, gibberellic acid, gibberellin A₄ and A₇, harpin protein, mepiquat chloride, prohexadione calcium, prohydrojasmon, sodium nitrophenolate and trinexapac-methyl, and plant growth modifying organisms such as *Bacillus cereus* strain BP01.

General references for agricultural protectants (i.e. herbicides, herbicide safeners, insecticides, fungicides, nematocides, acaricides and biological agents) include *The Pesticide Manual, 13th Edition*, C. D. S. Tomlin, Ed., British Crop Protection Council, Farnham, Surrey, U.K., 2003 and *The BioPesticide Manual, 2nd Edition*, L. G. Copping, Ed., British Crop Protection Council, Farnham, Surrey, U.K., 2001.

For embodiments where one or more of these various mixing partners are used, the mixing partners are typically used in the amounts similar to amounts customary when the mixture partners are used alone. More particularly in mixtures, active ingredients are often applied at an application rate between one-half and the full application rate specified on product labels for use of active ingredient alone. These amounts are listed in references such as *The Pesticide Manual* and *The BioPesticide Manual*. The weight ratio of these various mixing partners (in total) to the compound of Formula 1 is typically between about 1:3000 and about 3000:1. Of note are weight ratios between about 1:300 and about 300:1 (for example ratios between about 1:30 and about 30:1). One skilled in the art can easily determine through simple experimentation the biologically effective amounts of active

ingredients necessary for the desired spectrum of biological activity. It will be evident that including these additional components may expand the spectrum of weeds controlled beyond the spectrum controlled by the compound of Formula 1 alone.

5 In certain instances, combinations of a compound of this invention with other biologically active (particularly herbicidal) compounds or agents (i.e. active ingredients) can result in a greater-than-additive (i.e. synergistic) effect on weeds and/or a less-than-additive effect (i.e. safening) on crops or other desirable plants. Reducing the quantity of active ingredients released in the environment while ensuring effective pest control is always desirable. Ability to use greater amounts of active ingredients to provide more effective
10 weed control without excessive crop injury is also desirable. When synergism of herbicidal active ingredients occurs on weeds at application rates giving agronomically satisfactory levels of weed control, such combinations can be advantageous for reducing crop production cost and decreasing environmental load. When safening of herbicidal active ingredients occurs on crops, such combinations can be advantageous for increasing crop protection by
15 reducing weed competition.

Of note is a combination of a compound of the invention with at least one other herbicidal active ingredient. Of particular note is such a combination where the other herbicidal active ingredient has different site of action from the compound of the invention. In certain instances, a combination with at least one other herbicidal active ingredient having
20 a similar spectrum of control but a different site of action will be particularly advantageous for resistance management. Thus, a composition of the present invention can further comprise (in a herbicidally effective amount) at least one additional herbicidal active ingredient having a similar spectrum of control but a different site of action.

Compounds of this invention can also be used in combination with herbicide safeners
25 such as allidochlor, benoxacor, cloquintocet-mexyl, cumyluron, cyometrinil, cyprosulphonamide, daimuron, dichlormid, dicyclonon, dietholate, dimepiperate, fenchlorazole-ethyl, fenclorim, flurazole, fluxofenim, furilazole, isoxadifen-ethyl, mefenpyr-diethyl, mephenate, methoxyphenone naphthalic anhydride (1,8-naphthalic anhydride), oxabetrinil, *N*-(aminocarbonyl)-2-methylbenzenesulfonamide, *N*-(aminocarbonyl)-
30 2-fluorobenzenesulfonamide, 1-bromo-4-[(chloromethyl)sulfonyl]benzene (BCS), 4-(dichloroacetyl)-1-oxa-4-azospiro[4.5]decane (MON 4660), 2-(dichloromethyl)-2-methyl-1,3-dioxolane (MG 191), ethyl 1,6-dihydro-1-(2-methoxyphenyl)-6-oxo-2-phenyl-5-pyrimidinecarboxylate, 2-hydroxy-*N,N*-dimethyl-6-(trifluoromethyl)pyridine-3-carboxamide, and 3-oxo-1-cyclohexen-1-yl 1-(3,4-dimethylphenyl)-1,6-dihydro-6-oxo-2-
35 phenyl-5-pyrimidinecarboxylate, 2,2-dichloro-1-(2,2,5-trimethyl-3-oxazolidinyl)-ethanone and 2-methoxy-*N*-[[4-[[[(methylamino)carbonyl]amino]phenyl]sulfonyl]-benzamide to increase safety to certain crops. Antidotally effective amounts of the herbicide safeners can be applied at the same time as the compounds of this invention, or applied as seed

treatments. Therefore an aspect of the present invention relates to a herbicidal mixture comprising a compound of this invention and an antidotally effective amount of a herbicide safener. Seed treatment is particularly useful for selective weed control, because it physically restricts antidoting to the crop plants. Therefore a particularly useful embodiment of the present invention is a method for selectively controlling the growth of undesired vegetation in a crop comprising contacting the locus of the crop with a herbicidally effective amount of a compound of this invention wherein seed from which the crop is grown is treated with an antidotally effective amount of safener. Antidotally effective amounts of safeners can be easily determined by one skilled in the art through simple experimentation.

Compounds of the invention can also be mixed with: (1) polynucleotides including but not limited to DNA, RNA, and/or chemically modified nucleotides influencing the amount of a particular target through down regulation, interference, suppression or silencing of the genetically derived transcript that render a herbicidal effect; or (2) polynucleotides including but not limited to DNA, RNA, and/or chemically modified nucleotides influencing the amount of a particular target through down regulation, interference, suppression or silencing of the genetically derived transcript that render a safening effect.

Of note is a composition comprising a compound of the invention (in a herbicidally effective amount), at least one additional active ingredient selected from the group consisting of other herbicides and herbicide safeners (in an effective amount), and at least one component selected from the group consisting of surfactants, solid diluents and liquid diluents.

Table A1 lists specific combinations of a Component (a) with Component (b) illustrative of the mixtures, compositions and methods of the present invention. Compound 6 in the Component (a) column is identified in Index Table A. The second column of Table A1 lists the specific Component (b) compound (e.g., "2,4-D" in the first line). The third, fourth and fifth columns of Table A1 lists ranges of weight ratios for rates at which the Component (a) compound is typically applied to a field-grown crop relative to Component (b) (i.e. (a):(b)). Thus, for example, the first line of Table A1 specifically discloses the combination of Component (a) (i.e. Compound 6 in Index Table A) with 2,4-D is typically applied in a weight ratio between 1:192 – 6:1. The remaining lines of Table A1 are to be construed similarly.

TABLE A1

Component (a) (Compound #)	Component (b)	Typical Weight Ratio	More Typical Weight Ratio	Most Typical Weight Ratio
6	2,4-D	1:192 – 6:1	1:64 – 2:1	1:24 – 1:3
6	Acetochlor	1:768 – 2:1	1:256 – 1:2	1:96 – 1:11
6	Acifluorfen	1:96 – 12:1	1:32 – 4:1	1:12 – 1:2
6	Aclonifen	1:857 – 2:1	1:285 – 1:3	1:107 – 1:12

Component (a) (Compound #)	Component (b)	Typical Weight Ratio	More Typical Weight Ratio	Most Typical Weight Ratio
6	Alachlor	1:768 – 2:1	1:256 – 1:2	1:96 – 1:11
6	Ametryn	1:384 – 3:1	1:128 – 1:1	1:48 – 1:6
6	Amicarbazone	1:192 – 6:1	1:64 – 2:1	1:24 – 1:3
6	Amidosulfuron	1:6 – 168:1	1:2 – 56:1	1:1 – 11:1
6	Aminocyclopyrachlor	1:48 – 24:1	1:16 – 8:1	1:6 – 2:1
6	Aminopyralid	1:20 – 56:1	1:6 – 19:1	1:2 – 4:1
6	Amitrole	1:768 – 2:1	1:256 – 1:2	1:96 – 1:11
6	Anilofos	1:96 – 12:1	1:32 – 4:1	1:12 – 1:2
6	Asulam	1:960 – 2:1	1:320 – 1:3	1:120 – 1:14
6	Atrazine	1:192 – 6:1	1:64 – 2:1	1:24 – 1:3
6	Azimsulfuron	1:6 – 168:1	1:2 – 56:1	1:1 – 11:1
6	Beflubutamid	1:342 – 4:1	1:114 – 2:1	1:42 – 1:5
6	Benfuresate	1:617 – 2:1	1:205 – 1:2	1:77 – 1:9
6	Bensulfuron-methyl	1:25 – 45:1	1:8 – 15:1	1:3 – 3:1
6	Bentazone	1:192 – 6:1	1:64 – 2:1	1:24 – 1:3
6	Benzobicyclon	1:85 – 14:1	1:28 – 5:1	1:10 – 1:2
6	Benzofenap	1:257 – 5:1	1:85 – 2:1	1:32 – 1:4
6	Bicyclopyrone	1:42 – 27:1	1:14 – 9:1	1:5 – 2:1
6	Bifenox	1:257 – 5:1	1:85 – 2:1	1:32 – 1:4
6	Bispyribac-sodium	1:10 – 112:1	1:3 – 38:1	1:1 – 7:1
6	Bromacil	1:384 – 3:1	1:128 – 1:1	1:48 – 1:6
6	Bromobutide	1:384 – 3:1	1:128 – 1:1	1:48 – 1:6
6	Bromoxynil	1:96 – 12:1	1:32 – 4:1	1:12 – 1:2
6	Butachlor	1:768 – 2:1	1:256 – 1:2	1:96 – 1:11
6	Butafenacil	1:42 – 27:1	1:14 – 9:1	1:5 – 2:1
6	Butylate	1:1542 – 1:2	1:514 – 1:5	1:192 – 1:22
6	Cafenstrole	1:192 – 6:1	1:64 – 2:1	1:24 – 1:3
6	Carfentrazone-ethyl	1:128 – 9:1	1:42 – 3:1	1:16 – 1:2
6	Chlorimuron-ethyl	1:8 – 135:1	1:2 – 45:1	1:1 – 9:1
6	Chlorotoluron	1:768 – 2:1	1:256 – 1:2	1:96 – 1:11
6	Chlorsulfuron	1:6 – 168:1	1:2 – 56:1	1:1 – 11:1
6	Cinosulfuron	1:17 – 68:1	1:5 – 23:1	1:2 – 5:1
6	Cinidon-ethyl	1:384 – 3:1	1:128 – 1:1	1:48 – 1:6
6	Cinmethylin	1:34 – 34:1	1:11 – 12:1	1:4 – 3:1
6	Clacyfos	1:34 – 34:1	1:11 – 12:1	1:4 – 3:1

Component (a) (Compound #)	Component (b)	Typical Weight Ratio	More Typical Weight Ratio	Most Typical Weight Ratio
6	Clethodim	1:48 – 24:1	1:16 – 8:1	1:6 – 2:1
6	Clodinafop-propargyl	1:20 – 56:1	1:6 – 19:1	1:2 – 4:1
6	Clomazone	1:384 – 3:1	1:128 – 1:1	1:48 – 1:6
6	Clomeprop	1:171 – 7:1	1:57 – 3:1	1:21 – 1:3
6	Clopyralid	1:192 – 6:1	1:64 – 2:1	1:24 – 1:3
6	Cloransulam-methyl	1:12 – 96:1	1:4 – 32:1	1:1 – 6:1
6	Cumyluron	1:384 – 3:1	1:128 – 1:1	1:48 – 1:6
6	Cyanazine	1:384 – 3:1	1:128 – 1:1	1:48 – 1:6
6	Cyclopyrimorate	1:17 – 68:1	1:5 – 23:1	1:2 – 5:1
6	Cyclosulfamuron	1:17 – 68:1	1:5 – 23:1	1:2 – 5:1
6	Cycloxydim	1:96 – 12:1	1:32 – 4:1	1:12 – 1:2
6	Cyhalofop	1:25 – 45:1	1:8 – 15:1	1:3 – 3:1
6	Daimuron	1:192 – 6:1	1:64 – 2:1	1:24 – 1:3
6	Desmedipham	1:322 – 4:1	1:107 – 2:1	1:40 – 1:5
6	Dicamba	1:192 – 6:1	1:64 – 2:1	1:24 – 1:3
6	Dichlobenil	1:1371 – 1:2	1:457 – 1:4	1:171 – 1:20
6	Dichlorprop	1:925 – 2:1	1:308 – 1:3	1:115 – 1:13
6	Diclofop-methyl	1:384 – 3:1	1:128 – 1:1	1:48 – 1:6
6	Diclosulam	1:10 – 112:1	1:3 – 38:1	1:1 – 7:1
6	Difenzoquat	1:288 – 4:1	1:96 – 2:1	1:36 – 1:4
6	Diflufenican	1:857 – 2:1	1:285 – 1:3	1:107 – 1:12
6	Diflufenzopyr	1:12 – 96:1	1:4 – 32:1	1:1 – 6:1
6	Dimethachlor	1:768 – 2:1	1:256 – 1:2	1:96 – 1:11
6	Dimethametryn	1:192 – 6:1	1:64 – 2:1	1:24 – 1:3
6	Dimethenamid-P	1:384 – 3:1	1:128 – 1:1	1:48 – 1:6
6	Dithiopyr	1:192 – 6:1	1:64 – 2:1	1:24 – 1:3
6	Diuron	1:384 – 3:1	1:128 – 1:1	1:48 – 1:6
6	EPTC	1:768 – 2:1	1:256 – 1:2	1:96 – 1:11
6	Esprocarb	1:1371 – 1:2	1:457 – 1:4	1:171 – 1:20
6	Ethalfuralin	1:384 – 3:1	1:128 – 1:1	1:48 – 1:6
6	Ethametsulfuron-methyl	1:17 – 68:1	1:5 – 23:1	1:2 – 5:1
6	Ethoxyfen	1:8 – 135:1	1:2 – 45:1	1:1 – 9:1
6	Ethoxysulfuron	1:20 – 56:1	1:6 – 19:1	1:2 – 4:1
6	Etobenzanid	1:257 – 5:1	1:85 – 2:1	1:32 – 1:4
6	Fenoxaprop-ethyl	1:120 – 10:1	1:40 – 4:1	1:15 – 1:2

Component (a) (Compound #)	Component (b)	Typical Weight Ratio	More Typical Weight Ratio	Most Typical Weight Ratio
6	Fenoxasulfone	1:85 – 14:1	1:28 – 5:1	1:10 – 1:2
6	Fenquinotrione	1:17 – 68:1	1:5 – 23:1	1:2 – 5:1
6	Fentrazamide	1:17 – 68:1	1:5 – 23:1	1:2 – 5:1
6	Flazasulfuron	1:17 – 68:1	1:5 – 23:1	1:2 – 5:1
6	Florasulam	1:2 – 420:1	1:1 – 140:1	2:1 – 27:1
6	Fluazifop-butyl	1:192 – 6:1	1:64 – 2:1	1:24 – 1:3
6	Flucarbazone	1:8 – 135:1	1:2 – 45:1	1:1 – 9:1
6	Flucetosulfuron	1:8 – 135:1	1:2 – 45:1	1:1 – 9:1
6	Flufenacet	1:257 – 5:1	1:85 – 2:1	1:32 – 1:4
6	Flumetsulam	1:24 – 48:1	1:8 – 16:1	1:3 – 3:1
6	Flumiclorac-pentyl	1:10 – 112:1	1:3 – 38:1	1:1 – 7:1
6	Flumioxazin	1:25 – 45:1	1:8 – 15:1	1:3 – 3:1
6	Fluometuron	1:384 – 3:1	1:128 – 1:1	1:48 – 1:6
6	Flupyrsulfuron-methyl	1:3 – 336:1	1:1 – 112:1	2:1 – 21:1
6	Fluridone	1:384 – 3:1	1:128 – 1:1	1:48 – 1:6
6	Fluroxypyr	1:96 – 12:1	1:32 – 4:1	1:12 – 1:2
6	Flurtamone	1:857 – 2:1	1:285 – 1:3	1:107 – 1:12
6	Fluthiacet-methyl	1:48 – 42:1	1:16 – 14:1	1:3 – 3:1
6	Fomesafen	1:96 – 12:1	1:32 – 4:1	1:12 – 1:2
6	Foramsulfuron	1:13 – 84:1	1:4 – 28:1	1:1 – 6:1
6	Glufosinate	1:288 – 4:1	1:96 – 2:1	1:36 – 1:4
6	Glyphosate	1:288 – 4:1	1:96 – 2:1	1:36 – 1:4
6	Halauxifen	1:20 – 56:1	1:6 – 19:1	1:2 – 4:1
6	Halauxifen-methyl	1:20 – 56:1	1:6 – 19:1	1:2 – 4:1
6	Halosulfuron-methyl	1:17 – 68:1	1:5 – 23:1	1:2 – 5:1
6	Haloxypyr-methyl	1:34 – 34:1	1:11 – 12:1	1:4 – 3:1
6	Hexazinone	1:192 – 6:1	1:64 – 2:1	1:24 – 1:3
6	Imazamox	1:13 – 84:1	1:4 – 28:1	1:1 – 6:1
6	Imazapic	1:20 – 56:1	1:6 – 19:1	1:2 – 4:1
6	Imazapyr	1:85 – 14:1	1:28 – 5:1	1:10 – 1:2
6	Imazaquin	1:34 – 34:1	1:11 – 12:1	1:4 – 3:1
6	Imazethabenz-methyl	1:171 – 7:1	1:57 – 3:1	1:21 – 1:3
6	Imazethapyr	1:24 – 48:1	1:8 – 16:1	1:3 – 3:1
6	Imazosulfuron	1:27 – 42:1	1:9 – 14:1	1:3 – 3:1
6	Indanofan	1:342 – 4:1	1:114 – 2:1	1:42 – 1:5

Component (a) (Compound #)	Component (b)	Typical Weight Ratio	More Typical Weight Ratio	Most Typical Weight Ratio
6	Indaziflam	1:25 – 45:1	1:8 – 15:1	1:3 – 3:1
6	Iodosulfuron-methyl	1:3 – 336:1	1:1 – 112:1	2:1 – 21:1
6	Ioxynil	1:192 – 6:1	1:64 – 2:1	1:24 – 1:3
6	Ipfencarbazono	1:85 – 14:1	1:28 – 5:1	1:10 – 1:2
6	Isoproturon	1:384 – 3:1	1:128 – 1:1	1:48 – 1:6
6	Isoxaben	1:288 – 4:1	1:96 – 2:1	1:36 – 1:4
6	Isoxaflutole	1:60 – 20:1	1:20 – 7:1	1:7 – 2:1
6	Lactofen	1:42 – 27:1	1:14 – 9:1	1:5 – 2:1
6	Lenacil	1:384 – 3:1	1:128 – 1:1	1:48 – 1:6
6	Linuron	1:384 – 3:1	1:128 – 1:1	1:48 – 1:6
6	MCPA	1:192 – 6:1	1:64 – 2:1	1:24 – 1:3
6	MCPB	1:288 – 4:1	1:96 – 2:1	1:36 – 1:4
6	Mecoprop	1:768 – 2:1	1:256 – 1:2	1:96 – 1:11
6	Mefenacet	1:384 – 3:1	1:128 – 1:1	1:48 – 1:6
6	Mefluidide	1:192 – 6:1	1:64 – 2:1	1:24 – 1:3
6	Mesosulfuron-methyl	1:5 – 224:1	1:1 – 75:1	1:1 – 14:1
6	Mesotrione	1:42 – 27:1	1:14 – 9:1	1:5 – 2:1
6	Metamifop	1:42 – 27:1	1:14 – 9:1	1:5 – 2:1
6	Metazachlor	1:384 – 3:1	1:128 – 1:1	1:48 – 1:6
6	Metazosulfuron	1:25 – 45:1	1:8 – 15:1	1:3 – 3:1
6	Methabenzthiazuron	1:768 – 2:1	1:256 – 1:2	1:96 – 1:11
6	Metolachlor	1:768 – 2:1	1:256 – 1:2	1:96 – 1:11
6	Metosulam	1:8 – 135:1	1:2 – 45:1	1:1 – 9:1
6	Metribuzin	1:192 – 6:1	1:64 – 2:1	1:24 – 1:3
6	Metsulfuron-methyl	1:2 – 560:1	1:1 – 187:1	3:1 – 35:1
6	Molinate	1:1028 – 2:1	1:342 – 1:3	1:128 – 1:15
6	Napropamide	1:384 – 3:1	1:128 – 1:1	1:48 – 1:6
6	Napropamide-M	1:192 – 6:1	1:64 – 2:1	1:24 – 1:3
6	Naptalam	1:192 – 6:1	1:64 – 2:1	1:24 – 1:3
6	Nicosulfuron	1:12 – 96:1	1:4 – 32:1	1:1 – 6:1
6	Norflurazon	1:1152 – 1:1	1:384 – 1:3	1:144 – 1:16
6	Orbencarb	1:1371 – 1:2	1:457 – 1:4	1:171 – 1:20
6	Orthosulfamuron	1:20 – 56:1	1:6 – 19:1	1:2 – 4:1
6	Oryzalin	1:514 – 3:1	1:171 – 1:2	1:64 – 1:8
6	Oxadiargyl	1:384 – 3:1	1:128 – 1:1	1:48 – 1:6

Component (a) (Compound #)	Component (b)	Typical Weight Ratio	More Typical Weight Ratio	Most Typical Weight Ratio
6	Oxadiazon	1:548 – 3:1	1:182 – 1:2	1:68 – 1:8
6	Oxasulfuron	1:27 – 42:1	1:9 – 14:1	1:3 – 3:1
6	Oxaziclomefone	1:42 – 27:1	1:14 – 9:1	1:5 – 2:1
6	Oxyfluorfen	1:384 – 3:1	1:128 – 1:1	1:48 – 1:6
6	Paraquat	1:192 – 6:1	1:64 – 2:1	1:24 – 1:3
6	Pendimethalin	1:384 – 3:1	1:128 – 1:1	1:48 – 1:6
6	Penoxsulam	1:10 – 112:1	1:3 – 38:1	1:1 – 7:1
6	Penthoamid	1:384 – 3:1	1:128 – 1:1	1:48 – 1:6
6	Pentoxazone	1:102 – 12:1	1:34 – 4:1	1:12 – 1:2
6	Phenmedipham	1:102 – 12:1	1:34 – 4:1	1:12 – 1:2
6	Picloram	1:96 – 12:1	1:32 – 4:1	1:12 – 1:2
6	Picolinafen	1:34 – 34:1	1:11 – 12:1	1:4 – 3:1
6	Pinoxaden	1:25 – 45:1	1:8 – 15:1	1:3 – 3:1
6	Pretilachlor	1:192 – 6:1	1:64 – 2:1	1:24 – 1:3
6	Primisulfuron-methyl	1:8 – 135:1	1:2 – 45:1	1:1 – 9:1
6	Prodiamine	1:384 – 3:1	1:128 – 1:1	1:48 – 1:6
6	Profoxydim	1:42 – 27:1	1:14 – 9:1	1:5 – 2:1
6	Prometryn	1:384 – 3:1	1:128 – 1:1	1:48 – 1:6
6	Propachlor	1:1152 – 1:1	1:384 – 1:3	1:144 – 1:16
6	Propanil	1:384 – 3:1	1:128 – 1:1	1:48 – 1:6
6	Propaquizafop	1:48 – 24:1	1:16 – 8:1	1:6 – 2:1
6	Propoxycarbazone	1:17 – 68:1	1:5 – 23:1	1:2 – 5:1
6	Propyrisulfuron	1:17 – 68:1	1:5 – 23:1	1:2 – 5:1
6	Propyzamide	1:384 – 3:1	1:128 – 1:1	1:48 – 1:6
6	Prosulfocarb	1:1200 – 1:2	1:400 – 1:4	1:150 – 1:17
6	Prosulfuron	1:6 – 168:1	1:2 – 56:1	1:1 – 11:1
6	Pyraclonil	1:42 – 27:1	1:14 – 9:1	1:5 – 2:1
6	Pyraflufen-ethyl	1:5 – 224:1	1:1 – 75:1	1:1 – 14:1
6	Pyrasulfotole	1:13 – 84:1	1:4 – 28:1	1:1 – 6:1
6	Pyrazolynate	1:857 – 2:1	1:285 – 1:3	1:107 – 1:12
6	Pyrazosulfuron-ethyl	1:10 – 112:1	1:3 – 38:1	1:1 – 7:1
6	Pyrazoxyfen	1:5 – 224:1	1:1 – 75:1	1:1 – 14:1
6	Pyribenzoxim	1:10 – 112:1	1:3 – 38:1	1:1 – 7:1
6	Pyributicarb	1:384 – 3:1	1:128 – 1:1	1:48 – 1:6
6	Pyridate	1:288 – 4:1	1:96 – 2:1	1:36 – 1:4

Component (a) (Compound #)	Component (b)	Typical Weight Ratio	More Typical Weight Ratio	Most Typical Weight Ratio
6	Pyrifthalid	1:10 – 112:1	1:3 – 38:1	1:1 – 7:1
6	Pyriminobac-methyl	1:20 – 56:1	1:6 – 19:1	1:2 – 4:1
6	Pyrimisulfan	1:17 – 68:1	1:5 – 23:1	1:2 – 5:1
6	Pyriothiobac	1:24 – 48:1	1:8 – 16:1	1:3 – 3:1
6	Pyroxasulfone	1:85 – 14:1	1:28 – 5:1	1:10 – 1:2
6	Pyroxulam	1:5 – 224:1	1:1 – 75:1	1:1 – 14:1
6	Quinclorac	1:192 – 6:1	1:64 – 2:1	1:24 – 1:3
6	Quizalofop-ethyl	1:42 – 27:1	1:14 – 9:1	1:5 – 2:1
6	Rimsulfuron	1:13 – 84:1	1:4 – 28:1	1:1 – 6:1
6	Saflufenacil	1:25 – 45:1	1:8 – 15:1	1:3 – 3:1
6	Sethoxydim	1:96 – 12:1	1:32 – 4:1	1:12 – 1:2
6	Simazine	1:384 – 3:1	1:128 – 1:1	1:48 – 1:6
6	Sulcotrione	1:120 – 10:1	1:40 – 4:1	1:15 – 1:2
6	Sulfentrazone	1:147 – 8:1	1:49 – 3:1	1:18 – 1:3
6	Sulfometuron-methyl	1:34 – 34:1	1:11 – 12:1	1:4 – 3:1
6	Sulfosulfuron	1:8 – 135:1	1:2 – 45:1	1:1 – 9:1
6	Tebuthiuron	1:384 – 3:1	1:128 – 1:1	1:48 – 1:6
6	Tefuryltrione	1:42 – 27:1	1:14 – 9:1	1:5 – 2:1
6	Tembotrione	1:31 – 37:1	1:10 – 13:1	1:3 – 3:1
6	Tepaloxymid	1:25 – 45:1	1:8 – 15:1	1:3 – 3:1
6	Terbacil	1:288 – 4:1	1:96 – 2:1	1:36 – 1:4
6	Terbutylazine	1:857 – 2:1	1:285 – 1:3	1:107 – 1:12
6	Terbutryn	1:192 – 6:1	1:64 – 2:1	1:24 – 1:3
6	Thenylchlor	1:85 – 14:1	1:28 – 5:1	1:10 – 1:2
6	Thiazopyr	1:384 – 3:1	1:128 – 1:1	1:48 – 1:6
6	Thiencarbazone	1:3 – 336:1	1:1 – 112:1	2:1 – 21:1
6	Thifensulfuron-methyl	1:5 – 224:1	1:1 – 75:1	1:1 – 14:1
6	Tiafenacil	1:17 – 68:1	1:5 – 23:1	1:2 – 5:1
6	Thiobencarb	1:768 – 2:1	1:256 – 1:2	1:96 – 1:11
6	Topramezone	1:6 – 168:1	1:2 – 56:1	1:1 – 11:1
6	Tralkoxydim	1:68 – 17:1	1:22 – 6:1	1:8 – 2:1
6	Triallate	1:768 – 2:1	1:256 – 1:2	1:96 – 1:11
6	Triasulfuron	1:5 – 224:1	1:1 – 75:1	1:1 – 14:1
6	Triaziflam	1:171 – 7:1	1:57 – 3:1	1:21 – 1:3
6	Tribenuron-methyl	1:3 – 336:1	1:1 – 112:1	2:1 – 21:1

Component (a) (Compound #)	Component (b)	Typical Weight Ratio	More Typical Weight Ratio	Most Typical Weight Ratio
6	Triclopyr	1:192 – 6:1	1:64 – 2:1	1:24 – 1:3
6	Trifloxysulfuron	1:2 – 420:1	1:1 – 140:1	2:1 – 27:1
6	Trifluralin	1:288 – 4:1	1:96 – 2:1	1:36 – 1:4
6	Triflusulfuron-methyl	1:17 – 68:1	1:5 – 23:1	1:2 – 5:1
6	Tritosulfuron	1:13 – 84:1	1:4 – 28:1	1:1 – 6:1

Table A2 is constructed the same as Table A1 above except that entries below the “Component (a)” column heading are replaced with the respective Component (a) Column Entry shown below. Compound 2 in the Component (a) column is identified in Index Table A. Thus, for example, in Table A2 the entries below the “Component (a)” column heading all recite “Compound 1” (i.e. Compound 1 identified in Index Table A), and the first line below the column headings in Table A2 specifically discloses a mixture of Compound 1 with 2,4-D. Tables A3 through A18 are constructed similarly.

<u>Table Number</u>	<u>Component (a) Column Entries</u>	<u>Table Number</u>	<u>Component (a) Column Entries</u>
A2	Compound 1	A13	Compound 13
A3	Compound 2	A14	Compound 14
A4	Compound 3	A15	Compound 15
A5	Compound 4	A16	Compound 16
A6	Compound 5	A17	Compound 17
A7	Compound 7	A18	Compound 18
A8	Compound 8	A19	Compound 19
A9	Compound 9	A20	Compound 20
A10	Compound 10	A21	Compound 21
A11	Compound 11	A22	Compound 22
A12	Compound 12		


Preferred for better control of undesired vegetation (e.g., lower use rate such as from synergism, broader spectrum of weeds controlled, or enhanced crop safety) or for preventing the development of resistant weeds are mixtures of a compound of this invention with a herbicide selected from the group consisting of chlorimuron-ethyl, nicosulfuron, mesotrione, thifensulfuron-methyl, flupyr-sulfuron-methyl, tribenuron, pyroxasulfone, pinoxaden, tembotrione, pyroxsulam, metolachlor and *S*-metolachlor.

The following Tests demonstrate the control efficacy of the compounds of this invention against specific weeds. The weed control afforded by the compounds is not limited, however, to these species. See Index Table A for compound descriptions. The following abbreviations are used in the Index Table which follows: CF₃ is trifluoromethyl and Ph is phenyl. (*R*) or (*S*) denotes the absolute chirality of the asymmetric carbon center.

The abbreviation “(d)” indicates that the compound appeared to decompose on melting. The abbreviation “Cmpd. No.” stands for “Compound Number”. The abbreviation “Ex.” stands for “Example” and is followed by a number indicating in which example the compound is prepared.

5

INDEX TABLE A (1)



Cmpd. No.	Q ¹	Q ²	R ⁵	m.p. (°C)
1	Ph(4-F)	Ph(2-Cl)	H	173–175
2	Ph(4-F)	Ph(2-F)	H	184–186
3	Ph(3,4-di-F)	Ph(2-Cl)	H	*
4	Ph(3,4-di-F)	Ph(2-F)	H	*
5	Ph(4-F)	Ph(2-CH ₃)	H	172–174
6 (Ex. 1)	Ph(4-F)	Ph(2-CF ₃)	H	144–146**
7	Ph(4-CF ₃)	Ph(2,3-di-F)	H	188–191
8	Ph(4-CF ₃)	Ph(2-F)	H	180–183
9	Ph(4-CF ₃)	Ph(2-CF ₃)	H	142–145
10	Ph(4-CF ₃)	Ph(3-F,2-CF ₃)	H	146–150
11	Ph(3-CF ₃)	Ph(2-F)	H	92–95
12	Ph(3-CF ₃)	Ph(2,3-di-F)	H	103–107
13	Ph(3-CF ₃)	Ph(3-F,2-CF ₃)	H	99–103
14	Ph(3-CF ₃)	Ph(2-CF ₃)	H	73–75
15	Ph(4-F,3-CF ₃)	Ph(2,3-di-F)	H	133–136
16	Ph(4-F,3-CF ₃)	Ph(2-F)	H	154–157
17	Ph(4-F,3-CF ₃)	Ph(2-CF ₃)	H	104–106
18	Ph(4-F,3-CF ₃)	Ph(3-F,2-CF ₃)	H	118–121
19	Ph(4-F)	Ph(2-F)	CH ₃	104–108
20	Ph(4-F)	Ph(2-F)	CH ₃	152–156
21	Ph(4-F)	Ph(2-CF ₃)	CH ₃	382 (M+H)
22	Ph(4-F)	Ph(2,3-di-F)	CH ₃	96–100

(1) Substituents in the 3 and 4 positions of the butyrolactone ring, i.e. C(O)N(Q²)H and Q¹, respectively, are predominately in the *trans* configuration. In some instances the presence of minor

amounts of the *cis* isomer can be detected by NMR. In this structure of Formula 1, each R², R³ and R⁴ is H.

* See Index Table B for ¹H NMR data.

** See Synthesis Example for ¹H NMR data.

5

INDEX TABLE B

Cmpd. No.	¹ H NMR Data (DMSO-d ₆ solution unless indicated otherwise) ^a
3	δ 10.03 (s, 1H), 7.72–7.69 (m, 1H), 7.64–7.59 (m, 1H), 7.51–7.43 (m, 2H), 7.35–7.19 (m, 3H), 4.74–4.70 (t, <i>J</i> = 16.4Hz, 1H), 4.32–4.19 (m, 3H).
4	δ 10.23 (s, 1H), 7.91–7.86 (m, 1H), 7.63–7.57 (m, 1H), 7.49–7.42 (m, 1H), 7.29–7.16 (m, 4H), 4.74–4.70 (t, <i>J</i> = 8.4 Hz, 1H), 4.35–4.22 (m, 3H).

^a ¹H NMR data are in ppm downfield from tetramethylsilane. Couplings are designated by (s)-singlet, (d)-doublet, (t)-triplet, (m)-multiplet.

BIOLOGICAL EXAMPLES OF THE INVENTION

TEST A

10 Seeds of plant species selected from barnyardgrass (*Echinochloa crus-galli*), crabgrass (large crabgrass, *Digitaria sanguinalis*), kochia (*Kochia scoparia*), ragweed (common ragweed, *Ambrosia elatior*), morningglory (*Ipomoea* spp.), velvetleaf (*Abutilon theophrasti*), ryegrass, Italian (Italian ryegrass, *Lolium multiflorum*), foxtail, giant (giant foxtail, *Setaria faberii*), wheat (*Triticum aestivum*), corn (*Zea mays*), and pigweed (*Amaranthus retroflexus*),
 15 were planted into a blend of loam soil and sand and treated preemergence with a directed soil spray using test chemicals formulated in a non-phytotoxic solvent mixture which included a surfactant.

At the same time, plants selected from these crop and weed species and also blackgrass (*Alopecurus myosuroides*), and galium (catchweed bedstraw, *Galium aparine*), were planted
 20 in pots containing the same blend of loam soil and sand and treated with postemergence applications of test chemicals formulated in the same manner. Plants ranged in height from 2 to 10 cm and were in the one- to two-leaf stage for the postemergence treatment. Treated plants and untreated controls were maintained in a greenhouse for approximately 10 d, after which time all treated plants were compared to untreated controls and visually evaluated for
 25 injury. Plant response ratings, summarized in Table A, are based on a 0 to 100 scale where 0 is no effect and 100 is complete control. A dash (–) response means no test result.

	Table A	Compounds				
		1	2	4	5	6
	1000 g ai/ha					
	Postemergence					
30	Barnyardgrass	20	0	70	50	20
	Corn	0	0	–	0	0
	Crabgrass	60	40	90	80	80

	Foxtail, Giant	40	0	70	60	80
	Morningglory	0	0	0	0	0
	Pigweed	20	0	0	0	0
	Velvetleaf	0	0	0	0	0
5	Wheat	0	0	0	0	0

Table A		Compounds													
500 g ai/ha		3	7	8	9	10	11	12	13	14	15	16	17	18	19
Postemergence															
	Barnyardgrass	0	20	20	0	0	40	40	0	20	70	50	0	0	80
10	Blackgrass	-	0	0	0	0	0	0	0	30	20	0	0	0	0
	Corn	-	0	0	0	0	0	0	0	0	20	0	0	0	0
	Crabgrass	70	-	-	-	-	-	-	-	-	-	-	-	-	-
	Foxtail, Giant	60	50	40	70	70	60	70	40	70	70	40	20	30	40
	Galium	-	0	0	0	0	0	0	0	30	20	0	0	0	0
15	Kochia	-	0	0	0	0	0	0	0	0	0	0	0	0	0
	Morningglory	0	-	-	-	-	-	-	-	-	-	-	-	-	-
	Pigweed	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	Ragweed	-	0	0	0	0	0	0	0	0	0	0	0	0	0
	Ryegrass, Italian	-	0	0	0	0	0	0	0	0	20	0	0	0	0
20	Velvetleaf	0	-	-	-	-	-	-	-	-	-	-	-	-	-
	Wheat	0	0	0	0	0	20	0	0	0	20	0	0	0	0

Table A		Compounds	
500 g ai/ha		20	21
Postemergence			
25	Barnyardgrass	0	30
	Blackgrass	0	30
	Corn	0	0
	Crabgrass	-	-
	Foxtail, Giant	0	0
30	Galium	0	30
	Kochia	0	0
	Morningglory	-	-
	Pigweed	0	0
	Ragweed	0	40
35	Ryegrass, Italian	0	0
	Velvetleaf	-	-
	Wheat	0	0

Table A		Compounds													
125 g ai/ha		3	7	8	9	10	11	12	13	14	15	16	17	18	19
Postemergence															
5	Barnyardgrass	0	0	0	0	0	10	40	0	0	30	20	0	0	20
	Blackgrass	-	0	0	0	0	0	0	0	0	0	0	0	0	0
	Corn	-	0	0	0	0	0	0	0	0	0	0	0	0	0
	Crabgrass	60	-	-	-	-	-	-	-	-	-	-	-	-	-
	Foxtail, Giant	50	20	20	30	30	20	30	20	20	30	20	0	0	20
10	Galium	-	0	0	0	0	0	0	0	0	0	0	0	0	0
	Kochia	-	0	0	0	0	0	0	0	0	0	0	0	0	0
	Morningglory	0	-	-	-	-	-	-	-	-	-	-	-	-	-
	Pigweed	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	Ragweed	-	0	0	0	0	0	0	0	0	0	0	0	0	0
15	Ryegrass, Italian	-	0	0	0	0	0	0	0	0	0	0	0	0	0
	Velvetleaf	0	-	-	-	-	-	-	-	-	-	-	-	-	-
	Wheat	0	0	0	0	0	0	0	0	0	0	0	0	0	0

Table A	Compounds			Table A	Compound
125 g ai/ha	20	21	22	31 g ai/ha	22
Postemergence					
Barnyardgrass	0	0	90	Barnyardgrass	40
Blackgrass	0	0	0	Blackgrass	0
Corn	0	0	0	Corn	0
Crabgrass	-	-	-	Foxtail, Giant	30
Foxtail, Giant	0	0	60	Galium	0
Galium	0	0	20	Kochia	0
Kochia	0	0	0	Pigweed	0
Morningglory	-	-	-	Ragweed	0
Pigweed	0	0	0	Ryegrass, Italian	20
Ragweed	0	0	0	Wheat	0
Ryegrass, Italian	0	0	20		
Velvetleaf	-	-	-		
Wheat	0	0	0		

Table A		Compounds				
1000 g ai/ha		1	2	4	5	6
Preemergence						
20	Barnyardgrass	0	30	70	60	20
	Corn	0	0	0	0	0

70

	Crabgrass	70	80	90	80	80									
	Foxtail, Giant	40	50	80	70	80									
	Morningglory	0	0	0	0	0									
	Pigweed	0	0	0	0	0									
5	Velvetleaf	0	0	0	0	0									
	Wheat	0	0	0	0	0									

Table A Compounds

	500 g ai/ha	3	7	8	9	10	11	12	13	14	15	16	17	18	19
	Preemergence														
10	Barnyardgrass	30	30	20	0	0	80	30	0	20	70	60	0	0	90
	Corn	0	-	-	-	-	-	-	-	-	-	-	-	-	-
	Crabgrass	50	-	-	-	-	-	-	-	-	-	-	-	-	-
	Foxtail, Giant	80	80	40	80	50	80	60	70	90	90	80	60	40	90
	Kochia	-	0	0	20	0	0	20	0	20	0	0	0	0	0
15	Morningglory	0	-	-	-	-	-	-	-	-	-	-	-	-	-
	Pigweed	0	0	0	0	0	0	0	0	0	0	0	0	0	20
	Ragweed	-	0	0	0	0	0	0	0	0	0	0	0	0	0
	Ryegrass, Italian	-	0	0	20	0	0	20	0	20	40	0	0	0	0
	Velvetleaf	0	-	-	-	-	-	-	-	-	-	-	-	-	-
20	Wheat	0	-	-	-	-	-	-	-	-	-	-	-	-	-

Table A Compounds

	125 g ai/ha	3	7	8	9	10	11	12	13	14	15	16	17	18	19
	Preemergence														
	Barnyardgrass	0	0	0	0	0	30	20	0	20	40	30	0	0	30
25	Corn	0	-	-	-	-	-	-	-	-	-	-	-	-	-
	Crabgrass	40	-	-	-	-	-	-	-	-	-	-	-	-	-
	Foxtail, Giant	70	30	20	30	40	70	30	40	70	70	50	20	0	20
	Kochia	-	0	0	0	0	0	0	0	0	0	0	0	0	-
	Morningglory	0	-	-	-	-	-	-	-	-	-	-	-	-	-
30	Pigweed	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	Ragweed	-	0	0	0	0	0	0	0	0	0	0	0	0	0
	Ryegrass, Italian	-	0	0	0	0	0	0	0	0	0	0	0	0	0
	Velvetleaf	0	-	-	-	-	-	-	-	-	-	-	-	-	-
	Wheat	0	-	-	-	-	-	-	-	-	-	-	-	-	-


35 TEST B

Plant species in the flooded paddy test selected from rice (*Oryza sativa*), sedge, umbrella (small-flower umbrella sedge, *Cyperus difformis*), duck salad (*Heteranthera*

CLAIMS

The Claims defining the invention are as follows:

- 5 1. A compound selected from Formula 1, *N*-oxides and salts thereof,



wherein

- 10 Q^1 is a phenyl ring; or a 5- to 6-membered heteroaromatic ring containing ring members selected from carbon atoms and 1 to 4 heteroatoms independently selected from up to 2 O, up to 2 S and up to 4 N atoms;
- Q^2 is a phenyl ring; or a 5- to 6-membered heteroaromatic ring containing ring members selected from carbon atoms and 1 to 4 heteroatoms independently selected from up to 2 O, up to 2 S and up to 4 N atoms;
- 15 Y^1 and Y^2 are each independently O, S or NR^6 ;
- R^1 is H, hydroxy, amino, C_1 - C_4 alkyl, C_1 - C_4 haloalkyl, C_2 - C_4 alkenyl, C_3 - C_4 alkynyl, C_2 - C_4 alkoxyalkyl, C_2 - C_4 haloalkoxyalkyl, C_2 - C_4 alkylthioalkyl, C_2 - C_4 alkylsulfanylalkyl, C_2 - C_4 alkylsulfonylalkyl, C_2 - C_4 alkylcarbonyl, C_2 - C_4 haloalkylcarbonyl, C_4 - C_6 cycloalkylcarbonyl, C_2 - C_4 alkoxy carbonyl, C_2 - C_4 haloalkoxy carbonyl or C_4 - C_6 cycloalkoxy carbonyl;
- 20 R^2 and R^3 are each independently H, halogen, hydroxy, C_1 - C_4 alkyl, C_1 - C_4 haloalkyl or C_1 - C_4 alkoxy; or
- R^2 and R^3 are taken together with the carbon atom to which they are bonded to form a C_3 - C_7 cycloalkyl ring;
- 25 R^4 and R^5 are each independently H, halogen, hydroxy, C_1 - C_4 alkyl or C_1 - C_4 alkoxy;
- each R^6 is independently H, cyano, C_1 - C_4 alkyl, C_1 - C_4 haloalkyl, C_1 - C_4 alkoxy, C_1 - C_4 haloalkoxy, $-(C=O)CH_3$ or $-(C=O)CF_3$;
- each R^7 and R^{10} is independently halogen, cyano, nitro, C_1 - C_4 alkyl, C_1 - C_4 haloalkyl, C_2 - C_4 alkenyl, C_2 - C_4 haloalkenyl, C_2 - C_4 alkynyl, C_2 - C_4 haloalkynyl, C_1 - C_4 nitroalkyl, C_2 - C_4 nitroalkenyl, C_2 - C_4 alkoxyalkyl, C_2 - C_4 haloalkoxyalkyl, C_3 - C_4 cycloalkyl, C_3 - C_4 halocycloalkyl, cyclopropylmethyl, methylcyclopropyl, C_1 - C_4 alkoxy, C_1 - C_4 haloalkoxy, C_2 - C_4 alkenyloxy, C_2 - C_4 haloalkenyloxy,
- 30

C₃-C₄ alkynyloxy, C₃-C₄ haloalkynyloxy, C₃-C₄ cycloalkoxy, C₁-C₄ alkylthio, C₁-C₄ haloalkylthio, C₁-C₄ alkylsulfinyl, C₁-C₄ haloalkylsulfinyl, C₁-C₄ alkylsulfonyl, C₁-C₄ haloalkylsulfonyl, hydroxy, formyl, C₂-C₄ alkylcarbonyl, C₂-C₄ alkylcarbonyloxy, C₁-C₄ alkylsulfonyloxy, C₁-C₄ haloalkylsulfonyloxy, amino, C₁-C₄ alkylamino, C₂-C₄ dialkylamino, formylamino, C₂-C₄ alkylcarbonylamino, -SF₅, -SCN, C₃-C₄ trialkylsilyl, trimethylsilylmethyl or trimethylsilylmethoxy; and
each R⁹ and R¹¹ is independently C₁-C₂ alkyl or C₂-C₃ alkoxy carbonyl.

2. The compound according to Claim 1 wherein Q¹ is a phenyl ring and the phenyl ring is substituted with 1 to 4 substituents independently selected from R⁷.

3. The compound according to Claim 1 wherein Q¹ is a 5- to 6 membered heteroaromatic ring and the 5- to 6 membered heteroaromatic ring is substituted with up to 4 substituents independently selected from R⁷ on carbon atom ring members and selected from R⁹ on nitrogen atom ring members.

4. The compound according to any one of the preceding claims wherein Q² is a phenyl ring and the phenyl ring is substituted with up to 5 substituents independently selected from R¹⁰.

5. The compound according to any one of Claims 1 to 3 wherein Q² is a 5- to 6 membered heteroaromatic ring and the 5- to 6 membered heteroaromatic ring is substituted with up to 5 substituents independently selected from R¹⁰ on carbon atom ring members and selected from R¹¹ on nitrogen atom ring members.

6. The compound according to any one of the preceding claims wherein

Y¹ is O;

Y² is O;

R¹ is H;

R² is H;

R³ is H;

R⁴ is H; and

R⁵ is H or CH₃.

7. The compound according to Claim 6 wherein

Q¹ is a phenyl ring substituted with 1 to 3 substituents independently selected from R⁷; and

Q² is a phenyl ring substituted with 1 to 3 substituents independently selected from R¹⁰.

8. The compound according to Claim 7 wherein
each R⁷ is independently halogen, cyano, C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₁-C₄
alkoxy or C₁-C₄ haloalkoxy; and
5 each R¹⁰ is independently halogen, cyano, C₁-C₄ alkyl, C₁-C₄ haloalkyl or C₁-C₄
alkylsulfonyl.
9. The compound according to Claim 8 wherein
Q¹ is a phenyl ring substituted with 1 substituent selected from R⁷ at the para
10 position or substituted with 2 substituents independently selected from R⁷
wherein one substituent is at the para position and the other substituent is at a
meta position; and
Q² is a phenyl ring substituted with 1 substituent selected from R¹⁰ at an ortho
15 position or substituted with 2 substituents independently selected from R¹⁰
wherein one substituent is at an ortho position and the other substituent is at the
adjacent meta position.
10. The compound according to Claim 9 wherein
each R⁷ is independently F or CF₃; and
20 each R¹⁰ is F.
11. The compound according to Claim 1 selected from the group consisting of
4-(4-fluorophenyl)tetrahydro-2-oxo-*N*-[2-(trifluoromethyl)phenyl]-3-
furancarboxamide;
25 (3*R*,4*S*)-4-(4-fluorophenyl)tetrahydro-2-oxo-*N*-[2-(trifluoromethyl)phenyl]-3-
furancarboxamide; and
(3*S*,4*R*)-4-(4-fluorophenyl)tetrahydro-2-oxo-*N*-[2-(trifluoromethyl)phenyl]-3-
furancarboxamide.
- 30 12. A herbicidal composition comprising a compound according to any one of the preceding
Claims and at least one component selected from the group consisting of surfactants, solid
diluent and liquid diluent.
- 35 13. A herbicidal composition comprising a compound according to any one of the preceding
Claims, at least one additional active ingredient selected from the group consisting of other
herbicides and herbicide safeners, and at least one component selected from the group
consisting of surfactants, solid diluent and liquid diluent.
- 40 14. A herbicidal mixture comprising (a) a compound according to any one of the preceding
Claims, and (b) at least one additional active ingredient selected from (b1) photosystem II
inhibitors, (b2) acetohydroxy acid synthase (AHAS) inhibitors, (b3) acetyl-CoA carboxylase
(ACCase) inhibitors, (b4) auxin mimics, (b5) 5-enol-pyruvylshikimate-3-phosphate (EPSP)
synthase inhibitors, (b6) photosystem I electron diverters, (b7) protoporphyrinogen oxidase

(PPO) inhibitors, (b8) glutamine synthetase (GS) inhibitors, (b9) very long chain fatty acid (VLCFA) elongase inhibitors, (b10) auxin transport inhibitors, (b11) phytoene desaturase (PDS) inhibitors, (b12) 4-hydroxyphenyl-pyruvate dioxygenase (HPPD) inhibitors, (b13) homogentisate solanesyltransferase (HST) inhibitors, (b14) cellulose biosynthesis inhibitors, (b15) other herbicides including mitotic disruptors, organic arsenicals, asulam, bromobutide, cinmethylin, cumyluron, dazomet, difenzoquat, dymron, etobenzanid, flurenol, fosamine, fosamine-ammonium, hydantocidin, metam, methyl dymron, oleic acid, oxaziclomefone, pelargonic acid and pyributicarb, (b16) herbicide safeners, and salts of compounds of (b1) through (b16).

15. A method for controlling the growth of undesired vegetation comprising contacting the vegetation or its environment with a herbicidally effective amount of a compound according to any one of the preceding Claims.