



HU000029932T2

(19) **HU**(11) Lajstromszám: **E 029 932**(13) **T2****MAGYARORSZÁG**
Szellemi Tulajdon Nemzeti Hivatala

EURÓPAI SZABADALOM SZÖVEGÉNEK FORDÍTÁSA

(21) Magyar ügyszám: **E 12 824674**(51) Int. Cl.: **A01N 43/60**

(2006.01)

(22) A bejelentés napja: **2012. 08. 15.****A01N 43/90**

(2006.01)

A01P 13/00

(2006.01)

(96) Az európai bejelentés bejelentési száma:

EP 20120824674

(86) A nemzetközi (PCT) bejelentési szám:

PCT/US 12/050862

(97) Az európai bejelentés közzétételi adatai:

EP 2744337 A1 2013. 02. 21.

(87) A nemzetközi közzétételi szám:

WO 13025754

(97) Az európai szabadalom megadásának meghirdetési adatai:

EP 2744337 B1 2016. 05. 18.

(30) Elsőbbségi adatok:

201161523884 P**2011. 08. 16.****US**

(73) Jogosult(ak):

**Dow AgroSciences LLC, Indianapolis, IN
46268 (US)**

(72) Feltaláló(k):

MANN, Richard K., Franklin, IN 46131 (US)

(74) Képviselő:

**Danubia Szabadalmi és Jogi Iroda Kft.,
Budapest**(54) **Penoxszulamot és floraszulamot tartalmazó szinergetikus herbicid készítmény**

Az európai szabadalom ellen, megadásának az Európai Szabadalmi Közlönyben való meghirdetésétől számított kilenc hónapon belül, felszólalást lehet benyújtani az Európai Szabadalmi Hivatalnál. (Európai Szabadalmi Egyezmény 99. cikk(1))

A fordítást a szabadalmas az 1995. évi XXXIII. törvény 84/H. §-a szerint nyújtotta be. A fordítás tartalmi helyességét a Szellemi Tulajdon Nemzeti Hivatala nem vizsgálta.



(11)

EP 2 744 337 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:
18.05.2016 Bulletin 2016/20

(51) Int Cl.:
A01N 43/60 (2006.01) **A01N 43/90** (2006.01)
A01P 13/00 (2006.01)

(21) Application number: **12824674.1**

(86) International application number:
PCT/US2012/050862

(22) Date of filing: **15.08.2012**

(87) International publication number:
WO 2013/025754 (21.02.2013 Gazette 2013/08)

(54) **SYNERGISTIC HERBICIDAL COMPOSITION CONTAINING PENOXSULAM AND FLORASULAM**
SYNERGISTISCHE HERBIZIDZUSAMMENSETZUNG MIT PENOXSULAM UND FLORASULAM
COMPOSITION HERBICIDE SYNERGIQUE CONTENANT DU PÉNOXSULAME ET DU
FLORASULAM

(84) Designated Contracting States:
**AL AT BE BG CH CY CZ DE DK EE ES FI FR GB
GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO
PL PT RO RS SE SI SK SM TR**

(30) Priority: **16.08.2011 US 201161523884 P**

(43) Date of publication of application:
25.06.2014 Bulletin 2014/26

(73) Proprietor: **Dow AgroSciences LLC**
Indianapolis, IN 46268 (US)

(72) Inventor: **MANN, Richard K.**
Franklin, IN 46131 (US)

(74) Representative: **f & e patent**
Fleischer, Engels & Partner mbB, Patentanwälte
Braunsberger Feld 29
51429 Bergisch Gladbach (DE)

(56) References cited:
US-A1- 2010 099 564 **US-A1- 2010 099 564**
US-A1- 2011 098 182 **US-A1- 2011 098 182**
US-A1- 2011 190 134 **US-A1- 2011 190 134**

• **JOHNSON T C ET AL:**
"Penoxsulam-Structure-activity relationships of
triazolopyrimidine sulfonamides", BIOORGANIC
& MEDICINAL CHEMISTRY, PERGAMON, GB, vol.
17, no. 12, 15 June 2009 (2009-06-15) , pages
4230-4240, XP026152235, ISSN: 0968-0896, DOI:
10.1016/J.BMC.2009.02.010 [retrieved on
2009-02-14]

EP 2 744 337 B1

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

DescriptionField of the Invention

5 [0001] This invention concerns a synergistic herbicidal composition containing (a) penoxsulam and (b) florasulam for controlling growth of undesirable vegetation, particularly in multiple crops, including rice, cereal and grain crops (including wheat, barley and corn), turf, industrial vegetation management (IVM), sugar cane, range and pasture, and tree and vine orchards.

10 Background of the Invention

[0002] The protection of crops from weeds and other vegetation which inhibit crop growth is a constantly recurring problem in agriculture. To help combat this problem, researchers in the field of synthetic chemistry have produced an extensive variety of chemicals and chemical formulations effective in the control of such unwanted growth. Chemical 15 herbicides of many types have been disclosed in the literature and a large number are in commercial use.

[0003] In some cases, herbicidal active ingredients have been shown to be more effective in combination than when applied individually and this is referred to as "synergism." As described in the Herbicide Handbook of the Weed Science Society of America, Ninth Edition, 2007, p. 429, "'synergism' [is] an interaction of two or more factors such that the effect 20 when combined is greater than the predicted effect based on the response to each factor applied separately." The present invention is based on the discovery that florasulam and penoxsulam, already known individually for their herbicidal efficacy, display a synergistic effect when applied in combination.

Summary of the Invention

25 [0004] The present invention concerns a synergistic herbicidal mixture comprising an herbicidally effective amount of (a) penoxsulam and (b) florasulam. The compositions may also contain an agriculturally acceptable adjuvant or carrier.

[0005] The present invention also concerns a method of controlling the growth of undesirable vegetation in multiple crops, including rice, cereal and grain crops, turf, IVM, sugar cane and tree and vine orchards, and the use of this synergistic composition.

30 [0006] The species spectrum of the compounds of the synergistic mixture, i.e., the weed species which the respective compounds control, are broad and highly complementary. These synergistic mixtures are particularly useful for the control of key weeds, e.g., chamomile, (*Anthemis cotula* L., ANTOR), field marigold (*Calendula arvensis*, CLDAR), henbit (*Lamium amplexicaule* L., LAMAM), annual sowthistle (*Sonchus oleraceus* L., SONOL), common mallow (*Malva neglecta* Wallr., MALNE), corn salad (*Valerianella echinata*, VLLEC), horseweed (*Conyza canadensis* (L.) Cronq., ERICA), field 35 madder (*Sherardia arvensis* L., SHRAR), and sowthistle (*Sonchus* species, SONSS) at application rates equal to or lower than the rates of the individual compounds.

Detailed Description of the Invention

40 [0007] Penoxsulam is the common name for (2-(2,2-difluoroethoxy)-N-(5,8-dimethoxy-[1,2,4]triazolo[1,5-c]pyrimidin-2-yl)-6-(trifluoromethyl)benzenesulfonamide. Its herbicidal activity is described in The Pesticide Manual, Fifteenth Edition, 2009. Penoxsulam controls barnyard grass, as well as many broadleaf and sedge weeds in rice, turf, tree nut and vineyard crops, cereal and grain crops, and IVM.

[0008] Florasulam is the common name for N-(2,6-difluorophenyl)-8-fluoro-5-methoxy[1,2,4]triazolo[1,5-c]pyrimidine-2-sulfonamide. Its herbicidal activity is described in The Pesticide Manual, Fifteenth Edition, 2009. Florasulam provides postemergent control of broadleaf weeds and cruciferae in cereals and maize.

[0009] The term herbicide is used herein to mean an active ingredient that kills, controls or otherwise adversely modifies the growth of plants. An herbicidally effective or vegetation controlling amount is an amount of active ingredient which causes an adversely modifying effect and includes deviations from natural development, killing, regulation, desiccation, retardation, and the like. The terms plants and vegetation include germinant seeds, emerging seedlings, plants emerging 50 from vegetative propagules, and established vegetation.

[0010] Herbicidal activity is exhibited by the compounds of the synergistic mixture when they are applied directly to the plant or to the locus of the plant at any stage of growth or before planting or emergence. The effect observed depends upon the plant species to be controlled, the stage of growth of the plant, the application parameters of dilution and spray drop size, the particle size of solid components, the environmental conditions at the time of use, the specific compound employed, the specific adjuvants and carriers employed, the soil type, and the like, as well as the amount of chemical applied. These and other factors can be adjusted as is known in the art to promote non-selective or selective herbicidal action. Generally, it is preferred to apply the composition of the present invention preemergence to early postemergence

to relatively immature undesirable vegetation to achieve the maximum control of weeds.

[0011] In the composition of this invention, the active ingredient ratio (weight-to-weight, wt:wt) of penoxsulam to florasulam at which the herbicidal effect is synergistic lies within the range of between about 17:1 to 1:1, with a ratio of about 7:1 being preferred.

[0012] The rate at which the synergistic composition is applied will depend upon the particular type of weed to be controlled, the degree of control required, and the timing and method of application. In general, the composition of the invention can be applied at an application rate of between about 10 grams active ingredient per hectare (gai/ha) and about 110 gai/ha based on the total amount of active ingredients in the composition. An application rate between about 10 gai/ha and about 70 gai/ha is preferred. In an especially preferred embodiment of the invention, florasulam is applied at a rate between about 3 gai/ha and about 10 gai/ha, and penoxsulam is applied at a rate between about 10 gai/ha and about 50 gai/ha.

[0013] The components of the synergistic mixture of the present invention can be applied either separately or as part of a multipart herbicidal system.

[0014] The synergistic mixture of the present invention can be applied in conjunction with one or more other herbicides to control a wider variety of undesirable vegetation. When used in conjunction with other herbicides, the composition can be formulated with the other herbicide or herbicides, tank mixed with the other herbicide or herbicides or applied sequentially with the other herbicide or herbicides. Some of the herbicides that can be employed in conjunction with the synergistic composition of the present invention include: 4-CPA, 4-CPB, 4-CPP, 2,4-D, 3,4-DA, 2,4-DB, 3,4-DB, 2,4-DEB, 2,4-DEP, 3,4-DP, 2,3,6-TBA, 2,4,5-T, 2,4,5-TB, acetochlor, acifluorfen, aclonifen, acrolein, alachlor, allidochlor, aloxydim, allyl alcohol, alorac, ametridione, ametryn, amibuzin, amicarbazone, amidosulfuron, aminocyclopyrachlor, aminopyralid, amiprofos-methyl, amitrole, ammonium sulfamate, anilofos, anisuron, asulam, atraton, atrazine, azafenidin, azimsulfuron, aziprotryne, barban, BCPC, beflubutamid, benazolin, bencarbazone, benfluralin, benfuresate, benzulfuron, bensulide, bentazone, benzadox, benzfendizone, benzipram, benzobicyclon, benzofenap, benzofluor, benzoylprop, benzthiazuron, bicyclopyrone, bifenoxy, bilanafos, bispyribac, borax, bromacil, bromobonil, bromobutide, bromfenoxim, bromoxynil, brompyrazon, butachlor, butafenacil, butamifos, butenachlor, buthidazole, buthiuron, butralin, butoxydim, buturon, butylate, cacodylic acid, cafenstrole, calcium chloride, calcium cyanamide, cambendichlor, carbasulam, carbetamide, carboxazole chlorprocarb, carfentrazone, CDEA, CEPC, chlomethoxyfen, chloramben, chloranocryl, chlorazifop, chlorazine, chlorbromuron, chlorbufam, chloretruron, chlorfenac, chlorfenprop, chlorflurazole, chlorflurenol, chloridazon, chlorimuron, chlornitrofen, chlorpon, chlorotoluron, chloroxuron, chloroxynil, chlorpropham, chlorsulfuron, chlorthal, chlorthiamid, cinidon-ethyl, cinmethylin, cinosulfuron, cisanilide, clethodim, cioldinate, clodinafop, clofop, clomazone, clomeprop, cloprop, cloproxydim, clopyralid, cloransulam, CMA, copper sulfate, CPMF, CPPC, credazine, cresol, cumyluron, cyanatry, cyanazine, cycloate, cyclosulfamuron, cycloxydim, cycluron, cyhalofop, cyperquat, cyprazine, cyprazole, cypromid, daimuron, dalapon, dazomet, delachlor, desmedipham, desmetryn, di-allate, dicamba, dichlobenil, dichloralurea, dichlormate, dichlorprop, dichlorprop-P, diclofop, diclosulam, diethamquat, diethylatyl, difenopenten, difenoxyuron, difenzoquat, diflufenican, diflufenzopyr, dimefuron, dimepiperate, dimethachlor, dimethametryn, dimethenamid, dimethenamid-P, dimexano, dimidazon, dinitramine, dinofenate, dinoprop, dinosam, dinoseb, dinoterb, diphenamid, dipropetryn, diquat, disul, dithiopyr, diuron, DMPA, DNOC, DSMA, EBEP, eglinazine, endothal, erpronaz, EPTC, erbon, esprocarb, ethafluralin, ethamsulfuron, ethidimuron, ethiolate, ethofumesate, ethoxyfen, ethoxysulfuron, etinofen, etnipromid, etobenzanid, EXD, fenasulam, fenoprop, fenoxaprop, fenoxaprop-P, fenoxasulfone, fenteracol, fenthiaprop, fentrazamide, fenuron, ferrous sulfate, flamprop, flamprop-M, fluazifop, fluazifop-P, fluazolate, flucarbazone, flucetosulfuron, fluchloralin, flufenacet, flufenican, flufenpyr, flumetsulam, flumezin, flumiclorac, flumioxazin, flumipropyn, fluometuron, fluorodifen, fluoroglycofen, fluoromidine, fluoronitrofen, fluothiuron, flupoxam, flupropacil, flupropanate, fluprysulfuron, fluridone, flurochloridone, fluroxypyr, flurtamone, fluthiacet, fomesafen, foramsulfuron, fosamine, furyloxyfen, glufosinate, glufosinate-P, glyphosate, halosafen, halosulfuron, haloxydine, haloxyfop, haloxyfop-P, hexachloroacetone, hexaflurate, hexazinone, imazamethabenz, imazamox, imazapic, imazapyr, imazaquin, imazethapyr, imazosulfuron, indanofan, indaziflam, iodobonil, iodomethane, iodosulfuron, iofensulfuron, ioxynil, ipazine, ipfencarbazone, iprymidam, isocarbamid, isocil, isomethiozin, isonoruron, isopolinate, isopropalin, isoproturon, isouron, isoxaben, isoxachlortole, isoxaflutole, isoxapryifop, karbutilate, ketospiradox, lactofen, lenacil, linuron, MAA, MAMA, MCPA, MCPA-thioethyl, MCPB, mecoprop, mecoprop-P, medinoterb, mefenacet, mefluidide, mesoprazine, mesosulfuron, mesotriione, metam, metamifop, metamitron, metazachlor, metazosulfuron, metflurazon, methabenzthiazuron, methalpropalin, methazole, methiobencarb, methiozolin, methiuron, methometon, methoprotyne, methyl bromide, methyl isothiocyanate, methyldymron, metobenzuron, metobromuron, metolachlor, metosulam, metoxuron, metribuzin, metsulfuron, molinate, monalide, monisuron, monochloroacetic acid, monolinuron, monuron, morfamquat, MSMA, naproanilide, napropamide, naptalam, neburon, nicosulfuron, nipyrapclofen, nitralin, nitrofen, nitrofluorfen, norflurazon, noruron, OCH, orbencarb, *ortho*-dichlorobenzene, orthosulfamuron, oryzalin, oxadiargyl, oxadiazon, oxapyrazon, oxasulfuron, oxaziclofene, oxyfluorfen, parafluron, paraquat, pebulate, pelargonic acid, pendimethalin, pentachlorophenol, pentanochlor, pentozacone, perfluidone, pethoxamid, phenisopham, phenmedipham, phenmedipham-ethyl, phenobenzuron, phenylmercury acetate, picloram, picolinafen, pinoxaden, piperophos, potassium arsenite, potassium azide,

potassium cyanate, pretilachlor, primisulfuron, procyzazine, prodiamine, profluazol, profluralin, profoxydim, progliazine, prometon, prometryn, propachlor, propanil, propaquazafop, propazine, prophan, propisochlor, propoxycarbazone, propisulfuron, propyzamide, prosulfalin, prosulfocarb, prosulfuron, proxan, prynachlor, pydanon, pyraclonil, pyraflufen, pyrasulfotole, pyrazolynate, pyrazosulfuron, pyrazoxyfen, pyribenzoxim, pyributicarb, pyriclor, pyridafol, pyridate, pyriflalid, pyriminobac, pyrimisulfan, pyriothiobac, pyroxasulfone, pyroxslam, quinclorac, quinmerac, quinoclamine, quinonamid, quizalofop, quizalofop-P, rhodethanil, rimsulfuron, saflufenacil, S-metolachlor, sebutylazine, secbumeton, sethoxydim, siduron, simazine, simeton, simetryn, SMA, sodium arsenite, sodium azide, sodium chloride, sulcotrione, sulfallate, sulfentrazone, sulfometuron, sulfosulfuron, sulfuric acid, sulglycapin, swep, TCA, tebutam, tebuthiuron, tefuryltrione, tembotrione, tepraloxydim, terbacil, terbucarb, terbuchlor, terbumeton, terbutylazine, terbutryn, tetrafluron, thenylchlor, thiazafluron, thiazopyr, thidiazimin, thidiazuron, thiencarbazone-methyl, thifensulfuron, thiobencarb, tiocarbazil, tioclorim, topramezone, tralkoxydim, triafamone, tri-allate, triasulfuron, triaziflam, tribenuron, tricamba, triclopyr, tridiphane, trietazine, trifloxysulfuron, trifluralin, triflusulfuron, trifop, trifopsime, trihydroxytriazine, trimeturon, tripropindan, tritac tritosulfuron, vernolate and xylachlor.

[0015] The synergistic mixture of the present invention can additionally be employed to control undesirable vegetation in many crops that have been made tolerant to or resistant to them or to other herbicides by genetic manipulation or by mutation and selection. The synergistic composition of the present invention can, further, be used in conjunction with 2,4-D, glyphosate, glufosinate, dicamba, sulfonylureas or imidazolinones on 2,4-D tolerant, glyphosate-tolerant, glufosinate-tolerant, dicamba-tolerant, sulfonylurea-tolerant crops or imidazolinone-tolerant crops.

[0016] It is generally preferred to use the synergistic composition of the present invention in combination with herbicides that are selective for the crop being treated and which complement the spectrum of weeds controlled by these compounds at the application rate employed. It is further generally preferred to apply the synergistic composition of the present invention and other complementary herbicides at the same time, either as a combination formulation or as a tank mix.

[0017] The synergistic composition of the present invention can generally be employed in combination with known herbicide safeners, such as benoxacor, benthiocarb, brassinolide, cloquintocet (mexyl), cyometrinil, cyrosulfamate, daimuron, dichlormid, dicyclonon, dietholate, dimepiperate, disulfoton, fenchlorazole-ethyl, fenclorim, flurazole, fluxofenim, furilazole, harpin proteins, isoxadifen-ethyl, mefenpyr-diethyl, mephenate, MG 191, MON 4660, naphthalic anhydride (NA), oxabentrinil, R29148 and N-phenyl-sulfonylbenzoic acid amides, to enhance their selectivity.

[0018] In practice, it is preferable to use the synergistic composition of the present invention in mixtures containing an herbicidally effective amount of the herbicidal components along with at least one agriculturally acceptable adjuvant or carrier. Suitable adjuvants or carriers should not be phytotoxic to valuable crops, particularly at the concentrations employed in applying the compositions for selective weed control in the presence of crops, and should not react chemically with herbicidal components or other composition ingredients. Such mixtures can be designed for application directly to weeds or their locus or can be concentrates or formulations that are normally diluted with additional carriers and adjuvants before application. They can be solids, such as, for example, dusts, granules, water dispersible granules, or wettable powders, or liquids, such as, for example, emulsifiable concentrates, solutions, emulsions or suspensions.

[0019] Suitable agricultural adjuvants and carriers that are useful in preparing the herbicidal mixtures of the invention are well known to those skilled in the art. Some of these adjuvants include, but are not limited to, crop oil concentrate (mineral oil (85%) + emulsifiers (15%)); nonylphenol ethoxylate; benzylcocoalkyldimethyl quaternary ammonium salt; blend of petroleum hydrocarbon, alkyl esters, organic acid, and anionic surfactant; C₉-C₁₁ alkylpolyglycoside; phosphated alcohol ethoxylate; natural primary alcohol (C₁₂-C₁₆) ethoxylate; di-sec-butylphenol EO-PO block copolymer; polysiloxane-methyl cap; nonylphenol ethoxylate + urea ammonium nitrate; emulsified methylated seed oil; tridecyl alcohol (synthetic) ethoxylate (8EO); tallow amine ethoxylate (15 EO); PEG(400) dioleate-99.

[0020] Liquid carriers that can be employed include water and organic solvents. The organic solvents typically used include, but are not limited to, petroleum fractions or hydrocarbons such as mineral oil, aromatic solvents, paraffinic oils, and the like; vegetable oils such as soybean oil, rapeseed oil, olive oil, castor oil, sunflower seed oil, coconut oil, corn oil, cottonseed oil, linseed oil, palm oil, peanut oil, safflower oil, sesame oil, tung oil and the like; esters of the above vegetable oils; esters of monoalcohols or dihydric, trihydric, or other lower polyalcohols (4-6 hydroxy containing), such as 2-ethyl hexyl stearate, n-butyl oleate, isopropyl myristate, propylene glycol dioleate, di-octyl succinate, di-butyl adipate, di-octyl phthalate and the like; esters of mono, di and polycarboxylic acids and the like. Specific organic solvents include toluene, xylene, petroleum naphtha, crop oil, acetone, methyl ethyl ketone, cyclohexanone, trichloroethylene, perchloroethylene, ethyl acetate, amyl acetate, butyl acetate, propylene glycol monomethyl ether and diethylene glycol monomethyl ether, methyl alcohol, ethyl alcohol, isopropyl alcohol, amyl alcohol, ethylene glycol, propylene glycol, glycerine, N-methyl-2-pyrrolidinone, N,N-dimethyl alkylamides, dimethyl sulfoxide, liquid fertilizers and the like. Water is generally the carrier of choice for the dilution of concentrates.

[0021] Suitable solid carriers include talc, pyrophyllite clay, silica, attapulgus clay, kaolin clay, kieselguhr, chalk, diatomaceous earth, lime, calcium carbonate, bentonite clay, Fuller's earth, cottonseed hulls, wheat flour, soybean flour, pumice, wood flour, walnut shell flour, lignin, and the like.

[0022] It is usually desirable to incorporate one or more surface-active agents into the compositions of the present

invention. Such surface-active agents are advantageously employed in both solid and liquid compositions, especially those designed to be diluted with carrier before application. The surface-active agents can be anionic, cationic or nonionic in character and can be employed as emulsifying agents, wetting agents, suspending agents, or for other purposes. Surfactants conventionally used in the art of formulation and which may also be used in the present formulations are described, *inter alia*, in " McCutcheon's Detergents and Emulsifiers Annual," MC Publishing Corp., Ridgewood, New Jersey, 1998 and in "Encyclopedia of Surfactants," Vol. I-III, Chemical Publishing Co., New York, 1980-81. Typical surface-active agents include salts of alkyl sulfates, such as diethanolammonium lauryl sulfate; alkylarylsulfonate salts, such as calcium dodecylbenzenesulfonate; alkylphenol-alkylene oxide addition products, such as nonylphenol-C₁₈ ethoxylate; alcohol-alkylene oxide addition products, such as tridecyl alcohol-C₁₆ ethoxylate; soaps, such as sodium stearate; alkylnaphthalene-sulfonate salts, such as sodium dibutylnaphthalenesulfonate; dialkyl esters of sulfosuccinate salts, such as sodium di(2-ethylhexyl) sulfosuccinate; sorbitol esters, such as sorbitol oleate; quaternary amines, such as lauryl trimethylammonium chloride; polyethylene glycol esters of fatty acids, such as polyethylene glycol stearate; block copolymers of ethylene oxide and propylene oxide; salts of mono- and dialkyl phosphate esters; vegetable or seed oils such as soybean oil, rapeseed/canola oil, olive oil, castor oil, sunflower seed oil, coconut oil, corn oil, cottonseed oil, linseed oil, palm oil, peanut oil, safflower oil, sesame oil, tung oil and the like; and esters of the above vegetable oils, particularly methyl esters.

[0023] Oftentimes, some of these materials, such as vegetable or seed oils and their esters, can be used interchangeably as an agricultural adjuvant, as a liquid carrier or as a surface active agent.

[0024] Other adjuvants commonly used in agricultural compositions include compatibilizing agents, antifoam agents, sequestering agents, neutralizing agents and buffers, corrosion inhibitors, dyes, odorants, spreading agents, penetration aids, sticking agents, dispersing agents, thickening agents, freezing point depressants, antimicrobial agents, and the like. The compositions may also contain other compatible components, for example, other herbicides, plant growth regulators, fungicides, insecticides, and the like and can be formulated with liquid fertilizers or solid, particulate fertilizer carriers such as ammonium nitrate, urea and the like.

[0025] The concentration of the active ingredients in the synergistic composition of the present invention is generally from 0.001 to 98 percent by weight. Concentrations from 0.01 to 90 percent by weight are often employed. In compositions designed to be employed as concentrates, the active ingredients are generally present in a concentration from 1 to 98 weight percent, preferably 2 to 90 weight percent. Such compositions are typically diluted with an inert carrier, such as water, before application. The diluted compositions usually applied to weeds or the locus of weeds generally contain 0.0001 to 1 weight percent active ingredient and preferably contain 0.001 to 0.05 weight percent.

[0026] The present compositions can be applied to weeds or their locus by the use of conventional ground or aerial dusters, sprayers, and granule applicators, by addition to irrigation water, and by other conventional means known to those skilled in the art.

[0027] The following examples illustrate the present invention.

Examples

Evaluation of Preemergence Herbicidal Activity of Mixtures under Field Conditions

Methodology

[0028] Field trials were conducted in olive tree crops in Spain using standard herbicide small plot research methodology. Plot size was typical for small plot research in tree crops, varying from 4 to 10 meters (m) wide by 4 to 10 m long. There were 4 replicates per treatment typically utilizing randomized complete block statistical design. Soil types ranged from coarse to medium to fine soil texture. Perennial olive crops were transplanted by hand into orchards as per normal local cultural practices. The olive crop was grown using normal cultural practices for fertilization, watering and maintenance to ensure good growth of the crop and the weeds.

[0029] Treatments were applied by backpack sprayer using either compressed air or CO₂, at spray pressures at 300 kilopascals (kPa). Spray tips were typically Flat Fan Teejet nozzles, such as FanJet 120 or Teejet 11003 VP. Spray volumes were approximately 300 liters per hectare (L/ha). The weed spectrum included, but was not limited to, chamomile, (*Anthemis cotula* L., ANTOR), field marigold (*Calendula arvensis*, CLDAR), henbit (*Lamium amplexicaule* L., LAMAM), annual sowthistle (*Sonchus oleraceus* L., SONOL), common mallow (*Malva neglecta* Wallr., MALNE), corn salad (*Valerianella echinata*, VLLEC), horseweed (*Conyza canadensis* (L.) Cronq., ERICA), field madder (*Sherardia arvensis* L., SHRAR), and sowthistle (*Sonchus* species, SONSS). Treatments were applied preemergence or early postemergence to the stage of the weeds. Olive tree spacings were typically rows that were 7 to 8 m apart, and trees in the row were from 4 to 6 m apart.

[0030] For each treatment, the appropriate formulated product amount to treat the plot area, to achieve the desired application rate, based on unit area of application (hectare), was calculated, measured, and mixed in water prior to

applying with the backpack sprayer

[0031] Treatments were rated as compared to the untreated control plots.

Evaluation

[0032] The treated plots and control plots were rated blind at various intervals after application. Ratings were based of Percent (%) Visual weed control, where 0 corresponds to no injury and 100 corresponds to complete kill.

[0033] Data were collected for all trials and analyzed using various statistical methods.

[0034] Colby's equation was used to determine the herbicidal effects expected from the mixtures (Colby, S. R. Calculation of the synergistic and antagonistic response of herbicide combinations. Weeds 1967, 15, 20-22).

[0035] The following equation was used to calculate the expected activity of mixtures containing two active ingredients, A and B:

$$\text{Expected} = A + B - (A \times B / 100)$$

A = observed efficacy of active ingredient A at the same concentration as used in the mixture;

B = observed efficacy of active ingredient B at the same concentration as used in the mixture.

[0036] The results are summarized in Tables 1 through 3.

Table 1. Control of ANTCO, CLDAR, LAMAM and SONOL by Penoxsulam plus Florasulam at 30-45 Days after Application (DAA) in the Field.

		% Control							
Application Rate (gai/ha)		ANTCO		CLDAR		LAMAM		SONOL	
Penoxsulam	Florasulam	Obs	Ex	Obs	Ex	Obs	Ex	Obs	Ex
10	0	0	-	5	-	0	-	0	-
0	5	50	-	40	-	5	-	50	-
10	5	100	50	90	43	50	5	98	50
13	0	5	-	-	-	0	-	2	-
0	6.3	69	-	-	-	30	-	60	-
13	6.3	100	70	-	-	58	30	98	61
15	0	5	-	-	-	4	-	5	-
0	7.5	95	-	-	-	30	-	60	-
15	7.5	100	95	-	-	80	32	99	62

ANTCO = chamomile, mayweed (*Anthemis cotula* L.)

CLDAR = field marigold (*Calendula arvensis*)

LAMAM = henbit (*Lamium amplexicaule* L.)

SONOL = annual sowthistle (*Sonchus oleraceus* L.)

Table 2. Control of MALNE and VLLEC by Penoxsulam plus Florasulam at 90 DAA in the Field.

		% Control			
Application Rate (gai/ha)		MALNE		VLLEC	
Penoxsulam	Florasulam	Obs	Ex	Obs	Ex
10	0	8	-	80	-
0	5	50	-	10	-
10	5	85	54	100	82

(continued)

		% Control				
Application Rate (gai/ha)		MALNE		VLLEC		
	Penoxsulam	Florasulam	Obs	Ex	Obs	Ex
5	13	0	-	-	80	-
10	0	6.3	-	-	10	-
15	13	6.3	-	-	99	82
MALNE = common mallow (<i>Malva neglecta</i> Wallr.)						
VLLEC = corn salad (<i>Valerianella echinata</i>)						

Table 3. Control of ERICA, SHRAR and SONSS by Penoxsulam plus Florasulam at 99-175 DAA in the Field.

		% Control						
Application Rate (gai/ha)		ERICA		SHRAR		SONSS		
	Penoxsulam	Florasulam	Obs	Ex	Obs	Ex	Obs	Ex
20	20	0	23	-	0	-	53	-
25	0	7.5	51	-	0	-	8	-
30	20	7.5	96	60	50	0	96	61
ERICA = horseweed (<i>Conyza canadensis</i> (L.) Cronq.)								
SHRAR = field madder (<i>Sherardia arvensis</i> L.)								
SONSS = sowthistle (<i>Sonchus</i> species)								

Claims

1. A synergistic herbicidal composition comprising a herbicidally effective amount of (a) penoxsulam and (b) florasulam.
2. The synergistic herbicidal composition of claim 1, wherein the weight ratio of penoxsulam to florasulam is from 17:1 to 1:1.
3. The synergistic herbicidal composition of claim 2, wherein the weight ratio is 7:1.
4. The synergistic herbicidal composition of claim 2, wherein the weight ratio is from 2.7:1 to 2.0:1.
5. The herbicidal composition of any of claims 1 to 4 further comprising an agriculturally acceptable adjuvant or carrier.
6. A method of controlling undesirable vegetation which comprises contacting the vegetation or the locus thereof with a herbicidally effective amount of (a) penoxsulam and (b) florasulam.
7. The method of claim 6, wherein the weight ratio of penoxsulam to florasulam is from 17:1 to 1:1.
8. The method of claim 7, wherein the weight ratio is 7:1.
9. The method of claim 7, wherein the weight ratio is from 2.7:1 to 2.0:1.
10. The method of claim 6, wherein the vegetation or the locus thereof is contacted with a herbicidally effective amount of the synergistic herbicidal composition of any of claims 1 to 4.
11. The method of claim 10, wherein the undesirable vegetation is controlled in rice, cereal and grain crops, turf, industrial vegetation management, sugar cane, range and pasture, or tree and vine orchards.

12. The method of claim 10 in which the synergistic herbicidal composition is applied preemergence to early postemergence.
- 5 13. The method of claim 10, wherein the undesirable vegetation is *Anthemis cotula* (ANTCO), *Calendula arvensis* (CLDAR), *Lamium amplexicaule* (LMAM), *Sonchus oleraceus* (SONOL), *Malva neglecta* (MALNE), or *Valerianella echinata* (VLLEC).

Patentansprüche

- 10 1. Eine synergistische herbizide Zusammensetzung umfassend eine herbizid wirksame Menge von (a) Penoxsulam und (b) Florasulam.
- 15 2. Die synergistische herbizide Zusammensetzung gemäß Anspruch 1, in welcher das Gewichtsverhältnis von Penoxsulam zu Florasulam von 17:1 bis 1:1 beträgt.
- 20 3. Die synergistische herbizide Zusammensetzung gemäß Anspruch 2, in welcher das Gewichtsverhältnis 7:1 beträgt.
- 25 4. Die synergistische herbizide Zusammensetzung gemäß Anspruch 2, in welcher das Gewichtsverhältnis von 2,7:1 bis 2,0:1 beträgt.
- 30 5. Die herbizide Zusammensetzung gemäß einem der Ansprüche 1 bis 4, weiterhin umfassend einen landwirtschaftlich akzeptablen Hilfs- oder Trägerstoff.
- 35 6. Ein Verfahren zur Bekämpfung unerwünschter Vegetation, welches das In-Kontakt-Bringen der Vegetation oder des Ortes derselben mit einer herbizid wirksamen Menge von (a) Penoxsulam und (b) Florasulam umfasst.
- 40 7. Das Verfahren gemäß Anspruch 6, in welchem das Gewichtsverhältnis von Penoxsulam zu Florasulam von 17:1 bis 1:1 beträgt.
- 45 8. Das Verfahren gemäß Anspruch 7, in welchem das Gewichtsverhältnis 7:1 beträgt.
- 50 9. Das Verfahren gemäß Anspruch 7, in welchem das Gewichtsverhältnis von 2,7:1 bis 2,0:1 beträgt.
- 55 10. Das Verfahren gemäß Anspruch 6, in welchem die Vegetation oder der Ort derselben mit einer herbizid wirksamen Menge der synergistischen herbiziden Zusammensetzung gemäß einem der Ansprüche 1 bis 4 in Kontakt gebracht wird.
- 60 11. Das Verfahren gemäß Anspruch 10, in welchem die unerwünschte Vegetation in Reis-, Getreide- und Kornnutzpflanzen, Rasen, im industriellen Vegetationsmanagement, in Zuckerrohr, Weide- und Wiesenland oder Baum- und Weinplantagen bekämpft wird.
- 65 12. Das Verfahren gemäß Anspruch 10, in welchem die synergistische herbizide Zusammensetzung im Vorlauf bis zum frühen Nachlauf angewandt wird.
- 70 13. Das Verfahren gemäß Anspruch 10, in welchem die unerwünschte Vegetation *Anthemis cotula* (ANTCO), *Calendula arvensis* (CLDAR), *Lamium amplexicaule* (LMAM), *Sonchus oleraceus* (SONOL), *Malva neglecta* (MALNE) oder *Valerianella echinata* (VLLEC) ist.

Revendications

1. Composition herbicide synergique comprenant, en quantité à effet herbicide,
- 55 a) du pénosulame
 b) et du florasulame.
2. Composition herbicide synergique conforme à la revendication 1, dans laquelle le rapport pondéral du pénosulame

au florasulame vaut de 17/1 à 1/1.

3. Composition herbicide synergique conforme à la revendication 2, dans laquelle ledit rapport pondéral vaut 7/1.

5 4. Composition herbicide synergique conforme à la revendication 2, dans laquelle ledit rapport pondéral vaut de 2,7/1
à 2,0/1.

10 5. Composition herbicide conforme à l'une des revendications 1 à 4, qui comprend en outre un adjuvant ou véhicule
admissible en agriculture.

15 6. Procédé de lutte contre des végétaux indésirables, comportant le fait de mettre ces végétaux, ou l'endroit où ils
poussent, en contact avec, en quantité à effet herbicide,

a) du pénoxsulame

15 b) et du florasulame.

7. Procédé conforme à la revendication 6, dans lequel le rapport pondéral du pénoxsulame au florasulame vaut de
17/1 à 1/1.

20 8. Procédé conforme à la revendication 6, dans lequel ledit rapport pondéral vaut 7/1.

9. Procédé conforme à la revendication 6, dans lequel ledit rapport pondéral vaut de 2,7/1 à 2,0/1.

25 10. Procédé conforme à la revendication 6, dans lequel on met les végétaux ou l'endroit où ils poussent en contact
avec une quantité à effet herbicide d'une composition herbicide synergique conforme à l'une des revendications 1 à 4.

30 11. Procédé conforme à la revendication 10, par lequel on lutte contre des végétaux indésirables dans des cultures de
riz, de céréales ou de grains, sur du gazon, dans la gestion industrielle de végétation, dans des cultures de canne
à sucre, dans des prairies et pâturages, ou dans des vergers et des vignes.

35 12. Procédé conforme à la revendication 10, dans lequel on applique la composition herbicide synergique depuis avant
la levée jusqu'peu après la levée.

13. Procédé conforme à la revendication 10, dans lequel les végétaux indésirables sont *Anthemis cotula* (ANTCO),
35 *Calendula arvensis* (CLDAR), *Lamium Amplexicaule* (LMAM), *Sonchus Oleraceus* (SONOL), *Malva neglecta* (MAL-
NE) ou *Valerianella echinata* (VLLEC).

40

45

50

55

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Non-patent literature cited in the description

- Herbicide Handbook of the Weed Science Society of America. 2007, 429 [0003]
- The Pesticide Manual. 2009 [0007] [0008]
- McCutcheon's Detergents and Emulsifiers Annual. MC Publishing Corp, 1998 [0022]
- Encyclopedia of Surfactants. Chemical Publishing Co, 1980, vol. I-III [0022]
- COLBY, S. R. Calculation of the synergistic and antagonistic response of herbicide combinations. *Weeds*, 1967, vol. 15, 20-22 [0034]

Szabadalmi igénypontok

1. Színergetikus herbicid készítmény, amely tartalmaz (a) penoxszulamot és (b) floraszulamot gyomirtásra hatásos mennyiségben.
2. Az 1. igénypont szerinti színergetikus herbicid készítmény, ahol a penoxszulammak floraszulamhoz mérő tömegaránya 17:1-től 1:1-ig terjed.
 3. A 2. igénypont szerinti színergetikus herbicid készítmény, ahol a tömegarány 7:1.
 4. A 2. igénypont szerinti színergetikus herbicid készítmény, ahol a tömegarány 2,7:1-től 2,0:1-ig terjed.
 5. Az 1-4. igénypontok bármelyike szerinti herbicid készítmény, amely tartalmaz továbbá egy mezőgazdaságilag elfogadható adjuvánt vagy hordozóanyagot.
 6. Eljárás nem kívánatos vegetáció irányára, amely tartalmazza a vegetációnak vagy annak helyének érintkezetét az (a) penoxszulam és (b) floraszulam gyomirtásra hatásos mennyiségével.
 7. A 6. igénypont szerinti eljárás, ahol a penoxszulammak floraszulamhoz mérő tömegaránya 17:1-től 1:1-ig terjed.
 8. A 7. igénypont szerinti eljárás, ahol a tömegarány 7:1.
 9. A 7. igénypont szerinti eljárás, ahol a tömegarány 2,7:1-től 2,0:1-ig terjed.
 10. A 6. igénypont szerinti eljárás, ahol a vegetációt vagy annak helyét érintkeztetjük az 1-4. igénypontok bármelyike szerinti színergetikus herbicid készítmény gyomirtásra hatásos mennyiségével.
 11. A 10. igénypont szerinti eljárás, ahol a nem kívánatos vegetációt a következőkben írjuk: rizs, gabona és szemes termények, gyep, ipari növényzet, cukornád, szabad terület és legelő, vagy fa és száló ültetvények.
 12. A 10. igénypont szerinti eljárás, amelyben a színergetikus herbicid készítményt preemergens ~ korai posztemergens időszakban alkalmazzuk.
 13. A 10. igénypont szerinti eljárás, ahol a nem kívánatos vegetáció *Anthemis cotula* (ANTCO), *Calendula arvensis* (CLOAR), *Lamium amplexicaule* (LMAM), *Sonchus oleraceus* (SONOL), *Malva neglecta* (MALNE) vagy *Valerianella echinata* (VLLEC).