

(19)



(11)

EP 3 795 667 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention of the grant of the patent:
02.11.2022 Bulletin 2022/44

(51) International Patent Classification (IPC):
C11D 3/39 ^(2006.01) **C11D 3/16** ^(2006.01)
C11D 7/26 ^(2006.01) **C11D 7/32** ^(2006.01)

(21) Application number: **19198748.6**

(52) Cooperative Patent Classification (CPC):
C11D 3/391; C11D 3/168; C11D 3/392;
C11D 7/267; C11D 7/3281

(22) Date of filing: **20.09.2019**

(54) NON-ENZYMATIC REMOVAL OF PROTEINACEOUS SOILS

NICHTENZYMATISCHE ENTFERNUNG VON EIWEISSHALTIGEN BÖDEN

ÉLIMINATION DE SOLS PROTÉINIQUES NON ENZYMATIQUES

(84) Designated Contracting States:
AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR

- **Schaefer, Sascha**
40822 Mettmann (DE)
- **Umbreit, Christian**
41466 Neuss (DE)
- **Hegetschweiler, Kaspar**
66386 St. Ingbert (DE)

(43) Date of publication of application:
24.03.2021 Bulletin 2021/12

(73) Proprietor: **Henkel AG & Co. KGaA**
40589 Düsseldorf (DE)

(56) References cited:
EP-A2- 0 458 397 WO-A1-98/11186
CA-A1- 2 882 331

(72) Inventors:
• **Kropf, Christian**
40724 Hilden (DE)

EP 3 795 667 B1

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Description

[0001] The present invention relates to detergents, especially hard surface cleaning compositions, and among those especially dishwashing compositions, comprising metal complexes with certain heterocyclic ligands. In addition, it relates to the use of such complexes to remove baked-on soils from surfaces, especially hard surfaces, for example the surfaces of cookware, tableware, and kitchen surfaces.


[0002] Cooked-, baked- and burnt-on soils are amongst the most severe types of soils to remove from surfaces. Traditionally, the removal of cooked-, baked- and burnt-on soils from cookware and tableware requires soaking the soiled object prior to a mechanical action. Manual dishwashing process requires a tremendous rubbing effort to remove cooked-, baked- and burnt-on soils and this can be detrimental to the safety and condition of the cookware/tableware. Generally, all cooked-, baked- and burnt-on soils are difficult to remove, however of particular relevance herein are the protein-based soils.

[0003] From the European patent application EP 0 458 397 A2 bleach activating Mn-complexes with 1,4,7-triazanonane ligands are known. The use of cleaning compositions for removing cooked-, baked- and burnt-on solids is also known in the art. For example, European patent application EP 0 286 075 A2 provides a method for treating hard surfaces soiled with cooked-on, baked-on or dried-on food residues, comprising applying to the soiled article a pre-spotting composition that comprises 1-40 % of surfactant, 1-10 % of builder, 0.2-2 % of mono-, di- and/or triethanolamine, water, and 3-50 % of organic solvents. International patent application WO 97/44427 A1 provides an aqueous hard surface cleaning composition for removing hardened dried or baked-on grease soil deposits; the composition comprises 0.01-0.85 % amine oxide non-ionic surfactant, up to 1.5 % chelating agent, 0.01-2.5 % caustic, 3-9 % of a glycol ether solvent system, up to 5 % water soluble organic amine and up to 2.5 % of an anti-redeposition agent. The international patent application WO 2005/059076 A1 discloses compositions for removing burnt-on soils; they comprise alkyl esters as surface modifying agents, soil swelling agents such as monoethanolamine, and an enzyme. Further detergent compositions for removing proteinaceous soils from surfaces are disclosed in WO 98/11186 A1 and CA 2 882 331 A1.

[0004] While those compositions do have some efficacy in the removal of grease-based soils, they are not effective in the removal of burnt-on protein-based soils. Agents of choice to break up proteins in stains are proteases; however, the thermal treatment during cooking-on or baking-on changes the structure of proteins, partially degrades proteins and forms different bonds among the protein parts and possibly other components of the soil. Many proteases show low performance on aged proteins and proteins which suffered such reactions.

[0005] It is therefore an object of the present invention to improve the efficacy on proteinaceous soils when removing cooked-, baked- and burnt-on soils from surfaces.


[0006] According to the present invention there is provided the use of a complex of formula I, or of a detergent comprising a complex of formula I,



in which the metal ions M^{n+} are selected from Al^{3+} , Ti^{4+} , Y^{3+} , Zr^{4+} , Ce^{3+} , Ce^{4+} , Sc^{3+} , Yb^{3+} , La^{3+} , Ta^{5+} and their mixtures,

n stands for a number in the range of 1 to 5,

L stands for a ligand according to formula II,



in which

E is selected from O, NR^1 , and mixtures of them,

R^1 is selected from H, optionally substituted C_{1-20} -alkyl or heteroalkyl groups, optionally substituted C_{2-20} -alkenyl, heteroalkenyl or alkinyl groups, optionally substituted aryl or heteroaryl groups, and optionally substituted C_{1-20} -alkylaryl or heteroaryl groups,

m stands for a number in the range of from 0 to 3,

A stands for an alkali metal cation and q stands for 1, an earth alkali metal cation and q stands for 2, or an


ammonium cation and q stands for 1,

X^v stands for an anion selected from the group of F^- , Cl^- , Br^- , I^- , OH^- , HSO_3^- , SO_3^{2-} , SO_4^{2-} , HSO_4^- , NO_2^- , NO_3^- , PO_4^{3-} , HPO_4^{2-} , $H_2PO_4^-$, BF_4^- , PF_6^- , ClO_4^- , organic anions such as acetate, citrate, formate, glutarate, lactate, malate, malonate, oxalate, pyruvate, tartrate, methanesulfonate, methyl sulfate, tosylate, succinate and their mixtures, with v being 1, 2 or 3, respectively, and p and r, irrespective of each other, are numbers in the range of from 0 to 7, selected such that the charge of the complex according to formula I is 0, preferably with the proviso that not each E is O, and/or the proviso that not each R^1 is H, to remove cooked-, baked- and burnt-on proteinaceous soils from surfaces, preferably hard surfaces, or textile surfaces.


[0007] Preferred are ligands of formula (II) in which at most one E is O. Also preferred are ligands of formula (II) in which each R^1 is not H.

[0008] As every chemist knows, not each number in the range given for n is feasible for each metal M, as metals of group 3 of the periodic table of elements tend to occur in oxidation state +3, metals of group 4, group 7, group 8, group 9, group 10, and the lanthanoid metals in oxidation states +2, +3, or +4, metals of group 5 in oxidation states +2, +3, +4, or +5, metals of group 6 in oxidation states +2, or +3, metals of group 11 in oxidation states +1, +2, or +3, metals of group 12 in oxidation states +1, or +2, and Al in oxidation state +3. The metal ions M^{n+} are Al^{3+} , Ti^{4+} , Y^{3+} , Zr^{4+} , Ce^{3+} , Ce^{4+} , Sc^{3+} , Yb^{3+} , La^{3+} , Ta^{5+} and their mixtures. Preferred heteroatoms in group R^1 are selected from O, N, S, Si, P, and their mixtures. Preferred substituents in group R^1 are selected from $-COOH$, $-COO^-$, $-OR^2$, and their mixtures, with $R^2 = H$, or C_{1-6} alkyl. Said alkyl, alkenyl, and alkynyl groups R^1 and R^2 , if comprising more than 2 atoms in case of alkyl and alkenyl groups and more than 3 atoms in case of alkynyl groups, may be linear or branched.


[0009] Preferred groups R^1 are selected from C_{1-18} alkyl groups,



, and their mixtures; especially preferred are



and




[0010] The compositions comprising the complex of formula (I) are suitable to be used in conventional washing and cleaning operations, such as manual or automatic laundry processes, and, preferably, the manual or automatic cleaning of hard surfaces, such as floors, tiles, kitchen surfaces, and, especially, surfaces that came into contact with protein containing food while being hot. Among the latter are for example cooking pots and pans, baking tins and moulds, stove plates and the interior surfaces of baking ovens. The detergents comprising the complex of formula (I) may be in the form of solids, liquids, gels or pastes and may exist as, for example, powders, tablets, bars, homogeneous liquids, microemulsions or emulsions. They preferably comprise said complex of formula I in amounts of from 0.001 % by weight


to 10 % by weight, more preferably from 0.01 % by weight to 3 % by weight; percentages given by weight here and in each following case are based on the total weight of the detergent, unless otherwise defined. The detergent compositions can be used as manual or automatic dishwashing detergent compositions or as a component thereof. If cooked-, baked- or burnt-on proteinaceous soil is to be removed from textiles, the detergent compositions can also be used as manual or automatic laundry detergent compositions or as a component thereof.

[0011] Ligands that are useful for this invention can be prepared by known reactions of the amine functionality of an amine (II) with at least one E being NR¹ and R¹ = H like i) reductive amination using a suitable starting material containing a carbonyl moiety and subsequent hydrogenation of the intermediate imine, or ii) nucleophilic substitution reactions using a suitable starting material containing a halogen moiety as a leaving group.

[0012] For example, the group



can be introduced as substituent by reductive amination with salicylaldehyde or by substitution with 2-(halogenomethyl)phenol or derivatives (preferred halogen is Cl, Br or I).



can be introduced as substituent by substitution with halogenoacetic acid or alkyl halogenoacetate (preferred halogen is Cl, Br or I) followed by saponification of the ester.

[0013] In case multiple functionalities are present in one of the starting materials or mixed substituents in the ligand are intended, it may be necessary to employ protection group strategies known from the state-of-the-art.

[0014] Complexes useful for this invention may be prepared by adding a solution of the respective metal salt to a solution of the ligand in the presence of a base and heating the reaction solution for a time sufficient for the formation of the complex. The anion in the metal salt can be any anion as defined for X^{v-}. A preferable solvent for the complex forming reaction is a solvent in which the ligand as well as the metal salt can be dissolved, e.g. water, methanol, ethanol, propanol, iso-propanol, acetonitrile, dioxane, dimethylacetamide, dimethylsulfoxide, other commonly used solvents or mixtures thereof. In case the metal salts are hygroscopic or sensitive versus formation of metal hydroxides, it is preferred to avoid water and use dry organic solvents of the above kind. Preferred bases to be employed in the complex forming reaction are ammonia, monoalkyl-, dialkyl-, trialkylamines or alkali metal hydroxides.

[0015] The amount of complex or of the detergent composition comprising it necessary to provide the required functionality will depend upon the nature of the complex or composition and can be determined by routine experimentation. Other conditions effective for removing cooked-, baked-and burnt-on proteinaceous soils, such as pH, temperature and treatment time, can also be determined by routine experimentation. Preferred herein, however is to employ aqueous washing or cleaning liquids that comprise a complex with a ligand of formula I in concentrations of from 0.001 mmol/l to 10 mmol/l, more preferred from 0.01 mmol/l to 5 mmol/l, and especially from 0.05 mmol/l to 1 mmol/l. A contact of the aqueous liquid with the proteinaceous soil for time periods in the range of from 2 minutes to 24 hours, preferably from 10 minutes to 12 hours, more preferably from 30 minutes to 4 hours are normally quite sufficient to remove the soils. Especially for manual processes the aqueous washing or cleaning liquid may be of ambient temperature or may be heated to higher temperatures still comfortable for manual contact; automatic processes may be operated at temperatures common for the device employed, such as for example 60 °C, 40 °C, 30 °C, or 20 °C for laundry machines and for example 70 °C, 60 °C, 55 °C, 50°C, 45 °C or 40°C for dishwashing machines.

[0016] Another subject of the invention is a method of removing cooked-, baked- and burnt-on proteinaceous soils from surfaces, preferably hard surfaces, or textile surfaces, by contacting the soiled surface with an aqueous liquid comprising said complex. The features given above are characteristics of preferred variants of such a process. Also, the features of the detergents given below are characteristics of preferred variants of such a process

[0017] The detergent compositions comprising said complex are particularly effective in removing proteinaceous soils from metallic or glass surfaces. They are also particularly effective in removing proteinaceous soils which are or comprise a product of a thermal reaction of a protein with a carbohydrate, the so-called Maillard reaction. The expression "a protein" or "a carbohydrate" encompasses a single protein and a single carbohydrate as well as mixtures of proteins

and mixtures of carbohydrates.

[0018] The detergents usable according to the invention may comprise further components that do not unfavourably interact with said complex, preferably selected from the group consisting of surfactants, in particular non-ionic surfactants and/or anionic surfactants, builders, enzymes, sequestering agents, electrolytes, corrosion inhibitors, suds suppressors, dyes, and fragrances. More preferably the detergents at least comprise a surfactant in addition to said complex. They may also comprise peroxygen-based bleaching agents, although those are not necessarily required for the catalytic action of the complex. Without wishing to be bound by theory, it is believed that in the absence of peroxygen-based bleaching agents oxygen out of the surrounding air will be sufficiently effective in sustaining the catalytic action of the complex.

[0019] The detergents described herein preferably comprise at least one non-ionic surfactant. All non-ionic surfactants known to a person skilled in the art may be used. Suitable non-ionic surfactants are, for example, alkyl glycosides of the general formula $RO(G)_x$, in which R corresponds to a primary straight-chain or methyl-branched, in particular methyl-branched at the 2-position, aliphatic functional group having 8 to 22, and preferably 12 to 18 carbon atoms, and G is the symbol that denotes a glucose unit having 5 or 6 carbon atoms, and preferably glucose. The degree of oligomerization x, which indicates the distribution of monoglycosides and oligoglycosides, is an arbitrary number between 1 and 10; x is preferably 1.2 to 1.4.

[0020] Non-ionic surfactants of the amine oxide type, for example N-cocoalkyl-N,N-dimethylamine oxide and N-tal-lowalkyl-N,N-dihydroxyethylamine oxide, and of the fatty acid alkanolamide type may also be suitable. The quantity of these non-ionic surfactants is preferably no more than that of the ethoxylated fatty alcohols, in particular no more than half thereof.


[0021] Further suitable surfactants are polyhydroxy fatty acid amides.

[0022] Preferably, however, low-sudsing non-ionic surfactants are used, in particular alkoxyated, and especially ethoxylated, low-sudsing non-ionic surfactants. It is particularly preferred when the automatic dishwasher detergents comprise non-ionic surfactants from the group of alkoxyated alcohols.

[0023] One class of non-ionic surfactants that can be used, which can be used either as the sole non-ionic surfactant or in combination with other non-ionic surfactants, is thus alkoxyated, preferably ethoxylated or ethoxylated and propoxylated fatty acid alkyl esters, preferably comprising 1 to 4 carbon atoms in the alkyl chain.

[0024] Surfactants that should preferably be used come from the groups of the ethoxylated primary alcohols and mixtures of these surfactants with structurally more complicated surfactants, such as polyoxypropylene/polyoxyethylene/polyoxypropylene (PO/EO/PO) surfactants. Such (PO/EO/PO) non-ionic surfactants are characterized by good suds control.


[0025] Non-ionic surfactants comprising alternating ethylene oxide and alkylene oxide units may be preferred. Among these, in turn, surfactants comprising EO-AO-EO-AO blocks are preferred, wherein in each case one to ten EO or AO groups are bound to one another before a block from the respective other group follows. Here, non-ionic surfactants of the general formula



are preferred, in which R^1 denotes a straight-chain or branched, saturated, monounsaturated or polyunsaturated C_{6-24} alkyl functional group or alkenyl residue, each group R^2 and R^3 , independently of one another, is selected from $-CH_3$, $-CH_2CH_3$, $-CH_2CH_2-CH_3$, $CH(CH_3)_2$, and the subscripts w, x, y and z, independently of one another, denote integers from 1 to 6.

[0026] Thus, in particular, non-ionic surfactants are preferred that comprise a C_{9-15} alkyl functional group having 1 to 4 ethylene oxide units, followed by 1 to 4 propylene oxide units, followed by 1 to 4 ethylene oxide units, followed by 1 to 4 propylene oxide units.

[0027] Preferred non-ionic surfactants are therefore those of the general formula



in which R^1 denotes a straight-chain or branched, saturated, monounsaturated or polyunsaturated C_{6-24} alkyl functional group or alkenyl residue; R^2 denotes H or a linear or branched hydrocarbon functional group having 2 to 26 carbon atoms; A, A', A'' and A''', independently of one another, denote a functional group from the group consisting of $-CH_2CH_2$, $-CH_2CH_2-CH_2$, $-CH_2-CH(CH_3)$, $-CH_2-CH_2-CH_2-CH_2$, $-CH_2-CH(CH_3)-CH_2-$, $-CH_2-CH(CH_2-CH_3)$; and w, x, y and z denote values between 0.5 and 120, wherein x, y and/or z may also be 0.

[0028] Preferred are in particular end-capped poly(oxyalkylated) non-ionic surfactants that, according to formula

$R^1O[CH_2CH_2O]_xCH_2CH(OH)R^2$, in addition to a functional group R^1 , which denotes linear or branched, saturated or unsaturated, aliphatic or aromatic hydrocarbon functional groups having 2 to 30 carbon atoms, and preferably having 4 to 22 carbon atoms, also comprise a linear or branched, saturated or unsaturated aliphatic or aromatic hydrocarbon functional group R^2 having 1 to 30 carbon atoms, wherein x denotes values between 1 and 90, preferably values between 30 and 80, and in particular values between 30 and 60.

[0029] Particularly preferred are surfactants of formula $R^1O[CH_2CH(CH_3)O]_x[CH_2CH_2O]_yCH_2CH(OH)R^2$, in which R^1 denotes a linear or branched, aliphatic hydrocarbon functional group having 4 to 18 carbon atoms or mixtures thereof, R^2 denotes a linear or branched hydrocarbon functional group having 2 to 26 carbon atoms or mixtures thereof, x denotes values between 0.5 and 1.5, and y denotes a value of at least 15.

[0030] The group of these non-ionic surfactants includes, for example, the C_{2-26} fatty alcohol-(PO)₁-(EO)₁₅₋₄₀-2-hydroxyalkyl ethers, and in particular also the C_{8-10} fatty alcohol-(PO)₁-(EO)₂₂-2-hydroxydecyl ethers. Particularly preferred are furthermore those end-capped poly(oxyalkylated) non-ionic surfactants of formula $R^1O[CH_2CH_2O]_x[CH_2CH(R^3)O]_yCH_2CH(OH)R^2$, in which R^1 and R^2 , independently of one another, denote a linear or branched, saturated, monounsaturated or polyunsaturated hydrocarbon functional group having 2 to 26 carbon atoms, R^3 , independently of one another, is selected from $-CH_3$, $-CH_2CH_3$, $-CH_2CH_2-CH_3$, $-CH(CH_3)_2$, preferably however is $-CH_3$, and x and y , independently of one another, denote values between 1 and 32, wherein non-ionic surfactants where $R^3 = -CH_3$ and values for x are from 15 to 32 and for y from 0.5 to 1.5 are especially particularly preferred.

[0031] Further non-ionic surfactants that may preferably be used are the end-capped poly(oxyalkylated) non-ionic surfactants of formula $R^1O[CH_2CH(R^3)O]_x[CH_2]_kCH(OH)[CH_2]_jOR^2$, in which R^1 and R^2 denote linear or branched, saturated or unsaturated, aliphatic or aromatic hydrocarbon functional groups having 1 to 30 carbon atoms, R^3 denotes H or a methyl, ethyl, n-propyl, iso-propyl, n-butyl, 2-butyl or 2-methyl-2-butyl functional group, x denotes values between 1 and 30, and k and j denote values between 1 and 12, and preferably between 1 and 5. When the value $x \geq 2$, each R^3 in the above formula $R^1O[CH_2CH(R^3)O]_x[CH_2]_kCH(OH)[CH_2]_jOR^2$ may be different. R^1 and R^2 are preferably linear or branched, saturated or unsaturated, aliphatic or aromatic hydrocarbon functional groups having 6 to 22 carbon atoms, wherein functional groups having 8 to 18 carbon atoms are particularly preferred. H, CH_3 or $-CH_2CH_3$ are particularly preferred for the functional group R^3 . Particularly preferred values for x are in the range of 1 to 20, and in particular of 6 to 15.

[0032] As described above, each R^3 in the above formula may be different when $x \geq 2$. In this way, the alkylene oxide unit in the square brackets may be varied. For example, when x is 3, then the functional group R^3 may be selected so as to form ethylene oxide- ($R^3 = H$) or propylene oxide- ($R^3 = CH_3$) units, which may be joined to one another in any arbitrary order, for example (EO)(PO)(EO), (EO)(EO)(PO), (EO)(EO)(EO), (PO)(EO)(PO), (PO)(PO)(EO) and (PO)(PO)(PO). The value 3 for x has been selected by way of example here and may certainly also be larger, wherein the variation range increases with increasing x values and, for example, includes a large number of (EO) groups, combined with a low number of (PO) groups, or vice versa.

[0033] Particularly preferred end-capped poly(oxyalkylated) alcohols of the above formula have values of $k = 1$ and $j = 1$, whereby the above formula is simplified to $R^1O[CH_2CH(R^3)O]_x[CH_2CH(OH)CH_2OR^2$. In the last formula, R^1 , R^2 and R^3 are as defined above, and x denotes numbers from 1 to 30, preferably from 1 to 20, and in particular 6 to 18. Particularly preferred are surfactants in which the functional groups R^1 and R^2 comprise 9 to 14 carbon atoms, R^3 denotes H, and x takes on values from 6 to 15.

[0034] Non-ionic surfactants that have proven to be particularly effective are those of the general formula $R^1-CH(OH)CH_2O-(AO)_w-R^2$ in which R^1 denotes a straight-chain or branched, saturated, monounsaturated or polyunsaturated C_{6-24} alkyl functional group or alkenyl residue; R^2 denotes a linear or branched hydrocarbon functional group having 2 to 26 carbon atoms; A denotes a functional group from the group consisting of CH_2CH_2 , $CH_2CH_2CH_2$, $CH_2CH(CH_3)$, and preferably CH_2CH_2 , and w denotes values between 1 and 120, preferably 10 to 80, and in particular 20 to 40. The group of these non-ionic surfactants includes, for example, the C_{4-22} fatty alcohol-(EO)₁₀₋₈₀-2-hydroxyalkyl ethers, and in particular also the C_{8-12} fatty alcohol-(EO)₂₂-2-hydroxydecyl ethers and the C_{4-22} fatty alcohol-(EO)₄₀₋₈₀-2-hydroxyalkyl ethers.

[0035] In various embodiments of the invention, it is also possible to use the corresponding not end-capped hydroxy mixed ethers instead of the above-defined end-capped hydroxy mixed ethers. These can satisfy the above formulas, wherein, however, R^2 is hydrogen, and R^1 , R^3 , A, A', A'', A''', w, x, y and z are as defined above.

[0036] Different embodiments of the detergents comprise the surfactant in an amount of at least 2 wt.%, and preferably of at least 5 wt.%. The absolute amounts used per application may be in the range of 0.5 to 10 g/job, and preferably in the range of 1 to 5 g/job, for example.


[0037] All anionic surface-active substances are suitable in the detergents. These are characterized by a water-soluble-rendering anionic group, such as a carboxylate, sulfate, sulfonate or phosphate group and a lipophilic alkyl group having approximately 8 to 30 carbon atoms. In addition, glycol or polyglycol ether groups, ester, ether and amide groups and hydroxyl groups can be present in the molecule. Suitable anionic surfactants are preferably present in the form of the sodium, potassium and ammonium salts, and monoalkanol, dialkanol and trialkanol ammonium salts having 2 to 4 carbon atoms in the alkanol group.

[0038] Preferred anionic surfactants are alkyl sulfates, alkyl polyglycol ether sulfates and ether carboxylic acids having 10 to 18 carbon atoms in the alkyl group, and up to 12 glycol ether groups in the molecule.

[0039] In various embodiments, the detergents thus comprise at least one surfactant of formula $R^4-O-(AO)_n-SO_3^-X^+$.

[0040] In this formula, R^4 denotes a linear or branched, substituted or unsubstituted alkyl, aryl or alkyl-aryl functional group, preferably a linear, unsubstituted alkyl functional group, and particularly preferably a fatty alcohol functional group. Preferred functional groups R^4 are selected from decyl, undecyl, dodecyl, tridecyl, tetradecyl, pentadecyl, hexadecyl, heptadecyl, octadecyl, nonadecyl and eicosyl functional groups and the mixtures thereof, wherein representatives having an even number of carbon atoms are preferred. Particularly preferred functional groups R^4 are derived from C_{12} to C_{18} fatty alcohols, for example from coconut fatty alcohol, tallow fatty alcohol, lauryl, myristyl, cetyl or stearyl alcohol, or from C_{10} to C_{20} oxo alcohols. AO denotes an ethylene oxide (EO) or propylene oxide (PO) grouping, preferably an ethylene oxide grouping. The subscript n denotes an integer from 1 to 50, preferably from 1 to 20, and in particular from 2 to 10. It is especially particularly preferred if n denotes the numbers 2, 3, 4, 5, 6, 7, or 8. X denotes a monovalent cation or the nth part of an n-valent cation, alkali metal ions being preferred, and among these Na^+ or K^+ , Na^+ being extremely preferred. Further cations X^+ may be selected from NH_4^+ , $1/2 Zn^{2+}$, $1/2 Ca^{2+}$, $1/2 Mn^{2+}$ and mixtures thereof.


[0041] Particularly preferred anionic surfactants are selected from ether sulfates of formula A-1



where $k = 11$ to 19 , $n = 2, 3, 4, 5, 6, 7$ or 8 . Especially particularly preferred representatives are $Na-C_{12-14}$ fatty alcohol ether sulfates comprising 2 EO ($k = 11$ to 13 , $n = 2$ in formula A-1).

[0042] In addition or as an alternative, the detergents can furthermore comprise at least one surfactant of formula $R^5-A-SO_3^-Y^+$. In this formula, R^5 denotes a linear or branched, substituted or unsubstituted alkyl, aryl or alkyl-aryl functional group, and the grouping -A- denotes -O- or a chemical bond. In other words, the above-described formula can be used to describe sulfate- ($A = O$) or sulfonate- ($A =$ chemical bond) surfactants. Depending on the selection of the grouping A, certain functional groups R^5 are preferred. In the sulfate surfactants ($A = O$), R^5 is preferably a linear, unsubstituted alkyl functional group, and particularly preferably a fatty alcohol functional group. Preferred functional groups R^5 are selected from decyl, undecyl, dodecyl, tridecyl, tetradecyl, pentadecyl, hexadecyl, heptadecyl, octadecyl, nonadecyl and eicosyl functional groups and the mixtures thereof, wherein representatives having an even number of carbon atoms are preferred. Particularly preferred functional groups R^5 are derived from C_{12} to C_{18} fatty alcohols, for example from coconut fatty alcohol, tallow fatty alcohol, lauryl, myristyl, cetyl or stearyl alcohol, or from C_{10} to C_{20} oxo alcohols. Y denotes a monovalent cation or the nth part of an n-valent cation, alkali metal ions being preferred, and among these Na^+ or K^+ , Na^+ being extremely preferred. Further cations Y^+ can be selected from NH_4^+ , $1/2 Zn^{2+}$, $1/2 Mg^{2+}$, $1/2 Ca^{2+}$, $1/2 Mn^{2+}$, and the mixtures thereof.

[0043] Such particularly preferred surfactants are selected from fatty alcohol sulfates of formula



where $k = 11$ to 19 . Especially particularly preferred representatives are $Na-C_{12-14}$ fatty alcohol sulfates ($k = 11$ to 13).

[0044] In the sulfonate surfactants ($A =$ chemical bond), R^5 is preferably a linear or branched unsubstituted alkyl-aryl functional group. Again, X denotes a monovalent cation or the nth part of an n-valent cation, alkali metal ions being preferred, and among these Na^+ or K^+ , Na^+ being extremely preferred. Further cations X^+ can be selected from NH_4^+ , $1/2 Zn^{2+}$, $1/2 Mg^{2+}$, $1/2 Ca^{2+}$, $1/2 Mn^{2+}$, and the mixtures thereof. Such surfactants may be selected from linear or branched alkylbenzene sulfonates.

[0045] Cationic and/or amphoteric surfactants, such as betaines or quaternary ammonium compounds, can also be used instead of or in conjunction with the described surfactants. However, it is preferred that no cationic and/or amphoteric surfactants are used.

[0046] Builders that may be present in the detergent are in particular silicates, aluminum silicates (in particular zeolites), carbonates, organic dicarboxylic and polycarboxylic acids, and amino carboxylic acids and the salts thereof, and, where ecological bias against their use is absent, also phosphates. Naturally, mixtures of these substances may also be used.

[0047] For example, it is possible to use crystalline phyllosilicates of the general formula $NaMSi_xO_{2x+1} \cdot y H_2O$, in which M is sodium or hydrogen, x is a number from 1.9 to 22, and preferably from 1.9 to 4, wherein particularly preferred values for x are 2, 3 or 4, and y is a number from 0 to 33, and preferably from 0 to 20. The crystalline phyllosilicates of formula $NaMSi_xO_{2x+1} \cdot y H_2O$ are sold under the trade name Na-SKS by Clariant GmbH (Germany), for example. Examples of these silicates are Na-SKS-1 ($Na_2Si_{22}O_{45} \cdot x H_2O$, kenyaite), Na-SKS-2 ($Na_2Si_{14}O_{29} \cdot x H_2O$, magadiite),

Na-SKS-3 ($\text{Na}_2\text{Si}_8\text{O}_{17} \cdot x \text{H}_2\text{O}$) or Na-SKS-4 ($\text{Na}_2\text{Si}_4\text{O}_9 \cdot x \text{H}_2\text{O}$, makatite). For the purposes of the present invention, crystalline phyllosilicates of formula $\text{NaMSi}_x\text{O}_{2x+1} \cdot y \text{H}_2\text{O}$ in which x denotes 2 are particularly suitable. In particular, both β - and δ -sodium silicates $\text{Na}_2\text{Si}_2\text{O}_5 \cdot y \text{H}_2\text{O}$ are preferred, and furthermore especially Na-SKS-5 (α - $\text{Na}_2\text{Si}_2\text{O}_5$), Na-SKS-7 (β - $\text{Na}_2\text{Si}_2\text{O}_5$, natrosilite), Na-SKS-9 ($\text{NaHSi}_2\text{O}_5 \cdot \text{H}_2\text{O}$), Na-SKS-10 ($\text{NaHSi}_2\text{O}_5 \cdot 3 \text{H}_2\text{O}$, kanemite), Na-SKS-11 (t - $\text{Na}_2\text{Si}_2\text{O}_5$) and Na-SKS-13 (NaHSi_2O_5), in particular however Na-SKS-6 (δ - $\text{Na}_2\text{Si}_2\text{O}_5$).

[0048] It is also possible to use amorphous sodium silicates having a $\text{Na}_2\text{O}:\text{SiO}_2$ module of 1:2 to 1:3.3, preferably of 1:2 to 1:2.8, and in particular of 1:2 to 1:2.6, which preferably exhibit retarded dissolution and secondary washing properties. The retarded dissolution compared to conventional amorphous sodium silicates can have been caused in a variety of ways, for example by way of surface treatment, compounding, compacting/compression or over-drying. Within the scope of the present invention, the term "amorphous" shall be understood to mean that the silicates do not supply any sharp X-ray reflexes in X-ray diffraction experiments, such as those that are typical of crystalline substances, but at best evoke one or more maxima of the scattered X-rays, which have a width of several degree units of the diffraction angle. Within the scope of the present invention, it is preferred that this silicate or these silicates, preferably alkali silicates, and particularly preferably crystalline or amorphous alkali disilicates, is or are present in the detergents in amounts of 1 to 40 wt.%, and preferably of 2 to 35 wt.%.

[0049] It is also possible, of course, to use the generally known phosphates as builder substances, provided that such use should not be avoided for ecological reasons. Alkali metal phosphates is the term that covers all the alkali metal (in particular sodium and potassium) salts of the different phosphoric acids, in which a distinction can be made between metaphosphoric acids $(\text{HPO}_3)_n$ and orthophosphoric acid H_3PO_4 , in addition to higher molecular weight representatives. The phosphates combine several advantages: They act as alkali carriers, prevent limescale deposits on machine parts or lime scaling on woven fabrics, and additionally contribute to the cleaning performance. Technically particularly important phosphates are pentasodium triphosphate, $\text{Na}_5\text{P}_3\text{O}_{10}$ (sodium tripolyphosphate), and the corresponding potassium salt pentapotassium triphosphate, $\text{K}_5\text{P}_3\text{O}_{10}$ (potassium tripolyphosphate) and corresponding mixed salts (sodium potassium tripolyphosphates). Preferably, however, the detergents are phosphate-free. If phosphate is used as substances with cleaning action in the detergent, preferred ones comprise phosphate in amounts from 5 to 80 wt.%, preferably from 10 to 60 wt.%, and in particular from 18 to 45 wt.%.

[0050] The detergents can in particular also comprise phosphonates as a further builder. The phosphonate compound used is preferably a hydroxyalkane phosphonate and/or aminoalkane phosphonate. Among the hydroxyalkane phosphonates, 1-hydroxyethane-1,1-diphosphonate (HEDP) is of particular importance. Possible preferred aminoalkane phosphonates include ethylenediaminetetramethylene phosphonate (EDTMP), diethylenetriaminepentamethylene phosphonate (DTPMP) and the higher homologs thereof. Phosphonates are preferably present in the detergents in amounts of 0.1 to 10 wt.%, and in particular in amounts of 0.5 to 8 wt.%.

[0051] Other builders are also alkali carriers. For example, alkali metal hydroxides, alkali metal carbonates, alkali metal hydrogen carbonates, alkali metal sesquicarbonates, the described alkali silicates, alkali metasilicates and mixtures of the above-mentioned substances are considered alkali carriers, wherein within the meaning of the present invention preferably the alkali carbonates, in particular sodium carbonate, sodium hydrogen carbonate or sodium sesquicarbonate, can be used. A builder system containing a mixture of tripolyphosphate and sodium carbonate is particularly preferred. A builder system containing a mixture of tripolyphosphate and sodium carbonate and sodium silicate is likewise particularly preferred. Given the low chemical compatibility with the remaining ingredients of detergents compared to other builder substances, the optional alkali metal hydroxides are preferably used only in low amounts, preferably in amounts of less than 10 wt.%, especially less than 6 wt.%, particularly preferably less than 4 wt.%, and in particular less than 2 wt.%. Detergents that, based on the total weight thereof, comprise less than 0.5 wt.%, and in particular no alkali metal hydroxides, are particularly preferred.


[0052] The use of carbonate(s) and/or hydrogen carbonate(s), preferably alkali carbonate(s), particularly preferably sodium carbonate, in amounts from 2 to 50 wt.%, preferably from 5 to 40 wt.%, and in particular from 7.5 to 30 wt.%, is particularly preferred. Detergents that contain less than 20 wt.%, especially less than 17 wt.%, preferably less than 13 wt.%, and in particular less than 9 wt.% carbonate(s) and/or hydrogen carbonate(s), preferably alkali carbonate(s), particularly preferably sodium carbonate, are particularly preferred.

[0053] Usable organic builder substances are, for example, the polycarboxylic acids that can be used in the form of the free acid and/or of the sodium salts thereof, wherein polycarboxylic acids shall be understood to mean those carboxylic acids that carry more than one acid function. These include, for example, citric acid, adipic acid, succinic acid, glutaric acid, malic acid, tartaric acid, maleic acid, fumaric acid, saccharic acids, nitrilotriacetic acid (NTA), provided that such use is not objectionable for ecological reasons, and mixtures thereof. In addition to the builder effect, the free acids typically also have the property of being an acidifying component and are thus also used to set a lower and milder pH value of the detergents. In particular, citric acid, succinic acid, glutaric acid, adipic acid, gluconic acid and arbitrary mixtures of these shall be mentioned here.

[0054] The use of citric acid and/or citrates in detergents has proven to be particularly advantageous for the cleaning and rinsing performance. Amino carboxylic acids and/or the salts thereof are another significant class of phosphate-free




builders. Particularly preferred representatives of this class are methyl glycine diacetic acid (MGDA) or the salts thereof, and glutamine diacetic acid (GLDA) or the salts thereof or ethylenediamine diacetic acid (EDDS) or the salts thereof. The content of these amino carboxylic acids or of the salts thereof can amount to between 0.1 and 30 wt.%, preferably between 1 and 25 wt.%, and in particular between 5 and 20 wt.%, for example. Amino carboxylic acids and the salts thereof can be used together with the above-mentioned builders, in particular also with the phosphate-free builders.

[0055] The detergents according to the invention can furthermore comprise a sulfo polymer. The percent by weight of the sulfo polymer in the detergent is preferably 0.1 to 20 wt.%, in particular 0.5 to 18 wt.%, particularly preferably 1.0 to 15 wt.%, in particular 4 to 14 wt.%, and especially 6 to 12 wt.%. The sulfo polymer is typically used in the form of an aqueous solution, wherein the aqueous solutions typically comprise 20 to 70 wt.%, in particular 30 to 50 wt.%, and preferably approximately 35 to 40 wt.% sulfo polymers. Preferably, the sulfo polymer used is a copolymeric polysulfonate, and preferably a hydrophobically modified copolymeric polysulfonate. The copolymers can comprise two, three, four or more different monomer units. Preferred copolymeric polysulfonates comprise at least one monomer from the group of the unsaturated carboxylic acids, in addition to sulfonic group-containing monomer(s). Particularly preferably, unsaturated carboxylic acids of formula $R^1(R^2)C=C(R^3)COOH$ are used as unsaturated carboxylic acid(s), in which R^1 to R^3 , independently of one another, denote -H, -CH₃, a straight-chain or branched saturated alkyl functional group having 2 to 12 carbon atoms, a straight-chain or branched, monounsaturated or polyunsaturated alkenyl residue having 2 to 12 carbon atoms, alkyl functional groups or alkenyl residues substituted with -NH₂, -OH or -COOH as defined above, or -COOH or -COOR⁴, wherein R⁴ is a saturated or unsaturated, straight-chain or branched hydrocarbon functional group having 1 to 12 carbon atoms. Particularly preferred unsaturated carboxylic acids are acrylic acid, methacrylic acid, ethacrylic acid, α -chloroacrylic acid, α -cyanoacrylic acid, crotonic acid, α -phenylacrylic acid, maleic acid, maleic anhydride, fumaric acid, itaconic acid, citraconic acid, methylene malonic acid, sorbic acid, cinnamic acid or the mixtures thereof. It is also possible, of course, to use the unsaturated dicarboxylic acids. Among the sulfonic acid group-containing monomers, those of formula



are preferred, in which R^5 to R^7 , independently of one another, denote -H, -CH₃, a straight-chain or branched saturated alkyl functional group having 2 to 12 carbon atoms, a straight-chain or branched, monounsaturated or polyunsaturated alkenyl residue having 2 to 12 carbon atoms, alkyl functional groups or alkenyl residues substituted with -NH₂, OH or -COOH, or -COOH or COOR⁴, wherein R⁴ is a saturated or unsaturated, straight-chain or branched hydrocarbon functional group having 1 to 12 carbon atoms, and X denotes an optionally present spacer group which is selected from -(CH₂)_n- where n = 0 to 4, -COO-(CH₂)_k- where k = 1 to 6, -C(O)-NH-C(CH₃)₂-, -C(O)-NH-C(CH₃)₂-CH₂- and -C(O)-NH-CH(CH₃)-CH₂-.

[0056] Preferred among these monomers are those of formulas



in which R^6 and R^7 , independently of one another, are selected from -H, -CH₃, -CH₂CH₃, -CH₂CH₂CH₃ and -CH(CH₃)₂, and X denotes an optionally present spacer group, which is selected from -(CH₂)_n- where n = 0 to 4, -COO-(CH₂)_k- where k = 1 to 6, -C(O)-NH-C(CH₃)₂-, -C(O)-NH-C(CH₃)₂-CH₂- and -C(O)-NH-CH(CH₃)-CH₂-. Particularly preferred sulfonic acid group-containing monomers are 1-acrylamido-1-propanesulfonic acid, 2-acrylamido-2-propanesulfonic acid, 2-acrylamido-2-methyl-1-propanesulfonic acid, 2-methacrylamido-2-methyl-1-propanesulfonic acid, 3-methacrylamido-2-hydroxypropanesulfonic acid, allylsulfonic acid, methallylsulfonic acid, allyloxybenzenesulfonic acid, methallyloxybenzenesulfonic acid, 2-hydroxy-3-(2-propenyloxy)propanesulfonic acid, 2-methyl-2-propenesulfonic acid, styrenesulfonic acid, vinylsulfonic acid, 3-sulfopropylacrylate, 3-sulfopropylmethacrylate, sulfomethacrylamide, sulfomethylmethacrylamide, and mixtures of the described acids or the water-soluble salts thereof. The sulfonic acid groups can be present entirely or partially in neutralized form in the polymer, which is to say that, in some or all sulfonic acid groups, the acid hydrogen atom of the sulfonic acid group can be replaced with metal ions, preferably alkali metal ions, and in particular with sodium ions. The use of partially or fully neutralized sulfonic acid group-containing copolymers is preferred according to the invention. The monomer distribution of the preferably used copolymers is preferably 5 to 95 wt.% for copolymers that comprise only carboxylic acid group-containing monomers and sulfonic acid group-containing monomers, and particularly preferably the content of the sulfonic acid group-containing monomer is 50 to 90 wt.% and the content of the carboxylic acid group-containing monomer is 10 to 50 wt.%, the monomers preferably being selected from those described above. The molar mass of the preferably used sulfo copolymers can be varied so as to adapt the properties of the polymers to

the desired intended purpose. Preferred detergents are characterized in that the copolymers have molar masses of 2000 to 200,000 g/mol⁻¹, preferably of 4000 to 25,000 g/mol⁻¹, and in particular of 5000 to 15000 g/mol⁻¹.

[0057] The detergents can comprise further polymers. The group of suitable polymers includes in particular polymers with cleaning action, for example rinse polymers and/or polymers acting as softeners. Polymers that are preferably used come from the group of alkylacrylamide/acrylic acid copolymers, alkylacrylamide/methacrylic acid copolymers, alkylacrylamide/methylmethacrylic acid copolymers, alkylacrylamide/acrylic acid/alkylaminoalkyl(meth)acrylic acid copolymers, alkylacrylamide/methacrylic acid/alkylaminoalkyl(meth)acrylic acid copolymers, alkylacrylamide/methylmethacrylic acid/alkylaminoalkyl(meth)acrylic acid copolymers, alkylacrylamide/alkyl methacrylate/alkylaminoethyl methacrylate/alkyl methacrylate copolymers, and copolymers of unsaturated carboxylic acids, cationically derivatized unsaturated carboxylic acids, and optionally further ionic or non-ionogenic monomers. Further polymers that may be used come from the group of acrylamidoalkyltrialkylammonium chloride/acrylic acid copolymers and the alkali and ammonium salts thereof, acrylamidoalkyltrialkylammonium chloride/methacrylic acid copolymers and the alkali and ammonium salts thereof, and methacroylethylbetaine/methacrylate copolymers. Cationic polymers that may be used come from the groups of quaternized cellulose derivatives, polysiloxanes comprising quaternary groups, cationic guar derivatives, polymeric dimethyldiallylammonium salts and the copolymers thereof with acrylic acid and methacrylic acid and the esters and amides of acrylic acid and methacrylic acid, copolymers of vinylpyrrolidone with quaternized derivatives of dialkylaminoacrylate and -methacrylate, vinylpyrrolidone/methoimidazolium chloride copolymers, quaternized polyvinyl alcohols, or the polymers described by the INCI names Polyquaternium 2, Polyquaternium 17, Polyquaternium 18, and Polyquaternium 27.

[0058] The detergents preferably contain at least one enzyme preparation or enzyme composition, which contains one or more enzymes. Suitable enzymes include, but are not limited to, proteases, amylases, lipases, hemicellulases, cellulases, perhydrolases or oxidoreductases, and preferably the mixtures thereof. These enzymes are, in principle, of natural origin; proceeding from the natural molecules, improved variants are available for use in detergents and are used in correspondingly preferred fashion. The detergents preferably comprise enzymes in total amounts of 1 x 10⁻⁶ to 5 wt.%, based on active protein. The protein concentration can be determined using known methods, such as the BCA method or the biuret method.

[0059] Proteases are some of the technically most significant enzymes. They cause protein-containing soiling on the goods to be cleaned to decompose. Among these, in turn, proteases of the subtilisin type (subtilases, subtiloepitidases, EC 3.4.21.62) are particularly important, which due to the catalytically active amino acids are serine proteases. They act as non-specific endopeptidases and hydrolyze arbitrary acid amide bonds that lie in the interior of peptides or proteins. The pH optimum of these is usually in the distinctly alkaline range. Subtilases are formed naturally from microorganisms. Among these, in particular the subtilisins formed and secreted by the *Bacillus* species shall be mentioned as the most significant group within the subtilases. Examples of the proteases of the subtilisin type preferably used in detergents are the subtilisins BPN¹ and Carlsberg, the protease PB92, the subtilisins 147 and 309, the protease from *Bacillus lentus*, and in particular from *Bacillus lentus* DSM 5483, subtilisin DY, and the thermitase enzymes, which can be assigned to the subtilases, but not to the subtilisins in the narrower sense, proteinase K, and the proteases TW3 and TW7, and variants of the described proteases which have a modified amino acid sequence compared to the starting protease. Proteases are modified deliberately or randomly using methods known from the prior art and thus optimized for the use in detergents. These include point mutagenesis, deletion or insertion mutagenesis or fusion with other proteins or protein fragments. Appropriately optimized variants are known for the majority of proteases known from the prior art.

[0060] Examples of amylases that may be used include the α -amylases from *Bacillus licheniformis*, from *B. amyloliquefaciens* and from *B. stearothermophilus*, from *Aspergillus niger* and *A. oryzae*, and the refinements of the aforementioned amylases improved for the use in detergents. For this purpose, furthermore the α -amylase from *Bacillus sp.* A 7-7 (DSM 12368) and the cyclodextrin glucanotransferase (CGTase) from *B. agaradherens* (DSM 9948) shall be emphasized.

[0061] Furthermore, lipases or cutinases may be used, in particular for the triglyceride-liberating activities thereof, but also so as to create peroxy acids *in situ* from suitable precursors. These include, for example, the lipases which were originally obtainable or further developed from *Humicola lanuginosa* (*Thermomyces lanuginosus*), in particular those including the D96L amino acid substitution.

[0062] Furthermore, enzymes that are combined under the term hemicellulases may be used. These include, for example, mannanases, xanthan lyases, pectin lyases (= pectinases), pectin esterases, pectate lyases, xyloglucanases (= xylanases), pullulanases and β -glucanases

[0063] To enhance the bleaching action, oxidoreductases, for example oxidases, oxygenases, catalases, peroxidases, such as haloperoxidases, chloroperoxidases, bromoperoxidases, lignin peroxidases, glucose peroxidases or manganese peroxidases, dioxygenases or laccases (phenol oxidases, polyphenol oxidases) may be used. Advantageously, preferably organic, and particularly preferably aromatic, compounds that interact with the enzymes are additionally added so as to enhance the activity of the particular oxidoreductases (enhancers), or so as to ensure the electron flux in the event of large differences in the redox potentials of the oxidizing enzymes and the soils (mediators).

[0064] In particular during storage, an enzyme can be protected against damage, such as inactivation, denaturing or

disintegration, for example due to physical influences, oxidation or proteolytic cleavage. Inhibiting proteolysis is particularly preferred in the case of microbial production of the proteins and/or enzymes, in particular when the detergents comprise proteases. Detergents may comprise stabilizers for this purpose.

5 [0065] Proteases and amylases with cleaning action are generally not provided in form of the pure protein, but rather in the form of stabilized, storable and transportable preparations. These preformulated preparations include, for example, solid preparations obtained by way of granulation, extrusion or lyophilization or, in particular in the case of liquid or gel-like detergents, solutions of the enzymes, advantageously concentrated to the extent possible, low-hydrate and/or mixed with stabilizers or other auxiliary agents. Alternatively, the enzymes can be encapsulated, both for the solid and the liquid packaging format, for example by spray drying or extruding the enzyme solution together with a preferably natural polymer, or in the form of capsules, for example those in which the enzymes are enclosed as in a solidified gel, or in those of the core-shell type, in which an enzyme-containing core is coated with a protective layer impervious to water, air and/or chemicals. Further active ingredients, such as stabilizers, emulsifiers, pigments, bleaching agents or dyes can additionally be applied in superimposed layers. Such capsules are applied using methods that are known per se, for example agitation or roll granulation or in fluid bed processes. Such granules are advantageously low-dust, for example by applying polymeric film formers, and storage-stable due to the coating. It is furthermore possible to formulate two or more enzymes together, so that individual granules have multiple enzyme activities.

10 [0066] As is apparent from the comments above, the enzyme protein forms only a fraction of the total weight of customary enzyme preparations. Preferably used protease and amylase preparations contain between 0.1 and 40 wt.%, preferably between 0.2 and 30 wt.%, particularly preferably between 0.4 and 20 wt.%, and in particular between 0.8 and 20 wt.% of the enzyme protein. In particular, detergents that comprise 0.1 to 12 wt.%, preferably 0.2 to 10 wt.%, and in particular 0.5 to 8 wt.% enzyme preparations are preferred.

15 [0067] The compositions described herein can also comprise enzyme stabilizers. One group of stabilizers is that of reversible protease inhibitors. Frequently, benzamidine hydrochloride, borax, boric acids, boronic acids or the salts or esters thereof are used for this purpose, including especially derivatives with aromatic groups, such as ortho-, meta- or para-substituted phenylboronic acids, in particular 4-formylphenylboronic acid, or the salts or esters of the compounds mentioned. Peptide aldehydes, which is to say oligopeptides having a reduced C terminus, and in particular those formed from 2 to 50 monomers, are used for this purpose. The peptidic reversible protease inhibitors include ovomucoid and leupeptin. Specific reversible peptide inhibitors for the protease subtilisin and fusion proteins formed from proteases and specific peptide inhibitors are also suitable for this purpose.

25 [0068] Further enzyme stabilizers are amino alcohols such as mono-, di-, triethanol- and propanolamine and the mixtures thereof, aliphatic carboxylic acids up to C₁₂, such as succinic acid, other dicarboxylic acids or salts of the acids mentioned. End-capped fatty acid amide alkoxylates are also suitable for this purpose. Other enzyme stabilizers are known to a person skilled in the art from the prior art.

30 [0069] Peroxygen-based bleaching agents may also be present in the detergents. Among the compounds that serve as bleaching agents and yield H₂O₂ in water, sodium percarbonate, sodium perborate tetrahydrate, and sodium perborate monohydrate are of particular importance. Further usable bleaching agents are, for example, peroxyphosphates, citrate perhydrates, and peracid salts or peracids that yield H₂O₂, such as perbenzoates, peroxophthalates, diperazelaic acid, phthalimino peracid, or diperdodecanedioic acid. All further inorganic or organic peroxy bleaching agents known from the prior art to a person skilled in the art may also be used. Percarbonates, and in particular sodium percarbonate, are particularly preferred bleaching agents.

35 [0070] In various embodiments, the detergents may comprise 1 to 35 wt.%, preferably 2.5 to 30 wt.%, particularly preferably 3.5 to 20 wt.%, and in particular 5 to 15 wt.% bleaching agent, preferably sodium percarbonate.

40 [0071] In various embodiments of the invention, the detergents may additionally comprise at least one bleach activator. Compounds that, under perhydrolysis conditions, yield aliphatic peroxocarboxylic acids having preferably 1 to 10 carbon atoms, in particular 2 to 4 carbon atoms, and/or optionally substituted perbenzoic acid, can be used as bleach activators. Out of all bleach activators known to a person skilled in the art from the prior art, polyacylated alkylenediamines, in particular tetra acetyl ethylene diamine (TAED), acylated triazine derivatives, in particular 1,5-diacetyl-2,4-dioxohexahydro-1,3,5-triazine (DADHT), acylated glycolurils, in particular tetraacetyl glycoluril (TAGU), N-acylimides, in particular N-nonanoyl succinimide (NOSI), acylated phenolsulfonates, in particular n-nonanoyl or iso-nonanoyl oxybenzene sulfonate (n- or iso-NOBS), are particularly preferred. It is also possible to use combinations of conventional bleach activators. TAED, in particular in combination with a percarbonate bleaching agent, preferably sodium percarbonate, is an especially particularly preferred bleach activator.

45 [0072] These bleach activators are preferably used in amounts of up to 10 wt.%, in particular 0.1 wt.% to 8 wt.%, particularly 2 to 8 wt.%, and particularly preferably 2 to 6 wt.%.

50 [0073] In general, the pH value of the detergent can be set using customary pH regulators, wherein the pH value is selected depending on the desired usage purpose. In various embodiments, the pH value is in a range of from 5.5 to 10.5, preferably from 5.5 to 9.5, still more preferably from 7 to 9, in particular greater than 7, and especially in the range of from 7.5 to 8.5, if the detergent is an automatic dishwasher detergent. If the detergent is a laundry detergent, the pH

value may be higher and for example be in the range of from 8 to 12. Acids and/or alkalis, preferably alkalis, may be used for pH adjustment. Usable acids are in particular organic acids, such as acetic acid, citric acid, glycolic acid, lactic acid, succinic acid, adipic acid, malic acid, tartaric acid, and gluconic acid, or amidosulfonic acid. In addition, however, it is also possible to use the mineral acids hydrochloric acid, sulfuric acid and nitric acid, or the mixtures thereof. Suitable bases come from the group of the alkali metal and alkaline earth metal hydroxides and carbonates, in particular the alkali metal hydroxides, among which potassium hydroxide and especially sodium hydroxide is preferred. Volatile alkali, however, is particularly preferred, for example in the form of ammonia and/or alkanolamines, which can contain up to 9 carbon atoms in the molecule. The alkanolamine is preferably selected from the group consisting of mono-, di-, triethanol- and -propanolamine and the mixtures thereof.

[0074] So as to set and/or stabilize the pH value, the detergent according to the invention can also comprise one or more buffer substances (INCI Buffering Agents), usually in amounts of 0.001 wt.% to 5 wt.%. Buffering agents that simultaneously are complexing agents or even chelating agents (chelators, INCI Chelating Agents) are preferred. Particularly preferred buffering agents are citric acid or the citrates, and in particular sodium and potassium citrates, such as trisodium citrate·2H₂O and tripotassium citrate·H₂O.

[0075] Glass corrosion inhibitors prevent the appearance of clouding, streaking, and scratching, but also iridescence of the glass surface of automatically cleaned glassware. Preferred glass corrosion inhibitors come from the group of magnesium and zinc salts and of the magnesium and zinc complexes. Within the scope of the present invention, the content of zinc salt, especially in dishwasher detergents, is especially between 0.1 wt.% and 5 wt.%, preferably between 0.2 wt.% and 4 wt.%, and in particular between 0.4 wt.% and 3 wt.%, or the content of zinc (calculated as Zn²⁺) is between 0.01 and 1 wt.%, especially between 0.02 and 0.5 wt.%, and in particular between 0.04 and 0.2 wt.%.

[0076] Individual odorous substance compounds, such as synthetic products of the ester, ether, aldehyde, ketone, alcohol, and hydrocarbon types, can be used as perfume oils or fragrances. Preferably, however, mixtures of different odorants are used, which together produce an appealing odorous note. Such perfume oils can also contain natural odorous substance mixtures such as those accessible from plant sources, for example pine, citrus, jasmine, patchouli, rose, or ylang ylang oil.

[0077] Furthermore, preservatives can be present in detergents. Suitable preservatives are, for example, those from the groups of the alcohols, aldehydes, antimicrobial acids and/or the salts thereof, carboxylic acid esters, acid amides, phenols, phenol derivatives, diphenyls, diphenyl alkanes, urea derivatives, oxygen and nitrogen acetals and formals, benzamidines, isothiazoles and the derivatives thereof, such as isothiazolins and isothiazolinones, phthalimide derivatives, pyridine derivatives, antimicrobial surface-active compounds, guanidines, antimicrobial amphoteric compounds, quinolines, 1,2-dibromo-2,4-dicyanobutane, iodo-2-propynyl butyl carbamate, iodine, iodophores, and peroxides. Preferred antimicrobial active ingredients are preferably selected from the group consisting of ethanol, n-propanol, i-propanol, 1,3-butanediol, phenoxyethanol, 1,2-propylene glycol, glycerol, undecylenic acid, citric acid, lactic acid, benzoic acid, salicylic acid, thymol, 2-benzyl-4-chlorophenol, 2,2'-methylene-bis-(6-bromo-4-chlorophenol), 2,4,4'-trichloro-2'-hydroxydiphenyl ether, N-(4-chlorophenyl)-N-(3,4-dichlorophenyl) urea, N,N'-(10-decandiyldi-1-pyridinyl-4-ylidene)-bis-(1-octanamine)-dihydrochloride, N,N'-bis-(4-chlorophenyl)-3,12-diimino-2,4,11,13-tetraazatetra-decane diimidamide, antimicrobial quaternary surface-active compounds, and guanidines. Particularly preferred preservatives, however, are selected from the group consisting of salicylic acid, quaternary surfactants, and in particular benzalkonium chloride and isothiazoles, and the derivatives thereof, such as isothiazolines and isothiazolinones.

[0078] The detergents can be in solid or liquid presentation forms or in a combination of solid and liquid presentation forms. Suitable solid presentation forms are, in particular, powders, granules, extrudates, compactates, and in particular tablets. The liquid presentation forms based on water and/or organic solvents can be present in thickened form, in the form of gels. The detergents can be formulated in the form of single-phase or multi-phase products. The individual phases of multiphase detergents can have identical or different states of aggregation. The detergent can also be present in the form of shaped bodies. So as to facilitate the breakdown of such prefabricated shaped bodies, it is possible to incorporate disintegration auxiliaries, known as tablet disintegrants, into these agents in order to shorten breakdown times. Tablet disintegrants or disintegration accelerators are understood to mean auxiliaries that ensure a rapid breakdown of tablets in water or other media, and the quick release of the active ingredients. Disintegration auxiliaries can preferably be used in amounts from 0.5 wt.% to 10 wt.%, preferably 3 wt.% to 7 wt.%, and in particular 4 wt.% to 6 wt.%.


[0079] The detergents may be preformulated as dosing units. These dosing units preferably comprise the quantity of substances with cleaning action necessary for one cleaning cycle. Preferred dosing units have a weight between 12 g and 30 g, preferably between 14 g and 26 g, and in particular between 16 g and 22 g. The volume of the aforementioned dosing units and the three-dimensional shape thereof are particularly preferably selected such that dosability of the preformulated units via the dosing chamber of a washing machine or dishwashing machine is ensured. The volume of the dosing unit is thus preferably between 10 ml and 35 ml, and especially between 12 ml and 30 ml. The prefabricated dosing units preferably comprise a water-soluble wrapping. The water-soluble wrapping may, for example, be formed of a water-soluble film material selected from the group consisting of polymers or polymer mixtures. The wrapping can be formed of one layer, or of two or more layers of the water-soluble film material. The water-soluble film material of the

first layer and that of the further layers, if such are present, can be the same or different. Films that can be bonded and/or sealed, after they have been loaded with the detergent, to form packaging such as tubes or cushions, are particularly preferred. The water-soluble packaging can comprise one or more chambers. The detergent can be present in one or more chambers, if present, of the water-soluble wrapping. The amount of the detergent preferably corresponds to the full dose, or half the dose, that is required for one washing operation. It is preferable for the water-soluble wrapping to comprise polyvinyl alcohol or a polyvinyl alcohol copolymer. Water-soluble wrappings comprising polyvinyl alcohol, or a polyvinyl alcohol copolymer exhibit good stability and sufficiently high water solubility, in particular cold-water solubility. Suitable water-soluble films for producing the water-soluble wrapping are preferably based on a polyvinyl alcohol, or a polyvinyl alcohol copolymer, having a relative molar mass in the range from 10,000 g mol⁻¹ to 1,000,000 g mol⁻¹, preferably from 20,000 g mol⁻¹ to 500,000 g mol⁻¹, particularly preferably from 30,000 g mol⁻¹ to 100,000 g mol⁻¹, and in particular from 40,000 g mol⁻¹ to 80,000 g mol⁻¹. Polyvinyl alcohol is typically produced by the hydrolysis of polyvinyl acetate. The same applies to polyvinyl alcohol copolymers produced accordingly from polyvinyl acetate copolymers and subsequent hydrolysis. It is preferred if at least one layer of the water-soluble wrapping comprises a polyvinyl alcohol having a degree of hydrolysis of 70 mole % to 100 mole %, preferably 80 mole % to 90 mole %, particularly preferably 81 mole % to 89 mole %, and in particular 82 mole % to 88 mole %. Additionally, a polymer selected from the group consisting of (meth)acrylic acid-containing (co)polymers, polyacrylamides, oxazoline polymers, polystyrene sulfonates, polyurethanes, polyesters, polyethers, polylactic acid or mixtures of the above polymers can be added to a polyvinyl alcohol-containing film material that is suitable for producing the water-soluble wrapping. A preferred additional polymer is polylactic acids. In addition to vinyl alcohol, preferred polyvinyl alcohol copolymers comprise dicarboxylic acids as further monomers. Suitable dicarboxylic acids are itaconic acid, malonic acid, succinic acid and mixtures thereof, itaconic acid being preferred. Likewise, preferred polyvinyl alcohol copolymers include an ethylenically unsaturated carboxylic acid, the salt thereof, or the ester thereof, in addition to vinyl alcohol. In addition to vinyl alcohol, such polyvinyl alcohol copolymers particularly preferably comprise acrylic acid, methacrylic acid, acrylic acid esters, methacrylic acid esters or mixtures thereof. It may be preferred for the film material to contain further additives. For example, the film material may contain plasticizers such as dipropylene glycol, ethylene glycol, diethylene glycol, propylene glycol, glycerol, sorbitol, mannitol or mixtures thereof. Examples of further additives include release aids, fillers, cross-linking agents, surfactants, antioxidants, UV absorbers, antiblocking agents, non-stick agents or mixtures thereof. Suitable water-soluble films for use in the water-soluble wrappings of the water-soluble packagings according to the invention are films sold by MonoSol LLC, for example, by the designation M8630, C8400 or M8900. Other suitable films include films by the designation Solublon® PT, Solublon® GA, Solublon® KC or Solublon® KL from Aicello Chemical Europe GmbH, or the VF-HP films from Kuraray.

Examples

Example 1: Synthesis of complex K1 from 1,4,7-triazacyclononane-1,4,7-triacetic acid and Ce(III)

[0080]



[0081] To a solution of the tri sodium salt of commercially available 1,4,7-triazacyclononane-1,4,7-triacetic acid (1.5 mmol) in 30 ml of methanol were added 0.6 g of 7M ammonia solution in methanol, resulting in pH 8 to 9. The solution was heated to 60 °C and 1.5 ml of CeCl₃ (1.5 mmol) in 4 ml methanol were added dropwise while stirring. The reaction mixture was stirred for 44 h at 65 °C, then cooled to room temperature, and the solvent was removed in vacuum to yield the complex K1.

Example 2: Synthesis of complex K2 from N,N,N-Tris(2-hydroxybenzyl)-1,4,7-triazacyclononane and Ce(III)


[0082]

5

10

15

20



[0083] To a solution of N,N,N-Tris(2-hydroxybenzyl)-1,4,7-triazacyclononane (0.79 mmol) in 40 ml of methanol were added 6.3 ml of 0.5 M KOH solution in methanol, resulting in pH 10. After heating to 60 °C, a solution of CeCl₃ (0.79 mmol) in 5 ml methanol was slowly added while stirring. The reaction mixture was stirred for 48 h at 70 °C and then cooled to room temperature. The solvent was removed in vacuum to yield complex K2.

25

Example 3: Hydrolysis of burnt-in soil

[0084] A mixture of lactose and albumin from chicken egg, approx. 45 000 g/mol, in a weight ratio of 3:1 was dissolved in de-salinated water and stirred for 30 min at room temperature. The resulting solution was lyophilized using a commercial lab lyophilizer (Christ; Gamma 1-20) and homogenized. 0.5 g of the resulting powder was heated to 140°C for 20 min and was then allowed to cool to room temperature. Characterisation of the molecular weight distribution was performed by SDS-PAGE elektrophoresis; bands at -50-55 kDa and -100-110 kDa were detected, indicating modified or cross-linked proteins that were results of the Maillard reaction between lactose und albumin. An aqueous solution of the Maillard product (3 g/l) was incubated at 60 °C at pH 9.5 (hydrogen carbonate / carbonate buffer) with the complexes of examples 1 or 2 at concentrations of 1 mM and for amounts of time given in table 1. The progress of protein degradation was followed by SDS page gel analysis. The intensity value of the band at -50-55 kDa at time t=0, that is before the start of incubation, is set to 1. The smaller the value after incubation is, the stronger was the degradation of the soil.

30

35

40

Table 1: Intensity of the band at -50-55 kDa

Complex	t=0h	t=2h	t=5h
-	1	1.18	0.87
K1	1	0,83	0,44
K2	1	0,74	n.d.
n.d.: not determined			


45

50

Claims

1. Use of a complex of formula I, or of a detergent comprising a complex of formula I,

55




in which the metal ions Mⁿ⁺ are selected from Al³⁺, Ti⁴⁺, Y³⁺, Zr⁴⁺, Ce³⁺, Ce⁴⁺, Sc³⁺, Yb³⁺, La³⁺, Ta⁵⁺ and their mixtures,

EP 3 795 667 B1

n stands for a number in the range of 1 to 5,
L stands for a ligand according to formula II,

5



10

in which

15

E is selected from O, NR¹, and mixtures of them,
R¹ is selected from H, optionally substituted C₁₋₂₀-alkyl or heteroalkyl groups, optionally substituted C₂₋₂₀-alkenyl, heteroalkenyl or alkinyl groups, optionally substituted aryl or heteroaryl groups, and optionally substituted C₁₋₂₀-alkylaryl or heteroaryl groups,
m stands for a number in the range of from 0 to 3,

20

A stands for an alkali metal cation and q stands for 1, an earth alkali metal cation and q stands for 2, or an ammonium cation and q stands for 1,

25

X^v stands for an anion selected from the group of F⁻, Cl⁻, Br⁻, I⁻, OH⁻, HSO₃⁻, SO₃²⁻, SO₄²⁻, HSO₄⁻, NO₂⁻, NO₃⁻, PO₄³⁻, HPO₄²⁻, H₂PO₄⁻, BF₄⁻, PF₆⁻, ClO₄⁻, organic anions such as acetate, citrate, formate, glutarate, lactate, malate, malonate, oxalate, pyruvate, tartrate, methanesulfonate, methyl sulfate, tosylate, succinate and their mixtures, with v being 1, 2 or 3, respectively, and

p and r, irrespective of each other, are numbers in the range of from 0 to 7, selected such that the charge of the complex according to formula I is 0


to remove cooked-, baked- and burnt-on proteinaceous soils from surfaces, preferably hard surfaces, or textile surfaces.

30

2. Use according to claim 1, **characterized in that** the detergent comprises a complex of formula I in amounts of from 0.001 % by weight to 20 % by weight, preferably from 0.01 % by weight to 3 % by weight.

35

3. Method of removing cooked-, baked- and burnt-on proteinaceous soils from surfaces, preferably hard surfaces, or textile surfaces, by contacting the soiled surface with an aqueous liquid comprising a complex of formula I,




40

in which the metal ions Mⁿ⁺ are selected from Al³⁺, Ti⁴⁺, Y³⁺, Zr⁴⁺, Ce³⁺, Ce⁴⁺, Sc³⁺, Yb³⁺, La³⁺, Ta⁵⁺ and their mixtures,

n stands for a number in the range of 1 to 5,

L stands for a ligand according to formula II,

45



50

in which

55

E is selected from O, NR¹, and mixtures of them,
R¹ is selected from H, optionally substituted C₁₋₂₀-alkyl or heteroalkyl groups, optionally substituted C₂₋₂₀-alkenyl, heteroalkenyl or alkinyl groups, optionally substituted aryl or heteroaryl groups, and optionally substituted C₁₋₂₀-alkylaryl or heteroaryl groups,
m stands for a number in the range of from 0 to 3,

A stands for an alkali metal cation and q stands for 1, an earth alkali metal cation and q stands for 2, or an

ammonium cation and q stands for 1,

X^{v-} stands for an anion selected from the group of F⁻, Cl⁻, Br⁻, I⁻, OH⁻, HSO₃⁻, SO₃²⁻, SO₄²⁻, HSO₄⁻, NO₂⁻, NO₃⁻, PO₄³⁻, HPO₄²⁻, H₂PO₄⁻, BF₄⁻, PF₆⁻, ClO₄⁻, organic anions such as acetate, citrate, formate, glutarate, lactate, malate, malonate, oxalate, pyruvate, tartrate, methanesulfonate, methyl sulfate, tosylate, succinate and their mixtures, with v being 1, 2 or 3, respectively, and

p and r, irrespective of each other, are numbers in the range of from 0 to 7, selected such that the charge of the complex according to formula I is 0.


4. Method according to claim 3, **characterized in that** the aqueous liquid comprises the complex in concentrations of from 0.001 mmol/l to 10 mmol/l, preferably from 0.01 mmol/l to 5 mmol/l.

5. Method according to claim 3 or 4, **characterized in that** the contact of the aqueous liquid with the proteinaceous soil is for a time period in the range of from 2 minutes to 24 hours, preferably from 10 minutes to 12 hours.

6. Use according to claim 1 or 2, or method according to claim 3 to 5, **characterized in that** heteroatoms in group R¹ are selected from O, N, S, Si, P, and their mixtures.

7. Use according to claim 1 or 2 or 6, or method according to claim 3 to 6, **characterized in that** substituents in group R¹ are selected from -COOH, -COO⁻, -OR², and their mixtures, with R² = H, or C₁₋₆ alkyl.


8. Use according to claim 1 or 2 or 6 or 7, or method according to claim 3 to 7, **characterized in that** R¹ is selected from C₁₋₁₈ alkyl groups,



, and their mixtures.

Patentansprüche


1. Verwendung eines Komplexes von Formel I oder eines Waschmittels umfassend einen Komplex der Formel I,



in dem die Metallionen Mⁿ⁺ aus Al³⁺, Ti⁴⁺, Y³⁺, Zr⁴⁺, Ce³⁺, Ce⁴⁺, Sc³⁺, Yb³⁺, La³⁺, Ta⁵⁺ und ihren Mischungen ausgewählt sind,

n für eine Zahl in dem Bereich von 1 bis 5 steht,

L für einen Ligand gemäß Formel II steht,



wobei

EP 3 795 667 B1

E aus O, NR¹ und Mischungen davon ausgewählt ist,

R¹ ausgewählt ist aus H, gegebenenfalls substituiertem C₁₋₂₀-Alkyl- oder Heteroalkylgruppen, gegebenenfalls substituiertem C₂₋₂₀-Alkenyl-, Heteroalkenyl- oder Alkinygruppen, gegebenenfalls substituiertem Aryl- oder Heteroarylgruppen und gegebenenfalls substituiertem C₁₋₂₀-Alkylaryl oder Heteroarylgruppen,

m für eine Zahl in dem Bereich von 0 bis 3 steht,

A für ein Alkalimetallkation steht und q für 1 steht, ein Erdalkalimetallkation und q für 2 steht oder ein Ammoniumkation und q für 1 steht,


X^{v-} für ein Anion steht, das aus der Gruppe von F⁻, Cl⁻, Br⁻, I⁻, OH⁻, HSO₃⁻, SO₃²⁻, SO₄²⁻, HSO₄⁻, NO₂⁻, NO₃⁻, PO₄³⁻, HPO₄²⁻, H₂PO₄⁻, BF₄⁻, PF₆⁻, ClO₄⁻, organischen Anionen wie etwa Acetat, Citrat, Formiat, Glutarat, Lactat, Malat, Maionat, Oxalat, Pyruvat, Tartrat, Methansulfonat, Methylsulfat, Tosylat, Succinat und ihren Mischungen ausgewählt ist, wobei v 1, 2 beziehungsweise 3 ist, und

p und r unabhängig voneinander Zahlen in dem Bereich von 0 bis 7 sind, die derart ausgewählt sind, dass die Ladung des Komplexes gemäß Formel I 0 ist

zum Entfernen von gekochten, gebackenen und angebrannten proteinhaltigen Verschmutzungen von Oberflächen, vorzugsweise harten Oberflächen oder textilen Oberflächen.

2. Verwendung nach Anspruch 1, **dadurch gekennzeichnet, dass** das Waschmittel einen Komplex der Formel I in Mengen von 0,001 Gew.-% bis 20 Gew.-%, vorzugsweise von 0,01 Gew.-% bis 3 Gew.-%, umfasst.


3. Verfahren zum Entfernen von gekochten, gebackenen und gebrannten proteinhaltigen Verschmutzungen von Oberflächen, vorzugsweise harten Oberflächen oder textilen Oberflächen, durch Inberührungbringen der verschmutzten Oberfläche mit einer wässrigen Flüssigkeit, umfassend einen Komplex der Formel I,



in dem die Metallionen Mⁿ⁺ aus Al³⁺, Ti⁴⁺, Y³⁺, Zr⁴⁺, Ce³⁺, Ce⁴⁺, Sc³⁺, Yb³⁺, La³⁺, Ta⁵⁺ und ihre Mischungen ausgewählt sind,

n für eine Zahl in dem Bereich von 1 bis 5 steht,

L für einen Ligand gemäß Formel II steht,



wobei

E aus O, NR¹ und Mischungen davon ausgewählt ist,

R¹ ausgewählt ist aus H, gegebenenfalls substituiertem C₁₋₂₀-Alkyl- oder Heteroalkylgruppen, gegebenenfalls substituiertem C₂₋₂₀-Alkenyl-, Heteroalkenyl- oder Alkinygruppen, gegebenenfalls substituiertem Aryl- oder Heteroarylgruppen und gegebenenfalls substituiertem C₁₋₂₀-Alkylaryl oder Heteroarylgruppen,

m für eine Zahl in dem Bereich von 0 bis 3 steht,

A für ein Alkalimetallkation steht und q für 1 steht, ein Erdalkalimetallkation und q für 2 steht oder ein Ammoniumkation und q für 1 steht,

X^{v-} für ein Anion steht, das aus der Gruppe von F⁻, Cl⁻, Br⁻, I⁻, OH⁻, HSO₃⁻, SO₃²⁻, SO₄²⁻, HSO₄⁻, NO₂⁻, NO₃⁻, PO₄³⁻, HPO₄²⁻, H₂PO₄⁻, BF₄⁻, PF₆⁻, ClO₄⁻, organischen Anionen wie etwa Acetat, Citrat, Formiat, Glutarat, Lactat, Malat, Maionat, Oxalat, Pyruvat, Tartrat, Methansulfonat, Methylsulfat, Tosylat, Succinat und ihren Mischungen ausgewählt ist, wobei v 1, 2 beziehungsweise 3 ist, und


p und r unabhängig voneinander Zahlen in dem Bereich von 0 bis 7 sind, die derart ausgewählt sind, dass die Ladung des Komplexes gemäß Formel I 0 ist.

4. Verfahren nach Anspruch 3, **dadurch gekennzeichnet, dass** die wässrige Flüssigkeit den Komplex in Konzentrationen von 0,001 mmol/L bis 10 mmol/L, vorzugsweise von 0,01 mmol/L bis 5 mmol/L, umfasst.

5. Verfahren nach Anspruch 3 oder 4, **dadurch gekennzeichnet, dass** die Berührung der wässrigen Flüssigkeit mit

dem proteinhaltigen Verschmutzungen für einen Zeitraum in dem Bereich von 2 Minuten bis 24 Stunden, vorzugsweise von 10 Minuten bis 12 Stunden, erfolgt.


6. Verwendung nach Anspruch 1 oder 2 oder Verfahren nach Anspruch 3 bis 5, **dadurch gekennzeichnet, dass** Heteroatome in der Gruppe R¹ aus O, N, S, Si, P und ihren Mischungen ausgewählt sind.
7. Verwendung nach Anspruch 1 oder 2 oder 6 oder Verfahren nach Anspruch 3 bis 6, **dadurch gekennzeichnet, dass** Substituenten in der Gruppe R¹ aus -COOH, -COO⁻, -OR² und ihren Mischungen, mit R² = H oder C₁₋₆-Alkyl ausgewählt sind.
8. Verwendung nach Anspruch 1 oder 2 oder 6 oder 7 oder Verfahren nach Anspruch 3 bis 7, **dadurch gekennzeichnet, dass** R¹ aus C₁₋₁₈-Alkylgruppen,




und ihren Mischungen ausgewählt ist.

Revendications

1. Utilisation d'un complexe de formule I, ou d'un détergent comprenant un complexe de formule I,



dans laquelle les ions métalliques Mⁿ⁺ sont choisis parmi Al³⁺, Ti⁴⁺, Y³⁺, Zr⁴⁺, Ce³⁺, Ce⁴⁺, Sc³⁺, Yb³⁺, La³⁺, Ta⁵⁺ et leurs mélanges,
n représente un nombre compris entre 1 et 5,
L représente un ligand selon la formule II,



dans laquelle

E est choisi parmi O, NR¹, et leurs mélanges,
R¹ est choisi parmi H, des groupes alkyle ou hétéroalkyle en C₁₋₂₀ éventuellement substitués, des groupes alcényle, hétéroalcényle ou alcynyle en C₂₋₂₀ éventuellement substitués, des groupes aryle ou hétéroaryle éventuellement substitués et des groupes alkylaryle ou groupes hétéroaryle en C₁₋₂₀ éventuellement substitués,
m représente un nombre compris entre 0 et 3,
A représente un cation de métal alcalin et q représente 1, un cation de métal alcalino-terreux et q représente 2, ou un cation ammonium et q représente 1,

EP 3 795 667 B1


X^v représente un anion choisi dans le groupe de F⁻, Cl⁻, Br⁻, I⁻, OH⁻, HSO₃⁻, SO₃²⁻, SO₄²⁻, HSO₄⁻, NO₂⁻, NO₃⁻, PO₄³⁻, HPO₄²⁻, H₂PO₄⁻, BF₄⁻, PF₆⁻, ClO₄⁻, des anions organiques tels que acétate, citrate, formiate, glutarate, lactate, malate, maionate, oxalate, pyruvate, tartrate, méthanesulfonate, méthylsulfate, tosylate, succinate et leurs mélanges, v étant respectivement 1, 2 ou 3, et

p et r, indépendamment l'un de l'autre, sont des nombres compris entre 0 et 7, choisis de telle sorte que la charge du complexe selon la formule I est 0

pour éliminer les souillures protéiques cuites, cuites au four et brûlées de surfaces, de préférence des surfaces dures, ou des surfaces textiles.

2. Utilisation selon la revendication 1, **caractérisé en ce que** le détergent comprend un complexe de formule I en des quantités de 0,001 % en poids à 20 % en poids, de préférence de 0,01 % en poids à 3 % en poids.


3. Procédé d'élimination des souillures protéiques cuites, cuites au four et brûlées de surfaces, de préférence de surfaces dures, ou de surfaces textiles, en mettant en contact la surface souillée avec un liquide aqueux comprenant un complexe de formule I,



dans laquelle les ions métalliques Mⁿ⁺ sont choisis parmi Al³⁺, Ti⁴⁺, Y³⁺, Zr⁴⁺, Ce³⁺, Ce⁴⁺, Sc³⁺, Yb³⁺, La³⁺, Ta⁵⁺ et leurs mélanges,

n représente un nombre compris entre 1 et 5,

L représente un ligand selon la formule II,



dans laquelle

E est choisi parmi O, NR¹, et leurs mélanges,

R¹ est choisi parmi H, des groupes alkyle ou hétéroalkyle en C₁₋₂₀ éventuellement substitués, des groupes alcényle, hétéroalcényle ou alcynyle en C₂₋₂₀ éventuellement substitués, des groupes aryle ou hétéroaryle éventuellement substitués et des groupes alkylaryle ou groupes hétéroaryle en C₁₋₂₀ éventuellement substitués,

m représente un nombre compris entre 0 et 3,

A représente un cation de métal alcalin et q représente 1, un cation de métal alcalino-terreux et q représente 2, ou un cation ammonium et q représente 1,

X^v représente un anion choisi dans le groupe de F⁻, Cl⁻, Br⁻, I⁻, OH⁻, HSO₃⁻, SO₃²⁻, SO₄²⁻, HSO₄⁻, NO₂⁻, NO₃⁻, PO₄³⁻, HPO₄²⁻, H₂PO₄⁻, BF₄⁻, PF₆⁻, ClO₄⁻, des anions organiques tels que acétate, citrate, formiate, glutarate, lactate, malate, maionate, oxalate, pyruvate, tartrate, méthanesulfonate, méthylsulfate, tosylate, succinate et leurs mélanges, v étant respectivement 1, 2 ou 3, et

p et r, indépendamment l'un de l'autre, sont des nombres compris entre 0 et 7, choisis de telle sorte que la charge du complexe selon la formule I soit 0.

4. Procédé selon la revendication 3, **caractérisé en ce que** le liquide aqueux comprend le complexe à des concentrations de 0,001 mmol/l à 10 mmol/l, de préférence de 0,01 mmol/l à 5 mmol/l.

5. Procédé selon la revendication 3 ou 4, **caractérisé en ce que** le contact du liquide aqueux avec les souillures protéiques se produit pendant une durée comprise dans la plage de 2 minutes à 24 heures, de préférence de 10 minutes à 12 heures.


6. Utilisation selon la revendication 1 ou 2, ou procédé selon les revendications 3 à 5, **caractérisé en ce que** les hétéroatomes du groupe R¹ sont choisis parmi O, N, S, Si, P et leurs mélanges.

EP 3 795 667 B1


7. Utilisation selon la revendication 1 ou 2 ou 6, ou procédé selon les revendications 3 à 6, **caractérisé en ce que** les substituants du groupe R¹ sont choisis parmi -COOH, -COO⁻, -OR², et leurs mélanges, avec R² = H, ou alkyle en C₁₋₆.
8. Utilisation selon la revendication 1 ou 2 ou 6 ou 7, ou procédé selon la revendication 3 à 7, **caractérisé en ce que** R¹ est choisi parmi des groupes alkyles C₁₋₁₈,

5

10



15



20

et leurs mélanges.

25

30

35

40

45

50

55

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- EP 0458397 A2 [0003]
- EP 0286075 A2 [0003]
- WO 9744427 A1 [0003]
- WO 2005059076 A1 [0003]
- WO 9811186 A1 [0003]
- CA 2882331 A1 [0003]