

REPUBLIC OF SOUTH AFRICA
PATENTS ACT, 1978
PUBLICATION PARTICULARS AND ABSTRACT
[Section 32 (3) (a) - Regulations 22 (1) (g) and 31]

OFFICIAL APPLICATION NO.		LODGING DATE	ACCEPTANCE DATE
21	01 2006/03326	22 26 Apr 2006	43 09-05-2007

INTERNATIONAL CLASSIFICATION	
51	A01N C07D

Not for publication
Classified by:

FULL NAME(S) OF APPLICANT(S)	
71	BASF AKTIENGESELLSCHAFT

FULL NAME(S) OF INVENTOR(S)	
72	TORMO I BLASCO, Jordi; GROTE, Thomas; SCHERER, Maria; STIERL, Reinhard; STRATHMANN, Siegfried; SCHÖFL, Ulrich

TITLE OF INVENTION	
54	FUNGICIDAL MIXTURES

EARLIEST PRIORITY CLAIMED	COUNTRY	NUMBER	DATE
NB - Use International abbreviation for country (See Schedule 4)	33 DE	31 103 46 136.1	32 1 Oct 2003

57	Abstract (not more than 150 words) and figure of the drawings to which the abstract refers, are attached.
----	---

Number of sheets	31
------------------	----

(12) NACH DEM VERTRAG ÜBER DIE INTERNATIONALE ZUSAMMENARBEIT AUF DEM GEBIET DES PATENTWESENS (PCT) VERÖFFENTLICHTE INTERNATIONALE ANMELDUNG

(19) Weltorganisation für geistiges Eigentum
Internationales Büro



(43) Internationales Veröffentlichungsdatum
14. April 2005 (14.04.2005)

PCT

(10) Internationale Veröffentlichungsnummer
WO 2005/032256 A1

(51) Internationale Patentklassifikation⁷: A01N 43/90,
C07D 487/04

STIERL, Reinhard [DE/DE]; Jahnstr. 8, 67251 Freinsheim (DE). STRATHMANN, Siegfried [DE/DE]; Donnersbergstr. 9, 67117 Limburgerhof (DE). SCHÖFL, Ulrich [DE/DE]; Erlenstr. 8, 68782 Brühl (DE).

(21) Internationales Aktenzeichen: PCT/EP2004/010918

(22) Internationales Anmeldedatum:
30. September 2004 (30.09.2004)

(81) Bestimmungsstaaten (soweit nicht anders angegeben, für jede verfügbare nationale Schutzrechtsart): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(25) Einreichungssprache: Deutsch

(26) Veröffentlichungssprache: Deutsch

(30) Angaben zur Priorität:
103 46 136.1 1. Oktober 2003 (01.10.2003) DE

(71) Anmelder (für alle Bestimmungsstaaten mit Ausnahme von US): BASF AKTIENGESELLSCHAFT [—/DE]; 67056 Ludwigshafen (DE).

(84) Bestimmungsstaaten (soweit nicht anders angegeben, für jede verfügbare regionale Schutzrechtsart): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), eurasisches (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), europäisches (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPK (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(72) Erfinder; und

(75) Erfinder/Anmelder (nur für US): TORMO I BLASCO, Jordi [ES/DE]; Carl-Benz-Str. 10-3, 69514 Laudendach (DE). GROTE, Thomas [DE/DE]; Im Höhnhausen 18, 67157 Wachenheim (DE). SCHERER, Maria [DE/DE]; Hermann-Jürgens-Str. 30, 76829 Godramstein (DE).


Veröffentlicht:

— mit internationalem Recherchenbericht

[Fortsetzung auf der nächsten Seite]

(54) Title: FUNGICIDAL MIXTURES

(54) Bezeichnung: FUNGIZIDE MISCHUNGEN



(57) Abstract: The invention relates to fungicidal mixtures that contain as the active component 1) the triazolopyrimidine derivative of formula (I), and 2) tebuconazoles of formula (II), in a synergistically effective amount. The invention also relates to a method for controlling parasitic fungi from the class of oomycetes with mixtures of compound I with compound II for producing mixtures of the aforementioned kind and to agents that contain said mixtures..


(57) Zusammenfassung: Fungizide Mischungen, enthaltend als aktive Komponenten 1) das Triazolopyrimidinderivat der Formel (I), und 2) Tebuconazole der Formel (II), in einer synergistisch wirksamen Menge, Verfahren zur Bekämpfung von Schadpilzen aus der Klasse der Oomyceten mit Mischungen der Verbindung I mit der Verbindung II und die Verwendung der Verbindung I mit der Verbindung II zur Herstellung derartiger Mischungen sowie Mittel, die diese Mischungen enthalten.

WO 2005/032256 A1

Fungicidal mixtures

The present invention relates to fungicidal mixtures comprising, as active components,


- 5 1) the triazolopyrimidine derivative of the formula I,



I

and

- 2) tebuconazole of the formula II,



II

10

in a synergistically effective amount.

Moreover, the invention relates to a method for controlling harmful fungi from the class
of the *Oomycetes* using mixtures of the compound I with the compound II and to the
15 use of the compound I with the compound II for preparing such mixtures and
compositions comprising these mixtures.

The compound I, 5-chloro-7-(4-methylpiperidin-1-yl)-6-(2,4,6-trifluorophenyl)-[1,2,4]tri-
azolo[1,5-a]pyrimidine, its preparation and its action against harmful fungi are known
20 from the literature (WO 98/46607).

The compound II, (R,S)-1-(4-chlorophenyl)-4,4-dimethyl-3-(1H-1,2,4-triazol-1-
ylmethyl)pentan-3-ol, its preparation and its action against harmful fungi are also
known from the literature (EP-A 40 345; common name: tebuconazole). Tebuconazole
25 has been commercially established for a long time as a fungicide against pathogens of
cereals

Mixtures of triazolopyrimidine derivatives with tebuconazole are known in a general
manner from EP-A 988 790. The compound I is embraced by the general disclosure of
30 this publication, but not explicitly mentioned. Accordingly, the combination of compound
I with tebuconazole is novel.

The synergistic mixtures of triazolopyrimidines described in EP-A 988 790 are described as being fungicidally active against various diseases of cereals, fruit and vegetables, in particular mildew on wheat and barley or gray mold on apples. However,
5 the fungicidal action of these mixtures against harmful fungi from the class of the *Oomycetes* is unsatisfactory.

The biological behavior of *Oomycetes* is clearly different from that of the *Ascomycetes*, *Deuteromycetes* and *Basidiomycetes*, since *Oomycetes* are biologically closer related
10 to algae than to fungi. Accordingly, what is known about the fungicidal activity of active compounds against "true fungi" such as *Ascomycetes*, *Deuteromycetes* and *Basidiomycetes* can be applied only to a very limited extent to *Oomycetes*.

Oomycetes cause economically relevant damage to various crop plants. In many
15 regions, infections by *Phytophthora infestans* in the cultivation of potatoes and tomatoes are the most important plant diseases. In viticulture, considerable damage is caused by peronospora of grapevines.

There is a constant demand for novel compositions against *Oomycetes* in agriculture,
20 since there is already widespread resistance of the harmful fungi to the products established in the market, such as, for example, metalaxyl and active compounds of a similar structure.

Practical agricultural experience has shown that the repeated and exclusive application
25 of an individual active compound in the control of harmful fungi leads in many cases to a rapid selection of fungus strains which have developed natural or adapted resistance against the active compound in question. Effective control of these fungi with the active compound in question is then no longer possible.

30 To reduce the risk of selection of resistant fungus strains, mixtures of different active compounds are nowadays preferably employed for controlling harmful fungi. By combining active compounds having different mechanisms of action, it is possible to ensure a successful control over a relatively long period of time.

35 A need exists to provide, with a view to effective resistance management and an effective control of harmful fungi from the class of the *Oomycetes* at application rates which are as low as possible, mixtures which, at a reduced total amount of active compounds applied, have a satisfactory effect against the harmful fungi.

40

AMENDED SHEET

We have found that this need is fulfilled by the mixtures defined at the outset. Moreover, we have found that simultaneous, that is joint or separate, application of the compound I and the compound II or successive application of the compound I and the compound II allows better control of *Oomyces* than is possible with the individual
5 compounds (synergistic mixtures).

The mixtures of the compound I and the compound II or the simultaneous, that is joint or separate, use of the compound I and the compound II are distinguished by being very highly active against phytopathogenic fungi from the class of the *Oomyces*, in
10 particular of *Phytophthora infestans* on potatoes and tomatoes and *Plasmopara viticola* on grapevines. They can be used in crop protection as foliar- and soil-acting fungicides.

They are particularly important for controlling *Oomyces* on various crop plants such as vegetable plants (for example cucumbers, beans and cucurbits), potatoes,
15 tomatoes, grapevines and the corresponding seeds.

They are particularly suitable for controlling late blight on tomatoes and potatoes caused by *Phytophthora infestans* and downy mildew of grapevines (peronospora of grapevines) caused by *Plasmopara viticola*.
20

In addition, the combination according to the invention of the compounds I and II is also suitable for controlling other pathogens such as, for example, *Septoria* and *Puccinia* species in cereals and *Alternaria* and *Botrytis* species in vegetables, fruit and grapevines.
25

When preparing the mixtures, it is preferred to employ the pure active compounds I and II, to which further active compounds against harmful fungi or against other pests, such as insects, arachnids or nematodes, or else herbicidal or growth-regulating active compounds or fertilizers can be added according to need.
30

Other suitable active compounds in the above sense are in particular fungicides selected from the following groups:

- acylalanines, such as benalaxyl, metalaxyl, ofurace, oxadixyl,
- 35 • amine derivates, such as aldimorph, dodemorph, fenpropidin, guazatine, iminoctadine, tridemorph,
- antibiotics, such as cycloheximid, griseofulvin, kasugamycin, natamycin, polyoxin or streptomycin,
- azoles, such as bitertanol, bromoconazole, cyproconazole, difenoconazole,
40 dinitroconazole, enilconazole, fenbuconazole, fluquinconazole, flusilazole, flutriafol,

AMENDED SHEET

hexaconazole, imazalil, ipconazole, myclobutanil, penconazole, propiconazole, prochloraz, prothioconazole, simeconazole, tetraconazole, triadimefon, triadimenol, triflumizol, triticonazole,

- dicarboximides, such as myclozolin, procymidone,
- 5 • dithiocarbamates, such as ferbam, nabam, metam, propineb, polycarbamat, ziram, zineb,
- heterocyclic compounds, such as anilazin, boscalid, carbendazim, carboxin, oxycarboxin, cyazofamid, dazomet, famoxadon, fenamidon, fuberidazole, flutolanil, furametpyr, isoprothiolane, mepronil, nuarimol, probenazole, pyroquilon, silthiofam,
- 10 thiabendazol, thifluzamid, tiadinil, tricyclazole, triforine,
- nitrophenyl derivatives, such as binapacryl, dinocap, dinobuton, nitrophthal-isopropyl,
- phenylpyrroles, such as fencpiclonil or fludioxonil,
- sulfur,
- 15 • other fungicides, such as acibenzolar-S-methyl, carpropamid, chlorothalonil, cyflufenamid, cymoxanil, diclomezin, diclocymet, diethofencarb, edifenphos, ethaboxam, fenhexamid, fentin-acetate, fenoxanil, ferimzone, fluazinam, fosetyl, hexachlorobenzene, metrafenon, pencycuron, propamocarb, phthalide, toloclofos-methyl, quintozene, zoxamide,
- 20 • strobilurins, such as fluoxastrobin, metominostrobin, oryastrobin, pyraclostrobin or trifloxystrobin,
- sulfenic acid derivatives, such as captafol,
- cinnamides and analogous compounds, such as flumetover.

25 In one embodiment of the mixtures according to the invention, a further fungicide III or two fungicides III and IV are added to the compounds I and II. Preference is given to mixtures comprising the compounds I and II and a component III. Particular preference is given to mixtures of the compounds I and II.

30 The compound I and the compound II can be applied simultaneously, that is jointly or separately, or in succession, the sequence, in the case of separate application, generally not having any effect on the result of the control measures.

The compound I and the compound II are usually applied in a weight ratio of from
35 100:1 to 1:100, preferably from 10:1 to 1:50, in particular from 5:1 to 1:10.

The compounds III and, where appropriate IV, are, if desired, mixed with the compound I in a ratio of from 20:1 to 1:20.

40 Depending on the type of compound and the desired effect, the application rates of the

mixtures according to the invention are from 5 g/ha to 2 000 g/ha, preferably from 50 to 1 500 g/ha, in particular from 50 to 750 g/ha.

Correspondingly, the application rates for the compound I are generally from 1 to 1 000 g/ha, preferably from 10 to 750 g/ha, in particular from 20 to 500 g/ha.

Correspondingly, the application rates for the compound II are generally from 5 to 2 000 g/ha, preferably from 10 to 1 000 g/ha, in particular from 50 to 750 g/ha.

In the treatment of seed, application rates of mixture are generally from 1 to 1000 g/-100 kg of seed, preferably from 1 to 750 g/100 kg, in particular from 5 to 500 g/100 kg.

In the control of phytopalliogenic harmful fungi, the separate or joint application of the compound I and the compound II or of the mixtures of the compound I and the compound II is carried out by spraying or dusting the seeds, the plants or the soil before or after sowing of the plants or before or after emergence of the plants.

The mixtures according to the invention, or the compounds I and II, can be converted into the customary formulations, for example solutions, emulsions, suspensions, dusts, powders, pastes and granules. The use form depends on the particular intended purpose; in each case, it should ensure a fine and even distribution of the compound according to the invention.

The formulations are prepared in a known manner, for example by extending the active compound with solvents and/or carriers, if desired using emulsifiers and dispersants. Solvents/auxiliaries suitable for this purpose are essentially:

- water, aromatic solvents (for example Solvesso products, xylene), paraffins (for example mineral oil fractions), alcohols (for example methanol, butanol, pentanol, benzyl alcohol), ketones (for example cyclohexanone, gamma-butyrolactone), pyrrolidones (NMP, NOP), acetates (glycol diacetate), glycols, fatty acid dimethylamides, fatty acids and fatty acid esters. In principle, solvent mixtures may also be used,
- carriers such as ground natural minerals (for example kaolins, clays, talc, chalk) and ground synthetic minerals (for example highly disperse silica, silicates);
- emulsifiers such as nonionic and anionic emulsifiers (for example polyoxyethylene fatty alcohol ethers, alkylsulfonates and arylsulfonates) and dispersants such as lignin-sulfite waste liquors and methylcellulose.

Suitable surfactants used are alkali metal, alkaline earth metal and ammonium salts of lignosulfonic acid, naphthalenesulfonic acid, phenolsulfonic acid,

dibutyl naphthalenesulfonic acid, alkylarylsulfonates, alkyl sulfates, alkylsulfonates, fatty alcohol sulfates, fatty acids and sulfated fatty alcohol glycol ethers, furthermore condensates of sulfonated naphthalene and naphthalene derivatives with formaldehyde, condensates of naphthalene or of naphthalenesulfonic acid with phenol and formaldehyde, polyoxyethylene octylphenyl ether, ethoxylated isooctylphenol, 5 octylphenol, nonylphenol, alkylphenyl polyglycol ethers, tributylphenyl polyglycol ether, tristearylphenyl polyglycol ether, alkylaryl polyether alcohols, alcohol and fatty alcohol ethylene oxide condensates, ethoxylated castor oil, polyoxyethylene alkyl ethers, ethoxylated polyoxypropylene, lauryl alcohol polyglycol ether acetal, sorbitol esters, 10 lignosulfite waste liquors and methylcellulose.

Substances which are suitable for the preparation of directly sprayable solutions, emulsions, pastes or oil dispersions are mineral oil fractions of medium to high boiling point, such as kerosene or diesel oil, furthermore coal tar oils and oils of vegetable or 15 animal origin, aliphatic, cyclic and aromatic hydrocarbons, for example toluene, xylene, paraffin, tetrahydronaphthalene, alkylated naphthalenes or their derivatives, methanol, ethanol, propanol, butanol, cyclohexanol, cyclohexanone, isophorone, highly polar solvents, for example dimethyl sulfoxide, N-methylpyrrolidone or water.

20 Powders, materials for spreading and dustable products can be prepared by mixing or concomitantly grinding the active substances with a solid carrier.

Granules, for example coated granules, impregnated granules and homogeneous granules, can be prepared by binding the active compounds to solid carriers. Examples 25 of solid carriers are mineral earths such as silica gels, silicates, talc, kaolin, attaclay, limestone, lime, chalk, bole, loess, clay, dolomite, diatomaceous earth, calcium sulfate, magnesium sulfate, magnesium oxide, ground synthetic materials, fertilizers, such as, for example, ammonium sulfate, ammonium phosphate, ammonium nitrate, ureas, and products of vegetable origin, such as cereal meal, tree bark meal, wood meal and 30 nutshell meal, cellulose powders and other solid carriers.

In general, the formulations comprise from 0.01 to 95% by weight, preferably from 0.1 to 90% by weight, of the active compounds. In this case, the active compounds are employed in a purity of from 90% to 100%, preferably 95% to 100% (according to NMR 35 spectrum).

The following are examples of formulations: 1. Products for dilution with water

A) Water-soluble concentrates (SL)
40 10 parts by weight of the active compounds are dissolved in water or in a water-soluble

solvent. As an alternative, wetters or other auxiliaries are added. The active compound dissolves upon dilution with water.

B) Dispersible concentrates (DC)

- 5 20 parts by weight of the active compounds are dissolved in cyclohexanone with addition of a dispersant, for example polyvinylpyrrolidone. Dilution with water gives a dispersion.

C) Emulsifiable concentrates (EC)

- 10 15 parts by weight of the active compounds are dissolved in xylene with addition of calcium dodecylbenzenesulfonate and castor oil ethoxylate (in each case 5% strength). Dilution with water gives an emulsion.

D) Emulsions (EW, EO)

- 15 40 parts by weight of the active compounds are dissolved in xylene with addition of calcium dodecylbenzenesulfonate and castor oil ethoxylate (in each case 5% strength). This mixture is introduced into water by means of an emulsifier machine (Ultraturrax) and made into a homogeneous emulsion. Dilution with water gives an emulsion.

20 E) Suspensions (SC, OD)

In an agitated ball mill, 20 parts by weight of the active compounds are comminuted with addition of dispersants, wetters and water or an organic solvent to give a fine active compound suspension. Dilution with water gives a stable suspension of the active compound.

25

F) Water-dispersible granules and water-soluble granules (WG, SG)

50 parts by weight of the active compounds are ground finely with addition of dispersants and wetters and made as water-dispersible or water-soluble granules by means of technical appliances (for example extrusion, spray tower, fluidized bed).

- 30 Dilution with water gives a stable dispersion or solution of the active compound.

G) Water-dispersible powders and water-soluble powders (WP, SP)

75 parts by weight of the active compounds are ground in a rotor-stator mill with addition of dispersants, wetters and silica gel. Dilution with water gives a stable

- 35 dispersion or solution of the active compound.

2. Products to be applied undiluted

H) Dustable powders (DP)

- 40 5 parts by weight of the active compounds are ground finely and mixed intimately with

95% of finely divided kaolin. This gives a dustable product. = 2006/03326

I) Granules (GR, FG, GG, MG)

0.5 part by weight of the active compounds is ground finely and associated with 95.5%
5 carriers. Current methods are extrusion, spray-drying or the fluidized bed. This gives granules to be applied undiluted.

J) ULV solutions (UL)

10 parts by weight of the active compounds are dissolved in an organic solvent, for
10 example xylene. This gives a product to be applied undiluted.

The active compounds can be used as such, in the form of their formulations or the use
forms prepared therefrom, for example in the form of directly sprayable solutions,
15 powders, suspensions or dispersions, emulsions, oil dispersions, pastes, dustable products, materials for spreading, or granules, by means of spraying, atomizing, dusting, spreading or pouring. The use forms depend entirely on the intended purposes; they are intended to ensure in each case the finest possible distribution of the active compounds according to the invention.

20 Aqueous use forms can be prepared from emulsion concentrates, pastes or wettable powders (sprayable powders, oil dispersions) by adding water. To prepare emulsions, pastes or oil dispersions, the substances, as such or dissolved in an oil or solvent, can be homogenized in water by means of a wetter, tackifier, dispersant or emulsifier.
25 However, it is also possible to prepare concentrates composed of active substance, wetter, tackifier, dispersant or emulsifier and, if appropriate, solvent or oil, and such concentrates are suitable for dilution with water.

The active compound concentrations in the ready-to-use preparations can be varied
30 within relatively wide ranges. In general, they are from 0.0001 to 10%, preferably from 0.01 to 1%.

The active compounds may also be used successfully in the ultra-low-volume process (ULV), it being possible to apply formulations comprising over 95% by weight of active
35 compound, or even to apply the active compound without additives.

Oils of various types, wetters, adjuvants, herbicides, fungicides, other pesticides, or
bactericides may be added to the active compounds, even, if appropriate, not until
immediately prior to use (tank mix). These agents can be admixed with the
40 compositions according to the invention, usually done in a weight ratio from 1:10 to 10:1.

The compounds I or II, the mixtures or the corresponding formulations are applied by treating the harmful fungi, the plants, seeds, soils, areas, materials or spaces to be kept free from them with a fungicidally effective amount of the mixture or, in the case of
5 separate application, of the compounds I and II. Application can be carried out before or after infection by the harmful fungi.

The fungicidal action of the compound and of the mixtures can be demonstrated by the following experiments:
10

The active compounds, separately or jointly, were prepared as a stock solution comprising 0.25% by weight of active compound in acetone or DMSO. 1% by weight of the emulsifier Uniperol® EL (wetting agent having emulsifying and dispersant action based on ethoxylated alkylphenols) was added to this solution, and the mixture was
15 diluted with water to the desired concentration.

Use example – activity against late blight of tomato caused by *Phytophthora infestans* during protective treatment

20 Leaves of potted tomato plants were sprayed to runoff point with an aqueous suspension having the concentration of active compound stated below. The next day, the leaves were infected with an aqueous sporangiospore suspension of *Phytophthora infestans*. The plants were then initially placed in a water-vapor saturated chamber at temperatures of between 18°C and 20°C. After 6 days late blight on the untreated, but
25 infected, control plants had developed to such an extent that the disease could be determined visually in %.

The visually determined percentages of infected leaf areas were converted into efficacies in % of the untreated control:
30

The efficacy (E) is calculated as follows using Abbot's formula:

$$E = (1 - \alpha/\beta) \cdot 100$$

35 α corresponds to the fungicidal infection of the treated plants in % and
 β corresponds to the fungicidal infection of the untreated (control) plants in %

An efficacy of 0 means that the infection level of the treated plants corresponds to that of the untreated control plants; an efficacy of 100 means that the treated plants were
40 not infected.

The expected efficacies of mixtures of active compounds were determined using Colby's formula (Colby, S.R. "Calculating synergistic and antagonistic responses of herbicide combinations", Weeds, 15, 20-22, 1967) and compared with the observed efficacies.

Colby's formula:

$$E = x + y - x \cdot y / 100$$

10


E expected efficacy, expressed in % of the untreated control, when using the mixture of the active compounds A and B at the concentrations a and b

x efficacy, expressed in % of the untreated control, when using the active compound A at the concentration a


15

y efficacy, expressed in % of the untreated control, when using the active compound B at the concentration b

20 Compounds A and B known from the tebuconazole mixtures described in EP-A 988 790 are used as comparative compounds:



A



B

Table A – Individual active compounds

Ex-ample	Active compound	Concentration of active compound in the spray liquor [ppm]	Efficacy in % of the untreated control
1	Control (untreated)	-	(88 % infection)
2	I	63	9
3	II (tebuconazole)	16 250	0 0
4	comparative A	63	0
5	comparative B	63	9

Table B – Mixtures according to the invention

Ex-ample	Mixture of active compounds Concentration Mixing ratio	Observed efficacy	Calculated efficacy*)
6	I + II 63 + 16 ppm 4:1	43	9
7	I + II 63 + 250 ppm 1:4	55	9

*) efficacy calculated using Colby's formula

Table C – Comparative tests – mixtures known from EP-A 988 780

Ex-ample	Mixture of active compounds Concentration Mixing ratio	Observed efficacy	Calculated efficacy*)
8	A + II 63 + 16 ppm 4:1	0	0
9	A + II 63 + 250 ppm 1:4	0	0
10	B + II 63 + 16 ppm 4:1	9	9
11	B + II 63 + 250 ppm 1:4	21	9

*) efficacy calculated using Colby's formula

- 5 The test results show that, by virtue of strong synergism, the mixtures according to the invention are considerably more effective against late blight than the tebuconazole mixtures known from EP-A 988 780.

10 Use example 2 - Activity against peronospora of grapevines caused by *Plasmopara viticola*

Leaves of potted grapevines of the cultivar "Riesling" were sprayed to runoff point with an aqueous suspension having the concentration of active compounds stated below. The next day, the undersides of the leaves were inoculated with an aqueous zoospore suspension of *Plasmopara viticola*. The grapevines were then initially placed in a water-vapor-saturated chamber at 24°C for 48 hours and then in a greenhouse at temperatures between 20 and 30°C for 5 days. After this period of time, the plants were, to promote sporangiophore eruption, again placed in a humid chamber for 16 hours. The extent of the development of the infection on the undersides of the leaves was then determined visually.

Evaluation was carried out analogously to example 1.

Table D - Individual active compounds

Ex-ample	Active compound	Concentration of active compound in the spray liquor [ppm]	Efficacy in % of the untreated control
12	-	control (untreated)	(84 % infection)
13	I	4	52
		1	4
14	II (tebuconazole)	4	28
		1	0

Table E - Mixtures according to the invention

Ex-ample	Mixture of active compounds Concentration Mixing ratio	Observed efficacy	Calculated efficacy*)
15	I+II 4+1 ppm 4:1	76	52
	I+II 1+4 ppm 1:4		
16	I+II 1+4 ppm 1:4	76	31

*) Efficacy calculated using Colby's formula

5

The test results show that for all mixing ratios the observed efficacy of the mixtures according to the invention is considerably higher than that predicted using Colby's formula.


- 10 "Comprises/comprising" when used in this specification is taken to specify the presence of stated features, integers, steps or components but does not preclude the presence or addition of one or more other features, integers, steps or components or groups thereof.

AMENDED SHEET

We claim:

1. A fungicidal mixture comprising, as active components,


5 1) the triazolopyrimidine derivative of the formula I



I

and

2) tebuconazole of the formula II,



II

10

in a synergistically effective amount.

2. A fungicidal mixture comprising the compound of the formula I and the compound of the formula II in a weight ratio of from 100:1 to 1:100.

15

3. A fungicidal composition comprising a liquid or solid carrier and a mixture as claimed in claim 1 or 2.

4. A method for controlling harmful fungi from the class of the *Oomycetes*, which comprises treating the fungi, their habitat or the seed, the soil or the plants to be protected against fungal attack with an effective amount of the compound I and the compound II as set forth in claim 1.

20

5. A method according to claim 4, wherein the compounds I and II as set forth in claim 1 are applied simultaneously, that is jointly or separately, or in succession.

25

6. A method according to claim 4, wherein the mixture as claimed in claim 1 or 2 is applied to the soil or the plants to be protected against fungal attack in an amount of from 5 g/ha to 2 000 g/ha.


7. A method according to claim 4 or 5, wherein the mixture as claimed in claim 1 or 2 is applied in an amount of from 1 to 1000 g/100 kg of seed.
8. A method according to any one of claims 4 to 7, wherein the harmful fungus *Phytophthora infestans* is controlled.
9. Seed comprising the mixture as claimed in claim 1 or 2 in an amount of from 1 to 1000 g/100 kg.
10. The use of the compound I and the compound II as set forth in claim 1 for preparing a composition suitable for controlling *Oomycetes*.
11. A fungicidal mixture as claimed in claim 1 or 2, substantially as hereinbefore described or exemplified.
12. A fungicidal mixture including any new and inventive integer or combination of integers, substantially as herein described.
13. A fungicidal composition as claimed in claim 3, substantially as hereinbefore described or exemplified.
14. A fungicidal composition including any new and inventive integer or combination of integers, substantially as herein described.
15. A method according to the invention for controlling harmful fungi, substantially as hereinbefore described or exemplified.
16. A method for controlling harmful fungi including any new and inventive integer or combination of integers, substantially as herein described.
17. Seed as claimed in claim 9, substantially as hereinbefore described or exemplified.
18. Seed according to the invention including any new and inventive integer or combination of integers, substantially as herein described.
19. The use as claimed in claim 10, substantially as hereinbefore described or exemplified.

Fungicidal mixtures

Abstract

5 Fungicidal mixtures comprising, as active components,

1) the triazolopyrimidine derivative of the formula I,




I

and

10

2) tebuconazole of the formula II,



II

15 in a synergistically effective amount, methods for controlling harmful fungi from the class of the *Oomycetes* using mixtures of the compound I with the compound II, the use of the compound I with the compound II for preparing such mixtures and compositions comprising these mixtures are described.