



- (51) **International Patent Classification:**
C07K 14/415 (2006.01) A01H 5/12 (2006.01)
C12N 15/82 (2006.01)
- (21) **International Application Number:** PCT/JP2011/007337
- (22) **International Filing Date:** 28 December 2011 (28.12.2011)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:** 2010-293783 28 December 2010 (28.12.2010) JP
- (71) **Applicant (for all designated States except US):** TOYOTA JIDOSHA KABUSHIKI KAISHA [JP/JP]; 1, Toyota-cho, Toyota-shi, Aichi, 4718571 (JP).
- (72) **Inventors; and**
- (75) **Inventors/Applicants (for US only):** HATTORI, Etsuko [JP/JP]; c/o Toyota Jidosha Kabushiki Kaisha, 1, Toyota-cho, Toyota-shi, Aichi, 4718571 (JP). NISHIMURA, Satoru [JP/JP]; c/o Toyota Jidosha Kabushiki Kaisha, 1, Toyota-cho, Toyota-shi, Aichi, 4718571 (JP). ITO, Kazuyo [JP/JP]; c/o Toyota Jidosha Kabushiki Kaisha, 1, Toyota-cho, Toyota-shi, Aichi, 4718571 (JP).
- (74) **Agents:** HIRAKI, Yusuke et al.; Atago Green Hills MORI Tower 32F, 5-1, Atago 2-chome, Minato-ku, Tokyo, 1056232 (JP).

- (81) **Designated States (unless otherwise indicated, for every kind of national protection available):** AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- (84) **Designated States (unless otherwise indicated, for every kind of regional protection available):** ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- with sequence listing part of description (Rule 5.2(a))



WO 2012/090499 A1

(54) **Title:** MATURE LEAF - SPECIFIC PROMOTER

(57) **Abstract:** The present invention provides gene expression regulatory DNA having activity of promoting gene expression in a mature-leaf-specific manner. The following is provided: gene expression regulatory DNA having activity of promoting gene expression in a mature-leaf-specific manner, which comprises any one of the following (a) to (d): (a) DNA consisting of the nucleotide sequence shown in SEQ ID NO: 1 or SEQ ID NO: 4; (b) DNA consisting of a nucleotide sequence that has a deletion, a substitution, an addition, or an insertion of one or a plurality of nucleotides in the nucleotide sequence shown in SEQ ID NO: 1 or SEQ ID NO: 4 and having activity of promoting gene expression in a mature-leaf-specific manner; (c) DNA consisting of a nucleotide sequence having 90% or more sequence identity to the nucleotide sequence shown in SEQ ID NO: 1 or SEQ ID NO: 4 and having activity of promoting gene expression in a mature-leaf-specific manner; and (d) DNA hybridizing under stringent conditions to DNA consisting of a sequence complementary to a part or the entirety of the nucleotide sequence shown in SEQ ID NO: 1 or SEQ ID NO: 4 and having activity of promoting gene expression in a mature-leaf-specific manner.

Description

Title of Invention: MATURE LEAF - SPECIFIC PROMOTER

Technical Field

[0001] The present invention relates to gene expression regulatory DNA having activity of promoting gene expression in a mature-leaf-specific manner and the use thereof.

Background Art

[0002] Genetic information of an organism is transmitted through a series of processes referred to as "central dogma" in which the DNA information of a functional gene is transcribed into mRNA and a functional protein is synthesized via translation of mRNA information. As a result, a biofunction is expressed. Genomic DNA is an aggregate of genetic information formed via association of parent haploids. Also for plants, genomic DNA generally specifies the nature of cells containing the DNA and a plant derived from such cells.

[0003] For the purpose of inducing expression of the biofunction through transmission of functional genetic information of genomic DNA contained in cells in a correct manner, it is necessary to induce expression of an adequate gene with adequate timing at an adequate site and with an adequate intensity. Therefore, strict regulation of the expression of specific genes is required.

For the expression of a functional gene, the time, site, and intensity of expression are regulated by the gene expression regulatory DNA which presents in the 5' upstream region of the gene.

[0004] In recent years, as a result of the deciphering of genomic DNA using *Arabidopsis thaliana*, rice, and the like, it has become possible to readily obtain gene expression regulatory DNAs of individual functional genes of such plants (Non-Patent Documents 1 and 2).

[0005] Meanwhile, genomic DNA of sugarcane, which is a readily available crop, has not been deciphered yet. Therefore, it is not easy to obtain gene expression regulatory DNAs of individual functional genes of sugarcane. Hitherto, for gene introduction into sugarcane, gene expression regulatory DNA from a non-sugarcane plant has been used to regulate expression of a transgene (Non-Patent Documents 3 and 4).

[0006] However, when functional gene expression is induced in sugarcane using gene expression regulatory DNA of a non-sugarcane plant, it is observed in some cases that the time, site, intensity, and other conditions of expression cannot be strictly regulated. Therefore, acquisition of sugarcane-derived gene expression regulatory DNA, and particularly, tissue-specific gene expression regulatory DNA, has been awaited in the

related fields.

Citation List

Non Patent Literature

- [0007] NPL 1: Nature, 2000, 408:769-815
 NPL 2: Science, 2002, 296:92-100
 NPL 3: Plant Mol Biol. 1992, Feb; 18(4):675-89
 NPL 4: Planta, 1998, 206:20-27

Summary of Invention

Technical Problem

- [0008] Therefore, an object of the present invention is to provide gene expression regulatory DNA having activity of promoting gene expression in a mature-leaf-specific manner.

Solution to Problem

- [0009] As a result of intensive studies to achieve the above object, the present inventors found gene expression regulatory DNA having activity of promoting gene expression in a mature-leaf-specific manner in the 5' upstream region of a gene expressed in mature leaves of sugarcane. This has led to the completion of the present invention.

Specifically, the present invention encompasses the following features {1} to {4}.

- [0010] {1} A gene expression regulatory DNA having activity of promoting gene expression in a mature-leaf-specific manner, which comprises any one of the following (a) to (d):

(a) DNA consisting of the nucleotide sequence shown in SEQ ID NO: 1 or SEQ ID NO: 4;

(b) DNA consisting of a nucleotide sequence that has a deletion, a substitution, an addition, or an insertion of one or a plurality of nucleotides in the nucleotide sequence shown in SEQ ID NO: 1 or SEQ ID NO: 4 and having activity of promoting gene expression in a mature-leaf-specific manner;

(c) DNA consisting of a nucleotide sequence having 90% or more sequence identity to the nucleotide sequence shown in SEQ ID NO: 1 or SEQ ID NO: 4 and having activity of promoting gene expression in a mature-leaf-specific manner; and

(d) DNA hybridizing under stringent conditions to DNA consisting of a sequence complementary to a part or the entirety of the nucleotide sequence shown in SEQ ID NO: 1 or SEQ ID NO: 4 and having activity of promoting gene expression in a mature-leaf-specific manner.

- [0011] {2} A recombinant vector, which comprises the gene expression regulatory DNA according to {1}.

{3} A transformed plant obtained by transformation using the recombinant vector according to {2}.

{4} A transformed plant obtained from plant cells transformed by plant cell trans-

formation using the recombinant vector according to {2}.

A part or all of the content disclosed in the description and/or drawings of Japanese Patent Application No. 2010-293783, which is a priority document of the present application, is herein incorporated by reference.

Advantageous Effects of Invention

[0012] According to the present invention, the following can be provided: gene expression regulatory DNA having activity of promoting gene expression in a mature-leaf-specific manner; a recombinant vector comprising the DNA, which can cause a functional gene to be expressed in a mature-leaf-specific manner; and a transformed plant transfected with the recombinant vector.

Brief Description of Drawings

[0013] [fig.1]Fig. 1 shows the nucleotide sequence of Saccharum officinarum-derived ecc0001 EST (searched for using the DFCI Sugarcane Gene Index).

[fig.2]Fig. 2 shows results of ecc0001 EST expression level analysis for individual tissues of Saccharum spp. hybrids cv. NiF8. The strongest expression level confirmed for stalk epidermis is designated as 100.

[fig.3]Fig. 3 shows the nucleotide sequence of the gene expression regulatory region of the ecc0001 gene (SEQ ID NO: 1) into which an HindIII restriction enzyme recognition sequence and a BlnI restriction enzyme recognition sequence were inserted at the 5'- and 3'-ends, respectively.

[fig.4-1]Fig. 4-1 shows a predicted promoter region in the gene expression regulatory region of the ecc0001 gene (SEQ ID NO: 4) identified using a promoter analysis tool.

[fig.4-2]Fig. 4-2 (continued from fig. 4-1)

[fig.5]Fig. 5 schematically shows a gene expression vector containing the gene expression regulatory region of the ecc0001 gene and the beta-glucuronidase gene ligated to each other.

[fig.6]Fig. 6 shows results of beta-glucuronidase gene expression level analysis for individual tissues of a transgenic sugarcane in which the beta-glucuronidase gene expression was regulated by expression regulatory DNA of the ecc0001 gene. The strongest expression level confirmed for mature leaves of the transgenic sugarcane is designated as 100.

Description of Embodiments

[0014] The present invention is described in detail below.

First, the gene expression regulatory DNA having activity of promoting gene expression in a mature-leaf-specific manner of the present invention is described. The gene expression regulatory DNA of the present invention comprises any one of the following (a) to (d):

- (a) DNA consisting of the nucleotide sequence shown in SEQ ID NO: 1 or SEQ ID NO: 4;
- (b) DNA consisting of a nucleotide sequence that has a deletion, a substitution, an addition, or an insertion of one or a plurality of nucleotides in the nucleotide sequence shown in SEQ ID NO: 1 or SEQ ID NO: 4 and having activity of promoting gene expression in a mature-leaf-specific manner;
- (c) DNA consisting of a nucleotide sequence having 90% or more sequence identity to the nucleotide sequence shown in SEQ ID NO: 1 or SEQ ID NO: 4 and having activity of promoting gene expression in a mature-leaf-specific manner; and
- (d) DNA hybridizing under stringent conditions to DNA consisting of a sequence complementary to a part or the entirety of the nucleotide sequence shown in SEQ ID NO: 1 or SEQ ID NO: 4 and having activity of promoting gene expression in a mature-leaf-specific manner.

[0015] The gene expression regulatory DNA of the present invention can be obtained in the following manner: candidate genes expressed in mature leaves are obtained by gene expression analysis using total RNAs from individual sugarcane tissues (of stalks, mature leaves, young leaves, and the like) or cDNAs from such RNAs; expression characteristics of the candidate genes are evaluated; genes evaluated as being expressed in a mature-leaf specific manner are specified based on the evaluation results; and the nucleotide sequence of the 5' upstream region of each candidate gene is identified based on cDNA or genomic DNA of the relevant specified gene. Here, gene expression analysis can be carried out using exhaustive gene expression analysis techniques generally known to persons skilled in the art such as a DNA chip method and a differential display method.

[0016] Specifically, the nucleotide sequence of SEQ ID NO: 1 or SEQ ID NO: 4 exists in the 5' upstream region of the gene (hereinafter referred to as "ecc0001") which is expressed in mature sugarcane leaves. Examples of "sugarcane" plants described herein include (but are not particularly limited to) plants belonging to the genus *Saccharum* such as *Saccharum officinarum*, *Saccharum sinense*, *Saccharum barberi*, *Saccharum robustum*, *Saccharum spontaneum*, *Saccharum edule*, and *Saccharum* spp. hybrids cv. NiF8; and plants belonging to a genus/species closely related to the genus *Saccharum*/*Saccharum* species, such as, *Sorghum* or *Erianthus*. Of these, *Saccharum* spp. hybrids cv. NiF8 is preferable.

[0017] A method for isolating DNA present in the 5' upstream region is not particularly limited. DNA isolation can be conducted by a method generally known to persons skilled in the art. For instance, DNA can be isolated by a conventionally known method comprising cloning an unknown region (i.e., the 5' upstream region in the above case) based on the nucleotide sequence (SEQ ID NO: 2) of the ecc0001 gene. In

such a method, genomic DNA containing the 5' upstream region of the ecc0001 gene is subjected to restriction enzyme treatment such that an adopter consisting of a predetermined nucleotide sequence is ligated to the DNA. Primers are designated for the nucleotide sequence of the ecc0001 gene and the adopter, followed by PCR. Accordingly, an unknown nucleotide sequence adjacent to the 5' upstream region of the nucleotide sequence of the ecc0001 gene can be amplified. After the amplified nucleotide sequence is determined, another pair of primers is designed based on the determined nucleotide sequence. Thus, another unknown nucleotide sequence adjacent to the determined nucleotide sequence can be amplified in a similar manner. This method can be carried out using a commercially available cloning kit such as a RightWalk (registered trademark) kit (BEX Co., Ltd.). Alternatively, a method using inverse PCR can be suggested in addition to the above. In such case, a pair of primers is designed based on the nucleotide sequence information of the ecc0001 gene. PCR is performed using the pair of primers and a genomic DNA fragment obtained via treatment with a certain restriction enzyme and self-ligation. Thus, the upstream region of the ecc0001 gene can be amplified. Further, another method for isolating the upstream region of the ecc0001 gene from a genomic DNA library can be suggested. In such case, a genomic DNA library that has been prepared by a standard method is screened with the use of cDNA comprising the ecc0001 gene as a probe to obtain genomic DNA comprising the ecc0001 gene. Then, the nucleotide sequence of genomic DNA obtained by screening is determined. Accordingly, the 5' upstream region present in the upstream region of the ecc0001 gene can be specified. Further, the 5' upstream region alone can be amplified by PCR or the like.

[0018] As described above, unknown nucleotide sequences located upstream of the ecc0001 gene are sequentially amplified or screened for to determine the nucleotide sequence by a standard method. Accordingly, the nucleotide sequence shown in SEQ ID NO: 1 or SEQ ID NO: 4 can be specified. Once the nucleotide sequence shown in SEQ ID NO: 1 or SEQ ID NO: 4 is determined, it becomes possible to obtain the nucleotide sequence shown in SEQ ID NO: 1 or SEQ ID NO: 4 by PCR using genomic DNA extracted from sugarcane as a template and primers designed based on the nucleotide sequence shown in SEQ ID NO: 1 or SEQ ID NO: 4.

[0019] The nucleotide sequence shown in SEQ ID NO: 1 or SEQ ID NO: 4 functions as a gene expression regulatory region capable of inducing gene expression in a mature-leaf-specific manner. A gene expression regulatory region contains nucleotide sequences involved in gene transcription control, such as, a promoter region, an enhancer region, a TATA box, and/or a CAT box (although the contents of the region are not particularly limited thereto).

[0020] The word "specific" used herein refers to the following conditions: a gene expression

inducible function is exclusively present in mature leaf tissue, which is one of individual tissues constituting a plant; and the gene expression inducible function in mature leaf tissue is remarkably or statistically significantly (e.g., about 2-fold, 3-fold, 4-fold, 5-fold, 6-fold, 7-fold, 8-fold, 9-fold, or 10-fold or more) greater than the gene expression inducible function in a non-mature-leaf tissue (e.g., a stalk, young leaf, root, or meristem tissue).

[0021] The term "mature leaf (tissue)" used herein refers to a leaf (tissue) that contains chloroplasts accumulating in cells for photosynthesis and thus is tinged with green. It also refers to a leaf (tissue) other than a young leaf (tissue) containing no chloroplasts for photosynthesis.

[0022] The gene expression inducible function can be confirmed by reporter assay or the like generally known to persons skilled in the art. Upon reporter assay, a vector is prepared in which various reporter genes (e.g., the beta-glucuronidase gene (GUS), the luciferase gene (LUC), and the green fluorescent protein gene (GFP)) are ligated to the downstream region of a nucleotide sequence to be examined in terms of the gene expression inducible function such that the reporter genes are regulated by the nucleotide sequence. Gene introduction (or transient gene introduction) into the genome of a host is carried out using the vector. Then, the expression level of each reporter gene is determined. Thus, the gene expression inducible function can be confirmed. The reporter gene is not particularly limited as long as the expression thereof is detectable. Examples of such reporter gene include reporter genes conventionally used by persons skilled in the art such as the CAT gene, the lacZ gene, the luciferase (hereafter denoted by "LUC") gene, the beta-glucuronidase (hereafter denoted by "GUS") gene, and the green fluorescent protein (hereafter denoted by "GFP") gene.

[0023] The expression level of a reporter gene can be determined by a method generally known to persons skilled in the art depending on the type of the reporter gene. For instance, if the reporter gene is the CAT gene, the expression level of the reporter gene can be determined by detecting acetylation of chloramphenicol with the gene product. The expression level of an individual reporter gene can be determined by the following technique. In a case in which the reporter gene is the lacZ gene, color development of a dye compound induced by the catalytic action of the gene expression product is detected. In a case in which the reporter gene is the LUC gene, fluorescence emission from a fluorescent compound induced by the catalytic action of the gene expression product is detected. In a case in which the reporter gene is the GFP gene, fluorescence emission from the GFP protein is detected. For instance, if the reporter gene is GUS, GUS activity is determined as promoter activity in a host cell by one of both of the following methods: (i) a method involving histochemical GUS staining (EMBO J. 6, 3901-3907 (1987)) and/or (ii) the method of Castle & Morris involving the use of a

fluorescent substrate (Plant Molecular Biology Manual, B5, 1-16 (1994); S. B. Gelvin & R. A. Schilperoort, Kluwer Academic Publishers). Further, the protein amount is determined by the method of Bradford (Anal. Biochem. 72, 248-254 (1976)). GUS activity is converted based on the protein amount into units of nmole 4-MU/min/mg protein. Thus, the gene expression inducible function can be confirmed for each case.

[0024] In addition, if a gene other than the above is used as a reporter gene, the gene transcription level is determined by Northern hybridization, RT-PCR, DNA array technology, or the like. Alternatively, the expression level of the protein encoded by the gene is determined by electrophoresis such as SDS-PAGE, Western blotting, or the like. Thus, the gene expression inducible function can be confirmed.

[0025] The gene expression regulatory DNA of the present invention is not limited to the nucleotide sequence shown in SEQ ID NO: 1 or SEQ ID NO: 4. As described in (b) above, it may be a nucleotide sequence that has a deletion, a substitution, an addition, or an insertion of one or a plurality of nucleotides in the nucleotide sequence shown in SEQ ID NO: 1 or SEQ ID NO: 4 as long as it has activity of promoting gene expression in a mature-leaf-specific manner.

[0026] For instance, even a nucleotide sequence that has a deletion, a substitution, an addition, or an insertion of 1 to 100 nucleotides, preferably 1 to 50 nucleotides, and more preferably 1 to 10 nucleotides in the nucleotide sequence shown in SEQ ID NO: 1 or SEQ ID NO: 4 is included in the gene expression regulatory DNA of the present invention as long as it shows activity of promoting gene expression in a mature-leaf-specific manner.

[0027] In addition, the gene expression regulatory DNA of the present invention is not limited to the nucleotide sequence shown in SEQ ID NO: 1 or SEQ ID NO: 4. As described in (c) above, it may be a nucleotide sequence having 80% or more, more preferably 90% or more, further preferably 95% or more, and most preferably 99% or more sequence identity to the nucleotide sequence shown in SEQ ID NO: 1 or SEQ ID NO: 4 as long as it exhibits activity of promoting gene expression in a mature-leaf-specific manner. Nucleotide sequences can be compared by a generally known method. Comparison can be performed using, for example, BLAST (Basic Local Alignment Search Tool of the National Center for Biological Information in the U.S.) based on default setting.

Further, the gene expression regulatory DNA of the present invention is not limited to the nucleotide sequence shown in SEQ ID NO: 1 or SEQ ID NO: 4. As described in (d) above, it may be a nucleotide sequence that hybridizes under stringent conditions to DNA consisting of a sequence complementary to a part or the entirety of the nucleotide sequence shown in SEQ ID NO: 1 or SEQ ID NO: 4 as long as it shows activity of promoting gene expression in a mature-leaf-specific manner.

[0028] Here, the term "stringent conditions" refers to conditions under which namely a specific hybrid is formed, but a non-specific hybrid is never formed. For example, such conditions comprise hybridization at 42 degrees C to 55 degrees C in a solution containing 2-6 x SSC (1 x SSC composition: 0.15M NaCl, 0.015M sodium citrate, pH 7.0) and 0.1 to 0.5% SDS, followed by washing at 55 degrees C to 65 degrees C in a solution containing 0.1 to 0.2 x SSC and 0.1 to 0.5% SDS.

[0029] Moreover, the gene expression regulatory DNA of the present invention having activity of promoting gene expression in a mature-leaf-specific manner may be a DNA fragment that has a deletion of 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 1100, 1200, 1300, 1400, or 1500 or more consecutive nucleotides from the 5'-end and/or 3'-end in the nucleotide sequence shown in SEQ ID NO: 1 or SEQ ID NO: 4 as long as it exhibits activity of promoting gene expression in a mature-leaf-specific manner. Nucleotides can be deleted by a method generally known to persons skilled in the art (e.g., PCR or restriction enzyme treatment). The DNA fragment may be a promoter region of the gene expression regulatory DNA of the present invention. A promoter region of a predetermined gene expression regulatory DNA can be searched for using a promoter analysis tool generally known to persons skilled in the art (e.g., Bioinformatics and Molecular Analysis Section (<http://www-bimas.cit.nih.gov/molbio/proscan/>); Prestridge, D. S. (1995), Predicting Pol II Promoter Sequences Using Transcription Factor Binding Sites, *J. Mol. Biol.* 249: 923-32). An example of such fragment of the nucleotide sequence shown in SEQ ID NO: 1 and SEQ ID NO: 4 is a DNA consisting of the 1412rd to 1662rd nucleotides of the sequence shown in SEQ ID NO: 1 and a DNA consisting of the 1413rd to 1663rd nucleotides of the sequence shown in SEQ ID NO: 4, respectively. It can be confirmed whether or not the obtained fragment has the gene expression inducible function by the above reporter assay or the like.

[0030] Once the nucleotide sequence of the gene expression regulatory DNA of the present invention is determined, it becomes possible to obtain the gene expression regulatory DNA of the present invention by chemical synthesis, PCR using genomic DNA as a template, or hybridization using a DNA fragment having the nucleotide sequence as a probe. Additionally, a nucleotide sequence that has a mutation in the nucleotide sequence shown in SEQ ID NO: 1 or SEQ ID NO: 4 can be synthesized by site-specific mutagenesis or the like. Mutation can be introduced into the nucleotide sequence shown in SEQ ID NO: 1 or SEQ ID NO: 4 by a known technique such as the Kunkel method or the Gapped duplex method or a method based thereof. For example, mutation is introduced with a mutagenesis kit using site-directed mutagenesis (e.g., Mutant-K or Mutant-G (both are trade names of TAKARA Bio)) or the like, or a LA PCR in vitro Mutagenesis series kit (trade name, TAKARA Bio).

- [0031] Next, a recombinant vector containing the above gene expression regulatory DNA having activity of promoting gene expression in a mature-leaf-specific manner is described.
- [0032] The recombinant vector of the present invention can be constructed by introducing DNA comprising a desired functional gene operably ligated to the above gene expression regulatory DNA into an appropriate vector. The term "operably ligated" used herein refers to conditions under which the above vector contains the gene expression regulatory DNA and the functional gene ligated to each other such that the functional gene is correctly expressed under the regulation of the gene expression regulatory DNA in a host cell transfected with the above vector. Here, "ligated" may be direct ligation or indirect ligation via a spacer having an appropriate length and an appropriate sequence. Preferable examples of a vector used in the present invention include pBI vectors, pBII vectors, pPZP vectors (Hajdukiewicz P, Svab Z, Maliga P.: The small, versatile pPZP family of Agrobacterium binary vectors for plant transformation., *Plant Mol Biol.*, 25: 989-94, 1994), pCAMBIA vectors (http://www.cambia.org/main/r_et_camvec.htm), and pSMA vectors by which a functional gene can be introduced into a plant using Agrobacterium. Particularly preferably, pBI and pBII binary vectors or intermediate vectors are used. Examples of such vectors include pBI121, pBI101, pBI101.2, pBI101.3, pBII221, and pIG121. A binary vector is a shuttle vector replicable in *Escherichia coli* and Agrobacterium. When a plant is infected with Agrobacterium containing a binary vector, DNA corresponding to a region between border sequences, the LB sequence and the RB sequence, present on the vector can be incorporated into nuclear DNA of the plant (*EMBO Journal*, 10(3), 697-704 (1991)). Meanwhile, a gene can be directly introduced into a plant using a pUC vector. Examples of a pUC vector include pUC18, pUC19, and pUC9. In addition, plant virus vectors such as cauliflower mosaic virus (CaMV), bean golden mosaic virus (BGMV), and tobacco mosaic virus (TMV) can be used.
- [0033] In order to facilitate ligation and/or insertion into a vector, DNA comprising the gene expression regulatory DNA and/or DNA comprising the functional gene can be adequately modified with the substitution, insertion, or addition of a restriction enzyme recognition sequence. For insertion into a vector, it is possible to use, for example, a method comprising cleaving purified DNA comprising the gene expression regulatory DNA and/or the functional gene with an adequate restriction enzyme and inserting each obtained fragment into the restriction enzyme recognition site or the multicloning site of adequate vector DNA for ligation to the vector.
- [0034] The term "functional gene" refers to an arbitrary endogenous gene of a target plant or an arbitrary exogenous gene that is expected to cause expression of a gene product in mature leaves. Examples of such gene include, but are not limited to, a photosynthesis-

related gene, a translocation-related gene, a gene capable of producing a useful substance (e.g., a drug, dye, or aromatic component), a sugar metabolism-related gene, a gene with disease/insect resistance (e.g., insect damage resistance, mycotic (fungal) and bacterial disease resistance, or viral (disease) resistance), an environmental stress (e.g., low temperature, high temperature, dry, photolesion, or ultraviolet)-resistance-related gene, and a plant growth regulation (promotion/suppression) gene.

[0035] If necessary, an enhancer, an intron, a poly-A addition signal, a 5'-UTR sequence, a selection marker gene, and the like can be ligated upstream/downstream of or between the gene expression regulatory DNA and/or functional gene in the vector.

An enhancer is used to, for example, improve efficiency of functional gene expression. An example thereof is an enhancer region containing a sequence located upstream of a CaMV35S promoter.

[0036] A terminator may be a sequence that can terminate transcription of a gene caused by the above promoter. Examples thereof include a nopaline synthetase gene terminator, an octopine synthetase gene terminator, and a CaMV 35S RNA gene terminator.

[0037] Examples of a selection marker gene include a hygromycin-resistant gene, a kanamycin-resistant gene, a bialaphos-resistant gene, a blasticidin S-resistant gene, and an acetolactate synthase gene. A selection marker gene may be ligated together with a functional gene to an identical plasmid as described above for preparation of a recombinant vector. Alternatively, a recombinant vector obtained by ligating a selection marker gene to a plasmid and a recombinant vector obtained by ligating a functional gene to a plasmid may be separately prepared. When they are separately prepared, a host is cotransfected with both vectors.

A transformant can be produced using the thus prepared recombinant vector.

[0038] When a transformed plant is prepared, various methods that have been reported and established can be adequately used. Preferable examples of such methods include an Agrobacterium method, a PEG-calcium phosphate method, an electroporation method, a liposome method, a particle gun method, and a microinjection method. For an Agrobacterium method, a protoplast or a tissue section is used or a plant itself is used (in an in planta method). When a protoplast is used, a method comprising coculture of Agrobacterium having a Ti plasmid with a protoplast or a method comprising fusion of an Agrobacterium spheroplast and a protoplast (i.e., a spheroplast method) is used. When a tissue section is used, a method comprising infecting an aseptically cultured leaf section (leaf disc) of a target plant with Agrobacterium, a method comprising infecting calluses, or the like with Agrobacterium can be used. In addition, if an in planta method using a seed or plant (a system for which tissue culture with the addition of a plant hormone is unnecessary) is employed, water-absorbing seeds, a young plant

(young seedling), a potted plant, etc. is directly treated with *Agrobacterium*.

[0039] It is possible to confirm the occurrence or nonoccurrence of incorporation of the gene into a plant by a PCR method, a Southern hybridization method, a Northern hybridization method, a Western blotting method, or the like. For example, DNA is prepared from a transformed plant. DNA-specific primers are designed, followed by PCR. After PCR, an amplification product is subjected to agarose gel electrophoresis, polyacrylamide gel electrophoresis, capillary electrophoresis, or the like, followed by staining with ethidium bromide, SYBR Green liquid, or the like. Then, a single band derived from an amplification product is detected. Thus, the occurrence of transformation can be confirmed. In addition, an amplification product can be detected by PCR using primers preliminarily labeled with a fluorescence dye or the like. Further, it is also possible to carry out a method comprising allowing an amplification product to bind to a solid phase such as a microplate and confirming an amplification product via, for example, a fluorescence or enzyme reaction.

[0040] Examples of a plant used for transformation in the present invention include, but are not particularly limited to, plants belonging to, for example, the families Gramineae, Solanaceae, Brassicaceae, Leguminosae, Rosaceae, Asteraceae, Liliaceae, Apiaceae, Caryophyllaceae, Cucurbitaceae, Convolvulaceae, and Chenopodiaceae. Preferable examples thereof include the plants belonging to the family Gramineae from which the above gene expression regulatory DNA has been isolated, including sugarcane, rice, barley, wheat, maize, zoysia, Sorghum, millet, Japanese millet, napier grass, and switchgrass.

[0041] Examples of plant materials subjected to transformation in the present invention include plant tissue of root, stalk, leaf, seed, embryo, ovule, ovary, shoot apex (growth point at the tip of a plant bud), anther, pollen, or the like, a section of such plant tissue, undifferentiated callus, and cultured plant cells such as protoplasts obtained by subjecting the above examples to enzyme treatment for cell wall removal. In addition, when an in planta method is employed, water-absorbing seeds and a plant as a whole can be used.

[0042] In addition, the transformed plant in the present invention may be a plant as a whole, a plant organ (e.g., a root, stalk, leaf, petal, seed, or fruit), a plant tissue (e.g., tissue of epidermis, phloem, parenchyma, xylem, or vascular bundle), or a plant culture cell.

[0043] When plant culture cells are used, a plant organ or plant itself can be regenerated by a known tissue culture method for regeneration of a transformed plant from the obtained transformed cells. Such regeneration operation can be readily carried out by persons skilled in the art as long as a conventionally known method is used as a method for regeneration of a plant from plant cells. A plant can be regenerated from plant cells as described below.

- [0044] First, when a plant tissue or protoplast is used as a target plant material for transformation, it is cultured in a sterilized callus formation medium supplemented with inorganic element(s), vitamin(s), carbon source(s), sugar(s) used as energy source(s), plant growth regulator(s) (e.g., a plant hormone such as auxin or cytokinin), and the like for formation of a dedifferentiated callus capable of growing adventitiously (hereinafter referred to as "callus induction"). The thus formed callus is transferred to fresh medium containing a plant growth regulator such as auxin for further growth (subculture).
- [0045] Callus induction is carried out on a solid medium such as agar. Subculture is carried out via, for example, liquid culture. In such case, each culturing can be carried out efficiently and in large scale. Next, the callus grown by subculture described above is cultured under adequate conditions for induction of organ redifferentiation (hereinafter referred to as "redifferentiation induction"). This eventually results in the regeneration of a complete plant. Redifferentiation induction can be performed by determining the types and amounts of a plant growth regulator such as auxin or cytokinin and different components such as a carbon source to be added to a medium and setting light, temperature, and other conditions in an adequate manner. Such redifferentiation induction results in formation of adventitious embryo, root, bud, stalk leaf, or the like, followed by further cultivation for acquisition of a complete plant. Alternatively, for example, it is possible to preserve a redifferentiated product in a state that would allow it to become a complete plant (e.g., an encapsulated artificial seed, dry embryo, or lyophilization cell or tissue).
- [0046] According to the present invention, the term "transformed plant" refers to not only a "T1 generation plant" obtained as a first generation plant via redifferentiation after transformation but also a "T2 generation plant" obtained as a plant of the subsequent generation from seeds of the T1 generation plant and a progeny plant such as the next generation (T3 generation) plant obtained via self-pollination of a flower of a "T2 generation" plant that has been found to be a transgenic plant via drug selection, Southern analysis, or the like.
- [0047] The thus produced transformed plant expresses a functional gene introduced in a mature-leaf-specific manner.

The gene expression regulatory DNA of the present invention has activity of promoting gene expression in a mature-leaf-specific manner in plants, and particularly, sugarcane. A desired functional gene is ligated to the gene expression regulatory DNA of the present invention and the ligation product is introduced into a plant, thereby allowing the functional gene to be expressed in a mature-leaf-specific manner in the plant. Thus, desired properties are imparted to the plant. The gene expression regulatory DNA of the present invention can be appropriately used for sugarcane for

which not many gene expression regulatory systems have been developed. The use of the gene expression regulatory DNA of the present invention enables production of transgenic sugarcane having properties appropriate for biomass resources and other applications.

Examples

[0048] The present invention is hereafter described in greater detail with reference to the following examples, although the technical scope of the present invention is not limited thereto.

Example 1

[0049] Cloning of mature leaf expression gene

Total RNA was extracted and purified from mature leaf, young leaf, and stalk tissues of sugarcane (*Saccharum* spp. cv. NiF8) using RNeasy Plant Mini Kits (QIAGEN). A cDNA library was constructed according to a standard method and used for gene expression analysis. Gene expression analysis was carried out using a Sugarcane Genome Array (Affimetrix) according to the manufacturer's instructions.

[0050] As a result of gene expression analysis, a gene expressed in mature leaves, young leaves, stalk pith, stalk epidermis, roots, and meristem of *Saccharum* spp. cv. NiF8 was found. The gene was designated as "ecc0001." The nucleotide sequence of the *Saccharum* spp. cv. NiF8-derived ecc0001 gene is identical to the nucleotide sequence of *Saccharum officinarum*-derived ecc0001 EST shown in fig. 1. Total RNA was extracted and purified from mature leaves, young leaves, stalk pith, stalk leaves, roots, and meristem of *Saccharum* spp. cv. NiF8. cDNA was prepared according to a standard method. The ecc0001 gene expression level in each tissue was analyzed by a SYBRGreen method using an ABI7500 real-time PCR system (Applied Biosystems).

[0051] Fig. 2 shows the results. The results shown in fig. 2 are expressed in relative values with reference to the strongest gene expression level in stalk epidermis, which is designated as 100 (%). The expression levels in mature leaves, young leaves, stalk pith, stalk epidermis, roots, and meristem were found to be 12, 13, 90, 100, 17, and 16, respectively. The results revealed that the ecc0001 gene is expressed in mature leaves, young leaves, stalk pith, stalk epidermis, roots, and meristem of *Saccharum* spp. cv. NiF8.

Example 2

[0052] Acquisition of mature-leaf-specific promoter

Genomic DNA (approximately 300 ng) was extracted and purified from mature leaf tissue (0.5 g) of sugarcane (*Saccharum* spp. cv. NiF8) using DNeasy Plant Mini Kits (QIAGEN). The gene expression regulatory region located 5'-upstream of the ecc0001 gene was obtained from the above genomic DNA based on the nucleotide sequence of

the ecc0001 gene obtained in Example 1 using RightWallk (registered trademark) Kits (BEX). An HindIII restriction enzyme recognition sequence (AAGCTT) and a BlnI restriction enzyme recognition sequence (CCTAGG) were used as linker sequences. The former was introduced at the 5'-end of the obtained gene expression regulatory region and the latter was introduced on the 3' side of the translation initiation site (ATG) of the ecc0001 gene presented at the 3'-end of the gene expression regulatory region. Thus, DNA comprising the expression regulatory region of the ecc0001 gene was prepared (fig. 3) (SEQ ID NO: 3).

[0053] The above DNA comprising the expression regulatory region of the ecc0001 gene was analyzed using a generally known promoter analysis tool (BioInformatics and Molecular Analysis Section) (<http://www-bimas.cit.nih.gov/molbio/proscan/>). As a result, a region capable of functioning as a promoter was presumed to exist in the region comprising the 1419th to 1669th nucleotides (figs. 4-1 and 4-2; SEQ ID NO: 5). The region corresponds to the region comprising the 1418th to 1668th nucleotides in the nucleotide sequence shown in SEQ ID NO: 3.

Example 3

[0054] Construction of gene expression vector

(1) Beta-glucuronidase gene expression vector

A gene expression vector comprising the DNA comprising the gene expression regulatory region obtained in Example 2 (SEQ ID NO:5) ligated to UidAcDNA comprising the beta-glucuronidase (GUS) gene was constructed. A plant transformation vector (pBII221) was used for the gene expression vector. For ligation of the DNA comprising a gene expression regulatory region and UidAcDNA, the ATG sequence encoding the first methionine on the 5'-end side of UidAcDNA and the ATG sequence encoding the first methionine of the ecc0001 gene present on the 3'-end side of the DNA comprising a gene expression regulatory region were matched for ligation (translational fusion type). Fig. 5 schematically shows the gene expression vector.

Example 4

[0055] Production of transgenic plant

The gene expression vector produced in Example 3 was introduced into a host plant (*Saccharum* spp. cv. NiF8) using an *Agrobacterium* method. Thus, a transgenic sugarcane in which the GUS gene expression was regulated by the expression regulatory DNA of the ecc0001 gene was produced.

[0056] Total RNA was extracted and purified from mature leaves, young leaves, stalk pith, stalk epidermis, roots, and meristem of the transgenic sugarcane. cDNA was prepared according to a standard method. The GUS gene expression level in each tissue was analyzed by a SYBRGreen method using an ABI7500 real-time PCR system (Applied

Biosystems).

[0057] As comparisons, the GUS expression levels in mature leaves and young leaves of the transgenic sugarcane in which the GUS gene expression was regulated by a cauliflower mosaic virus (CaMV)35S promoter and the GUS expression level in meristem of non-transgenic sugarcane (control) were analyzed in the above manner.

[0058] Fig. 6 shows the results. The expression levels of the individual tissues are expressed in relative values with reference to the strongest GUS gene expression level confirmed in mature leaves among the analyzed tissues of transgenic sugarcane (in which the GUS gene expression was regulated by the ecc0001 gene expression regulatory DNA), which was designated as 100%.

[0059] The above results revealed that the ecc0001 gene expression regulatory DNA can induce gene expression in a mature-leaf-specific manner to an extent approximately 36-fold greater than the GUS gene expression level regulated by a CaMV35S promoter.

Industrial Applicability

[0060] The gene expression regulatory DNA of the present invention has activity of promoting gene expression in a mature-leaf-specific manner. Since the DNA has such characteristics, it can be used to produce a plant having desired properties by causing a desired gene to be expressed in a mature-leaf-specific manner.

All publications, patents, and patent applications cited herein are incorporated herein by reference in their entirety.


Claims

- [Claim 1] A gene expression regulatory DNA having activity of promoting gene expression in a mature-leaf-specific manner, which comprises any one of the following (a) to (d):
- (a) DNA consisting of the nucleotide sequence shown in SEQ ID NO: 1 or SEQ ID NO: 4;
 - (b) DNA consisting of a nucleotide sequence that has a deletion, a substitution, an addition, or an insertion of one or a plurality of nucleotides in the nucleotide sequence shown in SEQ ID NO: 1 or SEQ ID NO: 4 and having activity of promoting gene expression in a mature-leaf-specific manner;
 - (c) DNA consisting of a nucleotide sequence having 90% or more sequence identity to the nucleotide sequence shown in SEQ ID NO: 1 or SEQ ID NO: 4 and having activity of promoting gene expression in a mature-leaf-specific manner; and
 - (d) DNA hybridizing under stringent conditions to DNA consisting of a sequence complementary to a part or the entirety of the nucleotide sequence shown in SEQ ID NO: 1 or SEQ ID NO: 4 and having activity of promoting gene expression in a mature-leaf-specific manner.
- [Claim 2] A recombinant vector, which comprises the gene expression regulatory DNA according to claim 1.
- [Claim 3] A transformed plant obtained by transformation using the recombinant vector according to claim 2.
- [Claim 4] A transformed plant obtained from plant cells transformed by plant cell transformation using the recombinant vector according to claim 2.

[Fig. 1]

1 ACGCACCCAG C CACTCACTC A GAACGAAAG C CCCAGCAAG C TCAGCTCAG C TGCTCACAC
 61 CTCACACCCA C AGGCCACAC T CACAGCGGA G CAGAGTAGT G AGCACTGAG G CAGGGCAGCA
 121 GAGCGCGCGG T CAGTTCAGG T CGATCGTCA G CCATGAGCA A AGGATGACA A GCACCACCGC
 181 CAGGAGCAC A CCTGCGT C G CTGCTGCGG G TGCATGGCGT C GTGCCCTGCT G GGCGCTGGTC
 241 CTCATCGT C C TTCAATCG C G TGGTCATC T ACCTCGCCC T TCCGCCCTC G AAGCCCGTCC
 301 TTCTACCTGC A GGACCTGCA G CTGCGGG C CCCATCTGC T TGGCCGACCC G TTCGCTGACG
 361 GCGTCGGCG C AGGTGACGCT G GCGTCCCG A ACCCTC A A CG ACGTGG G CATCTTCTAC
 421 AAGCGCCTGG A CGTGTTCGT G ACCTACCAG A ACGAGGGCG G TCACGGTGCC C CGTGTGGCTG
 481 CCGCCGCAGT A CCAGGGCCA C CGGACGTC A CCGTCTGGT C CCCCGTGCT G TTCGGCCGAG
 541 TCCGTGCCCG T GGCCGGCTA C GTCCGGGAC G ACCTCAAGC G TGACGTCCG G GCCCGGCTTC
 601 GTCGCTCTGC A GGTCAAGGT G GACGGCCCG G GTCAAAGTGG A AGTTCGGCAG G CTGGGTCTCC
 661 GGGAGCTACC A CCCTTTCGT C AGCTGCC C GCCGTGCTCT C CCGCGGGTA C CCCGCGGCTC
 721 GTGGCGGGCG G CGGCAACAA C ACCGTGT C G TCCGCTCAAGT T CGCGCAGCC C CACGGGATGC
 781 AGCGTCGAGG G TGTGATGTA T CCACTGATG A CGCACTCAA A GARCAGTTGC C CAAGTACTGC
 841 TACTCAAAAT T CTTTGGGTA C ACCGTTAAT T ACGCTGTTA A ATTTGTTCA T GTCTCATTC
 901 TCAATTCCTCA T TCTCTTGTG G GGAAATAAGT T TCTGTCCGA G GCCATATAG C CTGTTCAAAA
 961 ACTTGAAGTC A ATTGTACCA A CATGTGTTT A TTCAAAGAA G AAAAGAGGG G CCACTGCAAA
 1021 AACATACATA G ATGCGCTAA C CCCGAGCAT G CTCAAAACT A AGAAGA A CTC C CACCGACCGC
 1081 CTATAGTGGC G ACTGAATAC T GCCACA A A C TCCGGGGGG G CGCCTAAAAG A ATTACATCT
 1141 AATTTCCCTG T GTGGAAGA A ATTAATCCTC T CGGAGGGGG C CAAAAATAA A GGCACCCGGG
 1201 CCCAGTTTAA A AGAGGATGA G GGGAGAAAC A CCGGCGTTC A CAAAATATA A ATGCCCTTGA
 1261 CAGTAAACCC T CCCTTCTAC C ATGGGGTGT A TAAACAATA A AAGGCCACA C CTATAAAGC
 1321 C

[Fig. 2]



[Fig. 3]

HindIII

```

AAGCTTGATC TGTGGTATTT TGGACTTCTA TTACAAGTTT ACAGACACAG GGCTCATCTG
CAGTCGGTGG GGGAGATAAG GAGCGAATGG AGGGGCACAA ATGTGACGCC ACGCCGGGGA
AGGTTCTCTG ATCGGAGTGT ATTACGGACG GGCCGTCTGC TCCGTCGGCG TTTTGGGCTT
TTGGGGTAGG GCGGATGGAC TCGACGTCGG AACTACCAAC ATGATGATTG ATGAACGCGC
GCCCGACGCA TTGGCGCGTG CGTCCGTGCC TCCGTGCTTG CTACGCCCCA CCAACCTTCT
CCACTACGCC CGGACAAATG CCGTGGCAAG GAGGGACAAC ACCCACGCTA ACTAACGCTC
ATATAATCAA CTTTGTACTA TTTGTATAGT TATTAACTG GCAGCATTC CCAAGAAATGAC
CGCACGTAA TTCCTTCTGA CTGCCTATTG CCAACCGTTC CTTTTCTAGA CCCCATCCAT
CCAGTCCCAA ACAAGATGTA TTAGAAATGC TTACATGTGT AAATATATGT ACATAAATAC
TGCTTGCCTC GTTTTTCAAT CATTATTTAA AACTATGACT GAGTTTGATA TGTAGTAGTT
TTGCATTGAT TGGCACACAA TATGTATAAT GAGAGTGTGT CAACTTTCC TCCTCATTTT
CACTCATCCA AATTCGATGC ACGAATCCCT ACGGCCACAA CCGACTATTA AGAATAGTCC
AATTTGGCAC TACTCTAAAT AGTTGCTGGA GTTTGATGTT GTCTCTCCAA CTCCAACACG
GATATCTTGT GCGCTCCAGC TGCAAAACCA TTTATCACTA CACATTAGCA TCCTCCTCAT
CCTTTTCTTC CAACCATCGC CTCTCATGGT TTGCGATAAG GATGGGGTGG AGCGATCCAT
GTATCTCCTC TAGTTCAACT TAATGAAATA TATGTGCGAA AACAAGGCAA GGAATTGTCT
AATTAGCTCT TTCAGTTGAA ACCAGAGGCG AACCACATAT ACTCATCGGA ACATCATCCC
CATCCTTAGT TTGGGAGACA TTGATCTCAT GATCTGTCAA GCGGTTAGAG CTAGCGTTGC
CTTGTTGAAC ATTTGGCTTTG ATGTTGCAAA GACTACATCG CTTCATTAAG GGGTCGAAGA
CGTGGAAATT GTGCAGGACG AAGCAAAAGC CACACACTGC GTTGTTTGAA TACTTCACCG
TAATTCCTGG TTCAAGCTTT GAGCATACAA CCAGAATCAA TCTTGCTTG AAGCCTTAGT
CTTGCAATCC ACCAACCTCA GCGACCTTGA CGCTTGCTTC TCGGAGGCAA AAGTTTTGTC
AGTGATCAAT GACATTTTCA CAGACAAGGC GACATTACCG GACGGGTTC CTGGGCTCTT
CTACCAACGC CGCATCAAAC AAGCCCCAGA GATGTAAAAC AAACAGTTTA TATACATAAT
TAAAACACTAGT AAATTCAAGC GTGGCCGAGT GAATCTCAAG CTGACGTGTG TAGTTCCCGC
CGTCGCACTC AAGAAAGGTC AAAGGTGGAC ACGAATCTCA AAGCAATGCA TATGCATATG
CACATGCACA AGCACAACCC GACGTCCCCG TGCCAGACCC GGTGGCCCAC CGGATGAAGT
ACGTACCGAA TTAGCAGATG GAGGAATCTT CTTGCGTTGA GACGTTGCTT TCCGCTACAC
CTTCTCTGCC AGTCTGCTGT CTGCTCTGCT GCAGGCTGTC TGCTAGCTGA GCCCTGCAG
GTGATGTGCT AGCTTTTGAG TGGCACATGC GCTGTGTTTG TGTAGTCTAG TGTAGCCGGC
GTAGCAACGC TTGGTGTGCC ACTACCGATT TTAAAAGCCA TTACGCTTTC CTACACCGCA
CGGCTCAGTC ACCCACCCAC CCACTCACTC AGAACGAAAG CCCAGCAAG CTCAGCTCAG
CTGCTCACAC CTCACACCCA CAGGCCACAC TCACAGCGGA GCAGAGTAGT GAGCACTGAG
CAGGGCAGCA GAGCGCGCGG TCAGTTCAGG TCGATCGTCA GCCATGCCCTA GG

```

Initiation Bln I
codon

[Fig. 4-1]

Processed sequence:

1 AAGCTTGATCTGTGGTATTTTGGACTTCTATTACAAGTTTACAGACACAG
51 GGCTCATCTGCAGTCGGTGGGGGAGATAAGGAGCGAATGGAGGGGCACAA
101 ATGTGACGCCACGCCGGGAAGGTTCTCTGATCGGAGTGTATTACGGACG
151 GGCCGTCTGCTCCGTCCGCTTTTGGGCTTTTGGGGTAGGGCGGATGGAC
201 TCGACGTGGAACTACCAACATGATGATTGATGAACGCGCGCCCGACGCA
251 TTGGCGCGTGGTCCGTGCCTCCGTGCTTGCTACGCCCGACCAACCTTCT
301 CCAGTACGCCCGGACAAATGCCGTGGCAAGGAGGGACAACCCACGCTA
351 AGTAACGCTCATATAATCAACTTTGTACTATTTGTATAGTTATTAACCTG
401 GCAGCATTCCAAGAAATGACCGCAGGTTAATTCCTTCTGACTGCCTATTG
451 CCAACCGTTCCTTTTCTAGACCCCATCCATGCCAGTTCAAAACAAGATGT
501 ATTAGAAATGCTTACATGTGTAATATATGTACATAAAATACTGCTTGCCT
551 CGTTTTTCATTCATTATTTAAACTATGACTGAGTTTGATATGTAGTAGT
601 TTTGCATTGATTGGCACACAATATGTATAATGAGAGTGTGTCAAACCTTC
651 CTCCTCATTTTCACTCATCCAAATTCGATGCACGAATCCCTACGGCCACA
701 ACCGACTATTAAGAATAGTCCAATTTGGCACTACTCTAAATAGTTGCTGG
751 AGTTTGATGTTGTCTCTCCAACCTCCAACACGGATATCTTGTGCGCTCCAG
801 CTGCAAAACCATTTATCACTACACATTAGCATCCTCCTCATCCTTTTCTT
851 CCAACCATCGCCTCTCATGGTTTTGCGATAAGGATGGGGTGGAGCGATCCA
901 TGTATCTCCTCTAGTTCAACTTAATGAAATATATGTGCGAAAACAAGGCA
951 AGGAATTGTCTAATTAGCTCTTTAGTTGAAACCAGAGGGGAACCACATA
1001 TACTCATCGGAACATCATCCCCATCCTTAGTTTGGGAGACATTGATCTCA
1051 TGATCTGTCAAGGCGTTAGAGCTAGCGTTGCCTTGTGAAACATTGGCTTT
1101 GATGTTGCAAAGACTACATCGCTTCATTAAGGGGTGGAAGACGTGGAAT
1151 TGTGCAGGACGAAGCAAAAGCCACACACTGCGTTGTTTGAATACTTCACC
1201 GTAATTCTTGGTTCAAGCTTTGAGCATACAACCAGAATCAATCTTGTCTT
1251 GAAGCCTTAGTCTTGCAATCCACCAACCTCAGCGACCTTGACGCTTGCTT
1301 CTCGGAGGCAAAAGTTTTGTCAGTGATCAATGACATTTTCATCAGACAAGG
1351 CGACATTACGGACGGGTTCACTGGGCTCTTCTACCAACGCCGCATCAAA
1401 CAAGCCCCAGAGATGTAAAACAAACAGTTTATATACATAATTAAACTAG
1451 TAAATTCAAGCGTGGCCGAGTGAATCTCAAGCTGACGTGTGTAGTTCCCG
1501 CCGTGCGACTCAAGAAAGGTCAAAGGTGGACACGAATCTCAAAGCAATGC
1551 ATATGCATATGCACATGCACAAGCACAACCCGACGTCCCGTGCCAGACC
1601 CGGTGGCCACCGGATGAAGTACGTACCGAATTAGCAGATGGAGGAATCT
1651 TCTTCGGTTGAGACGTTGCTTCCGCTACACCTTCTCTGCCAGTCTGCTG

[Fig. 4-2]

```

1701 TCTGCTCTGCTGCAGGCTGTCTGCTAGCTGAGCCCTGCAGCGAGGGTTAA
1751 CCAATCGTCCATTGCGTATGTGCTAGCTTTTGGAGTGGCACATGCGCTGTG
1801 TTTGTGTAGTCTAGTGTAGCCGGGTAGCAACGCTTGGTGTGCCACTACC
1851 GATTTTAAAAGCCATTACGCTTTCCTACACCGCACGGCTCAGTCACCCAC
1901 CCACCCACTCACTCAGAACGAAAGCCCCAGCAAGCTCAGCTCAGCTGCTC
1951 ACACCTCACACCCACAGGCCACACTCACAGCGGAGCAGAGTAGTGAGCAC
2001 TGAGCAGGGCAGCAGAGCGCGGGTCAGTTCAGGTCGATCGTCAGCCATG
2051 CCTAGGN

```

: Predicted promoter region

(An HindIII restriction enzyme recognition sequence (AAGCTT) and a BlnI restriction enzyme recognition sequence (CCTAGG) are present at the 5'-end and the 3'-end, respectively.)

Proscan: Version 1.7

Processed Sequence: 2057 Base Pairs

Promoter region predicted on reverse strand in 1669 to 1419


Promoter Score: 64.48 (Promoter Cutoff = 53.000000)

TATA found at 1437, Est.TSS = 1405


Significant Signals:

Name	Strand	Location	Weight
AP-2	+	1607	1.091000
Sp1	-	1502	3.191000
Sp1	+	1497	3.119000
ATF/CREB	-	1488	1.564000
E4F1	-	1488	1.201000
NF-S	-	1487	1.019000
CREB	+	1483	5.737000
ATF	+	1483	1.591000
UCE.2	+	1464	1.216000
TFIID	-	1435	1.971000
TFIID	-	1433	2.618000

[Fig. 5]



[Fig. 6]



INTERNATIONAL SEARCH REPORT

International application No

PCT/JP2011/007337

A. CLASSIFICATION OF SUBJECT MATTER
 INV. C07K14/415 C12N15/82 A01H5/12
 ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
 C12N C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, BIOSIS, Sequence Search, EMBASE, MEDLINE, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	DATABASE EMBL [Online] 24 September 2003 (2003-09-24), "SCEZAM2060G03.g AM2 Saccharum hybrid cultivar SP80-3280 cDNA clone SCEZAM2060G03 5', mRNA sequence.", XP002672582, retrieved from EBI accession no. EM_EST:CA085096 Database accession no. CA085096 the whole document ----- -/--	1-4



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance
 "E" earlier document but published on or after the international filing date
 "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
 "O" document referring to an oral disclosure, use, exhibition or other means
 "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
 "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
 "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
 "&" document member of the same patent family

Date of the actual completion of the international search

29 March 2012

Date of mailing of the international search report

16/04/2012

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040,
 Fax: (+31-70) 340-3016

Authorized officer

Kurz, Birgit

INTERNATIONAL SEARCH REPORT

International application No

PCT/JP2011/007337

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>DATABASE Geneseq [Online]</p> <p>12 June 2008 (2008-06-12), "Maize promoter sequence SEQ ID 6836.", XP002672583, retrieved from EBI accession no. GSN:AQE59106 Database accession no. AQE59106 the whole document</p>	1-4
A	<p>-& US 2007/130645 A1 (WU WEI [US] ET AL) 7 June 2007 (2007-06-07) Seq. ID No. 6836; paragraphs 236-239, 242-263</p>	1-4
A	<p>-----</p> <p>VAN DER MERWE J ET AL: "Isolation and evaluation of a developmentally regulated sugarcane promoter", PROCEEDINGS OF THE SOUTH AFRICAN SUGAR TECHNOLOGISTS' ASSOCIATION, X, XX, vol. 77, 1 January 2003 (2003-01-01), pages 146-149, XP009088003, abstract; pages 146 and 149</p>	1-4
A	<p>-----</p> <p>WEI H ET AL: "Comparative expression analysis of two sugarcane polyubiquitin promoters and flanking sequences in transgenic plants", JOURNAL OF PLANT PHYSIOLOGY, FISCHER, STUTTGART, DE, vol. 160, no. 10, 1 January 2003 (2003-01-01), pages 1241-1251, XP004955105, ISSN: 0176-1617, DOI: 10.1078/0176-1617-01086 abstract; pages 1244, 1249, 1250</p>	1-4
A	<p>-----</p> <p>STEPHEN R MUDGE ET AL: "Efficient silencing of reporter transgenes coupled to known functional promoters in sugarcane, a highly polyploid crop species", PLANTA ; AN INTERNATIONAL JOURNAL OF PLANT BIOLOGY, SPRINGER, BERLIN, DE, vol. 229, no. 3, 15 November 2008 (2008-11-15), pages 549-558, XP019715456, ISSN: 1432-2048 abstract; pages 550, 555-557</p> <p>-----</p>	1-4

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/JP2011/007337

Patent document cited in search report	Publication date	Patent family member(s)	Publication date	
US 2007130645	A1	07-06-2007	US 2007130645 A1	07-06-2007
			US 2009178163 A1	09-07-2009
			US 2011078833 A1	31-03-2011
