




- (72) KOJIMA, Koichi, JP
(72) SAKAI, Junichi, JP
(72) SAMATA, Naozumi, JP
(72) KOZUKA, Masao, JP
(72) YOSHIMI, Kenji, JP
(72) KANEKO, Tsugio, JP
(71) SANKYO COMPANY LIMITED, JP
(51) Int.Cl.⁶ C07D 261/12, A61K 31/425, A61K 31/42, C07D 413/04,
C07D 261/10, C07D 417/04
(30) 1996/02/27 (8/39819) JP
(54) **DERIVES D'ISOXAZOLE**
(54) **ISOXAZOLE DERIVATIVES**



(I)


(57) Cette invention concerne des dérivés d'isoxazole qui correspondent à la formule générale (I) où R¹ représente aryle éventuellement substitué ou hétéroaryle éventuellement substitué; R² représente hydrogène, halogéno, alkyle éventuellement substitué, alcényle, alcynyle, cycloalkyle, cycloalcényle, alcoxy, cyano, carboxy, alcanoyle, alcoxycarbonyle ou carbamoyle éventuellement substitué; R³ représente amino éventuellement substitué ou un groupe hétérocyclique saturé; X représente oxygène ou soufre; et n est un nombre entier allant de 2 à 6. Ces dérivés possèdent une excellente activité inhibitrice de monoamine oxydase, et peuvent être utilisés dans le traitement ou la prévention de maladies nerveuses telle que la dépression.

(57) Isoxazole derivatives represented by general formula (I), which have an excellent monoamine oxidase inhibitory activity and are useful for the treatment or prevention of nervous diseases such as depression, wherein R¹ represents optionally substituted aryl or optionally substituted heteroaryl; R² represents hydrogen, halogeno, optionally substituted alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, alkoxy, cyano, carboxy, alkanoyl, alcoxycarbonyl, or optionally substituted carbamoyl; R³ represents optionally substituted amino or saturated heterocyclic group; X represents oxygen or sulfur; and n is an integer of 2 to 6.



Abstract

The isoxazole derivatives having the following general formula:



(I)

[wherein R¹ represents an optionally substituted aryl group or aromatic heterocyclic group; R² represents a hydrogen atom, a halogen atom, an optionally substituted alkyl group, an alkenyl group, an alkynyl group, a cycloalkyl group, a cycloalkenyl group, an alkoxy group, a cyano group, a carboxyl group, an alkanoyl group, an alkoxy carbonyl group or an optionally substituted carbamoyl group; R³ represents an optionally substituted amino group or a saturated heterocyclic group; X represents an oxygen atom or a sulfur atom; n is an integer of from 2 to 6] has an excellent monoamine oxidase inhibitory activity, and is useful as a therapeutic agent or a preventive agent against nervous diseases such as depression.

Specification

Isoxazole Derivatives

[Technical Field]

The present invention relates to isoxazole derivatives and pharmaceutically acceptable salts thereof, which have excellent inhibition activities against type A-monoamine oxidase;

compositions containing the compounds for treating or preventing nervous diseases (particularly for depression), including depression, Parkinson's disease, Alzheimer's dementia (cognitive disorder owing to Alzheimer's disease) or cerebrovascular dementia (cognitive disorder owing to cerebrovascular dementia);

use of the compounds for producing pharmaceutical preparations for treating or preventing the above-mentioned diseases; and

a method for treating or preventing the above diseases by administering a pharmaceutically effective amount of the compounds to warm-blood animals.

[Background Art]

Depression is a disease which shows a typical condition of suppressed mood among mood disorders, and one of its causes is considered to be functional disorders in the central serotonergic and noradrenergic nervous systems. Serotonin and noradrenaline are decomposed and metabolized by monoamine oxidases (mainly by type A-monoamine oxidase) to lose their biological activities. Type A-monoamine oxidase inhibitors are considered to be useful as antidepressants and such inhibitors have been studied and developed intensively. Recently, Moclobemide has been supplied clinically as a selective type A-monoamine oxidase inhibitor.

[Disclosure of the Invention]

The present inventors made intensive studies for years on the synthesis of

isoxazole derivatives and their pharmacological activities with the aim of developing an excellent therapeutic agent for depression and found that isoxazole derivatives having a specific structure exhibit potent type A-monoamine oxidase inhibitory activities and have therapeutic or preventing effects (particularly therapeutic effect) on nervous diseases (particularly for depression) including depression, Parkinson's disease, Alzheimer's dementia (cognitive disorder owing to Alzheimer's disease) or cerebrovascular dementia (cognitive disorder owing to cerebrovascular dementia).

The present invention provides


isoxazole derivatives and pharmaceutically acceptable salts thereof, which have excellent inhibition activities to type A-monoamine oxidase;

compositions containing the compounds for treating or preventing nervous diseases (particularly for depression) including depression, Parkinson's disease, Alzheimer's dementia (cognitive disorder owing to Alzheimer's disease) or cerebrovascular dementia (cognitive disorder owing to cerebrovascular dementia);

use of the compounds for producing pharmaceutical preparations for treating or preventing the above-mentioned diseases; and

a method for treating or preventing the above diseases by administering a pharmaceutically effective amount of the compounds to warm-blood animals.

The isoxazole derivatives of the present invention have the general formula (I):



(I)

wherein

R¹ represents a C₆ - C₁₄ aryl group optionally having from 1 to 3 substituents which may be the same as or different from each other and selected from the following substituent group, or a 5- or 6-membered aromatic heterocyclic group optionally having from 1 to 3 substituents and having one or

two hetero atoms which may be the same as or different from each other and selected from the group consisting of nitrogen, oxygen and sulfur atoms [the substituent group is a halogen; a C₁ - C₆ alkyl; a C₁ - C₆ alkyl substituted with a halogen or a C₁ - C₆ alkoxy; a C₁ - C₆ alkoxy; a C₆ - C₁₄ aryl, a C₇ - C₁₈ aralkyl, a C₆ - C₁₄ aryloxy or a C₇ - C₁₈ aralkyloxy optionally having from 1 to 3 substituents which may be the same as or different from each other and selected from the following group (the substituent group is a halogen, a C₁ - C₆ alkyl or a C₁ - C₆ alkoxy); a cyano; a nitro; a hydroxyl group; a C₁ - C₇ alkanoyl; a C₁ - C₇ alkanoyloxy; a C₂ - C₇ alkoxy-carbonyl; an amino; a carbamoyl; a mono (C₁ - C₆ alkyl)carbamoyl; a di(C₁ - C₆ alkyl)carbamoyl or a mono C₇ - C₁₅ arylcarbonylamino group optionally having from 1 to 3 substituents (the substituent group is a halogen, a C₁ - C₆ alkyl or a C₁ - C₆ alkoxy)],

R² represents a hydrogen atom; a halogen atom; a C₁ - C₆ alkyl; a C₁ - C₆ alkyl substituted with a halogen or a C₁ - C₆ alkoxy; a C₂ - C₆ alkenyl; a C₂ - C₆ alkynyl; a C₃ - C₁₀ cycloalkyl; a C₃ - C₁₀ cycloalkenyl; a C₁ - C₆ alkoxy; a cyano; a carboxyl; a C₁ - C₇ alkanoyl; a C₂ - C₇ alkoxy-carbonyl; a carbamoyl; a mono (C₁ - C₆ alkyl)carbamoyl or a di(C₁ - C₆ alkyl)carbamoyl group,

R³ represents an amino, a mono C₁ - C₆ alkylamino, a di(C₁ - C₆ alkyl)amino, a mono C₁ - C₇ alkanoylamino, a mono C₂ - C₇ alkoxy-carbonylamino, a mono C₇ - C₁₅ arylcarbonylamino optionally having from 1 to 3 substituents which may be the same as or different from each other and selected from the following group (the substituent group is a halogen, a C₁ - C₆ alkyl or a C₁ - C₆ alkoxy group) or a 5- or 6-membered saturated heterocyclic group (attached through a ring nitrogen atom), which contains one nitrogen atom and optionally may contain further one nitrogen atom or oxygen atom.

X represents an oxygen or a sulfur atom, and

n represents an integer of 2 to 6.

Further, the active ingredient of the monoamine oxidase inhibitor of the present invention is the isoxazole derivative of general formula (I).

In the formula (I), "the halogen atom" in the definition of R^2 and the definition of the substituent included in R^1 may be, for example, a fluorine, chlorine, bromine or iodine atom, preferably the fluorine or chlorine atom, more preferably the chlorine atom.

In the formula (I), "the $C_1 - C_6$ alkyl group" in the definition of R^2 and in the definition of the substituent included in R^1 may be, for example, a straight or branched alkyl group having from 1 to 6 carbon atoms such as a methyl, ethyl, propyl, isopropyl, butyl, isobutyl, s-butyl, t-butyl, pentyl, isopentyl, 2-methylbutyl, neopentyl, 1-ethylpropyl, hexyl, 4-methylpentyl, 3-methylpentyl, 2-methylpentyl, 1-methylpentyl, 3,3-dimethylbutyl, 2,2-dimethylbutyl, 1,1-dimethylbutyl, 1,2-dimethylbutyl, 1,3-dimethylbutyl, 2,3-dimethylbutyl or 2-ethylbutyl group. The substituent included in R^1 is preferably the $C_1 - C_4$ alkyl group, more preferably the methyl or ethyl group, and most preferably the methyl group. Further, R^2 is preferably the $C_1 - C_4$ alkyl group, more preferably the ethyl, propyl, isopropyl, isobutyl or t-butyl group, most preferably the isopropyl group.

In the formula (I), "the $C_2 - C_6$ alkenyl group" in the definition of R^2 may be, for example, a straight or branched alkenyl group having from 2 to 6 carbon atoms with one or two double bonds such as a vinyl, 1-propenyl, allyl, 1-methyl-1-propenyl, 1-methyl-2-propenyl, 2-methyl-1-propenyl, isopropenyl, allenyl, 1-butenyl, 2-butenyl, 3-butenyl, 2-pentenyl, isoprenyl, 5-hexenyl or 1,4-hexadienyl group, preferably the vinyl, 1-propenyl, allyl, 1-methyl-1-propenyl, isopropenyl, 2-butenyl or 3-butenyl group, more preferably the allyl,

isopropenyl or 2-butenyl groups, most preferably the allyl group.

In the formula (I), "the C₂ - C₆ alkynyl group" in the definition of R² may be, for example, a straight or branched alkynyl group having from 2 to 6 carbon atoms such as an ethynyl, 1-propynyl, propargyl, 1-methyl-2-propynyl, 2-methyl-2-propynyl, 2-ethyl-2-propynyl, 2-butynyl, 1-methyl-2-butynyl, 2-methyl-2-butynyl, 3-butynyl, 2-pentynyl, 5-hexynyl or 2-methyl-4-pentynyl group, preferably the ethynyl, propargyl, 2-butynyl or 3-butynyl group, more preferably the propargyl group.

In the formula (I), "the C₃ - C₁₀ cycloalkyl group" in the definition of R² may be, for example, a 3- to 10-membered saturated cyclic hydrocarbon group which may form a condensed ring, such as a cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, norbornyl or adamantyl group, preferably the cyclopropyl, cyclopentyl or cyclohexyl group, more preferably the cyclopropyl group.

In the formula (I), "the C₃ - C₁₀ cycloalkenyl group" in the definition of R² may be, for example, a 3- to 10-membered unsaturated cyclic hydrocarbon group which may form a condensed ring having one double bond, such as a 2-cyclopropenyl, 2-cyclobutenyl, 1-cyclopentenyl, 2-cyclopentenyl, 3-cyclopentenyl, 1-cyclohexenyl, 2-cyclohexenyl, 3-cyclohexenyl, 2-cycloheptenyl, 3-norbornenyl or 3-adamantenyl group, preferably the 2-cyclopentenyl, 3-cyclopentenyl, 2-cyclohexenyl or 3-cyclohexenyl group, more preferably the 2-cyclopentenyl group.

In the formula (I), "C₁ - C₆ alkyl group substituted with a halogen or a C₁ - C₆ alkoxy" in the definition of R² and in the definition of the substituent included in R¹ represents a group in which 1 to 5 "halogen atoms" mentioned above are bonded to the above-mentioned "C₁ - C₆ alkyl group" or a group in

which "the C₁ - C₆ alkoxy" mentioned below is bonded to the above-mentioned C₁ - C₆ alkyl group. The group in which the halogen is bonded to the C₁ - C₆ alkyl group may be, for example, a fluoromethyl, difluoromethyl, trifluoromethyl, 2-fluoroethyl, 2-chloroethyl, 2,2,2-trifluoroethyl, 3-fluoropropyl, 3-chloropropyl, 3-bromopropyl, 4-fluorobutyl or 6-iodohexyl group. The group in which C₁ - C₆ alkoxy is bonded to the C₁ - C₆ alkyl group may be, for example, a methoxymethyl, ethoxymethyl, propoxymethyl, butoxymethyl, methoxyethyl, ethoxyethyl, propoxyethyl, butoxyethyl, propoxypropyl, butoxybutyl or hexyloxyhexyl group. The substituent included in R¹ is preferably a C₁ - C₆ alkyl group substituted with 1 to 3 halogens or a C₁ - C₄ alkoxy, more preferably the fluoromethyl, difluoromethyl, trifluoromethyl, 2-fluoroethyl, 2-chloroethyl, 2,2,2-trifluoroethyl, methoxymethyl or methoxyethyl groups, still more preferably the trifluoromethyl, 2,2,2-trifluoroethyl or methoxymethyl group, most preferably the trifluoromethyl group. Further, R² is preferably the fluoromethyl, difluoromethyl, trifluoromethyl, 2-fluoroethyl, 1-chloroethyl, 2-chloroethyl, 2,2,2-trifluoroethyl, methoxymethyl or methoxyethyl group, more preferably the trifluoromethyl, 2-fluoroethyl, 1-chloroethyl or 2-chloroethyl group, most preferably the 1-chloroethyl group.

In the formula (I), "the C₁ - C₆ alkoxy group" in the definition of the substituent included in R¹ and R² and in the definition of R² represents a group in which the above-mentioned "C₁ - C₆ alkyl group" is bonded to an oxygen atom and such group may be, for example, a methoxy, ethoxy, propoxy, isopropoxy, butoxy, isobutoxy, s-butoxy, t-butoxy, pentoxy, isopentoxy, 2-methylbutoxy, neopentoxy, 1-ethylpropoxy, hexyloxy, 4-methylpentoxy, 3-methylpentoxy, 2-methylpentoxy, 2-methylpentoxy, 1-methylpentoxy, 3,3-dimethylbutoxy, 2,2-dimethylbutoxy, 1,1-dimethylbutoxy, 1,2-dimethylbutoxy, 1,3-dimethylbutoxy, 2,3-dimethylbutoxy or 2-ethylbutoxy group, preferably the C₁ - C₄ alkoxy group, more preferably the methoxy or ethoxy group, most preferably the methoxy group.

In the formula (I), the “C₆ - C₁₄ aryl group optionally having from 1 to 3 substituents which may be the same as or different from each other and selected from the following group (the substituent group is a halogen, a C₁ - C₆ alkyl or a C₁ - C₆ alkoxy)” in the definition of the substituent included in R¹ represents an aromatic hydrocarbon group having from 6 to 14 carbon atoms optionally having from 1 to 3 substituents which may be the same as or different from each other and selected from the substituent group, and such group may be, for example, a phenyl, fluorophenyl, chlorophenyl, dichlorophenyl, methylphenyl, trimethylphenyl, methoxyphenyl, indenyl, methylindenyl, naphthyl, dichloronaphthyl, phenanthrenyl, hexylphenanthrenyl, anthracenyl, dimethylanthracenyl or hexyloxyanthracenyl group, preferably the phenyl group optionally having one or two substituents which may be the same as or different from each other and selected from the group consisting of fluorine, chlorine, methyl and methoxy, more preferably the phenyl, 4-fluorophenyl, 4-chlorophenyl, 2,4-dichlorophenyl, 4-methylphenyl or 4-methoxyphenyl group, most preferably the phenyl group.

In the formula (I), “the C₇ - C₁₈ aralkyl group optionally having from 1 to 3 substituents which may be the same as or different from each other and selected from the following group (the substituent group is a halogen, a C₁ - C₆ alkyl or a C₁ - C₆ alkoxy)” in the definition of the substituent included in R¹ represents a group in which one or two phenyl groups or one naphthyl group optionally having from 1 to 3 substituents which may be the same as or different from each other and selected from the substituent group is bonded to the above-mentioned “C₁ - C₆ alkyl”, and such group may be, for example, a benzyl, fluorobenzyl, difluorobenzyl, trifluorobenzyl, chlorobenzyl, dichlorobenzyl, trichlorobenzyl, bromobenzyl, methylbenzyl, dimethylbenzyl, trimethylbenzyl, ethylbenzyl, propylbenzyl, methoxybenzyl, dimethoxybenzyl, ethoxybenzyl, hexyloxybenzyl, diphenylmethyl, naphthylmethyl, fluoronaphthylmethyl, difluoronaphthylmethyl, chloronaphthylmethyl, dichloronaphthylmethyl,

methylnaphthylmethyl, dimethylnaphthylmethyl, ethylnaphthylmethyl, phenethyl, fluorophenethyl, difluorophenethyl, chlorophenethyl, dichlorophenethyl, methylphenethyl, trimethylphenethyl, naphthylethyl, fluoronaphthylethyl, chloronaphthylethyl, phenylpropyl, fluorophenylpropyl, chlorophenylpropyl, dichlorophenylpropyl, methylphenylpropyl, dimethylphenylpropyl, trimethylphenylpropyl, naphthylpropyl, iodonaphthylpropyl, hexylnaphthylpropyl, methoxynaphthylpropyl, hexyloxynaphthylpropyl, phenylbutyl, fluorophenylbutyl, difluorophenylbutyl, chlorophenylbutyl, dichlorophenylbutyl, trimethylphenylbutyl or naphthylbutyl group, preferably the benzyl group or phenethyl group optionally having from 1 to 3 substituents which may be the same as or different from each other and selected from the group consisting of halogen, C₁ - C₄ alkyl and C₁ - C₄ alkoxy, more preferably, the benzyl group optionally having one substituent selected from the group consisting of fluorine, chlorine, C₁ - C₄ alkyl and C₁ - C₄ alkoxy, still more preferably the benzyl, fluorobenzyl, chlorobenzyl, difluorobenzyl, dichlorobenzyl, methylbenzyl, dimethylbenzyl or methoxybenzyl group, most preferably the benzyl group.

In the formula (I), "the C₆ - C₁₄ aryloxy group optionally having from 1 to 3 substituents which may be the same as or different from each other and selected from the following group (the substituent group is a halogen, a C₁ - C₆ alkyl or a C₁ - C₆ alkoxy)" in the definition of the substituent included in R¹ represents a group in which the "C₆ - C₁₄ aryl group optionally having from 1 to 3 substituents which may be the same as or different from each other and selected from the following group (the substituent group is halogen, C₁ - C₆ alkyl or C₁ - C₆ alkoxy)" is bonded to an oxygen atom, and such group may be, for example, a phenoxy, fluorophenoxy, chlorophenoxy, dichlorophenoxy, methylphenoxy, trimethylphenoxy, methoxyphenoxy, indenylloxy, methylindenylloxy, naphthylloxy, dichloronaphthylloxy, phenanthrenylloxy, hexylphenanthrenylloxy, anthracenylloxy, dimethylanthracenylloxy or hexyloxyanthracenylloxy group; preferably the phenoxy group optionally having one or

two substituents which may be the same as or different from each other and selected from the group consisting of fluorine, chlorine, methyl and methoxy, more preferably the phenoxy, 4-fluorophenoxy, 4-chlorophenoxy, 2,4-dichlorophenoxy, 4-methylphenoxy or 4-methoxyphenoxy group; most preferably the phenoxy group.

In the formula (I), "the C₇ - C₁₈ aralkyloxy group optionally having from 1 to 3 substituents which may be the same as or different from each other and selected from the following group (the substituent group is halogen, C₁ - C₆ alkyl or C₁ - C₆ alkoxy)" in the definition of the substituent included in R¹ represents a group in which the "C₇ - C₁₈ aralkyl group optionally having from 1 to 3 substituents which may be the same as or different from each other and selected from the following group (the substituent group is halogen, C₁ - C₆ alkyl or C₁ - C₆ alkoxy)" is bonded to an oxygen atom, and such group may be, for example, a benzyloxy, fluorobenzyloxy, difluorobenzyloxy, trifluorobenzyl-oxy, chlorobenzyloxy, dichlorobenzyloxy, trichlorobenzyloxy, bromobenzyl-oxy, methylbenzyloxy, dimethylbenzyloxy, trimethylbenzyloxy, ethylbenzyl-oxy, propylbenzyloxy, methoxybenzyloxy, dimethoxybenzyloxy, ethoxybenzyl-oxy, hexyloxybenzyloxy, diphenylmethoxy, naphthylmethoxy, fluoronaphthyl-methoxy, difluoronaphthylmethoxy, chloronaphthylmethoxy, dichloronaphthyl-methoxy, methylnaphthylmethoxy, dimethylnaphthylmethoxy, ethylnaphthyl-methoxy, phenylethoxy, fluorophenylethoxy, difluorophenylethoxy, chloro-phenylethoxy, dichlorophenylethoxy, methylphenylethoxy, trimethylphenyl-ethoxy, naphthylethoxy, fluoronaphthylethoxy, chloronaphthylethoxy, phenyl-propoxy, fluorophenylpropoxy, chlorophenylpropoxy, dichlorophenylpropoxy, methylphenylpropoxy, dimethylphenylpropoxy, trimethylphenylpropoxy, naphthylpropoxy, iodonaphthylpropoxy, hexylnaphthylpropoxy, methoxy-naphthylpropoxy, hexyloxynaphthylpropoxy, phenylbutoxy, fluorophenyl-butoxy, difluorophenylbutoxy, chlorophenylbutoxy, dichlorophenylbutoxy, trimethylphenylbutoxy or naphthylbutoxy group; preferably the benzyloxy and phenylethoxy groups optionally having from 1 to 3 substituents which may be

the same as or different from each other and selected from the group consisting of halogen, C₁ - C₄ alkyl and C₁ - C₄ alkoxy, more preferably the benzyloxy group optionally having one substituent selected from the group consisting of fluorine, chlorine, C₁ - C₄ alkyl and C₁ - C₄ alkoxy, still more preferably the benzyloxy, 4-fluorobenzyloxy, 4-chlorobenzyloxy, 2,4-difluorobenzyloxy, 2,4-dichlorobenzyloxy, 4-methylbenzyloxy, 2,4-dimethylbenzyloxy or 4-methoxybenzyloxy group, most preferably the benzyloxy group.

In the formula (I), "the C₁ - C₇ alkanoyl group" in the definition of the substituent included in R¹ and in the definition of R² represents a group in which a hydrogen atom or the above-mentioned "C₁ - C₆ alkyl" is bonded to a carbonyl group, and such group may be, for example, a formyl, acetyl, propionyl, butyryl, isobutyryl, valeryl, isovaleryl, pivaloyl, hexanoyl or heptanoyl group, preferably the formyl or acetyl group, most preferably the acetyl group.

In the formula (I), "the C₁ - C₇ alkanoyloxy group" in the definition of the substituent included in R¹ represents a group in which the above-mentioned "the C₁ - C₇ alkanoyl group" is bonded to an oxygen atom, and such group may be, for example, a formyloxy, acetoxy, propionyloxy, butyryloxy, isobutyryloxy, valeryloxy, isovaleryloxy, pivaloyloxy, hexanoyloxy or heptanoyloxy group, preferably the formyloxy or acetoxy group, more preferably the acetoxy group.

In the formula (I), "the C₂ - C₇ alkoxy carbonyl group" in the definition of the substituent included in R¹ and in the definition of R² represents a group in which the above-mentioned "C₁ - C₆ alkoxy group" is bonded to a carbonyl group, and such group may be, for example, a methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, isopropoxycarbonyl, butoxycarbonyl, isobutoxycarbonyl, s-butoxycarbonyl, t-butoxycarbonyl, pentoxycarbonyl, isopentoxycarbonyl,

carbonyl, 2-methylbutoxycarbonyl, neopentoxycarbonyl, 1-ethylpropoxycarbonyl, hexyloxycarbonyl, 4-methylpentoxycarbonyl, 3-methylpentoxycarbonyl, 2-methylpentoxycarbonyl, 1-methylpentoxycarbonyl, 3,3-dimethylbutoxycarbonyl, 2,2-dimethylbutoxycarbonyl, 1,1-dimethylbutoxycarbonyl, 1,2-dimethylbutoxycarbonyl, 1,3-dimethylbutoxycarbonyl, 2,3-dimethylbutoxycarbonyl or 2-ethylbutoxycarbonyl group, preferably a C₂ - C₅ alkoxycarbonyl group, more preferably the methoxycarbonyl or ethoxycarbonyl group, most preferably the methoxycarbonyl group.

In the formula (I), "the mono C₁ - C₆ alkylamino group" in the definition of R³ represents a group in which the above-mentioned "C₁ - C₆ alkyl group" is bonded to an amino group, and such group may be, for example, a methylamino, ethylamino, propylamino, isopropylamino, butylamino, isobutylamino, s-butylamino, t-butylamino, pentylamino or hexylamino group, preferably a mono C₁ - C₄ alkylamino group, more preferably the methylamino or ethylamino groups, most preferably the methylamino group.

In the formula (I), "the di(C₁ - C₆ alkyl)amino" in the definition of R³ may be, for example, a N,N-dimethylamino, N-ethyl-N-methylamino, N-methyl-N-propylamino, N-isopropyl-N-methylamino, N-butyl-N-methylamino, N-isobutyl-N-methylamino, N-s-butyl-N-methylamino, N-t-butyl-N-methylamino, N,N-diethylamino, N-ethyl-N-propylamino, N-ethyl-N-isobutylamino, N,N-dipropylamino, N,N-dibutylamino, N,N-dipentylamino or N,N-dihexylamino group, preferably the di(C₁ - C₄ alkyl)amino group, more preferably the N,N-dimethylamino or N,N-diethylamino group, most preferably the N,N-dimethylamino group.

In the formula (I), "the C₁ - C₇ alkanoylamino group" in the definition of R³ represents a group in which the above-mentioned "C₁ - C₇ alkanoyl group" is bonded to an amino group, and such group may be, for example, a

formylamino, acetylamino, propionylamino, butyrylamino, isobutyrylamino, valerylamino, isovalerylamino, pivaloylamino, hexanoylamino or heptanoylamino group, preferably the formylamino or acetylamino group, particularly preferably the acetylamino group.

In the formula (I), "the C₂ - C₇ alkoxy-carbonylamino group" in the definition of R³ represents a group in which the above-mentioned "C₂ - C₇ alkoxy-carbonyl group" is bonded to an amino group, and such group may be, for example, a methoxycarbonylamino, ethoxycarbonylamino, propoxycarbonylamino, isopropoxycarbonylamino, butoxycarbonylamino, isobutoxycarbonylamino, s-butoxycarbonylamino, t-butoxycarbonylamino, pentoxycarbonylamino, isopentoxycarbonylamino, 2-methylbutoxycarbonylamino, neopentoxycarbonylamino, 1-ethylpropoxycarbonylamino, hexyloxycarbonylamino, 4-methylpentoxycarbonylamino, 3-methylpentoxycarbonylamino, 2-methylpentoxycarbonylamino, 1-methylpentoxycarbonylamino, 3,3-dimethylbutoxycarbonylamino, 2,2-dimethylbutoxycarbonylamino, 1,1-dimethylbutoxycarbonylamino, 1,2-dimethylbutoxycarbonylamino, 1,3-dimethylbutoxycarbonylamino, 2,3-dimethylbutoxycarbonylamino or 2-ethylbutoxycarbonylamino group, preferably the C₂ - C₅ alkoxy-carbonylamino group, more preferably the methoxycarbonylamino or ethoxycarbonylamino group, most preferably the methoxycarbonylamino group.

In the formula (I), "the mono(C₁ - C₆ alkyl)carbamoyl group" in the definition of the substituent included in R¹ and in the definition of R² may be, for example, a methylcarbamoyl, ethylcarbamoyl, propylcarbamoyl, isopropylcarbamoyl, butylcarbamoyl, isobutylcarbamoyl, s-butylcarbamoyl, t-butylcarbamoyl, pentylcarbamoyl or hexylcarbamoyl group; preferably the mono(C₁ - C₄ alkyl)carbamoyl group; more preferably the methylcarbamoyl or ethylcarbamoyl group.

In the formula (I), "the di(C₁ - C₆ alkyl)carbamoyl group" in the

definition of the substituent included in R¹ and in the definition of R² may be, for example, a N,N-dimethylcarbamoyl, N-ethyl-N-methylcarbamoyl, N,N-diethylcarbamoyl, N,N-dipropylcarbamoyl, N,N-diisopropylcarbamoyl, N,N-dibutylcarbamoyl, N,N-diisobutylcarbamoyl, N,N-di-s-butylcarbamoyl, N,N-di-t-butylcarbamoyl, N,N-dipentylcarbamoyl or N,N-dihexylcarbamoyl group, preferably the di(C₁ - C₄ alkyl)carbamoyl group, more preferably the N,N-dimethylcarbamoyl or N,N-diethylcarbamoyl group, most preferably the N,N-dimethylcarbamoyl group.

In the formula (I), “the mono C₇ - C₁₅ arylcarbonylamino group optionally having from 1 to 3 substituents which may be the same as or different from each other and selected from the following group (the substituent group is halogen, C₁ - C₆ alkyl or C₁ - C₆ alkoxy)” in the definition of the substituent included in R¹ and in the definition of R³ represents a group in which the above-mentioned “C₆ - C₁₄ aryl group optionally having from 1 to 3 substituents which may be the same as or different from each other and selected from the following group (the substituent group is halogen, C₁ - C₆ alkyl or C₁ - C₆ alkoxy)” is bonded to a carbonylamino group, and such group may be, for example, a benzoylamino, fluorobenzoylamino, chlorobenzoylamino, dichlorobenzoylamino, toluoylamino, trimethylbenzoylamino, anisoylamino, indenoylamino, methylindenoylamino, naphthoylamino, dichloronaphthoylamino, phenanthrenoylamino, hexylphenanthrenoylamino, anthracenoylamino, dimethylanthracenoylamino or hexyloxyanthracenoylamino group; preferably the benzoylamino group optionally having one or two substituents which may be the same as or different from each other and selected from the group consisting of fluorine, chlorine, methyl and methoxy; more preferably the benzoylamino, 4-fluorobenzoylamino, 4-chlorobenzoylamino, 2,4-dichlorobenzoylamino, 4-toluoylamino or 4-anisoylamino group; most preferably the benzoylamino group.

In the formula (I), “the 5- or 6-membered saturated heterocyclic group

(provided that the group is attached through a ring nitrogen atom) containing one nitrogen atom and optionally containing one nitrogen atom or oxygen atom” in the definition of R³ may be, for example, a pyrrolidinyl, imidazolidinyl, pyrazolidinyl, piperidyl, piperazinyl or morpholinyl group, preferably the piperidyl or morpholinyl group.

In the formula (I), “the C₆ - C₁₄ aryl group optionally having from 1 to 3 substituents which may be the same as or different from each other and selected from the following substituent group” in the definition of R¹ may be, for example, a phenyl, indenyl, naphthyl, phenanthrenyl or anthracenyl group which may be substituted with the following substituent; preferably a C₆ - C₁₄ aryl group optionally having from 1 to 3 substituents which may be the same as or different from each other and selected from the following substituent group [the substituent group is halogen; C₁ - C₆ alkyl; C₁ - C₆ alkyl substituted with halogen or C₁ - C₆ alkoxy; C₁ - C₆ alkoxy; C₆ - C₁₄ aryl optionally having from 1 to 3 substituents which may be the same as or different from each other and selected from the following group (the substituent group is halogen, C₁ - C₆ alkyl or C₁ - C₆ alkoxy); a benzyl, fluorobenzyl, chlorobenzyl, difluorobenzyl, dichlorobenzyl, methylbenzyl, dimethylbenzyl, methoxybenzyl; phenoxy, 4-fluorophenoxy, 4-chlorophenoxy, 2,4-dichlorophenoxy, 4-methylphenoxy, 4-methoxyphenoxy; benzyloxy, 4-fluorobenzyloxy, 4-chlorobenzyloxy, 2,4-difluorobenzyloxy, 2,4-dichlorobenzyloxy, 4-methylbenzyloxy, 2,4-dimethylbenzyloxy, 4-methoxybenzyloxy; cyano; nitro; a hydroxyl group; acetoxy; C₂ - C₇ alkoxy-carbonyl; amino; carbamoyl; mono-(C₁ - C₆ alkyl)carbamoyl; di(C₁ - C₆ alkyl)carbamoyl; benzoylamino, 4-fluorobenzoylamino, 4-chlorobenzoylamino, 2,4-dichlorobenzoylamino, 4-toluoylamino or 4-anisoylamino group]; more preferably a C₆ - C₁₄ aryl group optionally having from 1 to 3 substituents which may be the same as or different from each other and selected from the following substituent group [the substituent group is a halogen; a C₁ - C₆ alkyl; a C₁ - C₆ alkyl substituted with a halogen or a C₁ - C₆ alkoxy; a

C₁ - C₆ alkoxy; a C₆ - C₁₄ aryl optionally having from 1 to 3 substituents which may be the same as or different from each other and selected from the following group (the substituent group is a halogen, a C₁ - C₆ alkyl or a C₁ - C₆ alkoxy); cyano; C₂ - C₇ alkoxy-carbonyl; carbamoyl; mono-(C₁ - C₆ alkyl)carbamoyl; or di(C₁ - C₆ alkyl)carbamoyl group]; more preferably the C₆ - C₁₄ aryl group optionally having from 1 to 3 substituents which may be the same as or different from each other and selected from the following substituent group [the substituent group is the halogen, C₁ - C₄ alkyl, fluoromethyl, difluoromethyl, trifluoromethyl, 2-fluoroethyl, 2,2,2-trifluoroethyl, methoxymethyl, methoxyethyl, C₁ - C₄ alkoxy, phenyl, 4-fluorophenyl, 4-chlorophenyl, 2,4-dichlorophenyl, 4-methylphenyl, 4-methoxyphenyl, cyano, methoxycarbonyl, ethoxycarbonyl, carbamoyl, methylcarbamoyl, ethylcarbamoyl, N,N-dimethylcarbamoyl or N,N-diethylcarbamoyl group]; still more preferably the phenyl group optionally having from 1 to 3 substituents which may be the same as or different from each other and selected from the following substituent group [the substituent group is the halogen, methyl, ethyl, trifluoromethyl, methoxy, phenyl, cyano, methoxycarbonyl, carbamoyl, methylcarbamoyl, ethylcarbamoyl or N,N-dimethylcarbamoyl group], further still more preferably the phenyl group optionally having one or two substituents which may be the same as or different from each other and selected from the following substituent group [the substituent group is the fluorine, chlorine, methyl, ethyl, trifluoromethyl and methoxy group]; particularly preferably the phenyl, fluorophenyl, chlorophenyl, difluorophenyl, dichlorophenyl or methylphenyl group; most preferably the phenyl, 2-chlorophenyl, 4-chlorophenyl, 2,4-difluorophenyl or 2,4-dichlorophenyl group.

Incidentally, the number of substituents on the aryl group is preferably 1 to 3, more preferably 1 or 2, most preferably 2.

In the formula (I), "the 5- or 6-membered aromatic heterocyclic group optionally having from 1 to 3 substituents and having one or two hetero atoms

which may be the same as or different from each other and selected from the group consisting of nitrogen, oxygen and sulfur atoms" in the definition of R¹ may be, for example, a pyrrolyl, imidazolyl, pyrazolyl, furyl, thienyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, pyridyl, pyrazinyl, pyrimidinyl or pyridazinyl group which are optionally substituted with the following substituent, preferably 5- or 6-membered aromatic heterocyclic group optionally having from 1 to 3 substituents which may be the same as or different from each other and selected from the following substituent group and having one or two hetero atoms which may be the same as or different from each other and selected from the group consisting of nitrogen, oxygen and sulfur atoms [the substituent group is halogen; C₁ - C₆ alkyl; C₁ - C₆ alkyl substituted with halogen or C₁ - C₆ alkoxy; C₁ - C₆ alkoxy; C₆ - C₁₄ aryl optionally having from 1 to 3 substituents which may be the same as or different from each other and selected from the following group (the substituent group is halogen, C₁ - C₆ alkyl or C₁ - C₆ alkoxy); benzyl, fluorobenzyl, chlorobenzyl, difluorobenzyl, dichlorobenzyl, methylbenzyl, dimethylbenzyl, methoxybenzyl; phenoxy, 4-fluorophenoxy, 4-chlorophenoxy, 2,4-dichlorophenoxy, 4-methylphenoxy, 4-methoxyphenoxy; benzyloxy, 4-fluorobenzyloxy, 4-chlorobenzyloxy, 2,4-difluorobenzyloxy, 2,4-dichlorobenzyloxy, 4-methylbenzyloxy, 2,4-dimethylbenzyloxy, 4-methoxybenzyloxy; cyano; nitro; hydroxyl; acetoxy; C₂ - C₇ alkoxycarbonyl; amino; carbamoyl; mono-(C₁ - C₆ alkyl)carbamoyl; di(C₁ - C₆ alkyl)carbamoyl; benzoylamino, 4-fluorobenzoylamino, 4-chlorobenzoylamino, 2,4-dichlorobenzoylamino, 4-toluoylamino and 4-anisoylamino group]; preferably the 5- or 6-membered aromatic heterocyclic group optionally having from 1 to 3 substituents which may be the same as or different from each other and selected from the following substituent group and having one or two hetero atoms which may be the same as or different from each other and selected from the group consisting of nitrogen, oxygen and sulfur atoms [the substituent group is the halogen; C₁ - C₆ alkyl; C₁ - C₆ alkyl substituted with halogen or C₁ - C₆ alkoxy; C₁ - C₆ alkoxy; C₆ - C₁₄ aryl optionally having from 1 to 3 substituents which may be the same as or different from each other and selected

from the following substituent group and having one or two hetero atoms which may be the same as or different from each other and selected from the group consisting of nitrogen, oxygen and sulfur atoms (the substituent group is a halogen; a C₁ - C₆ alkyl or a C₁ - C₆ alkoxy); cyano; C₂ - C₇ alkoxy-carbonyl; carbamoyl; mono-(C₁ - C₆ alkyl)carbamoyl; or di(C₁ - C₆ alkyl)carbamoyl group]; more preferably the 5- or 6-membered aromatic heterocyclic group optionally having one or two substituents which may be the same as or different from each other and selected from the following substituent group [the substituent group is the halogen, C₁ - C₄ alkyl, fluoromethyl, difluoromethyl, trifluoromethyl, 2-fluoroethyl, 2,2,2-trifluoroethyl, methoxymethyl, methoxyethyl, C₁ - C₄ alkoxy, phenyl, 4-fluorophenyl, 4-chlorophenyl, 2,4-dichlorophenyl, 4-methylphenyl, 4-methoxyphenyl, cyano, methoxycarbonyl, ethoxycarbonyl, carbamoyl, methylcarbamoyl, ethylcarbamoyl, N,N-dimethylcarbamoyl or N,N-diethylcarbamoyl group], still more preferably furyl, thienyl or pyridyl group optionally having one or two substituents which may be the same as or different from each other and selected from the following substituent group [the substituent group is the halogen, methyl, ethyl, trifluoromethyl, methoxy, phenyl, cyano, methoxycarbonyl, carbamoyl, methylcarbamoyl, ethylcarbamoyl and N,N-dimethylcarbamoyl group], further still more preferably the furyl, thienyl or pyridyl group optionally having one substituent selected from the following substituent group [the substituent group is the fluorine, chlorine, methyl, ethyl, trifluoromethyl and methoxy group], particularly preferably the 2-furyl, 3-furyl, 2-thienyl or 3-thienyl group, most preferably the 2-furyl or 2-thienyl group.

Incidentally, the number of substituents on the aromatic heterocyclic group is preferably from 1 to 3, more preferably one or two, most preferably one.

Moreover, with respect to the bond of the aromatic heterocyclic group and the isoxazole ring, the bond is preferably formed on a carbon atom on the aromatic heterocycle.

In the formula (I), X is preferably an oxygen atom.

In the formula (I), n is preferably an integer from 2 to 4, more preferably 2.

The compound (I) of the present invention can be converted to an acid addition salt by a conventional method. For example, the salt can be obtained by treating the compound (I) with a corresponding acid in a solvent (for example, ethers, esters or alcohols, particularly ethers) at room temperature for 5 to 30 minutes and collecting the precipitated crystals by filtration or removing the solvent by evaporation. Such a salt includes mineral acid salt such as hydrofluoride, hydrochloride, hydrobromide, hydroiodide, nitrate, perchlorate, sulfate and phosphate; sulfonic acid salt such as methanesulfonate, trifluoromethanesulfonate, ethanesulfonate, benzenesulfonate and p-toluenesulfonate; carboxylic acid salt such as fumarate, succinate, citrate, tartrate, oxalate and maleate; and amino acid salt such as glutamate and aspartate, preferably the mineral acid salt (more preferably the hydrochloride).

The compound (I) of the present invention sometimes has an asymmetric carbon atom in the molecule, and stereoisomers of the R-configuration and the S-configuration sometimes exist. Each of the stereoisomers or a mixture containing the isomers in an optional proportion are all included in the present invention.

The compound (I) and salt thereof occasionally absorbs moisture when they are left to stand in the atmosphere and they occasionally form hydrates when they are recrystallized. Such products containing water are also included in the present invention.

Examples of preferable compounds of formula (I) include:

(1) compounds in which R¹ is the C₆ - C₁₄ aryl group optionally having from 1

to 3 substituents which may be the same as or different from each other and selected from the following substituent group, or a 5- or 6-membered aromatic heterocyclic group optionally having from 1 to 3 substituents and having one or two hetero atoms which may be the same as or different from each other and selected from the group consisting of nitrogen, oxygen and sulfur atoms [the substituent group is the halogen; C₁ - C₆ alkyl; C₁ - C₆ alkyl substituted with halogen or C₁ - C₆ alkoxy; C₁ - C₆ alkoxy; C₆ - C₁₄ aryl optionally having from 1 to 3 substituents which may be the same as or different from each other and selected from the following group (the substituent group is halogen, C₁ - C₆ alkyl or C₁ - C₆ alkoxy); benzyl, fluorobenzyl, chlorobenzyl, difluorobenzyl, dichlorobenzyl, methylbenzyl, dimethylbenzyl, methoxybenzyl; phenoxy, 4-fluorophenoxy, 4-chlorophenoxy, 2,4-dichlorophenoxy, 4-methylphenoxy, 4-methoxyphenoxy; benzyloxy, 4-fluorobenzyloxy, 4-chlorobenzyloxy, 2,4-difluorobenzyloxy, 2,4-dichlorobenzyloxy, 4-methylbenzyloxy, 2,4-dimethylbenzyloxy, 4-methoxybenzyloxy; cyano; nitro; a hydroxyl group; acetoxy; C₂ - C₇ alkoxy carbonyl; amino; carbamoyl; mono-(C₁ - C₆ alkyl)carbamoyl; di(C₁ - C₆ alkyl)carbamoyl; benzoylamino, 4-fluorobenzoylamino, 4-chlorobenzoylamino, 2,4-dichlorobenzoylamino, 4-toluoylamino and 4-anisoylamino group],

(2) compounds in which R¹ is the C₆ - C₁₄ aryl group optionally having from 1 to 3 substituents which may be the same as or different from each other and selected from the following substituent group, or a 5- or 6-membered aromatic heterocyclic group optionally having from 1 to 3 substituents and having one or two hetero atoms which may be the same as or different from each other and selected from the group consisting of nitrogen, oxygen and sulfur atoms [the substituent group is the halogen; C₁ - C₆ alkyl; C₁ - C₆ alkyl substituted with halogen or C₁ - C₆ alkoxy; C₁ - C₆ alkoxy; C₆ - C₁₄ aryl optionally having from 1 to 3 substituents which may be the same as or different selected from the following group (the substituent group is halogen, C₁ - C₆ alkyl or C₁ - C₆

alkoxy); cyano; C₂ - C₇ alkoxy-carbonyl; carbamoyl; mono-(C₁ - C₆ alkyl)carbamoyl; and di(C₁ - C₆ alkyl)carbamoyl group],

(3) compounds in which R¹ is the C₆ - C₁₄ aryl group optionally having from 1 to 3 substituents which may be the same as or different each other and selected from the following substituent group, or a 5- or 6-membered aromatic heterocyclic group optionally having one or two substituents and having one or two hetero atoms which may be the same as or different from each other and selected from the group consisting of nitrogen, oxygen and sulfur atoms [the substituent group is the halogen, C₁ - C₄ alkyl, fluoromethyl, difluoromethyl, trifluoromethyl, 2-fluoroethyl, 2,2,2-trifluoroethyl, methoxymethyl, methoxyethyl, C₁ - C₄ alkoxy, phenyl, 4-fluorophenyl, 4-chlorophenyl, 2,4-dichlorophenyl, 4-methylphenyl, 4-methoxyphenyl, cyano, methoxycarbonyl, ethoxycarbonyl, carbamoyl, methylcarbamoyl, ethylcarbamoyl, N,N-dimethylcarbamoyl and N,N-diethylcarbamoyl group],

(4) compounds in which R¹ is the phenyl group optionally having from 1 to 3 substituents which may be the same as or different from each other and selected from the following substituent group, or the furyl, thienyl or pyridyl group optionally having one or two substituents [the substituent group is the halogen, methyl, ethyl, trifluoromethyl, methoxy, phenyl, cyano, methoxycarbonyl, carbamoyl, methylcarbamoyl, ethylcarbamoyl and N,N-dimethylcarbamoyl group],

(5) compounds in which R¹ is the phenyl group optionally having one or two substituents which may be the same as or different from each other and selected from the following substituent group, or the furyl, thienyl or pyridyl group optionally having one substituent [the substituent group is fluorine, chlorine, methyl, ethyl, trifluoromethyl and methoxy group],

(6) compounds in which R¹ is the phenyl, fluorophenyl, chlorophenyl,

difluorophenyl, dichlorophenyl, methylphenyl, 2-furyl, 3-furyl, 2-thienyl or 3-thienyl group,

(7) compounds in which R^1 is the phenyl, 2-chlorophenyl, 4-chlorophenyl, 2,4-difluorophenyl, 2,4-dichlorophenyl, 2-furyl or 2-thienyl group,

(8) compounds in which R^2 is the hydrogen, halogen, $C_1 - C_6$ alkyl, fluoro-methyl, difluoromethyl, trifluoromethyl, 2-fluoroethyl, 1-chloroethyl, 2-chloroethyl, 2,2,2-trifluoroethyl, methoxymethyl, methoxyethyl, $C_2 - C_6$ alkenyl, $C_2 - C_6$ alkynyl, cyclopropyl, cyclopentyl, cyclohexyl, 2-cyclopentenyl, 3-cyclopentenyl, 2-cyclohexenyl, 3-cyclohexenyl, methoxy, ethoxy, cyano, carboxyl, formyl, acetyl, methoxycarbonyl, ethoxycarbonyl, carbamoyl, methylcarbamoyl, ethylcarbamoyl or N,N-dimethylcarbamoyl group,

(9) compounds in which R^2 is the hydrogen, halogen, $C_1 - C_6$ alkyl, $C_2 - C_6$ alkenyl or $C_2 - C_6$ alkynyl group,

(10) compounds in which R^2 is the hydrogen, halogen, $C_1 - C_4$ alkyl, allyl, isopropenyl, 2-butenyl or propargyl group,

(11) compounds in which R^2 is the hydrogen, chlorine, ethyl, propyl, isopropyl, isobutyl or t-butyl group,

(12) compounds in which R^2 is the hydrogen or isopropyl group,

(13) compounds in which R^3 is the amino, mono $C_1 - C_6$ alkylamino, di($C_1 - C_6$ alkyl)amino or 5- or 6-membered saturated heterocyclic group (provided that the group is attached thorough a ring nitrogen atom), having one nitrogen atom and further optionally having one nitrogen atom or oxygen atom

(14) compounds in which R^3 is the amino, methylamino, ethylamino, N,N-dimethylamino, piperidyl or morpholinyl group,

(15) compounds in which R^3 is the amino group,

(16) compounds in which X is the oxygen atom, and

(17) compounds in which n is 2.

The order of preference of R^1 increases in the ascending orders of (1) to (7), that of R^2 increases in the ascending orders of (8) to (12), and that of R^3 increases in the ascending orders of (13) to (15).

Further, compounds of formula (I) include combinations of from two to five selected from the group consisting of (1)-(7), (8)-(12), (13)-(15), (16) and (17) and preferable examples of such combinations are shown below,

(18) compounds in which R^1 is the $C_6 - C_{14}$ aryl group optionally having from 1 to 3 substituents which may be the same as or different from each other and selected from the following substituent group, or the 5- or 6-membered aromatic heterocyclic group optionally having from 1 to 3 substituents and having one or two hetero atoms which may be the same as or different from each other and selected from the group consisting of nitrogen, oxygen and sulfur atoms [the substituent group is the halogen; $C_1 - C_6$ alkyl; $C_1 - C_6$ alkyl substituted with halogen or $C_1 - C_6$ alkoxy; $C_1 - C_6$ alkoxy; $C_6 - C_{14}$ aryl optionally having from 1 to 3 substituents which may be the same as or different from each other and selected from the following group (the substituent group is halogen, $C_1 - C_6$ alkyl or $C_1 - C_6$ alkoxy); cyano; $C_2 - C_7$ alkoxy-carbonyl; carbamoyl; mono-($C_1 - C_6$ alkyl)carbamoyl; and di($C_1 - C_6$ alkyl)carbamoyl group],

R^2 is the hydrogen, halogen, $C_1 - C_6$ alkyl, $C_2 - C_6$ alkenyl or $C_2 - C_6$ alkynyl group, and

R^3 is the amino, mono $C_1 - C_6$ alkylamino, di($C_1 - C_6$ alkyl)amino group or the 5- or 6-membered saturated heterocyclic group (provided that the group attached thorough a ring nitrogen atom) containing one nitrogen atom and further one nitrogen or oxygen atom,

(19) compounds in which R^1 is the $C_6 - C_{14}$ aryl group optionally having from 1 to 3 substituents which may be the same as or different from each other and selected from the following substituent group, or the 5- or 6-membered aromatic heterocyclic group optionally having from 1 to 3 substituents and having one or two hetero atoms which may be the same as or different from each other and selected from the group consisting of nitrogen, oxygen and sulfur atoms [the substituent group is the halogen; $C_1 - C_6$ alkyl; $C_1 - C_6$ alkyl substituted with halogen or $C_1 - C_6$ alkoxy; $C_1 - C_6$ alkoxy; $C_6 - C_{14}$ aryl optionally having from 1 to 3 substituents which may be the same as or different from each other and selected from the following group (the substituent group is halogen, $C_1 - C_6$ alkyl or $C_1 - C_6$ alkoxy); cyano; $C_2 - C_7$ alkoxy carbonyl; carbamoyl; mono($C_1 - C_6$ alkyl)carbamoyl; and di($C_1 - C_6$ alkyl)carbamoyl group],

R^2 is the hydrogen, halogen, $C_1 - C_6$ alkyl, $C_2 - C_6$ alkenyl or $C_2 - C_6$ alkynyl group,

R^3 is the amino group,

X is the oxygen atom, and

n is 2,

(20) compounds in which R^1 is the $C_6 - C_{14}$ aryl group optionally having from 1 to 3 substituents which may be the same as or different from each other and selected from the following substituent group, or the 5- or 6-membered aromatic heterocyclic group optionally having one or two substituents and having one or two hetero atoms which may be the same as or different from

each other and selected from the group consisting of nitrogen, oxygen and sulfur atoms [the substituent group is the halogen, C₁ - C₄ alkyl, fluoromethyl, difluoromethyl, trifluoromethyl, 2-fluoroethyl, 2,2,2-trifluoroethyl, methoxymethyl, methoxyethyl, C₁ - C₄ alkoxy, phenyl, 4-fluorophenyl, 4-chlorophenyl, 2,4-dichlorophenyl, 4-methylphenyl, 4-methoxyphenyl, cyano, methoxycarbonyl, ethoxycarbonyl, carbamoyl, methylcarbamoyl, ethylcarbamoyl, N,N-dimethylcarbamoyl and N,N-diethylcarbamoyl group],

R² is the hydrogen, halogen, C₁ - C₄ alkyl, allyl, isopropenyl, 2-butenyl or propargyl group,

R³ is the amino group,

X is the oxygen atom, and

n is 2,

(21) compounds in which R¹ is the phenyl group optionally having from 1 to 3 substituents which may be the same as or different from each other and selected from the following substituent group, or the furyl, thienyl or pyridyl group optionally having one or two substituents [the substituent group is the halogen, methyl, ethyl, trifluoromethyl, methoxy, phenyl, cyano, methoxycarbonyl, carbamoyl, methylcarbamoyl, ethylcarbamoyl and N,N-dimethylcarbamoyl group],

R² is the hydrogen, halogen, C₁ - C₄ alkyl, allyl, isopropenyl, 2-butenyl or propargyl group,

R³ is the amino group,

X is the oxygen atom, and

n is 2,

(22) compounds in which R¹ is the phenyl group optionally having one or two substituents which may be the same as or different from each other and selected from the following substituent group, or the furyl, thienyl or pyridyl group optionally having one substituent [the substituent group is the fluorine, chlorine,

methyl, ethyl, trifluoromethyl and methoxy group],

R^2 is the hydrogen, chlorine, ethyl, propyl, isopropyl, isobutyl or t-butyl group,

R^3 is the amino group,

X is the oxygen atom, and

n is 2,

(23) compounds in which R^1 is the fluorophenyl, chlorophenyl, difluorophenyl, dichlorophenyl, methylphenyl, 2-furyl, 3-furyl, 2-thienyl or 3-thienyl group,

R^2 is the hydrogen, chlorine, ethyl, propyl, isopropyl, isobutyl or t-butyl group,

R^3 is the amino group,

X is the oxygen atom, and

n is 2,

(24) compounds in which R^1 is the phenyl, 2-chlorophenyl, 4-chlorophenyl, 2,4-difluorophenyl, 2,4-dichlorophenyl, 2-furyl or 2-thienyl group,

R^2 is the hydrogen atom or the isopropyl group,

R^3 is the amino group,

X is the oxygen atom, and

n is 2.

With respect to the compounds described above, the order of preference of the compounds increases in the ascending orders of (18) to (24).


The representative compounds of the present invention are illustrated in the following Table, but the present invention is not limited to them.

In the Table the following abbreviations are used.

Ac	:	Acetyl
All	:	Allyl
Bn	:	Benzyl
Bu	:	Butyl

Bu ⁱ	:	Isobutyl
Bu ^s	:	s-Butyl
Bu ^t	:	t-Butyl
Bun(2)	:	2-Butenyl
Bz	:	Benzoyl
Et	:	Ethyl
Fur(2)	:	2-Furyl
Hex	:	Hexyl
Imid(2)	:	2-Imidazolyl
Inde(1)	:	1-Indenyl
Isothiz(3)	:	3-Isothiazolyl
Isox(3)	:	3-Isoxazolyl
Me	:	Methyl
Moc	:	Methoxycarbonyl
Mor(4)	:	4-Morpholinyl
Np(1)	:	1-Naphthyl
Np(2)	:	2-Naphthyl
Oxa(2)	:	2-Oxazolyl
Pen ^c (2)	:	2-Cyclopentenyl
Ph	:	Phenyl
Pip(1)	:	1-Piperidyl
Piz(1)	:	1-Piperazinyl
Pn	:	Pentyl
Pn ^c	:	Cyclopentyl
Pn ⁱ	:	Isopentyl
Pr	:	Propyl
Pr ^c	:	Cyclopropyl
Pr ⁱ	:	Isopropyl
Pre ⁱ	:	Isopropenyl
Prg	:	Propargyl
Pym(2)	:	2-Pyrimidinyl

Pyr(2)	:	2-Pyridyl
Pyr(3)	:	3-Pyridyl
Pyr(4)	:	4-Pyridyl
Pyrd(1)	:	1-Pyrrolidinyl
Pyrr(3)	:	3-Pyrrolyl
Pyz(2)	:	2-Pyrazinyl
Pyza(1)	:	1-Pyrazolyl
Pyzn(3)	:	3-Pyridazinyl
Thi(2)	:	2-Thienyl
Thi(3)	:	3-Thienyl
Thiz(2)	:	2-Thiazolyl



[Table 1]

Compd. No.	R ¹	R ²	R ³	X	n
1	Ph	H	NH ₂	O	2
2	Ph	H	NH ₂	O	3
3	Ph	H	NH ₂	O	4
4	Ph	F	NH ₂	O	2
5	Ph	Cl	NH ₂	O	2
6	Ph	Me	NH ₂	O	2
7	Ph	Et	NH ₂	O	2
8	Ph	Pr	NH ₂	O	2
9	Ph	Pr ⁱ	NH ₂	O	2

- 28 -

10	Ph	Bu	NH ₂	O	2
11	Ph	Bu ⁱ	NH ₂	O	2
12	Ph	Bu ^s	NH ₂	O	2
13	Ph	Bu ^t	NH ₂	O	2
14	Ph	CF ₃	NH ₂	O	2
15	Ph	H	NH ₂	S	2
16	Ph	H	NH ₂	S	3
17	Ph	H	NH ₂	S	4
18	Ph	F	NH ₂	S	2
19	Ph	Cl	NH ₂	S	2
20	Ph	Me	NH ₂	S	2
21	Ph	Et	NH ₂	S	2
22	Ph	Pr	NH ₂	S	2
23	Ph	Pr ⁱ	NH ₂	S	2
24	Ph	Bu	NH ₂	S	2
25	Ph	H	NHMe	O	2
26	Ph	H	NHEt	O	2
27	Ph	H	N(Me) ₂	O	2
28	Ph	H	Pip(1)	O	2
29	Ph	H	Mor(4)	O	2
30	2-F-Ph	H	NH ₂	O	2
31	2-F-Ph	F	NH ₂	O	2
32	2-F-Ph	Cl	NH ₂	O	2
33	2-F-Ph	Me	NH ₂	O	2
34	2-F-Ph	Et	NH ₂	O	2
35	2-F-Ph	Pr	NH ₂	O	2
36	2-F-Ph	Pr ⁱ	NH ₂	O	2
37	2-F-Ph	Bu	NH ₂	O	2

- 29 -

38	2-F-Ph	Bu ⁱ	NH ₂	O	2
39	2-F-Ph	Bu ^s	NH ₂	O	2
40	2-F-Ph	Bu ^t	NH ₂	O	2
41	2-F-Ph	H	NH ₂	S	2
42	2-F-Ph	F	NH ₂	S	2
43	2-F-Ph	Cl	NH ₂	S	2
44	2-F-Ph	Me	NH ₂	S	2
45	2-F-Ph	Et	NH ₂	S	2
46	2-F-Ph	Pr	NH ₂	S	2
47	2-F-Ph	Pr ⁱ	NH ₂	S	2
48	3-F-Ph	H	NH ₂	O	2
49	3-F-Ph	F	NH ₂	O	2
50	3-F-Ph	Cl	NH ₂	O	2
51	3-F-Ph	Me	NH ₂	O	2
52	3-F-Ph	Et	NH ₂	O	2
53	3-F-Ph	Pr	NH ₂	O	2
54	3-F-Ph	Pr ⁱ	NH ₂	O	2
55	3-F-Ph	Bu	NH ₂	O	2
56	3-F-Ph	Bu ⁱ	NH ₂	O	2
57	3-F-Ph	Bu ^s	NH ₂	O	2
58	3-F-Ph	Bu ^t	NH ₂	O	2
59	3-F-Ph	H	NH ₂	S	2
60	3-F-Ph	F	NH ₂	S	2
61	3-F-Ph	Cl	NH ₂	S	2
62	3-F-Ph	Me	NH ₂	S	2
63	3-F-Ph	Et	NH ₂	S	2
64	3-F-Ph	Pr	NH ₂	S	2
65	3-F-Ph	Pr ⁱ	NH ₂	S	2

- 30 -

66	4-F-Ph	H	NH ₂	O	2
67	4-F-Ph	H	NH ₂	O	3
68	4-F-Ph	H	NH ₂	O	4
69	4-F-Ph	F	NH ₂	O	2
70	4-F-Ph	Cl	NH ₂	O	2
71	4-F-Ph	Me	NH ₂	O	2
72	4-F-Ph	Et	NH ₂	O	2
73	4-F-Ph	Pr	NH ₂	O	2
74	4-F-Ph	Pr ⁱ	NH ₂	O	2
75	4-F-Ph	Bu	NH ₂	O	2
76	4-F-Ph	Bu ⁱ	NH ₂	O	2
77	4-F-Ph	Bu ^s	NH ₂	O	2
78	4-F-Ph	Bu ^t	NH ₂	O	2
79	4-F-Ph	H	NH ₂	S	2
80	4-F-Ph	H	NH ₂	S	3
81	4-F-Ph	H	NH ₂	S	4
82	4-F-Ph	F	NH ₂	S	2
83	4-F-Ph	Cl	NH ₂	S	2
84	4-F-Ph	Me	NH ₂	S	2
85	4-F-Ph	Et	NH ₂	S	2
86	4-F-Ph	Pr	NH ₂	S	2
87	4-F-Ph	Pr ⁱ	NH ₂	S	2
88	4-F-Ph	H	NHMe	O	2
89	4-F-Ph	H	NHEt	O	2
90	4-F-Ph	H	N(Me) ₂	O	2
91	4-F-Ph	H	Pip(1)	O	2
92	4-F-Ph	H	Mor(4)	O	2
93	2,4-diF-Ph	H	NH ₂	O	2

- 31 -

94	2,4-diF-Ph	F	NH ₂	O	2
95	2,4-diF-Ph	Cl	NH ₂	O	2
96	2,4-diF-Ph	Me	NH ₂	O	2
97	2,4-diF-Ph	Et	NH ₂	O	2
98	2,4-diF-Ph	Pr	NH ₂	O	2
99	2,4-diF-Ph	Pr ⁱ	NH ₂	O	2
100	2,4-diF-Ph	Bu	NH ₂	O	2
101	2,4-diF-Ph	Bu ⁱ	NH ₂	O	2
102	2,4-diF-Ph	Bu ^s	NH ₂	O	2
103	2,4-diF-Ph	Bu ^t	NH ₂	O	2
104	2,4-diF-Ph	H	NH ₂	S	2
105	2,4-diF-Ph	F	NH ₂	S	2
106	2,4-diF-Ph	Cl	NH ₂	S	2
107	2,4-diF-Ph	Me	NH ₂	S	2
108	2,4-diF-Ph	Et	NH ₂	S	2
109	2,4-diF-Ph	Pr	NH ₂	S	2
110	2,4-diF-Ph	Pr ⁱ	NH ₂	S	2
111	2-Cl-Ph	H	NH ₂	O	2
112	2-Cl-Ph	F	NH ₂	O	2
113	2-Cl-Ph	Cl	NH ₂	O	2
114	2-Cl-Ph	Me	NH ₂	O	2
115	2-Cl-Ph	Et	NH ₂	O	2
116	2-Cl-Ph	Pr	NH ₂	O	2
117	2-Cl-Ph	Pr ⁱ	NH ₂	O	2
118	2-Cl-Ph	H	NH ₂	S	2
119	2-Cl-Ph	F	NH ₂	S	2
120	2-Cl-Ph	Cl	NH ₂	S	2
121	2-Cl-Ph	Me	NH ₂	S	2

- 32 -

122	2-Cl-Ph	Et	NH ₂	S	2
123	2-Cl-Ph	Pr	NH ₂	S	2
124	2-Cl-Ph	Pr ⁱ	NH ₂	S	2
125	3-Cl-Ph	H	NH ₂	O	2
126	3-Cl-Ph	F	NH ₂	O	2
127	3-Cl-Ph	Cl	NH ₂	O	2
128	3-Cl-Ph	Me	NH ₂	O	2
129	3-Cl-Ph	Et	NH ₂	O	2
130	3-Cl-Ph	Pr	NH ₂	O	2
131	3-Cl-Ph	Pr ⁱ	NH ₂	O	2
132	3-Cl-Ph	Bu	NH ₂	O	2
133	3-Cl-Ph	Bu ⁱ	NH ₂	O	2
134	3-Cl-Ph	Bu ^s	NH ₂	O	2
135	3-Cl-Ph	Bu ^t	NH ₂	O	2
136	3-Cl-Ph	H	NH ₂	S	2
137	3-Cl-Ph	F	NH ₂	S	2
138	3-Cl-Ph	Cl	NH ₂	S	2
139	3-Cl-Ph	Me	NH ₂	S	2
140	3-Cl-Ph	Et	NH ₂	S	2
141	3-Cl-Ph	Pr	NH ₂	S	2
142	3-Cl-Ph	Pr ⁱ	NH ₂	S	2
143	4-Cl-Ph	H	NH ₂	O	2
144	4-Cl-Ph	H	NH ₂	O	3
145	4-Cl-Ph	H	NH ₂	O	4
146	4-Cl-Ph	F	NH ₂	O	2
147	4-Cl-Ph	Cl	NH ₂	O	2
148	4-Cl-Ph	Me	NH ₂	O	2
149	4-Cl-Ph	Et	NH ₂	O	2

- 33 -

150	4-Cl-Ph	Pr	NH ₂	O	2
151	4-Cl-Ph	Pr ⁱ	NH ₂	O	2
152	4-Cl-Ph	Bu	NH ₂	O	2
153	4-Cl-Ph	Bu ⁱ	NH ₂	O	2
154	4-Cl-Ph	Bu ^s	NH ₂	O	2
155	4-Cl-Ph	Bu ^t	NH ₂	O	2
156	4-Cl-Ph	H	NH ₂	S	2
157	4-Cl-Ph	H	NH ₂	S	3
158	4-Cl-Ph	H	NH ₂	S	4
159	4-Cl-Ph	F	NH ₂	S	2
160	4-Cl-Ph	Cl	NH ₂	S	2
161	4-Cl-Ph	Me	NH ₂	S	2
162	4-Cl-Ph	Et	NH ₂	S	2
163	4-Cl-Ph	Pr	NH ₂	S	2
164	4-Cl-Ph	Pr ⁱ	NH ₂	S	2
165	4-Cl-Ph	H	NHMe	O	2
166	4-Cl-Ph	H	NHEt	O	2
167	4-Cl-Ph	H	N(Me) ₂	O	2
168	4-Cl-Ph	H	Pip(1)	O	2
169	4-Cl-Ph	H	Mor(4)	O	2
170	2,4-diCl-Ph	H	NH ₂	O	2
171	2,4-diCl-Ph	F	NH ₂	O	2
172	2,4-diCl-Ph	Cl	NH ₂	O	2
173	2,4-diCl-Ph	Me	NH ₂	O	2
174	2,4-diCl-Ph	Et	NH ₂	O	2
175	2,4-diCl-Ph	Pr	NH ₂	O	2
176	2,4-diCl-Ph	Pr ⁱ	NH ₂	O	2
177	2,4-diCl-Ph	Bu	NH ₂	O	2

- 34 -

178	2,4-diCl-Ph	Bu ⁱ	NH ₂	O	2
179	2,4-diCl-Ph	Bu ^s	NH ₂	O	2
180	2,4-diCl-Ph	Bu ^t	NH ₂	O	2
181	2,4-diCl-Ph	H	NH ₂	S	2
182	2,4-diCl-Ph	F	NH ₂	S	2
183	2,4-diCl-Ph	Cl	NH ₂	S	2
184	2,4-diCl-Ph	Me	NH ₂	S	2
185	2,4-diCl-Ph	Et	NH ₂	S	2
186	2,4-diCl-Ph	Pr	NH ₂	S	2
187	2,4-diCl-Ph	Pr ⁱ	NH ₂	S	2
188	2,6-diCl-Ph	H	NH ₂	O	2
189	2,6-diCl-Ph	F	NH ₂	O	2
190	2,6-diCl-Ph	Cl	NH ₂	O	2
191	2,6-diCl-Ph	Me	NH ₂	O	2
192	2,6-diCl-Ph	Et	NH ₂	O	2
193	2,6-diCl-Ph	Pr	NH ₂	O	2
194	2,6-diCl-Ph	Pr ⁱ	NH ₂	O	2
195	2,6-diCl-Ph	Bu	NH ₂	O	2
196	2,6-diCl-Ph	Bu ⁱ	NH ₂	O	2
197	2,6-diCl-Ph	Bu ^s	NH ₂	O	2
198	2,6-diCl-Ph	Bu ^t	NH ₂	O	2
199	2,6-diCl-Ph	H	NH ₂	S	2
200	2,6-diCl-Ph	F	NH ₂	S	2
201	2,6-diCl-Ph	Cl	NH ₂	S	2
202	2,6-diCl-Ph	Me	NH ₂	S	2
203	2,6-diCl-Ph	Et	NH ₂	S	2
204	2,6-diCl-Ph	Pr	NH ₂	S	2
205	2,6-diCl-Ph	Pr ⁱ	NH ₂	S	2

- 35 -

206	3,5-diCl-Ph	H	NH ₂	O	2
207	3,5-diCl-Ph	F	NH ₂	O	2
208	3,5-diCl-Ph	Cl	NH ₂	O	2
209	3,5-diCl-Ph	Me	NH ₂	O	2
210	3,5-diCl-Ph	Et	NH ₂	O	2
211	3,5-diCl-Ph	Pr	NH ₂	O	2
212	3,5-diCl-Ph	Pr ⁱ	NH ₂	O	2
213	3,5-diCl-Ph	Bu	NH ₂	O	2
214	3,5-diCl-Ph	Bu ⁱ	NH ₂	O	2
215	3,5-diCl-Ph	Bu ^s	NH ₂	O	2
216	3,5-diCl-Ph	Bu ^t	NH ₂	O	2
217	3,5-diCl-Ph	H	NH ₂	S	2
218	3,5-diCl-Ph	F	NH ₂	S	2
219	3,5-diCl-Ph	Cl	NH ₂	S	2
220	3,5-diCl-Ph	Me	NH ₂	S	2
221	3,5-diCl-Ph	Et	NH ₂	S	2
222	3,5-diCl-Ph	Pr	NH ₂	S	2
223	3,5-diCl-Ph	Pr ⁱ	NH ₂	S	2
224	2-Me-Ph	H	NH ₂	O	2
225	2-Me-Ph	F	NH ₂	O	2
226	2-Me-Ph	Cl	NH ₂	O	2
227	2-Me-Ph	Me	NH ₂	O	2
228	2-Me-Ph	Et	NH ₂	O	2
229	2-Me-Ph	Pr	NH ₂	O	2
230	2-Me-Ph	Pr ⁱ	NH ₂	O	2
231	2-Me-Ph	Bu	NH ₂	O	2
232	2-Me-Ph	Bu ⁱ	NH ₂	O	2
233	2-Me-Ph	Bu ^s	NH ₂	O	2

- 36 -

234	2-Me-Ph	Bu ^t	NH ₂	O	2
235	2-Me-Ph	H	NH ₂	S	2
236	2-Me-Ph	F	NH ₂	S	2
237	2-Me-Ph	Cl	NH ₂	S	2
238	2-Me-Ph	Me	NH ₂	S	2
239	2-Me-Ph	Et	NH ₂	S	2
240	2-Me-Ph	Pr	NH ₂	S	2
241	2-Me-Ph	Pr ⁱ	NH ₂	S	2
242	3-Me-Ph	H	NH ₂	O	2
243	3-Me-Ph	F	NH ₂	O	2
244	3-Me-Ph	Cl	NH ₂	O	2
245	3-Me-Ph	Me	NH ₂	O	2
246	3-Me-Ph	Et	NH ₂	O	2
247	3-Me-Ph	Pr	NH ₂	O	2
248	3-Me-Ph	Pr ⁱ	NH ₂	O	2
249	3-Me-Ph	Bu	NH ₂	O	2
250	3-Me-Ph	Bu ⁱ	NH ₂	O	2
251	3-Me-Ph	Bu ^s	NH ₂	O	2
252	3-Me-Ph	Bu ^t	NH ₂	O	2
253	3-Me-Ph	H	NH ₂	S	2
254	3-Me-Ph	F	NH ₂	S	2
255	3-Me-Ph	Cl	NH ₂	S	2
256	3-Me-Ph	Me	NH ₂	S	2
257	3-Me-Ph	Et	NH ₂	S	2
258	3-Me-Ph	Pr	NH ₂	S	2
259	3-Me-Ph	Pr ⁱ	NH ₂	S	2
260	4-Me-Ph	H	NH ₂	O	2
261	4-Me-Ph	F	NH ₂	O	2

- 37 -

262	4-Me-Ph	Cl	NH ₂	O	2
263	4-Me-Ph	Me	NH ₂	O	2
264	4-Me-Ph	Et	NH ₂	O	2
265	4-Me-Ph	Pr	NH ₂	O	2
266	4-Me-Ph	Pr ⁱ	NH ₂	O	2
267	4-Me-Ph	Bu	NH ₂	O	2
268	4-Me-Ph	Bu ⁱ	NH ₂	O	2
269	4-Me-Ph	Bu ^s	NH ₂	O	2
270	4-Me-Ph	Bu ^t	NH ₂	O	2
271	4-Me-Ph	H	NH ₂	S	2
272	4-Me-Ph	F	NH ₂	S	2
273	4-Me-Ph	Cl	NH ₂	S	2
274	4-Me-Ph	Me	NH ₂	S	2
275	4-Me-Ph	Et	NH ₂	S	2
276	4-Me-Ph	Pr	NH ₂	S	2
277	4-Me-Ph	Pr ⁱ	NH ₂	S	2
278	4-Et-Ph	H	NH ₂	O	2
279	4-Et-Ph	F	NH ₂	O	2
280	4-Et-Ph	Cl	NH ₂	O	2
281	4-Et-Ph	Me	NH ₂	O	2
282	4-Et-Ph	Et	NH ₂	O	2
283	4-Et-Ph	Pr	NH ₂	O	2
284	4-Et-Ph	Pr ⁱ	NH ₂	O	2
285	4-Et-Ph	Bu	NH ₂	O	2
286	4-Et-Ph	Bu ⁱ	NH ₂	O	2
287	4-Et-Ph	Bu ^s	NH ₂	O	2
288	4-Et-Ph	Bu ^t	NH ₂	O	2
289	4-Et-Ph	H	NH ₂	S	2

- 38 -

290	4-Et-Ph	F	NH ₂	S	2
291	4-Et-Ph	Cl	NH ₂	S	2
292	4-Et-Ph	Me	NH ₂	S	2
293	4-Et-Ph	Et	NH ₂	S	2
294	4-Et-Ph	Pr	NH ₂	S	2
295	4-Et-Ph	Pr ⁱ	NH ₂	S	2
296	2-CF ₃ -Ph	H	NH ₂	O	2
297	2-CF ₃ -Ph	F	NH ₂	O	2
298	2-CF ₃ -Ph	Cl	NH ₂	O	2
299	2-CF ₃ -Ph	Me	NH ₂	O	2
300	2-CF ₃ -Ph	Et	NH ₂	O	2
301	2-CF ₃ -Ph	Pr	NH ₂	O	2
302	2-CF ₃ -Ph	Pr ⁱ	NH ₂	O	2
303	2-CF ₃ -Ph	Bu	NH ₂	O	2
304	2-CF ₃ -Ph	Bu ⁱ	NH ₂	O	2
305	2-CF ₃ -Ph	Bu ^s	NH ₂	O	2
306	2-CF ₃ -Ph	Bu ^t	NH ₂	O	2
307	2-CF ₃ -Ph	H	NH ₂	S	2
308	2-CF ₃ -Ph	F	NH ₂	S	2
309	2-CF ₃ -Ph	Cl	NH ₂	S	2
310	2-CF ₃ -Ph	Me	NH ₂	S	2
311	2-CF ₃ -Ph	Et	NH ₂	S	2
312	2-CF ₃ -Ph	Pr	NH ₂	S	2
313	2-CF ₃ -Ph	Pr ⁱ	NH ₂	S	2
314	3-CF ₃ -Ph	H	NH ₂	O	2
315	3-CF ₃ -Ph	F	NH ₂	O	2
316	3-CF ₃ -Ph	Cl	NH ₂	O	2
317	3-CF ₃ -Ph	Me	NH ₂	O	2

- 39 -

318	3-CF ₃ -Ph	Et	NH ₂	O	2
319	3-CF ₃ -Ph	Pr	NH ₂	O	2
320	3-CF ₃ -Ph	Pr ⁱ	NH ₂	O	2
321	3-CF ₃ -Ph	Bu	NH ₂	O	2
322	3-CF ₃ -Ph	Bu ⁱ	NH ₂	O	2
323	3-CF ₃ -Ph	Bu ^s	NH ₂	O	2
324	3-CF ₃ -Ph	Bu ^t	NH ₂	O	2
325	3-CF ₃ -Ph	H	NH ₂	S	2
326	3-CF ₃ -Ph	F	NH ₂	S	2
327	3-CF ₃ -Ph	Cl	NH ₂	S	2
328	3-CF ₃ -Ph	Me	NH ₂	S	2
329	3-CF ₃ -Ph	Et	NH ₂	S	2
330	3-CF ₃ -Ph	Pr	NH ₂	S	2
331	3-CF ₃ -Ph	Pr ⁱ	NH ₂	S	2
332	4-CF ₃ -Ph	H	NH ₂	O	2
333	4-CF ₃ -Ph	F	NH ₂	O	2
334	4-CF ₃ -Ph	Cl	NH ₂	O	2
335	4-CF ₃ -Ph	Me	NH ₂	O	2
336	4-CF ₃ -Ph	Et	NH ₂	O	2
337	4-CF ₃ -Ph	Pr	NH ₂	O	2
338	4-CF ₃ -Ph	Pr ⁱ	NH ₂	O	2
339	4-CF ₃ -Ph	Bu	NH ₂	O	2
340	4-CF ₃ -Ph	Bu ⁱ	NH ₂	O	2
341	4-CF ₃ -Ph	Bu ^s	NH ₂	O	2
342	4-CF ₃ -Ph	Bu ^t	NH ₂	O	2
343	4-CF ₃ -Ph	H	NH ₂	S	2
344	4-CF ₃ -Ph	F	NH ₂	S	2
345	4-CF ₃ -Ph	Cl	NH ₂	S	2

- 40 -

346	4-CF ₃ -Ph	Me	NH ₂	S	2
347	4-CF ₃ -Ph	Et	NH ₂	S	2
348	4-CF ₃ -Ph	Pr	NH ₂	S	2
349	4-CF ₃ -Ph	Pr ⁱ	NH ₂	S	2
350	4-MeO-Ph	H	NH ₂	O	2
351	4-MeO-Ph	F	NH ₂	O	2
352	4-MeO-Ph	Cl	NH ₂	O	2
353	4-MeO-Ph	Me	NH ₂	O	2
354	4-MeO-Ph	Et	NH ₂	O	2
355	4-MeO-Ph	Pr	NH ₂	O	2
356	4-MeO-Ph	Pr ⁱ	NH ₂	O	2
357	2-MeO-Ph	H	NH ₂	O	2
358	2-MeO-Ph	F	NH ₂	O	2
359	2-MeO-Ph	Cl	NH ₂	O	2
360	2-MeO-Ph	Me	NH ₂	O	2
361	2-MeO-Ph	Et	NH ₂	O	2
362	2-MeO-Ph	Pr ⁱ	NH ₂	O	2
363	3-MeO-Ph	H	NH ₂	O	2
364	3-MeO-Ph	Cl	NH ₂	O	2
365	3-MeO-Ph	Me	NH ₂	O	2
366	3-MeO-Ph	Et	NH ₂	O	2
367	3-MeO-Ph	Pr ⁱ	NH ₂	O	2
368	4-Ph-Ph	H	NH ₂	O	2
369	4-Ph-Ph	F	NH ₂	O	2
370	4-Ph-Ph	Cl	NH ₂	O	2
371	4-Ph-Ph	Me	NH ₂	O	2
372	4-Ph-Ph	Et	NH ₂	O	2
373	4-Ph-Ph	Pr	NH ₂	O	2

- 41 -

374	4-Ph-Ph	Pr ⁱ	NH ₂	O	2
375	4-Ph-Ph	Bu	NH ₂	O	2
376	4-Ph-Ph	Bu ⁱ	NH ₂	O	2
377	4-Ph-Ph	Bu ^s	NH ₂	O	2
378	4-Ph-Ph	Bu ^t	NH ₂	O	2
379	2-Ph-Ph	H	NH ₂	S	2
380	3-Ph-Ph	F	NH ₂	S	2
381	4-Ph-Ph	Cl	NH ₂	S	2
382	2-Ph-Ph	Me	NH ₂	S	2
383	2-Ph-Ph	Et	NH ₂	S	2
384	2-Ph-Ph	Pr	NH ₂	S	2
385	3-Ph-Ph	Pr ⁱ	NH ₂	S	2
386	4-CN-Ph	H	NH ₂	O	2
387	4-CN-Ph	F	NH ₂	O	2
388	4-CN-Ph	Cl	NH ₂	O	2
389	4-CN-Ph	Me	NH ₂	O	2
390	4-CN-Ph	Et	NH ₂	O	2
391	4-CN-Ph	Pr	NH ₂	O	2
392	4-CN-Ph	Pr ⁱ	NH ₂	O	2
393	4-CN-Ph	Bu	NH ₂	O	2
394	4-CN-Ph	Bu ⁱ	NH ₂	O	2
395	4-CN-Ph	Bu ^s	NH ₂	O	2
396	4-CN-Ph	Bu ^t	NH ₂	O	2
397	4-MeOCO-Ph	H	NH ₂	O	2
398	4-MeOCO-Ph	F	NH ₂	O	2
399	4-MeOCO-Ph	Cl	NH ₂	O	2
400	4-MeOCO-Ph	Me	NH ₂	O	2
401	4-MeOCO-Ph	Et	NH ₂	O	2

- 42 -

402	4-MeOCO-Ph	Pr	NH ₂	O	2
403	4-MeOCO-Ph	Pr ⁱ	NH ₂	O	2
404	4-MeOCO-Ph	Bu	NH ₂	O	2
405	4-MeOCO-Ph	Bu ⁱ	NH ₂	O	2
406	4-MeOCO-Ph	Bu ^s	NH ₂	O	2
407	4-MeOCO-Ph	Bu ^t	NH ₂	O	2
408	2-MeOCO-Ph	H	NH ₂	S	2
409	3-MeOCO-Ph	F	NH ₂	S	2
410	4-MeOCO-Ph	Cl	NH ₂	S	2
411	2-MeOCO-Ph	Me	NH ₂	S	2
412	2-MeOCO-Ph	Et	NH ₂	S	2
413	2-MeOCO-Ph	Pr	NH ₂	S	2
414	3-MeOCO-Ph	Pr ⁱ	NH ₂	S	2
415	4-H ₂ NCO-Ph	H	NH ₂	O	2
416	4-H ₂ NCO-Ph	F	NH ₂	O	2
417	4-H ₂ NCO-Ph	Cl	NH ₂	O	2
418	4-H ₂ NCO-Ph	Me	NH ₂	O	2
419	4-H ₂ NCO-Ph	Et	NH ₂	O	2
420	4-H ₂ NCO-Ph	Pr	NH ₂	O	2
421	4-H ₂ NCO-Ph	Pr ⁱ	NH ₂	O	2
422	4-H ₂ NCO-Ph	Bu	NH ₂	O	2
423	4-H ₂ NCO-Ph	Bu ⁱ	NH ₂	O	2
424	4-H ₂ NCO-Ph	Bu ^s	NH ₂	O	2
425	4-H ₂ NCO-Ph	Bu ^t	NH ₂	O	2
426	2-H ₂ NCO-Ph	H	NH ₂	S	2
427	3-H ₂ NCO-Ph	F	NH ₂	S	2
428	4-H ₂ NCO-Ph	Cl	NH ₂	S	2
429	2-H ₂ NCO-Ph	Me	NH ₂	S	2

- 43 -

430	2-H ₂ NCO-Ph	Et	NH ₂	S	2
431	2-H ₂ NCO-Ph	Pr	NH ₂	S	2
432	3-H ₂ NCO-Ph	Pr ⁱ	NH ₂	S	2
433	4-MeNHCO-Ph	H	NH ₂	O	2
434	4-MeNHCO-Ph	F	NH ₂	O	2
435	4-MeNHCO-Ph	Cl	NH ₂	O	2
436	4-MeNHCO-Ph	Me	NH ₂	O	2
437	4-MeNHCO-Ph	Et	NH ₂	O	2
438	4-MeNHCO-Ph	Pr	NH ₂	O	2
439	4-MeNHCO-Ph	Pr ⁱ	NH ₂	O	2
440	4-MeNHCO-Ph	Bu	NH ₂	O	2
441	4-MeNHCO-Ph	Bu ⁱ	NH ₂	O	2
442	4-MeNHCO-Ph	Bu ^s	NH ₂	O	2
443	4-MeNHCO-Ph	Bu ^t	NH ₂	O	2
444	2-MeNHCO-Ph	H	NH ₂	S	2
445	3-MeNHCO-Ph	F	NH ₂	S	2
446	4-MeNHCO-Ph	Cl	NH ₂	S	2
447	2-MeNHCO-Ph	Me	NH ₂	S	2
448	2-MeNHCO-Ph	Et	NH ₂	S	2
449	2-MeNHCO-Ph	Pr	NH ₂	S	2
450	3-MeNHCO-Ph	Pr ⁱ	NH ₂	S	2
451	4-(Me) ₂ NCO-Ph	H	NH ₂	O	2
452	4-(Me) ₂ NCO-Ph	F	NH ₂	O	2
453	4-(Me) ₂ NCO-Ph	Cl	NH ₂	O	2
454	4-(Me) ₂ NCO-Ph	Me	NH ₂	O	2
455	4-(Me) ₂ NCO-Ph	Et	NH ₂	O	2
456	4-(Me) ₂ NCO-Ph	Pr	NH ₂	O	2
457	4-(Me) ₂ NCO-Ph	Pr ⁱ	NH ₂	O	2

- 44 -

458	4-(Me) ₂ NCO-Ph	Bu	NH ₂	O	2
459	4-(Me) ₂ NCO-Ph	Bu ⁱ	NH ₂	O	2
460	4-(Me) ₂ NCO-Ph	Bu ^s	NH ₂	O	2
461	4-(Me) ₂ NCO-Ph	Bu ^t	NH ₂	O	2
462	2-(Me) ₂ NCO-Ph	H	NH ₂	S	2
463	3-(Me) ₂ NCO-Ph	F	NH ₂	S	2
464	4-(Me) ₂ NCO-Ph	Cl	NH ₂	S	2
465	2-(Me) ₂ NCO-Ph	Me	NH ₂	S	2
466	2-(Me) ₂ NCO-Ph	Et	NH ₂	S	2
467	2-(Me) ₂ NCO-Ph	Pr	NH ₂	S	2
468	3-(Me) ₂ NCO-Ph	Pr ⁱ	NH ₂	S	2
469	Inde(1)	H	NH ₂	O	2
470	Inde(1)	Cl	NH ₂	O	2
471	Inde(1)	Me	NH ₂	O	2
472	Inde(1)	Et	NH ₂	O	2
473	Inde(1)	Pr	NH ₂	O	2
474	Inde(1)	Pr ⁱ	NH ₂	O	2
475	Np(1)	H	NH ₂	O	2
476	Np(1)	Cl	NH ₂	O	2
477	Np(1)	Me	NH ₂	O	2
478	Np(1)	Et	NH ₂	O	2
479	Np(1)	Pr	NH ₂	O	2
480	Np(1)	Pr ⁱ	NH ₂	O	2
481	Np(2)	H	NH ₂	O	2
482	Np(2)	Cl	NH ₂	O	2
483	Np(2)	Me	NH ₂	O	2
484	Np(2)	Et	NH ₂	O	2
485	Np(2)	Pr	NH ₂	O	2

- 45 -

486	Np(2)	Pr ⁱ	NH ₂	O	2
487	Pyrr(3)	H	NH ₂	O	2
488	Pyrr(3)	Cl	NH ₂	O	2
489	Pyrr(3)	Me	NH ₂	O	2
490	Pyrr(3)	Et	NH ₂	O	2
491	Pyrr(3)	Pr	NH ₂	O	2
492	Pyrr(3)	Pr ⁱ	NH ₂	O	2
493	Imid(2)	H	NH ₂	O	2
494	Imid(2)	Cl	NH ₂	O	2
495	Imid(2)	Me	NH ₂	O	2
496	Imid(2)	Et	NH ₂	O	2
497	Imid(2)	Pr	NH ₂	O	2
498	Imid(2)	Pr ⁱ	NH ₂	O	2
499	Pyza(1)	H	NH ₂	O	2
500	Pyza(1)	Cl	NH ₂	O	2
501	Pyza(1)	Me	NH ₂	O	2
502	Pyza(1)	Et	NH ₂	O	2
503	Pyza(1)	Pr	NH ₂	O	2
504	Pyza(1)	Pr ⁱ	NH ₂	O	2
505	Fur(2)	H	NH ₂	O	2
506	Fur(2)	Cl	NH ₂	O	2
507	Fur(2)	Me	NH ₂	O	2
508	Fur(2)	Et	NH ₂	O	2
509	Fur(2)	Pr	NH ₂	O	2
510	Fur(2)	Pr ⁱ	NH ₂	O	2
511	Oxa(2)	H	NH ₂	O	2
512	Oxa(2)	Cl	NH ₂	O	2
513	Oxa(2)	Me	NH ₂	O	2

- 46 -

514	Oxa(2)	Et	NH ₂	O	2
515	Oxa(2)	Pr	NH ₂	O	2
516	Oxa(2)	Pr ⁱ	NH ₂	O	2
517	Isox(3)	H	NH ₂	O	2
518	Isox(3)	Cl	NH ₂	O	2
519	Isox(3)	Me	NH ₂	O	2
520	Isox(3)	Et	NH ₂	O	2
521	Isox(3)	Pr	NH ₂	O	2
522	Isox(3)	Pr ⁱ	NH ₂	O	2
523	Thiz(2)	H	NH ₂	O	2
524	Thiz(2)	Cl	NH ₂	O	2
525	Thiz(2)	Me	NH ₂	O	2
526	Thiz(2)	Et	NH ₂	O	2
527	Thiz(2)	Pr	NH ₂	O	2
528	Thiz(2)	Pr ⁱ	NH ₂	O	2
529	Isothiz(3)	H	NH ₂	O	2
530	Isothiz(3)	Cl	NH ₂	O	2
531	Isothiz(3)	Me	NH ₂	O	2
532	Isothiz(3)	Et	NH ₂	O	2
533	Isothiz(3)	Pr	NH ₂	O	2
534	Isothiz(3)	Pr ⁱ	NH ₂	O	2
535	Thi(2)	H	NH ₂	O	2
536	Thi(2)	H	NH ₂	O	3
537	Thi(2)	H	NH ₂	O	4
538	Thi(2)	F	NH ₂	O	2
539	Thi(2)	Cl	NH ₂	O	2
540	Thi(2)	Me	NH ₂	O	2
541	Thi(2)	Et	NH ₂	O	2

- 47 -

542	Thi(2)	Pr	NH ₂	O	2
543	Thi(2)	Pr ⁱ	NH ₂	O	2
544	Thi(2)	Bu	NH ₂	O	2
545	Thi(2)	Bu ⁱ	NH ₂	O	2
546	Thi(2)	Bu ^s	NH ₂	O	2
547	Thi(2)	Bu ^t	NH ₂	O	2
548	Thi(2)	H	NH ₂	S	2
549	Thi(2)	H	NH ₂	S	3
550	Thi(2)	H	NH ₂	S	4
551	Thi(2)	F	NH ₂	S	2
552	Thi(2)	Cl	NH ₂	S	2
553	Thi(2)	Me	NH ₂	S	2
554	Thi(2)	Et	NH ₂	S	2
555	Thi(2)	Pr	NH ₂	S	2
556	Thi(2)	Pr ⁱ	NH ₂	S	2
557	Thi(2)	H	NHMe	O	2
558	Thi(2)	H	NHEt	O	2
559	Thi(2)	H	N(Me) ₂	O	2
560	Thi(2)	H	Pip(1)	O	2
561	Thi(2)	H	Mor(4)	O	2
562	3-F-Thi(2)	H	NH ₂	O	2
563	3-F-Thi(2)	Cl	NH ₂	O	2
564	3-F-Thi(2)	Me	NH ₂	O	2
565	3-F-Thi(2)	Et	NH ₂	O	2
566	3-F-Thi(2)	Pr	NH ₂	O	2
567	3-F-Thi(2)	Pr ⁱ	NH ₂	O	2
568	4-F-Thi(2)	H	NH ₂	O	2
569	4-F-Thi(2)	Cl	NH ₂	O	2

- 48 -

570	4-F-Thi(2)	Me	NH ₂	O	2
571	4-F-Thi(2)	Et	NH ₂	O	2
572	4-F-Thi(2)	Pr	NH ₂	O	2
573	4-F-Thi(2)	Pr ⁱ	NH ₂	O	2
574	5-F-Thi(2)	H	NH ₂	O	2
575	5-F-Thi(2)	Cl	NH ₂	O	2
576	5-F-Thi(2)	Me	NH ₂	O	2
577	5-F-Thi(2)	Et	NH ₂	O	2
578	5-F-Thi(2)	Pr	NH ₂	O	2
579	5-F-Thi(2)	Pr ⁱ	NH ₂	O	2
580	3-Cl-Thi(2)	H	NH ₂	O	2
581	3-Cl-Thi(2)	Cl	NH ₂	O	2
582	3-Cl-Thi(2)	Me	NH ₂	O	2
583	3-Cl-Thi(2)	Et	NH ₂	O	2
584	3-Cl-Thi(2)	Pr	NH ₂	O	2
585	3-Cl-Thi(2)	Pr ⁱ	NH ₂	O	2
586	4-Cl-Thi(2)	H	NH ₂	O	2
587	4-Cl-Thi(2)	Cl	NH ₂	O	2
588	4-Cl-Thi(2)	Me	NH ₂	O	2
589	4-Cl-Thi(2)	Et	NH ₂	O	2
590	4-Cl-Thi(2)	Pr	NH ₂	O	2
591	4-Cl-Thi(2)	Pr ⁱ	NH ₂	O	2
592	5-Cl-Thi(2)	H	NH ₂	O	2
593	5-Cl-Thi(2)	Cl	NH ₂	O	2
594	5-Cl-Thi(2)	Me	NH ₂	O	2
595	5-Cl-Thi(2)	Et	NH ₂	O	2
596	5-Cl-Thi(2)	Pr	NH ₂	O	2
597	5-Cl-Thi(2)	Pr ⁱ	NH ₂	O	2

- 49 -

598	3-Br-Thi(2)	H	NH ₂	O	2
599	3-Br-Thi(2)	Cl	NH ₂	O	2
600	3-Br-Thi(2)	Me	NH ₂	O	2
601	3-Br-Thi(2)	Et	NH ₂	O	2
602	3-Br-Thi(2)	Pr	NH ₂	O	2
603	3-Br-Thi(2)	Pr ⁱ	NH ₂	O	2
604	3-Me-Thi(2)	H	NH ₂	O	2
605	3-Me-Thi(2)	Cl	NH ₂	O	2
606	3-Me-Thi(2)	Me	NH ₂	O	2
607	3-Me-Thi(2)	Et	NH ₂	O	2
608	3-Me-Thi(2)	Pr	NH ₂	O	2
609	3-Me-Thi(2)	Pr ⁱ	NH ₂	O	2
610	4-Me-Thi(2)	H	NH ₂	O	2
611	4-Me-Thi(2)	Cl	NH ₂	O	2
612	4-Me-Thi(2)	Me	NH ₂	O	2
613	4-Me-Thi(2)	Et	NH ₂	O	2
614	4-Me-Thi(2)	Pr	NH ₂	O	2
615	4-Me-Thi(2)	Pr ⁱ	NH ₂	O	2
616	4-Et-Thi(2)	H	NH ₂	O	2
617	4-Et-Thi(2)	Cl	NH ₂	O	2
618	4-Et-Thi(2)	Me	NH ₂	O	2
619	4-Et-Thi(2)	Et	NH ₂	O	2
620	4-Et-Thi(2)	Pr	NH ₂	O	2
621	4-Et-Thi(2)	Pr ⁱ	NH ₂	O	2
622	4-CF ₃ -Thi(2)	H	NH ₂	O	2
623	4-CF ₃ -Thi(2)	Cl	NH ₂	O	2
624	4-CF ₃ -Thi(2)	Me	NH ₂	O	2
625	4-CF ₃ -Thi(2)	Et	NH ₂	O	2

- 50 -

626	4-CF ₃ -Thi(2)	Pr	NH ₂	O	2
627	4-CF ₃ -Thi(2)	Pr ⁱ	NH ₂	O	2
628	4-MeO-Thi(2)	H	NH ₂	O	2
629	4-MeO-Thi(2)	Cl	NH ₂	O	2
630	4-MeO-Thi(2)	Me	NH ₂	O	2
631	4-MeO-Thi(2)	Et	NH ₂	O	2
632	4-MeO-Thi(2)	Pr	NH ₂	O	2
633	4-MeO-Thi(2)	Pr ⁱ	NH ₂	O	2
634	3-Ph-Thi(2)	H	NH ₂	O	2
635	3-Ph-Thi(2)	Cl	NH ₂	O	2
636	3-Ph-Thi(2)	Me	NH ₂	O	2
637	3-Ph-Thi(2)	Et	NH ₂	O	2
638	3-Ph-Thi(2)	Pr	NH ₂	O	2
639	3-Ph-Thi(2)	Pr ⁱ	NH ₂	O	2
640	4-Ph-Thi(2)	H	NH ₂	O	2
641	4-Ph-Thi(2)	Cl	NH ₂	O	2
642	4-Ph-Thi(2)	Me	NH ₂	O	2
643	4-Ph-Thi(2)	Et	NH ₂	O	2
644	4-Ph-Thi(2)	Pr	NH ₂	O	2
645	4-Ph-Thi(2)	Pr ⁱ	NH ₂	O	2
646	5-Ph-Thi(2)	H	NH ₂	O	2
647	5-Ph-Thi(2)	Cl	NH ₂	O	2
648	5-Ph-Thi(2)	Me	NH ₂	O	2
649	5-Ph-Thi(2)	Et	NH ₂	O	2
650	5-Ph-Thi(2)	Pr	NH ₂	O	2
651	5-Ph-Thi(2)	Pr ⁱ	NH ₂	O	2
652	3-MeOCO-Thi(2)	H	NH ₂	O	2
653	3-MeOCO-Thi(2)	Cl	NH ₂	O	2

- 51 -

654	3-MeOCO-Thi(2)	Me	NH ₂	O	2
655	3-MeOCO-Thi(2)	Et	NH ₂	O	2
656	3-MeOCO-Thi(2)	Pr	NH ₂	O	2
657	3-MeOCO-Thi(2)	Pr ⁱ	NH ₂	O	2
658	4-MeOCO-Thi(2)	H	NH ₂	O	2
659	4-MeOCO-Thi(2)	Cl	NH ₂	O	2
660	4-MeOCO-Thi(2)	Me	NH ₂	O	2
661	4-MeOCO-Thi(2)	Et	NH ₂	O	2
662	4-MeOCO-Thi(2)	Pr	NH ₂	O	2
663	4-MeOCO-Thi(2)	Pr ⁱ	NH ₂	O	2
664	5-MeOCO-Thi(2)	H	NH ₂	O	2
665	5-MeOCO-Thi(2)	Cl	NH ₂	O	2
666	5-MeOCO-Thi(2)	Me	NH ₂	O	2
667	5-MeOCO-Thi(2)	Et	NH ₂	O	2
668	5-MeOCO-Thi(2)	Pr	NH ₂	O	2
669	5-MeOCO-Thi(2)	Pr ⁱ	NH ₂	O	2
670	3-H ₂ NCO-Thi(2)	H	NH ₂	O	2
671	3-H ₂ NCO-Thi(2)	Cl	NH ₂	O	2
672	3-H ₂ NCO-Thi(2)	Me	NH ₂	O	2
673	3-H ₂ NCO-Thi(2)	Et	NH ₂	O	2
674	3-H ₂ NCO-Thi(2)	Pr	NH ₂	O	2
675	3-H ₂ NCO-Thi(2)	Pr ⁱ	NH ₂	O	2
676	4-H ₂ NCO-Thi(2)	H	NH ₂	O	2
677	4-H ₂ NCO-Thi(2)	Cl	NH ₂	O	2
678	4-H ₂ NCO-Thi(2)	Me	NH ₂	O	2
679	4-H ₂ NCO-Thi(2)	Et	NH ₂	O	2
680	4-H ₂ NCO-Thi(2)	Pr	NH ₂	O	2
681	4-H ₂ NCO-Thi(2)	Pr ⁱ	NH ₂	O	2

- 52 -

682	5-H ₂ NCO-Thi(2)	H	NH ₂	O	2
683	5-H ₂ NCO-Thi(2)	Cl	NH ₂	O	2
684	5-H ₂ NCO-Thi(2)	Me	NH ₂	O	2
685	5-H ₂ NCO-Thi(2)	Et	NH ₂	O	2
686	5-H ₂ NCO-Thi(2)	Pr	NH ₂	O	2
687	5-H ₂ NCO-Thi(2)	Pr ⁱ	NH ₂	O	2
688	3-MeNHCO-Thi(2)	H	NH ₂	O	2
689	3-MeNHCO-Thi(2)	Cl	NH ₂	O	2
690	3-MeNHCO-Thi(2)	Me	NH ₂	O	2
691	3-MeNHCO-Thi(2)	Et	NH ₂	O	2
692	3-MeNHCO-Thi(2)	Pr	NH ₂	O	2
693	3-MeNHCO-Thi(2)	Pr ⁱ	NH ₂	O	2
694	4-MeNHCO-Thi(2)	H	NH ₂	O	2
695	4-MeNHCO-Thi(2)	Cl	NH ₂	O	2
696	4-MeNHCO-Thi(2)	Me	NH ₂	O	2
697	4-MeNHCO-Thi(2)	Et	NH ₂	O	2
698	4-MeNHCO-Thi(2)	Pr	NH ₂	O	2
699	4-MeNHCO-Thi(2)	Pr ⁱ	NH ₂	O	2
700	5-MeNHCO-Thi(2)	H	NH ₂	O	2
701	5-MeNHCO-Thi(2)	Cl	NH ₂	O	2
702	5-MeNHCO-Thi(2)	Me	NH ₂	O	2
703	5-MeNHCO-Thi(2)	Et	NH ₂	O	2
704	5-MeNHCO-Thi(2)	Pr	NH ₂	O	2
705	5-MeNHCO-Thi(2)	Pr ⁱ	NH ₂	O	2
706	3-(Me) ₂ NCO-Thi(2)	H	NH ₂	O	2
707	3-(Me) ₂ NCO-Thi(2)	Cl	NH ₂	O	2
708	3-(Me) ₂ NCO-Thi(2)	Me	NH ₂	O	2
709	3-(Me) ₂ NCO-Thi(2)	Et	NH ₂	O	2

- 53 -

710	3-(Me) ₂ NCO-Thi(2)	Pr	NH ₂	O	2
711	3-(Me) ₂ NCO-Thi(2)	Pr ⁱ	NH ₂	O	2
712	4-(Me) ₂ NCO-Thi(2)	H	NH ₂	O	2
713	4-(Me) ₂ NCO-Thi(2)	Cl	NH ₂	O	2
714	4-(Me) ₂ NCO-Thi(2)	Me	NH ₂	O	2
715	4-(Me) ₂ NCO-Thi(2)	Et	NH ₂	O	2
716	4-(Me) ₂ NCO-Thi(2)	Pr	NH ₂	O	2
717	4-(Me) ₂ NCO-Thi(2)	Pr ⁱ	NH ₂	O	2
718	5-(Me) ₂ NCO-Thi(2)	H	NH ₂	O	2
719	5-(Me) ₂ NCO-Thi(2)	Cl	NH ₂	O	2
720	5-(Me) ₂ NCO-Thi(2)	Me	NH ₂	O	2
721	5-(Me) ₂ NCO-Thi(2)	Et	NH ₂	O	2
722	5-(Me) ₂ NCO-Thi(2)	Pr	NH ₂	O	2
723	5-(Me) ₂ NCO-Thi(2)	Pr ⁱ	NH ₂	O	2
724	Thi(3)	H	NH ₂	O	2
725	Thi(3)	H	NH ₂	O	3
726	Thi(3)	H	NH ₂	O	4
727	Thi(3)	H	NH ₂	O	5
728	Thi(3)	F	NH ₂	O	2
729	Thi(3)	Cl	NH ₂	O	2
730	Thi(3)	Me	NH ₂	O	2
731	Thi(3)	Et	NH ₂	O	2
732	Thi(3)	Pr	NH ₂	O	2
733	Thi(3)	Pr ⁱ	NH ₂	O	2
734	Thi(3)	Bu	NH ₂	O	2
735	Thi(3)	Bu ⁱ	NH ₂	O	2
736	Thi(3)	Bu ^s	NH ₂	O	2
737	Thi(3)	Bu ^t	NH ₂	O	2

738	Thi(3)	H	NH ₂	S	2
739	Thi(3)	H	NH ₂	S	3
740	Thi(3)	H	NH ₂	S	4
741	Thi(3)	F	NH ₂	S	2
742	Thi(3)	Cl	NH ₂	S	2
743	Thi(3)	Me	NH ₂	S	2
744	Thi(3)	Et	NH ₂	S	2
745	Thi(3)	Pr	NH ₂	S	2
746	Thi(3)	Pr ⁱ	NH ₂	S	2
747	Thi(3)	H	NHMe	O	2
748	Thi(3)	H	NHEt	O	2
749	Thi(3)	H	N(Me) ₂	O	2
750	Thi(3)	H	Pip(1)	O	2
751	Thi(3)	H	Mor(4)	O	2
752	2-F-Thi(3)	H	NH ₂	O	2
753	2-F-Thi(3)	Cl	NH ₂	O	2
754	2-F-Thi(3)	Me	NH ₂	O	2
755	2-F-Thi(3)	Et	NH ₂	O	2
756	2-F-Thi(3)	Pr	NH ₂	O	2
757	2-F-Thi(3)	Pr ⁱ	NH ₂	O	2
758	2-F-Thi(3)	H	NH ₂	S	2
759	2-F-Thi(3)	Cl	NH ₂	S	2
760	2-F-Thi(3)	Me	NH ₂	S	2
761	2-F-Thi(3)	Et	NH ₂	S	2
762	2-F-Thi(3)	Pr	NH ₂	S	2
763	2-F-Thi(3)	Pr ⁱ	NH ₂	S	2
764	4-F-Thi(3)	H	NH ₂	O	2
765	4-F-Thi(3)	Cl	NH ₂	O	2

- 55 -

766	4-F-Thi(3)	Me	NH ₂	O	2
767	4-F-Thi(3)	Et	NH ₂	O	2
768	4-F-Thi(3)	Pr	NH ₂	O	2
769	4-F-Thi(3)	Pr ⁱ	NH ₂	O	2
770	4-F-Thi(3)	H	NH ₂	S	2
771	4-F-Thi(3)	Cl	NH ₂	S	2
772	4-F-Thi(3)	Me	NH ₂	S	2
773	4-F-Thi(3)	Et	NH ₂	S	2
774	4-F-Thi(3)	Pr	NH ₂	S	2
775	4-F-Thi(3)	Pr ⁱ	NH ₂	S	2
776	5-F-Thi(3)	H	NH ₂	O	2
777	5-F-Thi(3)	Cl	NH ₂	O	2
778	5-F-Thi(3)	Me	NH ₂	O	2
779	5-F-Thi(3)	Et	NH ₂	O	2
780	5-F-Thi(3)	Pr	NH ₂	O	2
781	5-F-Thi(3)	Pr ⁱ	NH ₂	O	2
782	5-F-Thi(3)	H	NH ₂	S	2
783	5-F-Thi(3)	Cl	NH ₂	S	2
784	5-F-Thi(3)	Me	NH ₂	S	2
785	5-F-Thi(3)	Et	NH ₂	S	2
786	5-F-Thi(3)	Pr	NH ₂	S	2
787	5-F-Thi(3)	Pr ⁱ	NH ₂	S	2
788	2-Cl-Thi(3)	H	NH ₂	O	2
789	2-Cl-Thi(3)	Cl	NH ₂	O	2
790	2-Cl-Thi(3)	Me	NH ₂	O	2
791	2-Cl-Thi(3)	Et	NH ₂	O	2
792	2-Cl-Thi(3)	Pr	NH ₂	O	2
793	2-Cl-Thi(3)	Pr ⁱ	NH ₂	O	2

- 56 -

794	4-Cl-Thi(3)	H	NH ₂	O	2
795	4-Cl-Thi(3)	Cl	NH ₂	O	2
796	4-Cl-Thi(3)	Me	NH ₂	O	2
797	4-Cl-Thi(3)	Et	NH ₂	O	2
798	4-Cl-Thi(3)	Pr	NH ₂	O	2
799	4-Cl-Thi(3)	Pr ⁱ	NH ₂	O	2
800	5-Cl-Thi(3)	H	NH ₂	O	2
801	5-Cl-Thi(3)	Cl	NH ₂	O	2
802	5-Cl-Thi(3)	Me	NH ₂	O	2
803	5-Cl-Thi(3)	Et	NH ₂	O	2
804	5-Cl-Thi(3)	Pr	NH ₂	O	2
805	5-Cl-Thi(3)	Pr ⁱ	NH ₂	O	2
806	4-Me-Thi(3)	H	NH ₂	O	2
807	4-Me-Thi(3)	Cl	NH ₂	O	2
808	4-Me-Thi(3)	Me	NH ₂	O	2
809	4-Me-Thi(3)	Et	NH ₂	O	2
810	4-Me-Thi(3)	Pr	NH ₂	O	2
811	4-Me-Thi(3)	Pr ⁱ	NH ₂	O	2
812	4-Et-Thi(3)	H	NH ₂	O	2
813	4-Et-Thi(3)	Cl	NH ₂	O	2
814	4-Et-Thi(3)	Me	NH ₂	O	2
815	4-Et-Thi(3)	Et	NH ₂	O	2
816	4-Et-Thi(3)	Pr	NH ₂	O	2
817	4-Et-Thi(3)	Pr ⁱ	NH ₂	O	2
818	4-CF ₃ -Thi(3)	H	NH ₂	O	2
819	4-CF ₃ -Thi(3)	Cl	NH ₂	O	2
820	4-CF ₃ -Thi(3)	Me	NH ₂	O	2
821	4-CF ₃ -Thi(3)	Et	NH ₂	O	2

- 57 -

822	4-CF ₃ -Thi(3)	Pr	NH ₂	O	2
823	4-CF ₃ -Thi(3)	Pr ⁱ	NH ₂	O	2
824	4-MeO-Thi(3)	H	NH ₂	O	2
825	4-MeO-Thi(3)	Cl	NH ₂	O	2
826	4-MeO-Thi(3)	Me	NH ₂	O	2
827	4-MeO-Thi(3)	Et	NH ₂	O	2
828	4-MeO-Thi(3)	Pr	NH ₂	O	2
829	4-MeO-Thi(3)	Pr ⁱ	NH ₂	O	2
830	2-Ph-Thi(3)	H	NH ₂	O	2
831	2-Ph-Thi(3)	Cl	NH ₂	O	2
832	2-Ph-Thi(3)	Me	NH ₂	O	2
833	2-Ph-Thi(3)	Et	NH ₂	O	2
834	2-Ph-Thi(3)	Pr	NH ₂	O	2
835	2-Ph-Thi(3)	Pr ⁱ	NH ₂	O	2
836	4-Ph-Thi(3)	H	NH ₂	O	2
837	4-Ph-Thi(3)	Cl	NH ₂	O	2
838	4-Ph-Thi(3)	Me	NH ₂	O	2
839	4-Ph-Thi(3)	Et	NH ₂	O	2
840	4-Ph-Thi(3)	Pr	NH ₂	O	2
841	4-Ph-Thi(3)	Pr ⁱ	NH ₂	O	2
842	5-Ph-Thi(3)	H	NH ₂	O	2
843	5-Ph-Thi(3)	Cl	NH ₂	O	2
844	5-Ph-Thi(3)	Me	NH ₂	O	2
845	5-Ph-Thi(3)	Et	NH ₂	O	2
846	5-Ph-Thi(3)	Pr	NH ₂	O	2
847	5-Ph-Thi(3)	Pr ⁱ	NH ₂	O	2
848	2-MeOCO-Thi(3)	H	NH ₂	O	2
849	2-MeOCO-Thi(3)	Cl	NH ₂	O	2

- 58 -

850	2-MeOCO-Thi(3)	Me	NH ₂	O	2
851	2-MeOCO-Thi(3)	Et	NH ₂	O	2
852	2-MeOCO-Thi(3)	Pr	NH ₂	O	2
853	2-MeOCO-Thi(3)	Pr ⁱ	NH ₂	O	2
854	4-MeOCO-Thi(3)	H	NH ₂	O	2
855	4-MeOCO-Thi(3)	Cl	NH ₂	O	2
856	4-MeOCO-Thi(3)	Me	NH ₂	O	2
857	4-MeOCO-Thi(3)	Et	NH ₂	O	2
858	4-MeOCO-Thi(3)	Pr	NH ₂	O	2
859	4-MeOCO-Thi(3)	Pr ⁱ	NH ₂	O	2
860	5-MeOCO-Thi(3)	H	NH ₂	O	2
861	5-MeOCO-Thi(3)	Cl	NH ₂	O	2
862	5-MeOCO-Thi(3)	Me	NH ₂	O	2
863	5-MeOCO-Thi(3)	Et	NH ₂	O	2
864	5-MeOCO-Thi(3)	Pr	NH ₂	O	2
865	5-MeOCO-Thi(3)	Pr ⁱ	NH ₂	O	2
866	2-H ₂ NCO-Thi(3)	H	NH ₂	O	2
867	2-H ₂ NCO-Thi(3)	Cl	NH ₂	O	2
868	2-H ₂ NCO-Thi(3)	Me	NH ₂	O	2
869	2-H ₂ NCO-Thi(3)	Et	NH ₂	O	2
870	2-H ₂ NCO-Thi(3)	Pr	NH ₂	O	2
871	2-H ₂ NCO-Thi(3)	Pr ⁱ	NH ₂	O	2
872	4-H ₂ NCO-Thi(3)	H	NH ₂	O	2
873	4-H ₂ NCO-Thi(3)	Cl	NH ₂	O	2
874	4-H ₂ NCO-Thi(3)	Me	NH ₂	O	2
875	4-H ₂ NCO-Thi(3)	Et	NH ₂	O	2
876	4-H ₂ NCO-Thi(3)	Pr	NH ₂	O	2
877	4-H ₂ NCO-Thi(3)	Pr ⁱ	NH ₂	O	2

- 59 -

878	5-H ₂ NCO-Thi(3)	H	NH ₂	O	2
879	5-H ₂ NCO-Thi(3)	Cl	NH ₂	O	2
880	5-H ₂ NCO-Thi(3)	Me	NH ₂	O	2
881	5-H ₂ NCO-Thi(3)	Et	NH ₂	O	2
882	5-H ₂ NCO-Thi(3)	Pr	NH ₂	O	2
883	5-H ₂ NCO-Thi(3)	Pr ⁱ	NH ₂	O	2
884	2-MeNHCO-Thi(3)	H	NH ₂	O	2
885	2-MeNHCO-Thi(3)	Cl	NH ₂	O	2
886	2-MeNHCO-Thi(3)	Me	NH ₂	O	2
887	2-MeNHCO-Thi(3)	Et	NH ₂	O	2
888	2-MeNHCO-Thi(3)	Pr	NH ₂	O	2
889	2-MeNHCO-Thi(3)	Pr ⁱ	NH ₂	O	2
890	4-MeNHCO-Thi(3)	H	NH ₂	O	2
891	4-MeNHCO-Thi(3)	Cl	NH ₂	O	2
892	4-MeNHCO-Thi(3)	Me	NH ₂	O	2
893	4-MeNHCO-Thi(3)	Et	NH ₂	O	2
894	4-MeNHCO-Thi(3)	Pr	NH ₂	O	2
895	4-MeNHCO-Thi(3)	Pr ⁱ	NH ₂	O	2
896	5-MeNHCO-Thi(3)	H	NH ₂	O	2
897	5-MeNHCO-Thi(3)	Cl	NH ₂	O	2
898	5-MeNHCO-Thi(3)	Me	NH ₂	O	2
899	5-MeNHCO-Thi(3)	Et	NH ₂	O	2
900	5-MeNHCO-Thi(3)	Pr	NH ₂	O	2
901	5-MeNHCO-Thi(3)	Pr ⁱ	NH ₂	O	2
902	2-(Me) ₂ NCO-Thi(3)	H	NH ₂	O	2
903	2-(Me) ₂ NCO-Thi(3)	Cl	NH ₂	O	2
904	2-(Me) ₂ NCO-Thi(3)	Me	NH ₂	O	2
905	2-(Me) ₂ NCO-Thi(3)	Et	NH ₂	O	2

- 60 -

906	2-(Me) ₂ NCO-Thi(3)	Pr	NH ₂	O	2
907	2-(Me) ₂ NCO-Thi(3)	Pr ⁱ	NH ₂	O	2
908	4-(Me) ₂ NCO-Thi(3)	H	NH ₂	O	2
909	4-(Me) ₂ NCO-Thi(3)	Cl	NH ₂	O	2
910	4-(Me) ₂ NCO-Thi(3)	Me	NH ₂	O	2
911	4-(Me) ₂ NCO-Thi(3)	Et	NH ₂	O	2
912	4-(Me) ₂ NCO-Thi(3)	Pr	NH ₂	O	2
913	4-(Me) ₂ NCO-Thi(3)	Pr ⁱ	NH ₂	O	2
914	5-(Me) ₂ NCO-Thi(3)	H	NH ₂	O	2
915	5-(Me) ₂ NCO-Thi(3)	Cl	NH ₂	O	2
916	5-(Me) ₂ NCO-Thi(3)	Me	NH ₂	O	2
917	5-(Me) ₂ NCO-Thi(3)	Et	NH ₂	O	2
918	5-(Me) ₂ NCO-Thi(3)	Pr	NH ₂	O	2
919	5-(Me) ₂ NCO-Thi(3)	Pr ⁱ	NH ₂	O	2
920	Pyr(2)	H	NH ₂	O	2
921	Pyr(2)	H	NH ₂	O	2
922	Pyr(2)	H	NH ₂	O	3
923	Pyr(2)	H	NH ₂	O	4
924	Pyr(2)	F	NH ₂	O	2
925	Pyr(2)	Cl	NH ₂	O	2
926	Pyr(2)	Me	NH ₂	O	2
927	Pyr(2)	Et	NH ₂	O	2
928	Pyr(2)	Pr	NH ₂	O	2
929	Pyr(2)	Pr ⁱ	NH ₂	O	2
930	Pyr(2)	Bu	NH ₂	O	2
931	Pyr(2)	Bu ⁱ	NH ₂	O	2
932	Pyr(2)	Bu ^s	NH ₂	O	2
933	Pyr(2)	Bu ^t	NH ₂	O	2

- 61 -

934	Pyr(2)	H	NH ₂	S	2
935	Pyr(2)	H	NH ₂	S	3
936	Pyr(2)	H	NH ₂	S	4
937	Pyr(2)	F	NH ₂	S	2
938	Pyr(2)	Cl	NH ₂	S	2
939	Pyr(2)	Me	NH ₂	S	2
940	Pyr(2)	Et	NH ₂	S	2
941	Pyr(2)	Pr	NH ₂	S	2
942	Pyr(2)	Pr ⁱ	NH ₂	S	2
943	Pyr(2)	H	NHMe	O	2
944	Pyr(2)	H	NHEt	O	2
945	Pyr(2)	H	N(Me) ₂	O	2
946	Pyr(2)	H	Pip(1)	O	2
947	Pyr(2)	H	Mor(4)	O	2
948	3-F-Pyr(2)	H	NH ₂	O	2
949	3-F-Pyr(2)	Cl	NH ₂	O	2
950	3-F-Pyr(2)	Me	NH ₂	O	2
951	3-F-Pyr(2)	Et	NH ₂	O	2
952	3-F-Pyr(2)	Pr	NH ₂	O	2
953	3-F-Pyr(2)	Pr ⁱ	NH ₂	O	2
954	4-F-Pyr(2)	H	NH ₂	O	2
955	4-F-Pyr(2)	Cl	NH ₂	O	2
956	4-F-Pyr(2)	Me	NH ₂	O	2
957	4-F-Pyr(2)	Et	NH ₂	O	2
958	4-F-Pyr(2)	Pr	NH ₂	O	2
959	4-F-Pyr(2)	Pr ⁱ	NH ₂	O	2
960	3-Cl-Pyr(2)	H	NH ₂	O	2
961	3-Cl-Pyr(2)	Cl	NH ₂	O	2

- 62 -

962	3-Cl-Pyr(2)	Me	NH ₂	O	2
963	3-Cl-Pyr(2)	Et	NH ₂	O	2
964	3-Cl-Pyr(2)	Pr	NH ₂	O	2
965	3-Cl-Pyr(2)	Pr ⁱ	NH ₂	O	2
966	4-Cl-Pyr(2)	H	NH ₂	O	2
967	4-Cl-Pyr(2)	Cl	NH ₂	O	2
968	4-Cl-Pyr(2)	Me	NH ₂	O	2
969	4-Cl-Pyr(2)	Et	NH ₂	O	2
970	4-Cl-Pyr(2)	Pr	NH ₂	O	2
971	4-Cl-Pyr(2)	Pr ⁱ	NH ₂	O	2
972	4-Me-Pyr(2)	H	NH ₂	O	2
973	4-Me-Pyr(2)	Cl	NH ₂	O	2
974	4-Me-Pyr(2)	Me	NH ₂	O	2
975	4-Me-Pyr(2)	Et	NH ₂	O	2
976	4-Me-Pyr(2)	Pr	NH ₂	O	2
977	4-Me-Pyr(2)	Pr ⁱ	NH ₂	O	2
978	4-Et-Pyr(2)	H	NH ₂	O	2
979	4-Et-Pyr(2)	Cl	NH ₂	O	2
980	4-Et-Pyr(2)	Me	NH ₂	O	2
981	4-Et-Pyr(2)	Et	NH ₂	O	2
982	4-Et-Pyr(2)	Pr	NH ₂	O	2
983	4-Et-Pyr(2)	Pr ⁱ	NH ₂	O	2
984	4-CF ₃ -Pyr(2)	H	NH ₂	O	2
985	4-CF ₃ -Pyr(2)	Cl	NH ₂	O	2
986	4-CF ₃ -Pyr(2)	Me	NH ₂	O	2
987	4-CF ₃ -Pyr(2)	Et	NH ₂	O	2
988	4-CF ₃ -Pyr(2)	Pr	NH ₂	O	2
989	4-CF ₃ -Pyr(2)	Pr ⁱ	NH ₂	O	2

- 63 -

990	4-MeO-Pyr(2)	H	NH ₂	O	2
991	4-MeO-Pyr(2)	Cl	NH ₂	O	2
992	4-MeO-Pyr(2)	Me	NH ₂	O	2
993	4-MeO-Pyr(2)	Et	NH ₂	O	2
994	4-MeO-Pyr(2)	Pr	NH ₂	O	2
995	4-MeO-Pyr(2)	Pr ⁱ	NH ₂	O	2
996	3-Ph-Pyr(2)	H	NH ₂	O	2
997	3-Ph-Pyr(2)	Cl	NH ₂	O	2
998	3-Ph-Pyr(2)	Me	NH ₂	O	2
999	3-Ph-Pyr(2)	Et	NH ₂	O	2
1000	3-Ph-Pyr(2)	Pr	NH ₂	O	2
1001	3-Ph-Pyr(2)	Pr ⁱ	NH ₂	O	2
1002	4-Ph-Pyr(2)	H	NH ₂	O	2
1003	4-Ph-Pyr(2)	Cl	NH ₂	O	2
1004	4-Ph-Pyr(2)	Me	NH ₂	O	2
1005	4-Ph-Pyr(2)	Et	NH ₂	O	2
1006	4-Ph-Pyr(2)	Pr	NH ₂	O	2
1007	4-Ph-Pyr(2)	Pr ⁱ	NH ₂	O	2
1008	3-MeOCO-Pyr(2)	H	NH ₂	O	2
1009	3-MeOCO-Pyr(2)	Cl	NH ₂	O	2
1010	3-MeOCO-Pyr(2)	Me	NH ₂	O	2
1011	3-MeOCO-Pyr(2)	Et	NH ₂	O	2
1012	3-MeOCO-Pyr(2)	Pr	NH ₂	O	2
1013	3-MeOCO-Pyr(2)	Pr ⁱ	NH ₂	O	2
1014	4-MeOCO-Pyr(2)	H	NH ₂	O	2
1015	4-MeOCO-Pyr(2)	Cl	NH ₂	O	2
1016	4-MeOCO-Pyr(2)	Me	NH ₂	O	2
1017	4-MeOCO-Pyr(2)	Et	NH ₂	O	2

- 64 -

1018	4-MeOCO-Pyr(2)	Pr	NH ₂	O	2
1019	4-MeOCO-Pyr(2)	Pr ⁱ	NH ₂	O	2
1020	3-H ₂ NCO-Pyr(2)	H	NH ₂	O	2
1021	3-H ₂ NCO-Pyr(2)	Cl	NH ₂	O	2
1022	3-H ₂ NCO-Pyr(2)	Me	NH ₂	O	2
1023	3-H ₂ NCO-Pyr(2)	Et	NH ₂	O	2
1024	3-H ₂ NCO-Pyr(2)	Pr	NH ₂	O	2
1025	3-H ₂ NCO-Pyr(2)	Pr ⁱ	NH ₂	O	2
1026	4-H ₂ NCO-Pyr(2)	H	NH ₂	O	2
1027	4-H ₂ NCO-Pyr(2)	Cl	NH ₂	O	2
1028	4-H ₂ NCO-Pyr(2)	Me	NH ₂	O	2
1029	4-H ₂ NCO-Pyr(2)	Et	NH ₂	O	2
1030	4-H ₂ NCO-Pyr(2)	Pr	NH ₂	O	2
1031	4-H ₂ NCO-Pyr(2)	Pr ⁱ	NH ₂	O	2
1032	3-MeNHCO-Pyr(2)	H	NH ₂	O	2
1033	3-MeNHCO-Pyr(2)	Cl	NH ₂	O	2
1034	3-MeNHCO-Pyr(2)	Me	NH ₂	O	2
1035	3-MeNHCO-Pyr(2)	Et	NH ₂	O	2
1036	3-MeNHCO-Pyr(2)	Pr	NH ₂	O	2
1037	3-MeNHCO-Pyr(2)	Pr ⁱ	NH ₂	O	2
1038	4-MeNHCO-Pyr(2)	H	NH ₂	O	2
1039	4-MeNHCO-Pyr(2)	Cl	NH ₂	O	2
1040	4-MeNHCO-Pyr(2)	Me	NH ₂	O	2
1041	4-MeNHCO-Pyr(2)	Et	NH ₂	O	2
1042	4-MeNHCO-Pyr(2)	Pr	NH ₂	O	2
1043	4-MeNHCO-Pyr(2)	Pr ⁱ	NH ₂	O	2
1044	3-(Me) ₂ NCO-Pyr(2)	H	NH ₂	O	2
1045	3-(Me) ₂ NCO-Pyr(2)	Cl	NH ₂	O	2

- 65 -

1046	3-(Me) ₂ NCO-Pyr(2)	Me	NH ₂	O	2
1047	3-(Me) ₂ NCO-Pyr(2)	Et	NH ₂	O	2
1048	3-(Me) ₂ NCO-Pyr(2)	Pr	NH ₂	O	2
1049	3-(Me) ₂ NCO-Pyr(2)	Pr ⁱ	NH ₂	O	2
1050	4-(Me) ₂ NCO-Pyr(2)	H	NH ₂	O	2
1051	4-(Me) ₂ NCO-Pyr(2)	Cl	NH ₂	O	2
1052	4-(Me) ₂ NCO-Pyr(2)	Me	NH ₂	O	2
1053	4-(Me) ₂ NCO-Pyr(2)	Et	NH ₂	O	2
1054	4-(Me) ₂ NCO-Pyr(2)	Pr	NH ₂	O	2
1055	4-(Me) ₂ NCO-Pyr(2)	Pr ⁱ	NH ₂	O	2
1056	Pyr(3)	H	NH ₂	O	2
1057	Pyr(3)	H	NH ₂	O	2
1058	Pyr(3)	H	NH ₂	O	3
1059	Pyr(3)	H	NH ₂	O	4
1060	Pyr(3)	F	NH ₂	O	2
1061	Pyr(3)	Cl	NH ₂	O	2
1062	Pyr(3)	Me	NH ₂	O	2
1063	Pyr(3)	Et	NH ₂	O	2
1064	Pyr(3)	Pr	NH ₂	O	2
1065	Pyr(3)	Pr ⁱ	NH ₂	O	2
1066	Pyr(3)	Bu	NH ₂	O	2
1067	Pyr(3)	Bu ⁱ	NH ₂	O	2
1068	Pyr(3)	Bu ^s	NH ₂	O	2
1069	Pyr(3)	Bu ^t	NH ₂	O	2
1070	Pyr(3)	H	NH ₂	S	2
1071	Pyr(3)	H	NH ₂	S	3
1072	Pyr(3)	H	NH ₂	S	4
1073	Pyr(3)	F	NH ₂	S	2

- 66 -

1074	Pyr(3)	Cl	NH ₂	S	2
1075	Pyr(3)	Me	NH ₂	S	2
1076	Pyr(3)	Et	NH ₂	S	2
1077	Pyr(3)	Pr	NH ₂	S	2
1078	Pyr(3)	Pr ⁱ	NH ₂	S	2
1079	Pyr(3)	H	NHMe	O	2
1080	Pyr(3)	H	NHEt	O	2
1081	Pyr(3)	H	N(Me) ₂	O	2
1082	Pyr(3)	H	Pip(1)	O	2
1083	Pyr(3)	H	Mor(4)	O	2
1084	2-F-Pyr(3)	H	NH ₂	O	2
1085	2-F-Pyr(3)	Cl	NH ₂	O	2
1086	2-F-Pyr(3)	Me	NH ₂	O	2
1087	2-F-Pyr(3)	Et	NH ₂	O	2
1088	2-F-Pyr(3)	Pr	NH ₂	O	2
1089	2-F-Pyr(3)	Pr ⁱ	NH ₂	O	2
1090	4-F-Pyr(3)	H	NH ₂	O	2
1091	4-F-Pyr(3)	Cl	NH ₂	O	2
1092	4-F-Pyr(3)	Me	NH ₂	O	2
1093	4-F-Pyr(3)	Et	NH ₂	O	2
1094	4-F-Pyr(3)	Pr	NH ₂	O	2
1095	4-F-Pyr(3)	Pr ⁱ	NH ₂	O	2
1096	2-Cl-Pyr(3)	H	NH ₂	O	2
1097	2-Cl-Pyr(3)	Cl	NH ₂	O	2
1098	2-Cl-Pyr(3)	Me	NH ₂	O	2
1099	2-Cl-Pyr(3)	Et	NH ₂	O	2
1100	2-Cl-Pyr(3)	Pr	NH ₂	O	2
1101	2-Cl-Pyr(3)	Pr ⁱ	NH ₂	O	2

- 67 -

1102	4-Cl-Pyr(3)	H	NH ₂	O	2
1103	4-Cl-Pyr(3)	Cl	NH ₂	O	2
1104	4-Cl-Pyr(3)	Me	NH ₂	O	2
1105	4-Cl-Pyr(3)	Et	NH ₂	O	2
1106	4-Cl-Pyr(3)	Pr	NH ₂	O	2
1107	4-Cl-Pyr(3)	Pr ⁱ	NH ₂	O	2
1108	4-Me-Pyr(3)	H	NH ₂	O	2
1109	4-Me-Pyr(3)	Cl	NH ₂	O	2
1110	4-Me-Pyr(3)	Me	NH ₂	O	2
1111	4-Me-Pyr(3)	Et	NH ₂	O	2
1112	4-Me-Pyr(3)	Pr	NH ₂	O	2
1113	4-Me-Pyr(3)	Pr ⁱ	NH ₂	O	2
1114	4-Et-Pyr(3)	H	NH ₂	O	2
1115	4-Et-Pyr(3)	Cl	NH ₂	O	2
1116	4-Et-Pyr(3)	Me	NH ₂	O	2
1117	4-Et-Pyr(3)	Et	NH ₂	O	2
1118	4-Et-Pyr(3)	Pr	NH ₂	O	2
1119	4-Et-Pyr(3)	Pr ⁱ	NH ₂	O	2
1120	4-CF ₃ -Pyr(3)	H	NH ₂	O	2
1121	4-CF ₃ -Pyr(3)	Cl	NH ₂	O	2
1122	4-CF ₃ -Pyr(3)	Me	NH ₂	O	2
1123	4-CF ₃ -Pyr(3)	Et	NH ₂	O	2
1124	4-CF ₃ -Pyr(3)	Pr	NH ₂	O	2
1125	4-CF ₃ -Pyr(3)	Pr ⁱ	NH ₂	O	2
1126	4-MeO-Pyr(3)	H	NH ₂	O	2
1127	4-MeO-Pyr(3)	Cl	NH ₂	O	2
1128	4-MeO-Pyr(3)	Me	NH ₂	O	2
1129	4-MeO-Pyr(3)	Et	NH ₂	O	2

- 68 -

1130	4-MeO-Pyr(3)	Pr	NH ₂	O	2
1131	4-MeO-Pyr(3)	Pr ⁱ	NH ₂	O	2
1132	2-Ph-Pyr(3)	H	NH ₂	O	2
1133	2-Ph-Pyr(3)	Cl	NH ₂	O	2
1134	2-Ph-Pyr(3)	Me	NH ₂	O	2
1135	2-Ph-Pyr(3)	Et	NH ₂	O	2
1136	2-Ph-Pyr(3)	Pr	NH ₂	O	2
1137	2-Ph-Pyr(3)	Pr ⁱ	NH ₂	O	2
1138	4-Ph-Pyr(3)	H	NH ₂	O	2
1139	4-Ph-Pyr(3)	Cl	NH ₂	O	2
1140	4-Ph-Pyr(3)	Me	NH ₂	O	2
1141	4-Ph-Pyr(3)	Et	NH ₂	O	2
1142	4-Ph-Pyr(3)	Pr	NH ₂	O	2
1143	4-Ph-Pyr(3)	Pr ⁱ	NH ₂	O	2
1144	2-MeOCO-Pyr(3)	H	NH ₂	O	2
1145	2-MeOCO-Pyr(3)	Cl	NH ₂	O	2
1146	2-MeOCO-Pyr(3)	Me	NH ₂	O	2
1147	2-MeOCO-Pyr(3)	Et	NH ₂	O	2
1148	2-MeOCO-Pyr(3)	Pr	NH ₂	O	2
1149	2-MeOCO-Pyr(3)	Pr ⁱ	NH ₂	O	2
1150	4-MeOCO-Pyr(3)	H	NH ₂	O	2
1151	4-MeOCO-Pyr(3)	Cl	NH ₂	O	2
1152	4-MeOCO-Pyr(3)	Me	NH ₂	O	2
1153	4-MeOCO-Pyr(3)	Et	NH ₂	O	2
1154	4-MeOCO-Pyr(3)	Pr	NH ₂	O	2
1155	4-MeOCO-Pyr(3)	Pr ⁱ	NH ₂	O	2
1156	2-H ₂ NCO-Pyr(3)	H	NH ₂	O	2
1157	2-H ₂ NCO-Pyr(3)	Cl	NH ₂	O	2

- 69 -

1158	2-H ₂ NCO-Pyr(3)	Me	NH ₂	O	2
1159	2-H ₂ NCO-Pyr(3)	Et	NH ₂	O	2
1160	2-H ₂ NCO-Pyr(3)	Pr	NH ₂	O	2
1161	2-H ₂ NCO-Pyr(3)	Pr ⁱ	NH ₂	O	2
1162	4-H ₂ NCO-Pyr(3)	H	NH ₂	O	2
1163	4-H ₂ NCO-Pyr(3)	Cl	NH ₂	O	2
1164	4-H ₂ NCO-Pyr(3)	Me	NH ₂	O	2
1165	4-H ₂ NCO-Pyr(3)	Et	NH ₂	O	2
1166	4-H ₂ NCO-Pyr(3)	Pr	NH ₂	O	2
1167	4-H ₂ NCO-Pyr(3)	Pr ⁱ	NH ₂	O	2
1168	2-MeNHCO-Pyr(3)	H	NH ₂	O	2
1169	2-MeNHCO-Pyr(3)	Cl	NH ₂	O	2
1170	2-MeNHCO-Pyr(3)	Me	NH ₂	O	2
1171	2-MeNHCO-Pyr(3)	Et	NH ₂	O	2
1172	2-MeNHCO-Pyr(3)	Pr	NH ₂	O	2
1173	2-MeNHCO-Pyr(3)	Pr ⁱ	NH ₂	O	2
1174	4-MeNHCO-Pyr(3)	H	NH ₂	O	2
1175	4-MeNHCO-Pyr(3)	Cl	NH ₂	O	2
1176	4-MeNHCO-Pyr(3)	Me	NH ₂	O	2
1177	4-MeNHCO-Pyr(3)	Et	NH ₂	O	2
1178	4-MeNHCO-Pyr(3)	Pr	NH ₂	O	2
1179	4-MeNHCO-Pyr(3)	Pr ⁱ	NH ₂	O	2
1180	2-(Me) ₂ NCO-Pyr(3)	H	NH ₂	O	2
1181	2-(Me) ₂ NCO-Pyr(3)	Cl	NH ₂	O	2
1182	2-(Me) ₂ NCO-Pyr(3)	Me	NH ₂	O	2
1183	2-(Me) ₂ NCO-Pyr(3)	Et	NH ₂	O	2
1184	2-(Me) ₂ NCO-Pyr(3)	Pr	NH ₂	O	2
1185	2-(Me) ₂ NCO-Pyr(3)	Pr ⁱ	NH ₂	O	2

- 70 -

1186	4-(Me) ₂ NCO-Pyr(3)	H	NH ₂	O	2
1187	4-(Me) ₂ NCO-Pyr(3)	Cl	NH ₂	O	2
1188	4-(Me) ₂ NCO-Pyr(3)	Me	NH ₂	O	2
1189	4-(Me) ₂ NCO-Pyr(3)	Et	NH ₂	O	2
1190	4-(Me) ₂ NCO-Pyr(3)	Pr	NH ₂	O	2
1191	4-(Me) ₂ NCO-Pyr(3)	Pr ⁱ	NH ₂	O	2
1192	Pyr(4)	H	NH ₂	O	2
1193	Pyr(4)	H	NH ₂	O	2
1194	Pyr(4)	H	NH ₂	O	3
1195	Pyr(4)	H	NH ₂	O	4
1196	Pyr(4)	F	NH ₂	O	2
1197	Pyr(4)	Cl	NH ₂	O	2
1198	Pyr(4)	Me	NH ₂	O	2
1199	Pyr(4)	Et	NH ₂	O	2
1200	Pyr(4)	Pr	NH ₂	O	2
1201	Pyr(4)	Pr ⁱ	NH ₂	O	2
1202	Pyr(4)	Bu	NH ₂	O	2
1203	Pyr(4)	Bu ⁱ	NH ₂	O	2
1204	Pyr(4)	Bu ^s	NH ₂	O	2
1206	Pyr(4)	Bu ^t	NH ₂	O	2
1207	Pyr(4)	H	NH ₂	S	2
1208	Pyr(4)	H	NH ₂	S	3
1209	Pyr(4)	H	NH ₂	S	4
1210	Pyr(4)	F	NH ₂	S	2
1211	Pyr(4)	Cl	NH ₂	S	2
1212	Pyr(4)	Me	NH ₂	S	2
1213	Pyr(4)	Et	NH ₂	S	2
1214	Pyr(4)	Pr	NH ₂	S	2

- 71 -

1215	Pyr(4)	Pr ⁱ	NH ₂	S	2
1216	Pyr(4)	H	NHMe	O	2
1217	Pyr(4)	H	NHEt	O	2
1218	Pyr(4)	H	N(Me) ₂	O	2
1219	Pyr(4)	H	Pip(1)	O	2
1220	Pyr(4)	H	Mor(4)	O	2
1221	2-F-Pyr(4)	H	NH ₂	O	2
1222	2-F-Pyr(4)	Cl	NH ₂	O	2
1223	2-F-Pyr(4)	Me	NH ₂	O	2
1224	2-F-Pyr(4)	Et	NH ₂	O	2
1225	2-F-Pyr(4)	Pr	NH ₂	O	2
1226	2-F-Pyr(4)	Pr ⁱ	NH ₂	O	2
1227	3-F-Pyr(4)	H	NH ₂	O	2
1228	3-F-Pyr(4)	Cl	NH ₂	O	2
1229	3-F-Pyr(4)	Me	NH ₂	O	2
1230	3-F-Pyr(4)	Et	NH ₂	O	2
1231	3-F-Pyr(4)	Pr	NH ₂	O	2
1232	3-F-Pyr(4)	Pr ⁱ	NH ₂	O	2
1233	2-Cl-Pyr(4)	H	NH ₂	O	2
1234	2-Cl-Pyr(4)	Cl	NH ₂	O	2
1235	2-Cl-Pyr(4)	Me	NH ₂	O	2
1236	2-Cl-Pyr(4)	Et	NH ₂	O	2
1237	2-Cl-Pyr(4)	Pr	NH ₂	O	2
1238	2-Cl-Pyr(4)	Pr ⁱ	NH ₂	O	2
1239	3-Cl-Pyr(4)	H	NH ₂	O	2
1240	3-Cl-Pyr(4)	Cl	NH ₂	O	2
1241	3-Cl-Pyr(4)	Me	NH ₂	O	2
1242	3-Cl-Pyr(4)	Et	NH ₂	O	2

- 72 -

1243	3-Cl-Pyr(4)	Pr	NH ₂	O	2
1244	3-Cl-Pyr(4)	Pr ⁱ	NH ₂	O	2
1245	3-Me-Pyr(4)	H	NH ₂	O	2
1246	3-Me-Pyr(4)	Cl	NH ₂	O	2
1247	3-Me-Pyr(4)	Me	NH ₂	O	2
1248	3-Me-Pyr(4)	Et	NH ₂	O	2
1249	3-Me-Pyr(4)	Pr	NH ₂	O	2
1250	3-Me-Pyr(4)	Pr ⁱ	NH ₂	O	2
1251	3-Et-Pyr(4)	H	NH ₂	O	2
1252	3-Et-Pyr(4)	Cl	NH ₂	O	2
1253	3-Et-Pyr(4)	Me	NH ₂	O	2
1254	3-Et-Pyr(4)	Et	NH ₂	O	2
1255	3-Et-Pyr(4)	Pr	NH ₂	O	2
1256	3-Et-Pyr(4)	Pr ⁱ	NH ₂	O	2
1257	3-CF ₃ -Pyr(4)	H	NH ₂	O	2
1258	3-CF ₃ -Pyr(4)	Cl	NH ₂	O	2
1259	3-CF ₃ -Pyr(4)	Me	NH ₂	O	2
1260	3-CF ₃ -Pyr(4)	Et	NH ₂	O	2
1261	3-CF ₃ -Pyr(4)	Pr	NH ₂	O	2
1262	3-CF ₃ -Pyr(4)	Pr ⁱ	NH ₂	O	2
1263	3-MeO-Pyr(4)	H	NH ₂	O	2
1264	3-MeO-Pyr(4)	Cl	NH ₂	O	2
1265	3-MeO-Pyr(4)	Me	NH ₂	O	2
1266	3-MeO-Pyr(4)	Et	NH ₂	O	2
1267	3-MeO-Pyr(4)	Pr	NH ₂	O	2
1268	3-MeO-Pyr(4)	Pr ⁱ	NH ₂	O	2
1269	2-Ph-Pyr(4)	H	NH ₂	O	2
1270	2-Ph-Pyr(4)	Cl	NH ₂	O	2

- 73 -

1271	2-Ph-Pyr(4)	Me	NH ₂	O	2
1272	2-Ph-Pyr(4)	Et	NH ₂	O	2
1273	2-Ph-Pyr(4)	Pr	NH ₂	O	2
1274	2-Ph-Pyr(4)	Pr ⁱ	NH ₂	O	2
1275	3-Ph-Pyr(4)	H	NH ₂	O	2
1276	3-Ph-Pyr(4)	Cl	NH ₂	O	2
1277	3-Ph-Pyr(4)	Me	NH ₂	O	2
1278	3-Ph-Pyr(4)	Et	NH ₂	O	2
1279	3-Ph-Pyr(4)	Pr	NH ₂	O	2
1280	3-Ph-Pyr(4)	Pr ⁱ	NH ₂	O	2
1281	2-MeOCO-Pyr(4)	H	NH ₂	O	2
1282	2-MeOCO-Pyr(4)	Cl	NH ₂	O	2
1283	2-MeOCO-Pyr(4)	Me	NH ₂	O	2
1284	2-MeOCO-Pyr(4)	Et	NH ₂	O	2
1285	2-MeOCO-Pyr(4)	Pr	NH ₂	O	2
1286	2-MeOCO-Pyr(4)	Pr ⁱ	NH ₂	O	2
1287	3-MeOCO-Pyr(4)	H	NH ₂	O	2
1288	3-MeOCO-Pyr(4)	Cl	NH ₂	O	2
1289	3-MeOCO-Pyr(4)	Me	NH ₂	O	2
1290	3-MeOCO-Pyr(4)	Et	NH ₂	O	2
1291	3-MeOCO-Pyr(4)	Pr	NH ₂	O	2
1292	3-MeOCO-Pyr(4)	Pr ⁱ	NH ₂	O	2
1293	2-H ₂ NCO-Pyr(4)	H	NH ₂	O	2
1294	2-H ₂ NCO-Pyr(4)	Cl	NH ₂	O	2
1295	2-H ₂ NCO-Pyr(4)	Me	NH ₂	O	2
1296	2-H ₂ NCO-Pyr(4)	Et	NH ₂	O	2
1297	2-H ₂ NCO-Pyr(4)	Pr	NH ₂	O	2
1298	2-H ₂ NCO-Pyr(4)	Pr ⁱ	NH ₂	O	2

- 74 -

1299	3-H ₂ NCO-Pyr(4)	H	NH ₂	O	2
1300	3-H ₂ NCO-Pyr(4)	Cl	NH ₂	O	2
1301	3-H ₂ NCO-Pyr(4)	Me	NH ₂	O	2
1302	3-H ₂ NCO-Pyr(4)	Et	NH ₂	O	2
1303	3-H ₂ NCO-Pyr(4)	Pr	NH ₂	O	2
1304	3-H ₂ NCO-Pyr(4)	Pr ⁱ	NH ₂	O	2
1305	2-MeNHCO-Pyr(4)	H	NH ₂	O	2
1306	2-MeNHCO-Pyr(4)	Cl	NH ₂	O	2
1307	2-MeNHCO-Pyr(4)	Me	NH ₂	O	2
1308	2-MeNHCO-Pyr(4)	Et	NH ₂	O	2
1309	2-MeNHCO-Pyr(4)	Pr	NH ₂	O	2
1310	2-MeNHCO-Pyr(4)	Pr ⁱ	NH ₂	O	2
1311	3-MeNHCO-Pyr(4)	H	NH ₂	O	2
1312	3-MeNHCO-Pyr(4)	Cl	NH ₂	O	2
1313	3-MeNHCO-Pyr(4)	Me	NH ₂	O	2
1314	3-MeNHCO-Pyr(4)	Et	NH ₂	O	2
1315	3-MeNHCO-Pyr(4)	Pr	NH ₂	O	2
1316	3-MeNHCO-Pyr(4)	Pr ⁱ	NH ₂	O	2
1317	2-(Me) ₂ NCO-Pyr(4)	H	NH ₂	O	2
1318	2-(Me) ₂ NCO-Pyr(4)	Cl	NH ₂	O	2
1319	2-(Me) ₂ NCO-Pyr(4)	Me	NH ₂	O	2
1320	2-(Me) ₂ NCO-Pyr(4)	Et	NH ₂	O	2
1321	2-(Me) ₂ NCO-Pyr(4)	Pr	NH ₂	O	2
1322	2-(Me) ₂ NCO-Pyr(4)	Pr ⁱ	NH ₂	O	2
1323	3-(Me) ₂ NCO-Pyr(4)	H	NH ₂	O	2
1324	3-(Me) ₂ NCO-Pyr(4)	Cl	NH ₂	O	2
1325	3-(Me) ₂ NCO-Pyr(4)	Me	NH ₂	O	2
1326	3-(Me) ₂ NCO-Pyr(4)	Et	NH ₂	O	2

- 75 -

1327	3-(Me) ₂ NCO-Pyr(4)	Pr	NH ₂	O	2
1328	3-(Me) ₂ NCO-Pyr(4)	Pr ⁱ	NH ₂	O	2
1329	Pyrazinyl(2)	H	NH ₂	O	2
1330	Pyrazinyl(2)	Cl	NH ₂	O	2
1331	Pyrazinyl(2)	Me	NH ₂	O	2
1332	Pyrazinyl(2)	Et	NH ₂	O	2
1333	Pyrazinyl(2)	Pr	NH ₂	O	2
1334	Pyrazinyl(2)	Pr ⁱ	NH ₂	O	2
1335	Pyrimidinyl(2)	H	NH ₂	O	2
1336	Pyrimidinyl(2)	Cl	NH ₂	O	2
1337	Pyrimidinyl(2)	Me	NH ₂	O	2
1338	Pyrimidinyl(2)	Et	NH ₂	O	2
1339	Pyrimidinyl(2)	Pr	NH ₂	O	2
1340	Pyrimidinyl(2)	Pr ⁱ	NH ₂	O	2
1341	Pyridazinyl(3)	H	NH ₂	O	2
1342	Pyridazinyl(3)	Cl	NH ₂	O	2
1343	Pyridazinyl(3)	Me	NH ₂	O	2
1344	Pyridazinyl(3)	Et	NH ₂	O	2
1345	Pyridazinyl(3)	Pr	NH ₂	O	2
1346	Pyridazinyl(3)	Pr ⁱ	NH ₂	O	2
1347	Ph	H	NHAc	O	2
1348	Ph	H	NHMoc	O	2
1349	Ph	H	NHBz	O	2
1350	Ph	H	Pyrd(1)	O	2
1351	Ph	H	Piz(1)	O	2
1352	Ph	Cl	NHAc	O	2
1353	Ph	Cl	NHMoc	O	2
1354	Ph	Cl	NHBz	O	2
1355	Ph	Cl	Pyrd(1)	O	2

- 76 -

1356	Ph	Cl	Piz(1)	O	2
1357	Ph	Br	NH ₂	O	2
1358	Ph	Br	NH ₂	S	2
1359	Ph	I	NH ₂	O	2
1360	Ph	I	NH ₂	S	2
1361	Ph	Et	NHAc	O	2
1362	Ph	Et	NHMoc	O	2
1363	Ph	Et	NHBz	O	2
1364	Ph	Et	Pyrd(1)	O	2
1365	Ph	Et	Piz(1)	O	2
1366	Ph	Pr	NHAc	O	2
1367	Ph	Pr	NHMoc	O	2
1368	Ph	Pr	NHBz	O	2
1369	Ph	Pr	Pyrd(1)	O	2
1370	Ph	Pr	Piz(1)	O	2
1371	Ph	Pr ⁱ	NHAc	O	2
1372	Ph	Pr ⁱ	NHMoc	O	2
1373	Ph	Pr ⁱ	NHBz	O	2
1374	Ph	Pr ⁱ	Pyrd(1)	O	2
1375	Ph	Pr ⁱ	Piz(1)	O	2
1376	Ph	Bu ⁱ	NHAc	O	2
1377	Ph	Bu ⁱ	NHMoc	O	2
1378	Ph	Bu ⁱ	NHBz	O	2
1379	Ph	Bu ⁱ	Pyrd(1)	O	2
1380	Ph	Bu ⁱ	Piz(1)	O	2
1381	Ph	Bu ^t	NHAc	O	2
1382	Ph	Bu ^t	NHMoc	O	2
1383	Ph	Bu ^t	NHBz	O	2
1384	Ph	Bu ^t	Pyrd(1)	O	2

- 77 -

1385	Ph	Bu ^t	Piz(1)	O	2
1386	Ph	Pn	NH ₂	O	2
1387	Ph	Pn	NH ₂	S	2
1388	Ph	Hex	NH ₂	O	2
1389	Ph	Hex	NH ₂	S	2
1390	Ph	1-Cl-Et	NH ₂	O	2
1391	Ph	1-Cl-Et	NH ₂	S	2
1392	Ph	All	NH ₂	O	2
1393	Ph	All	NH ₂	S	2
1394	Ph	Pre ⁱ	NH ₂	O	2
1395	Ph	Pre ⁱ	NH ₂	S	2
1396	Ph	Bun(2)	NH ₂	O	2
1397	Ph	Bun(2)	NH ₂	S	2
1398	Ph	Prg	NH ₂	O	2
1399	Ph	Prg	NH ₂	S	2
1400	Ph	Pr ^c	NH ₂	O	2
1401	Ph	Pr ^c	NH ₂	S	2
1402	Ph	Pn ^c	NH ₂	O	2
1403	Ph	Pn ^c	NH ₂	S	2
1404	Ph	Pen ^c (2)	NH ₂	O	2
1405	Ph	Pen ^c (2)	NH ₂	S	2
1406	Ph	CN	NH ₂	O	2
1407	Ph	CN	NH ₂	S	2
1408	Ph	COOH	NH ₂	O	2
1409	Ph	COOH	NH ₂	S	2
1410	Ph	Ac	NH ₂	O	2
1411	Ph	Ac	NH ₂	S	2
1412	Ph	COOMe	NH ₂	O	2

- 78 -

1413	Ph	COOMe	NH ₂	S	2
1414	Ph	CONH ₂	NH ₂	O	2
1415	Ph	CONH ₂	NH ₂	S	2
1416	Ph	CONHMe	NH ₂	O	2
1417	Ph	CONHMe	NH ₂	S	2
1418	2,4-diF-Ph	Br	NH ₂	O	2
1419	2,4-diF-Ph	I	NH ₂	O	2
1420	2,4-diF-Ph	Pn	NH ₂	O	2
1421	2,4-diF-Ph	Hex	NH ₂	O	2
1422	2,4-diF-Ph	1-Cl-Et	NH ₂	O	2
1423	2,4-diF-Ph	All	NH ₂	O	2
1424	2,4-diF-Ph	Pre ⁱ	NH ₂	O	2
1425	2,4-diF-Ph	Bun(2)	NH ₂	O	2
1426	2,4-diF-Ph	Prg	NH ₂	O	2
1427	2,4-diF-Ph	Pr ^c	NH ₂	O	2
1428	2,4-diF-Ph	Pn ^c	NH ₂	O	2
1429	2,4-diF-Ph	Pen ^c (2)	NH ₂	O	2
1430	2,4-diF-Ph	CN	NH ₂	O	2
1431	2,4-diF-Ph	COOH	NH ₂	O	2
1432	2,4-diF-Ph	Ac	NH ₂	O	2
1433	2,4-diF-Ph	COOMe	NH ₂	O	2
1434	2,4-diF-Ph	CONH ₂	NH ₂	O	2
1435	2,4-diF-Ph	CONHMe	NH ₂	O	2
1436	2,4-diF-Ph	Br	NH ₂	S	2
1437	2,4-diF-Ph	I	NH ₂	S	2
1438	2,4-diF-Ph	Pn	NH ₂	S	2
1439	2,4-diF-Ph	Hex	NH ₂	S	2
1440	2,4-diF-Ph	1-Cl-Et	NH ₂	S	2

- 79 -

1441	2,4-diF-Ph	All	NH ₂	S	2
1442	2,4-diF-Ph	Pre ⁱ	NH ₂	S	2
1443	2,4-diF-Ph	Bun(2)	NH ₂	S	2
1444	2,4-diF-Ph	Prg	NH ₂	S	2
1445	2,4-diF-Ph	Pr ^c	NH ₂	S	2
1446	2,4-diF-Ph	Pn ^c	NH ₂	S	2
1447	2,4-diF-Ph	Pen ^c (2)	NH ₂	S	2
1448	2,4-diF-Ph	CN	NH ₂	S	2
1449	2,4-diF-Ph	COOH	NH ₂	S	2
1450	2,4-diF-Ph	Ac	NH ₂	S	2
1451	2,4-diF-Ph	COOMe	NH ₂	S	2
1452	2,4-diF-Ph	CONH ₂	NH ₂	S	2
1453	2,4-diF-Ph	CONHMe	NH ₂	S	2
1454	2-Cl-Ph	Br	NH ₂	O	2
1455	2-Cl-Ph	I	NH ₂	O	2
1456	2-Cl-Ph	Pn	NH ₂	O	2
1457	2-Cl-Ph	Hex	NH ₂	O	2
1458	2-Cl-Ph	1-Cl-Et	NH ₂	O	2
1459	2-Cl-Ph	All	NH ₂	O	2
1460	2-Cl-Ph	Bun(2)	NH ₂	O	2
1461	2-Cl-Ph	Pre ⁱ	NH ₂	O	2
1462	2-Cl-Ph	Prg	NH ₂	O	2
1463	2-Cl-Ph	Pr ^c	NH ₂	O	2
1464	2-Cl-Ph	Pn ^c	NH ₂	O	2
1465	2-Cl-Ph	Pen ^c (2)	NH ₂	O	2
1466	2-Cl-Ph	CN	NH ₂	O	2
1467	2-Cl-Ph	COOH	NH ₂	O	2
1468	2-Cl-Ph	Ac	NH ₂	O	2

- 80 -

1469	2-Cl-Ph	COOMe	NH ₂	O	2
1470	2-Cl-Ph	CONH ₂	NH ₂	O	2
1471	2-Cl-Ph	CONHMe	NH ₂	O	2
1472	2-Cl-Ph	Br	NH ₂	S	2
1473	2-Cl-Ph	I	NH ₂	S	2
1474	2-Cl-Ph	Pn	NH ₂	S	2
1475	2-Cl-Ph	Hex	NH ₂	S	2
1476	2-Cl-Ph	1-Cl-Et	NH ₂	S	2
1477	2-Cl-Ph	All	NH ₂	S	2
1478	2-Cl-Ph	Bun(2)	NH ₂	S	2
1479	2-Cl-Ph	Pre ⁱ	NH ₂	S	2
1480	2-Cl-Ph	Prg	NH ₂	S	2
1481	2-Cl-Ph	Pr ^c	NH ₂	S	2
1482	2-Cl-Ph	Pn ^c	NH ₂	S	2
1483	2-Cl-Ph	Pen ^c (2)	NH ₂	S	2
1484	2-Cl-Ph	CN	NH ₂	S	2
1485	2-Cl-Ph	COOH	NH ₂	S	2
1486	2-Cl-Ph	Ac	NH ₂	S	2
1487	2-Cl-Ph	COOMe	NH ₂	S	2
1488	2-Cl-Ph	CONH ₂	NH ₂	S	2
1489	2-Cl-Ph	CONHMe	NH ₂	S	2
1490	4-Cl-Ph	Br	NH ₂	O	2
1491	4-Cl-Ph	I	NH ₂	O	2
1492	4-Cl-Ph	Pn	NH ₂	O	2
1493	4-Cl-Ph	Hex	NH ₂	O	2
1494	4-Cl-Ph	1-Cl-Et	NH ₂	O	2
1495	4-Cl-Ph	All	NH ₂	O	2
1496	4-Cl-Ph	Bun(2)	NH ₂	O	2

- 81 -

1497	4-Cl-Ph	Pre ⁱ	NH ₂	O	2
1498	4-Cl-Ph	Prg	NH ₂	O	2
1499	4-Cl-Ph	Pr ^c	NH ₂	O	2
1500	4-Cl-Ph	Pn ^c	NH ₂	O	2
1501	4-Cl-Ph	Pen ^c (2)	NH ₂	O	2
1502	4-Cl-Ph	CN	NH ₂	O	2
1503	4-Cl-Ph	COOH	NH ₂	O	2
1504	4-Cl-Ph	Ac	NH ₂	O	2
1505	4-Cl-Ph	COOMe	NH ₂	O	2
1506	4-Cl-Ph	CONH ₂	NH ₂	O	2
1507	4-Cl-Ph	CONHMe	NH ₂	O	2
1508	4-Cl-Ph	Br	NH ₂	S	2
1509	4-Cl-Ph	I	NH ₂	S	2
1510	4-Cl-Ph	Pn	NH ₂	S	2
1511	4-Cl-Ph	Hex	NH ₂	S	2
1512	4-Cl-Ph	1-Cl-Et	NH ₂	S	2
1513	4-Cl-Ph	All	NH ₂	S	2
1514	4-Cl-Ph	Bun(2)	NH ₂	S	2
1515	4-Cl-Ph	Pre ⁱ	NH ₂	S	2
1516	4-Cl-Ph	Prg	NH ₂	S	2
1517	4-Cl-Ph	Pr ^c	NH ₂	S	2
1518	4-Cl-Ph	Pn ^c	NH ₂	S	2
1519	4-Cl-Ph	Pen ^c (2)	NH ₂	S	2
1520	4-Cl-Ph	CN	NH ₂	S	2
1521	4-Cl-Ph	COOH	NH ₂	S	2
1522	4-Cl-Ph	Ac	NH ₂	S	2
1523	4-Cl-Ph	COOMe	NH ₂	S	2
1524	4-Cl-Ph	CONH ₂	NH ₂	S	2

- 82 -

1525	4-Cl-Ph	CONHMe	NH ₂	S	2
1526	2,3-diCl-Ph	H	NH ₂	O	2
1527	2,3-diCl-Ph	Cl	NH ₂	O	2
1528	2,3-diCl-Ph	Et	NH ₂	O	2
1529	2,3-diCl-Ph	Pr	NH ₂	O	2
1530	2,3-diCl-Ph	Pr ⁱ	NH ₂	O	2
1531	2,3-diCl-Ph	Bu ⁱ	NH ₂	O	2
1532	2,3-diCl-Ph	Bu ^t	NH ₂	O	2
1533	2,3-diCl-Ph	H	NH ₂	S	2
1534	2,3-diCl-Ph	Cl	NH ₂	S	2
1535	2,3-diCl-Ph	Et	NH ₂	S	2
1536	2,3-diCl-Ph	Pr	NH ₂	S	2
1537	2,3-diCl-Ph	Pr ⁱ	NH ₂	S	2
1538	2,3-diCl-Ph	Bu ⁱ	NH ₂	S	2
1539	2,3-diCl-Ph	Bu ^t	NH ₂	S	2
1540	2,4-diCl-Ph	Br	NH ₂	O	2
1541	2,4-diCl-Ph	I	NH ₂	O	2
1542	2,4-diCl-Ph	Pn	NH ₂	O	2
1543	2,4-diCl-Ph	Hex	NH ₂	O	2
1544	2,4-diCl-Ph	1-Cl-Et	NH ₂	O	2
1545	2,4-diCl-Ph	All	NH ₂	O	2
1546	2,4-diCl-Ph	Bun(2)	NH ₂	O	2
1547	2,4-diCl-Ph	Pre ⁱ	NH ₂	O	2
1548	2,4-diCl-Ph	Prg	NH ₂	O	2
1549	2,4-diCl-Ph	Pr ^c	NH ₂	O	2
1550	2,4-diCl-Ph	Pn ^c	NH ₂	O	2
1551	2,4-diCl-Ph	Pen ^c (2)	NH ₂	O	2
1552	2,4-diCl-Ph	CN	NH ₂	O	2

- 83 -

1553	2,4-diCl-Ph	COOH	NH ₂	O	2
1554	2,4-diCl-Ph	Ac	NH ₂	O	2
1555	2,4-diCl-Ph	COOMe	NH ₂	O	2
1556	2,4-diCl-Ph	CONH ₂	NH ₂	O	2
1557	2,4-diCl-Ph	CONHMe	NH ₂	O	2
1558	2,4-diCl-Ph	Br	NH ₂	S	2
1559	2,4-diCl-Ph	I	NH ₂	S	2
1560	2,4-diCl-Ph	Pn	NH ₂	S	2
1561	2,4-diCl-Ph	Hex	NH ₂	S	2
1562	2,4-diCl-Ph	1-Cl-Et	NH ₂	S	2
1563	2,4-diCl-Ph	All	NH ₂	S	2
1564	2,4-diCl-Ph	Bun(2)	NH ₂	S	2
1565	2,4-diCl-Ph	Pre ⁱ	NH ₂	S	2
1566	2,4-diCl-Ph	Prg	NH ₂	S	2
1567	2,4-diCl-Ph	Pr ^c	NH ₂	S	2
1568	2,4-diCl-Ph	Pn ^c	NH ₂	S	2
1569	2,4-diCl-Ph	Pen ^c (2)	NH ₂	S	2
1570	2,4-diCl-Ph	CN	NH ₂	S	2
1571	2,4-diCl-Ph	COOH	NH ₂	S	2
1572	2,4-diCl-Ph	Ac	NH ₂	S	2
1573	2,4-diCl-Ph	COOMe	NH ₂	S	2
1574	2,4-diCl-Ph	CONH ₂	NH ₂	S	2
1575	2,4-diCl-Ph	CONHMe	NH ₂	S	2
1576	2,4-diCl-3-Me-Ph	H	NH ₂	O	2
1577	2,4-diCl-3-Me-Ph	Cl	NH ₂	O	2
1578	2,4-diCl-3-Me-Ph	Et	NH ₂	O	2
1579	2,4-diCl-3-Me-Ph	Pr	NH ₂	O	2
1580	2,4-diCl-3-Me-Ph	Pr ⁱ	NH ₂	O	2

- 84 -

1581	2,4-diCl-3-Me-Ph	Bu ⁱ	NH ₂	O	2
1582	2,4-diCl-3-Me-Ph	Bu ^t	NH ₂	O	2
1583	2,4-diCl-3-Me-Ph	H	NH ₂	S	2
1584	2,4-diCl-3-Me-Ph	Cl	NH ₂	S	2
1585	2,4-diCl-3-Me-Ph	Et	NH ₂	S	2
1586	2,4-diCl-3-Me-Ph	Pr	NH ₂	S	2
1587	2,4-diCl-3-Me-Ph	Pr ⁱ	NH ₂	S	2
1588	2,4-diCl-3-Me-Ph	Bu ⁱ	NH ₂	S	2
1589	2,4-diCl-3-Me-Ph	Bu ^t	NH ₂	S	2
1590	2,4-diCl-3-Et-Ph	H	NH ₂	O	2
1591	2,4-diCl-3-Et-Ph	Cl	NH ₂	O	2
1592	2,4-diCl-3-Et-Ph	Et	NH ₂	O	2
1593	2,4-diCl-3-Et-Ph	Pr	NH ₂	O	2
1594	2,4-diCl-3-Et-Ph	Pr ⁱ	NH ₂	O	2
1595	2,4-diCl-3-Et-Ph	Bu ⁱ	NH ₂	O	2
1596	2,4-diCl-3-Et-Ph	Bu ^t	NH ₂	O	2
1597	2,4-diCl-3-Et-Ph	H	NH ₂	S	2
1598	2,4-diCl-3-Et-Ph	Cl	NH ₂	S	2
1599	2,4-diCl-3-Et-Ph	Et	NH ₂	S	2
1600	2,4-diCl-3-Et-Ph	Pr	NH ₂	S	2
1601	2,4-diCl-3-Et-Ph	Pr ⁱ	NH ₂	S	2
1602	2,4-diCl-3-Et-Ph	Bu ⁱ	NH ₂	S	2
1603	2,4-diCl-3-Et-Ph	Bu ^t	NH ₂	S	2
1604	3,4-diCl-Ph	H	NH ₂	O	2
1605	3,4-diCl-Ph	Cl	NH ₂	O	2
1606	3,4-diCl-Ph	Et	NH ₂	O	2
1607	3,4-diCl-Ph	Pr	NH ₂	O	2
1608	3,4-diCl-Ph	Pr ⁱ	NH ₂	O	2

- 85 -

1609	3,4-diCl-Ph	Bu ⁱ	NH ₂	O	2
1610	3,4-diCl-Ph	Bu ^t	NH ₂	O	2
1611	3,4-diCl-Ph	H	NH ₂	S	2
1612	3,4-diCl-Ph	Cl	NH ₂	S	2
1613	3,4-diCl-Ph	Et	NH ₂	S	2
1614	3,4-diCl-Ph	Pr	NH ₂	S	2
1615	3,4-diCl-Ph	Pr ⁱ	NH ₂	S	2
1616	3,4-diCl-Ph	Bu ⁱ	NH ₂	S	2
1617	3,4-diCl-Ph	Bu ^t	NH ₂	S	2
1618	4-Pr ⁱ -Ph	H	NH ₂	O	2
1619	4-Pr ⁱ -Ph	Cl	NH ₂	O	2
1620	4-Pr ⁱ -Ph	Et	NH ₂	O	2
1621	4-Pr ⁱ -Ph	Pr	NH ₂	O	2
1622	4-Pr ⁱ -Ph	Pr ⁱ	NH ₂	O	2
1623	4-Pr ⁱ -Ph	Bu ⁱ	NH ₂	O	2
1624	4-Pr ⁱ -Ph	Bu ^t	NH ₂	O	2
1625	4-Pr ⁱ -Ph	H	NH ₂	S	2
1626	4-Pr ⁱ -Ph	Cl	NH ₂	S	2
1627	4-Pr ⁱ -Ph	Et	NH ₂	S	2
1628	4-Pr ⁱ -Ph	Pr	NH ₂	S	2
1629	4-Pr ⁱ -Ph	Pr ⁱ	NH ₂	S	2
1630	4-Pr ⁱ -Ph	Bu ⁱ	NH ₂	S	2
1631	4-Pr ⁱ -Ph	Bu ^t	NH ₂	S	2
1632	4-PhO-Ph	H	NH ₂	O	2
1633	4-PhO-Ph	Cl	NH ₂	O	2
1634	4-PhO-Ph	Et	NH ₂	O	2
1635	4-PhO-Ph	Pr	NH ₂	O	2
1636	4-PhO-Ph	Pr ⁱ	NH ₂	O	2

- 86 -

1637	4-PhO-Ph	Bu ⁱ	NH ₂	O	2
1638	4-PhO-Ph	Bu ^t	NH ₂	O	2
1639	4-PhO-Ph	H	NH ₂	S	2
1640	4-PhO-Ph	Cl	NH ₂	S	2
1641	4-PhO-Ph	Et	NH ₂	S	2
1642	4-PhO-Ph	Pr	NH ₂	S	2
1643	4-PhO-Ph	Pr ⁱ	NH ₂	S	2
1644	4-PhO-Ph	Bu ⁱ	NH ₂	S	2
1645	4-PhO-Ph	Bu ^t	NH ₂	S	2
1646	4-BnO-Ph	H	NH ₂	O	2
1647	4-BnO-Ph	Cl	NH ₂	O	2
1648	4-BnO-Ph	Et	NH ₂	O	2
1649	4-BnO-Ph	Pr	NH ₂	O	2
1650	4-BnO-Ph	Pr ⁱ	NH ₂	O	2
1651	4-BnO-Ph	Bu ⁱ	NH ₂	O	2
1652	4-BnO-Ph	Bu ^t	NH ₂	O	2
1653	4-BnO-Ph	H	NH ₂	S	2
1654	4-BnO-Ph	Cl	NH ₂	S	2
1655	4-BnO-Ph	Et	NH ₂	S	2
1656	4-BnO-Ph	Pr	NH ₂	S	2
1657	4-BnO-Ph	Pr ⁱ	NH ₂	S	2
1658	4-BnO-Ph	Bu ⁱ	NH ₂	S	2
1659	4-BnO-Ph	Bu ^t	NH ₂	S	2
1660	4-NO ₂ -Ph	H	NH ₂	O	2
1661	4-NO ₂ -Ph	Cl	NH ₂	O	2
1662	4-NO ₂ -Ph	Et	NH ₂	O	2
1663	4-NO ₂ -Ph	Pr	NH ₂	O	2
1664	4-NO ₂ -Ph	Pr ⁱ	NH ₂	O	2

- 87 -

1665	4-NO ₂ -Ph	Bu ⁱ	NH ₂	O	2
1666	4-NO ₂ -Ph	Bu ^t	NH ₂	O	2
1667	4-NO ₂ -Ph	H	NH ₂	S	2
1668	4-NO ₂ -Ph	Cl	NH ₂	S	2
1669	4-NO ₂ -Ph	Et	NH ₂	S	2
1670	4-NO ₂ -Ph	Pr	NH ₂	S	2
1671	4-NO ₂ -Ph	Pr ⁱ	NH ₂	S	2
1672	4-NO ₂ -Ph	Bu ⁱ	NH ₂	S	2
1673	4-NO ₂ -Ph	Bu ^t	NH ₂	S	2
1674	4-OH-Ph	H	NH ₂	O	2
1675	4-OH-Ph	Cl	NH ₂	O	2
1676	4-OH-Ph	Et	NH ₂	O	2
1677	4-OH-Ph	Pr	NH ₂	O	2
1678	4-OH-Ph	Pr ⁱ	NH ₂	O	2
1679	4-OH-Ph	Bu ⁱ	NH ₂	O	2
1680	4-OH-Ph	Bu ^t	NH ₂	O	2
1681	4-OH-Ph	H	NH ₂	S	2
1682	4-OH-Ph	Cl	NH ₂	S	2
1683	4-OH-Ph	Et	NH ₂	S	2
1684	4-OH-Ph	Pr	NH ₂	S	2
1685	4-OH-Ph	Pr ⁱ	NH ₂	S	2
1686	4-OH-Ph	Bu ⁱ	NH ₂	S	2
1687	4-OH-Ph	Bu ^t	NH ₂	S	2
1688	4-AcO-Ph	H	NH ₂	O	2
1689	4-AcO-Ph	Cl	NH ₂	O	2
1690	4-AcO-Ph	Et	NH ₂	O	2
1691	4-AcO-Ph	Pr	NH ₂	O	2
1692	4-AcO-Ph	Pr ⁱ	NH ₂	O	2

- 88 -

1693	4-AcO-Ph	Bu ⁱ	NH ₂	O	2
1694	4-AcO-Ph	Bu ^t	NH ₂	O	2
1695	4-AcO-Ph	H	NH ₂	S	2
1696	4-AcO-Ph	Cl	NH ₂	S	2
1697	4-AcO-Ph	Et	NH ₂	S	2
1698	4-AcO-Ph	Pr	NH ₂	S	2
1699	4-AcO-Ph	Pri	NH ₂	S	2
1700	4-AcO-Ph	Bu ⁱ	NH ₂	S	2
1701	4-AcO-Ph	Bu ^t	NH ₂	S	2
1702	4-NH ₂ -Ph	H	NH ₂	O	2
1703	4-NH ₂ -Ph	Cl	NH ₂	O	2
1704	4-NH ₂ -Ph	Et	NH ₂	O	2
1705	4-NH ₂ -Ph	Pr	NH ₂	O	2
1706	4-NH ₂ -Ph	Pri	NH ₂	O	2
1707	4-NH ₂ -Ph	Bu ⁱ	NH ₂	O	2
1708	4-NH ₂ -Ph	Bu ^t	NH ₂	O	2
1709	4-NH ₂ -Ph	H	NH ₂	S	2
1710	4-NH ₂ -Ph	Cl	NH ₂	S	2
1711	4-NH ₂ -Ph	Et	NH ₂	S	2
1712	4-NH ₂ -Ph	Pr	NH ₂	S	2
1713	4-NH ₂ -Ph	Pri	NH ₂	S	2
1714	4-NH ₂ -Ph	Bu ⁱ	NH ₂	S	2
1715	4-NH ₂ -Ph	Bu ^t	NH ₂	S	2
1716	4-BzHN-Ph	H	NH ₂	O	2
1717	4-BzHN-Ph	Cl	NH ₂	O	2
1718	4-BzHN-Ph	Et	NH ₂	O	2
1719	4-BzHN-Ph	Pr	NH ₂	O	2
1720	4-BzHN-Ph	Pri	NH ₂	O	2

- 89 -

1721	4-BzHN-Ph	Bu ⁱ	NH ₂	O	2
1722	4-BzHN-Ph	Bu ^t	NH ₂	O	2
1723	4-BzHN-Ph	H	NH ₂	S	2
1724	4-BzHN-Ph	Cl	NH ₂	S	2
1725	4-BzHN-Ph	Et	NH ₂	S	2
1726	4-BzHN-Ph	Pr	NH ₂	S	2
1727	4-BzHN-Ph	Pr ⁱ	NH ₂	S	2
1728	4-BzHN-Ph	Bu ⁱ	NH ₂	S	2
1729	4-BzHN-Ph	Bu ^t	NH ₂	S	2
1730	Fur(2)	Br	NH ₂	O	2
1731	Fur(2)	I	NH ₂	O	2
1732	Fur(2)	Pn	NH ₂	O	2
1733	Fur(2)	Hex	NH ₂	O	2
1734	Fur(2)	1-Cl-Et	NH ₂	O	2
1735	Fur(2)	All	NH ₂	O	2
1736	Fur(2)	Pre ⁱ	NH ₂	O	2
1737	Fur(2)	Bun(2)	NH ₂	O	2
1738	Fur(2)	Prg	NH ₂	O	2
1739	Fur(2)	Pr ^c	NH ₂	O	2
1740	Fur(2)	Pn ^c	NH ₂	O	2
1741	Fur(2)	Pen ^c (2)	NH ₂	O	2
1742	Fur(2)	CN	NH ₂	O	2
1743	Fur(2)	COOH	NH ₂	O	2
1744	Fur(2)	Ac	NH ₂	O	2
1745	Fur(2)	COOMe	NH ₂	O	2
1746	Fur(2)	CONH ₂	NH ₂	O	2
1747	Fur(2)	CONHMe	NH ₂	O	2
1748	Fur(2)	Br	NH ₂	S	2

- 90 -

1749	Fur(2)	I	NH ₂	S	2
1750	Fur(2)	Pn	NH ₂	S	2
1751	Fur(2)	Hex	NH ₂	S	2
1752	Fur(2)	1-Cl-Et	NH ₂	S	2
1753	Fur(2)	All	NH ₂	S	2
1754	Fur(2)	Pre ⁱ	NH ₂	S	2
1755	Fur(2)	Bun(2)	NH ₂	S	2
1756	Fur(2)	Prg	NH ₂	S	2
1757	Fur(2)	Pr ^c	NH ₂	S	2
1758	Fur(2)	Pn ^c	NH ₂	S	2
1759	Fur(2)	Pen ^c (2)	NH ₂	S	2
1760	Fur(2)	CN	NH ₂	S	2
1761	Fur(2)	COOH	NH ₂	S	2
1762	Fur(2)	Ac	NH ₂	S	2
1763	Fur(2)	COOMe	NH ₂	S	2
1764	Fur(2)	CONH ₂	NH ₂	S	2
1765	Fur(2)	CONHMe	NH ₂	S	2
1766	Thi(2)	Br	NH ₂	O	2
1767	Thi(2)	I	NH ₂	O	2
1768	Thi(2)	Pn	NH ₂	O	2
1769	Thi(2)	Hex	NH ₂	O	2
1770	Thi(2)	1-Cl-Et	NH ₂	O	2
1771	Thi(2)	All	NH ₂	O	2
1772	Thi(2)	Pre ⁱ	NH ₂	O	2
1773	Thi(2)	Bun(2)	NH ₂	O	2
1774	Thi(2)	Prg	NH ₂	O	2
1775	Thi(2)	Pr ^c	NH ₂	O	2
1776	Thi(2)	Pn ^c	NH ₂	O	2

- 91 -

1777	Thi(2)	Pen ^C (2)	NH ₂	O	2
1778	Thi(2)	CN	NH ₂	O	2
1779	Thi(2)	COOH	NH ₂	O	2
1780	Thi(2)	Ac	NH ₂	O	2
1781	Thi(2)	COOMe	NH ₂	O	2
1782	Thi(2)	CONH ₂	NH ₂	O	2
1783	Thi(2)	CONHMe	NH ₂	O	2
1784	Thi(2)	Br	NH ₂	S	2
1785	Thi(2)	I	NH ₂	S	2
1786	Thi(2)	Pn	NH ₂	S	2
1787	Thi(2)	Hex	NH ₂	S	2
1788	Thi(2)	1-Cl-Et	NH ₂	S	2
1789	Thi(2)	All	NH ₂	S	2
1790	Thi(2)	Pre ⁱ	NH ₂	S	2
1791	Thi(2)	Bun(2)	NH ₂	S	2
1792	Thi(2)	Prg	NH ₂	S	2
1793	Thi(2)	Pr ^C	NH ₂	S	2
1794	Thi(2)	Pn ^C	NH ₂	S	2
1795	Thi(2)	Pen ^C (2)	NH ₂	S	2
1796	Thi(2)	CN	NH ₂	S	2
1797	Thi(2)	COOH	NH ₂	S	2
1798	Thi(2)	Ac	NH ₂	S	2
1799	Thi(2)	COOMe	NH ₂	S	2
1800	Thi(2)	CONH ₂	NH ₂	S	2
1801	Thi(2)	CONHMe	NH ₂	S	2
1802	Ph	1-Cl-Pr	NH ₂	O	2
1803	Ph	1-Cl-Pr	NH ₂	S	2
1804	Ph	1-Cl-Bu ⁱ	NH ₂	O	2

1805	Ph	1-Cl-Bu ⁱ	NH ₂	S	2
1806	Ph	1-Cl-Pn ⁱ	NH ₂	O	2
1807	Ph	1-Cl-Pn ⁱ	NH ₂	S	2
1808	2,4-diMe-Ph	H	NH ₂	O	2
1809	2,4-diMe-Ph	Pr ⁱ	NH ₂	O	2
1810	3,5-diMe-Ph	H	NH ₂	O	2
1811	3,5-diMe-Ph	Pr ⁱ	NH ₂	O	2

Of the isoxazole derivatives illustrated in the Table invention, preferred compounds are 1, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 25, 27, 28, 29, 30, 32, 36, 48, 50, 54, 66, 70, 74, 93, 95, 99, 111, 112, 113, 114, 115, 116, 117, 125, 127, 131, 143, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 188, 190, 194, 206, 208, 212, 224, 226, 230, 242, 244, 248, 260, 262, 263, 264, 266, 278, 296, 298, 314, 316, 332, 334, 350, 357, 363, 368, 379, 386, 397, 408, 469, 475, 481, 505, 510, 511, 517, 523, 535, 538, 539, 540, 541, 542, 543, 544, 545, 546, 562, 568, 574, 580, 586, 592, 598, 604, 610, 616, 622, 628, 724, 728, 729, 730, 731, 732, 733, 752, 764, 776, 788, 794, 800, 806, 812, 818, 824, 1056, 1061, 1347, 1348, 1349, 1350, 1351, 1357, 1359, 1386, 1388, 1390, 1392, 1394, 1396, 1398, 1400, 1402, 1404, 1406, 1408, 1410, 1412, 1414, 1416, 1459, 1495, 1499, 1500, 1526, 1545, 1549, 1550, 1576, 1590, 1604, 1618, 1632, 1646, 1660, 1674, 1688, 1702, 1716, 1809 or 1811,

more preferred compounds are 1, 4, 5, 6, 7, 8, 9, 11, 13, 14, 15, 30, 32, 36, 48, 50, 54, 66, 70, 74, 93, 95, 99, 111, 113, 117, 125, 127, 131, 143, 147, 149, 150, 151, 170, 172, 176, 188, 190, 194, 206, 208, 212, 224, 226, 230, 242, 244, 248, 260, 262, 263, 264, 266, 278, 296, 298, 314, 316, 332, 334, 350, 357, 363, 368, 379, 386, 397, 408, 469, 475, 481, 505, 510, 511, 517, 523, 535, 538, 539, 540, 541, 542, 543, 544, 545, 546, 562, 568, 574, 580, 586, 592, 598, 604, 610, 616, 622, 628, 724, 728, 729, 730, 731, 732, 733, 752, 764, 776, 788, 794, 800, 806, 812, 818, 824, 1056, 1061, 1392, 1394, 1398, 1809 or 1811,

still more preferred compounds are 1, 4, 5, 6, 7, 8, 9, 11, 13, 14, 15, 30, 48, 66, 74, 93, 111, 117, 125, 143, 149, 150, 151, 170, 176, 188, 206, 224, 242, 260, 296, 314, 332, 350, 368, 386, 408, 469, 475, 481, 505, 510, 511, 517, 523, 535, 538, 539, 543, 568, 586, 598, 604, 622, 724, 733, 1392, 1394 or 1398, and

particularly preferred compounds are 1, 4, 5, 6, 7, 8, 9, 11, 13, 66, 93, 111, 117, 125, 143, 149, 150, 151, 170, 176, 224, 260, 332, 386, 510, 535, 539, 543, 604, 1392, 1394 or 1398.

The most preferred compounds are :

Compound list No. 1: 3-(2-aminoethoxy)-5-phenylisoxazole,

Compound list No. 5: 3-(2-aminoethoxy)-4-chloro-5-phenylisoxazole,

Compound list No. 7: 3-(2-aminoethoxy)-4-ethyl-5-phenylisoxazole,

Compound list No. 8: 3-(2-aminoethoxy)-5-phenyl-4-propylisoxazole,

Compound list No. 9: 3-(2-aminoethoxy)-4-isopropyl-5-phenylisoxazole,

Compound list No. 11: 3-(2-aminoethoxy)-4-isobutyl-5-phenylisoxazole,

Compound list No. 117: 3-(2-aminoethoxy)-5-(2-chlorophenyl)-4-isopropylisoxazole,

Compound list No. 143: 3-(2-aminoethoxy)-5-(4-chlorophenyl)isoxazole,

Compound list No. 151: 3-(2-aminoethoxy)-5-(4-chlorophenyl)-4-isopropylisoxazole,

Compound list No. 176: 3-(2-aminoethoxy)-5-(2,4-dichlorophenyl)-4-isopropylisoxazole,

Compound list No. 510: 3-(2-aminoethoxy)-5-(2-furyl)-4-isopropylisoxazole,

Compound list No. 535: 3-(2-aminoethoxy)-5-(2-thienyl)isoxazole,

Compound list No. 539: 3-(2-aminoethoxy)-4-chloro-5-(2-thienyl)isoxazole,


Compound list No. 543: 3-(2-aminoethoxy)-4-isopropyl-5-(2-thienyl)isoxazole,

or


Compound list No. 1392: 4-allyl-3-(2-aminoethoxy)-5-phenylisoxazole.

Methods for preparing the compounds of the present invention are illustrated below.


Method A



Method B



Method C



Wherein R¹, R², R³, X and n have the same meanings as defined above,

R^3_a has the same meanings as defined for R^3 with the proviso that the amino group in R^3 is protected, Y represents a hydroxyl group or a leaving group, and Z represents a halogen atom.

The protecting group of the amino group or the mono $C_1 - C_6$ alkylamino group of R^3_a can be used without particular limitation so long as it is a group generally used as a protecting group for an amino group, and may be, for example, a $C_1 - C_6$ alkanoyl group such as the formyl, acetyl, propionyl, butyryl, isobutyryl, pentanoyl, pivaloyl, valeryl, isovaleryl or hexanoyl groups; a $C_1 - C_4$ alkanoyl group substituted with halogen or $C_1 - C_4$ alkoxy, such as the chloroacetyl, dichloroacetyl, trichloroacetyl, trifluoroacetyl, 3-fluoropropionyl, 4,4-dichlorobutyryl, methoxyacetyl, butoxyacetyl, ethoxypropionyl or propoxybutyryl groups; an unsaturated $C_1 - C_4$ alkanoyl group such as the acryloyl, propioloyl, methacryloyl, crotonoyl or isocrotonoyl groups; a $C_6 - C_{10}$ arylcarbonyl group optionally substituted with halogen, $C_1 - C_4$ alkyl, $C_1 - C_4$ alkoxy, $C_1 - C_4$ alkoxy carbonyl, $C_6 - C_{10}$ aryl or nitro such as the benzoyl, α -naphthoyl, β -naphthoyl, 2-fluorobenzoyl, 2-bromobenzoyl, 2,4-dichlorobenzoyl, 6-chloro- α -naphthoyl, 4-toluoyl, 4-propylbenzoyl, 4-t-butylbenzoyl, 2,4,6-trimethylbenzoyl, 6-ethyl- α -naphthoyl, 4-anisoyl, 4-propoxybenzoyl, 4-t-butoxybenzoyl, 6-ethoxy- α -naphthoyl, 2-ethoxycarbonylbenzoyl, 4-t-butoxycarbonylbenzoyl, 6-methoxycarbonyl- α -naphthoyl, 4-phenylbenzoyl, 4-phenyl- α -naphthoyl, 6- α -naphthylbenzoyl, 4-nitrobenzoyl, 2-nitrobenzoyl group or 6-nitro- α -naphthoyl groups; a $C_1 - C_4$ alkoxy carbonyl group optionally substituted with halogen or tri- $C_1 - C_4$ alkylsilyl such as the methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, isopropoxycarbonyl, butoxycarbonyl, isobutoxycarbonyl, s-butoxycarbonyl, t-butoxycarbonyl, chloromethoxycarbonyl, 2,2,2-trichloroethoxycarbonyl, 2-fluoropropoxycarbonyl, 2-bromo-t-butoxycarbonyl, 2,2-dibromo-t-butoxycarbonyl, triethylsilylmethoxycarbonyl, 2-trimethylsilylethoxycarbonyl, 4-tripropylsilylbutoxycarbonyl or t-butyl-dimethylsilylpropoxycarbonyl groups; a $C_2 - C_5$ alkenyloxycarbonyl group

such as the vinyloxy carbonyl, allyloxy carbonyl, 1,3-butadienyloxy carbonyl or 2-pentyloxy carbonyl groups; an aryldicarbonyl group such as a phthaloyl group; an aralkyl group such as the benzyl, phenethyl, 3-phenylpropyl, 4-phenylbutyl, α -naphthylmethyl, β -naphthylmethyl, diphenylmethyl, triphenylmethyl, or α -naphthyl diphenylmethyl group or 9-anthrylmethyl groups; or a C₇ - C₁₅ aralkyloxy carbonyl group optionally substituted with methoxy or nitro such as the benzyloxy carbonyl, (1-phenyl)benzyloxy carbonyl, α -naphthylmethoxy carbonyl, β -naphthylmethoxy carbonyl, 9-anthrylmethoxy carbonyl, p-methoxybenzyloxy carbonyl or p-nitrobenzyloxy carbonyl groups, preferably, the C₁ - C₄ alkanoyl; trifluoroacetyl; methoxyacetyl; benzoyl group; α -naphthoyl; β -naphthoyl; anisoyl; nitrobenzoyl; the C₁ - C₄ alkoxy carbonyl; methoxy carbonyl; ethoxy carbonyl; t-butoxy carbonyl; 2,2,2-trichloroethoxy carbonyl; triethylsilylmethoxy carbonyl; 2-trimethylsilylethoxy carbonyl; vinyloxy carbonyl group; allyloxy carbonyl; phthaloyl; benzyl; benzyloxy carbonyl; or nitrobenzyloxy carbonyl groups, more preferably the formyl, acetyl, benzoyl, 4-anisoyl, 4-nitrobenzoyl, methoxy carbonyl, ethoxy carbonyl, butoxy carbonyl, t-butoxy carbonyl, phthaloyl, benzyl, benzyloxy carbonyl or p-nitrobenzyloxy carbonyl groups, most preferably the t-butoxy carbonyl group.

The leaving group of Y is not particularly limited so long as it is a usual leaving group as a nucleophilic residual group, and may be, for example, a halogen atom such as the chlorine, bromine or iodine atoms; a C₁ - C₄ alkanesulfonyloxy group such as the methanesulfonyloxy, ethanesulfonyloxy, propanesulfonyloxy or butanesulfonyloxy groups; a halogeno C₁ - C₄ alkanesulfonyloxy group such as the trifluoromethanesulfonyloxy, 2,2,2-trichloroethanesulfonyloxy, 3,3,3-tribromopropanesulfonyloxy or 4,4,4-trifluorobutanesulfonyloxy groups; or a C₆ - C₁₀ arylsulfonyloxy group optionally having from one to three C₁ - C₄ alkyls such as the benzenesulfonyloxy, α -naphthylsulfonyloxy, β -naphthylsulfonyloxy, p-toluenesulfonyloxy, 4-t-butylbenzenesulfonyloxy, mesitylenesulfonyloxy or 6-ethyl- α -naphthylsulfonyloxy groups, and preferably the chlorine, bromine or iodine

atoms; the methanesulfonyloxy, ethanesulfonyloxy, trifluoromethanesulfonyloxy, 2,2,2-trichloroethanesulfonyloxy, benzenesulfonyloxy, toluenesulfonyloxy or mesitylenesulfonyloxy groups, more preferably the chlorine, bromine or iodine atoms, the methanesulfonyloxy, trifluoromethanesulfonyloxy, benzenesulfonyloxy, p-toluenesulfonyloxy or mesitylenesulfonyloxy groups.

The halogen atom of Z may be, for example, fluorine, chlorine, bromine or iodine atoms, and preferably the chlorine atom.

Method A is a method for synthesizing the compound of general formula (I).

In step A1 a compound of formula (III) is prepared by reacting a compound (II) with a halogenating agent in an inert solvent or in the absence of a solvent (preferably in an inert solvent) in the presence or absence of a base (preferably in the presence of a base).

The solvent to be employed here is not particularly limited so long as it does not interfere with the reaction and can dissolve a certain amount of the starting material. Examples of suitable solvents may be aliphatic hydrocarbons such as hexane, heptane, ligroin or petroleum ether; aromatic hydrocarbons such as benzene, toluene or xylene; halogenated hydrocarbons such as methylene chloride, chloroform, carbon tetrachloride, dichloroethane, chlorobenzene or dichlorobenzene; or ethers such as diethyl ether, diisopropyl ether, tetrahydrofuran, dioxane, dimethoxyethane or diethylene glycol dimethyl ether, preferably halogenated hydrocarbons (particularly methylene chloride) or ethers (particularly tetrahydrofuran or dioxane).

The base to be employed here may be, for example, alkali metal carbonates such as sodium carbonate, potassium carbonate or lithium carbonate; alkali metal hydrogencarbonates such as sodium hydrogencarbonate, potassium hydrogencarbonate or lithium hydrogencarbonate; or organic amines such as triethylamine, tributylamine, diisopropylethylamine, N-methylmorpholine, pyridine, 4-(N,N-dimethylamino)pyridine, N,N-dimethylaniline, N,N-diethyl-

aniline, 1,5-diazabicyclo[4.3.0]non-5-ene (DBN), 1,4-diazabicyclo[2.2.2]octane (DABCO) or 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU), preferably alkali metal carbonates or organic amines, more preferably organic amines (particularly triethylamine or pyridine).

The halogenating agent to be employed here may be, for example, phosphorus oxychloride, phosphorus oxybromide, phosphorus oxyiodide or phosphorus pentachloride, or a mixture thereof, preferably phosphorus oxychloride, phosphorus pentachloride or a mixture thereof.

The reaction temperature may be varied depending on the nature of the starting material and reagent, and is usually from 0°C to 150°C, preferably from 10°C to 100°C.

The reaction time may be varied depending on the nature of the starting material, reagent and reaction temperature and is usually from 30 minutes to 10 hours, preferably from 1 to 5 hours.

After the completion of the reaction, the target compound of this step is isolated from the reaction mixture according to a conventional method. For example, the target compound is obtained by removing the solvent by evaporation, adding water to the reaction mixture, adding a hydrophobic solvent (for example, benzene, ether, ethyl acetate, etc.) to the resulting mixture to extract the target compound, washing the extracted organic layer with water, drying it over anhydrous magnesium sulfate and removing the solvent by evaporation. The target compound thus obtained may be further purified, if necessary, according to a conventional method, for example, recrystallization, reprecipitation or chromatography.

In step A2 a compound of formula (V) is prepared by reacting the compound (III) with a compound of the general formula (IV) in an inert solvent in the presence of a base.

The solvent to be employed here is not particularly limited so long as it does not interfere with the reaction and can dissolve a certain amount of the starting material. Examples of suitable solvents may be aliphatic hydrocarbons such as hexane, heptane, ligroin or petroleum ether; aromatic hydrocarbons such

as benzene, toluene or xylene; halogenated hydrocarbons such as methylene chloride, chloroform, carbon tetrachloride, dichloroethane, chlorobenzene or dichlorobenzene; ethers such as diethyl ether, diisopropyl ether, tetrahydrofuran, dioxane, dimethoxyethane or diethylene glycol dimethyl ether; amides such as formamide, dimethylformamide, dimethylacetamide or hexamethylphosphoric triamide; or sulfoxides such as dimethyl sulfoxide or sulfolane, preferably the ethers, amides or sulfoxides, more preferably the ethers (particularly diethyl ether, tetrahydrofuran or dioxane) or the amides (particularly dimethylformamide).

The base to be employed here may be, for example, alkali metal carbonates such as sodium carbonate, potassium carbonate or lithium carbonate; alkali metal hydrogencarbonates such as sodium hydrogencarbonate, potassium hydrogencarbonate or lithium hydrogencarbonate; alkali metal hydrides such as lithium hydride, sodium hydride or potassium hydride; alkali metal hydroxides such as sodium hydroxide, potassium hydroxide or lithium hydroxide; organic amines such as triethylamine, tributylamine, diisopropylethylamine, N-methylmorpholine, pyridine, 4-(N,N-dimethylamino)pyridine, N,N-dimethylaniline, N,N-diethylaniline, 1,5-diazabicyclo[4.3.0]non-5-ene (DBN), 1,4-diazabicyclo[2.2.2]octane (DABCO) or 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU); alkylolithiums such as methyllithium, ethyllithium or butyllithium; or lithium alkylamides such as lithium diisopropylamide or lithium dicyclohexylamide, preferably the alkali metal carbonates, alkali metal hydrides or organic amines, more preferably the alkali metal carbonates (particularly sodium carbonate or potassium carbonate) or the alkali metal hydrides (particularly sodium hydride).

Incidentally, in order to effectively carry out the reaction, crown ethers such as dibenzo-18-crown-6 can be also added to the reaction mixture.

The reaction temperature may be varied depending on the nature of the starting material and reagent, and is usually from -10°C to 150°C , preferably from 0°C to 80°C .

The reaction time may be varied depending on the nature of the starting material, reagent and reaction temperature and is usually from 30 minutes to 30 hours, preferably from 1 to 10 hours.

After the completion of the reaction, the target compound of this step is isolated from the reaction mixture according to a conventional method. For example, the target compound is obtained by appropriately neutralizing the reaction mixture, removing the insolubles by filtration, if necessary, in the case where the insolubles exist, removing the solvent by evaporation, adding water to the reaction mixture, adding a hydrophobic solvent (for example, benzene, ether, ethyl acetate, etc.) to the resulting mixture to extract the target compound, washing the extracted organic layer with water, drying the organic layer containing the desired product over anhydrous magnesium sulfate and removing the solvent by evaporation. The target compound thus obtained may be further purified, if necessary, according to a conventional method, for example, recrystallization, reprecipitation or chromatography.

In step A3 a compound of general formula (I) is prepared by removing the protecting group of the amino group or the alkylamino group, if necessary.

Removal of the protecting group of the amino group varies depending on the kind thereof and is generally carried out as follows according to a conventional method in organic chemistry.

In the case where the protecting group of the amino group is a C₁ - C₆ alkanoyl group (preferably formyl or acetyl groups); a C₆ - C₁₀ arylcarbonyl group (preferably a benzoyl group); a C₁ - C₄ alkoxy carbonyl group optionally substituted with halogen or a tri-C₁ - C₄ alkylsilyl group (preferably the methoxycarbonyl, ethoxycarbonyl, t-butoxycarbonyl, 2-trimethylsilylethoxycarbonyl, 2-bromo-t-butoxycarbonyl or 2,2-dibromo-t-butoxycarbonyl groups); a C₂ - C₅ alkenyloxy carbonyl group (preferably a vinyloxy carbonyl group); or a C₇ - C₁₅ aralkyloxy carbonyl group optionally substituted with methoxy or nitro (preferably the benzyloxy carbonyl, (1-phenyl)benzyloxy carbonyl, 9-anthrylmethoxycarbonyl, p-methoxybenzyloxy carbonyl or p-nitrobenzyloxy carbonyl groups), it can be removed by treating it with an acid in an inert solvent or an

aqueous solvent. Further, in this case, the target compound can be also obtained as a salt. The acid to be employed here may be, for example, an acid such as hydrochloric acid, sulfuric acid, phosphoric acid, hydrobromic acid or trifluoroacetic acid, preferably hydrochloric acid, sulfuric acid, hydrobromic acid or trifluoroacetic acid.

The solvent to be employed here is not particularly limited so long as it does not interfere with the reaction and can dissolve a certain amount of the starting material. Examples of suitable solvents may be aliphatic hydrocarbons such as hexane, heptane, ligroin or petroleum ether; aromatic hydrocarbons such as benzene, toluene or xylene; halogenated hydrocarbons such as methylene chloride, chloroform, carbon tetrachloride, dichloroethane, chlorobenzene or dichlorobenzene; ethers such as diethyl ether, diisopropyl ether, tetrahydrofuran, dioxane, dimethoxyethane or diethylene glycol dimethyl ether; esters such as methyl acetate or ethyl acetate; alcohols such as methanol, ethanol, propanol, isopropanol or butanol; amides such as formamide, dimethylformamide, dimethylacetamide or hexamethylphosphoric triamide; sulfoxides such as dimethyl sulfoxide or sulfolane; aliphatic acids such as formic acid or acetic acid; water; or an aqueous mixture of the above-mentioned solvent, preferably the halogenated hydrocarbons, ethers, alcohols, aliphatic acids or an aqueous mixture of the above-mentioned solvent, more preferably the halogenated hydrocarbons (particularly methylene chloride), ethers (particularly tetrahydrofuran or dioxane), aliphatic acids (particularly acetic acid), water; or an aqueous mixture of the above-mentioned solvent.

The reaction temperature may be varied depending on the nature of the starting material, solvent or acid used and is usually from -10°C to 150°C , preferably from 0°C to 60°C .

The reaction time may be varied depending on the nature of the starting material, solvent or acid used, and is usually from 5 minutes to 20 hours, preferably from 10 minutes to 5 hours.

After the completion of the reaction, the target compound of this step is isolated from the reaction mixture according to a conventional method. For example, the target compound is obtained by collecting the precipitated target

compound in the reaction mixture by filtration or appropriately neutralizing the reaction mixture, removing the solvent by evaporation, adding water to the reaction mixture, adding a hydrophobic solvent (for example, benzene, ether, ethyl acetate, etc.) to the resulting mixture to extract the target compound, washing the extracted organic layer with water, drying the organic layer containing the target compound over anhydrous magnesium sulfate and removing the solvent by evaporation. The target compound thus obtained can be further purified, if necessary, by a conventional method, for example, recrystallization, reprecipitation or chromatography.

In the case where the protecting group of the amino group is an alkanoyl, arylcarbonyl, alkoxycarbonyl, alkenyloxycarbonyl, aryldicarbonyl, aralkyl or aralkyloxycarbonyl, it can be removed by treating it with a base in an inert solvent or an aqueous solvent.

The base to be employed here may be, for example, an alkali metal carbonate such as sodium carbonate, potassium carbonate or lithium carbonate; an alkali metal hydrogencarbonate such as sodium hydrogencarbonate, potassium hydrogencarbonate or lithium hydrogencarbonate; an alkali metal hydride such as lithium hydride, sodium hydride or potassium hydride; an alkali metal hydroxide such as sodium hydroxide, potassium hydroxide or lithium hydroxide; an alkali metal alkoxide such as sodium methoxide, sodium ethoxide, potassium t-butoxide or lithium methoxide; or an alkali metal mercaptan such as methyl mercaptan sodium or ethyl mercaptan sodium, preferably the alkali metal carbonates (particularly the sodium carbonate or potassium carbonate), the alkali metal hydroxides (particularly the sodium hydroxide or potassium hydroxide), the alkali metal alkoxides (particularly the sodium methoxide, sodium ethoxide or potassium t-butoxide) or organic amines (particularly hydrazine or methylamine).

The solvent to be employed here is not particularly limited so long as it does not interfere with the reaction and can dissolve a certain amount of the starting material. Examples of suitable solvents may be aliphatic hydrocarbons such as hexane, heptane, ligroin or petroleum ether; aromatic hydrocarbons such

as benzene, toluene or xylene; ethers such as diethyl ether, diisopropyl ether, tetrahydrofuran, dioxane, dimethoxyethane or diethylene glycol dimethyl ether; alcohols such as methanol, ethanol, propanol, isopropanol or butanol; amides such as dimethylacetamide or hexamethylphosphoric triamide; sulfoxides such as dimethyl sulfoxide or sulfolane; or an aqueous mixture of the above-mentioned solvent, preferably the halogenated hydrocarbons, ethers, alcohols, or an aqueous mixture of the above-mentioned solvent, more preferably the ethers (particularly the tetrahydrofuran or dioxane), the alcohols (particularly the methanol or ethanol) or an aqueous mixture of the above-mentioned solvent.

The reaction temperature may be varied depending on the nature of the starting material, solvent or base used and is usually from -10°C to 150°C , preferably from 0°C to 50°C .

The reaction time may be varied depending on the nature of the starting material, solvent or base used, and is usually from 30 minutes to 20 hours, preferably from 1 to 5 hours.

After completion of the reaction, the target compound of this step is isolated from the reaction mixture according to a conventional method. For example, the target compound is obtained by separating the precipitated target compound from the reaction mixture by filtration or removing the solvent by evaporation, adding water to the reaction mixture, separating the precipitate from the mixture by filtration, or adding a hydrophobic solvent (for example, benzene, ether, ethyl acetate, etc.) to the resulting mixture to extract the target compound, washing the extracted organic layer containing the target compound with water, drying it over anhydrous magnesium sulfate and removing the solvent by evaporation. The target compound thus obtained may be further purified, if necessary, according to a conventional method, for example, recrystallization, reprecipitation or chromatography.

Further, in the case where the protecting group of the amino group is a tert-butoxycarbonyl group, it can be removed by treating it with a silyl compound or an acid particularly in an inert solvent.

The silyl compound to be employed here may be, for example,

trimethylsilyl chloride, trimethylsilyl iodide or trimethylsilyl trifluoromethanesulfonate.

The acid employed here may be, for example, aluminum chloride, hydrochloric acid or trifluoroacetic acid.

The solvent to be employed here is not particularly limited so long as it does not interfere with the reaction and can dissolve a certain amount of the starting material. Examples of suitable solvents may be halogenated hydrocarbons such as methylene chloride, chloroform or carbon tetrachloride; ethers such as diethyl ether, tetrahydrofuran or dioxane; and nitriles such as acetonitrile, preferably the halogenated hydrocarbons (particularly methylene chloride or chloroform) or the nitriles (particularly acetonitrile).

The reaction temperature may be varied depending on the nature of the starting material, reagent or solvent and is usually from -20°C to 100°C , preferably from 0°C to 50°C .

The reaction time may be varied depending on the nature of the starting material, reagent, solvent or reaction temperature, and is usually from 10 minutes to 10 hours, preferably from 30 minutes to 3 hours.

After the completion of the reaction, the target compound of this step is isolated from the reaction mixture according to a conventional method. For example, the target compound is obtained by separating the precipitated desired compound in the reaction mixture by filtration, or adding water to the reaction mixture, making the aqueous layer alkaline to separate the precipitated substance from the mixture by filtration or adding a hydrophobic solvent (for example, benzene, ether, ethyl acetate, etc.) to the resulting mixture to extract the desired compound, washing the organic layer containing the desired compound with water, drying it over anhydrous magnesium sulfate and removing the solvent by evaporation. The target compound thus obtained can be further purified, if necessary, by a usual method, for example, recrystallization, reprecipitation or chromatography.

Further, in the case where the protecting group of the amino group is an allyloxycarbonyl group, it can be removed using palladium and

triphenylphosphine or nickel tetracarbonyl under similar reaction condition such as solvent, reaction temperature, reaction time and the like to those of the removal reaction of the aralkyl group, etc. by catalytic reduction reagent.

In the case where the protecting group of the amino group is an aralkyl group or a C₇ - C₁₁ aralkyloxycarbonyl group, the protecting group can be removed easily by contacting it with a reductant (preferably a catalytic hydrogenation reaction in the presence of a catalyst) in an inert solvent or by a removal reaction using an oxidant.

In the case of the removal reaction of the protecting group by a catalytic hydrogenation reaction in a catalytic reduction, the solvent to be employed here is not particularly limited so long as it does not interfere with the present reaction. Examples of suitable solvents may be aliphatic hydrocarbons such as hexane or cyclohexane; aromatic hydrocarbons such as toluene, benzene or xylene; ethers such as diethyl ether, tetrahydrofuran or dioxane; esters such as ethyl acetate or propyl acetate; alcohols such as methanol, ethanol or isopropanol; aliphatic acids such as formic acid or acetic acid; or an aqueous mixture of the above solvent, preferably the aliphatic hydrocarbons, aromatic hydrocarbons, ethers, esters, alcohols, aliphatic acids or an aqueous mixture of the above solvent, more preferably the alcohols (particularly the methanol or ethanol), the aliphatic acids (particularly the formic acid or acetic acid) or an aqueous mixture of the above solvent.

The catalyst to be employed here is not particularly limited so long as it is used for usual catalytic reduction reaction. Examples of suitable catalysts may be palladium black, palladium-carbon, Raney nickel, rhodium-aluminum oxide or palladium-barium sulfate, preferably the palladium-carbon or Raney nickel.

The pressure of hydrogen is not particularly limited, and is usually between 1 and 10 atmospheric pressure, preferably it is 1 atmospheric pressures.

The reaction temperature may be varied depending on the nature of the starting material, solvent or reductant used and is usually from 0°C to 100°C,

preferably from 10°C to 50°C.

The reaction time may be varied depending on the nature of the starting material, solvent, reductant or reaction temperature, and is usually from 15 minutes to 10 hours, preferably from 30 minutes to 3 hours.

After the completion of the reaction, the target compound of this step is isolated from the reaction mixture according to a conventional method. For example, the target compound is obtained by removing the catalyst by filtration, removing the solvent by evaporation, adding water to the reaction mixture, making the aqueous layer alkaline to separate the precipitate from the mixture by filtration or adding a hydrophobic solvent (for example, benzene, ether, ethyl acetate, etc.) to the resulting mixture to extract the desired compound, washing the organic layer containing the desired compound with water, drying it over anhydrous magnesium sulfate and removing the solvent by evaporation. The target compound thus obtained can be further purified, if necessary, by a conventional method, for example, recrystallization, reprecipitation or chromatography.

In a removal reaction of protecting group by oxidation the solvent to be employed is not particularly limited so long as it does not interfere with the present reaction. Examples of suitable solvents may be ketones such as acetone; halogenated hydrocarbons such as methylene chloride, chloroform or carbon tetrachloride; nitriles such as acetonitrile; ethers such as diethyl ether, tetrahydrofuran or dioxane; amides such as dimethylformamide, dimethylacetamide or hexamethylphosphoric triamide; sulfoxides such as dimethyl sulfoxide or an aqueous mixture of the above solvent, preferably the ketones, halogenated hydrocarbons, nitriles, ethers, amides, sulfoxides or an aqueous mixture of the above solvent, more preferably the ketones (particularly the acetone), the halogenated hydrocarbons (particularly the methylene chloride), the nitriles (particularly the acetonitrile), the amides (particularly the hexamethylphosphoric triamide), the sulfoxides (particularly the dimethyl sulfoxide) or an aqueous mixture of the above solvent.

The oxidant to be employed here may be, for example, potassium

persulfate, sodium persulfate, ammonium cerium nitrate (CAN) or 2,3-dichloro-5,6-dicyano-p-benzoquinone (DDQ), preferably ammonium cerium nitrate (CAN) or 2,3-dichloro-5,6-dicyano-p-benzoquinone (DDQ).

The reaction temperature may be varied depending on the nature of the starting material, solvent or oxidant used and is usually from 0°C to 150°C, preferably from 10°C to 50°C.

The reaction time may be varied depending on the nature of the starting material, solvent and oxidant used and is usually from 15 minutes to 24 hours, preferably from 30 minutes to 5 hours.

After the completion of the reaction, the target compound of this step is isolated from the reaction mixture according to a conventional method. For example, the target compound is obtained by removing the oxidant by filtration, removing the solvent by evaporation, adding water to the reaction mixture, making the aqueous layer alkaline to separate the precipitate by filtration or adding a hydrophobic solvent (for example, benzene, ether, ethyl acetate, etc.) to the resulting mixture to extract the desired compound, washing the organic layer containing the desired compound with water, drying it over anhydrous magnesium sulfate and removing the solvent by evaporation. The target compound thus obtained can be further purified, if necessary, by a conventional method, for example, recrystallization, reprecipitation or chromatography.

Further, in the case where the protecting group of the amino group is removed using an acid, the desired compound is usually obtained in the form of a salt, but the amino group of the desired compound can be made to a free base by removing the acid used according to a conventional method in organic chemistry.

Method B is an alternative method for synthesizing the compound (Va) in which X is an oxygen atom in the intermediate compound (V) of Process A.

In step B1 a compound of formula (Va) is prepared by reacting the compound of formula (II) with the compound of formula (IVa).

In the case where Y is a hydroxyl group, the reaction is carried out by dehydration-condensation between the compound (II) and the corresponding compound (IVa) in an inert solvent in the presence of phosphine-compound and azo-compound as conducted based on the Mitsunobu reaction reported in Bull. Chem. Soc. Jap., 40, 2380 (1967).

The solvent to be employed here is not particularly limited so long as it does not interfere with the reaction and can dissolve a certain amount of the starting material. Examples of suitable solvents may be aliphatic hydrocarbons such as hexane, heptane, ligroin or petroleum ether; aromatic hydrocarbons such as benzene, toluene or xylene; halogenated hydrocarbons such as methylene chloride, chloroform, carbon tetrachloride, dichloroethane, chlorobenzene or dichlorobenzene; or ethers such as diethyl ether, diisopropyl ether, tetrahydrofuran, dioxane, dimethoxyethane or diethylene glycol dimethyl ether, preferably the aliphatic hydrocarbons, aromatic hydrocarbons or ethers, more preferably the ethers (particularly the diethyl ether or tetrahydrofuran).

The phosphine-compound to be employed here may be, for example, a tri-C₁ - C₆ alkylphosphine such as trimethylphosphine, triethylphosphine, tripropylphosphine, tributylphosphine, tripentylphosphine or trihexylphosphine; a tri-C₆ - C₁₀ arylphosphine such as triphenylphosphine, triindenylphosphine or trinaphthylphosphine; or a tri-C₆ - C₁₀ arylphosphine optionally having C₁ - C₄ alkyl as a substituent such as tolyldiphenylphosphine, tritolylphosphine, trimesitylphosphine, tributylphenylphosphine or tri-6-ethyl-2-naphthylphosphine, preferably the tri-C₁ - C₆ alkylphosphines (particularly trimethylphosphine, triethylphosphine, tripropylphosphine or tributylphosphine) or tri-C₆ - C₁₀ arylphosphine (particularly triphenylphosphine, triindenylphosphine or trinaphthylphosphine), more preferably the tri-C₆ - C₁₀ arylphosphines (particularly triphenylphosphine).

The azo-compound to be employed here is not particularly limited so long as it is a known azodicarboxylic acid derivative, and may be, for example, a di-C₁ - C₄ alkyl azodicarboxylate such as dimethyl azodicarboxylate, diethyl

azodicarboxylate, dipropyl azodicarboxylate or dibutyl azodicarboxylate, preferably dimethyl azodicarboxylate or diethyl azodicarboxylate.

The reaction temperature may be varied depending on the nature of the starting material and reagent, and is usually from -10°C to 100°C , preferably from 0°C to 50°C .

The reaction time may be varied depending on the nature of the starting material, reagent and reaction temperature, and is usually from 15 minutes to 48 hours, preferably from 30 minutes to 24 hours.

After the completion of the reaction, the target compound of this step is isolated from the reaction mixture according to a conventional method. For example, the target compound is obtained by filtering off the insolubles in the case where they exist, removing the solvent by evaporation, or adding water to the residue obtained by removing the solvent by evaporation, adding a hydrophobic solvent (for example, benzene, ether, ethyl acetate, etc.) to the resulting mixture to effect extraction, washing the extract with water, drying over anhydrous magnesium sulfate and removing the solvent by evaporation. The target compound thus obtained can be further purified, if necessary, by a conventional method, for example, recrystallization, reprecipitation or chromatography.

In the case where Y is a leaving group, the compound (Va) can be prepared by reacting the compound (II) with the corresponding compound (IVa) in the presence of a base in an inert solvent.

The solvent to be employed here is not particularly limited so long as it does not interfere with the reaction and can dissolve a certain amount of the starting material. Examples of suitable solvents may be aliphatic hydrocarbons such as hexane, heptane, ligroin or petroleum ether; aromatic hydrocarbons such as benzene, toluene or xylene; halogenated hydrocarbons such as methylene chloride, chloroform, carbon tetrachloride, dichloroethane, chlorobenzene or dichlorobenzene; ethers such as diethyl ether, diisopropyl ether, tetrahydrofuran, dioxane, dimethoxyethane or diethylene glycol dimethyl ether; amides such as formamide, dimethylformamide, dimethylacetamide or hexamethylphosphoric

triamide; or sulfoxides such as dimethyl sulfoxide or sulfolane, preferably the amides and sulfoxides, more preferably the amides (particularly dimethylformamide, dimethylacetamide or hexamethylphosphoric triamide).

The base to be employed here may be, for example, an alkali metal carbonate such as sodium carbonate, potassium carbonate or lithium carbonate; an alkali metal hydrogencarbonate such as sodium hydrogencarbonate, potassium hydrogencarbonate or lithium hydrogencarbonate; an alkali metal hydride such as lithium hydride, sodium hydride or potassium hydride; an alkali metal hydroxide such as sodium hydroxide, potassium hydroxide or lithium hydroxide; an alkali metal alkoxide such as sodium methoxide, sodium ethoxide, potassium t-butoxide or lithium methoxide; an alkali metal mercaptan such as methylmercaptan sodium or ethylmercaptan sodium; an organic amine such as triethylamine, tributylamine, diisopropylethylamine, N-methylmorpholine, pyridine, 4-(N,N-dimethylamino)pyridine, N,N-dimethylaniline, N,N-diethylaniline, 1,5-diazabicyclo[4.3.0]non-5-ene, 1,4-diazabicyclo[2.2.2]octane (DABCO) or 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU); an alkyllithium such as methyllithium, ethyllithium or butyllithium; or a lithium alkylamide such as lithium diisopropylamide or lithium dicyclohexylamide, preferably the alkali metal carbonates, alkali metal hydrides or alkali metal hydroxides, more preferably the alkali metal hydrides (particularly sodium hydroxide).

A crown ether such as dibenzo-18-crown-6 may be added to enhance the reaction.

The reaction temperature may be varied depending on the nature of the starting material, the reagent, etc., and is usually from -10°C to 100°C, preferably from 0°C to 50°C.

The reaction time may be varied depending on the nature of the starting material, the reagent and the reaction temperature, and is usually from 30 minutes to 20 hours, preferably from 1 to 5 hours.

After the completion of the reaction, the target compound of this step is isolated from the reaction mixture according to a conventional method. For example, the target compound is obtained by appropriately neutralizing the reaction mixture, removing the insolubles by filtration in the case where they

exist, removing the solvent by evaporation, adding water to the reaction mixture, adding a hydrophobic solvent (for example, benzene, ether, ethyl acetate, etc.) to the resulting mixture to effect extraction, washing the organic layer including the target compound with water, drying it over anhydrous magnesium sulfate and removing the solvent by evaporation. The target compound thus obtained can be further purified, if necessary, by a conventional method, for example, recrystallization, reprecipitation or chromatography.

Method C is an alternative method for synthesizing a compound (I) of method A.

Step C1 is carried out, if necessary, and includes

reaction (a): the reaction in which an alkyl group, an alkoxy group or a carboxyl group is introduced to an isoxazole ring or an aromatic ring included in R¹,

reaction (b): the reaction in which a hydroxyalkyl group is introduced to an isoxazole ring or an aromatic ring included in R¹,

reaction (c): the reaction in which a hydroxyl group contained in the hydroxyalkyl group produced by the reaction (b) is converted to a halogen atom,

reaction (d): the reaction in which a hydroxyl group contained in the hydroxyalkyl group produced by the reaction (b) is subjected to a 1,2-elimination reaction (β elimination),

reaction (e): the reaction in which the hydroxyalkyl group produced by the reaction (b) is converted to a carbonyl group,

reaction (f): the reaction in which a carboxyl group is esterified,

reaction (g): the reaction in which an alkoxy carbonyl group is converted to a carbamoyl group,

reaction (h): the reaction in which a carboxyl group is converted to a carbamoyl group,

reaction (i): the reaction in which a carbamoyl group is converted to a cyano group,

- reaction (j): the reaction in which an alkoxy group on the aromatic ring is converted to a hydroxyl group,
- reaction (k): the reaction in which a hydroxyl group or an amino group is subjected to an acylation reaction,
- reaction (l): the reaction in which a hydroxyl group or an amino group is subjected to an aralkylation reaction,
- reaction (m): the reaction in which a nitro group is converted to an amino group, and
- reaction (n): the reaction in which the protecting group of the amino group included in R^3_a is removed.

These reaction are carried out by appropriately in any order.

Reaction (a):

The reaction in which an alkyl group, an alkoxy group or a carboxyl group is introduced to an isoxazole ring or an aromatic ring in R^1 is carried out according to a conventional method in organic chemistry. For example, the reaction is carried out by reacting a halo $C_1 - C_6$ alkane, di- $C_1 - C_6$ alkylcarbonate or carbon dioxide (preferably a halo $C_1 - C_6$ alkane or carbon dioxide) in the presence of a base in an inert solvent.

The halo $C_1 - C_6$ alkane to be employed here may be, for example, methyl chloride, methyl bromide, methyl iodide, ethyl chloride, ethyl iodide, propyl bromide, butyl iodide, pentyl iodide or hexyl iodide, preferably methyl bromide or methyl iodide, more preferably methyl iodide.

The di- $C_1 - C_6$ alkylcarbonate may be, for example, dimethyl carbonate, diethyl carbonate, dipropyl carbonate, diisopropyl carbonate, dibutyl carbonate, di-sec-butyl carbonate, di-tert-butyl carbonate, dipentyl carbonate or dihexyl carbonate, preferably dimethyl carbonate or diethyl carbonate.

The solvent to be employed here is not particularly limited so long as it does not interfere with the reaction and can dissolve a certain amount of the starting material. Examples of suitable solvents may be aliphatic hydrocarbons

such as hexane, heptane, ligroin or petroleum ether; aromatic hydrocarbons such as benzene, toluene or xylene; halogenated hydrocarbons such as methylene chloride, chloroform, carbon tetrachloride, dichloroethane, chlorobenzene or dichlorobenzene; ethers such as diethyl ether, diisopropyl ether, tetrahydrofuran, dioxane, dimethoxyethane or diethylene glycol dimethyl ether; diamines such as N,N,N',N'-tetramethylethylenediamine; amides such as formamide, dimethylformamide, dimethylacetamide or hexamethylphosphoric triamide or hexamethylphosphorous triamide; or sulfoxides such as dimethyl sulfoxide or sulfolane, preferably the ethers, amides or sulfoxides, more preferably the ethers (particularly tetrahydrofuran).

The base to be employed here may be, for example, an alkali metal hydride such as lithium hydride, sodium hydride or potassium hydride; an alkyl lithium such as methyl lithium, ethyl lithium, butyl lithium or sec-butyl lithium; or a lithium alkylamide such as lithium diisopropylamide, lithium dicyclohexylamide, lithium bis(trimethylsilyl)amide, potassium bis(trimethylsilyl)amide or sodium bis(trimethylsilyl)amide, preferably the alkyl lithiums (particularly butyl lithium) or lithium alkylamides (particularly lithium diisopropylamide).

The reaction temperature may be varied depending on the nature of the starting material, the reagent, etc., and is usually from -100°C to 30°C, preferably from -70°C to 0°C.

The reaction time may be varied depending on the nature of the starting material, the reagent and the reaction temperature, and is usually from 5 minutes to 10 hours, preferably from 10 minutes to 5 hours.

After the completion of the reaction, the target compound of this step is isolated from the reaction mixture according to a conventional method. For example, the target compound is obtained by removing the solvent by evaporation, or adding water to the residue obtained by removing the solvent by evaporation, making the aqueous layer acidic, if desired, adding a hydrophobic solvent (for example, benzene, ether, ethyl acetate, etc.) to the resulting mixture to extract the desired compound, washing the extracted organic layer with water, drying it over anhydrous magnesium sulfate and removing the solvent by

evaporation. The target compound thus obtained can be further purified, if necessary, by a conventional method, for example, recrystallization, reprecipitation or chromatography.

Reaction (b):

The reaction in which a hydroxylalkyl group is introduced on an isoxazole ring or an aromatic ring in R¹ is carried out according to a conventional method in organic chemistry. For example, the reaction is carried out by reacting the compounds with aldehydes or ketones in the presence of a base in an inert solvent.

The aldehydes may be, for example, a straight or branched alkanal having from 2 to 6 carbon atoms such as acetaldehyde, propionaldehyde, butylaldehyde, isobutylaldehyde, valeraldehyde, isovaleraldehyde or hexaldehyde, preferably a C₂ - C₄ alkanal, more preferably acetaldehyde.

The ketones may be, for example, a straight or branched alkanone having from 3 to 6 carbon atoms such as acetone, 2-butanone, 2-pentanone, 3-pentanone, 3-methyl-2-butanone, 2-hexanone, 3-hexanone, 3-methyl-2-pentanone, 4-methyl-2-pentanone or 3,3-dimethyl-2-butanone, preferably the acetone, 2-butanone or 3-butanone, more preferably acetone.

The solvent to be employed here is not particularly limited so long as it does not interfere with the reaction and can dissolve a certain amount of the starting material. Examples of suitable solvent may be aliphatic hydrocarbons such as hexane, heptane, ligroin or petroleum ether; aromatic hydrocarbons such as benzene, toluene or xylene; halogenated hydrocarbons such as methylene chloride, chloroform, carbon tetrachloride, dichloroethane, chlorobenzene or dichlorobenzene; ethers such as diethyl ether, diisopropyl ether, tetrahydrofuran, dioxane, dimethoxyethane or diethylene glycol dimethyl ether; diamines such as N,N,N',N'-tetramethylethylenediamine; amides such as formamide, dimethylformamide, dimethylacetamide, hexamethylphosphoric triamide or hexamethylphosphorous triamide; or sulfoxides such as dimethyl sulfoxide or sulfolane, preferably the ethers, amides or sulfoxides, more preferably the ethers (particularly tetrahydrofuran).

The base to be employed here may be, for example, an alkali metal hydride such as lithium hydride, sodium hydride or potassium hydride; an alkyllithium such as methyllithium, ethyllithium, butyllithium or sec-butyl-lithium; or a lithium alkylamide such as lithium diisopropylamide, lithium dicyclohexylamide, lithium bis(trimethylsilyl)amide, potassium bis(trimethylsilyl)amide or sodium bis(trimethylsilyl)amide, preferably the alkyllithiums (particularly butyllithium) or lithium alkylamides (particularly lithium diisopropylamide).

The reaction temperature may be varied depending on the nature of the starting material, the reagent, etc., and is usually from -100°C to 30°C , preferably from -70°C to 0°C .

The reaction time may be varied depending on the nature of the starting material, the reagent and the reaction temperature, and is usually from 5 minutes to 10 hours, preferably from 10 minutes to 5 hours.

After the completion of the reaction, the target compound of this step is isolated from the reaction mixture according to a conventional method. For example, the target compound is obtained by removing the solvent by evaporation, or adding water to the residue obtained by removing the solvent by evaporation, adding a hydrophobic solvent (for example, benzene, ether, ethyl acetate, etc.) to the resulting mixture to extract the desired compound, washing the extracted organic layer with water, drying it over anhydrous magnesium sulfate and removing the solvent by evaporation. The target compound thus obtained can be further purified, if necessary, by a conventional method, for example, recrystallization, reprecipitation or chromatography.

Reaction (c):

The reaction in which a hydroxyl group in a hydroxyalkyl group produced by the reaction (b) is converted to a halogen atom is carried out according to a conventional method in organic chemistry. For example, the reaction is carried out by reacting a hydroxyl group with a hydrohalogenic acid in an inert solvent.

The hydrohalogenic acid to be employed may be, for example,

hydrofluoric acid, hydrochloric acid, hydrobromic acid or hydroiodic acid, preferably hydrochloric acid.

The solvent to be employed here is not particularly limited so long as it does not interfere with the reaction and can dissolve a certain amount of the starting material. Examples of suitable solvents may be aliphatic hydrocarbons such as hexane, heptane, ligroin or petroleum ether; aromatic hydrocarbons such as benzene, toluene or xylene; halogenated hydrocarbons such as methylene chloride, chloroform, carbon tetrachloride, dichloroethane, chlorobenzene or dichlorobenzene; ethers such as diethyl ether, diisopropyl ether, tetrahydrofuran, dioxane, dimethoxyethane or diethylene glycol dimethyl ether; esters such as methyl acetate or ethyl acetate; water, or an aqueous mixture of the above-mentioned solvent, preferably the ethers (particularly dioxane) or a mixture of ethers and water.

The reaction temperature may be varied depending on the nature of the starting material and reagent, and is usually from -50°C to 80°C , preferably from 0°C to 50°C .

The reaction time may be varied depending on the nature of the starting material, reagent and reaction temperature, and is usually from 5 minutes to 10 hours, preferably from 10 minutes to 5 hours.

After the completion of the reaction, the target compound of this step is isolated from the reaction mixture according to a conventional method. For example, after the reaction, the target compound is obtained by removing the solvent by evaporation, or adding water to the residue obtained by removing the solvent by evaporation, adding a hydrophobic solvent (for example, benzene, ether, ethyl acetate, etc.) to the resulting mixture to extract the desired compound, washing the extracted organic layer with water, drying it over anhydrous magnesium sulfate and removing the solvent by evaporation. The target compound thus obtained can be further purified, if necessary, by a conventional method, for example, recrystallization, reprecipitation or chromatography.

Reaction (d):

The reaction in which a hydroxyl group in the hydroxyalkyl group produced by the reaction (b) is subjected to 1,2-elimination reaction (β elimination) is carried out according to a conventional method in organic chemistry. For example, the reaction is carried out by reacting the hydroxyl group with an acid in an inert solvent.

The acid to be employed here may be, for example, a mineral acid such as hydrofluoric acid, hydrochloric acid, hydrobromic acid, hydroiodic acid, nitric acid, perchloric acid, sulfuric acid or phosphoric acid; a sulfonic acid such as methanesulfonic acid, trifluoromethanesulfonic acid, ethanesulfonic acid, benzenesulfonic acid or p-toluenesulfonic acid; or a carboxylic acid such as trifluoroacetic acid, fumaric acid, succinic acid, citric acid, tartaric acid, oxalic acid or maleic acid, preferably the mineral acids (particularly hydrochloric acid).

The solvent to be employed here is not particularly limited so long as it does not interfere with the reaction and can dissolve a certain amount of the starting material. Examples of suitable solvents may be aliphatic hydrocarbons such as hexane, heptane, ligroin or petroleum ether; aromatic hydrocarbons such as benzene, toluene or xylene; halogenated hydrocarbons such as methylene chloride, chloroform, carbon tetrachloride, dichloroethane, chlorobenzene or dichlorobenzene; ethers such as diethyl ether, diisopropyl ether, tetrahydrofuran, dioxane, dimethoxyethane or diethylene glycol dimethyl ether; or esters such as methyl acetate or ethyl acetate, preferably the ethers or esters, more preferably the ethers (particularly dioxane).

The reaction temperature may be varied depending on the nature of the starting material and reagent, and is usually from 0°C to 150°C, preferably from 50°C to 130°C.

The reaction time may be varied depending on the nature of the starting material, reagent and reaction temperature, and is usually from 5 minutes to 10 hours, preferably from 10 minutes to 5 hours.

After the completion of the reaction, the target compound of this step is isolated from the reaction mixture according to a conventional method. For example, after the reaction, the target compound is obtained by removing the solvent by evaporation, or adding water to the residue obtained by removing the

solvent by evaporation, adding a hydrophobic solvent (for example, benzene, ether, ethyl acetate, etc.) to the resulting mixture to extract the desired compound, washing the extracted organic layer with water, drying it over anhydrous magnesium sulfate and removing the solvent by evaporation. The target compound thus obtained can be further purified, if necessary, by a conventional method, for example, recrystallization, reprecipitation or chromatography.

Reaction (e):

The reaction in which the hydroxyalkyl group produced by the reaction (b) is converted to a carbonyl group is carried out according to a conventional method in organic chemistry. For example, the reaction is carried out by reacting the hydroxyalkyl group with an oxidant in an inert solvent.

The oxidant to be employed here is not particularly limited so long as it is used for an oxidation reaction, and may be, for example, an inorganic metal oxidant such as manganese oxides, e.g. potassium permanganate or manganese dioxide; a ruthenium oxide, e.g. ruthenium tetroxide; a silver compound such as silver oxide; a chromic acid compound, e.g. potassium chromate, chromic acid-sulfuric acid complex or chromic acid-pyridine complex, or a cerium compound such as ammonium cerium nitrate (CAN); a reagent used for DMSO oxidation (a complex of dimethyl sulfoxide and dicyclohexylcarbodiimide, oxalyl chloride, acetic anhydride or phosphorus pentoxide or a complex of pyridine-sulfuric anhydride); a succinimide such as N-bromosuccinimide; or a quinone compound such as 2,3-dichloro-5,6-dicyano-p-benzoquinone (DDQ), preferably the inorganic metal oxidants or the reagents used for DMSO oxidation, more preferably the inorganic metal oxidants (particularly the chromic acid-pyridine complex).

The solvent to be employed here is not particularly limited so long as it does not interfere with the reaction and can dissolve a certain amount of the starting material. Examples of suitable solvents may be aliphatic hydrocarbons such as hexane, heptane, ligroin or petroleum ether; aromatic hydrocarbons such as benzene, toluene or xylene; halogenated hydrocarbons such as methylene

chloride, chloroform, carbon tetrachloride, dichloroethane, chlorobenzene or dichlorobenzene; ethers such as diethyl ether, diisopropyl ether, tetrahydrofuran, dioxane, dimethoxyethane or diethylene glycol dimethyl ether; ketones such as acetone; esters such as methyl acetate or ethyl acetate; nitriles such as acetonitrile; amides such as dimethylformamide, dimethylacetamide or hexamethylphosphoric triamide; sulfoxides such as dimethyl sulfoxide; or an aqueous mixture of the above solvent, preferably the halogenated hydrocarbons, ethers, ketones, nitriles, amides or sulfoxides, more preferably the halogenated hydrocarbons (particularly methylene chloride).

The reaction temperature may be varied depending on the nature of the starting material and reagent, and is usually from 0°C to 100°C, preferably from 20°C to 50°C.

The reaction time may be varied depending on the nature of the starting material, reagent and reaction temperature, and is usually from 30 minutes to 48 hours, preferably from 1 to 30 hours.

After the completion of the reaction, the target compound of this step is isolated from the reaction mixture according to a conventional method. For example, the target compound is obtained by filtering off the oxidant, removing the solvent by evaporation, or adding water to the residue obtained by removing the solvent by evaporation, adding a hydrophobic solvent (for example, benzene, ether, ethyl acetate, etc.) to the resulting mixture to extract the desired compound, washing the extracted organic layer with water, drying it over anhydrous magnesium sulfate and removing the solvent by evaporation. The target compound thus obtained can be further purified, if necessary, by a conventional method, for example, recrystallization, reprecipitation or chromatography.

Reaction (f):

The reaction in which a carboxyl group is esterified is carried out according to a conventional method in organic chemistry. For example, the reaction is carried out by

(1) reaction with an esterifying agent in an inert solvent,

(2) reaction with an active esterifying agent in an inert solvent to prepare an active ester and then reaction of the active ester with alcohol in an inert solvent, or

(3) reaction with a halogenating agent in an inert solvent to prepare an acid halide and then reaction of the acid halide with alcohol.

The esterifying agent used in reaction (f1) is not particularly limited so long as it is usually used in organic chemistry, and may be, for example, a diazoalkane or trialkylsilyldiazoalkane, preferably a C₁ - C₆ diazoalkane such as diazomethane, diazoethane, diazopropane, diazobutane, diazopentane or diazohexane; or trimethylsilyldiazomethane, more preferably a C₁ - C₄ diazoalkane or trimethylsilyldiazomethane, particularly preferably diazomethane.

The solvent used in the reaction with the diazo C₁ - C₆ alkane is not particularly limited so long as it does not interfere with the reaction and can dissolve a certain amount of the starting material. Examples of suitable solvents may be aliphatic hydrocarbons such as hexane, heptane, ligroin or petroleum ether; aromatic hydrocarbons such as benzene, toluene or xylene; halogenated hydrocarbons such as methylene chloride, chloroform, carbon tetrachloride, dichloroethane, chlorobenzene or dichlorobenzene; ethers such as diethyl ether, diisopropyl ether, tetrahydrofuran, dioxane, dimethoxyethane or diethylene glycol dimethyl ether; esters such as methyl acetate or ethyl acetate; or a mixture of the above-mentioned solvents, preferably the halogenated hydrocarbons, ethers, esters or the mixture of the above-mentioned solvents, more preferably the ethers (particularly diethyl ether), esters (particularly the ethyl acetate) or a mixture of the above-mentioned solvents.

The solvent used in the reaction with the trimethylsilyldiazomethane is not particularly limited so long as it does not interfere with the reaction and can dissolve a certain amount of the starting material. Examples of suitable solvents may be alcohols such as methanol, ethanol, propanol, isopropanol, butanol, isobutanol, t-butanol, pentanol or hexanol; or a mixture of the solvent selected

from the group consisting of aliphatic hydrocarbons such as hexane, heptane, ligroin or petroleum ether; aromatic hydrocarbons such as benzene, toluene or xylene; halogenated hydrocarbons such as methylene chloride, chloroform, carbon tetrachloride, dichloroethane, chlorobenzene or dichlorobenzene; ethers such as diethyl ether, diisopropyl ether, tetrahydrofuran, dioxane, dimethoxyethane or diethylene glycol dimethyl ether; and esters such as methyl acetate or ethyl acetate and the above-mentioned alcohols, preferably the alcohols (particularly methanol), and a mixture of aromatic hydrocarbons (particularly benzene) and the alcohols (particularly methanol).

The reaction temperature may be varied depending on the nature of the starting material and reagent, and is usually from -10°C to 100°C , preferably from 10°C to 50°C .

The reaction time may be varied depending on the nature of the starting material and reagent and reaction temperature, and is usually from 10 minutes to 10 hours, preferably from 15 minutes to 2 hours.

After the completion of the reaction, the target compound of this step is isolated from the reaction mixture according to a conventional method. For example, after the reaction, the target compound is obtained by removing the solvent by evaporation. The target compound thus obtained can be further purified, if necessary, by a conventional method, for example, recrystallization, reprecipitation or chromatography.

The active esterifying agent used in reaction (f2) is not particularly limited so long as it is generally used in the technology of organic chemistry, and may be, for example, an N-hydroxy compound such as ethyl chloroformate, N-hydroxysuccinimide, 1-hydroxybenzotriazole or N-hydroxy-5-norbornene-2,3-dicarboximide; or a disulfide compound such as dipyridyldisulfide.

The esterification reaction is preferably carried out in the presence of a condensing agent such as dicyclohexylcarbodiimide, carbonyldiimidazole or triphenylphosphine.

The solvent used in both reactions is not particularly limited so long as it does not interfere with the reaction and can dissolve a certain amount of the

starting material. Examples of suitable solvents may be halogenated hydrocarbons such as methylene chloride, chloroform, carbon tetrachloride, dichloroethane, chlorobenzene or dichlorobenzene; ethers such as diethyl ether, diisopropyl ether, tetrahydrofuran, dioxane, dimethoxyethane or diethylene glycol dimethyl ether; amides such as formamide, dimethylformamide, dimethylacetamide or hexamethylphosphoric triamide; or nitriles such as acetonitrile, preferably the ethers (particularly tetrahydrofuran) or amides (particularly dimethylformamide).

The reaction temperature may be varied depending on the nature of the starting materials and reagents, and is usually from -70°C to 150°C (preferably from -10°C to 100°C) in the active esterification reaction and from -20°C to 100°C (preferably from 0°C to 50°C) in the reaction of the active ester compound with alcohols.

The reaction time may be varied depending on the nature of the starting materials, reagents and reaction temperatures, and is usually from 30 minutes to 80 hours (preferably from 1 to 48 hours) in both reactions.

After completion of the reaction, the target compound of this reaction is isolated from the reaction mixture according to a conventional method. For example, after the reaction, the target compound is obtained by removing the solvent by evaporation, or adding water to the residue obtained by removing the solvent by evaporation, adding a hydrophobic solvent (for example, benzene, ether, ethyl acetate, etc.) to the resulting mixture to extract the desired compound, washing the extracted organic layer with water, drying it over anhydrous magnesium sulfate and removing the solvent by evaporation. The target compound thus obtained can be further purified, if necessary, by a conventional method, for example, recrystallization, reprecipitation or chromatography.

The halogenating agent used in reaction (f3) is not particularly limited so long as it is generally used in the technology of organic chemistry, and may be, for example, oxalyl chloride, thionyl chloride, phosphorus oxychloride or phosphorus pentachloride, preferably thionyl chloride.

The solvent used in both reactions is not particularly limited so long as it does not interfere with the reaction and can dissolve a certain amount of the starting material. Examples of suitable solvents may be aromatic hydrocarbons such as benzene, toluene or xylene; halogenated hydrocarbons such as methylene chloride, chloroform, carbon tetrachloride, dichloroethane, chlorobenzene or dichlorobenzene; and ethers such as diethyl ether, diisopropyl ether, tetrahydrofuran, dioxane, dimethoxyethane or diethylene glycol dimethyl ether, preferably the ethers (particularly tetrahydrofuran).

The reaction temperature varies depending on the starting materials and reagents, and is usually from -70°C to 150°C (preferably from -10°C to 100°C) in the halogenation reaction and from -20°C to 100°C (preferably from 0°C to 50°C) in the reaction of an acid halide with alcohol.

The time required for both reactions vary depending on the nature of the starting materials and reagents and reaction temperatures, and is usually from 30 minutes to 80 hours (preferably from 1 to 48 hours).

After completion of the reaction, the target compound of this reaction is isolated from the reaction mixture according to a conventional method. For example, after the reaction, the target compound is obtained by removing the solvent by evaporation, or adding water to the residue obtained by removing the solvent by evaporation, adding a hydrophobic solvent (for example, benzene, ether, ethyl acetate, etc.) to the resulting mixture to extract the desired compound, washing the extracted organic layer with water, drying it over anhydrous magnesium sulfate and removing the solvent by evaporation. The target compound thus obtained can be further purified, if necessary, by a conventional method, for example, recrystallization, reprecipitation or chromatography.

Reaction (g):

The reaction in which an alkoxycarbonyl group is converted to a carbamoyl group in reaction (g) is carried out according to a conventional method in organic chemistry. For example, the reaction is carried out by reacting the alkoxycarbonyl group with ammonia gas or conc. ammonia water in

an inert solvent.

The solvent to be employed here is not particularly limited so long as it does not interfere with the reaction and can dissolve a certain amount of the starting material. Examples of suitable solvents may be aliphatic hydrocarbons such as hexane, heptane, ligroin or petroleum ether; aromatic hydrocarbons such as benzene, toluene or xylene; halogenated hydrocarbons such as methylene chloride, chloroform, carbon tetrachloride, dichloroethane, chlorobenzene or dichlorobenzene; ethers such as diethyl ether, diisopropyl ether, tetrahydrofuran, dioxane, dimethoxyethane or diethylene glycol dimethyl ether; alcohols such as methanol, ethanol, propanol, isopropanol, butanol or isobutanol; diamines such as N,N,N',N'-tetramethylethylenediamine; amides such as formamide, dimethylformamide, dimethylacetamide, hexamethylphosphoric triamide or hexamethylphosphorous triamide; or sulfoxides such as dimethyl sulfoxide or sulfolane, preferably the ethers or alcohols, more preferably the ethers (particularly tetrahydrofuran).

The reaction temperature may be varied depending on the nature of the starting material and reagent, and is usually from -10°C to 100°C, preferably from 0°C to 50°C.

The reaction time may be varied depending on the nature of the starting material, reagent and reaction temperature, and is usually from 10 minutes to 10 hours, preferably from 30 minutes to 3 hours.

After completion of the reaction, the target compound of this reaction is isolated from the reaction mixture according to a conventional method. For example, after the reaction, the target compound is obtained by removing the solvent by evaporation, or adding water to the residue obtained by removing the solvent by evaporation, adding a hydrophobic solvent (for example, benzene, ether, ethyl acetate, etc.) to the resulting mixture to extract the desired compound, washing the extracted organic layer with water, drying it over anhydrous magnesium sulfate and removing the solvent by evaporation. The target compound thus obtained can be further purified, if necessary, by a conventional method, for example, recrystallization, reprecipitation or chromatography.

Reaction (h):

The reaction in which a carboxyl group is converted to a carbamoyl group is carried out according to a conventional method in organic chemistry. For example, the reaction is carried out by condensing the carboxyl group with ammonia in an inert solvent according to a conventional method in a peptide synthesis method, for example, an azide method, an active ester method, a mixed acid anhydride method or a condensation method (preferably the mixed acid anhydride method).

In the above-mentioned method, the azide method is carried out by reacting an amino acid hydrazide, which is obtained by reacting with hydrazine in an inert solvent (for example, an amide such as formamide, dimethylformamide, dimethylacetamide or hexamethylphosphoric triamide, preferably dimethylformamide) at from -10°C to 100°C (preferably from 0°C to 50°C), with a nitrite compound to afford an azide compound and then treating it with ammonia.

The nitrite compound to be employed here may be, for example, an alkali metal nitrite such as sodium nitrite or an alkyl nitrite such as isoamyl nitrite.

The reaction is preferably carried out in an inert solvent, and the solvent to be employed here may be, for example, an amide such as formamide, dimethylformamide, dimethylacetamide or hexamethylphosphoric triamide; a sulfoxide such as dimethyl sulfoxide or sulfolane; or a pyrrolidone such as N-methylpyrrolidone, preferably the amides (particularly dimethylformamide).

Further, the two steps of the present reaction (azidation and the reaction with ammonia) are usually carried out in one reaction solution.

The reaction temperature may be varied depending on the nature of the starting materials and reagents, and is usually from -70°C to 50°C (preferably from -50°C to 0°C) in the step of azidation and usually from -70°C to 50°C (preferably from -10°C to 10°C) in the reaction with ammonia.

The time required for the reaction varies depending on the nature of the

starting material, reagent and reaction temperature. It is usually from 5 minutes to 3 hours (preferably from 10 minutes to one hour) in the step of azidation. It is from 5 hours to 7 days (preferably 10 hours to 5 days) in the reaction with ammonia.

After completion of the reaction, the target compound of this reaction is isolated from the reaction mixture according to a conventional method. For example, after the reaction, the target compound is obtained by removing the solvent by evaporation, or adding water to the residue obtained by removing the solvent by evaporation, adding a hydrophobic solvent (for example, benzene, ether, ethyl acetate, etc.) to the resulting mixture to extract the desired compound, washing the extracted organic layer with water, drying it over anhydrous magnesium sulfate and removing the solvent by evaporation. The target compound thus obtained can be further purified, if necessary, by a conventional method, for example, recrystallization, reprecipitation or chromatography.

The active ester method is carried out by reacting with a reagent to prepare an active ester in an inert solvent and then reacting the active ester with ammonia in an inert solvent.

The solvent used in both reactions is not particularly limited so long as it does not interfere with the reaction and can dissolve a certain amount of the starting material, and may be, for example, a halogenated hydrocarbon such as methylene chloride, chloroform, carbon tetrachloride, dichloroethane, chlorobenzene or dichlorobenzene; an ether such as diethyl ether, diisopropyl ether, tetrahydrofuran, dioxane, dimethoxyethane or diethylene glycol dimethyl ether; an amide such as formamide, dimethylformamide, dimethylacetamide or hexamethylphosphoric triamide; or a nitrile such as acetonitrile, preferably the ethers (particularly tetrahydrofuran) or the amides (particularly dimethylformamide).

The reagent to prepare an active ester employed here may be, for example, an N-hydroxy compound such as N-hydroxysuccinimide, 1-hydroxybenzotriazole or N-hydroxy-5-norbornene-2,3-dicarboxyimide; or a

disulfide compound such as dipyridyldisulfide. The active esterification reaction is preferably carried out in the presence of a condensing agent such as dicyclohexylcarbodiimide, carbonyldiimidazole or triphenylphosphine.

The reaction temperatures may be varied depending on the nature of the starting materials and reagent, and are usually from -70°C to 150°C (preferably from -10°C to 100°C) in the active esterification reaction and from -20°C to 100°C (preferably from 0°C to 50°C) in the reaction of the active ester compound with ammonia.

The times required for the both reactions may be varied depending on the nature of the starting materials, reagents and reaction temperatures, and are usually from 30 minutes to 80 hours (preferably from 1 to 48 hours).

After completion of the reaction, the target compound of this reaction is isolated from the reaction mixture according to a conventional method. For example, after the reaction, the target compound is obtained by removing the solvent by evaporation, or adding water to the residue obtained by removing the solvent by evaporation, adding a hydrophobic solvent (for example, benzene, ether, ethyl acetate, etc.) to the resulting mixture to extract the desired compound, washing the extracted organic layer with water, drying it over anhydrous magnesium sulfate and removing the solvent by evaporation. The target compound thus obtained can be further purified, if necessary, by a conventional method, for example, recrystallization, reprecipitation or chromatography.

The mixed acid anhydride method is carried out by reacting with a reagent to prepare a mixed acid anhydride in the presence of a base in an inert solvent and then reacting the mixed acid anhydride with ammonia in an inert solvent.

The solvent used in the reaction for preparing the mixed acid anhydride is not particularly limited so long as it does not interfere with the reaction and can dissolve a certain amount of the starting material, and may be, for example, a halogenated hydrocarbon such as methylene chloride, chloroform, carbon tetrachloride, dichloroethane, chlorobenzene or dichlorobenzene; an ether such

as diethyl ether, diisopropyl ether, tetrahydrofuran, dioxane, dimethoxyethane or diethylene glycol dimethyl ether; or an amide such as formamide, dimethylformamide, dimethylacetamide or hexamethylphosphoric triamide, preferably the ethers (particularly tetrahydrofuran).

Examples of reagents to prepare mixed acid anhydrides may be, for example, C₁ - C₄ alkyl haloformates such as ethyl chloroformate or isobutyl chloroformate; C₁ - C₅ alkanoyl halide such as pivaloyl chloride or C₁ - C₄ alkyl; or di-C₆ - C₁₄ arylcyanophosphoric acid such as diethylcyanophosphoric acid or diphenylcyanophosphoric acid, preferably the C₁ - C₄ alkyl haloformate (particularly isobutyl chloroformate).

The base employed here may be, for example, an alkali metal carbonate such as sodium carbonate, potassium carbonate or lithium carbonate; or an organic amine such as triethylamine, tributylamine, diisopropylethylamine, N-methylmorpholine, pyridine, 4-(N,N-dimethylamino)pyridine, N,N-dimethylaniline, N,N-diethylaniline, 1,5-diazabicyclo[4.3.0]non-5-ene, 1,4-diazabicyclo[2.2.2]octane (DABCO) or 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU), preferably the organic amines (particularly triethylamine).

The reaction temperature in the reaction for preparing the mixed acid anhydride may be varied depending on the nature of the starting material and reagent, and is usually from -50°C to 100°C (preferably from -10°C to 50°C).

The reaction time in the reaction for preparing the mixed acid anhydride may be varied depending on the nature of the starting material, reagent and reaction temperature, and is usually from 5 minutes to 20 hours (preferably from 10 minutes to 10 hours).

The solvent used in the reaction of the mixed acid anhydride with ammonia is not particularly limited so long as it does not interfere with the reaction and can dissolve a certain amount of the starting material, and may be, for example, an ether such as diethyl ether, diisopropyl ether, tetrahydrofuran, dioxane, dimethoxyethane or diethylene glycol dimethyl ether; or an amide such as formamide, dimethylformamide, dimethylacetamide or hexamethylphosphoric triamide, preferably the ethers (particularly

tetrahydrofuran).

The reaction temperature in the reaction of the mixed acid anhydride with ammonia may be varied depending on the nature of the starting material and reagent, and is usually from -30°C to 100°C (preferably from 0°C to 80°C).

The reaction time in the reaction of the mixed acid anhydride with ammonia may be varied depending on the nature of the starting material, reagent and reaction temperature, and is usually from 5 minutes to 24 hours (preferably from 10 minutes to 5 hours).

After completion of the reaction, the target compound of this reaction is isolated from the reaction mixture according to a conventional method. For example, after the reaction, the target compound is obtained by removing the solvent by evaporation, or adding water to the residue obtained by removing the solvent by evaporation, adding a hydrophobic solvent (for example, benzene, ether, ethyl acetate, etc.) to the resulting mixture to extract the desired compound, washing the extracted organic layer with water, drying it over anhydrous magnesium sulfate and removing the solvent by evaporation. The target compound thus obtained can be further purified, if necessary, by a conventional method, for example, recrystallization, reprecipitation or chromatography.

The condensation method is carried out by directly reacting a carboxyl group with ammonia in the presence of a condensing agent in an inert solvent.

The condensing agent employed here may be, for example, dicyclohexylcarbodiimide, carbonyldiimidazole or 1-methyl-2-chloro-pyridiniumiodo-triethylamine, preferably dicyclohexylcarbodiimide.

The present reaction can be carried out under similar conditions to those described in the preparation of active ester.

After completion of the reaction, the target compound of this reaction is isolated from the reaction mixture according to a conventional method. For example, after the reaction, the target compound is obtained by removing the solvent by evaporation, or adding water to the residue obtained by removing the solvent by evaporation, adding a hydrophobic solvent (for example, benzene,

ether, ethyl acetate, etc.) to the resulting mixture to extract the desired compound, washing the extracted organic layer with water, drying it over anhydrous magnesium sulfate and removing the solvent by evaporation. The target compound thus obtained can be further purified, if necessary, by a conventional method, for example, recrystallization, reprecipitation or chromatography.

Reaction (i):

The reaction in which a carbamoyl group is converted to a cyano group is carried out according to a conventional method in organic chemistry. For example, the reaction is carried out by reacting a carbamoyl group with a dehydrating agent in an inert solvent.

The solvent to be employed here is not particularly limited so long as it does not interfere with the reaction and can dissolve a certain amount of the starting material, and may be, for example, an aliphatic hydrocarbon such as hexane, heptane, ligroin or petroleum ether; an aromatic hydrocarbon such as benzene, toluene or xylene; a halogenated hydrocarbon such as methylene chloride, chloroform, carbon tetrachloride, dichloroethane, chlorobenzene or dichlorobenzene; an ether such as diethyl ether, diisopropyl ether, tetrahydrofuran, dioxane, dimethoxyethane or diethylene glycol dimethyl ether; an ester such as methyl acetate or ethyl acetate; a ketone such as acetone; an amide such as formamide, dimethylformamide, dimethylacetamide, hexamethylphosphoric triamide or hexamethylphosphorous triamide; or a sulfoxide such as dimethyl sulfoxide or sulfolane, preferably the ethers, amides, or sulfoxides, more preferably the amides (particularly dimethylformamide).

The dehydrating agent employed here may be, for example, phosphorus oxychloride, trifluoroacetic anhydride, methanesulfonyl chloride, para-toluenesulfonyl chloride or phosphorus pentoxide, preferably phosphorus oxychloride.

The reaction temperature may be varied depending on the nature of the starting material and reagent, and is usually from -30°C to 100°C, preferably from 0°C to 50°C.

The reaction time may be varied depending on the nature of the starting material, reagent and the reaction temperature, and is usually from 5 minutes to 10 hours, preferably from 10 minutes to 3 hours.

After completion of the reaction, the target compound of this reaction is isolated from the reaction mixture according to a conventional method. For example, the target compound is obtained by removing the solvent by evaporation, or adding water to the reaction mixture, adding a hydrophobic solvent (for example, benzene, ether, ethyl acetate, etc.) to the resulting mixture to extract the desired compound, washing the organic layer with water, drying it over anhydrous magnesium sulfate and removing the solvent by evaporation. The target compound thus obtained can be further purified, if necessary, by a conventional method, for example, recrystallization, reprecipitation or chromatography.

Reaction (j):

The reaction in which the alkoxy group on an aromatic ring is converted to a hydroxyl group is carried out according to a conventional method in organic chemistry. For example, the reaction is carried out by reacting the alkoxy group with aluminum chloride in an inert solvent.

The solvent employed here is not particularly limited so long as it does not interfere with the reaction and can dissolve a certain amount of the starting material, and may be, for example, an aliphatic hydrocarbon such as hexane, heptane, ligroin or petroleum ether; an aromatic hydrocarbon such as benzene, toluene or xylene; a halogenated hydrocarbon such as methylene chloride, chloroform, carbon tetrachloride, dichloroethane, chlorobenzene or dichlorobenzene; an ether such as diethyl ether, diisopropyl ether, tetrahydrofuran, dioxane, dimethoxyethane or diethylene glycol dimethyl ether; or an ester such as methyl acetate or ethyl acetate, preferably the halogenated hydrocarbons (particularly methylene chloride).

The reaction temperature may be varied depending on the nature of the starting material and reagent, and is usually from -10°C to 100°C, preferably from 10°C to 50°C.

The reaction time may be varied depending on the nature of the starting material, reagent and reaction temperature, and is usually from 1 to 72 hours, preferably from 2 to 30 hours.

After completion of the reaction, the target compound of this reaction is isolated from the reaction mixture according to a conventional method. For example, the target compound is obtained by removing the solvent by evaporation, adding water to the reaction mixture, neutralizing it, if desired, adding a hydrophobic solvent (for example, benzene, ether, ethyl acetate, etc.) to the resulting mixture to extract the desired compound, washing the organic layer with water, drying it over anhydrous magnesium sulfate and removing the solvent by evaporation. The target compound thus obtained can be further purified, if necessary, by a conventional method, for example, recrystallization, reprecipitation or chromatography.

Reaction (k):

The reaction in which a hydroxyl group or an amino group is acylated is carried out according to a conventional method in organic chemistry. For example, the reaction is carried out by reacting a hydroxyl group or amino group with an acylating agent (preferably alkanoyl halide, a mixed acid anhydride of formic acid and acetic acid, alkanecarboxylic acid anhydride, arylcarbonyl halide or arylcarboxylic acid anhydride) in the presence or absence of a base (preferably in the presence of a base), in an inert solvent.

The alkanoyl halide employed here may be, for example, a straight or branched alkanoyl halide having from 2 to 6 carbon atoms such as acetyl chloride, propionyl chloride, butyryl chloride, butyryl bromide, isobutyryl chloride, valeryl chloride, pivaloyl chloride or hexanoyl chloride, preferably a C₂ - C₄ alkanoyl chloride, more preferably acetyl chloride.

The alkanecarboxylic acid anhydride employed here may be, for example, a straight or branched alkanecarboxylic acid anhydride having from 4 to 12 carbon atoms such as acetic anhydride, propionic anhydride, butanoic anhydride, valeric anhydride, pivalic anhydride, pentanoic anhydride or hexanoic anhydride, preferably a C₄-C₈ alkanecarboxylic acid anhydride, more

preferably acetic anhydride.

The arylcarbonyl halide employed here may be, for example, a C₆ - C₁₀ arylcarboxylic acid halide such as benzoyl chloride, benzoyl bromide, fluorobenzoyl chloride, chlorobenzoyl chloride, dichlorobenzoyl chloride, toluoyl chloride, anisoyl chloride, indenoyl chloride, indenoyl bromide, naphthoyl chloride, naphthoyl bromide, phenanthrenoyl chloride or anthracenoyl chloride, preferably benzoyl chloride.

The arylcarboxylic acid anhydride employed here may be, for example, a C₆ - C₁₀ arylcarboxylic acid anhydride such as benzoic anhydride, fluorobenzoic anhydride, chlorobenzoic anhydride, methylbenzoic anhydride, methoxybenzoic anhydride, indenylcarboxylic anhydride or naphthylcarboxylic anhydride, preferably benzoic anhydride.

The base employed here may be, for example, an alkali metal carbonate such as sodium carbonate, potassium carbonate or lithium carbonate; or an organic amine such as triethylamine, tributylamine, diisopropylethylamine, N-methylmorpholine, pyridine, 4-(N,N-dimethylamino)pyridine, N,N-dimethylaniline, N,N-diethylaniline, 1,5-diazabicyclo[4.3.0]non-5-ene, 1,4-diazabicyclo[2.2.2]octane (DABCO) or 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU), preferably the organic amines (particularly triethylamine, diisopropylethylamine, pyridine or 4-(N,N-dimethylamino)pyridine).

The solvent employed here is not particularly limited so long as it does not interfere with the reaction and can dissolve a certain amount of the starting material, and may be, for example, an aliphatic hydrocarbon such as hexane, heptane, ligroin or petroleum ether; an aromatic hydrocarbon such as benzene, toluene or xylene; a halogenated hydrocarbon such as methylene chloride, chloroform, carbon tetrachloride, dichloroethane, chlorobenzene or dichlorobenzene; an ether such as diethyl ether, diisopropyl ether, tetrahydrofuran, dioxane, dimethoxyethane or diethylene glycol dimethyl ether; a ketone such as acetone and methyl ethyl ketone; or an amide such as formamide, dimethylformamide, dimethylacetamide or hexamethylphosphoric triamide, preferably the ethers (particularly tetrahydrofuran).

The reaction temperature may be varied depending on the nature of the

starting material and the reagent, and is usually from -50°C to 100°C (preferably from 0°C to 50°C).

The reaction time may be varied depending on the nature of the starting material, reagent and reaction temperature, and is usually from 5 minutes to 20 hours (preferably from 10 minutes to 10 hours).

After completion of the reaction, the target compound of this reaction is isolated from the reaction mixture according to a conventional method. For example, after the reaction, the target compound is obtained by removing the solvent by evaporation, or adding water to the residue obtained by removing the solvent by evaporation, adding a hydrophobic solvent (for example, benzene, ether, ethyl acetate, etc.) to the resulting mixture to extract the desired compound, washing the extracted organic layer with water, drying it over anhydrous magnesium sulfate and removing the solvent by evaporation. The target compound thus obtained can be further purified, if necessary, by a conventional method, for example, recrystallization, reprecipitation or chromatography.

Reaction (1):

The reaction in which a hydroxyl group or amino group is aralkylated is carried out according to a conventional method in organic chemistry. For example, the reaction is carried out by reacting a hydroxyl group or amino group with an aralkyl halide in the presence or absence of a base (preferably in the presence of a base) in an inert solvent.

The $\text{C}_6 - \text{C}_{48}$ aralkyl halide to be employed here may be, for example, a $\text{C}_6 - \text{C}_{48}$ aralkyl halide which has from 1 to 3 substituents which may be the same as or different from each other and selected from the group consisting of halogen, $\text{C}_1 - \text{C}_6$ alkyl and $\text{C}_1 - \text{C}_6$ alkoxy such as benzyl chloride, benzyl bromide, 4-chlorobenzyl chloride, 4-chlorobenzyl bromide, 4-bromobenzyl chloride, 4-bromobenzyl bromide, 2,4-difluorobenzyl chloride, 2,4-dichlorobenzyl chloride, 2,4-dichlorobenzyl bromide, 4-methoxybenzyl chloride, 4-methoxybenzyl bromide, trityl chloride, trityl bromide, dimethoxytrityl chloride

or α -naphthyldiphenylmethyl chloride, preferably benzyl chloride or benzyl bromide which may have from 1 to 3 substituents which may be the same as or different from each other and selected from the group consisting of halogen, C₁ - C₄ alkyl and C₁ - C₄ alkoxy, more preferably benzyl chloride or benzyl bromide.

The base employed here may be, for example, an alkali metal carbonate such as sodium carbonate, potassium carbonate or lithium carbonate; an alkali metal hydrogencarbonate such as sodium hydrogencarbonate, potassium hydrogencarbonate or lithium hydrogencarbonate; an alkali metal hydride such as lithium hydride, sodium hydride or potassium hydride; an alkali metal hydroxide such as sodium hydroxide, potassium hydroxide or lithium hydroxide; an organic amine such as triethylamine, tributylamine, diisopropylethylamine, N-methylmorpholine, pyridine, 4-(N,N-dimethylamino)pyridine, N,N-dimethylaniline, N,N-diethylaniline, 1,5-diazabicyclo[4.3.0]non-5-ene (DBN), 1,4-diazabicyclo[2.2.2]octane (DABCO) or 1,8-diazabicyclo[5.4.0]-undec-7-ene (DBU); an alkyllithium such as methyllithium, ethyllithium or butyllithium; or a lithium alkylamide such as lithium diisopropylamide or lithium dicyclohexylamide, preferably the alkali metal carbonates, alkali metal hydrides or organic amines, more preferably the alkali metal carbonates (particularly sodium carbonate or potassium carbonate) or alkali metal hydrides (particularly sodium hydride).

The solvent employed here is not particularly limited so long as it does not interfere with the reaction and can dissolve a certain amount of the starting material, and may be, for example, an aliphatic hydrocarbon such as hexane, heptane, ligroin or petroleum ether; an aromatic hydrocarbon such as benzene, toluene or xylene; a halogenated hydrocarbon such as methylene chloride, chloroform, carbon tetrachloride, dichloroethane, chlorobenzene or dichlorobenzene; an ether such as diethyl ether, diisopropyl ether, tetrahydrofuran, dioxane, dimethoxyethane or diethylene glycol dimethyl ether; a ketone such as acetone or methyl ethyl ketone; or an amide such as formamide, dimethylformamide, dimethylacetamide or hexamethylphosphoric triamide, preferably the amides (particularly dimethylformamide).

The reaction temperature may be varied depending on the nature of the starting material and reagent, and is usually from -50°C to 100°C (preferably from 0°C to 50°C).

The reaction time may be varied depending on the nature of the starting material, reagent and reaction temperature, and is usually from 5 minutes to 24 hours (preferably from 10 minutes to 5 hours).

After completion of the reaction, the target compound of this reaction is isolated from the reaction mixture according to a conventional method. For example, after the reaction, the target compound is obtained by removing the solvent by evaporation, or adding water to the residue obtained by removing the solvent by evaporation, neutralizing it, if desired, adding a hydrophobic solvent (for example, benzene, ether, ethyl acetate, etc.) to the resulting mixture to extract the desired compound, washing the extracted organic layer with water, drying it over anhydrous magnesium sulfate and removing the solvent by evaporation. The target compound thus obtained can be further purified, if necessary, by a conventional method, for example, recrystallization, reprecipitation or chromatography.

Reaction (m):

The reaction in which a nitro group is converted to an amino group is carried out according to a conventional method in organic chemistry. For example, the reaction is carried out by reacting a nitro group with zinc in the presence of acetic acid in an inert solvent.

The solvent employed here is not particularly limited so long as it does not interfere with the reaction and can dissolve a certain amount of the starting material, and may be, for example, an aliphatic hydrocarbon such as hexane, heptane, ligroin or petroleum ether; an aromatic hydrocarbon such as benzene, toluene or xylene; a halogenated hydrocarbon such as methylene chloride, chloroform, carbon tetrachloride, dichloroethane, chlorobenzene or dichlorobenzene; an ether such as diethyl ether, diisopropyl ether, tetrahydrofuran, dioxane, dimethoxyethane or diethylene glycol dimethyl ether; an alcohol such as methanol, ethanol, propanol, isopropanol, butanol or

isobutanol; a diamine such as N,N,N',N'-tetramethylethylenediamine; an amide such as formamide, dimethylformamide, dimethylacetamide, hexamethylphosphoric triamide or hexamethylphosphorous triamide; a sulfoxide such as dimethyl sulfoxide or sulfolane; water or an aqueous mixture of the above-mentioned solvent, preferably water.

The reaction temperature may be varied depending on the nature of the starting material and reagent, and is usually from -10°C to 100°C, preferably from 0°C to 50°C.

The reaction time may be varied depending on the nature of the starting material, reagent and reaction temperature, and is usually from 10 minutes to 10 hours, preferably from 30 minutes to 3 hours.

After completion of the reaction, the target compound of this reaction is isolated from the reaction mixture according to a conventional method. For example, after the reaction, the target compound is obtained by removing the solvent by evaporation, or adding water to the residue obtained by removing the solvent by evaporation, neutralizing it, adding a hydrophobic solvent (for example, benzene, ether, ethyl acetate, etc.) to the resulting mixture to extract the desired compound, washing the extracted organic layer with water, drying it over anhydrous magnesium sulfate and removing the solvent by evaporation. The target compound thus obtained can be further purified, if necessary, by a conventional method, for example, recrystallization, reprecipitation or chromatography.

Reaction (n):

The reaction for removal of the protecting group of the amino group included in R³_a, etc. is carried out under similar conditions to those described in Step A3.


The compound of formula (II) which is a starting material of the present invention is a known compound or can be prepared according to a known method [for example, Chem. Abstr., 74, 125521 (1970), Annual report of Sankyo Research Laboratories, 22, 215 (1970), Agric. Biol. Chem., EN, 50,

1831 (1986), Can. J. Chem., 48, 1371 (1970), Japanese Patent Application (Kokai) No. Sho 59-216881, Japanese Patent Publication (Kokoku) No. Sho 43-14704, etc.]


Further, the compound of formula (IV) or (IVa) is a known compound or can be prepared according to a known method [for example, Synthesis, 366 (1990), J. Med. Chem., 34, 1258 (1991), etc.]

Furthermore, the compound (II) which is a starting material of the present invention is a known compound or can be prepared according to a known method. The compound of the general formula (VI), (X) or (XIII) can be prepared by reacting according to the methods described below.


Method D



Method E



Method F



Wherein R^1 , R^2 and Z have the same meanings as defined above, R^{2a} represents a specific substituent group in R^2 (the substituent group consists of a halogen atom, $C_1 - C_6$ alkyl, $C_1 - C_6$ alkyl substituted with halogen or $C_1 - C_6$ alkoxy, $C_2 - C_6$ alkenyl, $C_2 - C_6$ alkynyl, $C_3 - C_{10}$ cycloalkyl, $C_3 - C_{10}$ cycloalkenyl, $C_2 - C_6$ alkanoyl and $C_1 - C_6$ alkoxy carbonyl groups), and R^4 represents a $C_1 - C_6$ alkyl group.

The alkyl group in R⁴ may be, for example, a straight or branched alkyl group having from 1 to 6 carbon atoms such as the methyl, ethyl, propyl, isopropyl, butyl, isobutyl, s-butyl, t-butyl, pentyl, isopentyl, 2-methylbutyl, neopentyl, 1-ethylpropyl, hexyl, 4-methylpentyl, 3-methylpentyl, 2-methylpentyl, 1-methylpentyl, 3,3-dimethylbutyl, 2,2-dimethylbutyl, 1,1-dimethylbutyl, 1,2-dimethylbutyl, 1,3-dimethylbutyl, 2,3-dimethylbutyl or 2-ethylbutyl groups, preferably the C₁ - C₄ alkyl group, more preferably the methyl or ethyl groups.

Method D is a process for preparing the compound (II) which is a starting material in Process A or Process B.

In step D1 the compound of formula (VIII) is prepared

- (a) by reacting the compound of general formula (VI) in an inert solvent with a reagent to prepare an active ester, followed by reacting the active ester with the compound of formula (VII) in an inert solvent,
- (b) by reacting the compound of general formula (VI) with a halogenating agent in an inert solvent, followed by reacting the halogenated compound with the compound (VII), or
- (c) by reacting the compound of formula (VI) with the compound of formula (VII) in the presence of an acid in an inert solvent.

Step D1(a) and Step D1(b) are carried out under similar conditions to those described in reaction (f2) and reaction (f3) in Step C1, respectively.

The acid used in Step D1(c) is not particularly limited so long as it is conventionally used in the technology of organic chemistry, and may be a mineral acid such as hydrofluoric acid, hydrochloric acid, hydrobromic acid, hydroiodic acid, nitric acid, perchloric acid, sulfuric acid or phosphoric acid; a sulfonic acid such as methanesulfonic acid, trifluoromethanesulfonic acid, ethanesulfonic acid, benzenesulfonic acid or p-toluenesulfonic acid; or a carboxylic acid such as fumaric acid, succinic acid, citric acid, tartaric acid, oxalic acid or maleic acid; preferably the mineral acids (particularly sulfuric

acid).

The solvent employed here is not particularly limited so long as it does not interfere with the reaction and can dissolve a certain amount of the starting material, and may be, for example, an aliphatic hydrocarbon such as hexane, heptane, ligroin or petroleum ether; an aromatic hydrocarbon such as benzene, toluene or xylene; a halogenated hydrocarbon such as methylene chloride, chloroform, carbon tetrachloride, dichloroethane, chlorobenzene or dichlorobenzene; an ether such as diethyl ether, diisopropyl ether, tetrahydrofuran, dioxane, dimethoxyethane or diethylene glycol dimethyl ether; a ketone such as acetone or methyl ethyl ketone; an alcohol such as methanol, ethanol, propanol, isopropanol, butanol or isobutanol; or an amide such as formamide, dimethylformamide, dimethylacetamide or hexamethylphosphoric triamide, preferably the alcohols (particularly methanol or ethanol).

The reaction temperature may be varied depending on the nature of the starting material and reagent, and is usually from -50°C to 150°C (preferably from 20°C to 100°C).

The reaction time may be varied depending on the nature of the starting material, reagent and reaction temperature, and is usually from 5 minutes to 24 hours (preferably from 10 minutes to 5 hours).

After completion of the reaction, the target compound of this reaction is isolated from the reaction mixture according to a conventional method. For example, after the reaction, the target compound is obtained by removing the solvent by evaporation, or adding water to the residue obtained by removing the solvent by evaporation, adding a hydrophobic solvent (for example, benzene, ether, ethyl acetate, etc.) to the resulting mixture to extract the desired compound, washing the extracted organic layer with water, drying it over anhydrous magnesium sulfate and removing the solvent by evaporation. The target compound thus obtained can be further purified, if necessary, by a conventional method, for example, recrystallization, reprecipitation or chromatography.

In step D2 the compound of formula (IX) is prepared by reacting the

compound (VIII) with a halogen molecule in an inert solvent.

The halogen molecule employed here may be, for example, a chlorine molecule or a bromine molecule, preferably the bromine molecule.

The solvent employed here is not particularly limited so long as it does not interfere with the reaction and can dissolve a certain amount of the starting material, and may be, for example, an aliphatic hydrocarbon such as hexane, heptane, ligroin or petroleum ether; an aromatic hydrocarbon such as benzene, toluene or xylene; a halogenated hydrocarbon such as methylene chloride, chloroform, carbon tetrachloride, dichloroethane, chlorobenzene or dichlorobenzene; an ether such as diethyl ether, diisopropyl ether, tetrahydrofuran, dioxane, dimethoxyethane or diethylene glycol dimethyl ether; a ketone such as acetone and methyl ethyl ketone; an alcohol such as methanol, ethanol, propanol, isopropanol, butanol or isobutanol; or an amide such as formamide, dimethylformamide, dimethylacetamide or hexamethylphosphoric triamide, preferably the halogenated hydrocarbons (particularly chloroform or carbon tetrachloride).

The reaction temperature may be varied depending on the nature of the starting material and reagent, and is usually from -50°C to 100°C (preferably from 0°C to 50°C).

The reaction time may be varied depending on the nature of the starting material, reagent and reaction temperature, and is usually from 5 minutes to 12 hours (preferably from 10 minutes to 5 hours).

After completion of the reaction, the target compound of this reaction is isolated from the reaction mixture according to a conventional method. For example, after the reaction, the target compound is obtained by removing the solvent by evaporation, or adding water to the residue obtained by removing the solvent by evaporation, adding a hydrophobic solvent (for example, benzene, ether, ethyl acetate, etc.) to the resulting mixture to extract the desired compound, washing the extracted organic layer with water, drying it over anhydrous magnesium sulfate and removing the solvent by evaporation. The target compound thus obtained can be further purified, if necessary, by a conventional method, for example, recrystallization, reprecipitation or

chromatography.

In step D3 the compound of formula (II) is prepared by reacting the compound of formula (IX) with hydroxylamine or a mineral acid salt of hydroxylamine (preferably the mineral acid salt of hydroxylamine) in the presence or absence of a base (preferably in the presence of a base) in an inert solvent.

The mineral acid salt of hydroxylamine employed here may be, for example, hydroxylamine hydrofluoride, hydroxylamine hydrochloride, hydroxylamine hydrobromide, hydroxylamine hydroiodide, hydroxylamine nitric acid salt, hydroxylamine perchloric acid salt, hydroxylamine sulfuric acid salt or hydroxylamine phosphoric acid salt, preferably hydroxylamine hydrochloride.

Method E is an alternative method for preparing the compound (II) which is a starting material in Process A or Process B.

In step E1 the compound of formula (XII) is prepared by reacting the compound of formula (X) with the compound of formula (XI) in the presence of a base in an inert solvent.

The solvent employed here is not particularly limited so long as it does not interfere with the reaction and can dissolve a certain amount of the starting material, and may be, for example, an aliphatic hydrocarbon such as hexane, heptane, ligroin or petroleum ether; an aromatic hydrocarbon such as benzene, toluene or xylene; a halogenated hydrocarbon such as methylene chloride, chloroform, carbon tetrachloride, dichloroethane, chlorobenzene or dichlorobenzene; an ether such as diethyl ether, diisopropyl ether, tetrahydrofuran, dioxane, dimethoxyethane or diethylene glycol dimethyl ether; or a sulfoxide such as dimethyl sulfoxide or sulfolane, preferably the aromatic hydrocarbons, halogenated hydrocarbons or ethers, more preferably the aromatic hydrocarbons (particularly benzene) or ethers (particularly tetrahydrofuran and dioxane).

The base employed here may be, for example, an alkali metal carbonate such as sodium carbonate, potassium carbonate or lithium carbonate; an alkali metal hydrogencarbonate such as sodium hydrogencarbonate, potassium hydrogencarbonate or lithium hydrogencarbonate; an alkali metal hydride such as lithium hydride, sodium hydride or potassium hydride; an alkali metal hydroxide such as sodium hydroxide, potassium hydroxide or lithium hydroxide; an alkali metal alkoxide such as sodium methoxide, sodium ethoxide, potassium t-butoxide or lithium methoxide; a mercaptan alkali metal such as methylmercaptan sodium or ethylmercaptan sodium; an organic amine such as triethylamine, tributylamine, diisopropylethylamine, N-methylmorpholine, pyridine, 4-(N,N-dimethylamino)pyridine, N,N-dimethylaniline, N,N-diethylaniline, 1,5-diazabicyclo[4.3.0]non-5-ene, 1,4-diazabicyclo[2.2.2]octane (DABCO) or 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU); an alkyl lithium such as methyllithium, ethyllithium or butyllithium; or a lithium alkylamide such as lithium diisopropylamide or lithium dicyclohexylamide, preferably the lithium alkylamides (particularly lithium diisopropylamide) or organic amines (particularly 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU)).

The reaction temperature may be varied depending on the nature of the starting material and reagent, and is usually from -100°C to 100°C, preferably from -70°C to 50°C.

The reaction time may be varied depending on the nature of the starting material, reagent and reaction temperature, and is usually from 5 minutes to 48 hours, preferably from 10 minutes to 24 hours.

After the completion of the reaction, the target compound of this step is isolated from the reaction mixture according to a conventional method. For example, the target compound is obtained by removing the solvent by evaporation, adding water to the reaction mixture, making the reaction mixture weakly acidic, adding a hydrophobic solvent (for example, benzene, ether, ethyl acetate, etc.) to the resulting mixture to extract the desired compound, washing the organic layer with water, drying it over anhydrous magnesium sulfate and removing the solvent by evaporation. The target compound thus obtained can

be further purified, if necessary, by a conventional method, for example, recrystallization, reprecipitation or chromatography.

In step E2 the compound of formula (II) is prepared by reacting the compound of formula (XII) with hydroxylamine or a mineral acid salt of hydroxylamine (preferably the mineral acid salt of hydroxylamine) in the presence or absence of a base (preferably in the presence of a base) in an inert solvent, and is carried out under similar conditions to those described in Step D3.

Method F is a process for preparing the compound (XIIa) in which R² is a halogen atom, C₁ - C₆ alkyl, C₁ - C₆ alkyl substituted with halogen or C₁ - C₆ alkoxy, C₂ - C₆ alkenyl, C₂ - C₆ alkynyl, C₃ - C₁₀ cycloalkyl, C₃ - C₁₀ cycloalkenyl, C₂ - C₆ alkanoyl or C₁ - C₆ alkoxy carbonyl in the intermediate (XII) of Method E.

In step F1 the compound of formula (XIIa) is prepared by reacting the compound of formula (XIII) with the compound of formula (XIV) in the presence of a base in an inert solvent, and is carried out under similar conditions to those described in Step E1.

The isoxazole derivative (I) of the present invention has excellent type A-monoamine oxidase inhibiting activities and is weakly toxic. Therefore, it is useful as an agent for treating or preventing (particularly an agent for treating) nervous diseases including depression, Parkinson's disease, Alzheimer's dementia (cognitive disorder owing to Alzheimer's disease) or cerebrovascular dementia (cognitive disorder owing to cerebrovascular dementia), (particularly for depression).

[Best mode for carrying out the invention]

In the following, the present invention will be described in more detail by way of Examples, Reference examples, Test example and Preparation examples, but the present invention is not limited thereto.

Example 1

3-(2-Aminoethoxy)-5-phenylisoxazole hydrochloride (Compound list No. 1)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-phenylisoxazole

Triphenylphosphine (0.87 g) was dissolved in tetrahydrofuran (10 ml), and diethyl azodicarboxylate (0.57 g) was added dropwise to the solution under ice-cooling with stirring, and the mixture was stirred at the same temperature for 10 minutes. 3-Hydroxy-5-phenylisoxazole (0.48 g) and 2-(N-tert-butoxycarbonylamino)ethanol (0.48 g) were added to the reaction mixture, and the resulting mixture was stirred under ice-cooling for 10 minutes and at room temperature for 24 hours. The solvent was evaporated under reduced pressure and the residue was purified by silica gel column chromatography (eluent: cyclohexane/ethyl acetate = 4/1) and was crystallized from isopropyl ether to obtain the title compound (0.63 g, 69%) as colorless crystals.

m.p.: 125-126°C;

IR spectrum (KBr) ν_{\max} cm⁻¹: 3322, 1721, 1710, 1619;

NMR spectrum (CDCl₃) δ ppm: 1.46(9H, s), 3.56(2H, q, J=5.1Hz), 4.35(2H, t, J=5.1Hz), 4.94(1H, brs), 6.14(1H, s), 7.43-7.51(3H, m), 7.71-7.74(2H, m).

(b) 3-(2-Aminoethoxy)-5-phenylisoxazole hydrochloride

A solution of 4N hydrochloric acid/1,4-dioxane (4.0 ml) was added to 3-(2-(N-tert-butoxycarbonylamino)ethoxy)-5-phenylisoxazole (0.50 g), and the mixture was stirred at room temperature for 15 minutes. The precipitate was separated from the mixture by filtration and washed with ethyl acetate to obtain the title compound (0.39 g, 99%) as colorless crystals.

m.p.: 215-218°C (decomposed);

- 147 -

IR spectrum (KBr) ν_{\max} cm^{-1} : 3132, 2963, 2810, 2756, 1620, 1597, 1579;

NMR spectrum (DMSO- d_6) δ ppm: 3.26(2H, t, $J=5.1\text{Hz}$), 4.45(2H, t, $J=5.1\text{Hz}$), 6.85(1H, s), 7.51-7.57(3H, m), 7.84-7.87(2H, m), 8.25(3H, brs).

Example 2

3-(2-Aminoethoxy)-4-chloro-5-phenylisoxazole hydrochloride (Compound list No.: 5)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-chloro-5-phenylisoxazole

4-Chloro-3-hydroxy-5-phenylisoxazole (0.58 g) and 2-(N-tert-butoxycarbonylamino)ethanol (0.48 g) were subjected to reaction and post-treatment in a similar manner to that described in Example 1(a) to obtain the title compound (0.73 g, 72%) as colorless crystals.

m.p.: 115-116°C;

IR spectrum (KBr) ν_{\max} cm^{-1} : 3346, 1720, 1709, 1616;

NMR spectrum (CDCl_3) δ ppm: 1.46(9H, s), 3.61(2H, q, $J=5.1\text{Hz}$), 4.42(2H, t, $J=5.1\text{Hz}$), 4.97(1H, brs), 7.46-7.53(3H, m), 7.94-8.00(2H, m).

(b) 3-(2-Aminoethoxy)-4-chloro-5-phenylisoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-chloro-5-phenylisoxazole (0.54 g) was subjected to reaction and post-treatment in a similar manner to that described in Example 1(b) to obtain the title compound (0.41 g, 93%) as colorless crystals.

m.p.: 204-207°C (decomposed);

IR spectrum (KBr) ν_{\max} cm^{-1} : 2971, 2905, 2848, 2775, 1606, 1575, 1534;

NMR spectrum (DMSO- d_6) δ ppm: 3.31(2H, t, $J=5.1\text{Hz}$), 4.56(2H, t, $J=5.1\text{Hz}$), 7.59-7.67(3H, m), 7.92-7.97(2H, m), 8.27(3H, brs).

Example 3

3-(2-Aminoethoxy)-5-(4-chlorophenyl)isoxazole hydrochloride (Compound list No.: 143)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(4-chlorophenyl)isoxazole

5-(4-Chlorophenyl)-3-hydroxyaminoisoxazole (0.58 g) and 2-(N-tert-butoxycarbonylamino)ethanol (0.48 g) were subjected to reaction and post-treatment in a similar manner to that described in Example 1(a) to obtain the title compound (0.69 g, 68%) as colorless crystals.

m.p.: 128-129°C;

IR spectrum (KBr) ν_{\max} cm^{-1} : 3378, 1683, 1622;

NMR spectrum (CDCl_3) δ ppm: 1.46(9H, s), 3.56(2H, q, J=5.1Hz), 4.35(2H, t, J=5.1Hz), 4.93(1H, brs), 6.14(1H, s), 7.43(2H, d, J=8.7Hz), 7.66(2H, d, J=8.7Hz).

(b) 3-(2-Aminoethoxy)-5-(4-chlorophenyl)isoxazole hydrochloride

3-(2-(N-tert-butoxycarbonylamino)ethoxy)-5-(4-chlorophenyl)isoxazole (0.54 g) was subjected to reaction and post-treatment in a similar manner to that described in Example 1(b) to obtain the title compound (0.43 g, 98%) as colorless crystals.

m.p.: 218-223°C (decomposed);

IR spectrum (KBr) ν_{\max} cm^{-1} : 3135, 2998, 2809, 1618, 1603, 1594, 1575, 1567;

NMR spectrum (DMSO-d_6) δ ppm: 3.26(2H, t, J=5.1Hz), 4.45(2H, t, J=5.1Hz), 6.91(1H, s), 7.63(2H, d, J=8.7Hz), 7.89(2H, d, J=8.7Hz), 8.25(3, brs).

Example 4

3-(2-Aminoethoxy)-4-isopropyl-5-phenylisoxazole hydrochloride (Compound list No.: 9)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-isopropyl-5-phenylisoxazole

3-Hydroxy-4-isopropyl-5-phenylisoxazole (0.50 g) and 2-(N-tert-butoxycarbonylamino)ethanol (0.40 g) were subjected to reaction and post-treatment in a similar manner to that described in Example 1(a) to obtain the

title compound (0.60 g, 69%) as colorless crystals.

m.p.: 98-99°C;

IR spectrum (KBr) ν_{\max} cm^{-1} : 3386, 1686, 1642;

NMR spectrum (CDCl_3) δ ppm: 1.29(6H, d, $J=6.8$), 1.46(9H, s), 3.06(1H, q, $J=6.8$), 3.60(2H, q, $J=5.1\text{Hz}$), 4.38(2H, t, $J=5.1\text{Hz}$), 4.85(1H, brs), 7.42-7.50(3H, m), 7.55-7.60(2H, m).

(b) 3-(2-Aminoethoxy)-4-isopropyl-5-phenylisoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-isopropyl-5-phenylisoxazole (0.50 g) was subjected to reaction and post-treatment in a similar manner to that described in Example 1(b) to obtain the title compound (0.39 g, 95%) as colorless crystals.

m.p.: 202-210°C (decomposed);

IR spectrum (KBr) ν_{\max} cm^{-1} : 2975, 2939, 1642, 1599, 1575;

NMR spectrum (DMSO-d_6) δ ppm: 1.26(6H, d, $J=6.8\text{Hz}$), 3.04(1H, q, $J=6.8\text{Hz}$), 3.28(2H, t, $J=5.1\text{Hz}$), 4.46(2H, t, $J=5.1\text{Hz}$), 7.50-7.64(5H, m), 8.26(3H, brs).

Example 5

3-(2-Aminoethoxy)-5-(2-thienyl)isoxazole hydrochloride (Compound list No.: 535)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(2-thienyl)isoxazole

3-Hydroxy-5-(2-thienyl)isoxazole (0.42 g) and 2-(N-tert-butoxycarbonylamino)ethanol (0.40 g) were subjected to reaction and post-treatment in a similar manner to that described in Example 1(a) to obtain the title compound (0.63 g, 82%) as colorless crystals.

m.p.: 129-130°C (decomposed);

IR spectrum (KBr) ν_{\max} cm^{-1} : 3323, 1708, 1694, 1618;

NMR spectrum (CDCl_3) δ ppm: 1.46(9H, s), 3.56(2H, q, $J=5.1\text{Hz}$), 4.34(2H, t, $J=5.1\text{Hz}$), 4.93(1H, brs), 6.05(1H, s), 7.11(1H, dd, $J=5.1\text{Hz}$, $J=3.7\text{Hz}$), 7.43-7.48(2H, m).

(b) 3-(2-Aminoethoxy)-5-(2-thienyl)isoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(2-thienyl)isoxazole (0.05 g) was subjected to reaction and post-treatment in a similar manner to that described in Example 1(b) to be obtain the title compound (0.37g, 95%) as colorless crystals.

m.p.: 278-283°C (decomposed);

IR spectrum (KBr) ν_{\max} cm^{-1} : 3108, 3086, 2993, 2978, 2913, 1613, 1596;

NMR spectrum (DMSO- d_6) δ ppm: 3.25(2H, t, J=5.1Hz), 4.44(2H, t, J=5.1Hz), 6.69(1H, s), 7.25(1H, dd, J=5.5Hz, J=3.7Hz), 7.71(1H, d, J=3.7Hz), 7.84(1H, d, J=5.5Hz), 8.25(3H, brs).

Example 6

3-(2-Aminoethoxy)-4-chloro-5-(2-thienyl)isoxazole hydrochloride (Compound list No.: 539)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(2-thienyl)isoxazole

4-Chloro-3-hydroxy-5-(2-thienyl)isoxazole (0.50 g) and 2-(N-tert-butoxycarbonylamino)ethanol (0.40 g) were subjected to reaction and post-treatment in a similar manner to that described in Example 1(a) to obtain the title compound (0.57 g, 66%) as colorless crystals.

m.p.: 94-95°C;

IR spectrum (KBr) ν_{\max} cm^{-1} : 3342, 1718, 1708, 1622;

NMR spectrum (CDCl₃) δ ppm: 1.46(9H, s), 3.60(2H, q, J=5.1Hz), 4.40(2h, t, J=5.1Hz), 4.96(1H, brs), 7.19(1H, dd, J=5.2Hz, J=3.6Hz), 7.56(1H, d, J=5.2Hz), 7.74(1H, d, 3.6Hz).

(b) 3-(2-Aminoethoxy)-4-chloro-5-(2-thienyl)isoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-chloro-5-(2-thienyl)-isoxazole (0.40 g) was subjected to reaction and post-treatment in a similar manner to that described in Example 1(b) to obtain the title compound (0.31 g, 95%) as colorless crystals.

m.p.: 278-283°C (decomposed);

IR spectrum (KBr) ν_{\max} cm^{-1} : 3109, 2960, 2897, 1626, 1596, 1579;

NMR spectrum (DMSO- d_6) δ ppm: 3.30(2H, t, $J=5.1\text{Hz}$), 4.54(2H, t, $J=5.1\text{Hz}$), 7.34(1H, dd, $J=5.1\text{Hz}$, $J=3.6\text{Hz}$), 7.83(1H, d, $J=3.6\text{Hz}$), 8.01(1H, d, $J=5.1\text{Hz}$), 8.26(3H, brs).

Example 7

3-(2-Aminoethoxy)-5-(3-pyridyl)isoxazole hydrochloride (Compound list No.: 1056)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(3-pyridyl)isoxazole

3-Hydroxy-5-(3-pyridyl)isoxazole (0.41 g) and 2-(N-tert-butoxycarbonylamino)ethanol (0.40 g) were subjected to reaction and post-treatment in a similar manner to that described in Example 1(a) to obtain the title compound (0.50 g, 65%) as colorless crystals.

m.p.: 97-98°C;

IR spectrum (KBr) ν_{\max} cm^{-1} : 3249, 3145, 1712, 1626;

NMR spectrum (CDCl₃) δ ppm: 1.46(9H, s), 3.57(2H, q, $J=5.1\text{Hz}$), 4.37(2H, t, $J=5.1\text{Hz}$), 4.94(1H, brs), 6.25(1H, s), 7.42(1H, dd, $J=8.0\text{Hz}$, $J=4.9\text{Hz}$), 8.04(1H, d, $J=8.0\text{Hz}$), 8.64(1H, d, $J=4.9\text{Hz}$), 8.97(1H, s).

(b) 3-(2-Aminoethoxy)-5-(3-pyridyl)isoxazole dihydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(3-pyridyl)isoxazole (0.48 g) was subjected to reaction and post-treatment in a similar manner to that described in Example 1(b) to obtain the title compound (0.41 g, 92%) as colorless crystals.

m.p.: 222-227°C;

IR spectrum (KBr) ν_{\max} cm^{-1} : 3096, 3068, 3043, 2967, 2886, 2813, 1641, 1597, 1539;

NMR spectrum (DMSO- d_6) δ ppm: 3.26(2H, q, $J=5.1\text{Hz}$), 4.48(2H, t, $J=5.1\text{Hz}$), 7.11(1H, s), 7.77(1H, dd, $J=8.0\text{Hz}$, $J=5.1\text{Hz}$), 8.38(3H, brs), 8.46(1H, d, $J=8.0\text{Hz}$), 8.79(1H, d, $J=5.1\text{Hz}$), 9.21(1H, s).

Example 8

3-(2-Aminoethoxy)-4-chloro-5-(3-pyridyl)isoxazole hydrochloride (Compound list No.: 1061)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(3-pyridyl)isoxazole

4-Chloro-3-hydroxy-5-(3-pyridyl)isoxazole (0.49 g) and 2-(N-tert-butoxycarbonylamino)ethanol (0.40 g) were subjected to reaction and post-treatment in a similar manner to that described in Example 1(a) to obtain the title compound (0.54 g, 63%) as colorless crystals.

m.p. : 76-77°C;

IR spectrum (KBr) ν_{\max} cm^{-1} : 3353, 3248, 1754, 1721, 1709, 1616;

NMR spectrum (CDCl_3) δ ppm: 1.47(9H, s), 3.62(2H, q, $J=5.1\text{Hz}$), 4.43(2H, t, $J=5.1\text{Hz}$), 4.97(1H, brs), 7.45(1H, dd, $J=8.0\text{Hz}$, $J=5.1\text{Hz}$), 8.24(1H, ddd, $J=8.0\text{Hz}$, $J=2.0\text{Hz}$, $J=1.5\text{Hz}$), 8.72(1H, dd, $J=5.1\text{Hz}$, $J=1.5\text{Hz}$), 9.24(1H, d, $J=2.0\text{Hz}$).

(b) 3-(2-Aminoethoxy)-4-chloro-5-(3-pyridyl)isoxazole dihydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-chloro-5-(3-pyridyl)-isoxazole (0.40 g) was subjected to reaction and post-treatment in a similar manner to that described in Example 1(b) to obtain the title compound (0.35 g, 96%) as colorless crystals.

m.p.: 205-210°C (decomposed);

IR spectrum (KBr) ν_{\max} cm^{-1} : 3103, 3053, 2937, 2899, 2875, 2823, 2800, 1634, 1607, 1590, 1541;

NMR spectrum (DMSO-d_6) δ ppm: 3.31(2H, q, $J=5.1\text{Hz}$), 4.59(2H, t, $J=5.1\text{Hz}$), 7.74(1H, dd, $J=8.0\text{Hz}$, $J=5.1\text{Hz}$), 8.40(1H, ddd, $J=8.0\text{Hz}$, $J=2.0\text{Hz}$, $J=1.5\text{Hz}$), 8.40(3H, brs), 8.82(1H, dd, $J=5.1\text{Hz}$, $J=1.5\text{Hz}$), 9.15(1H, d, $J=2.0\text{Hz}$).

Example 9

3-(2-Aminoethoxy)-5-(2-methoxyphenyl)isoxazole hydrochloride (Compound list No.: 357)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(2-methoxyphenyl)isoxazole

Triphenylphosphine (0.45 g) was dissolved in tetrahydrofuran (5 ml), and diethyl azodicarboxylate (0.27 ml) was added dropwise to the solution under ice-cooling with stirring, and the resulting mixture was stirred at the same temperature for 30 minutes. Then, 2-(N-tert-butoxycarbonylamino)ethanol (0.20 g) and 3-hydroxy-5-(2-methoxyphenyl)isoxazole (0.22 g) were added to the reaction mixture, followed by stirring of the resulting mixture under ice-cooling for 10 minutes and at room temperature for 24 hours. The solvent was evaporated under reduced pressure and the residue was purified by silica gel column chromatography (eluent: hexane/ethyl acetate = 4/1) to obtain the title compound (0.33 g, 86%) as colorless crystals.

m.p.: 131-133°C;

IR spectrum (KBr) ν_{\max} cm^{-1} : 3309, 1712, 1620;

NMR spectrum (CDCl_3) δ ppm: 1.46(9H, s), 3.57(2H, q, $J=5.2\text{Hz}$), 3.95(3H, s), 4.36(2H, t, $J=5.2\text{Hz}$), 4.96(1H, brs), 6.43(1H, s), 7.00(1H, d, $J=7.8\text{Hz}$), 7.06(1H, t, $J=7.8\text{Hz}$), 7.41(1H, ddd, $J=7.8\text{Hz}$, $J=7.8\text{Hz}$, $J=1.7\text{Hz}$), 7.91(1H, dd, $J=7.8\text{Hz}$, $J=1.7\text{Hz}$).

(b) 3-(2-Aminoethoxy)-5-(2-methoxyphenyl)isoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(2-methoxyphenyl)isoxazole (0.31 g) was dissolved in a solution of 4N hydrochloric acid/1,4-dioxane (2.3 ml), and the solution was stirred at room temperature for 30 minutes. The solvent was evaporated under reduced pressure and the residue was recrystallized from a mixture of ethanol and isopropanol to obtain the title compound (0.21 g, 84%) as colorless crystals.

m.p.: 160-162°C (decomposed);

IR spectrum (KBr) ν_{\max} cm^{-1} : 3000, 2959, 2837, 1621, 1614;

NMR spectrum (DMSO-d_6) δ ppm: 3.25(2H, t, $J=5.1\text{Hz}$), 3.94(3H, s), 4.45(2H,

t, J=5.1Hz), 6.56(1H, s), 7.11(1H, t, J=7.8Hz), 7.23(1H, d, J=7.8Hz), 7.52(1H, ddd, J=7.8Hz, J=7.8Hz, J=1.7Hz), 7.81(1H, dd, J=7.8Hz, J=1.7Hz), 8.25(3H, brs).

Example 10

3-(2-Aminoethoxy)-5-(3-methoxyphenyl)isoxazole hydrochloride (Compound list No.: 363)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(3-methoxyphenyl)isoxazole

3-Hydroxy-5-(3-methoxyphenyl)isoxazole (0.22 g) and 2-(N-tert-butoxycarbonylamino)ethanol (0.20 g) were subjected to reaction and post-treatment in a similar manner to that described in Example 9(a) to obtain the title compound (0.31 g, 82%) as a colorless powder.

m.p.: 89-90°C;

IR spectrum (KBr) ν_{\max} cm^{-1} : 3312, 1710;

NMR spectrum (CDCl_3) δ ppm: 1.46(9H, s), 3.56(2H, q, J=5.2Hz), 3.86(3H, s), 4.35(2H, t, J=5.2Hz), 4.93(1H, brs), 6.13(1H, s), 6.96-7.00(1H, m), 7.26-7.42(3H, m).

(b) 3-(2-Aminoethoxy)-5-(3-methoxyphenyl)isoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(3-methoxyphenyl)-isoxazole (0.29 g) was allowed to react in a similar manner to that described in Example 9(b) and the reaction product was recrystallized from a mixture of ethanol and isopropanol to obtain the title compound (0.19 g, 83%) as colorless crystals.

m.p.: 180-182°C (decomposed);

IR spectrum (KBr) ν_{\max} cm^{-1} : 2995, 2976, 2914, 1591;

NMR spectrum (DMSO-d_6) δ ppm: 3.26(2H, t, J=5.1Hz), 3.84(3H, s), 4.45(2H, t, J=5.1Hz), 6.88(1H, s), 7.09-7.11(1H, m), 7.38-7.48(3H, m), 8.28(3H, brs).

Example 11

3-(2-Aminoethoxy)-5-(4-methoxyphenyl)isoxazole hydrochloride (Compound list No.: 350)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(4-methoxyphenyl)isoxazole

3-Hydroxy-5-(4-methoxyphenyl)isoxazole (0.22 g) and 2-(N-tert-butoxycarbonylamino)ethanol (0.20 g) were subjected to reaction and post-treatment in a similar manner to that described in Example 9(a) to obtain the title compound (0.32 g, 84%) as a colorless powder.

m.p.: 117-118°C;

IR spectrum (KBr) ν_{\max} cm^{-1} : 3344, 1719, 1623;

NMR spectrum (CDCl_3) δ ppm: 1.46(9H, s), 3.56(2H, q, J=5.2Hz), 3.86(3H, s), 4.34(2H, t, J=5.2Hz), 4.95(1H, brs), 6.02(1H, s), 6.96(2H, d, J=8.9Hz), 7.66(2H, d, J=8.9Hz).

(b) 3-(2-Aminoethoxy)-5-(4-methoxyphenyl)isoxazole hydrochloride

3-2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(4-methoxyphenyl)-isoxazole (0.30 g) was reacted in a similar manner to that described in Example 9(b) and the reaction product was recrystallized from a mixture of ethanol and isopropanol to obtain the title compound (0.17 g, 71%) as colorless crystals.

m.p.: 190-193°C(decomposed);

IR spectrum (KBr) ν_{\max} cm^{-1} : 2990, 2969, 2950, 2900, 1617;

NMR spectrum (DMSO-d_6) δ ppm: 3.25(2H, t, J=5.1Hz), 3.83(3H, s), 4.43(2H, t, J=5.1Hz), 6.69(1H, s), 7.09(2H, d, J=8.9Hz), 7.79(2H, d, J=8.9Hz), 8.25(3H, brs).

Example 12

3-(2-Aminoethoxy)-5-(2-chlorophenyl)isoxazole hydrochloride (Compound list No.: 111)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(2-chlorophenyl)isoxazole

5-(2-Chlorophenyl)-3-hydroxy-isoxazole (0.23 g) and 2-(N-tert-butoxycarbonylamino)ethanol (0.20 g) were subjected to reaction and post-treatment in a similar manner to that described in Example 9(a) to obtain the

title compound (0.35 g, 88%) as a colorless powder.

m.p.: 125-127°C;

IR spectrum (KBr) ν_{\max} cm^{-1} : 3333, 1710, 1697, 1618;

NMR spectrum (CDCl_3) δ ppm: 1.46(9H, s), 3.57(2H, q, $J=5.2\text{Hz}$), 4.37(2H, t, $J=5.2\text{Hz}$), 4.96(1H, brs), 6.59(1H, s), 7.33-7.43(2H, m), 7.46-7.54(1H, m), 7.87-7.94(1H, m).

(b) 3-(2-Aminoethoxy)-5-(2-chlorophenyl)isoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(2-chlorophenyl)isoxazole (0.33 g) was reacted in a similar manner to that described in Example 9(b) and the reaction product was recrystallized from isopropanol to obtain the title compound (0.20 g, 74%) as colorless crystals.

m.p.: 141-144°C (decomposed);

IR spectrum (KBr) ν_{\max} cm^{-1} : 3003, 2965, 1610;

NMR spectrum (DMSO-d_6) δ ppm: 3.278(2H, t, $J=5.1\text{Hz}$), 4.48(2H, t, $J=5.1\text{Hz}$), 6.80(1H, s), 7.52-7.60(2H, m), 7.68(1H, dd, $J=7.5\text{Hz}$, $J=1.9\text{Hz}$), 7.87(1H, dd, $J=7.5\text{Hz}$, $J=1.9\text{Hz}$), 8.27(3H, brs).

Example 13

3-(2-Aminoethoxy)-5-(3-chlorophenyl)isoxazole hydrochloride (Compound list No.: 125)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(3-chlorophenyl)isoxazole

5-(3-Chlorophenyl)-3-hydroxyisoxazole (0.23 g) and 2-(N-tert-butoxycarbonylamino)ethanol (0.20 g) were subjected to reaction and post-treatment in a similar manner to that described in Example 9(a) to obtain the title compound (0.34 g, 85%) as a colorless powder.

m.p.: 117-119°C;

IR spectrum (KBr) ν_{\max} cm^{-1} : 3385, 1685;

NMR spectrum (CDCl_3) δ ppm: 1.46(9H, s), 3.57(2H, q, $J=5.2\text{Hz}$), 4.36(2H, t, $J=5.2\text{Hz}$), 4.93(1H, brs), 6.16(1H, s), 7.36-7.44(2H, m), 7.57-7.67(1H, m), 7.70(1H, s).

(b) 3-(2-Aminoethoxy)-5-(3-chlorophenyl)isoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(3-chlorophenyl)isoxazole (0.33 g) was reacted in a similar manner to that described in Example 9(b) and the reaction product was recrystallized from a mixture of ethanol and isopropanol to obtain the title compound (0.21 g, 78%) as colorless crystals.

m.p.: 204-208°C (decomposed);

IR spectrum (KBr) ν_{\max} cm^{-1} : 2996, 2980, 2920, 1619;

NMR spectrum (DMSO- d_6) δ ppm: 3.26(2H, t, J=5.1Hz), 4.45(2H, t, J=5.1Hz), 6.98(1H, s), 7.56-7.63(2H, m), 7.80-7.84(1H, m), 7.96(1H, s), 8.22(3H, brs).

Example 14

3-(2-Aminoethoxy)-4-methyl-5-phenylisoxazole hydrochloride (Compound list

No.: 6)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-methyl-5-phenylisoxazole

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-phenylisoxazole (0.30 g) was dissolved in tetrahydrofuran (10 ml), and a solution of n-butyllithium/n-hexane (1.56M, 1.4 ml) was added dropwise thereto at -70°C under a nitrogen atmosphere, followed by stirring of the resulting mixture for 10 minutes. Then, methyl iodide (0.09 ml) was added dropwise to the reaction mixture, and the resulting mixture was stirred for 10 minutes. After the temperature of the reaction mixture was raised to 0°C, the reaction mixture was poured into ice-cold water, and an aqueous potassium dihydrogenphosphate solution was added to the mixture to adjust the pH to a value of 6. Then, the mixture was extracted with ethyl acetate and the organic layer was washed with a saturated aqueous NaCl solution, followed by drying over anhydrous magnesium sulfate. After filtration, the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (eluent: hexane/ethyl acetate = 4/1) to obtain the title compound (0.29 g, 94%) as a colorless powder.

m.p.: 118-120°C;

IR spectrum (KBr) ν_{\max} cm^{-1} : 3334, 2974, 1719, 1708;

NMR spectrum (CDCl₃) δ ppm: 1.46(9H, s), 2.12(3H, s), 3.59(2H, q, J=5.1Hz), 4.38(2H, t, J=5.1Hz), 4.94(1H, brs), 7.40-7.53(3H, m), 7.69(2H, d, J=7.9Hz).

(b) 3-(2-Aminoethoxy)-4-methyl-5-phenylisoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-methyl-5-phenylisoxazole (0.29 g) was allowed to react in a similar manner to that described in Example 9(b) and the reaction product was recrystallized from a mixture of methanol and ethanol to obtain the title compound (0.18g, 78%) as colorless crystals.

m.p.: 245-250°C (decomposed);

IR spectrum (KBr) ν_{\max} cm⁻¹: 3003, 2892, 1516;

NMR spectrum (DMSO-d₆) δ ppm: 2.13(3H, s), 3.28(2H, t, J=5.1Hz), 4.46(2H, t, J=5.1Hz), 7.50-7.59(3H, m), 7.73(2H, d, J=7.2Hz), 8.21(3H, brs).

Example 15

3-(2-Aminoethoxy)-4-ethyl-5-phenylisoxazole hydrochloride (Compound list No.: 7)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-ethyl-5-phenylisoxazole

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-phenylisoxazole (0.3 g) and ethyl iodide (0.12 ml) were subjected to reaction and post-treatment in a similar manner to that described in Example 14(a) to obtain the title compound (0.24 g, 73%) as a colorless oil.

IR spectrum (CHCl₃) ν_{\max} cm⁻¹: 3460, 2980, 1713;

NMR spectrum (CDCl₃) δ ppm: 1.21(3H, t, J=7.5Hz), 1.46(9H, s), 2.56(2H, q, J=7.5Hz), 3.59(2H, q, J=5.2Hz), 4.38(2H, t, J=5.2Hz), 4.90(1H, brs), 7.37-7.51(3H, m), 7.64-7.68(2H, m).

(b) 3-(2-Aminoethoxy)-4-ethyl-5-phenylisoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-ethyl-5-phenylisoxazole (0.21 g) was subjected to reaction and post-treatment in a similar manner to that described in Example 9(b) to obtain the title compound (0.12 g, 71%) as colorless crystals.

- 159 -

m.p.: 210-215°C (decomposed);

IR spectrum (KBr) ν_{\max} cm^{-1} : 2968, 2886, 1518;

NMR spectrum (DMSO- d_6) δ ppm: 1.15(3H, t, $J=7.5\text{Hz}$), 2.58(2H, q, $J=7.5\text{Hz}$), 3.29(2H, t, $J=5.1\text{Hz}$), 4.46(2H, t, $J=5.1\text{Hz}$), 7.51-7.59(3H, m), 7.67-7.70(2H, m), 8.25(3H, brs).

Example 16

3-(2-Aminoethoxy)-5-phenyl-4-propylisoxazole hydrochloride (Compound list No.: 8)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-phenyl-4-propylisoxazole

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-phenylisoxazole (0.3 g) and propyl iodide (0.29 ml) were subjected to reaction and post-treatment in a similar manner to that described in Example 14(a) to obtain the title compound (0.22 g, 65%) as a colorless oil.

IR spectrum (CHCl_3) ν_{\max} cm^{-1} : 3460, 2966, 1713;

NMR spectrum (CDCl_3) δ ppm: 0.97(3H, t, $J=7.4\text{Hz}$), 1.46(9H, s), 1.56-1.69(2H, m), 2.51(2H, t, $J=7.6\text{Hz}$), 3.58(2H, q, $J=5.2\text{Hz}$), 4.37(2H, t, $J=5.2\text{Hz}$), 4.90(1H, brs), 7.43-7.53(3H, m), 7.65-7.68(2H, m).

(b) 3-(2-Aminoethoxy)-5-phenyl-4-propylisoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-phenyl-4-propylisoxazole (0.18 g) was subjected to reaction and post-treatment in a similar manner to that described in Example 9(b) to obtain the title compound (0.10 g, 71%) as colorless crystals.

m.p.: 119-121°C (decomposed);

IR spectrum (KBr) ν_{\max} cm^{-1} : 2960, 2933, 2872, 1518;

NMR spectrum (DMSO- d_6) δ ppm: 0.91(3H, t, $J=7.3\text{Hz}$), 1.51-1.61(2H, m), 2.54(2H, t, $J=7.7\text{Hz}$), 3.28(2H, t, $J=5.1\text{Hz}$), 4.45(2H, t, $J=5.1\text{Hz}$), 7.50-7.59(3H, m), 7.68-7.70(2H, m), 8.20(3H, brs).

Example 17

3-(2-Aminoethoxy)-4-butyl-5-phenylisoxazole hydrochloride (Compound list No.: 10)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-butyl-5-phenylisoxazole

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-phenylisoxazole (0.3 g) and butyl iodide (0.17 ml) were subjected to reaction and post-treatment in a similar manner to that described in Example 14(a) to obtain the title compound (0.23 g, 66%) as a colorless oil.

IR spectrum (CHCl₃) ν_{\max} cm⁻¹: 3460, 2962, 1713;

NMR spectrum (CDCl₃) δ ppm: 0.93(3H, t, J=7.3Hz), 1.31-1.63(4H, m), 1.46(9H, s), 2.53(2H, t, J=7.6Hz), 3.58(2H, q, J=5.3Hz), 4.38(2H, t, J=5.3Hz), 4.90(1H, brs), 7.42-7.51(3H, m), 7.65-7.68(2H, m).

(b) 3-(2-Aminoethoxy)-4-butyl-5-phenylisoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-butyl-5-phenylisoxazole (0.20 g) was subjected to reaction and post-treatment in a similar manner to that described in Example 9(b) to obtain the title compound (0.12 g, 75%) as colorless crystals.

m.p.: 104-106°C;

IR spectrum (KBr) ν_{\max} cm⁻¹: 3006, 2963, 2951, 2931, 2869, 1516;

NMR spectrum (DMSO-d₆) δ ppm: 0.87(3H, t, J=7.3Hz), 1.27-1.37(2H, m), 1.47-1.55(2H, m), 2.57(2H, t, J=7.7Hz), 3.28(2H, t, J=5.1Hz), 4.46(2H, t, J=5.1Hz), 7.51-7.59(3H, m), 7.68-7.70(2H, m), 8.23(3H, brs).

Example 18

3-(2-Aminoethoxy)-4-hexyl-5-phenylisoxazole hydrochloride (Compound list No.: 1388)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-hexyl-5-phenylisoxazole

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-phenylisoxazole (0.4 g) and hexyl iodide (0.23 ml) were subjected to reaction and post-treatment in a similar manner to that described in Example 14(a) to obtain the title compound

(0.31 g, 61%) as a colorless powder.

IR spectrum (KBr) ν_{\max} cm^{-1} : 3387, 2936, 1715;

NMR spectrum (CDCl_3) δ ppm: 0.88(3H, t, $J=6.6\text{Hz}$), 1.23-1.42(6H, m), 1.46(9H, s), 1.53-1.70(2H, m), 2.52(2H, t, $J=7.7\text{Hz}$), 3.58(2H, q, $J=5.1\text{Hz}$), 4.37(2H, t, $J=5.1\text{Hz}$), 4.90(1H, brs), 7.40-7.51(3H, m), 7.65-7.68(2H, m).

(b) 3-(2-Aminoethoxy)-4-hexyl-5-phenylisoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-hexyl-5-phenylisoxazole (0.2 g) was subjected to reaction and post-treatment in a similar manner to that described in Example 9(b) to obtain the title compound (0.13 g, 81%) as colorless crystals.

m.p.: 99-101°C;

IR spectrum (KBr) ν_{\max} cm^{-1} : 2954, 2930, 1515;

NMR spectrum (DMSO-d_6) δ ppm: 0.83(3H, t, $J=6.9\text{Hz}$), 1.18-1.32(6H, m), 1.48-1.55(2H, m), 2.56(2H, t, $J=7.7\text{Hz}$), 3.28(2H, t, $J=5.1\text{Hz}$), 4.45(2H, t, $J=5.1\text{Hz}$), 7.50-7.59(3H, m), 7.67-7.69(2H, m), 8.16(3H, brs).

Example 19

3-(2-Aminoethoxy)-4-carboxy-5-phenylisoxazole hydrochloride (Compound list No.: 1408)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-carboxy-5-phenylisoxazole

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-phenylisoxazole (2.00 g) was dissolved in tetrahydrofuran (20 ml), and butyllithium (1.56M hexane solution, 9.3 ml) was added dropwise thereto at -70°C under a nitrogen atmosphere, and the resulting mixture was stirred for 10 minutes. Then, carbon dioxide gas was bubbled into the reaction mixture for 10 minutes and the temperature of the reaction mixture was raised to 0°C . The reaction mixture was poured into ice-cold water and the pH of the mixture was adjusted to a value of 6 with an aqueous potassium dihydrogenphosphate solution, followed by extraction with ethyl acetate. The organic layer was washed with a saturated aqueous NaCl solution and dried over anhydrous magnesium sulfate. After

filtration, the solvent was evaporated under reduced pressure. The residue was washed with ether to obtain the title compound (2.18 g, 95%) as a colorless powder.

IR spectrum (KBr) ν_{\max} cm^{-1} : 2982, 1706;

NMR spectrum (DMSO- d_6) δ ppm: 1.38(9H, s), 3.38(2H, q, $J=5.7\text{Hz}$), 4.29(2H, t, $J=5.7\text{Hz}$), 7.01(1H, brs), 7.51-7.62(3H, m), 7.84-7.91(2H, m).

(b) 3-(2-Aminoethoxy)-4-carboxy-5-phenylisoxazole hydrochloride

3-(2-(N-tert-butoxycarbonylamino)ethoxy)-4-carboxy-5-phenylisoxazole (0.12 g) was subjected to reaction and post-treatment in a similar manner to that described in Example 9(b) to obtain the title compound (0.06 g, 60%) as colorless crystals.

m.p.: 180-183°C (decomposed);

IR spectrum (KBr) ν_{\max} cm^{-1} : 3149, 2873, 2820, 1755, 1709;

NMR spectrum (DMSO- d_6) δ ppm: 3.30(2H, t, $J=5.3\text{Hz}$), 4.54(2H, t, $J=5.3\text{Hz}$), 7.54-7.65(3H, m), 7.85-7.87(2H, m), 8.22(3H, brs).

Example 20

3-(2-Aminoethoxy)-4-carbamoyl-5-phenylisoxazole hydrochloride (Compound list No.: 1414)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-carbamoyl-5-phenylisoxazole

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-carboxy-5-phenylisoxazole (0.6 g) was dissolved in tetrahydrofuran (6 ml) and carbonyl-diimidazole (0.31 g) was added thereto under ice-cooling with stirring, followed by stirring of the mixture at room temperature for 30 minutes. Aqueous ammonia (1 ml) was added dropwise to the reaction mixture and the mixture was stirred at room temperature for one hour. The reaction mixture was poured into ice-cold water and extracted with ethyl acetate. The organic layer was washed with a saturated aqueous NaCl solution and dried using anhydrous magnesium sulfate. After filtration, the solvent was evaporated under reduced pressure to obtain the title compound (0.60 g, quantitative) as a colorless

powder.

IR spectrum (KBr) ν_{\max} cm^{-1} : 3439, 3371, 3149, 1694, 1680;

NMR spectrum (DMSO- d_6) δ ppm: 1.39(9H, s), 3.39(2H, q, $J=5.0\text{Hz}$), 4.30(2H, t, $J=5.0\text{Hz}$), 7.38(1H, brs), 7.51-7.57(3H, m), 7.71(1H, brs), 7.91-7.94(2H, m).

(b) 3-(2-Aminoethoxy)-4-carbamoyl-5-phenylisoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-carbamoyl-5-phenylisoxazole (0.22 g) was subjected to reaction and post-treatment in a similar manner to that described in Example 9(b) to obtain the title compound (0.13 g, 72%) as colorless crystals.

m.p.: 225-230°C (decomposed);

IR spectrum (KBr) ν_{\max} cm^{-1} : 3407, 3213, 2963, 2878, 1662;

NMR spectrum (DMSO- d_6) δ ppm: 3.32(2H, t, $J=4.9\text{Hz}$), 4.54(2H, t, $J=4.9\text{Hz}$), 7.52-7.60(4H, m), 7.77(1H, brs), 7.92-7.94(2H, m), 8.29(3H, brs).

Example 21

3-(2-Aminoethoxy)-4-cyano-5-phenylisoxazole hydrochloride (Compound list No.: 1406)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-cyano-5-phenylisoxazole

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-carbamoyl-5-phenylisoxazole (0.38 g) was dissolved in dimethylformamide (4 ml), and phosphorus oxychloride (0.11 ml) was added dropwise thereto at 5°C under a nitrogen atmosphere, followed by stirring of the mixture at room temperature for 30 minutes. The reaction mixture was poured into ice-cold water and extracted with ethyl acetate. The organic layer was washed with a saturated aqueous NaCl solution and dried over anhydrous magnesium sulfate. After filtration, the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (eluent: hexane/ethyl acetate = 3/1) to obtain the title compound (0.28 g, 78%) as a colorless solid.

IR spectrum (KBr) ν_{\max} cm^{-1} : 3384, 2237, 1690, 1680;

NMR spectrum (CDCl₃) δ ppm: 1.47(9H, s), 3.61(2H, q, $J=5.2\text{Hz}$), 4.43(2H, t,

J=5.2Hz), 4.97(1H, brs), 7.52-7.64(3H, m), 8.01-8.05(2H, m).

(b) 3-(2-Aminoethoxy)-4-cyano-5-phenylisoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-cyano-5-phenylisoxazole (0.25 g) was subjected to reaction and post-treatment in a similar manner to that described in Example 9(b) to obtain the title compound (0.13 g, 65%) as colorless crystals.

m.p.: 200-205°C (decomposed);

IR spectrum (KBr) ν_{\max} cm^{-1} : 2967, 2236, 1611;

NMR spectrum (DMSO- d_6) δ ppm: 3.32(2H, t, J=5.1Hz), 4.60(2H, t, J=5.1Hz), 7.68-7.76(3H, m), 7.98-8.01(2H, m), 8.29(3H, brs).

Example 22

3-(2-Aminoethoxy)-4-methoxycarbonyl-5-phenylisoxazole hydrochloride

(Compound list No.: 1412)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-methoxycarbonyl-5-phenylisoxazole

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-carboxy-5-phenylisoxazole (0.2 g) was dissolved in a mixture of methanol and benzene (1:5, 10 ml), and trimethylsilyldiazomethane (0.6 ml, 2.0M hexane solution) was added dropwise thereto under ice-cooling, followed by stirring of the mixture at room temperature for 30 minutes. The reaction mixture was poured into an ice-cold water and extracted with ethyl acetate. The organic layer was washed with a saturated aqueous NaCl solution and dried over anhydrous magnesium sulfate. After filtration, the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (eluent: hexane/ethyl acetate = 3/1) to obtain the title compound (0.17 g, 81%) as a colorless solid.

IR spectrum (KBr) ν_{\max} cm^{-1} : 3355, 1725, 1691;

NMR spectrum (CDCl_3) δ ppm: 1.46(9H, s), 3.62(2H, q, J=5.1Hz), 3.85(3H, s), 4.43(3H, t, J=5.1Hz), 5.03(1H, brs), 7.46-7.54(3H, m), 7.87-7.91(2H, m).

(b) 3-(2-Aminoethoxy)-4-methoxycarbonyl-5-phenylisoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-methoxycarbonyl-5-phenylisoxazole (0.16 g) was subjected to reaction and post-treatment in a similar manner to that described in Example 9(b) to obtain the title compound (0.08 g, 62%) as colorless crystals.

m.p.: 195-198°C (decomposed);

IR spectrum (KBr) ν_{\max} cm^{-1} : 3148, 2870, 2848, 2821, 1708;

NMR spectrum (DMSO- d_6) δ ppm: 3.30(2H, t, J=5.2Hz), 3.77(3H, s), 4.54(2H, t, J=5.2Hz), 7.56-7.66(3H, m), 7.85-7.87(2H, m), 8.24(3H, brs).

Example 23

3-(2-Aminoethoxy)-4-methylaminocarbonyl-5-phenylisoxazole hydrochloride

(Compound list No.: 1416)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-methylaminocarbonyl-5-phenylisoxazole

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-carboxy-5-phenylisoxazole (0.2 g) and methylamine (30% methanol solution, 0.12 ml) were subjected to reaction and post-treatment in a similar manner to that described in Example 20(a) to obtain the title compound (0.18 g, 87%) as a colorless powder.

IR spectrum (KBr) ν_{\max} cm^{-1} : 3405, 3369, 1677;

NMR spectrum (CDCl₃) δ ppm: 1.45(9H, s), 2.97(3H, d, J=5.0Hz), 3.64(2H, q, J=5.1Hz), 4.44(2H, t, J=5.1Hz), 4.90(1H, brs), 7.30(1H, brs), 7.46-7.50(3H, m), 8.05-8.09(2H, m).

(b) 3-(2-Aminoethoxy)-4-methylaminocarbonyl-5-phenylisoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-methylaminocarbonyl-5-phenylisoxazole (0.17 g) was subjected to reaction and post-treatment in a similar manner to that described in Example 14(a) to obtain the title compound (0.08 g, 57%) as colorless crystals.

m.p.: 223-226°C (decomposed);

IR spectrum (KBr) ν_{\max} cm^{-1} : 3102, 2935, 2879, 1646;

NMR spectrum (DMSO- d_6) δ ppm: 2.78(3H, d, $J=4.4\text{Hz}$), 3.32(2H, t, $J=4.8\text{Hz}$), 4.53(2H, t, $J=4.8\text{Hz}$), 7.51-7.60(3H, m), 7.89-7.92(2H, m), 8.18(1H, d, $J=4.4\text{Hz}$), 8.31(3H, brs).

Example 24

3-(2-Aminoethoxy)-5-(4-chlorophenyl)-4-methylisoxazole hydrochloride

(Compound list No.: 148)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(4-chlorophenyl)-4-methylisoxazole

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(4-chlorophenyl)isoxazole (0.25 g) and methyl iodide (0.06 ml) were subjected to reaction and post-treatment in a similar manner to that described in Example 14(a) to obtain the title compound (0.23 g, 89%) as a colorless powder.

IR spectrum (KBr) ν_{\max} cm^{-1} : 3344, 2980, 1682;

NMR spectrum (CDCl_3) δ ppm: 1.46(9H, s), 2.10(3H, s), 3.58(2H, q, $J=5.2\text{Hz}$), 4.37(2H, t, $J=5.2\text{Hz}$), 4.90(1H, brs), 7.45(2H, d, $J=8.6\text{Hz}$), 7.63(2H, d, $J=8.6\text{Hz}$).

(b) 3-(2-Aminoethoxy)-5-(4-chlorophenyl)-4-methylisoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(4-chlorophenyl)-4-methylisoxazole (0.22 g) was subjected to reaction and post-treatment in a similar manner to that described in Example 9(b) to obtain the title compound (0.09 g, 50%) as colorless crystals.

m.p.: 248-253°C (decomposed);

IR spectrum (KBr) ν_{\max} cm^{-1} : 3020, 2991, 2884, 1513;

NMR spectrum (DMSO- d_6) δ ppm: 2.12(3H, s), 3.27(2H, t, $J=5.1\text{Hz}$), 4.45(2H, t, $J=5.1\text{Hz}$), 7.63(2H, d, $J=8.6\text{Hz}$), 7.75(2H, d, $J=8.6\text{Hz}$), 8.23(3H, brs).

Example 25

3-(2-Aminoethoxy)-5-(4-chlorophenyl)-4-ethylisoxazole hydrochloride

(Compound list No.: 149)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(4-chlorophenyl)-4-ethylisoxazole

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(4-chlorophenyl)isoxazole (0.4 g) and ethyl iodide (0.11 ml) were subjected to reaction and post-treatment in a similar manner to that described in Example 14(a) to obtain the title compound (0.26 g, 61%) as a colorless oil.

IR spectrum (CHCl₃) ν_{\max} cm⁻¹: 3460, 2980, 1713;

NMR spectrum (CDCl₃) δ ppm: 1.20(3H, t, J=7.4Hz), 1.46(9H, s), 2.54(2H, q, J=7.4Hz), 3.59(2H, q, J=5.2Hz), 4.37(2H, t, J=5.2Hz), 4.90(1H, brs), 7.45(2H, d, J=8.5Hz), 7.59(2H, d, J=8.5Hz).

(b) 3-(2-Aminoethoxy)-5-(4-chlorophenyl)-4-ethylisoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(4-chlorophenyl)-4-ethylisoxazole (0.22 g) was subjected to reaction and post-treatment in a similar manner to that described in Example 9(b) to obtain the title compound (0.13 g, 72%) as colorless crystals.

m.p.: 217-220°C (decomposed);

IR spectrum (KBr) ν_{\max} cm⁻¹: 2972, 2892, 1515;

NMR spectrum (DMSO-d₆) δ ppm: 1.14(3H, t, J=7.5Hz), 2.58(2H, q, J=7.5Hz), 3.28(2H, t, J=5.2Hz), 4.46(2H, t, J=5.2Hz), 7.64(2H, d, J=8.6Hz), 7.71(2H, d, J=8.6Hz), 8.20(3H, brs).

Example 26

3-(2-Aminoethoxy)-5-(4-chlorophenyl)-4-propylisoxazole hydrochloride

(Compound list No.: 150)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(4-chlorophenyl)-4-propylisoxazole

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(4-chlorophenyl)isoxazole

(0.4 g) and propyl iodide (0.14 ml) were subjected to reaction and post-treatment in a similar manner to that described in Example 14(a) to obtain the title compound (0.21 g, 47%) as a colorless powder.

IR spectrum (KBr) ν_{\max} cm^{-1} : 3392, 2963, 1685;

NMR spectrum (CDCl_3) δ ppm: 0.96(3H, t, $J=7.4\text{Hz}$), 1.46(9H, s), 1.57-1.69(2H, m), 2.49(2H, t, $J=7.6\text{Hz}$), 3.58(2H, q, $J=5.2\text{Hz}$), 4.37(2H, t, $J=5.2\text{Hz}$), 4.87(1H, brs), 7.45(2H, d, $J=8.6\text{Hz}$), 7.60(2H, d, $J=8.6\text{Hz}$).

(b) 3-(2-Aminoethoxy)-5-(4-chlorophenyl)-4-propylisoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(4-chlorophenyl)-4-propylisoxazole (0.19 g) was subjected to reaction and post-treatment in a similar manner to that described in Example 9(b) to obtain the title compound (0.13 g, 81%) as colorless crystals.

m.p.: 146-149°C (decomposed);

IR spectrum (KBr) ν_{\max} cm^{-1} : 2958, 2871, 2829, 1516;

NMR spectrum (DMSO-d_6) δ ppm: 0.90(3H, t, $J=7.3\text{Hz}$), 1.50-1.59(2H, m), 2.54(2H, t, $J=7.3\text{Hz}$), 3.28(2H, t, $J=5.2\text{Hz}$), 4.45(2H, t, $J=5.2\text{Hz}$), 7.63(2H, d, $J=8.7\text{Hz}$), 7.72(2H, d, $J=8.7\text{Hz}$), 8.21(3H, brs).

Example 27

3-(2-Aminoethoxy)-5-(4-methylphenyl)isoxazole hydrochloride (Compound list No.: 260)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(4-methylphenyl)isoxazole

3-Hydroxy-5-(4-methylphenyl)isoxazole (1.5 g) and 2-(N-tert-butoxycarbonylamino)ethanol (1.5 g) were subjected to reaction and post-treatment in a similar manner to that described in Example 9(a) to obtain the title compound (2.2 g, 82%) as a colorless powder.

IR spectrum (KBr) ν_{\max} cm^{-1} : 3375, 3356, 3332, 1719, 1684;

NMR spectrum (CDCl_3) δ ppm: 1.46(9H, s), 2.40(3H, s), 3.56(2H, q, $J=5.1\text{Hz}$), 4.35(2H, t, $J=5.1\text{Hz}$), 4.94(1H, brs), 6.09(1H, s), 7.26(2H, d, $J=8.1\text{Hz}$), 7.61(2H, d, $J=8.1\text{Hz}$).

(b) 3-(2-Aminoethoxy)-5-(4-methylphenyl)isoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(4-methylphenyl)isoxazole (0.3 g) was subjected to reaction and post-treatment in a similar manner to that described in Example 9(b) to obtain the title compound (0.16 g, 67%) as colorless crystals.

m.p.: 215-220°C (decomposed);

IR spectrum (KBr) ν_{\max} cm^{-1} : 2993, 2979, 1622;

NMR spectrum (DMSO- d_6) δ ppm: 2.37(3H, s), 3.25(2H, t, J=5.0Hz), 4.44(2H, t, J=5.0Hz), 6.76(1H, s), 7.35(2H, d, J=8.1Hz), 7.74(2H, d, J=8.1Hz), 8.26(3H, brs).

Example 28

3-(2-Aminoethoxy)-5-(4-trifluoromethylphenyl)isoxazole hydrochloride

(Compound list No.: 332)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(4-trifluoromethylphenyl)isoxazole

3-Hydroxy-5-(4-trifluoromethylphenyl)isoxazole (0.2 g) and 2-(N-tert-butoxycarbonylamino)ethanol (0.15 g) were subjected to reaction and post-treatment in a similar manner to that described in Example 9(a) to obtain the title compound (0.26 g, 81%) as a colorless powder.

IR spectrum (KBr) ν_{\max} cm^{-1} : 3376, 1679, 1608;

NMR spectrum (CDCl₃) δ ppm: 1.46(9H, s), 3.58(2H, q, J=5.1Hz), 4.37(2H, t, J=5.1Hz), 4.92(1H, brs), 6.25(1H, s), 7.72(2H, d, J=8.2Hz), 7.84(2H, d, J=8.2Hz).

(b) 3-(2-Aminoethoxy)-5-(4-trifluoromethylphenyl)isoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(4-trifluoromethylphenyl)isoxazole (0.24 g) was subjected to reaction and post-treatment in a similar manner to that described in Example 9(b) to obtain the title compound (0.14 g, 70%) as colorless crystals.

m.p.: 226-232°C (decomposed);

IR spectrum (KBr) ν_{\max} cm^{-1} : 2993, 2965, 2914, 1608;

NMR spectrum (DMSO-d₆) δ ppm: 3.27(2H, t, J=5.1Hz), 4.47(2H, t, J=5.1Hz), 7.07(1H, s), 7.93(2H, d, J=8.3Hz), 8.09(2H, d, J=8.3Hz), 8.26(3H, brs).

Example 29

3-(2-Aminoethoxy)-5-(4-fluorophenyl)isoxazole hydrochloride (Compound list No.: 66)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(4-fluorophenyl)isoxazole

5-(4-Fluorophenyl)-3-hydroxyisoxazole (0.06 g) and 2-(N-tert-butoxycarbonylamino)ethanol (0.06 g) were subjected to reaction and post-treatment in a similar manner to that described in Example 9(a) to obtain the title compound (0.08 g, 74%) as a colorless powder.

IR spectrum (KBr) ν_{\max} cm⁻¹: 3375, 1682;

NMR spectrum (CDCl₃) δ ppm: 1.46(9H, s), 3.56(2H, q, J=5.1Hz), 4.35(2H, t, J=5.1Hz), 4.93(1H, brs), 6.09(1H, s), 7.15(2H, t, J=8.6Hz), 7.71(2H, dd, J=8.6Hz, J=5.3Hz).

(b) 3-(2-Aminoethoxy)-5-(4-fluorophenyl)isoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(4-fluorophenyl)isoxazole (0.05 g) was subjected to reaction and post-treatment in a similar manner to that described in Example 9(b) to obtain the title compound (0.02 g, 56%) as colorless crystals.

m.p.: 232-238°C (decomposed);

IR spectrum (KBr) ν_{\max} cm⁻¹: 2993, 2978, 2911, 1621;

NMR spectrum (DMSO-d₆) δ ppm: 3.26(2H, t, J=5.0Hz), 4.44(2H, t, J=5.0Hz), 6.84(1H, s), 7.40(2H, t, J=8.8Hz), 7.93(2H, dd, J=8.8Hz, J=5.3Hz), 8.25(3H, brs).

Example 30

3-(2-Aminoethoxy)-5-(1-naphthyl)isoxazole hydrochloride (Compound list No.: 475)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(1-naphthyl)isoxazole

3-Hydroxy-5-(1-naphthyl)isoxazole (0.20 g) and 2-(N-tert-butoxy-

- 171 -

carbonylamino)ethanol (0.17 g) were subjected to reaction and post-treatment in a similar manner to that described in Example 9(a) to obtain the title compound (0.30 g, 89%) as a colorless powder.

IR spectrum (KBr) ν_{\max} cm^{-1} : 3318, 1712;

NMR spectrum (CDCl_3) δ ppm: 1.47(9H, s), 3.61(2H, q, $J=5.3\text{Hz}$), 4.42(2H, t, $J=5.3\text{Hz}$), 4.97(1H, brs), 6.25(1H, s), 7.52-7.61(3H, m), 7.77-7.79(1H, m), 7.90-7.99(2H, m), 8.27-8.31(1H, m).

(b) 3-(2-Aminoethoxy)-5-(1-naphthyl)isoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(1-naphthyl)isoxazole (0.28 g) was subjected to reaction and post-treatment in a similar manner to that described in Example 9(b) to obtain the title compound (0.18 g, 78%) as colorless crystals.

m.p.: 201-205°C (decomposed);

IR spectrum (KBr) ν_{\max} cm^{-1} : 3002, 2968, 2913, 1602;

NMR spectrum (DMSO-d_6) δ ppm: 3.31(2H, t, $J=5.1\text{Hz}$), 4.52(2H, t, $J=5.1\text{Hz}$), 6.83(1H, s), 7.63-7.70(3H, m), 7.87-7.89(1H, m), 8.07-8.25(3H, m), 8.30(3H, brs).

Example 31

3-(2-Aminoethoxy)-4-bromo-5-phenylisoxazole hydrochloride (Compound list No.: 1357)

(a) 4-Bromo-3-(2-(N-tert-butoxycarbonylamino)ethoxy)-5-phenylisoxazole

4-Bromo-3-hydroxy-5-phenylisoxazole (0.2 g) and 2-(N-tert-butoxycarbonylamino)ethanol (0.15 g) were subjected to reaction and post-treatment in a similar manner to that described in Example 9(a) to obtain the title compound (0.26 g, 81%) as a colorless powder.

IR spectrum (KBr) ν_{\max} cm^{-1} : 3352, 2976, 1719, 1709;

NMR spectrum (CDCl_3) δ ppm: 1.47(9H, s), 3.61(2H, q, $J=5.1\text{Hz}$), 4.41(2H, t, $J=5.1\text{Hz}$), 4.97(1H, brs), 7.47-7.55(3H, m), 7.98-8.04(2H, m).

(b) 3-(2-Aminoethoxy)-4-bromo-5-phenylisoxazole hydrochloride

4-Bromo-3-(2-(N-tert-butoxycarbonylamino)ethoxy)-5-phenylisoxazole (0.24 g) was subjected to reaction and post-treatment in a similar manner to that described in Example 9(b) to obtain the title compound (0.13 g, 65%) as colorless crystals.

m.p.: 192-198°C (decomposed);

IR spectrum (KBr) ν_{\max} cm^{-1} : 2991, 2921, 2894, 1614, 1594, 1574;

NMR spectrum (DMSO- d_6) δ ppm: 3.32(2H, t, J=5.1Hz), 4.54(2H, t, J=5.1Hz), 7.60-7.66(3H, m), 7.95-7.98(2H, m), 8.19(3H, brs).

Example 32

3-(2-Aminoethoxy)-4-iodo-5-phenylisoxazole hydrochloride (Compound list No.: 1359)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-iodo-5-phenylisoxazole

3-Hydroxy-4-iodo-5-phenylisoxazole (0.2 g) and 2-(N-tert-butoxycarbonylamino)ethanol (0.12 g) were subjected to reaction and post-treatment in a similar manner to that described in Example 9(a) to obtain the title compound (0.22 g, 73%) as a colorless powder.

IR spectrum (KBr) ν_{\max} cm^{-1} : 3328, 2977, 1696;

NMR spectrum (CDCl₃) δ ppm: 1.47(9H, s), 3.61(2H, q, J=5.2Hz), 4.41(2H, t, J=5.2Hz), 4.97(1H, brs), 7.47-7.55(3H, m), 7.99-8.06(2H, m).

(b) 3-(2-Aminoethoxy)-4-iodo-5-phenylisoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-iodo-5-phenylisoxazole (0.20 g) was subjected to reaction and post-treatment in a similar manner to that described in Example 9(b) to obtain the title compound (0.12 g, 71%) as colorless crystals.

m.p.: 201-206°C (decomposed);

IR spectrum (KBr) ν_{\max} cm^{-1} : 2961, 2912, 1591;

NMR spectrum (DMSO- d_6) δ ppm: 3.30(2H, t, J=5.3Hz), 4.51(2H, t, J=5.3Hz), 7.58-7.64(3H, m), 7.96-8.01(2H, m), 8.22(3H, brs).

Example 33

3-(2-Aminoethoxy)-5-(4-isopropylphenyl)isoxazole hydrochloride (Compound list No.: 1618)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(4-isopropylphenyl)isoxazole

3-Hydroxy-5-(4-isopropylphenyl)isoxazole (0.2 g) and 2-(N-tert-butoxycarbonylamino)ethanol (0.17 g) were subjected to reaction and post-treatment in a similar manner to that described in Example 9(a) to obtain the title compound (0.26 g, 77%) as a colorless powder.

IR spectrum (KBr) ν_{\max} cm^{-1} : 3326, 2973, 1713, 1697, 1623;

NMR spectrum (CDCl_3) δ ppm: 1.27(6H, d, $J=6.9\text{Hz}$), 1.46(9H, s), 2.95(1H, qq, $J=6.9\text{Hz}$, $J=6.9\text{Hz}$), 3.56(2H, q, $J=5.1\text{Hz}$), 4.35(2H, t, $J=5.1\text{Hz}$), 4.94(1H, brs), 6.09(1H, s), 7.31(2H, d, $J=8.3\text{Hz}$), 7.65(2H, d, $J=8.3\text{Hz}$).

(b) 3-(2-Aminoethoxy)-5-(4-isopropylphenyl)isoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(4-isopropylphenyl)-isoxazole (0.25 g) was subjected to reaction and post-treatment in a similar manner to that described in Example 9(b) to obtain the title compound (0.11 g, 55%) as colorless crystals.

m.p.: 162-166°C (decomposed);

IR spectrum (KBr) ν_{\max} cm^{-1} : 3000, 2959, 2924, 1623, 1604;

NMR spectrum (DMSO-d_6) δ ppm: 1.23(6H, d, $J=6.9\text{Hz}$), 2.96(1H, qq, $J=6.9\text{Hz}$, $J=6.9\text{Hz}$), 3.26(2H, t, $J=5.1\text{Hz}$), 4.44(2H, t, $J=5.1\text{Hz}$), 6.77(1H, s), 7.41(2H, d, $J=8.2\text{Hz}$), 7.77(2H, d, $J=8.2\text{Hz}$), 8.20(3H, brs).

Example 34

3-(2-Aminoethoxy)-5-(2-methylphenyl)isoxazole hydrochloride (Compound list No.: 224)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(2-methylphenyl)isoxazole

3-Hydroxy-5-(2-methylphenyl)isoxazole (0.2 g) and 2-(N-tert-butoxycarbonylamino)ethanol (0.2 g) were subjected to reaction and post-treatment in

a similar manner to that described in Example 9(a) to obtain the title compound (0.31 g, 86%) as a colorless powder.

IR spectrum (KBr) ν_{\max} cm^{-1} : 3315, 1710, 1619;

NMR spectrum (CDCl_3) δ ppm: 1.46(9H, s), 2.49(3H, s), 3.57(2H, q, $J=5.2\text{Hz}$), 4.37(2H, t, $J=5.2\text{Hz}$), 4.95(1H, brs), 6.06(1H, s), 7.19-7.39(3H, m), 7.65-7.69(1H, m).

(b) 3-(2-Aminoethoxy)-5-(2-methylphenyl)isoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(2-methylphenyl)isoxazole (0.30 g) was subjected to reaction and post-treatment in a similar manner to that described in Example 9(b) to obtain the title compound (0.18 g, 75%) as colorless crystals.

m.p.: 165-167°C;

IR spectrum (KBr) ν_{\max} cm^{-1} : 2996, 2976, 2911, 1613;

NMR spectrum (DMSO-d_6) δ ppm: 2.46(3H, s), 3.26(2H, t, $J=5.2\text{Hz}$), 4.46(2H, t, $J=5.2\text{Hz}$), 6.59(1H, s), 7.34-7.46(3H, m), 7.67-7.69(1H, m), 8.21(3H, brs).

Example 35

3-(2-Aminoethoxy)-5-(4-phenylphenyl)isoxazole hydrochloride (Compound list No.: 368)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(4-phenylphenyl)isoxazole

3-Hydroxy-5-(4-phenylphenyl)isoxazole (0.2 g) and 2-(N-tert-butoxycarbonylamino)ethanol (0.15 g) were subjected to reaction and post-treatment in a similar manner to that described in Example 9(a) to obtain the title compound (0.22 g, 69%) as a colorless powder.

IR spectrum (KBr) ν_{\max} cm^{-1} : 3345, 1694;

NMR spectrum (CDCl_3) δ ppm: 1.46(9H, s), 3.58(2H, q, $J=5.1\text{Hz}$), 4.37(2H, t, $J=5.1\text{Hz}$), 4.95(1H, brs), 6.18(1H, s), 7.37-7.51(3H, m), 7.61-7.70(4H, m), 7.79-7.82(2H, m).

(b) 3-(2-Aminoethoxy)-5-(4-phenylphenyl)isoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(4-phenylphenyl)isoxazole

(0.2 g) was subjected to reaction and post-treatment in a similar manner to that described in Example 9(b) to obtain the title compound (0.1 g, 63%) as colorless crystals.

m.p.: 212-218°C (decomposed);

IR spectrum (KBr) ν_{\max} cm^{-1} : 2996, 2966, 2909, 1619, 1602;

NMR spectrum (DMSO- d_6) δ ppm: 3.27(2H, t, J=5.1Hz), 4.46(2H, t, J=5.1Hz), 6.90(1H, s), 7.41-7.55(3H, m), 7.74-7.76(2H, m), 7.84-7.87(2H, m), 7.94-7.96(2H, m), 8.18(3H, brs).

Example 36

3-(2-Aminoethoxy)-5-(4-phenoxyphenyl)isoxazole hydrochloride (Compound list No.: 1632)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(4-phenoxyphenyl)isoxazole

3-Hydroxy-5-(4-phenoxyphenyl)isoxazole (0.2 g) and 2-(N-tert-butoxycarbonylamino)ethanol (0.14 g) were subjected to reaction and post-treatment in a similar manner to that described in Example 9(a) to obtain the title compound (0.23 g, 74%) as a colorless powder.

IR spectrum (KBr) ν_{\max} cm^{-1} : 3331, 1720;

NMR spectrum (CDCl₃) δ ppm: 1.46(9H, s), 3.56(2H, q, J=5.2Hz), 4.34(2H, t, J=5.2Hz), 4.95(1H, brs), 6.06(1H, s), 7.02-7.08(4H, m), 7.15-7.20(1H, m), 7.35-7.42(2H, m), 7.65-7.70(2H, m).

(b) 3-(2-Aminoethoxy)-5-(4-phenoxyphenyl)isoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(4-phenoxyphenyl)-isoxazole (0.18 g) was subjected to reaction and post-treatment in a similar manner to that described in Example 9(b) to obtain the title compound (0.1 g, 67%) as colorless crystals.

m.p.: 207-213°C (decomposed) ;

IR spectrum (KBr) ν_{\max} cm^{-1} : 3000, 2959, 2909, 1625 ;

NMR spectrum (DMSO- d_6) δ ppm : 3.26(2H, t, J=5.1Hz), 4.43(2H, t, J=5.1Hz), 6.76(1H, s), 7.10-7.14(4H, m), 7.21-7.25(1H, m), 7.44-7.48(2H, m), 7.85-7.88(2H,

m), 8.18(3H, brs).

Example 37

3-(2-Aminoethoxy)-5-(2-trifluoromethylphenyl)isoxazole hydrochloride

(Compound list No.: 296)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(2-trifluoromethylphenyl)isoxazole

3-Hydroxy-5-(2-trifluoromethylphenyl)isoxazole (0.3 g) and 2-(N-tert-butoxycarbonylamino)ethanol (0.23 g) were subjected to reaction and post-treatment in a similar manner to that described in Example 9(a) to obtain the title compound (0.36 g, 74%) as a colorless oil.

IR spectrum (CHCl₃) ν_{\max} cm⁻¹: 3457, 2983, 1712;

NMR spectrum (CDCl₃) δ ppm: 1.46(9H, s), 3.58(2H, q, J=5.2Hz), 4.37(2H, t, J=5.2Hz), 4.96(1H, brs), 6.21(1H, s), 7.56-7.69(2H, m), 7.75-7.82(2H, m).

(b) 3-(2-Aminoethoxy)-5-(2-trifluoromethylphenyl)isoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(2-trifluoromethylphenyl)-isoxazole (0.35 g) was subjected to reaction and post-treatment in a similar manner to that described in Example 9(b) to obtain the title compound (0.22 g, 76%) as colorless crystals.

m.p. : 118-122°C (decomposed);

IR spectrum (KBr) ν_{\max} cm⁻¹: 2996, 2917, 1622;

NMR spectrum (DMSO-d₆) δ ppm: 3.28(2H, t, J=5.1Hz), 4.46(2H, t, J=5.1Hz), 6.63(1H, s), 7.79-7.89(3H, m), 7.96-7.98(1H, m), 8.22(3H, brs).

Example 38

3-(2-Aminoethoxy)-5-(4-hydroxyphenyl)isoxazole hydrochloride (Compound list No.: 1674)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(4-hydroxyphenyl)isoxazole

3-Hydroxy-5-(4-hydroxyphenyl)isoxazole (0.1 g) and 2-(N-tert-butoxycarbonylamino)ethanol (0.1 g) were subjected to reaction and post-treatment in

a similar manner to that described in Example 9(a) to obtain the title compound (0.12 g, 67%) as a colorless powder.

IR spectrum (KBr) ν_{\max} cm^{-1} : 3331, 3247, 1698, 1666, 1645, 1619;

NMR spectrum (DMSO- d_6) δ ppm: 1.38(9H, s), 3.32(2H, q, $J=5.6\text{Hz}$), 4.17(2H, t, $J=5.6\text{Hz}$), 6.53(1H, s), 6.87(2H, d, $J=8.8\text{Hz}$), 7.05(1H, brs), 7.63(2H, d, $J=8.8\text{Hz}$), 8.98(1H, brs).

(b) 3-(2-Aminoethoxy)-5-(4-hydroxyphenyl)isoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(4-hydroxyphenyl)-isoxazole (0.12 g) was subjected to reaction and post-treatment in a similar manner to that described in Example 9(b) to obtain the title compound (0.05 g, 56%) as colorless crystals.

m.p.: 240-245°C (decomposed);

IR spectrum (KBr) ν_{\max} cm^{-1} : 3145, 3056, 1620;

NMR spectrum (DMSO- d_6) δ ppm: 3.24(2H, t, $J=5.1\text{Hz}$), 4.41(2H, t, $J=5.1\text{Hz}$), 6.58(1H, s), 6.90(2H, d, $J=8.7\text{Hz}$), 7.67(2H, d, $J=8.7\text{Hz}$), 8.13(3H, brs).

Example 39

3-(2-Aminoethoxy)-5-(2,4-dichlorophenyl)isoxazole hydrochloride (Compound list No.: 170)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(2,4-dichlorophenyl)isoxazole

5-(2,4-Dichlorophenyl)-3-hydroxyisoxazole (0.2 g) and 2-(N-tert-butoxycarbonylamino)ethanol (0.17 g) were subjected to reaction and post-treatment in a similar manner to that described in Example 9(a) to obtain the title compound (0.26 g, 81%) as a colorless powder.

IR spectrum (KBr) ν_{\max} cm^{-1} : 3386, 1681, 1606;

NMR spectrum (CDCl_3) δ ppm: 1.46(9H, s), 3.57(2H, q, $J=5.2\text{Hz}$), 4.37(2H, t, $J=5.2\text{Hz}$), 4.94(1H, brs), 6.58(1H, s), 7.38(1H, dd, $J=8.5\text{Hz}$, $J=2.1\text{Hz}$), 7.53(1H, d, $J=2.1\text{Hz}$), 7.85(1H, d, $J=8.5\text{Hz}$).

(b) 3-(2-Aminoethoxy)-5-(2,4-dichlorophenyl)isoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(2,4-dichlorophenyl)-isoxazole (0.24 g) was subjected to reaction and post-treatment in a similar manner to that described in Example 9(b) to obtain the title compound (0.12 g, 60%) as colorless crystals.

m.p.: 192-195°C (decomposed);

IR spectrum (KBr) ν_{\max} cm^{-1} : 3069, 3005, 2967, 1611;

NMR spectrum (DMSO- d_6) δ ppm: 3.27(2H, t, J=5.1Hz), 4.47(2H, t, J=5.1Hz), 6.84(1H, s), 7.64(1H, dd, J=8.7Hz, J=2.1Hz), 7.89(1H, d, J=2.1Hz), 7.90(1H, d, J=8.7Hz), 8.20(3H, brs).

Example 40

3-(2-Aminoethoxy)-5-(3,4-dichlorophenyl)isoxazole hydrochloride (Compound list No.: 1604)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(3,4-dichlorophenyl)isoxazole

5-(3,4-Dichlorophenyl)-3-hydroxyisoxazole (0.3 g) and 2-(N-tert-butoxycarbonylamino)ethanol (0.23 g) were subjected to reaction and post-treatment in a similar manner to that described in Example 9(a) to obtain the title compound (0.41 g, 85%) as a colorless powder.

IR spectrum (KBr) ν_{\max} cm^{-1} : 3369, 1687;

NMR spectrum (CDCl₃) δ ppm: 1.46(9H, s), 3.56(2H, q, J=5.1Hz), 4.35(2H, t, J=5.1Hz), 4.91(1H, brs), 6.16(1H, s), 7.54(2H, s), 7.80(1H, s).

(b) 3-(2-Aminoethoxy)-5-(3,4-dichlorophenyl)isoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(3,4-dichlorophenyl)-isoxazole (0.4 g) was subjected to reaction and post-treatment in a similar manner to that described in Example 9(b) to obtain the title compound (0.22 g, 67%) as colorless crystals.

m.p.: 202-210°C (decomposed);

IR spectrum (KBr) ν_{\max} cm^{-1} : 2993, 2977, 2915, 1617;

NMR spectrum (DMSO- d_6) δ ppm: 3.26 (2H, t, J=5.1Hz), 4.45 (2H, t,

J=5.1Hz), 7.02 (1H, s), 7.84 (1H, d, J=8.9Hz), 7.85 (1H, d, J=8.9Hz), 8.17 (4H, brs).

Example 41

3-(2-Aminoethoxy)-5-(2,3-dichlorophenyl)isoxazole hydrochloride (Compound list No.: 1526)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(2,3-dichlorophenyl)isoxazole
5-(2,3-Dichlorophenyl)-3-hydroxyisoxazole (0.3 g) and 2-(N-tert-butoxycarbonylamino)ethanol (0.23 g) were subjected to reaction and post-treatment in a similar manner to that described in Example 9(a) to obtain the title compound (0.37 g, 77%) as a colorless powder.

IR spectrum (KBr) ν_{\max} cm^{-1} : 3381, 1688, 1607;

NMR spectrum (CDCl_3) δ ppm: 1.46 (9H, s), 3.58 (2H, q, J=5.1Hz), 4.37 (2H, t, J=5.1Hz), 4.95 (1H, brs), 6.60 (1H, s), 7.34 (1H, t, J=7.9Hz), 7.58 (1H, dd, J=7.9Hz, J=1.4Hz), 7.81 (1H, dd, J=7.9Hz, J=1.4Hz).

(b) 3-(2-Aminoethoxy)-5-(2,3-dichlorophenyl)isoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(2,3-dichlorophenyl)-isoxazole (0.36 g) was subjected to reaction and post-treatment in a similar manner to that described in Example 9(b) to obtain the title compound (0.20 g, 67%) as colorless crystals.

m.p.: 183-186°C (decomposed);

IR spectrum (KBr) ν_{\max} cm^{-1} : 3010, 2979, 2908, 1604;

NMR spectrum (DMSO-d_6) δ ppm: 3.27 (2H, t, J=5.1Hz), 4.48 (2H, t, J=5.1Hz), 6.87 (1H, s), 7.57 (1H, t, J=7.8Hz), 7.82-7.87 (2H, m), 8.19 (3H, brs).

Example 42

3-(2-Aminoethoxy)-5-(2,6-dichlorophenyl)isoxazole hydrochloride (Compound list No.: 188)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(2,6-dichlorophenyl)isoxazole
5-(2,6-Dichlorophenyl)-3-hydroxyisoxazole (0.3 g) and 2-(N-tert-butoxycarbonylamino)ethanol (0.23 g) were subjected to reaction and post-

treatment in a similar manner to that described in Example 9(a) to obtain the title compound (0.31 g, 65%) as a colorless powder.

IR spectrum (KBr) ν_{\max} cm^{-1} : 3358, 1703, 1626;

NMR spectrum (CDCl_3) δ ppm: 1.46 (9H, s), 3.58 (2H, q, $J=5.1\text{Hz}$), 4.39 (2H, t, $J=5.1\text{Hz}$), 4.96 (1H, brs), 6.11 (1H, s), 7.32-7.44 (3H, m).

(b) 3-(2-Aminoethoxy)-5-(2,6-dichlorophenyl)isoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(2,6-dichlorophenyl)-isoxazole (0.16 g) was subjected to reaction and post-treatment in a similar manner to that described in Example 9(b) to obtain the title compound (0.07 g, 54%) as colorless crystals.

m.p.: 148-151°C (decomposed);

IR spectrum (KBr) ν_{\max} cm^{-1} : 3005, 2966, 2935, 1631;

NMR spectrum (DMSO-d_6) δ ppm: 3.27 (2H, t, $J=5.0\text{Hz}$), 4.47 (2H, t, $J=5.0\text{Hz}$), 6.69 (1H, s), 7.61-7.71 (3H, m), 8.15 (3H, brs).

Example 43

3-(2-Aminoethoxy)-5-(2,4-difluorophenyl)isoxazole hydrochloride (Compound list No.: 93)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(2,4-difluorophenyl)isoxazole

5-(2,4-difluorophenyl)-3-hydroxyisoxazole (0.3 g) and 2-(N-tert-butoxycarbonylamino)ethanol (0.27 g) were subjected to reaction and post-treatment in a similar manner to that described in Example 9(a) to obtain the title compound (0.44 g, 85%) as a colorless powder.

IR spectrum (KBr) ν_{\max} cm^{-1} : 3382, 1694;

NMR spectrum (CDCl_3) δ ppm: 1.46 (9H, s), 3.57 (2H, q, $J=5.1\text{Hz}$), 4.36 (2H, t, $J=5.1\text{Hz}$), 4.93 (1H, brs), 6.30 (1H, s), 6.90-7.05 (2H, m), 7.85-7.94 (1H, m).

(b) 3-(2-Aminoethoxy)-5-(2,4-difluorophenyl)isoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(2,4-difluorophenyl)-isoxazole (0.42 g) was subjected to reaction and post-treatment in a similar

manner to that described in Example 9(b) to obtain the title compound (0.32 g, 94%) as colorless crystals.

m.p.: 226-232°C (decomposed);

IR spectrum (KBr) ν_{\max} cm^{-1} : 2997, 2980, 2922, 1626;

NMR spectrum (DMSO- d_6) δ ppm: 3.26 (2H, t, $J=5.1\text{Hz}$), 4.47 (2H, t, $J=5.1\text{Hz}$), 6.63 (1H, s), 7.29-7.34 (1H, m), 7.53-7.59 (1H, m), 7.94-8.00 (1H, m), 8.21 (3H, brs).

Example 44

3-(2-Aminoethoxy)-4-(1-chloroethyl)-5-phenylisoxazole hydrochloride

(Compound list No.: 1390)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-(1-hydroxyethyl)-5-phenylisoxazole

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-phenylisoxazole (0.8 g) was dissolved in tetrahydrofuran (16 ml), and butyllithium (1.6M hexane solution, 3.7 ml) was added dropwise thereto at -70°C under a nitrogen atmosphere, and the mixture was stirred for 10 minutes. After acetaldehyde (0.22 ml) was added dropwise to the reaction mixture, the resulting mixture was stirred for 10 minutes and the temperature of the mixture was raised to 0°C . The mixture was then poured into ice-cold water and the pH of the mixture was adjusted to a value of 6 with an aqueous potassium dihydrogenphosphate solution. After the reaction mixture was extracted with ethyl acetate and the organic layer was washed with a saturated aqueous NaCl solution, the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (eluent: hexane/ethyl acetate = 2/1) to obtain the title compound (0.90 g, 98%) as a colorless oil.

IR spectrum (CHCl_3) ν_{\max} cm^{-1} : 3602, 3459, 2982, 2936, 1712;

NMR spectrum (CDCl_3) δ ppm: 1.45 (9H, s), 1.60 (3H, d, $J=6.7\text{Hz}$), 2.64 (1H, brs), 3.58 (2H, q, $J=5.0\text{Hz}$), 4.39 (2H, t, $J=5.0\text{Hz}$), 4.90-5.10 (2H, m), 7.45-7.50 (3H, m), 7.64-7.69 (2H, m).

(b) 3-(2-Aminoethoxy)-4-(1-chloroethyl)-5-phenylisoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-(1-hydroxyethyl)-5-phenylisoxazole (0.2 g) was subjected to reaction and post-treatment in a similar manner to that described in Example 1(b) to obtain the title compound (0.13 g, 75%) as colorless crystals.

m.p.: 200-204°C (decomposed);

IR spectrum (KBr) ν_{\max} cm^{-1} : 2975, 1640;

NMR spectrum (DMSO- d_6) δ ppm: 1.86 (3H, d, $J=7.0\text{Hz}$), 3.27 (2H, t, $J=5.2\text{Hz}$), 4.52 (2H, t, $J=5.2\text{Hz}$), 5.46 (1H, q, $J=7.0\text{Hz}$), 7.60-7.68 (3H, m), 7.71-7.77 (2H, m), 8.20 (3H, brs).

Example 45

4-Acetyl-3-(2-aminoethoxy)-5-phenylisoxazole hydrochloride (Compound list No.: 1410)

(a) 4-Acetyl-3-(2-(N-tert-butoxycarbonylamino)ethoxy)-5-phenylisoxazole

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-(1-hydroxyethyl)-5-phenylisoxazole (0.3 g) was dissolved in methylene chloride (3 ml), and pyridinium dichromate (0.49 g) was added thereto at room temperature, followed by stirring of the resulting mixture at the same temperature for 24 hours. After the reaction, insolubles were filtered and the solvent of the filtrate was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (eluent: hexane/ethyl acetate = 2/1) to obtain the title compound (0.26 g, 87%) as a colorless powder.

IR spectrum (KBr) ν_{\max} cm^{-1} : 3359, 1685;

NMR spectrum (CDCl_3) δ ppm: 1.47 (9H, s), 2.51 (3H, s), 3.65 (2H, q, $J=5.2\text{Hz}$), 4.48 (2H, t, $J=5.2\text{Hz}$), 4.87 (1H, brs), 7.40-7.63 (3H, m), 7.93-7.96 (2H, m).

(b) 4-Acetyl-3-(2-aminoethoxy)-5-phenylisoxazole hydrochloride

4-Acetyl-3-(2-(N-tert-butoxycarbonylamino)ethoxy)-5-phenylisoxazole (0.2 g) was subjected to reaction and post-treatment in a similar manner to that described in Example 9(b) to obtain the title compound (0.12 g, 75%) as

colorless crystals.

m.p.: 150-152°C (decomposed);

IR spectrum (KBr) ν_{\max} cm^{-1} : 3006, 2909, 1682;

NMR spectrum (DMSO- d_6) δ ppm: 2.52 (3H, s), 3.35 (2H, t, $J=5.0\text{Hz}$), 4.59 (2H, t, $J=5.0\text{Hz}$), 7.54-7.64 (3H, m), 7.85-7.88 (2H, m), 8.30 (3H, brs).

Example 46

3-(2-Aminoethoxy)-4-isopropenyl-5-phenylisoxazole hydrochloride (Compound list No.: 1394)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-(1-hydroxyisopropyl)-5-phenylisoxazole

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-phenylisoxazole (0.8 g) was dissolved in tetrahydrofuran (16 ml), butyllithium (1.6M hexane solution, 3.7 ml) was added dropwise thereto at -70°C under a nitrogen atmosphere, and the resulting mixture was stirred at the same temperature for 10 minutes. After acetone (0.3 ml) was added dropwise to the reaction mixture and the mixture was stirred for 10 minutes, the temperature of the mixture was raised to 0°C . The reaction mixture was poured into ice-cold water and the pH of the mixture was adjusted to a value of 6 with an aqueous potassium dihydrogenphosphate solution. The mixture was extracted with ethyl acetate and the organic layer was washed with a saturated aqueous NaCl solution, followed by evaporation of the solvent under reduced pressure. The residue was purified by silica gel column chromatography (eluent: hexane/ethyl acetate = 3/1) to obtain the title compound (0.42 g, 44%) as a colorless oil.

IR spectrum (CHCl_3) ν_{\max} cm^{-1} : 3460, 2982, 2936, 1713;

NMR spectrum (CDCl_3) δ ppm: 1.46 (9H, s), 1.50 (6H, s), 2.52 (1H, brs), 3.60 (2H, q, $J=5.3\text{Hz}$), 4.41 (2H, t, $J=5.3\text{Hz}$), 4.87 (1H, brs), 7.41-7.49 (3H, m), 7.52-7.56 (2H, m).

(b) 3-(2-Aminoethoxy)-4-isopropenyl-5-phenylisoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-(1-hydroxyisopropyl)-5-

phenylisoxazole (0.13 g) was dissolved in a 4N hydrochloric acid/1,4-dioxane solution (10 ml), and the solution was stirred at 100°C for one hour. After the reaction, the reaction mixture was left to be cooled, followed by evaporation of the solvent under reduced pressure. The residue was washed with ethyl acetate to obtain the title compound (0.10 g, 83%) as colorless crystals.

m.p.: 130-133°C (decomposed);

IR spectrum (KBr) ν_{\max} cm^{-1} : 2974, 1591, 1572;

NMR spectrum (DMSO- d_6) δ ppm: 1.95 (3H, s), 3.28 (2H, t, $J=5.2\text{Hz}$), 4.48 (2H, t, $J=5.2\text{Hz}$), 5.23 (1H, s), 5.33 (1H, s), 7.53-7.58 (3H, m), 7.62-7.71 (2H, m), 8.20 (3H, brs).

Example 47

3-(2-Aminoethoxy)-5-(4-nitrophenyl)isoxazole hydrochloride (Compound list No.: 1660)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(4-nitrophenyl)isoxazole

3-Hydroxy-5-(4-nitrophenyl)isoxazole (0.64 g) and 2-(N-tert-butoxycarbonylamino)ethanol (0.55 g) were subjected to reaction and post-treatment in a similar manner to that described in Example 9(a) to obtain the title compound (0.86 g, 80%) as a colorless powder.

IR spectrum (KBr) ν_{\max} cm^{-1} : 3385, 1681;

NMR spectrum (CDCl₃) δ ppm: 1.46 (9H, s), 3.57 (2H, q, $J=5.2\text{Hz}$), 4.38 (2H, t, $J=5.2\text{Hz}$), 4.90 (1H, brs), 6.33 (1H, s), 7.90 (2H, d, $J=8.6\text{Hz}$), 8.33 (2H, d, $J=8.6\text{Hz}$).

(b) 3-(2-Aminoethoxy)-5-(4-nitrophenyl)isoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(4-nitrophenyl)isoxazole (0.28 g) was subjected to reaction and post-treatment in a similar manner to that described in Example 9(b) to obtain the title compound (0.12 g, 52%) as colorless crystals.

m.p.: 216-220°C (decomposed);

IR spectrum (KBr) ν_{\max} cm^{-1} : 2994, 2977, 2913, 1620, 1608;

NMR spectrum (DMSO- d_6) δ ppm: 3.28 (2H, t, $J=5.1\text{Hz}$), 4.47 (2H, t,

J=5.1Hz), 7.16 (1H, s), 8.15 (2H, d, J=8.9Hz), 8.22 (3H, brs), 8.39 (2H, d, J=8.9Hz).

Example 48

3-(2-Aminoethoxy)-5-(4-aminophenyl)isoxazole dihydrochloride (Compound list No.: 1702)

(a) 5-(4-Aminophenyl)-3-(2-(N-tert-butoxycarbonylamino)ethoxy)isoxazole

3-(2-(N-tert-butoxycarbonylamino)ethoxy)-5-(4-nitrophenyl)isoxazole (0.55 g) was dissolved in a mixture of acetic acid and water (9:1, 5.5 ml), and zinc powder (0.55 g) was added thereto at room temperature, followed by stirring of the resulting mixture at the same temperature for 2 hours. After the reaction, zinc powder was filtered off and the filtrate was extracted with ethyl acetate. The organic layer was washed with a saturated aqueous NaCl solution and dried over anhydrous magnesium sulfate. After filtration, the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (eluent: hexane/ethyl acetate = 2/1) to obtain the title compound (0.45 g, 90%) as a colorless powder.

IR spectrum (KBr) ν_{\max} cm^{-1} : 3391, 3378, 3300, 1712, 1615;

NMR spectrum (CDCl_3) δ ppm: 1.46 (9H, s), 3.55 (2H, q, J=5.1Hz), 4.00 (2H, brs), 4.33 (2H, t, J=5.1Hz), 4.96 (1H, brs), 5.94 (1H, s), 6.70 (2H, d, J=8.7Hz), 7.52 (2H, d, J=8.7Hz).

(b) 3-(2-Aminoethoxy)-5-(4-aminophenyl)isoxazole dihydrochloride

5-(4-Aminophenyl)-3-(2-(N-tert-butoxycarbonylamino)ethoxy)isoxazole (0.25 g) was subjected to reaction and post-treatment in a similar manner to that described in Example 9(b) to obtain the title compound (0.15 g, 65%) as colorless crystals.

m.p.: 242-248°C (decomposed);

IR spectrum (KBr) ν_{\max} cm^{-1} : 3039, 2939, 2584, 1625, 1597;

NMR spectrum (DMSO-d_6) δ ppm: 3.25 (2H, t, J=5.0Hz), 4.21 (2H, brs), 4.42 (2H, t, J=5.0Hz), 6.56 (1H, s), 6.94 (2H, d, J=8.3Hz), 7.65 (2H, d, J=8.3Hz), 8.25 (3H, brs).

Example 49

3-(2-Aminoethoxy)-5-(4-benzoylaminophenyl)isoxazole hydrochloride

(Compound list No.: 1716)

(a) 5-(4-Benzoylaminophenyl)-3-(2-(N-tert-butoxycarbonylamino)ethoxy)isoxazole

5-(4-Aminophenyl)-3-(2-(N-tert-butoxycarbonylamino)ethoxy)isoxazole (0.1 g) was dissolved in tetrahydrofuran (1 ml), and triethylamine (0.05 ml) and benzoyl chloride (0.04 ml) were successively added dropwise thereto at 5°C under a nitrogen atmosphere, followed by stirring of the resulting mixture at room temperature for 30 minutes. The reaction mixture was poured into ice-cold water and extracted with ethyl acetate. The organic layer was washed with a saturated aqueous NaCl solution and dried over anhydrous magnesium sulfate. After filtration, the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (eluent: hexane/ethyl acetate = 3/1) to obtain the title compound (0.08 g, 62%) as a colorless powder.

IR spectrum (KBr) ν_{\max} cm^{-1} : 3375, 1712;

NMR spectrum (CDCl_3) δ ppm: 1.46 (9H, s), 3.57 (2H, q, $J=5.1\text{Hz}$), 4.35 (2H, t, $J=5.1\text{Hz}$), 4.96 (1H, brs), 6.12 (1H, s), 7.52-7.61 (3H, m), 7.72-7.80 (4H, m), 7.87-7.95 (3H, m).

(b) 3-(2-Aminoethoxy)-5-(4-benzoylaminophenyl)isoxazole hydrochloride

5-(4-Benzoylaminophenyl)-3-(2-(N-tert-butoxycarbonylamino)ethoxy)-isoxazole (0.07 g) was subjected to reaction and post-treatment in a similar manner to that described in Example 9(b) to obtain the title compound (0.04 g, 68%) as colorless crystals.

m.p.: 242-248°C (decomposed);

IR spectrum (KBr) ν_{\max} cm^{-1} : 3322, 2959, 2910, 1645, 1621;

NMR spectrum (DMSO-d_6) δ ppm: 3.26 (2H, t, $J=5.1\text{Hz}$), 4.44 (2H, t, $J=5.1\text{Hz}$), 6.75 (1H, s), 7.53-7.65 (3H, m), 7.83-8.00 (6H, m), 8.20 (3H, brs), 10.55 (1H, s).

Example 50

3-(2-Aminoethoxy)-5-(2,4-dichloro-3-methylphenyl)isoxazole hydrochloride

(Compound list No.: 1576)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(2,4-dichloro-3-methylphenyl)isoxazole

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(2,4-dichlorophenyl)-isoxazole (0.3 g) and methyl iodide (0.08 ml) were subjected to reaction and post-treatment in a similar manner to that described in Example 14(a) to obtain the title compound (0.14 g, 45%) as a colorless powder.

IR spectrum (KBr) ν_{\max} cm^{-1} : 3342, 1710;

NMR spectrum (CDCl_3) δ ppm: 1.46 (9H, s), 2.56 (3H, s), 3.58 (2H, q, $J=5.1\text{Hz}$), 4.38 (2H, t, $J=5.1\text{Hz}$), 4.95 (1H, brs), 6.54 (1H, s), 7.39 (1H, d, $J=8.5\text{Hz}$), 7.76 (1H, d, $J=8.5\text{Hz}$).

(b) 3-(2-Aminoethoxy)-5-(2,4-dichloro-3-methylphenyl)isoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(2,4-dichloro-3-methylphenyl)isoxazole (0.13 g) was subjected to reaction and post-treatment in a similar manner to that described in Example 9(b) to obtain the title compound (0.06 g, 55%) as colorless crystals.

m.p.: 197-200°C (decomposed);

IR spectrum (KBr) ν_{\max} cm^{-1} : 3130, 2969, 2897, 1607;

NMR spectrum (DMSO-d_6) δ ppm: 2.52 (3H, s), 3.27 (2H, t, $J=5.1\text{Hz}$), 4.47 (2H, t, $J=5.1\text{Hz}$), 6.82 (1H, s), 7.65 (1H, d, $J=8.5\text{Hz}$), 7.71 (1H, d, $J=8.5\text{Hz}$), 8.17 (3H, brs).

Example 51

3-(2-Aminoethoxy)-5-(2,4-dichloro-3-ethylphenyl)isoxazole hydrochloride

(Compound list No.: 1590)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(2,4-dichloro-3-ethylphenyl)isoxazole

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(2,4-dichlorophenyl)isoxazole (0.3 g) and ethyl iodide (0.1 ml) were subjected to reaction and post-treatment in a similar manner to that described in Example 14(a) to obtain the title compound (0.25 g, 78%) as a colorless powder.

IR spectrum (KBr) ν_{\max} cm^{-1} : 3342, 1702;

NMR spectrum (CDCl_3) δ ppm: 1.21 (3H, t, $J=7.5\text{Hz}$), 1.46 (9H, s), 3.05 (2H, q, $J=7.5\text{Hz}$), 3.57 (2H, q, $J=5.1\text{Hz}$), 4.36 (2H, t, $J=5.1\text{Hz}$), 4.94 (1H, brs), 6.54 (1H, s), 7.40 (1H, d, $J=8.5\text{Hz}$), 7.65 (1H, d, $J=8.5\text{Hz}$).

(b) 3-(2-Aminoethoxy)-5-(2,4-dichloro-3-ethylphenyl)isoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(2,4-dichloro-3-ethylphenyl)isoxazole (0.23 g) was subjected to reaction and post-treatment in a similar manner to that described in Example 9(b) to obtain the title compound (0.14 g, 74%) as colorless crystals.

m.p.: 173-176°C (decomposed);

IR spectrum (KBr) ν_{\max} cm^{-1} : 2970, 2935, 2876, 1608;

NMR spectrum (DMSO-d_6) δ ppm: 1.16 (3H, t, $J=7.4\text{Hz}$), 3.00 (2H, q, $J=7.4\text{Hz}$), 3.27 (2H, t, $J=5.1\text{Hz}$), 4.47 (2H, t, $J=5.1\text{Hz}$), 6.82 (1H, s), 7.65 (1H, d, $J=8.4\text{Hz}$), 7.70 (1H, d, $J=8.4\text{Hz}$), 8.21 (3H, brs).

Example 52

5-(4-Acetoxyphenyl)-3-(2-aminoethoxy)isoxazole hydrochloride (Compound list No.: 1688)(a) 5-(4-Acetoxyphenyl)-3-(2-(N-tert-butoxycarbonylamino)ethoxy)isoxazole

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(4-hydroxyphenyl)-isoxazole (0.15 g) was dissolved in tetrahydrofuran (1.5 ml), and triethylamine

(0.07 ml) and acetyl chloride (0.04 ml) were added dropwise thereto at 5°C under a nitrogen atmosphere, followed by stirring of the resulting mixture at room temperature for 30 minutes. The reaction mixture was poured into ice-cold water and extracted with ethyl acetate. The organic layer was washed with a saturated aqueous NaCl solution and dried over anhydrous magnesium sulfate. After filtration, the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (eluent: hexane/ethyl acetate = 2/1) to obtain the title compound (0.12 g, 71%) as a colorless powder.

IR spectrum (KBr) ν_{\max} cm^{-1} : 3345, 1696;

NMR spectrum (CDCl_3) δ ppm: 1.46 (9H, s), 2.33 (3H, s), 3.56 (2H, q, $J=5.2\text{Hz}$), 4.35 (2H, t, $J=5.2\text{Hz}$), 4.94 (1H, brs), 6.12 (1H, s), 7.20 (2H, d, $J=8.8\text{Hz}$), 7.74 (2H, d, $J=8.8\text{Hz}$).

(b) 5-(4-Acetoxyphenyl)-3-(2-aminoethoxy)isoxazole hydrochloride

5-(4-Acetoxyphenyl)-3-(2-(N-tert-butoxycarbonylamino)ethoxy)-isoxazole (0.1 g) was subjected to reaction and post-treatment in a similar manner to that described in Example 9(b) to obtain the title compound (0.05 g, 63%) as colorless crystals.

m.p.: 202-212°C (decomposed);

IR spectrum (KBr) ν_{\max} cm^{-1} : 2996, 2977, 2912, 1755, 1621;

NMR spectrum (DMSO-d_6) δ ppm: 2.30 (3H, s), 3.26 (2H, t, $J=5.1\text{Hz}$), 4.45 (2H, t, $J=5.1\text{Hz}$), 6.84 (1H, s), 7.32 (2H, d, $J=8.6\text{Hz}$), 7.91 (2H, d, $J=8.6\text{Hz}$), 8.22 (3H, brs).

Example 53

3-(2-Aminoethoxy)-5-(4-benzyloxyphenyl)isoxazole hydrochloride (Compound list No.: 1646)

(a) 5-(4-Benzyloxyphenyl)-3-(2-(N-tert-butoxycarbonylamino)ethoxy)isoxazole

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(4-hydroxyphenyl)-isoxazole (0.2 g) was dissolved in dimethylformamide (2 ml), and 55% sodium hydride (oil, 0.03 g) was added thereto at 5°C under a nitrogen atmosphere,

followed by stirring of the resulting mixture at the same temperature for 10 minutes. Then, benzyl bromide (0.08 ml) was added dropwise thereto and the mixture was stirred at room temperature for 30 minutes. The reaction mixture was poured into ice-cold water and extracted with ethyl acetate. The organic layer was washed with a saturated aqueous NaCl solution and dried over anhydrous magnesium sulfate. After filtration, the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (eluent: hexane/ethyl acetate = 3/1) to obtain the title compound (0.14 g, 54%) as a colorless powder.

IR spectrum (KBr) ν_{\max} cm^{-1} : 3338, 1698;

NMR spectrum (CDCl_3) δ ppm: 1.46 (9H, s), 3.56 (2H, q, $J=5.0\text{Hz}$), 4.34 (2H, t, $J=5.0\text{Hz}$), 4.94 (1H, brs), 5.12 (2H, s), 6.02 (1H, s), 7.04 (2H, d, $J=8.7\text{Hz}$), 7.28-7.47 (5H, m), 7.66 (2H, d, $J=8.7\text{Hz}$).

(b) 3-(2-Aminoethoxy)-5-(4-benzyloxyphenyl)isoxazole hydrochloride

5-(4-Benzyloxyphenyl)-3-(2-(N-tert-butoxycarbonylamino)ethoxy)-isoxazole (0.13 g) was subjected to reaction and post-treatment in a similar manner to that described in Example 9(b) to obtain the title compound (0.07 g, 64%) as colorless crystals.

m.p.: 205-210°C (decomposed);

IR spectrum (KBr) ν_{\max} cm^{-1} : 2997, 2966, 2912, 1621, 1604;

NMR spectrum (DMSO-d_6) δ ppm: 3.25 (2H, t, $J=5.1\text{Hz}$), 4.43 (2H, t, $J=5.1\text{Hz}$), 5.19 (2H, s), 6.69 (1H, s), 7.17 (2H, d, $J=9.0\text{Hz}$), 7.33-7.48 (5H, m), 7.79 (2H, d, $J=9.0\text{Hz}$), 8.21 (3H, brs).

Example 54

3-(2-Aminoethoxy)-5-(2-furyl)-4-isopropylisoxazole hydrochloride (Compound list No.: 510)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(2-furyl)-4-isopropylisoxazole

5-(2-Furyl)-3-hydroxy-4-isopropylisoxazole (0.2 g) and 2-(N-tert-butoxycarbonylamino)ethanol (0.18 g) were subjected to reaction and post-

treatment in a similar manner to that described in Example 9(a) to obtain the title compound (0.29 g, 83%) as a colorless powder.

IR spectrum (KBr) ν_{\max} cm^{-1} : 3317, 2985, 1690;

NMR spectrum (CDCl_3) δ ppm: 1.28 (6H, d, $J=7.2\text{Hz}$), 1.46 (9H, s), 3.33 (1H, qq, $J=7.2\text{Hz}$, $J=7.2\text{Hz}$), 3.58 (2H, q, $J=5.1\text{Hz}$), 4.35 (2H, t, $J=5.1\text{Hz}$), 4.84 (1H, brs), 6.52 (1H, dd, $J=3.4\text{Hz}$, $J=1.8\text{Hz}$), 6.80 (1H, d, $J=3.4\text{Hz}$), 7.55 (1H, d, $J=1.8\text{Hz}$).

(b) 3-(2-Aminoethoxy)-5-(2-furyl)-4-isopropylisoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(2-furyl)-4-isopropylisoxazole (0.27 g) was subjected to reaction and post-treatment in a similar manner to that described in Example 1(b) to obtain the title compound (0.17 g, 77%) as colorless crystals.

m.p.: 137-139°C;

IR spectrum (KBr) ν_{\max} cm^{-1} : 2972, 2898, 1560, 1513;

NMR spectrum (DMSO-d_6) δ ppm: 1.26 (6H, d, $J=7.0\text{Hz}$), 3.24 (1H, qq, $J=7.0\text{Hz}$, $J=7.0\text{Hz}$), 3.28 (2H, t, $J=5.2\text{Hz}$), 4.44 (2H, t, $J=5.2\text{Hz}$), 6.73 (1H, dd, $J=3.4\text{Hz}$, $J=1.8\text{Hz}$), 7.01 (1H, d, $J=3.4\text{Hz}$), 7.97 (1H, d, $J=1.8\text{Hz}$), 8.21 (3H, brs).

Example 55

3-(2-Aminoethoxy)-4-(tert-butyl)-5-phenylisoxazole hydrochloride (Compound list No.: 13)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-(tert-butyl)-5-phenylisoxazole

4-(tert-Butyl)-3-hydroxy-5-phenylisoxazole (0.15 g) and 2-(N-tert-butoxycarbonylamino)ethanol (0.12 g) were subjected to reaction and post-treatment in a similar manner to that described in Example 9(a) to obtain the title compound (0.16 g, 64%) as a colorless powder.

IR spectrum (KBr) ν_{\max} cm^{-1} : 3374, 2974, 1683;

NMR spectrum (CDCl_3) δ ppm: 1.18 (9H, s), 1.46 (9H, s), 3.60 (2H, q, $J=5.2\text{Hz}$), 4.38 (2H, t, $J=5.2\text{Hz}$), 4.85 (1H, brs), 7.36-7.55 (5H, m).

(b) 3-(2-Aminoethoxy)-4-(tert-butyl)-5-phenylisoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-(tert-butyl)-5-phenylisoxazole (0.14 g) was subjected to reaction and post-treatment in a similar manner to that described in Example 1(b) to obtain the title compound (0.09 g, 82%) as colorless crystals.

m.p.: 230-234°C (decomposed);

IR spectrum (KBr) ν_{\max} cm^{-1} : 2961, 2910, 2890, 1516;

NMR spectrum (DMSO- d_6) δ ppm: 1.16 (9H, s), 3.30 (2H, t, J=5.3Hz), 4.46 (2H, t, J=5.3Hz), 7.44-7.57 (5H, m), 8.20 (3H, brs).

Example 56

3-(2-Aminoethoxy)-4-cyclopropyl-5-phenylisoxazole hydrochloride

(Compound list No.: 1400)

(a) 3-(2-(N-tert-butoxycarbonylamino)ethoxy)-4-cyclopropyl-5-phenylisoxazole

4-Cyclopropyl-3-hydroxy-5-phenylisoxazole (0.2 g) and 2-(N-tert-butoxycarbonylamino)ethanol (0.18 g) were subjected to reaction and post-treatment in a similar manner to that described in Example 9(a) to obtain the title compound (0.25 g, 74%) as a colorless powder.

IR spectrum (KBr) ν_{\max} cm^{-1} : 3369, 1692, 1536, 1520;

NMR spectrum (CDCl₃) δ ppm: 0.66-0.75 (2H, m), 0.88-0.95 (2H, m), 1.46 (9H, s), 1.67-1.73 (1H, m), 3.58 (2H, q, J=5.1Hz), 4.35 (2H, t, J=5.1Hz), 4.90 (1H, brs), 7.40-7.51 (3H, m), 7.84-7.87 (2H, m).

(b) 3-(2-Aminoethoxy)-4-cyclopropyl-5-phenylisoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-cyclopropyl-5-phenylisoxazole (0.15 g) was subjected to reaction and post-treatment in a similar manner to that described in Example 1(b) to obtain the title compound (0.06 g, 49%) as colorless crystals.

m.p.: 180-182°C;

IR spectrum (KBr) ν_{\max} cm^{-1} : 2919, 2851, 1713, 1519;

NMR spectrum (DMSO- d_6) δ ppm: 0.68-0.76 (2H, m), 0.83-0.92 (2H, m), 1.71-

1.78 (1H, m), 3.27 (2H, t, J=5.2Hz), 4.44 (2H, t, J=5.2Hz), 7.51-7.60 (3H, m), 7.83-7.86 (2H, m), 8.20 (3H, brs).

Example 57

3-(2-Aminoethoxy)-5-(2,4-dichlorophenyl)-4-isopropylisoxazole hydrochloride

(Compound list No.: 176)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(2,4-dichlorophenyl)-4-isopropylisoxazole

5-(2,4-Dichlorophenyl)-3-hydroxy-4-isopropylisoxazole (0.06 g) and 2-(N-tert-butoxycarbonylamino)ethanol (0.04 g) were subjected to reaction and post-treatment in a similar manner to that described in Example 9(a) to obtain the title compound (0.078 g, 86%) as a colorless oil.

IR spectrum (CHCl₃) ν_{\max} cm⁻¹: 3460, 2975, 2936, 1713;

NMR spectrum (CDCl₃) δ ppm: 1.18 (6H, d, J=7.0Hz), 1.46 (9H, s), 2.69 (1H, qq, J=7.0Hz, J=7.0Hz), 3.59 (2H, q, J=5.2Hz), 4.38 (2H, t, J=5.2Hz), 4.85 (1H, brs), 7.30 (1H, d, J=8.3Hz), 7.35 (1H, dd, J=8.3Hz, J=1.9Hz), 7.52 (1H, d, J=1.9Hz).

(b) 3-(2-Aminoethoxy)-5-(2,4-dichlorophenyl)-4-isopropylisoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(2,4-dichlorophenyl)-4-isopropylisoxazole (0.07 g) was subjected to reaction and post-treatment in a similar manner to that described in Example 1(b) to obtain the title compound (0.04 g, 68%) as colorless crystals.

m.p.: 171-173°C;

IR spectrum (KBr) ν_{\max} cm⁻¹: 2968, 2934, 2875, 1514;

NMR spectrum (DMSO-d₆) δ ppm: 1.13 (6H, d, J=7.0Hz), 2.69 (1H, qq, J=7.0Hz, J=7.0Hz), 3.29 (2H, t, J=5.3Hz), 4.46 (2H, t, J=5.3Hz), 7.56 (1H, d, J=8.5Hz), 7.62 (1H, dd, J=8.5Hz, J=2.0Hz), 7.89 (1H, d, J=2.0Hz), 8.18 (3H, brs).

Example 58

3-(2-Aminoethoxy)-5-(2-chlorophenyl)-4-isopropylisoxazole hydrochloride

(Compound list No.: 117)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(2-chlorophenyl)-4-isopropylisoxazole

5-(2-Chlorophenyl)-3-hydroxy-4-isopropylisoxazole (0.13 g) and 2-(N-tert-butoxycarbonylamino)ethanol (0.1 g) were subjected to reaction and post-treatment in a similar manner to that described in Example 9(a) to obtain the title compound (0.15 g, 72%) as a colorless powder.

IR spectrum (KBr) ν_{\max} cm^{-1} : 3382, 2974, 1691;

NMR spectrum (CDCl_3) δ ppm: 1.18 (6H, d, $J=6.9\text{Hz}$), 1.46 (9H, s), 2.71 (1H, qq, $J=6.9\text{Hz}$, $J=6.9\text{Hz}$), 3.60 (2H, q, $J=5.2\text{Hz}$), 4.39 (2H, t, $J=5.2\text{Hz}$), 4.87 (1H, brs), 7.34-7.51 (4H, m).

(b) 3-(2-Aminoethoxy)-5-(2-chlorophenyl)-4-isopropylisoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(2-chlorophenyl)-4-isopropylisoxazole (0.13 g) was subjected to reaction and post-treatment in a similar manner to that described in Example 1(b) to obtain the title compound (0.08 g, 73%) as colorless crystals.

m.p.: 143-145°C;

IR spectrum (KBr) ν_{\max} cm^{-1} : 2970, 2937, 2896, 2879, 1513;

NMR spectrum (DMSO-d_6) δ ppm: 1.13 (6H, d, $J=7.0\text{Hz}$), 2.69 (1H, qq, $J=7.0\text{Hz}$, $J=7.0\text{Hz}$), 3.29 (2H, t, $J=5.2\text{Hz}$), 4.47 (2H, t, $J=5.2\text{Hz}$), 7.50-7.68 (4H, m), 8.24 (3H, brs).

Example 59

3-(2-Aminoethoxy)-5-(4-chlorophenyl)-4-isopropylisoxazole hydrochloride

(Compound list No.: 151)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(4-chlorophenyl)-4-isopropylisoxazole

5-(4-Chlorophenyl)-3-hydroxy-4-isopropylisoxazole (0.15 g) and 2-(N-

tert-butoxycarbonylamino)ethanol (0.11 g) were subjected to reaction and post-treatment in a similar manner to that described in Example 9(a) to obtain the title compound (0.19 g, 79%) as a colorless powder.

IR spectrum (KBr) ν_{\max} cm^{-1} : 3382, 2966, 1685;

NMR spectrum (CDCl_3) δ ppm: 1.29 (6H, d, $J=7.0\text{Hz}$), 1.46 (9H, s), 3.03 (1H, qq, $J=7.0\text{Hz}$, $J=7.0\text{Hz}$), 3.59 (2H, q, $J=5.1\text{Hz}$), 4.36 (2H, t, $J=5.1\text{Hz}$), 7.44 (2H, d, $J=8.7\text{Hz}$), 7.52 (2H, d, $J=8.7\text{Hz}$).

(b) 3-(2-Aminoethoxy)-5-(4-chlorophenyl)-4-isopropylisoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(4-chlorophenyl)-4-isopropylisoxazole (0.18 g) was subjected to reaction and post-treatment in a similar manner to that described in Example 1(b) to obtain the title compound (0.12 g, 80%) as colorless crystals.

m.p.: 224-227°C;

IR spectrum (KBr) ν_{\max} cm^{-1} : 2971, 2919, 2851, 1639;

NMR spectrum (DMSO-d_6) δ ppm: 1.26 (6H, d, $J=7.0\text{Hz}$), 3.02 (1H, qq, $J=7.0\text{Hz}$, $J=7.0\text{Hz}$), 3.28 (2H, t, $J=5.3\text{Hz}$), 4.45 (2H, t, $J=5.3\text{Hz}$), 7.62 (2H, d, $J=9.0\text{Hz}$), 7.63 (2H, d, $J=9.0\text{Hz}$), 8.18 (3H, brs).

Example 60

4-Allyl-3-(2-aminoethoxy)-5-phenylisoxazole hydrochloride (Compound list No.: 1392)

(a) 4-Allyl-3-(2-(N-tert-butoxycarbonylamino)ethoxy)-5-phenylisoxazole

4-Allyl-3-hydroxy-5-phenylisoxazole (1.00 g) and 2-(N-tert-butoxycarbonylamino)ethanol (0.96 g) were subjected to reaction and post-treatment in a similar manner to that described in Example 1(a) to obtain the title compound (1.57 g, 91%) as colorless crystals.

m.p.: 78-79°C;

IR spectrum (KBr) ν_{\max} cm^{-1} : 3327, 1708, 1644, 1526, 1518;

NMR spectrum (CDCl_3) δ ppm: 1.46 (9H, s), 3.27-3.29 (2H, m), 3.57 (2H, q, $J=5.1\text{Hz}$), 4.38 (2H, t, $J=5.1\text{Hz}$), 4.90 (1H, brs), 5.05-5.15 (2H, m), 5.91-6.01 (1H, m),

7.41-7.49 (3H, m), 7.63-7.68 (2H, m).

(b) 4-Allyl-3-(2-aminoethoxy)-5-phenylisoxazole hydrochloride

4-Allyl-3-(2-(N-tert-butoxycarbonylamino)ethoxy)-5-phenylisoxazole (200 mg) was subjected to reaction and post-treatment in a similar manner to that described in Example 1(b) to obtain the title compound (160 mg, 98%) as colorless crystals.

m.p.: 95-96°C;

IR spectrum (KBr) ν_{\max} cm^{-1} : 2967, 2912, 2885, 2817, 2694, 1643, 1601, 1577, 1514, 1495;

NMR spectrum (DMSO- d_6) δ ppm: 3.25-3.30 (2H, m), 3.35 (2H, t, J=5.1Hz), 4.46 (2H, t, J=5.1Hz), 5.03-5.09 (2H, m), 5.91-6.00 (1H, m), 7.52-7.58 (3H, m), 7.67-7.70 (2H, m), 8.20 (3H, brs).

Example 61

3-(2-Aminoethoxy)-5-phenyl-4-propargylisoxazole hydrochloride (Compound list No.: 1398)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-phenyl-4-propargylisoxazole

3-Hydroxy-5-phenyl-4-propargylisoxazole (1.00 g) and 2-(N-tert-butoxycarbonylamino)ethanol (0.96 g) were subjected to reaction and post-treatment in a similar manner to that described in Example 1(a) to obtain the title compound (1.44 g, 84%) as a colorless oil.

IR spectrum (KBr) ν_{\max} cm^{-1} : 3352, 3302, 1711, 1643, 1600, 1577, 1520, 1499;

NMR spectrum (CDCl₃) δ ppm: 1.46 (9H, s), 2.09 (1H, t, J=2.8Hz), 3.43 (2H, d, J=2.8Hz), 3.60 (2H, q, J=5.1Hz), 4.41 (2H, t, J=5.1Hz), 5.02 (1H, brs), 7.45-7.55 (3H, m), 7.74-7.76 (2H, m).

(b) 3-(2-Aminoethoxy)-5-phenyl-4-propargylisoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-phenyl-4-propargylisoxazole (200 mg) was subjected to reaction and post-treatment in a similar

manner to that described in Example 1(b) to obtain the title compound (152 mg, 94%) as colorless crystals.

m.p.: 210-212°C (decomposed);

IR spectrum (KBr) ν_{\max} cm^{-1} : 3258, 2959, 2899, 2830, 1632, 1601, 1577, 1523;

NMR spectrum (DMSO- d_6) δ ppm: 2.97 (1H, t, $J=2.8\text{Hz}$), 3.30 (1H, t, $J=5.1\text{Hz}$), 3.59 (2H, d, $J=2.8\text{Hz}$), 4.47 (2H, t, $J=5.1\text{Hz}$), 7.56-7.62 (3H, m), 7.79-7.82 (2H, m), 8.13 (3H, brs).

Example 62

3-(2-Aminoethoxy)-4-isobutyl-5-phenylisoxazole hydrochloride (Compound list No.: 11)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-isobutyl-5-phenylisoxazole

3-Hydroxy-4-isobutyl-5-phenylisoxazole (217 mg) and 2-(N-tert-butoxycarbonylamino)ethanol (193 mg) were subjected to reaction and post-treatment in a similar manner to that described in Example 1(a) to obtain the title compound (303 mg, 84%) as colorless crystals.

m.p.: 80-81°C;

IR spectrum (KBr) ν_{\max} cm^{-1} : 3377, 1683, 1637, 1516, 1498;

NMR spectrum (CDCl_3) δ ppm: 0.93 (6H, d, $J=6.8\text{Hz}$), 1.46 (9H, s), 1.89-1.96 (1H, m), 2.42 (2H, d, $J=7.3\text{Hz}$), 3.58 (2H, q, $J=5.1\text{Hz}$), 4.37 (2H, t, $J=5.1\text{Hz}$), 4.88 (1H, brs), 7.40-7.49 (3H, m), 7.66-7.70 (2H, m).

(b) 3-(2-Aminoethoxy)-4-isobutyl-5-phenylisoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-isobutyl-5-phenylisoxazole (200 mg) was subjected to reaction and post-treatment in a similar manner to that described in Example 1(b) to obtain the title compound (160 mg, 98%) as colorless crystals.

m.p.: 202-210°C (decomposed);

IR spectrum (KBr) ν_{\max} cm^{-1} : 3005, 2957, 2869, 1629, 1601, 1576, 1514, 1495;

NMR spectrum (DMSO-d₆) δ ppm: 0.86 (6H, d, J=6.8Hz), 2.47 (2H, d, J=7.3Hz), 3.27 (2H, t, J=5.1Hz), 4.45 (2H, t, J=5.1Hz), 7.49-7.58 (3H, m), 7.70-7.74 (2H, m), 8.19 (3H, brs).

Example 63

3-(2-Aminoethoxy)-4-cyclopentyl-5-phenylisoxazole hydrochloride (Compound list No.: 1402)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-cyclopentyl-5-phenylisoxazole

3-Hydroxy-4-cyclopentyl-5-phenylisoxazole (229 mg) and 2-(N-tert-butoxycarbonylamino)ethanol (193 mg) were subjected to reaction and post-treatment in a similar manner to that described in Example 1(a) to obtain the title compound (308 mg, 83%) as colorless crystals.

m.p.: 112-113°C;

IR spectrum (KBr) ν_{\max} cm⁻¹: 3391, 1691, 1655, 1643, 1531, 1515;

NMR spectrum (CDCl₃) δ ppm: 1.46 (9H, s), 1.56-1.67 (2H, m), 1.71-1.92 (6H, m), 3.03-3.12 (1H, m), 3.58 (2H, q, J=5.1Hz), 4.38 (2H, t, J=5.1Hz), 4.82 (1H, brs), 7.41-7.49 (3H, m), 7.55-7.61 (2H, m).

(b) 3-(2-Aminoethoxy)-4-cyclopentyl-5-phenylisoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-cyclopentyl-5-phenylisoxazole (200 mg) was subjected to reaction and post-treatment in a similar manner to that described in Example 1(b) to obtain the title compound (161 mg, 97%) as colorless crystals.

m.p.: 197-199°C;

IR spectrum (KBr) ν_{\max} cm⁻¹: 2954, 2869, 1637, 1599, 1576, 1511, 1490;

NMR spectrum (DMSO-d₆) δ ppm: 1.57-1.63 (2H, m), 1.72-1.91 (6H, m), 2.98-3.07 (1H, m), 3.27 (2H, t, J=5.1Hz), 4.45 (2H, t, J=5.1Hz), 7.51-7.61 (5H, m), 8.07 (3H, brs).

Example 64

3-(2-Aminoethoxy)-4-(2-cyclopentenyl)-5-phenylisoxazole hydrochloride

(Compound list No.: 1404)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-(2-cyclopentenyl)-5-phenylisoxazole

3-Hydroxy-4-(2-cyclopentenyl)-5-phenylisoxazole (227 mg) and 2-(N-tert-butoxycarbonylamino)ethanol (193 mg) were subjected to reaction and post-treatment in a similar manner to that described in Example 1(a) to obtain the title compound (295 mg, 80%) as colorless crystals.

m.p.: 81-82°C;

IR spectrum (KBr) ν_{\max} cm^{-1} : 3394, 1690, 1636, 1515, 1495;

NMR spectrum (CDCl_3) δ ppm: 1.46 (9H, s), 1.88-1.98 (1H, m), 2.24-2.34 (1H, m), 2.39-2.49 (1H, m), 2.52-2.61 (1H, m), 3.47-3.59 (2H, m), 4.01-4.06 (1H, m), 4.30-4.36 (2H, m), 4.86 (1H, brs), 5.63-5.67 (1H, m), 5.89-5.92 (1H, m), 7.42-7.49 (3H, m), 7.60-7.63 (2H, m).

(b) 3-(2-Aminoethoxy)-4-(2-cyclopentenyl)-5-phenylisoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-(2-cyclopentenyl)-5-phenylisoxazole (200 mg) was subjected to reaction and post-treatment in a similar manner to that described in Example 1(b) to obtain the title compound (156 mg, 94%) as colorless crystals.

m.p.: 157-160°C (decomposed);

IR spectrum (KBr) ν_{\max} cm^{-1} : 3051, 2954, 2901, 2852, 1635, 1599, 1575, 1516, 1492;

NMR spectrum (DMSO-d_6) δ ppm: 1.79-1.99 (1H, m), 2.21-2.41 (2H, m), 2.44-2.53 (1H, m), 3.25 (2H, brs), 3.95-4.02 (1H, m), 4.38-4.48 (2H, m), 5.56-5.71 (1H, m), 5.85-5.89 (1H, m), 7.51-7.57 (3H, m), 7.59-7.63 (2H, m), 8.12 (3H, brs).

Example 65

3-(2-Aminoethoxy)-4-pentyl-5-phenylisoxazole hydrochloride (Compound list No.: 1386)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-pentyl-5-phenylisoxazole

3-Hydroxy-4-pentyl-5-phenylisoxazole (231 mg) and 2-(N-tert-butoxycarbonylamino)ethanol (193 mg) were subjected to reaction and post-treatment in a similar manner to that described in Example 1(a) to obtain the title compound (305 mg, 82%) as a colorless oil.

IR spectrum (KBr) ν_{\max} cm^{-1} : 3461, 1713, 1640, 1510, 1496;

NMR spectrum (CDCl_3) δ ppm: 0.89 (3H, t, $J=7.0\text{Hz}$), 1.26-1.38 (2H, m), 1.46 (9H, s), 1.49-1.61 (2H, m), 2.52 (2H, t, $J=7.7\text{Hz}$), 3.59 (2H, q, $J=5.1\text{Hz}$), 4.37 (2H, t, $J=5.1\text{Hz}$), 4.89 (1H, brs), 7.40-7.49 (3H, m), 7.64-7.69 (2H, m).

(b) 3-(2-Aminoethoxy)-4-pentyl-5-phenylisoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-pentyl-5-phenylisoxazole (100 mg) was subjected to reaction and post-treatment in a similar manner to that described in Example 1(b) to obtain the title compound (80 mg, 95%) as colorless crystals.

m.p.: 107-109°C;

IR spectrum (KBr) ν_{\max} cm^{-1} : 3008, 2955, 2930, 2869, 1641, 1601, 1576, 1566, 1516, 1496;

NMR spectrum (DMSO-d_6) δ ppm: 0.83 (3H, t, $J=7.0\text{Hz}$), 1.25-1.30 (4H, m), 1.48-1.56 (2H, m), 2.55 (2H, t, $J=7.7\text{Hz}$), 3.28 (2H, t, $J=5.1\text{Hz}$), 4.45 (2H, t, $J=5.1\text{Hz}$), 7.51-7.59 (3H, m), 7.66-7.69 (2H, m), 8.10 (3H, brs).

Example 66

3-(2-Aminoethoxy)-4-(2-butenyl)-5-phenylisoxazole hydrochloride (Compound list No.: 1396)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-(2-butenyl)-5-phenylisoxazole

3-Hydroxy-4-(2-butenyl)-5-phenylisoxazole (215 mg) and 2-(N-tert-butoxycarbonylamino)ethanol (193 mg) were subjected to reaction and post-

treatment in a similar manner to that described in Example 1(a) to obtain the title compound (292 mg, 82%) as colorless crystals.

m.p.: 62-63°C;

IR spectrum (KBr) ν_{\max} cm^{-1} : 3378, 1682, 1634, 1514, 1497;

NMR spectrum (CDCl_3) δ ppm: 1.46 (9H, s), 1.68-1.74 (3H, m), 3.19-3.21 (2H, m), 3.57 (2H, q, $J=5.1\text{Hz}$), 4.37 (2H, t, $J=5.1\text{Hz}$), 4.89 (1H, brs), 5.46-5.60 (2H, m), 7.40-7.52 (3H, m), 7.64-7.68 (2H, m).

(b) 3-(2-Aminoethoxy)-4-(2-butenyl)-5-phenylisoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-(2-butenyl)-5-phenylisoxazole (200 mg) was subjected to reaction and post-treatment in a similar manner to that described in Example 1(b) to obtain the title compound (162 mg, 98%) as colorless crystals.

m.p.: 123-125°C;

IR spectrum (KBr) ν_{\max} cm^{-1} : 3028, 2961, 2934, 2916, 2855, 1641, 1601, 1577, 1570, 1515, 1494;

NMR spectrum (DMSO-d_6) δ ppm: 1.60-1.70 (3H, m), 3.26-3.28 (2H, m), 3.33 (2H, t, $J=5.1\text{Hz}$), 4.45 (2H, t, $J=5.1\text{Hz}$), 5.43-5.59 (2H, m), 7.50-7.58 (3H, m), 7.65-7.70 (2H, m), 8.10 (3H, brs).

Example 67

3-(2-Aminoethoxy)-4-isopropyl-5-(2-thienyl)isoxazole hydrochloride

(Compound list No.: 543)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-isopropyl-5-(2-thienyl)isoxazole

3-Hydroxy-4-isopropyl-5-(2-thienyl)isoxazole (209 mg) and 2-(N-tert-butoxycarbonylamino)ethanol (193 mg) were subjected to reaction and post-treatment in a similar manner to that described in Example 1(a) to obtain the title compound (301 mg, 86%) as colorless crystals.

m.p.: 94-95°C;

IR spectrum (KBr) ν_{\max} cm^{-1} : 3325, 1711, 1693, 1635, 1542, 1528, 1508;

NMR spectrum (CDCl₃) δ ppm: 1.29 (6H, d, J=7.1Hz), 1.46 (9H, s), 3.19 (1H, m), 3.58 (2H, q, J=5.1Hz), 4.36 (2H, t, J=5.1Hz), 4.83 (1H, brs), 7.14 (1H, dd, J=5.3Hz, J=3.7Hz), 7.42 (1H, dd, J=3.4Hz, J=1.4Hz), 7.46 (1H, d, J=5.3Hz).

(b) 3-(2-Aminoethoxy)-4-isopropyl-5-(2-thienyl)isoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-isopropyl-5-(2-thienyl)-isoxazole (200 mg) was subjected to reaction and post-treatment in a similar manner to that described in Example 1(b) to obtain the title compound (150 mg, 93%) as colorless crystals.

m.p.: 192-194°C (decomposed);

IR spectrum (KBr) ν_{\max} cm⁻¹: 3105, 3087, 2973, 1644, 1579, 1527, 1498;

NMR spectrum (DMSO-d₆) δ ppm: 1.27 (6H, d, J=7.2Hz), 3.16 (1H, m), 3.28 (2H, t, J=5.1Hz), 4.44 (2H, t, J=5.1Hz), 7.26 (1H, dd, J=5.2Hz, J=1.4Hz), 7.54 (1H, dd, J=4.7Hz, J=1.0Hz), 7.87 (1H, d, J=5.2Hz), 8.13 (3H, brs).

Example 68

3-(2-Aminoethoxy)-4-fluoro-5-phenylisoxazole hydrochloride (Compound list No.: 4)

(a) 3-Methoxymethoxy-5-phenylisoxazole

3-Hydroxy-5-phenylisoxazole (8.05 g) was dissolved in dimethylformamide (80 ml), and sodium methoxide (28% methanol solution, 3.24 g) was added dropwise thereto, followed by stirring of the resulting mixture at room temperature for one hour. While the reaction mixture was cooled to 5°C, chloromethyl methyl ether (4.83 g) was added thereto, followed by stirring of the resulting mixture at the same temperature for one hour. The reaction mixture was poured into ice-cold water (200 ml) and extracted with ether (200 ml x 2), and the organic layer was dried over anhydrous magnesium sulfate. After filtration, the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (eluent: hexane/ethyl acetate = 4/1) to obtain the title compound (6.30 g, 61%) as a colorless oil.

IR spectrum (KBr) ν_{\max} cm^{-1} : 1621, 1596, 1577, 1512;

NMR spectrum (CDCl_3) δ ppm: 3.58 (3H, s), 5.37 (2H, s), 7.40-7.48 (3H, m), 7.71-7.95 (2H, m).

(b) 4-Fluoro-3-methoxymethoxy-5-phenylisoxazole

3-Methoxymethoxy-5-phenylisoxazole (2.05 g) was dissolved in anhydrous tetrahydrofuran (30 ml), and the solution was cooled to -78°C . Butyllithium (1.68M hexane solution, 7.1 ml) was added dropwise thereto, and the resulting mixture was stirred at the same temperature for 15 minutes. Then, N-fluorobenzenesulfonimide (3.15 g) was added to the reaction mixture, followed by stirring of the resulting mixture at the same temperature for 15 minutes. The cooling bath was removed and the temperature of the resulting mixture was raised to room temperature. The reaction mixture was poured into ice-cold water (200 ml) and extracted with ether (200 ml x 2). The organic layer was dried over anhydrous magnesium sulfate. After filtration, the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (eluent: hexane/ethyl acetate = 4/1) to obtain the title compound (1.78 g, 80%) as a colorless oil.

IR spectrum (KBr) ν_{\max} cm^{-1} : 1672, 1545;

NMR spectrum (CDCl_3) δ ppm: 3.62 (3H, s), 5.45 (2H, s), 7.43-7.52 (3H, m), 7.70-7.80 (2H, m).

(c) 4-Fluoro-3-hydroxy-5-phenylisoxazole

4-Fluoro-3-methoxymethoxy-5-phenylisoxazole (0.44 g) was dissolved in a solution of 4N hydrochloric acid/dioxane (5.0 ml), and the solution was stirred at room temperature for one hour. After the reaction, the solvent was evaporated under reduced pressure and the crystal thus obtained was washed with dichloromethane (10 ml) to obtain the title compound (0.30 g, 83%) as colorless crystals.

IR spectrum (KBr) ν_{\max} cm^{-1} : 3059, 3014, 2995, 2919, 2851, 2822, 2747, 2645, 2565, 1674, 1582, 1527;

NMR spectrum (DMSO- d_6) δ ppm: 7.51-7.61 (3H, m), 7.73-7.74 (2H, m), 12.6 (1H, brs).

(d) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-fluoro-5-phenylisoxazole

4-Fluoro-3-hydroxy-5-phenylisoxazole (100 mg) and 2-(N-tert-butoxycarbonylamino)ethanol (108 mg) were subjected to reaction and post-treatment in a similar manner to that described in Example 1(a) to obtain the title compound (132 mg, 73%) as colorless crystals.

m.p.: 103-104°C;

IR spectrum (KBr) ν_{\max} cm^{-1} : 3325, 1718, 1667, 1549, 1532;

NMR spectrum (CDCl₃) δ ppm: 1.46 (9H, s), 3.60 (2H, q, J=5.1Hz), 4.41 (2H, t, J=5.1Hz), 4.97 (1H, brs), 7.41-7.52 (3H, m), 7.76-7.79 (2H, m).

(e) 3-(2-Aminoethoxy)-4-fluoro-5-phenylisoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-fluoro-5-phenylisoxazole (110 mg) was subjected to reaction and post-treatment in a similar manner to that described in Example 1(b) to obtain the title compound (82 mg, 93%) as colorless crystals.

m.p.: 207-211°C (decomposed);

IR spectrum (KBr) ν_{\max} cm^{-1} : 2999, 2965, 2909, 2845, 1665, 1602, 1581, 1555, 1532;

NMR spectrum (DMSO- d_6) δ ppm: 3.32 (2H, t, J=5.1Hz), 4.55 (2H, t, J=5.1Hz), 7.55-7.64 (3H, m), 7.75-7.78 (2H, m), 8.17 (3H, brs).

Example 69

3-(2-Dimethylaminoethoxy)-5-phenylisoxazole hydrochloride (Compound list No.: 27)

(a) 3-(2-Bromoethoxy)-5-phenylisoxazole

Triphenylphosphine (15.7 g) was dissolved in toluene (200 ml), and the solution was cooled to 5°C. Diethyldiazodicarboxylate (10.4 g) was added to the solution, and the resulting mixture was stirred at the same temperature for 10

minutes. Then, 3-hydroxy-5-phenylisoxazole (8.0 g) and 2-bromoethanol (7.5 g) were added in this order to the reaction mixture, followed by stirring of the resulting mixture at room temperature for 10 minutes and further at room temperature for 2 hours. Insolubles were filtered off from the mixture, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (eluent: hexane/ethyl acetate = 4/1) to obtain the title compound (12.1 g, 90%) as colorless crystals.

IR spectrum (KBr) ν_{\max} cm^{-1} : 3148, 1623, 1597, 1576, 1510;

NMR spectrum (CDCl_3) δ ppm: 3.71 (2H, t, $J=5.9\text{Hz}$), 4.62 (2H, t, $J=5.9\text{Hz}$), 6.19 (1H, s), 7.42-7.49 (3H, m), 7.70-7.77 (2H, m).

(b) 3-(2-Dimethylaminoethoxy)-5-phenylisoxazole

After dimethylamine hydrochloride (815 mg) was dissolved in methanol (50 ml), 3-(2-bromoethoxy)-5-phenylisoxazole (268 mg) and triethylamine (2.8 ml) were added thereto, followed by reflux of the resulting mixture for 8 hours. After the reaction, the solvent was evaporated under reduced pressure. The residue was dissolved in ether (20 ml), and the solution was washed with a diluted aqueous NaCl solution (20 ml), followed by drying of the organic layer over anhydrous magnesium sulfate. After filtration, the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (eluent: ethyl acetate) to obtain the title compound (146 mg, 63%) as colorless crystals.

m.p.: 28-29°C;

IR spectrum (KBr) ν_{\max} cm^{-1} : 2980, 2951, 1625, 1597, 1577, 1514;

NMR spectrum (CDCl_3) δ ppm: 2.34 (6H, s), 2.75 (2H, t, $J=5.4\text{Hz}$), 4.40 (2H, t, $J=5.4\text{Hz}$), 6.18 (1H, s), 7.40-7.48 (3H, m), 7.69-7.73 (2H, m).

(c) 3-(2-Dimethylaminoethoxy)-5-phenylisoxazole hydrochloride

3-(2-Dimethylaminoethoxy)-5-phenylisoxazole (130 mg) was dissolved in dioxane (1.0 ml), and a solution of 4N hydrochloric acid/dioxane (0.2 ml) was added thereto, and the resulting mixture was allowed to stand at room

temperature for 15 minutes. The solvent was evaporated under reduced pressure and the residue was washed with ethyl acetate (5 ml) to obtain the title compound (148 mg, 98%) as colorless crystals.

m.p.: 163-164°C (decomposed);

IR spectrum (KBr) ν_{\max} cm^{-1} : 3133, 1624, 1597, 1577, 1513;

NMR spectrum (DMSO- d_6) δ ppm: 2.84 (6H, s), 3.55 (2H, t, J=5.0Hz), 4.61 (2H, t, J=5.0Hz), 7.50-7.58 (3H, m), 7.81-7.87 (2H, m), 10.31 (1H, brs).

Example 70

5-Phenyl-3-(2-(1-piperidyl)ethoxy)isoxazole hydrochloride (Compound list No.: 28)

(a) 5-Phenyl-3-(2-(1-piperidyl)ethoxy)isoxazole

3-(2-Bromoethoxy)-5-phenylisoxazole (268 mg) was dissolved in methanol (1.0 ml), and piperidine (426 mg) was added thereto, followed by reflux of the resulting mixture for 3 hours. After the reaction mixture was added to ice-cold water (20 ml) and extracted with ether (20 ml x 2), the organic layer was dried over anhydrous magnesium sulfate. After filtration, the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (eluent: ethyl acetate) to obtain the title compound (251 mg, 92%) as colorless crystals.

m.p.: 75-76°C;

IR spectrum (KBr) ν_{\max} cm^{-1} : 3149, 1625, 1598, 1577, 1513;

NMR spectrum (CDCl₃) δ ppm: 1.42-1.48 (2H, m), 1.54-1.67 (4H, m), 2.50 (4H, brs), 2.79 (2H, t, J=5.8Hz), 4.42 (2H, t, J=5.8Hz), 6.16 (1H, s), 7.41-7.48 (3H, m), 7.69-7.75 (2H, m).

(b) 5-Phenyl-3-(2-(1-piperidyl)ethoxy)isoxazole hydrochloride

5-Phenyl-3-(2-piperidylethoxy)isoxazole (200 mg) was subjected to reaction and post-treatment in a similar manner to that described in Example 1(b) to obtain the title compound (181 mg, 80%) as colorless crystals.

m.p.: 190-192°C (decomposed);

IR spectrum (KBr) ν_{\max} cm^{-1} : 2950, 2938, 1620, 1595, 1576, 1511;

NMR spectrum (DMSO- d_6) δ ppm: 1.32-1.44 (2H, m), 1.68-1.86 (4H, m), 2.95-3.04 (2H, m), 3.45-3.58 (4H, m), 4.64 (2H, t, $J=4.7\text{Hz}$), 6.87 (1H, s), 7.50-7.58 (3H, m), 7.81-7.85 (2H, m), 9.94-10.05 (1H, brs).

Example 71

5-Phenyl-3-(2-(1-pyrrolidinyl)ethoxy)isoxazole hydrochloride (Compound list No.: 1350)

(a) 5-Phenyl-3-(2-(1-pyrrolidinyl)ethoxy)isoxazole

Piperidine (710 mg) was added to 3-(2-bromoethoxy)-5-phenylisoxazole (268 mg), and the mixture was stirred at 100°C for one hour. After ice-cold water (20 ml) was added to the reaction mixture and the mixture was extracted with ether (20 ml x 2), the organic layer was dried over anhydrous magnesium sulfate. After filtration, the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (eluent: ethyl acetate) to obtain the title compound (211 mg, 82%) as colorless crystals.

m.p.: 61-62°C;

IR spectrum (KBr) ν_{\max} cm^{-1} : 1623, 1596, 1575, 1509;

NMR spectrum (CDCl₃) δ ppm: 1.77-1.87 (4H, m), 2.57-2.65 (4H, m), 2.92 (2H, t, $J=5.5\text{Hz}$), 4.43 (2H, t, $J=5.5\text{Hz}$), 6.18 (1H, s), 7.40-7.48 (3H, m), 7.69-7.75 (2H, m).

(b) 5-Phenyl-3-(2-(1-pyrrolidinyl)ethoxy)isoxazole hydrochloride

5-Phenyl-3-(2-(1-pyrrolidinyl)ethoxy)isoxazole (200 mg) was subjected to reaction and post-treatment in a similar manner to that described in Example 1(b) to obtain the title compound (211 mg, 93%) as colorless crystals.

m.p.: 182-184°C (decomposed);

IR spectrum (KBr) ν_{\max} cm^{-1} : 2982, 2950, 2884, 2848, 2671, 2653, 2607, 2560, 2480, 1619, 1595, 1576, 1511;

NMR spectrum (DMSO- d_6) δ ppm: 1.83-2.08 (4H, m), 3.01-3.21 (2H, m), 3.50-3.68 (4H, m), 4.60 (2H, t, $J=5.1\text{Hz}$), 6.87 (1H, s), 7.50-7.58 (3H, m), 7.81-7.87 (2H,

m), 10.63 (1H, brs).

Example 72

3-(2-(4-Morpholinyl)ethoxy)-5-phenylisoxazole hydrochloride (Compound list No.: 29)

(a) 3-(2-(4-Morpholinyl)ethoxy)-5-phenylisoxazole

Morpholine (871 mg) was added to 3-(2-bromoethoxy)-5-phenylisoxazole (268 mg), and the mixture was stirred at 100°C for one hour. After ice-cold water (20 ml) was added to the reaction mixture and the resulting mixture was extracted with ether (20 ml x 2), the organic layer was dried over anhydrous magnesium sulfate. After filtration, the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (eluent: ethyl acetate) to obtain the title compound (250 mg, 91%) as colorless crystals.

m.p.: 66-67°C;

IR spectrum (KBr) ν_{\max} cm^{-1} : 3152, 1623, 1596, 1575, 1512;

NMR spectrum (CDCl_3) δ ppm: 2.57 (2H, t, J=5.4Hz), 2.82 (2H, t, J=5.4Hz), 3.75 (2H, t, J=5.5Hz), 4.44 (2H, t, J=5.5Hz), 6.17 (1H, s), 7.41-7.50 (3H, m), 7.69-7.76 (2H, m).

(b) 3-(2-(4-Morpholinyl)ethoxy)-5-phenylisoxazole hydrochloride

3-(2-(4-Morpholinyl)ethoxy)-5-phenylisoxazole (200 mg) was subjected to reaction and post-treatment in a similar manner to that described in Example 1(b) to obtain the title compound (202 mg, 89%) as colorless crystals.

m.p.: 180-182°C (decomposed);

IR spectrum (KBr) ν_{\max} cm^{-1} : 2981, 2925, 2897, 2878, 2683, 2646, 2584, 2512, 2469, 2429, 1620, 1595, 1576, 1512;

NMR spectrum (DMSO-d_6) δ ppm: 3.10-3.25 (2H, m), 3.40-3.55 (2H, m), 3.55-3.65 (2H, m), 3.70-3.85 (2H, m), 3.90-4.05 (2H, m), 4.67 (2H, brs), 6.86 (1H, s), 7.50-7.58 (3H, m), 7.81-7.87 (2H, m), 10.96 (1H, brs).

Example 73

5-Phenyl-3-(2-(1-piperazinyl)ethoxy)isoxazole dihydrochloride (Compound list No.: 1351)

(a) 3-(2-(4-N-tert-Butoxycarbonyl-1-piperazinyl)ethoxy)-5-phenylisoxazole

3-(2-Bromoethoxy)-5-phenylisoxazole (268 mg) was dissolved in methanol (1.0 ml), and piperazine (861 mg) was added thereto, followed by reflux of the resulting mixture for 3 hours. After the reaction mixture was added to a diluted aqueous NaCl solution (40 ml) and extracted with dichloromethane (40 ml x 2), the organic layer was dried over anhydrous magnesium sulfate. After filtration, the solvent was evaporated under reduced pressure. The residue was dissolved in dichloromethane (5 ml), and di-tert-butyl dicarbonate (1.09 g) was added to the mixture, followed by stirring of the resulting mixture at room temperature for 30 minutes. After the reaction, the solvent was evaporated under reduced pressure and the residue was purified by silica gel column chromatography (eluent: hexane/ethyl acetate = 2/1) to obtain the title compound (310 mg, 83%) as colorless crystals.

m.p.: 110-111°C;

IR spectrum (KBr) ν_{\max} cm^{-1} : 3146, 1696, 1626, 1598, 1577, 1514;

NMR spectrum (CDCl_3) δ ppm: 1.46 (9H, s), 2.51 (2H, t, J=5.0Hz), 2.83 (2H, t, J=5.5Hz), 3.47 (2H, t, J=5.0Hz), 4.43 (2H, t, J=5.5Hz), 6.16 (1H, s), 7.41-7.49 (3H, m), 7.69-7.75 (2H, m).

(b) 5-Phenyl-3-(2-(1-piperazinyl)ethoxy)isoxazole dihydrochloride

3-(2-(4-tert-Butoxycarbonyl-1-piperazinyl)ethoxy)-5-phenylisoxazole (250 mg) was subjected to reaction and post-treatment in a similar manner to that described in Example 1(b) to obtain the title compound (222 mg, 95%) as colorless crystals.

m.p.: 216-222°C (decomposed);

IR spectrum (KBr) ν_{\max} cm^{-1} : 3443, 3189, 3123, 3001, 2972, 2950, 2915, 2771, 2715, 2623, 2527, 2422, 1704, 1648, 1621, 1596, 1578, 1565, 1514;

NMR spectrum (DMSO-d_6) δ ppm: 3.00-3.85 (10H, brs), 4.64 (2H, brs), 6.86

(1H, s), 7.50-7.57 (3H, m), 7.82-7.85 (2H, m), 9.10-9.80 (1H, brs).

Example 74

3-(2-N-Methylaminoethoxy)-5-phenylisoxazole hydrochloride (Compound list No.: 25)

(a) 3-(2-(4-N-tert-Butoxycarbonyl-N-methylamino)ethoxy)-5-phenylisoxazole

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-phenylisoxazole (304 mg) was dissolved in dimethylformamide (3 ml), and sodium hydride [$>55\%$ (oily), 52 mg] was added thereto, followed by stirring of the resulting mixture at room temperature for one hour. The reaction mixture was cooled to 5°C , and methyl iodide (220 mg) was added thereto, followed by stirring of the resulting mixture at the same temperature for 15 minutes and further at room temperature for one hour. At the end of this time, the reaction mixture was added to ice-cold water (20 ml) and extracted with ether ($20\text{ml}\times 2$), the organic layer was dried over anhydrous magnesium sulfate. After filtration, the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (eluent: hexane/ethyl acetate = 4/1) to obtain the title compound (296 mg, 93%) as colorless crystals.

m.p.: $72-73^{\circ}\text{C}$;

IR spectrum (KBr) $\nu_{\text{max}} \text{ cm}^{-1}$: 3125, 1680, 1626, 1596, 1578, 1514;

NMR spectrum (CDCl_3) δ ppm: 1.49 (9H, s), 2.97 (3H, s), 3.64 (2H, brs), 4.41 (2H, brs), 6.14 (1H, s), 7.41-7.52 (3H, m), 7.69-7.75 (2H, m).

(b) 3-(2-Methylaminoethoxy)-5-phenylisoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonyl-N-methylamino)ethoxy)-5-phenylisoxazole (250 mg) was subjected to reaction and post-treatment in a similar manner to that described in Example 1(b) to obtain the title compound (197 mg, 99%) as colorless crystals.

m.p.: $219-221^{\circ}\text{C}$ (decomposed);

IR spectrum (KBr) $\nu_{\text{max}} \text{ cm}^{-1}$: 3134, 3022, 2981, 2954, 2883, 2869, 2843, 2803, 2785, 2734, 1621, 1597, 1578, 1516;

NMR spectrum (DMSO-d₆) δ ppm: 2.62 (3H, s), 3.37 (2H, t, J=5.1Hz), 4.50 (1H, t, J=5.1Hz), 6.85 (1H, s), 7.50-7.58 (3H, m), 7.82-7.88 (2H, m), 8.79 (2H, brs).

Example 75

3-(2-Acetylaminoethoxy)-5-phenylisoxazole (Compound list No.: 1347)

3-(2-Aminoethoxy)-5-phenylisoxazole hydrochloride (240 mg) was suspended in anhydrous tetrahydrofuran (5 ml), and the suspension was cooled to 5°C. Acetyl chloride (94 mg) and triethylamine (243 mg) were added thereto, and the resulting mixture was stirred at the same temperature for 30 minutes. At the end of this time, the reaction mixture was added to ice-cold water (40 ml) and extracted with ethyl acetate (40 ml x 2). Then, the organic layer was dried over anhydrous magnesium sulfate. After filtration, the solvent was evaporated under reduced pressure. The crystals thus obtained were recrystallized from ethyl acetate to obtain the title compound (225 mg, 91%) as colorless crystals.

m.p.: 144-145°C;

IR spectrum (KBr) ν_{\max} cm⁻¹ : 3316, 1641, 1624, 1597, 1578, 1547, 1509;

NMR spectrum (CDCl₃) δ ppm: 2.03(3H, s), 3.70 (2H, q, J=5.1Hz), 4.39 (2H, q, J=5.1Hz), 6.16 (1H, brs), 6.26 (1H, s), 7.41-7.49 (3H, m), 7.70-7.75 (2H, m).

Example 76

3-(2-Benzoylaminoethoxy)-5-phenylisoxazole (Compound list No.: 1349)

3-(2-Aminoethoxy)-5-phenylisoxazole hydrochloride (240 mg) was suspended in anhydrous tetrahydrofuran (5 ml), and the suspension was cooled to 5°C. Benzoyl chloride (168 mg) and triethylamine (243 mg) were added thereto, followed by stirring of the resulting mixture at the same temperature for 30 minutes. At the end of this time, the reaction mixture was added to ice-cold water (40 ml) and extracted with ethyl acetate (40 ml x 2). Then, the organic layer was dried over anhydrous magnesium sulfate. After filtration, the solvent was evaporated under reduced pressure. The residue was recrystallized from ethyl acetate to obtain the title compound (276 mg, 90%) as colorless crystals.

m.p.: 129-130°C;

IR spectrum (KBr) ν_{\max} cm^{-1} : 3359, 1641, 1626, 1597, 1578, 1534, 1518;

NMR spectrum (CDCl_3) δ ppm: 3.92 (2H, q, $J=5.1\text{Hz}$), 4.52 (2H, q, $J=5.1\text{Hz}$), 6.17 (1H, s), 6.74 (1H, brs), 7.42-7.53 (6H, m), 7.71-7.74 (2H, m), 7.79-7.88 (2H, m).

Example 77

3-(2-Methoxycarbonylaminoethoxy)-5-phenylisoxazole (Compound list No.: 1348)

3-(2-Aminoethoxy)-5-phenylisoxazole hydrochloride (240 mg) was suspended in anhydrous tetrahydrofuran (5 ml), and the suspension was cooled to 5°C . Then, methyl chloroformate (113 mg) and triethylamine (243 mg) were added thereto, and the resulting mixture was stirred at the same temperature for 30 minutes. At the end of this time, the reaction mixture was added to ice-cold water (40 ml) and extracted with ethyl acetate (40 ml x 2). Then, the organic layer was dried over anhydrous magnesium sulfate. After filtration, the solvent was evaporated under reduced pressure. The crystals thus obtained were recrystallized from a mixture of ethyl acetate and ether to obtain the title compound (233 mg, 89%) as colorless crystals.

m.p.: $95\text{-}96^\circ\text{C}$;

IR spectrum (KBr) ν_{\max} cm^{-1} : 3293, 1729, 1624, 1596, 1577, 1552, 1516;

NMR spectrum (CDCl_3) δ ppm: 3.63 (2H, q, $J=5.1\text{Hz}$), 3.70 (3H, s), 4.37 (2H, q, $J=5.1\text{Hz}$), 5.11 (1H, brs), 6.14 (1H, s), 7.40-7.49 (3H, m), 7.69-7.75 (2H, m).

Example 78

3-(2-Aminoethylthio)-5-phenylisoxazole hydrochloride (Compound list No.: 15)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethylthio)-5-phenylisoxazole

2-(N-tert-Butoxycarbonylamino)ethanethiol (300 mg) was dissolved in dimethylformamide (3.0 ml), and the mixture was cooled to 5°C , followed by addition of sodium hydride [$>55\%$ (oil), 73 mg]. The resulting mixture was stirred at the same temperature for 30 minutes. Then, 3-chloro-5-phenylisoxazole (300 mg) was added to the reaction mixture, followed by stirring of the mixture at the same temperature for 30 minutes and further at room

temperature for 3 days. At the end of this time, the reaction mixture was added to ice-cold water (40 ml) and extracted with ethyl acetate (40 ml x 2), and the organic layer was dried over anhydrous magnesium sulfate. After filtration, the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (eluent: hexane/ethyl acetate = 4/1) to obtain the title compound (130 mg, 24%) as colorless crystals.

m.p.: 87-88°C;

IR spectrum (KBr) ν_{\max} cm^{-1} : 3376, 1682, 1613, 1572, 1521, 1494;

NMR spectrum (CDCl_3) δ ppm: 1.44 (9H, s), 3.27 (2H, t, $J=6.3\text{Hz}$), 3.53 (2H, q, $J=6.3\text{Hz}$), 5.01 (1H, brs), 6.44 (1H, s), 7.42-7.50 (3H, m), 7.71-7.76 (2H, m).

(b) 3-(2-Aminoethylthio)-5-phenylisoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethylthio)-5-phenylisoxazole (64 mg) was subjected to reaction and post-treatment in a similar manner to that described in Example 1(b) to obtain the title compound (40 mg, 78%) as colorless crystals.

m.p.: 196-198°C (decomposed);

IR spectrum (KBr) ν_{\max} cm^{-1} : 3141, 2988, 2951, 2923, 1609, 1590, 1569, 1492;

NMR spectrum (DMSO-d_6) δ ppm: 3.18 (2H, t, $J=7.2\text{Hz}$), 3.36 (2H, t, $J=7.2\text{Hz}$), 7.23 (1H, s), 7.53-7.59 (3H, m), 7.83-7.97 (2H, m), 7.97 (3H, brs).

Example 79

3-(2-Aminoethoxy)-4-isopropyl-5-(3-pyridyl)isoxazole dihydrochloride

(Compound list No.: 1065)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-isopropyl-5-(3-pyridyl)isoxazole

3-Hydroxy-4-isopropyl-5-(3-pyridyl)isoxazole (0.12 g) and 2-(N-tert-butoxycarbonylamino)ethanol (0.09 g) were subjected to reaction and post-treatment in a similar manner to that described in Example 9(a) to obtain the title compound (0.14 g, 74%) as a colorless powder.

IR spectrum (KBr) ν_{\max} cm^{-1} : 3323, 3247, 2979, 1753, 1690;

NMR spectrum (CDCl_3) δ ppm: 1.31 (6H, d, $J=6.8\text{Hz}$), 1.46 (9H, s), 3.06 (1H, qq, $J=6.8\text{Hz}$, $J=6.8\text{Hz}$), 3.60 (2H, q, $J=5.2\text{Hz}$), 4.39 (2H, t, $J=5.2\text{Hz}$), 4.84 (1H, brs), 7.43 (1H, dd, $J=8.0\text{Hz}$, $J=4.7\text{Hz}$), 7.90 (1H, ddd, $J=8.0\text{Hz}$, $J=2.0\text{Hz}$, $J=1.4\text{Hz}$), 8.69 (1H, dd, $J=4.7\text{Hz}$, $J=1.4\text{Hz}$), 8.83 (1H, d, $J=2.0\text{Hz}$).

(b) 3-(2-Aminoethoxy)-4-isopropyl-5-(3-pyridyl)isoxazole dihydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-isopropyl-5-(3-pyridyl)-isoxazole (0.13 g) was subjected to reaction and post-treatment in a similar manner to that described in Example 9(b) to obtain the title compound (0.07 g, 64%) as colorless crystals.

m.p.: 193-197°C (decomposed);

IR spectrum (KBr) ν_{\max} cm^{-1} : 3049, 2963, 2874, 1549;

NMR spectrum (DMSO-d_6) δ ppm: 1.27 (6H, d, $J=7.0\text{Hz}$), 3.03 (1H, qq, $J=7.0\text{Hz}$, $J=7.0\text{Hz}$), 3.29 (2H, t, $J=5.4\text{Hz}$), 4.48 (2H, t, $J=5.4\text{Hz}$), 7.68 (1H, dd, $J=8.0\text{Hz}$, $J=5.0\text{Hz}$), 8.12 (1H, ddd, $J=8.0\text{Hz}$, $J=2.0\text{Hz}$, $J=1.4\text{Hz}$), 8.28 (3H, brs), 8.78 (1H, dd, $J=5.0\text{Hz}$, $J=1.4\text{Hz}$), 8.85 (1H, d, $J=2.0\text{Hz}$).

Example 80

3-(2-Aminoethoxy)-4-(1-chloropropyl)-5-phenylisoxazole hydrochloride

(Compound list No.: 1802)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-(1-hydroxypropyl)-5-phenylisoxazole

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-phenylisoxazole (0.2 g), butyllithium (1.6M hexane solution, 0.9 ml) and propionaldehyde (0.06 ml) were subjected to reaction and post-treatment in a similar manner to that described in Example 44(a) to obtain the title compound (0.14 g, 58%) as a colorless oil.

IR spectrum (CHCl_3) ν_{\max} cm^{-1} : 3602, 3459, 2980, 2937, 1712;

NMR spectrum (CDCl_3) δ ppm: 0.95 (3H, t, $J=7.4\text{Hz}$), 1.45 (9H, s), 1.83-2.08 (2H, m), 2.51 (1H, brs), 3.57 (2H, q, $J=5.2\text{Hz}$), 4.39 (2H, t, $J=5.2\text{Hz}$), 4.68 (1H, t,

J=7.2Hz), 4.94 (1H, brs), 7.43-7.52 (3H, m), 7.64-7.75 (2H, m).

(b) 3-(2-Aminoethoxy)-4-(1-chloropropyl)-5-phenylisoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-(1-hydroxypropyl)-5-phenylisoxazole (0.13 g) was subjected to reaction and post-treatment in a similar manner to that described in Example 1(b) to obtain the title compound (0.10 g, 91%) as colorless crystals.

m.p.: 122-124°C;

IR spectrum (KBr) ν_{\max} cm^{-1} : 2974, 2936, 2878, 1631, 1600, 1577;

NMR spectrum (DMSO- d_6) δ ppm: 0.94 (3H, t, J=7.3Hz), 2.11-2.28 (2H, m), 3.31 (2H, t, J=5.1Hz), 4.50 (2H, t, J=5.1Hz), 5.14 (1H, t, J=6.5Hz), 7.61-7.65 (3H, m), 7.72-7.75 (2H, m).

Example 81

3-(2-Aminoethoxy)-4-(1-chloroisobutyl)-5-phenylisoxazole hydrochloride

(Compound list No.: 1804)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-(1-hydroxyisobutyl)-5-phenylisoxazole

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-phenylisoxazole (0.2 g), butyllithium (1.6M hexane solution, 0.9 ml) and isobutylaldehyde (0.07 ml) were subjected to reaction and post-treatment in a similar manner to that described in Example 44(a) to obtain the title compound (0.21 g, 84%) as a colorless oil.

IR spectrum (CHCl_3) ν_{\max} cm^{-1} : 3606, 3459, 2981, 2937, 1712, 1637;

NMR spectrum (CDCl_3) δ ppm: 0.79 (3H, d, J=6.6Hz), 1.11 (3H, d, J=6.6Hz), 1.45 (9H, s), 2.16-2.29 (1H, m), 2.56 (1H, brs), 3.58 (2H, q, J=5.1Hz), 4.30-4.48 (3H, m), 4.94 (1H, brs), 7.46-7.51 (3H, m), 7.70-7.74 (2H, m).

(b) 3-(2-Aminoethoxy)-4-(1-chloroisobutyl)-5-phenylisoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-(1-hydroxyisobutyl)-5-phenylisoxazole (0.20 g) was subjected to reaction and post-treatment in a

similar manner to that described in Example 1(b) to obtain the title compound (0.11 g, 65%) as colorless crystals.

m.p.: 165-167°C;

IR spectrum (KBr) ν_{\max} cm^{-1} : 2967, 2902, 2870, 1636, 1600;

NMR spectrum (DMSO- d_6) δ ppm: 0.78 (3H, d, J=6.6Hz), 1.15 (3H, d, J=6.6Hz), 2.53-2.63 (1H, m), 3.30 (2H, t, J=5.2Hz), 4.50 (2H, t, J=5.2Hz), 4.85 (1H, d, J=10.7Hz), 7.60-7.65 (3H, m), 7.71-7.75 (2H, m), 8.25 (3H, brs).

Example 82

3-(2-Aminoethoxy)-4-(1-chloroisopentyl)-5-phenylisoxazole hydrochloride

(Compound list No.: 1806)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-(1-hydroxyisopentyl)-5-phenylisoxazole

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-phenylisoxazole (0.2 g), butyllithium (1.6M hexane solution, 0.9 ml) and isovaleraldehyde (0.09 ml) were subjected to reaction and post-treatment in a similar manner to that described in Example 44(a) to obtain the title compound (0.2 g, 77%) as a colorless oil.

IR spectrum (CHCl_3) ν_{\max} cm^{-1} : 3601, 3459, 2981, 2961, 2936, 1712, 1639;

NMR spectrum (CDCl_3) δ ppm: 0.87 (3H, d, J=6.3Hz), 0.92 (3H, d, J=6.3Hz), 1.45 (9H, s), 1.63-1.78 (2H, m), 1.83-1.98 (1H, m), 3.58 (2H, q, J=5.2Hz), 4.40 (2H, t, J=5.2Hz), 4.75-5.00 (2H, m), 7.44-7.50 (3H, m), 7.66-7.71 (2H, m).

(b) 3-(2-Aminoethoxy)-4-(1-chloroisopentyl)-5-phenylisoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-(1-hydroxyisopentyl)-5-phenylisoxazole (0.19 g) was subjected to reaction and post-treatment in a similar manner to that described in Example 1(b) to obtain the title compound (0.11 g, 69%) as colorless crystals.

m.p.: 134-136°C;

IR spectrum (KBr) ν_{\max} cm^{-1} : 2961, 2905, 2873, 1630;

NMR spectrum (DMSO- d_6) δ ppm: 0.79 (3H, d, J=6.6Hz), 0.82 (3H, d,

J=6.6Hz), 1.57 (1H, qq, J=6.6Hz, J=6.6Hz), 2.00-2.12 (2H, m), 3.31 (2H, t, J=5.4Hz), 4.51 (2H, t, J=5.4Hz), 5.21 (1H, t, J=8.0Hz), 7.62-7.65 (3H, m), 7.69-7.72 (2H, m), 8.22 (3H, brs).

Example 83

3-(2-Aminoethoxy)-5-(2,4-difluorophenyl)-4-isopropylisoxazole hydrochloride

(Compound list No.: 99)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(2,4-difluorophenyl)-4-isopropylisoxazole

5-(2,4-Difluorophenyl)-3-hydroxy-4-isopropylisoxazole (0.2 g) and 2-(N-tert-butoxycarbonylamino)ethanol (0.16 g) were subjected to reaction and post-treatment in a similar manner to that described in Example 1(a) to obtain the title compound (0.25 g, 78%) as a colorless powder.

IR spectrum (KBr) ν_{\max} cm^{-1} : 3400, 2980, 2936, 1683;

NMR spectrum (CDCl_3) δ ppm: 1.22 (6H, d, J=6.9Hz), 1.46 (9H, s), 2.78 (1H, qq, J=6.9Hz, J=6.9Hz), 3.59 (2H, q, J=5.2Hz), 4.38 (2H, t, J=5.2Hz), 4.86 (1H, brs), 6.90-7.03 (2H, m), 7.42-7.50 (1H, m).

(b) 3-(2-Aminoethoxy)-5-(2,4-difluorophenyl)-4-isopropylisoxazole hydrochloride

3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-5-(2,4-difluorophenyl)-4-isopropylisoxazole (0.24 g) was subjected to reaction and post-treatment in a similar manner to that described in Example 1(b) to obtain the title compound (0.17 g, 85%) as colorless crystals.

m.p.: 177-179°C;

IR spectrum (KBr) ν_{\max} cm^{-1} : 2973, 2936, 1642, 1614, 1590;

NMR spectrum (DMSO-d_6) δ ppm: 1.18 (6H, d, J=6.9Hz), 2.76 (1H, qq, J=6.9Hz, J=6.9Hz), 3.29 (2H, t, J=5.3Hz), 4.46 (2H, t, J=5.3Hz), 7.28-7.33 (1H, m), 7.50-7.56 (1H, m), 7.60-7.66 (1H, m), 8.12 (3H, brs).

Example 84

3-(2-Aminoethoxy)-4-isopropyl-5-(4-methylphenyl)isoxazole hydrochloride

(Compound list No.: 266)

(a) 3-(2-(N-tert-Butoxycarbonylamino)ethoxy)-4-isopropyl-5-(4-methylphenyl)isoxazole

3-Hydroxy-4-isopropyl-5-(4-methylphenyl)isoxazole (0.2 g) and 2-(N-tert-butoxycarbonylamino)ethanol (0.16 g) were subjected to reaction and post-treatment in a similar manner to that described in Example 1(a) to obtain the title compound (0.27 g, 82%) as a colorless powder.

IR spectrum (KBr) ν_{\max} cm^{-1} : 3383, 2965, 1686;

NMR spectrum (CDCl_3) δ ppm: 1.28 (6H, d, $J=7.0\text{Hz}$), 1.46 (9H, s), 2.41 (3H, s), 3.06 (1H, qq, $J=7.0\text{Hz}$, $J=7.0\text{Hz}$), 3.59 (2H, q, $J=5.2\text{Hz}$), 4.37 (2H, t, $J=5.2\text{Hz}$), 4.84 (1H, brs), 7.27 (2H, d, $J=8.1\text{Hz}$), 7.46 (2H, d, $J=8.1\text{Hz}$).

(b) 3-(2-Aminoethoxy)-4-isopropyl-5-(4-methylphenyl)isoxazole hydrochloride

3-(2-(N-tert-butoxycarbonylamino)ethoxy)-4-isopropyl-5-(4-methylphenyl)isoxazole (0.25 g) was subjected to reaction and post-treatment in a similar manner to that described in Example 1(b) to obtain the title compound (0.13 g, 65%) as colorless crystals.

m.p.: 208-210°C;

IR spectrum (KBr) ν_{\max} cm^{-1} : 2971, 2877, 1641, 1522;

NMR spectrum (DMSO-d_6) δ ppm: 1.25 (6H, d, $J=7.0\text{Hz}$), 2.38 (3H, s), 3.02 (1H, qq, $J=7.0\text{Hz}$, $J=7.0\text{Hz}$), 3.28 (2H, t, $J=5.2\text{Hz}$), 4.44 (2H, t, $J=5.2\text{Hz}$), 7.37 (2H, d, $J=8.1\text{Hz}$), 7.47 (2H, d, $J=8.1\text{Hz}$), 8.20 (3H, brs).

Reference example 1

4-Chloro-5-(4-chlorophenyl)-3-hydroxyisoxazole(a) 4-Chlorocinnamic acid ethyl ester

4-Chlorocinnamic acid (300 g) was suspended in benzene (1200 ml), and ethanol (340 g) and conc. sulfuric acid (14 ml) were added thereto, followed by reflux of the resulting mixture for 15 hours. After the reaction mixture had been

washed successively with a diluted aqueous NaCl solution (500 ml), a saturated aqueous sodium hydrogencarbonate solution (500 ml) and a dilute aqueous NaCl solution (500 ml), the organic layer was dried over anhydrous magnesium sulfate. After filtration, the solvent was evaporated under reduced pressure. The residue was evaporated under reduced pressure to obtain the title compound (334 g, 97%) as a colorless liquid.

b.p.: 147-148°C (4mmHg);

NMR spectrum (CDCl₃) δ ppm: 1.33 (3H, t, J=7.0Hz), 4.30 (2H, q, J=7.0Hz), 6.47 (1H, d, J=16.0Hz), 7.63 (1H, d, J=16.0Hz), 7.63 (4H, ca.s).

(b) α,β-Dibromo-4-chlorocinnamic acid ethyl ester

4-Chlorocinnamic acid ethyl ester (330 g) was dissolved in carbon tetrachloride (1200 ml), and bromine (251 g) was added dropwise thereto under stirring at room temperature. The resulting mixture was stirred at room temperature for 4 hours and allowed to stand overnight. The solvent was evaporated under reduced pressure to obtain the title compound (572 g, 99%) as colorless crystals.

m.p.: 67-68°C;

NMR spectrum (CDCl₃) δ ppm: 1.34 (3H, t, J=7.0Hz), 4.37 (2H, q, J=7.0Hz), 4.80 (1H, d, J=12.0Hz), 5.32 (1H, d, J=12.0Hz), 7.37 (4H, ca.s).

(c) 5-(4-Chlorophenyl)-3-hydroxyisoxazole

Sodium hydroxide (151 g) was dissolved in methanol (600 ml), and an aqueous hydroxylamine hydrochloride (45 g) solution (50 ml) was added dropwise thereto under stirring at 0°C, and then a solution of α,β-dibromo-4-chlorocinnamic acid ethyl ester (200 g) in dioxane (200 ml) was added dropwise to the mixture. The resulting mixture was stirred at room temperature for 4 hours and further refluxed for 5 hours. The reaction mixture was cooled to 5°C, and the pH of the mixture was adjusted to a value of 2 with conc. hydrochloric acid. Then, the reaction mixture was added to water (5 L). Precipitate was separated from the mixture by filtration and washed with water (2 L) and then

with ethanol (1 L) to obtain 90 g of the title compound (85%) as a pale yellow powder.

m.p.: 215-220°C (decomposed);

NMR spectrum (DMF-d₇) δ ppm: 6.66 (1H, s), 7.52-8.20 (4H, m), 11.5-12.0 (1H, brs).

(d) 4-Chloro-5-(4-chlorophenyl)-3-hydroxyisoxazole

To a solution of 5-(4-chlorophenyl)-3-hydroxyisoxazole (50.0 g) in dry tetrahydrofuran (300 ml), a solution of sulfonyl chloride (34.5 g) in dry benzene (50 ml) was added dropwise with stirring at 5°C. The resulting mixture was stirred at the same temperature for 30 minutes and then at room temperature for one hour and further refluxed for 3 hours. The solvent of the reaction mixture was evaporated under reduced pressure, and the solid thus obtained was recrystallized from ethanol to obtain the title compound (44.2 g, 76.8%) as colorless needle-like crystals.

m.p.: 235-238°C (decomposed);

NMR spectrum (DMF-d₇) δ ppm: 7.60-8.20 (4H, m), 12.6-13.6 (1H, brs).

Reference example 2

3-Hydroxy-5-(2-thienyl)isoxazole

2-Thiophenacrylic acid was subjected to reaction and post-treatment in a similar manner to that described in Reference example 1(a), Reference example 1(b) and Reference example 1(c) to obtain the title compound.

m.p.: 163-165°C;

NMR spectrum (DMSO-d₆) δ ppm: 6.38 (1H, s), 7.20-7.25 (1H, m), 7.60-7.80 (2H, m), 11.2-11.6 (1H, brs).

Reference example 3

4-Chloro-3-hydroxy-5-(2-thienyl)isoxazole

3-Hydroxy-5-(2-thienyl)isoxazole was subjected to reaction and post-treatment in a similar manner to that described in Reference example 1(d) to

obtain the title compound.

m.p.: 191-194°C;

NMR spectrum (DMF-d₇) δ ppm: 7.21-7.45 (1H, m), 7.70-8.20 (2H, m), 10.0-13.0 (1H, brs).

Reference example 4

3-Hydroxy-5-(3-pyridyl)isoxazole

3-Pyridineacrylic acid was subjected to reaction and post-treatment in a similar manner to that described in Reference example 1(a), Reference example 1(b) and Reference example 1(c) to obtain the title compound.

m.p.: 212-214°C (decomposed);

NMR spectrum (DMF-d₇) δ ppm: 6.76 (1H, s), 7.40-7.80 (1H, m), 8.10-8.50 (1H, m), 8.66-9.00 (1H, m), 9.05-9.33 (1H, m).

Reference example 5

4-Chloro-3-hydroxy-5-(2-pyridyl)isoxazole

3-Hydroxy-5-(2-pyridyl)isoxazole was subjected to reaction and post-treatment in a similar manner to that described in Reference example 1(d) to obtain the title compound.

Reference example 6

3-Hydroxy-4-isopropyl-5-phenylisoxazole

Ethyl benzoylacetate was subjected to reaction and post-treatment in a similar manner to that described in Agric. Biol. Chem., EN. 50, 1831 (1986) to obtain the title compound.

m.p.: 203-205°C;

NMR spectrum (DMSO-d₆) δ ppm: 1.24 (6H, d, J=7.1Hz), 3.01 (1H, q, J=7.1Hz), 7.47-7.61 (5H, m), 11.2-11.6 (1H, brs).

Reference example 7

2-(N-tert-Butoxycarbonylamino)ethanol

2-Aminoethanol (6.1 g) was dissolved in a mixture of tetrahydrofuran and water (1:1, 100 ml), and di-tert-butyl dicarbonate (21.8 g) was added dropwise thereto under ice-cooling with stirring, followed by stirring of the resulting mixture at the same temperature for one hour and further at room temperature for 5 hours. After ethyl acetate (200 ml) was added to the reaction mixture and the mixture was washed with water, the organic layer was dried over anhydrous magnesium sulfate. After filtration, the solvent was evaporated under reduced pressure to obtain the title compound (15.3 g, 95%) as a colorless oil.

Rf value: 0.35 (Developing solvent: cyclohexane/ethyl acetate = 1/1)

NMR spectrum (CDCl₃) δ ppm: 1.45 (9H, s), 2.35-2.50 (1H, brs), 3.29 (2H, q, J=5.3Hz), 3.71 (2H, q, J=5.3Hz), 4.85-5.05 (1H, brs).

Reference example 8

4-(tert-Butyl)-3-hydroxy-5-phenylisoxazole(a) 2-Benzoyl-3,3-dimethylbutyric acid ethyl ester

Diisopropylamine (5.6 ml) was dissolved in tetrahydrofuran (56 ml), butyllithium (1.6M hexane solution, 25 ml) was added dropwise thereto at 5°C with stirring under a nitrogen atmosphere, and the resulting mixture was stirred for 15 minutes. The reaction mixture was cooled to -70°C, and 3,3-dimethylbutyric acid ethyl ester (6.7 ml) was added dropwise thereto, followed by stirring of the resulting mixture for 10 minutes. Then, benzoyl chloride (2.3 ml) was added dropwise to the reaction mixture and the resulting mixture was stirred at the same temperature for 10 minutes. After the reaction, a saturated aqueous ammonium chloride solution was added to the reaction mixture, and the mixture was extracted with ethyl acetate. The organic layer was washed with a saturated aqueous NaCl solution and dried over anhydrous magnesium sulfate. After filtration, the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (eluent: hexane/ethyl acetate = 20/1) to obtain the title compound (4.5 g, 91%) as a colorless oil.

(b) 4-(tert-Butyl)-3-hydroxy-5-phenylisoxazole

2-Benzoyl-3,3-dimethylbutyric acid ethyl ester (2.0 g) was dissolved in methanol (20 ml), and sodium methoxide (28% methanol solution, 1.6 ml) was added dropwise thereto at 5°C under a nitrogen atmosphere, followed by stirring of the resulting mixture for 10 minutes. The reaction mixture was cooled to -30°C, and a suspension of hydroxylamine hydrochloride (1.1 g) and sodium methoxide (28% methanol solution, 6.2 ml) in methanol (10 ml) was added dropwise to the reaction mixture. The resulting mixture was stirred at the same temperature for 30 minutes, and 6N hydrochloric acid (14 ml) was added to the reaction mixture, followed by stirring of the resulting mixture at 80°C for one hour. After the reaction, the solvent was evaporated under reduced pressure. After the residue was poured into ice-cold water and extracted with ethyl acetate, the organic layer was washed with a saturated aqueous NaCl solution and dried over anhydrous magnesium sulfate. After filtration, the solvent was evaporated under reduced pressure. The residue was crystallized from isopropyl ether to obtain the title compound (0.35 g, 20%) as colorless crystals.

IR spectrum (KBr) ν_{\max} cm^{-1} : 3027, 2993, 2960, 2936, 2869, 2790, 2697, 2623, 2574, 1644, 1600;

NMR spectrum (CDCl_3) δ ppm: 1.22 (9H, s), 7.39-7.50 (5H, m).

Reference example 9

5-(4-Chlorophenyl)-3-hydroxy-4-isopropylisoxazole

(a) 2-(4-Chlorobenzoyl)isovaleric acid ethyl ester

4-Chlorobenzoyl chloride (2.5 ml) and isovaleric acid ethyl ester (1.5 ml) were subjected to reaction and post-treatment in a similar manner to that described in Reference example 8(a) to obtain the title compound (2.2 g, 82%) as a colorless oil.

(b) 5-(4-Chlorophenyl)-3-hydroxy-4-isopropylisoxazole

2-(4-Chlorobenzoyl)isovaleric acid ethyl ester (2.1 g), hydroxylamine

hydrochloride (1.1 g) and sodium methoxide (28% methanol solution, 7.5 ml) were subjected to reaction and post-treatment in a similar manner to that described in Reference example 8(b) to obtain the title compound (1.3 g, 71%) as colorless crystals.

IR spectrum (KBr) ν_{\max} cm^{-1} : 3065, 3018, 2970, 2935, 2875, 2820, 2768, 2695, 2609, 1646;

NMR spectrum (CDCl_3) δ ppm: 1.35 (6H, d, $J=7.0\text{Hz}$), 3.06 (1H, qq, $J=7.0\text{Hz}$, $J=7.0\text{Hz}$), 7.46 (2H, d, $J=8.6\text{Hz}$), 7.54 (2H, d, $J=8.6\text{Hz}$).

Reference example 10

5-(2,4-Difluorophenyl)-3-hydroxyisoxazole

(a) 2,4-Difluorocinnamic acid ethyl ester

2,4-Difluorocinnamic acid (10.1 g) was dissolved in ethanol (100 ml), and conc. sulfuric acid (1 ml) was added thereto, and the resulting mixture was refluxed for 3 hours. The solvent was evaporated under reduced pressure and the residue was poured into ice-cold water and extracted with ethyl acetate. The extract was washed successively with 5% aqueous sodium hydrogencarbonate and a saturated aqueous NaCl solution, and the organic layer was dried over anhydrous magnesium sulfate. After filtration, the solvent was evaporated under reduced pressure to obtain the title compound (11.2 g, 97%) as a colorless oil.

(b) α,β -Dibromo-2,4-difluorocinnamic acid ethyl ester

2,4-Difluorocinnamic acid ethyl ester (11.2 g) was dissolved in carbon tetrachloride (110 ml), and bromine (2.7 ml) was added dropwise thereto at room temperature with stirring, followed by stirring of the mixture at room temperature for 3 hours. After the reaction, the solvent was evaporated under reduced pressure to obtain the title compound (19.6 g, quantitative) as a colorless powder.

(c) 5-(2,4-Difluorophenyl)-3-hydroxyisoxazole

Sodium hydroxide (10.9 g) was dissolved in methanol (110 ml), and a solution of hydroxylamine hydrochloride (4.2 g) in water (10 ml) was added dropwise thereto at 5°C with stirring. A solution of α,β -dibromo-2,4-difluorocinnamic acid ethyl ester (19.6 g) in tetrahydrofuran (20 ml) was added dropwise to the mixture, and the resulting mixture was stirred at room temperature for 2 hours, followed by reflux for 5 hours. The solvent was evaporated under reduced pressure and the residue was poured into ice-cold water. The pH of the reaction mixture was adjusted to a value of 2 with conc. hydrochloric acid, and the reaction mixture was extracted with ethyl acetate. The extract was washed with a saturated aqueous NaCl solution, and the organic layer was dried over anhydrous magnesium sulfate. After filtration, the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (eluent: cyclohexane/ethyl acetate = 1/1) to obtain the title compound (8.2 g, 79%) as colorless crystals.

IR spectrum (KBr) ν_{\max} cm^{-1} : 3170, 3090, 3028, 2848, 2806, 2689, 2655, 2603, 1630;

NMR spectrum (DMSO- d_6) δ ppm: 6.38 (1H, s), 7.25-7.32 (1H, m), 7.47-7.56 (1H, m), 7.89-7.99 (1H, m), 11.62 (1H, brs).

Reference example 11

3-Hydroxy-5-(2-trifluoromethylphenyl)isoxazole

(a) 2-Trifluoromethylcinnamic acid ethyl ester

2-Trifluoromethylcinnamic acid (10.1 g), conc. sulfuric acid (1 ml) and ethanol (100 ml) were subjected to reaction and post-treatment in a similar manner to that described in Reference example 10(a) to obtain the title compound (10.8 g, 95%) as a colorless oil.

(b) α,β -Dibromo-2-trifluoromethylcinnamic acid ethyl ester

2-Trifluoromethylcinnamic acid ethyl ester (10.8 g) and bromine (2.3 ml) were subjected to reaction and post-treatment in a similar manner to that described in Reference example 10(b) to obtain the title compound (17.9 g,

quantitative) as a colorless powder.

(c) 3-Hydroxy-5-(2-trifluoromethylphenyl)isoxazole

α,β -Dibromo-2-trifluoromethylcinnamic acid ethyl ester (17.9 g), hydroxylamine hydrochloride (3.8 g) and sodium hydroxide (9.1 g) were subjected to reaction and post-treatment in a similar manner to that described in Reference example 10(c) to obtain the title compound (7.6 g, 76%) as colorless crystals.

IR spectrum (KBr) ν_{\max} cm^{-1} : 3176, 3096, 3022, 2950, 2836, 2796, 2669, 1620, 1600;

NMR spectrum (DMSO- d_6) δ ppm: 6.34 (1H, s), 7.73-7.87 (3H, m), 7.92-7.95 (1H, m), 11.58 (1H, brs).

Reference example 12

3-Hydroxy-5-(4-trifluoromethylphenyl)isoxazole

(a) Ethyl 4-trifluoromethylcinnamate

4-Trifluoromethylcinnamic acid (10.2 g), conc. sulfuric acid (1 ml) and ethanol (100 ml) were subjected to reaction and post-treatment in a similar manner to that described in Reference example 10(a) to obtain the title compound (10.9 g, 95%) as a colorless oil.

(b) α,β -Dibromo-4-trifluoromethylcinnamic acid ethyl ester

4-Trifluoromethylcinnamic acid ethyl ester (10.8 g) and bromine (2.4 ml) were subjected to reaction and post-treatment in a similar manner to that described in Reference example 10(b) to obtain the title compound (17.9 g, quantitative) as a colorless powder.

(c) 3-Hydroxy-5-(4-trifluoromethylphenyl)isoxazole

α,β -Dibromo-4-trifluoromethylcinnamic acid ethyl ester (17.9 g), hydroxylamine hydrochloride (3.8 g) and sodium hydroxide (9.1 g) were subjected to reaction and post-treatment in a similar manner to that described in Reference example 10(c) to obtain the title compound (8.3 g, 83%) as colorless crystals.

IR spectrum (KBr) ν_{\max} cm^{-1} : 3154, 3018, 2987, 2838, 2788, 2673, 2637,

2607, 2547, 1631, 1614;

NMR spectrum (DMSO- d_6) δ ppm: 6.77 (1H, s), 7.88 (2H, d, $J=8.4\text{Hz}$), 8.03 (2H, d, $J=8.4\text{Hz}$), 11.60 (1H, brs).

Reference example 13

3-Hydroxy-5-(4-isopropylphenyl)isoxazole

(a) 4-Isopropylcinnamic acid ethyl ester

4-Isopropylcinnamic acid (5.0 g), conc. sulfuric acid (0.5 ml) and ethanol (50 ml) were subjected to reaction and post-treatment in a similar manner to that described in Reference example 10(a) to obtain the title compound (5.5 g, 97%) as a colorless oil.

(b) α,β -Dibromo-4-isopropylcinnamic acid ethyl ester

4-Isopropylcinnamic acid ethyl ester (5.5 g) and bromine (1.3 ml) were subjected to reaction and post-treatment in a similar manner to that described in Reference example 10(b) to obtain the title compound (9.5 g, quantitative) as a colorless powder.

(c) 3-Hydroxy-5-(4-isopropylphenyl)isoxazole

α,β -Dibromo-4-isopropylcinnamic acid ethyl ester (9.5 g), hydroxylamine hydrochloride (2.2 g) and sodium hydroxide (5.2 g) were subjected to reaction and post-treatment in a similar manner to that described in Reference example 10(c) to obtain the title compound (3.6 g, 71%) as colorless crystals.

IR spectrum (KBr) ν_{max} cm^{-1} : 3013, 2964, 2934, 2893, 2872, 2793, 2665, 2631, 2542, 1624;

NMR spectrum (DMSO- d_6) δ ppm: 1.22 (6H, d, $J=6.9\text{Hz}$), 2.94 (1H, qq, $J=6.9\text{Hz}$, $J=6.9\text{Hz}$), 6.48 (1H, s), 7.39 (2H, d, $J=8.3\text{Hz}$), 7.72 (2H, d, $J=8.3\text{Hz}$).

Reference example 14

3-Hydroxy-5-(4-phenoxyphenyl)isoxazole

(a) 4-Phenoxybenzaldehyde

4-Phenoxybenzaldehyde (10.0 g) and potassium acetate (9.8 g) were

suspended in acetic anhydride (9.5 ml), followed by reflux at 180°C for 5 hours. After the reaction, the pH of the reaction mixture was adjusted to a value of 2 with hydrochloric acid and the resulting mixture was extracted with ethyl acetate. The extract was washed with a saturated aqueous NaCl solution, and the organic layer was dried over anhydrous magnesium sulfate. After filtration, the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (eluent: hexane/ethyl acetate = 2/1) to obtain the title compound (6.5 g, 54%) as colorless crystals.

(b) 4-Phenoxy-cinnamic acid ethyl ester

4-Phenoxy-cinnamic acid (4.0 g), conc. sulfuric acid (0.4 ml) and ethanol (40 ml) were subjected to reaction and post-treatment in a similar manner to that described in Reference example 10(b) to obtain the title compound (4.4 g, 98%) as a colorless oil.

(c) α,β -Dibromo-4-phenoxy-cinnamic acid ethyl ester

4-Phenoxy-cinnamic acid ethyl ester (4.4 g), and bromine (0.84 ml) were subjected to reaction and post-treatment in a similar manner to that described in Reference example 10(b) to obtain the title compound (7.0 g, quantitative) as a colorless powder.

(d) 3-Hydroxy-5-(4-phenoxyphenyl)isoxazole

α,β -Dibromo-4-phenoxy-cinnamic acid ethyl ester (7.0 g), hydroxyl-amine hydrochloride (1.4 g) and sodium hydroxide (3.3 g) were subjected to reaction and post-treatment in a similar manner to that described in Reference example 10(c) to obtain the title compound (3.4 g, 83%) as colorless crystals.

IR spectrum (KBr) ν_{\max} cm^{-1} : 3147, 3013, 2951, 2852, 2785, 2614, 2557, 1627;

NMR spectrum (DMSO- d_6) δ ppm: 6.47 (1H, s), 7.07-7.25 (5H, m), 7.42-7.48 (2H, m), 7.79-7.83 (2H, m), 11.36 (1H, brs).

Reference example 15

3-Hydroxy-5-(1-naphthyl)isoxazole(a) 1-Naphthylacrylic acid

1-Naphthaldehyde (30.5 g), potassium acetate (38.3 g) and acetic anhydride (36.9 ml) were subjected to reaction and post-treatment in a similar manner to that described in Reference example 14(a) to obtain the title compound (22.4 g, 58%) as a colorless powder.

(b) 1-Naphthylacrylic acid ethyl ester

1-Naphthylacrylic acid (9.5 g), conc. sulfuric acid (1 ml) and ethanol (100 ml) were subjected to reaction and post-treatment in a similar manner to that described in Reference example 10(a) to obtain the title compound (10.2 g, 94%) as a colorless oil.

(c) α,β -Dibromo-1-naphthylacrylic acid ethyl ester

1-Naphthylacrylic acid ethyl ester (10.1 g) and bromine (2.5 ml) were subjected to reaction and post-treatment in a similar manner to that described in Reference example 10(b) to obtain the title compound (17.2 g, quantitative) as a colorless powder.

(d) 3-Hydroxy-5-(1-naphthyl)isoxazole

α,β -Dibromo-1-naphthylacrylic acid ethyl ester (6.0 g), hydroxylamine hydrochloride (1.3 g) and sodium hydroxide (7.2 g) were subjected to reaction and post-treatment in a similar manner to that described in Reference example 10(c) to obtain the title compound (2.6 g, 78%) as colorless crystals.

IR spectrum (KBr) ν_{\max} cm^{-1} : 3136, 3045, 3017, 2790, 2710, 2640, 2569, 1628;

NMR spectrum (DMSO- d_6) δ ppm: 6.53 (1H, s), 7.61-7.70 (3H, m), 7.84-7.87 (1H, m), 8.03-8.13 (2H, m), 8.22-8.28 (1H, m).

Reference example 16

3-Hydroxy-5-(4-hydroxyphenyl)isoxazole

3-Hydroxy-5-(4-methoxyphenyl)isoxazole (5.0 g) was suspended in dichloromethane (50 ml), and aluminum chloride (7.0 g) was added thereto,

followed by reflux of the resulting mixture for 66 hours. At the end of this time, the reaction mixture was poured into ice-cold water, and 6N hydrochloric acid was added thereto, followed by extraction with ethyl acetate. After the extract was washed with a saturated aqueous NaCl solution, the organic layer was dried over anhydrous magnesium sulfate. After filtration, the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (eluent: cyclohexane/ethyl acetate = 2/1) to obtain the title compound (3.8 g, 83%) as colorless crystals.

IR spectrum (KBr) ν_{\max} cm^{-1} : 3140, 3074, 1624, 1588;

NMR spectrum (DMSO- d_6) δ ppm: 6.24 (1H, s), 6.86 (2H, d, $J=8.2\text{Hz}$), 7.62 (2H, d, $J=8.2\text{Hz}$), 10.03 (1H, s), 11.24 (1H, brs).

Reference example 17

5-(2,4-Difluorophenyl)-3-hydroxy-4-isopropylisoxazole

(a) 2-(2,4-Difluorobenzoyl)isovaleric acid ethyl ester

2,4-Difluorobenzoyl chloride (4.9 ml) and isovaleric acid ethyl ester (3.0 ml) were subjected to reaction and post-treatment in a similar manner to that described in Reference example 8(a) to obtain the title compound (3.65 g, 68%) as a colorless liquid.

(b) 5-(2,4-Difluorophenyl)-3-hydroxy-4-isopropylisoxazole

2-(2,4-Difluorobenzoyl)isovaleric acid ethyl ester (1.7 g), hydroxylamine hydrochloride (0.9 g) and sodium methoxide (28% methanol solution, 7.5 ml) were subjected to reaction and post-treatment in a similar manner to that described in Reference example 8(b) to obtain the title compound (0.59 g, 39%) as colorless crystals.

IR spectrum (KBr) ν_{\max} cm^{-1} : 3082, 3033, 2980, 2968, 2935, 2913, 2877, 2836, 2795, 2699, 2636, 2604, 1660, 1610;

NMR spectrum (DMSO- d_6) δ ppm: 1.28 (6H, d, $J=7.0\text{Hz}$), 2.81 (1H, qq, $J=7.0\text{Hz}$, $J=7.0\text{Hz}$), 6.91-7.05 (2H, m), 7.43-7.52 (1H, m).

Reference example 18

4-Cyclopropyl-3-hydroxy-5-phenylisoxazole(a) 2-Benzoyl-2-cyclopropylacetic acid methyl ester

Benzoyl chloride (1.0 ml) and 2-cyclopropylacetic acid methyl ester were subjected to reaction and post-treatment in a similar manner to that described in Reference example 8(a) to obtain the title compound (1.05 g, 28%) as a colorless liquid.

(b) 4-Cyclopropyl-3-hydroxy-5-phenylisoxazole

2-Benzoyl-2-cyclopropylacetic acid methyl ester (0.9 g), hydroxylamine hydrochloride (0.6 g) and sodium methoxide (28% methanol solution, 4.0 ml) were subjected to reaction and post-treatment in a similar manner to that described in Reference example 8(b) to obtain the title compound (0.32 g, 39%) as colorless crystals.

IR spectrum (KBr) ν_{\max} cm^{-1} : 3085, 3066, 3012, 2972, 2909, 2851, 2778, 2716, 2652, 2601, 1646;

NMR spectrum (CDCl_3) δ ppm: 0.79-0.85 (2H, m), 0.91-0.98 (2H, m), 1.67-1.77 (1H, m), 7.43-7.53 (3H, m), 7.86-7.90 (2H, m).

Reference example 19

3-Hydroxy-4-isopropyl-5-(3-pyridyl)isoxazole(a) 2-Nicotinoylisovaleric acid ethyl ester

Nicotinic chloride hydrochloride (0.9 g), isovaleric acid ethyl ester (2.3 ml), diisopropylamine (2.1 ml) and butyllithium (1.6M hexane solution, 9.4 ml) were subjected to reaction and post-treatment in a similar manner to that described in Reference example 8(a) to obtain the title compound (0.78 g, 44%) as a colorless liquid.

(b) 3-Hydroxy-4-isopropyl-5-(3-pyridyl)isoxazole

2-Nicotinoylisovaleric acid ethyl ester (0.76 g), hydroxylamine hydrochloride (0.46 g) and sodium methoxide (28% methanol solution, 3.1 ml) were subjected to reaction and post-treatment in a similar manner to that described in Reference example 8(b) to obtain the title compound (0.15 g, 23%)

as colorless crystals.

IR spectrum (KBr) ν_{max} cm^{-1} : 3069, 3035, 2970, 2934, 2876, 2809, 2770, 2709, 2665, 2605, 2576, 2519, 1511;

NMR spectrum (CDCl_3) δ ppm: 1.38 (6H, d, $J=7.0\text{Hz}$), 3.10 (1H, qq, $J=7.0\text{Hz}$, $J=7.0\text{Hz}$), 7.45 (1H, dd, $J=7.9\text{Hz}$, $J=4.4\text{Hz}$), 7.94 (1H, d, $J=7.9\text{Hz}$), 8.72 (1H, d, $J=4.4\text{Hz}$), 8.88 (1H, s).

Reference example 20

4-Chloro-3-hydroxy-5-(3-pyridyl)isoxazole

3-Hydroxy-5-(3-pyridyl)isoxazole was subjected to reaction and post-treatment in a similar manner to that described in Reference example 1(d) to obtain the title compound.

m.p.: 220-224°C (decomposed);

NMR spectrum (DMF-d_7) δ ppm: 7.60-8.20 (4H, m), 12.6-13.6 (1H, brs).

Test example 1

Type A-monoamine oxidase inhibitory activity

Type A-monoamine oxidase inhibitory activity was determined according to the method described in Biochem. Pharmacol., **12**, 1439 (1963) and J. Neurochem., **35**, 109 (1980). To 30 μl of a sample of mouse brain crude mitochondria (30 μg protein) were added 210 μl of a phosphate buffer (pH 7.4) and a test compound (dissolved in a mixture of 10% DMSO and water), and the resulting mixture was preincubated at 38°C for 20 minutes. Subsequently, ^{14}C -5-hydroxytryptamine (5-HT, final concentration: 100 μM) was added to the preincubated mixture to effect reaction at 38°C for 20 minutes. After the reaction had been quenched by adding 2N-hydrochloric acid (200 μl), a ^{14}C -labelled metabolite obtained by the enzymatic reaction was extracted with a solvent (ethyl acetate : toluene = 1 : 1), and the ^{14}C radioactivity was determined using a liquid scintillation counter to obtain the concentration of the compound which decreases the ^{14}C radioactivity of the control by 50%.

The results of the tests showed that the compounds of Examples 1, 2, 3, 4, 5,

6, 7, 8, 54, 57, 58, 59, 60, 62 and 67 have particularly excellent activities of $IC_{50} \leq 28$ nM.

Preparation example 1

Hard capsule preparation

50 mg of the powdery compound of Example 4, 128.7 mg of lactose, 70 mg of cellulose and 1.3 mg of magnesium stearate were mixed and passed through a 60 mesh sieve, and the resulting powder was filled into 250 mg No. 3 gelatin capsules to obtain a capsule preparation.

Preparation example 2

Tablet preparation

50 mg of the powdery compound of Example 4, 124 mg of lactose, 25 mg of cellulose and 1 mg of magnesium stearate were mixed and formed into tablets with a tableting machine to obtain 200 mg tablets. These tablets may be coated with a sugar coating as necessary.

[Industrial applicability]

The isoxazole derivatives having general formula (I) of the present invention have excellent type A-monoamine oxidase inhibitory activity and are relatively free from undesirable side effects. Therefore, they are useful as agents for treating or preventing (particularly agents for treating) nervous diseases including depression, Parkinson's disease, Alzheimer's dementia (cognitive disorder owing to Alzheimer's disease) or cerebrovascular dementia (cognitive disorder owing to cerebrovascular dementia), (particularly for depression).

In the case where the compounds (I) or the pharmaceutically acceptable salts thereof of the present invention are used as a therapeutic or preventive agent for a nervous disease, the compounds (I) or the salts thereof as such or a mixture obtained appropriately blending the compound (I) with pharmaceutically acceptable excipients, diluents, etc. can be administered in an


oral administration by tablets, capsules, granules, powders or syrups, or in a parenteral administration by injection or suppository, etc.

Such pharmaceutical preparations are prepared by a conventional method using additives such as excipients (for example, sugar derivatives such as lactose, sucrose, glucose, mannitol and sorbitol; starch derivatives such as corn starch, potato starch, α -starch, dextrine and carboxymethyl starch; cellulose derivatives such as crystalline cellulose, low-substituted hydroxypropylcellulose, hydroxypropylmethylcellulose, carmellose, carmellose calcium and internally croscarmellose sodium; acacia; dextran; pullulan; silicate derivatives such as light anhydrous silicic acid, synthetic aluminum silicate or magnesium aluminometasilicate; phosphate salts such as calcium phosphate; carbonate salts such as calcium carbonate; and sulfate salts such as calcium sulfate), binders (for example, the above mentioned excipients; gelatin; polyvinylpyrrolidone; macrogol and the like), decay agents (for example, the above mentioned excipients; croscarmellose sodium, sodium carboxymethyl starch, starch which is chemically modified like crospovidone, cellulose derivatives and the like), lubricants (for example, talc; stearic acid; metal stearates such as calcium stearate and magnesium stearate; colloidal silica; beegum; waxes such as spermaceti; boric acid; glycol; carboxylic acids such as fumaric acid and adipic acid; sodium carboxylates such as sodium benzoate; sulfates such as sodium sulfate; leucine; lauryl sulfates such as sodium lauryl sulfate and magnesium lauryl sulfate; silicic acids such as silicic acid anhydride and silicic acid hydrate; and the starch derivatives in the above mentioned excipients), fungicides (for example, p-hydroxybenzoates such as methylparaben and propylparaben; alcohols such as chlorobutanol, benzyl alcohol and phenylethyl alcohol; benzalkonium chloride; phenols such as phenol and cresol; thimerosal; acetic acid anhydride; and sorbic acid), taste or odor-masking agents (for example, generally used sweeteners, acidulants and flavors and the like), diluents, and solvents for injection (for example, water, ethanol, glycerine and the like).

The dose used differs depending on the symptoms and age of the patient. For example, in the case of oral administration, it is desirable to administer 1 mg (preferably 10 mg) as a lower limit and 2000 mg (preferably 400 mg) as an upper limit, and in the case of intravenous administration, it is desirable to administer 0.1 mg (preferably 1 mg) as a lower limit and 500 mg (preferably 300 mg) as an upper limit for an adult once to 6 times a day depending on the symptoms.

Claims:

1. Isoxazole compound of formula (I):



(I)

wherein

R^1 represents a $C_6 - C_{14}$ aryl group optionally having from 1 to 3 substituents which may be the same as or different from each other and selected from the following substituent group, or a 5- or 6-membered aromatic heterocyclic group optionally having from 1 to 3 substituents and having one or two hetero atoms which may be the same as or different from each other and selected from the group consisting of nitrogen, oxygen and sulfur atoms [the substituent group is a halogen; a $C_1 - C_6$ alkyl; a $C_1 - C_6$ alkyl substituted with a halogen or a $C_1 - C_6$ alkoxy; a $C_1 - C_6$ alkoxy; a $C_6 - C_{14}$ aryl, a $C_7 - C_{18}$ aralkyl, a $C_6 - C_{14}$ aryloxy or a $C_7 - C_{18}$ aralkyloxy optionally having from 1 to 3 substituents which may be the same as or different from each other and selected from the following group (the substituent group is a halogen, a $C_1 - C_6$ alkyl or a $C_1 - C_6$ alkoxy); a cyano; a nitro; a hydroxyl; a $C_1 - C_7$ alkanoyl; a $C_1 - C_7$ alkanoyloxy; a $C_2 - C_7$ alkoxy carbonyl; an amino; a carbamoyl; a mono($C_1 - C_6$ alkyl)carbamoyl; a di($C_1 - C_6$ alkyl)carbamoyl or a mono $C_7 - C_{15}$ arylcarbonylamino (the substituent group is a halogen, a $C_1 - C_6$ alkyl or a $C_1 - C_6$ alkoxy)],

R^2 represents a hydrogen; a halogen; a $C_1 - C_6$ alkyl; a $C_1 - C_6$ alkyl substituted with a halogen or a $C_1 - C_6$ alkoxy; a $C_2 - C_6$ alkenyl; a $C_2 - C_6$ alkynyl; a $C_3 - C_{10}$ cycloalkyl; a $C_3 - C_{10}$ cycloalkenyl; a $C_1 - C_6$ alkoxy; a cyano; a carboxyl; a $C_1 - C_7$ alkanoyl; a $C_2 - C_7$ alkoxy carbonyl; a carbamoyl;

a mono (C₁ - C₆ alkyl)carbamoyl or a di(C₁ - C₆ alkyl)carbamoyl group,

R³ represents an amino, a mono C₁ - C₆ alkylamino, a di(C₁ - C₆ alkyl)amino, a mono C₁ - C₇ alkanoylamino, a mono-C₂ - C₇ alkoxy-carbonylamino, a mono C₇ - C₁₅ arylcarbonylamino optionally having from 1 to 3 substituents which may be the same as or different from each other and selected from the following group (the substituent group is a halogen, a C₁ - C₆ alkyl or a C₁ - C₆ alkoxy) or a 5- or 6-membered saturated heterocyclic group (attached thorough a ring nitrogen atom), which contains one nitrogen atom and optionally contains further one nitrogen atom or oxygen atom,

X represents an oxygen atom or a sulfur atom, and

n represents an integer of 2 to 6,

or pharmaceutically acceptable salts thereof.

2. The isoxazole compounds of Claim 1, wherein R¹ is a C₆ - C₁₄ aryl group optionally having from 1 to 3 substituents which may be the same as or different from each other and selected from the following substituent group, or a 5- or 6-membered aromatic heterocyclic group optionally having from 1 to 3 substituents and having one or two hetero atoms which may be the same as or different from each other and selected from the group consisting of nitrogen, oxygen and sulfur atoms [the substituent group is a halogen; a C₁ - C₆ alkyl; a C₁ - C₆ alkyl substituted with a halogen or a C₁ - C₆ alkoxy; a C₁ - C₆ alkoxy; a C₆ - C₁₄ aryl optionally having from 1 to 3 substituents which may be the same as or different from each other and selected from the following group (the substituent group is a halogen, a C₁ - C₆ alkyl or a C₁ - C₆ alkoxy); a benzyl, a fluorobenzyl, a chlorobenzyl, a difluorobenzyl, a dichlorobenzyl, a methylbenzyl, a dimethylbenzyl, a methoxybenzyl; a phenoxy, a 4-fluorophenoxy, a 4-chlorophenoxy, a 2,4-dichlorophenoxy, a 4-methylphenoxy, a 4-methoxyphenoxy; a benzyloxy, a 4-fluorobenzyloxy, a 4-chlorobenzyloxy, a 2,4-difluorobenzyloxy, a 2,4-dichlorobenzyloxy, a 4-methylbenzyloxy, a 2,4-dimethylbenzyloxy, a 4-methoxybenzyloxy; a cyano; a nitro; a hydroxyl; an

acetoxyl; a C₂ - C₇ alkoxy carbonyl; an amino; a carbamoyl; a mono(C₁ - C₆ alkyl)carbamoyl; a di(C₁ - C₆ alkyl)carbamoyl; a benzoylamino, a 4-fluorobenzoylamino, a chlorobenzoylamino, a 2,4-dichlorobenzoylamino, a 4-toluoylamino and a 4-anisoylamino group], or pharmaceutically acceptable salts thereof.

3. The isoxazole compound of Claim 1, wherein R¹ is a C₆ - C₁₄ aryl group optionally having from 1 to 3 substituents which may be the same as or different from each other and selected from the following substituent group, or a 5- or 6-membered aromatic heterocyclic group optionally having from 1 to 3 substituents and having one or two hetero atoms which may be the same as or different from each other and selected from the group consisting of nitrogen, oxygen and sulfur atoms [the substituent group is a halogen; a C₁ - C₆ alkyl; a C₁ - C₆ alkyl substituted with a halogen or a C₁ - C₆ alkoxy; a C₁ - C₆ alkoxy; a C₆ - C₁₄ aryl optionally having from 1 to 3 substituents which may be the same as or different from each other and selected from the following group (the substituent group is a halogen, a C₁ - C₆ alkyl or a C₁ - C₆ alkoxy); a cyano; a C₂ - C₇ alkoxy carbonyl; a carbamoyl; a mono-(C₁ - C₆ alkyl)carbamoyl; and a di(C₁ - C₆ alkyl)carbamoyl group], or pharmaceutically acceptable salts thereof.

4. The isoxazole compound of Claim 1, wherein R¹ is a C₆ - C₁₄ aryl group optionally having from 1 to 3 substituents which may be the same as or different from each other and selected from the following substituent group, or a 5- or 6-membered aromatic heterocyclic group optionally having one or two substituents and having one or two hetero atoms which may be the same as or different from each other and selected from the group consisting of nitrogen, oxygen and sulfur atoms [the substituent group is a halogen, a C₁ - C₄ alkyl, a fluoromethyl, a difluoromethyl, a trifluoromethyl, a 2-fluoroethyl, a 2,2,2-trifluoroethyl, a methoxymethyl, a methoxyethyl, a C₁ - C₄ alkoxy, a phenyl, a 4-fluorophenyl, a 4-chlorophenyl, a 2,4-dichlorophenyl, a 4-methylphenyl, a 4-

methoxyphenyl, a cyano, a methoxycarbonyl, an ethoxycarbonyl, a carbamoyl, a methylcarbamoyl, an ethylcarbamoyl, an N,N-dimethylcarbamoyl and an N,N-diethylcarbamoyl group], or pharmaceutically acceptable salts thereof.

5. The isoxazole compound of Claim 1, wherein R¹ is a phenyl group optionally having from 1 to 3 substituents which may be the same as or different from each other and selected from the following substituent group, or a furyl, a thienyl or a pyridyl group optionally having one or two substituents [the substituent group is a halogen, a methyl, an ethyl, a trifluoromethyl, a methoxy, a phenyl, a cyano, a methoxycarbonyl, a carbamoyl, a methylcarbamoyl, an ethylcarbamoyl and an N,N-dimethylcarbamoyl group], or pharmaceutically acceptable salts thereof.

6. The isoxazole compound of Claim 1, wherein R¹ is a phenyl group optionally having one or two substituents which may be the same as or different from each other and selected from the following substituent group, or a furyl, a thienyl or a pyridyl group optionally having one substituent [the substituent group is a fluorine, a chlorine, a methyl, an ethyl, a trifluoromethyl and a methoxy group], or pharmaceutically acceptable salts thereof.

7. The isoxazole compound of Claim 1, wherein R¹ is a phenyl, a fluorophenyl, a chlorophenyl, a difluorophenyl, a dichlorophenyl, a methylphenyl, a 2-furyl, a 3-furyl, a 2-thienyl or a 3-thienyl group, or pharmaceutically acceptable salts thereof.

8. The isoxazole compound of Claim 1, wherein R¹ is a phenyl, a 2-chlorophenyl, a 4-chlorophenyl, a 2,4-difluorophenyl, a 2,4-dichlorophenyl, a 2-furyl or a 2-thienyl group, or pharmaceutically acceptable salts thereof.

9. The isoxazole compound of Claim 1, wherein R² is a hydrogen, a halogen, a C₁ - C₆ alkyl, a fluoromethyl, a difluoromethyl, a trifluoromethyl, a 2-

fluoroethyl, a 1-chloroethyl, a 2-chloroethyl, a 2,2,2-trifluoroethyl, a methoxymethyl, a methoxyethyl, a C₂ - C₆ alkenyl, a C₂ - C₆ alkynyl, a cyclopropyl, a cyclopentyl, a cyclohexyl, a 2-cyclopentenyl, a 3-cyclopentenyl, a 2-cyclohexenyl, a 3-cyclohexenyl, a methoxy, an ethoxy, a cyano, a carboxyl, a formyl, an acetyl, a methoxycarbonyl, an ethoxycarbonyl, a carbamoyl, a methylcarbamoyl, an ethylcarbamoyl or an N,N-dimethylcarbamoyl group, or pharmaceutically acceptable salts thereof.

10. The isoxazole compound of Claim 1, wherein R² is a hydrogen, a halogen, a C₁ - C₆ alkyl, a C₂ - C₆ alkenyl or a C₂ - C₆ alkynyl group, or pharmaceutically acceptable salts thereof.

11. The isoxazole compound of Claim 1, wherein R² is a hydrogen, a halogen, a C₁ - C₄ alkyl, an allyl, an isopropenyl, a 2-butenyl or a propargyl group, or pharmaceutically acceptable salts thereof

12. The isoxazole compound of Claim 1, wherein R² is a hydrogen, a chlorine, an ethyl, a propyl, an isopropyl, an isobutyl or a t-butyl group, or pharmaceutically acceptable salts thereof.

13. The isoxazole compound of Claim 1, wherein R² is a hydrogen or an isopropyl group, or pharmaceutically acceptable salts thereof.

14. The isoxazole compound of Claim 1, wherein R³ is an amino, a mono C₁ - C₆ alkylamino, a di(C₁ - C₆ alkyl)amino or a 5- or 6-membered saturated heterocyclic group (provided that the group attached thorough a ring nitrogen atom) having one nitrogen atom and further optionally having one nitrogen atom or oxygen atom, or pharmaceutically acceptable salts thereof.

15. The isoxazole compound of Claim 1, wherein R³ is an amino, a

methylamino, an ethylamino, an N,N-dimethylamino, a piperidyl or a morpholinyl group, or pharmaceutically acceptable salts thereof.

16. The isoxazole compound of Claim 1, wherein R³ is an amino group, or pharmaceutically acceptable salts thereof.

17. The isoxazole compound of Claim 1, wherein X is an oxygen atom, or pharmaceutically acceptable salts thereof.

18. The isoxazole compound of Claim 1, wherein n is 2, or pharmaceutically acceptable salts thereof.

19. The isoxazole compound of Claim 1, wherein R¹ is a C₆ - C₁₄ aryl group optionally having from 1 to 3 substituents which may be the same as or different from each other and selected from the following substituent group, or a 5- or 6-membered aromatic heterocyclic group optionally having from 1 to 3 substituents and having one or two hetero atoms which may be the same as or different from each other and selected from the group consisting of nitrogen, oxygen and sulfur atoms [the substituent group is a halogen; a C₁ - C₆ alkyl; a C₁ - C₆ alkyl substituted with a halogen or a C₁ - C₆ alkoxy; a C₁ - C₆ alkoxy; a C₆ - C₁₄ aryl optionally having from 1 to 3 substituents which may be the same as or different from each other and selected from the following group (the substituent group is a halogen, a C₁ - C₆ alkyl or a C₁ - C₆ alkoxy); a cyano; a C₂ - C₇ alkoxy carbonyl; a carbamoyl; a mono-(C₁ - C₆ alkyl)carbamoyl; and a di(C₁ - C₆ alkyl)carbamoyl group],

R² is a hydrogen, a halogen, a C₁ - C₆ alkyl, a C₂ - C₆ alkenyl or a C₂ - C₆ alkynyl group, and

R³ is an amino, a mono C₁ - C₆ alkylamino, a di(C₁ - C₆ alkyl)amino group or a 5- or 6-membered saturated heterocyclic group (provided that the group attached through a ring nitrogen atom) containing one nitrogen atom and

further one nitrogen or oxygen atom,
or pharmaceutically acceptable salts thereof.

20. The isoxazole compound of Claim 1, wherein R¹ is a C₆ - C₁₄ aryl group optionally having from 1 to 3 substituents which may be the same as or different from each other and selected from the following substituent group, or a 5- or 6-membered aromatic heterocyclic group optionally having from 1 to 3 substituents and having one or two hetero atoms which may be the same as or different from each other and selected from the group consisting of nitrogen, oxygen and sulfur atoms [the substituent group is a halogen; a C₁ - C₆ alkyl; a C₁ - C₆ alkyl substituted with a halogen or a C₁ - C₆ alkoxy; a C₁ - C₆ alkoxy; a C₆ - C₁₄ aryl optionally having from 1 to 3 substituents which may be the same as or different from each other and selected from the following group (the substituent group is a halogen, a C₁ - C₆ alkyl or a C₁ - C₆ alkoxy); a cyano; a C₂ - C₇ alkoxy carbonyl; a carbamoyl; a mono(C₁ - C₆ alkyl)carbamoyl; and a di(C₁ - C₆ alkyl)carbamoyl group],

R² is a hydrogen, a halogen, a C₁ - C₆ alkyl, a C₂ - C₆ alkenyl or a C₂ - C₆ alkynyl group,

R³ is an amino group,

X is a oxygen atom, and

n is 2,

or pharmaceutically acceptable salts thereof.

21. The isoxazole compound of Claim 1, wherein R¹ is a C₆ - C₁₄ aryl group optionally having from 1 to 3 substituents which may be the same as or different from each other and selected from the following substituent group, or a 5- or 6-membered aromatic heterocyclic group optionally having one or two substituents and having one or two hetero atoms which may be the same as or different from each other and selected from the group consisting of nitrogen, oxygen and sulfur atoms [the substituent group is a halogen, a C₁ - C₄ alkyl, a

fluoromethyl, a difluoromethyl, a trifluoromethyl, a 2-fluoroethyl, a 2,2,2-trifluoroethyl, a methoxymethyl, a methoxyethyl, a C₁ - C₄ alkoxy, a phenyl, a 4-fluorophenyl, a 4-chlorophenyl, a 2,4-dichlorophenyl, a 4-methylphenyl, a 4-methoxyphenyl, a cyano, a methoxycarbonyl, an ethoxycarbonyl, a carbamoyl, a methylcarbamoyl, an ethylcarbamoyl, an N,N-dimethylcarbamoyl and an N,N-diethylcarbamoyl group],

R² is a hydrogen, a halogen, a C₁ - C₄ alkyl, an allyl, an isopropenyl, a 2-butenyl or a propargyl group,

R³ is an amino group,

X is an oxygen atom, and

n is 2,

or pharmaceutically acceptable salts thereof.

22. The isoxazole compound of Claim 1, wherein R¹ is a phenyl group optionally having from 1 to 3 substituents which may be the same as or different from each other and selected from the following substituent group, or a furyl, a thienyl or a pyridyl group optionally having one or two substituents [the substituent group is a halogen, a methyl, an ethyl, a trifluoromethyl, a methoxy, a phenyl, a cyano, a methoxycarbonyl, a carbamoyl, a methylcarbamoyl, an ethylcarbamoyl and an N,N-dimethylcarbamoyl group],

R² is a hydrogen, a halogen, a C₁ - C₄ alkyl, an allyl, an isopropenyl, a 2-butenyl or a propargyl group,

R³ is an amino group,

X is an oxygen atom, and

n is 2,

or pharmaceutically acceptable salts thereof.

23. The isoxazole compound of Claim 1, wherein R¹ is a phenyl group optionally having one or two substituents which may be the same as or different from each other and selected from the following substituent group, or a furyl, a thienyl or a pyridyl group optionally having one substituent [the substituent

group is a fluorine, a chlorine, a methyl, an ethyl, a trifluoromethyl and a methoxy group],

R^2 is a hydrogen, a chlorine, an ethyl, a propyl, an isopropyl, an isobutyl or a t-butyl group,

R^3 is an amino group,

X is an oxygen atom, and

n is 2,

or pharmaceutically acceptable salts thereof.

24. The isoxazole compound of Claim 1, wherein R^1 is a fluorophenyl, a chlorophenyl, a difluorophenyl, a dichlorophenyl, a methylphenyl, a 2-furyl, a 3-furyl, a 2-thienyl or a 3-thienyl group,

R^2 is a hydrogen, a chlorine, an ethyl, a propyl, an isopropyl, an isobutyl or a t-butyl group,

R^3 is an amino group,

X is an oxygen atom, and

n is 2,

or pharmaceutically acceptable salts thereof.

25. The isoxazole compound of Claim 1, wherein R^1 is a phenyl, a 2-chlorophenyl, a 4-chlorophenyl, a 2,4-difluorophenyl, a 2,4-dichlorophenyl, a 2-furyl or a 2-thienyl group,

R^2 is a hydrogen atom or a propyl group,

R^3 is an amino group,

X is an oxygen atom, and

n is 2,

or pharmaceutically acceptable salts thereof.

26. The isoxazole compound of Claim 1 selected from the group consisting of 3-(2-aminoethoxy)-5-phenylisoxazole, 3-(2-aminoethoxy)-4-chloro-5-phenylisoxazole,

3-(2-aminoethoxy)-4-ethyl-5-phenylisoxazole,
3-(2-aminoethoxy)-5-phenyl-4-propylisoxazole,
3-(2-aminoethoxy)-4-isopropyl-5-phenylisoxazole,
3-(2-aminoethoxy)-4-isobutyl-5-phenylisoxazole,
3-(2-aminoethoxy)-5-(2-chlorophenyl)-4-isopropylisoxazole,
3-(2-aminoethoxy)-5-(4-chlorophenyl)isoxazole,
3-(2-aminoethoxy)-5-(4-chlorophenyl)-4-isopropylisoxazole,
3-(2-aminoethoxy)-5-(2,4-dichlorophenyl)-4-isopropylisoxazole,
3-(2-aminoethoxy)-5-(2-furyl)-4-isopropylisoxazole,
3-(2-aminoethoxy)-5-(2-thienyl)isoxazole,
3-(2-aminoethoxy)-4-chloro-5-(2-thienyl)isoxazole,
3-(2-aminoethoxy)-4-isopropyl-5-(2-thienyl)isoxazole, and
4-allyl-3-(2-aminoethoxy)-5-phenylisoxazole,
and pharmaceutically acceptable salts thereof.

27. A pharmaceutical composition for a medicament in the prophylaxis or treatment of nervous diseases comprising an effective amount of an isoxazole derivative or pharmaceutically acceptable salts thereof as defined in any one of Claims 1 to 26.

28. A pharmaceutical composition for a medicament in the prophylaxis or treatment of depression comprising an effective amount of an isoxazole derivative or pharmaceutically acceptable salts thereof as defined in any one of Claims 1 to 26.


29. A use of an isoxazole derivative or pharmaceutically acceptable salts thereof as defined in any one of Claims 1 to 26 for the production of pharmaceutical preparations for treatment or prevention of nervous diseases.

30. A use of an isoxazole derivative or pharmaceutically acceptable salts thereof as defined in any one of Claims 1 to 26 for the production of pharmaceutical preparations for treatment or prevention of depression.

31. A method for the prophylaxis or treatment of nervous diseases, which method

comprises administering to a warm blood animals a composition comprising a pharmaceutically effective amount of an isoxazole derivative as defined in any one of Claims 1 to 26 or pharmaceutically acceptable salts thereof.

32. A method for the prophylaxis or treatment of depression, which method comprises administering to a warm blood animals a composition comprising a pharmaceutically effective amount of an isoxazole derivative as defined in any one of Claims 1 to 26 or pharmaceutically acceptable salts thereof.



(I)