

(19) **DANMARK**

(10) **DK/EP 2699698 T3**



(12) **Oversættelse af  
europæisk patentskrift**

Patent- og  
Varemærkestyrelsen

- 
- (51) Int.Cl.: **C 12 Q 1/68 (2006.01)** **C 12 N 15/11 (2006.01)** **C 12 P 19/34 (2006.01)**  
**C 12 Q 1/48 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2017-04-10**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2017-01-04**
- (86) Europæisk ansøgning nr.: **12773717.9**
- (86) Europæisk indleveringsdag: **2012-04-20**
- (87) Den europæiske ansøgnings publiceringsdag: **2014-02-26**
- (86) International ansøgning nr.: **US2012034589**
- (87) Internationalt publikationsnr.: **WO2012145725**
- (30) Prioritet: **2011-04-20 US 201161477357 P** **2011-04-20 US 201161477437 P**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **Mesa Biotech, Inc., 6181 Cornerstone Court , Suite 101, San Diego, California 92121-4727, USA**
- (72) Opfinder: **Cai, Hong, 2306 Canyon Glen Road, Los Alamos, NM 87544, USA**  
**COBB, Nathan, J., 403 N.Gregson Street, Durham, NC 27701, USA**
- (74) Fuldmægtig i Danmark: **PATRADE A/S, Fredens Torv 3A, 8000 Århus C, Danmark**
- (54) Benævnelse: **OSCILLERENDE AMPLIFIKATIONS-REAKTION FOR NUKLEINSYRER**
- (56) Fremdragne publikationer:  
**WO-A1-94/23055**  
**WO-A1-2010/105074**  
**WO-A2-2012/083189**  
**US-A1- 2001 019 825**  
**US-A1- 2005 136 443**  
**US-A1- 2007 231 798**  
**US-A1- 2010 203 532**  
**US-B1- 6 300 069**  
**IAKOBASHVILI R ET AL: "Low temperature cycled PCR protocol for Klenow fragment of DNA polymerase I in the presence of proline", NUCLEIC ACIDS RESEARCH, OXFORD UNIVERSITY PRESS, GB, vol. 27, no. 6, 1 March 1999 (1999-03-01), pages 1566-1568, XP002104331, ISSN: 0305-1048, DOI: 10.1093/NAR/27.6.1566**  
**KEMPITIYA A ET AL: "Localized microwave heating in microwells for parallel DNA amplification applications", APPLIED PHYSICS LETTERS, AMERICAN INSTITUTE OF PHYSICS, US, vol. 94, no. 6, 10 February 2009 (2009-02-10), pages 64106-64106, XP012119049, ISSN: 0003-6951, DOI: 10.1063/1.3078273**  
**ZHANG CHUNSUN ET AL: "PCR microfluidic devices for DNA amplification", BIOTECHNOLOGY ADVANCES, ELSEVIER PUBLISHING, BARKING, GB, vol. 24, no. 3, 1 May 2006 (2006-05-01), pages 243-284, XP002446769, ISSN: 0734-9750, DOI: 10.1016/J.BIOTECHADV.2005.10.002**

Fortsættes ...

**WEIGHARDT F ET AL: "A SIMPLE PROCEDURE FOR ENHANCING PCR SPECIFICITY", PCR METHODS & APPLICATIONS, COLD SPRING HARBOR LABORATORY PRESS, US, vol. 3, no. 1, 1 August 1993 (1993-08-01), pages 77-81, XP000609125, ISSN: 1054-9803**

**GOBINDA SARKAR ET AL: 'Formamide can dramatically improve the specificity of PCR' NUCLEIC ACIDS RESEARCH vol. 18, no. 24, 01 January 1990, GB, page 7465, XP055237925 DOI: 10.1093/nar/18.24.7465 ISSN: 0305-1048**

**BRAUN DIETER ET AL: 'Exponential DNA replication by laminar convection' PHYSICAL REVIEW LETTERS, AMERICAN PHYSICAL SOCIETY, US vol. 91, no. 15, 10 October 2003, pages 158103-1 - 158103-4, XP002609588 DOI: 10.1103/PHYSREVLETT.91.158103 ISSN: 0031-9007**

## DESCRIPTION

### REFERENCE TO A SEQUENCE LISTING, A TABLE, OR COMPUTER PROGRAM

[0001] Applicant hereby submits a sequence listing as a text file titled 041812\_ST25.txt created on April 20, 2012 having 10K bytes that is ASCII compliant and is incorporated herein by reference.

### BACKGROUND OF THE INVENTION

#### Field of the Invention (Technical Field):

[0002] Embodiments of the present invention relate to methods and apparatuses for template-dependent amplification of nucleic acid target sequences by oscillating reaction temperature in a small range, preferably no more than 20 °C during any given thermal polymerization cycle.

#### Background Art:

[0003] Note that the following discussion refers to a number of publications and references. Discussion of such publications herein is given for more complete background of the scientific principles and is not to be construed as an admission that such publications are prior art for patentability determination purposes.

[0004] Amplification of nucleic acids is among the most indispensable techniques in molecular biology, widely used in research, genetic testing, agriculture, and forensics. The most common amplification method is the polymerase chain reaction (PCR) in which the prevalence of a specific nucleic acid target sequence is increased exponentially in solution (See U.S. Pat. Nos. 4,683,195, 4,683,202, and 4,800,159). A PCR reaction employs two oligonucleotide primers that hybridize to opposite strands of the DNA double helix either upstream (5') or downstream (3') of the target sequence to be amplified. A (generally thermostable) DNA polymerase is used to extend hybridized primers in the 5' → 3' direction by adding deoxynucleoside-triphosphates (dNTPs) in order to 'copy' the target sequence and generate double-stranded DNA products. By cycling the temperature of the reaction mixture (typically 95°C Celsius), the two strands of DNA can be separated at high temperature allowing them to serve as templates for further primer binding and polymerization at lower temperatures (e.g. 55 °C and 60 °C). After repeating this process many times, a single target sequence can be amplified into billions of copies.

[0005] While PCR is the gold-standard amplification methodology in the well-equipped laboratory, it is rather complex, requiring both expensive and sophisticated thermal cycling instrumentation with active heating and cooling heating elements and precise temperature control, and trained technicians to gather meaningful results. For instance, most PCR reaction requires rapid and precise cycling between at least two temperatures (e.g. 95 deg and 57 deg), that typically results in the use of an expensive and energy-inefficient Peltier engine (thermal electric cooling mechanism) and precise temperature control elements. Such inherent limitations make PCR incompatible with the development of cost-effective, point-of-care nucleic acid diagnostics-useful where a supporting laboratory infrastructure is absent. In an effort to eliminate some of the resource-intensive requirements of PCR, various 'isothermal' amplification techniques have been developed in the past decades. In such reactions, nucleic acids may be amplified at a single temperature, removing the requirement of the costly thermocycler, and making them more amenable for use in low-cost diagnostic devices. Examples include nucleic acid sequence-based amplification (NASBA), helicase-mediated amplification, strand displacement amplification, loop-mediated isothermal amplification (LAMP) etc. However, these isothermal amplification methods often require 60-90 minute amplification time (due to slow kinetic enzymatic process in vitro) and precise temperature control at the single temperature point to accommodate the extremely stringent amplification reactions, again lacking the robustness and speed desired for the point of use diagnostic application.

[0006] Template-dependent nucleic acid amplification is the cornerstone of the nucleic acid-based molecular diagnostics. Robust, low cost, rapid, point-of-care nucleic acid diagnostics are a pressing need in health care, agriculture, and in the context of biological terrorism and warfare. However, the assay chemistry strategies associated with the existing PCR or isothermal amplification posse significant engineering and robustness limitations rendering such amplification approaches expensive and impractical for the resource-limited settings where nucleic acid-based molecular could make the most impact for emerging disease

prevention and control. Considerable improvements in nucleic acid amplification must yet be made to bring affordable and robust diagnostics to settings devoid of dedicated laboratory infrastructure.

**[0007]** Conventional PCR relies on highly specific and rapid thermal cycling, commonly varying temperature by as much as 40 °C. Such an amplification methodology requires expensive instrumentation in order to rapidly heat and (particularly) cool the PCR reaction mixture, in addition to accurately maintaining solution temperatures. Isothermal nucleic acid amplification procedures, while eliminating the need for complex thermal cycling instrumentation, are generally slow (at least 60 minute reaction time), unreliable, and require precise temperature calibration.

**[0008]** In a PCR thermal cycling process, a PCR cycler must have a good temperature control to maintains temperature uniformity within the sample and a typical sample heating (and/or cooling) rate of at least 2 °C per second. Temperature control is typically achieved by a feedback loop system, while temperature uniformity is achieved by highly thermally conductive but bulky materials such as copper. A high heating rate is accomplished by the implementation of a proportional integrated derivative (PID) control method limited by maximum dissipated power and heat capacitance. A high cooling rate is rather difficult to achieve and bulky systems require forced cooling by either a thermoelectric element (P.Wilding, M. A. Shoffner and L. J. Kricka, Clin. Chem., 1994, 40, 1815-1817.) (often called a Peltier element) or by other means, such as water (J. B. Findlay, S. M. Atwood, L. Bergmeyer, J. Chemelli, K. Christy, T. Cummins, W. Donish, T. Ekeze, J. Falvo, D. Patterson, J. Puskas, J. Quenin, J. Shah, D. Sharkey, J. W. H. Sutherland, R. Sutton, H. Warren and J. Wellman, Clin. Chem., 1993, 39, 9, 1927-1933 ). These PCR machines are complicated and power hungry devices. As the systems are bulky, their thermal time constants are in minutes rather than seconds which result in long transition times and unwanted by-products of the PCR. The high power consumption eliminates the possibility of making a battery operated and portable PCR system.

Document WO1994/023055 A1 recites a method for performing a cycled primer extension reaction by contacting a template DNA with a primer in the presence of sufficient glycerol or ethylene glycol to lower the melting temperature of template DNA and primer hybrids below 70 DEG C and a DNA polymerase under conditions in which the DNA polymerase can cause primer extensions and is stable to the temperature at which the reaction mixture is heated to denature the primer extension product from the template nucleic acid; and a kit suitable for use in cycle primer extension reaction including the necessary primers, buffers and enzymes required for the procedure, and glycerol.

**[0009]** With the recent advancement of silicon technology based micromachining and biological micro-electromechanical systems (bioMEMS), many groups around the world have started the development of microPCRs ( $\mu$ PCR), which are a central part of a lab-on-a-chip or micro total analysis systems ( $\mu$ TAS). Researchers follow two basic approaches: a stationary system with cycling temperature a flow system with three zones at different temperatures. Both systems have their advantages and disadvantages. Stationary systems cycle the temperature of the chamber in order to modify the temperature of the PCR solution. They do not require a pumping system or other means to move the PCR sample around. The flow-through systems typically have zones at three constant temperatures. Only the sample changes temperature by moving between zones of different temperatures. This type of PCR system is faster than the first one but it requires an implementation of a mechanism to move the sample around. In both cases, the heaters are integrated with the PCR system, so it is not economical to dispose the device to avoid cross-contamination after performing only a single test. The major advantages demonstrated by these two formats are reduced cycle time with the use of reduced sample volume compared to a conventional device. However, these PCR chips use substrate materials such as silicon that require the employment of expensive and sophisticated fabrication process, leading to a very high unit price. Furthermore, as a result of extreme small reaction volume ( $<\mu$ l) to achieve increased surface to volume ratio and the type of materials used in the  $\mu$ PCR chips, some effects not very common with the conventional PCRs become significant, including nonspecific adsorption of biological samples, inhibition, sample evaporation, and formation of bubbles. Other current effort also involves the development of a temperature cycling reaction microchip that integrates stationary chamber and continuous flow PCRs to perform efficient temperature cycling of the flow-through microchannel PCR chip while the flexibility of varying the-cycle number and the number of temperature zones in the stationary chamber PCR chip. However, the efficiency of the hybrid PCR device is still being validated and issues related to sample inhibition, adsorption, and bubble formation associated with such  $\mu$ PCR chip approach remain to poses significant stringency to all the upfront sample preparation/nucleic acid isolation process, and amplification reagents and reaction conditions e.g. ultra high polymerase concentration, PCR primer concentrations etc.

## **SUMMARY OF THE INVENTION**

**[0010]** The present invention is defined in the appended claims. One embodiment of the present invention provides for a method for amplifying a template of nucleic acid target sequence contained in a sample. The method includes contacting the sample with an amplification reaction mixture containing a primer complementary to the template of nucleic acid target sequence. A

temperature of the reaction is oscillated between an upper temperature and a lower temperature wherein the change in temperature is no greater than about 20°C during a plurality of temperature cycles. The template of nucleic acid target sequence is amplified.

**[0011]** One embodiment of the present invention provides that the change in temperature is no greater than about 15°C during a plurality of temperature cycles. Another embodiment provides that the change in temperature is no greater than about 10°C during a plurality of temperature cycles. Yet another embodiment provides that the change in temperature is no greater than about 5°C during a plurality of temperature cycles. The temperature may fluctuate by (+/-2°C) for a given temperature and/or range according to one embodiment of the present invention.

Another embodiment of the present invention provides that upon reaching the upper temperature or the lower temperature, the temperature is maintained for a set period of time within a temperature fluctuation. Alternatively, upon reaching an upper or lower temperature within the temperature range, the temperature is varied to the other temperature. In one embodiment, the lower temperature is no less than about 50°C. In another embodiment, the upper temperature is no greater than about 85°C. The upper and lower temperature may vary by about +/- 5°C according to one embodiment.

**[0012]** According to one embodiment of the present invention the template of nucleic acid target sequence may be single stranded DNA or RNA, double stranded DNA or RNA, RNA, DNA or any combination thereof. The length of the target nucleic acid may be less than 1000bp, less than 250bp, less than 150bp or less than 100bp.

**[0013]** One or more of the embodiments may comprise a pair of primers which bind to opposite strands of the template of nucleic acid. The pair of primers may have a length and a GC content so that the melting temperature is  $\geq 65^{\circ}\text{C}$ . In another embodiment, the pair of primers have a length and a GC content so that the melting temperature is  $\geq 70^{\circ}\text{C}$ . For example, each primer of the pair of primers independently has a length of between 35-70 base pairs. According to one embodiment of the present invention, the melting temperature of each primer of the primer pair is between 70-80°C. In a preferred embodiment, the pair of primers include a forward primer and a reverse primer each having a length of between 40-47 base pairs.

**[0014]** Yet another embodiment of the present invention provides a method for amplifying a template of nucleic acid target sequence contained in a sample. The method includes contacting the sample with an amplification reaction mixture comprising a primer or a primer pair having a length of between 35-70 base pairs and complementary to the template of the nucleic acid target sequence and wherein the melting temperature of each primer of the primer pair is between 70-80°C. The amplification reaction mixture also includes DMSO, monovalent cation, divalent cation, dNTPs, and DNA Polymerase. A temperature of the reaction is oscillated between an upper temperature and a lower temperature wherein the change in temperature is no greater than about 20°C during a plurality of temperature cycles and amplifying the template of nucleic acid target sequence. In a preferred embodiment, the divalent cation is selected from the group consisting of magnesium, manganese, copper, zinc or any combination thereof and the monovalent cation is selected from the group consisting of sodium, potassium, lithium, rubidium, cesium, ammonium or any combination thereof. In another preferred embodiment, the amplification reaction mixture comprises a nucleic acid destabilizing agent. In a more preferred embodiment the reaction comprises a DNA polymerase which may be a thermostable DNA polymerase. The DNA polymerase may be selected from the group consisting of TAQ DNA polymerase, VentR DNA polymerase, and DeepVentR DNA polymerase but is not limited thereto as other polymerases disclosed herein and known in the art may be included. The DNA polymerase may include a strand displacing activity. In another embodiment, the DNA polymerase does not have 3'-> 5' exonuclease activity. In another embodiment, the method for amplifying a template further comprises adding a reverse transcriptase and a DNA polymerase. For example, the reverse transcriptase is a thermostable reverse transcriptase. The reverse transcriptase may be selected from AMV-RT, Superscript II reverse transcriptase, Superscript III reverse transcriptase, or MMLV-RT but is not limited thereto as other reverse transcriptases known in the art may be used. Another embodiment of the present invention further comprises the addition of a single stranded binding protein to the reaction mixture as disclosed. For example, the single stranded binding protein is a thermal stable single stranded binding protein or the single stranded binding protein is a non-thermal stable single stranded binding protein.

**[0015]** Yet another embodiment of the present disclosure provides a mixture which includes a single strand or double strand nucleic acid destabilizing agent. For example dimethylsulfoxide (DMSO) or formamide but not limited thereto as other agents such as glycerol may be added for the same purpose.

**[0016]** Another embodiment of the present disclosure provides a method wherein the sample is not alcohol free and or the sample is not salt free.

**[0017]** Yet another embodiment of the present invention provides a method for amplifying a template of nucleic acid target sequence contained in a sample wherein the amplification reaction mixture comprises single strand or double strand nucleic acid

destabilizer; monovalent cation ;divalent cation; dNTPs and DNA Polymerase buffered at a pH to support activity.

**[0018]** Another embodiment of the present disclosure provides an amplification reaction mixture buffer comprising one or more of the following: single strand or double strand nucleic acid destabilizer; monovalent cation; divalent cation; dNTPs ;and DNA Polymerase buffered at a pH to support activity. The DNA polymerase may be a thermostable DNA polymerase. The DNA polymerase may be selected from the group consisting of TAQ DNA polymerase, VentR DNA polymerase, and DeepVentR DNA polymerase but not limited thereto. The DNA polymerase may have a strand displacing activity. The DNA polymerase may be selected which does not have 3'-> 5' exonuclease activity. The mixture may also include one or more of the following: a single stranded binding protein, a destabilizing agent is dimethylsulfoxide (DMSO) or formamide; a divalent cation which may be a salt selected from the group consisting of magnesium, manganese, copper, zinc or any combination thereof, and a monovalent cation which is a salt selected from the group consisting of sodium, potassium, lithium, rubidium, cesium, ammonium or any combination thereof.

**[0019]** Objects, advantages and novel features, and further scope of applicability of the present invention will be set forth in part in the detailed description to follow, taken in conjunction with the accompanying drawings, and in part will become apparent to those skilled in the art upon examination of the following, or may be learned by practice of the invention. The objects and advantages of the invention may be realized and attained by means of the instrumentalities and combinations particularly pointed out in the appended claims.

### **BRIEF DESCRIPTION OF THE DRAWINGS**

**[0020]** The accompanying drawings, which are incorporated into and form a part of the specification, illustrate an embodiment of the present invention and, together with the description, serve to explain the principles of the invention. The drawings are only for the purpose of illustrating a preferred embodiment of the invention and are not to be construed as limiting the invention. In the drawings:

FIG. 1. Illustrates a schematic diagram of an Oscillating PCR Amplification Reaction according to one embodiment of the present invention with panel A, a representative trace of the thermal fluctuations observed during several cycles of OPCRar (gray line), in comparison with a conventional two-stage PCR reaction (black line) and panel B illustrating a method for nucleotide amplification according to one embodiment of the present invention.

FIG. 2. Panels A-E represents a series of photos of acrylamide gels illustrating different polymerases used to generate a product of 153 base pair (bp) according to one embodiment of the present invention.

FIG. 3. Panels A-B represents a series of photos of an acrylamide gels illustrating an effect of ethanol on nucleic acid amplification according to one embodiment of the present invention.

FIG. 4. is a photo of an acrylamide gel illustrating differences in temperature of annealing and primer melting temperature to support efficient amplification according to one embodiment of the present invention.

FIG. 5. is a photo of an acrylamide gel illustrating an effect of hot-start DNA polymerase on primer dimer formation according to one embodiment of the present invention.

FIG. 6. is a photo of an acrylamide gel illustrating an effect of GC and AT clamps on primer-dimer formation according to one embodiment of the present invention.

FIG. 7. is a photo of an acrylamide gel illustrating an effect of single stranded binding protein on product formation according to one embodiment of the present invention.

FIG. 8. Panels A-B represent photos of acrylamide gels illustrating a reduction in the amount of primer-dimer formation by T4 gene protein 32 according to one embodiment of the present invention.

FIG. 9. is a photo of an acrylamide gel illustrating a specific target sequence present in double stranded DNA amplified according to one embodiment of the present invention.

FIG. 10. is a photo of an acrylamide gel illustrating amplification of a specific target sequence present in ssDNA according to one embodiment of the present invention.

FIG. 11. is a photo of an acrylamide gel illustrating amplification of a specific target sequence present in plasmid DNA according to one embodiment of the present invention.

FIG. 12. is a photo of an acrylamide gel illustrating amplification of a single stranded RNA according to one embodiment of the present invention.

FIG. 13. is a photo of an acrylamide gel illustrating amplification of a specific target sequence in bacterial genomic DNA according to one embodiment of the present invention.

FIG. 14. is a photo of an acrylamide gel illustrating amplification of a specific target sequence present in chloroplast DNA according to one embodiment of the present invention.

FIG. 15. is a photo of an acrylamide gel illustrating amplification of two target sequences according to one embodiment of the present invention.

FIG. 16 panel A-B represents photos of acrylamide gels illustrating amplification of a target sequence in the presence of SSB at lower melting temperatures according to one embodiment of the present invention.

FIG. 17 panel A-B represents photos of acrylamide gels illustrating amplification of a target with precise temperature control and/or rapid ramping parameters as required in a typical PCR thermocycler.

FIG. 18 is a photo of an acrylamide gel illustrating amplification of a target RbcL amplified in the low cost heater without ramping or precise temperature control.

### **DETAILED DESCRIPTION OF THE INVENTION**

**[0021]** As used throughout the specification and claims, the term '*nucleic acid*' means single stranded or double stranded DNA, RNA, or DNA/RNA hybrid molecules. Single stranded nucleic acids may have secondary structure such as hairpin, loop, and stem elements. Double stranded or single stranded nucleic acids may be linear or circular. Double stranded nucleic acids may be intact or nicked. Double stranded molecules may be blunt-ended or have single strand overhanging ends. Nucleic acid samples may be isolated from cells or viruses and may include chromosomal DNA, extra-chromosomal DNA including plasmid DNA, recombinant DNA, DNA fragments, messenger RNA, ribosomal RNA, transfer RNA, double stranded RNA or other RNAs that occur in cells or viruses. Nucleic acid may be isolated from any number of sources such as agriculture, food, environmental, fermentations, or biological fluids such as saliva, blood, nasal or lung aspirates, cerebrospinal fluid, sputum, stool, milk, swabs of mucosal tissues, tissue samples, or cells. Nucleic acid may be isolated, cloned or synthesized in vitro. Within the described nucleic acids above, individual nucleotides may be subject to modification or chemical alterations such as methylation. These modifications or alterations may arise naturally or by in vitro synthesis.

**[0022]** As used throughout the specification and claims, the terms '*target nucleic acid*' or '*template nucleic acid*' mean a single stranded or double stranded DNA or RNA fragment or sequence that is intended to be selectively amplified. The size of the nucleic acid to be amplified is defined by upstream (5') and downstream (3') boundaries and may be less than 500 bp, preferably less than 250 bp, and more preferably less than 150 bp and more preferably less than 100bp. The target nucleic acid may be a fragment contained within a longer double stranded or single stranded nucleic acid or may be an entire double stranded or single stranded nucleic acid.

**[0023]** As used throughout the specification and claims, the term '*duplex*' means a DNA or RNA nucleic acid molecule that is double stranded in whole or in part.

**[0024]** As used throughout the specification and claims, the term '*thermal cycle*' means the repeated temperature fluctuation necessary for nucleic acid amplification to occur. The thermal cycle may include, but is not limited to, a high temperature melting or denaturation step, and a low temperature annealing or hybridization step.

**[0025]** As used throughout the specification and claims, the terms '*melting*' or '*denaturation*' mean separating all or part of two complementary strands of a nucleic acid duplex with high temperature. The melting or denaturation temperature may be influenced by the length and sequence of the oligonucleotide primer, the concentration of duplex destabilizing reagents such as DMSO and formamide, and the ionic strength or pH of the solution.

**[0026]** As used throughout the specification and claims, the terms '*annealing*' or '*hybridization*' mean the sequence-specific binding of an oligonucleotide primer to a single-stranded nucleic acid template. The primer may bind only to its complementary sequence on one of the template strands and no other region of the template. The specificity of annealing or hybridization may be

influenced by the length and sequence of the oligonucleotide primer, the temperature at which binding is performed, the concentration of duplex destabilizing reagents such as DMSO and formamide, and the ionic strength or pH of the solution.

**[0027]** As used throughout the specification and claims, the term '*primer*' means a single stranded nucleic acid or oligonucleotide capable of binding to a single stranded region on a target nucleic acid in a sequence-specific manner that allows polymerase-dependent replication of the target nucleic acid.

**[0028]** As used throughout the specification and claims, the term '*OPCRar*' means Oscillating PCR Amplification Reaction which is an in vitro technique for amplifying nucleic acids using variations in temperature less than the typical amplification techniques, for example less than 20 °C, preferably less than 15 °C, and more preferably less than 10 °C between the denaturation temperature and the annealing temperature.

**[0029]** As used throughout the specification and claims, the term '*accessory protein*' refers to any protein capable of stimulating activity, for example, a thermostable single stranded binding protein (SSB), for example rec A or RPA (Replication Protein A but not limited thereto).

**[0030]** In an embodiment of the invention, a method is provided for exponentially amplifying a specific nucleic acid target by thermal cycling where temperature variation is preferably less than 20 °C, more preferably less than 15 °C, and even more preferably less than 10 °C. This includes the steps of providing a single-stranded template of the nucleic acid to be amplified, oligonucleotide primers for hybridization to the nucleic acid template, using the hybridized oligonucleotide primers to synthesize a double-stranded extension product which is complementary to the template strand by means of a DNA polymerase, and repeating of the above steps to exponentially amplify a select nucleic acid target.

**[0031]** Referring now to **figure 1A**, a schematic diagram of an Oscillating PCR Amplification Reaction (OPCRar) according to one embodiment of the present invention is illustrated with panel A showing a representative trace of the thermal fluctuations observed during several cycles of OPCRar (gray line), in comparison with a conventional two-stage PCR reaction (black line). Note the dramatic reduction in the temperature variation in OPCRar. **Figure 1B** illustrates, double stranded target nucleic acid enters the melt stage where, depending on the temperatures, may result in either partial or complete denaturation of the duplex according to one embodiment of the present invention. Unwinding of the duplex begins at the ends of the target and single stranded nucleic acid is bound and stabilized by single stranded binding protein (circles). The reaction is cooled and enters the hybridization/polymerization stage where primers hybridize in a specific manner to the 5' ends of each strand of the target duplex. After primer hybridization, DNA polymerase (squares) binds to the template/primer duplex and extends the primer in the 5' → 3' direction by incorporation of dNTPs, copying the template stand of DNA. If the polymerase used has strand displacement activity, it will be able to displace the opposing strand in the partially denatured complex. Upon generation of new duplex DNA, the thermal cycle is repeated many times to result in exponential amplification of the target nucleic acid sequence.

**[0032]** In additional embodiments of the invention, thermal cycling involves temperature oscillation or cycling between two temperatures with a  $\Delta T$  of preferably no more than 20 °C, more preferably no more than 15 °C, and even more preferably less than 10 °C. The higher of the two temperatures may be sufficient to denature the double stranded target DNA, or preferably result in only partial denaturation of the double stranded DNA target. Upon reaching either the high or low temperature, said temperature is maintained for a set period of time or, preferably, immediately varied to the other temperature.

**[0033]** In additional embodiments of the invention, the nucleic acid target may be a double stranded nucleic acid such as double stranded DNA, or a single stranded nucleic acid such as single stranded RNA or DNA. If the target nucleic acid is double stranded, it must be denatured either entirely or partially by heat, or enzymatically, to form a single stranded template or template region necessary for DNA polymerase activity and amplification. The length of the target nucleic acid may be less than 1000 bp, preferably less than 250 bp, and more preferably less than 150 bp.

**[0034]** In additional embodiments of the invention, the oligonucleotide primers used for target nucleic acid amplification are a pair of primers which bind to opposite strands of a specific double stranded target nucleic acid, where one primer binds upstream at the 5' end, and one primer binds downstream at the 3' end of the target. Under multiplexing conditions, more than one oligonucleotide primer pair may be used to simultaneously amplify multiple nucleic acid targets in the same reaction mixture. The oligonucleotide primers may have a length and GC content so that the melting temperature is greater than 65 °C under universally accepted PCR buffer conditions, preferably greater than 70 °C.

**[0035]** In additional embodiments of the invention, the DNA polymerase used is preferably selected from Taq DNA polymerase, VentR DNA polymerase, DeepVentR DNA polymerase, and similar thermostable DNA polymerases. Preferably, the DNA



polymerase possesses a strand-displacing activity and does not contain a 3' → 5' exonuclease activity (see **figure 1B**). In addition, if the template nucleic acid is single stranded RNA, the reverse transcriptase used will be selected from AMV-RT, Superscript II reverse transcriptase (Invitrogen, Carlsbad, CA), Superscript III reverse transcriptase (Invitrogen), and similar thermostable enzymes.

## **Other - Thermophilic polymerase possibilities**

### **[0036]**

Thermophilic DNA Polymerase

Polymerase and (Vender)

VentR® (NEB)

VentR (exo-)® (NEB)

Deep Vent (NEB)

Deep VentR (exo-) (NEB)

Tag (NEB)

PyroScript (Lucigen)

PyroPhage® 3173, Wildtype (Lucigen)

LongAmp Tag (NEB)

Bst Polymerase

Phire Hot Start II (NEB)

Phusion High Fidelity DNA Polymerase (NEB)

Phusion (NEB)

Phusion® Flash (NEB)

9 Nm (NEB)

DyNAzyme II Hot Start (NEB)

DyNAzyme EXT (NEB)

DreamTag (Fermentas)

Tag (native) (Fermentas)

Maxima® Hot Start Tag (Fementas)

Pfu (recombinant), (Fermentas)

Bsm (large fragment), (Fermentas)

TrueStart™ Hot Start Tag (Fermentas)

Tfi (invitrogen)

AmpiTag® (Invitrogen)

AmpliTag Gold® (Invitrogen)

Platinum ® Pfx

[0037] In additional embodiments of the invention, the reaction mixture preferably comprises a single stranded binding protein (SSB) such as T4 gene 32 protein, or thermal stable SSB isolated and cloned from a thermophilic organism.

[0038] Additionally, the enzyme preparation includes a single or double stranded nucleic acid destabilizing agent such as dimethylsulfoxide (DMSO) or formamide, preferably at a concentration of 8-15% of the total reaction volume. Alternatively other reagents such as glycerol deaza-dGTP, 3dazopurein, dITP may be utilized alone or in combination with each other or the prior list of agents.

[0039] Embodiments of this invention are ideally suited for use in low cost, point-of-care nucleic acid diagnostics in which a microfluidic layer is positioned over a heating element. By reducing temperature range cycling requirements, relatively simple heating with passive cooling mechanisms can be used to rapidly cycle temperature of a reaction solution. Lower maximum temperatures during thermal cycling minimize fluid evaporation which may negatively impact the overall amplification reaction. More importantly, the robustness of the amplification is greatly improved comparing to the conventional PCR process giving the new method is able to accommodate temperature fluctuation (imprecise temperature control) during an amplification process. The specific reaction chemistry of the invention was shown to work over a wide range of melting (e.g. 70-105 °C, essentially insensitive to bubbling) and hybridization temperatures eliminating the need for uniform temperature throughout the entire reaction volume. Finally, embodiments of the invention perform well in the presence of alcohol, and salt (e.g. ~10% ethanol), greatly reducing the stringency of up-front nucleic acid isolation methodologies through the elimination of a centrifugation, heat-dry or vacuum step between alcohol-based washing (e.g. ethanol or isopropanol) and nucleic acid elution step involved conventional nucleic acid extraction chemistry.

[0040] Embodiments of this invention include the detection of pathogens in a biological sample where a nucleic acid of the pathogen is the target nucleic acid. Alternatively, the invention may be used to detect differences in chromosomal DNA, where a fragment of chromosomal DNA is the target nucleic acid. In this way, single nucleotide polymorphisms may be detected in the target nucleic acid from the same or different sources.

[0041] Embodiments of the amplification technology of the present invention are referred to as an 'Oscillating PCR Amplification Reaction' (OPCRar). OPCRar is based upon, but not limited to, the combined use of double stranded destabilizing agents which lower the reaction melting temperature, and oligonucleotide primers of unusually high melting temperature ( $T_m$ ) to raise the annealing temperature in a given thermal cycle. In this way, in vitro amplification of a target nucleic acid may be performed by rapid thermal cycling between temperatures preferably differing by 20 °C, more preferably less than 15 °C, and even more preferably less than 10 °C (**figure 1A**). Oligonucleotide primers specific for upstream (5') and downstream (3') regions of the target nucleic acid hybridize to the template allowing for extension by DNA polymerase to amplify the target. If the DNA polymerase used is a strand displacing polymerase without exonuclease activity, complete thermal denaturation of the double stranded target nucleic acid is unnecessary, working in conjunction with duplex destabilizing agents to lower the required melting temperature. The temperature cycling process is repeated and results in exponential amplification of the specific nucleic acid target sequence (**figure 1B**). In OPCRar, double stranded target nucleic acid enters the melt stage where, depending on the temperatures, may result in either partial or complete denaturation of the duplex. Unwinding of the duplex begins at the ends of the target and single stranded nucleic acid is bound and stabilized by single stranded binding protein (circles). The reaction is cooled and enters the hybridization/polymerization stage where primers hybridize in a specific manner to the 5' ends of each strand of the target duplex. After primer hybridization, DNA polymerase (squares) binds to the template/primer duplex and extends the primer in the 5' → 3' direction by incorporation of dNTPs, copying the template strand of DNA. If the polymerase used has strand displacement activity, it will be able to displace the opposing strand in the partially denatured complex. Upon generation of new duplex DNA, the thermal cycle is repeated many times to result in exponential amplification of the target nucleic acid sequence.

[0042] The OPCRar method is based upon, but not limited to, the combined use of nucleic acid destabilizing agents which lower the reaction melting temperature, and two oligonucleotide primers of unusually high melting temperature ( $T_m$ ) to raise the annealing temperature during thermal cycling. For a given target nucleic acid, one oligonucleotide primer preferably hybridizes to the 5'-end of the sense strand containing the target sequence, and one primer preferably hybridizes to the 5'-end of the anti-sense strand containing the reverse-complementary target sequence. OPCRar preferably utilizes, but is not limited to, the use of a strand displacing DNA polymerase without exonuclease activity to further lower the melting or denaturation temperature necessary for efficient target nucleic acid amplification. OPCRar may amplify a target nucleic acid in the presence or absence of an accessory protein. Any specific OPCRar system may be optimized by addition, subtraction, or substitution of components within the mixture.

[0043] This amplification technology has improved characteristics over other amplification methodologies reported in prior art in

the context of low-cost, rapid, point-of-care nucleic acid diagnostics. Unlike the above described nucleic acid amplification methodologies, the OPCRar system, enabled by its robust enzymatic process is robust, fast and tolerant to temperature fluctuations of a low cost heating device, thus ideally suited for low-cost, point-of-care applications. By minimizing the temperature differentials encountered during thermal cycling, OPCRar combines the speed and reliability of PCR with the lowered instrumentation requirements of isothermal amplification methodologies.

**[0044]** The OPCRar system's simplified thermal cycling requirement is ideally suited for passive-cooling instrumentation, where heat may be applied to one surface of a chamber and cooling occurs through heat dissipation to the atmosphere on the opposing surface. Such passive-cooling dramatically lowers the cost and complexity of any nucleic acids diagnostics device. Passive-cooling has been previously reported for use in diagnostics devices, however, these devices have employed conventional PCR cycling assay chemistry to amplify target nucleic acids limiting the rate of reaction (Luo et. al. *Nuc Acids Res.* 2005; Wilding et. al., *Nuc. Acids. Res.* 1996; Burke et. al., *Science* 1998;). Another advantage of OPCRar is that efficient nucleic acid amplification may occur over a wide range of melting and annealing temperatures and consequently requires less stringent temperature control mechanism. In the construction of miniaturized nucleic acid diagnostics, maintenance of uniform temperature throughout the entire reaction volume can be challenging, with a particularly high temperature gradient occurring between the heated and unheated sides of the reaction chamber. Such temperature variation may result in inefficient amplification using conventional PCR or isothermal reaction chemistries. OPCRar, through the use of the combination of robust polymerase, destabilizing reagent and other polymerase accessory factors, is designed to minimize problems associated with precise temperature regulation and maintenance; so long as the coolest regions of the reaction chamber observe the minimal possible melt temperature and the maximal possible annealing temperature for a given nucleic acid target the reaction will progress efficiently, even if other regions of the reaction volume vary by >10 °C. Moreover, the robust OPCRar nature of amplification chemistry along with the minimal power/energy consumption enables rapid and efficient amplification reaction at much large volume (e.g. 20 µl instead sub µl in a typical µPCR chips) greatly relaxed the stringency of the up-front sample preparation/nucleic acid isolation process (in terms of obtain sub µl of input template that is both highly concentrated and ultra-pure nucleic acid free of any trace contaminant e.g. salt and ethanol carry over and inhibitory substance) and requirement of ultra-high concentration and ultra-pure of PCR enzymes and bioagents.

#### **Solvent Reagents**

**[0045]** Solvents such as DMSO and formamide are known to lower the melting temperature of duplex nucleic acids by ~0.5-0.7 °C for every 1 % volume added. These reagents are often used to improve amplification efficiency of target nucleic acids containing high GC content and, thus, high T<sub>m</sub> to facilitate complete denaturation of difficult-to-melt double stranded templates. Commonly, PCR thermal cycling temperatures are kept constant upon incorporation of duplex destabilizing agents into PCR reactions. In contrast, OPCRar preferably utilizes the addition of uniquely high concentrations of DMSO to dramatically lower the melting temperature of the thermal cycle. In conventional PCR, DMSO is rarely used above 10% v/v due to the loss of polymerase activity associated with high concentrations of these reagents in conjunction with the high temperatures (generally greater than 90 °C) of the melting stage. The OPCRar system and method, on the other hand, preferably uses DMSO concentrations between 10 and 15%. Unexpectedly, this amount of DMSO does not produce significant loss of polymerase activity.

**[0046]** Referring now to **figure 2**, amplification according to one embodiment of the present invention is compatible with a variety of DNA polymerases, depending on the conditions. Ribonucleic acid isolated from nasal aspirate containing Influenza A virus (0.3 ng/µL) was used as a nucleic acid template under multiple Reverse Transcription-OPCRar conditions using VentR polymerase. The OPCRar primers FP3 and RP4 were used to generate a product of 153 bp which was visualized by electrophoresis on a 12% polyacrylamide gel stained with ethidium bromide. All reactions contained Superscript III Reverse Transcriptase, where an initial cDNA generation stage was performed for 5 minutes at 55 °C prior to thermal cycling. Gel lanes are labeled with concentration of DMSO, and the melt and hybridization stage temperatures used; reactions were cycled 40 times.

**[0047]** The conventional PCR enzyme, e.g. Taq DNA polymerase reaction mixture is extremely sensitive to any trace amount of alcohol, e.g. ethanol, whereas in one embodiment of the invention the novel reaction mixture is exceedingly resistant to inhibition by ethanol. Referring now to **figure 3**, ethanol effect on nucleic acid amplification according to one embodiment of the present invention is demonstrated. Total nucleic acid isolated from *Candidatus Liberibacter asiaticus*-infected leaf tissue (3.4 ng/µL) was used as a starting template. Reactions were performed either under OPCRar conditions using VentR (exo-) DNA polymerase, Et SSB, and the primers *hyv1\_For* and *hyv1\_Rev*, in the presence of 15% DMSO (**figure 3A**), or conventional PCR conditions using Taq polymerase and the primers HLBas-P2 and HLBs-P1 (**figure 3B**). OPCRar solutions were heated at 85 °C for 2 minutes to denature the template and then cycled 40 times, oscillating between 76 °C for 10 seconds, and 60 °C for 10 seconds to generate a product of 139 bp. Conventional PCR reactions were heated to 95 °C for 2 minutes and then cycled 40 times, varying between

95 °C for 10 seconds and 58 °C for 40 seconds to generate a product of 130 bp. The amplified products were visualized on a 12% acrylamide gel, stained with ethidium bromide. Ethanol concentrations included in the amplification reaction mixture are shown. It is evident that the OPCRar formulation (**figure 3A**) is dramatically more resistant to inactivation by ethanol as compared to conventional PCR (**figure 3B**).

**[0048]** Use of VentR(exo-) DNA polymerase and Et SSB under typical OPCRar conditions results in no loss of activity in up to 10% ethanol. This is a significant yet surprising discovery regarding the application of OPCRar to low cost point-of-care devices. Since the conventional wisdom of PCR and isothermal amplification typically advise users to provide the highly purified nucleic acid input that is free of alcohol and salt. As the result, almost all the researchers are employing some kind of vacuum dry, air dry, spin down or heating steps between the alcohol-based washes and eluting the target nucleic acid from nucleic acid- affinity microbeads, glass frit, matrix or filter etc before they run the sample through the PCR amplification process. For instance of an integrated point of care diagnostic device, in addition to amplification and detection of nucleic acids, the devices must also rapidly isolate target nucleic acids. Generally this is performed by binding nucleic acid to a glass fiber matrix and washing in the presence of significant concentrations of salt and ethanol, subsequently eluting in buffer containing minimal salt and no ethanol. Before elution, wash buffer retained on the binding matrix is removed to prevent carry-over to the elution volume; in commercial nucleic acid isolation kits this is typically performed by centrifugation. The specific enzyme mixture of OPCRar eliminates this need for careful removal of ethanol during nucleic acid isolation, making this embodiment of the invention tailored for low cost integrated diagnostics that does not require a vacuum, centrifuge, air dry or heating dry component.

### **Primers**

**[0049]** Oligonucleotide primers as described here can be synthesized and purified by methods known in the art. (See, for example U. S. Patent No. 6,214, 587). In present embodiments, two sequence-specific primers, representing a primer pair are used to exponentially amplify a target nucleic acid sequence. The first primer hybridizes to the upstream 5' region of the target nucleic acid, and the second primer hybridizes to the downstream, 3' region of the target sequence. The primers hybridize to the 5' end of one strand present in the target duplex, and the primers are extended by a polymerase in a 5' to 3' direction using the target nucleotide sequence as a template (**figure 1B**). Conditions of hybridization are standard as described in "Molecular Cloning and Laboratory Manual", 2nd ed. Sambrook, Rich and Maniatis, pub. Cold Spring Harbor (2003). To achieve specific amplification of a given target sequence a homologous primer is preferred, where every nucleotide in the primer is complementary to the target sequence. Primers may, however, include bases that are non-homologous with respect to the template nucleic acid, or 5' sequences which are non complementary to the target nucleotide sequence(s). Multiple pairs of primers can be used in a single OPCRar experiment in order to amplify multiple nucleic acid targets simultaneously in the same reaction mixture. So-called multiplexing is a commonly used technique in single nucleotide polymorphism analysis, in detection of pathogens, and for incorporation of internal controls into an individual reaction. Higher level of multiplexing may also be achieved through a use of 5' universal tag sequence introduced to each target-specific 3'-region that allows the universal amplification of all the target sequences with different internal pathogen sequences.

**[0050]** Oligonucleotide primer design involves several parameters such as melting temperature and intra- and inter-primer sequence alignment. Melting temperature is governed by factors such as the primer length and GC content. Inter-primer sequence complements can result in hairpin structures, which can impede efficient amplification, whereas intra-primer homology can result in unwanted amplification products dubbed primer-dimers. When designing a primer, it is important to select a sequence within the target which is specific to the nucleic acid molecule to be amplified and will minimally interact with either itself or other primers present in the amplification reaction.

**[0051]** In most nucleic acid amplification strategies, the melting temperature of a primer is preferably about 10 to 30°C higher than the temperature at which the hybridization and amplification takes place. With the temperature of the annealing/polymerization stage(s) being 55-60 °C in a PCR reaction, primers are typically 18-30 base pairs in length. This specific oligonucleotide length is minimized to allow for easy primer binding without loss of sequence specificity. In the OPCRar system, however, primers are preferably designed to be unusually long at 35-55 base pairs, with a melting temperature preferably between 70-80 °C in order to raise the temperature of the annealing stage. Considering the levels of the duplex destabilizing agent, DMSO, used in a typical OPCRar reaction (~10-15%), the calculated T<sub>m</sub> of OPCRar primers is preferably only <10 °C above the annealing temperature used during thermal cycling. In experiments and with the extreme length of OPCRar primers, efficient amplification occurs despite a minimal difference in primer T<sub>m</sub> (compensating for the concentration of DMSO) and the annealing/elongation temperature.

**[0052]** Referring now to **figure 4**, OPCRar primers require minimal difference between annealing stage temperature and primer

$T_m$  to support efficient amplification according to one embodiment of the present invention. A plasmid containing the *hyvI* gene sequence (12 ng/ $\mu$ L) was amplified using the primers *hyvI*\_For and *hyvI*\_Rev to generate a product of 139 bp, which was visualized by electrophoresis on a 12% acrylamide gel stained with ethidium bromide. Following an initial 2 minute 85 °C melt step, all reactions were cycled 40 times, 10 seconds at each of the indicated melt and hybridization temperatures. The calculated  $T_m$  for primers *hyvI*\_For and *hyvI*\_Rev are 72.2 °C and 70.9 °C, respectively. The reaction was performed in 10% DMSO, lowering the effective  $T_m$  by 7 °C, assuming a reduction of 0.7 °C per 1% volume. Even with a negligible difference between primer  $T_m$  and hybridization temperature, the amplification reaction is observed to proceed as efficiently as if the temperature difference is much lower.

**[0053]** Referring now to **figure 5**, the effect of hot-start DNA polymerase on primer dimer formation using OPCRar. Control reactions containing no nucleic acid template were performed with Superscript III RT, and either Taq or Platinum Taq DNA polymerase in the presence of primer pair FP3 and RP4 (8  $\mu$ M each) and 15 % DMSO. The cycling parameters were 55 °C for 5 minutes, 85 °C for 2 minutes, and 40 cycles of 80 °C for 15 seconds and 65 °C for 15 seconds. After electrophoresis on a 12% acrylamide gel, OPCRar products were visualized by staining with ethidium bromide. Platinum Taq is a commercially available hot-start enzyme (Invitrogen, Carlsbad, CA) that is conjugated to an antibody which dissociates upon heating the reaction solution to 94 °C under normal PCR conditions. In the presence of template nucleic acid, this primer pair will produce a product of 153 bp. It is clear that this hot-start preparation is no better than conventional Taq in reducing the formation of <110 bp primer dimers during OPCRar. One complication of long primers used in the OPCRar reaction is that they are more prone to result in non-specific and unwanted amplification products known as primer dimers. Primer dimers are formed when the 3' ends of primer oligonucleotides transiently bind one another during the initial increase of temperature at the start of an amplification reaction. During this critical time period, DNA polymerase may extend these transient complexes resulting in products that compete with specific target amplification during thermal cycling, particularly if the starting template nucleic acid concentration is very low. A commonly employed technique to reduce primer dimer formation during PCR is to utilize so-called 'hot-start' DNA polymerases. These commercially available enzymes are non-covalently bound to an inhibitory molecule such as an antibody. When the reaction temperature increases above 90 °C, the inhibitory molecule dissociates freeing the polymerase to perform formally. However, surprisingly, in our hands the hot-start enzyme Platinum Taq DNA Polymerase (Invitrogen, Carlsbad, CA) failed to appreciably reduce the abundance of primer dimer amplification, indicating that this commonly used methodology is insufficient for OPCRar, indicating that OPCRar process is a much more efficient amplification process than traditional PCR that enables dimer formation originated from transient kinetic collisions of homo and hetero-dimers that don't typically occur in the conventional PCR process.

**[0054]** Referring now to **figure 6**, a gel illustrating the effect of GC and AT clamps on primer dimer formation during OPCRar according to one embodiment of the present invention is presented. Primers designed to amplify the elongation factor gene of *C. Liberibacter asiaticus* were used in the absence of a starting template (3.5 ng/ $\mu$ L) to determine the propensity for primer dimer formation. In the presence of template the product sizes for the various primer sets would range from between 140-155 bp. Primer sequences can be seen to the right, with the AT or GC clamps in gray. For 'mod' primer sets; each primer contains a non-homologous base with respect to the template (underline), that increases homology with the second primer. All reactions were performed using VentR(exo-) DNA polymerase, Et SSB, in the presence of 15% DMSO. Solutions were heated at 85 °C for 2 minutes to denature the template and then cycled either 30 or 40 times, oscillating between 80 °C for 15 seconds, and 65 °C for 15 seconds. The amplified products were visualized on a 12% acrylamide gel, stained with ethidium bromide. As is clearly observed, GC clamp primer sets result in significant primer dimer formation while AT clamp primers result in no primer dimer formation.

**[0055]** In order to minimize the potential for primer dimer formation, OPCRar primers may be designed to employ several strategies differing from those used to generate conventional PCR primers. First, PCR primers generally possess a GC rich 3' end called a 'GC clamp', which results in greater specific binding to the target sequence. In OPCRar primers, however, it has been observed that a high GC content in the 3' region of the primer results in greater primer dimer formation, thus, OPCRar primers are made to contain AT rich 3' regions to energetically reduce the affinity of 3'-3' primer interactions resulting in these unwanted amplification products (**figure 6**). A second strategy for OPCRar primer design is to design primers that contain complementary 5' or internal sequences of at least 5 consecutive nucleotides. Oligonucleotides designed in this way will steer any primer hybridization during the initial increase in reaction temperature towards duplex structures that are not competent for polymerase extension. If suitable complementary sequences cannot be found within the target nucleic acid sequence, non-homologous, or mutated, bases can be used to generate them within the OPCRar primers. The exceptional length of OPCRar primers overcomes minor mispairing between primer and target during the early cycles of an amplification reaction. Primer sets are

EU AT

Forward

GTTCTTGTAG CGTTGCAGTC TTCTGCGGAA GATAAGGAAT TGCTTT (SEQ ID NO 21)

Reverse

GGGCACGTTT ATTAGCAACA ATAGAAGGAT CAAGCATCTG CACAGAAAT (SEQ ID NO 22)

EU GC

Forward

CTTGTAGCGT TGCAGTCTTC TGC GGAAGAT AAGGAATTGC TTTCTGCG (SEQ ID NO 23)

Reverse

CACGTTTATT AGCAACAATA GAAGGATCAA GCATCTGCAC AGAAATCACCG (SEQ ID NO 24)

EU-Atmod

Forward

GGTGTTCTTG TATCGTTGCA GTCTTCTGCG GAAGATAAGG AATTGCTTT (SEQ ID NO 25)

Reverse

GTAATGGGCA CGTTTATTAG CAACGATAGA AGGATCAAGC AACTGCACAG AAAT (SEQ ID NO 26)

EU GCmod

Forward

CTTGTATCGT TGCAGTCTTC TGC GGAAGAT AAGGAATTGC TTTCTGCG (SEQ ID NO 27)

Reverse GGCACGTTTA TTAGCAACGA TAGAAGGATC AAGCATCTGC ACAGAAATCA CCG (SEQ ID NO 28)

**[0056]** OPCRar primers may include any of the deoxyribonucleotide bases adenine "A", thymine "T", guanine "G" or cytosine "C" and/or one or more ribonucleotide bases, A, C, uracil "U", G. Furthermore, OPCRar primers may contain one or more modified deoxyribonucleotide or ribonucleotide bases where the modification does not prevent primer hybridization to the target nucleic acid, primer elongation by polymerase, or denaturation of duplex nucleic acid. OPCRar primers may be modified with chemical groups such as methylphosphonates or phosphorothioates, with non-nucleotide linkers, with biotin, or with fluorescent labels such as the amine-reactive fluorescein ester of carboxyfluorescein. Such modifications may enhance primer performance or facilitate the detection and characterization of amplification products.

## Polymerases

**[0057]** After single stranded template nucleic acid region has hybridized with a primer during OPCRar, a polymerization step occurs. If the target nucleic acid is DNA, a DNA polymerase is selected which acts on the target to extend the hybridized primers along the nucleic acid template in the presence of the four dNTP nucleotide bases to form a double stranded product where the newly synthesized strand is complementary to the nucleotide sequence of the template (**figure 1**). If the initial target is RNA, a reverse transcriptase is first used to copy the RNA template into a cDNA molecule, which is further amplified during OPCRar by a DNA polymerase.

**[0058]** A variety of DNA polymerases may be selected for OPCRar on the basis of thermostability and processivity, especially in the presence of the destabilizing agent, and alcohol (**figure 2**). Although not required, polymerases displaying strand displacement activity and lacking an exonuclease activity are found to significantly improve OPCRar reactions (**figure 2**). Examples of suitable DNA polymerases include Taq polymerase, KlenTaq DNA polymerase (AB Peptides, (St Louis, MO)), Bst DNA polymerase Large fragment (New England Biolabs, Beverly, MA), VentR or VentR (exo-) (New England BioLabs), DeepVentR or DeepVentR (exo-) (New England BioLabs), and similar enzymes. Suitable thermostable reverse transcriptases include Superscript II (Invitrogen, Carlsbad, CA), Superscript III (Invitrogen), and similar enzymes. It should be noted that the published conventional PCR amplification polymerase mixture fail to perform OPCRar due to the unique robustness requirement of OPCRar amplification. All the selected polymerase and bioreagent components should be carefully evaluated and experimentally tested before use.

## Single-stranded Binding Proteins

[0059] The OPCRar system preferably minimizes the temperature differential between melting and annealing thermal cycling stages, where this temperature differential is lowest if complete denaturation of duplex nucleic acid is unnecessary. While a strand-displacing DNA polymerase is helpful in this regard, accessory proteins may be used to further lower the thermal requirements for efficient amplification. Single-stranded binding proteins (SSBs) are known to stabilize single stranded nucleic acid to prevent the annealing of double stranded duplex formation, and have been shown to increase the efficiency of nucleic acid amplification reactions. The addition of a thermostable SSB to OPCRar methods according to an embodiment of the present invention is found to result in improved activity (**figure 7**). As an example, Et SSB (BioHelix Corporation, Beverly, MA), although the choice of SSB is not limited to a specific protein and may include SSBs isolated and cloned from a thermophilic organism, or engineered from a non-thermostable precursor SSB.

[0060] Referring now to **figure 7**, a gel showing the effect of single stranded binding protein on OPCRar product formation according to one embodiment of the present invention is illustrated. A Universal Influenza A single stranded DNA template (1E6 copies/ $\mu$ L, Biosearch Technologies, Inc., Novato, CA) was amplified using the OPCRar primers FP3 and RP3 to generate a product of 133 bp, which was visualized by electrophoresis on a 12% acrylamide gel stained with ethidium bromide. OPCRar according to one embodiment of the present invention was performed in the presence or absence of thermostable SSB under the following conditions: *i*) 15% DMSO; *ii*) 15% DMSO, 5% Glycerol; *iii*) 15% DMSO, 0.25 M Betaine. The cycling parameters used for all reactions were 75 °C for 15 sec and 65 °C for 15 sec, repeated 45 times.

[0061] In addition to thermostable SSBs that aid OPCRar, non-thermostable SSBs such as T4 bacteriophage SSB (New England BioLabs) may be used to reduce primer dimer formation in the initial heating of the OPCRar solution (**figure 8**). By preincubating OPCRar primers in the presence of a molar excess of T4 gene 32 protein and then adding them to the reaction mixture, it has been observed that the unwanted amplification of primer dimers is minimized during OPCRar. These SSBs presumably bind to the single stranded oligonucleotide primers, reducing the potential for 3'-3' pairing and, thus, primer dimer formation. Upon heating the solution above 65 °C, the T4 SSB is denatured and releases the primers for normal reactivity during thermal cycling. Referring now to **figure 8**, a gel illustrates the reduction in the amount of primer-dimer formed during one embodiment of the present invention with pre-incubation of primers with T4 gene 32 protein. Before addition to the reaction mixture, OPCRar primers were incubated with the indicated stoichiometric excess of active units of T4 gene 32 protein (T4 SSB) at 25 °C for 5 minutes in the presence of 1X ThermoPol buffer (New England BioLabs, Beverly, MA). A synthetic Universal Flu A DNA template (1 E6 copies/ $\mu$ L, Biosearch Technologies, Inc.) was amplified using primers FP3 and RP3 to generate a product sequence of 133 bp. The reactions were held at 85 °C for 2 minutes followed by 50 cycles of 75 °C for 15 sec and 65 °C for 15 seconds. Reaction products were visualized by electrophoresis on a 12% acrylamide gel stained with ethidium bromide. The experiment described in **figure 7** was repeated on a different day by a different researcher to assess the reproducibility of the data and shown in **figure 8**. It can be clearly observed that the pre-incubation of primers with T4 SSB both increases the amount of amplified product and decreases the intensity of the primer-dimer band (~100 bp), relative to no pre-incubation (rightmost lanes).

[0062] The OPCRar method is particularly well suited for use with a device such as that described in the commonly owned provisional patent application filed on the same date hereof, entitled "INTEGRATED DEVICE FOR NUCLEIC ACID DETECTION AND IDENTIFICATION". The configuration of certain embodiments of that device enables the temperature of a solution to rapidly cycle while the solution remains in the same chamber, preferably without active cooling. For example, the temperature could increase or decrease sufficiently to perform OPCRar in less than or equal to 20 seconds, or more preferably less than or equal to 15 seconds, or more preferably less than or equal to about 8 seconds, or more preferably less than or equal to about 4 seconds. Thus and OPCRar temperature cycle could be performed in as little as, or even faster than, 8 seconds.

#### **EXAMPLE 1: Method of Amplification of a DNA Target Duplex by OPCRar**

[0063] To demonstrate that OPCRar is capable of amplifying a specific target sequence present in a double stranded DNA analyte, we used two OPCRar primers, primer HLB (Huang Long Bing) ForSh and primer HLBRevSh, to generate a 140 bp sequence from a PCR-amplified fragment of the C. Liberibacter asiaticus elongation factor gene by the OPCRar system. OPCRar Buffer (10X) was premade and contained 400 mM Tris-HCl (pH 8.4), 10 mM ammonium sulfate, 100 mM potassium chloride, and 0.25% Triton X-100. A 20  $\mu$ L OPCRar solution was set up by mixing:

8.4  $\mu$ L water

2.0  $\mu$ L 10x OPCRar Buffer

3.0  $\mu\text{L}$  DMSO  
0.4  $\mu\text{L}$  potassium chloride (2 M)  
0.5  $\mu\text{L}$  magnesium chloride (100 mM)  
0.5  $\mu\text{L}$  dithiothreitol (100 mM)  
0.5  $\mu\text{L}$  dNTPs (10 mM)  
2.0  $\mu\text{L}$  Primer set HLBForSh and HLBRevSh (4  $\mu\text{M}$  each)  
0.5  $\mu\text{L}$  VentR (exo-) DNA Polymerase (2 U/  $\mu\text{L}$ )  
0.2  $\mu\text{L}$  Et SSB, Extreme Thermostable Single Stranded Binding Protein (500  $\mu\text{g}/\text{mL}$ )  
2.0  $\mu\text{L}$  of PCR product dilution (0.6 to 0.0006 ng/  $\mu\text{L}$  starting concentration)

**[0064]** The reaction was heated at 85 °C for 2 minutes to denature the template and then cycled 40 times, oscillating between 80 °C for 5 seconds, and 65 °C for 5 seconds. After the reactions were complete, 5  $\mu\text{L}$  of OPCRar product was mixed with 2  $\mu\text{L}$  of 6X Sample Loading Buffer (New England BioLabs) and 1  $\mu\text{L}$  of formamide, run on a 12% acrylamide gel, and visualized with ethidium bromide. A 140 bp product was clearly observed at all dilutions shown, and matches the predicted length of the OPCRar target sequence (**figure 9**).

**[0065]** Referring now to **figure 9**, amplification of a specific target sequence present in a double stranded DNA according to one embodiment of the present invention is shown in a gel. Serial dilutions of a PCR-amplified fragment of the *C. Liberibacter asiaticus* elongation factor gene (0.6 to 0.0006 ng/ $\mu\text{L}$ ) were used as the starting template. The reaction was carried out using VentR(exo-) DNA polymerase, Et SSB, and the primers HLBForSh and HLBRevSh to generate a 140 bp sequence in the presence of 15% DMSO. The reactions were heated at 85 °C for 2 minutes to denature the template and then cycled 40 times, oscillating between 80 °C for 5 seconds, and 65 °C for 5 seconds. The OPCRar products were visualized on a 12% acrylamide gel, stained with ethidium bromide. A 140 bp product matching the predicted length of the target sequence was clearly observed at all dilutions shown.

**EXAMPLE 2: Method of Amplification of a Single Stranded DNA Target by OPCRar**

**[0066]** To demonstrate that OPCRar is capable of amplifying a specific target sequence from a single stranded DNA template, we used OPCRar primers FP3 and RP4 to generate a 153 bp sequence from a commercially available Universal Influenza A template (Biosearch Technologies, Inc.) by the OPCRar system. 10xOPCRar Buffer contains 400 mM Tris-HCl (pH 8.4), 10 mM ammonium sulfate, 100 mM potassium chloride, and 0.25% Triton X-100. A 20  $\mu\text{L}$  OPCRar solution was set up by mixing:

8.4  $\mu\text{L}$  water  
2.0  $\mu\text{L}$  10x OPCRar Buffer  
3.0  $\mu\text{L}$  DMSO  
0.4  $\mu\text{L}$  potassium chloride (2 M)  
0.5  $\mu\text{L}$  magnesium chloride (100 mM)  
0.5  $\mu\text{L}$  dithiothreitol (100 mM)  
0.5  $\mu\text{L}$  dNTPs (10 mM)  
2.0  $\mu\text{L}$  Primer set FP3 and RP4 (8  $\mu\text{M}$  each)  
0.5  $\mu\text{L}$  VentR (exo-) DNA Polymerase (2 U/  $\mu\text{L}$ )  
0.2  $\mu\text{L}$  Et SSB, Extreme Thermostable Single Stranded Binding Protein (500  $\mu\text{g}/\text{mL}$ )  
2.0  $\mu\text{L}$  of single stranded DNA template (1 E9 to 1 E2 copies/ $\mu\text{L}$ ).



**[0067]** As a comparison of sensitivity, a real time PCR reaction was run using identical template concentrations as that used for the above OPCRars. 10x ThermoPol (New England BioLabs) contains 200 mM Tris-HCl (pH 8.8), 100 mM ammonium sulfate, 100 mM potassium chloride, 20 mM magnesium sulfate, and 1% Triton X-100. A 15  $\mu$ L RT-PCR solution was set up by combining:

9.7  $\mu$ L water

1.5  $\mu$ L 10x ThermoPol Buffer

0.4  $\mu$ L dNTPs (10 mM)

1.5  $\mu$ L Primer set UniAfCDC/UniArCDC (4  $\mu$ M each) including TaqMan probe UniApCDC (1  $\mu$ M)

0.4  $\mu$ L Taq Polymerase (5 U/  $\mu$ L)

1.5  $\mu$ L single stranded DNA template (1 E9 to 1 E2 copies/ $\mu$ L)

**[0068]** A serial dilution of universal Influenza A single stranded DNA template between 1 E9 to 1 E2 copies/ $\mu$ L was amplified by OPCRar in the presence of 15% DMSO, and real time PCR using a TaqMan probe. OPCRar reactions were first heated to 85  $^{\circ}$ C for 2 minutes, then cycled between 80  $^{\circ}$ C for 15 sec and 65  $^{\circ}$ C for 15 sec, repeated 40 times. RT-PCR reactions were heated to 95  $^{\circ}$ C for 2 minutes, then cycled 45 times between 95  $^{\circ}$ C for 10 sec and 58  $^{\circ}$ C for 40 sec. After the reactions were complete, 5  $\mu$ L of OPCRar product was mixed with 2  $\mu$ L of 6X Sample Loading Buffer (New England BioLabs) and 1  $\mu$ L of formamide, run on a 12% acrylamide gel, and visualized with ethidium bromide. A 153 bp product was clearly observed for all samples (**figure 10**), and matches the predicted length of the OPCRar target sequence.

**[0069]** Referring now to **figure 10**, a gel showing a target sequence present in a single stranded DNA template amplified according to one embodiment of the present invention is presented. A serial dilution of universal Influenza A single stranded DNA template (1 E9 to 1E2 copies/ $\mu$ L, Biosearch Technologies, Inc.) was amplified by OPCRar using the primers FP3 and RP4 in the presence of 15% DMSO to generate a product of 153 bp, which was visualized by electrophoresis on a 12% acrylamide gel stained with ethidium bromide (Left panel). As a comparison, identical dilutions were used as a starting template for real time PCR using the primers set UniAfCDC/UniArCDC and TaqMan probe UniApCDC (Right panel). OPCRar reactions were first heated to 85  $^{\circ}$ C for 2 minutes, then cycled between 80  $^{\circ}$ C for 15 sec and 65  $^{\circ}$ C for 15 sec, repeated 40 times. RT-PCR reactions were heated to 95  $^{\circ}$ C for 2 minutes, then cycled 45 times between 95  $^{\circ}$ C for 10 sec and 58  $^{\circ}$ C for 40 sec. It is evident that OPCRar has a similar sensitivity to conventional PCR reactions when properly optimized.

### **EXAMPLE 3: Method of Amplification of a Specific Sequence Present on Plasmid DNA by OPCRar**

**[0070]** To demonstrate that OPCRar is capable of amplifying a specific target sequence present in a double stranded plasmid DNA, we used two OPCRar primers, primer *hyvl\_For* and primer *hyvl\_Rev*, to generate a 139 bp sequence from a plasmid containing the *C. Liberibacter asiaticus hyvl* gene by the OPCRar system. OPCRar Buffer (10X) was premade and contained 400 mM Tris-HCl (pH 8.4), 10 mM ammonium sulfate, 100 mM potassium chloride, and 0.25% Triton X-100. A 20  $\mu$ L OPCRar solution was set up by mixing:

9.4  $\mu$ L water

2.0  $\mu$ L 10x OPCRar Buffer

2.0  $\mu$ L DMSO

0.4  $\mu$ L potassium chloride (2 M)

0.5  $\mu$ L magnesium chloride (100 mM)

0.5  $\mu$ L dithiothreitol (100 mM)

0.5  $\mu$ L dNTPs (10 mM)

2.0  $\mu$ L Primer set *hyvl\_For* and *hyvl\_Rev* (8  $\mu$ M each)

0.5  $\mu$ L VentR (exo-) DNA Polymerase (2 U/  $\mu$ L)

0.2  $\mu$ L Et SSB, Extreme Thermostable Single Stranded Binding Protein (500  $\mu$ g/mL)

2.0  $\mu$ L of DNA extracted from healthy and *C. Liberibacter* infected tissue (17.2 ng/  $\mu$ L)

**[0071]** A titration with DMSO was performed from 13-8% v/v. The reaction was heated at 85 °C for 2 minutes to denature the template and then cycled 40 times, oscillating between 80 °C for 10 seconds, and 65 °C for 10 seconds. After the reactions were complete, 5  $\mu$ L of OPCRar product was mixed with 2  $\mu$ L of 6X Sample Loading Buffer (New England BioLabs) and 1  $\mu$ L of formamide, run on a 12% acrylamide gel, and visualized with ethidium bromide. A 139 bp product was clearly observed for all samples (**figure 11**), and matches the predicted length of the OPCRar target sequence.

**[0072]** Referring now to **figure 11**, a gel of a specific target sequence present in plasmid DNA amplified according to one embodiment of the present invention is presented. A plasmid containing the *C. Liberibacter asiaticus hyvI* gene (17.2 ng/ $\mu$ L) was used as a starting template, and the concentration of DMSO was titrated from 13-8% v/v. All reactions were performed using VentR(exo-) DNA polymerase, Et SSB, and the primers *hyvI\_For* and *hyvI\_Rev*. The reactions were heated at 85 °C for 2 minutes to denature the template and then cycled 40 times, oscillating between 80 °C for 10 seconds, and 65 °C for 10 seconds. The OPCRar products were visualized on a 12% acrylamide gel, stained with ethidium bromide. A 139 bp product matching the predicted length of the target sequence was clearly observed at all DMSO concentrations tested.

**EXAMPLE 4: Method of Amplification of a RNA Target Sequence of a Human Pathogenic Virus Present in Nasal Aspirate by OPCRar**

**[0073]** To demonstrate that OPCRar is capable of amplifying a specific target sequence present in a single stranded RNA template, we used the OPCRar primer pair, FP3 and RP4, to generate a 153 bp sequence from ribonucleic acid isolated from clinical nasal aspirates either infected or uninfected with Influenza A virus. OPCRar Buffer (10X) was premade and contained 400 mM Tris-HCl (pH 8.4), 10 mM ammonium sulfate, 100 mM potassium chloride, and 0.25% Triton X-100. A 20  $\mu$ L OPCRar solution was set up by combining:

9.3  $\mu$ L water

2.0  $\mu$ L 10x OPCRar Buffer

3.0  $\mu$ L DMSO

0.4  $\mu$ L potassium chloride (2 M)

0.5  $\mu$ L magnesium chloride (100 mM)

0.5  $\mu$ L dithiothreitol (100 mM)

0.5  $\mu$ L dNTPs (10 mM)

2.0  $\mu$ L Primer set FP3 and RP4 (8  $\mu$ M each)

0.5  $\mu$ L VentR (exo-) DNA Polymerase (2 U/ $\mu$ L)

0.2  $\mu$ L Et SSB, Extreme Thermostable Single Stranded Binding Protein (500  $\mu$ g/mL)

0.1  $\mu$ L Superscript III Reverse Transcriptase (200 U/ $\mu$ L)

2.0  $\mu$ L of Nucleic acid isolated from clinical Nasal Aspirate (0.3 ng/  $\mu$ L)

**[0074]** The reaction was incubated at 55 °C for 5 minutes to generate cDNA, heated to 85 °C for 2 minutes to denature the template and then cycled 40 times, oscillating between 80 °C for 10 seconds, and 65 °C for 10 seconds. After the reactions were complete, 5  $\mu$ L of OPCRar product was mixed with 2  $\mu$ L of 6X Sample Loading Buffer (New England BioLabs) and 1  $\mu$ L of formamide, run on a 12% acrylamide gel, and visualized with ethidium bromide. A 153 bp product was clearly observed in the positive, but not negative clinical sample (**figure 12**), and matches the predicted length of the OPCRar target sequence.

**[0075]** Referring now to **figure 12**, a gel illustrating a specific target sequence present in single stranded RNA amplified according to one embodiment of the present invention is illustrated. Ribonucleic acid isolated from nasal aspirates either infected or uninfected with Influenza A virus (0.3 ng/ $\mu$ L) was used as template. All reactions were performed using Superscript III reverse transcriptase, VentR(exo-) DNA polymerase, Et SSB, and the primers FP3 and RP4, in the presence of 15% DMSO. The reaction was incubated at 55 °C for 5 minutes to generate cDNA, heated to 85 °C for 2 minutes to denature the template and then cycled 40 times, oscillating between 80 °C for 10 seconds, and 65 °C for 10 seconds. The OPCRar products were visualized on a 12% acrylamide gel, stained with ethidium bromide. A 153 bp product matching the predicted length of the target sequence was clearly observed in the positive, but not negative, clinical sample.

**EXAMPLE 5: Method of Amplification of a Target Sequence from a Pathogenic Plant Bacteria by OPCRar**

**[0076]** To demonstrate that OPCRar is capable of amplifying a specific target sequence present in a pathogenic bacterial genome, we used the OPCRar primer pair, EU523377-F-57 and EU523377-R-56, to generate a 213 bp fragment of the *C. Liberibacter asiaticus* elongation factor gene from total nucleic acid isolated from infected plant tissue. OPCRar Buffer (10X) was premade and contained 400 mM Tris-HCl (pH 8.4), 10 mM ammonium sulfate, 100 mM potassium chloride, and 0.25% Triton X-100. A 20  $\mu$ L OPCRar solution was set up by combining:

8.4  $\mu$ L water

2.0  $\mu$ L 10x OPCRar Buffer

3.0  $\mu$ L DMSO

0.4  $\mu$ L potassium chloride (2 M)

0.5  $\mu$ L magnesium chloride (100 mM)

0.5  $\mu$ L dithiothreitol (100 mM)

0.5  $\mu$ L dNTPs (10 mM)

2.0  $\mu$ L Primer set EU523377-F-57 and EU523377-R-56 (4  $\mu$ M each), primer EU523377-F-57 was

either biotinylated (5') or non-biotinylated

0.5  $\mu$ L VentR (exo-) DNA Polymerase (2 U/ $\mu$ L)

0.2  $\mu$ L Et SSB, Extreme Thermostable Single Stranded Binding Protein (500  $\mu$ g/mL)

2.0  $\mu$ L of Total nucleic acid isolated from *C. Liberibacter asiaticus* infected plant tissue (1.1 ng/  $\mu$ L)

**[0077]** To demonstrate that primer modifications are compatible with OPCRar, the forward primer EU523377-F-57 was biotinylated at the 5' end in some reactions. OPCRar solutions were heated at 85 °C for 2 minutes to denature the template and then cycled 40 times, oscillating between 80 °C for 15 seconds, and 65 °C for 15 seconds. After the reactions were complete, 5  $\mu$ L of OPCRar product was mixed with 2  $\mu$ L of 6X Sample Loading Buffer (New England BioLabs) and 1  $\mu$ L of formamide, run on a 12% acrylamide gel, and visualized with ethidium bromide. A 213 bp product was clearly observed for all samples, and matches the predicted length of the OPCRar target sequence (**figure 13**).

**[0078]** Referring now to **figure 13**, a gel of a specific target sequence present in bacterial genomic DNA as amplified according to one embodiment of the present invention is presented. Total nucleic acid isolated from *C. Liberibacter asiaticus*-infected leaf tissue (1.1 ng/ $\mu$ L) was used as a starting template. All reactions were performed using VentR(exo-) DNA polymerase, Et SSB, and the primers EU523377-F-57 and EU523377-R-56, in the presence of 15% DMSO. Primer EU523377-F-57 was either biotinylated at the 5' end of the oligonucleotide or unmodified. OPCRar solutions were heated at 85 °C for 2 minutes to denature the template and then cycled 40 times, oscillating between 80 °C for 15 seconds, and 65 °C for 15 seconds. The amplified products were visualized on a 12% acrylamide gel, stained with ethidium bromide. A 213 bp product was clearly observed for all positive samples and no negative samples, matching the predicted length of the OPCRar target sequence.

**EXAMPLE 6: Method of Amplification of a Specific Sequence on Organelle DNA**

**[0079]** To demonstrate that OPCRar is capable of amplifying a specific target sequence present in organelle DNA, in this case chloroplast DNA, we used the OPCRar primer pair, *rbcL\_For* and *rbcL\_Rev*, to generate a 137 bp fragment of the *rbcL* gene of plant from total nucleic acid isolated from plant tissue either infected or uninfected with *C. Liberibacter asiaticus*. OPCRar Buffer (10X) was premade and contained 400 mM Tris-HCl (pH 8.4), 10 mM ammonium sulfate, 100 mM potassium chloride, and 0.25% Triton X-100. A 20  $\mu$ L OPCRar solution was set up by combining:

8.4  $\mu$ L water  
 2.0  $\mu$ L 10x OPCRar Buffer  
 3.0  $\mu$ L DMSO  
 0.4  $\mu$ L potassium chloride (2 M)  
 0.5  $\mu$ L magnesium chloride (100 mM)  
 0.5  $\mu$ L dithiothreitol (100 mM)  
 0.5  $\mu$ L dNTPs (10 mM)  
 2.0  $\mu$ L Primer set *rbcL\_For* and *rbcL\_Rev* (2 or 4  $\mu$ M each)  
 0.5  $\mu$ L VentR (exo-) DNA Polymerase (2 U/ $\mu$ L)  
 0.2  $\mu$ L Et SSB, Extreme Thermostable Single Stranded Binding Protein (500  $\mu$ g/mL)  
 2.0  $\mu$ L of Total nucleic acid isolated from leaf tissue (3.3 ng/  $\mu$ L)

**[0080]** Two different concentrations of the primer pair *rbcL\_For* and *rbcL\_Rev* were used to determine a threshold for the primer concentration necessary to efficiently amplify the *rbcL* gene fragment. The reaction was heated at 85 °C for 2 minutes to denature the template and then cycled 40 times, oscillating between 76 °C for 10 seconds, and 60 °C for 10 seconds. After the reactions were complete, 5  $\mu$ L of OPCRar product was mixed with 2  $\mu$ L of 6X Sample Loading Buffer (New England Biolabs) and 1  $\mu$ L of formamide, run on a 12% acrylamide gel, and visualized with ethidium bromide. A 137 bp product was clearly observed for both infected and uninfected samples, and matches the predicted length of the OPCRar target sequence (**figure 14**).

**[0081]** Referring now to **figure 14**, amplification of a specific target sequence present in chloroplast DNA according to one embodiment of the present invention is presented in a gel. Total nucleic acid isolated from either *C. Liberibacter asiaticus*-infected (*i*) or uninfected (*ii*) leaf tissue (3.3 ng/ $\mu$ L) was used as a starting template. All reactions were performed using VentR(exo-) DNA polymerase, Et SSB, and the primers *rbcL\_For* and *rbcL\_Rev*, in the presence of 10% DMSO. OPCRar solutions were heated at 85 °C for 2 minutes to denature the template and then cycled 40 times, oscillating between 76 °C for 10 seconds, and 60 °C for 10 seconds. The amplified products were visualized on a 12% acrylamide gel, stained with ethidium bromide. A 137 bp product was clearly observed for all positive samples, and matches the predicted length of the OPCRar target sequence.

**EXAMPLE 7: Method of Multiplex Amplification of a Target Sequence and Positive Control by OPCRar**

**[0082]** To demonstrate that OPCRar is capable of amplifying multiple specific target sequences, we amplified the nucleic acid extracted from *C. Liberibacter asiaticus* infected plant tissue with the OPCRar primer pairs, *hyvI\_For/hyvI\_Rev*, and *rbcL\_For/rbcL\_Rev*. These primer sets generate products of 139 and 137 bp, respectively. OPCRar Buffer (10X) was premade and contained 400 mM Tris-HCl (pH 8.4), 10 mM ammonium sulfate, 100 mM potassium chloride, and 0.25% Triton X-100. A 20  $\mu$ L OPCRar solution was set up by combining:

6.4  $\mu$ L water  
 2.0  $\mu$ L 10x OPCRar Buffer  
 3.0  $\mu$ L DMSO  
 0.4  $\mu$ L potassium chloride (2 M)

0.5  $\mu$ L magnesium chloride (100 mM)  
 0.5  $\mu$ L dithiothreitol (100 mM)  
 0.5  $\mu$ L dNTPs (10 mM)  
 2.0  $\mu$ L Primer set *rbcL*\_For and *rbcL*\_Rev (2 or 3  $\mu$ M each)  
 2.0  $\mu$  Primer set *hyvI*\_For and *hyvI*\_Rev (8  $\mu$ M each)  
 0.5  $\mu$ L VentR (exo-) DNA Polymerase (2 U/ $\mu$ L)  
 0.2  $\mu$ L Et SSB, Extreme Thermostable Single Stranded Binding Protein (500  $\mu$ g/mL)  
 2.0  $\mu$ L of Total nucleic acid isolated from leaf tissue (3.3 ng/  $\mu$ L)

**[0083]** Two different concentrations of the primer pair *rbcL*\_For and *rbcL*\_Rev were used to determine a threshold for the primer concentration necessary to efficiently amplify the *rbcL* gene fragment in the presence of 800 nM primers specific for the *hyvI* gene fragment. The reaction was heated at 85 °C for 2 minutes to denature the template and then cycled 40 times, oscillating between 76 °C for 10 seconds, and 60 °C for 10 seconds. After the reactions were complete, 5  $\mu$ L of OPCRar product was mixed with 2  $\mu$ L of 6X Sample Loading Buffer (New England BioLabs) and 1  $\mu$ L of formamide, run on a 12% acrylamide gel, and visualized with ethidium bromide. Both 139 bp and 137 bp products are clearly observed, matching the predicted length of the OPCRar target sequences. For clarity, OPCRar products generated using both primer pairs alone (Examples 3 and 6) were run alongside the multiplex reactions (**figure 15**).

**[0084]** Referring now to **figure 15**, a gel illustrating amplification products of multiplex of two target sequences according to one embodiment of the present invention is presented. Total nucleic acid isolated from *C. Liberibacter asiaticus*-infected leaf tissue (3.3 ng/ $\mu$ L) was used as a starting template. All reactions were performed using VentR(exo-) DNA polymerase, Et SSB, and the primers sets *hyvI*\_For/*hyvI*\_Rev and *rbcL*\_For/*rbcL*\_Rev at the indicated concentrations, in the presence of 10% DMSO. Multiplex OPCRar solutions were heated at 85 °C for 2 minutes to denature the template and then cycled 40 times, oscillating between 76 °C for 10 seconds, and 60 °C for 10 seconds. The multiplex products were visualized on a 12% acrylamide gel, stained with ethidium bromide. When compared to OPCRar products generated from either primer set alone (See **figures 11** and **15**), both 139 and 137 bp products were clearly observed.

**[0085]** Referring now to **figure 16**, an amplification product resulting from a reaction in the presence of Et SSB according to one embodiment of the present invention is illustrated. SSB enhances the amplification efficiency at lower melting temperatures using the system and method of the present invention. We investigated the use of extreme thermostable single strand binding protein "ET SSB" to show it aids the performance of OPCRar at certain melting temperatures. OPCRar reaction was performed for a HLB pathogen target with purified leaf sample DNA containing HLB disease genes. Template was DNA purified from citrus leaf of a *C. Liberibacter* infected tree. Primers were *hyvI*\_For/ and *hyvI*\_Rev. Thermocycler conditions were: Initial melt of 85°C for 2 minutes, followed by 40 cycles denaturation of either 76°C or 74°C for 10 seconds and annealing at 60°C for 10 seconds. The results indicated in the presence of Et SSB resulted in the amplification of the *C. Liberibacter* target at all assayed temperature regimens. The comparative OPCRar experiment was done in duplicates in the presence of ET SSB protein and no ET SSB. A standard PCR Thermocycler with precise temperature control was used to allow the evaluation of amplification performance with and without SSB. Results indicated that in the presence of ET SSB we picked up amplified HLB at 74°C and without ET SSB it was unobservable.

**[0086]** Referring now to **figure 17**, gel electrophoresis of OPCRar reactions conducted using *hyvI*\_For and *hyvI*\_Rev primers and purified *C. Liberibacter* DNA isolated from the leaf of an infected tree is shown according to one embodiment of the present invention wherein the reaction does not include ramping parameters involved in a typical PCR Thermocycler **figure 17A** or does include ramping time to cycling temperature **figure 17B**. Ramping as used herein is a reference to the heating up (e.g. in each thermal cycling, the heating process to raise the amplification reaction temperature from annealing to denaturing temperature is called ramping up or cooling down from denaturation temperature to annealing temperature is called ramp down. All the conventional PCR cyclers and methods, have controlled heating design where ramping up is about > 3 deg/second, and ramping down rate is about >1 deg/second. Ramping time is not included in the typical cycling profiles, the instrument does not start counting time duration during denaturation, annealing and extension stage until the desired temperature is reached. For instance, denaturation time is 10 second at 90 deg, the instrument will not starting counting the 10 second time until 90 deg is reached. In

contrast, a system and method of the present invention provides for a low cost heater device which is designed for example the 80 deg for 10 second means the time starts counting when the heating process starts (rather than waiting until the desired denaturing temperature is reached). **Figure 17 A:** we examined the OPCRar amplification of a HLB disease (a citrus greening disease pathogen, Huang Long Bing) target in a low cost, thermal heater engine (without active cooling and precise temperature control,  $\pm 2$  degree fluctuation, (see for example systems and apparatus as disclosed in 61/477,357). OPCRar 20  $\mu$ L reactions were conducted in microheater reaction chambers under microprocessor control in a device developed by Mesa Tech International, Inc. (MTI Device) or in a traditional PCR thermocycler (PCR Thermocycler) such that the 10-second dwell times for each temperature segment of the program were calculated starting immediately after command to change temperatures was executed by the microprocessor (i.e. no ramping time **figure 17A**). This is in contrast, to 10-second dwell times for each temperature segment of the program calculated starting immediately after the target temperature was detected by the temperature sensor located at the reaction well (i.e. ramping time **figure 17B**). The PCR Thermocycler conditions were as follows: Initial Melt: 85°C for 2 minutes, followed by 40 cycles of 80°C for 10 seconds and 60°C for 10 seconds. The MTI Device conditions were as follows: Initial Melt: 85°C for 2 minutes, 40 cycles of 82°C for 10 seconds and 59°C for 20 seconds.

**[0087] Figure 17A:** First Lane, OPCRar amplification (20  $\mu$ l reaction) without separate ramping. From the left: 50 bp standard DNA size ladder; second and third lanes: OPCRar amplification reaction (in duplicate, 2<sup>nd</sup> and 3<sup>rd</sup> lanes) in a standard PCR thermal cycler engine where ramp step and precise temperature control are accommodated. Initial Melt: 85°C for 2 minutes, Cycling between denaturation of 80°C for 10 seconds and annealing of 60°C for 10 seconds with 40 cycles. The 4<sup>th</sup> and 5<sup>th</sup> lanes: OPCRar amplification performed in a low cost thermal engine without active cooling and/or a separate ramping control. The thermal engine ramps to either 80° C denaturation or ramping down to 60°C annealing temperature from the ambient field temperatures (~ 25 deg). as compared to not having any ramping stage for either denaturation or annealing temperature. HLB OPCRar was performed with purified plant DNA containing HLB disease target sequences in a 20  $\mu$ L reaction. We used the PCR Thermocycler for a positive control for the MTI Device test. The PCR Thermocycler conditions were the following: The MTI Device Thermocycler conditions for no ramping were the following: Initial Melt: 85°C for 2 minutes, Cycling between denaturation of 82°C for 10 seconds and annealing of 59°C for 20 seconds. The cycling was repeated 40 times. The MTI device thermocycler for ramping up stages were the same conditions except there was ramp up stage < 10 seconds and ramp down strage < 20 seconds. Data suggested HLB OPCRar can still amplify up HLB amplicon even without ramp up and ramp down stages. Comparison of 17A and 17B reveals a significant improvement in the yield of amplification product when ramp time is provided.

**[0088]** Referring now to **figure 18**, multiplexed OPCRar reactions according to one embodiment of the present invention containing two primer sets in MTI's thermal resistor-based low cost amplification device as described above: 1) *hylV*\_For and *hylV*\_Rev primers specific for a *C. Liberibacter* DNA target and 2) primers *rbcl*\_(Ribulose-1,5-bisphosphate carboxylase oxygenase) For and *rbcl*\_Rev specific for a citrus house keeping, gene *rbcl*. amplification product are illustrated in the gel. The reaction products resulting from PCR amplification as described and performed in a low cost heater without ramping (e.g. 5 deg/second in PCR device) and without precise temperature control were run on a gel. We tested the RbCL internal positive control with HLB primers in the PCR thermocycler and the MTI Device (with ramping or no ramping program). We performed a 40  $\mu$ L reaction with purified HLB sample. The PCR thermocycler conditions were the following: Initial Melt: 85°C for 2 minutes, Cycling between denaturation of 80°C for 10 seconds and annealing of 60°C for 10 seconds. The MTI Device amplification conditions were the following: Initial Melt: 90°C for 2 minutes, cycling between denaturation of 82°C for 10 seconds and annealing of 59°C for 20 seconds. We discovered without ramping it was still able to amplify up both HLB and RbCL primer sequences. HLB product around - 147 bp and RbCL product -140 bp. The MTI Device for both conditions was comparable to the PCR Thermocycler amplified product. The forward primer for this reaction is ccagcctga tcgtacaaa gggcgatgct acaacatt (SEQ ID NO 9) (Tm of about 73.9C (10% DMSO , 76/60C) and the reverse primer iscatgttagta acagaacctt ctcaaaaag gtctaacggg taa (SEQ ID NO 10) (Tm of about 71.2C (10% DMSO, 76/60 C). OPCRar reactions were conducted with or without the inclusion of ramp times in the temperature dwell times (as described for Figures 16 and 17) as indicated. 40  $\mu$ L reactions were conducted using purified citrus leaf DNA from a *C. Liberibacter* infected tree. The PCR Thermocycler control conditions were as follows: Initial Melt: 85°C for 2 minutes, 40 cycles of 80°C for 10 seconds and 60°C for 10 seconds. The MTI Device amplification conditions were as follows: Initial Melt: 90°C for 2 minutes, 40 cycles of 82°C for 10 seconds and 59°C for 20 seconds. The results revealed without the inclusion of ramp times in the dwell time calculation, the MTI device was able to amplify up both *C. Liberibacter* and *rbcl* sequences. The *C. Liberibacter* product is approximately 147 bp (HLB) and *rbcl* (RbCL) product is approximately 140 bp.

**[0089]** All primer melting temperatures (Tm) calculated using IDT OligoAnalyzer 3.1 (Integrated DNA Technologies, Inc., Coralville, IA) using the Primer 3 Tm calculating software where salt, dNTP, Mg, primer concentration parameters are considered using the following parameters:

Oligonucleotide Concentration: 0.25  $\mu$ M; Na<sup>+</sup> Concentration: 50mM; Mg<sup>++</sup> Concentration = 2.5 mM; dNTPs Concentration = 0.25  $\mu$ M. Symbol "a" means adenine, "g" means guanine, "c" means cytosine, "t" means thymine, "u" means uracil, "r" means purine,

"y" means pyrimidine, "m" means amino, "k" means keto, "n" means any of a or g or c or t/u, unknown, or other.

(SEQ ID NO 1)

**ACCESSION NUMBER:** CY087034

**TYPE:** Viral RNA

**LENGTH:** 987

**ORGANISM:** Influenza A Virus (H1N1)

**OTHER INFORMATION:** matrix protein 2 (M2) and matrix protein 1 (M1) genes

(SEQ ID NO 2)

**TYPE:** Forward Primer

**NAME:** FP3

**LENGTH:** 46

**Tm:** mean 75 °C

(SEQ ID NO 3)

**TYPE:** Reverse Primer

**NAME:** RP3

**LENGTH:** 45

**Tm:** mean 77.8 °C

(SEQ ID NO 4)

**TYPE:** Reverse Primer

**NAME:** RP4

**LENGTH:** 46

**Tm:** mean 74.7 °C

(SEQ ID NO 5)

**TYPE:** Forward Primer

**NAME:** UhiAfCDC

**LENGTH:** 22

**Tm:** mean 65.0 °C

(SEQ ID NO 6)

**TYPE:** Reverse Primer

**NAME:** UhiArCDC

**LENGTH:** 24

**Tm:** mean 66.6 °C

(SEQ ID NO 7) FAM-tgcagtcctc gctcactggg cacg-BHQ

**TYPE:** TaqMan Probe

**NAME:** UhiApCDC

**LENGTH:** 24

**Tm:** 73.4 °C

(SEQ ID NO 8)

**ACCESSION NUMBER:** AB505957

**TYPE:** Chloroplast DNA

**LENGTH:** 1326

**ORGANISM:** Citrus sinensis

**OTHER INFORMATION:** *rbcL*, ribulose-1,5-bisphosphate carboxylase/oxygenase large subunit

(SEQ ID NO 9)

**TYPE:** Forward Primer

**NAME:** rbcL\_For

**LENGTH:** 38

**Tm:** 73.9 °C

(SEQ ID NO 10)

**TYPE:** Reverse Primer

**NAME:** rbcL\_Rev

**LENGTH:** 43

**Tm:** 71.2 °C

(SEQ ID NO 11)

**ACCESSION NUMBER:** From EU523377

**TYPE:** Bacterial DNA

**LENGTH:** 890

**ORGANISM:** *Candidatus Liberibacter asiaticus*

**OTHER INFORMATION:** elongation factor Ts

(SEQ ID NO 12)

**TYPE:** Forward Primer

**NAME:** NBEU523377-F-57

**LENGTH:** 57

**Tm:** 75.8 °C

(SEQ ID NO 13) [Biotin-5]tcttcgtatc ttcacgcttc tcctctcgag ggtaggat cgattggtgt tcttgta

**TYPE:** Biotinylated Forward Primer

**NAME:** EU523377-F-57

**LENGTH:** 57

**Tm:** 75.8 °C

(SEQ ID NO 14)

**TYPE:** Forward Primer

**NAME:** HLBForSh

**LENGTH:** 47

**Tm:** 75.6 °C

(SEQ ID NO 15)

**TYPE:** Reverse Primer

**NAME:** EU523377-R-56

**LENGTH:** 56

**Tm:** 75.8 °C

(SEQ ID NO 16)

**TYPE:** Reverse Primer

**NAME:** HLBRevSh

**LENGTH:** 49

**Tm:** 75.5 °C

(SEQ ID NO 17)

**TARGET:** *Candidatus Liberibacter asiaticus* 16S ribosomal RNA

**TYPE:** Forward Primer (underlined), contains 5' detection sequence

**NAME:** HLBas-P2

**LENGTH:** 39

**Tm:** 62.7 °C

(SEQ ID NO 18)

**TARGET:** *Candidatus Liberibacter asiaticus* 16S ribosomal RNA

**TYPE:** Reverse Primer (underlined), contains 5' T7 promoter

**NAME:** HLBr-P1

**LENGTH:** 56

**Tm:** 64.5°C

(SEQ ID NO 19)

**TYPE:** Forward Primer

**NAME:** hyvl\_For

**LENGTH:** 45

**Tm:** 72.2 °C



(SEQ ID NO 20)  
**TYPE:** Reverse Primer  
**NAME:** hyvl\_Rev  
**LENGTH:** 51  
**Tm:** 70.9 °C

**[0090]** Although the invention has been described in detail with particular reference to the described embodiments, other embodiments can achieve the same results. Variations and modifications of the present invention will be obvious to those skilled in the art and it is intended to cover all such modifications and equivalents. The entire disclosures of all patents and publications cited above are hereby incorporated by reference.

**SEQUENCE LISTING**

**[0091]**

<110> Mesa Tech International, Inc.  
<120> Oscillating Amplification Reaction for Nucleic Acids  
<130> 33106-PCT2  
<140> 61/477,357  
<141> 2011-04-20  
<150> 61/477,437  
<151> 2011-04-20  
<160> 28  
<170> PatentIn version 3.5  
<210> 1  
<211> 1010  
<212> DNA  
<213> Influenza A virus  
<220>  
<221> misc\_feature  
<222> (993)..(993)  
<223> n is a, c, g, or t  
<220>  
<221> misc\_feature  
<222> (1002)..(1002)  
<223> n is a, c, g, or t  
<400> 1

```

taaagatgag tcttctaacc gaggtcgaaa cgtacgttct ttctatcatc ccgtcaggcc      60
ccctcaaagc cgagatcgcg cagagactgg aaagtgtctt tgcaggaaa aacacagatc      120
ttgaggctct catggaatgg ctaaagacaa gaccaatctt gtcacctctg actaagggaa      180
ttttaggatt tgtgttcacg ctcaccgtgc ccaagtgcg aggactgcag cgtagacgct      240
ttgtccaaaa tgccctaaat gggaatgggg acccgaacaa catggataga gcagttaaac      300
tatacaagaa gctcaaagga gaaataacgt tccatggggc caagcaggtg tccactaagct      360
ttcaactggt gcacttgcca gttgcatggg cctcatatac aacaggatgg gaacagtgac      420
cacagaagct gcttttggto tagtgtgtgc cacttgtgaa cagattgctg attcacagca      480
tcggtctcac agacaaatgg ctactaccac caatccacta atcaggcatg aaaacagaat      540
ggtgctggct agcactacgg caaaggctat ggaacagatg gctggatcga gtgaacaggc      600
agcagaggcc atggaggttg ctaatcagac taggcagatg gtacatgcaa tgagaactat      660
tgggactcat cctagctcca gtgctggtct gaaagatgac cttcttgaaa atttgaggc      720
ctaccagaag cgaatgggag tgcagatgca gcgattcaag tgatcctctc gtcattgcag      780
caaatatcat tgggatcttg cacctgatat tgtggattac tgatcgtctt tttttcaaat      840

gtatztatcg tcgctttaa tacggtttga aaagagggcc ttctacggaa ggagtgcctg      900
agtccatgag ggaagaatat caacaggaac agcagagtgc tgtggatgtt gacgatggtc      960
attttgtcaa catagagcta gagtaataga cgntttgtcc anaatgcctt      1010

```

<210> 2

<211> 46

<212> DNA

<213> Artificial Sequence

<220>

<223> forward primer designed for SEQ ID NO 1

<400> 2

ywctcatgga rtggctaaag acaagaccra tctgtcacc tctgac 46

<210> 3

<211> 40

<212> DNA

<213> Artificial Sequence

<220>

<223> reverse primer designed for SEQ ID NO 1

<400> 3

agggcattyt ggacaaakcg tctacgytgc agtccycgyt 40

<210> 4

<211> 46

<212> DNA

<213> Artificial Sequence

<220>

<223> reverse primer for SEQ ID NO 1

<400> 4

ttggtrctc cattyccatt tagggcatty tggacaaakc gtclat 46

<210> 5

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> forward primer UniAfCDC for SEQ ID NO 1

<400> 5  
gaccratcct gtcacctctg ac 22

<210> 6

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> reverse primer UniArCDC designed for SEQ ID NO 1

<400> 6

agggcattyt ggacaakcg tcta 24

<210> 7

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> TAQman probe UniApCDC

<400> 7

tcgagtcctc gctcactggg cacg 24

<210> 8

<211> 1326

<212> DNA

<213> Citrus sinensis

<400> 8

tgttggattc aaggccggtg ttaagatta taaattgact tattatactc ctgactatgt	60
aaccaaagat actgatatct tggcagcatt ccgagtaact cctcagcccg gagttccacc	120
cgaggaagcg ggggctgcgg tagctgcgga atcttctact ggtacctgga cagctgtgtg	180
gacogatggg cttaccagcc ttgatcgtta caaaggcga tgctacaaca ttgagcccgt	240
tgctggagaa gagaatcaat atatatgta tgtagcttac ccgttagacc tttttgaaga	300
aggttctgtt actaacatgt ttaactccat tgtgggtaat gtatttgggt tcaaagcaact	360
gcgcgctcta cgtctagagg atctacgaat ccctcctgcg tatactaaaa ctttccaagg	420
cccgcctcac ggcattccaag ttgagagaga taaattgaac aagtatggcc gtcocctgtt	480
gggatgtact attaaacctt aactgggggt atccgcgaag aattatgta gggcggttta	540
tgaatgtcta cgtggtggac ttgactttac caaagatgat gagaactgta actcccaacc	600
atztatgcgt tggagggacc gtttcttatt ttgtgcggaa gctctttata aagcgcgaagc	660
tgaaacaggt gaaatcaaag gtcattactt gaatgctact gcagggacat gcgaagaaat	720
gctaaaaagg gctgtctttg ccagagagtt gggagttcct atcgtaatgc atgactactt	780
aacaggggga ttcaccgcaa atactacctt ggctcattat tgccgagata atggtctact	840
tcttcacatc caccgtgcaa tgcattgcagt tattgataga cagaagaatc atggtatgca	900
ctttcgtgta ctagctaaag ctttgcgtct gtctggtgga gatcatatc acgcccgtac	960
agtagtaggt aaacttgagg gggaaagaga cataaccttg ggatttgttg atttactacg	1020
tgatgatatt gttgaaaaag atcgaagccg cggatattat ttcactcaag attgggtctc	1080
tataccaggt gttataacct tggcttccgg gggtattcac gtttggcata tgcctgcggt	1140
gacagagatc tttggagatg attccgtatt acaatttggg ggaggaactt taggacacc	1200
ttggggaaat gcacccggcg ctgtagctaa tcgagtagct ctagaagcat gtgtacaage	1260
tcgtaatgaa ggaagcgatc ttgctcgtga aggtaatgaa attatccggg aggctagcaa	1320
atggag	1326

<210> 9

<211> 38  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> forward primer rbcL\_For

<400> 9  
 ccagccttga tegtacaaa gggcgtatgct acaacatt 38

<210> 10  
 <211> 43  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> reverse primer rbcL\_Rev

<400> 10  
 catgttagta acagaacctt ctcaaaaag gtctaacggg taa 43

<210> 11  
 <211> 890  
 <212> DNA  
 <213> Candidatus Liberibacter asiaticus

<400> 11  
 atgagtaagg tatctgctgt tgcggtaaag gagttacgtg ggaaaactgg tgcaggatc 60  
 ttgactgca agaatgocct tttggaggct aagggatgata gtgaattagc gattgatatt 120  
 ttgcgcacta aaggagctat ggcagctagt aagagagagg gttagaaagt ttcagagggg 180  
 cttatagtaa ttgctcgtga tggatataag aagcctcga ttgtagaagt caatgttgag 240  
 accgatagtt tggcaaaaaa cactgatatt cagagtcctg tttctaataat tgcgggtatt 300  
 gctctttcca cogatggttc tttggacaat gttcttgcga tgcatttga coatagtggg 360  
 attactgtgg gggatggaat taagcagcag attgctatca cgggtgaatg tattaagctg 420  
 agcggttccg ctcttctgtg tgtttcggaa ggggtcatct ctctctatct tcatgcttct 480  
 ccttctgagg gtttaggato gattggtgtt cttgtagcgt tgcagctctc tgcggaagat 540  
 aaggaattgc tttctgcgat tggagagaag attgcagtgc atgtaatgct ggctctcct 600  
 tctgtgattt ctgtgcagat gcttgatcct tctattgttg ctaataaacg tgcaccattac 660  
 atgacagaag cacttgatcc tgggaaatca ggtaaatattg ttgaaaagat cgtaaatgga 720  
 aagatgcaaa gcttttgtaa ggaatgtgtt cttttgcctc aggggtttgt tgttgatcct 780  
 tccaaaactg tatcagattt tttgaaagaa tctgaaaaat caatcggtgc ttctattgaa 840  
 gttgtggtg talctcatt tttgtgggt aaagaaaatg atgatggta 890

<210> 12  
 <211> 57  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> forward primer NBEU523377-F-57

<400> 12  
 tctctgatac tcatgcttc tcctctgag ggttaggatc cgattggtgt tcttgta 57

<210> 13  
 <211> 57  
 <212> DNA  
 <213> Artificial Sequence

<220>

<223> biotinylated forward primer EU523377-F-57

<400> 13

tcttcgtatc tcatgctc tcctctgag ggtttaggat cgattggtg tcttga 57

<210> 14

<211> 47

<212> DNA

<213> Artificial Sequence

<220>

<223> forward primer HLBForSh

<400> 14

cgattggtg tctttagcg ttgcagctt ctgcggaaga taaggaa 47

<210> 15

<211> 56

<212> DNA

<213> Artificial Sequence

<220>

<223> reverse primer EU523377-R-56

<400> 15

tgtctctgc atglaatggg cacgttatt agcaacaata gaaggalcaa gcatct 56

<210> 16

<211> 49

<212> DNA

<213> Artificial Sequence

<220>

<223> reverse primer HLBRevSh

<400> 16

aacaatagaa ggatcaagca tctgcacaga aatcaccgaa ggagaagcc 49

<210> 17

<211> 39

<212> DNA

<213> Artificial Sequence

<220>

<223> Forward Primer (underlined), contains 5' detection sequence HLBas-P2

<400> 17

gatgcaaggt cgcatatgag gagcgcgtat gcaatacga 39

<210> 18

<211> 56

<212> DNA

<213> Artificial Sequence

<220>

<223> Reverse Primer (underlined), contains 5' T7 promoter HLBBr-P1

<400> 18

aattctaata cgactcacta tagggagaag ggcggtatcc cgtagaaaaa ggtaga 56

<210> 19

<211> 45

<212> DNA  
<213> Artificial Sequence

<220>  
<223> forward primer hyv1\_For

<400> 19  
ggccgtttta acacaaaaga tgaatatcat agaiggggta gtcaa 45

<210> 20  
<211> 51  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> reverse primer hyvl\_Rev

<400> 20  
ggccatttt agataaatca attgttcta gtttagatac atcaattgt t 51

<210> 21  
<211> 46  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> forward primer

<400> 21  
gttctgtag cgttcagtc ttctcggaa gataaggaat tgcttt 46

<210> 22  
<211> 49  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> reverse primer

<400> 22  
gggcacgttt attagcaaca atagaaggat caagcatctg cacagaaat 49

<210> 23  
<211> 48  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> forward primer

<400> 23  
ctgtagcgt tgcagtctc tgcggaagat aaggaattgc ttctgcg 48

<210> 24  
<211> 51  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> reverse primer

<400> 24  
cacgtttatt agcaacaata gaaggatcaa gcatctgcac agaaatcacc g 51

<210> 25  
<211> 49  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> forward primer

<400> 25  
ggtgttcttg tatcgttgca gtctctgcg gaagataagg aattgctt 49

<210> 26  
<211> 54  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> reverse primer

<400> 26  
gtaatggca cgttattag caacgataga aggatcaagc aactgcacag aaat 54

<210> 27  
<211> 48  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> forward primer

<400> 27  
ctgtatcgt tgcagtctc tgcggaagat aaggaattgc ttctgcg 48

<210> 28  
<211> 53  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> reverse primer

<400> 28  
ggcagttta tttagcaacga tagaaggatc aagcatctgc acagaaatca ccg 53

## REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

### Patent documents cited in the description

- [US4683195A \[0004\]](#)
- [US4683202A \[0004\]](#)
- [US4800158A \[0004\]](#)
- [WO1994023055A1 \[0008\]](#)

- [US9214587A](#) [0049]
- [WO61477357A](#) [0091]
- [WO61477437A](#) [0091]

**Non-patent literature cited in the description**

- **P.WILDINGM. A. SHOFFNERL. J. KRICKA**Clin. Chem., 1994, vol. 40, 1815-1817 [0008]
- **J. B. FINDLAYS. M. ATWOODL. BERGMEYERJ. CHEMELLIK. CHRISTYT. CUMMINSW. DONISHT. EKEZEJ. FALVO D. PATTERSON**Clin. Chem., 1993, vol. 39, 91927-1933 [0008]
- **LUO**Nuc Acids Res, 2005, [0044]
- **WILDING**Nuc. Acids. Res., 1996, [0044]
- **BURKE**Science, 1998, [0044]
- **SAMBROOKRICHMANIATIS**Molecular Cloning and Laboratory ManualCold Spring Harbor20030000 [0049]



## P A T E N T K R A V

1. Fremgangsmåde til amplifikation af en template-nukleinsyre-targetsekvens, hvilken er indeholdt i en prøve, som omfatter:  
5  
i berøring bringning af prøven med en amplifikations-reaktionsblanding omfattende en primer, der er komplementær til template-nukleinsyre-targetsekvensen og et nukleinsyre-destabiliserende middel omfattende i det mindste en DMSO, formamid og betain i en koncentration andragende 8 – 15 vol%, hvor primeren udviser  
10 en smeltetemperatur andragende  $>55^{\circ}\text{C}$ ;  
oscillering af en reaktions-temperatur mellem en øvre denaturerings-temperatur og en lavere renaturerings-temperatur (fra engelsk: annealing-temperatur), hvor en ændring i temperatur inden for et flertal af temperaturocyklings-forløb ikke andrager mere end  $20^{\circ}\text{C}$ , og hvor den øvre denaturerings-temperatur er til-  
15 strækkelig høj til fuldstændig denaturering af templatens; og  
amplifikation af template-nukleinsyre-target-sekvensen.
2. Fremgangsmåde ifølge krav 1, hvor ændringen i temperatur ikke andrager mere end  $15^{\circ}\text{C}$ .  
20
3. Fremgangsmåde ifølge krav 1, hvor temperaturen for en fastlagt tidsperiode, efter at have nået den øvre temperatur eller den lavere temperatur, opretholdes inden for en fluktuation af temperaturen.
- 25 4. Fremgangsmåde ifølge krav 1, hvor temperaturen efter opnåelse af en øvre eller nedre temperatur inden for temperaturområdet underkastes variation til den anden temperatur.
5. Fremgangsmåde ifølge krav 1, hvor den nedre temperatur ikke er mindre end  
30  $50^{\circ}\text{C}$ .
6. Fremgangsmåde ifølge krav 1, hvor den øvre temperatur ikke er højere end  $85^{\circ}\text{C}$ .
7. Fremgangsmåde ifølge krav 1, hvor nukleinsyre-target-sekvens-templatens er enkelt- eller dobbelt-streng DNA eller RNA.  
35
8. Fremgangsmåde ifølge krav 1, hvor target-nukleinsyrens længde kan være mindre end 1000bp.

9. Fremgangsmåde ifølge et hvilket som helst af de forudgående krav, hvor amplifikations-reaktionsblandingen omfatter et par primere, som binder til modsatte streng af nukleinsyre-templatet.
- 5
10. Fremgangsmåde ifølge krav 9, hvor primerparret udviser en længde og et GC-indhold, således at smeltetemperaturen andrager  $\geq 65$  °C.
11. Fremgangsmåde ifølge krav 9, hvor primerparret udviser en længde andragende mellem 35 – 70 basepar, fortrinsvis en længde andragende mellem 40 – 47 basepar.
- 10
12. Fremgangsmåde ifølge et hvilket som helst af de forudgående krav, hvor amplifikations-reaktionsblandingen omfatter:
- 15     DMSO som destabiliserende middel;  
          monovalent kation;  
          divalent kation;  
          dNTP'er; og  
          DNA-polymerase.
- 20
13. Fremgangsmåde ifølge krav 12, hvor den divalente kation er et salt udvalgt blandt gruppen omfattende magnesium, mangan, kobber, zink eller en hvilken som helst kombination deraf.
- 25
14. Fremgangsmåde ifølge krav 12, hvor den monovalente kation er et salt udvalgt blandt gruppen omfattende natrium, kalium, lithium, rubidium, cæsium, ammonium eller en hvilken som helst kombination deraf.
- 30
15. Fremgangsmåde ifølge krav 12, hvor DNA-polymerasen er en termostabil DNA-polymerase.

**DRAWINGS**




FIG. 1A




FIG. 1B




FIG. 2A




FIG. 2B




FIG. 2C




FIG. 2D






FIG. 2E

VentR (exo-) + ET SSB




FIG. 3A




FIG. 3B




FIG. 4




FIG. 5




FIG. 6




FIG. 7




FIG. 8A






FIG. 8B




FIG. 9




FIG. 10




FIG. 11




FIG. 12




FIG. 13




FIG. 14




FIG. 15






FIG. 16A




FIG. 16B




FIG. 17A




FIG. 17B




FIG. 18