

(19) **DANMARK**

(10) **DK/EP 2074205 T4**



(12) **Oversættelse af ændret
europæisk patentskrift**

Patent- og
Varemærkestyrelsen

-
- (51) Int.Cl.: **C 11 D 3/386 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2017-02-06**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om opretholdelse af patentet i ændret form: **2016-11-23**
- (86) Europæisk ansøgning nr.: **07821004.4**
- (86) Europæisk indleveringsdag: **2007-10-08**
- (87) Den europæiske ansøgnings publiceringsdag: **2009-07-01**
- (86) International ansøgning nr.: **EP2007060631**
- (87) Internationalt publikationsnr.: **WO2008040818**
- (30) Prioritet: **2006-10-06 DK 200601307**
- (84) Designerede stater: **AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IS IT LI LT LU LV MC MT NL PL PT RO SE SI SK TR**
- (73) Patenthaver: **Novozymes A/S, Krogshøjvej 36, 2880 Bagsværd, Danmark**
- (72) Opfinder: **MIKKELSEN, Mikael, Valmuehaven 44, 2765 Smørum, Danmark**
RYOM, Niels, Munk, Johannevej 27 st., 2920 Charlottenlund, Danmark
LADEFOGED, Claus, Fruerhoej 29, 2970 Hørsholm, Danmark
FRIIS-JENSEN, Sandra, Ordrupvej 98D, 4. th., 2920 Charlottenlund, Danmark
- (74) Fuldmægtig i Danmark: **Novozymes A/S Patents, Krogshøjvej 36, 2880 Bagsværd, Danmark**
- (54) Benævnelse: **CELLULOTISKE ENZYMESAMMENSÆTNINGER OG ANVENDELSER DERAFF**
- (56) Fremdragne publikationer:
US-A1- 2003 180 933

DESCRIPTION

[0001] The present invention relates to aqueous liquid or gel type detergent compositions comprising specific combinations of enzymes. The detergent compositions may further comprise a combination of boric acid or a boron compound capable of forming boric acid in the composition, a polyhydroxy compound, preferably propanediol, and relatively high level of calcium ion to stabilize a selected combination of a protease enzyme and other enzymes. The invention also relates to a process for enhancing stability of the non protease enzymes in combination of a protease enzyme with other enzymes in a liquid or gel detergent composition. The invention further relates to the use of specific protease enzymes in detergent compositions

BACKGROUND ART

[0002] Proteases have been used in detergent compositions for about 50 years and a number of such proteases have in the past 10 years been developed by protein engineering of a number of precursor proteases.

[0003] The most successful precursor protease on the market is subtilisin 309 - or Savinase®. Protein engineering of Savinase was first disclosed in 1989 in WO 89/06279. Subsequently a high number of patent applications relating to protein engineering of Savinase have been filed by the applicant and other companies, such as Genencor International, Inc., Procter & Gamble, Unilever NV, etc. Also, a number of Savinase variants have been marketed by Novozymes A/S and Genencor International, Inc.

[0004] The specific Savinase variant comprising the modifications Y167A+R170S+A194P was disclosed in WO 98/20115. In the present application we designate this variant subtilisin KL.

[0005] Aqueous liquid and gel detergent compositions containing enzymes, including proteases, are well known in the art. The major problem encountered with such compositions is that of ensuring a sufficient storage stability of the enzymes in the compositions. It is particularly difficult to stabilize amylases in the presence of proteases, which can readily degrade amylases in aqueous liquid or gel detergent compositions but also other enzymes, such as lipases, cellulases, etc. are frequently degraded by the proteases.

[0006] High-alkaline amylases such as alpha amylases are described in British Specification No. 1,296,839. The use of an enzyme stabilizing system comprising a mixture of boric acid or an alkali metal borate with calcium ion, and preferably with a polyol, is disclosed in U.S. Patent 4,537,706, Severson. Certain α -amylases that provide improved cleaning and stain removal are disclosed in WO97/32961, Baeck et al., and in WO 96/23873 and U.S. Patent 6,093,562.

[0007] US2003/180933 describes liquid and or gel detergent compositions comprising a subtilase variant comprising the mutations A167A + R170S + A194P in combination with other enzymes.

DISCLOSURE OF THE INVENTION

[0008] The present invention relates to a liquid or gel composition comprising subtilisin KL or variants thereof in combination with at least one lipase; amylase; cellulase; or mannanase, wherein the weight ratio between the content of subtilisin KL or variants thereof to the content of lipase, amylase, cellulase or mannanase is from 0.001 to 100, preferably from 0.01 to 10, especially from 0.5 to 5, especially from 1 to 3, wherein the subtilisin KL variant is one of the group defined in claim 1. In a particular embodiment the content of subtilisin KL or variants thereof is from 0.001 to 5 weight% and if present the content of each of the following lipase, amylase, cellulase, or mannanase, is from 0.001 to 5 weight%.

[0009] Another embodiment of the invention relates to the use of subtilisin KL or variants thereof in combination with at least one, lipase, amylase, cellulase or mannanase, for the preparation of aqueous liquid or gel type detergent compositions having enhanced stability of the non protease enzymes, wherein the subtilisin KL variant is one of the group defined in claim 6.

[0010] Yet another embodiment of the invention relates to a process for enhancing stability of the non protease enzymes in combination of a protease enzyme with other enzymes in a liquid or gel detergent composition comprising a protease and at least one non protease enzyme, wherein the liquid or gel detergent composition is prepared using subtilisin KL or a variant thereof as the protease enzyme and wherein the at least one non protease enzyme is selected among lipase, amylase, cellulase or mannanase and wherein the subtilisin KL variant is one of the group defined in claim 7.

[0011] In particular embodiment of the invention concerns

[0012] The amylases to be used in the detergent compositions of the invention are the amylase from *B. licheniformis* and other amylases, such as those disclosed in WO 2001/066712, WO 2006/002643, WO 2000/60060.

[0013] The cellulases to be used in the detergent compositions of the invention are such as those disclosed in WO 1995/024471, WO 91/17244, WO 2002/099091.

[0014] The lipases to be used in the detergent compositions of the invention are such as those disclosed in WO 2000/060063.

[0015] The mannanases to be used in the detergent compositions of the invention are such as those disclosed in WO 99/64619, e.g. SEQ ID NO: 2.

[0016] The endoglucanase to be used in the detergent compositions of the invention are such as those disclosed in WO 91/17244

[0017] The subtilisin KL variants of the present invention are those indicated in Table 1, which are also disclosed in WO 98/20115:

Table 1

Mutations in subtilisin KL
None
*36D
P14T
N18K
N62D
V83L
A133P
E136Q
E136R
E136K
N140R
N140K
S141E
S141N
S141Y
S141R
T143R
T143K
S153R
S156R
A160R
S162R
S162K
I165R
I165K
Y171R
Y171K
A172R

Mutations in subtilisin KL
A172K
A174R
N173R
N173K
A174K
N76D
Y176R
Y176K
A187R
A187K
S188P
S190P
Q191R
Y192R
Y192R
Q191P
Y192A
Y192P
D197N
D197R
D197E
D197K
D197G
A228V
A230V
T260R
T260K
G264R
G264K
S265T
S265R
S265K
N218S
M222S
M222A
M222G
M222T
M222V
M222S
N243R
V244R
N248R
K251R

Mutations in subtilisin KL
N252R
N261R
Combinations
S9R+A15T+T22A+N218S+K251R
S9R+A15T+T22A+V841+N218S
V30H+V139L +N218S
V84H+V139L+N218S
N76D+N218S
N76D+A228V
N76D+A230V
N76D+N218S+A230V
N76D+A228V+A230V
N218S+R247Q
N218S+R247H
N218S+R247E
N218S+R247K
D181N+N218S
N218S+A230V
K251R+S265K
P14T+N18K
T274H+R275H+*275aH+*275bH+*275cH+*275dH=
T274H+R275HHHHH
T274H+R275H+*275aH+*275bH+*275cH=T274H+R275HHHH
S87N+S101G,V104N
*36D+N76D+H120D+G195E+K235L
A133P+M222S
Insertions and combinations therewith
*96aA
*96aA+A98T
*96aA+A133P
*96aA+A98T+A133P
*96aA+A98T+N218S
*97aP+A98T+N218S
*98aT,
*98aT+S99N+N218S
G97D+*98aT+N218S
*99aE=S99SE
*99aD=S99SD
*99aD+M222S=S99SD+M222S
N76D+s99A+*99aE=N76D+S99AE
N76D+*99aD+A230V=N76D+S99SD+A230V
S99A+*99aD=S99AD
S99A+*99aD+M222S=S99AD+M222S

Mutations in subtilisin KL
S99A+*99aD+N218S=S99AD+N218S
S99A+*99aE+A230V=S99AE+A230V
A228V+A230V
*130aL+P194A

[0018] It has surprisingly been found that subtilisin KL and the above variants thereof exhibit a remarkable compatibility to other enzymes used in liquid detergent compositions such as lipases, amylases, cellulases, peroxidases/oxidases and hemicellulases. This property results in a substantial increase in the residual activity of these enzymes in combination with subtilisin KL and the above variants thereof as compared to the residual activity in the presence of other proteases, even after long periods of storage. In the end the result is an improved performance of the detergent composition or that similar results can be obtained with reduced amounts of enzyme

NOMENCLATURE AND CONVENTIONS FOR DESIGNATION OF VARIANTS

[0019] In describing the various subtilisin KL enzyme variants produced or contemplated according to the invention, the following nomenclatures and conventions have been adapted for ease of reference: A frame of reference is first defined by aligning the parent enzyme with subtilisin BPN (BASBPN).

[0020] The alignment can be obtained by the GAP routine of the GCG package version 9.1 to number the variants using the following parameters: gap creation penalty = 8 and gap extension penalty = 8 and all other parameters kept at their default values.

[0021] Another method is to use known recognized alignments between subtilases, such as the alignment indicated in WO 91/00345. In most cases the differences will not be of any importance.

[0022] Thereby a number of deletions and insertions will be defined in relation to BASBPN (SEQ ID NO.1). For a detailed description of the nomenclature of modifications introduced in a polypeptide by genetic manipulation we refer to WO 00/71691 page 7-12.

[0023] Numbering of amino acid positions/residues If nothing else is mentioned the amino acid numbering used herein correspond to that of the subtilase BPN' (BASBPN) sequence. For further description of the BPN' sequence, see Siezen et al., Protein Engng. 4 (1991) 719-737.

[0024] "SAVINASE®" Savinase® is marketed by Novozymes A/S. It is subtilisin 309 from B. Lentus.

[0025] Modification(s) of a subtilisin KL variant. The term "modification(s)" used herein is defined to include chemical modification as well as genetic manipulation of the DNA encoding subtilisin KL. The modification(s) can be replacement(s) of the amino acid side chain(s), substitution(s), deletion(s) and/or insertions in or at the amino acid(s) of interest.

[0026] Subtilase variant. In the context of this invention, the term subtilase variant or mutated subtilase means a subtilase that has been produced by an organism which is expressing a mutant gene derived from a parent microorganism which possessed an original or parent gene and which produced a corresponding parent enzyme, the parent gene having been mutated in order to produce the mutant gene from which said mutated subtilase protease is produced when expressed in a suitable host.

[0027] Homologous subtilase sequences. The homology between two amino acid sequences is in this context described by the parameter "identity". In order to determine the degree of identity between two subtilases the GAP routine of the GCG package version 9.1 can be applied (infra) using the same settings. The output from the routine is besides the amino acid alignment the calculation of the "Percent Identity" between the two sequences. Based on this description it is routine for a person skilled in the art to identify suitable homologous subtilases, which can be modified according to the invention.

[0028] Isolated polynucleotide. The term "isolated", when applied to a polynucleotide, denotes that the polynucleotide has been removed from its natural genetic milieu and is thus free of other extraneous or unwanted coding sequences, and is in a form suitable for use within genetically engineered protein production systems. Such isolated molecules are those that are separated

from their natural environment and include cDNA and genomic clones. Isolated DNA molecules of the present invention are free of other genes with which they are ordinarily associated, but may include naturally occurring 5' and 3' untranslated regions such as promoters and terminators. The identification of associated regions will be evident to one of ordinary skill in the art (see for example, Dynan and Tjian, Nature 316:774-78, 1985). The term "an isolated polynucleotide" may alternatively be termed "a cloned polynucleotide".

[0029] Isolated protein. When applied to a protein, the term "isolated" indicates that the protein has been removed from its native environment. In a preferred form, the isolated protein is substantially free of other proteins, particularly other homologous proteins (i.e. "homologous impurities" (see below)). An isolated protein is more than 10% pure, preferably more than 20% pure, more preferably more than 30% pure, as determined by SDS-PAGE. Further it is preferred to provide the protein in a highly purified form, i.e., more than 40% pure, more than 60% pure, more than 80% pure, more preferably more than 95% pure, and most preferably more than 99% pure, as determined by SDS-PAGE. The term "isolated protein" may alternatively be termed "purified protein".

[0030] Homologous impurities. The term "homologous impurities" means any impurity (e.g. another polypeptide than the subtilase of the invention), which originate from the homologous cell where the subtilase of the invention is originally obtained from.

[0031] Obtained from. The term "obtained from" as used herein in connection with a specific microbial source, means that the polynucleotide and/or subtilase produced by the specific source, or by a cell in which a gene from the source has been inserted.

[0032] Substrate. The term "substrate" used in connection with a substrate for a protease should be interpreted in its broadest form as comprising a compound containing at least one peptide (amide) bond susceptible to hydrolysis by a subtilisin protease.

[0033] Product. The term "product" used in connection with a product derived from a protease enzymatic reaction should, in the context of the present invention, be interpreted to include the products of a hydrolysis reaction involving a subtilase protease. A product may be the substrate in a subsequent hydrolysis reaction.

[0034] Wash Performance. In the present context the term "wash performance" is used as an enzyme's ability to remove proteinaceous or organic stains present on the object to be cleaned during e.g. wash or hard surface cleaning.

[0035] The detergent composition of the invention may for example be formulated as a hand or machine laundry detergent composition including a laundry additive composition suitable for pre-treatment of stained fabrics and a rinse added fabric softener composition, or be formulated as a detergent composition for use in general household hard surface cleaning operations, or be formulated for hand or machine dishwashing operations.

[0036] In general the properties of the chosen enzyme(s) should be compatible with the selected detergent, (i.e. pH-optimum, compatibility with other enzymatic and non-enzymatic ingredients, etc.), and the enzyme(s) should be present in effective amounts.

[0037] Lipases: Suitable lipases include those of bacterial or fungal origin. Chemically modified or protein engineered mutants are included. Examples of useful lipases include lipases from *Humicola* (synonym *Thermomyces*), e.g. from *H. insolens* as described in WO 96/13580, a *Pseudomonas* lipase, e.g. from *Pseudomonas* sp. strain SD 705 (WO 95/06720 and WO 96/27002), *P. wisconsinensis* (WO 96/12012), or a *Bacillus* lipase as disclosed in WO 2000/060063.

[0038] Other examples are lipase variants such as those described in WO 92/05249, WO 94/01541, EP 407225, EP 260105, WO 95/35381, WO 96/00292, WO 95/30744, WO 94/25578, WO 95/14783, WO 95/22615, WO 97/04079 and WO 97/07202. Preferred commercially used lipase enzymes include Lipolase®, Lipolase Ultra® and Lipex® (Novozymes A/S).

[0039] Amylases: Suitable amylases (α and/or β) include those of bacterial or fungal origin. Chemically modified or protein engineered mutants are included. Amylases include, for example, α -amylases obtained from *Bacillus*. Examples of useful amylases are the variants described in WO 94/02597, WO 94/18314, WO 96/23873, WO 2000/60060, and WO 97/43424, especially the variants with substitutions in one or more of the following positions: 15, 23, 105, 106, 124, 128, 133, 154, 156, 181, 188, 190, 197, 202, 208, 209, 243, 264, 304, 305, 391, 408, and 444. Commercially used amylases are Duramyl®, Termamyl®, Stainzyme®, Stainzyme Plus®, Stainzyme ultra®, Fungamyl® and BAN® (Novozymes A/S), Rapidase™, Purastar™ and Purastar OxAm™ (from Genencor International Inc.).

[0040] Cellulases: Suitable cellulases include those of bacterial or fungal origin. Chemically modified or protein engineered

mutants are included. Suitable cellulases include cellulases from the genera *Bacillus*, *Pseudomonas*, *Humicola*, *Fusarium*, *Thielavia*, *Acremonium*, e.g. the fungal cellulases produced from *Humicola insolens*, *Myceliophthora thermophila* and *Fusarium oxysporum* disclosed in US 5,648,263, US 5,691,178, US 5,776,757 and WO 89/09259. Especially suitable cellulases are the alkaline or neutral cellulases having colour care and whiteness maintenance benefits. Examples of such cellulases are cellulases described in EP 0 531 372, WO 96/11262, WO 96/29397, WO 98/08940. Other examples are cellulase variants such as those described in WO 94/07998, EP 0 531 315, US 5,457,046, US 5,686,593, US 5,763,254, WO 95/24471, WO 98/12307 and PCT/DK98/00299. Commercially used cellulases include Renozyme®, Celluzyme®, Celluclean®, Endolase® and Carezyme® (Novozymes A/S), Clazinase™, and Puradax HA™ (Genencor Int. Inc.), and KAC-500(B)™ (Kao Corporation).

[0041] Peroxidases/Oxidases: Suitable peroxidases/oxidases include those of plant, bacterial or fungal origin. Chemically modified or protein engineered mutants are included. Examples of useful peroxidases include peroxidases from *Coprinus*, e.g. from *C. cinereus*, and variants thereof as those described in WO 93/24618, WO 95/10602, and WO 98/15257. Commercially used peroxidases include Guardzyme™ (Novozymes A/S).

[0042] Hemicellulases: Suitable hemicellulases include those of bacterial or fungal origin. Chemically modified or protein engineered mutants are included. Suitable hemicellulases include mannanase, lichenase, xylanase, arabinase, galactanase, acetyl xylan esterase, glucuronidase, ferulic acid esterase, coumaric acid esterase and arabinofuranosidase as described in WO 95/35362. Suitable mannanases are described in WO 99/64619. Commercially used hemicellulases include Mannaway® (Novozymes A/S).

[0043] The detergent enzyme(s) may be included in a detergent composition by adding separate additives containing one or more enzymes, or by adding a combined additive comprising all of these enzymes. A detergent additive of the invention, i.e. a separate additive or a combined additive, can be formulated e.g. as a gel, a liquid, a slurry, etc. Preferred detergent additive formulations are liquids, in particular stabilized liquids, or slurries.

[0044] Liquid enzyme preparations may, for instance, be stabilized by adding a polyol such as propylene glycol, a sugar or sugar alcohol, lactic acid or boric acid according to established methods. Protected enzymes may be prepared according to the method disclosed in EP 238,216.

[0045] The detergent composition of the invention is in the form of a gel or a liquid. A liquid detergent may be aqueous, typically containing up to 70 % water and 0-30 % organic solvent, or non-aqueous.

[0046] The detergent composition comprises one or more surfactants, which may be non-ionic including semi-polar and/or anionic and/or cationic and/or zwitterionic. The surfactants are typically present at a level of from 0.1 % to 60% by weight.

[0047] When included therein the detergent will usually contain from about 1% to about 40% of an anionic surfactant such as linear alkylbenzenesulfonate, alpha-olefinsulfonate, alkyl sulfate (fatty alcohol sulfate), alcohol ethoxysulfate, secondary alkanesulfonate, alpha-sulfo fatty acid methyl ester, alkyl- or alkenylsuccinic acid or soap.

[0048] When included therein the detergent will usually contain from about 0.2% to about 40% of a non-ionic surfactant such as alcohol ethoxylate, nonylphenol ethoxylate, alkylpolyglycoside, alkyldimethylamineoxide, ethoxylated fatty acid monoethanolamide, fatty acid monoethanolamide, polyhydroxy alkyl fatty acid amide, or N-acyl N-alkyl derivatives of glucosamine ("glucamides").

[0049] The detergent may contain 0-65 % of a detergent builder or complexing agent such as zeolite, diphosphate, triphosphate, phosphonate, carbonate, citrate, nitrilotriacetic acid, ethylenediaminetetraacetic acid, diethylenetriaminepentaacetic acid, alkyl- or alkenylsuccinic acid, soluble silicates or layered silicates (e.g. SKS-6 from Hoechst).

[0050] The detergent may comprise one or more polymers. Examples are carboxymethylcellulose, poly(vinylpyrrolidone), poly(ethylene glycol), poly(vinyl alcohol), poly(vinylpyridine-N-oxide), poly(vinylimidazole), polycarboxylates such as polyacrylates, maleic/acrylic acid copolymers and lauryl methacrylate/acrylic acid copolymers.

[0051] The detergent may contain a bleaching system which may comprise a H₂O₂ source such as perborate or percarbonate which may be combined with a peracid-forming bleach activator such as tetraacetylenediamine or nonanoyloxybenzenesulfonate. Alternatively, the bleaching system may comprise peroxyacids of e.g. the amide, imide, or sulfone type.

[0052] The enzyme(s) of the detergent composition of the invention may be stabilized using conventional stabilizing agents, e.g.,

a polyol such as propylene glycol, diethylene glycol, methylpropanediol, or glycerol, a sugar or sugar alcohol, lactic acid, boric acid, or a boric acid derivative, e.g., an aromatic borate ester, or a phenyl boronic acid derivative such as 4-formylphenyl boronic acid or mono- or triethanolamine, and the composition may be formulated as described in e.g. WO 92/19709, WO 92/19708, US 5,972,873 or EP 0832174.

[0053] The detergent may also contain other conventional detergent ingredients such as e.g. fabric conditioners including clays, foam boosters, suds suppressors, anti-corrosion agents, soil-suspending agents, anti-soil redeposition agents, dyes, bactericides, optical brighteners, hydrotropes, tarnish inhibitors, or perfumes.

[0054] It is at present contemplated that in the detergent compositions any enzyme, in particular the enzyme of the invention, may be added in an amount corresponding to 0.01-100 mg of enzyme protein per litre of wash liquor, preferably 0.05-5 mg of enzyme protein per litre of wash liquor, in particular 0.1-1 mg of enzyme protein per litre of wash liquor.

[0055] Variations in local and regional conditions, such as water hardness and wash temperature call for regional detergent compositions. Detergent Examples 1 provide ranges for the composition of a liquid detergent.

Materials and Methods

Enzymes

[0056] In the examples below the following commercial available enzymes are used. Alcalase® and Savinase® are used as standards for comparison:

Name	Enzyme type	Derived from or disclosed in
Alcalase®	Protease, subtilisin Carlsberg	B. licheniformis
Savinase®	Protease, subtilisin 309	B. lentus
Termamyl®	amylase	B.licheniformis
Novozym 342®		H. Insolens
Amylase A	amylase	The amylase variant D183*+G184*+R118K+N195F+R458K. WO 01/66712
Mannan A	Mannanase	WO 99/64619
Lipase A	Lipase	T231 R+N233R variant of T. lanuginosus lipase, WO00/60063
Cellulase A	Cellulase	H. Insolens, WO 91/17244

Also the protease designated subtilisin KL and variants thereof are used. Subtilisin KL is a Y167A+R170S+A194P variant of Savinase (using BPN numbering)

Assays

Protease Compatibility:

[0057] The protease compatibility of the enzymes is determined by preparing the detergent compositions as indicated in each Example and measuring the residual activity of the other enzyme activities after the periods indicated in the Examples.

Enzyme Activity:

[0058] Enzyme activities are measured using well known recognized standard methods.

Detergent Compositions

[0059] The detergent compositions used in the examples are either a model detergent according to the compositions provided below or commercial liquid laundry detergents e.g. Tide, Era, Gain, Cheer, Wisk, All, Purex, Arm & Hammer, Sun, Great Value, Ariel, Persil, Total, Skip, Dash, Dixan, Ava or any other brand extension or concentrated versions for the liquid detergent. If the commercial laundry detergent used comprises enzymes these are inactivated prior to use by heating the detergent in a microwave oven at 85°C for 5 minutes. Model detergent composition A - Detergent Example 1

Group	Subname	Content
Surfactants		5-60%
	Sulphonates	0-30%
	Sulphates	0-15%
	Soaps	0-15%
	Non-ionics	0-15%
	Cationics	0-15%
	Amine oxides	0-10%
	FAGA	0-10%
Solvents		5-35%
	Ethanol	0-10%
	MPG - monopropylene glycol	0-20%
	DEG- Diethylene glycol	0-15%
	MPD - methylpropanediol	0-15%
	MEA - Monoethanolamine	0-10%
	TEA - Triethanolamine	0-10%
	Hydrotropes like SXS, SCS, etc	
	Sodium Cumene Sulfonate	
	Sodium Xylene Sulfonates	0-10%
	Other solvents	0-10%
Builders		0-20%
	NaCitrate	0-15%
	Other builders	0-15%
Others		0-20%
	Polymers	0-5%
	Enzymes	0-10%
	Boric acid and derivatives thereof	0-5%
Builders		0-20%
	Foam Regulators	0-10%
	Others	0-10%
Water is added to the balance of 100%		

Example 1

[0060] A commercial liquid detergent for laundry was added commercial proteases, amylases, Lipase, and cellulases as listed below (if the detergent already contains enzymes then these can be inactivated by heating the detergent in a microwave oven up to 85°C for 5 minutes). When Subtilisin KL was used in comparison with commercial protease, same amount of activity units was used.

[0061] The stability of the enzymes as determined by % residual enzyme activity after storage at 20°C for 1, 2 and 4 weeks is shown in table 2-5.

[0062] Storage conditions: 20°C for 1, 2, 4 weeks in closed glass vessels

Table 2 Residual amylase activity

Weeks	1	2	3	4
0.5% Alcalase Ultra 2.5 L				
0.3% Termamyl 300L	93	92	89	87
Subtilisin KL				
0.3% Termamyl 300 L	96	98	95	92
0.5% Alcalase Ultra 2.5 L				
0.3% Amylase A 12L	34	16	10	7
Subtilisin KL				
0.3% Amylase A 12 L	90	86	82	78

Table 3 Residual lipase activity

Weeks	1	2	3	4
0.5% Alcalase Ultra 2.5 L				
0.3% Lipase A 100 L	12	11	8	9
Subtilisin KL				
0.3% Lipase A 100 L	72	54	46	38

Table 4 Residual cellulase activity

Weeks	1	2	3	4
0.5% Alcalase Ultra 2.5 L				
0.3% Cellulase A 5000 L		85	76	68
Subtilisin KL				
0.3% Cellulase A 5000 L		99	87	88

Table 5 Residual protease activity

Weeks	1	2	3	4
0.5% Alcalase Ultra 2.5 L				
0.3% Cellulase A 5000 L	86	64	57	50
Subtilisin KL				
0.3% Cellulase A 5000 L	84	74	65	56

[0063] As can be seen above the enzyme compatibility of the present invention is clearly improved when Subtilisin KL is selected as the protease instead of Alcalase 2.5L. The enzyme stability of Cellulase A 5000L, Lipase A 100L, Termamyl 300L and Amylase A 12L after 1, 2, 3 and 4 weeks at 30°C is clearly improved if Subtilisin KL is the protease. The Subtilisin KL protease is just as stable as the reference protease, Alcalase 2.5L, used.

Example 2

[0064] The commercial liquid detergent for laundry of Example 1 was added commercial proteases, amylases, Lipase, and cellulases as listed below (if the detergent already contains enzymes then these are inactivated by heating the detergent in a micro oven up to 85°C for 5 minutes). When Subtilisin KL was used in comparison with commercial protease, same amount of activity units was used.

[0065] The stability of the enzymes as determined by % residual enzyme activity after storage at 30°C for 1, 2 and 4 weeks in shown in table 6-9.

Table 6 Residual amylase activity

Weeks	1	2	3	4
0.5% Alcalase Ultra 2.5 L				
0.3% Termamyl 300L	85	78	71	66
Subtilisin KL				
0.3% Termamyl 300 L	93	87	83	73
0.5% Alcalase Ultra 2.5 L				
0.3% Amylase A 12L	10	5	4	4
Subtilisin KL				
0.3% Amylase A 12 L	81	74	63	59

Table 7 Residual lipase activity

Weeks	1	2	3	4
0.5% Alcalase Ultra 2.5 L				
0.3% Lipase A 100 L	9	8	5	6
Subtilisin KL				
0.3% Lipase A 100 L	35	17	11	6

Table 8 Residual cellulase activity

Weeks	1	2	3	4
0.5% Alcalase Ultra 2.5 L				
0.3% Cellulase A 5000 L	47	24	16	13
Subtilisin KL				
0.3% Cellulase A 5000 L	67	66	55	55

Table 9 Residual protease activity

Weeks	1	2	3	4
0.5% Alcalase Ultra 2.5 L	57	36	29	21
Subtilisin KL	55	36	24	16

[0066] As can be seen above the enzyme compatibility of the present invention is clearly improved when Subtilisin KL is selected as the protease instead of Alcalase 2.5L. The enzyme stability of Cellulase A 5000L, Lipase A 100L, Termamyl 300L and Amylase A 12L after 1, 2, 3 and 4 weeks at 30°C is clearly improved if Subtilisin KL is selected as protease. The Subtilisin KL protease is just as stable as the reference protease, Alcalase 2.5L, used.

Example 3

[0067] A commercial liquid detergent for laundry was added commercial proteases, amylases, and lipases as listed below (if the detergent already contains enzymes then these can be inactivated by heating the detergent in a micro oven up to 85°C for 5 minutes). When Subtilisin KL was used in comparison with commercial protease, same amount of activity units was used.

[0068] The stability of the enzymes as determined by % residual enzyme activity after storage at 30°C for 1, 2, 4 and 8 weeks is shown in table 10-11.

Table 10 Residual amylase activity

Weeks	1	2	4	8
0.4% Alcalase 2.5 L				
0.4% Amylase A 12 L	42	36	19	9
0.4% Savinase 16 L				
0.4% Amylase A 12 L	48	41	24	9
Subtilisin KL				

Weeks	1	2	4	8
0.4% Amylase A	77	73	63	42
0.4% Amylase A 12 L				
(without protease)	88	89	82	62

Table 11 Residual lipase activity

Weeks	1	2
0.4% Alcalase 2.5 L		
0.4% Lipase A 100 L	9	8
Subtilisin KL		
0.4% Lipase A 100 L	33	22
0.4% Lipase A 100 L		
(without protease)	86	81

[0069] As can be seen above the enzyme compatibility of the present invention is clearly improved when Subtilisin KL is selected as the protease instead of Savinase 16L and Alcalase 2.5L. The enzyme stability of Lipase A 100L and Amylase A 12L after 2 and 8 weeks is improved significantly if Subtilisin KL is selected as the preferred protease.

Example 4

[0070] A liquid detergent with the following formulation as shown in table 13 is prepared.

Table 13 Detergent formulation

Subname	Content
Calcium Chloride	0,1%
LAS-Sodium Salt	11,81%
Soya sebacic acid - sodium salt	5,94%
Propyleneglycol	5,05%
C-13-Oxoalcohol ethoxylat, 8EO	9,45%
Phosphonate	1,00%
Coconut sebacic acid - Triethanolamine salt	6,50%
Sodium citrate	1,00%
Ethanol	4,63%
Opacifier	0,12%
Perfume	0,35%
Colour	-
Water to 100%	

Enzymes used

[0071]

Protease:	Savinase 16L
	Alcalase 2.5L
	Subtilisin KL
	Subtilisin KL M222S
	Subtilisin KL *36D

	Subtilisin KL N76D+S99SE+A230V
	Subtilisin KL S162R
	Subtilisin KL S99SE+N76D
	Subtilisin KL N76D
	Subtilisin KL A228V
	Subtilisin KL A230V
	Subtilisin KL A228V+A230V
Lipase:	Lipase A 100L
Amylase:	Termamyl 300L
Mannase:	Mannan A4,0L

Test set-up I**[0072]**

Addition of enzymes:	I) Savinase 16L (0,17mg EP/g)
	II) Subtilisin KL (0,17mg EP/g)
	III) Alcalase 2,5L(0,17mg EP/g)
Amylase :	Termamyl 300L (0,4%)

The amounts of protease are given in enzyme protein (active) per grammes [EP/g].

[0073] The detergent formulations are stored in 2, and 4 weeks at 30°C in closed glass vessels. After storage the residual protease and amylase activities are determined.

Table 14 % Residual Protease activity

Weeks	2	4
0,17mg Savinase 16L + 0.4% Termamyl 300L	21	15
0,17mg Alcalase 2,5L + 0.4% Termamyl 300L	23	16
0,17mg Subtilisin KL + 0.4% Termamyl 300L	16	10

Table 15 % Residual Amylase activity

Weeks	2	4
0,17mg Savinase 16L + 0.4% Termamyl 300L	90	92
0,17mg Alcalase 2,5L + 0.4% Termamyl 300L	94	95
0,17mg Subtilisin KL + 0.4% Termamyl 300L	97	97

Test set-up II**[0074]**

Addition of enzymes:	I) Savinase 16L (0,07mg EP/g)
	II) Subtilisin KL (0,07mg EP/g)
	III) Alcalase 2,5L (0,07mg EP/g)
	IV) Subtilisin 2,5KL M222S (0,07mg EP/g)
	V) Subtilisin 2,5KL *36D (0,07mg EP/g)
	VI) Subtilisin KL N76D+S99SE, A230V
Lipase :	Lipase A 100L (0,2%)

Amylase:	Termamyl 300L (0,2%)
Mannase:	Mannan A 4,0L (0,2%)

[0075] The detergent formulations are stored in 2, and 4 weeks at 30°C in closed glass vessels. After storage the residual protease, lipase (Lip.), mannase (Man.) and amylase (Ter.) activities are determined.

Table 16 % Residual Protease activity

Weeks	2	4
0,07mg Savinase 16L		
0.2% Ter., 0.2% Lip. and 0.2% Man.	21	13
0,07mg Alcalase 2,5L		
0.2% Ter., 0.2% Lip. and 0.2% Man.	24	22
0,07mg Subtilisin KL		
0.2% Ter., 0.2% Lip. and 0.2% Man.	18	13
0,07mg Subtilisin KL M222S		
0.2% Ter., 0.2% Lip. and 0.2% Man.	50	50
0,07mg Subtilisin KL *36D		
0.2% Ter., 0.2% Lip. and 0.2% Man.	59	19
0,07mg Subtilisin KL N76D+S99SE+A230V		
0.2% Ter., 0.2% Lip. and 0.2% Man.	84	77

Table 17 % Residual Amylase activity

Weeks	2	4
0,07mg Savinase 16L		
0.2% Ter., 0.2% Lip. and 0.2% Man.	97	96
0,07mg Alcalase 2,5L		
0.2% Ter., 0.2% Lip. and 0.2% Man.	87	89
0,07mg Subtilisin KL		
0.2% Ter., 0.2% Lip. and 0.2% Man.	97	97
0,07mg Subtilisin KL M222S		
0.2% Ter., 0.2% Lip. and 0.2% Man.	98	101
0,07mg Subtilisin KL *36D		
0.2% Ter., 0.2% Lip. and 0.2% Man.	97	98
0,07mg Subtilisin KL N76D+S99SE+A230V		
0.2% Ter., 0.2% Lip. and 0.2% Man.	98	98

Table 18 % Residual Lipase activity

Weeks	2	4
0,07mg Savinase 16L		
0.2% Ter., 0.2% Lip. and 0.2% Man.	5	5
0,07mg Alcalase 2,5L		
0.2% Ter., 0.2% Lip. and 0.2% Man.	5	5
0,07mg Subtilisin KL		
0.2% Ter., 0.2% Lip. and 0.2% Man.	4	4
0,07mg Subtilisin KL M222S		
0.2% Ter., 0.2% Lip. and 0.2% Man.	20	15
0,07mg Subtilisin KL *36D		
0.2% Ter., 0.2% Lip. and 0.2% Man.	6	6

Weeks	2	4
0,07mg Subtilisin KL N76D+S99SE+A230V		
0.2% Ter., 0.2% Lip. and 0.2% Man.	22	17

Table 19 % Residual Mannase activity

Weeks	2	4
0,07mg Savinase 16L		
0.2% Ter., 0.2% Lip. and 0.2% Man.	38	25
0,07mg Alcalase 2,5L		
0.2% Ter., 0.2% Lip. and 0.2% Man.	14	13
0,07mg Subtilisin KL		
0.2% Ter., 0.2% Lip. and 0.2% Man.	62	48
0,07mg Subtilisin KL M222S		
0.2% Ter., 0.2% Lip. and 0.2% Man.	89	84
0,07mg Subtilisin KL *36D		
0.2% Ter., 0.2% Lip. and 0.2% Man.	63	54
0,07mg Subtilisin KL N76D+S99SE+A230V		
0.2% Ter., 0.2% Lip. and 0.2% Man.	99	95

Test set-up III**[0076]**

Addition of enzymes:	I) Savinase 16L (0,05mg EP/g det.)
	II) Subtilisin KL (0,05mg EP/g det.)
	III) Alcalase 2,5L (0,05mg EP/g det.)
	VII) Subtilisin 2,5KL S162R (0,05mg EP/g det.)
	VIII) Subtilisin KL S99SE+N76D (0,05mg EP/g det.)
	IX) Subtilisin KL N76D (0,05mg EP/g det.)
	X) Subtilisin KL A228V (0,05mg EP/g det.)
	XI) Subtilisin KL A230V (0,05mg EP/g det.)
	XII) Subtilisin KL A228V, A230V (0,05mg EP/g det.)
	EP = Enzyme Protein
	det = detergent
Lipase:	Lipase A 100L (0,2%)
Amylase:	Termamyl 300L (0,2%)
Mannase:	Mannan A 4.0L (0,2%)

[0077] The detergent formulations are stored in 1, 2 and 3 weeks at 30°C in closed glass vessels. After storage the residual protease, lipase (Lip.), mannase (Man.) and amylase (Ter.) activities are determined.

Table 20 % Residual Protease activity

Weeks	1	2	3
0,05mg Savinase 16L			
0.2% Ter., 0.2% Lip. and 0.2% Man.	89	20	12

Weeks	1	2	3
0,05mg Alcalase 2,5L			
0.2% Ter., 0,2% Lip. and 0,2% Man.	85	37	37
0,05mg Subtilisin KL			
0.2% Ter., 0,2% Lip. and 0,2% Man.	70	17	17
0,05mg Subtilisin KL S162R			
0.2% Ter., 0,2% Lip. and 0,2% Man.	45	12	12
0,05mg Subtilisin KL S99SE+N76D			
0.2% Ter., 0,2% Lip. and 0,2% Man.	100	75	77
0,05mg Subtilisin KL N76D			
0.2% Ter., 0,2% Lip. and 0,2% Man.	94	95	89
0,05mg Subtilisin KL A228V			
0.2% Ter., 0,2% Lip. and 0,2% Man.	85	83	78
0,05mg Subtilisin KL A230V			
0.2% Ter., 0,2% Lip. and 0,2% Man.	99	87	80
0,05mg Subtilisin KL A228V+A230V			
0.2% Ter., 0,2% Lip. and 0,2% Man.	100	98	89

Table 21 % Residual Amylase activity

Weeks	1	2	3
0,05mg Savinase 16L			
0.2% Ter., 0,2% Lip. and 0,2% Man.	100	98	96
0,05mg Alcalase 2,5L			
0.2% Ter., 0,2% Lip. and 0,2% Man.	100	96	97
0,05mg Subtilisin KL			
0.2% Ter., 0,2% Lip. and 0,2% Man.	100	98	97
0,05mg Subtilisin KL S162R			
0.2% Ter., 0,2% Lip. and 0,2% Man.	99	97	97
0,05mg Subtilisin KL S99SE+N76D			
0.2% Ter., 0,2% Lip. and 0,2% Man.	99	98	98
0,05mg Subtilisin KL N76D			
0.2% Ter., 0,2% Lip. and 0,2% Man.	100	100	100
0,05mg Subtilisin KL A228V			
0.2% Ter., 0,2% Lip. and 0,2% Man.	100	100	100
0,05mg Subtilisin KL A230V			
0.2% Ter., 0,2% Lip. and 0,2% Man.	100	100	100
0,05mg Subtilisin KL A228V+A230V			
0.2% Ter., 0,2% Lip. and 0,2% Man.	100	100	100

Table 22 % Residual Lipase activity

Weeks	1	2	3
0,05mg Savinase 16L			
0.2% Ter., 0,2% Lip. and 0,2% Man.	30	5	5
0,05mg Alcalase 2,5L			
0.2% Ter., 0,2% Lip. and 0,2% Man.	10	6	6
0,05mg Subtilisin KL			
0.2% Ter., 0,2% Lip. and 0,2% Man.	59	8	5

Weeks	1	2	3
0,05mg Subtilisin KL S162R			
0.2% Ter., 0.2% Lip. and 0.2% Man.	82	14	6
0,05mg Subtilisin KL S99SE+N76D			
0.2% Ter., 0.2% Lip. and 0.2% Man.	81	15	20
0,05mg Subtilisin KL N76D			
0.2% Ter., 0.2% Lip. and 0.2% Man.	49	49	57
0,05mg Subtilisin KL A228V			52
0.2% Ter., 0.2% Lip. and 0.2% Man.	53		47
0,05mg Subtilisin KL A230V			59
0.2% Ter., 0.2% Lip. and 0.2% Man.	65		52
0,05mg Subtilisin KL A228V+A230V			55
0.2% Ter., 0.2% Lip. and 0.2% Man.	61		48

Table 23 % Residual Mannase activity

Weeks	1	2	3
0,05mg Savinase 16L			
0.2% Ter., 0.2% Lip. and 0.2% Man.	93	44	27
0,05mg Alcalase 2,5L			
0.2% Ter., 0.2% Lip. and 0.2% Man.	81	29	24
0,05mg Subtilisin KL			
0.2% Ter., 0.2% Lip. and 0.2% Man.	98	71	58
0,05mg Subtilisin KL S162R			
0.2% Ter., 0.2% Lip. and 0.2% Man.	105	77	73
0,05mg Subtilisin KL S99SE+N76D			
0.2% Ter., 0.2% Lip. and 0.2% Man.	98	98	100
0,05mg Subtilisin KL N76D			
0.2% Ter., 0.2% Lip. and 0.2% Man.	89	96	90
0,05mg Subtilisin KL A228V			
0.2% Ter., 0.2% Lip. and 0.2% Man.	95	96	92
0,05mg Subtilisin KL A230V			
0.2% Ter., 0.2% Lip. and 0.2% Man.	107	90	89
0,05mg Subtilisin KL A228V+A230V			
0.2% Ter., 0.2% Lip. and 0.2% Man.	97	88	84

REFERENCES CITED IN THE DESCRIPTION

[0078] This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

[0079]

- WO8906279A [0003]

- WO9820115A [0004] [0017]
- GB1296839A [0006]
- US4537706A [0006]
- WO9732961 [0006]
- WO9623873A [0006] [0039]
- US8093562A [0006]
- US2003180933A [0007]
- WO2001066712A [0012]
- WO2006002643A [0012]
- WO2006006060A [0012] [0039]
- WO1995024471A [0013]
- WO9117244A [0013] [0016] [0056]
- WO2002099091A [0013]
- WO2000060063A [0014] [0037]
- WO9964619A [0015] [0042] [0056]
- WO9100345A [0021]
- WO0071691A [0022]
- WO9613580A [0037]
- WO9506720A [0037]
- WO9627002A [0037]
- WO9612012A [0037]
- WO9205249A [0038]
- WO9401541A [0038]
- EP407225A [0038]
- EP260105A [0038]
- WO9535381A [0038]
- WO9600292A [0038]
- WO9530744A [0038]
- WO9425578A [0038]
- WO9514783A [0038]
- WO9522615A [0038]
- WO9704079A [0038]
- WO9707202A [0038]
- WO9402597A [0039]
- WO9418314A [0039]
- WO9743424A [0039]
- US5648263A [0040]
- US5691178A [0040]
- US5776757A [0040]
- WO8909259A [0040]
- EP0531372A [0040]
- WO9611262A [0040]
- WO9629397A [0040]
- WO9808940A [0040]
- WO9407998A [0040]
- EP0531315A [0040]
- US5457046A [0040]
- US5686593A [0040]
- US5763254A [0040]
- WO9524471A [0040]
- WO9812307A [0040]
- DK9800299W [0040]
- WO9324618A [0041]
- WO9510602A [0041]
- WO9815257A [0041]
- WO9535362A [0042]
- EP238216A [0044]
- WO9219709A [0052]

- WO9219708A [0052]
- US5972873A [0052]
- EP0832174A [0052]
- WO0166712A [0056]
- WO0060063A [0056]

Non-patent literature cited in the description

[0080]

- SIEZEN et al. Protein Engng., 1991, vol. 4, 719-737 [0023]
- DYNANTIJAN Nature, 1985, vol. 316, 774-78 [0028]

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- WO8906279A [0003] [0079]
- WO9820115A [0004] [0017] [0079]
- GB1296839A [0006] [0079]
- US4537706A [0005] [0079]
- WO9732961A [0006] [0079]
- WO9623873A [0006] [0039] [0079]
- US6093562A [0005] [0079]
- US2003180933A [0007] [0079]
- WO2001066712A [0012] [0079]
- WO2006002643A [0012] [0079]
- WO2006006060A [0012] [0039] [0079]
- WO1995024471A [0013] [0079]
- WO9117244A [0013] [0016] [0056] [0079]
- WO2002099091A [0013] [0079]
- WO2000060063A [0014] [0037] [0079]
- WO9964619A [0015] [0042] [0056] [0079]
- WO9100345A [0021] [0079]
- WO0071691A [0022] [0079]
- WO9613580A [0037] [0079]
- WO9506720A [0037] [0079]
- WO9627002A [0037] [0079]
- WO9612012A [0037] [0079]
- WO9205249A [0038] [0079]
- WO9401541A [0038] [0079]
- EP407225A [0038] [0079]
- EP260105A [0038] [0079]
- WO9535381A [0038] [0079]
- WO9600292A [0038] [0079]

- [WO9530744A](#) [0038] [0079]
- [WO9425579A](#) [0038] [0079]
- [WO9514783A](#) [0038] [0079]
- [WO9522615A](#) [0038] [0079]
- [WO9704079A](#) [0038] [0079]
- [WO9707202A](#) [0038] [0079]
- [WO9402597A](#) [0039] [0079]
- [WO9418314A](#) [0039] [0079]
- [WO9743424A](#) [0039] [0079]
- [US5648263A](#) [0040] [0079]
- [US5691178A](#) [0040] [0079]
- [US5776757A](#) [0040] [0079]
- [WO8909259A](#) [0040] [0079]
- [EP0531372A](#) [0040] [0079]
- [WO9611262A](#) [0040] [0079]
- [WO9629397A](#) [0040] [0079]
- [WO9808940A](#) [0040] [0079]
- [WO9407998A](#) [0040] [0079]
- [EP0531315A](#) [0040] [0079]
- [US5457046A](#) [0040] [0079]
- [US5686593A](#) [0040] [0079]
- [US5763254A](#) [0040] [0079]
- [WO9624471A](#) [0040] [0079]
- [WO9812307A](#) [0040] [0079]
- [DK9600299W](#) [0040]
- [WO9324618A](#) [0041] [0079]
- [WO9510602A](#) [0041] [0079]
- [WO9815257A](#) [0041] [0079]
- [WO9535362A](#) [0042] [0079]
- [EP238216A](#) [0044] [0079]
- [WO9219709A](#) [0052] [0079]
- [WO9219708A](#) [0052] [0079]
- [US5972873A](#) [0052] [0079]
- [EP0832174A](#) [0052] [0079]
- [WO0166712A](#) [0056] [0079]
- [WO0006063A](#) [0056] [0079]

Non-patent literature cited in the description

- **SIEZEN et al.** Protein Engng., 1991, vol. 4, 719-737 [0023] [0080]
- **DYNANTIJAN** Nature, 1985, vol. 316, 774-78 [0028]
- **DYNANTIJAN** Nature, 1985, vol. 316, 774-78 [0080]

PATENTKRAV

1. Væske- eller gelformig detergentsammensætning, der omfatter subtilisin KL eller varianter deraf kombineret med mindst én lipase; amylase; cellulase; eller mannanase, hvor vægtforholdet mellem indholdet af subtilisin KL eller varianter deraf og indholdet af lipase, amylase, cellulase eller mannanase er fra 0,001 til 100, fortrinsvis fra 0,01 til 10, især fra 0,5 til 5, særligt fra 1 til 3, og
- 5

hvor subtilisin KL varianten er valgt fra gruppen bestående af

*36D

P14T

N18K

N62D

V83L

A133P

E136Q

E136R

E136K

N140R

N140K

S141E

S141N

S141Y

S141R

T143R

T143K

S153R

S156R

A160R

S162R

S162K

I165R

I165K

Y171R

Y171K

A172R

A172K

A174R

N173R

N173K

A174K

N76D

Y176R

Y176K

A187R

A187K

S188P

S190P

Q191R

Y192R

Y192R

Q191P

Y192A

Y192P

D197N

D197R

D197E

D197K

D197G

A228V

A230V

T260R

T260K

G264R

G264K

S265T

S265R

S265K

N218S

M222S

M222A

M222G

M222T

M222V

M222S

N243R

V244R

N248R

K251R

N252R

N261R

S9R+A15T+T22A+N218S+K251R

S9R+A15T+T22A+V84I+N218S

V30I+V139L+N218S

V84I+V139L+N218S

N76D+N218S

N76D+A228V

N76D+A230V

N76D+N218S+A230V

N76D+A228V+A230V

N218S+R247Q

N218S+R247H

N218S+R247E

N218S+R247K

D181N+N218S

N218S+A230V

K251R+S265K

P14T+N18K

T274H+R275H+*275aH+*275bH+*275cH+*275dH=

T274H+R275HHHHH

T274H+R275H+*275aH+*275bH+*275cH=T274H+R275HHHH

S87N+S101G,V104N

*36D+N76D+H120D+G195E+K235L

A133P+M222S

*96aA

*96aA+A98T

*96aA+A133P

*96aA+A98T+A133P

*96aA+A98T+N218S

*97aP+A98T+N218S

*98aT,

*98aT+S99N+N218S

G97D+*98aT+N218S

*99aE=S99SE

*99aD=S99SD

*99aD+M222S=S99SD+M222S

N76D+s99A+*99aE=N76D+S99AE

N76D+*99aD+A230V=N76D+S99SD+A230V

S99A+*99aD=S99AD

S99A+*99aD+M222S=S99AD+M222S

S99A+*99aD+N218S=S99AD+N218S

S99A+*99aE+A230V=S99AE+A230V

A228V+A230V

*130aL+P194A

2. Væske- eller gelformig detergentsammensætning ifølge krav 1, hvor lipasen er udvalgt fra gruppen, der omfatter lipaser fra *Humicola (Thermomyces)*, f.eks. fra *H. lanuginosa (T. lanuginosus)* eller fra *H. insolens*, *Pseudomonas*-lipaser, f.eks. fra *P. alcaligenes* eller *P. pseudoalcaligenes*, *P. cepacia*, *P. stutzeri*, *P. fluorescens*, *Pseudomonas* sp.-stamme SD 705, *P. wisconsinensis*, *Bacillus*-lipaser, f.eks. fra *B. subtilis*, *B. stearothermophilus* eller *B. pumilus* og kemisk eller proteinmodificerede varianter deraf.
- 5
3. Væske- eller gelformig detergentsammensætning ifølge krav 1 eller 2, hvor amylasen er udvalgt fra gruppen, der omfatter amylaser fra *Bacillus*, f.eks. *B. licheniformis*.
- 10
4. Væske- eller gelformig detergentsammensætning ifølge et hvilket som helst af kravene 1 eller 2, hvor cellulase er udvalgt fra slægterne *Bacillus*, *Pseudomonas*, *Myceliophthora*, *Humicola*, *Fusarium*, *Thielavia*, *Acremonium*, f.eks. fra *Humicola insolens*, *Myceliophthora thermophila* og *Fusarium oxysporum*.
- 15
5. Væske- eller gelformig detergentsammensætning ifølge et hvilket som helst af kravene 1 til 4, hvor indholdet af subtilisin KL eller varianter deraf er fra 0,001 til 5 vægt-%, og, hvis de er til stede, indholdet af hver af følgende lipase, amylase, cellulase eller mannanase er fra 0,001 til 5 vægt-%.
- 20
6. Anvendelse af subtilisin KL eller varianter deraf kombineret med mindst én lipase, amylase, cellulase eller mannanase til fremstilling af vandige væske- eller gelformige typer af detergentsammensætninger, der har øget stabilitet over for non-proteaseenzymerne,

hvor subtilisin KL varianten er valgt fra gruppen bestående af

*36D

P14T

N18K

N62D

V83L

A133P
E136Q
E136R
E136K
N140R
N140K
S141E
S141N
S141Y
S141R
T143R
T143K
S153R
S156R
A160R
S162R
S162K
I165R
I165K
Y171R
Y171K
A172R
A172K
A174R
N173R
N173K
A174K
N76D
Y176R
Y176K
A187R
A187K
S188P
S190P
Q191R
Y192R
Y192R

Q191P
Y192A
Y192P
D197N
D197R
D197E
D197K
D197G
A228V
A230V
T260R
T260K
G264R
G264K
S265T
S265R
S265K
N218S
M222S
M222A
M222G
M222T
M222V
M222S
N243R
V244R
N248R
K251R
N252R
N261R
S9R+A15T+T22A+N218S+K251R
S9R+A15T+T22A+V84I+N218S
V30I+V139L+N218S
V84I+V139L+N218S
N76D+N218S
N76D+A228V
N76D+A230V

N76D+N218S+A230V
N76D+A228V+A230V
N218S+R247Q
N218S+R247H
N218S+R247E
N218S+R247K
D181N+N218S
N218S+A230V
K251R+S265K
P14T+N18K
T274H+R275H+*275aH+*275bH+*275cH+*275dH=
T274H+R275HHHHH
T274H+R275H+*275aH+*275bH+*275cH=T274H+R275HHHH
S87N+S101G,V104N
*36D+N76D+H120D+G195E+K235L
A133P+M222S
*96aA
*96aA+A98T
*96aA+A133P
*96aA+A98T+A133P
*96aA+A98T+N218S
*97aP+A98T+N218S
*98aT,
*98aT+S99N+N218S
G97D+*98aT+N218S
*99aE=S99SE
*99aD=S99SD
*99aD+M222S=S99SD+M222S
N76D+s99A+*99aE=N76D+S99AE
N76D+*99aD+A230V=N76D+S99SD+A230V
S99A+*99aD=S99AD
S99A+*99aD+M222S=S99AD+M222S
S99A+*99aD+N218S=S99AD+N218S
S99A+*99aE+A230V=S99AE+A230V
A228V+A230V
*130aL+P194A

7. Fremgangsmåde til forøgelse af stabiliteten af non-proteaseenzymerne i en kombination af et proteaseenzym med andre enzymer i en væske- eller gelformig detergentsammensætning, der omfatter en protease og mindst ét non-proteaseenzym, hvor den væske- eller gelformige detergentsammensætning fremstilles ved brug af subtilisin KL eller en variant deraf som proteaseenzymet, og hvor det mindst ene non-proteaseenzym er udvalgt blandt lipase, amylase, cellulase eller mannanase,
- 5

hvor subtilisin KL varianten er valgt fra gruppen bestående af

*36D

P14T

N18K

N62D

V83L

A133P

E136Q

E136R

E136K

N140R

N140K

S141E

S141N

S141Y

S141R

T143R

T143K

S153R

S156R

A160R

S162R

S162K

I165R

I165K

Y171R

Y171K

A172R

A172K

A174R

N173R

N173K

A174K

N76D

Y176R

Y176K

A187R

A187K

S188P

S190P

Q191R

Y192R

Y192R

Q191P

Y192A

Y192P

D197N

D197R

D197E

D197K

D197G

A228V

A230V

T260R

T260K

G264R

G264K

S265T

S265R

S265K

N218S

M222S

M222A

M222G

M222T

M222V

M222S

N243R

V244R

N248R

K251R

N252R

N261R

S9R+A15T+T22A+N218S+K251R

S9R+A15T+T22A+V84I+N218S

V30I+V139L+N218S

V84I+V139L+N218S

N76D+N218S

N76D+A228V

N76D+A230V

N76D+N218S+A230V

N76D+A228V+A230V

N218S+R247Q

N218S+R247H

N218S+R247E

N218S+R247K

D181N+N218S

N218S+A230V

K251R+S265K

P14T+N18K

T274H+R275H+*275aH+*275bH+*275cH+*275dH=

T274H+R275HHHHH

T274H+R275H+*275aH+*275bH+*275cH=T274H+R275HHHH

S87N+S101G,V104N

*36D+N76D+H120D+G195E+K235L

A133P+M222S

*96aA

*96aA+A98T

*96aA+A133P

*96aA+A98T+A133P

*96aA+A98T+N218S

*97aP+A98T+N218S

*98aT,

*98aT+S99N+N218S

G97D+*98aT+N218S

*99aE=S99SE

*99aD=S99SD

*99aD+M222S=S99SD+M222S

N76D+s99A+*99aE=N76D+S99AE

N76D+*99aD+A230V=N76D+S99SD+A230V

S99A+*99aD=S99AD

S99A+*99aD+M222S=S99AD+M222S

S99A+*99aD+N218S=S99AD+N218S

S99A+*99aE+A230V=S99AE+A230V

A228V+A230V

*130aL+P194A