

(12) PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. AU 199896048 B2
(10) Patent No. 740846

(54) Title
Novel fungicide composition containing a 2-imidazoline-5-one

(51)⁷ International Patent Classification(s)
A01N 043/50

(21) Application No: **199896048**

(22) Application Date: **1997.12.02**

(87) WIPO No: **WO99/27788**

(43) Publication Date : **1999.06.16**

(43) Publication Journal Date : **1999.08.19**

(44) Accepted Journal Date : **2001.11.15**

(71) Applicant(s)
Rhone-Poulenc Agro

(72) Inventor(s)
Maurice Chazalet; Marie-Pascale Latorse; Richard Mercer



(74) Agent/Attorney
DAVIES COLLISON CAVE, 1 Little Collins Street, MELBOURNE VIC 3000



PCT

 ORGANISATION MONDIALE DE LA PROPRIETE INTELLECTUELLE
 Bureau international

DEMANDE INTERNATIONALE PUBLIEE EN VERTU DU TRAITE DE COOPERATION EN MATIERE DE BREVETS (PCT)


(51) Classification internationale des brevets ⁶ : A01N 43/50 // (A01N 43/50, 47:12, 43:54, 43:50, 43:30, 37:40)	A1	(11) Numéro de publication internationale: WO 99/27788 (43) Date de publication internationale: 10 juin 1999 (10.06.99)
<p>(21) Numéro de la demande internationale: PCT/FR97/02170</p> <p>(22) Date de dépôt international: 2 décembre 1997 (02.12.97)</p> <p>(71) Déposant (pour tous les Etats désignés sauf US): RHONE POULENC AGROCHIMIE [FR/FR]; 14-20, rue Pierre Baizet, F-69009 Lyon (FR).</p> <p>(72) Inventeurs; et</p> <p>(75) Inventeurs/Déposants (US seulement): CHAZALET, Maurice [FR/FR]; Le Mont Lucenay, F-69480 Anse (FR). LATORSE, Marie-Pascale [FR/FR]; Lieu-dit "Le Jannt", F-69210 Sourcieux les Mines (FR). MERCER, Richard [FR/FR]; 14, rue du Domaine, F-69130 Ecully (FR).</p> <p>(74) Représentant commun: RHONE POULENC AGROCHIMIE; Boîte postale 9163, F-69263 Lyon Cedex 09 (FR).</p>		<p>(81) Etats désignés: AL, AU, BA, BB, BG, BR, CA, CN, CU, CZ, EE, GE, HU, IL, IS, JP, KP, KR, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, SL, TR, TT, UA, US, UZ, VN, YU, brevet ARIPO (GH, KE, LS, MW, SD, SZ, UG, ZW), brevet européen (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), brevet OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p style="text-align: center;">Publiée <i>Avec rapport de recherche internationale.</i></p> <div style="text-align: center;">  </div>
<p>(54) Title: NOVEL FUNGICIDE COMPOSITION CONTAINING A 2-IMIDAZOLINE-5-ONE</p>		
<p>(54) Titre: NOUVELLE COMPOSITION FONGICIDE COMPRENANT UNE 2-IMIDAZOLINE-5-ONE</p>		
<div style="display: flex; justify-content: space-around; align-items: center;"> <div style="text-align: center;">  </div> <div style="text-align: center;"> <p>(IIB)</p> </div> </div>		
<p>(57) Abstract</p> <p>The invention concerns fungicide compositions comprising a compound (I) which is (4-S)-4-methyl-2-methylthio-4-phenyl-1-phenylamino-2-imidazoline-5-one and a compound (II) selected among the group comprising: (IIA) propamocarb; (IIB) S- methyl ester of 1,2,3-benzothiadiazole-7-carbothioic acid, of formula (IIB); (IIC) cyprodinil; (IID) 2-hydroxy-benzoic acid or salicylic acid, its esters and its salts, in particular of alkaline and alkaline-earth metals; compound (IIF) or 8-t-butyl-2-(N-ethyl-N-n-propylamino)-methyl-1,4-dioxaspiro[4.5]decane, also called spiroxamine; compound (IIG) or isopropyl ester of [2-methyl-1-(1-p-tolyl-ethylcarbamoyl)-propyl] acid, also called fencaramide; compound (IIH) or 4-chloro-2-cyano-1-dimethylsulphamoyl-5-(4-methylphenyl)imidazole; the compound (I)/compound (II) ratio ranging between 0.01 and 50, preferably between 0.1 and 10. The invention also concerns a method, curative or preventive, for fighting against phytopathogenic fungi of crops, characterised in that it consists in applying on the aerial parts of plants an efficient and non-phytotoxic amount of one of said fungicide compositions.</p>		

ABSTRACT

1) Fungicidal compositions comprising a compound (I) which is (4S)-4-methyl-2-methylthio-4-phenyl-1-phenylamino-2-imidazolin-5-one and a compound (II) chosen from the group comprising:

(IIA) propamocarb;

(IIB) the S-methyl ester of 1,2,3-benzothiadiazole-7-carbothioic acid, of formula:



(IIB)

- (IIC) cyprodinil;

- (IID) 2-hydroxybenzoic acid or salicylic acid, its esters and its salts, in particular the alkali-metal salts and the alkaline-earth metal salts;

- the compound (IIF) or 8-t-butyl-2-(N-ethyl-N-n-propylamino)methyl-1,4-dioxaspiro[4.5]decane, also known as spiroxamine;

- the compound (IIG) or isopropyl ester of [2-methyl-1-(1-p-tolyloethylcarbonyl)propyl] acid, also known as fencaramide;



- the compound (IIH) or 4-chloro-2-cyano-1-dimethylsulphamoyl-5-(4-methylphenyl)imidazole;

the compound (I)/compound (II) ratio being between 0.01 and 50, preferably between 0.01 and 10.

2) Process for curatively or preventively combating the phytopathogenic fungi of crops, characterized in that an effective and non-phytotoxic amount of one of these fungicidal compositions is applied to the aerial parts of the vegetation.



Novel fungicidal composition comprising a
2-imidazolin-5-one

The present invention relates to novel
fungicidal compositions comprising a 2-imidazolin-
5 5-one, which are intended in particular for protecting
crops. The invention also relates to a process for
protecting crops against fungal diseases.

Compounds derived from 2-imidazolin-5-ones
with fungicidal action are known, in particular from
10 European patent application EP 551,048, these compounds
making it possible to prevent the growth and
development of phytopathogenic fungi which attack or
are liable to attack crops.

International patent application WO 96/03044
15 also discloses a certain number of fungicidal composi-
tions comprising a 2-imidazolin-5-one in combination
with one or more fungicidal active materials.

However, it is always desirable to improve
the products which can be used by growers in order to
20 combat fungal diseases of crops, and in particular
mildews.

It is also always desirable to reduce the
doses of chemical products spread into the environment
to combat fungal attacks on crops, in particular by
25 reducing the application doses of the products.

Lastly, it is always desirable to increase
the number of antifungal products available to growers
in order that they will find, among these products, the



one which is best suited to the grower's specific use.

One aim of the invention is thus to provide a novel fungicidal composition which is useful for the problems outlined above.


5 Another aim of the invention is to propose a novel fungicidal composition which is useful in the preventive and curative treatment of Solanacea diseases.

10 Another aim of the invention is to propose a novel fungicidal composition which is of improved efficacy against mildew and/or early blight in Solanaceae.

15 Another aim of the invention is to propose a novel fungicidal composition which is of improved efficacy against mildew and/or oidium and/or botrytis in grapevine.

It has now been found that these aims may be achieved, partly or totally, by means of the fungicidal compositions according to the present invention.

20 The subject of the present invention is thus, firstly, fungicidal compositions comprising a compound (I) of formula:



(I)




in which:

- M represents an oxygen or sulphur atom;
- n is an integer equal to 0 or 1;
- Y is a fluorine or chlorine atom or a methyl radical;

and a compound (II) chosen from the group comprising:

- compound (IIA) or propamocarb, also known as propyl 3-(dimethylamino)propyl carbamate;
- a compound (IIB) of formula (IIB)



(IIB)

in which:

- R₁ is a nitrogen atom or the -CH group, and
- R₂ is a thiomethyl SCH₃ group or a diethylamino N(C₂H₅)₂ group,
- the compound (IIC) or cyprodinil, also known as 2-phenylamino-4-cyclopropyl-6-methylpyrimidine;
- the compound (IID), which is 2-hydroxybenzoic acid or salicylic acid, its esters and its salts, in particular the alkali-metal and alkaline-earth metal salts;
- the compound (IIF) or 8-t-butyl-2-(N-ethyl-



N-n-propylamino)methyl-1,4-dioxaspiro[4.5]decane, also known as spiroxamine;

- the compound (IIG) or isopropyl ester of [2-methyl-1-(1-p-tolyloethylcarbonyl)propyl] acid, also
5 known as fencaramide;

- the compound (IIH) or 4-chloro-2-cyano-1-dimethylsulphamoyl-5-(4-methylphenyl)imidazole;

the compound (I)/compound (II) ratio being
between 0.01 and 50, preferably between 0.1 and 10 and
wherein the compound (I)/compound (II) ratio is chosen
so as to produce a synergistic effect.
10

It is clearly understood that the said fungicidal compositions may contain a single compound (II) or more than one such compound, for example 1, 2 or 3 compounds (II) depending on the use for which they
15 are intended.

The composition according to the invention are advantageous for combating, in particular, mildews of the Solanaceae, such as potatoes or tomatoes, as well as for combating mildew and oidium of grapevine.
20

Compound (I) is known, in particular from patent application EP 629,616.

Compound (IIA), or propamocarb, is a fungicide described in the Pesticide Manual 10th Edition, published by the British Crop Protection
25 Council page 843.

Compound (IIB) and its use are known in particular from European patent applications

EP 313,512, EP 420,803 and EP 690,061.

Compound (IIC), or cyprodinil, is described



in European patent application EP 310,550.

Compound (IIF) and its use as a fungicide are described in European patent application EP 0,281,842.

Compound (IIG) and its use as a fungicide are described in at least one of the European patent applications EP 0,398,072 or EP 0,472,996.

Compound (IIH) and its use as a fungicide are described in at least one of the European patent applications EP 0,298,196 or EP 0,705,823.

The compound (I)/compound (II) ratio is defined as being the ratio of the weight of these 2 compounds. This is likewise the case for any ratio of 2 chemical compounds, mentioned below in the present text, insofar as a definition different from this ratio is not expressly indicated.

These compositions generally appreciably improve the respective and isolated action of compound (I) and of compound (II) for a certain number of fungi that are particularly harmful in crops, in particular for Solanaceae, more particularly for mildew of Solanaceae while at the same time retaining an absence of phytotoxicity towards these crops. This therefore results in an improvement in the spectrum of activity and the possibility of decreasing the respective dose of each active material used, the latter quality being particularly advantageous for readily appreciated ecological reasons.

The fungicidal compositions according to the



invention for which the compound (I) is the compound of formula (I) in which M is a sulphur atom and n is equal to 0, also known as (4S)-4-methyl-2-methylthio-4-phenyl-1-phenylamino-2-imidazolin-5-one, are preferred.

5 In the compositions according to the invention, the compound (I)/compound (II) ratio is advantageously chosen so as to produce a synergistic effect. The term synergistic effect is understood to refer in particular to that defined by Colby S.R. in an
10 article entitled "Calculation of the synergistic and antagonistic responses of herbicide combinations" published in the journal Weeds, 1967, 15, p. 20-22. The said article uses the formula:

$$E = X + Y - XY/100$$

15 in which E represents the expected percentage of inhibition of the disease for the combination of the two fungicides at defined doses (for example equal to x and y respectively), X is the percentage of inhibition observed for the disease by the compound (I)
20 at a defined dose (equal to x), Y is the percentage of inhibition observed for the disease by the compound (II) at a defined dose (equal to y). When the percentage of inhibition observed for the combination is greater than E, there is a synergistic effect.

25 The term synergistic effect is also understood to refer to that defined by application of the Tammes method, "Isoboles, a graphic representation of synergism in pesticides" Netherlands Journal of



Plant Pathology, 70 (1964), p. 73-80.

The ranges of the compound (I)/compound (II) ratio indicated above do not in any way limit the scope of the invention, but are rather mentioned as a guide,
5 a person skilled in the art being entirely capable of carrying out complementary tests in order to find other values of the ratio of doses of these 2 compounds for which a synergistic effect is observed.

According to a first preferred variant of the
10 compositions according to the invention, the compound (II) is chosen from the group comprising compounds (IIA), (IIB), (IIC) and (IID). These compositions in particular have advantageous synergistic properties.

According to a more preferred aspect of this
15 first variant of the compositions according to the invention, the compound (II) is compound (IIA) or propamocarb. In this case, particularly improved protection against mildew of Solanaceae is obtained.

In a preferred manner, when the compound (II)
20 is propamocarb, the compound (I)/compound (II) ratio is between 0.01 and 10, preferably between 0.05 and 1 and even more preferably between 0.1 and 1.

According to another more preferred aspect of
this first variant of the compositions according to the
25 invention, the compound (II) is compound (IIB).

In this case, the compound of formula (IIB) in which R_1 represents $-CH$ and R_2 represents $-SCH_3$ is more particularly preferred. The latter compound is the



S-methyl ester of 1,2,3-benzothiadiazole-7-carbothioic acid (also known as CGA 245704).

According to another aspect of the first variant of the compositions according to the invention,
5 the compound (II) is compound (IID).

In a preferred manner, when the compound (II) is compound (IIB) or (IID), the compound (I)/compound (II) ratio is between 0.05 and 50, preferably between 0.1 and 10 and even more preferably between 0.1 and 5.

10 According to another aspect of the first variant of the compositions according to the invention, the compound (II) is compound (IIC).

In a preferred manner, when the compound (II) is compound (IIC), the compound (I)/compound (II) ratio
15 is between 0.05 and 50, preferably between 0.1 and 10 and even more preferably between 0.2 and 1.

According to a third variant of the compositions according to the invention, the compound (II) is compound (IIF).

20 When the compound (II) is compound (IIF), the compound (I)/compound (II) ratio is preferably between 0.05 and 10, preferably between 0.1 and 5.

According to a fourth variant of the compositions according to the invention, the compound
25 (II) is compound (IIG).

When the compound (II) is compound (IIG), the compound (I)/compound (II) ratio is preferably between 0.25 and 5, preferably between 0.5 and 4.



According to an even more preferred embodiment of the compositions according to the invention, they advantageously comprise, in the latter case, besides the compounds (I) and (IIG), a compound (IIG') chosen from:

- the salts of a monoalkyl phosphite and of a mono-, di- or trivalent metal cation, such as fosetyl-Al, or
- phosphorous acid and its alkali-metal or alkaline-earth metal salts.

The (IIG)/(IIG') molar ratio of the compounds (IIG) and (IIG') is generally between 0.037 and 0.37, preferably between 0.018 and 1.8. For the purposes of the present invention, the expression (IIG)/(IIG') molar ratio is understood to refer to the ratio calculated in the following way. The numerator of this ratio is equal to the number of moles of the compound (IIG). The denominator of this ratio is equal to the number of moles of the compound (IIG') multiplied by the number of moles of phosphorous acid resulting from the hydrolysis of one mole of compound (IIG'). Phosphorous acid is the compound of empirical formula H_3PO_3 .

Fosetyl-Al is preferably used as compound (IIG'). The (IIG)/(IIG') weight ratio of the compounds (IIG) and (IIG') is, in this case, between 0.01 and 1, preferably between 0.05 and 5.

According to a fifth variant of the



compositions according to the invention, the compound (II) is compound (IIH). In this case, the compound (I)/compound (II) ratio is between 0.05 and 5, preferably between 0.5 and 2.

5 Besides the compound (I) and the compound (II), the compositions according to the invention comprise an agriculturally suitable inert support and optionally an agriculturally suitable surfactant. In the following text, the term active material denotes
10 the combination of the compound (I) with the compound (II), and the percentages quoted are, except where otherwise mentioned, weight/weight percentages.

 In the present description, the term "support" denotes an organic or inorganic, natural or
15 synthetic material with which the active material is combined in order to facilitate its application onto the plant or onto the soil. This support is thus generally inert and should be agriculturally acceptable, in particular on the crop treated. The
20 support may be solid (in particular: clays, natural or synthetic silicates, silica, resins, waxes, solid fertilizers) or liquid (in particular: water, alcohols, ketones, petroleum fractions, aromatic or paraffinic hydrocarbons, chlorohydrocarbons, liquefied gases).

25 The surfactant may be an emulsifying, dispersing or wetting agent of ionic or nonionic type. Mention may be made, for example, of polyacrylic acid salts, liginosulphonic acid salts, phenolsulphonic or



naphthalenesulphonic acid salts, polycondensates of ethylene oxide with fatty alcohols or with fatty acids or with fatty amines, substituted phenols (in particular alkylphenols or arylphenols), salts of
5 sulphosuccinic acid esters, taurine derivatives (in particular alkyltaurates) and phosphoric esters of polyoxyethylated phenols or alcohols. The presence of at least one surfactant is often required since the active material and/or the inert support are not water-
10 soluble and the vector agent for the application is water.

These compositions may also contain other ingredients of any kind such as, for example, protective colloids, adhesives, thickeners, thixotropic
15 agents, penetration agents, stabilizers, sequestering agents, pigments, dyes or polymers.

More generally, the compositions according to the invention may be combined with any solid or liquid additives corresponding to the usual techniques of
20 agrochemical formulation.

The application techniques are well known to those skilled in the art and they may be used without inconvenience in the context of the present invention. Mention may be made, for example, of spraying.

25 Among the compositions, solid or liquid compositions may be mentioned generally.

As forms of solid compositions, mention may be made of powders for dusting or dispersion (with an



active material content which may be up to 100%) and granules, in particular those obtained by extrusion, by compacting, by impregnation of a granular support or by granulation of a powder (the active material content in these granules being between 1 and 80% for the latter cases).

The compositions may also be used in the form of a powder for dusting; a composition comprising 50 g of active material, 10 g of finely divided silica, 10 g of organic pigment and 970 g of talc may thus be used; these constituents are mixed together and ground and the mixture is applied by dusting.

As liquid composition forms or forms intended to constitute liquid compositions during application, mention may be made of solutions, in particular water-soluble concentrates, emulsifiable concentrates, emulsions, concentrated suspensions, aerosols, wetttable powders (or powder to be sprayed), pastes and dispersible granules.

The emulsifiable or soluble concentrates usually comprise 10 to 80% of active material, the ready-to-apply emulsions or solutions themselves containing 0.01 to 20% of active material.

For example, in addition to the solvent, the emulsifiable concentrates may contain, whenever necessary, 2 to 20% of suitable additives such as the stabilizers, surfactants, penetration agents, corrosion inhibitors, dyes or adhesives mentioned above.



Starting with these concentrates, emulsions of any desired concentration may be obtained by dilution with water.

The concentrated suspensions, which may also
5 be applied by spraying, are prepared so as to obtain a stable fluid product which does not separate on settling and they usually contain from 10 to 75% of active material, from 0.5 to 15% of surfactants, from 0.1 to 10% of thixotropic agents, from 0 to 10% of
10 suitable additives, such as pigments, dyes, antifoaming agents, corrosion inhibitors, stabilizers, penetration agents and adhesives and, as support, water or an organic liquid in which the active material is sparingly soluble or insoluble: certain solid organic
15 materials or inorganic salts may be dissolved in the support in order to help prevent sedimentation, or as antifreezes for the water.

The wettable powders (or powder to be sprayed) are usually prepared such that they contain 20
20 to 95% of active material, and they usually contain, in addition to the solid support, from 0 to 5% of a wetting agent, from 3 to 10% of a dispersing agent and, whenever necessary, from 0 to 10% of one or more stabilizers and/or other additives, such as pigments,
25 dyes, penetration agents, adhesives, anticaking agents, etc.

In order to obtain these powders to be sprayed or wettable powders, the active materials are



mixed intimately in suitable mixers with additional substances and are ground using mills or other suitable grinders. Powders to be sprayed whose wettability and placing in suspension are advantageous are thus
5 obtained; they may be placed in suspension with water at any desired concentration.

In place of wettable powders, pastes may be prepared. The conditions and modes of preparation and of use of these pastes are similar to those for the
10 wettable powders or powders to be sprayed.

The dispersible granules are usually prepared by agglomeration, in suitable granulation systems, of the compositions of wettable powder type.

As has already been stated, the aqueous
15 emulsions and dispersions, for example the compositions obtained by diluting a wettable powder or an emulsifiable concentrate according to the invention with water, are included within the general scope of the present invention. The emulsions may be of the
20 water-in-oil or oil-in-water type and they may have a thick consistency such as that of a "mayonnaise".

The fungicidal compositions according to the invention usually contain from 0.5 to 95% of the combination of compound (I) and compound (II).

25 This may be the concentrated composition, that is to say the commercial product combining compound (I) and compound (II). It may also be the dilute composition ready to be applied to the crops to



be treated. In the latter case, the dilution with water may be carried out either using a commercial concentrated composition containing compound (I) and compound (II) (this mixture is referred to as the "ready-to-use" mixture or "ready mix"), or using a mixture prepared at the time of use (known as the "tank mix") of two commercial concentrated compositions each containing compound (I) and compound (II).


Lastly, the subject of the invention is a process for curatively or preventively combating the phytopathogenic fungi of crops, characterized in that an effective and non-phytotoxic amount of a fungicidal composition according to the invention is applied to the vegetation to be treated.

The phytopathogenic fungi of the crops which may be combated by this process are, in particular, those:

- of the group of oomycetes:

- of the genus *Phytophthora* such as *Phytophthora infestans* (mildew of Solanaceae, in particular late blight of potato or tomato mildew),

- of the family of Peronosporaceae, in particular *Plasmopara viticola* (downy mildew of grapevine), *Plasmopara halstedei* (sunflower mildew), *Pseudoperonospora* sp (in particular cucurbit mildew and downy mildew of hop), *Bremia lactucae* (mildew of lettuce), *Peronospora tabacinae* (downy mildew of tobacco) and *Peronospora parasitica* (downy mildew of



cabbage),

- of the group of adelomycetes:

- of the genus *Alternaria*, for example
Alternaria solani (early blight of Solanaceae and in
5 particular of tomato and potato),


- of the genus *Guignardia*, in particular
Guignardia bidwelli (black rot of grapevine),

- of the genus *Oidium*, for example
powdery mildew of grapevine (*Uncinula necator*); powdery
10 mildew of leguminous crops, for example *Erysiphe*
polygوني (powdery mildew of Cruciferae); *Leveillula*
taurica, *Erysiphe cichoracearum*, *Sphaerotheca fuligena*;
(powdery mildew of cucurbits, of composites and of
tomato); *Erysiphe communis* (powdery mildew of beetroot
15 and cabbage); *Erysiphe pisi* (powdery mildew of pea and
alfalfa); *Erysiphe polyphaga* (powdery mildew of bean
and cucumber mildew); *Erysiphe umbelliferarum* (powdery
mildew of umbellifera, in particular of carrot);
Sphaerotheca humuli (hop mildew).

20 The expression "are applied to the vegetation
to be treated" is understood to mean, for the purposes
of the present text, that the fungicidal compositions
which form the subject of the invention may be applied
by means of various treatment processes such as:

25 - spraying a liquid comprising one of the
said compositions onto the aerial parts of the said
vegetation,

- dusting, incorporation of granules or




powders into the soil, watering around the said vegetation and, in the case of trees, injection or sprinkling.

The spraying of a liquid onto the aerial parts of the crops to be treated is the preferred treatment process.

The expression "effective and non-phytotoxic amount" is understood to refer to an amount of composition according to the invention which is sufficient to allow the control or destruction of the fungi present or liable to appear on the crops, this amount entailing no symptoms of phytotoxicity for the said crops. Such an amount is liable to vary within a wide range depending on the fungus to be combated, the type of crop, the climatic conditions and the nature of the compound (II) included in the fungicidal composition according to the invention. This amount may be determined by systematic field trials, which are within the capabilities of those skilled in the art.

Under the usual conditions of agricultural practice, an amount of fungicidal composition according to the invention corresponding to a dose of compound (I) of between 10 and 500 g/ha, preferably between 20 and 300 g/ha, generally gives good results.

When the compound (II) is compound (IIA), the amount of fungicidal composition according to the invention advantageously corresponds to a dose of compound (II) of between 50 and 2500 g/ha, preferably



between 200 and 1500 g/ha.

When the compound (II) is compound (IIB), the amount of fungicidal composition according to the invention advantageously corresponds to a dose of
5 compound (II) of between 2 and 100 g/ha, preferably between 5 and 50 g/ha.


When the compound (II) is compound (IIC), the amount of fungicidal composition according to the invention advantageously corresponds to a dose of
10 compound (II) of between 20 and 2000 g/ha, preferably between 100 and 550 g/ha.

When the compound (II) is compound (IID), the amount of fungicidal composition according to the invention advantageously corresponds to a dose of
15 compound (II) of between 20 and 5000 g/ha, preferably between 50 and 1000 g/ha.

When the compound (II) is compound (IIF), the amount of fungicidal composition according to the invention advantageously corresponds to a dose of
20 compound (II) of between 100 and 1000 g/ha, preferably between 300 and 800 g/ha.

When the compound (II) is compound (IIG), the amount of fungicidal composition according to the invention advantageously corresponds to a dose of
25 compound (II) of between 50 and 250 g/ha, preferably between 100 and 200 g/ha.

When the compound (II) is compound (IIH), the amount of fungicidal composition according to the



invention advantageously corresponds to a dose of compound (II) of between 100 and 400 g/ha, preferably between 150 and 350 g/ha.

The examples which follow are given purely by way of illustration of the invention, which they do not limit in any way.


In these examples, as well as in the figures to which reference is made, compound (I) is understood to denote (4S)-4-methyl-2-methylthio-4-phenyl-1-phenyl-10 amino-2-imidazolin-5-one and compound (IIB) is understood to denote the S-methyl ester of 1,2,3-benzothiadiazole-7-carbothioic acid (or CGA 245704). Similarly, compound (IIG') is understood to refer to Fosetyl-Al.

15 Example 1: Field trial of a composition comprising compounds (I) and (IIA) against late blight of potato (*Phytophthora infestans*):

A composition is used comprising compound (I) in the form of a concentrated suspension at 500 g/l and 20 a composition comprising compound (IIA) or propamocarb in the form of a soluble concentrate at 724 g/l.

These 2 compositions are mixed so as to obtain a compound (I)/compound (IIA) ratio equal to 0.15 to 0.10.

25 The mixture is applied, after dilution with water, at a rate of 500 l/ha to potatoes 2 months after planting the tubers. The doses applied are indicated in



the table below. This application is repeated 5 times every 6 days.

After the 2nd application, contamination is carried out by spraying the *Phytophthora infestans* 5 spores.

The results are observed 3 days after the 5th application. For this, the contamination C (or degree of attack) expressed by the number of blackish marks (caused by the disease) per plot is evaluated visually 10 (relative to an untreated plot which is also contaminated). The efficacy E is calculated according to the so-called Abbott formula:

$$E \text{ (in \%)} = [(C_{\text{untreated control}} - C_{\text{treated plot}}) / C_{\text{untreated control}}] \times 100$$

The results are collated in the following 15 table:

Composition tested	Doses (in g/ha)	(I) / (IIA) ratio	Efficacy (in %)
(I) + (IIA)	100 + 1000	0.1	96
(I) + (IIA)	100 + 666	0.15	95

20 A neighbouring plot treated with mancozeb at a rate of 1600 g/ha gave rise to an efficacy of 90%.



Example 2:

Example 1 is repeated on a potato field located in another region. The results are observed 12 days after the 5th treatment. For this, and on account of the intensity of the attack of the disease, the percentage of foliar surface destroyed is evaluated.

The efficacy is calculated according to the same formula as above.

For the 0.10 ratio, an efficacy of 79% is obtained and for the 0.15 ratio an efficacy of 77% is obtained.

A neighbouring plot treated with mancozeb at a rate of 1600 g/ha gave rise to an efficacy of 59%.

Example 3: Greenhouse test of a composition comprising compounds (I) and (IIB) against downy mildew of grapevine (*Plasmopara viticola*):

A suspension of 60 mg comprising compounds (I) and (IIB) in a liquid mixture consisting of 0.3 ml of a surfactant (oleate of polyoxyethylenated derivative of sorbitan) diluted to 10% in water and 60 ml of water is prepared.

The compound (I)/compound (IIB) ratio in the suspensions prepared is equal to:

0.125; 0.25; 0.5; 2.

Cuttings of grapevine (*Vitis vinifera*), Chardonnay variety, are cultivated in pots. When these



plants are 2 months old (8- to 10-leaf stage, 10 to 15 cm in height), they are treated by spraying with the above suspensions.

Plants used as controls are treated with a similar suspension but which contains no active material ("formulation blank").


After drying for 4 days, each plant is contaminated by spraying with an aqueous suspension of *Plasmopara viticola* spores obtained from sporulated leaves contaminated 7 days beforehand. These spores are placed in suspension in a proportion of 100,000 units per cm³.

The contaminated plants are then incubated for two days at about 18°C, in an atmosphere of saturated humidity and then for 5 days at 20-22°C under 90-100% relative humidity.

The results are observed 7 days after contamination, by comparison with the control plants. The contamination C is measured by visual evaluation of the percentage of contaminated foliar surface (whitish appearance). Using C, the efficacy is calculated with the same formula as in Example 1.

The IC90 is calculated from the efficacy.

The IC90 is defined as being the weight of mixture (defined for a fixed ratio of the compounds (I) and (IIB)) which it is necessary to apply in order to obtain an efficacy of 90%. The IC90 is expressed in the form of the corresponding weight of the compound (I) in



the mixture per ml of liquid sprayed onto the plants.

The results are collated in the table below:


Ratio : compound (I)/compound (IIB)	IC90 (mg/l)
0.125	17
0.25	17
0.5	20
1	35
2	18

Example 4: Greenhouse test of a composition comprising compound (I) and the sodium salt of salicylic acid (compound IID) against late blight of potato (*Phytophthora infestans*):

A concentrated suspension of compound (I) and an aqueous solution of the sodium salt of salicylic acid are used.

10 These 2 liquids are mixed together so as to obtain a compound (I)/compound (IID) ratio equal to: 0.25 - 0.5 - 1 and 2.

15 Potato plants (Bintje variety) are cultivated in pots. When these plants are one month old (5- to 6-leaf stage, 12 to 15 cm in height), they are treated by spraying with a liquid comprising one of the



compounds (I) and (IID) either alone or as a mixture in the ratio indicated above.

After 4 days, each plant is contaminated by spraying with an aqueous suspension of spores
5 (30,000 sp/cm³) of *Phytophthora infestans*.


After this contamination, the potato plants are incubated for 5 days at about 18°C in an atmosphere of saturated humidity.

The results are obtained 5 days after placing
10 in incubation, by comparison with untreated control plants which are also contaminated.

The efficacy results obtained are reported in the form of points, corresponding to 70% destruction of the parasite, and placed in a Tammes diagram which
15 includes the doses of compound (I) expressed in mg/l on the x-axis and the doses of compound (IID), also in mg/l, on the y-axis.

The diagram of Figure 1 is obtained, in which it appears that, when it is applied alone, the sodium
20 salt of salicylic acid has no efficacy under the test conditions. It appears, however, that the addition of this salt makes it possible, entirely unexpectedly, to lower the dose of compound (I) required to destroy 70% of the parasite to below 36 mg/l, which corresponds to
25 the dose of compound (I) alone which needs to be applied in order to obtain this same percentage of destruction.

The arrangement of the points obtained thus



indicates a unilateral effect, which is termed according to the Tammes method mentioned above as a "one-sided effect". This arrangement corresponds to a type-II isobole according to the said method (page 74 of the corresponding bibliographic reference already mentioned) and is characteristic of synergism.

Example 5: in vivo greenhouse test of the combination of compound (I) with compound (IIA), propamocarb, on *Plasmopara viticola* (downy mildew of grapevine) by preventive treatment at 4 days:

A soluble concentrate containing 722 g/l of propamocarb and a 70% dispersible granule of compound (I) are used.

Dilute suspensions corresponding to a spraying volume of 500 l of spraying liquid per ha are prepared from these compositions by dilution with water.

These 2 liquids are mixed together so as to obtain a ratio: compound (I)/propamocarb, equal to: 0.015 - 0.125 - 0.5.

Cuttings of grapevine (*Vitis vinifera*), Chardonnay variety, are cultivated in pots. When these plants are 2 months old, they are treated by spraying with the above liquids, taken individually or as a mixture.

After 4 days, each plant is contaminated by spraying with an aqueous suspension of spores of



Plasmopara viticola obtained from sporulated leaves contaminated 7 days beforehand. These spores are placed in suspension in a proportion of 100,000 units per cm³.

The contaminated plants are then incubated
5 for two days at about 18°C in an atmosphere of saturated humidity and then for 5 days at 20-22°C under 90-100% relative humidity.

The results are read 7 days after the contamination, by comparison with the control plants,
10 that is to say untreated but contaminated plants.

The surface area of leaves showing a whitish appearance on their underside, corresponding to attack by the fungus, is assessed visually and the degree of attack is obtained by dividing by the total foliar
15 surface.

The efficacy is then calculated using the Abbott formula and using the result on the control plant.

The efficacy results are reported in the form
20 of points, corresponding to an efficacy of 90%, and placed in a Tammes isobole diagram which includes the doses of the compound (I) expressed in mg/l on the x-axis and the doses of propamocarb, also in mg/l, on the y-axis.

25 The diagram of Figure 2 is obtained, in which it appears that, when it is applied alone, the propamocarb has no efficacy under the test conditions. It appears, however, that the addition of propamocarb



makes it possible, entirely unexpectedly, to lower the dose of compound (I) needed to destroy 90% of the parasite to below 8.2 mg/l, which corresponds to the dose of compound (I) alone which needs to be applied to
5 obtain the same efficacy.

The arrangement of the points obtained thus indicates a unilateral effect, termed according to the Tammes method mentioned above as a "one-sided effect". This arrangement corresponds to a type-II isobole
10 according to the said method (page 74 of the corresponding bibliographic reference already mentioned) and is characteristic of synergism.


Example 6: in vivo greenhouse test of a composition comprising compounds (I) and (IIA)
15 (propamocarb) on *Phytophthora infestans* (late blight of potato) by preventive treatment at 4 days:

A soluble concentrate containing 722 g/l of propamocarb and a 70% dispersible granule of compound (I) are used.

20 Dilute suspensions corresponding to a spraying volume of 1000 l of spraying liquid per ha are prepared from these compositions by dilution with water.

25 These 2 liquids are mixed together so as to obtain a compound (I)/propamocarb ratio equal to:
0.125 - 0.25 - 1.

Potato plants (Bintje variety) are cultivated



in pots. When these plants are one and a half months old, they are treated with the above liquids, taken individually or as a mixture.

4 days after the treatment, lamellae are cut
5 off and placed on moistened filter paper in a Petri dish, and these lamellae are contaminated by applying 10 drops of an aqueous suspension of spores (containing 30,000 sp/cm³) of *Phytophthora infestans*.

Next, the potato lamellae are incubated for
10 3 days at about 18°C in an atmosphere of saturated humidity.

The results are observed 4 days after the contamination, by comparison with the control plants.

The surface area of the lamellae having a
15 greyish appearance, corresponding to attack by the fungus, is assessed visually and the degree of attack is obtained by dividing this surface area by the total surface of the lamellae.

The efficacy is then calculated using the
20 Abbott formula and using the result on the lamellae used as control.

The results obtained are reported in the form of points, corresponding to 90% destruction of the parasite, and are placed in a Tammes diagram which
25 includes the doses of compound (I) expressed in mg/l on the x-axis and the doses of propamocarb, also in mg/l, on the y-axis.

The diagram of Figure 3 is obtained, in which



it appears that the addition of a dose of compound (I) of less than 8.7 mg/l (which corresponds to the dose of compound (I) alone which needs to be applied in order to obtain 90% destruction of the parasite) makes it possible, entirely unexpectedly, to lower the dose of propamocarb needed to destroy 90% of the parasite to below 553 mg/l (this value corresponding to the dose of propamocarb alone which needs to be applied in order to obtain this same percentage of destruction).

10 The arrangement of the points obtained thus indicates a bilateral effect, termed according to the Tammes method mentioned above as a "two-sided effect". This arrangement corresponds to a type-III isobole according to the said method (page 75 of the corresponding bibliographic reference already mentioned) and is characteristic of synergism.

Example 7: in vivo greenhouse test of a composition comprising compounds (I) and (IIB) (CGA 245704) against late blight of potato (*Phytophthora infestans*):

20 A suspension of 60 mg comprising compounds (I) and (IIB) in a liquid mixture consisting of 0.3 ml of a surfactant (oleate of polyoxyethylenated derivative of sorbitan) diluted to 10% in water and 25 60 ml of water is prepared.

 The compound (I)/compound (IIB) ratio in the suspensions prepared is equal to 0.125; 0.25; 0.5; 1.



Potato plants (Bintje variety) are cultivated in pots. When these plants are one and a half months old, they are treated with the above liquids, taken individually or as a mixture.

5 4 days after the treatment, the plants are contaminated by spraying with an aqueous suspension of spores (containing 30,000 sp/cm³) of *Phytophthora infestans*.

Next, the potato plants are incubated for
10 5 days at about 18°C in an atmosphere of saturated humidity.


The results are observed 5 days after the contamination, in comparison with the control plants.

The foliar surface having a greyish
15 appearance, corresponding to attack by the fungus, is assessed visually and the degree of attack is obtained by dividing this surface area by the total surface of the leaves.

The efficacy is then calculated using the
20 Abbott formula and using the result on the control plant.

The results obtained are reported in the form of points, corresponding to 70% efficacy against the parasite, and are placed in a Tammes diagram which
25 includes the doses of compound (I) expressed in mg/l on the x-axis and the doses of CGA 245704, also in mg/l, on the y-axis.

The diagram of Figure 4 is obtained, in which



it appears that, when it is applied alone, CGA 245704 has no efficacy under the test conditions. It appears, however, that the addition of CGA 245704 makes it possible, entirely unexpectedly, to lower the dose of compound (I) needed to destroy 70% of the parasite to below 177 mg/l, which corresponds to the dose of compound (I) alone which needs to be applied in order to obtain the same efficacy.

The arrangement of the points obtained thus indicates a unilateral effect, termed according to the Tammes method mentioned above as a "one-sided effect". This arrangement corresponds to a type-II isobole according to the said method (page 74 of the corresponding bibliographic reference already mentioned) and is characteristic of synergism.

Example 8: in vivo greenhouse test of a composition comprising compound (I) and compound (IIC) (cyprodinil) against grey rot of grapevine (*Botrytis cinerea*):

A 70% dispersible granule of compound (I) and a 75% dispersible granule of cyprodinil are used.

Dilute suspensions corresponding to a spraying volume of 1000 l of spraying liquid per ha are prepared from these compositions by dilution with water.

These 2 liquids are mixed together so as to obtain a compound (I)/cyprodinil ratio equal to 0.2.



Grapevine plants (Chardonnay variety) are cultivated in pots. When these plants are 2 months old, they are treated by spraying with a liquid comprising one of the compounds (I) and (IIC), either individually
5 or as a mixture in the ratio indicated above.

After 24 hours, leaves are cut off the treated plants, placed on moistened filter paper in a Petri dish and are contaminated by applying 10 drops of an aqueous suspension of spores (containing
10 150,000 sp/cm³) of *Botrytis cinerea*.

After this contamination, the treated and contaminated grapevine leaves are placed for 6 days at about 20°C in an atmosphere of saturated humidity.

The results are then observed by comparison
15 with the untreated control plants which are also contaminated.

For this, the number of drops of *Botrytis cinerea* suspension from which a grey mark, corresponding to attack by the phytopathogenic fungus,
20 developed is counted for a grapevine leaf. This number is divided by the number of drops applied (equal to 10) and the degree of attack (expressed in %) is thus obtained.

The efficacy of the treatment (expressed in
25 %) is calculated according to the so-called Abbott formula.

The efficacy results obtained are reported in the form of points, corresponding to 90% destruction of



the parasite, and are placed in a Tammes diagram which includes the doses of cyprodinil expressed in mg/l on the x-axis and the doses of compound (I), also in mg/l, on the y-axis.

5 The diagram of Figure 5 is obtained, in which it appears that, when it is applied alone, compound (I) has no efficacy under the test conditions. It appears, however, that the addition of cyprodinil makes it possible, entirely unexpectedly, to lower the dose of
10 compound (I) needed to destroy 90% of the parasite to below 73 mg/l, which corresponds to the dose of cyprodinil alone which needs to be applied in order to obtain this same percentage of destruction.

 The arrangement of the points obtained thus
15 indicates a unilateral effect, termed according to the Tammes method mentioned above as a "one-sided effect". This arrangement corresponds to a type-II isobole according to the said method (page 74 of the corresponding bibliographic reference already
20 mentioned) and is characteristic of synergism.

Example 9: in vivo greenhouse test of a composition comprising compound (I) and compound (IIC) (cyprodinil) against alternaria rot in radish (*Alternaria brassicae*):

25 A 70% dispersible granule of compound (I) and a 75% dispersible granule of cyprodinil are used.

Dilute suspensions corresponding to a



spraying volume of 1000 l of spraying liquid per ha are prepared from these compositions by dilution with water.

5 These 2 liquids are mixed together so as to obtain a compound (I)/cyprodinil ratio equal to 0.2 - 0.5 - 1.

10 Radish plantules (Pernot variety) are cultivated in pots. When these plantules have reached the 2-leaf cotyledon stage, they are treated by spraying with a liquid comprising compounds (I) and (IIC), either individually or as a mixture in the ratios indicated above.

15 24 hours after treatment, contamination is carried out by spraying with an aqueous suspension of spores (containing 40,000 sp/cm³) of *Alternaria brassicae*.

After this contamination, the treated and contaminated plantules are placed for 10 days at about 20°C in an atmosphere of saturated humidity.

20 The results are then observed, by comparison with untreated control plantules which are also contaminated.

25 For this, the percentage of attack of the cotyledon leaves by the fungus is assessed visually, the said attack being recognizable by brown necrosis marks. From the degree of attack thus obtained (expressed in %), the efficacy is calculated according to the Abbott formula with reference to the degree of



attack of the control.

The efficacy results obtained are reported in the form of points, corresponding to 70% destruction of the parasite, and are placed in a Tammes diagram which includes the doses of compound (I) expressed in mg/l on the x-axis and the doses of cyprodinil, also expressed in mg/l, on the y-axis.

The diagram of Figure 6 is obtained, in which it appears that the addition of a dose of compound (I) of less than 167 mg/l (which corresponds to the dose of compound (I) alone which needs to be applied in order to obtain 70% destruction of the parasite) makes it possible, entirely unexpectedly, to lower the dose of cyprodinil needed to destroy 70% of the parasite to below 178 mg/l (this value corresponding to the dose of cyprodinil alone which needs to be applied in order to obtain this same percentage of destruction).

The arrangement of the points obtained thus indicates a bilateral effect, termed according to the Tammes method mentioned above as a "two-sided effect". This arrangement corresponds to a type-III isobole according to the said method (page 75 of the corresponding bibliographic reference already mentioned) and is characteristic of synergism.

Example 10: *in vivo* greenhouse test of a composition comprising compound (I) and compound (IIF) (spiroxamine) against powdery mildew of grapevine



(*Uncinula necator*):

A suspension of 60 mg comprising compound (I) in a liquid mixture consisting of 0.3 ml of a surfactant (oleate of polyoxyethylenated derivative of sorbitan) diluted to 10% in water and 60 ml of water is prepared.

An emulsifiable concentrate containing 500 g/l of spiroxamine is used.

Dilute suspensions corresponding to a spraying volume of 250 l of spraying liquid per ha are prepared from these compositions by dilution with water.

These 2 liquids are mixed together so as to obtain a compound (I)/spiroxamine ratio equal to 0.33:1:3.

Grapevine plants (Chardonnay variety) are cultivated in pots. When these plants are 2 months old, they are treated by spraying with a liquid comprising one of the compounds (I) and (IIF), either individually or as a mixture in one of the ratios indicated above.

After 24 hours, the grapevine leaves are contaminated by dusting them with powdery mildew spores (*Uncinula necator*) obtained from naturally contaminated leaves.

After this contamination, the treated and contaminated grapevine plants are placed for 15 days at 20°C at a relative humidity of about 70%.

The results are then observed by comparison



with the untreated control plants which are also contaminated.

For this, the area of white marks corresponding to growth of the fungus is estimated for all of the leaves of a plant, relative to an untreated
5 contaminated plant. A level of contamination is thus obtained.

The efficacy of the treatment (expressed in %) is calculated according to the so-called Abbott
10 formula.

The efficacy results obtained are reported in the form of points, corresponding to 70% destruction of the parasite, and are placed in a Tammes diagram which includes the doses of spiroxamine expressed in g/ha on
15 the x-axis and the doses of compound (I), also in g/ha, on the y-axis.

The diagram of Figure 7 is obtained, in which it appears that, when it is applied alone, compound (I) has no efficacy under the test conditions. It appears,
20 however, that the addition of compound (I) makes it possible, entirely unexpectedly, to lower the dose of spiroxamine needed to destroy 70% of the parasite to below 6.6 g/ha, which corresponds to the dose of spiroxamine alone which needs to be applied in order to
25 obtain this same percentage of destruction.

The arrangement of the points obtained thus indicates a unilateral effect, termed according to the Tammes method mentioned above as a "one-sided effect".



This arrangement corresponds to a type-II isobole according to the said method (page 74 of the corresponding bibliographic reference already mentioned) and is characteristic of synergism.

5 Example 11: in vivo greenhouse test of a composition comprising compound (I) and compound (IIF) (spiroxamine) against powdery mildew of grapevine (*Uncinula necator*).

 Example 10 is repeated, setting the compound
10 (I)/spiroxamine ratio to 0.11:0.33 and 1 and observing the results 22 days after contamination.

 The diagram of Figure 8 is obtained, in which it appears that, when it is applied alone, compound (I) has no efficacy under the test conditions. It appears,
15 however, that the addition of compound (I) makes it possible, entirely unexpectedly, to lower the dose of spiroxamine needed to destroy 70% of the parasite to below 61 g/ha, which corresponds to the dose of
20 spiroxamine alone which needs to be applied in order to obtain this same percentage of destruction.

 The arrangement of the points obtained thus indicates a unilateral effect, termed according to the Tammes method mentioned above as a "one-sided effect". This arrangement corresponds to a type-II isobole
25 according to the said method (page 74 of the corresponding bibliographic reference already mentioned) and is characteristic of synergism.



Example 12: in vivo greenhouse test of a ternary composition comprising compounds (I), IIG (fencaramide) and IIG' (Fosetyl-Al) against downy mildew of grapevine (*Plasmopara viticola*):

5 A 70% dispersible granule of compound I, a 50% dispersible granule of fencaramide and an 80% dispersible granule of Fosetyl-Al are used.

 Dilute suspensions corresponding to a spraying volume of 1000 l/ha are prepared from these
10 compositions by dilution with water. The dilute suspensions of fencaramide and of Fosetyl-Al are mixed together so as to obtain a stock suspension comprising these 2 active materials in a fixed Fencaramide/
Fosetyl-Al ratio equal to 0.1.

15 The dilute suspension of compound (I) and this stock suspension are mixed together so as to obtain a mixture of the 3 active materials in the compound (I)/Fencaramide + Fosetyl-Al ratio equal to $1/(1+10)$ (i.e. 0.09) and $2/(1+10)$ (i.e. 0.18).

20 Grapevine (Chardonnay variety) cuttings are cultivated in pots. When these plants are 2 months old (8- to 10-leaf growth stage), they are treated by spraying with a liquid comprising either compound (I) alone or the mixture Fencaramide + Fosetyl-Al, or the
25 mixture of the 3 active materials. The spraying volume of 1000 l/ha allows coverage of the lower and upper faces of the leaves.

The treated plants were contaminated



beforehand (24 hours before the fungicidal treatment) by spraying with an aqueous suspension of *Plasmopara viticola* spores (100,000 sp/cm³). After contamination, the plants are left at room temperature for 1 hour at
5 70% relative humidity, then for 24 hours at 90 to 100% relative humidity, before receiving the fungicidal treatment.

After the fungicidal treatment, the plants are maintained for 1 hour in an atmosphere at 70%
10 relative humidity and are then left for 7 days at 90 to 100% relative humidity.

The results are then observed by comparison with the untreated control plants which are also contaminated. For this, the contaminated fraction
15 (estimated in %) of the lower surface of the leaves is determined by visual evaluation.

The efficacy results obtained are reported in the form of points, corresponding to 70% destruction of the parasite, and are placed in a Tammes diagram which
20 includes the doses of compound (I) expressed in mg/l on the x-axis and the doses of the fencaramide + fosetyl-Al mixture (in a 1/10 ratio), expressed in the form of a single dose of fencaramide in the mixture, in mg/l, on the y-axis.

25 The diagram of Figure 9 is obtained, in which it appears that the addition of a dose of compound (I) of less than 36 mg/l (which corresponds to the dose of compound (I) alone which needs to be applied in order



to obtain 70% destruction of the parasite) makes it possible, entirely unexpectedly, to lower the dose of the fencaramide/fosetyl-Al mixture (in the ratio 1/10) to well below 34 mg/l (this value corresponding to the equivalent dose of fencaramide in the said mixture which needs to be applied in order to obtain this same percentage of destruction).

The arrangement of the points obtained thus indicates a bilateral effect, termed according to the Tammes method mentioned above as a "two-sided effect". This arrangement corresponds to a type-III isobole according to the said method (page 75 of the corresponding bibliographic reference already mentioned) and is characteristic of synergism.


Throughout this specification and the claims which follow, unless the context requires otherwise, the word "comprise", and variations such as "comprises" and "comprising", will be understood to imply the inclusion of a stated integer or step or group of integers or steps but not the exclusion of any other integer or step or group of integers or steps.

The reference to any prior art in this specification is not, and should not be taken as, an acknowledgment or any form of suggestion that the prior art forms part of the common general knowledge in Australia.



THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. Fungicidal compositions comprising a compound (I) of formula:



(I)

in which:

5

- M represents an oxygen or sulphur atom;

- n is an integer equal to 0 or 1;


- Y is a fluorine or chlorine atom or a methyl radical;

10

and a compound (II) chosen from the group comprising:

- compound (IIA) or propamocarb, also known as propyl 3-(dimethylamino)propyl carbamate;

- a compound (IIB) of formula (IIB)



(IIB)

15

in which:

- R₁ is a nitrogen atom or the -CH group, and



- R₂ is a thiomethyl SCH₃ group or a diethylamino N(C₂H₅)₂ group,

- the compound (IIC) or cyprodinil, also known as 2-phenylamino-4-cyclopropyl-6-methyl-
5 pyrimidine;

- the compound (IID), which is 2-hydroxybenzoic acid or salicylic acid, its esters and its salts, in particular the alkali-metal and alkaline-earth metal salts;

10 - the compound (IIF) or 8-t-butyl-2-(N-ethyl-N-n-propylamino)methyl-1,4-dioxaspiro[4.5]decane, also known as spiroxamine;

- the compound (IIG) or isopropyl ester of [2-methyl-1-(1-p-tolyloethylcarbamoyl)propyl] acid,
15 also known as fencaramide;

- the compound (IIH) or 4-chloro-2-cyano-1-dimethylsulphamoyl-5-(4-methylphenyl)imidazole;

the compound (I)/compound (II) ratio being between 0.01 and 50, wherein the compound
20 (I)/compound (II) ratio is chosen so as to produce a synergistic effect.

2. Fungicidal compositions according to claim 1, wherein the compound (I)/compound (II) ratio is between 0.1 and 10.

3. Fungicidal compositions according to claim 1, or claim 2, wherein the compound (I) is (4S)-4-methyl-2-methylthio-4-phenyl-1-phenylamino-2-
25 imidazolin-5-one.

4. Fungicidal compositions according to any one



- 44 -

of claims 1 to 3, wherein the compound (II) is chosen from the group comprising compounds (IIA), (IIB), (IIC) and (IID).

5 5. Fungicidal compositions according to any one of claims 1 to 4, wherein the compound (II) is compound (IIA) (or propamocarb) and the compound (I)/compound (II) ratio is between 0.01 and 10.

10 6. Fungicidal compositions according to claim 5 wherein the compound (I)/compound (II) ratio is between 0.05 and 1.

7. Fungicidal compositions according to claim 5 or claim 6, wherein the compound (I)/compound (II) ratio is between 0.1 and 1.

15 8. Fungicidal compositions according to any one of claims 1 to 4, wherein the compound (II) is compound (IIB).

9. Fungicidal compositions according to claim 8, wherein the compound (IIB) is the S-methyl ester of 1,2,3-benzothiadiazole-7-carbothioic acid.

20 10. Fungicidal compositions according to any one of claims 1 to 4, wherein the compound (II) is compound (IID).

25 11. Fungicidal compositions according to any one of claims 8 to 10, wherein the compound (I)/compound (II) ratio is between 0.05 and 50.

12. Fungicidal compositions according to claim 11 wherein the compound (I)/compound (II) ratio is between 0.1 and 10.

30 13. Fungicidal compositions according to claim 11 or 12 wherein the compound (I)/compound (II) ratio is between 0.1 and 5.

35 14. Fungicidal compositions according to any one of claims 1 to 4, wherein the compound (II) is compound (IIC) and the compound (I)/compound (II) ratio is between 0.05 and 50.



- 45 -

15. Fungicidal compositions according to claim 14, wherein the compound (I)/compound (II) ratio is between 0.1 and 10.

5 16. Fungicidal compositions according to claim 14 or claim 15 wherein the compound (I)/compound (II) ratio is between 0.2 and 1.

17. Fungicidal compositions according to any one of claims 1 to 3, wherein the compound (II) is compound (IIF).

10 18. Fungicidal compositions according to claim 17, wherein the compound (I)/compound (IIF) ratio is between 0.05 and 10.

15 19. Fungicidal compositions according to claim 18 wherein the compound (I)/compound (IIF) ratio is between 0.1 and 5.

20. Fungicidal compositions according to any one of claims 1 to 3, wherein the compound (II) is compound (IIG).

20 21. Fungicidal compositions according to claim 20, wherein the compound (I)/compound (IIG) ratio is between 0.25 and 5.

22. Fungicidal compositions according to claim 21, wherein the compound (I)/compound (IIG) ratio is between 0.5 and 4.

25 23. Fungicidal compositions according to any one of claims 20 to 22, further comprising a compound (IIG') chosen from:

30 - the salts of a monoalkyl phosphite and of a mono-, di- or trivalent metal cation, such as fosetyl-Al, or

- phosphorous acid and its alkali-metal or alkaline-earth metal salts;

the (IIG)/(IIG') molar ratio being between 0.037 and 0.37.



24. Fungicidal compositions according to any one of claims 20 to 22, further comprising a compound (IIG') chosen from:

5 - the salts of a monoalkyl phosphite and of a mono-, di- or trivalent metal cation, such as fosetyl-Al, or

- phosphorous acid and its alkali-metal or alkaline-earth metal salts;

10 the (IIG)/(IIG') molar ratio being between 0.018 to 1.8.

25. Fungicidal compositions according to any one of claims 20 to 22, further comprising a compound (IIG'), wherein the compound (IIG') is fosetyl-Al and the (IIG)/(IIG') weight ratio is between 0.01 and 1.

15 26. Fungicidal compositions according to any one of claims 20 to 22, further comprising a compound (IIG'), wherein the compound (IIG') is fosetyl-Al and the (IIG)/(IIG') weight ratio is between 0.05 and 5.

20 27. Fungicidal compositions according to any one of claims 1 to 3, wherein the compound (II) is compound (IIH).

28. Fungicidal compositions according to claim 27, wherein the compound (I)/compound (II) ratio is between 0.05 and 5.

25 29. Fungicidal compositions according to claim 28 wherein the compound (I)/compound (II) ratio is between 0.5 and 2.

30 30. Fungicidal compositions according to any one of claims 1 to 29, further comprising besides the compounds (I) and (II), an agriculturally suitable inert support and optionally an agriculturally suitable surfactant.

31. Fungicidal compositions according to any one of claims 1 to 30, wherein the compositions comprise from 0.5 to 95% of the combination of compound (I) and compound (II).



32. Process for curatively or preventively combating the phytopathogenic fungi of crops, wherein an effective and non-phytotoxic amount of a fungicidal composition according to one of claims 1 to 31 is applied to the vegetation to be treated.

33. Process according to claim 32, wherein the fungicidal composition is applied by spraying a liquid onto the aerial parts of the crops to be treated.

34. Process according to claim 32 or claim 33 wherein the amount of fungicidal composition corresponds to a dose of compound (I) of between 10 and 500 g/ha.

35. Process according to claim 34 where the amount of fungicidal composition corresponds to a dose of compound (I) of between 20 and 300 g/ha.

36. Fungicidal compositions according to claim 1, substantially as hereinbefore described with reference to the Examples.

37. Process for curatively or preventively combating the phytopathogenic fungi of crops, substantially as hereinbefore described with reference to the Examples.

25

DATED this 18th day of September, 2001


Rhone Poulenc Agro

30 By DAVIES COLLISON CAVE

Patent Attorneys for the Applicants



Figure 1



Dose in mg/l
Propamocarb (IC90 >2000)





Figure 2

II/IX

Figure 3






Figure 4

IV/IX

V/IX

Figure 5




Figure 6




Figure 7




Figure 8




Figure 9

