




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

| | | | | | | | |
|--|-------------------------|---|----|-----------|-------------------------|----|---|
| (51) International Patent Classification ⁶ : C07D 401/04, A61K 31/47, C07D 401/14 | A1 | (11) International Publication Number: WO 99/64411 (43) International Publication Date: 16 December 1999 (16.12.99) | | | | | |
| <p>(21) International Application Number: PCT/EP99/03936</p> <p>(22) International Filing Date: 8 June 1999 (08.06.99)</p> <p>(30) Priority Data:</p> <table border="0"> <tr> <td>9812410.0</td> <td>10 June 1998 (10.06.98)</td> <td>GB</td> </tr> <tr> <td>9812408.4</td> <td>10 June 1998 (10.06.98)</td> <td>GB</td> </tr> </table> <p>(71) Applicant (for all designated States except US): GLAXO WELLCOME SPA [IT/IT]; Via Alessandro Fleming, 2, I-37100 Verona (IT).</p> <p>(72) Inventor; and (75) Inventor/Applicant (for US only): DI FABIO, Romano [IT/IT]; Glaxo Wellcome S.p.A., Via Alessandro Fleming, 2, I-37100 Verona (IT).</p> <p>(74) Agent: FILLER, Wendy, Anne; Glaxo Wellcome plc, Glaxo Wellcome House, Berkeley Avenue, Greenford, Middlesex UB6 0NN (GB).</p> | 9812410.0 | 10 June 1998 (10.06.98) | GB | 9812408.4 | 10 June 1998 (10.06.98) | GB | <p>(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published With international search report.</p> |
| 9812410.0 | 10 June 1998 (10.06.98) | GB | | | | | |
| 9812408.4 | 10 June 1998 (10.06.98) | GB | | | | | |

(54) Title: TETRAHYDROQUINOLINE DERIVATIVES AS GLYCINE ANTAGONISTS



(57) Abstract

Compounds of formula (I), or a salt or a non toxic metabolically labile esters thereof, wherein Y represents a carbon atom; Z is the group CH which is linked to the group Y via a double bond and X is CH or Z is methylene or NR₁₁ and X is a carbon atom linked to the group Y via a double bond; A represents a C₁₋₂alkylene chain and which chain may be substituted by one or two groups selected from C₁₋₆alkyl optionally substituted by hydroxy, amino, C₁₋₄alkyl amino or C₁₋₄dialkyl amino or which chain may be substituted by the group =O; R represents a halogen atom or C₁₋₄alkyl group; R₁ represents a hydrogen, a halogen atom or C₁₋₄alkyl group; R₂ represents optionally substituted phenyl, a 5 membered heteroaryl group containing 1 to 3 heteroatoms selected from oxygen, sulphur and nitrogen or 6 membered heteroaryl group containing 1 to 3 nitrogen atoms, processes for their preparation and their use as glycine antagonists.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

| | | | | | | | |
|----|--------------------------|----|--|----|--|----|--------------------------|
| AL | Albania | ES | Spain | LS | Lesotho | SI | Slovenia |
| AM | Armenia | FI | Finland | LT | Lithuania | SK | Slovakia |
| AT | Austria | FR | France | LU | Luxembourg | SN | Senegal |
| AU | Australia | GA | Gabon | LV | Latvia | SZ | Swaziland |
| AZ | Azerbaijan | GB | United Kingdom | MC | Monaco | TD | Chad |
| BA | Bosnia and Herzegovina | GE | Georgia | MD | Republic of Moldova | TG | Togo |
| BB | Barbados | GH | Ghana | MG | Madagascar | TJ | Tajikistan |
| BE | Belgium | GN | Guinea | MK | The former Yugoslav Republic of Macedonia | TM | Turkmenistan |
| BF | Burkina Faso | GR | Greece | | | TR | Turkey |
| BG | Bulgaria | HU | Hungary | ML | Mali | TT | Trinidad and Tobago |
| BJ | Benin | IE | Ireland | MN | Mongolia | UA | Ukraine |
| BR | Brazil | IL | Israel | MR | Mauritania | UG | Uganda |
| BY | Belarus | IS | Iceland | MW | Malawi | US | United States of America |
| CA | Canada | IT | Italy | MX | Mexico | UZ | Uzbekistan |
| CF | Central African Republic | JP | Japan | NE | Niger | VN | Viet Nam |
| CG | Congo | KE | Kenya | NL | Netherlands | YU | Yugoslavia |
| CH | Switzerland | KG | Kyrgyzstan | NO | Norway | ZW | Zimbabwe |
| CI | Côte d'Ivoire | KP | Democratic People's Republic of Korea | NZ | New Zealand | | |
| CM | Cameroon | | | PL | Poland | | |
| CN | China | KR | Republic of Korea | PT | Portugal | | |
| CU | Cuba | KZ | Kazakistan | RO | Romania | | |
| CZ | Czech Republic | LC | Saint Lucia | RU | Russian Federation | | |
| DE | Germany | LI | Liechtenstein | SD | Sudan | | |
| DK | Denmark | LK | Sri Lanka | SE | Sweden | | |
| EE | Estonia | LR | Liberia | SG | Singapore | | |

TETRAHYDROQUINOLINE DERIVATIVES AS GLYCINE ANTAGONISTS

This invention relates to 1, 2, 3, 4 tetrahydroquinoline derivatives, to processes for their preparation, to pharmaceutical compositions containing them and to their use in medicine. In particular, it relates to 1, 2, 3, 4 tetrahydroquinoline derivatives which are potent and specific antagonists of excitatory amino acids.


Carling *et al.*, Bioorganic and Medicinal Chemistry Letters, Vol. 13 pp. 65-70, 1993, teaches 4-substituted-2-carboxy tetrahydroquinolines having good *in vitro* affinity for the glycine modulatory site of the NMDA receptor complex but at best only weak *in vivo* activity. More particularly, it teaches that such derivatives substituted at the 4 position by the group CH₂CO₂H or CH₂CONHPh have little or no *in vivo* activity when administered systemically (ip).

WO 97/12870 and WO 98/07704 describe novel 4-substituted-2-carboxy-tetrahydroquinoline derivatives which not only have a good *in vitro* affinity for the strychnine insensitive glycine binding site associated with the NMDA receptor complex but also have good *in vivo* activity when administered intravenously (iv).

We have now discovered a novel group of 4-substituted-2-carboxy tetrahydroquinoline having a particularly useful profile of activity as selective antagonists for the strychnine insensitive glycine binding site associated with the NMDA receptor complex.

Thus the present invention provides a compound of formula (I)

2



(I)

- 5 or a salt or a non toxic metabolically labile esters thereof, wherein
 Y represents a carbon atom;
 Z is the group CH which is linked to the group Y via a double bond and
 X is CH or Z is methylene or NR₁₁, and X is a carbon atom linked to the
 group Y via a double bond ;
- 10 A represents a C₁₋₂ alkylene chain and which chain may be substituted by
 one or two groups selected from C₁₋₆ alkyl optionally substituted by
 hydroxy, amino, C₁₋₄ alkyl amino or C₁₋₄ dialkyl amino or which chain may
 be substituted by the group =O;
- R represents a halogen atom or C₁₋₄ alkyl group;
- 15 R₁ represents a hydrogen, a halogen atom or C₁₋₄ alkyl group;
 R₂ represents phenyl which may be substituted with up to 3 groups
 selected from halogen, hydrogen, or (CH₂)_nR₃ wherein R₃ is COR₄,
 NR₆R₅, NHCOR₇, NHCONR₉R₈ or NH SO₂ R₁₀ group or R₂ is a 5
 membered heteroaryl group containing 1 to 3 heteroatoms selected from
 20 oxygen, sulphur and nitrogen; or 6 membered heteroaryl group
 containing 1 to 3 nitrogen atoms
- R₄ represents an amino, a hydroxyl or C₁₋₄ alkoxy group;
 R₅ and R₆ each independently represents hydrogen or C₁₋₄ alkyl group
 or
- 25 R₅ and R₆ together with the nitrogen atom to which they are attached
 represent a saturated 5-7 membered heterocyclic group optionally

containing an additional heteroatom selected from oxygen, sulphur or nitrogen

R₇ represents a hydrogen atom or C₁₋₄ alkyl, C₁₋₄ alkoxy, or phenyl;

R₈ represents hydrogen or C₁₋₄ alkyl group;

5 R₉ represents hydrogen, optionally substituted C₁₋₄ alkyl (optionally substituted by one or more hydroxy, carboxyl and amino group), phenyl;

R₁₁ represents hydrogen or C₁₋₄ alkyl group;

R₁₀ represents hydrogen, C₁₋₄ alkyl or a nitrogen protecting group.

n is zero or an integer from 1 to 2;

10

A further embodiment of the invention provides compounds of formula(I) or a salt or a non toxic metabolically labile esters thereof, wherein

Y represents a carbon atom;

15 Z is the group CH which is linked to the group Y via a double bond and X is CH or Z is methylene or NR₁₁ and X is a carbon atom linked to the group Y via a double bond ;

A represents a C₁₋₂ alkylene chain and which chain may be substituted by one or two groups selected from C₁₋₆ alkyl optionally substituted by hydroxy, amino, C₁₋₄ alkyl amino or C₁₋₄ dialkyl amino or which chain may be substituted by the group =O;

20

R represents a halogen atom;

R₁ represents a hydrogen or a halogen atom;

25 R₂ represents phenyl which may be substituted with up to 3 groups selected from halogen, hydrogen, or (CH₂)_nR₃ wherein R₃ is COR₄, NR₆R₅, NHCOR₇, NHCONR₉R₈ or NH SO₂ R₁₀ group or R₂ is a 5 membered heteroaryl group containing 1 to 3 heteroatoms selected from oxygen, sulphur and nitrogen; or 6 membered heteroaryl group containing 1 to 3 nitrogen atoms

30 R₄ represents an amino or a hydroxyl;

R₅ and R₆ each independently represents hydrogen or C₁₋₄ alkyl group or

R₅ and R₆ together with the nitrogen atom to which they are attached represent a saturated 5-7 membered heterocyclic group optionally

containing an additional heteroatom selected from oxygen, sulphur or nitrogen

R₇ represents a hydrogen atom or C₁₋₄ alkyl, C₁₋₄ alkoxy, or phenyl;

R₈ represents hydrogen or C₁₋₄ alkyl group;

5 R₉ represents hydrogen, optionally substituted C₁₋₄ alkyl (optionally substituted by one or more hydroxy carboxyl and amino group), phenyl;

R₁₁ represents hydrogen or C₁₋₄ alkyl group;

R₁₀ represents hydrogen, C₁₋₄ alkyl or a nitrogen protecting group;

10 n is zero or an integer from 1 to 2 with the proviso that when X is a carbon atom linked to the group Y via a double bond then R₁ is hydrogen;

For use in medicine the salts of the compounds of formula (I) will be physiologically acceptable thereof. Other salts however may be useful in the preparation of the compounds of formula (I) or physiologically acceptable salts thereof. Therefore, unless otherwise stated, references to salts include both physiologically acceptable salts and non-physiologically acceptable salts of compounds of formula (I).

20 Suitable physiologically acceptable salts of compounds of the invention include base addition salts and, where appropriate, acid addition salts. Suitable physiologically acceptable base addition salts of compounds of formula (I) include alkali metal or alkaline metal salts such as sodium, potassium, calcium, magnesium and ammonium salts, formed with amino acids (e.g. lysine and arginine) and organic bases (e.g. procaine, phenylbenzylamine, ethanolamine diethanolamine and N-methyl glucosamine).

30 The compounds of formula (I) and/or salts thereof may form solvates (e.g. hydrates) and the invention includes all such solvates.

The term halogen refers to a fluorine, chlorine, bromine or iodine atom.

The term C₁₋₄ alkyl as used herein as a group or part of a group refers to a straight or branched chain alkyl group containing from 1 to 4 carbon atom, examples of such groups include methyl, ethyl, propyl, isopropyl, n-butyl, isobutyl, secondary butyl or tertiary butyl.

5

When R₂ is a 5 or 6 membered heteroaryl group this may be for example furanyl, thiophenyl, imidazolyl, thiazolyl, oxazolyl, pyridyl or pyrimidinyl.

10

When R₅ and R₆ together with the nitrogen atom to which they are attached form a saturated 5-7 membered heterocyclic group optionally containing an additional heteroatom selected from oxygen, sulphur or nitrogen this may be morpholino, 2,6 dimethylmorpholino, thiomorpholino, piperidino, pyrrolidino, piperazino or N-methylpiperazino.

15

When R₂ is a substituted phenyl group this is conveniently a mono substituted phenyl group. The substituent is conveniently in the meta position or more conveniently in the para position.





20

When X-Y represents a double bond, the compounds of formula (I) possess at least one asymmetric carbon atom (namely the carbon atom occupying the 2 position of the 1, 2, 3, 4 tetrahydroquinoline ring system) and other asymmetric carbon atoms are possible in the group R₂. It is to be understood that all enantiomers and diastereomers and mixtures thereof are encompassed within the scope of the present invention.

25

When X-Y represents a single bond, the compounds of formula (I) possess at least two asymmetric carbon atoms (namely the carbon atom occupying the 2 and 4 position of the 1, 2, 3, 4 tetrahydroquinoline ring system) and these may be represented by the formulae (1a, 1b, 1c and 1d).

30



5 The solid wedge shaped bond indicates that the bond is above the plane of the paper and is referred to as the β configuration. The broken indicates that the bond is below the plane of the paper and is referred to as α configuration.

10 Further other asymmetric carbon atoms are possible in the groups R_2 . It is to be understood that all enantiomers and diastereomers and mixtures thereof are encompassed within the scope of the present invention.

15 Non-toxic metabolically labile esters of compound of formula (I) are esters of compounds of formula (I) that are hydrolysed in vivo to afford said compound of formula I and a physiologically acceptable alcohol. Non toxic metabolically esters of compound of formula (I) may be formed by esterification, for example of any of the carboxylic acid groups in the parent compound of general formula (I) with, where appropriate, prior protection of any other reactive groups present in the molecule, followed by deprotection if required. Examples of such metabolically labile esters include C_{1-4} alkyl esters e.g. methyl or ethyl esters, substituted or

20

unsubstituted aminoalkyl esters (e.g. aminoethyl, 2-(N,N- diethylamino)
 ethyl, or 2-(4-morpholino)ethyl esters or acloxyalkyl esters such as,
 acyloxymethyl or 1-acyloxyethyl e.g. pivaloyloxymethyl, 1-
 5 pivaloyloxyethyl, acetoxymethyl, 1- acetoxyethyl, 1-(1-methoxy-1-
 methyl)ethylcarbonyloxyethyl, 1- benzoyloxyethyl,
 isopropoxycarbonyloxymethyl, 1-isopropoxycarbonyloxyethyl,
 cyclohexylcarbonyloxymethyl, 1-cyclohexylcarbonyloxyethyl ester,
 cyclohexyloxycarbonyloxymethyl, 1-cyclohexyloxycarbonyloxyethyl, 1-
 (4-tetrahydropyranyloxy)carbonyloxyethyl or 1-(4-
 10 tetrahydropyranyl)carbonyloxyethyl.

The group R is conveniently chlorine.

The group R₁ is conveniently a hydrogen or a chlorine atom.

15 A preferred class of compounds of formula(I) is that wherein R is chlorine and R₁ is a hydrogen or a chlorine atom.

A further preferred class of compounds of formula(I) is that wherein R is chlorine and R₁ is a hydrogen atom.

20

When X-Y is a single bound, a preferred class of compounds of formula
 (I) is that in which the carbon atom in 4 position is β configuration and
 the carbon atom in 2 position is in α configuration (1a) and that in which
 the carbon atom in 4 position is α configuration and the carbon atom in 2
 25 position is in β configuration (1c).

When A is an optionally substituted C₁₋₂ alkylene chain this may be, for example, methylene, ethylene or C=O.

30 A preferred class of compounds of formula (I) includes those wherein A is a chain selected from -CH₂-, -(CH₂)₂-, C=O.

When Z is a group NR₁₁ this is conveniently the group NH.

A preferred class of compounds of formula (I) includes those wherein Z is CH which is linked to the group Y via a double bond, a methylene or a NH group.

5 When R_2 is an optionally substituted phenyl group this is conveniently phenyl substituted by a single substituent selected from $(CH_2)_nNR_6R_5$ in which R_5 is hydrogen and R_6 is hydrogen, C_{1-4} alkyl (e.g. methyl, ethyl) or NR_6R_5 represents a saturated 6 membered ring containing oxygen e.g. morpholino; $(CH_2)_nNHCOR_7$ wherein R_7 is hydrogen, alkyl e.g. methyl, isopropyl, isobutyl, phenyl; $(CH_2)_nNHCONHR_9$ wherein R_9 is hydrogen; $(CH_2)_nNH SO_2 R_{10}$ in which R_{10} is alkyl e.g. methyl. n is zero or an integer from 1 to 2; Examples of such R_2 groups include phenyl (optionally substituted by amino, t-butoxycarbonylamino, acetylamino or methanesulphonylamino)

15

When R_2 is substituted phenyl the substituents are conveniently in the meta or more preferably in the para position.

20 When R_2 is a 5 or 6 membered heteroaryl group as above defined this is conveniently pyridyl e.g. 3-pyridyl.

A preferred class of compounds of formula (I) is that wherein R_2 represents phenyl (optionally substituted by acetylamino, methanesulphonylamino) or 3-pyridyl. Within this class those wherein R_2 is phenyl are particularly preferred.

25

A further preferred class of compounds of formula (I) is that wherein X is a carbon atom linked to the group Y via a double bond.

30 A preferred group of compounds of formula(I) is that wherein A is a chain selected from $-CH_2-$ or $-(CH_2)_2-$, Z is a group CH which is linked to the group Y via a double bond or a methylene group, or A is the chain CO and Z is an NH group, R is chlorine, R_1 is chlorine or hydrogen and

R₂ is phenyl (optionally substituted by acetamino or methanesulphonylamino) or 3-pyridyl.

5 Specific preferred compounds of the invention include:

(±)7-chloro-4-(2-oxo-1-phenyl-3-pyrrolidinylidene)-1,2,3,4-tetrahydro-2-quinoline carboxylic acid ,

(±)7-chloro-4-(1-phenyl-Δ³-pyrrolin-2-one-3yl)-1,2,3,4-tetrahydro-2-quinoline carboxylic acid ,

10 and physiologically acceptable salts (e.g. sodium salt) non-toxic metabolically labile esters or enantiomers thereof.

(-)-Sodium 7-chloro-4-(2-oxo-1-phenyl-3-pyrrolidinylidene)-1,2,3,4-tetrahydro-2-quinolinecarboxylate.

(-)-Sodium 7-chloro-4-(1-phenyl-Δ³-pyrrolin-2-one-3yl)-1,2,3,4-tetrahydroquinoline-2-carboxylate,

15 (+)-Sodium 7-chloro-4-(1-phenyl-Δ³-pyrrolin-2-one-3yl)-1,2,3,4-tetrahydroquinoline-2-carboxylate.

Further specific preferred compounds of the invention include:

20 (±)-7-chloro-4-(1-(3-pyridin)-Δ³-pyrrolin-2-one-3yl)-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

(±)-7-chloro-4-(1-phenylΔ³-5,6-dihydro-pyridin-2-one-3yl)-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

25 (±)-5,7-dichloro-4-(1-phenyl-Δ³-pyrrolin-2-one-3yl)-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

(+/-)-7-chloro-4-(1-(4-acetylamino)-1-phenyl-Δ³-pyrrolin-2-one-3yl)-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

(+/-)-7-chloro-4-(1-(4-methanesulfonylamino)-1-phenyl-Δ³-pyrrolin-2-one-3yl)-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

30 (±)-7-chloro-4-(2-oxo-1-phenyl-3-piperidinylidene)-1,2,3,4-tetrahydro-2-quinolinecarboxylic acid,

(±)-7-chloro-4-(2,5-dioxo-1-phenyl-imidazolidin-4-ylidene)-1,2,3,4-tetrahydro-2-quinolinecarboxylic acid,
(±)-7-chloro-4-(2-oxo-1-(pyridin-3-yl)-pyrrolidin-3-ylidene)-1,2,3,4-tetrahydro-2-quinolinecarboxylate,
5 (±)-7-chloro-4-(2-oxo-1-(4-acetylamino)phenyl-pyrrolidin-3-ylidene)-1,2,3,4-tetrahydro-2-quinolinecarboxylic acid,
(±)-7-chloro-4-(2-oxo-1-((4-methanesulfonyl amino)phenyl-pyrrolidin-3-ylidene)-1,2,3,4-tetrahydro-2-quinolinecarboxylic acid,
10 5,7-dichloro-4-(2-oxo-1-(phenyl)-pyrrolidin-3-ylidene)-1,2,3,4-tetrahydro-2-quinoline carboxylic acid (enantiomer A);
5,7-dichloro-4-(2-oxo-1-phenyl- Δ^3 -pyrrolin-2-one-3-yl)-1,2,3,4-tetrahydro-quinoline-2-carboxylic acid (enantiomer A);
and physiologically acceptable salts (e.g. sodium salts), non-toxic metabolically labile esters or enantiomers thereof.

15 The compounds of formula (I) and/or physiologically acceptable salts thereof are excitatory amino acid antagonists. More particularly they are potent antagonists at the strychnine insensitive glycine binding site associated with the NMDA receptor complex. As such they are potent
20 antagonists of the NMDA receptor complex. These compounds are therefore useful in the treatment or prevention of neurotoxic damage or neurodegenerative diseases. Thus the compounds are useful for the treatment of neurotoxic injury which follows cerebral stroke, thromboembolic stroke, hemorrhagic stroke, cerebral ischemia, cerebral
25 vasospasm, hypoglycemia, amnesia, hypoxia, anoxia, perinatal asphyxia cardiac arrest. The compounds are useful in the treatment of chronic neurodegenerative diseases such as: Huntingdon's disease, Alzheimer's senile dementia, amyotrophic lateral sclerosis, Glutaric Acidaemia type, multi-infarct dementia, status epilepticus, contusive injuries (e.g. spinal
30 cord injury and head injury), viral infection induced neurodegeneration (e.g. AIDS, encephalopathies), Down syndrome, ocular neurodegeneration (e.g. glaucoma), epilepsy, schizophrenia, depression, migraine, headaches including cluster headaches and or tension headaches, anxiety, pain (e.g. inflammatory pain and neuropathic pain), neurogenic bladder, emesis

,irritative bladder disturbances, drug dependency, including withdrawal symptoms from alcohol, cocaine, opiates, nicotine (e.g. smoking cessation) benzodiazepines and. inhibition of tolerance induced by opioids (i.e morphine).

5

The potent and selective action of the compound of the invention at the strychnine-insensitive glycine binding site present on the NMDA receptor complex may be readily determined using conventional test procedures. Thus the ability to bind at the strychnine insensitive glycine binding site was determined using the procedure of Kishimoto H *et al.*, J Neurochem 10 1981, 37, 1015-1024. The selectivity of the action of compounds of the invention for the strychnine insensitive glycine site was confirmed in studies at other ionotropic known excitatory amino acid receptors. Thus compounds of the invention were found to show little or no affinity for the 15 kainic acid (kainate) receptor, α -amino-3-hydroxy-5-methyl-4-isoxazole-propionic acid (AMPA) receptor or at the NMDA binding site.

Compounds of the invention may be found found to inhibit NMDA induced convulsions in mice using the procedure Chiamulera C *et al.*, 20 Psychopharmacology (1990), 102, 551-552.

The neuroprotective activity of the compounds of the invention may be demonstrated in the middle cerebral artery occlusion preparation in mice, using the procedure described by Chiamulera C. et al., European 25 Journal of Pharmacology, 216 (1992) pp. 335-336.

The ability of compounds of the invention to alleviate withdrawal symptoms from nicotine following smoking cessation may be demonstrated in conventional tests of nicotine induced relapse using the 30 procedure described in C. Chiamulera et al., Arch. Pharmacol., 358, 1998.

The invention therefore provides for the use of a compound of formula (I) and/or physiologically acceptable salts or non-toxic metabolically labile

esters thereof for use in therapy and in particular use as medicine for antagonising the effects of excitatory amino acids upon the NMDA receptor complex.

5 The ability of compounds of the invention to inhibit pain may be demonstrated in conventional analgesic screen such as those described by Dubuisson and Dennis, *Pain*, 1977, 4:161-174; J.J. Bennett and J.K Xue, *Pain*, 1988, 41, 87-107.

10 The invention also provides for the use of a compound of formula (I) and/or a physiologically acceptable salt or non-toxic metabolically labile esters thereof for the manufacture of a medicament for antagonising the effects of excitatory amino acids upon the NMDA receptor complex.

15 According to a further aspect, the invention also provides for a method for antagonising the effects of excitatory amino acids upon the NMDA receptor complex, comprising administering to a patient in need thereof an antagonistic amount of a compound of formula (I) and/or a physiologically acceptable salt thereof.

20 It will be appreciated by those skilled in the art that reference herein to treatment extends to prophylactics as well as the treatment of established diseases or symptoms.

25 It will further be appreciated that the amount of a compound of the invention required for use in treatment will vary with the nature of the condition being treated, the route of administration and the age and the condition of the patient and will be ultimately at the discretion of the attendant physician. In general however doses employed for adult

30 human treatment will typically be in the range of 2 to 800mg per day, dependent upon the route of administration.

Thus for parenteral administration a daily dose will typically be in the range 20-100mg, preferably 60-80mg per day. For oral administration a

daily dose will typically be within the range 200-800mg, e.g. 400-600mg per day.

5 The desired dose may conveniently be presented in a single dose or as divided doses administered at appropriate intervals, for example as two, three, four or more sub-doses per day.

10 While it is possible that, for use in therapy, a compound of the invention may be administered as the raw chemical, it is preferable to present the active ingredient as a pharmaceutical formulation.

15 The invention thus further provides a pharmaceutical formulation comprising a compound of formula (I) or a pharmaceutically acceptable salt or non-toxic metabolically labile esters thereof together with one or more pharmaceutically acceptable carriers thereof and, optionally, other therapeutic and/or prophylactic ingredients. The carrier(s) must be 'acceptable' in the sense of being compatible with the other ingredients of the formulation and not deleterious to the recipient thereof.

20 The compositions of the invention include those in a form especially formulated for oral, buccal, parenteral, inhalation or insufflation, implant or rectal administration.

25 Tablets and capsules for oral administration may contain conventional excipients such as binding agents, for example, syrup, acacia, gelatin, sorbitol, tragacanth, mucilage of starch or polyvinylpyrrolidone; fillers, for example, lactose, sugar, microcrystalline cellulose, maize-starch, calcium phosphate or sorbitol; lubricants, for example, magnesium stearate, stearic acid, talc, polyethylene glycol or silica; disintegrants, for
30 example, potato starch or sodium starch glycollate, or wetting agents such as sodium lauryl sulphate. The tablets may be coated according to methods well known in the art. Oral liquid preparations may be in the form of, for example, aqueous or oily suspensions, solutions emulsions, syrups or elixirs, or may be presented as a dry product for constitution

with water or other suitable vehicle before use. Such liquid preparations may contain conventional additives such as suspending agents, for example, sorbitol syrup, methyl cellulose, glucose/sugar syrup, gelatin, hydroxyethylcellulose, carboxymethyl cellulose, aluminium stearate gel
5 or hydrogenated edible fats; emulsifying agents, for example, lecithin, sorbitan mono-oleate or acacia; non-aqueous vehicles (which may include edible oils), for example, almond oil, fractionated coconut oil, oily esters, propylene glycol or ethyl alcohol; solubilizers such as surfactants for example polysorbates or other agents such as cyclodextrins; and
10 preservatives, for example, methyl or propyl p- hydroxybenzoates or ascorbic acid. The compositions may also be formulated as suppositories, e.g. containing conventional suppository bases such as cocoa butter or other glycerides.

15 For buccal administration the composition may take the form of tablets or lozenges formulated in conventional manner.

The composition according to the invention may be formulated for parenteral administration by injection or continuous infusion.
20 Formulations for injection may be presented in unit dose form in ampoules, or in multi-dose containers with an added preservative. The compositions may take such forms as suspensions, solutions, or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilising and/or dispersing agents.
25 Alternatively the active ingredient may be in powder form for constitution with a suitable vehicle, e.g. sterile, pyrogen-free water, before use.

For administration by inhalation the compounds according to the invention are conveniently delivered in the form of an aerosol spray
30 presentation from pressurised packs, with the use of a suitable propellant, such as dichlorodifluoromethane, trichlorofluoromethane, dichloro-tetrafluoroethane, carbon dioxide or other suitable propellants, such as dichlorodifluoromethane, trichlorofluoromethane, dichloro-tetrafluoroethane, carbon dioxide or other suitable gases, or from a

nebuliser. In the case of a pressurised aerosol the dosage unit may be determined by providing a valve to deliver a metered amount.

5 Alternatively, for administration by inhalation or insufflation, the compounds according to the invention may take the form of a dry powder composition, for example a powder mix of the compound and a suitable carrier such as lactose or starch. The powder composition may be presented in unit dosage form in, for example, capsules or cartridges of e.g. gelatin, or blister packs from which the powder may be administered
10 with the aid of an inhaler or insufflator.


The composition according to the invention may also be formulated as a depot preparation. Such long acting formulations may be administered by implantation (for example subcutaneously or intramuscularly) or by
15 intramuscular injection. Thus for example, the compounds of the invention may be formulated with suitable polymeric or hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

20 The compositions according to the invention may contain between 0.1 - 99% of the active ingredient, conveniently from 30- 95% for tablets and capsules and 3-50% for liquid preparations.

25 Compounds of general formula (I) enantiomers and salts thereof may be prepared by the general methods outlined hereinafter. In the following description, the groups R, R₁, R₂, A, Z, X and Y are as defined for the compounds of formula (I) unless otherwise stated.

30 Compounds of formula (I) and enantiomers thereof may be prepared by the cyclisation of a compound of formula (II) in which R₁₂ is a carboxylic protecting group, R₁₃ represents a bromine or iodine atom, R₁₄ represents hydrogen or a nitrogen protecting group.

16



5 followed where necessary or desired by removal of one or more protecting groups.

10 In one embodiment of this process the reaction may be carried out using a catalytic amount of a Palladium (O) complex such as tetrakis(triphenylphosphine)palladium and a suitable organic base such as trialkylamine e.g. triethylamine or inorganic base, e.g. potassium carbonate.

15 The reaction is conveniently carried out in an aprotic polar solvent such as acetonitrile, dimethylformamide or in aprotic apolar solvent such as hydrocarbon (ie toluene, xylene, hexane) at a temperature within the range of 60°C to 150°C followed, where necessary or desired, by subsequent removal of the carboxyl protecting group R₁₂ and any protecting group R₁₄.

20 In a further embodiment of the process the reaction is carried out using a catalytic amount of a Pd(II) salt such as: palladium acetate or palladium dichloride in the presence of a suitable organic base such as trialkyl amine e.g. triethylamine and of a triarylphosphine such as triphenylphosphine.

25 The reaction is carried out in an aprotic solvent such as acetonitrile or dimethylformamide and preferably with heating followed, where

necessary or desired, by subsequent removal of the carboxyl protecting group R₁₂ and any nitrogen protecting group R₁₄.

5 Compounds of formula (I) wherein X-Y is a double bond may be regioselectively prepared by carrying out the cyclisation reaction in an aprotic apolar solvent such as toluene in the presence of catalytic amount of a Palladium (0) complex such as tetrakis(triphenylphosphine)palladium and a suitable organic base such as trialkylamine e.g. triethylamine or inorganic base, e.g. potassium
10 carbonate.

Compounds of formula (I) wherein X-Y is a single bond may be prepared by carrying out the reaction the cyclisation reaction in an aprotic polar solvent (such as acetonitrile, dimethylformamide) in the presence of a catalytic amount of a Pd(II) salt such as: palladium acetate or palladium
15 dichloride in the presence of a suitable organic base such as trialkyl amine e.g. triethylamine and of a triarylphosphine such as triphenylphosphine.


20 Suitable carboxyl protecting groups R₁₂ for use in this reaction include alkyl, such as ethyl, trichloroalkyl, trialkylsilylalkyl, or arylmethyl groups such as benzyl, nitrobenzyl or trityl.

Further convenient carboxyl protecting groups are those having a chiral group derived from chiral alcohols such as (+)-S-indanol, (+)-S-methyl mandelate, chiral (C₁₋₄)alkyl lactate: i.e. (+)-R- or (-)-S-methyl lactate, (+)-
25 R-t-butyl lactate, (+)-R- or (-)-S-ethyl lactate, (-)-S-isopropyl lactate, (-)-S-butyl lactate, (+)-R-isobutyl lactate or chiral aralkyl lactate (i.e. benzyl lactate), (-)-S-perillyl alcohol, (-)-methyl-(R)-3-hydroxy-2-methylpropionate, (-)-R-2-butanol, (-)-S-2-methyl-1-butanol.

R₁₂ is preferably an ethyl, benzyl group or a group derived from a
30 chiral(C₁₋₄) alkyl lactate alcohol (eg (+)-R-t-butyl lactate (-)-S-butyl lactate, (+)-R-isobutyl lactate alcohol).

When R_{14} is nitrogen protecting examples of suitable groups include alkoxy carbonyl e.g. t-butoxycarbonyl, arylsulphonyl e.g. phenylsulphonyl or 2-trimethylsilylethoxymethyl.


5 Compounds of formula (II) may be prepared from compound of formula (III) in which R_{12} is a carboxyl protecting group and R_{14} is hydrogen or a nitrogen protecting group as defined in formula (II) and R_{13} represents a bromine or iodine atom.



(III)


10

by reaction with an appropriate phosphorus reagent capable of converting the group CHO into the group :



15 followed, where necessary or desired, by removal of the carboxyl protecting group R_{12} and nitrogen protecting group R_{13} .

In one embodiment of this process the reaction may be carried out using a phosphorus ylide of formula (IV)



(IV)


20

wherein R_{15} is an alkyl or phenyl group.

The reaction is carried out in an aprotic solvent such as acetonitrile or dimethylformamide at a temperature ranging from -20°C to the reflux temperature of the solvent.

5 Compounds of formula (III) and (IV) are either known compounds or may be prepared by analogous methods to those used for known compounds.


A convenient method for preparing compounds of formula (III) is reacting
 10 compound of formula (V) in which R_{12} is a carboxyl protecting group and R_{14} is hydrogen or a nitrogen protecting group as defined in formula (II) and R_{13} represents a bromine or iodine atom with an allyltintri-
 halide.(VI) followed by ozonization reaction



15 The reaction conveniently takes place in a solvent such as hydrocarbon e.g. Toluene or halogenated hydrocarbon (e.g. dichloro methane at a temperature ranging from -78°C to room temperature).

The ozonization may be carried out by passing a stream of ozone into a solution in the presence of dimethyl sulphide or triphenylphosphine in a
 20 suitable solvent such as halohydrocarbon (e.g dichloromethane) at low temperature e.g. -78°C .

Alternatively compounds (III) may be prepared by aldolic reaction of the
 25 imino compound (V), with the enol ether (VII), wherein R_{16} is a C_{1-4} alkyl group.



The reaction may be carried out in a solvent such as methylene chloride or acetonitrile in the presence of a Lewis acid such as Ytterbium triflate.

In any of the above reactions the carboxyl protecting group may be removed by conventional procedures known for removing such groups. Thus compounds where R₁₂ is a benzyl, ethyl or (+)-R- or (-)-S-t-butyl lactate group may be removed by hydrolysis using an alkali metal hydroxide e.g. lithium hydroxide or sodium hydroxide in a suitable solvent such as ethanol or isopropanol, water or mixtures thereof, followed, where desired or necessary, by that addition of a suitable acid e.g. hydrochloric acid to give the corresponding free carboxylic acid.

10 In any of the above reactions the nitrogen protecting group may be removed by conventional procedures known for removing such groups, for example by acid or base hydrolysis. Thus when R₁₄ is alkoxycarbonyl e.g. t-butoxycarbonyl or phenylsulphonyl it may be removed by alkaline hydrolysis using for example lithium hydroxide in a
15 suitable solvent such as tetrahydrofuran or an alkanol e.g. isopropanol. Alternatively the alkoxycarbonyl group may be removed by acid hydrolysis.

20 Physiologically acceptable salts of compounds of formula (I) may be prepared by treating the corresponding acid with an appropriate base in a suitable solvent. For example the sodium or potassium salt may be prepared by treating a solution of the corresponding acid of a compound of formula (I) with sodium or potassium 2-ethylhexanoate with alkali or alkaline metal hydroxide, or the corresponding carbonate or bicarbonate
25 thereof. Alternatively alkali or alkaline metal salts may be prepared by direct hydrolysis of carboxyl protected derivatives of compounds of formula (I) with the appropriate alkali or alkaline metal hydroxide.

30 Metabolically labile esters of compounds of formula (I) may be prepared by esterification of the carboxylic acid group or a salt thereof or by trans esterification using conventional procedures. Thus, for example, acyloxyalkyl esters may be prepared by reacting the free carboxylic acid or a salt thereof with the appropriate acyloxylalkyl halide in a suitable solvent such as dimethylformamide. For the esterification of the free

carboxyl group this reaction is preferably carried out in the presence of a quaternary ammonium halide such as tetrabutylammonium chloride or benzyltriethylammonium chloride.

- 5 Specific enantiomers of the compounds of formula (I) may also be obtained from corresponding racemic compounds of formula (I) using chiral HPLC procedure.

10 Alternatively the enantiomers may be prepared by esterification of the corresponding racemic compounds of formula (I) with a suitable chiral alcohol, separating the resultant diastereomeric esters by conventional means e.g. chromatography or crystallisation followed by hydrolysis of the diastereomeric esters.

- 15 Suitable chiral alcohols for use in the process include (+)-S-indanol, (+)-S-methyl mandelate, chiral (C₁₋₄)alkyl lactate: i.e. (+)-R- or (-)-S-methyl lactate, (+)-R-t-butyl lactate, (+)-R- or (-)-S-ethyl lactate, (-)-S-isopropyl lactate, (-)-S-butyl lactate, (+)-R-isobutyl lactate or chiral aralkyl lactate (i.e. benzyl lactate), (-)-S-perillyl alcohol, (-)-methyl-(R)-3-hydroxy-2-
20 methylpropionate, (-)-R-2-butanol, (-)-S-2-methyl-1-butanol.

The diastereomeric esters of compounds of formula (I) may be prepared by conventional means such as reaction of the chiral alcohol with an activated derivative of a compound of formula (I) in an aprotic solvent
25 such as ether e.g. tetrahydrofuran.

The activated derivative of a compound of formula (I) may be prepared from compounds (I) using conventional means for preparing activated derivatives of a carboxylic acid groups such as those conveniently used in peptide synthesis.

30

A particularly convenient method of preparing the diastereomeric esters of compounds (I) is to prepare the activated derivative of compounds (I) in the presence of the chiral alcohol.

Thus for example racemic mixture of compounds (I) may be treated with the Mitsunobu combination of reagents, i.e. a dialkyl azo-dicarboxylate such as diethylazodicarboxylate and a triarylphosphine e.g. triphenylphosphine or trialkylphosphine (i.e. tributylphosphine) in the presence of the chiral alcohol.

The reaction conveniently takes place in the presence of a suitable solvent such as an ether (e.g. diethylether or tetrahydrofuran), a halohydrocarbon (e.g. dichloromethane) or a nitrile (e.g. acetonitrile) or a mixture thereof at a temperature ranging from 0-30°C.


The required single diastereomeric ester of compounds (I) may be obtained from the mixture thereof by conventional means, for example by the use of conventional chromatographic procedures such as preparative HPLC or by fractional crystallization.

Alternatively the required single diastereomeric ester of compound of formula (I) may be obtained using a suitable chiral protecting group R_{12} as defined in formula (II).

Specific enantiomers of compounds of formula (I) may be prepared from the corresponding single diastereomeric ester of compounds (I) by hydrolysis e.g. alkaline hydrolysis. Thus, for example, the hydrolysis may be carried using an alkali metal hydroxide e.g. sodium hydroxide or lithium hydroxide in a solvent such as an ether e.g. tetrahydrofuran and water.

Alternatively specific enantiomers of compounds of formula (I) may be prepared by stereoselective enzymatic hydrolysis of compounds of formula (VIII)

23



Wherein R_{17} is a carboxyl protecting group

Suitable carboxyl protecting group R_{17} for use in this reaction include C_{1-4} alkyl such as methyl, ethyl, propyl, butyl, or arylmethyl groups such as benzyl, nitrobenzyl or trityl.

Suitable enzymes for use in this reaction are lipase enzymes such as *Aspergillus niger* (AP-12) lipase-DS (*Aspergillus niger*, Amano), *Candida rugosa* lipase (Amano), *Candida cylindracea* lipase (Amano), *Alcaligenes* sp. lipase, *Rhizopus arrizus* lipase (Biotal), Wheat germ lipase (Sigma), *Rhizopus niveus* lipase (Amano), Promod 215-P protease (Biocatalyst), lipase E-7 (Themogen), lipase E-17 (Themogen). Further suitable enzymes which may be used in this reaction are porcine pancreatic lipase, alpha-chymotrypsin or trypsin.

A particular preferred enzyme for use in this reaction is *Aspergillus niger* (AP-12).

Resting cells of the following organisms may also be used in this reaction *Aspergillus ochraceus*, *Aspergillus niger*, *Aspergillus chevalieri* & *Aspergillus cervinus*.

The reaction is conveniently carried out in an aprotic solvent such as DMSO, tetrahydrofuran in the presence of a suitable aqueous buffer (i.e. phosphate buffer or $CaCl_2$). If required a solubilising agent such as Tween-80 may be added to the reaction mixture.

In a further process the enzyme may be immobilized and the reaction is carried out in essentially "neat" water-saturated organic solvents such as methyl *tert*-butyl ether or *tert*-amyl alcohol.

In order that the invention may be more fully understood the following examples are given by way of illustration only.

5 In the Intermediates and Examples unless otherwise stated:

Melting points (m.p.) were determined on a Gallenkamp m.p. apparatus and are uncorrected. All temperatures refer to °C. Infrared spectra were measured on a FT-IR instrument. Proton Magnetic Resonance (¹H-NMR) spectra were recorded at 400 MHz, chemical shifts are reported
10 in ppm downfield (δ) from Me₄Si, used as internal standard, and are assigned as singlets (s), doublets (d), doublets of doublets (dd), triplets (t), quartets (q) or multiplets (m). Column chromatography was carried out over silica gel (Merck AG Darmstadt, Germany). The following abbreviations are used in text: EA = ethyl acetate, CH = cyclohexane,
15 DCM = dichloromethane, THF = tetrahydrofuran, TFA = trifluoroacetic acid, TEA = triethylamine, DMF = dimethylformamide, Ac₂O = acetic anhydride, PPA = polyphosphoric acid, DBU = 1,8-diazobicyclo [5,4,0]undec-7-ene, DMSO = dimethylsulphoxide, IMS=mixture of Ethanol with 5% of methanol, LHDMS=Lithiumbis(trimethylsilyl)amide.
20 DIPEA=diisopropylethylamine Tlc refers to thin layer chromatography on silica plates, and dried refers to a solution dried over anhydrous sodium sulphate; r.t. (RT) refers to room temperature.
Enantiomer A or diastereoisomer A refer to a single enantiomer or a single diastereoisomer respectively whose absolute stereochemistry was not
25 characterized.

Intermediate 1

(±)-Ethyl 2-(5-chloro-2-iodoanilino)-4-pentenoate

To a solution of 2-iodo 4 chloro aniline (9.1g) in dry toluene (150 ml) ethyl glyoxylate (50% solution in toluene, 14.6 ml) and MgSO₄ (2 g) were
30 added and the resulting suspension was refluxed overnight. It was then filtered and concentrated to dryness under high vacuum at 50°C for 1.5 h. The resulting brown oil was dissolved in dichloromethane (150 ml) cooled to -78°C and TiCl₄ (99.995% purity, 4 ml) was added via syringe.

The suspension was stirred 15 min at -78°C, then allowed to warm to rt over 15 min before being cooled again to -78°C. Allyltributyltin (17 ml) was then added and the reaction allowed to proceed for 1 h. The black solution was poured into 200 ml of ethyl acetate and washed first with a saturated solution of NH₄Cl (2 x 150 ml), then with water and brine. The organic phase was dried and concentrated to give the crude product, which was purified by column chromatography (cyclohexane, then cyclohexane/ethyl acetate 98/2) to give the title compound (10.4 g) as a colourless oil.

NMR (CDCl₃) δ (ppm) 7.57 (d, 1H), 6.49 (dd, 1H), 6.45 (dd, 1H), 5.79 (m, 1H), 5.25 (dd, 1H) 5.24 (dd, 1H), 4.83 (d, 1H), 4.25 (q, 2H), 4.13 (m, 1H), 2.66 (m, 2H), 1.30 (t, 3H)

Intermediate 2

15 (±)-Ethyl 2-(5-chloro-2-iodoanilino)-4-oxobutanoate

A solution of intermediate 1 (5.2g) in dichloromethane (150 ml) was cooled to -78°C and ozone was bubbled through it until the clear solution became brick-red. At this point the flux of ozone was interrupted and the solution was purged with nitrogen for a few minutes. Triphenyl phosphine (7.1g) was added and stirring continued for 1.5 h, without control of the temperature. The resulting solution was poured into 200 ml of ethyl acetate and washed first with a saturated solution of NH₄Cl (2 x 150 ml), then with water and brine. The organic phase was dried and concentrated to give the crude product, which was purified by column chromatography (cyclohexane/ethyl acetate 80/20) to give the title compound (2.4g) as a colourless oil.

NMR (DMSO) δ (ppm) 9.80 (t, 1H), 7.57 (d, 1H), 6.55 (d, 1H), 6.51 (dd, 1H), 4.99 (d, 1H), 4.46 (m, 1H), 4.24 (q, 2H), 3.08 (m, 2H), 1.28 (t, 3H)

30 Intermediate 2a

(±) Ethyl 2-(3,5-dichloro-2-iodoanilino)-4-oxobutanoate

A solution of ethyl glyoxylate (50% solution in toluene, 1ml) and MgSO₄ (7 g) in toluene (30 ml) was refluxed in Dean-Stark apparatus for 0.5 hrs.

Then, 3,5,-chloro-2iodoaniline was added, and the mixture refluxed for 1 hr. Then mixture was cooled, filtered through celite to eliminate the MgSO₄, and concentrated. The resulting brown oil was dissolved in dichloromethane (15ml) cooled to -78°C and Yb(OTf)₃·xH₂O (0.186g) was added. The suspension was stirred for 5 mins at -78°C, then the vinyloxytrimethylsilane (0.29g) was added and the temperature was risen to 20°C. After 1 hr at that temperature a saturated solution of NH₄Cl (20 cc) was added followed by ethyl acetate (30ml). The organic phase was washed with brine (20 ml) and dried over sodium sulphate and concentrated to give the crude product, which was purified by column chromatography (cyclohexane, then cyclohexane/ethyl acetate 85/15) to give the title compound (0.562 g) as a colourless oil.

NMR (CDCl₃) δ (ppm) 9.65 (s, 1H), 7.00 (d, 1H), 6.70 (d, 1H), 5.60 (d, 1H), 4.80 (m, 1H) 4.10 (q, 2H), 3.10 (m, 2H), 1.15 (t, 3H).

15

Intermediate 3

Tributyl (2-oxo-1-phenylpyrrolidin-1-yl) phosphonium bromide

N,N,N'¹N'-Tetramethylethylene diamine (23.3ml) was added to a solution of N-phenylpyrrolidinone (5g) in dichloromethane (50ml). The solution was cooled to 0-5° and trimethylsilyl triflate (8.4ml) was added over ca 20 mins maintaining the temperature in the range 0-5°. The resultant solution was stirred for 10 mins and a solution of pyridinium bromide perbromide (13g) in acetonitrile (20ml) was added over ca 20 mins maintaining the temperature in the range 0-10°. The resultant suspension was stirred at 0-5° for ca 60 mins. Aqueous sodium bicarbonate solution (50ml) was added, cautiously. The mixture was stirred for ca 5 mins and the layers are separated. The aqueous phase was diluted with water (20ml) and back extracted with dichloromethane (20ml). The combined organic phases were washed with further sodium bicarbonate solution (50ml), 2M hydrochloric acid (2x50ml) and water (50ml), back extracting each wash with dichloromethane (10ml). The organic solution was dried (MgSO₄) and concentrated on a rotavapor. The red/brown solid was stirred with ethyl acetate (50ml) and warmed to give a solution which was then cooled and tributylphosphine (8.5ml) was

30

added. The solution was heated to reflux and maintained at reflux for 2.5 hours. The solution was allowed to cool to room temperature and was then cooled to 0-5°. The resulting suspension was aged at 0-5° for ca 60 min. The product was isolated by vacuum filtration and then washed with ethyl acetate:t-butylmethylether (1:1, 40ml) and dried in a vacuum oven at 45° to give the title compound as a white crystalline solid (10.12g), mp 127-128°.

Intermediate 4

10 (±) E-Ethyl 2-(5-chloro-2-iodoanilino)-4-(2-oxo-1-phenyl-3-pyrrolidinylidene) butanoate (4a); (±)-Z-Ethyl 2-(5-chloro-2-iodoanilino)-4-(2-oxo-1-phenyl-3-pyrrolidinylidene) butanoate(4b)

To a solution of intermediate 2 (2.4g) in acetonitrile (100 ml) at r.t. intermediate 3(3.7 g) and DBU (13 ml) were added and stirring was continued overnight at -20°C. The crude solution was poured into 200 ml of ethyl acetate and washed with a saturated solution of NH₄Cl (2 x 150 ml), then with water and brine. The organic phase was dried and concentrated to give the crude product as a 4/1 mixture of 4a/4b compounds. Purification by column chromatography (cyclohexane/ethyl acetate 80/20) gave the title 4a (2.16 g) and the 4b (0.5g) compounds as colourless oils.

Intermediate 4a

25 NMR (CDCl₃) δ (ppm) 7.72 (d, 2H), 7.56 (d, 1H), 7.38 (t, 2H), 7.16 (t, 1H), 6.6 (m, 1H), 6.50 (dd, 1H), 6.49 (d, 1H), 4.88 (d, 1H), 4.26 (m, 3H), 3.87 (t, 2H), 2.79 (m, 4H), 1.30 (t, 3H)

Intermediate 4b

30 NMR (CDCl₃) δ (ppm) 7.69 (d, 2H), 7.52 (d, 1H), 7.38 (t, 2H), 7.17 (t, 1H), 6.47 (d, 1H), 6.44 (dd, 1H), 5.98 (m, 1H), 5.00 (d, 1H), 4.22 (m, 2H), 4.13 (m, 1H), 3.84 (t, 2H), 3.2-3.6 (m, 2H), 2.85 (m, 2H), 1.26 (t, 3H)

Intermediate 5

(1R)-2-(tert-butoxy)-1-methyl-2-oxoethyl -2-(5-chloro-2-iodoanilino)-4-pentenoate (5a) and (1R)-2-(tert-butoxy)-1-methyl-2-oxoethyl -2-(5-chloro-2-iodoanilino)-4-pentenoate (5b)

5 A solution of intermediate 1-tert-butyl-(R)-2(oxoacetoxy)-2-methyl acetate (4.1 g) in toluene (200 ml) was refluxed in a Dean-Stark apparatus for 2 hrs. After cooling to room temperature, 5-chloro-2-iodoaniline (4.3 g) was added, and the solution refluxed in the presence of MgSO₄ for 3 hrs. The clear solution was cooled, filtered through cotton to eliminate the MgSO₄, concentrated to dryness and re-dissolved in
10 dichloromethane (150 ml). The solution was cooled to -78°C, and TiCl₄ (1.9 ml) was added slowly from a syringe. After 15 min, allyl tributyltin (7.9 ml) was added, and the resulting black suspension was stirred for 1 hr. It was then poured onto ethyl acetate (300 ml), and saturated NH₄Cl (150 ml) was added. The organic phase was separated, washed with
15 water and brine, dried and concentrated. Final purification by column chromatography (cyclohexane/ethyl acetate 95/5) afforded the title compound (4.1g) (65/35 mixture of diastereomers) as a colourless oil (7.01 g)

20 NMR (CDCl₃) δ (ppm) 7.54 (1H), 6.46 (dd, 1H), 5.86 (m, 1H), 5.3-5.2 (m, 2H), 5.03 (m, 1H), 4.77 (bd, 1H), 4.16 (m, 1H), 2.8-2.68 (m, 2H), 1.50 (d, 3H), 1.45 (s, 9H)

Intermediate 5a

(1R)-2-(tert-butoxy)-1-methyl-2-oxoethyl -2-(5-chloro-2-iodoanilino)-4-pentenoate

25 To a solution of allyltributyl tin (3.3g) in dry DCM (100ml) a 1M solution in DCM of SnCl₄ (10ml) was added at -78C. The mixture was stirred for 20 min, then intermediate 2-[2-(5-Chloro-2-iodo-phenylimino)-acetoxy]-1-(R)-methyl-acetic acid tertbutyl ester (2.39g) in dry DCM (50ml) was added. The reaction was allowed to react at -78C for 20min, then a
30 saturated solution of NH₄Cl was added and the resulting mixture was extracted with ethyl acetate (2x200ml). The organic layer was washed with a solution of KF 10% in water, then diethyl ether was added and the resulting solid was filtered.

The solution was dried and evaporated under vacuum. Final purification by flash chromatography (CH/EA 95:5) give the title compound as pure diastereomer as a colourless oil (1.3g).

5 NMR (CDCl₃) : 7.55 (d, 1H); 6.47 (d, 1H); 6.43 (d, 1H); 5.88 (m, 1H); 5.27 (m, 2H); 5.05 (q, 1H); 4.78 (d, 1H); 4.18 (m, 1H); 2.74 (m, 2H); 1.52 (d, 3H); 1.67 (s, 9H).

IR (CDCl₃): 3379, 1740

Intermediate 6

10 (1R)-2-(tert-butoxy)-1-methyl-2-oxoethyl-2-(5-chloro-2-iodoanilino)-4-oxobutanoate (6a) and (1R)-2-(tert-butoxy)-1-methyl-2-oxoethyl-2-(5-chloro-2-iodoanilino)-4-oxobutanoate (6b)

15 A solution of intermediate 5 (7.1 g) in dichloromethane (200 ml) was cooled to -78°C and ozone was bubbled through it until the solution turned red. Triphenylphosphine (8 g) was then added, and the reaction allowed to stir for 2 hrs, without control of the temperature. The crude mixture was evaporated to dryness and purified repeatedly by column chromatography ((cyclohexane/ethyl acetate 85/15) to afford the title compound 6a (2.75 g) and 6b(0.87g) as colourless oils.

20 compound 6a:

NMR (CDCl₃) δ (ppm) 9.85 (t, 1H), 7.57 (d, 1H), 6.58 (d, 1H), 6.51 (dd, 1H), 5.04 (q, 1H), 4.96 (d, 1H), 4.62 (m, 1H), 3.13 (dd, 2H), 1.55-1.42 (m, 12 H)

IR (CDCl₃) (cm⁻¹) 1740

25 compound 6b:

NMR (CDCl₃) δ (ppm) 9.81 (t, 1H), 7.57 (d, 1H), 6.60 (d,1H), 6.52 (dd, 1H), 5.02 (q, 1H), 4.95 (d, 1H), 4.55 (m, 1H), 3.11 (m, 2H), 1.55-1.43 (m, 12H).

IR (CDCl₃) (cm⁻¹) 1740

30 Intermediate 6a

(1R)-2-(tert-butoxy)-1-methyl-2-oxoethyl-2-(5-chloro-2-iodoanilino)-4-oxobutanoate

The title compound was obtained starting from intermediate 5a following the same procedure described for intermediate 6.

Intermediate 7

5 (E)-(1R)-2-(tert-butoxy)-1-methyl-2-oxoethyl 2-(5-chloro-2-iodoanilino)-4-(2-oxo-1-phenyl -3-pyrrolidinylidene)butanoate
(diastereoisomer A)

To a solution of intermediate 6a (2.7 g) in acetonitrile (60 ml) 2b(3 g) and DBU (1 ml) were added and the mixture was reacted at -20°C overnight.
10 It was then taken up with ethyl acetate (300 ml) and washed with 1N HCl, water and brine, dried and concentrated. Final purification by column chromatography (cyclohexane/ethyl acetate 85/15) afforded the title compound (2.1g) as a white solid .

m.p. 36-39°, $[\alpha]_D^{22}$ 22° (c=0.160% w/v in DMSO)

15 NMR (CDCl₃) δ (ppm) 7.72 (d, 2H), 7.55 (d, 1H), 7.38 (t, 2H), 7.15 (t, 1H), 6.66 (m, 1H), 6.49 (dd, 1H), 6.48 (d, 1H), 5.05 (m, 1H), 4.81 (d, 1H), 4.30 (m, 1H), 3.87 (t, 2H), 3.0 (m, 2H), 2.75 (m, 2H), 1.51 (d, 3H), 1.45 (s, 9H).

20 Intermediate 8

(-)-(1R)-2-(tert-butoxy)-1-methyl-2-oxoethyl 7-chloro-4-(1-phenyl-Δ³-pyrrolin-2-one-3yl)-1,2,3,4-tetrahydroquinoline-2-carboxylate (8a)
(-)-7-chloro-4-(2-oxo-1-phenyl-3-pyrrolidinylidene)-1,2,3,4-tetrahydro-2-quinoline-2-carboxylic acid, [1-(R)-(1-tert-butoxycarbonyl)]ethyl ester (8b)
25

To a solution of intermediate 7 (2.1 g) in DMF (40 ml) Pd(PPh₃)₄ (0.393 g) and triethylamine (0.95 ml) were added and the mixture was heated to 150°C for 1 hr. The crude solution was taken up with ethyl acetate and washed with 1N HCl, water and brine, dried and evaporated.
30 purification by column chromatography (cyclohexane/dichloromethane/ethyl acetate 50/40/10) afforded the title compound 8a (0.7 g) as a white solid.

m.p.=69-73°C

$[\alpha]_D$ -70.1° (c=0.190% w/v in DMSO)

NMR (DMSO) δ (ppm) 7.80 (m, 2H), 7.39 (m, 2H), 7.12 (m, 1H), 6.82 (d, 1H), 6.77 (d, 1H), 6.70 (m, 1H), 6.49 (dd, 1H), 6.46 (bs, 1H), 4.93 (q, 1H), 4.49 (m, 2H), 4.02 (m, 1H), 3.87 (m, 1H), 2.44 (m, 1H), 2.00 (m, 1H), 1.39 (s, 9H), 1.38 (d, 3H).

IR (Nujol) (cm⁻¹) 3380, 1741, 1681, 1601

and the title compound 8b (0.8 g) as a yellow solid.

m.p.=59-64°C

$[\alpha]_D$ -76.2° (c=0.510% w/v in DMSO)

NMR (DMSO) δ (ppm) 7.73 (m, 2H), 7.36 (m, 2H), 7.21 (d, 1H), 7.11 (m, 1H), 6.98 (da, 1H), 6.75 (d, 1H), 6.57 (dd, 1H), 4.70 (q, 1H), 4.24 (m, 2H), 3.84 (m, 1H), 3.75 (m, 1H), 3.18 (m, 1H), 3.05 (m, 1H), 2.94 (m, 1H), 1.25 (s, 9H), 1.23 (d, 3H)

15 Intermediate 9

(±)-Ethyl 2-(5-chloro-2-iodoanilino)-4-(2-oxo-1-phenyl piperidinylidene) butanoate.

To a solution of tributyl-3-(1-phenyl-2-piperidinone) phosphonium bromide (0.83g) in acetonitrile (20 ml) DBU (0.27ml) was added and after 15 min a solution of the intermediate 2 (0.35g) in acetonitrile (20 ml). The reaction mixture was stirred for 30 min, then diluted with ethyl acetate and washed with a 1N solution of HCl and brine. The organic phase was dried and concentrated to give the crude product which was purified by flash column chromatography to obtain the title compound (0.29g) as pale yellow foam.

NMR (CDCl₃) δ (ppm) 7.56 (dd, 1H), 7.38 (dd, 2H), 7.27 (dd, 2H), 7.24 (t, 1H), 6.93 (t, 1H), 6.50-6.47 (m, 2H), 4.85 (d, 1H), 4.25 (q, 2H), 4.22 (m, 1H), 3.71 (m, 2H), 2.76, (m, 2H), 2.59 (m, 2H), 2.01 (m, 2H), 1.29 (t, 3H)

Intermediate 10

30 (±)-Ethyl 2-(5-chloro-2-iodoanilino)-4-(2-oxo-1-(pyridin-3-yl)-pyrrolidin-3-ylidene)butanoate.

To a solution of the (1-(pyridin-3-yl)-2-oxo-pyrrolidin-3-yl) tributylphosphonium bromide (0.93g) in acetonitrile (10 ml) DBU (0.22ml)

was added and after 10 min a solution of the intermediate 2 (0.46g) in acetonitrile (10 ml). The reaction mixture was stirred for 3 hr, then diluted with ethyl acetate and washed with a saturated solution of NH_4Cl and brine. The organic phase was dried and concentrated to give the crude product which was purified by flash column chromatography to obtain the

5 title compound (0.47g) as a mixture of E/Z isomer (80/20)
MS (m/z) 526

Intermediate 11

10 (±)-E-Ethyl 2-(3,5-dichloro-2-iodoanilino)-4-(2-oxo-1-phenyl-3-pyrrolidinylidene) butanoate (11a);(±)-Z-Ethyl 2-(3,5-dichloro-2-iodoanilino)-4-(2-oxo-1-phenyl-3-pyrrolidinylidene) butanoate (11b)

To a solution of intermediate 2a in acetonitrile (10ml) at r.t 2b (0.726 g) and DBU (0.33 ml) were added and stirring was continued overnight at -20°C. The crude solution was poured into 20 ml of ethyl acetate and washed first with a saturated solution of NH_4Cl (2 x 15 ml), then with water and brine. The organic phase was dried and concentrated to give the crude product as a 4/1 mixture of Z/E isomers. Purification by column chromatography (cyclohexane/ethyl acetate 85/15) gave the

20 title compound 11a (0.498 g) and the title compound 11b (0.122g) as colourless oils.

intermediate 11a

NMR (CDCl_3) δ (ppm) 7.78 (d, 2H), 7.39 (t, 2H), 7.16 (t, 1H), 6.90 (d, 1H), 6.58 (m, 1H), 6.36 (d, 1H), 5.22 (d, 1H), 4.26 (m, 3H), 3.87 (t, 2H), 2.79 (m, 4H), 1.30 (t, 3H)

25 IR (CDCl_3) (cm^{-1}) 3370, 1738, 1697, 1671.

MS (m/z) 559.

intermediate 11b

30 NMR (CDCl_3) δ (ppm) 7.69 (d, 2H), 7.38 (t, 2H), 7.17 (t, 1H), 6.84 (d, 1H), 6.34 (d, 1H), 5.96 (m, 1H), 5.34 (d, 1H), 4.22 (m, 2H), 4.12 (m, 1H), 3.84 (t, 2H), 3.63-3.27 (m, 2H), 2.85 (t, 2H), 1.26 (t, 3H)

IR (CDCl_3) (cm^{-1}) 1733, 1685.

MS (m/z) 559.

Intermediate 12

-(1R)-2-(tert-butoxy)-1-methyl-2-oxoethyl 2-(5-chloro-2-iodoanilino)-4-(2-oxo-1-phenyl -3-pyrrolidinylidene)butanoate (diastereoisomer B)

5 To a solution of intermediate 6b (0.87 g) in acetonitrile (20 ml) tributyl-3-(N-phenyl-1-pyrrolidonyl)phosphonium bromide (1.6 g) and DBU (0.33 ml) were added and the mixture was reacted at -20°C overnight. It was then taken up with ethyl acetate (100 ml) and washed with 1N HCl, water
10 and brine, dried and concentrated. Final purification by column chromatography (cyclohexane/ethyl acetate 85/15) afforded the title compound (0.47 g) as a white solid oil.

m.p.=38-42°C

15 NMR (CDCl₃) δ (ppm) 7.72 (d, 2H), 7.55 (d, 1H), 7.38 (t, 2H), 7.16 (t, 1H), 6.60 (m, 1H), 6.56 (d, 1H), 6.49 (dd, 1H), 5.03 (q, 1H), 4.80 (d, 1H), 4.33 (m, 1H), 3.88 (t, 2H), 2.9 (m, 2H), 2.75 (m, 2H), 1.48 (d, 3H), 1.44 (s, 9H).

IR (CDCl₃) (cm⁻¹) 3375, 1738, 1693, 1665

20 **Intermediate 13**

-(1R)-2-(tert-butoxy)-1-methyl-2-oxoethyl 7-chloro-4-(1-phenyl-Δ³-pyrrolin-2-one-3yl)-1,2,3,4-tetrahydroquinoline-2-carboxylate (diastereoisomer B)

25 To a solution of intermediate 12 (0.46 g) in DMF (8 ml) Pd(PPh₃)₄ (0.043 g) and triethylamine (0.21 ml) were added and the mixture was heated to 150°C for 1 hr. The crude solution was taken up with ethyl acetate and washed with 1N HCl, water and brine, dried and evaporated. Final purification by column chromatography (cyclohexane/dichloromethane/ethyl acetate 50/40/10) afforded the title
30 compound (0.114 g) as a white solid.

m.p.=62-67°C

NMR (DMSO) δ (ppm) 7.79 (m, 2H), 7.38 (m, 2H), 7.11 (t, 1H), 6.81 (d, 1H), 6.77 (d, 1H), 6.70 (d, 1H), 6.55 (bs, 1H), 6.48 (dd, 1H), 4.90 (q, 1H), 4.5 (m, 2H), 3.99 (m, 1H), 3.84 (t, 1H), 2.35 (m, 1H), 2.02 (m, 1H), 1.39 (s, 12H).

5 **Intermediate 14**

2,4-dibromo-N-(4-(tert-butoxycarbonylamino)phenyl)-butyramide

To the derivative 2,4- dibromobutyl bromide (3.1 g) in dry dichloromethane (60 ml) pyridine(3.2 ml) was added, the mixture was kept at 0°C under a nitrogen atmosphere for 10 minutes and then the N-
10 t-butoxy carbonyl-1,4 phenylene diamine (2.08 g) was dropped. After 1 hour the mixture was poured into a saturated solution of NH₄Cl (200 ml) extracted with EA (3x150ml) and the organic phase washed with brine (200 ml), dried and concentrated *in vacuum*, the crude was purified by flash chromatography (eluting with CH/EA 80: 20) to give of the title
15 compound as a yellow foam (3.5 g). T.l.c. CH/EA 8:2, R_f = 0.53.
1H-NMR: 7.89 (sa); 7.44 (d); 7.35 (d); 6.46 (sa); 4.66 (dd); 3.60 (m); 2.76(m); 2.55(m); 1.51(s).

Intermediate 15

20 **3-bromo-1-(4-(tert-butoxycarbonylamino)phenyl)-2-oxo-pyrrolidine**

To a solution of intermediate 14 (3.5 g) in dry THF (50ml) cooled (0°C), a solution of LHMDS (9.6 ml of a 1M solution in tetrahydrofuran) was added drop-wise. The mixture was stirred under nitrogen until the temperature reached r.t for 2 hours. Then it was quenched into a
25 saturated solution of NH₄Cl (200 ml) extracted with EA (3x150ml) and the organic extracts were washed with brine (200 ml), dried and concentrated *in vacuum*. The mixture was purified by flash chromatography (eluting with CH/EA 8:2) to give the title compound (2.6 g). T.l.c. CH/EA 8:2, R_f =0.31. 1H-NMR: 7.57 (d); 7.39 (d); 6.49 (sa);
30 4.59 (m); 4.03 (m); 3.81 (m); 2.73 (m); 2.46 (m); 1.53(s).

Intermediate 16

(+/-)-Z-Ethyl2-(5-chloro-2-iodoanilino)-4-(2-oxo-1-(4-tert-butoxycarbonylamino)phenyl-pyrrolidin-3-ylidene)butanoate

A solution of intermediate 15 (2.6 g) in dry DMF (100ml) and tributylphosphine was refluxed at 110 °C under a nitrogen atmosphere for 4h, until reaction completion (TLC). The mixture was concentrated *in vacuum* to give the crude 1-(4-tert-butoxycarbonylamino)phenyl-2-oxo-pyrrolidin-3-yl-tributylphosphonium bromide (1.75g) which was dissolved in dry CH₃CN (100ml) was cooled at -30 °C and stirred under a nitrogen atmosphere, then DBU (0.44 ml) and intermediate 2 (1.0 g) were added. The mixture was stirred for 1h then was poured into a saturated solution of NH₄Cl (200 ml) extracted with EA (3x150ml) and the organic extracts were washed with brine (200 ml), dried and concentrated *in vacuum* to give a yellow oil which was purified by flash chromatography (eluting with CH/EA 80:20) to give the title compound (0.085 g) as a white solid. T.l.c. CH-EA (7:3), R_f=0.23 IR: 1727 and 1695 (C=O) cm⁻¹. ¹H-NMR: 7.64 (d); 7.53 (d); 7.38 (d); 6.48 (d); 6.47 (sa); 6.45 (dd); 5.97(m); 5.02(d); 4.23 (m); 4.14 (m); 3.8(t); 3.6 (m); 3.3 (m); 2.85 (m); 1.53(s); 1.27(t).

Intermediate 17

(±)-Z-Benzyl 2-(5-chloro-2-iodoanilino)-4-(2,5-dioxo-1-phenyl-imidazolidin-4-ylidene)butanoate.

To a solution of the derivative N-(phenylaminocarbonyl)α-phosphonoglycine-trimethyl ester (0.1g) in acetonitrile (10 ml) DBU (0.1ml) was added and after 10 min a solution of the (+/-)- 2- (5- Chloro-2-iodo-phenylamino)-4-oxo-butyric acid benzyl ester (0.1g) in acetonitrile (2 ml). The reaction mixture was stirred for 1 ½ hr, then diluted with ethyl acetate and washed with a 1N solution of HCl and brine. The organic phase was dried and concentrated to give the crude product which was purified by flash column chromatography to obtain the title compound (0.065g)

NMR (DMSO) δ (ppm) 10.80 (s, 1H), 7.65 (d, 1H), 7.7-7.3 (m, 10H), 6.75 (d, 1H), 6.55 (dd, 1H), 5.70 (t, 1H), 5.20 (s, 2H), 5.07 (d, 1H), 4.72 (m, 1H), 2.86 (t, 2H)

IR (Nujol) (cm⁻¹) 3339, 3160, 1768, 1721, 1691

Intermediate 18**(±)-Benzyl 7-chloro-4-(2,5-dioxo-1-phenyl-imidazolidin-4-ylidene)-1,2,3,4-tetrahydro-2-quinolinecarboxylate.**

5 To a solution of intermediate 17 (0.065 g) in DMF (5 ml) Pd(PPh₃)₄ (16 mg) and TEA (0.05 ml) were added and the resulting solution was heated to 110°C for 1 h. The crude solution was poured into ethyl acetate and washed with a 1N solution of HCl and brine. The organic phase was dried and concentrated to give the crude product which was
10 purified by flash column chromatography to obtain the title compound (0.015g) as yellow powder.

m.p.>220 °C

NMR (DMSO) δ (ppm) 10.5 (s, 1H), 7.5-7.2 (m, 11H), 7.16 (bd, 1H), 6.75 (d, 1H), 6.58 (dd, 1H), 5.2-5.01 (dd, 2H), 4.40 (m, 1H), 4.25 (dd, 1H),
15 2.83 (dd, 1H).

IR (Nujol) (cm⁻¹) 3378, 1752, 1728, 1704

Intermediate 19**2-[2-(5-Chloro-2-iodo-phenylimino)-acetoxy]-1-(R)-methyl-acetic acid isobutyl ester**

To a solution of acrylic acid 1-isobutoxycarbonyl-1-(R)-methyl-methyl ester (3.7g) in THF/H₂O OsO₄ 4% in H₂O (4ml) was added. The black
25 suspension was then treated with NaIO₄ (10.5g) by portions.

After 5hrs, the solution was taken up with ethyl acetate (2x50ml) and washed with water (2x50ml). The organic phase evaporated under vacuum and the crude mixture was purified by flash chromatography (CH₂Cl₂/EA 1:1) to afford 2-(2-Oxo-acetoxy)-1-(R)-methyl -acetic acid
30 isobutyl ester as colourless oil (3g). 24.8g of 2-(2-Oxo-acetoxy)-1-(R)-methyl -acetic acid isobutyl ester was dissolved in toluene (1000 ml) and refluxed in a Dean-Stark apparatus for 2 hrs. After cooling to room temperature, 5-chloro-2-iodoaniline (22 g) was added, and the solution

refluxed in the presence of $MgSO_4$ for 3 hrs. The clear solution was cooled, filtered through cotton to eliminate the $MgSO_4$, concentrated to dryness to obtain the title compound (38 g) as a yellow oil.

NMR ($CDCl_3$) δ (ppm) 7.83 (1H, d), 7.79 (s 1H), 7.02(dd,1H),6.96 (d, 1H),
5 5.373 (q 1H), 4.00 (m, 2H), 2.00 (m, 1H), 1.67 (d, 3H), 0.96 (2d, 6H)
IR ($CDCl_3$): 1749, 1730

Intermediate 20

10 2-(5-Chloro-2-iodo-phenylamino)-4-oxo-butyric acid 1-isobutoxycarbonyl- 1(R)-methyl-methyl ester (20a and 20b)

A solution of intermediate 19 (38 g) in toluene (1 ml) was cooled to $-20^\circ C$ and $Yb(OTf)_3$ (16.5g) was added and , after a few minutes, vinyloxy trimethylsilane (12.5g) dissolved in toluene (50ml) was added drop-wise. The bath was removed and the reaction allowed to stir for 2 hrs. The
15 crude mixture was taken up with ethyl acetate (500ml) and the organic phase was washed with a saturated solution of ammonium chloride (300ml) and evaporated. Then, the mixture was purified by column chromatography (cyclohexane/ethyl acetate 85/15) to afford the title compounds 20a (14 g) and 20b (4g) as colourless oils.

20 Intermediate 20a

NMR ($CDCl_3$) δ (ppm) 9.85 (s, 1H), 7.57 (d, 1H), 6.58 (d, 1H), 6.51 (dd, 1H), 5.19 (m, 1H), 4.97 (d, 1H), 4.63 (m, 1H), 3.93 (m, 2H), 3.24-3.04 (m, 2 H), 1.94 (m, 1H), 1.53 (d, 3H), 0.93 (dt, 3H); 0.91 (d, 3H).
IR ($CDCl_3$) (cm^{-1}) 1742, 1740

25 Intermediate 20b

NMR ($CDCl_3$) δ (ppm) 9.81 (s, 1H), 7.57 (d, 1H), 6.60 (d,1H), 6.52 (dd, 1H), 5.17 (m, 1H), 4.95 (d, 1H), 4.57 (m, 1H), 3.92 (m, 2H), 3.11 (m, 2H); 1.92 (m, 1H); 1.50 (d, 3H); 0.90 (d, 6H) .
IR ($CDCl_3$) (cm^{-1}) 3375, 1734

30

Intermediate 21

(E)-2-(5-Chloro-2-iodo-phenylamino)-4-(2-oxo-1-phenyl-pyrrolidin-3-ylidene)-butyric acid 1-isobutoxycarbonyl-1-(R) methyl-methyl ester

To a solution of intermediate 3 (14.45g) in acetonitrile (200 ml) DBU (4.43 ml) was added at room temperature and the mixture was stirred for 10 min. The mixture was then cooled at -25°C and intermediate 31a (12.98g) in 60 ml of CH_3CN was added drop-wise in 15 min. Then the reaction was stirred at this temperature for 2h. Then the mixture was taken up with ethyl acetate (100ml) and the organic phase washed with a saturated solution of NH_4Cl (150ml), and HCl 2% (200ml) and brine (2x200ml). The solution was then dried and concentrated. Final purification by column chromatography (cyclohexane/ethyl acetate/ CH_2Cl_2 7/0.5/2.5) afforded the title compound (11.04) as a white foam.

NMR (CDCl_3) δ (ppm) 7.73 (m, 2H), 7.56 (d, 1H), 7.38 (t, 2H), 7.16 (m, 1H), 6.67 (m, 1H), 6.50(dd, 1H), 6.49 (s, 1H), 5.20 (q, 1H), 4.81 (d, 1H), 4.33 (m, 1H), 3.94 (d, 2H), 3.88 (t, 2H), 3.0-2.74 (m, 4H), 1.94 (m, 1H), 1.57 (d, 3H); 0.91 (d, 6H).
IR (CDCl_3); 1696, 1670 cm^{-1}

Intermediate 22

7-Chloro-4-(2-oxo-1-phenyl-pyrrolidin-3-ylidene)-1,2,3,4-tetrahydro-quinoline-2-carboxylic acid, [1-(R)-methyl-1-isobutoxycarbonyl]-methyl ester (diastereoisomer A)

To a solution of intermediate 21 (9.55g) in toluene (130 ml), $\text{Pd}(\text{PPh}_3)_4$ (3.52 g) and triethylamine (5.1 ml) were added in portions and the mixture was heated to 110°C for 3.5 hr. The crude solution was taken up with ethyl acetate (600ml) and washed with NH_4Cl and brine, dried and evaporated. Purification by column chromatography (cyclohexane/dichloromethane/ethyl acetate 6.5/3/0.5) afforded the title compound (6.08 g) as a yellow foam.

NMR (DMSO) δ (ppm) 7.71 (d, 2H), 7.35 (t, 2H), 7.20 (d, 1H), 7.11 (t, 1H), 7.00 (s, 1H), 6.74 (d, 1H), 6.57 (dd, 1H), 4.89 (q, 1H), 4.24 (m, 2H), 3.84-3.60 (m, 4H), 3.2-2.8 (m, 3H), 1.70 (m, 1H), 1.24 (d, 3H); 0.73 (d, 6H).
IR (nujol): 3377, 1746, 1670

Intermediate 23**5** 7-Chloro-4-(2-oxo-1-phenyl-2,5-dihydro-1H-pyrrol-3-yl)-1,2,3,4-tetrahydro-quinoline-2-carboxylic acid, [1-(R)-methyl-1-isobutoxycarbonyl] -methyl ester

To a solution of intermediate 22 (3.67g) in DMF (50ml) Pd(PPh₃)₄ (0.340 g) and triethylamine (2 ml) were added and the mixture was heated to 110°C for 2 hrs. The crude solution was taken up with ethyl acetate (2x200ml) and washed with NH₄Cl and brine, dried and evaporated.
10 Final purification by column chromatography (cyclohexane/dichloromethane/ethyl acetate 6.5/3/0.5) afforded the title compound (1.289 g) as a yellow foam.

15 NMR (DMSO) δ (ppm) 7.79 (d, 2H), 7.38(t, 2H), 7.11 (t, 1H), 6.79 (d, 1H), 6.57 (d, 1H), 6.74 (d, 1H); 6.47 (dd, 1H); 6.47 (m, 1H); 5.10 (q, 1H); 4.49 (m, 2H); 4.06 (m, 1H); 3.92-3.82 (m, 3H); 2.45 (m, 1H); 2.019 (m, 1H); 1.84 (m, 1H); 1.42 (d, 3H); 0.84 (d, 6H).

IR (nujol): 3375, 1749, 1683.

20 Intermediate 24**2-(3,5-Dichloro-2-iodo-phenylamino)-4-oxo-butyric acid 1-n-butoxycarbonyl- 1(S)-methyl-methyl ester (24a and 24b)**

To a solution of intermediate Acrylic acid 1-n-butoxycarbonyl-1-(S)-methyl -methyl ester(4.9g) in THF/H₂O (100ml, 2/1) OsO₄ 4% in H₂O
25 (2.8g) was added. The black suspension was then treated with NaIO₄ (13g) by portions. After 5hrs, the solution was taken up with ethyl acetate (2x50ml) and washed with water (2x50ml). The organic phase evaporated under vacuum and the crude mixture was purified by flash chromatography (CH/EA 1:1) to afford the 2-(2-oxo-acetoxy)-1-(S)-
30 methyl -acetic acid n-butyl ester_ as a colourless oil (4.85g). (2.5g) of which was dissolved in toluene (200 ml) and refluxed in a Dean-Stark apparatus for 2 hrs. After cooling to room temperature, 3,5-dichloro-2-iodoaniline (2.46 g) was added, and the solution refluxed in the presence of MgSO₄ for 3 hrs. The clear solution was cooled, filtered

through cotton to eliminate the MgSO_4 , concentrated to dryness to obtain (2-[2-(5-chloro-2-iodo-phenylimino)acetoxyl]-1- (S)-methyl-acetic acid n butyl ester(4 g,) as a yellow oil.

5 A solution of such a yellow oil in CH_3CN (70ml) was cooled to -30°C and $\text{Yb}(\text{OTf})_3$ (2.1g) was added and, after a few minutes, vinyloxy trimethylsilane (1.1g) dissolved in CH_3CN (20ml) was added, drop-wise. The reaction was stirred for 10min. The crude mixture was taken up with ethyl acetate (500ml) and the organic phase was washed with a saturated solution of ammonium chloride (2x50ml) and evaporated.
10 Then, the mixture was purified by column chromatography (cyclohexane/ethyl acetate 90/10) to afford the title compounds 24a (1.4 g) and 24b (0.7g) as colourless oils.

Intermediate 24a: .

15 NMR (CDCl_3) δ (ppm) 9.84 (t, 1H), 6.92 (d, 1H); 6.45 (d, 1H); 5.33 (da, 1H); 5.17 (q, 1H); 4.60 (m, 1H); 4.14 (m, 2H); 3.34-3.06 (m, 2H); 1.6 (m, 2H); 1.52 (d, 3H); 1.37 (m, 2H); 0.93 (t, 3H).

IR (CDCl_3) (cm^{-1}) 3370,1742

Intermediate 24b:

20 NMR (CDCl_3) δ (ppm) 9.80 (s, 1H), 6.92 (d, 1H); 6.47 (d, 1H); 5.3 (da, 1H); 5.15 (q, 1H); 4.55 (m, 1H); 4.14 (m, 2H); 3.13 (m, 2H); 1.57 (m, 2H); 1.49 (d, 3H); 1.34 (m, 2H); 0.91 (t, 3H).

IR (CDCl_3) (cm^{-1}) 3370, 1744.

Intermediate 25

25 (E)-2-(3,5-Dichloro-2-iodo-phenylamino)-4-(2-oxo-1-phenyl-pyrrolidin-3-ylidene)-butyric acid 1-n-butoxycarbonyl-1-(S) methyl-methyl ester (diastereoisomerA)

To a solution of intermediate 2a (0.893) in acetonitrile (20 ml) DBU (0.25 ml) was added at room temperature and the mixture was stirred for 10
30 min. The mixture was then cooled at -25°C and intermediate 6b (0.8g) in 10 ml of CH_3CN was added drop-wise in 15 min. Then the reaction was stirred at this temperature for 30 min. Then the mixture was taken up with ethyl acetate (50ml) and the organic phase washed with a saturated

solution of NH₄Cl (50ml), and HCl 2% (10ml) and brine (2x20ml). The solution was then dried and concentrated. Final purification by column chromatography (cyclohexane/ethyl acetate 8/2) afforded the title product (0.734g) as a white foam.

5 NMR (CDCl₃) δ (ppm) 7.72 (d, 2H), 7.39 (t, 2H), 7.17 (t, 1H); 6.92 (d, 1H); 6.60 (m, 1H); 6.43 (d, 1H); 5.16 (q, 1H); 5.14 (d, 1H); 4.34 (d, 1H); 4.15 (m, 2H); 3.89 (t, 2H); 2.75-2.4 (m, 4H); 1.60 (m, 2H); 1.53 (d, 3H); 1.34 (m, 2H); 0.91 (t, 3H).

IR (CDCl₃); 3377, 1744, 1697, 1672 cm⁻¹

10

Intermediate 26

5,7-Dichloro-4-(2-oxo-1-phenyl-2,5-dihydro-1H-pyrrol-3-yl)-1,2,3,4-tetrahydro-quinoline-2-carboxylic acid , [1-(S)-methyl-1-n-butoxycarbonyl]-methyl ester (26a)
15 5,7-Dichloro-4-(2-oxo-1-phenyl-pyrrolidin-3-ylidene)-1,2,3,4-tetrahydro-quinoline-2-carboxylic acid , [1-(S)-methyl-1-n-butoxycarbonyl]-methyl ester (26b)

To a solution of intermediate 25 (0.734g) in DMF (20ml) Pd(OAc)₂ (0.110 g) and triethylamine (0.37 ml) were added in portions, and the mixture was heated to 120°C for 3hr. The crude solution was taken up with ethyl acetate (1000ml) and washed with NH₄Cl and brine, dried and evaporated. Final purification by column chromatography (cyclohexane/dichloromethane/ethyl acetate 7/2.5/0.5) afforded the title compound 26 a (0.35 g) and 26b(0.06g) as a yellow foam.

25

Intermediate 26a

NMR (DMSO) δ (ppm) 7.80(d, 2H); 7.38 (t, 2H); 7.11 (t, 1H); 6.89 (d, 1H); 6.83 (s, 1H); 6.68 (d, 1H); 6.47 (d, 1H); 5.07 (q, 1H); 4.48 (m, 2H); 4.11 (m, 1H); 4.06 (t, 2H); 3.8 (dd, 1H); 2.3-1.8 (m, 2H); 1.52 (m, 2H); 1.40 (d, 3H); 1.54 (m, 2H); 1.3 (m, 2H); 0.84 (t, 3H).

30

IR (nujol): 3374, 1740, 1683 cm⁻¹

Intermediate 26b

NMR (DMSO) δ (ppm) 7.69 (d, 2H); 7.39 (t, 2H); 7.33 (d, 1H); 7.15 (t, 1H); 6.71 (d, 1H); 6.62 (d, 1H); 4.72 (d, 1H); 4.40 (q, 1H); 4.40 (m, 1H);

3.94 (t, 2H); 3.76 (t, 1H); 3.60 (q, 1H); 3.12 (m, 1H); 2.35 (m, 1H); 2.21 (dd, 1H); 1.42 (m, 2H); 1.21 (m, 2H); 0.97 (d, 3H); 0.82 (t, 3H).

IR (nujol): 3377, 1746, 1684, 1594 cm^{-1}

5 **Example 1**

(±)-Sodium 7-chloro-4-(1-phenyl- Δ^3 -pyrrolin-2-one-3yl)-1,2,3,4-tetrahydroquinoline-2-carboxylate

To a solution of example 31a (540 mg) in IMS (5% methanol in absolute ethanol, 7 ml) NaOH (1N, 1.4 ml) was added and stirring continued for 2
10 hrs. The resulting solution was dried on the rotary evaporator and the resulting solid was triturated with diethyl ether. After filtration and drying the title compound (440 mg) was obtained as an off-white solid.

m.p. >200 °C

15 NMR (DMSO) δ (ppm) 7.80 (m, 2H), 7.39 (m, 2H), 7.11 (m, 1H), 6.80 (d, 1H), 6.72 (d, 1H), 6.36 (d, 1H), 6.34 (dd, 1H), 5.71 (s, 1H), 4.42 (m, 2H), 3.77 (m, 1H), 3.13 (m, 1H), 2.29 (m, 1H), 1.44 (m, 1H).

IR (Nujol) (cm^{-1}) 1672, 1600.

Example 2

20 **(-)-Sodium 7-chloro-4-(1-phenyl- Δ^3 -pyrrolin-2-one-3yl)-1,2,3,4-tetrahydroquinoline-2-carboxylate**

To a solution of intermediate 8a (690 mg) in THF/ H_2O (1/1) (14 ml) LiOH (65 mg) was added and stirring continued for 1 h. The resulting solution was concentrated to dryness, taken up with ethyl acetate and 1N HCl
25 was added. After vigorous stirring, the organic phase was separated, washed with water and brine and concentrated. The resulting solid was dissolved in THF (15 ml) and treated with sodium ethylhexanoate (232 mg) for 30 min. After drying, the resulting solid was triturated with hot diethyl ether and filtered, to afford the title compound (160 mg) as a
30 white solid.

e.e. = 99%

$[\alpha]_D = -102.3^\circ$ (c = 0.09% w/v in DMSO)

m.p. >200 °C

NMR (DMSO) δ (ppm) 7.80 (m, 2H), 7.39 (m, 2H), 7.11 (m, 1H), 6.80 (d, 1H), 6.72 (d, 1H), 6.36 (d, 1H), 6.34 (dd, 1H), 5.71 (s, 1H), 4.42 (m, 2H), 3.77 (m, 1H), 3.13 (m, 1H), 2.29 (m, 1H), 1.44 (m, 1H).

IR (Nujol) (cm^{-1}) 1672, 1600.

5

Example 3

(\pm)-Ethyl 7-chloro-4-(1-phenyl- Δ^3 -5,6-dihydro-pyridin-2-one-3yl)-1,2,3,4-tetrahydroquinoline-2-carboxylate (3a)

10 (\pm)-Ethyl 7-chloro-4-(2-oxo-1-phenyl-3-piperidinylidene)-1,2,3,4-tetrahydro-2-quinolinecarboxylate (3b)

To a solution of intermediate 9 (0.2 g) in DMF (5 ml) $\text{Pd}(\text{PPh}_3)_4$ (41 mg) and TEA (0.1 ml) were added and the resulting solution was heated to 110°C for 2 hrs. The crude solution was poured into ethyl acetate and washed with a 1N solution of HCl and brine. The organic phase was

15 dried and concentrated to give the crude product which was purified by flash column chromatography to obtain the title compound 3a (0.085g) as a white powder.

m.p.=131-133 °C

20 NMR (DMSO) δ (ppm) 7.4-7.3 (m, 4H), 7.20 (t, 1H), 6.78 (d, 1H), 6.75 (d, 1H), 6.48 (dd, 1H), 6.34 (bs, 1H), 5.99 (t, 1H), 4.13 (m, 2H), 3.97 (t, 1H), 3.93 (dd, 1H), 3.77 (m, 2H), 2.45 (m, 2H), 2.15 (m, 1H), 1.85 (m, 1H), 1.19 (t, 3H).

IR (Nujol) (cm^{-1}) 3392, 1723, 1659

and the title compound 3b (0.055g) as pale yellow powder.

25 m.p.=99-101 °C

NMR (DMSO) δ (ppm) 7.4-7.2 (m, 5H), 7.01 (d, 1H), 6.93 (d, 1H), 6.68 (d, 1H), 6.52 (dd, 1H), 4.20 (m, 1H), 4.16-3.96 (m, 2H), 3.74-3.60, 3.40 (m, 2H), 2.9-2.5 (m, 3H), 2.0-1.6 (m, 2H), 1.14 (t, 3H).

30 Example 4

(\pm)-Ethyl 7-chloro-4-(1-pyridin- Δ^3 -pyrrolin-2-one-3yl)-1,2,3,4-tetrahydroquinoline-2-carboxylate (4a)

(±)-Ethyl 7-chloro-4-(2-oxo-1-(pyridin3-yl)-pyrrolidin-3-ylidene)-1,2,3,4-tetrahydro-2-quinolinecarboxylate (4b)

To a solution of example 3 (0.47 g) in DMF (20 ml) Pd(PPh₃)₄ (100 mg) and TEA (0.38 ml) were added and the resulting solution was heated to 110°C for 1 ½ h. The crude solution was poured into ethyl acetate and washed with a saturated solution of NH₄Cl and brine. The organic phase was dried and concentrated to give the crude mixture which was dissolved in ethyl acetate (2ml) and treated with petroleum (2ml) the solid was filtered to give the title compound 4a (0.08 g) as a white powder.

m.p.=132-134 ° C

NMR (DMSO) δ (ppm) 8.99 (d, 1H), 8.32 (dd, 1H), 8.21 (m, 1H), 7.41 (dd, 1H), 6.80 (d, 1H), 6.77 (m, 1H), 6.75 (d, 1H), 6.47 (dd, 1H), 6.45 (m, 1H), 4.56 (m, 1H), 4.50 (m, 1H), 4.2-4.02 (m, 2H), 3.99 (m, 1H), 3.81 (t, 1H), 2.31 (m, 1H), 1.97 (m, 1H), 1.18 (t, 3H).

IR (Nujol) (cm⁻¹) 3391, 1728, 1679

The mother liquor was purified by flash chromatography to obtain a product which was triturated in cyclohexane to obtain title compound 4b (0.067 g, yellow powder).

NMR (DMSO) δ (ppm) 8.94 (d, 1H), 8.34 (dd, 1H), 8.14 (m, 1H), 7.41 (dd, 1H), 7.19 (d, 1H), 7.00 (d, 1H), 6.73 (d, 1H), 6.56 (dd, 1H), 4.27 (m, 1H), 4.20 (m, 1H), 4.00 (m, 1H), 3.89 (m, 1H), 3.85 (m, 1H), 3.72 (m, 1H), 3.21 (m, 1H), 2.93 (m, 1H), 2.84 (m, 1H), 0.90 (t, 3H).

IR (Nujol) (cm⁻¹) 3366, 1734, 1676 .

Example 5

(±)-Ethyl 5,7-dichloro-4-(1-phenyl-Δ³-pyrrolin-2-one-3yl)-1,2,3,4-tetrahydroquinoline-2-carboxylate.

To a solution of intermediate 11 a (0.430 g) in DMF (10 ml) Pd(OAc)₂ (11.6mg) and TEA (0.12 ml) were added and the resulting solution was heated to 130°C for 2 h. The crude solution was poured into 20 ml of ethyl acetate and washed first with a saturated solution of NH₄Cl (2 x 15 ml), then with water and brine. The organic phase was dried with Na₂SO₄ and concentrated to give the crude product. Purification by column

chromatography (cyclohexane/dichloromethane/ethyl acetate 60/30/10) gave the title compound (0.087g) as an off-white solid.

NMR (DMSO) δ (ppm) 7.81 (m, 2H), 7.40 (m, 2H), 7.13 (m, 1H), 6.91 (d, 1H), 6.75 (Sa, 1H), 6.68 (d, 1H), 6.45 (m, 1H), 4.46 (m, 2H), 4.17-4.10
5 (m, 3H), 3.79 (dd, 1H), 2.31 (m, 1H), 1.84 (m, 1H), 1.20 (t, 3H)
IR (Nujol) (cm^{-1}) 3390, 1724, 1678.

Example 6

(+/-)-Ethyl 7-chloro-4-(1-(4-tert-butoxycarbonylamino)-phenyl- Δ^3 -pyrrolin-2-one-3yl)-1,2,3,4-tetrahydroquinoline-2-carboxylate

A solution of intermediate 16 (0.085 g) in dry DMF (5ml) was stirred in the presence of TEA (0.018 ml) and Pd(OAc)₂ (0.0015 g) under a nitrogen atmosphere at 110°C for 1h. The mixture was diluted with a saturated aqueous ammonium chloride solution (100ml) and EA (100ml);
15 the organic layer was washed with brine (100ml), dried and concentrated *in vacuo*. The crude mixture was purified by flash chromatography (eluting with CH/EA 8:2) to give the title compound as a yellow solid (0.050 g).

T.l.c. CH-EA (8:2) R_f=0.30. ¹H-NMR:9.30 (sa); 7.64 (d); 7.43 (d); 6.80
20 (d); 6.75 (d); 6.63 (m); 6.46 (dd); 6.42(sa); 4.40(m); 4.13 (m); 3.92 (m); 3.78(m); 2.31 (m); 1.94 (m); 1.45 (s); 1.18(t).

Example 7

(+/-)-Ethyl 7-chloro-4-(1-(4-amino)-phenyl- Δ^3 -pyrrolin-2-one-3yl)-1,2,3,4-tetrahydroquinoline-2-carboxylate

To a solution of example 6 (0.070 g) in ethyl acetate (35 ml) HCl conc.(2.0 ml) was added.The mixture was stirred a r.t. under nitrogen atmosphere for 1h. The mixture was poured into a saturated aqueous solution of NaHCO₃ (100ml) and extracted with EA (200ml); the organic
30 layer was dried and concentrated *in vacuo*. The crude mixture was purified by flash chromatography (eluting with CH/ EA 1:1) to give the title compound as a yellow solid (0.043 g).

T.l.c. EA R_f = 0.289. IR:3388 (NH), 3161 (NH₂), 1718 and 1670 (C=O) cm⁻¹.

¹H-NMR: 7.36 (d); 6.80 (d); 6.75 (d); 6.56 (m); 6.47 (dd); 6.41(sa); 4.97(m); 4.32 (m); 4.14 (m); 3.91 (m); 3.77(m); 2.31 (m); 1.94 (m); 1.19(t).

5

Example 8

(+/-)-Ethyl 7-chloro-4-(1-(4-acetylamino)-phenyl- Δ^3 -pyrrolin-2-one-3yl)-1,2,3,4-tetrahydroquinoline-2-carboxylate

10 To a solution of example 7 (0.030 g) in pyridine dry (1 ml) Ac₂O (0.012 ml) was added. The mixture was stirred at r.t. under nitrogen atmosphere for 30 minutes. The mixture was poured into a saturated aqueous solution of NH₄Cl (50 ml) and extracted with EA (50 ml), the organic layer was dried and concentrated *in vacuo*. The crude mixture was triturated with EA to give the title compound as a white solid (0.025g)

15

T.l.c.CH/ EA (1:1) R_f =0.33. IR:3401(NH), 1730, 1675, 1651 (C=O) cm⁻¹.
d

¹H-NMR:9.9 (s); 7.69 (d); 7.56 (d); 6.80 (d); 6.75 (d); 6.65 (m); 6.47 (dd); 6.43 (sa); 4.5-4.37 (m); 4.13 (m); 3.93 (m); 3.79 (m); 2.3-1.94 (m); 2.03 (s); 1.19 (t).

20

Example 9

(+/-)-Ethyl 7-chloro-4-(1-(4-methanesulfonylamino)-phenyl- Δ^3 -pyrrolin-2-one-3yl)-1,2,3,4-tetrahydroquinoline-2-carboxylate

25 To a solution of example 7 (0.040 g) in CH₂Cl₂ dry (10 ml) DIPEA (0.021 ml) and CH₃SO₂Cl (0.008 ml) were added. The mixture was stirred at r.t. under nitrogen atmosphere for 1 h. The mixture was poured onto a saturated aqueous solution of NH₄Cl (50 ml) and extracted with EA (50 ml), the organic layer was dried and concentrated *in vacuo*. The crude mixture was purified by flash chromatography (eluting with CH/EA (1:1) to give the title compound as a yellow solid (0.027g) .

30

T.l.c.CH/ EA (1:1) R_f =0.63. IR:3394(NH), 1726, 1680, 1635 (C=O), (C=C) cm⁻¹.

¹H-NMR: 7.89 (d); 7.52 (d); 6.81 (d); 6.76 (d); 6.76 (s); 6.47 (dd); 6.45 (sa); 4.52 (m); 4.13 (m); 3.94 (m); 3.81(m); 3.51 (s); 2.3-1.97 (m); 1.19 (t)

Example 10

5 (+)-Ethyl 7-chloro-4-(2-oxo-1-((4-tert-butoxycarbonylamino) phenyl-pyrrolidin-3-ylidene)-1,2,3,4-tetrahydro-2-quinolinecarboxylate

A solution of intermediate 16 (1.02 g) in dry DMF (100 ml) was stirred in the presence of TEA (0.018 ml) and Pd(PPh₃)₄ (0.184 g) under a nitrogen atmosphere at 110°C for 2h until reaction completion (TLC).

10 The mixture was diluted with a saturated aqueous ammonium chloride solution (100ml) and EA (200ml); the organic layer was washed with brine (200ml), dried and concentrated *in vacuo*. The crude mixture was purified by flash chromatography (eluting with CH/DCM/EA 5:4:1) to give the title compound (280 mg).

15 IR: 3350 (NH), 1718 and 1670 (C=O) cm⁻¹. ¹H-NMR: 9.32 (sa); 7.59 (d); 7.43 (d); 7.17 (d); 6.94 (d); 6.72 (m); 6.55 (dd); 4.26(dd); 4.19(m); 4.04-3.88 (m); 3.8-3.6 (m); 3.18(m); 2.94-2.86 (m); 1.46 (s); 0.92 (t).

Example 11

20 (+)-Ethyl 7-chloro-4-(2-oxo-1-(4-amino)phenyl-pyrrolidin-3-ylidene)-1,2,3,4-tetrahydro-2-quinolinecarboxylate

To a solution of example 10 (0.280 g) in ethyl acetate (100 ml) HCl conc.(9.5 ml) was added. The mixture was stirred at r.t. under nitrogen atmosphere for 1h until reaction completion(Tlc). The mixture was
25 poured onto a saturated aqueous solution of NaHCO₃ (100ml) and extracted with EA (200ml); the organic layer was dried and concentrated *in vacuo*. The crude mixture was triturated with CH/ EA 1:1 to give the title compound as a yellow solid (0.191 g).

30 T.l.c. EA R_f = 0.33. IR: 3464-3406(NH), 3364(NH₂), 1730, 1658 and 1633 (C=O) cm⁻¹.

¹H-NMR: 7.31 (d); 7.16 (d); 6.91 (da); 6.71 (d); 6.55 (d); 6.54 (dd); 5.01(s); 4.26 (dd); 4.17 (m); 4.04-3.9 (m); 3.74-3.54(m); 3.14 (m); 2.87 (m); 0.96(t).

Example 12

(+)-Ethyl 7-chloro-4-(2-oxo-1-(4-acetylamino)phenyl-pyrrolidin-3-ylidene)-1,2,3,4-tetrahydro-2-quinolinecarboxylate

To a solution of intermediate 19 dry pyridine (1 ml) Ac₂O (0.010 ml) was added. The mixture was stirred a r.t. under nitrogen atmosphere for 30
5 minutes. The mixture was poured onto a saturated aqueous solution of NH₄Cl (50 ml) and extracted with EA (50 ml), the organic layer was dried and concentrated *in vacuum*. The crude mixture was triturated with EA to give the title compound as a yellow solid (0.027g).

T.l.c.CH/ EA (1:1) R_f=0.63 IR:3396-3325(NH), 1724-1685 (C=O) cm⁻¹.
10 ¹H-NMR:9.92 (s); 7.62 (d); 7.55 (d);7.16 (d); 6.95 (da); 6.71 (d); 655 (dd); 5.01(s); 4.25 (dd); 4.18 (m);4.1-3.85 (m); 3.77(m); 3.64 (m); 3.18 (m); 2.88 (m); 2.01 (s); 0.91 (t).

Example 13

15 **(+)-Ethyl 7-chloro-4-(2-oxo-1-((4-methanesulfonyl-amino)phenyl-pyrrolidin-3-ylidene)-1,2,3,4-tetrahydro-2-quinolinecarboxylate**

To a solution of example 12 (0.040 g) in dry CH₂Cl₂ (10 ml) DIPEA (0.021 ml) and CH₃SO₂Cl (0.008 ml) were added. The mixture was stirred a r.t. under nitrogen atmosphere for 1 h (Tlc). The mixture was
20 poured onto a saturated aqueous solution of NH₄Cl (50 ml) and extracted with EA (50 ml), the organic layer was dried and concentrated *in vacuum*. The crude mixture was crystallised with CH/EA (1:1) to give the title compound as a yellow solid (0.023g) . T.l.c.CH/ EA (1:1) R_f=0.63. IR:3384(NH), 1734, 1683 (C=O), 1600 (C=C) cm⁻¹.

25 ¹H-NMR:7.83 (d); 7.53 (d); 7.21 (d);7.00 (d); 6.75 (d); 6.57 (dd); 4.2-4.3 (m); 4.01 (m); 3.93 (m); 3.87(m); 3.73 (m);3.52(s); 3.22 (m); 3.0-2.9 (m); 0.95 (t).

Example 14

30 **(±)-Sodium 7-chloro-4-(1-(3-pyridin)--Δ³-pyrrolin-2-one-3yl)-1,2,3,4-tetrahydroquinoline-2-carboxylate.**

To a solution of example 4a (70mg) in IMS (5% of methanol in ethanol) (10ml) a 1N solution of NaOH (0.18ml) was added and reaction mixture was stirred for 1 ½ hr. The solvent was evaporated and the crude

product was first triturated in methanol/ethyl acetate 05ml/2ml then in isopropyl alcohol (3ml) to yield the title compound (40 mg) as a pale yellow solid.

m.p.>220°C

5 NMR (DMSO) δ (ppm) 8.98 (d, 1H), 8.31 (dd, 1H), 8.21 (m, 1H), 7.41 (m, 1H), 6.79 (d, 1H), 6.72 (d, 1H), 6.42 (d, 1H), 6.33 (dd, 1H), 5.71 (s, 1H), 4.50 (m, 1H), 4.44 (m, 1H), 3.76 (m, 1H), 3.11 (m, 1H), 2.27 (m, 1H), 1.43 (m, 1H).

IR (Nujol) (cm⁻¹) 3300, 1684.

10

Example 15

(±)-Sodium 7-chloro-4-(1-phenyl- Δ^3 -5,6-dihydro-pyridin-2-one-3yl)-1,2,3,4-tetrahydroquinoline-2-carboxylate.

To a solution of example 3a (80mg) in IMS (5% of methanol in ethanol) (6ml) a 0.1N solution of NaOH (2.9ml) was added and reaction mixture was stirred for 1 hr. The solution was poured into ethyl acetate and washed with a 1N solution of HCl and brine. The organic phase was dried with Na₂SO₄ and concentrated to give the crude acid compound. The latter was suspended in ethyl acetate (2ml) and sodium 2-ethylhexanoate (35mg) was added obtaining a solution. Diethyl ether (4ml) and petroleum (3ml) was added to precipitate the title compound (42 mg) as a white solid.

20

m.p.>163-166°C

25 NMR (DMSO) δ (ppm) 7.4-7.34 (m, 4H), 7.19 (m, 1H), 6.72 (d, 1H), 6.67 (d, 1H), 6.32 (d, 1H), 6.32 (dd, 1H), 5.71 (t, 1H), 5.64 (s, 1H), 3.96 (m, 1H), 3.8-3.65 (m, 2H), 3.17 (dd, 1H), 2.4 (m, 2H), 2.08 (1H), 1.3 (m, 1H)

IR (Nujol) (cm⁻¹) 3373, 1658, 1653

Example 16

30 (±)-Sodium 5,7-dichloro-4-(1-phenyl- Δ^3 -pyrrolin-2-one-3yl)-1,2,3,4-tetrahydroquinoline-2-carboxylate

To a solution of example 5 (87 mg) in IMS (5% methanol in absolute ethanol, 5 ml) NaOH (1N, 0.22 ml) was added and stirring continued for.

3 hrs. The resulting solution was dried on the rotary evaporator and the resulting solid was triturated with diethyl ether. After filtration and drying the title compound (78 mg) was obtained as an off-white solid.

5 NMR (DMSO) δ (ppm) 7.80 (m, 2H), 7.38 (t, 2H), 7.10 (t, 1H), 6.82 (d, 1H), 6.46 (d, 1H), 6.37 (s, 1H), 6.11 (s, 1H), 4.42 (s, 2H), 3.98 (d, 1H), 3.05 (dd, 1H), 2.24 (dd, 1H), 1.34 (m, 1H).

IR (Nujol) (cm^{-1}) 3385, 1663, 1591, 1555

Example 17

10 (+)**Sodium 7-chloro-4-(1-phenyl- Δ^3 -pyrrolin-2-one-3yl)-1,2,3,4-tetrahydroquinoline-2-carboxylate**

Method A

15 To a solution of intermediate 13 (110 mg) in THF/ H_2O (1/1) (3 ml) LiOH (11 mg) was added and stirring continued for 1 h. The resulting solution was concentrated to dryness, taken up with ethyl acetate and 1N HCl was added. After vigorous stirring, the organic phase was separated, washed with water and brine and concentrated. The resulting solid was dissolved in THF (15 ml) and treated with sodium ethylhexanoate (39 mg) for 30 min. After drying, the resulting solid was triturated with hot diethyl ether and filtered, to afford the title compound (69 mg) as a white solid.

e.r.=98%

$[\alpha]_{\text{D}}=92.5^\circ$ (c=0.420% w/v in DMSO)

m.p.>200 °C

25 NMR (DMSO) δ (ppm) 7.80 (m, 2H), 7.39 (m, 2H), 7.11 (m, 1H), 6.80 (d, 1H), 6.72 (d, 1H), 6.36 (d, 1H), 6.34 (dd, 1H), 5.71 (s, 1H), 4.42 (m, 2H), 3.77 (m, 1H), 3.13 (m, 1H), 2.29 (m, 1H), 1.44 (m, 1H).

IR (Nujol) (cm^{-1}) 1672, 1600.

Method B

30 Starting from Example 28 using the procedure as described for Example 21 (Method B).

Example 18

(+/-)-7-chloro-4-(1-(4-acetylamino)-phenyl- Δ^3 -pyrrolin-2-one-3yl)-1,2,3,4-tetrahydroquinoline-2-carboxylic acid

To a solution of example 8 (0.023 g) in IMS (5 ml) NaOH (0.150 ml) was added and the mixture was stirred at r.t. for 1 h.

- 5 The mixture was poured onto a solution of HCl 6 N (50 ml) and extracted with EA (50 ml), the organic layer was washed with brine (30 ml), dried and concentrated *in vacuum*. The crude mixture was triturated with Et₂O to give the title compound as a yellow solid (0.019g). T.l.c. EA R_f=0.2. IR:3401(NH, OH), 1734,1651 (C=O) cm⁻¹.
- 10 ¹H-NMR:12.84 (bs); 9.9 (s); 7.69(d); 7.56 (d); 6.80 (d); 6.76 (d);6.6 (d); 6.45 (dd);6.33 (sa); 4.42(m); 3.84-3.78(m); 3.70 (m); 2.3 (m); 2.017 (s); 1.9 (m).

Example 19

- 15 **(+/-)7-chloro-4-(1-(4-methanesulfonylamino)-phenyl- Δ^3 -pyrrolin-2-one-3yl)-1,2,3,4-tetrahydroquinoline-2-carboxylic acid**

To a solution of example 9 (0.027 g) in IMS (5 ml) was added NaOH (0.142 ml). The mixture was stirred a r.t. for 2 h.

- 20 The mixture was poured onto a solution of HCl 6N (50 ml) and extracted with EA (50 ml), the organic layer was washed with brine (30 ml), dried and concentrated *in vacuum*. The crude mixture was crystallized with CH/EA (1:1) to give the title compound as a yellow solid (0.015g) . T.l.c. EA R_f=0.2. IR:3446(NH,), 1732- (C=O), 1337-1154 (SO₂) cm⁻¹.
- 25 ¹H-NMR:13-12 (broad); 9.61 (s); 7.75 (d); 7.21 (d); 6.80 (d); 6.76 (d); 6.63 (dd); 6.46 (dd); 6.34 (dd); 4.43(m); 3.85-3.78 (m);2.93 (s); 2.3 (m); 1.92 (m).

Example 20

- 30 **(±)-Sodium 7-chloro-4-(2-oxo-1-phenyl-3-pyrrolidinyliidene)-1,2,3,4-tetrahydro-2-quinolinecarboxylate**

To a solution of example 31b (540 mg) in IMS (5% methanol in absolute ethanol, 7 ml) NaOH (1N,1.4 ml) was added and stirring continued for 2 hrs. The resulting suspension was filtered and the solid was washed with

small portions of diethyl ether. After drying, the title compound (450 mg) was obtained as a yellow solid.

m.p.>200 °C

5 NMR (DMSO) δ (ppm) 7.74 (d, 2H), 7.37 (t, 2H), 7.11 (t, 1H), 7.12 (d, 1H), 6.77 (d, 1H), 6.38 (dd, 1H), 6.13 (bs, 1H), 4.48 (dd, 1H), 3.78 (m, 2H), 3.2-3.4 (m, 2H), 2.90 (m, 1H), 1.98 (m, 1H)

Example 21

10 (-)-Sodium 7-chloro-4-(2-oxo-1-phenyl-3-pyrrolidinylidene)-1,2,3,4-tetrahydro-2-quinolinecarboxylate

Method A

To a solution of intermediate 8b (790 mg) in THF/H₂O (1/1) (16 ml) LiOH (73 mg) was added and stirring continued for 1 hr. The resulting solution was concentrated to dryness, taken up with ethyl acetate and 1N HCl was added. After vigorous stirring, the organic phase was separated, washed with water and brine and concentrated. The resulting solid was dissolved in THF (15 ml) and treated with sodium ethylhexanoate (265 mg) for 30 mins. After drying, the resulting solid was triturated with hot ethyl acetate and filtered to afford the title compound (400 mg) as a yellow solid.

ee=88.8%

$[\alpha]_D -603.7^\circ$ (c=0.316% w/v in DMSO)

m.p.>200 °C

25 NMR (DMSO) δ (ppm) 7.74 (d, 2H), 7.37 (t, 2H), 7.11 (t, 1H), 7.12 (d, 1H), 6.77 (d, 1H), 6.38 (dd, 1H), 6.13 (bs, 1H), 4.48 (dd, 1H), 3.78 (m, 2H), 3.2-3.4 (m, 2H), 2.90 (m, 1H), 1.98 (m, 1H)

IR (Nujol) (cm⁻¹) 3425, 1666, 1592

Method B

30 To a solution of example 27 (3.18g) in IMS (5% of methanol in ethanol) (100ml) a 1N solution of NaOH (8.64ml) was added: the sodium salt precipitates after 5 min. To the resulting suspension diethyl ether was added (50ml) and the solid was filtered.

The solution was evaporated and the solid obtained was mixed to the previous one and triturated with diethylether to afford the title sodium salt (3.2g) as yellow solid.

m.p.>220°C

5 NMR (DMSO) δ (ppm) 7.74 (d, 2H), 7.37 (t, 2H), 7.11 (t, 1H); 7.11 (d, 1H); 6.76 (d, 1H); 6.38 (dd, 1H); 6.11 (s, 1H); 4.48 (dd, 1H); 3.78 (m, 2H); 3.4-3.2 (m, 2H); 2.9 (m, 1H); 1.95 (m, 1H).

IR (Nujol) (cm⁻¹) 3392, 1669.

[α] -603.7 ° (c=0.316%w/v in DMSO)

10 e.e.: 96%

Method C

125g of *Aspergillus niger* lipase (Amano AP12) were suspended in 650 ml of 100mM calcium chloride solution in a stirred reactor. The suspension was cooled to 15° C. 50g of example 31b were then dissolved in dimethyl sulphoxide (350 ml) and this solution added to the reactor. The reactor was then heated to 37° C and the mixture stirred for 24 hours.

The reactor temperature was then lowered to 20° C and 1 litre of 0.2M hydrochloric acid was slowly added to the reactor. The reactor was then emptied and 50g of filter aid (Dicalite) were added to the reaction mixture. The mixture was then filtered and the filter cake washed with water, before being dried. A 20 g sample of dried filter cake was dispersed in 390 ml of methyl t-butyl ether and 10 ml of 2M hydrochloric acid were added. This was stirred for 3 hours and filtered, the filter cake was washed with 100ml of methyl t-butyl ether. The product was back extracted from the methyl t-butyl ether 500 into ml of 0.05M sodium hydroxide solution. The aqueous layer was then separated, acidified with 6ml of 5M hydrochloric acid and the product extracted into 500ml ethyl acetate. The ethyl acetate was removed by evaporation and the residue dissolved in IMS (80ml). The title compound was identified in this solution by HPLC assay as follows:

0.5 ml reaction mixture diluted into 2mls DMSO and mixed to dissolve. 5
ul of this further diluted into 1ml of mobile phase (70% acetonitrile in
20mM Ammonium acetate pH 3.0), Column: Spherisorb C6 50x4.6mm,
Flow rate: 1ml/min, Detection: uv adsorbance at 254 nm, Injection vol:
5 10ul, Retention time: 0.8 min.

The solution was diluted to 96 ml with IMS and stirred while 10ml of 1 M
sodium hydroxide were added drop-wise over 15 minutes. 40 ml of
diethyl ether were added over 10 minutes and stirring continued for 1
hour. The mixture was then placed in the fridge for 1 hour and the
10 product filtered, washed with 50ml of cold diethyl ether before being
dried overnight under vacuum to obtain the title compound (3.3 g).

HPLC analyses: the title compound was dissolved in DMSO at 1mg/ml.
10 ul of this diluted into 990ul of mobile phase.

Column: Phenomenex Luna Phenyl hexyl 150x4.6 mm, Injection vol: 50
15 ul, Retention time: 3.4 min.

Example 22

**(±)-Sodium 7-chloro-4-(2-oxo-1-(pyridin-3-yl)-pyrrolidin-3-ylidene)-
1,2,3,4-tetrahydro-2-quinolinecarboxylate.**

To a solution of example 4b (55mg) in IMS (5% of methanol in ethanol)
20 (10ml) a 1N solution of NaOH (0.145ml) was added and reaction mixture
was stirred for 1 ½ hr. The solvent was evaporated and the crude
product was triturated in ethyl acetate 2ml to yield the title compound (38
mg) as a yellow solid.

m.p. >220°C

25 NMR (DMSO) δ (ppm) 8.96 (d, 1H), 8.32 (dd, 1H), 8.18 (m, 1H), 7.40 (m,
1H), 7.12 (d, 1H), 6.78 (d, 1H), 6.38 (dd, 1H), 6.15 (s, 1H), 4.46 (m, 1H),
3.83 (m, 2H), 3.3-3.2 (m, 2H), 2.92 (m, 1H), 1.97 (m, 1H).

IR (Nujol) (cm⁻¹) 3361, 1669.

30 Example 23

**(±)-7-chloro-4-(2-oxo-1-phenyl-3-piperidinylidene)-1,2,3,4-tetrahydro-
2-quinolinecarboxylic acid.**

To a solution of example 3b (48mg) in IMS (5% of methanol in ethanol) (2ml) a 0.1N solution of NaOH (1.2ml) was added and reaction mixture was stirred for 2 ½ hr. The solution was poured into ethyl acetate and washed with a 1N solution of HCl and brine. The organic phase was dried and concentrated to give the crude product which was triturated in ethyl acetate/petroleum 2ml/5ml, to yield the title compound (14 mg) as a yellow solid.

5

m.p.>130-133°C

NMR (DMSO) δ (ppm) 12.64 (s, 1H), 7.38 (t, 2H), 7.30 (d, 2H), 7.22 (t, 1H), 6.99 (d, 1H), 6.87 (bd, 1H), 6.67 (d, 1H), 6.50 (dd, 1H), 4.08 (m, 1H), 3.54 (m, 2H), 3.43 (m, 1H), 2.83 (m, 1H), 2.72 (m, 1H), 2.58 (1H), 1.93-1.8 (m, 2H)

10

IR (Nujol) (cm⁻¹) 3348, 1732, 1717

MS (*m/z*) 383

15

Example 24

(±)-7-chloro-4-(2,5-dioxo-1-phenyl-imidazolidin-4-ylidene)-1,2,3,4-tetrahydro-2-quinolinecarboxylic acid.

To a solution of intermediate 18 (10mg) in CH₂Cl₂ (5 ml) a 1M solution of BCl₃ in hexane (0.1 ml) was added at -78°C and reaction mixture was stirred for 1 ½ hr maintaining the temperature between -20 and -10°C. The solution was poured into ethyl acetate and washed with a 3N solution of HCl and brine. The organic phase was dried with Na₂SO₄ and concentrated to give the crude product which was triturated in diethyl ether/petroleum (1ml/3ml), to yield the title compound (6 mg) as a yellow solid.

20

25

m.p.>190°C deg.

NMR (DMSO) δ (ppm) 12.75 (bs, 1H), 10.50 (bs, 1H), 7.50-7.39 (m, 6H), 6.99 (bs, 1H), 6.76 (d, 1H), 6.57 (dd, 1H), 4.15 (m, 1H), 3.77.(m, 1H), 3.17 (dd, 1H),.

30

IR (Nujol) (cm⁻¹) 3400, 2800, 1746, 1701

Example 25

(+/-)-7-chloro-4-(2-oxo-1-(4-acetylamino)phenyl-pyrrolidin-3-ylidene)-1,2,3,4-tetrahydro-2-quinolinecarboxylic acid

To a solution of example 12 (0.027 g) in THF H₂O (3:1) (10 ml) was added LiOH (0.010 g). The mixture was stirred a r.t. for 1 h. The mixture
5 was poured onto a saturated aqueous solution of NH₄Cl (50 ml) and extracted with EA (50 ml), the organic layer was washed with brine (30 ml), dried and concentrated *in vacuum*. The crude mixture was triturated with EA to give the title compound as a yellow solid (0.020g) . T.l.c.CH/ EA (1:1) R_f=0.2. IR:3400-2700(NH, OH), 1660 (C=O) cm⁻¹.
10 ¹H-NMR:12.63 (sa); 9.94 (sa); 7.65(d); 7.58 (d); 7.20 (d); 6.83 (sa); 6.74 (d); 6.54 (dd); 4.03(m); 3.78(m); 3.70 (m); 3.2-2.6 (m); 2.03 (s).

Example 26

(+/-)-7-chloro-4-(2-oxo-1-((4-methanesulfonyl amino)phenyl-pyrrolidin-3-ylidene)-1,2,3,4-tetrahydro-2-quinolinecarboxylic acid

To a solution of example 13 (0.023 g) in IMS (5 ml) NaOH (0.120 ml) was added. The mixture was stirred a r.t. for 2 h. The mixture was poured onto a solution of HCl 6N (50 ml) and extracted with EA (50 ml), the organic layer was washed with brine (30 ml), dried and concentrated
20 *in vacuum*. The crude mixture was chromatographed with Et₂O to give the title compound as a yellow solid (0.007g). T.l.c.CH/ EA (1:1) R_f=0.2. IR:3411(NH,), 1692,1651-1583 (C=O), (C=C), 1306-1154 (SO₂) cm⁻¹.
¹H-NMR: 9.65 (s); 7.69(d); 7.22 (d); 7.20 (d); 6.73 (d); 6.55 (dd); 4.03(m); 3.8-3.5 (m); 3.3-2.9 (m); 2.9 (s).

Example 27

7-chloro-4-(2-oxo-1-phenyl-3-pyrrolidinylidene)-1,2,3,4-tetrahydro-2-quinolinecarboxylic acid (enantiomer A)

To a solution of intermediate 22 (6.2g) in THF/H₂O (100ml, 3/1) at room temperature LiOH (1g) was added and stirring continued for 1 hr. The
30 THF was evaporated and H₂O (100ml) was added. The resulting solution was washed with diethylether (2x50ml). The aqueous phase was acidified until pH=4 with HCl 10% and the product extracted with ethyl acetate (2x100ml) The organic phase was washed with water and

brine, dried and evaporated to afford the title compound (4.2g) as a yellow solid. m.p.>200 °C

NMR (DMSO) δ (ppm) 12.62 (bs, 1H); 7.72 (d, 2H), 7.38 (t, 2H), 7.20 (d, 1H), 7.13 (t, 1H), 6.86 (d, 1H), 6.74 (d, 1H), 6.54 (dd, 1H), 4.06 (m, 1H),
5 3.86-3.68 (m, 3H), 3.3 (m, 1H), 3.18-2.88 (m, 2H).

IR (nujol): 3356, 1724

Example 28

10 7-Chloro-4-(2-oxo-1-phenyl- Δ^3 -pyrrolin-2-one-3-yl)-1,2,3,4-tetrahydro-quinoline-2-carboxylic acid; (enantiomer A)

To a solution of intermediate 10 (1.289g) in THF/H₂O (30ml, 3/1) at room temperature LiOH (0.24g) was added and stirring continued for 1 hr. The THF was evaporated and H₂O (80ml) was added. The resulting solution was washed with diethylether (2x50ml). The aqueous phase was
15 acidified until pH=4 with HCl 10% and the product was filtered and washed with water (10ml). The product was dried under vacuum at 60C for 12hrs to obtain 0.734g as white solid.

m.p.: 190°C

e.e.: 100%

20 NMR (DMSO) δ (ppm) 12.86 (bs, 1H); 7.79 (d, 2H), 7.38 (t, 2H), 7.11 (d, 1H), 6.81 (d, 1H), 6.77 (d, 1H), 6.64 (s, 1H), 6.46 (dd, 1H); 6.34 (s, 1H); 4.46 (m, 1H), 3.82-3.79 (m, 2H), 2.34 (m, 1H); 1.92 (m, 1H).

IR (nujol): 3356, 1724

Example 29

25 Sodium, 5,7-Dichloro-4-(2-oxo-1-phenyl- Δ^3 -pyrrolin-2-one-3-yl)-1,2,3,4-tetrahydro-quinoline-2-carboxylic acid; (enantiomer A)

To a solution of intermediate 26a (0.35g) in THF/H₂O (10ml, 3/1) at room temperature LiOH (0.06g) was added and stirring continued for 30min. The THF was evaporated and H₂O (5ml) was added. The resulting
30 solution was washed with diethylether (2x50ml). The aqueous phase was acidified until pH=4 with HCl 10% and the product filtered and dried under vacuum at 60 C for 12 hrs to afford the title compound (0.134g) as white solid.

The solid was dissolved in IMS (5% of methanol in ethanol) (10ml) and a 1N solution of NaOH (0.33ml) was added. To the resulting suspension diethyl ether was added (10ml) and the solid was filtered, washed with diethyl ether (10ml) and dried under vacuum for 12 hrs to give the title compound (0.082g) as a white solid.

5

m.p.>220°C

NMR (D₂O) δ (ppm) 7.49 (d, 2H); 7.40(t, 2H); 7.23 (t, 1H); 6.74 (d, 1H); 6.70 (d, 1H); 6.51 (m, 1H); 4.40-4.35 (m, 2H); 4.11 (m, 1H); 3.53 (dd, 1H); 2.18 (m, 1H); 1.74 (td, 1H)

10

HPLC Column: Cyclobond I, R,S-Hydroxypropyl ether 25 cm × 4.6 mm; Mobile Phase: Methanol=50 20 mM Ammonium Acetate buffer pH 5 = 50% by volume; Flow rate: 1 ml/min; Retention time: 12mins.

Example 30

15

Sodium 5,7-dichloro-4-(2-oxo-1-(phenyl)-pyrrolidin-3-ylidene)-1,2,3,4-tetrahydro-2-quinoline carboxylate(enantiomer A)

To a solution of intermediate 26b (0.052g) in THF/H₂O (4ml, 3/1) at room temperature LiOH (0.01g) was added and stirring continued for 30min. The THF was evaporated and H₂O (2ml) was added. The resulting solution was washed with diethylether (2x50ml). The aqueous phase was acidified until pH=4 with HCl 10% and the product was filtered and washed with water (10ml) and was dried under vacuum at 60°C for 12hrs to obtain 5,7-dichloro-4-(2-oxo-1-(pyridn-3yl)-pyrrolidin-3-ylidene)-1,2,3,4-tetrahydro-2-quinoline carboxylic acid 0.033g as a yellow solid.

20

The solid was dissolved in IMS (5% of methanol in ethanol) (5ml) and a 1N solution of NaOH (0.08ml) was added. After 5min, the solvent was evaporated and the solid triturated with diethyl ether (5ml), filtered, dried under vacuum for 12 hrs to give the title compound (0.01g) as a yellow solid.

25

m.p.: >200°

NMR (DMSO) δ (ppm) 7.74 (d, 2H); 7.39 (t, 2H); 7.15 (t, 1H); 6.76 (d, 1H); 6.51 (d, 1H); 6.20 (m, 1H); 4.63 (dd, 1H); 3.78 (m, 2H); 3.41 (dd, 1H); 3.18 (m, 1H); 2.35 (dd, 1H); 1.81 (t, 1H).

30

IR (nujol): 3363, 1688, 1630, 1586 cm⁻¹

Example 31

(±)-Ethyl 7-chloro-4-(1-phenyl- Δ^3 -pyrrolin-2-one-3yl)-1,2,3,4-tetrahydroquinoline-2-carboxylate (31a)

5 **(±)-Ethyl 7-chloro-4-(2-oxo-1-phenyl-3-pyrrolidinylidene)-1,2,3,4-tetrahydro-2-quinolinecarboxylate (31b)**

To a solution of intermediate 4 (2.2 g) in DMF (50 ml) Pd(PPh₃)₄ (244 mg) and TEA (1.2 ml) were added and the resulting solution was heated to 110°C for 2 h. The crude solution was poured into 200 ml of ethyl acetate and washed first with a saturated solution of NH₄Cl (2 x 150 ml),
10 then with water and brine. The organic phase was dried and concentrated to give the crude product. Purification by column chromatography (cyclohexane/dichloromethane/ethyl acetate 50/40/10) Rf=0.41 gave the title compound 31a (540 mg) as an off-white solid.

m.p.=150-153°C

15 NMR (DMSO) δ (ppm) 7.80 (m, 2H), 7.39 (m, 2H), 7.12 (m, 1H), 6.83 (d, 1H), 6.77 (d, 1H), 6.69 (m, 1H), 6.48 (dd, 1H), 6.45 (s, 1H), 4.48 (m, 2H), 4.15 (m, 2H), 3.94 (m, 1H), 3.82 (m, 1H), 2.34 (m, 1H), 1.97 (m, 1H), 1.20 (t, 3H)

IR (Nujol) (cm⁻¹) 3385, 1728, 1680

20 and the title compound 31b (475 mg) Rf=0.29 as a yellow solid.

m.p.=152-156°C

25 NMR (DMSO) δ (ppm) 7.72 (m, 2H), 7.39 (m, 2H), 7.20 (d, 1H), 7.16 (m, 1H), 6.98 (d, 1H), 6.74 (d, 1H), 6.57 (dd, 1H), 4.29 (dd, 1H), 4.21 (m, 1H), 4.02 (m, 1H), 3.93 (m, 1H), 3.82 (m, 1H), 3.69 (m, 1H), 3.20 (m, 1H), 2.92 (m, 2H), 0.93 (t, 3H)

Example 31a

(±)-Ethyl 7-chloro-4-(1-phenyl- Δ^3 -pyrrolin-2-one-3yl)-1,2,3,4-tetrahydroquinoline-2-carboxylate

30 To a solution of intermediate 4a (0.1g) in dry DMF (5ml) Pd(OAc)₂ (10mg) and TEA (0.026ml) were added. The mixture was heated at 110C for 2 hrs, then diluted with a saturated solution of NH₄Cl and extracted with ethyl acetate (2x10ml).

The solvent was evaporated and the crude purified by flash chromatography (Cyclohexane/EA 8:2) to afford the title compound as a white solid (40mg).

5 **Example 31b**

(±)-Ethyl 7-chloro-4-(2-oxo-1-phenyl-3-pyrrolidinylidene)-1,2,3,4-tetrahydro-2-quinolinecarboxylate

To a solution of intermediate 4b (370g) in toluene (5.2lit), Triethylamine (248ml), Triphenylphosphine (7.4g) and PdCl₂ (2.52g) were added. The resulting solution was warmed to 100 °C and stirred for 2h. The suspension was chilled to 20-25 °C and toluene (2.6ml) was added.

The reaction mixture was washed with NH₄Cl 8% (3x5.2lit) and water (5.2lit). The organic layer was filtered over a celite pad and it was washed with toluene (1lit); then it was distilled under vacuum (T=50°C; P=60mbar) to reach 6.3lit. After cooling to T=20-25°C, isooctane (5.2lit) was dropped over 30 min. The precipitate was stirred for 2h 30min then it was filtered and washed with a mixture toluene/isooctane 1/1 (1.85lit). The yellow solid was dried in vacuum at T= 40°C for 18h to obtain the title compound as a yellow solid 210g.

20 m.p. 160-162°C

NMR (DMSO): 7.72 (m, 2H); 7.39 (m, 2H); 7.20 (d, 2H); 7.15 (m, 2H); 6.96 (dd, 1H); 6.74 (d, 1H); 6.57 (dd, 1H); 4.29 (dd, 1H); 4.21 (m, 1H); 4.02 (m, 1H); 3.93 (m, 1H); 3.82 (m, 1H); 3.69 (m, 1H); 3.20 (m, 1H). 2.92 (m, 2H); 2.92 (m, 2H); 0.93 (t, 3H).

25

Pharmacy Examples

A. Capsules/ Tablets

| | |
|-----------------------|---------|
| Active ingredient | 20.0mg |
| Starch 1500 | 2.5mg |
| Microcrystalline | 200.0mg |
| Cellulose | |
| Croscarmellose Sodium | 6.0mg |
| Magnesium Stearate | 1.5mg |

5 The active ingredient is blended with the other excipients. The blend can be used to fill gelatin capsules or compressed to form tablets using appropriate punches. The tablets can be coated using conventional techniques and coatings.

B. Tablets

| | |
|----------------------------|--------|
| Active ingredient | 20.0mg |
| Lactose | 200.0m |
| | g |
| Microcrystalline Cellulose | 70.0mg |
| Povidone | 25.0mg |
| Croscarmellose | 6.0mg |
| Sodium Magnesium Stearate | 1.5mg |

10 The active ingredient is blended with lactose, microcrystalline cellulose and part of the croscarmellose sodium. The blend is granulated with povidone after dispersing in a suitable solvent (i.e. water). The granule, after drying and comminution is blended with the remaining excipients. The blend can be compressed using appropriate punches and the
15 tablets coated using conventional techniques and coatings.

c) Bolus

| | |
|---------------------------|---------------|
| 20 Active ingredient | 0.1-32mg/ml |
| Sodium phosphate | 1.0-50.0mg/ml |
| water for injection qs to | 1ml |

25 The formulation may be packed in glass ampoules or vials and syringes with a rubber stopper and a plastic/metal overseal (vials only).

D) Infusion

Active ingredient 0.01-3.2mg/ml
5 5% dextrose injection qs to 100ml

The formulation may be packed in glass vials or plastic bag.

10 The affinity of the compound of the invention for the strychnine insensitive glycine binding site was determined using procedure of Kishimoto H. *et al* J.

The pki values obtained with representative compounds of the invention are given in the following table:

15

| Example No | pki |
|------------|------|
| 1 | 8.1 |
| 14 | 7.9 |
| 15 | 7.73 |
| 16 | 7.8 |
| 17 | 8.7 |
| 18 | 7.78 |
| 19 | 8.9 |
| 21 | 7.1 |
| 22 | 7.9 |
| 24 | 7.8 |
| 25 | 7.15 |
| 30 | 7.7 |
| 29 | 8.7 |

The ability of compounds of the invention to inhibit pain in mouse has been assessed in the formalin test as described by Dubuisson and

Dennis (*Pain*, 1977, 4:161-174). In this test 20 μ l of 1 % formalin was injected into the plantar surface of the mouse left hind paw. The amount of time, in seconds, the animals spent licking the injected paw for the first 5 minutes (early phase) and then from 20 to 60 minutes (late phase) after formalin was used as measurement of the intensity of pain.

5 The compounds of the invention were administered orally 1 hour before formalin injection.

10 From these results the dose required to reduce the licking time by 50% expressed as mg/kg is referred to as the ED_{50s} value.


Representative results obtained for compounds of the invention when given by oral administration are given in the following table:

| Ex No | ED ₅₀ (mg/kg po) |
|-------|-----------------------------|
| 21 | 0.14 |
| 17 | 0.3 |
| 2 | 0.03 |

15 No untoward effects have been observed when compounds of the invention have been administered to mice at pharmacologically active doses.

Claims

1. A compound of formula(I)



5

or a salt or a non toxic metabolically labile esters thereof, wherein
Y represents a carbon atom;

- 10 Z is the group CH which is linked to the group Y via a double bond
and X is CH or Z is methylene or NR_{11} and X is a carbon atom
linked to the group Y via a double bond;

- A represents a C_{1-2} alkylene chain and which chain may be substituted
by one or two groups selected from C_{1-6} alkyl optionally substituted by
15 hydroxy, amino, C_{1-4} alkyl amino or C_{1-4} dialkyl amino or which chain
may be substituted by the group =O;

R represents a halogen atom or C_{1-4} alkyl group;

R_1 represents a hydrogen, a halogen atom or C_{1-4} alkyl group;

- 20 R_2 represents phenyl which may be substituted with up to 3 groups
selected from halogen, hydrogen, or $(\text{CH}_2)_n\text{R}_3$ wherein R_3 is COR_4 ,
 NR_6R_5 , NHCOR_7 , $\text{NHCONR}_9\text{R}_8$ or $\text{NH SO}_2 \text{R}_{10}$ group or R_2 is a 5
membered heteroaryl group containing 1 to 3 heteroatoms selected
from oxygen, sulphur and nitrogen; or 6 membered heteroaryl group
containing 1 to 3 nitrogen atoms

- 25 R_4 represents an amino, a hydroxyl or C_{1-4} alkoxy group;

R_5 and R_6 each independently represents hydrogen or C_{1-4} alkyl
group or

R₅ and R₆ together with the nitrogen atom to which they are attached represent a saturated 5-7 membered heterocyclic group optionally containing an additional heteroatom selected from oxygen, sulphur or nitrogen

- 5 R₇ represents a hydrogen atom or C₁₋₄ alkyl, C₁₋₄ alkoxy, or phenyl;
R₈ represents hydrogen or C₁₋₄ alkyl group;
R₉ represents hydrogen, optionally substituted C₁₋₄ alkyl (optionally substituted by one or more hydroxy carboxyl and amino group), phenyl;
- 10 R₁₁ represents hydrogen or C₁₋₄ alkyl group;
R₁₀ represents hydrogen, C₁₋₄ alkyl or a nitrogen protecting group.
n is zero or an integer from 1 to 2;

2. A compound as claimed in claim 1 wherein R is chlorine and
15 R₁ is hydrogen or a chlorine atom.

3. A compound as claimed in Claims 1 or 2 wherein A is a chain selected from -CH₂-, -(CH₂)₂- or C=O.

20 4. A compound as claimed in any of the claims 1-3 wherein Z is CH which is linked to the group Y via a double bond, a methylene or a NH group.

25 5. A compound as claimed in any of the claims 1-4 wherein R₂ is a group selected from phenyl (optionally substituted by acetylamino, methanesulphonylamino) or 3-pyridyl.

6. A compound as claimed in any of the claims 1-5 wherein R₂ represents phenyl.

30 7. A compound as claimed in any of the claims 1-6 wherein A is a chain selected from -CH₂-, -(CH₂)₂-, and Z is the group CH which is linked to the group Y via a double bond, or a methylene group, or A is C=O and Z is a NH group, R is chlorine, R₁ is chlorine or hydrogen

and R₂ is phenyl (optionally substituted by acetylamino or methanesulphonylamino) or 3-pyridyl.


- 5 8. (±)7-chloro-4-(2-oxo-1-phenyl-3-pyrrolidinylidene)-1,2,3,4-tetrahydro-2-quinoline carboxylic acid, physiologically acceptable salts or non toxic metabolically labile esters thereof.
9. Sodium (±) 7-chloro-4-(2-oxo-1-phenyl-3-pyrrolidinylidene)
10 1,2,3,4-tetrahydro-2-quinolinecarboxylate.
10. (-) Sodium 7-chloro-4-(2-oxo-1-phenyl-3-pyrrolidinylidene)
1,2,3,4- tetrahydro-2-quinolinecarboxylate.
- 15 11. (±)7-chloro-4-(1-phenyl-Δ³-pyrrolin-2-one-3yl)-1,2,3,4-tetrahydro-2-quinoline carboxylic acid, physiologically acceptable salts or non toxic metabolically labile esters thereof.
12. Sodium (±)7-chloro-4-(1-phenyl-Δ³-pyrrolin-2-one-3yl)-
20 1,2,3,4-tetrahydro-2-quinolinecarboxylate.
13. (-)Sodium 7-chloro-4-(1-phenyl-Δ³-pyrrolin-2-one-3yl)-
1,2,3,4-tetrahydro-2-quinolinecarboxylate.
- 25 14. (+) Sodium 7-chloro-4-(1-phenyl-Δ³-pyrrolin-2-one-3yl)-
1,2,3,4-tetrahydro-2-quinolinecarboxylate
15. A compound selected from:
- 30 (±)-7-chloro-4-(1-(3-pyridin)-Δ³-pyrrolin-2-one-3yl)-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,
(±)-7-chloro-4-(1-phenylΔ³-5,6-dihydro-pyridin-2-one-3yl)-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,
(±)-5,7-dichloro-4-(1-phenyl-Δ³-pyrrolin-2-one-3yl)-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

- (+/-)-7-chloro-4-(1-(4-acetylamino)-1-phenyl- Δ^3 -pyrrolin-2-one-3yl)-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,
(+/-)-7-chloro-4-(1-(4-methanesulfonylamino)-1-phenyl- Δ^3 -pyrrolin-2-one-3yl)-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,
5 (±)-7-chloro-4-(2-oxo-1-phenyl-3-piperidinylidene)-1,2,3,4-tetrahydro-2-quinolinecarboxylic acid,
(±)-7-chloro-4-(2,5-dioxo-1-phenyl-imidazolidin-4-ylidene)-1,2,3,4-tetrahydro-2-quinolinecarboxylic acid,
(±)-7-chloro-4-(2-oxo-1-(pyridin-3yl)-pyrrolidin-3-ylidene)-1,2,3,4-
10 tetrahydro-2-quinolinecarboxylate,
(±)-7-chloro-4-(2-oxo-1-(4-acetylamino)phenyl-pyrrolidin-3-ylidene)-1,2,3,4-tetrahydro-2-quinolinecarboxylic acid,
(±)-7-chloro-4-(2-oxo-1-((4-methanesulfonyl amino)phenyl-pyrrolidin-3-ylidene)-1,2,3,4-tetrahydro-2-quinolinecarboxylic acid,
15 5,7-dichloro-4-(2-oxo-1-(phenyl)-pyrrolidin-3-ylidene)-1,2,3,4-tetrahydro-2-quinoline carboxylic acid,
5,7-dichloro-4-(2-oxo-1-phenyl- Δ^3 -pyrrolin-2-one-3-yl)-1,2,3,4-tetrahydro-quinoline-2-carboxylic acid;
and physiologically acceptable salts (e.g. sodium salts), non-toxic
20 metabolically labile esters or enantiomers thereof.

16. A compound as claimed in any of claims 1-15 for use in therapy.
- 25 17. The use of a compound as claimed in any of claims 1-15 in the manufacturing of a therapeutic agent for antagonising the effects of excitatory amino acids on the NMDA receptor complex.
- 30 18. A pharmaceutical composition comprising a compound as claimed in any of claims 1-15 in admixture with one or more physiologically acceptable carriers or excipients.
- 35 19. A method of treatment of a mammal including man for conditions where antagonising the effects of excitatory amino acids on the NMDA receptor complex is of therapeutic benefit comprising

administration of an effective amount of a compound as claimed in any of claims 1 to 16.

20. A process for the preparation of compounds as claimed in any of the claims 1-15 which comprises cyclising a compound of formula (II) in which R, R₁, R₂, A, X, Y, Z have the meanings defined in claim 1 and R₁₂ is a carboxylic protecting group, R₁₃ represents a bromine or iodine atom, R₁₄ represents hydrogen or a nitrogen protecting group,



(II)

15 followed where necessary or desired by one or more of the following steps:

(i) removal of a protecting group.

(ii) isolation of the compound as a salt thereof;

20 (iii) conversion of a compound of formula (I) or a salt thereof into a metabolically labile ester thereof.

(iv) separation of a compound of formula (I) or a derivative thereof into the enantiomers thereof.

25