

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
9 September 2011 (09.09.2011)

PCT

(10) International Publication Number
WO 2011/108753 A2

(51) International Patent Classification:
A01N 37/38 (2006.01)

(21) International Application Number:
PCT/JP2011/055432

(22) International Filing Date:
2 March 2011 (02.03.2011)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
2010-046373 3 March 2010 (03.03.2010) JP

(71) Applicant (for all designated States except US): **SUMITOMO CHEMICAL COMPANY, LIMITED** [JP/JP]; 27-1, Shinkawa 2-chome, Chuo-ku, Tokyo, 1048260 (JP).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **KIGUCHI, So** [JP/JP]; 2-11-8-204, Sonehigashino-cho, Toyonaka-shi, Osaka, 5610802 (JP). **TANAKA, Soichi** [JP/JP]; 9-19-304, Matsuyama-cho, Nishinomiya-shi, Hyogo, 6638101 (JP). **OZAWA, Mayuko** [JP/JP]; 8-7-20-228, Honmachi, Toyonaka-shi, Osaka, 5600021 (JP). **IWATA, Atsushi** [JP/JP]; 3-12-23-306, Isoshi, Takarazuka-shi, Hyogo, 6650033 (JP).

(74) Agents: **TANAKA, Mitsuo** et al.; AOYAMA & PARTNERS, IMP Building, 3-7, Shiromi 1-chome, Chuo-ku, Osaka-shi, Osaka, 5400001 (JP).


(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report (Rule 48.2(g))

(54) Title: PLANT DISEASE CONTROLLING COMPOSITION AND METHOD FOR CONTROLLING PLANT DISEASE



(1)

(57) Abstract: The present invention provides a composition having an excellent controlling activity on plant disease. The composition comprising the compound represented by the formula (1) and one or more antibiotic fungicidal compound selected from the group (A) shows an excellent controlling activity on a plant disease. group (A): a group consisting of kasugamycin, polyoxins, streptomycin, and validamycin

WO 2011/108753 A2

DESCRIPTION

PLANT DISEASE CONTROLLING COMPOSITION AND METHOD FOR
CONTROLLING PLANT DISEASE

5

Technical Field

The present invention relates to a plant disease
controlling composition and a method for controlling a
plant disease.


10

Background Art

Hitherto, there has been provided compounds as an
active ingredient for a composition for controlling plant
disease (see e.g., The Pesticide Manual - 15th edition
15 (BCPC published) ISBN 1901396188; and SHIBUYA INDEX 13th
Edition (SHIBUYA-INDEX RESEARCH GROUP published)).

Also there has been provided a compound of the formula

(1):



20 (see e.g., WO 95/27693 pamphlet and WO 02/10101 pamphlet).


Disclosure of Invention

An object of the present invention is to provide a composition having an excellent control effect on a plant disease.

The present inventors have intensively studied to find out a composition having an excellent control effect on a plant disease. As a result, they have found that a composition comprising a compound represented by the formula (1) and one or more antibiotic fungicidal compound selected from the following group (A) shows a synergistic activity, and thus has an excellent control effect on a plant disease, and therefore the present invention has been completed.

The present invention provides:

[1] A plant disease controlling composition comprising a compound represented by the formula (1):



and one or more antibiotic fungicidal compound selected from the following group (A):


group (A): a group consisting of kasugamycin, polyoxins, streptomycin, and validamycin.

[2] The plant disease controlling composition according to the above [1], wherein a weight ratio of the compound

represented by the formula (1) to the antibiotic fungicidal compound is that of the compound represented by the formula (1)/the antibiotic fungicidal compound = 0.0125/1 to 500/1.

[3] The plant disease controlling composition according to the above [1] or [2], wherein the compound represented by the formula (1) is that represented by the formula (1) having R- absolute configuration.

[4] A method for controlling a plant disease which comprises applying each effective amount of the compound of the formula (1):



and one or more antibiotic fungicidal compound selected from the following group (A) to a plant or a soil for cultivating the plant,

group (A): a group consisting of kasugamycin, polyoxins, streptomycin, and validamycin.


[5] The method for controlling a plant disease according to the above [4], wherein the plant or the soil for cultivating the plant is a seed.

[6] The method for controlling a plant disease according to the above [4] or [5], wherein a weight ratio of the compound represented by the formula (1) to the antibiotic

fungicidal compound is that of the compound represented by the formula (1)/the antibiotic fungicidal compound = 0.0125/1 to 500/1.

[7] The method for controlling a plant disease according to any one of the above [4] to [6], wherein the compound represented by the formula (1) is that represented by the formula (1) having R- absolute configuration.

[8] A use of a combination of the compound represented by the formula (1):



10

and one or more antibiotic fungicidal compound selected from the following group (A) for controlling a plant disease,

group (A): a group consisting of kasugamycin, polyoxins, streptomycin, and validamycin.


15

The present invention enables to control a plant disease.

20 Best Mode for Carrying Out the Invention

A plant disease controlling composition of the present invention (hereinafter, referred to as a composition of the

present invention) comprises a compound represented by the formula (1):



(hereinafter, referred to as an amide compound of the present invention) and one or more antibiotic compound selected from the following group (A) (hereinafter, referred to as a antibiotic compound of the present invention),

group (A): a group consisting of kasugamycin, polyoxins, streptomycin, and validamycin.

The present amide compounds are those described in for example, WO 95/27693 pamphlet and WO 02/10101 pamphlet, and thus can be prepared according to the method described therein.

The present amide compound has one asymmetric carbon. Herein, a compound represented by the formula (1) wherein an enantiomer having R- absolute configuration is enriched is referred to as the amide compound having R- absolute configuration.

The present amide compound encompasses the following compounds:

a compound represented by the formula (1) which is

contained an enantiomer having R- absolute configuration in 70% and more;

a compound represented by the formula (1) which is contained an enantiomer having R- absolute configuration in
5 90% and more;

a compound represented by the formula (1) which is contained an enantiomer having R- absolute configuration in 95% and more.

Kasugamycin, polyoxins, streptomycin, and validamycin
10 that used in the present invention are all known compounds, and are described in for example, ''The PESTICIDE MANUAL - 15th EDITION (BCPC published) ISBN 1901396188'', pages 685, 920, 1053 and 1187 respectively. These compounds are either commercially available, or can be prepared by a
15 known method.

The weight ratio of the present amide compound to the present antibiotic compound in the composition of the present invention is usually that of the present
20 compound/the present antibiotic compound = 0.0125/1 to 500/1, preferably 0.025/1 to 100/1, and more preferably 0.1/1 to 10/1.

Although the composition of the present invention may be a mixture as itself of the present amide compound and the present antibiotic compound, the composition of the
25 present invention is usually prepared by mixing the present

amide compound, the present antibiotic compound and an inert carrier, and if necessary, adding a surfactant or other pharmaceutical additives, and then formulating into the form of oil solution, emulsifiable concentrate, flowable formulation, wettable powder, granulated wettable powder, dust formulation, granules and so on. Such formulations can be used by itself or with an addition of other inert components as an agent for controlling a plant disease.

10 Usually, the composition of the present invention can contain 0.1 to 99 % by weight, preferably 0.2 to 90 % by weight, and more preferably 1 to 80 % by weight of the present amide compound and the present antibiotic compound in total.

15 Examples of a solid carrier used on the formulation include finely-divided powder or particles of clay consisting of minerals (e.g., kaolin clay, attapulgite clay, bentonite, montmorillonite, acid clay, pyrophyllite, talc, diatomaceous earth, or calcite), natural organic substances (e.g., corncob powder, or walnut shell powder), synthetic organic substances (e.g., urea), salts (e.g., calcium carbonate, or ammonium sulfate), synthetic inorganic substances (e.g., synthetic hydrous silicon oxide) and so on. Examples of a liquid carrier include aromatic hydrocarbons (e.g., xylene, alkyl benzene, or

20

25

methylnaphtalene), alcohols (e.g., 2-propanol, ethylene glycol, propylene glycol, or ethylene glycol monoethyl ether), ketones (e.g., acetone, cyclohexanone, or isophorone), vegetable oils (e.g., soybean oil, or cotton
5 oils), petroleum-derived aliphatic hydrocarbons, esters, dimethylsulfoxide, acetonitrile and water.

Examples of the surfactant include anionic surfactant (e.g., alkyl sulfate salts, alkylaryl sulfate salts, dialkyl sulfosuccinate salts, polyoxyethylene alkylaryl
10 ether phosphates, lignin sulfonate, or naphthalenesulfonate formaldehyde polycondensation), nonionic surfactant (e.g., polyoxyethylene alkylaryl ether, polyoxyethylene alkyl polyoxypropylene block copolymer, or sorbitan fatty acid ester) and cationic surfactant (e.g., alkyltrimethyl
15 ammonium salts).

Examples of the other pharmaceutical additives include water-soluble polymer (e.g., polyvinyl alcohol, or polyvinyl pyrrolidone), polysaccharides (e.g. arabic gum, alginic acid and salts thereof, CMC (carboxymethyl-
20 cellulose), or xanthan gum), inorganic substances (e.g, aluminum magnesium silicate, or alumina-sol), antiseptic agent, coloring agent, and PAP (isopropyl acid phosphate), and stabilizing agent (e.g., BHT).

The composition of the present invention can also be
25 prepared by separately formulating the present amide

compound and the present antibiotic compound into different formulations by the above procedures, if necessary, further diluting each of them with water, thereafter, mixing the separately prepared different formulations or the dilute solutions.

The composition of the present invention may further contain one or more other fungicide and/or insecticide.

The composition of the present invention is used to control a plant disease by applying it to a plant or a soil for cultivating the plant.

The plant disease which can be controlled by the present invention is exemplified below:

Rice diseases: blast (*Magnaporthe oryzae*), helminthosporium leaf spot (*Cochliobolus miyabeanus*), sheath blight (*Rhizoctonia solani*) and bakanae disease (*Gibberella fujikuroi*);

Diseases of barley, wheat, oats and rye: powdery mildew (*Erysiphe graminis*), Fusarium head blight (*Fusarium graminearum*, *F. avenaceum*, *F. culmorum*, *F. asiaticum*, *Microdochium nivale*), rust (*Puccinia striiformis*, *P. graminis*, *P. recondite*, *P. hordei*), snow blight (*Typhula* sp., *Micronectriella nivalis*), loose smut (*Ustilago tritici*, *U. nuda*), bunt (*Tilletia caries*), eyespot (*Pseudocercospora herpotrichoides*), scald (*Rhynchosporium secalis*), leaf blotch (*Septoria tritici*),

glume blotch (*Leptosphaeria nodorum*) and net blotch (*Pyrenophora teres Drechsler*) ;

Citrus diseases: melanose (*Diaporthe citri*) , scab (*Elsinoe fawcetti*) and Penicillium rot (*Penicillium digitatum, P. italicum*) ;

Apple diseases: blossom blight (*Monilinia mali*) , canker (*Valsa ceratosperma*), powdery mildew (*Podosphaera leucotricha*) , Alternaria leaf spot (*Alternaria alternata apple pathotype*) , scab (*Venturia inaequalis*) , bitter rot (*Colletotrichum acutatum*) and late blight (*Phytophthora cactorum*) ;

Pear diseases: scab (*Venturia nashicola, V. pirina*) , black spot (*Alternaria alternate Japanese pear pathotype*) , rust (*Gymnosporangium haraeaeum*) and late blight (*Phytophthora cactorum*) ;

Peach diseases: brown rot (*Monilinia fructicola*) , scab (*Cladosporium carpophilum*) and Phomopsis rot (*Phomopsis sp.*) ;

Grapes diseases: anthracnose (*Elsinoe ampelina*) , ripe rot (*Glomerella cingulata*) , powdery mildew (*Uncinula necator*) , rust (*Phakopsora ampelopsidis*) , black rot (*Guignardia bidwellii*) , downy mildew (*Plasmopara viticola*) and Gray mold (*Botrytis cinerea*) ;

Diseases of Japanese persimmon: anthracnose (*Gloeosporiura kaki*) and leaf spot (*Cercospora kaki*,

Mycosphaerella nawae) ;

Diseases of gourd family: anthracnose (*Colletotrichum lagenarium*) , powdery mildew (*Sphaerotheca fuliginea*) , gummy stem blight (*Mycosphaerella melonis*) , Fusarium wilt
5 (*Fusarium oxysporum*) , downy mildew (*Pseudoperonospora cubensis*) , Phytophthora rot (*Phytophthora* sp.) and damping-off (*Pythium* sp.) ;

Tomato diseases: early blight (*Alternaria solani*) , leaf mold (*Cladosporium fulvum*) and late blight
10 (*Phytophthora infestans*) ;

Egg plant disease: brown spot (*Phomopsis vexans*) and powdery mildew (*Erysiphe cichoracearum*) ;

Diseases of Cruciferous Vegetables: *Alternaria* leaf spot (*Alternaria japonica*) , white spot (*Cercospora brassicae*) , clubroot (*Plasmodiophora brassicae*), and downy
15 mildew (*Peronospora parasitica*) ;

Rapeseed diseases: *Sclerotinia* rot (*Sclerotinia sclerotiorum*), black spot (*Alternaria brassicae*), powdery mildew (*Erysiphe cichoracearum*), blackleg (*Leptosphaeria maculans*) ;
20

Welsh onion diseases: rust (*Puccinia allii*) ;

Soybean diseases: purple seed stain (*Cercospora kikuchii*) , Sphaceloma scad (*Elsinoe glycines*) , pod and stem blight (*Diaporthe phaseolorum* var. *sojae*) , rust
25 (*Phakopsora pachyrhizi*) and phytophthora stem rot

(*Phytophthora sojae*) ;

Adzuki-bean diseases: Gray mold (*Botrytis cinerea*) ,
Sclerotinia rot (*Sclerotinia sclerotiorum*) ;

Kindney bean diseases: Gray mold (*Botrytis cinerea*) ,
5 Sclerotinia rot (*Sclerotinia sclero tiorum*) , anthracnose
(*Colletotrichum lindemthianum*) ;

Peanut diseases: leaf spot (*Cercospora personata*) ,
brown leaf spot (*Cercospora arachidicola*) and southern
blight (*Sclerotium rolfsii*) ;

10 Garden pea diseases: powdery mildew (*Erysiphe pisi*) ;

Potato diseases: early blight (*Alternaria solani*) and
late blight (*Phytophthora infestans*) ;

Strawberry diseases: powdery mildew (*Sphaerotheca
humuli*) ;

15 Tea diseases: net blister blight (*Exobasidium
reticulatum*) , white scab (*Elsinoe leucospila*) , gray
blight (*Pestalotiopsis* sp.) and anthracnose (*Colletotrichum
theae-sinensis*) ;

Cotton diseases: fusarium wilt (*Fusarium oxysporum*) ,
20 damping-off (*Rhizoctonia solani*);

Tabacco diseases: brown spot (*Alternaria longipes*) ,
powdery mildew (*Erysiphe cichoracearum*) , anthracnose
(*Colletotrichum tabacum*) , downy mildew (*Peronospora
tabacina*) and late blight (*Phytophthora nicotianae*) ;

25 Sugar beet diseases: Cercospora leaf spot (*Cercospora*

beticola) , leaf blight (*Thanatephorus cucumeris*) , Root rot (*Aphanidermatum cochlioides*) ;

Rose diseases: black spot (*Diplocarpon rosae*) and powdery mildew (*Sphaerotheca pannosa*) ;

5 Chrysanthemum diseases: leaf blight (*Septoria chrysanthemi-indici*) and white rust (*Puccinia horiana*) ;

Various plants diseases: diseases caused by *Pythium* spp. (*Pythium aphanidermatum*, *Pythium debarianum*, *Pythium graminicola*, *Pythium irregulare*, *Pythium ultimum*) , Gray mold (*Botrytis cinerea*) , Sclerotinia rot (*Sclerotinia sclerotiorum*) ,

10

Japanese radish diseases: *Alternaria* leaf spot (*Alternaria brassicicola*) ;

Turfgrass diseases: dollar spot (*Sclerotinia homeocarpa*) , brown patch and large patch (*Rhizoctonia solani*) ; and

15

Banana diseases: Sigatoka disease (*Mycosphaerella fijiensis*, *Mycosphaerella musicola*, *Pseudocercospora musae*).

Examples of the plants to which the composition of the present invention can be applied are as follows:

20

Crops: corn, rice, wheat, barley, rye, oat, sorghum, cotton, soybean, adzuki-bean, kidney bean, peanut, buckwheat, beet, rapeseed, sunflower, sugar cane, and tobacco, etc.;

25 Vegetables: solanaceous vegetables (eggplant, tomato,

pimento, pepper, and potato, etc.) , cucurbitaceous vegetables (cucumber, pumpkin, zucchini, water melon, melon, and squash, etc.) , cruciferous vegetables (Japanese radish, white turnip, horseradish, kohlrabi, Chinese cabbage, cabbage, leaf mustard, broccoli, and cauliflower, etc.) ,
5 asteraceous vegetables (burdock, crown daisy, artichoke, and lettuce, etc.) , liliaceous vegetables (green onion, onion, garlic, and asparagus) , ammiaceous vegetables (carrot, parsley, celery, and parsnip, etc.) ,
10 chenopodiaceous vegetables (spinach, and Swiss chard, etc.) , lamiaceous vegetables (Perilla frutescens, mint, and basil, etc.) , strawberry, sweet potato, Dioscorea japonica, and colocasia, etc.;

Flowers;

15 Foliage plants;

Turfgrass;

Fruits: pomaceous fruits (apple, pear, Japanese pear, Chinese quince, and quince, etc.) , stone fleshy fruits (peach, plum, nectarine, Prunus mume, cherry fruit, apricot, and prune, etc.) , citrus fruits (Citrus unshiu, orange, lemon, lime, and grapefruit, etc.) , nuts (chestnut, walnuts, hazelnuts, almond, pistachio, cashew nuts, and macadamia nuts, etc.) , berrys (blueberry, cranberry, blackberry, and raspberry, etc.) , grape, kaki fruit, olive,
20 Japanese plum, banana, coffee, date palm, and coconuts,
25

etc.; and

Trees other than fruit trees: tea, mulberry, flowering plant, roadside trees (ash, birch, dogwood, Eucalyptus, Ginkgo biloba, lilac, maple, Quercus, poplar, Judas tree, Liquidambar formosana, plane tree, zelkova, Japanese arborvitae, fir wood, hemlock, juniper, Pinus, Picea, and Taxus cuspidate), etc.

The aforementioned "plants" include plants which resistance has been imparted by genetic recombination.

Exemplary embodiments of the composition of the present invention are as follows:

a composition comprising the present amide compound and kasugamycin wherein a weight ratio thereof is that of the present amide compound/kasugamycin = 0.0125/1 to 500/1;

a composition comprising the present amide compound and kasugamycin wherein a weight ratio thereof is that of the present amide compound/kasugamycin = 0.025/1 to 100/1;

a composition comprising the present amide compound and kasugamycin wherein a weight ratio thereof is that of the present amide compound/kasugamycin = 0.1/1 to 10/1;

a composition comprising the present amide compound and polyoxins wherein a weight ratio thereof is that of the present amide compound/polyoxins = 0.0125/1 to 500/1;

a composition comprising the present amide compound and polyoxins wherein a weight ratio thereof is that of the

present amide compound/polyoxins = 0.025/1 to 100/1;

a composition comprising the present amide compound and polyoxins wherein a weight ratio thereof is that of the present amide compound/polyoxins = 0.1/1 to 10/1;

5 a composition comprising the present amide compound and streptomycin wherein a weight ratio thereof is that of the present amide compound/streptomycin = 0.0125/1 to 500/1;

10 a composition comprising the present amide compound and streptomycin wherein a weight ratio thereof is that of the present amide compound/streptomycin = 0.025/1 to 100/1;

a composition comprising the present amide compound and streptomycin wherein a weight ratio thereof is that of the present amide compound/streptomycin = 0.1/1 to 10/1;

15 a composition comprising the present amide compound and validamycin wherein a weight ratio thereof is that of the present amide compound/validamycin = 0.0125/1 to 500/1;

20 a composition comprising the present amide compound and validamycin wherein a weight ratio thereof is that of the present amide compound/validamycin = 0.025/1 to 100/1; and

a composition comprising the present amide compound and validamycin wherein a weight ratio thereof is that of the present amide compound/validamycin = 0.1/1 to 10/1.

25 The method for controlling a plant disease of the

present invention (hereinafter, referred to as the method
for controlling of the present invention) is carried out by
applying each effective amount of the present amide
compound and the present antibiotic compound to the plants
5 or the soil for cultivating the plant.

Such the plants may be, for example, foliages of plant,
seeds of plant, or bulbs of plant. The bulbs herein are
intended to mean bulb, corm, rootstock, tubera, tuberous
root and rhizophore.

10 In the method for controlling of the present invention,
the present amide compound and the present antibiotic
compound may be applied separately around the same time to
the plant or the soil for cultivating the plant, but is
usually applied as the composition of the present invention
15 in terms of a convenience on applying.

In the method for controlling of the present invention,
examples of the method of applying the present amide
compound and the antibiotic compound include foliage
treatment, soil treatment, root treatment and seed
20 treatment.

Such the foliage treatment includes for example, a
method of applying the composition of the present invention
to a surface of the plant to be cultivated by a foliage
application or a stem application.

25 Such the root treatment includes for example, a method

of soaking a whole or a root of the plant into a medicinal solution comprising the present amide compound and the present antibiotic compound, and a method of attaching a solid formulation comprising the present amide compound, the present antibiotic compound and the solid carrier to a root of the plant.

Such the soil treatment includes for example, soil broadcast, soil incorporation, and irrigation of the medicinal solution to a soil.

Such the seed treatment includes for example, an applying of the composition of the present invention to a seed or a bulb of the plant to be prevented from the plant disease, specifically, for example, a spray treatment by spraying a suspension of the composition of the present invention in a mist form to a surface of a seed or a surface of a bulb, a smear treatment by smearing the wettable powder, the emulsifiable concentrate or the flowable formulation of the composition of the present invention with added by small amounts of water or as itself to a seed or a bulb, an immerse treatment of a seed into a solution of the composition of the present invention for a given time, a film-coating treatment, and a pellet-coating treatment.

Each dose of the present amide compound and the present antibiotic compound in the method for controlling

of the present invention may be varied depending on a kind of plant to be treated, a kind or a frequency of an occurrence of a plant disease as a control subject, a dosage form, a treatment period, a treatment method, a treatment site, a climate condition, etc. In case of an application to a foliage of the plant or a soil for cultivating the plant, a total amount of the present amide compound and the antibiotic compound is usually 1 to 500 g, preferably 2 to 200 g, and more preferably 10 to 100 g, per 1000 m². Each dose of the present amide compound and the present antibiotic compound in the treatment for seed is usually 0.001 to 10 g, and preferably 0.01 to 1 g, per 1kg of seeds.

The emulsifiable concentrate, the wettable powder or the flowable formulation, etc., is usually applied by diluting them with water, and then spreading them. In this case, usually, each concentration of the present amide compound and the present antibiotic compound contain 0.0005 to 2% by weight, and preferably 0.005 to 1% by weight of the present amide compound and the present antibiotic compound in total. The dust formulation or the granular formulation, etc, is usually applied as itself without diluting them.

25 EXAMPLES

Next, the present invention is described in more detail below by the following examples including formulation examples and a test example, but the present invention should not be construed to be limited thereto.

5 The formulation examples are given below. It is to be noted that in the formulation examples, the term "'part'" indicates "'part by weight'".

Formulation 1

10 5 parts of the present amide compound, 5 parts of kasugamycin, 35 parts of a mixture of white carbon and polyoxyethylene alkylether sulfate ammonium salts (weight ratio 1:1), and 55 parts of water were mixed and the resulting solution was then subjected to fine grinding
15 according to a wet grinding method, so as to obtain a flowable formulation. The same above operations were carried out with polyoxins, streptomycin, or validamycin instead of kasugamycin, so as to obtain various types of flowable formulations.

20

Formulation 2

10 parts of the present amide compound, 5 parts of kasugamycin and 1.5 parts of sorbitan trioleate were mixed into 28 parts of an aqueous solution that contained 2 parts
25 of polyvinyl alcohol, and the mixed solution was then

subjected to fine grinding according to a wet grinding method. Thereafter, 45.50 parts of an aqueous solution that contained 0.05 parts of xanthan gum and 0.1 part of aluminum magnesium silicate was added to the resultant, and 5 10 parts of propylene glycol was further added thereto. The obtained mixture was blended by stirring, so as to obtain the flowable formulation. The same above operations were carried out with polyoxins, streptomycin, or validamycin instead of kasugamycin, so as to obtain various 10 types of flowable formulations.

Formulation 3

10 parts of the present amide compound, 40 parts of kasugamycin, 3 parts of calcium lignosulfonate, 2 parts of 15 sodium lauryl sulfate, and 45 parts of synthetic hydrous silicon oxide were fully crushed and mixed, so as to obtain wettable powders. The same above operations were carried out with polyoxins, streptomycin, or validamycin instead of kasugamycin, so as to obtain various types of wettable 20 powders.

The test examples are given below.

Test Example 1

To a SUS beaker are added 30 parts of the present 25 compound (racemate), 5 parts of Soprophor FLK

(polyoxyethylene tristyrylphenyl ether phosphate, produced by Rhodia Nicca Ltd.), 65 parts of ion-exchange water and 150 parts of glass beads (1.0mm diameter). The mixture is subjected to grinding at a velocity of 1000 rpm with
5 cooling to 10°C, and thereafter the glass beads are separated, so as to obtain a flowable formulation comprising the present compound (racemate).

True leaf of cucumber is punched out with cork borer to 13mm in diameter to prepare a leaf disk. In 24 well
10 microwell plate that is dispensed with 1ml 0.8% water agar, the leaf disk is placed such that the upper side of the leaf is in an upward direction. Thereto is added 20 micro liter a testing solution prepared by mixing the flowable formulation of the present compound (racemate) prepared by
15 the above procedure and a commercial liquid formulation of validamycin or a commercial wettable powders of polyoxins to a predetermined concentration to treat the leaf disk.

After confirming that the testing medical solution is dried, conidium of gray mold fungus (*Botrytis cinerea*) is
20 suspended into potato dextrose broth (DIFCO) in a density of about 10^5 conidium/mL and is then subjected to a spray inoculation. After leaving to stand the leaf disk in a growth chamber set up at 15°C for four days, an onset area on the leaf is measured and then calculated an onset area
25 rate (hereinafter, referred to as an onset area rate of

treated group).

The same operation is carried out with 20 micro liter water instead of 20 micro liter a testing medicine solution to calculate an onset area rate (hereinafter, referred to an onset area rate of non-treated group).

A preventive value is calculated from the above onset area rate of treated group and the onset area rate of non-treated group by the following equation:

$$\text{Preventive value (\%)} = 100 \times (A-B)/A$$

10 wherein

A: an onset area rate of treated group

B: an onset area rate of non-treated group

The results are shown in Tables 1 and 2.

Table 1

	treatment concentration (ppm)		preventive value (%)
	the present amide compound	validamycin	
1	2.5	0.5	100
2	1.0	5.0	100

15

Table 2

	treatment concentration (ppm)		preventive value (%)
	the present amide compound	polyoxins	
1	2.5	0.5	100
2	1.0	5.0	100

Test Example 2

The same operations as described in Test Example 1 are

carried out with kasugamycin or streptomycin instead of validamycin or polyoxins, so as to calculate respective preventive values.

Also for comparison, the same operations as described in Test Example 1 are carried out with the exception that the testing medicine solution is substituted with a predetermined concentration of either a flowable formulation of the present amide compound or a commercial wettable powder of kasugamycin or streptomycin, so as to calculate respective preventive values.

The results are shown in Tables 3 to 4.

Table 3

	treatment concentration (ppm)		preventive value (%)
	the present amide compound	kasugamycin	
1	2.5	0.5	100
2	1.0	5.0	100
	2.5	-	56
	1.0	-	46
	-	0.5	0
	-	5.0	10

Table 4

	treatment concentration (ppm)		preventive value (%)
	the present amide compound	streptomycin	
1	2.5	0.5	100
2	1.0	5.0	100
	2.5	-	56
	1.0	-	46
	-	0.5	0
	-	5.0	10

Next, the Reference Examples are given below.

Reference Examples

For comparison, the same operations as described in Test Example 1 are carried out with the exception that the testing medicine solution is substituted with a predetermined concentration of a solution of a wettable powder of validamycin or polyoxins, so as to calculate a preventive value.

The results are shown in Tables 5 to 6.

10 Table 5


	treatment concentration (ppm)	preventive value (%)
	validamycin	
	0.5	5
	5.0	10

Table 6

	treatment concentration (ppm)	preventive value (%)
	polyoxins	
	0.5	11
	5.0	15

CLAIMS

1. A plant disease controlling composition comprising a compound represented by the formula (1):



5


and one or more antibiotic fungicidal compound selected from the following group (A):

group (A): a group consisting of kasugamycin, polyoxins, streptomycin, and validamycin.

10 2. The plant disease controlling composition according to claim 1, wherein a weight ratio of the compound represented by the formula (1) to the antibiotic fungicidal compound is that of the compound represented by the formula (1)/the antibiotic fungicidal compound = 0.0125/1 to 500/1.

15 3. The plant disease controlling composition according to claim 1 or 2, wherein the compound represented by the formula (1) is that represented by the formula (1) having R- absolute configuration.

20 4. A method for controlling a plant disease which comprises applying each effective amount of the compound of the formula (1):



and one or more antibiotic fungicidal compound selected from the following group (A) to a plant or a soil for cultivating the plant,


5 group (A): a group consisting of kasugamycin, polyoxins, streptomycin, and validamycin.

5. The method for controlling a plant disease according to claim 4, wherein the plant or the soil for cultivating the plant is a seed.

10 6. The method for controlling a plant disease according to claim 4 or 5, wherein a weight ratio of the compound represented by the formula (1) to the antibiotic fungicidal compound is that of the compound represented by the formula (1)/the antibiotic fungicidal compound = 0.0125/1 to 500/1.

15 7. The method for controlling a plant disease according to any one of claims 4 to 6, wherein the compound represented by the formula (1) is that represented by the formula (1) having R- absolute configuration.

8. A use of a combination of the compound represented by
20 the formula (1):



and one or more antibiotic fungicidal compound selected from the following group (A) for controlling a plant disease,

5 group (A): a group consisting of kasugamycin, polyoxins, streptomycin, and validamycin.