



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) **EP 1 090 022 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

- (45) Date of publication and mention of the grant of the patent:
06.08.2003 Bulletin 2003/32
- (51) Int Cl.7: **C07H 19/167**, A61K 31/70,
C07H 15/04
- (21) Application number: **99931113.7**
- (86) International application number:
PCT/EP99/04269
- (22) Date of filing: **23.06.1999**
- (87) International publication number:
WO 99/067265 (29.12.1999 Gazette 1999/52)

(54) **2-(PURIN-9-YL)-TETRAHYDROFURAN-3,4-DIOL DERIVATIVES**

2-(PURIN-9-YL)-TETRAHYDROFURAN-3,4-DIOL DERIVATE
DERIVES DU 2-(PURIN-9-YL)-TETRAHYDROFURAN-3,4 DIOL

- (84) Designated Contracting States:
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE**
Designated Extension States:
AL LT LV MK RO SI
- (30) Priority: **23.06.1998 GB 9813538**
23.04.1999 GB 9909482
- (43) Date of publication of application:
11.04.2001 Bulletin 2001/15
- (73) Proprietor: **GLAXO GROUP LIMITED**
Greenford, Middlesex UB6 ONN (GB)
- (72) Inventors:
• **ALLEN, David George**
Stevenage, Hertfordshire SG1 2NY (GB)
• **CHAN, Chuen**
Stevenage, Hertfordshire SG1 2NY (GB)
• **COUSINS, Richard Peter Charles**
Stevenage, Hertfordshire SG1 2NY (GB)
• **COX, Brian**
Stevenage, Hertfordshire SG1 2NY (GB)
- **GEDEN, Joanna Victoria**
Aston Science Park, Birmingham B7 4EJ (GB)
• **HOBBS, Heather**
Stevenage, Hertfordshire SG1 2NY (GB)
• **KEELING, Suzanne Elaine**
Stevenage, Hertfordshire SG1 2NY (GB)
• **REDGRAVE, Alison Judith**
Stevenage, Hertfordshire SG1 2NY (GB)
• **ROPER, Thomas Davis IV**
Research Triangle Park, NC 27709 (US)
• **XIE, Shiping**
Research Triangle Park, NC 27709 (US)
- (74) Representative: **Teuten, Andrew John et al**
GlaxoSmithKline
Corporate Intellectual Property (CN9.25.1)
980 Great West Road
Brentford, Middlesex TW8 9GS (GB)
- (56) References cited:
WO-A-94/17090 **WO-A-98/01459**
WO-A-98/28319 **WO-A-99/38877**
WO-A-99/41267

EP 1 090 022 B1

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Description

[0001] This invention relates to new chemical compounds, processes for their preparation, pharmaceutical formulations containing them and their use in therapy.

[0002] Inflammation is a primary response to tissue injury or microbial invasion and is characterised by leukocyte adhesion to the endothelium, diapedesis and activation within the tissue. Leukocyte activation can result in the generation of toxic oxygen species (such as superoxide anion), and the release of granule products (such as peroxidases and proteases). Circulating leukocytes include neutrophils, eosinophils, basophils, monocytes and lymphocytes. Different forms of inflammation involve different types of infiltrating leukocytes, the particular profile being regulated by the profile of adhesion molecule, cytokine and chemotactic factor expression within the tissue.

[0003] The primary function of leukocytes is to defend the host from invading organisms such as bacteria and parasites. Once a tissue is injured or infected a series of events occurs which causes the local recruitment of leukocytes from the circulation into the affected tissue. Leukocyte recruitment is controlled to allow for the orderly destruction and phagocytosis of foreign or dead cells, followed by tissue repair and resolution of the inflammatory infiltrate. However in chronic inflammatory states, recruitment is often inappropriate, resolution is not adequately controlled and the inflammatory reaction causes tissue destruction.

[0004] There is evidence from both *in vitro* and *in vivo* studies to suggest that compounds active at the adenosine A2a receptor will have anti-inflammatory actions. The area has been reviewed by Cronstein (1994). Studies on isolated neutrophils show an A2 receptor-mediated inhibition of superoxide generation, degranulation, aggregation and adherence (Cronstein et al, 1983 and 1985; Burkey and Webster, 1993; Richter, 1992; Skubitz et al, 1988. When agents selective for the A2a receptor over the A2b receptor (eg CGS21680) have been used, the profile of inhibition appears consistent with an action on the A2a receptor subtype (Dianzani et al, 1994). Adenosine agonists may also down-regulate other classes of leucocytes (Elliot and Leonard, 1989; Peachell et al, 1989). Studies on whole animals have shown the anti-inflammatory effects of methotrexate to be mediated through adenosine and A2 receptor activation (Asako et al, 1993; Cronstein et al, 1993 and 1994). Adenosine itself, and compounds that raise circulating levels of adenosine also show anti-inflammatory effects *in vivo* (Green et al, 1991; Rosengren et al, 1995). In addition raised levels of circulating adenosine in man (as a result of adenosine deaminase deficiency) results in immunosuppression (Hirschorn, 1993).

[0005] Certain substituted 4'-carboxamido and 4'-thioamido adenosine derivatives which are useful for the treatment of inflammatory diseases are described in International Patent Application Nos. WO94/17090, WO96/02553, WO96/02543 (Glaxo Group). Substituted 4'-carboxamidoadenosine derivatives useful in the treatment of dementia are described in AU 8771946 (Hoechst Japan). Substituted 4'-hydroxymethyl adenosine derivatives which are useful for the treatment of gastrointestinal motility disorders are described in EP-A-423776 and EP-A-423777 (Searle). Substituted 4'-hydroxymethyl adenosine derivatives which are useful as platelet aggregation inhibitors are described in BE-768925 (Takeda). 4'-Hydroxymethyl adenosine derivatives and 4'-esters thereof which are useful as anti-hypertensive agents or have other cardiovascular activity are described in US 4663313, EP 139358 and US 4767747 (Warner Lambert), US 4985409 (Nippon Zoki) and US 5043325 (Whitby Research). 4-Hydroxymethyladenosine derivatives useful in the treatment of autoimmune disorders are described in US 5106837 (Scripps Research Institute). 4'-Hydroxymethyladenosine derivatives useful as anti-allergic agents are described in US 4704381 (Boehringer Mannheim). Certain 4'-tetrazolylalkyl adenosine derivatives which are useful in the treatment of heart and circulatory disorders are generically described in DT-A-2621470 (Pharma-Waldhof). Other 4'-carboxamidoadenosine derivatives useful in the treatment of cardiovascular conditions are described in US 5219840, GB 2203149 and GB 2199036 (Sandoz), WO94/02497 (US Dept. Health), US 4968697 and EP 277917 (Ciba Geigy), US 5424297 (Univ. Virginia) and EP 232813 (Warner Lambert). Other 4'-carboxamidoadenosine derivatives lacking substitution on the purine ring in the 2-position are described in DT 2317770, DT 2213180, US 4167565, US 3864483 and US 3966917 (Abbott Labs), DT 2034785 (Boehringer Mannheim), JP 58174322 and JP 58167599 (Tanabe Seiyaku), WO92/05177 and US 5364862 (Rhone Poulenc Rorer), EP 66918 (Procter and Gamble), WO86/00310 (Nelson), EP 222330, US 4962194, WO88/03147 and WO88/03148 (Warner Lambert) and US 5219839, WO95/18817 and WO93/14102 (Lab UPSA). 4'-Hydroxymethyladenosine derivatives lacking substitution on the purine ring in the 2-position are described in WO95/11904 (Univ Florida). 4'-Substituted adenosine derivatives useful as adenosine kinase inhibitors are described in WO94/18215 (Gensia). Other 4'-halomethyl, methyl, thioalkylmethyl or alkoxymethyl adenosine derivatives are described in EP 161128 and EP 181129 (Warner Lambert) and US 3983104 (Schering). Other 4'-carboxamidoadenosine derivatives are described in US 7577528 (NIH), WO91/13082 (Whitby Research) and WO95/02604 (US Dept Health).

[0006] Certain tetrazole containing deoxynucleotides which were found to lack antiinfective activity are described in Baker et al (1974) Tetrahedron 30, 2939-2942. Other tetrazole containing adenosine derivatives which show activity as platelet aggregation inhibitors are described in Mester and Mester (1972) Pathologie-Biologie, 20 (Suppl) 11-14. Certain nitrile containing ribose derivatives are described in Schmidt et al (1974) Liebigs. Ann. Chem. 1856-1863.


[0007] Other publications include: WO 98/16539 (Novo Nordisk A/S) which describes adenosine derivatives for the

treatment of myocardial and cerebral ischaemia and epilepsy; WO 98/01426 (Rhone-Poulenc Rorer Pharmaceuticals Inc.) which relates to adenosine derivatives possessing antihypertensive, cardioprotective, anti-ischaemic and antilipolytic properties; and WO 98/01459 (Novo Nordisk A/S) which describes *N*,9-disubstituted adenine derivatives which are substituted in the 4' position by unsubstituted oxazolyl or isoxazolyl and the use of such compounds for the treatment of disorders involving cytokines in humans. WO 98/28319 (Glaxo Group Limited) was published subsequent to the earliest priority date of this application and describes 4'-substituted tetrazole 2-(purin-9-yl)-tetrahydrofuran-3,4-diol derivatives.

[0008] We have now found a novel group of compounds with broad anti-inflammatory properties which inhibit leucocyte recruitment and activation and which are agonists of the adenosine 2a receptor. The compounds are therefore of potential therapeutic benefit in providing protection from leukocyte-induced tissue damage in diseases where leukocytes are implicated at the site of inflammation. The compounds of the invention may also represent a safer alternative to corticosteroids in the treatment of inflammatory diseases, whose uses may be limited by their side-effect profiles.


[0009] More particularly, the compounds of this invention may show an improved profile over known A2a-selective agonists in that they generally lack significant agonist activity at the human A3 receptor. This profile can be considered of benefit as A3 receptors are also found on leucocytes (eg eosinophil) and other inflammatory cells (eg mast cell) and activation of these receptors may have pro-inflammatory effects (Kohno et al, 1996; Van Schaick et al 1996). It is even considered that the bronchoconstrictor effects of adenosine in asthmatics may be mediated via the adenosine A3 receptor (Kohno et al, 1996).

[0010] Thus, according to the invention we provide compounds of formula (I):




wherein R¹ and R² independently represent a group selected from:

- (i) C₃₋₈cycloalkyl-;
- (ii) hydrogen;
- (iii) aryl₂CHCH₂-;
- (iv) C₃₋₈cycloalkylC₁₋₆alkyl-;
- (v) C₁₋₈alkyl-;
- (vi) arylC₁₋₆alkyl-;
- (vii) R⁴R⁵N-C₁₋₆alkyl-;
- (viii) C₁₋₆alkyl-CH(CH₂OH)-;
- (ix) arylC₁₋₅alkyl-CH(CH₂OH)-;
- (x) arylC₁₋₅alkyl-C(CH₂OH)₂-;
- (xi) C₃₋₈cycloalkyl independently substituted by one or more (e.g. 1, 2 or 3) -(CH₂)_pR⁶ groups;
- (xii) H₂NC(=NH)NHC₁₋₆alkyl-;
- (xiii) a group of formula



- or such a group in which one methylene carbon atom adjacent to X, or both if such exist, is substituted by methyl;
- (xiv) -C₁₋₆alkyl-OH;
- (xv) -C₁₋₈haloalkyl;

(xvi) a group of formula



(xvii) aryl; and

(xviii) $-(\text{CH}_2)_f \text{SO}_2 \text{NH}_g (\text{C}_{1-4} \text{alkyl})_{2-g}$ or $-(\text{CH}_2)_f \text{SO}_2 \text{NH}_g (\text{arylC}_{1-4} \text{alkyl})_{2-g}$;

R^3 represents methyl, ethyl, $-\text{CH}=\text{CH}_2$, n-propyl, $-\text{CH}_2\text{CH}=\text{CH}_2$, $-\text{CH}=\text{CHCH}_3$, isopropyl, isopropenyl, cyclopropyl, cyclopropenyl, cyclopropylmethyl, cyclopropenylmethyl, cyclobutyl, cyclobutenyl, $-(\text{CH}_2)_q$ halogen,

$-(\text{CH}_2)_h \text{Y} (\text{CH}_2)_i \text{H}$, $-(\text{CH}_2)_h \text{COOCH}_3$, $-(\text{CH}_2)_h \text{OCOCH}_3$,

$-(\text{CH}_2)_h \text{CON} (\text{CH}_2)_m \text{H} ((\text{CH}_2)_n \text{H})$, $-(\text{CH}_2)_h \text{CO} (\text{CH}_2)_o \text{H}$ or

$-\text{CH}_2\text{C} ((\text{CH}_2)_u \text{H}) = \text{NO} (\text{CH}_2)_v \text{H}$;

Y represents O, S or $\text{N}(\text{CH}_2)_j$;

a and b independently represent an integer 0 to 4 provided that $a + b$ is in the range 3 to 5;

c, d and e independently represent an integer 0 to 3 provided that $c + d + e$ is in the range 2 to 3;

f represents 2 or 3 and g represents an integer 0 to 2;

p represents 0 or 1;

q represents 2 or 3;

h represents 2 or 3;

i represents an integer 0 to 2 such that $h+i$ is in the range 2 to 4

j represents an integer 0 to 2 such that $h+i+j$ is in the range 2 to 4

m and n independently represent an integer 0 to 2 such that $m+n$ is in the range 0 to 2;

o represents an integer 0 to 2 such that $h+o$ is in the range 2 to 3;

u and v independently represent 0 or 1 such that $u+v$ is in the range 0 to 1;

R^4 and R^5 independently represent hydrogen, C_{1-6} alkyl, aryl, arylC_{1-6} alkyl- or NR^4R^5 together may represent pyrrolidinyl, piperidinyl, morpholinyl, azetidiny, azepinyl, piperazinyl or N-C_{1-6} alkylpiperazinyl.

R^6 represents OH, NH_2 , NHCOCH_3 or halogen;

R^7 represents hydrogen, C_{1-6} alkyl, C_{1-6} alkylaryl or $-\text{COC}_{1-6}$ alkyl;


X represents NR^7 , O, S, SO or SO_2 ;

provided that when R^3 represents methyl, ethyl or isopropyl then R^1 and/or R^2 independently must represent:

(a) $-(\text{CH}_2)_f \text{SO}_2 \text{NH}_g (\text{C}_{1-4} \text{alkyl})_{2-g}$ or $-(\text{CH}_2)_f \text{SO}_2 \text{NH}_g (\text{arylC}_{1-4} \text{alkyl})_{2-g}$ where f is 2 or 3 and g is an integer 0 to 2;


(b) C_{3-8} cycloalkyl independently substituted by one or more $-(\text{CH}_2)_p \text{NHCOCH}_3$ groups;

(c) a group of formula



or such a group in which one methylene carbon atom adjacent to X, or both if such exist, is substituted by methyl;

(d) a group of formula



and salts and solvates thereof.

[0011] References to C₁₋₆alkyl include references to an aliphatic hydrocarbon grouping containing 1 to 6 carbon atoms which may be straight chain or branched and may be saturated or unsaturated although will be preferably saturated. References to C₁₋₄alkyl, C₁₋₅alkyl, C₂₋₄alkyl and C₁₋₈alkyl may be interpreted similarly.

[0012] References to aryl include references to mono- and bicyclic carbocyclic aromatic rings (e.g. phenyl, naphthyl) and heterocyclic aromatic rings containing 1-3 hetero atoms selected from N, O and S (e.g. pyridinyl, pyrimidinyl, thiophenyl, imidazolyl, quinolinyl, furanyl, pyrrolyl, oxazolyl) all of which may be optionally substituted, e.g. by C₁₋₆alkyl, halogen, hydroxy, nitro, C₁₋₆alkoxy, cyano, amino, SO₂NH₂ or -CH₂OH.

[0013] Examples of C₃₋₈cycloalkyl for R¹ and R² include monocyclic alkyl groups (e.g. cyclopentyl, cyclohexyl) and bicyclic alkyl groups (e.g. norbornyl such as exonorborn-2-yl).

Examples of (aryl)2CHCH₂- for R¹ and R² include Ph₂CHCH₂- or such a group in which one or both phenyl moieties is substituted, e.g. by halogen or C₁₋₄alkyl.

Examples of C₃₋₈cycloalkylC₁₋₆alkyl- for R¹ and R² include ethylcyclohexyl.


Examples of C₁₋₈alkyl for R¹ and R² include -(CH₂)₂C(Me)₃, -CH(Et)₂ and CH₂=C(Me)CH₂CH₂-.

Examples of arylC₁₋₆alkyl- for R¹ and R² include -(CH₂)₂Ph, -CH₂Ph or either in which Ph is substituted (one or more times) by halogen (e.g. iodine), amino, methoxy, hydroxy, -CH₂OH or SO₂NH₂; -(CH₂)₂ pyridinyl (e.g. -(CH₂)₂pyridin-2-yl) optionally substituted by amino; (CH₂)₂imidazolyl or this group in which imidazolyl is N-substituted by C₁₋₆alkyl (especially methyl).

Examples of R⁴R⁵N-C₁₋₆alkyl- for R¹ and R² include ethyl-piperidin-1-yl, ethyl-pyrrolidin-1-yl, ethyl-morpholin-1-yl, -(CH₂)₂NH(pyridin-2-yl) and -(CH₂)₂NH₂.

Examples of C₁₋₆alkyl-CH(CH₂OH)- for R¹ and R² include Me₂CHCH(CH₂OH)-.

Examples of arylC₁₋₅alkyl-CH(CH₂OH)- for R¹ and R² include PhCH₂CH(CH₂OH)-especially




Examples of aryl C₁₋₅alkyl-O(CH₂OH)₂- for R¹ and R² include PhCH₂C(CH₂OH)₂-.

Examples of C₃₋₈ cycloalkyl independently substituted by one or more -(CH₂)_pR⁶ groups (eg 1, 2 or 3 such groups) for R¹ and R² include 2-hydroxy-cyclopentyl and 4-aminocyclohexyl (especially trans-4-amino-cyclohexyl).

Examples of H₂NC(=NH)NHC₁₋₆alkyl for R¹ and R² include H₂NC(=NH)NH(CH₂)₂-

Examples of groups of formula




for R¹ and R² include pyrrolidin-3-yl, piperidin-3-yl, piperidin-4-yl or a derivative in which the ring nitrogen is substituted by C₁₋₆alkyl (e.g. methyl) or benzyl, tetrahydro-1,1-dioxide thiophen-3-yl, tetrahydropyran-4-yl, tetrahydrothiopyran-4-yl and 1,1-dioxo-hexahydro-1.lamda.6-thiopyran-4-yl.

Examples of -C₁₋₆alkyl-OH groups for R¹ and R² include -CH₂CH₂OH.

Examples of C₁₋₈haloalkyl for R¹ and R² include -CH₂CH₂Cl and (CH₃)₂CIC(CH₂)₃-.

Examples of groups of formula



for R¹ and R² include 2-oxopyrrolidin-4-yl, 2-oxo-pyrrolidin-5-yl or a derivative in which the ring nitrogen is substituted by C₁₋₆alkyl (e.g. methyl) or benzyl.

Examples of aryl for R¹ and R² include phenyl optionally substituted by halogen (e.g. fluorine, especially 4-fluorine). Examples of C₁₋₆alkyl for R⁷ include methyl and C₁₋₆alkylaryl for R⁷ include benzyl. Examples of COC₁₋₆ alkyl for R⁷ include -COCH₃.

[0014] Examples of C₁₋₅alkyl for R³ include n-propyl and allyl. An example of C₃₋₄ cycloalkyl for R³ includes cyclobutyl. An example of -(CH₂)_hO(CH₂)_iH for R³ includes -(CH₂)₂OMe. Examples of C₂₋₄alkyl substituted by halogen or hydroxy include -(CH₂)₂Cl, -(CH₂)₂OH and -(CH₂)₃OH.

[0015] We prefer that R¹ and R² do not both represent hydrogen.

We prefer R¹ to represent aryl₂CHCH₂-, C₁₋₈alkyl, hydrogen or aryl C₁₋₆alkyl-.

We particularly prefer R¹ to represent Ph₂CHCH₂-, -CH(Et)₂, hydrogen or phenylethyl-, especially Ph₂CHCH₂-.

[0016] We prefer R² to represent R⁴R⁵N-C₁₋₆alkyl-, arylC₁₋₆alkyl-, arylC₁₋₅alkylCH(CH₂OH)-, aryl C₁₋₆ alkyl or C₁₋₆ alkyl-CH(CH₂OH)-.

We particularly prefer R² to represent (CH₂)₂(piperidin-1-yl), 2-(1-methyl-1H-imidazol-4-yl)ethyl, 1S-hydroxymethyl-2-phenylethyl, phenylethyl or 1S-hydroxymethyl-2-methyl propyl, especially -(CH₂)₂(piperidin-1-yl).

[0017] We prefer R³ to represent C₁₋₃alkyl (including n-propyl and 2-propenyl), cyclobutyl, cyclopropylmethyl, -(CH₂)₂OCOCH₃, -(CH₂)₂₋₃OH or -(CH₂)₂halogen. More preferably R³ represents n-propyl, 2-propenyl, cyclobutyl, cyclopropylmethyl, -(CH₂)₂OCOCH₃, or -(CH₂)₂₋₃OH.

We particularly prefer R³ to represent -(CH₂)₂OCOCH₃, -(CH₂)₂OH or (CH₂)₃OH, especially -(CH₂)₂OCOCH₃ or -(CH₂)₂OH, most especially -(CH₂)₂OH.

[0018] We prefer R⁴ and R⁵ independently to represent hydrogen, C₁₋₆alkyl or aryl or NR⁴R⁵ together to represent pyrrolidinyl, piperidinyl, morpholinyl, azetidiny, azepinyl, piperazinyl or N-methylpiperazinyl;

We prefer X to represent NR⁷, O, S or SO₂, particularly NR⁷ or SO₂, especially NR⁷.

[0019] We prefer that a and b both represent 2 or that a represents 1 and b represents 2.

We prefer that R⁷ represents hydrogen.

We prefer that p represents 0. We prefer q to represent 2. We prefer h to represent 2. We prefer i to represent 0 or 1, especially 0. We prefer j to represent 1. We prefer m and n to represent 0 or 1. We prefer o to represent 1.

We prefer u and v to represent 0.


We prefer that R⁶ represents OH or NH₂ especially NH₂.

We prefer that c represents 0 and either d represents 2 and e represents 0 or d represents 1 and e represents 1.

[0020] The representation of formula (I) indicates the absolute stereochemistry at positions around the tetrahydrofuran ring. When sidechains contain chiral centres the invention extends to mixtures of enantiomers (including racemic mixtures) and diastereoisomers as well as to individual enantiomers. Generally it is preferred to use a compound of formula (I) in the form of a purified single enantiomer.

[0021] We also provide a process for preparation of compounds of formula (I) which comprises:

(a) reacting a corresponding compound of formula (II)




wherein L represents a leaving group

or a protected derivative thereof

with a compound of formula R²NH₂ or a protected derivative thereof;

(b) preparing a compound of formula (I) in which R¹ represents hydrogen by reducing a compound of formula (III)



15 or a protected derivative thereof; or

(c) deprotecting a compound of formula (I) which is protected; and where desired or necessary converting a compound of formula (I) or a salt thereof into another salt thereof.

20 **[0022]** In process (a) L represents a leaving group such as halogen eg chlorine or fluorine. The reaction of process (a) will generally be carried out on heating the reagents to a temperature of 50 °C-150 °C in the presence of a solvent such as DMSO. Preferably an organic base, e.g. a trisubstituted organic amine (such as diisopropylethylamine) is also present for the reaction. Under these conditions we particularly prefer that Hal represents fluorine (especially when R¹ represents hydrogen) since the reaction has a tendency to proceed rapidly with high efficiency.

[0023] In process (b) the reduction reaction may be performed by catalytic hydrogenation, e.g. over Pd/C under standard conditions.


25 **[0024]** In process (c) examples of protecting groups and the means for their removal can be found in T W Greene "Protective Groups in Organic Synthesis" (J Wiley and Sons, 1991). Suitable hydroxyl protecting groups include alkyl (e.g. methyl), acetal (e.g. acetonide) and acyl (e.g. acetyl or benzoyl) which may be removed by hydrolysis, and arylalkyl (e.g. benzyl) which may be removed by catalytic hydrogenolysis. Suitable amine protecting groups include sulphonyl (e.g. tosyl), acyl e.g. benzyloxycarbonyl or t-butoxycarbonyl) and arylalkyl (e.g. benzyl) which may be removed by hydrolysis or hydrogenolysis as appropriate.

30 **[0025]** Suitable salts of the compounds of formula (I) include physiologically acceptable salts such as acid addition salts derived from inorganic or organic acids, for example hydrochlorides, hydrobromides, 1-hydroxy-2-naphthoates, mesylates, sulphates, phosphates, acetates, benzoates, citrates, succinates, lactates, tartrates, fumarates and maleates, and if appropriate, inorganic base salts such as alkali metal salts, for example sodium salts. Other salts of the compounds of formula (I) include salts which may not be physiologically acceptable but may be useful in the preparation of compounds of formula (I) and physiologically acceptable salts thereof. Examples of such salts include trifluoroacetates and formates.

35 **[0026]** Examples of suitable solvates of the compounds of formula (I) include hydrates.

40 **[0027]** Acid-addition salts of compounds of formula (I) may be obtained by treating a free-base of formula (I) with an appropriate acid.


[0028] The compounds of formula (II) or a protected derivative thereof may be prepared by reacting a compound of IV



or a protected derivative thereof with a compound of formula R¹NH₂. L¹ and L² independently represent a leaving group such as halogen eg chlorine or fluorine. This reaction will preferably be performed in the presence of a base

such as an organic amine base (e.g. diisopropyl ethylamine) in a solvate such as an alcohol (e.g. isopropanol) at elevated temperature (e.g. reflux).


[0029] Compounds of formula (III) or a protected derivative thereof may be prepared by reacting a compound of formula (IIIA)



wherein L represents a leaving group such as halogen eg chlorine or fluorine, or a protected derivative thereof, with a compound of formula R^2NH_2 under conventional conditions.

[0030] Compounds of formula (IIIA), or a protected derivative thereof, may be prepared by reacting a compound of formula (IV), or a protected derivative thereof, with an azide, e.g. sodium azide under conventional conditions.


[0031] The compound of formula (IV) or a protective derivative thereof may be prepared by reacting a compound of formula (V)



wherein L represents a leaving group or a protected derivative thereof with a 2,6-dihalopurine, e.g. 2,6-dichloropurine.

[0032] We prefer to use the compound of formula (V) wherein the ribose 2- and 3-hydroxy groups are protected, e.g. by acetyl. Leaving group L may represent OH but will preferably represent C_{1-6} alkoxy (e.g. methoxy or ethoxy), an ester moiety (e.g. acetyloxy or benzoyloxy) or halogen. The preferred group L is acetyloxy. The reaction may be performed by combining the reactants in an inert solvent such as MeCN in the presence of a Lewis Acid (eg TMSOTf) and DBU and warming to, say, 70-80 °C.

[0033] Compounds of formula (V) may be prepared from a compound of formula (VI)




wherein alk represents C_{1-6} alkyl eg methyl by treating the compound of formula (VI) with trifluoroacetic acid in water followed by reprotection, e.g. by reaction with acetic anhydride in pyridine.

[0034] Compounds of formula (V) in which L represents halogen, may be prepared from the corresponding 1-alcohol or a 1-ester such as the acetate. Reaction will generally occur on treatment with anhydrous HCl or HBr. 1-iodides may be prepared directly on treatment with trimethylsilyliodide and 1-fluorides may be prepared on treatment with DAST.

An inert solvent eg diethylether, DCM, THF or CCl_4 will generally be suitable.


[0035] Compound of formula (VI) may be prepared following Scheme 1:



[0036] General conditions for Stages 1-6 will be known to persons skilled in the art. It will also be appreciated that the reagents and conditions set out in Scheme 1 are example conditions and alternative reagents and conditions for achieving the same chemical transformation may be known to persons skilled in the art. For example an alternative alcohol, e.g. a C_{1-6} alkyl alcohol may be used in Stage 1 to give a different C_{1-6} alkyloxy leaving group in compounds of formula (VII) and (VI). Stage 1 may also be modified in that the use of HCl can be substituted by perchloric acid (HClO_4) and 2,2 dimethoxypropane, or alternatively acetyl chloride (which has the advantage that it maintains high yield and avoids use of perchlorate salts). Alternative reaction conditions can be used in Stage 3, which may utilise ethyl acetate, thionyl chloride and gaseous ammonia (which has the advantage that it avoids chlorinated solvent and the synthesis of troublesome ammonium pivaloate impurity). Stage 4 may also be performed using POCl_3 , TEA, DMF and ethyl acetate in the reaction conditions (which avoids the use of hazardous DMAP). Compounds of formula (VII) wherein a leaving group besides OMe is desired may be prepared by analogy with the method described above for preparation of compounds of formula (V). Alternative groups may be used to protect the 2' and 3' hydroxy groups on the ribose in Stage 1. We have also found that Stage 5 may desirably be performed using azidotrimethylsilane and dibutyltin oxide in toluene.

[0037] Following stage 6, the impure product may be purified using conventional techniques, and especially using flash chromatography conditions under nitrogen pressure. We have found that satisfactory conditions include loading the impure product in a minimum volume of dichloromethane onto a Keiselgel 60 (Merck 9385) column and eluting using a gradient solvent system with ethyl acetate (10-40%) in cyclohexane.

[0038] Compounds of formula (II), and protected derivatives thereof, may also be prepared by reacting a compound of formula (V), or a protected derivative thereof with a compound of formula (VIII)



10 wherein L represents a leaving group such as halogen, e.g. chlorine or fluorine optionally followed by a deprotection or deprotection and reprotection reaction.


[0039] We prefer to use compounds of formula (V) in protected form. In particular we prefer that at least the hydroxy group in the 2- position on the ribose is protected as an ester group, e.g. by acetyl or benzoyl since this has a tendency to result in greater stereoselectivity in the coupling reaction. We prefer that the 2- and 3-position hydroxy groups are protected by acetyl. Suitable leaving groups L are as described previously. The preferred leaving group L is acetyloxy.

15 [0040] This process is particularly preferred when L represents fluorine (and most especially when R¹ represents hydrogen) since the reaction is generally fast and efficient and the reaction has a tendency to produce products of high crystallinity.

[0041] The product of this reaction may be deprotected if desired under conventional conditions eg on treatment with an alcohol (eg isopropanol) under mild basic conditions (eg in the presence of potassium carbonate)

20 [0042] The reaction of compounds of formula (V) (in protected form) and compounds of formula (VIII) may be performed in the presence of a Lewis Acid (eg TMSOTf) and optionally a silylating agent (eg BSA) in an inert solvent such as acetonitrile followed by work-up eg with water. When L represents halogen the Lewis Acid can generally be omitted when a silylating agent is present.

25 [0043] Certain compounds of formula (VIII) are known. Other compounds of formula (VIII) may be prepared by reaction of a compound of formula (IX)



wherein L¹ and L² independently represent a leaving group such as halogen, e.g. chlorine or fluorine, with R¹NH₂ under conventional conditions.

40 [0044] Compounds of formula R¹NH₂, R²NH₂ and IX are either known or may be prepared by conventional methods known *per se*.


[0045] As a further aspect of the invention we also provide new process which may be used to provide compounds of formula (I) without the proviso.

45 [0046] Thus we provide a process for preparation of a compound of formula (I) without the proviso which comprises

(d) reacting a corresponding compound of formula (X)

50


55



15 with a compound of formula (XI)



wherein L is a leaving group; or
(e) reacting a corresponding compound of formula (XII)




with a compound of formula (V) or a protected derivative thereof.

[0047] Process (d) will generally take place on combining the two reagents in the presence of a mild base e.g. K_2CO_3 and an inert organic solvent eg DMF. Typical leaving groups L include halogen (e.g. Br).

[0048] Process (e) will generally take place in the presence of a Lewis Acid (e.g. TMSOTf) and optionally a silylating agent (e.g. BSA) in an inert solvent such as MeCN followed by work-up e.g. with water. We prefer L to represent acetyloxy and the two hydroxy groups to be protected as the acetyl ester. A deprotection step (using mild base e.g. K_2CO_3) will then be necessary to generate the compound of formula (I).

Compounds of formula (X) may be prepared by analogous methods to those described above for the preparation of compounds of formula (I). When compounds of formula (X) are prepared via the analogues of compounds of formula (II), (III), (IIIA) and/or (IV) in which R^3 is replaced by hydrogen, such compounds are preferably protected in the N2-position of the tetrazole. A suitable protecting group is benzyl which may be incorporated by treating the unprotected tetrazole with a benzyl halide (e.g. benzyl bromide) in the presence of base (e.g. K_2CO_3). An illustrative process for the preparation of compounds of formula (X) is given in Scheme 2:


Scheme 2



[0049] Compounds of formula (XI) are known or may be prepared by known methods.

[0050] Compounds of formula (XII) may be prepared, for example following Scheme 3:

Scheme 3



30 [0051] Processes (d) and (e) are particularly suitable for preparing the compound (2R,3R,4S,5R)-2-[6-Amino-2-(1S-hydroxymethyl-2-phenyl-ethyl-amino)-purin-9-yl]-5-(2-ethyl-2H-tetrazol-5-yl)-tetrahydro-furan-3,4-diol and salts and solvates thereof, especially the maleate salt.

[0052] We prefer process (e).

35 [0053] The potential for compounds of formula (I) to inhibit leukocyte function may be demonstrated, for example, by their ability to inhibit superoxide (O_2^-) generation from neutrophils stimulated with chemoattractants such as N-formylmethionylleucyl-phenylalanine (fMLP). Accordingly, compounds of formula (I) are of potential therapeutic benefit in providing protection from leukocyte-induced tissue damage in diseases where leukocytes are implicated at the site of inflammation.

40 [0054] Examples of disease states in which the compounds of the invention have potentially beneficial anti-inflammatory effects include diseases of the respiratory tract such as adult respiratory distress syndrome (ARDS), bronchitis (including chronic bronchitis), cystic fibrosis, asthma (including allergen-induced asthmatic reactions), chronic obstructive pulmonary disease (COPD), emphysema, rhinitis and septic shock. Other relevant disease states include diseases of the gastrointestinal tract such as intestinal inflammatory diseases including inflammatory bowel disease (e.g. Crohn's disease or ulcerative colitis), *Helicobacter-pylori* induced gastritis and intestinal inflammatory diseases secondary to radiation exposure or allergen exposure, and non-steroidal anti-inflammatory drug-induced gastropathy. Furthermore, compounds of the invention may be used to treat skin diseases such as psoriasis, allergic dermatitis and hypersensitivity reactions and diseases of the central nervous system which have an inflammatory component eg Alzheimer's disease and multiple sclerosis.

45 [0055] Further examples of disease states in which compounds of the invention have potentially beneficial effects include cardiac conditions such as peripheral vascular disease, post-ischæmic reperfusion injury and idiopathic hypereosinophilic syndrome.

50 [0056] Compounds of the invention which inhibit lymphocyte function may be useful as immunosuppressive agents and so have use in the treatment of auto-immune diseases such as rheumatoid arthritis and diabetes.

[0057] Compounds of the invention may also be useful in inhibiting metastasis.

[0058] Diseases of principal interest include asthma and COPD.

55 [0059] It will be appreciated by those skilled in the art that reference herein to treatment extends to prophylaxis as well as the treatment of established conditions.

[0060] As mentioned above, compounds of formula (I) are useful in human or veterinary medicine, in particular as anti-inflammatory agents.

[0061] There is thus provided as a further aspect of the invention a compound of formula (I) or a physiologically acceptable salt or solvate thereof for use in human or veterinary medicine, particularly in the treatment of patients with inflammatory conditions who are susceptible to leukocyte-induced tissue damage.

5 **[0062]** According to another aspect of the invention, there is provided the use of a compound of formula (I) or a physiologically acceptable salt or solvate thereof for the manufacture of a medicament for the treatment of patients with inflammatory conditions who are susceptible to leukocyte-induced tissue damage.

10 **[0063]** In a further or alternative aspect there is provided a method for the treatment of a human or animal subject with an inflammatory condition who is susceptible to leukocyte-induced tissue damage, which method comprises administering to said human or animal subject an effective amount of a compound of formula (I) or a physiologically acceptable salt or solvate thereof.

[0064] The compounds according to the invention may be formulated for administration in any convenient way, and the invention therefore also includes within its scope pharmaceutical compositions for use in anti-inflammatory therapy, comprising a compound of formula (I) or a physiologically acceptable salt or solvate thereof together, if desirable, with one or more physiologically acceptable diluents or carriers.

15 **[0065]** There is also provided a process for preparing such a pharmaceutical formulation which comprises mixing the ingredients.

[0066] The compounds according to the invention may, for example, be formulated for oral, buccal, parenteral, topical or rectal administration, preferably for parenteral or topical (e.g. by aerosol) administration.

20 **[0067]** Tablets and capsules for oral administration may contain conventional excipients such as binding agents, for example syrup, acacia, gelatin, sorbitol, tragacanth, mucilage of starch, cellulose or polyvinyl pyrrolidone; fillers, for example, lactose, microcrystalline cellulose, sugar, maize- starch, calcium phosphate or sorbitol; lubricants, for example, magnesium stearate, stearic acid, talc, polyethylene glycol or silica; disintegrants, for example, potato starch, croscarmellose sodium or sodium starch glycolate; or wetting agents such as sodium lauryl sulphate. The tablets may be coated according to methods well known in the art. Oral liquid preparations may be in the form of, for example, 25 aqueous or oily suspensions, solutions, emulsions, syrups or elixirs, or may be presented as a dry product for constitution with water or other suitable vehicle before use. Such liquid preparations may contain conventional additives such as suspending agents, for example, sorbitol syrup, methyl cellulose, glucose/sugar syrup, gelatin, hydroxymethyl cellulose, carboxymethyl cellulose, aluminium stearate gel or hydrogenated edible fats; emulsifying agents, for example, lecithin, sorbitan mono-oleate or acacia; non-aqueous vehicles (which may include edible oils), for example almond 30 oil, fractionated coconut oil, oily esters, propylene glycol or ethyl alcohol; or preservatives, for example, methyl or propyl *p*- hydroxybenzoates or sorbic acid. The preparations may also contain buffer salts, flavouring, colouring and/or sweetening agents (e.g. mannitol) as appropriate.

[0068] For buccal administration the compositions may take the form of tablets or lozenges formulated in conventional manner.

35 **[0069]** The compounds may also be formulated as suppositories, e.g. containing conventional suppository bases such as cocoa butter or other glycerides.

[0070] The compounds according to the invention may also be formulated for parenteral administration by bolus injection or continuous infusion and may be presented in unit dose form, for instance as ampoules, vials, small volume infusions or pre-filled syringes, or in multi-dose containers with an added preservative. The compositions may take 40 such forms as solutions, suspensions, or emulsions in aqueous or non-aqueous vehicles, and may contain formulatory agents such as anti-oxidants, buffers, antimicrobial agents and/or tonicity adjusting agents. Alternatively, the active ingredient may be in powder form for constitution with a suitable vehicle, e.g. sterile, pyrogen-free water, before use. The dry solid presentation may be prepared by filling a sterile powder aseptically into individual sterile containers or by filling a sterile solution aseptically into each container and freeze-drying.

45 **[0071]** By topical administration as used herein, we include administration by insufflation and inhalation. Examples of various types of preparation for topical administration include ointments, creams, lotions, powders, pessaries, sprays, aerosols, capsules or cartridges for use in an inhaler or insufflator, solutions for nebulisation or drops (e.g. eye or nose drops).

50 **[0072]** Ointments and creams may, for example, be formulated with an aqueous or oily base with the addition of suitable thickening and/or gelling agents and/or solvents. Such bases may thus, for example, include water and/or an oil such as liquid paraffin or a vegetable oil such as arachis oil or castor oil or a solvent such as a polyethylene glycol. Thickening agents which may be used include soft paraffin, aluminium stearate, cetostearyl alcohol, polyethylene glycols, microcrystalline wax and beeswax.

55 **[0073]** Lotions may be formulated with an aqueous or oily base and will in general also contain one or more emulsifying agents, stabilising agents, dispersing agents, suspending agents or thickening agents.

[0074] Powders for external application may be formed with the aid of any suitable powder base, for example, talc, lactose or starch. Drops may be formulated with an aqueous or non-aqueous base also comprising one or more dispersing agents, solubilising agents or suspending agents.

[0075] Spray compositions may be formulated, for example, as aqueous solutions or suspensions or as aerosols delivered from pressurised packs, with the use of a suitable propellant, e.g. dichlorodifluoromethane, trichlorofluoromethane, dichlorotetra-fluoroethane, 1,1,1,2,3,3,3-heptafluoropropane, 1,1,1,2-tetrafluoroethane, carbon dioxide or other suitable gas.

[0076] Intranasal sprays may be formulated with aqueous or non-aqueous vehicles with the addition of agents such as thickening agents, buffer salts or acid or alkali to adjust the pH, isotonicity adjusting agents or anti-oxidants. Capsules and cartridges of for example gelatin, or blisters of for example laminated aluminium foil, for use in an inhaler or insufflator may be formulated containing a powder mix of a compound of the invention and a suitable powder base such as lactose or starch.

[0077] Solutions for inhalation by nebulation may be formulated with an aqueous vehicle with the addition of agents such as acid or alkali, buffer salts, isotonicity adjusting agents or antimicrobials. They may be sterilised by filtration or heating in an autoclave, or presented as a non-sterile product.

[0078] The pharmaceutical compositions according to the invention may also be used in combination with other therapeutic agents, for example anti-inflammatory agents (such as corticosteroids (eg fluticasone propionate, beclomethasone dipropionate, mometasone furoate, triamcinolone acetonide or budesonide) or NSAIDs (eg sodium cromoglycate)) or beta adrenergic agents (such as salmeterol, salbutamol, formoterol, fenoterol or terbutaline and salts thereof) or anti-infective agents (eg antibiotics, antivirals).

[0079] The invention thus provides, in a further aspect, a combination comprising a compound of formula (I) or a physiologically acceptable salt or solvate thereof together with another therapeutically active agent, for example an anti-inflammatory agent such as a corticosteroid or NSAID.

[0080] The combination referred to above may conveniently be presented for use in the form of a pharmaceutical formulation and thus pharmaceutical formulations comprising a combination as defined above together with a physiologically acceptable diluent or carrier thereof represent a further aspect of the invention.

[0081] The individual components of such combinations may be administered either sequentially or simultaneously in separate or combined pharmaceutical formulations. Appropriate doses of known therapeutic agents will be readily appreciated by those skilled in the art.

[0082] Compounds of the invention may conveniently be administered in amounts of, for example, 0.01 to 500mg/kg body weight, preferably 0.01 to 100mg/kg body weight, 1 to 4 times daily. The precise dose will of course depend on the age and condition of the patient and the particular route of administration chosen.

[0083] The compounds of the invention have the advantage that they may be more efficacious, show greater selectivity, have fewer side effects, have a longer duration of action, be more bioavailable by the preferred route, show less systemic activity when administered by inhalation or have other more desirable properties than similar known compounds.

[0084] In particular the compounds of the invention have the advantage that they may show greater selectivity for the adenosine 2a receptor subtype over other adenosine receptor subtypes (especially the A1 and A3 receptor subtypes) than hitherto known compounds.

[0085] As a further aspect of the invention we provide certain compounds as new and useful intermediates.

[0086] Compounds of the invention may be tested for in vitro and in vivo biological activity in accordance with the following screens:

(1) Agonist activity against adenosine 2a, adenosine 1 and adenosine 3 receptor subtypes.

Agonist selectivity of compounds against other human adenosine receptors is determined using Chinese hamster ovary (CHO) cells transfected with the gene for the relevant human adenosine receptor following a method based on that of Castanon and Spevak, 1994. The CHO cells are also transfected with cyclic AMP response elements promoting the gene for secreted placental alkaline phosphatase (SPAP) (Wood, 1995). The effect of test compounds is determined by their effects on basal levels of cAMP (A2a) or on forskolin-enhanced cAMP (A1 and A3) as reflected by changes in levels of SPAP. EC₅₀ values for compounds are then determined as a ratio to that of the non-selective agonist N-ethyl carboxamide adenosine (NECA).

(2) Antigen-induced lung eosinophil accumulation in sensitised guinea pigs.

Ovalbumin sensitised guinea pigs are dosed with mepyramine (1mg/kg ip) to protect against anaphylactic bronchospasm. A compound of the invention is then given by the inhaled route (30min breathing of an aerosol of the compound) immediately prior to ovalbumin challenge (30min breathing of an aerosol generated from a 50ug/ml solution of ovalbumin). Twenty four hours after challenge, the guinea pigs are killed and the lungs lavaged. Total and differential leukocyte counts are then obtained for the bronchoalveolar lavage fluid and the dose of test compound giving a 50% reduction in eosinophil accumulation (ED₅₀) is determined (Sanjar et al. 1992).

References:

[0087]

- 5 Asako H, Wolf, RE, Granger, DN (1993), Gastroenterology 104, pp 31-37;
 Burkey TH, Webster, RO, (1993), Biochem. Biophys Acta 1175, pp 312-318;
 Castanon MJ, Spevak W, (1994), Biochem. Biophys Res. Commun. 198, pp 626-631;
 Cronstein BN, Kramer SB, Weissmann G, Hirschhorn R, (1983), Trans. Assoc. Am. Physicians 96, pp 384-91;
 10 Cronstein BN, Kramer SB, Rosenstein ED, Weissmann G, Hirschhorn R, (1985), Ann N.Y. Acad. Sci. 451, pp
 291-301;
 Cronstein BN, Naime D, Ostad E, (1993), J. Clin. Invest. 92, pp 2675-82;
 Cronstein BN, Naime D, Ostad E, (1994), Adv. Exp. Med. Biol., 370, pp 411-6;
 Cronstein BN, (1994), J. Appl. Physiol. 76, pp 5-13;
 Dianzani C, Brunelleschi S, Viano I, Fantozzi R, (1994), Eur. J. Pharmacol 263, pp 223-226;
 15 Elliot KRF, Leonard EJ, (1989), FEBS Letters 254, pp 94-98;
 Green PG, Basbaum AI, Helms C, Levine JD, (1991), Proc. Natl. Acad Sci. 88, pp 4162-4165;
 Hirschhorn R, (1993), Pediatr. Res 33, pp S35-41;
 Kohno Y; Xiao-duo J; Mawhorter SD; Koshiha M; Jacobson KA. (1996). Blood 88 p3569-3574.
 Peachell PT, Lichtenstein LM, Schleimer RP, (1989), Biochem Pharmacol 38, pp 1717-1725;
 20 Richter J, (1992), J. Leukocyte Biol. 51, pp 270-275;
 Rosengren S, Bong GW, Firestein GS, (1995), J. Immunol. 154, pp 5444-5451;
 Sanjar S, McCabe PJ, Fattah D, Humbles AA, Pole SM, (1992), Am. Rev. Respir. Dis. 145, A40;
 Skubitz KM, Wickman NW, Hammerschmidt DE, (1988), Blood 72, pp 29-33
 Van Schaick EA; Jacobson KA; Kim HO; Ijzerman AP; Danhof M. (1996) Eur J Pharmacol 308 p311-314.
 25 Wood KV. (1995) Curr Opinion Biotechnology 6 p50-58.

[0088] The invention is illustrated by the following Examples:

Examples

30

General experimental details

[0089] Where products were purified by column chromatography, 'flash silica' refers to silica gel for chromatography, 0.040 to 0.063mm mesh (e.g. Merck Art 9385), where column elution was accelerated by an applied pressure of nitrogen at up to 5 p.s.i. Where thin layer chromatography (TLC) has been used it refers to silica gel TLC using 5 x 10 cm silica gel 60 F₂₅₄ plates (e.g. Merck Art 5719).

35

[0090] Where products were purified by preparative HPLC, this was carried out on a C18-reverse-phase column (1" Dynamax), eluting with a gradient of acetonitrile (containing 0.1% trifluoroacetic acid) in water (containing 0.1% trifluoroacetic acid) and the compounds isolated as their trifluoroacetate salts unless otherwise specified.

40

Standard Automated Preparative HPLC column, conditions & eluent

[0091] Automated preparative high performance liquid chromatography (autoprep. HPLC) was carried out using a Supelco ABZ+ 5µm 100mmx22mm i.d. column eluted with a mixture of solvents consisting of i) 0.1% formic acid in water and ii) 0.05% formic acid in acetonitrile, the eluent being expressed as the percentage of ii) in the solvent mixture, at a flow rate of 4ml per minute. Unless otherwise stated the eluent was used as a gradient of 5-95 % over 20 minutes.

45

LC/MS System

[0092] The Liquid Chromatography Mass Spectroscopy (LC/MS) systems used:
 LC/MS System A - A Supelco ABZ+, 3.3cm x 4.6mm i.d. column eluting with solvents: A - 0.1%v/v formic acid + 0.077% w/v ammonium acetate in water, and B - 95:5 acetonitrile:water + 0.05% v/v formic acid. The following gradient protocol was used: 100% A for 0.7 mins; A+B mixtures, gradient profile 0- 100% B over 3.5mins; hold at 100% B for 3.5mins; return to 0% B over 0.3mins. Positive and negative electrospray ionization was employed.
 50
 LC/MS System B - A Supelco ABZ+, 5cm x 2.1mm i.d. column eluting with solvents: A - 0.1%v/v formic acid + 0.077% w/v ammonium acetate in water, and B - 95:5 acetonitrile:water + 0.05% v/v formic acid. The following gradient protocol was used: 0 - 100% B over 3.5mins; hold at 100% B for 1.50mins; return to 0% B over 0.50mins. Positive and negative electrospray ionization was employed.
 55

EP 1 090 022 B1

LC/MS System C - A Supelco ABZ+, 3.3cm x 4.6mm i.d. column eluting with solvents: A - 0.1%v/v formic acid + 10mmol ammonium acetate in water, and B - 95:5 acetonitrile:water + 0.05% v/v formic acid. The following gradient protocol was used: 100% A for 0.7 mins; A+B mixtures, gradient profile 0- 100% B over 3.7mins; hold at 100% B for 0.9mins; return to 0% B over 0.2mins. Positive and negative electrospray ionization was employed.

5

Example of Novel Process

Intermediate A: 2-Bromohypoxanthine

10 **[0093]** This compound is prepared from 2-thioxanthine** through oxidation of the mercapto group by bromine and *in situ* displacement by hydrobromine. For reference on the oxidation and displacement, see Beaman, A.G.; Gerster, J.F.; Robins, R.K, J. *Org. Chem*, 1962, 27, 986.1.

**Elion, G.B.; Lange, H.L., Hitchings, G.H., J. *Am. Chem. Soc.*, 1956, 78, 217.

15 Intermediate B: 2-[(1S)-1-benzyl-2-hydroxyethyl]amino-1,9-dihydro-6H-purin-6-one

[0094] A mixture of 10.0 g (46.5 mmol) of Intermediate A and 14.1 g (93.0 mmol) of L-phenylalaninol in 30 mL of 2-methoxyethanol in an 100 mL round bottom flask was heated to reflux overnight (>12 h). The mixture was cooled to ambient temperature and gave rise to precipitation of solids. Additional precipitation was generated by addition of 150 mL of water. After being stirred for 1 h, the suspension was filtered and the filter cake was washed with 50 mL of water and dried under vacuum to afford 7.40 g (56%) of title compound as a yellow solid. The product was a mixture of two tautomers by ¹H NMR. The combined filtrate and washing were allowed to stand at ambient temperature for two days. The resultant solids were filtered and dried to give 1.12 g (8.4%) of product as a white solid. The total yield was 64%. TLC (silica gel, 50% MeOH in CH₂Cl₂, 254 nm visualisation): Rf 0.9; 2-bromohypoxanthine Rf 0.6. MS (ES⁻): m/z 284 (M-1).

20 ¹H NMR (major tautomer, 300 Mhz) δ 2.76-2.98 (m, 2H), 3.49 (m, 2H), 4.09 (br s, 1H), 5.04 (2, 1H), 6.36 (d, J = 8.1 Hz, 1H), 7.19-7.38 (m, 5H), 7.66 (s, 1H), 10.5 (s, 1H), 12.5 (s, 1H).

25

30 Intermediate C: (2S)-2-[(9-acetyl-6-oxo-6,9-dihydro-1H-purin-2-yl)amino]-3-phenylpropyl acetate

[0095] To a suspension of 500 mg (1.75 mmol) of Intermediate B in 3.5 mL of DMF in a 25 mL round bottom flask was successively added 0.66 mL (7.02 mmol) of acetic anhydride, 5 mg (catalytic) of N,N-dimethylpyridine and 0.98 (7.02 mmol) of triethylamine at ambient temperature. The mixture was stirred at ambient temperature overnight. After being quenched with 15 mL of water and stirred for 2 h, the suspension was filtered and the filtering cake was washed with 10 mL of water, dried under vacuum at 70-100 °C to afford 470 mg (73%) of title compound as an off-white powder. TLC (silica gel, 10% MeOH in CH₂Cl₂, 254 nm visualisation): Rf 0.45. MS (ES⁻): m/z 368 (M-1)⁻, 326 (M-1-Ac)⁻, ¹H NMR (300 Mhz) δ 1.96 (s, 3H), 2.80 (s, 3H), 2.90 (m, 2H), 4.16 (m, 2H), 4.37 (m, 1H), 6.70 (br s, 1H), 7.16-7.31 (m, 5H), 8.16 (s, 1H), 10.8 (s, 1H).

35

40 Intermediate D: (2S)-2-[(6-chloro-9H-purin-2-yl)amino]-3-phenylpropyl acetate

[0096] To 16.7 mL (179 mmol) of phosphorus oxychloride in a 100 mL round bottom flask was added 2.27 mL (17.9mmol) of N,N-dimethylaniline at ambient temperature. The mixture was stirred for 10 min; then 4.40 g (11.9 mmol) of 6-hydroxypurine Intermediate C was added in two equal portions over 15 min. The mixture was heated at reflux for 15 min. After being cooled to ambient temperature, the mixture was slowly added to 550 mL of ice water with stirring. The aqueous mixture was neutralised to pH 3.5 by addition of solid NaOAc and extracted with CH₂Cl₂ (3X). The combined organic layers were washed with aqueous NaHCO₃ (2X), dried over anhydrous Na₂SO₄ and concentrated under vacuum. The resultant brown oil was chromatographed on silica gel. Elution with 5-10% MeOH in CH₂Cl₂ afforded 3.08 g (75%) of title compound as a brown solid. TLC (silica gel, 10% MeOH in CH₂Cl₂, 254 nm visualisation): Rf 0.50. MS (ES⁻): m/z 344 (M-1)⁻, 346 (M-1, isotope)⁻. ¹H NMR (300 MHz) δ 2.15 (s, 3H), 2.98 (m, 2H), 4.08-4.35 (m, 2H), 4.49 (m, 1H), 7.26-7.53 (m, 5H), 7.64 (br s, 1H), 8.25 (s, 1H), 13.1 (s, 1H).

45

50

Intermediate E: (2S)-2-[(6-amino-9H-purin-2-yl)amino]-3-phenyl-1-propanol

55 **[0097]** A 200 mL glass liner in a Parr pressure reactor was charged with 288 mg (0.834 mmol) of Intermediate D and 25 mL of 2M NH₃ in methanol. The reactor was sealed and heated at 90-100 °C for 16 h. After being cooled to ambient temperature, the solvent and excess reagent were evaporated under vacuum. Despite incomplete reaction as indicated by TLC, the resultant oil was chromatographed on silica gel. Elution with 10-15% MeOH in CH₂Cl₂ afforded 48 mg

(20%) of title compound as a solid. Further elution afforded 156 mg (62%) of recovered starting material as the deacetylated form. TLC (silica gel, 10% MeOH in CH₂Cl₂, 254 nm visualisation): R_f 0.22. ¹H NMR (300 MHz) δ 2.61-2.80 (m, 2H), 3.30-3.45 (m, 2H), 3.95 (m, 1H), 4.70 (s, 1H), 5.65 (d, J = 8.0 Hz), 6.41 (s, 2H), 6.99-7.26 (m, 5H), 7.52 (s, 1H), 12.1 (s, 1H).

5 Intermediate F: (3aS,4S,6R,6aR)-Methoxy-2,2-dimethyl-tetrahydro-furo[3,4-d][1,3]dioxole-4-carboxylic acid

[0098] To a 1L three neck round bottom flask equipped with an addition funnel, thermocouple probe and nitrogen inlet was added D-ribose (50 g) and acetone (400 mL). The mixture was cooled to -5 °C and then 2,2-dimethoxypropane (100 mL) followed by perchloric acid (20 mL) were added. The reaction mixture was allowed to warm to room temperature and then stirred for a brief period. Methanol (70 mL) was added and the reaction mixture was stirred overnight. The reaction solution was cooled to ca. 5 °C and ca. 95 mL of 30% sodium carbonate was added dropwise. The mixture was allowed to warm then filtered.

15 [0099] The resulting cake was washed with ethyl acetate (50 mL). The filtrate was concentrated *in vacuo* at ca. 200 mbar until 250 mL of residual volume remained, diluted with ethyl acetate (200 mL) and reconcentrated to a residual volume of 170 mL. Ethyl acetate (200 mL) and water (200 mL) were added and the phases were mixed and separated. The aqueous phase was washed twice with ethyl acetate (200 mL) and the layers were separated. The combined organic extracts were concentrated to a residual volume of 200 mL and rediluted with ethyl acetate (200 mL) to provide an ethyl acetate solution of 6R-methoxy-2,2-dimethyl-tetrahydro-(3aR,6aR)-furo[3,4-d][1,3]dioxol-4R-yl)-methanol.

20 [0100] To a 2L three neck round bottom flask was added the ethyl acetate solution of 6R-methoxy-2,2-dimethyl-tetrahydro-(3aR,6aR)-furo[3,4-d][1,3]dioxol-4R-yl)-methanol, 6% sodium bicarbonate (158 mL) potassium bromide (2.3 g), and TEMPO (0.167 g). The reaction mixture was cooled to -7 °C. Meanwhile, sodium bicarbonate (6.8 g) was dissolved into 10-13% sodium hypochlorite (400.5 mL). The bleach solution was added dropwise over ca. 40 minutes, keeping the temperature below 15 °C. The reaction mixture was stirred for ca. 2 hours and 10% aqueous sodium sulfite solution (47 mL) was added. The reaction mixture was stirred for 15 minutes, the phases separated and the aqueous phase adjusted to pH 2 with 4M HCl and extracted twice with ethyl acetate (225 mL). The ethyl acetate extracts were concentrated *in vacuo* to provide a white residue which was triturated with cyclohexane (90 mL). The solids were filtered and dried *in vacuo* at 45 °C to provide title product (33.6 g) (46% yield re D-ribose) as a white solid: m.p. 126-129 °C.

30 Intermediate G: (3aS,4S,6R,6aR)-6-Methoxy-2,2-dimethyl-tetrahydro-furo[3,4-d][1,3]dioxole-4-carboxylic acid amide

[0101] To a 500 mL three neck round bottom flask was added Intermediate F (20 g) and ethyl acetate (160 mL) followed by thionyl chloride (9.4 mL). The reaction solution was warmed at 50 °C for 2 hours. Gaseous ammonia (16 g) was added at such a rate that the temperature remains between 40-60 °C. Water (120 mL) was added. The layers were separated and the aqueous layer was washed twice with ethyl acetate (80 mL). The combined organic washes were concentrated *in vacuo* to dryness. The residue was triturated with cyclohexane (40 mL) and the solids filtered. The cake was washed with cyclohexane (40 mL) and the solids dried *in vacuo* at 45 °C to provide the title product (16.7 g) (83.9% yield) as a light tan solid: m.p. = 134-136 °C; TLC (95/5 chloroform/methanol/~5 drops TFA per 50 mL/phosphomolybdic acid spray) R_f=0.49.

40 Intermediate H: (3aS,4S,6R,6aR)-6-Methoxy-2,2-dimethyl-tetrahydro-furo[3,4-d][1,3]dioxole-4-carbonitrile

[0102] To a 22L three neck round bottom flask was added Intermediate G (643 g), ethyl acetate (7.72L), N,N-dimethylformamide (1.26L), and triethylamine (2.15L). The reaction solution was cooled to ca. 0 °C and of then phosphorus oxychloride (1.38L) was added at such a rate that the temperature was maintained below 25 °C. The reaction was stirred for one and one-half hours. Aqueous potassium hydrogen carbonate (20%, 6.5L) was added dropwise maintaining the temperature at or below 20 °C. The layers were separated and the aqueous layer re-extracted with ethyl acetate (3.5 L). The combined organic layers were washed twice with 20% potassium hydrogen carbonate (3.5L) and concentrated to a residual volume of ca. 1L. Activated carbon (15 grams) was added to the thin oil and the mixture was filtered through celite (80 g). The cake was washed with ethyl acetate (100 mL). The filtrate was concentrated *in vacuo* to provide title product (519 g) (88% yield) as a reddish-orange oil: TLC (1:1 Ethyl acetate/cyclohexane; phosphomolybdic acid reagent development) R_f = 0.73.

55 Intermediate I: 5-(6R-Methoxy-2,2-dimethyl-tetrahydro-(3aR,6aR)-furo[3,4 d][1,3]dioxol-4R-yl)-1H-tetrazole

[0103] To a 3L three neck round bottom flask was added Intermediate H (200 g), toluene (2L), azidotrimethylsilane (332 mL) and dibutyltin oxide (24.9 g). The reaction mixture was heated to 60 °C for 15 hours. The reaction mixture was concentrated *in vacuo* to a residual volume of ca. 300 mL. Toluene (1L) was added and the solution was recon-

EP 1 090 022 B1

centrated to a residual volume of ca. 470 mL. Toluene (400 mL) and water (19.8 mL) were added and the mixture was stirred at room temperature for approximately 2 hours. The mixture was concentrated to provide ca. 250 mL of residue. The residue was dissolved in toluene (800 mL) with warming then was allowed to cool to room temperature and was stirred for >3 days. The solids are filtered and washed twice with toluene (250 mL). The product was dried *in vacuo* to provide title product (135 g) (55% yield) as a white solid: mp 130°C.

Intermediate J: 2-Ethyl-5-(6R-methoxy-2,2-dimethyl-tetrahydro-(3aR,6aR)-furo [3,4-d][1,3]dioxol-4R-yl)-2H-tetrazole

[0104] To a 1L three neck round bottom flask was added Intermediate I (31.8 g), potassium carbonate (12.7 g) and acetone (238 mL). Ethyl iodide (14.1 mL) of was added *via* syringe and the reaction mixture was warmed at 42 °C for 2.5-3 hours. The reaction mixture was allowed to cool to room temperature and then cyclohexane (238 mL) was added. The resulting precipitate was filtered and the cake was washed three times with cyclohexane (65 mL). The filtrate was concentrated to a residual volume of 195 mL and then rediluted with cyclohexane (238 mL). The cyclohexane solution was cooled at 0-5 °C for 3 days and the resulting crystalline solid (N1 alkylation product) was filtered and washed three times with cyclohexane (65 mL). The combined filtrates was concentrated *in vacuo* to provide intermediate grade title product as an oil. The oil was dissolved in cyclohexane (200 mL) at 60 °C and the solution allowed to cool to room temperature and filtered. The resulting crystalline solid was filtered and washed three times with cyclohexane (65 mL). The combined filtrate was concentrated to provide title product as a yellow oil: TLC (1:1 Ethyl acetate/hexanes; phosphomolybdic acid reagent visualisation) *r_f* = 0.68.

Intermediate K: rel-Acetic acid 4R,5-diacetoxy-2R-(2-ethyl-2H-tetrazol-5-yl)-tetrahydro-furan-3R-yl ester

[0105] To a round bottom flask was added Intermediate J (5.0 g). A solution of acetyl chloride (0.73 g) in methanol (50 mL) was added to the flask and the reaction solution was heated to reflux at 300 mbar pressure. The reaction was distilled over an 8-9 hour period and methanol (135 mL) was added portionwise during this time to replenish the reaction volume. The reaction mixture was allowed to cool to room temperature and pyridine (15 mL) was added. The mixture was concentrated *in vacuo* and rediluted with pyridine. Ethyl acetate (25 mL) and acetic anhydride (6.6 g) were added to the pyridine solution and the resulting mixture stirred overnight at room temperature. The reaction mixture was cooled to 5-10 °C and approximately 2M sulfuric acid (ca 45 mL) was added dropwise over 20 minutes while maintaining the temperature below 10 °C. The layers were separated and the organic layer was washed with approximately 0.7M sulfuric acid (ca 25 mL). The organic layer was washed with sat. sodium bicarbonate and brine and then concentrated *in vacuo* to provide a pale yellow oil that was dissolved in 50 mL of ethyl acetate. Acetic anhydride (3.04 g) and of concentrated sulfuric acid (0.65 g) were added and the reaction mixture was warmed to 50 °C for ca. 3.5 hours. The reaction was quenched with saturated sodium bicarbonate solution (25 mL). The organic layer was concentrated *in vacuo* to provide title product ((5.1 g) (82% yield) as a yellow oil: TLC (1:1 Ethyl acetate/hexanes; phosphomolybdic acid reagent visualisation) *r_f* = 0.44.

Intermediate L: (2R,3S,4S,5R)-4-(acetyloxy)-2-(6-amino-2-[(1S)-1-benzyl-2-hydroxyethyl]amino-9H-purin-9-yl)-5-(2-ethyl-2H-1,2,3,4-tetraazol-5-yl)tetrahydro-3-furanyl acetate

[0106] To a mixture of 65 mg (0.19 mmol) of Intermediate K and 45 mg (0.16 mmol) of Intermediate E in 2.5 mL of MeCN in a 10 mL round bottom flask was successively added 88mL (0.36 mmol) of N,O-bis(trimethylsilyl)acetamide and 34 mL (0.19 mmol) of trimethylsilyl trifluoromethanesulfonate at ambient temperature. The yellow suspension was heated to reflux and became a dark yellow solution. After being heated at reflux for 5 h, the mixture was cooled to ambient temperature, quenched with 2 mL of 10% KHCO₃ and extracted with CH₂Cl₂ (2X8 mL). The combined organic layers were washed with 10% brine (3 mL) and evaporated under vacuum. The resultant yellow foam was chromatographed on silica gel. Elution with 5% MeOH in CH₂Cl₂ afforded 70 mg (78%) of title compound as a solid. Despite low purity, the material was used for the next step without further purification. TLC (silica gel, 10% MeOH in CH₂Cl₂, 254 nm visualisation): *R_f* 0.54.

Example A: (2R,3R,4S,5R)-2-[6-Amino-2-(1S-hydroxymethyl-2-phenyl-ethylamino)-purin-9-yl]-5-(2-ethyl-2H-tetrazol-5-yl)-tetrahydro-furan-3,4-diol

[0107] A mixture of 70 mg (0.12 mmol) of Intermediate L and 20 mg (0.15 mmol) of anhydrous K₂CO₃ in 5 mL of methanol was stirred at ambient temperature for 2.5 h. The mixture was evaporated to near dryness, diluted with 2 mL of water and extracted with EtOAc (3X5 mL). The combined organic layers were dried with Na₂SO₄ and evaporated under vacuum. The resultant crude product was chromatographed on silica gel. Elution with 10% MeOH in CH₂Cl₂ afforded 24.5 mg (41%) of title compound as a solid.

TLC (silica gel, 10% MeOH in CH₂Cl₂, 254 nm visualisation): Rf 0.35.

Novel Examples

5 Intermediates

Intermediate 1: 2-Benzyl-5-(6-methoxy-2,2-dimethyl-tetrahydro-(3aR,6aR)-furo[3,4-d][1,3]dioxol-4R-yl)-2H-tetrazole

10 **[0108]** To a stirred solution of Intermediate 10 (10g, 41.3mM) in dimethylformamide(50ml) under nitrogen was added potassium carbonate(5.7g, 41.3mM) followed by benzyl bromide (6ml,49.6mM). The mixture was stirred at room temperature 18hours. Water(100ml) was added and the mixture was extracted with ethyl acetate(2x100ml). The organic phases were combined, washed with water, brine and dried with magnesium sulphate. The residue, obtained after evaporation under reduced pressure was purified by column chromatography on flash silica eluted with 20%ethyl acetate/cyclohexane yielding the title compound as a waxy solid (2.98g).

15 TLC SiO₂ (20% ethyl acetate in cyclohexane) Rf = 0.45

Intermediate 2: Acetic acid 4R,5-diacetoxy-2R-(2-benzyl-2H-tetrazol-5-yl)-tetrahydro-furan-3R-yl ester

20 **[0109]** To Intermediate 1 (2.98g,8.9mM) was added a mixture of TFA/Water(40ml/4ml) at room temperature and stirred for 1 hour. The mixture was evaporated under reduced pressure, azeotroped with toluene (3x20ml). The residue was taken up into dichloromethane (100ml) and dimethylaminopyridine(catalytic) and triethylamine(40ml,356mM) was added. The mixture was cooled to 0 °C and acetic anhydride (17ml, 166mM) was added dropwise over 15m.. The mixture was allowed to warm to room temperature and stirred for 16h.. The mixture was evaporated under reduced pressure and purified by column chromatography on flash silica eluted with 50% ethyl acetate/cyclohexane yielding the title compound as a as an oil (2.44g).

25 LC/MS System A Rt = 3.39min, *m/z* = 279 (MH⁺).

Intermediate 3: Acetic acid 4R-acetoxy-5R-(2-benzyl-2H-tetrazol-5-yl)-2R-(2,6-dichloro-purin-9-yl)-tetrahydro-furan-3R-yl ester

30 **[0110]** To a stirred solution of Intermediate 2 (2.43g, 6mM) under nitrogen in acetonitrile (18ml) was added 1,8-diazobicyclo[5,4,0]undec-7ene(1.35ml, 9mM) followed by 2,6-dichloropurine(1.5g). The mixture was cooled to 0°C and trimethylsilyltriflate (1.87ml, 10.2mM) was added dropwise over 15m., allowed to warm to 20 C and stirred for 38 h. The reaction mixture was quenched with aqueous saturated sodium hydrogen carbonate (35ml) and extracted with ethyl acetate (3x 50ml). The organics were combined and washed with water (50ml), dried with magnesium sulphate and evaporated under reduced pressure. The residue obtained was purified by column chromatography on flash silica eluted with 30% ethyl acetate/cyclohexane yielding the title compound as a as an oil (2.36g). LC/MS System B Rt = 3.43 min, *m/z* = 535 (MH⁺).

35

Intermediate 4: Acetic acid 4R-acetoxy-5R-(2-benzyl-2H-tetrazol-5-yl)-2R-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-tetrahydro-furan-3R-yl ester

40 **[0111]** To a stirred solution of Intermediate 3 (2.3g, 4.3mM) under nitrogen in isopropanol (40ml) was added diisopropylethylamine(1.12ml, 6.5mM) followed by diphenylethyl amine(1.02g, 5.2mM), the resultant mixture was heated to 50°C for 18 hours. The reaction mixture was evaporated under reduced pressure and the residue obtained purified by column chromatography on flash silica eluted with 50% ethyl acetate/cyclohexane yielding the title compound as a as an off white solid (2.9g). LC/MS System B Rt = 3.68 min, *m/z* = 694 (MH⁺).

45

Intermediate 5: (2R,3S,4R,5R)-2-(2-Benzyl-2H-tetrazol-5-yl)-5-[6-(2,2-diphenyl-ethylamino)-2-(2-piperidin-1-yl-ethylamino)-purin-9-yl]-tetrahydro-furan-3,4-diol

50 **[0112]** A solution of Intermediate 4 (2.9g, 4.2mM) and 2-piperidinoethylamine (3ml, 20.9mM) in dimethylsulfoxide (1ml) under nitrogen was heated to 90°C for 72 h.. The mixture was allowed to cool and purified by column chromatography on flash silica eluted with 20% methanol,79% chloroform and 1% ammonia yielding the title compound as a as an oil (1.6g). LC/MS System A Rt = 3.86 min, *m/z* = 702 (MH⁺).

55

Intermediate 6: (2R,3S,4R,5R)-2-(2H-tetrazol-5-yl)-5-[6-(2,2-diphenyl-ethylamino)-2-(2-piperidin-1-yl-ethylamino)-purin-9-yl]-tetrahydro-furan-3,4-diol

[0113] To 10% palladium on carbon (1.6g) under nitrogen was added a solution of Intermediate 5 (1.67g, 2.38mM) in ethanol (50ml) followed by ammonium formate (0.72g, 11.9mM). The mixture was heated to 50 °C for 4 h., filtered through a pad of Harborlite®. The filtrate was evaporated under reduced pressure to yield the title compound as a pale yellow solid (1.45g).

LC/MS System A Rt = 3.66 min, *m/z* = 612 (MH⁺).

Intermediate 7: (3aS,4S,6R,6aR)-Methoxy-2,2-dimethyl-tetrahydro-furo[3,4-d][1,3]dioxole-4-carboxylic acid

[0114] To a 1L three neck round bottom flask equipped with an addition funnel, thermocouple probe and nitrogen inlet was added D-ribose (50 g) and acetone (400 mL). The mixture was cooled to -5°C and then 2,2-dimethoxypropane (100 mL) followed by perchloric acid (20 mL) were added. The reaction mixture was allowed to warm to room temperature and then stirred for a brief period. Methanol (70 mL) was added and the reaction mixture was stirred overnight. The reaction solution was cooled to ca. 5°C and ca. 95 mL of 30% sodium carbonate was added dropwise. The mixture was allowed to warm then filtered. The resulting cake was washed with ethyl acetate (50 mL). The filtrate was concentrated *in vacuo* at ca. 200 mbar until 250 mL of residual volume remained, diluted with ethyl acetate (200 mL) and reconcentrated to a residual volume of 170 mL. Ethyl acetate (200 mL) and water (200 mL) were added and the phases were mixed and separated. The aqueous phase was washed twice with ethyl acetate (200 mL) and the layers were separated. The combined organic extracts were concentrated to a residual volume of 200 mL and rediluted with ethyl acetate (200 mL) to provide an ethyl acetate solution of 6R-methoxy-2,2-dimethyl-tetrahydro-(3aR,6aR)-furo[3,4-d][1,3]dioxol-4R-yl)-methanol.

[0115] To a 2L three neck round bottom flask was added the ethyl acetate solution of 6R-methoxy-2,2-dimethyl-tetrahydro-(3aR,6aR)-furo[3,4-d][1,3]dioxol-4R-yl)-methanol, 6% sodium bicarbonate (158 mL) potassium bromide (2.3 g), and TEMPO (0.167 g). The reaction mixture was cooled to -7°C. Meanwhile, sodium bicarbonate (6.8 g) was dissolved into 10-13% sodium hypochlorite (400.5 mL). The bleach solution was added dropwise over ca. 40 minutes, keeping the temperature below 15 °C. The reaction mixture was stirred for ca. 2 hours and 10% aqueous sodium sulfite solution (47 mL) was added. The reaction mixture was stirred for 15 minutes, the phases separated and the aqueous phase adjusted to pH 2 with 4M HCl and extracted twice with ethyl acetate (225 mL). The ethyl acetate extracts were concentrated *in vacuo* to provide a white residue which was triturated with cyclohexane (90 mL). The solids were filtered and dried *in vacuo* at 45 °C to provide title product (33.6 g) (46% yield re D-ribose) as a white solid: m.p. 126-129 °C.

Intermediate 8 (3aS,4S,6R,6aR)-6-Methoxy-2,2-dimethyl-tetrahydro-furo[3,4-d][1,3]dioxole-4-carboxylic acid amide

[0116] To a 500 mL three neck round bottom flask was added Intermediate 1 (20 g) and ethyl acetate (160 mL) followed by thionyl chloride (9.4 mL). The reaction solution was warmed at 50 °C for 2 hours. Gaseous ammonia (16 g) was added at such a rate that the temperature remains between 40-60 °C. Water (120 mL) was added. The layers were separated and the aqueous layer was washed twice with ethyl acetate (80 mL). The combined organic washes were concentrated *in vacuo* to dryness. The residue was triturated with cyclohexane (40 mL) and the solids filtered. The cake was washed with cyclohexane (40 mL) and the solids dried *in vacuo* at 45 °C to provide the title product (16.7 g) (83.9% yield) as a light tan solid: m.p. = 134-136 °C; TLC (95/5 chloroform/methanol/~5 drops TFA per 50 mL/phosphomolybdic acid spray) *rf*=0.49.

Intermediate 9: (3aS,4S,6R,6aR)-6-Methoxy-2,2-dimethyl-tetrahydro-furo[3,4-d][1,3]dioxole-4-carbonitrile

[0117] To a 22L three neck round bottom flask was added Intermediate 2 (643 g), ethyl acetate (7.72L), N,N-dimethylformamide (1.26L), and triethylamine (2.15L). The reaction solution was cooled to ca. 0 °C and of then phosphorus oxychloride (1.38L) was added at such a rate that the temperature was maintained below 25 °C. The reaction was stirred for one and one-half hours. Aqueous potassium hydrogen carbonate (20%, 6.5L) was added dropwise maintaining the temperature at or below 20 °C. The layers were separated and the aqueous layer re-extracted with ethyl acetate (3.5 L). The combined organic layers were washed twice with 20% potassium hydrogen carbonate (3.5L) and concentrated to a residual volume of ca. 1L. Activated carbon (15 grams) was added to the thin oil and the mixture was filtered through celite (80 g). The cake was washed with ethyl acetate (100 mL). The filtrate was concentrated *in vacuo* to provide title product (519 g) (88% yield) as a reddish-orange oil: TLC (1:1 Ethyl acetate/cyclohexane; phosphomolybdic acid reagent development) *rf* = 0.73.

EP 1 090 022 B1

Intermediate 10: 5-(6R-Methoxy-2,2-dimethyl-tetrahydro-(3aR,6aR)-furo[3,4-d][1,3]dioxol-4R-yl)-1H-tetrazole

[0118] To a 3L three neck round bottom flask was added Intermediate 3 (200 g), toluene (2L), azidotrimethylsilane (332 mL) and dibutyltin oxide (24.9 g). The reaction mixture was heated to 60 °C for 15 hours. The reaction mixture was concentrated *in vacuo* to a residual volume of ca. 300 mL. Toluene (1L) was added and the solution was re-concentrated to a residual volume of ca. 470 mL.

[0119] Toluene (400 mL) and water (19.8 mL) were added and the mixture was stirred at room temperature for approximately 2 hours. The mixture was concentrated to provide ca. 250 mL of residue. The residue was dissolved in toluene (800 mL) with warming then was allowed to cool to room temperature and was stirred for >3 days. The solids are filtered and washed twice with toluene (250 mL). The product was dried *in vacuo* to provide title product (135 g) (55% yield) as a white solid: mp 130 °C.

Examples

Example 1: 2R,3R,4S,5R)-2-[6-(2,2-Diphenyl-ethylamino)-2-(2-piperidin-1-yl-ethylamino)-purin-9-yl]-5-[2-(3-hydroxy-propyl)-2H-tetrazol-5-yl]-tetrahydro-furan-3,4-diol bis(trifluoroacetate)

[0120] To a solution of Intermediate 6 (0.06g, 0.098mM) in dimethylformamide(1ml) was added potassium carbonate (0.023g, 0.167mM) followed by 3-bromo-propanol (0.013ml, 0.147mM) in a sealed vial (eg Reacti-vial™), the reaction mixture stirred for 18h.. The mixture was filtered and the filtrate was evaporated under reduced pressure and purified using preparative HPLC, (using a Capital column ODS2-IK5 15mm x 20mm i.d, on a 30min gradient of 5% to 95% acetonitrile containing 0.1% trifluoroacetic acid) to yield after freeze drying the title compound as a white solid (0.022g) LC/MS System A Rt = 3.59min, *m/z* = 670 (MM⁺).

Example 2: 2R,3R,4S,5R)-2-[6-(2,2-Diphenyl-ethylamino)-2-(2-piperidin-1-yl-ethylamino)-purin-9-yl]-5-(2-propyl-2H-tetrazol-5-yl)-tetrahydro-furan-3,4-diol bis(trifluoroacetate)

[0121] Example 2 was prepared in an analogous manner to Example 1 using 1-Bromopropane (0.013ml, 0.147mM) to yield after freeze drying the title compound as a white solid (0.026g). LC/MS System A Rt = 3.76min, *m/z* = 654 (MH⁺).

Example 3: Acetic acid 2-(5-{5R-[6-(2,2-diphenyl-ethylamino)-2-(2-piperidin-1-yl-ethylamino)-purin-9-yl]-3S,4R-dihydroxy-tetrahydro-furan-2R-yl}-tetrazol-2-yl)-ethyl ester bis(trifluoroacetate)

[0122] Example 3 was prepared in an analogous manner to Example 1 using 2-bromoethyl acetate (0.017ml, 0.147mM) to yield after freeze drying the title compound as a white solid (0.029g). LC/MS System A Rt = 3.68min, *m/z* = 698 (MH⁺).

Example 4: (2R,3S,4R,5R)-2-(2-Cyclopropylmethyl-2H-tetrazol-5-yl)-5-[6-(2,2-diphenyl-ethylamino)-2-(2-piperidin-1-yl-ethylamino)-purin-9-yl]-tetrahydro-furan-3,4-diol bis(trifluoroacetate)

[0123] Example 4 was prepared in an analogous manner to Example 1 using (Bromomethyl)cyclopropane (0.0165ml, 0.147mM) to yield after freeze drying the title compound as a white solid (0.023g). LC/MS System A = 3.74min, *m/z* = 666 (MH⁺).

Example 5: (2R,3R,4S,5R)-2-[6-(2,2-Diphenyl-ethylamino)-2-(2-piperidin-1-yl-ethylamino)-purin-9-yl]-5-[2-(2-hydroxy-ethyl)-2H-tetrazol-5-yl]-tetrahydro-furan-3,4-diol bis(trifluoroacetate)

[0124] To Example 3 (0.01g) in methanol under nitrogen was added a solution of sodium methoxide (0.005ml), the mixture was stirred at room temperature for 18 hours. The reaction mixture was evaporated under reduced pressure purified using preparative HPLC, (using a Capital column ODS2-IK5 15mm x 20mm i.d, on a 30min gradient of 5% to 95% acetonitrile) yielding the title compound as a gum (0.006g). LC/MS System C Rt = 2.54min, *m/z* = 656 (MH⁺).

Example 6: (2R,3S,4R,5R)-2-[2-(2-Chloro-ethyl)-2H-tetrazol-5-yl]-5-[6-(2,2-diphenyl-ethylamino)-2-(2-piperidin-1-yl-ethylamino)-purin-9-yl]-tetrahydro-furan-3,4-diol bis(trifluoroacetate)

[0125] Example 6 was prepared in an analogous manner to Example 1 using 1-Bromo-2-chloroethane (0.012ml, 0.147mM) to yield after freeze drying the title compound as a white solid (0.004g). LC/MS System A = 3.79min, *m/z* = 674 (MH⁺).

EP 1 090 022 B1

Example 7: (2R,3S,4R,5R)-2-(2-Cyclobutyl-2H-tetrazol-5-yl)-5-[6-(2,2-diphenyl-ethylamino)-2-(2-piperidin-1-yl-ethylamino)-purin-9-yl]-tetrahydro-furan-3,4-diol bis (trifluoroacetate)

[0126] Example 7 was prepared in an analogous manner to Example 1 using Bromocyclobutane (0.014ml, 0.147mM) to yield after freeze drying the title compound as a white solid (0.008g). LC/MS System C = 2.76min, m/z = 666 (MH⁺).

Example 8: (2R,3S,4R,5R)-2-(2-Allyl-2H-tetrazol-5-yl)-5-[6-(2,2-diphenyl-ethylamino)-2-(2-piperidin-1-yl-ethylamino)-purin-9-yl]-tetrahydro-furan-3,4-diol bis(trifluoroacetate)

[0127] Example 8 was prepared in an analogous manner to Example 1 using allyl bromide (0.014ml, 0.147 mM) at 0 °C and the mixture was then stirred at 0 °C for 3 h. to yield the title compound as a clear gum (0.004g). LC/MS System C R_t = 2.68min, m/z = 652 (MH⁺).

Biological Data

(A) Agonist activity against receptor sub-types

[0128] The compounds of the Examples were tested in screen (1) (agonist activity against receptor sub-types) and the results obtained were as follows:

Example No	A2a	A1	A3
1	22.64	434.8	>93
2	30.95	755.4	>93
3	16.59	310.5	>93
4	37.24	1318.09	>93
5	10.54	159.5	>94
6	24.05	411.9	>97
7	22.98	597.82	>95
8	26.38	>6131	>165

[0129] Data are minimum values since preparation was found, after testing to contain an inactive impurity at around 20%.

Values given in the Table are EC₅₀ values as a ratio of that of NECA.


ABBREVIATIONS

[0130]

TMS	trimethylsilyl
TFA	trifluoroacetic acid
DMF	N,N-dimethylformamide
NECA	N-ethylcarboxamideadenosine
DMAP	4-dimethylaminopyridine
TEMPO	2,2,6,6-tetramethyl-1-piperidinyloxy, free radical
TMSOTf	Trimethylsilyltrifluoromethylsulphonate
DBU	1,8-diazabicyclo[5.4.0]undec-7-ene
BSA	bistrimethylsilylacetamide
DCM	dichloromethane
DAST	diethylaminosulphur trifluoride
Ph	phenyl
CDI	carbonyldiimidazole
NSAID	non-steroidal antiinflammatory drug
Bn	benzyl


Claims

1. A compound of formula (I):



wherein R¹ and R² independently represent a group selected from:


- 20
- (i) C₃₋₈cycloalkyl-;
- (ii) hydrogen;
- (iii) aryl₂CHCH₂-;
- (iv) C₃₋₈cycloalkylC₁₋₆alkyl-;
- (v) C₁₋₈alkyl-;
- 25
- (vi) arylC₁₋₆alkyl-;
- (vii) R⁴R⁵N-C₁₋₆alkyl-;
- (viii) C₁₋₆alkyl-CH(CH₂OH)-;
- (ix) arylC₁₋₅alkyl-CH(CH₂OH)-;
- (x) arylC₁₋₅alkyl-C(CH₂OH)₂-;
- 30
- (xi) C₃₋₈cycloalkyl independently substituted by one or more (e.g. 1, 2 or 3) -(CH₂)_pR⁶ groups;
- (xii) H₂NC(=NH)NHC₁₋₆alkyl-;
- (xiii) a group of formula



40

or such a group in which one methylene carbon atom adjacent to X, or both if such exist, is substituted by methyl;

- (xiv) -C₁₋₆alkyl-OH;
- (xv) -C₁₋₈haloalkyl;
- 45
- (xvi) a group of formula



- (xvii) aryl; and
- (xviii) -(CH₂)_fSO₂NH_g(C₁₋₄alkyl-)_{2-g} or -(CH₂)_fSO₂NH_g(arylC₁₋₄alkyl-)_{2-g};
- 55
- R³ represents methyl, ethyl, -CH=CH₂, n-propyl, -CH₂CH=CH₂, -CH=CHCH₃, isopropyl, isopropenyl, cyclopropyl, cyclopropenyl, cyclopropylmethyl, cyclopropenylmethyl, cyclobutyl, cyclobutenyl, -(CH₂)_qhalogen, -(CH₂)_hY(CH₂)_iH, -(CH₂)_hCOOCH₃, -(CH₂)_hOCOCH₃, -(CH₂)_hCON(CH₂)_mH((CH₂)_nH), -(CH₂)_hCO(CH₂)_oH or

$-\text{CH}_2\text{C}((\text{CH}_2)_u\text{H})=\text{NO}(\text{CH}_2)_v\text{H}$;

Y represents O, S or $\text{N}(\text{CH}_2)_j$;

a and b independently represent an integer 0 to 4 provided that $a + b$ is in the range 3 to 5;

c, d and e independently represent an integer 0 to 3 provided that $c + d + e$ is in the range 2 to 3;

f represents 2 or 3 and g represents an integer 0 to 2;

p represents 0 or 1;

q represents 2 or 3;

h represents 2 or 3;

i represents an integer 0 to 2 such that $h+i$ is in the range 2 to 4

j represents an integer 0 to 2 such that $h+i+j$ is in the range 2 to 4

m and n independently represent an integer 0 to 2 such that $m+n$ is in the range 0 to 2;

o represents an integer 0 to 2 such that $h+o$ is in the range 2 to 3;

u and v independently represent 0 or 1 such that $u+v$ is in the range 0 to 1;

R^4 and R^5 independently represent hydrogen, C_{1-6} alkyl, aryl, aryl C_{1-6} alkyl- or NR^4R^5 together may represent pyrrolidinyl, piperidinyl, morpholinyl, azetidiny, azepinyl, piperazinyl or N-C_{1-6} alkylpiperazinyl.

R^6 represents OH, NH_2 , NHCOCH_3 or halogen;

R^7 represents hydrogen, C_{1-6} alkyl, C_{1-6} alkylaryl or $-\text{COC}_{1-6}$ alkyl;


X represents NR^7 , O, S, SO or SO_2 ;

provided that when R^3 represents methyl, ethyl or isopropyl then R^1 and/or R^2 independently must represent:

(a) $-(\text{CH}_2)_f\text{SO}_2\text{NH}_g(\text{C}_{1-4}\text{alkyl})_{2-g}$ or $-(\text{CH}_2)_f\text{SO}_2\text{NH}_g(\text{arylC}_{1-4}\text{alkyl})_{2-g}$ where f is 2 or 3 and g is an integer 0 to 2;


(b) C_{3-8} cycloalkyl independently substituted by one or more $-(\text{CH}_2)_p\text{NHCOCH}_3$ groups;

(c) a group of formula



or such a group in which one methylene carbon atom adjacent to X, or both if such exist, is substituted by methyl;

(d) a group of formula



and salts and solvates thereof.

2. A compound of formula (I) according to claim 1 wherein R^1 and R^2 do not both represent hydrogen.
3. A compound according to claim 1 or claim 2 wherein R^1 represents aryl $_2\text{CHCH}_2-$, C_{1-8} alkyl, hydrogen or aryl C_{1-6} alkyl-.
4. A compound according to any one of claims 1 to 3 wherein R^1 represents $\text{Ph}_2\text{CHCH}_2-$.
5. A compound according to any one of claims 1 to 4 wherein R^2 represents $\text{R}^4\text{R}^5\text{N-C}_{1-6}$ alkyl-, aryl C_{1-6} alkyl-, aryl C_{1-5} alkyl $\text{CH}(\text{CH}_2\text{OH})-$, aryl C_{1-6} alkyl or C_{1-6} alkyl- $\text{CH}(\text{CH}_2\text{OH})-$.
6. A compound according to any one of claims 1 to 5 wherein R^2 represents $-(\text{CH}_2)_2(\text{piperidin-1-yl})$.
7. A compound according to any one of claims 1 to 6 wherein R^3 represents C_{1-3} alkyl, cyclobutyl, cyclopropylmethyl,

$-(\text{CH}_2)_2\text{OCOCH}_3$, $-(\text{CH}_2)_{2-3}\text{OH}$ or $-(\text{CH}_2)_2\text{halogen}$.

8. A compound according to any one of claims 1 to 7 wherein R^3 represents n-propyl, 2-propenyl, cyclobutyl, cyclopropylmethyl, $-(\text{CH}_2)_2\text{OCOCH}_3$, or $-(\text{CH}_2)_{2-3}\text{OH}$.

9. A compound according to any one of claims 1 to 8 wherein R^4 and R^5 independently represent hydrogen, C_{1-6} alkyl or aryl or NR^4R^5 together may represent pyrrolidinyl, piperidinyl, morpholinyl, azetidiny, azepinyl, piperazinyl or N-methylpiperazinyl.

10. A compound according to any one of claims 1 to 9 wherein R^6 represents OH or NH_2 .

11. A compound according to any one of claims 1 to 10 wherein X represents NR^7 , O, S or SO_2 .

12. A compound of the formula (I) which is

2R,3R,4S,5R)-2-[6-(2,2-Diphenyl-ethylamino)-2-(2-piperidin-1-yl-ethylamino)-purin-9-yl]-5-[2-(3-hydroxy-propyl)-2H-tetrazol-5-yl]-tetrahydro-furan-3,4-diol;

2R,3R,4S,5R)-2-[6-(2,2-Diphenyl-ethylamino)-2-(2-piperidin-1-yl-ethylamino)-purin-9-yl]-5-(2-propyl-2H-tetrazol-5-yl)-tetrahydro-furan-3,4-diol;

Acetic acid 2-(5-[5R-[6-(2,2-diphenyl-ethylamino)-2-(2-piperidin-1-yl-ethylamino)-purin-9-yl]-3S,4R-dihydroxy-tetrahydro-furan-2R-yl]-tetrazol-2-yl)-ethyl ester;

(2R,3S,4R,5R)-2-(2-Cyclopropylmethyl-2H-tetrazol-5-yl)-5-[6-(2,2-diphenyl-ethylamino)-2-(2-piperidin-1-yl-ethylamino)-purin-9-yl]-tetrahydro-furan-3,4-diol;

(2R,3R,4S,5R)-2-[6-(2,2-Diphenyl-ethylamino)-2-(2-piperidin-1-yl-ethylamino)-purin-9-yl]-5-[2-(2-hydroxy-ethyl)-2H-tetrazol-5-yl]-tetrahydro-furan-3,4-diol;

(2R,3S,4R,5R)-2-[2-(2-Chloro-ethyl)-2H-tetrazol-5-yl]-5-[6-(2,2-diphenyl-ethylamino)-2-(2-piperidin-1-yl-ethylamino)-purin-9-yl]-tetrahydro-furan-3,4-diol;

(2R,3S,4R,5R)-2-(2-Cyclobutyl-2H-tetrazol-5-yl)-5-[6-(2,2-diphenyl-ethylamino)-2-(2-piperidin-1-yl-ethylamino)-purin-9-yl]-tetrahydro-furan-3,4-diol;

(2R,3S,4R,5R)-2-(2-Allyl-2H-tetrazol-5-yl)-5-[6-(2,2-diphenyl-ethylamino)-2-(2-piperidin-1-yl-ethylamino)-purin-9-yl]-tetrahydro-furan-3,4-diol;

or a salt or solvate of any thereof.


13. A pharmaceutical composition comprising a compound of formula (I) as defined in any one of claims 1 to 12 or a pharmaceutically acceptable salt or solvate thereof in admixture with one or more pharmaceutically acceptable diluents or carriers.

14. A compound of formula (I) as defined in any one of claims 1 to 12 or a pharmaceutically acceptable salt or solvate thereof for use as a pharmaceutical.

15. Use of a compound of formula (I) as defined in any one of claims 1 to 12 or a pharmaceutically acceptable salt or solvate thereof in the manufacture of a medicament for the treatment of inflammatory diseases.

16. A process for preparation of a compound of formula (I) as defined in any one of claims 1 to 12 which comprises

(a) reacting a corresponding compound of formula (II)




wherein L represents a leaving group or a protected derivative thereof with a compound of formula R^2NH_2 or a protected derivative thereof, wherein R^1 , R^2 and R^3 are as defined in any one of claims 1 to 12; or
 (b) preparing a compound of formula (I) in which R^1 represents hydrogen by reducing a compound of formula (III)

5

10

15



20

or a protected derivative thereof, wherein R^2 and R^3 are as defined in any one of claims 1 to 12; or
 (c) deprotecting a compound of formula (I) which is protected; and where desired or necessary converting a compound of formula (I) or a salt thereof into another salt thereof,


17. A process for preparation of a compound of formula (I) as defined in any one of claims 1 to 12 without the proviso which comprises

25

(a) reacting a corresponding compound of formula (X)

30

35



40

with a compound of formula (XI)




45

wherein L is a leaving group and R^1 , R^2 and R^3 are as defined in any one of claims 1 to 12; or
 (b) reacting a corresponding compound of formula (XII)

50


55




with a compound of formula (V) or a protected derivative thereof, wherein R¹ and R² are as defined in any one of claims 1 to 12.

18. A process according to claim 17 which is a process for preparing the compound (2R,3R,4S,5R)-2-[6-Amino-2-(1S-hydroxymethyl-2-phenyl-ethylamino)-purin-9-yl]-5-(2-ethyl-2H-tetrazol-5-yl)-tetrahydro-furan-3,4-diol and salts and solvates thereof.

19. A process according to claim 18 which comprises reacting a corresponding compound of formula




with a compound of formula



or a protected derivative thereof, wherein L is a leaving group.


20. A compound of formula (II)



wherein L represents a leaving group and R¹ is as defined in any one of claims 1 to 12, and R³ represents n-propyl, 2-propenyl, cyclobutyl, cyclopropylmethyl, -(CH₂)₂OCOCH₃, or -(CH₂)_{2,3}OH or a protected derivative thereof.

21. A compound of formula (III)

5




10

15

wherein R^2 is as defined in any one of claims 1 to 12 and R^3 represents n-propyl, 2-propenyl, cyclobutyl, cyclopropylmethyl, $-(CH_2)_2OCOCH_3$, or $-(CH_2)_{2-3}OH$.

22. A compound of formula (IIIA)

20



25


30

wherein L represents a leaving group and R^3 represents n-propyl, 2-propenyl, cyclobutyl, cyclopropylmethyl, $-(CH_2)_2OCOCH_3$, or $-(CH_2)_{2-3}OH$ or a protected derivative thereof.

35

23. A compound of formula (IV)

40




45

50

wherein L^1 and L^2 independently represent a leaving group and R^3 represents n-propyl, 2-propenyl, cyclobutyl, cyclopropylmethyl, $-(CH_2)_2OCOCH_3$, or $-(CH_2)_{2-3}OH$ or a protected derivative thereof.


55

24. A compound of formula (V)



10 wherein L represents a leaving group and R³ represents n-propyl, 2-propenyl, cyclobutyl, cyclopropylmethyl, -(CH₂)₂OCOCH₃, or -(CH₂)₂₋₃OH or a protected derivative thereof.


25. A compound of formula (VI)



20 wherein alk represents C₁₋₆ alkyl eg methyl and R³ represents n-propyl, 2-propenyl, cyclobutyl, cyclopropylmethyl, -(CH₂)₂OCOCH₃, or -(CH₂)₂₋₃OH.

30 Patentansprüche

1. Verbindung der Formel (I):




worin R¹ und R² unabhängig eine Gruppe darstellen, die aus den folgenden ausgewählt ist:

- 50
- (i) C₃₋₈-Cycloalkyl-;
 - (ii) Wasserstoff;
 - (iii) Aryl₂CHCH₂-;
 - (iv) C₃₋₈-Cycloalkyl-C₁₋₆-alkyl-;
 - (v) C₁₋₈-Alkyl-;

55


 - (vi) Aryl-C₁₋₆-alkyl-;
 - (vii) R⁴R⁵N-C₁₋₆-Alkyl-;
 - (viii) C₁₋₆-Alkyl-CH(CH₂OH)-;
 - (ix) Aryl-C₁₋₅-alkyl-CH(CH₂OH)-;

- (x) Aryl-C₁₋₅-alkyl-C(CH₂OH)₂;
 (xi) C₃₋₈-Cycloalkyl, das unabhängig mit einer oder mehreren (z.B. 1, 2 oder 3) -(CH₂)_pR⁶-Gruppen substituiert ist;
 (xii) H₂NC(=NH)NHC₁₋₆-Alkyl-;
 (xiii) eine Gruppe der Formel



oder eine solche Gruppe, in der ein zu X benachbartes Methylen-Kohlenstoffatom oder beide, falls solche existieren, mit Methyl substituiert ist;

- (xiv) -C₁₋₆-Alkyl-OH;
 (xv) -C₁₋₈-Halogenalkyl;
 (xvi) eine Gruppe der Formel



- (xvii) Aryl; und
 (xviii) -(CH₂)_fSO₂NH_g(C₁₋₄-Alkyl)_{2-g} oder -(CH₂)_fSO₂NH_g(Aryl-C₁₋₄-alkyl)_{2-g};
 R³ Methyl, Ethyl, -CH=CH₂, n-Propyl, -CH₂CH=CH₂, -CH=CHCH₃, Isopropyl, Isopropenyl, Cyclopropyl, Cyclopropenyl, Cyclopropylmethyl, Cyclopropenylmethyl, Cyclobutyl, Cyclobutenyl,
 -(CH₂)_q-Halogen, -(CH₂)_hY(CH₂)_iH, -(CH₂)_hCCOCH₃,
 -(CH₂)_hOCOCH₃, -(CH₂)_hCON(CH₂)_mH((CH₂)_nH),
 -(CH₂)_hCO(CH₂)_oH oder -CH₂C((CH₂)_uH)=NO(CH₂)_vH darstellt;
 Y O, S oder N(CH₂)_j darstellt;

a und b unabhängig eine ganze Zahl von 0 bis 4 darstellen, mit der Massgabe, dass a + b im Bereich von 3 bis 5 ist;

c, d und e unabhängig eine ganze Zahl von 0 bis 3 darstellen, mit der Massgabe, dass c + d + e im Bereich von 2 bis 3 ist;

f 2 oder 3 darstellt und g eine ganze Zahl von 0 bis 2 darstellt;

p 0 oder 1 darstellt;

q 2 oder 3 darstellt;

h 2 oder 3 darstellt;

i eine ganze Zahl von 0 bis 2 darstellt, so dass h + i im Bereich von 2 bis 4 ist;

j eine ganze Zahl von 0 bis 2 darstellt, so dass h + i + j im Bereich von 2 bis 4 ist;

m und n unabhängig eine ganze Zahl von 0 bis 2 darstellen, so dass m + n im Bereich von 0 bis 2 ist;

o eine ganze Zahl von 0 bis 2 darstellt, so dass h + o im Bereich von 2 bis 3 ist;

u und v unabhängig 0 oder 1 darstellen, so dass u + v im Bereich von 0 bis 1 ist;

R⁴ und R⁵ unabhängig Wasserstoff, C₁₋₆-Alkyl, Aryl oder Aryl-C₁₋₆-alkyl- darstellen oder NR⁴R⁵ zusammen Pyrrolidinyl, Piperidinyl, Morpholinyl, Azetidiny, Azepiny, Piperazinyl oder N-C₁₋₆-Alkylpiperazinyl darstellen können;

R⁶ OH, NH₂, NHCOCH₃ oder Halogen darstellt;

R⁷ Wasserstoff, C₁₋₆-Alkyl, C₁₋₆-Alkylaryl oder -COC₁₋₆-Alkyl darstellt;

X NR⁷, O, S, SO oder SO₂ darstellt;


mit der Massgabe, dass dann, wenn R³ Methyl, Ethyl oder Isopropyl darstellt, R¹ und/oder R² unabhängig darstellen müssen:

EP 1 090 022 B1

(a) $-(\text{CH}_2)_f\text{SO}_2\text{NH}_g(\text{C}_{1-4}\text{-Alkyl-})_{2-g}$ oder $-(\text{CH}_2)_f\text{SO}_2\text{NH}_g(\text{Aryl-C}_{1-4}\text{-alkyl-})_{2-g}$, worin f 2 oder 3 ist und g eine ganze Zahl von 0 bis 2 ist;


(b) C_{3-8} -Cycloalkyl, das unabhängig mit einer oder mehreren $-(\text{CH}_2)_p\text{NHCOCH}_3$ -Gruppen substituiert ist ;

(c) eine Gruppe der Formel



oder eine solche Gruppe, in der ein zu X benachbartes Methylen-Kohlenstoffatom oder beide, falls solche existieren, mit Methyl substituiert ist;

(d) eine Gruppe der Formel



und Salze und Solvate davon.

2. Verbindung der Formel (I) gemäss Anspruch 1, worin R^1 und R^2 nicht beide Wasserstoff darstellen.
3. Verbindung gemäss Anspruch 1 oder 2, worin R^1 $\text{Aryl}_2\text{CHCH}_2$ -, C_{1-6} -Alkyl, Wasserstoff oder Aryl- C_{1-6} alkyl- darstellt.
4. Verbindung gemäss einem der Ansprüche 1 bis 3, worin R^1 Ph_2CHCH_2 - darstellt.
5. Verbindung gemäss einem der Ansprüche 1 bis 4, worin R^2 $\text{R}^4\text{R}^5\text{N-C}_{1-6}$ -Alkyl-, Aryl- C_{1-6} -alkyl-, Aryl- C_{1-5} alkyl-CH (CH_2OH) -, Aryl- C_{1-6} -alkyl oder C_{1-6} -Alkyl-CH (CH_2OH) - darstellt.
6. Verbindung gemäss einem der Ansprüche 1 bis 5, worin R^2 $-(\text{CH}_2)_2(\text{Piperidin-1-yl})$ darstellt.
7. Verbindung gemäss einem der Ansprüche 1 bis 6, worin R^3 C_{1-3} -Alkyl, Cyclobutyl, Cyclopropylmethyl, $-(\text{CH}_2)_2\text{OCOCH}_3$, $-(\text{CH}_2)_{2-3}\text{OH}$ oder $-(\text{CH}_2)_2$ -Halogen darstellt.
8. Verbindung gemäss einem der Ansprüche 1 bis 7, worin R^3 n-Propyl, 2-Propenyl, Cyclobutyl, Cyclopropylmethyl, $-(\text{CH}_2)_2\text{OCOCH}_3$ oder $-(\text{CH}_2)_{2-3}\text{OH}$ darstellt.
9. Verbindung gemäss einem der Ansprüche 1 bis 8, worin R^4 und R^5 unabhängig Wasserstoff, C_{1-6} -Alkyl oder Aryl darstellen oder NR^4R^5 zusammen Pyrrolidinyl, Piperidinyl, Morpholinyl, Azetidiny, Azepiny, Piperazinyl oder N-Methylpiperazinyl darstellen kann.
10. Verbindung gemäss einem der Ansprüche 1 bis 9, worin R^6 OH oder NH_2 darstellt.
11. Verbindung gemäss einem der Ansprüche 1 bis 10, worin X NR^7 , O, S oder SO_2 darstellt.
12. Verbindung der Formel (I), die
 $(2\text{R},3\text{R},4\text{S},5\text{R})$ -2-[6-(2,2-Diphenyl-ethylamino)-2-(2-piperidin-1-yl-ethylamino)-purin-9-yl]-5-[2-(3-hydroxy-propyl)-2H-tetrazol-5-yl]-tetrahydro-furan-3,4-diol;

(2R,3R,4S,5S)-2-[6(2,2-Diphenyl-ethylamino)-2-(2-piperidin-1-yl-ethylamino)-purin-9-yl]-5-(2-propyl-2H-tetrazol-5-yl)-tetrahydro-furan-3,4-diol;

Essigsäure-2-(5R-[5-[6-(2,2-diphenyl-ethylamino)-2-(2-piperidin-1-yl-ethylamino)-purin-9-yl]-3S,4Rdihydroxy-tetrahydro-furan-2R-yl]-tetrazol-2-yl)-ethylester;

(2R,3S,4R,5R)-2-(2-Cyclopropylmethyl-2H-tetrazol-5-yl)-5-[6-(2,2-diphenyl-ethylamino)-2-(2-piperidin-1-yl-ethylamino)-purin-9-yl]-tetrahydro-furan-3,4-diol; (2R,3R,4S,5R)-2-[6-(2,2-Diphenyl-ethylamino)-2-(2-piperidin-1-yl-ethylamino)-purin-9-yl]-5-[2-(2-hydroxy-ethyl)-2H-tetrazol-5-yl]-tetrahydro-furan-3,4-diol;

(2R,3S,4R,5R)-2-[2-(2-Chlor-ethyl)-2H-tetrazol-5-yl]-5-[6-(2,2-diphenyl-ethylamino)-2-(2-piperidin-1-ylethylamino)-purin-9-yl]-tetrahydro-furan-3,4-diol; (2R,3S,4R,5R)-2-(2-Cyclobutyl-2H-tetrazol-5-yl)-5-[6-(2,2-diphenyl-ethylamino)-2-(2-piperidin-1-ylethylamino)-purin-9-yl]-tetrahydro-furan-3,4-diol; (2R,3S,4R,5R)-2-(2-Allyl-2H-tetrazol-5-yl)-5-[6-(2,2-diphenyl-ethylamino)-2-(2-piperidin-1-yl-ethylamino)-purin-9-yl]-tetrahydro-furan-3,4-diol; oder ein Salz oder Sivat von jedem davon ist.


13. Pharmazeutische Zusammensetzung, die eine Verbindung der Formel (I) wie in einem der Ansprüche 1 bis 12 definiert oder ein pharmazeutisch akzeptables Salz oder Sivat davon im Gemisch mit einem oder mehreren pharmazeutisch akzeptablen Verdünnungstoffen oder Trägern umfasst.

14. Verbindung der Formel (I) wie in einem der Ansprüche 1 bis 12 definiert oder ein pharmazeutisch akzeptables Salz oder Sivat davon zur Verwendung als Pharmazeutikum.

15. Verwendung einer Verbindung der Formel (I) wie in einem der Ansprüche 1 bis 12 definiert oder eines pharmazeutisch akzeptablen Salzes oder Sivats davon in der Herstellung eines Medikaments zur Behandlung von entzündlichen Erkrankungen.

16. Verfahren zur Herstellung einer Verbindung der Formel (I) wie in einem der Ansprüche 1 bis 12 definiert, welches umfasst:

(a) Umsetzen einer entsprechenden Verbindung der Formel (II) :




worin L eine Abgangsgruppe darstellt, oder eines geschützten Derivats davon mit einer Verbindung der Formel R²NH₂ oder einem geschützten Derivat davon, worin R¹, R² und R³ wie in einem der Ansprüche 1 bis 12 definiert sind; oder

(b) Herstellen einer Verbindung der Formel (I), worin R¹ Wasserstoff darstellt, durch Reduzieren einer Verbindung der Formel (III) :

5

10

15



oder eines geschützten Derivats davon, worin R^2 und R^3 wie in einem der Ansprüche 1 bis 12 definiert sind; oder

20

(c) Entschützen einer Verbindung der Formel (I), die geschützt ist; und nach Wunsch oder Bedarf Umwandeln einer Verbindung der Formel (I) oder eines Salzes davon zu einem anderen Salz davon.

17. Verfahren zur Herstellung einer Verbindung der Formel (I) wie in einem der Ansprüche 1 bis 12 ohne die Massgabe definiert, welches umfasst:


25

(a) Umsetzen einer entsprechenden Verbindung der Formel (X) :

30

35

40



mit einer Verbindung der Formel (XI) :

45




worin L eine Abgangsgruppe ist und R^1 , R^2 und R^3 wie in einem der Ansprüche 1 bis 12 definiert sind; oder

50

(b) Umsetzen einer entsprechenden Verbindung der Formel (XII):

55




mit einer Verbindung der Formel (V) oder einem geschützten Derivat davon, worin R¹ und R² wie in einem der Ansprüche 1 bis 12 definiert sind.

18. Verfahren gemäss Anspruch 17, das ein Verfahren zur Herstellung der Verbindung (2R,3R,4S,5R)-2-[6-Amino-2-(1S-hydroxymethyl-2-phenyl-ethylamino)-purin-9-yl]-5-(2-ethyl-2H-tetrazol-5-yl)-tetrahydro-furan-3,4-diol und von Salzen und Solvaten davon ist.


15

19. Verfahren gemäss Anspruch 18, welches das Umsetzen einer entsprechenden Verbindung der Formel:

20



mit einer Verbindung der Formel:



oder einem geschützten Derivat davon umfasst, worin L eine Abgangsgruppe ist.


20. Verbindung der Formel (II):

45

50


55

55




worin L eine Abgangsgruppe darstellt und R¹ wie in einem der Ansprüche 1 bis 12 definiert ist und R³ n-Propyl, 2-Propenyl, Cyclobutyl, Cyclopropylmethyl, -(CH₂)₂OCOCH₃ oder -(CH₂)₂₋₃OH darstellt, oder ein geschütztes Derivat davon.

20 **21.** Verbindung der Formel (III) :



worin R² wie in einem der Ansprüche 1 bis 12 definiert ist und R³ n-Propyl, 2-Propenyl, Cyclobutyl, Cyclopropylmethyl, -(CH₂)₂OCOCH₃ oder -(CH₂)₂₋₃OH darstellt.

40 **22.** Verbindung der Formel (IIIA) :



worin L eine Abgangsgruppe darstellt und R³ n-Propyl, 2-Propenyl, Cyclobutyl, Cyclopropylmethyl,


$-(\text{CH}_2)_2\text{OCOCH}_3$ oder $-(\text{CH}_2)_{2-3}\text{OH}$ darstellt, oder ein geschütztes Derivat davon.

23. Verbindung der Formel (IV) :

5

10

15




20

worin L^1 und L^2 unabhängig eine Abgangsgruppe darstellen und R^3 n-Propyl, 2-Propenyl, Cyclobutyl, Cyclopropylmethyl, $-(\text{CH}_2)_2\text{OCOCH}_3$ oder $-(\text{CH}_2)_{2-3}\text{OH}$ darstellt, oder ein geschütztes Derivat davon.

24. Verbindung der Formel (V):

25

30




35

worin L eine Abgangsgruppe darstellt und R^3 n-Propyl, 2-Propenyl, Cyclobutyl, Cyclopropylmethyl, $-(\text{CH}_2)_2\text{OCOCH}_3$ oder $-(\text{CH}_2)_{2-3}\text{OH}$ darstellt, oder ein geschütztes Derivat davon.

25. Verbindung der Formel (VI):

40

45




50

worin alk C_{1-6} -Alkyl, z.B. Methyl, darstellt und R^3 n-Propyl, 2-Propenyl, Cyclobutyl, Cyclopropylmethyl, $-(\text{CH}_2)_2\text{OCOCH}_3$ oder $-(\text{CH}_2)_{2-3}\text{OH}$ darstellt.

55


Revendications

1. Composé de la formule (I)




15 où R¹ et R² représentent indépendamment un groupe choisi parmi :

- 20 (i) C₃₋₈-cycloalkyle
 (ii) l'hydrogène
 (iii) aryl₂CHCH₂
 (iv) C₃₋₈cycloalkyl-C₁₋₆alkyle
 (v) C₁₋₈ alkyle
 (vi) Aryl C₁₋₆ alkyle
 (vii) R⁴R⁵N-C₁₋₆ alkyle
 (viii) C₁₋₆ alkyl-CH(CH₂OH)
 (ix) Aryl C₁₋₅alkyl-CH(CH₂OH)-
 (x) Aryl C₁₋₅alkyl-C(CH₂OH)₂-
 (xi) C₃₋₈cycloalkyle indépendamment substitué par un ou plusieurs (par ex. 1, 2, ou 3) groupes de -(CH₂)_pR⁶
 (xii) H₂NC(=NH)NHC₁₋₆alkyle
 (xiii) Un groupe de formule



ou un tel groupe dans lequel un atome de carbone de méthylène adjacent à X, ou les deux s'ils existent, est substitué par du méthyle ;

- 40 (xiv) -C₁₋₆alkyl-OH
 (xv) -C₁₋₈haloalkyle
 (xvi) un groupe de formule



- (xvii) un aryle et
 (xviii) - (CH₂)_fSO₂NH_g(C₁₋₄alkyle)₂₋₉ ou -(CH₂)_fSO₂NH_g(aryl C₁₋₄-alkyl)₂₋₉
 R³ représente du méthyle, de l'éthyle, du -CH=CH₂, du n-propyle, du -CH₂CH=CH₂, du -CH=CHCH₃, de l'isopropyle, de l'isopropényle, du cyclopropyle, du cyclopropényle, du cyclopropylméthyle, du cyclopropénylméthyle, du cyclobutyle, du cyclobutényle, du -(CH₂)_q halogène,
 55 -(CH₂)_hY(CH₂)_iH, -(CH₂)_hCOOCH₃, -(CH₂)_hOCOCH₃,
 -(CH₂)_hCON(CH₂)_mH((CH₂)_nH), -(CH₂)_hCO(CH₂)_oH ou
 -CH₂C((CH₂)_uH)=NO(CH₂)_vH ;
 Y représente O, S ou N(CH₂)

EP 1 090 022 B1

a et b représentent indépendamment un nombre entier de 0 à 4 à condition que a + b soit entre 3 et 5
 c, d et e représentent indépendamment un nombre entier de 0 à 3 à condition que c + d + e soit entre 2 et 3.
 f représente 2 ou 3 et g représente un nombre entier de 0 à 2

p représente 0 ou 1

q représente 2 ou 3

h représente 2 ou 3

i représente un nombre entier de 0 à 2 tel que h+i soit entre 2 et 4

j représente un nombre entier de 0 à 2 tel que h+i+j soit entre 2 et 4

m et n représentent indépendamment un nombre entier de 0 à 2 tel que m + n soit entre 0 et 2

o représente un nombre entier de 0 à 2 tel que h + o soit entre 2 et 3

u et v représentent indépendamment 0 ou 1 tel que u+v soit entre 0 et 1.

R⁴ et R⁵ représentent indépendamment de l'hydrogène, un C₁₋₆ alkyle, un aryle, un aryle C₁₋₆ alkyle ou NR⁴R⁵ ensemble peuvent représenter le pyrrolidinyle, le pipéridinyle, le morpholinyle, l'azétidinyle, l'azépinyle, le pipérazinyle ou le N-C₁₋₆alkylpipérazinyle.

R⁶ représente du OH, NH₂, NHCOCH₃ ou un halogène.

R⁷ représente de l'hydrogène, du C₁₋₆alkyle, du C₁₋₆ alkylaryle ou du -COC₁₋₆ alkyle


X représente du NR⁷, O, S, SO ou SO₂

à condition que lorsque R³ représente du méthyle, de l'éthyle ou de l'isopropyle, alors R¹ et/ou R² doivent représenter indépendamment :

(a) du -(CH₂)_fSO₂NH_g(C₁₋₄-alkyle)_{2-g} ou -(CH₂)_fSO₂NH_g(arylC₁₋₄ alkyle)_{2-g} où f est égal à 2 ou 3 et g est un nombre entier de 0 à 2.


(b) C₃₋₈ cycloalkyle indépendamment substitué par un ou plusieurs groupes de -(CH₂)_pNHCOCH₃

(c) un groupe de formule



ou un tel groupe dans lequel un atome de carbone de méthylène adjacent à X, ou les deux s'ils existent, est substitué par du méthyle ;

(d) un groupe de formule



et des sels et solvates de celui-ci.

2. Composé de la formule (I) selon la revendication 1, **caractérisé en ce que** R¹ et R² ne représentent pas tous deux de l'hydrogène.

3. Composé selon la revendication 1 ou 2 **caractérisé en ce que** R¹ représente un aryl₂CHCH₂-, un C₁₋₈-alkyle, de l'hydrogène ou un aryle C₁₋₆alkyle.


4. Composé selon l'une quelconque des revendications 1 à 3, **caractérisé en ce que** R¹ représente du Ph₂CHCH₂-.

5. Composé selon l'une quelconque des revendications 1 à 4, **caractérisé en ce que** R² représente du R⁴R⁵N-C₁₋₆-alkyle-, un arylC₁₋₆alkyle, un arylC₁₋₅alkyleCH(CH₂OH)-, un arylC₁₋₆alkyle ou un C₁₋₆ alkyle-CH(CH₂OH).

6. Composé selon l'une quelconque des revendications 1 à 5, **caractérisé en ce que** R² représente du -(CH₂)₂


(pipéridin-1-yl).

- 5
7. Composé selon l'une quelconque des revendications 1 à 6, **caractérisé en ce que** R³ représente un C₁₋₃alkyle, du cyclobutyle, du cyclopropylméthyle, du -(CH₂)₂OCOCH₃, du -(CH₂)₂₋₃OH ou un -(CH₂)₂ halogène.
8. Composé selon l'une quelconque des revendications 1 à 7, **caractérisé en ce que** R³ représente du n-propyle, du 2-propényle, du cyclobutyle, du cyclopropylméthyle, du -(CH₂)₂OCOCH₃, ou du -(CH₂)₂₋₃OH.
- 10
9. Composé selon l'une quelconque des revendications 1 à 8, **caractérisé en ce que** R⁴ et R⁵ représentent indépendamment de l'hydrogène, un C₁₋₆ alkyle, ou un aryle ou NR⁴R⁵ ensemble peuvent représenter le pyrrolidinyle, le pipéridinyle, le morpholinyle, l'azétidinyle, l'azépinyle, le pipérazinyle ou le N-méthylpipérazinyle.
- 15
10. Composé selon l'une quelconque des revendications 1 à 9, **caractérisé en ce que** R⁶ représente du OH ou NH₂.
- 15
11. Composé selon l'une quelconque des revendications 1 à 10, **caractérisé en ce que** X représente du NR⁷, O, S ou SO₂.
- 20
12. Composé de formule (I) qui est du
 (2R,3R,4S,5R)-2-[6-(2,2-diphényl-éthylamino)-2-(2-pipéridin-1-yl-éthylamino)-purin-9-yl]-5-[2-(3-hydroxypropyl)-2H-tétrazol-5-yl]-tétrahydro-furan-3,4-diol ;
 (2R,3R,4S,5R)-2-[6-(2,2-diphényl-éthylamino)-2-(2-pipéridin-1-yl-éthylamino)-purin-9-yl]-5-(2-propyl-2H-tétrazol-5-yl)-tétrahydrofuran-3,4-diol ;
 Acide acétique 2-(5-{5R-[6-(2,2-diphényl-éthylamino)-2-(2-pipéridin-1-yl-éthylamino)-purin-9-yl]-3S,4R-dihydroxy-tétrahydrofuran-2R-yl}-tétrazol-2-yl)-éthylester ;
 25 (2R,3S,4R,5R)-2-(2-cyclopropylméthyl-2H-tétrazol-5-yl)-5-[6-(2,2-diphényl-éthylamino)-2-(2-pipéridin-1-yl-éthylamino)-purin-9-yl]-tétrahydro-furan-3,4-diol
 (2R,3R,4S,5R)-2-[6-(2,2-diphényl-éthylamino)-2-(2-pipéridin-1-yl-éthylamino)-purin-9-yl]-5-[2-(2-hydroxyéthyl)-2H-tétrazol-5-yl]-tétrahydro-furan-3,4-diol
 (2R,3S,4R,5R)-2-[2-(2-chloro-éthyl)-2H-tétrazol-5-yl]-5-[6-(2,2-diphényl-éthylamino)-2-(2-pipéridin-1-yl-éthylamino)-purin-9-yl]-tétrahydro-furan-3,4-diol
 30 (2R,3S,4R,5R)-2-(2-cyclobutyl-2H-tétrazol-5-yl)-5-[6-(2,2-diphényl-éthylamino)-2-(2-pipéridin-1-yl-éthylamino)-purin-9-yl]-tétrahydro-furan-3,4-diol.
 (2R,3S,4R,5R)-2-(2-allyl-2H-tétrazol-5-yl)-5-[6-(2,2-diphényl-éthylamino)-2-(2-pipéridin-1-yl-éthylamino)-purin-9-yl]-tétrahydro-furan-3,4-diol ;
 35 ou un sel ou solvate de l'un quelconque de ces composés.
- 40
13. Composition pharmaceutique comprenant un composé de formule (I), comme défini dans l'une quelconque des revendications 1 à 12, ou sel ou solvate de celui-ci pharmaceutiquement acceptable en mélange avec un ou plusieurs diluants ou supports pharmaceutiquement acceptables.
- 45
14. Composé de formule (I), comme défini dans l'une quelconque des revendications 1 à 12, ou sel ou solvate pharmaceutiquement acceptable de celui-ci à utiliser comme produit pharmaceutique.
- 50
15. Utilisation d'un composé de formule (I) comme défini dans l'une quelconque des revendications 1 à 12, ou sel ou solvate pharmaceutiquement acceptable de celui-ci dans la fabrication d'un médicament pour le traitement de maladies inflammatoires.
- 55
16. Procédé pour la préparation d'un composé de formule (I), comme défini dans l'une quelconque des revendications 1 à 12 qui comprend :
- (a) la réaction d'un composé correspondant de formule (II)



15 à L représente un groupe partant ou un dérivé protégé de celui-ci avec un composé de formule R²NH₂ ou un dérivé protégé de celui-ci, dans lequel R¹, R² et R³ sont comme définis dans l'une quelconque des revendications 1 à 12 ou

(b) la préparation d'un composé de formule (I) dans lequel R¹ représente de l'hydrogène en réduisant un composé de formule (III)




ou un dérivé protégé de celui-ci, dans laquelle R² et R³ sont comme définis dans l'une quelconque des revendications 1 à 12 ou

(c) la déprotection d'un composé de formule (I) qui est protégé ; et quand cela est souhaité ou nécessaire, la conversion d'un composé de formule (I) ou d'un sel de celui-ci en un autre sel de celui-ci.

35 **17.** Procédé pour la préparation d'un composé de formule (I) comme défini dans l'une quelconque des revendications 1 à 12, sans la condition qui comprend


40 (a) la réaction d'un composé correspondant de formule (X)



55 avec un composé de formule (XI)




où L est un groupe partant et R¹, R² et R³ sont comme définis dans l'une quelconque des revendications 1 à 12, ou
 (b) la réaction d'un composé correspondant de formule (XII)




avec un composé de formule (V) ou un dérivé protégé de celui-ci, où R¹ et R² sont comme définis dans l'une quelconque des revendications 1 à 12.

18. Procédé selon la revendication 17, qui est un procédé de préparation du composé (2R,3R,4S,5R)-2-[6-amino-2-(1S-hydroxyméthyl-2-phényl-éthylamino)-purin-9-yl]-5-(2-éthyl-2H-tétrazol-5-yl)-tétrahydro-furan-3,4-diol et des sels et solvates de celui-ci.

19. Procédé selon la revendication 18, qui comprend la réaction d'un composé correspondant de formule




avec un composé de formule




ou un dérivé protégé de celui-ci, où L est un groupe partant.

20. Composé de formule (II)




15 où L représente un groupe partant et R¹ est comme défini dans l'une quelconque des revendications 1 à 12, et R³ représente du n-propyle, du 2-propényle, du cyclobutyle, du cyclopropylméthyle, du $-(CH_2)_2OCOCH_3$ ou du $-(CH_2)_{2-3}OH$, ou un dérivé protégé de celui-ci.

21. Composé de formule (III)



35 où R² est comme défini dans l'une quelconque des revendications 1 à 12 et R³ représente du n-propyle, du 2-propényle, du cyclobutyle, du cyclopropylméthyle, du $-(CH_2)_2OCOCH_3$ ou du $-(CH_2)_{2-3}OH$.

22. Composé de formule (IIIA)




55 où L représente un groupe partant et R³ représente du n-propyle, du 2-propényle, du cyclobutyle, du cyclopropylméthyle, du $-(CH_2)_2OCOCH_3$ ou du $-(CH_2)_{2-3}OH$, ou un dérivé protégé de celui-ci.

23. Composé de formule (IV)

5

10




15

où L¹ et L² représentent indépendamment un groupe partant et R³ représente du n-propyle, du 2-propényle, du cyclobutyle, du cyclopropylméthyle, du -(CH₂)₂OCOCH₃ ou du -(CH₂)₂₋₃OH, ou un dérivé protégé de celui-ci.

24. Composé de formule (V)

20

25




30

où L représente un groupe partant et R³ représente du n-propyle, du 2-propényle, du cyclobutyle, du cyclopropylméthyle, du -(CH₂)₂OCOCH₃ ou du -(CH₂)₂₋₃OH, ou un dérivé protégé de celui-ci.

25. Composé de formule (VI)

35

40



45

où alk représente un C₁₋₆ alkyle, par exemple du méthyle et R³ représente du n-propyle, du 2-propényle, du cyclobutyle, du cyclopropylméthyle, du -(CH₂)₂OCOCH₃ ou du -(CH₂)₂₋₃OH.

50

55