




(86) Date de dépôt PCT/PCT Filing Date: 2002/07/12  
 (87) Date publication PCT/PCT Publication Date: 2003/01/23  
 (45) Date de délivrance/Issue Date: 2011/04/19  
 (85) Entrée phase nationale/National Entry: 2004/01/12  
 (86) N° demande PCT/PCT Application No.: JP 2002/007129  
 (87) N° publication PCT/PCT Publication No.: 2003/006439  
 (30) Priorités/Priorities: 2001/07/13 (JP2001-213418);  
 2001/07/13 (JP2001-213417);  
 2001/10/09 (JP2001-310900);  
 2001/11/27 (JP2001-360339); 2002/01/16 (JP2002-7015);  
 2002/02/19 (JP2002-42076)

(51) Cl.Int./Int.Cl. *C07D 239/42* (2006.01),  
*C07D 239/22* (2006.01), *C07D 239/30* (2006.01),  
*C07D 239/34* (2006.01), *C07D 239/36* (2006.01)  
 (72) Inventeurs/Inventors:  
 MATSUSHITA, AKIO, JP;  
 ODA, MIZUHO, JP;  
 KAWACHI, YASUHIRO, JP;  
 CHIKA, JUN-ICHI, JP  
 (73) Propriétaire/Owner:  
 ASTRAZENECA UK LIMITED, GB  
 (74) Agent: FETHERSTONHAUGH & CO.

(54) Titre : PREPARATION DE COMPOSES D'AMINOPYRIMIDINE  
 (54) Title: PREPARATION OF AMINOPYRIMIDINE COMPOUNDS




(57) Abrégé/Abstract:

A 2-(N-methyl-N-methanesulfonylamino)pyrimidine compound of the formula (3): [R is a hydrocarbonyl group], is prepared by the steps of: (I) reacting an isobutyrylacetate ester with 4-fluorobenzaldehyde and urea in the presence of a protonic compound and a metal salt; (II) oxidizing the reaction product of the step (I); (III) reacting the oxidation product of the step (II) with an organic sulfonyl halide or an organic sulfonyl anhydride; and (IV) reacting the reaction product of the step (III) with N-methyl-N-methanesulfonamide. (see formula 3)

23940-1566

**ABSTRACT**

A 2-(N-methyl-N-methanesulfonylamino)pyrimidine compound of the formula (3): [R is a hydrocarbyl group], is prepared by the steps of: (I) reacting an isobutyrylacetate ester with 4-fluorobenzaldehyde and urea in the presence of a protonic compound and a metal salt; (II) oxidizing the reaction product of the step (I); (III) reacting the oxidation product of the step (II) with an organic sulfonyl halide or an organic sulfonyl anhydride; and (IV) reacting the reaction product of the step (III) with N-methyl-N-methanesulfonamide.




23940-1566

-1-

PREPARATION OF AMINOPYRIMIDINE COMPOUNDSFIELD OF THE INVENTION

5 The present invention relates to the preparation of aminopyrimidine compounds having the following formula (8):


10



15 [in the formula (8), R is a hydrocarbyl group, and each of R<sup>1</sup> and R<sup>2</sup> independently is a hydrogen atom, an alkyl group, an alkylsulfonyl group, or an arylsulfonyl group], more particularly to the preparation of a 2-(N-methyl-N-methanesulfonylamino)pyrimidine compound having the following formula (3):

20

25



wherein R represents a hydrocarbyl group.

30


BACKGROUND OF THE INVENTION

Bioorg. Med. Chem., 5, 437(1997) describes that the 2-(N-methyl-N-methanesulfonylamino)pyrimidine compound is employable as an intermediate compound for producing a cholesterol reducing agent (HMG-CoA reductase inhibitor: S-4522) having the following formula:

35

23940-1566

-2-



and which is now generally known as the calcium salt of  
10 rosuvastatin or rosuvastatin calcium.

WO 01/04100 describes a process for preparing the 2-  
(N-methyl-N-methanesulfonylamino)pyrimidine compound  
which comprises the steps of:

15 reacting methyl isobutyrylacetate with 4-fluoro-  
benzonitrile to produce methyl 2-[1-amino-1-(4-fluoro-  
phenyl)methylene]-4-methyl-3-oxopentanoate; and

20 reacting the 2-[1-amino-1-(4-fluorophenyl)methyl-  
ene]-4-methyl-3-oxopentanoate with N-cyano-N-methyl-  
methanesulfonamide which is obtained by reaction between  
N-methylmethanesulfonamide and cyanogen chloride, to pro-  
duce 4-(4-fluorophenyl)-6-isopropyl-5-methoxycarbonyl-2-  
(N-methanesulfonyl-N-methylamino)pyrimidine.

It is described that the total yield (based on the  
amount of methyl isobutyrylacetate) is 45.5%.

25 It appears that the process described in WO 01/04100  
is disadvantageous for the industrial preparation, be-  
cause the yield is not high and it is necessary to employ  
toxic cyanogen chloride as one of the starting compounds.

#### 30 SUMMARY OF THE INVENTION


The invention provides a novel  
process for preparing a 2-(N-methyl-N-  
methanesulfonylamino)pyrimidine or an analogous amino-  
pyrimidine compound thereof, more particularly to provide  
35 a novel process which provides the compound more conve-  
niently and/or without employing a toxic compound and/or  
provides the compound in high yield and/or high purity.

23940-1566

-3-


The invention provides a novel process for preparing a 2-(N-methyl-N-methanesulfonylamino)pyrimidine compound or an analogous amino-pyrimidine compound thereof which is favorably employable  
5 in the industrial preparation.

The present invention resides in a process for preparing a 2-(N-methyl-N-methanesulfonylamino)pyrimidine having the formula (3):  
10



[R is a hydrocarbyl group],  
which comprises the steps of:

20 reacting a hydroxypyrimidine compound having the formula (1):



in which R is the same as above,  
with an organic sulfonyl halide having the formula (2):  
30



in which R' is a hydrocarbyl group and X is a halogen atom, or an organic sulfonic anhydride having the formula  
35 (2a):

-4-




in which R' is the same as above, and

5 reacting the resulting reaction product with N-methyl-N-methanesulfonamide.

The invention also resides in a hydroxypyrimidine compound having the above-identified formula (1).


10 The invention further resides in a method for preparing a hydroxypyrimidine compound of the formula (1), which comprises oxidizing a dihydropyrimidinone compound having the formula (4):



wherein R is a hydrocarbyl group.

20 The invention further resides in a dihydropyrimidinone compound having the formula (4).


The invention furthermore resides in a method for preparing a dihydropyrimidinone compound of the formula (4), which comprises reacting an isobutyrylacetate ester  
25 having the formula (5):



30 in which R is a hydrocarbyl group, with 4-fluorobenzaldehyde and urea in the presence of a protonic compound and a metal salt.

The invention furthermore resides in a method for preparing an aminopyrimidine compound having the formula  
35 (8):


-5-



5

wherein R is a hydrocarbyl group, and each of R<sup>1</sup> and R<sup>2</sup> independently is hydrogen atom, an alkyl group, an alkyl-sulfonyl group, or an arylsulfonyl group, which comprises reacting a 2-substituted pyrimidine compound having the formula (6):


10



15

wherein R is the same as above, and X is a halogen atom or an organic sulfonyloxy group, with an amine compound having the formula (7):

20




25

wherein each of R<sup>1</sup> and R<sup>2</sup> is the same as above.

The invention furthermore resides in a halogenopyrimidine compound having the formula (9):

30




35

wherein R is a hydrocarbyl group, and Hal is a halogen atom.

-6-

The invention furthermore resides in a method for preparing the halogenopyrimidine compound of the formula (9), which comprises reacting a hydroxypyrimidine compound of the aforementioned formula (1) with a halogenating agent.

The invention furthermore resides in an organic sulfonyloxypyrimidine compound having the formula (10):



15 wherein each of R and R' independently is a hydrocarbyl group.

The invention furthermore resides in a method for preparing an organic sulfonyloxypyrimidine compound of the formula (10), which comprises reacting a hydroxypyrimidine compound of the aforementioned formula (1) with an organic sulfonyl halide having the formula (2):



25 wherein R' is a hydrocarbyl group, and X is a halogen atom, or an organic sulfonic anhydride having the formula (2a):



30 in which R' is the same as above.

The invention furthermore resides in a process for preparing a 2-(N-methyl-N-methanesulfonylamino)pyrimidine of the formula (3) which comprises the steps of:

35 (I) reacting an isobutyrylacetate ester of the formula (5) with 4-fluorobenzaldehyde and urea in the presence of a protonic compound and a metal salt;



23940-1566

-7-


(II) oxidizing the reaction product of the step (I);  
 (III) reacting the oxidation product of the step  
 (II) with an organic sulfonyl halide of the formula (2)  
 or an organic sulfonic anhydride of the formula (2a); and  
 5 (IV) reacting the reaction product of the step (III)  
 with N-methyl-N-methanesulfonamide.

In the above-mentioned process, the steps (III) and  
 (IV) can be carried out continuously in the same reaction  
 mixture.

10

DETAILED DESCRIPTION OF THE INVENTION

The representative process for the preparation of 2-  
 (N-methyl-N-methanesulfonylamino)pyrimidine of the for-  
 mula (3) according to the present invention is schemati-  
 15 cally illustrated as follows:




-8-

Each step in the above-illustrated reaction scheme is described below in more detail.

Step (I)

5 In the step (I), an isobutyrylacetate ester of the following formula (5):



10

[R is a hydrocarbyl group].

is reacted with 4-fluorobenzaldehyde and urea in the presence of a protonic compound and a metal salt.

15 The hydrocarbyl group (i.e., hydrocarbon group) represented by R in the formulas of the compounds involved in the reactions of the invention can be an alkyl group such as methyl, ethyl, propyl, butyl, pentyl, hexyl, heptyl, octyl, nonyl, or decyl, more particularly an alkyl group having 1-6 carbon atoms and especially an  
20 alkyl group having 1-4 carbon atoms; a cycloalkyl group such as cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl; an aralkyl group such as benzyl, phenylethyl, or phenylpropyl; or an aryl group such as phenyl or methylphenyl. The hydrocarbyl group can take any isomer configurations such as normal, iso, and tertiary. The  
25 hydrocarbyl group can have one or more substituents, provided that the substituents do not disturb the reaction involved.

30 The protonic compound can be an inorganic acid or its salt such as hydrochloric acid, sulfuric acid, potassium hydrogensulfate, sodium hydrogen sulfate, nitric acid, or phosphoric acid; an organic sulfonic acid such as methanesulfonic acid, ethanesulfonic acid, benzenesulfonic acid, p-toluenesulfonic acid, or p-bromobenzenesulfonic acid; an organic carboxylic acid such as acetic  
35 acid, propionic acid, butyric acid, or benzoic acid; an

-9-

alcohol such as methanol, ethanol, or propanol. Preferred are protonic acids such as hydrochloric acid, sulfuric acid, p-toluenesulfonic acid, and acetic acid. Most preferred is sulfuric acid. The protonic compounds  
5 can be employed singly or in combination.

The protonic compound can be employed in an amount of, preferably, 0.01 to 3 mol., more preferably 0.1 to 1 mol., per one mol. of the isobutyrylacetate ester.

The metal salt employed in the reaction can be copper(I) chloride, copper(II) chloride, copper(II) acetate,  
10 iron(II) chloride, iron(III) chloride, aluminum chloride, nickel(II) bromide, tin(IV) chloride, titanium tetrachloride, or magnesium bromide. Preferred are copper(I) chloride, copper(II) chloride, iron(III) chloride and  
15 nickel(II) bromide. Most preferred is copper(I) chloride. The metal salts may contain water of crystallization. The metal salts can be employed singly or in combination.

The metal salt can be employed in an amount of,  
20 preferably, 0.001 to 5 mol., more preferably 0.01 to 0.1 mol., per one mol. of the isobutyrylacetate ester.

The 4-fluorobenzaldehyde can be employed in an amount of, preferably, 0.5 to 10 mol., more preferably 0.9 to 1.1 mol., per one mol. of the isobutyrylacetate  
25 ester.

The urea can be employed in an amount of, preferably, 0.5 to 10 mol., more preferably 1.5 to 2 mol., per one mol. of the isobutyrylacetate ester.

The reaction can be performed in the presence or  
30 absence of a solvent. There are no specific limitations with respect to the solvent employed, so far as the solvent does not disturb the desired reaction. Examples of the employable solvents include alcohols such as methanol, ethanol, n-propyl alcohol, isopropyl alcohol, n-butyl alcohol, isobutyl alcohol, sec-butyl alcohol, and  
35 t-butyl alcohol; ethers such as diethyl ether, diisopro-

-10-

pyl ether, tetrahydrofuran, and dimethoxyethane; nitriles such as acetonitrile, propionitrile, butyronitrile, and isobutyronitrile; halogenated aliphatic hydrocarbons such as dichloromethane, dichloroethane, chloroform, and carbon tetrachloride; aromatic hydrocarbons such as benzene, toluene, and xylene; halogenated aromatic hydrocarbons such as chlorobenzene; and nitrated aromatic hydrocarbons such as nitrobenzene. Preferred are methanol, ethanol, n-propyl alcohol, isopropyl alcohol, n-butyl alcohol, diisopropyl ether, tetrahydrofuran, dimethoxyethane, acetonitrile, butyronitrile, isobutyronitrile, dichloromethane, dichloroethane, chloroform, toluene, xylene, and chlorobenzene. Especially preferred are methanol, ethanol, and isopropyl alcohol. The solvents can be employed singly or in combination.

The solvent can be employed in an amount of, preferably 0.1 to 10 liters, more preferably 0.3 to 2 liters, per one mole of the isobutyrylacetate ester. The amount may vary depending on homogeneity and dispersability of the reaction mixture.

The reaction can be conducted by reacting the isobutyrylacetate ester, 4-fluorobenzaldehyde, and urea, in a solvent in the presence of a protonic compound and a metal salt under inert gas atmosphere. The reaction can be carried out at a temperature of, preferably -10 to 200°C, more preferably 30 to 100°C. There are no specific limitations with respect to the surrounding pressure.

The resulting product of the reaction, that is, a dihydropyrimidinone compound of the formula (4), can be isolated and purified according to the conventional procedures such as distillation, crystallization, recrystallization, and column chromatography.

#### Step (II)

In the step (II), a dihydropyrimidinone compound of the formula (4), that is, the reaction product of the

-11-

step (I), is oxidized to give a hydroxypyrimidine compound of the formula (1).

The oxidation (or dehydrogenation oxidation) can be performed in various conventional manners. Preferred is  
5 oxidation utilizing nitric acid, because this oxidation procedure is easily carried out and the post-treatment of the reaction product is easy.

The nitric acid can be employed in an amount of, preferably 1 to 20 mol., more preferably 3 to 15 mol.,  
10 per one mole of the dihydropyrimidinone compound of the formula (4). The nitric acid of a concentration of, preferably 40 to 80%, more preferably 50 to 70%, can be preferably employed.

The oxidation can be performed in the presence or  
15 absence of a solvent. There are no specific limitations with respect to the solvent employed, so far as the solvent does not disturb the desired reaction. Examples of the preferred solvents include carboxylic acids such as acetic acid, propionic acid, and butyric acid. The sol-  
20 vents can be employed singly or in combination.

The solvent can be employed in an amount of, preferably 0.1 to 7 mL, more preferably 0.5 to 3 mL, per 1 g of the dihydropyrimidinone compound. The amount may vary depending on homogeneity and dispersability of the reac-  
25 tion mixture.

The oxidation can be conducted by reacting the dihydropyrimidinone compound and nitric acid in a solvent under inert gas atmosphere. The oxidation can be carried out at a temperature of, preferably -10 to 100°C, more  
30 preferably 0 to 50°C. There are no specific limitations with respect to the surrounding pressure. A reaction initiator such as sodium nitrite may be incorporated into the reaction system to accelerate the oxidation rate.

The resulting product of the reaction, that is, the  
35 hydroxypyrimidine compound of the formula (1), can be isolated and purified according to the conventional pro-

-12-


cedures such as distillation, crystallization, recrystallization, and column chromatography.

Steps (III) and (IV)

5 In the steps (III) and (IV), a hydroxypyrimidine compound of the formula (1), that is, the reaction product of the step (II), is reacted with an organic sulfonyl halide of the formula (2):



or an organic sulfonic anhydride of the formula (2a):



and

reacting the resulting reaction product with N-methyl-N-methanesulfonamide.

20 In the formulas (2) and (2a), R' is a hydrocarbyl group which can have one or more substituents. Examples of the hydrocarbyl groups include alkyl groups such as methyl, ethyl, propyl, butyl, pentyl, heptyl, octyl, nonyl, and decyl, more particularly an alkyl group having 1-6 carbon atoms and especially an alkyl group having 1-4  
25 carbon atoms; fluorinated alkyl groups such as trifluoromethyl, nonafluorobutyl, tridecafluorohexyl, heptadecafluorooctyl, and uncosafluorodecyl; cycloalkyl groups such as cyclopropyl, cyclobutyl, cyclopentyl, and cyclohexyl; aralkyl groups such as benzyl, phenylethyl, and  
30 phenylpropyl; and aryl groups, including unsubstituted and substituted phenyl or naphthyl groups, such as phenyl, naphthyl, tolyl, xylyl, mesityl, triisopropylphenyl, methoxyphenyl, chlorophenyl, and nitrophenyl. Thus, the hydrocarbyl group can have one or more substituents, pro-  
35 vided that the substituents do not disturb the reaction involved. The hydrocarbyl group can take any isomer con-

-13-

figurations such as normal, iso, and tertiary. A particularly suitable value for R' when it is aryl includes, for example, a phenyl or naphthyl group (particularly phenyl) which is unsubstituted or bears 1, 2 or 3 substituents. The substituents may be independently selected from, for example, alkyl having 1-4 carbon atoms, alkoxy having 1-4 carbon atoms, halogeno, and nitro.

In the formula (2), X is a halogen atom such as fluorine, chlorine, bromine, and iodine.

Examples of the sulfonyl halides include methanesulfonyl fluoride, methanesulfonyl chloride, ethanesulfonyl chloride, 1-propanesulfonyl chloride, 2-propanesulfonyl chloride, trifluoromethanesulfonyl fluoride, trifluoromethanesulfonyl chloride, nonafluorobutanesulfonyl fluoride, tridecafluorohexanesulfonyl fluoride, heptadecafluorooctanesulfonyl fluoride, uncosafluorodecanesulfonyl fluoride, benzenesulfonyl chloride, 1-naphthalenesulfonyl chloride, 2-naphthalenesulfonyl chloride, p-toluenesulfonyl fluoride, p-toluenesulfonyl chloride, 2,4,6-trimethylbenzenesulfonyl chloride, 2,4,6-triisopropylbenzenesulfonyl chloride, p-methoxybenzenesulfonyl chloride, p-chlorobenzenesulfonyl chloride, and 2-nitrobenzenesulfonyl chloride. Preferred are trifluoromethanesulfonyl fluoride, benzenesulfonyl chloride, 1-naphthalenesulfonyl chloride, 2-naphthalenesulfonyl chloride, p-toluenesulfonyl chloride, 2,4,6-trimethylbenzenesulfonyl chloride, 2,4,6-triisopropylbenzenesulfonyl chloride, p-methoxybenzenesulfonyl chloride, and p-chlorobenzenesulfonyl chloride. Particularly preferred are p-toluenesulfonyl chloride, 2,4,6-trimethylbenzenesulfonyl chloride, 2,4,6-triisopropylbenzenesulfonyl chloride, and p-methoxybenzenesulfonyl chloride.

Examples of the sulfonic anhydrides include methanesulfonic anhydride, trifluoromethanesulfonic anhydride, benzenesulfonic anhydride, and p-toluenesulfonic anhydride. Preferred are trifluoromethanesulfonic anhydride,

-14-

benzenesulfonic anhydride, and p-toluenesulfonic anhydride. Particularly preferred are trifluoromethanesulfonic anhydride and p-toluenesulfonic anhydride.

5 The sulfonyl halide or sulfonic anhydride can be employed in an amount of, preferably 0.1 to 20 mol., more preferably 0.5 to 5 mol., most preferably 1 to 2 mol., per one mole of the hydroxypyrimidine compound.

10 In the subsequent step, N-methylmethanesulfonamide can be employed in an amount of, preferably 0.1 to 30 mol., more preferably 1 to 5 mol., per one mol. of the hydroxypyrimidine compound.

15 The reactions of the steps (III) and (IV) can be preferably performed in the presence of a base. Examples of the bases include alkali metal carbonates such as sodium carbonate and potassium carbonate; alkali metal hydrogencarbonates such as sodium hydrogencarbonate; alkali metal hydroxides such as lithium hydroxide, sodium hydroxide and potassium hydroxide; alkali metal alkoxides such as sodium methoxide, sodium t-butoxide, potassium t-butoxide, and sodium t-pentoxide; and tertiary amines  
20 such as triethylamine, triisopropylamine, diisopropylethylamine, and pyridine. Preferred are sodium carbonate, potassium carbonate, potassium t-butoxide, sodium t-pentoxide, triethylamine, and pyridine. Particularly preferred are potassium carbonate, sodium t-pentoxide, and triethylamine. Most preferred are potassium carbonate and sodium t-pentoxide. The bases can be employed singly or in combination.

25 The base can be employed in an amount of, preferably 0.1 to 30 mol., more preferably 1 to 5 mol., per one mol. of the hydroxypyrimidine compound. The whole amount of the base can be incorporated in the reaction system before the reaction begins, or the base can be portionwise added to the reaction system after the reaction begins.

35 The reaction can be performed in the presence or absence of a solvent. There are no specific limitations



-15-

with respect to the solvent, so far as the solvent does not disturb the reaction. Examples of the solvents include water; ketones such as acetone, methyl ethyl ketone, and diethyl ketone; ethers such as diethyl ether and tetrahydrofuran; esters such as ethyl acetate, propyl acetate, and butyl acetate; nitriles such as acetonitrile and propionitrile; amides such as N,N-dimethylformamide and N-methylpyrrolidone; sulfoxides such as dimethylsulfoxide; ureas such as N,N'-dimethylimidazolinone. Preferred are acetone, tetrahydrofuran, ethyl acetate, butyl acetate, acetonitrile, N,N-dimethylformamide, and dimethylsulfoxide. Particularly preferred are ethyl acetate, butyl acetate and acetonitrile. Most preferred are butyl acetate and acetonitrile. The solvents can be employed singly or in combination.

The solvent can be employed in an amount of, preferably 0.01 to 100 liters, more preferably 0.5 to 5 liters, per one mole of the hydroxypyrimidine compound. The amount may vary depending on homogeneity and dispersability of the reaction mixture.


The reaction can be performed by reacting the hydroxypyrimidine compound and the organic sulfonyl halide or sulfonic anhydride in a solvent in the presence of a base with stirring under inert gas atmosphere. The base can be added portionwise. The reaction can be carried out at a temperature of, preferably -30 to 250°C, more preferably 0 to 150°C. There are no specific limitations with respect to the surrounding pressure.

The resulting product of the reaction, that is, the 2-(N-methyl-N-methanecarbonylamino)pyrimidine compound of the formula (3), can be isolated and purified according to the conventional procedures such as distillation, crystallization, recrystallization, and column chromatography.

The 2-(N-methyl-N-methanesulfonylamino)pyrimidine compound of the formula (3) and other pyrimidine com-

-16-

pounds of the formula (8) can be prepared from a hydroxypyrimidine compound of the formula (1) via a 2-substituted pyrimidine compound of the formula (6) in the following steps (V) and (VI):



In the formula (8), R has the same meaning as described above, and each of R<sup>1</sup> and R<sup>2</sup> independently is a hydrogen atom, an alkyl group, an alkylsulfonyl group, or arylsulfonyl group.

#### Step (V)

In the step (V), a hydroxypyrimidine compound of the formula (1) is reacted with a halogenating agent such as a chlorinating agent, an organic sulfonyl halide of the formula (2):



in which R' has the same meaning as above and X is a halogen atom, or an organic sulfonic anhydride of the formula (2a):



-17-

in which R' has the same meaning as above.

Examples of the halogenating agents include phosphorus oxychloride and thionyl chloride. The halogenating agents can be employed singly or in combination.

5 The halogenating agent can be employed in an amount of, preferably 0.1 to 50 mol., more preferably 1 to 20 mol., most preferably 1.5 to 10 mol., per one mol. of the hydroxypyrimidine compound.

10 Examples of the organic sulfonyl halides and sulfonic anhydrides are those described hereinbefore.

The organic sulfonyl halide or sulfonic anhydride can be employed in an amount of, preferably 0.1 to 20 mol., more preferably 0.5 to 5 mol., most preferably 1 to 2 mol., per one mol. of the hydroxypyrimidine compound.

15 The reaction can be performed in the presence or absence of a solvent. There are no specific limitations with respect to the solvent, so far as the solvent does not disturb the reaction. Examples of the solvents include aromatic hydrocarbons such as toluene; halogenated  
20 aromatic hydrocarbons such as chlorobenzene; nitrated hydrocarbons such as nitrobenzene; halogenated aliphatic hydrocarbons such as methylene chloride and 1,2-dichloroethane; amides such as N,N-dimethylformamide; water (not for a halogenating agent); nitriles such as acetonitrile  
25 and propionitrile; carboxylic acid esters such as ethyl acetate, propyl acetate, butyl acetate; ketones such as acetone, methyl ethyl ketone, diethyl ketone; and ethers such as diethyl ether and tetrahydrofuran. Preferred are butyl acetate, toluene, methylene chloride, acetonitrile,  
30 chlorobenzene, nitrobenzene, and N,N-dimethylformamide. The solvents can be employed singly or in combination.

The solvent can be employed in the reaction utilizing the halogenating agent in an amount of, preferably 0.01 to 10 liters, more preferably 0.1 to 2 liters, per  
35 one mole of the hydroxypyrimidine compound. The amount may vary depending on homogeneity and dispersability of

-18-

the reaction mixture.

The solvent can be employed in the reaction utilizing the sulfonyl chloride or sulfonic anhydride in an amount of, preferably 0.1 to 50 liters, more preferably 5 0.5 to 2 liters, per one mole of the hydroxypyrimidine compound. The amount may vary depending on homogeneity and dispersability of the reaction mixture.

The reaction can be carried out by reacting the hydroxypyrimidine compound and the halogenating agent, in 10 a solvent with stirring under inert gas atmosphere. The reaction can be carried out at a temperature of, preferably 0 to 200°C, more preferably 50 to 120°C. There are no specific limitations with respect to the surrounding pressure.

The reaction can be carried out by reacting the 15 hydroxypyrimidine compound and the sulfonyl halide or sulfonyl anhydride in a solvent with stirring under inert gas atmosphere. The reaction can be carried out at a temperature of, preferably -30 to 200°C, more preferably 0 20 to 50°C. There are no specific limitations with respect to the surrounding pressure.

The resulting product of the reaction, that is, a 2-substituted pyrimidine compound such as a chloropyrimidine compound or a sulfonyloxypyrimidine compound, can be 25 isolated and purified according to the conventional procedures such as distillation, crystallization, recrystallization, and column chromatography.

#### Step (VI)

30 In the step (VI), the 2-substituted pyrimidine compound, such as a chloropyrimidine compound or a sulfonyloxypyrimidine compound prepared in the step (V) is reacted with an amine compound having the formula (7):



-19-

wherein each of R<sup>1</sup> and R<sup>2</sup> is the same as above.

Examples of the groups of R<sup>1</sup> and R<sup>2</sup> include a hydrogen atom, alkyl groups such as methyl, ethyl, propyl, butyl, pentyl and hexyl; alkylsulfonyl groups such as methanesulfonyl; and arylsulfonyl groups such as benzene-sulfonyl and p-toluenesulfonyl.

The amine compound can be employed in an amount of, preferably 0.1 to 30 mol., more preferably 1 to 5 mol., per one mol. of the 2-substituted pyrimidine compound.

The reaction is preferably performed in the presence of a base. Examples of the bases are those described hereinbefore.

The base can be preferably employed in an amount of, preferably 0.1 to 30 mol., more preferably 1 to 5 mol., per one mol. of the 2-substituted pyrimidine compound.

The reaction can be performed in the presence or absence of a solvent. There are no specific limitations with respect to the solvent, so far as the solvent does not disturb the reaction. Examples of the solvents include water; ketones such as acetone, methyl ethyl ketone, and diethyl ketone; ethers such as diethyl ether and tetrahydrofuran; esters such as ethyl acetate, propyl acetate, and butyl acetate; nitriles such as acetonitrile and propionitrile; amides such as N,N-dimethylformamide and N-methylpyrrolidone; sulfoxides such as dimethylsulfoxide; ureas such as N,N'-dimethylimidazolidinone. Preferred are acetone, tetrahydrofuran, ethyl acetate, butyl acetate, acetonitrile, N,N-dimethylformamide, and dimethylsulfoxide. Particularly preferred are ethyl acetate, butyl acetate and acetonitrile. The solvents can be employed singly or in combination.

The solvent can be employed in an amount of, preferably 0.01 to 100 liters, more preferably 0.5 to 5 liters, per one mole of the 2-substituted pyrimidine compound.

The amount may vary depending on homogeneity and dispersability of the reaction mixture.

-20-

The reaction can be conducted by reacting the 2-substituted pyrimidine compound and the amine compound in a solvent in the presence of a base with stirring under inert gas atmosphere. The reaction can be carried out at a temperature of, preferably -20 to 250°C, more preferably 25 to 150°C. There are no specific limitations with respect to the surrounding pressure.

The reaction can be conducted in two separate liquid phases in the presence of a phase transfer catalyst. Examples of the phase transfer catalysts include tetramethylammonium chloride, tetramethylammonium bromide, tetraethylammonium fluoride, tetraethylammonium chloride, tetraethylammonium bromide, tetrapropylammonium bromide, tetrapropylammonium iodide, tetrabutylammonium fluoride, tetrabutylammonium chloride, tetrabutylammonium bromide, tetrabutylammonium iodide, tetrapentylammonium bromide, tetrahexylammonium bromide, tetraheptylammonium bromide, tetraoctylammonium bromide, benzyltrimethyltetradecylammonium chloride, benzyltriethylammonium chloride, phenyltrimethylammonium chloride, phenyltrimethylammonium iodide, and hexadecyltrimethylammonium chloride. Preferred are tetrabutylammonium chloride, tetrabutylammonium bromide, tetrabutylammonium iodide, benzyltriethylammonium chloride, and hexadecyltrimethylammonium chloride. Most preferred are tetrabutylammonium bromide, benzyltriethylammonium chloride, and hexadecyltrimethylammonium chloride.

The phase transfer catalyst can be employed in an amount of 0.01 to 0.5 mol., preferably 0.05 to 0.2 mol., per one mol. of the 2-substituted pyrimidine compound.

The resulting product of the reaction, that is, a 2-(N-methyl-N-methanesulfonylamino)pyrimidine compound of the formula (3) or other aminopyrimidine compounds of formula (8), can be isolated and purified according to the conventional procedures such as distillation, crystallization, recrystallization, or column chromatography.

-21-

The present invention is further described by the following non-limiting examples.

[Example 1] Preparation of 4-(4-fluorophenyl)-6-isopropyl-5-methoxycarbonyl-3,4,2(1H)-dihydropyrimidinone

In a 500 mL-volume glass flask equipped with a stirrer, a thermometer and a reflux condenser were placed 28.8 g (0.2 mol.) of methyl isobutyrylacetate, 24.8 g (0.2 mol.) of 4-fluorobenzaldehyde, 21.0 g (0.35 mol.) of urea, 200 mg (2 mmol.) of copper(I) chloride, 2 mL of sulfuric acid, and 200 mL of methanol. The content of the flask was heated to 64-65°C for 24 hours under reflux with stirring, to carry out the reaction. There was precipitated crystalline product. The crystalline product was collected on a filter paper and washed with methanol to obtain 49.7 g of 4-(4-fluorophenyl)-6-isopropyl-5-methoxycarbonyl-3,4,2(1H)-dihydropyrimidinone as a colorless crystalline product having the below-mentioned characteristics. The yield was 85% (based on the amount of methyl isobutyrylacetate).

m.p.: 223-225°C

UV  $\lambda_{\max}$  (CH<sub>3</sub>CN, nm): 194.3, 278.6

IR (KBr, cm<sup>-1</sup>): 3296, 3229, 3137, 2963, 1685, 1629, 1504, 1225, 1097.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>,  $\delta$  (ppm)): 1.14 (6H, dd, J=6.8, 6.9Hz), 3.52 (3H, s), 4.0-4.2 (1H, m), 5.15 (1H, d, J=3.4Hz), 7.1-7.2 (2H, m), 7.2-7.3 (2H, m), 7.76 (1H, d, J=3.2Hz), 8.91 (1H, s).

HRMS: 292.1247 (theoretical value (C<sub>15</sub>H<sub>17</sub>FN<sub>2</sub>O<sub>3</sub> (M<sup>+</sup>)) 292.1223)

[Example 2] Preparation of 4-(4-fluorophenyl)-6-isopropyl-5-methoxycarbonyl-3,4,2(1H)-dihydropyrimidinone

The procedures of Example 1 were repeated except for

-22-

replacing 200 mg (2 mmol.) of copper(I) chloride with 5.41 g (20 mmol.) of iron(III) chloride·hexahydrate. There was obtained 35.6 g of 4-(4-fluorophenyl)-6-isopropyl-5-methoxycarbonyl-3,4-2(1H)-dihydropyrimidinone. The yield was 61% (based on the amount of methyl isobutyryl-  
5 acetate).

[Example 3] Preparation of 4-(4-fluorophenyl)-2-hydroxy-6-isopropyl-5-methoxycarbonylpyrimidine

10 In a 50 mL-volume glass flask equipped with a stirrer and a thermometer was placed 11 mL (144 mmol.) of nitric acid (60-61%, sp.gr.: 1.38). To the nitric acid was slowly added at room temperature 4.00 g (13.7 mmol.) of 4-(4-fluorophenyl)-6-isopropyl-5-methoxycarbonyl-3,4-  
15 2(1H)-dihydropyrimidinone prepared in the same manner as in Example 1, and the mixture was subjected to reaction for 30 minutes at room temperature. After the reaction was complete, the reaction mixture was neutralized by placing the mixture in 140 mL of saturated aqueous sodium  
20 hydrogen carbonate solution. The reaction mixture was then extracted with ethyl acetate. The organic liquid portion was separated and concentrated under reduced pressure. The residue was crystallized from toluene. The crystalline product was collected on a filter and  
25 washed with toluene to obtain 3.64 g of 4-(4-fluorophenyl)-2-hydroxy-6-isopropyl-5-methoxycarbonylpyrimidine as a colorless crystalline product having the below-mentioned characteristics. The yield was 92% (based on the amount of 4-(4-fluorophenyl)-6-isopropyl-5-  
30 methoxycarbonyl-3,4-2(1H)-dihydropyrimidinone).

m.p.: 193°C (decomposed)

UV  $\lambda_{\max}$  (CH<sub>3</sub>CN, nm): 196.6, 243.2, 317.9

IR (KBr, cm<sup>-1</sup>): 2991, 2887, 1717, 1653, 1589, 1433,  
35 1280, 1223.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>,  $\delta$  (ppm)): 1.23 (6H, d, J=6.8Hz),



-23-

3.0-3.2 (1H, m), 3.56 (3H, s), 7.3-7.4  
(2H, m), 7.5-7.6 (2H, m), 12.25 (1H, brs).

HRMS: 290.1054 (theoretical value ( $C_{15}H_{15}FN_2O_3$  (M+))  
290.1067)

5

[Example 4] Preparation of 4-(4-fluorophenyl)-2-hydroxy-6-isopropyl-5-methoxycarbonylpyrimidine

In a 50 mL-volume glass flask equipped with a stirrer and a thermometer were placed 2.92 g (10 mmol.) of 4-(4-fluorophenyl)-6-isopropyl-5-methoxycarbonyl-3,4-dihydropyrimidinone prepared in the same manner as in Example 1 and 5 mL of acetic acid. To the mixture was slowly added 3.74 mL (50 mmol.) of nitric acid (60-61%, sp.gr.: 1.38). To the mixture was further added 0.07 g (1 mmol.) of sodium nitrite, and the reaction was carried out for one hour at room temperature. After the reaction was complete, the reaction mixture was neutralized by placing the mixture in 50 mL of saturated aqueous sodium hydrogen carbonate solution. The reaction mixture was then extracted with ethyl acetate. The organic liquid portion was separated and concentrated under reduced pressure. The residue was crystallized from toluene. The crystalline product was collected on a filter and washed with toluene to obtain 2.61 g of 4-(4-fluorophenyl)-2-hydroxy-6-isopropyl-5-methoxycarbonylpyrimidine as a colorless crystalline product. The yield was 90% (based on the amount of 4-(4-fluorophenyl)-6-isopropyl-5-methoxycarbonyl-3,4-dihydropyrimidinone).

30 [Example 5] Preparation of 4-(4-fluorophenyl)-2-hydroxy-6-isopropyl-5-methoxycarbonylpyrimidine

In a 200 mL-volume glass flask equipped with a stirrer and a thermometer was placed 54.0 g (735 mmol.) of nitric acid (60-61%, sp.gr.: 1.38). To the nitric acid was slowly added at room temperature 30.6 g (105 mmol.) of 4-(4-fluorophenyl)-6-isopropyl-5-methoxycarbonyl-3,4-

-24-

2 (1H)-dihydropyrimidinone prepared in the same manner as in Example 1, and the mixture was subjected to reaction for 30 minutes at room temperature. After the reaction was complete, the reaction mixture was poured into 162 mL of water. The aqueous mixture was neutralized by adding 61 g of aqueous sodium hydroxide solution (48 wt.%) to precipitate a crystalline product. The crystalline product was collected by filtration and dried to obtain 27.6 g of 4-(4-fluorophenyl)-2-hydroxy-6-isopropyl-5-methoxycarbonylpyrimidine as a colorless crystalline product. The yield was 91% (based on the amount of 4-(4-fluorophenyl)-6-isopropyl-5-methoxycarbonyl-3,4-2(1H)-dihydropyrimidinone).

[Example 6] Preparation of 4-(4-fluorophenyl)-2-hydroxy-6-isopropyl-5-methoxycarbonylpyrimidine

In a 2 L-volume glass flask equipped with a stirrer and a thermometer was placed 323.3 g (3.09 mol.) of nitric acid (60-61%, sp.gr.: 1.38). The concentrated nitric acid was then cooled to 10°C. To the nitric acid was added 2.36 g (34.2 mmol.) of sodium nitrite, and was further added slowly 100 g (342 mmol.) of 4-(4-fluorophenyl)-6-isopropyl-5-methoxycarbonyl-3,4-2(1H)-dihydropyrimidinone prepared in the same manner as in Example 1. The mixture was subjected to reaction for 2 hours at a temperature of 10-12°C. After the reaction was complete, 970 mL of water was poured into the reaction mixture. The aqueous mixture was then neutralized by adding 257 g of aqueous sodium hydroxide solution (48 wt.%) to precipitate a crystalline product. The crystalline product was collected by filtration and dried to obtain 93.3 g of 4-(4-fluorophenyl)-2-hydroxy-6-isopropyl-5-methoxycarbonylpyrimidine as a colorless crystalline product. The yield was 94% (based on the amount of 4-(4-fluorophenyl)-6-isopropyl-5-methoxycarbonyl-3,4-2(1H)-

-25-

dihydropyrimidinone).

[Example 7] Preparation of 4-(4-fluorophenyl)-6-isopropyl-5-methoxycarbonyl-2-(N-methyl-N-methanesulfonylamino)pyrimidine

5 In a 200 mL-volume glass flask equipped with a stirrer, a thermometer and a reflux condenser were placed 5.81 g (20 mmol.) of 4-(4-fluorophenyl)-2-hydroxy-6-isopropyl-5-methoxycarbonylpyrimidine, 3.59 g (26 mmol.) of  
10 potassium carbonate (available from Asahi Glass Works, Co., Ltd., Lot No. 1111632, particle size distribution: 75-250  $\mu\text{m}$ : 14%, 75  $\mu\text{m}$  pass: 86%), and 40 mL of butyl acetate. To the mixture was slowly added 4.19 g (22 mmol.) of p-toluenesulfonyl chloride under stirring, and the  
15 reaction was carried out at 40°C for 4 hours. Subsequently, the reaction mixture was cooled to room temperature. To the cooled reaction mixture were added 2.84 g (26 mmol.) of N-methylmethanesulfonamide and 4.15 g (30  
20 mmol.) of potassium carbonate (same as above). The mixture was heated to 110-125°C for 2 hours under refluxing to carry out a reaction. After the reaction was complete, the mixture was cooled to room temperature. To the cooled mixture were added 25 mL of water and 20 mL of acetone, and the organic liquid portion was separated.  
25 The organic liquid portion was washed with a saturated aqueous sodium chloride solution and dried over anhydrous magnesium sulfate. The dry organic liquid portion was filtered and concentrated under reduced pressure. The residue was crystallized from heptane, to obtain 6.58 g  
30 of 4-(4-fluorophenyl)-6-isopropyl-5-methoxycarbonyl-2-(N-methyl-N-methanesulfonylamino)pyrimidine as a pale yellow crystalline product. The yield was 86% (based on the amount of 4-(4-fluorophenyl)-2-hydroxy-6-isopropyl-5-methoxycarbonylpyrimidine).

35

[Example 8] Preparation of 4-(4-fluorophenyl)-6-isopro-

-26-

pyl-5-methoxycarbonyl-2-(N-methyl-N-methanesulfonyl-amino)pyrimidine

In a 1000 mL-volume glass flask equipped with a stirrer, a thermometer and a reflux condenser were placed  
5 50.0 g (172 mmol.) of 4-(4-fluorophenyl)-2-hydroxy-6-isopropyl-5-methoxycarbonylpyrimidine, 20.8 g (189 mmol.) of sodium t-pentoxide, and 344 mL of acetonitrile, and the resulting mixture was stirred at 0-10°C for 30 minutes. To the mixture was slowly added 36.1 g (189 mmol.) of p-  
10 toluenesulfonyl chloride, and the reaction was carried out at for 5 hours at room temperature. Subsequently, the reaction mixture was cooled to a temperature of 0-10°C. To the cooled reaction mixture were added 28.2 g (258 mmol.) of N-methylmethanesulfonamide and 26.5 g (241  
15 mmol.) of sodium t-pentoxide. The mixture was kept at 0-10°C for one hour and then heated to 75-82°C for 2 hours under refluxing, to carry out a reaction. After the reaction was complete, the mixture was cooled to room temperature. To the cooled mixture was added 344 mL of  
20 water. The aqueous mixture was cooled to 0-10°C and stirred for one hour, precipitating a crystalline product. The crystalline product was collected by filtration and dried, to obtain 45.3 g of 4-(4-fluorophenyl)-6-isopropyl-5-methoxycarbonyl-2-(N-methyl-N-methanesulfonyl-  
25 ylamino)pyrimidine as a pale yellow crystalline product. The yield was 68% (based on the amount of 4-(4-fluorophenyl)-2-hydroxy-6-isopropyl-5-methoxycarbonyl-pyrimidine).

30 [Example 9] Preparation of 4-(4-fluorophenyl)-6-isopropyl-5-methoxycarbonyl-2-(N-methyl-N-methanesulfonyl-amino)pyrimidine from methyl isobutyrylacetate, 4-fluorobenzaldehyde and urea

1) In a 200 L-volume glass-lined reaction vessel  
35 equipped with a stirrer, a thermometer and a reflux condenser were placed 24.4 kg (169 mol.) of methyl iso-

-27-

butyrylacetate, 20.0 kg (161 mol.) of 4-fluorobenzaldehyde, 16.9 kg (282 mol.) of urea, 0.2 kg (2 mol.) of copper(I) chloride, 3.0 kg of sulfuric acid, and 80.4 kg of methanol. The mixture was heated to 64-66°C for 20  
5 hours under refluxing, to carry out reaction. After the reaction was complete, the reaction mixture was cooled to room temperature, to precipitate a crystalline product. The crystalline product was collected on a filter paper and washed with methanol to obtain 43.4 kg of 4-(4-  
10 fluorophenyl)-6-isopropyl-5-methoxycarbonyl-3,4-2(1H)-dihydropyrimidinone as a colorless crystalline product.

2) In a 200 L-volume glass-lined reaction vessel equipped with a stirrer and a thermometer were placed 62.5 kg (615.6 mol.) of diluted nitric acid and 0.5 kg  
15 (6.8 mol.) of sodium nitrite. To the mixture was slowly added under chilling 20.0 kg (68.4 mmol.) of the 4-(4-fluorophenyl)-6-isopropyl-5-methoxycarbonyl-3,4-2(1H)-dihydropyrimidinone prepared as above. The resulting mixture was subjected to reaction at a low temperature  
20 (10°C). After the reaction was complete, the reaction mixture was neutralized by addition of an aqueous methanol solution of sodium hydroxide. Subsequently, an aqueous sodium hydroxide solution was added to the mixture. The resulting mixture was placed under reduced pressure  
25 to distill methanol off. To the residue were added 96.5 kg of acetone and 96.5 kg of water. The aqueous residue was then neutralized by addition of acetic acid to precipitate a crystalline product. The crystalline product was collected on a filter paper and washed with a acetone/water mixture, to give 17.9 kg of 4-(4-fluorophenyl)-2-hydroxy-6-isopropyl-5-methoxycarbonylpyrimidine.  
30

3) In a 200 L-volume glass-lined reaction vessel equipped with a stirrer, a thermometer and a reflux condenser were placed 17.9 kg (62.0 mol.) of 4-(4-fluorophenyl)-2-hydroxy-6-isopropyl-5-methoxycarbonylpyrimidine  
35 prepared as above, 107.7 kg of butyl acetate, 11.1 kg

-28-

(80.3 mol.) of potassium carbonate (available from Asahi Glass Works, Co., Ltd., Lot No. 1111632, particle size distribution: 75-250  $\mu\text{m}$ : 14%, 75  $\mu\text{m}$  pass: 86%), and 12.9 kg (67.7 mol.) of p-toluenesulfonyl chloride. The mixture was heated at 60°C for 2 hours, to carry out reaction. Subsequently, the reaction mixture was cooled to room temperature. To the cooled mixture were added 8.8 kg (80.6 mol.) of N-methylmethanesulfonamide and 12.9 kg (93.3 mol.) of potassium carbonate, and the resulting mixture was heated at 122-125°C for 3 hours, for carrying reaction. After the reaction was complete, the reaction mixture was cooled to room temperature. To the cooled mixture were added acetone and water, and the organic liquid portion was separated. The organic liquid portion was then washed successively with aqueous sodium hydroxide solution (3 wt.%) and a saturated aqueous sodium chloride solution. The washed organic liquid portion was concentrated under reduced pressure. Isopropyl alcohol and water were added to the residue, resulting in precipitation of a crystalline product. The crystalline product was filtered on a filter paper and washed with isopropyl alcohol. The washed crystalline product and 85.7 kg of acetone were placed in a 200 L-volume glass lined reaction vessel equipped with a stirrer, a thermometer and a reflux condenser. The mixture was stirred at 50-55°C, to dissolve the crystalline product in acetone. The insoluble was removed with a line filter. Subsequently, 58.3 kg of water was added to the solution, to precipitate a crystalline product. The crystalline product was collected on a filter paper and washed with an acetone/water mixture, to give 19.5 kg of 4-(4-fluorophenyl)-6-isopropyl-5-methoxycarbonyl-2-(N-methyl-N-methanesulfonylamino)pyrimidine.

[Example 10] Preparation of 2-chloro-4-(4-fluorophenyl)-6-isopropyl-5-methoxycarbonylpyrimidine

-29-

In a 25 mL-volume glass flask equipped with a stirrer, a thermometer and a reflux condenser were placed 1.00 g (3.43 mmol.) of 4-(4-fluorophenyl)-2-hydroxy-6-isopropyl-5-methoxycarbonylpyrimidine and 3.4 mL (3.7 mmol.) of phosphorus oxychloride. The mixture was heated to 100-106°C for 1.5 hours under refluxing, to carry out reaction. After the reaction was complete, the reaction mixture was cooled to room temperature, and poured into an ice/water mixture. The resulting aqueous mixture was neutralized with a saturated aqueous sodium hydrogen carbonate solution. The neutralized aqueous mixture was extracted with ethyl acetate. The ethyl acetate portion was separated, washed with a saturated aqueous sodium chloride solution, and dried over anhydrous magnesium sulfate. The dried ethyl acetate portion was filtered and concentrated under reduced pressure, to obtain 1.03 g of 2-chloro-4-(4-fluorophenyl)-6-isopropyl-5-methoxycarbonylpyrimidine as a colorless crystalline product having the below-mentioned characteristics. The yield was 97% (based on the amount of 4-(4-fluorophenyl)-2-hydroxy-6-isopropyl-5-methoxycarbonylpyrimidine).

m.p.: 99-101°C

UV  $\lambda_{\max}$  (CH<sub>3</sub>CN, nm): 194.7, 276.5

IR (KBr, cm<sup>-1</sup>): 2980, 1728, 1542, 1508, 1227, 1086.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>,  $\delta$  (ppm)): 1.33 (6H, d, J=6.8Hz),  
3.1-3.2 (1H, m), 3.76 (3H, s), 7.15 (2H, t, J=8.5Hz), 7.6-7.7 (2H, m).

HRMS: 308.0695 (theoretical value (C<sub>15</sub>H<sub>14</sub>ClFN<sub>2</sub>O<sub>2</sub> (M<sup>+</sup>))  
308.0728)

[Example 11] Preparation of 2-chloro-4-(4-fluorophenyl)-6-isopropyl-5-methoxycarbonylpyrimidine

In a 25 mL-volume glass flask equipped with a stirrer, a thermometer and a reflux condenser were placed 1.00 g (3.43 mmol.) of 4-(4-fluorophenyl)-2-hydroxy-6-

-30-

isopropyl-5-methoxycarbonylpyrimidine, 0.5 mL (3.9 mmol.) of thionyl chloride, 3.44 mL of toluene, and 0.11 mL of N,N-dimethylformamide. The mixture was heated to 80°C for 3 hours, to carry out reaction. After the reaction was complete, the reaction mixture was cooled to room temperature, and poured into an ice/water mixture. The resulting aqueous mixture was neutralized with a saturated aqueous sodium hydrogen carbonate solution. The neutralized aqueous mixture was extracted with ethyl acetate. The ethyl acetate portion was separated, washed with a saturated aqueous sodium chloride solution, and dried over anhydrous magnesium sulfate. The dried ethyl acetate portion was filtered and concentrated under reduced pressure, to obtain 0.80 g of 2-chloro-4-(4-fluorophenyl)-6-isopropyl-5-methoxycarbonylpyrimidine as a colorless crystalline product. The yield was 76% (based on the amount of 4-(4-fluorophenyl)-2-hydroxy-6-isopropyl-5-methoxycarbonylpyrimidine).

[Example 12] Preparation of 4-(4-fluorophenyl)-6-isopropyl-5-methoxycarbonyl-2-(N-methyl-N-methanesulfonylamino)pyrimidine

In a 25 mL-volume glass flask equipped with a stirrer, a thermometer and a reflux condenser were placed 546 mg (5 mmol.) of N-methylmethanesulfonamide, 551 mg (5 mmol.) of sodium t-pentoxide, 10 mL of acetonitrile, and 309 mg (1 mmol.) of 2-chloro-4-(4-fluorophenyl)-6-isopropyl-5-methoxycarbonylpyrimidine. The mixture was heated to 81-82°C for 3 hours under refluxing, to carry out reaction. After the reaction was complete, the reaction mixture was cooled to room temperature. To the cooled mixture was added 10 mL of water, and the aqueous mixture was extracted with ethyl acetate. The ethyl acetate portion was separated, and dried over anhydrous magnesium sulfate. The dried ethyl acetate portion was filtered and concentrated under reduced pressure. The residue was



-31-

purified by silica gel column chromatography (column: Wako Gel C-200, eluent: hexane/ethyl acetate (2:1, volume ratio)). There was obtained 339 mg of 4-(4-fluorophenyl)-6-isopropyl-5-methoxycarbonyl-2-(N-methyl-N-methanesulfonylamino)pyrimidine. The yield was 89% (based on the amount of 2-chloro-4-(4-fluorophenyl)-6-isopropyl-5-methoxycarbonylpyrimidine).

[Example 13] Preparation of 4-(4-fluorophenyl)-6-isopropyl-5-methoxycarbonyl-2-methanesulfonyloxypyrimidine

In a 100 mL-volume glass flask were placed 10.0 g (34.4 mmol.) of 4-(4-fluorophenyl)-2-hydroxy-6-isopropyl-5-methoxycarbonylpyrimidine, 5.22 g (58.5 mmol.) of triethylamine, and 34 mL of acetonitrile. The mixture in the flask was chilled to 0-5°C in an ice bath. To the chilled mixture was slowly added 5.12 g (44.7 mmol.) of methanesulfonyl chloride, and the resulting mixture was subjected to reaction at 20-25°C for 2 hours. After the reaction was complete, to the reaction mixture was added 60 mL of water. The aqueous reaction mixture was extracted with toluene and the toluene portion was separated. The toluene portion was washed with a saturated aqueous sodium chloride solution and dried over anhydrous magnesium sulfate. The dried mixture was filtered and concentrated under reduced pressure. The residue was crystallized from methanol, to give 11.3 g of 4-(4-fluorophenyl)-6-isopropyl-5-methoxycarbonyl-2-methanesulfonyloxypyrimidine as a colorless crystalline product having the below-mentioned characteristics. The yield was 89% (based on the amount of 4-(4-fluorophenyl)-2-hydroxy-6-isopropyl-5-methoxycarbonylpyrimidine).

m.p.: 110-111°C

UV  $\lambda_{\max}$  (CH<sub>3</sub>CN, nm): 193.7, 276.8

IR (KBr, cm<sup>-1</sup>): 2980, 1724, 1562, 1391, 1250, 1175, 1079, 971.

-32-

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ ,  $\delta$  (ppm)): 1.33 (6H, d,  $J=6.6\text{Hz}$ ),  
3.20 (1H, m), 3.60 (3H, s), 7.1-7.2  
(2H, s), 7.6-7.8 (2H, m).

HRMS: 368.0842 (theoretical value ( $\text{C}_{15}\text{H}_{17}\text{FN}_2\text{O}_5\text{S}(\text{M}^+)$ )  
5 368.0892)

[Example 14] Preparation of 4-(4-fluorophenyl)-6-isopropyl-5-methoxycarbonyl-2-(p-toluenesulfonyloxy)pyrimidine

In a 200 mL-volume glass flask were placed 27.6 g  
10 (95.1 mmol.) of 4-(4-fluorophenyl)-2-hydroxy-6-isopropyl-5-methoxycarbonylpyrimidine, 12.5 g (123 mmol.) of triethylamine, and 95 mL of acetonitrile. The mixture of the flask was chilled to 0-5°C in an ice bath. To the chilled mixture was slowly added 20.0 g (105 mmol.) of p-  
15 toluenesulfonyl chloride, and the resulting mixture was subjected to reaction at 20-25°C for one hour. After the reaction was complete, to the reaction mixture was added 95 mL of water. The aqueous reaction mixture was extracted with toluene and the toluene portion was separated.  
20 The toluene portion was washed with a saturated aqueous sodium chloride solution and dried over anhydrous magnesium sulfate. The dried mixture was filtered and concentrated under reduced pressure. The residue was crystallized from methanol, to give 35.9 g of 4-(4-  
25 fluorophenyl)-6-isopropyl-5-methoxycarbonyl-2-(p-toluenesulfonyloxy)pyrimidine as a colorless crystalline product having the below-mentioned characteristics. The yield was 85% (based on the amount of 4-(4-fluorophenyl)-2-hydroxy-6-isopropyl-5-methoxycarbonylpyrimidine).

30

m.p.: 94-96°C

UV  $\lambda_{\text{max}}$  ( $\text{CH}_3\text{CN}$ , nm): 194.9, 275.2

IR (KBr,  $\text{cm}^{-1}$ ): 2961, 1734, 1539, 1389, 1352, 1247,  
1090, 980.

35

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ ,  $\delta$  (ppm)): 1.23 (6H, d,  $J=6.8\text{Hz}$ ),  
2.45 (3H, s), 3.0-3.2 (1H, m), 3.74

-33-

J=8.5Hz).

HRMS: 444.1155 (theoretical value ( $C_{32}H_{21}FN_2O_5S$  (M+))  
444.1194)

5 [Example 15] Preparation of 4-(4-fluorophenyl)-6-isopropyl-5-methoxycarbonyl-2-benzenesulfonyloxypyrimidine

The procedures of Example 13 were repeated except for replacing p-toluenesulfonyl chloride with 18.5 g (105 mmol.) of benzenesulfonyl chloride.

There was obtained 39.3 g of 4-(4-fluorophenyl)-6-isopropyl-5-methoxycarbonyl-2-benzenesulfonyloxypyrimidine as a pale yellow crystalline product having the below-mentioned characteristics. The yield was 96% (based on the amount of 4-(4-fluorophenyl)-2-hydroxy-6-isopropyl-5-methoxycarbonylpyrimidine).

$^1H$ -NMR ( $CDCl_3$ ,  $\delta$  (ppm)): 1.21 (6H, d, J=6.4Hz),  
3.0-3.1 (1H, m), 3.73 (3H, s), 7.1-7.2  
(2H, m), 7.5-7.7 (5H, m), 8.1-8.2 (2H, m).

[Example 16] Preparation of 4-(4-fluorophenyl)-6-isopropyl-5-methoxycarbonyl-2-(2,4,6-trimethylbenzenesulfonyloxy)pyrimidine

The procedures of Example 13 were repeated except for replacing p-toluenesulfonyl chloride with 23.0 g (105 mmol.) of 2,4,6-trimethylbenzenesulfonyl chloride.

There was obtained 37.7 g of 4-(4-fluorophenyl)-6-isopropyl-5-methoxycarbonyl-2-(2,4,6-trimethylbenzenesulfonyloxy)pyrimidine as a pale yellow crystalline product having the below-mentioned characteristics. The yield was 84% (based on the amount of 4-(4-fluorophenyl)-2-hydroxy-6-isopropyl-5-methoxycarbonylpyrimidine).

$^1H$ -NMR ( $CDCl_3$ ,  $\delta$  (ppm)): 1.17 (6H, d, J=6.8Hz),

-34-

2.34 (3H, s), 2.67 (6H, s), 3.0-3.1 (1H, m), 3.73 (3H, s), 7.00 (2H, s), 7.0-7.2 (2H, m), 7.4-7.5 (2H, m).

5 [Example 17] Preparation of 4-(4-fluorophenyl)-6-isopropyl-5-methoxycarbonyl-2-(2,4,6-triisopropylbenzenesulfonyloxy)pyrimidine

The procedures of Example 13 were repeated except for replacing p-toluenesulfonyl chloride with 31.8 g (105  
10 mmol.) of 2,4,6-triisopropylbenzenesulfonyl chloride.

There was obtained 47.1 g of 4-(4-fluorophenyl)-6-isopropyl-5-methoxycarbonyl-2-(2,4,6-triisopropylbenzenesulfonyloxy)pyrimidine as a pale yellow crystalline product having the below-mentioned characteristics. The  
15 yield was 89% (based on the amount of 4-(4-fluorophenyl)-2-hydroxy-6-isopropyl-5-methoxycarbonylpyrimidine).

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ ,  $\delta$  (ppm)): 1.12 (6H, d,  $J=6.6\text{Hz}$ ),  
1.19 (12H, d,  $J=6.8\text{Hz}$ ), 1.27 (6H, d,  
20  $J=7.1\text{Hz}$ ), 2.8-2.95 (1H, m), 2.95-3.1 (1H, m), 3.73 (3H, s), 4.1-4.3 (2H, m), 7.0-7.1 (2H, m), 7.20 (2H, s), 7.4-7.5 (2H, m).

[Example 18] Preparation of 4-(4-fluorophenyl)-6-isopropyl-5-methoxycarbonyl-2-(p-methoxybenzenesulfonyloxy)pyrimidine  
25

The procedures of Example 13 were repeated except for replacing p-toluenesulfonyl chloride with 21.7 g (105 mmol.) of p-methoxybenzenesulfonyl chloride.

30 There was obtained 39.9 g of 4-(4-fluorophenyl)-6-isopropyl-5-methoxycarbonyl-2-(p-methoxybenzenesulfonyloxy)pyrimidine as a colorless crystalline product having the below-mentioned characteristics. The yield was 91% (based on the amount of 4-(4-fluorophenyl)-2-hydroxy-6-isopropyl-5-methoxycarbonylpyrimidine).  
35

-35-

<sup>1</sup>H-NMR (CDCl<sub>3</sub>, δ (ppm)): 1.25 (6H, d, J=6.8Hz),  
3.0-3.2 (1H, m), 3.74 (3H, s), 3.88 (3H,  
s), 6.99 (2H, dd, J=2.0, 9.0Hz), 7.0-7.2  
(2H, m), 7.5-7.7 (2H, m), 8.07 (2H, dd,  
J=2.2, 9.0Hz).

[Example 19] Preparation of 4-(4-fluorophenyl)-6-isopro-  
pyl-5-methoxycarbonyl-2-(p-chlorobenzenesulfonyloxy)-  
pyrimidine

The procedures of Example 13 were repeated except  
for replacing p-toluenesulfonyl chloride with 22.2 g (105  
mmol.) of p-chlorobenzenesulfonyl chloride.

There was obtained 39.9 g of 4-(4-fluorophenyl)-6-  
isopropyl-5-methoxycarbonyl-2-(p-chlorobenzenesulfonyl-  
oxy)pyrimidine as a colorless crystalline product having  
the below-mentioned characteristics. The yield was 89%  
(based on the amount of 4-(4-fluorophenyl)-2-hydroxy-6-  
isopropyl-5-methoxycarbonylpyrimidine).

<sup>1</sup>H-NMR (CDCl<sub>3</sub>, δ (ppm)): 1.23 (6H, d, J=6.6Hz),  
3.0-3.2 (1H, m), 3.74 (3H, s), 7.1-7.2  
(2H, m), 7.5-7.7 (4H, m), 8.0-8.1 (2H, m).

[Example 20] Preparation of 4-(4-fluorophenyl)-6-isopro-  
pyl-5-methoxycarbonyl-2-(2-nitrobenzenesulfonyloxy)-  
pyrimidine

The procedures of Example 13 were repeated except  
for replacing p-toluenesulfonyl chloride with 23.3 g (105  
mmol.) of 2-nitrobenzenesulfonyl chloride.

There was obtained 28.0 g of 4-(4-fluorophenyl)-6-  
isopropyl-5-methoxycarbonyl-2-(2-nitrobenzenesulfonyl-  
oxy)pyrimidine as an opaque crystalline product having  
the below-mentioned characteristics. The yield was 62%  
(based on the amount of 4-(4-fluorophenyl)-2-hydroxy-6-  
isopropyl-5-methoxycarbonylpyrimidine).

-36-

<sup>1</sup>H-NMR (CDCl<sub>3</sub>, δ (ppm)): 1.17 (6H, d, J=6.8Hz),  
3.0-3.2 (1H, m), 3.75 (3H, s), 7.1-7.2  
(2H, m), 7.5-7.6 (2H, m), 7.7-8.0 (3H, m),  
8.33 (1H, dd, J=1.7, 8.1Hz).

5

[Example 21] Preparation of 4-(4-fluorophenyl)-6-isopropyl-5-methoxycarbonyl-2-(N-methyl-N-methanesulfonylamino)pyrimidine

In a 25 mL-volume glass flask equipped with a stirrer, a thermometer and a reflux condenser were placed 196 mg (1.8 mmol.) of N-methylmethanesulfonamide, 198 mg (1.8 mmol.) of sodium t-pentoxide, 7.5 mL of acetonitrile, and 667 mg (1.5 mmol.) of 4-(4-fluorophenyl)-6-isopropyl-5-methoxycarbonyl-2-(p-toluenesulfonyloxy)pyrimidine. The mixture was heated to 81-82°C for 1.5 hours under refluxing, to carry out reaction. After the reaction was complete, the reaction mixture was cooled to room temperature. To the cooled mixture was added 10 mL of water, and the aqueous mixture was extracted with ethyl acetate. The ethyl acetate portion was separated, and dried over anhydrous magnesium sulfate. The dried ethyl acetate portion was filtered and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (column: Wako Gel C-200, eluent: hexane/ethyl acetate (2:1, volume ratio)). There was obtained 428 mg of 4-(4-fluorophenyl)-6-isopropyl-5-methoxycarbonyl-2-(N-methyl-N-methanesulfonylamino)pyrimidine. The yield was 75% (based on the amount of 4-(4-fluorophenyl)-6-isopropyl-5-methoxycarbonyl-2-(p-toluenesulfonyloxy)pyrimidine).

[Example 22] Preparation of (2-amino-4-(4-fluorophenyl)-6-isopropyl-5-methoxycarbonyl)pyrimidine

In a 25 mL-volume glass flask equipped with a stirrer, a thermometer and a gas inlet were placed under ice-chilling 1.00 g (2.71 mmol.) of 4-(4-fluorophenyl)-6-iso-

-37-

propyl-5-methoxycarbonyl-2-methanesulfonyloxypyrimidine and 8.1 mL of tetrahydrofuran. The mixture was stirred at room temperature for 12 hours under gaseous ammonia atmosphere, for carrying out reaction. After the reaction was complete, 10 mL of water was added to the reaction mixture. The aqueous mixture was then subjected to extraction with toluene. The toluene portion was separated, washed with a saturated aqueous sodium chloride solution, and dried over anhydrous magnesium sulfate. The dried toluene portion was filtered and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (column: Wako Gel C-200, eluent: hexane/ethyl acetate (2:1, volume ratio)). There was obtained 0.63 g of 2-amino-4-(4-fluorophenyl)-6-isopropyl-5-methoxycarbonylpyrimidine. The yield was 80% (based on the amount of 4-(4-fluorophenyl)-6-isopropyl-5-methoxycarbonyl-2-methanesulfonyloxypyrimidine).

6-isopropyl-5-methoxycarbonylpyrimidine

[Example 23] Preparation of (4-(4-fluorophenyl)-6-isopropyl-5-methoxycarbonyl-2-N-methylaminopyrimidine

In a 50 mL-volume glass flask equipped with a stirrer, a thermometer and a dropping funnel was placed 6.00 g (16.3 mmol.) of 4-(4-fluorophenyl)-6-isopropyl-5-methoxycarbonyl-2-methanesulfonyloxypyrimidine. Into the flask was slowly dropped under ice-chilling 5.06 g (65.2 mmol) of aqueous 40 wt.% methylamine solution. The resulting mixture was stirred for one hour at the same temperature for carrying out reaction. After the reaction was complete, 16 mL of water was added to the reaction mixture. The aqueous mixture was then subjected to extraction with toluene. The toluene portion was separated, washed with a saturated aqueous sodium chloride solution, and dried over anhydrous magnesium sulfate. The dried toluene portion was filtered and concentrated under reduced pressure to give 4.81 g of 4-(4-fluorophenyl)-6-

-38-

isopropyl-5-methoxycarbonyl-2-N-methylaminopyrimidine. The yield was 97% (based on the amount of 4-(4-fluorophenyl)-6-isopropyl-5-methoxycarbonyl-2-methanesulfonyloxy-pyrimidine).

5

[Example 24] Preparation of 4-(4-fluorophenyl)-6-isopropyl-5-methoxycarbonyl-2-trifluoromethanesulfonyloxy-pyrimidine

In a 300 mL-volume glass flask equipped with a stirrer, a thermometer and a reflux condenser were placed 8.7 g (30.0 mmol.) of 4-(4-fluorophenyl)-2-hydroxy-6-isopropyl-5-methoxycarbonylpyrimidine, 3.0 g (30.0 mmol.) of triethylamine, and 150 mL of toluene. The mixture in the flask was chilled to 0°C in an ice bath. To the chilled mixture was slowly added 8.46 g (30.0 mmol.) of trifluoromethanesulfonic anhydride, and the resulting mixture was subjected to reaction for 3 hours at the same temperature. After the reaction was complete, to the reaction mixture was added 90 mL of water. From the aqueous reaction mixture, an organic liquid portion was separated. The organic liquid portion was concentrated under reduced pressure. The residue was purified by silica gel column chromatography (column: Wako Gel C-200, eluent: hexane/ ethyl acetate (8:2, volume ratio)). There was obtained 8.46 g of 4-(4-fluorophenyl)-6-isopropyl-5-methoxycarbonyl-2-trifluoromethanesulfonyloxy-pyrimidine having the below-mentioned characteristics as a colorless oil. The yield was 74% (based on the amount of 4-(4-fluorophenyl)-2-hydroxy-6-isopropyl-5-methoxycarbonylpyrimidine).

30

IR (KBr,  $\text{cm}^{-1}$ ): 3421, 2978, 1737, 1570, 1429, 1222, 1136, 973, 851

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ ,  $\delta$  (ppm)): 1.33 (6H, d,  $J=6.6\text{Hz}$ ), 3.1-3.2 (1H, m), 3.80 (3H, s), 7.1-7.2 (2H, m), 7.7-7.8 (2H, m)

35



-39-

HRMS: 422.0585 (theoretical value ( $C_{16}H_{14}F_4N_2O_5S$  (M+))  
422.0560)

5 [Example 25] Preparation of 4-(4-fluorophenyl)-6-isopro-  
pyl-5-methoxycarbonyl-2-trifluoromethanesulfonyloxy-  
pyrimidine

In a 300 mL-volume glass flask equipped with a stir-  
rer, a thermometer and a reflux condenser were placed 2.9  
g (10.0 mmol.) of 4-(4-fluorophenyl)-2-hydroxy-6-isopro-  
10 pyl-5-methoxycarbonylpyrimidine, 1.7 g (16.8 mmol.) of  
triethylamine, and 50 mL of toluene. The mixture in the  
flask was chilled to 0°C in an ice bath. To the chilled  
mixture was slowly added 2.4 g (14.1 mmol.) of tri-  
fluoromethanesulfonyl chloride, and the resulting mixture  
15 was subjected to reaction for 3 hours at the same temper-  
ature. After the reaction was complete, to the reaction  
mixture was added 30 mL of water. From the aqueous reac-  
tion mixture, an organic liquid portion was separated.  
The organic liquid portion was concentrated under reduced  
20 pressure. The residue was purified by silica gel column  
chromatography (column: Wako Gel C-200, eluent: hexane/  
ethyl acetate (8:2, volume ratio)). There was obtained  
2.8 g of 4-(4-fluorophenyl)-6-isopropyl-5-methoxy-  
carbonyl-2-trifluoromethanesulfonyloxypyrimidine having  
25 the below-mentioned characteristics as a colorless oil.  
The yield was 66% (based on the amount of 4-(4-fluoro-  
phenyl)-2-hydroxy-6-isopropyl-5-methoxycarbonylpyrimi-  
dine).

30 [Example 26] Preparation of 4-(4-fluorophenyl)-6-iso-  
propyl-5-methoxycarbonyl-2-(N-methyl-N-methanesulfonyl-  
amino)pyrimidine

In a 50 mL-volume glass flask equipped with a stir-  
rer, a thermometer and a reflux condenser were placed 3.0  
35 g (7 mmol.) of 4-(4-fluorophenyl)-6-isopropyl-5-methoxy-  
carbonyl-2-trifluoromethanesulfonyloxypyrimidine, 1.45 g

-40-

(10.5 mmol.) of potassium carbonate (available from Wako Junyaku Co., Ltd., special grade), and 14 mL of butyl acetate. The mixture was heated to 122-125°C for 3 hours under refluxing, to carry out reaction. After the reaction was complete, the reaction mixture was cooled to room temperature. To the reaction mixture were added 10 mL of water and 7 mL of acetone, and the organic liquid portion was separated. The organic liquid portion was washed with a saturated aqueous sodium chloride solution and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (column: Wako Gel C-200, eluent: hexane/ ethyl acetate (5:1, volume ratio)). There was obtained 2.1 g of 4-(4-fluorophenyl)-6-isopropyl-5-methoxycarbonyl-2-(N-methyl-N-methanesulfonylamino)pyrimidine as a white crystalline product. The yield was 78% (based on the amount of 4-(4-fluorophenyl)-6-isopropyl-5-methoxycarbonyl-2-trifluoromethanesulfonyloxypyrimidine).

[Example 27] Preparation of 4-(4-fluorophenyl)-6-isopropyl-5-methoxycarbonyl-2-(N-methyl-N-methanesulfonylamino)pyrimidine

In a 50 mL-volume glass flask equipped with a stirrer, a thermometer and a reflux condenser were placed 1.1 g (2.5 mmol.) of 4-(4-fluorophenyl)-6-isopropyl-5-methoxycarbonyl-2-(p-toluenesulfonyloxy)pyrimidine, 0.55 g (5.0 mmol.) of N-methylmethanesulfonamide, 0.69 g (5.0 mmol.) of potassium carbonate (available from Wako Junyaku Co., Ltd., special grade), 0.32 g (1.0 mmol.) of tetrabutylammonium bromide, 20 mL of toluene and 5 mL of water. The mixture was heated to 85°C for 28 hours under refluxing, to carry out reaction. After the reaction was complete, the reaction mixture was cooled to room temperature. To the reaction mixture were added 10 mL of water and 7 mL of acetone, and the organic liquid portion was separated. The organic liquid portion was analyzed by

23940-1566

-41-

high performance liquid chromatography. It was confirmed that 0.6 g of 4-(4-fluorophenyl)-6-isopropyl-5-methoxycarbonyl-2-(N-methyl-N-methanesulfonylamino)pyrimidine was produced. The yield was 63% (based on the amount of 4-(4-fluorophenyl)-6-isopropyl-5-methoxycarbonyl-2-(p-toluenesulfonyloxy)pyrimidine).

[Industrial Utility]


The pyrimidine compound, particularly, 2-(N-methyl-N-methanesulfonylamino)pyrimidine compound, prepared by the invention is of value as an intermediate compound for the production of a cholesterol reducing agent (HMG-CoA reductase agent). The compound of formula (3) can be converted to an HMG CoA reductase inhibitor by the processes disclosed in European Patent Application Publication No. 0521471, Bioorg. Med. Chem., 5, 437 (1997) and International Patent Application No. WO 00/49014. The disclosures of these references demonstrate how a compound of formula (3) or formula (8) can be converted to an HMG CoA reductase inhibitor, in particular, rosuvastatin or a pharmaceutically acceptable salt thereof, such as rosuvastatin calcium.

23940-1566

-42-


CLAIMS:

1. A process for preparing a 2-(N-methyl-N-methanesulfonylamino)pyrimidine compound having the formula (3):




in which R is a hydrocarbyl group, which comprises the steps of:

15 reacting a hydroxypyrimidine compound having the formula (1):



in which R is the same as above, with an organic sulfonyl halide having the formula (2):



in which R' is a hydrocarbyl group and X is a halogen atom, or an organic sulfonic anhydride having the formula (2a):




in which R' has the same meaning as above, and reacting the resulting reaction product with N-methyl-N-methanesulfonamide.

35

-43-

2. The process of claim 1, wherein both of the reaction of the hydroxypyrimidine compound with the organic sulfonyl halide or the organic sulfonic anhydride and the reaction of the resulting reaction product with N-methyl-N-methanesulfonamide are performed in the presence of a base.

3. The process of claim 1, wherein the hydroxypyrimidine compound is prepared by oxidizing a dihydropyrimidinone compound having the formula (4):




wherein R is the same as defined in claim 1.

20

4. The process of claim 3, wherein the dihydropyrimidinone compound is oxidized using nitric acid.

5. The process of claim 3, wherein the dihydropyrimidinone compound is prepared by reacting an isobutyrylacetate ester having the formula (5):

30



in which R is the same as defined in claim 1, with 4-fluorobenzaldehyde and urea in the presence of a protonic compound and a metal salt.

6. The process of claim 5, wherein the protonic compound is a protonic acid.

35


-44-

7. The process of claim 6, wherein the protonic acid is sulfuric acid.

8. The process of claim 5, wherein the metal salt is copper(I) chloride.

9. A hydroxypyrimidine compound having the formula (1):

10




15 in which R is a hydrocarbyl group.

10. The hydroxypyrimidine compound of claim 9, wherein R is an alkyl group having 1 to 10 carbon atoms, a cycloalkyl group having 3 to 6 carbon atoms, an aryl-alkyl group having an alkyl moiety of 1-3 carbon atoms, or an aryl group.

11. A method for preparing the hydroxypyrimidine compound of claim 9, which comprises oxidizing a dihydropyrimidinone compound having the formula (4):

30




wherein R is a hydrocarbyl group.

12. The method of claim 11, wherein the dihydropyrimidinone compound is oxidized using nitric acid.

-45-

13. The method of claim 11, wherein the dihydropyrimidinone compound is prepared by reacting an isobutyrylacetate ester having the formula (5):

5



in which R is the same as defined in claim 11, with 4-fluorobenzaldehyde and urea in the presence of a protonic compound and a metal salt.

10

14. The method of claim 13, wherein the protonic compound is a protonic acid.

15


15. The method of claim 14, wherein the protonic acid is sulfuric acid.

16. The method of claim 13, wherein the metal salt is copper(I) chloride.

20

17. A dihydropyrimidinone compound having the formula (4):

25



30

wherein R is a hydrocarbyl group.


18. The hydroxypyrimidinone compound of claim 17, wherein R is an alkyl group having 1 to 10 carbon atoms, a cycloalkyl group having 3 to 6 carbon atoms, an aryl-alkyl group having an alkyl moiety of 1-3 carbon atoms, or an aryl group.

35

-46-

19. A method for preparing the dihydropyrimidinone compound of claim 17, which comprises reacting an isobutyrylacetate ester having the formula (5):

5



10 in which R is the same as defined in claim 17, with 4-fluorobenzaldehyde and urea in the presence of a protonic compound and a metal salt.


15 20. The method of claim 19, wherein the protonic compound is a protonic acid.

21. The method of claim 19, wherein the protonic acid is sulfuric acid.

20 22. The method of claim 19, wherein the metal salt is copper(I) chloride.

23. A method for preparing an aminopyrimidine compound having the formula (8):

25



30


wherein R is a hydrocarbyl group, and each of R<sup>1</sup> and R<sup>2</sup> independently is hydrogen atom, an alkyl group, an alkylsulfonyl group, or an arylsulfonyl group,

35 which comprises reacting a 2-substituted pyrimidine compound having the formula (6):




-47-

5



wherein R is the same as above, and X is a halogen atom  
 or an organic sulfonyloxy group,  
 10 with an amine compound having the formula (7):




15 wherein each of R<sup>1</sup> and R<sup>2</sup> is the same as above.

24. The method of claim 23, wherein R<sup>1</sup> is methyl and  
 R<sup>2</sup> is methanesulfonyl.

20 25. The method of claim 23, wherein the reaction of  
 the 2-substituted pyrimidine compound with the amine com-  
 pound is performed in the presence of a base.

25 26. A halogenopyrimidine compound having the formu-  
 la (9):

30




wherein R is a hydrocarbyl group, and Hal is a halogen  
 35 atom.

-48-

27. The halogenopyrimidine compound of claim 26, wherein R is an alkyl group having 1 to 10 carbon atoms, a cycloalkyl group having 3 to 6 carbon atoms, an aryl-alkyl group having an alkyl moiety of 1-3 carbon atoms, or an aryl group.

28. The halogenopyrimidine compound of claim 26, wherein Hal is a chlorine atom.


29. A method for preparing the halogenopyrimidine compound of claim 26, which comprises reacting a hydroxypyrimidine compound having the formula (1):



wherein R is a hydrocarbyl group, with a halogenating agent.

30. The method of claim 29, wherein the halogenating agent is phosphorus oxychloride or thionyl chloride.

31. An organic sulfonyloxypyrimidine compound having the formula (10):




wherein each of R and R' independently is a hydrocarbyl group.

23940-1566


-49-

32. The organic sulfonyloxy-pyrimidine compound of claim 31, wherein each of R and R' independently is an alkyl group having 1 to 10 carbon atoms, a cycloalkyl group having 3 to 6 carbon atoms, an arylalkyl group having an alkyl moiety of 1-3 carbon atoms, or an aryl group.


33. A method for preparing the organic sulfonyloxy-pyrimidine compound of claim 31, which comprises reacting a hydroxypyrimidine compound having the formula (1):



wherein R is a hydrocarbyl group, with an organic sulfonyl halide having the formula (2):




wherein R' is a hydrocarbyl group, and X is a halogen atom, or an organic sulfonic anhydride having the formula (2a):



in which R' has the same meaning as above.

34. A process for preparing a 2-(N-methyl-N-methanesulfonylamino)pyrimidine compound having the formula (3):




-50-

in which R is a hydrocarbyl group,  
which comprises the steps of:

(I) reacting an isobutyrylacetate ester of the following formula (5):

5



wherein R is a hydrocarbyl group.

10 with 4-fluorobenzaldehyde and urea in the presence of a protonic compound and a metal salt;

(II) oxidizing the reaction product of the step (I);

(III) reacting the oxidation product of the step (II) with an organic sulfonyl halide having the formula

15 (2):



20 in which R' is a hydrocarbyl group, and X is a halogen atom, or an organic sulfonic anhydride having the formula (2a):



25 in which R' has the same meaning as above; and

(IV) reacting the reaction product of the step (III) with N-methyl-N-methanesulfonamide.

30 35. The process of claim 1, wherein R' is a substituted or unsubstituted aryl group.

36. The process as claimed in claim 1, followed by conversion of the compound of formula (3) to rosuvastatin or a pharmaceutically acceptable salt thereof.

35

23940-1566

- 51 -

37. The process as claimed in claim 34, followed by conversion of the compound of formula (3) to rosuvastatin or a pharmaceutically acceptable salt thereof.

38. The method as claimed in claim 23, followed by conversion of the  
5 compound of formula (8) to an HMG CoA reductase inhibitor.

39. A method as claimed in claim 38, wherein the HMG CoA reductase inhibitor is rosuvastatin or a pharmaceutically acceptable salt thereof.

