



US 20190300614A1

(19) **United States**

(12) **Patent Application Publication**
DIMOUDIS et al.

(10) **Pub. No.: US 2019/0300614 A1**

(43) **Pub. Date: Oct. 3, 2019**

(54) **ANTIBODIES AGAINST HUMAN CSF-1R AND USES THEREOF**

application No. 15/292,710, filed on Oct. 13, 2016, now Pat. No. 9,663,580, which is a division of application No. 14/640,183, filed on Mar. 6, 2015, now Pat. No. 9,499,625, which is a division of application No. 12/962,445, filed on Dec. 7, 2010, now Pat. No. 8,999,327.

(71) Applicant: **HOFFMANN-LA ROCHE INC.**,
Little Falls, NJ (US)

(72) Inventors: **Nikolaos DIMOUDIS**, Wielenbach (DE); **Georg FERTIG**, Penzberg (DE); **Alexander FIDLER**, Penzberg (DE); **Guy GEORGES**, Habach (DE); **Klaus KALUZA**, Weilheim (DE); **Martin LANZENDOERFER**, Tutzing (DE); **Carola RIES**, Penzberg (DE); **Stefan SEEBER**, Penzberg (DE); **Marlene THOMAS**, Penzberg (DE)

(30) **Foreign Application Priority Data**

Dec. 10, 2009 (EP) 09015310.7
Aug. 19, 2010 (EP) 10173407.7

Publication Classification

(51) **Int. Cl.**
C07K 16/28 (2006.01)
(52) **U.S. Cl.**
CPC **C07K 16/2866** (2013.01); **C07K 2317/76** (2013.01); **C07K 2317/73** (2013.01); **C07K 2317/567** (2013.01); **A61K 2039/505** (2013.01); **C07K 2317/56** (2013.01); **C07K 2317/21** (2013.01); **C07K 2317/92** (2013.01); **C07K 2317/565** (2013.01)

(73) Assignee: **HOFFMANN-LA ROCHE INC.**,
Little Falls, NJ (US)

(21) Appl. No.: **16/362,962**

(22) Filed: **Mar. 25, 2019**

Related U.S. Application Data

(62) Division of application No. 16/055,947, filed on Aug. 6, 2018, now Pat. No. 10,287,358, which is a division of application No. 15/875,530, filed on Jan. 19, 2018, now Pat. No. 10,077,314, which is a division of application No. 15/489,519, filed on Apr. 17, 2017, now Pat. No. 9,879,085, which is a division of

(57) **ABSTRACT**

The present invention relates to antibodies against human CSF-1R (anti-CSF-1R antibody), methods for their production, pharmaceutical compositions containing said antibodies, and uses thereof.

Specification includes a Sequence Listing.




FIG. 1A




FIG. 1B




FIG. 2A




FIG. 2B




FIG. 2C




FIG. 2D




FIG. 2E




FIG. 2F




FIG. 3A




FIG. 3B




FIG. 3C




FIG. 3D




FIG. 4

ANTIBODIES AGAINST HUMAN CSF-1R AND USES THEREOF

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application is a Divisional of U.S. patent application Ser. No. 15/292,710, filed Oct. 13, 2016, which is a Divisional of U.S. patent application Ser. No. 14/640,183, filed Mar. 6, 2015, now U.S. Pat. No. 9,499,625, issued Nov. 22, 2016, which is a Divisional of U.S. patent application Ser. No. 12/962,445, filed Dec. 7, 2010, now U.S. Pat. No. 8,999,327, issued Apr. 7, 2015, which claims the benefit of EP Patent Application Nos. 09 015 310.7, filed Dec. 10, 2009, and 10 173 407.7, filed Aug. 19, 2010, the entire disclosures of which are expressly incorporated by reference herein.

SUBMISSION OF SEQUENCE LISTING ON ASCII TEXT FILE

[0002] The content of the following submission on ASCII text file is incorporated herein by reference in its entirety: a computer readable form (CRF) of the Sequence Listing (file name: 146392029613SEQLIST.txt, date recorded: Apr. 14, 2017, size: 68 KB).

FIELD OF THE INVENTION

[0003] The present invention relates to antibodies against human CSF-1R (anti-CSF-1R antibodies), methods for their production, pharmaceutical compositions containing said antibodies, and uses thereof.

BACKGROUND OF THE INVENTION

[0004] The human CSF-1 receptor (CSF-1R; colony stimulating factor 1 receptor; synonyms: M-CSF receptor; Macrophage colony-stimulating factor 1 receptor. *Fms* proto-oncogene, *c-fms*. SEQ ID NO: 62) is known since 1986 (Coussens, L., et al., *Nature* 320 (1986) 277-280). CSF-1R is a growth factor and encoded by the *c-fms* proto-oncogene (reviewed e.g. in Roth, P., and Stanley, E. R., *Curr. Top. Microbiol. Immunol.* 181 (1992) 141-67).

[0005] CSF-1R is the receptor for CSF-1 (colony stimulating factor 1, also called M-CSF, macrophage colony-stimulating factor) and mediates the biological effects of this cytokine (Sherr, C. J., et al., *Cell* 41 (1985) 665-676). The cloning of the colony stimulating factor-1 receptor (CSF-1R) (also called *c-fms*) was described for the first time in Roussel, M. F., et al., *Nature* 325 (1987) 549-552. In that publication, it was shown that CSF-1R had transforming potential dependent on changes in the Cterminal tail of the protein including the loss of the inhibitory tyrosine 969 phosphorylation which binds Cbl and thereby regulates receptor down regulation (Lee, P. S., et al., *Embo J.* 18 (1999) 3616-3628). Recently a second ligand for CSF-1R termed interleukin-34 (IL-34) was identified (Lin, H., et al. *Science* 320 (2008) 807-811).

[0006] The cytokine CSF-1 (colony stimulating factor 1, also called M-CSF, macrophage) is found extracellularly as a disulfide-linked homodimer (Stanley, E. R. et al.,

[0007] *Journal of Cellular Biochemistry* 21 (1983) 151-159; Stanley, E. R. et al., *Stem Cells* 12 Suppl. 1 (1995) 15-24).

[0008] The main biological effects of CSF-1R signaling are the differentiation, proliferation, migration, and survival

of hematopoietic precursor cells to the macrophage lineage (including osteoclast). Activation of CSF-1R is mediated by its ligands, CSF-1 (M-CSF) and IL-34. Binding of CSF-1 (M-CSF) to CSF-1R induces the formation of homodimers and activation of the kinase by tyrosine phosphorylation (Li, W. et al, *EMBO Journal.* 10 (1991) 277-288; Stanley, E. R., et al., *Mol. Reprod. Dev.* 46 (1997) 4-10).

[0009] The biologically active homodimer CSF-1 binds to the CSF-1R within the subdomains D1 to D3 of the extracellular domain of the CSF-1 receptor (CSF-1R-ECD). The CSF-1R-ECD comprises five immunoglobulin-like subdomains (designated D1 to D5). The subdomains D4 to D5 of the extracellular domain (CSF-1R-ECD) are not involved in the CSF-1 binding. (Wang, Z., et al *Molecular and Cellular Biology* 13 (1993) 5348-5359). The subdomain D4 is involved in dimerization (Yeung, Y-G., et al *Molecular & Cellular Proteomics* 2 (2003) 1143-1155; Pixley, F. J., et al., *Trends Cell Biol* 14 (2004) 628-638).

[0010] Further signaling is mediated by the p85 subunit of PI3K and Grb2 connecting to the PI3K/AKT and Ras/MAPK pathways, respectively. These two important signaling pathways can regulate proliferation, survival and apoptosis. Other signaling molecules that bind the phosphorylated intracellular domain of CSF-1R include STAT1, STAT3, PLC γ , and Cbl (Bourette, R. P. and Rohrschneider, L. R., *Growth Factors* 17 (2000) 155-166).

[0011] CSF-1R signaling has a physiological role in immune responses, in bone remodeling and in the reproductive system. The knockout animals for either CSF-1 (Polard, J. W., *Mol. Reprod. Dev.* 46 (1997) 54-61) or CSF-1R (Dai, X. M., et al., *Blood* 99 (2002) 111-120) have been shown to have osteopetrotic, hematopoietic, tissue macrophage, and reproductive phenotypes consistent with a role for CSF-1R in the respective cell types.

[0012] Sherr, C. J., et al., *Blood* 73 (1989) 1786-1793 relates to some antibodies against CSF-1R that inhibit the CSF-1 activity (see Sherr, C. J. et al., *Blood* 73 (1989) 1786-1793). Ashmun, R. A., et al., *Blood* 73 (1989) 827-837 relates to CSF-1R antibodies. Lenda, D., et al., *Journal of Immunology* 170 (2003) 3254-3262 relates to reduced macrophage recruitment, proliferation, and activation in CSF-1-deficient mice results in decreased tubular apoptosis during renal inflammation. Kitaura, H., et al., *Journal of Dental Research* 87 (2008) 396-400 refers to an anti-CSF-1 antibody which inhibits orthodontic tooth movement. WO 2001/030381 mentions CSF-1 activity inhibitors including antisense nucleotides and antibodies while disclosing only CSF-1 antisense nucleotides. WO 2004/045532 relates to metastases and bone loss prevention and treatment of metastatic cancer by a CSF-1 antagonist disclosing as antagonist anti-CSF-1-antibodies only. WO 2005/046657 relates to the treatment of inflammatory bowel disease by anti-CSF-1-antibodies. US 2002/0141994 relates to inhibitors of colony stimulating factors. WO 2006/096489 relates to the treatment of rheumatoid arthritis by anti-CSF-1-antibodies. WO 2009/026303 and WO 2009/112245 relate to certain anti-CSF-1R antibodies binding to CSF-1R within the first three subdomains (D1 to D3) of the Extracellular Domain (CSF-1R-ECD).

SUMMARY OF THE INVENTION

[0013] The invention comprises an isolated antibody binding to human CSF-1R, wherein the antibody binds to human

CSF-1R fragment delD4 (SEQ ID NO: 65) and to human CSF-1R Extracellular Domain (SEQ ID NO: 64) with a ratio of 1:50 or lower.

[0014] The invention further comprises an isolated antibody wherein

[0015] a) the heavy chain variable domain comprises SEQ ID NO:7 and the light chain variable domain comprises SEQ ID NO:8,

[0016] b) the heavy chain variable domain comprises SEQ ID NO:15 and the light chain variable domain comprises SEQ ID NO: 16;

[0017] c) the heavy chain variable domain comprises SEQ ID NO:75 and the light chain variable domain comprises SEQ ID NO:76;

[0018] d) the heavy chain variable domain comprises SEQ ID NO:83 and the light chain variable domain comprises SEQ ID NO:84;

[0019] or a humanized version thereof.

[0020] The invention further comprises an isolated antibody wherein

[0021] a) the heavy chain variable domain comprises SEQ ID NO:7 and the light chain variable domain comprises SEQ ID NO:8,

[0022] b) the heavy chain variable domain comprises SEQ TD NO:15 and the light chain variable domain comprises SEQ ID NO: 16;

[0023] or a humanized version thereof.

[0024] In one embodiment the isolated antibody comprises

[0025] a) a heavy chain variable domain comprising SEQ ID NO:23 and a light chain variable domain comprising SEQ ID NO:24, or

[0026] b) a heavy chain variable domain comprising SEQ ID NO:31 and a light chain variable domain comprising SEQ ID NO:32, or

[0027] c) a heavy chain variable domain comprising SEQ ID NO:39 and a light chain variable domain comprising SEQ ID NO:40, or

[0028] d) a heavy chain variable domain comprising SEQ ID NO:47 and a light chain variable domain comprising SEQ ID NO:48, or

[0029] e) a heavy chain variable domain comprising SEQ ID NO:55 and a light chain variable domain comprising SEQ ID NO:56.

[0030] The invention further comprises an isolated antibody, wherein

[0031] a) the heavy chain variable domain comprises a CDR3 region comprising SEQ ID NO: 1, a CDR2 region comprising SEQ ID NO: 2, and a CDR1 region comprising SEQ ID NO:3, and the light chain variable domain comprises a CDR3 region comprising SEQ ID NO: 4, a CDR2 region comprising SEQ ID NO:5, and a CDR1 region comprising SEQ ID NO:6, or

[0032] b) the heavy chain variable domain comprises a CDR3 region comprising SEQ ID NO: 9, a CDR2 region comprising SEQ ID NO: 10, and a CDR1 region comprising SEQ ID NO: 11, and the light chain variable domain comprises a CDR3 region comprising SEQ ID NO:12, a CDR2 region comprising SEQ ID NO: 13, and a CDR1 region comprising SEQ ID NO: 14, or

[0033] c) the heavy chain variable domain comprises a CDR3 region comprising SEQ ID NO: 17, a CDR2 region comprising SEQ ID NO: 18, and a CDR1 region

comprising SEQ ID NO:19, and the light chain variable domain comprises a CDR3 region comprising SEQ ID NO: 20, a CDR2 region comprising SEQ ID NO:21, and a CDR1 region comprising SEQ ID NO:22, or

[0034] d) the heavy chain variable domain comprises a CDR3 region comprising SEQ ID NO: 25, a CDR2 region comprising SEQ ID NO: 26, and a CDR1 region comprising SEQ ID NO: 27, and the light chain variable domain comprises a CDR3 region comprising SEQ ID NO:28, a CDR2 region comprising SEQ ID NO: 29, and a CDR1 region comprising SEQ ID NO: 30, or

[0035] e) the heavy chain variable domain comprises a CDR3 region comprising SEQ ID NO: 33, a CDR2 region comprising SEQ ID NO: 34, and a CDR1 region comprising SEQ ID NO: 35, and the light chain variable domain comprises a CDR3 region comprising SEQ ID NO:36, a CDR2 region comprising SEQ ID NO: 37, and a CDR1 region comprising SEQ ID NO: 38, or

[0036] f) the heavy chain variable domain comprises a CDR3 region comprising SEQ ID NO:41, a CDR2 region comprising SEQ ID NO: 42, and a CDR1 region comprising SEQ ID NO:43, and the light chain variable domain comprises a CDR3 region comprising SEQ ID NO: 44, a CDR2 region comprising SEQ ID NO:45, and a CDR1 region comprising SEQ ID NO:46, or

[0037] g) the heavy chain variable domain comprises a CDR3 region comprising SEQ ID NO: 49, a CDR2 region comprising SEQ ID NO: 50, and a CDR1 region comprising SEQ ID NO: 51, and the light chain variable domain comprises a CDR3 region comprising SEQ ID NO:52, a CDR2 region comprising SEQ ID NO: 53, and a CDR1 region comprising SEQ ID NO: 54; or

[0038] h) the heavy chain variable domain comprises a CDR3 region comprising SEQ ID NO:69, a CDR2 region comprising SEQ ID NO: 70, and a CDR1 region comprising SEQ ID NO:71, and the light chain variable domain comprises a CDR3 region comprising SEQ ID NO: 72, a CDR2 region comprising SEQ ID NO:73, and a CDR1 region comprising SEQ ID NO:74, or

[0039] i) the heavy chain variable domain comprises a CDR3 region comprising SEQ ID NO: 77, a CDR2 region comprising SEQ ID NO: 78, and a CDR1 region comprising SEQ ID NO: 79, and the light chain variable domain comprises a CDR3 region comprising SEQ ID NO:80, a CDR2 region comprising SEQ ID NO: 81, and a CDR1 region comprising SEQ ID NO: 82.

[0040] Preferably the antibody according to the invention is of human IgG1 subclass or of human IgG4 subclass.

[0041] A further embodiment of the invention is a pharmaceutical composition comprising any of the antibodies described herein.

[0042] The invention further comprises the use an of an antibody according to the invention for the manufacture of a medicament for treatment of a CSF-1R mediated disease.

[0043] The invention further comprises the use an of an antibody according to the invention for the manufacture of a medicament for treatment of cancer.

[0044] The invention further comprises the use an of an antibody according to the invention for the manufacture of a medicament for treatment of bone loss.

[0045] The invention further comprises the use of an antibody according to the invention for the manufacture of a medicament for treatment of metastasis.

[0046] The invention further comprises the use of an antibody according to the invention for the manufacture of a medicament for treatment of inflammatory diseases.

[0047] The invention further comprises an antibody according to the invention for treatment of a CSF-1R mediated disease.

[0048] The invention further comprises an antibody according to the invention for treatment of cancer.

[0049] The invention further comprises an antibody according to the invention for treatment of bone loss.

[0050] The invention further comprises an antibody according to the invention for treatment of metastasis.

[0051] The invention further comprises an antibody according to the invention for treatment of inflammatory diseases.

[0052] A further embodiment of the invention is a nucleic acid encoding an antibody wherein

[0053] a) the heavy chain variable domain comprises a CDR3 region comprising SEQ ID NO: 1, a CDR2 region comprising SEQ ID NO: 2, and a CDR1 region comprising SEQ ID NO:3, and the light chain variable domain comprises a CDR3 region comprising SEQ ID NO: 4, a CDR2 region comprising SEQ ID NO:5, and a CDR1 region comprising SEQ ID NO:6, or

[0054] b) the heavy chain variable domain comprises a CDR3 region comprising SEQ ID NO: 9, a CDR2 region comprising SEQ ID NO: 10, and a CDR1 region comprising SEQ ID NO: 11, and the light chain variable domain comprises a CDR3 region comprising SEQ ID NO:12, a CDR2 region comprising SEQ ID NO: 13, and a CDR1 region comprising SEQ ID NO: 14, or

[0055] c) the heavy chain variable domain comprises a CDR3 region comprising SEQ ID NO: 17, a CDR2 region comprising SEQ ID NO: 18, and a CDR1 region comprising SEQ ID NO:19, and the light chain variable domain comprises a CDR3 region comprising SEQ ID NO: 20, a CDR2 region comprising SEQ ID NO:21, and a CDR1 region comprising SEQ ID NO:22, or

[0056] d) the heavy chain variable domain comprises a CDR3 region comprising SEQ ID NO: 25, a CDR2 region comprising SEQ ID NO: 26, and a CDR1 region comprising SEQ ID NO: 27, and the light chain variable domain comprises a CDR3 region comprising SEQ ID NO:28, a CDR2 region comprising SEQ ID NO: 29, and a CDR1 region comprising SEQ ID NO: 30, or

[0057] e) the heavy chain variable domain comprises a CDR3 region comprising SEQ ID NO: 33, a CDR2 region comprising SEQ ID NO: 34, and a CDR1 region comprising SEQ ID NO: 35, and the light chain variable domain comprises a CDR3 region comprising SEQ ID NO:36, a CDR2 region comprising SEQ ID NO: 37, and a CDR1 region comprising SEQ ID NO: 38, or

[0058] f) the heavy chain variable domain comprises a CDR3 region comprising SEQ ID NO:41, a CDR2 region comprising SEQ ID NO: 42, and a CDR1 region comprising SEQ ID NO:43, and the light chain variable domain comprises a CDR3 region comprising SEQ ID

NO: 44, a CDR2 region comprising SEQ ID NO:45, and a CDR1 region comprising SEQ ID NO:46, or

[0059] g) the heavy chain variable domain comprises a CDR3 region comprising SEQ ID NO: 49, a CDR2 region comprising SEQ ID NO: 50, and a CDR1 region comprising SEQ ID NO: 51, and the light chain variable domain comprises a CDR3 region comprising SEQ ID NO:52, a CDR2 region comprising SEQ ID NO: 53, and a CDR1 region comprising SEQ ID NO: 54, or

[0060] h) the heavy chain variable domain comprises a CDR3 region comprising SEQ ID NO:69, a CDR2 region comprising SEQ ID NO: 70, and a CDR1 region comprising SEQ ID NO:71, and the light chain variable domain comprises a CDR3 region comprising SEQ ID NO: 72, a CDR2 region comprising SEQ ID NO:73, and a CDR1 region comprising SEQ ID NO:74, or

[0061] i) the heavy chain variable domain comprises a CDR3 region comprising SEQ ID NO: 77, a CDR2 region comprising SEQ ID NO: 78, and a CDR1 region comprising SEQ ID NO: 79, and the light chain variable domain comprises a CDR3 region comprising SEQ ID NO:80, a CDR2 region comprising SEQ ID NO: 81, and a CDR1 region comprising SEQ ID NO: 82.

[0062] A further embodiment of the invention is a nucleic acid encoding an antibody wherein

[0063] a) the heavy chain variable domain comprises SEQ ID NO:7 and the light chain variable domain comprises SEQ ID NO:8,

[0064] b) the heavy chain variable domain comprises SEQ ID NO:15 and the light chain variable domain comprises SEQ ID NO: 16;

[0065] c) the heavy chain variable domain comprises SEQ ID NO:75 and the light chain variable domain comprises SEQ ID NO:76;

[0066] d) the heavy chain variable domain comprises SEQ ID NO:83 and the light chain variable domain comprises SEQ ID NO:84;

[0067] or a humanized version thereof.

[0068] A further embodiment of the invention is a nucleic acid encoding an antibody wherein

[0069] a) the heavy chain variable domain comprises SEQ ID NO:23 and the light chain variable domain comprises SEQ ID NO:24, or

[0070] b) the heavy chain variable domain comprises SEQ ID NO:31 and the light chain variable domain comprises SEQ ID NO:32, or

[0071] c) the heavy chain variable domain comprises SEQ ID NO:39 and the light chain variable domain comprises SEQ ID NO:40, or

[0072] d) the heavy chain variable domain comprises SEQ ID NO:47 and the light chain variable domain comprises SEQ ID NO:48, or

[0073] e) the heavy chain variable domain comprises SEQ ID NO:55 and the light chain variable domain comprises SEQ ID NO:56.

[0074] The invention further provides expression vectors containing nucleic acid according to the invention capable of expressing said nucleic acid in a prokaryotic or eukaryotic host cell, and host cells containing such vectors for the recombinant production of an antibody according to the invention.

[0075] The invention further comprises a prokaryotic or eukaryotic host cell comprising a vector according to the invention.

[0076] The invention further comprises a method for the production of a recombinant human or humanized antibody according to the invention, the method comprising expressing a nucleic acid according to the invention in a prokaryotic or eukaryotic host cell and recovering said antibody from said cell or the cell culture supernatant.

[0077] The invention further comprises the antibody obtained by such a recombinant method.

[0078] Antibodies according to the invention show benefits for patients in need of a CSF-1R targeting therapy. The antibodies according to the invention show efficient antiproliferative activity against ligand-independent and ligand-dependant proliferation and are therefore especially useful in the treatment of cancer and metastasis.

[0079] The invention further provides a method for treating a patient suffering from cancer, the method comprising administering to a patient diagnosed as having such a disease (and therefore being in need of such a therapy) an effective amount of an antibody according to the invention. The antibody is administered preferably in a pharmaceutical composition.

[0080] A further embodiment of the invention is a method for treating a patient suffering from cancer the method comprising administering to the patient an antibody according to the invention.

[0081] These and other embodiments of the invention are further described in the detailed description that follows.

DESCRIPTION OF THE SEQUENCES

- [0082]** SEQ ID NO: 1 heavy chain CDR3, Mab 2F11
[0083] SEQ ID NO: 2 heavy chain CDR2, Mab 2F11
[0084] SEQ ID NO: 3 heavy chain CDR1, Mab 2F11
[0085] SEQ ID NO: 4 light chain CDR3, Mab 2F11
[0086] SEQ ID NO: 5 light chain CDR2, Mab 2F11
[0087] SEQ ID NO: 6 light chain CDR1, Mab 2F11
[0088] SEQ ID NO: 7 heavy chain variable domain, Mab 2F11
[0089] SEQ ID NO: 8 light chain variable domain, Mab 2F11
[0090] SEQ ID NO: 9 heavy chain CDR3, Mab 2E10
[0091] SEQ ID NO: 10 heavy chain CDR2, Mab 2E10
[0092] SEQ ID NO: 11 heavy chain CDR1, Mab 2E10
[0093] SEQ ID NO: 12 light chain CDR3, Mab 2E10
[0094] SEQ ID NO: 13 light chain CDR2, Mab 2E10
[0095] SEQ ID NO: 14 light chain CDR1, Mab 2E10
[0096] SEQ ID NO: 15 heavy chain variable domain, Mab 2E10
[0097] SEQ ID NO: 16 light chain variable domain, Mab 2E10
[0098] SEQ ID NO: 17 heavy chain CDR3, hMab 2F11-c11
[0099] SEQ ID NO: 18 heavy chain CDR2, hMab 2F11-c11
[0100] SEQ ID NO: 19 heavy chain CDR1, hMab 2F11-c11
[0101] SEQ ID NO: 20 light chain CDR3, hMab 2F11-c11
[0102] SEQ ID NO: 21 light chain CDR2, hMab 2F11-c11
[0103] SEQ ID NO: 22 light chain CDR1, hMab 2F11-c11
[0104] SEQ ID NO: 23 heavy chain variable domain, hMab 2F11-c11
[0105] SEQ ID NO: 24 light chain variable domain, hMab 2F11-c11
[0106] SEQ ID NO: 25 heavy chain CDR3, hMab 2F11-d8
[0107] SEQ ID NO: 26 heavy chain CDR2, hMab 2F11-d8
[0108] SEQ ID NO: 27 heavy chain CDR1, hMab 2F11-d8
[0109] SEQ ID NO: 28 light chain CDR3, hMab 2F11-d8
[0110] SEQ ID NO: 29 light chain CDR2, hMab 2F11-d8
[0111] SEQ ID NO: 30 light chain CDR1, hMab 2F11-d8
[0112] SEQ ID NO: 31 heavy chain variable domain, hMab 2F11-d8
[0113] SEQ ID NO: 32 light chain variable domain, hMab 2F11-d8
[0114] SEQ ID NO: 33 heavy chain CDR3, hMab 2F11-e7
[0115] SEQ ID NO: 34 heavy chain CDR2, hMab 2F11-e7
[0116] SEQ ID NO: 35 heavy chain CDR1, hMab 2F11-e7
[0117] SEQ ID NO: 36 light chain CDR3, hMab 2F11-e7
[0118] SEQ ID NO: 37 light chain CDR2, hMab 2F11-e7
[0119] SEQ ID NO: 38 light chain CDR1, hMab 2F11-e7
[0120] SEQ ID NO: 39 heavy chain variable domain, hMab 2F11-e7
[0121] SEQ ID NO: 40 light chain variable domain, hMab 2F11-e7
[0122] SEQ ID NO: 41 heavy chain CDR3, hMab 2F11-f12
[0123] SEQ ID NO: 42 heavy chain CDR2, hMab 2F11-f12
[0124] SEQ ID NO: 43 heavy chain CDR1, hMab 2F11-f12
[0125] SEQ ID NO: 44 light chain CDR3, hMab 2F11-f12
[0126] SEQ ID NO: 45 light chain CDR2, hMab 2F11-f12
[0127] SEQ ID NO: 46 light chain CDR1, hMab 2F11-f12
[0128] SEQ ID NO: 47 heavy chain variable domain, hMab 2F11-f12
[0129] SEQ ID NO: 48 light chain variable domain, hMab 2F11-f12
[0130] SEQ ID NO: 49 heavy chain CDR3, hMab 2F11-g1
[0131] SEQ ID NO: 50 heavy chain CDR2, hMab 2F11-g1
[0132] SEQ ID NO: 51 heavy chain CDR1, hMab 2F11-g1
[0133] SEQ ID NO: 52 light chain CDR3, hMab 2F11-g1
[0134] SEQ ID NO: 53 light chain CDR2, hMab 2F11-g1
[0135] SEQ ID NO: 54 light chain CDR1, hMab 2F11-g1

- [0136] SEQ ID NO: 55 heavy chain variable domain, hMab 2F11-g1
- [0137] SEQ ID NO: 56 light chain variable domain, hMab 2F11-g1
- [0138] SEQ ID NO: 57 human kappa light chain constant region
- [0139] SEQ ID NO: 58 human heavy chain constant region derived from IgG
- [0140] SEQ ID NO: 59 human heavy chain constant region derived from IgG1 mutated on L234A and L235A
- [0141] SEQ ID NO: 60 human heavy chain constant region derived from IgG4
- [0142] SEQ ID NO: 61 human heavy chain constant region derived from IgG4 mutated on S228P
- [0143] SEQ ID NO: 62 human wildtype CSF-1R (wt CSF-1R)
- [0144] SEQ ID NO: 63 human mutant CSF-1R L301S Y969F
- [0145] SEQ ID NO: 64 human CSF-1R Extracellular Domain
- [0146] SEQ ID NO: 65 human CSF-1R fragment delD4
- [0147] SEQ ID NO: 66 human CSF-1R fragment D1-D3
- [0148] SEQ ID NO: 67 signal peptide
- [0149] SEQ ID NO: 68 Primer
- [0150] SEQ ID NO: 69 heavy chain CDR3, Mab 1G10
- [0151] SEQ ID NO: 70 heavy chain CDR2, Mab 1G10
- [0152] SEQ ID NO: 71 heavy chain CDR1, Mab 1G10
- [0153] SEQ ID NO: 72 light chain CDR3, Mab 1G10
- [0154] SEQ ID NO: 73 light chain CDR2, Mab 1G10
- [0155] SEQ ID NO: 74 light chain CDR1, Mab 1G10
- [0156] SEQ ID NO: 75 heavy chain variable domain, Mab 1G10
- [0157] SEQ ID NO: 76 light chain variable domain, Mab 1G10
- [0158] SEQ ID NO: 77 heavy chain CDR3, Mab 2H7
- [0159] SEQ ID NO: 78 heavy chain CDR2, Mab 2H7
- [0160] SEQ ID NO: 79 heavy chain CDR1, Mab 2H7
- [0161] SEQ ID NO: 80 light chain CDR3, Mab 2H7
- [0162] SEQ ID NO: 81 light chain CDR2, Mab 2H7
- [0163] SEQ ID NO: 82 light chain CDR1, Mab 2H7
- [0164] SEQ ID NO: 83 heavy chain variable domain, Mab 2H7
- [0165] SEQ ID NO: 84 light chain variable domain, Mab 2H7

DESCRIPTION OF THE FIGURES

[0166] FIG. 1A-1B Growth inhibition of BeWo tumor cells in 3D culture under treatment with different anti-CSF-1R monoclonal antibodies at a concentration of 10 μ g/ml.

[0167] X axis: viability normalized mean relative light units (RLU) corresponding to the ATP-content of the cells (CellTiterGlo assay).

[0168] Y axis: tested probes: Minimal Medium (0.5% FBS), mouse IgG1 (mIgG1, 10 μ g/ml), mouse IgG2a (mIgG2a 10 μ g/ml), CSF-1 only, Mab 2F11, Mab 2E10, Mab 2H7, Mab 1G10 and SC 2-4A5. Highest inhibition of CSF-1 induced growth was observed with the anti-CSF-1R antibodies according to the invention.

[0169] FIG. 2A Biacore sensogram of binding of different anti-CSF-1R antibodies to immobilized human CSF-1R fragment delD4 (comprising the extracellular subdomains D1-D3 and D5) (SEQ ID NO: 65) (y-axis: binding signal in

Response Units (RU), baseline=0 RU, x-axis: time in seconds (s)): While the antibodies Mab 3291 and sc 2-4A5 clearly show binding to this delD4 fragment, the antibodies according to the invention e.g. Mab 2F11, and Mab 2E10, did not bind to the CSF-1R fragment delD4. The control anti-CCR5 antibody m<CCR5>Pz03.1C5 did also not bind to the CSF-1R fragment delD4.

[0170] FIG. 2B Biacore sensogram of binding of different anti-CSF-1R antibodies to immobilized human CSF-1R Extracellular Domain (CSF-1R-ECD) (comprising the extracellular subdomains D1-D5) (SEQ ID NO: 64) (y-axis: binding signal in Response Units (RU), baseline=0 RU, x-axis: time in seconds (s)): All anti-CSF-1R antibodies show binding to CSF-1R-ECD. The control anti-CCR5 antibody m<CCR5>Pz03.1C5 did not bind to the CSF-1R-ECD.

[0171] FIG. 2C Biacore sensogram of binding of different anti-CSF-1R antibodies to immobilized human CSF-1R fragment delD4 (comprising the extracellular subdomains D1-D3 and D5) (SEQ ID NO: 65) (y-axis: binding signal in Response Units (RU), baseline=0 RU, x-axis: time in seconds (s)): Mab 1G10, Mab 2H7 and humanized hMab 2F11-e7 did not bind to the CSF-1R fragment delD4. The control anti-CCR5 antibody m<CCR5>Pz03.1C5 did also not bind to the CSF-1R fragment delD4.

[0172] FIG. 2D Biacore sensogram of binding of different anti-CSF-1R antibodies to immobilized human CSF-1R Extracellular Domain (CSF-1R-ECD) (comprising the extracellular subdomains D1-D5) (SEQ ID NO: 64) (y-axis: binding signal in Response Units (RU), baseline=0 RU, x-axis: time in seconds (s)): All anti-CSF-1R antibodies Mab 1G10, Mab 2H7 and humanized hMab 2F11-e7 showed binding to CSF-1R-ECD. The control anti-CCR5 antibody m<CCR5>Pz03.1C5 did not bind to the CSF-1R-ECD.

[0173] FIG. 2E Biacore sensogram of binding of different anti-CSF-1R antibodies to immobilized human CSF-1R fragment delD4 (comprising the extracellular subdomains D1-D3 and D5) (SEQ ID NO: 65) (y-axis: binding signal in Response Units (RU), baseline=0 RU, x-axis: time in seconds (s)): All anti-CSF-1R antibodies 1.2.SM, CXIIG6, ab10676 and MAB3291 show binding to the CSF-1R fragment delD4. The control anti-CCR5 antibody m<CCR5>Pz03.1C5 did also not bind to the CSF-1R fragment delD4.

[0174] FIG. 2F Biacore sensogram of binding of different anti-CSF-1R antibodies to immobilized human CSF-1R Extracellular Domain (CSF-1R-ECD) (comprising the extracellular subdomains D1-D5) (SEQ ID NO: 64) (y-axis: binding signal in Response Units (RU), baseline=0 RU, x-axis: time in seconds (s)): All anti-CSF-1R antibodies 1.2.SM, CXIIG6, ab10676 and MAB3291 show binding to CSF-1R-ECD. The control anti-CCR5 antibody m<CCR5>Pz03.1C5 did not bind to the CSF-1R-ECD.

[0175] FIG. 3A-3D CSF-1 levels in Cynomolgus monkey after application of different dosages of anti-CSF-1R antibody according to the invention

[0176] FIG. 4 In vivo efficacy—tumor growth inhibition of anti-CSF-1R antibodies according to the invention in breast cancer BT20 xenograft

DETAILED DESCRIPTION OF THE INVENTION

I. Introduction

[0177] The present invention provides compositions that bind human CSF-1R or a portion thereof, kits and articles of manufacture comprising such compositions, and methods of using such compositions, including, e.g., methods for modulating ligand binding to CSF-1R and for modulating biological/physiological activities associated with ligand binding to CSF-1R. The invention is based in part on the identification of a variety of anti-CSF-1R antibodies that bind CSF-1R. The anti-CSF-1R antibodies of the invention can conveniently be used as therapeutic and diagnostic agents for use in targeting CSF-1R-mediated diseases including, e.g., cancer (including, for example, metastatic cancer), bone disorders (including, for example, bone loss), and inflammatory disorders.

[0178] Surprisingly it has been found that, using a human CSF-1R fragment delD4 in which the D4 subdomain of human CSF-1R-ECD was deleted (SEQ ID NO:65), the new anti-CSF-1R antibodies according to the invention could be selected. These antibodies show valuable properties like excellent ligand-dependant cell growth inhibition and at the same time ligand independent cell growth inhibition of NIH 3T3 cell, retrovirally infected with either an expression vector for full-length wildtype CSF-1R (SEQ ID NO:62) or mutant CSF-1R L301S Y969F (SEQ ID NO:63) whereby mutant CSF-1R recombinant cells are able to form spheroids independent of the CSF-1 ligand. Furthermore the antibodies according to the invention inhibit (both) human and cynomolgous macrophage differentiation, as they inhibit survival of human and cynomolgous monocytes.

II. Definitions

[0179] As used herein, the following terms have the meanings ascribed to them below unless otherwise specified.

[0180] The human CSF-1R (CSF-1 receptor; synonyms: M-CSF receptor; Macrophage colony-stimulating factor 1 receptor. *Fms* proto-oncogene, *c-fms*, SEQ ID NO: 22) is known since 1986 (Coussens, L., et al., *Nature* 320 (1986) 277-280). CSF-1R is a growth factor and encoded by the *c-fms* proto-oncogene (reviewed e.g. in Roth, P. and Stanley, E. R., *Curr. Top. Microbiol. Immunol.* 181 (1992) 141-67).

[0181] CSF-1R is the receptor for CSF-1 (macrophage colony stimulating factor, also called M-CSF) and IL-34 and mediates the biological effects of these cytokines (Sherr, C. J., et al., *Cell* 41 (1985) 665-676 (Lin, H., et al., *Science* 320 (2008) 807-811). The cloning of the colony stimulating factor-1 receptor (also called *c-fms*) was described for the first time in Roussel, M. F., et al., *Nature* 325 (1987) 549-552. In that publication, it was shown that CSF-1R had transforming potential dependent on changes in the C-terminal tail of the protein including the loss of the inhibitory tyrosine 969 phosphorylation which binds Cbl and thereby regulates receptor down regulation (Lee, P. S., et al., *Embo J.* 18 (1999) 3616-3628). CSF-1R is a single chain, transmembrane receptor tyrosine kinase (RTK) and a member of the family of immunoglobulin (Ig) motif containing RTKs characterized by 5 repeated Ig-like subdomains D1-D5 in the extracellular domain (ECD) of the receptor (Wang, Z., et al *Molecular and Cellular Biology* 13 (1993) 5348-5359). The human CSF-1R Extracellular Domain (CSF-1R-ECD)

(SEQ ID NO: 64) comprises all five extracellular Ig-like subdomains D1-D5. The human CSF-1R fragment delD4 (SEQ ID NO: 65) comprises the extracellular Ig-like subdomains D1-D3 and D5, but is missing the D4 subdomain. The human CSF-1R fragment D1-D3 (SEQ ID NO: 66) comprises the respective subdomains D1-D3. The sequences are listed without the signal peptide MGSQPGVLLLVATAWHGQ G (SEQ ID NO: 67).

[0182] The intracellular protein tyrosine kinase domain is interrupted by a unique insert domain that is also present in the other related RTK class III family members that include the platelet derived growth factor receptors (PDGFR), stem cell growth factor receptor (*c-Kit*) and fins-like cytokine receptor (FLT3). In spite of the structural homology among this family of growth factor receptors, they have distinct tissue-specific functions.

[0183] CSF-1R is mainly expressed on cells of the monocytic lineage and in the female reproductive tract and placenta. In addition expression of CSF-1R has been reported in Langerhans cells in skin, a subset of smooth muscle cells (Inaba, T., et al., *J. Biol. Chem.* 267 (1992) 5693-5699), B cells (Baker, A. H., et al., *Oncogene* 8 (1993) 371-378) and microglia (Sawada, M., et al., *Brain Res.* 509 (1990) 119-124). Cells with mutant human CSF-1R (SEQ ID NO: 23) are known to proliferate independently of ligand stimulation.

[0184] As used herein, “binding to human CSF-1R” or “specifically binding to human CSF-1R” refers to an antibody specifically binding to the human CSF-1R antigen with a binding affinity of KD-value of 1.0×10^{-8} mol/l or lower at 35° C., in one embodiment of a KD-value of 1.0×10^{-9} mol/l or lower at 35° C. The binding affinity is determined with a standard binding assay at 35° C., such as surface plasmon resonance technique (BIAcore®, GE-Healthcare Uppsala, Sweden) A method for determining the KD-value of the binding affinity is described in Example 9. Thus an “antibody binding to human CSF-1R” as used herein refers to an antibody specifically binding to the human CSF-1R antigen with a binding affinity of KD 1.0×10^{-8} mol/l or lower (preferably $1.0 \times 10^{0.8}$ mol/l- 1.0×10^{-12} mol/l) at 35° C., preferably of a KD 1.0×10^{-9} mol/l or lower at 35° C. (preferably 1.0×10^{-9} mol/l- 1.0×10^{-2} mol/l).

[0185] The “binding to human CSF-1R fragment delD4 (SEQ ID NO: 65) and to human CSF-1R Extracellular Domain (SEQ ID NO: 64)” as used herein is measured by a Surface Plasmon Resonance assay (Biacore assay) as described in Example 4. The human CSF-1R fragment delD4 (SEQ ID NO: 65) or human CSF-1R Extracellular Domain (SEQ ID NO: 64), respectively, are captured to the surface (each to a separate surface) and the test antibodies were added (each in a separate measurement) and the respective binding signals (Response Units (RU)) were determined. Reference signals (blank surface) were subtracted. If signals of nonbinding test antibodies were slightly below 0 the values were set as 0. Then the ratio of the respective binding signals (binding signal (RU) to human CSF-1R fragment delD4/binding signal (RU) to human CSF-1R Extracellular Domain (CSF-1R-ECD)) is determined. The antibodies according to the invention have a ratio of the binding signals (RU(delD4)/RU(CSF-1R-ECD)) of 1:50 or lower, preferably of 1:100 or lower (the lower included end is 0 (e.g. if the RU is 0, then the ratio is 0:50 or 0:100)).

[0186] This means that such anti-CSF-1R antibodies according to the invention do not bind to the human CSF-1R fragment delD4 (like the anti-CCR5 antibody m<CCR5>Pz03.1C5 (deposited as DSM ACC 2683 on 18 Aug. 2004 at DSMZ) and have binding signals for binding to the human CSF-1R fragment delD4 in the range of the anti-CCR5 antibody m<CCR5>Pz03.1C5, which are below 20 RU (Response Units), preferably below 10 RU in a Surface Plasmon Resonance (BIAcore) assay as shown in Example 4.

[0187] The term “binding to human CSF-1R fragment D1-D3” refers to a binding affinity determination by a Surface Plasmon Resonance assay (Biacore assay). The test antibody is captured to the surface and the human CSF-1R fragment D1-D3 (SEQ ID NO: 66) was added and the respective binding affinities were determined. The term “not binding to human CSF-1R fragment D1-D3” denotes that in such an assay the detected signal was in the area of no more than 1.2 fold of background signal and therefore no significant binding could be detected and no binding affinity could be determined (see Example 10).

[0188] The term “antibody” encompasses the various forms of antibodies including but not being limited to whole antibodies, antibody fragments, human antibodies, humanized antibodies, chimeric antibodies, T cell epitope depleted antibodies, and further genetically engineered antibodies as long as the characteristic properties according to the invention are retained. “Antibody fragments” comprise a portion of a full length antibody, preferably the variable domain thereof, or at least the antigen binding site thereof. Examples of antibody fragments include diabodies, single-chain antibody molecules, and multispecific antibodies formed from antibody fragments, scFv antibodies are, e.g., described in Houston, J. S., *Methods in Enzymol.* 203 (1991) 46-88). In addition, antibody fragments comprise single chain polypeptides having the characteristics of a V_H domain binding to CSF-1R, namely being able to assemble together with a V_L domain, or of a V_L domain binding to CSF-1R, namely being able to assemble together with a V_H domain to a functional antigen binding site and thereby providing the property.

[0189] The terms “monoclonal antibody” or “monoclonal antibody composition” as used herein refer to a preparation of antibody molecules of a single amino acid composition.

[0190] The term “chimeric antibody” refers to a monoclonal antibody comprising a variable region, i.e., binding region, from mouse and at least a portion of a constant region derived from a different source or species, usually prepared by recombinant DNA techniques. Chimeric antibodies comprising a mouse variable region and a human constant region are especially preferred. Such rat/human chimeric antibodies are the product of expressed immunoglobulin genes comprising DNA segments encoding rat immunoglobulin variable regions and DNA segments encoding human immunoglobulin constant regions. Other forms of “chimeric antibodies” encompassed by the present invention are those in which the class or subclass has been modified or changed from that of the original antibody. Such “chimeric” antibodies are also referred to as “class-switched antibodies.” Methods for producing chimeric antibodies involve conventional recombinant DNA and gene transfection techniques now well known in the art. See, e.g., Morrison, S. L., et al., *Proc. Natl. Acad. Sci. USA* 81 (1984) 6851-6855; U.S. Pat. Nos. 5,202,238 and 5,204,244.

[0191] The term “humanized antibody” refers to antibodies in which the framework or “complementarity determining regions” (CDR) have been modified to comprise the CDR of an immunoglobulin of different specificity as compared to that of the parent immunoglobulin. In a preferred embodiment, a murine CDR is grafted into the framework region of a human antibody to prepare the “humanized antibody.”

[0192] See e.g. Riechmann, L., et al., *Nature* 332 (1988) 323-327; and Neuberger, M. S., et al., *Nature* 314 (1985) 268-270. Optionally the framework region can be modified by further mutations. Also the CDRs can be modified by one or more mutations to generate antibodies according to the invention e.g. by mutagenesis based upon molecular modeling as described by Riechmann, L., et al., *Nature* 332 (1988) 323-327 and Queen, C., et al., *Proc. Natl. Acad. Sci. USA* 86 (1989) 10029-10033, or others. Particularly preferred CDRs correspond to those representing sequences recognizing the antigens noted above for chimeric antibodies. A “humanized version of an antibody according to the invention” (which is e.g. of mouse origin) refers to an antibody, which is based on the mouse antibody sequences in which the V_H and V_L are humanized by standard techniques (including CDR grafting and optionally subsequent mutagenesis of certain amino acids in the framework region and the CDRs). Preferably such humanized version is chimerized with a human constant region (see e.g. Sequences SEQ ID NO:57-61).

[0193] Other forms of “humanized antibodies” encompassed by the present invention are those in which the constant region has been additionally modified or changed from that of the original antibody to generate the properties according to the invention, especially in regard to C1q binding and/or Fc receptor (FcR) binding.

[0194] In the following examples the terms “Mab” or “muMab” refer to murine monoclonal antibodies such as Mab 2F11 or Mab 2E10, whereas the term “hMab” refers to humanized monoclonal versions of such murine antibodies such as hMab 2F11-c11, hMab 2F11-d8, hMab 2F11-e7, hMab 2F11-f12, etc.

[0195] The term “human antibody”, as used herein, is intended to include antibodies having variable and constant regions derived from human germ line immunoglobulin sequences. Human antibodies are well-known in the state of the art (van Dijk, M. A., and van de Winkel, J. G., *Curr. Opin. Chem. Biol.* 5 (2001) 368-374). Human antibodies can also be produced in transgenic animals (e.g., mice) that are capable, upon immunization, of producing a full repertoire or a selection of human antibodies in the absence of endogenous immunoglobulin production. Transfer of the human germ-line immunoglobulin gene array in such germ-line mutant mice will result in the production of human antibodies upon antigen challenge (see, e.g., Jakobovits, A., et al., *Proc. Natl. Acad. Sci. USA* 90 (1993) 2551-2555; Jakobovits, A., et al., *Nature* 362 (1993) 255-258; Brueggemann, M., et al., *Year Immunol.* 7 (1993) 33-40). Human antibodies can also be produced in phage display libraries (Hoogenboom, H. R., and Winter, G. *J. Mol. Biol.* 227 (1992) 381-388; Marks, J. D., et al., *J. Mol. Biol.* 222 (1991) 581-597). The techniques of Cole, et al., and Boerner, et al., are also available for the preparation of human monoclonal antibodies (Cole, S. P. C., et al., *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, p. 77 (1985); and Boerner, P., et al., *J. Immunol.* 147 (1991) 86-95). As already mentioned

for chimeric and humanized antibodies according to the invention the term “human antibody” as used herein also comprises such antibodies which are modified in the constant region to generate the properties according to the invention, especially in regard to C1q binding and/or FcR binding, e.g. by “class switching” i.e. change or mutation of Fc parts (e.g. from IgG1 to IgG4 and/or IgG1/IgG4 mutation).

[0196] The term “recombinant human antibody”, as used herein, is intended to include all human antibodies that are prepared, expressed, created or isolated by recombinant means, such as antibodies isolated from a host cell such as a NS0 or CHO cell or from an animal (e.g. a mouse) that is transgenic for human immunoglobulin genes or antibodies expressed using a recombinant expression vector transfected into a host cell. Such recombinant human antibodies have variable and constant regions in a rearranged form. The recombinant human antibodies according to the invention have been subjected to *in vivo* somatic hypermutation. Thus, the amino acid sequences of the VH and VL regions of the recombinant antibodies are sequences that, while derived from and related to human germ line VH and VL sequences, may not naturally exist within the human antibody germ line repertoire *in vivo*.

[0197] The antibodies according to the invention include, in addition, such antibodies having “conservative sequence modifications”, nucleotide and amino acid sequence modifications which do not affect or alter the above-mentioned characteristics of the antibody according to the invention. Modifications can be introduced by standard techniques known in the art, such as site-directed mutagenesis and PCR-mediated mutagenesis. Conservative amino acid substitutions include ones in which the amino acid residue is replaced with an amino acid residue having a similar side chain. Families of amino acid residues having similar side chains have been defined in the art. These families include amino acids with basic side chains (e.g., lysine, arginine, histidine), acidic side chains (e.g., aspartic acid, glutamic acid), uncharged polar side chains (e.g. glycine, asparagine, glutamine, serine, threonine, tyrosine, cysteine, tryptophan), nonpolar side chains (e.g., alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine), beta-branched side chains (e.g., threonine, valine, isoleucine) and aromatic side chains (e.g., tyrosine, phenylalanine, tryptophan, histidine). Thus, a predicted nonessential amino acid residue in a human anti-CSF-1R antibody can be preferably replaced with another amino acid residue from the same side chain family.

[0198] Amino acid substitutions can be performed by mutagenesis based upon molecular modeling as described by Riechmann, L., et al., *Nature* 332 (1988) 323-327 and Queen, C., et al., *Proc. Natl. Acad. Sci. USA* 86 (1989) 10029-10033.

[0199] The term “epitope” denotes a protein determinant of human CSF-1R capable of specifically binding to an antibody. Epitopes usually consist of chemically active surface groupings of molecules such as amino acids or sugar side chains and usually epitopes have specific three dimensional structural characteristics, as well as specific charge characteristics. Conformational and nonconformational epitopes are distinguished in that the binding to the former but not the latter is lost in the presence of denaturing solvents. Preferably an antibody according to the invention binds specifically to native and to denatured CSF-1R.

[0200] The “variable domain” (variable domain of a light chain (V_L), variable domain of a heavy chain (V_H)) as used herein denotes each of the pair of light and heavy chain domains which are involved directly in binding the antibody to the antigen. The variable light and heavy chain domains have the same general structure and each domain comprises four framework (FR) regions whose sequences are widely conserved, connected by three “hypervariable regions” (or complementary determining regions, CDRs). The framework regions adopt a β -sheet conformation and the CDRs may form loops connecting the β -sheet structure. The CDRs in each chain are held in their three-dimensional structure by the framework regions and form together with the CDRs from the other chain the antigen binding site. The antibody’s heavy and light chain CDR3 regions play a particularly important role in the binding specificity/affinity of the antibodies according to the invention and therefore provide a further object of the invention.

[0201] The term “antigen-binding portion of an antibody” when used herein refer to the amino acid residues of an antibody which are responsible for antigen-binding. The antigen-binding portion of an antibody comprises amino acid residues from the “complementary determining regions” or “CDRs”. “Framework” or “FR” regions are those variable domain regions other than the hypervariable region residues as herein defined. Therefore, the light and heavy chain variable domains of an antibody comprise from N- to C-terminus the domains FR1, CDR1, FR2, CDR2, FR3, CDR3, and FR4. Especially, CDR3 of the heavy chain is the region which contributes most to antigen binding and defines the antibody’s properties. CDR and FR regions are determined according to the standard definition of Kabat et al., *Sequences of Proteins of Immunological Interest*, 5th ed., Public Health Service, National Institutes of Health, Bethesda, Md. (1991) and/or those residues from a “hypervariable loop”.

[0202] The “Fc part” or “Fc portion” of an antibody is not involved directly in binding of an antibody to an antigen, but exhibit various effector functions. A “Fc part of an antibody” is a term well known to the skilled artisan and defined on the basis of papain cleavage of antibodies. Depending on the amino acid sequence of the constant region of their heavy chains, antibodies or immunoglobulins are divided in the classes: IgA, IgD, IgE, IgG and IgM, and several of these may be further divided into subclasses (isotypes), e.g. IgG1, IgG2, IgG3, and IgG4, IgA1, and IgA2. According to the heavy chain constant regions the different classes of immunoglobulins are called α , δ , ϵ , γ , and μ , respectively. The Fc part of an antibody is directly involved in ADCC (antibody-dependent cell-mediated cytotoxicity) and CDC (complement-dependent cytotoxicity) based on complement activation, C1q binding and Fc receptor binding. Complement activation (CDC) is initiated by binding of complement factor C1q to the Fc part of most IgG antibody subclasses. While the influence of an antibody on the complement system is dependent on certain conditions, binding to C1q is caused by defined binding sites in the Fc part. Such binding sites are known in the state of the art and described e.g. by Boackle, R. J., et al., *Nature* 282 (1979) 742-743, Lukas, T. J., et al., *J. Immunol.* 127 (1981) 2555-2560, Brunhouse, R., and Cebra, J. J., *Mol. Immunol.* 16 (1979) 907-917, Burton, D. R., et al., *Nature* 288 (1980) 338-344, Thommesen, J. E., et al., *Mol. Immunol.* 37 (2000) 995-1004, Idusogie, E. E., et al., *J. Immunol.* 164 (2000) 4178-4184, Hezareh, M., et

al., *J. Virology* 75 (2001) 12161-12168, Morgan, A., et al., *Immunology* 86 (1995) 319-324, EP 0307434. Such binding sites are e.g. L234, L235, D270, N297, E318, K320, K322, P331 and P329 (numbering according to EU index of Kabat, E. A., see below). Antibodies of subclass IgG1, IgG2 and IgG3 usually show complement activation and C1q and C3 binding, whereas IgG4 do not activate the complement system and do not bind C1q and C3.

[0203] In one embodiment the antibody according to the invention comprises a Fc part derived from human origin and preferably all other parts of the human constant regions. As used herein the term "Fc part derived from human origin" denotes a Fc part which is either a Fc part of a human antibody of the subclass IgG1, IgG2, IgG3 or IgG4, preferably a Fc part from human IgG1 subclass, a mutated Fc part from human IgG1 subclass (preferably with a mutation on L234A+L235A), a Fc part from human IgG4 subclass or a mutated Fc part from human IgG4 subclass (preferably with a mutation on S228P). Mostly preferred are the human heavy chain constant regions of SEQ ID NO: 58 (human IgG1 subclass), SEQ ID NO: 59 (human IgG1 subclass with mutations L234A and L235A), SEQ ID NO: 60 human IgG4 subclass), or SEQ ID NO: 61 (human IgG4 subclass with mutation S228P).

[0204] The terms "nucleic acid" or "nucleic acid molecule", as used herein, are intended to include DNA molecules and RNA molecules. A nucleic acid molecule may be single-stranded or double-stranded, but preferably is double-stranded DNA.

[0205] Nucleic acid is "operably linked" when it is plated into a functional relationship with another nucleic acid sequence. For example, DNA for a presequence or secretory leader is operably linked to DNA for a polypeptide if it is expressed as a preprotein that participates in the secretion of the polypeptide; a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to facilitate translation. Generally, "operably linked" means that the DNA sequences being linked are contiguous, and, in the case of a secretory leader, contiguous and in reading frame. However, enhancers do not have to be contiguous. Linking is accomplished by ligation at convenient restriction sites. If such sites do not exist, the synthetic oligonucleotide adaptors or linkers are used in accordance with conventional practice.

[0206] The term "amino acid" as used within this application denotes the group of naturally occurring carboxy α -amino acids comprising alanine (three letter code: ala, one letter code: A), arginine (arg, R), asparagine (asn, N), aspartic acid (asp, D), cysteine (cys, C), glutamine (gin, Q), glutamic acid (glu, E), glycine (gly, G), histidine (his, H), isoleucine (ile, I), leucine (leu, L), lysine (lys, K), methionine (met, M), phenylalanine (phe, F), proline (pro, P), serine (ser, S), threonine (thr, T), tryptophan (trp, W), tyrosine (tyr, Y), and valine (val, V).

[0207] As used herein, the expressions "cell", "cell line", and "cell culture" are used interchangeably and all such designations include progeny. Thus, the words "transformants" and "transformed cells" include the primary subject cell and cultures derived therefrom without regard for the number of transfers. It is also understood that all progeny may not be precisely identical in DNA content, due to deliberate or inadvertent mutations. Variant progeny that

have the same function or biological activity as screened for in the originally transformed cell are included.

[0208] An "effective amount" refers to an amount effective, at dosages and for periods of time necessary, to achieve the desired therapeutic or prophylactic result.

[0209] A "therapeutically effective amount" of a substance/molecule of the invention may vary according to factors such as the disease state, age, sex, and weight of the individual, and the ability of the substance/molecule, to elicit a desired response in the individual. A therapeutically effective amount encompasses an amount in which any toxic or detrimental effects of the substance/molecule are outweighed by the therapeutically beneficial effects. A therapeutically effective amount also encompasses an amount sufficient to confer benefit, e.g., clinical benefit.

[0210] A "prophylactically effective amount" refers to an amount effective, at dosages and for periods of time necessary, to achieve the desired prophylactic result. Typically, but not necessarily, since a prophylactic dose is used in subjects prior to or at an earlier stage of disease, the prophylactically effective amount would be less than the therapeutically effective amount. A prophylactically effective amount encompasses an amount sufficient to confer benefit, e.g., clinical benefit.

[0211] In the case of pre-cancerous, benign, early or late-stage tumors, the therapeutically effective amount of the angiogenic inhibitor may reduce the number of cancer cells; reduce the primary tumor size; inhibit (i.e., slow to some extent and preferably stop) cancer cell infiltration into peripheral organs; inhibit (i.e., slow to some extent and preferably stop) tumor metastasis; inhibit or delay, to some extent, tumor growth or tumor progression; and/or relieve to some extent one or more of the symptoms associated with the disorder. To the extent the drug may prevent growth and/or kill existing cancer cells, it may be cytostatic and/or cytotoxic. For cancer therapy, efficacy in vivo can, for example, be measured by assessing the duration of survival, time to disease progression (TTP), the response rates (RR), duration of response, and/or quality of life.

[0212] To "reduce" or "inhibit" is to decrease or reduce an activity, function, and/or amount as compared to a reference. In certain embodiments, by "reduce" or "inhibit" is meant the ability to cause an overall decrease of 20% or greater. In another embodiment, by "reduce" or "inhibit" is meant the ability to cause an overall decrease of 50% or greater. In yet another embodiment, by "reduce" or "inhibit" is meant the ability to cause an overall decrease of 75%, 85%, 90%, 95%, or greater. Reduce or inhibit can refer to the symptoms of the disorder being treated, the presence or size of metastases, the size of the primary tumor, or the size or number of the blood vessels in angiogenic disorders.

[0213] A "disorder" is any condition that would benefit from treatment including, but not limited to, chronic and acute disorders or diseases including those pathological conditions which predispose the mammal to the disorder in question. Disorders include cancer (including metastatic cancer), bone disorders (including, e.g., bone loss such as osteoporosis), and inflammatory disorders.

III. Antibodies and Methods of the Invention

[0214] The invention comprises an antibody binding to human CSF-1R, wherein the antibody binds to human CSF-1R fragment delD4 (comprising the extracellular subdomains D1-D3 and D5) (SEQ ID NO: 65) and to human

CSF-1R Extracellular Domain (CSF-1R-ECD) (comprising the extracellular subdomains D1-D5) (SEQ ID NO: 64) with a ratio of 1:50 or lower.

[0215] The invention further comprises an antibody comprising a heavy chain variable domain CDR3 region comprising SEQ ID NO: 1, SEQ ID NO: 9, SEQ ID NO:23, SEQ ID NO:31, SEQ ID NO:39, SEQ ID NO:47 or SEQ ID NO:55.

[0216] The invention further comprises an antibody wherein

[0217] a) the heavy chain variable domain comprises SEQ ID NO:7 and the light chain variable domain comprises SEQ ID NO:8,

[0218] b) the heavy chain variable domain comprises SEQ ID NO:15 and the light chain variable domain comprises SEQ ID NO: 16;

[0219] or a humanized version thereof.

[0220] The invention further comprises an antibody wherein

[0221] a) the heavy chain variable domain comprises SEQ ID NO:7 and the light chain variable domain comprises SEQ ID NO:8,

[0222] b) the heavy chain variable domain comprises SEQ ID NO:15 and the light chain variable domain comprises SEQ ID NO: 16;

[0223] c) the heavy chain variable domain comprises SEQ ID NO:75 and the light chain variable domain comprises SEQ ID NO:76;

[0224] d) the heavy chain variable domain comprises SEQ ID NO:83 and the light chain variable domain comprises SEQ ID NO:84;

[0225] or a humanized version thereof.

[0226] The invention further comprises an antibody wherein

[0227] the heavy chain variable domain comprises SEQ ID NO:7 and the light chain variable domain comprises SEQ ID NO:8, or a humanized version thereof.

[0228] In one embodiment the antibody comprises

[0229] a) a heavy chain variable domain comprising SEQ ID NO:23 and a light chain variable domain comprising SEQ ID NO:24, or

[0230] b) a heavy chain variable domain comprising SEQ ID NO:31 and a light chain variable domain comprising SEQ ID NO:32, or

[0231] c) a heavy chain variable domain comprising SEQ ID NO:39 and a light chain variable domain comprising SEQ ID NO:40, or

[0232] d) a heavy chain variable domain comprising SEQ ID NO:47 and a light chain variable domain comprising SEQ ID NO:48, or

[0233] e) a heavy chain variable domain comprising SEQ ID NO:55 and a light chain variable domain comprising SEQ ID NO:56.

[0234] In one embodiment the antibody comprises

[0235] a) a heavy chain variable domain comprising SEQ ID NO:23 and a light chain variable domain comprising SEQ ID NO:24, or

[0236] b) a heavy chain variable domain comprising SEQ ID NO:31 and a light chain variable domain comprising SEQ ID NO:32, or

[0237] c) a heavy chain variable domain comprising SEQ ID NO:39 and a light chain variable domain comprising SEQ ID NO:40, or

[0238] d) a heavy chain variable domain comprising SEQ ID NO:47 and a light chain variable domain comprising SEQ ID NO:48.

[0239] In one embodiment the antibody comprises

[0240] a heavy chain variable domain comprising SEQ ID NO:23 and a light chain variable domain comprising SEQ ID NO:24, or

[0241] In one embodiment the antibody comprises

[0242] a heavy chain variable domain comprising SEQ ID NO:31 and a light chain variable domain comprising SEQ ID NO:32.

[0243] In one embodiment the antibody comprises

[0244] a heavy chain variable domain comprising SEQ ID NO:39 and a light chain variable domain comprising SEQ ID NO:40.

[0245] In one embodiment the antibody comprises

[0246] a heavy chain variable domain comprising SEQ ID NO:47 and a light chain variable domain comprising SEQ ID NO:48.

[0247] The invention further comprises an antibody wherein

[0248] the heavy chain variable domain comprises SEQ ID NO:15 and the light chain variable domain comprises SEQ ID NO: 16, or a humanized version thereof.

[0249] The invention further comprises an antibody wherein

[0250] the heavy chain variable domain comprises SEQ ID NO:75 and the light chain variable domain comprises SEQ ID NO:76;

[0251] or a humanized version thereof.

[0252] The invention further comprises an antibody wherein

[0253] the heavy chain variable domain comprises SEQ ID NO:83 and the light chain variable domain comprises SEQ ID NO:84;

[0254] or a humanized version thereof.

[0255] The invention further comprises an antibody, wherein

[0256] a) the heavy chain variable domain comprises a CDR3 region comprising SEQ ID NO:1, a CDR2 region comprising SEQ ID NO: 2, and a CDR1 region comprising SEQ ID NO:3, and the light chain variable domain comprises a CDR3 region comprising SEQ ID NO: 4, a CDR2 region comprising SEQ ID NO:5, and a CDR1 region comprising SEQ ID NO:6, or,

[0257] b) the heavy chain variable domain comprises a CDR3 region comprising SEQ ID NO: 9, a CDR2 region comprising SEQ ID NO: 10, and a CDR1 region comprising SEQ ID NO: 11, and the light chain variable domain comprises a CDR3 region comprising SEQ ID NO:12, a CDR2 region comprising SEQ ID NO: 13, and a CDR1 region comprising SEQ ID NO: 14, or

[0258] c) the heavy chain variable domain comprises a CDR3 region comprising SEQ ID NO: 17, a CDR2 region comprising SEQ ID NO: 18, and a CDR1 region comprising SEQ ID NO:19, and the light chain variable domain comprises a CDR3 region comprising SEQ ID NO: 20, a CDR2 region comprising SEQ ID NO:21, and a CDR1 region comprising SEQ ID NO:22, or

[0259] d) the heavy chain variable domain comprises a CDR3 region comprising SEQ ID NO: 25, a CDR2 region comprising SEQ ID NO: 26, and a CDR1 region comprising SEQ ID NO: 27, and the light chain vari-

- SEQ ID NO:28, a CDR2 region comprising SEQ ID NO: 29, and a CDR1 region comprising SEQ ID NO: 30, or
- [0279] c) the heavy chain variable domain comprises a CDR3 region comprising SEQ ID NO: 33, a CDR2 region comprising SEQ ID NO: 34, and a CDR1 region comprising SEQ ID NO: 35, and the light chain variable domain comprises a CDR3 region comprising SEQ ID NO:36, a CDR2 region comprising SEQ ID NO: 37, and a CDR1 region comprising SEQ ID NO: 38, or
- [0280] d) the heavy chain variable domain comprises a CDR3 region comprising SEQ ID NO:41, a CDR2 region comprising SEQ ID NO: 42, and a CDR1 region comprising SEQ ID NO:43, and the light chain variable domain comprises a CDR3 region comprising SEQ ID NO: 44, a CDR2 region comprising SEQ ID NO:45, and a CDR1 region comprising SEQ ID NO:46, or
- [0281] e) the heavy chain variable domain comprises a CDR3 region comprising SEQ ID NO: 49, a CDR2 region comprising SEQ ID NO: 50, and a CDR1 region comprising SEQ ID NO: 51, and the light chain variable domain comprises a CDR3 region comprising SEQ ID NO:52, a CDR2 region comprising SEQ ID NO: 53, and a CDR1 region comprising SEQ ID NO: 54.
- [0282] In one embodiment the antibody comprises
- [0283] a) the heavy chain variable domain comprises a CDR3 region comprising SEQ ID NO: 17, a CDR2 region comprising SEQ ID NO: 18, and a CDR1 region comprising SEQ ID NO:19, and the light chain variable domain comprises a CDR3 region comprising SEQ ID NO: 20, a CDR2 region comprising SEQ ID NO:21, and a CDR1 region comprising SEQ ID NO:22, or
- [0284] b) the heavy chain variable domain comprises a CDR3 region comprising SEQ ID NO: 25, a CDR2 region comprising SEQ ID NO: 26, and a CDR1 region comprising SEQ ID NO: 27, and the light chain variable domain comprises a CDR3 region comprising SEQ ID NO:28, a CDR2 region comprising SEQ ID NO: 29, and a CDR1 region comprising SEQ ID NO: 30, or
- [0285] c) the heavy chain variable domain comprises a CDR3 region comprising SEQ ID NO: 33, a CDR2 region comprising SEQ ID NO: 34, and a CDR1 region comprising SEQ ID NO: 35, and the light chain variable domain comprises a CDR3 region comprising SEQ ID NO:36, a CDR2 region comprising SEQ ID NO: 37, and a CDR1 region comprising SEQ ID NO: 38, or
- [0286] d) the heavy chain variable domain comprises a CDR3 region comprising SEQ ID NO:41, a CDR2 region comprising SEQ ID NO: 42, and a CDR1 region comprising SEQ ID NO:43, and the light chain variable domain comprises a CDR3 region comprising SEQ ID NO: 44, a CDR2 region comprising SEQ ID NO:45, and a CDR1 region comprising SEQ ID NO:46.
- [0287] In one embodiment the antibody comprises
- [0288] the heavy chain variable domain comprises a CDR3 region comprising SEQ ID NO: 17, a CDR2 region comprising SEQ ID NO: 18, and a CDR1 region comprising SEQ ID NO: 19, and the light chain variable domain comprises a CDR3 region comprising SEQ ID NO: 20, a CDR2 region comprising SEQ ID NO:21, and a CDR1 region comprising SEQ ID NO:22.
- [0289] In one embodiment the antibody comprises
- [0290] the heavy chain variable domain comprises a CDR3 region comprising SEQ ID NO: 25, a CDR2 region comprising SEQ ID NO: 26, and a CDR1 region comprising SEQ ID NO: 27, and the light chain variable domain comprises a CDR3 region comprising SEQ ID NO:28, a CDR2 region comprising SEQ ID NO: 29, and a CDR1 region comprising SEQ ID NO: 30.
- [0291] In one embodiment the antibody comprises
- [0292] the heavy chain variable domain comprises a CDR3 region comprising SEQ ID NO: 33, a CDR2 region comprising SEQ ID NO: 34, and a CDR1 region comprising SEQ ID NO: 35, and the light chain variable domain comprises a CDR3 region comprising SEQ ID NO:36, a CDR2 region comprising SEQ ID NO: 37, and a CDR1 region comprising SEQ ID NO: 38.
- [0293] In one embodiment the antibody comprises
- [0294] the heavy chain variable domain comprises a CDR3 region comprising SEQ ID NO:41, a CDR2 region comprising SEQ ID NO: 42, and a CDR1 region comprising SEQ ID NO:43, and the light chain variable domain comprises a CDR3 region comprising SEQ ID NO: 44, a CDR2 region comprising SEQ ID NO:45, and a CDR1 region comprising SEQ ID NO:46.
- [0295] In one embodiment the antibody binding to human CSF-1R binds to human CSF-1R fragment delD4 (SEQ ID NO: 65) and to human CSF-1R-ECD (SEQ ID NO: 64) with a ratio of 1:50 or lower, further does not bind to human CSF-1R fragment D1-D3 (SEQ ID NO: 66).
- [0296] One embodiment of the invention is a screening method for selecting antibodies according to the invention comprising the following steps:
- [0297] a) determining the binding signal (Response Units (RU)) of anti-CSF-1R antibodies to human CSF-1R fragment delD4 (SEQ ID NO: 65) and to human CSF-1R Extracellular Domain (CSF-1R-ECD) (SEQ ID NO: 64) by a Surface Plasmon Resonance assay (Biacore assay).
- [0298] b) selecting antibodies with ratio of the binding signals (human CSF-1R fragment delD4/human CSF-1R Extracellular Domain (CSF-1R-ECD)) of 50:1 or lower.
- [0299] In one embodiment the determination is performed at 25° C.
- [0300] In one embodiment the screening method comprises as further steps the measuring of the binding of anti-CSF-1R antibodies to human CSF-1R fragment D1-D3 (SEQ ID NO: 66) (D1-D3) and the selecting of antibodies which show no binding to said fragment.
- [0301] In one embodiment the antibodies according to the invention inhibit CSF-1 binding to CSF-1R. In one embodiment with an IC50 of 200 ng/ml or lower, in one embodiment with an IC50 of 50 ng/ml or lower. The IC50 of inhibition of CSF-1 binding to CSF-1R can be determined as shown in Example 2.
- [0302] In one embodiment the antibodies according to the invention inhibit CSF-1-induced CSF-1R phosphorylation (in NIH3T3-CSF-1R recombinant cells).
- [0303] In one embodiment with an IC50 of 800 ng/ml or lower, in one embodiment with an IC50 of 600 ng/ml or

lower, in one embodiment with an IC₅₀ of 250 ng/ml or lower. The IC₅₀ of CSF-1-induced CSF-1R phosphorylation can be determined as shown in Example 3.

[0304] In one embodiment the antibodies according to the invention inhibit the growth of recombinant NIH3T3 cells expressing human CSF-1R (SEQ ID No: 62). In one embodiment with an IC₅₀ of 10 µg/ml or lower, in one embodiment with an IC₅₀ of 5 µg/ml or lower, in one embodiment with an IC₅₀ of 2 µg/ml or lower. In one embodiment with an IC₃₀ of 10 µg/ml or lower, in one embodiment with an IC₃₀ of 5 µg/ml or lower, in one embodiment with an IC₃₀ of 2 µg/ml or lower. The IC₅₀ value, the IC₃₀ value or the % growth inhibition is determined as shown in Example 5.

[0305] In one embodiment the antibodies according to the invention inhibit the growth of recombinant NIH3T3 cells expressing human mutant CSF-1R L301S Y969F (SEQ ID No: 63). In one embodiment with an IC₅₀ of 15 µg/ml or lower, in one embodiment with an IC₅₀ of 10 µg/ml or lower. In one embodiment with an IC₃₀ of 10 µg/ml or lower, in one embodiment with an IC₅₀ of 5 µg/ml or lower, in one embodiment with an IC₅₀ of 2 µg/ml or lower. The IC₅₀ value, the IC₃₀ value or the % growth inhibition is determined as shown in Example 5.

[0306] In one embodiment the antibodies according to the invention inhibit the growth of BeWo tumor cells (ATCC CCL-98) by 65% or more (at an antibody concentration of 10 µg/ml; and as compared to the absence of antibody). The % growth inhibition is determined as shown in Example 8. E.g. Mab 2F11 shows a growth inhibition of BeWo tumor cells of 70%.

[0307] In one embodiment the antibodies according to the invention inhibit (both) human and cynomolgous macrophage differentiation (which is indicated by the inhibition of the survival of human and cynomolgous monocytes as shown in Examples 7 and 8). In one embodiment the antibodies according to the invention inhibit the survival of human monocytes with an IC₅₀ of 0.15 µg/ml or lower, in one embodiment with an IC₅₀ of 0.10 µg/ml or lower. The inhibition of the survival of human monocytes is determined as shown in Example 7. In one embodiment the antibodies according to the invention inhibit the survival of cynomolgous monocytes by 80% or more, in one embodiment by 90% or more (at an antibody concentration of 5 µg/ml; and as compared to the absence of antibody). The inhibition of the survival of human monocytes is determined as shown in Example 8.

[0308] A further embodiment of the invention is a method for the production of an antibody against CSF-1R wherein the sequence of a nucleic acid encoding the heavy chain of a human IgG1 class antibody binding to human CSF-1R according to the invention said modified nucleic acid and the nucleic acid encoding the light chain of said antibody are inserted into an expression vector, said vector is inserted in a eukaryotic host cell, the encoded protein is expressed and recovered from the host cell or the supernatant.

[0309] The antibodies according to the invention are preferably produced by recombinant means. Therefore the antibody is preferably an isolated monoclonal antibody. Such recombinant methods are widely known in the state of the art and comprise protein expression in prokaryotic and eukaryotic cells with subsequent isolation of the antibody polypeptide and usually purification to a pharmaceutically acceptable purity. For the protein expression, nucleic acids

encoding light and heavy chains or fragments thereof are inserted into expression vectors by standard methods. Expression is performed in appropriate prokaryotic or eukaryotic host cells like CHO cells, NS0 cells, SP2/0 cells, HEK293 cells, COS cells, yeast, or *E. coli* cells, and the antibody is recovered from the cells (supernatant or cells after lysis).

[0310] Recombinant production of antibodies is well-known in the state of the art and described, for example, in the review articles of Makrides, S. C., *Protein Expr. Purif.* 17 (1999) 183-202; Geisse, S., et al., *Protein Expr. Purif.* 8 (1996) 271-282; Kaufman, R. J., *Mol. Biotechnol.* 16 (2000) 151-161; Werner, R. G., *Drug Res.* 48 (1998) 870-880.

[0311] The antibodies may be present in whole cells, in a cell lysate, or in a partially purified or substantially pure form. Purification is performed in order to eliminate other cellular components or other contaminants, e.g. other cellular nucleic acids or proteins, by standard techniques, including alkaline/SDS treatment, CsCl banding, column chromatography, agarose gel electrophoresis, and others well known in the art. See Ausubel, F., et al., ed. *Current Protocols in Molecular Biology*, Greene Publishing and Wiley Interscience, New York (1987).

[0312] The monoclonal antibodies are suitably separated from the culture medium by conventional immunoglobulin purification procedures such as, for example, protein A-Sepharose, hydroxylapatite chromatography, gel electrophoresis, dialysis, or affinity chromatography. DNA and RNA encoding the monoclonal antibodies are readily isolated and sequenced using conventional procedures. The hybridoma cells can serve as a source of such DNA and RNA. Once isolated, the DNA may be inserted into expression vectors, which are then transfected into host cells such as HEK 293 cells, CHO cells, or myeloma cells that do not otherwise produce immunoglobulin protein, to obtain the synthesis of recombinant monoclonal antibodies in the host cells.

[0313] Expression in NS0 cells is described by, e.g., Barnes, L. M., et al., *Cytotechnology* 32 (2000) 109-123; and Barnes, L. M., et al., *Biotech. Bioeng.* 73 (2001) 261-270. Transient expression is described by, e.g., Durocher, Y., et al., *Nucl. Acids. Res.* 30 (2002) E9. Cloning of variable domains is described by Orlandi, R., et al., *Proc. Natl. Acad. Sci. USA* 86 (1989) 3833-3837; Carter, P., et al., *Proc. Natl. Acad. Sci. USA* 89 (1992) 4285-4289; and Norderhaug, L., et al., *J. Immunol. Methods* 204 (1997) 77-87. A preferred transient expression system (HEK 293) is described by Schlaeger, E.-J., and Christensen, K., in *Cytotechnology* 30 (1999) 71-83 and by Schlaeger, E.-J., in *J. Immunol. Methods* 194 (1996) 191-199.

[0314] The control sequences that are suitable for prokaryotes, for example, include a promoter, optionally an operator sequence, and a ribosome binding site. Eukaryotic cells are known to utilize promoters, enhancers and polyadenylation signals.

[0315] In some embodiments, the antibody according to the invention is of human IgG1 subclass or of human IgG4 subclass. In one embodiment the antibody according to the invention is of human IgG1 subclass. In one embodiment the antibody according to the invention is of human IgG4 subclass.

[0316] In one embodiment the constant chains are of human origin. Such constant chains are well known in the state of the art and e.g. described by Kabat, E. A., (see e.g. Johnson, G. and Wu, T. T., *Nucleic Acids Res.* 28 (2000)

214-218). For example, a useful human heavy chain constant region comprises an amino acid sequence of SEQ ID NO: 58. For example, a useful human light chain constant region comprises an amino acid sequence of a kappa-light chain constant region of SEQ ID NO: 57.

[0317] Another aspect of the invention is an antibody binding to human CSF-1R, wherein

[0318] a) the heavy chain variable domain comprises SEQ ID NO:7 and the light chain variable domain comprises SEQ ID NO:8,

[0319] b) the heavy chain variable domain comprises SEQ ID NO: 15 and the light chain variable domain comprises SEQ ID NO: 16;

[0320] or a humanized version thereof.

[0321] Another aspect of the invention is an antibody binding to human CSF-1R, wherein

[0322] a) the heavy chain variable domain comprises SEQ ID NO:7 and the light chain variable domain comprises SEQ ID NO:8,

[0323] b) the heavy chain variable domain comprises SEQ ID NO: 15 and the light chain variable domain comprises SEQ ID NO: 16;

[0324] c) the heavy chain variable domain comprises SEQ ID NO:75 and the light chain variable domain comprises SEQ ID NO:76;

[0325] d) the heavy chain variable domain comprises SEQ ID NO:83 and the light chain variable domain comprises SEQ ID NO:84;

[0326] or a humanized version thereof.

[0327] Another aspect of the invention is an antibody binding to human CSF-1R, wherein the heavy chain variable domain comprises SEQ ID NO:7 and the light chain variable domain comprises SEQ ID NO:8, or a humanized version thereof.

[0328] Another aspect of the invention is an antibody binding to human CSF-1R, wherein

[0329] a) the heavy chain variable domain comprises SEQ ID NO:23 and the light chain variable domain comprises SEQ ID NO:24, or

[0330] b) the heavy chain variable domain comprises SEQ ID NO:31 and the light chain variable domain comprises SEQ ID NO:32, or

[0331] c) the heavy chain variable domain comprises SEQ ID NO:39 and the light chain variable domain comprises SEQ ID NO:40, or

[0332] d) the heavy chain variable domain comprises SEQ ID NO:47 and the light chain variable domain comprises SEQ ID NO:48, or

[0333] e) the heavy chain variable domain comprises SEQ ID NO:55 and the light chain variable domain comprises SEQ ID NO:56.

[0334] Another aspect of the invention is an antibody binding to human CSF-1R, wherein

[0335] a) the heavy chain variable domain comprises SEQ ID NO:23 and the light chain variable domain comprises SEQ ID NO:24, or

[0336] b) the heavy chain variable domain comprises SEQ ID NO:31 and the light chain variable domain comprises SEQ ID NO:32, or

[0337] c) the heavy chain variable domain comprises SEQ ID NO:39 and the light chain variable domain comprises SEQ ID NO:40, or

[0338] d) the heavy chain variable domain comprises SEQ ID NO:47 and the light chain variable domain comprises SEQ ID NO:48.

[0339] Another aspect of the invention is an antibody binding to human CSF-1R, wherein the heavy chain variable domain comprises SEQ ID NO:23 and the light chain variable domain comprises SEQ ID NO:24, or Another aspect of the invention is an antibody binding to human CSF-1R, wherein the heavy chain variable domain comprises SEQ ID NO:31 and the light chain variable domain comprises SEQ ID NO:32.

[0340] Another aspect of the invention is an antibody binding to human CSF-1R, wherein the heavy chain variable domain comprises SEQ ID NO:39 and the light chain variable domain comprises SEQ ID NO:40.

[0341] Another aspect of the invention is an antibody binding to human CSF-1R, wherein the heavy chain variable domain comprises SEQ ID NO:47 and the light chain variable domain comprises SEQ ID NO:48.

[0342] Another aspect of the invention is an antibody binding to human CSF-1R, wherein

[0343] the heavy chain variable domain comprises SEQ ID NO:15 and the light chain variable domain comprises SEQ ID NO:16, or a humanized version thereof.

[0344] Another aspect of the invention is an antibody binding to human CSF-1R, wherein

[0345] the heavy chain variable domain comprises SEQ ID NO:75 and the light chain variable domain comprises SEQ ID NO:76;

[0346] or a humanized version thereof.

[0347] Another aspect of the invention is an antibody binding to human CSF-1R, wherein

[0348] the heavy chain variable domain comprises SEQ ID NO:83 and the light chain variable domain comprises SEQ ID NO:84;

[0349] or a humanized version thereof.

[0350] Another aspect of the invention is an antibody binding to human CSF-1R, wherein

[0351] a) the heavy chain variable domain comprises a CDR3 region comprising SEQ ID NO:1, a CDR2 region comprising SEQ ID NO: 2, and a CDR1 region comprising SEQ ID NO:3, and the light chain variable domain comprises a CDR3 region comprising SEQ ID NO: 4, a CDR2 region comprising SEQ ID NO:5, and a CDR1 region comprising SEQ ID NO:6, or,

[0352] b) the heavy chain variable domain comprises a CDR3 region comprising SEQ ID NO: 9, a CDR2 region comprising SEQ ID NO: 10, and a CDR1 region comprising SEQ ID NO: 11, and the light chain variable domain comprises a CDR3 region comprising SEQ ID NO:12, a CDR2 region comprising SEQ ID NO: 13, and a CDR1 region comprising SEQ ID NO: 14, or

[0353] c) the heavy chain variable domain comprises a CDR3 region comprising SEQ ID NO: 17, a CDR2 region comprising SEQ ID NO: 18, and a CDR1 region comprising SEQ ID NO:19, and the light chain variable domain comprises a CDR3 region comprising SEQ ID NO: 20, a CDR2 region comprising SEQ ID NO:21, and a CDR1 region comprising SEQ ID NO:22, or

[0354] d) the heavy chain variable domain comprises a CDR3 region comprising SEQ ID NO: 25, a CDR2 region comprising SEQ ID NO: 26, and a CDR1 region comprising SEQ ID NO: 27, and the light chain vari-

ID NO:28, a CDR2 region comprising SEQ ID NO: 29, and a CDR1 region comprising SEQ ID NO: 30.

[0373] Another aspect of the invention is an antibody binding to human CSF-1R, wherein

[0374] the heavy chain variable domain comprises a CDR3 region comprising SEQ ID NO: 33, a CDR2 region comprising SEQ ID NO: 34, and a CDR1 region comprising SEQ ID NO: 35, and the light chain variable domain comprises a CDR3 region comprising SEQ ID NO:36, a CDR2 region comprising SEQ ID NO: 37, and a CDR1 region comprising SEQ ID NO: 38.

[0375] Another aspect of the invention is an antibody binding to human CSF-1R, wherein

[0376] the heavy chain variable domain comprises a CDR3 region comprising SEQ ID NO:41, a CDR2 region comprising SEQ ID NO: 42, and a CDR1 region comprising SEQ ID NO:43, and the light chain variable domain comprises a CDR3 region comprising SEQ ID NO: 44, a CDR2 region comprising SEQ ID NO:45, and a CDR1 region comprising SEQ ID NO:46.

[0377] The invention comprises a method for the treatment of a patient in need of therapy, comprising administering to the patient a therapeutically effective amount of an antibody according to the invention.

[0378] The invention comprises the use of an antibody according to the invention for therapy.

[0379] One preferred embodiment of the invention are the CSF-1R antibodies of the present invention for use in the treatment of "CSF-1R mediated diseases" or the CSF-1R antibodies of the present invention for use for the manufacture of a medicament in the treatment of "CSF-1R mediated diseases", which can be described as follows:

[0380] There are 3 distinct mechanisms by which CSF-1R signaling is likely involved in tumor growth and metastasis. The first is that expression of CSF-ligand and receptor has been found in tumor cells originating in the female reproductive system (breast, ovarian, endometrium, cervical) (Scholl, S. M., et al., *J. Natl. Cancer Inst.* 86 (1994) 120-126; Kacinski, B. M., *Mol. Reprod. Dev.* 46 (1997) 71-74; Ngan, H. Y., et al., *Eur. J. Cancer* 35 (1999) 1546-1550; Kirma, N., et al., *Cancer Res* 67 (2007) 1918-1926) and the expression has been associated with breast cancer xenograft growth as well as poor prognosis in breast cancer patients. Two point mutations were seen in CSF-1R in about 10-20% of acute myelocytic leukemia, chronic myelocytic leukemia and myelodysplasia patients tested in one study, and one of the mutations was found to disrupt receptor turnover (Ridge, S. A., et al., *Proc. Natl. Acad. Sci USA* 87 (1990) 1377-1380). However the incidence of the mutations could not be confirmed in later studies (Abu-Duhier, F. M., et al., *Br. J. Haematol.* 120 (2003) 464-470). Mutations were also found in some cases of hepatocellular cancer (Yang, D. H., et al., *Hepatobiliary Pancreat. Dis. Int.* 3 (2004) 86-89) and idiopathic myelofibrosis (Abu-Duhier, F. M., et al., *Br. J. Haematol.* 120 (2003) 464-470). Recently, in the GDM-1 cell line derived from a patient with myelomonoblastic leukemia the Y571D mutation in CSF-1R was identified (Chase, A., et al., *Leukemia* 23 (2009) 358-364).

[0381] Pigmented villonodular synovitis (PVNS) and Tenosynovial Giant cell tumors (TGCT) can occur as a result of a translocation that fuses the M-CSF gene to a collagen gene COL6A3 and results in overexpression of M-CSF (West, R. B., et al., *Proc. Natl. Acad. Sci. USA* 103 (2006)

690-695). A landscape effect is proposed to be responsible for the resulting tumor mass that consists of monocytic cells attracted by cells that express M-CSF. TGCTs are smaller tumors that can be relatively easily removed from fingers where they mostly occur. PVNS is more aggressive as it can recur in large joints and is not as easily controlled surgically.

[0382] The second mechanism is based on blocking signaling through M-CSF/CSF-1R at metastatic sites in bone which induces osteoclastogenesis, bone resorption and osteolytic bone lesions. Breast, multiple myeloma and lung cancers are examples of cancers that have been found to metastasize to the bone and cause osteolytic bone disease resulting in skeletal complications. M-CSF released by tumor cells and stroma induces the differentiation of hematopoietic myeloid monocyte progenitors to mature osteoclasts in collaboration with the receptor activator of nuclear factor kappa-B ligand-RANKL. During this process, M-CSF acts as a permissive factor by giving the survival signal to osteoclasts (Tanaka, S., et al., *J. Clin. Invest.* 91 (1993) 257-263). Inhibition of CSF-1R activity during osteoclast differentiation and maturation with a anti-CSF-1R antibody is likely to prevent unbalanced activity of osteoclasts that cause osteolytic disease and the associated skeletal related events in metastatic disease. Whereas breast, lung cancer and multiple myeloma typically result in osteolytic lesions, metastasis to the bone in prostate cancer initially has an osteoblastic appearance in which increased bone forming activity results in 'woven bone' which is different from typical lamellar structure of normal bone. During disease progression bone lesions display a significant osteolytic component as well as high serum levels of bone resorption and suggests that anti-resorptive therapy may be useful. Bisphosphonates have been shown to inhibit the formation of osteolytic lesions and reduced the number of skeletal-related events only in men with hormone-refractory metastatic prostate cancer but at this point their effect on osteoblastic lesions is controversial and bisphosphonates have not been beneficial in preventing bone metastasis or hormone responsive prostate cancer to date. The effect of anti-resorptive agents in mixed osteolytic/osteoblastic prostate cancer is still being studied in the clinic (Choueiri, M. B., et al., *Cancer Metastasis Rev.* 25 (2006) 601-609; Vessella, R. L. and Corey, E., *Clin. Cancer Res.* 12 (20 Pt 2) (2006) 6285s-6290s).

[0383] The third mechanism is based on the recent observation that tumor associated macrophages (TAM) found in solid tumors of the breast, prostate, ovarian and cervical cancers correlated with poor prognosis (Bingle, L., et al., *J. Pathol.* 196 (2002) 254-265; Pollard, J. W., *Nat. Rev. Cancer* 4 (2004) 71-78). Macrophages are recruited to the tumor by M-CSF and other chemokines. The macrophages can then contribute to tumor progression through the secretion of angiogenic factors, proteases and other growth factors and cytokines and may be blocked by inhibition of CSF-1R signaling. Recently it was shown by Zins et al (Zins, K., et al., *Cancer Res.* 67 (2007) 1038-1045) that expression of siRNA of Tumor necrosis factor alpha (TNF alpha), M-CSF or the combination of both would reduce tumor growth in a mouse xenograft model between 34% and 50% after intratumoral injection of the respective siRNA. siRNA targeting the TNF alpha secreted by the human SW620 cells reduced mouse M-CSF levels and led to reduction of macrophages in the tumor. In addition treatment of MCF7 tumor xenografts with an antigen binding fragment directed against M-CSF

did result in 40% tumor growth inhibition, reversed the resistance to chemotherapeutics and improved survival of the mice when given in combination with chemotherapeutics (Paulus, P., et al., *Cancer Res.* 66 (2006) 4349-4356).

[0384] TAMs are only one example of an emerging link between chronic inflammation and cancer. There is additional evidence for a link between inflammation and cancer as many chronic diseases are associated with an increased risk of cancer, cancers arise at sites of chronic inflammation, chemical mediators of inflammation are found in many cancers; deletion of the cellular or chemical mediators of inflammation inhibits development of experimental cancers and long-term use of anti-inflammatory agents reduce the risk of some cancers. A link to cancer exists for a number of inflammatory conditions among those *H. pylori* induced gastritis for gastric cancer, Schistosomiasis for bladder cancer, HHVX for Kaposi's sarcoma, endometriosis for ovarian cancer and prostatitis for prostate cancer (Balkwill, F., et al., *Cancer Cell* 7 (2005) 211-217). Macrophages are key cells in chronic inflammation and respond differentially to their microenvironment. There are two types of macrophages that are considered extremes in a continuum of functional states: M1 macrophages are involved in Type 1 reactions. These reactions involve the activation by microbial products and consequent killing of pathogenic microorganisms that result in reactive oxygen intermediates. On the other end of the extreme are M2 macrophages involved in Type 2 reactions that promote cell proliferation, tune inflammation and adaptive immunity and promote tissue remodeling, angiogenesis and repair (Mantovani, A., et al., *Trends Immunol.* 25 (2004) 677-686). Chronic inflammation resulting in established neoplasia is usually associated with M2 macrophages. A pivotal cytokine that mediates inflammatory reactions is TNF alpha that true to its name can stimulate anti-tumor immunity and hemorrhagic necrosis at high doses but has also recently been found to be expressed by tumor cells and acting as a tumor promoter (Zins, K., et al., *Cancer Res.* 67 (2007) 1038-1045; Balkwill, F., *Cancer Metastasis Rev.* 25 (2006) 409-416). The specific role of macrophages with respect to the tumor still needs to be better understood including the potential spatial and temporal dependence on their function and the relevance to specific tumor types.

[0385] Thus one embodiment of the invention are the CSF-1R antibodies of the present invention for use in the treatment of cancer. The term "cancer" as used herein may be, for example, lung cancer, non small cell lung (NSCL) cancer, bronchioloalveolar cell lung cancer, bone cancer, pancreatic cancer, skin cancer, cancer of the head or neck, cutaneous or intraocular melanoma, uterine cancer, ovarian cancer, rectal cancer, cancer of the anal region, stomach cancer, gastric cancer, colon cancer, breast cancer, uterine cancer, carcinoma of the fallopian tubes, carcinoma of the endometrium, carcinoma of the cervix, carcinoma of the vagina, carcinoma of the vulva, Hodgkin's Disease, cancer of the esophagus, cancer of the small intestine, cancer of the endocrine system, cancer of the thyroid gland, cancer of the parathyroid gland, cancer of the adrenal gland, sarcoma of soft tissue, cancer of the urethra, cancer of the penis, prostate cancer, cancer of the bladder, cancer of the kidney or ureter, renal cell carcinoma, carcinoma of the renal pelvis, mesothelioma, hepatocellular cancer, biliary cancer, neoplasms of the central nervous system (CNS), spinal axis tumors, brain stem glioma, glioblastoma multiforme, astrocytomas, schwannomas, ependymomas, medulloblastomas, menin-

giomas, squamous cell carcinomas, pituitary adenoma, lymphoma, lymphocytic leukemia, including refractory versions of any of the above cancers, or a combination of one or more of the above cancers. Preferably such cancer is a breast cancer, ovarian cancer, cervical cancer, lung cancer or prostate cancer. Preferably such cancers are further characterized by CSF-1 or CSF-1R expression or overexpression. One further embodiment the invention are the CSF-1R antibodies of the present invention for use in the simultaneous treatment of primary tumors and new metastases.

[0386] Thus another embodiment of the invention are the CSF-1R antibodies of the present invention for use in the treatment of periodontitis, histiocytosis X, osteoporosis, Paget's disease of bone (PDB), bone loss due to cancer therapy, periprosthetic osteolysis, glucocorticoid-induced osteoporosis, rheumatoid arthritis, psoriatic arthritis, osteoarthritis, inflammatory arthritides, and inflammation.

[0387] Rabello, D., et al., *Biochem. Biophys. Res. Commun.* 347 (2006) 791-796 has demonstrated that SNPs in the CSF1 gene exhibited a positive association with aggressive periodontitis: an inflammatory disease of the periodontal tissues that causes tooth loss due to resorption of the alveolar bone.

[0388] Histiocytosis X (also called Langerhans cell histiocytosis, LCH) is a proliferative disease of Langerhans dendritic cells that appear to differentiate into osteoclasts in bone and extra osseous LCH lesions. Langerhans cells are derived from circulating monocytes. Increased levels of M-CSF that have been measured in scra and lesions were found to correlate with disease severity (da Costa, C. E., et al., *J. Exp. Med.* 201 (2005) 687-693). The disease occurs primarily in a pediatric patient population and has to be treated with chemotherapy when the disease becomes systemic or is recurrent.

[0389] The pathophysiology of osteoporosis is mediated by loss of bone forming osteoblasts and increased osteoclast dependent bone resorption. Supporting data has been described by Cenci et al showing that an anti-M-CSF antibody injection preserves bone density and inhibits bone resorption in ovariectomized mice (Cenci, S., et al., *J. Clin. Invest.* 105 (2000) 1279-1287). Recently a potential link between postmenopausal bone loss due to estrogen deficiency was identified and found that the presence of TNF alpha producing T-cell affected bone metabolism (Roggia, C., et al., *Minerva Med.* 95 (2004) 125-132). A possible mechanism could be the induction of M-CSF by TNF alpha in vivo. An important role for M-CSF in TNF-alpha-induced osteoclastogenesis was confirmed by the effect of an antibody directed against M-CSF that blocked the TNF alpha induced osteolysis in mice and thereby making inhibitors of CSF-1R signaling potential targets for inflammatory arthritis (Kitaura, H., et al., *J. Clin. Invest.* 115 (2005) 3418-3427).

[0390] Paget's disease of bone (PDB) is the second most common bone metabolism disorder after osteoporosis in which focal abnormalities of increased bone turnover lead to complications such as bone pain, deformity, pathological fractures and deafness. Mutations in four genes have been identified that regulate normal osteoclast function and predispose individuals to PDB and related disorders: insertion mutations in TNFRSF11A, which encodes receptor activator of nuclear factor (NF) kappaB (RANK)—a critical regulator of osteoclast function, inactivating mutations of TNFRSF11B which encodes osteoprotegerin (a decoy receptor for RANK ligand), mutations of the sequestosome

1 gene (SQSTM1), which encodes an important scaffold protein in the NF κ B pathway and mutations in the valosin-containing protein (VCP) gene. This gene encodes VCP, which has a role in targeting the inhibitor of NF κ B for degradation by the proteasome (Daroszewska, A. and Ralston, S. H., *Nat. Clin. Pract. Rheumatol.* 2 (2006) 270-277). Targeted CSF-1R inhibitors provide an opportunity to block the deregulation of the RANKL signaling indirectly and add an additional treatment option to the currently used bisphosphonates.

[0391] Cancer therapy induced bone loss especially in breast and prostate cancer patients is an additional indication where a targeted CSF-1R inhibitor could prevent bone loss (Lester, J. E., et al., *Br. J. Cancer* 94 (2006) 30-35). With the improved prognosis for early breast cancer the long-term consequences of the adjuvant therapies become more important as some of the therapies including chemotherapy, irradiation, aromatase inhibitors and ovary ablation affect bone metabolism by decreasing the bone mineral density, resulting in increased risk for osteoporosis and associated fractures (Lester, J. E., et al., *Br. J. Cancer* 94 (2006) 30-35). The equivalent to adjuvant aromatase inhibitor therapy in breast cancer is androgen ablation therapy in prostate cancer which leads to loss of bone mineral density and significantly increases the risk of osteoporosis-related fractures (Stoch, S. A., et al., *J. Clin. Endocrinol. Metab.* 86 (2001) 2787-2791).

[0392] Targeted inhibition of CSF-1R signaling is likely to be beneficial in other indications as well when targeted cell types include osteoclasts and macrophages e.g. treatment of specific complications in response to joint replacement as a consequence of rheumatoid arthritis. Implant failure due to periprosthetic bone loss and consequent loosening of prostheses is a major complication of joint replacement and requires repeated surgery with high socioeconomic burdens for the individual patient and the health-care system. To date, there is no approved drug therapy to prevent or inhibit periprosthetic osteolysis (Drees, P., et al., *Nat. Clin. Pract. Rheumatol.* 3 (2007) 165-171).

[0393] Glucocorticoid-induced osteoporosis (GIOP) is another indication in which a CSF-1R inhibitor could prevent bone loss after longterm glucocorticocosteroid use that is given as a result of various conditions among those chronic obstructive pulmonary disease, asthma and rheumatoid arthritis (Guzman-Clark, J. R., et al., *Arthritis Rheum.* 57 (2007) 140-146; Feldstein, A. C., et al., *Osteoporos. Int.* 16 (2005) 2168-2174).

[0394] Rheumatoid arthritis, psoriatic arthritis and inflammatory arthritides are in itself potential indications for CSF-1R signaling inhibitors in that they consist of a macrophage component and to a varying degree bone destruction (Ritchlin, C. T., et al., *J. Clin. Invest.* 111 (2003) 821-831). Osteoarthritis and rheumatoid arthritis are inflammatory autoimmune disease caused by the accumulation of macrophages in the connective tissue and infiltration of macrophages into the synovial fluid, which is at least partially mediated by M-CSF. Campbell, I., K., et al., *J. Leukoc. Biol.* 68 (2000) 144-150, demonstrated that M-CSF is produced by human-joint tissue cells (chondrocytes, synovial fibroblasts) in vitro and is found in synovial fluid of patients with rheumatoid arthritis, suggesting that it contributes to the synovial tissue proliferation and macrophage infiltration which is associated with the pathogenesis of the disease. Inhibition of CSF-1R signaling is likely to control the number of macrophages in the joint and alleviate the pain

from the associated bone destruction. In order to minimize adverse affects and to further understand the impact of the CSF-1R signaling in these indications, one method is to specifically inhibit CSF-1R without targeting a myriad other kinases, such as Raf kinase.

[0395] Recent literature reports correlate increased circulating M-CSF with poor prognosis and atherosclerotic progression in chronic coronary artery disease (Saitoh, T., et al., *J. Am. Coll. Cardiol.* 35 (2000) 655-665; Ikonomidis, I., et al., *Eur. Heart. J.* 26 (2005) p. 1618-1624); M-CSF influences the atherosclerotic process by aiding the formation of foam cells (macrophages with ingested oxidized LDL) that express CSF-1R and represent the initial plaque (Murayama, T., et al., *Circulation* 99 (1999) 1740-1746).

[0396] Expression and signaling of M-CSF and CSF-1R is found in activated microglia. Microglia, which are resident macrophages of the central nervous system, can be activated by various insults, including infection and traumatic injury. M-CSF is considered a key regulator of inflammatory responses in the brain and M-CSF levels increase in HIV-1, encephalitis, Alzheimer's disease (AD) and brain tumors. Microgliosis as a consequence of autocrine signaling by M-CSF/CSF-1R results in induction of inflammatory cytokines and nitric oxides being released as demonstrated by e.g. using an experimental neuronal damage model (Hao, A. J., et al., *Neuroscience* 112 (2002) 889-900; Murphy, G. M., Jr., et al., *J. Biol. Chem.* 273 (1998) 20967-20971). Microglia that have increased expression of CSF-1R are found to surround plaques in AD and in the amyloid precursor protein V717F transgenic mouse model of AD (Murphy, G. M., Jr., et al., *Am. J. Pathol.* 157 (2000) 895-904). On the other hand op/op mice with fewer microglia in the brain resulted in fibrillar deposition of A-beta and neuronal loss compared to normal control suggesting that microglia do have a neuroprotective function in the development of AD lacking in the op/op mice (Kaku, M., et al., *Brain Res. Brain Res. Protoc.* 12 (2003) 104-108).

[0397] Expression and signaling of M-CSF and CSF-1R is associated with inflammatory bowel disease (IBD) (WO 2005/046657). The term "inflammatory bowel disease" refers to serious, chronic disorders of the intestinal tract characterized by chronic inflammation at various sites in the gastrointestinal tract, and specifically includes ulcerative colitis (UC) and Crohn's disease.

[0398] The invention comprises an antibody binding to human CSF-1R comprising the above mentioned epitope binding properties or alternatively by the above mentioned amino acid sequences and amino acid sequence fragments for the treatment of cancer.

[0399] The invention comprises an antibody binding to human CSF-1R comprising the above mentioned epitope binding properties or alternatively by the above mentioned amino acid sequences and amino acid sequence fragments for the treatment of bone loss.

[0400] The invention comprises an antibody binding to human CSF-1R comprising the above mentioned epitope binding properties or alternatively by the above mentioned amino acid sequences and amino acid sequence fragments for the prevention or treatment of metastasis.

[0401] The invention comprises an antibody binding to human CSF-1R comprising the above mentioned epitope binding properties or alternatively by the above mentioned amino acid sequences and amino acid sequence fragments for treatment of inflammatory diseases.

- [0402] The invention comprises the use of an antibody comprising the antibody binding to human CSF-1R, wherein the antibody comprises the above mentioned epitope binding properties or alternatively by the above mentioned amino acid sequences and amino acid sequence fragments for the treatment of cancer or alternatively for the manufacture of a medicament for the treatment of cancer.
- [0403] The invention comprises the use of an antibody comprising the antibody binding to human CSF-1R, wherein the antibody comprises the above mentioned epitope binding properties or alternatively by the above mentioned amino acid sequences and amino acid sequence fragments for the treatment of bone loss or alternatively for the manufacture of a medicament for the treatment of bone loss.
- [0404] The invention comprises the use of an antibody comprising the antibody binding to human CSF-1R, wherein the antibody comprises the above mentioned epitope binding properties or alternatively by the above mentioned amino acid sequences and amino acid sequence fragments for the prevention or treatment of metastasis or alternatively for the manufacture of a medicament for the prevention or treatment of metastasis.
- [0405] The invention comprises the use of an antibody comprising the antibody binding to human CSF-1R, wherein the antibody comprises the above mentioned epitope binding properties or alternatively by the above mentioned amino acid sequences and amino acid sequence fragments for treatment of inflammatory diseases or alternatively for the manufacture of a medicament for the treatment of inflammatory diseases.
- [0406] A further embodiment of the invention is a method for the production of an antibody against CSF-1R wherein the sequence of a nucleic acid encoding the heavy chain of a human IgG1 class antibody binding to human CSF-1R according to the invention said modified nucleic acid and the nucleic acid encoding the light chain of said antibody are inserted into an expression vector, said vector is inserted in a eukaryotic host cell, the encoded protein is expressed and recovered from the host cell or the supernatant.
- [0407] The antibodies according to the invention are preferably produced by recombinant means. Such methods are widely known in the state of the art and comprise protein expression in prokaryotic and eukaryotic cells with subsequent isolation of the antibody polypeptide and usually purification to a pharmaceutically acceptable purity. For the protein expression nucleic acids encoding light and heavy chains or fragments thereof are inserted into expression vectors by standard methods. Expression is performed in appropriate prokaryotic or eukaryotic host cells, such as CHO cells, NS0 cells, SP2/0 cells, HEK293 cells, COS cells, yeast, or *E. coli* cells, and the antibody is recovered from the cells (from the supernatant or after cells lysis).
- [0408] Recombinant production of antibodies is well-known in the state of the art and described, for example, in the review articles of Makrides, S. C., *Protein Expr. Purif.* 17 (1999) 183-202; Geisse, S., et al., *Protein Expr. Purif.* 8 (1996) 271-282; Kaufman, R. J., *Mol. Biotechnol.* 16 (2000) 151-161; Werner, R. G., *Drug Res.* 48 (1998) 870-880.
- [0409] The antibodies may be present in whole cells, in a cell lysate, or in a partially purified, or substantially pure form. Purification is performed in order to eliminate other cellular components or other contaminants, e.g. other cellular nucleic acids or proteins, by standard techniques, including alkaline/SDS treatment, CsCl banding, column chromatography, agarose gel electrophoresis, and others well known in the art. See Ausubel, F., et al., ed. *Current Protocols in Molecular Biology*, Greene Publishing and Wiley Interscience, New York (1987).
- [0410] Expression in NS0 cells is described by, e.g., Barnes, L. M., et al., *Cytotechnology* 32 (2000) 109-123; Barnes, L. M., et al., *Biotech. Bioeng.* 73 (2001) 261-270. Transient expression is described by, e.g., Durocher, Y., et al., *Nucl. Acids. Res.* 30 (2002) E9. Cloning of variable domains is described by Orlandi, R., et al., *Proc. Natl. Acad. Sci. USA* 86 (1989) 3833-3837; Carter, P., et al., *Proc. Natl. Acad. Sci. USA* 89 (1992) 4285-4289; Norderhaug, L., et al., *J. Immunol. Methods* 204 (1997) 77-87. A preferred transient expression system (HEK 293) is described by Schlaeger, E.-J. and Christensen, K., in *Cytotechnology* 30 (1999) 71-83, and by Schlaeger, E.-J., in *J. Immunol. Methods* 194 (1996) 191-199.
- [0411] Nucleic acid molecules encoding amino acid sequence variants of anti-CSF-1R antibody are prepared by a variety of methods known in the art. These methods include, but are not limited to, isolation from a natural source (in the case of naturally occurring amino acid sequence variants) or preparation by oligonucleotide-mediated (or site-directed) mutagenesis, PCR mutagenesis, and cassette mutagenesis of an earlier prepared variant or a non-variant version of humanized anti-CSF-1R antibody.
- [0412] The heavy and light chain variable domains according to the invention are combined with sequences of promoter, translation initiation, constant region, 3' untranslated region, polyadenylation, and transcription termination to form expression vector constructs. The heavy and light chain expression constructs can be combined into a single vector, co-transfected, serially transfected, or separately transfected into host cells which are then fused to form a single host cell expressing both chains.
- [0413] In another aspect, the present invention provides a composition, e.g. a pharmaceutical composition, containing one or a combination of monoclonal antibodies, or the antigen-binding portion thereof, of the present invention, formulated together with a pharmaceutically acceptable carrier.
- [0414] As used herein, "pharmaceutically acceptable carrier" includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption/resorption delaying agents, and the like that are physiologically compatible. Preferably, the carrier is suitable for injection or infusion.
- [0415] A composition of the present invention can be administered by a variety of methods known in the art. As will be appreciated by the skilled artisan, the route and/or mode of administration will vary depending upon the desired results.
- [0416] Pharmaceutically acceptable carriers include sterile aqueous solutions or dispersions and sterile powders for the preparation of sterile injectable solutions or dispersion. The use of such media and agents for pharmaceutically active substances is known in the art. In addition to water, the carrier can be, for example, an isotonic buffered saline solution.
- [0417] Regardless of the route of administration selected, the compounds of the present invention, which may be used in a suitable hydrated form, and/or the pharmaceutical

compositions of the present invention, are formulated into pharmaceutically acceptable dosage forms by conventional methods known to those of skill in the art.

[0418] Actual dosage levels of the active ingredients in the pharmaceutical compositions of the present invention may be varied so as to obtain an amount of the active ingredient which is effective to achieve the desired therapeutic response for a particular patient, composition, and mode of administration, without being toxic to the patient (effective amount). The selected dosage level will depend upon a variety of pharmacokinetic factors including the activity of the particular compositions of the present invention employed, or the ester, salt or amide thereof, the route of administration, the time of administration, the rate of excretion of the particular compound being employed, other drugs, compounds and/or materials used in combination with the particular compositions employed, the age, sex, weight, condition, general health and prior medical history of the patient being treated, and like factors well known in the medical arts.

[0419] The invention comprises the use of the antibodies according to the invention for the treatment of a patient suffering from cancer, especially from colon, lung or pancreas cancer.

[0420] The invention comprises also a method for the treatment of a patient suffering from such disease.

[0421] The invention further provides a method for the manufacture of a pharmaceutical composition comprising an effective amount of an antibody according to the invention together with a pharmaceutically acceptable carrier and the use of the antibody according to the invention for such a method.

[0422] The invention further provides the use of an antibody according to the invention in an effective amount for the manufacture of a pharmaceutical agent, preferably together with a pharmaceutically acceptable carrier, for the treatment of a patient suffering from cancer.

[0423] The invention also provides the use of an antibody according to the invention in an effective amount for the manufacture of a pharmaceutical agent, preferably together with a pharmaceutically acceptable carrier, for the treatment of a patient suffering from cancer.

EXAMPLES

[0424] The examples and figures are provided to aid the understanding of the present invention, the true scope of which is set forth in the appended claims. It is understood that modifications can be made in the procedures set forth without departing from the spirit of the invention.

Example 1

[0425] Generation of a Hybridoma Cell Line Producing Anti-CSF-1R Antibodies

[0426] Immunization Procedure of NMR1 Mice

[0427] NMR1 mice were immunized with an expression vector pDisplay™ (Invitrogen, USA) encoding the extracellular domain of huCSF-1R by utilizing electroporation. Every mouse was 4 times immunized with 100 µg DNA. When serum titers of anti-huCSF-1R were found to be sufficient, mice were additionally boosted once with 50 µg of a 1:1 mixture huCSF-1R ECD/huCSF-1R ECDhuFc chimera in 200 µl PBS intravenously (i.v.) 4 and 3 days before fusion.

[0428] Antigen Specific ELISA

[0429] Anti-CSF-1R titers in sera of immunized mice were determined by antigen specific ELISA.

[0430] 0.3 µg/ml huCSF-1R-huFc chimera (soluble extracellular domain) was captured on a streptavidin plate (MaxiSorb; MicroCoat, DE, Cat. No. 11974998/MC1099) with 0.1 mg/ml biotinylated anti Fcγ (Jackson ImmunoResearch, Cat. No. 109-066-098) and horse radish peroxidase (HRP)-conjugated F(ab')₂ anti mouse IgG (GE Healthcare, UK, Cat. No. NA9310V) diluted 1/800 in PBS/0.05% Tween20/0.5% BSA was added. Sera from all taps were diluted 1/40 in PBS/0.05% Tween20/0.5% BSA and serially diluted up to 1/1638400. Diluted sera were added to the wells. Pre-tap serum was used as negative control. A dilution series of mouse anti-human CSF-1R Mab3291 (R&D Systems, UK) from 500 ng/ml to 0.25 ng/ml was used as positive control. All components were incubated together for 1.5 hours. Wells were washed 6 times with PBST (PBS/0.2% Tween20) and assays were developed with freshly prepared ABTS solution (1 mg/ml) (ABTS: 2,2'-azino bis (3-ethylbenzthiazoline-6-sulfonic acid) for 10 minutes at RT. Absorbance was measured at 405 nm.

[0431] Hybridoma Generation

[0432] The mouse lymphocytes can be isolated and fused with a mouse myeloma cell line using PEG based standard protocols to generate hybridomas. The resulting hybridomas are then screened for the production of antigen-specific antibodies. For example, single cell suspensions of splenic derived lymphocytes from immunized mice are fused to Ag8 non-secreting mouse myeloma cells P3X63Ag8.653 (ATCC, CRL-1580) with 50% PEG. Cells are plated at approximately 10⁴ in flat bottom 96 well micro titer plate, followed by about two weeks incubation in selective medium. Individual wells are then screened by ELISA for human anti-CSF-1R monoclonal IgM and IgG antibodies. Once extensive hybridoma growth occurs, the antibody secreting hybridomas are replated, screened again, and if still positive for human IgG, anti-CSF-1R monoclonal antibodies, can be subcloned by FACS. The stable subclones are then cultured in vitro to produce antibody in tissue culture medium for characterization. Antibodies according to the invention could be selected using the determination of the binding of anti-CSF-1R antibodies to human CSF-1R fragment delD4 and to human CSF-1R Extracellular Domain (CSF-1R-ECD) as described in Example 4, as well as the determination of growth inhibition of NIH3T3 cells transfected with wildtype CSF-1R (ligand dependent signalling) or mutant CSF-1R L301S Y969F (ligand independent signalling) under treatment with anti-CSF-1R monoclonal antibodies as described in Example 5.

[0433] Culture of Hybridomas

[0434] Generated muMAb hybridomas were cultured in RPMI 1640 (PAN—Catalogue No. (Cat. No.) PO4-17500) supplemented with 2 mM L-glutamine (GIBCO—Cat. No. 35050-038), 1 mM Na-Pyruvat (GIBCO—Cat. No. 11360-039), 1×NEAA (GIBCO—Cat. No. 11140-035), 10% FCS (PAA—Cat. No. A15-649), 1× Pen Strep (Roche—Cat. No. 1074440), 1× Nutridoma CS (Roche—Cat. No. 1363743), 50 µM Mercaptoethanol (GIBCO—Cat. No. 31350-010) and 50 U/ml IL 6 mouse (Roche—Cat. No. 1 444 581) at 37° C. and 5% CO₂. Some of the resulting mouse antibodies have been humanized (e.g. Mab 2F11) and been expressed recombinantly.

Example 2

[0435] Inhibition of CSF-1 Binding to CSF-1R (ELISA)

[0436] By setting-up this assay to first allow for anti-CSF-1R antibody binding to the CSF-1R-ECD followed by detection of ligand not bound to the receptor both-ligand displacing antibodies and dimerization inhibitor anti-CSF-1R antibodies—can be tested. The test was performed on 384 well microtiter plates (MicroCoat, DE, Cat. No. 464718) at RT. After each incubation step plates were washed 3 times with PBST.

[0437] At the beginning, plates were coated with 0.5 mg/ml goat F(ab')₂ biotinylated anti Fc γ (Jackson ImmunoResearch, Cat. No. 109-006-170) for 1 hour (h).

[0438] Thereafter the wells were blocked with PBS supplemented with 0.2% Tween®-20 and 2% BSA (Roche Diagnostics GmbH, DE) for 0.5 h. 75 ng/ml of huCSF-1R-huFc chimera (which forms the dimeric soluble extracellular domain of huCSF-1R) was immobilized to plate for 1 h. Then dilutions of purified antibodies in PBS/0.05% Tween20/0.5% BSA were incubated for 1 h. After adding a mixture of 3 ng/ml CSF-1 (Biomol, DE, Cat. No. 60530), 50 ng/ml biotinylated anti CSF-1 clone BAF216 (R&D Systems, UK) and 1:5000 diluted streptavidin HRP (Roche Diagnostics GmbH, DE, Cat. No. 11089153001) for 1 h the plates were washed 6 times with PBST. Anti CSF-1R SC 2-4A5 (Santa Cruz Biotechnology, US), which inhibits the ligand-receptor interaction, was used as positive control. Plates were developed with freshly prepared BM Blue® POD substrate solution (BM Blue®: 3,3'-5,5'-Tetramethylbenzidine, Roche Diagnostics GmbH, DE, Cat. No. 11484281001) for 30 minutes at RT. Absorbance was measured at 370 nm. A decrease of absorbance is found, if the anti-CSF-1R antibody causes a release of CSF-1 from the dimeric complex. All anti-CSF-1R antibodies showed significant inhibition of the CSF-1 interaction with CSF-1R (see Table 1). Anti CSF-1R SC 2-4A5 (Santa Cruz Biotechnology, US see also Sherr, C. J. et al., Blood 73 (1989) 1786-1793), which inhibits the ligand-receptor interaction, was used as reference control.

TABLE 1

Calculated IC₅₀ values for the inhibition of the CSF-1/CSF-1R interaction

CSF-1R Mab	IC ₅₀ CSF-1/CSF-1R Inhibition [ng/ml]
Mab 2F11	19.3
Mab 2E10	20.6
Mab 2H7	18.2
Mab 1G10	11.8
SC-2-4A5	35.2

Example 3

[0439] Inhibition of CSF-1-Induced CSF-1R Phosphorylation in NIH3T3-CSF-1R Recombinant Cells

[0440] 4.5×10³ NIH 3T3 cells, retrovirally infected with an expression vector for full-length CSF-1R, were cultured in DMEM (PAA Cat. No. E15-011), 2 mM L-glutamine (Sigma, Cat. No. G7513, 2 mM Sodium pyruvate, 1× nonessential aminoacids, 10% FKS (PAA, Cat. No. A15-649) and 100 g/ml PenStrep (Sigma, Cat. No. P4333 [10 mg/ml]) until they reached confluency. Thereafter cells were washed with serum-free DMEM media (PAA Cat. No.

E15-011) supplemented with sodium selenite [5 ng/ml] (Sigma, Cat. No. S9133), transferrin [10 μg/ml] (Sigma, Cat. No. T8158), BSA [400 g/ml] (Roche Diagnostics GmbH, Cat. No. 10735078), 4 mM L-glutamine (Sigma, Cat. No. G7513), 2 mM sodium pyruvate (Gibco, Cat. No. 11360), 1× nonessential aminoacids (Gibco, Cat: 11140-035), 2-mercaptoethanol [0.05 mM] (Merck, Cat. No. M7522), 100 μg/ml and PenStrep (Sigma, Cat. No. P4333) and incubated in 30 μl of the same medium for 16 hours to allow for receptor up-regulation. 10 μl of diluted anti-CSF-1R antibodies were added to the cells for 1.5 h. Then cells were stimulated with 10 μl of 100 ng/ml huM-CSF-1 (Biomol Cat. No. 60530) for 5 min. After the incubation, supernatant was removed, cells were washed twice with 80 μl of ice-cold PBS and 50 μl of freshly prepared ice-cold lysis buffer (150 mM NaCl/20 mM Tris pH 7.5/1 mM EDTA/1 mM EGTA/1% Triton X-100/1 protease inhibitor tablet (Roche Diagnostics GmbH Cat. No. 1 836 170) per 10 ml buffer, 10 μl/ml phosphatase inhibitor cocktail 1 (Sigma Cat. No. P-2850, 100× Stock)/10 μl/ml protease inhibitor 1 (Sigma Cat. No. P-5726, 100× Stock)/10 μl/ml 1 M NaF) was added. After 30 minutes on ice the plates were shaken vigorously on a plateshaker for 3 minutes and then centrifuged 10 minutes at 2200 rpm (Heraeus Megafuge 10).

[0441] The presence of phosphorylated and total CSF-1 receptor in the cell lysate was analyzed with Elisa. For detection of the phosphorylated receptor the kit from R&D Systems (Cat. No. DYC3268-2) was used according to the instructions of the supplier. For detection of total CSF-1R 10 μl of the lysate was immobilized on plate by use of the capture antibody contained in the kit. Thereafter 1:750 diluted biotinylated anti CSF-1R antibody BAF329 (R&D Systems) and 1:1000 diluted streptavidin-HRP conjugate was added. After 60 minutes plates were developed with freshly prepared ABTS® solution and the absorbance was detected. Data were calculated as % of positive control without antibody and the ratio value phospho/total receptor expressed. The negative control was defined without addition of M-CSF-1. Anti CSF-1R SC 2-4A5 (Santa Cruz Biotechnology, US, see also Sherr, C. J. et al., Blood 73 (1989) 1786-1793), which inhibits the ligand-receptor interaction, was used as reference control.

TABLE 2

Calculated IC₅₀ values for the inhibition of CSF-1 receptor phosphorylation.

CSF-1R Mab	IC ₅₀ CSF-1R Phosphorylation [ng/ml]
Mab 2F11	219.4
Mab 2E10	752.0
Mab 2H7	703.4
Mab 1G10	56.6
SC-2-4A5	1006.6

Example 4

[0442] Determination of the Binding of Anti-CSF-1R Antibodies to Human CSF-1R Fragment delD4 and to Human CSF-1R Extracellular Domain (CSF-1R-ECD)

[0443] Preparation of Human CSF-1R Extracellular Domain (CSF-1R-ECD) (Comprising the Extracellular Subdomains D1-D5, hCSF-1R-ECD) of SEQ ID NO: 64:

[0444] pCMV-preS-Fc-hCSF-1R-ECD (7836 bp) encodes the complete ECD of human CSF-1R (SEQ ID NO: 64)

C-terminally fused to a PreScission protease cleavage site, followed by aa100-330 of human IgG1 and a 6×His-Tag, under the control of CMV promoter. The natural signal peptide has been varied by insertion of amino acids G and S after the first M, in order to create a BamHI restriction site.

[0445] Preparation of Human CSF-1R Fragment delD4 (Comprising the Extracellular Subdomains D1-D3 and D5, hCSF-1R-delD4) of SEQ ID NO: 65:

[0446] hCSF1R-delD4-V1-PreSc-hFc-His was cloned from pCMV-preS-Fc-hCSF-1R-ECD by means of the Stratagene QuikChange XL site-directed mutagenesis protocol, using delD4—for with sequence CACCTCCATGT-TCTTCCGGTACCCCCAGAGGTAAG (SEQ ID NO: 68) as the forward primer and delD4-rev with the reverse complement sequence as the reverse primer. A protocol variation published in *BioTechniques* 26 (1999) 680 was used to extend both primers in separate reactions in three cycles preceding the regular Stratagene protocol:

[0447] Two separate 50 μ l reaction mixtures were set up according to the manufacturer's manual, each containing 10 ng plasmid pCMV-preS-Fc-hCSF1R-ECD as the template and 10 μ M of one of the primers delD4—for or delD4-rev, and 0.5 μ l Pfu DNA polymerase as provided with the kit. Three PCR cycles 95° C. 30 sec/55° C. 60 sec/68° C. 8 min were run, then 25 μ l each of both reaction mixtures were combined in a new tube and 0.5 μ l fresh Pfu DNA polymerase were added. The regular PCR protocol with 18 temperature cycles as specified by Stratagene in the kit manual was carried out, followed by 2 hrs final digestion with the Dpn restriction enzyme provided with the kit. Clones bearing the deletion were detected by digestion with *Cel II* and *Not I* and verified by sequencing.

[0448] Protein was prepared by transient transfection in the Hek293 FreeStyle suspension cell system (Invitrogen) according to the manufacturer's specifications. After 1 week 500 ml supernatant was filtered and loaded onto a 1 ml HiTrap MabSelect Xtra (GE healthcare) protein A column (0.2 ml/min). The column was washed first with PBS, then with 50 mM Tris/150 mM NaCl/1 mM EDTA/pH 7.3. 75 μ l PreScission Protease (GE #27-0843-01) diluted in 375 μ l of the same buffer were loaded onto the column and the closed column was incubated over night at 4° C. with rolling. The column was mounted on top of a 1 ml GStrap FF column (GE healthcare) and the desired protein was eluted (0.2 ml/min, 0.2 ml fractions). Pooled fractions were concentrated from 1.8 ml to 0.4 ml by centrifugal ultrafiltration via a 3 k Nanosep and chromatographed over an S200 HR SEC in PBS (0.5 ml/min).

[0449] Human CSF-1R fragment delD4 was obtained in two fractions as a dimeric molecule (pool1, V=1.5 ml; c=0.30 mg/ml; apparent mass on SDS page 83 kDa, reduced 62 kDa) and as the monomer (pool 2, V=1.4 ml; c=0.25 mg/ml apparent mass on SDS page 62 kDa). The dimeric form was used for all experiments.

[0450] Determination of the Binding of Anti-CSF-1R Antibodies to Human CSF-1R Fragment delD4 and to Human CSF-1R Extracellular Domain (CSF-1R-ECD) (Binding Signals as Response Units (RU):

[0451] Instrument: Biacore T00 (GE Healthcare)

[0452] Software: T100 Control, Version 2.0.1

[0453] T100 Evaluation, Version 2.0.2

[0454] Assayformat Chip: CM5

[0455] Temperature: 25° C.

[0456] CSF-1R fragments were immobilized via amine coupling. To compare the binding of different anti-CSF-1R

antibodies according to the invention one concentration of the test antibody was injected. Anti CSF-1R Mab3291 (R&D-Systems) and SC 2-4A5 (Santa Cruz Biotechnology, US— see also Sherr, C. J. et al., *Blood* 73 (1989) 1786-1793), was used as reference control, anti-CCR5 m<CCR5>Pz03.1C5 (deposited as DSM ACC 2683 on 18 Aug. 2004 at DSMZ) as negative control, all under the same conditions as the anti-CSF-1R antibodies according to the invention.

[0457] Amine Coupling of CSF-1R Fragments

[0458] Standard amine coupling according to the manufacturer's instructions: running buffer: PBS-T (Roche: 11 666 789+0.05% Tween20: 11 332 465), activation by mixture of EDC/NHS, injection of human CSF-1R fragment delD4 (comprising the extracellular subdomains D1-D3 and D5) (SEQ ID NO: 65) and human CSF-1R Extracellular Domain (CSF-1R-ECD) (comprising the extracellular subdomains D1-D5) (SEQ ID NO: 64) for 600 seconds at flow rate 10 μ l/min; diluted in coupling buffer NaAc, pH 5.0, c=10 μ g/ml; finally remaining activated carboxyl groups were blocked by injection of 1 M Ethanolamin.

[0459] Binding of <CSF-1R>Mab 2F11, Mab 2E10, Mab 3291 and Sc2-4A5 and Other Anti-CSF-1R Antibodies to Human CSF-1R Fragment delD4 and Human CSF-1R Extracellular Domain (CSF-1R-ECD) at 25° C.

[0460] Running buffer: PBS-T (Roche: 11 666 789+0.05% Tween20: 11 332 465)

[0461] Analyte Sample:

[0462] Binding was measured at a flow rate of 30 μ l/min by one injection of the analyte with concentration c=10 nM. (for Mab 1G10, Mab 2H7 and humanized hMab 2F11-e7 in second experiment) Each injection was 700 seconds long, followed by a dissociation phase of 180 seconds. Final regeneration was performed after each cycle using 50 mM NaOH, contact time 60 seconds, flow rate 30 μ l/min.

[0463] Signals were measured by a report point 10 seconds after end of injection. Reference signals (signals from a blank reference flow cell (treated with EDC/NHS and ethanolamine, only) were subtracted to give the binding signals (as RU). If binding signals of nonbinding antibodies were slightly below 0 (Mab 2F11=-3; Mab 2E10=-2; Mab 1G10=-6, Mab 2H7=-9; and humanized hMab 2F11-e7=-7) the values were set as 0.

TABLE 3a

Binding of <CSF-1R> MAb to human CSF-1R fragment delD4 and CSF-1R-ECD and ratio at 25° C., measured by SPR			
	Binding to delD4 [RU]	Binding to CSF-1R-ECD [RU]	Ratio of binding of anti-CSF1R antibodies to CSF1R fragment delD4/ to CSF-1R-ECD
Mab 3291	1015	627	1015/627 = 1.61
sc2-4A5	374	249	374/249 = 1.50
Mab 2F11	0	176	0/176 = 0
hMab 2F11-e7	0	237	0/237 = 0
Mab 2E10	0	120	0/120 = 0
Mab 1G10	0	2708	0/2708 = 0
Mab 2H7	0	147	0/147 = 0
m<CCR5>Pz03.1C5	2	5	—

[0464] Mab 2F11 and Mab 2E10 showed binding to the human CSF-1R Extracellular Domain (CSF-1R-ECD) (see FIG. 2b); however no binding was detected to CSF-1R fragment delD4. (see FIG. 2a).

[0465] Sc2-4A5 and MAB3291 showed binding to CSF-1R-ECD and to del D4 (see FIGS. 2b and 2a).

[0466] Thus the ratio of binding of anti-CSF1R antibodies Mab 2F11 and Mab 2E10 to CSF1R fragment delD4/to CSF-1R-ECD was clearly below 1:50 ($=0.02$), while the binding ratio of MAB3291 and Sc2-4A5 were 1.61 and 1.50, respectively and were highly above 1:50 ($=0.02$). Negative control antibody m<CCR5>Pz03.1C5 did not show any binding (as expected).

[0467] Mab 1G10, Mab 2H7 and humanized hMab 2F11-e7 showed binding to the human CSF-1R Extracellular Domain (CSF-1R-ECD) (see FIG. 2d); however no binding was detected to CSF-1R fragment delD4. (see FIG. 2c). Thus the ratio of binding of anti-CSF1R antibodies Mab 1G10, Mab 2H7 and humanized hMab 2F11-e7 to CSF1R fragment delD4/to CSF-1R-ECD was clearly below 1:50 ($=0.02$).

[0468] In a further experiment anti-CSF-1R antibodies 1.2.SM (ligand displacing CSF-1R antibody described in WO2009026303), CXIIG6 (ligand displacing CSF-1R antibody described in WO 2009/112245), the goat polyclonal anti-CSF-1R antibody ab10676 (abcam) were investigated. Anti-CSF-1R antibody Mab3291 (R&D-Systems) was used as reference control. Anti-CCR5 m<CCR5>Pz03.1C5 (deposited as DSM ACC 2683 on 18 Aug. 2004 at DSMZ) was used as negative control.

TABLE 3b

Binding of <CSF-1R> MAbs to human CSF-1R fragment delD4 and CSF-1R-ECD and ratio at 25° C., measured by SPR			
	Binding to delD4 [RU]	Binding to CSF-1R-ECD [RU]	Ratio of binding of anti-CSF1R antibodies to CSF1R fragment delD4/to CSF-1R-ECD
MAB3291	1790	1222	1790/1222 = 1.47
1.2.SM	469	704	469/704 = 0.67
CXIIG6	1983	1356	1983/1356 = 1.46
ab10676	787	547	787/547 = 1.44
m<CCR5>Pz03.1C5	0	0	—

[0469] 1.2.SM, CXIIG6, ab10676 and MAB3291 showed binding to CSF-1R-ECD and to del D4 (see FIGS. 2f and 2e).

[0470] The binding ratio of 1.2.SM, CXIIG6, ab10676 and MAB3291 was highly above 1:50 ($=0.02$). Negative control antibody m<CCR5>Pz03.1C5 did not show any binding (as expected).

Example 5

[0471] Growth Inhibition of NIH3T3-CSF-1R Recombinant Cells in 3D Culture Under Treatment with Anti-CSF-1R Monoclonal Antibodies (CellTiterGlo-Assay)

[0472] NIH 3T3 cells, retrovirally infected with either an expression vector for full-length wildtype CSF-1R (SEQ ID NO: 62) or mutant CSF-1R L301S Y969F (SEQ ID NO: 63), were cultured in DMEM high glucose media (PAA, Pasching, Austria) supplemented with 2 mM L-glutamine, 2 mM sodium pyruvate and non-essential amino acids and 10% fetal bovine serum (Sigma, Taufkirchen, Germany) on poly-HEMA (poly(2-hydroxyethylmethacrylate)) (Polysciences, Warrington, Pa., USA) coated dishes to prevent adherence to the plastic surface. Cells are seeded in medium replacing serum with 5 ng/ml sodium selenite, 10 ng/ml transferrin,

400 g/ml BSA and 0.05 mM 2-mercaptoethanol. When treated with 100 ng/ml huCSF-1 (Biomol, Hamburg, Germany) wtCSF-1R (expressing cells form dense spheroids that grow three dimensionally, a property that is called anchorage independence. These spheroids resemble closely the three dimensional architecture and organization of solid tumors in situ. Mutant CSF-1R recombinant cells are able to form spheroids independent of the CSF-1 ligand. Spheroid cultures were incubated for 3 days in the presence of different concentrations of antibody in order to determine an IC50 (concentration with 50 percent inhibition of cell viability). The CellTiterGlo assay was used to detect cell viability by measuring the ATP-content of the cells.

TABLE 5a

CSF-1R Mab	wtCSF-1R IC ₅₀ [µg/ml]	Mutant CSF-1R IC ₅₀ [µg/ml]
Mab 2F11	1.1	8.0
Mab 2E10	0.49	4.9
Mab 2H7	0.31	5.3
Mab 1G10	0.29	14.2
SC 2-4A5	10.0	10.0

[0473] Reference control Mab R&D-Systems 3291 did not show inhibition of mutant CSF-1R recombinant cell proliferation.

[0474] In a further experiment the anti-CSF-1R antibody according to the invention hMab 2F11-e7 and the anti-CSF-1R antibodies 1.2.SM (ligand displacing CSF-1R antibody described in WO2009026303), CXIIG6 (ligand displacing CSF-1R antibody described in WO 2009/112245), the goat polyclonal anti-CSF-1R antibody ab10676 (abcam), and SC 2-4A5 (Santa Cruz Biotechnology, US— see also Sherr, C. J. et al., Blood 73 (1989) 1786-1793) were investigated.

[0475] Spheroid cultures were incubated for 3 days in the presence of different concentrations of antibody in order to determine an IC30 (concentration with 30 percent inhibition of cell viability). Maximum concentration was 20 µg/ml The CellTiterGlo assay was used to detect cell viability by measuring the ATP-content of the cells.

TABLE 5b

CSF-1R Mab	wtCSF-1R IC ₃₀ [µg/ml]	Mutant CSF-1R IC ₃₀ [µg/ml]
hMab 2F11-e7	4.91	0.54
1.2.SM	1.19	>20 µg/ml (-19% inhibition at 20 µg/ml = 19% stimulation)
CXIIG6	>20 µg/ml (21% inhibition at 20 µg/ml)	>20 µg/ml (-36% inhibition at 20 µg/ml = 36% stimulation)
ab10676	14.15	>20 µg/ml (0% inhibition at 20 µg/ml)
SC 2-4A5	16.62	2.56

Example 6

[0476] Growth Inhibition of BeWo Tumor Cells in 3D Culture Under Treatment with Anti-CSF-1R Monoclonal Antibodies (CellTiterGlo-Assay)

[0477] BeWo choriocarcinoma cells (ATCC CCL-98) were cultured in F12K media (Sigma, Steinheim, Germany) supplemented with 10% FBS (Sigma) and 2 mM L-glutamine. 5×10^4 cells/well were seeded in 96-well poly-HEMA

(poly(2-hydroxyethylmethacrylate)) coated plates containing F12K medium supplemented with 0.5% FBS and 5% BSA. Concomitantly, 200 ng/ml huCSF-1 and 10 µg/ml of different anti-CSF-1R monoclonal antibodies were added and incubated for 6 days. The CellTiterGlo assay was used to detect cell viability by measuring the ATP-content of the cells in relative light units (RLU). When BeWo spheroid cultures were treated with different anti-CSF-1R antibodies (10 µg/ml) inhibition of CSF-1 induced growth was observed. To calculate antibody-mediated inhibition the mean RLU value of unstimulated BeWo cells was subtracted from all samples. Mean RLU value of CSF-1 stimulated cells was set arbitrarily to 100%. Mean RLU values of cells stimulated with CSF-1 and treated with anti-CSF-1R antibodies were calculated in % of CSF-1 stimulated RLU. The Table 6 shows the calculated data of growth inhibition of BeWo tumor cells in 3D culture under treatment with anti-CSF-1R monoclonal antibodies; FIGS. 1a and b depicts normalized mean RLU values.

TABLE 6

CSF-1R Mab	% inhibition 10 µg/ml antibody concentration
CSF-1 only	0
Mab 2F11	70
Mab 2E10	102
Mab 2H7	103
Mab 1G10	99
SC 2-4A5	39

Example 7

[0478] Inhibition of Human Macrophage Differentiation Under Treatment with Anti-CSF-1R Monoclonal Antibodies (CellTiterGlo-Assay)

[0479] Human monocytes were isolated from peripheral blood using the RosetteSep™ Human Monocyte Enrichment Cocktail (StemCell Tech. —Cat. No. 15028). Enriched monocyte populations were seeded into 96 well microtiterplates (2.5×10⁴ cells/well) in 100 µl RPMI 1640 (Gibco—Cat. No. 31870) supplemented with 10% FCS (GIBCO—Cat. No. 011-090014M), 4 mM L-glutamine (GIBCO—Cat. No. 25030) and 1× PenStrep (Roche Cat. No. 1 074 440) at 37° C. and 5% CO₂ in a humidified atmosphere. When 150 ng/ml huCSF-1 was added to the medium, a clear differentiation into adherent macrophages could be observed. This differentiation could be inhibited by addition of anti-CSF-1R antibodies. Furthermore, the monocyte survival is affected and could be analyzed by CellTiterGlo (CTG) analysis. From the concentration dependent inhibition of the survival of monocytes by antibody treatment, an IC₅₀ was calculated (see Table 7).

TABLE 7

CSF-1R Mab	IC ₅₀ [µg/ml]
Mab 2F11	0.08
Mab 2E10	0.06
Mab 2H7	0.03
Mab 1G10	0.06
SC 2-4A5	0.36

[0480] In a separate test series humanized versions of Mab 2 F11, e.g. hMab 2F11-c11, hMab 2F11-d8, hMab 2F11-e7,

hMab 2F11-f12, showed TC50 values of 0.07 µg/ml (hMab 2F11-c11), 0.07 µg/ml (hMab 2F11-d8), 0.04 g g/ml (hMab 2F11-e7) and 0.09 µg/ml (hMab 2F11-f12).

Example 8

[0481] Inhibition of Cynomolgous Macrophage Differentiation Under Treatment with Anti-CSF-1R Monoclonal Antibodies (CellTiterGlo-Assay)

[0482] Cynomolgous monocytes were isolated from peripheral blood using the CD14 MicroBeads non-human primate kit (Miltenyi Biotec—Cat. No. 130-091-097) according to the manufacturers description. Enriched monocyte populations were seeded into 96 well microtiterplates (1-3×10³ cells/well) in 100 µl RPMI 1640 (Gibco—Cat. No. 31870) supplemented with 10% FCS (GIBCO—Cat. No. 011-090014M), 4 mM L-glutamine (GIBCO—Cat. No. 25030) and 1× PenStrep (Roche Cat. No. 1 074 440) at 37° C. and 5% CO₂ in a humidified atmosphere. When 150 ng/ml huCSF-1 was added to the medium, a clear differentiation into adherent macrophages could be observed. This differentiation could be inhibited by addition of anti-CSF-1R antibodies. Furthermore, the monocyte survival is affected and could be analyzed by CellTiterGlo (CTG) analysis. The viability was analyzed at a concentration of 5 µg/ml antibody treatment (see Table 8).

TABLE 8

CSF-1R Mab	% survival	% inhibition (of survival) = (100% - % survival)
Mab 2F11	4*	96
Mab 2E10	17**	83
Mab 2H7	8	92
Mab 1G10	2	98
SC 2-4A5	31	69

*mean of four experiments (3 expts. using the murine, 1 expt. using the chimeric mAb)
**mean of two experiments using the murine mAb only

Example 9

[0483] Determination of the Binding Affinity of Anti-CSF-1R Antibodies to Human CSF-1R

[0484] Instrument: BIACORE® A100

[0485] Chip: CM5 (Biacore BR-1006-68)

[0486] Coupling: amine coupling

[0487] Buffer: PBS (Biacore BR-1006-72), pH 7.4, 35° C.

[0488] For affinity measurements 36 µg/ml anti mouse Fcγ antibodies (from goat, Jackson Immuno Research JIR115-005-071) have been coupled to the chip surface for capturing the antibodies against CSF-1R. Human CSF-1R Extracellular Domain (CSF-1R-ECD) (comprising the extracellular subdomains D1-D5) (SEQ ID NO: 64) (R&D-Systems 329-MR or subcloned pCMV-presS-HisAvitag-hCSF-1R-ECD) was added in various concentrations in solution. Association was measured by an CSF-1R-injection of 1.5 minutes at 35° C.; dissociation was measured by washing the chip surface with buffer for 10 minutes at 35° C. For calculation of kinetic parameters the Langmuir 1:1 model was used.

TABLE 9

Affinity data measured by SPR				
CSF-1R Mab	K_D (nM)	k_a (1/Ms)	k_d (1/s)	$t_{1/2}$ (min)
Mab 2F11	0.29	1.77E+05	5.18E-05	223
Mab 2E10	0.2	1.52E+05	2.97E-05	389
Mab 2H7	0.21	1.47E+05	3.12E-05	370
Mab 1G10	0.36	1.75E+05	6.28E-05	184

[0489] In a separate biacore binding assay using the CSF-1R ECD (data not shown) some competition of the antibodies Mab 2F11 and Mab 2E10 with the antibody Ab SC-2-4A5 was shown. However Mab 2F11/Mab 2E10 do not bind to the human CSF-1R fragment delD4, whereas Ab SC-2-4A5 binds to this delD4 fragment (see Example 4 and FIG. 2a). Thus the binding region of Mab 2F11/Mab 2E10 is clearly distinct from the binding region of Ab SC-2-4A5, but probably located in a vicinity area. In such competition assay both antibodies Mab 2F11 and Mab 2E10 did not compete with Mab3291 from R&D-Systems (data not shown).

Example 10

[0490] Determination of the Binding of Anti-CSF-1R Antibodies to Human CSF-1R Fragment D1-D3

[0491] Instrument: Biacore T100 (GE Healthcare)

[0492] Software: T100 Control, Version 1.1.11

[0493] B3000 Evaluation, Version 4.01

[0494] Scrubber, Version 2.0a

[0495] Assayformat Chip: CM5-Chip

[0496] Antibodies against CSF-1R were captured via amine coupled capture molecules. Using the single cycle kinetics five increasing concentrations of human CSF-1R fragment D1-D3 (SEQ ID NO: 66) were injected. Human CSF-1R fragment D1-D3 was subcloned into pCMV-presS-HisAvitag expression vector.

[0497] Anti CSF-1R SC 2-4A5 (Santa Cruz Biotechnology, US; Sherr. C. J. et al., Blood 73 (1989) 1786-1793) which inhibits the ligand-receptor interaction, and Mab 3291 (R&D-Systems) were used as reference controls.

[0498] Capture molecules: Anti mouse Fc γ antibodies (from goat, Jackson Immuno Reasearch JIR115-005-071) for antibodies according to the invention and the R&D-Systems control Mab 3291 and Anti rat Fc γ antibodies (from goat, Jackson Immuno Reasearch JIR112-005-071) for the reference control anti CSF-1R SC 2-4A5.

[0499] Amine Coupling of Capture Molecules

[0500] Standard amine coupling according to the manufacturer's instructions: running buffer: HBS-N buffer, activation by mixture of EDC/NHS, aim for ligand density of 2000 RU; the capture-Abs were diluted in coupling buffer NaAc, pH 4.5, c=10 g/mL; finally remaining activated carboxyl groups were blocked by injection of 1 M Ethanolamin.

[0501] Kinetic Characterization of Human CSF-1R Fragments D1-D3 Binding to MAbs <CSF-1R> at 37° C.

[0502] Running buffer: PBS (Biacore BR-1006-72)

[0503] Capturing of Mabs <CSF-1R> on flow cells 2 to 4: Flow 20 μ L/min, contact time 90 seconds, c(Abs<CSF-1R>) =50 nM, diluted with running buffer+1 mg/mL BSA;

[0504] Analyte Sample:

[0505] Single Cycle Kinetics was measured at a flow rate of 30 μ L/min by five consecutive injections of the analyte

with concentrations, c=7.8, 31.25, 125 500 and 2000 nM, without regeneration. Each injection was 30 seconds long and followed by a dissociation phase of 120 Seconds for the first four injections, and finally 1200 seconds for the highest concentration (=last injection).

[0506] Final regeneration was performed after each cycle using 10 mM Glycin pH 1.5 (Biacore BR-1003-54), contact time 60 seconds, flow rate 30 μ L/min.

[0507] Kinetic parameters were calculated by using the usual double referencing (control reference: binding of analyte to capture molecule; Flow Cell: subdomain CSF-1R concentration "0" as Blank) and calculation with model 'titration kinetics 1:1 binding with draft'.

TABLE 10

Affinity data for binding of human CSF-1R fragment D1-D3 measured by SPR					
CSF-1R Mab	Sub domain	K_D (nM)	k_a (1/Ms)	k_d (1/s)	$t_{1/2}$ (min)
Mab 2F11	D1-D3	no binding			
Mab 2E10	D1-D3	no binding			
Mab 2H7	D1-D3	not determined			
Mab 1G10	D1-D3	no binding			
SC-2-4A5	D1-D3	no binding			
R&D-Systems 3291	D1-D3	5.4	2.2E+5	1.2E-3	9.6

[0508] The antibodies Mab 2F11, Mab 2E10 and Mab 1G10 showed no binding to human CSF-1R fragment D1-D3

[0509] Also reference control-Ab SC-2-4A5 did not bind to human CSF-1R fragment D1-D3.

[0510] The reference control Mab R&D-Systems 3291 showed binding to the human CSF-1R fragment D1-D3.

Example 11

[0511] CSF-1 level increase during CSF-1R inhibition in Cynomolgus monkey Serum CSF-1 levels provide a pharmacodynamic marker of CSF-1R neutralizing activity of anti-human CSF-1R dimerization inhibitor hMab 2F11-e7. One male and one female cynomolgus monkey per dosage group (1 and 10 mg/kg) were intravenously administered anti-CSF1R antibody hMab 2F11-e7. Blood samples for analysis of CSF-1 levels were collected 1 week before treatment (pre-dose), 2, 24, 48, 72, 96, 168 hours post-dose and weekly for two additional weeks. CSF-1 levels were determined using a commercially available ELISA kit (Quantikine® human M-CSF) according to the manufacturer's instructions (R&D Systems, UK). Monkey CSF-1 level were determined by comparison with CSF-1 standard curve samples provided in the kit.

[0512] Administration of hMab 2F11-e7 induced a dramatic increase in CSF-1 by ~1000-fold, which depending on the dose administered lasted for 48 hr (1 mg/kg) or 15 days (10 mg/kg). Hence, a dimerization inhibitor for CSF-1R offers the advantage to not directly compete with the dramatically upregulated ligand for binding to the receptor in contrast to a ligand displacing antibody.

Example 12

[0513] In Vivo Efficacy—Tumor Growth Inhibition of Anti-CSF-1R Antibodies in Breast Cancer BT20 Xenograft Tumor Cells in SCID Beige Mice

[0514] The human breast cancer cell line BT-20 expresses human CSF-1R but lacks CSF-1 expression (Sapi, E. et al Cancer Res 59 (1999) 5578-5585). Since the mouse derived CSF-1 fails to activate human CSF-1R on the tumor cells recombinant human CSF-1 (Biomol, Hamburg, Germany) was supplemented via osmotic minipumps (ALZET, Cupertino, Calif.) providing a continuous CSF-1 infusion rate of 2 µg/day (Martin, T. A., Carcinogenesis 24 (2003) 1317-1323).

[0515] To directly compare the efficacy of an antibody interfering with dimerization of CSF-1R with a ligand displacing CSF-1R antibody we tested the chimeric anti-CSF-1R Mab 2F11 (antibody interfering with dimerization of CSF-1R) and 1.2.SM (ligand displacing CSF-1R antibody described in WO2009026303) in the BT-20 xenograft model.

[0516] SCID beige mice (Charles River, Sulzfeld, Germany) were subcutaneously coinjected with 1×10⁷ cells BT-20 cells (ATCC HTB-19) and 100l of Matrigel. Treatment of animals started at day of randomization at a mean

tumor volume of 100 mm³. Mice are treated once weekly i.p. with the respective antibodies (see FIG. 4) in 20 mM Histidine, 140 mM NaCl pH 6.0 buffer. The tumor dimensions are measured by caliper beginning on the staging day and subsequently 2 times per week during the whole treatment period. Tumor volume is calculated according to NCI protocol (Tumor weight= $\frac{1}{2}ab^2$, where “a” and “b” are the long and the short diameters of the tumor, respectively).

[0517] Tumor growth analysis is shown in FIG. 4. Inhibition of human CSF-1R on tumor cells with the chimeric anti-CSF-1R Mab 2F11 was statistically more efficacious in mediating tumor growth inhibition than anti-CSF-1R antibody 1.2.SM (CSF-1R antibody described in WO2009026303).

[0518] Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, the descriptions and examples should not be construed as limiting the scope of the invention. The disclosures of all patents, patent applications, scientific references, and Genbank Accession Nos. cited herein are expressly incorporated by reference in their entirety for all purposes as if each patent, patent application, scientific reference, and Genbank Accession No. were specifically and individually incorporated by reference.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 84

<210> SEQ ID NO 1

<211> LENGTH: 8

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 1

Asp Gln Arg Leu Tyr Phe Asp Val
1 5

<210> SEQ ID NO 2

<211> LENGTH: 16

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 2

Val Ile Trp Thr Asp Gly Gly Thr Asn Tyr Asn Ser Pro Phe Met Ser
1 5 10 15

<210> SEQ ID NO 3

<211> LENGTH: 5

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 3

Thr Tyr Asp Ile Ser
1 5

<210> SEQ ID NO 4

<211> LENGTH: 8

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 4

Gly Gln Ser Phe Ser Tyr Pro Thr

-continued

1 5

<210> SEQ ID NO 5
 <211> LENGTH: 7
 <212> TYPE: PRT
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 5

Gly Ala Ser Asn Arg Tyr Thr
 1 5

<210> SEQ ID NO 6
 <211> LENGTH: 11
 <212> TYPE: PRT
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 6

Lys Ala Ser Glu Asp Val Asn Thr Tyr Val Ser
 1 5 10

<210> SEQ ID NO 7
 <211> LENGTH: 116
 <212> TYPE: PRT
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 7

Gln Val Gln Leu Lys Glu Ser Gly Pro Gly Leu Val Ala Pro Ser Gln
 1 5 10 15

Ser Leu Ser Ile Thr Cys Thr Val Ser Gly Phe Ser Leu Thr Thr Tyr
 20 25 30

Asp Ile Ser Trp Ile Arg Gln Ser Pro Gly Lys Gly Leu Glu Trp Leu
 35 40 45

Gly Val Ile Trp Thr Asp Gly Gly Thr Asn Tyr Asn Ser Pro Phe Met
 50 55 60

Ser Arg Leu Ser Ile Arg Lys Asp Asn Ser Lys Ser Gln Val Phe Leu
 65 70 75 80

Lys Met Asn Arg Leu Gln Thr Asp Asp Thr Ala Ile Tyr Tyr Cys Val
 85 90 95

Arg Asp Gln Arg Leu Tyr Phe Asp Val Trp Gly Ala Gly Thr Thr Val
 100 105 110

Thr Val Ser Ser
 115

<210> SEQ ID NO 8
 <211> LENGTH: 106
 <212> TYPE: PRT
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 8

Asn Ile Val Met Thr Gln Ser Pro Lys Ser Met Ser Met Ser Val Gly
 1 5 10 15

Glu Arg Val Thr Leu Asn Cys Lys Ala Ser Glu Asp Val Asn Thr Tyr
 20 25 30

Val Ser Trp Tyr Gln Gln Gln Pro Glu Gln Ser Pro Lys Leu Leu Ile
 35 40 45

Tyr Gly Ala Ser Asn Arg Tyr Thr Gly Val Pro Asp Arg Phe Thr Gly
 50 55 60

-continued

Gly Gly Ser Thr Thr Asp Phe Thr Leu Thr Ile Ser Ser Val Gln Ala
65 70 75 80

Glu Asp Leu Ala Asp Tyr Phe Cys Gly Gln Ser Phe Ser Tyr Pro Thr
85 90 95

Phe Gly Thr Gly Thr Lys Leu Glu Ile Lys
100 105

<210> SEQ ID NO 9
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 9

Asp Pro Arg Leu Tyr Phe Asp
1 5

<210> SEQ ID NO 10
<211> LENGTH: 16
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 10

Val Ile Trp Thr Gly Gly Gly Thr Asn Tyr Asn Ser Gly Phe Met Ser
1 5 10 15

<210> SEQ ID NO 11
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 11

Ser Phe Asp Ile Ser
1 5

<210> SEQ ID NO 12
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 12

Gly Gln Thr Phe Ser Tyr Pro Thr
1 5

<210> SEQ ID NO 13
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 13

Gly Ala Ser Asn Arg Tyr Thr
1 5

<210> SEQ ID NO 14
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 14

Lys Ala Ser Glu Asp Val Val Thr Tyr Val Ser
1 5 10

-continued

<210> SEQ ID NO 15
 <211> LENGTH: 116
 <212> TYPE: PRT
 <213> ORGANISM: Mus musculus

 <400> SEQUENCE: 15

 Gln Val Gln Leu Lys Glu Ser Gly Pro Gly Leu Val Ala Pro Ser Lys
 1 5 10 15

 Ser Leu Ser Ile Thr Cys Thr Val Ser Gly Ser Ser Leu Asp Ser Phe
 20 25 30

 Asp Ile Ser Trp Ile Arg Gln Ser Pro Gly Lys Gly Leu Glu Trp Leu
 35 40 45

 Gly Val Ile Trp Thr Gly Gly Gly Thr Asn Tyr Asn Ser Gly Phe Met
 50 55 60

 Ser Arg Leu Arg Ile Thr Lys Asp Asn Ser Lys Ser Gln Val Leu Leu
 65 70 75 80

 Lys Met Asn Ser Leu Gln Ser Asp Asp Thr Ala Ile Tyr Tyr Cys Val
 85 90 95

 Arg Asp Pro Arg Leu Tyr Phe Asp Val Trp Gly Ala Gly Thr Thr Val
 100 105 110

 Thr Val Ser Ser
 115

<210> SEQ ID NO 16
 <211> LENGTH: 106
 <212> TYPE: PRT
 <213> ORGANISM: Mus musculus

 <400> SEQUENCE: 16

 Asn Ile Val Met Thr Gln Ser Pro Lys Ser Met Ser Met Ser Val Gly
 1 5 10 15

 Glu Arg Val Thr Leu Ser Cys Lys Ala Ser Glu Asp Val Val Thr Tyr
 20 25 30

 Val Ser Trp Tyr Gln Gln Lys Pro Asp Gln Ser Pro Lys Leu Leu Ile
 35 40 45

 Tyr Gly Ala Ser Asn Arg Tyr Thr Gly Val Pro Asp Arg Phe Thr Gly
 50 55 60

 Ser Gly Ser Ala Thr Asp Phe Thr Leu Thr Ile Ser Ser Val Gln Ala
 65 70 75 80

 Glu Asp Leu Ala Asp Tyr Tyr Cys Gly Gln Thr Phe Ser Tyr Pro Thr
 85 90 95

 Phe Gly Thr Gly Thr Lys Leu Glu Ile Lys
 100 105

<210> SEQ ID NO 17
 <211> LENGTH: 8
 <212> TYPE: PRT
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: heavy chain CDR3, hMab 2F11-c11

 <400> SEQUENCE: 17

Asp Gln Arg Leu Tyr Phe Asp Val
 1 5

<210> SEQ ID NO 18
 <211> LENGTH: 16

-continued

<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: heavy chain CDR2, hMab 2F11-c11

<400> SEQUENCE: 18

Val Ile Trp Thr Asp Gly Gly Thr Asn Tyr Asn Ser Pro Phe Met Ser
1 5 10 15

<210> SEQ ID NO 19
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: heavy chain CDR1, hMab 2F11-c11

<400> SEQUENCE: 19

Thr Tyr Asp Ile Ser
1 5

<210> SEQ ID NO 20
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: light chain CDR3, hMab 2F11-c11

<400> SEQUENCE: 20

Gly Gln Ser Phe Ser Tyr Pro Thr
1 5

<210> SEQ ID NO 21
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: light chain CDR2, hMab 2F11-c11

<400> SEQUENCE: 21

Gly Ala Ser Asn Arg Tyr Thr
1 5

<210> SEQ ID NO 22
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: light chain CDR1, hMab 2F11-c11

<400> SEQUENCE: 22

Arg Ala Ser Glu Asp Val Asn Thr Tyr Val Ser
1 5 10

<210> SEQ ID NO 23
<211> LENGTH: 116
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: heavy chain variable domain, hMab 2F11-c11

<400> SEQUENCE: 23

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ser
1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Phe Ser Leu Thr Thr Tyr

-continued

```

                20                25                30
Asp Ile Ser Trp Ile Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
   35                40                45
Gly Val Ile Trp Thr Asp Gly Gly Thr Asn Tyr Asn Ser Pro Phe Met
   50                55                60
Ser Arg Val Thr Ile Thr Lys Asp Glu Ser Thr Ser Thr Ala Tyr Met
   65                70                75                80
Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys Val
   85                90                95
Arg Asp Gln Arg Leu Tyr Phe Asp Val Trp Gly Gln Gly Thr Thr Val
   100                105                110
Thr Val Ser Ser
   115

```

```

<210> SEQ ID NO 24
<211> LENGTH: 106
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: light chain variable domain, hMab 2F11-c11

```

<400> SEQUENCE: 24

```

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1                5                10                15
Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Glu Asp Val Asn Thr Tyr
20                25                30
Val Ser Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile
35                40                45
Tyr Gly Ala Ser Asn Arg Tyr Thr Gly Val Pro Ser Arg Phe Ser Gly
50                55                60
Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
65                70                75                80
Glu Asp Phe Ala Thr Tyr Tyr Cys Gly Gln Ser Phe Ser Tyr Pro Thr
85                90                95
Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
100                105

```

```

<210> SEQ ID NO 25
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: heavy chain CDR3, hMab 2F11-d8

```

<400> SEQUENCE: 25

```

Asp Gln Arg Leu Tyr Phe Asp Val
1                5

```

```

<210> SEQ ID NO 26
<211> LENGTH: 16
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: heavy chain CDR2, hMab 2F11-d8

```

<400> SEQUENCE: 26

```

Val Ile Trp Thr Asp Gly Gly Ala Asn Tyr Ala Gln Lys Phe Gln Gly
1                5                10                15

```

-continued

<210> SEQ ID NO 27
 <211> LENGTH: 5
 <212> TYPE: PRT
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: heavy chain CDR1, hMab 2F11-d8

<400> SEQUENCE: 27

Thr Tyr Asp Ile Ser
 1 5

<210> SEQ ID NO 28
 <211> LENGTH: 8
 <212> TYPE: PRT
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: light chain CDR3, hMab 2F11-d8

<400> SEQUENCE: 28

Gly Gln Ser Phe Ser Tyr Pro Thr
 1 5

<210> SEQ ID NO 29
 <211> LENGTH: 7
 <212> TYPE: PRT
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: light chain CDR2, hMab 2F11-d8

<400> SEQUENCE: 29

Gly Ala Ser Asn Arg Tyr Thr
 1 5

<210> SEQ ID NO 30
 <211> LENGTH: 11
 <212> TYPE: PRT
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: light chain CDR1, hMab 2F11-d8

<400> SEQUENCE: 30

Lys Ala Ser Glu Asp Val Asn Thr Tyr Val Ser
 1 5 10

<210> SEQ ID NO 31
 <211> LENGTH: 116
 <212> TYPE: PRT
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: heavy chain variable domain, hMab 2F11-d8

<400> SEQUENCE: 31

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ser
 1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Phe Ser Leu Thr Thr Tyr
 20 25 30

Asp Ile Ser Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
 35 40 45

Gly Val Ile Trp Thr Asp Gly Gly Ala Asn Tyr Ala Gln Lys Phe Gln
 50 55 60

Gly Arg Val Thr Ile Thr Ala Asp Glu Ser Thr Ser Thr Ala Tyr Met

-continued

```

65              70              75              80
Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys Ala
      85              90              95
Arg Asp Gln Arg Leu Tyr Phe Asp Val Trp Gly Gln Gly Thr Thr Val
      100              105              110
Thr Val Ser Ser
      115

```

```

<210> SEQ ID NO 32
<211> LENGTH: 106
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: light chain variable domain, hMab 2F11-d8

```

```

<400> SEQUENCE: 32

```

```

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1              5              10              15
Asp Arg Val Thr Ile Thr Cys Lys Ala Ser Glu Asp Val Asn Thr Tyr
      20              25              30
Val Ser Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile
      35              40              45
Tyr Gly Ala Ser Asn Arg Tyr Thr Gly Val Pro Ser Arg Phe Ser Gly
      50              55              60
Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
      65              70              75              80
Glu Asp Phe Ala Thr Tyr Tyr Cys Gly Gln Ser Phe Ser Tyr Pro Thr
      85              90              95
Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
      100              105

```

```

<210> SEQ ID NO 33
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: heavy chain CDR3, hMab 2F11-e7

```

```

<400> SEQUENCE: 33

```

```

Asp Gln Arg Leu Tyr Phe Asp Val
1              5

```

```

<210> SEQ ID NO 34
<211> LENGTH: 16
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: heavy chain CDR2, hMab 2F11-e7

```

```

<400> SEQUENCE: 34

```

```

Val Ile Trp Thr Asp Gly Gly Thr Asn Tyr Ala Gln Lys Leu Gln Gly
1              5              10              15

```

```

<210> SEQ ID NO 35
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: heavy chain CDR1, hMab 2F11-e7

```

-continued

<400> SEQUENCE: 35

Ser Tyr Asp Ile Ser
 1 5

<210> SEQ ID NO 36

<211> LENGTH: 8

<212> TYPE: PRT

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: light chain CDR3, hMab 2F11-e7

<400> SEQUENCE: 36

Gln Gln Ser Phe Ser Tyr Pro Thr
 1 5

<210> SEQ ID NO 37

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: light chain CDR2, hMab 2F11-e7

<400> SEQUENCE: 37

Ala Ala Ser Asn Arg Tyr Thr
 1 5

<210> SEQ ID NO 38

<211> LENGTH: 11

<212> TYPE: PRT

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: light chain CDR1, hMab 2F11-e7

<400> SEQUENCE: 38

Arg Ala Ser Glu Asp Val Asn Thr Tyr Val Ser
 1 5 10

<210> SEQ ID NO 39

<211> LENGTH: 116

<212> TYPE: PRT

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: heavy chain variable domain, hMab 2F11-e7

<400> SEQUENCE: 39

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
 1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Ser Tyr
 20 25 30

Asp Ile Ser Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
 35 40 45

Gly Val Ile Trp Thr Asp Gly Gly Thr Asn Tyr Ala Gln Lys Leu Gln
 50 55 60

Gly Arg Val Thr Met Thr Thr Asp Thr Ser Thr Ser Thr Ala Tyr Met
 65 70 75 80

Glu Leu Arg Ser Leu Arg Ser Asp Asp Thr Ala Val Tyr Tyr Cys Ala
 85 90 95

Arg Asp Gln Arg Leu Tyr Phe Asp Val Trp Gly Gln Gly Thr Thr Val
 100 105 110

Thr Val Ser Ser

-continued

115

<210> SEQ ID NO 40
 <211> LENGTH: 106
 <212> TYPE: PRT
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: light chain variable domain, hMab 2F11-e7

<400> SEQUENCE: 40

```

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1           5           10           15
Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Glu Asp Val Asn Thr Tyr
          20           25           30
Val Ser Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile
          35           40           45
Tyr Ala Ala Ser Asn Arg Tyr Thr Gly Val Pro Ser Arg Phe Ser Gly
50           55           60
Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
65           70           75           80
Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Ser Phe Ser Tyr Pro Thr
          85           90           95
Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
          100           105

```

<210> SEQ ID NO 41
 <211> LENGTH: 8
 <212> TYPE: PRT
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: heavy chain CDR3, hMab 2F11-f12

<400> SEQUENCE: 41

```

Asp Gln Arg Leu Tyr Phe Asp Val
1           5

```

<210> SEQ ID NO 42
 <211> LENGTH: 16
 <212> TYPE: PRT
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: heavy chain CDR2, hMab 2F11-f12

<400> SEQUENCE: 42

```

Val Ile Trp Thr Asp Gly Gly Thr Asn Tyr Asn Ser Pro Phe Met Ser
1           5           10           15

```

<210> SEQ ID NO 43
 <211> LENGTH: 5
 <212> TYPE: PRT
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: heavy chain CDR1, hMab 2F11-f12

<400> SEQUENCE: 43

```

Thr Tyr Asp Ile Ser
1           5

```

<210> SEQ ID NO 44
 <211> LENGTH: 8
 <212> TYPE: PRT

-continued

```

<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: light chain CDR3, hMab 2F11-f12

<400> SEQUENCE: 44
Gly Gln Ser Phe Ser Tyr Pro Thr
1                5

<210> SEQ ID NO 45
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: light chain CDR2, hMab 2F11-f12

<400> SEQUENCE: 45
Gly Ala Ser Ser Leu Gln Ser
1                5

<210> SEQ ID NO 46
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: light chain CDR1, hMab 2F11-f12

<400> SEQUENCE: 46
Arg Ala Ser Glu Asp Val Asn Thr Tyr Val Ser
1                5                10

<210> SEQ ID NO 47
<211> LENGTH: 116
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: heavy chain variable domain, hMab 2F11-f12

<400> SEQUENCE: 47
Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ser
1                5                10                15
Ser Val Lys Val Ser Cys Lys Ala Ser Gly Phe Ser Leu Thr Thr Tyr
20                25                30
Asp Ile Ser Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
35                40                45
Gly Val Ile Trp Thr Asp Gly Gly Thr Asn Tyr Asn Ser Pro Phe Met
50                55                60
Ser Arg Val Thr Ile Thr Lys Asp Glu Ser Thr Ser Thr Ala Tyr Met
65                70                75                80
Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys Val
85                90                95
Arg Asp Gln Arg Leu Tyr Phe Asp Val Trp Gly Gln Gly Thr Thr Val
100                105                110

Thr Val Ser Ser
115

<210> SEQ ID NO 48
<211> LENGTH: 106
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: light chain variable domain, hMab 2F11-f12

```

-continued

<400> SEQUENCE: 48

```

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1           5           10           15
Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Glu Asp Val Asn Thr Tyr
          20           25           30
Val Ser Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile
          35           40           45
Tyr Gly Ala Ser Ser Leu Gln Ser Gly Val Pro Ser Arg Phe Ser Gly
          50           55           60
Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
65           70           75           80
Glu Asp Phe Ala Thr Tyr Tyr Cys Gly Gln Ser Phe Ser Tyr Pro Thr
          85           90           95
Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
          100           105

```

<210> SEQ ID NO 49

<211> LENGTH: 8

<212> TYPE: PRT

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: heavy chain CDR3, hMab 2F11-g1

<400> SEQUENCE: 49

```

Asp Gln Arg Leu Tyr Phe Asp Val
1           5

```

<210> SEQ ID NO 50

<211> LENGTH: 16

<212> TYPE: PRT

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: heavy chain CDR2, hMab 2F11-g1

<400> SEQUENCE: 50

```

Val Ile Trp Thr Asp Gly Gly Thr Asn Tyr Asn Ser Pro Leu Lys Ser
1           5           10           15

```

<210> SEQ ID NO 51

<211> LENGTH: 5

<212> TYPE: PRT

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: heavy chain CDR1, hMab 2F11-g1

<400> SEQUENCE: 51

```

Thr Tyr Asp Ile Ser
1           5

```

<210> SEQ ID NO 52

<211> LENGTH: 8

<212> TYPE: PRT

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: light chain CDR3, hMab 2F11-g1

<400> SEQUENCE: 52

```

Gly Gln Ser Phe Ser Tyr Pro Thr
1           5

```


-continued

<210> SEQ ID NO 53
 <211> LENGTH: 7
 <212> TYPE: PRT
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: light chain CDR2, hMab 2F11-g1

<400> SEQUENCE: 53

Gly Ala Ser Ser Arg Ala Thr
 1 5

<210> SEQ ID NO 54
 <211> LENGTH: 11
 <212> TYPE: PRT
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: light chain CDR1, hMab 2F11-g1

<400> SEQUENCE: 54

Arg Ala Ser Glu Asp Val Asn Thr Tyr Leu Ala
 1 5 10

<210> SEQ ID NO 55
 <211> LENGTH: 116
 <212> TYPE: PRT
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: heavy chain variable domain, hMab 2F11-g1

<400> SEQUENCE: 55

Gln Val Gln Leu Gln Glu Ser Gly Pro Gly Leu Val Lys Pro Ser Glu
 1 5 10 15

Thr Leu Ser Leu Thr Cys Thr Val Ser Gly Phe Ser Leu Thr Thr Tyr
 20 25 30

Asp Ile Ser Trp Ile Arg Gln Pro Pro Gly Lys Gly Leu Glu Trp Ile
 35 40 45

Gly Val Ile Trp Thr Asp Gly Gly Thr Asn Tyr Asn Ser Pro Leu Lys
 50 55 60

Ser Arg Val Thr Ile Ser Val Asp Thr Ser Lys Asn Gln Phe Ser Leu
 65 70 75 80

Lys Leu Ser Ser Val Thr Ala Ala Asp Thr Ala Val Tyr Tyr Cys Ala
 85 90 95

Arg Asp Gln Arg Leu Tyr Phe Asp Val Trp Gly Gln Gly Thr Thr Val
 100 105 110

Thr Val Ser Ser
 115

<210> SEQ ID NO 56
 <211> LENGTH: 106
 <212> TYPE: PRT
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: light chain variable domain, hMab 2F11-g1

<400> SEQUENCE: 56

Glu Ile Val Leu Thr Gln Ser Pro Gly Thr Leu Ser Leu Ser Pro Gly
 1 5 10 15

Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Glu Asp Val Asn Thr Tyr
 20 25 30

-continued

Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu Ile
 35 40 45
 Tyr Gly Ala Ser Ser Arg Ala Thr Gly Ile Pro Asp Arg Phe Ser Gly
 50 55 60
 Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Arg Leu Glu Pro
 65 70 75 80
 Glu Asp Phe Ala Val Tyr Tyr Cys Gly Gln Ser Phe Ser Tyr Pro Thr
 85 90 95
 Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
 100 105

<210> SEQ ID NO 57
 <211> LENGTH: 107
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 57

Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu
 1 5 10 15
 Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe
 20 25 30
 Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln
 35 40 45
 Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser
 50 55 60
 Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu
 65 70 75 80
 Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser
 85 90 95
 Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
 100 105

<210> SEQ ID NO 58
 <211> LENGTH: 330
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 58

Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys
 1 5 10 15
 Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
 20 25 30
 Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
 35 40 45
 Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
 50 55 60
 Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr
 65 70 75 80
 Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys
 85 90 95
 Lys Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys
 100 105 110
 Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro
 115 120 125

-continued

```

Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys
 130                135                140

Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp
145                150                155                160

Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu
                165                170                175

Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu
                180                185                190

His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn
                195                200                205

Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly
                210                215                220

Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu
225                230                235                240

Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr
                245                250                255

Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn
                260                265                270

Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe
                275                280                285

Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn
290                295                300

Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr
305                310                315                320

Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
                325                330

```

<210> SEQ ID NO 59

<211> LENGTH: 330

<212> TYPE: PRT

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: human heavy chain constant region derived from IgG1 mutated on L234A and L235A

<400> SEQUENCE: 59

```

Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys
 1          5          10          15

Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
                20          25          30

Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
                35          40          45

Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
50          55          60

Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr
65          70          75          80

Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys
                85          90          95

Lys Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys
                100         105         110

Pro Ala Pro Glu Ala Ala Gly Gly Pro Ser Val Phe Leu Phe Pro Pro
                115         120         125

Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys
130                135                140

```

-continued

Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp
 145 150 155 160
 Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu
 165 170 175
 Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu
 180 185 190
 His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn
 195 200 205
 Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly
 210 215 220
 Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu
 225 230 235 240
 Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr
 245 250 255
 Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn
 260 265 270
 Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe
 275 280 285
 Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn
 290 295 300
 Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr
 305 310 315 320
 Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
 325 330

<210> SEQ ID NO 60

<211> LENGTH: 327

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 60

Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Cys Ser Arg
 1 5 10 15
 Ser Thr Ser Glu Ser Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
 20 25 30
 Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
 35 40 45
 Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
 50 55 60
 Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Lys Thr
 65 70 75 80
 Tyr Thr Cys Asn Val Asp His Lys Pro Ser Asn Thr Lys Val Asp Lys
 85 90 95
 Arg Val Glu Ser Lys Tyr Gly Pro Pro Cys Pro Ser Cys Pro Ala Pro
 100 105 110
 Glu Phe Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys
 115 120 125
 Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val
 130 135 140
 Asp Val Ser Gln Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr Val Asp
 145 150 155 160
 Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Phe

-continued

	165		170		175															
Asn	Ser	Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp					
			180						185					190						
Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Gly	Leu					
		195					200					205								
Pro	Ser	Ser	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg					
	210					215					220									
Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Gln	Glu	Glu	Met	Thr	Lys					
225					230					235					240					
Asn	Gln	Val	Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp					
			245						250						255					
Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys					
		260						265						270						
Thr	Thr	Pro	Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser					
		275						280					285							
Arg	Leu	Thr	Val	Asp	Lys	Ser	Arg	Trp	Gln	Glu	Gly	Asn	Val	Phe	Ser					
	290					295					300									
Cys	Ser	Val	Met	His	Glu	Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser					
305					310					315					320					
Leu	Ser	Leu	Ser	Leu	Gly	Lys														
				325																

<210> SEQ ID NO 61
 <211> LENGTH: 327
 <212> TYPE: PRT
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: human heavy chain constant region derived from
 IgG4 mutated onS228P

<400> SEQUENCE: 61

Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu	Ala	Pro	Cys	Ser	Arg
1				5					10					15	
Ser	Thr	Ser	Glu	Ser	Thr	Ala	Ala	Leu	Gly	Cys	Leu	Val	Lys	Asp	Tyr
			20					25					30		
Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser	Gly	Ala	Leu	Thr	Ser
		35					40				45				
Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser	Ser	Gly	Leu	Tyr	Ser
		50				55					60				
Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser	Ser	Leu	Gly	Thr	Lys	Thr
65					70					75					80
Tyr	Thr	Cys	Asn	Val	Asp	His	Lys	Pro	Ser	Asn	Thr	Lys	Val	Asp	Lys
			85						90					95	
Arg	Val	Glu	Ser	Lys	Tyr	Gly	Pro	Pro	Cys	Pro	Pro	Cys	Pro	Ala	Pro
		100						105					110		
Glu	Phe	Leu	Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys
		115					120					125			
Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val
		130				135					140				
Asp	Val	Ser	Gln	Glu	Asp	Pro	Glu	Val	Gln	Phe	Asn	Trp	Tyr	Val	Asp
145					150					155					160
Gly	Val	Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Phe
				165					170						175

-continued

```

Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp
      180                               185                190

Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu
      195                               200                205

Pro Ser Ser Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg
      210                               215                220

Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Gln Glu Glu Met Thr Lys
      225                               230                235                240

Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp
      245                               250                255

Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys
      260                               265                270

Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser
      275                               280                285

Arg Leu Thr Val Asp Lys Ser Arg Trp Gln Glu Gly Asn Val Phe Ser
      290                               295                300

Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser
      305                               310                315                320

Leu Ser Leu Ser Leu Gly Lys
      325

```

<210> SEQ ID NO 62

<211> LENGTH: 972

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 62

```

Met Gly Pro Gly Val Leu Leu Leu Leu Val Ala Thr Ala Trp His
  1      5      10      15

Gly Gln Gly Ile Pro Val Ile Glu Pro Ser Val Pro Glu Leu Val Val
  20      25      30

Lys Pro Gly Ala Thr Val Thr Leu Arg Cys Val Gly Asn Gly Ser Val
  35      40      45

Glu Trp Asp Gly Pro Pro Ser Pro His Trp Thr Leu Tyr Ser Asp Gly
  50      55      60

Ser Ser Ser Ile Leu Ser Thr Asn Asn Ala Thr Phe Gln Asn Thr Gly
  65      70      75      80

Thr Tyr Arg Cys Thr Glu Pro Gly Asp Pro Leu Gly Gly Ser Ala Ala
  85      90      95

Ile His Leu Tyr Val Lys Asp Pro Ala Arg Pro Trp Asn Val Leu Ala
  100     105     110

Gln Glu Val Val Val Phe Glu Asp Gln Asp Ala Leu Leu Pro Cys Leu
  115     120     125

Leu Thr Asp Pro Val Leu Glu Ala Gly Val Ser Leu Val Arg Val Arg
  130     135     140

Gly Arg Pro Leu Met Arg His Thr Asn Tyr Ser Phe Ser Pro Trp His
  145     150     155     160

Gly Phe Thr Ile His Arg Ala Lys Phe Ile Gln Ser Gln Asp Tyr Gln
  165     170     175

Cys Ser Ala Leu Met Gly Gly Arg Lys Val Met Ser Ile Ser Ile Arg
  180     185     190

Leu Lys Val Gln Lys Val Ile Pro Gly Pro Pro Ala Leu Thr Leu Val
  195     200     205

```

-continued

Pro Ala Glu Leu Val Arg Ile Arg Gly Glu Ala Ala Gln Ile Val Cys
 210 215 220
 Ser Ala Ser Ser Val Asp Val Asn Phe Asp Val Phe Leu Gln His Asn
 225 230 235 240
 Asn Thr Lys Leu Ala Ile Pro Gln Gln Ser Asp Phe His Asn Asn Arg
 245 250 255
 Tyr Gln Lys Val Leu Thr Leu Asn Leu Asp Gln Val Asp Phe Gln His
 260 265 270
 Ala Gly Asn Tyr Ser Cys Val Ala Ser Asn Val Gln Gly Lys His Ser
 275 280 285
 Thr Ser Met Phe Phe Arg Val Val Glu Ser Ala Tyr Leu Asn Leu Ser
 290 295 300
 Ser Glu Gln Asn Leu Ile Gln Glu Val Thr Val Gly Glu Gly Leu Asn
 305 310 315 320
 Leu Lys Val Met Val Glu Ala Tyr Pro Gly Leu Gln Gly Phe Asn Trp
 325 330 335
 Thr Tyr Leu Gly Pro Phe Ser Asp His Gln Pro Glu Pro Lys Leu Ala
 340 345 350
 Asn Ala Thr Thr Lys Asp Thr Tyr Arg His Thr Phe Thr Leu Ser Leu
 355 360 365
 Pro Arg Leu Lys Pro Ser Glu Ala Gly Arg Tyr Ser Phe Leu Ala Arg
 370 375 380
 Asn Pro Gly Gly Trp Arg Ala Leu Thr Phe Glu Leu Thr Leu Arg Tyr
 385 390 395 400
 Pro Pro Glu Val Ser Val Ile Trp Thr Phe Ile Asn Gly Ser Gly Thr
 405 410 415
 Leu Leu Cys Ala Ala Ser Gly Tyr Pro Gln Pro Asn Val Thr Trp Leu
 420 425 430
 Gln Cys Ser Gly His Thr Asp Arg Cys Asp Glu Ala Gln Val Leu Gln
 435 440 445
 Val Trp Asp Asp Pro Tyr Pro Glu Val Leu Ser Gln Glu Pro Phe His
 450 455 460
 Lys Val Thr Val Gln Ser Leu Leu Thr Val Glu Thr Leu Glu His Asn
 465 470 475 480
 Gln Thr Tyr Glu Cys Arg Ala His Asn Ser Val Gly Ser Gly Ser Trp
 485 490 495
 Ala Phe Ile Pro Ile Ser Ala Gly Ala His Thr His Pro Pro Asp Glu
 500 505 510
 Phe Leu Phe Thr Pro Val Val Val Ala Cys Met Ser Ile Met Ala Leu
 515 520 525
 Leu Leu Leu Leu Leu Leu Leu Leu Tyr Lys Tyr Lys Gln Lys Pro
 530 535 540
 Lys Tyr Gln Val Arg Trp Lys Ile Ile Glu Ser Tyr Glu Gly Asn Ser
 545 550 555 560
 Tyr Thr Phe Ile Asp Pro Thr Gln Leu Pro Tyr Asn Glu Lys Trp Glu
 565 570 575
 Phe Pro Arg Asn Asn Leu Gln Phe Gly Lys Thr Leu Gly Ala Gly Ala
 580 585 590
 Phe Gly Lys Val Val Glu Ala Thr Ala Phe Gly Leu Gly Lys Glu Asp
 595 600 605

-continued

Ala Val Leu Lys Val Ala Val Lys Met Leu Lys Ser Thr Ala His Ala
610 615 620

Asp Glu Lys Glu Ala Leu Met Ser Glu Leu Lys Ile Met Ser His Leu
625 630 635 640

Gly Gln His Glu Asn Ile Val Asn Leu Leu Gly Ala Cys Thr His Gly
645 650 655

Gly Pro Val Leu Val Ile Thr Glu Tyr Cys Cys Tyr Gly Asp Leu Leu
660 665 670

Asn Phe Leu Arg Arg Lys Ala Glu Ala Met Leu Gly Pro Ser Leu Ser
675 680 685

Pro Gly Gln Asp Pro Glu Gly Gly Val Asp Tyr Lys Asn Ile His Leu
690 695 700

Glu Lys Lys Tyr Val Arg Arg Asp Ser Gly Phe Ser Ser Gln Gly Val
705 710 715 720

Asp Thr Tyr Val Glu Met Arg Pro Val Ser Thr Ser Ser Asn Asp Ser
725 730 735

Phe Ser Glu Gln Asp Leu Asp Lys Glu Asp Gly Arg Pro Leu Glu Leu
740 745 750

Arg Asp Leu Leu His Phe Ser Ser Gln Val Ala Gln Gly Met Ala Phe
755 760 765

Leu Ala Ser Lys Asn Cys Ile His Arg Asp Val Ala Ala Arg Asn Val
770 775 780

Leu Leu Thr Asn Gly His Val Ala Lys Ile Gly Asp Phe Gly Leu Ala
785 790 795 800

Arg Asp Ile Met Asn Asp Ser Asn Tyr Ile Val Lys Gly Asn Ala Arg
805 810 815

Leu Pro Val Lys Trp Met Ala Pro Glu Ser Ile Phe Asp Cys Val Tyr
820 825 830

Thr Val Gln Ser Asp Val Trp Ser Tyr Gly Ile Leu Leu Trp Glu Ile
835 840 845

Phe Ser Leu Gly Leu Asn Pro Tyr Pro Gly Ile Leu Val Asn Ser Lys
850 855 860

Phe Tyr Lys Leu Val Lys Asp Gly Tyr Gln Met Ala Gln Pro Ala Phe
865 870 875 880

Ala Pro Lys Asn Ile Tyr Ser Ile Met Gln Ala Cys Trp Ala Leu Glu
885 890 895

Pro Thr His Arg Pro Thr Phe Gln Gln Ile Cys Ser Phe Leu Gln Glu
900 905 910

Gln Ala Gln Glu Asp Arg Arg Glu Arg Asp Tyr Thr Asn Leu Pro Ser
915 920 925

Ser Ser Arg Ser Gly Gly Ser Gly Ser Ser Ser Ser Glu Leu Glu Glu
930 935 940

Glu Ser Ser Ser Glu His Leu Thr Cys Cys Glu Gln Gly Asp Ile Ala
945 950 955 960

Gln Pro Leu Leu Gln Pro Asn Asn Tyr Gln Phe Cys
965 970

<210> SEQ ID NO 63

<211> LENGTH: 972

<212> TYPE: PRT

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: mutant CSF-1R L301S Y969F

-continued

<400> SEQUENCE: 63

Met Gly Pro Gly Val Leu Leu Leu Leu Val Ala Thr Ala Trp His
1 5 10 15
Gly Gln Gly Ile Pro Val Ile Glu Pro Ser Val Pro Glu Leu Val Val
20 25 30
Lys Pro Gly Ala Thr Val Thr Leu Arg Cys Val Gly Asn Gly Ser Val
35 40 45
Glu Trp Asp Gly Pro Pro Ser Pro His Trp Thr Leu Tyr Ser Asp Gly
50 55 60
Ser Ser Ser Ile Leu Ser Thr Asn Asn Ala Thr Phe Gln Asn Thr Gly
65 70 75 80
Thr Tyr Arg Cys Thr Glu Pro Gly Asp Pro Leu Gly Gly Ser Ala Ala
85 90 95
Ile His Leu Tyr Val Lys Asp Pro Ala Arg Pro Trp Asn Val Leu Ala
100 105 110
Gln Glu Val Val Val Phe Glu Asp Gln Asp Ala Leu Leu Pro Cys Leu
115 120 125
Leu Thr Asp Pro Val Leu Glu Ala Gly Val Ser Leu Val Arg Val Arg
130 135 140
Gly Arg Pro Leu Met Arg His Thr Asn Tyr Ser Phe Ser Pro Trp His
145 150 155 160
Gly Phe Thr Ile His Arg Ala Lys Phe Ile Gln Ser Gln Asp Tyr Gln
165 170 175
Cys Ser Ala Leu Met Gly Gly Arg Lys Val Met Ser Ile Ser Ile Arg
180 185 190
Leu Lys Val Gln Lys Val Ile Pro Gly Pro Pro Ala Leu Thr Leu Val
195 200 205
Pro Ala Glu Leu Val Arg Ile Arg Gly Glu Ala Ala Gln Ile Val Cys
210 215 220
Ser Ala Ser Ser Val Asp Val Asn Phe Asp Val Phe Leu Gln His Asn
225 230 235 240
Asn Thr Lys Leu Ala Ile Pro Gln Gln Ser Asp Phe His Asn Asn Arg
245 250 255
Tyr Gln Lys Val Leu Thr Leu Asn Leu Asp Gln Val Asp Phe Gln His
260 265 270
Ala Gly Asn Tyr Ser Cys Val Ala Ser Asn Val Gln Gly Lys His Ser
275 280 285
Thr Ser Met Phe Phe Arg Val Val Glu Ser Ala Tyr Ser Asn Leu Ser
290 295 300
Ser Glu Gln Asn Leu Ile Gln Glu Val Thr Val Gly Glu Gly Leu Asn
305 310 315 320
Leu Lys Val Met Val Glu Ala Tyr Pro Gly Leu Gln Gly Phe Asn Trp
325 330 335
Thr Tyr Leu Gly Pro Phe Ser Asp His Gln Pro Glu Pro Lys Leu Ala
340 345 350
Asn Ala Thr Thr Lys Asp Thr Tyr Arg His Thr Phe Thr Leu Ser Leu
355 360 365
Pro Arg Leu Lys Pro Ser Glu Ala Gly Arg Tyr Ser Phe Leu Ala Arg
370 375 380
Asn Pro Gly Gly Trp Arg Ala Leu Thr Phe Glu Leu Thr Leu Arg Tyr

-continued

```

Arg Asp Ile Met Asn Asp Ser Asn Tyr Ile Val Lys Gly Asn Ala Arg
      805                               810                       815
Leu Pro Val Lys Trp Met Ala Pro Glu Ser Ile Phe Asp Cys Val Tyr
      820                               825                       830
Thr Val Gln Ser Asp Val Trp Ser Tyr Gly Ile Leu Leu Trp Glu Ile
      835                               840                       845
Phe Ser Leu Gly Leu Asn Pro Tyr Pro Gly Ile Leu Val Asn Ser Lys
      850                               855                       860
Phe Tyr Lys Leu Val Lys Asp Gly Tyr Gln Met Ala Gln Pro Ala Phe
      865                               870                       875                       880
Ala Pro Lys Asn Ile Tyr Ser Ile Met Gln Ala Cys Trp Ala Leu Glu
      885                               890                       895
Pro Thr His Arg Pro Thr Phe Gln Gln Ile Cys Ser Phe Leu Gln Glu
      900                               905                       910
Gln Ala Gln Glu Asp Arg Arg Glu Arg Asp Tyr Thr Asn Leu Pro Ser
      915                               920                       925
Ser Ser Arg Ser Gly Gly Ser Gly Ser Ser Ser Ser Glu Leu Glu Glu
      930                               935                       940
Glu Ser Ser Ser Glu His Leu Thr Cys Cys Glu Gln Gly Asp Ile Ala
      945                               950                       955                       960
Gln Pro Leu Leu Gln Pro Asn Asn Phe Gln Phe Cys
      965                               970

```

```

<210> SEQ ID NO 64
<211> LENGTH: 493
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: human CSF-1R Extracellular Domain

```

```

<400> SEQUENCE: 64

```

```

Ile Pro Val Ile Glu Pro Ser Val Pro Glu Leu Val Val Lys Pro Gly
1      5      10      15
Ala Thr Val Thr Leu Arg Cys Val Gly Asn Gly Ser Val Glu Trp Asp
20     25     30
Gly Pro Pro Ser Pro His Trp Thr Leu Tyr Ser Asp Gly Ser Ser Ser
35     40     45
Ile Leu Ser Thr Asn Asn Ala Thr Phe Gln Asn Thr Gly Thr Tyr Arg
50     55     60
Cys Thr Glu Pro Gly Asp Pro Leu Gly Gly Ser Ala Ala Ile His Leu
65     70     75
Tyr Val Lys Asp Pro Ala Arg Pro Trp Asn Val Leu Ala Gln Glu Val
85     90     95
Val Val Phe Glu Asp Gln Asp Ala Leu Leu Pro Cys Leu Leu Thr Asp
100    105   110
Pro Val Leu Glu Ala Gly Val Ser Leu Val Arg Val Arg Gly Arg Pro
115    120   125
Leu Met Arg His Thr Asn Tyr Ser Phe Ser Pro Trp His Gly Phe Thr
130    135   140
Ile His Arg Ala Lys Phe Ile Gln Ser Gln Asp Tyr Gln Cys Ser Ala
145    150   155   160
Leu Met Gly Gly Arg Lys Val Met Ser Ile Ser Ile Arg Leu Lys Val
165    170   175

```

-continued

Gln Lys Val Ile Pro Gly Pro Pro Ala Leu Thr Leu Val Pro Ala Glu
 180 185 190
 Leu Val Arg Ile Arg Gly Glu Ala Ala Gln Ile Val Cys Ser Ala Ser
 195 200 205
 Ser Val Asp Val Asn Phe Asp Val Phe Leu Gln His Asn Asn Thr Lys
 210 215 220
 Leu Ala Ile Pro Gln Gln Ser Asp Phe His Asn Asn Arg Tyr Gln Lys
 225 230 235 240
 Val Leu Thr Leu Asn Leu Asp Gln Val Asp Phe Gln His Ala Gly Asn
 245 250 255
 Tyr Ser Cys Val Ala Ser Asn Val Gln Gly Lys His Ser Thr Ser Met
 260 265 270
 Phe Phe Arg Val Val Glu Ser Ala Tyr Leu Asn Leu Ser Ser Glu Gln
 275 280 285
 Asn Leu Ile Gln Glu Val Thr Val Gly Glu Gly Leu Asn Leu Lys Val
 290 295 300
 Met Val Glu Ala Tyr Pro Gly Leu Gln Gly Phe Asn Trp Thr Tyr Leu
 305 310 315 320
 Gly Pro Phe Ser Asp His Gln Pro Glu Pro Lys Leu Ala Asn Ala Thr
 325 330 335
 Thr Lys Asp Thr Tyr Arg His Thr Phe Thr Leu Ser Leu Pro Arg Leu
 340 345 350
 Lys Pro Ser Glu Ala Gly Arg Tyr Ser Phe Leu Ala Arg Asn Pro Gly
 355 360 365
 Gly Trp Arg Ala Leu Thr Phe Glu Leu Thr Leu Arg Tyr Pro Pro Glu
 370 375 380
 Val Ser Val Ile Trp Thr Phe Ile Asn Gly Ser Gly Thr Leu Leu Cys
 385 390 395 400
 Ala Ala Ser Gly Tyr Pro Gln Pro Asn Val Thr Trp Leu Gln Cys Ser
 405 410 415
 Gly His Thr Asp Arg Cys Asp Glu Ala Gln Val Leu Gln Val Trp Asp
 420 425 430
 Asp Pro Tyr Pro Glu Val Leu Ser Gln Glu Pro Phe His Lys Val Thr
 435 440 445
 Val Gln Ser Leu Leu Thr Val Glu Thr Leu Glu His Asn Gln Thr Tyr
 450 455 460
 Glu Cys Arg Ala His Asn Ser Val Gly Ser Gly Ser Trp Ala Phe Ile
 465 470 475 480
 Pro Ile Ser Ala Gly Ala His Thr His Pro Pro Asp Glu
 485 490

<210> SEQ ID NO 65

<211> LENGTH: 388

<212> TYPE: PRT

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: human CSF-1R fragment delD4

<400> SEQUENCE: 65

Ile Pro Val Ile Glu Pro Ser Val Pro Glu Leu Val Val Lys Pro Gly
 1 5 10 15
 Ala Thr Val Thr Leu Arg Cys Val Gly Asn Gly Ser Val Glu Trp Asp
 20 25 30

-continued

Gly Pro Pro Ser Pro His Trp Thr Leu Tyr Ser Asp Gly Ser Ser Ser
 35 40 45
 Ile Leu Ser Thr Asn Asn Ala Thr Phe Gln Asn Thr Gly Thr Tyr Arg
 50 55 60
 Cys Thr Glu Pro Gly Asp Pro Leu Gly Gly Ser Ala Ala Ile His Leu
 65 70 75 80
 Tyr Val Lys Asp Pro Ala Arg Pro Trp Asn Val Leu Ala Gln Glu Val
 85 90 95
 Val Val Phe Glu Asp Gln Asp Ala Leu Leu Pro Cys Leu Leu Thr Asp
 100 105 110
 Pro Val Leu Glu Ala Gly Val Ser Leu Val Arg Val Arg Gly Arg Pro
 115 120 125
 Leu Met Arg His Thr Asn Tyr Ser Phe Ser Pro Trp His Gly Phe Thr
 130 135 140
 Ile His Arg Ala Lys Phe Ile Gln Ser Gln Asp Tyr Gln Cys Ser Ala
 145 150 155 160
 Leu Met Gly Gly Arg Lys Val Met Ser Ile Ser Ile Arg Leu Lys Val
 165 170 175
 Gln Lys Val Ile Pro Gly Pro Pro Ala Leu Thr Leu Val Pro Ala Glu
 180 185 190
 Leu Val Arg Ile Arg Gly Glu Ala Ala Gln Ile Val Cys Ser Ala Ser
 195 200 205
 Ser Val Asp Val Asn Phe Asp Val Phe Leu Gln His Asn Asn Thr Lys
 210 215 220
 Leu Ala Ile Pro Gln Gln Ser Asp Phe His Asn Asn Arg Tyr Gln Lys
 225 230 235 240
 Val Leu Thr Leu Asn Leu Asp Gln Val Asp Phe Gln His Ala Gly Asn
 245 250 255
 Tyr Ser Cys Val Ala Ser Asn Val Gln Gly Lys His Ser Thr Ser Met
 260 265 270
 Phe Phe Arg Tyr Pro Pro Glu Val Ser Val Ile Trp Thr Phe Ile Asn
 275 280 285
 Gly Ser Gly Thr Leu Leu Cys Ala Ala Ser Gly Tyr Pro Gln Pro Asn
 290 295 300
 Val Thr Trp Leu Gln Cys Ser Gly His Thr Asp Arg Cys Asp Glu Ala
 305 310 315 320
 Gln Val Leu Gln Val Trp Asp Asp Pro Tyr Pro Glu Val Leu Ser Gln
 325 330 335
 Glu Pro Phe His Lys Val Thr Val Gln Ser Leu Leu Thr Val Glu Thr
 340 345 350
 Leu Glu His Asn Gln Thr Tyr Glu Cys Arg Ala His Asn Ser Val Gly
 355 360 365
 Ser Gly Ser Trp Ala Phe Ile Pro Ile Ser Ala Gly Ala His Thr His
 370 375 380
 Pro Pro Asp Glu
 385

<210> SEQ ID NO 66

<211> LENGTH: 292

<212> TYPE: PRT

<213> ORGANISM: Artificial

<220> FEATURE:

-continued

<223> OTHER INFORMATION: human CSF-1R fragment D1-D3

<400> SEQUENCE: 66

Ile Pro Val Ile Glu Pro Ser Val Pro Glu Leu Val Val Lys Pro Gly
 1 5 10 15

Ala Thr Val Thr Leu Arg Cys Val Gly Asn Gly Ser Val Glu Trp Asp
 20 25 30

Gly Pro Pro Ser Pro His Trp Thr Leu Tyr Ser Asp Gly Ser Ser Ser
 35 40 45

Ile Leu Ser Thr Asn Asn Ala Thr Phe Gln Asn Thr Gly Thr Tyr Arg
 50 55 60

Cys Thr Glu Pro Gly Asp Pro Leu Gly Gly Ser Ala Ala Ile His Leu
 65 70 75 80

Tyr Val Lys Asp Pro Ala Arg Pro Trp Asn Val Leu Ala Gln Glu Val
 85 90 95

Val Val Phe Glu Asp Gln Asp Ala Leu Leu Pro Cys Leu Leu Thr Asp
 100 105 110

Pro Val Leu Glu Ala Gly Val Ser Leu Val Arg Val Arg Gly Arg Pro
 115 120 125

Leu Met Arg His Thr Asn Tyr Ser Phe Ser Pro Trp His Gly Phe Thr
 130 135 140

Ile His Arg Ala Lys Phe Ile Gln Ser Gln Asp Tyr Gln Cys Ser Ala
 145 150 155 160

Leu Met Gly Gly Arg Lys Val Met Ser Ile Ser Ile Arg Leu Lys Val
 165 170 175

Gln Lys Val Ile Pro Gly Pro Pro Ala Leu Thr Leu Val Pro Ala Glu
 180 185 190

Leu Val Arg Ile Arg Gly Glu Ala Ala Gln Ile Val Cys Ser Ala Ser
 195 200 205

Ser Val Asp Val Asn Phe Asp Val Phe Leu Gln His Asn Asn Thr Lys
 210 215 220

Leu Ala Ile Pro Gln Gln Ser Asp Phe His Asn Asn Arg Tyr Gln Lys
 225 230 235 240

Val Leu Thr Leu Asn Leu Asp Gln Val Asp Phe Gln His Ala Gly Asn
 245 250 255

Tyr Ser Cys Val Ala Ser Asn Val Gln Gly Lys His Ser Thr Ser Met
 260 265 270

Phe Phe Arg Val Val Glu Ser Ala Tyr Leu Asn Leu Ser Ser Glu Gln
 275 280 285

Asn Leu Ile Gln
 290

<210> SEQ ID NO 67

<211> LENGTH: 21

<212> TYPE: PRT

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: signal peptide

<400> SEQUENCE: 67

Met Gly Ser Gly Pro Gly Val Leu Leu Leu Leu Val Ala Thr Ala
 1 5 10 15

Trp His Gly Gln Gly
 20

-continued

<210> SEQ ID NO 68
<211> LENGTH: 36
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: Primer

<400> SEQUENCE: 68

cacctccatg ttcttcgggt acccccaga ggtaag

36

<210> SEQ ID NO 69
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 69

Asp Leu Arg Leu Tyr Phe Asp Val
1 5

<210> SEQ ID NO 70
<211> LENGTH: 16
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 70

Val Ile Trp Ser Gly Gly Gly Thr Asn Tyr Asn Ser Pro Phe Met Ser
1 5 10 15

<210> SEQ ID NO 71
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 71

Gly Phe Ser Leu Thr Ser Tyr Asp Ile Ser
1 5 10

<210> SEQ ID NO 72
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 72

Gly Gln Ser Phe Thr Tyr Pro Thr
1 5

<210> SEQ ID NO 73
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 73

Gly Ser Ser Asn Arg Tyr Thr
1 5

<210> SEQ ID NO 74
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 74

-continued

Lys Ala Ser Glu Asp Val Gly Thr Tyr Val Ser
1 5 10

<210> SEQ ID NO 75
<211> LENGTH: 116
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 75

Arg Val Gln Leu Lys Glu Ser Gly Pro Gly Leu Val Ala Pro Ser Gln
1 5 10 15
Ser Leu Ser Ile Thr Cys Thr Val Ser Gly Phe Ser Leu Thr Ser Tyr
20 25 30
Asp Ile Ser Trp Ile Arg Gln Ser Pro Gly Lys Gly Leu Glu Trp Leu
35 40 45
Gly Val Ile Trp Ser Gly Gly Thr Asn Tyr Asn Ser Pro Phe Met
50 55 60
Ser Arg Leu Arg Ile Ser Lys Asp Asp Ser Arg Ser Gln Val Phe Leu
65 70 75 80
Lys Val Asn Arg Leu Gln Thr Asp Asp Thr Ala Ile Tyr Tyr Cys Val
85 90 95
Arg Asp Leu Arg Leu Tyr Phe Asp Val Trp Gly Ala Gly Thr Thr Val
100 105 110
Thr Val Ser Ser
115

<210> SEQ ID NO 76
<211> LENGTH: 106
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 76

Lys Ile Val Met Thr Gln Ser Pro Lys Ser Met Ser Val Ser Val Gly
1 5 10 15
Glu Arg Val Ser Leu Ser Cys Lys Ala Ser Glu Asp Val Gly Thr Tyr
20 25 30
Val Ser Trp Tyr Gln Gln Lys Pro Glu Gln Ser Pro Lys Leu Leu Ile
35 40 45
Tyr Gly Ser Ser Asn Arg Tyr Thr Gly Val Pro Asp Arg Phe Thr Gly
50 55 60
Ser Gly Ser Ala Thr Asp Phe Thr Leu Thr Ile Ser Ser Val Gln Ala
65 70 75 80
Glu Asp Leu Ala Asp Tyr Ser Cys Gly Gln Ser Phe Thr Tyr Pro Thr
85 90 95
Phe Gly Thr Gly Thr Lys Leu Glu Ile Lys
100 105

<210> SEQ ID NO 77
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 77

Asp Pro Arg Leu Tyr Phe Asp Val
1 5

-continued

<210> SEQ ID NO 78
 <211> LENGTH: 16
 <212> TYPE: PRT
 <213> ORGANISM: Mus musculus
 <400> SEQUENCE: 78
 Val Ile Trp Thr Gly Gly Gly Thr Asn Tyr Asn Ser Gly Phe Met Ser
 1 5 10 15

<210> SEQ ID NO 79
 <211> LENGTH: 10
 <212> TYPE: PRT
 <213> ORGANISM: Mus musculus
 <400> SEQUENCE: 79
 Gly Ser Ser Leu Asp Ser Phe Asp Ile Ser
 1 5 10

<210> SEQ ID NO 80
 <211> LENGTH: 8
 <212> TYPE: PRT
 <213> ORGANISM: Mus musculus
 <400> SEQUENCE: 80
 Gly Gln Thr Phe Ser Tyr Pro Thr
 1 5

<210> SEQ ID NO 81
 <211> LENGTH: 7
 <212> TYPE: PRT
 <213> ORGANISM: Mus musculus
 <400> SEQUENCE: 81
 Gly Ala Ser Asn Arg Tyr Thr
 1 5

<210> SEQ ID NO 82
 <211> LENGTH: 11
 <212> TYPE: PRT
 <213> ORGANISM: Mus musculus
 <400> SEQUENCE: 82
 Lys Ala Ser Glu Asp Val Val Thr Tyr Val Ser
 1 5 10

<210> SEQ ID NO 83
 <211> LENGTH: 116
 <212> TYPE: PRT
 <213> ORGANISM: Mus musculus
 <400> SEQUENCE: 83
 Gln Val Gln Leu Lys Glu Ser Gly Pro Gly Leu Val Ala Pro Ser Lys
 1 5 10 15
 Ser Leu Ser Ile Thr Cys Thr Val Ser Gly Ser Ser Leu Asp Ser Phe
 20 25 30
 Asp Ile Ser Trp Ile Arg Gln Pro Pro Gly Lys Gly Leu Glu Trp Leu
 35 40 45
 Gly Val Ile Trp Thr Gly Gly Gly Thr Asn Tyr Asn Ser Gly Phe Met
 50 55 60
 Ser Arg Leu Arg Ile Ser Lys Asp Asn Ser Lys Ser Gln Val Phe Leu
 65 70 75 80

-continued

```

Lys Met Ser Ser Leu Gln Ser Asp Asp Thr Ala Ile Tyr Tyr Cys Val
           85                               90                       95
Arg Asp Pro Arg Leu Tyr Phe Asp Val Trp Gly Ala Gly Thr Thr Val
           100                             105                       110
Thr Val Ser Ser
           115

```

```

<210> SEQ ID NO 84
<211> LENGTH: 106
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

```

```

<400> SEQUENCE: 84

```

```

Asn Ile Val Met Thr Gln Ser Pro Lys Ser Met Ser Met Ser Val Gly
 1           5                               10                       15
Glu Arg Val Thr Leu Ser Cys Lys Ala Ser Glu Asp Val Val Thr Tyr
           20                             25                       30
Val Ser Trp Tyr Gln Gln Lys Pro Glu Gln Ser Pro Lys Leu Leu Ile
           35                             40                       45
Tyr Gly Ala Ser Asn Arg Tyr Thr Gly Val Pro Asp Arg Phe Thr Gly
           50                             55                       60
Ser Gly Ser Ala Thr Asp Phe Thr Leu Thr Ile Ser Ser Ile Gln Ala
           65                             70                       75                       80
Glu Asp Leu Ala Asp Tyr Tyr Cys Gly Gln Thr Phe Ser Tyr Pro Thr
           85                             90                       95
Phe Gly Thr Gly Thr Lys Leu Glu Ile Lys
           100                             105

```

1. An isolated antibody binding to human CSF-1R, wherein the antibody comprises a heavy chain variable domain and a light chain variable domain, and wherein the heavy chain variable domain comprises a CDR3 region comprising SEQ ID NO: 41, a CDR2 region comprising SEQ ID NO: 42, and a CDR1 region comprising SEQ ID NO: 43, and the light chain variable domain comprises a CDR3 region comprising SEQ ID NO: 44, a CDR2 region comprising SEQ ID NO: 45, and a CDR1 region comprising SEQ ID NO: 46.

2. A pharmaceutical composition comprising the antibody according to claim 1.

3. The pharmaceutical composition according to claim 2, wherein the antibody is formulated in a pharmaceutically acceptable carrier.

4. The antibody according to claim 1, wherein the antibody is of human IgG1 subclass or is of human IgG4 subclass.

5. The antibody according to claim 1, wherein the antibody is a chimeric, single chain, multispecific, or humanized antibody.

6. A pharmaceutical composition comprising the antibody according to claim 5.

7. The antibody according to claim 1, wherein the antibody is an antigen binding fragment.

8. A pharmaceutical composition comprising the antibody according to claim 7.

9. The antibody according to claim 1, wherein the antibody is a monoclonal antibody.

10. A pharmaceutical composition comprising the antibody according to claim 9.

11. An isolated antibody binding to human CSF-1R, wherein the antibody comprises a heavy chain variable domain and a light chain variable domain, and wherein the heavy chain variable domain comprises SEQ ID NO: 47 and the light chain variable domain comprises SEQ ID NO: 48.

12. A pharmaceutical composition comprising the antibody according to claim 11.

13. The pharmaceutical composition according to claim 12, wherein the antibody is formulated in a pharmaceutically acceptable carrier.

14. The antibody according to claim 11, wherein the antibody is of human IgG1 subclass or is of human IgG4 subclass.

15. The antibody according to claim 11, wherein the antibody is a chimeric, single chain, multispecific, or humanized antibody.

16. A pharmaceutical composition comprising the antibody according to claim 15.

17. The antibody according to claim 11, wherein the antibody is an antigen binding fragment.

18. A pharmaceutical composition comprising the antibody according to claim 17.

19. The antibody according to claim 11, wherein the antibody is a monoclonal antibody.

20. A pharmaceutical composition comprising the antibody according to claim 19.

* * * * *