

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 2003210400 B8**

(54) Title
Form of presentation for 3-[(2-[[4-hexyloxycarbonylamino-imino-methyl]-phenylamino]-methyl]-1-methyl-1H-benzimidazol-5-carbonyl)-pyridin-2-yl-amino] propionic acid ethyl ester to be administered orally

(51) International Patent Classification(s)
A61K 31/4439 (2006.01) **A61K 47/38** (2006.01)
A61K 9/48 (2006.01) **A61P 7/02** (2006.01)
A61K 47/12 (2006.01) **A61P 43/00** (2006.01)
A61K 47/16 (2006.01) **C07D 401/12** (2006.01)
A61K 47/32 (2006.01)

(21) Application No: **2003210400** (22) Date of Filing: **2003.03.03**

(87) WIPO No: **WO03/074056**

(30) Priority Data

(31) Number	(32) Date	(33) Country
102 45 624.0	2002.09.30	DE
102 09 985.5	2002.03.07	DE

(43) Publication Date: **2003.09.16**

(43) Publication Journal Date: **2003.10.23**

(44) Accepted Journal Date: **2008.08.07**

(48) Corrigenda Journal Date: **2008.09.25**

(71) Applicant(s)
Boehringer Ingelheim Pharma GmbH & Co. KG

(72) Inventor(s)
Brauns, Ulrich;Hael, Norbert

(74) Agent / Attorney
Davies Collison Cave, Level 15 1 Nicholson Street, Melbourne, VIC, 3000

(56) Related Art
WO 1998/037075

(12) NACH DEM VERTRAG ÜBER DIE INTERNATIONALE ZUSAMMENARBEIT AUF DEM GEBIET DES
PATENTWESENS (PCT) VERÖFFENTLICHTE INTERNATIONALE ANMELDUNG

(19) Weltorganisation für geistiges Eigentum
Internationales Büro



(43) Internationales Veröffentlichungsdatum
12. September 2003 (12.09.2003)

PCT

(10) Internationale Veröffentlichungsnummer
WO 03/074056 A1

(51) Internationale Patentklassifikation⁷: A61K 31/4439,
A61P 7/02, C07D 235/14

(21) Internationales Aktenzeichen: PCT/EP03/02141

(22) Internationales Anmeldedatum:
3. März 2003 (03.03.2003)

(25) Einreichungssprache: Deutsch

(26) Veröffentlichungssprache: Deutsch

(30) Angaben zur Priorität:
102 09 985.5 7. März 2002 (07.03.2002) DE
102 45 624.0 30. September 2002 (30.09.2002) DE

(81) Bestimmungsstaaten (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SI, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Bestimmungsstaaten (*regional*): ARIPO-Patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), eurasisches Patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), europäisches Patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI-Patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Anmelder (*für alle Bestimmungsstaaten mit Ausnahme von US*): BOEHRINGER INGELHEIM PHARMA GMBH & CO. KG [DE/DE]; Binger Strasse 173, 55216 Ingelheim am Rhein (DE).

(72) Erfinder; und

(75) Erfinder/Anmelder (*nur für US*): BRAUNS, Ulrich [DE/DE]; Alleenstrasse 54, 88400 Biberach (DE).

(74) Gemeinsamer Vertreter: BOEHRINGER INGELHEIM PHARMA GMBH & CO. KG; Binger Strasse 173, 55216 Ingelheim am Rhein (DE).

Veröffentlicht:

mit internationalem Recherchenbericht

Zur Erklärung der Zweibuchstaben-Codes und der anderen Abkürzungen wird auf die Erklärungen ("Guidance Notes on Codes and Abbreviations") am Anfang jeder regulären Ausgabe der PCT-Gazette verwiesen.

(54) Title: FORM OF PRESENTATION FOR 3-[(2-[[4-(HEXYLOXYCARBONYLAMINO-IMINO-METHYL)-PHENYLAMINO]-METHYL]-1-METHYL-1H-BENZIMIDAZOL-5-CARBONYL)-PYRIDIN-2-YL-AMINO] PROPIONIC ACID ETHYL ESTER TO BE ADMINISTERED ORALLY

(54) Bezeichnung: ORAL ZU APPLIZIERENDE DARREICHUNGSFORM FÜR 3-[(2-[[4-(HEXYLOXYCARBONYLAMINO-IMINO-METHYL)-PHENYLAMINO]-METHYL]-1-METHYL-1H-BENZIMIDAZOL-5-CARBONYL)-PYRIDIN-2-YL-AMINO]-PROPIONSÄURE-ETHYLESTER UND DESSEN SALZE

(57) Abstract: The invention relates to a novel, orally administered form of presentation for the agent 3-[(2-[[4-(hexyloxy-carbonylamino-imino-methyl)-phenylamino]-methyl]-1-methyl-1H-benzimidazol-5-carbonyl)-pyridin-2-yl-amino] propionic acid ethyl ester and the pharmaceutically compatible salts thereof.


(57) Zusammenfassung: Die Erfindung betrifft eine neue, oral zu applizierende Darreichungsform für den Wirkstoff 3-[(2-[[4-(Hexyloxy-carbonylamino-imino-methyl)-phenylamino]-methyl]-1-methyl-1H-benzimidazol-5-carbonyl)-pyridin-2-yl-amino]-propionsäure-ethylester und dessen pharmakologisch verträglichen Salze.

WO 03/074056 A1


80433pct.210

Administration form for the oral application of 3-[(2-[(4-(hexyloxycarbonylamino-imino-methyl)-phenylamino)-methyl]-1-methyl-1*H*-benzimidazol-5-carbonyl)-pyridin-2-yl-amino]-propionic acid ethyl ester and the salts thereof

The invention relates to an administration form for the oral application of the active substance ethyl 3-[(2-[(4-(hexyloxycarbonylamino-imino-methyl)-phenylamino)-methyl]-1-methyl-1*H*-benzimidazole-5-carbonyl)-pyridin-2-yl-amino]-propionate and the pharmacologically acceptable salts thereof. This active substance having the chemical formula



is already known from WO 98/37075, which discloses compounds with a thrombin-inhibiting effect and the effect of prolonging the thrombin time, under the name 1-methyl-2-[*N*-[4-(*N*-*n*-hexyloxycarbonylamidino)phenyl]-amino-methyl]-benzimidazole-5-yl-carboxylic acid-*N*-(2-pyridyl)-*N*-(2-ethoxycarbonyl-ethyl)-amides. The compound of formula I is a double prodrug of the compound



- 2 -

i.e. the compound of formula I is only converted into the active compound, namely the compound of formula II, after entering the body. The main indication for the compound of chemical formula I is the post-operative prevention of deep-vein thrombosis.

The aim of the invention is to provide an improved formulation for oral use of the compound of formula I (which is also referred to hereinafter as the "active substance").

Surprisingly it has now been found that the use of pharmaceutically acceptable organic acids with a water solubility of $> 1 \text{ g} / 250 \text{ ml}$ at 20°C , preferably $> 1 \text{ g} / 160 \text{ ml}$ at 25°C , in solid oral preparations leads to a significantly improved formulation of ethyl 3-[(2-[[4-(hexyloxycarbonylamino-imino-methyl)-phenylamino]-methyl]-1-methyl-1*H*-benzimidazole-5-carbonyl)-pyridin-2-yl-amino]-propionate as well as the pharmaceutically acceptable salts thereof.

Accordingly, in a first aspect, the invention provides a pharmaceutical composition for oral administration comprising at least

- a) a substantially spherical core material which consists of or contains one or more pharmaceutically acceptable organic acids with a water solubility of $> 1 \text{ g} / 250 \text{ ml}$ at 20°C and
- b) an active substance layer containing binder and optionally separating agent, which encloses the core material, wherein said active substance is ethyl 3-[(2-[[4-(hexyloxycarbonylamino-imino-methyl)-phenylamino]-methyl]-1-methyl-1*H*-benzimidazole-5-carbonyl)-pyridin-2-yl-amino]-propionate or one of the pharmaceutically acceptable salts thereof.

In a second aspect, the invention provides a process for preparing a pharmaceutical composition for oral administration containing ethyl 3-[(2-[[4-(hexyloxycarbonylamino-imino-methyl)-phenylamino]-methyl]-1-methyl-1*H*-benzimidazole-5-carbonyl)-pyridin-2-yl-amino]-propionate or one of the physiologically acceptable salts thereof, comprising the steps of:

- 2A -

- a) forming the core material from one or more pharmaceutically acceptable organic acid(s) with a water solubility of > 1 g / 250 ml at 20°C, optionally with the addition of binders or other technological adjuvants, by pan methods, on pelleting plates or by extrusion / spheronisation,
- b) applying to the core material an insulating layer consisting of one or more water-soluble, pharmaceutically acceptable polymers optionally with the addition of plasticisers, separating agents and/or pigments,
- c) applying the active substance from a dispersion which contains a binder and optionally separating agents and simultaneously and/or subsequently drying to eliminate the dispersant,
- d) optionally applying a coating of film-forming agents, plasticisers and optionally pigments and
- e) packing the active substance-containing pellets thus obtained into hard capsules.

In a third aspect, the invention provides Ethyl 3-[(2-[[4-(hexyloxycarbonylamino-imino-methyl)-phenylamino]-methyl]-1-methyl-1*H*-benzimidazole-5-carbonyl)-pyridin-2-yl-amino]-propionate methanesulphonate.

A fourth aspect of the invention provides a pharmaceutical composition comprising the methanesulfonate salt of ethyl 3-[(2-[[4-(hexyloxycarbonylamino-imino-methyl)-phenylamino]-methyl]-1-methyl-1*H*-benzimidazole-5-carbonyl)-pyridin-2-yl-amino]-propionate.

Pharmaceutically suitable acids for the purposes of this invention are for example tartaric acid, fumaric acid, succinic acid, citric acid, malic acid, glutamic acid and aspartic acid including the hydrates and acid salts thereof. Particularly suitable for the purposes of this invention are tartaric acid, fumaric acid, succinic acid and citric acid.

2003210400 21 Jul 2008

- 2B -

A preferred embodiment of the invention is a multiparticulate preparation in which the individual particles are constructed as in Figure 1.

Figure 1 shows the diagrammatic structure of the pharmaceutical composition by means of a section through a pellet suitable for the preparation of the pharmaceutical composition according to the invention. The roughly bead-shaped / spherical core region of this pellet contains/consists of the pharmaceutically acceptable organic acid. Then follows a layer, the so-called insulating layer, which separates the acid core from the layer containing the active substance. The insulating layer is in turn surrounded by the equally spherically shaped layer of active substance which may in turn be enclosed in a coating which increases the abrasion resistance and shelf life of the pellets.

One advantage of the formulation thus constructed is the spatial separation of the organic acid and active substance by the insulating layer. A further advantage of the construction of the pellets as described above is the fact that the organic acid does not go into solution until after the preparation has been taken and then produces an acid microclimate in which the active substance can dissolve.

The core material used is a pharmaceutically acceptable organic acid with a water solubility of $> 1 \text{ g} / 250 \text{ ml}$ at 20°C , such as e.g. tartaric acid, fumaric acid, succinic acid, citric acid, malic acid, glutamic acid and aspartic acid including the hydrates and acid salts thereof, to which a small amount of 1 to 10 % by weight, preferably 3 to 6 % by weight of a suitable binder is optionally added. The use of a binder may be necessary, for example, if the starting acids are produced by a pan build-up process. If the method used is extrusion or spheronisation, other technological adjuvants such as microcrystalline cellulose will be needed instead of binders. It is also possible to use pure (100 %) acid as the starting material if it can be obtained in a sufficiently narrow range of particle sizes. The pharmaceutically acceptable organic acids used are preferably tartaric acid, fumaric acid, succinic acid or citric acid; tartaric acid is particularly preferred. As binder, it is possible to use gum arabic or a partially or totally synthetic polymer selected from among the hydroxypropyl-celluloses, hydroxypropylmethylcelluloses, methylcelluloses, hydroxyethylcelluloses, carboxymethylcelluloses, polyvinylpyrrolidone, the copolymers of N-vinylpyrrolidone and vinyl acetate, or combinations of these polymers; gum arabic is preferred. The spherical core material preferably has an average diameter of 0.4 – 1.5 mm. The content of the pharmaceutically acceptable organic acid is usually between 30 and 100 % in the core material, corresponding to an amount of between 20 and 90 %, preferably between 20 and 80 % in the finished pellet (i.e. in the pharmaceutical composition).

To increase the durability of the finished product it is advantageous to coat the core material before the application of the active substance with an insulating layer based on a water-soluble, pharmaceutically acceptable polymer.

Examples of such water-soluble polymers include for example gum arabic or a partially or totally synthetic polymer selected from among the hydroxypropyl-celluloses, hydroxypropylmethylcelluloses, methylcelluloses, hydroxyethylcelluloses, carboxymethylcelluloses, polyvinylpyrrolidone, the copolymers of N-vinylpyrrolidone and vinyl acetate, or combinations of these polymers. Gum arabic or a hydroxypropylmethylcellulose is preferably used. If desired, the coating with the water-soluble, pharmaceutically acceptable polymer may be carried out with the addition of suitable plasticisers, separating agents and pigments, such as for example triethylcitrate, tributylcitrate, triacetin, polyethyleneglycols (plasticisers), talc, silicic acid (separating agents), titanium dioxide or iron oxide pigments (pigments).

The active substance layer contains the active substance ethyl 3-[(2-[[4-(hexyloxycarbonylamino-imino-methyl)-phenylamino]-methyl]-1-methyl-1H-benzimidazole-5-carbonyl)-pyridin-2-yl-amino]-propionate (BIBR 1048) or one of the pharmaceutically acceptable salts thereof as well as binders and optionally separating agents. A preferred salt of the active substance is the mesylate (methanesulphonate) of the compound of formula I. Suitable binders include for example hydroxypropylcellulose, hydroxypropylmethylcellulose, methylcellulose, hydroxyethylcellulose, carboxymethylcellulose, polyvinylpyrrolidone, copolymers of N-vinylpyrrolidone and vinyl acetate or combinations of these polymers. Preferably, hydroxypropylcellulose or copolymers of N-vinylpyrrolidone and vinyl acetate are used. The addition of separating agents such as e.g. talc or silicic acid serves to prevent the particles from aggregating during the process. The active substance content is 5 to 60 %, preferably 10 to 50 % of the pharmaceutical composition.

The optional outermost layer, which serves to reduce any increased abrasion during packing into capsules and/or to increase the shelf life, consists of pharmaceutically conventional film-forming agents, plasticisers and optionally pigments. Suitable film-forming agents include for example hydroxypropylcellulose, hydroxypropylmethylcellulose, methylcellulose, polymers and copolymers of acrylic and methacrylic acid and the esters thereof, or combinations of these polymers. Suitable plasticisers include *inter alia*

triethylcitrate, tributylcitrate, triacetin or polyethyleneglycols. The pigments used may be e.g. titanium dioxide or iron oxide pigments. Preferably, the outer coating consists of hydroxypropylmethylcellulose and/or methylcellulose, optionally with the addition of polyethyleneglycols as plasticisers.

The pellets may be prepared by the method described hereinafter:

The acid-containing core material consists either of crystals of the particular organic acid used or, more advantageously, of roughly spherical particles of the desired size containing a large amount of organic acid, which can be produced by methods known and established in pharmaceutical technology. The core material may be produced, in particular, by pan methods, on pelleting plates or by extrusion/spheronisation. Then the core material thus obtained may be divided into fractions of the desired diameter by screening. Suitable core material has an average diameter of 0.4 to 1.5 mm, preferably 0.6 to 0.8 mm.

First, the insulating layer is applied to this acid-containing core material. This can be done by conventional methods, e.g. by applying an aqueous dispersion of the water-soluble, pharmaceutically acceptable polymer, optionally with the addition of plasticisers, separating agents and/or pigments, in a fluidised bed, in coating pans or in conventional film coating apparatus. If necessary the product can then be screened again.

Then the active substance is applied from a dispersion containing binder and optionally separating agent. The volatile dispersant is removed during or after the process by drying. Suitable binders in the dispersion may be for example hydroxypropylcellulose, hydroxypropylmethylcellulose, methylcellulose, hydroxyethylcellulose, carboxymethylcellulose, polyvinylpyrrolidone, copolymers of N-vinylpyrrolidone and vinyl acetate or combinations of these polymers. Preferably, hydroxypropylcellulose or copolymers of N-vinylpyrrolidone and vinyl acetate are used. Suitable separating agents include e.g. talc or silicic acid; preferably, talc is used. The dispersants may

be for example ethanol, 2-propanol, acetone or mixtures of these solvents with one another or with water, preferably 2-propanol. The application of active substance to the core material may be carried out by established methods known in pharmaceutical technology, e.g. in coating pans, conventional film coating apparatus or by the fluidised bed method. Then a further screening process may be carried out.

To reduce any increased abrasion during transfer into capsules or to increase the shelf life the system may finally be coated with a coating of a pharmaceutically conventional film forming agent, plasticiser and optionally pigment. This may be done by conventional methods as mentioned earlier in the description of the application of the insulating layer.

When core material with an average diameter of 0.4 – 1.5 mm is used, the process described above produces pellets containing active substance, which can then be packed into hard capsules, for example. To do this, a number of these units corresponding to the required dosage are packed into hard capsules in a standard capsule filling machine. Suitable hard capsules include, for example, hard gelatine capsules or hard capsules of hydroxypropylmethylcellulose (HPMC); HPMC capsules are preferred. The active substance content of the pharmaceutical composition is 5 to 60 %, preferably 10 to 50 %; the content of the pharmaceutically acceptable organic acid is usually between 20 and 90 %, preferably between 20 and 80 %.

Unless otherwise stated, percentages specified are always percent by weight. All the data on the active substance content relate to the active substance base of formula I (not to a specific salt) unless otherwise stated.

Clinical trials

In preliminary tests on test subjects with conventional tablets containing the compound of formula I it had been established that highly variable plasma levels occurred, with individual cases of malabsorption. The variability of the plasma level patterns is significantly lower after the administration of the

compound of formula I as an orally administered solution; there were no cases of malabsorption under these circumstances.

Tests have shown that the compound of formula I dissolves relatively well in water at low pH levels, whereas at pH levels above 5 in accordance with the definition of the European Pharmacopoeia it is virtually insoluble. Therefore the volunteers in one branch of the clinical trials were given pantoprazole, which serves to produce an elevated gastric pH.

For example, the pharmaceutical compositions according to Examples 1 and 2 were tested for their bioavailability by comparison with a conventional tablet.

To do this, the formulation prepared according to Example 1 containing 50 mg of active substance base per capsule was clinically tested for its bioavailability on a total of 15 volunteers. In one branch of the treatment, the volunteers were given the composition by mouth (= orally) on an empty stomach without any pre-treatment. In another branch of the treatment the same volunteers were pre-treated, prior to the oral administration of the composition, with 40 mg of pantoprazole b.i.d. (= twice a day) for three days by mouth to increase the gastric pH; the treatment with pantoprazole was continued during the administration of the formulation according to the invention.

The degree of absorption was determined by measuring the quantity of active metabolite of formula II excreted in the urine.

The relative bioavailability after pre-treatment with pantoprazole was 94% on average compared with administration without any pre-treatment.

Under comparable conditions of administration, the relative bioavailability (based on the area under the plasma concentration/time curve) of a tablet containing 50 mg of active substance, developed and produced according to the prior art and containing no water-soluble organic acid, after corresponding pre-treatment with pantoprazole, is 18 %. The following list shows the precise composition of the tablet used:

	Ingredient	mg/tablet
Core	mesylate of the compound of form. I	57.7
	lactose monohydrate	58.0
	microcrystalline cellulose	48.3
	crospovidone	3.4
	magnesium stearate	2.6
Film coating	polyethyleneglycol 6000	0.56
	titanium dioxide	0.80
	talc	0.64
	hydroxypropylmethylcellulose	1.92
	iron oxide yellow	0.08
	Total	174.0

The relative bioavailability was thus improved by about a factor of 5 by using the formulation according to the invention.

The formulation prepared according to Example 2 containing 50 mg of active substance base per capsule was also clinically tested for its bioavailability on a total of 15 volunteers. In one branch of the treatment, the volunteers were given the composition by mouth on an empty stomach without any pre-treatment. In another branch of the treatment the same volunteers were pre-treated, prior to the oral administration of the composition, with 40 mg of pantoprazole b.i.d. for three days by mouth to increase the gastric pH; the treatment with pantoprazole was continued during the administration of the formulation according to the invention.

The degree of absorption was determined by measuring the quantity of the active metabolite of formula II excreted in the urine.

The relative bioavailability after pre-treatment with pantoprazole was 76% on average compared with administration without any pre-treatment.

Under comparable conditions of administration, the relative bioavailability (based on the area under the plasma concentration/ time curve) of a tablet containing 50 mg of active substance, developed and produced according to the prior art and containing no water-soluble organic acid, after

corresponding pre-treatment with pantoprazole, is 18 %. The following list shows the precise composition of the tablet used:

	Ingredient	mg/tablet
Core	mesylate of the compound of form. I	57.7
	lactose monohydrate	58.0
	microcrystalline cellulose	48.3
	crospovidone	3.4
	magnesium stearate	2.6
Film coating	polyethyleneglycol 6000	0.56
	titanium dioxide	0.80
	talc	0.64
	hydroxypropylmethylcellulose	1.92
	iron oxide yellow	0.08
	Total	174.0

The relative bioavailability of the active substance compared with conventional formulations was thus improved by about a factor of 4 by using the formulation according to the invention. The bioavailability of the two formulations according to the invention compared with the tablet described above with and without the simultaneous administration of pantoprazole is graphically illustrated in Figure 2.

The clinical trials show another advantage of the preparation according to the invention containing the compound of formula I, which is that it ensures adequate bioavailability of the active substance, better than that of a conventional pharmaceutical preparation and largely independent of the gastric pH, it reduces fluctuations in the bioavailability of the active substance and it prevents malabsorption. Another advantageous property of the pharmaceutical composition according to the invention is the fact that it is suitable for all patients, i.e. including those in whom the gastric pH is increased by normal physiological variability, by disease or by co-medication with drugs which raise the gastric pH.

The dosage for oral use is expediently 25 to 300 mg of the active substance base (per capsule), preferably 50 to 200 mg, most preferably 75 to 150 mg of the active substance base, in each case once or twice a day.

The preferred ratio of acid to active substance is about 0.9 : 1 to about 4 : 1, most preferably between about 1:1 and 3:1. Preferably, at least one equivalent of acid is used per mol of the compound of formula I. The upper limit of about 4:1 (acid to active substance) is generally determined by the maximum acceptable size of the preparation in the desired dosages (number of pellets per capsule).

The Examples that follow are intended to illustrate the invention:

Example 1

	percentage composition				per capsule [mg]	per capsule [mg]
	core material	insulating layer	active substance layer	total		
tartaric acid	61.3	-	-	61.3	176.7	353.4
gum arabic	3.1	2.8		5.9	17.0	34.0
talc	-	5.6	3.2	8.8	25.4	50.7
hydroxypropylcellulose	-	-	4.0	4.0	11.5	23.1
active substance (mesylate of the compound of formula I)	-	-	20.0	20.0	57.7*	115.3**
total				100.0	288.3	576.5

*) corresponds to 50 mg of the compound of formula 1 (active substance base)

***) corresponds to 100 mg of the compound of formula 1 (active substance base)

a) Production of core material containing tartaric acid

Composition:

gum arabic 1 part by weight
tartaric acid 20 parts by weight

1 part by weight of gum arabic is dissolved in 4 parts by weight of purified water at 50°C with stirring. Then 5 parts by weight of tartaric acid are dissolved in this solution with stirring.

8.3 parts by weight of tartaric acid crystals with an average particle size of 0.4 to 0.6 mm are placed in a suitable coating apparatus fitted with an air inlet and exhaust, and the pan is set in rotation. At an air inlet temperature of 60° – 80°C the tartaric acid crystals are sprayed at intervals with the solution of

tartaric acid and gum arabic and sprinkled with a total of 6.7 parts by weight of powdered tartaric acid, so that roughly spherical particles are formed.

The spherical tartaric acid core material is then dried in the rotating pan at an air inlet temperature of 60° - 80° C.

The core material is fractionated using a tumbler screening machine with perforated plates with a nominal mesh size of 0.6 and 0.8 mm. The product fraction between 0.6 and 0.8 mm is used in the rest of the process.

b) Insulation of the core material containing tartaric acid

Composition:

core material containing tartaric acid	23 parts by weight
gum arabic	1 part by weight
talc	2 parts by weight

1 part by weight of gum arabic is dissolved in a mixture of 6.7 parts by weight of 96 % ethanol and 13.5 parts by weight of purified water with stirring. Then 2 parts by weight of talc are dispersed in the solution with stirring.

In a fluidised bed processing apparatus, 23 parts by weight of core material containing tartaric acid are sprayed at an air inlet temperature of 35° – 40° C with the dispersion of gum arabic and talc by the under-bed spraying process.

The insulated core material containing tartaric acid is then dried in the circulating air drier at 40°C for 8 hours.

To remove any lumps the dried insulated core material containing tartaric acid is screened through a screen with a nominal mesh size of 1.0 mm. The fraction of material with a particle size of < 1 mm is further processed.

c) Production of the active substance layer

Composition:

insulated core material containing tartaric acid	91 parts by weight
hydroxypropylcellulose	5 parts by weight
talc	4 parts by weight
active substance (mesylate of BIBR 1048)	25 parts by weight

Hydroxypropylcellulose is dissolved in 168 parts by weight of 2-propanol with stirring and then the active substance and talc are dispersed in this solution with stirring.

In a fluidised bed processing apparatus, 91 parts by weight of insulated core material containing tartaric acid are sprayed at an air inlet temperature of 20° - 30°C with the dispersion containing the active substance by the under-bed spraying process.

The pellets containing the active substance are then dried in the circulating air drier at 35°C for 8 hours.

To remove any lumps the pellets containing the active substance are screened through a screen with a nominal mesh size of 1.25 mm. The fraction of material with a particle size of < 1.25 mm is further processed.

d) Packing into capsules

A quantity of active substance pellets containing in each case 50 or 100 mg of active substance base is packed into size 1 or size 0 elongated hard gelatine capsules or HPMC capsules by means of a capsule filling machine.

Example 2

	percentage composition				per capsule [mg]	per capsule [mg]
	core material	insulating layer	active substance layer	total		
tartaric acid	38.5	-	-	38.5	55.5	166.5
gum arabic	1.9	1.7		3.6	5.2	15.6
talc	-	3.5	6.4	9.9	14.3	42.8
hydroxypropylcellulose	-	-	8.0	8.0	11.5	34.6
active substance (mesylate of the compound of formula I)	-	-	40.0	40.0	57.7*	173.0**
total				100.0	144.2	432.5

*) corresponds to 50 mg of the compound of formula 1 (active substance base)

***) corresponds to 150 mg of the compound of formula 1 (active substance base)

a) Production of core material containing tartaric acid

Composition:

gum arabic 1 part by weight
tartaric acid 20 parts by weight

1 part by weight of gum arabic is dissolved in 4 parts by weight of purified water at 50°C with stirring. Then 5 parts by weight of tartaric acid are dissolved in this solution with stirring.

8.3 parts by weight of tartaric acid crystals with an average particle size of 0.4 to 0.6 mm are placed in a suitable coating apparatus fitted with an air inlet and exhaust, and the pan is set in rotation. At an air inlet temperature of 60° – 80°C the tartaric acid crystals are sprayed at intervals with the solution of

tartaric acid and gum arabic and sprinkled with a total of 6.7 parts by weight of powdered tartaric acid, so that roughly spherical particles are formed.

The spherical tartaric acid core material is then dried in the rotating pan at an air inlet temperature of 60° - 80° C.

The core material is fractionated using a tumbler screening machine with perforated plates with a nominal mesh size of 0.6 and 0.8 mm. The product fraction between 0.6 and 0.8 mm is used in the rest of the process.

b) Insulation of the core material containing tartaric acid

Composition:

core material containing tartaric acid	23 parts by weight
gum arabic	1 part by weight
talc	2 parts by weight

1 part by weight of gum arabic is dissolved in a mixture of 6.7 parts by weight of 96 % ethanol and 13.5 parts by weight of purified water with stirring. Then 2 parts by weight of talc are dispersed in the solution with stirring.

In a fluidised bed processing apparatus, 23 parts by weight of core material containing tartaric acid are sprayed at an air inlet temperature of 35° – 40°C with the dispersion of gum arabic and talc by the under-bed spraying process.

The insulated core material containing tartaric acid is then dried in the circulating air drier at 40°C for 8 hours.

To remove any lumps the dried insulated core material containing tartaric acid is screened through a screen with a nominal mesh size of 1.0 mm. The fraction of material with a particle size of < 1 mm is further processed.

c) Production of the active substance layer

Composition:

insulated core material containing tartaric acid	57 parts by weight
hydroxypropylcellulose	10 parts by weight
talc	8 parts by weight
active substance (mesylate of BIBR 1048)	50 parts by weight

Hydroxypropylcellulose is dissolved in 335 parts by weight of 2-propanol with stirring and then the active substance and talc are dispersed in this solution with stirring.

In a fluidised bed processing apparatus, 91 parts by weight of insulated core material containing tartaric acid are sprayed at an air inlet temperature of 20° - 30°C with the dispersion containing the active substance by the under-bed spraying process.

The pellets containing the active substance are then dried in the circulating air drier at 35°C for 8 hours.


To remove any lumps the pellets containing the active substance are screened through a screen with a nominal mesh size of 1.25 mm. The fraction of material with a particle size of < 1.25 mm is further processed.

d) Packing into capsules

A quantity of active substance pellets containing in each case 50 or 150 mg of active substance base is packed into size 2 or size 0 hard gelatine capsules or HPMC capsules by means of a capsule filling machine.

Example 3

Preparation of ethyl 3-[(2-[[4-(hexyloxycarbonylamino-imino-methyl)-phenylamino]-methyl]-1-methyl-1*H*-benzimidazole-5-carbonyl)-pyridin-2-yl-amino]-propionate methanesulphonate



A solution of 5.0 mmol of methanesulphonic acid in 25 ml ethyl acetate was added dropwise, with stirring, to a solution of 3139 mg (5.0 mmol) of ethyl 3-[(2-[[4-(hexyloxycarbonylamino-imino-methyl)-phenylamino]-methyl]-1-methyl-1*H*-benzimidazole-5-carbonyl)-pyridin-2-yl-amino]-propionate base (prepared as described in WO 98/37075) in 250 ml ethyl acetate, at ambient temperature. After a few minutes the product began to crystallise out. It was stirred for another hour at ambient temperature and then for one more hour while cooling with ice, the precipitate was suction filtered, washed with about 50 ml of ethyl acetate and 50 ml of diethyl ether and dried at 50°C in a circulating air drier.

Yield: 94% of theory

melting point: 178 - 179°C

$C_{34}H_{41}N_7O_5 \times CH_4SO_3$ (723.86)

Elemental analysis: calc.:	C 58.07%	H 6.27%	N 13.55%	S 4.43%
found:	58.11%	6.30%	13.50%	4.48%

2003210400 21 Jul 2008

- 17A -

Throughout this specification and the claims which follow, unless the context requires otherwise, the word "comprise", and variations such as "comprises" and "comprising", will be understood to imply the inclusion of a stated integer or step or group of integers or steps but not the exclusion of any other integer or step or group of integers or steps.

The reference in this specification to any prior publication (or information derived from it), or to any matter which is known, is not, and should not be taken as an acknowledgment or admission or any form of suggestion that that prior publication (or information derived from it) or known matter forms part of the common general knowledge in the field of endeavour to which this specification relates.

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. Pharmaceutical composition for oral administration comprising at least
 - a) a substantially spherical core material which consists of or contains one or
5 more pharmaceutically acceptable organic acids with a water solubility of > 1 g / 250 ml at 20° C and
 - b) an active substance layer containing binder and optionally separating agent, which encloses the core material, wherein said active substance is ethyl 3-[(2-
10 {4-(hexyloxycarbonylamino-imino-methyl)-phenylamino]-methyl}-1-methyl-1*H*-benzimidazole-5-carbonyl)-pyridin-2-yl-amino]-propionate or one of the pharmaceutically acceptable salts thereof.
2. Pharmaceutical composition according to claim 1, wherein the
15 pharmaceutically acceptable organic acid is tartaric acid, fumaric acid, succinic acid, citric acid, malic acid, glutamic acid or aspartic acid or one of the hydrates or acid salts thereof.
3. Pharmaceutical composition according to claim 2, characterised in that the
20 pharmaceutically acceptable organic acid is tartaric acid, fumaric acid, citric acid or succinic acid.
4. Pharmaceutical composition according to claim 3, characterised in that the pharmaceutically acceptable organic acid is tartaric acid.
- 25 5. Pharmaceutical composition according to any one of claims 1 to 4, wherein the content of ethyl 3-[(2-[[4-(hexyloxycarbonylamino-imino-methyl)-phenylamino]-methyl]-1-methyl-1*H*-benzimidazole-5-carbonyl)-pyridin-2-yl-amino]-propionate or the salts thereof in the pharmaceutical composition is 5 to 60 %.
- 30 6. Pharmaceutical composition according to any one of claims 1 to 5, wherein the content of pharmaceutically acceptable organic acid is 20 to 90 %.

7. Pharmaceutical composition according to any one of claims 1 to 6, wherein the binder is selected from the group of hydroxypropylcelluloses, hydroxypropylmethylcelluloses, methylcelluloses, hydroxyethylcelluloses, carboxymethylcelluloses, the polyvinylpyrrolidones, the copolymers of N-vinylpyrrolidone and vinyl acetate or combinations of these polymers.
8. Pharmaceutical composition according to any one of claims 1 to 6, wherein the core material has an average particle size of 0.4 to 1.5 mm.
9. Pharmaceutical composition according to any one of claims 1 to 6, wherein the core material and the active substance layer are separated from one another by an insulating layer consisting of a water-soluble polymer, optionally with the addition of suitable plasticisers, separating agents and pigments.
10. Pharmaceutical composition according to claim 9, wherein the water-soluble polymer consists of gum arabic or a partially or totally synthetic polymer from the group of hydroxypropylcelluloses, hydroxypropylmethylcelluloses, methylcelluloses, hydroxyethylcelluloses, carboxymethylcelluloses, the polyvinylpyrrolidones, the copolymers of N-vinylpyrrolidone and vinyl acetate, or of combinations of these polymers.
11. Pharmaceutical composition according to any one of claims 1 to 6, wherein the product containing the active substance is packed into hard capsules.
12. Pharmaceutical composition according to claim 11, wherein the composition containing the active substance is packed into hydroxypropylmethylcellulose capsules.
13. Pharmaceutical composition according to any one of claims 1 to 12, wherein ethyl 3-[(2-[[4-(hexyloxycarbonylamino-imino-methyl)-phenylamino]-methyl]-1-

methyl-1*H*-benzimidazole-5-carbonyl)-pyridin-2-yl-amino]-propionate mesylate is used as the active substance.

14. Process for preparing a pharmaceutical composition for oral administration
5 containing ethyl 3-[(2-[[4-(hexyloxycarbonylamino-imino-methyl)-phenylamino]-methyl]-1-methyl-1*H*-benzimidazole-5-carbonyl)-pyridin-2-yl-amino]-propionate or one of the physiologically acceptable salts thereof, comprising the steps of:
- 10 a) forming the core material from one or more pharmaceutically acceptable organic acid(s) with a water solubility of > 1 g / 250 ml at 20°C, optionally with the addition of binders or other technological adjuvants, by pan methods, on pelleting plates or by extrusion / spheronisation,
b) applying to the core material an insulating layer consisting of one or more water-soluble, pharmaceutically acceptable polymers optionally with the addition of plasticisers, separating agents and/or pigments,
15 c) applying the active substance from a dispersion which contains a binder and optionally separating agents and simultaneously and/or subsequently drying to eliminate the dispersant,
d) optionally applying a coating of film-forming agents, plasticisers and optionally pigments and
20 e) packing the active substance-containing pellets thus obtained into hard capsules.
15. Ethyl 3-[(2-[[4-(hexyloxycarbonylamino-imino-methyl)-phenylamino]-methyl]-1-methyl-1*H*-benzimidazole-5-carbonyl)-pyridin-2-yl-amino]-propionate
25 methanesulphonate.
16. A pharmaceutical composition comprising the methanesulfonate salt of ethyl 3-[(2-[[4-(hexyloxycarbonylamino-imino-methyl)-phenylamino]-methyl]-1-methyl-1*H*-benzimidazole-5-carbonyl)-pyridin-2-yl-amino]-propionate.
30

2003210400 21 Jul 2008

- 21 -

17. The pharmaceutical composition of claim 16 for oral administration further comprising one or more pharmaceutically acceptable organic acids with a water solubility of > 1 g / 250 ml at 20°C.
- 5 18. Pharmaceutical composition prepared according to the process of claim 17.
19. Pharmaceutical composition according to claim 1, 16 or 18; or a process according to claim 14, substantially as hereinbefore described and/or exemplified and/or illustrated in the Figures.

10

Figure 1:

Schematic structure of the pharmaceutical composition:




Figure 2:

