



(12) **EUROPEAN PATENT SPECIFICATION**

- (45) Date of publication and mention of the grant of the patent:  
**11.08.2004 Bulletin 2004/33**
- (21) Application number: **95904020.5**
- (22) Date of filing: **28.12.1994**
- (51) Int Cl.7: **C12N 15/82, C12N 9/02, C12N 5/10, A01H 1/00, A01H 5/00 // C12N9/00**
- (86) International application number:  
**PCT/JP1994/002288**
- (87) International publication number:  
**WO 1995/018222 (06.07.1995 Gazette 1995/29)**

(54) **GENE FOR FATTY ACID DESATURASE, VECTOR CONTAINING SAID GENE, PLANT CONTAINING SAID GENE TRANSFERRED THEREINTO, AND PROCESS FOR CREATING SAID PLANT**

GEN FÜR DIE FETTSÄURE-DESATURASE, BESAGTES GEN ENTHALTENDER VEKTOR, EINE PFLANZE, IN DIE BESAGTES GEN EINGEBRACHT IST, UND EIN VERFAHREN ZUR SCHAFFUNG BESAGTER PFLANZE

GENE POUR ACIDE GRAS-DESATURASE, VECTEUR CONTENANT LEDIT GENE, VEGETAL CONTENANT LEDIT GENE LUI AYANT ETE TRANSFERE, ET PROCEDE POUR CREER LEDIT VEGETAL

(84) Designated Contracting States:  
**AT BE CH DE DK ES FR GB GR IE IT LI LU MC NL PT SE**

(30) Priority: **28.12.1993 JP 35285893**

(43) Date of publication of application:  
**09.10.1996 Bulletin 1996/41**

(73) Proprietor: **KIRIN BEER KABUSHIKI KAISHA Chuo-Ku, Tokyo 104 (JP)**

(72) Inventors:  
• **NISHIZAWA, Osamu**  
**Kirin Beer Kabushiki Kaisha Kiban ku, Yokohama-shi, Kanagawa 236 (JP)**  
• **TOGURI, Toshihiro**  
**Kirin Beer Kabushiki Kaisha azawa-ku, Yokohama-shi, Kanagawa 236 (JP)**

(74) Representative: **Polz, Leo, Dr. et al Hoffmann Eitle**  
**Postfach 81 04 20**  
**81904 München (DE)**

(56) References cited:  
**EP-A- 0 550 162 EP-A- 0 561 569**  
**EP-A- 0 644 263 WO-A-91/13972**  
**NL-A- 9 002 130**

- **SHANKLIN J ET AL:**  
**"STEAROYL-ACYL-CARRIER-PROTEIN DESATURASE FROM HIGHER PLANTS IS STRUCTURALLY UNRELATED TO THE ANIMAL AND FUNGAL HOMOLOGS" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, vol. 88, no. 6, March 1991, pages 2510-2514, XP000651759**
- **ISHIZAKI-NISHIZAWA, O., ET AL. :**  
**"low-temperature resistance of higher plants is significantly enhanced by a nonspecific cyanobacterial desaturase" NATURE BIOTECHNOLOGY, vol. 14, August 1996, pages 1003-1006, XP002057130**
- **JOURNAL OF BIOLOGICAL CHEMISTRY, Vol. 269, No. 41, (1994), T. SAKAMOTO et al., "Delta, 9 Acyl-Lipid Desaturases of Cyanobacteria. Molecular Cloning and Substrate Specificities in Terms of Fatty Acids, sn-Positions, and Polar Head Groups", p. 25576-25580.**
- **JOURNAL OF BIOLOGICAL CHEMISTRY, Vol. 265, No. 33, (1990), J.E. STUKEY et al., "The OLE1 Gene of Saccharomyces Cerevisiae Encodes the Delta-9 Fatty Acid Desaturase and can be Functionally Replaced by the Rat StearoylCoA Desaturase Gene", p. 20144-20149.**

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

- WADA H. ET AL: 'THE DESA GENE OF THE CYANOBACTERIUM SYNECHOCYSTIS SP. STRAIN PCC6803 IS THE STRUCTURAL GENE FOR DELTA12 DESATURASE' JOURNAL OF BACTERIOLOGY, WASHINGTON, DC, US vol. 175, no. 18, 1993, pages 6056 - 6058
- HIGASHI S.; MURATA N.: 'AN IN VIVO STUDY OF SUBSTRATE SPECIFICITIES OF ACYL-LIPID DESATURASES AND ACYLTRANSFERASES IN LIPID SYNTHESIS IN SYNECHOCYSTIS PCC6803' PLANT PHYSIOLOGY, AMERICAN SOCIETY OF PLANT PHYSIOLOGISTS, ROCKVILLE, MD, US vol. 102, no. 4, 1993, pages 1275 - 1278

**Description**

## TECHNICAL FIELD


5 **[0001]** The present invention relates to genes encoding proteins having an activity of desaturating lipid-bound fatty acids at the  $\Delta 9$  position (hereinafter referred to as " $\Delta 9$  desaturases"), vectors containing said genes, plants transformed with said genes and process for creating said plants.

## BACKGROUND ART

10 **[0002]** The condition of membrane lipids which compose the biomembranes of organisms changes from liquid crystal to solid in accordance with the decrease in external temperatures. This change is called "phase separation". The phase separation involves the change in properties of biomembranes. It is believed that membrane lipids lose the selectivity of mass permeability in solid conditions, making it impossible for biomembranes to carry out their essential functions and that, as a result, cells receive an injury (low temperature injury).

15 **[0003]** The phase transition temperatures of membrane lipids, at which the condition of the membrane lipids changes from liquid crystal to solid, are chiefly dependent on the degree of unsaturation (the number of double bonds in carbon chains) of fatty acid acyl groups bound to lipids. A lipid molecular species in which two bound fatty acid acyl groups are both saturated fatty acid residues has a higher phase transition temperature than room temperature, whereas a  
20 lipid molecular species having at least one double bond in bound fatty acid acyl groups has a phase transition temperature below about 0°C (Santaren, J.F. et al., *Biochim. Biophys. Acta*, 687:231, 1982).

**[0004]** In general, the position of a double bond in a fatty acid is indicated after the symbol " $\Delta$ " by the number of carbons from the carboxyl terminus to the carbon having the double bond. The total number of double bonds is indicated after a colon following the total number of carbons. For example, linoleic acid is designated as 18:2  $\Delta 9,12$ , which is  
25 represented by the following structural formula:



30 In some cases, the position of a double bond is indicated after the symbol " $\omega$ " by the number of carbons from the methyl terminus of a fatty acid to the carbon having the double bond.

**[0005]** Among the membrane lipids of higher plants, only phosphatidylglycerol (PG) contains a relatively large number of saturated molecular species and it has been strongly suggested that the phase transition of PG is responsible for low temperature injury in plants (Murata, N. et al., *Plant Cell Physiol.*, 23:1071, 1982; Roughan, P.G., *Plant Physiol.*,  
35 77:740, 1985) and that the molecular species composition of PG is determined by the substrate specificity of glycerol-3-phosphate acyl transferase (hereinafter referred to as "ATase") present in chloroplasts (Frentzen, M. et al., *Eur. J. Biochem.*, 129:629, 1983; Murata, N., *Plant Cell Physiol.*, 24:81, 1983; Frentzen, M. et al., *Plant Cell Physiol.*, 28:1195, 1988).

**[0006]** Nishizawa et al. showed that if an ATase gene obtained from *Arabidopsis thaliana* Heynhold, a plant resistant to chilling, was introduced and expressed in tobacco, the content of saturated molecular species of PG decreased, thereby imparting a higher chilling resistance to the tobacco than when it was of a wild type (PCT/JP92/00024, 1992). However, the ATase exists originally in plants and even if a large amount of an exogenous ATase is expressed in plants, it will compete inevitably with the endogenous ATase and its effect is therefore likely to be diluted. For example, the content of saturated molecular species of PG was about 28% in the leaf of a clone which expressed the largest amount  
45 of ATase from *Arabidopsis thaliana* Heynhold out of the created tobacco transformants, which content was lower by about 8% than in the wild-type tobacco and higher by about 8% than in the wild-type *Arabidopsis thaliana* Heynhold (PCT/JP92/00024, 1992).

**[0007]** In general, the majority of acyl-ACP produced in plastids consists of 16:0-ACP and 18:1-ACP, and their proportions are believed to be equal. In some tissues, the proportions of 16:0-ACP and 18:0-ACP may be higher than that  
50 of 18:1-ACP (Toriyama, S. et al., *Plant Cell Physiol.*, 29:615, 1988). In these tissues, it may be difficult to reduce satisfactorily the content of saturated molecular species by using an exogenous ATase.

**[0008]** The composition of membrane lipids in photosynthetic cyanobacteria (blue-green algae) is similar to that of lipids in membrane systems composing higher plant's chloroplasts (Murata, N. et al., in "The Biochemistry of Plants", Academic Press, 1987). In blue-green algae, the degree of unsaturation of fatty acids bound to membrane lipids is  
55 controlled by enzymes capable of desaturating lipid-bound fatty acids. It is known that *Anacystis nidulans* (*Synechococcus* PCC 7942) which can introduce only one double bond into lipid-bound fatty acids is sensitive to chilling (Ono, T. et al., *Plant Physiol.*, 67:176, 1981), whereas *Synechocystis* PCC6803 which can introduce at least two double bonds is resistant to chilling (Wada, H. et al., *Plant Cell Physiol.*, 30:971, 1989).

[0009] All the desaturases of fatty acids in blue-green algae react with lipids as substrates to introduce a double bond into lipid-bound fatty acids. Therefore, a cis-type double bond can be introduced into fatty acids such as PG, SQDG, MGDG and DGDG in membrane lipids of blue-green algae, which are composed of 16:0/16:0- and 18:0/16:0-saturated molecular species (Murata, N. et al., in "The Biochemistry of Plants", Academic Press, 1987). In this respect, blue-green algae are very much different from higher plants that have fatty acid desaturases capable of introducing a double bond into stearyl-ACP (18:0-ACP) at the  $\Delta$  9 position and which never introduce a cis-type double bond into PG or SQDG after the synthesis of these lipids, which are composed of 16: 0/16:0-(and a little amount of 18:0/16:0-) as saturated molecular species.

[0010] At present, it is known that the introduction and expression of the  $\Delta$  12 desaturase gene of *Synechocystis* PCC6803 in *Anacystis nidulans* enables the production of 16:2  $\Delta$  9,12 which is not inherently present in the *Anacystis nidulans*, thereby imparting a chilling resistance to the *Anacystis nidulans* which is essentially chilling-sensitive (Wada, H. et al., Nature, 347:200, 1990).

[0011] Genes so far obtained for desaturases of blue-green algae include a  $\Delta$  6 desaturase gene (Reddy, A.S. et al., Plant Mol. Biol., 27:293, 1993) and a  $\Delta$  12 desaturase gene (Wada, H. et al., Nature, 347:200, 1990). However, the  $\Delta$  6 and  $\Delta$  12 desaturases cannot desaturate fatty acids at the  $\Delta$  6 and  $\Delta$  12 positions, respectively, unless a double bond is introduced at the  $\Delta$  9 position. Moreover,  $\Delta$  15 desaturase cannot desaturate fatty acids at the  $\Delta$  15 position unless the fatty acids are desaturated at both the  $\Delta$  9 and 12 positions. Hence, if the genes for enzymes capable of desaturating fatty acids at the  $\Delta$  9 position are introduced and expressed in higher plants, they should be able to reduce the content of saturated molecular species in higher plants and thereby impart a chilling resistance to the higher plants. However, no gene for enzymes capable of desaturating fatty acids at the  $\Delta$  9 position has been obtained until now.

[0012] WO 91/13972 describes the isolation of a plant  $\Delta$ -9 desaturase obtained from *C. tinctorius* and its use for the transformation of plants. It also discloses genes encoding for further higher plant  $\Delta$ -9 desaturases such as from *R. communites* and *B. campestris*. Said transformation may result in chill tolerance.

[0013] Therefore, an object of the present invention is to provide further genes for enzymes capable of desaturating fatty acids at the  $\Delta$  9 position and polynucleotides containing part of said genes.

[0014] Another object of the present invention is to provide vectors which contain said genes for enzymes capable of desaturating fatty acids at the  $\Delta$ -9 position or polynucleotides containing part of the genes.

[0015] A further object of the present invention is to provide plant cells and plants which are transformed with said genes for enzymes capable of desaturating fatty acids at the  $\Delta$ -9 position or polynucleotides containing part of the genes.

#### DISCLOSURE OF THE INVENTION

[0016] As a result of various studies conducted to achieve the above objects, the inventors succeeded in cloning a gene encoding a  $\Delta$ -9 desaturase from genomic DNA of blue-green alga belonging to the genus *Anacystis*, obtaining vector DNA incorporating the gene, transforming a plant cell with the vector DNA, differentiating the cell to regenerate the plant, thereby imparting a chilling resistance to the plant. The present invention was accomplished on the basis of these findings. The subject matters of the present invention are as follows:

- (1) Genes encoding proteins having an activity of desaturating lipidbound fatty acids at the  $\Delta$ -9 position.
- (2) The genes of (1) or polynucleotides containing part thereof, wherein the proteins having an activity of desaturating lipid-bound fatty acids at the  $\Delta$ -9 position contain the amino acid sequence substantially shown in SEQ ID NO: 4.
- (3) The genes of (1) or polynucleotides containing part of thereof, wherein the genes encoding proteins having an activity of desaturating lipid-bound fatty acids at the  $\Delta$  9 position are DNA chains containing the base sequence of SEQ ID NO: 3.
- (4) Vectors which contain the genes or polynucleotides containing part of thereof according to any one of (1)-(3).
- (5) Plant cells transformed with the genes or polynucleotides containing part of thereof according to any one of (1)-(3).
- (6) A method for creating plants by differentiating the plant cells of (5) to regenerate plants.
- (7) Plants transformed with the genes or polynucleotides containing part of thereof according to any one of (1)-(3).

#### BRIEF DESCRIPTION OF THE DRAWINGS

[0017]

Figure 1 shows the amino acid sequence encoded by des 9 var fragment as compared with the amino acid sequence of mouse stearyl-CoA desaturase (MSCD2). In Figure 1, : indicates that the two sequences have the

same amino acids and indicates that the two sequences have amino acids having similar properties. X shows a range of high homology.

Figure 2 is an electropherogram showing an autoradiogram in a Southern analysis of genomic DNA of *Anacystis nidulans* with des 9 var fragment used as a probe.

Figure 3 shows the relationship among inserted DNA fragments,  $\lambda$  5,  $\lambda$  15 and p15X. Thick arrows show the positions in which a protein is encoded and their directions. Thin arrows show the positions which were sequenced and their directions.

Figure 4 shows the amino acid sequences of des 9 nid as compared with the amino acid sequence of mouse stearoyl-CoA desaturase (MSCD2). The comparison was made in the same manner as in Figure 1.

Figure 5 is a pair of photographs showing in biological morphology the influence of a low temperature treatment on transformed tobacco plants. The left photo shows the result of the low temperature treatment of tobacco transformed with a desaturase gene and the right photo shows the result of the low temperature treatment of tobacco transformed with pBI121.

## PREFERRED EMBODIMENTS OF CARRYING OUT THE INVENTION

**[0018]** The  $\Delta$  9 desaturase of the present invention is an enzyme which is inherently present in blue-green algae as described above in the BACKGROUND ART. The chemical structure of the  $\Delta$  9 desaturase has local similarities to those of the stearoyl-CoA desaturases of mouse (Kaestner, K. H. et al., J. Biol. Chem., 264:14755, 1989), rat (Mihara, K., J. Biochem., 108:1022, 1990) and yeast (Stukey, J. E. et al., J. Biol. Chem., 265:20144, 1990) but it is very different taken as a whole. In addition, it is entirely dissimilar to the chemical structures of known blue-green algal enzymes capable of desaturating lipid-bound fatty acids at the  $\Delta$  6 and  $\Delta$  12 positions and that of a higher plant enzyme capable of desaturating lipid-bound fatty acids at the  $\omega$  3 position (Yadav, N.S. et al., Plant Physiol., 103:467, 1993). In the case where the genes of the present invention are produced from natural materials, blue-green algae may be used as raw materials. The blue-green algae to be used include but are not limited to those belonging to the genera *Anacystis*, *Synechocystis*, *Anabaena* and the like. In order to desaturate saturated molecular species in higher plants, *Anacystis*-type  $\Delta$  9 desaturase (Murata, N. et al., Plant Cell Physiol., 33:933, 1992, Group 1-type blue-green algae belong to the genus *Anacystis*) is preferred to *Anabaena*- and *Synechocystis*-types for the following reason.

**[0019]** The sn-1 and sn-2 positions of almost all membrane lipids of *Synechocystis* PCC6803 and *Anabaena variabilis* are occupied by fatty acids of 18 carbon atoms (C18) and 16 carbon atoms (C16), respectively (Sato, N. et al., Biochim. Biophys. Acta, 710:279, 1982; Wada, H. et al., Plant Cell Physiol., 30:971, 1989). On the other hand, the both sn-1 and sn-2 positions of almost all membrane lipids of *Anacystis nidulans* are occupied by C16 (Bishop, D. G. et al., Plant Cell Physiol., 27:1593, 1986). Hence, it is believed that the  $\Delta$  9 desaturases of *Anabaena* and *Synechocystis* have an activity of reacting with the substrate 18:0/16:0-molecular species to desaturate the 18:0 at the position of sn-1 into 18:1  $\Delta$  9, whereas the  $\Delta$  9 desaturase of *Anacystis* has an activity of reacting with the substrate 16:0/16:0-molecular species to desaturate the 16:0 at the position of sn-1 into 16:1  $\Delta$  9. Considering the fact that higher plants contain a larger amount of 16:0/16:0-saturated molecular species, the *Anacystis*-type  $\Delta$  9 desaturase is more suitable than the *Anabaena*- and *Synechocystis*-type enzymes for the purpose of desaturating saturated molecular species in higher plants.

**[0020]** As will be shown below in the Examples, the genes of the present invention include genes encoding  $\Delta$  9 desaturase which contains the amino acid sequence substantially shown in SEQ ID NO: 4 and their degenerate isomers capable of encoding the same polypeptide except for degenerate codons. The genes of the present invention are predominantly in the form of DNA chains. The "amino acid sequence substantially shown in SEQ ID NO: 4" includes not only the amino acid sequence shown in SEQ ID NO: 4 but also those amino acid sequences in which part of the amino acid sequence of SEQ ID NO: 4 may be modified by deletions, substitutions and/or additions, provided that a  $\Delta$  9 desaturase activity is retained.

**[0021]** The genes of the present invention can be prepared by any conventional known techniques from the aforementioned blue-green algal cells.

**[0022]** In brief, the genes of the present invention can be prepared by culturing blue-green algal cells, collecting them, preparing genomic DNA from the blue-green algal cells by a conventional known technique such as ethanol precipitation or the like, preparing a gene library based on the genomic DNA, selecting clones containing the desired gene from the library and amplifying the clone DNA.

**[0023]** Vectors to be used for the preparation of the gene library include any conventional vectors and specific examples include phages such as  $\lambda$  DASH II (Stratagene), cosmids such as pWE15 (Stratagene), phagemids such as pBluescript II (Stratagene) and the like. The genes of the present invention can be introduced into these vectors by a conventional known method selected for a specific kind of vectors.

**[0024]** Clones in which the gene of the present invention has been introduced is selected from the thus prepared gene library.

[0025] The methods for clone selection include any conventional known selection methods, for example, immunological methods such as plaque hybridization and colony hybridization in which antibodies are used, as well as plaque hybridization and colony hybridization in which nucleotide probes are used. A preferred criterion for the nucleotide probe selection is to use as a probe part of base sequences which are estimated to be similar to the genes of the present invention (eg., base sequences encoding part of the 260-295 amino acid sequence of MSCD2 in Figure 1).

[0026] The base sequence of the gene of the present invention in the selected clone can be determined and confirmed by any conventional known methods such as the Maxam-Gilbert method (Maxam-Gilbert, *Methods Enzymol.*, 65:499, 1980), the dideoxynucleotide chain terminator technique using M13 phage (Messing, J. et al., *Gene*, 19:269, 1982) and the like.

[0027] The actual expression of  $\Delta 9$  desaturase can be confirmed by, for example, the method of Wada et al. (*J. Bacteriol.*, 175:6056, 1993).

[0028] Once the genes of the present invention are sequenced, they can be synthesized by any conventional known methods such as a phosphite method using a commercially available DNA synthesizer.

[0029] The gene of the present invention or a polynucleotide that contains part of the gene and which has the  $\Delta 9$  desaturase activity is separated from the selected clones, incorporated into a vector for gene introduction into a plant host, introducing the vector into a plant cell and expressing  $\Delta 9$  desaturase in the plant, thereby imparting a chilling resistance to the plant.

[0030] The kind of plants into which the genes can be introduced is not particularly limited.

[0031] The vectors for gene introduction should be constructed such that the  $\Delta 9$  desaturase gene can be expressed stably in plants. More specifically, a promoter, a DNA chain encoding a translation control region, a DNA chain encoding a peptide for transfer into chloroplasts, the gene of the present invention or a polynucleotide that contains part of the gene and which has the  $\Delta 9$  desaturase activity, a DNA chain encoding a termination codon and a terminator should be incorporated in appropriate positions. As construction elements for the introduction of genes other than the gene of the present invention, any conventional known elements can be used. Preferred examples of the DNA chain encoding a peptide for transfer into chloroplasts include a gene for the small subunit of pea ribulose-1,5-bisphosphate carboxylase. Examples of the promoter include a cauliflower mosaic viral 35S promoter. Examples of the terminator include a terminator of nopaline synthase.

[0032] The methods for gene introduction into plant cells include any conventional known methods such as the methods described in "Plant genetic transformation and gene expression; a laboratory manual", Draper, J. et al. eds., Blackwell Scientific Publications, 1988. Exemplary methods include biological methods using viruses and Agrobacterium, physicochemical methods such as electroporation, a polyethylene glycol method and microinjection, and the like. Among these, the use of Agrobacterium is preferred for dicotyledonous plants such as tobacco since it secures stable transformation. The methods using Agrobacterium include an intermediate vector method using a wild-type tumor plasmid (*Nature*, 287 (1980) p.654; *Cell*, 32 (1983) p.1033; *EMBO J.*, 3 (1984) p.1525), an intermediate vector method using a vector deficient of a tumor formation gene region on T-DNA (*EMBO J.*, 2 (1983) p.2143; *Bio/Technology*, 3 (1985) p.629), a binary vector method (*Bio/Technology*, 1 (1983) p. 262; *Nature*, 303 (1983) p. 179; *Nucl. Acids Res.*, 12 (1984) p.8711) and the like. Any of these methods can be used. Examples of the methods in which plants are infected with Agrobacterium include direct inoculation to cultured cells, protoplast co-cultivation, a leaf-disk method and the like. A leaf-disk method is preferred in terms of the ability to produce a large number of transformed plants in a direct and easy way.

[0033] In order to regenerate plants, the transformed plant cells may be cultured in known media such as Murashige-Skoog medium that are supplemented with selection antibiotics, plant growth hormones and any other agents. The rooted seedlings are transplanted into the soil and cultured for growth to complete plants.

[0034] The transformed plants which have been grown to mature plants can be tested for chilling resistance by the following procedure:

[0035] A test plant is cultured at a temperature (eg., 25 °C) at which it does not receive a low temperature injury and then cultured temporarily (eg., for a week) at a low temperature (eg., 4°C ). The injury to the plant, for example, chlorosis and the reduction of fertility are measured. Alternatively, the amount of the plant growth may be compared with that of a control plant.

[0036] The present invention will now be explained in greater detail with reference to the following examples which are by no means intended to limit the scope of the present invention.

[Example 1] The cloning of a DNA fragment of open reading frame flanking at the upstream side of the  $\Delta 12$  desaturase gene (des A) of Anabaena variabilis

[0037] Anabaena variabilis IAM M-3 provided by Institute of Molecular and Cellular Biosciences, University of Tokyo, was cultured in about 100 ml of a BG-11 medium ("Plant Molecular Biology", Shaw, C. H. ed., p.279, IRL PRESS, 1988). The culture was shaken at 25 °C under 1,000-lux fluorescent light with a 120-rpm shaking rate to grow bacteria

thoroughly. The culture solution was centrifuged at room temperature at 5,000 g for 10 minutes to collect the precipitated cells.

[0038] For the purpose of preparation of genomic DNA, the cells were suspended in 50 ml of solution A (50 mM Tris-HCl, 1mM EDTA, pH 8.0), washed and centrifuged to collect the precipitated cells. Subsequently, the resulting cells were suspended in 15 ml of solution B (50 mM Tris-HCl, 20 mM EDTA, 50 mM NaCl, 0.25 M sucrose, pH 8.0). To the suspension, 40 mg of lysozyme (Sigma) dissolved in solution B was added and the mixture was shaken at 37 °C for 1 hour. To the shaken culture, proteinase K (15 mg) and SDS (final concentration of 1%) were added and the mixture was shaken at 37 °C overnight. To the shaken culture, NaClO<sub>4</sub> was added to a final concentration of 1M and 20 ml of chloroform/isoamyl alcohol (24:1) was also added. The resulting mixture was shaken for 10 minutes and then centrifuged to collect the aqueous layer. After re-extraction with chloroform/isoamyl alcohol (24: 1), 50 ml of ethanol was added to the aqueous layer and a genomic DNA preparation was collected by winding the DNA on a glass rod. The DNA preparation was dissolved in 20 ml of solution A and NaCl was added at a final concentration of 0.1 M. RNase was added at a final concentration of 50 mg/ml and the resulting mixture was incubated at 37 °C for 1 hour. Subsequently, extraction with an equal amount of phenol saturated with solution A was performed twice and ethanol was then added to the aqueous layer, thereby collecting the precipitated genomic DNA. The genomic DNA was washed with 70% ethanol and dissolved in 1 ml of solution A to prepare an *Anabaena variabilis* genomic DNA solution.

[0039] Sakamoto et al., reporting the presence of an open reading frame (ORF) flanking at the upstream side of  $\Delta$  12 desaturase gene, suggested that it may have something to do with the desaturase when they made a speech on the cloning of a membrane lipid-bound fatty acid  $\Delta$  12 desaturase gene derived from *Anabaena variabilis* (Lecture Abstract No. 3aF04 in the 1993 annual meeting of The Japanese Society of Plant Physiologists). However, the function of the ORF was not identified. The inventors took interest in the ORF and its function. Thus, they synthesized four primers (SEQ ID NOs: 5-8) giving attention to three base sequences in the DNA chain of the ORF and used the primers in PCR with *Anabaena variabilis* genomic DNA used as a template.

[0040] Of the four primers, those having the base sequences of SEQ ID NOs: 5 and 6 encode sense strands and those having the base sequences of SEQ ID NOs: 7 and 8 encode antisense strands. The base sequences of SEQ ID NOs: 6 and 7 are derived from the same amino acid sequence. A primer set was selected from each of the sense and antisense strands and PCR was performed with 4 combinations of primers in total. The primers (20  $\mu$ M, each) and the *Anabaena variabilis* genomic DNA (1  $\mu$ g) were added to the reaction solution (100  $\mu$ l) and reaction was performed with a GeneAmp PCR Kit (TAKARA SHUZO CO., LTD.). The PCR was performed through 35 cycles, each cycle consisting of 95 °C (1 minute), 45 °C (1 minute) and 72°C (2 minutes). In the first cycle, the temperature of 95 °C was maintained for 3 minutes. After the end of the reaction, the reaction solution (10  $\mu$ l) was electrophoresed on a 2% agarose gel to separate the synthesized DNA for analysis. As a result of the analysis, a DNA fragment having an expected size (about 190 bp) was detected as a main band in the DNAs synthesized using a combination of primers having the base sequences of SEQ ID NOs: 6 and 8. The DNA fragment (hereinafter referred to as "des 9 var") was treated with a Klenow fragment to make blunt ends and then cloned into plasmid pTZ18R (Pharmacia) at a SmaI site, followed by determination of the base sequence with a fluorescent DNA sequencer (Applied Biosystems). The determined base sequence is shown in SEQ ID NO: 1. The amino acid sequence (SEQ ID NO: 2) deduced from this base sequence had a significant homology with mouse stearyl-CoA desaturase (see Figure 1 showing the amino acid sequence encoded by the des 9 var fragment as compared with that of mouse stearyl-CoA desaturase (MSCD2)).

[0041] Subsequently, *Anacystis nidulans* genomic DNA was analyzed by Southern blotting technique with the des 9 var fragment used as a probe. Restriction enzymes XhoI, PstI and BamHI were used individually to cleave *Anacystis nidulans* genomic DNA (about 0.1  $\mu$ g). The resulting DNA fragments were separated by electrophoresis on a 0.8% agarose gel and then blotted on a nylon membrane (Hybond-N<sup>+</sup>; Amersham). Probe DNAs were labelled with [ $\alpha$ -<sup>32</sup>P] dCTP using a Multiprime DNA labelling Kit (Amersham). The probe DNAs were reacted with the membrane by incubation in a solution consisting of a 6xSSPE [1xSSPE was a mixture of a 10 mM phosphate buffer (pH 7.0), 1 mM EDTA and 0.15 M NaCl], 0.2% SDS and 100  $\mu$ g/ml herring sperm DNA at 55 °C for 16 hours. The resulting membrane was then washed by shaking it twice in 2xSSC [1xSSC was a mixture of 0.15 M NaCl and 15 mM sodium citrate] at room temperature for 15 minutes and twice in 0.1xSSC at 40 °C for 15 minutes and analyzed by autoradiography. As a result, only one DNA fragment was detected in the case where the genomic DNA had been cleaved with any of the restriction enzymes (see Figure 2. In Figure 2, "Non" means that the genomic DNA was not cleaved with any restriction enzymes.).

[Example 2] Cloning of a DNA chain in *Anacystis nidulans* genomic DNA having high homology with the des 9 var fragment

[0042] The cultivation of *Anacystis nidulans* R2-SPc provided by Institute of Molecular and Cellular Biosciences, University of Tokyo and the preparation of genomic DNA were conducted by the same method as in the case of *Anabaena variabilis*. The genomic DNA (about 100  $\mu$ g) was partially digested with Sau3AI and DNA fragments of about 9-23 kbp were then collected by ultracentrifugation in a sucrose density gradient according to the method described

in "Molecular Cloning, 2nd edition, pp.2.85-2.87 (Sambrook, J. et al., eds., Cold Spring Harbor Laboratory, 1989). The collected DNA fragments were cloned into lambda phage vector  $\lambda$  DASH II (Stratagene) cleaved with BamHI and HindIII and then packaged in phage particles to yield an *Anacystis nidulans* genomic DNA library. *E. coli* P2392 cells were infected with the phage library and cultivated in NZYM-agar plates of about 15 cm in diameter. A total of about 100,000 plaques were formed and then blotted on nylon membranes (Hybond-N<sup>+</sup>; Amersham). In the same manner as in the above Southern analysis, the des 9 var fragment labelled with [ $\alpha$ -<sup>32</sup>P]dCTP was reacted with the resulting membranes and positive phages detected by autoradiography were screened to yield about 30 phage clones having different signal intensities, from which 12 clones were selected. Phage DNA was obtained from each clone by a conventional method. The obtained phage DNA was cleaved with several kinds of restriction enzymes and separated by electrophoresis on a 0.8% agarose gel, followed by blotting on a nylon membrane. The resulting membrane was analyzed by Southern blotting technique under the same conditions as in the above screening and comparison was made on the lengths and signal intensities of DNA fragments hybridizing with the probe DNAs. As a result, 2 clones,  $\lambda$  5 and  $\lambda$  15, showed the strongest signals. Since these two clones had inserts DNA of 11 and 15 kbp, it was estimated that they were sufficiently long to contain the entire region of the desired ORF. The insert DNA of each of the two clone was cleaved with several restriction enzymes and analysed by Southern blotting technique. As a result, a DNA fragment of about 5 kbp was detected in both clones when they were cleaved with XhoI and hybridized. This DNA fragment was subcloned into pBluescript SK-(Stratagene) at a XhoI site to yield plasmids p5X and p15X which contained the DNA fragments derived from  $\lambda$  5 and  $\lambda$  15, respectively. The detailed restriction maps of p5X and p15X were prepared and compared. These maps show that both p5X and p15X contain the same genomic DNA fragment (see Figure 3 showing the correlativity between the DNA inserts of  $\lambda$  5 and  $\lambda$  15. The hatched rectangles show DNA fragments hybridizing with the probe des 9 var fragment in the process of screening. Thick arrows show the region of des 9 nid (to be described below) and the direction of a sense strand. Thin arrows show the sequencing direction of a region containing des 9 nid. The bars of 5 kbp, 1.25 kbp and 0.5 kbp show size markers for the left maps. Abbreviations of restriction enzymes have the following meanings: B, BamHI; H, HindIII; N, NotI; Hp, HpaI; RI, EcoRI; RV, EcoRV; S, Sall; P, PstI; and X, XhoI).

**[0043]** Deletion plasmids were prepared from p15X with restriction enzymes or ExoIII and the base sequence of a DNA fragment of about 2 kbp which contained a region hybridizing with the des 9 var fragment was determined with a fluorescent DNA sequencer (see Figure 3). As a result, it was estimated that the DNA fragment would contain ORF (des 9 nid) of 834 bp (SEQ ID NO: 3) which would encode 278 amino acids (SEQ ID NO: 4). The amino acid sequence had about 80% homology with the amino acid sequence (SEQ ID NO: 2) of the previously cloned des 9 var fragment derived from *Anabaena variabilis*. A search for highly homologous amino acid sequences was conducted with nucleic acid and amino acid sequence analysis software GENETYX (Software Development) and nucleic acid and amino acid sequence data bases EMBL and DDBJ. The results of the search revealed that des 9 nid had about 30% general homology with mouse stearyl-CoA desaturase but that part of des 9 nid had a very high homology with the mouse desaturase (see Figure 4 showing the amino acid sequence of des 9 nid as compared with that of mouse stearyl-CoA desaturase (MSCD2)), suggesting that the obtained des 9 nid should encode an enzyme capable of desaturating fatty acids.

[Example 3] Determination of activity by the expression of the des 9 nid gene in *E. coli*

**[0044]** *Anacystis nidulans* desaturase has only a  $\Delta$  9 desaturase activity or the ability to desaturate lipid-bound saturated fatty acids at the 9 position (Bishop, D.G. et al., Plant Cell Physiol., 27:1593, 1986). For this reason, an attempt was made to express the polypeptide encoded by des 9 nid in *E. coli* and measure its activity.

**[0045]** Since direct expression from p15X is difficult, a vector for expression in *E. coli* was prepared. Specifically, pET3a (Novagen) was selected as a vector and des 9 nid was cloned into pET3a between the sites of NdeI and BamHI such that extra amino acids would not be added to the amino terminus of the polypeptide encoded by des 9 nid by the following procedure:

**[0046]** In order to insert a BamHI site just downstream of the C terminus of the des 9 nid-encoded protein, PCR was performed with the base sequences of two parts having the C terminus therebetween. Specifically, the PCR was performed with the following two primers and using p15X as a template to yield a product of about 140 bp.

**Sense primer:** 5'-ACGTCATGGCCTGCAGT (a PstI site is underlined) (SEQ ID NO: 9)



Antisense primer: 5'-CGCGGATCCTTAGTTGTTTGGAGACG (a single line is drawn under a BamHI site and a double line is drawn under a stop codon) (SEQ ID NO: 10)

[0047] The product was subcloned into pUC19 at a SmaI site and the accuracy of the base sequence was confirmed. As a result, an EcoRI site was created downstream of the BamHI site in the resulting plasmid. The plasmid was cleaved sequentially with EcoRI and PstI. In the meantime, p15X was cleaved with the same restriction enzymes to introduce a BamHI site just downstream of the stop codon. The plasmid was cleaved with Sall and a fill-in reaction was then performed with a DNA polymerase Klenow fragment in the presence of four kinds of dNTP, followed by cleavage with HindIII. The adaptor consisting of the following two synthesized DNAs was inserted into the resulting plasmid, thereby introducing an NdeI site into the amino terminus. The adaptor was a mixture of equal molar amounts of the following two DNAs.

5'-CATATGACCCTTGCTATCCGACCCA (an NdeI site is underlined) (SEQ ID NO: 11) and

5'-AGCTTGGGTCGGATAGCAAGGGTCATATG (a single line is drawn under an NdeI site and a double line is drawn under part of a HindIII site) (SEQ ID NO: 12)

[0048] The resulting plasmid (pDes9Nde) was introduced into competent cells of *E. coli* strain BL21 (DE3) (Novagen) which were prepared by a conventional method (Molecular cloning, pp.250-251, 1982). Transformed strain BLDES1 was obtained by selection for ampicillin resistance.

[0049] The BLDES1 strain and pET3a-containing BL21 strain (BL1) were inoculated in a 100-ml M9 medium (containing 200 µg/ml ampicillin, 4 mg/ml glucose, 10 µM FeCl<sub>3</sub>, 0.5 µg/ml vitamin B1, 1 mg/ml casamino acids) and cultured at 37 °C. The cultivation was continued until the turbidity of the culture solution reached 0.5 O.D. at a wavelength of 600 nm. Isopropylthiogalactoside (IPTG) was added at a final concentration of 1 mM to the culture solution. The cells were cultured for an additional 1 hour to induce the expression of the Δ 9 desaturase gene. The *E. coli* pellets were collected and washed with 1.2% NaCl, followed by lipid extraction. Lipids were extracted by the method of Bligh and Dyer (Can J. Biochem. Physiol., 37:911, 1959) and reacted with 5% methanolic hydrochloride (2.5 ml) at 85°C for 2.5 hours under a completely sealed condition to yield methylated fatty acids. The produced fatty acid methyl esters were extracted 4 times with hexane (2.5 ml) and concentrated by removing the solvent with nitrogen gas. The fatty acid methyl esters were analyzed by gas chromatography. The fatty acids were identified by comparison of retention times with standard fatty acid methyl esters. The quantitative analysis was conducted with a Chromatopack C-R7A plus (Shimadzu Corp.). The results are shown in the following Table 1.

Table 1.

Fatty Acid Composition in <i>E. coli</i>				
Strain	16:0	16:1	18:1 Δ 11	Others
BL1 (0 hour)	47	20	29	4
BL1 (1 hour)	50	17	29	4
BLDES1 (0 hour)	44	22	30	4
BLDES1 (1 hour)	40	28	28	4

The hours indicate the time of protein induction with IPTG.

[0050] The results revealed that 16:1 increased in the BLDES1 strain, showing that the gene of the present invention has an activity of desaturating 16:0.

**[0051]** The two strains were cultured in an M9 medium supplemented with 0.1 mM stearic acid and a comparison was made in the same manner as above. In contrast to the BL strain, the BLDES1 strain produced not only 16:1 but also 18:1  $\Delta$  9. This indicates that the polypeptide encoded by des 9 nid can use not only 16:0 but also 18:0 as a substrate to produce unsaturated fatty acids.

[Example 4] Introduction of the des 9 nid gene into tobacco plants

**[0052]** The Anacystis nidulans-derived des 9 nid gene was incorporated into tobacco plants as follows:

(1) Construction of vector plasmid for expression in plants

**[0053]** Plasmid pDes9Nde was cleaved with SacI and Sall to yield a des 9 nid gene fragment held between the sites of the two restriction enzymes. A chloroplast-transit sequence was cut off from the clone pSNIP9 containing a pea RuBisCO gene (Schreicher et al., EMBO J. 4, 25 (1985)) with HindIII and SphI and cloned into pUC118 cleaved with the same restriction enzymes, thereby yielding plasmid pTRA3 containing a multicloning site downstream of the transit sequence. The HindIII site of this plasmid was cleaved and filled in with a Klenow enzyme, followed by insertion of a XbaI linker (pTRA3X). The plasmid pTRA3X was cleaved with Sall and SacI and the des 9 nid fragment which had been obtained by cleavage with the same restriction enzymes was inserted (pTRA3Xdes9). In the pTRA3Xdes9, the des 9 nid gene would be translated, following the transit sequence of RuBisCO, in the same reading frame. This plasmid was cleaved with SacI and XbaI and inserted into the following vector for plants. Plant expression-type binary plasmid pBI121 (Clontech) was cleaved with SacI and XbaI to yield  $\beta$ -glucuronidase gene (GUS gene)-free plasmid pBI(-GUS). The transgene prepared above was inserted into the plasmid pBI(-GUS) between the cauliflower mosaic viral 35S promoter and the nopaline synthase (NOS) terminator to yield a vector (pBI121(-GUS)Rbcs-des9) for introduction into plants.

(2) Introduction of pBI121(-GUS)Rbcs-des9 into Agrobacterium

**[0054]** Agrobacterium tumefaciens LBA4404 (Clontech) was inoculated in a 50-ml YEB medium (5g beef extract, 1g yeast extract, 1g peptone and 5g sucrose per liter supplemented with 2 mM MgSO<sub>4</sub> (pH 7.4)) and cultured at 28 °C for 24 hours. The cultured solution was centrifuged at 3,000 rpm at 4 °C for 20 minutes to collect the cells. The cells were washed 3 times with 10 ml of 1 mM Hepes-KOH (pH 7.4) and once with 3 ml of 10% glycerol. The cells were suspended in 3 ml of 10% glycerol to prepare Agrobacterium cells for DNA introduction.

**[0055]** The thus obtained cell solution (50  $\mu$ l) and the plasmid pBI121(-GUS)Rbcs-des9 (1  $\mu$ g) were placed in a cuvette and treated with electric pulses using an electroporation apparatus Gene Pulser (BioRad) under the conditions of 25  $\mu$ f, 2500 V and 200  $\Omega$ , thereby introducing the plasmid DNA into the Agrobacterium. The cell solution was transferred into an Eppendorf tube and 800  $\mu$ l of an SOC medium (20g tryptone, 5g yeast extract, and 0.5g NaCl per liter supplemented with 2.5 mM KCl, 10 mM MgSO<sub>4</sub>, 10 mM MgCl<sub>2</sub> and 20 mM glucose (pH 7.0)) was added. The cells were cultured statically at 28 °C for 1.5 hours. The culture solution (50  $\mu$ l) was inoculated in a YEB agar medium (agar 1.2%) supplemented with 100 ppm kanamycin and cultured at 28 °C for 2 days.

**[0056]** Well separated colonies were selected from the resulting colonies and plasmid DNA was prepared from the selected colonies by an alkaline method. The plasmid DNA was digested with appropriate restriction enzymes. The resulting DNA fragments were then separated by electrophoresis on a 1% agarose gel and analyzed by Southern blotting technique using <sup>32</sup>P-labelled des 9 nid gene fragment as a probe; it was confirmed that the Agrobacterium cells contained the plasmid pBI121(-GUS)Rbcs-des9. This Agrobacterium tumefaciens cell line is abbreviated as "ALBBSDES".

(3) Transformation of Tobacco

**[0057]** The cell line ALBBSDES was cultured with shaking in an LB liquid medium supplemented with 50 ppm kanamycin at 28 °C for 2 days. The culture solution (1.5 ml) was centrifuged at 10,000 rpm for 3 minutes to collect the cells. The cells were then washed with a 1-ml LB medium to remove kanamycin. Additionally, the cell solution was centrifuged at 10,000 rpm for 3 minutes to collect the cells. The cells were then re-suspended in a 1.5-ml LB medium to prepare a cell solution for infection.

**[0058]** For infection of tobacco, a young leaf of tobacco was taken and immersed into a 0.5% sodium hypochlorite solution in water for 10 minutes. Subsequently, the leaf was washed 3 times with sterilized water and wiped with sterilized filter paper to prepare a leaf sample to be infected. The leaf was cut off aseptically with a knife in pieces of 1 cm<sup>2</sup> each. The pieces were placed on the solution of Agrobacterium such that the back of each piece faced up and shaken gently for 2 minutes. The pieces were then placed on sterilized filter paper to remove excess Agrobacterium.

## EP 0 736 598 B1

Whatman No.1 filter paper ( $\phi$  7.0 cm) was placed on an MS-B5 medium (containing 1.0 ppm benzyladenine, 0.1 ppm naphthaleneacetic acid, and 0.8% agar) (Murashige, T. and Skoog, F. *Plant Physiol.*, 15:473, (1962)) in a plate and each piece of the sample leaves was placed on the filter paper such that the back of the leaf faced up. The plate was sealed with a parafilm and the leaf was cultured through cycles of 16 hours under light and 8 hours in the dark at 25 °C for 2 days. Subsequently, the leaf was transferred into an MS-B5 medium supplemented with 250 ppm calforan and cultured under the same conditions to remove the *Agrobacterium*. Additionally, the leaf was placed in an MS-B5 medium supplemented with 250 ppm calforan and 100 ppm kanamycin on the bottom of a plate and cultured under the same conditions for 7 days. In the mean time, the calluses were formed around the leaf and shoots were initiated. After cultivation for an additional 10 days, the extended shoot was placed in an MS-HF medium (MS-B5 medium free from benzyladenine and naphthaleneacetic acid) supplemented with 250 ppm calforan and 100 ppm kanamycin. After cultivation for 10 days, the rooted shoot was selected as a kanamycin-tolerant transformant and transplanted into an MS-HF medium supplemented with 250 ppm calforan in a plant box.

[Example 5] Southern and Northern analyses of the transformed tobacco genome

**[0059]** DNA was extracted from the kanamycin-tolerant tobacco and analyzed by the Southern and Northern blotting techniques to confirm the introduction and the expression of the desired gene. The extraction of the genomic DNA was conducted by a CTAB method according to a manual (Rogers, S. O. & Bendich, A. J.: *Plant Molecular Biology Manual A6*; 1 (1988)). In brief, tobacco leaves (2 g) were ground in liquid nitrogen and genomic DNA was extracted with a CTAB extraction buffer. The DNA (10  $\mu$ g) was cleaved with EcoRI and XbaI and then electrophoresed on a 0.7% agarose gel, followed by blotting of the separated DNA fragments with 0.4 N NaOH on a nylon membrane (Hybond N+; Amersham). The transit-containing desaturase gene derived from pTRA3Xdes9 was used as a probe for hybridization with the membrane at 65 °C for 16 hours, thereby confirming that the desired gene was introduced into the tobacco genome.

**[0060]** RNA was extracted from the tobacco leaves (about 2 g) and analyzed to confirm the expression of the transgene. The procedure consisted of extracting poly(A)+RNA with guanidium thiocyanic acid (Nagy, F. et al., *Plant Molecular Biology Manual B4*; 1 (1988)) and electrophoresing on a formaldehyde-containing agarose gel. The RNA was blotted on a nylon membrane (Hybond N; Amersham) and analyzed under the same hybridization condition as in the Southern method. Among the transformants expressing various amounts of RNA, those expressing large amounts of RNA were selected and analyzed for fatty acids.

[Example 6] Fatty acid analysis of the lipids in the transformed tobacco plants

**[0061]** Lipids such as phosphatidylglycerol (PG) and sulfoquinovosyldiacylglycerol (SQDG) were prepared by the following method from the leaves of the transformed tobacco plants which were verified for high RNA expression in Example 5 and those of control tobacco plants transformed with pBI121. Their fatty acid compositions were analyzed. The root lipids were also analyzed in regard to part of the transformants and non-transformants.

(1) Extraction of total lipids

**[0062]** Lipids were extracted by the Bligh-Dyer method (*Can J. Biochem. Physiol.*, 37:911, 1959). Two grams by wet weight of the leaves (1 g, when part of the root was used as a sample) were sliced with a knife and 20 ml of chloroform/methanol (1:2, volume ratio) was added. The leaves were disrupted with a homogenizer and left to stand for 15 minutes. Chloroform (12 ml) and distilled water (12 ml) were added to the disrupted leaves and the mixture was stirred vigorously. The mixture was centrifuged at 3,000 rpm at 4 °C for 30 minutes to separate the aqueous and organic layers. The organic (low) layer was collected and an appropriate amount of ethanol was added. The solvents were distilled off at 30 °C under a reduced pressure with a rotary evaporator. The residue was dissolved in 2 ml of chloroform/methanol (1:4, volume ratio) to prepare a total lipid extract. Part of the total lipid extract was treated with 5% methanolic hydrochloric acid by the following method to yield methylated fatty acids.

(2) Fractionation of lipids

**[0063]** A suspension (2.5 ml) of DEAE-Toyopearl 650C (TOSOH) was mixed with 25 ml of a 1 M sodium acetate solution in water (pH 7.0) to prepare an acetic acid form. The resulting suspension was washed sequentially with distilled water and methanol and suspended in methanol. The resulting suspension was packed in a column (i.d. 2 cm) to 1.5 cm in height and washed with 50 ml of chloroform/methanol (1:4, volume ratio).

**[0064]** Subsequently, the total lipid extract was applied to the column. Monogalactosyldiacylglycerol (MGDG), digalactosyldiacylglycerol (DGDG), phosphatidylethanolamine (PE) and phosphatidylcholine (PC) were eluted with 50 ml

of chloroform/methanol (1:4, volume ratio) to prepare neutral lipid (MGDG, DGDG, PE and PC) fractions. Then, phosphatidylserine (PS) was eluted out with 5 ml of acetic acid and the acetic acid was washed away with 20 ml of chloroform/methanol (1:4, volume ratio). A fraction containing PG, SQDG and phosphatidylinositol (PI) was obtained by extraction with 50 ml of an aqueous solution of chloroform, methanol and 10 M ammonium acetate (20:80:0.2, volume ratio). Ethanol (15 ml) was added to this fraction and the solvents were distilled off under a reduced pressure. The residue was dissolved in 0.2 ml of chloroform/methanol (2:1, volume ratio) to prepare acidic lipid (PG, SQDG and PI) fractions. **[0065]** The MGDG, DGDG, PE and PC fractions were further fractionated by silicic acid column chromatography (Iatrobeads, Iatoron Laboratories Inc.). More specifically, the samples dissolved in chloroform (1 ml) were applied to a column equilibrated with chloroform and eluted sequentially with chloroform/acetone (4:1), acetone and methanol, whereby glycolipids (MGDG and DGDG) were eluted with acetone and phospholipids (PC and PE) with methanol.

### (3) Isolation and purification of PG by thin layer chromatography (TLC) and fatty acid analysis

**[0066]** The fractions obtained in step (2) were separated with a silica gel-TLC plate #5721 (Merck). As developing solvents, chloroform/acetone/methanol/acetic acid/water (50:20:10:15:5, volume ratio) was used for the separation of acidic lipids and chloroform/methanol/water (70:21:3, volume ratio) was used for the separation of neutral lipids. After separation by TLC, primulin (80% solution in acetone) was sprayed to develop fluorescence under ultraviolet light. The various classes of the lipid fractions were estimated by comparison of mobility with standard lipids. The fluorescence-developing lipids were scraped together with the silica gel and placed in test tubes equipped with screw caps. When the fatty acid compositions of the lipids were to be estimated, 3 ml of methanolic 5% hydrochloric acid was added to the lipids and the mixtures were reacted at 85 °C for 2 hours under a completely sealed condition to yield methylated fatty acids. In the meantime, in order to determine the fatty acid compositions at the sn-1 and sn-2, the lipids were collected from the scraped silica gel with 5 ml of a chloroform/methanol (2:1) mixed solution and dried. To the lipids, 1 ml of 50 mM TrisCl (pH 7.2) and 0.05% Triton X-100 were added and the mixture was stirred vigorously to disperse the lipids. *Rhizopus deleamar*-derived lipase (2500U; Boehringer) was added to the dispersion and the mixture was held at 37°C for 30 minutes to liberate fatty acids selectively at the sn-1 position. After the reaction products were concentrated, unreacted lipids, Lysolipids and free fatty acids were separated by TLC using chloroform/acetone/methanol/acetic acid/water (10:4:2:3:1) as a developing solvent. These substances were collected from the gel and reacted with methanolic hydrochloric acid by the same method as described above to yield methylated fatty acids. The produced fatty acid methyl esters were extracted 4 times with 3 ml of hexane and concentrated by distilling off the solvent under a reduced pressure. The fatty acid methyl esters were analyzed by gas chromatography. The fatty acids were identified by comparison of retention time with standard fatty acid methyl esters. The quantitative analysis was conducted with a Chromatopack C-R7A plus (Shimadzu Corp.). The results for the total lipids, PG and other representative lipids are shown in Tables 2, 3 and 4, respectively. These tables show average analytical values for 2 controls, and 2 or 3 independent transformants.

Table 2

Results of fatty acid analysis on the total lipids in leaves									
Plant	16:0	16:1	16:2	16:3	18:0	18:1	18:2	18:3	Σ 16:0+18:0
Control plant	17	3	1	4	3	1	9	63	20
Transformed plant	10	12	1	5	1	2	14	56	11

Table 3

Results of fatty acid analysis on PG									
	Plant	16:0	16:1t	16:1c	18:0	18:1	18:2	18:3	Σ 16:0+18:0+16:1t
PG	Control plant	32	37	0	1	5	10	14	70
	Transformed plant	18	37	8	0	10	12	15	55

Table 4

Results of fatty acid analysis on other lipids											
	Plant	16:0	16:1	16:2	16:3	18:0	18:1	18:2	18:3	Σ 16:0+18:0	
5	SQDG	Control plant	51	1	0	0	3	2	7	36	54
		Transformed plant	36	22	0	0	0	4	9	28	36
10	MGDG	Control plant	7	0	1	9	1	2	4	76	8
		Transformed plant	3	9	1	10	0	1	5	69	3
15	DGDG	Control plant	19	0	0	0	3	1	4	73	22
		Transformed plant	9	13	0	1	0	1	5	70	9
20	PC	Control plant	28	0	0	0	5	1	21	44	33
		Transformed plant	19	12	0	0	3	4	40	23	22
25	PE	Control plant	20	0	0	0	3	1	6	70	23
		Transformed plant	18	10	0	0	2	2	31	38	20
30	PI	Control plant	48	1	0	0	2	1	11	37	50
		Transformed plant	44	7	0	0	1	2	18	28	45

**[0067]** The results of the analysis of PG-bound fatty acids revealed that in the transformed tobacco plants expressing *Anacystis nidulans*-derived fatty acid desaturase, 16:0 (palmitic acid) decreased greatly whereas 16:1 cis increased and 18:0 became almost zero (a small amount of 18:0 was present in the control) whereas 18:1 increased. Hence, the content of saturated fatty acids (16:0+16:1trans+18:0(stearic acid)) was 70% in the control tobacco plants whereas it was significantly reduced to 55% in the tobacco plants transformed with the desaturase gene. The results of the respective analyses on PG at the sn-1 and sn-2 positions revealed that more than 98% of the sn-2 position was occupied with saturated fatty acids (16:0 or 16:1trans) and that all of the 16:1 newly produced by gene introduction was present at the sn-1 position. Hence, it is clear that the amount of saturated fatty acids at the sn-1 position of PG became extremely small in the tobacco transformed with the desaturase gene. In conclusion, it was found that the amount of saturated molecular species consisting of saturated fatty acids at both sn-1 and 2 positions decreased greatly and that, therefore, the tobacco plants changed into a significantly chilling-resistant type as estimated from the composition of molecular species of the lipids.

**[0068]** In regard to the other lipids, MGDG, DGDG, SQDG, PC, PE and PI, it is clear that 16:0 decreased with a corresponding increase of about 10% in 16:1 and that the desaturation of 18:0 was enhanced. In MGDG and DGDG, 16:1 was produced mainly at the sn-1 position but the production of a small amount of 16:1 was also detected at the sn-2 position. It was found that the desaturation of MGDG, DGDG, SQDG and PI, which were lipids present predominantly in chloroplasts, was surprisingly enhanced by expressing the desaturase of the blue-green alga *Anacystis nidulans* in the chloroplasts of higher plants. There was a high possibility that these four kinds of lipids which were present in the membranes of *Anacystis nidulans* could be used as substrates for the desaturase. It was quite surprising that palmitic acid and stearic acid were desaturated in PC, PE and PI because these lipids are absent in the membranes of *Anacystis nidulans* but present predominantly outside the chloroplasts in higher plants.

**[0069]** In this example, the results of lipid analysis of the transformed tobacco plants demonstrated that the *Anacystis nidulans*-derived fatty acid desaturase could desaturate 16:0 and 18:0 in almost all lipids with an extremely high efficiency in the transformed tobacco which is a higher plant.

**[0070]** The results of fatty acid analysis on total root lipids are shown in Table 5.

Table 5

Results of fatty acid analysis on total root lipids									
Plant	16:0	16:1	16:2	16:3	18:0	18:1	18:2	18:3	Σ 16:0+18:0
Non-transformed plant	26	0	0	0	5	2	47	21	31
Transformed plant	17	13	0	0	2	4	58	6	19

**[0071]** The results revealed that surprisingly enough, the *Anacystis nidulans*-derived fatty acid desaturase catalyzed

the desaturation of 16:0 and 18:0 in not only leaves but also roots, suggesting that the fatty acid desaturase gene of the present invention should have not only the possibility of improving the chilling resistance of plants but also the possibility of increasing the content of unsaturated fatty acids and that it is useful in industries where plants are used as raw materials for oil production.

5

[Example 7] Testing the transformed tobacco plants for chilling resistance

**[0072]** The transformants estimated to be promising in the RNA expression assay and the lipid analyses were self-pollinated and the seeds of next generation were collected. Part of the seeds were planted in an MS-HF medium supplemented with 800 ppm kanamycin and cultured at 25 °C for 2 weeks under the conditions of 16 hours under light and 8 hours in the dark. Kanamycin-tolerant seedlings were selected. The seedlings were transplanted in plant boxes and cultured for an additional 4 weeks. In regard to the plant transformed with pBI121, the above procedure was repeated (control).

10

**[0073]** The plants were subjected to a low temperature treatment at 4°C under continuous light for 11 days and then cultured at 25 °C for 2 days. As a result, marked leaf waving and chlorosis were observed in the control plants (the plant transformed with pBI121) but no injury was observed in the transformed plants. Hence, it was assumed that the chilling resistance was improved by introduction of the desaturase gene.

15

#### INDUSTRIAL APPLICABILITY

20

**[0074]** Chilling resistance can be imparted to plants and their content of unsaturated fatty acids can be increased by introducing the gene of the present invention which encodes the  $\Delta 9$  desaturase.

#### SEQUENCE LISTING

25

##### **[0075]**

##### (1) GENERAL INFORMATION:

30

##### (i) APPLICANT:

- (A) NAME: KIRIN BEER KABUSHIKI KAISHA
- (B) STREET: 13-5, Fukuura 1-chome, Kanazawa-ku
- (C) CITY: Yokohama-shi
- (D) STATE: Kanagawa 236
- (E) COUNTRY: Japan
- (F) POSTAL CODE (ZIP): None

35

(ii) TITLE OF INVENTION: Gene for fatty acid desaturase, vector containing said gene, plant containing said gene transferred thereto, and process for creating said plant

40

(iii) NUMBER OF SEQUENCES: 12

##### (iv) COMPUTER READABLE FORM:

45

- (A) MEDIUM TYPE: Floppy disk
- (B) COMPUTER: IBM PC compatible
- (C) OPERATING SYSTEM: PC-DOS/MS-DOS
- (D) SOFTWARE: PatentIn Release #1.0, Version #1.30 (EPO)

50

##### (v) CURRENT APPLICATION DATA:

APPLICATION NUMBER: EP 95904020.5-2105

##### (2) INFORMATION FOR SEQ ID NO: 1:

55

##### (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 196 base pairs

EP 0 736 598 B1

- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

5 (ii) MOLECULE TYPE: DNA (genomic)

(vi) ORIGINAL SOURCE:

- 10 (A) ORGANISM: Anabaena variabilis
- (B) STRAIN: IAM M-3

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

15 GCTCTGGGGT TGTTCGCTGTT ATATCTAGGC GGGTGGTCTT TTGTGGTCTG GGGAGTTTTTC 60  
TTTCGCATCG TTTGGGTTTA CCACTGTACT TGGTTGGTAA ACAGCGCTAC CCATAAGTTT 120  
20 GGCTACCGCA CCTATGATGC TGGTGACAGA TCCACTAACT GTTGGTGGGT AGCTGTCCTA 180  
GTGTTTGGTG AAGGTT 196

25

(2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:

30

- (A) LENGTH: 65 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

35

(ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:

- 40 (A) ORGANISM: Anabeana variabilis
- (B) STRAIN: IAM M-3

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

45

Ala Leu Gly Leu Leu Leu Tyr Leu Gly Gly Trp Ser Phe Val Val  
1 5 10 15

50

55

EP 0 736 598 B1

5 Trp Gly Val Phe Phe Arg Ile Val Trp Val Tyr His Cys Thr Trp Leu  
20 25 30

10 Val Asn Ser Ala Thr His Lys Phe Gly Tyr Arg Thr Tyr Asp Ala Gly  
35 40 45

15 Asp Arg Ser Thr Asn Cys Trp Trp Val Ala Val Leu Val Phe Gly Glu  
50 55 60

20 Gly  
65

20 (2) INFORMATION FOR SEQ ID NO: 3:

(i) SEQUENCE CHARACTERISTICS:

- 25 (A) LENGTH: 837 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

30 (ii) MOLECULE TYPE: DNA (genomic)

(vi) ORIGINAL SOURCE:

- 35 (A) ORGANISM: Anacystis nidulans
- (B) STRAIN: R2-SPc

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

40 ATGACCCTTG CTATCCGACC CAAGCTTGCC TTCAACTGGC CGACCGCCCT GTTCATGGTC 60

GCCATTCACA TTGGAGCACT GTTAGCGTTC CTGCCGGCCA ACTTTAACTG GCCCGCTGTG 120

45 GGCGTGATGG TTGCGCTGTA TTACATTACC GGTGTGTTTTG GCATCACCCCT AGGCTGGCAC 180

50

55



EP 0 736 598 B1

CGGCTAATTT CGCACCGTAG CTTTGAAGTT CCCAAATGGC TGAATACGT GCTGGTGTTTC 240  
5 TGTGGCACCT TGGCCATGCA GCACGGCCCC ATCGAATGGA TCGGTCTGCA CCGCCACCAT 300  
CACCTCCACT CTGACCAAGA TGTCGATCAC CACGACTCCA ACAAGGGTTT CCTCTGGAGT 360  
10 CACTTCCTGT GGATGATCTA CGAAATTCCG GCCCGTACGG AAGTAGACAA GTTCACGCGC 420  
GATATCGCTG GCGACCCTGT CTATCGCTTC TTTAACAAAT ATTTCTTCGG TGTCCAAGTC 480  
15 CTACTGGGGG TACTTTTGTA CGCCTGGGGC GAGGCTGGG TTGGCAATGG CTGGTCTTTC 540  
GTCGTTTGGG GGATCTTCGC CCGCTTGGTG GTGGTCTACC ACGTCACTTG GCTGGTGAAC 600  
AGTGCTACCC ACAAGTTTGG CTACCGCTCC CATGAGTCTG GCGACCAGTC CACCAACTGC 660  
25 TGGTGGGTTG CCCTTCTGGC CTTTGGTGAA GGCTGGCACA ACAACCACCA CGCCTACCAG 720  
TACTCGGCAC GTCATGGCCT GCAGTGGTGG GAATTTGACT TGACTTGGTT GATCATCTGC 780  
30 GGCCTGAAGA AGGTGGGTCT GGCTCGCAAG ATCAAAGTGG CGTCTCCAAA CAACTAA 837

35 (2) INFORMATION FOR SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:

- 40 (A) LENGTH: 278 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

45 (ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:

- 50 (A) ORGANISM: *Anacystis nidulans*  
(B) STRAIN: R2-SPc

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

55

EP 0 736 598 B1

Met Thr Leu Ala Ile Arg Pro Lys Leu Ala Phe Asn Trp Pro Thr Ala  
 1                    5                    10                    15  
 5  
 Leu Phe Met Val Ala Ile His Ile Gly Ala Leu Leu Ala Phe Leu Pro  
                   20                    25                    30  
 10  
 Ala Asn Phe Asn Trp Pro Ala Val Gly Val Met Val Ala Leu Tyr Tyr  
                   35                    40                    45  
 15  
 Ile Thr Gly Cys Phe Gly Ile Thr Leu Gly Trp His Arg Leu Ile Ser  
                   50                    55                    60  
 20  
 His Arg Ser Phe Glu Val Pro Lys Trp Leu Glu Tyr Val Leu Val Phe  
 65                    70                    75                    80  
 25  
 Cys Gly Thr Leu Ala Met Gln His Gly Pro Ile Glu Trp Ile Gly Leu  
                   85                    90                    95  
 30  
 His Arg His His His Leu His Ser Asp Gln Asp Val Asp His His Asp  
                   100                    105                    110  
 35  
 Ser Asn Lys Gly Phe Leu Trp Ser His Phe Leu Trp Met Ile Tyr Glu  
                   115                    120                    125  
 40  
 Ile Pro Ala Arg Thr Glu Val Asp Lys Phe Thr Arg Asp Ile Ala Gly  
                   130                    135                    140  
 45  
 Asp Pro Val Tyr Arg Phe Phe Asn Lys Tyr Phe Phe Gly Val Gln Val  
 145                    150                    155                    160  
 50  
 Leu Leu Gly Val Leu Leu Tyr Ala Trp Gly Glu Ala Trp Val Gly Asn  
                   165                    170                    175  
 55  
 Gly Trp Ser Phe Val Val Trp Gly Ile Phe Ala Arg Leu Val Val Val

EP 0 736 598 B1

	180	185	190
5	Tyr His Val Thr Trp Leu Val Asn Ser Ala Thr His Lys Phe Gly Tyr		
	195	200	205
10	Arg Ser His Glu Ser Gly Asp Gln Ser Thr Asn Cys Trp Trp Val Ala		
	210	215	220
15	Leu Leu Ala Phe Gly Glu Gly Trp His Asn Asn His His Ala Tyr Gln		
	225	230	235 240
20	Tyr Ser Ala Arg His Gly Leu Gln Trp Trp Glu Phe Asp Leu Thr Trp		
	245	250	255
25	Leu Ile Ile Cys Gly Leu Lys Lys Val Gly Leu Ala Arg Lys Ile Lys		
	260	265	270
30	Val Ala Ser Pro Asn Asn		
	275		

(2) INFORMATION FOR SEQ ID NO: 5:

35 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 18 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- 40 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "synthetic DNA"

45

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

50

ATGACAATTG CTA CTTCA

18

(2) INFORMATION FOR SEQ ID NO: 6:

55 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 15 base pairs
- (B) TYPE: nucleic acid

- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "synthetic DNA"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

GCTCTGGGGT TGTTG

15

(2) INFORMATION FOR SEQ ID NO: 7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 15 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "synthetic DNA"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

CAACAACCCC AGAGC

15

(2) INFORMATION FOR SEQ ID NO: 8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 13 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "synthetic DNA"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

TGTGTTTTTG CCA

13

(2) INFORMATION FOR SEQ ID NO: 9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 17 base pairs

- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

5 (ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "synthetic DNA"

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

ACGTCATGGC CTGCAGT

17

15

(2) INFORMATION FOR SEQ ID NO: 10:

(i) SEQUENCE CHARACTERISTICS:

- 20 (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

25 (ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "synthetic DNA"

30 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:

CGCGGATCCT TAGTTGTTTG GAGACG

26

35

(2) INFORMATION FOR SEQ ID NO: 11:

(i) SEQUENCE CHARACTERISTICS:

- 40 (A) LENGTH: 25 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

45 (ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "synthetic DNA"

50 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:

CATATGACCC TTGCTATCCG ACCCA

25

55

(2) INFORMATION FOR SEQ ID NO: 12:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 29 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

5

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "synthetic DNA"

10

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12:

AGCTTGGGTC GGATAGCAAG GGTCATATG

29

15

### Claims

20

1. A polynucleotide molecule encoding a protein having the following (1) to (4):

- (1) having activities of desaturating lipid-bound fatty acids at the  $\Delta 9$  position,
- (2) reacting with the 16:0- and 18:0-saturated molecular species,
- (3) reacting at the position of sn-1 and sn-2,
- (4) and comprising the amino acid sequence shown in SEQ ID NO:4 or the amino and sequence shown in SEQ ID NO:4 in which part of the amino acid sequence has been deleted, substituted and/or added.

25

2. The polynucleotide molecule of claim 1, wherein said polynucleotide molecule is present in the genus *Anacystis*.

30

3. The polynucleotide molecule of any one of claims 1 or 2, wherein said protein comprises the amino acid sequence shown in SEQ ID NO:4.

4. The polynucleotide molecule of any one of claims 1 or 2, wherein the polynucleotide molecule comprises the nucleotide sequence of SEQ ID NO:3.

35

5. A vector which contains the polynucleotide according to any one of claims 1-4:

6. A plant cell transformed with the polynucleotide molecule according to any one of claims 1-4.

40

7. A method for creating a plant by regenerating the plant cell of claim 6 to produce a mature plant.

8. A plant transformed with the polynucleotide molecule according to any one of claims 1-4.

45

9. A method for creating a plant having an improved chilling-resistance, which comprises introducing a polynucleotide molecule according to any one of claims 1-4.

### Patentansprüche

50

1. Polynukleotidmolekül kodierend ein Protein, das die Merkmale (1) bis (4) aufweist:

- (1) fettgebundene Fettsäuren an der  $\Delta 9$ -Position entsättigt,
- (2) mit 16:0- und 18:0-gesättigten molekularen Arten reagiert,
- (3) an der Position sn-1 und sn-2 reagiert und
- (4) die in SEQ ID NO: 4 gezeigte Aminosäuresequenz oder die in SEQ ID NO: 4 gezeigte Aminosäuresequenz, wobei Teile der Aminosäuresequenz deletiert, substituiert und/oder zugegeben wurden, umfaßt.

55

2. Polynukleotidmolekül nach Anspruch 1, wobei das Polynukleotidmolekül in der Gattung *Anacystis* vorhanden ist.

3. Polynukleotidmolekül nach einem der Ansprüche 1 oder 2, wobei das Protein die in SEQ ID NO: 4 gezeigte Aminosäuresequenz umfaßt.
- 5 4. Polynukleotidmolekül nach einem der Ansprüche 1 oder 2, wobei das Polynukleotidmolekül die Nukleotidsequenz SEQ ID NO: 3 umfaßt.
- 10 5. Vektor, der das Polynukleotid nach einem der Ansprüche 1 bis 4 enthält.
6. Pflanzenzelle, die mit dem Polynukleotidmolekül nach einem der Ansprüche 1 bis 4 transformiert ist.
7. Verfahren zum Erzeugen einer Pflanze durch Wiedergewinnung der Pflanzenzelle nach Anspruch 6, um eine reife Pflanze zu erzeugen.
- 15 8. Pflanze, die mit dem Polynukleotidmolekül nach einem der Ansprüche 1 bis 4 transformiert ist.
9. Verfahren zum Erzeugen einer Pflanze mit einer verbesserten Kälteresistenz, das das Einführen eines Polynukleotidmoleküls nach einem der Ansprüche 1 bis 4 umfaßt.

20 **Revendications**

1. Molécule polynucléotide codant une protéine présentant les points (1) à (4) suivants :
- 25 (1) possède des activités de désaturation des acides gras liés aux lipides à la position  $\Delta 9$ ,
- (2) réagit avec les espèces moléculaires saturées 16 : 0 - et 18 : 0,
- (3) réagit sur la position de sn-1 et de sn-2,
- 30 (4) et comprend la séquence d'acides aminés montrée en SEQ ID NO : 4 ou la séquence d'acides aminés montrée en SEQ ID NO : 4 dans laquelle, une partie de la séquence des acides aminés a été supprimée, substituée et/ou ajoutée.
- 35 2. Molécule nucléotide de la revendication 1, dans laquelle ladite molécule nucléotide est présente dans le genre *Anacystis*.
3. Molécule polynucléotide de l'une quelconque des revendications 1 ou 2, dans laquelle ladite protéine comprend la séquence d'acides aminés montrée en SEQ ID NO : 4.
- 40 4. Molécule polynucléotide de l'une quelconque des revendications 1 ou 2, dans laquelle la molécule polynucléotide comprend la séquence nucléotide de SEQ ID NO : 3.
5. Vecteur qui contient le polynucléotide selon l'une quelconque des revendications 1-4.
- 45 6. Cellule végétale transformée par la molécule polynucléotide selon l'une quelconque des revendications 1-4.
7. Procédé pour créer une plante en régénérant la cellule végétale de la revendication 6 pour produire une plante adulte.
- 50 8. Plante transformée par la molécule polynucléotide selon l'une quelconque des revendications 1-4.
9. Procédé pour créer une plante ayant une résistance améliorée au refroidissement, qui comprend l'introduction d'une molécule polynucléotide selon l'une quelconque des revendications 1-4.

FIG. 1

	10	20	30	40	50	60
des9var	ALGLLLLYLGGWSFVWGVFFRIVWVYHCTWLVNSATHKFGYRTYDAGDRSTNCWVAVL					
MSCD2	LVPWYCWGETFVNSLCVSTFLRYAVVLNATWLVNSAAHLYGYRYPYDKNISSRENILVSMG					
	240	250	260	270	280	290
des9var	VFGEG					
	:X					
MSCD2	AVGER					



FIG. 2




FIG. 3




FIG. 4

```

                10      20      30      40      50      60
des9nid      MTLAIRPKLAFNWP T ALFMVAIHIGALLAFLPANFNWPAVGVMVALYYITGCGGITLGWH
                X::.. ..: .:>::: .. . . . . :::.. .::: : :
MSCD2        DDEGPPPKLEYVWRNIILMALLHLGALYGITLVPSCKLYTCLFAYLYYVISALGITAGAH
60           70           80           90           100          110
                70           80           90           100          110          120
des9nid      RLISHRSFEVPKWLEYVLFVCGTLAMQHGPIEWIGLHRHHHLHSDQDQVDHSDSNXGFLWS
                :::..... : ..:>::: ..:>::: ::: ::: : ..:>:::.....:
MSCD2        RLWSHRTYKARLPLRLFLIIANTMAFQNDVYEWARHRAHHKFSETHADPHNSRRGFFFS
120          130          140          150          160          170
                130          140          150          160          170
des9nid      HFLW-MIYEIPA-RTEVDKPTRDIAGDPVYRFFNKYFFGVQVLLGVLLYAWGEAWVGNW
                X : . . . :: . . . : . . . . . : . . . . . : . . . : . :
MSCD2        HVGWLLVRKHPAVKEXGGKLDMSDLKAEKLYMFQRRYKPDLLMCFVLP TLPWYCWGE
180          190          200          210          220          230
                180          190          200          210          220          230
MSCD2        SFV---VWGI FARLVVYHYVTLVNSATHKFGYRSHESGDQSTNCWWVALLAFGEGWHNN
                .:: . . : : .:: . : . . . . . : : . . . . . : . . . : : .::
MSCD2        TFVNSLCVSTFLRYAVVLNATWLVNSAAHLYGYRKYDKNISSRENI LVS MGAVGERFHNY
240          250          260          270          280          290
                240          250          260          270
des9nid      HHAYQYSARHGLOWWEOFDTWLIICGLKKVGLARKIKVASPNN
                :::..: . . : . . . : . . : . . : . . : . . : . . : . . : . . : . . :
MSCD2        HHAFPYDYSASEYRWHINFTTFIDCMALLGLAYDRKRVSRAA
300          310          320          330          340

```

FIG. 5

DESATURASE GENE

CONTROL

