

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 2005319279 B2**

- (54) Title
RNAi inhibition of CTGF for treatment of ocular disorders
- (51) International Patent Classification(s)
A61K 48/00 (2006.01)
- (21) Application No: **2005319279** (22) Date of Filing: **2005.12.19**
- (87) WIPO No: **WO06/069037**
- (30) Priority Data
- | | | |
|-------------------|-------------------|--------------|
| (31) Number | (32) Date | (33) Country |
| 60/638,705 | 2004.12.23 | US |
- (43) Publication Date: **2006.06.29**
- (44) Accepted Journal Date: **2011.08.18**
- (71) Applicant(s)
Alcon, Inc.
- (72) Inventor(s)
Shepard, Allan R.
- (74) Agent / Attorney
Phillips Ormonde Fitzpatrick, 367 Collins Street, Melbourne, VIC, 3000
- (56) Related Art
WO2003/049773 A1
US 2004/0235031 A1

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
29 June 2006 (29.06.2006)

PCT

(10) International Publication Number
WO 2006/069037 A1

(51) International Patent Classification:
A61K 48/00 (2006.01)

(21) International Application Number:
PCT/US2005/046064

(22) International Filing Date:
19 December 2005 (19.12.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/638,705 23 December 2004 (23.12.2004) US

(71) Applicants (for all designated States except US): **ALCON, INC.** [CH/CH]; Bosch 69, PO Box 62, CH-6331 Hunenberg (CH). **PANG, Iok-Hou** [US/US]; 125 Starbridge Lane, Grand Prairie, Texas 75052 (US).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **SHEPARD, Allan R.** [US/US]; 6404 High Brook Drive, Fort Worth, Texas 76132 (US).

(74) Agents: **NORBERG, Gloria L.** et al.; Winstead Sechrest & Minick P.C., P.O. Box 50784, Dallas, Texas 75201 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,

CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:


- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))
- of inventorship (Rule 4.17(iv))

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: RNAI INHIBITION OF CTGF FOR TREATMENT OF OCULAR DISORDERS



(57) Abstract: RNA interference is provided for inhibition of connective tissue growth factor mRNA expression in ocular disorders involving CTGF expression. Ocular disorders involving aberrant CTGF expression include glaucoma, macular degeneration, diabetic retinopathy, choroidal neovascularization, proliferative vitreoretinopathy and wound healing. Such disorders are treated by administering interfering RNAs of the present invention.

WO 2006/069037 A1

**RNAi INHIBITION OF CTGF
FOR TREATMENT OF OCULAR DISORDERS**

FIELD OF THE INVENTION

[0001] The present invention relates to the field of interfering RNA compositions for inhibition of expression of connective tissue growth factor (CTGF) in ocular disorders.

BACKGROUND OF THE INVENTION

[0002] Most ocular disorders are associated with cellular processes including cell proliferation, survival, migration, differentiation, and angiogenesis. CTGF is a secreted cytokine and a central mediator in these cellular processes. In particular, CTGF is known to increase extracellular matrix production primarily via increased deposition of collagen I and fibronectin. Overexpression of CTGF has been implicated as a major causative factor in conditions such as scleroderma, fibroproliferative diseases, and scarring in which there is an overaccumulation of extracellular matrix components.

[0003] An overaccumulation of extracellular matrix materials in the region of the trabecular meshwork (TM) is a hallmark of many forms of glaucoma; such increases are believed to lead to increased resistance to aqueous outflow and, therefore, elevated intraocular pressures. International Patent Application No. PCT/US2003/012521 to Fleenor *et al.* published November 13, 2003 as WO 03/092584 and assigned to Alcon, Inc. describes the elevated presence of CTGF mRNA in glaucomatous TM cells vs. normal TM cells. Thus, it is believed that CTGF plays a role in extracellular matrix production by the trabecular meshwork cells.

[0004] Macular degeneration is the loss of photoreceptors in the portion of the central retina, termed the macula, responsible for high-acuity vision. Degeneration of the macula is associated with abnormal deposition of extracellular matrix components in the membrane between the retinal pigment epithelium and the vascular choroid. This debris-like material is termed drusen. Drusen is observed using a funduscopy eye examination. Normal eyes may have maculas free of drusen, yet drusen may be abundant in the retinal periphery. The presence of soft drusen in the macula, in the absence of any loss of macular vision, is considered an early stage of AMD.

[0005] Choroidal neovascularization commonly occurs in macular degeneration in addition to other ocular disorders and is associated with proliferation of choroidal endothelial cells, overproduction of extracellular matrix, and formation of a fibrovascular subretinal membrane. Retinal pigment epithelium cell proliferation and production of angiogenic factors appears to effect choroidal neovascularization.

[0006] Diabetic retinopathy is an ocular disorder that develops in diabetes due to thickening of capillary basement membranes and lack of contact between pericytes and endothelial cells of the capillaries. Loss of pericytes increases leakage of the capillaries and leads to breakdown of the blood-retina barrier.

[0007] Proliferative vitreoretinopathy is associated with cellular proliferation of cellular and fibrotic membranes within the vitreous membranes and on the surfaces of the retina. Retinal pigment epithelium cell proliferation and migration is common with this ocular disorder. The membranes associated with proliferative vitreoretinopathy contain extracellular matrix components such as collagen types I, II, and IV and fibronectin, and become progressively fibrotic.

[0008] Wound healing disorders may lead to severe ocular tissue damage via activation of inflammatory cells, release of growth factors and cytokines, proliferation and differentiation of ocular cells, increased capillary permeability, alterations in basement membrane matrix composition, increased deposition of extracellular matrix, fibrosis, neovascularization, and tissue remodeling.

[0009] Overexpression of CTGF therefore has been implicated as a major causative factor in these ocular disorders. Current therapies do not directly address the pathogenic mechanism of these disorders.

[0009a] Throughout the description and claims of the specification, the word "comprise" and variations of the word, such as "comprising" and "comprises", is not intended to exclude other additives, components, integers or steps.

[0009b] A reference herein to a patent document or other matter which is given as prior art is not to be taken as an admission that that document or matter was known or that the information it contains was part of the common general knowledge as at the priority date of any of the claims.

SUMMARY OF THE INVENTION

[0010] The present invention is directed to interfering RNAs that target CTGF mRNA and thereby interfere with CTGF mRNA expression. The interfering RNAs of the invention are useful for treating CTGF-related ocular disorders such as glaucoma, macular degeneration, diabetic retinopathy, choroidal neovascularization, proliferative vitreoretinopathy and aberrant wound healing.

[0011] An embodiment of the present invention provides a method of attenuating expression of connective tissue growth factor mRNA in an eye of a subject. The method comprises administering to the eye of the subject a composition comprising an effective amount of interfering RNA such as double-stranded (ds) siRNA or single-stranded (ss) siRNA having a length of 19 to 49 nucleotides and a pharmaceutically acceptable carrier.

[0011a] An embodiment of the present invention provides a method of attenuating expression of connective tissue growth factor mRNA in an eye of a subject, comprising:

administering to the eye of the subject a composition comprising an effective amount of interfering RNA having a length of 19 to 49 nucleotides and a pharmaceutically acceptable carrier, the interfering RNA comprising:

a sense nucleotide sequence, an antisense nucleotide sequence, and a region of at least near-perfect contiguous complementarity of at least 19 nucleotides;

wherein the antisense sequence comprises 3'-TTcccguuuucacguaggca-5' (SEQ ID NO:33), 3'-TTccggagaagacacugaag-5' (SEQ ID NO:31), or 3'-TTccaucauaguagucuauc-5' (SEQ ID NO:28), and

wherein the expression of connective tissue growth factor mRNA is attenuated.

[0011b] An embodiment of the present invention provides a method of attenuating expression of connective tissue growth factor mRNA in an eye of a subject, comprising:

administering to the eye of the subject a composition comprising an effective amount of single-stranded interfering RNA having a length of 19 to 49 nucleotides and a pharmaceutically acceptable carrier,

wherein the single stranded interfering RNA hybridizes under physiological conditions to a portion of mRNA corresponding to SEQ ID NO:1 beginning at nucleotide 379, 691, 801, 901, 932, 937, 969, 986, 1119, 1170, 1201, 1346, 1473, 1478, 1481, 1488, 1626, 1660, or 1666, and the interfering RNA has a region of at least near-perfect complementarity with the hybridizing portion of mRNA corresponding to SEQ ID NO:1,

wherein the expression of connective tissue growth factor mRNA is attenuated.

[0011c] An embodiment of the present invention provides a method of treating a connective tissue growth factor-associated ocular disorder in a subject in need thereof, comprising:

administering to the eye of the subject a composition comprising an effective amount of interfering RNA having a length of 19 to 49 nucleotides and a pharmaceutically acceptable carrier, the interfering RNA comprising:

a sense nucleotide sequence, an antisense nucleotide sequence, and a region of at least near-perfect contiguous complementarity of at least 19 nucleotides;

wherein the antisense sequence hybridizes under physiological conditions to a portion of mRNA corresponding to SEQ ID NO:1, and has a region of at least near-perfect contiguous complementarity of at least 19 nucleotides with the hybridizing portion of mRNA corresponding to SEQ ID NO:1, wherein the antisense sequence is designed to target a nucleotide sequence of mRNA corresponding to SEQ ID NO:1 comprising nucleotide 379, 691, 801, 901, 932, 937, 969, 986, 1119, 1170, 1201, 1346, 1473, 1478, 1481, 1488, 1626, 1660, or 1666, and

wherein the connective tissue growth factor-associated ocular disorder is treated thereby.

[0011d] An embodiment of the present invention provides a method of treating glaucoma, macular degeneration, diabetic retinopathy, choroidal neovascularization, proliferative vitreoretinopathy or aberrant wound healing in a subject in need thereof, comprising:

administering to the eye of the subject a composition comprising an effective amount of single-stranded interfering RNA having a length of 19 to 49 nucleotides and a pharmaceutically acceptable carrier,

wherein the single stranded interfering RNA hybridizes under physiological conditions to a portion of mRNA corresponding to SEQ ID NO:1 beginning at nucleotide

379, 691, 801, 901, 932, 937, 969, 986, 1119, 1170, 1201, 1346, 1473, 1478, 1481, 1488, 1626, 1660, or 1666, and the interfering RNA has a region of at least near-perfect complementarity with the hybridizing portion of mRNA corresponding to SEQ ID NO:1,

wherein the glaucoma, macular degeneration, diabetic retinopathy, choroidal neovascularization, proliferative vitreoretinopathy or aberrant wound healing is treated thereby.

[0012] The double stranded siRNA comprises a sense nucleotide sequence, an antisense nucleotide sequence and a region of at least near-perfect contiguous complementarity of at least 19 nucleotides. Further, the antisense sequence hybridizes under physiological conditions to a portion of mRNA corresponding to SEQ ID NO:1 (the sense strand sequence of DNA for connective tissue growth factor for humans, GenBank reference no. NM_001901), and has a region of at least near-perfect contiguous complementarity of at least 19 nucleotides with the hybridizing portion of mRNA corresponding to SEQ ID NO:1. The administration of such a composition attenuates the expression of connective tissue growth factor mRNA of the eye of the subject.

[0013] The single-stranded siRNA has a length of 19 to 49 nucleotides, hybridizes under physiological conditions to a portion of mRNA corresponding to SEQ ID NO:1 beginning at nucleotide 379, 691, 801, 901, 932, 937, 969, 986, 1119, 1170, 1201, 1346, 1473, 1478, 1481, 1488, 1626, 1660, or 1666, and has a region of at least near-perfect complementarity with the hybridizing portion of mRNA corresponding to SEQ ID NO:1.

[0014] In an embodiment of the invention, the antisense sequence of a double-stranded interfering RNA is designed to target a nucleotide sequence of mRNA corresponding to SEQ ID NO:1 beginning at or

comprising nucleotide 379, 691, 801, 901, 932, 937, 969, 986, 1119, 1170, 1201, 1346, 1473, 1478, 1481, 1488, 1626, 1660, or 1666.

[0015] A further embodiment of the invention is a method of treating a connective tissue growth factor-associated ocular disorder in a subject in need thereof. The method comprises administering to the eye of the subject a composition comprising an effective amount of interfering RNA having a length of 19 to 49 nucleotides and a pharmaceutically acceptable carrier, the interfering RNA comprising a sense nucleotide sequence, an antisense nucleotide sequence, and a region of at least near-perfect contiguous complementarity of at least 19 nucleotides. The antisense sequence hybridizes under physiological conditions to a portion of mRNA corresponding to SEQ ID NO:1, and has a region of at least near-perfect contiguous complementarity of at least 19 nucleotides with the hybridizing portion of mRNA corresponding to SEQ ID NO:1. The connective tissue growth factor-associated ocular disorder is treated thereby.

BRIEF DESCRIPTION OF THE DRAWINGS

[0016] FIG. 1A shows SITOX™ data demonstrating that transfection efficiency of trabecular meshwork cells was not rate-limiting when taken together with data of FIG. 1B. GTM3 cells were transfected with SITOX™ (Dharmacon) transfection control. After 24 hr, trypan blue exclusion was used to determine the number of viable cells remaining in the SITOX™ culture, which reflects the relative transfection efficiency. Open bars: no transfection; Solid bars: with SITOX™.

[0017] FIG. 1B shows a SIGLO™ image of siRNA uptake in GTM3 cells demonstrating that transfection efficiency was not rate-limiting when taken together with data of FIG. 1A. GTM3 cells were transfected with SIGLO™ siRNA (Dharmacon) using LIPOFECTAMINE 2000™. SIGLO™ siRNA uptake was determined after 24 hr using fluorescence microscopy (red irregular shapes). Individual cell nuclei were identified by DAPI (4',6-diamidino-2-phenylindole), a stain for double stranded DNA (blue round areas). As the data of FIG. 1A and the image of FIG. 1B show, nearly all cells were either dead (SITOX™) or fluorescent (SIGLO™).

[0018] FIG. 2A is a schematic showing the CTGF gene exon (boxes) and intron (lines) structure and location of siRNAs S1, S2, and S3 and QPCR primer/probe sets Q1 and Q2 in relation to the GenBank CTGF sequence NM_001901, the sequence of which is provided as SEQ ID NO:1. The sequences of the siRNAs and primer/probe sets are provided in Example 1.

[0019] FIG. 2B shows QPCR amplification of CTGF mRNA using the exon 5 primer/probe set Q2. Using the S1 and S4 siRNAs, no significant knock-down of the CTGF mRNA levels was detected.

[0020] FIG. 2C shows QPCR amplification of CTGF mRNA using the exon 4/5 spanning primer/probe set Q1. Knock-down of CTGF mRNA was observed by each of the siRNAs and an ~90% knock-down was observed with the S2 siRNA.

[0021] FIG. 3 shows a titration study in which various concentrations of the S2 siRNA were tested for efficacy of knock-down of CTGF mRNA levels. CTGF mRNA knock-down was assessed by QPCR

amplification using primer/probe set Q1. An IC_{50} of ~2.5nM was observed in GTM3 cells after 24 hour treatment with 0, 1, 3, 10, 30, and 100 nM S2 siRNA as described in Example 1.

DETAILED DESCRIPTION OF THE INVENTION

[0022] RNA interference, termed "RNAi," is a method for reducing the expression of a target gene that is effected by small single- or double-stranded RNA molecules. Interfering RNAs include small interfering RNAs, either double-stranded or single-stranded (ds siRNAs or ss siRNAs), microRNAs (miRNAs), small hairpin RNAs (shRNAs), and others. While not wanting to be bound by theory, RNA interference appears to occur *in vivo* with the cleavage of dsRNA precursors into small RNAs of about 20 to 25 nucleotides in length. Cleavage is accomplished by RNaseIII-RNA helicase Dicer. The "sense" strand of an siRNA, i.e., the strand that has exactly the same sequence as a target mRNA sequence, is removed, leaving the 'antisense' strand which is complementary to the target mRNA to function in reducing expression of the mRNA. The antisense strand of the siRNA appears to guide a protein complex known as RISC (RNA-induced silencing complex) to the mRNA, which complex then cleaves the mRNA by the Argonaute protein of the RISC, thereby reducing protein production by that mRNA. Interfering RNAs are catalytic and reduction in expression of mRNA can be achieved with substoichiometric amounts of interfering RNAs in relation to mRNA. Reduction in mRNA expression may also occur via transcriptional and translational mechanisms.

[0023] The present invention relates to the use of interfering RNA for inhibition of expression of connective tissue growth factor (CTGF) in ocular disorders. According to the present invention, tissues of the eye, in particular, trabecular meshwork cells of the eye, carry out siRNA silencing, and exogenously provided siRNAs effect silencing. Further, aspects of the present invention have determined that, when using a PCR-based approach to determine the efficacy of siRNA knock-down, the PCR amplification primers should be designed to encompass the siRNA targeting sequence to accurately measure silencing.

[0024] Nucleic acid sequences cited herein are written in a 5' to 3' direction unless indicated otherwise. The term "nucleic acid," as used herein, refers to either DNA or RNA or a modified form thereof comprising the purine or pyrimidine bases present in DNA (adenine "A," cytosine "C," guanine "G," thymine "T") or in RNA (adenine "A," cytosine "C," guanine "G," uracil "U"). Interfering RNAs provided herein may comprise "T" bases, particularly at 3' ends, even though "T" bases do not naturally occur in RNA. "Nucleic acid" includes the terms "oligonucleotide" and "polynucleotide" and can refer to a single stranded molecule or a double stranded molecule. A double stranded molecule is formed by Watson-Crick base pairing between A and T bases, C and G bases, and A and U bases. The strands of a double stranded molecule may have partial, substantial or full complementarity to each other and will form a duplex hybrid, the strength of bonding of which is dependent upon the nature and degree of complementarity of the sequence of bases. A mRNA sequence is readily determined by knowing the sense or antisense strand sequence of DNA encoding therefor. For example, SEQ ID NO:1 provides the sense strand sequence of DNA corresponding to the mRNA for connective tissue growth factor. The

sequence of mRNA is identical to the sequence of the sense strand of DNA with the "T" bases replaced with "U" residues. Therefore, the mRNA sequence of connective tissue growth factor is known from SEQ ID NO:1.

[0025] *Connective tissue growth factor mRNA*: The GenBank database of the National Center for Biotechnology Information at ncbi.nlm.nih.gov provides the corresponding DNA sequence for the messenger RNA of human connective tissue growth factor as reference no. NM_001901, provided below as SEQ ID NO:1. The coding sequence for connective tissue growth factor is from nucleotides 146-1195. SEQ ID NO:1:

```

1 tccagtgacg gagccgcccg gccgacagcc ccgagacgac agcccggcgc gtcccgggcc
61 ccacctccga ccaccgccag cgctccagc cccgcgctcc ccgctcgccg ccaccgcgcc
121 ctccgctccg cccgcagtc caaccatgac cgccgccagt atgggccccg tccgcgtcgc
181 ctctgtggtc ctctcgccc tctgcagccg gccggccgtc ggccagaact gcagcggggc
241 gtgccggtgc ccggacgagc cggcgccgcg ctgccggcg ggcgtgagcc tctgtctgga
301 cggctcgccg tctgcccgc tctgcgcaa gcagctggc gagctgtgca ccgagcgcga
361 cccctcgac ccgacaagg gcctctctg tgaactcggc tccccggca accgcaagat
421 cggcgtgtgc accgcaaag atggtctcc ctgcatctc ggtggtacgg tgtaccgag
481 cggagagtcc ttccagagca gctgcaagta ccagtgcacg tgcctggacg gggcgggtgg
541 ctgcatgccc ctgtgcagca tggacgttc tctgccagc cctgactgcc cctccccgag
601 gaggggtcaag ctgcccggga aatgctgca ggagtgggtg tgtgacgagc ccaaggacca
661 aaccgtggtt gggcctgccc tgcggctta ccgactggaa gacacgttg gccagacc
721 aactatgatt agagccaact gcctgtcca gaccacagag tggagcgcct gttcaagac
781 ctgtgggatg ggcattcca cccgggttac caatgacaac gcctcctgca ggtagagaa
841 gcagagcccg ctgtgatgg ttagccttg cgaagctgac ctggaagaga acattaagaa
901 gggcaaaaag tgcaccgta ctccaaaat ctccaagcct atcaagttg agctttctg
961 ctgaccagc atgaagacat accgagtaa attctgtgga gtatgtaccg acggccgatg
1021 ctgaccccc cacagaacca ccacctgcc ggtggagttc aatgcccctg acggcgaggt
1081 catgaagaag aacatgatgt tcatcaagac ctgtgcctgc cattacaact gtcccggaga
1141 caatgacatc ttgaatcgc tgaactacag gaagatgtac ggagacatgg catgaagcca
1201 gagagtgaga gacattaact cattagactg gaactgaaac tgattcacat ctcatcttc
1261 cgtaaaaatg attcagtag cacaagttat taaatctgt tttctaact gggggaaaag
1321 attccacc aattcaaac attgtgcat tcaaaacaa tagtctatct tcccagaca
1381 ctggttgaa gaatgtaag actgacagt ggaactacat tagtacacag caccagaatg
1441 tatattaagg tgtggctta ggagcagtg gaggtacca gcagaaaggt tagtatcatc
1501 agatagctct tatacagta atatgcctgc tatttgaagt gtaattgaga aggaaaatt
1561 tagcgtgctc actgacctgc ctgtagccc agtgacagct aggatgtgca ttctccagcc
1621 atcaagagac tgagtcaagt tttccttaa gtcagaacag cagactcagc tctgacattc
1681 tgattcgaat gacctgttc agaatcgga atcctgtcga ttagactgga cagctgtgg
1741 caagtgaatt tctgtaaca agccagattt ttaaaaatt atattgaaa tattgtgtg
1801 gtgtgtgtgt gtgtatata atatataat gtacagttat ctaagtaat taaagttgt
1861 ttgtgccttt ttattttgt tttaatgct ttgatattc aatgtagcc tcaattctg
1921 aacaccatag gtagaatgta aagctgtct gatcgtcaa agcatgaaat ggatacttat
1981 atgaaattc tctcagatag aatgacagtc cgtcaaaaca gattgttgc aaaggggagg
2041 catcagtgct ctggcagc tgatttctag gtaggaaatg tggtagtca cgctcacttt
2101 taatgaacaa atggcctta taaaaactg agtgactcta tatagctgat cagtttttc
2161 acctggaagc attgtttct acttgatata gactgtttt cggacagttt atttgtgag
2221 agtgtgacca aaagttacat gttgcacct ttctagttga aaataaagta tatttttct
2281 aaaaaaaaaa aaaaacgaca gcaacggaat tc.

```

Equivalents of the above cited CTGF mRNA sequence are alternative splice forms, allelic forms, or a cognate thereof. A cognate is a connective tissue growth factor mRNA from another mammalian species that is homologous to SEQ ID NO:1. CTGF nucleic acid sequences related to SEQ ID NO:1 are those having GenBank accession numbers AK092280, AK125220, AY395801, AY550024, BT019794, BT019795, CR541759, M92934, U14750, and X78947, and the sequence of SEQ ID NO:1 of U.S. 5,585,270, incorporated by reference herein.

[0026] *Attenuating expression of an mRNA:* The phrase, "attenuating expression of an mRNA," as used herein, means administering an amount of interfering RNA to effect a reduction of the full mRNA transcript levels of a target gene in a cell, thereby decreasing translation of the mRNA into protein as compared to a control RNA having a scrambled sequence. The reduction in expression of the mRNA is commonly referred to as "knock-down" of mRNA. Knock-down of expression of an amount of between and including an amount of 50% and 100% is contemplated by embodiments herein. However, it is not necessary that such knock-down levels be achieved for purposes of the present invention. Further, two sets of interfering RNAs may be mildly effective at knock-down individually, however, when administered together may be significantly more effective. In one embodiment, an individual ds siRNA is effective at knock-down at an amount of at least up to 70%. In another embodiment, two or more ds siRNAs are together effective at knock-down at an amount of at least up to 70%.

[0027] Knock-down is commonly measured by determining the mRNA levels by Quantitative Polymerase Chain Reaction (QPCR) amplification or by determining protein levels by Western Blot or enzyme linked immunosorbent assay (ELISA). Analyzing the protein level provides an assessment of both mRNA degradation by the RNA Induced Silencing Complex (RISC) as well as translation inhibition. Further techniques for measuring knock-down include RNA solution hybridization, nuclease protection, Northern hybridization, reverse transcription, gene expression monitoring with a microarray, antibody binding, radioimmunoassay, and fluorescence activated cell analysis. A further method of measurement includes overexpressing TGF β 2 which induces CTGF, adding back CTGF siRNA, and then measuring CTGF mRNA/protein knockdown by any of the above-cited methods.

[0028] Inhibition of CTGF is also inferred in a human or mammal by observing an improvement in an ocular disorder. For example, in age related macular degeneration a slowing or reversal of vision loss indicates an inhibition of CTGF and silencing of CTGF mRNA in glaucoma patients leads to lowered intraocular pressure and a delay or prevention of the onset of symptoms in a subject at risk for developing glaucoma..

[0029] Interfering RNA of embodiments of the invention act in a catalytic manner, i.e., interfering RNA is able to effect inhibition of target mRNA in substoichiometric amounts. As compared to antisense therapies, significantly less interfering RNA is required to provide a therapeutic effect.

[0030] *Double-stranded interfering RNA:* Double stranded interfering RNA (also referred to as ds siRNA), as used herein, has a sense nucleotide sequence and an antisense nucleotide sequence, the sense and antisense sequence comprising a region of at least near-perfect contiguous complementarity of at least 19 nucleotides. The length of the interfering RNA comprises 19 to 49 nucleotides, and may

comprise a length of 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, or 49 nucleotides. The antisense sequence of the ds siRNA hybridizes under physiological conditions to a portion of mRNA corresponding to SEQ ID NO:1, and has a region of at least near-perfect contiguous complementarity of at least 19 nucleotides with the hybridizing portion of mRNA corresponding to SEQ ID NO:1.

[0031] The antisense strand of the siRNA is the active guiding agent of the siRNA in that the antisense strand binds to a RISC complex within a cell, and guides the bound complex to bind with specificity to the mRNA at a sequence complementary to the sequence of the antisense RNA, thereby allowing subsequent cleavage of the mRNA by the bound complex.

[0032] Techniques for selecting target sequences for siRNAs are provided by Tuschl, T. *et al.*, "The siRNA User Guide," revised May 6, 2004, available on the Rockefeller University web site, by Technical Bulletin #506, "siRNA Design Guidelines," Ambion Inc. at Ambion's web site, by the Invitrogen web site using search parameters of min 35%, max 55% G/C content, and by the Dharmacon web site. The target sequence may be located in the coding region or a 5' or 3' untranslated region of the mRNA.

[0033] An embodiment of a DNA target sequence for CTGF is present at nucleotides 1488 to 1506 of SEQ ID NO:1:

5'- ggtagtagtatcatcagatag -3' SEQ ID NO:18. nt 1488.

A double stranded siRNA of the invention for targeting a corresponding mRNA sequence of SEQ ID NO:18 and having a 3'UU overhang on each strand is:

5'- gguuaguaucaucagauagUU-3' SEQ ID NO:25
3' - UUccaucauaguagucuauc -5' SEQ ID NO:26.

The 3' overhang may have a number of "U" residues, for example, a number of "U" residues between and including 2, 3, 4, 5, and 6. The 5' end may also have a 5' overhang of nucleotides. A double stranded siRNA of the invention for targeting a corresponding mRNA sequence of SEQ ID NO:18 and having a 3'TT overhang on each strand is:

5'- gguuaguaucaucagauagTT-3' SEQ ID NO:27
3' - TTccaucauaguagucuauc -5' SEQ ID NO:28.

The strands of a double-stranded siRNA may be connected by a hairpin loop to form a single stranded siRNA as follows:

5'- gguuaguaucaucagauagUUNNN \ N
3' - UUccaucauaguagucuaucNNNNN / SEQ ID NO:29.

N is a nucleotide A, T, C, G, U, or a modified form known by one of ordinary skill in the art. The number of nucleotides N is a number between and including 3 to 23, or 5 to 15, or 7 to 13, or 4 to 9, or 9 to 11, or the number of nucleotides N is 9.

[0034] Table 1 lists examples of CTGF DNA target sequences of SEQ ID NO:1 from which siRNAs of the present invention are designed in a manner as set forth above.

Table 1. CTGF Target Sequences for siRNAs

Target Sequence	# of Starting Nucleotide with reference to SEQ ID NO:1	SEQ ID NO:
GGGCCTCTTCTGTGACTTC	379	2
CCGACTGGAAGACACGTTT	691	3
CCCGGGTTACCAATGACAA	801	4
GGGCAAAAAGTGCATCCGT	901	5
TCCAAGCCTATCAAGTTTGAGCTTT	932	6
GCCTATCAAGTTTGAGCTT	937	7
GCATGAAGACATACCGAGCTAAATT	969	8
GCTAAATTCTGTGGAGTAT	986	9
GCCATTACAACGTGCCCGGAGACAA	1119	10
GGAAGATGTACGGAGACAT	1170	11
GAGAGTGAGAGACATTAACCTATTA	1201	12
GCCATGTCAAACAAATAGTCTATCT	1346	13
GGGTACCAGCAGAAAGGTT	1473	14
CCAGCAGAAAGGTTAGTAT	1478	15
GCAGAAAGGTTAGTATCAT	1481	16
GCAGAAAGGTTAGTATCATCAGATA	1481	17
GGTTAGTATCATCAGATAG	1488	18
GGTTAGTATCATCAGATAGCTCTTA	1488	19
GAGACTGAGTCAAGTTGTTCTTAA	1626	20
GCAGACTCAGCTCTGACAT	1660	21
TCAGCTCTGACATTCTGATTCTGAAT	1666	22
TCCTGTCGATTAGACTGGACAGCTT	1712	23
GCTTGTGGCAAGTGAATTT	1733	24

As cited in the examples above, one of skill in the art is able to use the target sequence information provided in Table 1 to design interfering RNAs having a length shorter or longer than the sequences provided in Table 1 by referring to the sequence position in SEQ ID NO:1 and adding or deleting nucleotides complementary or near complementary to SEQ ID NO:1.

[0035] The target RNA cleavage reaction guided by ds or ss siRNAs is highly sequence specific. In general, siRNA containing a sense nucleotide sequence identical to a portion of the target mRNA and an antisense portion exactly complementary to the sense sequence are siRNA embodiments for inhibition of CTGF mRNA. However, 100% sequence complementarity between the antisense strand of siRNA and the target mRNA is not required to practice the present invention. Thus the invention allows for sequence variations that might be expected due to genetic mutation, strain polymorphism, or evolutionary divergence. For example, siRNA sequences with insertions, deletions, or single point mutations relative to the target sequence are effective for inhibition.

[0036] The antisense sequence of the siRNA has at least near-perfect contiguous complementarity of at least 19 nucleotides with the target sequence of the mRNA. "Near-perfect," as used herein, means the antisense sequence of the siRNA is "substantially complementary to," and the sense sequence of the

siRNA is "substantially identical" to at least a portion of the target mRNA. "Identity," as known by one of ordinary skill in the art, is the degree of sequence relatedness between nucleotide sequences as determined by matching the order of nucleotides between the sequences. In one embodiment, antisense RNA having 80% and between 80% up to 100% complementarity to the target mRNA sequence are considered near-perfect complementarity and may be used in the present invention. "Perfect" contiguous complementarity is standard Watson-Crick base pairing of adjacent base pairs. "At least near-perfect" contiguous complementarity includes "perfect" complementarity as used herein. Computer methods for determining identity or complementarity are designed to provide the greatest degree of matching of nucleotide sequences, for example, BLASTP and BLASTN (Altschul, S.F., *et al.* (1990) *J. Mol. Biol.* 215:403-410), and FASTA.

[0037] The target sequence of SEQ ID NO:1 may be in the 5' or 3' untranslated regions of the mRNA as well as in the coding region of the mRNA.

[0038] One or both of the strands of double-stranded interfering RNA may have a 3' overhang of from 1 to 6 nucleotides which may be ribonucleotides or deoxyribonucleotides or a mixture thereof. The nucleotides of the overhang are not base-paired. In one embodiment of the invention, the interfering ds RNA comprises a 3' overhang of TT or UU.

[0039] The sense and antisense strands of the double stranded siRNA may be in a duplex formation of two single strands as described above or may be a single molecule where the regions of complementarity are base-paired and are covalently linked by a hairpin or loop so as to form a single strand. It is believed that the hairpin is cleaved intracellularly by a protein termed Dicer to form an interfering RNA of two individual base-paired RNA molecules.

[0040] Interfering RNAs may differ from naturally-occurring RNA by the addition, deletion, substitution or modification of one or more nucleotides. Non-nucleotide material may be bound to the interfering RNA, either at the 5' end, the 3' end, or internally. Such modifications are commonly designed to increase the nuclease resistance of the interfering RNAs, to improve cellular uptake, to enhance cellular targeting, to assist in tracing the interfering RNA, or to further improve stability. For example, interfering RNAs may comprise a purine nucleotide at the ends of overhangs. Conjugation of cholesterol to the 3' end of the sense strand of a ds siRNA molecule by means of a pyrrolidine linker, for example, also provides stability to an siRNA. Further modifications include a 3' terminal biotin molecule, a peptide known to have cell-penetrating properties, a nanoparticle, a peptidomimetic, a fluorescent dye, or a dendrimer, for example.

[0041] Nucleotides may be modified on their base portion, on their sugar portion, or on the phosphate portion of the molecule and function in embodiments of the present invention. Modifications include substitutions with alkyl, alkoxy, amino, deaza, halo, hydroxyl, thiol groups, or a combination thereof, for example. Nucleotides may be substituted with analogs with greater stability such as replacing U with 2'-deoxy-U, or having a sugar modification such as a 2'-OH replaced by a 2' amino or 2' methyl group, 2'-methoxyethyl groups, or a 2'-O, 4'-C methylene bridge, for example. Examples of a purine or pyrimidine analog of nucleotides include a xanthine, a hypoxanthine, an azapurine, a methylthioadenine,

7-deaza-adenosine and O- and N-modified nucleotides. The phosphate group of the nucleotide may be modified by substituting one or more of the oxygens of the phosphate group with nitrogen or with sulfur (phosphorothioates).

[0042] There may be a region of the antisense siRNA that is not complementary to a portion of mRNA corresponding to SEQ ID NO:1. Non-complementary regions may be at the 3', 5' or both ends of a complementary region.

[0043] Interfering RNAs may be synthetically generated, generated by *in vitro* transcription, siRNA expression vectors, or PCR expression cassettes, for example. Interfering RNAs that function well as transfected siRNAs also function well as siRNAs expressed *in vivo*.

[0044] Interfering RNAs are chemically synthesized using protected ribonucleoside phosphoramidites and a conventional DNA/RNA synthesizer and may be obtained from commercial suppliers such as Ambion Inc. (Austin, Texas), Invitrogen (Carlsbad, CA), or Dharmacon (Lafayette, Colo., USA), for example. Interfering RNAs are purified by extraction with a solvent or resin, precipitation, electrophoresis, chromatography, or a combination thereof, for example. Alternatively, interfering RNA may be used with little if any purification to avoid losses due to sample processing.

[0045] Interfering RNA may be provided to a subject by expression from a recombinant plasmid using a constitutive or inducible promoter such as the U6 or H1 RNA pol III promoter, the cytomegalovirus promoter, SP6, T3, or T7 promoter, known to those of ordinary skill in the art. For example, the psiRNA™ from InvivoGen (San Diego, CA) allows production of siRNAs within cells from an RNA pol III promoter. Interfering RNA expressed from recombinant plasmids may be isolated by standard techniques.

[0046] A viral vector for expression of interfering RNA may be derived from adenovirus, adeno-associated virus, vaccinia virus, retroviruses (lentiviruses, Rhabdoviruses, murine leukemia virus, for example), herpes virus, or the like, using promoters as cited above, for example, for plasmids. Selection of viral vectors, methods for expressing the interfering RNA by the vector and methods of delivering the viral vector are within the ordinary skill of one in the art.

[0047] Expression of interfering RNAs is also provided by use of SILENCER EXPRESS™ (Ambion, Austin, Texas) via expression cassettes (SECs) with a human H1, human U6 or mouse U6 promoter by PCR. Silencer expression cassettes are PCR products that include promoter and terminator sequences flanking a hairpin siRNA template. Upon transfection into cells, the hairpin siRNA is expressed from the PCR product and induces specific silencing.

[0048] *Hybridization under Physiological Conditions:* "Hybridization" refers to a technique where single-stranded nucleic acids (DNA or RNA) are allowed to interact so that hydrogen-bonded complexes called hybrids are formed by those nucleic acids with complementary or near-complementary base sequences. Hybridization reactions are sensitive and selective so that a particular sequence of interest is identified in samples in which it is present at low concentrations. The specificity of hybridization (i.e., stringency) is controlled by the concentrations of salt or formamide in the prehybridization and hybridization solutions *in vitro*, for example, and by the hybridization temperature, and are well known in

the art. In particular, stringency is increased by reducing the concentration of salt, increasing the concentration of formamide, or raising the hybridization temperature.

[0049] For example, high stringency conditions could occur at about 50% formamide at 37 °C to 42 °C. Reduced stringency conditions could occur at about 35% to 25% formamide at about 30 °C to 35 °C. Examples of stringency conditions for hybridization are provided in Sambrook, J., 1989, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. Further examples of stringent hybridization conditions include 400 mM NaCl, 40 mM PIPES pH 6.4, 1 mM EDTA, 50 °C or 70 °C for 12-16 hours followed by washing, or hybridization at 70 °C in 1XSSC or 50 °C in 1XSSC, 50% formamide followed by washing at 70 °C in 0.3XSSC, or hybridization at 70 °C in 4XSSC or 50 °C in 4XSSC, 50% formamide followed by washing at 67 °C in 1XSSC. The temperature for hybridization is about 5-10 °C less than the melting temperature (T_m) of the hybrid where T_m is determined for hybrids between 19 and 49 base pairs in length using the following calculation: T_m °C = $81.5 + 16.6(\log_{10}[\text{Na}^+]) + 0.41 (\% \text{ G+C}) - (600/N)$ where N is the number of bases in the hybrid, and $[\text{Na}^+]$ is the concentration of sodium ions in the hybridization buffer.

[0050] In embodiments of the present invention, an antisense strand of an interfering RNA that hybridizes with CTGF mRNA *in vitro* under high stringency conditions will bind specifically *in vivo* under physiological conditions. Identification or isolation of a related nucleic acid that does not hybridize to a nucleic acid under highly stringent conditions is carried out under reduced stringency.

[0051] *Single stranded interfering RNA:* As cited above, interfering RNAs ultimately function as single strands. SS siRNA has been found to effect mRNA silencing, albeit less efficiently than double-stranded RNA. Therefore, embodiments of the present invention also provide for administration of ss siRNA where the single stranded siRNA hybridizes under physiological conditions to a portion of mRNA corresponding to SEQ ID NO:1, and has a region of at least near-perfect contiguous complementarity of at least 19 nucleotides with the hybridizing portion of mRNA corresponding to SEQ ID NO:1. The ss siRNA has a length of 19 to 49 nucleotides as for the ds siRNA cited above. The ss siRNA has a 5' phosphate or is phosphorylated *in situ* or *in vivo* at the 5' position. The term "5' phosphorylated" is used to describe, for example, polynucleotides or oligonucleotides having a phosphate group attached via ester linkage to the C5 hydroxyl of the 5' sugar (e.g., the 5' ribose or deoxyribose, or an analog of same). The ss siRNA may have a mono-, di-, or triphosphate group.

[0052] SS siRNAs are synthesized chemically or via vectors as for ds siRNAs. 5' Phosphate groups may be added via a kinase, or a 5' phosphate may be the result of nuclease cleavage of an RNA. Delivery is as for ds siRNAs. In one embodiment, ss siRNAs having protected ends and nuclease resistant modifications are administered for silencing. SS siRNAs may be dried for storage or dissolved in an aqueous solution. The solution may contain buffers or salts to inhibit annealing or for stabilization.

[0053] *Hairpin interfering RNA:* A hairpin interfering RNA is single-stranded and contains both the sense and antisense sequence within the one strand. For expression by a DNA vector, the corresponding DNA oligonucleotides of at least 19-nucleotides corresponding to the sense siRNA sequence are linked to

its reverse complementary antisense sequence by a short spacer. If needed for the chosen expression vector, 3' terminal T's and nucleotides forming restriction sites may be added. The resulting RNA transcript folds back onto itself to form a stem-loop structure.

[0054] *Mode of administration:* Interfering RNA may be delivered directly to the eye by ocular tissue injection such as periocular, conjunctival, sub-Tenons, intracameral, intravitreal, sub-retinal, retrobulbar, or intracanalicular injections; by direct application to the eye using a catheter or other placement device such as a retinal pellet, intraocular insert, suppository or an implant comprising a porous, non-porous, or gelatinous material; by topical ocular drops or ointments; by a slow release device in the cul-de-sac or implanted adjacent to the sclera (transscleral) or within the eye. Intracameral injection may be through the cornea into the anterior chamber to allow the agent to reach the trabecular meshwork. Intracanalicular injection may be into the venous collector channels draining Schlemm's canal or into Schlemm's canal.

[0055] *Subject:* A subject in need of treatment for an ocular disorder or at risk for developing an ocular disorder is a human or other mammal having a condition or at risk of having a condition associated with expression or activity of CTGF, i.e., a CTGF-associated ocular disorder. Such an ocular disorder may include, for example, glaucoma, macular degeneration, diabetic retinopathy, choroidal neovascularization, proliferative vitreoretinopathy, wound healing, and conditions with excessive scarring, with endothelial cell proliferation, or fibroproliferation. Ocular structures associated with such disorders may include the retina, choroid, lens, cornea, trabecular meshwork, rod, cone, ganglia, macula, iris, sclera, aqueous chamber, vitreous chamber, ciliary body, optic disc, papilla, or fovea, for example.

[0056] *Formulations and Dosage:* Pharmaceutical formulations comprise an interfering RNA, or salt thereof, of the invention up to 99% by weight mixed with a physiologically acceptable ophthalmic carrier medium such as water, buffer, saline, glycine, hyaluronic acid, mannitol, and the like.

[0057] Interfering RNAs of the present invention are administered as solutions, suspensions, or emulsions. The following are examples of possible formulations embodied by this invention.

	Amount in weight %
Interfering RNA	up to 99; 0.1-99; 0.1 – 50; 0.5 – 10.0
Hydroxypropylmethylcellulose	0.5
Sodium chloride	.8
Benzalkonium Chloride	0.01
EDTA	0.01
NaOH/HCl	qs pH 7.4
Purified water	qs 100 mL

	Amount in weight %
Interfering RNA	up to 99; 0.1-99; 0.1 – 50; 0.5 – 10.0
Phosphate Buffered Saline	1.0
Benzalkonium Chloride	0.01
Polysorbate 80	0.5
Purified water	q.s. to 100%

Interfering RNA	up to 99; 0.1-99; 0.1 – 50; 0.5 – 10.0
Monobasic sodium phosphate	0.05
	Amount in weight %
Dibasic sodium phosphate (anhydrous)	0.15
Sodium chloride	0.75
Disodium EDTA	0.05
Cremophor EL	0.1
Benzalkonium chloride	0.01
HCl and/or NaOH	pH 7.3-7.4
Purified water	q.s. to 100%

	Amount in weight %
Interfering RNA	up to 99; 0.1-99; 0.1 – 50; 0.5 – 10.0
Phosphate Buffered Saline	1.0
Hydroxypropyl- β -cyclodextrin	4.0
Purified water	q.s. to 100%

[0058] Generally, an effective amount of the interfering RNA of embodiments of the invention comprises an intercellular concentration at or near the ocular site of from 200pM to 100 nM, or from 1 nM to 50 nM, or from 5 nM to about 25 nM. Topical compositions are delivered to the surface of the eye one to four times per day according to the routine discretion of a skilled clinician. The pH of the formulation is about pH 4-9, or pH 4.5 to pH 7.4.

[0059] While the precise regimen is left to the discretion of the clinician, interfering RNA may be administered by placing one drop in each eye one to four times a day, or as directed by the clinician. An effective amount of a formulation may depend on factors such as the age, race, and sex of the subject, or the severity of the ocular disorder, for example. In one embodiment, the interfering RNA is delivered topically to the eye and reaches the trabecular meshwork, retina or optic nerve head at a therapeutic dose thereby ameliorating a CTGF-associated disease process.

[0060] *Acceptable carriers:* An ophthalmically acceptable carrier refers to those carriers that cause at most, little to no ocular irritation, provide suitable preservation if needed, and deliver one or more interfering RNAs of the present invention in a homogenous dosage. An acceptable carrier for administration of interfering RNA of embodiments of the present invention include the Mirus TransIT[®]-TKO siRNA Transfection Reagent (Mirus Corporation, Madison, Wisconsin), LIPOFECTIN[®], lipofectamine, OLIGOFECTAMINE[™] (Invitrogen, Carlsbad, CA), CELLFECTIN[®], DHARMAFECT[™] (Dharmacon, Chicago, IL) or polycations such as polylysine, liposomes, or fat-soluble agents such as cholesterol. Liposomes are formed from standard vesicle-forming lipids and a sterol, such as cholesterol, and may include a targeting molecule such as a monoclonal antibody having binding affinity for endothelial cell surface antigens, for example. Further, the liposomes may be PEGylated liposomes.

[0061] For ophthalmic delivery, an interfering RNA may be combined with ophthalmologically acceptable preservatives, co-solvents, surfactants, viscosity enhancers, penetration enhancers, buffers, sodium chloride, or water to form an aqueous, sterile ophthalmic suspension or solution. Ophthalmic solution formulations may be prepared by dissolving the inhibitor in a physiologically acceptable isotonic

aqueous buffer. Further, the ophthalmic solution may include an ophthalmologically acceptable surfactant to assist in dissolving the inhibitor. Viscosity building agents, such as hydroxymethyl cellulose, hydroxyethyl cellulose, methylcellulose, polyvinylpyrrolidone, or the like, may be added to the compositions of the present invention to improve the retention of the compound.

[0062] In order to prepare a sterile ophthalmic ointment formulation, the interfering RNA is combined with a preservative in an appropriate vehicle, such as mineral oil, liquid lanolin, or white petrolatum. Sterile ophthalmic gel formulations may be prepared by suspending the interfering RNA in a hydrophilic base prepared from the combination of, for example, CARBOPOL®-940 (BF Goodrich, Charlotte, NC), or the like, according to methods known in the art for other ophthalmic formulations. VISCOAT® (Alcon Laboratories, Inc., Fort Worth, TX) may be used for intraocular injection, for example. Other compositions of the present invention may contain penetration enhancing agents such as cremephor and TWEEN® 80 (polyoxyethylene sorbitan monolaureate, Sigma Aldrich, St. Louis, MO), in the event the interfering RNA is less penetrating in the eye.

[0063] *Kits:* Embodiments of the present invention provide a kit that includes reagents for attenuating the expression of a CTGF mRNA in a cell. The kit contains a DNA template that has two different promoters such as a T7 promoter, a T3 promoter or an SP6 promoter, each operably linked to a nucleotide sequence that encodes two complementary single-stranded RNAs corresponding to an interfering RNA. RNA is transcribed from the DNA template and is annealed to form a double-stranded RNA effective to attenuate expression of the target mRNA. The kit optionally contains amplification primers for amplifying the DNA sequence from the DNA template and nucleotide triphosphates (i.e., ATP, GTP, CTP and UTP) for synthesizing RNA. Optionally, the kit contains two RNA polymerases, each capable of binding to a promoter on the DNA template and effecting transcription of the nucleotide sequence to which the promoter is operably linked, a purification column for purifying single-stranded RNA, such as a size exclusion column, one or more buffers, for example, a buffer for annealing single-stranded RNAs to yield double stranded RNA, and RNase A or RNase T for purifying double stranded RNA.

Example 1 **Interfering RNA for Silencing CTGF in Trabecular Meshwork Cells** **and Criteria for Measuring Silencing**

[0064] The present study examines the ability of CTGF interfering RNA to knock-down the levels of endogenous CTGF expression in human trabecular meshwork (TM) cells. The present study also provides criteria for determining the efficacy of interfering RNA on mRNA levels when QPCR primers are used for measurement.

[0065] Transfection of a transformed human TM cell line designated GTM3 or HTM-3 (see Pang, I.H. *et al.*, 1994. *Curr. Eye Res.* 13:51-63) was accomplished using standard *in vitro* concentrations of CTGF interfering RNA (100 nM) and LIPOFECTAMINE™ 2000 (Invitrogen, Carlsbad, California) at a 1:1 (w/v) ratio. A pool of commercially designed interfering RNAs of unknown sequence (*siGENOME*

SMARTPOOL[®] CTGF interfering RNA (designated siRNA S4 herein), Dharmacon, Lafayette, Colorado) was used to target CTGF. Scrambled and lamin A/C siRNA (Dharmacon) were used as controls.

[0066] Control experiments resulted in close to 90% knock-down efficiency of lamin A/C using lamin A/C interfering RNA when compared to the scrambled interfering RNA control. Initial studies showed an efficiency of knock-down of CTGF of about 20-30% when using *siGENOME SMARTPOOL*[®] CTGF siRNA M-012633-00-0020 (siRNA S4) using primer/probe set Q2 directed to the CTGF mRNA 3'UTR in exon 5 (FIG. 2B). Q2 is a QPCR TAQMAN[®] primer/probe sets from ABI (Applied Biosystems, Foster City, CA).

[0067] To determine the reason for the poor CTGF siRNA efficacy, several variables were tested. Dose response with the CTGF interfering RNA was tested to determine if a suboptimal interfering RNA concentration or a suboptimal interfering RNA:lipid ratio was being used. Resultant data indicated poor CTGF mRNA knock-down regardless of the interfering RNA concentration or interfering RNA:lipid ratio employed. Given the importance of cellular uptake on siRNA activity and the inherent difficulty of transfecting TM cells, the TM cell transfection efficiency was determined under the above-cited conditions. Transfection efficiency was examined as a reflection of either cell death induced by SITOXTM (Dharmacon) delivery to the cell cytoplasm or cell fluorescence as measured by cytoplasmic fluorescence with SIGLO[™] (Dharmacon). In both cases, nearly all cells were either dead (FIG. 1A; SITOXTM) or fluorescent (FIG. 1B; SIGLO[™]), suggesting that the transfection efficiency was nearly quantitative and not the rate-limiting step in the process.

[0068] Further, three additional individual CTGF siRNA sequences from Ambion Inc. (Austin, Texas) designated siRNA S1, S2, and S3 were tested in combination with two different QPCR TAQMAN[®] primer/probe sets designated Q2 and Q1 (ABI, Applied Biosystems, Foster City, CA). The target sequences for Ambion siRNAs are as follows using GenBank reference sequence number NM_001901 for nucleotides (nts) of CTGF:

target for S1: (nts 379-397):	gggcctctctgtgacttc	SEQ ID NO:2
target for S2: (nts 901-919):	gggcaaaaagtgcacccgt	SEQ ID NO:5
target for S3: (nts 1488-1506):	ggttagtatcatcagatag	SEQ ID NO:18

[0069] Double stranded siRNA with a 3'TT overhang on each strand for each of the above targeted sequences are:

siRNA S1:

5'-gggccucuucugugacuucTT-3'	SEQ ID NO:30
3'-Ttcccgagaagacacugaag-5'	SEQ ID NO:31

siRNA S2:

5'-gggcaaaaagugcauccguTT-3'	SEQ ID NO:32
3'-TTcccguuuuucacguaggca-5'	SEQ ID NO:33

siRNA S3:

5'-gguuaguucaucauagTT-3' SEQ ID NO:27

3'-TTccaucauaguagucuauc-5' SEQ ID NO:28

[0070] The QPCR Q1 primer is a proprietary sequence from ABI ASSAY ON DEMAND™ Hs00170014_m1 (Applied BioSystems).

The QPCR Q2 forward primer has the sequence:

5'-CAGCTCTGACATTCTGATTGAA-3' SEQ ID NO:34

and the Q2 reverse primer has the sequence:

5'-TGCCACAAGCTGTCCAGTCT-3' SEQ ID NO:35

The Q2 probe has the sequence:

5'-AATCGACAGGATTCCGATTCCCTGAACAGTG-3' SEQ ID NO:36 and has an FAM group at the 5' end (6-carboxyfluorescein) and a TAMRA group at the 3' end (Applied Biosystems).

[0071] The location of the primer/probe sets in relation to the siRNA target sites for the individual siRNAs is shown in FIG. 2A. Also shown in the schematic of FIG. 2A are the CTGF gene exon (boxes) and intron (lines) structure and location of siRNAs S1, S2, and S3 and QPCR primer/probe sets Q1 and Q2 in relation to the GenBank CTGF sequence NM_001901, the sequence of which is provided as SEQ ID NO:1.

[0072] FIG. 2B shows QPCR amplification of CTGF mRNA using the exon 5 primer/probe set Q2 and siRNAs S1-S4. Using the S1 and S4 siRNAs, no significant knock-down of the CTGF mRNA levels was detected with the Q2 primer/probe set. Knock-down was demonstrated by siRNAs S2 and S3. The primer/probe set Q2 has closer proximity to the targets of the S2 and S3 siRNAs as compared to the target of the S1 siRNA.

[0073] FIG. 2C shows QPCR amplification of CTGF mRNA using the exon 4/5 spanning primer/probe set Q1. Knock-down of CTGF mRNA was demonstrated by each of the siRNAs and an ~90% knock-down was observed with the S2 siRNA using the Q1 primer/probe set for detection. The primer/probe set Q1 appears to be more efficient at demonstrating knock-down by the siRNAs as compared to the Q2 primer probe set.

[0074] The data of FIG. 2B and FIG. 2C suggest that the particular region amplified using the 3'-UTR-directed primer/probe set Q2 may be relatively stable and thus a poor choice for assessing the cleavage and degradation of the CTGF mRNA by the targeting siRNA. Therefore, siRNA efficacy may be underreported in specific cases where the QPCR amplification region lies outside the siRNA targeting region.

[0075] To reduce the chance of non-specific, off-target effects, the lowest possible siRNA concentration for inhibiting CTGF mRNA expression was determined. CTGF mRNA knock-down was assessed by QPCR amplification using primer/probe set Q1. A dose response of CTGF S2 siRNA in GTM3 cells is

shown in FIG. 3. An IC_{50} of ~ 2.5 nM was observed in GTM3 cells after 24 hour treatment with 0, 1, 3, 10, 30, and 100 nM dose range of S2 siRNA. Data were fitted using GraphPad Prism 4 software (GraphPad Software, Inc., San Diego, CA) with a variable slope, sigmoidal dose response algorithm and a top constraint of 100%.

[0076] The results of this example demonstrate that i) trabecular meshwork cells carry out siRNA silencing, ii) all of the siRNAs cited herein effect a degree of silencing, and iii) when using a PCR-based approach to determine the efficacy of siRNA knock-down, the PCR amplification primers are designed to encompass the siRNA targeting sequence for optimum detection of silencing.

[0077] Cleavage of target mRNA by the RISC endonuclease has been shown to occur near the center of the siRNA targeting sequence (Elbashir, S.M., *et al.*, 2001. *Genes Dev* 15:188-200) and is accomplished by Argonaute RNaseH activity (Liu, J., *et al.*, 2004. *Science* 305:1437-1441). However, complete degradation of the remaining mRNA appears not to be guaranteed. Stable fragments of mRNA may remain following Argonaute cleavage and amplification of one of these fragments by QPCR may underreport the siRNA efficacy as shown herein. The present invention provides an embodiment where QPCR primer sets encompass the siRNA target sequence to ensure optimum siRNA efficiency readout.

[0078] The references cited herein, to the extent that they provide exemplary procedural or other details supplementary to those set forth herein, are specifically incorporated by reference.

[0079] Those of skill in the art, in light of the present disclosure, will appreciate that obvious modifications of the embodiments disclosed herein can be made without departing from the spirit and scope of the invention. All of the embodiments disclosed herein can be made and executed without undue experimentation in light of the present disclosure. The full scope of the invention is set out in the disclosure and equivalent embodiments thereof. The specification should not be construed to unduly narrow the full scope of protection to which the present invention is entitled.

[0080] As used herein and unless otherwise indicated, the terms "a" and "an" are taken to mean "one", "at least one" or "one or more".

The claims defining the invention are as follows:

1. A method of attenuating expression of connective tissue growth factor mRNA in an eye of a subject, comprising:
 - administering to the eye of the subject a composition comprising an effective amount of interfering RNA having a length of 19 to 49 nucleotides and a pharmaceutically acceptable carrier, the interfering RNA comprising:
 - a sense nucleotide sequence, an antisense nucleotide sequence, and a region of at least near-perfect contiguous complementarity of at least 19 nucleotides;
 - wherein the antisense sequence comprises 3'-TTcccguuuucacguaggca-5' (SEQ ID NO:33), 3'-TTcccgagaagacacugaag-5' (SEQ ID NO:31), or 3'-TTccaucauaguagucuauc-5' (SEQ ID NO:28), and
 - wherein the expression of connective tissue growth factor mRNA is attenuated.
2. The method of Claim 1 wherein the subject has a connective tissue growth factor-associated ocular disorder.
3. The method of Claim 1 wherein the subject is at risk of developing a connective tissue growth factor-associated ocular disorder.
4. The method of Claim 2 wherein the connective tissue growth factor-associated ocular disorder is glaucoma, macular degeneration, diabetic retinopathy, choroidal neovascularization, proliferative vitreoretinopathy or wound healing.
5. The method of Claim 1 wherein the antisense sequence has a region of at least near-perfect contiguous complementarity of at least 21 to 23 nucleotides with the hybridizing portion of mRNA corresponding to SEQ ID NO:1 and comprises an additional TT sequence at the 3' end of each of the sense and the antisense sequence.
6. The method of Claim 1 wherein the sense nucleotide sequence and the antisense nucleotide sequence are connected by a loop nucleotide sequence.
7. The method of Claim 1 wherein the composition is administered via a topical, intravitreal, or transcleral route.

8. The method of Claim 1 wherein the antisense sequence comprises:
3'-TTcccguuuuucacguaggca-5' SEQ ID NO:33.
9. The method of Claim 1 wherein the antisense sequence comprises:
3'-TTcccgagaagacacugaag-5' SEQ ID NO:31.
10. The method of Claim 1 wherein the antisense sequence comprises:
3'-TTccaucauaguagucuauc-5' SEQ ID NO:28.
11. The method of Claim 1 wherein the interfering RNA comprises:
5'-gggccucucugugacuucTT-3' SEQ ID NO:30 and
3'-TTcccgagaagacacugaag-5' SEQ ID NO:31.
12. The method of Claim 1 wherein the interfering RNA comprises:
5'-gggcaaaaagugcauccguTT-3' SEQ ID NO:32 and
3'-TTcccguuuuucacguaggca-5' SEQ ID NO:33.
13. The method of Claim 1 wherein the interfering RNA comprises:
5'-gguuagaucaucagauagTT-3' SEQ ID NO:27 and
3'-TTccaucauaguagucuauc-5' SEQ ID NO:28.
14. The method of Claim 1 further comprising administering to the eye of the subject a second interfering RNA having a length of 19 to 49 nucleotides, and comprising
a sense nucleotide sequence, an antisense nucleotide sequence, and a region of at least near-perfect complementarity of at least 19 nucleotides;
wherein the antisense sequence of the second interfering RNA hybridizes under physiological conditions to a second portion of mRNA corresponding to SEQ ID NO:1, and the antisense sequence has a region of at least near-perfect contiguous complementarity of at least 19 nucleotides with the second hybridizing portion of mRNA corresponding to SEQ ID NO:1.
15. A method of attenuating expression of connective tissue growth factor mRNA in an eye of a subject, comprising:
administering to the eye of the subject a composition comprising an effective amount of single-stranded interfering RNA having a length of 19 to 49 nucleotides and a pharmaceutically acceptable carrier,

wherein the single stranded interfering RNA hybridizes under physiological conditions to a portion of mRNA corresponding to SEQ ID NO:1 beginning at nucleotide 379, 691, 801, 901, 932, 937, 969, 986, 1119, 1170, 1201, 1346, 1473, 1478, 1481, 1488, 1626, 1660, or 1666, and the interfering RNA has a region of at least near-perfect complementarity with the hybridizing portion of mRNA corresponding to SEQ ID NO:1,

wherein the expression of connective tissue growth factor mRNA is attenuated.

16. A method of treating a connective tissue growth factor-associated ocular disorder in a subject in need thereof, comprising:

administering to the eye of the subject a composition comprising an effective amount of interfering RNA having a length of 19 to 49 nucleotides and a pharmaceutically acceptable carrier, the interfering RNA comprising:

a sense nucleotide sequence, an antisense nucleotide sequence, and a region of at least near-perfect contiguous complementarity of at least 19 nucleotides;

wherein the antisense sequence hybridizes under physiological conditions to a portion of mRNA corresponding to SEQ ID NO:1, and has a region of at least near-perfect contiguous complementarity of at least 19 nucleotides with the hybridizing portion of mRNA corresponding to SEQ ID NO:1, wherein the antisense sequence is designed to target a nucleotide sequence of mRNA corresponding to SEQ ID NO:1 comprising nucleotide 379, 691, 801, 901, 932, 937, 969, 986, 1119, 1170, 1201, 1346, 1473, 1478, 1481, 1488, 1626, 1660, or 1666, and

wherein the connective tissue growth factor-associated ocular disorder is treated thereby.

17. The method of Claim 16 wherein the connective tissue growth factor-associated ocular disorder is glaucoma, macular degeneration, diabetic retinopathy, choroidal neovascularization, proliferative vitreoretinopathy or aberrant wound healing.

18. The method of Claim 16 wherein the antisense sequence has a region of at least near-perfect contiguous complementarity of at least 21 to 23 nucleotides with the hybridizing portion of mRNA corresponding to SEQ ID NO:1 and comprises an additional TT sequence at the 3' end of each of the sense and the antisense sequence.

19. The method of Claim 16 wherein the sense nucleotide sequence and the antisense nucleotide sequence are connected by a loop nucleotide sequence.

20. The method of Claim 16 wherein the composition is administered via a topical, intravitreal, or transcleral route.
21. The method of Claim 16 wherein the antisense sequence is designed to target a nucleotide sequence of mRNA corresponding to SEQ ID NO:1 beginning at nucleotide 379, 691, 801, 901, 932, 937, 969, 986, 1119, 1170, 1201, 1346, 1473, 1478, 1481, 1488, 1626, 1660, or 1666.
22. The method of Claim 16 wherein the antisense sequence is designed to target a nucleotide sequence of mRNA corresponding to SEQ ID NO:1 beginning at nucleotide 379, 901, or 1488.
23. The method of Claim 16 wherein the antisense sequence is designed to target a nucleotide sequence of mRNA corresponding to SEQ ID NO:1 comprising nucleotide 379, 901, or 1488.
24. The method of Claim 16 further comprising administering to the eye of the subject a second interfering RNA having a length of 19 to 49 nucleotides, and comprising
a sense nucleotide sequence, an antisense nucleotide sequence, and a region of at least near-perfect complementarity of at least 19 nucleotides;
wherein the antisense sequence of the second interfering RNA hybridizes under physiological conditions to a second portion of mRNA corresponding to SEQ ID NO:1, and the antisense sequence has a region of at least near-perfect contiguous complementarity of at least 19 nucleotides with the second hybridizing portion of mRNA corresponding to SEQ ID NO:1.
25. A method of treating glaucoma, macular degeneration, diabetic retinopathy, choroidal neovascularization, proliferative vitreoretinopathy or aberrant wound healing in a subject in need thereof, comprising:
administering to the eye of the subject a composition comprising an effective amount of single-stranded interfering RNA having a length of 19 to 49 nucleotides and a pharmaceutically acceptable carrier,
wherein the single stranded interfering RNA hybridizes under physiological conditions to a portion of mRNA corresponding to SEQ ID NO:1 beginning at nucleotide 379, 691, 801, 901, 932, 937, 969, 986, 1119, 1170, 1201, 1346, 1473, 1478, 1481, 1488, 1626, 1660, or 1666, and the interfering RNA has a region of at least near-perfect complementarity with the hybridizing portion of mRNA corresponding to SEQ ID NO:1,
wherein the glaucoma, macular degeneration, diabetic retinopathy, choroidal neovascularization, proliferative vitreoretinopathy or aberrant wound healing is treated thereby.

26. A method according to any one of claims 1, 15, 16, or 25, substantially as hereinbefore described with reference to any of the Examples and/or Figures.

FIG. 1A





FIG. 1B



siRNA uptake in GTM3 cells

FIG. 2A




FIG. 2B




FIG. 2C





FIG. 3



34576.40.ST25.txt
SEQUENCE LISTING

<110> Shepard, Allan R.
Pang, Iok-Hou

<120> RNAi INHIBITION OF CTGF FOR TREATMENT OF OCULAR DISORDERS

<130> 34576.40

<150> 60/638,705

<151> 2004-12-23

<160> 36

<170> PatentIn version 3.3

<210> 1

<211> 2312

<212> DNA

<213> Homo sapiens

<400> 1

tccagtgacg gagccgcccg gccgacagcc ccgagacgac agcccggcgc gtcccgggtcc	60
ccacctccga ccaccgccag cgctccaggc cccgcgctcc ccgctcgccg ccaccgcgcc	120
ctccgctccg cccgcagtgc caaccatgac cgccgccagt atgggccccg tccgcgtcgc	180
cttcgtggtc ctctcgccc tctgcagccg gccggccgtc ggccagaact gcagcgggcc	240
gtgccgggtgc ccggacgagc cggcgcccg9 ctgcccggcg ggcgtgagcc tcgtgctgga	300
cggctgcggc tgctgcccgc tctgcgcaa gcagctgggc gagctgtgca ccgagcgcga	360
cccctgcgac ccgcacaagg gcctcttctg tgacttcggc tccccggcca accgcaagat	420
cggcgtgtgc accgcaaag atggtgctcc ctgcatctc ggtggtacgg tgtaccgcag	480
cggagagtcc ttccagagca gctgcaagta ccagtgcacg tgcctggacg gggcgggtggg	540
ctgcatgccc ctgtgcagca tggacgttcg tctgcccagc cctgactgcc cttccccgag	600
gagggtaag ctgcccggga aatgctgcga ggagtgggtg tgtgacgagc ccaaggacca	660
aaccgtggtt gggcctgccc tcgcggtta ccgactggaa gacacgtttg gcccagacct	720
aactatgatt agagccaact gcctggtcca gaccacagag tggagcgcct gttccaagac	780
ctgtgggatg ggcattcca cccgggttac caatgacaac gcctcctgca ggctagagaa	840
gcagagccgc ctgtgcatgg tcaggccttg cgaagctgac ctggaagaga acattaagaa	900
gggcaaaaag tgcatccgta ctccaaaat ctccaagcct atcaagtttg agctttctgg	960
ctgcaccagc atgaagacat accgagctaa attctgtgga gtatgtaccg acggccgatg	1020
ctgcaccccc cacagaacca ccaccctgcc ggtggagttc aagtgccctg acggcgaggt	1080
catgaagaag aacatgatgt tcatcaagac ctgtgcctgc cattacaact gtcccggaga	1140
caatgacatc tttgaatcgc tgtactacag gaagatgtac ggagacatgg catgaagcca	1200
gagagtgaga gacattaact cattagactg gaacttgaac tgattcacat ctcatTTTTc	1260
cgtaaaaatg atttcagtag cacaagttat ttaaactctgt ttttctaact gggggaaaag	1320
attccccacc aattcaaac attgtgccat gtcaaacaaa tagtctatct tccccagaca	1380

34576.40.ST25.txt

```

ctggtttgaa gaatgttaag acttgacagt ggaactacat tagtacacag caccagaatg 1440
tatattaagg tgtggcttta ggagcagtgg gagggtacca gcagaaaggt tagtatcatc 1500
agatagctct tatacgagta atatgcctgc tatttgaagt gtaattgaga aggaaaattt 1560
tagcgtgctc actgacctgc ctgtagcccc agtgacagct aggatgtgca ttctccagcc 1620
atcaagagac tgagtcaagt tgttccttaa gtcagaacag cagactcagc tctgacattc 1680
tgattcgaat gacactgttc aggaatcgga atcctgtcga ttagactgga cagcttgtgg 1740
caagtgaatt tcctgtaaca agccagattt tttaaaattt atattgtaa tattgtgtgt 1800
gtgtgtgtgt gtgtatatat atatatatat gtacagttat ctaagttaat ttaaagttgt 1860
ttgtgccttt ttatttttgt ttttaatgct ttgatatttc aatgttagcc tcaatttctg 1920
aacaccatag gtagaatgta aagcttgtct gatcgttcaa agcatgaaat ggatacttat 1980
atggaaattc tctcagatag aatgacagtc cgtaaaaaca gattgtttgc aaaggggagg 2040
catcagtgtc cttggcaggc tgatttctag gtaggaaatg tggtagctca cgctcacttt 2100
taatgaaca atggccttta ttaaaaactg agtgactcta tatagctgat cagttttttc 2160
acctggaagc atttgtttct actttgatat gactgttttt cggacagttt atttgttgag 2220
agtgtgacca aaagttacat gtttgcacct ttctagttga aaataaagta tattttttct 2280
aaaaaaaaa aaaaacgaca gcaacggaat tc 2312
    
```

```

<210> 2
<211> 19
<212> DNA
<213> Artificial Sequence
    
```

```

<220>
<223> TARGETING SEQUENCE
    
```

```

<400> 2
gggcctcttc tgtgacttc 19
    
```

```

<210> 3
<211> 19
<212> DNA
<213> Artificial Sequence
    
```

```

<220>
<223> Targeting Sequence
    
```

```

<400> 3
ccgactggaa gacacgttt 19
    
```

```

<210> 4
<211> 19
<212> DNA
<213> Artificial Sequence
    
```

```

<220>
<223> Targeting Sequence
    
```

```

<400> 4
cccgggttac caatgacaa 19
    
```

34576.40.ST25.txt

<210> 5
 <211> 25
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Targeting Sequence

<400> 5
 tccaagccta tcaagtttga gcttt 25

<210> 6
 <211> 25
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Targeting Sequence

<400> 6
 tccaagccta tcaagtttga gcttt 25

<210> 7
 <211> 19
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Targeting Sequence

<400> 7
 gcctatcaag tttgagctt 19

<210> 8
 <211> 25
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Targeting Sequence

<400> 8
 gcatgaagac ataccgagct aaatt 25

<210> 9
 <211> 19
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Targeting Sequence

<400> 9
 gctaaattct gtggagtat 19

<210> 10
 <211> 25
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Targeting Sequence

34576.40.ST25.txt

<400> 10
gccattacaa ctgtcccgga gacaa 25

<210> 11
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Targeting Sequence

<400> 11
ggaagatgta cggagacat 19

<210> 12
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Targeting Sequence

<400> 12
gagagtgaga gacattaact catta 25

<210> 13
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Targeting Sequence

<400> 13
gccatgtcaa acaaatagtc tatct 25

<210> 14
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Targeting Sequence

<400> 14
gggtaccagc agaaagggtt 19

<210> 15
<211> 19
<212> DNA
<213> Artificial

<220>
<223> Targeting Sequence

<400> 15
ccagcagaaa ggtagtat 19

<210> 16
<211> 19
<212> DNA

34576.40.ST25.txt

<213> Artificial Sequence

<220>

<223> Targeting Sequence

<400> 16

gcagaaaggt tagtatcat

19

<210> 17

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Targeting Sequence

<400> 17

gcagaaaggt tagtatcatc agata

25

<210> 18

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Targeting Sequence

<400> 18

ggtagtatc atcagatag

19

<210> 19

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Targeting Sequence

<400> 19

ggtagtatc atcagatagc tctta

25

<210> 20

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Targeting Sequence

<400> 20

gagactgagt caagtggtc cttaa

25

<210> 21

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Targeting Sequence

<400> 21

gcagactcag ctctgacat

19

34576.40.ST25.txt

<210> 22
 <211> 25
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Targeting Sequence

<400> 22
 tcagctctga cattctgatt cgaat 25

<210> 23
 <211> 25
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Targeting Sequence

<400> 23
 tcctgtcgat tagactggac agctt 25

<210> 24
 <211> 19
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Targeting Sequence

<400> 24
 gcttgtggca agtgaattt 19

<210> 25
 <211> 21
 <212> RNA
 <213> Artificial Sequence

<220>
 <223> Sense Strand

<400> 25
 gguuaguauc aucagauagu u 21

<210> 26
 <211> 21
 <212> RNA
 <213> Artificial Sequence

<220>
 <223> Antisense Strand

<400> 26
 cuaucugaug auacuaaccu u 21

<210> 27
 <211> 21
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Sense Strand with 3'TT

34576.40.ST25.txt

<220>
 <221> misc_RNA
 <222> (1)..(19)
 <223> ribonucleotides

<220>
 <221> misc_feature
 <222> (20)..(21)
 <223> deoxyribonucleotides

<400> 27
 gguuaguaucaucagauagt t 21

<210> 28
 <211> 21
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Antisense Strand with 3'TT

<220>
 <221> misc_RNA
 <222> (1)..(19)
 <223> ribonucleotides

<220>
 <221> misc_feature
 <222> (20)..(21)
 <223> deoxyribonucleotides

<400> 28
 cuaucugaug auacuaacct t 21

<210> 29
 <211> 51
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Hairpin Duplex with Loop

<220>
 <221> misc_RNA
 <222> (1)..(21)
 <223> ribonucleotides

<220>
 <221> misc_feature
 <222> (22)..(30)
 <223> any, A, T/U, C, G

<220>
 <221> misc_feature
 <222> (31)..(51)
 <223> ribonucleotides

<400> 29
 gguuaguaucaucagauagu unnnnnnnnn cuaucugaug auacuaaccu u 51

<210> 30

34576.40.ST25.txt

<211> 21
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Sense Strand with 3'TT

<220>
 <221> misc_RNA
 <222> (1)..(19)
 <223> ribonucleotides

<220>
 <221> misc_feature
 <222> (20)..(21)
 <223> deoxyribonucleotides

<400> 30
 gggccucuuc ugugacuuct t 21

<210> 31
 <211> 21
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Antisense Strand with 3'TT

<220>
 <221> misc_RNA
 <222> (1)..(19)
 <223> ribonucleotides

<220>
 <221> misc_feature
 <222> (20)..(21)
 <223> deoxyribonucleotides

<400> 31
 gaagucacag aagaggcct t 21

<210> 32
 <211> 21
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Sense Strand with 3'TT

<220>
 <221> misc_RNA
 <222> (1)..(19)
 <223> ribonucleotides

<220>
 <221> misc_feature
 <222> (20)..(21)
 <223> deoxyribonucleotides

<400> 32
 gggcaaaaag ugcauccgut t 21

34576.40.ST25.txt

<210> 33
 <211> 21
 <212> DNA
 <213> Artificial

<220>
 <223> Antisense Strand with 3'TT

<220>
 <221> misc_RNA
 <222> (1)..(19)
 <223> ribonucleotides

<220>
 <221> misc_feature
 <222> (20)..(21)
 <223> deoxyribonucleotides

<400> 33
 acggaugcac uuuuugcct t 21

<210> 34
 <211> 23
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Probe/Primer Sequence

<400> 34
 cagctctgac attctgattc gaa 23

<210> 35
 <211> 20
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Probe/Primer Sequence

<400> 35
 tgccacaagc tgtccagtct 20

<210> 36
 <211> 30
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Probe/Primer Sequence

<400> 36
 aatcgacagg attccgattc ctgaacagtg 30