



(86) Date de dépôt PCT/PCT Filing Date: 2000/01/10  
(87) Date publication PCT/PCT Publication Date: 2000/07/20  
(85) Entrée phase nationale/National Entry: 2001/07/09  
(86) N° demande PCT/PCT Application No.: US 2000/000580  
(87) N° publication PCT/PCT Publication No.: 2000/041524  
(30) Priorité/Priority: 1999/01/11 (60/115,498) US

(51) Cl.Int.<sup>7</sup>/Int.Cl.<sup>7</sup> C12Q 1/68  
(71) Demandeur/Applicant:  
PRESIDENT AND FELLOWS OF HARVARD COLLEGE,  
US  
(72) Inventeurs/Inventors:  
RICHARDSON, CHARLES, US;  
TABOR, STANLEY, US  
(74) Agent: SMART & BIGGAR

(54) Titre : AMPLIFICATION ISOTHERME D'ADN  
(54) Title: ISOTHERMAL AMPLIFICATION OF DNA

10	20	30	40	50	60	70	80
ATGGACAAT	CGCAGGATTC	CGATAGTGT	TTTCTTTACC	ACATTCTCTG	TGCAACTGT	GGGAGTAGTG	ATGGGAAGTC 80
GCTGTCTCT	GACGGACACA	CGTTCTGCTA	CGTATGCGAG	AAGTGGACTG	CTGGTAATGA	AGACACTAAA	GAGAGGGCTT 160
CAAAACGGAA	ACCCTCGG	GGAAAGCCG	GGACTTACAA	CGTGTGGAAC	TTCCGGGAAT	CCAATGGACG	CTACTCCGCG 240
TTAACTGCGA	GAGGAATCTC	CAAGGAAACC	TGTCASAAGG	CTGGCTACTG	GATTGCCAAA	GTPAGCGGTG	TGATGTACCA 320
AGTGGCTGAC	TATCGGGACC	AGAACGGCAA	CATTGTGAGT	CAGAAGGTT	GAGATAAGA	TAAGAAGTTT	AAGACCACTG 400
GTAGTCACAA	GAGTACGCT	CTGTTCCGGA	AGCACTTGTG	GAATGGTGGT	ARGAAGATTG	TCGTTACAGA	AGGTGAAATC 480
GACATGCTTA	CCGTGATGGA	ACTTCAAGAC	TGTAAGTATC	CTGTAGTGTG	GTTGGGTGAC	GGTCCCTCTG	CCGCTAAGAA 560
GACATGCGCT	GCCAACTAGC	AATACTTTGA	CCAGTTCGAA	CAGATTATCT	TAATGTTGGA	TATGGACGAA	GCAGGGCGCA 640
AAGCAATCGA	AGAGGCTGCA	CAGGTCTTAC	CTGGCTGTAA	GGTACGAGTG	CCAGTCTTCT	CGTSTAAGGA	TGCAAAACGAG 720
TGTCACTTAA	ATGGTCAACG	CCGTGAAATC	ATGGAGCAAG	TGTGGAATGC	TGGTCCCTGG	ATCCCTGATG	GTCTGGTATC 800
GGCTCTTTG	TTACGTGAAC	GAATCCGTGA	GCACCTATCG	TCCGAGGAAT	CAGTAGGTTT	ACTTTTCAGT	GGCTGCACCTG 880
GTATCAACGA	TAAGACCTTA	GGTCCCGTGG	GTGGTGAAGT	CATTATGCTC	ACTTCCGGTT	CCGGTATGGG	TAAGTCAACG 960
TTCGTCCGTC	AACAAGCTCT	ACAATGGGGC	ACAACGATGG	GCAAGAAGGT	AGGCTTAGCG	ATGCTTGAGG	AGTCCGTGTA 1040
GGAGACCGCT	GAGGACCTTA	TAGGTCTACA	CAACCGTCTC	CGACTGAGAC	AATCCGACTC	ACTAAACAGA	GAGATTATTG 1120
AGAACGGTAA	GTTCCGACCA	TGGTTCGATG	AATGTTCCGG	CAACGATACG	TTCCATCTAT	ATGACTCATT	CGCCGAGGCT 1200
GAGACGGATA	CACTGCTCGC	TAAGCTGGCC	TACATGCGCT	CAGGCTTGGG	CTGTGACGTA	ATCATCTAG	ACCACATCTC 1280
AATCGTGGTA	TCCGCTTCTG	GTGAATCCGA	TGAGCGTAAG	ATGATTGACA	ACCTGATGAC	CAAGCTCAAA	GGTTCGCTA 1360
AGTCAACTGG	GGTGGTGGCT	GTGTAATTT	GTCACTTAA	GAACCCAGAC	AAAGTAAAG	CACATGAGGA	AGGTCCGCCC 1440
GTTCCTATTA	CTGACCTACG	TGGTCTGGC	GCACCTACCC	AACATCTGA	TACTATTATT	GCCTTGAGC	GTAATCAGCA 1520
AGGCGATATG	CCTAACCTTG	TCCCTGTTCC	TATTCTCAAG	TGCCGCTTTA	CTGGTGATAC	TGGTATCGCT	GGCTACATGG 1600
AATACAACAA	GGAAACCGGA	TGGCTTGAAC	CATCAAGTTA	CTCAGGGGAA	GAAGACTCAC	ACTCAGATC	AACAGACTGG 1680
TCCAACGACA	CTGACTTCTG	ACAGGATCTT	TGATGACTTT	CCAGACGACT	ACGAGAAGTT	TCCGCTGGAGA	GTCCCATCTT 1760
AATACGACTC	ACTAAAGGAG	ACACACCATG	TTCAAAGTGA	TTAAGAAGTT	AGGCCAACTG	CTGGTTCGTA	TGTACAACGT 1840
CGAAGCCAA	CGACTGAACG	ATGAGGCTCG	TAAAGAGGCG	ACACAGTCA	CGCTCTGGC	GATTCGCTCC	AAAAGTGGTT 1920
TTGCGCTTAC	CCCAACCAAC	AGGGGATTTG	CTGCTTTCCA	TTGAGCTGT	TTCTCTGGC	GACGTTCCGG	GGCGCGTGT 2000
TGTGCATCCA	TCTGGATTCT	CCTGTCAATT	AGCTTTGGTG	GTGTGTGGCA	GTTGTACTCC	TGAACGAAAA	CCCCCGCGA 2080
TTGGCACATT	GGCAGCTAAT	CCGGAATCGC	ACTTACCGCC	AATGCTTCT	TTCTTATCAC	ACACCCCAAA	GCCTTCTGCT 2160
TTGAATGCTG	CCCTTCTTCA	GGGCTTAATT	TTTAAGAGCG	TCACCTTCAT	GGTGGTCACT	GGCTCTGCT	GATGTGCTCA 2240
GTATCACCGC	CAGTGGTATT	TATGTCAACA	CCGCCAGAGA	TAATTTATCA	CCGCAGATGG	TTATCTGTAT	GTTTTTTATA 2320
TGAATTTATT	TTTTGCAGG	GGGCATTGTT	TGGTAGGTGA	GAGATCCGGC	TGTAACAAA	GGCCGAAAGG	AAGCTGAGTT 2400
GGCTGCTGCC	ACCGCTGAGC	AATAACTAGC	ATAACCCCTT	GGGCCTCTA	AACGGTCTT	GAGGGTTTTT	TTGCTGAAAG 2480
GAGGAACAT	ATCCGGATAT	CCCGCAAGAG	GGCCGGCAGT	ACCGGCATAA	CCAAGCCTAT	GCCTACAGCA	TCCAGGGTGA 2560
CGGTCCCGAG	GATGACGATG	AGCGCATGCT	TAGATTTCAT	ACACGGTGGC	TGACTGCGTT	AGCAATTTAA	CTGTGATAAA 2640
CTACCGCAT	AAAGCTTGGC	SCCGCACTCG	ACGAACCCCT	CGGATCTCGA	TCCCGCGAAA	TTAATACGAC	TCACTATAGG 2720
GAGACCACAA	CGTTTCCCT	CTAGAAATAA	TTTTGTTTAA	CTTTAAGAA	GAGATATACA	TATGCGTGA	CGAATCCGTC 2800
AGCACCTATC	GTCCGAGGAA	TCAGTAGGTT	TACTTTTCAG	TGGCTGCACT	GGTATCAACG	ATAAGACCTT	AGGTGCCCGT 2880
GGTGGTGAAG	TCATTATGCT	CACTTCCGGT	TCCGGTATGG	GTAAAGTCAAC	GTTGCTCCGT	CAACAAGCTC	TACAATGGGG 2960
CACAGCGATG	GGCAAGAAGG	TAGGCTTAGC	GATGCTTGAG	GAGTCCGTTG	AGGAGACCGC	TGAGGACCTT	ATAGGTCTAC 3040
ACAACCGTGT	CCGACTGAGA	CAATCCGACT	CACTAAAGAG	AGAGATTATT	GAGAACGGTA	AGTTCCGACCA	ATGGTTCGAT 3120
GAAGTGTTCG	GCARCGATAC	GTTCCATCTA	TATGACTCAT	TCCCGAGGC	TGAGACGGAT	AGACTGCTCG	CTAAGCTGGC 3200
CTACATGCGC	TCAGGCTTGG	GCTGTGACGT	AATCAATCTA	GACCCACATC	CAATCGTCTG	ATCCGCTTCT	GGTGAATCCG 3280
ATGAGCGTAA	GATGATTGAC	AACCTGATGA	CCAAGCTCAA	AGGGTTCGCT	AAGTCAACTG	GGGTGGTGT	GGTCTGTAAT 3360
TGTCACCTTA	AGAACCAGCA	CAAAAGGTA	GCACATGAGG	AAGGTCCGCC	CGTTCTATT	ACTGACCTAC	GTGGTCTTGG 3440
CGCACTACGC	CAACTATCTG	ATACTATTAT	TGCCCTTGAG	CGTAATCAGC	AAGCCGATAT	GCCTAACCTT	GTCCCTGCTC 3520

(57) Abrégé/Abstract:

A method of amplifying a template DNA molecule comprising incubating the template DNA molecule in a reaction mixture comprising a DNA polymerase and at least one accessory protein at a constant temperature to produce amplified product, wherein production of amplified product does not require exogenously-added oligonucleotide primers and the template DNA molecule does not have terminal protein covalently bound to either 5' end.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
20 July 2000 (20.07.2000)

PCT

(10) International Publication Number  
WO 00/41524 A3

- (51) International Patent Classification<sup>7</sup>: C12Q 1/68 [US/US]; 9a Lowell Street, Cambridge, MA 02138 (US). RICHARDSON, Charles [US/US]; 78 Chestnut Hill Road, Chestnut Hill Road, MA 02167 (US).
- (21) International Application Number: PCT/US00/00580
- (22) International Filing Date: 10 January 2000 (10.01.2000) (74) Agent: LYON & LYON LLP; 633 West Fifth Street, Suite 4700, Los Angeles, CA 90071-2066 (US).
- (25) Filing Language: English
- (26) Publication Language: English (81) Designated States (*national*): AU, CA, JP, US.
- (30) Priority Data: 60/115,498 11 January 1999 (11.01.1999) US (84) Designated States (*regional*): European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).
- (71) Applicant (*for all designated States except US*): PRESIDENT AND FELLOWS OF HARVARD COLLEGE [US/US]; 17 Quincy Street, Cambridge, MA 02138 (US).  
Published:  
— With international search report.  
— Before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments.
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): TABOR, Stanley

[Continued on next page]

(54) Title: ISOTHERMAL AMPLIFICATION OF DNA

10	20	30	40	50	60	70	80
ATGGACAATT	CGCAGGATTC	CGATAGTGT	TTTCTTTACC	ACATTCCTTG	TGACAACGTG	GGGAGTAGTG	ATGGGAACCTC 80
GCTGTTCTCT	GACGGACACA	CGTTCGTGTA	CGTATGCGAG	AAGTGGACTG	CTGGTAATGA	AGACACTAAA	GAGAGGGCTT 160
CAAAACGGAA	ACCCTCcgge	GGaAagCCcg	gGACTTACAA	CGTGTGGAAc	TTCGGGGAAT	CCAATGGAGc	CTACTCCGCG 240
TTAACTGGGA	GAGGAATCTC	CAAGGAACC	TGTGAGAGG	CTGGCTACTG	GATTGCCAAA	GTAGACGGTG	TGATGTACCA 320
AGTGGCTGAC	TATCGGGACC	AGAACGGCAA	CATTGTGAGT	CAGAAGGTTc	GAGATAAAGA	TAAGAACCTT	AAGACCACCTG 400
GTAGTCACAA	GAGTGACGCT	CTGTTCCGGA	AGCACTTGTG	GAATGGTGGT	AAGAAGATTG	TCGTTACAGA	AGGTGAAATC 480
GACATGCTTA	CCGTGATGGA	ACTTCAAGAC	TGTAAGTATC	CTGTAGTGTc	GTTGGGTCAc	GGTGCCTCTG	CCGCTAAGAA 560
GACATGGGCT	GCCAACTACG	AATACTTTGA	CCAGTTCGAA	CAGATTATCT	TAATGTTCTG	TATGGACGAA	GCAGGGCCGCA 640
AAGCAGTCGA	AGAGSCTGCA	CAGSTTCTAC	CTGCTGGTAA	GGTACGAGTG	GCAGTTCCTC	CGTGTAAAGG	TGCAAAACGAG 720
TGTCACCTAA	ATGGTCACGA	CCGTGAAATC	ATGGAGCAAG	TCTGGAAATG	TGGTCCCTGG	ATTCCTGATG	GTGTGGTATC 800
GGCTCTTTGc	TTACGTGAAC	GAATCCGTGA	GCACCTATCG	TCCGAGGAAT	CAGTAGGTTT	ACTTTTCAGT	GGCTGCACCTG 880
GTATCAACGA	TAAACCTTA	GGTCCCGGTG	GTGGTGAAGT	CATTATGGTC	ACTTCCGGTT	CCGGTATGGG	TAAGTCAACG 960
TTCTGCTGTC	AACAAGCTCT	ACAAATGGGG	ACAGCGATGG	GCAAGAAAGT	AGGCTTAGCG	ATGCTTGAGG	AGTCCGTTCA 1040
GGAGACGGCT	GAGGACCTTA	TAGGTCTACA	CAACCGTGC	CGACTGAGAC	AATCCGACTC	ACTAAACAGA	GAGATTATTG 1120
AGAACGGTAA	GTTCGACCAA	TGGTTCGATG	AACCTGTCGG	CAACGATACG	TCCATCTAT	ATGACTCAT	CGCCGAGGCT 1200
GAGACGGATA	CACTGCTGCG	TAACTGGCC	TACATGCGCT	CAGGCTTGGG	CTGTGACGTA	ATCATTCTAG	ACCACATCTC 1280
AATCGTCTGA	TCCGTTGTCG	GTGAATCCGA	TGAGCGTAAG	ATGATTGACA	ACCTGATGAC	CAAGCTCAA	GGGTTCCGTA 1360
AGTCAACTGG	GSTGGTGTG	GTCTGAATTT	GTCACTTAA	GAACCCAGAC	AAAGGTAAAG	CACATGAGGA	AGGTCCGCCC 1440
GTTCTTATTA	CTGACCTACG	TGGTTCGGC	GCACCTACCC	AACTATCTGA	TACTATTATT	GCCCTTGAGC	GTAATCAGCA 1520
AGCGGATATG	CCTAACCTTG	TCTCTGTTCC	TATCTCAAG	TGCGCTTTA	CTGGTATAC	TGGTATCGCT	GGCTACATGG 1600
AATACAACA	GGAAACCGGA	TGGCTTGAAC	CATCAAGTTA	CTCAGGGGAA	GAAGACTCAC	ACTCAGAGTC	AACAGACTGG 1680
TCCACCGACA	CTGACTTCTG	ACAGGATTCT	TGATGACTTT	CCAGACGACT	ACGAGAAGTT	TCCGTGGAGA	GTCCCATCTC 1760
AATACGACTC	ACTAAAGGAG	ACACACCATG	TTCAAATGA	TTAAGAAGTT	AGGCCAAGTG	CTGGTTCGTA	TGTACAACGT 1840
GGAAAGCCAG	CGACTGAACG	ATGAGGCTCG	TAAAGAGGCC	ACACAGTCAC	GCGCTCTGGC	GATTCTGCTC	AAACTGGGTT 1920
TTGCGCTTAC	CCCAACCAAC	AGGGCAATTC	CTGCTTCCCA	TTGAGCCTGT	TTCTCTGCCC	GACGTTCCGG	GGGGCGTGT 2000
TGTGCATCCA	TCTGGATTCT	CCTCTCAGTT	AGCTTTGGTG	GTGTGTGCCA	GTTGTAGTCC	TGAACGAAA	CCCCCGCGA 2080
TTGGCACATT	GGCAGCTAAT	CCGGAATCCG	ACTTACGGCC	AAATGCTTCT	TTCGTATCAC	ACACCCCAA	GCCTTCTGCT 2160
TTGAATGCTG	CCCTTCTTCA	GGGCTTAAAT	TTAAGAGCG	TCACCTTCAT	GGTGGTCAGT	GCGTCTGCT	GATGTGCTCA 2240
GTATCAGCGC	CAGTGGTATT	TATGTCAACA	CCGCCAGAGA	TAATTTATCA	CCGCAGATGG	TTATCTGTAT	GTTTTTTATA 2320
TGAATTTATT	TTTTGCAGGG	GGGCATTGTT	TGTTAGGTGA	GAGATCCGGC	TGCTAACAAA	GCCCGAAAG	AAGCTGAGTT 2400
GGCTGCTGCC	ACCCGTGAGC	AATAACTAGC	ATAACCCCTT	GGGGCTCTA	AACGGGCTTT	GAGGGGTTTT	TTGCTGAAAG 2480
GAGGAATAT	ATCCGGATAT	CCCGCAAGAG	GCCCCGAGT	ACCGGCATA	CCAAGCCTAT	GCCTACAGCA	TCCAGGGTGA 2560
CGGTCCCGAG	GATGACGATG	AGCGCATTGT	TAGATTTCA	ACACGGTSCC	TGACTCGGTT	AGCAATTTAA	CTGTGATAAA 2640
CTACCCGATT	AAAGCTTGGC	GCCGCACTCG	ACGAAACCTT	CGGATCTCGA	TCCCAGGAAA	TTAATACGAC	TCACTATAGG 2720
GAGACCACAA	CGGTTCCCTT	CTAGAAATAA	TTTTGTTTAA	CTTTAAGAA	GAGATATACA	TATGCGTGA	CGAATCCGTC 2800
AGCACCTATC	GTCCGAGGAA	TCAGTAGGTT	TACTTTTTCAG	TGGCTGCACT	GATATCAACG	ATAAGACCTT	AGGTGCCCGT 2880
GTTGGTGAAG	TCATTATGGT	CACCTCCGGT	TCCGGTATGG	GTAAGTCAAC	GTTCTGCTGT	CAACAAGCTC	TACAAATGGG 2960
CACAGCGATG	GGCAAGAAGG	TAGGCTTAGC	GATGCTTAGC	GAGTCCGTTG	AGGAGACCGC	TGAGGACCTT	ATAGGTCTAC 3040
ACAACCGTGT	CEGACTGAGA	CAATCCGACT	CACTAAGAG	AGAGATTATT	GAGAACGGTA	AGTTCGACCA	ATGGTTCGAT 3120
GAAGTGTTCG	GCAACGATAC	GTCCATCTA	TATGACTCAT	TCCCGAGGC	TGAGACGGAT	AGACTGCTCG	CTAAGCTGGC 3200
CTACATGCGC	TCAGGCTTGG	GCTGTGACGT	AATCATCTA	GACACATCT	CAATCGTCTG	ATCCGCTTCT	GGTGAATCCG 3280
ATGAGCGTAA	GATGATTGAC	AACCTGATGA	CCAAGCTCA	AGGGTTCGCT	AAGTCAACTG	GGGTGGTGT	GCTCGTAAAT 3360
TGTCACCTTA	AGAACCCAGA	CAAAGGTAAA	GCACATGAGG	AAGGTCCGCC	CGTTCTTATT	ACTGACTTAC	GTGGTCTGTC 3440
CGCACTACGC	CAACTATCTG	ATACTATTAT	TGCCCTTGAG	CGTAATCAGC	AAGGCGATAT	GCCTAACCTT	GTCCCTGTTT 3520

[Continued on next page]

WO 00/41524 A3



---

**(88) Date of publication of the international search report:**  
8 February 2001

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

---

**(57) Abstract:** A method of amplifying a template DNA molecule comprising incubating the template DNA molecule in a reaction mixture comprising a DNA polymerase and at least one accessory protein at a constant temperature to produce amplified product, wherein production of amplified product does not require exogenously-added oligonucleotide primers and the template DNA molecule does not have terminal protein covalently bound to either 5' end.

## ISOTHERMAL AMPLIFICATION OF DNA

### FIELD OF THE INVENTION

The present invention relates to the field of amplification of DNA and RNA.

### BACKGROUND OF THE INVENTION

5 This invention was made with government support including a grant from the U.S. Dept. of Energy, contract number DE-FG02-96ER62251. The U.S. government may have certain rights in the invention.

The following is a discussion of the relevant art, none of which is admitted to be prior art to the appended claims.

10 A variety of methods have been developed to efficiently amplify DNA using purified proteins. These generally can be classified as ones that use thermocycling of the reaction mixtures and ones in which the reactions are isothermal. Methods that use thermocycling of the reaction mixture are based on the polymerase chain reaction, or PCR (Saiki *et al.*, Science 230:1350, 1985). In PCR, two primers are added to the target  
15 DNA, such that the two primers are complementary to opposite strands of the target sequence with their 3' ends oriented towards one another. Multiple cycles of denaturation of target DNA, annealing of the two primers, and then extension of the primers with a DNA polymerase, results in an exponential synthesis of the target DNA sequence located between the two primers. This procedure requires the use of a DNA  
20 polymerase that is thermostable in order to survive the high temperatures required to denature the product DNA each cycle. The most common polymerase used for PCR is from *Thermus aquaticus*, or Taq DNA polymerase (Saiki *et al.*, Science 239:487, 1988). Modification of the reaction conditions and the enzyme mixture used can allow the amplification of DNA up to 50 kbp in length (Barnes, Proc. Natl. Acad. Sci. USA  
25 91:2216, 1994 and Cheng *et al.*, Proc. Natl. Acad. Sci. USA 91:5695, 1994).

Isothermal procedures for the amplification of nucleic acid include (1) Q $\beta$  replicase, (2) self-sustained sequence replication (3SR), (3) strand displacement amplification (SDA), (4) terminal protein-primed DNA amplification using  $\phi$ 29 DNA polymerase, and (5) rolling circle amplification (RCA).

5 The RNA-dependent RNA polymerase Q $\beta$  polymerase has been used for gene detection in a strategy that uses two probes that hybridize to adjacent positions on a target sequence that are then ligated to form an amplifiable reporter RNA (Tyagi *et al.*, Proc. Natl. Acad. Sci. USA 93:5395, 1996).

10 Self-sustained sequence replication (3SR) is a method of amplifying RNA that depends on the action of three enzymes; reverse transcriptase, DNA-dependent RNA polymerase and ribonuclease H (Guatelli *et al.*, Proc. Natl. Acad. Sci. USA 87:1874, 1990).

15 Strand displacement amplification (SDA) is a DNA amplification system that uses a restriction enzyme to introduce specific nicks in a target to be amplified and a DNA polymerase that is capable of strand displacement synthesis (Walker *et al.*, Proc. Natl. Acad. Sci. USA 89:392, 1992).

20 Terminal protein-primed DNA amplification exponentially amplifies linear  $\phi$ 29 DNA using the  $\phi$ 29 proteins DNA polymerase, terminal protein, double-stranded DNA binding protein, and single-stranded DNA binding protein (Blanco *et al.*, Proc. Natl. Acad. Sci. USA 91:12198, 1994). To date this method has only been successful using linear  $\phi$ 29 DNA as template.

25 Rolling circle amplification (RCA) can either be linear or exponential amplification of circular oligonucleotide probes that use two primers that anneal to each of the two strands, and a DNA polymerase that strand displaces (Lizardi *et al.*, Nature Genetics 19:225, 1998 and Zhange *et al.*, Gene 211:277, 1998).

In order to be used as a generalized amplification system, all of these methods require the addition of specific oligonucleotide primers that are extended by a

polymerase; the primers serve to fix the limits of the segment of nucleic acid to be amplified.

### SUMMARY OF THE INVENTION

The present invention provides an *in vitro* method to amplify DNA exponentially at a constant temperature and without the use of exogenously added primers by using a DNA polymerase and various accessory proteins. The accessory proteins function with the DNA polymerase to carry out amplification under these conditions.

In one aspect, the method involves the addition of a mixture of enzymes including a DNA polymerase, a helicase and a primase and single-stranded binding protein to template DNA, which can result in amplification of the template DNA of at least 10-fold and up to and greater than 1 million-fold amplification. Such an amplification system is useful for common molecular biology procedures, including but not limited to, the preparation of plasmid DNA from cells for DNA sequencing, cloning, providing substrates for transcription/translation reactions, and mutagenesis. In addition, this generalized system for DNA amplification can serve as a useful way to prepare large amounts of human genomic DNA from small samples for genotype analysis. This could be important both for preserving or "immortalizing" small amounts of rare DNA samples, and for more general use in clinical applications as a way to avoid removing large amounts of blood from patients to obtain DNA for genotyping. Finally, this amplification system provides a very sensitive detection system for DNA contamination in samples, for example in enzymes that have been purified under good manufacturing protocol (GMP) for use in human therapeutics.

One way in which the present invention is distinguished from other exponential procedures for amplification is that it does not require the use of exogenous oligonucleotide primers. Exogenous oligonucleotide primers refer to small nucleic acid molecules that are generally in the range of 6 to 50 bases in length, although longer primers can work. They are normally synthesized by chemical as opposed to enzymatic methods. The primers are complementary to certain regions of the template and are

utilized by the polymerase to initiate DNA synthesis. The region of the template that is amplified is defined by the primers utilized. Primers are required in the present invention, however they are made by one of the components of the system, the primase, *e.g.*, gene 4 protein, rather than being synthesized in advance and then added to the system. The addition of primers to the system of the present invention has little effect on increasing the rate of DNA synthesis (generally less than two-fold), as the system works very well even without exogenous primers. Thus, even if the replication system of the present invention contains exogenous primers, there is still substantial DNA synthesis that is exponential in nature that is independent of the primers. One of ordinary skill in the art would readily be able to distinguish DNA synthesis carried out by the claimed method from exogenously added primer dependent synthesis. In the situation where DNA synthesis is dependent on specific exogenously added primers, a new set of primers will be required for each new DNA template amplified. In the situation where DNA synthesis is dependent on the exogenous addition of random primers (*e. g.* random hexamers) the DNA synthesis will not be exponential. This invention describes a system whereby DNA synthesis is exponential even in the absence of any exogenously added primers.

The template DNA used in this invention also does not need have a terminal protein bound to either 5' end. A terminal protein is a protein like the type used in the bacteriophage  $\phi$  29 replication system, *e.g.*, phage  $\phi$ 29 the gene 3 protein (Salas, Ann. Rev. Biochem. 60:39, 1991). Such a terminal protein may be present but is not functionally required for the amplification system to work.

The term "constant temperature" refers to an amplification reaction that is carried out under isothermal conditions as opposed to thermocycling conditions. Thermocycling conditions are used by polymerase chain reaction methods in order to denature the DNA and anneal new primers after each cycle. Constant temperature procedures rely on other methods to denature the DNA, such as the strand displacement ability of some polymerases or of DNA helicases that act as accessory proteins for some DNA polymerases. Thus, the term "constant temperature" does not mean that no temperature

fluctuation occurs, but rather indicates that the temperature variation during the amplification process is not sufficiently great to provide the predominant mechanism to denature product/template hybrids. Preferably the constant temperature is less than 60°C, more preferably less than 50°, still more preferably less than 45°, and can even be less than 40°C.

The constant temperature amplification is carried out "*in vitro*", meaning that the reaction is not carried out in cells. Generally the amplification is performed using purified or at least substantially purified proteins.

In a preferred embodiment the DNA polymerase is from a bacteriophage. Further preferred is that the DNA polymerase is from bacteriophage T7. An even further preferred embodiment is that the DNA polymerase is a mixture of enzymes including both the wild-type DNA polymerase from bacteriophage T7 as well as a mutant T7 DNA polymerase that lacks its 3' to 5' exonuclease activity. It is also preferred that the reaction mixture contains a helicase or a primase. The helicase and primase are preferably the gene 4 protein of bacteriophage T7, most preferably the 63-kDa T7 gene 4 protein that encodes the helicase/primase complex. Also preferred in the reaction mixture is a single-stranded DNA binding protein, preferably a single-stranded DNA binding protein from *Escherichia coli*.

Applicant has surprisingly found that such a combination of proteins results in an unexpectedly large amount of DNA synthesis, and that the kinetics of this DNA synthesis is exponential rather than linear in nature. By exponential it is meant that at some period of time during the reaction the rate of DNA synthesis increases. In other words, the amount of DNA synthesized at a particular time will be greater than twice the amount of DNA synthesized at half the time. For example, if the amount of DNA synthesized after 20 minutes is ten times the amount of DNA synthesized after 10 minutes, then the kinetics of DNA synthesis is exponential. On the other hand, if the amount of DNA synthesized after 20 minutes is only twice the amount of DNA synthesized after 10 minutes, then the kinetics of DNA synthesis is linear. In order to test for whether DNA synthesis is linear or exponential, a small amount of radioactive



deoxynucleoside triphosphate can be added to the mixture; *e.g.* example, one million cpm of [ $\alpha$ - $^{32}$ P]dATP in a 50  $\mu$ l reaction, or a specific activity of 40 cpm per pmol of dATP if the dATP concentration is 500  $\mu$ M. Once the reaction is initiated by the addition of the enzyme mixture, aliquots are removed at varying times and the amount of DNA synthesized is determined, as described in Example 1. For example, 5  $\mu$ l aliquots could be removed at 0, 1, 2, 4, 8, 16 and 32 min (although the amount of each aliquot removed and the times can vary). The amount of DNA synthesized at the different times is plotted as a function of the time of the reaction. If DNA synthesis is exponential, then the shape of the plot will be sigmoidal with the slope of the curve increasing over some portion of the time interval. A reaction will be exponential only for a portion of the time; the dNTPs will be rapidly used and the rate of new DNA synthesis will then rapidly decrease. Thus to determine if a reaction is exponential it is important to inspect the kinetics of DNA synthesis over a broad range of time (for example, from 0 to 30 min), and determine if the rate of DNA synthesis increases over any period of the time of the reaction. Also, the total amount of DNA synthesized is quite large, typically at least 10-fold more than the amount of template added to the reaction mixture, and preferably is at least 100-fold greater, or at least 1000-fold greater, or at least 10,000-fold greater, or at least 100,000-fold greater, or at least 1,000,000-fold greater, or at least 10,000,000-fold greater or even more. Thus, one unique feature of the present invention is that synthesis is both exponential and results in the production of a large amount of DNA, *e.g.* 10-100 times or more greater than the amount of template. Furthermore as the present invention does not require the use of exogenously added primers specific for each template being amplified, it provides a generalized amplification method not limited to the region of the DNA template delineated by primers.

Another very effective method of monitoring the amount of DNA synthesized using the isothermal exponential amplification system is to use fluorescent probes or fluorescent dyes such as SYBR Green II (Molecular Probes, Eugene, OR) to continuously report the amount of DNA in the reaction in real time. The basic principle of this method has been used successfully to determine the initial concentrations of specific RNAs and DNAs in a PCR reaction (Gibson *et al.*, Genome Research 6:995, 1996). Fluorescence can be monitored at intervals of 15 seconds using fluorescence-

based real-time PCR instruments such as the PE Biosystems 5700 (PE Biosystems, Foster City, CA), the Roche LightCycler (Indianapolis, IN), or fluorescence microtiter plate readers that can maintain a constant temperature and can carry out kinetic measurements such as the SPECTRAmax (Molecular Devices, Sunnyvale, CA).

5 A preferred embodiment for amplification of a plasmid template utilizes a polymerase in the reaction mixture such that the production of amplified product does not require exogenously-added oligonucleotide primers, the amplification is exponential, and the amount of amplified product is at least 10-fold greater than the amount of plasmid template DNA put into the reaction mixture. More preferably the amplified  
10 product is at least 100-fold greater or 1,000-fold greater, even more preferably at least 10,000-fold greater or 100,000-fold greater, and most preferably at least 1,000,000-fold greater or 10,000,000-fold greater.

The present invention also concerns the use of other reagents that further enhance amplification carried out by a core set of enzymes (the DNA polymerase, the helicase,  
15 the primase and the single-stranded binding protein). Thus, in further preferred embodiments the amplification reaction mixture also includes one or more of the following: a single-stranded binding protein of bacteriophage T7 (gene 2.5 protein), a nucleoside diphosphokinase such as that from *E. coli*, inorganic pyrophosphatase such as that from *E. coli*, an ATP regeneration system such as the combination of creatine  
20 kinase and phosphocreatine, a 5' to 3' exonuclease, preferably the 5' to 3' exonuclease is the bacteriophage T7 gene 6 exonuclease, and a ligase, preferably the ligase is the bacteriophage T7 DNA ligase. In another preferred embodiment the invention features the addition of chemical additives to the reaction mixture that increase the efficiency of the amplification reaction. Three examples of such additives are potassium glutamate,  
25 DMSO, and dextran polymer.

In another preferred embodiment the invention features the treatment of the enzymes prior to inclusion in the reaction mixture with UV irradiation in order to reduce the amplification that arises from the contaminating DNA in the enzyme preparations. Preferably the ultraviolet light is a dose from 10 to 1000  $\mu\text{W}/\text{cm}^2$ , or preferably 100 to  
30 1000  $\mu\text{W}/\text{cm}^2$ , for from 15 sec to 15 min.

In other preferred embodiments, the amplification of the DNA is exponential, the DNA molecule to be amplified can range in size from 1,000 bp, to 2,000 bp, 3,000 bp, 5,000 bp, 10,000 bp, or even larger, up to a complete eukaryotic genome. Both single-stranded and double-stranded DNAs are amplified efficiently. When the DNA fragments being amplified are less than one thousand bases, they can be very poor substrates for amplification unless they are circular. Amplification of the DNA molecule is such that the amount of amplified product is preferably at least about 10-fold amplified, more preferably the amplified product is 100-fold greater, or at least 1000-fold greater, or at least 10,000-fold greater, or at least 100,000-fold greater, or at least 1,000,000-fold greater, or at least 10,000,000-fold greater or even more than the DNA put into the reaction mixture. The constant temperature at which the amplification reaction is carried out is preferably less than 60 °C, more preferably less than 45 °C, and more preferably at 37 °C.

In yet another preferred embodiment the reaction mixture contains ATP and CTP in addition to the 4 dNTPs.

A preferred source of the DNA molecule to be amplified is lysed *E. coli* cells and the DNA is either a plasmid, a bacterial artificial chromosome (BAC) or phage DNA. Preferably, the added plasmid DNA template is less than 100 ng and the reaction mixture is 10 to 200 µl.

The present method of amplification is especially useful to prepare a DNA molecule for use in a DNA sequencing reaction. Preferably, the reaction mixture containing the amplified DNA is treated with a phosphatase to remove dNTPs that may be present prior to use in a sequencing reaction. A preferred phosphatase is from arctic shrimp. The sequencing reaction preferably uses fluorescent primers or fluorescent dideoxynucleotides. A preferred embodiment for preparation of a DNA molecule for use in a sequencing reaction utilizes a polymerase in a reaction mixture such that the production of amplified product does not require exogenously-added oligonucleotide primers, the amplification is exponential, and the amount of amplified product is at least 10-fold greater than the amount of template DNA put into the mixture. More preferably the amplified product is 100-fold greater, or at least 1000-fold greater, or at least 10,000-

fold greater, or at least 100,000-fold greater, or at least 1,000,000-fold greater, or at least 10,000,000-fold greater or even more than the DNA put into the reaction mixture.

In another embodiment, the invention features a method of isolating plasmid, phage, or bacterial artificial chromosomes (BAC) from bacterial cells, such as from *E. coli* cells and then amplifying that DNA in a reaction mixture that does not require exogenous oligonucleotide primers. The bacterial cells are lysed to form a lysate which is then added to the amplification reaction mixture. Lysis can be carried out by any method known to those who practice the art, such as the use of lysozyme or detergents. A preferred embodiment for amplifying DNA directly from a bacterial cell utilizes a polymerase in a reaction mixture such that the production of amplified product does not require exogenously-added oligonucleotide primers, the amplification is exponential, and the amount of amplified product is at least 10-fold greater than the amount of DNA in the lysate that was put into the reaction mixture. More preferably the amplified product is 100-fold greater, or at least 1000-fold greater, or at least 10,000-fold greater, or at least 100,000-fold greater, or at least 1,000,000-fold greater, or at least 10,000,000-fold greater or even more.

Other types of cells can also be used. In preferred embodiments, the cells are eukaryotic microbial cells, e.g., yeast cells, fungal cells or multi-nucleate structure, or mammalian cells, e.g., human cells.

In another embodiment the invention features a method of amplifying genomic DNA in order to preserve small amounts of DNA for the purposes of long-term storage. A preferred embodiment is the amplification of human genomic DNA from a small sample from a patient (e.g. a blood sample or cheek swab) that can then be used for genotype analysis.

In another embodiment the invention features a method of determining the amount of contaminating DNA in a sample by incubating the sample potentially containing contaminating DNA with a reaction mixture comprising a DNA polymerase such that if the contaminating DNA is present the DNA molecule is amplified to produce amplified product. At least one pre-determined amount of control DNA is incubated with the same reaction mixture to produce amplified control product. The amount of amplified product in the sample is compared with the amount of amplified control DNA

as an indication of the amount of contaminating DNA in the sample. Alternatively, the rate of DNA synthesis is monitored in real-time using a probe that fluoresces only when bound to DNA; the greater the amount of input DNA, the shorter the lag period before the exponential phase of DNA synthesis can be observed (see Example 4).

5           Contaminating DNA refers to any DNA that may be contaminating a protein preparation. In general it is most likely to arise from the host organism in which the protein was obtained, and not purified adequately to remove it, or could be acquired during the purification of the protein from columns or reagents. Thus, the most likely source of contaminating DNA is *E. coli*, if the protein is a recombinant protein produced  
10 in *E. coli*, or it is human DNA if the protein is a recombinant protein produced in human tissue culture cells.

By "predetermined amount of control DNA" is meant an amount of DNA such as a supercoiled plasmid (e.g., pUC18 or pUC19) for example in the range of one femtogram to one nanogram in a 50 ul reaction. The synthesis obtained using this  
15 amount of DNA would be compared to an analogous reaction that was carried out in the absence of added DNA. The amount of synthesis at varying times specific for the added plasmid DNA would be determined by subtracting the amount of DNA synthesized in the absence of any DNA from that obtained in the presence of plasmid DNA. In further preferred embodiments the DNA polymerase is from bacteriophage T7, the reaction  
20 mixture further comprises a helicase, a primase and a single stranded binding protein, the control DNA is amplified at least about 10-fold, the reaction does not require exogenously-added oligonucleotide primers, the reaction mixture is carried out at a constant temperature, the control DNA is amplified at least 10-fold, and the kinetics of amplification of the control DNA is exponential.

25           In another aspect, the invention features a kit for amplification. The kit preferably contains a DNA polymerase, a primase, a helicase and a single-stranded binding protein. In further preferred embodiments, the components of the kit comprise those independently selected from the group consisting of a T7 DNA polymerase, the gene 4 protein from bacteriophage (the primase and the helicase), a DNA polymerase mixture of  
30 a wild-type T7 DNA polymerase and a T7 DNA polymerase modified to have reduced 3' to 5' exonuclease activity, and the single-stranded binding protein from *E. coli*. More

preferably, these components, a T7 DNA polymerase, the gene 4 protein from bacteriophage (the primase and the helicase), a DNA polymerase mixture of a wild-type T7 DNA polymerase and a T7 DNA polymerase modified to have reduced 3' to 5' exonuclease activity, and the single-stranded binding protein from *E. coli*, are selected as a group. Still more preferably, the gene 4 protein is the 63-kDa form of the protein. The kit can also contain one or more of the following components: a nucleoside diphosphokinase, an inorganic pyrophosphatase, an ATP regeneration system, preferably consisting of phosphocreatine and creatine kinase, a T7 gene 6 exonuclease, a T7 DNA ligase (gene 1.3 protein) and a single-stranded DNA binding protein of bacteriophage T7 (gene 2.5 protein).

In a further aspect the invention features a method of purifying the T7 helicase/primase (63-kDa gene 4 protein) by overproducing the protein in an *E. coli* cell preferably along with a carboxy terminal fragment of the T7 gene 4 protein, preferably the peptide comprises the carboxy terminus of the protein after residue 221 (glutamine), more preferably the peptide comprises the carboxy terminus of the protein between residues 260 (tryptophan) and residue 280 (serine), even more preferably the peptide initiates at residue 271 (arginine). This gene 4 protein peptide reduces the toxicity of the full-length gene 4 protein to the cells and increases the expression level of the full-length gene by at least 10-fold.

In another embodiment the amplification reaction is carried out in a solid matrix such as agarose. Under these conditions, in the presence of a fluorescent dye such as SYBR Green II (Molecular Probes, Inc., Eugene, OR) the amplification of single DNA molecules can be observed in a fluorescent microscope as focal centers of fluorescence that develop with time and are dependent on added DNA. In preferred embodiment, this amplification in a solid support is used to generate a library of individual "subclones" of DNA molecules, without having to passage the DNA through *E. coli* cells. More preferably these DNA clones are used as templates for DNA sequencing reactions, either by isolating individual clones and amplifying each in solution, or by carrying out the sequencing reaction using fluorescent dideoxy terminators on all the clones together simultaneously on the agarose support. This latter approach would allow a tremendous reduction in the amount (and thus the cost) of fluorescent sequencing reagents. Recently,

Mitra and Church (Nucleic Acids Research 27:e34, 1999) have described a PCR-based system to amplify and detect individual DNA molecules embedded in a polyacrylamide support.

The invention also provides a method for sequencing a DNA sequence by providing amplified DNA, that has been amplified using the constant temperature amplification method described above. The resulting products are sequenced by any of the conventional methods, such as dideoxy termination methods, preferably in an automated sequencer. Preferably the amplification is carried out in a solid matrix, preferably a gel such as an agarose or polyacrylamide gel. Preferably a circular nucleic acid construct with primer sites suitable for extension through the sequence of interest is utilized, e.g., as described in Example 10. Thus, the invention provides an efficient method of sequencing nucleic acids.

As indicated above, the source of the template sequence to be amplified can be from any of a variety of sources, including, for example, nucleic acid molecules from blood from human or non-human, cheek swabs, other tissue samples, microbial cells such as bacterial cells, viruses, plasmids or other cloning vector sequences, or other purified sequences from any source including synthetic sequences.

Other features and advantages of the invention will be apparent from the following description of the preferred embodiments thereof and from the claims.

All articles, publications and patents cited in this application are hereby incorporated by reference, in their entirety.

#### BRIEF DESCRIPTION OF THE DRAWING

Fig. 1 represents the nucleotide sequence of pGP4A/E-1 (SEQ. ID. NO. 1). Nucleotides 1-3 (ATG) correspond to the start codon for the 63-kDa gene 4 protein, while nucleotides 1699-1701 (TGA) correspond to its termination codon. Nucleotides 2785-2787 correspond to the start codon for the truncated gene 4 protein beginning at residue 271 (encoded for by the codon at nucleotides 2785-2787), while nucleotides 3670-3672 (TGA) correspond to the termination codon for this truncated gene.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

The following description and examples are provided for further illustrating various aspects and embodiments of the present invention and are in no way intended to be limiting in scope.

5

DNA polymerase

In general a polymerase for use in this invention is one that carries out extensive DNA synthesis on both strands of a DNA template, with the synthesized DNA in turn being capable of being used as a template for new DNA synthesis. This results in an exponential increase in the amount of DNA synthesized with time. Many replicative DNA polymerases have accessory proteins such as processivity factors, helicases, primases and DNA binding proteins that are specific for each DNA polymerase (for review, see Kornberg and Baker, DNA Replication, Freeman and Co., New York, 1992). Many such combinations will result in efficient DNA synthesis. The bacteriophage T7 replication system provides one example of a polymerase and accessory proteins. Those of skill in the art will appreciate that combinations of polymerases and accessory proteins from other systems (*e.g.* bacteriophage T4 or *E. coli*) will also be useful in the present method of amplification. Such polymerases and accessory proteins can be used with all components being from one system, or can be used in other combinations with functionally compatible proteins from two or more different systems being utilized. Those of skill in the art would be able to determine such combinations of polymerases and accessory based on the Examples and descriptions provided.

In the T7 DNA replication system, a combination of two forms of the T7 DNA polymerase result in the most efficient DNA synthesis. One is the unmodified or native T7 DNA polymerase that has high 3' to 5' exonuclease activity (Tabor *et al.*, J. Biol. Chem. 262:16212, 1987). The other is a form of T7 DNA polymerase that is deficient in its 3' to 5' exonuclease activity. An example of such a DNA polymerase is the  $\Delta 28$  T7 DNA polymerase, missing residues from lysine 118 to arginine 145 (Tabor and Richardson, J. Biol. Chem. 264:6647, 1989; U.S. Patent Nos. 4,942,130 and 4,946,786). Both these forms of T7 DNA polymerase have the processivity factor thioredoxin bound

30



to them in a one-to-one complex. DNA synthesis is most efficient when the exonuclease-deficient form is in excess over native T7 DNA polymerase; the optimum molar ratio is about 20:1. Native and exonuclease deficient polymerases are known for other replication systems, such as bacteriophage T4, bacteriophage Ø29, and *E. coli* DNA polymerases I, II and III.

#### DNA helicase

Most replicative DNA polymerases require a DNA helicase for strand displacement leading strand DNA synthesis. In each replication system the helicase interacts specifically with the DNA polymerase from that system. The energy for helicase activity is obtained by the hydrolysis of nucleoside triphosphates.

The helicase of bacteriophage T7 is the gene 4 protein. Its preferred substrate for hydrolysis is dTTP. The phage makes two forms of the gene 4 protein of molecular weight 56,000 and 63,000; the two forms arise from two in-frame start codons. Efficient amplification requires the 63-kDa form of the gene 4 protein, since, as discussed below, this form also provides primase activity (Bernstein and Richardson, J. Biol. Chem. 264:13066, 1989). Altered forms, substitutions, insertions, deletions, of the 63-kDa protein are also suitable for the present invention. One example is the 63-kDa gene 4 protein in which the methionine at residue 64 is changed to a glycine (G4A<sub>G64</sub>). (Mendelman *et al.*, Proc. Natl. Acad. Sci. USA 89:10638, 1992; Mendelman *et al.*, J. Biol. Chem. 268:27208, 1993). This form of the 63-kDa gene 4 protein is useful because the mutation prevents the initiation of synthesis of the 56-kDa gene 4 protein. All enzymatic properties of the G4A<sub>G64</sub> form of the gene 4 protein that have been examined are comparable to those of the wild-type 63-kDa gene 4 protein, including its use as a primase and helicase for amplification as described in the current invention.

#### DNA primase

In most replication systems, synthesis of the lagging strand is initiated from short oligoribonucleotide primers that are synthesized at specific sites by primases. Specific interactions between a primase and the DNA polymerase allow the DNA polymerase to initiate DNA synthesis from the oligoribonucleotide resulting in the synthesis of the lagging strand.

A unique primase mechanism is found in the terminal protein exemplified by adenovirus and bacteriophage  $\phi$ 29 (Salas, Ann. Rev. Biochem. 60:39, 1991). In these organisms, duplex linear DNA is replicated by the covalent binding of the terminal protein to each 5' end, followed by initiation of DNA synthesis from a mononucleotide bound to the terminal protein by a DNA polymerase. While this system can provide a very efficient form of amplification of the phage genomes such as  $\phi$ 29 DNA (Blanco *et al.*, Proc. Natl. Acad. Sci. USA 91:12198, 1994), the nature of this priming mechanism makes it a difficult system to modify for general amplification of DNAs.

In the T7 replication system, the gene 4 protein, in addition to being a helicase, is also the primase (Bernstein and Richardson, Proc. Natl. Acad. USA 85:396, 1988; Bernstein and Richardson, J. Biol. Chem. 264:13066, 1989). Only the 63-kDa form of the gene 4 protein has primase activity. At specific pentanucleotide recognition sites, the 63-kDa gene 4 protein synthesizes tetranucleotides that are used as primers by T7 DNA polymerase to initiate lagging strand DNA synthesis. The synthesis of RNA primers by 63-kDa gene 4 protein requires ATP and CTP. However, the addition of exogenous short oligonucleotides (7 bases or less) can be used by the gene 4 protein to initiate DNA synthesis primers by T7 DNA polymerase (Scherzinger *et al.*, Nucleic Acids Research 4:4151, 1977). The 63-kDa gene 4 protein is essential for amplification using the T7 replication system. While amplification is most efficient when ATP and CTP are present, it is possible that short oligonucleotides could be added to the reaction mixture to provide priming for amplification in the absence of ATP and CTP. While the native T7 DNA polymerase is capable of interacting with the 63-kDa gene 4 protein to initiate DNA synthesis from RNA primers, the  $\Delta$ 28 T7 DNA polymerase (missing residues 118 to 145) does so much less efficiently. This may be one explanation why a small amount of native T7 DNA polymerase is critical for efficient amplification.

#### Single-stranded DNA binding protein

Most replication systems require single-stranded DNA binding proteins. They serve a number of roles (for review, see Kornberg and Baker, DNA Replication, Freeman and Co., New York, 1992). For example, they remove secondary structure from single-stranded DNA to allow efficient DNA synthesis.

The single-stranded DNA binding protein (ssb) from *E. coli* has been extensively characterized. It has a dramatic effect on improving the processivity of T7 DNA polymerase, particularly at temperatures below 30° C (Tabor *et al.*, J. Biol. Chem. 262:16212, 1987). A low level of ssb is required for efficient amplification using the T7 replication proteins described in this invention. Optimally, the amount of ssb for a 50 µl reaction is from 0.01 to 1 µg.

Bacteriophage T7 also encodes its own single-stranded DNA binding protein, the gene 2.5 protein (Kim *et al.*, J. Biol. Chem. 267:15022, 1992) The presence of this protein is less important for amplification than is the *E. coli* single-stranded DNA binding protein. However, it does stimulate the rate of DNA synthesis several-fold at a concentration of from 0.05 to 5 µg in a 50 µl reaction containing the core amplification components.

#### Nucleoside Diphosphokinase

Nucleoside diphosphokinase rapidly transfers the terminal phosphate from a nucleoside triphosphate to a nucleoside diphosphate. It is relatively nonspecific for the nucleoside, recognizing all four ribo- and deoxyribonucleosides. Thus it efficiently equilibrates the ratio of nucleoside diphosphates and nucleoside triphosphates among all the nucleotides in the mixture. This enzyme can increase the amount of DNA synthesis if one of the required nucleoside triphosphates is preferentially hydrolyzed during the reaction.

In the bacteriophage T7 replication system, the T7 helicase (gene 4 protein) hydrolyzes dTTP to dTDP to provide energy (Matson and Richardson, J. Biol. Chem. 258:14009, 1983). Thus the concentration of dTTP will decrease disproportionately during DNA synthesis. The addition of nucleoside diphosphokinase will re-equilibrate the level of triphosphates of all the nucleosides present. This results in an increase of several-fold in the amount of DNA synthesized. Nucleoside diphosphokinase from Baker's Yeast can be purchased from Sigma Chemical Co. (St. Louis, MO) and the overproduction and purification of nucleoside diphosphokinase from *E. coli* has been described (Almaula *et al.* J. Bact. 177:2524, 1995). Other nucleoside diphosphokinases are known to those who practice the art and are useful in the present invention.

### Inorganic Pyrophosphatase

During a DNA synthesis reaction, inorganic pyrophosphate will accumulate as a product of the reaction. If the concentration becomes too high, it can reduce the amount of DNA synthesis due to product inhibition. This can be prevented by the addition of inorganic pyrophosphatase. Yeast inorganic pyrophosphatase can be purchased from Sigma Chemical Co. (St. Louis, MO). Other inorganic pyrophosphatases are known to those who practice the art and are useful in the present invention.

### ATP-regenerating system

During DNA synthesis in many DNA replication systems, some of the deoxynucleoside triphosphates will be degraded to deoxynucleoside diphosphates due to hydrolysis by the helicase present. This can be minimized by the use of an ATP regeneration system which, in the presence of nucleoside diphosphokinase, will convert any nucleoside diphosphate in the reaction mixture to the triphosphate. In the T7 replication system, the helicase very rapidly degrades dTTP to dTDP for energy. The presence of an ATP-regeneration system will increase the amount of nucleotides capable of serving as precursors for DNA synthesis.

A number of ATP regeneration systems are available and have been widely used. For example, the combination of phosphocreatine and creatine kinase will push the equilibrium between ADP and ATP towards ATP, at the expense of the phosphocreatine. Both phosphocreatine and creatine kinase are available from Sigma Chemical Co. (St. Louis, MO).

### Joining lagging strand fragments

In most DNA replication systems, the lagging strand is synthesized as a series of short, "Okazaki" fragments, that are initiated by a short RNA primer at each 5' end. In order to form a continuous strand, a 5' to 3' exonuclease is needed to remove the RNA primer, and then, after a DNA polymerase fills in the gap, a DNA ligase is needed to seal the nick. In the T7 DNA replication system, these functions are accomplished by the action of the gene 6 protein, a 5' to 3' exonuclease, and the gene 1.3 protein, a DNA ligase (Engler and Richardson, *J. Biol. Chem.* 258:11197, 1983). Purification of the gene 6 protein (Kerr and Sadowski, *J. Biol. Chem.* 247:305, 1972; Engler and

Richardson, J. Biol. Chem. 258:11197, 1983) and gene 1.3 protein (Doherty *et al.*, J. Biol. Chem. 271:11083, 1996) have been described. Exonucleases and ligases from other replication systems are known to those who practice the art and are useful in the present invention.

5                                    Removal of contaminating DNA from enzymes

Since the amplification system described in this invention is nonspecific, contaminating DNA will potentially be a serious source of background. This is particularly a problem in enzyme preparations in which the enzymes bind DNA, such as DNA polymerases, helicases and DNA binding proteins. Most DNA can be removed  
10 from enzyme preparations during purification. For example, an anion exchange column such as DEAE will bind DNA much tighter than it binds proteins.

The residual DNA present in enzyme preparations can be inactivated as a template for DNA synthesis by treatment of the enzymes with ultraviolet light. Effective doses of ultraviolet light are 10 to 1000  $\mu\text{W}/\text{cm}^2$ . In the T7 replication system, treatment  
15 of the enzymes with an ultraviolet dose of 200  $\mu\text{W}/\text{cm}^2$  for 2 min reduces the background synthesis due to contaminating DNA by 99%, while it reduces the activity of the enzymes by less than 20%. This dose of ultraviolet light is equivalent to the dose obtained using a standard, hand-held short-wave ultraviolet light source 10 cm from the enzyme mixture. The enzyme mixture is kept on ice during the treatment with ultraviolet  
20 light to help maintain its activity.

Removal of unincorporated nucleoside triphosphates from reaction mixture

In order to be used for DNA sequencing reactions, the template DNA added to the reactions can only have low levels of unincorporated dNTPs. An efficient method of removing dNTPs is to treat the amplified product with alkaline phosphatase such as that  
25 from arctic shrimp (Amersham Pharmacia Biotech, Inc., Piscataway, NJ). This phosphatase is rapidly inactivated by treatment at elevated temperatures. Thus treatment of a 50  $\mu\text{l}$  reaction mixture with 0.1 to 10 units of shrimp alkaline phosphatase at 20-37°C for 10 to 60 min, followed by treatment at 65-95°C for 10 to 60 min, results in template DNA free of dNTPs that can be added directly to a DNA sequencing reaction  
30 such as the dye terminator reactions using AmpliTaq FS enzyme (Applied Biosystems,

Foster City, CA) or ThermoSequenase enzyme (Amersham Pharmacia Biotech Inc., Piscataway, N.J).

Reaction mixture for amplification using T7 replication proteins

Reaction mixtures contain a buffer at a concentration of 5 mM to 50 mM at pH 7.0 to 8.5, preferably pH 7.5; *e.g.*, 20-40 mM Tris-glutamate, pH 7.5. 2 mM to 15 mM  $Mg^{2+}$  is required, preferably 5 to 10 mM; *e.g.*, 9 mM  $MgCl_2$ . A reducing agent is present, such as 1 to 50 mM dithiothreitol or 2-mercaptoethanol; *e.g.* 5 mM dithiothreitol. The reaction mixture contains 50 to 200 mM potassium glutamate; *e.g.* 100 mM potassium glutamate, which stimulates the reaction. In contrast chloride salts generally inhibit the reaction; if NaCl or KCl are present, preferably their concentration is below 50 mM. The 4 dNTPs are present at a concentration preferably between 20 and 2,000  $\mu M$ , most preferably at a concentration of 500-1000  $\mu M$  of each dNTP. rATP and rCTP are each present at a concentration between 20 and 2000  $\mu M$ , most preferably at a concentration of 500 to 1000  $\mu M$ . Other chemicals may be added that enhance the amplification reaction. For example, DMSO between 0.5 and 8% gives an increased amount of DNA synthesis, most preferably a concentration of 4%. Similarly, a dextran such as Dextran T-10 or Dextran T-500 polysaccharide (Amersham Pharmacia Biotech Inc., Piscataway, NJ) between 0.5 and 20% enhances the amount of DNA synthesis, most preferably a concentration of 5%. After the addition of the DNA to be amplified and the enzyme mixture containing at least some of the T7 replication enzymes described above, the reaction is allowed to proceed at a temperature between 10 ° C and 50° C, preferably 37° C, for 10 to 60 min, preferably 20 min. The reaction can be stopped using standard procedures such as the addition of EDTA to a final concentration of 25 mM, or heating the mixture at 70° C for 20 minutes.

25 Sequencing from Single Molecule Amplification

The amplification reaction can be used to provide DNA for sequencing. The amplification is carried out in a solid matrix such as agarose. Under these conditions, in the presence of a fluorescent dye such as SYBR Green II (Molecular Probes, Inc., Eugene, OR) the amplification of single DNA molecules can be observed in a fluorescent microscope as focal centers of fluorescence that develop with time and are

30

dependent on added DNA. Low-melt agarose such as SeaPlaque Agarose (FMC Products, Rockland, ME) is preferred as the matrix, since the DNA can be easily removed from the agarose when desired by heat. It may be helpful to use a combination of agaroses or other matrices to optimize the stability of the individual foci in the solid support while allowing the DNA to be readily extracted when desired.

If the single molecules being amplified are from a large DNA fragment that has been digested with a restriction enzyme (e.g. EcoRI), and then ligated intramolecularly to form circular molecules, the collection of foci within the agarose support will correspond to a population of "subclones" of that original DNA molecule, comparable to subclones obtained more conventionally by transforming these molecules into *E. coli* and then plating the cells to obtain individual clones. The advantage of the approach described here is that the entire library of clones is generated without the need to transform the DNA into bacteria, and that each "clone" observed as a foci in agarose is pure DNA that can be used directly for sequencing or other applications. Recently, Mitra and Church (Nucleic Acids Research 27:e34, 1999) have described a PCR-based system to amplify and detect individual DNA molecules embedded in a polyacrylamide support.

Several different approaches can be used to sequence the DNA amplified in the agarose support. One would be to pick the DNA from each foci into a tube, heat at 70 °C to melt the agarose, and then add an aliquot to a new amplification reaction mixture to amplify the DNA in solution. This amplified DNA could then be used for DNA sequence analysis as described in Example 5.

Alternatively, the DNA on the agarose support could be sequenced directly. The unincorporated nucleotides from the amplification would first be removed by soaking the agarose in a suitable buffer, e.g., 10 mM Tris-HCl, pH 7.5, 1 mM EDTA. The DNA in the agarose would then be denatured by treatment with 10 mM NaOH for 5 min, followed by neutralization by several washes in 10 mM Tris-HCl, pH 7.5, 1 mM EDTA. The primer used for DNA sequencing would then be soaked into the agarose and allowed to anneal to the denatured DNA. Excess primer would be removed by several washes in 10 mM Tris-HCl, pH 7.5, 1 mM EDTA, and then the DNA sequencing reactions would be carried out by the addition of sequencing reaction buffer, fluorescent nucleotides (e.g. BigDye nucleotides from PE Biosystems (Foster City, CA)), and a DNA polymerase

such as AmpliTaq FS (PE Biosystems, Foster City, CA). The reaction would be initiated by heating the sample to 70 °C. After a 10 min reaction, the sample would be chilled to 4 °C, and the product DNA would be denatured by treatment again with 10 mM NaOH for 5 min. Finally, the salt would be removed by extensive washing in water.

5           The samples are now ready to be directly loaded onto a capillary DNA sequencing instrument. Injection of the capillaries would be a modification of that used in existing instruments such as the PE Biosystems 3700 Genetic Analyzer (Foster City, CA). One direct-load approach would be to insert a 100-1000 capillary array into the sample, heat to 70 °C to melt the agarose, and then apply a voltage to electrosmotically  
10 inject the DNA into each capillary. While only a subset of the capillaries would contain DNA from a single foci, and thus produce unique DNA sequence, the advantages of this procedure would be the ease of carrying out multiple sequencing reactions simultaneously and the low reagent cost of carrying out such parallel reactions.

15

**Example 1: Amplification of purified plasmid DNA using a minimal number of T7 replication proteins**

The reaction mixture (45 µl) contains 20 mM Tris-glutamate, pH 7.5, 9 mM MgCl<sub>2</sub>, 6 mM dithiothreitol, 100 mM potassium glutamate, 3.5% dimethylsulfoxide  
20 (DMSO), 7% Dextran T-500 polysaccharide (Amersham Pharmacia Biotech, Inc., Piscataway, NJ), 550 µM each dGTP, dATP, dTTP and dCTP, 330 µM ATP, 440 µM CTP, and DNA. The DNA samples range from 0.1 pg to 10 ng of supercoiled double-stranded plasmid DNA.

Although circular DNA is a preferred substrate for amplification, linear DNA can  
25 also be used. The DNA can be either single- or double-stranded. In addition to plasmid DNA, other DNAs such as phage lambda DNA, phage M13 DNA bacterial artificial chromosomal (BAC) DNA and genomic DNA (e.g., bacterial or human) are effective templates for amplification. Synthetic DNA such as oligonucleotides 100 nucleotides in length can be used if they are first ligated to form circular molecules. Data suggest that



for circular DNA molecules the initial stages of amplification involve a rolling circle mechanism.

One method of initiating rolling circle DNA synthesis on a plasmid DNA from a specific site that has been described is to use the site-specific single-stranded  
5 endonuclease gene II protein from filamentous phage such as M13 or fd. This protein recognizes a sequence of approximately 50 bp (the plus strand replication origin) and introduces a specific nick into one of the strands (Higashitani *et al.*, Nucleic Acids Research 20:2685, 1992). While the wild-type gene II protein requires supercoiled DNA for activity, mutant proteins have been described (*e.g.* glycine 73 changed to alanine) that  
10 reduce this superhelicity requirement, so that the protein now recognizes relaxed duplex DNA (Higashitani *et al.*, Nucleic Acids Research 20:2685, 1992). The purification of the wild-type and mutant gene II proteins have been described (Greenstein and Horiuchi, J. Molec. Biol. 197:157, 1987; Higashitani *et al.*, Nucleic Acids Research 20:2685, 1992). It has been demonstrated that cleavage of supercoiled DNA with the gene II  
15 protein results in a substrate that the T7 DNA polymerase and T7 gene 4 protein can use to initiate lagging strand DNA synthesis (Harth *et al.*, Eur. J. Biochem. 119:663, 1981). It has also been demonstrated that these molecules promote rolling circle DNA synthesis using the replication proteins from bacteriophage T4 (Meyer *et al.*, J. Biol. Chem. 256:5810, 1981). This class of site-specific nicking proteins may be useful for  
20 stimulating initiation of the amplification process described in this invention. However, in the invention described here the inventors have shown that surprisingly nicking of the circular DNA is not necessary for efficient initiation of DNA synthesis; the combination of T7 gene 4 protein and T7 DNA polymerase are able to efficiently initiate and synthesize DNA on covalently supercoiled plasmid DNAs whether they are supercoiled  
25 or relaxed.

For diagnostic characterization of the extent of DNA synthesis, one of the dNTPs can be radioactively labeled; for example, [<sup>3</sup>H] dTTP can be used at a specific activity of 20 cpm/pmol. The synthesized DNA can also be detected by fluorescence or chemiluminescence by incorporating the appropriate modified nucleotides using standard  
30 techniques (Current Protocols in Molecular Biology, Ausubel *et al.*, eds., John Wiley &

Sons, Inc., 1996). ). In addition, the rate of DNA synthesis can be monitored in real time by the addition of a fluorescent probe to the reaction mixture (see Example 4).

The enzyme mixture (5  $\mu$ l per reaction) is prepared in a buffer containing 20 mM Tris-glutamate, pH 7.5, 10 mM 2-mercaptoethanol and 0.5 mg/ml bovine serum albumin. 5  $\mu$ l of this mixture contains 1  $\mu$ g of  $\Delta$ 28 T7 DNA polymerase, 0.03  $\mu$ g of native T7 DNA polymerase, 1  $\mu$ g of T7 63-kDa gene 4 protein, and 0.3  $\mu$ g of *E. coli* single-stranded DNA binding protein.  $\Delta$ 28 T7 DNA polymerase is a one-to-one mixture of *E. coli* thioredoxin and T7 gene 5 protein with a deletion of residues 118 to 145, as described in Tabor and Richardson, J. Biol. Chem. 264:6647, 1989, and U.S. Patents 10 Nos. 4,942,130 and 4,946,786. It is the same as Sequenase Version 2.0 enzyme sold by Amersham Pharmacia Biotech, Inc. (Piscataway, NJ). Native T7 DNA polymerase is a one-to-one complex of thioredoxin and wild-type T7 gene 5 protein. It can be purchased from Amersham Pharmacia Biotech, Inc. (Piscataway, NJ) or New England Biolabs (Beverly, MA). T7 63-kDa gene 4 protein is the G4A<sub>G64</sub> protein described in 15 Mendelman *et al.*, Proc. Natl. Acad. Sci. USA 89:10638, 1992 and Mendelman *et al.*, J. Biol. Chem. 268:27208, 1993; it is the wild-type gene 4 63 kDa protein except that the methionine at residue 64 has been replaced with a glycine to prevent initiation of synthesis of the 56-kDa form of the gene 4 protein. *E. coli* single-stranded DNA binding protein (ssb) is overproduced and purified as described in Lohman *et al.*, Biochemistry 20 25:21 (1986). It can be purchased from Amersham Pharmacia Biotech, Inc. (Piscataway, NJ).

In order to reduce the level of background DNA synthesis resulting from contaminating DNA, the enzyme mixture is treated with ultraviolet light at a dose of 200  $\mu$ W/cm<sup>2</sup> for 2 min on ice prior to adding to the reaction mixture. This dose corresponds 25 to a distance of about 10 cm from a standard short-wave ultraviolet lamp. Times of 10 sec to 15 min are also effective at reducing the amount of contaminating DNA that can be replicated without affecting significantly the activity of the enzymes.

To initiate the amplification reaction, 5  $\mu$ l of the enzyme mixture is added to the 45  $\mu$ l reaction mixture. The reaction is then allowed to proceed at 37° C for 20 min. The

reaction can be carried out at a temperature of 10°C to 45°C, for 1 min to 2 hr. The reaction is stopped by the addition of 5 µl of 200 mM EDTA.

There are many published procedures to measure the amount of DNA synthesized during the reaction. For example, if [<sup>3</sup>H]dTTP is present, the amount of radioactivity incorporated into DNA can be measured by binding the reaction mixture to a DE81 filter disk, then washing the disk with 0.3 M ammonium formate to remove unincorporated nucleotides, and then measuring the amount of radioactivity retained on the disk using a scintillation counter, as described in Kim *et al.*, J. Biol. Chem. 267:15032 (1992). Alternatively, the DNA can be analyzed by agarose gel electrophoresis and staining with ethidium bromide, either with or without digestion with restriction enzymes, following standard procedures (Current Protocols in Molecular Biology, Ausubel *et al.*, eds., John Wiley & Sons, Inc., 1996).

This procedure should result in the incorporation of about 5% of the total dNTPs into DNA, or about 1.5 µg of DNA synthesized in the 50 µl reaction mixture. If the reaction mixture contains 10 pg of plasmid DNA, this corresponds to a 150,000-fold amplification.

**Example 2: Amplification of purified plasmid DNA using a T7 replication system optimized for maximum rate and amount of DNA synthesis**

The rate and amount of DNA synthesis obtained using the conditions defined in Example 1 can be improved by using the modifications described below. All other components and conditions are identical to those outlined in Example 1.

Phosphocreatine (11 mM) is added to the 45 µl reaction mixture. This in combination with creatine kinase provides an efficient ATP regeneration system. The phosphocreatine that is most effective is synthetic phosphocreatine, Sigma catalog number P6502 (Sigma Chemical Co., St. Louis, MO), dissolved in H<sub>2</sub>O to a concentration of 500 mM.

The enzyme mixture contains at least some of the following six enzymes. These are in addition to the Δ28 T7 DNA polymerase, native T7 DNA polymerase, 63-kDa

gene 4 protein and *E. coli* single-stranded DNA binding protein, which are present in the amounts defined in Example 1.

(1) Creatine kinase, rabbit muscle (Boehringer catalog number 127566) (Boehringer Mannheim, Indianapolis, IN). A stock solution is prepared at 100 mg/ml in H<sub>2</sub>O and stored at -40° C. Each enzyme mixture (5 µl) for a 1X reaction (50 µl total volume) contains 2 µg of creatine kinase. Creatine kinase, in conjunction with phosphocreatine and nucleoside diphosphokinase, provides an ATP regeneration system that converts nucleoside diphosphates that arise during the reaction to nucleoside triphosphates.

(2) Nucleoside diphosphokinase, Baker's yeast (Sigma catalog number N0379) (Sigma Chemical Co., St. Louis, MO). A stock solution is prepared at 1 mg/ml in H<sub>2</sub>O and stored at -40° C. Each enzyme mixture (5 µl) for a 1X reaction (50 µl total volume) contains 50 ng of nucleoside diphosphokinase. Alternatively, nucleoside diphosphokinase can be overproduced and purified from *E. coli* (Almaula *et al.*, J. Bact. 177:2524, 1995). 50 ng of the *E. coli* enzyme is used for a 1X reaction. Nucleoside diphosphokinase is added to maintain an equal ratio of all four dNTPs.

(3) Inorganic pyrophosphatase, Baker's yeast (Sigma catalog number I1891) (Sigma Chemical Co., St. Louis, MO). A stock solution is prepared at 1 mg/ml in H<sub>2</sub>O and stored at -40° C. Each enzyme mixture (5 µl) for a 1X reaction (50 µl total volume) contains 20 ng of inorganic pyrophosphatase. Inorganic pyrophosphatase is added to degrade the pyrophosphate that will build up during polymerization and could inhibit DNA synthesis by product inhibition.

(4) T7 single-stranded DNA binding protein (T7 gene 2.5 protein). T7 single-stranded DNA binding protein is overproduced and purified as described by Kim *et al.* (J. Biol. Chem., 267:15022, 1992). Each enzyme mixture (5 µl) for a 1X reaction (50 µl total volume) contains 1 µg of T7 single-stranded DNA binding protein. The T7 single-stranded DNA binding protein will increase the rate of DNA synthesis by several-fold. One possible explanation for this is that the T7 single-stranded DNA binding protein is necessary to have efficient coupling of leading and lagging strand DNA synthesis by the

T7 DNA polymerase and helicase/primase complex (Park *et al.*, J. Biol. Chem. 273:5260, 1998; Lee *et al.*, Mol. Cell. 1:1001, 1998).

5 (5) T7 gene 6 exonuclease. The combination of T7 gene 6 exonuclease and T7 DNA ligase (T7 gene 1.3 protein) are used to remove the RNA primers from the 5' ends of the lagging strand fragments and then, after the gaps are filled in by the wild-type T7 DNA polymerase, seal the resulting nicks. The T7 gene 6 exonuclease is purified as described in Kerr and Sadowski, J. Biol. Chem. 247:305, 1972 and Engler and Richardson, J. Biol. Chem. 258:11197, 1983. Each enzyme mixture (5  $\mu$ l) for a 1X reaction (50  $\mu$ l total volume) contains 50 ng of T7 gene 6 exonuclease.

10 (6) T7 DNA ligase (gene 1.3 protein). The T7 ligase is added to seal any nicks that are present in the synthesized DNA, in particular those that occur on the lagging strand. T7 DNA ligase is overproduced and purified as described in Doherty *et al.*, J. Biol. Chem. 271:11083, 1996. Each enzyme mixture (5  $\mu$ l) for a 1X reaction (50  $\mu$ l total volume) contains 0.1  $\mu$ g of T7 DNA ligase.

15 As in Example 1, the enzyme mixture is treated with an ultraviolet dose of 200  $\mu$ W/cm<sup>2</sup> for 2 min on ice prior to adding to the reaction mixture. The amplification reaction is then initiated by the addition of 5  $\mu$ l of the enzyme mixture to the 45  $\mu$ l reaction mixture. The reaction is allowed to proceed at 37° C for 20 min, and then stopped by the addition of 5  $\mu$ l of 200 mM EDTA. The products are analyzed as described in Example 1.

20 This procedure should result in the incorporation of up to 50% of the total dNTPs into DNA, or about 15  $\mu$ g of DNA synthesized in the 50  $\mu$ l reaction mixture. If the initial reaction mixture contains 10 pg of plasmid DNA, this corresponds to a 1,500,000-fold amplification.

25

Example 3: Assay to determine whether amplification is exponential

Reactions as described in Examples 1 and 2 are carried out using varying amounts of plasmid DNA and for varying lengths of time. A radioactively labeled dNTP is used in order to determine the amount of DNA synthesis; for example, 50 cpm/pmole [3H]dTTP or [ $\alpha$ -<sup>32</sup>P]dATP. 100  $\mu$ l reactions are carried out in the absence of added

30

DNA, and in the presence of 1, 10, 100 and 1,000 pg of supercoiled pUC18 plasmid DNA. Reactions are carried out at 37° C and 20 µl aliquots are removed at 5, 10, 15, 20 and 30 min and stopped by the addition of 5 µl of 200 mM EDTA, pH 8.0. The amount of radioactivity incorporated into DNA is determined using standard methods. For example, the amount of radioactivity that can be precipitated by trichloroacetic acid, which will precipitate DNA but not free nucleotide, can be determined as described in Tabor and Richardson, J. Biol. Chem. 264:6647, 1989. Alternatively, the amount of radioactivity that is retained by DE81 filter disks in the presence of 0.3 M ammonium formate, pH 8.0, can be determined (Kim *et al.*, J. Biol. Chem. 267:15032, 1992); DNA but not dNTPs is retained by the disks under these conditions.

For each time point, the amount of DNA synthesized in the absence of added DNA is subtracted from the amount synthesized in the presence of DNA. Up to 20 min, the amount of DNA synthesized in the absence of added input DNA should be very low (for example, less than 1% the amount synthesized in the presence of 100 pg of added input DNA).

For each amount of input DNA, the amount of DNA synthesized above the background level synthesized in the absence of input DNA is plotted as a function of the time of the reaction. If amplification is exponential, then the rate of synthesis of DNA will increase during some portion of the time course (i.e., the plot will be sigmoidal). For example, the amount of DNA synthesized after 15 min might be 50-fold higher than the amount synthesized after 10 min.

#### Example 4: Real-time amplification assay using fluorescence

A very effective method of monitoring the amount of DNA synthesized using the isothermal exponential amplification system is to use fluorescent probes to continuously report the amount of DNA in the reaction in real time. The basic principle of this method has been used successfully to determine the initial concentrations of specific RNAs and DNAs in a PCR reaction (Gibson *et al.*, Genome Research 6:995, 1996). The higher the initial DNA concentration, the shorter the lag period before the exponential phase of DNA synthesis. To measure the amount of DNA synthesis in real time, reactions were carried out as described in Example 3 except that radioactivity was

omitted and the fluorescent dye SYBR Green II (Molecular Probes, Eugene, OR) was added at a final concentration of 1/80,000 the stock solution. The dye was added after the reaction mixture was treated with UV light to destroy contaminating DNA. SYBR Green II has virtually no fluorescence on its own but fluoresces very strongly when  
5 bound to nucleic acid. At concentrations higher than 1/80,000 of the initial stock it is inhibitory to the amplification reaction described in Example 3. Other dyes that fluoresce specifically when bound to nucleic acid will also work in the present invention (e.g. SYBR Green I, Pico Green, Oligreen, SYTO 11, SYTO 12, SYTO 13, SYTO 14, SYTO 15, SYTO 16 (Molecular Probes, Eugene, OR), and SYBR Green (PE  
10 Biosystems, Foster City, CA)) or any other dye which shows significantly enhanced fluorescence when bound to DNA. Preferably, the significantly enhanced fluorescence refers to at least a 10-fold increase in fluorescence upon binding to DNA. More preferably, DNA binding will result in at least a 30-fold, 100-fold, 300-fold or 1000-fold increase in fluorescence.

15 In the example, reactions are carried out at 37 °C. Fluorescence is monitored once every 15 sec for 30 min. Exemplary instruments that can be used to measure fluorescence include real-time PCR instruments such as the PE Biosystems 5700 (PE Biosystems, Foster City, CA) and the Roche LightCycler (Indianapolis, IN), and fluorescence microtiter plate readers that can maintain a constant temperature and can  
20 carry out kinetic measurements, such as the SPECTRAmax (Molecular Devices, Sunnyvale, CA). Ideally the SYBR Green II fluorescence is monitored using an excitation wavelength of 480 nM and an emission wavelength of 530 nM.

In order to measure the DNA concentration in unknown samples, a standard curve is determined for each experiment by carrying out reactions either in the absence  
25 of DNA (blank) or in the presence of known concentrations of a standard DNA such as supercoiled pUC18 (e. g. ten-fold dilutions from 0.1 pg to 10 ng). A standard curve is determined by plotting the time required for fluorescence to reach half-maximum (or alternatively the time required for the relative fluorescence to reach some threshold value) versus the log of the initial concentration of DNA. This plot yields a straight line  
30 over at least three orders of magnitude of initial DNA concentration. A comparison of

the values obtained using unknown samples to the standard curve allows one to extrapolate the amount of DNA present at the outset in each sample.

Example 5: Use of amplified DNA as a template for DNA sequencing reactions

5 The following example shows how the amplification reaction can be used to generate enough template from a small amount of DNA to be sequenced using conventional radioactive and nonradioactive dideoxy terminator sequencing strategies. While the example shown is starting from a small amount of plasmid DNA, this procedure will also work using other DNAs such as those from bacterial artificial  
10 chromosomes (BACs), phage lambda, phage M13, or genomic DNA (*e.g.*, bacterial or human). Also, while the example shown uses purified DNA, much less pure samples, such as that from freshly lysed *E. coli* cells, can also be used.

Supercoiled pUC18 DNA (100 pg) is added to a 50  $\mu$ l reaction mixture as described in Example 2, and amplified at 37°C for 20 min. The reaction is stopped by  
15 heating the mixture at 70°C for 5 min. 0.5 units of arctic shrimp alkaline phosphatase (Amersham catalog number E70092) (Amersham Pharmacia Biotech, Inc., Piscataway, NJ) is added, and the mixture is incubated at 37°C for 30 min. The phosphatase is then inactivated by incubation at 80°C for 15 min. The amplified DNA can now be directly added to a DNA sequencing reaction. Other methods can also be used to purify the  
20 template DNA away from unincorporated dNTPs. These include precipitating the DNA with ethanol or isopropanol, and purifying the DNA by binding it to a silica matrix, for example using a commercial kit such as the QIAquick PCR Purification Kit, product number 28104 (Qiagen, Inc., Valencia, CA).

In order to sequence the amplified DNA using the PE Biosystems BigDye  
25 Terminators (catalog number 4303149 (PE Biosystems, Foster City, CA), 1  $\mu$ l of the amplified DNA (500 ng) that had been treated with arctic shrimp phosphatase was added to 8  $\mu$ l of the BigDye Reaction Mixture, 1.5  $\mu$ l (3 pmoles) of the "universal" forward primer, and 9.5  $\mu$ l of water. Cycle sequencing was carried out using 25 cycles consisting of 96 °C for 10 sec, 50 °C for 5 sec, and then 60 °C for 4 min. After the reaction was  
30 complete the unincorporated dye terminators were removed by gel filtration centrifugation using Centri-Sep columns (Princeton Separations, Inc., Princeton, NJ) and



the eluent was used directly for capillary sequence analysis on the PE Biosystems 310 single-capillary Genetic Analyzer or the PE Biosystems 3700 96-capillary Genetic Analyzer. The accuracy of the sequence was analyzed by determining the number of bases that had a Phred score greater than 20 (Ewing et al., Genome Research 8:186, 5 1998). In general, three times less amplified DNA was required compared with supercoiled plasmid DNA to produce DNA sequence of a given quality as analyzed by their relative Phred scores. This is probably due to the fact that primers will anneal more efficiently to the amplified DNA since it is linear compared with the supercoiled plasmid DNA.

10 The amplified DNA can also be sequenced using other methods of labeling the fragments. For example, it can be used as a template for reactions containing radioactively labeled dideoxyterminators; 2  $\mu$ l of the DNA can be added directly to the 20  $\mu$ l reaction mixture from a ThermoSequenase radiolabeled terminator cycle sequencing kit, Amersham Pharmacia product number US 79750 (Amersham Pharmacia Biotech, Inc., 15 Piscataway, NJ). The DNA can also be used for DNA sequencing techniques that use either radioactive or fluorescently labeled primers.

Example 6: Amplification and sequencing of plasmid DNA directly from a bacterial colony

20 The procedure below describes the amplification *in vitro* of DNA obtained from direct lysis of a bacterial colony, followed by DNA sequencing of the amplified DNA. While the example is using cells containing a high copy plasmid (pUC18), one skilled in the art will recognize that cells containing other types of DNAs, such as bacterial artificial chromosome (BAC) DNAs, could also be used in this procedure.

25 Bacterial cells containing the plasmid to be amplified and sequenced, for example a pUC18 derivative, are plated on LB plates containing 50  $\mu$ g/ml ampicillin at 37° C overnight. A single colony is placed in 500  $\mu$ l of a solution containing 20 mM Tris-HCl, pH 7.5, 1 mM EDTA and 100  $\mu$ g lysozyme. After incubation at room temperature for 15 min, the solution is centrifuged at 14,000 rpm for 10 min. 1  $\mu$ l of the supernatant is then 30 added to a 50  $\mu$ l amplification reaction as described in Example 2. The resulting

amplified DNA is then sequenced as described in Example 4. While the centrifugation step removes genomic DNA and thus improves the purity of the sample, it is possible to avoid this step and use the lysed sample directly for amplification and sequencing.

While the lysis method described here uses lysozyme, other methods could be used, such as the addition of nonionic detergents (e.g. Triton X-100 or Tween 20) or the expression of the T7 lysozyme gene (gene 3.5 protein) within the *E. coli* cell (pLysS and pLysE, Novagen, Inc., Madison, WI). It is possible that the method of lysis will have to be optimized for plasmids of varying size in order to minimize the amount of contaminating chromosomal DNA released.

10

Example 7: BAC end sequencing by selective amplification of the ends of the insert

In this example the ends of BAC clones are sequenced by first preferentially amplifying DNA fragments containing the ends of each insert. This is accomplished by digesting the BAC DNA and then carrying out a ligation in the presence of a splint that will result in circularization of only the two end fragments. Since the amplification system described in Example 2 preferentially amplifies circular DNA when the DNA fragments are less than several kbp, this circularization results in an enrichment of the end fragments in the final amplified mixture. (This is not necessarily the case for other amplification systems of the invention.) While the example below uses the vector pBELO-11 (Research Genetics, Huntsville, AL), which has a single Bae I site and no Mlu I sites, the strategy used can readily be applied to other BAC vectors by choosing the appropriate restriction enzymes and designing the appropriate splint oligonucleotides.

pBELO-11 BAC DNA containing inserts is isolated from *E. coli* cells using standard procedures. 50 ng of the DNA is treated with the enzymes Bae I, Mlu I and T4 DNA ligase (New England Biolabs, Beverly, MA) and a 50-fold molar excess of the following three oligonucleotides: BAC-1 (5'-CGCGGTACACCGACGTCAA-3'), BAC-2 (5'-CGCGGTACACCGACTTAAT-3') and BAC-3 (5'-GTCGGTGTAC-3'). BAC-1 and BAC-3 will anneal to form a splint that will result in circularization and ligation of one end of the pBELO-11 DNA, while BAC-2 and BAC-3 will anneal to form a splint that will result in circularization and ligation of the other end of the pBELO-11 DNA. The

30

reactions are carried out in 20  $\mu$ l in 20 mM Tris-HCl, pH 7.5, 10 mM MgCl<sub>2</sub>, 5 mM DTT, 20  $\mu$ M S-adenosylmethionine and 500  $\mu$ M ATP at 37 °C for 60 min. 5  $\mu$ l of the ligated products are amplified as described in Example 2 in a 50  $\mu$ l reaction. After removal of the unincorporated nucleotides, the amplified DNA is sequenced using the  
5 universal T7 (TAATACGACTCACTATAGGGCGA) or SP6 (CATACGATTTAGGTGACACTATAG) primers that anneal upstream of each of the two ends of the insert in pBELO-11.

Example 8: Generic amplification of human genomic DNA

10 A cheek swab is taken of a human subject and applied to FTA Card (GIBCO BRL, Rockville, MD). The DNA bound to the paper is purified using FTA Purification Reagent as described by the distributor (GIBCO BRL, Rockville, MD). The DNA is eluted from the paper in 10 mM Tris-HCl, pH 7.5, 1 mM EDTA by heating at 80 °C for 5 min. The concentration of the eluted DNA is approximately 1 ng/ $\mu$ l. One  $\mu$ l of this  
15 DNA is amplified in a 50  $\mu$ l reaction as described in Example 2, incubating at 37 °C for 20 min. The concentration of the DNA in the completed reaction is 0.5  $\mu$ g/ $\mu$ l, a 25,000-fold amplification. Aliquots of this DNA can be used directly for genotyping using PCR reactions; e.g. 1  $\mu$ l in a 25  $\mu$ l PCR reaction.

20 Example 9: Detection of contaminating DNA by amplification reaction

In order to characterize the level of contamination of DNA in an unknown sample, reactions are carried out as described in Example 2 using varying amounts of the unknown sample. As a control, a standard series of reactions are also carried out both in the absence of DNA and in the presence of 1, 10, 100 and 1,000 pg of a standard  
25 supercoiled plasmid DNA such as pUC18. The reactions are carried out at 37°C for 15 min. The reaction mixtures are carried out in the presence of 20 cpm/pmole [<sup>3</sup>H]TTP, and the amount of DNA synthesis in each sample is determined by measuring the amount of radioactivity taken up into DNA, as described in Example 1. Alternatively, DNA synthesis can be monitored in real time using a fluorescent probe (see Example 4).  
30 The amounts of DNA synthesized in the unknown samples are compared with the

amounts of DNA synthesized in the standard reactions containing different initial concentrations of the known DNA in order to extrapolate the relative amount of contaminating DNA in the unknown sample.

5 Example 10: Single molecule amplification in agarose

In accordance with the description above, in this example, single DNA molecules are amplified as individual foci embedded in a solid matrix, such as agarose. The amplified DNA is detected by fluorescence in the presence of the dye SYBR Green II (Molecular Probes, Eugene, OR).

10 Reactions are carried out as described in Example 4. Two-fold concentrated reaction mixtures are prepared, and then diluted two-fold with a solution containing liquid 1% SeaPlaque agarose (FMC Products, Rockland, ME). The agarose is prepared in water and dissolved at 90 °C, then cooled to 37 °C before mixing with the reaction mixtures. The reaction mixtures contain a final 1/80,000 X concentration of SYBR  
15 Green II fluorescent dye, and varying amounts of supercoiled pUC18 DNA, ranging from none up to 100,000 molecules per  $\mu\text{l}$ . After mixing the agarose and the reaction mixture, the mixture is plated on a microscope slide, covered with a cover slip, and placed on ice to solidify the mixture. The slide is then heated at 37 °C and the progression of foci is monitored using a fluorescence microscope, exciting at a  
20 wavelength of 480 nM and observing the emission at 530 nM. In the absence of added DNA, approximately 20 foci are observed on a one  $\text{cm}^2$  region, that correspond to the amplification of contaminating DNA. In the presence of increasing amounts of pUC18 DNA, the number of foci increasing proportionally, and thus correspond to amplification of the added DNA. Each foci corresponds to the amplification of a single DNA  
25 molecule, and thus is "clonal".

While the example below uses supercoiled pUC18 DNA as a template, one skilled in the art will recognize that other circular DNAs could be used. For example, if BAC DNA was digested with EcoRI, and then ligated, a population of circular fragments, or a "library", would be generated. If a splint oligonucleotide was ligated  
30 between the EcoRI sites, these could then serve as primer sites to sequence each insert in both directions. Using this strategy, all the amplified foci embedded on an agarose bed

could be sequenced in a single dye-terminator sequencing reaction. After carrying out the sequencing reactions, the unincorporated dye terminators and the salts could be washed out of the agarose and each foci could be sequenced directly by injection into a capillary-based sequencing instrument such as the PE Biosystems 3700. This would allow one to generate a library of subclones of a large DNA fragment rapidly without the use of *E. coli* cells, and then to sequence all the fragments in a single sequencing reaction, that could be directly loaded onto a capillary sequencing instrument. This would represent a tremendous reduction in the scale of required reagents and human labor involved in cloning and sequencing fragments.

10

Example 11: Overproduction and purification of the 63-kDa T7 gene 4 protein

A preferred T7 gene 4 protein used in this invention is the 63-kDa form referred to as G4A<sub>G64</sub> in Mendelman *et al.*, (Proc. Natl. Acad. Sci. USA 89:10638, 1992) and Mendelman *et al.*, (J. Biol. Chem. 268:27208, 1993). It is the wild-type 63-kDa gene 4 protein except that the methionine at residue 64 has been replaced with a glycine to prevent initiation of synthesis of the 56-kDa form of the gene 4 protein. Expression of the T7 gene 4 in *E. coli* is toxic to the cells and must be kept tightly repressed when uninduced. Furthermore, when induced, the toxicity prevents efficient overproduction of the protein. It is known that mutants of gene 4 protein such as those in the nucleotide binding site eliminate this toxicity and result in much greater overproduction of the protein (Notarnicola and Richardson, J. Biol. Chem. 268:27198, 1993). This observation suggests that it is the potent TTPase activity of the wild-type gene 4 protein that is responsible for the toxicity. Thus one method to increase the amount of wild-type gene 4 protein overproduced in cells is to coexpress another molecule that will bind to the wild-type gene 4 protein and inhibit its TTPase activity. Three possible classes of molecules that may accomplish this are protein, RNA and DNA aptamers (Cohen *et al.*, Proc. Natl. Acad. Sci. USA 95:14272, 1998; Famulok and Jenne, Curr. Opin. Chem. Biol. 2:320, 1998), which can be selected for from libraries on the basis of either binding to wild-type gene 4 protein *in vitro*, or inhibit the toxicity of its production *in vivo*.

25

An alternative approach is to coexpress a fragment of gene 4 that interferes with the activity of the wild-type gene 4 protein. The 295 residue carboxyl-terminal fragment of gene 4 protein from arginine 271 to the end of the gene has this property; *in vitro* it inhibits the wild-type gene 4 protein's TTPase activity, and *in vivo* it reduces the toxicity of the wild-type gene 4 and increases the overproduction of the wild-type gene 4 protein by at least 10-fold (Guo *et al.*, J. Biol. Chem. 274:30303, 1999). It is likely that other carboxyl-terminal fragments of the gene 4 will have a similar effect.

To overproduce the G4A<sub>64A</sub> form of the gene 4 protein, the plasmid pGP4A/4E-1 whose sequence is shown in Figure 1 (SEQ. ID. NO. 1), was used. It expresses both the G4A<sub>64A</sub> gene 4 and the gene 4 fragment initiating at codon 271, both from T7 RNA polymerase promoters. The two genes are in tandem, with the G4A<sub>64A</sub> expressed upstream of the gene 4 fragment that starts at codon 271 (see Figure 1). pGP4A/4E-1 was used to transform the *E. coli* cells BL21/DE3 (Novagen, Inc., Madison, WI) and selected for by plating on agar plates containing 50 µg/ml kanamycin. The cells were grown in LB media containing 50 µg/ml kanamycin at 30°C. When the cell density reached A<sub>590</sub>=1, the gene 4 proteins were induced by the addition of 0.5 mM IPTG. After inducing for 3 hours at 30°C, the cells were harvested and frozen at -80°C.

The 63-kDa gene 4 protein was purified by the procedure described by Notarnicola *et al.* (J. Biol. Chem. 270:20215, 1995). Briefly, the cells were lysed by the addition of lysozyme and by sonication. After the cell debris was removed by centrifugation, the 63-kDa gene 4 protein was precipitated by the addition of 15% polyethylene glycol 4000. After resuspension of the precipitate, the gene 4 protein was purified by phosphocellulose chromatography. Finally, the pool of the fractions containing gene 4 protein from the phosphocellulose column was purified by ATP-agarose affinity chromatography. The gene 4 protein fragment present in the extract separates from the G4A<sub>64A</sub> gene 4 protein both in the PEG precipitation, the phosphocellulose chromatography, and the ATP affinity chromatography. From 10 liters of induced cell culture, 100 mg of pure G4A<sub>64A</sub> gene 4 protein were obtained.

Other embodiments within the scope of the following claims.

CLAIMS

What is claimed is:

1. A method of amplifying a template DNA molecule comprising:  
incubating said template DNA molecule with a reaction mixture  
5 comprising a DNA polymerase and at least one accessory protein at a constant  
temperature to produce amplified product, wherein production of amplified product  
does not require exogenously-added oligonucleotide primers and said template DNA  
molecule does not have a terminal protein covalently bound to either 5' end.
2. The method of claim 1, wherein said DNA polymerase is a  
10 bacteriophage DNA polymerase.
3. The method of claim 1, wherein said DNA polymerase is  
bacteriophage T7 DNA polymerase.
4. The method of claim 3, wherein said DNA polymerase is a mixture of  
wild-type T7 DNA polymerase and a T7 DNA polymerase modified to have reduced  
15 3' to 5' exonuclease activity.
5. The method of claim 1, wherein said accessory protein is a helicase.
6. The method of claim 1, wherein said accessory protein is a primase.
7. The method of claim 1, wherein said accessory protein is the  
helicase/primase of bacteriophage T7.
- 20 8. The method of claim 7, wherein said helicase/primase is the 63-kDa  
form of the protein from bacteriophage T7.
9. The method of claim 1, wherein said accessory protein is a single-  
stranded DNA binding protein.
- 25 10. The method of claim 9, wherein said single-stranded DNA binding  
protein is from *Escherichia coli*.

11. A method of amplifying a template DNA molecule comprising:  
incubating said template DNA molecule with an *in vitro* reaction  
mixture comprising a DNA polymerase, a helicase, and a primase at a constant  
temperature to produce amplified product.
- 5 12. The method of claim 11, wherein said reaction mixture does not  
require exogenously-added oligonucleotide primers.
13. The method of claim 11, wherein said polymerase is from  
bacteriophage T7.
14. The method of claim 13, wherein said DNA polymerase is a mixture of  
10 wild-type T7 DNA polymerase and a T7 DNA polymerase modified to have reduced  
3' to 5' exonuclease activity.
15. The method of claim 11, wherein said helicase and said primase is the  
helicase/primase of bacteriophage T7 .
16. The method of claim 15, wherein said helicase/primase is the 63-kDa  
15 form of the protein from bacteriophage T7.
17. The method of claim 11, wherein said reaction mixture further  
comprises a single-stranded DNA binding protein.
18. The method of claim 17, wherein said single-stranded binding protein  
is from *Escherichia coli*.
- 20 19. The method of claim 11, where the amount of DNA produced is a  
1,000-fold increase over the amount of the template DNA.
20. The method of claim 11, where the amount of DNA produced is a  
10,000-fold increase over the amount of the template DNA.
21. The method of claim 11, where the amount of DNA produced is a  
25 100,000-fold increase over the amount of the template DNA.



22. The method of claim 11, where the amount of DNA produced is a 1,000,000-fold increase over the amount of the template DNA.
23. The method of claim 11, where the amount of DNA produced is a 10,000,000-fold increase over the amount of the template DNA.
- 5 24. A method of amplifying a template DNA molecule comprising:  
incubating said template DNA molecule in an *in vitro* reaction mixture comprising a wild-type T7 DNA polymerase and a T7 DNA polymerase modified to have reduced 3' to 5' exonuclease activity, a 63-kDa form of a gene 4 protein from bacteriophage T7 and a single-stranded binding protein from *Escherichia coli* at a  
10 constant temperature to produce amplified product.
25. The method of claim 24, wherein said reaction mixture does not require exogenously-added oligonucleotide primers.
26. The method of claims 1, 11, or 24, wherein said amplification of said template DNA is exponential.
- 15 27. The method of claims 1, 11, or 24, wherein said reaction mixture further comprises a single-stranded DNA binding protein of bacteriophage T7.
28. The method of claims 1, 11 or 24, wherein said reaction mixture further comprises a nucleoside diphosphokinase.
29. The method of claim 28, wherein said nucleoside diphosphokinase is  
20 from *Escherichia coli*.
30. The method of claims 1, 11 or 24, wherein said reaction mixture further comprises an inorganic pyrophosphatase.
31. The method of claims 1, 11 or 24, wherein said reaction mixture further comprises an ATP regeneration system.

32. The method of claim 31, wherein said ATP regeneration system consists of phosphocreatine and creatine kinase.

33. The method of claims 1, 11 or 24, wherein said reaction mixture further comprises a 5' to 3' exonuclease.

5 34. The method of claim 33, wherein said exonuclease is T7 gene 6 exonuclease.

35. The method of claims 1, 11 or 24, wherein said reaction mixture further comprises a DNA ligase.

36. The method of claim 35, wherein said ligase is T7 DNA ligase.

10 37. The method of claims 1, 11 or 24, wherein said template DNA molecule has a length of at least 1,000 bp.

38. The method of claims 1, 11 or 24, wherein said template DNA molecule has a length of at least 1,500 bp.

15 39. The method of claims 1, 11 or 24, where in said template DNA molecule has a length of at least 2,000 bp.

40. The method of claims 1, 11 or 24, wherein said template DNA molecule has a length of at least 3,000 bp.

41. The method of claims 1, 11 or 24, wherein said template DNA molecule is amplified at least about 10-fold.

20 42. The method of claims 1, 11 or 24, wherein said template DNA molecule is amplified at least about 100-fold.

43. The method of claims 1, 11 or 24, wherein said template DNA molecule is amplified at least about 1000-fold.

44. The method of claims 1, 11 or 24, wherein said constant temperature is less than 60° C.
45. The method of claims 1, 11 or 24, wherein said constant temperature of less than 45° C.
- 5 46. The method of claims 1, 11 or 24, wherein said amplification requires the presence of ATP and CTP.
47. The method of claims 1, 11 or 24, wherein said reaction mixture further comprises 0.5 to 5% DMSO.
48. The method of claims 1, 11 or 24, wherein said reaction mixture  
10 contains 0.5-5% dextran.
49. The method of claims 1, 11 or 24, wherein said template DNA molecule is from lysed *E. coli* cells.
50. The method of claim 49, wherein said DNA is a plasmid.
51. The method of claim 49, wherein said DNA is a bacterial artificial  
15 chromosome (BAC).
52. The method of claim 1, further comprising the step of treating said polymerase and at least one accessory protein with ultraviolet light prior to incubating with said template DNA molecule.
53. The method of claim 11, further comprising the step of treating said  
20 polymerases, said helicase, and said primase with ultraviolet light prior to incubating with said template DNA molecule.
54. The method of claim 24, further comprising the step of treating said polymerases, said gene 4 protein and said single-stranded binding protein with ultraviolet light prior to incubating with said template DNA molecule.

55. The method of claims 52, 53, or 54, wherein said ultraviolet light is a dose from 10 to 1000  $\mu\text{W}/\text{cm}^2$  for from 15 sec to 5 min.

56. The method of claim 1, 11 or 24, wherein said template DNA is less than 100 ng of a plasmid and said reaction mixture is 10 to 200  $\mu\text{l}$ .

5 57. A method for preparing a DNA molecule for use in a DNA sequencing reaction comprising the steps of:

incubating said DNA molecule in an *in vitro* reaction mixture comprising a DNA polymerase, a helicase, and a primase at a constant temperature so as to produce amplified product; and

10 providing said amplified product for use in a DNA sequencing reaction.

58. The method of claim 57, wherein said reaction mixture further comprises a single-stranded binding protein.

15 59. The method of claim 57, wherein said reaction mixture does not require exogenously-added oligonucleotide primers.

60. The method of claim 57, wherein production of said amplified product is exponential.

61. The method of claim 57, wherein the amount of said amplified product is at least 10-fold greater than said DNA molecule put into said reaction mixture.

20 62. A method for preparing a DNA molecule for use in a DNA sequencing reaction comprising the steps of:

incubating said DNA molecule in an *in vitro* reaction mixture comprising a T7 polymerase and a T7 gene 4 protein at a constant temperature so as to produce amplified product; and

25 providing said amplified product for use in a DNA sequencing reaction.

63. The method of claim 62, wherein said polymerase is a mixture of wild-type T7 DNA polymerase and a T7 DNA polymerase modified to have reduced 3' to 5' exonuclease activity.

64. The method of claim 62, wherein said T7 gene 4 protein is 63-kDa form of the protein.

65. The method of claim 62, wherein said reaction mixture further comprises a single-stranded binding protein from *Escherichia coli*.

66. The method of claim 62, wherein the amount of said amplified product is amplified is at least 10-fold greater than said DNA molecule put into said reaction mixture.

67. The method of claims 57 or 62, wherein said production of said amplified product is exponential.

68. The method of claim 62, wherein said reaction does not require exogenously-added oligonucleotide primers

69. The method of claims 57 or 62, wherein said the DNA sequencing reaction uses fluorescent primers.

70. The method of claims 57 or 62, wherein said DNA sequencing reaction uses fluorescent dideoxynucleotides.

71. The method of claims 57 or 62, wherein dNTPs present in said reaction mixture are removed after amplification by treatment with phosphatase.

72. The method of claim 71, wherein said phosphatase is from arctic shrimp.

73. The method of claims 57 or 62, wherein said DNA molecule is from lysed *E. coli* cells.

74. The method of claim 73, wherein said DNA is a plasmid.
75. The method of claim 73, wherein said DNA is a bacterial artificial chromosome (BAC).
76. The method of claim 73, wherein said DNA is from a human patient.
- 5 77. The method of claims 57 or 62, wherein said reaction mixture further comprises DMSO at a concentration of 0.5 to 8%.
78. The method of claims 57 or 62, wherein said reaction mixture further comprises dextran at a concentration of between 0.5 and 20%.
79. A kit for amplification, comprising:  
10 a DNA polymerase;  
a primase;  
a helicase; and  
a single-stranded binding protein.
80. The kit of claim 79, wherein said polymerase is T7 DNA polymerase.
- 15 81. The kit of claim 79, wherein said primase and helicase are a gene 4 protein from bacteriophage T7.
82. The kit of claim 79, wherein said single-stranded binding protein is from *Escherichia coli*.
83. The kit of claim 80, wherein said T7 polymerase is a mixture of a wild-  
20 type T7 DNA polymerase and a T7 DNA polymerase modified to have reduced 3' to 5' exonuclease activity.
84. The kit of claim 81, wherein said gene 4 protein is the 63-kDa form of the protein.
85. The kit of claim 84, further comprising a nucleoside diphosphokinase.

86. The kit of claim 85, wherein said nucleoside diphosphokinase is from *Escherichia coli*.
87. The kit of claim 79, further comprising inorganic pyrophosphatase.
88. The kit of claim 79, further comprises an ATP regeneration system.
- 5 89. The kit of claim 88, wherein said ATP regeneration system consists of phosphocreatine and creatine kinase.
90. The kit of claim 79, further comprising a T7 gene 6 exonuclease.
91. The kit of claim 79, further comprising a T7 DNA ligase.
92. The kit of claim 79, further comprising a single-stranded DNA binding  
10 protein of bacteriophage T7.
93. Method for overproducing the gene 4 protein of bacteriophage T7 comprising:  
coproducing the gene 4 protein in *E. coli* with an agent that increases its level of production.
- 15 94. The method of claim 93, wherein said agent is a carboxy terminal fragment of gene 4 protein.
95. The method of claim 94, wherein said carboxy terminal fragment of gene 4 comprises the carboxy terminus of the protein after residue 221 (glutamine).
96. The method of claim 95, wherein said carboxy terminal fragment of  
20 gene 4 comprises the carboxy terminus of the protein between residues 260 (tryptophan) and residue 280 (serine).
97. Method of determining the amount of contaminating DNA in a sample comprising the steps of:

incubating said sample potentially containing contaminating DNA with a reaction mixture comprising a DNA polymerase such that if said contaminating DNA is present said DNA molecule is amplified to produce amplified product;

5 incubating at least one pre-determined amount of control DNA with said reaction mixture comprising a DNA polymerase such that said control DNA molecule is amplified to produce amplified control product; and

comparing the amount of amplified product in said sample with the amount of amplified control product as an indication of the amount of contaminating DNA in said sample.

10 98. The method of claim 97, wherein said DNA polymerase is from bacteriophage T7.

99. The method of claim 97, wherein said reaction mixture is at constant temperature.

15 100. The method of claim 97, wherein said reaction mixture further comprises a helicase, a primase and a single-stranded binding protein.

101. The method of claim 97, wherein said control DNA is amplified at least 10-fold.

102. The method of claim 97, wherein said reaction does not require exogenously-added oligonucleotide primers.

20 103. Method of claim 97, wherein amplification of said control DNA is exponential.

104. Method for amplifying DNA directly from cells, comprising the steps of:

25 lysing said cells to form a lysate; and  
incubating said lysate in a reaction mixture comprising a DNA polymerase, a helicase, and a primase at a constant temperature, such that DNA present in said lysate is amplified to produce amplified product, wherein production



of amplified DNA product does not require exogenously-added oligonucleotide primers.

105. The method of claim 104, wherein said cells are bacterial cells.

106. The method of claim 104, wherein said cells are human cells.

5 107. The method of claim 104, wherein said reaction mixture further comprises a single-stranded binding protein.

108. Method for amplifying DNA directly from cells comprising the steps of:

lysing said cells to form a lysate; and

10 incubating said lysate in a reaction mixture comprising a DNA polymerase such that DNA present in said lysate is amplified to produce amplified product, wherein production of amplified product does not require exogenously-added oligonucleotide primers, said amplification is exponential and wherein the amount of amplified product is at least ten-fold greater than the amount of DNA present in said  
15 lysate put into said reaction mixture.

109. The method of claim 108, wherein said cells are bacterial cells.

110. The method of claim 108, wherein said cells are human cells.

111. The method of claim 49, wherein said DNA is from bacteriophage.

20 112. A method of amplifying a plasmid template DNA molecule, comprising incubating said template DNA molecule in a reaction mixture comprising a DNA polymerase to produce amplified product, wherein production of amplified product does not require exogenously-added oligonucleotide primers, is exponential, and the amount of amplified product is at least 10-fold greater than the amount of template DNA put into the reaction mixture.

113. A method for preparing a DNA molecule for use in a DNA sequencing reaction comprising the steps of:

incubating said DNA molecule in a reaction mixture comprising a DNA polymerase to produce amplified product, wherein production of amplified product does not require exogenously-added oligonucleotide primers, is exponential, the amount of amplified product is at least 10-fold greater than the amount of DNA molecule put into the reaction mixture; and

providing said amplified product for use in a DNA sequencing reaction.

10 114. The method of claim 108, wherein said amount of said amplified product

is at least 100-fold greater than the amount of DNA present in said lysate added to said reaction mixture.

15 115. The method of claim 108, wherein said amount of amplified product is at

least 1000-fold greater than the amount of DNA present in said lysate added to said reaction mixture.

116. The method of claim 112, wherein said amount of amplified product is at

20 least 100-fold greater than the amount of template DNA added to said reaction mixture.

117. The method of claim 112, wherein said amount of amplified product is at

25 least 1000-fold greater than the amount of template DNA present added to said reaction mixture.

118. The method of claim 113, wherein said amount of amplified product is  
at

least 100-fold greater than the amount of DNA added to said reaction mixture.

119. The method of claim 113, wherein said amount of amplified product is  
5 at

least 1000-fold greater than the amount of DNA added to said reaction mixture.

120. The method of claim 96, wherein said carboxy terminal fragment of  
gene 4 initiates at residue 271 (arginine).

121. A method for sequencing a nucleic acid sequence, wherein the DNA  
10 molecule to be sequenced is prepared in a constant temperature amplification reaction  
comprising the steps of:

incubating a single DNA molecule to be sequenced in a reaction  
mixture comprising a DNA polymerase to produce amplified product, wherein  
production of amplified product is done isothermally, is exponential, the amount of  
15 amplified product is at least 10-fold greater than the amount of DNA molecule put  
into the reaction mixture.

122. A method for sequencing wherein the DNA molecule to be sequenced  
is prepared in a reaction comprising the steps of:

incubating a single DNA molecule to be sequenced in a reaction  
20 mixture comprising a DNA polymerase to produce amplified product, wherein  
production of amplified product does not require exogenously-added oligonucleotide  
primers, is exponential, the amount of amplified product is at least 10-fold greater  
than the amount of DNA molecule put into the reaction mixture.

123. The method of claim 121, wherein said method of DNA preparing a  
25 DNA molecule for sequencing is performed isothermally.

Figure 1.

	10	20	30	40	50	60	70	80	
ATGGACAATT	CGCACGATTC	CGATAGTGTA	TTTCTTTACC	ACATTCCTTG	TGACAACCTGT	GGGAGTAGTG	ATGGGAACCTC	80	
GCTGTTCTCT	GACGGACACA	CGTTCTGCTA	CGTATGCGAG	AAGTGGACTG	CTGGTAATGA	AGACACTAAA	GAGAGGGCTT	160	
CAAAACGGAA	ACCTTCcGGt	GGaAAgCCcg	gGACTTACAA	CGTGTGGAAC	TTCGGGGAAT	CCAATGGACG	CTACTCCGCG	240	
TTAACTGCGA	GAGGAATCTC	CAAGGAAACC	TGTCAGAAGG	CTGGCTACTG	GATTGCCAAA	GTAGACGGTG	TGATGTACCA	320	
AGTGGCTGAC	TATCGGGACC	AGAACGGCAA	CATTGTGAGT	CAGAAGGTTT	GAGATAAAGA	TAAGAACCTT	AAGACCCTG	400	
GTAGTCACAA	GAGTGACGCT	CTGTTCGGGA	AGCACTTGTG	GAATGGTGGT	AAGAAGATTG	TCGTTACAGA	AGGTGAAATC	480	
GACATGCTTA	CCGTGATGGA	ACTTCAAGAC	TGTAAGTATC	CTGTAGTGTG	GTTGGGTCCAC	GGTGCCTCTG	CCGCTAAGAA	560	
GACATGCGCT	GCCAACTACG	AATACTTTGA	CCAGTTCGAA	CAGATTATCT	TAATGTTCTG	TATGGACGAA	GCAGGGCGCA	640	
AAGCAGTCGA	AGAGGCTGCA	CAGGTTCTAC	CTGCTGGTAA	GGTACGAGTG	GCAGTTCCTC	CGTGTAAAGG	TGCAAACGAG	720	
TGTCACCTAA	ATGGTCACGA	CCGTGAAATC	ATGGAGCAAG	TGTGGAATGC	TGGTCCTTGG	ATTCCTGATG	GTGTGGTATC	800	
GGCTCTTTTCG	TTACGTGAAC	GAATCCGTGA	GCACCTATCG	TCCGAGGAAT	CAGTAGGTTT	ACTTTTCAGT	GGCTGCACTG	880	
GTATCAACGA	TAAGACCTTA	GGTGCCCGTG	GTGGTGAAGT	CATTATGGTC	ACTTCCGGTT	CCGGTATGGG	TAAGTCAACG	960	
TTCGTCCGTC	AACAAGCTCT	ACAATGGGGC	ACAGCGATGG	GCAAGAAGGT	AGGCTTAGCG	ATGCTTGAGG	AGTCCGTTGA	1040	
GGAGACCGCT	GAGGACCTTA	TAGGTCTACA	CAACCGTGTG	CGACTGAGAC	AATCCGACTC	ACTAAACAGA	GAGATTATTG	1120	
AGAACGGTAA	GTTTCGACCA	TGGTTCGATG	AACGTTCGG	CAACGATACG	TTCCATCTAT	ATGACTCATT	CGCCGAGGCT	1200	
GAGACGGATA	GACTGCTCGC	TAAGCTGGCC	TACATGCGCT	CAGGCTTGGG	CTGTGACGTA	ATCATTCTAG	ACCACATCTC	1280	
AATCGTCGTA	TCCGCTTCTG	GTGAATCCGA	TGAGCGTAAAG	ATGATTGACA	ACCTGATGAC	CAAGCTCAA	GGGTTCCGTA	1360	
AGTCAACTGG	GGTGGTGCTG	GTCGTAATTT	GTCACCTTAA	GAACCCAGAC	AAAGGTAAAG	CACATGAGGA	AGGTCGCCCC	1440	
GTTTCTATTA	CTGACCTACG	TGGTTCCTGGC	GCACTACGCC	AACATCTGTA	TACTATTATT	GCCCTTGAGC	GTAATCAGCA	1520	
AGGCGATATG	CCTAACCTTG	TCCTCGTTCCG	TATTCTCAAG	TGCCGCTTTA	CTGGTGATAC	TGGTATCGCT	GGCTACATGG	1600	
AATACAACAA	GGAAACCGGA	TGGCTTGAAC	CATCAAGTTA	CTCAGGGGAA	GAAGACTCAC	ACTCAGAGTC	AACAGACTGG	1680	
TCCAACGACA	CTGACTTCTG	ACAGGATTCT	TGATGACTTT	CCAGACGACT	ACGAGAAGTT	TCCGCTGGAGA	GTCCCATCT	1760	
AATACGACTC	ACTAAAGGAG	ACACACCATG	TTCAAACCTGA	TTAAGAAGTT	AGGCCAACTG	CTGGTTCGTA	TGTACAACGT	1840	
GGAAGCCAAG	CGACTGAACG	ATGAGGCTCG	TAAAGAGGCC	ACACAGTCAC	GCGCTCTGGC	GATTCGCTCC	AAAACCTGGT	1920	
TTGCGCTTAC	CCCAACCAAC	AGGGGATTTG	CTGCTTTCCA	TTGAGCCTGT	TTCTCTGCCG	GACGTTCCGG	GCGGCGTGT	2000	
TGTGCATCCA	TCTGGATTCT	CCTGTCAGTT	AGCTTTGGTG	GTGTGTGGCA	GTTGTAGTCC	TGAACGAAA	CCCCCGCGA	2080	
TTGGCACATT	GGCAGCTAAT	CEGGAATCGC	ACTTACGGCC	AATGCTTCGT	TTCGTATCAC	ACACCCCAA	GCCTTCTGCT	2160	
TTGAATGCTG	CCCTTCTTCA	GGGCTTAATT	TTAAGAGCC	TCACCTTCAT	GGTGGTCAGT	GCGTCTGCT	GATGTGCTCA	2240	
GTATCACCGC	CAGTGGTATT	TATGTCAACA	CCGCCAGAGA	TAATTTATCA	CCGCAGATGG	TTATCTGTAT	GTTTTTTATA	2320	
TGAATTTATT	TTTTGCAGGG	GGGCATTGTT	TGGTAGGTGA	GAGATCCGGC	TGCTAACAAA	GCCCGAAGG	AAGCTGAGTT	2400	
GGCTGCTGCC	ACCGCTGAGC	AATAACTAGC	ATAACCCCTT	GGGGCCTCTA	AACGGGTCTT	GAGGGGTTTT	TTGCTGAAAG	2480	
GAGGAACTAT	ATCCGGATAT	CCCACAAGAG	GCCCCGCAGT	ACCGGCATAA	CCAAGCCTAT	GCCTACAGCA	TCCAGGGTGA	2560	
CGGTGCCGAG	GATGACGATG	AGCGCATTGT	TAGATTTTCT	ACACGGTGCC	TGACTGCGTT	AGCAATTTAA	CTGTGATAAA	2640	
CTACCGCATT	AAAGCTTGGC	GCCGCACTCG	ACGAACCCCTT	CGGATCTCGA	TCCCGCGAAA	TTAATACGAC	TCACTATAGG	2720	
GAGACCACAA	CGGTTTCCCT	CTAGAAATAA	TTTTGTTTAA	CTTTAAGAAG	GAGATATACA	TATGCGTGAA	CGAATCCGTG	2800	
AGCACCTATC	GTCCGAGGAA	TCAGTAGGTT	TACTTTTTCAG	TGGCTGCACT	GGTATCAACG	ATAAGACCTT	AGGTGCCCGT	2880	
GGTGGTGAAG	TCATTATGGT	CACTTCCGGT	TCCGGTATGG	GTAAGTCAAC	GTTCTCCGT	CAACAAGCTC	TACAATGGGG	2960	
CACAGCGATG	GGCAAGAAGG	TAGGCTTAGC	GATGCTTGAG	GAGTCCGTTG	AGGAGACCGC	TGAGGACCTT	ATAGGTCTAC	3040	
ACAACCGTGT	CCGACTGAGA	CAATCCGACT	CACTAAAGAG	AGAGATTATT	GAGAACGGTA	AGTTCGACCA	ATGGTTCGAT	3120	
GAACGTGTCG	GCAACGATAC	GTTCCATCTA	TA'GACTCAT	TCGCCGAGGC	TGAGACGGAT	AGACTGCTCG	CTAAGCTGGC	3200	
CTACATGCGC	TCAGGCTTGG	GCTGTGACGT	AATCATTCTA	GACCACATCT	CAATCGTCGT	ATCCGCTTCT	GGTGAATCCG	3280	
ATGAGCGTAA	GATGATTGAC	AACCTGATGA	CCAAGCTCAA	AGGGTTCGCT	AAGTCAACTG	GGGTGGTGCT	GGTCGTAATT	3360	
TGTCACCTTA	AGAACCCAGA	CAAAGGTAAA	GCACATGAGG	AAGGTGCCCC	CGTTTCTATT	ACTGACCTAC	GTGGTTCCTG	3440	
CGCACTACGC	CAACTATCTG	ATACTATTAT	TGCCCTTGAG	CGTAATCAGC	AAGGCGATAT	GCCTAACCTT	GTCCTCGTTC	3520	

GTATTCTCAA GTGCCGCTTT ACTGGTGATA CTGGTATCGC TGGCTACATG GAATACAACA AGGAAACCGG ATGGCTTGAA 3600  
 CCATCAAGTT ACTCAGGGGA AGAAGAGTCA CACTCAGAGT CAACAGACTG GTCCAACGAC ACTGACTTCT GAGGATCCAC 3680  
 TAGTAACGGC CGCCAGTGTG CTGGAATTCT GCAGATATCC ATCACACTGG CGGCCGCTCG AGCACCACCA CCACCACCAC 3760  
 TGAGATCCGG CTGCTAACAA AGCCCGAAAG GAAGCTGAGT TGGCTGCTGC CACCGCTGAG CAATAACTAG CATAACCCCT 3840  
 TGGGGCCTCT AAACGGGTCT TGAGGGGTTT TTTGCTGAAA GGAGGAACTA TATCCGATT GCGGAATGGG ACGCGCCCTG 3920  
 TAGCGGCGCA TTAAGCGCGG CGGGTGTGGT GGTTACGCGC AGCGTGACCG CTACACTTGC CAGCGCCCTA GCGCCCGCTC 4000  
 CTTTCGCTTT CTCCCTTCC TTTCTCGCCA CGTTCGCCGG CTTTCCCGT CAAGCTCTAA ATCGGGGGCT CCCTTTAGGG 4080  
 TTCCGATTTA GTGCTTTACG GCACCTCGAC CCCAAAAAC TTGATTAGGG TGATGGTTCA CGTAGTGGC CATCGCCCTG 4160  
 ATAGACGGTT TTTCCGCTT TGACGTTGGA GTCCACGTTT TTAATAGTG GACTCTTGT CCAAAGTGA ACAACTCA 4240  
 ACCCTATCTC GGTCTATTCT TTTGATTTAT AAGGGATTTT GCCGATTCG GCCTATTGGT TAAAAATGA GCTGATTTAA 4320  
 CAAAAATTA ACGCGAATTT TAACAAAATA TTAACGTTTA CAATTCAGG TGGCACTTTT CGGGGAAATG TGCGCGGAAC 4400  
 CCCTATTTGT TTATTTTCT AAATACATTC AAATATGTAT CCGCTCATGA ATTAATTCTT AGAAAACTC ATCGAGCATC 4480  
 AAATGAACT GCAATTTATT CATATCAGGA TTATCAATAC CATATTTTGG AAAAAGCCGT TTCTGTAATG AAGGAGAAAA 4560  
 CTCACCGAGG CAGTTCCATA GGATGGCAAG ATCCTGGTAT CCGTCTGCGA TTCCGACTCG TCCAACATCA ATACAACCTA 4640  
 TTAATTTCCC CTCGTCAAAA ATAAGGTTAT CAAGTGAGAA ATCACCATGA GTGACGACTG AATCCGGTGA GAATGGCAA 4720  
 AGTTTATGCA TTTCTTTCCA GACTTGTTCA ACAGGCCAGC CATTACGCTC GTCATCAAAA TCACTCGCAT CAACCAAACC 4800  
 GTTATTCATT CGTGATTGCG CCTGAGCGAG ACGAAATACG CGATCGCTGT TAAAAGGACA ATTACAACA GGAATCGAAT 4880  
 GCACCAGCGG CAGGAACACT GCCAGCGCAT CAACAATATT TTCACCTGAA TCAGGATATT CTTCTAATAC CTGGAATGCT 4960  
 GTTTTCCCGG GGATCGCAGT GGTGAGTAAC CATGCATCAT CAGGAGTACG GATAAATGC TTGATGGTCG GAAGAGGCAT 5040  
 AAATCCGCTC AGCCAGTTTA GTCTGACCAT CTCATCTGTA ACATCATTGG CAACGCTACC TTTGCCATGT TTCAGAAACA 5120  
 ACTCTGGCGC ATCGGGCTTC CCATACAATC GATAGATTGT CGCACCTGAT TGCCCGACAT TATCGCGAGC CCATTTATAC 5200  
 CCATA'AAAA' CAGCATCCAT G'ITGGAA'IT' AA'CGCGGCC 'TAGAGCAAGA CG'IT'CCCGT TGAATATGGC TCATAACACC 5280  
 CTTTCTATTA CTCTTTATCT ACCAGACAG TTTTATTGTT CATGACCAA ATCCCTAAC CTCACTTTTC GTTCCACTGA 5360  
 GCGTCAGACC CCGTAGAAAA GATCAAAGGA TCTTCTTGAG ATCCTTTTTT TCTGCGGTA ATCTGCTGCT TGCAACAAA 5440  
 AAAACCACCG CTACCAGCGG TGGTTTGTTT GCCGGATCAA GAGCTACCAA CTCTTTTTCC GAAGGTAAT GGCTTCAGCA 5520  
 GAGCGCAGAT ACCAAACT GTCCCTCTAG TGTAGCCGTA GTTAGGCCAC CACTTCAAGA ACTCTGTAGC ACCGCCTACA 5600  
 TAGCTCGCTC TGCTAATCCT GTTACCAGTG GCTGCTGCCA GTGGCGATAA GTCGTGTCTT ACCGGGTTGG ACTCAAGACG 5680  
 ATAGTTACCG GATAAGCCGC AGCGGTCCGG CTGAAACGGG GGTTCGTGCA CACAGCCCAG CTTGGAGCGA ACGACCTACA 5760  
 CCGAAGTGA ATACCTACAG CGTGAGCTAT GAGAAAGCGC CACGCTTCCC GAAGGGAGAA AGGCGGACAG GTATCCGGTA 5840  
 AGCGGCAGGG TCGGAACAGG AGAGCGCAG AGGGAGCTTC CAGGGGAAA CGCCTGGTAT CTTTATAGTC CTGTCGGTT 5920  
 TCGCCACCTC TGACTTGAGC GTCGATTTT GTGATGCTCG TCAGGGGGG GAGCCTATG GAAAACGCC AGCAACGCGG 6000  
 CCTTTTTACG GTTCTTGCC TTTTGCTGGC CTTTGTCTCA CATGTTCTTT CCTGCGTTAT CCCCTGATTC TGTGGATAAC 6080  
 CGTATTACCG CCTTTGAGTG AGCTGATACC GCTCGCCGCA GCCGAACGAC CGAGCGCAGC GAGTCAGTGA GCGAGGAAGC 6160  
 GGAAGAGCGC CTGATGCGGT ATTTTCTCCT TACGCATCTG TCGGGTATT CACACCGCAT ATATGGTGCA CTCTCAGTAC 6240  
 AATCTGCTCT GATGCCGCAT AGTTAAGCCA GTATACACTC CGCTATCGCT ACGTGACTGG GTCATGGCTG CGCCCGACA 6320  
 CCCGCCAACA CCCGCTGAGC CGCCCTGACG GGCTTGTCTG CTCCCGCAT CCGCTTACAG ACAAGCTGTG ACCGTCTCCG 6400  
 GGAGCTGCAT GTGTCAGAGG TTTTACCCT CATCACCGAA ACGCGCGAG CAGCTGCGGT AAAGCTCATC AGCGTGGTCG 6480  
 TGAAGCGATT CACAGATGTC TGCTGTTC TCCCGTCCA GCTCGTTGAG TTTCTCCAGA AGCGTTAATG TCTGGCTTCT 6560  
 GATAAAGCGG GCCATGTTAA GGGCGGTTT TTCTGTGTTG GTCACTGATG CCTCCGTGTA AGGGGATTT CTGTTTATGG 6640  
 GGGTAATGAT ACCGATGAAA CGAGAGAGGA TGCTACAGAT ACGGGTACT GATGATGAAC ATGCCCAGTT ACTGGAACGP 6720  
 TGTGAGGGTA ACAAAGTGG GGTATGGATG CGCGGGACC AGAGAAAAAT CACTCAGGGT CAATGCCAGC GCTTCGTTAA 6800  
 TACAGATGTA GGTGTTCCAC AGGGTAGCCA GCAGCATCCT GCGATGCAGA TCCGGAACAT AA'GGT'GAG GCGCTGACT 6880  
 TCCCGCTTTC CAGACTTTAC GAAACACGGA AACCAGAGAC CATTCATGTT GTTGCTCAGG TCGCAGACGT TTTGCAGCAG 6960  
 CAGTCGCTTC ACGTTCGCTC GCGTATCGGT GATTCATTCT GCTAACCAGT AAGGCAACCC CGCCAGCCTA GCCGGTCTCT 7040  
 CAACGACACC AGCAGGATCA TGCGCACCCG TGGGGCCGCC ATGCCGGCGA TAATGGCCTG CTTCTCGCCG AAACGTTTGG 7120  
 TGGCGGGACC AGTGACGAAG GCTTGAGCGA GGGCGTGCAA GATTCCGAAT ACCGCAAGCG ACAGGCCGAT CATCGTCGCG 7200  
 CTCCAGCGAA AGCGGTCTTC GCCGAAAATG ACCCAGAGCG CTGCCGGCAC CTGTCTACG AGTTGCATGA TAAAGAAGAC 7280  
 AGTCATAAGT GCGGCGACGA TAGTCATGCC CCGCGCCAC CGGAAGGAGC TGAAGGGTT GAAGGCTCTC AAGGGCATCG 7360  
 GTCGAGATCC CCGTGCCTAA TGAGTGAGCT AACTTACATT AATTGCGTTC CGCTCACTGC CCGCTTTCCA GTCGGGAAAC 7440  
 CTGTGCTGCC AGCTGCATTA ATGAATCGGC CAACGCGCGG GGAGAGGCGG TTTGCGTATT GGGCGCCAGG GTGGTTTTTC 7520  
 TTTTACCAG TGAGACGGGC AACAGCTGAT TGCCCTTAC CGCTGGCCC TGAGAGAGTT GCAGCAAGCG GTCCACGCTG 7600  
 GTTTGCCCA GCAGGCGAAA ATCCTGTTT ATGGTGGTTA ACGCGGGAT ATAACATGAG CTGTCTTCGG TATCGTCTGTA 7680  
 TCCCACTACC GAGATATCCG CACCAACGCG CAGCCCGGAC TCGGTAATGG CGCGCATTC GCCCAGCGCC ATCTGATCGT 7760  
 TGGCAACCAG CATCGCAGTG GGAACGATGC CCTCATTCAG CATTTGCATG GTTTGTTGAA AACCGACAT GGCCTCCAG 7840

TCGCCTTCCC GTTCCGCTAT CGGCTGAATT TGATTGCGAG TGAGATATTT ATGCCAGCCA GCCAGACGCA GACGCGCCGA 7920  
GACAGAACTT AATGGGCCCC CTAACAGCGC GATTTGCTGG TGACCCAATG CGACCAGATG CTCCACGCCC AGTCGCGTAC 8000  
CGTCTTCATG GGAGAAAATA ATACTGTTGA TGGGTGTCTG GTCAGAGACA TCAAGAAATA ACGCCGGAAC ATTAGTGCAG 8090  
GCAGCTTCCA CAGCAATGGC ATCCTGGTCA TCCAGCGGAT AGTTAATGAT CAGCCCACTG ACGCGTTGCG CGAGAAGATT 8160  
GTGCACCGCC GCTTTACAGG CTTCGACGCC GCTTCGTTCT ACCATCGACA CCACCACGCT GGCACCCAGT TGATCGGCGC 8240  
GAGATTTAAT CGCCGCGACA ATTTGCGACG GCGCGTGCAG GGCCAGACTG GAGGTGGCAA CGCCAATCAG CAACGACTGT 8320  
TTGCCCCCA GTTGTTGTGC CACGCGGTTG GGAATGTAAT TCAGCTCCGC CATCGCCGCT TCCACTTTTT CCCGCGTTT 8400  
CGCAGAAACG TGGCTGGCCT GGTCACCAC GCGGGAAACG GTCTGATAAG AGACACCGGC ATACTCTGCG ACATCGTATA 8480  
ACGTTACTGG TTTACATTC ACCACCCTGA ATTGACTCTC TTCCGGGCGC TATCATGCCA TACCGCGAAA GGTTTTGCGC 8560  
CATTGATGG TGTCCGGGAT CTCGACGCTC TCCCTTATGC GACTCCTGCA TTAGGAAGCA GCCCAGTAGT AGGTTGAGGC 8640  
CGTTGAGCAC CGCCGCCGCA AGGAATGGTG CATGCAAGGA GATGGGCGCC AACAGTCCCC CGGCCACGGG GCCTGCCACC 8720  
ATACCCACGC CGAAACAAGC GTCATGAGC CCGAAGTGGC GAGCCCGATC TTCCCCATCG GTGATGTCCG CGATATAGGC 8800  
GCCAGCAACC GCACCTGTGG CGCCGGTGAT GCCGGCCACG ATGCGTCCGG CGTAGAGGAT CGAGATCTCG ATCCC CGCAA 8890  
ATTAATACGA CTCACTATAG GGGAAATGTTG AGCGGATAAC AATCCCCCTC TAGAAATAAT TTTGTTAAC TTTAAGAAGG 8960  
AGATATACAT 8970

WO 00/41524

PCT/US00/00580

1

## SEQUENCE LISTING

<110> TABOR, STANLEY  
RICHARDSON, CHARLES

<120> ISOTHERMAL AMPLIFICATION OF DNA

<130> 249/107 WO

<140> To Be Assigned

<141> Filed Herewith

<150> 60/115,498

<151> 1999-01-11

<160> 6

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 8970

<212> DNA

<213> Bacteriophage

<400> 1

atggacaatt	cgcacgattc	cgatagtgta	tttctttacc	acattccttg	tgacaactgt	60
gggagtagtg	atgggaactc	gctgttctct	gacggacaca	cgttctgcta	cgtatgcgag	120
aagtggactg	ctggtaatga	agacactaaa	gagagggctt	caaaacggaa	accctccggt	180
ggaaagcccg	ggacttacia	cgtgtggaac	ttcgggggat	ccaatggacg	ctactccgcg	240
ttaactgcga	gaggaatctc	caaggaaacc	tgtcagaagg	ctggctactg	gattgccaaa	300
gtagacggtg	tgatgtacca	agtggctgac	tatcgggacc	agaacggcaa	cattgtgagt	360
cagaaggttc	gagataaaga	taagaacttt	aagaccactg	gtagtcacaa	gagtgacgct	420
ctgttcggga	agcacttgtg	gaatgggtgg	aagaagattg	tcggtacaga	aggtgaaatc	480
gacatgctta	ccgtgatgga	acttcaagac	tgtaagtatc	ctgtagtgtc	gttgggtcac	540
ggtgcctctg	ccgctaagaa	gacatgcgct	gccaactacg	aatactttga	ccagttcgaa	600
cagattatct	taatgttcga	tatggacgaa	gcagggcgca	aagcagtcga	agaggctgca	660
caggttctac	ctgctggtaa	ggtacgagtg	gcagttcttc	cgtgtaagga	tgcaaacgag	720
tgtcacctaa	atggtcacga	ccgtgaaatc	atggagcaag	tgtggaatgc	tggtccttgg	780
attcctgatg	gtgtggtatc	ggctctttcg	ttacgtgaac	gaatccgtga	gcacctatcg	840
tccgaggaat	cagtaggttt	acttttcagt	ggctgcactg	gtatcaacga	taagacctta	900
ggtgcccgtg	gtggtgaagt	cattatggtc	acttccgggt	ccggtatggg	taagtcaacg	960
ttcgtccgtc	aacaagctct	acaatggggc	acagcgatgg	gcaagaaggt	aggcttagcg	1020
atgcttgagg	agtccggtga	ggagaccgct	gaggacctta	taggtctaca	caaccgtgtc	1080
cgactgagac	aatccgactc	actaaagaga	gagattattg	agaacggtaa	gttcgaccaa	1140
tggttcgatg	aactgttcgg	caacgatacg	ttccatctat	atgactcatt	cgccgaggct	1200
gagacggata	gactgctcgc	taagctggcc	tacatgcgct	caggcttggg	ctgtgacgta	1260

WO 00/41524

PCT/US00/00580

2

atcattctag	accacatctc	aatcgtcgtg	tccgcttctg	gtgaatccga	tgagcgtgtaag	1320
atgattgaca	acctgatgac	caagctcaaa	gggttcgcta	agtcaactgg	gggtgggtgctg	1380
gtcgttaatt	gtcaccttaa	gaaccagac	aaaggtaaag	cacatgagga	aggctcgcccc	1440
gtttctatta	ctgacctacg	tggttctggc	gcactacgcc	aactatctga	tactattatt	1500
gcccttgagc	gtaatcagca	aggcgatatg	cctaaccttg	tcctcgttcg	tattctcaag	1560
tgccgcttta	ctgggtgatac	tggtatcgct	ggctacatgg	aatacaacaa	ggaaaccgga	1620
tggcttgaac	catcaagtta	ctcaggggaa	gaagagtcac	actcagagtc	aacagactgg	1680
tccaacgaca	ctgacttctg	acaggattct	tgatgacttt	ccagacgact	acgagaagtt	1740
tcgctggaga	gtcccattct	aatacagctc	actaaaggag	acacaccatg	ttcaaactga	1800
ttaagaagtt	aggccaactg	ctggttcgtg	tgtacaacgt	ggaagccaag	cgactgaacg	1860
atgaggctcg	taaagaggcc	acacagtcac	gcgctctggc	gattcgctcc	aaaactgggt	1920
ttgcgcttac	cccaaccaac	aggggatttg	ctgctttcca	ttgagcctgt	ttctctgcgc	1980
gacgttcgcg	gcggcgtggt	tgtgcatcca	tctggattct	cctgtcagtt	agctttgggtg	2040
gtgtgtggca	gttgtagtcc	tgaacgaaaa	ccccccgca	ttggcacatt	ggcagctaat	2100
ccggaatcgc	acttacggcc	aatgcttcgt	ttcgtatcac	acaccccaa	gccttctgct	2160
ttgaatgctg	cccttcttca	gggcttaatt	tttaagagcg	tcaccttcat	gggtggctcagt	2220
gcgtcctgct	gatgtgctca	gtatcacccg	cagtgggtatt	tatgtcaaca	ccgccagaga	2280
taatttatca	ccgcagatgg	ttatctgtat	gttttttata	tgaatttatt	ttttgcaggg	2340
gggcattggt	tggtaggtga	gagatccggc	tgctaacaaa	gcccgaagg	aagctgagtt	2400
ggctgctgcc	accgctgagc	aataactagc	ataaccctt	ggggcctcta	aacgggtctt	2460
gaggggtttt	ttgctgaaag	gaggaactat	atccggatat	cccgcgaag	gcccggcagt	2520
accggcataa	ccaagcctat	gcctacagca	tccaggggtg	cggtgccgag	gatgacgatg	2580
agcgcattgt	tagatttcat	acacgggtgcc	tgactgcggt	agcaatttaa	ctgtgataaa	2640
ctaccgcatt	aaagcttgcg	gccgcactcg	acgaaccctt	cggatctcga	tcccgcgaaa	2700
ttaatacagc	tcactatagg	gagaccacaa	cggtttccct	ctagaaataa	ttttgtttaa	2760
ctttaagaag	gagatataca	tatgcgtgaa	cgaatccgtg	agcacctatc	gtccgaggaa	2820
tcagtaggtt	tacttttcag	tggctgcact	ggtatcaacg	ataagacctt	agggtgccgt	2880
gggtgggtgaag	tcattatggt	cacttccggt	tccggtatgg	gtaagtcaac	gttcgctcgt	2940
caacaagctc	tacaatgggg	cacagcgatg	ggcaagaagg	taggcttagc	gatgcttgag	3000
gagtccgctg	aggagaccgc	tgaggacctt	ataggtctac	acaaccgtgt	ccgactgaga	3060
caatccgact	cactaaagag	agagattatt	gagaaccggt	agttcgacca	atggttcgat	3120
gaactgttcg	gcaacgatac	gttccatcta	tatgactcat	tcgccgaggc	tgagacggat	3180
agactgctcg	ctaagctggc	ctacatgcgc	tcaggcttgg	gctgtgacgt	aatcattcta	3240
gaccacatct	caatcgtcgt	atccgcttct	ggtgaatccg	atgagcgtaa	gatgattgac	3300
aacctgatga	ccaagctcaa	agggttcgtc	aagtcaactg	gggtgggtgct	ggctcgtaatt	3360
tgtcacctta	agaaccagca	caaaggtaaa	gcacatgagg	aaggctgccc	cgtttctatt	3420
actgacctac	gtggttctgg	cgactacgc	caactatctg	atactattat	tgcccttgag	3480
cgtaatcagc	aaggcgatat	gcctaacctt	gtcctcgttc	gtattctcaa	gtgccgcttt	3540
actgggtgata	ctgggtatcg	tggctacatg	gaatacaaca	aggaaaccgg	atggcttgaa	3600
ccatcaagtt	actcagggga	agaagagtca	cactcagagt	caacagactg	gtccaacgac	3660
actgacttct	gaggatccac	tagtaacggc	cgccagtgtg	ctggaattct	gcagatatcc	3720
atcacactgg	cggccgctcg	agcaccacca	ccaccaccac	tgagatccgg	ctgctaacaa	3780
agcccgaag	gaagctgagt	tggctgctgc	caccgctgag	caataactag	cataaccctt	3840
tggggcctct	aaacgggtct	tgaggggttt	tttgctgaaa	ggaggaacta	tatccggatt	3900
ggcgaatggg	acgcgcctcg	tagcggcgca	ttaaagcgcg	cgggtgtggt	ggttacgcgc	3960
agcgtgaccg	ctacacttgc	cagcgcccta	gcgcccgcct	ctttcgtttt	cttcccttcc	4020
tttctcgcca	cgttcgccgg	ctttccccgt	caagctctaa	atcgggggct	ccctttaggg	4080
ttccgattta	gtgctttacg	gcacctcgac	cccaaaaaac	ttgattaggg	tgatggttca	4140
cgtagtgggc	catcgccctg	atagacgggt	tttcgcccct	tgacgttgga	gtccacgctt	4200
tttaatagtg	gactcttggt	ccaaactgga	acaacactca	accctatctc	ggctctattct	4260
tttgatttat	aagggtttt	gocgatttct	gcctattggt	taaaaaatga	gctgatttaa	4320
caaaaattta	acgcgaattt	taacaaaata	ttaacgttta	caatttcagg	tggcactttt	4380
cggggaaatg	tgccgcggaac	ccctatttgt	ttatttttct	aaatacatc	aaatatgtat	4440
ccgctcatga	attaattctt	agaaaaactc	atcgagcatc	aatgaaact	gcaatttatt	4500
catatcagga	ttatcaatac	catatttttg	aaaaagccgt	ttctgtaatg	aaggagaaaa	4560
ctcaccgagg	cagttccata	ggatggcaag	atcctgggat	cggtctgcga	ttccgactcg	4620
tccaacatca	atacaaccta	ttaatttccc	ctcgtcaaaa	ataaggttat	caagtgagaa	4680



atcaccatga	gtgacgactg	aatccggtga	gaatggcaaa	agtttatgca	tttctttcca	4740
gacttgttca	acaggccagc	cattacgctc	gtcatcaaaa	tcactcgcac	caaccaaacc	4800
gttattcatt	cgtgattgcg	cctgagcgag	acgaaatacg	cgatcgctgt	taaaaggaca	4860
attacaaaca	ggaatcgaat	gcaaccggcg	caggaacact	gccagcgcat	caacaatatt	4920
ttcacctgaa	tcaggatatt	cttctaatac	ctggaatgct	gttttcccgg	ggatcgcagt	4980
ggtgagtaac	catgcatcat	caggagtacg	gataaaatgc	ttgatggctg	gaagaggcat	5040
aaattccgtc	agccagttta	gtctgacat	ctcatctgta	acatcattgg	caacgctacc	5100
tttgccatgt	ttcagaaaca	actctggcgc	atcgggcttc	ccatacaatc	gatagattgt	5160
cgcacctgat	tgcccagacat	tatcgcgagc	ccatthatac	ccatataaat	cagcatccat	5220
gttggaattt	aatcgcggcc	tagagcaaga	cgtttcccgt	tgaatatggc	tcataacacc	5280
ccttgattta	ctgtttatgt	aagcagacag	ttttattggt	catgaccaa	atcccttaac	5340
gtgagttttc	gttccactga	gcgtcagacc	ccgtagaaaa	gatcaaagga	tcttcttgag	5400
atcctttttt	tctgcgcgta	atctgctgct	tgcaaaaaaa	aaaaccaccg	ctaccagcgg	5460
tggtttggtt	gccggatcaa	gagctaccaa	ctctttttcc	gaaggtaact	ggcttcagca	5520
gagcgcagat	accaaatact	gtccttctag	tgtagccgta	gttaggccac	cacttcaaga	5580
actctgtagc	accgectaca	tacctcgcct	tgctaactct	gttaccagtg	gctgctgcca	5640
gtggcgataa	gtcgtgtctt	accgggttgg	actcaagacg	atagttaccg	gataaggcgc	5700
agcggtcggg	ctgaacgggg	ggttcgtgca	cacagcccag	cttgagcga	acgacctaca	5760
ccgaactgag	atacctacag	cgtgagctat	gagaaagcgc	cacgcttccc	gaagggagaa	5820
aggcggacag	gtatccggta	agcggcaggg	tcggaacagg	agagcgcacg	agggagcttc	5880
cagggggaaa	cgcttggtat	ctttatagtc	ctgtcgggtt	tcgccacctc	tgacttgagc	5940
gtcgattttt	gtgatgctcg	tcaggggggc	ggagcctatg	gaaaaacgcc	agcaacgcgg	6000
ccttttttacg	gttcctggcc	ttttgctggc	cttttgctca	catgttcttt	cctgcgttat	6060
ccctgatctc	tgtggataac	cgtattaccg	cctttgagtg	agctgatacc	gctcgcgcga	6120
gccgaacgac	cgagcgcagc	gagtcagtga	gcgaggaagc	ggaagagcgc	ctgatgcggt	6180
atthttctct	tacgcactctg	tgcggtatth	cacaccgcat	atatggtgca	ctctcagtac	6240
aatctgctct	gatgcgcgat	agttaagcca	gtatacactc	cgctatcgct	acgtgactgg	6300
gtcatggctg	cgccccgaca	cccgccaaaca	cccgctgacg	cgccctgacg	ggcttgctctg	6360
ctcccggcat	ccgcttacag	acaagctgtg	accgtctccg	ggagctgcat	gtgtcagagg	6420
ttttcaccgt	catcaccgaa	acgcgcgagg	cagctgcggt	aaagctcatc	agcgtggctg	6480
tgaagcgatt	cacagatgtc	tgcttggtca	tccgcgtcca	gctcgttgag	tttctccaga	6540
agcgttaatg	tctggcttct	gataaagcgg	gccatgttaa	ggcgggtttt	ttcctgtttg	6600
gtcactgatg	cctccgtgta	agggggatth	ctgttcatgg	gggtaatgat	accgatgaaa	6660
cgagagagga	tgctcacgat	acgggttact	gatgatgaac	atgcccggtt	actggaacgt	6720
tgtgagggta	aacaactggc	ggtatggatg	cggcgggacc	agagaaaaat	cactcagggt	6780
caatgccagc	gcttcgtaa	tacagatgta	ggtgttccac	agggtagcca	gcagcatcct	6840
gcgatgcaga	tccggaacat	aatgggtgcag	ggcgcctgact	tccgcgtttc	cagactttac	6900
gaaacacgga	aaccgaagac	cattcatgth	gthgctcagg	tcgcagacgt	tttgcagcag	6960
cagtcgcttc	acgttcgctc	gcgtatcggt	gattcattct	gctaaccagt	aaggcaaccc	7020
cgccagccta	gccgggtcct	caacgacagg	agcacgatca	tgccgacccg	tggggccgcc	7080
atgccggcga	taatggcctg	cttctcgcgc	aaacgthtg	tggcgggacc	agtgacgaag	7140
gcttgagcga	ggcgtgcaa	gattccgaat	accgcaagcg	acaggccgat	catcgtcgcg	7200
ctccagcga	agcggctcct	gccgaaaatg	accagagcgc	ctgccggcac	ctgtcctacg	7260
agttgcatga	taaagaagac	agtcataagt	gcggcgacga	tagtcatgcc	ccgcgcccac	7320
cggaaggagc	tgactgggtt	gaaggctctc	aaggcctcgc	gtcagatccc	cggtgcctaa	7380
tgagtgagct	aacttacatt	aattgctgtg	cgctcactgc	ccgctttcca	gtcgggaaac	7440
ctgtcgtgcc	agctgcatta	atgaatcggc	caacgcgcgc	ggagaggcgg	tttgcgtatt	7500
gggcgccagg	gtggthtttc	ttttcaccag	tgagacgggc	aacagctgat	tgcccttcac	7560
cgcttgcccc	tgagagagtt	gcagcaagcg	gtccacgctg	gthtgcccc	gcaggcga	7620
atcctgthtg	atggthgth	acggcgggat	ataacatgag	ctgtcttcgg	tatcgtcgt	7680
tcccactacc	gagatatccg	caccaacgcg	cagcccggac	tcggthaatg	cgcgcatthc	7740
gccagcgcgc	atctgatcgt	tggcaaccag	catcgcagtg	ggaacgatgc	cctcattcag	7800
catttgcatg	gthtgthgaa	aaccggacat	ggcactccag	tcgccttccc	gttcgcgtat	7860
cggctgaatt	tgattgcgag	tgagatath	atgccagcca	gccagacgca	gacgcgccga	7920
gacagaactt	aatgggcccg	ctaacagcgc	gatttgctgg	tgacccaatg	cgaccagatg	7980
ctccacgccc	agtcgcgtac	cgtcttcatg	ggagaaaata	atactgthga	tgggtgctg	8040
gtcagagaca	tcaagaaata	acgccggaac	attagthcag	gcagcttcca	cagcaatggc	8100

WO 00/41524

PCT/US00/00580

4

atcctgggtca	tccagcggat	agttaatgat	cagcccactg	acgcgttgcg	cgagaagatt	8160
gtgcaccgcc	gctttacagg	cttcgacgcc	gcttcggttct	accatcgaca	ccaccacgct	8220
ggcaccagcgt	tgatcggcgc	gagatttaat	cgccgcgaca	atttgcgacg	gcgcggtgcag	8280
ggccagactg	gaggtggcaa	cgccaatcag	caacgactgt	ttgcccgcca	gttggttgtc	8340
cacgcggttg	ggaatgtaat	tcagctccgc	catcgccgct	tccacttttt	cccgcgtttt	8400
cgcagaaacg	tggtggcct	ggttcaccac	gcgggaaacg	gtctgataag	agacaccggc	8460
atactctgcg	acatcgtata	acgttactgg	tttcacattc	accaccctga	attgactctc	8520
ttccgggcg	tatcatgcca	taccgcgaaa	ggttttgcg	cattcgatgg	tgtccgggat	8580
ctcgacgctc	tcccttatgc	gactcctgca	ttaggaagca	gccagtagt	aggttgaggc	8640
cgttgagcac	cgccgccgca	aggaatggtg	catgcaagga	gatggcgccc	aacagtcccc	8700
cggccacggg	gctgcccacc	ataccacgc	cgaaacaagc	gctcatgagc	ccgaagtggc	8760
gagcccgatc	ttccccatcg	gtgatgtcgg	cgatataggc	gccagcaacc	gcacctgtgg	8820
cgcggtgat	gccggccacg	atgcgtccgg	cgtagaggat	cgagatctcg	atcccgcgaa	8880
attaatacga	ctcactatag	gggaattgtg	agcggataac	aattcccctc	tagaaataat	8940
ttgtttaac	tttaagaagg	agatatacat				8970

<210> 2  
 <211> 19  
 <212> DNA  
 <213> Bacteriophage

<220>  
 <223> BAC-1

<400> 2

cgcggtacac cgacgtcaa 19

<210> 3  
 <211> 19  
 <212> DNA  
 <213> Bacteriophage

<220>  
 <223> BAC-2

<400> 3

cgcggtacac cgacttaat 19

<210> 4  
 <211> 10  
 <212> DNA  
 <213> Bacteriophage

<220>  
 <223> BAC-3

<400> 4

gtcgggtgtac 10

WO 00/41524

PCT/US00/00580

5

<210> 5  
<211> 23  
<212> DNA  
<213> Bacteriophage

<220>  
<223> T7

<400> 5

taatagact cactataggg cga

23

<210> 6  
<211> 24  
<212> DNA  
<213> Bacteriophage

<220>  
<223> SP6

<400> 6

catagattt aggtgacact atag

24

10	20	30	40	50	60	70	80	
ATGGACAATT	CGCACGATTC	CGATAGTGTA	TTTCTTTACC	ACATTCCTTG	TGACAACCTGT	GGGAGTAGTG	ATGGGAACCTC	80
GCTGTTCTCT	GACGGACACA	CGTTCCTGCTA	CGTATGCGAG	AAGTGGACTG	CTGGTAATGA	AGACACTAAA	GAGAGGGCTT	160
CAAAACGGAA	ACCCTCcgGe	GGaAAgCCcg	gGACTTACAA	CGTGTGGAAC	TTCGGGGGAAT	CCAATGGACG	CTACTCCGCG	240
TTAACTGCGA	GAGGAATCTC	CAAGGAAACC	TGTCAGAAGG	CTGGCTACTG	GATTGCCAAA	GTAGACGGTG	TGATGTACCA	320
AGTGGCTGAC	TATCGGGACC	AGAACGGCAA	CATTGTGAGT	CAGAAGGTTT	GAGATAAAGA	TAAGAACCTT	AAGACCACTG	400
GTAGTCACAA	GAGTGACGCT	CTGTTCCGGGA	AGCACTTGTG	GAATGGTGGT	AAGAAGATTG	TCGTTACAGA	AGGTGAAATC	480
GACATGCTTA	CCGTGATGGA	ACTTCAAGAC	TGTAAGTATC	CTGTAGTGTC	GTTGGGTCAE	GGTGCCTCTG	CCGCTAAGAA	560
GACATGCGCT	GCCAACCTACG	AATACTTTGA	CCAGTTCGAA	CAGATTATCT	TAATGTTTGA	TATGGACGAA	GCAGGGCGCA	640
AAGCAGTCGA	AGAGGCTGCA	CAGGTTCTAC	CTGCTGGTAA	GGTACGAGTG	GCAGTTCCTC	CGTSTAAGGA	TGCAAACGAG	720
TGTCACCTAA	ATGGTCACGA	CCGTGAAATC	ATGGAGCAAG	TGTGGAATGC	TGGTCCCTGG	ATTCCTGATG	GTGTGGTATC	800
GGCTCTTTTCG	TTACGTGAAC	GAATCCGTGA	GCACCTATCG	TCCGAGGAAT	CAGTAGGTTT	ACTTTTCAGT	GGCTGCACTG	880
GTATCAACGA	TAAGACCTTA	GGTGCCCGTG	GTGGTGAAGT	CATTATGGTC	ACTTCCGGTT	CCGGTATGGG	TAAGTCAACG	960
TTCGTCCGTC	AACAAGCTCT	ACAATGGGGC	ACAGCCGATGG	GCAAGAAGGT	AGGCTTAGCG	ATGCTTGAGG	AGTCCGTTGA	1040
GGAGACCGCT	GAGGACCTTA	TAGGTCTACA	CAACCCGTGTC	CGACTGAGAC	AATCCGACTC	ACTAAACAGA	GAGATTATTG	1120
AGAACGGTAA	GTTCCGACCAA	TGGTTCGATG	AACTGTTCCG	CAACGATACG	TTCATCTAT	ATGACTCATT	CGCCGAGGCT	1200
GAGACGGATA	GACTGCTCGC	TAAGCTGGCC	TACATGCGCT	CAGGCTTGGG	CTGTGACGTA	ATCATTCTAG	ACCACATCTC	1280
AATCGTCGTA	TCCGCTTCTG	GTGAATCCGA	TGAGCGTAAG	ATGATTGACA	ACCTGATGAC	CAAGCTCAA	GGGTTCGCTA	1360
AGTCAACTGG	GGTGGTGCCTG	GTCGTAATTT	GTCACCTTAA	GAACCCAGAC	AAAGGTAAAG	CACATGAGGA	AGGTCCGCCC	1440
GTTTCTATTA	CTGACCTACG	TGGTTCCTGG	GCACTACGCC	AACTATCTGA	TACTATTATT	GCCCTTGAGC	GTAATCAGCA	1520
AGGCGATATG	CCTAACCTTG	TCCTCGTTTCG	TATTCTCAAG	TGCCGCTTTA	CTGGTGATAC	TGGTATCGCT	GGCTACATGG	1600
AATACAACAA	GGAAACCGGA	TGGCTTGAAC	CATCAAGTTA	CTCAGGGGAA	GAAGAGTCAC	ACTCAGAGTC	AACAGACTGG	1680
TCCAACGACA	CTGACTTCTG	ACAGGATTCT	TGATGACTTT	CCAGACGACT	ACGAGAAGTT	TGGCTGGAGA	GTCCCATTCT	1760
AATACGACTC	ACTAAAGGAG	ACACACCATG	TTCAAACTGA	TTAAGAAGTT	AGGCCAACTG	CTGGTTCGTA	TGTACAACGT	1840
GGAAGCCAAG	CGACTGAACG	ATGAGGCTCG	TAAAGAGGCC	ACACAGTCAC	GCGCTCTGGC	GATTCGCTCC	AAACTGGT	1920
TTGCGCTTAC	CCCAACCAAC	AGGGGATTTG	CTGCTTTCCA	TTGAGCCTGT	TTCTCTGGCG	GACGTTCCGG	GCGGCGTGTT	2000
TGTGCATCCA	TCTGGATTCT	CCTGTCAAGT	AGCTTTGGTG	GTGTGTGGCA	GTTGTAGTCC	TGAACGAAA	CCCCCGCGA	2080
TTGGCACATT	GGCAGCTAAT	CCGGAACTCG	ACTTACCGCC	AATGCTTCGT	TTCGTATCAC	ACACCCCAA	GCCTTCTGCT	2160
TTGAATGCTG	CCCTTCTTCA	GGGCTTAATT	TTAAGAGCG	TCACCTTCAT	GGTGGTCAGT	GCGTCCCTGCT	GATGTGCTCA	2240
GTATEACCGC	CAGTGGTATT	TATGTCAACA	CCGCCAGAGA	TAATTTATCA	CCGCAGATGG	TTATCTGTAT	GTTTTTTATA	2320
TGAATTTATT	TTTTGCAGGG	GGGCATTGTT	TGGTAGGTGA	GAGATCCGGC	TGCTAACAAA	GCCCGAAAGG	AAGCTGAGTT	2400
GGCTGCTGCC	ACCGCTGAGC	AATAACTAGC	ATAACCCCTT	GGGGCCTCTA	AACGGGTCTT	GAGGGGTTTT	TTGCTGAAAG	2480
GAGGAACTAT	ATCCGGATAT	CCCGCAAGAG	GCCCGGCAGT	ACCGGCATAA	CCAAGCCTAT	GCCTACAGCA	TCCAGGGTGA	2560
CGGTGCCGAG	GATGACGATG	AGCGCATTGT	TAGATTTTAT	ACACGGTGCC	TGACTGCGTT	AGCAATTTAA	CTGTGATAAA	2640
CTACCGCATT	AAAGCTTGCG	GCCGCACTCG	ACGAACCCTT	CGGATCTCGA	TCCCGCGAAA	TTAATACGAC	TCACTATAGG	2720
GAGACCACAA	CGGTTTCCCT	CTAGAAATAA	TTTTGTTTAA	CTTAAGAAG	GAGATATACA	TATCGTGAA	CGAATCCGTC	2800
AGCACCTATC	GTCCGAGGAA	TCAGTAGGTT	TACTTTTTCAG	TGGCTGCACT	GGTATCAACG	ATAAGACCTT	AGGTGCCCGT	2880
GGTGGTGAAG	TCATTATGGT	CACTTCCGGT	TCCGGTATGG	GTAAGTCAAC	GTTCTGTCCT	CAACAAGCTC	TACAATGGGG	2960
CACAGCGATG	GGCAAGAAGG	TAGGCTTAGC	GATGCTTGAG	GAGTCCGTTG	AGGAGACCGC	TGAGGACCTT	ATAGGTCTAC	3040
ACAACCGTGT	CCGACTGAGA	CAATCCGACT	CACTAAAGAG	AGAGATTATT	GAGAACGGTA	AGTTCCGACCA	ATGGTTCGAT	3120
GAACTGTTTCG	GCAACGATAC	GTTCATCTA	TATGACTCAT	TCGCCGAGGC	TGAGACGGAT	AGACTGCTCG	CTAAGCTGGC	3200
CTACATGCGC	TCAGGCTTGG	GCTGTGACGT	AATCATTCTA	GACCACATCT	CAATCGTCTG	ATCCGCTTCT	GGTGAATCCG	3280
ATGAGCGTAA	GATGATTGAC	AACCTGATGA	CCAAGCTCAA	AGGGTTCGCT	AAGTCAACTG	GGGTGGTGCT	GGTCGTAATT	3360
TGTCACCTTA	AGAACCCAGA	CAAAGGTA	GCACATGAGG	AAGGTCCGCC	CGTTTCTATT	ACTGACCTAC	GTGGTTCCTGG	3440
CGCACTACGC	CAACTATCTG	ATACTATTAT	TGECCTTGAG	CGTAATCAGC	AAGGCCGATAT	GCCTAACCTT	GTCCCTCGTTC	3520