



(86) Date de dépôt PCT/PCT Filing Date: 2004/05/28
 (87) Date publication PCT/PCT Publication Date: 2004/12/23
 (45) Date de délivrance/Issue Date: 2010/10/12
 (85) Entrée phase nationale/National Entry: 2005/11/21
 (86) N° demande PCT/PCT Application No.: US 2004/017037
 (87) N° publication PCT/PCT Publication No.: 2004/110366
 (30) Priorité/Priority: 2003/05/29 (US60/474,362)

(51) Cl.Int./Int.Cl. *A61K 8/37* (2006.01),
A61K 8/18 (2006.01), *A61K 8/34* (2006.01),
A61K 8/35 (2006.01), *A61Q 17/04* (2006.01)
 (72) Inventeurs/Inventors:
 SANOGUEIRA, JAMES P., US;
 DUEVA-KOGANOV, OLGA V., US
 (73) Propriétaire/Owner:
 SUN PHARMACEUTICALS CORPORATION, US
 (74) Agent: GOWLING LAFLEUR HENDERSON LLP

(54) Titre : COMPOSITIONS DE PROTECTION SOLAIRE
 (54) Title: SUNSCREEN COMPOSITION

(57) **Abrégé/Abstract:**

There is provided a composition comprising one or more photoactive compounds and one or more optimization agents. Surprisingly, the composition requires a small amount of optimization agent to efficiently optimize the polarity, critical wavelength, SPF, PFA, Star Rating, photostability, or any combinations thereof, of the composition. Subsequently, an efficient sunscreen composition is achieved.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
23 December 2004 (23.12.2004)

PCT

(10) International Publication Number
WO 2004/110366 A3

(51) International Patent Classification⁷: **A61K 7/00**,
7/42, 7/44

(21) International Application Number:
PCT/US2004/017037

(22) International Filing Date: 28 May 2004 (28.05.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/474,362 29 May 2003 (29.05.2003) US

(71) Applicant: **SUN PHARMACEUTICALS CORPORA-
TION** [US/US]; 50 N. Dupont Highway, Dover, DE 19903
(US).

(72) Inventors: **DUEVA, Olga, V.**; 30 Lake Street, Apt. 2D,
White Plains, NY 10603 (US). **SANOGUEIRA, James,
P.**; 4 Kimmissy Court, Suffern, NY 10901 (US).

(74) Agent: **RUGGIERO, Charles, N., J.**; Ohlandt, Greeley
& Ruggiero & Perle, L.L.P., 10th Floor, One Landmark
Square, Stamford, CT 06901 (US).

(81) Designated States (*unless otherwise indicated, for every
kind of national protection available*): AE, AG, AL, AM,

AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,
TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every
kind of regional protection available*): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI,
SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

(88) Date of publication of the international search report:
2 June 2005

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: SUNSCREEN COMPOSITION

(57) Abstract: There is provided a composition comprising one or more photoactive compounds and one or more optimization agents. Surprisingly, the composition requires a small amount of optimization agent to efficiently optimize the polarity, critical wavelength, SPF, PFA, Star Rating, photostability, or any combinations thereof, of the composition. Subsequently, an efficient sun-screen composition is achieved.

WO 2004/110366 A3

SUNSCREEN COMPOSITION

BACKGROUND OF THE INVENTION

5 1. Field of the Invention

The present invention relates generally to a photoprotective composition. More particularly, the present invention relates to a sunscreen composition having optimized polarity, critical wavelength, SPF
10 PFA, photostability, Star Rating, or any combinations thereof. The present invention also relates generally to a method of optimizing photoprotective compositions.

15 2. Description of the Prior Art

Sunscreen compositions are applied to the skin to protect the skin from the sun's ultraviolet rays that can lead to erythema, a reddening of the skin also known as sunburn. Sunlight or ultraviolet radiation in the UV-B range has a wavelength of 290nm to 320nm and is known to be the primary
20 cause of sunburn. Ultraviolet rays at a wavelength of 320nm to 400nm, known as UV-A radiation, produces tanning of the skin. However, in the process of doing so, the UV-A rays can damage or harm the skin.

Besides the immediate malady of sunburn, excessive sunlight
25 exposure can lead to skin disorders. For instance, prolonged and constant

exposure to the sun may lead to actinic keratoses and carcinomas.

Another long-term effect is premature aging of the skin. This condition is characterized by skin that is wrinkled, cracked and has lost its elasticity.

5 As stated above, sunscreens are typically formulated with the goal of inhibiting skin damage from the sun's rays. The sunscreen composition filters or blocks the harmful UV-A and UV-B rays that can damage and harm the skin. It is believed that sunscreen agents accomplish this by absorbing the UV-A and/or UV-B rays.

10

Typically, the above-described UV-B filters are combined with the above-described UV-A filters in a solution with other lipophilic or oily ingredients and solvents to form an oil phase. The solvents are used to dissolve the sunscreen actives into the oil phase. Typically, but not
15 necessarily, the oil phase is dispersed with the help of emulsifiers and stabilizers into an aqueous solution composed primarily of water, to make an emulsion, which becomes the final sunscreen composition.

One problem associated with the use of UV filters, and especially
20 those that are rapidly-degrading photoactive compounds, is that they are not photostable and will degrade rapidly and exponentially when exposed to UV radiation. The organic UV-A filters most commonly used in commercial sunscreen compositions are the dibenzoylmethane derivatives, particularly 4-(1,1-dimethylethyl)-4'-methoxydibenzoylmethane (also called
25 avobenzene, sold under the brand name PARSOL 1789). Other

dibenzoylmethane derivatives described as UV-A filters are disclosed in U.S. Pat. Nos. 4,489,057; 4,387,089 and 4,562,067. It is also well known that the above described UV-A filters, particularly the dibenzoylmethane derivatives, can suffer in rapid photochemical degradation, when used
5 alone or when combined with the above-described most commercially used UV-B filters. Thus, the efficiency of the sunscreen composition (i.e., SPF, PFA, critical wavelength, Star Rating) containing these photoactive compounds is compromised, unless the photodegradation is controlled by improving the photostability of the system -in UVA and/or
10 UVB regions.

By controlling the polarity of the solvent used in a sunscreen composition, the rate of photodecay of the photoactive compounds in the composition can be controlled. By controlling the polarity, greater
15 stability is imparted to the photoactive compounds, thus resulting in a more stable overall composition. The dielectric constant, for example, is a good indicator or measure of polarity in a composition. This is due to the fact that the dielectric constant is a measure of both inherent and inducible dipole moments.

20

In addition, polar solvents tend to decrease the energy required to excite a pi-bonding electron, and increase the energy required to excite a non-bonding electron. This phenomenon is called "state switching" and is a mechanism by which photoactive compounds absorb UV
25 radiation. By

enhancing the state switching in a photoprotective or sunscreen composition, a more efficient UV absorbing composition can result.

It is also known that the use of different solvents in sunscreen formulations may increase or decrease the effectiveness of a sunscreen chemical. The shifts (hypsochromic to the lower wavelength or bathochromic to higher wavelength) in the UV spectrum are due to the relative degrees of solvation by the solvent of the ground state and the excited state of the chemical.

10

It has been found in the prior art that as the polarity of a solvent system including a dissolved, rapidly-photodegradable compound is increased, the rate of photodecay initially decreases, but then increases again as the polarity is further increased. Thus, a photodegradable compound in solution will degrade as a second-order function of the overall polarity of the solution. Currently accepted photochemical theory provides the possibility that the mechanism by which a photodegradable compound is stabilized is the transfer of a photonicly-excited electron to a nearby molecule of the same or different species (see, e.g., N. J. Turro, *Modern Molecular Photochemistry*, Chapter 9, Benjamin/Cummings Publ. Co., Menlo Park, Calif. (1991)). Additional photochemical theory is believed to coincide with the electron transfer theory of Professor Rudolph A. Marcus of the California Institute of Technology, for which he received the 1992 Nobel Prize in Chemistry.

25

U.S. Patents No. 6,485,713 and 6,537,529 to Bonda et al., consistent with the above-described theory, discloses the use of amides, malates and bis-urethanes in a solvent system to control the polarity of the solvent system in a sunscreen composition. The use of these specific components results in an oil phase having a dielectric constant no greater than about 12. The named components are used in an oil-in-water sunscreen composition in an amount about 0.1% to about 40% by weight of the total weight of the composition, and more preferably about 3 wt.% to about 20 wt.%.

In addition to the above, U.S. Patent Application Publication No. 2004/0057916 A1 to Bonda et al. discloses polymers and compounds including a diphenylmethylene or a 9H-fluorene moiety for use in sunscreen compositions to photostabilize UVA sunscreen actives.

Critical wavelength is another important aspect in optimizing the performance of a photoprotective composition. In 1994, Diffey described the Critical Wavelength in vitro method, which is based on the absorption spectrum of a sunscreen product obtained via UV substrate spectrophotometry (Diffey BL (1994) A Method for Broad-Spectrum Classification of Sunscreens. Intl J Cosmet Sci, 16: 47-52), which is incorporated by reference herein. The absorption spectrum of a sunscreen is characterized by an index, namely critical wavelength, which is the wavelength where the integral of the spectral absorbance curve reached

90% of the integral from 290 nm to 400 nm. The critical wavelength method is used to determine the breadth of UV protection and is the recommended method for the evaluation of long wave efficacy of sunscreen products. Therefore, by optimizing the critical wavelength properties of a photoprotective composition, enhanced photoprotection may result.

Another measure of a sunscreen composition's efficiency is the Star Rating (UVA/UVB Ratio) according to the Boots Star Rating System (4-star that was recently revised to 5 star category). The Star Rating is calculated as an indicator of the UVA absorbance properties of a sunscreen product, relative to UVB as described in the Revised Guidelines to the practical measurement of UVA:UVB ratios according to Boots Star Rating System. The calculation of the UVA:UVB absorbance ratio will typically yield values from zero (equal to no UVA absorbance) up to 1.0 (UVA absorbance equal to UVB).

What is absent in the prior art is a photoprotective composition having one or more agents that differ from the prior art that are capable of optimizing at least one of the following properties: polarity, critical wavelength, SPF, PFA, Star Rating, or any combinations thereof, in the oil phase, water phase, both phases, or the final sunscreen formulation; thus resulting in a more efficient and photostable photoprotective composition.

The present invention addresses this shortcoming by providing an efficient photoprotective composition having one or more optimization agents capable of optimizing at least one of the following properties: polarity, critical wavelength, SPF, PFA, Star Rating, photostability or any combinations thereof, in the oil phase, water phase, both phases of the composition, or the final sunscreen formulation.

SUMMARY OF THE INVENTION

It is an object of the present invention to provide an efficient photoprotective composition.

It is another object of the present invention to provide such a composition that is a sunscreen composition.

It is still another object of the present invention to provide such a composition having one or more optimization agents capable of optimizing the polarity, critical wavelength, SPF, PFA, Star Rating, photostability, or any combinations thereof of the oil phase, water phase, both oil and water phases, of the composition.

It is another object of the present invention to provide such a composition where the one or more optimization agents are lipophilic, hydrophilic, or both.

It is still another object of the present invention to provide such a composition where the one or more optimization agents have a dielectric constant greater than about 10.5.

5 It is still another object of the present invention to provide such a composition where the one or more optimization agents have a dielectric constant greater than about 13.

It is a further object of the present invention to provide such a
10 composition where the one or more optimization agents are one or more alcohols, such as, for example, glycols, diols, or any derivatives thereof.

These and other objects of the present invention are achieved by a composition comprising one or more photoactive compounds and one or
15 more optimization agents. Surprisingly, the composition requires a small amount of optimization agent to efficiently optimize the polarity, critical wavelength, SPF, PFA, Star Rating, photostability, or any combinations thereof, of the composition. Subsequently, an efficient sunscreen composition is achieved.

20

BRIEF DESCRIPTION OF THE FIGURES

Figure 1 is a graph depicting the UVA photostability of compositions
25 according to the present invention and comparative examples;

Figure 2 is a graph depicting the UVB photostability of compositions according to the present invention and comparative examples;

5 Figure 3 is a graph depicting the critical wavelength of compositions according to the present invention and comparative examples;

Figure 4 is a graph depicting the UVA photostability of compositions with a synergistic combination of optimizing agent and octocrylene
10 according to the present invention;

Figure 5 is a graph depicting the critical wavelength of compositions with a synergistic combination of optimizing agent and octocrylene
according to the present invention;

15

Figure 6 is a fusion graph depicting the photostability of avobenzene in a photoprotective composition according to the present invention; and

Figure 7 is another fusion graph depicting the photostability of
20 avobenzene in a photoprotective composition according to the present invention.

DETAILED DESCRIPTION OF THE INVENTION

The present invention provides photoprotective compositions that
25 are uniquely formulated with one or more optimization agents that results in

the optimization of one or more of the following properties: polarity, critical wavelength, SPF, PFA, Star Rating (UVA/UVB Ratio), photostability, or any combinations thereof, of the composition. Photoprotective compositions according to the present invention include, but are not limited to,

5 sunscreens, cosmetics, paints, coatings, and the like.

A photoactive compound, as used herein, is a compound that responds to UV radiation photoelectrically. Examples of photoactive compounds include, but are not limited to, UV filters, pigments, or dyes.

10

By way of example, the present invention is illustrated herein by reference to a sunscreen composition for use on mammalian hair and/or skin. It is to be understood, however, that the principles set forth below apply equally to any photoprotective composition.

15 According to the present invention, a sunscreen composition is provided having one or more photoactive agents and one or more optimization agents. The resulting sunscreen composition is photostable and possesses optimized polarity, critical wavelength, SPF, PFA, Star

20 Rating, photostability, or any combinations thereof, which results in a more efficient UV radiation-absorbing composition.

The one or more photoactive agents suitable for use in the sunscreen composition of the present invention include one or more UV

25 filters. Suitable UV filters may include, but are not limited to, one or more

compounds selected from the following categories (with specific examples):

p-aminobenzoic acid, its salts and its derivatives (ethyl, isobutyl, glyceryl esters; p-dimethylaminobenzoic acid); anthranilates (o-aminobenzoates; methyl, menthyl, phenyl, benzyl, phenylethyl, linalyl, terpinyl, and cyclohexenyl esters); salicylates (octyl, amyl, phenyl, benzyl, menthyl (homosalate), glyceryl, and dipropyleneglycol esters); cinnamic acid derivatives (menthyl and benzyl esters, alpha-phenyl cinnamitrile; butyl cinnamoyl pyruvate); dihydroxycinnamic acid derivatives (umbelliferone, methylumbelliferone, methylaceto-umbelliferone); camphor derivatives (3-benzylidene, 4-methylbenzylidene, polyacrylamidomethyl benzylidene, benzalkonium methosulfate, benzylidene camphor sulfonic acid, and terephthalylidene dicamphor sulfonic acid); trihydroxycinnamic acid derivatives (esculetin, methylesculetin, daphnetin, and the glucosides, esculin and daphnin); hydrocarbons (diphenylbutadiene, stilbene); dibenzalacetone and benzalacetophenone; naphtholsulfonates (sodium salts of 2-naphthol-3,6-disulfonic and of 2-naphthol-6,8-disulfonic acids); dihydroxy-naphthoic acid and its salts; o- and p-hydroxydiphenyldisulfonates; coumarin derivatives (7-hydroxy, 7-methyl, 3-phenyl); diazoles (2-acetyl-3-bromoindazole, phenyl benzoxazole, methyl naphthoxazole, various aryl benzothiazoles); quinine salts (bisulfate, sulfate, chloride, oleate, and tannate); quinoline derivatives (8-hydroxyquinoline salts, 2-phenylquinoline); hydroxy- or methoxy-substituted benzophenones; uric and vilouric acids; tannic acid and its derivatives; hydroquinone; benzophenones (oxybenzone, sulisobenzene, dioxybenzone, benzoescorcinol, 2,2',4,4'-tetrahydroxybenzophenone, 2,2'-

dihydroxy-4,4'-dimethoxybenzophenone, octabenzene), dibenzoylmethane derivatives, avobenzene, 4-isopropylidibenzoylmethane, butylmethoxydibenzoylmethane, 4-isopropyl-dibenzoylmethane, octocrylene, drometrizole trisiloxane, and metal oxides (titanium dioxide, 5 zinc oxide).

In one embodiment of the invention a photoactive compound is selected from the group consisting of UV-A filters, UV-B filters, or any combinations thereof. In a cosmetically-acceptable sunscreen embodiment 10 for use on human skin, a photoactive compound preferably is selected from approved (if regulated), cosmetically-acceptable UV-A filters, UV-B filters, or any combinations thereof.

For example, for a product marketed in the United States, preferred 15 cosmetically-acceptable photoactive compounds and concentrations (by way of example, reported as a percentage by weight of the total cosmetic sunscreen composition) include: aminobenzoic acid (also called para-aminobenzoic acid and PABA; 15% or less), avobenzene (also called butyl methoxy dibenzoylmethane; 3% or less), cinoxate (also called 2- 20 ethoxyethyl p-methoxycinnamate; 3% or less), dioxybenzone (also called benzophenone-8; 3% or less), homosalate (15% or less), menthyl anthranilate (also called menthyl 2-aminobenzoate; 5% or less), octocrylene (also called 2-ethylhexyl-2-cyano-3,3 diphenylacrylate; 10% or less), octyl methoxycinnamate (7.5% or less), octyl salicylate (also called 2- 25 ethylhexyl salicylate; 5% or less), oxybenzone (also called benzophenone-

3; 6% or less), padimate O (also called octyl dimethyl PABA; 8% or less),
phenylbenzimidazole sulfonic acid (water soluble; 4% or less),
sulisobenzone (also called benzophenone-4; 10% or less), titanium dioxide
(25% or less), trolamine salicylate (also called triethanolamine salicylate;
5 12% or less), and zinc oxide (25% or less).

Other preferred cosmetically-acceptable photoactive compounds
and concentrations (by way of example, percent by weight of the total
cosmetic sunscreen composition) include diethanolamine
10 methoxycinnamate (10% or less), ethyl-[bis(hydroxypropyl)]aminobenzoate
(5% or less), glyceryl aminobenzoate (3% or less), 4-isopropyl
dibenzoylmethane (5% or less), 4-methylbenzylidene camphor (6% or
less), terephthalylidene dicamphor sulfonic acid (10% or less), and
sulisobenzone (also called benzophenone-4, 10% or less).

15

For a product marketed in the European Union, preferred
cosmetically-acceptable photoactive compounds and concentrations (by
way of example, reported as a percentage by weight of the total cosmetic
sunscreen composition) include: PABA (5% or less), camphor
20 benzalkonium methosulfate (6% or less), homosalate (10% or less),
benzophenone-3 (10% or less), phenylbenzimidazole sulfonic acid (8% or
less, expressed as acid), terephthalidene dicamphor sulfonic acid (10% or
less, expressed as acid), butyl methoxydibenzoylmethane (5% or less),
benzylidene camphor sulfonic acid (6% or less, expressed as acid),
25 octocrylene (10% or less, expressed as acid), polyacrylamidomethyl

benzylidene camphor (6% or less), ethylhexyl methoxycinnamate (10% or less), PEG-25 PABA (10% or less), isoamyl p-methoxycinnamate (10% or less), ethylhexyl triazone (5% or less), drometrizole trielloxane (15% or less), diethylhexyl butamido triazone (10% or less), 4-methylbenzylidene
5 camphor (4% or less), 3-benzylidene camphor (2% or less), ethylhexyl salicylate (5% or less), ethylhexyl dimethyl PABA (8% or less), benzophenone-4 (5%, expressed as acid), methylene bis-benzotriazolyl tetramethylbutylphenol (10% or less), disodium phenyl dibenzimidazole tetrasulfonate (10% or less, expressed as acid), bis-ethylhexyloxyphenol
10 methoxyphenol triazine (10% or less), methylene bisbenzotriazolyl tetramethylbutylphenol (10% or less, also called TINOSORB M), and bisethylhexyloxyphenol methoxyphenyl triazine.(10% or less, also called TINOSORB S), Mexoryl XL (also called drometrizole trisiloxane, 15% or less), Mexoryl SX (15% or less).

15

The one or more photoactive compounds are present in the composition in an amount about 1% to about 40% by weight of the total weight of the sunscreen composition. The amount of sunscreen agent in the composition can vary in the above range depending on the sun
20 protection factor (SPF) desired. Usually, the higher the SPF, the greater the total amount of sunscreen agent used in the composition. However, as demonstrated herein, the present invention provides the possibility of formulating a photoprotective composition with an increased or boosted SPF without the inclusion of additional sunscreen agent or increasing the
25 total amount of sunscreen agent in the composition.

Preferably, the one or more sunscreen agents are included at about 2 wt.% to about 35 wt.% to achieve a SPF of about 2 to about 50. More preferably, the one or more sunscreen agents are included in an amount
5 about 4 wt.% to about 30 wt.% to achieve a SPF value of about 4 to about 45.

The composition of the present invention must include one or more optimization agents. It has been unexpectedly found that the inclusion of
10 one or more optimization agents according to the present invention in a sunscreen composition results in a stable, efficient composition.

The one or more optimization agents may be present in a photoprotective composition according to the present invention from about
15 0.1 wt.% to about 40 wt.%, based on the total weight of the composition. Preferably, the one or more optimization agents are present from about 0.5 wt.% to about 15 wt.%, and more preferably from about 1 wt.% to about 11 wt.%, based on the total weight of the composition.

20 Surprisingly, it has been found that only small amounts of the one or more optimization agents, on the order of about 0.1 wt.% to about 10 wt.%, based on the total weight of the composition, are required to effect the desired optimization of polarity, critical wavelength, SPF, PFA, Star Rating (UVA/UVB Ratio), photostability, or any combinations thereof, in the
25 composition. Preferably, the one or more optimization agents, taken alone

or in combination, have a dielectric constant greater than about 10.5. More preferably, the one or more optimization agents, taken alone or in combination, have a dielectric constant greater than about 13.

5 Suitable optimization agents for use in the present invention may include, but are not limited to, diols, alcohols, glycols, polyhydric alcohols, polyhydric alcohol derivatives having one or more hydroxyl groups, or any combinations thereof. Preferably, the one or more optimization agents are one or more alcohols. More preferably, the one or more optimization
10 agents are one or more diols, glycols, or any combinations thereof. Most preferably, diols are 1,2-diols.

 Suitable glycols for use in the invention include, but are not limited to, pentylene glycol (1,2-pentanediol), neopentyl glycol (neopentanediol),
15 caprylyl glycol (1,2-octanediol), etoxydiglycol, butylene glycol monopropionate, diethylene glycol monobutyl ether, PEG-7 methyl ether, octacosanyl glycol, arachidyl glycol, benzyl glycol, cetyl glycol (1,2-hexanediol), C₁₄₋₁₈ glycol, C₁₅₋₁₈ glycol, lauryl glycol (1,2-dodecanediol), butoxydiglycol, 1,10-decanediol, ethyl hexanediol, or any combinations
20 thereof. It is believed that glycols, such as those listed above, have a dielectric constant greater than about 10.5.

 Determining the polarity of a mixture or an emulsion can be performed in various ways. For example, determining a polarity can
25 include measuring a property that is a function of polarity, such as a

dielectric constant. Measurement of a dielectric constant of a liquid can be performed by various sensors, such as immersion probes, flow-through probes, and cup-type probes, attached to various meters, such as those available from the Brookhaven Instruments Corporation of Holtsville, N.Y. (e.g., model BI-870) and the Scientifica Company of Princeton, N.J. (e.g. models 850 and 870). For consistency of comparison, preferably all measurements for a particular filter system are performed at substantially the same sample temperature, e.g., by use of a water bath. Generally, the measured dielectric constant of a substance will increase at lower temperatures and decrease at higher temperatures.

Dielectric Constants of the following glycols and their derivatives have been measured (see Table 1).

15

Table 1. Dielectric Constants

glycols/derivatives 20	Trade Name/Manufacturer	Dielectric Constant, 23 °C
ethylene Glycol	Hydrolite-5/Symrize	18.2
ethylene Glycol Monobutyl Ether	™ DB Solvent/Eastman	10.59
Hexanediol and 1,2- Octanediol (50:50 ture)	Symdiol 68/Symrize	13.07
oxydiglycol	Educol-421/MMP	13.69
ethylene Glycol Monopropionate	BG Monopropionate/ MMP	11.72
Octanediol (Caprylyl Glycol)	LexGard O/Inolex	10.61 (measured at 30 °C)*
Hexanediol		15.1
ethylene Glycol-7 Methyl Ether	Sasol	13.1

*Solid at room temperature (23 C).

It has been surprisingly found that the inclusion of one or more optimization agents in a composition according to the present invention results in a SPF boost on the order of at least 25% as compared to a composition without one or more optimization agents. Particularly, a SPF boost on the order of at least about 30% is experienced by a composition according to the present invention.

In order to evaluate the unexpected SPF boost in photoprotective compositions by the one or more optimization agents according to the present invention, by way of example, the performance of 1,2-octanediol (caprylyl glycol) and pentylene glycol *in vivo* was evaluated. The following formulations outlined in Table 2 were tested in SPF studies (static and Very Water Resistant (VWR)). The SPF was determined using the method outlined in the Food and Drug Administration (FDA) Final Monograph for sunscreen testing published in the Federal Register, Vol. 64, No. 98, May 21, 1999.

Referring to Table 2 below, the differences among the formulations are the presence or absence of 1,2- octanediol (caprylyl glycol) and/or octocrylene.

Table 2. Sunscreen Compositions

		MF2822-1	MF2822-2	MF2770-114
Chemical/INCI/USP Name	Trade Name	% w/w	% w/w	% w/w
Octinoxate	Neo Heliopan AV	7.50	7.50	7.50
Octyl Salicylate	Neo Heliopan OS	5.00	5.00	5.00
Homosalate	Escalol 567	5.00	5.00	5.00
Avobenzone	Parsol 1789	3.00	3.00	3.00
Octocrylene		NONE	NONE	2.50
Purified Water		60.03	65.03	62.53
Neopentyl Glycol Diheptanoate	Lexfeel 7	5.00	5.00	5.00
1,2-Octane diol	LexGard O	5.00	NONE	NONE
Tapioca Starch	Stabilix T	2.50	2.50	2.50
Acrylates/C12-22 Alkylmethacrylate Copolymer	Allianz OPT	2.50	2.50	2.50
Glycerin	Emery 917	2.00	2.00	2.00
Phenoxyethanol (and) Methylparaben (and) Butylparaben (and) Ethylparaben (and) Propylparaben (and) Isopropylparaben	Phenonip	1.00	1.00	1.00
Triethanolamine	TEA 99%	0.53	0.53	0.53
PEG-20 Almond Glycerides	Crovol A-40	0.35	0.35	0.35
Acrylates/C10-30 AlkylAcrylate Crosspolymer	Pemulen TR-2	0.27	0.27	0.27
Carbomer	Ultrez 10	0.15	0.15	0.15
Tocopheryl Acetate (USP)	Tocopheryl Acetate	0.05	0.05	0.05
Disodium EDTA	Dissolvine Na2S	0.07	0.07	0.07
Aloe Barbadensis Leaf Extract	Aloe Oil Extract 101	0.05	0.05	0.05
		100.00	100.00	100.00

5

The results of the SPF tests are outlined below in Tables 3 through 5.

Table 3. Results for MF2822-1

	SPF Static	SPF VWR
Average SPF (N=5)	37.06	35.00
Standard Deviation	2.20	0.00
Standard Error	0.98	0.00
t (one-tail)	2.132	2.132
A	2.10	0.00
SPF Label	34.96	35.00

10

Table 4. Results for MF2822-2

	SPF Static	SPF VWR
Average SPF (N=5)	28.38	28.38
Standard Deviation	1.65	1.65
Standard Error	0.74	0.74
t (one-tail)	2.132	2.132
A	1.57	1.57
SPF Label	26.81	26.81

Table 5. Results for MF2770-114

	SPF Static	SPF VWR
Average SPF (N=21)	37.02	36.25
Standard Deviation	2.98	2.29
Standard Error	0.65	0.50
t (one-tail)	1.725	1.725
A	1.12	0.86
SPF Label	35.00	35.00

5

The results of SPF tests demonstrate that the addition of 5 wt.% of 1,2-octanediol (caprylyl glycol) to the sunscreen formulation provided an SPF boost of more than 8 SPF units (about 30.5%) when compared to the formulation without 1,2-octanediol (caprylyl glycol). Sunscreen formulation with 5 wt.% of 1,2-octanediol (caprylyl glycol) and without octocrylene has a SPF VWR of 35.0, which is the same as the formulation with 2.5 wt.% of octocrylene and without 1,2-octanediol. Overall, the inclusion of one or more optimization agents according to the present invention, such as, for example, 1,2-octanediol (caprylyl glycol), in a sunscreen formulation results in a significant SPF boost. In addition, the inclusion of one or more

15

optimization agents in a sunscreen formulation can result in a composition having a desired SPF with less amount of sunscreen active in the composition.

5

EXAMPLE 2

The following additional example demonstrates the effectiveness of the present invention in providing a stable and efficient photoprotective composition. A comparison of a composition according to the present invention (Comp. B) having a optimizing agent (in this case pentylene glycol) to one not having an optimization agent (Comp. A) is set forth below. The test methodology used to determine SPF is the same as that described above with respect to Example 1.

15

Table 6. Sunscreen Compositions With and Without Pentylene Glycol

<u>INCI Adopted Name</u>	<u>Weight%</u>	
	<u>Comp. A</u>	<u>Comp. B</u>
Water	62.3773	60.8773
Octyl Methoxycinnamate	7.5000	7.5000
Ethylhexyl Salicylate	5.0000	5.0000
C12-15 Alkyl Benzoate	5.0000	5.0000
Benzophenone-3	4.1000	4.1000
Isopropyl Myristate	4.0000	4.0000
Cocoglycerides	3.5000	3.5000
Glycerin	2.0000	2.0000
Acrylates/C12-22 Alkylmethacrylate Copolymer	2.0000	2.0000
Butyl methoxydibenzoylmethane	1.5000	1.5000

Pentylene Glycol	NONE	1.5000
Phenoxyethanol (and) Methylparaben (and) Butylparaben (and) Ethylparaben (and) Propylparaben (and) Isobutylparaben	1.0000	1.0000
Triethanolamine	0.6500	0.6500
PEG-20 Almond Glycerides	0.3500	0.3500
Acrylates C10-30 Alkyl Acrylate Crosspolymer	0.3000	0.3000
Fragrance	0.2000	0.2000
Carbomer	0.1500	0.1500
Xanthan Gum	0.1000	0.1000
Disodium EDTA	0.0700	0.0700
Tocopheryl Acetate	0.0500	0.0500
Aloe Barbadensis Leaf Extract	0.0500	0.0500
Carthamus Tinctorius (Safflower) Seed Oil (and) Chamomile Recutita (Matricaria) Extract	0.0500	0.0500
Carthamus Tinctorius (Safflower) Seed Oil (and) Lavandula Angustifolia (Lavender) Extract	0.0500	0.0500
Yellow 5	0.0020	0.0020
Green 5	0.0007	0.0007
TOTAL:	100.0000	100.0000
Average SPF Static <i>in vivo</i>	< 25.68	31.76
Average SPF VWR <i>in vivo</i>	< 24.12	31.22

As is evident from this example, the inclusion of optimization agent according to the present invention, such as 1.5 wt.% pentylene glycol (1,2-pentenediol) in the sunscreen formulation (Comp. B), results in an increase in SPF by more than 6 units (about 29%) due to the increased polarity of the oil phase.

While, as noted above, it has been unexpectedly found that one or more optimization agents according to the present invention boost SPF, it has also been unexpectedly found that the one or more optimization agents also photostabilize UVA sunscreens, such as avobenzone and its derivatives, and UVB sunscreens. As a result, both UVA and UVB protection is optimized. This is demonstrated by way of Example 3 below.

Test compositions were prepared with octocrylene (Positive control), glycols, glycol derivatives, or isopropyl myristate (Negative control) added to the following sunscreen components: AVOBENZONE-3 g; HOMOSALATE - 15 g; OCTYL SALICYLATE - 5 g, thus imitating the oil phase of a sunscreen formulation (see Table 7):

Table 7. Compositions

#	ADDITIVE:	g	DEC, 23 C
Positive Control 1	Octocrylene	5	7.89
2	Pentylene Glycol	5	10.5
3	Ethoxydiglycol (Diethylene glycol monoethyl ether)	5	8.59
4	BG Monopropionate	5	8
5	1,2-Octanediol (Caprylyl Glycol)	5	8.1
Negative Control 6	Isopropyl Myristate	5	6.15
7	1,2-Hexanediol + 1,2-Octanediol	5	7.8
8	Diethylene Glycol Monobutyl Ether	5	7.6
9	1,2- Hexanediol	5	8
10	Octocrylene + 1,2-Octanediol	2.5 +2 .5	7.8
Negative Control 11	No additives	0	7.03
12	PEG-7 Methyl Ether	5	7.69
13	Octocrylene	2.5	7.79
14	Octocrylene and PEG-7 Methyl Ether	2.5 + 2.5	8.36
15	Octocrylene + 1,2-Octanediol + PEG-7 Methyl Ether	2.5+1.25+ 1.25	8.39

Referring to Table 7 above, the compositions were applied on Vitro-Skin (0.6 mg/cm²) and irradiated. The irradiation dose was 5 MEDs repeated 4 times (=20 MEDs total). Retained Absorbance at 370 nm (UVA photostability, Figure 1), 310 nm (UVB photostability, Figure 2) and critical wavelength (Table 8, Figure 3) were determined for each composition before and after each irradiation dose in order to determine their photostability.

For each sunscreen composition, the filter system was blended with the solvent system to form an oil phase. Next, the dielectric constant of the oil phase was measured. Dielectric constant measurements were performed with a Scientifica Model 850 dielectric constant meter.

5

The resulting sunscreen oil phases were tested for photostability by measuring absorbance on a Labsphere UV-1000S Ultraviolet Transmittance Analyzer before and after irradiation with a Solar Light Company model 16S solar simulator (equipped with a WG 320 filter to transmit radiation greater than 290 nm) in 5 MED (105 mJ/cm^{sup.2}) increments up to 20 MED cumulative dose. Output was monitored by a PMA 2105 UV-B DCS Detector (biologically weighted) and controlled by a PMA 2100 Automatic Dose Controller (Solar Light Co.).

15

A synthetic skin substrate was used for testing the sunscreen compositions (VITRO-SKIN substrate by IMS, Inc. of Milford, Conn.). To prepare the substrate, a 300 g solution of 44 g of glycerin and 256 g of deionized water was added to an IMS hydrating chamber, and a sheet of VITRO-SKIN was placed in the hydrating chamber for approx. 16 hours. Several 6.5 cm squares were cut from the hydrated VITRO-SKIN and used for absorbance measurements.

20

To prepare slides for testing, sunscreen composition is drawn or placed into a pipette. The test composition is uniformly applied to VITRO-SKIN square (0.6 mg/cm²). This application dose of the oil phase, 0.6

25

mg/cm², corresponds to the application dose of 2 mg/cm² of sunscreen formulation (typically the concentration of the oil phase in the sunscreen formulation is about 30%). The VITRO-SKIN square was then placed on a foam block, and the test material was spread by finger (covered with a latex
 5 glove or finger cot), first in a circular motion, then by a side-to-side motion during which the VITRO-SKIN is deformed by the pressure. The square was then mounted in a slide holder and allowed to dry for about 15 to 20 minutes.

10 To test photostability, a slide was positioned on the UV transmittance analyzer using registration marks, and a scan of 1 sq. cm spot on the slide was performed. The slide was then transferred to a holder placed adjacent to the solar simulator and, using a calipers, was positioned such that the beam of UV radiation exiting the solar simulator illuminated
 15 the same 1 cm spot on the slide. The following software settings were used: UV-B--290-320 nm; UV-A--320-400 nm; SPF--290-400 nm; Spectral Irradiance; Noon, July 3, Albuquerque, N.Mex.; SPF Spectral Irradiance and Erythermal Effectiveness settings as set by manufacturer. Following an exposure of 5 MED, the slide was again placed in position on the UV
 20 transmittance analyzer, and a scan of the exposed spot was performed. The procedure was repeated on the same 1 cm spot on the slide until the desired total radiation dosage was achieved.

Table 8. Critical Wavelength

Critical Wavelength, CW							
					MED		

ADITIVE:	Amount, g	#	0	5	10	15	20
Acrylene (Positive Control)	5	1	380	377	377	377	377
Pentylene Glycol	5	2	381	378	376	369	355
Ethoxydiglycol	5	3	379	378	373	369	367
BG Monopropionate	5	4	378	376	373	365	363
1,2-Octanediol	5	5	382	381	379	378	376
Propyl Myristate (Negative Control)	5	6	378	375	370	355	350
1,2-Hexanediol + 1,2-Octanediol	5	7	379	376	370	358	348
Pentylene Glycol Monobutyl Ether	5	8	379	377	373	364	357
1,2-Hexanediol	5	9	378	375	367	356	349
Acrylene + 1,2-Octanediol	2.5 + 2.5	10	381	380	380	379	379
Propyl Myristate (Negative Control)	0	11	380	377	370	364	362
PEG-7 Methyl Ether	5	12	380	377	370	361	351
Acrylene (Positive Control)	2.5	13	381	378	378	377	376
Acrylene and PEG-7 Methyl Ether	2.5 + 2.5	14	380	378	378	378	378
Acrylene+ 1,2-Octanediol+PEG-7 Methyl Ether	2.5 + 1.25 + 1.25	15	381	380	379	379	379

As is evident by the results outlined in the graphs of Figures 1 and 2, as well as Table 8 and the graph of Figure 3, it was unexpectedly found that the polarity of an oil phase alone is not just a single factor that affects the photostability of UVA sunscreens, such as avobenzone and its derivatives. Glycol or glycol derivative structure may also play a very important role in photostability. For example, sunscreen compositions containing 1,2-octanediol, BG monopropionate and 1,2-hexanediol have very similar dielectric constants (8.0-8.1). However, only 1,2 –octanediol most effectively stabilizes avobenzone. It stabilizes avobenzone by increasing the retained absorbance at 370 nm, 310 nm, and minimizing the decrease of Critical Wavelength (CW) after UV irradiation.

It was also found unexpectedly that certain glycols, for example, pentylene glycol, ethoxydiglycol, BG monopropionate and 1,2- hexanediol can improve the photostability of UVB sunscreens at 310 nm without the significant improvement of the photostability of avobenzone at 370 nm and also critical wavelength.

Mechanisms of photostabilizing UVA and UVB sunscreens by diols, glycols and their derivatives may include, but are not limited to, external stabilization by hydrogen bonding. In addition, the one or more optimization agents, such as, for example, caprylyl glycol and pentylene glycol (most preferred optimizing agents) do not have σ - π bonds and unsaturated groups in their molecular structures. In addition, most preferred optimizing agents (caprylyl glycol and pentylene glycol) are 1,2-diols.

EXAMPLE 4

Referring to Table 2 above, the PFA of MF2822-1 and MF2822-2 compositions was measured by JCIA Method (methods described in co-pending patent application Serial No. 10/779,314, filed February 13, 2004, which is incorporated in its entirety by reference herein). PFA value is an indicator of a sunscreen's protection in UVA region, which is associated with photoaging, as opposed to the UVB region, which is associated with sunburn. PFA is also related to the degree of photostabilization of UVA sunscreens, such as avobenzene and its derivatives. So a PFA boost to a sunscreen composition will directly correlate with higher UVA efficacy of a sunscreen as well as a higher degree of photostabilization of a UVA sunscreen.

It has been unexpectedly found that the inclusion of one or more optimization agents in a photoprotective composition according to the

present invention results in a PFA boost to the composition, as compared to a composition without one or more optimization agents. A PFA boost on the order on at least 10% is experienced by the compositions according to the present invention. Particularly, a PFA boost on the order of greater than about 50% is experienced, and more particularly a PFA boost of greater than about 85% is experienced by a composition according to the present invention, as compared to a composition without one or more optimization agents.

PFA values for MF2822-1 and MF 2822-2 are outlined in Table 9 below.

Table 9. Average PFA Values

MF2822-1	MF2822-2
10.34	5.51

15

The results of PFA tests have shown that the addition of 5 wt.% of 1,2-octanediol to the sunscreen formulation provided a PFA (PPD, JCIA) boost by almost 5 PFA units (about 88%) when compared to the formulation without 1,2-octanediol.

20

In addition to the above-described unexpected features of the present invention, it has also unexpectedly been found that there exists a synergistic effect on the photostability of UVA sunscreen, such as

avobenzone and its derivatives, when octocrylene and one or more optimization agents, such as 1,2-octanediol (caprylyl glycol) are both present in a sunscreen composition. As a result of this synergistic effect, a photoprotective composition according to the present invention possesses at least a 10% increase in UVA photostability, at least a 10% increase in UVB photostability and a decrease in critical wavelength, as compared to a composition without the synergistic combination according to the present invention.

The one or more optimization agents to octocrylene may be present in a photoprotective composition in a weight ratio of about 0.01 to about 100. Preferably, the synergistic combination is present in a ratio between about 0.1 to about 10, and more preferably about 0.5 to about 5.

EXAMPLE 5

Referring to Table 10 below, octocrylene and/or 1,2-octanediol were added to the following sunscreen components: AVOBENZONE-3 g; HOMOSALATE-15 g; OCTYL SALICYLATE-5 g; which are intended to resemble an oil phase of a sunscreen composition. It is noted that the base composition #'s 1, 5, 10 and 11 are described in Table 2 above.

Table 10. Compositions

ADDITIVE:	g	Table 2 Comp.
Octocrylene (Positive Control)	5	1
1,2-Octanediol	5	5

Octocrylene + 1,2-Octanediol	2.5 + 2.5	10
No additives (Negative Control)	0	11

The above compositions were applied on Vitro-Skin (0.6 mg/cm²) and irradiated. The irradiation dose was 5 MEDs repeated 4 times (=20 MEDs total). Retained Absorbance at 370 nm (Table 11 and Figure 4), 310 nm (Table 12) and critical wavelength (Table 13 and Figure 5) were determined for each composition before and after each irradiation dose in order to determine their photostability. The methodology is described above in more detail in Example 3.

10

Table 11. UVA Photostability

UVA Photostability							
370 nm							
MED							
ADDITIVE:	Amount, g	#	0	5	10	15	20
Octocrylene (Positive Control)	5	1	100	75	66	63	60
1,2-Octanediol	5	5	100	89.9	72.9	58.9	45.9
Octocrylene + 1,2-Octanediol	2.5+2.5	10	100	83	74	65	60
No additives (Negative Control)	0	11	100	64	29	16	10

15

Table 12. UVB Photostability

UVB Photostability, 310 nm							
MED							
ADDITIVE:	Amount, g	#	0	5	10	15	20
Octocrylene (Positive Control)	5	1	100	82	72.5	67	60
1,2-Octanediol	5	5	100	90.4	82.8	80.4	77.4
Octocrylene + 1,2-Octanediol	2.5+2.5	10	100	97.4	93.4	85.3	80.4
No additives (Negative Control)	0	11	100	86	78.3	72	70

Table 13. Critical Wavelength

Critical Wavelength, CW							
					MED		
ADITIVE:	Amount, g	#	0	5	10	15	20
Octocrylene (Positive Control)	5	1	380	377	377	377	377
1,2-Octanediol	5	5	382	381	379	378	378
Octocrylene + 1,2-Octanediol	2.5 + 2.5	10	381	380	380	379	379
Other additives (Negative Control)	0	11	380	377	370	364	364

As is evident from the above tables, in conjunction with graphs set forth in Figures 4 and 5, the composition with 2.5 g octocrylene and 2.5 g of 1,2 octanediol, provides better photostabilization of avobenzene than 5 g of octocrylene alone. Therefore, there is an unexpected synergistic effect on the photostabilization of avobenzene by using both octocrylene and 1,2 octanediol. Surprisingly, the UVB photostabilization is also achieved by the synergistic effect of the combination of octocrylene and 1,2 octanediol.

10

In addition, unexpectedly, the critical wavelength of the composition including the synergistic combination of octocrylene and 1,2 octanediol has been stabilized.

15

In another embodiment of the present invention, it has been surprisingly found that certain molar ratios of the one or more optimization agents according to the present invention to one or more dibenzoylmethane derivatives, such as avobenzene, results in a highly photostable composition.

20

Suitable molar ratio ranges for dibenzoylmethane derivative, according to the present invention are between about 0.016M to about 0.193M. Preferably, the molar range is between about 0.048M to about 0.096M. The dibenzoylmethane derivative is present in the composition in
5 an amount about 0.5 wt.% to about 6 wt.%.

Suitable molar ratios for the one or more optimization agents to dibenzoylmethane derivative is between about 0.5 to about 400. Preferable, the molar ratio of one or more optimization agent to
10 dibenzoylmethane derivative is about 0.5 to about 100 and more preferably about 0.5 to 10.

This embodiment of the present invention is demonstrated below in Example 6, with avobenzene as the dibenzoylmethane derivative.
15

EXAMPLE 6

In order to determine the impact of the concentration of avobenzene, from 0.04842 M to 0.0968 M (Factor 1) and different molar ratios of 1,2-
20 octanediol (caprylyl glycol) to avobenzene, from 0 to 10 (Factor 2), on the photostability of avobenzene in the different sunscreen compositions, numerous studies were conducted. Three designs of experiments that represent central composite designs (CCDs) and D.O.E. Fusion 7.2.2 software were employed.

25

Experimental Set-Up

The CCD type of design was used to determine the effect of two factors: concentration of Avobenzene and the molar ratio 1, 2-octanediol/avobenzene on the photostability of avobenzene in the different sunscreen compositions. Two responses were measured: the retained absorbance (%) of a sunscreen composition at 370 nm (UVA) and the same measurement at 310 nm (UVB) after irradiation dose of 15 MED.

10 Table 14. Ranges for the Two Experimental Factors

<i>Factor</i>	<i>Range</i>
Avobenzene Concentration	0.04842M to 0.0968M
Glycol/Avo Molar Ratio	0 to 10

15 Table 15 below presents the experimental CCD set-up used. The second column presents the experimental order in which the experiments were carried out, since order randomization is necessary

Table 15. Experimental CCD

5

		Factor 1	Factor 2
<i>Std</i>	<i>Run</i>	<i>A:Avobenzone</i> <i>M</i>	<i>B:Molar Ratio</i> <i>Glycol:Avo</i>
1	8	0.055505	1.4645
2	10	0.089715	1.4645
3	7	0.055505	8.5355
4	12	0.089715	8.5355
5	1	0.04842	5
6	13	0.0968	5
7	6	0.07261	0
8	11	0.07261	10
9	3	0.07261	5
10	9	0.07261	5
11	4	0.07261	5
12	2	0.07261	5
13	5	0.07261	5

The same order of experiments was run in the different set-ups. As

a result, two categorical factors were added: Oxybenzone (absent or present at a predetermined constant level) and Octocrylene (absent or

10 present at a predetermined constant level). Table 16 presents the three

set-ups as well as the numbers of the formulation tables and response tables.

Table 16. Full Experimental Runs

5

	Avobenzone and Glycol/Avo CCD	Oxybenzone	Octocrylene	Formulation Table	Results Table
DOE 1	Yes	Absent	Absent	Table 17	Table 18
DOE 2	Yes	Present	Absent	Table 19	Table 20
DOE 3	Yes	Present	Present	Table 21	Table 22

Response surface equalization were fitted to both responses (retained absorbance at 370 nm and retained absorbance at 310 nm, both after an irradiation dose of 15 MED). The significant factors were selected.

10 Response 1: Retained absorbance at 370 nm after irradiation at 15 MED

The response surface model indicates a high fit ($R^2 = 0.9534$). The presence of Oxybenzone in the system (model oil phase) gives the highest

15 boost in retain absorbance at 370 nm and 15 MED exposure, as

demonstrated by the graph in Figure 6. An interaction between the presence of oxybenzone and the level of the glycol/avobenzone ratio was observed. This may suggest that at a high level of glycol/avobenzone oxybenzone may interact with the glycol and affect the stability of avobenzone.

There is a significant effect of the glycol/avobenzone ratio. This effect shows a statistically significant improvement in the measured response and it is stronger for lower concentration of avobenzone.

Higher concentrations of avobenzone help a better retention. The presence of octocrylene helps the absorbance retention 370 nm, but not as significant as in the case of oxybenzone. This effect seems to be enhanced at higher level glycol/avobenzone ratios. The graph of Figure 7 demonstrates these results.

Response 2: Retain absorbance at 310 nm after irradiation at 15 MED

The response surface model indicates a moderate fit ($R^2 = 0.6668$). Oxybenzone helps retention at 310 nm where a mild jump in response especially at the lower levels of glycol/avobenzone ratio. Octocrylene also gives mild a gain in absorbency especially at higher levels of glycol/avobenzone and low levels of avobenzone. The higher concentration of avobenzone has a positive impact on the response, especially in the presence of octocrylene.

Table 17. DOE 1 Formulations

DOE 1													
RUN #:	1	2	3	4	5	6	7	8	9	10	11	12	13
MODEL OF OIL PHASE:													
Homosalate	11	11	11	11	11	11	11	11	11	11	11	11	11
Octyl Salicylate	5	5	5	5	5	5	5	5	5	5	5	5	5
Avobenzene	1.5	2.3	2.3	2.3	2.3	2.3	1.7	1.7	2.3	2.8	2.3	2.8	3
1,2-Octanediol	3.5	5.3	5.3	5.3	5.3	0	6.9	1.2	5.3	1.9	10.6	11	7
Neopentyl Glycol Diheptanoate QS to 30 g	9	6.4	6.4	6.4	6.4	11.7	5.4	11.1	6.4	9.3	1.1	0	4
	30	30	30	30	30	30	30	30	30	30	30	30	30

Table 18. UVA and UVB Photostabilization for DOE 1

5 DOE 1	Factor 1	Factor 2	Dielectric Constant	Retained Abs, %	Retained Abs, %
Run	A:Avo, M	B:Molar Ratio Octanediol:Avo	23 C	at 370 nm after 15 MED	at 310 nm after 15 MED
1	0.04842	5	6.1	20	65
2	0.07261	5	6.64	25	73
3	0.07261	5	6.64	30	76
4	0.07261	5	6.64	29	79
10 5	0.07261	5	6.64	26	77
6	0.07261	0	5.58	13	61
7	0.055505	8.5355	6.78	26	67
8	0.055505	1.4645	5.69	17	62
9	0.07261	5	6.64	27	74
10	0.089715	1.4645	6.11	17	66
11	0.07261	10	7.8	33	77
12	0.089715	8.5355	8.08	24	72
13	0.0968	5	7.24	31	76

15 Table 19. DOE 2

DOE 2	1	2	3	4	5	6	7	8	9	10	11	12	13
MODEL OF OIL PHASE:													
Cholesterol	11	11	11	11	11	11	11	11	11	11	11	11	11
ethyl Salicylate	5	5	5	5	5	5	5	5	5	5	5	5	5
Hydrobenzoin	6	6	6	6	6	6	6	6	6	6	6	6	6
Hydrobenzoin	1.5	2.3	2.3	2.3	2.3	2.3	1.7	1.7	2.3	2.8	2.3	2.8	3
1-Octanediol	3.5	5.3	5.3	5.3	5.3	0	6.9	1.2	5.3	1.9	10.6	11	7
Caprylyl Glycol													
Caprylate QS to 36 g	9	6.4	6.4	6.4	6.4	11.7	5.4	11.1	6.4	9.3	1.1	0	4
	36	36	36	36	36	36	36	36	36	36	36	36	36

Table 20. UVA and UVB Photostabilization for DOE 2

DOE 2 ₅	Factor 1	Factor 2	Dielectric Constant	Retained Abs, %	Retained Abs, %
Run	A:Avo, M	B:Molar Ratio Octanediol:Avo	23 C	at 370 after 15 MED	at 370 after 15 MED
1	0.04842	5	7.2	47	70
2	0.07261	5	7.7	62	81
3	0.07261	5	7.7	60	79
4	0.07261	5	7.7	62	80
5	0.07261	5	7.7	62	81
106	0.07261	0	6.7	44	63
7	0.055505	8.5355	7.9	56	75
8	0.055505	1.4645	6.7	51	74
9	0.07261	5	7.7	61	79
10	0.089715	1.4645	7.2	61	79
11	0.07261	10	8.7	58	78
12	0.089715	8.5355	8.9	51	66
13	0.0968	5	8.1	63	85

15

Table 21. DOE 3

DOE 3													
RUN #:	1	2	3	4	5	6	7	8	9	10	11	12	13
MODEL OF OIL PHASE:													
Homosalate	11	11	11	11	11	11	11	11	11	11	11	11	11
Octyl Salicylate	5	5	5	5	5	5	5	5	5	5	5	5	5
Oxybenzone	6	6	6	6	6	6	6	6	6	6	6	6	6
Octocrylene	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5
Avobenzonone	1.5	2.3	2.3	2.3	2.3	2.3	1.7	1.7	2.3	2.8	2.3	2.8	3
1,2-Octanediol	3.5	5.3	5.3	5.3	5.3	0	6.9	1.2	5.3	1.9	10.6	11	7
Neopentyl Glycol Diheptanoate QS to 38.5 g	9	6.4	6.4	6.4	6.4	11.7	5.4	11.1	6.4	9.3	1.1	0	4
	38.5	38.5	38.5	38.5	38.5	38.5	38.5	38.5	38.5	38.5	38.5	38.5	38.5

10 Table 22. UVA and UVB Photostabilization for DOE 3

DOE 3	Factor 1	Factor 2	Dielectric Constant	Retained Abs, % at 370 nm after 15 MED	Retained Abs, % at 310 nm after 15 MED
n	A:Avo, M	B:Molar Ratio Octanediol:Avo	23 C		
1	0.04842	5	7.40	54	78
2	0.07261	5	8.10	61	85
3	0.07261	5	8.10	63	79
4	0.07261	5	8.10	63	81
5	0.07261	5	8.10	63	81
6	0.07261	0	7.10	60	72
7	0.055505	8.5355	8.10	77	97
8	0.055505	1.4645	7.10	47	72
9	0.07261	5	8.10	63	81
10	0.089715	1.4645	7.40	60	70
11	0.07261	10	8.90	63	81
12	0.089715	8.5355	9.20	66	82
13	0.0968	5	8.50	62	75

1,2-octanediol (caprylyl glycol) significantly improves the photostability of sunscreens at 310 nm (UVB sunscreens) and at 370 nm (UVA sunscreens such as Avobenzene). The concentration of avobenzene, molar ratio of 1,2-octanediol (caprylyl glycol) to avobenzene and also other sunscreens
5 that are present in the sunscreen composition significantly influence the degree of photostabilization at 310 nm and 370 nm provided by 1,2-octanediol.

It should be understood that while many of the embodiments of the present invention were exemplified above through the testing and analysis
10 of oil phases of a photoprotective composition, the unexpected results are even more enhanced in final photoprotective compositions (versus just an oil phase alone), due in part to the inclusion of additional components, such as those described below.

15

ADDITIONAL COMPONENTS

The one or more optimization agents are preferably included in a solvent system used to dissolve the one or more photoactive compounds.
20 As used herein, a solvent system includes all of the compounds used to dissolve the one or more photoactive compounds. By way of example, the solvent system for cosmetic sunscreen compositions will include lipophilic compounds for dissolving oil miscible photoactive compounds and hydrophilic compounds for dissolving water miscible photoactive
25 compounds. In a preferred embodiment, the solvent system includes one

or more optimization agents that are both oil miscible and water miscible. Such agents offer a distinct advantage when formulating the sunscreen composition, as the need for separate oil and water miscible solvents is reduced or eliminated.

5

In addition to the one or more optimization agents, the solvent system may include, but is not limited to, one or more solvents, such as C₁₂-C₁₅ alkyl benzoates, capric triglycerides, caprylic triglycerides, diethylhexyl adipate, diethylhexyl malate, diethylhexyl 2,6-naphthalate, ethylhexyl palmitate, ethylhexyl stearate, isoeicosane, isopropyl myristate, isopropyl palmitate, mineral oil, octyldodecyl neopentanoate, polyisobutene, PPG-2 myristyl ether propionate, cocoglycerides, isostearyl linoleate, diisopropyl adipate, myristyl ether myristate, octyl palmitate, propylene glycol ricinoleate, cetyl esters, propylene glycol laurate, or any combinations thereof. Although not necessary, it is preferred that the one or more solvents, when used with photoactive compounds such as those identified above, have a dielectric constant of about 1 to about 12. By using highly polar solvents, in addition to dissolving the one or more photoactive compounds, the solvents may contribute to the overall polarity of the composition.

20

In addition to the one or more photoactive compounds and the one or more optimizing agents, any well-known cosmetically-acceptable additives, such as, for example, water, emulsifier, thickening agent, emollient, SPF booster, moisturizer, humectant, film former/waterproofing

25

agent, bio-active (functional) ingredient, pH adjuster/chelating agent, preservative, fragrance, effect pigment, color additive, lubricant, elastomer, or any combinations thereof, and/or other common cosmetic formulation additives for solubility of sunscreen active compounds, emulsification, thickening and to provide other skin enhancement, e.g., moisturizing properties, may be included in the sunscreen composition.

The compositions of the present invention preferably include water. The water is present in the compositions of the present invention in an amount about 40 wt.% to about 90 wt.%, and preferably about 50 wt.% to about 80 wt.%, of the total weight of the compositions.

The compositions of the present invention may include one or more emulsifiers. The one or more emulsifiers suitable for use in the present invention include, but are not limited to, acrylates crosspolymer, acrylates/C₁₀₋₃₀ alkylacrylate crosspolymer, acrylates/vinyl isodecanoate crosspolymer, polyacrylic acid, sodium polymethacrylate, sodium polyacrylate, polyacrylates, cetyl alcohol, cetearyl alcohol, oleth-10, diethylhexyl esters, sorbitan oleate, sorbitan sesquioleate, sorbitan isostearate, sorbitan trioleate, PEG-20 almond glycerides, polyglyceryl-3-diisostearate, polyglycerol esters of oleic/isostearic acid, polyglyceryl-6 hexaricinolate, polyglyceryl-4-oleate, polyglyceryl-4 oleate/PEG-8 propylene glycol cocoate, sodium glyceryl oleate phosphate, hydrogenated vegetable glycerides phosphate, or any combinations thereof.

The amount of emulsifier present in the compositions of the present invention is about 0.01 wt.% to about 10 wt.% of the total weight of the composition. Preferably, the emulsifier is present in an amount about 0.1 wt.% to about 5 wt.% of the total weight of the composition.

5

In an alternative embodiment, the sunscreen composition of the present invention may be formulated to be emulsifier free, which can improve water resistance of the sunscreen formulation and reduce its irritation potential.

10

One or more thickening agents that may be used in the compositions of the present invention. Suitable thickening agent includes, but is not limited to, one or more stabilizers, synthetic and natural gum or polymer products, polysaccharide thickening agents, associative
15 thickeners, anionic associative rheology modifiers, nonionic associative rheology modifiers, oil-thickening agents, acrylates/C10-30 alkyl acrylate crosspolymer, acrylates/aminoacrylates/C10-30 alkyl PEG-20 itaconate copolymer, acrylates copolymer, acrylates/steareth-20 methacrylate copolymer, acrylates/beheneth-25 methacrylate copolymer, PEG-150/decyl
20 alcohol/SMDI copolymer, PVP, PVM/MA decadiene crosspolymer, carbomer, PEG crosspolymer, acrylates/palmeth-25 acrylates copolymer, polysaccharides, polyacrylates, polyether-1, sodium magnesium silicate, sodium carbomer, sodium polyacrylate, sodium polymethacrylate, sodium polyacryloyldimethyl taurate, sodium acryloyldimethyl taurate copolymer,
25 sodium carragenan, sodium carboxymethyl dextran, hydroxyethylcellulose,

hydroxypropyl cyclodextran, bentonites, trihydroxystearin, aluminum-magnesium hydroxide stearate, xanthan gum, or any combinations thereof.

The amount of thickening agent present in the compositions of the present invention is about 0.01 wt.% to about 10 wt.% of the total weight of the composition. Preferably, the thickener is present in an amount about 0.1 wt.% to about 5 wt.% of the total weight of the composition.

The present compositions may include one or more emollients. An emollient provides a softening, protective or soothing effect on the skin surface and is generally considered safe for topical use. It also helps control the rate of evaporation and the tackiness of the compositions.

Suitable emollients include, for example, cocoglycerides, cyclomethicone, dimethicone, dicapryl maleate, caprylic/capric triglyceride, isopropyl myristate, octyl stearate, isostearyl linoleate, lanolin oil, coconut oil, cocoa butter, olive oil, avocado oil, aloe extracts, jojoba oil, castor oil, fatty acid such as oleic and stearic, fatty alcohol such as cetyl and hexadecyl, diisopropyl adipate, hydroxybenzoate esters, benzoic acid esters of C₉-C₁₅ alcohols, isononyl iso-nonanoate, alkanes such as mineral oil, silicone such as dimethyl polysiloxane, ether such as polyoxypropylene butyl ether and polyoxypropylene cetyl ether, C₁₂-C₁₅ alkyl benzoate, or any combinations thereof.

The total amount of emollient present in the compositions is typically about 0.1 wt.% to about 30 wt.% of the total weight of the composition. Preferably, emollient is present in an amount about 1 wt.% to about 20 wt.% of the total weight of the composition.

5

The pH of the compositions of the present invention may be adjusted by one or more basic pH adjusters and/or chelating agents. For example, sodium hydroxide, triethanolamine, EDTA salts, or any combinations thereof are suitable pH adjusters/chelating agents that may be included in the sunscreen compositions of the present invention.

10

An effective amount of a pH adjuster and/or chelating agent is included to adjust the pH of the final compositions to about 3 to about 9. Preferably, the pH is adjusted to about 5 to about 8 and more preferably about 6 to about 7.

15

One or more humectants may be used in the compositions of the present invention. Suitable humectants include, but are not limited to, glycerin, pentylene glycol, hexylene glycol, propylene glycol, butylene glycol, sorbitol, PEG-4, or any combinations thereof.

20

One or more humectants may be included in the compositions of the present invention in an amount about 0.1 wt.% to about 15 wt.% of the total weight of the composition. Preferably, humectant is present in an amount about 1 wt.% to about 5 wt.% of the total weight of the composition.

25

The present compositions may include one or more SPF boosters. SPF booster itself is not an active ingredient, but is designed to enhance the effectiveness of the sunscreen actives present in the formulation. Suitable SPF boosters include, but are not limited to, styrene/acrylates copolymer, sodium bentonites, highly purified white sodium bentonites, montmorillonite, hydrogels, fluorene derivatives, ester derivatives of cyano(9H-fluoren-9-ylidene), amides, malates, bis-urethanes, or any combinations thereof. A preferred styrene/acrylates copolymer for use in the present invention is sold under the trade name SunSpheres[®] by Rohm and Haas Company.

When present, the one or more SPF boosters may be included in the compositions of the present invention in an amount about 1 wt.% to about 6 wt.% of the total weight of the composition. Preferably, SPF booster is present in an amount about 2 wt.% to about 3 wt.% of the total weight of the composition.

Another component that may be used in the compositions of the present invention is a film former/waterproofing agent. The film former/waterproofing agent is a hydrophobic material that imparts film forming and waterproofing characteristics to the emulsion. Suitable film former/waterproofing agent for use in the compositions of the present invention include, but is not limited to, acrylates/acrylamide copolymer, acrylates copolymer, acrylates/C₁₂₋₂₂ alkylmethacrylate copolymer, polyethylene, waxes, VP/Dimethiconylacrylate/Polycarbamyl polyglycol

ester, butylated PVP, PVP/Hexadecene copolymer, octadecene/MA
copolymer, PVP/Eicosene copolymer, tricontanyl PVP, Brassica
Campestris/Aleuritis Fordi Oil copolymer, decamethyl cyclopentasiloxane
(and) trimethylsiloxysilicate, or any combinations thereof.

5

One or more film formers/waterproofing agents may be present in
the compositions of the present invention in an amount about 0.1 wt.% to
about 5 wt.% of the total weight of the composition. Preferably, the one or
more film formers/waterproofing agents is present in the compositions of
10 the present invention in an amount about 1 wt.% to about 3 wt.% of the
total weight of the composition.

In addition, it has been unexpectedly found that the acrylates/ C_{12-22}
alkylmethacrylate copolymer may protect the lipids in a user's skin by
15 imparting structure to the epidermal lipids in the skin (stratum corneum)
and sebaceous lipids (sebum) and preventing them from depletion. As a
result, it is believed that the acrylates/ C_{12-22} alkylmethacrylate copolymer
may enhance and help to maintain the barrier properties of the lipid barrier
in stratum corneum.

20

Therefore, the compositions of the present invention may exhibit
moisturizing and anti-inflammatory properties without a need for including
moisturizers and/or anti-inflammatory agents in the compositions.

One or more preservatives may be included in the compositions of the present invention. The preservative protects the compositions from microbial contamination and/or oxidation. As such, the preservative can include an antioxidant. Preservatives, such as diazolidinyl urea, 5 iodopropynyl butylcarbamate, chloromethylisothiazolinone, methylisothiazolinone, vitamin E and its derivatives including vitamin E acetate, vitamin C, butylated hydroxytoluene, butylparaben, ethylparaben, methylparaben, propylparaben, isobutylparaben, phenoxyethanol, or any mixtures thereof, may be included as a preservative in a composition of the 10 present invention.

About 0.01 wt.% to about 2 wt.% of preservative may be included in a composition of the present invention. Preferably, one or more preservatives total about 0.5 wt.% to about 1.5 wt.% of the total weight of 15 the composition.

The compositions of the present invention may also have other optional additives including bio-active (functional) ingredients. For instance, one or more plant extracts, fruit extracts, vegetable extracts, 20 algae extracts, sugars, polysaccharides, lipids, proteins, peptides, aminoacids, aminoacid derivatives, absorbents, elastomers, for example DC 9011 silicone elastomer blend (Dow Corning) (cyclopentasiloxane (and) PEG-12 (and) dimethicone crosspolymer), salicylic acid, alpha and beta hydroxy acids, oil and water soluble vitamins including vitamins A, C, and E

and their derivatives, or any mixtures thereof, may be included in the
sunscreen compositions.

When present, the optional additives may be included in the present
5 composition in an amount about 0.001 wt.% to about 10 wt.%, based on
the total weight of the composition.

The compositions can be produced as lotions, creams, ointments,
gels, solid sticks, emulsions, aerosols, solutions, dispersions, or any other
10 forms of cosmetic compositions.

Further aspects of the invention may become apparent to those
skilled in the art from a review of the detailed description set forth above. It
should be understood that the disclosure is merely illustrative, and is not
15 intended to limit the invention to the specific embodiments described.

WE CLAIM:

1. A photoprotective composition comprising:

5 one or more sunscreen agents; and

one or more optimizing agents selected from the group consisting of diol, alcohol, glycol, polyhydric alcohol, any derivatives thereof, and any combinations thereof,

10

wherein said photoprotective composition has one or more optimized properties selected from the group consisting of polarity, critical wavelength, SPF, PFA, Star Rating, photostability, and any combinations thereof, as compared to a composition without said one or more optimizing agents.

15

2. The photoprotective composition of claim 1, wherein said one or more sunscreen agents is selected from the group consisting of p-aminobenzoic acid, p-aminobenzoic acid salts, p-aminobenzoic acid derivatives, anthranilates, salicylates, glyceryl ester, dipropyleneglycol esters, cinnamic acid derivatives, dihydroxycinnamic acid derivatives, camphor, camphor derivatives, trihydroxycinnamic acid derivatives, hydrocarbons, dibenzalacetone, benzalacetophenone, naphtholsulfonates, dihydroxy-naphthoic acid, dihydroxy-naphthoic acid salts; o-

20

25 hydroxydiphenyldisulfonates, p-hydroxydiphenyldisulfonates, coumarin

derivatives, diazoles, quinine salts, quinoline derivatives, hydroxy-substituted benzophenones, methoxy-substituted benzophenones, uric acids, vilouric acids, tannic acid, tannic acid derivatives, hydroquinone, benzophenones, avobenzene, 4-isopropylidibenzoylmethane,
5 butylmethoxydibenzoylmethane, octocrylene, 4-isopropyl-
dibenzoylmethane, metal oxides, titanium dioxide, zinc oxide, and any combinations thereof.

3. The photoprotective composition of claim 1, wherein said one or
10 more sunscreen agents are present in an amount about 1 wt.% to about 40 wt.%, based on the total weight of the composition.

4. The photoprotective composition of claim 1, wherein said one or
15 more optimizing agents are one or more diols, glycols, and any combinations thereof.

5. The photoprotective composition of claim 1, wherein said one or
more optimizing agents are one or more glycols selected from the group
consisting of pentylene glycol (1,2-pentanediol), neopentyl glycol
20 (neopentanediol), caprylyl glycol (1,2-octanediol), ethoxydiglycol, butylene glycol monopropionate, diethylene glycol monobutyl ether, PEG-7 methyl ether, octacosanyl glycol, arachidyl glycol, benzyl glycol, cetyl glycol (1,2-hexanediol), C₁₄₋₁₈ glycol, C₁₅₋₁₈ glycol, lauryl glycol (1,2-dodecanediol), butoxydiglycol, 1,10-decanediol, ethyl hexanediol, or any combinations
25 thereof.

6. The photoprotective composition of claim 1, wherein said one or more optimizing agents are one or more glycols selected from the group consisting of 1,2-pentanediol, 1,2-octanediol, and any combination thereof.
- 5 7. The photoprotective composition of claim 1, wherein said one or more optimizing agents is present in said composition in an amount about 0.1 wt.% to about 40 wt.%, based on the total weight of the composition.
8. The photoprotective composition of claim 1, wherein said one or
10 more optimizing agents is present in said composition in an amount about 0.1 wt.% to about 10 wt.%, based on the total weight of the composition.
9. The photoprotective composition of claim 1, wherein said one or more optimizing agents, taken alone or in combination, has a dielectric
15 constant greater than about 10.5.
10. The photoprotective composition of claim 1, wherein said one or more optimizing agents, taken alone or in combination, has a dielectric
20 constant greater than about 13.
11. The photoprotective composition of claim 1, wherein said composition further comprises one or more components selected from the group consisting of solvent, water, emulsifier, thickening agent, emollient, SPF booster, moisturizer, humectant, film former/waterproofing agent, bio-

active, pH adjuster/chelating agent, preservative, fragrance, effect pigment, color additive, lubricant, elastomer, and any combinations thereof.

12. A photoprotective composition comprising:

5

one or more sunscreen agents; and

one or more optimizing agents selected from the group consisting of diol, alcohol, glycol, polyhydric alcohol, any derivatives thereof, and any combinations thereof,

10

wherein said one or more optimizing agents, taken alone or in combination, has a dielectric constant greater than about 13.

15 13. The photoprotective composition of claim 12, wherein said one or more sunscreen agents is selected from the group consisting of p-aminobenzoic acid, p-aminobenzoic acid salts, p-aminobenzoic acid derivatives, anthranilates, salicylates, glyceryl ester, dipropyleneglycol esters, cinnamic acid derivatives, dihydroxycinnamic acid derivatives,

20 camphor, camphor derivatives, trihydroxycinnamic acid derivatives, hydrocarbons, dibenzalacetone, benzalacetophenone, naphtholsulfonates, dihydroxy-naphthoic acid, dihydroxy-naphthoic acid salts; o-hydroxydiphenyldisulfonates, p-hydroxydiphenyldisulfonates, coumarin derivatives, diazoles, quinine salts, quinoline derivatives, hydroxy-

25 substituted benzophenones, methoxy-substituted benzophenones, uric

acids, vilouric acids, tannic acid, tannic acid derivatives, hydroquinone, benzophenones, avobenzene, 4-isopropylidibenzoylmethane, butylmethoxydibenzoylmethane, octocrylene, 4-isopropylidibenzoylmethane, metal oxides, titanium dioxide, zinc oxide, and any
5 combinations thereof.

14. The photoprotective composition of claim 12, wherein said one or more sunscreen agents are present in an amount about 1 wt.% to about 40 wt.%, based on the total weight of the composition.

10

15. The photoprotective composition of claim 12, wherein said one or more optimizing agents are one or more glycols selected from the group consisting of pentylene glycol, neopentyl glycol, caprylyl glycol, 1,2-octanediol, etoxydiglycol, butylene glycol monopropionate, diethylene
15 glycol monobutyl ether, PEG-7 methyl ether, octacosanyl glycol, arachidyl glycol, benzyl glycol, cetyl glycol, C₁₄₋₁₈ glycol, C₁₅₋₁₈ glycol, lauryl glycol, butoxydiglycol, 1,10-decanediol, ethyl hexanediol, and any combinations thereof.

20 16. The photoprotective composition of claim 12, wherein said one or more optimizing agents are one or more glycols selected from the group consisting of 1,2-pentanediol, 1,2-octanediol, and any combination thereof.

17. The photoprotective composition of claim 12, wherein said one or more optimizing agents is present in said composition in an amount about 0.1 wt.% to about 40 wt.%, based on the total weight of the composition.

5 18. The photoprotective composition of claim 12, wherein said one or more optimizing agents is present in said composition in an amount about 0.1 wt.% to about 10 wt.%, based on the total weight of the composition.

19. The photoprotective composition of claim 12, wherein said
10 composition further comprises one or more components selected from the group consisting of solvent, water, emulsifier, thickening agent, emollient, SPF booster, moisturizer, humectant, film former/waterproofing agent, bio-active, pH adjuster/chelating agent, preservative, fragrance, effect pigment, color additive, lubricant, elastomer, and any combinations thereof.

15

20. A photoprotective composition comprising:

one or more sunscreen agents; and

20 one or more optimizing agents selected from the group consisting of diol, alcohol, glycol, polyhydric alcohol, any derivatives thereof, and any combinations thereof,

wherein said one or more optimizing agents increase a SPF of said composition by at least 25% compared to a composition without said one or more optimizing agents.

- 5 21. The photoprotective composition of claim 20, wherein said one or more optimizing agents increase a SPF of said composition by at least about 30% compared to a composition without said one or more optimizing agents.
- 10 22. The photoprotective composition of claim 20, wherein said one or more sunscreen agents is selected from the group consisting of p-aminobenzoic acid, p-aminobenzoic acid salts, p-aminobenzoic acid derivatives, anthranilates, salicylates, glyceryl ester, dipropylene glycol esters, cinnamic acid derivatives, dihydroxycinnamic acid derivatives,
- 15 camphor, camphor derivatives, trihydroxycinnamic acid derivatives, hydrocarbons, dibenzalacetone, benzalacetophenone, naphtholsulfonates, dihydroxy-naphthoic acid, dihydroxy-naphthoic acid salts; o-
- hydroxydiphenyldisulfonates, p-hydroxydiphenyldisulfonates, coumarin derivatives, diazoles, quinine salts, quinoline derivatives, hydroxy-
- 20 substituted benzophenones, methoxy-substituted benzophenones, uric acids, vilouric acids, tannic acid, tannic acid derivatives, hydroquinone, benzophenones, avobenzene, 4-isopropylidibenzoylmethane, butylmethoxydibenzoylmethane, octocrylene, 4-isopropyl-
- dibenzoylmethane, metal oxides, titanium dioxide, zinc oxide, and any
- 25 combinations thereof.

23. The photoprotective composition of claim 20, wherein said one or more sunscreen agents are present in an amount about 1 wt.% to about 40 wt.%, based on the total weight of the composition.
- 5 24. The photoprotective composition of claim 20, wherein said one or more optimizing agents are one or more glycols selected from the group consisting of pentylene glycol (1,2-pentanediol), neopentyl glycol (neopentanediol), caprylyl glycol (1,2-octanediol), ethoxydiglycol, butylene glycol monopropionate, diethylene glycol monobutyl ether, PEG-7 methyl ether, octacosanyl glycol, arachidyl glycol, benzyl glycol, cetyl glycol (1,2-
10 hexanediol), C₁₄₋₁₈ glycol, C₁₅₋₁₈ glycol, lauryl glycol (1,2-dodecanediol), butoxydiglycol, 1,10-decanediol, ethyl hexanediol, or any combinations thereof.
- 15 25. The photoprotective composition of claim 20, wherein said one or more optimizing agents are one or more glycols selected from the group consisting of 1,2-pentanediol, 1,2-octanediol, and any combination thereof.
26. The photoprotective composition of claim 20, wherein said one or
20 more optimizing agents is present in said composition in an amount about 0.1 wt.% to about 40 wt.%, based on the total weight of the composition.
27. The photoprotective composition of claim 20, wherein said one or
25 more optimizing agents is present in said composition in an amount about 0.1 wt.% to about 10 wt.%, based on the total weight of the composition.

28. The photoprotective composition of claim 20, wherein said one or more optimizing agents, taken alone or in combination, has a dielectric constant greater than about 10.5.

5

29. The photoprotective composition of claim 20, wherein said one or more optimizing agents, taken alone or in combination, has a dielectric constant greater than about 13.

10 30. The photoprotective composition of claim 20, wherein said composition further comprises one or more components selected from the group consisting of solvent, water, emulsifier, thickening agent, emollient, SPF booster, moisturizer, humectant, film former/waterproofing agent, bio-active, pH adjuster/chelating agent, preservative, fragrance, effect pigment,
15 color additive, lubricant, elastomer, and any combinations thereof.

31. A photoprotective composition comprising:

one or more sunscreen agents; and

20

one or more optimizing agents selected from the group consisting of diol, alcohol, glycol, polyhydric alcohol, any derivatives thereof, and any combinations thereof,

wherein said one or more optimizing agents increase a PFA of said composition by at least 10% compared to a composition without said one or more optimizing agents.

5 32. The photoprotective composition of claim 31, wherein said one or more optimizing agents increase a PFA of said composition by at least about 85% compared to a composition without said one or more optimizing agents.

10 33. The photoprotective composition of claim 31, wherein said one or more sunscreen agents is selected from the group consisting of p-aminobenzoic acid, p-aminobenzoic acid salts, p-aminobenzoic acid derivatives, anthranilates, salicylates, glyceryl ester, dipropylene glycol esters, cinnamic acid derivatives, dihydroxycinnamic acid derivatives,
15 camphor, camphor derivatives, trihydroxycinnamic acid derivatives, hydrocarbons, dibenzalacetone, benzalacetophenone, naphtholsulfonates, dihydroxy-naphthoic acid, dihydroxy-naphthoic acid salts; o-hydroxydiphenyldisulfonates, p-hydroxydiphenyldisulfonates, coumarin derivatives, diazoles, quinine salts, quinoline derivatives, hydroxy-
20 substituted benzophenones, methoxy-substituted benzophenones, uric acids, vilouric acids, tannic acid, tannic acid derivatives, hydroquinone, benzophenones, avobenzene, 4-isopropyl dibenzoylmethane, butylmethoxydibenzoylmethane, octocrylene, 4-isopropyl-
dibenzoylmethane, metal oxides, titanium dioxide, zinc oxide, and any
25 combinations thereof.

34. The photoprotective composition of claim 31, wherein said one or more sunscreen agents are present in an amount about 1 wt.% to about 40 wt.%, based on the total weight of the composition.

5

35. The photoprotective composition of claim 31, wherein said one or more optimizing agents are one or more glycols selected from the group consisting of pentylene glycol (1,2-pentanediol), neopentyl glycol (neopentanediol), caprylyl glycol (1,2-octanediol), ethoxydiglycol, butylene glycol monopropionate, diethylene glycol monobutyl ether, PEG-7 methyl ether, octacosanyl glycol, arachidyl glycol, benzyl glycol, cetyl glycol (1,2-hexanediol), C₁₄₋₁₈ glycol, C₁₅₋₁₈ glycol, lauryl glycol (1,2-dodecanediol), butoxydiglycol, 1,10-decanediol, ethyl hexanediol, or any combinations thereof.

15

36. The photoprotective composition of claim 31, wherein said one or more optimizing agents are one or more glycols selected from the group consisting of 1,2-pentanediol, pentylene glycol, 1,2-octanediol, and any combination thereof.

20 37. The photoprotective composition of claim 31, wherein said one or more optimizing agents is present in said composition in an amount about 0.1 wt.% to about 40 wt.%, based on the total weight of the composition.

38. The photoprotective composition of claim 31, wherein said one or more optimizing agents is present in said composition in an amount about 0.1 wt.% to about 10 wt.%, based on the total weight of the composition.

5 39. The photoprotective composition of claim 31, wherein said one or more optimizing agents, taken alone or in combination, has a dielectric constant greater than about 10.5.

40. The photoprotective composition of claim 31, wherein said one or
10 more optimizing agents, taken alone or in combination, has a dielectric constant greater than about 13.

41. The photoprotective composition of claim 31, wherein said
composition further comprises one or more components selected from the
15 group consisting of solvent, water, emulsifier, thickening agent, emollient, SPF booster, moisturizer, humectant, film former/waterproofing agent, bio-active, pH adjuster/chelating agent, preservative, fragrance, effect pigment, color additive, lubricant, elastomer, and any combinations thereof.

20 42. A photoprotective composition comprising:

a synergistic combination of octocrylene and one or more optimizing agents selected from the group consisting of diol, alcohol, glycol, polyhydric alcohol, any derivatives thereof, and any combinations thereof,

25

wherein, as a result of said synergistic combination, said photoprotective composition has an increase in a UVA photostability of at least about 10%, an increase in a UVB photostability of at least about 10% and a decrease in a critical wavelength as compared to a composition
5 without said synergistic combination of octocrylene and one or more optimizing agents.

43. The photoprotective composition of claim 42, wherein said synergistic combination of one or more optimizing agents and octocrylene
10 is present in a weight ratio of one or more optimizing agents to octocrylene of about 0.1 to about 10.

44. The photoprotective composition of claim 42, further comprising one or more sunscreen agents selected from the group consisting of p-
15 aminobenzoic acid, p-aminobenzoic acid salts, p-aminobenzoic acid derivatives, anthranilates, salicylates, glyceryl ester, dipropyleneglycol esters, cinnamic acid derivatives, dihydroxycinnamic acid derivatives, camphor, camphor derivatives, trihydroxycinnamic acid derivatives, hydrocarbons, dibenzalacetone, benzalacetophenone, naphtholsulfonates,
20 dihydroxy-naphthoic acid, dihydroxy-naphthoic acid salts; o-hydroxydiphenyldisulfonates, p-hydroxydiphenyldisulfonates, coumarin derivatives, diazoles, quinine salts, quinoline derivatives, hydroxy-substituted benzophenones, methoxy-substituted benzophenones, uric acids, vilouric acids, tannic acid, tannic acid derivatives, hydroquinone,
25 benzophenones, avobenzene, 4-isopropylidibenzoylmethane,

butylmethoxydibenzoylmethane, 4-isopropyl-dibenzoylmethane, metal oxides, titanium dioxide, zinc oxide, and any combinations thereof.

45. The photoprotective composition of claim 44, wherein said one or
5 more sunscreen agents are present in an amount about 1 wt.% to about 40 wt.%, based on the total weight of the composition.

46. The photoprotective composition of claim 42, wherein said one or
more optimizing agents are one or more glycols selected from the group
10 consisting of pentylene glycol (1,2-pentanediol), neopentyl glycol (neopentanediol), caprylyl glycol (1,2-octanediol), ethoxydiglycol, butylene glycol monopropionate, diethylene glycol monobutyl ether, PEG-7 methyl ether, octacosanyl glycol, arachidyl glycol, benzyl glycol, cetyl glycol (1,2-hexanediol), C₁₄₋₁₈ glycol, C₁₅₋₁₈ glycol, lauryl glycol (1,2-dodecanediol),
15 butoxydiglycol, 1,10-decanediol, ethyl hexanediol, or any combinations thereof.

47. The photoprotective composition of claim 42, wherein said one or
more optimizing agents are one or more glycols selected from the group
20 consisting of 1,2-pentanediol, 1,2-octanediol, and any combination thereof.

48. The photoprotective composition of claim 42, wherein said one or
more optimizing agents is present in said composition in an amount about
0.1 wt.% to about 40 wt.%, based on the total weight of the composition.

25

49. The photoprotective composition of claim 42, wherein said one or more optimizing agents is present in said composition in an amount about 0.1 wt.% to about 10 wt.%, based on the total weight of the composition.

5 50. The photoprotective composition of claim 42, wherein said composition further comprises one or more components selected from the group consisting of solvent, water, emulsifier, thickening agent, emollient, SPF booster, moisturizer, humectant, film former/waterproofing agent, bio-
10 active, pH adjuster/chelating agent, preservative, fragrance, effect pigment, color additive, lubricant, elastomer, and any combinations thereof.

51. A photoprotective composition comprising:

one or more dibenzoylmethane derivatives;

15

one or more additional sunscreen agents; and

one or more optimizing agents selected from the group consisting of diol, alcohol, glycol, polyhydric alcohol, any derivatives thereof, and any
20 combinations thereof,

wherein said one or more dibenzoylmethane derivatives is present in a molar concentration between about 0.016M to about 0.193M , and

wherein said one or more optimizing agents to said one or more dibenzoylmethane derivatives are present in said composition in a molar ratio of about 0.5 to about 400.

5 52. The photoprotective composition of claim 51, wherein said one or more dibenzoylmethane derivatives is present in a molar concentration between about 0.048M to about 0.096M.

53. The photoprotective composition of claim 51, wherein said one or
10 more optimizing agents to said one or more dibenzoylmethane derivatives are present in said composition in a molar ratio of about 0.5 to about 10.

54. The photoprotective composition of claim 51, wherein said one or more dibenzoylmethane derivatives is selected from the group consisting of
15 avobenzene, 4-isopropylidibenzoylmethane, butylmethoxydibenzoylmethane, 4-isopropyl-dibenzoylmethane, and any combinations thereof.

55. The photoprotective composition of claim 51, wherein said one or
20 more additional sunscreen agents are selected from the group consisting of p-aminobenzoic acid, p-aminobenzoic acid salts, p-aminobenzoic acid derivatives, anthranilates, salicylates, glyceryl ester, dipropyleneglycol esters, cinnamic acid derivatives, dihydroxycinnamic acid derivatives, camphor, camphor derivatives, trihydroxycinnamic acid derivatives,
25 hydrocarbons, dibenzalacetone, benzalacetophenone, naphtholsulfonates,

dihydroxy-naphthoic acid, dihydroxy-naphthoic acid salts; o-hydroxydiphenyldisulfonates, p-hydroxydiphenyldisulfonates, coumarin derivatives, diazoles, quinine salts, quinoline derivatives, hydroxy-substituted benzophenones, methoxy-substituted benzophenones, uric acids, vilouric acids, tannic acid, tannic acid derivatives, hydroquinone, benzophenones, octocrylene, metal oxides, titanium dioxide, zinc oxide, and any combinations thereof.

56. The photoprotective composition of claim 51, wherein said one or more dibenzoylmethane derivatives are present in an amount about 0.5 wt.% to about 6 wt.%, based on the total weight of the composition.

57. The photoprotective composition of claim 51, wherein said one or more additional sunscreen agents are present in an amount about 1 wt.% to about 40 wt.%, based on the total weight of the composition.

58. The photoprotective composition of claim 51, wherein said one or more optimizing agents are one or more glycols selected from the group consisting of pentylene glycol (1,2-pentanediol), neopentyl glycol (neopentanediol), caprylyl glycol (1,2-octanediol), ethoxydiglycol, butylene glycol monopropionate, diethylene glycol monobutyl ether, PEG-7 methyl ether, octacosanyl glycol, arachidyl glycol, benzyl glycol, cetyl glycol (1,2-hexanediol), C₁₄₋₁₈ glycol, C₁₅₋₁₈ glycol, lauryl glycol (1,2-dodecanediol), butoxydiglycol, 1,10-decanediol, ethyl hexanediol, or any combinations thereof.

59. The photoprotective composition of claim 51, wherein said one or more optimizing agents are one or more glycols selected from the group consisting of 1,2-pentanediol, 1,2-octanediol, and any combination thereof.

5

60. The photoprotective composition of claim 51, wherein said one or more optimizing agents is present in said composition in an amount about 0.1 wt.% to about 40 wt.%, based on the total weight of the composition.

10 61. The photoprotective composition of claim 51, wherein said one or more optimizing agents is present in said composition in an amount about 0.1 wt.% to about 10 wt.%, based on the total weight of the composition.

62. The photoprotective composition of claim 51, wherein said
15 composition further comprises one or more components selected from the group consisting of solvent, water, emulsifier, thickening agent, emollient, SPF booster, moisturizer, humectant, film former/waterproofing agent, bio-active, pH adjuster/chelating agent, preservative, fragrance, effect pigment, color additive, lubricant, elastomer, and any combinations thereof.

20

63. A photoprotective composition comprising:

one or more sunscreen agents; and

one or more optimizing agents selected from the group consisting of 1,2-octanediol, 1,2-pentanediol, and combinations thereof,

wherein said one or more optimizing agents optimizes one or more
5 properties selected from the group consisting of polarity, critical wavelength, SPF, PFA, Star Rating, photostability, and any combinations thereof, of said composition, as compared to a composition without said one or more optimizing agents.

10 64. The photoprotective composition of claim 63, wherein said one or more sunscreen agents is selected from the group consisting of p-aminobenzoic acid, p-aminobenzoic acid salts, p-aminobenzoic acid derivatives, anthranilates, salicylates, glyceryl ester, dipropylene glycol esters, cinnamic acid derivatives, dihydroxycinnamic acid derivatives,
15 camphor, camphor derivatives, trihydroxycinnamic acid derivatives, hydrocarbons, dibenzalacetone, benzalacetophenone, naphtholsulfonates, dihydroxy-naphthoic acid, dihydroxy-naphthoic acid salts; o-hydroxydiphenyldisulfonates, p-hydroxydiphenyldisulfonates, coumarin derivatives, diazoles, quinine salts, quinoline derivatives, hydroxy-
20 substituted benzophenones, methoxy-substituted benzophenones, uric acids, vilouric acids, tannic acid, tannic acid derivatives, hydroquinone, benzophenones, avobenzene, 4-isopropylidibenzoylmethane, butylmethoxydibenzoylmethane, octocrylene, 4-isopropyl-
dibenzoylmethane, metal oxides, titanium dioxide, zinc oxide, and any
25 combinations thereof.

65. The photoprotective composition of claim 63, wherein said one or more sunscreen agents are present in an amount about 1 wt.% to about 40 wt.%, based on the total weight of the composition.

5

66. The photoprotective composition of claim 63, wherein said one or more optimizing agents is present in said composition in an amount about 0.1 wt.% to about 40 wt.%, based on the total weight of the composition.

10 67. The photoprotective composition of claim 63, wherein said one or more optimizing agents is present in said composition in an amount about 0.1 wt.% to about 10 wt.%, based on the total weight of the composition.

68. The photoprotective composition of claim 63, wherein said
15 composition further comprises one or more components selected from the group consisting of solvent, water, emulsifier, thickening agent, emollient, SPF booster, moisturizer, humectant, film former/waterproofing agent, bio-active, pH adjuster/chelating agent, preservative, fragrance, effect pigment, color additive, lubricant, elastomer, and any combinations thereof.

20

69. A method of providing an efficient photoprotective composition comprising the steps of:

formulating a photoprotective composition with one or more
25 sunscreen agents and one or more optimizing agents selected from the

group consisting of diol, alcohol, glycol, polyhydric alcohol, any derivatives thereof, and any combinations thereof,

wherein said photoprotective composition has one or more
5 optimized properties selected from the group consisting of polarity, critical wavelength, SPF, PFA, Star Rating, photostability, and any combinations thereof, as compared to a composition without said one or more optimizing agents.

10 70. The method of claim 69, wherein said one or more sunscreen agents is selected from the group consisting of p-aminobenzoic acid, p-aminobenzoic acid salts, p-aminobenzoic acid derivatives, anthranilates, salicylates, glyceryl ester, dipropyleneglycol esters, cinnamic acid
15 derivatives, dihydroxycinnamic acid derivatives, camphor, camphor derivatives, trihydroxycinnamic acid derivatives, hydrocarbons, dibenzalacetone, benzalacetophenone, naphtholsulfonates, dihydroxy-naphthoic acid, dihydroxy-naphthoic acid salts; o-
20 hydroxydiphenyldisulfonates, p-hydroxydiphenyldisulfonates, coumarin derivatives, diazoles, quinine salts, quinoline derivatives, hydroxy-substituted benzophenones, methoxy-substituted benzophenones, uric
25 acids, vilouric acids, tannic acid, tannic acid derivatives, hydroquinone, benzophenones, avobenzene, 4-isopropylidibenzoylmethane, butylmethoxydibenzoylmethane, octocrylene, 4-isopropylidibenzoylmethane, metal oxides, titanium dioxide, zinc oxide, and any combinations thereof.

71. The method of claim 69, wherein said one or more sunscreen agents are present in an amount about 1 wt.% to about 40 wt.%, based on the total weight of the composition.

5

72. The method of claim 69, wherein said one or more optimizing agents are one or more glycols selected from the group consisting of pentylene glycol (1,2-pentanediol), neopentyl glycol (neopentanediol), caprylyl glycol (1,2-octanediol), ethoxydiglycol, butylene glycol monopropionate, diethylene glycol monobutyl ether, PEG-7 methyl ether, octacosanyl glycol, arachidyl glycol, benzyl glycol, cetyl glycol (1,2-hexanediol), C₁₄₋₁₈ glycol, C₁₅₋₁₈ glycol, lauryl glycol (1,2-dodecanediol), butoxydiglycol, 1,10-decanediol, ethyl hexanediol, or any combinations thereof.

73. The method of claim 69, wherein said one or more optimizing agents are one or more glycols selected from the group consisting of pentylene glycol, 1,2-octanediol, and any combination thereof.

74. The method of claim 69, wherein said one or more optimizing agents is present in said composition in an amount about 0.1 wt.% to about 40 wt.%, based on the total weight of the composition.

75. The method of claim 69, wherein said one or more optimizing agents is present in said composition in an amount about 0.1 wt.% to about 10 wt.%, based on the total weight of the composition.

76. The method of claim 69, wherein said one or more optimizing agents, taken alone or in combination, has a dielectric constant greater than about 10.5.

5

77. The method of claim 69, wherein said one or more optimizing agents, taken alone or in combination, has a dielectric constant greater than about 13.

10 78. The method of claim 69, wherein said composition further comprises one or more components selected from the group consisting of solvent, water, emulsifier, thickening agent, emollient, SPF booster, moisturizer, humectant, film former/waterproofing agent, bio-active, pH adjuster/chelating agent, preservative, fragrance, effect pigment, color
15 additive, lubricant, elastomer, and any combinations thereof.

79. A method for optimizing a polarity, critical wavelength, SPF, PFA, Star Rating, photostability, or any combinations thereof, in a photoprotective composition comprising the step of:

20

formulating said photoprotective composition with one or more sunscreen agents and one or more optimizing agents selected from the group consisting of diol, alcohol, glycol, polyhydric alcohol, any derivatives thereof, and any combinations thereof,

25

wherein said photoprotective composition has one or more optimized properties selected from the group consisting of polarity, critical wavelength, SPF, PFA, Star Rating, photostability, and any combinations thereof, as compared to a composition without said one or more optimizing agents.

80. The method of claim 79, wherein said one or more sunscreen agents is selected from the group consisting of p-aminobenzoic acid, p-aminobenzoic acid salts, p-aminobenzoic acid derivatives, anthranilates, salicylates, glyceryl ester, dipropyleneglycol esters, cinnamic acid derivatives, dihydroxycinnamic acid derivatives, camphor, camphor derivatives, trihydroxycinnamic acid derivatives, hydrocarbons, dibenzalacetone, benzalacetophenone, naphtholsulfonates, dihydroxy-naphthoic acid, dihydroxy-naphthoic acid salts; o-hydroxydiphenyldisulfonates, p-hydroxydiphenyldisulfonates, coumarin derivatives, diazoles, quinine salts, quinoline derivatives, hydroxy-substituted benzophenones, methoxy-substituted benzophenones, uric acids, vilouric acids, tannic acid, tannic acid derivatives, hydroquinone, benzophenones, avobenzene, 4-isopropylidibenzoylmethane, butylmethoxydibenzoylmethane, octocrylene, 4-isopropylidibenzoylmethane, metal oxides, titanium dioxide, zinc oxide, and any combinations thereof.

81. The method of claim 79, wherein said one or more sunscreen agents are present in an amount about 1 wt.% to about 40 wt.%, based on the total weight of the composition.

5 82. The method of claim 79, wherein said one or more optimizing agents are one or more glycols selected from the group consisting of pentylene glycol (1,2-pentanediol), neopentyl glycol (neopentanediol), caprylyl glycol (1,2-octanediol), ethoxydiglycol, butylene glycol monopropionate, diethylene glycol monobutyl ether, PEG-7 methyl ether, octacosanyl glycol,
10 arachidyl glycol, benzyl glycol, cetyl glycol (1,2-hexanediol), C₁₄₋₁₈ glycol, C₁₅₋₁₈ glycol, lauryl glycol (1,2-dodecanediol), butoxydiglycol, 1,10-decanediol, ethyl hexanediol, or any combinations thereof.

83. The method of claim 79, wherein said one or more optimizing agents
15 are one or more glycols selected from the group consisting of 1,2-pentanediol, pentylene glycol, 1,2-octanediol, and any combination thereof.

84. The method of claim 79, wherein said one or more optimizing agents is present in said composition in an amount about 0.1 wt.% to about 40
20 wt.%, based on the total weight of the composition.

85. The method of claim 79, wherein said one or more optimizing agents is present in said composition in an amount about 0.1 wt.% to about 10 wt.%, based on the total weight of the composition.

25

86. The method of claim 79, wherein said one or more optimizing agents, taken alone or in combination, has a dielectric constant greater than about 10.5.

5 87. The method of claim 79, wherein said one or more optimizing agents, taken alone or in combination, has a dielectric constant greater than about 13.

88. The method of claim 79, wherein said composition further comprises
10 one or more components selected from the group consisting of solvent, water, emulsifier, thickening agent, emollient, SPF booster, moisturizer, humectant, film former/waterproofing agent, bio-active, pH adjuster/chelating agent, preservative, fragrance, effect pigment, color additive, lubricant, elastomer, and any combinations thereof.

15

89. The photoprotective composition according to claim 1, further comprising an oil-phase, wherein said one or more optimizing agents are in said oil-phase.

20 90. The photoprotective composition according to claim 12, further comprising an oil-phase, wherein said one or more optimizing agents are in said oil-phase.

91. The photoprotective composition according to claim 20, further comprising an oil-phase, wherein said one or more optimizing agents are in said oil-phase.

5 92. The photoprotective composition according to claim 63, further comprising an oil-phase, wherein said one or more optimizing agents are in said oil-phase.

93. The photoprotective composition according to claim 69, further
10 comprising an oil-phase, wherein said one or more optimizing agents are in said oil-phase.

94. The photoprotective composition according to claim 79, further
15 comprising an oil-phase, wherein said one or more optimizing agents are in said oil-phase.

FIGURE 1




FIGURE 2




FIGURE 3




FIGURE 4




FIGURE 5




FIGURE 6

Retain Absorbance @ 370nm 15MED Response Surface
 Oxybenzone = present, Octocrylene = absent

D.D.E. FUSION GRAPH




FIGURE 7

Retain Absorbance @ 370nm 15MED Response Surface
 Oxybenzone = present, Octocrylene = present

D.o.E. FUSION GRAPH

