



(86) Date de dépôt PCT/PCT Filing Date: 2011/09/09
 (87) Date publication PCT/PCT Publication Date: 2012/03/22
 (45) Date de délivrance/Issue Date: 2020/01/14
 (85) Entrée phase nationale/National Entry: 2013/03/11
 (86) N° demande PCT/PCT Application No.: US 2011/051072
 (87) N° publication PCT/PCT Publication No.: 2012/036993
 (30) Priorité/Priority: 2010/09/14 (US61/403,407)

(51) Cl.Int./Int.Cl. *C12N 15/33* (2006.01),
C12N 15/44 (2006.01), *C12N 15/63* (2006.01)
 (72) Inventeurs/Inventors:
ROSS, TED M., US;
GILES, BRENDAN M., US
 (73) Propriétaire/Owner:
UNIVERSITY OF PITTSBURGH-OF THE
COMMONWEALTH SYSTEM OF HIGHER
EDUCATION, US
 (74) Agent: SMART & BIGGAR LLP

(54) Titre : ANTIGENES CONTRE LA GRIPPE A REACTIVITE LARGE OPTIMISES PAR ORDINATEUR
 (54) Title: COMPUTATIONALLY OPTIMIZED BROADLY REACTIVE ANTIGENS FOR INFLUENZA

(57) **Abrégé/Abstract:**

Described herein is the development of a computationally optimized influenza HA protein that elicits broadly reactive immune response to all H5N1 influenza virus isolates. The optimized HA protein was developed through a series of HA protein alignments, and subsequent generation of consensus sequences, for clade 2 H5N1 influenza virus isolates. The final consensus HA amino acid sequence was reverse translated and optimized for expression in mammalian cells. It is disclosed herein that influenza virus-like particles containing the optimized HA protein are an effective vaccine against H5N1 influenza virus infection in animals.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
22 March 2012 (22.03.2012)

PCT

(10) International Publication Number
WO 2012/036993 A1

(51) International Patent Classification:

C12N 15/33 (2006.01) C12N 15/63 (2006.01)
C12N 15/44 (2006.01)

(21) International Application Number:

PCT/US2011/051072

(22) International Filing Date:

9 September 2011 (09.09.2011)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/403,407 14 September 2010 (14.09.2010) US

(71) Applicant (for all designated States except US): **UNIVERSITY OF PITTSBURGH-OF THE COMMONWEALTH SYSTEM OF HIGHER EDUCATION** [US/US]; 200 Gardner Steel Conference Center, Thackeray And O'hara Streets, Pittsburgh, PA 15260 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **ROSS, Ted, M.** [US/US]; 1652 Heritage Drive, Pittsburgh, PA 15237 (US). **GILES, Brendan, M.** [US/US]; 1401 Delgany Street, Denver, CO 80202 (US).(74) Agent: **CONNOLLY, Jodi, L.**; Klarquist Sparkman, LLP, One World Trade Center, Suite 1600, 121 SW Salmon Street, Portland, OR 97204 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— of inventorship (Rule 4.17(iv))

Published:

— with international search report (Art. 21(3))

— with sequence listing part of description (Rule 5.2(a))

(54) Title: COMPUTATIONALLY OPTIMIZED BROADLY REACTIVE ANTIGENS FOR INFLUENZA

(57) Abstract: Described herein is the development of a computationally optimized influenza HA protein that elicits broadly reactive immune response to all H5N1 influenza virus isolates. The optimized HA protein was developed through a series of HA protein alignments, and subsequent generation of consensus sequences, for clade 2 H5N1 influenza virus isolates. The final consensus HA amino acid sequence was reverse translated and optimized for expression in mammalian cells. It is disclosed herein that influenza virus-like particles containing the optimized HA protein are an effective vaccine against H5N1 influenza virus infection in animals.



WO 2012/036993 A1

63198-1692

COMPUTATIONALLY OPTIMIZED BROADLY REACTIVE ANTIGENS FOR INFLUENZA

CROSS REFERENCE TO RELATED APPLICATIONS

This application claims priority from U.S. Provisional Application No. 61/403,407, filed September 14, 2010.

FIELD

This disclosure concerns an optimized influenza hemagglutinin protein that elicits broadly reactive immune responses to H5N1 virus isolates and its use as a vaccine.

BACKGROUND

Influenza virus is a member of *Orthomyxoviridae* family. There are three subtypes of influenza viruses, designated influenza A, influenza B, and influenza C. The influenza virion contains a segmented negative-sense RNA genome, which encodes the following proteins: hemagglutinin (HA), neuraminidase (NA), matrix (M1), proton ion-channel protein (M2), nucleoprotein (NP), polymerase basic protein 1 (PB1), polymerase basic protein 2 (PB2), polymerase acidic protein (PA), and nonstructural protein 2 (NS2). The HA, NA, M1, and M2 are membrane associated, whereas NP, PB1, PB2, PA, and NS2 are nucleocapsid associated proteins. The M1 protein is the most abundant protein in influenza particles. The HA and NA proteins are envelope glycoproteins, responsible for virus attachment and penetration of the viral particles into the cell, and the sources of the major immunodominant epitopes for virus neutralization and protective immunity. Both HA and NA proteins are considered the most important components for prophylactic influenza vaccines.

Each year, seasonal influenza causes over 300,000 hospitalizations and 36,000 deaths in the U.S. alone (Simonsen *et al.*, *Lancet Infect Dis* 7:658-66, 2007). The emergence of the novel H1N1 influenza virus in 2009 demonstrated how

quickly a new influenza pandemic can sweep across the world. The spread of highly pathogenic H5N1 viruses in birds and coincident infections in humans have raised the concerns that H5N1 viruses may cause a new pandemic in humans. Vaccination is an effective method to prevent influenza infection. There are two influenza vaccine approaches licensed in the United States; the inactivated, split vaccine and the live-attenuated virus vaccine. Inactivated vaccines can efficiently induce humoral immune responses but generally only poor cellular immune responses. Thus, a need exists for a broadly protective influenza virus vaccine.

SUMMARY

Disclosed herein is the development of an optimized influenza HA protein that elicits broadly reactive immune response to H5N1 influenza virus isolates. The optimized HA protein was developed through a series of HA protein alignments, and subsequent generation of consensus sequences for clade 2 H5N1 influenza virus isolates (FIG.1). The final consensus HA amino acid sequence was reverse translated and optimized for expression in mammalian cells. The optimized HA coding sequence is set forth herein as SEQ ID NO: 1, and the optimized HA protein sequence is set forth herein as SEQ ID NO: 2.

Provided herein is an isolated nucleic acid molecule comprising a nucleotide sequence encoding an optimized influenza HA polypeptide, wherein the nucleotide sequence encoding the HA polypeptide is at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99% identical to SEQ ID NO: 1. Optimized influenza HA polypeptides encoded by the nucleic acid molecule, vectors comprising the nucleic acid molecule, and host cells containing the disclosed vectors are also provided herein.

Further provided is an optimized influenza HA polypeptide, wherein the amino acid sequence of the polypeptide is at least 99% identical to SEQ ID NO: 2. Also provided are fusion proteins comprising the optimized HA polypeptide, virus-like particles (VLPs) containing the optimized HA polypeptides, and compositions comprising the optimized HA polypeptide.

63198-1692

Collections of plasmids are also provided herein. In some embodiments, the collections of plasmids include a plasmid encoding an influenza NA, a plasmid encoding an influenza MA, and a plasmid encoding the optimized HA protein disclosed herein.

Further provided is a method of eliciting an immune response to influenza virus in a subject by administering the optimized influenza HA protein, fusion proteins containing the optimized influenza HA, or VLPs containing the optimized influenza HA, as disclosed herein. Also provided is a method of immunizing a subject against influenza virus by administering to the subject VLPs containing the optimized influenza HA protein disclosed herein.

10 In one embodiment, the invention provides an isolated nucleic acid molecule comprising a nucleotide sequence encoding an influenza hemagglutinin (HA) polypeptide, wherein the nucleotide sequence encoding the HA polypeptide is at least 94% identical to SEQ ID NO: 1.

15 In another embodiment, the invention provides an influenza HA polypeptide encoded by the nucleic acid molecule as described herein.

In another embodiment, the invention provides a vector comprising the isolated nucleic acid molecule as described herein.

20 In another embodiment, the invention provides an influenza HA polypeptide produced by transfecting a host cell with the vector as described herein under conditions sufficient to allow for expression of the polypeptide.

In another embodiment, the invention provides an isolated cell comprising the vector as described herein.

25 In another embodiment, the invention provides an isolated influenza HA polypeptide, wherein the amino acid sequence of the polypeptide is at least 99% identical to SEQ ID NO: 2.

In another embodiment, the invention provides a fusion protein comprising the polypeptide as described herein.

In another embodiment, the invention provides an influenza virus-like particle (VLP) comprising the polypeptide as described herein.

5 In another embodiment, the invention provides an influenza VLP comprising the influenza HA polypeptide as described herein, produced by transfecting a host cell with a vector encoding the HA polypeptide, a vector encoding an influenza NA protein and a vector encoding an influenza M1 protein under conditions sufficient to allow for expression of the HA, M1 and NA proteins.

10 In another embodiment, the invention provides a collection of plasmids comprising: (i) a plasmid encoding an influenza NA (ii) a plasmid encoding an influenza M1; and (iii) a plasmid encoding a codon-optimized influenza HA, wherein the nucleotide sequence encoding the codon-optimized influenza HA is at least 97% identical to SEQ ID NO: 1.

15 In another embodiment, the invention provides a composition comprising the influenza HA protein as described herein, the fusion protein as described herein, or the VLP as described herein, and a pharmaceutically acceptable carrier.

In another embodiment, the invention provides use of a composition comprising the influenza HA protein of as described herein, the fusion protein as described
20 herein, or the VLP as described herein, for eliciting an immune response to influenza virus in a subject.

In another embodiment, the invention provides use of the composition as described herein for eliciting an immune response to influenza virus in a subject.

In another embodiment, the invention provides use of a composition
25 comprising the VLP as described herein and pharmaceutically acceptable carrier for immunization of a subject against influenza virus.

In another embodiment, the invention provides a method of generating an optimized influenza virus polypeptide sequence, comprising: (i) obtaining the amino acid sequences of the polypeptide from a group of influenza virus isolates, wherein the influenza virus isolates are from the same subtype; (ii) organizing the amino acid sequences of the polypeptide from the group of influenza virus isolates by clade or sub-clade and then by geographical region within each clade or sub-clade; (iii) aligning the amino acid sequences within each geographical region to generate primary consensus sequences, wherein each geographic region is represented by a primary consensus sequence; (iv) aligning the primary consensus sequences to generate secondary consensus sequences, wherein each clade or sub-clade is represented by a secondary consensus sequence; and (v) producing the optimized influenza virus polypeptide generated by aligning the secondary consensus sequences.

The foregoing and other objects and features of the disclosure will become more apparent from the following detailed description, which proceeds with reference to the accompanying figures.

15 BRIEF DESCRIPTION OF THE FIGURES

FIGS. 1A-1B: COBRA HA Design. (A) Schematic illustrating the design of the COBRA HA molecule. The phylogenetic tree was inferred from hemagglutinin amino acid sequences using the maximum likelihood method and clade/sub-clade groupings were identified. Primary consensus sequences were generated for each outbreak group. Secondary consensus sequences were then generated for each sub-clade using the primary sequences as input. The secondary consensus sequences were then aligned and the resulting consensus, designated COBRA, was generated. (B) Phylogenetic analysis of the COBRA HA. The unrooted phylogenetic tree was inferred from hemagglutinin amino acid sequences from human H5N1 infections isolated from 2004 to 2009 and the clade/sub-clade groupings are indicated. The star represents the COBRA HA sequence relative to human H5N1 infections.

FIGS. 2A-2C: COBRA HA Functional Characterization. COBRA HA was translated *in vitro* and the cell culture lysates were analyzed by SDS-PAGE (A). Lane designations: 1) H5N1 recombinant HA; 2) COBRA HA; 3) Expression

vector; 4) H5N1 reassortant virus. The COBRA HA (lane 2) migrates at its expected molecular weight confirming expression of the synthetic protein. COBRA HA VLPs were prepared in various amounts, serially diluted, and incubated with 1% erythrocytes to evaluate receptor binding (B). HA titer was determined as the last well in which the RBCs remained suspended in a lattice structure. COBRA HA and control lentiviral pseudoparticles packaging a CMV-Luc gene were generated in HEK 293T cells and used to infect MDCK cells with or without trypsin (C). Particle fusion was determined by luciferase production by infected cells.

FIGS. 3A-3F: COBRA HA Mouse Dosing Immunogenicity. BALB/c mice (n=5/group) were vaccinated at 0 and 3 weeks with blood collected at 14 to 21 days after each vaccination. Vaccines were formulated at high (1.5 μg HA), and low (0.03 μg HA) doses, with and without Imject® alum, and delivered intramuscularly. Total IgG at week 5 was determined via ELISA for each vaccine group (A and B). Values represent the geometric mean titer ($\pm 95\%$ confidence interval) of \log_{10} transformed endpoint titers. IgG isotypes were evaluated via ELISA for each vaccine group (C and D). Values represent the mean OD_{450} of a 1:200 dilution of serum. Hemagglutination inhibition (HAI) serum antibody titer for each vaccine group was determined at week 5 using representative reassortant viruses (E and F). Values represent the geometric mean titer ($\pm 95\%$ confidence interval) of \log_2 transformed titers. The dotted line represents the 1:40 titer. Significant differences were determined by two-way ANOVA with Bonferroni's post-test to evaluate differences between the vaccine formulations for each test antigen. A *p* value of less than 0.05 was considered significant.

FIGS. 4A-4D: COBRA HA Mouse Dosing Efficacy. BALB/c mice (n=5/group) were vaccinated with COBRA HA VLPs with or without adjuvant. Mice were infected with 5×10^3 PFU of the highly pathogenic clade 2.2 H5N1 virus A/Whooper Swan/Mongolia/244/2005. Mice were followed to monitor weight loss (A and B) and sickness (C and D). Sickness score was determined by evaluating activity (0=normal, 1=reduced, 2=severely reduced), hunched back (0=absent, 1=present) and ruffled fur (0=absent, 1=present). All mock vaccinated mice reached the experimental endpoint and required euthanasia by 6 days post infection.

FIGS. 5A-5B: Mouse Comparison Immunogenicity. BALB/c mice (n=20/group) were vaccinated at 0 and 3 weeks with blood collected at 14 to 21 days after each vaccination. Vaccines were formulated at a high dose (3 µg HA) with Imject® alum and delivered intramuscularly. Total IgG at week 5 was determined via ELISA for each vaccine group (A). Values represent the geometric mean titer ($\pm 95\%$ confidence interval) of \log_{10} transformed endpoint titers. Hemagglutination inhibition (HAI) serum antibody titer for each vaccine group was determined at week 5 using representative reassortant viruses (B). Values represent the geometric mean titer ($\pm 95\%$ confidence interval) of \log_2 transformed titers. The dotted line represents the 1:40 titer. Significant differences were determined by two-way ANOVA with Bonferroni's post-test to evaluate differences between the vaccine formulations for each test antigen. A *p* value of less than 0.05 was considered significant.

FIGS. 6A-6B: Mouse Comparison Efficacy. BALB/c mice (n=20/group) were vaccinated with VLPs and adjuvant. Mice were infected with 5×10^3 PFU of the highly pathogenic clade 2.2 H5N1 virus A/Whooper Swan/Mongolia/244/2005. Mice were followed to monitor weight loss (A) and sickness (B). Sickness score was determined by evaluating activity (0=normal, 1=reduced, 2=severely reduced), hunched back (0=absent, 1=present) and ruffled fur (0=absent, 1=present). All mock (adjuvant-only) vaccinated mice reached the experimental endpoint and required euthanasia by 6 days post infection.

FIGS. 7A-7B: Ferret Immunogenicity. Ferrets (n=9/group) were vaccinated with VLPs (15 µg HA) with Imject® alum at weeks 0 and 3 and serum collected at week 5. Total IgG at week 5 was determined via ELISA for each vaccine group (A). Values represent the geometric mean titer ($\pm 95\%$ confidence interval) of \log_{10} transformed endpoint titers. Hemagglutination inhibition (HAI) serum antibody titer for each vaccine group was determined at week 5 using representative reassortant viruses (B). Values represent the geometric mean titer ($\pm 95\%$ confidence interval) of \log_2 transformed titers. The dotted line represents the 1:40 titer. Significant differences were determined by two-way ANOVA with

Bonferroni's post-test to evaluate differences between the vaccine formulations for each test antigen. A p value of less than 0.05 was considered significant.

FIGS. 8A-8E: Ferret Efficacy. Ferrets (n=9/group) were vaccinated with VLPs formulated with adjuvant. Ferrets were challenged with 1×10^6 PFU of the highly pathogenic clade 2.2 H5N1 virus A/Whooper Swan/Mongolia/244/2005. Animals were monitored daily for weight loss (A), survival (B), temperature (C) and clinical symptoms (D). Relative sickness scores were determined by measuring lethargy (0-3), runny nose (0-1), sneezing (0-1), loss of appetite (0-1) and diarrhea (0-1). Animals reaching experimental endpoint were euthanized according to institutional guidelines. Nasal washes were collected serially post infection and virus titers determined via plaque assay (E). Statistical significance was determined using a two-way ANOVA with Bonferroni's post test. A p value of less than 0.05 was considered significant.

FIG. 9: Phylogenetic diversity of H5N1 influenza. The unrooted phylogenetic tree was inferred from HA amino acid sequences derived from 8 to 10 representative isolates in all clades and sub-clades and the COBRA HA using the maximum likelihood method. Clade/sub-clade clusters were identified and are indicated in the shaded ovals. The star identifies where the COBRA antigen is located relative to the various representative isolates. Sequences were aligned with MUSCLE 3.7 software and the alignment was refined by Gblocks 0.91b software. Phylogeny was determined using the maximum likelihood method with PhyML software. Trees were rendered using TreeDyn 198.3 software (Dereeper *et al.*, *Nucleic Acids Res* 36:W465-W469, 2008). The NCBI accession numbers for the HA sequences used in phylogeny inference were obtained through the Influenza Virus Resource (Bao *et al.*, *J Virol* 82:596-601, 2008).

FIGS. 10A-10F: Serology. Total IgG at week 3 (A) and week 6 (B) was determined via ELISA for each vaccine group. Each collected antiserum was assayed for antibody binding to representative HA molecules from clade 2.1 (A/Indonesia/5/2005), clade 2.2 (A/Whooper Swan/Mongolia/244/2005), and clade 2.3 (A/Anhui/1/2005). Values represent the geometric mean titer ($\pm 95\%$ confidence interval) of \log_{10} transformed endpoint titers. Statistical significance of the antibody

titer data was determined using a two-way analysis of variance (ANOVA) followed by Bonferroni's post-test to analyze differences between each vaccine group for each of the different antigens that were tested (multiparametric). Significance was defined as $p < 0.05$. Statistical analyses were performed with GraphPad Prism software. HAI titer for each vaccine group was determined at week 3 (C) and week 6 (D) using representative H5N1 influenza viruses: clade 2.1 (A/Indonesia/5/2005), clade 2.2 (A/Whooper swan/Mongolia/244/2005), and clade 2.3 (A/Anhui/1/2005). Values represent the geometric mean titer ($\pm 95\%$ confidence interval) of \log_2 transformed titers. The dotted line represents the 1:40 titer. Significant differences were determined by two-way ANOVA with Bonferroni's post-test to evaluate differences between the vaccine formulations for each test antigen. A p value of less than 0.05 was considered significant. The number of monkeys that responded with a titer greater than 1:40 is listed above each bar. Neutralizing antibody at week 3 (E) and week 6 (F) was determined via microneutralization assay for each vaccine group. Values represent the geometric mean titer ($\pm 95\%$ confidence interval).

FIG. 11: HAI serum antibody titers from vaccinated monkeys against a panel of clade 0, 1, 2, 4, and 7 isolates. HAI titer for each vaccine group was determined at week 9 using H5N1 influenza viruses. Values represent the geometric mean titer ($\pm 95\%$ confidence interval) of \log_2 transformed titers. Significant differences were determined by two-way ANOVA with Bonferroni's post-test to evaluate differences between the vaccine formulations for each test antigen. A p value of less than 0.05 was considered significant as described in FIG. 10.

FIGS. 12A-12D: Vaccine induced serum antibody responses. BALB/c mice ($n=30$ /group) or Fitch ferrets ($n=6$ /group) were vaccinated at 0 and 3 weeks with blood collected 14 to 21 days after each vaccination. Total IgG after the second vaccination was determined via ELISA for each vaccine group (A and C). Receptor blocking antibody titers after the second vaccination were determined via hemagglutination inhibition (HAI) for each vaccine group (B and D). Values represent the geometric mean of the reciprocal dilution ($\pm 95\%$ confidence interval) of the last positive well. Significant differences between COBRA and

polyvalent vaccines were determined by a two-tailed Student's T test and a *p* value of less than 0.05 was considered significant (*).

FIGS. 13A-13D: Highly pathogenic Clade 2.2 challenge. Vaccinated BALB/c mice (n=5/group) were infected with 5×10^3 PFU of the highly pathogenic clade 2.2 H5N1 virus A/Whooper Swan/Mongolia/244/2005 (WS/05). Mice were monitored daily for weight loss (A) and sickness (B). Vaccinated Fitch ferrets (n=6/group) were infected with 1×10^6 PFU of the highly pathogenic clade WS/05 virus. Ferrets were monitored daily for weight loss (C) and sickness (D). Values represent mean (+/- SEM) for each group.

FIGS. 14A-14B: Clade 2.2 viral loads. Vaccinated BALB/c mice (n=15/group) were infected with 5×10^3 PFU of the highly pathogenic clade 2.2 H5N1 virus A/Whooper Swan/Mongolia/244/2005 (WS/05). Cohorts of mice (n=5/group) were sacrificed at 1, 3 and 5 days post infection, lungs harvested, and viral load determined by plaque assay (A). Vaccinated Fitch ferrets (n=6/group) were infected with 1×10^6 PFU of the highly pathogenic WS/05 virus. Nasal washes were collected and viral load determined by plaque assay (B). Values represent mean (+/- SEM) viral titer for each group. Significant differences between COBRA and polyvalent vaccines were determined by a two-tailed Student's T test and a *p* value of less than 0.05 was considered significant (*).

FIGS. 15A-15B: Histopathology of infected lungs. Vaccinated BALB/c mice (n=15/group) were infected with 5×10^3 PFU of the highly pathogenic clade 2.2 H5N1 virus A/Whooper Swan/Mongolia/244/2005 (WS/05). Cohorts of mice (n=5/group) were sacrificed at 3 days post infection and *in situ* hybridization (ISH) for influenza matrix protein (MP) was performed on sections from paraffin embedded lung tissue (A). Severity of influenza ISH foci was assessed in the bronchi (B). Scoring: 0 = no definitive signal; 1 = occasional focus; 2 = focus in most fields; 3 = more than one focus per field.

FIGS. 16A-16C: Clade 1 challenge. Vaccinated BALB/c mice (n=4/group) were infected with 5×10^3 PFU of reassortant virus containing the HA and NA genes from the clade 1 H5N1 virus A/Vietnam/1203/2004 (VN/04). Mice were monitored daily for weight loss (A) and sickness (B). Values represent mean

63198-1692

(+/- SD) for each group. An additional cohort of vaccinated mice (n=3/group) were infected and lungs were harvested 3 days post infection for analysis of viral burden (C). Values represent mean (+/- SEM) viral titer for each group.

FIGS. 17A-17B: Post-challenge cellular immune responses. Vaccinated BALB/c mice (n=3/group) were infected with 5×10^3 PFU of reassortant virus containing the HA and NA genes from the clade 1 H5N1 virus A/Vietnam/1203/2004 (VN/04). Mice were sacrificed 6 days post infection, lungs were harvested and the numbers of antibody secreting cells (A) and IFN- γ producing cells (B) were determined by ELISpot assay. Values represent the mean (+/- SEM) spots for each group.

FIGS. 18A-18B: Passive transfer clade 1 challenge. BALB/c mice (n=10/group) were vaccinated at 0 and 3 weeks with blood collected 14 to 21 days after each vaccination. Serum collected after the second vaccination was pooled for each vaccine group and administered to naïve recipient mice (n=5/group). One day after passive transfer, recipient mice were infected with 5×10^3 PFU of reassortant virus containing the HA and NA genes from the clade 1 H5N1 virus A/Vietnam/1203/2004 (VN/04). Mice were monitored daily for weight loss (A) and sickness (B). Values represent mean (+/- SD) for each group. Significant differences were determined by two-way ANOVA with Bonferroni's post-test to evaluate differences between vaccines at each day. A *p* value of less than 0.05 was considered significant (*).

SEQUENCE LISTING

The nucleic and amino acid sequences listed in the accompanying sequence listing are shown using standard letter abbreviations for nucleotide bases, and three letter code for amino acids. Only one strand of each nucleic acid sequence is shown, but the complementary strand is understood as included by any reference to the displayed strand.

In the accompanying sequence listing:

SEQ ID NOs: 1 and 2 are the nucleotide and amino acid sequences, respectively, of a codon-optimized influenza HA (designated COBRA).

SEQ ID NOS: 3 and 4 are the nucleotide and amino acid sequences, respectively, of a codon-optimized influenza NA.

SEQ ID NOS: 5 and 6 are the nucleotide and amino acid sequences, respectively, of a codon-optimized influenza M1.

SEQ ID NO: 7 is the nucleotide sequence of a plasmid encoding a codon-optimized influenza HA.

SEQ ID NO: 8 is the nucleotide sequence of a plasmid encoding a codon-optimized influenza NA.

SEQ ID NO: 9 is the nucleotide sequence of a plasmid encoding a codon-optimized influenza M1.

SEQ ID NO: 10 is the amino acid sequence of a T cell epitope in H5 HA (HA₅₃₃).

SEQ ID NO: 11 is the amino acid sequence of an ovalbumin T cell epitope (Ova₂₅₇).

DETAILED DESCRIPTION

I. Abbreviations

ASC: antibody secreting cell

DPI: days post infection

HA: hemagglutinin or hemagglutination assay

HAI: hemagglutination inhibition

hRBC: horse red blood cell

IFU: infectious unit

LD₅₀: lethal dose 50

M1: matrix protein 1

MN: microneutralization

MOI: multiplicity of infection

NA: neuraminidase

PFU: plaque form unit

RDE: receptor destroying enzyme

TCID: tissue culture infectious dose

tRBC: turkey red blood cell

VLP: virus-like particle

II. Terms and Methods

Unless otherwise noted, technical terms are used according to conventional usage. Definitions of common terms in molecular biology may be found in Benjamin Lewin, *Genes V*, published by Oxford University Press, 1994 (ISBN 0-19-854287-9); Kendrew *et al.* (eds.), *The Encyclopedia of Molecular Biology*, published by Blackwell Science Ltd., 1994 (ISBN 0-632-02182-9); and Robert A. Meyers (ed.), *Molecular Biology and Biotechnology: a Comprehensive Desk Reference*, published by VCH Publishers, Inc., 1995 (ISBN 1-56081-569-8).

In order to facilitate review of the various embodiments of the disclosure, the following explanations of specific terms are provided:

Adjuvant: A substance or vehicle that non-specifically enhances the immune response to an antigen. Adjuvants can include a suspension of minerals (alum, aluminum hydroxide, or phosphate) on which antigen is adsorbed; or water-in-oil emulsion in which antigen solution is emulsified in mineral oil (for example, Freund's incomplete adjuvant), sometimes with the inclusion of killed mycobacteria (Freund's complete adjuvant) to further enhance antigenicity. Immunostimulatory oligonucleotides (such as those including a CpG motif) can also be used as adjuvants (for example, see U.S. Patent Nos. 6,194,388; 6,207,646; 6,214,806; 6,218,371; 6,239,116; 6,339,068; 6,406,705; and 6,429,199). Adjuvants also include biological molecules, such as costimulatory molecules. Exemplary biological adjuvants include IL-2, RANTES, GM-CSF, TNF- α , IFN- γ , G-CSF, LFA-3, CD72, B7-1, B7-2, OX-40L and 41 BBL.

Administer: As used herein, administering a composition to a subject means to give, apply or bring the composition into contact with the subject. Administration can be accomplished by any of a number of routes, such as, for example, topical, oral, subcutaneous, intramuscular, intraperitoneal, intravenous, intrathecal and intramuscular.

Antibody: An immunoglobulin molecule produced by B lymphoid cells with a specific amino acid sequence. Antibodies are evoked in humans or other animals by a specific antigen (immunogen). Antibodies are characterized by reacting specifically with the antigen in some demonstrable way, antibody and antigen each being defined in terms of the other. “Eliciting an antibody response” refers to the ability of an antigen or other molecule to induce the production of antibodies.

Antigen: A compound, composition, or substance that can stimulate the production of antibodies or a T-cell response in an animal, including compositions that are injected or absorbed into an animal. An antigen reacts with the products of specific humoral or cellular immunity, including those induced by heterologous immunogens. In some embodiments of the disclosed compositions and methods, the antigen is an influenza HA protein.

Attenuated: In the context of a live virus, the virus is attenuated if its ability to infect a cell or subject and/or its ability to produce disease is reduced (for example, eliminated) compared to a wild-type virus. Typically, an attenuated virus retains at least some capacity to elicit an immune response following administration to an immunocompetent subject. In some cases, an attenuated virus is capable of eliciting a protective immune response without causing any signs or symptoms of infection. In some embodiments, the ability of an attenuated virus to cause disease in a subject is reduced at least about 10%, at least about 25%, at least about 50%, at least about 75% or at least about 90% relative to wild-type virus.

Clade: Refers to the different categorizations of the known influenza viruses, such as influenza A H5N1 viruses. Viruses in an H5N1 clade are genetically related, but do not share the exact viral genome. There are at least ten different clades of H5N1 subtypes designated in the art: clade 0 clade 1, clade 2, clade 3, clade 4, clade 5, clade 6, clade 7, clade 8 and clade 9 (Abdel-Ghafar *et al.*, *N Engl J Med* 358:261-273, 2008). Clade 2 is further divided into sub-clades (including clade 2.1, clade 2.2, clade 2.3, clade 2.4 and clade 2.5).

Codon-optimized: A “codon-optimized” nucleic acid refers to a nucleic acid sequence that has been altered such that the codons are optimal for expression

in a particular system (such as a particular species or group of species). For example, a nucleic acid sequence can be optimized for expression in mammalian cells. Codon optimization does not alter the amino acid sequence of the encoded protein.

Fusion protein: A protein generated by expression of a nucleic acid sequence engineered from nucleic acid sequences encoding at least a portion of two different (heterologous) proteins. To create a fusion protein, the nucleic acid sequences must be in the same reading frame and contain no internal stop codons. For example, a fusion protein includes an influenza HA fused to a heterologous protein.

Geographical location or geographical region: Refers to preselected divisions of geographical areas of the earth, for example, by continent or other preselected territory or subdivision (*e.g.*, the Middle East, which spans more than one continent). Examples of different geographical regions include countries (*e.g.*, Turkey, Egypt, Iraq, Azerbaijan, China, United States), continents (*e.g.*, Asia, Europe, North America, South America, Oceania, Africa), and recognized geopolitical subdivisions (such as the Middle East).

Hemagglutinin (HA): An influenza virus surface glycoprotein. HA mediates binding of the virus particle to a host cell and subsequent entry of the virus into the host cell. The nucleotide and amino acid sequences of numerous influenza HA proteins are known in the art and are publically available, such as those deposited with GenBank (see Table 1 for a list of GenBank Accession Nos. of H5N1 HA sequences). HA (along with NA) is one of the two major influenza virus antigenic determinants.

Immune response: A response of a cell of the immune system, such as a B-cell, T-cell, macrophage or polymorphonucleocyte, to a stimulus such as an antigen or vaccine. An immune response can include any cell of the body involved in a host defense response, including for example, an epithelial cell that secretes an interferon or a cytokine. An immune response includes, but is not limited to, an innate immune response or inflammation. As used herein, a protective immune response refers to an immune response that protects a subject from infection (prevents

infection or prevents the development of disease associated with infection).

Methods of measuring immune responses are well known in the art and include, for example, measuring proliferation and/or activity of lymphocytes (such as B or T cells), secretion of cytokines or chemokines, inflammation, antibody production and the like.

Immunogen: A compound, composition, or substance which is capable, under appropriate conditions, of stimulating an immune response, such as the production of antibodies or a T-cell response in an animal, including compositions that are injected or absorbed into an animal. As used herein, as “immunogenic composition” is a composition comprising an immunogen (such as an HA polypeptide).

Immunize: To render a subject protected from an infectious disease, such as by vaccination.

Influenza virus: A segmented negative-strand RNA virus that belongs to the *Orthomyxoviridae* family. There are three types of Influenza viruses, A, B and C. Influenza A viruses infect a wide variety of birds and mammals, including humans, horses, marine mammals, pigs, ferrets, and chickens. In animals, most influenza A viruses cause mild localized infections of the respiratory and intestinal tract. However, highly pathogenic influenza A strains, such as H5N1, cause systemic infections in poultry in which mortality may reach 100%. H5N1 is also referred to as "avian influenza."

Isolated: An “isolated” biological component (such as a nucleic acid, protein or virus) has been substantially separated or purified away from other biological components (such as cell debris, or other proteins or nucleic acids). Biological components that have been “isolated” include those components purified by standard purification methods. The term also embraces recombinant nucleic acids, proteins or viruses, as well as chemically synthesized nucleic acids or peptides.

Linker: One or more amino acids that serve as a spacer between two polypeptides of a fusion protein.

Matrix (M1) protein: An influenza virus structural protein found within the viral envelope. M1 is thought to function in assembly and budding.

Neuraminidase (NA): An influenza virus membrane glycoprotein. NA is involved in the destruction of the cellular receptor for the viral HA by cleaving terminal sialic acid residues from carbohydrate moieties on the surfaces of infected cells. NA also cleaves sialic acid residues from viral proteins, preventing aggregation of viruses. NA (along with HA) is one of the two major influenza virus antigenic determinants.

Operably linked: A first nucleic acid sequence is operably linked with a second nucleic acid sequence when the first nucleic acid sequence is placed in a functional relationship with the second nucleic acid sequence. For instance, a promoter is operably linked to a coding sequence if the promoter affects the transcription or expression of the coding sequence. Generally, operably linked DNA sequences are contiguous and, where necessary to join two protein-coding regions, in the same reading frame.

Optimized influenza HA protein: As used herein, “optimized influenza HA protein” refers to the HA protein consensus sequence generated by sequence alignments of clade 2 H5N1 influenza viruses (as described in Example 1 below). The nucleotide sequence encoding the optimized HA protein was further optimized for expression in mammalian cells via codon-optimization and RNA optimization (such as to increase RNA stability). The optimized influenza HA protein disclosed herein (and set forth herein as SEQ ID NO: 2) is also referred to as “COBRA.”

ORF (open reading frame): A series of nucleotide triplets (codons) coding for amino acids without any termination codons. These sequences are usually translatable into a peptide.

Outbreak: As used herein, an influenza virus “outbreak” refers to a collection of virus isolates from within a single country in a given year.

Pharmaceutically acceptable vehicles: The pharmaceutically acceptable carriers (vehicles) useful in this disclosure are conventional. *Remington's Pharmaceutical Sciences*, by E. W. Martin, Mack Publishing Co., Easton, PA, 15th Edition (1975), describes compositions and formulations suitable for pharmaceutical delivery of one or more therapeutic compositions, such as one or more influenza vaccines, and additional pharmaceutical agents.

In general, the nature of the carrier will depend on the particular mode of administration being employed. For instance, parenteral formulations usually comprise injectable fluids that include pharmaceutically and physiologically acceptable fluids such as water, physiological saline, balanced salt solutions, aqueous dextrose, glycerol or the like as a vehicle. For solid compositions (for example, powder, pill, tablet, or capsule forms), conventional non-toxic solid carriers can include, for example, pharmaceutical grades of mannitol, lactose, starch, or magnesium stearate. In addition to biologically-neutral carriers, pharmaceutical compositions to be administered can contain minor amounts of non-toxic auxiliary substances, such as wetting or emulsifying agents, preservatives, and pH buffering agents and the like, for example sodium acetate or sorbitan monolaurate.

Plasmid: A circular nucleic acid molecule capable of autonomous replication in a host cell.

Polypeptide: A polymer in which the monomers are amino acid residues which are joined together through amide bonds. When the amino acids are alpha-amino acids, either the L-optical isomer or the D-optical isomer can be used. The terms “polypeptide” or “protein” as used herein are intended to encompass any amino acid sequence and include modified sequences such as glycoproteins. The term “polypeptide” is specifically intended to cover naturally occurring proteins, as well as those which are recombinantly or synthetically produced. The term “residue” or “amino acid residue” includes reference to an amino acid that is incorporated into a protein, polypeptide, or peptide.

Conservative amino acid substitutions are those substitutions that, when made, least interfere with the properties of the original protein, that is, the structure and especially the function of the protein is conserved and not significantly changed by such substitutions. Examples of conservative substitutions are shown below.

Original Residue	Conservative Substitutions
Ala	Ser
Arg	Lys
Asn	Gln, His
Asp	Glu

Cys	Ser
Gln	Asn
Glu	Asp
His	Asn; Gln
Ile	Leu, Val
Leu	Ile; Val
Lys	Arg; Gln; Glu
Met	Leu; Ile
Phe	Met; Leu; Tyr
Ser	Thr
Thr	Ser
Trp	Tyr
Tyr	Trp; Phe
Val	Ile; Leu

Conservative substitutions generally maintain (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain.

The substitutions which in general are expected to produce the greatest changes in protein properties will be non-conservative, for instance changes in which (a) a hydrophilic residue, for example, seryl or threonyl, is substituted for (or by) a hydrophobic residue, for example, leucyl, isoleucyl, phenylalanyl, valyl or alanyl; (b) a cysteine or proline is substituted for (or by) any other residue; (c) a residue having an electropositive side chain, for example, lysyl, arginyl, or histadyl, is substituted for (or by) an electronegative residue, for example, glutamyl or aspartyl; or (d) a residue having a bulky side chain, for example, phenylalanine, is substituted for (or by) one not having a side chain, for example, glycine.

Preventing, treating or ameliorating a disease: “Preventing” a disease refers to inhibiting the full development of a disease. “Treating” refers to a therapeutic intervention that ameliorates a sign or symptom of a disease or pathological condition after it has begun to develop. “Ameliorating” refers to the reduction in the number or severity of signs or symptoms of a disease.

Promoter: An array of nucleic acid control sequences which direct transcription of a nucleic acid. A promoter includes necessary nucleic acid sequences near the start site of transcription. A promoter also optionally includes

distal enhancer or repressor elements. A “constitutive promoter” is a promoter that is continuously active and is not subject to regulation by external signals or molecules. In contrast, the activity of an “inducible promoter” is regulated by an external signal or molecule (for example, a transcription factor). In some embodiments herein, the promoter is a CMV promoter.

Purified: The term “purified” does not require absolute purity; rather, it is intended as a relative term. Thus, for example, a purified peptide, protein, virus, or other active compound is one that is isolated in whole or in part from naturally associated proteins and other contaminants. In certain embodiments, the term “substantially purified” refers to a peptide, protein, virus or other active compound that has been isolated from a cell, cell culture medium, or other crude preparation and subjected to fractionation to remove various components of the initial preparation, such as proteins, cellular debris, and other components.

Recombinant: A recombinant nucleic acid, protein or virus is one that has a sequence that is not naturally occurring or has a sequence that is made by an artificial combination of two otherwise separated segments of sequence. This artificial combination is often accomplished by chemical synthesis or by the artificial manipulation of isolated segments of nucleic acids, for example, by genetic engineering techniques.

Sequence identity: The similarity between amino acid or nucleic acid sequences is expressed in terms of the similarity between the sequences, otherwise referred to as sequence identity. Sequence identity is frequently measured in terms of percentage identity (or similarity or homology); the higher the percentage, the more similar the two sequences are. Homologs or variants of a given gene or protein will possess a relatively high degree of sequence identity when aligned using standard methods.

Methods of alignment of sequences for comparison are well known in the art. Various programs and alignment algorithms are described in: Smith and Waterman, *Adv. Appl. Math.* 2:482, 1981; Needleman and Wunsch, *J. Mol. Biol.* 48:443, 1970; Pearson and Lipman, *Proc. Natl. Acad. Sci. U.S.A.* 85:2444, 1988; Higgins and Sharp, *Gene* 73:237-244, 1988; Higgins and Sharp, *CABIOS* 5:151-153, 1989; Corpet *et al.*, *Nucleic Acids Research* 16:10881-10890, 1988; and Pearson and Lipman, *Proc. Natl. Acad. Sci. U.S.A.* 85:2444, 1988. Altschul *et al.*, *Nature Genet.* 6:119-129, 1994.

The NCBI Basic Local Alignment Search Tool (BLASTTM) (Altschul *et al.*, *J. Mol. Biol.* 215:403-410, 1990) is available from several sources, including the National Center for Biotechnology Information (NCBI, Bethesda, MD) and on the Internet, for use in connection with the sequence analysis programs blastp, blastn, blastx, tblastn and tblastx.

Subject: Living multi-cellular vertebrate organisms, a category that includes both human and non-human mammals, such as non-human primates. In one example, a subject is one who is infected with H5N1 or is susceptible to such infection.

Therapeutically effective amount: A quantity of a specified agent sufficient to achieve a desired effect in a subject being treated with that agent. For example, this may be the amount of an influenza virus vaccine useful for eliciting an immune response in a subject and/or for preventing infection by influenza virus. Ideally, in the context of the present disclosure, a therapeutically effective amount of an influenza vaccine is an amount sufficient to increase resistance to, prevent, ameliorate, and/or treat infection caused by influenza virus in a subject without causing a substantial cytotoxic effect in the subject. The effective amount of an influenza vaccine useful for increasing resistance to, preventing, ameliorating, and/or treating infection in a subject will be dependent on, for example, the subject being treated, the manner of administration of the therapeutic composition and other factors.

Transformed: A transformed cell is a cell into which has been introduced a nucleic acid molecule by molecular biology techniques. As used herein, the term transformation encompasses all techniques by which a nucleic acid molecule might be introduced into such a cell, including transfection with viral vectors, transformation with plasmid vectors, and introduction of naked DNA by electroporation, lipofection, and particle gun acceleration.

Vaccine: A preparation of immunogenic material capable of stimulating an immune response, administered for the prevention, amelioration, or treatment of disease, such as an infectious disease. The immunogenic material may include, for example, attenuated or killed microorganisms (such as attenuated viruses), or antigenic proteins, peptides or DNA derived from them. Vaccines may elicit both

prophylactic (preventative) and therapeutic responses. Methods of administration vary according to the vaccine, but may include inoculation, ingestion, inhalation or other forms of administration. Inoculations can be delivered by any of a number of routes, including parenteral, such as intravenous, subcutaneous or intramuscular. Vaccines may be administered with an adjuvant to boost the immune response.

Vector: A vector is a nucleic acid molecule allowing insertion of foreign nucleic acid without disrupting the ability of the vector to replicate and/or integrate in a host cell. A vector can include nucleic acid sequences that permit it to replicate in a host cell, such as an origin of replication. An insertional vector is capable of inserting itself into a host nucleic acid. A vector can also include one or more selectable marker genes and other genetic elements. An expression vector is a vector that contains the necessary regulatory sequences to allow transcription and translation of inserted gene or genes. In some embodiments of the present disclosure, the vector encodes an influenza HA, NA or M1 protein. In some embodiments, the vector is the pTR600 expression vector (U.S. Patent Application Publication No. 2002/0106798; Ross *et al.*, *Nat Immunol.* 1(2):102-103, 2000; Green *et al.*, *Vaccine* 20:242-248, 2001).

Virus-like particle (VLP): Virus particles made up of one or more viral structural proteins, but lacking the viral genome. Because VLPs lack a viral genome, they are non-infectious. In addition, VLPs can often be produced by heterologous expression and can be easily purified. Most VLPs comprise at least a viral core protein that drives budding and release of particles from a host cell. One example of such a core protein is influenza M1. In some embodiments herein, an influenza VLP comprises the HA, NA and M1 proteins. As described herein, influenza VLPs can be produced by transfection of host cells with plasmids encoding the HA, NA and M1 proteins. After incubation of the transfected cells for an appropriate time to allow for protein expression (such as for approximately 72 hours), VLPs can be isolated from cell culture supernatants. Example 1 provides an exemplary protocol for purifying influenza VLPs from cell supernatants. In this example, VLPs are isolated by low speed centrifugation (to remove cell debris), vacuum filtration and ultracentrifugation through 20% glycerol.

63198-1692

Unless otherwise explained, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this disclosure belongs. The singular terms “a,” “an,” and “the” include plural referents unless context clearly indicates otherwise. Similarly, the word “or” is intended to include “and” unless the context clearly indicates otherwise. Hence “comprising A or B” means including A, or B, or A and B. It is further to be understood that all base sizes or amino acid sizes, and all molecular weight or molecular mass values, given for nucleic acids or polypeptides are approximate, and are provided for description. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present disclosure, suitable methods and materials are described below. In addition, the materials, methods, and examples are illustrative only and not intended to be limiting.

III. Overview of Several Embodiments

Disclosed herein is the development of a computationally optimized influenza HA protein that elicits broadly reactive immune responses to H5N1 influenza virus isolates, such as the isolates listed in Table 1. The optimized HA protein was developed through a series of HA protein alignments, and subsequent generation of consensus sequences, for clade 2 H5N1 influenza virus isolates (described in detail in Example 1 below; see also FIG. 1). The final consensus HA amino acid sequence was reverse translated and optimized for expression in mammalian cells. Optimization of the nucleic acid sequence included optimization of the codons for expression in mammalian cells and RNA optimization (such as RNA stability). The optimized HA coding sequence is set forth herein as SEQ ID NO: 1, and the optimized HA protein sequence is set forth herein as SEQ ID NO: 2.

Thus, provided herein is an isolated nucleic acid molecule comprising a nucleotide sequence encoding an influenza HA polypeptide. In some embodiments, the nucleotide sequence encoding the HA polypeptide is at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99% identical to SEQ ID NO: 1.

In some examples, the nucleotide sequence encoding the influenza HA polypeptide that is at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% identical to SEQ ID NO: 1 lacks the start codon (nucleotides 1-3 of SEQ ID NO: 1), encoding a N-terminal methionine. In particular examples, the nucleotide sequence encoding the influenza HA polypeptide is at least 94% identical to nucleotides 4-1707 of SEQ ID NO: 1. In other examples, the nucleotide sequence encoding the HA polypeptide comprises or consists of nucleotides 4-1707 of SEQ ID NO: 1.

In some examples, the nucleotide sequence encoding the HA polypeptide comprises SEQ ID NO: 1. In particular examples, the nucleotide sequence encoding the HA polypeptide consists of SEQ ID NO: 1. Also provided herein are influenza HA polypeptides encoded by the disclosed nucleic acid molecules.

Further provided are vectors containing a nucleotide sequence encoding an optimized HA polypeptide. In some embodiments of the vectors provided herein, the nucleotide sequence encoding the HA polypeptide is at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99% identical to SEQ ID NO: 1. In some examples, the vector further includes a promoter operably linked to the nucleotide sequence encoding the HA polypeptide. In particular examples, the promoter is a cytomegalovirus (CMV) promoter. In some embodiments, the nucleotide sequence of the vector is at least 85%, at least 90%, at least 95%, at least 98% or at least 99% identical to the nucleotide sequence of SEQ ID NO: 7. In some examples, the nucleotide sequence of the vector comprises the nucleotide sequence of SEQ ID NO: 7. In particular examples, the nucleotide sequence of the vector consists of the nucleotide sequence of SEQ ID NO: 7.

Also provided herein are influenza HA polypeptides produced by transfecting a host cell with a vector provided herein under conditions sufficient to

allow for expression of the HA polypeptide. Isolated cells containing the disclosed vectors are also provided.

Also provided herein are optimized influenza HA polypeptides. In some embodiments, the amino acid sequence of the polypeptide is at least 99% identical to SEQ ID NO: 2. In some examples, the amino acid sequence of the influenza HA polypeptide that is at least 99% identical to SEQ ID NO: 2 lacks the N-terminal methionine residue. In particular examples, the amino acid sequence of the influenza HA polypeptide is at least 99% identical to amino acids 2-568 of SEQ ID NO: 2. In other examples, the amino acid sequence of the HA polypeptides comprises or consists of amino acids 2-568 of SEQ ID NO: 2.

In some examples, the amino acid sequence of the polypeptide comprises the amino acid sequence of SEQ ID NO: 2. In particular examples, the amino acid sequence of the polypeptide consists of the amino acid sequence of SEQ ID NO: 2. Fusion proteins comprising the influenza HA polypeptides disclosed herein are also provided. The influenza HA polypeptide can be fused to any heterologous amino acid sequence to form the fusion protein.

Further provided herein are influenza virus-like particles (VLPs) containing an optimized influenza HA protein disclosed herein. In some embodiments, the HA protein of the VLP is at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% or 100% identical to SEQ ID NO: 2. The influenza VLPs can further include any additional influenza proteins necessary to form the virus particle. In some embodiments, the influenza VLPs further include influenza neuraminidase (NA) protein, influenza matrix (M1) protein, or both.

In some embodiments of the influenza VLPs, the amino acid sequence of the influenza NA protein is at least 85%, at least 90%, at least 95%, at least 98% or at least 99% identical to SEQ ID NO: 4. In some examples, the amino acid sequence of the influenza NA protein comprises SEQ ID NO: 4. In particular examples, the amino acid sequence of the influenza NA protein consists of SEQ ID NO: 4. In some embodiments, the amino acid sequence of the influenza M1 protein is at least 85%, at least 90%, at least 95%, at least 98% or at least 99% identical to SEQ ID NO: 6. In some examples, the amino acid sequence of the influenza M1 protein

63198-1692

comprises SEQ ID NO: 6. In particular examples, the amino acid sequence of the influenza M1 protein consists of SEQ ID NO: 6.

Also provided is an influenza VLP containing an influenza HA polypeptide as described herein, produced by transfecting a host cell with a vector encoding the HA polypeptide, a vector encoding an influenza NA protein and a vector encoding an influenza M1 protein, under conditions sufficient to allow for expression of the HA, M1 and NA proteins.

The vectors used to express the HA, NA and M1 proteins can be any suitable expression vectors known in the art. The vectors can be, for example, mammalian expression vectors, or viral vectors. In some embodiments, the vector is the pTR600 expression vector (U.S. Patent Application Publication No. 2002/0106798; Ross *et al.*, *Nat Immunol.* 1(2):102-103, 2000; Green *et al.*, *Vaccine* 20:242-248, 2001).

In some embodiments, the nucleotide sequence of the vector encoding the HA polypeptide is at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99% identical to SEQ ID NO: 7. In some examples, the nucleotide sequence of the vector encoding the HA polypeptide comprises SEQ ID NO: 7. In particular examples, the nucleotide sequence of the vector encoding the HA polypeptide consists of SEQ ID NO: 7.

In some embodiments, the nucleotide sequence of the vector encoding the NA protein is at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99% identical to SEQ ID NO: 8. In some examples, the nucleotide sequence of the vector encoding the NA protein comprises SEQ ID NO: 8. In particular examples, the nucleotide sequence of the vector encoding the NA protein consists of SEQ ID NO: 8.

In some embodiments, the nucleotide sequence of the vector encoding the M1 protein is at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99% identical to SEQ ID NO: 9. In some examples, the nucleotide sequence of the vector encoding the M1 protein comprises SEQ ID NO: 9. In particular examples, the nucleotide sequence of the vector encoding the M1 protein consists of SEQ ID NO: 9.

Collections of plasmids are also provided herein. In some embodiments, the collection of plasmids includes a plasmid encoding an influenza NA, a plasmid encoding an influenza MA, and a plasmid encoding the optimized HA protein disclosed herein. In some embodiments, the nucleotide sequence encoding the codon-optimized influenza HA of the HA-encoding plasmid is at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99% identical to SEQ ID NO: 1. Also provided are kits comprising the collection of plasmids.

In some embodiments of the collections of plasmids, the influenza NA is codon-optimized and/or the influenza M1 is codon-optimized. In some examples, the nucleotide sequence encoding the codon-optimized influenza NA is at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99% identical to SEQ ID NO: 3. In particular examples, the nucleotide sequence encoding the codon-optimized influenza NA comprises, or consists of, SEQ ID NO: 3. In some examples, the nucleotide sequence encoding the codon-optimized influenza M1 is at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99% identical to SEQ ID NO: 5. In particular examples, the nucleotide sequence encoding the codon-optimized influenza M1 comprises, or consists of, SEQ ID NO: 5.

In one non-limiting example, the plasmid encoding influenza NA comprises SEQ ID NO: 8; the plasmid encoding influenza M1 comprises SEQ ID NO: 9; and the plasmid encoding influenza HA comprises SEQ ID NO: 10.

In some embodiments, transfection of the collection of plasmids into host cells under conditions sufficient to allow for expression of the HA, NA and M1 proteins produces influenza VLPs comprising the HA, NA and M1 proteins.

Also provided herein are compositions comprising an optimized influenza HA protein as disclosed herein, or a fusion protein or VLP comprising the optimized influenza HA protein. In some embodiments, the compositions further comprise a pharmaceutically acceptable carrier and/or an adjuvant. For example, the adjuvant can be alum, Freund's complete adjuvant, a biological adjuvant or immunostimulatory oligonucleotides (such as CpG oligonucleotides).

Further provided is a method of eliciting an immune response to influenza virus in a subject by administering an influenza HA protein disclosed herein, fusion proteins containing the influenza HA, or VLPs containing the influenza HA, as disclosed herein. In some embodiments, the influenza virus is an H5N1 influenza virus. In some embodiments, the HA protein, HA fusion protein or VLP can be administered using any suitable route of administration, such as, for example, intramuscular. In some embodiments, the HA protein, fusion protein or VLP is administered as a composition further comprising a pharmaceutically acceptable carrier and/or an adjuvant. For example, the adjuvant can be alum, Freund's complete adjuvant, a biological adjuvant or immunostimulatory oligonucleotides (such as CpG oligonucleotides).

Also provided is a method of immunizing a subject against influenza virus by administering to the subject VLPs containing the optimized influenza HA protein disclosed herein, or administering a composition thereof. In some embodiments of the method, the composition further comprises a pharmaceutically acceptable carrier and/or an adjuvant. For example, the adjuvant can be alum, Freund's complete adjuvant, a biological adjuvant or immunostimulatory oligonucleotides (such as CpG oligonucleotides). In some embodiments, the VLPs (or compositions thereof) are administered intramuscularly.

In some embodiments of the methods of eliciting an immune response or immunizing a subject, the subject is administered at least 1 μg of the VLPs containing the optimized HA protein, such as at least 5 μg , at least 10 μg , at least 15 μg , at least 20 μg , at least 25 μg , at least 30 μg , at least 40 μg or at least 50 μg of the VLPs containing the optimized HA protein, for example about 1 to about 50 μg or about 1 to about 25 μg of the VLPs containing the optimized HA protein. In particular examples, the subject is administered about 5 to about 20 μg of the VLPs, or about 10 to about 15 μg of the VLPs. In one specific non-limiting example, the subject is administered about 15 μg of the VLPs. However, one of skill in the art is capable of determining a therapeutically effective amount (for example an amount

that provides protection against H5N1 influenza virus infection) of VLPs to administer to a subject.

It is disclosed herein that administration of VLPs comprising the COBRA HA disclosed herein (SEQ ID NO: 2) elicits protective levels of HAI antibodies to a number of representative clade 2 isolates and provides complete protection against lethal challenge with a clade 2.2 H5N1 virus. In some embodiments, administration of VLPs containing an optimized influenza HA results in production of high HAI titers ($\geq 1:40$) to H5N1 clade 1, clade 2.1, clade 2.2 and clade 2.3 isolates. In some examples, the VLPs containing an optimized influenza HA elicit high HAI titers against clade 1 and/or clade 7 viruses. The VLPs containing an optimized influenza HA disclosed herein elicit a broader immune response (*e.g.*, elicit neutralizing antibodies to a broader range of H5N1 virus isolates, such as those from clade 1, sub-clades of clade 2, and clade 7) than a polyvalent influenza virus vaccine.

Also provided herein is a method of optimizing an influenza protein sequence to elicit broadly reactive immune responses in a subject. In the context of the present disclosure, "broadly reactive" means the protein sequence elicits an immune response in a subject that is sufficient to inhibit, neutralize or prevent infection of a broad range of influenza viruses (such as most or all influenza viruses within a specific subtype). In some embodiments, the influenza protein is influenza HA or influenza NA. In one example, the optimized influenza protein is capable of eliciting a protective immune response against most or all known H5N1 influenza virus isolates (such as those listed in Table 1), such as about 80%, about 85%, about 90% or about 95% of known H5N1 influenza virus isolates.

In some embodiments, the method of optimizing an influenza protein sequence includes obtaining the amino acid sequences of a group of influenza virus isolates. For example, the group can consist of influenza virus isolates from a specific subtype (such as, for example, H5N1 or H1N1), and/or from one or more clades/sub-clades of a specific influenza subtype (for example, one or more of clades/sub-clades 0, 1, 2.1, 2.2, 2.3, 2.4, 3, 4, 5, 6, 7, 8 and 9 of H5N1). Amino acid sequences of the group of influenza viruses are first organized by clade or sub-clade and then by geographic location within each clade or sub-clade. The amino acid

sequences for each geographic location are aligned to generate a primary consensus sequence for each geographical region. Grouping virus isolates by geographical region controls for single outbreak dominance and incomplete reporting and sequencing. The primary consensus sequence can be generated, for example, by multiple alignment analysis using AlignX (Vector NTI), or by any other method known in the art. The primary geographically-based consensus sequences for each clade or sub-clade are then aligned, and a secondary consensus sequence is generated for each clade or sub-clade. The secondary consensus sequences for each clade or sub-clade are then aligned to generate the optimized, broadly reactive, consensus sequence (see FIG. 1). In some embodiments, the optimized influenza virus polypeptide sequence is further optimized for expression in mammalian cells. In some examples, optimization includes reverse translation of the optimized influenza virus polypeptide sequence to generate a coding sequence, followed by codon-optimization and/or optimization of the RNA (such as for stability).

In one non-limiting example, the method of optimization includes: (i) obtaining the amino acid sequences of the polypeptide from a group of influenza virus isolates, wherein the influenza virus isolates are from the same subtype; (ii) organizing the amino acid sequences of the polypeptide from the group of influenza virus isolates by clade or sub-clade and then by geographical region within each clade or sub-clade; (iii) aligning the amino acid sequences within each geographical region to generate primary consensus sequences, wherein each geographic region is represented by a primary consensus sequence; (iv) aligning the primary consensus sequences to generate secondary consensus sequences, wherein each clade or sub-clade is represented by a secondary consensus sequence; and (v) aligning the secondary consensus sequences to generate the optimized influenza virus polypeptide sequence. In some cases, the method further includes (i) reverse translating the optimized influenza virus polypeptide sequence to generate a coding sequence; and (ii) optimizing the coding sequence for expression in mammalian cells.

In an alternative embodiment, the primary consensus sequence is obtained by aligning influenza protein sequences (such as HA or NA sequences) from viral

isolates from a single outbreak (a collection of influenza virus isolates within a single country within a given year). Thus, in one non-limiting example, the method of optimization includes: (i) obtaining the amino acid sequences of the polypeptide from a group of influenza virus isolates, wherein the influenza virus isolates are from the same subtype; (ii) organizing the amino acid sequences of the polypeptide from the group of influenza virus isolates by clade or sub-clade and then by outbreak; (iii) aligning the amino acid sequences within each outbreak to generate primary consensus sequences, wherein each outbreak is represented by a primary consensus sequence; (iv) aligning the primary consensus sequences to generate secondary consensus sequences, wherein each clade or sub-clade is represented by a secondary consensus sequence; and (v) aligning the secondary consensus sequences to generate the optimized influenza virus polypeptide sequence. In some cases, the method further includes (i) reverse translating the optimized influenza virus polypeptide sequence to generate a coding sequence; and (ii) optimizing the coding sequence for expression in mammalian cells.

VI. Influenza

Influenza viruses are segmented negative-strand RNA viruses that belong to the *Orthomyxoviridae* family. There are three types of Influenza viruses, A, B and C. Influenza A viruses infect a wide variety of birds and mammals, including humans, horses, marine mammals, pigs, ferrets, and chickens. In animals, most influenza A viruses cause mild localized infections of the respiratory and intestinal tract. However, highly pathogenic influenza A strains, such as H5N1, cause systemic infections in poultry in which mortality may reach 100%. Animals infected with influenza A often act as a reservoir for the influenza viruses and certain subtypes have been shown to cross the species barrier to humans.

Influenza A viruses can be classified into subtypes based on allelic variations in antigenic regions of two genes that encode surface glycoproteins, namely, hemagglutinin (HA) and neuraminidase (NA) which are required for viral attachment and cellular release. Currently, sixteen subtypes of HA (H1-H16) and nine NA (N1-N9) antigenic variants are known for influenza A virus. Previously,

only three subtypes were known to circulate in humans (H1N1, H1N2, and H3N2). However, in recent years, the pathogenic H5N1 subtype of avian influenza A has been reported to cross the species barrier and infect humans as documented in Hong Kong in 1997 and 2003, leading to the death of several patients.

In humans, the avian influenza virus infects cells of the respiratory tract as well as the intestinal tract, liver, spleen, kidneys and other organs. Symptoms of avian influenza infection include fever, respiratory difficulties including shortness of breath and cough, lymphopenia, diarrhea and difficulties regulating blood sugar levels. In contrast to seasonal influenza, the group most at risk is healthy adults which make up the bulk of the population. Due to the high pathogenicity of certain avian influenza A subtypes, particularly H5N1, and their demonstrated ability to cross over to infect humans, there is a significant economic and public health risk associated with these viral strains, including a real epidemic and pandemic threat. Currently, no effective vaccines for H5N1 infection are available.

The influenza A virus genome encodes nine structural proteins and one nonstructural (NS1) protein with regulatory functions. The influenza virus segmented genome contains eight negative-sense RNA (nsRNA) gene segments (PB2, PB1, PA, NP, M, NS, HA and NA) that encode at least ten polypeptides, including RNA-directed RNA polymerase proteins (PB2, PB1 and PA), nucleoprotein (NP), neuraminidase (NA), hemagglutinin (subunits HA1 and HA2), the matrix proteins (M1 and M2) and the non-structural proteins (NS1 and NS2) (Krug *et al.*, In "The Influenza Viruses," R. M. Krug, ed., Plenum Press, N.Y., 1989, pp. 89 152).

Influenza virus' ability to cause widespread disease is due to its ability to evade the immune system by undergoing antigenic change, which is believed to occur when a host is infected simultaneously with both an animal influenza virus and a human influenza virus. During mutation and reassortment in the host, the virus may incorporate an HA and/or NA surface protein gene from another virus into its genome, thereby producing a new influenza subtype and evading the immune system.

HA is a viral surface glycoprotein generally comprising approximately 560 amino acids and representing 25% of the total virus protein. It is responsible for adhesion of the viral particle to, and its penetration into, a host cell in the early stages of infection. Cleavage of the virus HA0 precursor into the HA1 and HA2 sub-fragments is a necessary step in order for the virus to infect a cell. Thus, cleavage is required in order to convert new virus particles in a host cell into virions capable of infecting new cells. Cleavage is known to occur during transport of the integral HA0 membrane protein from the endoplasmic reticulum of the infected cell to the plasma membrane. In the course of transport, hemagglutinin undergoes a series of co- and post-translational modifications including proteolytic cleavage of the precursor HA into the amino-terminal fragment HA1 and the carboxy terminal HA2. One of the primary difficulties in growing influenza strains in primary tissue culture or established cell lines arises from the requirement for proteolytic cleavage activation of the influenza hemagglutinin in the host cell.

Although it is known that an uncleaved HA can mediate attachment of the virus to its neuraminic acid-containing receptors on a cell surface, it is not capable of the next step in the infectious cycle, which is fusion. It has been reported that exposure of the hydrophobic amino terminus of HA2 by cleavage is required so that it can be inserted into the target cell, thereby forming a bridge between virus and target cell membrane. This process is followed by fusion of the two membranes and entry of the virus into the target cell.

Proteolytic activation of HA involves cleavage at an arginine residue by a trypsin-like endoprotease, which is often an intracellular enzyme that is calcium dependent and has a neutral pH optimum. Since the activating proteases are cellular enzymes, the infected cell type determines whether the HA is cleaved. The HA of the mammalian influenza viruses and the nonpathogenic avian influenza viruses are susceptible to proteolytic cleavage only in a restricted number of cell types. On the other hand, HA of pathogenic avian viruses among the H5 and H7 subtypes are cleaved by proteases present in a broad range of different host cells. Thus, there are differences in host range resulting from differences in hemagglutinin cleavability which are correlated with the pathogenic properties of the virus.

Neuraminidase (NA) is a second membrane glycoprotein of the influenza viruses. The presence of viral NA has been shown to be important for generating a multi-faceted protective immune response against an infecting virus. For most influenza A viruses, NA is 413 amino acid in length, and is encoded by a gene of 1413 nucleotides. Nine different NA subtypes have been identified in influenza viruses (N1, N2, N3, N4, N5, N6, N7, N8 and N9), all of which have been found among wild birds. NA is involved in the destruction of the cellular receptor for the viral HA by cleaving terminal neuraminic acid (also called sialic acid) residues from carbohydrate moieties on the surfaces of infected cells. NA also cleaves sialic acid residues from viral proteins, preventing aggregation of viruses. Using this mechanism, it is hypothesized that NA facilitates release of viral progeny by preventing newly formed viral particles from accumulating along the cell membrane, as well as by promoting transportation of the virus through the mucus present on the mucosal surface. NA is an important antigenic determinant that is subject to antigenic variation.

In addition to the surface proteins HA and NA, influenza virus comprises six additional internal genes, which give rise to eight different proteins, including polymerase genes PB1, PB2 and PA, matrix proteins M1 and M2, nucleoprotein (NP), and non-structural proteins NS1 and NS2 (Horimoto *et al.*, *Clin Microbiol Rev.* 14(1):129-149, 2001).

In order to be packaged into progeny virions, viral RNA is transported from the nucleus as a ribonucleoprotein (RNP) complex composed of the three influenza virus polymerase proteins, the nucleoprotein (NP), and the viral RNA, in association with the influenza virus matrix 1 (M1) protein and nuclear export protein (Marsh *et al.*, *J Virol.* 82:2295-2304, 2008). The M1 protein that lies within the envelope is thought to function in assembly and budding. A limited number of M2 proteins are integrated into the virions (Zebedee, *J. Virol.* 62:2762-2772, 1988). They form tetramers having H⁺ ion channel activity, and when activated by the low pH in endosomes, acidify the inside of the virion, facilitating its uncoating (Pinto *et al.*, *Cell* 69:517-528, 1992). Amantadine is an anti-influenza drug that prevents viral

infection by interfering with M2 ion channel activity, thus inhibiting virus uncoating.

NS1, a nonstructural protein, has multiple functions, including regulation of splicing and nuclear export of cellular mRNAs as well as stimulation of translation. The major function of NS1 seems to be to counteract the interferon activity of the host, since an NS1 knockout virus was viable although it grew less efficiently than the parent virus in interferon-nondefective cells (Garcia-Sastre, *Virology* 252:324-330, 1998).

NS2 has been detected in virus particles (Richardson *et al.*, *Arch. Virol.* 116:69-80, 1991; Yasuda *et al.*, *Virology* 196:249-255, 1993). The average number of NS2 proteins in a virus particle was estimated to be 130-200 molecules. An *in vitro* binding assay shows direct protein-protein contact between M1 and NS2. NS2-M1 complexes have also been detected by immunoprecipitation in virus-infected cell lysates. The NS2 protein is thought to play a role in the export of RNP from the nucleus through interaction with M1 protein (Ward *et al.*, *Arch. Virol.* 140:2067-2073, 1995).

V. Influenza Proteins, VLPs and Administration Thereof

Optimized influenza HA polypeptides and influenza VLPs comprising an optimized HA (such as the HA having the sequence set forth as SEQ ID NO: 2) are provided herein. The optimized HA polypeptides can be administered to elicit an immune response against influenza. In particular examples, the optimized HA polypeptides are administered as part of a VLP.

The influenza VLPs are made up of the HA, NA and M1 proteins. The production of influenza VLPs has been described in the art and is within the abilities of one of ordinary skill in the art. As described herein, influenza VLPs can be produced by transfection of host cells with plasmids encoding the HA, NA and M1 proteins. After incubation of the transfected cells for an appropriate time to allow for protein expression (such as for approximately 72 hours), VLPs can be isolated from cell culture supernatants. Example 1 below provides an exemplary protocol for purifying influenza VLPs from cell supernatants. In this example, VLPs are

isolated by low speed centrifugation (to remove cell debris), vacuum filtration and ultracentrifugation through 20% glycerol.

The influenza VLPs disclosed herein can be used as influenza vaccines to elicit a protective immune response against H5N1 influenza viruses.

Influenza HA polypeptides and VLPs comprising HA polypeptides, or compositions thereof, can be administered to a subject by any of the routes normally used for introducing recombinant virus into a subject. Methods of administration include, but are not limited to, intradermal, intramuscular, intraperitoneal, parenteral, intravenous, subcutaneous, vaginal, rectal, intranasal, inhalation or oral. Parenteral administration, such as subcutaneous, intravenous or intramuscular administration, is generally achieved by injection. Injectables can be prepared in conventional forms, either as liquid solutions or suspensions, solid forms suitable for solution or suspension in liquid prior to injection, or as emulsions. Injection solutions and suspensions can be prepared from sterile powders, granules, and tablets of the kind previously described. Administration can be systemic or local.

Influenza VLPs, or compositions thereof, are administered in any suitable manner, such as with pharmaceutically acceptable carriers. Pharmaceutically acceptable carriers are determined in part by the particular composition being administered, as well as by the particular method used to administer the composition. Accordingly, there is a wide variety of suitable formulations of pharmaceutical compositions of the present disclosure.

Preparations for parenteral administration include sterile aqueous or non-aqueous solutions, suspensions, and emulsions. Examples of non-aqueous solvents are propylene glycol, polyethylene glycol, vegetable oils such as olive oil, and injectable organic esters such as ethyl oleate. Aqueous carriers include water, alcoholic/aqueous solutions, emulsions or suspensions, including saline and buffered media. Parenteral vehicles include sodium chloride solution, Ringer's dextrose, dextrose and sodium chloride, lactated Ringer's, or fixed oils. Intravenous vehicles include fluid and nutrient replenishers, electrolyte replenishers (such as those based on Ringer's dextrose), and the like. Preservatives and other additives may also be

present such as, for example, antimicrobials, anti-oxidants, chelating agents, and inert gases and the like.

Some of the compositions may potentially be administered as a pharmaceutically acceptable acid- or base-addition salt, formed by reaction with inorganic acids such as hydrochloric acid, hydrobromic acid, perchloric acid, nitric acid, thiocyanic acid, sulfuric acid, and phosphoric acid, and organic acids such as formic acid, acetic acid, propionic acid, glycolic acid, lactic acid, pyruvic acid, oxalic acid, malonic acid, succinic acid, maleic acid, and fumaric acid, or by reaction with an inorganic base such as sodium hydroxide, ammonium hydroxide, potassium hydroxide, and organic bases such as mono-, di-, trialkyl and aryl amines and substituted ethanolamines.

Administration can be accomplished by single or multiple doses. The dose administered to a subject in the context of the present disclosure should be sufficient to induce a beneficial therapeutic response in a subject over time, or to inhibit or prevent H5N1 influenza virus infection. The dose required will vary from subject to subject depending on the species, age, weight and general condition of the subject, the severity of the infection being treated, the particular composition being used and its mode of administration. An appropriate dose can be determined by one of ordinary skill in the art using only routine experimentation.

Provided herein are pharmaceutical compositions which include a therapeutically effective amount of the influenza VLPs alone or in combination with a pharmaceutically acceptable carrier. Pharmaceutically acceptable carriers include, but are not limited to, saline, buffered saline, dextrose, water, glycerol, ethanol, and combinations thereof. The carrier and composition can be sterile, and the formulation suits the mode of administration. The composition can also contain minor amounts of wetting or emulsifying agents, or pH buffering agents. The composition can be a liquid solution, suspension, emulsion, tablet, pill, capsule, sustained release formulation, or powder. The composition can be formulated as a suppository, with traditional binders and carriers such as triglycerides. Oral formulations can include standard carriers such as pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, and

magnesium carbonate. Any of the common pharmaceutical carriers, such as sterile saline solution or sesame oil, can be used. The medium can also contain conventional pharmaceutical adjunct materials such as, for example, pharmaceutically acceptable salts to adjust the osmotic pressure, buffers, preservatives and the like. Other media that can be used with the compositions and methods provided herein are normal saline and sesame oil.

The influenza VLPs described herein can be administered alone or in combination with other therapeutic agents to enhance antigenicity. For example, the influenza VLPs can be administered with an adjuvant, such as Freund incomplete adjuvant or Freund's complete adjuvant.

Optionally, one or more cytokines, such as IL-2, IL-6, IL-12, RANTES, GM-CSF, TNF- α , or IFN- γ , one or more growth factors, such as GM-CSF or G-CSF; one or more molecules such as OX-40L or 41 BBL, or combinations of these molecules, can be used as biological adjuvants (see, for example, Salgaller *et al.*, 1998, *J. Surg. Oncol.* 68(2):122-38; Lotze *et al.*, 2000, *Cancer J. Sci. Am.* 6(Suppl 1):S61-6; Cao *et al.*, 1998, *Stem Cells* 16(Suppl 1):251-60; Kuiper *et al.*, 2000, *Adv. Exp. Med. Biol.* 465:381-90). These molecules can be administered systemically (or locally) to the host.

A number of means for inducing cellular responses, both *in vitro* and *in vivo*, are known. Lipids have been identified as agents capable of assisting in priming CTL *in vivo* against various antigens. For example, as described in U.S. Patent No. 5,662,907, palmitic acid residues can be attached to the alpha and epsilon amino groups of a lysine residue and then linked (for example, via one or more linking residues, such as glycine, glycine-glycine, serine, serine-serine, or the like) to an immunogenic peptide. The lipidated peptide can then be injected directly in a micellar form, incorporated in a liposome, or emulsified in an adjuvant. As another example, *E. coli* lipoproteins, such as tripalmitoyl-S-glycerylcysteinylserine can be used to prime tumor specific CTL when covalently attached to an appropriate peptide (see, Deres *et al.*, *Nature* 342:561, 1989). Further, as the induction of neutralizing antibodies can also be primed with the same molecule conjugated to a

peptide which displays an appropriate epitope, two compositions can be combined to elicit both humoral and cell-mediated responses where that is deemed desirable.

Although administration of VLPs containing the optimized HA protein, one of skill in the art would understand that it is also possible to administer the optimized influenza HA protein itself (in the absence of a viral particle) or as a fusion protein to elicit an immune response in a subject.

The following examples are provided to illustrate certain particular features and/or embodiments. These examples should not be construed to limit the disclosure to the particular features or embodiments described.

EXAMPLES

Example 1: A computationally optimized broadly reactive antigen (COBRA) based H5N1 VLP vaccine elicits broadly reactive antibodies in mice and ferrets

This example describes the finding that mice and ferrets vaccinated with COBRA hemagglutinin (HA) H5N1 VLPs exhibited protective levels of HAI antibodies to representative isolates from each sub-clade of clade 2 and were completely protected from lethal challenge with a clade 2.2 H5N1 virus.

Materials and Methods

COBRA hemagglutinin (HA) construction and synthesis

Influenza A HA amino acid sequences isolated from human H5N1 infections were downloaded from the NCBI Influenza Virus Resource database (Bao *et al.*, *J Virol* 82:596-601, 2008; see Table 1 for a complete list of accession numbers and isolate descriptions). Nucleotide sequences were translated into protein sequences using the standard genetic code. All available full length sequences from H5N1 clade 2 human infections from 2004 to 2006 were acquired and used for subsequent consensus generations. For each round of consensus generation, multiple alignment analysis was applied and the consensus sequence was generated using AlignX (Vector NTI). The final amino acid sequence, termed computationally optimized broadly reactive antigen (COBRA), was reverse translated and optimized for

expression in mammalian cells, including codon usage and RNA optimization (GeneArt; Regensburg, Germany). This construct was then synthesized and inserted into the pTR600 expression vector (U.S. Patent Application Publication No. 2002/0106798; Ross *et al.*, *Nat Immunol.* 1(2):102-103, 2000; Green *et al.*, *Vaccine* 20:242-248, 2001).

Table 1: COBRA input sequences

Strain	Clade	Accession	Host	Country	Year
A/Indonesia/534H/2006	2.1.2	EU146737	Human	Indonesia	2006
A/Indonesia/535H/2006	2.1.2	EU146753	Human	Indonesia	2006
A/Indonesia/536H/2006	2.1.2	EU146754	Human	Indonesia	2006
A/Indonesia/538H/2006	2.1.2	EU146745	Human	Indonesia	2006
A/Indonesia/546bH/2006	2.1.2	EU146793	Human	Indonesia	2006
A/Indonesia/546H/2006	2.1.2	EU146755	Human	Indonesia	2006
A/Indonesia/560II/2006	2.1.2	EU146785	Human	Indonesia	2006
A/Indonesia/CDC594/2006	2.1.2	CY014272	Human	Indonesia	2006
A/Indonesia/CDC595/2006	2.1.2	CY014280	Human	Indonesia	2006
A/Indonesia/CDC596/2006	2.1.2	CY014288	Human	Indonesia	2006
A/Indonesia/CDC597/2006	2.1.2	CY014296	Human	Indonesia	2006
A/Indonesia/CDC599/2006	2.1.2	CY014303	Human	Indonesia	2006
A/Indonesia/CDC599N/2006	2.1.2	CY014477	Human	Indonesia	2006
A/Indonesia/CDC625/2006	2.1.2	CY014433	Human	Indonesia	2006
A/Indonesia/CDC625L/2006	2.1.2	CY014465	Human	Indonesia	2006
A/Indonesia/160H/2005	2.1.3	EU146648	Human	Indonesia	2005
A/Indonesia/175H/2005	2.1.3	EU146640	Human	Indonesia	2005
A/Indonesia/177H/2005	2.1.3	EU146680	Human	Indonesia	2005
A/Indonesia/195H/2005	2.1.3	EU146656	Human	Indonesia	2005
A/Indonesia/239H/2005	2.1.3	EU146664	Human	Indonesia	2005
A/Indonesia/245H/2005	2.1.3	EU146672	Human	Indonesia	2005
A/Indonesia/283H/2006	2.1.3	EU146681	Human	Indonesia	2006
A/Indonesia/286H/2006	2.1.3	EU146688	Human	Indonesia	2006
A/Indonesia/292H/2006	2.1.3	EU146713	Human	Indonesia	2006
A/Indonesia/298H/2006	2.1.3	EU146697	Human	Indonesia	2006
A/Indonesia/304H/2006	2.1.3	EU146705	Human	Indonesia	2006
A/Indonesia/321H/2006	2.1.3	EU146721	Human	Indonesia	2006
A/Indonesia/341H/2006	2.1.3	EU146729	Human	Indonesia	2006
A/Indonesia/5/2005	2.1.3	EF541394	Human	Indonesia	2005
A/Indonesia/542H/2006	2.1.3	EU146777	Human	Indonesia	2006
A/Indonesia/567H/2006	2.1.3	EU146801	Human	Indonesia	2006

Strain	Clade	Accession	Host	Country	Year
A/Indonesia/569H/2006	2.1.3	EU146809	Human	Indonesia	2006
A/Indonesia/583H/2006	2.1.3	EU146817	Human	Indonesia	2006
A/Indonesia/604H/2006	2.1.3	EU146825	Human	Indonesia	2006
A/Indonesia/7/2005	2.1.3	EU146632	Human	Indonesia	2005
A/Indonesia/CDC184/2005	2.1.3	CY014197	Human	Indonesia	2005
A/Indonesia/CDC194P/2005	2.1.3	CY014168	Human	Indonesia	2005
A/Indonesia/CDC287E/2005	2.1.3	CY014198	Human	Indonesia	2005
A/Indonesia/CDC287T/2005	2.1.3	CY014199	Human	Indonesia	2005
A/Indonesia/CDC292N/2005	2.1.3	CY014200	Human	Indonesia	2005
A/Indonesia/CDC292T/2005	2.1.3	CY014201	Human	Indonesia	2005
A/Indonesia/CDC326/2006	2.1.3	CY014204	Human	Indonesia	2006
A/Indonesia/CDC326N/2006	2.1.3	CY014202	Human	Indonesia	2006
A/Indonesia/CDC326N2/2006	2.1.3	CY014203	Human	Indonesia	2006
A/Indonesia/CDC326T/2006	2.1.3	CY014205	Human	Indonesia	2006
A/Indonesia/CDC329/2006	2.1.3	CY014206	Human	Indonesia	2006
A/Indonesia/CDC357/2006	2.1.3	CY014207	Human	Indonesia	2006
A/Indonesia/CDC370/2006	2.1.3	CY014209	Human	Indonesia	2006
A/Indonesia/CDC370E/2006	2.1.3	CY014210	Human	Indonesia	2006
A/Indonesia/CDC370P/2006	2.1.3	CY014211	Human	Indonesia	2006
A/Indonesia/CDC370T/2006	2.1.3	CY014212	Human	Indonesia	2006
A/Indonesia/CDC390/2006	2.1.3	CY014213	Human	Indonesia	2006
A/Indonesia/CDC523/2006	2.1.3	CY014311	Human	Indonesia	2006
A/Indonesia/CDC523E/2006	2.1.3	CY014368	Human	Indonesia	2006
A/Indonesia/CDC523T/2006	2.1.3	CY014376	Human	Indonesia	2006
A/Indonesia/CDC582/2006	2.1.3	CY014384	Human	Indonesia	2006
A/Indonesia/CDC610/2006	2.1.3	CY014393	Human	Indonesia	2006
A/Indonesia/CDC623/2006	2.1.3	CY014401	Human	Indonesia	2006
A/Indonesia/CDC623E/2006	2.1.3	CY014409	Human	Indonesia	2006
A/Indonesia/CDC624/2006	2.1.3	CY014417	Human	Indonesia	2006
A/Indonesia/CDC624E/2006	2.1.3	CY014425	Human	Indonesia	2006
A/Indonesia/CDC634/2006	2.1.3	CY014441	Human	Indonesia	2006
A/Indonesia/CDC634P/2006	2.1.3	CY014449	Human	Indonesia	2006
A/Indonesia/CDC634T/2006	2.1.3	CY014457	Human	Indonesia	2006
A/Indonesia/CDC644/2006	2.1.3	CY014518	Human	Indonesia	2006
A/Indonesia/CDC644T/2006	2.1.3	CY014510	Human	Indonesia	2006
A/Indonesia/CDC669/2006	2.1.3	CY014481	Human	Indonesia	2006
A/Indonesia/CDC669P/2006	2.1.3	CY014489	Human	Indonesia	2006
A/Indonesia/CDC699/2006	2.1.3	CY014497	Human	Indonesia	2006
A/Indonesia/CDC7/2005	2.1.3	CY014177	Human	Indonesia	2005
A/Indonesia/CDC739/2006	2.1.3	CY014529	Human	Indonesia	2006

Strain	Clade	Accession	Host	Country	Year
A/Indonesia/CDC759/2006	2.1.3	CY014543	Human	Indonesia	2006
A/Indonesia/CDC835/2006	2.1.3	CY017662	Human	Indonesia	2006
A/Indonesia/CDC836/2006	2.1.3	CY017670	Human	Indonesia	2006
A/Indonesia/CDC836T/2006	2.1.3	CY017678	Human	Indonesia	2006
A/Indonesia/CDC887/2006	2.1.3	CY017688	Human	Indonesia	2006
A/Indonesia/CDC938/2006	2.1.3	CY017638	Human	Indonesia	2006
A/Indonesia/CDC938E/2006	2.1.3	CY017646	Human	Indonesia	2006
A/Indonesia/CDC940/2006	2.1.3	CY017654	Human	Indonesia	2006
A/Indonesia/TLL001/2006	2.1.3	EU015403	Human	Indonesia	2006
A/Indonesia/TLL002/2006	2.1.3	EU015404	Human	Indonesia	2006
A/Indonesia/TLL003/2006	2.1.3	EU015405	Human	Indonesia	2006
A/Indonesia/TLL004/2006	2.1.3	EU015406	Human	Indonesia	2006
A/Indonesia/TLL005/2006	2.1.3	EU015407	Human	Indonesia	2006
A/Indonesia/TLL006/2006	2.1.3	EU015408	Human	Indonesia	2006
A/Indonesia/TLL007/2006	2.1.3	EU015409	Human	Indonesia	2006
A/Indonesia/TLL008/2006	2.1.3	EU015410	Human	Indonesia	2006
A/Indonesia/TLL009/2006	2.1.3	EU015411	Human	Indonesia	2006
A/Indonesia/TLL010/2006	2.1.3	EU015412	Human	Indonesia	2006
A/Indonesia/TLL011/2006	2.1.3	EU015413	Human	Indonesia	2006
A/Indonesia/TLL012/2006	2.1.3	EU015414	Human	Indonesia	2006
A/Indonesia/TLL013/2006	2.1.3	EU015415	Human	Indonesia	2006
A/Indonesia/TLL014/2006	2.1.3	EU015416	Human	Indonesia	2006
A/Djibouti/5691NAMRU3/2006	2.2	DQ666146	Human	Djibouti	2006
A/Egypt/7021-NAMRU3/2006	2.2	CY062439	Human	Egypt	2006
A/human/Iraq/207-NAMRU3/2006	2.2	DQ435202	Human	Iraq	2006
A/Iraq/1/2006	2.2	EU146870	Human	Iraq	2006
A/Iraq/659/2006	2.2	EU146876	Human	Iraq	2006
A/Iraq/754/2006	2.2	EU146877	Human	Iraq	2006
A/Iraq/755/2006	2.2	EU146869	Human	Iraq	2006
A/Iraq/756/2006	2.2	EU146878	Human	Iraq	2006
A/Turkey/12/2006	2.2	EF619982	Human	Turkey	2006
A/Turkey/15/2006	2.2	EF619989	Human	Turkey	2006
A/Turkey/651242/2006	2.2	EF619990	Human	Turkey	2006
A/Turkey/65596/2006	2.2	EF619998	Human	Turkey	2006
A/Xinjiang/1/2006	2.2	FJ492886	Human	China	2006
A/Egypt/14724-NAMRU3/2006	2.2.1	EF200512	Human	Egypt	2006
A/Egypt/14725-NAMRU3/2006	2.2.1	EF200513	Human	Egypt	2006
A/Egypt/2782-NAMRU3/2006	2.2.1	DQ464377	Human	Egypt	2006
A/Egypt/2991-NAMRU3/2006	2.2.1	EU095023	Human	Egypt	2006
A/Egypt/2992-NAMRU3/2006	2.2.1	EU095024	Human	Egypt	2006

Strain	Clade	Accession	Host	Country	Year
A/Egypt/902782/2006	2.2.1	EU146867	Human	Egypt	2006
A/Egypt/902786/2006	2.2.1	EU146868	Human	Egypt	2006
A/Anhui/1/2005	2.3.4	DQ371928	Human	China	2005
A/Anhui/2/2005	2.3.4	DQ371929	Human	China	2005
A/China/2006	2.3.4	EF624256	Human	China	2006
A/China/GD01/2006	2.3.4	DQ835313	Human	China	2006
A/Fujian/1/2005	2.3.4	FJ492882	Human	China	2005
A/Guangdong/1/2006	2.3.4	FJ492884	Human	China	2006
A/Guangxi/1/2005	2.3.4	DQ371930	Human	China	2005
A/human/China/GD02/2006	2.3.4	EU263981	Human	China	2006
A/Hunan/1/2006	2.3.4	FJ492879	Human	China	2006
A/Jiangxi/1/2005	2.3.4	FJ492885	Human	China	2005
A/Shanghai/1/2006	2.3.4	AB462295	Human	China	2006
A/Shenzhen/406H/2006	2.3.4	EF137706	Human	China	2006
A/Sichuan/1/2006	2.3.4	FJ492881	Human	China	2006
A/Vietnam/UT30850/2005	2.3.4	HM114537	Human	Viet Nam	2005
A/Zhejiang/1/2006	2.3.4	FJ492880	Human	China	2006
A/Zhejiang/16/2006	2.3.4	DQ643809	Human	China	2006

COBRA HA antigenic modeling

Influenza hemagglutinin (HA) protein sequences representing COBRA, A/Indonesia/5/2005 (Clade 2.1), A/Whooper Swan/Mongolia/244/2005 (Clade 2.2) and A/Anhui/1/2005 (Clade 2.3) were submitted to the 3D-JIGSAW Protein Comparative Modeling website for rendering (Bates *et al.*, *Proteins* 45(S5):39-46, 2001; Bates and Sternberg, *Proteins* 37(S3):47-54, 1999; Contreras-Moreira and Bates, *Bioinformatics* 18(8):1141-1142, 2002). Structures were overlaid and analyzed using Swiss-Pdb viewer software (Guex and Peitsch, *Electrophoresis* 18(15):2714-23, 1998). Antigenic regions and designations are based on the original description of the antigenic structure of the H3N2 virus A/Hong Kong/1/1968 (Wiley *et al.*, *Nature* 289(5796):373-378, 1981). No significant alterations were observed in region B of the COBRA HA relative to the primary influenza isolates; however, some divergent structures in HA regions A and C were identified in primary isolates.

***In vitro* expression**

COBRA HA protein expression was confirmed by transfecting mammalian cells. Human embryonic kidney (HEK) 293T cells (1×10^6) were transiently transfected with 3 μ g of DNA expressing COBRA. Cells were incubated for 72 hours at 37°C, supernatants were removed, the cells were lysed with 1% Triton-X 100 and cell lysates were collected. Cell lysates were electrophoresed on a 10% SDS-PAGE gel and transferred to a PVDF membrane. The blot was probed with mouse polyclonal antisera pooled from mice infected with 6:2 reassortant H5N1 viruses with the surface glycoproteins derived from either A/Vietnam/1203/2004 or A/Whooper Swan/244/2005, and the HA-antibody complexes were detected using a goat anti-mouse IgG conjugated to horse radish peroxidase (HRP) (Southern Biotech; Birmingham, AL, USA). HRP was detected by chemiluminescent substrate (Pierce Biotechnology; Rockford IL, USA) and exposed to X-ray film (ThermoFisher; Pittsburgh, PA, USA).

COBRA HA functional characterization

To determine receptor binding characteristics, virus-like particles (VLPs) containing COBRA HA proteins were purified from the supernatants of mammalian cell lines. HEK 293T cells were transiently transfected with plasmids expressing HIV Gag, COBRA HA and neuraminidase (NA, A/Thailand/1(KAN-1)/2004) and incubated for 72 hours at 37°C. Supernatants were collected and VLPs were purified via ultracentrifugation (100,000 X g through 20% glycerol, weight per volume) for 4 hours at 4°C. The pellets were subsequently resuspended in phosphate buffered saline PBS, pH 7.2 and stored at -80°C until use. Protein concentration was determined by Micro BCA™ Protein Assay Reagent Kit (Pierce Biotechnology, Rockford, IL, USA). COBRA HA VLPs were prepared in various amounts as measured by total protein and each individual preparation was two-fold serially diluted in v-bottom microtiter plates. An equal volume of either 1% turkey or 1% horse erythrocytes (RBC) (Lampire; Pipersville, PA, USA) in PBS was added to the diluted VLPs and incubated for 30-60 minutes at room temperature. The HA titer

was determined by the reciprocal dilution of the last well which contained agglutinated RBC.

To determine endosomal fusion characteristics, COBRA-pseudotyped lentiviral vectors expressing a luciferase reporter gene were produced as described (Yang *et al.*, *J Virol* 78(8):4029-4036). Briefly, 293T cells were co-transfected by using the following plasmids: 7µg of pCMVdeltaR8.2, 7µg of pHRCMV-Luc, 3µg pCMV/R N1(Kan-1) and 3 µg pTR600 COBRA. Cells were transiently transfected and incubated for 72 hours at 37°C. Supernatants were harvested, centrifuged to clear cell debris, filtered through a 0.22 µm syringe filter, aliquotted, and used immediately or frozen at -80°C. For fusion assays, 100 µl of pseudoviruses were added to confluent MDCK cells in 48-well plates (~30,000 cells per well). Plates were incubated at room temperature for 30 minutes, washed, and fresh medium added. Forty-eight hours after infection, cells were lysed in mammalian cell lysis buffer (Promega; Madison, WI, USA). A standard quantity of cell lysate was used in a luciferase assay with luciferase assay reagent (Promega; Madison, WI, USA) according to the manufacturer's protocol.

Vaccine Preparation and Dose Determination

HEK 293T cells were transiently transfected with plasmids expressing M1 (A/Puerto Rico/8/1934, optimized for expression in mammalian cells; SEQ ID NO: 9), NA (A/Thailand/1(KAN-1)/2004, optimized for expression in mammalian cells; SEQ ID NO: 8) and COBRA HA (SEQ ID NO: 7) and incubated for 72 hours at 37°C. Supernatants were collected and cell debris removed by low speed centrifugation followed by vacuum filtration through a 0.22 µm sterile filter. VLPs were purified via ultracentrifugation (100,000 x g through 20% glycerol, weight per volume) for 4 hours at 4°C. The pellets were subsequently resuspended in PBS pH 7.2 and stored in single use aliquots at -80°C until use. Total protein concentration was determined by Micro BCA™ Protein Assay Reagent Kit (Pierce Biotechnology, Rockford, IL, USA).

HA specific content was determined by western blot and densitometry. Purified recombinant COBRA HA and purified VLPs were prepared in standard

total protein amounts and were electrophoresed on a 10% SDS-PAGE gel and transferred to a PVDF membrane. The blot was probed with mouse polyclonal antisera pooled from mice infected with 6:2 reassortant H5N1 viruses with the surface glycoproteins derived from either A/Vietnam/1203/2004 or A/Whooper Swan/244/2005 and the HA-antibody complexes were detected using a goat anti-mouse IgG conjugated to horse radish peroxidase (HRP) (Southern Biotech; Birmingham, AL, USA). HRP was detected by chemiluminescent substrate (Pierce Biotechnology; Rockford IL, USA) and exposed to X-ray film (ThermoFisher; Pittsburgh, PA, USA). Density of bands was determined using ImageJ software (NIH) (Abramoff *et al.*, *Biophotonics International* 11(7):36-42, 2004). Density of recombinant HA bands were used to calculate a standard curve and the density of the purified VLPs was interpolated using the results from the recombinant HA. Experiments were performed in triplicate and multiple exposure times were analyzed for all iterations.

Codon-optimized influenza HA, NA and M1 genes

The nucleotide sequences of the codon-optimized HA (SEQ ID NO: 1), codon-optimized NA (SEQ ID NO: 3) and codon-optimized M1 (SEQ ID NO: 5) genes are set forth in the Sequence Listing. The corresponding amino acid sequences of the encoded HA, NA and M1 proteins are set forth in the Sequence Listing as SEQ ID NO: 2, SEQ ID NO: 4 and SEQ ID NO: 6, respectively.

Mouse studies

BALB/c mice (*Mus musculus*, females, 6-8 weeks old) were purchased from Harlan Sprague Dawley (Indianapolis, IN, USA) and housed in microisolator units and allowed free access to food and water and were cared for under USDA guidelines for laboratory animals. Mice (5 mice per group) were vaccinated with one of three doses of purified COBRA HA VLPs (1.5 µg, 0.3 µg or 0.06 µg), based upon HA content from a densitometry assay, via intramuscular injection at week 0 and then boosted with the same dose at week 3. For comparison studies, mice (20 mice per group) were vaccinated with purified VLPs (3 µg) via intramuscular

injection at week 0 and then boosted with the same dose at week 3. Vaccines at each dose were formulated with Imject® alum adjuvant (Imject® Alum, Pierce Biotechnology; Rockford, IL, USA) according to the manufacturer's protocol or vehicle alone. Fourteen to twenty-one days after each vaccination, blood was collected from anesthetized mice via the retro-orbital plexus and transferred to a microfuge tube. Tubes were centrifuged and sera was removed and frozen at $-20 \pm 5^{\circ}\text{C}$.

Three weeks after final vaccination, mice were challenged intranasally with 5×10^3 plaque forming units (PFU) of the highly pathogenic H5N1 virus A/Whooper Swan/Mongolia/244/2005 (clade 2.2) in a volume of 50 μl . The challenge dose represents approximately 50LD₅₀ in mice. After infection, mice were monitored daily for weight loss, disease signs and death for 14 days after infection. Individual body weights, sickness scores and death were recorded for each group on each day after inoculation. Sickness score was determined by evaluating activity (0=normal, 1=reduced, 2=severely reduced), hunched back (0=absent, 1=present) and ruffled fur (0=absent, 1=present) (Toapanta and Ross, *Respiratory Res* 10(1):112, 2009). Experimental endpoint was defined as >20% weight loss or display of neurological disease such as hind limb paralysis. All H5N1 influenza virus studies were performed under high-containment biosafety level 3 enhanced conditions (BSL3+).

Ferret studies

Fitch ferrets (*Mustela putorius furo*, female, 6-12-months of age), influenza naïve and de-scented, were purchased from Marshall Farms (Sayre, PA, USA). Ferrets were pair housed in stainless steel cages (Shor-line, Kansas City, KS, USA) containing Sani-chips Laboratory Animal Bedding (P.J. Murphy Forest Products, Montville, NJ, USA). Ferrets were provided with Teklad Global Ferret Diet (Harlan Teklad, Madison, WI, USA) and fresh water *ad libitum*. The COBRA HA VLPs were diluted in PBS, pH 7.2 to achieve final concentration. Ferrets (n=3) were vaccinated with 15 μg of purified COBRA VLPs, based upon HA content as determined by densitometry assay, via intramuscular injection in the quadriceps

muscle in a volume of 0.25 ml at week 0 and then boosted with the same dose at week 3. Vaccines were stored at -80°C prior to use and formulated with Imject® alum adjuvant (Imject® Alum; Pierce Biotechnology, Rockford, IL, USA) immediately prior to use. Animals were monitored for adverse events including weight loss, temperature, loss of activity, nasal discharge, sneezing and diarrhea weekly during the vaccination regimen. Prior to vaccination, animals were confirmed by HAI assay to be seronegative for circulating influenza A (H1N1 and H3N2) and influenza B viruses. Fourteen to twenty-one days after each vaccination, blood was collected from anesthetized ferrets via the anterior vena cava and transferred to a microfuge tube. Tubes were centrifuged and sera was removed and frozen at $-20 \pm 5^{\circ}\text{C}$.

Three weeks after final vaccination, ferrets were challenged intranasally with 1×10^6 plaque forming units (PFU) of the highly pathogenic H5N1 virus A/Whooper Swan/Mongolia/244/2005 (clade 2.2) in a volume of 0.5 ml in each nostril for a total infection volume of 1 ml. After infection, ferrets were monitored daily for weight loss, disease signs and death for 14 days after infection. Individual body weights, sickness scores, and death were recorded for each group on each day after inoculation. Sickness score was determined by evaluating activity (0=normal, 1=alert and active with stimulation, 2=alert but not active after stimulation, 3=not alert or active after stimulation), nasal discharge (0=absent, 1=present), sneezing (0=absent, 1=present), decreased food intake (0=absent, 1=present), diarrhea (0=absent, 1=present), dyspnea (0=absent, 1=present) and neurological symptoms (0=absent, 1=present). Nasal washes were performed by instilling 3 ml of PBS into the nares of anesthetized ferrets each day for 14 days after inoculation. Washes were collected and stored at -80°C until use. Experimental endpoint was defined as $>20\%$ weight loss, development of neurological symptoms, or an activity score of 3 (not active or alert after stimulation). All H5N1 influenza virus studies were performed under high-containment biosafety level 3 enhanced conditions (BSL3+).

ELISA

The ELISA assay was used to assess total antibody titer and IgG isotype titer to the HA. High binding, 96-well polystyrene plates (Costar; Lowell, MA, USA) were coated overnight with 50 ng/well of recombinant HA. Coating antigens were derived from the following representative viral isolates: A/Vietnam/1203/2004 (clade 1), A/Indonesia/5/2005 (clade 2.1), A/Whooper Swan/244/2005 (clade 2.2) and A/Anhui/1/2005 (clade 2.3). Plates were blocked with 5% milk diluted in PBS with 0.05% Tween 20. Serum samples were diluted in blocking buffer and added to plates. Serum was two-fold serially diluted and allowed to incubate for 1 hour at room temperature. Plates were washed and species specific antibody against IgG, IgG1, IgG2a, IgG2b or IgG3 and linked to horseradish peroxidase (HRP) (Southern Biotech; Birmingham, AL, USA) were diluted in blocking buffer and added to plates. Plates were incubated for 1 hour at room temperature. Plates were washed and HRP was developed with TMB substrate (Sigma-Aldrich; St. Louis, MO, USA). Plates were incubated in the dark for 15 minutes and then the reaction was stopped with 2N H₂SO₄. Optical densities at a wavelength of 450 nm (OD₄₅₀) were read by a spectrophotometer (BioTek; Winooski, VT, USA) and end point dilution titers were determined. End point titers were determined as the reciprocal dilution of the last well which had an OD₄₅₀ above the mean OD₄₅₀ plus two standard deviations of naïve animal sera.

Hemagglutination Inhibition (HAI)

The HAI assay was used to assess functional antibodies to HA able to inhibit agglutination of horse erythrocytes. The protocol was adapted from the CDC laboratory-based influenza surveillance manual (Gillim-Ross and Subbarao, *Clin Microbiol Rev* 19(4):614-636, 2006). To inactivate non-specific inhibitors, sera were treated with receptor destroying enzyme (RDE; Denka Seiken, Co., Japan) prior to being tested (Bright *et al.*, *Lancet* 366(9492):1175-1181, 2005; Bright *et al.*, *Virology* 308(2):270-278, 2003; Bright *et al.*, *JAMA* 295(8):891-894, 2006; Mitchell *et al.*, *Vaccine* 21(9-10):902-914, 2004; Ross *et al.*, *Nat Immunol* 1(2):127-131, 2000). Briefly, three parts RDE was added to one part sera and incubated overnight

at 37°C. RDE was inactivated by incubation at 56°C for ~30 minutes. RDE-treated sera was two-fold serially diluted in v-bottom microtiter plates. An equal volume of reassortant virus, adjusted to approximately 8 HAU/50 µl, was added to each well. The reassortant viruses contained the internal genes from the mouse adapted strain A/Puerto Rico/8/1934 and the surface proteins HA and NA from the following representative viral isolates: A/Vietnam/1203/2004 (clade 1), A/Indonesia/5/2005 (clade 2.1), A/Whooper Swan/244/2005 (clade 2.2) and A/Anhui/1/2005 (clade 2.3). The plates were covered and incubated at room temperature for 20 minutes followed by the addition of 1% horse erythrocytes (hRBC) (Lampire Biologicals, Pipersville, PA, USA) in PBS. Red blood cells were stored at 4°C and used within 72 hours of preparation. The plates were mixed by agitation, covered, and the RBCs were allowed to settle for 1 hour at room temperature (Askonas B, McMichael A, Webster R. The immune response to influenza viruses and the problem of protection against infection. In: Beare AS, editor. Basic and applied influenza research: CRC Press 1982: 159-188). The HAI titer was determined by the reciprocal dilution of the last row which contained non-agglutinated RBCs. Positive and negative serum controls were included for each plate. All mice were negative (HAI ≤1:10) for pre-existing antibodies to currently circulating human influenza viruses prior to vaccination.

Plaque Assay

Madin-Darby Canine Kidney (MDCK) cells were plated (5×10^5) in each well of a 6-well plate. Samples were diluted (final dilution factors of 10^0 to 10^{-6}) and overlaid onto the cells in 100 µl of DMEM supplemented with penicillin-streptomycin and incubated for 1 hour. Samples were removed, cells were washed twice and media was replaced with 2 ml of L15 medium plus 0.8% agarose (Cambrex; East Rutherford, NJ, USA) and incubated for 72 hours at 37°C with 5% CO₂. Agarose was removed and discarded. Cells were fixed with 10% buffered formalin, and then stained with 1% crystal violet for 15 minutes. Following thorough washing in dH₂O to remove excess crystal violet, plates were allowed to dry, plaques counted, and the plaque forming units (PFU)/ml were calculated.

Statistical Analysis

Statistical significance of the antibody data was determined using a two-way analysis of variance (ANOVA) with Bonferroni's post-test to analyze differences between each vaccine group for the different test antigens (multiparametric). Differences in weight loss, sickness score, and viral titers were analyzed by two-way ANOVA, followed by Bonferroni's post test for each vaccine group at multiple time points. Significance was defined as $p < 0.05$. Statistical analyses were done using GraphPad Prism software.

Results

Computationally optimized broadly reactive antigen design

To address the challenge of antigenic diversity present in H5N1 influenza, a computationally optimized broadly reactive antigen (COBRA) was designed. For the first step of antigen generation, 129 unique hemagglutinin (HA) sequences were downloaded from the NCBI Influenza Virus Resource (IVR) sequence database (Bao *et al.*, *J Virol* 82:596-601, 2008) representing clade 2 H5N1 viruses isolated from human infections between 2004 and 2006. The sequences were first grouped into phylogenetic sub-clades and then further divided into individual outbreak groups within each sub-clade based upon geographic location and time of isolation. HA amino acid sequences for each individual outbreak group were aligned and the most common amino acid at each position was determined resulting in primary consensus sequences representing each outbreak group within each sub-clade (FIG. 1A). Primary consensus sequences within each sub-clade were then aligned and the most common amino acid was chosen resulting in secondary consensus sequences representing each sub-clade (FIG. 1A). The secondary consensus sequences were aligned and the most common amino acid at each position was selected resulting in the final consensus sequence referred to as clade 2 COBRA HA (FIG. 1A). Phylogenetic analysis of the COBRA HA with all human isolates of H5N1 HA proteins indicated that COBRA retained a clade 2-like sequence without being grouped specifically within any clade 2 sub-clade cluster (FIG. 1B). Furthermore, a

BLAST search using the COBRA HA sequence revealed that it is a unique sequence that has not been isolated from the environment.

Characterization of COBRA

Since COBRA is a fully synthetic protein, the retention of natural hemagglutinin function was confirmed. Initially, COBRA expression was verified by transient transfection of mammalian cells. Analysis of the total cell lysate demonstrated that the COBRA HA migrates at its predicted molecular weight of approximately 73kDa (FIG. 2A). Because the poly-basic cleavage site was retained in the COBRA HA sequence, both HA0 and the HA1 subunits were detected by immunoblot at similar molecular weights as recombinant HA and the HA on the H5N1 virion (FIG. 2A). Virus-like particles (VLPs) with COBRA HA on the surface bound sialic acid in a dose-dependent manner and this binding was specific to COBRA, since empty lentiviral core alone did not bind to the receptor (FIG. 2B).

To determine if the COBRA HA was functional, the protein was pseudotyped onto lentiviral Gag_{p24} to generate pseudoparticles (Nefkens *et al.*, *J Clin Virol* 39(1):27-33, 2007; Haynes *et al.*, *Vaccine* 27(4):530-541, 2009). COBRA HA containing pseudoparticles mediated cell fusion as efficiently as H5N1 control pseudoparticles without the requirement for trypsin. In contrast, H1N1 pseudoparticles did require trypsin and pseudoparticles without surface HA produced luciferase at similar levels as the cell only controls (FIG. 2C). Taken together, these results demonstrate that although the COBRA HA is a synthetic protein not found in nature, it retains all of the functions of a natural hemagglutinin protein.

Mouse Dosing Immunizations

Mice (BALB/c; n=5) were vaccinated (week 0 and 3) via intramuscular injection with purified COBRA VLPs at either a high dose (1.5 µg HA) or low dose (0.3 µg HA) with and without Imject® alum adjuvant. At week 5, all COBRA VLP-vaccinated animals had anti-HA antibodies that recognized heterologous recombinant HA derived from both clade 1 and also sub-clades of clade 2 (FIG. 3A

and 3B). Imject® alum significantly increased anti-HA antibody titers in both low and high dose groups as compared to the non-adjuvanted groups ($p < 0.01$). The IgG isotype subclasses elicited by the VLP vaccines against a clade 2.1 coating antigen consisted mainly of IgG1 and IgG2a, indicating a mixed T helper response (FIG. 3C and 3D). Similar results were found for additional coating antigens representing clade 1, clade 2.2 and clade 2.3. In addition to recognizing HA, antibodies were also evaluated for the ability to block virus from binding its receptor via inhibition of viral-induced agglutination of horse erythrocytes (HAI). All mice administered Imject® alum adjuvanted vaccines, regardless of dose, had HAI titers $\geq 1:40$ to viruses expressing HA from clades 2.1 and 2.2 and 90% of the mice had titers $\geq 1:40$ to a clade 2.3 representative virus (FIG. 3E and 3F). Non-adjuvanted vaccines elicited generally lower HAI antibody titers with 100% of high dose animals achieving titers $\geq 1:40$ only against clade 2.1 viruses. Imject® alum adjuvanted vaccines elicited significantly higher HAI antibody titers to clade 2.2 and clade 2.3 viruses regardless of dose as compared to non-adjuvanted vaccines ($p < 0.05$ for high dose and $p < 0.001$ for low dose, respectively). None of the vaccines elicited high HAI titer antibodies to a clade 1 virus.

Mouse Dosing Challenge

Mice that received the COBRA VLP vaccines or mock vaccinated control mice were challenged intranasally with a lethal dose of clade 2.2 H5N1 highly pathogenic avian influenza (A/Mongolia/whooper swan/244/2005) to evaluate the protective efficacy of the different COBRA vaccine formulations. All COBRA vaccinated mice, regardless of dose or the presence of adjuvant, were protected from weight loss and death following lethal challenge, while all mock vaccinated animals rapidly lost weight and required euthanasia by day 6 post infection (FIGS. 4A and 4B). Additionally, COBRA VLP vaccinated mice had no signs of disease, while mock vaccinated animals developed such symptoms as ruffled fur, hunched back, and lethargy (FIGS. 4C and 4D).

Mouse Comparison Immunizations

To determine if the COBRA HA vaccine elicits a broader antibody response compared to a vaccine derived from a primary isolate, an additional set of mice were vaccinated with either COBRA VLPs or clade 2.2 (A/Mongolia/whooper swan/244/2005) VLPs. Mice (BALB/c; n=20) were vaccinated (week 0 and 3) via intramuscular injection with either COBRA VLPs or clade 2.2 VLPs at a high dose (3 µg HA) with Imject® alum adjuvant. At week 5, all COBRA VLP-vaccinated mice and all clade 2.2 VLP-vaccinated mice had anti-HA antibodies that recognized heterologous recombinant HA derived from both clade 1 and various sub-clades of clade 2 (FIG. 5A). Although no significant differences were found in total IgG titers between vaccine groups, COBRA VLP-vaccinated animals had higher HAI antibody titers against all viruses tested as compared to clade 2.2 VLP-vaccinated animals ($p < 0.01$; FIG. 5B). Furthermore, COBRA VLP-vaccinated animals had an increased frequency of HAI titers of $\geq 1:40$ compared to clade 2.2 VLP-vaccinated animals (Table 2).

Table 2: Mouse seroconversion frequency

Vaccine Antigen	Clade 1 ^a	Clade 2.1 ^b	Clade 2.2 ^c	Clade 2.3 ^d
COBRA	45% (9/20)	100% (20/20)	100% (20/20)	100% (20/20)
Clade 2.2 ^c	0% (0/20)	0% (0/20)	0% (0/20)	0% (0/20)

Percentage of VLP-vaccinated animals achieving an HAI titer of $\geq 1:40$ to each test antigen.

^a A/Vietnam/1203/2004

^b A/Indonesia/5/2005

^c A/Whooper Swan/Mongolia/244/2005

^d A/Anhui/1/2005

Mouse Comparison Challenge

Mice that received the COBRA VLP vaccine, clade 2.2 VLP vaccine or mock vaccinated control mice were challenged intranasally with a lethal dose of clade 2.2 H5N1 highly pathogenic avian influenza (A/Mongolia/whooper swan/244/2005) to evaluate the protective efficacy of the VLP vaccines. All VLP-vaccinated mice were protected from weight loss and death following lethal challenge while all mock vaccinated animals rapidly lost weight and required

euthanasia by day 6 post infection (FIG. 6A). Additionally, VLP vaccinated mice did not show signs of disease, while mock vaccinated animals developed ruffled fur, hunched back, and lethargy (FIG. 6B). Even though the clade 2.2 VLP was matched to the challenge virus, no significant differences were found between COBRA VLP and clade 2.2 VLP vaccinated mice in any of the parameters analyzed indicating that the COBRA VLP vaccine protected animals as efficiently as the homologous vaccine.

Ferret Comparison Immunizations

Ferrets are the most relevant model for influenza disease and as such the COBRA vaccine was tested in this more rigorous animal model. Ferrets (Fitch; n=9) were vaccinated (week 0 and 3) via intramuscular injection with COBRA VLPs or clade 2.2 VLPs at a high dose (15 µg HA) with Imject® alum adjuvant. Serum was collected from ferrets at week 5 and antibody responses to the COBRA vaccines were evaluated. All vaccinated ferrets had anti-HA antibodies that recognized heterologous recombinant HA derived from both clade 1 and also sub-clades of clade 2 (FIG. 7A). No significant difference in anti-HA antibody was found between the COBRA VLP vaccine and the clade 2.2 VLP vaccine for any of the antigens tested ($p>0.05$). In addition to recognizing HA, antibodies were also evaluated for HAI activity. COBRA VLP-vaccinated animals had higher HAI antibody titers against clade 2.1 and clade 2.3 viruses as compared to clade 2.2 VLP-vaccinated animals ($p<0.01$ FIG. 7B). Similar to the mice, COBRA VLP-vaccinated ferrets displayed an increased rate of achieving HAI titers $\geq 1:40$ when compared to clade 2.2 VLP-vaccinated ferrets (Table 3).

Table 3: Ferret seroconversion frequency

Vaccine Antigen	Clade 1 ^a	Clade 2.1 ^b	Clade 2.2 ^c	Clade 2.3 ^d
COBRA	0% (0/9)	78% (7/9)	56% (5/9)	56% (5/9)
Clade 2.2 ^c	0% (0/9)	0% (0/9)	22% (2/9)	0% (0/9)

Percentage of VLP-vaccinated animals achieving an HAI titer of $\geq 1:40$ to each test antigen.

^a A/Vietnam/1203/2004

^b A/Indonesia/5/2005

^c A/Whooper Swan/Mongolia/244/2005

^d A/Anhui/1/2005

Ferret Comparison Challenge

Ferrets that received the COBRA VLP vaccines, clade 2.2 VLP vaccines or mock vaccinated control animals were challenged intranasally with clade 2.2 H5N1 highly pathogenic avian influenza (A/Mongolia/whooper swan/244/2005) to evaluate the protective efficacy of the COBRA vaccine in the ferret model of influenza infection. All VLP vaccinated ferrets were protected from weight loss and death following viral challenge, while all mock vaccinated animals rapidly lost weight and 78% (7/9) of mock vaccinated animals required euthanasia by day 7 post-infection (FIGS. 8A and 8B). Additionally, both COBRA VLP-vaccinated and clade 2.2-vaccinated ferrets were protected from acute fever and failed to develop significant signs of disease while mock vaccinated animals had an elevated body temperature and developed such symptoms as lethargy, diarrhea and decreased food and water intake (FIGS. 8C and 8D). In addition to monitoring outward signs of disease progression, nasal washes were collected for determination of viral replication in the upper respiratory tract. Ferrets vaccinated with either COBRA VLPs or clade 2.2 VLPs did not have detectable virus at any point after infection, while mock vaccinated animals had high levels of viral replication for the first five days of the infection (FIG. 8E). No significant differences were found between COBRA VLP and clade 2.2 VLP vaccinated ferrets in any of the challenge parameters analyzed confirming the findings in mice that the COBRA VLP vaccine protected animals as efficiently as the homologous vaccine.

The percent identity of COBRA HA and the test antigens used in the mouse and ferret studies described above are shown in Table 4.

Table 4: Percent Identity of Test Antigens

Vaccine Antigen	Clade 1^a	Clade 2.1^b	Clade 2.2^c	Clade 2.3^d
COBRA	97%	97%	95%	97%
Clade 2.2 ^c	94%	97%	100%	94%

HA amino acid sequences were aligned and percent identity across the entire protein was determined for the vaccine immunogens compared to the representative test antigens.

^a A/Vietnam/1203/2004

^b A/Indonesia/5/2005

^c A/Whooper Swan/Mongolia/244/2005

^d A/Anhui/1/2005

Example 2: A Computationally-Optimized HA VLP Vaccines Elicits Broadly-Reactive Antibodies that Protect Monkeys from H5N1 Infection

This example describes the finding that a COBRA clade 2 HA H5N1 VLP elicits broad humoral immunity against multiple H5N1 isolates from different clades.

Materials and Methods

Expression and purification of virus-like particles

The COBRA HA sequence is described above in Example 1. 293T cells were transiently transfected with plasmids expressing HA, M1, and NA in low serum media, incubated for 72h at 37^oC, and purified by ultracentrifugation through a 20% glycerol cushion as previously described (Giles and Ross, *Vaccine* 29:3043-3054, 2011). All VLP vaccines were engineered using the same NA from A/Thailand/1(KAN-1)/2004. HA content was quantified as previously described (Giles and Ross, *Vaccine* 29:3043-3054, 2011). Two different VLP preparations were purified, each containing one of the HA influenza gene products: WS/05 or the COBRA HA.

Primate immunizations and H5N1 challenges

Cynomolgus macaques (*Macaca fascicularis*, male, 3-5 years old) were vaccinated with 15 µg (based upon HA content) of purified COBRA HA VLPs (n=7) or WS/05 VLPs (n=7), via intramuscular injection at weeks 0, 3 and 6. Vaccines at each dose were formulated with alum adjuvant (Imject® Alum, Pierce Biotechnology; Rockford, IL, USA) immediately prior to use. Twenty-one days after each vaccination, blood was collected from anesthetized macaques. All

procedures were in accordance with the NRC Guide for the Care and Use of Laboratory Animals.

Three weeks after final vaccination, macaques were placed into BSL3+ isolator units (Bioqual, Inc., Rockville, MD) and then challenged by a multi-route of infection (ocular, nasal, tracheal) as previously described (Kobasa *et al.*, *Nature* 445:319-323, 2007; Kuiken *et al.*, *Vet Pathol* 40:304-310, 2003; Rimmelzwaan *et al.*, *Avian Dis* 47:931-933, 2003) using 1×10^6 plaque forming units (PFU) of the highly pathogenic H5N1 virus, A/Whooper Swan/Mongolia/244/2005 (clade 2.2), at each location. Monkeys were monitored daily for weight loss, signs of disease, and mortality until 7 days after infection. Individual body weights, sickness scores (based upon lethargy, temperature change, nasal discharge, lack of appetite, dehydration, lack of responsiveness), and death were recorded for each group.

Nasal and tracheal washes were performed at day 0, 1, 3, 5, and 7 post-infection. In addition, subsets of monkeys were sacrificed following administration of anesthesia and necropsies were performed according to standard procedures for assessment of gross pathologic and histopathologic changes, as well as the extent of virus replication.

Serological assays

A quantitative ELISA was performed to assess anti-HA specific IgG in immune serum as previously described (Bright *et al.*, *PLoS One* 3:e1501, 2008; Giles and Ross, *Vaccine* 29:3043-3054, 2011). The hemagglutination inhibition (HAI) assay was used on sera treated with receptor destroying enzyme (RDE; Denka Seiken, Co., Japan) prior to being tested (Bright *et al.*, *Vaccine* 25:3871-3878, 2007; Mitchell *et al.*, *Vaccine* 21:902-914, 2003; Bright *et al.*, *PLoS One* 3:e1501, 2008) to assess functional antibodies to the HA able to inhibit agglutination of horse red blood (Askonas B, McMichael A, Webster R. The immune response to influenza viruses and the problem of protection against infection. In: Beare AS, editor. Basic and applied influenza research: CRC Press 1982: 159-188). The protocol was adapted from the CDC laboratory-based influenza surveillance manual and performed as previously described (Gillim-Ross and Subbarao, *Clin Microbiol Rev*

19:614-636, 2006; Bright *et al.*, *PLoS One* 3:e1501, 2008). The HAI titer was determined by the reciprocal dilution of the last row which contained non-agglutinated RBC. Positive and negative serum controls were included for each plate. All monkeys were negative (HAI \leq 1:10) for pre-existing antibodies to currently circulating human influenza viruses prior to vaccination. Serum neutralizing antibody titers were determined by microneutralization (MN) assays performed on Madin Darby Canine Kidney (MDCK) cells following the procedure until CPE was observed (Rowe *et al.*, *J Clin Microbiol* 37:937-943, 1999). Cells were then fixed in 10% formalin and stained with 1% crystal violet to quantify CPE. The neutralizing antibody titers are expressed as the reciprocal of the highest dilution of serum that gave 50% neutralization of 100 TCID₅₀ of virus in MDCK cells. Geometric mean neutralizing antibody titers were calculated for each group.

Histopathologic evaluation and immunohistochemical analysis

Formalin-inflated lungs and trachea were fixed in 10% neutral buffered formalin. Cross-sections of upper and lower left and right lung lobes and trachea were made, concentrating on gross-lesions. Tissue was paraffin-embedded and 6- μ m sections were stained with hematoxylin and eosin for histologic evaluation. Sequential sections were processed for immunohistochemistry or *in situ* hybridization (ISH). Immunohistochemistry was performed as described previously (Bissel *et al.*, *Am J Pathol* 160:927-941, 2002) using an immunoperoxidase method with a polyclonal antibody (Maine Biotechnology Services, Portland, ME) directed against influenza A virus. A biotinylated donkey anti-goat IgG (Rockland Immunochemicals, Gilbertsville, PA) was used as the secondary antibody. ISH was performed as described previously (Fallert *et al.*, *J Virol Methods* 99:23-32, 2002) using a 35S-labeled riboprobe synthesized using templates derived from 760 bp of influenza A/California/04/2009 matrix protein.

Results

Vaccine induced antibody responses

Cynomolgus macaques were vaccinated with COBRA VLPs or WS/05 VLPs formulated with Imject® alum at 0, 3 and 6 weeks. At week 3 post-vaccination, all COBRA VLP-vaccinated animals had anti-HA antibodies that recognized recombinant HA derived from three sub-clades of clade 2, which were boosted at week 6 (FIGS. 10A and 10B). There was no statistical difference ($p>0.05$) in the anti-HA titers elicited against any of the HA proteins, except monkeys vaccinated with COBRA VLPs had a statistically higher titer against the Indo/05 HA (clade 2.1) compared with monkeys vaccinated with the WS/05 VLP (derived from clade 2.2) on week 6.

A single COBRA VLP vaccination induced high titer HAI and MN antibodies to clade 2 H5N1 viruses

Monkeys vaccinated with COBRA VLPs (but not with WS/05 VLPs) had HAI activity against viruses representing all three clade 2 sub-clades after a single vaccination (FIG. 10C). Four to six monkeys responded to the COBRA VLP vaccine with an HAI titer $\geq 1:40$ for all of the various test antigens. In contrast, 4 of 7 monkeys vaccinated with the WS/05 VLP responded to the homologous clade 2.2 virus, but none of these vaccinated monkeys responded to the clade 2.1 or 2.3 virus. Following a second vaccination, almost all the monkeys vaccinated with either vaccine responded to all three viruses (FIG. 11D). These results were confirmed by microneutralization assay (FIGS. 11E and 11F). However, monkeys vaccinated with COBRA VLPs showed boosted HAI titers to all three clade 2 viruses (FIG. 11).

COBRA VLPs induced HAI antibodies that recognize broader numbers of H5N1 viruses

In order to determine if the COBRA HA elicited antibodies that recognized a broader number of H5N1 isolates, serum was collected and tested for the ability to inhibit influenza virus induced hemagglutination of red blood cells *in vitro*.

Antisera collected from both vaccinated and unvaccinated monkeys were then tested against a broad panel of H5N1 viruses representing not only sub-clades of clade 2, but also non-clade 2 H5N1 virus strains (0, 1, 4, and 7) by HAI. Monkeys vaccinated with the COBRA VLP had high average HAI titers against all clade 2 isolates, regardless of sub-clade (FIG. 11). In general, all 7 monkeys responded to the COBRA VLP vaccine and seroconverted with an HAI titer $\geq 1:40$ against all the clade 2 viruses. In contrast, monkeys vaccinated with the WS/05 VLP vaccine had lower HAI titers against clade 2 viruses (FIG. 10) and fewer number of monkeys responded to the vaccine. Of the 10 clade 2 viruses tested in the HAI assay, WS/05 VLP vaccinated monkeys responded more poorly (fewer than 4 of 7 monkeys) to 4 of the isolates and none of these monkeys had antibodies that responded to the Dk/HU/02 (clade 2.1.1) or Eg/3300/08 (clade 2.2.1) isolates. The COBRA VLPs elicited significantly higher HAI titers against almost all of the clade 2 viruses than the WS/05 VLPs (FIG. 11).

In addition to clade 2 isolates, a minimum of five COBRA VLP vaccinated monkeys had HAI antibodies against both clade 1 and 7 virus isolates (FIG. 11). In comparison, almost none of the WS/05 VLP vaccinated monkeys had HAI antibodies against clade 1 and clade 7 viruses. None of the monkeys, regardless of the vaccine, had antibodies that responded to the clade 0 or 4 isolates. All mock vaccinated monkeys did not recognize any of the H5N1 isolates.

Challenge of vaccinated and unvaccinated primates with H5N1 clade 2.2 virus

Three weeks after final vaccination, both VLP vaccinated and mock-vaccinated monkeys were transferred to ABSL3+ isolator units and then challenged with highly pathogenic H5N1 virus, A/Whooper Swan/Mongolia/244/2005 (clade 2.2) (1×10^6 pfu), by a multi-route (ocular, nasal, tracheal, oral) of infection (Kobasa *et al.*, *Nature* 445:319-323, 2007; Kuiken *et al.*, *Vet Pathol* 40:304-310, 2003; Rimmelzwaan *et al.*, *Avian Dis* 47:931-933, 2003). There was no significant weight loss or mortality in any of the monkeys over the 7 day period of observation. Unvaccinated monkeys had an elevated temperature of $\sim 2^{\circ}$ C that was sustained for

5 days post-infection and higher gross pathology scores by day 3 post-infection (Table 5).

Table 5. Lung pathology, temperature and viral titer of vaccinated macaques

Vaccine	Lung Pathology Score (day 3)	Elevated temperature (days)	Peak Viral Titer (pfu/ml) (day)
Mock	5.3	1.9°C (1-5 DPI)	Nasal wash: 2.2-2.5 (5 DPI) Trachea wash: 2.0-4.4 (3 DPI)
WS/05 VLP	3.3	1.1°C–1.3°C (1-5 DPI)	Nasal wash: <2 Trachea wash: <2
COBRA VLP	2.1	1.3°C (2 DPI)	Nasal wash: <2 Trachea wash: <2

The lungs of unvaccinated monkeys had mild to moderate acute pneumonia with alveolar pulmonary exudate by day 3 post-infection by H&E staining. ISH showed focal collections of H5N1 infected cells present at day 3 post-infection in alveolar spaces and to a lesser extent in bronchial epithelium. These results were similar to unvaccinated monkeys infected with the clade 1 H5N1 virus, A/Vietnam/1203/2004. In contrast, monkeys vaccinated with either the COBRA VLP or the WS/05 VLP vaccine had a reduced gross pathology scores of 2.1-3.3 at day 3 post-infection with a milder increase in body temperature (1.1-1.3⁰C) that spiked between days 2-3 post-infection and then returned to pre-infection temperatures. Vaccinated animals had fewer H5N1 infected cells that were detected primarily on day 1 post-infection (Table 6).

Table 6. H5N1 lung infection scores

	Vaccine			Alveolar infection score			Submucosal infection score		
	1	3	5	1	3	5	1	3	5
Mock	1.00	0.05	0	1.10	0.48	0.25	0	0	0
WS/05 VLP	0.05	0	0	0.55	0.10	0	0	0	0
COBRA VLP	0	0	0	0.60	0.03	0.05	0	0	0

ISH for influenza was performed on tissue sections of from upper and lower left and right lung. A semi-quantitative scoring system was developed to evaluate the presence of influenza infected cells. Scores were then averaged: 0.2 = rare or occasional cells but <5% of fields; 1 = >1/2 to 1/4 low power fields; 2 = >1/4 low power fields; 3 = essentially all low power fields.

However, monkeys vaccinated with the COBRA VLP had little to no signs of lung inflammation by H&E staining, while animals vaccinated with the WS/05 VLP vaccine had similar signs of inflammation as non-vaccinated monkeys (Table 7). In addition, unvaccinated monkeys had high titers of virus in both the nasal and tracheal washes between days 3 and 5 post-infection. In contrast, no virus was detected in either vaccinated groups.

Table 7. Lung involvement and inflammation scores

Vaccine	% lung involvement ^a			Bronchial inflammation ^b			Alveolar inflammation ^b		
	1	3	5	1	3	5	1	3	5
Mock	0.38 (0-1)	1.13 (0-2)	1.25 (0-2)	0.63 (0-1)	0.75 (0-2)	1.25 (0-2)	0.63 (0-1)	1.00 (0-2)	1.25 (0-2)
WS/05 VLP	0.75 (0-2)	1.50 (0-3)	0.88 (0-3)	1.00 (1)	1.42 (1-2)	0.63 (0-2)	1.00 (0-2)	1.25 (0-2)	1.00 (0-2)
COBRA VLP	0.88 (0-2)	0.50 (0-2)	0.38 (0-2)	1.13 (1-2)	0.75 (0-2)	0.88 (0-2)	1.13 (0-2)	0.67 (0-2)	0.25 (0-1)

^a % Lung involvement. Tissue sections from upper and lower left and right lung were evaluated for percent area demonstrating pneumonia. Scores were then averaged. Range in parentheses. 0 = <10%, 1 = 10-24%, 2 = 25-50%, 3 = >50%.

^b Bronchial and alveolar inflammation scores. Tissue sections from upper and lower left and right lung were evaluated for presence of bronchial inflammation and denudation and alveolar immune cell infiltration. Scores were then averages: 0 = absent, 1 = present, 2 = abundant.

Example 3: Comparison of Protective Efficacy by Vaccination with Computationally Optimized HA and Polyvalent HA Based H5N1 VLP Vaccines

This example describes a comparison of the COBRA HA vaccine to a polyvalent H5N1 vaccine. The results demonstrate that a single COBRA antigen elicits broader antibodies and is more effective than a polyvalent mixture of primary antigens.

Materials and Methods

Vaccine antigens and preparation

The design and characterization of the computationally optimized broadly reactive antigen (COBRA) is described in Example 1. Polyvalent vaccine HA antigens were derived via reverse transcription from the following 6:2 reassortant H5N1 viruses: A/Indonesia/5/2005 (clade 2.1; IN/05), A/Whooper Swan/Mongolia/244/2005 (clade 2.2; WS/05) and A/Anhui/1/2005 (clade 2.3; AN/05). All HA antigens were cloned into the pTR600 expression vector.

Virus-like particles (VLPs) were generated by transiently transfecting HEK 293T cells with plasmids expressing M1 (A/Puerto Rico/8/1934), NA (A/Thailand/1(KAN-1)/2004), and a single HA for each preparation. Cells were incubated for 72h at 37°C after which supernatants were harvested. Cell debris was cleared by low speed centrifugation followed by vacuum filtration through a 0.22 µm sterile filter. VLPs were purified by ultracentrifugation (100,000 x g through 20% glycerol, weight to volume) for 4 hours at 4°C. Pellets were then resuspended in PBS pH 7.2 and stored in single use aliquots at -80°C until use. Total protein concentration was determined by MicroBCA™ Protein Assay Reagent Kit (Pierce Biotechnology, Rockford, IL, USA). HA specific content of each VLP was determined by scanning densitometry as described previously (Giles and Ross, *Vaccine* 29:3043-3054, 2011). Briefly, purified HA matched to each VLP was electrophoresed with purified VLPs, transferred to a PVDF membrane and probed by western blot with H5-specific antisera. The relative density of the HA band in the purified protein lanes was used to calculate a standard curve and the density of the HA in the VLP lanes was interpolated. In total, four different VLP preparations

were purified and HA content quantified independently, each containing one of the three wild-type influenza gene products (IN/05, WS/05, AN/05) or the COBRA HA.

Mouse studies

BALB/c mice (*Mus musculus*, females, 6–8 weeks) were purchased from Harlan Sprague Dawley, (Indianapolis, IN, USA) and housed in microisolator units and allowed free access to food and water and were cared for under USDA guidelines for laboratory animals. Mice were vaccinated with purified COBRA VLPs (3 µg HA) or a polyvalent formulation of VLPs consisting of 1µg HA each IN/05, WS/05 and AN/05 (3 µg HA total) via intramuscular injection at week 0 and then boosted at week 3. Vaccines were formulated with Imject® alum adjuvant (Imject® Alum, Pierce Biotechnology; Rockford, IL, USA) according to the manufacturer's protocol. Fourteen to twenty-one days after each vaccination, blood was collected from anesthetized mice via the retro-orbital plexus and transferred to a microfuge tube. Tubes were centrifuged and sera was removed and frozen at $-20 \pm 5^{\circ}\text{C}$.

Three weeks after final vaccination, mice were challenged intranasally with 5×10^3 plaque forming units (PFU) of either highly pathogenic wild type H5N1 virus A/Whooper Swan/Mongolia/244/2005 (n=20/group) or 6:2 reassortant virus with internal genes from the mouse adapted virus A/Puerto Rico/8/1934 and the surface proteins HA and NA from A/Vietnam/1203/2004 (n=10/group) in a total volume of 50 µl. Challenge doses for both viruses were established independently and represent approximately 50LD₅₀. After infection, mice were monitored daily for weight loss, disease signs and death for 14 days after infection. Individual body weights, sickness scores and death were recorded for each group on each day after inoculation. Sickness score was determined by evaluating activity (0=normal, 1=reduced, 2=severely reduced), hunched back (0=absent, 1=present) and ruffled fur (0=absent, 1=present) (Toapanta and Ross, *Respiratory Res* 10(1):112, 2009). Experimental endpoint was determined by >20% weight loss or display of neurological disease such as hind limb paralysis. All highly pathogenic wild type

H5N1 influenza virus studies were performed under high-containment biosafety level 3 enhanced conditions (BSL3+).

Ferret studies

Fitch ferrets (*Mustela putorius furo*, female, 6-12-months of age), influenza naïve and descended, were purchased from Marshall Farms (Sayre, PA, USA). Ferrets were pair housed in stainless steel cages (Shor-line, Kansas City, KS, USA) containing Sani-chips Laboratory Animal Bedding (P.J. Murphy Forest Products, Montville, NJ, USA). Ferrets were provided with Teklad Global Ferret Diet (Harlan Teklad, Madison, WI, USA) and fresh water *ad libitum*. The VLPs were diluted in PBS, pH 7.2 to achieve final concentration. Ferrets (n=6) were vaccinated with purified COBRA VLPs (15 µg HA) or a polyvalent formulation of VLPs consisting of 5 µg HA each IN/05, WS/05 and AN/05 (15 µg HA total) via intramuscular injection at week 0 and then boosted at week 3. Vaccines were formulated with Imject® alum adjuvant (Imject® Alum, Pierce Biotechnology; Rockford, IL, USA) immediately prior to use according to the manufacturer's protocol. Animals were monitored for adverse events including weight loss, temperature, loss of activity, nasal discharge, sneezing and diarrhea weekly during the vaccination regimen. Prior to vaccination, animals were confirmed by HAI assay to be seronegative for circulating influenza A (H1N1 and H3N2) and influenza B viruses. Fourteen to twenty-one days after each vaccination, blood was collected from anesthetized ferrets via the anterior vena cava and transferred to a microfuge tube. Tubes were centrifuged and sera was removed and frozen at $-20 \pm 5^{\circ}\text{C}$.

Three weeks after final vaccination, ferrets were challenged intranasally with 1×10^6 plaque forming units (PFU) of the highly pathogenic H5N1 virus A/Whooper Swan/Mongolia/244/2005 (clade 2.2) in a volume of 0.5 ml in each nostril for a total infection volume of 1 ml. After infection, ferrets were monitored daily for weight loss, disease signs and death for 14 days after infection. Individual body weights, sickness scores, and death were recorded for each group on each day after inoculation. Sickness score was determined by evaluating activity (0=normal, 1=alert and active after stimulation, 2=alert but not active after stimulation,

3=neither active nor alert after stimulation), nasal discharge (0=absent, 1=present), sneezing (0=absent, 1=present), decreased food intake (0=absent, 1=present), diarrhea (0=absent, 1=present), dyspnea (0=absent, 1=present) and neurological symptoms (0=absent, 1=present) as previously described (Giles and Ross, *Vaccine* 29:3043-3054, 2011). Experimental endpoint was defined as >20% weight loss, development of neurological disease or an activity score of 3 (neither active nor alert after stimulation). Nasal washes were performed by instilling 3 ml of PBS into the nares of anesthetized ferrets each day for 14 days after inoculation. Washes were collected and stored at -80°C until use. All highly pathogenic wild type H5N1 influenza virus studies were performed under high-containment biosafety level 3 enhanced conditions (BSL3+).

ELISA assay

The ELISA assay was used to assess total antibody titer to the HA. High binding, 96-well polystyrene plates (Costar; Lowell, MA, USA) were coated overnight with 50 ng/well of recombinant HA. Coating antigens were derived from the following representative viral isolates: A/Vietnam/1203/2004 (clade 1), A/Indonesia/5/2005 (clade 2.1), A/Whooper Swan/Mongolia/244/2005 (clade 2.2) and A/Anhui/1/2005 (clade 2.3). Plates were blocked with 5% milk diluted in PBS with 0.05% Tween 20. Serum samples were diluted in blocking buffer and added to plates. Serum was two-fold serially diluted and allowed to incubate for 1 hour at room temperature. Plates were washed and species specific antibody against IgG linked to horseradish peroxidase (HRP) was diluted in blocking buffer and added to plates. Plates were incubated for 1 hour at room temperature. Plates were washed and HRP was developed with TMB substrate (Sigma-Aldrich; St. Louis, MO, USA). Plates were incubated in the dark for 15 minutes and then the reaction was stopped with 2N H₂SO₄. Optical densities at a wavelength of 450 nm (OD₄₅₀) were read by a spectrophotometer (BioTek; Winooski, VT, USA) and end point dilution titers were determined as the reciprocal dilution of the last well which had an OD₄₅₀ above the mean OD₄₅₀ plus two standard deviations of naïve animal sera.

Hemagglutination inhibition (HAI) assay

The HAI assay was used to assess functional antibodies to the HA able to inhibit agglutination of horse erythrocytes. The protocol was adapted from the CDC laboratory-based influenza surveillance manual (Gillim-Ross and Subbarao, *Clin Microbiol Rev* 19(4):614-636, 2006). To inactivate non-specific inhibitors, sera were treated with receptor destroying enzyme (RDE; Denka Seiken, Co., Japan) prior to being tested. Briefly, three parts RDE was added to one part sera and incubated overnight at 37°C. RDE was inactivated by incubation at 56°C for ~30 min. RDE treated sera was two-fold serially diluted in v-bottom microtiter plates. An equal volume of reassortant virus, adjusted to approximately 8 HAU/50µl, was added to each well. The reassortant viruses contained the internal genes from the mouse adapted strain A/Puerto Rico/8/1934 and the surface proteins HA and NA from the following representative viral isolates: A/Vietnam/1203/2004 (clade 1), A/Indonesia/5/2005 (clade 2.1), A/Whooper Swan/Mongolia/244/2005 (clade 2.2) and A/Anhui/1/2005 (clade 2.3). The plates were covered and incubated at room temperature for 20 minutes followed by the addition of 1% horse erythrocytes (HRBC) (Lampire Biologicals, Pipersville, PA, USA) in PBS. Red blood cells were stored at 4°C and used within 72 h of preparation. The plates were mixed by agitation, covered, and the RBCs were allowed to settle for 1 h at room temperature (Askonas B, McMichael A, Webster R. The immune response to influenza viruses and the problem of protection against infection. In: Beare AS, editor. Basic and applied influenza research: CRC Press 1982: 159-188). The HAI titer was determined by the reciprocal dilution of the last well which contained non-agglutinated RBC. Positive and negative serum controls were included for each plate. All mice and ferrets were negative (HAI \leq 1:10) for pre-existing antibodies to currently circulating human influenza viruses prior to vaccination.

Plaque assay

For mouse infections, lung virus titers were evaluated. For ferret infections, nasal wash virus titers were used to assess viral burden. Both lungs and nasal wash virus titers were determined using a plaque assay (Tobita *et al.*, *Med Microbiol*

Immunol 162:23-27, 1975; Tobita *et al.*, *Med Microbiol Immunol* 162:9-14, 1975). Briefly, lungs from mice infected with virus were harvested post infection, snap-frozen and stored at -80°C until use. Samples were thawed, weighed and single cell suspensions were prepared via passage through a 70 µm mesh (BD Falcon, Bedford, MA, USA) in an appropriate volume of DMEM supplemented with penicillin-streptomycin (iDEME) as to achieve 100 mg/ml final concentration. Cell suspensions were centrifuged at 2000 rpm for 5 minutes and the supernatants were collected.

Madin-Darby Canine Kidney (MDCK) cells were plated (5×10^5) in each well of a 6 well plate. Samples (lung supernatants for mice and nasal washes for ferrets) were diluted (dilution factors of 1×10^1 to 10^6) and overlaid onto the cells in 100 µl of iDMEM and incubated for 1 hour. Virus-containing medium was removed and replaced with 2 ml of L15 medium plus 0.8% agarose (Cambrex, East Rutherford, NJ, USA) and incubated for 96 hours at 37°C with 5% CO₂. Agarose was removed and discarded. Cells were fixed with 10% buffered formalin, and then stained with 1% crystal violet for 15 minutes. Following thorough washing in dH₂O to remove excess crystal violet, plates were allowed to dry, plaques counted, and the plaque forming units (PFU)/g for or PFU/ml for nasal washes were calculated.

Histopathological analysis

Left lobes of lungs from infected mice were collected 4 days post-infection and placed into 10% buffered formalin. After fixation, lungs were paraffin embedded and 6 µm sections were prepared for histopathological analysis. For *in situ* hybridization (ISH), vectors containing 760 bp of Influenza/California/04/2009 matrix protein were linearized to create antisense and sense templates. ³⁵S-labeled riboprobes were generated using MAXIscript *in vitro* transcription kit (Ambion, Austin, TX). ISH was performed as described before (Bissel *et al.*, *Brain Pathol*, Accepted Article doi: 10.1111/j.1750-3639.2010.00514.x). Control riboprobes did not hybridize to lung tissue at any time point post-infection and non-infected tissue did not show hybridization with viral probes. Hybridized slides were assessed and scored for abundance of foci.

Cellular assays

The number of anti-influenza specific cells secreting interferon gamma (IFN- γ) was determined by enzyme-linked immunospot (ELISpot) assay (R&D systems, Minneapolis, MN, USA) following the manufacturer's protocol. Mice were sacrificed at 6 days post infection (DPI) and spleens and lungs were harvested and prepared in single cell suspensions. Briefly, pre-coated anti-IFN γ plates were blocked with RPMI plus 10% FCS and antibiotics (cRPMI) for 30 minutes at room temperature. Media was removed from wells and 10^5 cells were added to each well. Cells were stimulated with purified recombinant HA from A/Vietnam/1203/2004 (truncated at residue 530; 1 μ g/well), inactivated 6:2 reassortant virus A/Vietnam/1203/2004 (1:100 dilution of inactivated stock; 100 μ l/well) or the immunodominant H2-K^d CD8⁺ T cell epitope in H5 HA: HA₅₃₃ (IYSTVASSL; SEQ ID NO: 10; 1 μ g/well) (Pepscan Presto, Leystad, Netherlands). Additional wells were stimulated with PMA (50 ng/well) and ionomycin (500 ng/well) as positive controls or Ova₂₅₇ (SIINFEKL; SEQ ID NO: 11; 1 μ g/well) (Pepscan Presto, Leystad, Netherlands) as negative controls. Additionally, IL-2 (10 U/ml) was added to each well. Plates were incubated at 37°C for 48 hours. After incubation, plates were washed four times with R&D wash buffer and were incubated at 4°C overnight with biotinylated anti-mouse IFN γ . Plates were washed as before and incubated at room temperature for 2 hours with streptavidin conjugated to alkaline phosphatase. Plates were washed as before and spots were developed by incubating at room temperature for 1 hour in the dark with BCIP/NBT chromogen substrate. The plates were washed extensively with DI H₂O and allowed to dry overnight prior to spots being counted using an ImmunoSpot ELISpot reader (Cellular Technology Ltd., Cleveland, OH, USA).

The number of anti-HA and anti-NA specific antibody secreting cells was determined by B cell ELISpot assay as previously described (Joo *et al.*, *Vaccine* 28:2186-2194, 2009; Sasaki *et al.*, *PLoS ONE* 3:e2975, 2008; Sasaki *et al.*, *J Virol* 81:215-228, 2007). Mice were sacrificed at 6 DPI and spleens and lungs were

harvested and prepared in single cell suspensions. Briefly, 0.45µm PVDF membrane plates (Millipore, Billerica, MA, USA) were coated with either purified recombinant HA from A/Vietnam/1203/2004 or purified recombinant NA from A/Thailand/1(KAN-1)/2004 (250 ng/well) and incubated at 4°C overnight. Plates were washed three times with PBS and blocked with cRPMI for at 37°C for 3-4 hours. Media was removed from wells and 10⁵ cells were added to each well. Plates were incubated at 37°C for 48 hours. After incubation, plates were washed as before and incubated at room temperature for 2 hours with horse radish peroxidase conjugated anti-mouse IgG or IgA (Southern Biotech, Birmingham, AL, USA). Plates were washed as before and spots were developed at room temperature for 1 hour in the dark with detection substrate (NovaRED™; Vector Labs, Burlingame, CA, USA). The plates were washed extensively with DI H₂O and allowed to dry overnight prior to spots being counted using an ImmunoSpot ELISpot reader (Cellular Technology Ltd., Cleveland, OH, USA).

Passive transfer of sera

Serum from vaccinated mice was pooled and passively transferred into 9 week old recipient BALB/c mice (n=5/group). Equal amounts of serum from each mouse in a particular vaccine group were pooled and heat inactivated for 30 minutes at 56°C. 200 µl of pooled and inactivated serum was transferred to recipient mice via IP injection. 24 hours post transfer, mice were infected with 6:2 reassortant virus with internal genes from the mouse adapted virus A/Puerto Rico/8/1934 and surface antigens from A/Vietnam/1203/2004.

Statistical analysis

Statistical significance of the antibody and cellular immunology data was determined using a two-tailed Student's T test to analyze differences between COBRA and polyvalent vaccine groups for each of the different test antigens. Differences in weight loss and sickness score were analyzed by two-way ANOVA, followed by Bonferroni's post test for each vaccine group at multiple time points (multiparametric). Statistical significance of viral titer data was evaluated using a

two-tailed Student's T test on Log_{10} transformed values. Significance was defined as $p < 0.05$. Statistical analyses were done using GraphPad Prism software.

Results

Immunogenicity in mice and ferrets

BALB/c mice were vaccinated twice via intramuscular injection with either purified COBRA or polyvalent VLPs and two weeks after the second vaccination serum was analyzed for antibody responses. All vaccinated mice had high titer anti-HA antibodies that bound to recombinant HA derived from both clade 1 and various sub-clades of clade 2 (FIG. 12A). Although both COBRA and polyvalent vaccines elicited similar total IgG, COBRA vaccinated animals had higher HAI antibody titers for all viruses tested ($p < 0.001$; FIG. 12B). In addition to higher HAI titer, COBRA vaccinated mice had an increased frequency of HAI titers $\geq 1:40$ for all viruses tested, including those which were components of the polyvalent formulation (Table 8).

To confirm the results from mice in a more rigorous animal model, ferrets were vaccinated twice via intramuscular injection with either COBRA or polyvalent vaccines. Serum was collected two weeks after the second vaccination and antibody responses were evaluated. Similar to the mice, all vaccinated ferrets had anti-HA antibodies that bound to diverse recombinant HA and the relative total IgG titers were equivalent for both COBRA and polyvalent vaccines (FIG. 12C). COBRA vaccinated ferrets demonstrated increased HAI antibody titers compared to polyvalent vaccinated animals against all viruses tested, however only the antibodies to the clade 2.1 virus were significantly different ($p < 0.05$; FIG. 12D). Furthermore, COBRA vaccinated animals displayed an increased rate of achieving an HAI titer of $\geq 1:40$ in comparison to the polyvalent vaccinated ferrets for all test antigens (Table 8).

Table 8. Seroconversion frequency

Species	Vaccine antigen	Clade 1	Clade 2.1	Clade 2.2	Clade 2.3
Mouse	COBRA	60% (18/30)	100% (30/30)	100% (30/30)	100% (30/30)
	Polyvalent	3.3% (1/30)	70% (21/30)	50% (15/30)	53% (16/30)
Ferret	COBRA	33% (2/6)	67% (4/6)	50% (3/6)	50% (3/6)
	Polyvalent	0% (0/6)	33% (2/6)	0% (0/6)	0% (0/6)

Wild type clade 2.2 challenge

To confirm protective efficacy against highly pathogenic H5N1 infection, vaccinated animals were challenged with a lethal dose of the wild-type clade 2.2 isolate A/Whooper Swan/Mongolia/244/2005. All VLP vaccinated mice were protected from weight loss and death while mock vaccinated animals rapidly lost weight and reached experimental end-point by 6 days post infection (DPI; FIG. 13A). COBRA and polyvalent vaccinated mice both had a mean maximum weight loss of 4% at 12 and 13 DPI, respectively. Additionally, all VLP vaccinated mice failed to develop any overt signs of disease while mock vaccinated mice developed visible illness (FIG. 13B).

Similar to the mice, all VLP vaccinated ferrets were protected from death following a lethal challenge. Vaccinated ferrets demonstrated mild weight loss in response to the infection with COBRA vaccinated animals having mean maximum weight loss of 5.5% at 2 DPI and polyvalent vaccinated animals losing 6.8% at 3 DPI (FIG. 13C). Both groups rapidly recovered weight and failed to develop any significant signs of disease (FIG. 13D). Furthermore, VLP vaccinated animals did not demonstrate any temperature spikes while mock vaccinated animals had an elevated temperature of ~3°C for 1-3 DPI.

To evaluate vaccine efficacy with a more sensitive output than morbidity and mortality, the viral burden of infected animals was also determined. Both COBRA and polyvalent vaccinated mice had reduced lung viral titers as quickly as 1 DPI when compared to mock vaccinated animals. Furthermore, COBRA vaccinated mice did not have detectable virus by 3 DPI while polyvalent vaccinated mice demonstrated prolonged viral replication with 1.8×10^3 PFU/g at 3DPI ($p < 0.05$;

FIG. 14A). Additionally, both VLP vaccines prevented extra-pulmonary spread of the virus while mock vaccinated animals had detectable virus in both kidney and liver by 3 DPI. Control of virus replication in ferrets was similar to that observed in mice, although complete clearance of the virus was delayed (FIG. 14B). All VLP vaccinated animals had decreased recovery of virus in nasal washes compared to mock vaccinated ferrets at all timepoints tested ($p<0.05$). COBRA vaccinated animals did not have detectable virus by 5 DPI. In contrast, virus replication did not reach undetectable levels until 9DPI in polyvalent vaccinated ferrets.

Histopathology of infected lungs

To evaluate the location and severity of influenza viral antigen and viral replication, ISH for influenza A MP was scored on 3 DPI lung sections. COBRA vaccinated animals had rare bronchial epithelium infection (FIGS. 15A and 15B). Animals receiving polyvalent vaccines had occasional bronchial epithelium infection that was comparable to the COBRA vaccinated animals (FIGS. 15A and 15B). This was in contrast to significant bronchial epithelium infection and replication observed in mock animals (FIGS. 15A and 15B).

Reassortant clade 1 challenge

Having established the clade 2.2 protective profile of both the COBRA and polyvalent vaccines, the efficacy of these vaccines against a more divergent clade 1 challenge in mice was evaluated. COBRA and polyvalent vaccinated mice were challenged with 6:2 reassortant virus containing the HA and NA proteins from the clade 1 virus A/Vietnam/1203/2004. All VLP vaccinated animals were protected from weight loss and death while mock vaccinated animals rapidly lost weight and reached experimental endpoint by 7 DPI (FIG. 16A). Furthermore, vaccinated mice also did not develop any signs of disease throughout the course of the study (FIG. 16B). Lungs were harvested at 3 DPI for determination of viral burden (FIG. 16C). COBRA vaccinated animals did not have detectable virus while polyvalent animals had 1.1×10^3 PFU/g virus ($p=0.12$). Importantly, both vaccines had significantly less recoverable virus than mock vaccinated animals at 3 DPI ($p<0.01$).

Post-challenge cellular immune responses

The magnitude of influenza specific cellular immune responses in the lungs post-infection was evaluated via ELISpot assay for both antibody secreting cells (ASC) and IFN- γ producing cells. Vaccinated mice were infected with reassortant A/Vietnam/1203/2994 virus as before and lungs were harvested at 6 DPI. COBRA and polyvalent vaccinated animals had statistically equivalent numbers of both IgG and IgA ASC specific for HA from the challenge virus ($p>0.05$; FIG. 17A). No ASC were detected in mock vaccinated animals indicating that the 6 DPI time point is likely representative of a recall response. Additionally, the majority of the ASC response to infection was specific for HA as lower numbers of cells were detected for the NA component of the vaccines.

VLP vaccine primed IFN- γ secreting cells were also evaluated after infection. IFN- γ responses were equivalent between VLP vaccine groups regardless of stimulating antigen ($p>0.05$; FIG. 17B). Recombinant HA and inactivated virus were inefficient stimulators of IFN- γ production compared to the HA₅₃₃ peptide. HA₅₃₃ is the immunodominant CD8⁺ T cell epitope in BALB/c mice and is conserved in all HA vaccine antigens used in this study. Overlapping peptide pools spanning the entire HA molecule were also used to stimulate cells and no differences were observed between COBRA and polyvalent vaccines for any of the pools. Similar to the ASC data, no IFN- γ responses were detectable above background in mock vaccinated animals at 6 DPI.

Passive transfer of immune sera

The contribution of serum factors to protection from clade 1 challenge was evaluated using a passive transfer model. Nine-week old recipient mice were administered pooled sera via IP injection from COBRA, polyvalent and mock vaccinated mice. The next day, recipient mice were challenged with the clade 1 reassortant A/Vietnam/1203/2004 virus as before. Regardless of transferred serum, all recipient mice lost weight and became visibly ill (FIGS. 18A and 18B). COBRA serum recipient mice lost less weight than polyvalent recipient mice with maximum

losses of 5.2% (6 DPI) and 11.8% (7 DPI), respectively ($p < 0.05$ at 7 DPI). COBRA serum recipient mice also began to resolve the clinical symptoms more rapidly than polyvalent recipient mice ($p < 0.05$ at 7 DPI). Although COBRA serum prevented recipient mice from developing illness more efficiently than polyvalent serum, both COBRA and polyvalent serum protected all recipient mice from death. Conversely, all mice receiving serum from mock vaccinated mice rapidly lost weight, became visibly ill and reached experimental endpoint by 7 DPI.

In view of the many possible embodiments to which the principles of the disclosed invention may be applied, it should be recognized that the illustrated embodiments are only examples of the disclosure and should not be taken as limiting the scope of the invention. Rather, the scope of the invention is defined by the following claims. We therefore claim as our invention all that comes within the scope and spirit of these claims.

SEQUENCE LISTING IN ELECTRONIC FORM

In accordance with Section 111(1) of the Patent Rules, this description contains a sequence listing in electronic form in ASCII text format (file: 63198-1692 Seq 30-MAR-13 vl.txt).

A copy of the sequence listing in electronic form is available from the Canadian Intellectual Property Office.

The sequences in the sequence listing in electronic form are reproduced in the following table.

SEQUENCE TABLE

<110> UNIVERSITY OF PITTSBURGH - OF THE COMMONWEALTH SYSTEM OF
HIGHER EDUCATION
Ross, Ted M.
Giles, Brendan M.

<120> COMPUTATIONALLY OPTIMIZED BROADLY REACTIVE ANTIGENS FOR INFLUENZA

<130> 63198-1692

<140> CA national phase of PCT/US2011/051072

<141> 2011-09-09

<150> US 61/403,407

<151> 2010-09-14

<160> 11

<170> PatentIn version 3.5

<210> 1

<211> 1707

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic polynucleotide

<220>

<221> CDS

<222> (1)..(1707)

<400> 1

atg gaa aag atc gtg ctg ctg ctg gct atc gtg agc ctg gtg aag agc	48
Met Glu Lys Ile Val Leu Leu Leu Ala Ile Val Ser Leu Val Lys Ser	
1 5 10 15	

gac cag att tgc atc ggc tac cac gcc aac aac agc acc gag cag gtg	96
Asp Gln Ile Cys Ile Gly Tyr His Ala Asn Asn Ser Thr Glu Gln Val	
20 25 30	

gac acc atc atg gaa aag aac gtc acc gtg acc cac gcc cag gac atc	144
Asp Thr Ile Met Glu Lys Asn Val Thr Val Thr His Ala Gln Asp Ile	
35 40 45	

ctg gaa aag acc cac aac ggc aag ctg tgc gac ctg gac ggc gtg aag	192
Leu Glu Lys Thr His Asn Gly Lys Leu Cys Asp Leu Asp Gly Val Lys	
50 55 60	

ccc ctg atc ctg agg gac tgc agc gtg gcc ggc tgg ctg ctg ggc aac	240
Pro Leu Ile Leu Arg Asp Cys Ser Val Ala Gly Trp Leu Leu Gly Asn	
65 70 75 80	

ccc atg tgc gac gag ttc atc aac gtg ccc gag tgg agc tac atc gtg	288
Pro Met Cys Asp Glu Phe Ile Asn Val Pro Glu Trp Ser Tyr Ile Val	
85 90 95	

gag aag gcc aac ccc gcc aac gac ctg tgc tac ccc ggc aac ttc aac	336
Glu Lys Ala Asn Pro Ala Asn Asp Leu Cys Tyr Pro Gly Asn Phe Asn	
100 105 110	

gac tac gag gaa ctg aag cac ctg ctg tcc agg atc aac cac ttc gag	384
Asp Tyr Glu Glu Leu Lys His Leu Leu Ser Arg Ile Asn His Phe Glu	
115 120 125	

aag atc cag atc atc ccc aag agc agc tgg tcc gac cac gag gcc agc	432
Lys Ile Gln Ile Ile Pro Lys Ser Ser Trp Ser Asp His Glu Ala Ser	
130 135 140	

agc ggc gtg agc agc gcc tgc cca tac cag ggc agc ccc agc ttc ttc	480
Ser Gly Val Ser Ser Ala Cys Pro Tyr Gln Gly Ser Pro Ser Phe Phe	
145 150 155 160	
aga aac gtg gtg tgg ctg atc aag aag aac aac acc tac ccc acc atc	528
Arg Asn Val Val Trp Leu Ile Lys Lys Asn Asn Thr Tyr Pro Thr Ile	
165 170 175	
aag agg tcc tac aac aac acc aac cag gaa gat ctg ctg gtc ctg tgg	576
Lys Arg Ser Tyr Asn Asn Thr Asn Gln Glu Asp Leu Leu Val Leu Trp	
180 185 190	
ggc atc cac cac cct aat gac gcc gcc gaa cag acc agg ctg tac cag	624
Gly Ile His His Pro Asn Asp Ala Ala Glu Gln Thr Arg Leu Tyr Gln	
195 200 205	
aac ccc acc acc tac atc agc gtg ggc aca agc acc ctg aac cag agg	672
Asn Pro Thr Thr Tyr Ile Ser Val Gly Thr Ser Thr Leu Asn Gln Arg	
210 215 220	
ctg gtg ccc aag atc gcc acc agg tcc aag gtg aac gga cag tcc ggc	720
Leu Val Pro Lys Ile Ala Thr Arg Ser Lys Val Asn Gly Gln Ser Gly	
225 230 235 240	
agg atg gaa ttc ttc tgg acc atc ctg aag cct aac gac gcc atc aac	768
Arg Met Glu Phe Phe Trp Thr Ile Leu Lys Pro Asn Asp Ala Ile Asn	
245 250 255	
ttc gag agc aac ggc aac ttt atc gcc ccc gag tac gcc tac aag atc	816
Phe Glu Ser Asn Gly Asn Phe Ile Ala Pro Glu Tyr Ala Tyr Lys Ile	
260 265 270	
gtg aag aag ggc gac agc gcc atc atg aag agc gag ctg gaa tac ggc	864
Val Lys Lys Gly Asp Ser Ala Ile Met Lys Ser Glu Leu Glu Tyr Gly	
275 280 285	
aac tgc aac acc aag tgc cag acc ccc atc ggc gcc atc aac agc agc	912
Asn Cys Asn Thr Lys Cys Gln Thr Pro Ile Gly Ala Ile Asn Ser Ser	
290 295 300	
atg ccc ttc cac aac atc cac ccc ctg acc atc ggc gag tgc ccc aag	960
Met Pro Phe His Asn Ile His Pro Leu Thr Ile Gly Glu Cys Pro Lys	
305 310 315 320	
tac gtg aag agc aac agg ctg gtg ctg gcc acc ggc ctg agg aac agc	1008
Tyr Val Lys Ser Asn Arg Leu Val Leu Ala Thr Gly Leu Arg Asn Ser	
325 330 335	
ccc cag aga gag agc aga aga aag aag agg ggc ctg ttc ggc gct atc	1056
Pro Gln Arg Glu Ser Arg Arg Lys Lys Arg Gly Leu Phe Gly Ala Ile	
340 345 350	
gcc ggc ttc atc gag ggc ggc tgg cag ggc atg gtg gac ggg tgg tac	1104
Ala Gly Phe Ile Glu Gly Gly Trp Gln Gly Met Val Asp Gly Trp Tyr	
355 360 365	

ggc tac cac cac tct aac gag cag ggc agc ggc tac gcc gcc gac aaa 1152
 Gly Tyr His His Ser Asn Glu Gln Gly Ser Gly Tyr Ala Ala Asp Lys
 370 375 380

gag agc acc cag aag gcc atc gac ggc gtc acc aac aag gtg aac agc 1200
 Glu Ser Thr Gln Lys Ala Ile Asp Gly Val Thr Asn Lys Val Asn Ser
 385 390 395 400

atc atc gac aag atg aac acc cag ttc gag gcc gtg ggc aga gag ttc 1248
 Ile Ile Asp Lys Met Asn Thr Gln Phe Glu Ala Val Gly Arg Glu Phe
 405 410 415

aac aac ctg gaa agg cgg atc gag aac ctg aac aag aaa atg gaa gat 1296
 Asn Asn Leu Glu Arg Arg Ile Glu Asn Leu Asn Lys Lys Met Glu Asp
 420 425 430

ggc ttc ctg gac gtg tgg acc tac aac gcc gag ctg ctg gtg ctg atg 1344
 Gly Phe Leu Asp Val Trp Thr Tyr Asn Ala Glu Leu Leu Val Leu Met
 435 440 445

gaa aac gag agg acc ctg gac ttc cac gac agc aac gtg aag aac ctg 1392
 Glu Asn Glu Arg Thr Leu Asp Phe His Asp Ser Asn Val Lys Asn Leu
 450 455 460

tac gac aaa gtg cgg ctg cag ctg agg gac aac gcc aaa gag ctg ggc 1440
 Tyr Asp Lys Val Arg Leu Gln Leu Arg Asp Asn Ala Lys Glu Leu Gly
 465 470 475 480

aac ggc tgc ttc gag ttc tac cac aag tgc gac aac gag tgc atg gaa 1488
 Asn Gly Cys Phe Glu Phe Tyr His Lys Cys Asp Asn Glu Cys Met Glu
 485 490 495

agc gtg agg aac ggc acc tac gac tac ccc cag tac agc gag gaa gcc 1536
 Ser Val Arg Asn Gly Thr Tyr Asp Tyr Pro Gln Tyr Ser Glu Glu Ala
 500 505 510

agg ctg aag agg gaa gag atc agc gga gtg aag ctg gaa agc atc ggc 1584
 Arg Leu Lys Arg Glu Glu Ile Ser Gly Val Lys Leu Glu Ser Ile Gly
 515 520 525

acc tac cag atc ctg agc atc tac agc acc gtc gcc agc agc ctg gcc 1632
 Thr Tyr Gln Ile Leu Ser Ile Tyr Ser Thr Val Ala Ser Ser Leu Ala
 530 535 540

ctg gct atc atg gtg gcc gga ctg agc ctg tgg atg tgc agc aac ggc 1680
 Leu Ala Ile Met Val Ala Gly Leu Ser Leu Trp Met Cys Ser Asn Gly
 545 550 555 560

agc ctg cag tgc agg atc tgc atc tga 1707
 Ser Leu Gln Cys Arg Ile Cys Ile
 565

<210> 2
 <211> 568
 <212> PRT
 <213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 2

Met Glu Lys Ile Val Leu Leu Leu Ala Ile Val Ser Leu Val Lys Ser
 1 5 10 15
 Asp Gln Ile Cys Ile Gly Tyr His Ala Asn Asn Ser Thr Glu Gln Val
 20 25 30
 Asp Thr Ile Met Glu Lys Asn Val Thr Val Thr His Ala Gln Asp Ile
 35 40 45
 Leu Glu Lys Thr His Asn Gly Lys Leu Cys Asp Leu Asp Gly Val Lys
 50 55 60
 Pro Leu Ile Leu Arg Asp Cys Ser Val Ala Gly Trp Leu Leu Gly Asn
 65 70 75 80
 Pro Met Cys Asp Glu Phe Ile Asn Val Pro Glu Trp Ser Tyr Ile Val
 85 90 95
 Glu Lys Ala Asn Pro Ala Asn Asp Leu Cys Tyr Pro Gly Asn Phe Asn
 100 105 110
 Asp Tyr Glu Glu Leu Lys His Leu Leu Ser Arg Ile Asn His Phe Glu
 115 120 125
 Lys Ile Gln Ile Ile Pro Lys Ser Ser Trp Ser Asp His Glu Ala Ser
 130 135 140
 Ser Gly Val Ser Ser Ala Cys Pro Tyr Gln Gly Ser Pro Ser Phe Phe
 145 150 155 160
 Arg Asn Val Val Trp Leu Ile Lys Lys Asn Asn Thr Tyr Pro Thr Ile
 165 170 175
 Lys Arg Ser Tyr Asn Asn Thr Asn Gln Glu Asp Leu Leu Val Leu Trp
 180 185 190
 Gly Ile His His Pro Asn Asp Ala Ala Glu Gln Thr Arg Leu Tyr Gln
 195 200 205
 Asn Pro Thr Thr Tyr Ile Ser Val Gly Thr Ser Thr Leu Asn Gln Arg
 210 215 220
 Leu Val Pro Lys Ile Ala Thr Arg Ser Lys Val Asn Gly Gln Ser Gly
 225 230 235 240
 Arg Met Glu Phe Phe Trp Thr Ile Leu Lys Pro Asn Asp Ala Ile Asn
 245 250 255
 Phe Glu Ser Asn Gly Asn Phe Ile Ala Pro Glu Tyr Ala Tyr Lys Ile
 260 265 270
 Val Lys Lys Gly Asp Ser Ala Ile Met Lys Ser Glu Leu Glu Tyr Gly
 275 280 285
 Asn Cys Asn Thr Lys Cys Gln Thr Pro Ile Gly Ala Ile Asn Ser Ser
 290 295 300
 Met Pro Phe His Asn Ile His Pro Leu Thr Ile Gly Glu Cys Pro Lys
 305 310 315 320
 Tyr Val Lys Ser Asn Arg Leu Val Leu Ala Thr Gly Leu Arg Asn Ser
 325 330 335
 Pro Gln Arg Glu Ser Arg Arg Lys Lys Arg Gly Leu Phe Gly Ala Ile
 340 345 350
 Ala Gly Phe Ile Glu Gly Gly Trp Gln Gly Met Val Asp Gly Trp Tyr
 355 360 365
 Gly Tyr His His Ser Asn Glu Gln Gly Ser Gly Tyr Ala Ala Asp Lys
 370 375 380
 Glu Ser Thr Gln Lys Ala Ile Asp Gly Val Thr Asn Lys Val Asn Ser
 385 390 395 400
 Ile Ile Asp Lys Met Asn Thr Gln Phe Glu Ala Val Gly Arg Glu Phe
 405 410 415

Asn Asn Leu Glu Arg Arg Ile Glu Asn Leu Asn Lys Lys Met Glu Asp
 420 425 430
 Gly Phe Leu Asp Val Trp Thr Tyr Asn Ala Glu Leu Leu Val Leu Met
 435 440 445
 Glu Asn Glu Arg Thr Leu Asp Phe His Asp Ser Asn Val Lys Asn Leu
 450 455 460
 Tyr Asp Lys Val Arg Leu Gln Leu Arg Asp Asn Ala Lys Glu Leu Gly
 465 470 475 480
 Asn Gly Cys Phe Glu Phe Tyr His Lys Cys Asp Asn Glu Cys Met Glu
 485 490 495
 Ser Val Arg Asn Gly Thr Tyr Asp Tyr Pro Gln Tyr Ser Glu Glu Ala
 500 505 510
 Arg Leu Lys Arg Glu Glu Ile Ser Gly Val Lys Leu Glu Ser Ile Gly
 515 520 525
 Thr Tyr Gln Ile Leu Ser Ile Tyr Ser Thr Val Ala Ser Ser Leu Ala
 530 535 540
 Leu Ala Ile Met Val Ala Gly Leu Ser Leu Trp Met Cys Ser Asn Gly
 545 550 555 560
 Ser Leu Gln Cys Arg Ile Cys Ile
 565

<210> 3
 <211> 1350
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Synthetic polynucleotide

<220>
 <221> CDS
 <222> (1)..(1350)

<400> 3
 atg aat cct aat aag aag atc atc aca atc gga agc atc tgc atg gtg 48
 Met Asn Pro Asn Lys Lys Ile Ile Thr Ile Gly Ser Ile Cys Met Val
 1 5 10 15
 aca gga atg gtg agc ctg atg ctg cag atc gga aat ctg atc agc atc 96
 Thr Gly Met Val Ser Leu Met Leu Gln Ile Gly Asn Leu Ile Ser Ile
 20 25 30
 tgg gtg agc cac agc atc cac aca gga aat cag cac aag gcc gag cct 144
 Trp Val Ser His Ser Ile His Thr Gly Asn Gln His Lys Ala Glu Pro
 35 40 45
 atc agc aat aca aat ttt ctg aca gag aag gcc gtg gcc agc gtg aag 192
 Ile Ser Asn Thr Asn Phe Leu Thr Glu Lys Ala Val Ala Ser Val Lys
 50 55 60
 ctg gcc gga aat agc agc ctg tgc cct atc aat gga tgg gcc gtg tac 240
 Leu Ala Gly Asn Ser Ser Leu Cys Pro Ile Asn Gly Trp Ala Val Tyr
 65 70 75 80

agc aag gat aat agc atc aga atc gga agc aag gga gat gtg ttt gtg	288
Ser Lys Asp Asn Ser Ile Arg Ile Gly Ser Lys Gly Asp Val Phe Val	
85 90 95	
atc aga gag cct ttt atc agc tgc agc cac ctg gag tgc aga aca ttt	336
Ile Arg Glu Pro Phe Ile Ser Cys Ser His Leu Glu Cys Arg Thr Phe	
100 105 110	
ttt ctg aca cag gga gcc ctg ctg aat gat aag cac agc aat gga aca	384
Phe Leu Thr Gln Gly Ala Leu Leu Asn Asp Lys His Ser Asn Gly Thr	
115 120 125	
gtg aag gat aga agc cct cac aga aca ctg atg agc tgc cct gtg gga	432
Val Lys Asp Arg Ser Pro His Arg Thr Leu Met Ser Cys Pro Val Gly	
130 135 140	
gag gcc cct agc cct tac aat agc aga ttt gag agc gtg gcc tgg agc	480
Glu Ala Pro Ser Pro Tyr Asn Ser Arg Phe Glu Ser Val Ala Trp Ser	
145 150 155 160	
gcc agc gcc tgc cac gat gga aca agc tgg ctg aca atc gga atc agc	528
Ala Ser Ala Cys His Asp Gly Thr Ser Trp Leu Thr Ile Gly Ile Ser	
165 170 175	
gga cct gat aat gga gcc gtg gcc gtg ctg aag tac aat gga atc atc	576
Gly Pro Asp Asn Gly Ala Val Ala Val Leu Lys Tyr Asn Gly Ile Ile	
180 185 190	
aca gat aca atc aag agc tgg aga aat aat atc ctg aga aca cag gag	624
Thr Asp Thr Ile Lys Ser Trp Arg Asn Asn Ile Leu Arg Thr Gln Glu	
195 200 205	
agc gag tgc gcc tgc gtg aat gga agc tgc ttt aca gtg atg aca gat	672
Ser Glu Cys Ala Cys Val Asn Gly Ser Cys Phe Thr Val Met Thr Asp	
210 215 220	
gga cct agc aat gga cag gcc agc cac aag atc ttt aag atg gag aag	720
Gly Pro Ser Asn Gly Gln Ala Ser His Lys Ile Phe Lys Met Glu Lys	
225 230 235 240	
gga aag gtg gtg aag agc gtg gag ctg gat gcc cct aat tac cac tac	768
Gly Lys Val Val Lys Ser Val Glu Leu Asp Ala Pro Asn Tyr His Tyr	
245 250 255	
gag gag tgc agc tgc tac cct gat gcc gga gag atc aca tgc gtg tgc	816
Glu Glu Cys Ser Cys Tyr Pro Asp Ala Gly Glu Ile Thr Cys Val Cys	
260 265 270	
aga gat aat tgg cac gga agc aat aga cct tgg gtg agc ttt aat cag	864
Arg Asp Asn Trp His Gly Ser Asn Arg Pro Trp Val Ser Phe Asn Gln	
275 280 285	
aat ctg gag tac cag atc gga tac atc tgc agc gga gtg ttt gga gat	912
Asn Leu Glu Tyr Gln Ile Gly Tyr Ile Cys Ser Gly Val Phe Gly Asp	
290 295 300	

aat cct aga cct aat gat gga aca gga agc tgc gga cct gtg agc agc 960
 Asn Pro Arg Pro Asn Asp Gly Thr Gly Ser Cys Gly Pro Val Ser Ser
 305 310 315 320

aat gga gcc tac gga gtg aag gga ttt agc ttt aag tac gga aat gga 1008
 Asn Gly Ala Tyr Gly Val Lys Gly Phe Ser Phe Lys Tyr Gly Asn Gly
 325 330 335

gtg tgg atc gga aga aca aag agc aca aat agc aga agc gga ttt gag 1056
 Val Trp Ile Gly Arg Thr Lys Ser Thr Asn Ser Arg Ser Gly Phe Glu
 340 345 350

atg atc tgg gac cct aat gga tgg aca gag aca gat agc agc ttt agc 1104
 Met Ile Trp Asp Pro Asn Gly Trp Thr Glu Thr Asp Ser Ser Phe Ser
 355 360 365

gtg aag cag gat atc gtg gcc atc aca gat tgg agc gga tac agc gga 1152
 Val Lys Gln Asp Ile Val Ala Ile Thr Asp Trp Ser Gly Tyr Ser Gly
 370 375 380

agc ttt gtg cag cac cct gag ctg aca gga ctg gat tgc atc aga cct 1200
 Ser Phe Val Gln His Pro Glu Leu Thr Gly Leu Asp Cys Ile Arg Pro
 385 390 395 400

tgc ttt tgg gtg gag ctg atc aga gga aga cct aag gag agc aca atc 1248
 Cys Phe Trp Val Glu Leu Ile Arg Gly Arg Pro Lys Glu Ser Thr Ile
 405 410 415

tgg aca agc gga agc agc atc agc ttt tgc gga gtg aat agc gat aca 1296
 Trp Thr Ser Gly Ser Ser Ile Ser Phe Cys Gly Val Asn Ser Asp Thr
 420 425 430

gtg gga tgg agc tgg cct gat gga gcc gag ctg cct ttt aca atc gat 1344
 Val Gly Trp Ser Trp Pro Asp Gly Ala Glu Leu Pro Phe Thr Ile Asp
 435 440 445

aag tga 1350
 Lys

<210> 4
 <211> 449
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Synthetic Construct

<400> 4
 Met Asn Pro Asn Lys Lys Ile Ile Thr Ile Gly Ser Ile Cys Met Val
 1 5 10 15
 Thr Gly Met Val Ser Leu Met Leu Gln Ile Gly Asn Leu Ile Ser Ile
 20 25 30
 Trp Val Ser His Ser Ile His Thr Gly Asn Gln His Lys Ala Glu Pro
 35 40 45
 Ile Ser Asn Thr Asn Phe Leu Thr Glu Lys Ala Val Ala Ser Val Lys
 50 55 60

Leu Ala Gly Asn Ser Ser Leu Cys Pro Ile Asn Gly Trp Ala Val Tyr
 65 70 75 80
 Ser Lys Asp Asn Ser Ile Arg Ile Gly Ser Lys Gly Asp Val Phe Val
 85 90 95
 Ile Arg Glu Pro Phe Ile Ser Cys Ser His Leu Glu Cys Arg Thr Phe
 100 105 110
 Phe Leu Thr Gln Gly Ala Leu Leu Asn Asp Lys His Ser Asn Gly Thr
 115 120 125
 Val Lys Asp Arg Ser Pro His Arg Thr Leu Met Ser Cys Pro Val Gly
 130 135 140
 Glu Ala Pro Ser Pro Tyr Asn Ser Arg Phe Glu Ser Val Ala Trp Ser
 145 150 155 160
 Ala Ser Ala Cys His Asp Gly Thr Ser Trp Leu Thr Ile Gly Ile Ser
 165 170 175
 Gly Pro Asp Asn Gly Ala Val Ala Val Leu Lys Tyr Asn Gly Ile Ile
 180 185 190
 Thr Asp Thr Ile Lys Ser Trp Arg Asn Asn Ile Leu Arg Thr Gln Glu
 195 200 205
 Ser Glu Cys Ala Cys Val Asn Gly Ser Cys Phe Thr Val Met Thr Asp
 210 215 220
 Gly Pro Ser Asn Gly Gln Ala Ser His Lys Ile Phe Lys Met Glu Lys
 225 230 235 240
 Gly Lys Val Val Lys Ser Val Glu Leu Asp Ala Pro Asn Tyr His Tyr
 245 250 255
 Glu Glu Cys Ser Cys Tyr Pro Asp Ala Gly Glu Ile Thr Cys Val Cys
 260 265 270
 Arg Asp Asn Trp His Gly Ser Asn Arg Pro Trp Val Ser Phe Asn Gln
 275 280 285
 Asn Leu Glu Tyr Gln Ile Gly Tyr Ile Cys Ser Gly Val Phe Gly Asp
 290 295 300
 Asn Pro Arg Pro Asn Asp Gly Thr Gly Ser Cys Gly Pro Val Ser Ser
 305 310 315 320
 Asn Gly Ala Tyr Gly Val Lys Gly Phe Ser Phe Lys Tyr Gly Asn Gly
 325 330 335
 Val Trp Ile Gly Arg Thr Lys Ser Thr Asn Ser Arg Ser Gly Phe Glu
 340 345 350
 Met Ile Trp Asp Pro Asn Gly Trp Thr Glu Thr Asp Ser Ser Phe Ser
 355 360 365
 Val Lys Gln Asp Ile Val Ala Ile Thr Asp Trp Ser Gly Tyr Ser Gly
 370 375 380
 Ser Phe Val Gln His Pro Glu Leu Thr Gly Leu Asp Cys Ile Arg Pro
 385 390 395 400
 Cys Phe Trp Val Glu Leu Ile Arg Gly Arg Pro Lys Glu Ser Thr Ile
 405 410 415
 Trp Thr Ser Gly Ser Ser Ile Ser Phe Cys Gly Val Asn Ser Asp Thr
 420 425 430
 Val Gly Trp Ser Trp Pro Asp Gly Ala Glu Leu Pro Phe Thr Ile Asp
 435 440 445
 Lys

<210> 5
 <211> 759
 <212> DNA
 <213> Artificial Sequence

<220>

<223> Synthetic polynucleotide

<220>

<221> CDS

<222> (1)..(759)

<400> 5

atg agc ctg ctg acc gag gtg gag aca tac gtg ctg tcc atc atc ccc	48
Met Ser Leu Leu Thr Glu Val Glu Thr Tyr Val Leu Ser Ile Ile Pro	
1 5 10 15	
agc ggc cct ctg aag gcc gag atc gcc cag aga ctg gaa gat gtg ttc	96
Ser Gly Pro Leu Lys Ala Glu Ile Ala Gln Arg Leu Glu Asp Val Phe	
20 25 30	
gcc ggc aag aac acc gac ctg gaa gtg ctg atg gaa tgg ctg aaa acc	144
Ala Gly Lys Asn Thr Asp Leu Glu Val Leu Met Glu Trp Leu Lys Thr	
35 40 45	
aga ccc atc ctg agc cct ctg acc aag ggc atc ctg ggc ttc gtg ttc	192
Arg Pro Ile Leu Ser Pro Leu Thr Lys Gly Ile Leu Gly Phe Val Phe	
50 55 60	
acc ctg acc gtg ccc agc gag aga ggc ctg cag agg cgg aga ttc gtg	240
Thr Leu Thr Val Pro Ser Glu Arg Gly Leu Gln Arg Arg Arg Phe Val	
65 70 75 80	
cag aac gcc ctg aac ggc aac ggc gac ccc aac aac atg gac aag gcc	288
Gln Asn Ala Leu Asn Gly Asn Gly Asp Pro Asn Asn Met Asp Lys Ala	
85 90 95	
gtg aag ctg tac aga aag ctg aag cgg gag atc acc ttc cac ggc gcc	336
Val Lys Leu Tyr Arg Lys Leu Lys Arg Glu Ile Thr Phe His Gly Ala	
100 105 110	
aaa gag atc agc ctg agc tac agc gct ggc gcc ctg gcc agc tgc atg	384
Lys Glu Ile Ser Leu Ser Tyr Ser Ala Gly Ala Leu Ala Ser Cys Met	
115 120 125	
ggc ctg atc tac aac aga atg ggc gcc gtg acc acc gag gtg gcc ttc	432
Gly Leu Ile Tyr Asn Arg Met Gly Ala Val Thr Thr Glu Val Ala Phe	
130 135 140	
ggc ctg gtc tgc gcc acc tgc gag cag atc gcc gac agc cag cac aga	480
Gly Leu Val Cys Ala Thr Cys Glu Gln Ile Ala Asp Ser Gln His Arg	
145 150 155 160	
tcc cac aga cag atg gtc acc acc acc aac ccc ctg atc aga cac gag	528
Ser His Arg Gln Met Val Thr Thr Thr Asn Pro Leu Ile Arg His Glu	
165 170 175	
aac aga atg gtg ctg gcc tct acc acc gcc aag gcc atg gaa cag atg	576
Asn Arg Met Val Leu Ala Ser Thr Thr Ala Lys Ala Met Glu Gln Met	
180 185 190	

Ser Ser Ala Gly Leu Lys Asn Asp Leu Leu Glu Asn Leu Gln Ala Tyr
 225 230 235 240
 Gln Lys Arg Met Gly Val Gln Met Gln Arg Phe Lys
 245 250

<210> 7
 <211> 5511
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Synthetic construct

<400> 7
 cgacaatatt ggctattggc cattgcatac gttgtatcta tatcataata tgtacattta 60
 tattggctca tgtccaatat gaccgccatg ttgacattga ttattgacta gttattaata 120
 gtaatcaatt acggggtcat tagttcatag cccatatatg gagttccgcg ttacataact 180
 tacggtaaat ggcccgcctc gtgaccgccc aacgaccccc gccattgac gtcaataatg 240
 acgtatgttc ccatagtaac gccaataggg actttccatt gacgtcaatg ggtggagtat 300
 ttacggtaaa ctgccactt ggcagtacat caagtgtatc atatgccaaag tccgccccta 360
 ttgacgtcaa tgacggtaaa tggcccgctt ggcattatgc ccagtacatg accttacggg 420
 actttcctac ttggcagtac atctacgtat tagtcatcgc tattaccatg gtgatgcggt 480
 tttggcagta caccaatggg cgtggatagc gttttgactc acgggggattt ccaagtctcc 540
 accccattga cgtcaatggg agtttgtttt ggcacaaaaa tcaacggggac tttccaaaat 600
 gtcglataaa ccccggcccg ttgacgcaaa tgggocgtag gcgtgtacgg tgggaggtct 660
 atataagcag agctcgttta gtgaaccgtc agatocctcg gagaagccat ccaagctgtt 720
 ttgacctcca tagaagacac cgggaccgat ccagcctccg cggccgggaa cggtgcatg 780
 gaacgoggat tcccctgccc aagagtgacg taagtaccgc ctatagactc tataggcaca 840
 cccctttggc tcttatgcat getatactgt ttttgcttg gggcctatac acccccgctc 900
 cttatgclat aggtgatggg atagcttagc ctatagggtg gggttattga ccattattga 960
 ccactcccct attggtgacg atactttcca ttaactatcc ataacatggc tctttgccac 1020
 aactatctct attggctata tgccaatact ctgtccttca gagactgaca cggactctgt 1080
 atttttacag gatggggccc catttattat ttacaaaatc acatatacaa caacgcgcgc 1140
 ccccggtccc gcagttttta ttaaacatag cgtgggatct ccacgcgaat ctcgggtagc 1200
 tgtlccggac atgggctctt ctccggtagc ggocgagctt ccacatccga gccctggctc 1260
 catgctcca gcggctcatg gtcgctcgge agctccttgc tctaacagt ggaggccaga 1320
 cttaggcaca gcacaatgcc caccaccacc agtgtgcgcg acaaggccgt ggcggtaggg 1380
 tatgtgtctg aaaatgagct cggagattgg gctcgcaccg tgacgcagat ggaagactta 1440
 aggcagcggc agaagaagat gcaggcagct gagttgtgtt attctgataa gagtcagagg 1500
 taactcccgt tgcggtgctg ttaacggtgg aggcaggtg agtctgagca gtactcgttg 1560
 ctgcccgcgc cgcaccacaga cataataget gacagactaa cagactgttc ctttccatgg 1620
 gtctttctg cagtcaccgt ccaagcttat ggaaaagatc gtgctgctgc tggctalctg 1680
 gagcctggg aagagcgacc agatttgcac cggctaccac gccacaaca gcaccgagca 1740
 ggtggacacc atcatggaaa agaacgtcac cgtgacccac gcccaggaca tctgggaaa 1800
 gaccacaac ggcaagctgt gcgacctgga cggcgtgaag ccctgatcc tgagggactg 1860
 cagcgtggcc ggctggctgc tgggcaacc catgtgcgac gagtccatca acgtgcccga 1920
 gtggagctac atcgtggaga aggcacaacc cggcaacgac ctgtgctacc ccggcaactt 1980
 caacgactac gaggaactga agcacctgct gtccaggatc aaccacttcg agaagatcca 2040
 gatcatccc aagagcagct ggtccgacca ctagggcaag agcggcgtga gcagcgcctg 2100
 cccataccag ggcagcccca gcttcttcag aaactgggtg tggtgatca agaagaaca 2160
 cacctacccc accatcaaga ggtctcaaa caacaccaac caggaagatc tgctggctct 2220
 gtggggcacc caccacccta atgacgcgcg cgaacagacc aggtgtacc agaaccacc 2280
 caccacatc agcgtgggca caagcaccct gaaccagagg ctggtgccca agatcgccac 2340
 caggtccaag gtgaacggac agtcccgag gatggaatc tcttgacca tctgaagcc 2400
 taacgacgcc atcaacttcg agagcaacgg caactttatc gccccgagt acgcctacaa 2460
 gatcgtgaag aagggcgaca gcgccatcat gaagagcgag ctggaatacg gcaactgcaa 2520

caccaagtgc	cagaccccca	tggcgccat	caacagcagc	atgcccttcc	acaacatcca	2580
ccccctgacc	atcggcgagt	gccccaaagta	cgtgaagagc	aacaggctgg	tgctggccac	2640
cggcctgagg	aacagccccc	agagagagag	cagaagaaa	aagaggggcc	tgctcggcgc	2700
tatcgcgggc	ttcatcgagg	gcggclggca	ggcctatggt	gacgggtggt	acggctacca	2760
ccactctaac	gagcagggca	gcggctacgc	cgccgacaaa	gagagcacc	agaaggccat	2820
cgacggcgtc	accaacaagg	tgaacagcat	catcgacaag	atgaacacc	agttcgaggc	2880
cgtgggcaga	gagttcaaca	acctgaaaag	gcggatcgag	aacctgaaca	agaaaatgga	2940
agatggcttc	ctggacgtgt	ggacctacaa	cgccgagctg	ctggtgctga	tygaaaacga	3000
gaggaccctg	gacttccacg	acagcaacgt	gaagaacctg	tacgacaaag	tgcggtgca	3060
gctgagggac	aacgccaag	agctgggcaa	cggtctcttc	gagttctacc	acaagtgcga	3120
caacgagtg	atgaaagcg	tgaggaacgg	cacctacgac	tacccccagt	acagcgagga	3180
agccaggctg	aagaggaag	agatcagcgg	agtgaagctg	gaaagcatcg	gcacctacca	3240
gatcctgagc	atctacagca	ccgtcgccag	cagcctggcc	ctggctatca	tggtggccgg	3300
actgagcctg	tgatgtgca	gcaacggcag	cctgacgtgc	aggatctgca	tcggatcctc	3360
gcaatcccta	gggtctgtgc	ttctagttgc	cagccaaact	gttgtttgcc	cctccccctg	3420
gccttccttg	acctggaag	gtgccactcc	actgtcctt	tctaataaaa	atgaggaaat	3480
tgcatcgcat	tgtctgagta	ggtgtcattc	tattctgggg	ggtgggggtg	ggcaggacag	3540
caaggggag	gattgggaag	acaatagcag	gcatgctggg	gatgctgggtg	gctctatggt	3600
cagaacgctc	ggttgccgcc	gggcgttttt	tatctagagt	cgacaaattc	agaagaactc	3660
gtcaagaagg	cgatagaagg	cgatgcgctg	cgaatcgga	gcggcgatac	cgtaaagcac	3720
gaggaacgg	tcagcccatt	cgccgccaa	ctcttcagca	atatacggg	tagccaacgc	3780
tatgtcctga	tagcggctcg	ccacaccag	ccggccacag	tcgatgaatc	cagaaaagcg	3840
gccattttcc	accatgatat	tcggcaagca	ggcatcgcca	tggytcaacg	cgagatcctc	3900
gccgtcgggc	atgctcgct	tgagcctggc	gaacagttcc	gctggcgcca	gccccctgat	3960
ctcttcgtec	agatcatcct	gatcgacaag	accggcttcc	atccgagtac	gtgctcgctc	4020
gatcgatgt	ttcgcttgg	ggtcgaatgg	gcaggtagcc	ggatcaagcg	tatgcagccg	4080
ccgcattgca	tcagccatga	tggatacttt	ctcgccagga	gcaagggtgag	atgacaggag	4140
atcctgcccc	ggcacttcgc	ccaatagcag	ccagtcctt	cccgttccag	tgacaacgctc	4200
gagcacagct	gogcaaggaa	cgcccgtcgt	ggccagccac	gatagccgcg	ctgctcgtc	4260
ttgcagttca	ttcagggcac	cggacaggtc	ggtcttgaca	aaaagaaccg	ggcgccccctg	4320
cgctgacagc	cggaacacgg	cggcatcaga	gcagccgatt	gtctgttggtg	cccagtcata	4380
gccgaatagc	ctctccacc	aagcgcccg	agaacctgcg	tgcaatccat	cttgttcaat	4440
catgcgaaac	gatectcacc	ctgtctcttg	atcagatctt	galccccctgc	gccatcagat	4500
ccttggcggc	gagaaagcca	tcagtttac	tttgacgggc	ttcccaacct	taccagaggg	4560
cgccccagct	ggcaattccg	gttcgcttgc	tgtccataaa	accgcccagt	ctagctatcg	4620
ccatgtaagc	ccactgcaag	ctacctgctt	tctctttgcg	cttgogtttt	cccttgtcca	4680
gatagcccag	tagctgacat	tcctccgggg	tcagcaacctg	ttctgoggac	tggttttcta	4740
cgtgaaaagg	atctaggtga	agatcctttt	tgataatctc	atgacaaaaa	tcccttaacg	4800
tgagttttcg	ttccactgag	cgtcagaccc	cgtagaaaag	atcaaaggat	cttcttgaga	4860
tccttttttt	ctgcgcgtaa	tctgctgctt	gcaaacaaaa	aaaccaccgc	taccagcggg	4920
ggtttgtttg	ccggatcaag	agctaccaac	tctttttccg	aaggtaactg	gcttcagcag	4980
agcgcagata	ccaaatactg	tccttctagt	gtagccgtag	ttaggccacc	acttcaagaa	5040
ctctgtagca	ccgctacat	acctcgctct	gctaactctg	ttaccagtgg	ctgctgccag	5100
tggcgataag	tcgtgtctta	ccgggttgg	ctcaagacga	tagttaccgg	ataaggcgca	5160
gcggtcgggc	tgaacggggg	gttcgtgcac	acagcccagc	ttggagcgaa	cgacctacac	5220
ogaactgaga	tacctacagc	gtgagctatg	agaaaagccc	acgcttcccc	aaggagagaaa	5280
ggcggacagg	tatccggtaa	gcggcagggg	cggaacagga	gagcgcacga	gggagcttcc	5340
agggggaaac	gcctggtatc	tttatagtc	tgtcgggttt	cgccacctct	gacttgagcg	5400
tcgatttttg	tgatgctcgt	cagggggggc	gagcctatgg	aaaaacgcca	gcaacggcgg	5460
ctttttacgg	ttcctgggct	tttctggtcc	ttttgctcac	atgttgctga	c	5511

<210> 8

<211> 5769

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic construct

<400> 8

tgcgcggttt	oggtgatgac	ggtgaaaacc	tctgacacat	gcagctcccg	gagacgggtca	60
cagcttgtct	gtaagcggat	gccgggagca	gacaagcccg	tcagggcgcg	tcagcgggtg	120
ttggcgggtg	tcggggctgg	cttaactatg	cggcatcaga	gcagattgta	ctgagagtgc	180
accatatgcg	gtgtgaaata	ccgcacagat	gcgtaaggag	aaaataccgc	atcagattgg	240
ctattggcca	ttgcatacgt	tqtatccata	tcataatatg	tacatttata	ttggctcatg	300
tccaacatta	ccgccaatgt	gacattgatt	attgactagt	tattaatagt	aatcaattac	360
ggggtcatta	gttcatagcc	catatatgga	gttccgcggt	acataactta	cggtaaattg	420
cccgcctggc	tgaccgcca	acgacccccg	cccattgacg	tcaataatga	cgtatgttcc	480
catagtaacg	ccaatagggg	ctttccattg	acgtcaatgg	gtggagtatt	tacggtaaac	540
tgcccacttg	gcagtacatc	aagtgtatca	tatgccaagt	acgccccta	ttgacgtcaa	600
tgacggtaaa	ttgcccgcct	ggcattatgc	ccagtacatg	accttatggg	actttcctac	660
ttggcagtac	atctacgtat	tagtcatcgc	tattaccatg	gtgatgcggt	tttggcagta	720
catcaatggg	cgtggatagc	gglttgactc	acggggattt	ccaagtctcc	accccattga	780
cgtcaatggg	agtttgtttt	ggcaccaaaa	tcaacgggac	tttccaaaat	gtcgtaacaa	840
ctccgcccc	ttgacgcaa	tgggcggtag	gcgtgtacgg	tgggaggtct	atataagcag	900
agctcgttta	gtgaaccgtc	agatgcctcg	gagacgcoat	ccacgctggt	ttgacctcca	960
tagaagacac	cgggaccgat	ccagcctcca	tccgctcgca	tctctccttc	acgcgcccgc	1020
cgcctacct	gaggccgcca	tccaagccgg	ttgagtccgg	ttctgcggcc	tcccgcctgt	1080
ggtgcctcct	gaactgcgtc	cgccgtctag	gtaagtttaa	agctcaggtc	gagaccgggc	1140
ctttgtccgg	cgctcccttg	gagctacct	agactcagcc	ggctctccac	gctttgctcg	1200
accctgcttg	ctcaactcta	gttaacggtg	gagggcagtg	tagctgagc	agtactcgtt	1260
gctgcccgcg	gcgccaccag	acataatagc	tgacagacta	acagactggt	cctttccatg	1320
ggtcttttct	gcagtaccgg	tcgtcgacac	gatccgatat	cgcgcgccacc	atgaatccta	1380
ataagaagat	catcacaatc	ggaagcatct	gcattggtgac	aggaatggtg	agcctgatgc	1440
tgcagatcgg	aaatctgatc	agcatctggg	tgagccacag	catccacaca	ggaatcagc	1500
acaaggccga	gcctatcagc	aatacaaatt	ttctgacaga	gaaggccgtg	gccagcgtga	1560
agctggccgg	aaatagcagc	ctgtgcctca	tcaatggatg	ggcctgttac	agcaaggata	1620
atagcatcag	aatcgggaagc	aagggagatg	tgtttgtgat	cagagagcct	tttatcagct	1680
gcagccacct	ggagtgcaga	acattttttc	tgacacaggg	agccctgctg	aatgataagc	1740
acagagatgg	aacagtgaa	gatagaagcc	ctcacagaac	actgatgagc	tgcctgtggg	1800
gagaggcccc	tagcccttac	aatagcagat	ttgagagcgt	ggcctggagc	gccagcgcct	1860
gccacgatgg	aacaagctgg	ctgacaatcg	gaatcagcgg	acctgataat	ggagccgtgg	1920
ccgtgctgaa	gtacaatgga	atcatcacag	atacaatcaa	gagctggaga	aataatatcc	1980
tgagaacaca	ggagagcgag	tcgcctcgcg	tgaatggaag	ctgctttaca	gtgatgacag	2040
atggacctag	caatggacag	gccagccaca	agatctttaa	gatggagaag	gaaaggtgg	2100
tgaagagcgt	ggagctggat	gcccctaatt	accactacga	ggagtgcagc	tgctaccctg	2160
atgcccggaga	gatcacatgc	gtgtgcagag	ataattggca	cggaaagcaat	agacctggg	2220
tgagctttaa	tcagaatctg	gagtaccaga	tcggatacat	ctgcagcgga	gtgtttggag	2280
ataatcctag	acctaattgat	ggaacaggaa	gctgcggacc	tgtgagcagc	aatggagcct	2340
acggagtgaa	gggatttagc	tttaagtacg	gaaatggagt	gtggatcgga	agaacaaaga	2400
gcacaaatag	cagaagcggg	tttgagatga	tctgggaccc	taatggatgg	acagagacag	2460
atagcagctt	ttagcgtgaag	caggatatcg	tggccatcac	agattggagc	ggatacagcg	2520
gaagctttgt	gcagcaccct	gagctgacag	gactggattg	catcagacct	tgcttttggg	2580
tggagctgat	cagaggaaga	cctaaggaga	gcacaatctg	gacaagcgga	agcagcatca	2640
gcttttgccg	agtgaatagc	gatacagtgg	gatggagctg	gcctgatgga	gccgagctgc	2700
cttttacaat	cgataagtga	gcgccgcctc	tagaccaggg	cctggatcca	gatctgctgt	2760
gccttctagt	tgccagccat	ctgttgtttg	ccctccccc	gtgccttctc	tgacctgga	2820
aggtgccact	cccactgtcc	tttctaata	aaatgaggaa	attgcatcgc	attgtctgag	2880
taggtgcat	tctattctgg	gggggtgggt	ggggcaggac	agcaaggggg	aggattggga	2940
agacaatagc	aggcatgctg	gggatcggt	gggtctatg	ggtaccaggg	tgctgaagaa	3000
ttgaccgggt	tctcctggg	ccagaaagaa	gcaggcacat	cccttctct	gtgacacacc	3060
ctgtccacgc	ccctggttct	tagttccagc	cccactcata	ggactctcat	agctcaggag	3120
ggctccgcct	tcaatcccac	ccgctaaagt	acttggagcg	gtctctcct	ccctcatcag	3180

```

cccaccaaac caaacctagc ctccaagagt ggaagaat taaagcaaga taggetatta 3240
agtgacagagg gagagaaaat gcctccaaca tgtgaggaag taatgagaga aatcatagaa 3300
ttttaaggcc atgatttaag gccatcatgg ccttaatctt ccgcttcctc gctcactgac 3360
tcgctgcgct cggctcgttc gctgcggcga gcggtatcag ctcaactcaa ggcggtaata 3420
cggttatcca cagaatcagg ggataacgca ggaagaaca tgtgagcaaa aggccagcaa 3480
aaggccagga accgtaaaaa ggcccgcttg cttggcgttt tccatagget ccgccccct 3540
gacgagcatc acaaaaatcg acgctcaagt cagagggtgc gaaaccgac aggactataa 3600
agataccagg cgtttccccc tggaaagctc ctctgctgct ctctgttcc gaccctgccg 3660
cttacoggat acctgtccgc ctttctccct togggaagcg tggcgctttc tcatagctca 3720
cgctgtaggt atctcagttc ggtgtaggtc gttcgtcca agctgggctg tgtgcaagaa 3780
cccccgctt agcccgaccg ctgcgcctta tccggtaact atcgtcttga gtccaaccog 3840
gtaagacacg acttatcgcc actggcagca gccactggtg acaggattag cagagcaggg 3900
tatgtaggcg gtgctacaga gttcttgaag tgggtggcta actacggcta cactagaaga 3960
acagtatttg gtatctgcgc tctgctgaag ccagttacct tcggaaaaag agttggtagc 4020
tcttgatccg gcaaaaaaac caccgctggt agcgggtggt tttttgttg caagcagcag 4080
attaacgcca gaaaaaagg atctcaagaa gatccttga tctttctac ggggtctgac 4140
gctcagtgga acgaaaaactc acgttaaggy attttggtea tgagattatc aaaaagatc 4200
ttcacctaga tccttttaaa ttaaaaatga agttttaaat caatctaaag tatatatgag 4260
taaaacttgg ctgacagtta ccaatgetta atcagtgagg cacctatctc agcagatctg 4320
ctatttcggt catccatagt tgctgactc gggggggggg ggcgctgagg tctgctcgt 4380
gaagaagggt ttgctgactc ataccaggcc tgaatcgccc catcatccag ccagaaagtg 4440
agggagccac ggttgatgag agctttgttg taggtggacc agttggtgat tttgaacttt 4500
tgctttgcca cggaacggtc tgctgtgctg ggaagatgcg tgatctgate cttcaactca 4560
gcaaaagttc gatttattca acaaagccgc cgtccgctca agtcagcgtg atgetctgcc 4620
agtgttaca ccaattaacc aattctgatt agaaaaactc atcgagcacc aaatgaaact 4680
gcaatattatt catatcagga ttatcaatc catattttg aaaaagccgt ttctgtaatg 4740
aaggagaaaa ctaccggagg cagttccata ggatggcaag atcctggtat cggctcgcga 4800
ttccgactcg tccaacatca atacaacct ttaatttccc ctctcaaaa ataagttat 4860
caagtgagaa atcaccatga gtgacgactg aatccggtga gaatggcaaa agcttatgca 4920
tttctttcca gacttgttca acaggccagc cattacgctc gtcatacaaa tcaactcgc 4980
caaccaaacc gttattcatt cgtgattgcg cctgagcgag acgaaatacg cgatecctgt 5040
taaaaggaca attacaaaca ggaatogaat gcaaccggcg caggaacact gccagcgc 5100
caacaatatt ttcacctgaa tcaggatatt cttctaatac ctggaatgct gttttcccg 5160
ggatgcgagt ggtgagtaac catgcatcat caggagtacg gataaaatgc ttgatggtcg 5220
gaagaggcat aaattccgtc agccagttta gtctgacct ctcactgta acatcattgg 5280
caacgctacc tttgccatgt ttcagaaaca actctggcgc atcgggcttc ccatacaatc 5340
gatagattgt cgcacctgat tgcccagatc tatecgagc ccatttatac ccatataaat 5400
cagcatccat gttggaattt aatcgcggcc tcgagcaaga cgtttcccg tgaatatggc 5460
tcataacacc ccttgattta ctggttatgt aagcagacag ttttatggt catgatgata 5520
tattttatc ttgtgcaatg taacatcaga gattttgaga cacaacgtgg ctttcccccc 5580
cccccaatta ttgaagcatt taccagggtt attgtctcat gagcggatac atatttgaat 5640
gtatttagaa aaataaaca ataggggttc cgcgcacatt tccccgaaa gtgccacctg 5700
acgtctaaga aaccattatt atcatgacat taacctataa aaataggcgt atcacaggc 5760
cctttcgtc

```

```

<210> 9
<211> 4598
<212> DNA
<213> Artificial Sequence

```

```

<220>
<223> Artificial construct

```

```

<400> 9
cgacaatatt ggetattggc cattgcatac gttgtatcta tatcataata tgtacattta 60
tattggctca tgtccaatat gaccgccatg ttgacattga ttattgacta gttattaata 120

```

gtaatcaatt	acgggggtcat	tagttcatag	ccatataatg	gagttccgcg	ttacataact	180
tacggtaaat	ggcccgccctc	gtgaccgccc	aacgaccccc	gcccattgac	gtcaataatg	240
acgtatgttc	ccatagtaac	gccaataggg	actttccatt	gacgtcaatg	ggtggagtat	300
ttacggtaaa	ctgcccactt	ggcagtacat	caagtgtatc	atatgccaaag	tccgccccta	360
ttgacgtcaa	tgacggtaaa	tgcccgcct	ggcattatgc	ccagtacatg	accttacggg	420
actttccctac	ttggcagtac	atctacgtat	tagtcatcgc	tattaccatg	gtgatgcggt	480
tttgccagta	caccaatggg	cgtggatagc	ggtttgactc	acggggattt	ccaagtctcc	540
accccattga	cgtcaatggg	agtttgtttt	ggcaccaaaa	tcaacgggac	tttccaaaat	600
gtcgtaaata	ccccgccccg	ttgacgcaaa	tggcggtag	gcgtgtacgg	tgggaggtct	660
atataagcag	agctcgttta	gtgaaccgtc	agatcgctg	gagacgccat	ccacgctgtt	720
ttgacctcca	tagaagacac	cgggaccgat	ccagcctccg	cggccgggaa	cgggtgcattg	780
gaacgcggat	tccccgtgcc	aagagtgcg	taagtaccgc	ctatagactc	tataggcaca	840
cccccttggc	tcttatgcat	gctatactgt	ttttggcttg	gggcctatac	acccccgctc	900
cttatgctat	aggtgatggt	atagcttagc	ctataggtgt	gggttattga	ccattattga	960
ccactccctc	attggtgacg	atactttcca	ttactaatcc	ataacatggc	tctttgccac	1020
aactatctct	attggctata	tgccaatact	ctgctctca	gagactgaca	cggactctgt	1080
atthttacag	gatggggtcc	catttattat	ttacaaatcc	acatatacaa	caacgcctgc	1140
ccccgtgccc	gcagttttta	ttaaacatag	cgtgggatct	ccacgcgaat	ctcgggtacg	1200
tgttccggac	atgggctctt	ctccggtagc	ggcgagctt	ccacatccga	gccctggctc	1260
catgctcca	gcggtcctatg	gtcgtcggc	agctccttgc	tccaaacagt	ggaggccaga	1320
cttaggcaca	gcacaatgcc	caccaccacc	agtgtgccgc	acaaggccgt	ggcggtaggg	1380
tatgtgtctg	aaaaatgagct	cggagattgg	gctcgcaccg	tgacgcagat	ggaagactta	1440
aggcagcggc	agaagaagat	gcaggcagct	gagttggtgt	attctgataa	gagtcagagg	1500
taactcccg	tgcggtgctg	ttaacgggtg	agggcaggtg	agtctgagca	gtactcgttg	1560
ctgcgcggc	cgcaccacga	cataatagct	gacagactaa	cagactgttc	ctttccatgg	1620
gtctttctg	cagtcaccgt	ccaagcttag	atctgccacc	atgagcctgc	tgaccgaggt	1680
ggagacatac	gtgctgtcca	tcctccccag	cgccctctg	aaggccgaga	tcgccagag	1740
actggaagat	gtgttcgccc	gcaagaacac	cgacctggaa	gtgctgatgg	aatggctgaa	1800
aaccagaccc	atcctgagcc	ctctgaccaa	ggcctcctg	ggcttcgtgt	tcacctgac	1860
cgtgccacc	gagagaggcc	tgacagggcg	gagattcgtg	cagaacgccc	tgaaacggcaa	1920
cggcgacccc	aacaacatgg	acaaggccgt	gaagctgtac	agaaagctga	agcgggagat	1980
caccttccac	ggcgccaaag	agatcagcct	gagctacagc	gctggcgccc	tgcccagctg	2040
catgggcctg	atclacaaca	gaatggggcg	cgtgaccacc	gaggtggcct	tcggcctggt	2100
ctgcgccacc	tgccgacaga	tgccgacag	ccagcacaga	tcccacagac	agatggtcac	2160
caccaccaac	cccctgatca	gacacgagaa	cagaatggtg	ctggcctcta	ccaccgcca	2220
ggccatggaa	cagatggccg	gcagcagcga	gcagccgcgc	gaggetatgg	aagtcgcctc	2280
tcaggctagg	cagatggctc	aggccatgag	aaccatcggc	acccacccca	gcagctctgc	2340
tggcctgaag	aacgacctgc	tggaaaacct	gcaggcctac	cagaaaagaa	tgggcgtcca	2400
gatgcagaga	ttcaagtgat	gagatatacc	tcagagaggg	gatcctcgca	atccctaggg	2460
ctgtgccttc	tagttgccag	ccaaactggt	gtttgccctt	cccccgctgc	ttccttgacc	2520
ctggaagggt	ccactccca	tgtcctttcc	taataaaatg	aggaaattgc	atcgcttgt	2580
ctgagtaggt	gtcaattctat	tctggggggg	gggtggggc	aggacagcaa	gggggaggt	2640
tgggaagaca	atagcaggca	tgctggggat	gcggtgggct	ctatgttcag	aacgctcgg	2700
tgccgcccgg	cgttttttat	ctagagtoga	caaattcaga	agaactcgtc	aagaaggcga	2760
tagaaggcga	tgctgtcga	atcgggagcg	gcgataccgt	aaagcacgag	gaagcggctc	2820
gcccatctgc	cgccaagctc	ttcagcaata	tcacgggtag	ccaacgctat	gtcctgatag	2880
cggctcccca	caccacagcg	gccacagctg	atgaatccag	aaaagcggcc	atthtccacc	2940
atgatattcg	gcaagcaggc	atcgccatgg	gtcaccgacg	gatcctcgcc	gtcgggcatg	3000
ctcgccctga	gcctggcgaa	cagttcggct	ggcgcgagcc	cctgatgctc	ttcgtccaga	3060
tcactctgat	cgacaagacc	ggcttccatc	cgagtacgtg	ctcgtctgat	gcgatgtttc	3120
gcttggtgtg	cgaatgggca	ggtagccgga	tcaagcgtat	gcagccgccc	cattgcatca	3180
gccatgatgg	atactttctc	ggcaggagca	aggtgagatg	acaggagatc	ctgcccggc	3240
acttcgcca	atagcagcca	gtcccttccc	gcttcagtga	caacgctcag	cacagctcgc	3300
caaggaaacg	ccgtctgtgc	cagccacgat	agccgcgctg	cctcgtcttg	cagttcattc	3360
agggcacccg	acaggtcgg	cttgacaaaa	agaaccgggc	gcccctgcgc	tgacagccgg	3420
aacacggcgg	catcagagca	gccgattgtc	tggtgtgccc	agtcatagcc	gaatagcctc	3480
tccaccaag	cggccggaga	acctgcgtgc	aatccatctt	gttcaatcat	gcgaaacgat	3540

```

cctcaccctg  tctcttgatc  agatcttgat  cccctgcgcc  atcagatcct  tggcggcgag  3600
aaagccatcc  agtttacttt  gcagggcttc  ccaaccttac  cagagggcgc  cccagctggc  3660
aattccgggt  cgcttgctgt  ccataaaacc  gccagctcta  gctatcgcca  tgtaagccca  3720
ctgcaagcta  cctgctttct  ctttgcgctt  gcttttccc  ttgtccagat  agcccagtag  3780
ctgacattca  tccggggcca  gcaccgttcc  tgcggactgg  cttctacgt  gaaaaggatc  3840
taggtgaaga  tcctttttga  taatctcatg  accaaaatcc  cttaacgtga  gttttcgttc  3900
cactgagcgt  cagaccccg  agaaaagatc  aaaggatctt  cttgagatcc  ttttttctg  3960
cgcgtaatct  getgcttgca  aacaaaaaaa  ccaccgctac  cagcgggtgt  ttgtttgccg  4020
gatcaagagc  taccaactct  ttttcogaag  gtaactggct  tcagcagagc  gcagatacca  4080
aatactgtcc  ttctagtgt  gccgtagtta  ggccaccact  tcaagaactc  ttagcaccgc  4140
cctacatacc  tcgctctgct  aatcctgtta  ccagtggctg  ctgccagtgg  cgataagtcg  4200
tgtcttaccg  ggttgactc  aagacgatag  ttaccggata  aggcgcagcg  gtcgggctga  4260
acgggggggt  cgtgcacaca  gccagcttg  gagcgaacga  cctacaccga  actgagatac  4320
ctacagcgtg  agctatgaga  aagcgcacag  ctcccgaag  ggagaaaggc  ggacaggtat  4380
ccggttaagc  gcagggctcg  aacaggagag  cgcacgaggg  agcttcagg  gggaaacgcc  4440
tggatcttt  atagtcctgt  cgggtttcgc  cacctctgac  ttgagcgtcg  atttttgtga  4500
tgctcgtcag  gggggcggag  cctatggaaa  aacgccagca  acgcgccctt  tttacggttc  4560
ctgggctttt  getggccttt  tgctcacatg  ttgtcgac  4598

```

```

<210> 10
<211> 9
<212> PRT
<213> Artificial Sequence

```

```

<220>
<223> Synthetic peptide

```

```

<400> 10
Ile Tyr Ser Thr Val Ala Ser Ser Leu
1           5

```

```

<210> 11
<211> 8
<212> PRT
<213> Artificial Sequence

```

```

<220>
<223> Synthetic peptide

```

```

<400> 11
Ser Ile Ile Asn Phe Glu Lys Leu
1           5

```

CLAIMS:

1. An isolated nucleic acid molecule comprising a nucleotide sequence encoding an influenza hemagglutinin (HA) polypeptide, wherein the nucleotide sequence encoding the HA polypeptide is at least 97% identical to SEQ ID NO: 1.
- 5 2. The isolated nucleic acid molecule of claim 1, wherein the nucleotide sequence encoding the HA polypeptide comprises SEQ ID NO: 1.
3. The isolated nucleic acid molecule of claim 1, wherein the nucleotide sequence encoding the HA polypeptide consists of SEQ ID NO: 1.
4. An influenza HA polypeptide encoded by the nucleic acid molecule of any one
10 of claims 1-3.
5. A vector comprising the isolated nucleic acid molecule of any one of claims 1-3.
6. The vector of claim 5, further comprising a promoter operably linked to the nucleotide sequence encoding the HA polypeptide.
- 15 7. The vector of claim 5 or claim 6, wherein the nucleotide sequence of the vector comprises SEQ ID NO: 7.
8. An influenza HA polypeptide produced by transfecting a host cell with the vector of any one of claims 5-7 under conditions sufficient to allow for expression of the polypeptide.
- 20 9. An isolated cell comprising the vector of any one of claims 5-7.
10. An isolated influenza HA polypeptide, wherein the amino acid sequence of the polypeptide is at least 99% identical to SEQ ID NO: 2.
11. The influenza HA polypeptide of claim 10, wherein the amino acid sequence of the polypeptide comprises SEQ ID NO: 2.

12. The influenza HA polypeptide of claim 10, wherein the amino acid sequence of the polypeptide consists of SEQ ID NO: 2.
13. A fusion protein comprising the polypeptide of any one of claims 10-12.
14. An influenza virus-like particle (VLP) comprising the polypeptide of any one
5 of claims 10-12.
15. The influenza VLP of claim 14, further comprising an influenza neuraminidase (NA) protein and an influenza matrix (M1) protein.
16. The influenza VLP of claim 15, wherein the amino acid sequence of the influenza NA protein is at least 95% identical to SEQ ID NO: 4.
- 10 17. The influenza VLP of claim 15, wherein the amino acid sequence of the influenza NA protein comprises SEQ ID NO: 4.
18. The influenza VLP of any one of claims 15-17, wherein the amino acid sequence of the influenza M1 protein is at least 95% identical to SEQ ID NO: 6.
19. The influenza VLP of any one of claims 15-17, wherein the amino acid
15 sequence of the influenza M1 protein comprises SEQ ID NO: 6.
20. An influenza VLP comprising the influenza HA polypeptide of any one of claims 10-12, produced by transfecting a host cell with a vector encoding the HA polypeptide, a vector encoding an influenza NA protein and a vector encoding an influenza M1 protein under conditions sufficient to allow for expression of the HA, M1 and NA proteins.
- 20 21. The influenza VLP of claim 20, wherein the nucleotide sequence of the vector encoding the HA polypeptide is at least 95% identical to SEQ ID NO: 7.
22. The influenza VLP of claim 20, wherein the nucleotide sequence of the vector encoding the HA polypeptide comprises SEQ ID NO: 7.

23. The influenza VLP of any one of claims 20-22, wherein the nucleotide sequence of the vector encoding the NA protein is at least 95% identical to SEQ ID NO: 8.
24. The influenza VLP of any one of claims 20-22, wherein the nucleotide sequence of the vector encoding the NA protein comprises SEQ ID NO: 8.
- 5 25. The influenza VLP of any one of claims 20-24, wherein the nucleotide sequence of the vector encoding the M1 protein is at least 95% identical to SEQ ID NO: 9.
26. The influenza VLP of any one of claims 20-24, wherein the nucleotide sequence of the vector encoding the M1 protein comprises SEQ ID NO: 9.
27. A collection of plasmids comprising:
- 10 (i) a plasmid encoding an influenza NA
- (ii) a plasmid encoding an influenza M1; and
- (iii) a plasmid encoding a codon-optimized influenza HA, wherein the nucleotide sequence encoding the codon-optimized influenza HA is at least 94% identical to SEQ ID NO: 1.
- 15 28. The collection of claim 27, wherein the influenza NA is codon-optimized.
29. The collection of claim 28, wherein the nucleotide sequence encoding the codon-optimized influenza NA is at least 95% identical to SEQ ID NO: 3.
30. The collection of any one of claims 27-29, wherein the influenza M1 is codon-optimized.
- 20 31. The collection of claim 30, wherein the nucleotide sequence encoding the codon-optimized influenza M1 is at least 95% identical to SEQ ID NO: 5.
32. The collection of any one of claims 27-31, wherein:
- (i) the plasmid encoding influenza NA comprises SEQ ID NO: 8;

- (ii) the plasmid encoding influenza M1 comprises SEQ ID NO: 9;
- (iii) the plasmid encoding influenza HA comprises SEQ ID NO: 7; or
- (iv) any combination of (i) to (iii).

33. A composition comprising the influenza HA protein of any one of
5 claims 10-12, the fusion protein of claim 13, or the VLP of any one of claims 14-26, and a
pharmaceutically acceptable carrier.
34. Use of a composition comprising the influenza HA protein of any one of
claims 10-12, the fusion protein of claim 13, or the VLP of any one of claims 14-26, for
eliciting an immune response to influenza virus in a subject.
- 10 35. The use of claim 34, wherein the composition further comprises an adjuvant.
36. Use of the composition of claim 33 for eliciting an immune response to
influenza virus in a subject.
37. Use of a composition comprising the VLP of any one of claims 14-26 and
pharmaceutically acceptable carrier for immunization of a subject against influenza virus.
- 15 38. The use of claim 36 or claim 37, wherein the composition further comprises an
adjuvant.
39. The use of any one of claims 35-38, wherein the composition is formulated for
intramuscular administration.
40. The use of any one of claims 35-39, wherein the composition comprises about
20 1 to about 25 µg of the VLP.
41. The use of claim 40, wherein the composition comprises about 15 µg of the
VLP.

42. A method of generating an optimized influenza virus polypeptide sequence, comprising:

(i) obtaining the amino acid sequences of the polypeptide from a group of influenza virus isolates, wherein the influenza virus isolates are from the same subtype;

5 (ii) organizing the amino acid sequences of the polypeptide from the group of influenza virus isolates by clade or sub-clade and then by geographical region within each clade or sub-clade;

(iii) aligning the amino acid sequences within each geographical region to generate primary consensus sequences, wherein each geographic region is represented by a
10 primary consensus sequence;


(iv) aligning the primary consensus sequences to generate secondary consensus sequences, wherein each clade or sub-clade is represented by a secondary consensus sequence; and

(v) producing the optimized influenza virus polypeptide from an optimized
15 influenza virus polypeptide sequence generated by aligning the secondary consensus sequences.

43. The method of claim 42, further comprising:

(i) reverse translating the optimized influenza virus polypeptide sequence to generate a coding sequence; and

20 (ii) optimizing the coding sequence for expression in mammalian cells.









FIG. 1B







FIG. 3F

HAI Titer
Low Dose





FIG. 3E

HAI Titer
High Dose









FIG. 5A




FIG. 5B




FIG. 6A




FIG. 6B




FIG. 7B





FIG. 7A










FIG. 8E

Nasal Wash Titers






FIG. 9

★ COBRA

FIG. 10A

Week 3




FIG. 10B

Week 6




FIG. 10C




FIG. 10D




FIG. 10E

Week 3




FIG. 10F

Week 6





FIG. 11
Receptor Blocking Antibody (HAI Titers)
 ■ COBRA VLP
 □ Whooper Swan VLP
 ▨ Mock (Alum Only)

*** $p < 0.001$
 ** $p < 0.01$
 * $p < 0.05$







FIG. 13A Weight Loss




FIG. 13B Sickness




FIG. 13C




FIG. 13D




FIG. 14A




FIG. 14B




FIG. 15A




FIG. 15B




FIG. 16A




FIG. 16B




FIG. 16C




FIG. 17A

**Antibody Secreting Cells
(Lung 6DPI)**




FIG. 17B

**IFN- γ Production
(Lung 6DPI)**




FIG. 18B

Sickness




FIG. 18A

Weight Loss

