

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
8 July 2010 (08.07.2010)

PCT

(10) International Publication Number  
**WO 2010/077775 A1**

(51) International Patent Classification:  
C12N 9/24 (2006.01) C12P 21/00 (2006.01)

(21) International Application Number:  
PCT/US2009/067641

(22) International Filing Date:  
11 December 2009 (11.12.2009)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
61/122,839 16 December 2008 (16.12.2008) US

(71) Applicant (for all designated States except US):  
NOVOZYMES, INC. [US/US]; 1445 Drew Avenue,  
Davis, California 95618 (US).

(72) Inventor; and

(75) Inventor/Applicant (for US only): REY, Michael  
[US/US]; 605 Robin Place, Davis, California 95616 (US).

(74) Agent: STARNES, Robert; Novozymes, Inc., 1445  
Drew Avenue, Davis, California 95618 (US).

(81) Designated States (unless otherwise indicated, for every  
kind of national protection available): AE, AG, AL, AM,  
AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ,  
CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO,

DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,  
HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP,  
KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD,  
ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI,  
NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD,  
SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT,  
TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every  
kind of regional protection available): ARIPO (BW, GH,  
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,  
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ,  
TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE,  
ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,  
MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, SM,  
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,  
ML, MR, NE, SN, TD, TG).

**Declarations under Rule 4.17:**

— as to applicant's entitlement to apply for and be granted  
a patent (Rule 4.17(ii))

**Published:**

— with international search report (Art. 21(3))

— with sequence listing part of description (Rule 5.2(a))



WO 2010/077775 A1

(54) Title: POLYPEPTIDES HAVING ALPHA-MANNOSIDASE ACTIVITY AND POLYNUCLEOTIDES ENCODING SAME

(57) Abstract: The present invention relates to isolated polypeptides having alpha-mannosidase activity and isolated polynucleotides encoding the polypeptides. The invention also relates to nucleic acid constructs, vectors, and host cells comprising the polynucleotides as well as methods of producing and using the polypeptides.

**POLYPEPTIDES HAVING ALPHA-MANNOSIDASE ACTIVITY  
AND POLYNUCLEOTIDES ENCODING SAME**

5

**Reference to a Sequence Listing**

This application contains a Sequence Listing filed electronically by EFS, which is incorporated herein by reference.

10

**Reference to a Deposit of Biological Material**

This application contains a reference to a deposit of biological material, which deposit is incorporated herein by reference.

**Background of the Invention**

15

**Field of the Invention**

The present invention relates to isolated polypeptides having alpha-mannosidase activity and isolated polynucleotides encoding the polypeptides. The invention also relates to nucleic acid constructs, vectors, and host cells comprising the polynucleotides as well as methods of producing and using the polypeptides.

20

**Description of the Related Art**

When produced in an eukaryotic host cell, most proteins developed for pharmaceutical applications have oligosaccharides attached to their polypeptide backbone. Sugar chains of such glycoproteins may be attached by N-glycosidic bonds to the amide group of asparagine residues or O-glycosidic bonds to the hydroxyl group of serine or threonine residues.

25

O-Mannosylation of recombinant human proteins/polypeptides produced in a non-human host, such as a fungal host, can result in a mannosylation pattern, which is different from that seen in humans. Due to the possible antigenic properties of such recombinant proteins, the mannose residues have to be completely removed in proteins developed for therapeutic applications. It is, therefore, desirable to completely remove O-linked mannose residues from proteins developed for pharmaceutical applications.

30

Letourneur *et al.*, 2001, *Biotechnol. Appl. Biochem.* 33: 35-45, describe the enzymatic deglycosylation of a surface antigen I from *Pichia pastoris* using jack-bean (alpha1-2/1-3/1-6)-mannosidase. Neustroev *et al.*, 1993, *FEBS Letters* 316: 157-160,

35

describes enzymatic deglycosilation of O-linked mannose by treatment of a glycoamylase from *Aspergillus awamori* with alpha-mannosidase. It is reported that the treatment results in removal of 24-26 % of total mannose. WO 2005/033325 discloses methods of producing a recombinant polypeptide free of alpha-linked mannose residues.

5           The present invention provides polypeptides having alpha-mannosidase activity and polynucleotides encoding the polypeptides.

### Summary of the Invention

10           The present invention relates to isolated polypeptides having alpha-mannosidase activity selected from the group consisting of:

(a) a polypeptide comprising an amino acid sequence having at least 60% identity to SEQ ID NO: 2;

15           (b) a polypeptide encoded by a polynucleotide that hybridizes under at least medium stringency conditions with (i) SEQ ID NO: 1, (ii) the genomic DNA sequence comprising SEQ ID NO: 1, or (iii) a full-length complementary strand of (i) or (ii);

(c) a polypeptide encoded by a polynucleotide comprising a nucleotide sequence having at least 60% identity to SEQ ID NO: 1; and

20           (d) a variant comprising a substitution, deletion, and/or insertion of one or more (several) amino acids of SEQ ID NO: 2.

The present invention also relates to isolated polynucleotides encoding polypeptides having alpha-mannosidase activity, selected from the group consisting of:

(a) a polynucleotide encoding a polypeptide comprising an amino acid sequence having at least 60% identity to SEQ ID NO: 2;

25           (b) a polynucleotide that hybridizes under at least medium stringency conditions with (i) SEQ ID NO: 1, (ii) the genomic DNA sequence comprising SEQ ID NO: 1, or (iii) a full-length complementary strand of (i) or (ii);

(c) a polynucleotide comprising a nucleotide sequence having at least 60% identity to SEQ ID NO: 1; and

30           (d) a polynucleotide encoding a variant comprising a substitution, deletion, and/or insertion of one or more (several) amino acids of SEQ ID NO: 2.

The present invention also relates to nucleic acid constructs, recombinant expression vectors, recombinant host cells comprising the polynucleotides, and methods of producing a polypeptide having alpha-mannosidase activity.

35           The present invention also relates to methods of inhibiting the expression of a polypeptide having alpha-mannosidase activity in a cell, comprising administering to the cell

or expressing in the cell a double-stranded RNA (dsRNA) molecule, wherein the dsRNA comprises a subsequence of a polynucleotide of the present invention. The present also relates to such a double-stranded inhibitory RNA (dsRNA) molecule, wherein optionally the dsRNA is a siRNA or a miRNA molecule.

5 The present invention also relates to plants comprising an isolated polynucleotide encoding a polypeptide having alpha-mannosidase activity.

The present invention also relates to methods of producing a polypeptide having alpha-mannosidase, comprising: (a) cultivating a transgenic plant or a plant cell comprising a polynucleotide encoding the polypeptide having alpha-mannosidase activity under  
10 conditions conducive for production of the polypeptide; and (b) recovering the polypeptide.

The present invention also relates to methods for producing a demannosylated protein, comprising: treating a protein comprising O-linked mannose residues with a polypeptide having mannosidase activity, wherein the O-linked mannose residues on the protein are removed.

15

### Brief Description of the Figures

Figure 1 shows the cDNA sequence and the deduced amino acid sequence of a  
20 *Thielavia terrestris* NRRL 8126 alpha-mannosidase gene (SEQ ID NOs: 1 and 2, respectively).

Figure 2 shows a map of pTter08D4.

### Definitions

25 **Alpha-mannosidase activity:** The term "alpha-mannosidase activity" is defined herein as an alpha-D-mannohydrolase activity (EC 3.2.1.24, EC 3.2.1.113, and EC 3.2.1.114) that catalyzes the hydrolysis of terminal alpha-D-mannose residues. For purposes of the present invention, alpha-mannosidase activity is determined according to the procedure described by Li, 1966, *J. Biol. Chem.* 241: 1010-1012. One unit of alpha-  
30 mannosidase activity equals the amount of enzyme capable of releasing 1  $\mu$ mole of alpha-D-mannose per minute under conditions of optimal pH and temperature.

The polypeptides of the present invention have at least 20%, preferably at least 40%, more preferably at least 50%, more preferably at least 60%, more preferably at least 70%, more preferably at least 80%, even more preferably at least 90%, most preferably at least  
35 95%, and even most preferably at least 100% of the alpha-mannosidase activity of SEQ ID NO: 2.

**Isolated polypeptide:** The term "isolated polypeptide" as used herein refers to a polypeptide that is isolated from a source. In a preferred aspect, the polypeptide is at least 1% pure, preferably at least 5% pure, more preferably at least 10% pure, more preferably at least 20% pure, more preferably at least 40% pure, more preferably at least 60% pure, even more preferably at least 80% pure, and most preferably at least 90% pure, as determined by SDS-PAGE.

**Substantially pure polypeptide:** The term "substantially pure polypeptide" denotes herein a polypeptide preparation that contains at most 10%, preferably at most 8%, more preferably at most 6%, more preferably at most 5%, more preferably at most 4%, more preferably at most 3%, even more preferably at most 2%, most preferably at most 1%, and even most preferably at most 0.5% by weight of other polypeptide material with which it is natively or recombinantly associated. It is, therefore, preferred that the substantially pure polypeptide is at least 92% pure, preferably at least 94% pure, more preferably at least 95% pure, more preferably at least 96% pure, more preferably at least 97% pure, more preferably at least 98% pure, even more preferably at least 99% pure, most preferably at least 99.5% pure, and even most preferably 100% pure by weight of the total polypeptide material present in the preparation. The polypeptides of the present invention are preferably in a substantially pure form, *i.e.*, that the polypeptide preparation is essentially free of other polypeptide material with which it is natively or recombinantly associated. This can be accomplished, for example, by preparing the polypeptide by well-known recombinant methods or by classical purification methods.

**Polypeptide coding sequence:** The term "polypeptide coding sequence" is defined herein as a nucleotide sequence that encodes a polypeptide having beta-glucosidase activity. In a preferred aspect, the polypeptide coding sequence is nucleotides 89 to 1363 of SEQ ID NO: 1. The SignalP software program (Nielsen *et al.*, 1997, *Protein Engineering* 10: 1-6) predicts that SEQ ID NO: 1 does not encode a signal peptide.

**Identity:** The relatedness between two amino acid sequences or between two nucleotide sequences is described by the parameter "identity".

For purposes of the present invention, the degree of identity between two amino acid sequences is determined using the Needleman-Wunsch algorithm (Needleman and Wunsch, 1970, *J. Mol. Biol.* 48: 443-453) as implemented in the Needle program of the EMBOSS package (EMBOSS: The European Molecular Biology Open Software Suite, Rice *et al.*, 2000, *Trends in Genetics* 16: 276-277), preferably version 3.0.0 or later. The optional parameters used are gap open penalty of 10, gap extension penalty of 0.5, and the EBLOSUM62 (EMBOSS version of BLOSUM62) substitution matrix. The output of Needle

labeled "longest identity" (obtained using the `-nobrief` option) is used as the percent identity and is calculated as follows:

$$(\text{Identical Residues} \times 100) / (\text{Length of Alignment} - \text{Total Number of Gaps in Alignment})$$

For purposes of the present invention, the degree of identity between two deoxyribonucleotide sequences is determined using the Needleman-Wunsch algorithm (Needleman and Wunsch, 1970, *supra*) as implemented in the Needle program of the EMBOSS package (EMBOSS: The European Molecular Biology Open Software Suite, Rice *et al.*, 2000, *supra*), preferably version 3.0.0 or later. The optional parameters used are gap open penalty of 10, gap extension penalty of 0.5, and the EDNAFULL (EMBOSS version of NCBI NUC4.4) substitution matrix. The output of Needle labeled "longest identity" (obtained using the `-nobrief` option) is used as the percent identity and is calculated as follows:

$$(\text{Identical Deoxyribonucleotides} \times 100) / (\text{Length of Alignment} - \text{Total Number of Gaps in Alignment})$$

**Homologous sequence:** The term "homologous sequence" is defined herein as a predicted protein having an E value (or expectancy score) of less than 0.001 in a tfasty search (Pearson, W.R., 1999, in *Bioinformatics Methods and Protocols*, S. Misener and S. A. Krawetz, ed., pp. 185-219) with the *Thielavia terrestris* alpha-mannosidase of SEQ ID NO: 2 or the polypeptide thereof.

**Polypeptide fragment:** The term "polypeptide fragment" is defined herein as a polypeptide having one or more (several) amino acids deleted from the amino and/or carboxyl terminus of the polypeptide of SEQ ID NO: 2; or a homologous sequence thereof; wherein the fragment has alpha-mannosidase activity. In a preferred aspect, a fragment contains at least 380 amino acid residues, more preferably at least 400 amino acid residues, and most preferably at least 420 amino acid residues of the polypeptide of SEQ ID NO: 2 or a homologous sequence thereof.

**Subsequence:** The term "subsequence" is defined herein as a nucleotide sequence having one or more (several) nucleotides deleted from the 5' and/or 3' end of SEQ ID NO: 1; or a homologous sequence thereof; wherein the subsequence encodes a polypeptide fragment having alpha-mannosidase activity. In a preferred aspect, a subsequence contains at least 1140 nucleotides, more preferably at least 1200 nucleotides, and most preferably at least 1260 nucleotides of SEQ ID NO: 1 or a homologous sequence thereof.

**Allelic variant:** The term "allelic variant" denotes herein any of two or more alternative forms of a gene occupying the same chromosomal locus. Allelic variation arises naturally through mutation, and may result in polymorphism within populations. Gene mutations can be silent (no change in the encoded polypeptide) or may encode polypeptides having altered amino acid sequences. An allelic variant of a polypeptide is a polypeptide

encoded by an allelic variant of a gene.

**Isolated polynucleotide:** The term "isolated polynucleotide" as used herein refers to a polynucleotide that is isolated from a source. In a preferred aspect, the polynucleotide is at least 1% pure, preferably at least 5% pure, more preferably at least 10% pure, more preferably at least 20% pure, more preferably at least 40% pure, more preferably at least 60% pure, even more preferably at least 80% pure, and most preferably at least 90% pure, as determined by agarose electrophoresis.

**Substantially pure polynucleotide:** The term "substantially pure polynucleotide" as used herein refers to a polynucleotide preparation free of other extraneous or unwanted nucleotides and in a form suitable for use within genetically engineered protein production systems. Thus, a substantially pure polynucleotide contains at most 10%, preferably at most 8%, more preferably at most 6%, more preferably at most 5%, more preferably at most 4%, more preferably at most 3%, even more preferably at most 2%, most preferably at most 1%, and even most preferably at most 0.5% by weight of other polynucleotide material with which it is natively or recombinantly associated. A substantially pure polynucleotide may, however, include naturally occurring 5' and 3' untranslated regions, such as promoters and terminators. It is preferred that the substantially pure polynucleotide is at least 90% pure, preferably at least 92% pure, more preferably at least 94% pure, more preferably at least 95% pure, more preferably at least 96% pure, more preferably at least 97% pure, even more preferably at least 98% pure, most preferably at least 99% pure, and even most preferably at least 99.5% pure by weight. The polynucleotides of the present invention are preferably in a substantially pure form, *i.e.*, that the polynucleotide preparation is essentially free of other polynucleotide material with which it is natively or recombinantly associated. The polynucleotides may be of genomic, cDNA, RNA, semisynthetic, synthetic origin, or any combinations thereof.

**Coding sequence:** When used herein the term "coding sequence" means a nucleotide sequence, which directly specifies the amino acid sequence of its protein product. The boundaries of the coding sequence are generally determined by an open reading frame, which usually begins with the ATG start codon or alternative start codons such as GTG and TTG and ends with a stop codon such as TAA, TAG, and TGA. The coding sequence may be a DNA, cDNA, synthetic, or recombinant nucleotide sequence.

**cDNA:** The term "cDNA" is defined herein as a DNA molecule that can be prepared by reverse transcription from a mature, spliced, mRNA molecule obtained from a eukaryotic cell. cDNA lacks intron sequences that may be present in the corresponding genomic DNA. The initial, primary RNA transcript is a precursor to mRNA that is processed through a series of steps before appearing as mature spliced mRNA. These steps include the

removal of intron sequences by a process called splicing. cDNA derived from mRNA lacks, therefore, any intron sequences.

**Nucleic acid construct:** The term "nucleic acid construct" as used herein refers to a nucleic acid molecule, either single- or double-stranded, which is isolated from a naturally occurring gene or which is modified to contain segments of nucleic acids in a manner that would not otherwise exist in nature or which is synthetic. The term nucleic acid construct is synonymous with the term "expression cassette" when the nucleic acid construct contains the control sequences required for expression of a coding sequence of the present invention.

**Control sequences:** The term "control sequences" is defined herein to include all components necessary for the expression of a polynucleotide encoding a polypeptide of the present invention. Each control sequence may be native or foreign to the nucleotide sequence encoding the polypeptide or native or foreign to each other. Such control sequences include, but are not limited to, a leader, polyadenylation sequence, propeptide sequence, promoter, signal peptide sequence, and transcription terminator. At a minimum, the control sequences include a promoter, and transcriptional and translational stop signals. The control sequences may be provided with linkers for the purpose of introducing specific restriction sites facilitating ligation of the control sequences with the coding region of the nucleotide sequence encoding a polypeptide.

**Operably linked:** The term "operably linked" denotes herein a configuration in which a control sequence is placed at an appropriate position relative to the coding sequence of a polynucleotide sequence such that the control sequence directs the expression of the coding sequence of a polypeptide.

**Expression:** The term "expression" includes any step involved in the production of a polypeptide including, but not limited to, transcription, post-transcriptional modification, translation, post-translational modification, and secretion.

**Expression vector:** The term "expression vector" is defined herein as a linear or circular DNA molecule that comprises a polynucleotide encoding a polypeptide of the present invention and is operably linked to additional nucleotides that provide for its expression.

**Host cell:** The term "host cell", as used herein, includes any cell type that is susceptible to transformation, transfection, transduction, and the like with a nucleic acid construct or expression vector comprising a polynucleotide of the present invention.

**Modification:** The term "modification" means herein any chemical modification of the polypeptide comprising or consisting of SEQ ID NO: 2; or a homologous sequence thereof; as well as genetic manipulation of the DNA encoding such a polypeptide. The



modification can be a substitution, a deletion, and/or an insertion of one or more (several) amino acids as well as replacements of one or more (several) amino acid side chains.

**Artificial variant:** When used herein, the term "artificial variant" means a polypeptide having alpha-mannosidase activity produced by an organism expressing a modified polynucleotide sequence of SEQ ID NO: 1; or a homologous sequence thereof. The modified nucleotide sequence is obtained through human intervention by modification of the polynucleotide sequence disclosed in SEQ ID NO: 1; or a homologous sequence thereof.

## Detailed Description of the Invention

### Polypeptides Having Alpha-Mannosidase Activity

In a first aspect, the present invention relates to isolated polypeptides comprising amino acid sequences having a degree of sequence identity to the mature polypeptide of SEQ ID NO: 2 of preferably at least 60%, more preferably at least 65%, more preferably at least 70%, more preferably at least 75%, more preferably at least 80%, more preferably at least 85%, even more preferably at least 90%, most preferably at least 95%, and even most preferably at least 96%, at least 97%, at least 98%, or at least 99%, which have alpha-mannosidase activity (hereinafter "homologous polypeptides"). In a preferred aspect, the homologous polypeptides comprise amino acid sequences that differ by ten amino acids, preferably by five amino acids, more preferably by four amino acids, even more preferably by three amino acids, most preferably by two amino acids, and even most preferably by one amino acid from the mature polypeptide of SEQ ID NO: 2.

A polypeptide of the present invention preferably comprises the amino acid sequence of SEQ ID NO: 2 or an allelic variant thereof; or a fragment thereof having alpha-mannosidase activity. In a preferred aspect, the polypeptide comprises the amino acid sequence of SEQ ID NO: 2. In another preferred aspect, the polypeptide comprises SEQ ID NO: 2. In another preferred aspect, the polypeptide consists of the amino acid sequence of SEQ ID NO: 2 or an allelic variant thereof; or a fragment thereof having alpha-mannosidase activity. In another preferred aspect, the polypeptide consists of the amino acid sequence of SEQ ID NO: 2. In another preferred aspect, the polypeptide consists of SEQ ID NO: 2.

In a second aspect, the present invention relates to isolated polypeptides having alpha-mannosidase activity that are encoded by polynucleotides that hybridize under preferably very low stringency conditions, more preferably low stringency conditions, more preferably medium stringency conditions, more preferably medium-high stringency conditions, even more preferably high stringency conditions, and most preferably very high

stringency conditions with (i) SEQ ID NO: 1, (ii) the genomic DNA sequence comprising the mature polypeptide coding sequence of SEQ ID NO: 1, or (iii) a full-length complementary strand of (i) or (ii) (J. Sambrook, E.F. Fritsch, and T. Maniatis, 1989, *Molecular Cloning, A Laboratory Manual*, 2d edition, Cold Spring Harbor, New York).

5           The nucleotide sequence of SEQ ID NO: 1; or a subsequence thereof; as well as the amino acid sequence of SEQ ID NO: 2; or a fragment thereof; may be used to design nucleic acid probes to identify and clone DNA encoding polypeptides having alpha-mannosidase activity from strains of different genera or species according to methods well known in the art. In particular, such probes can be used for hybridization with the genomic  
10 or cDNA of the genus or species of interest, following standard Southern blotting procedures, in order to identify and isolate the corresponding gene therein. Such probes can be considerably shorter than the entire sequence, but should be at least 14, preferably at least 25, more preferably at least 35, and most preferably at least 70 nucleotides in length. It is, however, preferred that the nucleic acid probe is at least 100 nucleotides in  
15 length. For example, the nucleic acid probe may be at least 200 nucleotides, preferably at least 300 nucleotides, more preferably at least 400 nucleotides, or most preferably at least 500 nucleotides in length. Even longer probes may be used, e.g., nucleic acid probes that are preferably at least 600 nucleotides, more preferably at least 700 nucleotides, even more preferably at least 800 nucleotides, or most preferably at least 900 nucleotides in length.  
20 Both DNA and RNA probes can be used. The probes are typically labeled for detecting the corresponding gene (for example, with <sup>32</sup>P, <sup>3</sup>H, <sup>35</sup>S, biotin, or avidin). Such probes are encompassed by the present invention.

A genomic DNA or cDNA library prepared from such other strains may, therefore, be screened for DNA that hybridizes with the probes described above and encodes a  
25 polypeptide having alpha-mannosidase activity. Genomic or other DNA from such other strains may be separated by agarose or polyacrylamide gel electrophoresis, or other separation techniques. DNA from the libraries or the separated DNA may be transferred to and immobilized on nitrocellulose or other suitable carrier material. In order to identify a clone or DNA that is homologous with SEQ ID NO: 1, or a subsequence thereof, the carrier  
30 material is preferably used in a Southern blot.

For purposes of the present invention, hybridization indicates that the nucleotide sequence hybridizes to a labeled nucleic acid probe corresponding to SEQ ID NO: 1; the genomic DNA sequence comprising SEQ ID NO: 1; its full-length complementary strand; or a subsequence thereof; under very low to very high stringency conditions. Molecules to  
35 which the nucleic acid probe hybridizes under these conditions can be detected using, for example, X-ray film.

In a preferred aspect, the nucleic acid probe is SEQ ID NO: 1. In another preferred aspect, the nucleic acid probe is a polynucleotide sequence that encodes the polypeptide of SEQ ID NO: 2, or a subsequence thereof. In another preferred aspect, the nucleic acid probe is the polynucleotide sequence contained in plasmid pTter08D4 which is contained in *E. coli* NRRL B-50205, wherein the polynucleotide sequence thereof encodes a polypeptide having alpha-mannosidase activity.

For long probes of at least 100 nucleotides in length, very low to very high stringency conditions are defined as prehybridization and hybridization at 42°C in 5X SSPE, 0.3% SDS, 200 µg/ml sheared and denatured salmon sperm DNA, and either 25% formamide for very low and low stringencies, 35% formamide for medium and medium-high stringencies, or 50% formamide for high and very high stringencies, following standard Southern blotting procedures for 12 to 24 hours optimally.

For long probes of at least 100 nucleotides in length, the carrier material is finally washed three times each for 15 minutes using 2X SSC, 0.2% SDS preferably at 45°C (very low stringency), more preferably at 50°C (low stringency), more preferably at 55°C (medium stringency), more preferably at 60°C (medium-high stringency), even more preferably at 65°C (high stringency), and most preferably at 70°C (very high stringency).

For short probes of about 15 nucleotides to about 70 nucleotides in length, stringency conditions are defined as prehybridization, hybridization, and washing post-hybridization at about 5°C to about 10°C below the calculated  $T_m$  using the calculation according to Bolton and McCarthy (1962, *Proceedings of the National Academy of Sciences USA* 48:1390) in 0.9 M NaCl, 0.09 M Tris-HCl pH 7.6, 6 mM EDTA, 0.5% NP-40, 1X Denhardt's solution, 1 mM sodium pyrophosphate, 1 mM sodium monobasic phosphate, 0.1 mM ATP, and 0.2 mg of yeast RNA per ml following standard Southern blotting procedures for 12 to 24 hours optimally.

For short probes of about 15 nucleotides to about 70 nucleotides in length, the carrier material is washed once in 6X SSC plus 0.1% SDS for 15 minutes and twice each for 15 minutes using 6X SSC at 5°C to 10°C below the calculated  $T_m$ .

In a third aspect, the present invention relates to isolated polypeptides having alpha-mannosidase activity encoded by polynucleotides comprising or consisting of nucleotide sequences having a degree of sequence identity to the mature polypeptide coding sequence of SEQ ID NO: 1 of preferably at least 60%, more preferably at least 65%, more preferably at least 70%, more preferably at least 75%, more preferably at least 80%, more preferably at least 85%, even more preferably at least 90%, most preferably at least 95%, and even most preferably at least 96%, at least 97%, at least 98%, or at least 99%, which encode a polypeptide having alpha-mannosidase activity. See polynucleotide section herein.

In a fourth aspect, the present invention relates to artificial variants comprising a substitution, deletion, and/or insertion of one or more (or several) amino acids of SEQ ID NO: 2, or a homologous sequence thereof. Preferably, amino acid changes are of a minor nature, that is conservative amino acid substitutions or insertions that do not significantly affect the folding and/or activity of the protein; small deletions, typically of one to about 30 amino acids; small amino- or carboxyl-terminal extensions, such as an amino-terminal methionine residue; a small linker peptide of up to about 20-25 residues; or a small extension that facilitates purification by changing net charge or another function, such as a poly-histidine tract, an antigenic epitope or a binding domain.

Examples of conservative substitutions are within the group of basic amino acids (arginine, lysine and histidine), acidic amino acids (glutamic acid and aspartic acid), polar amino acids (glutamine and asparagine), hydrophobic amino acids (leucine, isoleucine and valine), aromatic amino acids (phenylalanine, tryptophan and tyrosine), and small amino acids (glycine, alanine, serine, threonine and methionine). Amino acid substitutions that do not generally alter specific activity are known in the art and are described, for example, by H. Neurath and R.L. Hill, 1979, *In, The Proteins*, Academic Press, New York. The most commonly occurring exchanges are Ala/Ser, Val/Ile, Asp/Glu, Thr/Ser, Ala/Gly, Ala/Thr, Ser/Asn, Ala/Val, Ser/Gly, Tyr/Phe, Ala/Pro, Lys/Arg, Asp/Asn, Leu/Ile, Leu/Val, Ala/Glu, and Asp/Gly.

In addition to the 20 standard amino acids, non-standard amino acids (such as 4-hydroxyproline, 6-*N*-methyl lysine, 2-aminoisobutyric acid, isovaline, and alpha-methyl serine) may be substituted for amino acid residues of a wild-type polypeptide. A limited number of non-conservative amino acids, amino acids that are not encoded by the genetic code, and unnatural amino acids may be substituted for amino acid residues. "Unnatural amino acids" have been modified after protein synthesis, and/or have a chemical structure in their side chain(s) different from that of the standard amino acids. Unnatural amino acids can be chemically synthesized, and preferably, are commercially available, and include pipercolic acid, thiazolidine carboxylic acid, dehydroproline, 3- and 4-methylproline, and 3,3-dimethylproline.

Alternatively, the amino acid changes are of such a nature that the physico-chemical properties of the polypeptides are altered. For example, amino acid changes may improve the thermal stability of the polypeptide, alter the substrate specificity, change the pH optimum, and the like.

Essential amino acids in the parent polypeptide can be identified according to procedures known in the art, such as site-directed mutagenesis or alanine-scanning mutagenesis (Cunningham and Wells, 1989, *Science* 244: 1081-1085). In the latter

technique, single alanine mutations are introduced at every residue in the molecule, and the resultant mutant molecules are tested for biological activity (*i.e.*, alpha-mannosidase activity) to identify amino acid residues that are critical to the activity of the molecule. See also, Hilton *et al.*, 1996, *J. Biol. Chem.* 271: 4699-4708. The active site of the enzyme or other biological interaction can also be determined by physical analysis of structure, as determined by such techniques as nuclear magnetic resonance, crystallography, electron diffraction, or photoaffinity labeling, in conjunction with mutation of putative contact site amino acids. See, for example, de Vos *et al.*, 1992, *Science* 255: 306-312; Smith *et al.*, 1992, *J. Mol. Biol.* 224: 899-904; Wlodaver *et al.*, 1992, *FEBS Lett.* 309: 59-64. The identities of essential amino acids can also be inferred from analysis of identities with polypeptides that are related to a polypeptide according to the invention.

Single or multiple amino acid substitutions, deletions, and/or insertions can be made and tested using known methods of mutagenesis, recombination, and/or shuffling, followed by a relevant screening procedure, such as those disclosed by Reidhaar-Olson and Sauer, 1988, *Science* 241: 53-57; Bowie and Sauer, 1989, *Proc. Natl. Acad. Sci. USA* 86: 2152-2156; WO 95/17413; or WO 95/22625. Other methods that can be used include error-prone PCR, phage display (*e.g.*, Lowman *et al.*, 1991, *Biochem.* 30: 10832-10837; U.S. Patent No. 5,223,409; WO 92/06204), and region-directed mutagenesis (Derbyshire *et al.*, 1986, *Gene* 46: 145; Ner *et al.*, 1988, *DNA* 7: 127).

Mutagenesis/shuffling methods can be combined with high-throughput, automated screening methods to detect activity of cloned, mutagenized polypeptides expressed by host cells (Ness *et al.*, 1999, *Nature Biotechnology* 17: 893-896). Mutagenized DNA molecules that encode active polypeptides can be recovered from the host cells and rapidly sequenced using standard methods in the art. These methods allow the rapid determination of the importance of individual amino acid residues in a polypeptide of interest, and can be applied to polypeptides of unknown structure.

The total number of amino acid substitutions, deletions and/or insertions of SEQ ID NO: 2 is 10, preferably 9, more preferably 8, more preferably 7, more preferably at most 6, more preferably 5, more preferably 4, even more preferably 3, most preferably 2, and even most preferably 1.

### **Sources of Polypeptides Having Alpha-Mannosidase Activity**

A polypeptide having alpha-mannosidase activity of the present invention may be obtained from microorganisms of any genus. For purposes of the present invention, the term "obtained from" as used herein in connection with a given source shall mean that the polypeptide encoded by a nucleotide sequence is produced by the source or by a strain in

which the nucleotide sequence from the source has been inserted. In a preferred aspect, the polypeptide obtained from a given source is secreted extracellularly.

A polypeptide having alpha-mannosidase activity of the present invention may be a bacterial polypeptide. For example, the polypeptide may be a gram positive bacterial polypeptide such as a *Bacillus*, *Streptococcus*, *Streptomyces*, *Staphylococcus*, *Enterococcus*, *Lactobacillus*, *Lactococcus*, *Clostridium*, *Geobacillus*, or *Oceanobacillus* polypeptide having alpha-mannosidase activity, or a Gram negative bacterial polypeptide such as an *E. coli*, *Pseudomonas*, *Salmonella*, *Campylobacter*, *Helicobacter*, *Flavobacterium*, *Fusobacterium*, *Ilyobacter*, *Neisseria*, or *Ureaplasma* polypeptide having alpha-mannosidase activity.

In a preferred aspect, the polypeptide is a *Bacillus alkalophilus*, *Bacillus amyloliquefaciens*, *Bacillus brevis*, *Bacillus circulans*, *Bacillus clausii*, *Bacillus coagulans*, *Bacillus firmus*, *Bacillus lautus*, *Bacillus lentus*, *Bacillus licheniformis*, *Bacillus megaterium*, *Bacillus pumilus*, *Bacillus stearothermophilus*, *Bacillus subtilis*, or *Bacillus thuringiensis* polypeptide having alpha-mannosidase activity.

In another preferred aspect, the polypeptide is a *Streptococcus equisimilis*, *Streptococcus pyogenes*, *Streptococcus uberis*, or *Streptococcus equi* subsp. *Zooepidemicus* polypeptide having alpha-mannosidase activity.

In another preferred aspect, the polypeptide is a *Streptomyces achromogenes*, *Streptomyces avermitilis*, *Streptomyces coelicolor*, *Streptomyces griseus*, or *Streptomyces lividans* polypeptide having alpha-mannosidase activity.

A polypeptide having alpha-mannosidase activity of the present invention may also be a fungal polypeptide, and more preferably a yeast polypeptide such as a *Candida*, *Kluyveromyces*, *Pichia*, *Saccharomyces*, *Schizosaccharomyces*, or *Yarrowia* polypeptide having alpha-mannosidase activity; or more preferably a filamentous fungal polypeptide such as an *Acremonium*, *Agaricus*, *Alternaria*, *Aspergillus*, *Aureobasidium*, *Botryosphaeria*, *Ceriporiopsis*, *Chaetomidium*, *Chrysosporium*, *Claviceps*, *Cochliobolus*, *Coprinopsis*, *Coptotermes*, *Corynascus*, *Cryphonectria*, *Cryptococcus*, *Diplodia*, *Exidia*, *Filibasidium*, *Fusarium*, *Gibberella*, *Holomastigotoides*, *Humicola*, *Irpex*, *Lentinula*, *Leptosphaeria*, *Magnaporthe*, *Melanocarpus*, *Meripilus*, *Mucor*, *Myceliophthora*, *Neocallimastix*, *Neurospora*, *Paecilomyces*, *Penicillium*, *Phanerochaete*, *Piromyces*, *Poitrasia*, *Pseudoplectania*, *Pseudotriconympha*, *Rhizomucor*, *Schizophyllum*, *Scytalidium*, *Talaromyces*, *Thermoascus*, *Thielavia*, *Tolypocladium*, *Trichoderma*, *Trichophaea*, *Verticillium*, *Volvariella*, or *Xylaria* polypeptide having alpha-mannosidase activity.

In a preferred aspect, the polypeptide is a *Saccharomyces carlsbergensis*, *Saccharomyces cerevisiae*, *Saccharomyces diastaticus*, *Saccharomyces douglasii*,

*Saccharomyces kluyveri*, *Saccharomyces norbensis*, or *Saccharomyces oviformis* polypeptide having alpha-mannosidase activity.

In another preferred aspect, the polypeptide is an *Acremonium cellulolyticus*, *Aspergillus aculeatus*, *Aspergillus awamori*, *Aspergillus fumigatus*, *Aspergillus foetidus*,  
5 *Aspergillus japonicus*, *Aspergillus nidulans*, *Aspergillus niger*, *Aspergillus oryzae*,  
*Chrysosporium keratinophilum*, *Chrysosporium lucknowense*, *Chrysosporium tropicum*,  
*Chrysosporium merdarium*, *Chrysosporium inops*, *Chrysosporium pannicola*, *Chrysosporium queenslandicum*, *Chrysosporium zonatum*, *Fusarium bactridioides*, *Fusarium cerealis*,  
10 *Fusarium crookwellense*, *Fusarium culmorum*, *Fusarium graminearum*, *Fusarium graminum*,  
*Fusarium heterosporum*, *Fusarium negundi*, *Fusarium oxysporum*, *Fusarium reticulatum*,  
*Fusarium roseum*, *Fusarium sambucinum*, *Fusarium sarcochroum*, *Fusarium sporotrichioides*, *Fusarium sulphureum*, *Fusarium torulosum*, *Fusarium trichothecioides*,  
*Fusarium venenatum*, *Humicola grisea*, *Humicola insolens*, *Humicola lanuginosa*, *Irpex lacteus*, *Mucor miehei*, *Myceliophthora thermophila*, *Neurospora crassa*, *Penicillium*  
15 *funiculosum*, *Penicillium purpurogenum*, *Phanerochaete chrysosporium*, *Trichoderma harzianum*, *Trichoderma koningii*, *Trichoderma longibrachiatum*, *Trichoderma reesei*, or  
*Trichoderma viride* polypeptide having alpha-mannosidase activity.

In another preferred aspect, the polypeptide is a *Thielavia achromatica*, *Thielavia albomyces*, *Thielavia albopilosa*, *Thielavia australeinsis*, *Thielavia fimeti*, *Thielavia*  
20 *microspora*, *Thielavia ovispora*, *Thielavia peruviana*, *Thielavia spededonium*, *Thielavia setosa*, *Thielavia subthermophila*, or *Thielavia terrestris* polypeptide having alpha-mannosidase activity.

In a more preferred aspect, the polypeptide is a *Thielavia terrestris* polypeptide having alpha-mannosidase activity. In a most preferred aspect, the polypeptide is a  
25 *Thielavia terrestris* NRRL 8126 polypeptide having alpha-mannosidase activity, e.g., the polypeptide comprising SEQ ID NO: 2.

It will be understood that for the aforementioned species the invention encompasses both the perfect and imperfect states, and other taxonomic equivalents, e.g., anamorphs, regardless of the species name by which they are known. Those skilled in the art will readily  
30 recognize the identity of appropriate equivalents.

Strains of these species are readily accessible to the public in a number of culture collections, such as the American Type Culture Collection (ATCC), Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH (DSM), Centraalbureau Voor Schimmelcultures (CBS), and Agricultural Research Service Patent Culture Collection, Northern Regional  
35 Research Center (NRRL).

Furthermore, such polypeptides may be identified and obtained from other sources

including microorganisms isolated from nature (e.g., soil, composts, water, etc.) using the above-mentioned probes. Techniques for isolating microorganisms from natural habitats are well known in the art. The polynucleotide may then be obtained by similarly screening a genomic or cDNA library of such a microorganism. Once a polynucleotide encoding a polypeptide has been detected with the probe(s), the polynucleotide can be isolated or cloned by utilizing techniques that are well known to those of ordinary skill in the art (see, e.g., Sambrook *et al.*, 1989, *supra*).

Polypeptides of the present invention also include fused polypeptides or cleavable fusion polypeptides in which another polypeptide is fused at the N-terminus or the C-terminus of the polypeptide or fragment thereof. A fused polypeptide is produced by fusing a nucleotide sequence (or a portion thereof) encoding another polypeptide to a nucleotide sequence (or a portion thereof) of the present invention. Techniques for producing fusion polypeptides are known in the art, and include ligating the coding sequences encoding the polypeptides so that they are in frame and that expression of the fused polypeptide is under control of the same promoter(s) and terminator.

A fusion polypeptide can further comprise a cleavage site. Upon secretion of the fusion protein, the site is cleaved releasing the polypeptide having alpha-mannosidase activity from the fusion protein. Examples of cleavage sites include, but are not limited to, a Kex2 site that encodes the dipeptide Lys-Arg (Martin *et al.*, 2003, *J. Ind. Microbiol. Biotechnol.* 3: 568-76; Svetina *et al.*, 2000, *J. Biotechnol.* 76: 245-251; Rasmussen-Wilson *et al.*, 1997, *Appl. Environ. Microbiol.* 63: 3488-3493; Ward *et al.*, 1995, *Biotechnology* 13: 498-503; and Contreras *et al.*, 1991, *Biotechnology* 9: 378-381), an Ile-(Glu or Asp)-Gly-Arg site, which is cleaved by a Factor Xa protease after the arginine residue (Eaton *et al.*, 1986, *Biochem.* 25: 505-512); a Asp-Asp-Asp-Asp-Lys site, which is cleaved by an enterokinase after the lysine (Collins-Racie *et al.*, 1995, *Biotechnology* 13: 982-987); a His-Tyr-Glu site or His-Tyr-Asp site, which is cleaved by Genenase I (Carter *et al.*, 1989, *Proteins: Structure, Function, and Genetics* 6: 240-248); a Leu-Val-Pro-Arg-Gly-Ser site, which is cleaved by thrombin after the Arg (Stevens, 2003, *Drug Discovery World* 4: 35-48); a Glu-Asn-Leu-Tyr-Phe-Gln-Gly site, which is cleaved by TEV protease after the Gln (Stevens, 2003, *supra*); and a Leu-Glu-Val-Leu-Phe-Gln-Gly-Pro site, which is cleaved by a genetically engineered form of human rhinovirus 3C protease after the Gln (Stevens, 2003, *supra*).

### Polynucleotides

The present invention also relates to isolated polynucleotides comprising or consisting of nucleotide sequences that encode polypeptides having alpha-mannosidase activity of the present invention.



In a preferred aspect, the nucleotide sequence comprises or consists of SEQ ID NO: 1. In another preferred aspect, the nucleotide sequence comprises or consists of the sequence contained in plasmid pTter08D4 which is contained in *E. coli* NRRL B-50205. The present invention also encompasses nucleotide sequences that encode polypeptides comprising or consisting of SEQ ID NO: 2, which differ from SEQ ID NO: 1 by virtue of the degeneracy of the genetic code. The present invention also relates to subsequences of SEQ ID NO: 1 that encode fragments of SEQ ID NO: 2 that have alpha-mannosidase activity.

The present invention also relates to mutant polynucleotides comprising or consisting of at least one mutation in SEQ ID NO: 1, in which the mutant nucleotide sequence encodes SEQ ID NO: 2.

The techniques used to isolate or clone a polynucleotide encoding a polypeptide are known in the art and include isolation from genomic DNA, preparation from cDNA, or a combination thereof. The cloning of the polynucleotides of the present invention from such genomic DNA can be effected, e.g., by using the well known polymerase chain reaction (PCR) or antibody screening of expression libraries to detect cloned DNA fragments with shared structural features. See, e.g., Innis *et al.*, 1990, *PCR: A Guide to Methods and Application*, Academic Press, New York. Other nucleic acid amplification procedures such as ligase chain reaction (LCR), ligated activated transcription (LAT) and nucleotide sequence-based amplification (NASBA) may be used. The polynucleotides may be cloned from a strain of *Thielavia*, or another or related organism and thus, for example, may be an allelic or species variant of the polypeptide encoding region of the nucleotide sequence.

The present invention also relates to isolated polynucleotides comprising or consisting of nucleotide sequences that have a degree of identity to SEQ ID NO: 1 of preferably at least 60%, more preferably at least 65%, more preferably at least 70%, more preferably at least 75%, more preferably at least 80%, more preferably at least 85%, even more preferably at least 90%, most preferably at least 95%, and even most preferably at least 96%, at least 97%, at least 98%, or at least 99%, which encode an active polypeptide.

Modification of a nucleotide sequence encoding a polypeptide of the present invention may be necessary for the synthesis of polypeptides substantially similar to the polypeptide. The term "substantially similar" to the polypeptide refers to non-naturally occurring forms of the polypeptide. These polypeptides may differ in some engineered way from the polypeptide isolated from its native source, e.g., artificial variants that differ in specific activity, thermostability, pH optimum, or the like. The variant sequence may be constructed on the basis of the nucleotide sequence presented as SEQ ID NO: 1, e.g., a subsequence thereof, and/or by introduction of nucleotide substitutions that do not give rise

to another amino acid sequence of the polypeptide encoded by the nucleotide sequence, but which correspond to the codon usage of the host organism intended for production of the enzyme, or by introduction of nucleotide substitutions that may give rise to a different amino acid sequence. For a general description of nucleotide substitution, see, e.g., Ford *et al.*, 1991, *Protein Expression and Purification 2*: 95-107.

It will be apparent to those skilled in the art that such substitutions can be made outside the regions critical to the function of the molecule and still result in an active polypeptide. Amino acid residues essential to the activity of the polypeptide encoded by an isolated polynucleotide of the invention, and therefore preferably not subject to substitution, may be identified according to procedures known in the art, such as site-directed mutagenesis or alanine-scanning mutagenesis (see, e.g., Cunningham and Wells, 1989, *supra*). In the latter technique, mutations are introduced at every positively charged residue in the molecule, and the resultant mutant molecules are tested for alpha-mannosidase activity to identify amino acid residues that are critical to the activity of the molecule. Sites of substrate-enzyme interaction can also be determined by analysis of the three-dimensional structure as determined by such techniques as nuclear magnetic resonance analysis, crystallography or photoaffinity labeling (see, e.g., de Vos *et al.*, 1992, *supra*; Smith *et al.*, 1992, *supra*; Wlodaver *et al.*, 1992, *supra*).

The present invention also relates to isolated polynucleotides encoding polypeptides of the present invention, which hybridize under preferably very low stringency conditions, more preferably low stringency conditions, more preferably medium stringency conditions, more preferably medium-high stringency conditions, even more preferably high stringency conditions, and most preferably very high stringency conditions with (i) the mature polypeptide coding sequence of SEQ ID NO: 1, (ii) the genomic DNA sequence comprising the mature polypeptide coding sequence of SEQ ID NO: 1, or (iii) a full-length complementary strand of (i) or (ii); or allelic variants and subsequences thereof (Sambrook *et al.*, 1989, *supra*), as defined herein.

The present invention also relates to isolated polynucleotides obtained by (a) hybridizing a population of DNA under preferably very low stringency conditions, more preferably low stringency conditions, more preferably medium stringency conditions, more preferably medium-high stringency conditions, even more preferably high stringency conditions, and most preferably very high stringency conditions with (i) the mature polypeptide coding sequence of SEQ ID NO: 1, (ii) the genomic DNA sequence comprising the mature polypeptide coding sequence of SEQ ID NO: 1, or (iii) a full-length complementary strand of (i) or (ii); and (b) isolating the hybridizing polynucleotide, which encodes a polypeptide having alpha-mannosidase activity.

### Nucleic Acid Constructs

The present invention also relates to nucleic acid constructs comprising an isolated polynucleotide of the present invention operably linked to one or more (several) control sequences that direct the expression of the coding sequence in a suitable host cell under conditions compatible with the control sequences.

An isolated polynucleotide encoding a polypeptide of the present invention may be manipulated in a variety of ways to provide for expression of the polypeptide. Manipulation of the polynucleotide's sequence prior to its insertion into a vector may be desirable or necessary depending on the expression vector. The techniques for modifying polynucleotide sequences utilizing recombinant DNA methods are well known in the art.

The control sequence may be an appropriate promoter sequence, a nucleotide sequence that is recognized by a host cell for expression of a polynucleotide encoding a polypeptide of the present invention. The promoter sequence contains transcriptional control sequences that mediate the expression of the polypeptide. The promoter may be any nucleotide sequence that shows transcriptional activity in the host cell of choice including mutant, truncated, and hybrid promoters, and may be obtained from genes encoding extracellular or intracellular polypeptides either homologous or heterologous to the host cell.

Examples of suitable promoters for directing the transcription of the nucleic acid constructs of the present invention, especially in a bacterial host cell, are the promoters obtained from the *E. coli lac* operon, *Streptomyces coelicolor* agarase gene (*dagA*), *Bacillus subtilis* levansucrase gene (*sacB*), *Bacillus licheniformis* alpha-amylase gene (*amyL*), *Bacillus stearothermophilus* maltogenic amylase gene (*amyM*), *Bacillus amyloliquefaciens* alpha-amylase gene (*amyQ*), *Bacillus licheniformis* penicillinase gene (*penP*), *Bacillus subtilis xylA* and *xylB* genes, and prokaryotic beta-lactamase gene (Villa-Kamaroff *et al.*, 1978, *Proceedings of the National Academy of Sciences USA* 75: 3727-3731), as well as the *tac* promoter (DeBoer *et al.*, 1983, *Proceedings of the National Academy of Sciences USA* 80: 21-25). Further promoters are described in "Useful proteins from recombinant bacteria" in *Scientific American*, 1980, 242: 74-94; and in Sambrook *et al.*, 1989, *supra*.

Examples of suitable promoters for directing the transcription of the nucleic acid constructs of the present invention in a filamentous fungal host cell are promoters obtained from the genes for *Aspergillus oryzae* TAKA amylase, *Rhizomucor miehei* aspartic proteinase, *Aspergillus niger* neutral alpha-amylase, *Aspergillus niger* acid stable alpha-amylase, *Aspergillus niger* or *Aspergillus awamori* glucoamylase (*glaA*), *Rhizomucor miehei* lipase, *Aspergillus oryzae* alkaline protease, *Aspergillus oryzae* triose phosphate isomerase,

*Aspergillus nidulans* acetamidase, *Fusarium venenatum* amyloglucosidase (WO 00/56900), *Fusarium venenatum* Daria (WO 00/56900), *Fusarium venenatum* Quinn (WO 00/56900), *Fusarium oxysporum* trypsin-like protease (WO 96/00787), *Trichoderma reesei* beta-glucosidase, *Trichoderma reesei* cellobiohydrolase I, *Trichoderma reesei* cellobiohydrolase II, *Trichoderma reesei* endoglucanase I, *Trichoderma reesei* endoglucanase II, *Trichoderma reesei* endoglucanase III, *Trichoderma reesei* endoglucanase IV, *Trichoderma reesei* endoglucanase V, *Trichoderma reesei* xylanase I, *Trichoderma reesei* xylanase II, *Trichoderma reesei* beta-xylosidase, as well as a NA2-tpi promoter (a modified promoter including a gene encoding a neutral alpha-amylase in *Aspergilli* in which the untranslated leader has been replaced by an untranslated leader from a gene encoding triose phosphate isomerase in *Aspergilli*; non-limiting examples include modified promoters including the gene encoding neutral alpha-amylase in *Aspergillus niger* in which the untranslated leader has been replaced by an untranslated leader from the gene encoding triose phosphate isomerase in *Aspergillus nidulans* or *Aspergillus oryzae*); and mutant, truncated, and hybrid promoters thereof.

In a yeast host, useful promoters are obtained from the genes for *Saccharomyces cerevisiae* enolase (ENO-1), *Saccharomyces cerevisiae* galactokinase (GAL1), *Saccharomyces cerevisiae* alcohol dehydrogenase/glyceraldehyde-3-phosphate dehydrogenase (ADH1, ADH2/GAP), *Saccharomyces cerevisiae* triose phosphate isomerase (TPI), *Saccharomyces cerevisiae* metallothionein (CUP1), and *Saccharomyces cerevisiae* 3-phosphoglycerate kinase. Other useful promoters for yeast host cells are described by Romanos *et al.*, 1992, *Yeast* 8: 423-488.

The control sequence may also be a suitable transcription terminator sequence, a sequence recognized by a host cell to terminate transcription. The terminator sequence is operably linked to the 3' terminus of the nucleotide sequence encoding the polypeptide. Any terminator that is functional in the host cell of choice may be used in the present invention.

Preferred terminators for filamentous fungal host cells are obtained from the genes for *Aspergillus oryzae* TAKA amylase, *Aspergillus niger* glucoamylase, *Aspergillus nidulans* anthranilate synthase, *Aspergillus niger* alpha-glucosidase, and *Fusarium oxysporum* trypsin-like protease.

Preferred terminators for yeast host cells are obtained from the genes for *Saccharomyces cerevisiae* enolase, *Saccharomyces cerevisiae* cytochrome C (CYC1), and *Saccharomyces cerevisiae* glyceraldehyde-3-phosphate dehydrogenase. Other useful terminators for yeast host cells are described by Romanos *et al.*, 1992, *supra*.

The control sequence may also be a suitable leader sequence, a nontranslated region of an mRNA that is important for translation by the host cell. The leader sequence is

operably linked to the 5' terminus of the nucleotide sequence encoding the polypeptide. Any leader sequence that is functional in the host cell of choice may be used in the present invention.

Preferred leaders for filamentous fungal host cells are obtained from the genes for *Aspergillus oryzae* TAKA amylase and *Aspergillus nidulans* triose phosphate isomerase.

Suitable leaders for yeast host cells are obtained from the genes for *Saccharomyces cerevisiae* enolase (ENO-1), *Saccharomyces cerevisiae* 3-phosphoglycerate kinase, *Saccharomyces cerevisiae* alpha-factor, and *Saccharomyces cerevisiae* alcohol dehydrogenase/glyceraldehyde-3-phosphate dehydrogenase (ADH2/GAP).

The control sequence may also be a polyadenylation sequence, a sequence operably linked to the 3' terminus of the nucleotide sequence and, when transcribed, is recognized by the host cell as a signal to add polyadenosine residues to transcribed mRNA. Any polyadenylation sequence that is functional in the host cell of choice may be used in the present invention.

Preferred polyadenylation sequences for filamentous fungal host cells are obtained from the genes for *Aspergillus oryzae* TAKA amylase, *Aspergillus niger* glucoamylase, *Aspergillus nidulans* anthranilate synthase, *Fusarium oxysporum* trypsin-like protease, and *Aspergillus niger* alpha-glucosidase.

Useful polyadenylation sequences for yeast host cells are described by Guo and Sherman, 1995, *Molecular Cellular Biology* 15: 5983-5990.

The control sequence may also be a signal peptide coding sequence that encodes a signal peptide linked to the amino terminus of a polypeptide and directs the encoded polypeptide into the cell's secretory pathway. The 5' end of the coding sequence of the nucleotide sequence may inherently contain a signal peptide coding sequence naturally linked in translation reading frame with the segment of the coding sequence that encodes the secreted polypeptide. Alternatively, the 5' end of the coding sequence may contain a signal peptide coding sequence that is foreign to the coding sequence. The foreign signal peptide coding sequence may be required where the coding sequence does not naturally contain a signal peptide coding sequence. Alternatively, the foreign signal peptide coding sequence may simply replace the natural signal peptide coding sequence in order to enhance secretion of the polypeptide. However, any signal peptide coding sequence that directs the expressed polypeptide into the secretory pathway of a host cell of choice, *i.e.*, secreted into a culture medium, may be used in the present invention.

Effective signal peptide coding sequences for bacterial host cells are the signal peptide coding sequences obtained from the genes for *Bacillus* NCIB 11837 maltogenic amylase, *Bacillus stearothermophilus* alpha-amylase, *Bacillus licheniformis* subtilisin,

*Bacillus licheniformis* beta-lactamase, *Bacillus stearothermophilus* neutral proteases (*nprT*, *nprS*, *nprM*), and *Bacillus subtilis* *prsA*. Further signal peptides are described by Simonen and Palva, 1993, *Microbiological Reviews* 57: 109-137.

5 Effective signal peptide coding sequences for filamentous fungal host cells are the signal peptide coding sequences obtained from the genes for *Aspergillus oryzae* TAKA amylase, *Aspergillus niger* neutral amylase, *Aspergillus niger* glucoamylase, *Rhizomucor miehei* aspartic proteinase, *Humicola insolens* cellulase, *Humicola insolens* endoglucanase V, and *Humicola lanuginosa* lipase.

10 Useful signal peptides for yeast host cells are obtained from the genes for *Saccharomyces cerevisiae* alpha-factor and *Saccharomyces cerevisiae* invertase. Other useful signal peptide coding sequences are described by Romanos *et al.*, 1992, *supra*.

15 The control sequence may also be a propeptide coding sequence that encodes a propeptide positioned at the amino terminus of a polypeptide. The resultant polypeptide is known as a proenzyme or propolypeptide (or a zymogen in some cases). A propeptide is generally inactive and can be converted to a mature active polypeptide by catalytic or autocatalytic cleavage of the propeptide from the propolypeptide. The propeptide coding sequence may be obtained from the genes for *Bacillus subtilis* alkaline protease (*aprE*), *Bacillus subtilis* neutral protease (*nprT*), *Saccharomyces cerevisiae* alpha-factor, *Rhizomucor miehei* aspartic proteinase, and *Myceliophthora thermophila* laccase (WO  
20 95/33836).

Where both signal peptide and propeptide sequences are present at the amino terminus of a polypeptide, the propeptide sequence is positioned next to the amino terminus of a polypeptide and the signal peptide sequence is positioned next to the amino terminus of the propeptide sequence.

25 It may also be desirable to add regulatory sequences that allow the regulation of the expression of the polypeptide relative to the growth of the host cell. Examples of regulatory systems are those that cause the expression of the gene to be turned on or off in response to a chemical or physical stimulus, including the presence of a regulatory compound. Regulatory systems in prokaryotic systems include the *lac*, *tac*, and *trp* operator systems. In  
30 yeast, the ADH2 system or GAL1 system may be used. In filamentous fungi, the TAKA alpha-amylase promoter, *Aspergillus niger* glucoamylase promoter, and *Aspergillus oryzae* glucoamylase promoter may be used as regulatory sequences. Other examples of regulatory sequences are those that allow for gene amplification. In eukaryotic systems, these regulatory sequences include the dihydrofolate reductase gene that is amplified in the  
35 presence of methotrexate, and the metallothionein genes that are amplified with heavy metals. In these cases, the nucleotide sequence encoding the polypeptide would be

operably linked with the regulatory sequence.

### Expression Vectors

5 The present invention also relates to recombinant expression vectors comprising a polynucleotide of the present invention, a promoter, and transcriptional and translational stop signals. The various nucleic acids and control sequences described herein may be joined together to produce a recombinant expression vector that may include one or more (several) convenient restriction sites to allow for insertion or substitution of the nucleotide sequence encoding the polypeptide at such sites. Alternatively, a polynucleotide sequence  
10 of the present invention may be expressed by inserting the nucleotide sequence or a nucleic acid construct comprising the sequence into an appropriate vector for expression. In creating the expression vector, the coding sequence is located in the vector so that the coding sequence is operably linked with the appropriate control sequences for expression.

The recombinant expression vector may be any vector (*e.g.*, a plasmid or virus) that  
15 can be conveniently subjected to recombinant DNA procedures and can bring about expression of the nucleotide sequence. The choice of the vector will typically depend on the compatibility of the vector with the host cell into which the vector is to be introduced. The vectors may be linear or closed circular plasmids.

The vector may be an autonomously replicating vector, *i.e.*, a vector that exists as an  
20 extrachromosomal entity, the replication of which is independent of chromosomal replication, *e.g.*, a plasmid, an extrachromosomal element, a minichromosome, or an artificial chromosome. The vector may contain any means for assuring self-replication. Alternatively, the vector may be one that, when introduced into the host cell, is integrated into the genome and replicated together with the chromosome(s) into which it has been  
25 integrated. Furthermore, a single vector or plasmid or two or more vectors or plasmids that together contain the total DNA to be introduced into the genome of the host cell, or a transposon, may be used.

The vectors of the present invention preferably contain one or more (several) selectable markers that permit easy selection of transformed, transfected, transduced, or  
30 the like cells. A selectable marker is a gene the product of which provides for biocide or viral resistance, resistance to heavy metals, prototrophy to auxotrophs, and the like.

Examples of bacterial selectable markers are the *dal* genes from *Bacillus subtilis* or *Bacillus licheniformis*, or markers that confer antibiotic resistance such as ampicillin, kanamycin, chloramphenicol, or tetracycline resistance. Suitable markers for yeast host  
35 cells are ADE2, HIS3, LEU2, LYS2, MET3, TRP1, and URA3. Selectable markers for use in a filamentous fungal host cell include, but are not limited to, *amdS* (acetamidase), *argB*

(ornithine carbamoyltransferase), *bar* (phosphinothricin acetyltransferase), *hph* (hygromycin phosphotransferase), *niaD* (nitrate reductase), *pyrG* (orotidine-5'-phosphate decarboxylase), *sC* (sulfate adenylyltransferase), and *trpC* (anthranilate synthase), as well as equivalents thereof. Preferred for use in an *Aspergillus* cell are the *amdS* and *pyrG* genes of *Aspergillus nidulans* or *Aspergillus oryzae* and the *bar* gene of *Streptomyces hygrosopicus*.

The vectors of the present invention preferably contain an element(s) that permits integration of the vector into the host cell's genome or autonomous replication of the vector in the cell independent of the genome.

For integration into the host cell genome, the vector may rely on the polynucleotide's sequence encoding the polypeptide or any other element of the vector for integration into the genome by homologous or nonhomologous recombination. Alternatively, the vector may contain additional nucleotide sequences for directing integration by homologous recombination into the genome of the host cell at a precise location(s) in the chromosome(s). To increase the likelihood of integration at a precise location, the integrational elements should preferably contain a sufficient number of nucleic acids, such as 100 to 10,000 base pairs, preferably 400 to 10,000 base pairs, and most preferably 800 to 10,000 base pairs, which have a high degree of sequence identity to the corresponding target sequence to enhance the probability of homologous recombination. The integrational elements may be any sequence that is homologous with the target sequence in the genome of the host cell. Furthermore, the integrational elements may be non-encoding or encoding nucleotide sequences. On the other hand, the vector may be integrated into the genome of the host cell by non-homologous recombination.

For autonomous replication, the vector may further comprise an origin of replication enabling the vector to replicate autonomously in the host cell in question. The origin of replication may be any plasmid replicator mediating autonomous replication that functions in a cell. The term "origin of replication" or "plasmid replicator" is defined herein as a nucleotide sequence that enables a plasmid or vector to replicate *in vivo*.

Examples of bacterial origins of replication are the origins of replication of plasmids pBR322, pUC19, pACYC177, and pACYC184 permitting replication in *E. coli*, and pUB110, pE194, pTA1060, and pAM $\beta$ 1 permitting replication in *Bacillus*.

Examples of origins of replication for use in a yeast host cell are the 2 micron origin of replication, ARS1, ARS4, the combination of ARS1 and CEN3, and the combination of ARS4 and CEN6.

Examples of origins of replication useful in a filamentous fungal cell are AMA1 and ANS1 (Gems *et al.*, 1991, *Gene* 98: 61-67; Cullen *et al.*, 1987, *Nucleic Acids Research* 15: 9163-9175; WO 00/24883). Isolation of the AMA1 gene and construction of plasmids or



vectors comprising the gene can be accomplished according to the methods disclosed in WO 00/24883.

More than one copy of a polynucleotide of the present invention may be inserted into a host cell to increase production of the gene product. An increase in the copy number of the polynucleotide can be obtained by integrating at least one additional copy of the sequence into the host cell genome or by including an amplifiable selectable marker gene with the polynucleotide where cells containing amplified copies of the selectable marker gene, and thereby additional copies of the polynucleotide, can be selected for by cultivating the cells in the presence of the appropriate selectable agent.

The procedures used to ligate the elements described above to construct the recombinant expression vectors of the present invention are well known to one skilled in the art (see, *e.g.*, Sambrook *et al.*, 1989, *supra*).

### Host Cells

The present invention also relates to recombinant host cells, comprising an isolated polynucleotide of the present invention operably linked to one or more (several) control sequences that direct the production of a polypeptide having alpha-mannosidase activity. A construct or vector comprising a polynucleotide of the present invention is introduced into a host cell so that the construct or vector is maintained as a chromosomal integrant or as a self-replicating extra-chromosomal vector as described earlier. The term "host cell" encompasses any progeny of a parent cell that is not identical to the parent cell due to mutations that occur during replication. The choice of a host cell will to a large extent depend upon the gene encoding the polypeptide and its source.

The host cell may be any cell useful in the recombinant production of a polypeptide of the present invention, *e.g.*, a prokaryote or a eukaryote.

The prokaryotic host cell may be any Gram positive bacterium or a Gram negative bacterium. Gram positive bacteria include, but not limited to, *Bacillus*, *Streptococcus*, *Streptomyces*, *Staphylococcus*, *Enterococcus*, *Lactobacillus*, *Lactococcus*, *Clostridium*, *Geobacillus*, and *Oceanobacillus*. Gram negative bacteria include, but not limited to, *E. coli*, *Pseudomonas*, *Salmonella*, *Campylobacter*, *Helicobacter*, *Flavobacterium*, *Fusobacterium*, *Ilyobacter*, *Neisseria*, and *Ureaplasma*.

The bacterial host cell may be any *Bacillus* cell. *Bacillus* cells useful in the practice of the present invention include, but are not limited to, *Bacillus alkalophilus*, *Bacillus amyloliquefaciens*, *Bacillus brevis*, *Bacillus circulans*, *Bacillus clausii*, *Bacillus coagulans*, *Bacillus firmus*, *Bacillus lautus*, *Bacillus lentus*, *Bacillus licheniformis*, *Bacillus megaterium*, *Bacillus pumilus*, *Bacillus stearothermophilus*, *Bacillus subtilis*, and *Bacillus thuringiensis*

cells.

In a preferred aspect, the bacterial host cell is a *Bacillus amyloliquefaciens* cell. In another preferred aspect, the bacterial host cell is a *Bacillus clausii* cell. In another preferred aspect, the bacterial host cell is a *Bacillus lentus* cell. In another preferred aspect, the bacterial host cell is a *Bacillus licheniformis* cell. In another preferred aspect, the bacterial host cell is a *Bacillus stearothermophilus* cell. In another preferred aspect, the bacterial host cell is a *Bacillus subtilis* cell.

The bacterial host cell may also be any *Streptococcus* cell. *Streptococcus* cells useful in the practice of the present invention include, but are not limited to, *Streptococcus equisimilis*, *Streptococcus pyogenes*, *Streptococcus uberis*, and *Streptococcus equi* subsp. *Zooepidemicus* cells.

In a preferred aspect, the bacterial host cell is a *Streptococcus equisimilis* cell. In another preferred aspect, the bacterial host cell is a *Streptococcus pyogenes* cell. In another preferred aspect, the bacterial host cell is a *Streptococcus uberis* cell. In another preferred aspect, the bacterial host cell is a *Streptococcus equi* subsp. *Zooepidemicus* cell.

The bacterial host cell may also be any *Streptomyces* cell. *Streptomyces* cells useful in the practice of the present invention include, but are not limited to, *Streptomyces achromogenes*, *Streptomyces avermitilis*, *Streptomyces coelicolor*, *Streptomyces griseus*, and *Streptomyces lividans* cells.

In a preferred aspect, the bacterial host cell is a *Streptomyces achromogenes* cell. In another preferred aspect, the bacterial host cell is a *Streptomyces avermitilis* cell. In another preferred aspect, the bacterial host cell is a *Streptomyces coelicolor* cell. In another preferred aspect, the bacterial host cell is a *Streptomyces griseus* cell. In another preferred aspect, the bacterial host cell is a *Streptomyces lividans* cell.

The introduction of DNA into a *Bacillus* cell may, for instance, be effected by protoplast transformation (see, e.g., Chang and Cohen, 1979, *Molecular General Genetics* 168: 111-115), by using competent cells (see, e.g., Young and Spizizen, 1961, *Journal of Bacteriology* 81: 823-829, or Dubnau and Davidoff-Abelson, 1971, *Journal of Molecular Biology* 56: 209-221), by electroporation (see, e.g., Shigekawa and Dower, 1988, *Biotechniques* 6: 742-751), or by conjugation (see, e.g., Koehler and Thorne, 1987, *Journal of Bacteriology* 169: 5271-5278). The introduction of DNA into an *E coli* cell may, for instance, be effected by protoplast transformation (see, e.g., Hanahan, 1983, *J. Mol. Biol.* 166: 557-580) or electroporation (see, e.g., Dower et al., 1988, *Nucleic Acids Res.* 16: 6127-6145). The introduction of DNA into a *Streptomyces* cell may, for instance, be effected by protoplast transformation and electroporation (see, e.g., Gong et al., 2004, *Folia Microbiol. (Praha)* 49: 399-405), by conjugation (see, e.g., Mazodier et al., 1989, *J. Bacteriol.* 171:

3583-3585), or by transduction (see, e.g., Burke *et al.*, 2001, *Proc. Natl. Acad. Sci. USA* 98: 6289-6294). The introduction of DNA into a *Pseudomonas* cell may, for instance, be effected by electroporation (see, e.g., Choi *et al.*, 2006, *J. Microbiol. Methods* 64: 391-397) or by conjugation (see, e.g., Pinedo and Smets, 2005, *Appl. Environ. Microbiol.* 71: 51-57).

5 The introduction of DNA into a *Streptococcus* cell may, for instance, be effected by natural competence (see, e.g., Perry and Kuramitsu, 1981, *Infect. Immun.* 32: 1295-1297), by protoplast transformation (see, e.g., Catt and Jollick, 1991, *Microbios.* 68: 189-207, by electroporation (see, e.g., Buckley *et al.*, 1999, *Appl. Environ. Microbiol.* 65: 3800-3804) or by conjugation (see, e.g., Clewell, 1981, *Microbiol. Rev.* 45: 409-436). However, any  
10 method known in the art for introducing DNA into a host cell can be used.

The host cell may also be a eukaryote, such as a mammalian, insect, plant, or fungal cell.

In a preferred aspect, the host cell is a fungal cell. "Fungi" as used herein includes the phyla Ascomycota, Basidiomycota, Chytridiomycota, and Zygomycota (as defined by  
15 Hawksworth *et al.*, In, *Ainsworth and Bisby's Dictionary of The Fungi*, 8th edition, 1995, CAB International, University Press, Cambridge, UK) as well as the Oomycota (as cited in Hawksworth *et al.*, 1995, *supra*, page 171) and all mitosporic fungi (Hawksworth *et al.*, 1995, *supra*).

In a more preferred aspect, the fungal host cell is a yeast cell. "Yeast" as used  
20 herein includes ascosporogenous yeast (Endomycetales), basidiosporogenous yeast, and yeast belonging to the Fungi Imperfecti (Blastomycetes). Since the classification of yeast may change in the future, for the purposes of this invention, yeast shall be defined as described in *Biology and Activities of Yeast* (Skinner, F.A., Passmore, S.M., and Davenport, R.R., eds, *Soc. App. Bacteriol. Symposium Series No. 9*, 1980).

25 In an even more preferred aspect, the yeast host cell is a *Candida*, *Hansenula*, *Kluyveromyces*, *Pichia*, *Saccharomyces*, *Schizosaccharomyces*, or *Yarrowia* cell.

In a most preferred aspect, the yeast host cell is a *Saccharomyces carlsbergensis* cell. In another most preferred aspect, the yeast host cell is a *Saccharomyces cerevisiae* cell. In another most preferred aspect, the yeast host cell is a *Saccharomyces diastaticus*  
30 cell. In another most preferred aspect, the yeast host cell is a *Saccharomyces douglasii* cell. In another most preferred aspect, the yeast host cell is a *Saccharomyces kluyveri* cell.

In another most preferred aspect, the yeast host cell is a *Saccharomyces norbensis* cell. In another most preferred aspect, the yeast host cell is a *Saccharomyces oviformis* cell. In another most preferred aspect, the yeast host cell is a *Kluyveromyces lactis* cell. In another  
35 most preferred aspect, the yeast host cell is a *Yarrowia lipolytica* cell.

In another more preferred aspect, the fungal host cell is a filamentous fungal cell.

“Filamentous fungi” include all filamentous forms of the subdivision Eumycota and Oomycota (as defined by Hawksworth *et al.*, 1995, *supra*). The filamentous fungi are generally characterized by a mycelial wall composed of chitin, cellulose, glucan, chitosan, mannan, and other complex polysaccharides. Vegetative growth is by hyphal elongation and carbon catabolism is obligately aerobic. In contrast, vegetative growth by yeasts such as *Saccharomyces cerevisiae* is by budding of a unicellular thallus and carbon catabolism may be fermentative.

In an even more preferred aspect, the filamentous fungal host cell is an *Acremonium*, *Aspergillus*, *Aureobasidium*, *Bjerkandera*, *Ceriporiopsis*, *Chrysosporium*, *Coprinus*, *Coriolus*, *Cryptococcus*, *Filibasidium*, *Fusarium*, *Humicola*, *Magnaporthe*, *Mucor*, *Myceliophthora*, *Neocallimastix*, *Neurospora*, *Paecilomyces*, *Penicillium*, *Phanerochaete*, *Phlebia*, *Piromyces*, *Pleurotus*, *Schizophyllum*, *Talaromyces*, *Thermoascus*, *Thielavia*, *Tolyocladium*, *Trametes*, or *Trichoderma* cell.

In a most preferred aspect, the filamentous fungal host cell is an *Aspergillus awamori*, *Aspergillus fumigatus*, *Aspergillus foetidus*, *Aspergillus japonicus*, *Aspergillus nidulans*, *Aspergillus niger* or *Aspergillus oryzae* cell. In another most preferred aspect, the filamentous fungal host cell is a *Fusarium bactridioides*, *Fusarium cerealis*, *Fusarium crookwellense*, *Fusarium culmorum*, *Fusarium graminearum*, *Fusarium graminum*, *Fusarium heterosporum*, *Fusarium negundi*, *Fusarium oxysporum*, *Fusarium reticulatum*, *Fusarium roseum*, *Fusarium sambucinum*, *Fusarium sarcochroum*, *Fusarium sporotrichioides*, *Fusarium sulphureum*, *Fusarium torulosum*, *Fusarium trichothecioides*, or *Fusarium venenatum* cell. In another most preferred aspect, the filamentous fungal host cell is a *Bjerkandera adusta*, *Ceriporiopsis aneirina*, *Ceriporiopsis aneirina*, *Ceriporiopsis caregiea*, *Ceriporiopsis gilvescens*, *Ceriporiopsis pannocinta*, *Ceriporiopsis rivulosa*, *Ceriporiopsis subrufa*, *Ceriporiopsis subvermispora*, *Chrysosporium keratinophilum*, *Chrysosporium lucknowense*, *Chrysosporium tropicum*, *Chrysosporium merdarium*, *Chrysosporium inops*, *Chrysosporium pannicola*, *Chrysosporium queenslandicum*, *Chrysosporium zonatum*, *Coprinus cinereus*, *Coriolus hirsutus*, *Humicola insolens*, *Humicola lanuginosa*, *Mucor miehei*, *Myceliophthora thermophila*, *Neurospora crassa*, *Penicillium purpurogenum*, *Phanerochaete chrysosporium*, *Phlebia radiata*, *Pleurotus eryngii*, *Thielavia terrestris*, *Trametes villosa*, *Trametes versicolor*, *Trichoderma harzianum*, *Trichoderma koningii*, *Trichoderma longibrachiatum*, *Trichoderma reesei*, or *Trichoderma viride* cell.

In another most preferred aspect, the filamentous fungal host cell is an *Aspergillus niger* cell. In another most preferred aspect, the filamentous fungal host cell is an *Aspergillus oryzae* cell. In another most preferred aspect, the filamentous fungal host cell is a *Chrysosporium lucknowense* cell. In another most preferred aspect, the filamentous

5 fungal host cell is a *Fusarium venenatum* cell. In another most preferred aspect, the filamentous fungal host cell is a *Myceliophthora thermophila* cell. In another most preferred aspect, the filamentous fungal host cell is a *Trichoderma reesei* cell.

5 Fungal cells may be transformed by a process involving protoplast formation, transformation of the protoplasts, and regeneration of the cell wall in a manner known *per se*. Suitable procedures for transformation of *Aspergillus* and *Trichoderma* host cells are described in EP 238 023 and Yelton *et al.*, 1984, *Proceedings of the National Academy of Sciences USA* 81: 1470-1474. Suitable methods for transforming *Fusarium* species are described by Malardier *et al.*, 1989, *Gene* 78: 147-156, and WO 96/00787. Yeast may be  
10 transformed using the procedures described by Becker and Guarente, *In* Abelson, J.N. and Simon, M.I., editors, *Guide to Yeast Genetics and Molecular Biology, Methods in Enzymology*, Volume 194, pp 182-187, Academic Press, Inc., New York; Ito *et al.*, 1983, *Journal of Bacteriology* 153: 163; and Hinnen *et al.*, 1978, *Proceedings of the National Academy of Sciences USA* 75: 1920.

#### 15 **Methods of Production**

The present invention also relates to methods of producing a polypeptide of the present invention, comprising: (a) cultivating a cell, which in its wild-type form produces the polypeptide, under conditions conducive for production of the polypeptide; and (b)  
20 recovering the polypeptide. In a preferred aspect, the cell is of the genus *Thielavia*. In a more preferred aspect, the cell is *Thielavia terrestris*. In a most preferred aspect, the cell is *Thielavia terrestris* NRRL 8126.

The present invention also relates to methods of producing a polypeptide of the present invention, comprising: (a) cultivating a recombinant host cell, as described herein,  
25 under conditions conducive for production of the polypeptide; and (b) recovering the polypeptide.

The present invention also relates to methods of producing a polypeptide of the present invention, comprising: (a) cultivating a recombinant host cell under conditions conducive for production of the polypeptide, wherein the host cell comprises a mutant  
30 nucleotide sequence having at least one mutation in the mature polypeptide coding sequence of SEQ ID NO: 1, wherein the mutant nucleotide sequence encodes a polypeptide that comprises or consists of the mature polypeptide of SEQ ID NO: 2; and (b) recovering the polypeptide.

In the production methods of the present invention, the cells are cultivated in a  
35 nutrient medium suitable for production of the polypeptide using methods well known in the art. For example, the cell may be cultivated by shake flask cultivation, and small-scale or

5 large-scale fermentation (including continuous, batch, fed-batch, or solid state fermentations) in laboratory or industrial fermentors performed in a suitable medium and under conditions allowing the polypeptide to be expressed and/or isolated. The cultivation takes place in a suitable nutrient medium comprising carbon and nitrogen sources and inorganic salts, using procedures known in the art. Suitable media are available from commercial suppliers or may be prepared according to published compositions (*e.g.*, in catalogues of the American Type Culture Collection). If the polypeptide is secreted into the nutrient medium, the polypeptide can be recovered directly from the medium. If the polypeptide is not secreted into the medium, it can be recovered from cell lysates.

10 The polypeptides may be detected using methods known in the art that are specific for the polypeptides. These detection methods may include use of specific antibodies, formation of an enzyme product, or disappearance of an enzyme substrate. For example, an enzyme assay may be used to determine the activity of the polypeptide as described herein.

15 The resulting polypeptide may be recovered using methods known in the art. For example, the polypeptide may be recovered from the nutrient medium by conventional procedures including, but not limited to, centrifugation, filtration, extraction, spray-drying, evaporation, or precipitation.

20 The polypeptides of the present invention may be purified by a variety of procedures known in the art including, but not limited to, chromatography (*e.g.*, ion exchange, affinity, hydrophobic, chromatofocusing, and size exclusion), electrophoretic procedures (*e.g.*, preparative isoelectric focusing), differential solubility (*e.g.*, ammonium sulfate precipitation), SDS-PAGE, or extraction (see, *e.g.*, *Protein Purification*, J.-C. Janson and Lars Ryden, editors, VCH Publishers, New York, 1989) to obtain substantially pure polypeptides.

## 25 **Plants**

30 The present invention also relates to plants, *e.g.*, a transgenic plant, plant part, or plant cell, comprising an isolated polynucleotide encoding a polypeptide having alpha-mannosidase activity of the present invention so as to express and produce the polypeptide in recoverable quantities. The polypeptide may be recovered from the plant or plant part. Alternatively, the plant or plant part containing the recombinant polypeptide may be used as such for improving the quality of a food or feed, *e.g.*, improving nutritional value, palatability, and rheological properties, or to destroy an antinutritive factor.

35 The transgenic plant can be dicotyledonous (a dicot) or monocotyledonous (a monocot). Examples of monocot plants are grasses, such as meadow grass (blue grass, *Poa*), forage grass such as *Festuca*, *Lolium*, temperate grass, such as *Agrostis*, and

cereals, e.g., wheat, oats, rye, barley, rice, sorghum, and maize (corn).

Examples of dicot plants are tobacco, legumes, such as lupins, potato, sugar beet, pea, bean and soybean, and cruciferous plants (family Brassicaceae), such as cauliflower, rape seed, and the closely related model organism *Arabidopsis thaliana*.

5           Examples of plant parts are stem, callus, leaves, root, fruits, seeds, and tubers as well as the individual tissues comprising these parts, e.g., epidermis, mesophyll, parenchyme, vascular tissues, meristems. Specific plant cell compartments, such as chloroplasts, apoplasts, mitochondria, vacuoles, peroxisomes and cytoplasm are also considered to be a plant part. Furthermore, any plant cell, whatever the tissue origin, is  
10 considered to be a plant part. Likewise, plant parts such as specific tissues and cells isolated to facilitate the utilisation of the invention are also considered plant parts, e.g., embryos, endosperms, aleurone and seeds coats.

Also included within the scope of the present invention are the progeny of such plants, plant parts, and plant cells.

15           The transgenic plant or plant cell expressing a polypeptide of the present invention may be constructed in accordance with methods known in the art. In short, the plant or plant cell is constructed by incorporating one or more (several) expression constructs encoding a polypeptide of the present invention into the plant host genome or chloroplast genome and propagating the resulting modified plant or plant cell into a transgenic plant or  
20 plant cell.

The expression construct is conveniently a nucleic acid construct that comprises a polynucleotide encoding a polypeptide of the present invention operably linked with appropriate regulatory sequences required for expression of the nucleotide sequence in the plant or plant part of choice. Furthermore, the expression construct may comprise a  
25 selectable marker useful for identifying host cells into which the expression construct has been integrated and DNA sequences necessary for introduction of the construct into the plant in question (the latter depends on the DNA introduction method to be used).

The choice of regulatory sequences, such as promoter and terminator sequences and optionally signal or transit sequences, is determined, for example, on the basis of when,  
30 where, and how the polypeptide is desired to be expressed. For instance, the expression of the gene encoding a polypeptide of the present invention may be constitutive or inducible, or may be developmental, stage or tissue specific, and the gene product may be targeted to a specific tissue or plant part such as seeds or leaves. Regulatory sequences are, for example, described by Tague *et al.*, 1988, *Plant Physiology* 86: 506.

35           For constitutive expression, the 35S-CaMV, the maize ubiquitin 1, and the rice actin 1 promoter may be used (Franck *et al.*, 1980, *Cell* 21: 285-294; Christensen *et al.*, 1992,

*Plant Mol. Biol.* 18: 675-689; Zhang *et al.*, 1991, *Plant Cell* 3: 1155-1165). Organ-specific promoters may be, for example, a promoter from storage sink tissues such as seeds, potato tubers, and fruits (Edwards and Coruzzi, 1990, *Ann. Rev. Genet.* 24: 275-303), or from metabolic sink tissues such as meristems (Ito *et al.*, 1994, *Plant Mol. Biol.* 24: 863-878), a seed specific promoter such as the glutelin, prolamin, globulin, or albumin promoter from rice (Wu *et al.*, 1998, *Plant and Cell Physiology* 39: 885-889), a *Vicia faba* promoter from the legumin B4 and the unknown seed protein gene from *Vicia faba* (Conrad *et al.*, 1998, *Journal of Plant Physiology* 152: 708-711), a promoter from a seed oil body protein (Chen *et al.*, 1998, *Plant and Cell Physiology* 39: 935-941), the storage protein *napA* promoter from *Brassica napus*, or any other seed specific promoter known in the art, *e.g.*, as described in WO 91/14772. Furthermore, the promoter may be a leaf specific promoter such as the *rbcS* promoter from rice or tomato (Kyoizuka *et al.*, 1993, *Plant Physiology* 102: 991-1000, the chlorella virus adenine methyltransferase gene promoter (Mitra and Higgins, 1994, *Plant Molecular Biology* 26: 85-93), or the *aldP* gene promoter from rice (Kagaya *et al.*, 1995, *Molecular and General Genetics* 248: 668-674), or a wound inducible promoter such as the potato *pin2* promoter (Xu *et al.*, 1993, *Plant Molecular Biology* 22: 573-588). Likewise, the promoter may be inducible by abiotic treatments such as temperature, drought, or alterations in salinity or induced by exogenously applied substances that activate the promoter, *e.g.*, ethanol, oestrogens, plant hormones such as ethylene, abscisic acid, and gibberellic acid, and heavy metals.

A promoter enhancer element may also be used to achieve higher expression of a polypeptide of the present invention in the plant. For instance, the promoter enhancer element may be an intron that is placed between the promoter and the nucleotide sequence encoding a polypeptide of the present invention. For instance, Xu *et al.*, 1993, *supra*, disclose the use of the first intron of the rice actin 1 gene to enhance expression.

The selectable marker gene and any other parts of the expression construct may be chosen from those available in the art.

The nucleic acid construct is incorporated into the plant genome according to conventional techniques known in the art, including *Agrobacterium*-mediated transformation, virus-mediated transformation, microinjection, particle bombardment, biolistic transformation, and electroporation (Gasser *et al.*, 1990, *Science* 244: 1293; Potrykus, 1990, *Bio/Technology* 8: 535; Shimamoto *et al.*, 1989, *Nature* 338: 274).

Presently, *Agrobacterium tumefaciens*-mediated gene transfer is the method of choice for generating transgenic dicots (for a review, see Hooykas and Schilperoort, 1992, *Plant Molecular Biology* 19: 15-38) and can also be used for transforming monocots, although other transformation methods are often used for these plants. Presently, the



method of choice for generating transgenic monocots is particle bombardment (microscopic gold or tungsten particles coated with the transforming DNA) of embryonic calli or developing embryos (Christou, 1992, *Plant Journal* 2: 275-281; Shimamoto, 1994, *Current Opinion Biotechnology* 5: 158-162; Vasil *et al.*, 1992, *Bio/Technology* 10: 667-674). An  
5 alternative method for transformation of monocots is based on protoplast transformation as described by Omirulleh *et al.*, 1993, *Plant Molecular Biology* 21: 415-428. Additional transformation methods for use in accordance with the present disclosure include those described in U.S. Patent Nos. 6,395,966 and 7,151,204 (both of which are herein incorporated by reference in their entirety).

10 Following transformation, the transformants having incorporated the expression construct are selected and regenerated into whole plants according to methods well-known in the art. Often the transformation procedure is designed for the selective elimination of selection genes either during regeneration or in the following generations by using, for example, co-transformation with two separate T-DNA constructs or site specific excision of  
15 the selection gene by a specific recombinase.

The present invention also relates to methods of producing a polypeptide of the present invention comprising: (a) cultivating a transgenic plant or a plant cell comprising a polynucleotide encoding the polypeptide having alpha-mannosidase activity of the present invention under conditions conducive for production of the polypeptide; and (b) recovering  
20 the polypeptide.

In embodiments, in addition to direct transformation of a particular plant genotype with a construct prepared according to the present invention, transgenic plants may be made by crossing a plant having a construct of the present invention to a second plant lacking the construct. For example, a construct encoding a polypeptide having alpha-mannosidase activity or a portion thereof can be introduced into a particular plant variety by  
25 crossing, without the need for ever directly transforming a plant of that given variety. Therefore, the present invention not only encompasses a plant directly regenerated from cells which have been transformed in accordance with the present invention, but also the progeny of such plants. As used herein, progeny may refer to the offspring of any generation of a parent plant prepared in accordance with the present invention. Such progeny may include a DNA construct prepared in accordance with the present invention, or a portion of a DNA construct prepared in accordance with the present invention. In  
30 embodiments, crossing results in a transgene of the present invention being introduced into a plant line by cross pollinating a starting line with a donor plant line that includes a transgene of the present invention. Non-limiting examples of such steps are further  
35 articulated in U.S. Patent No: 7,151,204.

It is envisioned that plants including a polypeptide having alpha-mannosidase activity of the present invention include plants generated through a process of backcross conversion. For examples, plants of the present invention include plants referred to as a backcross converted genotype, line, inbred, or hybrid.

5 In embodiments, genetic markers may be used to assist in the introgression of one or more transgenes of the invention from one genetic background into another. Marker assisted selection offers advantages relative to conventional breeding in that it can be used to avoid errors caused by phenotypic variations. Further, genetic markers may provide data regarding the relative degree of elite germplasm in the individual progeny of a particular  
10 cross. For example, when a plant with a desired trait which otherwise has a non-agronomically desirable genetic background is crossed to an elite parent, genetic markers may be used to select progeny which not only possess the trait of interest, but also have a relatively large proportion of the desired germplasm. In this way, the number of generations required to introgress one or more traits into a particular genetic background is minimized.

#### 15 **Removal or Reduction of Alpha-Mannosidase Activity**

The present invention also relates to methods of producing a mutant of a parent cell, which comprises disrupting or deleting a polynucleotide, or a portion thereof, encoding a polypeptide of the present invention, which results in the mutant cell producing less of the  
20 polypeptide than the parent cell when cultivated under the same conditions.

The mutant cell may be constructed by reducing or eliminating expression of a nucleotide sequence encoding a polypeptide of the present invention using methods well known in the art, for example, insertions, disruptions, replacements, or deletions. In a preferred aspect, the nucleotide sequence is inactivated. The nucleotide sequence to be  
25 modified or inactivated may be, for example, the coding region or a part thereof essential for activity, or a regulatory element required for the expression of the coding region. An example of such a regulatory or control sequence may be a promoter sequence or a functional part thereof, *i.e.*, a part that is sufficient for affecting expression of the nucleotide sequence. Other control sequences for possible modification include, but are not limited to,  
30 a leader, polyadenylation sequence, propeptide sequence, signal peptide sequence, transcription terminator, and transcriptional activator.

Modification or inactivation of the nucleotide sequence may be performed by  
35 subjecting the parent cell to mutagenesis and selecting for mutant cells in which expression of the nucleotide sequence has been reduced or eliminated. The mutagenesis, which may be specific or random, may be performed, for example, by use of a suitable physical or chemical mutagenizing agent, by use of a suitable oligonucleotide, or by subjecting the DNA

sequence to PCR generated mutagenesis. Furthermore, the mutagenesis may be performed by use of any combination of these mutagenizing agents.

5 Examples of a physical or chemical mutagenizing agent suitable for the present purpose include ultraviolet (UV) irradiation, hydroxylamine, N-methyl-N'-nitro-N-nitrosoguanidine (MNNG), O-methyl hydroxylamine, nitrous acid, ethyl methane sulphonate (EMS), sodium bisulphite, formic acid, and nucleotide analogues.

10 When such agents are used, the mutagenesis is typically performed by incubating the parent cell to be mutagenized in the presence of the mutagenizing agent of choice under suitable conditions, and screening and/or selecting for mutant cells exhibiting reduced or no expression of the gene.

15 Modification or inactivation of the nucleotide sequence may be accomplished by introduction, substitution, or removal of one or more (several) nucleotides in the gene or a regulatory element required for the transcription or translation thereof. For example, nucleotides may be inserted or removed so as to result in the introduction of a stop codon, the removal of the start codon, or a change in the open reading frame. Such modification or inactivation may be accomplished by site-directed mutagenesis or PCR generated mutagenesis in accordance with methods known in the art. Although, in principle, the modification may be performed *in vivo*, *i.e.*, directly on the cell expressing the nucleotide sequence to be modified, it is preferred that the modification be performed *in vitro* as exemplified below.

20 An example of a convenient way to eliminate or reduce expression of a nucleotide sequence by a cell is based on techniques of gene replacement, gene deletion, or gene disruption. For example, in the gene disruption method, a nucleic acid sequence corresponding to the endogenous nucleotide sequence is mutagenized *in vitro* to produce a defective nucleic acid sequence that is then transformed into the parent cell to produce a defective gene. By homologous recombination, the defective nucleic acid sequence replaces the endogenous nucleotide sequence. It may be desirable that the defective nucleotide sequence also encodes a marker that may be used for selection of transformants in which the nucleotide sequence has been modified or destroyed. In a particularly preferred aspect, the nucleotide sequence is disrupted with a selectable marker such as those described herein.

35 Alternatively, modification or inactivation of the nucleotide sequence may be performed by established anti-sense or RNAi techniques using a sequence complementary to the nucleotide sequence. More specifically, expression of the nucleotide sequence by a cell may be reduced or eliminated by introducing a sequence complementary to the nucleotide sequence of the gene that may be transcribed in the cell and is capable of

hybridizing to the mRNA produced in the cell. Under conditions allowing the complementary anti-sense nucleotide sequence to hybridize to the mRNA, the amount of protein translated is thus reduced or eliminated.

5 The present invention further relates to a mutant cell of a parent cell that comprises a disruption or deletion of a nucleotide sequence encoding the polypeptide or a control sequence thereof, which results in the mutant cell producing less of the polypeptide or no polypeptide compared to the parent cell.

10 The polypeptide-deficient mutant cells so created are particularly useful as host cells for the expression of native and/or heterologous polypeptides. Therefore, the present invention further relates to methods of producing a native or heterologous polypeptide, comprising: (a) cultivating the mutant cell under conditions conducive for production of the polypeptide; and (b) recovering the polypeptide. The term "heterologous polypeptides" is defined herein as polypeptides that are not native to the host cell, a native protein in which modifications have been made to alter the native sequence, or a native protein whose  
15 expression is quantitatively altered as a result of a manipulation of the host cell by recombinant DNA techniques.

In a further aspect, the present invention relates to a method of producing a protein product essentially free of alpha-mannosidase activity by fermentation of a cell that produces both a polypeptide of the present invention as well as the protein product of  
20 interest by adding an effective amount of an agent capable of inhibiting alpha-mannosidase activity to the fermentation broth before, during, or after the fermentation has been completed, recovering the product of interest from the fermentation broth, and optionally subjecting the recovered product to further purification.

25 In a further aspect, the present invention relates to a method of producing a protein product essentially free of alpha-mannosidase activity by cultivating the cell under conditions permitting the expression of the product, subjecting the resultant culture broth to a combined pH and temperature treatment so as to reduce the alpha-mannosidase activity substantially, and recovering the product from the culture broth. Alternatively, the combined pH and temperature treatment may be performed on an enzyme preparation recovered from the  
30 culture broth. The combined pH and temperature treatment may optionally be used in combination with a treatment with an alpha-mannosidase inhibitor.

In accordance with this aspect of the invention, it is possible to remove at least 60%, preferably at least 75%, more preferably at least 85%, still more preferably at least 95%, and most preferably at least 99% of the alpha-mannosidase activity. Complete removal of  
35 alpha-mannosidase activity may be obtained by use of this method.

The combined pH and temperature treatment is preferably carried out at a pH in the

range of 2-4 or 9-11 and a temperature in the range of at least 60-70°C for a sufficient period of time to attain the desired effect, where typically, 30 to 60 minutes is sufficient.

The methods used for cultivation and purification of the product of interest may be performed by methods known in the art.

5           The methods of the present invention for producing an essentially alpha-mannosidase-free product is of particular interest in the production of eukaryotic polypeptides, in particular fungal proteins such as enzymes. The enzyme may be selected from, *e.g.*, an amylolytic enzyme, lipolytic enzyme, proteolytic enzyme, cellulolytic enzyme, oxidoreductase, or plant cell-wall degrading enzyme. Examples of such enzymes include an  
10   aminopeptidase, amylase, amyloglucosidase, carbohydrase, carboxypeptidase, catalase, cellobiohydrolase, cellulase, chitinase, cutinase, cyclodextrin glycosyltransferase, deoxyribonuclease, endoglucanase, esterase, galactosidase, beta-galactosidase, glucoamylase, glucose oxidase, glucosidase, haloperoxidase, hemicellulase, invertase, isomerase, laccase, ligase, lipase, lyase, mannosidase, oxidase, pectinolytic enzyme,  
15   peroxidase, phytase, phenoloxidase, polyphenoloxidase, proteolytic enzyme, ribonuclease, transferase, transglutaminase, or xylanase. The alpha-mannosidase-deficient cells may also be used to express heterologous proteins of pharmaceutical interest such as hormones, growth factors, receptors, and the like.

It will be understood that the term "eukaryotic polypeptides" includes not only native  
20   polypeptides, but also those polypeptides, *e.g.*, enzymes, which have been modified by amino acid substitutions, deletions or additions, or other such modifications to enhance activity, thermostability, pH tolerance and the like.

In a further aspect, the present invention relates to a protein product essentially free from alpha-mannosidase activity that is produced by a method of the present invention.

25

### **Methods of Inhibiting Expression of a Polypeptide Having Alpha-Mannosidase Activity**

The present invention also relates to methods of inhibiting the expression of a polypeptide having alpha-mannosidase activity in a cell, comprising administering to the cell  
30   or expressing in the cell a double-stranded RNA (dsRNA) molecule, wherein the dsRNA comprises a subsequence of a polynucleotide of the present invention. In a preferred aspect, the dsRNA is about 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25 or more duplex nucleotides in length.

The dsRNA is preferably a small interfering RNA (siRNA) or a micro RNA (miRNA).  
35   In a preferred aspect, the dsRNA is small interfering RNA (siRNAs) for inhibiting

transcription. In another preferred aspect, the dsRNA is micro RNA (miRNAs) for inhibiting translation.

The present invention also relates to such double-stranded RNA (dsRNA) molecules, comprising a portion of the mature polypeptide coding sequence of SEQ ID NO: 1 for  
5 inhibiting expression of a polypeptide having alpha-mannosidase activity in a cell. While the present invention is not limited by any particular mechanism of action, the dsRNA can enter a cell and cause the degradation of a single-stranded RNA (ssRNA) of similar or identical sequences, including endogenous mRNAs. When a cell is exposed to dsRNA, mRNA from the homologous gene is selectively degraded by a process called RNA interference (RNAi).

10 The dsRNAs of the present invention can be used in gene-silencing. In one aspect, the invention provides methods to selectively degrade RNA using the dsRNAs of the present invention. The process may be practiced *in vitro*, *ex vivo* or *in vivo*. In one aspect, the dsRNA molecules can be used to generate a loss-of-function mutation in a cell, an organ or an animal. Methods for making and using dsRNA molecules to selectively degrade  
15 RNA are well known in the art; see, for example, U.S. Patent No. 6,506,559; U.S. Patent No. 6,511,824; U.S. Patent No. 6,515,109; and U.S. Patent No. 6,489,127.

### Compositions

20 The present invention also relates to compositions comprising a polypeptide of the present invention. Preferably, the compositions are enriched in such a polypeptide. The term "enriched" indicates that the alpha-mannosidase activity of the composition has been increased, *e.g.*, with an enrichment factor of at least 1.1.

The composition may comprise a polypeptide of the present invention as the major enzymatic component, *e.g.*, a mono-component composition. Alternatively, the composition  
25 may comprise multiple enzymatic activities, such as an aminopeptidase, amylase, carbohydrase, carboxypeptidase, catalase, cellulase, chitinase, cutinase, cyclodextrin glycosyltransferase, deoxyribonuclease, esterase, alpha-galactosidase, beta-galactosidase, glucoamylase, alpha-glucosidase, beta-glucosidase, haloperoxidase, invertase, laccase, lipase, mannosidase, oxidase, pectinolytic enzyme, peptidoglutaminase, peroxidase,  
30 phytase, polyphenoloxidase, proteolytic enzyme, ribonuclease, transglutaminase, or xylanase. The additional enzyme(s) may be produced, for example, by a microorganism belonging to the genus *Aspergillus*, preferably *Aspergillus aculeatus*, *Aspergillus awamori*, *Aspergillus fumigatus*, *Aspergillus foetidus*, *Aspergillus japonicus*, *Aspergillus nidulans*, *Aspergillus niger*, or *Aspergillus oryzae*; *Fusarium*, preferably *Fusarium bac-tridioides*,  
35 *Fusarium cerealis*, *Fusarium crookwellense*, *Fusarium culmorum*, *Fusarium graminearum*, *Fusarium graminum*, *Fusarium heterosporum*, *Fusarium negundi*, *Fusarium oxysporum*,

*Fusarium reticulatum*, *Fusarium roseum*, *Fusarium sambucinum*, *Fusarium sarcochroum*, *Fusarium sulphureum*, *Fusarium toruloseum*, *Fusarium trichothecioides*, or *Fusarium venenatum*; *Humicola*, preferably *Humicola insolens* or *Humicola lanuginosa*; or *Trichoderma*, preferably *Trichoderma harzianum*, *Trichoderma koningii*, *Trichoderma longibrachiatum*, *Trichoderma reesei*, or *Trichoderma viride*.

The polypeptide compositions may be prepared in accordance with methods known in the art and may be in the form of a liquid or a dry composition. For instance, the polypeptide composition may be in the form of a granulate or a microgranulate. The polypeptide to be included in the composition may be stabilized in accordance with methods known in the art.

Examples are given below of preferred uses of the polypeptide compositions of the invention. The dosage of the polypeptide composition of the invention and other conditions under which the composition is used may be determined on the basis of methods known in the art.

### Uses

The present invention also relates to methods for producing a demannosylated protein, comprising: treating a protein comprising O-linked mannose residues with a polypeptide having alpha-mannosidase activity, wherein the O-linked mannose residues on the protein are removed. The methods of the present invention are particularly useful for producing a therapeutic polypeptide free of O-linked mannose residues. See, for example, WO 2005/033325.

The present invention also relates to use of a therapeutic recombinant polypeptide expressed in a fungal host for the preparation of a medicament, wherein the therapeutic polypeptide is free of O-linked mannose residues by treatment with a polypeptide having alpha-mannosidase activity of the present invention.

The present invention also relates to use of a polypeptide having alpha-mannosidase activity for complete demannosylation of a therapeutic recombinant polypeptide expressed in a fungal host.

The present invention is further described by the following examples that should not be construed as limiting the scope of the invention.

## Examples

### Materials

Chemicals used as buffers and substrates were commercial products of at least reagent grade.

### Strains

*Thielavia terrestris* NRRL 8126 was used as the source of a gene encoding a polypeptide with identity to an alpha-mannosidase.

### Media

NNCYPmod medium was composed of 1.0 g of NaCl, 5.0 g of  $\text{NH}_4\text{NO}_3$ , 0.2 g of  $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ , 0.2 g of  $\text{CaCl}_2$ , 2.0 g of citric acid, 1.0 g of Bacto Peptone, 5.0 g of yeast extract, 1 ml of COVE trace metals solution, sufficient  $\text{K}_2\text{HPO}_4$  to achieve the final pH of approximately 5.4, and deionized water to 1 liter.

COVE trace metals solution was composed of 0.04 g of  $\text{Na}_2\text{B}_4\text{O}_7 \cdot 10\text{H}_2\text{O}$ , 0.4 g of  $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ , 1.2 g of  $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ , 0.7 g of  $\text{MnSO}_4 \cdot \text{H}_2\text{O}$ , 0.8 g of  $\text{Na}_2\text{MoO}_2 \cdot 2\text{H}_2\text{O}$ , and 10 g of  $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$ , and deionized water to 1 liter.

LB plates were composed of 10 g of tryptone, 5 g of yeast extract, 5 g of sodium chloride, 15 g of Bacto Agar, and deionized water to 1 liter.

SOC medium was composed of 2% tryptone, 0.5% yeast extract, 10 mM NaCl, 2.5 mM KCl, 10 mM  $\text{MgCl}_2$ , 10 mM  $\text{MgSO}_4$ , and deionized water to 1 liter; sterilized by autoclaving and then filter-sterilized glucose was added to 20 mM.

Freezing medium was composed of 60% SOC and 40% glycerol.

### Example 1: Expressed sequence tags (EST) cDNA library construction

*Thielavia terrestris* NRRL 8126 was cultivated in 50 ml of NNCYPmod medium supplemented with 1% glucose in a 250 ml flask at 45°C for 24 hours with shaking at 200 rpm. A two ml aliquot from the 24-hour liquid culture was used to seed a 500 ml flask containing 100 ml of NNCYPmod medium supplemented with 2% SIGMACELL® 20 (cellulose; Sigma Chemical Co., Inc., St. Louis, MO, USA). The culture was incubated at 45°C for 3 days with shaking at 200 rpm. The mycelia were harvested by filtration through a funnel with a glass fiber prefilter (Nalgene, Rochester, NY, USA), washed twice with 10 mM Tris-HCl-1 mM EDTA pH 8 (TE), and quick frozen in liquid nitrogen.

Total RNA was isolated using the following method. Frozen mycelia of *Thielavia terrestris* NRRL 8126 were ground in an electric coffee grinder. The ground material was



mixed 1:1 v/v with 20 ml of FENAZOL™ (Ambion, Inc., Austin, TX, USA) in a 50 ml tube. Once the mycelia were suspended, they were extracted with chloroform and three times with a mixture of phenol-chloroform-isoamyl alcohol 25:24:1 v/v/v. From the resulting aqueous phase, the RNA was precipitated by adding 1/10 volume of 3 M sodium acetate pH 5.2 and 1.25 volumes of isopropanol. The precipitated RNA was recovered by centrifugation at 12,000 x g for 30 minutes at 4°C. The final pellet was washed with cold 70% ethanol, air dried, and resuspended in 500 µl of diethylpyrocarbonate treated water (DEPC-water).

The quality and quantity of the purified RNA was assessed with an AGILENT® 2100 Bioanalyzer (Agilent Technologies, Inc., Palo Alto, CA, USA). Polyadenylated mRNA was isolated from 360 µg of total RNA with the aid of a POLY(A)PURIST™ Magnetic Kit (Ambion, Inc., Austin, TX, USA) according to the manufacturer's instructions.

To create a cDNA library, a CLONEMINER™ Kit (Invitrogen Corp., Carlsbad, CA, USA) was employed to construct a directional library that does not require the use of restriction enzyme cloning, thereby reducing the number of chimeric clones and size bias.

To insure the successful synthesis of the cDNA, two reactions were performed in parallel with two different concentrations of mRNA (2.2 and 4.4 µg of poly (A)<sup>+</sup> mRNA). The mRNA samples were mixed with a Biotin-attB2-Oligo(dt) primer (Invitrogen Corp., Carlsbad, CA, USA), 1X first strand buffer (Invitrogen Corp., Carlsbad, CA, USA), 2 µl of 0.1 M dithiothreitol (DTT), 10 mM of each dNTP, and water to a final volume of 18 and 16 µl, respectively.

The reaction mixtures were mixed and then 2 and 4 µl of SUPERSCRIPT™ reverse transcriptase (Invitrogen Corp., Carlsbad, CA, USA), respectively, were added. The reaction mixtures were incubated at 45°C for 60 minutes to synthesize the first complementary strand. For second strand synthesis, to each first strand reaction was added 30 µl of 5X second strand buffer (Invitrogen Corp., Carlsbad, CA, USA), 3 µl of 10 mM of each dNTP, 10 units of *E. coli* DNA ligase (Invitrogen Corp., Carlsbad, CA, USA), 40 units of *E. coli* DNA polymerase I (Invitrogen Corp., Carlsbad, CA, USA), and 2 units of *E. coli* RNase H (Invitrogen Corp., Carlsbad, CA, USA) in a total volume of 150 µl. The mixtures were then incubated at 16°C for two hours. After the two-hour incubation 2 µl of T4 DNA polymerase (Invitrogen Corp., Carlsbad, CA, USA) were added to each reaction and incubated at 16°C for 5 minutes to create a blunt-ended cDNA. The cDNA reactions were extracted with a mixture of phenol-chloroform-isoamyl alcohol 25:24:1 v/v/v and precipitated in the presence of 20 µg of glycogen, 120 µl of 5 M ammonium acetate, and 660 µl of ethanol. After centrifugation at 12,000 x g for 30 minutes at 4°C, the cDNA pellets were washed with cold 70% ethanol, dried under vacuum for 2-3 minutes, and resuspended in 18 µl of DEPC-water. To each resuspended cDNA sample were added 10 µl of 5X adapted

buffer (Invitrogen, Carlsbad, CA), 10 µg of each *attB1* adapter (Invitrogen, Carlsbad, CA, USA), 7 µl of 0.1 M DTT, and 5 units of T4 DNA ligase (Invitrogen, Carlsbad, CA, USA).

5 Ligation reactions were incubated overnight at 16°C. Excess adapters were removed by size-exclusion chromatography in 1 ml of SEPHACRYL™ S-500 HR resin (Amersham Biosciences, Piscataway, NJ, USA). Column fractions were collected according to the CLONEMINER™ Kit's instructions and fractions 3 to 14 were analyzed with an AGILENT® 2100 Bioanalyzer to determine the fraction at which the *attB1* adapters started to elute. This analysis showed that the adapters started eluting around fraction 10 or 11. For the first library fractions 6 to 11 were pooled and for the second library fractions 4-11 were pooled.

10 Cloning of the cDNA was performed by homologous DNA recombination according to the GATEWAY® System protocol (Invitrogen Corp., Carlsbad, CA, USA) using BP CLONASE™ (Invitrogen Corp., Carlsbad, CA, USA) as the recombinase. Each BP CLONASE™ recombination reaction contained approximately 70 ng of *attB*-flanked-cDNA, 15 250 ng of pDONR™222, 2 µl of 5X BP CLONASE™ buffer, 2 µl of TE, and 3 µl of BP CLONASE™. All reagents were obtained from Invitrogen, Carlsbad, CA, USA. Recombination reactions were incubated at 25°C overnight.

Heat-inactivated BP recombination reactions were then divided into 6 aliquots and electroporated into ELECTROMAX™ DH10B electrocompetent cells (Invitrogen Corp., 20 Carlsbad, CA, USA) using a GENE PULSER™ (Bio-Rad Laboratories, Inc., Hercules, CA, USA) with the following parameters: Voltage: 2.0 kV; Resistance: 200 Ω; and Capacity: 25 µF. Electroporated cells were resuspended in 1 ml of SOC medium and incubated at 37°C for 60 minutes with constant shaking at 200 rpm. After the incubation period, the transformed cells were pooled and mixed 1:1 with freezing medium. A 200 µl aliquot was removed from each library for library titration and then the rest of each library was aliquoted into 1.8 ml cryovials (Wheaton Science Products, Millville, NJ, USA) and stored frozen at - 25 80°C.

Four serial dilutions of each library were prepared: 1/100, 1/1000, 1/10<sup>4</sup>, and 1/10<sup>5</sup>. From each dilution, 100 µl were plated onto 150 mm LB plates supplemented with 50 µg of 30 kanamycin per ml and incubated at 37°C overnight. The number of colonies on each dilution plate was counted and used to calculate the total number of transformants in each library.

The first library contained approximately 5.4 million independent clones and the second library contained approximately 9 million independent clones.

35

**Example 2: Template preparation and nucleotide sequencing of cDNA clones**

Aliquots from both libraries described in Example 1 were mixed and plated onto 25 x 25 cm LB plates supplemented with 50 µg of kanamycin per ml. Individual colonies were arrayed onto 96-well plates containing 100 µl of LB medium supplemented with 50 µg of kanamycin per ml with the aid of a QPix Robot (Genetix Inc., Boston, MA, USA). Forty-five 96-well plates were obtained for a total of 4320 individual clones. The plates were incubated overnight at 37°C with shaking at 200 rpm. After incubation, 100 µl of sterile 50% glycerol was added to each well. The transformants were replicated with the aid of a 96-pin tool (Boekel, Feasterville, PA, USA) into secondary, deep-dish 96-well microculture plates (Advanced Genetic Technologies Corporation, Gaithersburg, MD, USA) containing 1 ml of MAGNIFICENT BROTH™ (MacConnell Research, San Diego, CA, USA) supplemented with 50 µg of kanamycin per ml in each well. The primary microtiter plates were stored frozen at -80°C. The secondary deep-dish plates were incubated at 37°C overnight with vigorous agitation at 300 rpm on a rotary shaker. To prevent spilling and cross-contamination, and to allow sufficient aeration, each secondary culture plate was covered with a polypropylene pad (Advanced Genetic Technologies Corporation, Gaithersburg, MD, USA) and a plastic microtiter dish cover. Plasmid DNA was prepared with a Robot-Smart 384 (MWG Biotech Inc., High Point, NC, USA) and a MONTAGE™ Plasmid Miniprep Kit (Millipore, Billerica, MA, USA).

Sequencing reactions were performed using BIGDYE® (Applied Biosystems, Inc., Foster City, CA, USA) terminator chemistry (Giesecke *et al.*, 1992, *Journal of Virology Methods* 38: 47-60) and a M13 Forward (-20) sequencing primer:

5'-GTAAAACGACGGCCAG-3' (SEQ ID NO: 3)

The sequencing reactions were performed in a 384-well format with a Robot-Smart 384. Terminator removal was performed with a MULTISCREEN® Seq384 Sequencing Clean-up Kit (Millipore, Billerica, MA, USA). Reactions contained 6 µl of plasmid DNA and 4 µl of sequencing master-mix (Applied Biosystems, Foster City, CA, USA) containing 1 µl of 5X sequencing buffer (Millipore, Billerica, MA, USA), 1 µl of BIGDYE® terminator (Applied Biosystems, Inc., Foster City, CA, USA), 1.6 pmoles of M13 Forward primer, and 1 µl of water. Single-pass DNA sequencing was performed with an ABI PRISM Automated DNA Sequencer Model 3700 (Applied Biosystems, Foster City, CA, USA).

5'-GTAAAACGACGGCCAG-3' (SEQ ID NO: 3)

**Example 3: Analysis of DNA sequence data of cDNA clones**

Base calling, quality value assignment, and vector trimming were performed with the assistance of PHRED/PHRAP software (University of Washington, Seattle, WA, USA).

Clustering analysis of the ESTs was performed with a Transcript Assembler v. 2.6.2. (Paracel, Inc., Pasadena, CA, USA). Analysis of the EST clustering indicated the presence of 395 independent clusters.

Sequence homology analysis of the assembled EST sequences against various databases was performed with the Blastx program (Altschul *et. al.*, 1990, *J. Mol. Biol.* 215:403-410) on a 32-node Linux cluster (Paracel, Inc., Pasadena, CA, USA) using the BLOSUM 62 matrix (Henikoff, 1992, *Proc. Natl. Acad. Sci. USA* 89: 10915-10919).

#### **Example 4: Identification of cDNA clones encoding a *Thielavia terrestris* alpha-mannosidase**

A cDNA clone encoding a *Thielavia terrestris* alpha-mannosidase was initially identified by sequence homology to a characterized mannan endo-1,6-alpha-mannosidase DCW1 from *Saccharomyces cerevisiae* (Kitagaki, *et al.*, 2002, *Mol. Microbiol.* 46: 1011-1022, UniProt accession number P36091).

After this initial identification, the clone, designated Tter 08D4, was retrieved from the original frozen stock plate and streaked onto a LB plate supplemented with 50 µg of kanamycin per ml. The plate was incubated overnight at 37°C and the next day a single colony from the plate was used to inoculate 3 ml of LB medium supplemented with 150 µg of kanamycin per ml. The liquid culture was incubated overnight at 37°C and plasmid DNA was prepared with a BIOROBOT® 9600 (QIAGEN, Inc., Valencia, CA, USA). Using a primer walking strategy, the inserted cDNA in the Tter 08D4 plasmid was completely sequenced.

Analysis of the deduced protein sequence of Tter 08D4 with the Interproscan program (Zdobnov and Apweiler, 2001, *Bioinformatics* 17: 847-8.) showed that the gene encoded by Tter08D4 contained the glycosyl hydrolase Family 76 sequence signature known as the PF03663. This sequence signature is located at amino acids position 24 through 400 in the deduced peptide sequence (SEQ ID NO: 2).

The cDNA sequence (SEQ ID NO: 1) and deduced amino acid sequence (SEQ ID NO: 2) are shown in Figure 1. The cDNA clone encodes a polypeptide of 425 amino acids with a molecular mass of 45.9 kDa. The %G+C content of the coding sequence of the gene is 66.4%. Using the SignalP software program (Nielsen *et al.*, 1997, *Protein Engineering* 10: 1-6), a signal peptide was not predicted.

A comparative pairwise global alignment of amino acid sequences was determined using the Needleman-Wunsch algorithm (Needleman and Wunsch, 1970, *J. Mol. Biol.* 48: 443-453) as implemented in the Needle program of EMBOSS with gap open penalty of 10, gap extension penalty of 0.5, and the EBLOSUM62 matrix. The alignment showed that the

deduced amino acid sequence of the *Thielavia terrestris*  $\alpha$ -mannosidase gene shared 46% identity to a characterized mannan endo-1,6- $\alpha$ -mannosidase DCW1 from *Saccharomyces cerevisiae* (Kitagaki, et al., 2002, *Mol. Microbiol.* 46: 1011-1022; UniProt accession number P36091).

5           Once the identity of Tter08D4 was confirmed, a 0.5  $\mu$ l aliquot of plasmid DNA from this clone (pTte08D4, Figure 2) was transferred into a vial of *E. coli* TOP10 cells (Invitrogen Corp., Carlsbad, CA, USA), gently mixed, and incubated on ice for 10 minutes. The cells were then heat-shocked at 42°C for 30 seconds and incubated again on ice for 2 minutes. The cells were resuspended in 250  $\mu$ l of SOC medium and incubated at 37°C for 60 minutes  
10 with constant shaking (200 rpm). After the incubation period, two 30  $\mu$ l aliquots were plated onto LB plates supplemented with 50  $\mu$ g of kanamycin per ml and incubated overnight at 37°C. The next day a single colony was picked and streaked onto three 1.8 ml cryovials containing about 1.5 mls of LB agarose supplemented with 50  $\mu$ g of kanamycin per ml. The vials were sealed with PETRISEAL™ (Diversified Biotech, Boston MA, USA) and deposited  
15 with the Agricultural Research Service Patent Culture Collection, Northern Regional Research Center, Peoria, IL, USA, as NRRL B-50205, with a deposit date of December 11, 2008.

#### Deposit of Biological Material

20

The following biological material has been deposited under the terms of the Budapest Treaty with the Agricultural Research Service Patent Culture Collection (NRRL), Northern Regional Research Center, 1815 University Street, Peoria, IL, USA, and given the following accession number:

25   Deposit	Accession Number	Date of Deposit
<i>E. coli</i> pTter08D4	NRRL B-50205	December 11, 2008

30           The strain has been deposited under conditions that assure that access to the culture will be available during the pendency of this patent application to one determined by foreign patent laws to be entitled thereto. The deposit represents a substantially pure culture of the deposited strain. The deposit is available as required by foreign patent laws in countries wherein counterparts of the subject application, or its progeny are filed. However, it should be understood that the availability of a deposit does not constitute a license to practice the subject invention in derogation of patent rights granted by governmental action.

35           The present invention is further described by the following paragraphs:

[1] An isolated polypeptide having alpha-mannosidase activity, selected from the

group consisting of: (a) a polypeptide comprising an amino acid sequence having at least 60% identity to SEQ ID NO: 2; (b) a polypeptide encoded by a polynucleotide that hybridizes under at least medium stringency conditions with (i) SEQ ID NO: 1, (ii) the genomic DNA sequence comprising SEQ ID NO: 1, or (iii) a full-length complementary strand of (i) or (ii);  
5 (c) a polypeptide encoded by a polynucleotide comprising a nucleotide sequence having at least 60% identity to SEQ ID NO: 1; and (d) a variant comprising a substitution, deletion, and/or insertion of one or more (several) amino acids of SEQ ID NO: 2.

[2] The polypeptide of paragraph 1, comprising an amino acid sequence having at least 60% identity to SEQ ID NO: 2.

10 [3] The polypeptide of paragraph 2, comprising an amino acid sequence having at least 65% identity to SEQ ID NO: 2.

[4] The polypeptide of paragraph 3, comprising an amino acid sequence having at least 70% identity to SEQ ID NO: 2.

15 [5] The polypeptide of paragraph 4, comprising an amino acid sequence having at least 75% identity to SEQ ID NO: 2.

[6] The polypeptide of paragraph 5, comprising an amino acid sequence having at least 80% identity to SEQ ID NO: 2.

[7] The polypeptide of paragraph 6, comprising an amino acid sequence having at least 85% identity to SEQ ID NO: 2.

20 [8] The polypeptide of paragraph 7, comprising an amino acid sequence having at least 90% identity to SEQ ID NO: 2.

[9] The polypeptide of paragraph 8, comprising an amino acid sequence having at least 95% identity to SEQ ID NO: 2.

25 [10] The polypeptide of paragraph 9, comprising an amino acid sequence having at least 97% identity to SEQ ID NO: 2.

[11] The polypeptide of paragraph 1, comprising or consisting of the amino acid sequence of SEQ ID NO: 2; or a fragment thereof having alpha-mannosidase activity.

[12] The polypeptide of paragraph 11, comprising or consisting of the amino acid sequence of SEQ ID NO: 2.

30 [13] The polypeptide of paragraph 1, which is encoded by a polynucleotide that hybridizes under at least medium stringency conditions with (i) SEQ ID NO: 1, (ii) the genomic DNA sequence comprising SEQ ID NO: 1, or (iii) a full-length complementary strand of (i) or (ii).

35 [14] The polypeptide of paragraph 13, which is encoded by a polynucleotide that hybridizes under at least medium-high stringency conditions with (i) SEQ ID NO: 1, (ii) the genomic DNA sequence comprising SEQ ID NO: 1, or (iii) a full-length complementary

strand of (i) or (ii).

[15] The polypeptide of paragraph 14, which is encoded by a polynucleotide that hybridizes under at least high stringency conditions with (i) SEQ ID NO: 1, (ii) the genomic DNA sequence comprising SEQ ID NO: 1, or (iii) a full-length complementary strand of (i) or  
5 (ii).

[16] The polypeptide of paragraph 15, which is encoded by a polynucleotide that hybridizes under at least very high stringency conditions with (i) SEQ ID NO: 1, (ii) the genomic DNA sequence comprising SEQ ID NO: 1, or (iii) a full-length complementary strand of (i) or (ii).

10 [17] The polypeptide of paragraph 1, which is encoded by a polynucleotide comprising a nucleotide sequence having at least 60% identity to SEQ ID NO: 1.

[18] The polypeptide of paragraph 17, which is encoded by a polynucleotide comprising a nucleotide sequence having at least 65% identity to SEQ ID NO: 1.

15 [19] The polypeptide of paragraph 18, which is encoded by a polynucleotide comprising a nucleotide sequence having at least 70% identity to SEQ ID NO: 1.

[20] The polypeptide of paragraph 19, which is encoded by a polynucleotide comprising a nucleotide sequence having at least 75% identity to SEQ ID NO: 1.

[21] The polypeptide of paragraph 20, which is encoded by a polynucleotide comprising a nucleotide sequence having at least 80% identity to SEQ ID NO: 1.

20 [22] The polypeptide of paragraph 21, which is encoded by a polynucleotide comprising a nucleotide sequence having at least 85% identity to SEQ ID NO: 1.

[23] The polypeptide of paragraph 22, which is encoded by a polynucleotide comprising a nucleotide sequence having at least 90% identity to SEQ ID NO: 1.

25 [24] The polypeptide of paragraph 23, which is encoded by a polynucleotide comprising a nucleotide sequence having at least 95% identity to SEQ ID NO: 1.

[25] The polypeptide of paragraph 24, which is encoded by a polynucleotide comprising a nucleotide sequence having at least 97% identity to SEQ ID NO: 1.

30 [26] The polypeptide of paragraph 1, which is encoded by a polynucleotide comprising or consisting of the nucleotide sequence of SEQ ID NO: 1; or a subsequence thereof encoding a fragment having alpha-mannosidase activity.

[27] The polypeptide of paragraph 26, which is encoded by a polynucleotide comprising or consisting of the nucleotide sequence of SEQ ID NO: 1.

[28] The polypeptide of paragraph 1, wherein the polypeptide is a variant comprising a substitution, deletion, and/or insertion of one or more (several) amino acids of SEQ ID NO:  
35 2.

[29] The polypeptide of paragraph 1, which is encoded by the polynucleotide

contained in plasmid pTter08D4 which is contained in *E. coli* NRRL B-50205.

[30] An isolated polynucleotide comprising a nucleotide sequence that encodes the polypeptide of any of paragraphs 1-29.

5 [31] The isolated polynucleotide of paragraph 30, comprising at least one mutation in SEQ ID NO: 1, in which the mutant nucleotide sequence encodes SEQ ID NO: 2.

[32] A nucleic acid construct comprising the polynucleotide of paragraph 30 or 31 operably linked to one or more (several) control sequences that direct the production of the polypeptide in an expression host.

10 [33] A recombinant expression vector comprising the nucleic acid construct of paragraph 32.

[34] A recombinant host cell comprising the nucleic acid construct of paragraph 32.

[35] A method of producing the polypeptide of any of paragraphs 1-29, comprising: (a) cultivating a cell, which in its wild-type form produces the polypeptide, under conditions conducive for production of the polypeptide; and (b) recovering the polypeptide.

15 [36] A method of producing the polypeptide of any of paragraphs 1-29, comprising: (a) cultivating a host cell comprising a nucleic acid construct comprising a nucleotide sequence encoding the polypeptide under conditions conducive for production of the polypeptide; and (b) recovering the polypeptide.

20 [37] A method of producing a mutant of a parent cell, comprising disrupting or deleting a polynucleotide encoding the polypeptide, or a portion thereof, of any of paragraphs 1-29, which results in the mutant producing less of the polypeptide than the parent cell.

[38] A mutant cell produced by the method of paragraph 37.

25 [39] The mutant cell of paragraph 38, further comprising a gene encoding a native or heterologous protein.

[40] A method of producing a protein, comprising: (a) cultivating the mutant cell of paragraph 39 under conditions conducive for production of the protein; and (b) recovering the protein.

30 [41] The isolated polynucleotide of paragraph 30 or 31, obtained by (a) hybridizing a population of DNA under at least medium stringency conditions with (i) SEQ ID NO: 1, (ii) the genomic DNA sequence comprising SEQ ID NO: 1, or (iii) a full-length complementary strand of (i) or (ii); and (b) isolating the hybridizing polynucleotide, which encodes a polypeptide having alpha-mannosidase activity.

35 [42] The isolated polynucleotide of paragraph 41, obtained by (a) hybridizing a population of DNA under at least medium-high stringency conditions with (i) SEQ ID NO: 1, (ii) the genomic DNA sequence comprising SEQ ID NO: 1, or (iii) a full-length



complementary strand of (i) or (ii); and (b) isolating the hybridizing polynucleotide, which encodes a polypeptide having alpha-mannosidase activity.

[43] The isolated polynucleotide of paragraph 42, obtained by (a) hybridizing a population of DNA under at least high stringency conditions with (i) SEQ ID NO: 1, (ii) the genomic DNA sequence comprising SEQ ID NO: 1, or (iii) a full-length complementary strand of (i) or (ii); and (b) isolating the hybridizing polynucleotide, which encodes a polypeptide having alpha-mannosidase activity.

[44] The isolated polynucleotide of paragraph 43, obtained by (a) hybridizing a population of DNA under at least very high stringency conditions with (i) SEQ ID NO: 1, (ii) the genomic DNA sequence comprising SEQ ID NO: 1, or (iii) a full-length complementary strand of (i) or (ii); and (b) isolating the hybridizing polynucleotide, which encodes a polypeptide having alpha-mannosidase activity.

[45] A method of producing a polynucleotide comprising a mutant nucleotide sequence encoding a polypeptide having alpha-mannosidase activity, comprising: (a) introducing at least one mutation into SEQ ID NO: 1, wherein the mutant nucleotide sequence encodes a polypeptide comprising or consisting of the polypeptide of SEQ ID NO: 2; and (b) recovering the polynucleotide comprising the mutant nucleotide sequence.

[46] A mutant polynucleotide produced by the method of paragraph 45.

[47] A method of producing a polypeptide, comprising: (a) cultivating a cell comprising the mutant polynucleotide of paragraph 46 encoding the polypeptide under conditions conducive for production of the polypeptide; and (b) recovering the polypeptide.

[48] A method of producing the polypeptide of any of paragraphs 1-29, comprising: (a) cultivating a transgenic plant or a plant cell comprising a polynucleotide encoding the polypeptide under conditions conducive for production of the polypeptide; and (b) recovering the polypeptide.

[49] A transgenic plant, plant part or plant cell transformed with a polynucleotide encoding the polypeptide of any of paragraphs 1-29.

[50] A double-stranded inhibitory RNA (dsRNA) molecule comprising a subsequence of the polynucleotide of paragraph 30 or 31, wherein optionally the dsRNA is a siRNA or a miRNA molecule.

[51] The double-stranded inhibitory RNA (dsRNA) molecule of paragraph 50, which is about 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25 or more duplex nucleotides in length.

[52] A method of inhibiting the expression of a polypeptide having alpha-mannosidase activity in a cell, comprising administering to the cell or expressing in the cell a double-stranded RNA (dsRNA) molecule, wherein the dsRNA comprises a subsequence of the polynucleotide of paragraph 30 or 31.

[53] The method of paragraph 52, wherein the dsRNA is about 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25 or more duplex nucleotides in length.

[54] A composition comprising the polypeptide of any of paragraphs 1-29.

5 [55] A method for producing a demannosylated protein, comprising: treating a protein comprising O-linked mannose residues with the polypeptide of any of paragraphs 1-29, wherein the O-linked mannose residues on the protein are removed.

10 The invention described and claimed herein is not to be limited in scope by the specific aspects herein disclosed, since these aspects are intended as illustrations of several aspects of the invention. Any equivalent aspects are intended to be within the scope of this invention. Indeed, various modifications of the invention in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description. Such modifications are also intended to fall within the scope of the appended claims. In the case of conflict, the present disclosure including definitions will control.

15

## Claims

### What is claimed is:

- 5 1. An isolated polypeptide having alpha-mannosidase activity, selected from the group consisting of:
- (a) a polypeptide comprising an amino acid sequence having at least 60% identity to SEQ ID NO: 2;
  - 10 (b) a polypeptide encoded by a polynucleotide that hybridizes under at least medium stringency conditions with (i) SEQ ID NO: 1, (ii) the genomic DNA sequence comprising SEQ ID NO: 1, or (iii) a full-length complementary strand of (i) or (ii);
  - (c) a polypeptide encoded by a polynucleotide comprising a nucleotide sequence having at least 60% identity to SEQ ID NO: 1; and
  - 15 (d) a variant comprising a substitution, deletion, and/or insertion of one or more (several) amino acids of SEQ ID NO: 2.
2. The polypeptide of claim 1, comprising or consisting of the amino acid sequence of SEQ ID NO: 2; or a fragment thereof having alpha-mannosidase activity.
- 20 3. The polypeptide of claim 1, comprising or consisting of the amino acid sequence of SEQ ID NO: 2.
4. The polypeptide of claim 1, which is encoded by a polynucleotide comprising or consisting of the nucleotide sequence of SEQ ID NO: 1; or a subsequence thereof encoding  
25 a fragment having alpha-mannosidase activity.
5. The polypeptide of claim 1, which is encoded by a polynucleotide comprising or consisting of the nucleotide sequence of SEQ ID NO: 1.
- 30 6. The polypeptide of claim 1, which is encoded by the polynucleotide contained in plasmid pTter08D4 which is contained in *E. coli* NRRL B-50205.
7. An isolated polynucleotide comprising a nucleotide sequence that encodes the polypeptide of any of claims 1-6.
- 35 8. A nucleic acid construct comprising the polynucleotide of claim 7 operably linked to

one or more (several) control sequences that direct the production of the polypeptide in an expression host.

9. A recombinant host cell comprising the nucleic acid construct of claim 8.

5

10. A method of producing the polypeptide of any of claims 1-6, comprising: (a) cultivating a cell, which in its wild-type form produces the polypeptide, under conditions conducive for production of the polypeptide; and (b) recovering the polypeptide.

10 11. A method of producing the polypeptide of any of claims 1-6, comprising: (a) cultivating a host cell comprising a nucleic acid construct comprising a nucleotide sequence encoding the polypeptide under conditions conducive for production of the polypeptide; and (b) recovering the polypeptide.

15 12. A method of producing a mutant of a parent cell, comprising disrupting or deleting a polynucleotide encoding the polypeptide, or a portion thereof, of any of claims 1-6, which results in the mutant producing less of the polypeptide than the parent cell.

13. A mutant cell produced by the method of claim 12.

20

14. The mutant cell of claim 13, further comprising a gene encoding a native or heterologous protein.

15. A method of producing a protein, comprising: (a) cultivating the mutant cell of claim 14 under conditions conducive for production of the protein; and (b) recovering the protein.

25

16. A method of producing the polypeptide of any of claims 1-6, comprising: (a) cultivating a transgenic plant or a plant cell comprising a polynucleotide encoding the polypeptide under conditions conducive for production of the polypeptide; and (b) recovering the polypeptide.

30

17. A transgenic plant, plant part or plant cell transformed with a polynucleotide encoding the polypeptide of any of claims 1-6.

35 18. A double-stranded inhibitory RNA (dsRNA) molecule comprising a subsequence of the polynucleotide of claim 7, wherein optionally the dsRNA is a siRNA or a miRNA

molecule.

19. A method of inhibiting the expression of a polypeptide having alpha-mannosidase activity in a cell, comprising administering to the cell or expressing in the cell a double-  
5 stranded RNA (dsRNA) molecule, wherein the dsRNA comprises a subsequence of the polynucleotide of claim 7.

20. A method for producing a demannosylated protein, comprising: treating a protein comprising O-linked mannose residues with the polypeptide of any of claims 1-6, wherein  
10 the O-linked mannose residues on the protein are removed.

```

1   TGCTCGCTGGCGGCAACCGGATCCAATTGCTCTTCTCGAACATCTCGTAGAGACGAATTA
      M K F V K S L K P L A .
61  GCCACACCACCTCCCAGTACTGCCACCATGAAGTTTGTCAAGTCCCCTCAAACCACTGGC
      · V G T A A I S A T A P K D L N I N D P S .
121 GGTGGGGACAGCCGCCATCTCGGCCACTGCTCCCAAGGACCTGAACATCAACGACCCATC
      · S I K G V A K T I A A G A M S Y Y P G D .
181 GTCGATCAAGGGCGTGGCCAAGACCATCGCCGCCGCGGCCATGTCGTACTACCCCGGCGA
      · A Q K F V D L P Q P Y Y W W E A G A L M .
241 TGCCAGAAGTTCGTGCGACCTCCCTCAGCCGTACTACTGGTGGGAGGCTGGTGCCTGAT
      · G S M L D Y S H Y T G D T T Y D Q V V A .
301 GGGCTCGATGTTGGACTACTCCCCTACACCGGCGACACGACCTACGACCAGGTGGTCCG
      · T A L L A Q V G P N F D F M L P S H F G .
361 CACGGCCCTGCTGGCCAGGTGGCCCAACTTCGACTTCATGCTACCGAGCCACTTCGG
      · Q E G N D D Q S F W G F A V M A A A E R .
421 CCAGGAGGGCAACGACGACCAATCGTTTTGGGGCTTCGCCGTCATGGCGGCCCGGAGAG
      · N F P Q P D E N V P S W L Q L G A N I F .
481 GAACTTCCCGCAGCCGGACGAGAACGTGCCGTCTTGGCTGCAGCTGGGCGCCAACATCTT
      · N S L A S R W N T T A C G G G L L W Q I .
541 CAACTCGCTGGCTTCGAGATGGAACACGACGGCGTGCGGCGGTGGCCTGCTGTGGCAGAT
      · F A S N P N G L D Y K N T V S N G G L F .
601 CTTCGCCTCGAACCCGAACGGGCTGGACTACAAGAACACGGTGAGCAACGGCGGCCCTCTT
      · Q I A A R L A R A T G N N T Y L E W A E .
661 CCAGATCGCGGCGGGCTCGCCCGGCCACCGCAACAACACGTACCTCGAGTGGGCCGA
      · K V W D W T E A V G L I D S D F N V H D .
721 GAAGGTCTGGGACTGGACCGAGGCCGTGGGGCTCATCGACAGCGACTTCAACGTGCACGA
      · G A S S S H N C T D T N P V T F S Y S A .
781 CGGGGCCAGCTCCAGCCACAATTGCACCGATACGAACCCGGTCACTTCTCGTACAGCGC
      · A I Y L Y G S A V L A N H T G D Q K W V .
841 GGCCATCTACCTGTACGGCTCGGCCGTGCTGGCCAACCACACGGGCGACCAGAAGTGGGT
      · Q R T E K L L G A A R S F F G P F D N A .
901 GCAGCGGACCGAGAAGCTGCTCGGCGCCCGCCGCTCCTTCTTCGGGCCCTTCGACAACGC
      · T D I M Y E H A C C E R V G S C N V D M R .
961 GACCGACATCATGTACGAGCAGCGTGCAGCGGGTCGGCAGCTGCAACGTGCACATGCG
      · S F K A Y F S R F V Y A A S L F V P S I .
1021 CAGCTTCAAGGCCTACTTCTCGCGCTTCGTCTACGCCGCCAGCCTCTTCGTGCCCTCGAT
      · Q P V I D E L W H P S A L A A A K A C S .
1081 CCAGCCGGTTCATCGACGAGCTGTGGCACCCGTCGGCGCTGGCCGCCCGCAAGGCCTGCTC
      · G G A S G T D C G Q K W Y V G G Y D G I .
1141 CGGCGGCGCCAGCGGCACCGACTGCGGCCAGAAGTGGTACGTTCGGCGGCTACGACGGCAT
      · T G L G E E M C A L E T I Q G L L V G Q .
1201 CACCGCCCTCGGCGAGGAGATGTGCGCCCTCGAGACCATCCAGGGCCTGCTCGTGGCCA
      · A A P P L K G S D I K V V R E F A G S S .
1261 GGCCGCGCCCGCCGCTTAAGGGCAGCGACATCAAGGTCGTTTCGCGAGTTCGCTGGCTCCTC
      · S A S K R G G R Y D M R R A .
1321 CTCCGCTTCTAAGCGCGGGGGACGGTACGACATGCGCCGGGCTTAGGTGGGCGAGAGAGT
1381 GTGCTGGTAAGAGGAGGGGGGAACTTCACAGTCTCGGGGATTACCTTCTTGTCTTGTATA
1441 TAGTTTTGGGCATTTTTGATCTGCAGACGTGGATCACTGCATTTTGGCATAGCATTGGGC
1501 GGGCTGGCAAAATAGACAGACGTGGACAGCCAAAAAAAAAAAAAAAAAAAAAAAAAAAAA
1561 AAAAAAAAAA
    
```

Fig. 1

2/2




Fig. 2

## INTERNATIONAL SEARCH REPORT

International application No

PCT/US2009/067641

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> INV. C12N9/24 C12P21/00		
According to International Patent Classification (IPC) or to both national classification and IPC		
<b>B. FIELDS SEARCHED</b>		
Minimum documentation searched (classification system followed by classification symbols) C12N C12P		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used) EPO-Internal, Sequence Search, WPI Data		
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE UniProt [Online] 21 March 2006 (2006-03-21), "SubName: Full=Putative uncharacterized protein;" XP002573850 retrieved from EBI accession no. UNIPROT:Q2H2S1 Database accession no. Q2H2S1	1-5,7
Y	compound	20
X	DATABASE UniProt [Online] 20 May 2008 (2008-05-20), "SubName: Full=Predicted CDS Pa_1_22600;" XP002573851 retrieved from EBI accession no. UNIPROT:B2AS76 Database accession no. B2AS76	1-5,7
Y	compound	20
-/--		
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C.		<input checked="" type="checkbox"/> See patent family annex.
* Special categories of cited documents :		
<p>*A* document defining the general state of the art which is not considered to be of particular relevance</p> <p>*E* earlier document but published on or after the international filing date</p> <p>*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>*O* document referring to an oral disclosure, use, exhibition or other means</p> <p>*P* document published prior to the international filing date but later than the priority date claimed</p>		<p>*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>*&amp;* document member of the same patent family</p>
Date of the actual completion of the international search  22 March 2010		Date of mailing of the international search report  08/04/2010
Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016		Authorized officer  Brück, Marianne



## INTERNATIONAL SEARCH REPORT

International application No  
PCT/US2009/067641

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 02/086090 A2 (ELITRA PHARMACEUTICALS INC [US]; JIANG BO [CA]; TISHKOFF DANIEL [US];) 31 October 2002 (2002-10-31) whole document esp. seq id 220	1-5,7-11
Y	WO 2005/033325 A1 (NOVOZYMES AS [DK]; NOVO NORDISK AS [DK]; DIERS IVAN [DK]; FRISNER HENR) 14 April 2005 (2005-04-14) the whole document	20
X	ZHAO FT, LI J, SHI GX, LIU Y, ZHU LP.: "Inhibition of ER alpha-mannosidase expression causes reduction and shortening of microvilli on rat liver epithelial cell WB-F344" ZHONGGUO-YIXUE-KEXUEYUAN-XUEBAO: SHUANGYUEKAN = ACTA ACADEMIAE MEDICINAE SINICAE, vol. 25, no. 1, 1 February 2003 (2003-02-01), pages 52-55, XP009130974 abstract	1-5,7-11
X,P	KHAN JAVED MOHAMMED ET AL: "A multi-species comparative structural bioinformatics analysis of inherited mutations in $\alpha$ -D-Mannosidase reveals strong genotype-phenotype correlation" BMC GENOMICS, BIOMED CENTRAL, LONDON, GB, no. Suppl 3, 3 December 2009 (2009-12-03), page S33, XP021062280 ISSN: 1471-2164 the whole document	1-5,7-11
X,P	VENKATESAN MEENAKSHI ET AL: "Human lysosomal alpha-mannosidases exhibit different inhibition and metal binding properties" PROTEIN SCIENCE, CAMBRIDGE UNIVERSITY PRESS, CAMBRIDGE, GB, vol. 18, no. 11, 1 November 2009 (2009-11-01), pages 2242-2251, XP009130972 ISSN: 0961-8368 the whole document	1-5,7-11

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box II.2

Claims Nos.: 12-19(completely); 1-11(partially)

Present claims 12-19 relate to an extremely large number of possible mutant cells/inhibitory RNAs. Due to the lack of any structural characterization or any example in the description disclosure and support in the sense of Article 6 and 5 PCT is entirely lacking. The non-compliance with the substantive provisions is to such an extent, that the search had to be restricted to claims 1-11 and 20 which are--apart from the under section VIII objected parts--clear and supported (cf. to The PCT Guidelines 9.01, 9.26, 9.39 and 9.40).

The applicant's attention is drawn to the fact that claims relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure. If the application proceeds into the regional phase before the EPO, the applicant is reminded that a search may be carried out during examination before the EPO (see EPO Guideline C-VI, 8.2), should the problems which led to the Article 17(2) declaration be overcome.

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US2009/067641

## Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1.  Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
  
2.  Claims Nos.: 12-19(completely); 1-11(partially)  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:  
see FURTHER INFORMATION sheet PCT/ISA/210
  
3.  Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1.  As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
  
2.  As all searchable claims could be searched without effort justifying an additional fees, this Authority did not invite payment of additional fees.
  
3.  As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
  
4.  No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

### Remark on Protest

- The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2009/067641

Patent document cited in search report	Publication date	Patent family member(s)	Publication date	
WO 02086090	A2	31-10-2002	CA 2445179 A1	31-10-2002
			EP 1390468 A2	25-02-2004
			JP 2005522980 T	04-08-2005
<hr/>				
WO 2005033325	A1	14-04-2005	NONE	
<hr/>				