



(86) Date de dépôt PCT/PCT Filing Date: 2005/03/10
 (87) Date publication PCT/PCT Publication Date: 2005/09/22
 (85) Entrée phase nationale/National Entry: 2006/09/06
 (86) N° demande PCT/PCT Application No.: US 2005/008029
 (87) N° publication PCT/PCT Publication No.: 2005/087007
 (30) Priorité/Priority: 2004/03/10 (US60/552,065)

(51) Cl.Int./Int.Cl. *A01N 57/20* (2006.01),
A01N 37/40 (2006.01), *A01N 25/30* (2006.01),
A01N 39/04 (2006.01), *A01N 61/00* (2006.01)

(71) Demandeur/Applicant:
 MONSANTO TECHNOLOGY LLC, US

(72) Inventeurs/Inventors:
 BECHER, DAVID Z., US;
 AGBAJE, HENRY E., US;
 TRAVERS, JEFFREY N., US;
 BRINKER, RONALD J., US;
 XU, XIAODONG C., US;
 OTTENS, TIMOTHY S., US

(74) Agent: MCFADDEN, FINCHAM

(54) Titre : COMPOSITIONS HERBICIDES CONTENANT DE LA N-PHOSPHONOMETHYLGLYCINE ET UN HERBICIDE
 A BASE D'AUXINE
 (54) Title: HERBICIDAL COMPOSITIONS CONTAINING N-PHOSPHONOMETHYL GLYCINE AND AN AUXIN
 HERBICIDE

(57) **Abrégé/Abstract:**

Herbicidal compositions are provided which cause rapid symptomology while delivering long term control of regrowth of plants. The herbicidal concentrate compositions comprise N-phosphonomethylglycine or a herbicidal derivative thereof, an auxin herbicide or a herbicidal derivative thereof, and at least one surfactant. Also provided is a method for killing or controlling the growth of certain plants by contacting the foliage of the plants with the diluted concentrate composition.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
22 September 2005 (22.09.2005)

PCT

(10) International Publication Number
WO 2005/087007 A1

- (51) International Patent Classification⁷: **A01N 57/20** // (A01N 57/20, 61:00, 39:04, 37:40, 25:30)
- (74) Agents: **HARPER, James, D.** et al.; Senniger, Powers, Leavitt & Roedel, 1 Metropolitan Sq., 16th Floor, St. Louis, Missouri 63102 (US).
- (21) International Application Number:
PCT/US2005/008029
- (81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (22) International Filing Date: 10 March 2005 (10.03.2005)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/552,065 10 March 2004 (10.03.2004) US
- (84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- (71) Applicant (*for all designated States except US*): **MON-SANTO TECHNOLOGY LLC** [US/US]; 800 North Lindbergh Boulevard, St. Louis, Missouri 63167 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): **BECHER, David, Z.** [US/US]; 12829 Mariners Point Court, St. Louis, MO 63141 (US). **AGBAJE, Henry, E.** [US/US]; 11427 Daykin Drive, St. Louis, MO 63146 (US). **TRAVERS, Jeffrey, N.** [US/US]; 14573 Bexhill Ct., Chesterfield, MO 63017 (US). **BRINKER, Ronald, J.** [US/US]; 925 Oakgrove Park Court, Ellisville, MO 63021 (US). **XU, Xiaodong, C.** [CN/US]; 537 Emmanuel Court, Valley Park, MO 63088 (US). **OTTENS, Timothy, S.** [US/US]; P.O. Box 130, Stanton, MO 63079 (US).
- Published:**
- with international search report
 - before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: HERBICIDAL COMPOSITIONS CONTAINING N-PHOSPHONOMETHYL GLYCINE AND AN AUXIN HERBICIDE

(57) Abstract: Herbicidal compositions are provided which cause rapid symptomology while delivering long term control of re-growth of plants. The herbicidal concentrate compositions comprise N-phosphonomethylglycine or a herbicidal derivative thereof, an auxin herbicide or a herbicidal derivative thereof, and at least one surfactant. Also provided is a method for killing or controlling the growth of certain plants by contacting the foliage of the plants with the diluted concentrate composition.


WO 2005/087007 A1

HERBICIDAL COMPOSITIONS CONTAINING N-PHOSPHONOMETHYL GLYCINE AND AN AUXIN HERBICIDE

BACKGROUND


[0001] The present invention relates generally to herbicidal compositions or formulations, and to methods of using such compositions to kill or control the growth and proliferation of unwanted plants. In particular, the present invention relates to herbicidal compositions, as well as their methods of use, which comprise N-phosphonomethylglycine (glyphosate), or a herbicidal derivative thereof, and an auxin herbicide, or a herbicidal derivative thereof, optionally with one or more suitable surfactants. Such compositions cause early visual symptoms of treatment and/or enhanced effectiveness or control when applied to the foliage of plants.

[0002] Glyphosate is well known in the art as an effective post-emergent foliar-applied herbicide. In its acid form, glyphosate has a structure represented by the formula:



and is relatively insoluble in water (1.16% by weight at 25°C). For this reason it is typically formulated as a water-soluble salt.

[0003] Among the water soluble salts of glyphosate is the potassium salt, having a structure represented by the formula:



[0004] in the ionic form predominantly present in aqueous solution at a pH of about 4. Glyphosate potassium salt has a molecular weight of 207. This

salt is disclosed, for example, by Franz in U.S. Patent No. 4,405,531, as one of the "alkali metal" salts of glyphosate useful as herbicides, with potassium being specifically disclosed as one of the alkali metals, along with lithium, sodium, cesium and rubidium. Example C discloses the preparation of the monopotassium salt by reacting the specified amounts of glyphosate acid and potassium carbonate in an aqueous medium.

[0005] Herbicidal compositions comprising the herbicide N-phosphonomethyl-glycine or derivatives thereof ("glyphosate"), are useful for suppressing the growth of, or killing, unwanted plants such as grasses, weeds and the like. Glyphosate typically is applied to the foliage of the target plant. After application the glyphosate is absorbed by the foliar tissue of the plant and translocated throughout the plant. Glyphosate noncompetitively blocks an important biochemical pathway which is common to virtually all plants, but which is absent in animals. Although glyphosate is very effective in killing or controlling the growth of unwanted plants, the uptake (i.e., absorption) of glyphosate by the plant foliar tissue and translocation of glyphosate throughout the plant is relatively slow. Visual symptoms that a plant has been treated with glyphosate may not appear until one week or more after treatment.

[0006] There is a continuing need for herbicidal compositions which exhibit long-term control of unwanted plants and exhibit early visual symptoms of treatment. These compositions would be well suited to applications in cooler temperatures wherein the early visual symptoms may be readily seen while the long-term control would improve as temperatures increase.

[0007] As will be clear from the disclosure that follows, these and other benefits are provided by the present invention.

SUMMARY OF THE INVENTION

[0008] The present invention provides herbicidal compositions comprising glyphosate or a herbicidal derivative thereof, an auxin herbicide or a herbicidal derivative thereof, and at least one surfactant. The present invention also provides methods for killing or controlling the growth of plants by contacting

the foliage of the plants with the diluted concentrate composition.

[0009] One embodiment of the present invention is directed to an aqueous herbicidal concentrate composition comprising glyphosate or a herbicidal derivative thereof, an auxin comprising one or more auxin herbicides selected from the group consisting of 2,4-D, 2,4-DB, dichlorprop, MCPA, MCPB, mecoprop, dicamba, picloram, quinclorac and agriculturally acceptable salts or esters thereof and a surfactant component in solution or stable suspension, emulsion or dispersion, comprising one or more surfactants. The glyphosate (acid equivalent basis) and the auxin herbicide (acid equivalent basis) are present in a weight ratio of at least 32:1 and the composition has a cloud point of at least about 50°C and a crystallization point not higher than about 0°C.

[0010] Another embodiment of the present invention is directed to an aqueous herbicidal concentrate composition comprising glyphosate, predominantly in the form of the potassium salt thereof in a concentration of at least 65 grams acid equivalent per liter, and an auxin herbicide comprising one or more auxin herbicides selected from the group consisting of 2,4-D, 2,4-DB, dichlorprop, MCPA, MCPB, mecoprop, dicamba, picloram, quinclorac and agriculturally acceptable salts or esters thereof. The herbicidal concentrate composition further comprises a first surfactant component in solution or stable suspension, emulsion or dispersion comprising one or more surfactants selected from the group consisting of secondary or tertiary amines, dialkoxylated quaternary ammonium salts, monoalkoxylated quaternary ammonium salts, quaternary ammonium salts, ether amines, amine oxides, dialkoxylated amines, aminated alkoxyated alcohols, alkyl alkoxyated phosphates and alkylpolyglycosides.

[0011] Yet another embodiment of the present invention is directed to an aqueous herbicidal concentrate composition comprising glyphosate, predominantly in the form of the isopropylammonium salt thereof in a concentration of greater than 360 grams acid equivalent per liter, an auxin herbicide component comprising one or more auxin herbicides selected from the group consisting of 2,4-D, 2,4-DB, dichlorprop, MCPA, MCPB, mecoprop,

dicamba, picloram, quinclorac and agriculturally acceptable salts or esters thereof, and a surfactant component in solution or stable suspension, emulsion or dispersion, comprising one or more surfactants. The glyphosate (acid equivalent basis) and the auxin herbicide component (acid equivalent basis) are present in a weight ratio of at least 9.5:1 and the composition has a cloud point of at least about 50°C and a crystallization point not higher than about 0°C.

[0012] Another embodiment of the present invention is directed to a method of killing or controlling weeds or unwanted plants comprising diluting an aqueous herbicidal concentrate composition in an amount of water to form an application mixture and applying a herbicidally effective amount of the application mixture to foliage of the weeds or unwanted plants, wherein the weeds or unwanted plants comprise *Commelina* and the aqueous herbicidal concentrate composition comprises glyphosate or a herbicidal derivative thereof, an auxin herbicide component comprising one or more auxin herbicides selected from the group consisting of 2,4-D, 2,4-DB, dichlorprop, MCPA, MCPB, mecoprop, dicamba, picloram, quinclorac and agriculturally acceptable salts or esters thereof, and a surfactant component in solution or stable suspension, emulsion or dispersion, comprising one or more surfactants.

[0013] Other objects and features of the present invention will be in part apparent and in part pointed out hereinafter.

DETAILED DESCRIPTION

[0014] According to the present invention, herbicidal compositions containing glyphosate or a derivative thereof, an auxin herbicide or a derivative thereof, and a suitable surfactant, are provided that are advantageous for a number of reasons, including early visual symptoms of plant treatment, rapid uptake by the target plant, and control of a broad spectrum of plant species, as well as enhanced, more consistent control of unwanted plants. Although use of reduced application rates is not preferred, in at least some embodiments, lower application rates may be used without a significant loss of effectiveness of plant control.

[0015] Among the various aspects of the present invention is an aqueous herbicidal composition of N-phosphonomethyl glycine (glyphosate), predominantly in the form of the potassium salt thereof, and an auxin herbicide. The word "predominantly" in the above context means that at least about 50%, preferably at least about 55, 60, 65, 70, 75, 80, 85, 90 or about 95%, by weight of the glyphosate, expressed as a.e., is present as the potassium salt. Other salts of glyphosate which can make up the balance of the glyphosate component are agriculturally acceptable salts including the isopropylamine, di-ammonium, ammonium, sodium, monoethanolamine, n-propylamine, methylamine, ethylamine, hexamethylenediamine, dimethylamine or trimethylsulfonium salts. The second salt ion should be chosen so as not to adversely affect the viscosity, cloud point, non-crystallization and other stability properties of the composition.

[0016] Another aspect of the present invention is an aqueous herbicidal composition of N-phosphonomethyl glycine (glyphosate), predominantly in the form of the isopropylamine salt thereof, and an auxin herbicide. Other salts of glyphosate which can make up the balance of the glyphosate component are agriculturally acceptable salts including the di-ammonium, ammonium, sodium, potassium, monoethanolamine, n-propylamine, methylamine, ethylamine, hexamethylenediamine, dimethylamine or trimethylsulfonium salts.

[0017] The auxin herbicide is selected from the group consisting of 2,4-dichlorophenoxyacetic acid (2,4-D), 4-(2,4-dichlorophenoxy)butanoic acid (2,4-DB), dichloroprop, (4-chloro-2-methylphenoxy)acetic acid (MCPA), 4-(4-chloro-2-methylphenoxy)butanoic acid (MCPB), mecoprop, dicamba, picloram, quinclorac, agriculturally acceptable salts or esters of any of these herbicides, and mixtures thereof. In one embodiment, preferably, the auxin herbicide is selected from the group consisting of 2,4-dichlorophenoxyacetic acid (2,4-D), dicamba, salts or esters thereof, and mixtures thereof. Generally, the primary action of auxin herbicides appears to involve cell wall plasticity and nucleic acid metabolism. 2,4-D is thought to acidify the cell wall by stimulating the activity of a membrane-bound ATPase-driven proton pump. The reduction in apoplasmic

pH induces cell elongation by increasing the activity of certain enzymes responsible for cell wall loosening. Low concentrations of 2,4-D are reported to stimulate RNA polymerase, resulting in subsequent increases in RNA, DNA, and protein biosynthesis. Abnormal increases in these processes presumably lead to uncontrolled cell division and growth, which results in vascular tissue destruction. In contrast, high concentrations of 2,4-D and other auxin-type herbicides inhibit cell division and growth, usually in meristematic regions that accumulate photosynthate assimilates and herbicide from the phloem.

[0018] In another embodiment, preferably, the auxin herbicide is at least 1% soluble by weight in water at pH 6. The auxin herbicide can be present in the composition in the form of its acid, an agriculturally acceptable salt (e.g., isopropylamine, di-ammonium, ammonium, sodium, monoethanolamine, n-propylamine, methylamine, ethylamine, hexamethylenediamine, dimethylamine or trimethylsulfonium), or an agriculturally acceptable ester (e.g., methyl, ethyl, propyl, butyl, octyl, ethoxyethyl, butoxyethyl or methoxypropyl). The salt or ester ion of the auxin herbicide should be chosen to not affect the viscosity, cloud point, non-crystallization and other stability properties of the composition.

[0019] In another aspect of the present invention, the glyphosate and auxin herbicide compositions may contain 5 g a.e./L (grams acid equivalent per liter) to 600 g glyphosate a.e./L, preferably from 65 to about 600, from about 75 to about 600, from about 100 to about 600, from about 150 to about 600, from about 200 to about 600, from about 250 to about 600, from about 300 to about 600, from about 350 to about 600, from about 400 to about 600, from about 450 to about 600, or from about 480 to about 600 g glyphosate a.e./L. In this context, generally, the weight ratio of the glyphosate (acid equivalent basis) to the auxin herbicide (acid equivalent basis) varies depending on the activity of the auxin herbicide which is generally determined using the standard use rates. A person skilled in the art would know that a higher standard use rate indicates a lower activity and thus more of the auxin herbicide should be used to achieve acceptable results. With this relationship in mind, in one embodiment, typically, the weight ratio of glyphosate to 2,4-D, 2,4-DB, MCPA, or MCPB is about 10:1 to

about 100:1. In another embodiment, typically, the weight ratio of glyphosate to mecoprop is about 10:1 to about 50:1. In yet another embodiment, typically, the weight ratio of glyphosate to dicamba, or picloram is about 20:1 to about 200:1. In a further embodiment, preferably, the weight ratio of glyphosate to 2,4-D is about 20:1 to about 100:1; more preferably, about 20:1 to about 50:1; particularly, about 25:1 to about 50:1. In yet a further embodiment, preferably, the weight ratio of glyphosate to dicamba is about 40:1 to about 200:1; more preferably, about 40:1 to about 100:1; particularly, about 50:1 to about 100:1.

[0020] In another embodiment of the invention, the glyphosate in the glyphosate and auxin herbicide compositions is present in an amount of at least about 75, 100, 125, 150, 175, 200, 225, 250, 275, 300, 325, 350, 375, 400, 425, 450, 475, 480, 500, 525, 550, 575, 580 or 600 g a.e./L.

[0021] In another embodiment, the glyphosate (a.e. basis) and auxin herbicide component (a.e. basis) are present in compositions of the invention in a weight ratio of at least 33:1, 34:1, 35:1, 36:1, 37:1, 38:1, 39:1, 40:1, 45:1, 50:1, 55:1, 60:1, 65:1, 70:1, 75:1, 80:1, 85:1, 90:1, 95:1, 100:1, 110:1, 120:1, 130:1, 140:1, 150:1, 160:1, 170:1, 180:1, 190:1 or 200:1. In another embodiment, the glyphosate (a.e. basis) and auxin herbicide component (a.e. basis) are present in a weight ratio of from about 40:1 to about 200:1, from about 50:1 to about 200:1, from about 60:1 to about 200:1, from about 50:1 to about 150:1, from about 50:1 to about 100:1 or from 32:1 to about 50:1.

[0022] In another embodiment, the glyphosate (a.e. basis) and auxin herbicide component (a.e. basis) are present in compositions of the invention in a weight ratio of at least about 5:1, 6:1, 7:1, 8:1, 9:1, 10:1, 15:1, 20:1, 25:1, 30:1, 35:1, 40:1, 45:1, 50:1, 55:1, 60:1, 65:1, 70:1, 75:1, 80:1, 85:1, 90:1, 95:1, 100:1, 110:1, 120:1, 130:1, 140:1, 150:1, 160:1, 170:1, 180:1, 190:1 or 200:1. Preferably, the glyphosate is present in an amount of at least 65 g a.e./L. In one embodiment, the glyphosate concentration is between 360 and 445 g a.e./L, and the glyphosate (a.e. basis) and auxin herbicide component (a.e. basis) are present in a weight ratio of about 5:1 to about 50:1, about 5:1 to about 40:1, or about 8:1 to about 36:1. In a second embodiment, the glyphosate concentration

is between 445 and 480 g a.e./L, and the glyphosate (a.e. basis) and auxin herbicide component (a.e. basis) are present in a weight ratio of about 41:1, 42:1, 43:1, 44:1, 45:1, 46:1, 47:1, 48:1, 49:1, 50:1, 51:1, 52:1, 53:1, 54:1, 55:1, 56:1, 57:1, 58:1, 59:1, 60:1, 61:1, 62:1, 63:1, 64:1, or 65:1. In a third embodiment, the glyphosate concentration is between 360 and 525 g a.e./L, and the glyphosate (a.e. basis) and auxin herbicide component (a.e. basis) are present in a weight ratio of about 8:1 to about 80:1 or about 25:1 to about 56:1. In a fourth embodiment, the glyphosate concentration is at least 480 g a.e./L, and the glyphosate (a.e. basis) and auxin herbicide component (a.e. basis) are present in a weight ratio of about 25:1 to about 80:1, about 50:1 to about 80:1, about 63:1 to about 80:1, or about 25:1 to about 52:1.

[0023] The selection of application rates that are herbicidally effective for a composition of the invention is within the skill of the ordinary agricultural scientist. Those of skill in the art will likewise recognize that individual plant conditions, weather and growing conditions, as well as the specific active ingredients and their weight ratio in the composition, will influence the degree of herbicidal effectiveness achieved in practicing this invention. Typical application rates of the glyphosate and auxin herbicide compositions of the present invention can be determined from the label of each commercially available herbicide for a particular weed species. In general, the application rate of glyphosate is about 340 grams per acre. A person skilled in the art would understand that when the weed growth is heavy or dense or where weeds are growing in an undisturbed area, a higher application rate may be necessary to achieve acceptable weed control. In addition, for difficult-to-control weeds, a higher application rate may be necessary for adequate weed control.

[0024] The potassium glyphosate and auxin herbicide composition of the present invention is useful in controlling a variety of broadleaf weeds. These weeds include Velvetleaf, Redroot Pigweed, Pigweed Species, Tall Waterhemp, Giant Ragweed, Indian Mustard, Sicklepod, Lambsquarters, Wild Poinsettia, Common Mallow, Hemp Sesbania, Prickly Sida, Wild Mustard, Morningglory (Brazil), Morningglory, Ivyleaf Morningglory, Pitted Morningglory, Buckwheat,

Cutleaf Evening Primrose, Curly Dock, Common Chickweed, Common Dayflower and Tropical Spiderwort.

[0025] Also provided by the present invention is a method of killing or controlling weeds or unwanted vegetation comprising diluting with a suitable volume of water a herbicidally effective amount of a composition as provided herein to form an application mixture, and applying the application mixture to foliage of the weeds or unwanted vegetation. If desired, the user can mix one or more adjuvants with a composition of the invention and the water of dilution when preparing the application composition. Such adjuvants can include additional surfactant and/or an inorganic salt such as ammonium sulfate with the aim of further enhancing herbicidal efficacy. However, under most conditions a herbicidal method of use of the present invention gives acceptable efficacy in the absence of such adjuvants.

[0026] In a particular contemplated method of use of a composition of the invention, the composition, following dilution in water, is applied to foliage of crop plants genetically transformed or selected to tolerate glyphosate, and simultaneously to foliage of weeds or undesired plants growing in close proximity to such crop plants. This method of use results in control of the weeds or undesired plants while leaving the crop plants substantially unharmed. Crop plants genetically transformed or selected to tolerate glyphosate include those whose seeds are sold by Monsanto Company or under license from Monsanto Company bearing the Roundup Ready® trademark. These include varieties of wheat, turfgrass, and corn.

[0027] Plant treatment compositions can be prepared simply by diluting a concentrate composition of the invention in water. Application of plant treatment compositions to foliage is preferably accomplished by spraying, using any conventional means for spraying liquids, such as spray nozzles, atomizers or the like. Compositions of the invention can be used in precision farming techniques, in which apparatus is employed to vary the amount of pesticide applied to different parts of a field, depending on variables such as the particular plant species present, soil composition, etc. In one embodiment of such techniques,

a global positioning system operated with the spraying apparatus can be used to apply the desired amount of the composition to different parts of a field.

[0028] A plant treatment composition is preferably dilute enough to be readily sprayed using standard agricultural spray equipment. Useful spray volumes for the present invention can range from about 10 to about 1000 liters per hectare (l/ha) or higher, by spray application.

High load

[0029] In a further embodiment, the glyphosate and auxin herbicide compositions may contain about 300 to about 600 g a.e./L of glyphosate, predominantly in the form of the potassium salt thereof. For these compositions, the balance of the glyphosate component is made up of agriculturally acceptable salts including the isopropylamine, monoethanolamine, n-propylamine, methylamine, ethylamine, ammonium, diammonium, hexamethylenediamine, dimethylamine or trimethylsulfonium salts. In another embodiment, preferably, the glyphosate and auxin herbicide compositions may contain about 450 to about 600 g a.e./L of glyphosate, predominantly in the form of the potassium salt thereof. In general, as the concentration of glyphosate is increased in the composition, the concentration of the auxin herbicide may be decreased to achieve acceptable weed control. Typically, for potassium glyphosate and auxin herbicide compositions containing about 450 to about 600 g a.e./L of glyphosate, the weight ratio of the glyphosate to the auxin herbicide is about 25:1 to about 100:1. In particular, for potassium glyphosate and 2,4-D compositions containing about 540 to about 600 g a.e./L of glyphosate, the weight ratio of the glyphosate to 2,4-D is about 25:1 to about 50:1.

[0030] In another embodiment, the glyphosate and auxin herbicide compositions may contain about 360 to about 600 g a.e./L of glyphosate, predominantly in the form of the isopropylamine salt thereof. For these compositions, the balance of the glyphosate component is made up of agriculturally acceptable salts including the monoethanolamine, n-propylamine, methylamine, ethylamine, ammonium, diammonium, potassium, hexamethylenediamine, dimethylamine or trimethylsulfonium salts. In another


embodiment, preferably, the glyphosate and auxin herbicide compositions may contain about 360 to about 450 g a.e./L of glyphosate, predominantly in the form of the isopropylamine salt thereof. In general, as the concentration of glyphosate is increased in the composition, the concentration of the auxin herbicide may be decreased to achieve acceptable weed control. Typically, for isopropylamine glyphosate and auxin herbicide compositions containing about 360 to about 450 g a.e./L of glyphosate, the weight ratio of the glyphosate to the auxin herbicide is about 10:1 to about 20:1. In one embodiment, the glyphosate is present in an amount of at least about 370, 380, 390, 400, 410, 420, 430, 440, 450, 475, 480, 500, 525, 550, 575, 580 or 600 g a.e./L, and the glyphosate (a.e. basis) and auxin herbicide component (a.e. basis) are present in a weight ratio of at least 9.5:1, 9.6:1, 9.7:1, 9.8:1, 9.9:1, 10:1, 11:1, 12:1, 13:1, 14:1, 15:1, 16:1, 17:1, 18:1, 19:1, or 20:1. In a second embodiment, the glyphosate is present in an amount of from about 400 to about 600, from about 420 to about 600, from about 430 to about 600, from about 440 to about 600, from about 450 to about 600, or from about 480 to about 600 g a.e./L, and the glyphosate (a.e. basis) and auxin herbicide component (a.e. basis) are present in a weight ratio of at least 9.5:1, 9.6:1, 9.7:1, 9.8:1, 9.9:1, 10:1, 11:1, 12:1, 13:1, 14:1, 15:1, 16:1, 17:1, 18:1, 19:1, or 20:1.

Surfactants

[0031] Surfactants and cosurfactants effective in formulating glyphosate, such as potassium or isopropylamine glyphosate, with auxin herbicides include cationic, nonionic, anionic, and amphoteric surfactants and cosurfactants as described below and mixtures thereof, wherein the surfactant component is present in an amount of at least about 5 wt.% based on the total weight of the composition.


[0032] Cationic surfactants and cosurfactants effective in such glyphosate formulations include:

(a) a secondary or tertiary amine having the formula:



wherein R¹ is hydrocarbyl having from 1 to about 30 carbon atoms, and R² and R³ are hydrogen or hydrocarbyl having from 1 to about 30 carbon atoms. In this context, preferred R¹, R², and R³ hydrocarbyl groups are linear or branched alkyl, linear or branched alkenyl, linear or branched alkynyl, aryl, or aralkyl groups. Preferably, R¹ is a linear or branched alkyl or linear or branched alkenyl group having from about 8 to about 30 carbon atoms, and R² and R³ are independently hydrogen or a linear or branched alkyl or linear or branched alkenyl group having from 1 to about 6 carbon atoms. More preferably, R¹ is a linear or branched alkyl or alkenyl group having from about 12 to about 22 carbon atoms, and R² and R³ are independently hydrogen, methyl or ethyl. In one embodiment of the amine of formula (1), R¹ is a linear or branched alkyl group having from about 12 to about 22 carbon atoms, and R² and R³ are independently linear or branched hydroxyalkyl groups having from 1 to about 6 carbon atoms.

(b) dialkoxylated quaternary ammonium salt having the formula:



(2)


wherein R¹¹ is hydrocarbyl or substituted hydrocarbyl having from 1 to about 30 carbon atoms, R¹² in each of the x (R¹²O) and y (R¹²O) groups is independently C₂-C₄ alkylene, R¹³ is hydrogen, or a linear or branched alkyl group having from 1 to about 4 carbon atoms, R¹⁴ hydrocarbyl or substituted hydrocarbyl having from 1 to about 30 carbon atoms, x and y are independently an average number

from 1 to about 40, and X^- is an agriculturally acceptable anion. In this context, preferred R^{11} and R^{14} hydrocarbyl groups are linear or branched alkyl, linear or branched alkenyl, linear or branched alkynyl, aryl, or aralkyl groups. Preferably, R^{11} and R^{14} are independently a linear or branched alkyl or linear or branched alkenyl group having from 1 to about 25 carbon atoms, R^{12} in each of the x ($R^{12}O$) and y ($R^{12}O$) groups is independently C_2 - C_4 alkylene, R^{13} is hydrogen, methyl or ethyl, and the sum of x and y is an average number from about 2 to about 30. More preferably, R^{11} and R^{14} are independently a linear or branched alkyl group having from 1 to about 22 carbon atoms, R^{12} in each of the x ($R^{12}O$) and y ($R^{12}O$) groups is independently ethylene or propylene, R^{13} is hydrogen or methyl, and the sum of x and y is an average number from about 2 to about 20. Even more preferably, R^{11} is a linear or branched alkyl group having from about 8 to about 22 carbon atoms and R^{14} is a linear or branched alkyl group having from 1 to about 22 carbon atoms, R^{12} in each of the x ($R^{12}O$) and y ($R^{12}O$) groups is independently ethylene or propylene, R^{13} is hydrogen or methyl, and x is an average number from about 2 to about 20. Most preferably, R^{11} is a linear or branched alkyl group having from about 8 to about 22 carbon atoms and R^{14} is a linear or branched alkyl group having from 1 to about 6 carbon atoms, R^{12} in each of the x ($R^{12}O$) and y ($R^{12}O$) groups is independently ethylene or propylene, R^{13} is hydrogen or methyl, and x is an average number from about 2 to about 15, or R^{11} and R^{14} are independently a linear or branched alkyl group having from about 8 to about 22 carbon atoms, R^{12} in each of the x ($R^{12}O$) and y ($R^{12}O$) groups is independently ethylene or propylene, R^{13} is hydrogen or methyl, and x is an average number from about 5 to about 15. Preferred dialkoxylated quaternary ammonium surfactants include Ethoquad™ C12 (a PEG 2 coco methyl ammonium chloride from Akzo Nobel), PEG 5 coco methyl ammonium chloride, PEG 5 tallow methyl ammonium chloride, PEG 5 ditallow ammonium bromide, and PEG 10 ditallow ammonium bromide.

(c) monoalkoxylated quaternary ammonium salts having the formula:

about 16 to about 22 carbon atoms, R^{22} in each of the x^2 ($R^{22}O$) groups is independently ethylene or propylene, R^{23} is hydrogen or methyl, R^{24} and R^{25} are independently a linear or branched alkyl group having from 1 to about 3 carbon atoms, and x^2 is an average number from about 5 to about 25. Preferred monoalkoxylated quaternary ammonium surfactants include PEG 7 C_{18} dimethyl ammonium chloride and PEG 22 C_{18} dimethyl ammonium chloride.


(d) quaternary ammonium salts having the formula:



wherein R^{31} , R^{33} and R^{34} are independently hydrogen or hydrocarbyl or substituted hydrocarbyl having from 1 to about 30 carbon atoms, R^{32} is hydrocarbyl or substituted hydrocarbyl having from 1 to about 30 carbon atoms, and X^- is an agriculturally acceptable anion. In this context, preferred R^{31} , R^{32} , R^{33} , and R^{34} hydrocarbyl groups are linear or branched alkyl, linear or branched alkenyl, linear or branched alkynyl, aryl, or aralkyl groups. Preferably, R^{31} is a linear or branched alkyl or linear or branched alkenyl group having from about 8 to about 30 carbon atoms, and R^{32} , R^{33} and R^{34} are independently a linear or branched alkyl or linear or branched alkenyl group having from 1 to about 30 carbon atoms. More preferably, R^{31} is a linear or branched alkyl or linear or branched alkenyl group having from about 8 to about 22 carbon atoms, and R^{32} , R^{33} and R^{34} are independently a linear or branched alkyl or linear or branched alkenyl group having from 1 to about 6 carbon atoms. Even more preferably, R^{31} is a linear or branched alkyl group having from about 8 to about 16 carbon atoms, and R^{32} , R^{33} and R^{34} are independently a linear or branched alkyl group having from 1 to about 6 carbon atoms. Most preferably, R^{31} is a linear or branched alkyl group having from about 8 to about 14 carbon atoms, and R^{32} , R^{33} and R^{34} are methyl. Preferred commercially available quaternary ammonium surfactants include Arquad™ C-50 (a dodecyl trimethyl ammonium chloride from

Akzo Nobel) and Arquad™ T-50 (a tallow trimethyl ammonium chloride from Akzo Nobel).


(e) ether amines having the formula:



wherein R^{41} is hydrocarbyl or substituted hydrocarbyl having from 1 to about 30 carbon atoms; R^{42} is hydrocarbylene or substituted hydrocarbylene having from 2 to about 30 carbon atoms; R^{43} and R^{44} are independently hydrogen, hydrocarbyl or substituted hydrocarbyl having from 1 to about 30 carbon atoms, or $-(\text{R}^{45}\text{O})_x\text{R}^{46}$, R^{45} in each of the $x^4(\text{R}^{45}-\text{O})$ groups is independently $\text{C}_2\text{-C}_4$ alkylene, R^{46} is hydrogen, or a linear or branched alkyl group having from 1 to about 4 carbon atoms, and x^4 is an average number from 1 to about 50. In this context, preferred R^{41} , R^{42} , R^{43} , and R^{44} hydrocarbyl (hydrocarbylene) groups are linear or branched alkyl (alkylene), linear or branched alkenyl (alkenylene), linear or branched alkynyl (alkynylene), aryl (arylene), or aralkyl (aralkylene) groups. Preferably, R^{41} is a linear or branched alkyl, linear or branched alkenyl, linear or branched alkynyl, aryl, or aralkyl group having from 8 to about 25 carbon atoms, R^{42} is a linear or branched alkylene or alkenylene group having from 2 to about 30 carbon atoms, R^{43} and R^{44} are independently hydrogen, a linear or branched alkyl, linear or branched alkenyl, linear or branched alkynyl, aryl, or aralkyl group having from 1 to about 30 carbon atoms, or $-(\text{R}^{45}\text{O})_x\text{R}^{46}$, R^{45} in each of the $x^4(\text{R}^{45}\text{O})$ groups is independently $\text{C}_2\text{-C}_4$ alkylene, R^{46} is hydrogen, methyl or ethyl, and x^4 is an average number from 1 to about 30. More preferably, R^{41} is a linear or branched alkyl or alkenyl group having from 8 to about 22 carbon atoms, R^{42} is a linear or branched alkylene or alkenylene group having from 2 to about 6 carbon atoms, R^{43} and R^{44} are independently hydrogen, a linear or branched alkyl or alkenyl group having from 1 to about 6 carbon atoms, or $-(\text{R}^{45}\text{O})_x\text{R}^{46}$, R^{45} in each of the $x^4(\text{R}^{45}\text{O})$ groups is independently ethylene or propylene, R^{46} is hydrogen or methyl, and x^4 is an average number from 1 to about 15. Most

preferably, R^{41} is a linear or branched alkyl or alkenyl group having from 8 to about 18 carbon atoms, R^{42} is ethylene or propylene, R^3 and R^4 are independently hydrogen, methyl, or $-(R^{45}O)_xR^{46}$, R^{45} in each of the x^4 ($R^{45}O$) groups is independently ethylene or propylene, R^{46} is hydrogen, and x^4 is an average number from 1 to about 5.


(f) amine oxides having the formula:



wherein R^{51} , R^{52} and R^{53} are independently hydrogen, hydrocarbyl or substituted hydrocarbyl, $-(R^{54}O)_xR^{55}$, or $-R^{56}(OR^{54})_xOR^{55}$, R^{54} in each of the x^5 ($R^{54}O$) groups is independently C_2 - C_4 alkylene, R^{55} is hydrogen, or a linear or branched alkyl group having from 1 to about 30 carbon atoms, R^{56} is hydrocarbylene or substituted hydrocarbylene having from 2 to about 6 carbon atoms, x^5 is an average number from 1 to about 50, and the total number of carbon atoms in R^{51} , R^{52} and R^{53} is at least 8. In this context, preferred R^{51} , R^{52} , R^{53} , and R^{56} hydrocarbyl (hydrocarbylene) groups are linear or branched alkyl (alkylene), linear or branched alkenyl (alkenylene), linear or branched alkynyl (alkynylene), aryl (arylene), or aralkyl (aralkylene) groups. Preferably, R^{51} and R^{52} are independently hydrogen, a linear or branched alkyl or linear or branched alkenyl group having from 1 to about 30 carbon atoms, or $-(R^{54}O)_xR^{55}$, R^{53} is a linear or branched alkyl or linear or branched alkenyl group having from about 8 to about 30 carbon atoms, R^{54} in each of the x^5 ($R^{54}O$) groups is independently C_2 - C_4 alkylene; R^{55} is hydrogen, methyl or ethyl, and x^5 is an average number from 1 to about 30. More preferably, R^{51} and R^{52} are independently hydrogen, or a linear or branched alkyl group having from 1 to about 6 carbon atoms, and R^{53} is a linear or branched alkyl group having from about 8 to about 22 carbon atoms; or R^{51} and R^{52} are independently $-(R^{54}O)_xR^{55}$, R^{53} is a linear or branched alkyl group having from about 8 to about 22 carbon atoms, R^{54} in each of the x^5 ($R^{54}O$) groups is ethylene or propylene, R^{55} is hydrogen or methyl, and x^5 is an average number from 1 to about 10. Most preferably, R^{51} and R^{52} are

independently methyl, and R^{53} is a linear or branched alkyl group having from about 8 to about 18 carbon atoms; or R^{51} and R^{52} are independently - $(R^{54}O)_x R^{55}$, R^{53} is a linear or branched alkyl group having from about 8 to about 18 carbon atoms, R^{54} in each of the $x^5 (R^{54}O)$ groups is ethylene or propylene, R^{55} is hydrogen, and x^5 is an average number from 1 to about 5. Commercially available amine oxide surfactants include Chemoxide L70.


(g) dialkoxylated amines having the formula:



wherein R^{61} is a linear or branched alkyl, linear or branched alkenyl, linear or branched alkynyl, aryl, or aralkyl group having from about 6 to about 30 carbon atoms, R^{62} in each of the $x^6 (R^{62}O)$ and the $y^6 (R^{62}O)$ groups is independently C_2 - C_4 alkylene, R^{63} is hydrogen, or a linear or branched alkyl group having from 1 to about 4 carbon atoms and x^6 and y^6 are independently an average number from 1 to about 40. Preferably, R^{61} is a linear or branched alkyl or linear or branched alkenyl group having from about 8 to about 30 carbon atoms, R^{62} in each of the $x^6 (R^{62}O)$ and the $y^6 (R^{62}O)$ groups is independently C_2 - C_4 alkylene, R^{63} is hydrogen, methyl or ethyl, and x^6 and y^6 are independently an average number from 1 to about 20. More preferably, R^{61} is a linear or branched alkyl group having from about 8 to about 25 carbon atoms, R^{62} in each of the $x^6 (R^{62}O)$ and the $y^6 (R^{62}O)$ groups is independently ethylene or propylene, R^{63} is hydrogen or methyl, and x and y are independently an average number from 1 to about 10. Even more preferably, R^{61} is a linear or branched alkyl group having from about 8 to about 22 carbon atoms, R^{62} in each of the $x^6 (R^{62}O)$ and the $y^6 (R^{62}O)$ groups is independently ethylene or propylene, R^{63} is hydrogen or methyl, and x^6 and y^6 are independently an average number from 1 to about 5. Preferred commercially available dialkoxylated amines include Trymeen™ 6617 (from Cognis), Ethomeen™ C/12, C/15, C/20, C/25, T/12, T/15, T/20 and T/25 (from Akzo Nobel), and Genamin™ T-200 DG and T-200 NF (from Clariant).

(h) aminated alkoxyated alcohols having the following chemical

structure:



wherein R^{71} is hydrogen or hydrocarbyl or substituted hydrocarbyl having from 1 to about 30 carbon atoms; R^{72} in each of the x^7 ($R^{72}O$) and y^7 ($R^{72}O$) groups is independently C_2 - C_4 alkylene; R^{73} is hydrocarbylene or substituted hydrocarbylene having from 2 to about 30 carbon atoms; R^{74} and R^{75} are each independently hydrogen, hydrocarbyl or substituted hydrocarbyl having from 1 to about 30 carbon atoms, $-(R^{76})_n^7-(R^{72}O)_y^7R^{77}$, or R^{74} and R^{75} , together with the nitrogen atom to which they are attached, form a cyclic or heterocyclic ring; R^{76} is hydrocarbylene or substituted hydrocarbylene having from 1 to about 30 carbon atoms; R^{77} is hydrogen or a linear or branched alkyl group having 1 to about 4 carbon atoms, n^7 is 0 or 1, x^7 and y^7 are independently an average number from 1 to about 60. In this context, preferred R^{71} , R^{73} , R^{74} , R^{75} , and R^{76} hydrocarbyl (hydrocarbylene) groups are linear or branched alkyl (alkylene), linear or branched alkenyl (alkenylene), linear or branched alkynyl (alkynylene), aryl (arylene), or aralkyl (aralkylene) groups. Preferably, R^{71} is a linear or branched alkyl or linear or branched alkenyl group having from about 8 to about 25 carbon atoms, R^{72} in each of the x^7 ($R^{72}O$) groups is independently C_2 - C_4 alkylene, R^{73} is a linear or branched alkylene group having from 2 to about 20 carbon atoms, R^{74} and R^{75} are each independently hydrogen or a linear or branched alkyl group having from 1 to about 6 carbon atoms, and x^7 is an average number from 1 to about 30. More preferably, R^{71} is a linear or branched alkyl group having from about 12 to about 22 carbon atoms, R^{72} in each of the x^7 ($R^{72}O$) groups is independently ethylene or propylene, R^{73} is a linear or branched alkylene group having from 2 to about 6 carbon atoms, R^{74} and R^{75} are each independently hydrogen, methyl, or tris(hydroxymethyl)methyl, and x^7 is an average number from about 2 to about 30. Even more preferably, R^{71} is a linear or branched alkyl group having from about 12 to about 18 carbon atoms, R^{72} in


C_{1-20} hydrocarbyl or C_{2-6} acyl group, and $R_2, R_3, R_4, R_5, R_6, R_7, R_8, R_9, R_{10}$ groups are independently substituted or unsubstituted C_{1-20} hydrocarbyl or nitrogen containing groups. Generally, in preferred embodiments, n is 0 to 6, a is 1 to about 30, b is 0 to about 10, m is 0 to about 30, q is 0 to about 3, X is hydrogen or a C_{1-6} hydrocarbyl or C_{2-6} acyl group, and $R_2, R_3, R_4, R_5, R_6, R_7, R_8, R_9, R_{10}$ groups are independently substituted or unsubstituted C_{1-4} hydrocarbyl or nitrogen containing groups. In one preferred embodiment, the polysiloxane is a polyoxyethylene heptamethyl trisiloxane wherein R_1 is -

$C_nH_{2n}O(CH_2CH_2O)_m(CH_2CH(CH_3)O)_qX$, n is 3 or 4, a is 1, b is 0, m is 1 to about 30, q is 0, X is hydrogen or a methyl, ethyl or acetyl group, and $R_2, R_3, R_4, R_5, R_6, R_7, R_8, R_9, R_{10}$ groups are independently substituted or unsubstituted C_{1-4} hydrocarbyl or nitrogen containing groups. In another preferred embodiment, a is 1 to 5, b is 0 to 10, n is 3 or 4, m is 1 to about 30, q is 0, X is hydrogen or a methyl, ethyl or acetyl group, and $R_2, R_3, R_4, R_5, R_6, R_7, R_8, R_9,$ and R_{10} are methyl groups. In another preferred embodiment, a is 1 to 5, b is 0 to 10, n is 3 or 4, m is 4 to 12, q is 0, X is hydrogen or a methyl or acetyl group, $R_2, R_3, R_4, R_5, R_6, R_7, R_8, R_9,$ and R_{10} are methyl groups. In a more preferred embodiment, a is 1, b is 0, n is 3 or 4, m is 1 to about 30, b is 0, X is hydrogen or a methyl, ethyl or acetyl group, and $R_2, R_3, R_4, R_5, R_6, R_7, R_8, R_9,$ and R_{10} are methyl groups. In a further preferred embodiment, a is 1, b is 0, n is 3, m is 8, b is 0, X is methyl and $R_2, R_3, R_4, R_5, R_6, R_7, R_8, R_9,$ and R_{10} are methyl groups.


Trisiloxanes of the above formula are generally described in product literature of Crompton Corporation and in U.S. Patent No. 3,505,377. Several of such trisiloxanes are ethoxylated organosilicone wetting agents available from Crompton Corporation as Silwet® silicone glycol copolymers. Both liquid organosilicones and dry organosilicones can be used in the surfactant composition; both are included within the scope of the invention. More preferred trisiloxanes are those sold commercially in the United States or elsewhere by Crompton Corporation as Silwet® L-77, Silwet® 408 and Silwet® 800, by Dow-Corning as Sylgard® 309, by Exacto, Inc., as Qwikwet® 100, and by Goldschmidt as Breakthru S-240. In the most preferred polyoxyethylene heptamethyl trisiloxanes, R^2 is hydrogen.

[0035] Additionally, it has been found that the addition of a C₄ to C₁₆ alkyl or aryl amine compound, or the corresponding quaternary ammonium compound, greatly enhances the compatibility of certain glyphosate salts (e.g., potassium or isopropylamine) with surfactants that otherwise exhibit low or marginal compatibility at a given glyphosate loading. Suitable alkyl or aryl amine compounds may also contain 0 to about 5 EO groups. Preferred alkylamine compounds include C₆ to C₁₂ alkylamines having 0 to 2 EO groups. Similarly, etheramine compounds having 4 to 12 carbons and 0 to about 5 EO groups, as well as the corresponding quaternary ammonium compounds, also enhance the compatibility of such formulations. In one embodiment, the compounds which enhance the compatibility of such surfactants include:


(m) amines or quaternary ammonium salts having the formula:




or



or



or



wherein R^{111} is linear or branched alkyl or aryl having from about 4 to about 16 carbon atoms, R^{112} is hydrogen, methyl, ethyl, or $-(\text{CH}_2\text{CH}_2\text{O})_d\text{H}$, R^{113} is hydrogen, methyl, ethyl, or $-(\text{CH}_2\text{CH}_2\text{O})_e\text{H}$ wherein the sum of d and e is not more than about 5; R^{114} is hydrogen or methyl; R^{116} in each of the c (R^{116}O) groups is independently C_2 - C_4 alkylene; R^{115} is hydrocarbylene or substituted hydrocarbylene having from 2 to about 6 carbon atoms; and A^- is an agriculturally acceptable anion.

[0036] In aqueous concentrated formulations of the present invention, the ratio (by weight) of the glyphosate a.e. to the surfactant is typically in the range of from about 1:1 to about 20:1, preferably from about 2:1 to about 10:1, more preferably from about 2:1 to about 8:1, still more preferably from about 2:1 to about 6:1, still more preferably from about 3:1 to about 6:1, and still more preferably about 4.5:1 to 6:1.

[0037] In another embodiment, preferably, the surfactant of the compositions of the invention comprises a first surfactant component which includes one or more surfactants selected from the group consisting of:

(a) ether amines having formula (5) described above in paragraph (e);
 (b) dialkoxylated amines having formula (7) described above in paragraph (g); and

(c) aminated alkoxylated alcohols having formula (8) described above in paragraph (h). The most preferred surfactants are those which provide a cloud point greater than about 60°C in a composition having a glyphosate loading of at least about 480 g a.e./l.

[0038] In a further embodiment, preferably, the surfactant of the compositions of the invention comprises a first surfactant component as described in detail above and additionally a second surfactant component which includes one or more surfactant(s) selected from the group consisting of:

(a) secondary or tertiary amines having formula (1) described above in paragraph (a);

(b) dialkoxylated quaternary ammonium salts having formula (2) described above in paragraph (b);

(c) monoalkoxylated quaternary ammonium salts having formula (3) described above in paragraph (c);

(d) quaternary ammonium salts having formula (4) described above in paragraph (d);

(e) amine oxides having formula (6) described above in paragraph (e);

(f) alkyl alkoxyated phosphates having formula (9) described above in paragraph (i);

(g) alkyl alkoxyated phosphates having formula (10) described above in paragraph (j);

(h) alkylpolyglycosides having formula (11) described above in paragraph (k); and

(i) amines or quaternary ammonium salts having formulae (12) - (15) described above in paragraph (l).

[0039] In yet another embodiment, more preferably, the second surfactant component is selected from the group consisting of:

(a) alkylpolyglycosides having formula (11) described above in paragraph (k); and

(b) amines or quaternary ammonium salts having formulae (12) - (15) described above in paragraph (l).

[0040] In an embodiment of the invention, the density of the formulation of the invention is preferably at least 1.210 grams/liter, more preferably at least about 1.215, 1.220, 1.225, 1.230, 1.235, 1.240, 1.245, 1.250, 1.255, 1.260, 1.265, 1.270, 1.275, 1.280, 1.285, 1.290, 1.295, 1.300, 1.305, 1.310, 1.315, 1.320, 1.325, 1.330, 1.335, 1.340, 1.345, 1.350, 1.355, 1.360, 1.365, 1.370,

1.375, 1.380, 1.385, 1.390, 1.395, 1.400, 1.405, 1.410, 1.415, 1.420, 1.425, 1.430, 1.435, 1.440, 1.445, or 1.450 grams/liter.

[0041] As further discussed herein, other additives, adjuvants, or ingredients may be introduced into the formulations of the present invention to improve certain properties of the resulting formulations. Although the formulations of the present invention generally show good overall stability and viscosity properties without the addition of any further additives, the addition of a solubilizer (also commonly referred to as a cloud point enhancer or stabilizer) can significantly improve the properties of the formulations of the present invention. Suitable solubilizers for use with the novel formulations of the present invention include, for example, cocoamine (Armeen C), dimethylcocoamine (Arquad DMCD), cocoammonium chloride (Arquad C), PEG 2 cocoamine (Ethomeen C12), PEG 5 tallowamine (Ethomeen T15), and PEG 5 cocoamine (Ethomeen C15), all of which are manufactured by Akzo Nobel (California). Additional excipient ingredients may include conventional formulation additives such as dyes, thickeners, crystallization inhibitors, antifreeze agents (e.g., glycols, such as ethylene glycol, or polyethylene glycols such as polyethylene glycol 200, 400, 600, 1500, 4000 or 6000), foam moderating agents (e.g., Antifoam™ or Y-14088 Antifoam™, both available from Crompton Corporation), antidrift agents, compatibilizing agents, antioxidants (e.g., ascorbic acid and sodium sulfite, in order for example to prevent the formation of a nitrosamine), other co-solvents (e.g., N-methylpyrrolidone, DMSO, DMF, propylene carbonate, or ethylene glycol), or some other agent added to lessen or overcome antagonism associated with hard water (e.g., ammonium sulfate, EDTA or a polymeric water conditioner, such as a polyacrylic acid).

[0042] Other components such as solvents and organic acids may be added to enhance concentrate stability. These additives generally function to increase solubility or dispersability of the surfactants in the aqueous carrier thus enabling the formulation of robust concentrates exhibiting enhanced thermal and pH stability, reduced viscosity, and high glyphosate loading. Non-limiting examples of water soluble solvents include acetates, C₁₋₆ alkanols, C₁₋₆ diols, C₁₋₆ alkyl ethers of alkylene glycols and polyalkylene glycols, and mixtures thereof.

The alkanol can be selected from methanol, ethanol, n-propanol, isopropanol, the various positional isomers of butanol, pentanol, and hexanol, and mixtures thereof. It may also be possible to utilize in addition to, or in place of, said alkanols, the diols such as methylene, ethylene, diethylene, propylene, dipropylene, and butylene glycols, and mixtures thereof, and including polyalkylene glycols. These components are generally employed in dispersion-effective or solubilizing effective amounts.

Suitable organic acids include, among others, acetic, dichloroacetic, citric, malic, oxalic, salicylic and tartaric. Effective concentrations of organic acids are generally between about 0.1 wt% and 5 wt%.

[0043] Although additional herbicides can be included in the compositions of the invention other than the glyphosate and auxin herbicides, it is preferred that the glyphosate and the auxin herbicides are the only herbicides in the composition.

DEFINITIONS

[0044] The terms "hydrocarbon" and "hydrocarbyl" as used herein describe organic compounds or radicals consisting exclusively of the elements carbon and hydrogen. These moieties include alkyl, alkenyl, alkynyl, and aryl moieties. These moieties also include alkyl, alkenyl, alkynyl, and aryl moieties substituted with other aliphatic or cyclic hydrocarbon groups, such as alkaryl, alkenaryl and alkynaryl. Unless otherwise indicated, these moieties preferably comprise 1 to 30 carbon atoms.

[0045] The term "hydrocarbylene" as used herein describes radicals joined at two ends thereof to other radicals in an organic compound, and which consist exclusively of the elements carbon and hydrogen. These moieties include alkylene, alkenylene, alkynylene, and arylene moieties. These moieties also include alkyl, alkenyl, alkynyl, and aryl moieties substituted with other aliphatic or cyclic hydrocarbon groups, such as alkaryl, alkenaryl and alkynaryl. Unless otherwise indicated, these moieties preferably comprise 1 to 30 carbon atoms.

[0046] The “substituted hydrocarbyl” moieties described herein are hydrocarbyl moieties which are substituted with at least one atom other than carbon, including moieties in which a carbon chain atom is substituted with a hetero atom such as nitrogen, oxygen, silicon, phosphorous, boron, sulfur, or a halogen atom. These substituents include halogen, heterocyclo, alkoxy, alkenoxy, alkynoxy, aryloxy, hydroxy, protected hydroxy, ketal, acyl, acyloxy, nitro, amino, amido, cyano, thiol, acetal, sulfoxide, ester, thioester, ether, thioether, hydroxyalkyl, urea, guanidine, amidine, phosphate, amine oxide, and quaternary ammonium salt.

[0047] The “substituted hydrocarbylene” moieties described herein are hydrocarbylene moieties which are substituted with at least one atom other than carbon, including moieties in which a carbon chain atom is substituted with a hetero atom such as nitrogen, oxygen, silicon, phosphorous, boron, sulfur, or a halogen atom. These substituents include halogen, heterocyclo, alkoxy, alkenoxy, alkynoxy, aryloxy, hydroxy, protected hydroxy, ketal, acyl, acyloxy, nitro, amino, amido, cyano, thiol, acetal, sulfoxide, ester, thioester, ether, thioether, hydroxyalkyl, urea, guanidine, amidine, phosphate, amine oxide, and quaternary ammonium salt.

[0048] Unless otherwise indicated, the alkyl groups described herein are preferably lower alkyl containing from one to 18 carbon atoms in the principal chain and up to 30 carbon atoms. They may be straight or branched chain or cyclic and include methyl, ethyl, propyl, isopropyl, n-butyl, isobutyl, hexyl, 2-ethylhexyl, and the like.

[0049] Unless otherwise indicated, the alkenyl groups described herein are preferably lower alkenyl containing from two to 18 carbon atoms in the principal chain and up to 30 carbon atoms. They may be straight or branched chain or cyclic and include ethenyl, propenyl, isopropenyl, butenyl, isobutenyl, hexenyl, and the like.

[0050] Unless otherwise indicated, the alkynyl groups described herein are preferably lower alkynyl containing from two to 18 carbon atoms in the

principal chain and up to 30 carbon atoms. They may be straight or branched chain and include ethynyl, propynyl, butynyl, isobutynyl, hexynyl, and the like.

[0051] The terms "aryl" as used herein alone or as part of another group denote optionally substituted homocyclic aromatic groups, preferably monocyclic or bicyclic groups containing from 6 to 12 carbons in the ring portion, such as phenyl, biphenyl, naphthyl, substituted phenyl, substituted biphenyl or substituted naphthyl. Phenyl and substituted phenyl are the more preferred aryl.

[0052] The term "aralkyl" as used herein denotes a group containing both alkyl and aryl structures such as benzyl.

[0053] As used herein, the alkyl, alkenyl, alkynyl, aryl and aralkyl groups can be substituted with at least one atom other than carbon, including moieties in which a carbon chain atom is substituted with a hetero atom such as nitrogen, oxygen, silicon, phosphorous, boron, sulfur, or a halogen atom. These substituents include hydroxy, nitro, amino, amido, nitro, cyano, sulfoxide, thiol, thioester, thioether, ester and ether, or any other substituent which can increase the compatibility of the surfactant and/or its efficacy enhancement in the potassium glyphosate formulation without adversely affecting the storage stability of the formulation.

[0054] The terms "halogen" or "halo" as used herein alone or as part of another group refer to chlorine, bromine, fluorine, and iodine. Fluorine substituents are often preferred in surfactant compounds.

[0055] Unless otherwise indicated, the term "hydroxyalkyl" includes alkyl groups substituted with at least one hydroxy group, and includes bis(hydroxyalkyl)alkyl, tris(hydroxyalkyl)alkyl and poly(hydroxyalkyl)alkyl groups. Preferred hydroxyalkyl groups include hydroxymethyl (-CH₂OH), and hydroxyethyl (-C₂H₄OH), bis(hydroxymethyl)methyl (-CH(CH₂OH)₂), and tris(hydroxymethyl)methyl (-C(CH₂OH)₃).

[0056] The term "cyclic" as used herein alone or as part of another group denotes a group having at least one closed ring, and includes alicyclic, aromatic (arene) and heterocyclic groups.

[0057] The terms "heterocyclo" or "heterocyclic" as used herein alone or as part of another group denote optionally substituted, fully saturated or unsaturated, monocyclic or bicyclic, aromatic or nonaromatic groups having at least one heteroatom in at least one ring, and preferably 5 or 6 atoms in each ring. The heterocyclo group preferably has 1 or 2 oxygen atoms, 1 or 2 sulfur atoms, and/or 1 to 4 nitrogen atoms in the ring, and may be bonded to the remainder of the molecule through a carbon or heteroatom. Exemplary heterocyclo include heteroaromatics such as furyl, thienyl, pyridyl, oxazolyl, pyrrolyl, indolyl, quinoliny, or isoquinoliny and the like, and non-aromatic heterocyclics such as tetrahydrofuryl, tetrahydrothienyl, piperidiny, pyrrolidino, etc. Exemplary substituents include one or more of the following groups: hydrocarbyl, substituted hydrocarbyl, keto, hydroxy, protected hydroxy, acyl, acyloxy, alkoxy, alkenoxy, alkynoxy, aryloxy, halogen, amido, amino, nitro, cyano, thiol, thioester, thioether, ketal, acetal, ester and ether.

[0058] The term "heteroaromatic" as used herein alone or as part of another group denote optionally substituted aromatic groups having at least one heteroatom in at least one ring, and preferably 5 or 6 atoms in each ring. The heteroaromatic group preferably has 1 or 2 oxygen atoms, 1 or 2 sulfur atoms, and/or 1 to 4 nitrogen atoms in the ring, and may be bonded to the remainder of the molecule through a carbon or heteroatom. Exemplary heteroaromatics include furyl, thienyl, pyridyl, oxazolyl, pyrrolyl, indolyl, quinoliny, or isoquinoliny and the like. Exemplary substituents include one or more of the following groups: hydrocarbyl, substituted hydrocarbyl, keto, hydroxy, protected hydroxy, acyl, acyloxy, alkoxy, alkenoxy, alkynoxy, aryloxy, halogen, amido, amino, nitro, cyano, thiol, thioether, thioester, ketal, acetal, ester and ether.

[0059] The term "acyl," as used herein alone or as part of another group, denotes the moiety formed by removal of the hydroxyl group from the group -COOH of an organic carboxylic acid, e.g., RC(O)-, wherein R is R¹, R¹O-, R¹R²N-, or R¹S-, R¹ is hydrocarbyl, heterosubstituted hydrocarbyl, or heterocyclo and R² is hydrogen, hydrocarbyl or substituted hydrocarbyl.

[0060] The term "acyloxy," as used herein alone or as part of another group, denotes an acyl group as described above bonded through an oxygen linkage

(--O--), e.g., RC(O)O- wherein R is as defined in connection with the term "acyl."

[0061] The term "pesticide" includes chemicals and microbial agents used as active ingredients of products for control of crop and lawn pests and diseases, animal ectoparasites, and other pests in public health. The term also includes plant growth regulators, pest repellants, synergists, herbicide safeners (which reduce the phytotoxicity of herbicides to crop plants) and preservatives, the delivery of which to the target may expose dermal and especially ocular tissue to the pesticide. Such exposure can arise by drift of the pesticide from the delivery means to the person performing the application of the pesticide or being present in the vicinity of an application.

[0062] When a maximum or minimum "average number" is recited herein with reference to a structural feature such as oxyethylene units or glucoside units, it will be understood by those skilled in the art that the integer number of such units in individual molecules in a surfactant preparation typically varies over a range that can include integer numbers greater than the maximum or smaller than the minimum "average number". The presence in a composition of individual surfactant molecules having an integer number of such units outside the stated range in "average number" does not remove the composition from the scope of the present invention, so long as the "average number" is within the stated range and other requirements are met.

[0063] By "storage-stable," in the context of an aqueous concentrate composition of glyphosate salt further containing a surfactant and auxin herbicide, is meant not exhibiting phase separation on exposure to temperatures up to about 50 °C, and preferably not forming crystals of glyphosate or salt thereof on exposure to a temperature of about 0 °C for a period of up to about 7 days (i.e., the composition must have a crystallization point of 0 °C or lower). For aqueous solution concentrates, high temperature storage stability is often indicated by a cloud point of about 50 °C or more. Cloud point of a composition is normally determined by heating the composition until the solution becomes

cloudy, and then allowing the composition to cool, with agitation, while its temperature is continuously monitored. A temperature reading taken when the solution clears is a measure of cloud point. A cloud point of 50 °C or more is normally considered acceptable for most commercial purposes for a glyphosate SL formulation. Ideally the cloud point should be 60 °C or more, and the composition should withstand temperatures as low as about -10 °C, preferably as low as about -20 °C, more preferably as low as about -30°C, for up to about 7 days without phase separation (i.e., without separation of frozen water or solid insoluble surfactant from the composition) and without crystal growth (even in the presence of seed crystals of the glyphosate salt).

[0064] Herbicidal effectiveness is one of the biological effects that can be enhanced through this invention. "Herbicidal effectiveness," as used herein, refers to any observable measure of control of plant growth, which can include one or more of the actions of (1) killing, (2) inhibiting growth, reproduction or proliferation, and (3) removing, destroying, or otherwise diminishing the occurrence and activity of plants. The herbicidal effectiveness data set forth herein report "control" as a percentage following a standard procedure in the art which reflects a visual assessment of plant mortality and growth reduction by comparison with untreated plants, made by technicians specially trained to make and record such observations. In all cases, a single technician makes all assessments of percent control within any one experiment or trial. Such measurements are relied upon and regularly reported by Monsanto Company in the course of its herbicide business.

EXAMPLES

[0065] The spray compositions of the following examples contain an exogenous chemical, such as glyphosate salt as indicated, in addition to the excipient ingredients listed. The amount of exogenous chemical was selected to provide the desired rate in grams per hectare (g/ha) when applied in a spray volume of 93 l/ha. Several exogenous chemical rates were applied for each composition. Thus, except where otherwise indicated, when spray compositions were tested, the concentration of exogenous chemical varied in direct proportion

to exogenous chemical rate, but the concentration of excipient ingredients was held constant across different exogenous chemical rates.

[0066] In the following Examples illustrative of the invention, greenhouse and field tests were conducted to evaluate the relative herbicidal effectiveness of glyphosate compositions. Standard compositions included for comparative purposes include the following:

STD1: 725 g/l of glyphosate potassium salt in aqueous solution with no added surfactant.

STD2: 50% by weight of glyphosate IPA salt in aqueous solution together with a surfactant. This formulation is sold by Monsanto Company under the ROUNDUP ULTRAMAX® trademark.

STD3: 570 g/l of glyphosate IPA salt in an aqueous solution with no added surfactant.

[0067] Various excipients were used in compositions of the examples. They may be identified as follows.

Cationic Surfactants:

CIS1	Witcamine TAM105	Ethoxylated (10.5) tallowamine
CIS2	3151 blend	55% Ethoxylated (10.5) tallowamine and 45% Ethoxylated (2) cocoamine
CIS3	Surfonic T-15	PEG 15 tallow amine
CIS4	Witcamine 302	PEG 2 cocoamine
CIS5	Witcamine 305	PEG 5 cocoamine
CIS6	Formulation E1 of Reissue Patent No. RE 37,866	C ₁₂₋₁₄ alkoxyated (1 PO) propylamine (5EO) ethoxylate
CIS7	Armeen C	coco (C ₁₂ -C ₁₈ unsaturated) primary amine

CIS8	Ethoquad T25	tallow ethoxylate (15EO) quaternary ammonium chloride
CIS9	MON 0818	polyoxyethylene tallowamine
CIS10		C ₁₈ NMe(EO)7.5H
CIS11	7164 blend	54% 4.5EO tallowamine ethoxylate, 23% 10 EO tallowamine ethoxylate, and 23% dipropylene glycol
CIS12	Witcamine TAM 45	4.5 EO tallowamine ethoxylate
CIS13	Arquad T-50PG	tallowtrimethylammonium chloride in propylene glycol
CIS14	Arquad SV-60PG	soyaalkyltrimethyl ammonium chloride
CIS15	Tomah E-17-5	poly(5)oxyethylene isodecyl oxypropylamine

Nonionic Surfactants:

NIS1	Hetoxol CS20	C ₁₆₋₁₈ alcohol ethoxylate (20EO)
NIS2	Agrimul PG 2067	Alkylpolyglucoside (Cognis)
NIS3		C ₁₆₋₁₈ alcohol ethoxylate (20EO)
NIS4	Witconol IS 100	PEG 10EO iso C ₁₈ alcohol
NIS5	Silwet L-77	silicone-polyether copolymer
NIS6	Brij 56	stearyl alcohol ethoxylate (10EO)
NIS7	ADMOX SC1485	myristyl dimethyl amine oxide
NIS8		20 EO linear C ₁₆₋₁₈ alcohol ethoxylate

NIS9	Emulgin L	cetereth propoxylate (2PO) ethoxylate (9EO)
NIS10		alkoxylated alcohol
NIS11		alkoxylated alcohol

Other Components:

OTH1	Di-ammonium Oxalate	
OTH2	Propylene Glycol	
OTH3	Oxalic Acid	
OTH4	Sodium Sulfite	
OTH5	Agnique DF6889	Silicone dispersion antifoam
OTH6	octyl amine	
OTH7	tetrahydrofuryl alcohol	
OTH8	Isopar L	paraffinic oil
OTH9	dipropylene glycol	
OTH10	diethylene glycol	
OTH11	NaCl	
OTH12	KOH	
OTH13	glycerin	
OTH14	phosphoric acid	
OTH15	dimethyl amine	
OTH16	N-decyl amine	
OTH17	diethyl amine	
OTH18	isopropyl alcohol	

[0068] The following greenhouse testing procedure was used for evaluating compositions of the Examples to determine herbicidal effectiveness, except where otherwise indicated.

[0069] Seeds of the plant species indicated were planted in 88 mm square pots in a soil mix which was previously sterilized and prefertilized with a 14-14-14 NPK slow release fertilizer at a rate of 3.6 kg/m³. The pots were

placed in a greenhouse with sub-irrigation. About one week after emergence, seedlings were thinned as needed, including removal of any unhealthy or abnormal plants, to create a uniform series of test pots.

[0070] The plants were maintained for the duration of the test in the greenhouse where they received a minimum of 14 hours of light per day. If natural light was insufficient to achieve the daily requirement, artificial light with an intensity of approximately 475 microeinsteins was used to make up the difference. Exposure temperatures were not precisely controlled but averaged about 29 °C during the day and about 21 °C during the night. Plants were sub-irrigated throughout the test to ensure adequate soil moisture levels.

[0071] Pots were assigned to different treatments in a fully randomized experimental design with 6 replications. A set of pots was left untreated as a reference against which the effects of a treatment could later be evaluated.

[0072] Application of glyphosate compositions was made by spraying with a track sprayer fitted with a 9501E tapered flat fan spray tip calibrated to deliver a spray volume of 93 liters per hectare (l/ha) at a pressure of 165 kilopascals (kPa). After treatment, pots were returned to the greenhouse until ready for evaluation.

[0073] Treatments were made using dilute aqueous compositions. These could be prepared as spray compositions directly from their ingredients, or by dilution with water of preformulated concentrate compositions.

[0074] For evaluation of herbicidal effectiveness, all plants in the test were examined by a single practiced technician, who recorded percent control, a visual measurement of the effectiveness of each treatment by comparison with untreated plants. Control of 0% indicates no effect, and control of 100% indicates that all of the plants are completely dead. The reported % control values represent the average for all replicates of each treatment.

EXAMPLE 1

[0075] The effect of glyphosate, 2,4-D, combinations of 2,4-D and glyphosate and combinations of all of the above with oxalic acid on velvetleaf

was tested. Aqueous concentrate compositions were prepared containing potassium glyphosate salt, reported in g a.e./liter, and excipient ingredients as shown in Table 1a. The 806D0T, 806E7S, 806F4Q and 806G3B formulations contained 62 grams acid equivalent per liter. Formulations 806A2D, 806B9H, 806C5Z, 806F4Q and 806G3B contained the IPA salt of 2,4-D measured in grams acid equivalent per liter.

Table 1a

Comp.	Gly.	2,4-D	Cmpnt. 2	wt%	Cmpnt. 3	wt%	Cmpnt. 4	wt%
806A2D	---	60.0	---	---	---	---	---	---
806B9H	---	60.0	CIS6	0.75	NIS2	1.0	---	---
806C5Z	---	60.0	CIS6	0.75	NIS2	1.0	OTH3	0.30
806D0T	K	---	CIS6	0.75	NIS2	1.0	---	---
806E7S	K	---	CIS6	0.75	NIS2	1.0	OTH3	0.30
806F4Q	K	2.0	CIS6	0.75	NIS2	1.0	---	0.75
806G3B	K	2.0	CIS6	0.75	NIS2	1.0	OTH3	0.30
765K4S	K	---	CIS5	9.0	NIS4	4.0	CIS7	1.0

[0076] The compositions of Table 1a and comparative compositions STD1 and STD2, were applied to velvetleaf (*Abutilon theophrasti*, ABUTH) plants. Results, averaged for all replicates of each treatment, are shown in Table 1b.

Table 1b:

Composition	Glyphosate Application Rate (g a.e./ha)	ABUTH %inhibition
806A2D	100, 200, 300, 400	59.2, 75.8, 77.5, 80.0
806B9H	100, 200, 300, 400	66.7, 75.0, 80.0, 80.0
806C5Z	100, 200, 300, 400	75.0, 78.3, 80.0, 82.5
806D0T	100, 200, 300, 400	24.2, 59.2, 85.0, 88.3

806E7S	100, 200, 300, 400	77.5, 87.5, 96.0, 98.0
806F4Q	100, 200, 300, 400	25.0, 75.0, 80.8, 86.7
806G3B	100, 200, 300, 400	68.3, 89.2, 95.5, 97.3
765K4S	100, 200, 300, 400	16.7, 63.3, 85.0, 90.0
STD1	100, 200, 300, 400	0.0, 1.7, 44.2, 77.5
STD2	100, 200, 300, 400	13.3, 81.7, 90.0, 95.0

[0077] The order of efficacy for ABUTH %inhibition was 806E7S > 806G3B > 806C5Z > 806B9H > 806A2D > STD2 > 806F4Q > 806D0T > 765K4S > STD1.

EXAMPLE 2

[0078] The effect of combinations of potassium glyphosate and 2,4-D with or without oxalic acid on velvetleaf was tested. Aqueous concentrate compositions were prepared containing potassium glyphosate salt, reported in g a.e./liter, and excipient ingredients as shown in Table 2a. The 820A9T, 820C1Z, 820D6Q, 820E3F, 820F0G and 820H7D formulations contained 62 grams acid equivalent per liter. Formulation 820B4H contained the IPA salt of 2,4-D measured in grams acid equivalent per liter.

Table 2a

Comp.	Gly.	2,4-D	Cmpnt. 2	wt%	Cmpnt. 4	wt%	Cmpnt. 4	wt%
820A9T	K	---	CIS6	0.75	NIS2	1.00	---	---
820B4H	---	60.0	CIS6	0.75	NIS2	1.00	---	---
820C1Z	K	2.0	CIS6	0.75	NIS2	1.00	OTH3	0.30
820D6Q	K	2.0	CIS6	0.75	NIS2	1.00	---	---
820E3F	K	6.0	CIS6	0.75	NIS2	1.00	OTH3	0.30
820F0G	K	6.0	CIS6	0.75	NIS2	1.00	---	---
820G5J	K	4.0	CIS6	0.75	NIS2	1.00	OTH3	0.60

820H7D	K	6.0	CIS6	0.75	NIS2	1.00	OTH3	0.60
--------	---	-----	------	------	------	------	------	------

[0079] The compositions of Table 2a and comparative compositions STD1 and STD2 were applied to velvetleaf (*Abutilon theophrasti*, ABUTH) plants. Results at 4 days after treatment (4DAT) and 14 days after treatment (14DAT), averaged for all replicates of each treatment, are shown in Table 2b.

Table 2b.

Comp.	Glyphosate Application Rate (g a.e./ha)	ABUTH %inhibition (4DAT)	ABUTH %inhibition (14DAT)
820A9T	150, 250, 400, 800	5.0, 5.0, 10.0, 10.0	40.8, 75.0, 84.2, 93.8
820B4H	150, 250, 400, 800	20.0, 25.0, 35.0, 40.0	66.7, 76.7, 80.8, 81.7
820C1Z	150, 250, 400, 800	5.0, 5.0, 5.0, 10.0	79.2, 85.0, 95.0, 99.2
820D6Q	150, 250, 400, 800	5.0, 5.0, 5.0, 10.0	25.8, 76.7, 85.0, 91.7
820E3F	150, 250, 400, 800	5.0, 5.0, 10.0, 15.0	78.3, 80.8, 90.5, 97.3
820F0G	150, 250, 400, 800	5.0, 5.0, 10.0, 15.0	22.5, 65.0, 83.3, 94.8
820G5J	150, 250, 400, 800	10.0, 15.0, 20.0, 25.0	79.2, 85.0, 93.8, 100.0
820H7D	150, 250, 400, 800	10.0, 15.0, 20.0, 25.0	79.2, 85.0, 96.2, 99.3
STD1	150, 250, 400, 800	0.0, 5.0, 5.0, 5.0	0.0, 26.7, 70.8, 84.2
STD2	150, 250, 400, 800	5.0, 5.0, 10.0, 10.0	70.0, 85.0, 90.0, 98.7

[0080] The order of efficacy for ABUTH %inhibition was 820H7D > 820C1Z > 820G5J > 820E3F > STD2 > 820B4H > 820A9T > 765K4S > 820D6Q > 820F0G > STD1.

EXAMPLE 3

[0081] The effect of tank mixtures of NH₄-oxalate with glyphosate package premix formulations of RT Master™ and Field Master™ on velvetleaf and barnyardgrass was tested. Aqueous tank mix compositions were prepared containing Roundup® UltraMax, RT Master™ and Field Master™ along with NH₄-oxalate at three glyphosate a.e.:oxalate ratios (2:1, 10:1 and 30:1) these compositions and excipient ingredients are shown in Table 3a.

Table 3a.

Composition	Glyphosate (g a.e./L)	2,4-D (g a.e./L)	Gly:Oxalic Acid Ratio
UltraMax 21	445	---	2:1
UltraMax 101	445	---	10:1
UltraMax 301	445	---	30:1
RT Master 21	360	38.6	2:1
RT Master 101	360	38.6	10:1
RT Master 301	360	38.6	30:1
Field Master 21	68	---	2:1
Field Master 101	68	---	10:1
Field Master 301	68	---	30:1

[0082] The compositions of Table 3a and comparative compositions STD1, STD2, RT Master™ and Field Master™ were applied to velvetleaf (*Abutilon theophrasti*, ABUTH) and Japanese millet (*Echinochloa crus-galli* var. *frumentae*, ECHCF) plants. Results at 5 days after treatment (5DAT) and 16 days after treatment (16DAT), averaged for all replicates of each treatment, are

shown in Table 3b.

Table 3b.

Composition	Glyphosate App. Rate (g a.e./ha)	ABUTH %inhibition (5DAT)	ABUTH %inhibition (16DAT)	ECHCF %inhibition (5DAT)	ECHCF %inhibition (16DAT)
UltraMax 21	75, 100, 200	0.0, 0.0, 0.0	78.3, 88.3, 97.5	0.0, 0.0, 0.0	61.7, 80.8, 94.8
UltraMax 101	75, 100, 200	0.0, 0.0, 0.0	70.8, 85.8, 97.8	0.0, 0.0, 0.0	61.7, 79.2, 97.5
UltraMax 301	75, 100, 200	0.0, 0.0, 0.0	65.8, 80.8, 96.0	0.0, 0.0, 0.0	55.0, 80.8, 92.8
RT Master 21	75, 100, 200	0.0, 2.5, 5.0	75.0, 83.3, 94.7	0.0, 0.0, 0.0	65.0, 85.0, 96.2
RT Master 101	75, 100, 200	0.0, 3.3, 5.0	66.7, 77.5, 91.7	0.0, 0.0, 0.0	63.3, 82.5, 94.8
RT Master 301	75, 100, 200	0.0, 4.2, 5.0	54.2, 75.8, 90.0	0.0, 0.0, 0.0	60.0, 80.8, 93.5
Field Master 21	75, 100, 200	3.3, 13.3, 32.5	76.7, 88.3, 90.5	0.0, 7.5, 16.7	48.3, 60.0, 61.7
Field Master 101	75, 100, 200	2.5, 10.8, 26.7	65.0, 76.7, 80.0	0.0, 4.2, 11.7	35.0, 52.5, 54.2
Field Master 301	75, 100, 200	1.7, 6.7, 16.7	48.3, 74.2, 79.2	0.0, 1.7, 9.2	20.8, 47.5, 50.0
RT Master™	75, 100, 200	2.5, 4.2, 5.0	47.5, 61.7, 77.5	0.0, 0.0, 0.0	55.0, 78.3, 91.2
Field Master™	75, 100, 200	4.2, 16.7, 48.3	40.8, 54.2, 84.8	0.0, 8.3, 23.3	27.5, 49.2, 50.0
STD1	75, 100, 200	0.0, 0.0, 0.0	6.7, 45.8, 64.2	0.0, 0.0, 0.0	3.3, 30.0, 49.2

STD2	75, 100, 200	0.0, 0.0, 0.0	40.8, 75.8, 86.3	0.0, 0.0, 0.0	58.3, 77.5, 91.5
------	-----------------	------------------	---------------------	------------------	---------------------

[0083] The order of efficacy averaged across application rates for the ABUTH %inhibition was UltraMax 21 > Field Master 21 > UltraMax 101 > RT Master 21 > UltraMax 301 > RT Master 101 > Field Master 101 > RT Master 301 > STD2 > Field Master 301 > RT Master > Field Master > STD1. The order of efficacy averaged across application rates for ECHF %inhibition was RT Master 21 > RT Master 101 > UltraMax 101 > UltraMax 21 > RT Master 301 > UltraMax 301 > STD 3 > RT Master > Field Master 21 > Field Master 101 > Field Master > Field Master 301 > STD1. The order of efficacy averaged across application rates for both ABUTH and ECHCF combined was UltraMax 21 > RT Master 21 > UltraMax 101 > RT Master 101 > UltraMax 301 > RT Master 301 > STD2 > Field Master 21 > RT Master > Field Master 101 > Field Master 301 > Field Master > STD1.

EXAMPLE 4

[0084] The effect of tank mixtures of NH₄-oxalate with glyphosate premix formulations of Roundup® RTU and Fallow Master® on velvetleaf and barnyardgrass was tested. Aqueous tank mix compositions were prepared containing Roundup® UltraMax, Roundup® RTU and Fallow Master® along with NH₄-oxalate at three glyphosate a.e.:oxalate ratios (2:1, 10:1 and 30:1) these compositions and excipient ingredients are shown in Table 4a.

Table 4a

Composition	Gly.	Component 2 (g a.e./L)	Gly:Oxalic Acid Ratio
UltraMax 21	445 gae/L	---	2:1
UltraMax 101	445 gae/L	---	10:1

UltraMax 301	445 gae/L	---	30:1
RTU 21	1.9 wt% ae	(diquat)	2:1
RTU 101	1.9 wt% ae	(diquat)	10:1
RTU 301	1.9 wt% ae	(diquat)	30:1
Fallow Master® 21	197 gae/L	46 (dicamba)	2:1
Fallow Master 101	197 gae/L	46 (dicamba)	10:1
Fallow Master 301	197 gae/L	46 (dicamba)	30:1

[0085] The compositions of Table 4a and comparative compositions Roundup® RTU, Fallow Master®, STD1 and STD2 were applied to velvetleaf (*Abutilon theophrasti*, ABUTH) and Japanese millet (*Echinochloa crus-galli* var. *frumentae*, ECHCF) plants. Results at 14 days after treatment (14DAT), averaged for all replicates of each treatment, are shown in Table 4b.

Table 4b.

Composition	Glyphosate Application Rate (g a.e./ha)	ABUTH %inhibition (14DAT)	ECHCF %inhibition (14DAT)
UltraMax 21	75, 100, 200	51.7, 80.0, 90.0	67.5, 68.3, 75.8
UltraMax 101	75, 100, 200	51.7, 76.7, 85.8	64.2, 68.3, 73.3
UltraMax 301	75, 100, 200	46.7, 67.5, 85.0	60.0, 68.3, 71.7
RTU 21	75, 100, 200	23.3, 27.5, 38.3	1.7, 6.7, 56.7
RTU 101	75, 100, 200	5.0, 10.0, 38.3	0.8, 7.5, 50.0
RTU 301	75, 100, 200	4.2, 10.0, 32.5	4.2, 17.5, 63.3

Fallow Master® 21	75, 100, 200	68.3, 78.3, 87.5	65.0, 67.5, 89.7
Fallow Master 101	75, 100, 200	66.7, 81.7, 85.0	65.8, 67.5, 90.0
Fallow Master 301	75, 100, 200	66.7, 78.3, 85.8	65.0, 70.8, 91.8
Roundup RTU	75, 100, 200	5.8, 16.7, 35.0	5.8, 21.7, 61.7
Fallow Master	75, 100, 200	60.0, 70.0, 84.2	64.2, 66.7, 81.7
STD1	75, 100, 200	3.3, 7.5, 31.7	20.8, 40.8, 59.2
STD2	75, 100, 200	10.8, 40.0, 76.7	64.2, 65.0, 72.5

[0086] The order of efficacy averaged across application rates for the ABUTH %inhibition was FallowMaster 21 > FallowMaster 101 > FallowMaster 301 > UltraMax 21 > UltraMax 101 > FallowMaster > UltraMax 301 > STD2 > RTU 21 > Roundup RTU > RTU 101 > STU 301 > STD1. The order of efficacy averaged across application rates for ECHF %inhibition was FallowMaster 301 > FallowMaster 101 > FallowMaster 21 > FallowMaster > UltraMax 21 > UltraMax 101 > STD 3 > UltraMax 301 > STD1 > Roundup RTU > RTU 301 > RTU 21 > RTU 101. The order of efficacy averaged across application rates for both ABUTH and ECHCF combined was FallowMaster 301 > FallowMaster 101 > FallowMaster 21 > UltraMax 21 > FallowMaster > UltraMax 101 > UltraMax 301 > STD2 > STD1 > RTU 21 > Roundup RTU > RTU 301 > RTU 101.

EXAMPLE 5

[0087] The effect of combinations of potassium glyphosate and 2,4-D and isopropylamine glyphosate on pitted morningglory (IPOLA) and cocklebur (XANST) was tested. Aqueous concentrate compositions were prepared containing 360 g a.e./L potassium glyphosate salt, amounts of 2,4-D are reported in wt% a.e. and excipient ingredients as shown in Table 5a.

Table 5a.

Comp.	Gly.	2,4-D	Cmpnt. 2	wt%	Cmpnt. 3	wt%
100B2T	K	1.60	CIS6	9.2	---	---
085A9K	K	1.60	CIS6	9.1	---	---
501A0X	K	1.66	CIS2	9.2	---	---
501B4S	K	1.82	CIS2	9.1	---	---
047B7Z	K	3.13	CIS6	9.2	OTH6	2.16
059A3D	K	3.13	CIS2	9.2	OTH6	2.20

[0088] The compositions of Table 5a and comparative composition RT Master™ were applied to pitted morningglory (IPOLA) and cocklebur (XANST) plants. Results at 10 days after treatment (10DAT), averaged for all replicates of each treatment, are shown in Table 5b.

Table 5b.

Composition	Glyphosate Application Rate (g a.e./ha)	XANST %inhibition (10DAT)	IPOLA %inhibition (10DAT)
100B2T	100, 200, 300, 400	38.8, 68.8, 73.8, 70.0	60.0, 76.3, 81.3, 93.8
085A9K	100, 200, 300, 400	42.5, 65.0, 83.8, 83.8	60.0, 77.5, 78.8, 83.8
501A0X	100, 200, 300, 400	56.3, 71.3, 82.5, 91.3	65.0, 78.8, 82.5, 85.0
501B4S	100, 200, 300, 400	58.8, 65.0, 76.3, 85.0	62.5, 83.8, 80.0, 91.3
047B7Z	100, 200, 300, 400	58.8, 73.8, 87.5, 97.5	86.3, 86.3, 90.0, 93.8
059A3D	100, 200, 300, 400	70.8, 79.5, 88.0, 90.0	72.5, 88.8, 87.5, 97.5

RT Master™	100, 200, 300, 400	53.8, 69.5, 78.5, 83.3	75.0, 83.8, 92.5, 88.8
------------	-----------------------	---------------------------	---------------------------

[0089] The order of efficacy averaged across application rates for the XANST %inhibition was 059A3D > 047B7Z > 501A0X > 501B4S > RT Master™ > 085A9K > 100B2T. The order of efficacy averaged across application rates for IPOLA %inhibition was 047B7Z > 059A3D > RT Master™ > 501B4S > 100B2T > 501A0X > 085A9K. The order of efficacy averaged across application rates for both XANST and IPOLA combined was 059A3D > 047B7Z > RT Master™ > 501A0X > 501B4S > 085A9K > 100B2T.

EXAMPLE 6

[0090] The effect of combinations of potassium glyphosate and 2,4-D and isopropylamine glyphosate on pitted morningglory (IPOLA) plants was tested. Aqueous concentrate compositions were prepared containing 480 g a.e./L potassium glyphosate salt, 2,4-D reported in wt% a.e, and excipient ingredients as shown in Table 6a.

Table 6a.

Comp.	Gly.	2,4-D	Cmpnt. 2	wt%
506A2T	K	0.72	CIS2	7.25
506B9Z	K	0.71	CIS2	7.25
510A4H	K	0.60	CIS6	7.60
510B8V	K	0.60	CIS6	7.60
508A1B	K	0.72	CIS2	7.60
508B0G	K	0.72	CIS2	7.60
503B5P	K	0.66	CIS6	9.10
504A3L	K	0.65	CIS6	9.10
504B2I	K	0.74	CIS2	9.10
505A6S	K	0.72	CIS2	9.10

[0091] The compositions of Table 6a and comparative compositions RT Master™ and STD2 were applied to pitted morningglory (IPOLA). Results at 5 days after treatment (5DAT) and 12 days after treatment (12DAT), averaged for all replicates of each treatment, are shown in Table 6b.

Table 6b.

Comp.	Glyphosate App. Rate (g a.e./ha)	IPOLA %inhibition (5DAT)	IPOLA %inhibitio n (11DAT)	IPOLA %inhibitio n (12DAT)	XANST %inhibition (11DAT)
506A2T	100, 200, 400	30.0, 67.5, 75.0	25.0, 72.5, 81.7	30.0, 75.0, 90.0	60.0, 71.3, 86.3
506B9Z	100, 200, 400	33.8, 47.5, 80.0	66.3, 72.5, 86.7	33.8, 71.3, 90.0	65.0, 77.5, 81.3
510A4H	100, 200, 400	61.3, 58.8, 85.0	22.5, 72.5, 80.0	58.8, 73.8, 92.5	55.0, 77.5, 78.8
510B8V	100, 200, 400	41.3, 65.0, 78.8	30.0, 67.5, 83.4	45.0, 77.5, 90.5	45.0, 60.0, 92.5
508A1B	100, 200, 400	35.0, 77.5, 80.0	56.3, 78.8, 86.7	42.5, 85.0, 90.0	60.0, 63.8, 95.7
508B0 G	100, 200, 400	2.5, 70.0, 75.0	57.5, 73.8, 78.4	2.5, 80.0, 90.0	61.3, 78.8, 95.0
503B5P	100, 200, 400	30.0, 52.5, 75.0	41.3, 67.5, 86.0	42.5, 70.0, 90.0	55.0, 75.0, 82.5
504A3L	100, 200, 400	31.3, 71.3, 80.0	56.3, 67.5, 90.0	28.8, 78.8, 91.3	57.5, 76.3, 88.8
504B2I	100, 200, 400	26.3, 43.8, 73.8	47.5, 78.8, 90.0	41.3, 63.8, 82.5	61.3, 78.8, 91.3
505A6S	100, 200, 400	28.8, 62.5, 80.0	58.8, 71.3, 88.4	35.0, 75.0, 88.8	63.8, 76.3, 81.3

RT Master	100, 200, 400	72.5, 85.0, 85.0	63.8, 87.5, 93.4	77.5, 96.3, 98.0	70.0, 76.3, 92.0
STD2	100, 200, 400	36.3, 36.3, 41.3	23.8, 62.5, 78.4	23.8, 50.0, 67.5	45.0, 70.0, 97.5

[0092] The order of efficacy averaged across application rates for the IPOLA %inhibition averaged over both 5 and 12 days after treatment was RT Master > 510A4H > 508A1B > 510B8V > 504A3L > 505A6S > 506A2T > 503B5P > 506B9Z > 504B2I > 508B0G > STD2. The order of efficacy for the XANST %inhibition was RT Master > 508A1B > 504B2I > 506B9Z > 504A3L > 505A6S > 506A2T > 508B0G > 503B5P > STD2 > 510A4H > 510B8V.

EXAMPLE 7

[0093] The effect of 128A5X and composition 139H2K on *zebrina pendula* (ZEBPE) plants to determine the appropriate rates for commercial control was tested. Aqueous concentrate compositions were prepared containing the indicated amount of glyphosate salt measured in g a.e./L and excipient ingredients as shown in Table 7a.

Table 7a.

Comp.	Gly.	2,4-D	Cmpnt. 1	wt%
139H2K	IPA (570)	---	NIS5	0.05
128A5X	MEA (480)	---	CIS6	9.6

[0094] The compositions of Table 7a and comparative composition 128A5X were applied to *Zebrina pendula* (ZEBPE). Results at 29 days after treatment (29DAT), averaged for all replicates of each treatment, are shown in Table 7b.

Table 7b.

Composition	Glyphosate Application Rate (g a.e./ha)	ZEBPE %inhibition (29DAT)
139H2K	1000, 2000, 3000, 4000, 5000, 6000, 7000, 8000	53.3, 72.7, 87.0, 84.3, 91.7, 90.0, 89.3, 93.3
128A5X	1000, 2000, 3000, 4000, 5000, 6000, 7000, 8000	43.3, 45.0, 41.7, 48.3, 72.7, 79.0, 81.7, 85.0

[0095] From the data, application rates of 2000, 3000, 4000 and 5000 g a.e./ha were used for the next set of experiments on *Zebrina pendula* (ZEBPE).

EXAMPLE 8

[0096] The effect of glyphosate compositions on *Zebrina pendula* (ZEBPE) plants was tested. Aqueous concentrate compositions were prepared containing the listed amount of glyphosate salt in g a.e./L and excipient ingredients as shown in Table 8a.

Table 8a.

Comp.	Gly.	Cmpnt 1	wt%	Cmpnt 2	wt%	Cmpnt 3	wt%	Compnt 4/5	wt%
553I3Z	IPA (360)	CIS8	9.6	NIS6	6.4	NIS7	1.0	OTH7/ OTH8	1.5/1.0
239K5X	K (480)	CIS15	9.2	---	---	---	---	---	---
128A5X	MEA (480)	CIS6	9.6	---	---	---	---	---	---

[0097] The compositions of Table 8a and comparative composition 128A5X were applied to *Zebrina pendula* (ZEBPE). Results at 28 days after treatment (28DAT), averaged for all replicates of each treatment, are shown in Table 8b.

Table 8b.

Composition	Glyphosate Application Rate (g a.e./ha)	ZEBPE %inhibition (28DAT)
55313Z	2000, 3000, 4000, 5000	73.8, 89.5, 88.0, 91.3
239K5X	2000, 3000, 4000, 5000	67.5, 82.8, 86.0, 86.0
128A5X	2000, 3000, 4000, 5000	82.0, 86.3, 86.5, 88.5

[0098] The most active composition was 128A5X.

EXAMPLE 9

[0099] The effect of glyphosate compositions on *Zebrina pendula* (ZEBPE) plants was tested. Aqueous concentrate compositions were prepared containing the listed amount of glyphosate salt in wt% and excipient ingredients as shown in Table 9a.

Table 9a.

Comp.	Gly.	Cmpt 1	wt%	Cmpt 2	wt%	Cmpt 3	wt %	Cmpt 4	wt %
128A5X	MEA (38.2)	CIS6	9.6	---	---	---	---	---	---
128B1T	MEA (38.2)	CIS6	9.6	OTH 3	3.82	---	---	---	---
318A9H	K (36.9)	CIS1	7.4	NIS9	4.9	OTH 9	6.5	---	---

318B2V	K (36.9)	CIS1	7.4	NIS9	4.9	OTH 9	6.5	OTH3	3.7
265A4C	K (31)	CIS10	126 g/L	---	---	---	---	---	---
265B0E	K (31)	CIS10	126 g/L	OTH 3	3.1	---	---	---	---
683A7T	Amm (68)	CIS1	9.5	NIS8	11.6	OTH 4	0.4	OTH 18	0.1

[0100] The compositions of Table 9a and comparative composition 128A5X were applied to *Zebrina pendula* (ZEBPE). Results at 27 days after treatment (27DAT), averaged for all replicates of each treatment, are shown in Table 9b.

Table 9b.

Composition	Glyphosate Application Rate (g a.e./ha)	ZEBPE %inhibition (27DAT)
128M1T	2000, 3500, 5000, 6500	70.5, 82.8, 84.3, 91.3
318A9H	2000, 3500, 5000, 6500	73.0, 75.0, 82.8, 92.3
318B2V	2000, 3500, 5000, 6500	78.3, 79.3, 83.8, 88.5
265A4C	2000, 3500, 5000, 6500	81.8, 85.5, 84.3, 93.5
265B0E	2000, 3500, 5000, 6500	67.5, 75.0, 75.5, 77.5
683A7T	2000, 3500, 5000, 6500	76.3, 83.8, 84.0, 90.8
128A5X	2000, 3500, 5000, 6500	73.5, 82.5, 84.3, 87.0

[0101] The most active composition was 265A4C. 265B0E showed lower rates of control with added oxalic acid. There were compatibility problems between oxalic acid and the other ingredients in 265B0E and 318B2V.

EXAMPLE 10

[0102] The effect of glyphosate compositions on *commelina benghalensis* (COMBE) plants was tested. Aqueous concentrate compositions

were prepared containing the listed amount of glyphosate salt in wt% and excipient ingredients as shown in Table 10a.

Table 10a.

Comp.	Gly.	Cmpt 1	wt%	Cmpt 2	wt%	Cmpt 3	wt %	Cmpt 4/5	wt%
128A5X	MEA (38.2)	CIS 11	9.6	---	---	---	---	---	---
633F3J	Amm (68)	CIS3	9.5	NIS3	11.6	OTH 17	0.4	OTH 18	0.1
483H8Q	Amm (68)	CIS 11	5.7	NIS1	8.0	OTH1	8.3	OTH4 / OTH5	0.4/ 0.1
050A6B	K (30)	CIS 12	0.8	---	---	---	---	---	---

[0103] The compositions of Table 10a and comparative composition 128A5X were applied to *commelina* (COMBE). Results at 33 days after treatment (33DAT), averaged for all replicates of each treatment, are shown in Table 10b.

Table 10b.

Composition	Glyphosate Application Rate (g a.e./ha)	COMBE %inhibition (33DAT)
128A5X	600, 800, 1000, 1200	34.0, 50.0, 58.0, 63.0
633F3J	600, 800, 1000, 1200	29.0, 59.0, 62.0, 59.0
483H8Q	600, 800, 1000, 1200	27.0, 48.0, 52.0, 56.0
050A6B	600, 800, 1000, 1200	54.0, 65.0, 68.0, 72.0

[0104] The most active composition was 050A6B.

EXAMPLE 11

[0105] The effect of glyphosate compositions on *commelina benghalensis* (COMBE) plants was tested. Aqueous concentrate compositions were prepared containing the listed amount of glyphosate salt in wt% and excipient ingredients as shown in Table 11a.

Table 11a.

Comp.	Gly.	Cmpt 1	wt %	Cmpt 2	wt%	Cmpt 3	wt %	Cmpt 4/5	wt%
128A5X	MEA (38.2)	CIS 6	9.6	---	---	---	---	---	---
483H8Q	Amm (68)	CIS 11	5.7	NIS1	8.0	OTH1	8.3	OTH4 / OTH5	0.4/ 0.1
633F3J	Amm (68)	CIS 11	9.5	NIS3	11.6	OTH4	0.4	OTH5	0.1
634T9P	Amm (65)	CIS1	11. 0	NIS8	13.4	OTH4	0.4	OTH5	0.1
765K4S	K (36.3)	CIS5	9.0	NIS4	4.0	CIS7	1.0	---	---

[0106] The compositions of Table 11a and comparative composition 128A5X were applied to *commelina* (COMBE). Results at 20 days after treatment (20DAT), averaged for all replicates of each treatment, are shown in Table 11b.

Table 11b.

Composition	Glyphosate Application Rate (g a.e./ha)	COMBE %inhibition (20DAT)
128A5X	800, 1100, 1400, 1700	60.0, 75.0, 65.0, 83.8
483H8Q	800, 1100, 1400, 1700	26.3, 61.3, 53.8, 72.5
633F3J	800, 1100, 1400, 1700	35.0, 61.3, 72.5, 72.5
634T9P	800, 1100, 1400, 1700	41.3, 70.0, 80.0, 81.3
765K4S	800, 1100, 1400, 1700	52.5, 75.0, 74.3, 79.5

The most active composition was 128A5X.

EXAMPLE 12

[0107] The effect of glyphosate compositions on *commelina benghalensis* (COMBE) plants was tested. Aqueous concentrate compositions were prepared containing the listed amount of glyphosate salt in wt% and excipient ingredients as shown in Table 12a.

Table 12a.

Comp.	Gly.	Cmpt 1	wt %	Cmpt 2	wt%	Cmpt 3	wt %	Cmpt 4/5	wt%
128A5X	MEA (38.2)	CIS6	9.6	---	---	---	---	---	---
553I3Z	IPA (360)	CIS8	9.6	NIS6	6.4	NIS7	1.0	OTH7/OT H8	1.5/ 1.0
483H8Q	Amm (68)	CIS 11	5.7	NIS1	8.0	OTH1	8.3	OTH4 / OTH5	0.4/ 0.1
633F3J	Amm (68)	CIS3	9.5	NIS3	11.6	OTH4	0.4	OTH5	0.1
634T9P	Amm (65)	CIS1	11. 0	NIS8	13.4	OTH4	0.4	OTH5	0.1

765K4S	K (36.3)	CIS5	9.0	NIS4	4.0	CIS7	1.0	---	---
239K5X	K (480)	CIS 15	9.2	---	---	---	---	---	---

[0108] The compositions of Table 12a were applied to *commelina* (COMBE). Results at 22 days after treatment (22DAT), averaged for all replicates of each treatment, are shown in Table 12b.

Table 12b.

Composition	Glyphosate Application Rate (g a.e./ha)	COMBE %inhibition (22DAT)
128A5X	800, 1100, 1400, 1700	72.5, 73.3, 83.8, 80.8
553I3Z	800, 1100, 1400, 1700	75.5, 75.8, 87.2, 88.3
483H8Q	800, 1100, 1400, 1700	70.0, 75.8, 79.2, 84.2
633F3J	800, 1100, 1400, 1700	69.2, 74.2, 74.7, 71.7
634T9P	800, 1100, 1400, 1700	70.8, 74.2, 79.5, 80.0
765K4S	800, 1100, 1400, 1700	75.0, 70.0, 75.3, 79.2
239K5X	800, 1100, 1400, 1700	70.5, 77.5, 83.3, 80.0

[0109] Formulation 553I3Z was the most effective composition for *commelina*.

EXAMPLE 13

[0110] The effect of glyphosate compositions on *commelina benghalensis* (COMBE) plants was tested. Aqueous concentrate compositions were prepared containing the listed amount of glyphosate salt in wt% and excipient ingredients as shown in Table 13a.

Table 13a.

Comp.	Gly.	Cmpt 1	wt%	Cmpt 2	wt%	Cmpt 3	wt%	Cmpt 4/5	wt%
128A5X	MEA (38.2)	CIS6	9.6	---	---	---	---	---	---
481Z7Y	K (35.5)	carfent razone	0.18	---	---	---	---	---	---
633F3J	Amm (68)	CIS3	9.5	NIS3	11.6	OTH4	0.4	OTH5	0.1
634T9P	Amm (65)	CIS1	11.0	NIS8	13.4	OTH4	0.4	OTH5	0.1
765K4S	K (36.3)	CIS5	9.0	NIS4	4.0	CIS7	1.0	---	---
239K5X	K (480)	CIS 15	9.2	---	---	---	---	---	---

[0111] The compositions of Table 13a and RT Master were applied to *commelina* (COMBE). Results at 20 days after treatment (20DAT), averaged for all replicates of each treatment, are shown in Table 13b.

Table 13b.

Composition	Glyphosate Application Rate (g a.e./ha)	COMBE %inhibition (20DAT)
128A5X	600, 700, 800, 900, 1000	62.5, 69.2, 63.3, 70.5, 68.8
481Z7Y	600, 700, 800, 900, 1000	79.7, 91.3, 91.7, 90.5, 97.2
633F3J	600, 700, 800, 900, 1000	49.2, 64.2, 55.8, 61.7, 65.0
634T9P	600, 700, 800, 900, 1000	57.5, 70.5, 55.0, 68.5, 68.3
765K4S	600, 700, 800, 900, 1000	64.2, 69.2, 69.2, 77.5, 71.7
239K5X	600, 700, 800, 900, 1000	65.8, 66.3, 61.7, 81.3, 69.2
RT Master	600, 700, 800, 900, 1000	80.0, 89.2, 91.3, 89.3, 95.8

[0112] The most active compositions in this example were RT Master and 481Z7Y. These results indicate that a second active ingredient (2,4-D or carfentrazone) increases the activity of the composition against *commelina*.

EXAMPLE 14

[0113] The effect of glyphosate compositions on *commelina benghalensis* (COMBE) plants was tested. Aqueous concentrate compositions were prepared containing the listed amount of glyphosate salt in wt% and excipient ingredients as shown in Table 14a, as well as those shown in Table 13a.

Table 14a.

Comp.	Gly.	Cmpt 1	wt %	Cmpt 2	wt%	Cmpt 3	wt %	Cmpt 4/5	wt%
483H8Q	Amm (68)	CIS 11	5.7	NIS1	8.0	OTH1	8.3	OTH4 / OTH5	0.4/ 0.1
770X1C	K (36.7)	CIS5	7.0	NIS10	4.0	CIS4	3.0	---	---
772N5D	K (36.7)	CIS5	7.0	NIS11	3.0	CIS4	3.0	---	---
780Y4O	K (40)	CIS6	8.0	NIS2	3.0	OTH3	1.0	OTH2	1.0
822B9T	K (30.5)	2,4-D	1.0	CIS6	6.0	NIS2	2.5	---	---
822C6U	IPA (36)	2,4-D	1.8	CIS6	8.0	NIS2	3.0	---	---

[0114] The compositions of Tables 13a and 14a and RT Master were applied to *commelina* (COMBE). Results at 10 days after treatment (10DAT), 24 days after treatment (24DAT) and 41 days after treatment (41DAT), averaged for all replicates of each treatment, are shown in Table 14b.

Table 14b.

Composition	Glyphosate Application Rate (g a.e./ha)	COMBE %inhibition (10DAT)	COMBE %inhibition (24DAT)	COMBE %inhibition (41DAT)
128A5X	800, 1100, 1400, 1700, 2000	1.5, 1.7, 5.2, 6.0, 6.0	41.7, 46.7, 53.3, 44.2, 55.0	77.2, 96.7, 94.5, 93.0, 92.0
481Z7Y	800, 1100, 1400, 1700, 2000	84.2, 92.5, 92.5, 93.3, 96.1	80.8, 89.3, 91.7, 91.8, 92.7	15.0, 26.5, 35.0, 75.5, 62.6
633F3J	800, 1100, 1400, 1700, 2000	3.0, 1.7, 2.3, 2.3, 5.2	20.0, 22.5, 35.0, 34.2, 45.0	51.2, 74.2, 74.7, 77.7, 89.7
634T9P	800, 1100, 1400, 1700, 2000	1.5, 1.7, 5, 4.3, 7.3	18.3, 21.7, 35.8, 33.3, 47.9	50, 91.7, 94.2, 82.3, 94.6
765K4S	800, 1100, 1400, 1700, 2000	3.7, 3.0, 1.3, 3.0, 6.1	17.5, 38.3, 30.0, 35.0, 43.9	62.5, 90.5, 95.0, 96.0, 97.2
483H8Q	800, 1100, 1400, 1700, 2000	3.7, 1.7, 4.5, 1.2, 4.3	26.7, 19.2, 31.7, 24.2, 31.9	51.7, 73.0, 89.5, 90.5, 96.0
770X1C	800, 1100, 1400, 1700, 2000	0.5, 2.3, 2.8, 3.7, 7.1	21.7, 24.2, 37.5, 35.0, 44.9	55.8, 81.7, 90.5, 99.2, 100.6
772N5D	800, 1100, 1400, 1700, 2000	2.2, 1.7, 1.5, 3.8, 3.8	29.2, 50.8, 36.7, 39.2, 49.2	73.5, 86.8, 92.5, 95.5, 96.8

780Y4O	800, 1100, 1400, 1700, 2000	2.0, 3.0, 1.7, 3.7, 5.0	18.3, 31.7, 45.8, 35.0, 44.2	54.2, 88.3, 96.3, 90.8, 95.8
822B9T	800, 1100, 1400, 1700, 2000	34.2, 39.2, 36.7, 39.2, 40.0	70.8, 81.7, 79.7, 80.3, 84.7	98.3, 97.8, 100.0, 100.0, 100.0
822C6U	800, 1100, 1400, 1700, 2000	39.2, 42.5, 40.8, 41.7, 42.1	78.7, 85.0, 80.0, 91.8, 96.5	100.0, 100.0, 100.0, 100.0, 100.0
RT Master	800, 1100, 1400, 1700, 2000	40.0, 41.7, 41.7, 39.2, 38.3	77.7, 86.7, 90.7, 90.0, 91.5	100.0, 99.7, 100.0, 100.0, 100.0

[0115] In this experiment, the compositions that were most effective, particularly against *commelina* regrowth, were 822B9T, 822C6U and RT Master, which all contain 2,4-D as a second active ingredient.

EXAMPLE 15

[0116] The effect of glyphosate compositions on *commelina benghalensis* (COMBE) plants was tested. Aqueous concentrate compositions were prepared containing the listed amount of glyphosate salt in wt% and excipient ingredients as shown in Table 15a.

Table 15a.

Comp.	Gly.	Cmpt 1	wt %	Cmpt 2	wt%	Cmpt 3	wt %	Cmpt 4/5	wt%
128A5X	MEA (38.2)	CIS6	9.6	---	---	---	---	---	---

822B9T	K (30.5)	2,4-D	1.0	CIS6	6.0	NIS2	2.5	---	---
822C6U	IPA (36)	2,4-D	1.8	CIS6	8.0	NIS2	3.0	---	---

[0117] The compositions of Table 15a, Ultra Blazer, Cobra and RT Master were applied to *commelina* (COMBE). Results at 7 days after treatment (7DAT) and 24 days after treatment (24DAT), averaged for all replicates of each treatment, are shown in Table 15b.

Table 15b.

Composition	Glyphosate Application Rate (g a.e./ha)	COMBE %inhibition (7DAT)	COMBE %inhibition (24DAT)
128A5X	100, 200, 300, 400	0.2, 0.2, 0.5, 3.7	29.2, 30.8, 28.3, 33.3
822B9T	100, 200, 300, 400	42.5, 40.0, 43.3, 43.3	49.2, 56.7, 72.5, 74.2
822C6U	100, 200, 300, 400	40.8, 44.2, 45.0, 46.7	63.3, 76.7, 79.2, 83.3
Ultra Blazer	18, 35, 70, 140, 280, 420	0.0, 1.7, 4.3, 4.3, 6.7, 8.3	10.0, 10.0, 10.8, 15.0, 41.7, 45.0
Cobra	9, 18, 35, 70, 140, 210	7.5, 8.3, 13.3, 13.3, 20.0, 21.7	33.3, 46.7, 41.7, 42.5, 44.2, 47.5
RT Master	100, 200, 300	42.5, 41.7, 49.2	73.3, 78.3, 82.5

[0118] This experiment shows that mixtures of glyphosate and 2,4-D, such as 822B9T, 822C6U and RT Master are more efficacious against *commelina* than single ingredient formulations.

EXAMPLE 16

[0119] The effect of glyphosate compositions on *commelina benghalensis* (COMBE) plants was tested. Aqueous concentrate compositions were prepared containing the listed amount of glyphosate salt in wt% and excipient ingredients as shown in Table 16a.

Table 16a.

Comp.	Gly.	2,4-D	Cmpnt. 1	wt%	Cmpnt. 2	wt%	Cmpnt. 3	wt%
085Z5F	IPA (30.5)	---	CIS9	7.5	---	---	---	---
714V9J	IPA (30.4)	---	2,4-DB	3.3	---	---	---	---

[0120] The compositions of Table 16a, Assure II and RT Master were applied to *commelina* (COMBE). Results at 7 days after treatment (7DAT) and 31 days after treatment (31DAT), averaged for all replicates of each treatment, are shown in Table 16b.

Table 16b.

Composition	Glyphosate Application Rate (g a.e./ha)	COMBE %inhibition (7DAT)	COMBE %inhibition (31DAT)
085Z5F	300, 400, 500, 600, 700, 800, 900, 1000, 1100, 1200	6.0, 5.0, 5.0, 8.0, 9.0, 11.0, 12.0, 16.0, 24.0, 26.0	9.0, 22.0, 24.0, 36.0, 46.0, 51.0, 53.0, 57.0, 56.0, 49.0
714V9J	300, 400, 500, 600, 700, 800, 900, 1000, 1100, 1200	41.0, 44.0, 46.0, 50.0, 53.0, 53.0, 53.0, 58.0, 60.0, 62.0	2.0, 20.0, 23.0, 59.0, 54.0, 79.0, 80.0, 78.0, 74.0, 85.0

Assure II	5,10, 20, 30, 40, 80, 160	9.0, 10.0, 15.0, 12.0, 15.0, 13.0, 19.0	5.0, 5.0, 5.0, 5.0, 5.0, 0.0, 0.0
RT Master	300, 400, 500, 600, 700, 800, 900, 1000, 1100, 1200	63.0, 61.0, 70.0, 66.0, 68.0, 74.0, 77.0, 80.0, 80.0, 80.0	18.0, 24.0, 64.0, 85.0, 72.0, 94.0, 100.0, 95.0, 96.0, 100.0

[0121] Both 714V9J and RT Master were effective against *commelina* in this experiment, however, RT Master was the most efficacious.

EXAMPLE 17

[0122] The effect of glyphosate compositions on morningglory (IPOSS) plants was tested. Aqueous concentrate compositions were prepared containing the listed amount of glyphosate salt in wt% and excipient ingredients as shown in Table 17a.

Table 17a.

Comp.	Gly.	2,4-D	Cmpnt. 1	wt%	Cmpnt. 2	wt%	Cmpnt. 3	wt%
085Z5F	IPA (30.5)	---	CIS9	7.5	---	---	---	---
714V9J	(30.4)	---	2,4-DB	3.3	---	---	---	---

[0123] The compositions of Table 17a and RT Master were applied to morningglory (IPOSS). Results at 7 days after treatment (7DAT) and 15 days after treatment (15DAT), averaged for all replicates of each treatment, are shown in Table 17b.

Table 17b.

Compositio n	Glyphosate Application Rate (g a.e./ha)	IPOSS %inhibition (7DAT)	IPOSS %inhibition (15DAT)
-----------------	---	-----------------------------	------------------------------

085Z5F	100, 150, 200, 250, 300, 350, 400, 450, 500, 550, 600, 650, 700	12.0, 50.0, 43.0, 50.0, 48.0, 48.0, 51.0, 50.0, 49.0, 52.0, 59.0, 54.0, 60.0	8.0, 53.0, 58.0, 63.0, 76.6, 76.0, 79.0, 81.2, 82.8, 80.6, 84.2, 83.2, 84.2
714V9J	100, 150, 200, 250, 300, 350, 400, 450, 500, 550, 600, 650, 700	17.0, 31.0, 43.0, 39.0, 78.0, 77.0, 66.0, 77.0, 66.0, 77.0, 76.0, 75.0, 82.0, 77.0, 78.0	8.0, 20.0, 23.0, 27.0, 30.0, 26.0, 29.0, 34.0, 30.0, 29.0, 30.0, 33.0, 34.0
RT Master	100, 150, 200, 250, 300, 350, 400, 450, 500, 550, 600, 650, 700	54.0, 62.0, 66.0, 70.0, 70.0, 75.0, 76.0, 79.0, 80.0, 84.0, 80.0, 83.0, 81.0	42.0, 68.0, 96.0, 97.0, 96.0, 100.0, 100.0, 100.0, 100.0, 100.0, 100.0, 100.0, 100.0

[0124] RT Master was the most effective composition for controlling morningglory at 7 and 15 days after treatment.

EXAMPLE 18

[0125] The effect of glyphosate compositions on morningglory (IPOSS) plants was tested. Aqueous concentrate compositions were prepared containing the listed amount of glyphosate salt in wt% and excipient ingredients as shown in Table 18a.

Table 18a.

Comp.	Gly.	2,4-D	Cmpnt. 1	wt%	Cmpnt. 2	wt%	Cmpnt. 3	wt%
085Z5F	IPA (30.5)	---	CIS9	7.5	---	---	---	---
714V9J	(30.4)	---	2,4-DB	3.3	---	---	---	---

[0126] The compositions of Table 18a, Pursuit and RT Master were applied to morningglory (IPOSS). Results at 14 days after treatment (14DAT),

averaged for all replicates of each treatment, are shown in Table 18b.

Table 18b.

Composition	Glyphosate Application Rate (g a.e./ha)	IPOSS %inhibition (14DAT)
085Z5F	100, 125, 150, 175, 200, 225, 250, 275, 300, 325, 350, 375, 400	7.0, 11.0, 32.0, 44.0, 53.0, 53.0, 55.0, 61.0, 61.0, 61.0, 63.0, 68.0, 76.0
714V9J	100, 125, 150, 175, 200, 225, 250, 275, 300, 325, 350, 375, 400	6.0, 29.0, 36.0, 31.0, 51.0, 49.0, 68.0, 74.0, 66.0, 80.0, 77.0, 74.0, 79.0
Pursuit	4, 8, 16, 35, 70, 105	0.0, 6.0, 36.0, 41.0, 77.0, 80.0
RT Master	100, 125, 150, 175, 200, 225, 250, 275, 300, 325, 350, 375, 400	47.0, 53.0, 66.0, 69.0, 82.0, 75.0, 89.0, 90.0, 78.0, 89.0, 82.0, 89.0, 94.0

[0127] RT Master was the most efficacious composition at the application levels of the experiment.

EXAMPLE 19

[0128] Aqueous compositions were prepared containing potassium glyphosate salt, IPA 2,4-D salt and excipient ingredients as shown in Table 19a. The formulations were prepared by mixing the 40.5% w/w a.e. aqueous solution of IPA 2,4-D to a concentration in w/w% as indicated by [2,4-D] in Table 19a, surfactant(s), glycol followed by addition of 47.8 (47.4)% w/w a.e. aqueous solution of potassium glyphosate to a concentration in w/w% as indicated by [gly] in Table 19a and then taking the total volume to 100% with water. Formulations were tested for cloud point and for density.

Table 19a.

Comp.	[Gly]	[2,4-D]	Cmpt. 1	wt%	Cmpt. 2	wt%	Cmpt. 3	wt%	Cloud pt. °C
612A7G	37.2	0.76	CIS6	4.72	NIS2	4.59	OTH2	4.01	72
613A9L	37.3	1.08	CIS6	4.74	NIS2	4.60	OTH2	3.99	52
638A1J	36.5	0.73	CIS6	4.63	CIS14	4.57	---	---	87
638B4T	36.8	0.75	CIS6	6.19	CIS14	3.05	---	---	72
639A5Z	36.6	0.75	CIS6	6.95	CIS14	2.30	---	---	67
639B3X	37.1	0.76	CIS6	3.95	NIS2	4.57	CIS14	0.78	52
640B7Q	36.6	0.73	CIS6	9.14	---	---	---	---	55
641A9V	36.7	0.74	CIS6	6.95	CIS13	2.30	---	---	68
641B8D	37.3	0.76	CIS6	4.73	NIS2	3.25	OTH10	5.00	57
645A7S	36.6	0.71	CIS13	8.91	---	---	---	---	>85
645B2B	35.8	0.65	CIS6	8.09	OTH2	5.51	---	---	56
645C1I	34.7	0.72	CIS6	8.70	OTH2	6.01	---	---	60
646A8K	36.6	0.72	CIS6	4.58	CIS13	4.57	OTH2	1.43	>85
646B3Z	34.8	0.69	CIS6	4.35	CIS13	4.35	OTH2	1.67	>85
654B9U	35.8	0.64	CIS6	7.97	---	---	OTH2	5.51	56
656A1T	36.6	0.74	CIS6	6.08	CIS13	3.07	OTH2	2.96	71
656B8Y	36.6	0.73	CIS6	7.33	CIS13	1.83	OTH2	4.18	54
656C3G	36.6	0.73	CIS6	7.31	CIS13	1.83	OTH2	3.17	70
665A2T	34.3	0.69	CIS6	8.53	OTH2	6.01	---	---	70
665B3O	34.2	0.68	CIS6	8.55	OTH2	5.00	---	---	73
667B6Z	34.2	0.68	CIS2	8.54	OTH2	4.99	---	---	79
668A5V	34.2	0.69	CIS6	8.56	OTH10	6.02	---	---	71
669B2O	36.6	0.74	CIS6	7.33	CIS13	1.83	OTH10	3.17	58
669C9X	36.6	0.73	CIS6	7.31	CIS13	1.83	OTH2	2.18	60
670A4F	36.6	0.74	CIS6	7.31	CIS13	1.83	OTH10	2.18	61
670B9G	36.6	0.73	CIS6	7.32	CIS13	1.82	OTH2	1.17	62

670C6L	36.6	0.74	CIS6	7.33	CIS13	1.83	OTH10	1.19	63
682A0M	36.6	0.73	CIS6	6.86	CIS13	2.29	OTH2	0.71	71
682B5V	36.6	0.72	CIS6	6.85	CIS13	2.29	---	---	69
682C7P	36.6	0.73	CIS6	6.10	CIS13	3.05	---	---	77
684A4O	36.6	0.73	CIS6	6.53	CIS13	2.61	OTH2	0.40	73
694A9Y	36.6	---	CIS6	13.8	---	---	---	---	---
695A2D	36.6	---	CIS6	8.12	CIS13	4.07	---	---	---
697A3U	36.6	0.71	CIS6	6.86	CIS14	2.29	---	---	70
697B5Y	36.6	0.71	CIS6	6.85	CIS14	2.29	OTH2	1.48	72
697C2T	36.6	0.71	CIS6	6.11	CIS14	3.05	---	---	76
698A8R	36.6	0.72	CIS6	6.09	CIS14	3.05	OTH2	3.97	74
312A6E	36.6	0.74	CIS6	4.57	NIS2	4.59	OTH2	4.01	72
313A1V	36.6	1.05	CIS6	4.59	NIS2	4.59	OTH2	3.99	52
316A5G	36.2	0.72	CIS6	4.53	NIS2	4.53	OTH10	4.00	66
316B7Y	36.2	0.72	CIS6	4.53	NIS2	4.53	OTH10	3.00	61
317A0J	36.2	0.72	CIS6	4.53	NIS2	4.53	OTH10	2.02	57
318A4B	35.8	0.72	CIS6	4.47	NIS2	4.50	OTH10	4.02	71
338A2W	36.6	0.73	CIS6	4.57	CIS14	4.58	---	---	87
338B4F	36.7	0.74	CIS6	6.09	CIS14	3.04	---	---	72
339A3Q	36.6	0.74	CIS6	6.84	CIS14	2.29	---	---	67
339B9P	36.6	0.74	CIS6	3.83	CIS14	0.77	NIS2	4.57	52
341A7H	36.6	0.73	CIS6	6.84	CIS13	2.29	---	---	68
341B5Z	36.6	0.74	CIS6	4.57	NIS2	3.24	OTH10	5.00	57
346B6T	36.6	0.73	CIS6	6.10	NIS2	3.05	OTH10	4.99	55
346C8X	35.8	0.72	CIS6	4.48	CIS14	0.74	NIS2	3.72	58
351A9M	35.8	0.73	CIS6	4.49	NIS2	4.50	OTH10	5.00	74
351B2V	35.8	0.72	CIS6	5.96	NIS2	2.99	OTH10	5.02	62
352A6G	35.8	0.72	CIS6	3.73	CIS13	0.75	NIS2	4.50	63
352B4N	36.6	0.74	CIS6	4.48	CIS4	4.49	OTH10	5.01	>90

352C5Z	36.6	0.73	CIS6	6.73	CIS4	2.24	OTH10	5.01	80
355A9K	36.6	1.46	CIS6	4.58	CIS4	4.59	OTH10	4.99	73
328D3J	34.3	0.69	CIS6	8.60	OTH10	5.99	---	---	72
331H1K	35.9	0.72	CIS6	4.48	NIS2	4.50	OTH10	4.99	74
074A2E	35.4	0.67	CIS2	6.11	---	---	---	---	60
074B9O	36.3	0.51	CIS2	5.97	---	---	---	---	62
075A3Q	36.2	0.54	CIS2	7.55	---	---	---	---	66
077C5Y	39.9	0.15	CIS2	6.78	---	---	---	---	61
078A8U	38.8	0.33	CIS6	6.11	OTH18	0.28	---	---	60
083A6B	24.3	1.99	CIS2	8.94	---	---	---	---	64
083B0V	27.8	1.47	CIS2	6.10	---	---	---	---	57
084A6G	41.2	0.17	CIS2	5.67	---	---	---	---	60
084B4R	39.8	0.24	CIS2	6.64	---	---	---	---	62
084C2W	36.2	0.57	CIS6	7.25	---	---	---	---	63
085A8I	28.3	1.59	CIS6	9.09	---	---	---	---	61
085B3S	38.9	0.29	CIS6	6.79	---	---	---	---	61
085C6H	39.2	0.25	CIS6	6.60	---	---	---	---	60
096A5F	39.3	0.25	CIS6	6.63	---	---	---	---	61
098B2X	36.4	0.57	CIS6	6.05	---	---	---	---	60
100A6T	28.2	1.62	CIS2	9.25	---	---	---	---	77
100B5G	28.7	1.60	CIS6	9.23	---	---	---	---	67
501A8V	28.7	1.66	CIS2	9.18	---	---	---	---	66
503A3S	36.1	0.64	CIS2	9.09	---	---	---	---	64
503C5A	35.8	0.65	CIS6	9.30	---	---	---	---	60
505A4R	36.1	0.72	CIS2	9.05	---	---	---	---	60
505C7P	35.9	0.75	CIS2	7.40	---	---	---	---	61
506B1V	36.3	0.71	CIS2	7.27	---	---	---	---	60
508B2M	36.1	0.72	CIS2	7.53	---	---	---	---	60
508E9C	35.9	0.66	CIS2	7.70	---	---	---	---	58

509B0K	36.0	0.62	CIS2	7.53	---	---	---	---	61
510B7L	36.2	0.59	CIS6	7.59	---	---	---	---	61
548B1Z	28.4	1.59	CIS2	9.12	---	---	---	---	56
564B0Y	28.3	1.66	CIS6	9.13	OTH3	1.02	---	---	64
569B2W	28.6	2.49	CIS6	7.23	OTH6	1.47	---	---	64
580B6G	28.3	3.09	CIS6	7.60	OTH6	2.55	OTH13	2.27	65
581A8J	28.4	3.21	CIS6	6.87	OTH6	2.58	OTH13	2.27	61
581B3E	27.9	3.15	CIS6	3.88	OTH6	4.81	OTH13	5.14	>90
405A8N	28.8	2.69	CIS6	9.61	OTH6	1.49	---	---	65
406B7V	28.8	3.08	CIS6	9.22	OTH6	1.65	---	---	57

[0129] Additional aqueous compositions were prepared containing potassium glyphosate salt, 2,4-D acid and excipient ingredients as shown in Table 19b. The formulations were prepared by mixing the 98% w/w a.e. aqueous solution of 2,4-D acid to a concentration in w/w% as indicated by [2,4-D] in Table 19b, surfactant(s), glycol followed by addition of 47.8 (47.4)% w/w a.e. aqueous solution of potassium glyphosate to a concentration in w/w% as indicated by [gly] in Table 19b and then taking the total volume to 100% with water. Formulations were tested for cloud point and for density.

Table 19b.

Comp.	[Gly]	[2,4-D]	Cmpt. 1	wt%	Cmpt. 2	wt%	Cmpt. 3	wt%	Cloud pt. °C
447A6T	23.2	2.64	CIS12	8.44	---	---	---	---	58
448A1Z	23.6	2.75	CIS5	9.83	---	---	---	---	64
448C5G	24.6	2.96	CIS5	10.3	---	---	---	---	71
451A0I	25.5	2.99	CIS5	9.83	---	---	---	---	50
472A2W	26.2	2.04	CIS5	1.72	CIS12	5.16	---	---	56
473B8K	30.4	2.99	CIS5	8.42	---	---	---	---	61
474A3R	28.1	2.70	CIS5	11.5	---	---	---	---	56

489A6H	28.2	2.00	CIS4	9.28	---	---	---	---	59
489B9M	29.1	2.90	CIS4	8.55	---	---	---	---	72
489C5V	28.3	2.97	CIS4	9.05	---	---	---	---	74
489D1Q	28.3	2.95	CIS4	8.99	---	---	---	---	74
009C4N	29.8	3.41	CIS11	8.72	OTH6	4.15	---	---	69
012A7O	27.9	3.09	CIS11	8.96	OTH6	2.13	---	---	57
012B2X	28.1	3.10	CIS11	8.85	OTH5	2.55	---	---	65
013A6K	28.3	3.10	CIS11	9.76	OTH6	2.83	---	---	66
013B6T	28.4	3.12	CIS11	9.27	OTH6	2.76	---	---	63
021B0U	28.4	3.14	CIS15	9.17	OTH6	2.77	---	---	62
026A8V	26.0	2.87	CIS6	8.59	CIS7	2.96	---	---	64
026B9Y	26.7	2.84	CIS2	9.19	CIS7	2.54	---	---	65
028A3Q	27.1	2.95	CIS6	10.4	CIS7	1.83	---	---	53
028B0H	27.5	2.99	CIS6	8.74	CIS7	5.81	---	---	53
029A4L	26.8	3.08	CIS6	11.1	CIS7	4.78	---	---	60
029B3V	28.0	3.08	CIS2	9.55	CIS7	5.81	---	---	58
034A6P	28.3	2.92	CIS2	9.95	OTH16	2.68	---	---	58
034B7Y	27.9	2.94	CIS2	9.16	OTH16	2.68	---	---	60
044A1L	28.8	3.08	CIS2	9.00	OTH6	2.39	---	---	67
044B5T	28.1	3.04	CIS2	9.21	OTH6	2.19	---	---	66
045A0X	38.2	3.09	CIS2	9.28	OTH6	2.24	---	---	65
045B8Q	28.5	3.10	CIS2	9.19	OTH6	2.17	---	---	65
046A2W	28.8	3.10	CIS6	9.13	OTH6	2.24	OTH14	1.15	75
047A6F	28.1	3.09	CIS6	9.09	OTH6	1.51	OTH14	1.87	62
047B7K	28.4	3.13	CIS2	9.21	OTH6	2.16	---	---	67
059A0U	28.4	3.13	CIS2	9.21	OTH6	2.19	---	---	65
066B1V	26.4	2.73	CIS6	7.70	CIS7	4.55	OTH14	8.70	60
071A3P	32.6	0.92	CIS6	7.74	---	---	---	---	60
072A5S	34.6	0.60	CIS2	7.84	---	---	---	---	60

072C9W	36.6	0.51	CIS2	7.69	---	---	---	---	62
073A4G	34.6	0.54	CIS2	5.89	---	---	---	---	60
073B0M	36.3	0.47	CIS2	6.08	---	---	---	---	62
501B2U	29.0	1.70	CIS2	9.23	---	---	---	---	74
501C9H	28.7	1.82	CIS2	9.07	---	---	---	---	64
502A6G	33.1	1.04	CIS6	9.26	---	---	---	---	62
502B1J	34.7	0.78	CIS2	9.26	---	---	---	---	62
502C7K	36.2	0.61	CIS2	9.10	---	---	---	---	61
503B3L	36.1	0.66	CIS6	9.08	---	---	---	---	61
504A8T	36.1	0.65	CIS6	9.08	---	---	---	---	60
504B4P	36.1	0.74	CIS2	9.09	---	---	---	---	61
505B0X	36.1	0.88	CIS2	7.23	---	---	---	---	50
506A4R	36.1	0.30	CIS2	7.24	---	---	---	---	61
508A5J	36.1	0.30	CIS2	7.62	---	---	---	---	60
508C1P	36.0	0.34	CIS2	7.52	---	---	---	---	58
508D2W	35.7	0.31	CIS6	7.50	---	---	---	---	59
509A9I	35.8	0.28	CIS2	7.67	---	---	---	---	60
510A7Z	36.3	0.26	CIS6	7.53	---	---	---	---	61
546A0V	28.4	1.57	CIS2	8.46	OTH9	1.92	---	---	60
548A2D	28.5	1.57	CIS2	9.23	OTH9	2.24	---	---	60
549B3X	28.5	1.45	CIS2	9.19	OTH9	2.22	---	---	66
549C1K	28.5	1.45	CIS2	9.23	OTH9	2.03	---	---	71
551B8D	28.5	1.47	CIS2	9.22	OTH9	2.11	---	---	66
553A7U	28.2	1.44	CIS2	9.60	OTH9	2.87	---	---	58
553B2F	28.4	1.44	CIS2	9.19	OTH9	2.55	---	---	63
564A6Y	28.4	1.60	CIS6	9.21	OTH3	1.00	---	---	68

[0130] Additional aqueous compositions were prepared containing potassium glyphosate salt, octyl amine 2,4-D salt and excipient ingredients as shown in Table 19c. The formulations were prepared by mixing an aqueous solution of 2,4-D octyl amine salt to a concentration in w/w% as indicated by [2,4-D] in Table 19b, surfactant(s), glycol followed by addition of 47.8 (47.4)% w/w a.e. aqueous solution of potassium glyphosate to a concentration in w/w% as indicated by [gly] in Table 19b and then taking the total volume to 100% with water. Formulations were tested for cloud point and for density.

Table 19c.

Comp.	[Gly]	[2,4-D]	Cmpt. 1	wt%	Cmpt. 2	wt%	Cmpt. 3	wt%	Cloud pt. °C
009A5T	31.3	3.18	CIS2	9.95	OTH6	5.17	---	---	81
009B9Z	30.4	2.94	CIS11	9.33	OTH6	4.76	---	---	71

EXAMPLE 20

[0131] The effect of glyphosate and combinations of 2,4-D and glyphosate on Roundup® ready soy was tested at 1 day, 3 days and 7 days after treatment. Aqueous concentrate compositions were prepared containing potassium glyphosate salt, reported in wt.% a.e. and excipient ingredients as shown for formulations in Table 19a above. The formulations were compared to RT Master® and Roundup Weathermax®. The compositions and comparative compositions RT Master® and Roundup Weathermax®, were applied to Roundup® ready soy plants. Results, averaged for all replicates of each treatment, are shown in Table 20a.

Table 20a.

Composition	Glyphosate Application Rate (g a.e./ha)	RR Soy %inhibition (1DAT)	RR Soy %inhibition (3DAT)	RR Soy %inhibition (7DAT)

656A1T	841, 1681, 3362	15.0, 19.2, 25.0	14.2, 19.2, 30.0	12.2, 20.0, 44.2
665A2T	841, 1681, 3362	10.8, 23.3, 25	10.8, 26.7, 31.7	10.5, 27.5, 48.3
667B6Z	841, 1681, 3362	13.3, 18.3, 21.7	13.3, 18.3, 23.3	11.0, 18.3, 39.2
668A5V	841, 1681, 3362	15.0, 20.8, 21.7	15.8, 23.3, 25.0	15.5, 25.0, 47.5
682A0M	841, 1681, 3362	15.0, 18.3, 29.2	15.0, 20.8, 30.8	13.3, 20.8, 44.2
646A8K	841, 1681, 3362	12.5, 15.0, 20.8	10.0, 17.5, 25.0	12.3, 19.2, 35.0
694A9Y	841, 1681, 3362	4.7, 7.5, 9.5	5.0, 7.0, 15.8	5.0, 10.0, 15.8
695A2D	841, 1681, 3362	1.0, 3.0, 5.0	3.0, 3.7, 15.0	3.0, 3.7, 13.3
Weathermax	841, 1681, 3362	1.0, 1.7, 3.0	1.0, 1.3, 4.3	0.7, 3.0, 3.7
RT Master	841, 1681, 3362	23.3, 20.0, 17.5	23.3, 23.3, 20.8	25.8, 41.7, 56.7

EXAMPLE 21

[0132] The effect of glyphosate and combinations of 2,4-D and glyphosate on velvetleaf was tested. Aqueous concentrate compositions were prepared containing potassium glyphosate salt, reported in wt.% a.e. and excipient ingredients as shown for the formulations in Table 19a above. The formulations were compared to RT Master® and Roundup Weathermax®. The compositions comparative compositions RT Master® and Roundup Weathermax®, were applied to velvetleaf (*Abutilon theophrasti*, ABUTH) plants. Results, averaged for all replicates of each treatment, are shown in Table 21a.

Table 21a.

Composition	Glyphosate Application Rate (g a.e./ha)	ABUTH %inhibition (14DAT)
656A1T	100, 200, 300, 400	14.2, 67.5, 83.8, 87.5
665A2T	100, 200, 300, 400	36.7, 75.8, 85, 91.7
667B6Z	100, 200, 300, 400	38.3, 74.2, 84.3, 90.3
668A5V	100, 200, 300, 400	48.3, 78.3, 87.5, 89.5
682A0M	100, 200, 300, 400	37.5, 72.5, 83.7, 89.8
646A8K	100, 200, 300, 400	40.8, 80, 85, 90.3
694A9Y	100, 200, 300, 400	67.5, 82.5, 90.8, 93.3
695A2D	100, 200, 300, 400	58.3, 79.2, 88, 90.8
Weathermax	100, 200, 300, 400	35.8, 73.3, 85.8, 90.8
RT Master	100, 200, 300, 400	18.3, 70, 80, 89.2

[0133] The order of efficacy for ABUTH %inhibition averaged over all application rates was 694A9Y > 695A2D > 668A5V > 646A8K > 665A2T > 667B6Z > Weathermax > 682A0M > RT Master > 656A1T.

EXAMPLE 22

[0134] The effect of glyphosate and combinations of 2,4-D and glyphosate on velvetleaf was tested at 16 days after treatment. Aqueous concentrate compositions were prepared containing potassium glyphosate salt, reported in wt.% a.e. and excipient ingredients as shown for formulations in Table 22a. The formulations were compared to RT Master® and Roundup Weathermax®. The 681C4J composition in Table 19a, compositions in Table 22a and comparative compositions RT Master® and Roundup Weathermax®, were applied to velvetleaf (*Abutilon theophrasti*, ABUTH) plants. Results, averaged for all replicates of each treatment, are shown in Table 22b.

Table 22a.

Comp.	[Gly]	[2,4-D]	Cmpt. 1	wt%	Cmpt. 2	wt%	Cmpt. 3	wt%
937C2V	40.0	---	CIS6	6.65	CIS14	3.34	---	---
936D9G	36.6	---	CIS6	13.6	---	---	---	---
974B3X	34.3	0.69	CIS6	8.60	OTH10	5.99	---	---
935A8Z	35.9	0.72	CIS6	4.48	NIS2	6.42	OTH10	4.99
342B6V	36.2	---	CIS6	9.05	NIS2	6.52	---	---
346A4F	36.6	---	CIS6	13.7	---	---	---	---
353A1S	36.6	—	CIS6	9.15	CIS4	4.58	---	---

Table 22b.

Composition	Glyphosate Application Rate (g a.e./ha)	ABUTH %inhibition (16DAT)
937C2V	100, 200, 300, 400	30.8, 70.8, 80.8, 91.2
936D9G	100, 200, 300, 400	44.2, 77.5, 85.5, 94.3
974B3X	100, 200, 300, 400	41.7, 70.8, 87.5, 92.3
935A8Z	100, 200, 300, 400	31.7, 62.5, 81.3, 89.0
342B6V	100, 200, 300, 400	45.8, 73.3, 90.0, 92.0
346A4F	100, 200, 300, 400	41.7, 70.0, 84.7, 91.8
353A1S	100, 200, 300, 400	43.3, 77.5, 91.7, 98.3
681C4J	100, 200, 300, 400	38.3, 70.8, 83.2, 89.3
Weathermax	100, 200, 300, 400	28.3, 72.5, 80.8, 90.0
RT Master	100, 200, 300, 400	25.0, 65.0, 84.2, 92.2

[0135] The order of efficacy for ABUTH %inhibition averaged over all application rates was 353A1S > 936D9G > 342B6V > 974B3X > 346A4F > 681C4J > 937C2V > Weathermax > RT Master > 935A8Z.

EXAMPLE 23

[0136] The effect of glyphosate and combinations of 2,4-D and glyphosate on velvetleaf was tested at 15 days after treatment. Aqueous concentrate compositions were prepared containing potassium glyphosate salt, reported in wt.% a.e. and excipient ingredients as shown for formulations in Tables 22a. The formulations were compared to RT Master® and Roundup Weathermax®. The 338A2W, 338B4F, 339A3Q, 341A7H, 352C5Z and 355A9K compositions in Table 19a, the 974B3X and 935A8Z compositions in Table 23a and comparative compositions RT Master® and Roundup Weathermax®, were applied to velvetleaf (*Abutilon theophrasti*, ABUTH) plants. Results, averaged for all replicates of each treatment, are shown in Table 23a.

Table 23a.

Composition	Glyphosate Application Rate (g a.e./ha)	ABUTH %inhibition (15DAT)
338A2W	100, 200, 300, 400	41.7, 74.2, 86.5, 90.5
338B4F	100, 200, 300, 400	43.3, 74.2, 88.0, 90.2
339A3Q	100, 200, 300, 400	46.7, 70.8, 89.2, 91.3
341A7H	100, 200, 300, 400	38.3, 73.3, 87.5, 91.7
352C5Z	100, 200, 300, 400	50.0, 72.5, 85.8, 90.7
355A9K	100, 200, 300, 400	45.8, 63.3, 84.7, 89.3
974B3X	100, 200, 300, 400	52.5, 76.7, 88.8, 94.0
935A8Z	100, 200, 300, 400	43.3, 65.0, 84.2, 88.2
Weathermax	100, 200, 300, 400	35.0, 67.5, 82.5, 96.2
RT Master	100, 200, 300, 400	32.5, 69.2, 84.5, 94.0

[0137] The order of efficacy for ABUTH %inhibition averaged over all application rates was 974B3X > 352C5Z > 339A3Q > 338B4F > 338A2W > 341A7H > 355A9K > Weathermax > 935A8Z > RT Master.

EXAMPLE 24

[0138] The effect of glyphosate and combinations of 2,4-D and glyphosate on Roundup ready soybean plants was tested at 1 day and 3 days after treatment. Aqueous concentrate compositions were prepared containing potassium glyphosate salt, reported in wt.% a.e. and excipient ingredients as shown for formulations in Tables 22a. The formulations were compared to RT Master® and Roundup Weathermax®. The 681C4J, 342B2H, 342C3A, 346A7C, 353A8Q, 338A2W, 338B4F, 339A3Q, 341A7H, 352C5Z and 355A9K compositions in Table 19a, the 937C2V, 936D9G, 974B3X and 935A8Z compositions in Table 22a and comparative compositions RT Master® and Roundup Weathermax®, were applied to Roundup ready soybean (GLXMG) plants using an AI nozzle. Results, averaged for all replicates of each treatment, are shown in Table 24a.

Table 24a.

Composition	Glyphosate Application Rate (g a.e./ha)	RR Soy %injury (1DAT)	RR Soy %injury (3DAT)
681C4J	841, 1681	2.0, 4.0	2.8, 6.5
342B2H	841, 1681	2.8, 10.3	5.0, 13.3
342C3A	841, 1681	1.8, 10.0	4.8, 11.3
346A7C	841, 1681	6.5, 14.0	5.0, 14.0
353A8Q	841, 1681	2.3, 10.8	6.0, 12.8
338A2W	841, 1681	25.0, 26.3	31.3, 33.8
338B4F	841, 1681	23.8, 23.8	30.0, 35.0
339A3Q	841, 1681	27.5, 26.3	30.0, 35.0
341A7H	841, 1681	18.8, 30.0	21.3, 38.8
352C5Z	841, 1681	25.0, 31.3	27.5, 36.3
355A9K	841, 1681	21.3, 26.3	20.0, 31.3
937C2V	841, 1681	2.0, 4.0	2.3, 6.8

936D9G	841, 1681	14.5, 20.0	9.3, 22.5
974B3X	841, 1681	28.8, 32.5	33.8, 38.8
935A8Z	841, 1681	25.0, 23.8	23.8, 27.5
Weatherma x	841, 1681	2.0, 10.0	2.0, 5.0
RT Master	841, 1681	21.3, 18.8	22.5, 27.5

[0139] The order of efficacy for %control in Roundup ready soybeans averaged over all application rates using an AI nozzle was 974B3X > 352C5Z > 339A3Q > 338A2W > 338B4F > 341A7H > 935A8Z > 355A9K > RT Master > 936D9G > 346A7C > 342B2H > 353A8Q > 342C3A > Weathermax > 681C4J > 937C2V.

EXAMPLE 25

[0140] The experiment in Example 24 was repeated using an TT nozzle to apply the formulations to the Roundup ready soybean plants. Results, averaged for all replicates of each treatment, are shown in Table 25a.

Table 25a.

Composition	Glyphosate Application Rate (g a.e./ha)	RR Soy %injury (1DAT)	RR Soy %injury (3DAT)
681C4J	841, 1681	1.0, 3.5	1.0, 2.8
342B2H	841, 1681	2.3, 6.0	2.0, 6.5
342C3A	841, 1681	3.0, 5.0	1.5, 5.5
346A7C	841, 1681	2.5, 4.5	2.3, 5.3
353A8Q	841, 1681	2.0, 6.0	2.3, 6.8
338A2W	841, 1681	20.0, 26.3	16.3, 28.8
338B4F	841, 1681	22.5, 26.3	15.0, 26.3
339A3Q	841, 1681	22.5, 22.5	20.0, 25.0

341A7H	841, 1681	20.0, 25.0	15.0, 26.3
352C5Z	841, 1681	26.3, 23.8	25.0, 30.0
355A9K	841, 1681	28.8, 26.3	27.5, 31.3
937C2V	841, 1681	1.8, 3.0	1.3, 3.5
936D9G	841, 1681	3.3, 7.3	3.3, 12.0
974B3X	841, 1681	20.0, 26.3	16.3, 28.8
935A8Z	841, 1681	21.3, 26.3	18.8, 27.5
Weatherma x	841, 1681	2.0, 4.5	2.0, 5.0
RT Master	841, 1681	18.8, 20.0	22.5, 27.5

[0141] The order of efficacy for %control in Roundup ready soybeans averaged over all application rates using a TT nozzle was 355A9K > 352C5Z > 935A8Z > 974B3X > 338A2W > 338B4F > 339A3Q > RT Master > 341A7H > 936D9G > 353A8Q > 342B2H > 342C3A > 346A7C > Weathermax > 937C2V > 681C4J.

EXAMPLE 26

[0142] Aqueous compositions were prepared containing potassium glyphosate salt, dicamba and excipient ingredients as shown in Table 26a. Formulations were tested for cloud point and for density.

Table 26a.

Comp.	[Gly]	[dica mba]	Cmpt. 1	wt%	Cmpt. 2	wt%	Cmpt. 3	wt%	Cloud pt. °C
561A2J	39.8	0.80	CIS6	6.63	CIS4	3.32	---	---	51
561B5M	39.8	0.79	CIS6	6.40	CIS4	3.56	---	---	53
561C3A	39.8	0.79	CIS6	5.97	CIS4	3.98	---	---	56
561D7Y	39.8	0.79	CIS6	5.62	CIS4	4.33	---	---	58
561E8P	39.8	0.50	CIS6	6.64	CIS4	3.32	---	---	66

WHAT IS CLAIMED IS:

1. An aqueous herbicidal concentrate composition comprising:
 - (a) glyphosate or a herbicidal derivative thereof;
 - (b) an auxin herbicide component comprising one or more auxin herbicides selected from the group consisting of 2,4-D, 2,4-DB, dichlorprop, MCPA, MCPB, mecoprop, dicamba, picloram, quinclorac and agriculturally acceptable salts or esters thereof; and
 - (c) a surfactant component in solution or stable suspension, emulsion or dispersion, comprising one or more surfactants;wherein the glyphosate (acid equivalent basis) and the auxin herbicide component (acid equivalent basis) are present in a weight ratio of at least 32:1 and the composition has a cloud point of at least about 50°C and a crystallization point not higher than about 0°C.
2. The composition of claim 1 wherein the auxin herbicide component comprises one or more auxin herbicides selected from the group consisting of 2,4-D, dicamba and agriculturally acceptable salts or esters thereof.
3. The composition of claim 1 wherein the auxin herbicide component comprises 2,4-D or an agriculturally acceptable salt or ester thereof.
4. The composition of claim 1 wherein the auxin herbicide component comprises dicamba or an agriculturally acceptable salt or ester thereof.
5. The composition of claim 1 wherein the glyphosate is present predominantly in the form of the potassium salt thereof.
6. The composition of claim 5 wherein at least about 90 percent by weight of the glyphosate is present in the form of the potassium salt thereof.

7. The composition of claim 1 wherein the glyphosate is present in an amount of at least about 325 grams acid equivalent per liter.

8. The composition of claim 1 wherein the glyphosate is present in an amount of at least about 425 grams acid equivalent per liter.

9. The composition of claim 1 wherein the glyphosate is present in an amount of at least about 525 grams acid equivalent per liter.

10. The composition of claim 1 wherein the glyphosate (acid equivalent basis) and auxin herbicide component (acid equivalent basis) are present in a weight ratio of at least 38:1.

11. The composition of claim 1 wherein the glyphosate (acid equivalent basis) and auxin herbicide component (acid equivalent basis) are present in a weight ratio of at least 50:1.

12. The composition of claim 1 wherein the surfactant component is present in an amount of at least about 3 wt.% based on the total weight of the composition.

13. The composition of claim 1 wherein the glyphosate (acid equivalent basis) and the surfactant component are present in a weight ratio of from about 1.9:1 to about 6:1.

14. The composition of claim 1 wherein the herbicide content consists of glyphosate or a herbicidal derivative thereof and the auxin herbicide component.

15. The composition of claim 1 wherein the composition further comprises an organic solvent, said composition containing not more than 10 weight percent of said organic solvent based on the total weight of the composition.

16. A method of killing or controlling weeds or unwanted vegetation comprising:

diluting the composition of claim 1 in an amount of water to form an application mixture; and

applying a herbicidally effective amount of the application mixture to foliage of the weeds or unwanted vegetation.

17. The method of claim 16 wherein the weeds or undesired plants comprise a dicot.

18. The method of claim 17 wherein the dicot comprises at least one broadleaf weed.

19. The method of claim 16 wherein the application mixture is applied to foliage of crop plants genetically transformed or selected to tolerate glyphosate and said weeds or said unwanted vegetation growing in close proximity to the crop plants to control the weeds or unwanted vegetation while leaving the crop plants substantially unharmed.

20. The method of claim 19 wherein the weeds or undesired plants comprise a dicot.

21. The method of claim 20 wherein the dicot comprises at least one broadleaf weed.


22. An aqueous herbicidal concentrate composition comprising:

(a) glyphosate, predominantly in the form of the potassium salt thereof, in a concentration of at least 65 grams acid equivalent per liter;

(b) an auxin herbicide component comprising one or more auxin herbicides selected from the group consisting of 2,4-D, 2,4-DB, dichlorprop, MCPA, MCPB, mecoprop, dicamba, picloram, quinclorac and agriculturally acceptable salts or esters thereof; and


(c) a first surfactant component in solution or stable suspension, emulsion or dispersion and comprising one or more surfactants selected from the group consisting of:

(i) a secondary or tertiary amine having the formula:




wherein R^1 is hydrocarbyl having from 1 to about 30 carbon atoms, and R^2 and R^3 are hydrogen or hydrocarbyl having from 1 to about 30 carbon atoms;

(ii) dialkoxylated quaternary ammonium salt having the formula:




wherein R^{11} is hydrocarbyl or substituted hydrocarbyl having from 1 to about 30 carbon atoms, R^{12} in each of the x (R^{12}O) and y (R^{12}O) groups is independently $\text{C}_2\text{-C}_4$ alkylene, R^{13} is hydrogen, or a linear or branched alkyl group having from 1 to about 4 carbon atoms, R^{14} hydrocarbyl or substituted hydrocarbyl having from 1 to about 30 carbon atoms, x and y are independently an average number from 1 to about 40, and X^- is an agriculturally acceptable anion;

(iii) monoalkoxylated quaternary ammonium salts having the formula:




wherein R^{21} and R^{25} are independently hydrogen or hydrocarbyl or substituted hydrocarbyl having from 1 to about 30 carbon atoms, R^{24} is hydrocarbyl or substituted hydrocarbyl having from 1 to about 30 carbon atoms, R^{22} in each of the x^2 ($R^{22}O$) groups is independently C_2 - C_4 alkylene, R^{23} is hydrogen, or a linear or branched alkyl group having from 1 to about 30 carbon atoms, x is an average number from 1 to about 60, and X^- is an agriculturally acceptable anion;

(iv) quaternary ammonium salts having the formula:




wherein R^{31} , R^{33} and R^{34} are independently hydrogen or hydrocarbyl or substituted hydrocarbyl having from 1 to about 30 carbon atoms, R^{32} is hydrocarbyl or substituted hydrocarbyl having from 1 to about 30 carbon atoms, and X^- is an agriculturally acceptable anion;

(v) ether amines having the formula:




wherein R^{41} is hydrocarbyl or substituted hydrocarbyl having from 1 to about 30 carbon atoms; R^{42} is hydrocarbylene or substituted hydrocarbylene having from 2 to about 30 carbon atoms; R^{43} and R^{44} are independently hydrogen, hydrocarbyl or substituted hydrocarbyl having from 1 to about 30 carbon atoms, or - $(R^{45}O)_x R^{46}$, R^{45} in each of the x^4 ($R^{45}-O$) groups is independently C_2 - C_4 alkylene, R^{46} is hydrogen, or a linear or branched alkyl group having from 1 to about 4 carbon atoms, and x^4 is an average number from 1 to about 50;

(vi) amine oxides having the formula:




wherein R^{51} , R^{52} and R^{53} are independently hydrogen, hydrocarbyl or substituted hydrocarbyl, $-(\text{R}^{54}\text{O})_x\text{R}^{55}$, or $-\text{R}^{56}(\text{OR}^{54})_x\text{OR}^{55}$, R^{54} in each of the x^5 (R^{54}O) groups is independently C_2 - C_4 alkylene, R^{55} is hydrogen, or a linear or branched alkyl group having from 1 to about 30 carbon atoms, R^{56} is hydrocarbylene or substituted hydrocarbylene having from 2 to about 6 carbon atoms, x^5 is an average number from 1 to about 50, and the total number of carbon atoms in R^{51} , R^{52} and R^{53} is at least 8;

(vii) dialkoxylated amines having the formula:



wherein R^{61} is a linear or branched alkyl, linear or branched alkenyl, linear or branched alkynyl, aryl, or aralkyl group having from about 6 to about 30 carbon atoms, R^{62} in each of the x^6 (R^{62}O) and the y^6 (R^{62}O) groups is independently C_2 - C_4 alkylene, R^{63} is hydrogen, or a linear or branched alkyl group having from 1 to about 4 carbon atoms and x^6 and y^6 are independently an average number from 1 to about 40;


(viii) aminated alkoxylated alcohols having the formula:



wherein R^{71} is hydrogen or hydrocarbyl or substituted hydrocarbyl having from 1 to about 30 carbon atoms; R^{72} in each of the x^7 (R^{72}O) and y^7 (R^{72}O) groups is independently C_2 - C_4 alkylene; R^{73} is hydrocarbylene or substituted hydrocarbylene having from 2 to about 30 carbon atoms; R^{74} and R^{75} are each


independently hydrogen, hydrocarbyl or substituted hydrocarbyl having from 1 to about 30 carbon atoms, $-(R^{76})_n^7-(R^{72}O)_y^7R^{77}$, or R^{74} and R^{75} , together with the nitrogen atom to which they are attached, form a cyclic or heterocyclic ring; R^{76} is hydrocarbylene or substituted hydrocarbylene having from 1 to about 30 carbon atoms; R^{77} is hydrogen or a linear or branched alkyl group having 1 to about 4 carbon atoms, n^7 is 0 or 1, x^7 and y^7 are independently an average number from 1 to about 60;

(ix) alkyl alkoxyated phosphates having the formula:



wherein R^{81} and R^{83} are independently a linear or branched alkyl, linear or branched alkenyl, linear or branched alkynyl, aryl, or aralkyl group having from about 4 to about 30 carbon atoms; R^{82} in each of the m ($R^{82}O$) and the n ($R^{82}O$) groups is independently C_2 - C_4 alkylene; and m and n are independently from 1 to about 30;

(x) alkyl alkoxyated phosphates having the formula:



wherein R^{91} is a linear or branched alkyl, linear or branched alkenyl, linear or branched alkynyl, aryl, or aralkyl group having from about 8 to about 30 carbon atoms; R^{92} in each of the a ($R^{92}O$) groups is independently C_2 - C_4 alkylene; and a is from 1 to about 30; and

(xi) alkylpolyglycosides having the formula:




where R^{101} is hydrogen or C_{1-18} hydrocarbyl, R^{104} is hydrogen or C_{1-4} hydrocarbyl, q is 0 or 1, *sug* is (i) an open or cyclic structure derived from sugars or (ii) a hydroxyalkyl, polyhydroxyalkyl or poly(hydroxyalkyl)alkyl group, u is an average number from 1 to about 2, and v is an integer from 1 to 3.

23. The composition of claim 22 wherein the first surfactant comprises
(i) a secondary or tertiary amine having the formula:



wherein R^1 is hydrocarbyl having from 1 to about 30 carbon atoms, and R^2 and R^3 are hydrogen or hydrocarbyl having from 1 to about 30 carbon atoms;

- (ii) monoalkoxylated quaternary ammonium salts having the formula:




wherein R^{21} and R^{25} are independently hydrogen or hydrocarbyl having from 1 to about 30 carbon atoms, R^{24} is hydrocarbyl having from 1 to about 30 carbon atoms, R^{22} in each of the $x^2 (R^{22}O)$ groups is independently C_2-C_4 alkylene, R^{23} is hydrogen, or a linear or branched alkyl group having from 1 to about 30 carbon atoms, x is an average number from 1 to about 60, and X^- is an agriculturally acceptable anion;

- (iii) quaternary ammonium salts having the formula:


wherein R^{81} and R^{83} are independently a linear or branched alkyl, linear or branched alkenyl, linear or branched alkynyl, aryl, or aralkyl group having from about 4 to about 30 carbon atoms; R^{82} in each of the m ($R^{82}O$) and the n ($R^{82}O$) groups is independently C_2 - C_4 alkylene; and m and n are independently from 1 to about 30;

(vi) alkyl alkoxyated phosphates having the formula:



wherein R^{91} is a linear or branched alkyl, linear or branched alkenyl, linear or branched alkynyl, aryl, or aralkyl group having from about 8 to about 30 carbon atoms; R^{92} in each of the a ($R^{92}O$) groups is independently C_2 - C_4 alkylene; and a is from 1 to about 30; and

(vii) alkylpolyglycosides having the formula:




where R^{101} is hydrogen or C_{1-18} hydrocarbyl, R^{104} is hydrogen or C_{1-4} hydrocarbyl, q is 0 or 1, *sug* is (i) an open or cyclic structure derived from sugars or (ii) a hydroxyalkyl, polyhydroxyalkyl or poly(hydroxyalkyl)alkyl group, u is an average number from 1 to about 2, and v is an integer from 1 to 3.

24. The composition of claim 22 wherein the auxin herbicide component comprises one or more auxin herbicides selected from the group consisting of 2,4-D, dicamba and agriculturally acceptable salts or esters thereof.


25. The composition of claim 22 wherein the auxin herbicide component comprises 2,4-D or an agriculturally acceptable salt thereof and the surfactant component comprises one or more surfactants selected from the group consisting of:

(i) ether amines having the formula:




wherein R^{41} is hydrocarbyl or substituted hydrocarbyl having from 1 to about 30 carbon atoms; R^{42} is hydrocarbylene or substituted hydrocarbylene having from 2 to about 30 carbon atoms; R^{43} and R^{44} are independently hydrogen, hydrocarbyl or substituted hydrocarbyl having from 1 to about 30 carbon atoms, or - $(\text{R}^{45}\text{O})_x\text{R}^{46}$, R^{45} in each of the $x^4(\text{R}^{45}\text{-O})$ groups is independently $\text{C}_2\text{-C}_4$ alkylene, R^{46} is hydrogen, or a linear or branched alkyl group having from 1 to about 4 carbon atoms, and x^4 is an average number from 1 to about 50;

(ii) dialkoxylated amines having the formula:



wherein R^{61} is a linear or branched alkyl, linear or branched alkenyl, linear or branched alkynyl, aryl, or aralkyl group having from about 6 to about 30 carbon atoms, R^{62} in each of the $x^6(\text{R}^{62}\text{O})$ and the $y^6(\text{R}^{62}\text{O})$ groups is independently $\text{C}_2\text{-C}_4$ alkylene, R^{63} is hydrogen, or a linear or branched alkyl group having from 1 to about 4 carbon atoms and x^6 and y^6 are independently an average number from 1 to about 40; and

(iii) aminated alkoxylated alcohols having the formula:




wherein R^{71} is hydrogen or hydrocarbyl or substituted hydrocarbyl having from 1 to about 30 carbon atoms; R^{72} in each of the $x^7(\text{R}^{72}\text{O})$ and $y^7(\text{R}^{72}\text{O})$ groups is independently $\text{C}_2\text{-C}_4$ alkylene; R^{73} is hydrocarbylene or substituted

hydrocarbylene having from 2 to about 30 carbon atoms; R^{74} and R^{75} are each independently hydrogen, hydrocarbyl or substituted hydrocarbyl having from 1 to about 30 carbon atoms, $-(R^{76})_n^7-(R^{72}O)_y^7R^{77}$, or R^{74} and R^{75} , together with the nitrogen atom to which they are attached, form a cyclic or heterocyclic ring; R^{76} is hydrocarbylene or substituted hydrocarbylene having from 1 to about 30 carbon atoms; R^{77} is hydrogen or a linear or branched alkyl group having 1 to about 4 carbon atoms, n^7 is 0 or 1, x^7 and y^7 are independently an average number from 1 to about 60.


26. The composition of claim 25 further comprising a second surfactant component comprising one or more surfactants selected from the group consisting of:

(a) a secondary or tertiary amine having the formula:



wherein R^1 is hydrocarbyl having from 1 to about 30 carbon atoms, and R^2 and R^3 are hydrogen or hydrocarbyl having from 1 to about 30 carbon atoms;


(b) dialkoxylated quaternary ammonium salt having the formula:



(2)


wherein R^{11} is hydrocarbyl or substituted hydrocarbyl having from 1 to about 30 carbon atoms, R^{12} in each of the x $(R^{12}O)$ and y $(R^{12}O)$ groups is independently C_2-C_4 alkylene, R^{13} is hydrogen, or a linear or branched alkyl group having from 1 to about 4 carbon atoms, R^{14} hydrocarbyl or substituted hydrocarbyl having from 1 to about 30 carbon atoms, x and y are independently an average number from 1 to about 40, and X^- is an agriculturally acceptable anion;

(c) monoalkoxylated quaternary ammonium salts having the formula:




wherein R^{21} and R^{25} are independently hydrogen or hydrocarbyl or substituted hydrocarbyl having from 1 to about 30 carbon atoms, R^{24} is hydrocarbyl or substituted hydrocarbyl having from 1 to about 30 carbon atoms, R^{22} in each of the x^2 (R^{22}O) groups is independently C_2 - C_4 alkylene, R^{23} is hydrogen, or a linear or branched alkyl group having from 1 to about 30 carbon atoms, x is an average number from 1 to about 60, and X^- is an agriculturally acceptable anion;

(d) quaternary ammonium salts having the formula:



wherein R^{31} , R^{33} and R^{34} are independently hydrogen or hydrocarbyl or substituted hydrocarbyl having from 1 to about 30 carbon atoms, R^{32} is hydrocarbyl or substituted hydrocarbyl having from 1 to about 30 carbon atoms, and X^- is an agriculturally acceptable anion;


(e) amine oxides having the formula:



wherein R^{51} , R^{52} and R^{53} are independently hydrogen, hydrocarbyl or substituted hydrocarbyl, $-(\text{R}^{54}\text{O})_x \text{R}^{55}$, or $-\text{R}^{56}(\text{OR}^{54})_x \text{OR}^{55}$, R^{54} in each of the x^5 (R^{54}O) groups


is independently C₂-C₄ alkylene, R⁵⁵ is hydrogen, or a linear or branched alkyl group having from 1 to about 30 carbon atoms, R⁵⁶ is hydrocarbylene or substituted hydrocarbylene having from 2 to about 6 carbon atoms, x⁵ is an average number from 1 to about 50, and the total number of carbon atoms in R⁵¹, R⁵² and R⁵³ is at least 8;

(f) alkyl alkoxyated phosphates having the formula:



wherein R⁸¹ and R⁸³ are independently a linear or branched alkyl, linear or branched alkenyl, linear or branched alkynyl, aryl, or aralkyl group having from about 4 to about 30 carbon atoms; R⁸² in each of the m (R⁸²O) and the n (R⁸²O) groups is independently C₂-C₄ alkylene; and m and n are independently from 1 to about 30;

(g) alkyl alkoxyated phosphates having the formula:




wherein R⁹¹ is a linear or branched alkyl, linear or branched alkenyl, linear or branched alkynyl, aryl, or aralkyl group having from about 8 to about 30 carbon atoms; R⁹² in each of the a (R⁹²O) groups is independently C₂-C₄ alkylene; and a is from 1 to about 30;

(h) alkylpolyglycosides having the formula:




where R^{101} is hydrogen or C_{1-18} hydrocarbyl, R^{104} is hydrogen or C_{1-4} hydrocarbyl, q is 0 or 1, *sug* is (i) an open or cyclic structure derived from sugars or (ii) a hydroxyalkyl, polyhydroxyalkyl or poly(hydroxyalkyl)alkyl group, u is an average number from 1 to about 2, and v is an integer from 1 to 3; and


(i) amines or quaternary ammonium salts having the formula:




or



or



or



wherein R^{111} is linear or branched alkyl or aryl having from about 4 to about 16 carbon atoms, R^{112} is hydrogen, methyl, ethyl, or $-(CH_2CH_2O)_dH$, R^{113} is hydrogen, methyl, ethyl, or $-(CH_2CH_2O)_eH$ wherein the sum of d and e is not more than about 5; R^{114} is hydrogen or methyl; R^{116} in each of the c ($R^{116}O$) groups is independently C_2-C_4 alkylene; R^{115} is hydrocarbylene or substituted hydrocarbylene having from 2 to about 6 carbon atoms; and A^- is an agriculturally acceptable anion.

27. The composition of claim 26 wherein the second surfactant component comprises said alkylpolyglycoside.

28. The composition of claim 22 wherein the glyphosate (acid equivalent basis) and the auxin herbicide component (acid equivalent basis) are present in a weight ratio of at least about 5:1.

29. The composition of claim 22 wherein the glyphosate (acid equivalent basis) and the auxin herbicide component (acid equivalent basis) are present in a weight ratio of at least about 30:1.

30. The composition of claim 22 wherein the glyphosate (acid equivalent basis) and the auxin herbicide component (acid equivalent basis) are present in a weight ratio of at least about 60:1.

31. The composition of claim 22 wherein at least about 90 percent by weight of the glyphosate is present in the form of the potassium salt thereof.

32. The composition of claim 22 wherein the glyphosate is present in an amount of at least about 325 grams acid equivalent per liter.

33. The composition of claim 22 wherein the glyphosate is present in an amount of at least about 425 grams acid equivalent per liter.

34. The composition of claim 22 wherein the glyphosate is present in an amount of at least about 525 grams acid equivalent per liter.

35. The composition of claim 22 wherein total surfactant present in the composition is in an amount of at least about 5 wt.% based on the total weight of the composition.

36. The composition of claim 22 wherein the glyphosate (acid equivalent basis) and the first surfactant component are present in a weight ratio of from about 1.9:1 to about 6:1.

37. The composition of claim 22 wherein the composition has a cloud point of at least about 50°C.

38. The composition of claim 22 wherein the composition has a crystallization point not higher than about 0°C.

39. The composition of claim 22 wherein the herbicide content consists of glyphosate or a herbicidal derivative thereof and the auxin herbicide component.

40. The composition of claim 22 wherein the composition further comprises an organic solvent, said composition containing not more than 10 weight percent of said organic solvent based on the total weight of the composition.

41. The composition of claim 26 wherein the first surfactant component and the second surfactant component are present in a weight ratio of from about 1:1 to about 6:1.

42. A method of killing or controlling weeds or unwanted vegetation comprising:

diluting the composition of claim 22 in an amount of water to form an application mixture; and

applying a herbicidally effective amount of the application mixture to foliage of the weeds or unwanted vegetation.

43. The method of claim 42 wherein the weeds or undesired plants comprise a dicot.

44. The method of claim 43 wherein the dicot comprises at least one broadleaf weed.

45. The method of claim 42 wherein the application mixture is applied to foliage of crop plants genetically transformed or selected to tolerate glyphosate and said weeds or said unwanted vegetation growing in close proximity to the crop plants to control the weeds or unwanted vegetation while leaving the crop plants substantially unharmed.

46. The method of claim 45 wherein the weeds or undesired plants comprise a dicot.

47. The method of claim 46 wherein the dicot comprises at least one broadleaf weed.

48. An aqueous herbicidal concentrate composition comprising:

(a) glyphosate, predominantly in the form of the isopropylammonium salt thereof, in a concentration of greater than 360 grams acid equivalent per liter;

(b) an auxin herbicide component comprising one or more auxin herbicides selected from the group consisting of 2,4-D, 2,4-DB, dichlorprop, MCPA, MCPB, mecoprop, dicamba, picloram, quinclorac and agriculturally acceptable salts or esters thereof; and

(c) a surfactant component in solution or stable suspension, emulsion or dispersion, comprising one or more surfactants;

wherein the glyphosate (acid equivalent basis) and the auxin herbicide component (acid equivalent basis) are present in a weight ratio of at least 9.5:1 and the composition has a cloud point of at least about 50°C and a crystallization point not higher than about 0°C.

49. The composition of claim 48 wherein the auxin herbicide component comprises one or more auxin herbicides selected from the group consisting of 2,4-D, dicamba and agriculturally acceptable salts or esters thereof.

50. The composition of claim 48 wherein the auxin herbicide component comprises 2,4-D or an agriculturally acceptable salt or ester thereof.

51. The composition of claim 48 wherein at least about 60 percent by weight of the glyphosate is present in the form of the isopropylammonium salt thereof.

52. The composition of claim 48 wherein at least about 75 percent by weight of the glyphosate is present in the form of the isopropylammonium salt thereof.

53. The composition of claim 48 wherein at least about 90 percent by weight of the glyphosate is present in the form of the isopropylammonium salt thereof.

54. The composition of claim 48 wherein the glyphosate is present in an amount of at least about 400 grams acid equivalent per liter.

55. The composition of claim 48 wherein the glyphosate is present in an amount of at least about 440 grams acid equivalent per liter.

56. The composition of claim 48 wherein the glyphosate is present in an amount of at least about 480 grams acid equivalent per liter.

57. The composition of claim 48 wherein the glyphosate (acid equivalent basis) and auxin herbicide component (acid equivalent basis) are present in a weight ratio of at least 10:1.

58. The composition of claim 48 wherein the glyphosate (acid equivalent basis) and auxin herbicide component (acid equivalent basis) are present in a weight ratio of at least 15:1.

59. The composition of claim 48 wherein the glyphosate (acid equivalent basis) and auxin herbicide component (acid equivalent basis) are present in a weight ratio of at least 20:1.

60. The composition of claim 48 wherein the surfactant component is present in an amount of at least about 5 weight percent based on the total weight of the composition.

61. The composition of claim 48 wherein the glyphosate (acid equivalent basis) and the surfactant component are present in a weight ratio of from about 1.9:1 to about 6:1.

62. The composition of claim 48 wherein the herbicide content consists of glyphosate or a herbicidal derivative thereof and the auxin herbicide component.

63. The composition of claim 48 wherein the composition further comprises an organic solvent, said composition containing not more than 10 weight percent of said organic solvent based on the total weight of the composition.

64. A method of killing or controlling weeds or unwanted vegetation comprising:

diluting the composition of claim 48 in an amount of water to form an application mixture; and

applying a herbicidally effective amount of the application mixture to foliage of the weeds or unwanted vegetation.

65. The method of claim 64 wherein the weeds or undesired plants comprise a dicot.

66. The method of claim 65 wherein the dicot comprises at least one broadleaf weed.

67. The method of claim 64 wherein the application mixture is applied to foliage of crop plants genetically transformed or selected to tolerate glyphosate and said weeds or said unwanted vegetation growing in close proximity to the crop plants to control the weeds or unwanted vegetation while leaving the crop plants substantially unharmed.

68. The method of claim 67 wherein the weeds or undesired plants comprise a dicot.

69. The method of claim 68 wherein the dicot comprises at least one broadleaf weed.

70. A method of killing or controlling weeds or unwanted plants comprising:

diluting an aqueous herbicidal concentrate composition in an amount of water to form an application mixture; and

applying a herbicidally effective amount of the application mixture to foliage of the weeds or unwanted plants, wherein the weeds or unwanted plants comprise *Commelina* and the aqueous herbicidal concentrate composition comprises:

- (a) glyphosate or a herbicidal derivative thereof;
- (b) an auxin herbicide component comprising one or more auxin herbicides selected from the group consisting of 2,4-D, 2,4-DB, dichlorprop,

MCPA, MCPB, mecoprop, dicamba, picloram, quinclorac and agriculturally acceptable salts or esters thereof; and

(c) a surfactant component in solution or stable suspension, emulsion or dispersion, comprising one or more surfactants.

71. The method of claim 70 wherein the glyphosate (acid equivalent basis) and the auxin herbicide component (acid equivalent basis) are present in a weight ratio of at least about 5:1.