

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
21 July 2005 (21.07.2005)

PCT

(10) International Publication Number
WO 2005/066123 A1

(51) International Patent Classification⁷: **C07D 207/06**,
223/02, 207/04

Shao-Ying; 5F, No. 42, Lane 30, Sindong Street, Songsha
Dist., Taipei City, 105 (TW).

(21) International Application Number:
PCT/US2004/044004

(74) Agent: **TSAO, Y., Rocky**; Fish & Richardson P.C., 225
Franklin Street, Boston, MA 02110-2804 (US).

(22) International Filing Date:
28 December 2004 (28.12.2004)

(81) Designated States (*unless otherwise indicated, for every
kind of national protection available*): AE, AG, AL, AM,
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,
ZW.

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/533,779 31 December 2003 (31.12.2003) US

(71) Applicant (*for all designated States except US*): **TAIGEN
BIOTECHNOLOGY**; 7F, 138 Shin Ming Rd., Neihu
Dist., Taipei 114 (TW).

(84) Designated States (*unless otherwise indicated, for every
kind of regional protection available*): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO,
SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN,
GQ, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **YANG, Syaulan**
[US/—]; 7F, 138 Shin Ming Rd., Neihu Dist., Taipei
114 (TW). **WU, Jen-Dar**; No. 76, Hudi, Minghua
Village, Dalin Township, Chiayi County, 62243 (TW).
SU, Feng-Yih; No. 201 Jhongshan Rd., Benfu Village,
Liouciou Township, Pingtung County, 92941 (TW). **KUO,
Chun-Wei**; 4F, No. 692 Jhongjheng Rd., Shihlin District,
Taipei City, 11173 (TW). **CHEN, Wen-Chang**; 4F, No.
37, Lane 6, Sec. 5, Chengde Rd., Shihlin District, Taipei
City, 11166 (TW). **XIANG, Yibin** [US/US]; 24 Marshall
Path, Acton, MA 01720 (US). **WANG, Ching-Cheng**; 7F,
138 Shin Ming Rd, Neihu Dist., Taipei 114 (TW). **LIAO,**

Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: PROTEASE INHIBITORS

(57) Abstract: This invention relates to treating an infection with a virus using protease inhibitors. Examples of the protease inhibitors include compounds of formula (I). Each variable is defined in the specification.



WO 2005/066123 A1

PROTEASE INHIBITORS

CROSS REFERENCE TO RELATED APPLICATION

Pursuant to 35 U.S.C. § 119(e), this application claims priority to U.S. Provisional Application Serial No. 60/533,779, filed December 31, 2003.

BACKGROUND


5 Coronavirus is a family of viruses that have the appearance of a corona when viewed under a microscope. Members of the coronavirus family can cause hepatitis in mice, gastroenteritis in pigs, and respiratory infections in birds and humans. Coronavirus was first isolated from chickens in 1937 by Beaudette and Hudson. In 1965, Tyrrell and Bynoe used cultures of human ciliated embryonic trachea to propagate the first human
10 coronavirus in vitro.

 Among the more than 30 strains of coronavirus isolated so far, three or four infect humans. For example, the severe acute respiratory syndrome, a newly emerged infectious disease, is associated with a novel coronavirus (Ksiazek et al., New England Journal Medicine, 2003, 348(20): 1953-1966). This life-threatening respiratory disease
15 caused worldwide outbreaks in 2003. Vaccines and drugs against severe acute respiratory syndrome virus are being vigorously sought. Nevertheless, the progress is rather slow due to safety concerns. Thus, there exists a need to develop drugs that are effective in treating infections with coronaviruses, as well as infections with other viruses.

SUMMARY

20 This invention is based on the unexpected discovery that certain peptide-like compounds are effective in treating viral infections by inhibiting viral proteases (e.g., coronaviral 3CL proteases, picornaviral proteases, or hepatitis C viral proteases).


 In one aspect, this invention features a method for treating an infection with a virus. The method includes administering to a subject in need thereof an effective
25 amount of a compound of formula (I):




In this formula, X is N(R_{a1}), O, or CH₂; or X and one of R₂ and R₃, together with the atom or atoms to which they are bonded, are C₃-C₂₀ heterocycloalkyl; in which R_{a1} is H or C₁-C₁₅ alkyl; R₁ is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, aryl, C(O)R_{b1}, CO₂R_{b1}, C(O)NR_{b1}R_{b2}, C(O)-N(R_{b1})-OR_{b2}, C(S)R_{b1}, C(S)NR_{b1}R_{b2}, S(O)R_{b1}, SO₂R_{b1}, S(O)NR_{b1}R_{b2}, S(O)-N(R_{b1})-OR_{b2}, SO₂NR_{b1}R_{b2}, or SO₃R_{b1}; in which each of R_{b1} and R_{b2}, independently, is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, or aryl; each of R₂ and R₃, independently, is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, aryl, OR_{c1}, SR_{c1}, or NR_{c1}R_{c2}; or X and one of R₂ and R₃, together with the atom or atoms to which they are bonded, are C₃-C₂₀ heterocycloalkyl; in which each of R_{c1} and R_{c2}, independently, is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, or aryl; each of R₄ and R₅, independently, is H, halogen, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, or aryl; one of R₆ and R₇ is C₁-C₁₅ alkyl substituted with C₃-C₂₀ heterocycloalkyl, heteroaryl, or aryl; and the other of R₆ and R₇ is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, or aryl; R₈ is H, halogen, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, aryl, OR_{d1}, or SR_{d1}; in which R_{d1} is H and C₁-C₁₅ alkyl; and each of R₉ and R₁₀, independently, is H, halogen, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, aryl, C(O)R_{e1}, CO₂R_{e1}, C(O)NR_{e1}R_{e2}, C(O)-N(R_{e1})-OR_{e2}, C(S)R_{e1}, C(S)NR_{e1}R_{e2}, CN, NO₂, S(O)R_{e1}, SO₂R_{e1}, S(O)NR_{e1}R_{e2}, S(O)-N(R_{e1})-OR_{e2}, SO₂NR_{e1}R_{e2}, SO₃R_{e1}, PO(OR_{e1})(OR_{e2}), PO(R_{e1})(R_{e2}), PO(NR_{e1}R_{e2})(OR_{e3}), PO(NR_{e1}R_{e2})(NR_{e3}R_{e4}), C(O)-N(R_{e1})-NR_{e2}R_{e3}, or C(S)-N(R_{e1})-NR_{e2}R_{e3}; or R₈ and R₁₀, taken together, is C₃-C₂₀ cycloalkyl or C₃-C₂₀ heterocycloalkyl; or R₉ and R₁₀, taken together, is C₃-C₂₀ cycloalkyl or C₃-C₂₀ heterocycloalkyl; in which each of R_{e1}, R_{e2}, R_{e3}, and R_{e4}, independently, is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, or aryl; or any two of R_{e1}, R_{e2}, R_{e3}, and R_{e4}, together with the atom or atoms to which they are bonded, is C₃-C₂₀ heterocycloalkyl; and the virus is a coronavirus or a hepatitis C virus.

In particular, this invention features a method for treating an infection with a severe acute respiratory syndrome virus, a human coronavirus 229E, or a hepatitis C virus, by administering to a subject in need thereof an effective amount of a compound of formula (I) shown above.

5 For example, one can administer to a subject infected with a virus (e.g., a severe acute respiratory syndrome virus) a compound of formula (I), in which X is N(R_{a1}) or O; or X and one of R₂ and R₃, together with the atom or atoms to which they are bonded, are C₃-C₂₀ heterocycloalkyl; R₁ is H, C(O)R_{b1}, CO₂R_{b1}, C(O)NR_{b1}R_{b2}, C(S)NR_{b1}R_{b2}, or SO₂R_{b1}; each of R₂ and R₃, independently, is H or C₁-C₁₅ alkyl; or X and one of R₂ and R₃, together with the atom or atoms to which they are bonded, are C₃-C₂₀ heterocycloalkyl; each of R₄ and R₅, independently, is H or C₁-C₁₅ alkyl; one of R₆ and R₇ is C₁-C₁₅ alkyl substituted with C₃-C₂₀ heterocycloalkyl, and the other of R₆ and R₇ is H; R₈ is H; and each of R₉ and R₁₀, independently, is H or CO₂R_{e1}. In this compound,



one of R₆ and R₇ can be ; one of R₂ and R₃ can be H or C₁-C₁₅ alkyl optionally substituted with halogen, heteroaryl, aryl, OR_{c1}, SR_{c1}, OC(O)R_{c1}, CO₂R_{c1}, C(O)NR_{c1}R_{c2}, NR_{c1}R_{c2}, N(R_{c1})-CO₂R_{c2}, N(R_{c1})-C(O)R_{c2}, N(R_{c1})-C(O)-NR_{c2}R_{c3}, N(R_{c1})-SO₂R_{c2}, SO₂R_{c1}, or O-SO₂-R_{c1}; or X and one of R₂ and R₃, together with the atom or atoms to which they are bonded, are C₃-C₂₀ heterocycloalkyl; in which R_{c3} is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, or aryl; and one of R₄ and R₅ can be H or C₁-C₁₅ alkyl optionally substituted with halogen, aryl, OR_{f1}, SR_{f1}, CO₂R_{f1}, C(O)NR_{f1}R_{f2}, SO₂R_{f1}, SO₃R_{f1}, or NR_{f1}R_{f2}; in which each of R_{f1} and R_{f2}, independently, is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, or aryl.

The term "treating" refers to administering one or more compounds of the invention to a subject, who has an infection with a virus, a symptom of such an infection, or a predisposition toward such an infection, with the purpose to confer a therapeutic effect, e.g., to cure, relieve, alter, affect, ameliorate, or prevent the infection with a virus, the symptom of it, or the predisposition toward it. The term "an effective amount" refers to the amount of one or more active compounds of the invention that is required to confer a therapeutic effect on a treated subject.

The term "alkyl" refers to a saturated or unsaturated, linear or branched, non-aromatic hydrocarbon moiety, such as -CH₃, -CH₂-, -CH₂-CH=CH₂-, or branched -C₃H₇. The term "cycloalkyl" refers to a saturated or unsaturated, non-aromatic, cyclic hydrocarbon moiety, such as cyclohexyl or cyclohexen-3-yl. The term

5 "heterocycloalkyl" refers to a saturated or unsaturated, non-aromatic, cyclic moiety having at least one ring heteroatom, such as 4-tetrahydropyranyl or 4-pyranyl. The term "aryl" refers to a hydrocarbon moiety having one or more aromatic rings. Examples of an aryl moiety include phenyl, phenylene, naphthyl, naphthylene, pyrenyl, anthryl, and phenanthryl. The term "heteroaryl" refers to a moiety having one or more aromatic rings

10 that contain at least one heteroatom. Examples of a heteroaryl moiety include furyl, furylene, fluorenyl, pyrrolyl, thienyl, oxazolyl, imidazolyl, thiazolyl, pyridyl, pyrimidinyl, quinazoliny, quinolyl, isoquinolyl and indolyl.

Alkyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl mentioned herein include both substituted and unsubstituted moieties, unless specified otherwise. Examples of

15 substituents on cycloalkyl, heterocycloalkyl, aryl, and heteroaryl include C₁-C₁₀ alkyl, C₂-C₁₀ alkenyl, C₂-C₁₀ alkynyl, C₃-C₈ cycloalkyl, C₅-C₈ cycloalkenyl, C₁-C₁₀ alkoxy, aryl, aryloxy, heteroaryl, heteroaryloxy, amino, C₁-C₁₀ alkylamino, C₁-C₂₀ dialkylamino, arylamino, diarylamino, hydroxyl, halogen, thio, C₁-C₁₀ alkylthio, arylthio, C₁-C₁₀ alkylsulfonyl, arylsulfonyl, cyano, nitro, acyl, acyloxy, carboxyl, and carboxylic ester.

20 On the other hand, examples of substituents on alkyl include all of the above-recited substituents except C₁-C₁₀ alkyl, C₂-C₁₀ alkenyl, and C₂-C₁₀ alkynyl. Cycloalkyl, heterocycloalkyl, aryl, and heteroaryl also include fused groups.

In another aspect, this invention features a method for inhibiting a viral protease (e.g., a coronaviral 3CL protease or a hepatitis C viral protease) in a cell. The method


25 includes contacting the cell with an effective amount of a compound of formula (I) shown above. In particular, this invention features a method for inhibiting a severe acute respiratory syndrome viral 3CL protease, a human coronaviral 229E protease, or a hepatitis C viral protease.

In still another aspect, this invention features a compound of formula (I) shown

30 above except that R₁ is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, aryl, C(O)R_{b1}, CO₂R_{b2}, C(O)NR_{b3}R_{b4}, C(O)-N(R_{b3})-OR_{b4}, C(S)R_{b3},

C(S)NR_{b3}R_{b4}, S(O)R_{b3}, SO₂R_{b3}, S(O)NR_{b3}R_{b4}, S(O)-N(R_{b3})-OR_{b4}, SO₂NR_{b3}R_{b4}, or SO₃R_{b3}; in which R_{b1} is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, 6-membered heteroaryl, fused heteroaryl, aryl, or NHCO₂R_{b5}; R_{b2} is H, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, aryl, heteroaryl, or C₁-C₁₅ alkyl optionally substituted with halogen, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, OR_{b5}, CO₂R_{b5}, or S(O)₂R_{b5}; and each of R_{b3} and R_{b4}, independently, is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, or aryl; R_{b5} being H, C₁-C₁₅ alkyl, heteroaryl, or aryl.

Referring to formula (I), a subset of the just-described compounds are those in which X is N(R_{a1}); or X and one of R₂ and R₃, together with the atom or atoms to which they are bonded, are C₃-C₂₀ heterocycloalkyl; R₁ is H, C(O)R_{b1}, CO₂R_{b2}, C(O)NR_{b1}R_{b2}, C(S)NR_{b1}R_{b2}, or SO₂R_{b1}, in which R_{b1} is H, aryl, or C₁-C₁₅ alkyl optionally substituted with halogen, aryl, or NHCO₂R_{b5}; and R_{b2} is aryl or C₁-C₁₅ alkyl optionally substituted with halogen, OR_{b5}, or CO₂R_{b5}; each of R₂ and R₃, independently, is H or C₁-C₁₅ alkyl optionally substituted with OR_{c1}, C(O)-NR_{c1}R_{c2}, NR_{c1}R_{c2}, N(R_{c1})-CO₂R_{c2}, N(R_{c1})-SO₂R_{c2}, or O-SO₂-R_{c1}; or X and one of R₂ and R₃, together with the atom or atoms to which they are bonded, are C₃-C₂₀ heterocycloalkyl; each of R₄ and R₅, independently, is H or C₁-C₁₅ alkyl; one of R₆ and R₇ is C₁-C₁₅ alkyl substituted with C₃-C₂₀ heterocycloalkyl, and the other of R₆ and R₇ is H; R₈ is H; and each of R₉ and R₁₀, independently, is H or CO₂R_{e1}. In these compounds, one of R₆ and R₇ can be





In still another aspect, this invention features a compound of formula (I) shown above except that R₁ is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, aryl, C(O)R_{b1}, CO₂R_{b2}, C(O)NR_{b2}R_{b3}, C(O)-N(R_{b2})-OR_{b3}, C(S)R_{b2}, C(S)NR_{b2}R_{b3}, S(O)R_{b2}, SO₂R_{b2}, S(O)NR_{b2}R_{b3}, S(O)-N(R_{b2})-OR_{b3}, SO₂NR_{b2}R_{b3}, or SO₃R_{b2}; in which R_{b1} is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, aryl, 6-membered heteroaryl, or fused heteroaryl; and each of R_{b2} and R_{b3}, independently, is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, aryl, or heteroaryl; and one of R₂ and R₃ is C₁-C₁₅ alkyl substituted with halogen, OR_{c1}, SR_{c1}, CO₂R_{c1}, OC(O)R_{c1},

C(O)NR_{c1}R_{c2}, SO₂R_{c1}, O-SO₂-R_{c1}, NR_{c1}R_{c2}, N(R_{c1})-C(O)R_{c2}, N(R_{c1})-CO₂R_{c2}, N(R_{c1})-SO₂R_{c2}, N(R_{c1})-C(O)-N(R_{c2}R_{c3}); the other of R₂ and R₃ is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, aryl, OR_{c1}, SR_{c1}, or NR_{c1}R_{c2}; in which each of R_{c1}, R_{c2}, and R_{c3}, independently, is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, or aryl.

Referring to formula (I), a subset of the just-described compounds are those in which X is N(R_{a1}) or O; or X and one of R₂ and R₃, together with the atom or atoms to which they are bonded, are C₃-C₂₀ heterocycloalkyl; R₁ is CO₂R_{b2}, in which R_{b2} is alkyl substituted with aryl; one of R₂ and R₃ is C₁-C₁₅ alkyl substituted with OR_{c1}, SR_{c1},

OC(O)R_{e1}, CO₂R_{c1}, C(O)NR_{c1}R_{c2}, SO₂R_{c1}, O-SO₂-R_{c1}, NR_{c1}R_{c2}, N(R_{c1})-C(O)R_{c2}, N(R_{c1})-CO₂R_{c2}, N(R_{c1})-SO₂R_{c2}, N(R_{c1})-C(O)-N(R_{c2}R_{c3}), and the other of R₂ and R₃ is H; or X and one of R₂ and R₃, together with the atom or atoms to which they are bonded, are C₃-C₂₀ heterocycloalkyl; each of R₄ and R₅, independently, is H or C₁-C₁₅ alkyl; one of R₆ and R₇ is C₁-C₁₅ alkyl substituted with C₃-C₂₀ heterocycloalkyl, and the other of R₆ and R₇ is H; R₈ is H; and each of R₉ and R₁₀, independently, is H, CN, C(O)R_{e1}, CO₂R_{e1}, or C(O)NR_{e1}R_{e2}; or R₉ and R₁₀, taken together, are C₃-C₂₀ heterocycloalkyl. In these




compounds, one of R₆ and R₇ can be ; and one of R₄ and R₅ can be H or C₁-C₁₅ alkyl optionally substituted with aryl, C₃-C₂₀ cycloalkyl, OR_{f1}, SR_{f1}, NR_{f1}R_{f2}, or C(O)NR_{f1}R_{f2}; in which each of R_{f1} and R_{f2}, independently, is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, or aryl.

In still another aspect, this invention features a compound of formula (I) shown above except that R₁ is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, aryl, C(O)R_{b1}, CO₂R_{b2}, C(O)NR_{b2}R_{b3}, C(O)-N(R_{b2})-OR_{b3}, C(S)R_{b2}, C(S)NR_{b2}R_{b3}, S(O)R_{b2}, SO₂R_{b2}, S(O)NR_{b2}R_{b3}, S(O)-N(R_{b2})-OR_{b3}, SO₂NR_{b2}R_{b3}, or SO₃R_{b2}; in which R_{b1} is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, aryl, 6-membered heteroaryl, or fused heteroaryl; and each of R_{b2} and R_{b3}, independently, is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, aryl, or heteroaryl; one of R₄ and R₅ is C₁-C₁₅ alkyl substituted with halogen, OR_{d1}, SR_{d1}, CO₂R_{d1}, C(O)NR_{d1}R_{d2}, SO₂R_{d1}, SO₃R_{d1}, or NR_{d1}R_{d2}; the other of R₄ and R₅ is H, halogen, C₁-C₁₅ alkyl, C₃-C₂₀


cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, or aryl; in which each of R_{d1} and R_{d2}, independently, is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, or aryl; R₈ is H, halogen, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, aryl, OR_{e1}, or SR_{e1}; in which R_{e1} is H and C₁-C₁₅ alkyl; and each of R₉ and R₁₀, independently, is H, halogen, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, aryl, C(O)R_{f1}, CO₂R_{f1}, C(O)NR_{f1}R_{f2}, C(O)-N(R_{f1})-OR_{f2}, C(S)R_{f1}, C(S)NR_{f1}R_{f2}, CN, NO₂, S(O)R_{f1}, SO₂R_{f1}, S(O)NR_{f1}R_{f2}, S(O)-N(R_{f1})-OR_{f2}, SO₂NR_{f1}R_{f2}, SO₃R_{f1}, PO(OR_{f1})(OR_{f2}), PO(R_{f1})(R_{f2}), PO(NR_{f1}R_{f2})(OR_{f3}), PO(NR_{f1}R_{f2})(NR_{f3}R_{f4}), C(O)-N(R_{f1})-NR_{f2}R_{f3}, or C(S)-N(R_{f1})-NR_{f2}R_{f3}; or R₈ and R₁₀, taken together, is C₃-C₂₀ cycloalkyl or C₃-C₂₀ heterocycloalkyl; or R₉ and R₁₀, taken together, is C₃-C₂₀ cycloalkyl or C₃-C₂₀ heterocycloalkyl; in which each of R_{f1}, R_{f2}, R_{f3}, and R_{f4}, independently, is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, or aryl; or any two of R_{f1}, R_{f2}, R_{f3}, and R_{f4}, together with the atom or atoms to which they are bonded, is C₃-C₂₀ heterocycloalkyl.

Referring to formula (I), a subset of the just-described compounds are those in which X is N(R_{a1}) or O; R₁ is CO₂R_{b2}, in which R_{b2} is alkyl substituted with aryl; each of R₂ and R₃, independently, is H or C₁-C₁₅ alkyl optionally substituted with aryl; one of R₄ and R₅ is C₁-C₁₅ alkyl substituted with SR_{d1} or C(O)NR_{d1}R_{d2}, and the other of R₄ and R₅ is H; one of R₆ and R₇ is C₁-C₁₅ alkyl substituted with C₃-C₂₀ heterocycloalkyl, and the other of R₆ and R₇ is H; R₈ is H; and each of R₉ and R₁₀, independently, is H or CO₂R_{f1}.



In these compounds, one of R₆ and R₇ can be

In a further aspect, this invention features a method for treating an infection with a virus. The method includes administering to a subject in need thereof an effective amount of a compound of formula (II):




In this formula, X is N(R_{a1}), O, or CH₂; or X and one of R₂ and R₃, together with the atom or atoms to which they are bonded, are C₃-C₂₀ heterocycloalkyl; in which R_{a1} is H

or C₁-C₁₅ alkyl; R₁ is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, aryl, C(O)R_{b1}, CO₂R_{b1}, C(O)NR_{b1}R_{b2}, C(O)-N(R_{b1})-OR_{b2}, C(S)R_{b1}, C(S)NR_{b1}R_{b2}, S(O)R_{b1}, SO₂R_{b1}, S(O)NR_{b1}R_{b2}, S(O)-N(R_{b1})-OR_{b2}, SO₂NR_{b1}R_{b2}, or SO₃R_{b1}; in which each of R_{b1} and R_{b2}, independently, is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, or aryl; each of R₂ and R₃, independently, is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, aryl, OR_{c1}, SR_{c1}, or NR_{c1}R_{c2}; or X and one of R₂ and R₃, together with the atom or atoms to which they are bonded, are C₃-C₂₀ heterocycloalkyl; in which each of R_{c1} and R_{c2}, independently, is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, or aryl; each of R₄ and R₅, independently, is H, halogen, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, or aryl; and one of R₆, R₇, and R₈ is C₁-C₁₅ alkyl substituted with C₃-C₂₀ heterocycloalkyl, heteroaryl, or aryl; and the others of R₆, R₇, and R₈, independently, is H, halogen, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, aryl, OR_{d1}, SR_{d1}, C(O)R_{d1}, CO₂R_{d1}, C(O)NR_{d1}R_{d2}, or C(O)-N(R_{d1})-OR_{d2}; in which R_{d1} and R_{d2}, independently, is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, or aryl; and the virus is a coronavirus or a hepatitis C virus.

In particular, this invention features a method for treating an infection with a severe acute respiratory syndrome virus, a human coronavirus 229E, or a hepatitis C virus, by administering to a subject in need thereof an effective amount of a compound of formula (II) shown above.

For example, one can administer to a subject infected with a virus (e.g., a severe acute respiratory syndrome virus) a compound of formula (II), in which X is N(R_{a1}) or O; R₁ is CO₂R_{b1}, in which R_{b1} is alkyl substituted with aryl; each of R₂ and R₃, independently, is H or C₁-C₁₅ alkyl optionally substituted with aryl, OR_{c1}, CO₂R_{c1}, N(R_{c1})-C(O)R_{c2}, or OC(O)-NR_{c1}R_{c2}; each of R₄ and R₅, independently, is H or C₁-C₁₅ alkyl; R₆ is C₁-C₁₅ alkyl substituted with C₃-C₂₀ heterocycloalkyl; R₇ is H; and R₈ is C(O)R_{d1}, CO₂R_{d1}, C(O)NR_{d1}R_{d2}, or C₁-C₁₅ alkyl optionally substituted with halogen or

OH. In these compounds, one of R₆ and R₇ can be



In a further aspect, this invention features a method for inhibiting a viral protease (e.g., a coronaviral 3CL protease or a hepatitis C viral protease) in a cell, by contacting the cell with an effective amount of a compound of formula (II) shown above. In particular, this invention features a method for inhibiting a severe acute respiratory syndrome viral 3CL protease, a human coronaviral 229E protease, or a hepatitis C viral protease.


In a further aspect, this invention features a compound of formula (II) shown above.

In still a further aspect, this invention features a method for treating an infection with a picornavirus (e.g., an enterovirus or a rhinovirus). The method includes administering to a subject in need thereof an effective amount of a compound of formula (I) or a compound of formula (II).

In still a further aspect, this invention features a method for inhibiting a picornaviral protease (e.g., an enteroviral protease or a rhinoviral 3C protease) in a cell. The method includes contacting the cell with an effective amount of a compound of formula (I) or a compound of formula (II).


In yet a further aspect, this invention features a chemical synthetic method. The

method includes reducing a compound of formula (III):



(III), to form an

alcohol, followed by reacting the alcohol with $\text{Ph}_3\text{P}=\text{COOR}$ in the presence of pyridine




sulfur trioxide to give a compound of formula (IV):

(IV), in which


P_1 is an amino-protecting group; P_2 is a carboxyl-protecting group; and R is C_1 - C_{15} alkyl, C_3 - C_{20} cycloalkyl, C_3 - C_{20} heterocycloalkyl, heteroaryl, or aryl. In formulas (III) and (IV), P_1 can be t-butoxycarbonyl, benzyloxycarbonyl, acetyl, phenylcarbonyl, or trialkylsilyl; P_2 can be C_1 - C_{15} alkyl; and R can be C_1 - C_{15} alkyl.

In particular, the chemical synthetic method can further include removing the amino-protecting group P_1 of the compound of formula (IV) to form a first de-protected intermediate, and then reacting the first de-protected intermediate with $P_3\text{HN-CH(R}_5\text{)-}$



COOH to give a compound of formula (V): (V), in which P_3

5 is an amino-protecting group and R_5 is H, halogen, $C_1\text{-}C_{15}$ alkyl, $C_3\text{-}C_{20}$ cycloalkyl, $C_3\text{-}C_{20}$ heterocycloalkyl, heteroaryl, or aryl. The method can further include removing the amino-protecting group P_3 of the compound of formula (V) to form a second de-protected intermediate, and then reacting the second de-protected intermediate with $R_1\text{HN-CH(R}_2\text{)-}$





COOH to give a compound of formula (VI): (VI), in

10 which R_1 is H, $C_1\text{-}C_{15}$ alkyl, $C_3\text{-}C_{20}$ cycloalkyl, $C_3\text{-}C_{20}$ heterocycloalkyl, heteroaryl, aryl, $C(O)R_{b1}$, CO_2R_{b1} , $C(O)NR_{b1}R_{b2}$, $C(O)\text{-}N(R_{b1})\text{-}OR_{b2}$, $C(S)R_{b1}$, $C(S)NR_{b1}R_{b2}$, $S(O)R_{b1}$, SO_2R_{b1} , $S(O)NR_{b1}R_{b2}$, $S(O)\text{-}N(R_{b1})\text{-}OR_{b2}$, $SO_2NR_{b1}R_{b2}$, or SO_3R_{b1} ; in which each of R_{b1} and R_{b2} , independently, is H, $C_1\text{-}C_{15}$ alkyl, $C_3\text{-}C_{20}$ cycloalkyl, $C_3\text{-}C_{20}$ heterocycloalkyl, heteroaryl, or aryl; and R_2 is H, $C_1\text{-}C_{15}$ alkyl, $C_3\text{-}C_{20}$ cycloalkyl, $C_3\text{-}C_{20}$ heterocycloalkyl, heteroaryl, aryl, OR_{c1} , SR_{c1} , or $NR_{c1}R_{c2}$; or X and one of R_2 and R_3 , together with the atom or atoms to which they are bonded, are $C_3\text{-}C_{20}$ heterocycloalkyl; in which each of R_{c1} and R_{c2} , independently, is H, $C_1\text{-}C_{15}$ alkyl, $C_3\text{-}C_{20}$ cycloalkyl, $C_3\text{-}C_{20}$ heterocycloalkyl, heteroaryl, or aryl. In formulas (V) and (VI), P_3 can be t-butoxycarbonyl, benzyloxycarbonyl, acetyl, phenylcarbonyl, or trialkylsilyl; and R can be

20 $C_1\text{-}C_{15}$ alkyl.

Alternatively, the chemical synthetic method mentioned above can further include removing the amino-protecting group P_1 of the compound of formula (IV) to form a first de-protected intermediate, and then reacting the first de-protected intermediate with a

compound of formula (VII):  (VII), to give a compound of

formula (VI):  (VI), in which R₁ is H, C₁-C₁₅ alkyl,


C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, aryl, C(O)R_{b1}, CO₂R_{b1},


C(O)NR_{b1}R_{b2}, C(O)-N(R_{b1})-OR_{b2}, C(S)R_{b1}, C(S)NR_{b1}R_{b2}, S(O)R_{b1}, SO₂R_{b1},

- 5 S(O)NR_{b1}R_{b2}, S(O)-N(R_{b1})-OR_{b2}, SO₂NR_{b1}R_{b2}, or SO₃R_{b1}; in which each of R_{b1} and R_{b2}, independently, is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, or aryl; R₂ is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, aryl, OR_{c1}, SR_{c1}, or NR_{c1}R_{c2}; or X and one of R₂ and R₃, together with the atom or atoms to which they are bonded, are C₃-C₂₀ heterocycloalkyl; in which each of R_{c1} and R_{c2},
- 10 independently, is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, or aryl; and R₅ is H, halogen, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, or aryl.


In addition, the chemical synthetic method can further include removing the amino-protecting group P₁ of the compound of formula (IV) to form a first de-protected intermediate, and then reacting the first de-protected intermediate with a compound of

15

formula (VIII):  (VIII), in which n is 1, 2, or 3; P₄ is an amino-protecting group; and R₅ is H, halogen, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, or aryl; followed by removing the amino-protecting group

P₄ to give a compound of formula (IX):  (IX). The

20 compound of formula (IX) can further react with R₁'COOH or R₁'COCl to give a



compound of formula (X):

(X), in which R_1' is

H, C_1-C_{15} alkyl, C_3-C_{20} cycloalkyl, C_3-C_{20} heterocycloalkyl, heteroaryl, aryl, or OR_{b1} ; R_{b1} being H, C_1-C_{15} alkyl, C_3-C_{20} cycloalkyl, C_3-C_{20} heterocycloalkyl, heteroaryl, or aryl. In formulas (VIII), (IX), and (X), P_4 can be t-butoxycarbonyl, benzyloxycarbonyl, acetyl, phenylcarbonyl, or trialkylsilyl; and R can be C_1-C_{15} alkyl.

In addition, this invention encompasses a pharmaceutical composition that contains an effective amount of at least one of the above-mentioned compounds and a pharmaceutically acceptable carrier.


The compounds of the invention include the compounds themselves, as well as their salts, prodrugs, and solvates, if applicable. A salt, for example, can be formed between an anion and a positively charged group (e.g., amino) on a compound of the invention. Examples of suitable anions include chloride, bromide, iodide, sulfate, nitrate, phosphate, citrate, methanesulfonate, trifluoroacetate, and acetate. Likewise, a salt can also be formed between a cation and a negatively charged group (e.g., carboxylate) on a compound of the invention. Examples of suitable cations include sodium ion, potassium ion, magnesium ion, calcium ion, and an ammonium cation such as tetramethylammonium ion. Examples of prodrugs include esters and other pharmaceutically acceptable derivatives, which, upon administration to a subject, are capable of providing active compounds of the invention. A solvate refers to a complex formed between an active compound of the invention and a pharmaceutically acceptable solvent. Examples of pharmaceutically acceptable solvents include water, ethanol, isopropanol, ethyl acetate, acetic acid, and ethanolamine.

Also within the scope of this invention is a composition containing one or more of the compounds described above for use in treating an infection with a virus, and the use of such a composition for the manufacture of a medicament for the just-mentioned treatment.


The details of one or more embodiments of the invention are set forth in the description below. Other features, objects, and advantages of the invention will be apparent from the description and from the claims.

DETAILED DESCRIPTION


Shown below are the structures of compounds 1-145, exemplary compounds of this invention:




Compound 1




Compound 2




Compound 3




Compound 4




Compound 5




Compound 6




Compound 7




Compound 8




Compound 9




Compound 10




Compound 11




Compound 12




Compound 13




Compound 14




Compound 15




Compound 16




Compound 17




Compound 18




Compound 19




Compound 20




Compound 21




Compound 22




Compound 23




Compound 24




Compound 25




Compound 26




Compound 27




Compound 28




Compound 29




Compound 30




Compound 31




Compound 32




Compound 33




Compound 34




Compound 35




Compound 36




Compound 37




Compound 38




Compound 39




Compound 40




Compound 41




Compound 42




Compound 43




Compound 44




Compound 45




Compound 46




Compound 47




Compound 48




Compound 49




Compound 50




Compound 51




Compound 52




Compound 53




Compound 54




Compound 55




Compound 56




Compound 57




Compound 58




Compound 59




Compound 60




Compound 61




Compound 62




Compound 63




Compound 64




Compound 65




Compound 66




Compound 67




Compound 68




Compound 69




Compound 70




Compound 71




Compound 72




Compound 73




Compound 74




Compound 75




Compound 76




Compound 77




Compound 78




Compound 79




Compound 80




Compound 81




Compound 82




Compound 83




Compound 84




Compound 85




Compound 86




Compound 87




Compound 88




Compound 89




Compound 90




Compound 91




Compound 92




Compound 93




Compound 94




Compound 95




Compound 96




Compound 97




Compound 98




Compound 99




Compound 100




Compound 101




Compound 102




Compound 103




Compound 104




Compound 105




Compound 106




Compound 107




Compound 108




Compound 109




Compound 110




Compound 111




Compound 112




Compound 113




Compound 114




Compound 115




Compound 116




Compound 117




Compound 118




Compound 119




Compound 120




Compound 121




Compound 122




Compound 123




Compound 124




Compound 125




Compound 126




Compound 127




Compound 128




Compound 129




Compound 130




Compound 131




Compound 132




Compound 133




Compound 134




Compound 135




Compound 136




Compound 137




Compound 138




Compound 139




Compound 140




Compound 141




Compound 142



Compound 143




Compound 144



Compound 145


The compounds described above can be prepared by methods well known to a skilled person in the art. For example, Schemes I-IV shown below depict typical synthetic routes for preparing exemplary compounds. In these schemes, R₁, R₁', R₂, and R₅ are as defined in the Summary section above. Details of preparation of these compounds are provided in Examples 1-145.

Scheme I:




5

Scheme II:



10


Scheme III:



5

10

Scheme IV:



As shown in Scheme I, L-glutamic acid can first be protected with *t*-butoxycarbonyl and methyl. The protected L-glutamic acid (intermediate 1) can then react with bromoacetonitrile and followed by a ring closure reaction to form an amino acid derivative that containing a 5-membered cyclic lactam (intermediate 3).

5 Intermediate 3 can subsequently be transformed to intermediate 5 that includes an additional double bond. Intermediate 5 thus formed can sequentially couple with two amino acid derivatives to prepare a compound of formula (I). Reagents other than those shown in Scheme I can also be used. For example, intermediate 5 mentioned above can couple with an acid containing hydroxyl, before coupling to an amino acid to prepare a
10 compound of formula (I) in which X is oxygen.

As shown in Scheme II, intermediate 5 can also couple with amino acid derivatives that contain a pyrrolidinone moiety to form certain compounds of formula (IX). The compounds thus obtained can be further modified (e.g., by reacting with an acid or an acyl chloride) to obtain compounds of formula (X).

15 As shown in Scheme III, intermediate 5 can also be hydrolyzed to form an acid, which in turn can react with O,N-dimethyl-hydroxylamine to form an amide. The amide can either sequentially couple with two amino acid derivatives or couple with a dipeptide derivative to form certain compounds of formula (I). The compounds thus obtained can further react with Grignard reagents to form other compounds of formula (I).

20 As shown in Scheme IV, intermediate 3 mentioned above can also sequentially couple with two amino acid derivatives (or couple with a dipeptide derivative) to form certain compounds of formula (I). The compounds thus obtained can be further reduced to form alcohols or oxidized to form acids. The alcohols can be further halogenated to form halides or oxidized to form aldehydes. The aldehydes can subsequently undergo
25 either Wittig reactions or addition-elimination reactions to form certain compounds of formula (II). The acids just-mentioned can undergo, e.g., esterification reactions, Grignard reactions, or amidation reactions to form other compounds of formula (II).

A compound synthesized by the methods described above can be purified by a
30 known method, such as column chromatography, high-pressure liquid chromatography, or recrystallization.

Other compounds of the invention can be prepared using other suitable starting materials following the synthetic routes disclosed herein and/or other synthetic methods known in the art. The methods described above may also additionally include steps, either before or after the steps described specifically herein, to add or remove suitable protecting groups in order to ultimately allow synthesis of the compounds of the invention. In addition, various synthetic steps may be performed in an alternate sequence or order to give the desired compounds. Synthetic chemistry transformations and protecting group methodologies (protection and deprotection) useful in synthesizing applicable compounds of the invention are known in the art and include, for example, those described in R. Larock, *Comprehensive Organic Transformations*, VCH Publishers (1989); T.W. Greene and P.G.M. Wuts, *Protective Groups in Organic Synthesis*, 2nd Ed., John Wiley and Sons (1991); L. Fieser and M. Fieser, *Fieser and Fieser's Reagents for Organic Synthesis*, John Wiley and Sons (1994); and L. Paquette, ed., *Encyclopedia of Reagents for Organic Synthesis*, John Wiley and Sons (1995) and subsequent editions thereof.

The compounds mentioned herein may contain a non-aromatic double bond and one or more asymmetric centers. Thus, they can occur as racemates and racemic mixtures, single enantiomers, individual diastereomers, diastereomeric mixtures, and cis- or trans- isomeric forms. All such isomeric forms are contemplated.

Also within the scope of this invention is a pharmaceutical composition contains an effective amount of at least one compound described above and a pharmaceutical acceptable carrier. Further, this invention covers a method of administering an effective amount of one or more of the compounds of the invention to a patient having an infection with a coronavirus. Effective doses will vary, as recognized by those skilled in the art, depending on the types of diseases treated, route of administration, excipient usage, and the possibility of co-usage with other therapeutic treatment.

To practice the method of the present invention, a composition having one or more compounds of the invention can be administered parenterally, orally, nasally, rectally, topically, or buccally. The term "parenteral" as used herein refers to subcutaneous, intracutaneous, intravenous, intramuscular, intraarticular, intraarterial,

intrasynovial, intrasternal, intrathecal, intralesional, or intracranial injection, as well as any suitable infusion technique.

A sterile injectable composition can be a solution or suspension in a non-toxic parenterally acceptable diluent or solvent, such as a solution in 1,3-butanediol. Among
5 the acceptable vehicles and solvents that can be employed are mannitol, water, Ringer's solution, and isotonic sodium chloride solution. In addition, fixed oils are conventionally employed as a solvent or suspending medium (e.g., synthetic mono- or diglycerides). Fatty acid, such as oleic acid and its glyceride derivatives are useful in the preparation of injectables, as are natural pharmaceutically acceptable oils, such as olive oil or castor oil,
10 especially in their polyoxyethylated versions. These oil solutions or suspensions can also contain a long chain alcohol diluent or dispersant, carboxymethyl cellulose, or similar dispersing agents. Other commonly used surfactants such as Tweens or Spans or other similar emulsifying agents or bioavailability enhancers which are commonly used in the manufacture of pharmaceutically acceptable solid, liquid, or other dosage forms can also
15 be used for the purpose of formulation.

A composition for oral administration can be any orally acceptable dosage form including capsules, tablets, emulsions and aqueous suspensions, dispersions, and solutions. In the case of tablets, commonly used carriers include lactose and corn starch. Lubricating agents, such as magnesium stearate, are also typically added. For oral
20 administration in a capsule form, useful diluents include lactose and dried corn starch. When aqueous suspensions or emulsions are administered orally, the active ingredient can be suspended or dissolved in an oily phase combined with emulsifying or suspending agents. If desired, certain sweetening, flavoring, or coloring agents can be added.

A nasal aerosol or inhalation composition can be prepared according to techniques
25 well known in the art of pharmaceutical formulation. For example, such a composition can be prepared as a solution in saline, employing benzyl alcohol or other suitable preservatives, absorption promoters to enhance bioavailability, fluorocarbons, and/or other solubilizing or dispersing agents known in the art. A composition having one or more compounds of the invention can also be administered in the form of suppositories
30 for rectal administration.

The carrier in the pharmaceutical composition must be "acceptable" in the sense that it is compatible with the active ingredient of the composition (and preferably, capable of stabilizing the active ingredient) and not deleterious to the subject to be treated. One or more solubilizing agents can be utilized as pharmaceutical excipients for delivery of a compound of the invention. Examples of other carriers include colloidal silicon oxide, magnesium stearate, cellulose, sodium lauryl sulfate, and D&C Yellow # 10.

The compounds of this invention can be preliminarily screened for their efficacy in treating an infection with a virus by an *in vitro* assay (See Example 146 below) and then confirmed by animal experiments and clinic trials. Other methods will also be apparent to those of ordinary skill in the art.

The specific examples below are to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever. Without further elaboration, it is believed that one skilled in the art can, based on the description herein, utilize the present invention to its fullest extent. All publications cited herein are hereby incorporated by reference in their entirety.

Example 1: Preparation of compound 1

TMSCl (190 mL, 4.4 eq) was slowly added to a stirred suspension of L-glutamic acid (50.0 g, 1 eq.) in dry MeOH (1100 mL, 0.3 M) surrounded by an ice-cold bath.

After the addition was complete, the ice-cold bath was removed and the reaction was stirred overnight until TLC showed completed conversion. Then, Et₃N (306 mL, 6.5 eq) and (Boc)₂O (82 g, 1.1 eq) were added sequentially to the above reaction mixture. The reaction mixture was then stirred until TLC showed complete protection. The solvent was removed under reduced pressure. The residue was then filtered and washed with Et₂O using a pad of celite. The organic layers were combined and concentrated. The resulting crude product was purified by silica gel column chromatography to afford intermediate 1, N-Boc-L-(+)-glutamic acid dimethyl ester, (88 g, 95%) as an oil: ¹H NMR (CDCl₃) δ 1.40 (s, 9H), 1.91 (m, 1H), 2.14 (m, 2H), 2.37 (m, 2H), 3.64 (s, 3H), 3.70 (s, 3H), 4.29 (br, s, 1H). ESI-MS (M+H⁺) = 276.

To a solution of intermediate 1 (20 g, 1 eq.) in THF(50 mL) was added dropwise a solution of lithium hexamethyldisilazide (2.2 eq.) in THF (250 mL) at -78°C under

nitrogen atmosphere. The resulting mixture was stirred at -78°C for another 1.5 hours. Bromoacetonitrile (13 g, 1.5 eq.) was added dropwise to the above solution over a period of 1 hour while maintaining the temperature below -70°C using a cooling bath. The reaction mixture was stirred at -78°C for additional for 1-2 hours and the disappearance
5 of the starting material was confirmed by TLC analysis. The reaction was then quenched with pre-cooled methanol (10 mL) stirred for 10 minutes. The resulting methoxide was then quenched with a pre-cooled acetic acid in THF solution (9 mL HOAc/60 mL THF). After stirred for another 10 minutes, the cooling bath was removed and replaced with water bath. The reaction mixture was allowed to warm up to $0\pm 5^{\circ}\text{C}$ and then poured into
10 brine solution (10 g of NaCl in 100 mL water) in a 1L extractor. The organic layer was separated and concentrated to afford a dark brown oil. Silica gel (25 g) and methylene chloride (60 mL) were added to the Rotovap flask and spun on a Rotovap for 1 hour without heat and vacuum. The slurry was then filtered and wash with another batch of methylene chloride (100 mL). The light brown filtrate was collected, concentrated, and
15 purified by silica gel column chromatography to afford intermediate 2 (19 g), 2-tert-butoxycarbonyl-amino-4-cyanomethyl-pentanedioic acid dimethyl ester. ^1H NMR (CDCl_3) δ 1.42 (s, 9H), 2.10-2.17 (m, 2H), 2.77-2.90 (m, 3H), 3.73 (s, 3H), 3.74 (s, 3H), 4.32-4.49 (m, 1H), 5.12 (d, $J = 6.0$ Hz, 1H). ESI-MS ($\text{M}+\text{H}^+$) = 315.

Intermediate 2 (10 g) was dissolved in HOAc (240 mL) and shaken with 10%
20 Pd/C (20 g) under H_2 gas (70 psi) for 2 days. The mixture was filtered over celite. The filtrate was evaporated under reduced pressure and the residue was repeatedly evaporated from methyl tert-butyl ether to yield a light pink solid. The crude product was dissolved in THF and then Et_3N (20 mL) was added to the solution. The resulting mixture was stirred at 60°C overnight. The reaction was quenched with by addition of H_2O (50 mL).
25 The organic layer was separated and the aqueous layer was further extracted with methylene chloride. The organic layers were then combined, concentrated, and purified by silica gel column chromatography to afford intermediate 3, 2-tert-butoxycarbonylamino-3-(2-oxo-pyrrolidin-3-yl)-propionic acid methyl ester. ^1H NMR (CDCl_3) δ 1.37 (s, 9H), 1.75-1.80 (m, 2H), 2.04-2.09 (m, 1H), 2.39-2.42 (m, 1H), 3.25-
30 3.29 (m, 2H), 3.67 (s, 3H), 4.23-4.26 (m, 1H), 5.47 (d, $J = 8.0$ Hz, 1H), 6.29 (s, 1H). ESI-MS ($\text{M}+\text{H}^+$) = 287.

Intermediate 3 (6.0 g, 18.4 mmol) was solved in THF (200 mL) surrounded by an ice bath, followed by addition of 2.0 M LiBH₄/THF (46 ml, 5.0 eq). The ice bath was then removed and the mixture was stirred for 2 hours at room temperature. To the mixture was sequentialled added water (200 mL), ethyl acetate (200 mL), and MgSO₄ (400 g). Then MgSO₄ was removed and the aqueous layer was extracted with ethyl acetate. The organic layers were combined and concentrated to afford intermediate 4, [2-hydroxy-1-(2-oxo-pyrrolidin-3-ylmethyl)-ethyl]-carbamic acid tert-butyl ester, as a white solid (5.2 g, 95%). ESI-MS (M+H⁺) = 259.

Triethylamine (0.7 mL) was added to a solution of intermediate 4 (0.59 g, 2.28 mmol, 1 eq.) in methylsulfoxide (10.5 mL). The resulting solution was cooled to 15 °C using an ice-water bath and then sulfur trioxide-pyridine complex (1.8g, 5eq.) was added. The reaction was stirred at that temperature for 1 hour.

(Carboethoxymethylenetriphenyl)-phosphorane (2.4 g, 3 eq.) was added and the reaction was stirred at ambient temperature for another 3 hours. The reaction was then quenched by saturated brine (150 mL) and extracted with ethyl acetate (3x50 mL). The organic layers were combined, dried over anhydrous MgSO₄, filtered, and concentrated to afford a dark red oil. The oil was purified though column chromatography (50% ethyl acetate in hexane) to afford intermediate 5, 4-tert-butoxycarbonylamino-5-(2-oxo-pyrrolidin-3-yl)-pent-2-enoic acid ethyl ester, as a white solid (0.34 g, 45.7%). ¹H NMR (CDCl₃) δ 1.21 (t, 3H, J=7.2), 1.36 (s, 9H), 1.48-1.57 (m, 1H), 1.66-1.79 (m, 1H), 1.90-1.97(m, 1H), 2.38-2.47 (m, 2H), 3.26-3.29 (m, 2H), 4.10(q, 2H, J=6.9), 4.27(s, br, 1H), 5.46(d, 1H, J=7.5), 5.87 (d, 1H, J=15.6), 6.78 (dd, 1H, J=15.3, J=5.4), 6.98(s, br, 1H). ESI-MS (M+H⁺) = 577.

Intermediate 5 (100 mg, 0.3 mmol) was added to a solution of HCl in 1,4-dioxane (4.0 M, 3 mL) and the solution was stirred at room temperature for 30 minutes. The resulting solution was concentrated by removing 1,4-dioxane. CH₂Cl₂ (3 mL) was then added to the residue thus obtained and the solution was cooled down to 0-5 °C. N-Methylmorpholine (0.13 mL, 4 eq.) was then added and the mixture was stirred for 10 minutes to form solution (a). Boc-L-Leu-OH (71 mg, 0.3 mmol) was mixed with 1,1-ethyl-3-(3-dimethylaminopropyl)carbodiimide (EDC, 70 mg, 1.2 eq.) and N-hydroxybenzotriazole (HOBT, 50 mg, 1.2 eq.) in CH₂Cl₂. The mixture was stirred for 20

minutes to form solution (b). Solution (a) was then added to solution (b) and the mixture was stirred at room temperature for 2 hours. The reaction residue was added with brine (25 mL) and extracted with ethyl acetate (3x15 mL). The organic layers were combined, dried over anhydrous MgSO₄, and concentrated. The residue was purified by flash column chromatography (3% MeOH in CH₂Cl₂) to afford intermediate 6, 4-(2-tert-butoxycarbonylamino-4-methyl-pentanoylamino)-5-(2-oxo-pyrrolidin-3-yl)-pent-2-enoic acid ethyl ester, as a white solid. ¹H NMR (CDCl₃) δ 0.94 (d, 6H, J=5.1 Hz), 1.26 (t, 3H, J=7.2 Hz), 1.43 (s, 9H), 1.66-1.73 (m, 5H), 2.41 (br s, 1H), 3.33 (d, 2H, J=8.1 Hz), 4.16 (q, 2H, J=6.9 Hz), 4.58 (br, s, 1H), 5.03 (br, s, 1H), 5.9 (d, 1H, J=15.6 Hz), 6.82 (dd, 1H, J=15.3 Hz, 5.1 Hz), 7.50 (br, s, 1H). ESI-MS (M+H⁺) = 440.

Intermediate 7 (compound 6) was prepared in 60% yield from 2-benzyloxycarbonyl-amino-3-tert-butoxy-butyric acid and intermediate 6 using the procedure similar to that described in the preceding paragraph. ¹H NMR (CDCl₃) δ 0.94-0.98 (m, 6H), 1.07 (d, 3H, J=6.3), 1.27 (s, 9H), 1.66-1.73 (m, 5H), 2.21-2.50 (m, 2H), 3.20-3.30 (m, 2H), 4.16 (q, 2H, J=6.9 Hz), 4.42 (br, s, 1H), 4.58 (br, s, 1H), 5.10 (s, 2H), 5.9 (d, 1H, J=15.6 Hz), 6.82 (dd, 1H, J=15.3 Hz, 5.1 Hz), 7.2-7.34 (m, 4H), 7.60 (d, 1H, J=7.5 Hz). ESI-MS (M+H⁺) = 631.

Compound 1 was prepared by treating intermediate 7 with trifluoroacetic acid (2 mL). ¹H NMR (CDCl₃) δ 0.89-0.93 (m, 6H), 1.13-1.15 (m, 3H), 1.22-1.27 (m, 3H), 1.55-1.76 (m, 4H), 1.95-2 (m, 1H), 2.03-2.46 (m, 2H), 3.28-3.30 (m, 2H), 4.11-4.18 (m, 3H), 4.33 (br s, 1H), 4.54-4.56 (br s, 2H), 5.07 (s, 2H), 5.84-5.94 (d, 1H, J=15.9 Hz), 6.03-6.06 (d, J= 7.2 Hz), 6.76-6.83 (dd, 1H, J=15.0 Hz, 5.4 Hz), 7.31 (br s, 5H), 8.02 (br s, 1H). ESI-MS (M+H⁺) = 575.

Example 2: Preparation of compound 2

Compound 2 was prepared in a manner similar to that described in Example 1.

¹H NMR (CDCl₃) δ 7.92 (d, J = 7.5 Hz, 1H), 7.24-7.38 (m, 5H), 6.78 (dd, J = 15, 2.7 Hz, 1H), 5.98 (d, J = 7.8 Hz, 1H), 5.89 (d, J = 15 Hz, 1H), 5.09 (d, J = 4.2 Hz, 1H), 5.07 (s, 2H), 4.60-4.78 (m, 1H), 4.43-4.53 (m, 2H), 4.21-4.38 (m, 1H), 4.17 (q, J = 5.1 Hz, 2H), 3.82-3.88 (m, 1H), 3.40-3.71 (m, 2H), 3.29 (m, 2H), 1.40-2.11 (m, 5H), 1.25 (t, J = 5.1 Hz), 0.90 (br s, 6H).

ESI-MS ($M+H^+$) = 561.

Example 3: Preparation of compound 3

Compound 3 was prepared in a manner similar to that described in Example 1.

5 1H NMR($CDCl_3$) δ 0.91(s, br, 6H), 1.27(t, 3H, $J=6.0$), 1.60-1.66(m, 5H), 1.97-2.44(m, 7H), 3.26(m, 2H), 4.17(q, 2H, $J=6.9$), 4.24(m, 1H), 4.60(m, 1H), 5.06(s, 2H), 5.92(d, 1H, $J=15.3$), 6.12-6.18(m, 2H), 6.62-6.70(m, 1H), 6.83(dd, 1H, $J=15.3$, 5.), 7.31(s, br, 5H), 7.56(m, 1H), 8.01(m, 1H).

ESI-MS ($M+Na^+$) = 623.9.

10

Example 4: Preparation of compound 4

Compound 4 was prepared in a manner similar to that described in Example 1.

1H NMR ($CDCl_3$) δ 7.82 (d, $J = 8.0$ Hz, 1H), 6.90 (d, $J = 7.5$ Hz, 1H), 6.82 (dd, $J = 15$, 5.4 Hz, 1H), 5.89 (m, 4H), 5.09 (m, 2H), 4.40-4.68 (m, 2H), 4.16 (q, $J = 7.5$ Hz, 2H), 3.32 (m, 2H), 2.03-2.51 (m, 2H), 1.50-2.03 (m, 10H), 1.43 (s, 9H), 1.69 (t, $J = 7.5$ Hz, 3H), 0.94 (d, $J = 6.0$ Hz, 3H), 0.92 (d, $J = 6.0$ Hz, 3H).

15

ESI-MS ($M+H^+$) = 659.

Example 5: Preparation of compound 5

Compound 5 was prepared in a manner similar to that described in Example 1.

20 1H NMR (CD_3OD) δ 0.86-1.04 (m, 6H), 1.23 (s, br, 3H), 1.60-1.71 (m, 6H), 2.06-2.85 (m, 6H), 3.38 (m, 2H), 4.10-4.28 (m, 2H), 4.34-4.45 (m, 1H), 4.60-4.72 (m, 1H), 5.13 (s, 2H), 5.91 (d, 1H, $J=15.9$), 6.82 (dd, 1H, $J=15.9$, 4.7), 7.27-7.54 (m, 5H).

ESI-MS ($M+H^+$) = 589.

25

Example 6: Preparation of compound 6

Compound 6 was prepared in a manner identical to that of intermediate 7 described in Example 1.

30 1H NMR ($CDCl_3$) δ 0.94-0.98 (m, 6H), 1.07 (d, 3H, $J=6.3$), 1.27 (s, 9H), 1.66-1.73 (m, 5H), 2.21-2.50 (m, 2H), 3.20-3.30 (m, 2H), 4.16 (q, 2H, $J=6.9$ Hz), 4.42 (br, s,

1H), 4.58 (br, s, 1H), 5.10 (s, 2H), 5.9 (d, 1H, J=15.6 Hz), 6.82 (dd, 1H, J=15.3 Hz, 5.1 Hz), 7.2-7.34 (m, 4H), 7.60 (d, 1H, J=7.5 Hz).

ESI-MS (M+H⁺) = 631.

5 Example 7: Preparation of compound 7

Compound 7 was prepared in a manner similar to that described in Example 1.

¹H NMR (CDCl₃) δ 0.86-0.87 (d, 6H, J=4.2), 1.09 (s, 9H), 1.18-1.22 (t, 3H, J=7.2), 1.66 (m, 5H), 2.30 (s, br, 2H), 3.21-3.24 (d, 2H, J=7.2), 3.35-3.40 (t, 1H, J=6.6), 3.74-3.76 (d, 1H, J=6.9), 4.10 (q, 2H, J=7.2), 4.16 (s, br, 1H), 4.51 (s, br, 2H), 5.03-5.05 (d, 2H, J=4.8), 5.64-5.65 (d, 1H, J=15.9), 6.72-6.77 (dd, 1H, J=4.5), 6.93-6.96 (d, 1H, J=6.9), 7.28 (s, br, 5H), 7.60 (s, br, 1H).

ESI-MS (M+H⁺) = 616.

Example 8: Preparation of compound 8

15 Compound 8 was prepared in a manner similar to that described in Example 1.

¹H NMR (CDCl₃) δ 7.74 (d, J = 7.5 Hz, 1H), 7.24-7.33 (m, 10H), 7.05 (d, J = 8.5 Hz, 1H), 6.80 (dd, J = 15.5, 5.5 Hz, 1H), 6.17 (s, 1H), 5.90 (d, J = 15.5 Hz, 1H), 5.73 (d, J = 6.5 Hz, 1H), 5.10 (s, 2H), 4.47-4.62 (m, 4H), 4.27-4.29 (m, 1H), 4.18-4.20 (m, 3H), 3.24-3.27 (m, 2H), 2.23-2.48 (m, 2H), 1.90-2.14 (m, 1H), 1.89 (s, 2H), 1.47-1.73 (m, 3H), 1.26 (t, J = 3.6 Hz, 3H), 1.21 (d, J = 10.2 Hz, 3H), 0.87 (d, J = 8.0 Hz, 3H), 0.85 (d, J = 8.0 Hz, 3H).

ESI-MS (M+H⁺) = 665.

Example 9: Preparation of compound 9

25 Compound 9 was prepared in a manner similar to that described in Example 1.

ESI-MS (M+H⁺) = 639.

Example 10: Preparation of compound 10

Compound 10 was prepared in a manner similar to that described in Example 1.

30 ¹H NMR (CDCl₃) δ 0.88-0.98 (m, 6 H), 1.19-1.86 (m, 15H), 2.0-2.08 (m, 1H), 2.15-2.39 (m, 2H), 2.47-2.59 (m, 1H), 2.94 (m, 2H), 3.98-4.20 (m, 3H), 4.30-4.36 (m,

1H), 4.60-4.64 (m, 1H), 5.09 (m, 2H), 5.89 (d, J = 15.6, 1H), 6.89 (dd, 1H, 15.6, 4.8),
7.26-7.33 (m, 5H).

ESI-MS (M+H⁺) = 639.

5 Example 11: Preparation of compound 11

Compound 11 was prepared in a manner similar to that described in Example 1.

¹H NMR (CDCl₃) δ 7.90 (d, J = 7.0 Hz, 1H), 7.31 (s, 5H), 6.77-6.85 (m, 2H), 6.41
9 (br, s, 1H), 5.90 (d, J = 15 Hz, 1H), 5.78 (d, J = 7 Hz, 1H), 5.07 (s, 2H), 4.87 (br, s,
1H), 4.41-4.62 (m, 2H), 4.07-4.14 (m, 4H), 3.21-3.37 (m, 2H), 2.94-3.18 (m, 2H), 2.28-
10 2.48 (m, 3H), 1.28-2.13 (m, 10H), 1.39 (s, 9H), 1.26 (t, 4.5 Hz, 3H), 0.92 (d, J = 6.3 Hz,
3H), 0.90 (d, J = 6.3 Hz, 3H)

ESI-MS (M+H⁺) = 702.

Example 12: Preparation of compound 12

15 Compound 12 was prepared in a manner similar to that described in Example 1.

ESI-MS (M+H⁺) = 680.

Example 13: Preparation of compound 13

Compound 13 was prepared in a manner similar to that described in Example 1.

20 ¹H NMR (CDCl₃) δ 7.94 (d, J = 7.0 Hz, 1H), 7.31 (s, 5H), 6.81 (dd, J = 15, 5 Hz,
1H), 6.38 (br s, 1H), 5.84-5.93 (m, 2H), 5.31 (br s, 1H), 5.07 (s, 2H), 4.41-4.64 (m, 2H),
4.11-4.16 (m, 4H), 3.62-3.64 (m, 2H), 3.60 (s, 3H), 3.43-3.51 (m, 2H), 3.23-3.29 (m,
2H), 3.06-3.20 (m, 2H), 2.28-2.48 (m, 2H), 1.20-1.96 (m, 8H), 1.26 (t, J = 4.5 Hz, 3H),
0.92 (d, J = 6.5 Hz, 3H), 0.90 (d, J = 6.5 Hz, 3H).

25 ESI-MS (M+H⁺) = 660.

Example 14: Preparation of compound 14

Compound 14 was prepared in a manner similar to that described in Example 1.

30 ¹H NMR (CDCl₃) δ 0.93 (m, 6H), 1.25-1.30 (m, 3H), 1.47-1.98 (m, 5H), 2.20-
2.60 (m, 5H), 2.93 (3, 3H), 3.11-3.45 (m, 4H), 4.12-4.19 (m, 2H), 4.45-4.48 (m, 3H),

5.09 (s, 2H), 5.86-5.92 (d, 1H, J=15.9 Hz), 6.75-6.82 (dd, 1H, J=15.9 Hz, 5.1 Hz), 7.33-7.36 (m, 5H), 8.12 (br, s, 1H), 8.23 (br, s, 1H).

ESI-MS (M+H⁺) = 637.

5 Example 15: Preparation of compound 15

Compound 15 was prepared in a manner similar to that described in Example 1.

ESI-MS (M+H⁺) = 631.

Example 16: Preparation of compound 16

10 Compound 16 was prepared in a manner similar to that described in Example 1.

¹H NMR (CDCl₃) δ 0.81-0.86 (m, 6H), 1.11-1.22 (m, 5H), 1.29-1.34 (m, 2H), 1.55-1.77 (m, 9H), 2.37 (s, br, 3H), 2.93 (s, 2H), 3.25 (s, br, 2H), 3.62-3.64 (m, 1H), 4.10 (q, 2H, J=6.6), 4.28-4.48 (m, 3H), 4.93-5.07 (m, 2H), 5.83 (d, 1H, J=15), 5.99 (s, 1H), 6.75 (dd, 1H, J=16.5, 6.3), 7.26 (s, br, 5H), 7.82 (m, 1H), 8.19 (s, 1H).

15

Example 17: Preparation of compound 17

Compound 17 was prepared in a manner similar to that described in Example 1.

¹H NMR (CD₃OD) δ 0.92-1.00 (m, 6H), 1.18-1.33 (m, 15H), 1.58-1.82 (m, 6H), 2.02-2.35 (m, 4H), 3.20-3.32 (m, 2H), 4.14-4.32 (m, 4H), 4.62-4.66 (m, 1H), 5.07-5.13 (m, 2H), 5.29 (d, 1H, J=5.4), 5.92 (d, 1H, J=15.6), 6.88 (dd, 1H, J=15.6, 5.7), 7.29-7.36 (m, 5H).

20

ESI-MS (M+H⁺) = 632.

Example 18: Preparation of compound 18

25 Compound 18 was prepared in a manner similar to that described in Example 1.

¹H NMR (CDCl₃) δ 0.90-0.92 (m, 12H), 3.65 (t, 3H, J=6.9), 1.50-1.61 (m, 4H), 1.77 (m, 1H), 2.01-2.07 (m, 3H), 2.37 (m, 2H), 3.27-3.36 (m, 2H), 3.65 (t, 2H, J=5.4), 3.96 (t, 1H, J=6.9), 4.15 (q, 2H, J=6.9), 4.30 (t, 2H, J=5.4), 4.60 (m, 2H), 5.60 (d, 1H, J=8.4), 5.89 (d, 1H, J=15.6), 6.64 (s, 1H), 6.80 (dd, 1H, J=15.6), 7.07 (d, 1H, J=7.8), 7.76 (d, 1H, J=7.5).

30

ESI-MS (M+H⁺) = 545.

Example 19: Preparation of compound 19

Compound 19 was prepared in a manner similar to that described in Example 1.

$^1\text{H NMR}$ (CDCl_3) δ 0.84-0.89 (m, 12H), 1.15-1.22 (m, 6H), 1.45-1.57 (m, 3H),
1.75 (s, br, 3H), 1.94-2.09 (m, 2H), 2.32 (s, br, 2H), 3.27 (d, 2H, $J=7.8$), 3.87 (t, 1H,
5 $J=6.9$), 4.00-4.13 (m, 4H), 4.53 (s, br, 2H), 5.20 (d, 1H, $J=8.4$), 5.83 (d, 1H, $J=15.3$), 6.42
(s, 1H), 6.75 (dd, 1H, $J=15.6, 5.4$), 6.92 (d, 1H, $J=8.1$), 7.67 (d, 1H, $J=7.2$).

ESI-MS ($\text{M}+\text{H}^+$) = 533.

Example 20: Preparation of compound 20

10 Compound 20 was prepared in a manner similar to that described in Example 1.

$^1\text{H NMR}$ (CDCl_3) δ 0.90-0.93 (m, 18H), 1.24-1.29 (t, 3H, $J=5.4\text{Hz}$), 1.52-2.20 (m,
4H), 2.40 (br, s, 2H), 3.32-3.35 (d, 2H), 3.82-3.84 (m, 2H), 3.91-3.96 (t, 1H, $J=7.8\text{Hz}$),
4.13-4.20 (q, 2H, $J=7.5\text{Hz}$), 4.58 (br s, 2H), 5.25-5.27 (d, 1H, $J=8.1\text{Hz}$), 5.89 (d, 1H,
 $J=15.3\text{ Hz}$), 6.32 (br, s, 1H), 6.81 (dd, 1H, $J=15.6\text{ Hz}, 5.1\text{ Hz}$), 6.92 (m, 1H), 7.77 (m,
15 1H).

ESI-MS ($\text{M}+\text{H}^+$) = 539.

Example 21: Preparation of compound 21

Compound 21 was prepared in a manner similar to that described in Example 1.

20 $^1\text{H NMR}$ (CDCl_3) δ 0.92-0.96 (m, 12H), 1.27 (t, 3H, $J=7.2$), 1.43 (s, 9H), 2.04-
2.11 (m, 2H), 2.38 (s, br, 2H), 3.23-3.35 (m, 2H), 3.83 (t, 1H, $J=7.2$), 4.16 (q, 2H, $J=7.2$),
4.61 (m, 2H), 5.00-5.03 (m, 1H), 5.89 (d, 1H, $J=15.3$), 6.48 (s, br, 1H), 6.81 (dd, 1H,
 $J=15.6, 5.1$), 7.00 (m, 1H), 7.68 (m, 1H).

ESI-MS ($\text{M}+\text{H}^+$) = 539.

Example 22: Preparation of compound 22

Compound 22 was prepared in a manner similar to that described in Example 1.

30 $^1\text{H NMR}$ (CDCl_3) δ 0.91 (s, br, 12H), 1.21-1.25 (m, 4H), 1.34-1.36 (m, 1H), 1.51-
1.75 (m, 1H), 2.00-2.13 (m, 2H), 2.34 (m, 2H), 3.26-3.29 (m, 2H), 3.64 (s, br, 2H), 3.94-
3.96 (m, 1H), 4.15 (q, 2H, $J=6.9$), 4.22 (m, 2H), 4.54 (s, 2H), 4.59 (s, br, 1H), 5.45 (d,

1H, J=7.5), 5.89 (d, 1H, J=15.6), 6.61 (s, br, 1H), 6.80 (dd, 1H, J=15.9, 4.5), 7.01 (d, 1H, J=7.2), 7.30 (m, 5H), 7.74 (d, 1H, J=7.2).

ESI-MS (M+H⁺) = 616.

5 Example 23: Preparation of compound 23

Compound 23 was prepared in a manner similar to that described in Example 1.

ESI-MS (M+H⁺) = 545.

Example 24: Preparation of compound 24

10 Compound 24 was prepared in a manner similar to that described in Example 1.

ESI-MS (M+H⁺) = 558.

Example 25: Preparation of compound 25

Compound 25 was prepared in a manner similar to that described in Example 1.

15 ¹H NMR (CD₃OD) δ 0.95-1.00 (m, 12 H), 1.26-1.29 (m, 6H), 1.57-1.85 (m, 5H), 2.03-2.14 (m, 3H), 2.27-2.32 (m, 1H), 2.45-2.52 (m, 1H), 3.93 (s, br, 2H), 4.03-4.04 (m, 1H), 4.16-4.20 (m, 5H), 4.37-4.41 (m, 1H), 4.59-4.63 (m, 1H), 5.94 (d, 1H, J=15.6), 6.89 (dd, 1H, J=15.6, 4.5).

ESI-MS (M+H⁺) = 568.

20

Example 26: Preparation of compound 26

Compound 26 was prepared in a manner similar to that described in Example 1.

25 ¹H NMR (CDCl₃) δ 0.86-0.96 (m, 12H), 1.27 (t, 3H, J=6.9), 1.49-1.82 (m, 10H), 1.95-2.02 (m, 1H), 2.40 (s, br, 2H), 3.32-3.34 (m, 2H), 4.16 (q, 2H, J=6.9), 4.55-4.67 (m, 3H), 5.89 (d, 1H, J=15.3), 6.39 (s, br, 1H), 6.72 (d, 1H, J=5.4), 6.81 (dd, 2H, J=15.9, 5.1), 7.31-7.76 (m, 5H).

ESI-MS (M+H⁺) = 557.

Example 27: Preparation of compound 27

30 Compound 27 was prepared in a manner similar to that described in Example 1.

ESI-MS (M+H⁺) = 690.

Example 28: Preparation of compound 28

Compound 28 was prepared in a manner similar to that described in Example 1.

^1H NMR (CDCl_3) δ 0.89-0.94 (m, 6H), 1.22-1.28 (m, 5H), 1.57-2.03 (m, 8H), 2.43 (m, 2H), 3.33 (d, 2H, $J=6.6$), 3.65 (s, 3H), 3.67 (s, 3H), 4.15 (q, 2H, $J=6.9$), 4.26 (m, 1H), 4.44 (m, 1H), 4.60 (m, 1H), 5.28 (s, 1H), 5.91 (d, 1H, $J=15.6$), 6.24-6.29 (m, 1H), 6.82 (dd, 1H, $J=15.9, 5.4$), 7.89 (m, 1H).

ESI-MS ($\text{M}+\text{H}^+$) = 542.

Example 29: Preparation of compound 29

Compound 29 was prepared in a manner similar to that described in Example 1.

^1H NMR (CDCl_3) δ 7.78 (d, $J = 7.5$ Hz, 1H), 7.28 (s, 5H), 7.01 (d, $J = 7.5$ Hz, 1H), 6.79 (dd, $J = 15.5, 5.5$ Hz, 1H), 6.50 (s, 1H), 5.89 (d, $J = 15.5$ Hz, 1H), 5.38 (d, $J = 7.0$ Hz, 1H), 4.46-4.61 (m, 4H), 4.06-4.21 (m, 4H), 3.29 (d, $J = 9.0$ Hz, 2H), 2.35 (br, s, 2H), 2.03 (s, 3H), 1.47-1.72 (m, 3H), 1.44 (s, 9H), 1.17-1.28 (m, 6H), 0.88 (d, $J = 8.4$ Hz, 3H), 0.86 (d, $J = 8.4$ Hz 3H).

ESI-MS ($\text{M}+\text{H}^+$) = 631.

Example 30: Preparation of compound 30

Compound 30 was prepared in a manner similar to that described in Example 1.

ESI-MS ($\text{M}+\text{H}^+$) = 663.

Example 31: Preparation of compound 31

Compound 31 was prepared in a manner similar to that described in Example 1.

^1H NMR (CDCl_3) δ 0.84-0.94 (m, 6H), 1.07 (m, 3H), 1.25 (m, 3H), 1.27 (s, 9H), 1.44 (s, 9H), 1.66-1.73 (m, 5H), 2.37 (br, s, 1H), 3.33 (d, 2H, $J=8.1$), 4.08-4.19 (m, 5H), 4.42-4.58 (m, 2H), 5.53-5.16 (d, 1H, $J=4.8\text{Hz}$), 5.87-5.92 (d, 1H, $J=15.6\text{Hz}$), 6.77-6.84 (dd, 1H, $J=15.3\text{Hz}, 5.1\text{Hz}$), 7.34-7.37 (d, 1H, $J=8.1\text{Hz}$), 7.650-7.60 (m, 1H).

ESI-MS ($\text{M}+\text{H}^+$) = 597.

Example 32: Preparation of compound 32

Compound 32 was prepared in a manner similar to that described in Example 1.

ESI-MS ($M+H^+$) = 561.

Example 33: Preparation of compound 33

Compound 33 was prepared in a manner similar to that described in Example 1.

5 $^1\text{H NMR}$ (CDCl_3) δ 0.83-0.88 (m, 6H), 1.10-1.20 (m, 9H), 1.30-1.35 (t, 2H, $J=7.5$), 1.57-1.96 (m, 5H), 2.38 (s, br, 2H), 3.02-3.09 (q, 1H, $J=7.8$), 3.27-3.39 (m, 4H), 4.10 (q, 2H, $J=6.9$), 4.34 (s, br, 2H), 4.50 (m, 1H), 4.60 (m, 1H), 4.95 (s, br, 1H), 5.80-5.85 (d, 1H, $J=15.6$), 6.25 (s, br, 1H), 6.73 (dd, 1H, $J=4.5$), 7.31 (s, br, 1H), 7.69 (s, br, 1H).

10 ESI-MS ($M+H^+$) = 527.

Example 34: Preparation of compound 34

Compound 34 was prepared in a manner similar to that described in Example 1.

15 $^1\text{H NMR}$ (CDCl_3) δ 0.93 (s, br, 6H), 1.28 (t, 3H, $J=6.9$), 1.60-2.00 (m, 9H), 2.29 (m, 1H), 2.49 (m, 1H), 3.21-3.34 (m, 4H), 3.65-3.69 (m, 6H), 4.17 (q, 2H, $J=6.6$), 4.30 (m, 1H), 4.50 (m, 1H), 4.78 (m, 1H), 5.14 (m, 1H), 5.87 (d, 1H, $J=15$), 6.49-6.56 (m, 1H), 6.80-6.86 (m, 2H).

ESI-MS ($M+H^+$) = 570.

20 Example 35: Preparation of compound 35

Compound 35 was prepared in a manner similar to that described in Example 1.

25 $^1\text{H NMR}$ (CDCl_3) δ 0.94 (dd, 6H, $J=12, 4.8$), 1.29 (t, 3H, $J=6.9$), 1.67-2.04 (m, 9H), 2.30-2.50 (m, 2H), 2.94 (s, 3H), 2.96 (s, 3H), 3.18 (s, br, 2H), 3.35 (s, br, 2H), 4.05 (m, 1H), 4.18 (q, 2H, $J=6.9$), 4.49-4.52 (m, 1H), 4.61-4.77 (m, 1H), 5.30 (s, 1H), 5.90 (d, 1H, $J=15.6$), 6.25-6.34 (m, 1H), 6.82 (m, 1H), 7.38 (m, 1H), 7.77 (d, 1H, $J=7.5$).

ESI-MS ($M+H^+$) = 610.

Example 36: Preparation of compound 36

Compound 36 was prepared in a manner similar to that described in Example 1.

30 $^1\text{H NMR}$ (CDCl_3) δ 0.838 (m, 12H), 1.18 (t, 3H, $J=6.3$), 1.44-1.68 (m, 5H), 2.04-2.10 (m, 3H), 2.28 (m, 2H), 3.20-3.23 (m, 2H), 3.90-3.95 (m, 1H), 4.07-4.10 (m, 2H),

4.54 (s, br, 1H), 5.01 (s, 2H), 5.52-5.55 (m, 1H), 5.84 (d, 1H, J=15.6), 6.63 (s, 1H), 6.75 (dd, 1H, J=15.6, J=4.8), 7.73 (d, 1H, J=6.9).

ESI-MS (M+H⁺) = 573.

5 Example 37: Preparation of compound 37

Compound 37 was prepared in a manner similar to that described in Example 1.

¹H NMR (CDCl₃) δ 0.89 (m, 6H), 1.23 (t, 3H, J=10.2), 1.45-1.99 (m, 6H), 2.36-2.58 (m, 2H), 2.94-3.09 (m, 2H), 3.26-3.34 (m, 2H), 4.10-4.20 (m, 3H), 4.44-4.54 (m, 2H), 5.04 (s, 2H), 5.89 (d, 1H, J=15.3), 6.79 (dd, 1H, J=15.3, 4.8), 7.16-7.30 (m, 10H).

10 ESI-MS (M+H⁺) = 621.

Example 38: Preparation of compound 38

Compound 38 was prepared in a manner similar to that described in Example 1.

15 ¹H NMR (CD₃OD) δ 0.79-0.91 (m, 6 H), 1.20-1.28 (m, 3H), 1.45-1.83 (m, 6H), 2.32-2.53 (m, 2H), 2.92-3.24 (m, 4H), 4.10-4.20 (m, 2H), 4.41-4.58 (m, 3H), 5.13 (s, 2H), 5.91 (dd, 1H, J=15.6, 1.5), 6.77-6.91 (m, 3H), 7.18-7.32 (m, 7H).

ESI-MS (M+H⁺) = 637.

Example 39: Preparation of compound 39

20 Compound 39 was prepared in a manner similar to that described in Example 1.

¹H NMR (CD₃OD) δ 0.79-0.91 (m, 6 H), 1.20-1.28 (m, 3H), 1.45-1.83 (m, 6H), 2.36-2.57 (m, 2H), 2.92-3.24 (m, 4H), 4.10-4.20 (m, 2H), 4.41-4.58 (m, 3H), 5.13 (s, 2H), 5.91 (dd, 1H, J=15.6, 1.5), 6.82-6.91 (m, 3H), 7.21-7.35 (m, 7H).

ESI-MS (M+H⁺) = 639.

25

Example 40: Preparation of compound 40

Compound 40 was prepared in a manner similar to that described in Example 1.

30 ¹H NMR (CD₃OD) δ 0.79-0.90 (m, 6 H), 1.20-1.29 (m, 3H), 1.45-1.83 (m, 6H), 2.36-2.57 (m, 2H), 2.94-3.10 (m, 2H), 3.26-3.34 (m, 2H), 4.10-4.20 (m, 2H), 4.44-4.61 (m, 3H), 5.10 (s, 2H), 5.90 (dd, 1H, J=15.6, 1.5), 6.83-6.94 (m, 4H), 7.24-7.32 (m, 5H).

ESI-MS (M+H⁺) = 657.

Example 41: Preparation of compound 41

Compound 41 was prepared in a manner similar to that described in Example 1.

^1H NMR (CDCl_3) δ 0.79 (s, br, 6H), 1.12-1.22 (m, 7H), 1.38-1.96 (m, 3H), 2.32 (s, br, 1H), 2.90-3.05 (m, 2H), 3.21 (m, 2H), 4.10 (q, 2H, $J=6.9$), 4.37-4.58 (m, 3H), 4.95-5.02 (m, 3H), 5.85 (d, 1H, $J=15.9$), 6.17-6.18 (m, 2H), 6.70-6.77 (m, 2H), 7.03 (m, 1H), 7.25 (s, br, 5H), 7.54 (s, 1H), 8.16 (d, 1H, $J=7.2$).

ESI-MS ($\text{M}+\text{H}^+$) = 610.

Example 42: Preparation of compound 42

Compound 42 was prepared in a manner similar to that described in Example 1.

ESI-MS ($\text{M}+\text{H}^+$) = 711.

Example 43: Preparation of compound 43

Compound 43 was prepared in a manner similar to that described in Example 1.

^1H NMR (CDCl_3) δ 0.69-0.83 (m, 6H), 1.21 (t, 3H, $J=6.9$), 1.29 (d, 1H, $J=5.1$), 1.44 (m, 1H), 1.86-2.23 (m, 6H), 2.97-2.99 (m, 2H), 3.13-3.21 (m, 2H), 3.88 (s, 1H), 4.09 (q, 2H, $J=6.9$), 4.43 (m, 1H), 4.71-4.77 (m, 1H), 5.02 (s, 2H), 5.36-5.40 (m, 1H), 5.65 (d, 1H, $J=15.9$), 6.61 (dd, 1H, $J=15.3$, $J=4.5$), 7.08-7.26 (m, 10H), 7.46-7.49 (m, 1H).

ESI-MS ($\text{M}+\text{H}^+$) = 607.

Example 44: Preparation of compound 44

Compound 44 was prepared in a manner similar to that described in Example 1.

ESI-MS ($\text{M}+\text{H}^+$) = 579.

Example 45: Preparation of compound 45

Compound 45 was prepared in a manner similar to that described in Example 1.

^1H NMR (CDCl_3) δ 7.83 (d, $J = 8$ Hz, 1H), 7.28-7.36 (m, 5H), 6.81 (dd, $J = 16, 6$ Hz, 1H), 6.47 (s, 1H), 5.91 (d, $J = 15$ Hz, 1H), 5.53 (d, $J = 8$ Hz, 1H), 5.07-5.13 (m, 2H), 4.60-4.68 (m, 2H), 4.13 (q, $J = 7$ Hz, 2H), 3.97 (dd, $J = 8, 3.6$ Hz, 1H), 3.28-3.31 (m, 2H), 2.49 (t, $J = 7$ Hz, 2H), 2.29-2.42 (m, 2H), 1.90-2.21 (m, 6H), 2.06 (s, 3H), 1.76 (m,

1H), 1.60 (m, 1H), 1.26 (t, J = 7 Hz, 3H), 0.94 (d, J = 6.9 Hz, 3H) 0.91 (d, J = 6.9 Hz, 3H).

ESI-MS (M+H⁺) = 591.

5 Example 46: Preparation of compound 46

Compound 46 was prepared in a manner similar to that described in Example 1.

¹H NMR (CDCl₃) δ 7.75 (d, J = 7.5 Hz, 1H), 7.50 (d, J = 8.4 Hz, 1H), 7.24-7.52 (m, 5H), 6.82 (dd, J = 15, 5 Hz, 1H), 6.54 (s, 1H), 5.89 (dd, J = 15, 1.8 Hz, 1H), 5.42 (d, J = 7 Hz, 1H), 5.02-5.12 (m, 2H), 4.60-4.67 (m, 2H), 4.13 (q, J = 7.2 Hz, 2H), 3.28-3.30 (m, 2H), 2.49 (t, J = 7.5 Hz, 2H), 2.33-2.43 (m, 2H), 1.90-2.14 (m, 6H), 2.06 (s, 3H), 1.46-1.79 (m, 3H), 1.26 (t, J = 7.2 Hz, 3H) 0.92 (d, J = 1.5 Hz, 6H).

ESI-MS (M+H⁺) = 605.

Example 47: Preparation of compound 47

15 Compound 47 was prepared in a manner similar to that described in Example 1.

ESI-MS (M+H⁺) = 639.

Example 48: Preparation of compound 48

Compound 48 was prepared in a manner similar to that described in Example 1.

20 ESI-MS (M+H⁺) = 546.

Example 49: Preparation of compound 49

Compound 49 was prepared in a manner similar to that described in Example 1.

25 ¹H NMR (CDCl₃) δ 1.26 (t, 3H, J=6.9), 1.55-1.63 (m, 1H), 1.75-1.82 (m, 1H), 1.94-2.02 (m, 1H), 2.22-2.31 (m, 1H), 2.41-2.51 (m, 1H), 3.59-3.60 (m, 1H), 3.68-3.73 (m, 1H), 3.82-3.87 (m, 2H), 4.15 (q, 2H, J=6.9), 4.27-4.29 (m, 1H), 4.44-4.50 (m, 1H), 4.63 (s, br, 1H), 5.10 (s, 2H), 5.95 (d, 1H, J=15.6), 6.85 (dd, 1H, J=15.6, 4.5), 7.2-7.34 (m, 4H), 8.16 (d, 1H, J=8.7).

ESI-MS (M+H⁺) = 571.

30

Example 50: Preparation of compound 50

Compound 50 was prepared in a manner similar to that described in Example 1.

ESI-MS ($M+H^+$) = 594.

5 Example 51: Preparation of compound 51

Compound 51 was prepared in a manner similar to that described in Example 1.

1H NMR ($CDCl_3$) δ 0.88 (s, br, 6H), 1.18-1.77 (m, 12H), 2.38 (m, 1H), 2.93 (s, 3H), 2.97 (s, 3H), 3.29 (m, 2H), 4.10 (q, 2H, $J=6.9$), 4.49 (m, 2H), 5.15-5.21 (m, 1H), 5.37 (q, 1H, $J=5.7$), 5.80-5.85 (m, 2H), 6.07 (s, br, 1H), 6.73 (dd, 1H, $J=15, 5.4$), 7.29 (d, 1H, $J=8.7$), 7.82 (m, 1H), 8.70 (m, 1H).

ESI-MS ($M+H^+$) = 596.

Example 52: Preparation of compound 52

Compound 52 was prepared in a manner similar to that described in Example 1.

15 1H NMR ($CDCl_3$) δ 0.83-0.87 (m, 6H), 1.14 (s, 6H), 1.16-1.20 (m, 6H), 1.42-1.67 (m, 6H), 2.22-2.26 (m, 2H), 3.32-4.10 (m, 4H), 4.47 (br, s, 2H), 5.80-5.85 (d, 1H, $J=15.3$ Hz), 6.70-6.77 (dd, 1H, $J=15.3$ Hz, 5.1 Hz), 7.79-7.87 (br, s, 2H).

ESI-MS ($M+H^+$)=497.

20 Example 53: Preparation of compound 53

Compound 53 was prepared in a manner similar to that described in Example 1.

1H NMR ($CDCl_3$) δ 0.96 (s, br, 6H), 1.16-1.30 (m, 15H), 1.62-2.02 (m, 6H), 2.33-2.46 (m, 2H), 2.98 (d, 3H, $J=8.7$), 3.34 (s, br, 2H), 3.88-3.91 (m, 1H), 4.14-4.19 (m, 3H), 4.40 (m, 1H), 4.60-4.76 (m, 1H), 5.76-6.14 (m, 3H), 6.81 (d, 1H, $J=15$), 7.60 (m, 1H).

25 ESI-MS ($M+H^+$) = 575.

Example 54: Preparation of compound 54

Compound 54 was prepared in a manner similar to that described in Example 1.

30 1H NMR ($CDCl_3$) δ 0.97 (t, 6H, $J=8$), 1.08 (d, 3H, $J=5.7$), 1.28 (s, b, 12H), 1.55-1.86 (m, 6H), 2.05 (m, 1H), 2.42 (m, 2H), 3.31 (d, 2H, $J=8.1$), 4.14-4.25 (m, 5H), 4.37-

4.45 (m, 3H), 4.60 (m, 1H), 5.91-6.10 (m, 3H), 6.83 (dd, 1H, J=15.6, 4.5), 7.26-7.76 (m, 9H).

ESI-MS (M+H⁺) = 719.

5 Example 55: Preparation of compound 55

Compound 55 was prepared in a manner similar to that described in Example 1.

¹H NMR (CDCl₃) δ 0.96-0.99 (m, 6H), 1.14 (d, 3H, J=6.6), 1.24 (s, 9H), 1.28 (t, 3H, J=6.9), 1.39 (t, 3H, J=6.9), 1.58-2.04 (m, 5H), 2.35-2.40 (m, 2H), 3.05 (q, 2H, J=7.5), 3.31-3.34 (m, 2H), 3.87-3.89 (m, 1H), 4.09-4.10 (m, 1H), 4.17 (q, 2H, J=7.2), 4.41 (m, 10 1H), 4.59 (m, 1H), 4.74 (m, 1H), 5.73 (d, 1H, J=6.0), 5.84-5.96 (m, 1H), 6.22 (s, 1H), 6.41 (q, 1H, J=15.9, 4.8), 7.57-7.60 (m, 1H), 7.69 (d, 1H, J=7.2).

ESI-MS (M+H⁺) = 589.

Example 56: Preparation of compound 56

15 Compound 56 was prepared in a manner similar to that described in Example 1.

ESI-MS (M+H⁺) = 603.

Example 57: Preparation of compound 57

Compound 57 was prepared in a manner similar to that described in Example 1.

20 ¹H NMR (CDCl₃) δ 0.85-0.90 (m, 6H), 1.27 (t, 3H, J=7.2 Hz), 1.60-1.66 (m, 5H), 2.33 (m, 1H), 2.79 (m, 1H), 2.94 (m, 1H), 3.20 (m, 2H), 4.17 (q, 2H, J=6.9 Hz), 4.39 (m, 1H), 4.55 (m, 2H), 5.10 (s, 2H), 5.91 (d, 1H, J=15.9 Hz), 6.40 (m, 1H), 6.85 (dd, 1H, J=15.9, 5.1 Hz), 7.31 (m, 20H), 7.84 (m, 1H).

ESI-MS (M+H⁺) = 637.

25

Example 58: Preparation of compound 58

Compound 58 was prepared in a manner similar to that described in Example 1.

30 ¹H NMR (CDCl₃) δ 0.88-0.94 (m, 6H), 1.27 (t, 3H, J=7.2 Hz), 1.60-1.66 (m, 5H), 2.07 (m, 1H), 2.43 (m, 2H), 2.72-2.80 (m, 2H), 3.30 (m, 2H), 4.17 (q, 2H, J=6.9 Hz), 4.39 (m, 1H), 4.55 (m, 2H), 5.10 (s, 2H), 5.91 (d, 1H, J=15.9 Hz), 6.10-6.20 (m, 2H), 6.30 (m, 2H), 6.85 (dd, 1H, J=15.9, 5.1 Hz), 7.31 (m, 5H), 7.60 (m, 1H).

ESI-MS ($M+H^+$) = 588.

Example 59: Preparation of compound 59

Compound 59 was prepared in a manner similar to that described in Example 1.

5 $^1\text{H NMR}$ (CD_3OD) δ 0.86-1.04 (m, 6H), 1.21-1.52 (m, 12H), 1.61-1.71 (m, 6H),
2.06-2.86 (m, 6H), 3.40 (m, 2H), 4.10-4.27 (m, 2H), 4.34-4.45 (m, 1H), 4.60-4.72 (m,
1H), 5.13 (s, 2H), 5.90 (d, 1H, $J=15.6$), 6.81 (dd, 1H, $J=15.6, 4.8$), 7.27-7.54 (m, 5H).

ESI-MS ($M+H^+$) = 645.

10 Example 60: Preparation of compound 60

Compound 60 was prepared in a manner similar to that described in Example 1.

$^1\text{H NMR}$ (CD_3OD) δ 0.92-1.00 (m, 6H), 1.19-1.23 (m, 6H), 1.57-1.80 (m, 6H),
2.03-2.36 (m, 4H), 3.19-3.31 (m, 2H), 4.16-4.34 (m, 4H), 4.58-4.62 (m, 1H), 5.04-5.11
15 (m, 2H), 5.24 (d, 1H, $J=5.1$), 5.91 (d, 1H, $J=15.3$), 6.89 (dd, 1H, $J=15.3, 5.4$), 7.30-7.35
(m, 5H).

ESI-MS ($M+H^+$) = 576.

Example 61: Preparation of compound 61

Compound 61 was prepared in a manner similar to that described in Example 1.

20 ESI-MS ($M+H^+$) = 576.

Example 62: Preparation of compound 62

Compound 62 was prepared in a manner similar to that described in Example 1.

ESI-MS ($M+H^+$) = 617.

25

Example 63: Preparation of compound 63

Compound 63 was prepared in a manner similar to that described in Example 1.

ESI-MS ($M+\text{Na}^+$) = 867.

30 Example 64: Preparation of compound 64

Compound 64 was prepared in a manner similar to that described in Example 1.

^1H NMR (CDCl_3) δ 0.94-0.99 (m, 6H), 1.07 (m, 3H), 1.28 (m, 15H), 1.60 (m, 3H), 2.10 (br, s, 2H), 2.29 (br, s, 2H), 3.32-3.35 (d, 2H), 4.09 (m, 6H), 4.32 (br, s, 1H), 4.60 (br, s, 1H), 5.76 (m, 1H), 5.84-5.89 (d, 1H, $J=14.7\text{Hz}$), 5.98 (br, s, 1H), 6.81 (dd, 1H, $J=15.9\text{ Hz}$, 5.4 Hz), 7.38 (m, 1H), 7.60 (m, 1H).

5 ESI-MS ($\text{M}+\text{H}^+$) = 569.

Example 65: Preparation of compound 65

Compound 65 was prepared in a manner similar to that described in Example 1.

^1H NMR (CDCl_3) δ 0.94-0.99 (m, 6H), 1.28 (t, 3H, $J=6.9$), 1.75-1.82 (m, 2H),
10 2.10-2.33 (m, 3H), 3.03-3.15 (m, 2H), 3.22-3.31 (m, 2H), 4.16 (q, 2H, $J=7.5$), 4.39-4.51
(m, 2H), 4.80-4.83 (m, 1H), 5.77 (d, 1H, $J=15.6$), 5.96 (s, 1H), 6.66-6.74 (m, 2H), 7.15-
7.62 (m, 10H), 7.7 (d, 2H, $J=8.1$).

ESI-MS ($\text{M}+\text{H}^+$) = 577.

15 Example 66: Preparation of compound 66

Compound 66 was prepared in a manner similar to that described in Example 1.

ESI-MS ($\text{M}+\text{H}^+$) = 630.

Example 67: Preparation of compound 67

20 Compound 67 was prepared in a manner similar to that described in Example 1.

^1H NMR (CDCl_3) δ 0.91 (s, br, 6H), 1.26 (t, 3H, $J=6.9$), 1.42 (s, 18H), 1.56-1.63
(m, 5H), 1.80-1.82 (m, 2H), 2.04 (m, 1H), 2.35-2.41 (m, 3H), 3.12 (s, br, 2H), 3.32 (s, br,
2H), 4.16 (m, 3H), 4.54 (s, br, 2H), 5.89 (d, 1H, $J=15.6$), 6.81(m, 2H).

ESI-MS ($\text{M}+\text{H}^+$) = 654.

25

Example 68: Preparation of compound 68

Compound 68 was prepared in a manner similar to that described in Example 1.

ESI-MS ($\text{M}+\text{H}^+$) = 809.

30 Example 69: Preparation of compound 69

Compound 69 was prepared in a manner similar to that described in Example 1.

ESI-MS ($M+H^+$) = 557.

Example 70: Preparation of compound 70

Compound 70 was prepared in a manner similar to that described in Example 1.

5 ESI-MS ($M+H^+$) = 547.

Example 71: Preparation of compound 71

Compound 71 was prepared in a manner similar to that described in Example 1.

ESI-MS ($M+H^+$) = 562.

10

Example 72: Preparation of compound 72

Compound 72 was prepared in a manner similar to that described in Example 1.

ESI-MS ($M+H^+$) = 605.

15 Example 73: Preparation of compound 73

Compound 73 was prepared in a manner similar to that described in Example 1.

ESI-MS ($M+H^+$) = 600.

Example 74: Preparation of compound 74

20 Compound 74 was prepared in a manner similar to that described in Example 1.

^1H NMR (CDCl_3) δ 0.88-0.98 (m, 6 H), 1.19-1.86 (m, 15H), 2.0-2.08 (m, 1H),
2.15-2.39 (m, 2H), 2.47-2.59 (m, 1H), 2.94 (m, 2H), 3.98-4.20 (m, 3H), 4.30-4.36 (m,
1H), 4.60-4.64 (m, 1H), 5.09 (m, 2H), 5.89 (d, 1H, $J=15.6$), 6.89 (dd, 1H, 15.6, 4.8),
7.26-7.33 (m, 5H).

25 ESI-MS ($M+H^+$) = 602.

Example 75: Preparation of compound 75

Compound 75 was prepared in a manner similar to that described in Example 1.

30 ^1H NMR(CDCl_3) δ 0.89-0.93(m, 6H), 1.13-1.15(m, 3H), 1.22-1.27(m, 3H), 1.55-
1.76(m, 4H), 1.95-2.-2(m, 1H), 2.03-2.46(m, 2H), 3.28-3.30(m, 2H), 4.11-4.18(m, 3H),

4.33(br s, 1H), 4.54-4.56(br s, 2H), 5.07(s, 2H), 5.84-5.94(d, 1H, J=15.9 Hz), 6.03-6.06(d, J=7.2 Hz), 6.76-6.83(dd, 1H, J=15.0 Hz, 5.4 Hz), 7.31(br s, 5H), 8.02(br s, 1H).

ESI-MS (M+H⁺)=575.

5 Example 76: Preparation of compound 76

Compound 76 was prepared as follows: Compound 74 (0.047 g, 0.11 mmol) was dissolved in CH₂Cl₂ (1.5 ml). Et₃N (0.037 ml, 3.0 eq.) was then added to the above solution. The resultant solution was stirred at room temperature for 30 minutes. Methyl chloroformate (0.013 ml, 1.5 eq.) was subsequently added. The reaction mixture was stirred at room temperature for another 20 hours and then concentrated to afford a residue. The residue was purified by flash column chromatography (5% MeOH in CH₂Cl₂) to afford compound 76 as a white solid (64 mg, 85%).

¹H NMR (CDCl₃) δ 0.88-0.98 (m, 6 H), 1.19-1.83 (m, 15H), 2.0-2.08 (m, 1H), 2.15-2.38 (m, 2H), 2.47-2.59 (m, 1H), 3.07 (m, 2H), 3.59 (s, 3H), 3.98-4.20 (m, 3H), 4.310-4.36 (m, 1H), 4.60-4.64 (m, 1H), 5.08 (m, 2H), 5.88 (d, 1H, J=15.6), 6.89 (dd, 1H, 15.6, 4.8), 7.27-7.32 (m, 5H).

ESI-MS (M+H⁺) = 660.

Example 77: Preparation of compound 77

20 Compound 77 was prepared in a manner similar to that described in Example 1.
ESI-MS (M+H⁺) = 523.

Example 78: Preparation of compound 78

25 Compound 78 was prepared in a manner similar to that described in Example 1.
ESI-MS (M+H⁺) = 557.

Example 79: Preparation of compound 79

30 Compound 79 was prepared in a manner similar to that described in Example 1.
ESI-MS (M+H⁺) = 423.

Example 80: Preparation of compound 80

Compound 80 was prepared as follows: Compound 59 (0.023 g, 0.036 mmol) was added to a solution of HCl in 1,4-dioxane (4.0 M, 2 ml). The solution was stirred at room temperature for 30 minutes and then concentrated to give compound 80 as a white solid (21 mg, 99%);

$^1\text{H NMR}$ (CD_3OD) δ 0.86-1.04 (m, 6H), 1.23 (s, br, 3H), 1.60-1.71 (m, 6H), 2.06-2.85 (m, 6H), 3.38 (m, 2H), 4.10-4.28 (m, 2H), 4.34-4.45 (m, 1H), 4.60-4.72 (m, 1H), 5.13 (s, 2H), 5.91 (d, 1H, $J=15.9$), 6.82 (dd, 1H, $J=15.9, 4.7$), 7.27-7.54 (m, 5H).

ESI-MS ($\text{M}+\text{H}^+$) = 589.

Example 81: Preparation of compound 81

Compound 81 was prepared in 20% yield from compound 52 and methyl chloroformate.

$^1\text{H NMR}$ (CDCl_3) δ 0.94-0.99 (m, 6H), 1.07 (m, 3H), 1.28 (m, 12H), 1.60 (m, 3H), 2.10 (br s, 2H), 2.29 (br s, 2H), 3.32-3.35 (d, 2H), 3.69 (s, 3H), 4.18 (m, 4H), 4.39 (br s, 1H), 4.61 (br s, 1H), 5.59-5.81 (m, 2H), 5.89-5.95 (d, 1H, $J=16.2\text{Hz}$), 6.82 (dd, 1H, $J=15.3\text{ Hz}, 5.7\text{ Hz}$), 7.34 (m, 1H), 7.56 (m, 1H).

ESI-MS ($\text{M}+\text{H}^+$) = 555.

Example 82: Preparation of compound 82

Compound 82 was prepared in a manner similar to that described in Example 1.

$^1\text{H NMR}$ (CDCl_3) δ 0.95 (m, 6H), 1.02-1.04 (m, 3H), 1.27 (s, 12H), 1.56-1.82 (m, 5H), 2.03 (s, 3H), 2.18 (m, 1H), 2.40 (m, 2H), 3.31-3.34 (m, 2H), 4.15-4.18 (m, 3H), 4.36 (m, 1H), 4.46 (m, 1H), 4.61 (m, 1H), 5.92 (d, 1H, $J = 15\text{ Hz}$), 6.38 (s, 1H), 6.63 (s, 1H), 6.82 (d, 1H, $J = 15\text{ Hz}$), 7.53 (d, 1H, $J = 6\text{ Hz}$), 7.65 (d, 1H, $J = 6\text{ Hz}$).

ESI-MS ($\text{M}+\text{H}^+$) = 539.

Example 83: Preparation of compound 83

Compound 83 was prepared in a manner similar to that described in Example 1.

$^1\text{H NMR}$ (CDCl_3) δ 0.96 (t, 6H, $J = 6\text{ Hz}$), 1.08 (d, 3H, $J = 6\text{ Hz}$), 1.26 (s, 9H), 1.54-2.42 (m, 8H), 3.35 (d, 2H, $J = 9\text{ Hz}$), 3.68 (t, 2H, $J = 6\text{ Hz}$), 4.14-4.18 (m, 4H), 4.27-

4.38 (m, 2H), 4.45 (br, 1H), 4.60 (br, 1H), 5.92 (d, 2H, J = 15 Hz), 6.32 (s, 1H), 6.83 (dd, 1H, J = 15.0 Hz, 6.0 Hz), 7.39 (d, 1H, J = 6 Hz), 7.66 (d, 1H, J = 6 Hz).

ESI-MS (M+Na⁺) = 625.

5 Example 84: Preparation of compound 84

Compound 84 was prepared in a manner similar to that described in Example 1.

¹H NMR (CDCl₃) δ 0.97 (q, 6H, J = 6 Hz), 1.13 (d, 3H, J = 6 Hz), 1.26 (s, 12H), 1.55-2.06 (m, 6H), 2.41 (br, 2H), 3.32 (d, 2H, J = 9 Hz), 4.04-4.09 (m, 2H), 4.18 (q, 2H, J = 6 Hz), 4.41 (br, 1H), 4.59 (br, 1H), 5.93 (d, 1H, J = 15 Hz), 6.32 (s, 1H), 6.83 (dd, 1H, J = 15.0 Hz, 6.0 Hz), 7.02 (d, 1H, J = 6 Hz), 7.58 (d, 1H, J = 6 Hz), 7.79 (d, 1H, J = 6 Hz).

ESI-MS (M+H⁺) = 629.

Example 85: Preparation of compound 85

Compound 85 was prepared in a manner similar to that described in Example 1 in a 68% yield.

¹H NMR (CDCl₃) δ 7.83 (d, J = 8 Hz, 1H), 7.28-7.36 (m, 5H), 6.81 (dd, J = 16, 6 Hz, 1H), 6.47 (s, 1H), 5.91 (d, J = 15 Hz, 1H), 5.53 (d, J = 8 Hz, 1H), 5.07-5.13 (m, 2H), 4.60-4.68 (m, 2H), 4.13 (m, 3H), 3.97 (dd, J = 8, 3.6 Hz, 1H), 3.28-3.31 (m, 2H), 2.49 (t, J = 7 Hz, 2H), 2.29-2.42 (m, 2H), 1.90-2.21 (m, 6H), 2.06 (s, 3H), 1.76 (m, 1H), 1.26 (m, 12H), 0.95 (d, J = 6.9 Hz, 3H).

ESI-MS (M+H⁺) = 649.

Example 86: Preparation of compound 86

Compound 86 was prepared in a manner similar to that described in Example 1 in a 57% yield.

¹H NMR (CDCl₃) δ 7.83 (d, J = 8 Hz, 1H), 7.28-7.36 (m, 5H), 6.81 (dd, J = 16, 6 Hz, 1H), 6.47 (s, 1H), 5.91 (d, J = 15 Hz, 1H), 5.53 (d, J = 8 Hz, 1H), 5.07-5.13 (m, 2H), 4.60-4.68 (m, 2H), 4.13 (q, J = 7 Hz, 2H), 3.97 (dd, J = 8, 3.6 Hz, 1H), 3.28-3.31 (m, 2H), 2.06-2.86 (m, 6H), 1.90-2.21 (m, 6H), 2.06 (s, 3H), 1.76 (m, 1H), 1.21-1.52 (m, 12H).

ESI-MS (M+H⁺) = 663.

Example 87: Preparation of compound 87

Compound 87 was prepared in a manner similar to that described in Example 1.

^1H NMR (CDCl_3) δ 0.98 (d, 6H, $J = 6$ Hz), 1.09 (s, 12H), 1.27 (t, 3H, $J = 6$ Hz), 1.50-2.49 (m, 8H), 3.32 (q, 2H, $J = 3$ Hz), 3.63 (q, 1H, $J = 3$ Hz), 3.87-3.90 (m, 1H), 4.16 (q, 2H, $J = 6$ Hz), 4.29-4.45 (m, 1H), 4.77 (m, 1H), 5.84 (br, 1H), 5.93 (d, 1H, $J = 15$ Hz), 6.12 (d, 1H, $J = 6$ Hz), 6.81 (dd, 1H, $J = 15.0$ Hz, 6.0 Hz), 7.33-7.44 (m, 5H), 8.48 (d, 1H, $J = 9$ Hz).

ESI-MS($\text{M}+\text{H}^+$)=651.

Example 88: Preparation of compound 88

Compound 88 was prepared in a manner similar to that described in Example 1.

^1H NMR (CDCl_3) δ 0.91-0.96 (m, 6H), 1.04-1.06 (m, 3H), 1.23 (m, 12H), 1.51-2.05 (m, 8H), 2.40 (br s, 2H), 3.30 (m, 2H), 4.15 (m, 4H), 4.39 (m, 1H), 4.54 (m, 2H), 5.19-5.32 (m, 2H), 5.84-5.97 (m, 3H), 6.82 (dd, 1H, $J=15.6$ Hz, 5.4 Hz), 7.35 (m, 1H), 7.60(m, 1H).

ESI-MS ($\text{M}+\text{H}^+$) = 581.

Example 89: Preparation of compound 89

Compound 89 was prepared in a 85% yield by reacting compound 1 with pivaloyl chloride in the presence of Et_3N and DMAP.

^1H NMR (CDCl_3) δ 0.94-0.98 (m, 6H), 1.08 (d, $J=6.3$ Hz, 3H), 1.25 (s, 9H), 1.66-1.73 (m, 5H), 2.21-2.50 (m, 2H), 3.20-3.30 (m, 2H), 4.16 (q, $J=6.9$ Hz, 2H), 4.42 (br, 1H), 4.58 (br, 1H), 5.10 (s, 2H), 5.9 (d, $J=15.6$ Hz, 1H), 6.82 (dd, $J=15.3$ Hz, 5.1 Hz, 1H), 7.2-7.34 (m, 4H), 7.60 (d, $J=7.5$ Hz, 1H).

ESI-MS ($\text{M}+\text{H}^+$) = 659.

Example 90: Preparation of compound 90

Compound 6 was dissolved in an 1N NaOH/EtOH solution. The mixture was stirred for 1 hour and concentrated to afford a residue. The residue was purified by column chromatography to give compound 90 in a 20% yield.

^1H NMR (CDCl_3) δ 0.83-0.93 (m, 6H), 1.02-1.04 (d, 4H, $J=5.7\text{Hz}$), 1.16-1.31 (m, 11H), 1.52-1.74 (m, 5H), 2.0 (m, 1H), 2.30-2.43 (m, 2H), 3.24-3.26 (m, 2H), 4.15 (m, 2H), 4.41 (br s, 1H), 4.63 (br s, 1H), 5.08 (s, 2H), 5.9 (m, 1H), 6.82 (m, 1H), 7.29-7.33 (m, 5H), 7.47 (d, 1H, $J=8.1\text{ Hz}$), 7.69 (m, 1H).

5 ESI-MS ($\text{M}+\text{H}^+$) = 603.

Example 91: Preparation of compound 91

Compound 91 was prepared in a manner similar to that described in Example 1.

10 ^1H NMR (CDCl_3) δ 0.93 (q, 6H, $J = 3\text{ Hz}$), 1.05 (d, 3H, $J = 6\text{ Hz}$), 1.21-1.24 (m, 15H), 1.42-2.05 (m, 6H), 2.38 (br, 2H), 3.26-3.29 (m, 2H), 4.15 (br, 2H), 4.39 (m, 1H), 4.57 (m, 1H), 4.97-5.14 (m, 3H), 5.85-5.90 (m, 3H), 6.78 (dd, 1H, $J = 15.6\text{ Hz}$, 5.4 Hz), 7.34 (m, 5H), 7.57 (d, 1H, $J = 7.2\text{ Hz}$).

ESI-MS ($\text{M}+\text{H}^+$) = 645.

15 Example 92: Preparation of compound 92

Compound 92 was prepared in a manner similar to that described in Example 1.

20 ^1H NMR (CDCl_3) δ 0.93-0.98 (m, 6H), 1.10 (d, 3H, $J = 6.3$), 1.25 (s, 9H), 1.55-1.68 (m, 9H), 2.40 (m, 3H), 3.30-3.39 (m, 2H), 4.10-4.20 (m, 4H), 4.26-4.36 (m, 1H), 4.25-4.58 (m, 1H), 5.10 (q, 2H, $J = 12$), 5.49 (d, 1H, $J = 16.5$), 5.64 (s, 1H), 5.84 (d, 1H, $J = 5.4$), 6.59 (dd, $J = 16.5$, 5.0H), 7.29-7.34 (m, 5H), 7.96 (d, $J = 6.0$).

ESI-MS ($\text{M}+\text{H}^+$) = 584.1.

Example 93: Preparation of compound 93

25 Compound 93 was prepared in a manner similar to that described in Example 1 in a 66% yield.

^1H NMR (CDCl_3) δ 0.86-0.95 (m, 6H), 1.22 (m, 4H), 1.41 (m, 11H), 1.55-1.90 (m, 9H), 2.32 (m, 2H), 3.25-3.35 (m, 3H), 3.39 (m, 1H), 4.11-4.19 (m, 2H), 4.34-4.47 (m, 2H), 4.85 (m, 1H), 5.08 (s, 2H), 5.84 (m, 2H), 6.09 (m, 1H), 6.73-6.75 (m, 1H), 7.73 (m, 1H).

30 ESI-MS($\text{M}+\text{H}^+$)=707

Example 94: Preparation of compound 94

Compound 94 was prepared as follows: Compound 93 was added into a solution of HCl in 1,4-dioxane. The mixture was stirred at room temperature for 30 minutes and concentrated. CH₂Cl₂ was then added to the resultant residue and the mixture was cooled
5 down to 0-5°C. N-methylmorpholine was added into the mixture, which was then stirred for 10 minutes. Benzochloromate was subsequently added to the solution, which was stirred at room temperature for another 2 hours to give a crude product. The crude product was purified by flash column chromatography to give compound 94.

¹H NMR (CDCl₃) δ 0.86-0.95 (m, 6H), 1.22 (m, 4H), 1.41 (m, 11H), 1.55-1.90 (m,
10 9H), 2.32 (m, 2H), 3.25-3.35 (m, 3H), 3.39 (m, 1H), 4.11-4.19 (m, 2H), 4.34-4.47 (m, 2H), 4.85 (m, 1H), 5.84 (m, 2H), 6.09 (m, 1H), 6.73-6.75 (m, 1H), 7.73 (m, 5H).

ESI-MS (M+H⁺) = 585.

Example 95: Preparation of compound 95

Compound 95 was prepared in a manner similar to that described in Example 1.
15

¹H NMR (CDCl₃) δ 0.93-0.98 (m, 6H), 1.07 (d, 3H, J = 5.4), 1.27 (s, 9H), 1.52-2.10 (m, 8H), 2.40 (m, 2H), 3.23 (s, 3H), 3.28-3.31 (m, 2H), 3.68 (s, 3H), 4.16 (m, 2H), 4.44-4.45 (m, 1H), 4.63-4.64 (m, 1H), 5.10 (q, 2H, J = 6), 5.90 (d, 1H, J = 4.5), 6.51 (d, 1H, J = 15.6), 6.83 (dd, 1H, J = 5.1, 15.3), 7.35 (m, 5H), 7.57-7.62 (m, 1H).
20

ESI-MS (M+H⁺) = 646.

Example 96: Preparation of compound 96

Compound 96 was prepared in a manner similar to that described in Example 1 in a 38% yield.

¹H NMR (CDCl₃) δ 0.95 (q, 6H, J = 8.4 Hz), 1.09 (d, 3H, J = 6.3 Hz), 1.17 (d, 6H, J = 6.6 Hz), 1.26 (s, 9H), 1.52-1.82 (m, 5H), 1.99-2.06 (m, 1H), 2.38 (br, 2H), 3.27-3.30 (m, 2H), 4.06-4.21 (m, 3H), 4.41 (m, 1H), 4.61 (m, 1H), 5.11 (q, 2H, J = 3 Hz), 5.65 (d, 1H, J = 7.8), 5.84-5.89 (m, 3H), 6.66 (dd, 1H, J = 14.7 Hz, 5.4 Hz), 7.29-7.35 (m, 5H), 7.47 (d, 1H, J = 7.8 Hz).
25

ESI-MS(M+H⁺)=644.
30

Example 97: Preparation of compound 97

Compound 97 was prepared in a manner similar to that described in Example 1 in a 71% yield.

$^1\text{H NMR}(\text{CDCl}_3)$ δ 0.25 (q, 2H, J = 5.4 Hz), 0.50-0.56 (m, 2H), 0.93 (q, 6H, J = 2.1 Hz), 1.03-1.18 (m, 4H), 1.24 (s, 9H), 1.50-2.07 (m, 7H), 2.36-2.41 (m, 2H), 3.25-3.28 (m, 2H), 3.92 (d, 2H, J = 7.2 Hz), 4.14-4.16 (m, 2H), 4.40-4.45 (m, 1H), 4.58 (m, 1H), 5.08 (q, 2H, J = 6.6 Hz), 5.87 (d, 1H, J = 5.1 Hz), 5.93 (d, 1H, J = 15.6 Hz), 6.08 (s, 1H), 6.82 (dd, 1H, J = 15.9 Hz, 5.4 Hz), 7.29-7.38 (m, 5H), 7.63 (d, 1H, J = 7.5 Hz).

ESI-MS ($\text{M}+\text{Na}^+$) = 679.

Example 98: Preparation of compound 98

Compound 98 was prepared in a manner similar to that described in Example 1 in a 50% yield.

$^1\text{H NMR}(\text{CDCl}_3)$ δ 0.85-0.98 (m, 6H), 1.06 (m, 4H), 1.26 (s, 9H), 1.44-1.63 (m, 9H), 1.75-1.80 (m, 7H), 2.05-2.34 (m, 2H), 3.27-3.29 (m, 2H), 3.70-3.72 (m, 1H), 4.14 (m, 2H), 4.42 (m, 1H), 4.59 (m, 1H), 5.08 (s, 2H), 5.88-5.98 (m, 4H), 6.09 (m, 1H), 6.80-6.87 (m, 1H), 7.25 (m, 10H), 7.65 (m, 1H).

ESI-MS ($\text{M}+\text{H}^+$) = 707.

Example 99: Preparation of compound 99

Compound 99 was prepared by the procedures described follows:

8 ml of a 1 N LiOH aqueous solution at 0°C was added to Intermediate 6 described in Example 1 (0.107 g, 0.243 mmol, 1 eq.) in THF (11 ml). The mixture was stirred at this temperature for 5 minutes, and then at room temperature for 1 hour. The reaction mixture was then partitioned between H₂O (10 ml) and EtOAc (10 ml). The aqueous layer was acidified to pH=2 with a 10% KHSO₄ aqueous solution and was extracted with EtOAc (2 X 10 ml). The organic layers were combined, dried over Na₂SO₄, and concentrated under vacuum to afford a crude corresponding carboxylic acid. Under an atmosphere of nitrogen, the carboxylic acid (0.100 g, 0.243 mmol, 1 eq.) was dissolved slowly in anhydrous DMF (1.2 ml, 0.2 M). After cesium carbonate (0.119 g, 0.365 mmol, 1.5 eq.) and benzyl bromide (0.062g, 0.040ml, 0.365 mmol, 1.5 eq.) were

sequentially added into the above solution, the reaction mixture was stirred at ambient temperature overnight under N₂. The reaction was then quenched by addition of a 10% KHSO₄ aqueous solution (5 ml) slowly. The resultant solution was extracted with CH₂Cl₂ (5 ml x 4). The organic layers were combined, washed with brine, dried over Na₂SO₄, and concentrated under vacuum to give a crude product. The crude product thus obtained was purified by flash chromatography (2% MeOH in CH₂Cl₂) on silica gel to afford Intermediate 8 (94 mg, 77%) as a white solid. ¹H NMR (CDCl₃) δ 0.85 (d, J=4.8 Hz, 6H), 1.19 (m, 1H), 1.32 (s, 9H), 1.38-1.71 (m, 4H), 1.95-2.08 (m, 2H), 2.25-2.30 (m, 2H), 3.19-3.27 (m, 2H), 4.15 (m, 1H), 4.51 (m, 1H), 5.08 (s, 2H), 5.88 (d, J=15.3 Hz, 1H), 6.57 (br s, 1H), 6.79 (dd, J=15.3, 4.8 Hz, 1H), 7.24 (m, 5H), 7.55(d, J=7.5 Hz, 1H). ESI-MS (M+H⁺) = 502.

A 4.0 M solution of HCl in 1,4-dioxane (1.78 ml) was added to a solution of Intermediate 8 (0.094 g, 0.187 mmol, 1eq.) in the same solvent (1.8 ml) at 23°C. After 2 hours of stirring, the reaction mixture was concentrated under reduced pressure. The residue thus obtained was dissolved in CH₂Cl₂ (15 ml). Z-Thr(*t*-Bu)-OH (0.058 g, 0.187 mmol, 1.0 eq.), 4-methylmorpholine (0.080 ml, 0.750 mmol, 4.0 eq.), HOBT (0.038 g, 0.281 mmol, 1.5 eq.), and EDC (0.054 g, 0.281 mmol, 1.5 eq.) were added sequentially. The resultant solution was stirred overnight at 23°C and then the solvent was removed under vacuum. The residue was partitioned between a 10% KHSO₄ aqueous solution (1 ml) and CH₂Cl₂ (7 ml). The organic layer was dried over MgSO₄, filtered, and concentrated. The crude product thus obtained was purified by flash chromatography (70% EtOAc in hexanes) to afford compound 99 (0.117 g, 90%) as a white foam. ¹H NMR (CDCl₃) δ 0.94 (dd, J=8.4, 6.0 Hz, 6H), 1.05 (d, J=6.3 Hz, 3H), 1.25 (s, 9H), 1.46-1.80 (m, 6H), 1.99-2.06 (m, 1H), 2.38 (m, 1H), 3.27-3.30 (m, 2H), 4.15 (m, 2H), 4.40 (m, 1H), 4.58 (m, 1H), 5.04-5.15 (m, 4H), 5.87 (s, 2H), 5.96 (dd, J=15.9, 1.2 Hz, 1H), 6.86 (dd, J=15.6, 5.4 Hz, 1H), 7.28-7.33 (m, 10H), 7.61(d, J=7.2 Hz, 1H). ESI-MS (M+H⁺) = 693.

Example 100: Preparation of compound 100

Compound 100 was prepared in a manner similar to that described in Example 1.

^1H NMR (CDCl_3) δ 0.90-0.96 (m, 6H), 1.02-1.04 (m, 3H), 1.24 (s, 9H), 1.49-1.69(m, 9H), 1.75-1.80 (m, 7H), 1.98-2.07 (m, 1H), 2.36 (m, 2H), 3.25-3.28 (m, 2H), 4.14 (m, 1H), 4.41 (m, 1H), 4.56 (m, 1H), 5.08 (s, 2H), 5.14-5.19 (m, 1H), 5.85-5.90 (m, 2H), 6.73-6.80 (m, 1H), 7.33 (m, 5H), 7.62 (m, 1H).

5 ESI-MS ($\text{M}+\text{H}^+$) = 671.

Example 101: Preparation of compound 101


Compound 101 was prepared in a manner similar to that described in Example 1.

^1H NMR (CDCl_3) δ 1.07 (d, $J=6.3$ Hz, 3H), 1.28 (s, 9H), 1.66-1.73 (m, 4H), 2.21-10 2.50 (m, 8H), 2.88 (m, 2H), 3.20-3.30 (m, 2H), 4.16 (q, $J=6.9$ Hz, 2H), 4.42 (br, 1H), 4.58 (br, 1H), 5.01-5.10 (m, 3H), 5.91 (d, $J=15.6$ Hz, 1H), 6.82 (dd, $J=15.3$ Hz, 5.1 Hz, 1H), 7.2-7.34 (m, 4H), 7.60 (d, $J=7.5$ Hz, 1H).

ESI-MS ($\text{M}+\text{H}^+$) = 632.

15 Example 102: Preparation of compound 102

Compound 102 was prepared in a 66% yield by reacting compound 112

with  (1.5 eq.) via a Wittig reaction for 24 hours.

^1H NMR(CDCl_3) δ 0.93-0.98(m, 6H), 1.26(s, 9H), 1.66-1.73(m, 5H), 2.22-2.51(m, 4H), 3.19-3.30(m, 2H), 4.26(t, $J=6.9$ Hz, 2H), 4.44(br, 1H), 4.57(br, 1H), 5.12(s, 20 2H), 6.82(d, $J= 5.1$ Hz, 1H), 7.22-7.34(m, 4H), 7.60(d, $J=7.5$ Hz, 1H).

ESI-MS ($\text{M}+\text{H}^+$) = 629.

Example 103: Preparation of compound 103

Compound 103 was prepared in a manner similar to that described in Example 1
25 in a 63% yield.

^1H NMR(CDCl_3) δ 0.94-0.98 (m, 6H), 1.06 (d, 3H, $J = 5.7$ Hz), 1.27 (s, 9H), 1.53-2.10 (m, 6H), 2.38-2.43 (m, 2H), 3.28-3.30 (m, 2H), 3.72 (s, 3H), 4.16 (br, 2H), 4.48 (m, 1H), 4.59 (m, 1H), 5.11 (q, 2H, $J = 6.6$ Hz), 5.93 (d, 2H, $J = 15.9$ Hz), 6.12 (br, 1H), 6.84 (dd, 1H, $J = 15.6$ Hz, 5.1 Hz), 7.35-7.39 (m, 5H), 7.65 (d, 1H, $J = 6.9$ Hz).

ESI-MS ($M+H^+$) = 617.

Example 104: Preparation of compound 104

Compound 104 was prepared in a manner similar to that described in Example 1
5 in a 35% yield.

1H NMR($CDCl_3$) δ 0.93-0.98 (m, 12H), 1.06 (d, 4H, $J = 5.7$ Hz), 1.27 (s, 9H),
1.53-2.16 (m, 7H), 2.39-2.43 (m, 2H), 3.28-3.34 (m, 2H), 3.90 (d, 2H, $J = 6.9$ Hz), 4.16-
4.18 (m, 2H), 4.47-4.49 (m, 1H), 4.60 (br, 1H), 5.11(q, 2H, $J = 6.9$ Hz), 5.89 (d, 1H, $J =$
5.4 Hz), 5.94 (d, 1H, $J = 15.6$ Hz), 6.08 (s, 1H), 6.83 (dd, 1H, $J = 15.6$ Hz, 5.1 Hz), 7.31-
10 7.40 (m, 5H), 7.63 (d, 1H, $J = 7.5$ Hz).

ESI-MS ($M+H^+$) = 659.

Example 105: Preparation of compound 105

Compound 105 was prepared in a manner similar to that described in Example 1
15 in a 79% yield.

1H NMR ($CDCl_3$) δ 0.93-0.98 (m, 6H), 1.05 (d, 3H, $J = 5.4$ Hz), 1.26 (s, 9H),
1.53-2.11 (m, 7H), 2.37 (br, 2H), 3.28-3.33 (m, 2H), 4.16 (br, 2H), 4.50-4.52 (m, 2H),
4.62 (d, 2H, $J = 5.7$ Hz), 5.10 (q, 2H, $J = 6.6$ Hz), 5.23 (d, 1H, $J = 10.2$ Hz), 5.31 (d, 1H, J
= 17.4 Hz), 5.85-5.98 (m, 3H), 6.86 (dd, 1H, $J = 15.6$ Hz, 5.4 Hz), 7.35-7.41 (m, 5H),
20 7.69-7.74 (m, 1H).

ESI-MS ($M+H^+$) = 643.

Example 106: Preparation of compound 106

Compound 106 was prepared in a manner similar to that described in Example 1
25 in a 92% yield.

1H NMR ($CDCl_3$) δ 0.92-0.97 (m, 6H), 1.1 (d, 3H, $J = 6$ Hz), 1.28 (s, 9H), 1.52-
2.19 (m, 6H), 2.43 (br, 2H), 3.19 (s, 3H), 3.28-3.30 (m, 2H), 3.78 (s, 3H), 4.17-4.19 (m,
2H), 4.38 (br, 1H), 4.90-4.95 (m, 1H), 5.10 (q, 2H, $J = 8.7$ Hz), 5.96 (m, 2H), 7.30-7.35
(m, 5H), 7.44 (d, 1H, $J = 7.8$ Hz).

30 ESI-MS ($M+H^+$) = 620

Example 107: Preparation of compound 107

Compound 107 was prepared in a 88 % yield by reducing compound 103 with LiBH₄ (1.5 eq.).

¹H NMR(CDCl₃) δ 0.92-0.99(m, 6H), 1.26(s, 9H), 1.66-1.73(m, 5H), 2.21-2.50(m, 2H), 3.20-3.32(m, 2H), 3.62(m, 1H), 3.75(d, J=7.5 Hz, 2H), 4.42(br, 1H), 4.58(br, 1H), 5.10(s, 2H), 7.21-7.34(m, 4H), 7.61(d, J=7.5 Hz, 1H).

ESI-MS (M+H⁺) = 563.

Example 108: Preparation of compound 108

Compound 108 was prepared in a manner similar to that described in Example 1.

ESI-MS (M+H⁺) = 612.

Example 109: Preparation of compound 109

Compound 109 was prepared as follows: Dry THF (5 ml) and 1-methyl-1-propenyl magnesium bromide (0.5 M in THF, 22 ml) were added to compound 106 (0.124 g, 0.2 mmol) under N₂. The mixture was stirred at room temperature under N₂ for 30 minutes, and then quenched by slowly adding a 10% HCl solution until the pH value was about 7. The mixture was then washed with brine and extracted with EtOAc (3x50 ml). The organic layers were combined, and dried with Na₂SO₄. The crude product was purified by silica gel chromatography using 2% MeOH/CH₂Cl₂ as an eluant to afford compound 109 as a white solid (0.085 mg, 69%).

¹H NMR (CDCl₃) δ 0.91-0.96 (m, 6H), 1.09 (d, 3H, J = 6 Hz), 1.28 (s, 9H), 1.49-1.83 (m, 5H), 1.79 (s, 3H), 1.89 (d, 3H, J = 6.9 Hz), 2.03-2.13 (m, 1H), 2.30-2.41 (m, 1H), 2.48 (m, 1H), 3.25-3.30 (m, 2H), 4.17-4.19 (m, 2H), 4.38-4.39 (m, 1H), 5.11 (q, 2H, J = 7.8 Hz), 5.33-5.38 (m, 1H), 5.92-5.97 (m, 2H), 6.85 (q, 1H, J = 6.3 Hz), 7.13-7.15 (m, 1H), 7.30-7.35 (m, 5H), 7.42 (d, 1H, J = 7.8 Hz).

ESI-MS (M+H⁺) = 615.

Example 110: Preparation of compound 110

Compound 110 was prepared in a manner similar to that of compound 109 in a 44% yield.

$^1\text{H NMR}$ (CDCl_3) δ 0.95 (m, 6H), 1.09 (d, 3H, $J = 6.6$ Hz), 1.27 (s, 9H), 1.52-1.85 (m, 5H), 2.05-2.14 (m, 1H), 2.38-2.42 (m, 2H), 3.28-3.33 (m, 2H), 4.17-4.19 (m, 2H), 4.40-4.42 (m, 1H), 4.79-4.84 (m, 1H), 5.11 (q, 2H, $J = 7.8$ Hz), 5.84-5.92 (m, 3H), 6.36-6.56 (m, 2H), 7.31-7.35 (m, 5H), 7.42 (d, 1H, $J = 7.8$ Hz), 7.61 (d, 1H, $J = 7.2$ Hz).

5 ESI-MS ($\text{M}+\text{H}^+$) = 587.

Example 111: Preparation of compound 111

Compound 111 was prepared in a manner similar to that of compound 109.

10 $^1\text{H NMR}$ (CDCl_3) δ 0.96 (m, 6H), 1.08 (d, 3H, $J = 5.7$ Hz), 1.27 (s, 9H), 1.54-1.78 (m, 5H), 2.03-2.07 (m, 1H), 2.25 (s, 3H), 2.40 (m, 2H), 3.28-3.32 (m, 2H), 4.16-4.19 (m, 2H), 4.60-4.61 (m, 1H), 5.11 (q, 2H, $J = 6.3$ Hz), 5.88 (d, 1H, $J = 4.8$ Hz), 6.02 (s, 1H), 6.16 (d, 1H, $J = 16.2$ Hz), 6.66 (dd, 1H, $J = 15.9$ Hz, 5.1 Hz), 7.35 (m, 5H), 7.73 (d, 1H, $J = 7.5$ Hz).

15 ESI-MS ($\text{M}+\text{H}^+$) = 601.

Example 112: Preparation of compound 112

Compound 112 was prepared in a 99% yield by reacting compound 107 with SO_3Py (2.5 eq.) via a Swern oxidation reaction for 1 hour.

20 $^1\text{H NMR}$ (CDCl_3) δ 0.92-0.98 (m, 6H), 1.26 (s, 9H), 1.66-1.73 (m, 5H), 2.21-2.50 (m, 2H), 3.20-3.32 (m, 2H), 4.42 (br, 2H), 4.59 (br, 1H), 5.11 (s, 2H), 7.21-7.34 (m, 4H), 7.61 (d, $J=7.5$ Hz, 1H), 9.82 (s, 1H).

ESI-MS ($\text{M}+\text{H}^+$) = 561.

Example 113: Preparation of compound 113


25 Compound 113 was prepared in a 68% yield in a manner similar to that of compound 89 using compound 102 as a starting material.

$^1\text{H NMR}$ (CDCl_3) δ 0.94-0.99 (m, 6H), 1.25 (s, 9H), 1.66-1.73 (m, 5H), 2.22-2.51 (m, 4H), 3.19-3.30 (m, 2H), 4.26 (t, $J=6.9$ Hz, 2H), 4.44 (br, 1H), 4.57 (br, 1H), 5.12 (s, 2H), 6.82 (d, $J = 5.1$ Hz, 1H), 7.22-7.34 (m, 4H), 7.61 (d, $J=7.5$ Hz, 1H).

30 ESI-MS ($\text{M}+\text{H}^+$) = 657.

Example 114: Preparation of compound 114

Methylsulfoxide and triethylamine were added to a solution of compound 107. The solution was cooled to 0-5°C with an ice-water bath, followed by addition of sulfur trioxide-pyridine complex. The reaction mixture was stirred at that temperature for 1



5 hour. After adding and stirring the reaction mixture at ambient temperature for another 3 hours, the reaction mixture was quenched by addition of saturated brine and subsequently extracted with ethyl acetate. The organic layers were combined, dried over MgSO₄, filtered, and concentrated to afford a dark red oil. The oil was purified through a chromatography (50% EA in hexane) to afford compound 114 as a white solid.

10 ESI-MS (M+H⁺) = 627.

Example 115: Preparation of compound 115

Compound 115 was prepared in a manner similar to that described in Example 1 in a 65% yield.

15 ¹H NMR (CDCl₃) δ 0.90-0.95 (m, 6H), 1.03-1.05 (m, 4H), 1.23 (s, 9H), 1.44-1.88 (m, 5H), 1.97 (m, 2H), 2.37 (m, 2H), 3.24-3.32 (m, 2H), 3.69 (s, 3H), 4.14 (m, 1H), 4.40-4.52 (m, 2H), 5.07 (s, 2H), 5.88-5.91 (m, 1H), 6.23 (m, 1H), 7.33 (m, 5H), 7.45 (m, 1H), 7.77 (m, 1H).

ESI-MS (M+H⁺) = 591.

20

Example 116: Preparation of compound 116

Compound 116 was prepared in a 77% yield in a manner similar to that of compound 109 using compound 106 as a starting material.

25 ¹H NMR (CDCl₃) δ 0.94-0.99 (m, 6H), 1.09 (d, 3H, J = 6 Hz), 1.27 (s, 9H), 1.53-1.84 (m, 5H), 2.02-2.12 (m, 1H), 2.20 (s, 3H), 2.32-2.43 (m, 2H), 3.24-3.35 (m, 2H), 5.11 (q, 2H, J = 6 Hz), 5.90 (d, 1H, J = 3 Hz), 5.98 (m, 1H), 7.32-7.36 (m, 5H), 7.42 (d, 1H, J = 6 Hz), 7.83 (d, 1H, J = 9 Hz).

ESI-MS (M+H⁺) = 575.

Example 117: Preparation of compound 117

Compound 117 was prepared by the procedures described below:

Triethylamine (0.25 ml, 1.757 mmol, 4.53 eq.) was added to a solution of Intermediate 4 described in Example 1 (0.100g, 0.387 mmol, 1 eq.) in DMSO (1.1 ml).

5 The solution was cooled to 0°C with an ice bath, followed by addition of sulfur trioxide-pyridine complex (0.568 g, 1.742 mmol, 4.5 eq.). The ice bath was removed after the addition and the reaction mixture was stirred at room temperature for 1 hour. 1-Acetyl-3-(triphenyl-15-phosphanylidene)-pyrrolidin-2-one (0.600g, 1.548 mmol, 4 eq.) was then added into the mixture. After stirring the mixture at room temperature overnight, the
10 reaction was quenched by saturated brine (7 ml) and then extracted with ethyl acetate (3 x 10 ml). The organic layers were combined, washed with saturated brine (3 x 15 ml), dried over MgSO₄, filtered, and concentrated to afford a dark red oil. The oil was purified through a column chromatography (2% MeOH in CH₂Cl₂) to afford Intermediate 9 as a white solid (75 mg, 53%). ESI-MS (M+H⁺) = 366.

15 Intermediate 9 (0.446 g, 1.220 mmol, 1eq.) was dissolved in a solution of HCl in 1,4-dioxane (4.0 M, 10 ml) was added to. The solution was stirred at room temperature for 30 minutes and 1,4-dioxane was removed under vacuum. CH₂Cl₂ (1.5 ml) was then added to the residue, followed by addition of *N*-methylmorpholine (0.540 ml, 4.880 mmol, 4 eq.), *Z*-Thr(*t*-Bu)-Leu-OH (0.515 g, 1.220 mmol, 1.0 eq.), HOBt (0.250 g, 1.830
20 mmol, 1.5 eq.), and EDC (0.351 g, 1.830 mmol, 1.5 eq.). The reaction mixture was stirred at room temperature overnight and then concentrated under vacuum. The residue was partitioned between a 10% KHSO₄ (6.5 ml) aqueous solution and CH₂Cl₂ (45 ml). The organic layer was collected, dried over MgSO₄, filtered, and concentrated. The crude product was purified by flash chromatography (2% MeOH in CH₂Cl₂) to afford
25 compound 117 (0.572 g, 70%) as a white solid. ¹H NMR (CDCl₃) δ 0.88-0.98 (m, 6H), 1.07 (d, J=6.0 Hz, 3H), 1.25(s, 9H), 1.48-1.62 (m, 3H), 1.74-1.90 (m, 3H), 2.06-2.16 (m, 3H), 2.32-2.47 (m, 3H), 2.54 (s, 3H), 2.71 (m, 1H), 3.03 (m, 1H), 3.27-3.30 (m, 2H), 3.77 (dd, J=7.5, 7.5 Hz, 2H), 4.14-4.18 (m, 2H), 4.53 (m, 1H), 4.60 (m, 1H), 5.00-5.18 (m, 2H), 5.87 (d, J=5.1 Hz, 1H), 6.02 (br s, 1H), 6.43 (d, J=9.0 Hz, 1H), 7.27-7.34 (m, 5H),
30 7.76(d, J=6.9 Hz, 1H). ESI-MS (M+H⁺) = 670.

Example 118: Preparation of compound 118

Compound 118 was prepared in a manner similar to that described in Example 1.

$^1\text{H NMR}$ (CDCl_3) δ 0.9 (m, 6H), 1.16 (s, 9H), 1.49-2.17 (m, 8H), 2.38-2.41 (m, 2H), 3.32-3.35 (m, 2H), 3.58-3.70 (m, 2H), 4.27-4.28 (m, 1H), 4.37 (t, 3H, $J = 7.5$ Hz), 4.57 (br, 1H), 5.10 (q, 2H, $J = 5.7$ Hz), 6.30 (s, 1H), 6.49-6.52 (m, 2H), 6.63-6.65 (m, 1H), 7.16 (d, 1H, $J = 8.4$ Hz), 7.34 (br, 5H), 8.17 (d, 1H, $J = 6.3$ Hz).

ESI-MS ($\text{M}+\text{H}^+$) = 642.

Example 119: Preparation of compound 119

Compound 119 was prepared in a manner similar to that described in Example 1.

ESI-MS ($\text{M}+\text{H}^+$) = 657.

Example 120: Preparation of compound 120

Compound 120 was prepared in a manner similar to that described in Example 1.

$^1\text{H NMR}$ (CDCl_3) δ 0.85-0.90 (m, 6H), 1.39 (s, 9H), 1.56-1.81 (m, 5H), 2.01-2.09 (m, 4H), 2.41 (m, 1H), 3.16 (s, 3H), 3.28-3.30 (m, 2H), 3.44-3.48 (m, 2H), 3.78 (s, 3H), 4.30-4.40 (m, 2H), 4.83-4.88 (m, 1H), 5.09 (s, 2H), 5.64 (br s, 1H), 6.32 (br s, 1H), 7.26-7.33 (m, 5H), 7.77-7.79 (m, 1H).

ESI-MS ($\text{M}+\text{H}^+$) = 645.

Example 121: Preparation of compound 121

Compound 121 was prepared in a manner similar to that described in Example 1 in a 10% yield.

$^1\text{H NMR}$ (CDCl_3) δ 0.87 (m, 6H), 1.26 (s, 9H), 1.53-1.76 (m, 5H), 1.99 (m, 6H), 2.38 (m, 2H), 2.88 (m, 1H), 3.14-3.48 (m, 4H), 4.14 (m, 1H), 4.35 (m, 2H), 4.56 (m, 1H), 5.07 (s, 2H), 6.40-6.46 (m, 1H), 7.04 (m, 1H), 7.31 (m, 5H), 8.08 (m, 1H).

ESI-MS ($\text{M}+\text{H}^+$) = 657.

Example 122: Preparation of compound 122

Compound 122 was prepared in a manner similar to that of compound 109 in a 60% yield.

^1H NMR (CDCl_3) δ 0.93 (m, 6H), 1.16 (s, 9H), 1.45-1.82 (m, 5H), 2.00-2.09 (m, 1H), 2.45 (s, 3H), 2.37-2.46 (m, 2H), 3.31-3.34 (m, 2H), 3.64-3.68 (m, 2H), 4.27-4.29 (m, 1H), 4.44-4.50 (m, 1H), 4.59-4.60 (m, 1H), 5.10 (q, 2H, $J = 5.1$ Hz), 6.10 (s, 1H), 6.17 (d, 1H, $J = 1.2$ Hz), 6.48 (br, 1H), 6.68 (dd, 2H, $J = 15.6$ Hz, 5.7 Hz), 7.13 (d, 1H, $J = 8.1$ Hz), 7.34 (br, 5H), 8.01 (d, 1H, $J = 6.9$ Hz).

ESI-MS ($\text{M}+\text{H}^+$) = 614.

Example 123: Preparation of compound 123

Compound 123 was prepared in a manner similar to that of compound 109 in a 32% yield.

^1H NMR (CDCl_3) δ 0.92-0.98 (m, 6H), 1.06-1.11 (m, 6H), 1.26 (s, 9H), 1.53-2.10 (m, 6H), 2.39 (m, 2H), 2.56 (q, 2H, $J = 7.2$ Hz), 3.28-3.31 (m, 2H), 4.15-4.18 (m, 2H), 4.44-4.45 (m, 1H), 4.58 (m, 1H), 5.11 (q, 2H, $J = 6.3$ Hz), 5.89 (d, 1H, $J = 4.8$ Hz), 6.04 (s, 1H), 6.19 (d, 1H, $J = 15.3$ Hz), 6.69 (dd, 1H, $J = 15.9$ Hz, 5.4 Hz), 7.35 (m, 5H), 7.71 (d, 1H, $J = 7.5$ Hz).

ESI-MS ($\text{M}+\text{H}^+$) = 615.

Example 124: Preparation of compound 124

Compound 124 was prepared in a manner similar to that described in Example 1.

ESI-MS ($\text{M}+\text{H}^+$) = 615.

Example 125: Preparation of compound 125

Compound 125 was prepared in a manner similar to that of compound 109 in a 55% yield.

^1H NMR (CDCl_3) δ 0.91(q, 6H, $J = 6$ Hz), 1.41 (s, 9H), 1.55-1.81 (m, 5H), 2.10 (m, 1H), 2.23 (s, 3H), 2.36-2.40 (m, 2H), 2.78 (m, 2H), 3.30-3.33 (m, 2H), 4.46-4.52 (m, 1H), 4.56-4.59 (m, 2H), 5.11 (q, 2H, $J = 6.6$ Hz), 6.01 (d, 1H, $J = 8.4$ Hz), 6.15 (d, 1H, $J = 15.6$ Hz), 6.30 (s, 1H), 6.65 (dd, 1H, $J = 16.2$ Hz, 5.4 Hz), 7.17 (d, 1H, $J = 8.4$ Hz), 7.34 (m, 5H), 7.89 (d, 1H, $J = 7.8$ Hz).

ESI-MS ($\text{M}+\text{Na}^+$) = 637.

Example 126: Preparation of compound 126

Compound 126 was prepared in a manner similar to that of compound 109 in an 82% yield.

^1H NMR (CDCl_3) δ 0.86-0.97 (m, 6H), 1.14-1.38 (m, 6H), 1.49-1.66 (m, 1H),
5 1.98-2.08 (m, 2H), 2.23 (s, 3H), 2.34-2.45 (m, 2H), 3.29-3.33 (m, 2H), 3.62-3.68 (m,
2H), 4.26 (dd, $J=9.9, 5.4$ Hz, 1H), 4.43 (m, 1H), 4.59 (m, 1H), 5.04-5.14 (m, 2H), 6.08-
6.18 (m, 2H), 6.37 (br s, 1H), 6.64-6.71 (dd, $J=16.2, 5.4$ Hz, 2H), 7.13 (d, $J=8.4$ Hz, 1H),
7.28-7.34 (m, 5H), 7.98 (d, $J=6.9$ Hz, 1H).

ESI-MS ($\text{M}+\text{H}^+$) = 640.

10

Example 127: Preparation of compound 127

Compound 127 was prepared in a manner similar to that of compound 109 in a 78% yield.

^1H NMR (CDCl_3) δ 0.85-0.93 (m, 6H), 0.98 (s, 9H), 1.48-1.64 (m, 4H), 1.75-1.85
15 (m, 1H), 1.91 (m, 1H), 1.98-2.08 (m, 1H), 2.03 (s, 3H), 2.23 (s, 3H), 2.37-2.46 (m, 2H),
3.29-3.36 (m, 2H), 3.54-3.72 (m, 2H), 4.24 (m, 1H), 4.44 (m, 1H), 4.58 (m, 1H), 5.02-
5.12 (m, 2H), 6.15 (d, $J=15.9$ Hz, 1H), 6.55-6.78 (m, 3H), 7.28-7.31 (m, 5H), 7.97 (d,
 $J=6.6$ Hz, 1H).

ESI-MS ($\text{M}+\text{H}^+$) = 628.

20

Example 128: Preparation of compound 128

Compound 128 was prepared in a manner similar to that of compound 109 in a 25% yield.

^1H NMR (CDCl_3) δ 0.89 (m, 6H), 1.39 (s, 9H), 1.53-2.14 (m, 7H), 2.22 (s, 3H),
25 2.39 (m, 2H), 3.29-3.30 (m, 2H), 3.46 (m, 2H), 4.21 (m, 2H), 4.43 (m, 1H), 4.58 (m, 1H),
5.09 (s, 2H), 6.18 (m, 2H), 6.46 (m, 1H), 6.62-6.69 (m, 1H), 7.32 (m, 5H), 7.93 (m, 1H).

ESI-MS ($\text{M}+\text{H}^+$) = 630.

Example 129: Preparation of compound 129

30 Compound 129 was prepared in a manner similar to that of compound 112 in a 38% yield.

$^1\text{H NMR}(\text{CDCl}_3) \delta$ 0.92-0.98(m, 6H), 1.27(s, 9H), 1.65-1.72(m, 5H), 2.22-2.51(m, 2H), 3.20-3.32(m, 2H), 4.43(br, 2H), 4.59(br, 1H), 5.12(s, 2H), 7.21-7.35(m, 4H), 7.60(d, $J=7.5$ Hz, 1H), 9.82(s, 1H)

ESI-MS ($\text{M}+\text{H}^+$) = 604.3.

5

Example 130: Preparation of compound 130

Compound 130 was prepared in a manner similar to that of compound 131.

$^1\text{H NMR}(\text{CDCl}_3) \delta$ 0.93-0.98(m, 6H), 1.26(s, 9H), 1.65-1.72(m, 5H), 2.22-2.51(m, 2H), 3.20-3.32(m, 2H), 3.58(d, $J=7.2$ Hz, 2H), 3.86(m, 1H), 4.44(br, 2H), 5.12(s, 2H), 7.21-7.35(m, 4H), 7.60(d, $J=7.5$ Hz, 1H).

10

ESI-MS ($\text{M}+\text{H}^+$) = 624.3.

Example 131: Preparation of compound 131

Compound 131 was prepared in a 51% yield by reacting compound 107 with thionyl chloride in CH_2Cl_2 at room temperature for 6 hours.

15

$^1\text{H NMR}(\text{CDCl}_3) \delta$ 0.93-0.98(m, 6H), 1.28(s, 9H), 1.65-1.72(m, 5H), 2.22-2.50(m, 2H), 3.20-3.32(m, 2H), 3.58(d, $J=7.2$ Hz, 2H), 3.86(m, 1H), 4.42(br, 2H), 5.11(s, 2H), 7.21-7.35(m, 4H), 7.60(d, $J=7.5$ Hz, 1H).

ESI-MS ($\text{M}+\text{H}^+$) = 581.3.

20

Example 132: Preparation of compound 132

1-(2,2,2-Trifluoro-1-methyl-ethylidene)-pyrrolidinium (1.5eq.) was prepared from methyltrifluoromethylketone and pyrrolidine at 0°C . Compound 112 (1.0eq.) was then added in situ to afford compound 132 in a 43% yield.

25

$^1\text{H NMR}(\text{CDCl}_3) \delta$ 0.94-0.98(m, 6H), 1.27(s, 9H), 1.66-1.73(m, 5H), 2.25-2.53(m, 2H), 3.22-3.31(m, 2H), 4.44(m, 1H), 4.63(m, 1H), 5.10(s, 2H), 6.21(d, $J=18.1$ Hz, 1H), 6.98(dd, $J=18.1$ Hz, 5.1 Hz, 1H), 7.22-7.35(m, 4H), 7.62(d, $J=7.5$ Hz, 1H).

ESI-MS ($\text{M}+\text{H}^+$) = 655.

30

Example 133: Preparation of compound 133

Compound 133 was prepared in a manner similar to that of compound 112 in a 35% yield.

$^1\text{H NMR}(\text{CDCl}_3)$ δ 0.89-0.95 (m, 6H), 1.42 (s, 9H), 1.55-2.06 (m, 6H), 2.37-2.47 (m, 2H), 2.64-2.94 (m, 2H), 3.32-3.34 (m, 2H), 4.45-4.58 (m, 3H), 5.12 (s, 2H), 6.00-6.16 (m, 2H), 7.35 (m, 5H), 9.49 (d, 1H, $J = 16.5$ Hz).

ESI-MS ($\text{M}+\text{H}^+$) = 575.

Example 134: Preparation of compound 134

Compound 134 was prepared in a manner similar to that of compound 112 in a 35% yield.

$^1\text{H NMR}(\text{CDCl}_3)$ δ 0.87-1.26 (m, 20H), 1.69-1.97 (m, 14H), 2.43 (m, 1H), 3.64 (m, 1H), 4.18 (m, 2H), 4.40 (m, 1H), 5.09 (s, 2H), 5.90 (m, 1H), 7.36 (m, 5H), 9.50 (s, 1H).

ESI-MS ($\text{M}+\text{H}^+$) = 601.

Example 135: Preparation of compound 135

Compound 135 was prepared in a manner similar to that of compound 109.

$^1\text{H NMR}(\text{CDCl}_3)$ δ 0.93 (q, 6H, $J = 5.7$ Hz), 1.03-1.08 (m, 9H), 1.23 (s, 9H), 1.49-1.83 (m, 5H), 2.00-2.03 (m, 1H), 2.36 (m, 2H), 2.70-2.79 (m, 1H), 3.26-3.29 (m, 2H), 4.13-4.17 (m, 2H), 4.43-4.44 (m, 1H), 4.58-4.60 (m, 1H), 5.09 (q, 2H, $J = 5.4$ Hz), 5.86 (d, 1H, $J = 5.1$ Hz), 6.01 (s, 1H), 6.26 (d, 1H, $J = 15.6$ Hz), 6.71 (dd, 1H, $J = 15.6$ Hz, 5.4 Hz), 7.30-7.34 (m, 5H), 7.60 (d, 1H, $J = 7.8$ Hz).

ESI-MS ($\text{M}+\text{H}^+$) = 629.

Example 136: Preparation of compound 136

Compound 136 was prepared in a manner similar to that of compound 109 in a 36% yield.

$^1\text{H NMR}(\text{CDCl}_3)$ δ 0.93 (q, 6H, $J = 5.7$ Hz), 1.06 (d, 3H, $J = 6.3$ Hz), 1.23 (s, 9H), 1.44-1.77 (m, 5H), 1.99-2.07 (m, 1H), 2.39 (m, 2H), 3.27-3.30 (m, 2H), 4.15-4.18 (m, 2H), 4.39-4.41 (m, 1H), 4.62 (m, 1H), 5.10 (q, 2H, $J = 6$ Hz), 5.82-5.87 (m, 3H), 6.28

(d, 1H, J = 16.5 Hz). 6.47-6.56 (m, 2H), 6.77 (dd, 1H, J = 15.3 Hz, 4.8 Hz), 7.29-7.34 (m, 5H), 7.66 (d, 1H, J = 7.5 Hz).

ESI-MS (M+Na⁺) = 635.

5 Example 137: Preparation of compound 137

Compound 137 was prepared in a manner similar to that of compound 109.

¹H NMR (CDCl₃) δ 0.94-0.97 (m, 6H), 1.08 (d, 3H, J = 6.3 Hz), 1.27 (s, 9H), 1.52-1.86 (m, 11H), 2.02-2.09 (m, 1H), 2.40 (m, 2H), 3.29-3.32 (m, 2H), 4.16 (m, 2H), 4.41 (m, 1H), 4.62 (m, 1H), 5.11 (q, 2H, J = 5.4 Hz), 5.81-5.89 (m, 2H), 6.61-6.79 (m, 10
3H), 7.35 (m, 5H), 7.53 (d, 1H, J = 7.2 Hz).

ESI-MS (M+H⁺) = 641.

Example 138: Preparation of compound 138

Compound 138 was prepared in a manner similar to that of compound 109 in a
15 26% yield.

¹H NMR (CDCl₃) δ 0.87-1.26 (m, 20H), 1.52-1.72 (m, 9H), 2.21-2.22 (m, 3H), 2.38 (m, 2H), 3.27-3.30 (m, 2H), 4.15 (m, 2H), 4.39 (m, 1H), 4.58 (m, 1H), 5.09 (s, 2H), 5.86 (m, 2H), 6.12-6.22 (m, 1H), 6.61-6.68 (m, 1H), 7.33 (m, 5H), 7.65 (m, 1H).

ESI-MS (M+H⁺) = 641.

20

Example 139: Preparation of compound 139

Compound 139 was prepared in a manner similar to that of compound 109 in a
28% yield.

¹H NMR (CDCl₃) δ 0.62-0.70 (m, 8H), 0.79-0.80 (m, 6H), 0.99 (s, 9H), 1.26-1.85
25 (m, 6H), 2.12 (m, 2H), 3.00-3.03 (m, 2H), 4.16 (m, 2H), 4.46 (m, 1H), 4.62 (m, 1H), 5.06-5.16 (m, 2H), 5.89 (s, 1H), 6.10 (s, 1H), 6.32 (d, 1H, J = 15.6 Hz), 6.74 (dd, 1H, J = 15.6 Hz, 5.1 Hz), 7.36 (m, 5H), 7.69 (m, 1H).

ESI-MS (M+Na⁺) = 649.

30

Example 140: Preparation of compound 140

Compound 140 was prepared in a manner similar to that of compound 109 in a 37% yield.

^1H NMR (CDCl_3) δ 0.93 (q, 6H, $J = 6$ Hz), 1.04 (d, 3H, $J = 6$ Hz), 1.23 (s, 9H),
5 1.49-1.83 (m, 5H), 1.89 (s, 3H), 1.99-2.09 (m, 1H), 2.37-2.42 (m, 2H), 3.27 (d, 2H, $J = 9$
Hz), 4.12-4.17 (m, 2H), 4.39-4.44 (m, 1H), 4.58-4.63 (m, 1H), 5.09 (q, 2H, $J = 5.4$ Hz),
5.78 (s, 1H), 5.86 (d, 1H, $J = 5.1$ Hz), 5.94 (s, 1H), 6.00 (m, 2H), 6.72-6.74 (m, 2H),
7.30-7.34 (m, 5H), 7.63 (d, 1H, $J = 7.5$ Hz).

ESI-MS ($\text{M}+\text{H}^+$) = 627.

10

Example 141: Preparation of compound 141

Compound 141 was prepared in a manner similar to that of compound 109 in a 82% yield.

^1H NMR (CDCl_3) δ 0.86-1.02 (m, 15H), 1.09-1.31 (m, 9H), 1.45-1.85 (m, 6H),
15 2.16-2.24 (m, 2H), 2.22 (s, 3H), 2.30-2.41 (m, 2H), 3.22-3.32 (m, 2H), 4.10 (dd, $J=5.4,$
5.4 Hz, 1H), 4.23 (ddd, $J=8.1, 6.0, 2.4$ Hz, 1H), 4.46 (m, 1H), 4.63(m, 1H), 4.99-5.14 (m,
2H), 5.38 (d, $J=5.1$ Hz, 1H), 5.88 (br s, 1H), 6.69 (dd, $J=15.9, 5.1$ Hz, 1H), 7.14 (d, $J=6.0$
Hz, 1H), 7.32-7.37 (m, 5H), 7.43 (m, 1H).

ESI-MS ($\text{M}+\text{H}^+$) = 700.

20

Example 142: Preparation of compound 142

Compound 142 was prepared in a manner similar to that described in Example 1.

^1H NMR (CDCl_3) δ 0.73-0.87 (m, 5H), 1.03 (d, $J = 6.0$ Hz, 3H), 1.05 (d, $J = 6.0$
Hz, 3H), 1.14 (s, 3H), 1.22-1.28 (m, 18H), 1.77-1.82 (m, 1H), 2.04-2.12 (m, 1H), 2.32-
25 2.48 (m, 2H), 3.28-3.32 (m, 2H), 4.13-4.19 (m, 6H), 4.32-4.35 (m, 1H), 4.69-4.78 (m,
1H), 5.05-5.16 (m, 2H), 5.50 (br s, 1H), 5.89 (d, $J = 18, 1\text{H}$), 6.13 (d, $J = 16$ Hz, 1H),
6.85 (dd, $J = 16.0, 5.5$ Hz, 1H), 7.08 (d, $J = 9.0$ Hz, 1H), 7.34 (m, 5H), 7.63 (d, $J = 6.0$
Hz, 1H).

ESI-MS ($\text{M}+\text{Na}^+$) = 675.

30

Example 143: Preparation of compound 143

Compound 143 was prepared in a manner similar to that of compound 109 in a 82 % yield.

$^1\text{H NMR}$ (CDCl_3) δ 0.84-0.94 (m, 6H), 1.10 (d, $J=6.0$ Hz, 3H), 1.15-1.24 (m, 5H),
5 1.49-1.79 (m, 10H), 2.03 (m, 1H), 2.22 (s, 3H), 2.38 (m, 2H), 3.28-3.31 (m, 2H), 3.39 (m,
1H), 4.15 (m, 2H), 4.45-4.58 (m, 2H), 5.09 (s, 2H), 5.75 (d, $J=5.7$ Hz, 1H), 5.90 (br
s, 1H), 6.15 (d, $J=15.9$ Hz, 1H), 6.65 (dd, $J=15.9, 5.4$ Hz, 1H), 7.12 (d, $J=7.8$ Hz, 1H),
7.30-7.34 (m, 5H), 7.75 (d, $J=7.5$ Hz, 1H).

ESI-MS ($\text{M}+\text{H}^+$) = 627.

10

Example 144: Preparation of compound 144

Compound 144 was prepared in a manner similar to that described in Example 1.

$^1\text{H NMR}$ (CDCl_3) δ 0.86-1.27 (m, 17H), 1.47-1.76 (m, 14H), 1.97-2.05 (m, 1H),
2.38 (br s, 2H), 3.28 (m, 2H), 4.15 (m, 4H), 4.38 (m, 1H), 4.58 (m, 1H), 5.04-5.14 (m,
15 2H), 5.79-5.93 (m, 3H), 6.80 (dd, 1H, $J=15.9$ Hz, 5.7 Hz), 7.34 (m, 5H), 7.60 (m, 1H).

ESI-MS ($\text{M}+\text{H}^+$) = 671.

Example 145: Preparation of compound 145

Compound 145 was prepared in a manner similar to that of compound 109 in a
20 25% yield.

$^1\text{H NMR}$ (CDCl_3) δ 0.73-0.87 (m, 5H), 0.95 (d, $J=6.0$ Hz, 3H), 1.14 (s, 9H),
1.47-1.69 (m, 13H), 1.90-1.99 (m, 3H), 2.00-2.36 (m, 3H), 3.17 (m, 2H), 4.05 (m, 2H),
4.30 (m, 1H), 4.51 (m, 1H), 4.94-5.04 (m, 2H), 5.77-5.81 (m, 2H), 6.21 (d, $J=16.0$ Hz
1H), 6.62 (dd, $J=16.0, 5.5$ Hz, 1H), 7.34 (m, 5H), 7.52 (d, $J=7.8$ Hz, 1H).

25

ESI-MS ($\text{M}+\text{Na}^+$) = 667.

Example 146: *in vitro* assay

A fusion protein, prepared by fusing a severe acute respiratory syndrome 3CL protease to *E. coli* maltose-binding protein (MBP), was expressed in *E. coli* BL21 (DE3)
30 pLys S cells (Novagen, Oakland, CA). Fusion protein thus obtained was purified by amylose-affinity chromatography and cleaved with factor Xa to release the severe acute

respiratory syndrome 3CL protease. Subsequently, the recombinant protease was purified to homogeneity using phenyl Sepharose CL-4B column (Pharmacia, Uppsala, Sweden) and was concentrated to form a 25 μ M solution.

The enzymatic activity of severe acute respiratory syndrome 3CL protease (75 nM) was determined by incubation with a solution containing 15 μ M of a substrate peptide (SITSAVLQSGFRKMA, SEQ ID No: 1) at 25°C for 30 minutes in a medium containing 20 mM Tris-HCl (pH 7.5), 200 mM NaCl, 1 mM EDTA, 1 mM dithiothreitol, and 1 mg/mL bovine serum albumin. The reaction was terminated by adding an equal volume of 0.2% trifluoroacetic acid. The reaction mixture was analyzed by reverse-phase HPLC using a C18 column. Cleaved products were resolved using a 5-95% linear gradient of acetonitrile in 0.9% trifluoroacetic acid. Quantification of peak areas was used to determine the extent of substrate conversion.

Compounds 1-145 were tested for their efficacy in inhibiting severe acute respiratory syndrome 3CL protease. Specifically, a test compound and the severe acute respiratory syndrome 3CL protease were pre-incubated at 25°C for 20 minutes before they were incubated with the substrate peptide. Unexpectedly, 77 compounds show IC_{50} values lower than 1 μ M, 32 compounds show IC_{50} values between 1 μ M and 10 μ M, and 12 compounds show IC_{50} values above 10 μ M.


OTHER EMBODIMENTS

All of the features disclosed in this specification may be combined in any combination. Each feature disclosed in this specification may be replaced by an alternative feature serving the same, equivalent, or similar purpose. Thus, unless expressly stated otherwise, each feature disclosed is only an example of a generic series of equivalent or similar features.

From the above description, one skilled in the art can easily ascertain the essential characteristics of the present invention, and without departing from the spirit and scope thereof, can make various changes and modifications of the invention to adapt it to various usages and conditions. Thus, other embodiments are also within the scope of the following claims.

WHAT IS CLAIMED IS:

1. A method for treating an infection with a virus, comprising administering to a subject in need thereof an effective amount of a compound of the formula (I):



wherein

X is N(R_{a1}), O, or CH₂; or X and one of R₂ and R₃, together with the atom or atoms to which they are bonded, are C₃-C₂₀ heterocycloalkyl; in which R_{a1} is H or C₁-C₁₅ alkyl;

R₁ is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, aryl, C(O)R_{b1}, CO₂R_{b1}, C(O)NR_{b1}R_{b2}, C(O)-N(R_{b1})-OR_{b2}, C(S)R_{b1}, C(S)NR_{b1}R_{b2}, S(O)R_{b1}, SO₂R_{b1}, S(O)NR_{b1}R_{b2}, S(O)-N(R_{b1})-OR_{b2}, SO₂NR_{b1}R_{b2}, or SO₃R_{b1}; in which each of R_{b1} and R_{b2}, independently, is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, or aryl;

each of R₂ and R₃, independently, is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, aryl, OR_{c1}, SR_{c1}, or NR_{c1}R_{c2}; or X and one of R₂ and R₃, together with the atom or atoms to which they are bonded, are C₃-C₂₀ heterocycloalkyl; in which each of R_{c1} and R_{c2}, independently, is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, or aryl;

each of R₄ and R₅, independently, is H, halogen, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, or aryl;

one of R₆ and R₇ is C₁-C₁₅ alkyl substituted with C₃-C₂₀ heterocycloalkyl, heteroaryl, or aryl; and the other of R₆ and R₇ is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, or aryl;


R₈ is H, halogen, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, aryl, OR_{d1}, or SR_{d1}; in which R_{d1} is H and C₁-C₁₅ alkyl; and

each of R₉ and R₁₀, independently, is H, halogen, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, aryl, C(O)R_{e1}, CO₂R_{e1}, C(O)NR_{e1}R_{e2}, C(O)-N(R_{e1})-OR_{e2}, C(S)R_{e1}, C(S)NR_{e1}R_{e2}, CN, NO₂, S(O)R_{e1}, SO₂R_{e1}, S(O)NR_{e1}R_{e2},

S(O)-N(R_{e1})-OR_{e2}, SO₂NR_{e1}R_{e2}, SO₃R_{e1}, PO(OR_{e1})(OR_{e2}), PO(R_{e1})(R_{e2}), PO(NR_{e1}R_{e2})(OR_{e3}), PO(NR_{e1}R_{e2})(NR_{e3}R_{e4}), C(O)-N(R_{e1})-NR_{e2}R_{e3}, or C(S)-N(R_{e1})-NR_{e2}R_{e3}; or R₈ and R₁₀, taken together, are C₃-C₂₀ cycloalkyl or C₃-C₂₀ heterocycloalkyl; or R₉ and R₁₀, taken together, are C₃-C₂₀ cycloalkyl or C₃-C₂₀ heterocycloalkyl; in which each of R_{e1}, R_{e2}, R_{e3}, and R_{e4}, independently, is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, or aryl; or any two of R_{e1}, R_{e2}, R_{e3}, and R_{e4}, together with the atom or atoms to which they are bonded, are C₃-C₂₀ heterocycloalkyl; and the virus is a coronavirus or a hepatitis C virus.

2. The method of claim 1, wherein X is N(R_{a1}) or O; or X and one of R₂ and R₃, together with the atom or atoms to which they are bonded, are C₃-C₂₀ heterocycloalkyl; R₁ is H, C(O)R_{b1}, CO₂R_{b1}, C(O)NR_{b1}R_{b2}, C(S)NR_{b1}R_{b2}, or SO₂R_{b1}; each of R₂ and R₃, independently, is H or C₁-C₁₅ alkyl; or X and one of R₂ and R₃, together with the atom or atoms to which they are bonded, are C₃-C₂₀ heterocycloalkyl; each of R₄ and R₅, independently, is H or C₁-C₁₅ alkyl; one of R₆ and R₇ is C₁-C₁₅ alkyl substituted with C₃-C₂₀ heterocycloalkyl, and the other of R₆ and R₇ is H; R₈ is H; and each of R₉ and R₁₀, independently, is H or CO₂R_{e1}.

3. The method of claim 2, wherein one of R₆ and R₇ is




4. The method of claim 3, wherein one of R₂ and R₃ is H or C₁-C₁₅ alkyl optionally substituted with halogen, heteroaryl, aryl, OR_{c1}, SR_{c1}, OC(O)R_{c1}, CO₂R_{c1}, C(O)NR_{c1}R_{c2}, NR_{c1}R_{c2}, N(R_{c1})-CO₂R_{c2}, N(R_{c1})-C(O)R_{c2}, N(R_{c1})-C(O)-NR_{c2}R_{c3}, N(R_{c1})-SO₂R_{c2}, SO₂R_{c1}, or O-SO₂-R_{c1}; or X and one of R₂ and R₃, together with the atom or atoms to which they are bonded, are C₃-C₂₀ heterocycloalkyl; in which R_{c3} is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, or aryl.

5. The method of claim 4, wherein one of R_4 and R_5 is H or C_1 - C_{15} alkyl optionally substituted with halogen, aryl, OR_{f1} , SR_{f1} , CO_2R_{f1} , $C(O)NR_{f1}R_{f2}$, SO_2R_{f1} , SO_3R_{f1} , or $NR_{f1}R_{f2}$; in which each of R_{f1} and R_{f2} , independently, is H, C_1 - C_{15} alkyl, C_3 - C_{20} cycloalkyl, C_3 - C_{20} heterocycloalkyl, heteroaryl, or aryl.

6. The method of claim 1, wherein the coronavirus is a severe acute respiratory syndrome virus or a human coronavirus 229E.

7. The method of claim 1, wherein the virus is a hepatitis C virus.

8. A method for inhibiting a viral protease, comprising contacting the protease with an effective amount of a compound of the formula (I):



wherein

X is $N(R_{a1})$, O, or CH_2 ; or X and one of R_2 and R_3 , together with the atom or atoms to which they are bonded, are C_3 - C_{20} heterocycloalkyl; in which R_{a1} is H or C_1 - C_{15} alkyl;

R_1 is H, C_1 - C_{15} alkyl, C_3 - C_{20} cycloalkyl, C_3 - C_{20} heterocycloalkyl, heteroaryl, aryl, $C(O)R_{b1}$, CO_2R_{b1} , $C(O)NR_{b1}R_{b2}$, $C(O)-N(R_{b1})-OR_{b2}$, $C(S)R_{b1}$, $C(S)NR_{b1}R_{b2}$, $S(O)R_{b1}$, SO_2R_{b1} , $S(O)NR_{b1}R_{b2}$, $S(O)-N(R_{b1})-OR_{b2}$, $SO_2NR_{b1}R_{b2}$, or SO_3R_{b1} ; in which each of R_{b1} and R_{b2} , independently, is H, C_1 - C_{15} alkyl, C_3 - C_{20} cycloalkyl, C_3 - C_{20} heterocycloalkyl, heteroaryl, or aryl;

each of R_2 and R_3 , independently, is H, C_1 - C_{15} alkyl, C_3 - C_{20} cycloalkyl, C_3 - C_{20} heterocycloalkyl, heteroaryl, aryl, OR_{c1} , SR_{c1} , or $NR_{c1}R_{c2}$; or X and one of R_2 and R_3 , together with the atom or atoms to which they are bonded, are C_3 - C_{20} heterocycloalkyl; in which each of R_{c1} and R_{c2} , independently, is H, C_1 - C_{15} alkyl, C_3 - C_{20} cycloalkyl, C_3 - C_{20} heterocycloalkyl, heteroaryl, or aryl;


each of R_4 and R_5 , independently, is H, halogen, C_1 - C_{15} alkyl, C_3 - C_{20} cycloalkyl, C_3 - C_{20} heterocycloalkyl, heteroaryl, or aryl;

one of R₆ and R₇ is C₁-C₁₅ alkyl substituted with C₃-C₂₀ heterocycloalkyl, heteroaryl, or aryl; and the other of R₆ and R₇ is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, or aryl;

R₈ is H, halogen, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, aryl, OR_{d1}, or SR_{d1}; in which R_{d1} is H and C₁-C₁₅ alkyl; and

each of R₉ and R₁₀, independently, is H, halogen, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, aryl, C(O)R_{e1}, CO₂R_{e1}, C(O)NR_{e1}R_{e2}, C(O)-N(R_{e1})-OR_{e2}, C(S)R_{e1}, C(S)NR_{e1}R_{e2}, CN, NO₂, S(O)R_{e1}, SO₂R_{e1}, S(O)NR_{e1}R_{e2}, S(O)-N(R_{e1})-OR_{e2}, SO₂NR_{e1}R_{e2}, SO₃R_{e1}, PO(OR_{e1})(OR_{e2}), PO(R_{e1})(R_{e2}), PO(NR_{e1}R_{e2})(OR_{e3}), PO(NR_{e1}R_{e2})(NR_{e3}R_{e4}), C(O)-N(R_{e1})-NR_{e2}R_{e3}, or C(S)-N(R_{e1})-NR_{e2}R_{e3}; or R₈ and R₁₀, taken together, are C₃-C₂₀ cycloalkyl or C₃-C₂₀ heterocycloalkyl; or R₉ and R₁₀, taken together, are C₃-C₂₀ cycloalkyl or C₃-C₂₀ heterocycloalkyl; in which each of R_{e1}, R_{e2}, R_{e3}, and R_{e4}, independently, is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, or aryl; or any two of R_{e1}, R_{e2}, R_{e3}, and R_{e4}, together with the atom or atoms to which they are bonded, are C₃-C₂₀ heterocycloalkyl; and the viral protease is a coronaviral 3CL protease or a hepatitis C viral protease.

9. The method of claim 8, wherein X is N(R_{a1}) or O; or X and one of R₂ and R₃, together with the atom or atoms to which they are bonded, are C₃-C₂₀ heterocycloalkyl; R₁ is H, C(O)R_{b1}, CO₂R_{b1}, C(O)NR_{b1}R_{b2}, C(S)NR_{b1}R_{b2}, or SO₂R_{b1}; each of R₂ and R₃, independently, is H or C₁-C₁₅ alkyl; or X and one of R₂ and R₃, together with the atom or atoms to which they are bonded, are C₃-C₂₀ heterocycloalkyl; each of R₄ and R₅, independently, is H or C₁-C₁₅ alkyl; one of R₆ and R₇ is C₁-C₁₅ alkyl substituted with C₃-C₂₀ heterocycloalkyl, and the other of R₆ and R₇ is H; R₈ is H; and each of R₉ and R₁₀, independently, is H or CO₂R_{e1}.



10. The method of claim 9, wherein one of R₆ and R₇ is


11. The method of claim 10, wherein one of R₂ and R₃ is H or C₁-C₁₅ alkyl optionally substituted with halogen, heteroaryl, aryl, OR_{c1}, SR_{c1}, OC(O)R_{c1}, CO₂R_{c1}, C(O)NR_{c1}R_{c2}, NR_{c1}R_{c2}, N(R_{c1})-CO₂R_{c2}, N(R_{c1})-C(O)R_{c2}, N(R_{c1})-C(O)-NR_{c2}R_{c3}, N(R_{c1})-SO₂R_{c2}, SO₂R_{c1}, or O-SO₂-R_{c1}; or X and one of R₂ and R₃, together with the atom or atoms to which they are bonded, are C₃-C₂₀ heterocycloalkyl; in which R_{c3} is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, or aryl.

12. The method of claim 11, wherein one of R₄ and R₅ is H or C₁-C₁₅ alkyl optionally substituted with halogen, aryl, OR_{f1}, SR_{f1}, CO₂R_{f1}, C(O)NR_{f1}R_{f2}, SO₂R_{f1}, SO₃R_{f1}, or NR_{f1}R_{f2}; in which each of R_{f1} and R_{f2}, independently, is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, or aryl.

13. The method of claim 8, wherein the coronaviral 3CL protease is a severe acute respiratory syndrome viral 3CL protease or a human coronaviral 229E protease.

14. The method of claim 8, wherein the viral protease is a hepatitis C viral protease.

15. A compound of formula (I):



wherein

X is N(R_{a1}), O, or CH₂; or X and one of R₂ and R₃, together with the atom or atoms to which they are bonded, are C₃-C₂₀ heterocycloalkyl; in which R_{a1} is H or C₁-C₁₅ alkyl;

R₁ is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, aryl, C(O)R_{b1}, CO₂R_{b2}, C(O)NR_{b3}R_{b4}, C(O)-N(R_{b3})-OR_{b4}, C(S)R_{b3}, C(S)NR_{b3}R_{b4}, S(O)R_{b3}, SO₂R_{b3}, S(O)NR_{b3}R_{b4}, S(O)-N(R_{b3})-OR_{b4}, SO₂NR_{b3}R_{b4}, or SO₃R_{b3}; in which R_{b1} is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, 6-membered heteroaryl, fused heteroaryl, aryl, or NHCO₂R_{b5}; R_{b2} is H, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, aryl,

heteroaryl, or C₁-C₁₅ alkyl optionally substituted with halogen, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, OR_{b5}, CO₂R_{b5}, S(O)₂R_{b5}; and each of R_{b3} and R_{b4}, independently, is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, or aryl; R_{b5} being H, C₁-C₁₅ alkyl, heteroaryl, or aryl;

each of R₂ and R₃, independently, is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, aryl, OR_{c1}, SR_{c1}, or NR_{c1}R_{c2}; or X and one of R₂ and R₃, together with the atom or atoms to which they are bonded, are C₃-C₂₀ heterocycloalkyl; in which each of R_{c1} and R_{c2}, independently, is H, C₁-C₆ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, or aryl;


each of R₄ and R₅, independently, is H, halogen, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, or aryl;

one of R₆ and R₇ is C₁-C₁₅ alkyl substituted with C₃-C₂₀ heterocycloalkyl, heteroaryl, or aryl; and the other of R₆ and R₇ is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, or aryl;

R₈ is H, halogen, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, aryl, OR_{d1}, or SR_{d1}; in which R_{d1} is H and C₁-C₁₅ alkyl; and

each of R₉ and R₁₀, independently, is H, halogen, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, aryl, C(O)R_{e1}, CO₂R_{e1}, C(O)NR_{e1}R_{e2}, C(O)-N(R_{e1})-OR_{e2}, C(S)R_{e1}, C(S)NR_{e1}R_{e2}, CN, NO₂, S(O)R_{e1}, SO₂R_{e1}, S(O)NR_{e1}R_{e2}, S(O)-N(R_{e1})-OR_{e2}, SO₂NR_{e1}R_{e2}, SO₃R_{e1}, PO(OR_{e1})(OR_{e2}), PO(R_{e1})(R_{e2}), PO(NR_{e1}R_{e2})(OR_{e3}), PO(NR_{e1}R_{e2})(NR_{e3}R_{e4}), C(O)-N(R_{e1})-NR_{e2}R_{e3}, or C(S)-N(R_{e1})-NR_{e2}R_{e3}; or R₈ and R₁₀, taken together, are C₃-C₂₀ cycloalkyl or C₃-C₂₀ heterocycloalkyl; or R₉ and R₁₀, taken together, are C₃-C₂₀ cycloalkyl or C₃-C₂₀ heterocycloalkyl; in which each of R_{e1}, R_{e2}, R_{e3}, and R_{e4}, independently, is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, or aryl; or any two of R_{e1}, R_{e2}, R_{e3}, and R_{e4}, together with the atom or atoms to which they are bonded, are C₃-C₂₀ heterocycloalkyl.

16. The compound of claim 15, wherein X is N(R_{a1}); or X and one of R₂ and R₃, together with the atom or atoms to which they are bonded, are C₃-C₂₀ heterocycloalkyl; R₁ is H, C(O)R_{b1}, CO₂R_{b2}, C(O)NR_{b1}R_{b2}, C(S)NR_{b1}R_{b2}, or SO₂R_{b1}, in which R_{b1} is H, aryl, or C₁-C₁₅ alkyl optionally substituted with halogen, aryl, or NHCO₂R_{b5}; and R_{b2} is aryl or C₁-C₁₅ alkyl optionally substituted with halogen, OR_{b5}, or CO₂R_{b5}; each of R₂ and R₃, independently, is H or C₁-C₁₅ alkyl optionally substituted with OR_{c1}, C(O)-NR_{c1}R_{c2}, NR_{c1}R_{c2}, N(R_{c1})-CO₂R_{c2}, N(R_{c1})-SO₂R_{c2}, or O-SO₂-R_{c1}; or X and one of R₂ and R₃, together with the atom or atoms to which they are bonded, are C₃-C₂₀ heterocycloalkyl; each of R₄ and R₅, independently, is H or C₁-C₁₅ alkyl; one of R₆ and R₇ is C₁-C₁₅ alkyl substituted with C₃-C₂₀ heterocycloalkyl, and the other of R₆ and R₇ is H; R₈ is H; and each of R₉ and R₁₀, independently, is H or CO₂R_{e1}.




17. The compound of claim 16, wherein one of R₆ and R₇ is

18. The compound of claim 17, wherein one of R₄ and R₅ is H or C₁-C₁₅ alkyl optionally substituted with aryl.

19. The compound of claim 18, wherein the compound is one of compounds 18-26, 28-35, 50-53, 55, 56, 64-69, 77, 79, 81-84, 87, 88, 93, and 141.

20. A pharmaceutical composition comprising a compound of claim 15 and a pharmaceutically acceptable carrier.

21. A compound of formula (I):



wherein

X is N(R_{a1}), O, or CH₂; or X and one of R₂ and R₃, together with the atom or atoms to which they are bonded, are C₃-C₂₀ heterocycloalkyl; in which R_{a1} is H or C₁-C₁₅ alkyl;

R₁ is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, aryl, C(O)R_{b1}, CO₂R_{b2}, C(O)NR_{b2}R_{b3}, C(O)-N(R_{b2})-OR_{b3}, C(S)R_{b2}, C(S)NR_{b2}R_{b3}, S(O)R_{b2}, SO₂R_{b2}, S(O)NR_{b2}R_{b3}, S(O)-N(R_{b2})-OR_{b3}, SO₂NR_{b2}R_{b3}, or SO₃R_{b2}; in which R_{b1} is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, aryl, 6-membered heteroaryl, or fused heteroaryl; and each of R_{b2} and R_{b3}, independently, is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, aryl, or heteroaryl;

one of R₂ and R₃ is C₁-C₁₅ alkyl substituted with halogen, OR_{c1}, SR_{c1}, CO₂R_{c1}, OC(O)R_{c1}, C(O)NR_{c1}R_{c2}, SO₂R_{c1}, O-SO₂-R_{c1}, NR_{c1}R_{c2}, N(R_{c1})-C(O)R_{c2}, N(R_{c1})-CO₂R_{c2}, N(R_{c1})-SO₂R_{c2}, N(R_{c1})-C(O)-N(R_{c2}R_{c3}); the other of R₂ and R₃ is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, aryl, OR_{c1}, SR_{c1}, or NR_{c1}R_{c2}; or X and one of R₂ and R₃, together with the atom or atoms to which they are bonded, are C₃-C₂₀ heterocycloalkyl; in which each of R_{c1}, R_{c2}, and R_{c3}, independently, is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, or aryl;

each of R₄ and R₅, independently, is H, halogen, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, or aryl;

one of R₆ and R₇ is C₁-C₁₅ alkyl substituted with C₃-C₂₀ heterocycloalkyl, heteroaryl, or aryl; and the other of R₆ and R₇ is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, or aryl;


R₈ is H, halogen, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, aryl, OR_{d1}, or SR_{d1}; in which R_{d1} is H and C₁-C₁₅ alkyl; and

each of R₉ and R₁₀, independently, is H, halogen, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, aryl, C(O)R_{e1}, CO₂R_{e1}, C(O)NR_{e1}R_{e2}, C(O)-N(R_{e1})-OR_{e2}, C(S)R_{e1}, C(S)NR_{e1}R_{e2}, CN, NO₂, S(O)R_{e1}, SO₂R_{e1}, S(O)NR_{e1}R_{e2}, S(O)-N(R_{e1})-OR_{e2}, SO₂NR_{e1}R_{e2}, SO₃R_{e1}, PO(OR_{e1})(OR_{e2}), PO(R_{e1})(R_{e2}), PO(NR_{e1}R_{e2})(OR_{e3}), PO(NR_{e1}R_{e2})(NR_{e3}R_{e4}), C(O)-N(R_{e1})-NR_{e2}R_{e3}, or C(S)-N(R_{e1})-NR_{e2}R_{e3}; or R₈ and R₁₀, taken together, are C₃-C₂₀ cycloalkyl or C₃-C₂₀ heterocycloalkyl; or R₉ and R₁₀, taken together, are C₃-C₂₀ cycloalkyl or C₃-C₂₀ heterocycloalkyl; in which each of R_{e1}, R_{e2}, R_{e3}, and R_{e4}, independently, is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-

C₂₀ heterocycloalkyl, heteroaryl, or aryl; or any two of R_{e1}, R_{e2}, R_{e3}, and R_{e4}, together with the atom or atoms to which they are bonded, are C₃-C₂₀ heterocycloalkyl.

22. The compound of claim 21, wherein X is N(R_{a1}) or O; or X and one of R₂ and R₃, together with the atom or atoms to which they are bonded, are C₃-C₂₀ heterocycloalkyl; R₁ is CO₂R_{b2}, in which R_{b2} is alkyl substituted with aryl; one of R₂ and R₃ is C₁-C₁₅ alkyl substituted with OR_{c1}, SR_{c1}, OC(O)R_{c1}, CO₂R_{c1}, C(O)NR_{c1}R_{c2}, SO₂R_{c1}, O-SO₂-R_{c1}, NR_{c1}R_{c2}, N(R_{c1})-C(O)R_{c2}, N(R_{c1})-CO₂R_{c2}, N(R_{c1})-SO₂R_{c2}, N(R_{c1})-C(O)-N(R_{c2}R_{c3}), and the other of R₂ and R₃ is H; or X and one of R₂ and R₃, together with the atom or atoms to which they are bonded, are C₃-C₂₀ heterocycloalkyl; each of R₄ and R₅, independently, is H or C₁-C₁₅ alkyl; one of R₆ and R₇ is C₁-C₁₅ alkyl substituted with C₃-C₂₀ heterocycloalkyl, and the other of R₆ and R₇ is H; R₈ is H; and each of R₉ and R₁₀, independently, is H, CN, C(O)R_{e1}, CO₂R_{e1}, or C(O)NR_{e1}R_{e2}; or R₉ and R₁₀, taken together, are C₃-C₂₀ heterocycloalkyl.

23. The compound of claim 22, wherein one of R₆ and R₇ is




24. The compound of claim 23, wherein one of R₄ and R₅ is H or C₁-C₁₅ alkyl optionally substituted with aryl, C₃-C₂₀ cycloalkyl, OR_{f1}, SR_{f1}, NR_{f1}R_{f2}, or C(O)NR_{f1}R_{f2}; in which each of R_{f1} and R_{f2}, independently, is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, or aryl.

25. The compound of claim 24, wherein the compound is one of compounds 1-17, 27, 49, 54, 57-63, 74, 78, 80, 85, 86, 89-92, 94-105, 108, 109, 110, 113, 114, 117-119, 121-128, 132, and 135-145.

26. A pharmaceutical composition comprising a compound of claim 21 and a pharmaceutically acceptable carrier.

27. A compound of formula (I):



wherein

X is N(R_{a1}), O, or CH₂; or X and one of R₂ and R₃, together with the atom or atoms to which they are bonded, are C₃-C₂₀ heterocycloalkyl; in which R_{a1} is H or C₁-C₁₅ alkyl;

R₁ is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, aryl, C(O)R_{b1}, CO₂R_{b2}, C(O)NR_{b2}R_{b3}, C(O)-N(R_{b2})-OR_{b3}, C(S)R_{b2}, C(S)NR_{b2}R_{b3}, S(O)R_{b2}, SO₂R_{b2}, S(O)NR_{b2}R_{b3}, S(O)-N(R_{b2})-OR_{b3}, SO₂NR_{b2}R_{b3}, or SO₃R_{b2}; in which R_{b1} is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, aryl, 6-membered heteroaryl, or fused heteroaryl; and each of R_{b2} and R_{b3}, independently, is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, aryl, or heteroaryl;

each of R₂ and R₃, independently, is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, aryl, OR_{c1}, SR_{c1}, or NR_{c1}R_{c2}; or X and one of R₂ and R₃, together with the atom or atoms to which they are bonded, are C₃-C₂₀ heterocycloalkyl; in which each of R_{c1} and R_{c2}, independently, is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, or aryl;

one of R₄ and R₅ is C₁-C₁₅ alkyl substituted with halogen, OR_{d1}, SR_{d1}, CO₂R_{d1}, C(O)NR_{d1}R_{d2}, SO₂R_{d1}, SO₃R_{d1}, or NR_{d1}R_{d2}; the other of R₄ and R₅ is H, halogen, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, or aryl; in which each of R_{d1} and R_{d2}, independently, is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, or aryl;

one of R₆ and R₇ is C₁-C₁₅ alkyl substituted with C₃-C₂₀ heterocycloalkyl, heteroaryl, or aryl; and the other of R₆ and R₇ is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, or aryl;


R₈ is H, halogen, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, aryl, OR_{e1}, or SR_{e1}; in which R_{e1} is H and C₁-C₁₅ alkyl; and

each of R₉ and R₁₀, independently, is H, halogen, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, aryl, C(O)R_{f1}, CO₂R_{f1}, C(O)NR_{f1}R_{f2},

C(O)-N(R_{f1})-OR_{f2}, C(S)R_{f1}, C(S)NR_{f1}R_{f2}, CN, NO₂, S(O)R_{f1}, SO₂R_{f1}, S(O)NR_{f1}R_{f2}, S(O)-N(R_{f1})-OR_{f2}, SO₂NR_{f1}R_{f2}, SO₃R_{f1}, PO(OR_{f1})(OR_{f2}), PO(R_{f1})(R_{f2}), PO(NR_{f1}R_{f2})(OR_{f3}), PO(NR_{f1}R_{f2})(NR_{f3}R_{f4}), C(O)-N(R_{f1})-NR_{f2}R_{f3}, or C(S)-N(R_{f1})-NR_{f2}R_{f3}; or R₈ and R₁₀, taken together, are C₃-C₂₀ cycloalkyl or C₃-C₂₀ heterocycloalkyl; or R₉ and R₁₀, taken together, are C₃-C₂₀ cycloalkyl or C₃-C₂₀ heterocycloalkyl; in which each of R_{f1}, R_{f2}, R_{f3}, and R_{f4}, independently, is H, C₁-C₁₅ alkyl, C₃-C₂₀ cycloalkyl, C₃-C₂₀ heterocycloalkyl, heteroaryl, or aryl; or any two of R_{f1}, R_{f2}, R_{f3}, and R_{f4}, together with the atom or atoms to which they are bonded, are C₃-C₂₀ heterocycloalkyl.

28. The compound of claim 27, wherein X is N(R_{a1}) or O; R₁ is CO₂R_{b2}, in which R_{b2} is alkyl substituted with aryl; each of R₂ and R₃, independently, is H or C₁-C₁₅ alkyl optionally substituted with aryl; one of R₄ and R₅ is C₁-C₁₅ alkyl substituted with SR_{d1} or C(O)NR_{d1}R_{d2}, and the other of R₄ and R₅ is H; one of R₆ and R₇ is C₁-C₁₅ alkyl substituted with C₃-C₂₀ heterocycloalkyl, and the other of R₆ and R₇ is H; R₈ is H; and each of R₉ and R₁₀, independently, is H or CO₂R_{f1}.

29. The compound of claim 28, wherein one of R₆ and R₇ is



30. The compound of claim 29, wherein the compound is one of compounds 45-48.

31. A pharmaceutical composition comprising a compound of claim 27 and a pharmaceutically acceptable carrier.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US04/44004

A. CLASSIFICATION OF SUBJECT MATTER
 IPC(7) : C07D 207/06, C07D 223/02, C07D 207/04
 US CL : 548/530, 518; 540/524
 According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED
 Minimum documentation searched (classification system followed by classification symbols)
 U.S. : 548/530, 518; 540/524

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
 STN (Registry, CAPLUS and Beilstein)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	CAULFIELD, et al, J. of Combinatorial. Chem. (2000), 2(6), 600-603.	19
A	US 5,962,487 (WEBBER et al) (4.10.1999)	19
A	JENWITHEESUK, E. Bioorganic and Medicinal Chemistry Letters October 2003, Vol. 13, No. 22, pages 3989-3992	19
A	WO 9743305 A1 (Webber et al) 20 Nov. 1997 (20.11.1997)	19

Further documents are listed in the continuation of Box C. See patent family annex.

Special categories of cited documents:		
"A" document defining the general state of the art which is not considered to be of particular relevance	"T"	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"E" earlier application or patent published on or after the international filing date	"X"	document of particular relevance. the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y"	document of particular relevance. the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O" document referring to an oral disclosure, use, exhibition or other means		
"P" document published prior to the international filing date but later than the priority date claimed	"&"	document member of the same patent family

Date of the actual completion of the international search: 08 June 2005 (08.06.2005)
 Date of mailing of the international search report: 24 JUN 2005

Name and mailing address of the ISA/US: Mail Stop PCT, Attn: ISA/US, Commissioner for Patents, P.O. Box 1450, Alexandria, Virginia 22313-1450, Facsimile No. (571) 273-8300
 Authorized officer: Nyeemah A. Grazie, Telephone No. (571) 272-8781

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US04/44004

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. Claims Nos.: 1-18 and 20-31
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
See Continuation Sheet.

3. Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
 2. As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
 3. As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

 4. No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
- Remark on Protest**
- | | |
|--------------------------|---|
| <input type="checkbox"/> | The additional search fees were accompanied by the applicant's protest. |
| <input type="checkbox"/> | No protest accompanied the payment of additional search fees. |