

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
30 March 2006 (30.03.2006)

PCT

(10) International Publication Number
WO 2006/033697 A1

- (51) International Patent Classification:
A23L 1/212 (2006.01) A23L 1/308 (2006.01)
- (21) International Application Number:
PCT/US2005/024497
- (22) International Filing Date: 12 July 2005 (12.07.2005)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/586,732 12 July 2004 (12.07.2004) US
- (71) Applicant (for all designated States except US):
CARGILL, INC. [US/US]; 15407 McGinty Road
West, Wayzata MN 55391-2399 (US).

3240 Yates Av. N., Crystal, MN 55422 (US). REEDER, David [US/US]; 6501 Troendle Circle, Chanhassen, MN 55317 (US).


(74) Agent: RIVARD, Paul, M.; Banner & Witcoff, Ltd., 1001 G Street, N.W., 11th Floor, Washington, D.C. 20001-4597 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

[Continued on next page]

(54) Title: PROCESS OF EXTRACTING CITRUS FIBER FROM CITRUS VESICLES



(57) Abstract: Citrus fiber is recovered from citrus vesicles to obtain a food additive for beverages, baked goods, meat or meat emulsions, confectionary, jams and jellies, dairy products, dressings, energy bars, and the like. Citrus vesicles optionally are washed with water, and water washed vesicles are recovered. The vesicles are contacted with an organic solvent to obtain organic solvent washed vesicles. The organic solvent washed vesicles are desolventized and dried citrus fiber is recovered.

WO 2006/033697 A1



Published:

- *with international search report*
- *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

PROCESS OF EXTRACTING CITRUS FIBER FROM CITRUS VESICLES

FIELD OF THE INVENTION

The present invention is directed to citrus fiber extracted from citrus vesicles.

- 5 The resulting dried citrus fiber is useful as a food additive for beverages, baked goods, meat or meat emulsions, confectionary, jams and jellies, dairy products, dressings, energy bars, and the like.

DESCRIPTION OF RELATED ART

Current processes for making citrus juice employ extractors for separating the
10 juice-containing inner part of the fruit from its outer peel. The juice extracted by the extractors contains the juice itself as well as a pectinaceous and cellulosic material called citrus vesicles. This material sometimes is also referred to as coarse pulp, floaters, citrus cells, floating pulp, juice sacs, or pulp.

Citrus vesicles typically are separated from the juice by means of filtration,
15 using equipment such as paddle finishers, screw finishers and turbo filters. The citrus vesicles contain a significant amount of juice-soluble solids. For economic reasons, the citrus vesicles often are subjected to a water washing step, *e.g.*, using blenders, screw mixers, inline static mixers, finishers, or turbo filters. The water washing step produces a stream of secondary grade juice, referred to as water extracted soluble
20 orange solids (WESOS) or simply "pulp wash." The washed cells, also called "washed pulp," are considered waste from this step. Typically, the washed cells are mixed with the peel (from the extraction process mentioned above), further processed (for example treated with lime to assist in dewatering) and dried for use as cattle feed.

Some efforts have been made to recover potentially valuable components from citrus waste. For example, U.S. Patent 6,183,806 B1 to Ficca *et al.* describes citrus peel extracts and flour prepared from the extracts. Citrus peel is extracted with an ethanol solvent and the solid residue is desolventized to recover orange flour. The citrus peel is provided in comminuted pieces or particles. The comminuted pieces or particles may be provided in the form of washed pulp (pomace). Ficca uses the term “washed pulp” somewhat unconventionally to refer to pomace, a component of the peel. Ficca indicates that the orange flour extracted from pomace is similar in composition to the flour obtained from orange peel.

Washed citrus cells contain citrus fiber, a valuable citrus component which has relatively high total dietary fiber content and a balanced ratio of soluble to insoluble dietary fiber. The balanced dietary fiber spectrum of insoluble (chiefly cellulose) and soluble (chiefly pectin) fiber is advantageous in physiological functionality over cereal-based fibers. Citrus fiber, particularly orange pulp fiber, has an extremely high water binding capacity, resulting in high viscosities compared to other citrus fibers such as Vitacel orange fiber (available from Rettenmaier).

It would be desirable to develop a process for recovering citrus fiber from citrus vesicles, especially a process which can recover citrus fiber in an efficient and relatively cost effective manner. It would be especially desirable to develop a process which can recover citrus fiber without the need for using potentially hazardous agents. It would be particularly desirable to develop a process which provides citrus fiber that can be used as a food and beverage ingredient.

SUMMARY OF THE INVENTION

The present invention, according to one aspect, is directed to a process of purifying citrus fiber in citrus vesicles to obtain a food additive suitable for human consumption. The process comprises an optional step of washing citrus vesicles with
5 water to remove unwanted flavors, odors, colors, sugars, acids and the like. The vesicles are contacted with an organic solvent to obtain organic solvent washed vesicles. The organic solvent washed vesicles are desolventized and dried citrus fiber is recovered therefrom.

In one preferred embodiment of the invention, the extraction process employs
10 an ethanol solvent and a solids-to-solvent weight ratio of at least about 0.25:1, preferably at least about 0.5:1. The solvent extraction is carried out in at least two counter current stages. At least a portion of the solvent, preferably at least about 70%, is recovered and reused.

In another aspect of the present invention, dried citrus fiber is prepared by the
15 extraction process as described above. The dried citrus fiber is useful as a food additive for such food products as beverages, baked goods, meat or meat emulsions, confectionary, jams and jellies, dairy products such as yogurt, dressings, energy bars, and the like.

In another aspect of the invention, a citrus fiber has a total dietary fiber
20 content of from about 60 to about 85 wt% and a water binding capacity from about 9 to about 25 (w/w).

In one embodiment, dried citrus fiber has a total dietary fiber content of from about 60 to about 80 wt% and a water binding capacity from about 7 to about 12

(w/w). Preferably the total dietary fiber content is at least about 70 wt% and the water binding capacity is at least about 8 (w/w).

BRIEF DESCRIPTION OF THE DRAWINGS

The objects, features, and advantages of the invention will be apparent from
5 the following more detailed description of certain embodiments of the invention and
as illustrated in the accompanying drawings in which:

FIG. 1 is a schematic illustration of a two-stage countercurrent solvent extraction process in accordance with a preferred embodiment of the present invention.

10 DETAILED DESCRIPTION OF THE INVENTION

The present invention will be described primarily with reference to extracting orange fiber from orange vesicles. It should be understood that the process may be used for extracting citrus fiber from citrus vesicles from a wide variety of other types of citrus fruit, non-limiting examples of which include tangerines, limes, lemons, and
15 grapefruit. The citrus vesicles typically have a water content of at least about 80 wt% and usually from about 90 to about 97 wt%.

The term "citrus vesicles," as used herein, refers to the pectinaceous and cellulosic material contained in the inner, juice-containing portion of citrus fruit. Citrus vesicles are sometimes also referred to as coarse pulp, floaters, citrus cells,
20 floating pulp, juice sacs, or pulp.

The term "water extracted soluble solids," as used herein, refers to secondary grade juice which is obtained by water-washing citrus vesicles. The term "water

extracted soluble solids” is particularly inclusive of water extracted soluble orange solids (WESOS). Water extracted soluble solids are sometimes also referred to as “pulp wash.”

The term “water washed vesicles” refers to citrus vesicles from which water
5 extracted soluble solids have been removed by washing with water. Water washed vesicles are also sometimes referred to as “washed cells” or “washed pulp.”

The term “citrus fiber,” as used herein, refers to a fibrous component obtained from water washed vesicles and isolated from waste components present therein. Citrus fiber is characterized by a high total dietary fiber content as well as a balanced
10 ratio of soluble to insoluble dietary fiber. Citrus fiber, particularly orange fiber, has a very high water binding capacity.

Citrus fiber, as compared to citrus flour obtained from citrus peel, is lighter in color and relatively free of taste and odor. In contrast, citrus flour obtained from citrus peel is characterized by an orange peel taste, odor and color, which severely
15 limit the product’s uses. Additional advantages of citrus fiber include a higher total dietary fiber content (e.g., greater than about 70% versus 58%); lower carbohydrate content (e.g., about 5% versus 15%); and higher water binding capacity (e.g., greater than about 8 grams of water per gram of fiber versus 5.5 g/g). The protein content of the citrus fiber typically ranges from about 8 to 12 wt%.

20 The ratio of soluble to insoluble dietary fiber is an important factor in the citrus fiber’s functionality. Preferably, the citrus fiber has a balanced ratio of soluble to insoluble dietary fiber. For example, the total dietary fiber preferably is made up of about 45-50% soluble dietary fiber and from 50-55% insoluble dietary fiber. Other important considerations include the degree of milling (granulometry) and drying

conditions (process of drying). Generally, a higher degree of milling (*i.e.*, a finer fiber granulometry) results in more smoothness of mouthfeel of the fiber in foods and beverages. Density and particle size may vary over a wide range depending on processing conditions. By way of example, density may range from about 80 to about 5 650 g/L, and average particle size may range from about 15 to about 600 microns. It should be understood that these ranges are merely exemplary. In some applications it may be desirable to employ significantly larger particle sizes, for example. In general, the citrus fiber may range from a very fine to a coarse powder structure.

As shown schematically in Fig. 1, orange vesicles can be separated from 10 orange juice by filtration, *e.g.*, using conventional filtration equipment 10 such as paddle finishers, screw finishers or turbo filters. The orange vesicles then can be subjected to an optional water washing step using suitable equipment 20 such as blenders, inline static mixers, finishers, or turbo filters. Water washing is desirable to eliminate undesirable flavors, odors, and acids present in the citrus vesicles. The 15 water washing step 20 produces a stream of water extracted soluble orange solids (WESOS). The water extracted soluble orange solids, together with the juice obtained by filtration 10, can be further processed using conventional juicing equipment 15.

An organic solvent is used to extract flavors, odors, colors and the like from the water washed vesicles. The solvent should be polar and water-miscible to 20 facilitate removal of the desired components. Preferred solvents include lower alcohols such as methanol, ethanol, propanol, isopropanol, or butanol. The solvent may be (and preferably is) provided in aqueous solution. The concentration of solvent in the solvent solution most often ranges from about 70 wt% to about 100 wt%. In one embodiment, a 75 wt% aqueous ethanol solution is used as solvent. In a preferred

embodiment, a 90 wt% aqueous ethanol solution is used as solvent. In general, solvents will remove water-soluble components at lower concentrations and oil-soluble components at higher concentrations. Optionally, a more non-polar co-solvent may be added to the aqueous alcohol to improve the recovery of oil-soluble components in the citrus vesicles. Examples of such non-polar solvents include ethyl acetate, methyl ethyl ketone, acetone, hexane, methyl isobutyl ketone and toluene. The more non-polar solvents may be added at up to 20% of the solvent mixture. Many solvents, such as ethanol, have a lower heat of vaporization than that of water, and therefore require less energy to volatilize than would be needed to volatilize an equivalent mass of water. The solvent preferably is removed and reclaimed for reuse.

Preferably, the water washed vesicles are contacted with organic solvent at a solids-to-solvent weight ratio of at least about 0.25:1, preferably at least about 0.5:1, and often at least about 0.75:1, from about 1:1 to about 5:1, or from about 1.5:1 to about 3:1, based on the wet weight of the solids. In one embodiment, the solids-to-solvent ratio is about 2:1.

Extraction can be accomplished using a single stage but preferably is performed using multi-stage extraction, *e.g.*, a two-, three-, or four-staged extraction process, and preferably using countercurrent extraction. There is no particular upper limit contemplated on the number of extraction stages that may be used. Fig. 1 schematically illustrates a preferred embodiment in which a two-stage countercurrent extraction process employs first and second solvent extractors 25a and 25b, respectively.

Water washed vesicles are fed into the second extractor 25b. An aqueous ethanol solvent is fed from a solvent tank 26 into the first solvent extractor 25a. Spent

solvent from the first solvent extractor 25a is fed into the second solvent extractor 25b, while the extracted citrus vesicles from the second solvent extractor 25b are fed into the first solvent extractor 25a. Spent solvent from the second solvent extractor 25b may be fed into an evaporator 35 (optional) to separate solids (*e.g.*, sugars, colors, 5 flavors, citrus oils, *etc.*) from the spent solvent, which can be condensed and returned to a still 24. Still bottoms (predominately water) are separated and removed.

After each extraction stage, a belt filter press (not shown) preferably is used to further remove liquids. A belt filter press may be of conventional construction which is known to persons skilled in the art. Details of suitable belt filter presses are found, 10 for example, in U.S. Patents 4,236,445, 4,297,215, and 5,022,989, the disclosures of which are hereby incorporated by reference.

Solvent washed vesicles from the first solvent extractor 25a are fed to a desolventizer 30. The desolventizer 30 removes solvent and water from the solids remaining after extraction, enabling the solvent to be reclaimed for future use and also 15 ensuring that the product is safe for milling and commercial use. The desolventizer 30 can employ indirect heat to remove significant amounts of solvent from the solid residue. Alternatively, direct heat can be provided for drying, *e.g.*, by providing hot air from flash dryers or fluidized bed dryers. Direct steam may be employed, if desired, to remove any trace amounts of solvent remaining in the solids. Vapors from 20 the desolventizer 30 preferably are recovered and fed to the still 24 to reclaim at least a portion of the solvent.

Retention time in each extraction step may vary over a wide range but usually is about 5 minutes or less and preferably is about 3 minutes or less. The temperature in the solvent extractor(s) depends on such factors as the type of solvent used but

most often ranges from about 40°F to about 180°F at atmospheric pressure. Temperatures can be appropriately increased or decreased for operation under super- or sub-atmospheric pressures. Optionally, techniques such as ultra-sound are used for enhancing efficiency of the extraction process. By maintaining a closed system, solvent losses during extraction, desolventizing, and distillation can be minimized. Preferably, at least about 70 wt% of the solvent is recovered and reused. A solvent make-up stream delivers fresh solvent into the solvent tank 26 to replenish any solvent that is not recovered.

The dried orange fiber product is generally a coarse flour. The flour can be milled if a powder is needed for a particular application. Optionally, the citrus fiber is subjected to one or more additional treatments needed or desired for a particular end use. For example, natural enzymes may be inactivated prior to the solvent extraction process or drying. Enzymes may be inactivated, *e.g.*, by ultrasound or by conventional thermal inactivation.

Unlike many typical commercial fibers, the orange fiber of the present invention has thickening power, creates pulpy textures (coarse fraction) and has good stability. For example, the citrus fiber can be used for 1% sucrose replacement in a standard bakery cream recipe with improved baking stability and good freeze/ thaw stability. This indicates an extremely high water binding capacity of the orange fiber, resulting in relatively high viscosities.

The high water binding capacity can be beneficial for many applications (*e.g.*, processed meat applications). In other applications, such as in bakery products, a high water binding capacity may not be desirable. Extremely fine milled fibers (*e.g.*, 20 to

30 μm particle size) generally will show superior sensory characteristics (high smoothness).

In addition to having utility as a food additive in beverages, baked goods, meat or meat emulsions (e.g., sausage, bologna, hot dogs), confectionary, jams and jellies, 5 dairy products, dressings, energy bars, and the like, the citrus fiber also potentially can be used as a natural thickening agent for food products, such as yogurt, or as a carrier for other components in beverage systems. Suitable amounts of citrus fiber for the various food products can be determined by persons skilled in the art in accordance with such factors as user preferences and with the aid of no more than 10 routine experimentation.

EXAMPLE 1

This example illustrates extracting citrus fiber from water washed vesicles using a 75 wt% aqueous ethanol solvent solution in a two stage countercurrent extraction process as described above. The resulting orange fiber was analyzed for 15 dietary fiber content using AOAC method 991.43 (Prosky). The citrus fiber was found to have the composition and properties set forth in Table 1:

TABLE 1

Total Dietary Fiber (TDF)	72.3 %
Soluble Dietary Fiber (SDF)	37.1 %
Insoluble Dietary Fiber (IDF)	34.4 %
Carbohydrates	5 %
Proteins (dry basis)	1.6 % N (9.7 % protein)
Fatty Acids (dry basis)	0.16 %
Ash	2.14 %
Sodium	102 mg/kg
Potassium	4960 mg/kg
Magnesium	648 mg/kg
Calcium	1850 mg/kg
Phytic Acid	479 ppm
Moisture	11%

EXAMPLE 2

This example illustrates preparing bread containing orange fiber. Table 2 lists
 5 the ingredients used to prepare the bread:

TABLE 2

Ingredient	Amount (g)
Duo Flour (Ceres)	950
Orange pulp fiber	30
Bread improver	100
Salt	17
Margarine	20
Water	551

All dry ingredients were blended in a spiral mixer (Veema type). Water was then added, and the resulting dough was mixed for 20 minutes. The dough was divided into 700 gram portions. Sheeting and proofing were carried out for 1.5 hours at 32°C and 82% relative humidity. The bread was baked in a floor oven at 220°C for 35 minutes.

EXAMPLE 3

This example illustrates preparing a dressing (30% oil type) containing orange fiber. Table 3 lists the ingredients used to prepare the dressing:

TABLE 3

Ingredient	Amount (wt%)
Sugar	3
Salt	2
Potassium sorbate	0.12
Water	49.8
Oil	12
Orange fiber	3
Egg yolk powder	1.5
Xanthan + guar gum	0.2
Sunflower oil	18
Vinegar	7.2
Mustard	3

10

The water and acid phase were added into a Fryma colloid mill. The dispersion phase was then added and homogenized for 30 seconds. The oil phase was added over the next one minute, and the mixture was emulsified for an additional 30 seconds. The dressing was then filled into jars.

15 EXAMPLE 4

This example illustrates preparing a beef burger containing orange fiber. Table 4 lists the ingredients used to prepare the beef burger:

TABLE 4

Ingredient	Amount (wt%)
Beef (grinded)	90
Spices	1.5
Orange fiber	1.5
Water	7

All ingredients (except spices) were blended in a Hobart Mixer Type N50CE at speed 1. The spices were added after 30 seconds. The mixture was blended up to 5 minutes, with the mass turn manually after 2.5 minutes. The mass was cooled in a refrigerator for 2 hours. The burgers were formed as 90g servings with an 80 mm diameter. The burgers can be cooked in frying pan until a core temperature of 74-75°C is obtained.

EXAMPLE 5

This example illustrates preparing a jam containing orange fiber. Table 5 lists the ingredients that can be used to prepare the jam:

TABLE 5

Ingredient	Amount (wt%)
Mixed Berries (red fruits, frozen) or strawberries	43
Sucrose	25
Fructose syrup	25
Orange fiber	2
Water	5
Citric acid solution (50%)	to bring pH to 3.4

The fruit, sweeteners and water can be blended and cooked together. The orange fiber then can be added and boiled to the desired dry substance. The citric acid solution then can be added until the mixture reaches a pH of 3.4. The jam can be hot-filled into glass jars.

EXAMPLE 6

This example illustrates preparing energy bars. Table 6 lists the ingredients used to prepare the energy bars:

TABLE 6

Ingredient	Amount (wt%)
Fructose syrup	14
Sucrose	8
Isomaltulose (Cerestar Cargill)	15
Water	3
Glycerol	3
Palm fat	7
Orange fiber	3
Red fruity nuggets	15
Lecithin	1
Oats flakes	13
Soy nuggets	11
Whey powder	7

5

The sweeteners and water were heated in a pan to 88°C. The molten fat and lecithin were blended in a Hobart mixer for 1 minute at speed 1. The sweetener was added to the fat and blended at speed 1 for 1 minute. The dry ingredients were added and mixed at speed 1 for 30 seconds. The mixture was deposited onto a sheet and cut to the desired size. The bars were then cooled and packaged.

10

While particular embodiments of the present invention have been described and illustrated, it should be understood that the invention is not limited thereto since modifications may be made by persons skilled in the art. The present application contemplates any and all modifications that fall within the spirit and scope of the underlying invention disclosed and claimed herein.

15

WHAT IS CLAIMED IS:

1. A process of recovering citrus fiber from citrus vesicles to obtain a food additive suitable for human consumption, the process comprising:

5 (i) washing citrus vesicles with water and recovering water washed vesicles therefrom;

(ii) contacting the water washed vesicles with an organic solvent to obtain organic solvent washed vesicles; and

(iii) desolventizing the organic solvent washed vesicles and recovering dried citrus fiber therefrom.

10

2. The process of claim 1 wherein the organic solvent is an alcohol.

3. The process of claim 2 wherein the organic solvent is ethanol.

15

4. The process of claim 1 wherein the citrus vesicles are obtained from citrus fruit selected from the group consisting of oranges, tangerines, limes, lemons, and grapefruit.

5. The process of claim 1 wherein the citrus vesicles have a water content
20 of at least about 80 wt%.

6. The process of claim 1 wherein the water washed vesicles are contacted with organic solvent at a solids-to-solvent weight ratio of at least about 0.25:1.

25

7. The process of claim 1 wherein the organic solvent extraction is carried out in at least two counter current stages.

8. The process of claim 1 wherein the organic solvent extraction is
5 carried out in a continuous process.

9. The process of claim 1 wherein the organic solvent is water-miscible.

10. The process of claim 1 wherein the organic solvent has a concentration
10 of at least about 70 wt%.

11. The process of claim 1 further comprising pressing the organic solvent extract on a filter press to remove liquid.

12. The process of claim 1 wherein at least a portion of the organic solvent
15 is re-cycled and recovered.

13. The process of claim 12 wherein at least about 70 wt% of the organic solvent is recovered.

20

14. The process of claim 1 wherein the dried citrus fiber has a total dietary fiber content of from about 60 to about 85 wt%.

15. The process of claim 1 wherein the dried fiber has a water binding
25 capacity of from about 7 to about 25 (w/w).

16. A process of recovering citrus fiber from citrus vesicles to obtain a food additive suitable for human consumption, the process comprising:

5 (i) providing citrus vesicles from citrus fruit selected from the group consisting of oranges, tangerines, limes, lemons, and grapefruit, wherein the citrus vesicles have a water content of at least about 80 wt%;

(ii) washing the citrus vesicles with water and recovering water washed vesicles therefrom;

10 (iii) contacting the water washed vesicles with an ethanol solvent at a solids-to-solvent weight ratio of at least about 0.25:1 to obtain solvent washed vesicles, wherein the solvent extraction is carried out in at least two countercurrent stages;

15 (iv) desolventizing the solvent washed vesicles and recovering dried citrus fiber therefrom, wherein the dried citrus fiber has a total dietary fiber content of from about 60 to about 85 wt% and a water binding capacity of from about 7 to about 25 (w/w); and

(v) recovering and recycling at least a portion of the ethanol solvent.

17. A dried citrus fiber food additive prepared by the process of claim 1.

20

18. A dried citrus fiber food additive prepared by the process of claim 16.

19. A food additive suitable for human consumption comprising dried citrus fiber having a total dietary fiber content of from about 60 to about 85 wt% and
25 a water binding capacity from about 7 to about 25 (w/w).

20. The food additive of claim 19 wherein the total dietary fiber content is about 60 to about 80 wt% and the water binding capacity is about 7 to about 12 (w/w).

21. A food product selected from the group consisting of beverages, baked
5 goods, meat, meat emulsions, confectionary, jams and jellies, dairy products, dressings, and energy bars, the food product containing the food additive of claim 19.

22. Citrus fiber isolated from water washed citrus vesicles, wherein the citrus fiber has a total dietary fiber content of from about 60 to about 85 wt% and a
10 water binding capacity from about 7 to about 25 (w/w).

23. The citrus fiber of claim 22 wherein the total dietary fiber content is at least about 70 wt% and the water binding capacity is at least about 8 (w/w).

15 24. The citrus fiber of claim 1 wherein the total dietary fiber comprises from about 45 to about 50 wt% soluble dietary fiber and from about 50 to about 55 wt% insoluble dietary fiber.

20 25. The citrus fiber of claim 22 which is obtained from citrus fruit selected from the group consisting of oranges, tangerines, limes, lemons, and grapefruit.

26. A process of recovering citrus fiber from citrus vesicles to obtain a food additive suitable for human consumption, the process comprising:

(i) contacting citrus vesicles with an organic solvent to obtain organic
25 solvent washed vesicles; and

(ii) desolventizing the organic solvent washed vesicles and recovering dried citrus fiber therefrom.




FIG. 1

INTERNATIONAL SEARCH REPORT

International application No
PCT 2005/024497

A. CLASSIFICATION OF SUBJECT MATTER
A23L1/212 A23L1/308

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
A23L

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, FSTA, PAJ

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 179 295 A (DANON, DAVID) 30 April 1986 (1986-04-30) page 2, paragraph 2 page 3, paragraph 3 - page 5, paragraph 5 page 5, line 35 - page 6, line 6; claims 1,2; example 3	1-26
X	----- DATABASE WPI Section Ch, Week 199110 Derwent Publications Ltd., London, GB; Class B04, AN 1991-071426 XP002364712 & SU 1 556 681 A (TBILISI MEDICINE IN) 15 April 1990 (1990-04-15) abstract -----	17-20, 22-26
	----- -/--	

Further documents are listed in the continuation of Box C.

See patent family annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

Date of mailing of the international search report

27 January 2006

28/02/2006

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Vuillamy, V

INTERNATIONAL SEARCH REPORT

Inter application No

PC 2005/024497

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DE 199 43 188 A1 (HERBSTREITH & FOX KG PEKTIN-FABRIK) 15 March 2001 (2001-03-15) page 3, lines 30-54 page 4, lines 1-12; claims -----	19-21
X	US 6 183 806 B1 (FICCA VINCENT G ET AL) 6 February 2001 (2001-02-06) cited in the application claims; figure; examples -----	19-21
X	G.R. GIBSON, C.M. WILLIAMS: "Functional Foods - Concept to Product" 2000, CRC, USA, XP002364675 Chapter 13, pp.315-326 column 324, paragraph 3; table 13.6 -----	19,20, 22,24,25
X	LARRAURI J A ET AL: "Water recycling in processing orange peel to a high dietary fibre powder." INTERNATIONAL JOURNAL OF FOOD SCIENCE & TECHNOLOGY 1997, vol. 32, no. 1, 1997, pages 73-76, XP002364674 Spain table 1 -----	19,20
A	WO 97/33688 A (GIST-BROCADES B.V) 18 September 1997 (1997-09-18) the whole document -----	
A	EP 0 223 703 A (SUCRE RECH DEV; SUCRE RECH DEV GRP INTERET ECONOM) 27 May 1987 (1987-05-27) the whole document -----	

INTERNATIONAL SEARCH REPORT

International application No
PCT 005/024497

Patent document cited in search report	A	Publication date	Patent family member(s)	Publication date
EP 0179295	A	30-04-1986	NONE	
SU 1556681	A	15-04-1990	NONE	
DE 19943188	A1	15-03-2001	AU 7507100 A WO 0117376 A1	10-04-2001 15-03-2001
US 6183806	B1	06-02-2001	NONE	
WO 9733688	A	18-09-1997	AU 2157497 A JP 2000507311 T	01-10-1997 13-06-2000
EP 0223703	A	27-05-1987	FR 2589683 A1	15-05-1987