



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification⁵ : A61K 31/44, 31/675, C07D 401/06, C07F 9/30, 9/38, 9/40, G01N 24/08</p>	A1	<p>(11) International Publication Number: WO 94/26275</p> <p>(43) International Publication Date: 24 November 1994 (24.11.94)</p>
<p>(21) International Application Number: PCT/US94/05009</p> <p>(22) International Filing Date: 5 May 1994 (05.05.94)</p> <p>(30) Priority Data: 057,588 6 May 1993 (06.05.93) US</p> <p>(71) Applicant: THE DOW CHEMICAL COMPANY [US/US]; 2030 Dow Center, Abbott Road, Midland, MI 48640 (US).</p> <p>(72) Inventors: KIEFER, Garry, E.; 114 Juniper Street, Lake Jackson, TX 77566 (US). KIM, Won, D.; 2400 Waterview Parkway #1122, Richardson, TX 75080 (US).</p> <p>(74) Agent: KIMBLE, Karen, L.; The Dow Chemical Company, Patent Dept., P.O. Box 1967, Midland, MI 48641-01967 (US).</p>		<p>(81) Designated States: AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, ES, FI, GB, HU, JP, KR, KZ, LK, LU, LV, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, UA, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>With international search report.</i></p>
<p>(54) Title: 2-PYRIDYLMETHYLENEPOLYAZAMACROCYCLOPHOSPHONIC ACIDS, COMPLEXES AND DERIVATIVES THEREOF, FOR USE AS CONTRAST AGENTS</p>		
<p>(57) Abstract</p> <p>2-Pyridylmethylenepolyazamacrocyclophosphonic acid compounds and their derivatives are disclosed which may form inert complexes with Gd, Mn or Fe ions. The overall charge of the complex can be varied to alter the <i>in vivo</i> biolocalization. The complexes are useful as contrast agents for diagnostic purposes.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

2-PYRIDYLMETHYLENEPOLYAZAMACROCYCLOPHOSPHONIC ACIDS, COMPLEXES AND
DERIVATIVES THEREOF, FOR USE AS CONTRAST AGENTS

This invention concerns ligands that are 2-pyridylmethylenepolyazamacrocyclo-
5 phosphonic acids, complexes and derivatives thereof, for use as contrast agents in magnetic
resonance imaging (MRI). To better understand this invention, a brief background on MRI is
provided in the following section.

MRI is a non-invasive diagnostic technique which produces well resolved cross-
sectional images of soft tissue within an animal body, preferably a human body. This technique
10 is based upon the property of certain atomic nuclei (e.g. water protons) which possess a
magnetic moment [as defined by mathematical equations; see G. M. Barrow, Physical
Chemistry, 3rd Ed., McGraw-Hill, NY (1973)] to align in an applied magnetic field. Once
aligned, this equilibrium state can be perturbed by applying an external radio frequency (RF)
pulse which causes the protons to be tilted out of alignment with the magnetic field. When the
15 RF pulse is terminated, the nuclei return to their equilibrium state and the time required for
this to occur is known as the relaxation time. The relaxation time consists of two parameters
known as spin-lattice (T1) and spin-spin (T2) relaxation and it is these relaxation measurements
which give information on the degree of molecular organization and interaction of protons
with the surrounding environment.

20 Since the water content of living tissue is substantial and variations in content and
environment exist among tissue types, diagnostic images of biological organisms are obtained
which reflect proton density and relaxation times. The greater the differences in relaxation
times (T1 and T2) of protons present in tissue being examined, the greater will be the contrast
in the obtained image [e.g., *J. Magnetic Resonance* 33, 83-106 (1979)].

25 It is known that paramagnetic chelates possessing a symmetric electronic ground
state can dramatically affect the T1 and T2 relaxation rates of juxtaposed water protons and
that the effectiveness of the chelate in this regard is related, in part, to the number of unpaired
electrons producing the magnetic moment [e.g., *Magnetic Resonance Annual*, 231-266, Raven
Press, NY (1985)]. It has also been shown that when a paramagnetic chelate of this type is
30 administered to a living animal, its effect on the T1 and T2 of various tissues can be directly
observed in the magnetic resonance (MR) images with increased contrast being observed in the
areas of chelate localization. It has therefore been proposed that stable, non-toxic
paramagnetic chelates be administered to animals in order to increase the diagnostic
information obtained by MRI [e.g., *Frontiers of Biol. Energetics* 1, 752-759 (1978); *J. Nucl. Med.*
35 25, 506-513 (1984); *Proc. of NMR Imaging Symp.* (Oct. 26-27, 1980); F. A. Cotton et al., *Adv.*
Inorg. Chem. 634-639 (1966)]. Paramagnetic metal chelates used in this manner are referred to
as contrast enhancement agents or contrast agents.


There are a number of paramagnetic metal ions which can be considered when undertaking the design of an MRI contrast agent. In practice, however, the most useful paramagnetic metal ions are gadolinium (Gd^{+3}), iron (Fe^{+3}), manganese (Mn^{+2}) and (Mn^{+3}), and chromium (Cr^{+3}), because these ions exert the greatest effect on water protons by virtue of their large magnetic moments. In a non-complexed form (e.g. $GdCl_3$), these metal ions are toxic to an animal, thereby precluding their use in the simple salt form. Therefore, a fundamental role of the organic chelating agent (also referred to as a ligand) is to render the paramagnetic metal non-toxic to the animal while preserving its desirable influence on T1 and T2 relaxation rates of the surrounding water protons.

Art in the MRI field is quite extensive, such that the following summary, not intended to be exhaustive, is provided only as a review of this area and other compounds that are possibly similar in structure. U.S. Patent 4,899,755 discloses a method of alternating the proton NMR relaxation times in the liver or bile duct of an animal using Fe^{+3} -ethylene-bis(2-hydroxyphenylglycine) complexes and its derivatives, and suggests among various other compounds the possible use of a pyridine macrocyclomethylenecarboxylic acid. U.S. Patent 4,880,008 (a CIP of U.S. Patent 4,899,755) discloses additional imaging data for liver tissue of rats, but without any additional complexes being shown. U.S. Patent 4,980,148 discloses gadolinium complexes for MRI which are non-cyclic compounds. C. J. Broan *et al.* [*J. Chem. Soc., Chem. Commun.*, 1739-1741 (1990)] describe some bifunctional macrocyclic phosphinic acid compounds. C. J. Broan *et al.* [*J. Chem. Soc., Chem. Commun.*, 1738-1739 (1990)] describe compounds that are triazabicyclo compounds. I. K. Adzamli *et al.* [*J. Med. Chem.* **32**, 139-144 (1989)] describe acyclic phosphonate derivatives of gadolinium complexes for NMR imaging.

At the present time, the only commercial contrast agents available in the U.S. are the complex of gadolinium with diethylenetriaminepentaacetic acid (DTPA- Gd^{+3} - Magnevist™ by Schering) and a DO3A derivative [1,4,7-tris(carboxymethyl)-10-(2-hydroxypropyl)-1,4,7,10-tetraazacyclododecanato]gadolinium (Prohance™ by Squibb). Magnevist™ and Prohance™ are considered as non-specific/perfusion agents since they freely distribute in extracellular fluid followed by efficient elimination through the renal system. Magnevist™ has proven to be extremely valuable in the diagnosis of brain lesions since the accompanying breakdown of the blood/brain barrier allows perfusion of the contrast agent into the affected regions. In addition to Magnevist™, Guerbet is commercially marketing a macrocyclic perfusion agent [1,4,7,10-tetrakis(carboxymethyl)-1,4,7,10-tetraazacyclododecanato]gadolinium (Dotarem™) which presently is only available in Europe. Prohance™ is shown to have fewer side effects than Magnevist™. A number of other potential contrast agents are in various stages of development.

The present invention is directed to novel ligands that are 2-pyridyl-methylenepolyazamacrocyclic compounds, and derivatives thereof, of the formula

5




(I)

10

wherein:


T is independently -CH₂-COOH,

15



or

20



where: R is OH, C₁-C₅ alkyl or -O-(C₁-C₅ alkyl);

R₁ is OH or OCH₃;


R₂ is NO₂, NH₂, isothiocyanato, semicarbazido, thiosemicarbazido, maleimido,

25

bromoacetamido or carboxyl;

with the proviso that only one T moiety is

30



where R₁ and R₂ are defined as before; or

pharmaceutically-acceptable salts thereof.

35

When the above ligands of Formula (I) have:

all T equal P(O)ROH, where R is OH, or the pharmaceutically-acceptable salts

thereof, the ligands are referred to herein as PD3P;

all T equal COOH, or the pharmaceutically-acceptable salts thereof, the ligands are referred to herein as PD3A;

all T equal P(O)ROH, where R is -O-(C₁-C₅ alkyl), or the pharmaceutically-acceptable salts thereof, the ligands are referred to herein as PD3(O-Alk)P; and

5 all T equal P(O)ROH, where R is C₁-C₅ alkyl, or the pharmaceutically-acceptable salts thereof, the ligands are referred to herein as PD3(Alk)P.

The complexes of this invention can be designed to provide a specific overall charge which advantageously influences the *in vivo* biolocalization and image contrast. For example, when the metal ion is + 3 the following can be obtained:

10 an overall charge of -3 - when Formula (I) is PD3P, which are useful as calcific tissue contrast agents;

an overall charge of -2 - when Formula (I) has two T equal to -CH₂-PO₃H₂ and one T has a COOH group present, which are useful as calcific tissue contrast agents;


15 an overall charge of -1 - when Formula (I) has one T equal to -CH₂-PO₃H₂ and the other T has a COOH group present, which are useful as calcific tissue contrast agents;

20 an overall neutral charge of 0 - when Formula (I) has one T equal to -CH₂-PO₃HR where R is -O-(C₁-C₅ alkyl) or C₁-C₅ alkyl and two T have a COOH group present; or all three T equal -CH₂-PO₃HR where R is -O-(C₁-C₅ alkyl) or C₁-C₅ alkyl; or all three T equal -CH₂-COOH (PD3A), which are useful as general perfusion, blood pool, brain or liver contrast agents.

The complexes may be formulated to be in a pharmaceutically acceptable form for administration to an animal.

When one T term is

25




30

where R¹ and R² are defined as before, then the compound is a bifunctional ligand/complex and may be linked through R² to a biologically active molecule.

The compounds of Formula (I) are numbered for nomenclature purposes as follows:

35



One aspect of the present invention concerns development of contrast agents having synthetic modifications to the paramagnetic chelate enabling site specific delivery of the contrast agent to a desired tissue. The advantage being increased contrast in the areas of interest based upon tissue affinity as opposed to contrast arising from non-specific perfusion which may or may not be apparent with an extracellular agent. The specificity of the ligand of Formula (I) may be controlled by adjusting the total charge and lipophilic character of the complex. The overall range of the charge of the complex is from -3 to 0 as indicated above. For example, for a complex having 3 PO_3H_2 groups (PD3P), the overall charge is highly negative and bone uptake is expected; whereas when the overall charge of the complex is 0 (thus neutral, PD3A, PD3(O-Alk)P, and PD3(Alk)P), the complex may have the ability to cross the blood brain barrier and normal brain uptake may be possible. Unexpectedly, for a complex having an overall charge of 0 [PD3(O-Pr)P], the complex displays liver uptake.

25 While not wishing to be bound by theory, it is believed that when a charged complex of the invention is made (e.g. possibly -3 for bone, -1 for liver, or + 1 for heart), the variations in that chelate ionic charge can influence biolocalization.

Tissue specificity may also be realized by ionic or covalent attachment of the chelate to a naturally occurring or synthetic molecule (e.g. through R^2) having specificity for a desired target tissue. One possible application of this approach is through the use of chelate conjugated monoclonal antibodies which would transport the paramagnetic chelate to diseased tissue enabling visualization by MRI. In addition, attachment of a paramagnetic chelate to a macromolecule can further increase the contrast agent efficiency resulting in improved contrast relative to the unbound chelate. Recent work by Lauffer (U.S. Patents 30 4,880,008 and 4,899,755) has demonstrated that variations in lipophilicity can result in tissue-specific agents and that increased lipophilic character favors non-covalent interactions with blood proteins resulting in enhancement of relaxivity.

Typical DO3A derivatives in which non-coordinating N-substituents are present generate neutral lanthanide chelates but have been found to be kinetically labile under acidic conditions. The DO3A derivative PROHANCE™ gains added kinetic stability by virtue of a pendant hydroxyl moiety which can coordinate to the metal ion but may alter the overall chelate ionic character.

In contrast, the present invention relates to a group of DO3A derivatives of Formula (I) which possess a 2-pyridyl pendant moiety which forms a coordinate bond without altering the overall chelate ionic character. In addition it has now been found that the 2-pyridyl substituent adds kinetic stability to the chelate under acidic conditions (pH below about 7). Chelates of this invention in which the ligation moieties are PO_3HR where R is $-\text{O}-(\text{C}_1-\text{C}_5$ alkyl) have demonstrated potential as liver imaging agents.

The terms used in Formula (I) and for this invention are further defined as follows:

- "C₁-C₅ alkyl", include both straight and branched chain alkyl groups.
- "Animal" includes a warmblooded mammal, preferably a human being.
- "Complex" refers to a complex of the compound of the invention, e.g. Formula (I), complexed with a metal ion, where at least one metal atom is chelated or sequestered.
- "Biologically active material" refers to a dextran, peptide, or molecules that have specific affinity for a receptor, or preferably antibodies or antibody fragments.
- "Antibody" refers to any polyclonal, monoclonal, chimeric antibody or heteroantibody, preferably a monoclonal antibody; "antibody fragment" includes Fab fragments and $\text{F}(\text{ab}')_2$ fragments, and any portion of an antibody having specificity toward a desired epitope or epitopes. When using the term "radioactive metal chelate/antibody conjugate" or "conjugate", the "antibody" is meant to include whole antibodies and/or antibody fragments, including semisynthetic or genetically engineered variants thereof. Possible antibodies are 1116-NS-19-9 (anti-colorectal carcinoma), 1116-NS-3d (anti-CEA), 703D4 (anti-human lung cancer), 704A1 (anti-human lung cancer), CC49, CC83 and B72.3. The hybridoma cell lines 1116-NS-19-9, 1116-NS-3d, 703D4, 704A1, CC49, CC83 and B72.3 are deposited with the American Type Culture Collection, having the accession numbers ATCC HB 8059, ATCC CRL 8019, ATCC HB 8301, ATCC HB 8302, ATCC HB 9459, ATCC HB 9453 and ATCC HB 8108, respectively.

The bifunctional chelating agents described herein [represented by Formula (I)] can be used to chelate or sequester the metal ions so as to form metal ion chelates (also referred to herein as "complexes"). The complexes, because of the presence of the functionalizing moiety [represented by R^2 in Formula (I)], can be covalently attached to biologically active materials, such as dextran, molecules that have specific affinity for a receptor, or preferably covalently attached to antibodies or antibody fragments. Thus the complexes described herein

may be covalently attached to an antibody or antibody fragment or have specific affinity for a receptor and are referred to herein as "conjugates".

As used herein, "pharmaceutically-acceptable salts" means any salt or mixtures of salts of a compound of Formula (I) which is sufficiently non-toxic to be useful in diagnosis of animals, preferably mammals. Thus, the salts are useful in accordance with this invention. Representative of those salts formed by standard reactions from both organic and inorganic sources include, for example, sulfuric, hydrochloric, phosphoric, acetic, succinic, citric, lactic, maleic, fumaric, palmitic, cholic, palmoic, mucic, glutamic, gluconic acid, d-camphoric, glutaric, glycolic, phthalic, tartaric, formic, lauric, steric, salicylic, methanesulfonic, benzenesulfonic, sorbic, picric, benzoic, cinnamic acids and other suitable acids. Also included are salts formed by standard reactions from both organic and inorganic sources such as ammonium or 1-deoxy-1-(methylamino)-D-glucitol, alkali metal ions, alkaline earth metal ions, and other similar ions. Particularly preferred are the salts of the compounds of Formula (I) where the salt is potassium, sodium, or ammonium. Also included are mixtures of the above salts.

The compounds of Formula (I) are prepared by various processes. Typical general synthetic approaches to such processes are provided by the reaction schemes given below.

The starting material of Formula (II) shown in these Schemes is prepared by alkylation of 1,4,7,10-tetraazacyclododecane ("cyclen") with 2-chloromethyl pyridine at room temperature in an inert organic solvent such as chloroform.


In Scheme 1, the compounds of Formula (I) are prepared wherein all T moieties are -CH₂-COOH.

25

30


35

Scheme 1



The aqueous base is alkali metal hydroxide, such as sodium hydroxide or potassium hydroxide. The pH of the reaction is maintained about 8-12. Temperature is between about 60-90°C, and pressure is not critical so that ambient pressure is used.


25 In Scheme 2, the compounds of Formula (I) are prepared wherein all T moieties are



30 where: R is OH.


35

Scheme 2



The hydrolysis in Scheme 2 is done by known aqueous acid conditions, such as
 25 using 3 to 12M hydrochloric acid. The pH of the reaction is maintained below 3. Temperature
 is at reflux. Pressure is not critical so that ambient pressure is used. Alternatively, when the
 reaction is run in one step, phosphorous acid, hydrochloric acid and excess formaldehyde are
 used. The pH of the reaction is below 2. Temperature is at reflux. Pressure is not critical so that
 ambient pressure is used.




30 In Scheme 3, the compounds of Formula (I) are prepared wherein all T moieties
 are



35

where: R is -O-(C₁-C₅ alkyl).

Scheme 3




wherein:

R is -O-(C₁-C₅ alkyl).

25 The hydrolysis in Scheme 3 is done under known aqueous basic conditions, such as using excess alkali metal hydroxide, such as sodium or potassium hydroxide. The pH of the reaction is above 9. Temperature is at reflux. Pressure is not critical so that ambient pressure is used.




30 In Scheme 4, the compounds of Formula (I) are prepared wherein all T moieties are



35 where: R is C₁-C₅ alkyl.

Scheme 4 shows the preparation of the compounds of Formula (I) when R is methyl.

Scheme 4



The hydrolysis in Scheme 4 is done under known aqueous basic conditions, such as using excess alkali metal hydroxide, such as sodium or potassium hydroxide. The pH of the reaction is above 9. Temperature is at reflux. Pressure is not critical so that ambient pressure is used.


25

Scheme 5 shows the preparation of the compounds of Formula (I) when R is ethyl.

30

35


Scheme 5



The reaction in Scheme 5 is under acidic conditions with a hydrochloric acid. The pH of the reaction is below 3. Temperature is at reflux. Pressure is not critical so that ambient pressure is used.

Scheme 6 shows the preparation of the compounds of Formula (I) when T has a bifunctional moiety.

Scheme 6




(next page)

25

30

35

Scheme 6 Cont'd




Scheme 7 shows the preparation of the compounds of Formula (I) when one T is PO_3HR where R is defined as in Formula (I) and the other two T are COOH .

30

35

Scheme 7



30

In Scheme 7 the hydrolysis steps are done as described in prior schemes. The formaldehyde/bisulfite addition complex reaction is done aqueous basic conditions to give selective alkylation at the 4,10-positions. The phosphonate group is then introduced selectively at the 7-position.

35

In the above Schemes, the general process description illustrates specific steps that may be used to accomplish a desired reaction step. The general description of these process steps follows.

The carboxylate derivatives described in Scheme 1 are prepared by conventional alkylation procedures utilizing chloro- or bromo-acetic acid under basic aqueous conditions.

The phosphonic acid derivatives outlined in Scheme 2 can be prepared by initial alkylation of the amine with a trialkyl phosphite and paraformaldehyde, resulting in an organic
5 soluble perester. This ester is then hydrolyzed under refluxing acid conditions to give the desired aminophosphonic acids. Alternatively, the phosphonic acid can be prepared under acidic conditions by employing phosphorous acid in combination with formaldehyde and hydrochloric acid.

Phosphonate half esters are prepared as shown in Scheme 3 by initial formation
10 of the dialkyl phosphonate ester, followed by hydrolysis under basic conditions. Base hydrolysis gives exclusive conversion to the half ester.

Scheme 4 illustrates the methodology for synthesizing the methyl phosphinate derivatives using diethoxymethylphosphine as the nucleophile and paraformaldehyde. Condensation can be conducted in solvents such as tetrahydrofuran (THF), dimethylformamide
15 (DMF), dioxane, acetonitrile, or alcoholic media. The resulting phosphinate ester is then hydrolyzed under acidic conditions (e.g. 6N HCl, 80-100°C) or basic conditions (excess base, 40-100°C) to give the corresponding methylphosphonic acid. Alternatively, as outlined in Scheme 5, the method devised by A. D. Sherry *et al.* (*Inorg. Chem.*, submitted 1991) using ethylphosphonic acid generated *in situ* can be used to obtain phosphinate derivatives having
20 increased lipophilic character.

Scheme 6 illustrates the preparation of bifunctional compounds of Formula (I) which may then be attached to a biologically active material.

Scheme 7 shows a general procedure for the selective alkylation of the 4,10-positions using the formaldehyde/bisulfite addition compound. The resulting intermediate is
25 converted to the corresponding nitrile, followed by introduction of the phosphonate moiety at the the 7-position. Acid or base hydrolysis yields the corresponding hydrolyzed final products.

The metal ions used to form the complexes of this invention are Gd^{+3} , Mn^{+2} , Fe^{+3} and available commercially, e.g. from Aldrich Chemical Company. The anion present is halide, preferably chloride, or salt free (metal oxide).

30 A "paramagnetic nuclide" of this invention means a metal ion which displays spin angular momentum and/or orbital angular momentum. The two types of momentum combine to give the observed paramagnetic moment in a manner that depends largely on the atoms bearing the unpaired electron and, to a lesser extent, upon the environment of such atoms. The paramagnetic nuclides found to be useful in the practice of the invention are gadolinium
35 (Gd^{+3}), iron (Fe^{+3}) and manganese (Mn^{+2}), with Gd^{+3} being preferred.

The complexes are prepared by methods well known in the art. Thus, for example, see Chelating Agents and Metal Chelates, Dwyer & Mellor, Academic Press (1964), Chapter 7. See also methods for making amino acids in Synthetic Production and Utilization of

Amino Acids, (edited by Kameko, et al.) John Wiley & Sons (1974). An example of the preparation of a complex involves reacting a bicyclopolyazamacrocyclophosphonic acid with the metal ion under aqueous conditions at a pH from 5 to 7. The complex formed is by a chemical bond and results in a stable paramagnetic nuclide composition, e.g. stable to the
5 disassociation of the paramagnetic nuclide from the ligand.

The complexes of the present invention are administered at a ligand to metal molar ratio of at least about 1:1, preferably from 1:1 to 3:1, more preferably from 1:1 to 1.5:1. A large excess of ligand is undesirable since uncomplexed ligand may be toxic to the animal or may result in cardiac arrest or hypocalcemic convulsions.

10 This invention is used with a physiologically acceptable carrier, excipient or vehicle therefore. The methods for preparing such formulations are well known. The formulations may be in the form of a suspension, injectable solution or other suitable formulations. Physiologically acceptable suspending media, with or without adjuvants, may be used.

15 An "effective amount" of the formulation is used for diagnosis. The dose will vary depending on the disease and physical parameters of the animal, such as weight. *In vivo* diagnostics are also contemplated using formulations of this invention.

Other uses of some of the chelants of the present invention may include the removal of undesirable metals (i.e. iron) from the body, attachment to polymeric supports for
20 various purposes, e.g. as diagnostic agents, and removal of metal ions by selective extraction. The ligands of Formula (I) having in at least two R terms T equal to $-\text{CH}_2-\text{P}(\text{O})\text{R}^1\text{OH}$ may be used for metal ion control as scale inhibitors. Some of these ligands can be used in less than stoichiometric amounts. Similar uses are known for compounds described in U.S. Patents 2,609,390; 3,331,773; 3,336,221; and 3,434,969.

25 The invention will be further clarified by a consideration of the following examples, which are intended to be purely exemplary of the present invention.

Some terms used in the following examples are defined as follows:

LC = liquid chromatography, purifications were carried out at low pressure using
30 Dionex 2010i system fitted with a hand-packed Q-Sepharose™ anion exchange column (23 x 2 cm).

DMF = dimethylformamide.

AcOH = acetic acid.

g = gram(s).

mg = milligrams.

35 kg = kilogram(s).

mL = milliliter(s).

μL = microliter(s).

pH Stability General Procedure

A stock $^{159}\text{GdCl}_3$ (or $^{153}\text{SmCl}_3$) solution was prepared by adding 2 μL of $3 \times 10^{-4}\text{M}$ $^{159}\text{GdCl}_3$ in 0.1N HCl to 2 mL of a $3 \times 10^{-4}\text{M}$ GdCl_3 carrier solution. Appropriate ligand solutions were then prepared in deionized water. The 1:1 ligand/metal complexes were then prepared by combining the ligands (dissolved in 100-500 μL of deionized water) with 2 mL of the stock $^{159}\text{GdCl}_3$ solution, followed by thorough mixing to give an acidic solution (pH = 2). The pH of the solution was then raised to 7.0 using 0.1N NaOH. The percent metal as a complex was then determined by passing a sample of the complex solution through a Sephadex™ G-50 column, eluting with 4:1 saline (85% NaCl/ NH_4OH) and collecting 2 x 3 mL fractions. The amount of radioactivity in the combined elutions was then compared with that left on the resin (non-complexed metal is retained on the resin). The pH stability profile was generated by adjusting the pH of an aliquot of the complex solution using 1M NaOH or 1M HCl and determining the percent of the metal existing as a complex using the ion exchange method described above. The Sm results are known by experimental comparison to be identical for complexation and biodistribution of the ligands of this invention.

SYNTHESIS OF LIGANDS.

General Materials and Methods.

All reagents were obtained from commercial suppliers and used as received without further purification. NMR spectra were recorded on a Bruker AC-250 MHz spectrometer equipped with a multi-nuclear quad probe (^1H , ^{13}C , ^{31}P , and ^{19}F) at 297°K unless otherwise indicated. ^1H spectra in D_2O were recorded by employing solvent suppression pulse sequence ("PRESAT", homo-nuclear presaturation). ^1H spectra are referenced to residual chloroform (in CDCl_3) at $\delta 7.26$ or external dioxane (in D_2O) at $\delta 3.55$. ^{13}C and ^{31}P spectra reported are proton decoupled (broad band). Assignments of ^{13}C { ^1H } chemical shifts were aided by DEPT (Distortionless Enhancement by Polarization Transfer) experiments. ^{13}C { ^1H } spectra are referenced to center peak of CDCl_3 at $\delta 77.00$ (in CDCl_3) and external dioxane at $\delta 66.66$ (in D_2O). ^{31}P { ^1H } spectra were referenced to external 85% H_3PO_4 at $\delta 0.00$. Melting points were determined by capillary melt methods and were uncorrected. Semipreparative ion-exchange chromatographic separations were performed at low pressure (< 600 psi) using a standard glass column fitted with hand-packed Q-Sepharose™ (anion exchange) or SP-Sepharose™ (cation exchange) glass column, and with on-line UV detector at 263 nm for eluent monitoring. GC/MS spectra were performed on a Hewlett Packard 5890A Gas Chromatograph/5970 Mass Selective Detector.

STARTING MATERIALS

35 Example A

Preparation of N-(2-pyridylmethyl)-1,4,7,10-tetraazacyclododecane.

A 2-picoyl chloride solution in chloroform was prepared by basifying 3 mL of an aqueous solution of 1.03 g (6.3 mmol) of 2-picoyl chloride hydrochloride to pH > 14 and

extracting into chloroform. To a stirred chloroform solution of 2.03 g (11.8 mmol) of 1,4,7,10-tetraazacyclododecane was added 10 mL of the prepared 2-picolyl chloride solution in chloroform in one portion. After stirring at temperature for 30 min, the reaction mixture was concentrated *in vacuo* to give a residue which was chromatographed on silica gel (column, 2.5 X 20 cm), eluting with $\text{CHCl}_3:\text{CH}_3\text{OH}:\text{NH}_4\text{OH}$ (10:4:1), $R_f = 0.29$ on SiO_2 plate. After concentration of eluent, the mono-alkylated product was isolated as a thick pale yellow liquid, which upon standing solidified to give 1.17 g of an off-white powder (72%, based on 2-picolyl chloride), and further characterized by:

^1H NMR (CDCl_3)
 10 δ 2.54 - 2.82 (m, 16H), 3.75 (s, 2H), 7.11 - 7.14 (m, 1H), 7.42 - 7.46 (m, 1H), 7.64 - 7.65 (m, 1H), 8.46 - 8.48 (m, 1H); and
 ^{13}C { ^1H } NMR (CDCl_3)
 δ 45.02, 46.31, 47.04, 51.51, 61.03, 122.04, 122.87, 136.61, 148.83, 150.53.

Example B

15 Preparation of N-(2-pyridylmethyl)-N',N'',N'''-tris(methylenediethylphosphonate)-1,4,7,10-tetraazacyclododecane.

To 331 mg (1.27 mmol) of N-(2-pyridylmethyl)-1,4,7,10-tetraazacyclododecane, prepared in Example A, was combined 229 mg (7.63 mmol, excess) of paraformaldehyde and 1.3 mL of 1.27 g (7.62 mmol, excess) of triethylphosphite. After the mixture was gently stirred for 10 min to obtain a well mixed slurry, it was heated to 90°C for 1 hr. The excess reagents and the byproducts were removed *in vacuo* (125°C/0.01 mmHg) to give 896 mg (99%) of the desired product as a yellow oil, and further characterized by:

^1H NMR (CDCl_3)
 δ 1.25 - 1.39 (m, 18H), 2.66 - 2.95 (m, 22H), 3.71 (s, 2H), 4.01 - 4.22 (m, 12H), 7.10 - 7.15 (m, 1H),
 25 7.57 - 7.65 (m, 2H), 8.46 - 8.52 (m, 1H);
 ^{13}C { ^1H } NMR (CDCl_3)
 δ 16.38, 16.46, 50.45, 50.67, 52.41, 53.19, 53.29, 53.48, 53.58, 61.37, 61.47, 61.52, 121.67, 123.28, 136.19, 148.61, 159.90; and
 ^{31}P { ^1H } NMR (CDCl_3 , 297°K)
 30 δ 26.21;
 ^{31}P { ^1H } NMR (CDCl_3 , 217°K)
 δ 24.18 (1P), 24.32 (2P).

Example C

35 Preparation of N-(2-pyridylmethyl)-N',N'',N'''-tris(methylenedipropylphosphonate)-1,4,7,10-tetraazacyclododecane.

To 445 mg (1.71 mmol) of N-(2-pyridylmethyl)-1,4,7,10-tetraazacyclododecane, prepared in Example A, was combined 154 mg (5.12 mmol, excess) of paraformaldehyde and 3.5 mL of 3.20 g (5.12 mmol, excess) of tripropylphosphite. After the mixture was gently stirred

for 10 min to obtain a well mixed slurry, it was heated to 100°C for 30 min. The excess reagents and the byproducts were removed *in vacuo* (4 hr. at 160°C/0.01 mmHg) to give 1.36 g (quantitative) of the desired product as a thick yellow oil, and further characterized by:

¹H NMR (CDCl₃)

5 δ 0.91 - 1.00 (m, 18H), 1.60 - 1.76 (m, 12H), 2.67 - 2.99 (m, 22H), 3.73 (s, 2H), 3.94 - 4.08 (m, 12H), 7.12 - 7.15 (m, 1H), 7.46 - 7.67 (m, 2H), 8.48 - 8.52 (m, 1H);

¹³C {¹H} NMR (CDCl₃)

δ 9.93, 10.21, 23.71, 23.80, 50.17, 50.44, 52.38, 53.09, 53.44, 61.44, 66.79, 66.83, 121.61, 123.23, 136.14, 148.54, 159.92; and

10 ³¹P {¹H} NMR (CDCl₃)

δ 26.20 (1P), 26.23 (2P).

FINAL PRODUCTS

Ligands: Preparation of dimethylenecarboxylic acids as shown in Scheme 1.

Example 1

15 Preparation of N-(2-pyridylmethyl)-N',N'',N'''-triacetic acid-1,4,7,10-tetraazacyclododecane (PD3A).

To 2 mL of an aqueous solution of 201.4 mg (0.51 mmol) of N-(2-pyridylmethyl)-1,4,7,10-tetraazacyclododecane, prepared by the procedure of Example A, was added 350 mg (2.52 mmol, 65% excess) of bromoacetic acid and the pH of the reaction mixture was maintained above 11 by adding small portions of concentrated sodium hydroxide until no more caustic was needed to keep pH > 11 (~30 min). The reaction mixture was then heated (60°C) for 1 hr. After the reaction mixture was cooled to room temperature, the pH of the reaction mixture was adjusted to 7 and the neutral solution chromatographed on a cation exchange (SP-Sepharose™) column (1.5 X 50 cm) eluting first with deionized water then with 20 1M hydrochloric acid. The acidic fraction which contained product was evaporated to dryness, followed by co-evaporation (azeotrope) with fresh deionized water (3 X 2 mL) to eliminate excess hydrochloric acid. The final product was isolated as white solid upon freeze-drying of the aqueous solution, and further characterized by:

¹H NMR (D₂O)

30 δ 2.79 - 4.13 (m, 24H), 7.88 - 8.01 (m, 2H), 8.35 - 8.42 (m, 1H), 8.63 - 8.66 (m, 1H); and

¹³C {¹H} NMR (D₂O)

δ 51.04, 51.25, 53.73, 55.47, 56.00, 56.90, 57.52, 129.86, 131.21, 145.49, 150.82, 153.54, 171.61, 178.76. Ligands: Preparation of dimethylenephosphonate half esters as shown in Scheme 3.

Example 2

35 Preparation of N-(2-pyridylmethyl)-N',N'',N'''-tris(methylenephosphonic acid ethyl ester)-1,4,7,10-tetraazacyclododecane (PD3EP).

N-(2-pyridylmethyl)-N',N'',N'''-tris(methylenediethylphosphonate)-1,4,7,10-tetraazacyclododecane, 102.7 mg (0.14 mmol), prepared in Example B, was combined with 1 mL of

0.1M of potassium hydroxide and heated at 90°C for 6 hr. After cooling to room temperature, the aqueous solution was freeze-dried to give a tan solid, which was chromatographed on an anion exchange (Q-Sepharose™) column (1.5 X 50 cm) eluting first with deionized water, then with 1M hydrochloric acid. Following freeze-drying of the eluent, the product was isolated as a brown solid and further characterized by:

¹H NMR (D₂O, 338°K)
δ 1.41 - 1.57 (m, 9H), 3.28 - 3.89 (m, 22H), 4.09 - 4.64 (m, 8H), 8.22 - 8.26 (m, 2H), 8.70 - 8.75 (m, 1H), 9.00 - 9.12 (m, 1H); and
¹³C {¹H} NMR (D₂O, 338°K)
δ 19.41, 19.51, 52.58, 53.00, 52.31, 53.75, 53.82, 56.04, 59.53, 64.60, 64.76, 129.86, 131.41, 147.31, 149.06, 154.34; and
³¹P {¹H} NMR (D₂O, 338°K)
δ 9.64 (2P), 19.79 (1P).

Example 3

Preparation of N-(2-pyridylmethyl)-N',N'',N'''-tris(methylenephosphonic acid propyl ester)-1,4,7,10-tetraazacyclododecane (PD3PP).

N-(2-pyridylmethyl)-N',N'',N'''-tris(methylenedipropylphosphonate)-1,4,7,10-tetraazacyclododecane, 0.71 g (0.89 mmol), prepared in Example C, was combined with 3 mL of 0.1M potassium hydroxide and heated at reflux for 18 hr. After cooling to room temperature, the aqueous solution was filtered, and freeze-dried to give a brown residue, which was redissolved into CH₂Cl₂/C₂H₅OH (95:5) and filtered. Following evaporation of solvent and concentration in vacuo, the product was isolated as a brown solid and further characterized by:

¹H NMR (D₂O, 353°K)
δ 1.24 - 1.36 (m, 9H), 1.95 - 2.04 (m, 6H), 3.03 - 3.29 (m, 22H), 4.10 - 4.25 (m, 8H), 7.74 - 7.92 (m, 2H), 8.23 - 8.29 (m, 1H), 8.87 - 8.96 (m, 1H); and
¹³C {¹H} NMR (D₂O, 353°K)
δ 13.15, 27.20, 50.43, 53.89, 54.48, 54.98, 55.42, 64.33, 69.41, 126.38, 128.30, 141.24, 152.46, 161.45; and
³¹P {¹H} NMR (D₂O, 353°K)
δ 21.61 (2P), 21.95 (1P).

Complexes: Preparation of metal/ligand complexes for biodistribution studies.

General Procedure

Metal ligand complexes were prepared by various methods. The methods included mixing of the metal and ligand in aqueous solution and adjusting the pH to the desired value. Complexation was done in solutions containing salts and/or buffers as well as water. Sometimes heated solutions were found to give higher complex yields than when the complexation was done at ambient temperatures.

For example, a solution of the ligand is prepared by dissolving the ligand in deionized water (about pH = 2). A ligand/metal complex was then prepared by combining the ligand solution with aqueous $\text{SmCl}_3 \cdot \text{H}_2\text{O}$ ($3 \times 10^{-4}\text{M}$ in 0.01N HCl) containing tracer $^{153}\text{SmCl}_3$. After thorough mixing, the percent metal as a complex was determined by passing a sample of the complex solution through a Sephadex™ column, eluting with 4:1 saline (0.85% NaCl/ NH_4OH), and collecting 2 x 3 mL fractions. The amount of radioactivity in the combined elutions was then compared with that left on the resin. Under these conditions, complex was removed with the eluent and non-complexed metal is retained on the resin. By this method complexation was determined to usually be about 95% or greater.

Using the above procedure, the complexes of samarium with N-(2-pyridylmethyl)-N',N'',N'''-triacetic acid-1,4,7,10-tetraazacyclododecane (PD3A);

N-(2-pyridylmethyl)-N',N'',N'''-tris(methylenephosphonic acid ethyl ester)-1,4,7,10-tetraazacyclododecane (PD3EP);

N-(2-pyridylmethyl)-N',N'',N'''-tris(methylenephosphonic acid propyl ester)-1,4,7,10-tetraazacyclododecane (PD3PP); and

N-(2-pyridylmethyl)-N',N'',N'''-tri(methylenephosphonic acid-1,4,7,10-tetraazacyclododecane (PD3P);

were made. Complexes with gadolinium are made in a similar manner.

The following tables illustrate the kinetic inertness of the Sm complexes derived from PD3A, PD3PP and PD3EP.

pH Stability Profile
of ^{153}Sm -PD3A

pH	% Complex
1	98.30
2	99.38
4	99.62
7	99.86
9	99.86
11	99.80
14	99.96

pH Stability Profile
of ¹⁵³Sm-PD3PP

	pH	% Complex
5	1	72.52
	3	87.56
	5	92.99
	7	95.06
10	9	95.22
	11	92.74
	14	87.65

pH Stability Profile
of ¹⁵³Sm-PD3EP

	pH	% Complex
15	1	78.95
	3	93.80
20	5	93.02
	7	93.77
	9	95.89
	11	95.77
25	14	94.07

BIODISTRIBUTION

General Procedure

30 Sprague Dawley rats were allowed to acclimate for five days then injected with 100 µL of the complex solution via a tail vein. The rats weighed between 150 and 200 g at the time of injection. After 30 min. the rats were killed by cervical dislocation and dissected. The amount of radioactivity in each tissue was determined by counting in a NaI scintillation counter coupled to a multichannel analyzer. The counts were compared to the counts in 100 µL standards in order to determine the percentage of the dose in each tissue or organ.

35 The percent dose in blood was estimated assuming blood to be 6.5% of the total body weight. The percent dose in bone was estimated by multiplying the percent dose in the femur by 25. The percent dose in muscle was estimated assuming muscle to be 43% of the total body weight. The number of rats used in the average is n.

In addition to organ biodistribution, chelates of the compounds of Formula (I) were evaluated for efficiency of bone localization since phosphonates are known for their ability to bind to hydroxyapatite. The results of these tests are given below.

EXAMPLE I

5 When the complex $^{153}\text{Sm}[\text{N}-(2\text{-pyridylmethyl})\text{-N}',\text{N}'',\text{N}''']\text{-tris(methylenephosphonic acid ethyl ester)-1,4,7,10-tetraazacyclododecane}$ ($^{153}\text{Sm-PD3EP}$) was evaluated, the results are given in Table I below. The numbers represent the average of a minimum of 2 rats per data point at 2 hours post injection.

10 **TABLE I**
BIODISTRIBUTION OF $^{153}\text{Sm-PD3EP}$
% INJECTED DOSE

ORGAN	AVERAGE
Bone	0.82
Liver	0.36
15 Kidney	1.24
Spleen	0.01
Muscle	0.62
Blood	0.45
20 Heart	0.01
Lung	0.02
Brain	0.01
Urine	97.00

25 EXAMPLE II

When the complex $^{153}\text{Sm}[\text{N}-(2\text{-pyridylmethyl})\text{-N}',\text{N}'',\text{N}''']\text{-tris(methylenephosphonic acid propyl ester)-1,4,7,10-tetraazacyclododecane}$ ($^{153}\text{Sm-PD3PP}$) was evaluated, the results are given in Table II below. The numbers represent the average of a minimum of 3 rats per data point at 2 and 24 hours post injection.

30

35


TABLE II
BIODISTRIBUTION OF ^{153}Sm -PD3PP
% INJECTED DOSE

ORGAN	AVERAGE 2 hrs	AVERAGE 24 hrs
Bone	5.93	12.39
Liver	20.36	2.31
Kidney	0.86	0.89
Spleen	1.19	0.22
Muscle	0.82	0.27
Blood	0.68	0.03
Heart	0.04	0.02
Lung	0.20	0.40
Brain	0.04	0.00
Stomach	1.39	0.04
Small Intestine	13.56	0.13
Large Intestine	0.18	0.16
Urine	50.00	83.00
Feces	-	5.33

Other embodiments of the invention will be apparent to those skilled in the art from a consideration of this specification or practice of the invention disclosed herein. It is intended that the specification and examples be considered as exemplary only, with the true scope and spirit of the invention being indicated by the following claims.



WHAT IS CLAIMED IS:

1. 2-Pyridylmethylenepolyazamacrocyclic compounds of the formula



wherein:

T is independently -CH₂-COOH,




where: R is OH, C₁-C₅ alkyl or -O-(C₁-C₅ alkyl);

25 R¹ is OH or OCH₃;

R² is NO₂, NH₂, isothiocyanato, semicarbazido, thiosemicarbazido, maleimido, bromoacetamido or carboxyl;

with the proviso that only one T moiety is



35 where R¹ and R² are defined as before; or pharmaceutically-acceptable salts thereof.

2. A compound of Claim 1 wherein T equal $-\text{CH}_2-\text{P}(\text{O})\text{ROH}$, where R is OH, or pharmaceutically-acceptable salts thereof, and is named as N-(2-pyridylmethyl)-N',N'',N'''-tri(methylenephosphonic acid-1,4,7,10-tetraazacyclododecane.

3. A compound of Claim 1 wherein T equal $-\text{CH}_2-\text{COOH}$, or pharmaceutically-acceptable salts thereof, and is named as N-(2-pyridylmethyl)-N',N'',N'''-triacetic acid-1,4,7,10-tetraazacyclododecane.


4. A compound of Claim 1 wherein T equal $-\text{CH}_2-\text{P}(\text{O})\text{ROH}$, where R is $-\text{O}-(\text{C}_1-\text{C}_5 \text{ alkyl})$, or pharmaceutically-acceptable salts thereof.

5. A compound of Claim 4 wherein R is ethoxy, or pharmaceutically-acceptable salts thereof, and is named as N-(2-pyridylmethyl)-N',N'',N'''-tris(methylenephosphonic acid ethyl ester)-1,4,7,10-tetraazacyclododecane.

6. A compound of Claim 4 wherein R is propoxy, or pharmaceutically-acceptable salts thereof, and is named as N-(2-pyridylmethyl)-N',N'',N'''-tris(methylenephosphonic acid propyl ester)-1,4,7,10-tetraazacyclododecane.


7. A compound of Claim 1 wherein T equal $-\text{CH}_2-\text{P}(\text{O})\text{ROH}$, where R is C_1-C_5 alkyl.

8. A compound of Claim 1 where one T is




where R¹ and R² are defined as in Claim 1; and the other T terms are defined as in Claim 1.

9. A complex which comprises a polyazamacrocyclic compound of the formula




wherein:

35 T is independently $-\text{CH}_2-\text{COOH}$,




5

10 where: R is OH, C₁-C₅ alkyl or -O-(C₁-C₅ alkyl);R¹ is OH or OCH₃;R² is NO₂, NH₂, isothiocyanato, semicarbazido, thiosemicarbazido, maleimido, bromoacetamido or carboxyl;

with the proviso that only one T moiety is

15



20

where R¹ and R² are defined as before; or


pharmaceutically-acceptable salts thereof;

complexed with a metal ion selected from Gd⁺³, Mn⁺² or Fe⁺³.25 10. A complex of Claim 9 wherein the metal is Gd⁺³.11. A complex of Claim 9 wherein T is -CH₂-P(O)ROH, where R is OH, and the metal is Gd⁺³.12. A complex of Claim 9 wherein T is -CH₂-COOH, and the metal is Gd⁺³.30 13. A complex of Claim 9 wherein T is -CH₂-P(O)ROH, where R is -O-(C₁-C₅ alkyl), and the metal is Gd⁺³.14. A complex of Claim 9 wherein T is -CH₂-P(O)ROH, where R is C₁-C₅ alkyl, and the metal is Gd⁺³.

15. A pharmaceutical formulation comprising a complex of Claim 9 with a pharmaceutically-acceptable carrier


35 16. A method for the diagnosis of a disease state in an animal which comprises administering to said animal an effective amount of the formulation of Claim 15.

17. A process for preparing polyazamacrocyclic compounds of the formula



10 wherein:

T is independently -CH₂-COOH or




where: R is OH, C₁-C₅ alkyl or -O-(C₁-C₅ alkyl); or

pharmaceutically-acceptable salts thereof;

which comprises any one of the following:

20 (A) reacting a compound of the formula




30 with X-CH₂-COOH, where X is chloro or bromo atom, in the presence of an aqueous alkali metal hydroxide, at a pH between about 8-10, at a temperature between about 60-90°C;

to provide a compound of Formula (I) above where T equal -CH₂-COOH; or

35


(B) reacting a compound of the formula



10 with phosphorous acid in hydrochloric acid and excess formaldehyde, at a pH below about 2, at reflux temperature;

to provide a compound of Formula (I) above where T equal $-\text{CH}_2-\text{PO}_3\text{H}_2$; or


(C) reacting a compound of the formula



25 with $\text{P}(\text{OR})_3$ where R is C_1-C_4 alkyl in formaldehyde, followed by hydrolysis using 3 to 12M hydrochloric acid, at a pH below about 3, at reflux temperature;

to provide a compound of Formula (I) above where T equal $-\text{CH}_2-\text{PO}_3\text{H}_2$; or

(D) reacting a compound of the formula




with $\text{P}(\text{OR})_3$ where R is C_1-C_4 alkyl in formaldehyde, followed by hydrolysis using excess alkali metal hydroxide, at a pH above about 12, at reflux temperature;

to provide a compound of Formula (I) above where T equal $-\text{CH}_2-\text{PO}_2\text{HR}$ where R is $-\text{O}-(\text{C}_1-\text{C}_5$ alkyl); or

(E) reacting a compound of the formula

5



(II)

10

with $\text{H}_3\text{C}-\text{P}(\text{OEt})_2$ in formaldehyde and tetrahydrofuran, followed by hydrolysis using excess alkali metal hydroxide, at a pH above about 12, at reflux temperature or with $\text{HP}(\text{O})\text{OH}-\text{C}_2\text{H}_5$ in formaldehyde and hydrochloric acid, at a pH below about 3, at reflux temperature;

15 to provide a compound of Formula (I) above where T equal $-\text{CH}_2-\text{PO}(\text{OH})\text{R}$ where R is C_1-C_5 alkyl.

20

25

30

35

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US94/05009

A. CLASSIFICATION OF SUBJECT MATTER

IPC(5) : Please See Extra Sheet.

US CL : 540/465, 474; 514/79, 340; 424/9

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 540/465, 474; 514/79, 340; 424/9

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Reviewed parent U.S. file

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CAS STN STRUCTURE (in parent U.S. file)

APS TEXT "tetrazacyclododecan?" (parent U.S. file)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO, A, 92/12978 (ROUSSEAU ET AL) 06 AUGUST 1992, see formula I in abstract; page 12, lines 20-25; pages 70-71.	1-16
X	INORGANIC CHEMISTRY, Volume 19, No. 5, issued 1980, J.F. Desreux, "Nuclear Magnetic Resonance Spectroscopy of Lanthanide Complexes with a Tetraacetic Tetraaza Macrocycle. Unusual Conformation Properties," pages 1319-1324, see page 1320, Experimental Section.	17
X	US, A, 3,434,969 (RALSTON) 25 MARCH 1969, see column 3, lines 13-75.	17

 Further documents are listed in the continuation of Box C.
 See patent family annex.

* Special categories of cited documents:	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E" earlier document published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"&" document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means	
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

09 AUGUST 1994

Date of mailing of the international search report

SEP 06 1994

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Authorized officer

PHILIP I. DATLOW

Facsimile No. (703) 305-3230

Telephone No. (703) 308-1235

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US94/05009

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	SYNTHESIS, Volume 12, issued 1987, I.A. Natchev, "Phosphono and Phosphino Analogues and Derivatives of the Natural Aminocarboxylic Acids and Peptides 1. Synthesis and Enzyme-Substrate Interactions of N-Phosphono-, and N-Phosphinomethylated Cyclic Amides," pages 1079-1084, see page 1082, left column, conversion of 4 to 5-8.	17
X,P	US, A, 5,247,077 (PARKER ET AL) 21 SEPTEMBER 1993, see columns 10-11, example 1.	17
X	INORGANIC CHEMISTRY, Volume 30, No. 26, issued 1991, I. Lazar et al, "Synthesis and Complexation Properties of a New Macrocyclic Polyaza Polyphosphinate Ligand, DOTE (1,4,7,10-Tetraazacyclododecane-1,4,7,10-tetrakis(methylene-ethylphosphinate))", pages 5016-5019, see page 5016.	17

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US94/05009

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

- 1. Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

- 2. Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

- 3. Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

- 1. As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
- 2. As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
- 3. As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

- 4. No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- The additional search fees were accompanied by the applicant's protest.
- No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US94/05009

A. CLASSIFICATION OF SUBJECT MATTER:

IPC (5):

A61K 31/44, 31/675; C07D 401/06; C07F 9/30, 9/38, 9/40; G01N 24/08

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

I. Claims 1-16 and 17(a), drawn to macrocyclic compounds, metal complexes thereof, compositions, method of use, and preparation thereof by reacting formula (II) with a haloalkyl acid, Class 540, Subclasses 465 and 475.

II. Claim 17(b), drawn to preparing macrocyclic compounds by reacting formula (II) with a phosphorus acid, Class 540, Subclass 475.

III. Claim 17(c) and 17(d), drawn to preparing macrocyclic compounds by reacting formula (II) with a phosphorus triester, Class 540, Subclass 475.

IV. Claim 17(e), drawn to preparing macrocyclic compounds by reacting formula (II) with a phosphorus diester, Class 540, Subclass 475.

This application lacks unity of invention since multiple distinct processes of preparation are presented. The first recited process in claim 17 is considered to be part of the main invention. All the other processes are considered separate inventions under PCT Rule 13.