



- (51) **International Patent Classification:**  
*C07K 16/18* (2006.01)    *C07K 16/00* (2006.01)  
*G01N 33/15* (2006.01)    *A61K 39/00* (2006.01)
- (21) **International Application Number:**  
PCT/US2017/014461
- (22) **International Filing Date:**  
20 January 2017 (20.01.2017)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**  
62/281,140    20 January 2016 (20.01.2016)    US  
62/350,105    14 June 2016 (14.06.2016)    US  
62/430,852    6 December 2016 (06.12.2016)    US
- (71) **Applicants:** **GENENTECH, INC.** [US/US]; 1 DNA Way, South San Francisco, California 94080 (US). **F. HOFFMANN-LA ROCHE AG** [CH/CH]; Grenzacherstrasse 124, 4070 Basel (CH).
- (72) **Inventors; and**
- (71) **Applicants :** **SMITH, Jillian** [GB/GB]; Hexagon Place, 6 Falcon Way, Shire Park, Welwyn Garden City Hertfordshire AL7 1TW (GB). **SMITH, Janice** [GB/GB]; Hexagon Place, 6 Falcon Way, Shire Park, Welwyn Garden City Hertfordshire AL7 1TW (GB).
- (72) **Inventor:** **KERCHNER, Geoff**; c/o Genentech, Inc., 1 DNA Way, South San Francisco, California 94080 (US).
- (74) **Agents:** **WAIS, Rebecca J.** et al.; c/o Genentech, Inc., 1 DNA Way, South San Francisco, California 94080 (US).
- (81) **Designated States** (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- (84) **Designated States** (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).
- Published:**  
— *with international search report (Art. 21(3))*  
— *with sequence listing part of description (Rule 5.2(a))*



WO 2017/127764 A1

(54) **Title:** HIGH DOSE TREATMENTS FOR ALZHEIMER'S DISEASE

(57) **Abstract:** Methods of treating Alzheimer's Disease (AD) in patients suffering from early AD, including amyloid positive patients, ApoE4 positive patients, and patients suffering from prodromal or mild AD are provided.

**HIGH DOSE TREATMENTS FOR ALZHEIMER'S DISEASE****CROSS REFERENCE TO RELATED APPLICATIONS**

[0001] This application claims priority to U.S. provisional patent application nos. 62/281140, filed 20 January 2016, and 62/350105, filed 14 June 2016, and 62/430852, filed 7 December 2016, which applications are hereby incorporated by reference in their entirety.

**FIELD**

[0002] Methods of treating patients suffering from Alzheimer's Disease using high doses of antibodies that target amyloid  $\beta$  are provided.

**BACKGROUND**

[0003] Alzheimer's Disease (AD) is the most common cause of dementia, affecting an estimated 4.5 million individuals in the United States and 26.6 million worldwide (Hebert *et al.*, Arch. Neurol. 2003; 60:1119-22; Brookmeyer *et al.*, Alzheimers Dement. 2007; 3:186-91). The disease is characterized pathologically by the accumulation of extracellular  $\beta$ -amyloid ("A $\beta$ ") plaques and intracellular neurofibrillary tangles in the brain. Diagnosis is made through the clinical assessment of the neurologic and neuropsychiatric signs and symptoms of AD and the exclusion of other causes of dementia. AD is commonly classified into stages based on cognitive screening examination tests, such as the Mini-Mental State Examination ("MMSE") or other tests. Currently, there are no approved therapies that modify progression of the disease: Approved medical therapies, such as those that inhibit acetylcholinesterase ("AChE") activity or antagonize N-methyl-D-aspartate receptors in the brain, may temporarily improve the symptoms of AD in some patients but do not modify the progression of the disease (Cummings, N. Engl. J. Med. 2004; 351:56-67).

[0004] A number of genetic factors in early- and late-onset familial AD have been documented. The ApoE4 allele is strongly associated with late-onset familial and sporadic AD, with a reported allele frequency of 50%-65% in patients with AD, which is approximately three times that in the general population and for other neurologic disorders (Saunders *et al.*, Neurology 1993; 43:1467-72; Prekumar *et al.*, Am. J. Pathol. 1996; 148:2083-95). In addition to AD, the ApoE4 allele has been implicated in other amyloid-forming disorders, including cerebral amyloid angiopathy ("CAA") (Prekumar *et al.*, Am. J. Pathol. 1996; 148:2083-95). Thus, patients who carry the ApoE4 allele may represent an

etiologically distinct population of patients with AD. Other genetic factors have also been identified.

**[0005]** The deposition of extracellular amyloid plaques in the brain is a hallmark pathologic finding in AD, first reported by Alois Alzheimer in 1906. These amyloid plaques are primarily composed of Abeta peptides (Haass and Selkoe, *Nat. Rev. Mol. Cell Biol.* 2007, 8(2): 101-112) generated by the sequential cleavage of amyloid precursor protein (“APP”) via  $\beta$  and  $\gamma$ -secretase activity. Techniques and tools have been developed to visualize the presence of plaques in patients. For example, position emission tomography (“PET”) scans using imaging agents, such as  $^{18}\text{F}$ -florbetapir, that detect amyloid-beta can be used to detect the presence of amyloid in the brain.

**[0006]** Abeta, particularly in its oligomerized forms, is toxic to neurons and is believed to be causative in AD. Therapies that reduce Abeta levels in the brain may alleviate cognitive dysfunction and block further synaptic loss, axon degeneration, and neuronal cell death. Abeta can be transported actively across the blood-brain barrier (Deane *et al.*, *Stroke* 2004; 35(Suppl D):2628-31). In murine models of AD, systemic delivery of antibodies to Abeta increases Abeta levels in plasma while reducing levels in the central nervous system (CNS) through several proposed mechanisms, including dissolution of brain Abeta plaque, phagocytic removal of opsonized Abeta, and finally via efflux of Abeta from the brain as a result of an equilibrium shift of Abeta resulting from circulating antibodies (Morgan, *Neurodegener. Dis.* 2005; 2:261-6).

**[0007]** Significant failures have marked the development of therapeutic antibodies for the treatment of AD. Large-scale phase three clinical trials of bapineuzumab, an antibody binding specifically to the N-terminal portion of Abeta, were halted when administration of the drug failed to arrest cognitive decline in treated patients (Miles *et al.*, *Scientific Reports* 2013; 3:1-4 Johnston & Johnson press release dated August 6, 2012, entitled “Johnson & Johnson Announces Discontinuation of Phase 3 Development of Bapineuzumab Intravenous (IV) in Mild-To-Moderate Alzheimer’s Disease”). Notably, bapineuzumab did appear to stabilize plaque levels and decreased phosphorylated tau levels in cerebrospinal fluid – suggesting that modification of these biomarkers alone is not necessarily predictive of clinical efficacy (Miles *et al.*, *Scientific Reports* 2013; 3:1-4). Similarly, in phase three clinical trials of solanezumab, an antibody specific for monomeric Abeta that binds in the middle portion of the peptide, the primary cognitive and functional endpoints were not met (Eli Lilly and Company press release dated August 24, 2012, “Eli Lilly and Company Announces Top-Line

Results on Solanezumab Phase 3 Clinical Trials in Patients with Alzheimer's Disease"; Eli Lilly and Company press release dated November 23, 2016, "Lilly Announces Top-Line Results of Solanezumab Phase 3 Clinical Trial", stating that "solanezumab did not meet the primary endpoint in the EXPEDITION3 clinical trial, a phase 3 study of solanezumab in people with mild dementia due to Alzheimer's disease"). Safety concerns have also been raised during the investigation of certain immunotherapies for AD; for example, incidence of amyloid-related imaging abnormalities (ARIA-E and ARIA-H) was over 20% among drug-treated patients in phase two clinical trials of bapineuzumab, an IgG1 isotype antibody (Sperling *et al.*, *The Lancet* 2012; 11:241-249). More recently, an IgG1 isotype anti-Abeta antibody binding to aggregated but not monomeric forms of amyloid beta, aducanumab, was reported to trigger ARIA-E, a form of edema in the brain, in subjects enrolled in a Phase I clinical trial. In a multiple-ascending-dose trial, ARIA-E was detected in an increasing percentage of subjects as the dose was increased and the percentage of subjects with ARIA-E was increased when looking at the subset of subjects carrying an ApoE4 allele, a risk factor for AD. Reportedly, 5% of subjects dosed at 1 and 3 mg/kg of the anti-Abeta antibody showed ARIA-E but 43% and 55% of subjects dosed at 6 mg/kg and 10 mg/kg respectively exhibited ARIA-E. Thus, at increasing doses, the incidence of ARIA-E adverse events also increased. *See* Press Coverage of 2015 Alzheimer's Association International Conference reporting by Gabrielle Strobel, Part 4 of 15, accessible at: [www.alzforum.org/news/conference-coverage/aducanumab-solanezumab-gantenerumab-data-lift-crenezumab-well](http://www.alzforum.org/news/conference-coverage/aducanumab-solanezumab-gantenerumab-data-lift-crenezumab-well) (accessed January 18, 2016). One third of the ARIA-E events led to symptoms in the subjects and some of the patients were reported to have discontinued or had their dose of anti-amyloid antibody reduced.

**[0008]** It is estimated that one in nine people over the age of 65 have AD -- the aggregated yearly costs for health care, long-term care and hospice care by and on behalf of individuals afflicted with AD are over \$200 billion in 2013, and are estimated to rise to \$1.2 trillion by 2050 (by and on behalf of affected individuals) (Alzheimer's Association 2013 Alzheimer's Disease Facts and Figures, Alzheimer's and Dementia 9:2). AD is the sixth-leading cause of death in the United States as of 2013 (*id.*). Current approved therapies treat only some of the symptoms of AD, and not the underlying degeneration. There is a tremendous unmet need for a safe and effective disease-modifying therapeutic for AD.

## SUMMARY

**[0009]** Crenezumab (also known as MABT5102A) is a fully humanized IgG4 monoclonal antibody to Abeta selected for its ability to bind both monomeric and oligomeric forms of Abeta *in vitro*. Crenezumab binds both Abeta1-40 and Abeta 1-42, inhibits Abeta aggregation, and promotes Abeta disaggregation. *See* Adolfsson et al., 2012, J Neurosci 32:9677-9689; *see also*, Ultsch et al., 2016, Sci Rep 6 Article number 35688. Because crenezumab is a human IgG4 backbone antibody, it has reduced Fc $\gamma$  receptor (“Fc $\lambda$ R”) binding affinity compared with human IgG1 or IgG2, which is predictive of reduced immune effector response. These properties, combined with the ability of systemically delivered crenezumab to decrease Abeta CNS levels in a murine model of AD, have suggested that this anti-Abeta therapeutic approach may offer clinical efficacy while having a reduced risk of toxicity, and, in particular, a lower risk of the potentially deleterious side effects, such as ARIA-E or cerebral vasogenic edema or hemorrhages, which have been seen in clinical trials of other Abeta antibody therapies.

**[0010]** The results of pre-clinical and clinical studies in AD patients described herein demonstrate that crenezumab can be administered at high doses without triggering dose-limiting adverse events such as ARIA-E. Furthermore, the effect is seen in patients having a brain amyloid load that is typically seen in patients diagnosed with AD and in patients who are ApoE4 positive, a trait associated with an increased incidence of ARIA-E. This application thus provides methods for treating and monitoring amyloid positive patients diagnosed with early AD, especially prodromal or mild AD, as well as ApoE4 positive patients. In particular, as exemplified herein, it has now been discovered that doses of approximately 2 or more grams of humanized monoclonal anti-amyloid beta antibody with a conformational epitope specific for the middle region of amyloid beta (A $\beta$ ) peptide (*i.e.*, within amino acids 13-24, such as crenezumab) can be administered to amyloid positive patients without an increased incidence of ARIA-E. Accordingly, this application provides high doses of therapeutic agents for modulating the severity of AD and improved methods of using the same, without increased risk of ARIA-E events.

**[0011]** Consequently, the present application provides methods of treating patients suffering from AD and other amyloidoses, comprising administering a humanized monoclonal anti-amyloid beta (A $\beta$  or Abeta) antibody, or antigen-binding fragment thereof, that binds within residues 13 and 24 of amyloid  $\beta$  (1-42)(SEQ ID NO:1) at doses of 2 grams

or more (such as about 50 mg/kg or more). In some embodiments, the antibody, or antigen-binding fragment thereof, is capable of binding fibrillar, oligomeric, and monomeric forms of Abeta. In some embodiments, the antibody binds to oligomeric forms of Abeta with higher affinity than it binds to monomeric forms of Abeta. In some embodiments, the antibody, or antigen-binding fragment thereof, binds to oligomers of Abeta with a 10-fold higher affinity, e.g., with a  $K_D$  of about 0.4 to about 0.6 nM for Abeta oligomers as compared to 3-5 nM for Abeta monomers. In some embodiments, the antibody is an IgG4 antibody. In particular embodiments, the antibody, or antigen-binding fragment thereof, comprises six hypervariable regions (HVRs) wherein HVR-H1 is SEQ ID NO:2, HVR-H2 is SEQ ID NO:3, HVR-H3 is SEQ ID NO:4, HVR-L1 is SEQ ID NO:6, HVR-L2 is SEQ ID NO:7, and HVR-L3 is SEQ ID NO:8. In some embodiments, the antibody comprises a heavy chain variable region having the amino acid sequence of SEQ ID NO:10, or antigen-binding fragment thereof and a light chain variable region, or antigen-binding fragment thereof, having the amino acid sequence of SEQ ID NO:11. In some embodiments, the antibody comprises a heavy chain having the amino acid sequence of SEQ ID NO:5, or antigen-binding fragment thereof, comprising a heavy chain variable region, and a light chain, or antigen-binding fragment thereof, having the amino acid sequence of SEQ ID NO:9, comprising a light chain variable region. In a specific example, the antibody is crenezumab.

**[0012]** The methods of treatment provided herein can be applied to patients suffering from AD or other amyloidosis, as described further herein. Suitable patients are amyloid-positive patients (patients having brain amyloid load consistent with that seen in patients diagnosed with AD) and include subjects suffering from mild cognitive impairment due to AD or having preclinical AD, prodromal AD, early or mild AD, subjects with an MMSE score of 20 or above (e.g., 20-30, 20-26, 24-30, 21-26, 22-26, 22-28, 23-26, 24-26, or 25-26) or with an MMSE score of 22 or above (e.g., 22-30, 23-30, 24-30, 22-26, 22-28, 23-26, 24-26, or 25-26), subjects with a Clinical Dementia Rating-Global Score (CDR-GS) of 0.5 or 1.0, and subjects with a Free and Cued Selective Reminding Test-Immediate Recall (FCSRT-IR) Cueing Index of 0.67 or above and a total free recall score of 27 or greater. In some embodiments, subjects are carriers of at least one ApoE4 allele ("ApoE4 carriers").

**[0013]** In some aspects, the methods provided herein are methods of reducing or slowing decline due to AD in patients suffering from early, mild, or mild to moderate AD. In some embodiments, the decline is one or more of: clinical decline, cognitive decline, and functional decline. In some embodiments, the decline is clinical decline. In some embodiments, the

decline is a decline in cognitive capacity or cognitive decline. In some embodiments, the decline comprises a decline in functional capacity or functional decline. Various tests and scales have been developed to measure cognitive capacity (including memory) and/or function. In various embodiments, one or more test is used to measure clinical, functional, or cognitive decline. A standard measurement of cognitive capacity is the Alzheimer's Disease Assessment Scale Cognitive (ADAS-Cog) test, for example, the 12-item ADAS-Cog or ADAS-Cog12, or the 13-item ADAS-Cog or ADAS-Cog-13. Thus, in some embodiments, the reduction or slowing in decline in cognitive capacity (or cognitive decline) in patients being treated with the antibodies of the invention is determined using the ADAS-Cog12 test. An increase in ADAS-Cog12 score is indicative of worsening in a patient's condition. In some embodiments, the reduction or slowing in cognitive decline (or decline in cognitive capacity) in patients being treated with the antibodies of the invention is determined by a Clinical Dementia Rating Scale / Sum of Boxes (CDR-SB) score. In some embodiments, reduction or slowing in functional decline (or decline in functional ability) in patients being treated with the antibodies of the invention is determined using the Instrumental Activities of Daily Living (iADL) scale. In some embodiments, decline of one or more types is assessed and one or more of the foregoing tests or scales is used to measure reduction or slowing in decline.

**[0014]** An antibody, or antigen-binding fragment thereof, of the invention is administered at a dose that is safe and effective to treat the AD or other amyloidosis, as described herein. Suitable dosages are, as described herein, multi-gram dosages and can range from about 1500 mg to about 24000 mg, or from about 45 mg/kg to about 200 mg/kg. In an exemplary embodiment, the dosage is 45 mg/kg. In a further exemplary embodiment, the dosage is 60 mg/kg. In a further exemplary embodiment, the dosage is 75 mg/kg. In a further exemplary embodiment, the dosage is 90 mg/kg. In a further exemplary embodiment, the dosage is 100 mg/kg. In a further exemplary embodiment, the dosage is 120 mg/kg. In some embodiments, the dosage is between 1500 mg and 24000 mg, such as about 1800 mg, about 2000 mg, about 2200 mg, about 2400 mg, about 2500 mg, about 5000 mg, or more. In the methods provided herein, a variety of dosage regimens are contemplated including dosage regimens in which the antibody is administered repeatedly, *e.g.*, on a weekly or monthly schedule, over an extended period of time, *e.g.*, months to years. In some embodiments, the antibody is administered once every 4 weeks, once every month, once every three weeks, or once every two weeks.

[0015] The humanized monoclonal anti- Abeta antibody of the present disclosure provides a significant benefit compared to other anti-Abeta antibodies in that, when administered in high doses, it does not increase the incidence of adverse events such as ARIA-E and ARIA-H. As shown herein, there was no increase in these adverse events in the treatment arm relative to the placebo arm. Thus, the present disclosure further provides methods of treating patients suffering from early, prodromal, or mild AD by administering high doses of an anti-Abeta antibody.

[0016] The present disclosure further provides pharmaceutical formulations suitable for use in the methods of treatment disclosed herein. The pharmaceutical formulations can be formulated for any convenient route of administration, *e.g.*, parenteral or intravenous injection, and will typically include, in addition to the anti-Abeta of the present disclosure, one or more acceptable carriers, excipients, and/or diluents suited to the desired mode of administration. In some embodiments, an antibody of the invention may be formulated for intravenous administration. In some embodiments, an antibody of the invention may be formulated in an arginine buffer, *e.g.*, an arginine succinate buffer. The buffer can contain one or more surfactants, *e.g.*, a polysorbate. In certain embodiments, the buffer concentration is 50 mM or greater. In some embodiments, the pH is between 4.5 and 7.0, *e.g.*, pH 5.5. Further embodiments are described herein. The pharmaceutical formulations can be package in unit dosage forms for ease of use.

[0017] Treatment with anti-Abeta antibodies for treatment of AD or other amyloidosis, as described herein, can be combined with other therapy, including one or more anti-Abeta antibodies other than crenezumab, or one or more therapeutic agents targeting Tau, for example an anti-Tau antibody. Non-limiting examples of other therapy include neurological drugs, corticosteroids, antibiotics, and antiviral agents. Non-limiting examples of anti-Abeta antibodies other than crenezumab include solanezumab, bapineuzumab, and aducanumab.

#### BRIEF DESCRIPTION OF THE DRAWINGS

[0018] FIG 1 provides the amino acid sequence of Abeta(1-42) (SEQ ID NO:1) with amino acids 13 to 24 underlined.

[0019] FIG 2 provides the amino acid sequence of three heavy chain hypervariable regions (HVR-H1, HVR-H2, and HVR-H3, respectively) and the amino acid sequence of three light chain regions (HVR-L1, HVR-L2, HVR-L3, respectively).



[0020] **FIG 3** provides the amino acid sequence of heavy chain (SEQ ID NO:5), comprising the heavy chain variable region spanning amino acids 1 to 112 of SEQ ID NO:5, and light chain (SEQ ID NO:9), comprising the light chain variable region spanning amino acids 1 to 112 of SEQ ID NO:9, of crenezumab. The underlining in SEQ ID NOs:5 and 9 shows the amino acid sequences of the three heavy chain HVR corresponding to SEQ ID NOs:2-4 and the three light chain HVR corresponding to SEQ ID NOs:6-8, respectively.

[0021] **FIG 4A-B** provides two depictions of the clinical study described in Example 1. **FIG 4A** shows the dosing schedule and assessment schedule, route of administration, and the number of participants in placebo versus treatment arms. **FIG 4B** shows the dose escalation scheme.

[0022] **FIG 5** provides a graph of mean serum concentration of crenezumab measured at three different doses (30 mg/kg, solid line; 45 mg/kg, dotted line; and 60 mg/kg, dashed line).

[0023] **FIG 6A-B** provides graphs of mean serum area under the curve ( $AUC_{INF}$ ) and mean peak or maximum serum concentration ( $C_{PEAK}$ ). **FIG 6A** shows mean  $AUC_{INF}$  at three doses of crenezumab. **FIG 6B** shows mean  $C_{PEAK}$  at three doses of crenezumab. Number of data points included in analysis is shown as “n” for each of the doses.

### DETAILED DESCRIPTION

[0024] Unless defined otherwise, technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Singleton *et al. et al.*, Dictionary of Microbiology and Molecular Biology 2nd ed., J. Wiley & Sons (New York, N.Y. 1994), and March, Advanced Organic Chemistry Reactions, Mechanisms and Structure 4th ed., John Wiley & Sons (New York, N.Y. 1992), provide one skilled in the art with a general guide to many of the terms used in the present application.

#### Certain Definitions and Abbreviations

[0025] For purposes of interpreting this specification, the following definitions will apply and whenever appropriate, terms used in the singular will also include the plural and vice versa. In the event that any definition set forth below conflicts with any document incorporated herein by reference, the definition set forth below shall control.

[0026] As used in this specification and the appended claims, the singular forms “a,” “an” and “the” include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to “a protein” or an “antibody” includes a plurality of proteins or antibodies, respectively; reference to “a cell” includes mixtures of cells, and the like.

**[0027]** Ranges provided in the specification and appended claims include both end points and all points between the end points. Thus, for example, a range of 2.0 to 3.0 includes 2.0, 3.0, and all points between 2.0 and 3.0.

**[0028]** The phrase “substantially similar,” or “substantially the same,” as used herein, denotes a sufficiently high degree of similarity between two numeric values (generally one associated with an antibody of the invention and the other associated with a reference/comparator antibody) such that one of skill in the art would consider the difference between the two values to be of little or no biological and/or statistical significance within the context of the biological characteristic measured by said values (*e.g.*, K<sub>d</sub> values). The difference between said two values is less than about 50%, less than about 40%, less than about 30%, less than about 20%, less than about 10% as a function of the value for the reference/comparator antibody.

**[0029]** The term “sample,” or “test sample” as used herein, refers to a composition that is obtained or derived from a subject of interest that contains a cellular and/or other molecular entity that is to be characterized and/or identified, for example based on physical, biochemical, chemical and/or physiological characteristics. In one embodiment, the definition encompasses blood and other liquid samples of biological origin and tissue samples such as a biopsy specimen or tissue cultures or cells derived therefrom. The source of the tissue sample may be solid tissue as from a fresh, frozen and/or preserved organ or tissue sample or biopsy or aspirate; blood or any blood constituents; bodily fluids; and cells from any time in gestation or development of the subject or plasma. The term “biological sample” as used herein includes, but is not limited to, blood, serum, plasma, sputum, tissue biopsies (*e.g.*, lung samples), and nasal samples including nasal swabs or nasal polyps.

**[0030]** The term “sample,” “biological sample,” or “test sample” includes biological samples that have been manipulated in any way after their procurement, such as by treatment with reagents, solubilization, or enrichment for certain components, such as proteins or polynucleotides, or embedding in a semi-solid or solid matrix for sectioning purposes. For the purposes herein a “section” of a tissue sample is meant a single part or piece of a tissue sample, *e.g.* a thin slice of tissue or cells cut from a tissue sample. Samples include, but are not limited to, whole blood, blood-derived cells, serum, plasma, lymph fluid, synovial fluid, cellular extracts, and combinations thereof. In one embodiment, the sample is a clinical sample. In another embodiment, the sample is used in a diagnostic assay.

**[0031]** In one embodiment, a sample is obtained from a subject or patient prior to treatment with an anti-Abeta antibody. In another embodiment, a sample is obtained from a subject or patient following at least one treatment with an anti-Abeta antibody.

**[0032]** A “reference sample,” as used herein, refers to any sample, standard, or level that is used for comparison purposes. In one embodiment, a reference sample is obtained from a healthy and/or non-diseased part of the body (e.g., tissue or cells) of the same subject or patient. In another embodiment, a reference sample is obtained from an untreated tissue and/or cell of the body of the same subject or patient. In yet another embodiment, a reference sample is obtained from a healthy and/or non-diseased part of the body (e.g., tissues or cells) of an individual who is not the subject or patient. In even another embodiment, a reference sample is obtained from an untreated tissue and/or cell part of the body of an individual who is not the subject or patient.

**[0033]** In certain embodiments, a reference sample is a single sample or combined multiple samples from the same subject or patient that are obtained at one or more different time points than when the test sample is obtained. For example, a reference sample is obtained at an earlier time point from the same subject or patient than when the test sample is obtained. In certain embodiments, a reference sample includes all types of biological samples as defined above under the term “sample” that is obtained from one or more individuals who is not the subject or patient. In certain embodiments, a reference sample is obtained from one or more individuals with amyloidosis, e.g., Alzheimer’s Disease, who is not the subject or patient.

**[0034]** In certain embodiments, a reference sample is a combined multiple samples from one or more healthy individuals who are not the subject or patient. In certain embodiments, a reference sample is a combined multiple samples from one or more individuals with a disease or disorder (e.g., amyloidosis such as, for example, Alzheimer’s Disease) who are not the subject or patient. In certain embodiments, a reference sample is pooled RNA samples from normal tissues or pooled plasma or serum samples from one or more individuals who are not the subject or patient.

**[0035]** The term “small molecule” refers to an organic molecule having a molecular weight between 50 Daltons to 2500 Daltons.

**[0036]** The terms “antibody” and “immunoglobulin” (“Ig”) are used interchangeably in the broadest sense and include, but are not limited to, monoclonal antibodies (for example, full length or intact monoclonal antibodies), polyclonal antibodies, multivalent antibodies,

antibodies with polyepitopic specificity, single chain antibodies, multi-specific antibodies (for example, bispecific antibodies, trispecific antibodies, tetraspecific antibodies), and fragments of antibodies, provided they exhibit the desired biological activity. Such antibodies can be chimeric, humanized, human, synthetic, and/or affinity matured. Such antibodies and methods of generating them are described in more detail herein.

**[0037]** “Antibody fragments” comprise only a portion of an intact antibody, wherein the portion preferably retains at least one, and typically most or all, of the functions normally associated with that portion when present in an intact antibody. In one embodiment, an antibody fragment comprises an antigen binding site of the intact antibody and thus retains the ability to bind antigen. In another embodiment, an antibody fragment, for example one that comprises the Fc region, retains at least one of the biological functions normally associated with the Fc region when present in an intact antibody, such as FcRn binding, antibody half life modulation, ADCC function and complement binding. In one embodiment, an antibody fragment is a monovalent antibody that has an *in vivo* half life substantially similar to an intact antibody. For example, such an antibody fragment may comprise an antigen binding arm linked to an Fc sequence capable of conferring *in vivo* stability to the fragment. Examples of antibody fragments include but are not limited to Fv, Fab, Fab', Fab'-SH, F(ab')<sub>2</sub>; diabodies; linear antibodies; single-chain antibody molecules (*e.g.* scFv); and multispecific antibodies formed from antibody fragments.

**[0038]** The term “target,” as used herein, refers to any native molecule from any vertebrate source, including mammals such as primates (*e.g.* humans) and rodents (*e.g.*, mice and rats), unless otherwise indicated. The term encompasses “full-length,” unprocessed target as well as any form of target that results from processing in the cell. The term also encompasses naturally occurring variants of targets, *e.g.*, splice variants or allelic variants.

**[0039]** The terms “amyloid beta,” “beta-amyloid,” “Abeta,” “amyloid $\beta$ ,” and “A $\beta$ ,” used interchangeably herein, refer to the fragment of amyloid precursor protein (“APP”) that is produced upon  $\beta$ -secretase 1 (“BACE1”) cleavage of APP, as well as modifications, fragments and any functional equivalents thereof, including, but not limited to, A $\beta$ 1-40, and A $\beta$ 1-42. A $\beta$  is known to exist in monomeric form, as well as to associate to form oligomers and fibril structures, which may be found as constituent members of amyloid plaque. The structure and sequences of such A $\beta$  peptides are well known to one of ordinary skill in the art and methods of producing said peptides or of extracting them from brain and other tissues are described, for example, in Glenner and Wong, *Biochem Biophys Res. Comm.* 129: 885-890

(1984). Moreover, A $\beta$  peptides are also commercially available in various forms. An exemplary amino acid sequence of human A $\beta$ 1-42 is DAEFRHDSGYEVHHQKLVFFAEDVGSNKGAIIGLMVGGVVIA (SEQ ID NO: 1).

**[0040]** The terms “anti-target antibody” and “an antibody that binds to target” refer to an antibody that is capable of binding the target with sufficient affinity such that the antibody is useful as a diagnostic and/or therapeutic agent in targeting the target. In one embodiment, the extent of binding of an anti-target antibody to an unrelated, non-target protein is less than about 10% of the binding of the antibody to target as measured, *e.g.*, by a radioimmunoassay (RIA) or biacore assay. In certain embodiments, an antibody that binds to a target has a dissociation constant (K<sub>d</sub>) of  $\leq 1\mu\text{M}$ ,  $\leq 100\text{ nM}$ ,  $\leq 10\text{ nM}$ ,  $\leq 1\text{ nM}$ ,  $\leq 0.1\text{ nM}$ ,  $\leq 0.01\text{ nM}$ , or  $\leq 0.001\text{ nM}$  (*e.g.*,  $10^{-8}\text{ M}$  or less, *e.g.*, from  $10^{-8}\text{ M}$  to  $10^{-13}\text{ M}$ , *e.g.*, from  $10^{-9}\text{ M}$  to  $10^{-13}\text{ M}$ ). In certain embodiments, an anti-target antibody binds to an epitope of a target that is conserved among different species.

**[0041]** “Anti-Abeta immunoglobulin,” “anti-Abeta antibody,” and “antibody that binds Abeta” are used interchangeably herein, and refer to an antibody that specifically binds to human Abeta. A nonlimiting example of an anti-Abeta antibody is crenezumab. Other non-limiting examples of anti-Abeta antibodies are solanezumab, bapineuzumab, aducanumab, and BAN2401.

**[0042]** The terms “crenezumab” and “MABT5102A” are used interchangeably herein, and refer to a specific anti-Abeta antibody that binds to monomeric, oligomeric, and fibril forms of Abeta, and which is associated with CAS registry number 1095207. In one embodiment, such antibody comprises HVR region sequences set forth in **FIG 2**. In another such embodiment, such antibody comprises: (1) an HVR-H1 comprising the amino acid sequence SEQ ID NO: 2; (2) an HVR-H2 sequence comprising the amino acid sequence SEQ ID NO: 3; (3) an HVR-H3 sequence comprising the amino acid sequence SEQ ID NO: 4; (4) an HVR-L1 sequence comprising the amino acid sequence SEQ ID NO: 6; (5) an HVR-L2 sequence comprising the amino acid sequence SEQ ID NO: 7; and (6) an HVR-L3 sequence comprising the amino acid sequence SEQ ID NO: 8. In another embodiment, the specific anti-Abeta antibody comprises heavy chain and light chain sequences, comprising VH and VL domains respectively, having the amino acid sequences set forth in **FIG 3**. In another such embodiment, such specific anti-Abeta antibody comprises a heavy chain comprising the amino acid sequence SEQ ID NO: 5 and a light chain comprising the amino acid sequence SEQ ID NO: 9. In another such embodiment, such specific anti-Abeta antibody comprises a

VH domain comprising the amino acid sequence SEQ ID NO: 10 and a VL domain comprising the amino acid sequence SEQ ID NO: 11. In another embodiment, the antibody is an IgG4 antibody. In another such embodiment, the IgG4 antibody comprises a mutation in its constant domain such that serine 228 is instead a proline.

**[0043]** The term “amyloidosis,” as used herein, refers to a group of diseases and disorders caused by or associated with amyloid or amyloid-like proteins and includes, but is not limited to, diseases and disorders caused by the presence or activity of amyloid-like proteins in monomeric, fibril, or polymeric state, or any combination of the three, including by amyloid plaques. Such diseases include, but are not limited to, secondary amyloidosis and age-related amyloidosis, such as diseases including, but not limited to, neurological disorders such as Alzheimer’s Disease (“AD”), diseases or conditions characterized by a loss of cognitive memory capacity such as, for example, mild cognitive impairment (MCI), Lewy body dementia, Down’s syndrome, hereditary cerebral hemorrhage with amyloidosis (Dutch type), the Guam Parkinson-Demential complex and other diseases which are based on or associated with amyloid-like proteins such as progressive supranuclear palsy, multiple sclerosis, Creutzfeldt Jacob disease, Parkinson’s disease, HIV-related dementia, ALS (amyotrophic lateral sclerosis), inclusion-body myositis (IBM), adult onset diabetes, endocrine tumor and senile cardiac amyloidosis, and various eye diseases including macular degeneration, drusen-related optic neuropathy, glaucoma, and cataract due to beta-amyloid deposition.

**[0044]** Glaucoma is a group of diseases of the optic nerve involving loss of retinal ganglion cells (RGCs) in a characteristic pattern of optic neuropathy. RGCs are the nerve cells that transmit visual signals from the eye to the brain. Caspase-3 and Caspase-8, two major enzymes in the apoptotic process, are activated in the process leading to apoptosis of RGCs. Caspase-3 cleaves amyloid precursor protein (APP) to produce neurotoxic fragments, including Abeta. Without the protective effect of APP, Abeta accumulation in the retinal ganglion cell layer results in the death of RGCs and irreversible loss of vision.

**[0045]** Glaucoma is often, but not always, accompanied by an increased eye pressure, which may be a result of blockage of the circulation of aqueous, or its drainage. Although raised intraocular pressure is a significant risk factor for developing glaucoma, no threshold of intraocular pressure can be defined which would be determinative for causing glaucoma. The damage may also be caused by poor blood supply to the vital optic nerve fibers, a weakness in the structure of the nerve, and/or a problem in the health of the nerve fibers

themselves. Untreated glaucoma leads to permanent damage of the optic nerve and resultant visual field loss, which can progress to blindness.

**[0046]** The different types of glaucomas are classified as open-angle glaucomas, if the condition is chronic, or closed-angle glaucomas, if acute glaucoma occurs suddenly.

Glaucoma usually affects both eyes, but the disease can progress more rapidly in one eye than in the other.

**[0047]** Chronic open-angle glaucoma (COAG), also known as primary open angle glaucoma (POAG), is the most common type of glaucoma. COAG is caused by microscopic blockage in the trabecular meshwork, which decreases the drainage of the aqueous outflow into the Schlemm's canal and raises the intraocular pressure (IOP). POAG usually affects both eyes and is strongly associated with age and a positive family history. Its frequency increases in elderly people as the eye drainage mechanism may gradually become clogged with aging. The increase in intraocular pressure in subjects affected by chronic open-angle glaucoma is not accompanied by any symptoms until the loss is felt on the central visual area.

**[0048]** Acute Angle Closure Glaucoma (AACG) or closed-angle glaucoma is a relatively rare type of glaucoma characterized by a sudden increase in intraocular pressure to 35 to 80 mmHg, leading to severe pain and irreversible loss of vision.. The sudden pressure increase is caused by the closing of the filtering angle and blockage of the drainage channels. Individuals with narrow angles have an increased risk for a sudden closure of the angle. AACG usually occurs monocularly, but the risk exists in both eyes. Age, cataract and pseudoexfoliation are also risk factors since they are associated with enlargement of the lens and crowding or narrowing of the angle. A sudden glaucoma attack may be associated with severe eye pain and headache, inflamed eye, nausea, vomiting, and blurry vision.

**[0049]** Mixed or Combined Mechanism Glaucoma is a mixture or combination of open and closed angle glaucoma. It affects patients with acute ACG whose angle opens after laser iridotomy, but who continue to require medications for IOP control, as well as patients with POAG or pseudoexfoliative glaucoma who gradually develop narrowing of the angle.

**[0050]** Normal tension glaucoma (NTG), also known as low tension glaucoma (LTG), is characterized by progressive optic nerve damage and loss of peripheral vision similar to that seen in other types of glaucoma; however, the intraocular pressure is the normal range or even below normal.

**[0051]** Congenital (infantile) glaucoma is a relatively rare, inherited type of open-angle glaucoma. Insufficient development of the drainage area results in increased pressure in the

eye that can lead to the loss of vision from optic nerve damage and to an enlarged eye. Early diagnosis and treatment are critical to preserve vision in infants and children affected by the disease.

**[0052]** Secondary glaucoma may result from an ocular injury, inflammation in the iris of the eye (iritis), diabetes, cataract, or use of steroids in steroid-susceptible individuals. Secondary glaucoma may also be associated with retinal detachment or retinal vein occlusion or blockage.

**[0053]** Pigmentary glaucoma is characterized by the detachment of granules of pigment from the iris. The granules cause blockage of the drainage system of the eye, leading to elevated intraocular pressure and damage to the optic nerve. Exfoliative glaucoma (pseudoexfoliation) is characterized by deposits of flaky material on the anterior capsule and in the angle of the eye. Accumulation of the flaky material blocks the drainage system and raises the eye pressure.

**[0054]** Diagnosis of glaucoma may be made using various tests. Tonometry determines the pressure in the eye by measuring the tone or firmness of its surface. Several types of tonometers are available for this test, the most common being the applanation tonometer. Pachymetry determines the thickness of the cornea which, in turn, measures intraocular pressure. Gonioscopy allows examination of the filtering angle and drainage area of the eye. Gonioscopy can also determine if abnormal blood vessels may be blocking the drainage of the aqueous fluid out of the eye. Ophthalmoscopy allows examination of the optic nerve and can detect nerve fiber layer drop or changes in the optic disc, or indentation (cupping) of this structure, which may be caused by increased intraocular pressure or axonal drop out. Gonioscopy is also useful in assessing damage to the nerve from poor blood flow or increased intraocular pressure. Visual Field testing maps the field of vision, subjectively, which may detect signs of glaucomatous damage to the optic nerve. This is represented by specific patterns of visual field loss. Ocular coherence tomography, an objective measure of nerve fiber layer loss, is carried out by looking at the thickness of the optic nerve fiber layer (altered in glaucoma) via a differential in light transmission through damaged axonal tissue.

**[0055]** An “antibody that binds to the same epitope” as a reference antibody refers to an antibody that blocks binding of the reference antibody to its antigen in a competition assay by 50% or more, and conversely, the reference antibody blocks binding of the antibody to its antigen in a competition assay by 50% or more. An exemplary competition assay is provided herein.



**[0056]** The term “therapeutic agent” refers to any agent that is used to treat a disease, including but not limited to an agent that treats a symptom of the disease.

**[0057]** As used herein, “treatment” (and grammatical variations thereof such as “treat” or “treating”) refers to clinical intervention in an attempt to alter the natural course of the individual being treated, and can be performed during the course of clinical pathology.

Desirable effects of treatment include, but are not limited to, alleviation or amelioration of one or more symptoms, diminishment of or delay in the appearance of or worsening of any direct or indirect pathological consequences of the disease, decrease of the rate of disease progression, and amelioration or palliation of the disease state. In some embodiments, antibodies are used to delay development of a disease or to slow the progression of a disease.

**[0058]** The term “treatment emergent” as used herein refers to an event that occurs after a first dose of a therapeutic agent is administered. For example, a “treatment emergent adverse event” is an event that is identified upon or after the first dose of a treatment in a clinical study.

**[0059]** “Treatment regimen” refers to a combination of dosage, frequency of administration, or duration of treatment, with or without addition of a second medication.

**[0060]** “Effective treatment regimen” refers to a treatment regimen that will offer beneficial response to a patient receiving the treatment.

**[0061]** “Modifying a treatment” refers to changing the treatment regimen including, changing dosage, frequency of administration, or duration of treatment, and/or addition of a second medication.

**[0062]** An “effective amount” or “effective dose” of an agent refers to an amount or dose effective, for periods of time necessary, to achieve the desired result. For example, a “therapeutically effective amount” is an amount effective, for periods of time necessary, to treat the indicated disease, condition, clinical pathology, or symptom, *i.e.*, to modify the course of progression of AD and/or to alleviate and/or prevent one or more symptoms of AD.

**[0063]** “Affinity” or “binding affinity” refers to the strength of the sum total of noncovalent interactions between a single binding site of a molecule (*e.g.*, an antibody) and its binding partner (*e.g.*, an antigen). Unless indicated otherwise, as used herein, “binding affinity” refers to intrinsic binding affinity which reflects a 1:1 interaction between members of a binding pair (*e.g.*, antibody and antigen binding arm). The affinity of a molecule X for its partner Y can generally be represented by the dissociation constant (K<sub>d</sub>). Affinity can be measured by common methods known in the art, including those described herein, any of

which can be used for purposes of the present invention. Specific illustrative and exemplary embodiments for measuring binding affinity are described herein.

**[0064]** An “affinity matured” antibody refers to an antibody with one or more alterations in one or more hypervariable regions (HVRs), compared to a parent antibody which does not possess such alterations, such alterations resulting in an improvement in the affinity of the antibody for antigen.

**[0065]** As used herein, the term “patient” refers to any single subject for which treatment is desired. In certain embodiments, the patient herein is a human.

**[0066]** A “subject” herein is typically a human. In certain embodiments, a subject is a non-human mammal. Exemplary non-human mammals include laboratory, domestic, pet, sport, and stock animals, *e.g.*, mice, cats, dogs, horses, and cows. Typically, the subject is eligible for treatment, *e.g.*, displays one or more indicia of disease. Generally, such subject or patient is eligible for treatment for amyloidosis, *e.g.*, AD. In one embodiment, such eligible subject or patient is one that is experiencing or has experienced one or more signs, symptoms, or other indicators of AD or has been diagnosed with AD, whether, for example, newly diagnosed, previously diagnosed or at risk for developing AD. Diagnosis of AD may be made based on clinical history, clinical examination, and established imaging modalities. A “patient” or “subject” herein includes any single human subject eligible for treatment who is experiencing or has experienced one or more signs, symptoms, or other indicators of AD. Intended to be included as a subject are any subjects involved in clinical research trials, or subjects involved in epidemiological studies, or subjects once used as controls. The subject may have been previously treated with an anti-Abeta antibody, or antigen-binding fragment thereof, or another drug, or not so treated. The subject may be naïve to an additional drug(s) being used when the treatment herein is started, *i.e.*, the subject may not have been previously treated with, for example, a therapy other than anti-Abeta at “baseline” (*i.e.*, at a set point in time before the administration of a first dose of anti-Abeta in the treatment method herein, such as the day of screening the subject before treatment is commenced). Such “naïve” subjects are generally considered to be candidates for treatment with such additional drug(s).

**[0067]** As used herein, “lifetime” of a subject refers to the remainder of the life of the subject after starting treatment.

**[0068]** The term “monoclonal antibody” as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, *i.e.*, the individual antibodies comprising the population are identical except for possible naturally occurring mutations that

may be present in minor amounts. Monoclonal antibodies are highly specific, being directed against a single antigen. Furthermore, in contrast to polyclonal antibody preparations that typically include different antibodies directed against different determinants (epitopes), each monoclonal antibody is directed against a single determinant on the antigen.

**[0069]** The monoclonal antibodies herein specifically include “chimeric” antibodies in which a portion of the heavy and/or light chain is identical with or homologous to corresponding sequences in antibodies derived from a particular species or belonging to a particular antibody class or subclass, while the remainder of the chain(s) is identical with or homologous to corresponding sequences in antibodies derived from another species or belonging to another antibody class or subclass, as well as fragments of such antibodies, so long as they exhibit the desired biological activity (U.S. Patent No. 4,816,567; and Morrison *et al.*, Proc. Natl. Acad. Sci. USA 81:6851-6855 (1984)).

**[0070]** The “class” of an antibody refers to the type of constant domain or constant region possessed by its heavy chain. There are five major classes of antibodies: IgA, IgD, IgE, IgG, and IgM, and several of these may be further divided into subclasses (or “isotypes”), *e.g.*, IgG1, IgG2, IgG3, IgG4, IgA1, and IgA2. The heavy chain constant domains that correspond to the different classes of immunoglobulins are called  $\alpha$ ,  $\delta$ ,  $\epsilon$ ,  $\gamma$ , and  $\mu$ , respectively.

**[0071]** “Humanized” forms of non-human (*e.g.*, murine) antibodies are chimeric antibodies that contain minimal sequence derived from non-human immunoglobulin. For the most part, humanized antibodies are human immunoglobulins (recipient antibody) in which residues from a hypervariable region of the recipient are replaced by residues from a hypervariable region of a non-human species (donor antibody) such as mouse, rat, rabbit or nonhuman primate having the desired specificity, affinity, and capacity. In some instances, framework region (FR) residues of the human immunoglobulin are replaced by corresponding non-human residues. Furthermore, humanized antibodies may comprise residues that are not found in the recipient antibody or in the donor antibody. These modifications are made to further refine antibody performance. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the hypervariable loops correspond to those of a non-human immunoglobulin and all or substantially all of the FRs are those of a human immunoglobulin sequence. The humanized antibody optionally will also comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin. For further details, see Jones *et al.*, Nature 321:522-525 (1986); Riechmann *et al.*, Nature 332:323-329

(1988); and Presta, *Curr. Op. Struct. Biol.* 2:593-596 (1992). See also the following review articles and references cited therein: Vaswani and Hamilton, *Ann. Allergy, Asthma & Immunol.* 1: 105-115 (1998); Harris, *Biochem. Soc. Transactions* 23:1035-1038 (1995); Hurle and Gross, *Curr. Op. Biotech.* 5:428-433 (1994).

**[0072]** A “human antibody” is one which comprises an amino acid sequence corresponding to that of an antibody produced by a human or a human cell and/or has been derived from a non-human source that utilizes human antibody repertoires or other human antibody-encoding sequences, for example made using any of the techniques for making human antibodies as disclosed herein. Such techniques include, but are not limited to, screening human-derived combinatorial libraries, such as phage display libraries (see, *e.g.*, Marks *et al.*, *J. Mol. Biol.*, 222: 581-597 (1991) and Hoogenboom *et al.*, *Nucl. Acids Res.*, 19: 4133-4137 (1991)); using human myeloma and mouse-human heteromyeloma cell lines for the production of human monoclonal antibodies (see, *e.g.*, Kozbor *J. Immunol.*, 133: 3001 (1984); Brodeur *et al.*, *Monoclonal Antibody Production Techniques and Applications*, pp. 55-93 (Marcel Dekker, Inc., New York, 1987); and Boerner *et al.*, *J. Immunol.*, 147: 86 (1991)); and generating monoclonal antibodies in transgenic animals (*e.g.*, mice) that are capable of producing a full repertoire of human antibodies in the absence of endogenous immunoglobulin production (see, *e.g.*, Jakobovits *et al.*, *Proc. Natl. Acad. Sci USA*, 90: 2551 (1993); Jakobovits *et al.*, *Nature*, 362: 255 (1993); Bruggermann *et al.*, *Year in Immunol.*, 7: 33 (1993)). This definition of a human antibody specifically excludes a humanized antibody comprising antigen-binding residues from a non-human animal.

**[0073]** An “isolated” antibody is one which has been identified and separated and/or recovered from a component of its natural environment. Contaminant components of its natural environment are materials which would interfere with diagnostic or therapeutic uses for the antibody, and may include enzymes, hormones, and other proteinaceous or nonproteinaceous solutes. In some embodiments, an antibody is purified to greater than 95% or 99% purity as determined by, for example, electrophoretic (*e.g.*, SDS-PAGE, isoelectric focusing (IEF), capillary electrophoresis) or chromatographic (*e.g.*, ion exchange or reverse phase HPLC). For review of methods for assessment of antibody purity, see, *e.g.*, Flatman *et al.*, *J. Chromatogr. B* 848:79-87 (2007).

**[0074]** The term “variable region” or “variable domain” refers to the domain of an antibody heavy or light chain that is involved in binding the antibody to antigen. The variable domains of the heavy chain and light chain (VH and VL, respectively) of a native antibody

generally have similar structures, with each domain comprising four conserved framework regions (FRs) and three hypervariable regions (HVRs). (See, *e.g.*, Kindt *et al.* Kuby Immunology, 6th ed., W.H. Freeman and Co., page 91 (2007).) A single VH or VL domain may be sufficient to confer antigen-binding specificity. Furthermore, antibodies that bind a particular antigen may be isolated using a VH or VL domain from an antibody that binds the antigen to screen a library of complementary VL or VH domains, respectively. *See, e.g.*, Portolano *et al.*, J. Immunol. 150:880-887 (1993); Clarkson *et al.*, Nature 352:624-628 (1991).

**[0075]** The term “hypervariable region,” “HVR,” or “HV,” when used herein refers to the regions of an antibody variable domain which are hypervariable in sequence and/or form structurally defined loops. Generally, antibodies comprise six hypervariable regions; three in the VH (H1, H2, H3), and three in the VL (L1, L2, L3). A number of hypervariable region delineations are in use and are encompassed herein. The Kabat Complementarity Determining Regions (CDRs) are based on sequence variability and are the most commonly used (Kabat *et al.*, Sequences of Proteins of Immunological Interest, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, Md. (1991)). Chothia refers instead to the location of the structural loops (Chothia and Lesk J. Mol. Biol. 196:901-917 (1987)). The AbM hypervariable regions represent a compromise between the Kabat CDRs and Chothia structural loops, and are used by Oxford Molecular's AbM antibody modeling software. The “contact” hypervariable regions are based on an analysis of the available complex crystal structures. The residues from each of these HVRs are noted below.

Loop	Kabat	AbM	Chothia	Contact
L1	L24-L34	L24-L34	L26-L32	L30-L36
L2	L50-L56	L50-L56	L50-L52	L46-L55
L3	L89-L97	L89-L97	L91-L96	L89-L96
H1	H31-H35B	H26-H35B	H26-H32	H30-H35B (Kabat Numbering)
H1	H31-H35	H26-H35	H26-H32	H30-H35 (Chothia Numbering)
H2	H50-H65	H50-H58	H53-H55	H47-H58
H3	H95-H102	H95-H102	H96-H101	H93-H101

**[0076]** Hypervariable regions may comprise “extended hypervariable regions” as follows: 24-36 or 24-34 (L1), 46-56 or 49-56 or 50-56 or 52-56 (L2) and 89-97 (L3) in the VL and 26-35 (H1), 50-65 or 49-65 (H2) and 93-102, 94-102 or 95-102 (H3) in the VH. The variable domain residues are numbered according to Kabat *et al.*, *supra* for each of these definitions.

[0077] “Framework” or “FR” residues are those variable domain residues other than the hypervariable region residues as herein defined. The FR of a variable domain generally consists of four FR domains: FR1, FR2, FR3, and FR4. Accordingly, the HVR and FR sequences generally appear in the following sequence in VH (or VL): FR1-H1(L1)-FR2-H2(L2)-FR3-H3(L3)-FR4.

[0078] An “acceptor human framework” for the purposes herein is a framework comprising the amino acid sequence of a light chain variable domain (VL) framework or a heavy chain variable domain (VH) framework derived from a human immunoglobulin framework or a human consensus framework, as defined below. An acceptor human framework “derived from” a human immunoglobulin framework or a human consensus framework may comprise the same amino acid sequence thereof, or it may contain amino acid sequence changes. In some embodiments, the number of amino acid changes are 10 or less, 9 or less, 8 or less, 7 or less, 6 or less, 5 or less, 4 or less, 3 or less, or 2 or less. In some embodiments, the VL acceptor human framework is identical in sequence to the VL human immunoglobulin framework sequence or human consensus framework sequence.

[0079] A “human consensus framework” is a framework which represents the most commonly occurring amino acid residue in a selection of human immunoglobulin VL or VH framework sequences. Generally, the selection of human immunoglobulin VL or VH sequences is from a subgroup of variable domain sequences. Generally, the subgroup of sequences is a subgroup as in Kabat *et al.* Sequences of Proteins of Immunological Interest, Fifth Edition, NIH Publication 91-3242, Bethesda MD (1991), vols. 1-3. *et al. et al.*

[0080] The term “Amyloid-Related Imaging Abnormality – Edema” or “ARIA-E” encompasses cerebral vasogenic edema and sulcal effusion.

[0081] The term “Amyloid-Related Imaging Abnormality – Hemorrhage” or “ARIA-H” encompasses microhemorrhage and superficial siderosis of the central nervous system.

[0082] “Apolipoprotein E4 carrier” or “ApoE4 carrier,” used interchangeably herein with “apolipoprotein E4 positive” or “ApoE4 positive,” refers to an individual having at least one apolipoprotein E4 (or “ApoE4”) allele. An individual with zero ApoE4 alleles is referred to herein as being “ApoE4 negative” or an “ApoE4 non-carrier.” *See also* Prekumar, *et al.*, 1996, *Am. J Pathol.* 148:2083–95.

[0083] The term “cerebral vasogenic edema” refers to an excess accumulation of intravascular fluid or protein in the intracellular or extracellular spaces of the brain. Cerebral vasogenic edema is detectable by, *e.g.*, brain MRI, including, but not limited to FLAIR MRI,

and can be asymptomatic (“asymptomatic vasogenic edema”) or associated with neurological symptoms, such as confusion, dizziness, vomiting, and lethargy (“symptomatic vasogenic edema”) (*see* Sperling *et al.* *Alzheimer’s & Dementia*, 7:367, 2011).

**[0084]** The term “cerebral macrohemorrhage” refers to an intracranial hemorrhage, or bleeding in the brain, of an area that is more than about 1 cm in diameter. Cerebral macrohemorrhage is detectable by, *e.g.*, brain MRI, including but not limited to T2\*-weighted GRE MRI, and can be asymptomatic (“asymptomatic macrohemorrhage”) or associated with symptoms such as transient or permanent focal motor or sensory impairment, ataxia, aphasia, and dysarthria (“symptomatic macrohemorrhage”) (*see, e.g., Chalela JA, Gomes J. Expert Rev. Neurother.* 2004 4:267, 2004 and Sperling *et al.* *Alzheimer’s & Dementia*, 7:367, 2011).

**[0085]** The term “cerebral microhemorrhage” refers to an intracranial hemorrhage, or bleeding in the brain, of an area that is less than about 1 cm in diameter. Cerebral microhemorrhage is detectable by, *e.g.*, brain MRI, including, but not limited to T2\*-weighted GRE MRI, and can be asymptomatic (“asymptomatic microhemorrhage”) or can potentially be associated with symptoms such as transient or permanent focal motor or sensory impairment, ataxia, aphasia, and dysarthria (“symptomatic microhemorrhage”). *See, e.g., Greenberg, et al., 2009, Lancet Neurol.* 8:165–74.

**[0086]** The term “sulcal effusion” refers to an effusion of fluid in the furrows, or sulci, of the brain. Sulcal effusions are detectable by, *e.g.*, brain MRI, including but not limited to FLAIR MRI. *See* Sperling *et al.* *Alzheimer’s & Dementia*, 7:367, 2011.

**[0087]** The term “superficial siderosis of the central nervous system” refers to bleeding or hemorrhage into the subarachnoid space of the brain and is detectable by, *e.g.*, brain MRI, including but not limited to T2\*-weighted GRE MRI. Symptoms indicative of superficial siderosis of the central nervous system include sensorineural deafness, cerebellar ataxia, and pyramidal signs. *See* Kumara-N, *Am J Neuroradiol.* 31:5, 2010.

**[0088]** The term “progression” as used herein refers to the worsening of a disease over time. The “progression rate” or “rate of progression” of a disease refers to how fast or slow a disease develops over time in a patient diagnosed with the disease. The progression rate of a disease can be represented by measurable changes over time of particular characteristics of the disease. A patient carrying particular genetic trait is said to have, or more likely to have, “increased progression rate” if her disease state progresses faster than those patients without such genetic trait. On the other hand, a patient responding to a therapy is said to have, or more likely to have, “decreased progression rate” if her disease progression slows down after

the therapy, when compared to her disease state prior to the treatment or to other patients without the treatment.

**[0089]** “More likely to respond” as used herein refers to patients that are most likely to demonstrate a slowing down or prevention of progression of amyloidosis, e.g., AD. With regard to AD, “more likely to respond” refers to patients that are most likely to demonstrate a reduction in loss of function or cognition with treatment. The phrase “responsive to” in the context of the present invention indicates that a patient suffering from, being suspected to suffer or being prone to suffer from, or diagnosed with a disorder as described herein, shows a response to anti-Abeta treatment.

**[0090]** The phrase “selecting a patient” or “identifying a patient” as used herein refers to using the information or data generated relating to the presence of an allele in a sample of a patient to identify or select the patient as more likely to benefit to benefit from a treatment comprising anti-Abeta antibody. The information or data used or generated may be in any form, written, oral or electronic. In some embodiments, using the information or data generated includes communicating, presenting, reporting, storing, sending, transferring, supplying, transmitting, dispensing, or combinations thereof. In some embodiments, communicating, presenting, reporting, storing, sending, transferring, supplying, transmitting, dispensing, or combinations thereof are performed by a computing device, analyzer unit or combination thereof. In some further embodiments, communicating, presenting, reporting, storing, sending, transferring, supplying, transmitting, dispensing, or combinations thereof are performed by a laboratory or medical professional. In some embodiments, the information or data includes an indication that a specific allele is present or absent in the sample. In some embodiments, the information or data includes an indication that the patient is more likely to respond to a therapy comprising anti-Abeta.

**[0091]** “Effector functions” refer to those biological activities attributable to the Fc region of an antibody, which vary with the antibody isotype. Examples of antibody effector functions include: C1q binding and complement dependent cytotoxicity (CDC); Fc receptor binding; antibody-dependent cell-mediated cytotoxicity (ADCC); phagocytosis; down regulation of cell surface receptors (e.g. B cell receptor); and B cell activation. It is known in the art that wild-type IgG4 antibodies have less effector function than wild-type IgG1 antibodies.

**[0092]** The term “Fc region” herein is used to define a C-terminal region of an immunoglobulin heavy chain that contains at least a portion of the constant region. The term



includes native sequence Fc regions and variant Fc regions. In one embodiment, a human IgG heavy chain Fc region extends from Cys226, or from Pro230, to the carboxyl-terminus of the heavy chain. However, the C-terminal lysine (Lys447) of the Fc region may or may not be present. Unless otherwise specified herein, numbering of amino acid residues in the Fc region or constant region is according to the EU numbering system, also called the EU index, as described in Kabat *et al.*, Sequences of Proteins of Immunological Interest, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD, 1991.

**[0093]** The terms “full length antibody,” “intact antibody,” and “whole antibody” are used herein interchangeably to refer to an antibody having a structure substantially similar to a native antibody structure or having heavy chains that contain an Fc region as defined herein.

**[0094]** The terms “host cell,” “host cell line,” and “host cell culture” are used interchangeably and refer to cells into which exogenous nucleic acid has been introduced, including the progeny of such cells. Host cells include “transformants” and “transformed cells,” which include the primary transformed cell and progeny derived therefrom without regard to the number of passages. Progeny may not be completely identical in nucleic acid content to a parent cell, but may contain mutations. Mutant progeny that have the same function or biological activity as screened or selected for in the originally transformed cell are included herein.

**[0095]** An “immunoconjugate” is an antibody conjugated to one or more heterologous molecule(s), including but not limited to a further therapeutic agent.

**[0096]** An “isolated” nucleic acid refers to a nucleic acid molecule that has been separated from a component of its natural environment. An isolated nucleic acid includes a nucleic acid molecule contained in cells that ordinarily contain the nucleic acid molecule, but the nucleic acid molecule is present extrachromosomally or at a chromosomal location that is different from its natural chromosomal location.

**[0097]** “Isolated nucleic acid encoding an anti-Aβ antibody” refers to one or more nucleic acid molecules encoding antibody heavy and light chains (or fragments thereof), including such nucleic acid molecule(s) in a single vector or separate vectors, and such nucleic acid molecule(s) present at one or more locations in a host cell.

**[0098]** The term “early Alzheimer’s Disease” or “early AD” as used herein (e.g., a “patient diagnosed with early AD” or a “patient suffering from early AD”) includes patients with mild cognitive impairment, such as a memory deficit, due to AD and patients having AD biomarkers, for example amyloid positive patients, as well as patients with prodromal

AD and mild AD. In some embodiments, patients with early AD have an MMSE of 22 or greater and a CDR-GS of 0.5 or 1.0.

**[0099]** A “naked antibody” refers to an antibody that is not conjugated to a heterologous moiety (e.g., a further therapeutic moiety) or radiolabel. The naked antibody may be present in a pharmaceutical formulation.

**[00100]** “Native antibodies” refer to naturally occurring immunoglobulin molecules with varying structures. For example, native IgG antibodies are heterotetrameric glycoproteins of about 150,000 daltons, composed of two identical light chains and two identical heavy chains that are disulfide-bonded. From N- to C-terminus, each heavy chain has a variable region (VH), also called a variable heavy domain or a heavy chain variable domain, followed by three constant domains (CH1, CH2, and CH3). Similarly, from N- to C-terminus, each light chain has a variable region (VL), also called a variable light domain or a light chain variable domain, followed by a constant light (CL) domain. The light chain of an antibody may be assigned to one of two types, called kappa ( $\kappa$ ) and lambda ( $\lambda$ ), based on the amino acid sequence of its constant domain.

**[00101]** The term “package insert” is used to refer to instructions customarily included in commercial packages of therapeutic products, that contain information about the indications, usage, dosage, administration, combination therapy, contraindications and/or warnings concerning the use of such therapeutic products. The term “package insert” is also used to refer to instructions customarily included in commercial packages of diagnostic products that contain information about the intended use, test principle, preparation and handling of reagents, specimen collection and preparation, calibration of the assay and the assay procedure, performance and precision data such as sensitivity and specificity of the assay.

**[00102]** “Percent (%) amino acid sequence identity” with respect to a reference polypeptide sequence is defined as the percentage of amino acid residues in a candidate sequence that are identical with the amino acid residues in the reference polypeptide sequence, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity, and not considering any conservative substitutions as part of the sequence identity. Alignment for purposes of determining percent amino acid sequence identity can be achieved in various ways that are within the skill in the art, for instance, using publicly available computer software such as BLAST, BLAST-2, ALIGN or Megalign (DNASTAR) software. Those skilled in the art can determine appropriate parameters for aligning sequences, including any algorithms needed to achieve maximal

alignment over the full length of the sequences being compared. For purposes herein, however, % amino acid sequence identity values are generated using the sequence comparison computer program ALIGN-2. The ALIGN-2 sequence comparison computer program was authored by Genentech, Inc., and the source code has been filed with user documentation in the U.S. Copyright Office, Washington D.C., 20559, where it is registered under U.S. Copyright Registration No. TXU510087. The ALIGN-2 program is publicly available from Genentech, Inc., South San Francisco, California, or may be compiled from the source code. The ALIGN-2 program should be compiled for use on a UNIX operating system, including digital UNIX V4.0D. All sequence comparison parameters are set by the ALIGN-2 program and do not vary.

**[00103]** In situations where ALIGN-2 is employed for amino acid sequence comparisons, the % amino acid sequence identity of a given amino acid sequence A to, with, or against a given amino acid sequence B (which can alternatively be phrased as a given amino acid sequence A that has or comprises a certain % amino acid sequence identity to, with, or against a given amino acid sequence B) is calculated as follows:

100 times the fraction (X/Y)

**[00104]** where X is the number of amino acid residues scored as identical matches by the sequence alignment program ALIGN-2 in that program's alignment of A and B, and where Y is the total number of amino acid residues in B. It will be appreciated that where the length of amino acid sequence A is not equal to the length of amino acid sequence B, the % amino acid sequence identity of A to B will not equal the % amino acid sequence identity of B to A. Unless specifically stated otherwise, all % amino acid sequence identity values used herein are obtained as described in the immediately preceding paragraph using the ALIGN-2 computer program.

**[00105]** The terms "pharmaceutical formulation" and "pharmaceutical composition" are used interchangeably herein and refer to a preparation which is in such form as to permit the biological activity of an active ingredient contained therein to be effective, and which contains no additional components which are unacceptably toxic to a subject to which the formulation would be administered.

**[00106]** A "pharmaceutically acceptable carrier" refers to an ingredient in a pharmaceutical formulation, other than an active ingredient, which is nontoxic to a subject. A pharmaceutically acceptable carrier includes, but is not limited to, a buffer, excipient, stabilizer, or preservative.

**[00107]** The term “vector,” as used herein, refers to a nucleic acid molecule capable of propagating another nucleic acid to which it is linked. The term includes the vector as a self-replicating nucleic acid structure as well as the vector incorporated into the genome of a host cell into which it has been introduced. Certain vectors are capable of directing the expression of nucleic acids to which they are operatively linked. Such vectors are referred to herein as “expression vectors.”

**[00108]** An “imaging agent” is a compound that has one or more properties that permit its presence and/or location to be detected directly or indirectly. Examples of such imaging agents include proteins and small molecule compounds incorporating a labeled moiety that permits detection.

**[00109]** A “label” is a marker coupled with a molecule to be used for detection or imaging. Examples of such labels include: a radiolabel, a fluorophore, a chromophore, or an affinity tag. In one embodiment, the label is a radiolabel used for medical imaging, for example <sup>99m</sup>Tc or <sup>1123</sup>I, or a spin label for nuclear magnetic resonance (NMR) imaging (also known as magnetic resonance imaging, MRI), such as iodine-123 again, iodine-131, indium-111, fluorine-19, carbon-13, nitrogen-15, oxygen-17, gadolinium, manganese, iron, etc.

#### METHODS AND COMPOSITIONS

**[00110]** The present disclosure provides compositions and methods for the treatment, prognosis, selection and/or identification of patients at risk for or having amyloidosis. In one aspect, the invention is based, in part, on improved methods of treatment.

**[00111]** In certain embodiments, antibodies that bind to Abeta are provided. Antibodies of the invention are useful, e.g., for the diagnosis or treatment of Alzheimer’s Disease (“AD”) and other diseases.

#### Exemplary Antibodies

**[00112]** In one aspect, the invention provides isolated antibodies that bind to Abeta. In certain embodiments, the invention provides an anti-Abeta antibody that can bind to monomeric, oligomeric and fibril forms of human Abeta with good affinity. In one embodiment, the anti-Abeta antibody is an antibody that binds to an epitope of Abeta within residues 13-24 of Abeta. In some embodiments, the anti-Abeta antibody specifically binds to residues 13-24 of Abeta in an extending conformation. While not intending to be bound by any theory of operation, binding Abeta in an extended conformation is thought to account for the ability of exemplary antibodies to bind to different forms of human Abeta, including

monomeric, oligomeric, and fibrillary forms. See Ultsch et al., 2016, supra. In one such embodiment, the antibody is crenezumab.

**[00113]** In one embodiment, the antibody comprises the heavy chain amino acid sequence set forth in SEQ ID NO:5 and the light chain amino acid sequence set forth in SEQ ID NO:9. In another embodiment, the antibody comprises the heavy chain variable region of amino acids 1 to 112 of the amino acid sequence set forth in SEQ ID NO:5 and the light chain variable region of amino acids 1 to 112 of the amino acid sequence set forth in SEQ ID NO:9. In some embodiments, the antibody comprises the heavy chain variable region sequence set forth in SEQ ID NO:10 and the light chain variable region sequence set forth in SEQ ID NO:11. In another embodiment, the antibody comprises the HVR sequences of SEQ ID NO:5 and SEQ ID NO:9. In another embodiment, the antibody comprises HVR sequences that are 95%, 96%, 97%, 98%, or 99% or more identical to the HVR sequences of SEQ ID NO:5 and SEQ ID NO:9.

**[00114]** In any of the above embodiments, an anti-Abeta antibody is humanized. In one embodiment, an anti-Abeta antibody comprises HVRs as in any of the above embodiments, and further comprises an acceptor human framework, e.g. a human immunoglobulin framework or a human consensus framework.

**[00115]** In another aspect, an anti-Abeta antibody comprises a heavy chain variable domain (VH) sequence having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to amino acids 1 to 112 of the amino acid sequence of SEQ ID NO:5. In certain embodiments, a VH sequence having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identity contains substitutions (e.g., conservative substitutions), insertions, or deletions relative to the reference sequence, but an anti-Abeta antibody comprising that sequence retains the ability to bind to Abeta. In certain embodiments, a total of 1 to 10 amino acids have been substituted, inserted and/or deleted in SEQ ID NO:5. In certain embodiments, substitutions, insertions, or deletions occur in regions outside the HVRs (i.e., in the FRs). Optionally, the anti-Abeta antibody comprises the VH sequence in SEQ ID NO:5, including post-translational modifications of that sequence.

**[00116]** In another aspect, an anti-Abeta antibody is provided, wherein the antibody comprises a light chain variable domain (VL) having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to amino acids 1 to 112 of the amino acid sequence of SEQ ID NO:9. In certain embodiments, a VL sequence having at least 90%,

91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identity contains substitutions (e.g., conservative substitutions), insertions, or deletions relative to the reference sequence, but an anti-Abeta antibody comprising that sequence retains the ability to bind to Abeta. In certain embodiments, a total of 1 to 10 amino acids have been substituted, inserted and/or deleted in SEQ ID NO:9. In certain embodiments, the substitutions, insertions, or deletions occur in regions outside the HVRs (i.e., in the FRs). Optionally, the anti-Abeta antibody comprises the VL sequence in SEQ ID NO:9, including post-translational modifications of that sequence.

**[00117]** In another aspect, an anti-Abeta antibody is provided, wherein the antibody comprises a VH as in any of the embodiments provided above, and a VL as in any of the embodiments provided above.

**[00118]** In a further aspect, the invention provides an antibody that binds to the same epitope as an anti-Abeta antibody provided herein. For example, in certain embodiments, an antibody is provided that binds to the same epitope as an anti-Abeta antibody comprising a VH sequence in SEQ ID NO:5 and a VL sequence in SEQ ID NO:9.

**[00119]** In a further aspect of the invention, an anti-Abeta antibody according to any of the above embodiments is a monoclonal antibody, including a chimeric, humanized or human antibody. In one embodiment, an anti-Abeta antibody is an antibody fragment, e.g., a Fv, Fab, Fab', scFv, diabody, or F(ab')<sub>2</sub> fragment. In another embodiment, the antibody is a full length antibody, e.g., an intact IgG4 antibody or other antibody class or isotype as defined herein. In another embodiment, the antibody is a bispecific antibody.

**[00120]** In a further aspect, an anti-Abeta antibody according to any of the above embodiments may incorporate any of the features, singly or in combination, as described in Sections 1-7 below.

**[00121]** In one embodiment, the anti-Abeta antibody comprises a HVR-L1 comprising amino acid sequence SEQ ID NO:6; an HVR-L2 comprising amino acid sequence SEQ ID NO:7; an HVR-L3 comprising amino acid sequence SEQ ID NO: 8; an HVR-H1 comprising amino acid sequence SEQ ID NO:2; an HVR-H2 comprising amino acid sequence SEQ ID NO: 3; and an HVR-H3 comprising amino acid sequence SEQ ID NO: 4.

**[00122]** In another embodiment, the antibody comprises the heavy and light sequences SEQ ID NO:5 and SEQ ID NO:9.

**[00123]** In another embodiment, the antibody comprises the variable region sequences in SEQ ID NO:5 and SEQ ID NO:9.

**[00124]** In another embodiment, the antibody comprises the variable region sequences SEQ ID NO:10 and SEQ ID NO:11.

**[00125]** In any of the above embodiments, an anti-Abeta antibody can be humanized. In one embodiment, an anti-Abeta antibody comprises HVRs as in any of the above embodiments, and further comprises an acceptor human framework, e.g. a human immunoglobulin framework or a human consensus framework.

#### 1. Antibody Affinity

**[00126]** In certain embodiments, an antibody provided herein has a dissociation constant (Kd) of  $\leq 1\mu\text{M}$ ,  $\leq 100\text{ nM}$ ,  $\leq 10\text{ nM}$ ,  $\leq 1\text{ nM}$ ,  $\leq 0.1\text{ nM}$ ,  $\leq 0.01\text{ nM}$ , or  $\leq 0.001\text{ nM}$  (e.g.  $10^{-8}\text{ M}$  or less, e.g. from  $10^{-8}\text{ M}$  to  $10^{-13}\text{ M}$ , e.g., from  $10^{-9}\text{ M}$  to  $10^{-13}\text{ M}$ ).

**[00127]** In one embodiment, Kd is measured by a radiolabeled antigen binding assay (RIA) performed with the Fab version of an antibody of interest and its antigen as described by the following assay. Solution binding affinity of Fabs for antigen is measured by equilibrating Fab with a minimal concentration of (125I)-labeled antigen in the presence of a titration series of unlabeled antigen, then capturing bound antigen with an anti-Fab antibody-coated plate (see, e.g., Chen et al. et al., J. Mol. Biol. 293:865-881(1999)). To establish conditions for the assay, MICROTITER® multi-well plates (Thermo Scientific) are coated overnight with 5  $\mu\text{g/ml}$  of a capturing anti-Fab antibody (Cappel Labs) in 50 mM sodium carbonate (pH 9.6), and subsequently blocked with 2% (w/v) bovine serum albumin in PBS for two to five hours at room temperature (approximately 23°C). In a non-adsorbent plate (Nunc #269620), 100 pM or 26 pM [125I]-antigen are mixed with serial dilutions of a Fab of interest (e.g., consistent with assessment of the anti-VEGF antibody, Fab-12, in Presta et al., Cancer Res. 57:4593-4599 (1997)). The Fab of interest is then incubated overnight; however, the incubation may continue for a longer period (e.g., about 65 hours) to ensure that equilibrium is reached. Thereafter, the mixtures are transferred to the capture plate for incubation at room temperature (e.g., for one hour). The solution is then removed and the plate washed eight times with 0.1% polysorbate 20 (TWEEN-20®) in PBS. When the plates have dried, 150  $\mu\text{l/well}$  of scintillant (MICROSCINT-20 TM; Packard) is added, and the plates are counted on a TOPCOUNT TM gamma counter (Packard) for ten minutes. Concentrations of each Fab that give less than or equal to 20% of maximal binding are chosen for use in competitive binding assays.

**[00128]** According to another embodiment, KD is measured using surface plasmon resonance assays using a BIACORE®-2000 or a BIACORE ®-3000 (BIAcore, Inc.,

Piscataway, NJ) at 25°C with immobilized antigen CM5 chips at ~10 response units (RU). Briefly, carboxymethylated dextran biosensor chips (CM5, BIACORE, Inc.) are activated with N-ethyl-N'- (3-dimethylaminopropyl)-carbodiimide hydrochloride (EDC) and N-hydroxysuccinimide (NHS) according to the supplier's instructions. Antigen is diluted with 10 mM sodium acetate, pH 4.8, to 5 µg/ml (~0.2 µM) before injection at a flow rate of 5 µl/minute to achieve approximately 10 response units (RU) of coupled protein. Following the injection of antigen, 1 M ethanolamine is injected to block unreacted groups. For kinetics measurements, two-fold serial dilutions of Fab (0.78 nM to 500 nM) are injected in PBS with 0.05% polysorbate 20 (TWEEN-20™) surfactant (PBST) at 25°C at a flow rate of approximately 25 µl/min. Association rates (kon) and dissociation rates (koff) are calculated using a simple one-to-one Langmuir binding model (BIACORE® Evaluation Software version 3.2) by simultaneously fitting the association and dissociation sensorgrams. The equilibrium dissociation constant (Kd) is calculated as the ratio koff/kon. See, e.g., Chen et al., *J. Mol. Biol.* 293:865-881 (1999). If the on-rate exceeds 10<sup>6</sup> M<sup>-1</sup> s<sup>-1</sup> by the surface plasmon resonance assay above, then the on-rate can be determined by using a fluorescent quenching technique that measures the increase or decrease in fluorescence emission intensity (excitation = 295 nm; emission = 340 nm, 16 nm band-pass) at 25°C of a 20 nM antigen antibody (Fab form) in PBS, pH 7.2, in the presence of increasing concentrations of antigen as measured in a spectrometer, such as a stop-flow equipped spectrophotometer (Aviv Instruments) or a 8000-series SLM-AMINCO™ spectrophotometer (ThermoSpectronic) with a stirred cuvette.

## 2. Antibody Fragments

**[00129]** In certain embodiments, an antibody provided herein is an antibody fragment. Antibody fragments include, but are not limited to, Fab, Fab', Fab'-SH, F(ab')<sub>2</sub>, Fv, and scFv fragments, and other fragments described below. For a review of certain antibody fragments, see Hudson et al. *Nat. Med.* 9:129-134 (2003). For a review of scFv fragments, see, e.g., Pluckthün, in *The Pharmacology of Monoclonal Antibodies*, vol. 113, Rosenberg and Moore eds., (Springer-Verlag, New York), pp. 269-315 (1994); see also WO 93/16185; and U.S. Patent Nos. 5,571,894 and 5,587,458. For discussion of Fab and F(ab')<sub>2</sub> fragments comprising salvage receptor binding epitope residues and having increased in vivo half-life, see U.S. Patent No. 5,869,046.

**[00130]** Diabodies are antibody fragments with two antigen-binding sites that may be bivalent or bispecific. See, for example, EP 404,097; WO 1993/01161; Hudson et al., *Nat.*



Med. 9:129-134 (2003); and Hollinger et al., Proc. Natl. Acad. Sci. USA 90: 6444-6448 (1993). Triabodies and tetrabodies are also described in Hudson et al., Nat. Med. 9:129-134 (2003).

**[00131]** Single-domain antibodies are antibody fragments comprising all or a portion of the heavy chain variable domain or all or a portion of the light chain variable domain of an antibody. In certain embodiments, a single-domain antibody is a human single-domain antibody (Domantis, Inc., Waltham, MA; see, e.g., U.S. Patent No. 6,248,516 B1). In certain embodiments, two or more single-domain antibodies may be joined together to form an immunoglobulin construct with multivalent affinity (i.e., the N- or C-terminus of a first single-domain antibody may be fused or otherwise joined to the N- or C-terminus of a second single-domain antibody).

**[00132]** Antibody fragments can be made by various techniques, including but not limited to proteolytic digestion of an intact antibody as well as production by recombinant host cells (e.g. E. coli or phage), as described herein.

### 3. Chimeric and Humanized Antibodies

**[00133]** In certain embodiments, an antibody provided herein is a chimeric antibody. Certain chimeric antibodies are described, e.g., in U.S. Patent No. 4,816,567; and Morrison et al., Proc. Natl. Acad. Sci. USA, 81:6851-6855 (1984)). In one example, a chimeric antibody comprises a non-human variable region (e.g., a variable region derived from a mouse, rat, hamster, rabbit, or non-human primate, such as a monkey) and a human constant region. In a further example, a chimeric antibody is a “class switched” antibody in which the class or subclass has been changed from that of the parent antibody. Chimeric antibodies include antigen-binding fragments thereof.

**[00134]** In certain embodiments, a chimeric antibody is a humanized antibody. Typically, a non-human antibody is humanized to reduce immunogenicity to humans, while retaining the specificity and affinity of the parental non-human antibody. Generally, a humanized antibody comprises one or more variable domains in which HVRs, e.g., CDRs, (or portions thereof) are derived from a non-human antibody, and FRs (or portions thereof) are derived from human antibody sequences. A humanized antibody optionally will also comprise at least a portion of a human constant region. In some embodiments, some FR residues in a humanized antibody are substituted with corresponding residues from a non-human antibody (e.g., the antibody from which the HVR residues are derived), e.g., to restore or improve antibody specificity or affinity.

**[00135]** Humanized antibodies and methods of making them are reviewed, e.g., in Almagro and Fransson, *Front. Biosci.* 13:1619-1633 (2008), and are further described, e.g., in Riechmann et al., *Nature* 332:323-329 (1988); Queen et al., *Proc. Nat'l Acad. Sci. USA* 86:10029-10033 (1989); US Patent Nos. 5, 821,337, 7,527,791, 6,982,321, and 7,087,409; Kashmiri et al., *Methods* 36:25-34 (2005) (describing SDR (a-CDR) grafting); Padlan, *Mol. Immunol.* 28:489-498 (1991) (describing "resurfacing"); Dall'Acqua et al., *Methods* 36:43-60 (2005) (describing "FR shuffling"); and Osbourn et al., *Methods* 36:61-68 (2005) and Klimka et al., *Br. J. Cancer*, 83:252-260 (2000) (describing the "guided selection" approach to FR shuffling).

**[00136]** Human framework regions that may be used for humanization include but are not limited to: framework regions selected using the "best-fit" method (see, e.g., Sims et al. *J. Immunol.* 151:2296 (1993)); framework regions derived from the consensus sequence of human antibodies of a particular subgroup of light or heavy chain variable regions (see, e.g., Carter et al. *Proc. Natl. Acad. Sci. USA*, 89:4285 (1992); and Presta et al. *J. Immunol.*, 151:2623 (1993)); human mature (somatically mutated) framework regions or human germline framework regions (see, e.g., Almagro and Fransson, *Front. Biosci.* 13:1619-1633 (2008)); and framework regions derived from screening FR libraries (see, e.g., Baca et al., *J. Biol. Chem.* 272:10678-10684 (1997) and Rosok et al., *J. Biol. Chem.* 271:22611-22618 (1996)).

#### 4. Human Antibodies

**[00137]** In certain embodiments, an antibody provided herein is a human antibody. Human antibodies can be produced using various techniques known in the art. Human antibodies are described generally in van Dijk and van de Winkel, *Curr. Opin. Pharmacol.* 5: 368-74 (2001) and Lonberg, *Curr. Opin. Immunol.* 20:450-459 (2008).

**[00138]** Human antibodies may be prepared by administering an immunogen to a transgenic animal that has been modified to produce intact human antibodies or intact antibodies with human variable regions in response to antigenic challenge. Such animals typically contain all or a portion of the human immunoglobulin loci, which replace the endogenous immunoglobulin loci, or which are present extrachromosomally or integrated randomly into the animal's chromosomes. In such transgenic mice, the endogenous immunoglobulin loci have generally been inactivated. For review of methods for obtaining human antibodies from transgenic animals, see Lonberg, *Nat. Biotech.* 23:1117-1125 (2005). See also, e.g., U.S. Patent Nos. 6,075,181 and 6,150,584 describing XENOMOUSE™

technology; U.S. Patent No. 5,770,429 describing HUMAB® technology; U.S. Patent No. 7,041,870 describing K-M MOUSE® technology, and U.S. Patent Application Publication No. US 2007/0061900, describing VELOCIMOUSE® technology). Human variable regions from intact antibodies generated by such animals may be further modified, e.g., by combining with a different human constant region.

**[00139]** Human antibodies can also be made by hybridoma-based methods. Human myeloma and mouse-human heteromyeloma cell lines for the production of human monoclonal antibodies have been described. (See, e.g., Kozbor *J. Immunol.*, 133: 3001 (1984); Brodeur et al., *Monoclonal Antibody Production Techniques and Applications*, pp. 51-63 (Marcel Dekker, Inc., New York, 1987); and Boerner et al., *J. Immunol.*, 147: 86 (1991).) Human antibodies generated via human B-cell hybridoma technology are also described in Li et al., *Proc. Natl. Acad. Sci. USA*, 103:3557-3562 (2006). Additional methods include those described, for example, in U.S. Patent No. 7,189,826 (describing production of monoclonal human IgM antibodies from hybridoma cell lines) and Ni, Xiandai Mianyixue, 26(4):265-268 (2006) (describing human-human hybridomas). Human hybridoma technology (Trioma technology) is also described in Vollmers and Brandlein, *Histology and Histopathology*, 20(3):927-937 (2005) and Vollmers and Brandlein, *Methods and Findings in Experimental and Clinical Pharmacology*, 27(3):185-91 (2005).

**[00140]** Human antibodies may also be generated by isolating Fv clone variable domain sequences selected from human-derived phage display libraries. Such variable domain sequences may then be combined with a desired human constant domain. Techniques for selecting human antibodies from antibody libraries are described below.

#### 5. Library-Derived Antibodies

**[00141]** Antibodies of the invention may be isolated by screening combinatorial libraries for antibodies with the desired activity or activities. For example, a variety of methods are known in the art for generating phage display libraries and screening such libraries for antibodies possessing the desired binding characteristics. Such methods are reviewed, e.g., in Hoogenboom et al. in *Methods in Molecular Biology* 178:1-37 (O'Brien et al., ed., Human Press, Totowa, NJ, 2001) and further described, e.g., in the McCafferty et al., *Nature* 348:552-554; Clackson et al., *Nature* 352: 624-628 (1991); Marks et al., *J. Mol. Biol.* 222: 581-597 (1992); Marks and Bradbury, in *Methods in Molecular Biology* 248:161-175 (Lo, ed., Human Press, Totowa, NJ, 2003); Sidhu et al., *J. Mol. Biol.* 338(2): 299-310 (2004); Lee

et al., *J. Mol. Biol.* 340(5): 1073-1093 (2004); Fellouse, *Proc. Natl. Acad. Sci. USA* 101(34): 12467-12472 (2004); and Lee et al., *J. Immunol. Methods* 284(1-2): 119-132(2004).

**[00142]** In certain phage display methods, repertoires of VH and VL genes are separately cloned by polymerase chain reaction (PCR) and recombined randomly in phage libraries, which can then be screened for antigen-binding phage as described in Winter et al., *Ann. Rev. Immunol.*, 12: 433-455 (1994). Phage typically display antibody fragments, either as single-chain Fv (scFv) fragments or as Fab fragments. Libraries from immunized sources provide high-affinity antibodies to the immunogen without the requirement of constructing hybridomas. Alternatively, the naive repertoire can be cloned (e.g., from human) to provide a single source of antibodies to a wide range of non-self and also self antigens without any immunization as described by Griffiths et al., *EMBO J.*, 12: 725-734 (1993). Finally, naive libraries can also be made synthetically by cloning unrearranged V-gene segments from stem cells, and using PCR primers containing random sequence to encode the highly variable CDR3 regions and to accomplish rearrangement in vitro, as described by Hoogenboom and Winter, *J. Mol. Biol.*, 227: 381-388 (1992). Patent publications describing human antibody phage libraries include, for example: US Patent No. 5,750,373, and US Patent Publication Nos. 2005/0079574, 2005/0119455, 2005/0266000, 2007/0117126, 2007/0160598, 2007/0237764, 2007/0292936, and 2009/0002360.

**[00143]** Antibodies or antibody fragments isolated from human antibody libraries are considered human antibodies or human antibody fragments herein.

## 6. Multispecific Antibodies

**[00144]** In certain embodiments, an antibody provided herein is a multispecific antibody, e.g. a bispecific antibody. Multispecific antibodies are monoclonal antibodies that have binding specificities for at least two different sites. In certain embodiments, one of the binding specificities is for Aβ and the other is for any other antigen. In certain embodiments, bispecific antibodies may bind to two different epitopes of Aβ. Bispecific antibodies may also be used to localize cytotoxic agents to cells. Bispecific antibodies can be prepared as full length antibodies or antibody fragments.

**[00145]** Techniques for making multispecific antibodies include, but are not limited to, recombinant co-expression of two immunoglobulin heavy chain-light chain pairs having different specificities (see Milstein and Cuellar, *Nature* 305: 537 (1983)), WO 93/08829, and Traunecker et al., *EMBO J.* 10: 3655 (1991)), and “knob-in-hole” engineering (see, e.g., U.S. Patent No. 5,731,168). Multi-specific antibodies may also be made by engineering

electrostatic steering effects for making antibody Fc-heterodimeric molecules (WO 2009/089004A1); cross-linking two or more antibodies or fragments (see, e.g., US Patent No. 4,676,980, and Brennan et al., *Science*, 229: 81 (1985)); using leucine zippers to produce bi-specific antibodies (see, e.g., Kostelny et al., *J. Immunol.*, 148(5):1547-1553 (1992)); using “diabody” technology for making bispecific antibody fragments (see, e.g., Hollinger et al., *Proc. Natl. Acad. Sci. USA*, 90:6444-6448 (1993)); and using single-chain Fv (sFv) dimers (see, e.g. Gruber et al., *J. Immunol.*, 152:5368 (1994)); and preparing trispecific antibodies as described, e.g., in Tutt et al. *J. Immunol.* 147: 60 (1991).

**[00146]** Engineered antibodies with three or more functional antigen binding sites, including “Octopus antibodies,” are also included herein (see, e.g. US 2006/0025576A1).

**[00147]** The antibody or fragment herein also includes a “Dual Acting FAb” or “DAF” comprising an antigen binding site that binds to Abeta as well as another, different antigen (see, US 2008/0069820, for example).

## 7. Antibody Variants

**[00148]** In certain embodiments, amino acid sequence variants of the antibodies provided herein are contemplated. For example, it may be desirable to improve the binding affinity and/or other biological properties of the antibody. Amino acid sequence variants of an antibody may be prepared by introducing appropriate modifications into the nucleotide sequence encoding the antibody, or by peptide synthesis. Such modifications include, for example, deletions from, and/or insertions into and/or substitutions of residues within the amino acid sequences of the antibody. Any combination of deletion, insertion, and substitution can be made to arrive at the final construct, provided that the final construct possesses the desired characteristics, e.g., antigen-binding.

### Substitution, Insertion, and Deletion Variants

**[00149]** In certain embodiments, antibody variants having one or more amino acid substitutions are provided. Sites of interest for substitutional mutagenesis include the HVRs and FRs. Conservative substitutions are shown in Table 1 under the heading of “conservative substitutions.” More substantial changes are provided in Table 1 under the heading of “exemplary substitutions,” and as further described below in reference to amino acid side chain classes. Amino acid substitutions may be introduced into an antibody of interest and the products screened for a desired activity, e.g., retained/improved antigen binding, decreased immunogenicity, or improved ADCC or CDC.

### **TABLE 1**

Original Residue	Exemplary Substitutions	Conservative Substitutions
Ala (A)	Val; Leu; Ile	Val
Arg (R)	Lys; Gln; Asn	Lys
Asn (N)	Gln; His; Asp, Lys; Arg	Gln
Asp (D)	Glu; Asn	Glu
Cys (C)	Ser; Ala	Ser
Gln (Q)	Asn; Glu	Asn
Glu (E)	Asp; Gln	Asp
Gly (G)	Ala	Ala
His (H)	Asn; Gln; Lys; Arg	Arg
Ile (I)	Leu; Val; Met; Ala; Phe; Norleucine	Leu
Leu (L)	Norleucine; Ile; Val; Met; Ala; Phe	Ile
Lys (K)	Arg; Gln; Asn	Arg
Met (M)	Leu; Phe; Ile	Leu
Phe (F)	Trp; Leu; Val; Ile; Ala; Tyr	Tyr
Pro (P)	Ala	Ala
Ser (S)	Thr	Thr
Thr (T)	Val; Ser	Ser
Trp (W)	Tyr; Phe	Tyr
Tyr (Y)	Trp; Phe; Thr; Ser	Phe
Val (V)	Ile; Leu; Met; Phe; Ala; Norleucine	Leu

**[00150]** Amino acids may be grouped according to common side-chain properties:

- (1) hydrophobic: Norleucine, Met, Ala, Val, Leu, Ile;
- (2) neutral hydrophilic: Cys, Ser, Thr, Asn, Gln;
- (3) acidic: Asp, Glu;
- (4) basic: His, Lys, Arg;
- (5) residues that influence chain orientation: Gly, Pro;
- (6) aromatic: Trp, Tyr, Phe.

**[00151]** Non-conservative substitutions will entail exchanging a member of one of these classes for another class.

**[00152]** One type of substitutional variant involves substituting one or more hypervariable region residues of a parent antibody (e.g. a humanized or human antibody). Generally, the

resulting variant(s) selected for further study will have modifications (e.g., improvements) in certain biological properties (e.g., increased affinity, reduced immunogenicity) relative to the parent antibody and/or will have substantially retained certain biological properties of the parent antibody. An exemplary substitutional variant is an affinity matured antibody. In certain embodiments, affinity matured antibodies will have nanomolar or even picomolar affinities for the target antigen. Affinity matured antibodies are produced by procedures known in the art, including, e.g., using phage display-based affinity maturation techniques such as those described herein. Briefly, one or more HVR residues are mutated and the variant antibodies displayed on phage and screened for a particular biological activity (e.g. binding affinity). Other procedures are also known. Marks et al. *Bio/Technology* 10:779-783 (1992) describes affinity maturation by VH and VL domain shuffling. Random mutagenesis of HVR and/or framework residues is described by: Barbas et al. *Proc Nat. Acad. Sci, USA* 91:3809-3813 (1994); Schier et al. *Gene* 169:147-155 (1996); Yelton et al. *J. Immunol.* 155:1994-2004 (1995); Jackson et al., *J. Immunol.* 154(7):3310-9 (1995); and Hawkins et al. *J. Mol. Biol.* 226:889-896 (1992).

**[00153]** Alterations (e.g., substitutions) may be made in HVRs, e.g., to improve antibody affinity. Such alterations may be made in HVR “hotspots,” i.e., residues encoded by codons that undergo mutation at high frequency during the somatic maturation process (see, e.g., Chowdhury, *Methods Mol. Biol.* 207:179-196 (2008)), and/or SDRs (a-CDRs), with the resulting variant VH or VL being tested for binding affinity. Affinity maturation by constructing and reselecting from secondary libraries has been described, e.g., in Hoogenboom et al. in *Methods in Molecular Biology* 178:1-37 (O’Brien et al., ed., Human Press, Totowa, NJ, (2001).) In some embodiments of affinity maturation, diversity is introduced into the variable genes chosen for maturation by any of a variety of methods (e.g., error-prone PCR, chain shuffling, or oligonucleotide-directed mutagenesis). A secondary library is then created. The library is then screened to identify any antibody variants with the desired affinity. Another method to introduce diversity involves HVR-directed approaches, in which several HVR residues (e.g., 4-6 residues at a time) are randomized. HVR residues involved in antigen binding may be specifically identified, e.g., using alanine scanning mutagenesis or modeling. CDR-H3 and CDR-L3 in particular are often targeted.

**[00154]** In certain embodiments, substitutions, insertions, or deletions may occur within one or more HVRs so long as such alterations do not substantially reduce the ability of the antibody to bind antigen. For example, conservative alterations (e.g., conservative

substitutions as provided herein) that do not substantially reduce binding affinity may be made in HVRs. Such alterations may be outside of HVR “hotspots” or SDRs. In certain embodiments of the variant VH and VL sequences provided above, each HVR either is unaltered, or contains no more than one, two or three amino acid substitutions.

**[00155]** A useful method for identification of residues or regions of an antibody that may be targeted for mutagenesis is called “alanine scanning mutagenesis” as described by Cunningham and Wells (1989) *Science*, 244:1081-1085. In this method, a residue or group of target residues (e.g., charged residues such as arg, asp, his, lys, and glu) are identified and replaced by a neutral or negatively charged amino acid (e.g., alanine or polyalanine) to determine whether the interaction of the antibody with antigen is affected. Further substitutions may be introduced at the amino acid locations demonstrating functional sensitivity to the initial substitutions. Alternatively, or additionally, a crystal structure of an antigen-antibody complex to identify contact points between the antibody and antigen. Such contact residues and neighboring residues may be targeted or eliminated as candidates for substitution. Variants may be screened to determine whether they contain the desired properties.

**[00156]** Amino acid sequence insertions include amino- and/or carboxyl-terminal fusions ranging in length from one residue to polypeptides containing a hundred or more residues, as well as intrasequence insertions of single or multiple amino acid residues. Examples of terminal insertions include an antibody with an N-terminal methionyl residue. Other insertional variants of the antibody molecule include the fusion to the N- or C-terminus of the antibody to an enzyme (e.g. for ADEPT) or a polypeptide which increases the serum half-life of the antibody.

#### Glycosylation variants

**[00157]** In certain embodiments, an antibody provided herein is altered to increase or decrease the extent to which the antibody is glycosylated. Addition or deletion of glycosylation sites to an antibody may be conveniently accomplished by altering the amino acid sequence such that one or more glycosylation sites is created or removed.

**[00158]** Where the antibody comprises an Fc region, the carbohydrate attached thereto may be altered. Native antibodies produced by mammalian cells typically comprise a branched, biantennary oligosaccharide that is generally attached by an N-linkage to Asn297 of the CH2 domain of the Fc region. See, e.g., Wright et al. *TIBTECH* 15:26-32 (1997). The oligosaccharide may include various carbohydrates, e.g., mannose, N-acetyl glucosamine



(GlcNAc), galactose, and sialic acid, as well as a fucose attached to a GlcNAc in the “stem” of the biantennary oligosaccharide structure. In some embodiments, modifications of the oligosaccharide in an antibody of the invention may be made in order to create antibody variants with certain improved properties.

**[00159]** In one embodiment, antibody variants are provided having a carbohydrate structure that lacks fucose attached (directly or indirectly) to an Fc region. For example, the amount of fucose in such antibody may be from 1% to 80%, from 1% to 65%, from 5% to 65% or from 20% to 40%. The amount of fucose is determined by calculating the average amount of fucose within the sugar chain at Asn297, relative to the sum of all glycostructures attached to Asn 297 (e. g. complex, hybrid and high mannose structures) as measured by MALDI-TOF mass spectrometry, as described in WO 2008/077546, for example. Asn297 refers to the asparagine residue located at about position 297 in the Fc region (Eu numbering of Fc region residues); however, Asn297 may also be located about  $\pm 3$  amino acids upstream or downstream of position 297, i.e., between positions 294 and 300, due to minor sequence variations in antibodies. Such fucosylation variants may have improved ADCC function. See, e.g., US Patent Publication Nos. US 2003/0157108 (Presta, L.); US 2004/0093621 (Kyowa Hakko Kogyo Co., Ltd). Examples of publications related to “defucosylated” or “fucose-deficient” antibody variants include: US 2003/0157108; WO 2000/61739; WO 2001/29246; US 2003/0115614; US 2002/0164328; US 2004/0093621; US 2004/0132140; US 2004/0110704; US 2004/0110282; US 2004/0109865; WO 2003/085119; WO 2003/084570; WO 2005/035586; WO 2005/035778; WO2005/053742; WO2002/031140; Okazaki et al. J. Mol. Biol. 336:1239-1249 (2004); Yamane-Ohnuki et al. Biotech. Bioeng. 87: 614 (2004). Examples of cell lines capable of producing defucosylated antibodies include Lec13 CHO cells deficient in protein fucosylation (Ripka et al. Arch. Biochem. Biophys. 249:533-545 (1986); US Pat Appl No US 2003/0157108 A1, Presta, L; and WO 2004/056312 A1, Adams et al., especially at Example 11), and knockout cell lines, such as alpha-1,6-fucosyltransferase gene, FUT8, knockout CHO cells (see, e.g., Yamane-Ohnuki et al. Biotech. Bioeng. 87: 614 (2004); Kanda, Y. et al., Biotechnol. Bioeng., 94(4):680-688 (2006); and WO2003/085107).

**[00160]** Antibodies variants are further provided with bisected oligosaccharides, e.g., in which a biantennary oligosaccharide attached to the Fc region of the antibody is bisected by GlcNAc. Such antibody variants may have reduced fucosylation and/or improved ADCC function. Examples of such antibody variants are described, e.g., in WO 2003/011878 (Jean-Mairet et al.); US Patent No. 6,602,684 (Umana et al.); and US 2005/0123546 (Umana et al.).

Antibody variants with at least one galactose residue in the oligosaccharide attached to the Fc region are also provided. Such antibody variants may have improved CDC function. Such antibody variants are described, e.g., in WO 1997/30087 (Patel et al.); WO 1998/58964 (Raju, S.); and WO 1999/22764 (Raju, S.).

#### Fc region variants

**[00161]** In certain embodiments, one or more amino acid modifications may be introduced into the Fc region of an antibody provided herein, thereby generating an Fc region variant. The Fc region variant may comprise a human Fc region sequence (e.g., a human IgG1, IgG2, IgG3 or IgG4 Fc region) comprising an amino acid modification (e.g. a substitution) at one or more amino acid positions.

**[00162]** In certain embodiments, the invention contemplates an antibody variant that possesses some but not all effector functions, which make it a desirable candidate for applications in which the half life of the antibody in vivo is important yet certain effector functions (such as complement and ADCC) are unnecessary or deleterious. In vitro and/or in vivo cytotoxicity assays can be conducted to confirm the reduction/depletion of CDC and/or ADCC activities. For example, Fc receptor (FcR) binding assays can be conducted to ensure that the antibody lacks Fc $\gamma$ R binding (hence likely lacking ADCC activity), but retains FcRn binding ability. The primary cells for mediating ADCC, NK cells, express Fc $\gamma$ RIII only, whereas monocytes express Fc $\gamma$ RI, Fc $\gamma$ RII and Fc $\gamma$ RIII. FcR expression on hematopoietic cells is summarized in Table 3 on page 464 of Ravetch and Kinet, *Annu. Rev. Immunol.* 9:457-492 (1991). Non-limiting examples of in vitro assays to assess ADCC activity of a molecule of interest is described in U.S. Patent No. 5,500,362 (see, e.g. Hellstrom, I. et al. *Proc. Nat'l Acad. Sci. USA* 83:7059-7063 (1986)) and Hellstrom, I et al., *Proc. Nat'l Acad. Sci. USA* 82:1499-1502 (1985); 5,821,337 (see Bruggemann, M. et al., *J. Exp. Med.* 166:1351-1361 (1987)). Alternatively, non-radioactive assays methods may be employed (see, for example, ACTITM non-radioactive cytotoxicity assay for flow cytometry (CellTechnology, Inc. Mountain View, CA; and CytoTox 96® non-radioactive cytotoxicity assay (Promega, Madison, WI). Useful effector cells for such assays include peripheral blood mononuclear cells (PBMC) and Natural Killer (NK) cells. Alternatively, or additionally, ADCC activity of the molecule of interest may be assessed in vivo, e.g., in a animal model such as that disclosed in Clynes et al. *Proc. Nat'l Acad. Sci. USA* 95:652-656 (1998). C1q

binding assays may also be carried out to confirm that the antibody is unable to bind C1q and hence lacks CDC activity. See, e.g., C1q and C3c binding ELISA in WO 2006/029879 and WO 2005/100402. To assess complement activation, a CDC assay may be performed (see, for example, Gazzano-Santoro et al., *J. Immunol. Methods* 202:163 (1996); Cragg, M.S. et al., *Blood* 101:1045-1052 (2003); and Cragg, M.S. and M.J. Glennie, *Blood* 103:2738-2743 (2004)). FcRn binding and in vivo clearance/half life determinations can also be performed using methods known in the art (see, e.g., Petkova, S.B. et al., *Int'l. Immunol.* 18(12):1759-1769 (2006)).

**[00163]** Antibodies with reduced effector function include those with substitution of one or more of Fc region residues 238, 265, 269, 270, 297, 327 and 329 (U.S. Patent No. 6,737,056). Such Fc mutants include Fc mutants with substitutions at two or more of amino acid positions 265, 269, 270, 297 and 327, including the so-called "DANA" Fc mutant with substitution of residues 265 and 297 to alanine (US Patent No. 7,332,581).

**[00164]** Certain antibody variants with improved or diminished binding to FcRs are described. (See, e.g., U.S. Patent No. 6,737,056; WO 2004/056312, and Shields et al., *J. Biol. Chem.* 9(2): 6591-6604 (2001).)

**[00165]** In certain embodiments, an antibody variant comprises an Fc region with one or more amino acid substitutions which improve ADCC, e.g., substitutions at positions 298, 333, and/or 334 of the Fc region (EU numbering of residues).

**[00166]** In some embodiments, alterations are made in the Fc region that result in altered (i.e., either improved or diminished) C1q binding and/or Complement Dependent Cytotoxicity (CDC), e.g., as described in US Patent No. 6,194,551, WO 99/51642, and Idusogie et al. *J. Immunol.* 164: 4178-4184 (2000).

**[00167]** Antibodies with increased half lives and improved binding to the neonatal Fc receptor (FcRn), which is responsible for the transfer of maternal IgGs to the fetus (Guyer et al., *J. Immunol.* 117:587 (1976) and Kim et al., *J. Immunol.* 24:249 (1994)), are described in US2005/0014934A1 (Hinton et al.). Those antibodies comprise an Fc region with one or more substitutions therein which improve binding of the Fc region to FcRn. Such Fc variants include those with substitutions at one or more of Fc region residues: 238, 256, 265, 272, 286, 303, 305, 307, 311, 312, 317, 340, 356, 360, 362, 376, 378, 380, 382, 413, 424 or 434, e.g., substitution of Fc region residue 434 (US Patent No. 7,371,826). See also Duncan & Winter, *Nature* 322:738-40 (1988); U.S. Patent No. 5,648,260; U.S. Patent No. 5,624,821; and WO 94/29351 concerning other examples of Fc region variants.

### Cysteine engineered antibody variants

**[00168]** In certain embodiments, it may be desirable to create cysteine engineered antibodies, e.g., “thioMAbs,” in which one or more residues of an antibody are substituted with cysteine residues. In particular embodiments, the substituted residues occur at accessible sites of the antibody. By substituting those residues with cysteine, reactive thiol groups are thereby positioned at accessible sites of the antibody and may be used to conjugate the antibody to other moieties, such as drug moieties or linker-drug moieties, to create an immunoconjugate, as described further herein. In certain embodiments, any one or more of the following residues may be substituted with cysteine: V205 (Kabat numbering) of the light chain; A118 (EU numbering) of the heavy chain; and S400 (EU numbering) of the heavy chain Fc region. Cysteine engineered antibodies may be generated as described, e.g., in U.S. Patent No. 7,521,541.

### Antibody Derivatives

**[00169]** In certain embodiments, an antibody provided herein may be further modified to contain additional nonproteinaceous moieties that are known in the art and readily available. The moieties suitable for derivatization of the antibody include but are not limited to water soluble polymers. Non-limiting examples of water soluble polymers include, but are not limited to, polyethylene glycol (PEG), copolymers of ethylene glycol/propylene glycol, carboxymethylcellulose, dextran, polyvinyl alcohol, polyvinyl pyrrolidone, poly-1, 3-dioxolane, poly-1,3,6-trioxane, ethylene/maleic anhydride copolymer, polyaminoacids (either homopolymers or random copolymers), and dextran or poly(n-vinyl pyrrolidone)polyethylene glycol, propylene glycol homopolymers, polypropylene oxide/ethylene oxide copolymers, polyoxyethylated polyols (e.g., glycerol), polyvinyl alcohol, and mixtures thereof. Polyethylene glycol propionaldehyde may have advantages in manufacturing due to its stability in water. The polymer may be of any molecular weight, and may be branched or unbranched. The number of polymers attached to the antibody may vary, and if more than one polymer is attached, they can be the same or different molecules. In general, the number and/or type of polymers used for derivatization can be determined based on considerations including, but not limited to, the particular properties or functions of the antibody to be improved, whether the antibody derivative will be used in a therapy under defined conditions, etc.

**[00170]** In another embodiment, conjugates of an antibody and nonproteinaceous moiety that may be selectively heated by exposure to radiation are provided. In one embodiment, the

nonproteinaceous moiety is a carbon nanotube (Kam et al., Proc. Natl. Acad. Sci. USA 102: 11600-11605 (2005)). The radiation may be of any wavelength, and includes, but is not limited to, wavelengths that do not harm ordinary cells, but which heat the nonproteinaceous moiety to a temperature at which cells proximal to the antibody-nonproteinaceous moiety are killed.

#### Recombinant Methods and Compositions

**[00171]** Antibodies may be produced using recombinant methods and compositions, e.g., as described in U.S. Patent No. 4,816,567. In one embodiment, isolated nucleic acid encoding an anti-Abeta antibody described herein is provided. Such nucleic acid may encode an amino acid sequence comprising the VL and/or an amino acid sequence comprising the VH of the antibody (e.g., the light and/or heavy chains of the antibody). In a further embodiment, one or more vectors (e.g., expression vectors) comprising such nucleic acid are provided. In a further embodiment, a host cell comprising such nucleic acid is provided. In one such embodiment, a host cell comprises (e.g., has been transformed with): (1) a vector comprising a nucleic acid that encodes an amino acid sequence comprising the VL of the antibody and an amino acid sequence comprising the VH of the antibody, or (2) a first vector comprising a nucleic acid that encodes an amino acid sequence comprising the VL of the antibody and a second vector comprising a nucleic acid that encodes an amino acid sequence comprising the VH of the antibody. In one embodiment, the host cell is eukaryotic, e.g. a Chinese Hamster Ovary (CHO) cell or lymphoid cell (e.g., Y0, NS0, Sp20 cell). In one embodiment, a method of making an anti-Abeta antibody is provided, wherein the method comprises culturing a host cell comprising a nucleic acid encoding the antibody, as provided above, under conditions suitable for expression of the antibody, and optionally recovering the antibody from the host cell (or host cell culture medium).

**[00172]** For recombinant production of an anti-Abeta antibody, nucleic acid encoding an antibody, e.g., as described above, is isolated and inserted into one or more vectors for further cloning and/or expression in a host cell. Such nucleic acid may be readily isolated and sequenced using conventional procedures (e.g., by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of the antibody).

**[00173]** Suitable host cells for cloning or expression of antibody-encoding vectors include prokaryotic or eukaryotic cells described herein. For example, antibodies may be produced in bacteria, in particular when glycosylation and Fc effector function are not needed. For expression of antibody fragments and polypeptides in bacteria, see, e.g., U.S. Patent Nos.

5,648,237, 5,789,199, and 5,840,523. (See also Charlton, *Methods in Molecular Biology*, Vol. 248 (B.K.C. Lo, ed., Humana Press, Totowa, NJ, 2003), pp. 245-254, describing expression of antibody fragments in *E. coli*.) After expression, the antibody may be isolated from the bacterial cell paste in a soluble fraction and can be further purified.

**[00174]** In addition to prokaryotes, eukaryotic microbes such as filamentous fungi or yeast are suitable cloning or expression hosts for antibody-encoding vectors, including fungi and yeast strains whose glycosylation pathways have been “humanized,” resulting in the production of an antibody with a partially or fully human glycosylation pattern. See Gerngross, *Nat. Biotech.* 22:1409-1414 (2004), and Li et al., *Nat. Biotech.* 24:210-215 (2006).

**[00175]** Suitable host cells for the expression of glycosylated antibody are also derived from multicellular organisms (invertebrates and vertebrates). Examples of invertebrate cells include plant and insect cells. Numerous baculoviral strains have been identified which may be used in conjunction with insect cells, particularly for transfection of *Spodoptera frugiperda* cells.

**[00176]** Plant cell cultures can also be utilized as hosts. See, e.g., US Patent Nos. 5,959,177, 6,040,498, 6,420,548, 7,125,978, and 6,417,429 (describing PLANTIBODIESTM technology for producing antibodies in transgenic plants).

**[00177]** Vertebrate cells may also be used as hosts. For example, mammalian cell lines that are adapted to grow in suspension may be useful. Other examples of useful mammalian host cell lines are monkey kidney CV1 line transformed by SV40 (COS-7); human embryonic kidney line (293 or 293 cells as described, e.g., in Graham et al., *J. Gen Virol.* 36:59 (1977)); baby hamster kidney cells (BHK); mouse sertoli cells (TM4 cells as described, e.g., in Mather, *Biol. Reprod.* 23:243-251 (1980)); monkey kidney cells (CV1); African green monkey kidney cells (VERO-76); human cervical carcinoma cells (HELA); canine kidney cells (MDCK; buffalo rat liver cells (BRL 3A); human lung cells (W138); human liver cells (Hep G2); mouse mammary tumor (MMT 060562); TRI cells, as described, e.g., in Mather et al., *Annals N.Y. Acad. Sci.* 383:44-68 (1982); MRC 5 cells; and FS4 cells. Other useful mammalian host cell lines include Chinese hamster ovary (CHO) cells, including DHFR-CHO cells (Urlaub et al., *Proc. Natl. Acad. Sci. USA* 77:4216 (1980)); and myeloma cell lines such as Y0, NS0 and Sp2/0. For a review of certain mammalian host cell lines suitable for antibody production, see, e.g., Yazaki and Wu, *Methods in Molecular Biology*, Vol. 248 (B.K.C. Lo, ed., Humana Press, Totowa, NJ), pp. 255-268 (2003).

## Assays

**[00178]** Anti-Abeta antibodies provided herein may be identified, screened for, or characterized for their physical/chemical properties and/or biological activities by various assays known in the art.

### Binding assays and other assays

**[00179]** In one aspect, an antibody of the invention is tested for its antigen binding activity, e.g., by known methods such as ELISA, Western blot, etc.

**[00180]** In another aspect, competition assays may be used to identify an antibody that competes with an anti-Abeta antibody of the invention for binding to Abeta. In certain embodiments, such a competing antibody binds to the same epitope (e.g., a linear or a conformational epitope) that is bound by crenezumab or another anti-Abeta antibody specified herein. Detailed exemplary methods for mapping an epitope to which an antibody binds are provided in Morris (1996) "Epitope Mapping Protocols," in *Methods in Molecular Biology* vol. 66 (Humana Press, Totowa, NJ).

**[00181]** In an exemplary competition assay, immobilized Abeta in the desired form (e.g., monomeric, oligomeric, or fibril) is incubated in a solution comprising a first labeled antibody that binds to Abeta (e.g., crenezumab) and a second unlabeled antibody that is being tested for its ability to compete with the first antibody for binding to Abeta. The second antibody may be present in a hybridoma supernatant. As a control, immobilized Abeta is incubated in a solution comprising the first labeled antibody but not the second unlabeled antibody. After incubation under conditions permissive for binding of the first antibody to Abeta, excess unbound antibody is removed, and the amount of label associated with immobilized Abeta is measured. If the amount of label associated with immobilized Abeta is substantially reduced in the test sample relative to the control sample, then that indicates that the second antibody is competing with the first antibody for binding to Abeta. See Harlow and Lane (1988) *Antibodies: A Laboratory Manual* ch.14 (Cold Spring Harbor Laboratory, Cold Spring Harbor, NY).

### Activity assays

**[00182]** In one aspect, assays are provided for identifying anti-Abeta antibodies thereof having biological activity, for example the biological activity of crenezumab. Biological activity may include, but is not limited to, e.g., prevention of aggregation of monomeric Abeta into oligomeric Abeta, or disaggregation of oligomeric Abeta into monomeric Abeta. Antibodies having such biological activity in vivo and/or in vitro are also provided.

**[00183]** In certain embodiments, an antibody of the invention is tested for such biological activity.

#### Methods and Compositions for Diagnostics and Detection

**[00184]** In certain embodiments, any of the anti-Abeta antibodies provided herein is useful for detecting the presence of Abeta in a biological sample. The term “detecting” as used herein encompasses quantitative or qualitative detection. In certain embodiments, a biological sample comprises a cell or tissue, such as serum, plasma, nasal swabs, sputum, cerebrospinal fluid, , aqueous humor of the eye and the like, or tissue or cell samples obtained from an organism such as samples containing neural or brain tissue.

**[00185]** In one embodiment, an anti-Abeta antibody for use in a method of diagnosis or detection is provided. In a further aspect, a method of detecting the presence of Abeta in a biological sample is provided. In certain embodiments, the method comprises contacting the biological sample with an anti-Abeta antibody as described herein under conditions permissive for binding of the anti-Abeta antibody to Abeta, and detecting whether a complex is formed between the anti-Abeta antibody and Abeta. Such method may be an in vitro or in vivo method.

**[00186]** Exemplary disorders that may be diagnosed using an antibody of the invention are diseases and disorders caused by or associated with amyloid or amyloid-like proteins. These include, but are not limited to, diseases and disorders caused by the presence or activity of amyloid-like proteins in monomeric, fibril, or polymeric state, or any combination of the three, including by amyloid plaques. Exemplary diseases include, but are not limited to, secondary amyloidosis and age-related amyloidosis, such as diseases including, but not limited to, neurological disorders such as Alzheimer’s Disease (“AD”), diseases or conditions characterized by a loss of cognitive memory capacity such as, for example, mild cognitive impairment (MCI), Lewy body dementia, Down’s syndrome, hereditary cerebral hemorrhage with amyloidosis (Dutch type), the Guam Parkinson-Demential complex and other diseases which are based on or associated with amyloid-like proteins such as progressive supranuclear palsy, multiple sclerosis, Creutzfeldt Jacob disease, Parkinson’s disease, HIV-related dementia, ALS (amyotrophic lateral sclerosis), inclusion-body myositis (IBM), adult onset diabetes, endocrine tumor and senile cardiac amyloidosis, and various eye diseases including macular degeneration, drusen-related optic neuropathy, glaucoma, and cataract due to beta-amyloid deposition.



**[00187]** In certain embodiments, labeled anti-Abeta antibodies are provided. Labels include, but are not limited to, labels or moieties that are detected directly (such as fluorescent, chromophoric, electron-dense, chemiluminescent, and radioactive labels), as well as moieties, such as enzymes or ligands, that are detected indirectly, e.g., through an enzymatic reaction or molecular interaction. Exemplary labels include, but are not limited to, the radioisotopes <sup>32</sup>P, <sup>14</sup>C, <sup>125</sup>I, <sup>3</sup>H, and <sup>131</sup>I, fluorophores such as rare earth chelates or fluorescein and its derivatives, rhodamine and its derivatives, dansyl, umbelliferone, luciferases, e.g., firefly luciferase and bacterial luciferase (U.S. Patent No. 4,737,456), luciferin, 2,3-dihydrophthalazinediones, horseradish peroxidase (HRP), alkaline phosphatase,  $\beta$ -galactosidase, glucoamylase, lysozyme, saccharide oxidases, e.g., glucose oxidase, galactose oxidase, and glucose-6-phosphate dehydrogenase, heterocyclic oxidases such as uricase and xanthine oxidase, coupled with an enzyme that employs hydrogen peroxide to oxidize a dye precursor such as HRP, lactoperoxidase, or microperoxidase, biotin/avidin, spin labels, bacteriophage labels, stable free radicals, and the like.

#### Pharmaceutical Formulations

**[00188]** Pharmaceutical formulations of an anti-Abeta antibody as described herein are prepared by mixing such antibody or molecule having the desired degree of purity with one or more optional pharmaceutically acceptable carriers (Remington's Pharmaceutical Sciences 16th edition, Osol, A. Ed. (1980)), in the form of lyophilized formulations or aqueous solutions. Pharmaceutically acceptable carriers are generally nontoxic to recipients at the dosages and concentrations employed, and include, but are not limited to: buffers such as phosphate, citrate, and other organic acids; antioxidants including ascorbic acid and methionine; preservatives (such as octadecyldimethylbenzyl ammonium chloride; hexamethonium chloride; benzalkonium chloride; benzethonium chloride; phenol, butyl or benzyl alcohol; alkyl parabens such as methyl or propyl paraben; catechol; resorcinol; cyclohexanol; 3-pentanol; and m-cresol); low molecular weight (less than about 10 residues) polypeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, histidine, arginine, or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugars such as sucrose, mannitol, trehalose or sorbitol; salt-forming counter-ions such as sodium; metal complexes (e.g. Zn-protein complexes); and/or non-ionic surfactants such as polyethylene glycol (PEG). Exemplary pharmaceutically acceptable carriers herein further include

interstitial drug dispersion agents such as soluble neutral-active hyaluronidase glycoproteins (sHASEGP), for example, human soluble PH-20 hyaluronidase glycoproteins, such as rHuPH20 (HYLENEX®, Baxter International, Inc.). Certain exemplary sHASEGPs and methods of use, including rHuPH20, are described in US Patent Publication Nos. 2005/0260186 and 2006/0104968. In one aspect, a sHASEGP is combined with one or more additional glycosaminoglycanases such as chondroitinases.

**[00189]** In one embodiment, an antibody of the invention may be formulated in an arginine buffer. In one aspect, the arginine buffer may be an arginine succinate buffer. In one such aspect, the concentration of the arginine succinate buffer may be 50 mM or greater. In another such aspect, the concentration of the arginine succinate buffer may be 100 mM or greater. In another such aspect, the concentration of the arginine succinate buffer may be 150 mM or greater. In another such aspect, the concentration of the arginine succinate buffer may be 200 mM or greater. In another aspect, the arginine buffer formulation may further contain a surfactant. In another such aspect, the surfactant is a polysorbate. In another such aspect, the polysorbate is polysorbate 20. In another such aspect, the concentration of polysorbate 20 in the formulation is 0.1% or less. In another such aspect, the concentration of polysorbate 20 in the formulation is 0.05% or less. In another aspect, the pH of the arginine buffer formulation is between 4.5 and 7.0. In another aspect, the pH of the arginine buffer formulation is between 5.0 and 6.5. In another aspect, the pH of the arginine buffer formulation is between 5.0 and 6.0. In another aspect, the pH of the arginine buffer formulation is 5.5. In any of the foregoing embodiments and aspects, the antibody of the invention may be crenezumab.

**[00190]** Exemplary lyophilized antibody formulations are described in US Patent No. 6,267,958. Aqueous antibody formulations include those described in US Patent No. 6,171,586 and WO2006/044908, the latter formulations including a histidine-acetate buffer.

**[00191]** The formulation herein may also contain more than one active ingredients as necessary for the particular indication being treated, preferably those with complementary activities that do not adversely affect each other. For example, it may be desirable to further provide one or more compounds to prevent or treat symptoms of Alzheimer's Disease. Such active ingredients are suitably present in combination in amounts that are effective for the purpose intended.

**[00192]** Active ingredients may be entrapped in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization, for example,

hydroxymethylcellulose or gelatin-microcapsules and poly-(methylmethacrylate) microcapsules, respectively, in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano-particles and nanocapsules) or in macroemulsions. Such techniques are disclosed in Remington's Pharmaceutical Sciences 16th edition, Osol, A. Ed. (1980).

**[00193]** Sustained-release preparations may be prepared. Suitable examples of sustained-release preparations include semipermeable matrices of solid hydrophobic polymers containing the antibody, which matrices are in the form of shaped articles, e.g. films, or microcapsules.

**[00194]** The formulations to be used for in vivo administration are generally sterile. Sterility may be readily accomplished, e.g., by filtration through sterile filtration membranes.  
Therapeutic Methods and Compositions

**[00195]** As shown herein, intravenous administration of high (multi-gram) doses of crenezumab did not trigger or increase the incidence of ARIA-E or any dose-limiting toxicity in patients suffering from AD. Specifically, patients with mild to moderate AD, including patients with mild AD and ApoE4 positive patients, as well as patients with brain amyloid load typically seen in patients diagnosed with AD, showed no increase in ARIA-E compared to a placebo at doses two to three times higher than doses tested in a Phase II clinical trial. These multi-gram doses exceed the doses reported for other anti-Abeta antibodies tested in the clinic, up to several fold higher than doses of anti-Abeta antibodies reported to increase the incidence of edema in the brain.

**[00196]** Therefore, in one embodiment, an antibody of the invention is administered in doses of 1500 mg or greater to treat AD, including mild to moderate AD, mild AD, and early AD, without increased risk of one or more adverse effects, such as ARIA-E. In another embodiment, an antibody of the invention is used to treat an amyloidosis. In one such embodiment, the amyloidosis is mild cognitive impairment. In another such embodiment, the amyloidosis is Down's syndrome. In another such embodiment, the amyloidosis is hereditary cerebral hemorrhage with amyloidosis (Dutch type). In another such embodiment, the amyloidosis is the Guam Parkinson-Dementia complex. In another such embodiment, the amyloidosis is an ocular disease related to drusen or other amyloid deposit in the eye. In one aspect, the ocular disease is macular degeneration. In another aspect, the ocular disease is a drusen-related optic neuropathy. In another aspect, the ocular disease is glaucoma. In

another aspect, the ocular disease is cataract. In any of the foregoing embodiments and aspects, the antibody of the invention may be crenezumab.

**[00197]** A patient is typically first assessed for the presence of one or more amyloidosis prior to determining the suitability of an antibody of the invention to treat such patient. As one nonlimiting example, AD may be diagnosed in a patient using the “NINCDS-ADRDA” (Neurological and Communicative Disorders and Stroke-Alzheimer’s Disease Related Disorders Assessment) criteria. See McKhann, et al., 1984, *Neurology* 34:939–44. Another exemplary method for diagnosing AD or prodromal AD relies on the criteria and guidelines set forth in the National Institute on Aging/Alzheimer’s Association (NIAAA) Diagnostic Criteria and Guidelines for AD (McKhann et al., 2011, *Alz & Dement* 7:263-269 (for mild AD); Albert et al., 2011, *Alz & Dement* 7:270-279 (for prodromal AD or mild cognitive impairment)). A potential patient to be administered one or more antibodies of the invention may also be tested for the presence or absence of one or more genetic markers which may predispose such patient either to (i) a higher or lower likelihood of such patient experiencing one or more amyloidoses, or (ii) a higher or lower likelihood of such patient experiencing one or more adverse events or side effects during the course of administration of an antibody of the invention. As one nonlimiting example, it is known that patients carrying the ApoE4 allele have a substantially higher risk of developing AD than those lacking the allele (Saunders et al., *Neurology* 1993; 43:1467-72; Prekumar et al., *Am. J. Pathol.* 1996; 148:2083-95), and that such patients were disproportionately represented in ARIA-type adverse events observed in the clinical trial of bapineuzumab, another anti-Abeta antibody (Sperling et al., *Alzheimer’s & Dementia* 2011, 7:367-385; Salloway et al., *N. Engl. J. Med.* 2014, 370:322-333).

**[00198]** In some embodiments, the antibody of the invention is used to treat mild to moderate AD in a patient. In some embodiments, the antibody of the invention is used to treat early AD in a patient. In some embodiments, the antibody of the invention is used to treat mild AD. In some embodiments, the antibody of the invention is used to treat prodromal AD in a patient. The patient can be ApoE4 positive or ApoE4 negative. In some embodiments, the antibody of the invention is used to treat an ApoE4 positive patient suffering from mild to moderate AD or early AD. In some embodiments, the antibody of the invention is used to treat a patient suffering from mild AD. In some embodiments, the antibody of the invention is used to treat a patient suffering from prodromal AD.

**[00199]** In some embodiments, the antibody of the invention is used to treat a patient having an MMSE score of between 20 and 30, between 20 and 26, between 24 and 30, between 21 and 26, between 22 and 26, between 22 and 28, between 23 and 26, between 24 and 26, or between 25 and 26. In some embodiments, the patient has an MMSE score between 22 and 26. As used herein, an MMSE score between two numbers includes the numbers at each end of the range. For example, an MMSE score between 22 and 26 includes MMSE scores of 22 and 26.

**[00200]** In some embodiments, the antibodies of the invention are used to treat a patient who is 'amyloid positive,' e.g., a patient having brain amyloid deposits that are typical of a patient diagnosed with AD or a patient having a positive florbetapir PET scan. In some embodiments, the antibodies of the invention are used to reduce the accumulation of brain amyloid deposits or neuritic plaques (i.e., to reduce an increase in brain amyloid burden or load).

**[00201]** The antibodies of the invention are useful for treating mild to moderate AD without increasing the incidence of ARIA-E or ARIA-H. In some embodiments, the patients are suffering from mild AD. In some embodiments, the patients are ApoE4 positive. In some embodiments, the patients are ApoE4 positive and suffering from mild AD.

**[00202]** As evidenced in the Examples herein, doses of 1500 mg or more can be used to treat patients with milder forms of AD without increasing the incidence of ARIA-E. Consequently, in some embodiments, the antibody of the invention is used to treat a patient with early AD. In certain embodiments, the patient to be treated has one or more of the following characteristics: (a) mild cognitive impairment (MCI) due to AD; (b) one or more biomarkers indicative of Alzheimer's Disease without a clinically detectable deficit; (c) an objective memory loss quantified using the Free and Cued Selective Reminding Test (FCSRT) as a score of 27 or greater; an MMSE of 24-30; (d) a global Clinical Dementia Rating (CDR) of 0.5; and (e) a positive amyloid PET scan (as determined by a qualified reader).

**[00203]** Antibodies of the invention are formulated, dosed, and administered in a fashion consistent with good medical practice. Factors for consideration in this context include the particular disorder being treated, the particular mammal being treated, the clinical condition of the individual subject, the cause of the disorder, the site of delivery of the agent, the method of administration, the scheduling of administration, and other factors known to medical practitioners.

### Routes of Administration

**[00204]** An antibody of the invention (and any additional therapeutic agent) can be administered by any suitable means, including parenteral, intrapulmonary, and intranasal, and, if desired for local treatment, intralesional administration. Parenteral infusions include intramuscular, intravenous, intraarterial, intraperitoneal, or subcutaneous administration. Dosing can be by any suitable route, e.g. by injections, such as intravenous or subcutaneous injections, depending in part on whether the administration is brief or chronic. In one embodiment, the antibody is injected subcutaneously. In another embodiment, the antibody is injected intravenously. In another embodiment, the antibody is administered using a syringe (e.g., prefilled or not) or an autoinjector. In another embodiment, the antibody is inhaled.

### Dosing

**[00205]** For the treatment of an amyloidosis, the appropriate dosage of an antibody of the invention (when used alone or in combination with one or more other additional therapeutic agents) will depend on the specific type of disease to be treated, the type of antibody, the severity and course of the disease, previous therapy, the patient's clinical history and response to the antibody, and the discretion of the attending physician. The antibody is suitably administered to the patient at one time or over a series of treatments. Various dosing schedules including, but not limited to, single or multiple administrations over various time-points, bolus administration, and pulse infusion are contemplated herein.

**[00206]** Depending on the type and severity of the disease, about 45 mg/kg to 200 mg/kg (e.g. 50 mg/kg-200 mg/kg, or any dosage within that range) of antibody can be an initial candidate dosage for administration to the patient, whether, for example, by one or more separate administrations, or by continuous infusion. One typical daily, weekly, bi-weekly, monthly, or quarterly dosage might range from about 45 mg/kg to 200 mg/kg or more, depending on the factors mentioned above. The dosage can be administered in a single dose or a divided dose (e.g., two doses of 30 mg/kg for a total dose of 60 mg/kg). For repeated administrations over several weeks or longer, depending on the condition, the treatment would generally be sustained until a desired suppression of disease symptoms occurs. One exemplary dosage of the antibody would be in the range from about 50 mg/kg to about 150 mg/kg. Thus, one or more doses of about 15 mg/kg, about 20 mg/kg, about 25 mg/kg, about 30 mg/kg, about 35 mg/kg, about 40 mg/kg, about 50 mg/kg, about 60 mg/kg, about 70 mg/kg, about 80 mg/kg, about 90 mg/kg, about 100 mg/kg, about 110 mg/kg, about 120 mg/kg, or about 130 mg/kg (or any combination thereof) may be administered to the patient.

In some embodiments, the total dose administered is in the range of 1500 mg to 24000 mg. An exemplary dose of about 1500 mg, about 1600 mg, about 1700 mg, about 1800 mg, about 2000 mg, about 3000 mg, about 4000 mg, about 5000 mg, about 6000 mg, about 7000 mg, about 7200 mg, about 10000 mg, about 10500 mg, about 11000 mg, about 12000 mg, about 13000 mg, about 14000 mg, about 15000 mg, about 16000 mg, about 17000 mg, about 18000 mg, about 19000 mg, about 20000 mg, about 20500 mg, about 21000 mg, about 22000 mg, about 23000 mg, or about 24000 mg (or any combination thereof) may be administered to the patient. Such doses may be administered intermittently, e.g. every week, every two weeks, every three weeks, every four weeks, every month, every two months, every three months, or every six months. In some embodiments, the patient receives from one to thirty five doses (e.g. about eighteen doses of the antibody). However, other dosage regimens may be useful. The progress of this therapy can be monitored by conventional techniques and assays.

**[00207]** In certain embodiments, an antibody of the invention is administered at a dose of 45 mg/kg, 50 mg/kg, 60 mg/kg, 70 mg/kg, 80 mg/kg, 90 mg/kg, 100 mg/kg, 110 mg/kg, 120 mg/kg, 130 mg/kg, 140 mg/kg, 150 mg/kg or a flat dose, e.g., 1500 mg, 1800 mg, 2000 mg, 2400 mg, 3000 mg, 3200 mg, 4000 mg, 5000 mg, 5400 mg, 6000 mg, 7000 mg, 7200 mg, 8000 mg, or higher. In some embodiments, the dose is administered by intravenous injection every 2 weeks or every 4 weeks for a period of time. In some embodiments, the dose is administered by subcutaneous injection every 2 weeks or every 4 weeks for a period of time. In certain embodiments, the period of time is 6 months, one year, eighteen months, two years, five years, ten years, 15 years, 20 years, or the lifetime of the patient.

#### Monitoring/Assessing Response to Therapeutic Treatment

**[00208]** As used in methods of the present disclosure, the antibody, or antigen-binding fragment hereof, provides therapeutic effect or benefit to the patient. In certain embodiments, the therapeutic benefit is a delay in, or inhibition of, progression of AD or a reduction in clinical, functional, or cognitive decline. In some embodiments, therapeutic effect or benefit is reflected in a “patient response” or “response” (and grammatical variations thereof). Patient response can be assessed using any endpoint indicating a benefit to the patient, including, without limitation, (1) inhibition, to some extent, of disease progression, including slowing down and complete arrest; (2) reduction in amount of plaque or reduction in brain amyloid accumulation; (3) improvement in one or more assessment metrics, including but not limited to ADAS-Cog, iADL, and CDR-SB scales; (4) improvement in daily functioning of the patient; (5) increase in concentration of one or more biomarkers, e.g., Abeta, in

cerebrospinal fluid; and (6) decrease in one or more biomarkers indicative of the presence of AD. An assessment of patient response may also include an assessment of any adverse events that may occur that may be correlated with the treatment.

**[00209]** In one embodiment, the cognitive ability and daily functioning of the patient is assessed prior to, during, and/or after a course of therapy with an antibody of the invention. A number of cognitive and functional assessment tools have been developed for use in assessing, diagnosing, and scoring mental function, cognition, and neurological deficit. These tools include, but are not limited to, the ADAS-Cog, including the 12 item ADAS-Cog (ADAS-Cog12), the 13-item ADAS-Cog (ADAS-Cog13), the 14-item ADAS-Cog (ADAS-Cog14); the CDR-SB, including CDR Judgment and Problem solving and CDR Memory components; the Instrumental Activities of Daily Living (iADL); and the MMSE.

**[00210]** “ADAS-Cog” refers to the Alzheimer’s Disease Assessment Scale Cognitive Subscale, a multi-part cognitive assessment. See Rosen et al., 1984, *Amer. J. Psych.* 141:1356-1364; Mohs et al., 1997, *Alzheimer’s Disease Assoc. Disorders* 11(2):S13-S21. The higher the numerical score on the ADAS-Cog, the greater the tested patient’s deficit or impairment relative to another individual with a lower score. The ADAS-Cog may be used as one measure for assessing whether a treatment for AD is therapeutically effective. An increase in ADAS-Cog score is indicative of worsening in the patient’s condition, whereas a decrease in ADAS-Cog score denotes improvement in the patient’s condition. As used herein, a “decline in ADAS-Cog performance” or an “increase in ADAS-Cog score” indicates a worsening in the patient’s condition and may reflect progression of AD. The ADAS-Cog is an examiner-administered battery that assesses multiple cognitive domains, including memory, comprehension, praxis, orientation, and spontaneous speech (Rosen et al. 1984, *Am J Psychiatr* 141:1356–64; Mohs et al. 1997, *Alzheimer Dis Assoc Disord* 11(S2):S13–S21). The ADAS-Cog is a standard primary endpoint in AD treatment trials (Mani 2004, *Stat Med* 23:305–14). The ADAS-Cog12 is the 70-point version of the ADAS-Cog plus a 10-point Delayed Word Recall item assessing recall of a learned word list. Other ADAS-Cog scales include the ADAS-Cog13 and ADAS-Cog14.

**[00211]** In some embodiments, the methods of treatment provided herein provide a reduction in cognitive decline as measured by an ADAS-Cog score that is at least about 30%, at least about 35%, at least about 40%, or at least about 45% lower relative to placebo.

**[00212]** “MMSE” refers to the Mini Mental State Examination, which provides a score between 1 and 30. See Folstein, et al., 1975, *J. Psychiatr. Res.* 12:189–98. Scores of 26 and



lower are generally considered to be indicative of a deficit. The lower the numerical score on the MMSE, the greater the tested patient's deficit or impairment relative to another individual with a lower score. An increase in MMSE score may be indicative of improvement in the patient's condition, whereas a decrease in MMSE score may denote worsening in the patient's condition.

**[00213]** "CDR-SB" refers to the Clinical Dementia Rating Scale / Sum of Boxes. See Hughes et al, 1982, Br J Psychiatry 140:566-72. CDR-assesses 6 components: memory, orientation, judgment/problem solving, community affairs, home and hobbies, and personal care. The test is administered to both the patient and the caregiver and each component (or each "box"), is scored on a scale of 0 to 3. A complete CDR-SB score is based on the sum of the scores across all 6 boxes. Subscores can be obtained for each of the boxes or components individually as well, e.g., CDR/ Memory or CDR/ Judgment and Problem solving. As used herein, a "decline in CDR-SB performance" or an "increase in CDR-SB score" indicates a worsening in the patient's condition and may reflect progression of AD. In some embodiments, the methods of treatment provided herein provide a reduction in decline in CDR-SB performance of at least about 30%, at least about 35%, or at least about 40% relative to placebo.

**[00214]** "iADL" refers to the Instrumental Activities of Daily Living scale. See Lawton, M.P., and Brody, E.M., 1969, Gerontologist 9:179-186. This scale measures the ability to perform typical daily activities such as housekeeping, laundry, operating a telephone, shopping, preparing meals, etc. The lower the score, the more impaired the individual is in conducting activities of daily living. In some embodiments, the methods of treatment provided herein provide a reduction in decline of at least about 10%, at least about 15%, or at least about 20% on the iADL scale relative to placebo.

**[00215]** Brain amyloid load or burden can be determined using neurological imaging techniques and tools, for example using PET (positron emission tomography) scanning. Serial PET scans of a patient taken over time, e.g., before and after administration of a treatment (or at one or more intervals throughout the course of a treatment regimen), can permit detection of increased, decreased, or unchanged amyloid burden in the brain. This technique can further be used to determine whether amyloid accumulation is increasing or decreasing. In some embodiments, detection of amyloid deposits in the brain is performed using florbetapir 18F. In some embodiments, a florbetapir PET scan is considered positive if, based on a

centralized visual read of the scan, it establishes the presence of moderate-to-frequent neuritic plaques.

#### Co-Administration

**[00216]** The antibody need not be, but is optionally formulated with one or more agents currently used to prevent or treat the disorder in question or one or more of its symptoms. The effective amount of such other agents depends on the amount of antibody present in the formulation, the type of disorder or treatment, and other factors discussed above. These are generally used in the same dosages and with administration routes as described herein, or about from 1 to 99% of the dosages described herein, or in any dosage and by any route that is empirically/clinically determined to be appropriate. It will be understood by one of ordinary skill in the art that an antibody of the invention may be co-administered simultaneously with any of the foregoing compounds, or may be administered prior to or subsequent to administration of any of the foregoing compounds.

**[00217]** When treating an amyloidosis with an antibody of the invention, a neurological drug may be co-administered. Such neurological drug may be selected from the group including, but not limited to, an antibody or other binding molecule (including, but not limited to a small molecule, a peptide, an aptamer, or other protein binder) that specifically binds to a target selected from: beta secretase, tau, presenilin, amyloid precursor protein or portions thereof, amyloid beta peptide or oligomers or fibrils thereof, death receptor 6 (DR6), receptor for advanced glycation endproducts (RAGE), parkin, and huntingtin; a cholinesterase inhibitor (i.e., galantamine, donepezil, rivastigmine and tacrine); an NMDA receptor antagonist (i.e., memantine), a monoamine depletor (i.e., tetrabenazine); an ergoloid mesylate; an anticholinergic antiparkinsonism agent (i.e., procyclidine, diphenhydramine, trihexylphenidyl, benztropine, biperiden and trihexyphenidyl); a dopaminergic antiparkinsonism agent (i.e., entacapone, selegiline, pramipexole, bromocriptine, rotigotine, selegiline, ropinirole, rasagiline, apomorphine, carbidopa, levodopa, pergolide, tolcapone and amantadine); a tetrabenazine; an anti-inflammatory (including, but not limited to, a nonsteroidal anti-inflammatory drug (i.e., indomethacin and other compounds listed above); a hormone (i.e., estrogen, progesterone and leuprolide); a vitamin (i.e., folate and nicotinamide); a dimebolin; a homotaurine (i.e., 3-aminopropanesulfonic acid; 3APS); a serotonin receptor activity modulator (i.e., xaliproden); an, an interferon, and a glucocorticoid or corticosteroid. In some embodiments, one or more anti-Abeta antibodies other than crenezumab are co-administered. Non-limiting examples of such anti-Abeta antibodies

include solanezumab, bapineuzumab, and aducanumab. In some embodiments, a therapeutic agent targeting tau is co-administered with an antibody of the invention. The term “corticosteroid” includes, but is not limited to fluticasone (including fluticasone propionate (FP)), beclometasone, budesonide, ciclesonide, mometasone, flunisolide, betamethasone and triamcinolone. “Inhalable corticosteroid” means a corticosteroid that is suitable for delivery by inhalation. Exemplary inhalable corticosteroids are fluticasone, beclomethasone dipropionate, budesonide, mometasone furoate, ciclesonide, flunisolide, and triamcinolone acetonide.

**[00218]** When treating an amyloidosis that is an ocular disease or disorder with an antibody of the invention, a neurological drug may be selected that is an anti-angiogenic ophthalmic agent (i.e., bevacizumab, ranibizumab and pegaptanib), an ophthalmic glaucoma agent (i.e., carbachol, epinephrine, demecarium bromide, apraclonidine, brimonidine, brinzolamide, levobunolol, timolol, betaxolol, dorzolamide, bimatoprost, carteolol, metipranolol, dipivefrin, travoprost and latanoprost), a carbonic anhydrase inhibitor (i.e., methazolamide and acetazolamide), an ophthalmic antihistamine (i.e., naphazoline, phenylephrine and tetrahydrozoline), an ocular lubricant, an ophthalmic steroid (i.e., fluorometholone, prednisolone, loteprednol, dexamethasone, difluprednate, rimexolone, fluocinolone, medrysone and triamcinolone), an ophthalmic anesthetic (i.e., lidocaine, proparacaine and tetracaine), an ophthalmic anti-infective (i.e., levofloxacin, gatifloxacin, ciprofloxacin, moxifloxacin, chloramphenicol, bacitracin/polymyxin b, sulfacetamide, tobramycin, azithromycin, besifloxacin, norfloxacin, sulfisoxazole, gentamicin, idoxuridine, erythromycin, natamycin, gramicidin, neomycin, ofloxacin, trifluridine, ganciclovir, vidarabine), an ophthalmic anti-inflammatory agent (i.e., nepafenac, ketorolac, flurbiprofen, suprofen, cyclosporine, triamcinolone, diclofenac and bromfenac), and an ophthalmic antihistamine or decongestant (i.e., ketotifen, olopatadine, epinastine, naphazoline, cromolyn, tetrahydrozoline, pemirolast, bepotastine, naphazoline, phenylephrine, nedocromil, lodoxamide, phenylephrine, emedastine and azelastine). It is understood that any of the above formulations or therapeutic methods may be carried out using an immunoconjugate of the invention in place of or in addition to an anti-Aβ antibody.

#### Articles of Manufacture

**[00219]** In another aspect of the invention, an article of manufacture containing materials useful for the treatment, prevention and/or diagnosis of the disorders described above is provided. The article of manufacture comprises a container and a label or package insert on

or associated with the container. Suitable containers include, for example, bottles, vials, syringes, IV solution bags, etc. The containers may be formed from a variety of materials such as glass or plastic. The container holds a composition which is by itself or combined with another composition effective for treating, preventing and/or diagnosing the condition and may have a sterile access port (for example the container may be an intravenous solution bag or a vial having a stopper pierceable by a hypodermic injection needle). At least one active agent in the composition is an antibody of the invention. The label or package insert indicates that the composition is used for treating the condition of choice. Moreover, the article of manufacture may comprise (a) a first container with a composition contained therein, wherein the composition comprises an antibody of the invention; and (b) a second container with a composition contained therein, wherein the composition comprises a further cytotoxic or otherwise therapeutic agent. The article of manufacture in this embodiment of the invention may further comprise a package insert indicating that the compositions can be used to treat a particular condition. Alternatively, or additionally, the article of manufacture may further comprise a second (or third) container comprising a pharmaceutically-acceptable buffer, such as bacteriostatic water for injection (BWFI), phosphate-buffered saline, Ringer's solution and dextrose solution. It may further include other materials desirable from a commercial and user standpoint, including other buffers, diluents, filters, needles, and syringes.

**[00220]** It is understood that any of the above articles of manufacture may include an immunoconjugate of the invention in place of or in addition to an anti-A $\beta$  antibody.

#### EXEMPLARY EMBODIMENTS

**[00221]** Provided herein are exemplary embodiments, for illustration.

1. A method of reducing the decline in functional or cognitive capacity in a patient diagnosed with early or mild to moderate Alzheimer's Disease (AD) comprising administering to a patient suffering from early or mild to moderate AD a humanized monoclonal anti-amyloid beta (A $\beta$ ) antibody that binds within residues 13 and 24 of amyloid  $\beta$  (1-42)(SEQ ID NO:1) in an amount effective to slow the decline in functional or cognitive capacity in the patient.
2. The method of **embodiment 1**, wherein the antibody is capable of binding oligomeric and monomeric forms of amyloid  $\beta$ .
3. The method of **claim 1**, wherein the antibody is an IgG4 antibody.

4. The method of **embodiment 2 or 3**, wherein the antibody comprises six hypervariable regions (HVRs), wherein:
  - (i) HVR-H1 is SEQ ID NO:2;
  - (ii) HVR-H2 is SEQ ID NO:3;
  - (iii) HVR-H3 is SEQ ID NO:4;
  - (iv) HVR-L1 is SEQ ID NO:6;
  - (v) HVR-L2 is SEQ ID NO:7; and
  - (vi) HVR-L3 is SEQ ID NO:8.
5. The method of **embodiment 4**, wherein the antibody comprises a heavy chain having the amino acid sequence of SEQ ID NO:5 and a light chain having the amino acid sequence of SEQ ID NO:9.
6. The method of **embodiment 5**, wherein the antibody is crenezumab.
7. The method of **any one of the preceding embodiments**, wherein decline in cognitive capacity is assessed by determining the patient's score before and after administration of said antibody using a 12-item Alzheimer's Disease Assessment Scale – Cognition (ADAS-Cog12), 13-item Alzheimer's Disease Assessment Scale – Cognition (ADAS-Cog13), or 14-item Alzheimer's Disease Assessment Scale – Cognition (ADAS-Cog12) test, optionally wherein the reduction in cognitive decline as measured by ADAS-Cog is at least 30%, at least 35%, at least 40%, or at least 45% relative to placebo.
8. The method of **embodiment 7**, wherein the patient is ApoE4 positive.
9. The method of **embodiment 7**, wherein the patient is suffering from mild AD.
10. The method of **embodiment 7**, wherein the patient is suffering from early AD.
11. The method of **any one of embodiments 1 to 8**, wherein the patient has an MMSE score of at least 20, between 20 and 30, between 20 and 26, between 24 and 30, between 21 and 26, between 22 and 26, between 22 and 28, between 23 and 26, between 24 and 26, or between 25 and 26 before initiation of treatment.
12. The method of **embodiment 11**, wherein the patient has an MMSE between 22 and 26.

13. The method of **any one of the preceding embodiments**, wherein the antibody is administered at a dose of 30 mg/kg to 200 mg/kg or 100 mg/kg to 200 mg/kg of patient body weight.
14. The method of **embodiment 13**, wherein the antibody is administered at a dose of at least 60 mg/kg.
15. The method of **embodiment 14**, wherein the antibody is administered at a dose of 60 mg/kg, 100 mg/kg, 120 mg/kg, or 150 mg/kg.
16. The method of **embodiment 13 or 14**, wherein the antibody is administered via intravenous injection.
17. The method of **any one of embodiments 13 to 16**, wherein the antibody is administered every 2 weeks, every 4 weeks, every month, every two months, or every six months.
18. A method of treating early or mild to moderate AD without increasing the risk of an adverse event comprising administering to a patient diagnosed with early or mild to moderate AD an amount of a humanized monoclonal anti-A $\beta$  antibody that binds within residues 13 and 24 of amyloid  $\beta$  (1-42)(SEQ ID NO:1) that is effective to treat the AD without increasing the risk of a treatment emergent adverse event, wherein the adverse event is selected from: (i) Amyloid-Related Imaging Abnormality—Edema (ARIA-E) and (ii) Amyloid-Related Imaging Abnormality—Hemorrhage (ARIA-H).
19. The method of **embodiment 18**, wherein the antibody is capable of binding oligomeric and monomeric forms of amyloid  $\beta$ .
20. The method of **embodiment 18**, wherein the antibody is an IgG4 antibody.
21. The method of **embodiment 19**, wherein the antibody comprises six hypervariable regions (HVRs), wherein:
  - (i) HVR-H1 is SEQ ID NO:2;
  - (ii) HVR-H2 is SEQ ID NO:3;
  - (iii) HVR-H3 is SEQ ID NO:4;
  - (iv) HVR-L1 is SEQ ID NO:6;
  - (v) HVR-L2 is SEQ ID NO:7; and

- (vi) HVR-L3 is SEQ ID NO:8.
22. The method of **embodiment 21**, wherein the antibody comprises a heavy chain having the amino acid sequence of SEQ ID NO:5 and a light chain having the amino acid sequence of SEQ ID NO:9.
23. The method of **embodiment 22**, wherein the antibody is crenezumab.
24. The method of any one of embodiments **18 to 23**, wherein the patient is ApoE4 positive.
25. The method of **any one of embodiments 18 to 23**, wherein the adverse event is ARIA-E.
26. The method of **embodiment 25**, wherein, if a treatment emergent ARIA-E is detected, administration of the antibody is halted and, optionally, treatment for ARIA-E is administered.
27. The method of **embodiment 26**, further comprising resuming administration of said antibody after the ARIA-E is resolved, wherein the antibody is administered at a lower dose than before administration was halted.
28. The method of **embodiment 18**, wherein if one or more new ARIA-Es is detected in the patient during treatment with said antibody, no more antibody is administered, and, optionally, a corticosteroid is administered to the patient.
29. The method of **embodiment 28**, wherein the patient is ApoE4 positive.
30. A method of reducing the decline in functional or cognitive capacity in a patient diagnosed with early or mild to moderate Alzheimer's Disease (AD) comprising administering to an ApoE4 positive patient suffering from early or mild to moderate AD a humanized monoclonal anti-amyloid beta (A $\beta$ ) antibody that binds within residues 13 and 24 of amyloid  $\beta$  (1-42)(SEQ ID NO:1) in an amount effective to slow the decline in functional or cognitive capacity in the patient.
31. The method of **embodiment 30**, wherein the antibody is capable of binding oligomeric and monomeric forms of amyloid  $\beta$ .
32. The method of **embodiment 30**, wherein the antibody is an IgG4 antibody.

33. The method of **embodiment 31 or 32**, wherein the antibody comprises six hypervariable regions (HVRs), wherein:
- (i) HVR-H1 is SEQ ID NO:2;
  - (ii) HVR-H2 is SEQ ID NO:3;
  - (iii) HVR-H3 is SEQ ID NO:4;
  - (iv) HVR-L1 is SEQ ID NO:6;
  - (v) HVR-L2 is SEQ ID NO:7; and
  - (vi) HVR-L3 is SEQ ID NO:8.
34. The method of **embodiment 33**, wherein the antibody comprises a heavy chain having the amino acid sequence of SEQ ID NO:5 and a light chain having the amino acid sequence of SEQ ID NO:9.
35. The method of **embodiment 34**, wherein the antibody is crenezumab.
36. The method of **any one of embodiments 30 to 36**, wherein decline in cognitive capacity is assessed by determining the patient's score before and after administration of said antibody using an ADAS-Cog12, ADAS-Cog13, or ADAS-Cog14 test, optionally wherein the reduction in cognitive decline as measured by ADAS-Cog is at least 30%, at least 35%, at least 40%, or at least 45% relative to placebo.
37. The method of **embodiment 36**, wherein the patient has mild AD.
38. The method of **embodiment 36**, wherein the patient has early AD.
39. The method of **any one of embodiments 30 to 37**, wherein the patient has an MMSE score of at least 20, between 20 and 30, between 20 and 26, between 24 and 30, between 21 and 26, between 22 and 26, between 22 and 28, between 23 and 26, between 24 and 26, or between 25 and 26 before initiation of treatment.
40. The method of **embodiment 39**, wherein the patient has an MMSE score between 22 and 26.
41. The method of **any one of embodiments 30 to 39**, wherein the antibody is administered at a dose of 30 mg/kg to 200 mg/kg or 100 mg/kg to 200 mg/kg of patient body weight.



42. The method of **embodiment 41**, wherein the antibody is administered at a dose of at least 60 mg/kg.
43. The method of **embodiment 42**, wherein the antibody is administered at a dose of 60 mg/kg, 100 mg/kg, 120 mg/kg, or 150 mg/kg.
44. The method of **embodiment 41 or 42**, wherein the antibody is administered via intravenous injection.
45. The method of **any one of embodiments 41 to 44**, wherein the antibody is administered every 2 weeks, every 4 weeks, every month, every two months, or every six months.
46. A method of treating early or mild to moderate AD without increasing the risk of an adverse event comprising administering to an ApoE4 positive patient diagnosed with early or mild to moderate AD an amount of a humanized monoclonal anti-A $\beta$  antibody that binds within residues 13 and 24 of amyloid  $\beta$  (1-42)(SEQ ID NO:1) that is effective to treat the AD without increasing the risk of a treatment emergent adverse event, wherein the adverse event is selected from: (i) Amyloid-Related Imaging Abnormality—Edema (ARIA-E) and (ii) Amyloid-Related Imaging Abnormality—Hemorrhage (ARIA-H).
47. The method of **embodiment 46**, wherein the antibody is capable of binding oligomeric and monomeric forms of amyloid  $\beta$ .
48. The method of **embodiment 46**, wherein the antibody is an IgG4 antibody.
49. The method of **embodiment 47**, wherein the antibody comprises six hypervariable regions (HVRs), wherein:
- (i) HVR-H1 is SEQ ID NO:2;
  - (ii) HVR-H2 is SEQ ID NO:3;
  - (iii) HVR-H3 is SEQ ID NO:4;
  - (iv) HVR-L1 is SEQ ID NO:6;
  - (v) HVR-L2 is SEQ ID NO:7; and
  - (vi) HVR-L3 is SEQ ID NO:8.

50. The method of **embodiment 49**, wherein the antibody comprises a heavy chain having the amino acid sequence of SEQ ID NO:5 and a light chain having the amino acid sequence of SEQ ID NO:9.
51. The method of **embodiment 50**, wherein the antibody is crenezumab.
52. The method of **any one of embodiments 46 to 51**, wherein the adverse event is ARIA-E.
53. The method of **embodiment 52**, wherein if a treatment emergent ARIA-E is detected, administration of the antibody is halted and, optionally, treatment for ARIA-E is administered.
54. The method of **embodiment 53**, further comprising resuming administration of said antibody after the ARIA-E is resolved, optionally comprising resuming administration of said antibody at a lower dose than before administration was halted.
55. The method of **embodiment 46**, wherein if one or more new ARIA-Es is detected in the patient during treatment with said antibody, no more antibody is administered, and, optionally, a corticosteroid is administered to the patient.
56. The method of **any one of the preceding embodiments**, wherein the patient is concurrently treated with one or more agents selected from the group consisting of: a therapeutic agent that specifically binds to a target; a cholinesterase inhibitor; an NMDA receptor antagonist; a monoamine depletor; an ergoloid mesylate; an anticholinergic antiparkinsonism agent; a dopaminergic antiparkinsonism agent; a tetrabenazine; an anti-inflammatory agent; a hormone; a vitamin; a dimebolin; a homotaurine; a serotonin receptor activity modulator; an interferon, and a glucocorticoid; an anti-Abeta antibody other than crenezumab; an antibiotic; an anti-viral agent.
57. The method of **embodiment 56**, wherein the agent is a cholinesterase inhibitor.
58. The method of **embodiment 57**, wherein the cholinesterase inhibitor is selected from the group consisting of galantamine, donepezil, rivastigmine and tacrine.
59. The method of **embodiment 56**, wherein the agent is an NMDA receptor antagonist.
60. The method of **embodiment 59**, wherein the NMDA receptor antagonist is memantine or a salt thereof.

61. The method of **embodiment 56**, wherein the agent is a therapeutic agent that specifically binds to a target and the target is selected from the group consisting of beta secretase, tau, presenilin, amyloid precursor protein or portions thereof, amyloid beta peptide or oligomers or fibrils thereof, death receptor 6 (DR6), receptor for advanced glycation endproducts (RAGE), parkin, and huntingtin.
62. The method of **embodiment 56**, wherein the agent is a monoamine depletory, optionally tetrabenazine.
63. The method of **embodiment 56**, wherein the agent is an anticholinergic antiparkinsonism agent selected from the group consisting of procyclidine, diphenhydramine, trihexylphenidyl, benztropine, biperiden and trihexyphenidyl.
64. The method of **embodiment 56**, wherein the agent is a dopaminergic antiparkinsonism agent selected from the group consisting of: entacapone, selegiline, pramipexole, bromocriptine, rotigotine, selegiline, ropinirole, rasagiline, apomorphine, carbidopa, levodopa, pergolide, tolcapone and amantadine.
65. The method of **embodiment 56**, wherein the agent is an anti-inflammatory agent selected from the group consisting of: a nonsteroidal anti-inflammatory drug and indomethacin.
66. The method of **embodiment 56**, wherein the agent is a hormone selected from the group consisting of: estrogen, progesterone and leuprolide.
67. The method of **embodiment 56**, wherein the agent is a vitamin selected from the group consisting of: folate and nicotinamide.
68. The method of **embodiment 56**, wherein the agent is a homotaurine, which is 3-aminopropanesulfonic acid or 3APS.
69. The method of **embodiment 56**, wherein the agent is xaliproden.
70. A method of slowing clinical decline in a patient diagnosed with early or mild to moderate Alzheimer's Disease (AD) comprising administering to a patient suffering from early or mild to moderate AD a humanized monoclonal anti-amyloid beta (A $\beta$ ) antibody that binds within residues 13 and 24 of amyloid  $\beta$  (1-42)(SEQ ID NO:1) in an amount effective to slow the decline in the patient.

71. The method of **embodiment 70**, wherein the antibody is capable of binding oligomeric and monomeric forms of amyloid  $\beta$ .
72. The method of **embodiment 70**, wherein the antibody is an IgG4 antibody.
73. The method of **embodiment 71 or 72**, wherein the antibody comprises six hypervariable regions (HVRs), wherein:
- (i) HVR-H1 is SEQ ID NO:2;
  - (ii) HVR-H2 is SEQ ID NO:3;
  - (iii) HVR-H3 is SEQ ID NO:4;
  - (iv) HVR-L1 is SEQ ID NO:6;
  - (v) HVR-L2 is SEQ ID NO:7; and
  - (vi) HVR-L3 is SEQ ID NO:8.
74. The method of **embodiment 73**, wherein the antibody comprises a heavy chain having the amino acid sequence of SEQ ID NO:5 and a light chain having the amino acid sequence of SEQ ID NO:9.
75. The method of **embodiment 74**, wherein the antibody is crenezumab.
76. The method of **any one of embodiments 70 to 75**, further comprising a decline in cognitive capacity assessed by determining the patient's score before and after administration of said antibody using a 12-item Alzheimer's Disease Assessment Scale – Cognition (ADAS-Cog12), a 13-item Alzheimer's Disease Assessment Scale – Cognition (ADAS-Cog13), or a 14-item Alzheimer's Disease Assessment Scale – Cognition (ADAS-Cog14) test, optionally wherein the reduction in cognitive decline as measured by ADAS-Cog is at least 30%, at least 35%, at least 40%, or at least 45% relative to placebo.
77. The method of **embodiment 76**, wherein the patient is ApoE4 positive.
78. The method of **embodiment 76**, wherein the patient is suffering from mild AD.
79. The method of **embodiment 76**, wherein the patient is suffering from early AD.
80. The method of **any one of embodiments 70 to 78**, wherein the patient has an MMSE score of at least 20, between 20 and 30, between 20 and 26, between 24 and 30, between 21

and 26, between 22 and 26, between 22 and 28, between 23 and 26, between 24 and 26, or between 25 and 26 before initiation of treatment.

81. The method of **embodiment 80**, wherein the patient has an MMSE score between 22 and 26.

82. The method of **any one of embodiments 70 to 80**, wherein the antibody is administered at a dose of 30 mg/kg to 200 mg/kg or 100 mg/kg to 200 mg/kg of patient body weight.

83. The method of **embodiment 82**, wherein the antibody is administered at a dose of at least 60 mg/kg.

84. The method of **embodiment 83**, wherein the antibody is administered at a dose of 60 mg/kg, 100 mg/kg, 120 mg/kg, or 150 mg/kg.

85. The method of **embodiment 82 or 83**, wherein the antibody is administered via intravenous injection.

86. The method of **any one of embodiments 82 to 85**, wherein the antibody is administered every 2 weeks, every 4 weeks, every month, every two months, or every six months.

87. A method of treating early or mild AD in a subject, comprising administering to a patient suffering from early or mild AD a humanized monoclonal anti-amyloid beta ( $A\beta$ ) antibody that binds within residues 13 and 24 of amyloid  $\beta$  (1-42)(SEQ ID NO:1) in an amount effective to treat the AD.

88. The method of **embodiment 87**, wherein the antibody is capable of binding oligomeric and monomeric forms of amyloid  $\beta$ .

89. The method of **embodiment 87**, wherein the antibody is an IgG4 antibody.

90. The method of **embodiment 88 or 89**, wherein the antibody comprises six hypervariable regions (HVRs), wherein:

- (i) HVR-H1 is SEQ ID NO:2;
- (ii) HVR-H2 is SEQ ID NO:3;
- (iii) HVR-H3 is SEQ ID NO:4;
- (iv) HVR-L1 is SEQ ID NO:6;

- (v) HVR-L2 is SEQ ID NO:7; and
- (vi) HVR-L3 is SEQ ID NO:8.

91. The method of **embodiment 90**, wherein the antibody comprises a heavy chain having the amino acid sequence of SEQ ID NO:5 and a light chain having the amino acid sequence of SEQ ID NO:9.
92. The method of **embodiment 91**, wherein the antibody is crenezumab.
93. The method of **any one of embodiments 87 to 92**, wherein the amount is effective to reduce decline in cognitive capacity, which is assessed by determining the patient's score before and after administration of said antibody using a 12-item Alzheimer's Disease Assessment Scale – Cognition (ADAS-Cog12), a 13-item Alzheimer's Disease Assessment Scale – Cognition (ADAS-Cog13), or a 14-item Alzheimer's Disease Assessment Scale – Cognition (ADAS-Cog14) test, optionally wherein the reduction in cognitive decline as measured by ADAS-Cog is at least 30%, at least 35%, at least 40%, or at least 45% relative to placebo.
94. The method of **embodiment 93**, wherein the patient is ApoE4 positive.
95. The method of **any one of embodiments 87 to 94**, wherein the patient has an MMSE score of at least 20, between 20 and 30, between 20 and 26, between 24 and 30, between 21 and 26, between 22 and 26, between 22 and 28, between 23 and 26, between 24 and 26, or between 25 and 26 before initiation of treatment.
96. The method of **embodiment 95**, wherein the patient has an MMSE score between 22 and 26.
97. The method of **any one of embodiments 87 to 95**, wherein the antibody is administered at a dose of 30 mg/kg to 200 mg/kg or 100 mg/kg to 200 mg/kg of patient body weight.
98. The method of **embodiment 97**, wherein the antibody is administered at a dose of at least 60 mg/kg.
99. The method of **embodiment 98**, wherein the antibody is administered at a dose of 60 mg/kg, 100 mg/kg, 120 mg/kg, or 150 mg/kg.

100. The method of **embodiment 97 or 98**, wherein the antibody is administered via intravenous injection.
101. The method of **any one of embodiments 97 to 100**, wherein the antibody is administered every 2 weeks, every 4 weeks, every month, every two months, or every six months.
102. The method of **any one of embodiments 70 to 101**, wherein the patient is concurrently treated with one or more agents selected from the group consisting of: a therapeutic agent that specifically binds to a target; a cholinesterase inhibitor; an NMDA receptor antagonist; a monoamine depletor; an ergoloid mesylate; an anticholinergic antiparkinsonism agent; a dopaminergic antiparkinsonism agent; a tetrabenazine; an anti-inflammatory agent; a hormone; a vitamin; a dimebolin; a homotaurine; a serotonin receptor activity modulator; an interferon, and a glucocorticoid; an anti-Abeta antibody; an antibiotic; an anti-viral agent.
103. The method of **embodiment 102**, wherein the agent is a cholinesterase inhibitor.
104. The method of **embodiment 103**, wherein the cholinesterase inhibitor is selected from the group consisting of galantamine, donepezil, rivastigmine and tacrine.
105. The method of **embodiment 102**, wherein the agent is an NMDA receptor antagonist.
106. The method of **embodiment 105**, wherein the NMDA receptor antagonist is memantine or a salt thereof.
107. The method of **embodiment 102**, wherein the agent is a therapeutic agent that specifically binds to a target and the target is selected from the group consisting of beta secretase, tau, presenilin, amyloid precursor protein or portions thereof, amyloid beta peptide or oligomers or fibrils thereof, death receptor 6 (DR6), receptor for advanced glycation endproducts (RAGE), parkin, and huntingtin.
108. The method of **embodiment 102**, wherein the agent is a monoamine depletory, optionally tetrabenazine.
109. The method of **embodiment 102**, wherein the agent is an anticholinergic antiparkinsonism agent selected from the group consisting of procyclidine, diphenhydramine, trihexylphenidyl, benzotropine, biperiden and trihexyphenidyl.

110. The method of **embodiment 102**, wherein the agent is a dopaminergic antiparkinsonism agent selected from the group consisting of: entacapone, selegiline, pramipexole, bromocriptine, rotigotine, selegiline, ropinirole, rasagiline, apomorphine, carbidopa, levodopa, pergolide, tolcapone and amantadine.
111. The method of **embodiment 102**, wherein the agent is an anti-inflammatory agent selected from the group consisting of: a nonsteroidal anti-inflammatory drug and indomethacin.
112. The method of **embodiment 102**, wherein the agent is a hormone selected from the group consisting of: estrogen, progesterone and leuprolide.
113. The method of **embodiment 102**, wherein the agent is a vitamin selected from the group consisting of: folate and nicotinamide.
114. The method of **embodiment 102**, wherein the agent is a homotaurine, which is 3-aminopropanesulfonic acid or 3APS.
115. The method of **embodiment 102**, wherein the agent is xaliproden.
116. The method of **embodiment 102**, wherein the agent is an anti-A $\beta$  antibody other than crenezumab.

### EXAMPLES

#### **EXAMPLE 1 -- Clinical study of the safety and tolerability of crenezumab, a humanized anti-A $\beta$ monoclonal antibody, administered to patients with mild to moderate Alzheimer's Disease**

**[00222]** A randomized, double blind Phase I trial was conducted, using a placebo control, to evaluate the safety, tolerability, and pharmacokinetics of the humanized monoclonal anti-amyloid beta ("A $\beta$ ") antibody crenezumab in patients diagnosed with mild to moderate Alzheimer's Disease (AD). The study was designed to assess doses up to 8 times the dose administered to patients in a Phase II clinical trial. Participants included in the study were, at the time of screening, between the ages of 50 and 90, with a Mini-Mental State Examination (MMSE) score of 18 to 28 points (inclusive), a Geriatric Depression Scale (GDS-15) score of less than 6, a Clinical Dementia Rating – Global Score (CDR-GS) of 0.5 or 1.0, and a diagnosis of probable mild-to-moderate Alzheimer's disease by NINCDS-ADRDA criteria. Participants were also required to have increased brain (cerebral) amyloid as measured by amyloid PET scan (e.g., florbetapir amyloid PET scan). The study was designed to ensure that



at least 50% of the enrolled participants in each dose level were ApoE4 positive (carrying at least one ApoE4 allele, also referred to as ApoE4 carriers).

**[00223]** Participants were eligible for the study regardless of whether they were receiving approved standard of care treatment for AD (i.e. ChEIs or memantine, or SOUVENAID) provided the standard of care treatment had been administered as a stable dose for at least 3 months prior to screening.

**[00224]** The study had a screening period lasting up to 6 weeks, followed by a double-blind treatment period and Dose Limiting Toxicity (“DLT”) assessment window of 13 weeks, with the final safety assessment, including MRI, following the last dose (i.e. fourth dose in Week 13), followed by an ongoing Open Label Extension phase during which patients previously receiving placebo were rolled over to the active treatment arm. See FIG. 4A-B (Study schematics). Treatment (or placebo) was administered via intravenous infusion, once every 4 weeks (Q4W).

**[00225]** For each dose studied, participants were enrolled in the trial and randomized into one of two arms, a treatment (i.e., crenezumab) arm and a placebo arm in a 5:1 (treatment arm:placebo arm) randomization, with at least 12 participants at each dose level tested (e.g., 10 participants per treatment arm and 2 participants per placebo arm). The safety and tolerability of crenezumab was assessed by measuring the frequency and severity of treatment emergent adverse events throughout the trial, especially instances of symptomatic or asymptomatic ARIA-E (including cerebral vasogenic edema), symptomatic or asymptomatic ARIA-H (including cerebral microhemorrhage), and cerebral macrohemorrhage. The presence and/or number of cerebral vasogenic edema cases was assessed by amyloid PET scan, using 18F florbetapir (AMYVID) as an amyloid imaging agent, and MRI. The presence and/or number of ARIA events was assessed during the screening period (Week 1-6), and during the double-blind treatment period, at Weeks 5 and 13, followed by further assessment during an Open Label Extension period or at Week 21 for participants not enrolling in an Open Label Extension. Blood samples were collected and serum concentration of crenezumab at each dose level was measured. Serum exposure (area under the curve and peak concentration) was also determined across the doses.

**[00226]** Three dose cohorts were studied. In a first cohort, two dose levels were studied: 30 mg/kg and 45 mg/kg. A total of 26 participants were enrolled in the first cohort. The participants received crenezumab (at least 4 doses) or placebo, based on the randomization scheme of 5:1 per dose level. In a second cohort, a 60 mg/kg dose level was studied in which

participants were randomized to either 60 mg/kg crenezumab or placebo in a 5:1 ratio, in a total of 26 participants. In a third cohort, a 120 mg/kg dose was studied in which participants were randomized to either 120 mg/kg crenezumab or placebo in a 5:1 ratio. Escalation from cohort 1 to cohort 2, and from cohort 2 to cohort 3, occurred after review by an internal unblinded safety monitoring committee of all available safety and tolerability data up to the date that the last participant in the previous cohort completed the second dose of study drug and subsequent MRI scan. All participants underwent regular brain MRI to monitor for ARIA-E and ARIA-H. Patient baseline characteristics for the first two cohorts are shown in Table 2 below.

**Table 2**

<b>Characteristics</b>	<b>Cohort 1 (n=26)</b>	<b>Cohort 2 (n=26)</b>
Age, mean (range)	73.5 (54–82)	72.7 (51–87)
Males, n (%)	14 (54)	15 (58)
ApoE status, n (%)		
E2/E3	1 (2)	0
E3/E3	5 (19)	5 (19)
E3/E4	17 (65)	18 (69)
E4/E4	3 (12)	3 (12)
Baseline MMSE, mean (range)	22.4 (18–28)	22.7 (18–29)
Median duration of exposure, weeks (range)	52.1 (4–64)	32.1 (12–40)

**[00227]** Based on observations and interim analyses during the 12-week double-blind study period of the first and second cohorts, the safety and tolerability profile of crenezumab at doses of 30 mg/kg, 45 mg/kg, and 60 mg/kg was unchanged from that reported for doses up to 15 mg/kg. No dose-limiting toxicity or drug-related serious adverse events were reported. In particular, in the period under review at the interim analysis, there was no instance of amyloid-related imaging abnormalities-edema/effusion, or ARIA-E, reported even at doses up to three times higher than those previously tested. A single instance of pneumonia, unrelated to study drug, was reported.

**[00228]** Ongoing results from the first and second cohorts are shown in the tables below. Data for Tables 3 and 4 were collected from patients in cohorts 1 and 2 as follows. From the first cohort, of the 26 patients enrolled, 23 patients reached Week 25, 22 patients reached Week 49, of these at least 3 patients reached Week 61. Five patients discontinued the trial.

From the second cohort, of the 26 patients enrolled, 23 patients reached Week 25, 22 patients reached Week 37, and 4 patients discontinued the trial.

**Table 3**

<b>Adverse Events (AEs), n (%)</b>	<b>Cohort 1 (n=26)</b>	<b>Cohort 2 (n=26)</b>
Total number of patients with at least one AE	21 (81)	20 (77)
AE related to study treatment (as assessed by investigator)	7 (27)	4 (15)
AE Grade $\geq$ 3 (severe, life-threatening or resulting in death)	1 (4)	1 (4)
Serious AE	1 (4)	2 (8)
Treatment withdrawal due to AE	1 (4)	2 (8)

**[00229]** Adverse events and their grading were defined according to Common Terminology Criteria for Adverse Events (CTCAE) version 4.0. The serious AEs observed at the time of interim analysis were as follows: in cohort 1, one patient had a malignant melanoma and in cohort 2, one patient suffered an accidental overdose, pneumonia and subdural hematoma, while the second patient had atypical chest pain. In cohort 1, a patient with malignant melanoma discontinued the study. In cohort 2, both patients who discontinued the study had non-serious events (one confusional state, one with atrial fibrillation).

**[00230]** Common and selected AEs are shown in Table 4 below. In cohort 1, 3 patients presented with cerebral microhemorrhage and one patient with cerebellar microhemorrhage.

**Table 4**

<b>AEs, n (%)</b>	<b>Cohort 1 (n=26)</b>	<b>Cohort 2 (n=26)</b>
<b>Common AEs</b>		
Headache	4 (15)	2 (8)
Anxiety	5 (19)	1 (4)
Fatigue	0	4 (15)
Muscle spasms	3 (12)	0
Cerebral microhemorrhage	4 (15)	2 (8)
<b>Selected events</b>		
Pneumonia	0	1 (4)
Infusion-related reactions	3 (12)	3 (12)

[00231] The majority of the AEs observed for crenezumab doses of 30, 45, and 60 mg/kg were low grade and non-serious. No dose-limiting toxicities were observed and no ARIA-E events were reported. There were no investigator assessed drug-related serious AEs. Few patients experienced ARIA-H (6 out of a total of 52). All ARIA-H events were asymptomatic and did not result in treatment discontinuation.

[00232] Preliminary data for the third cohort (120 mg/kg) dose were consistent with other cohorts. The data showed no significant change to the safety and tolerability of crenezumab, even at this highest dose tested.

[00233] In addition to assessing safety of increased crenezumab dosing, the study also confirmed a dose-proportional increase in serum concentration of crenezumab as the dose increased from 15 mg/kg to 30 mg/kg, 45 mg/kg, and 60 mg/kg. In particular serum concentrations up to four-fold higher relative to serum concentrations measured following 15 mg/kg doses given at the same interval, consistent with and confirming pharmacokinetics models based on Phase II data for crenezumab. See FIG. 5 and FIG. 6A-B.

[00234] These data establish that crenezumab can be administered at high doses to achieve higher serum concentration without increasing the incidence of a treatment emergent adverse event such as ARIA-E in amyloid positive patients suffering from mild to moderate AD.

**EXAMPLE 2 -- Clinical study of crenezumab, a humanized anti-A $\beta$  monoclonal antibody, in the treatment of prodromal to mild Alzheimer's Disease**

**Study Design and Objectives**

[00235] A multi-center, randomized, double-blind, placebo-controlled trial is conducted, to confirm the impact of the humanized monoclonal anti-amyloid beta ("A $\beta$ ") antibody crenezumab in amyloid positive patients diagnosed with prodromal to mild Alzheimer's Disease (AD). Participants in the study are, at the time of screening, between the ages of 50 and 85, with a weight between 40 kg and 120 kg (inclusive), having evidence of the AD pathological process by a positive amyloid assessment either on cerebrospinal fluid (CSF) amyloid beta 1-42 levels as measured on the Elecsys® beta-amyloid(1-42) test system or amyloid PET scan. Additional criteria for inclusion are: (1) a demonstrated abnormal memory function at screening with a Free and Cued Selective Reminding Test-Immediate Recall (FCSRT) cueing index less than or equal to 0.67 and free recall less than or equal to 27; (2) evidence of retrospective decline confirmed by a diagnosis verification form; (3) mild symptomatology, as defined by a screening mini mental state examination (MMSE) score of greater than or equal to 22 points and Clinical Dementia Rating-Global Score (CDR-GS) of

0.5 or 1.0; (4) meeting National Institute on Aging/Alzheimer's Association (NIAAA) core clinical criteria for probable AD dementia or prodromal AD (consistent with the NIAAA diagnostic criteria and guidelines for mild cognitive impairment (MCI)).

**[00236]** Participants are randomized 1:1 to receive either intravenous (IV) infusion of crenezumab or placebo every 4 weeks (q4w) for 100 weeks. Approximately 750 participants are enrolled in the trial and randomized to either the treatment arm or the placebo arm. The final efficacy and safety assessment is performed 4 weeks after the last dose of crenezumab administration (Week 105). In the treatment arm, participants receive a 30 mg/kg, 45 mg/kg, 60 mg/kg, or 120 mg/kg dose of crenezumab. Patients are stratified according to: ApoE4 status (carrier versus non-carrier) and MMSE score.

**[00237]** Data are collected for changes in: CDR-SB, ADAS-Cog13, CDR-GS, ADAS-Cog12, ADCS-ADL, MMSE, amyloid burden as measured using florbetapir-PET, and Abeta levels in cerebrospinal fluid (CSF), at intervals throughout the trial. In addition, adverse events such as ARIA-E, ARIA-H, infusion or injection reactions, pneumonia, and immunogenic reactions, are also monitored.

EXAMPLE 3 – Exposure response to crenezumab supports a dose of 60 mg/kg in the treatment of prodromal to mild Alzheimer's Disease

#### Methods and Objectives

**[00238]** Phase 2 studies of crenezumab demonstrated a consistent treatment benefit in the 15 mg/kg intravenous dose for patients suffering from milder forms of AD, while a low 300 mg q2wk subcutaneous dose level lacked a consistent treatment effect across endpoints, suggesting that higher doses are associated with greater efficacy signals. In both Phase 2 studies, crenezumab was safe and well-tolerated supporting that the therapeutic window has not been fully explored. A disease progression model for mild to moderate AD was established that described the longitudinal changes of the clinical endpoints ADAS-Cog and CDR sum-of-boxes (CDR-SB) simultaneously for patients in the Phase 2 studies. The model was extended to describe the effect of key demographic covariates on disease progression, and the effect of crenezumab on each endpoint as a hyperbolic function. Clinical trial simulations with 1000 replications of potential clinical study designs were conducted across a range of doses, describing the likelihood of achieving a percent relative reduction of disease progression in treated patients compared to placebo for ADAS-Cog and CDR-SB.

#### Results

**[00239]** Model validation demonstrated that the model replicated the available clinical longitudinal data accurately and is fit for purpose for simulation of the disease progression and crenezumab treatment effect in the population of interest (milder AD population, baseline MMSE 22-26). The analysis showed faster disease progression in patients with moderate AD disease (lower baseline MMSE), ApoE4 positive genotype, female gender, and younger age. A relationship was seen between crenezumab exposure and treatment effect, which appeared to asymptote at the higher end of the range of exposures measured in Phase 2 studies. Crenezumab treatment effect was associated with high baseline MMSE and ApoE4 positive genotype supporting better treatment effect in patients with mild AD. Based on the analysis of the model that has been developed, it is now envisioned that a 60 mg/kg dose administered once every 4 weeks could achieve a substantial improvement over the previously tested high dose of 15 mg/kg. In particular, it is now predicted that this increased dose could achieve a 41% greater relative reduction on ADAS-Cog12, and 44% on the CDR-SB in the milder AD population (baseline MMSE 22-26) relative to the effects observed with the 15 mg/kg dose.

**EXAMPLE 4 --** Clinical study of crenezumab, a humanized anti-A $\beta$  monoclonal antibody, in the treatment of prodromal to mild Alzheimer's Disease

#### Study Design and Objectives

**[00240]** A multi-center, randomized, double-blind, placebo-controlled trial is conducted, to confirm the impact of the humanized monoclonal anti-amyloid beta ("A $\beta$ ") antibody crenezumab in amyloid positive patients diagnosed with prodromal to mild Alzheimer's Disease (AD). Participants in the study are, at the time of screening, between the ages of 50 and 85, with a weight between 40 kg and 120 kg (inclusive), having evidence of the AD pathological process by a positive amyloid assessment either on cerebrospinal fluid (CSF) amyloid beta 1-42 levels as measured on the Elecsys® beta-amyloid(1-42) test system or amyloid PET scan. Additional criteria for inclusion are: (1) a demonstrated abnormal memory function at screening with a Free and Cued Selective Reminding Test-Immediate Recall (FCSRT) cueing index less than or equal to 0.67 and free recall less than or equal to 27; (2) evidence of retrospective decline confirmed by a diagnosis verification form; (3) mild symptomatology, as defined by a screening mini mental state examination (MMSE) score of greater than or equal to 22 points and Clinical Dementia Rating-Global Score (CDR-GS) of 0.5 or 1.0; (4) meeting National Institute on Aging/Alzheimer's Association (NIAAA) core clinical criteria for probable AD dementia or prodromal AD (consistent with the NIAAA diagnostic criteria and guidelines for mild cognitive impairment (MCI)). Patients are eligible

for the study regardless of whether or not they are receiving standard-of-care symptomatic medications for AD, such as memantine or cholinesterase inhibitors or combinations thereof.

**[00241]** The study consists of a screening period of eight weeks for each patient.

Participants are randomized 1:1 to receive either intravenous (IV) infusion of crenezumab or placebo every 4 weeks (q4w) for 100 weeks. A baseline visit is performed and referred to as “Week 1” of the study. Approximately 750 participants are enrolled in the trial and randomized to either the treatment arm or the placebo arm. The final efficacy and safety assessment is performed 4 weeks after the last dose of crenezumab administration (Week 105). Two follow-up visits are conducted at 16 and 52 weeks after the last dose. In the treatment arm, participants receive a 60 mg/kg dose of crenezumab. A total of 26 doses are given to patients who complete the study. Patients are stratified according to: ApoE4 status (carrier versus non-carrier), dementia status (prodromal AD versus mild AD), and presence or absence of anti-dementia medications at baseline.

**[00242]** Data are collected for changes in: CDR-SB, ADAS-Cog13, CDR-GS, ADAS-Cog12, ADCS-ADL, MMSE, amyloid burden as measured using florbetapir-PET, and Abeta levels in cerebrospinal fluid (CSF), at intervals throughout the trial. In addition, adverse events such as ARIA-E, ARIA-H, infusion or injection reactions, pneumonia, and immunogenic reactions, are also monitored.

**[00243]** Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, the descriptions and examples should not be construed as limiting the scope of the invention. The disclosures of all patent applications and publications and scientific literature cited herein are expressly incorporated in their entirety by reference for any purpose.

SEQUENCE LISTING KEY

SEQ ID NO:	Sequence
1	Human Aβ1-42 amino acid sequence: DAEFRHDSGYEVHHQKLVFFAEDVGSNKGAIIGLMVGGVVIA
2	Crenezumab HVR-H1 amino acid sequence: GFTFSSYGMS
3	Crenezumab HVR-H2 amino acid sequence: SINSNGGSTYYPDSVK
4	Crenezumab HVR-H3 amino acid sequence: GDY
5	Crenezumab heavy chain amino acid sequence (HVR regions marked in bold text): EVQLVESGGGLVQPGGSLRLSCAAS <b>GFTFSSYGMS</b> WVRQAPGK GLELVA <u>SINSNGGSTYYPDSVK</u> GRFTISRDNKNSLYLQMNSLR AEDTAVYYCAS <b>GDY</b> WGQGTTVTVSSASTKGPSVFPLAPCSRSTS ESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYS LSSVVTVPSSSLGTKYTCNVDHKPSNTKVDKRVESKYGPPCPPC PAPEFLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSQEDPEVQF NWYVDGVEVHNAKTKPREEQFNSTYRVVSVLTVLHQDWLNGK EYKCKVSNKGLPSSIEKTISKAKGQPREPQVYTLPPSQEEMTKNQ VSLTCLVKGFIYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLY SRLTVDKSRWQEGNVFSCSVMHEALHNHYTQKLSLSLGLG
6	Crenezumab HVR-L1 amino acid sequence: RSSQSLVYSNGDTYLH
7	Crenezumab HVR-L2 amino acid sequence: KVSNRFS
8	Crenezumab HVR-L3 amino acid sequence: SQSTHVPWT
9	Crenezumab light chain amino acid sequence (HVR regions marked in bold and underlined text): DIVMTQSPLSLPVTPGEPASISCR <b><u>RSSQSLVYSNGDTYLH</u></b> HWYLQKP GQSPQLLIY <b><u>KVSNRFS</u></b> GVDPDRFSGSGSGTDFTLKISRVEAEDVGV YYC <b><u>SQSTHVPWT</u></b> FGQGTKVEIKRTVAAPSVFIFPPSDEQLKSGTA SVCVCLLNFFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSTYS LSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC
10	Variable region heavy chain sequence: EVQLVESGGGLVQPGGSLRLSCAAS <b>GFTFSSYGMS</b> WVRQAPGKG LELVA <u>SINSNGGSTYYPDSVK</u> GRFTISRDNKNSLYLQMNSLRAE DTAVYYCAS <b>GDY</b> WGQGTTVTVSS
11	Variable region light chain sequence: DIVMTQSPLSLPVTPGEPASISCR <b><u>RSSQSLVYSNGDTYLH</u></b> HWYLQKPG QSPQLLIY <b><u>KVSNRFS</u></b> GVDPDRFSGSGSGTDFTLKISRVEAEDVGVYY CSQSTHVPWT <b><u>FGQGTKVEIK</u></b>



**CLAIMS:**

1. A method of treating early Alzheimer's Disease (AD) comprising: administering to a patient suffering from early AD between 1500 mg and 15000 mg of a humanized monoclonal anti-amyloid beta (A $\beta$ ) antibody that binds within residues 13 and 24 of amyloid  $\beta$  (1-42)(SEQ ID NO:1).
2. The method of **claim 1**, wherein the antibody is capable of binding oligomeric and monomeric forms of amyloid  $\beta$ .
3. The method of **claim 1**, wherein the antibody is an IgG4 antibody.
4. The method of **claim 2 or 3**, wherein the antibody comprises six hypervariable regions (HVRs), wherein:
  - (i) HVR-H1 is SEQ ID NO:2;
  - (ii) HVR-H2 is SEQ ID NO:3;
  - (iii) HVR-H3 is SEQ ID NO:4;
  - (iv) HVR-L1 is SEQ ID NO:6;
  - (v) HVR-L2 is SEQ ID NO:7; and
  - (vi) HVR-L3 is SEQ ID NO:8.
5. The method of **claim 4**, wherein the antibody comprises a heavy chain having the amino acid sequence of SEQ ID NO:5 and a light chain having the amino acid sequence of SEQ ID NO:9.
6. The method of **claim 5**, wherein the antibody is crenezumab.
7. The method of **any one of the preceding claims**, wherein the patient is amyloid positive.
8. The method of **claim 7**, wherein the patient is ApoE4 positive.
9. The method of **claim 7**, wherein the patient is suffering from mild AD.
10. The method of **claim 7**, wherein the patient is suffering from prodromal AD.
11. The method of **any one of claims 1 to 8**, wherein the patient has an MMSE score of at least 22, between 24 and 30, between 22 and 26, between 22 and 28, between 23 and 26, between 24 and 26, or between 25 and 26 before initiation of treatment.

12. The method of **claim 11**, wherein the patient has an MMSE between 22 and 26.
13. The method of **any one of the preceding claims**, wherein the antibody is administered at a dose between about 45 mg/kg and about 130 mg/kg of patient body weight.
14. The method of **claim 13**, wherein the antibody is administered at a dose of at least 50 mg/kg.
15. The method of **claim 14**, wherein the antibody is administered at a dose of 50 mg/kg, 60 mg/kg, 70 mg/kg, 80 mg/kg, 90 mg/kg, 100 mg/kg, 110 mg/kg, 120 mg/kg, or 130 mg/kg.
16. The method of **claim 13 or 14**, wherein the antibody is administered via intravenous injection.
17. The method of **any one of claims 13 to 16**, wherein the antibody is administered every 2 weeks, every 4 weeks, every month, every two months, or every six months.
18. The method of **any one of the preceding claims**, wherein the patient is concurrently treated with one or more agents selected from the group consisting of: a therapeutic agent that specifically binds to a target; a cholinesterase inhibitor; an NMDA receptor antagonist; a monoamine depletor; an ergoloid mesylate; an anticholinergic antiparkinsonism agent; a dopaminergic antiparkinsonism agent; a tetrabenazine; an anti-inflammatory agent; a hormone; a vitamin; a dimebolin; a homotaurine; a serotonin receptor activity modulator; an interferon, and a glucocorticoid; an anti-Abeta antibody other than crenezumab; an antibiotic; an anti-viral agent.
19. The method of **claim 18**, wherein the agent is a cholinesterase inhibitor.
20. The method of **claim 19**, wherein the cholinesterase inhibitor is selected from the group consisting of galantamine, donepezil, rivastigmine and tacrine.
21. The method of **claim 18**, wherein the agent is an NMDA receptor antagonist.
22. The method of **claim 21**, wherein the NMDA receptor antagonist is memantine or a salt thereof.
23. The method of **claim 18**, wherein the agent is a therapeutic agent that specifically binds to a target and the target is selected from the group consisting of beta secretase, tau, presenilin, amyloid precursor protein or portions thereof, amyloid beta peptide or oligomers

or fibrils thereof, death receptor 6 (DR6), receptor for advanced glycation endproducts (RAGE), parkin, and huntingtin.

24. The method of **claim 18**, wherein the agent is a monoamine depletory, optionally tetrabenazine.
25. The method of **claim 18**, wherein the agent is an anticholinergic antiparkinsonism agent selected from the group consisting of procyclidine, diphenhydramine, trihexylphenidyl, benztropine, biperiden and trihexyphenidyl.
26. The method of **claim 18**, wherein the agent is a dopaminergic antiparkinsonism agent selected from the group consisting of: entacapone, selegiline, pramipexole, bromocriptine, rotigotine, selegiline, ropinirole, rasagiline, apomorphine, carbidopa, levodopa, pergolide, tolcapone and amantadine.
27. The method of **claim 18**, wherein the agent is an anti-inflammatory agent selected from the group consisting of: a nonsteroidal anti-inflammatory drug and indomethacin.
28. The method of **claim 18**, wherein the agent is a hormone selected from the group consisting of: estrogen, progesterone and leuprolide.
29. The method of **claim 18**, wherein the agent is a vitamin selected from the group consisting of: folate and nicotinamide.
30. The method of **claim 18**, wherein the agent is a homotaurine, which is 3-aminopropanesulfonic acid or 3APS.
31. The method of **claim 18**, wherein the agent is xaliproden.
32. The method of **claim 18**, wherein the agent is an anti-Abeta antibody other than crenezumab

SEQ ID NO:1

1 DAEFRHDSGY EVHHQKLVFF AEDVGSNKGA IIGLMVGGVV IA

**FIG. 1**

HVR-H1 (SEQ ID NO:2): GFTFSSYGMS

HVR-H2 (SEQ ID NO:3): SINSNGGSTIYPDSVK

HVR-H3 (SEQ ID NO:4): GDY

HVR-L1 (SEQ ID NO:6): RSSQSLVYSNGDTYLH

HVR-L2 (SEQ ID NO:7): KVSNRFS

HVR-L3 (SEQ ID NO:8): SQSTHVPWT

**FIG. 2**

HC Sequence (SEQ ID NO:5)


1 EVQLVESGGG LVQPGGSLRL SCAASGFTFS SYGMSWVRQA PGKGLELVAS  
 51 INSNGGSTYY PDSVKGRFTI SRDNAKNSLY LQMNSLRAED TAVYYCASGD  
 101 YWGQGTTVTV SSASTKGPSV FPLAPCSRST SESTAALGCL VKDYFPEPVT  
 151 VSWNSGALTS GVHTFPAVLQ SSGLYSLSSV VTVPSSSLGT KTYTCNV<sup>D</sup>HK  
 201 PSNTKVDKRV ESKYGPPCPP CPAPEFLGGP SVFLFPPKPK DTLMISRTPE  
 251 VTCVVVDVSQ EDPEVQFNWY VDGVEVHNAK TKPREEQFNS TYRVVSVLTV  
 301 LHQDWLNGKE YKCKVSNKGL PSSIEKTISK AKGQPREPQV YTLPPSQEEM  
 351 TKNQVSLTCL VKGFYPSDIA VEWESNGQPE NNYKTTTPVL DSDGSFFLYS  
 401 RLTVDKSRWQ EGNVFSCSVM HEALHNHYTQ KSLSLSLG

LC Sequence (SEQ ID NO:9)

1 DIVMTQSPLS LPVTPGEPAS ISCRSSQSLV YSNGDTYLHW YLQKPGQSPQ  
 51 LLIYKVSNRF SGVPDRFSGS GSGTDFTLKI SRVEAEDVGV YYCSOSTHVP  
 101 WTFGQGTKVE IKRTVAAPSV FIFPPSDEQL KSGTASVVCL LNNFYPREAK  
 151 VQWKVDNALQ SGNSQESVTE QDSKDYSTYSL SSTLTLSKAD YEKHKVYACE  
 201 VTHQGLSSPV TKSFNRGEC


**FIG. 3**

2 / 4




**FIG. 4A**

Dose Escalation Schema




**FIG. 4B**

3 / 4




**FIG. 5**

4 / 4



**FIG. 6A**



**FIG. 6B**

INTERNATIONAL SEARCH REPORT

International application No  
PCT/US2017/014461

A. CLASSIFICATION OF SUBJECT MATTER  
INV. C07K16/18 G01N33/15 C07K16/00 A61K39/00  
ADD.  
According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED  
Minimum documentation searched (classification system followed by classification symbols)  
C07K G01N A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)  
EPO-Internal, BIOSIS, EMBASE, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 2010/150906 A1 (PFEIFER ANDREA [CH] ET AL) 17 June 2010 (2010-06-17) claims 171-180; sequence 52 -----	1-32
Y	EP 1 787 998 A1 (MITSUBISHI CHEM CORP [JP]; MITSUBISHI PHARMA CORP [JP]) 23 May 2007 (2007-05-23) [0063]discloses the application of 100mg/kg of an anti-amylospheroid antibody SEQ ID NO 1 is discloses as SEQ ID NO:2 in EP1787998; sequence 2 -----	1-32
Y	US 2014/328856 A1 (GELMONT DAVID M [US] ET AL) 6 November 2014 (2014-11-06) claims 79,188 ----- -/--	1-32

Further documents are listed in the continuation of Box C.

See patent family annex.

\* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search  22 March 2017	Date of mailing of the international search report  04/04/2017
--	--

Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer  Klee, Barbara
--	---



## INTERNATIONAL SEARCH REPORT

International application No  
PCT/US2017/014461

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2015/120280 A1 (GENENTECH INC [US]; HOFFMANN LA ROCHE [CH]) 13 August 2015 (2015-08-13) [0012], [0343] -----	1-32
X	WO 2015/120233 A1 (GENENTECH INC [US]; HOFFMANN LA ROCHE [CH]) 13 August 2015 (2015-08-13) paragraph [0251]; claims 1-116 -----	1-32

## INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2017/014461

Patent document cited in search report	Publication date	Patent family member(s)	Publication date	
US 2010150906	A1	17-06-2010	AR 062065 A1	15-10-2008
			AU 2007275467 A1	24-01-2008
			CA 2657681 A1	24-01-2008
			CL 2007002070 A1	08-02-2008
			CO 6150201 A2	20-04-2010
			CR 10556 A	03-04-2009
			EP 2046833 A2	15-04-2009
			EP 2074145 A2	01-07-2009
			EP 2468770 A1	27-06-2012
			HR P20090221 A2	30-11-2009
			JP 5641737 B2	17-12-2014
			JP 2010505415 A	25-02-2010
			KR 20090039790 A	22-04-2009
			KR 20090088869 A	20-08-2009
			MA 30632 B1	03-08-2009
			NZ 574188 A	25-05-2012
			RU 2009104769 A	27-08-2010
			SG 173385 A1	29-08-2011
			TW 200827372 A	01-07-2008
			US 2009093002 A1	09-04-2009
			US 2010080800 A1	01-04-2010
			US 2010150906 A1	17-06-2010
			US 2012329149 A1	27-12-2012
US 2013164278 A1	27-06-2013			
WO 2008011348 A2	24-01-2008			
EP 1787998	A1	23-05-2007	AU 2005272396 A1	16-02-2006
			CA 2576405 A1	16-02-2006
			CN 101080421 A	28-11-2007
			EP 1787998 A1	23-05-2007
			JP 4820757 B2	24-11-2011
			KR 20070073743 A	10-07-2007
			US 8168188 B1	01-05-2012
			WO 2006016644 A1	16-02-2006
US 2014328856	A1	06-11-2014	AU 2014262890 A1	26-11-2015
			EP 2994160 A1	16-03-2016
			JP 2016518411 A	23-06-2016
			US 2014328856 A1	06-11-2014
			WO 2014182631 A1	13-11-2014
WO 2015120280	A1	13-08-2015	AU 2015213741 A1	25-08-2016
			CA 2938731 A1	13-08-2015
			CN 106456729 A	22-02-2017
			EP 3102230 A1	14-12-2016
			KR 20160113722 A	30-09-2016
			SG 11201606490Y A	29-09-2016
			TW 201613637 A	16-04-2016
			WO 2015120280 A1	13-08-2015
			WO 2015120233	A1
CA 2938466 A1	13-08-2015			
CN 106163548 A	23-11-2016			
EP 3102231 A1	14-12-2016			
KR 20160111039 A	23-09-2016			
SG 11201606316X A	30-08-2016			
TW 201613636 A	16-04-2016			
US 2015246963 A1	03-09-2015			

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2017/014461

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
<p>WO 2015120233 A1 13-08-2015</p> <hr style="border-top: 1px dashed black;"/>			