



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification <sup>7</sup> : <b>A61K 38/48, A61P 7/02, 31/04 // (A61K 38/48, 38:17)</b></p>	<p><b>A1</b></p>	<p>(11) International Publication Number: <b>WO 00/23101</b></p> <p>(43) International Publication Date: 27 April 2000 (27.04.00)</p>
<p>(21) International Application Number: PCT/US99/24409</p> <p>(22) International Filing Date: 15 October 1999 (15.10.99)</p> <p>(30) Priority Data: 60/105,239 22 October 1998 (22.10.98) US</p> <p>(71) Applicant (for all designated States except US): ELI LILLY AND COMPANY [US/US]; Lilly Corporate Center, Indianapolis, IN 46285 (US).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): FISHER, Charles, Jack [US/US]; 12538 Pebblepoint Pass, Carmel, IN 46033 (US). YAN, Sau-Chi, Betty [US/US]; 8131 Menlo Court East Drive, Indianapolis, IN 46240 (US).</p>	<p>(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p><b>Published</b> <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>	
<p>(54) Title: METHODS FOR TREATING SEPSIS</p>		
<p>(57) Abstract</p> <p>The present invention provides a method of treatment for patients with sepsis. The claimed treatment is a combination therapy with protein C and BPI protein. Combining protein C, with its anti-coagulant/anti-inflammatory properties, and BPI, with its bactericidal and endotoxin neutralizing activities, provides an effective, synergistic therapy for sepsis that will reduce or ameliorate the adverse events and improve the clinical outcome of septic patients.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

**METHODS FOR TREATING SEPSIS**

This application claims priority of Provisional Application Serial No. 60/105,239 filed October 22, 1998.

5 This invention relates to medical science particularly the treatment of sepsis with protein C in combination with Bactericidal Permeability Increasing (BPI) Protein.

Protein C is a serine protease and naturally occurring anticoagulant that plays a role in the regulation of hemostasis by inactivating Factors Va and VIIIa in the coagulation cascade. Human protein C circulates as a 2-chain zymogen, but functions at the endothelial and platelet surface following conversion to activated protein C (aPC) by limited proteolysis with thrombin in complex with the cell surface membrane protein, thrombomodulin.

10 In conjunction with other proteins, aPC functions as perhaps the most important down-regulator of blood coagulation resulting in protection against thrombosis. In addition to its anti-coagulation functions, aPC has anti-inflammatory effects through its inhibition of cytokine generation (e.g. TNF and IL-1) and also exerts profibrinolytic properties that facilitate clot lysis. Thus, the protein C enzyme system represents a major physiological mechanism of anti-coagulation, anti-inflammation, and fibrinolysis.

20 Bactericidal permeability-increasing protein (BPI), is a protein isolated from the granules of mammalian polymorphonuclear neutrophils (PMNs). Human BPI has been isolated from PMNs by acid extraction combined with chromatography (Elsbach, 1979, *J. Biol. Chem.* 254:11000; Weiss et al. 1987, *Blood* 69:652), and has been shown to have potent bactericidal activity against a broad spectrum of Gram-negative bacteria. In addition to its bactericidal effect on Gram negative bacteria, BPI is also capable of binding to and neutralizing lipopolysaccharide (LPS) which is also known as endotoxin because of the inflammatory response that it stimulates.

Sepsis, which includes severe sepsis and septic shock, is a systemic inflammatory response to infection or trauma, associated with and mediated by the activation of a number of host defense mechanisms including the cytokine network, leukocytes, and the complement and coagulation/fibrinolysis systems. [Mesters, et al., Blood 88:881-886, 1996]. Disseminated intravascular coagulation [DIC], with widespread deposition of fibrin in the microvasculature of various organs, is an early manifestation of sepsis/septic shock. DIC is an important mediator in the development of the multiple organ failure syndrome and contributes to the poor prognosis of patients with septic shock. [Fourrier, et al., Chest 101:816-823, 1992].

Sepsis may be caused by bacterial (either Gram negative or Gram positive), fungal, viral and other infections as well as by non-infective stimuli such as multiple trauma, severe burns, and organ transplantation.

Although sepsis can follow any bacterial infection, it is often associated with a gram negative infection. Sepsis usually begins with tremor, fever, falling blood pressure, rapid breathing and heart beat, and skin lesions. Within hours or days it can progress to spontaneous clotting in the blood vessels, severe hypotension, multiple organ failure, and death.

Most of the damage comes not from the invading bacteria but from endotoxin. This effect by endotoxin is manifested by its binding to cells such as monocytes/macrophages or endothelial cells, and triggering them to produce various mediators such as tumor necrosis factor-alpha (TNF- ), and various interleukins (IL-1, IL-6, and IL-8). Production of excessive TNF- , IL-1, IL-6, and IL-8 can elicit septic shock.

There have been numerous recent attempts to treat sepsis in humans, for the most part using agents that block inflammatory mediators associated with the pathophysiology of this disease. However, clinical studies with a variety of agents that block inflammatory mediators have been

unsuccessful [reviewed in Natanson, et al., Ann. Intern. Med  
120:771-783, 1994; Gibaldi, Pharmacotherapy 13:302-308,  
1993]. Since many of the mediators involved in inflammation  
are compensatory responses, and therefore have salutary  
5 effects, some investigators have suggested that blocking  
their action may not be appropriate [e.g., Parrillo, N.  
Engl. J. Med. 328:1471-1477, 1993].

Several encouraging studies using protein C in various  
animal models of sepsis have been reported. A study in a  
10 baboon sepsis model by Taylor, et al., [J. Clin. Invest.  
79:918-25, 1987], used plasma-derived human activated  
protein C. The animals were treated prophylactically (i.e.,  
the aPC was given at the start of the two hour infusion of  
the LD<sub>100</sub> E. coli). Five out of five animals survived 7 days  
15 and were considered permanent survivors to the experimental  
protocol. In control animals receiving an identical  
infusion of E. coli, five out of five animals died in 24 to  
32 hours. In addition, plasma-derived human protein C  
zymogen has been used as a successful adjunct to aggressive  
20 conventional therapy in the management of human patients  
with purpura fulminans in bacterial sepsis (Gerson, et al.,  
Pediatrics 91:418-422, 1993; Smith, et al., Thromb.  
Haemost, PS1709, p419, 1997; Rintala, et al., Lancet  
347:1767, 1996; Rivard, et al., J. Pediatr. 126:646-652,  
25 1995).

Recombinant BPI protein has been shown to neutralize  
lethal and sublethal effects of endotoxin administered to  
mice, rats, and rabbits (Fisher, et al., Crit. Care Med.,  
22(4): 553-558, 1994). Because of this ability to  
30 neutralize endotoxin and its Gram-negative bacteriacidal  
activity, BPI can be utilized for the treatment of human  
patients suffering from diseases caused by gram-negative  
bacteria, including bacteremia, endotoxemia, and sepsis.

The present invention is the first to describe the  
35 combination of aPC with BPI in the treatment of sepsis. The  
combination of aPC and BPI results in a synergy that allows  
the reduction of the dosages of both aPC and BPI and an

improvement of clinical outcome of the patient being treated. The reduction of the dosages of the agents in combination therapy in turn results in reduced side effects that may occur with either agent. Therefore, combining aPC, with its anti-coagulant/anti-inflammatory properties, and BPI, with its bactericidal and endotoxin neutralizing activities will provide an effective synergistic therapy for sepsis that will reduce or ameliorate the adverse events and improve the clinical outcome of septic patients.

10

The present invention provides a method of treating a patient suffering from sepsis which comprises administering to said patient a pharmaceutically effective amount of protein C in combination with bactericidal permeability-increasing (BPI) protein.

15

The present invention further provides a method of treating sepsis in a patient in need thereof, which comprises administering to said patient a pharmaceutically effective amount of BPI protein and activated protein C such that an activated protein C plasma level of about 2 ng/ml to about 300 ng/ml is achieved.

20

For purposes of the present invention, as disclosed and claimed herein, the following terms are as defined below.

25

Protein C refers to a vitamin K dependent serine protease with anticoagulant, anti-inflammatory, and profibrinolytic properties which includes, but is not limited to, plasma derived and recombinant produced protein C. Protein C includes and is preferably human protein C although protein C may also include other species or derivatives having protein C proteolytic, amidolytic, esterolytic, and biological (anticoagulant, pro-fibrinolytic, and anti-inflammatory) activities. Examples of protein C derivatives are described by Gerlitz, et al., U.S. Patent No. 5,453,373, and Foster, et al., U.S. Patent No. 5,516,650, the entire teachings of which are hereby included by reference.

30  
35

Zymogen - an enzymatically inactive precursor of a proteolytic enzyme. Protein C zymogen, as used herein, refers to secreted, inactive forms, whether one chain or two chains, of protein C.

5 Activated protein C or aPC refers to protein C zymogen which has been converted by limited proteolysis to its activated form. aPC includes and is preferably human protein C although aPC may also include other species or derivatives having protein C proteolytic, amidolytic,  
10 esterolytic, and biological (anticoagulant or pro-fibrinolytic) activities. Examples of protein C derivatives are noted above in the description of protein C.

r-hPC - recombinant human protein C zymogen.

r-aPC - recombinant activated protein C, preferably  
15 produced by activating r-hPC *in vitro* or by direct secretion of the activated form of protein C from procaryotic cells, eukaryotic cells, and transgenic animals or plants, including, for example, secretion from human kidney 293 cells as a zymogen then purified and activated by techniques  
20 well known to the skilled artisan and demonstrated in Yan, U.S. Patent No. 4,981,952, and Cottingham, WO97/20043, the entire teachings of which are herein incorporated by reference.

Plasma derived activated protein C - activated protein  
25 C produced by activating plasma protein C as described in Eibl, U.S. Patent No. 5,478,558, the entire teaching of which is herein incorporated by reference.

Continuous infusion - continuing substantially  
uninterrupted the introduction of a solution into a vein for  
30 a specified period of time.

Bolus injection - the injection of a drug in a defined quantity (called a bolus) over a period of time up to about 120 minutes.

Suitable for administration - a lyophilized formulation  
35 or solution that is appropriate to be given as a therapeutic agent.

Unit dosage form - refers to physically discrete units suitable as unitary dosages for human subjects, each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect, in association with a suitable pharmaceutical excipient.

Pharmaceutically effective amount - represents an amount of protein C of the present invention that is capable of treating sepsis in humans. The particular dose of protein C administered according to this invention will, of course, be determined by the attending physician evaluating the particular circumstances surrounding the case.

BPI protein - includes naturally and recombinantly produced bactericidal permeability increasing (BPI) protein; natural, synthetic, and recombinant biologically active polypeptide fragments of BPI protein; biologically active polypeptide variants of BPI protein or fragments thereof, including hybrid fusion proteins and dimers; biologically active polypeptide analogs of BPI protein or fragments or variants thereof, including cysteine-substituted analogs; and BPI-derived peptides. The complete amino acid sequence of human BPI, as well as the nucleotide sequence of DNA encoding BPI have been elucidated by Gray et al., 1989, *J. Biol. Chem* 264:9505. Recombinant genes encoding and methods for expression of BPI proteins, including BPI holoprotein and fragments of BPI are disclosed in US Patent No. 5,198,541, herein incorporated by reference.

The present invention relates to the treatment of sepsis with protein C in combination with BPI protein. The combination of protein C and BPI results in a synergy that allows the reduction of the dosages of both protein C and BPI and an improvement of clinical outcome of the patient being treated. The reduction of the dosages of the agents in combination therapy in turn results in reduced side effects that may occur with either agent. Therefore, combining protein C, with its anti-coagulant/anti-inflammatory properties, and BPI, with its bactericidal and



endotoxin neutralizing activities will provide an effective synergistic therapy for sepsis that will reduce or ameliorate the adverse events and improve the clinical outcome of septic patients.

5           The protein C administered according to this invention may be generated and/or isolated by any means known in the art or as described in U.S. Patent No. 4,981,952, and U.S. Patent No. 5,550,036, herein incorporated by reference. For example, the invention provides a method for producing and  
10       secreting full-length, soluble protein C, or biologically active polypeptide variants of protein C from a cell which comprises (a) constructing a vector comprising DNA encoding protein C; (b) transfecting the cell with the vector; and (c) culturing the cell so transfected in culture medium  
15       under conditions such that full length soluble protein C or biologically active polypeptide variants of protein C, is secreted. Further, the cell is a eukaryotic cell, e.g. mammalian cell such as Syrian hamster AV12 cell, human embryonic 293 cell, or Baby Hamster Kidney cell.

20           The protein C used in such combination can be formulated according to known methods to prepare pharmaceutically useful compositions. For example, a desired formulation would be one that is a stable lyophilized product of high purity comprising a bulking  
25       agent such as sucrose, a salt such as sodium chloride, a buffer such as sodium citrate and protein C or aPC.

          The protein C will be administered parenterally to ensure its delivery into the bloodstream in an effective form by injecting the appropriate dose as continuous  
30       infusion for about 1 hour to about 240 hours.

          In conjunction with treatment with BPI protein, the amount of protein C administered will be from about 5.0  $\mu\text{g}/\text{kg}/\text{hr}$  to about 250  $\mu\text{g}/\text{kg}/\text{hr}$ . Preferably, the protein C administered in combination with BPI protein will be  
35       activated protein C. The aPC administered will be from about 1.0  $\mu\text{g}/\text{kg}/\text{hr}$  to about 50  $\mu\text{g}/\text{kg}/\text{hr}$ . More preferably the amount of aPC administered will be about 1.0  $\mu\text{g}/\text{kg}/\text{hr}$  to

about 40  $\mu\text{g}/\text{kg}/\text{hr}$ . While more preferably the amount of aPC administered will be about 1.0  $\mu\text{g}/\text{kg}/\text{hr}$  to 35  $\mu\text{g}/\text{kg}/\text{hr}$ . Even more preferably the amount of aPC administered will be about 5.0  $\mu\text{g}/\text{kg}/\text{hr}$  to 30  $\mu\text{g}/\text{kg}/\text{hr}$ . Yet even more preferably  
5 the amount of aPC administered will be about 15  $\mu\text{g}/\text{kg}/\text{hr}$  to 30  $\mu\text{g}/\text{kg}/\text{hr}$ . Still even more preferably the amount of aPC administered will be about 20  $\mu\text{g}/\text{kg}/\text{hr}$  to 30  $\mu\text{g}/\text{kg}/\text{hr}$ . The most preferable amount of aPC administered will be about 24  $\mu\text{g}/\text{kg}/\text{hr}$ . The appropriate dose of aPC administered with  
10 BPI protein results in either an improved efficacy or reduction in dose of either agent or both.

The plasma ranges obtained from the amount of aPC administered will be about 2 ng/ml to about 300 ng/ml. The preferred plasma ranges are from about 2 ng/ml to 200 ng/ml.  
15 Most preferably, plasma ranges are from about 30 ng/ml to about 150 ng/ml and still more preferably about 100 ng/ml.

Alternatively, the aPC will be administered by injecting one third of the appropriate dose per hour as a bolus injection followed by the remaining two thirds of the  
20 hourly dose as continuous infusion for one hour followed by continuous infusion of the appropriate dose for twenty-three hours which results in the appropriate dose administered over 24 hours. In addition, the bolus injection will be administered via an intravenous bag drip pump or syringe  
25 pump at about 2 times the normal rate for about 10 to 20 minutes followed by about 1.5 times the normal rate for about 40 to 50 minutes. The normal rate i.e. that rate which has been determined to administer the appropriate dose level of the therapeutic agent per time period, is then  
30 continued for up to 240 hours. over 24 hours.

BPI protein suitable for use under the present invention includes, but is not limited to, naturally and recombinantly produced BPI protein, for example, a  
35 recombinant BPI holoprotein as described in Gray *et al.* (1989) and U.S. Patent No. 5,733,872, herein incorporated by reference; natural, synthetic, and recombinant biologically

active polypeptide fragments of BPI protein, for example, as described in Ooi *et al.*, *J. Exp. Med.*, 174:649 (1991) and Gazzano-Santoro *et al.*, *Infect. Immun.* 60:4754-4761 (1992); biologically active polypeptide variants of BPI protein or  
5 fragments thereof, including hybrid fusion proteins and dimers; biologically active polypeptide analogs of BPI protein or fragments or variants thereof, including cysteine-substituted analogs; and BPI-derived peptides, examples of which are described in U.S. Patent Nos.  
10 5,733,872, 5,627,262, 5,753,620, 5,607,916 and 5,756,464, herein incorporated by reference.

Preferably, the BPI protein of the present invention includes biologically active molecules that have the same or similar amino acid sequence as a natural human BPI  
15 holoprotein. Nonlimiting examples of such BPI proteins are the 25 Kd N-terminal fragment of natural human BPI protein, described in Ooi *et al.*, (1991), and the recombinant expression product of DNA encoding N-terminal amino acids from 1 to about 193 or 199 of natural human BPI, described  
20 in Gazzano-Santoro *et al.*, (1992).

The BPI protein administered according to this invention may be generated and/or isolated by any means known in the art or as described in U.S. Patent No. 5,308,834, herein incorporated by reference. For example,  
25 the invention provides a method for producing and secreting full-length, soluble BPI holoprotein, biologically active polypeptide fragments, or biologically active polypeptide variants of BPI protein or fragments thereof from a cell which comprises (a) constructing a vector comprising DNA  
30 encoding BPI; (b) transfecting the cell with the vector; and (c) culturing the cell so transfected in culture medium under conditions such that full length soluble BPI protein, biologically active polypeptide fragments, or biologically active polypeptide variants of BPI protein or fragments  
35 thereof, is secreted. Further, the cell is a eukaryotic cell, e.g. mammalian cell such as Syrian hamster AV12 cell, human embryonic 293 cell, or Baby Hamster Kidney cell.

Alternatively, the cell is a prokaryotic cell, e.g. a yeast cell or a bacterial cell.

The phrase "in combination with" refers to the administration of BPI protein with protein C either  
5 simultaneously, sequentially or a combination thereof. The BPI protein utilized and the appropriate dose level is known in the art and described in U.S. Patent No. 5,756,464, herein incorporated by reference. A skilled artisan recognizes the appropriate dose level to use to achieve a  
10 pharmaceutically effective amount for treating sepsis. Pharmaceutically effective compositions comprising BPI protein may be administered systemically or topically. Systemic routes of administration include, intravenous, intramuscular or subcutaneous injection (including into a  
15 depot for long-term release), intraocular and retrobulbar, intrathecal, intraperitoneal (e.g. by intraperitoneal lavage), intrapulmonary using aerosolized or nebulized drug, or transdermal. The preferred route is intravenous administration. When given parenterally, BPI protein  
20 compositions are generally injected in doses ranging from about 0.04 ug/kg/hr to about 4 mg/kg/hr. Preferably, the BPI protein is administered at about 4 ug/kg/hr to about 420 ug/kg/hr. More preferably the BPI protein is administered at about 50 ug/kg/hr to about 300 ug/kg/hr.  
25 Even more preferably the BPI protein is administered at about 100 ug/kg/hr to about 200 ug/kg/hr. The treatment may continue by continuous infusion or intermittent injection or infusion, at the same, reduced or increased dose per day for, e.g. 24 hours to 240 hours, and additionally as  
30 determined by the treating physician. BPI protein is preferably administered intravenously by an initial bolus injection followed by a continuous infusion. A preferred dosing regimen is about 0.1 mg/kg to about 10 mg/kg intravenous bolus of BPI protein followed by intravenous  
35 infusion at about 4 ug/kg/hr to about 420 ug/kg/hr, continuing for up to 10 days. Those skilled in the art can readily optimize pharmaceutically effective dosages and

administration regimens for therapeutic compositions comprising BPI protein, as determined by good medical practice and the clinical condition of the individual patient.

5           The combination of the endotoxin neutralization and the Gram-negative bactericidal activity of BPI protein with the anti-coagulation and anti-inflammation activities of aPC results in enhanced efficacy in treating sepsis. The synergy results in the ability to reduce the dosages of the  
10 agents in combination therapy.

The following examples are provided merely to further illustrate the present invention. The scope of the invention shall not be construed as merely consisting of the  
15 following examples.

#### **Preparation 1**

##### **Preparation of Human Protein C**

Recombinant human protein C (r-hPC) was produced in Human Kidney 293 cells by techniques well known to the  
20 skilled artisan such as those set forth in Yan, U.S. Patent No. 4,981,952, the entire teaching of which is herein incorporated by reference. The gene encoding human protein C is disclosed and claimed in Bang, et al., U.S. Patent No. 4,775,624, the entire teaching of which is incorporated  
25 herein by reference. The plasmid used to express human protein C in 293 cells was plasmid pLPC which is disclosed in Bang, et al., U.S. Patent No. 4,992,373, the entire teaching of which is incorporated herein by reference. The construction of plasmid pLPC is also described in European  
30 Patent Publication No. 0 445 939, and in Grinnell, et al., 1987, Bio/Technology 5:1189-1192, the teachings of which are also incorporated herein by reference. Briefly, the plasmid was transfected into 293 cells, then stable transformants were identified, subcultured and grown in serum-free media.  
35 After fermentation, cell-free medium was obtained by microfiltration.

The human protein C was separated from the culture fluid by an adaptation of the techniques of Yan, U.S. Patent No. 4,981,952. The clarified medium was made 4 mM in EDTA before it was absorbed to an anion exchange resin (Fast-Flow Q, Pharmacia). After washing with 4 column volumes of 20 mM Tris, 200 mM NaCl, pH 7.4 and 2 column volumes of 20 mM Tris, 150 mM NaCl, pH 7.4, the bound recombinant human protein C zymogen was eluted with 20 mM Tris, 150 mM NaCl, 10 mM CaCl<sub>2</sub>, pH 7.4. The eluted protein was greater than 95% pure after elution as judged by SDS-polyacrylamide gel electrophoresis.

Further purification of the protein was accomplished by making the protein 3 M in NaCl followed by adsorption to a hydrophobic interaction resin (Toyopearl Phenyl 650 M, TosoHaas) equilibrated in 20 mM Tris, 3 M NaCl, 10 mM CaCl<sub>2</sub>, pH 7.4. After washing with 2 column volumes of equilibration buffer without CaCl<sub>2</sub>, the recombinant human protein C was eluted with 20 mM Tris, pH 7.4.

The eluted protein was prepared for activation by removal of residual calcium. The recombinant human protein C was passed over a metal affinity column (Chelex-100, Bio-Rad) to remove calcium and again bound to an anion exchanger (Fast Flow Q, Pharmacia). Both of these columns were arranged in series and equilibrated in 20 mM Tris, 150 mM NaCl, 5 mM EDTA, pH 7.4. Following loading of the protein, the Chelex-100 column was washed with one column volume of the same buffer before disconnecting it from the series. The anion exchange column was washed with 3 column volumes of equilibration buffer before eluting the protein with 0.4 M NaCl, 20 mM Tris-acetate, pH 6.5. Protein concentrations of recombinant human protein C and recombinant activated protein C solutions were measured by UV 280 nm extinction  $E^{0.1\%} = 1.81$  or  $1.85$ , respectively.

## Preparation 2

### Activation of Recombinant Human Protein C

Bovine thrombin was coupled to Activated CH-Sepharose 4B (Pharmacia) in the presence of 50 mM HEPES, pH 7.5 at  
5 4°C. The coupling reaction was done on resin already packed into a column using approximately 5000 units thrombin/mL resin. The thrombin solution was circulated through the column for approximately 3 hours before adding 2-amino-ethanol (MEA) to a concentration of 0.6 mL/L of circulating  
10 solution. The MEA-containing solution was circulated for an additional 10-12 hours to assure complete blockage of the unreacted amines on the resin. Following blocking, the thrombin-coupled resin was washed with 10 column volumes of 1 M NaCl, 20 mM Tris, pH 6.5 to remove all non-specifically  
15 bound protein, and was used in activation reactions after equilibrating in activation buffer.

Purified r-hPC was made 5 mM in EDTA (to chelate any residual calcium) and diluted to a concentration of 2 mg/mL with 20 mM Tris, pH 7.4 or 20 mM Tris-acetate, pH 6.5. This  
20 material was passed through a thrombin column equilibrated at 37°C with 50 mM NaCl and either 20 mM Tris pH 7.4 or 20 mM Tris-acetate pH 6.5. The flow rate was adjusted to allow for approximately 20 min. of contact time between the r-hPC and thrombin resin. The effluent was collected and  
25 immediately assayed for amidolytic activity. If the material did not have a specific activity (amidolytic) comparable to an established standard of aPC, it was recycled over the thrombin column to activate the r-hPC to completion. This was followed by 1:1 dilution of the  
30 material with 20 mM buffer as above, with a pH of either 7.4 or 6.5 to keep the aPC at lower concentrations while it awaited the next processing step.

Removal of leached thrombin from the aPC material was accomplished by binding the aPC to an anion exchange resin  
35 (Fast Flow Q, Pharmacia) equilibrated in activation buffer (either 20 mM Tris, pH 7.4 or 20 mM Tris-acetate, pH 6.5) with 150 mM NaCl. Thrombin does not interact with the anion

exchange resin under these conditions, but passes through the column into the sample application effluent. Once the aPC is loaded onto the column, a 2-6 column volume wash with 20 mM equilibration buffer is done before eluting the bound  
5 aPC with a step elution using 0.4 M NaCl in either 5 mM Tris-acetate, pH 6.5 or 20 mM Tris, pH 7.4. Higher volume washes of the column facilitated more complete removal of the dodecapeptide. The material eluted from this column was stored either in a frozen solution (-20°C) or as a  
10 lyophilized powder.

The anticoagulant activity of activated protein C was determined by measuring the prolongation of the clotting time in the activated partial thromboplastin time (APTT) clotting assay. A standard curve was prepared in dilution  
15 buffer (1 mg/mL radioimmunoassay grade bovine serum albumin [BSA], 20 mM Tris, pH 7.4, 150 mM NaCl, 0.02% NaN<sub>3</sub>) ranging in protein C concentration from 125-1000 ng/mL, while samples were prepared at several dilutions in this concentration range. To each sample cuvette, 50 μL of cold  
20 horse plasma and 50 μL of reconstituted activated partial thromboplastin time reagent (APTT Reagent, Sigma) were added and incubated at 37°C for 5 min. After incubation, 50 μL of the appropriate samples or standards were added to each cuvette. Dilution buffer was used in place of sample or  
25 standard to determine basal clotting time. The timer of the fibrometer (CoA Screener Hemostasis Analyzer, American Labor) was started immediately after the addition of 50 μL 37°C 30 mM CaCl<sub>2</sub> to each sample or standard. Activated protein C concentration in samples are calculated from the  
30 linear regression equation of the standard curve. Clotting times reported here are the average of a minimum of three replicates, including standard curve samples.

The above descriptions enable one with appropriate skill in the art to prepare aPC for utilization in  
35 combination therapy with BPI protein for the treatment of sepsis.



### Preparation 3

#### Expression of BPI protein

In order to produce BPI protein and /or BPI protein variants in mammalian cells, the cDNA sequences is inserted into a suitable plasmid vector as described in US Patent No. 5,171,739, herein incorporated by reference. A suitable vector for such an application is pSV-1, which contains the origin of replication and early and late promoters of SV40, followed by multiple insert coning sites, followed by termination sequences from the hepatitis B surface antigen gene. Also contained in the plasmid are an origin of bacterial DNA replication, and the genes encoding ampicillin resistance and dihydrofolate reductase. Similar vectors have been used to express other foreign genes [McGrogan, et al. Biotechnology, 6: 172-177]. Vector DNA is prepared for acceptance of BPI protein cDNA sequences by digestion with HindIII and Bam HI, and dephosphorylation with alkaline phosphatase.

A BPI protein cDNA containing insert is prepared for insertion into pSV-1 by inserting encoding full-length BPI protein prepared by digestion of the parent plasmid with appropriate restriction enzymes, for example EcoRI and Bgl II, yielding two DNA fragments containing portions of the BPI protein coding sequence. These two fragments are ligated together into prepared SV-1, and the recombinant clones obtained are screened by restriction enzyme digestion for the presence of the two inserts in the proper orientation.

The construct is verified by restriction digest analysis, and then prepared in amounts sufficient for transfection into CHO cell line DUXB11 cells. Transfection is performed using lipofectin, and the resulting transformed cells are selected in the presence of increasing amounts of methotrexate using standard protocols.

Supernatants from either transfected pools or clones derived from pools are assayed for the presence of endotoxin binding activity by inhibition of TNr release. BPI protein

is purified from the selected supernatant by standard procedures known in the art.

#### Preparation 4

##### Formulation of Activated Protein C

5 A stable lyophilized formulation of activated protein C was prepared by a process which comprises lyophilizing a solution comprising about 2.5 mg/mL activated protein C, about 15 mg/mL sucrose, about 20 mg/mL NaCl, and a sodium citrate buffer having a pH greater than 5.5 but less than  
10 6.5. Additionally, the stable lyophilized formulation of activated protein C comprises lyophilizing a solution comprising about 5 mg/mL activated protein C, about 30 mg/mL sucrose, about 38 mg/mL NaCl, and a citrate buffer having a pH greater than 5.5 but less than 6.5.

15 The ratio of aPC:salt:bulking agent (w:w:w) is an important factor in a formulation suitable for the freeze drying process. The ratio varies depending on the concentration of aPC, salt selection and concentration and bulking agent selection and concentration. Particularly, a  
20 ratio of about 1 part activated protein C to about 7.6 parts salt to about 6 parts bulking agent is preferred.

A unit dosage formulation of activated protein C suitable for administration by continuous infusion was prepared by mixing activated protein C, NaCl, sucrose, and  
25 sodium citrate buffer. After mixing, 4 mL of the solution was transferred to a unit dosage receptacle and lyophilized. The unit dosage receptacle containing about 5 mg to about 20 mg of activated protein C, suitable for administering a dosage of about 0.01 mg/kg/hr to about 0.05 mg/kg/hr to  
30 patients in need thereof, was sealed and stored until use.

#### Preparation 5

##### Pharmaceutical Composition of BPI Protein

A pharmaceutical composition of BPI protein is prepared by a process which comprises the BPI protein at a  
35 concentration of 1 mg/ml in citrate buffered saline (5 or 20 mM citrate, 150 mM NaCl, pH 5.0) comprising 0.1% by weight of poloxamer 188 (Pluronic F-68, BASF Wyandotte, Parsippany,

N.J.) and 0.002% by weight of polysorbate 80 (Tween 80, ICI Americas, Inc., Wilmington, Del.). Another pharmaceutical composition containing BPI protein comprises the BPI protein at a concentration of 2 mg/ml, in 5 mM citrate, 150 mM NaCl, 5 0.2% poloxamer 188 and 0.002% polysorbate 80. Such combinations are described in U.S. Patent No. 5,756,464.

### Example 1

#### Administering Activated Protein C in Combination with BPI protein in a Double-blinded Placebo-controlled 10 Trial in Human Patients With Sepsis

This protocol is a double-blinded placebo-controlled trial in patients with severe sepsis. Patients are treated with placebo, aPC alone, BPI protein alone, or a combination therapy of aPC and BPI protein. BPI protein is administered 15 by continuous infusion at about 100  $\mu\text{g}/\text{kg}/\text{hr}$  to about 200  $\mu\text{g}/\text{kg}/\text{hr}$  for about 48 hours. aPC is administered simultaneously by a continuous infusion of about 1  $\mu\text{g}/\text{kg}/\text{hr}$  to about 50  $\mu\text{g}/\text{kg}/\text{hr}$  for about 96 hours.

Entry criteria includes three of the four commonly 20 accepted criteria for sepsis (heart rate, respiratory effort, increased/decreased temperature, increase/decrease white blood cell count). The patients also demonstrate some degree of organ dysfunction defined as either shock, decreased urine output, or hypoxemia.

25 The primary endpoints of this study are: safety as a function of dose and dose duration; and, comparing treatment with aPC alone or BPI protein alone to the ability of aPC in combination with BPI protein to correct coagulopathy as a function of dose and dose duration. A 28-day all cause 30 mortality is the end-point in patients receiving placebo vs. patients receiving aPC in combination with BPI or either agent alone.

Combination therapy with aPC and BPI protein results in a synergy that is safer and more efficacious and reduces the 35 dosages of both aPC and BPI protein necessary to treat sepsis.

-18-

We Claim:

1. A method of treating a patient suffering from sepsis which comprises, administering to said patient a pharmaceutically effective amount of protein C in combination with bactericidal/permeability-increasing (BPI) protein.
2. The method of Claim 1, wherein said patient is human.
3. The method of Claim 1 wherein the protein C is human protein C zymogen.
4. The method of Claim 1 wherein the protein C is human activated protein C.
5. The method according to Claim 4, wherein the amount of human activated protein C is about 1  $\mu\text{g}/\text{kg}/\text{hr}$  to about 50  $\mu\text{g}/\text{kg}/\text{hr}$ .
6. The method of Claim 5, wherein the human activated protein C is administered by continuous infusion for about 1 to about 240 hours.
7. The method of Claim 1, wherein the BPI protein is administered by bolus injection.
8. The method of Claim 7, wherein the amount of BPI protein is about 0.1 mg/kg to about 10 mg/kg of body weight.

9. The method of Claim 1, wherein the BPI protein is administered by continuous infusion for about 1 to about 240 hours.

10. The method of Claim 9, wherein the amount of BPI protein is about 4  $\mu\text{g}/\text{kg}/\text{hr}$  to about 420  $\mu\text{g}/\text{kg}/\text{hr}$ .

11. A method of treating sepsis in a patient in need thereof, which comprises administering to said patient a pharmaceutically effective amount of BPI protein and activated protein C such that an activated protein C plasma level of about 2 ng/ml to about 300 ng/ml is achieved.

12. The method of Claim 11 wherein the activated protein C is administered in a bolus injection.

13. The method of Claim 11 wherein the activated protein C is administered by continuous infusion for about 1 to about 240 hours.

14. The method of Claim 11 wherein the activated protein C is administered first as a bolus then as a continuous infusion.

15. The method of Claim 14 wherein one third of the activated protein C required to achieve activated protein C plasma levels in the range of about 2 ng/ml to about 300 ng/ml is administered in a bolus injection followed by continuous infusion of the remaining two thirds of the activated protein C.