

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 2012201801 B2**

(54) Title
Sulfoperoxycarboxylic acids, their preparation and methods of use as bleaching and antimicrobial agents

(51) International Patent Classification(s)
C07C 309/12 (2006.01) **C07C 309/08** (2006.01)
A23L 3/3535 (2006.01) **C07D 303/16** (2006.01)
C07C 309/05 (2006.01) **C11D 3/48** (2006.01)

(21) Application No: **2012201801** (22) Date of Filing: **2012.03.27**

(43) Publication Date: **2012.04.19**

(43) Publication Journal Date: **2012.04.19**

(44) Accepted Journal Date: **2014.05.01**

(62) Divisional of:
2009230713

(71) Applicant(s)
Ecolab Inc.

(72) Inventor(s)
McSherry, David D.;Everts, Frank;Lange, Steven J.;Lascotte, Keith G.;Staub, Richard K.;Li, Junzhong


(74) Agent / Attorney
Spruson & Ferguson, L 35 St Martins Tower 31 Market St, Sydney, NSW, 2000

(56) Related Art
WO 1998/003513 A1
US 6,183,807 B1

SULFOPEROXYCARBOXYLIC ACIDS, THEIR PREPARATION AND METHODS OF USE AS BLEACHING AND ANTIMICROBIAL AGENTS

Abstract

Figure 1



The present invention relates to novel sulfoperoxydicarboxylic acid compounds, and methods for making and using them. The sulfoperoxydicarboxylic compounds of the invention are storage stable, water soluble and have low to no odor. Further, the compounds of the present invention can be formed from non-petroleum based renewable materials. The compounds of the present invention can be used as antimicrobials, and bleaching agents. The compounds of the present invention are also suitable for use as coupling agents.

AUSTRALIA

PATENTS ACT 1990

COMPLETE SPECIFICATION

FOR A STANDARD PATENT

Name and Address of Applicant :	Ecolab Inc., of 370 N. Wabasha Street, St. Paul, Minnesota, 5102, United States of America
Actual Inventor(s):	MCSHERRY, David D.; EVERTS, Frank; LANGE, Steven J.; LASCOTTE, Keith G.; STAUB, Richard K.; LI, Junzhong
Address for Service:	Spruson & Ferguson St Martins Tower Level 35 31 Market Street Sydney NSW 2000 (CCN 3710000177)
Invention Title:	Sulfoperoxydicarboxylic acids, their preparation and methods of use as bleaching and antimicrobial agents

The following statement is a full description of this invention, including the best method of performing it known to me/us:

SULFOPEROXYCARBOXYLIC ACIDS, THEIR PREPARATION AND METHODS OF USE AS BLEACHING AND ANTIMICROBIAL AGENTS

FIELD OF THE INVENTION

The present invention relates to novel sulfoperoxy-carboxylic acid compounds, compositions, and methods of making and using these compounds.


BACKGROUND

Peroxy-carboxylic acids are known for use as antimicrobials and bleaching agents. However, conventional peroxy-carboxylic acids have inherent disadvantages of limited storage stability, and water solubility. Further, most peroxy-carboxylic acids have an unpleasant odor. Thus, a need exists for storage stable, low or no odor, water soluble peroxy-carboxylic acid compounds and compositions that also possess antimicrobial and bleaching properties.

Summary

According to a first aspect of the present invention there is provided an aqueous, antimicrobial bleaching composition comprising:

- a) a compound according to Formula I:



(Formula I)

wherein:

- R₁ is a sulfonate substituted C₁ alkyl group;
- R₂ is a substituted or unsubstituted C_n alkyl group;
- X is hydrogen, a cationic group, or an ester forming moiety;
- n is 1 to 10;
- or salts or esters thereof,
- b) a C₁ to C₄ carboxylic acid;
- c) a C₅ to C₁₁ carboxylic acid; and
- d) an oxidizing agent wherein the pH of the composition is about 7 to about 14

According to a second aspect of the present invention there is provided a method for treating laundry, the method comprising:

- (a) applying the composition according to the first aspect to the laundry;
- (b) draining the composition from the laundry.

In some aspects, the present invention relates to novel sulfoperoxy-carboxylic acids, and methods for making them. The compounds of the invention are storage stable, have low or no-odor, and are water soluble. Further, the compounds of the present invention can be derived from non-petroleum based, renewable oils.

In some aspects, the present invention provides methods for using the compounds of the present invention as bleaching and/or antimicrobial agents. In some aspects, the present invention provides methods for using the compounds of the invention as coupling agents. In some aspects, the present invention provides methods for using the compounds of the present invention as low foaming bleach hydrotopes for tunnel washers, and for side loading washing machines.

In some embodiments, the compounds and compositions of the present invention are suitable for use as low temperature bleaches, e.g., at about 40 degrees Celsius. In some embodiments, the compounds of the present invention are suitable for use as pH optimized peroxygen bleaches, in combination with alkaline detergents. In

some embodiments, the present invention includes a method for using the compounds and compositions of the present invention as color safe, textile tolerant bleaches for textiles, e.g., wools and cotton.

5 BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a graphical depiction of the stability profile of peroxyoctanoic acid over time when contacted with different test solutions.

Figure 2 is a graphical depiction of the stability of an exemplary composition of the present invention over time at an elevated temperature.

10 Figure 3 is a graphical depiction of the ability of selected compositions of the present invention to stabilize percarboxylic acids over time.

Figure 4 is a graphical depiction of the bleaching performance of compositions of the present invention compared to commercially available bleaching agents.

15 Figure 5 is a graphical depiction of the stability profile of peroxyoctanoic acid in combination with exemplary compositions of the present invention.

Figure 6 is a graphical depiction of the coupling capabilities of a selected composition of the present invention.

DETAILED DESCRIPTION

20 The present invention relates to sulfoperoxydicarboxylic acids of Formula I, and methods of making and using them. Unlike conventional peroxycarboxylic acids, the sulfoperoxydicarboxylic acids of the present invention are low-odor, water soluble, and storage stable. The compounds of the present invention can be used as a pure solid powder, or blended with additional functional ingredients, for example, chelators,
25 buffers, or other cleaning agents. They can also be incorporated into liquid formulas. The compounds and compositions of the present invention have many uses including, but not limited to, antimicrobials, bleaches, and coupling agents.

So that the invention maybe more readily understood, certain terms are first defined.

As used herein, "weight percent," "wt-%," "percent by weight," "% by weight," and variations thereof refer to the concentration of a substance as the weight of that substance divided by the total weight of the composition and multiplied by 100. It is understood that, as used here, "percent," "%," and the like are intended to be
5 synonymous with "weight percent," "wt-%," etc.

As used herein, the term "about" refers to variation in the numerical quantity that can occur, for example, through typical measuring and liquid handling procedures used for making concentrates or use solutions in the real world; through inadvertent error in these procedures; through differences in the manufacture, source, or purity of
10 the ingredients used to make the compositions or carry out the methods; and the like. The term "about" also encompasses amounts that differ due to different equilibrium conditions for a composition resulting from a particular initial mixture. Whether or not modified by the term "about", the claims include equivalents to the quantities.

It should be noted that, as used in this specification and the appended claims, the
15 singular forms "a," "an," and "the" include plural referents unless the content clearly dictates otherwise. Thus, for example, reference to a composition containing "a compound" includes a composition having two or more compounds. It should also be noted that the term "or" is generally employed in its sense including "and/or" unless the content clearly dictates otherwise.

As used herein, the phrases "objectionable odor," "offensive odor," or
20 "malodor," refer to a sharp, pungent, or acrid odor or atmospheric environment from which a typical person withdraws if they are able to. Hedonic tone provides a measure of the degree to which an odor is pleasant or unpleasant. An "objectionable odor," "offensive odor," or "malodor" has an hedonic tone rating it as unpleasant as or more
25 unpleasant than a solution of 5 wt-% acetic acid, propionic acid, butyric acid, or mixtures thereof.

As used herein, the term "microorganism" refers to any noncellular or unicellular (including colonial) organism. Microorganisms include all prokaryotes. Microorganisms include bacteria (including cyanobacteria), spores, lichens, fungi,

protozoa, virinos, viroids, viruses, phages, and some algae. As used herein, the term "microbe" is synonymous with microorganism.

As used herein, the phrase "food product" includes any food substance that might require treatment with an antimicrobial agent or composition and that is edible
5 with or without further preparation. Food products include meat (e.g. red meat and pork), seafood, poultry, produce (e.g., fruits and vegetables), eggs, living eggs, egg products, ready to eat food, wheat, seeds, roots, tubers, leaves, stems, corns, flowers, sprouts, seasonings, or a combination thereof. The term "produce" refers to food
10 products such as fruits and vegetables and plants or plant-derived materials that are typically sold uncooked and, often, unpackaged, and that can sometimes be eaten raw.

As used herein, the phrase "plant" or "plant product" includes any plant substance or plant-derived substance. Plant products include, but are not limited to, seeds, nuts, nut meats, cut flowers, plants or crops grown or stored in a greenhouse, house plants, and the like. Plant products include many animal feeds.

15 As used herein, the phrase "meat product" refers to all forms of animal flesh, including the carcass, muscle, fat, organs, skin, bones and body fluids and like components that form the animal. Animal flesh includes, but is not limited to, the flesh of mammals, birds, fishes, reptiles, amphibians, snails, clams, crustaceans, other edible species such as lobster, crab, etc., or other forms of seafood. The forms of animal flesh
20 include, for example, the whole or part of animal flesh, alone or in combination with other ingredients. Typical forms include, for example, processed meats such as cured meats, sectioned and formed products, minced products, finely chopped products, ground meat and products including ground meat, whole products, and the like.

As used herein the term "poultry" refers to all forms of any bird kept, harvested,
25 or domesticated for meat or eggs, and including chicken, turkey, ostrich, game hen, squab, guinea fowl, pheasant, quail, duck, goose, emu, or the like and the eggs of these birds. Poultry includes whole, sectioned, processed, cooked or raw poultry, and encompasses all forms of poultry flesh, by-products, and side products. The flesh of poultry includes muscle, fat, organs, skin, bones and body fluids and like components
30 that form the animal. Forms of animal flesh include, for example, the whole or part of

animal flesh, alone or in combination with other ingredients. Typical forms include, for example, processed poultry meat, such as cured poultry meat, sectioned and formed products, minced products, finely chopped products and whole products.

As used herein, the phrase "poultry debris" refers to any debris, residue, material, dirt, offal, poultry part, poultry waste, poultry viscera, poultry organ, fragments or combinations of such materials, and the like removed from a poultry carcass or portion during processing and that enters a waste stream.

As used herein, the phrase "food processing surface" refers to a surface of a tool, a machine, equipment, a structure, a building, or the like that is employed as part of a food processing, preparation, or storage activity. Examples of food processing surfaces include surfaces of food processing or preparation equipment (e.g., slicing, canning, or transport equipment, including flumes), of food processing wares (e.g., utensils, dishware, wash ware, and bar glasses), and of floors, walls, or fixtures of structures in which food processing occurs. Food processing surfaces are found and employed in food anti-spoilage air circulation systems, aseptic packaging sanitizing, food refrigeration and cooler cleaners and sanitizers, ware washing sanitizing, blancher cleaning and sanitizing, food packaging materials, cutting board additives, third-sink sanitizing, beverage chillers and warmers, meat chilling or scalding waters, autodish sanitizers, sanitizing gels, cooling towers, food processing antimicrobial garment sprays, and non-to-low-aqueous food preparation lubricants, oils, and rinse additives.

As used herein, the term "ware" refers to items such as eating and cooking utensils, dishes, and other hard surfaces such as showers, sinks, toilets, bathtubs, countertops, windows, mirrors, transportation vehicles, and floors. As used herein, the term "warewashing" refers to washing, cleaning, or rinsing ware. Ware also refers to items made of plastic. Types of plastics that can be cleaned with the compositions according to the invention include but are not limited to, those that include polycarbonate polymers (PC), acrylonitrile-butadiene-styrene polymers (ABS), and polysulfone polymers (PS). Another exemplary plastic that can be cleaned using the compounds and compositions of the invention include polyethylene terephthalate (PET).

As used herein, the phrase "air streams" includes food anti-spoilage air circulation systems. Air streams also include air streams typically encountered in hospital, surgical, infirmity, birthing, mortuary, and clinical diagnosis rooms.

As used herein, the term "waters" includes food process or transport waters.
5 Food process or transport waters include produce transport waters (e.g., as found in flumes, pipe transports, cutters, slicers, blanchers, retort systems, washers, and the like), belt sprays for food transport lines, boot and hand-wash dip-pans, third-sink rinse waters, and the like. Waters also include domestic and recreational waters such as pools, spas, recreational flumes and water slides, fountains, and the like.

10 As used herein, the phrase "health care surface" refers to a surface of an instrument, a device, a cart, a cage, furniture, a structure, a building, or the like that is employed as part of a health care activity. Examples of health care surfaces include surfaces of medical or dental instruments, of medical or dental devices, of electronic apparatus employed for monitoring patient health, and of floors, walls, or fixtures of
15 structures in which health care occurs. Health care surfaces are found in hospital, surgical, infirmity, birthing, mortuary, and clinical diagnosis rooms. These surfaces can be those typified as "hard surfaces" (such as walls, floors, bed-pans, etc.), or fabric surfaces, e.g., knit, woven, and non-woven surfaces (such as surgical garments, draperies, bed linens, bandages, etc.), or patient-care equipment (such as respirators,
20 diagnostic equipment, shunts, body scopes, wheel chairs, beds, etc.), or surgical and diagnostic equipment. Health care surfaces include articles and surfaces employed in animal health care.

As used herein, the term "instrument" refers to the various medical or dental instruments or devices that can benefit from cleaning with a composition according to
25 the present invention.

As used herein, the phrases "medical instrument," "dental instrument," "medical device," "dental device," "medical equipment," or "dental equipment" refer to instruments, devices, tools, appliances, apparatus, and equipment used in medicine or dentistry. Such instruments, devices, and equipment can be cold sterilized, soaked or
30 washed and then heat sterilized, or otherwise benefit from cleaning in a composition of

the present invention. These various instruments, devices and equipment include, but are not limited to: diagnostic instruments, trays, pans, holders, racks, forceps, scissors, shears, saws (e.g. bone saws and their blades), hemostats, knives, chisels, rongeurs, files, nippers, drills, drill bits, rasps, burrs, spreaders, breakers, elevators, clamps, 5 needle holders, carriers, clips, hooks, gouges, curettes, retractors, straightener, punches, extractors, scoops, keratomes, spatulas, expressors, trocars, dilators, cages, glassware, tubing, catheters, cannulas, plugs, stents, scopes (e.g., endoscopes, stethoscopes, and arthoscopes) and related equipment, and the like, or combinations thereof.

As used herein, "agricultural" or "veterinary" objects or surfaces include animal 10 feeds, animal watering stations and enclosures, animal quarters, animal veterinarian clinics (e.g. surgical or treatment areas), animal surgical areas, and the like.

As used herein, the term "phosphorus-free" or "substantially phosphorus-free" refers to a composition, mixture, or ingredient that does not contain phosphorus or a phosphorus-containing compound or to which phosphorus or a phosphorus-containing 15 compound has not been added. Should phosphorus or a phosphorus-containing compound be present through contamination of a phosphorus-free composition, mixture, or ingredients, the amount of phosphorus shall be less than 0.5 wt %. More preferably, the amount of phosphorus is less than 0.1 wt-%, and most preferably the amount of phosphorus is less than 0.01 wt %.

20 For the purpose of this patent application, successful microbial reduction is achieved when the microbial populations are reduced by at least about 50%, or by significantly more than is achieved by a wash with water. Larger reductions in microbial population provide greater levels of protection.

As used herein, the term "sanitizer" refers to an agent that reduces the number of 25 bacterial contaminants to safe levels as judged by public health requirements. In an embodiment, sanitizers for use in this invention will provide at least a 99.999% reduction (5-log order reduction). These reductions can be evaluated using a procedure set out in *Germicidal and Detergent Sanitizing Action of Disinfectants*, Official Methods of Analysis of the Association of Official Analytical Chemists, paragraph 30 960.09 and applicable sections, 15th Edition, 1990 (EPA Guideline 91-2). According to

this reference a sanitizer should provide a 99.999% reduction (5-log order reduction) within 30 seconds at room temperature, 25±2°C, against several test organisms.

As used herein, the term "disinfectant" refers to an agent that kills all vegetative cells including most recognized pathogenic microorganisms. using the procedure
5 described in *A.O.A.C. Use Dilution Methods*, Official Methods of Analysis of the Association of Official Analytical Chemists, paragraph 955.14 and applicable sections, 15th Edition, 1990 (EPA Guideline 91-2). As used herein, the term "high level disinfection" or "high level disinfectant" refers to a compound or composition that kills
10 substantially all organisms, except high levels of bacterial spores, and is effected with a chemical germicide cleared for marketing as a sterilant by the Food and Drug Administration. As used herein, the term "intermediate-level disinfection" or "intermediate level disinfectant" refers to a compound or composition that kills mycobacteria, most viruses, and bacteria with a chemical germicide registered as a tuberculocide by the Environmental Protection Agency (EPA). As used herein, the term
15 "low-level disinfection" or "low level disinfectant" refers to a compound or composition that kills some viruses and bacteria with a chemical germicide registered as a hospital disinfectant by the EPA.

As used in this invention, the term "sporicide" refers to a physical or chemical agent or process having the ability to cause greater than a 90% reduction (1-log order
20 reduction) in the population of spores of *Bacillus cereus* or *Bacillus subtilis* within 10 seconds at 60° C. In certain embodiments, the sporicidal compositions of the invention provide greater than a 99% reduction (2-log order reduction), greater than a 99.99% reduction (4-log order reduction), or greater than a 99.999% reduction (5-log order reduction) in such population within 10 seconds at 60° C.

25 Differentiation of antimicrobial "-cidal" or "-static" activity, the definitions which describe the degree of efficacy, and the official laboratory protocols for measuring this efficacy are considerations for understanding the relevance of antimicrobial agents and compositions. Antimicrobial compositions can effect two kinds of microbial cell damage. The first is a lethal, irreversible action resulting in
30 complete microbial cell destruction or incapacitation. The second type of cell damage

is reversible, such that if the organism is rendered free of the agent, it can again multiply. The former is termed microbiocidal and the later, microbistatic. A sanitizer and a disinfectant are, by definition, agents which provide antimicrobial or microbiocidal activity. In contrast, a preservative is generally described as an inhibitor
5 or microbistatic composition

As used herein, the term "alkyl" or "alkyl groups" refers to saturated hydrocarbons having one or more carbon atoms, including straight-chain alkyl groups (e.g., methyl, ethyl, propyl, butyl, pentyl, hexyl, heptyl, octyl, nonyl, decyl, etc.), cyclic alkyl groups (or "cycloalkyl" or "alicyclic" or "carbocyclic" groups) (e.g., cyclopropyl,
10 cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, etc.), branched-chain alkyl groups (e.g., isopropyl, tert-butyl, sec-butyl, isobutyl, etc.), and alkyl-substituted alkyl groups (e.g., alkyl-substituted cycloalkyl groups and cycloalkyl-substituted alkyl groups).

Unless otherwise specified, the term "alkyl" includes both "unsubstituted alkyls" and "substituted alkyls." As used herein, the term "substituted alkyls" refers to alkyl
15 groups having substituents replacing one or more hydrogens on one or more carbons of the hydrocarbon backbone. Such substituents may include, for example, alkenyl, alkynyl, halogeno, hydroxyl, alkylcarbonyloxy, arylcarbonyloxy, alkoxy carbonyloxy, aryloxy, aryloxy carbonyloxy, carboxylate, alkylcarbonyl, arylcarbonyl, alkoxy carbonyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylthiocarbonyl, alkoxy,
20 phosphate, phosphonato, phosphinato, cyano, amino (including alkyl amino, dialkylamino, arylamino, diarylamino, and alkylarylamino), acylamino (including alkylcarbonylamino, arylcarbonylamino, carbamoyl and ureido), imino, sulfhydryl, alkylthio, arylthio, thiocarboxylate, sulfates, alkylsulfinyl, sulfonates, sulfamoyl, sulfonamido, nitro, trifluoromethyl, cyano, azido, heterocyclic, alkylaryl, or aromatic
25 (including heteroaromatic) groups.

In some embodiments, substituted alkyls can include a heterocyclic group. As used herein, the term "heterocyclic group" includes closed ring structures analogous to carbocyclic groups in which one or more of the carbon atoms in the ring is an element other than carbon, for example, nitrogen, sulfur or oxygen. Heterocyclic groups may be
30 saturated or unsaturated. Exemplary heterocyclic groups include, but are not limited to,

aziridine, ethylene oxide (epoxides, oxiranes), thiirane (episulfides), dioxirane, azetidene, oxetane, thietane, dioxetane, dithietane, dithiete, azolidine, pyrrolidine, pyrroline, oxolane, dihydrofuran, and furan.

5 *Compounds of the Invention*

The present invention relates, at least in part, to sulfoperoxycarboxylic acids, compositions thereof, and the use thereof in a variety of bleaching, disinfecting and cleaning applications. The sulfoperoxycarboxylic acids of the present invention are also useful as coupling agents. Further, certain compounds of the present invention can be
10 derived from non-petroleum based, renewable oils, e.g., castor, toll, soybean, canola, olive, peanut, tallow, rapeseed, and palm oils.


As used herein, the term "sulfoperoxycarboxylic acid" or "sulfonated peroxycarboxylic acid" refers to the peroxycarboxylic acid form of a sulfonated carboxylic acid. The sulfoperoxycarboxylic acids of the present invention can be used
15 alone, or can be combined with additional ingredients. In some embodiments, compositions of the present invention can include one or more of the sulfoperoxycarboxylic acids of the present invention.

Peroxy-carboxylic (or percarboxylic) acids generally have the formula $R(CO_3H)_n$, where, for example, R is an alkyl, arylalkyl, cycloalkyl, aromatic, or
20 heterocyclic group, and n is one, two, or three, and named by prefixing the parent acid with peroxy. Percarboxylic acids can be made by the direct, acid catalyzed equilibrium action of hydrogen peroxide with the carboxylic acid, by autooxidation of aldehydes, or from acid chlorides, and hydrides, or carboxylic anhydrides with hydrogen or sodium peroxide. The R group can be saturated or unsaturated as well as substituted or
25 unsubstituted.

The chemical structures herein are drawn according to the conventional standards known in the art. Thus, where an atom, such as a carbon atom, as drawn appears to have an unsatisfied valency, then that valency is assumed to be satisfied by a hydrogen atom, even though that hydrogen atom is not necessarily explicitly drawn.
30 The structures of some of the compounds of this invention include stereogenic carbon

atoms. It is to be understood that isomers arising from such asymmetry (e.g., all enantiomers and diastereomers) are included within the scope of this invention unless indicated otherwise. That is, unless otherwise stipulated, any chiral carbon center may be of either (R)- or (S)-stereochemistry. Such isomers can be obtained in substantially pure form by classical separation techniques and by stereochemically-controlled synthesis. Furthermore, alkenes can include either the E- or Z-geometry, where appropriate. In addition, the compounds of the present invention may exist in unsolvated as well as solvated forms with acceptable solvents such as water, THF, ethanol, and the like. In general, the solvated forms are considered equivalent to the unsolvated forms for the purposes of the present invention.

In some aspects, the present invention pertains to sulfoperoxycarboxylic acids of Formula I:



15

(Formula I)

wherein R_1 is hydrogen, or a substituted or unsubstituted alkyl group;

R_2 is a substituted or unsubstituted alkyl group;

X is hydrogen, a cationic group, or an ester forming moiety;

20 or salts or esters thereof.

In some embodiments, R_1 is a substituted or unsubstituted C_m alkyl group; X is hydrogen a cationic group, or an ester forming moiety; R_2 is a substituted or unsubstituted C_n alkyl group; $m=1$ to 10 ; $n = 1$ to 10 ; and $m+n$ is less than 18 , or salts, esters or mixtures thereof.

25 In some embodiments, R_1 is hydrogen. In other embodiments, R_1 is a substituted or unsubstituted alkyl group. In some embodiments, R_1 is a substituted or unsubstituted alkyl group that does not include a cyclic alkyl group. In some embodiments, R_1 is a substituted alkyl group. In some embodiments, R_1 is an unsubstituted C_1 - C_9 alkyl group. In some embodiments, R_1 is an unsubstituted C_7 or C_8 alkyl. In other embodiments, R_1 is a substituted $C_8 - C_{10}$ alkyl group. In some



30





embodiments, R_1 is a substituted C_9 - C_{10} alkyl group is substituted with at least 1, or at least 2 hydroxyl groups. In still yet other embodiments, R_1 is a substituted C_1 - C_9 alkyl group. In some embodiments, R_1 is a substituted C_1 - C_9 substituted alkyl group is substituted with at least 1 SO_3H group.

- 5 In other embodiments, R_1 is a C_9 - C_{10} substituted alkyl group. In some embodiments, R_1 is a substituted C_9 - C_{10} alkyl group wherein at least two of the carbons on the carbon backbone form a heterocyclic group. In some embodiments, the heterocyclic group is an epoxide group.




- 10 In some embodiments, R_2 is a substituted C_1 to C_{10} alkyl group. In some embodiments, R_2 is a substituted C_8 - C_{10} alkyl. In some embodiments, R_2 is an unsubstituted C_6 - C_9 alkyl. In other embodiments, R_2 is a C_8 to C_{10} alkyl group substituted with at least one hydroxyl group. In some embodiments, R_2 is a C_{10} alkyl group substituted with at least two hydroxyl groups. In other embodiments, R_2 is a C_8 alkyl group substituted with at least one SO_3H group. In some embodiments, R_2 is a
- 15 substituted C_9 group, wherein at least two of the carbons on the carbon backbone form a heterocyclic group. In some embodiments, the heterocyclic group is an epoxide group. In some embodiments, R_1 is a C_8 - C_9 substituted or unsubstituted alkyl, and R_2 is a C_7 - C_8 substituted or unsubstituted alkyl.

- 20 In some embodiments, the compound of the invention is selected from the group consisting of:






5




and mixtures and derivatives thereof.


10 In other embodiments, the compound of the invention is selected from the group consisting of:





5



10



15







, and mixtures and derivatives thereof.





Compounds of the invention are also shown in Table 1 below.

5 Table 1.

Sulfonated Peroxyacid Compounds	
I	Structure/Name of Compound
D	
A	$\text{CH}_3(\text{CH}_2)_6-\overset{\text{H}}{\underset{\text{H}}{\text{C}}}-\overset{\text{H}}{\underset{\text{OH}}{\text{C}}}-\overset{\text{H}}{\underset{\text{SO}_3\text{H}}{\text{C}}}-\text{(CH}_2)_7-\overset{\text{O}}{\parallel}{\text{C}}-\text{O}-\text{OH}$ <p style="text-align: center;">10-Hydroxy-9-sulfooctadecaneperoxoic acid</p>
B	$\text{CH}_3(\text{CH}_2)_7-\overset{\text{H}}{\underset{\text{OH}}{\text{C}}}-\overset{\text{H}}{\underset{\text{OH}}{\text{C}}}-\overset{\text{H}}{\underset{\text{SO}_3\text{H}}{\text{C}}}-\text{(CH}_2)_6-\overset{\text{O}}{\parallel}{\text{C}}-\text{O}-\text{OH}$ <p style="text-align: center;">9,10-Dihydroxy-8-sulfooctadecaneperoxoic acid</p>
C	

	$\text{CH}_3(\text{CH}_2)_7-\overset{\text{H}}{\underset{\text{H}}{\text{C}}}-\overset{\text{H}}{\underset{\text{SO}_3\text{H}}{\text{C}}}-\overset{\text{H}}{\text{C}}-(\text{CH}_2)_6-\overset{\text{O}}{\parallel}-\text{O}-\text{OH}$ <p style="text-align: center;">9-Sulfooctadecaneperoxoic acid</p>
D	$\overset{\text{H}}{\text{HC}}-\overset{\text{H}}{\underset{\text{SO}_3\text{H}}{\text{C}}}-\overset{\text{H}}{\text{C}}-(\text{CH}_2)_7-\overset{\text{O}}{\parallel}-\text{O}-\text{OH}$ <p style="text-align: center;">11-Sulfoundecaneperoxoic acid</p>
E	$\overset{\text{H}}{\text{HC}}-\overset{\text{H}}{\underset{\text{SO}_3\text{H}}{\text{C}}}-\overset{\text{H}}{\underset{\text{SO}_3\text{H}}{\text{C}}}-\overset{\text{H}}{\text{C}}-(\text{CH}_2)_7-\overset{\text{O}}{\parallel}-\text{O}-\text{OH}$ <p style="text-align: center;">10,11-Disulfoundecaneperoxoic acid</p>
F	<p style="text-align: center;">8-(3-octyloxiran-2-yl)-8-sulfooctaneperoxoic acid</p> $\text{CH}_3(\text{CH}_2)_7-\overset{\text{H}}{\underset{\text{O}}{\text{C}}}-\overset{\text{H}}{\text{C}}-\overset{\text{H}}{\underset{\text{SO}_3\text{H}}{\text{C}}}-\text{CH}_2(\text{CH}_2)_6-\overset{\text{O}}{\parallel}-\text{O}-\text{OH}$
G	$\text{CH}_3(\text{CH}_2)_6-\overset{\text{H}}{\underset{\text{SO}_3\text{H}}{\text{C}}}-\overset{\text{H}}{\underset{\text{OH}}{\text{C}}}-\overset{\text{H}}{\underset{\text{OH}}{\text{C}}}-\text{CH}_2(\text{CH}_2)_6-\overset{\text{O}}{\parallel}-\text{O}-\text{OH}$

	<p>9,10-Dihydroxy-11-sulfooctadecaneperoxoic acid</p>
H	 <p>8-(3-octyloxiran-2-yl)-8-sulfoctaneperoxoic acid</p>
I	<p>9-Hydroxy-10-sulfooctadecaneperoxoic acid</p> 
J	 <p>10-Sulfooctadecaneperoxoic acid</p>
K	

	9,10-Disulfoctadecaneperoxoic acid
L	
	10-Sulfoundecaneperoxoic acid
M	
	9-(3-heptyloxiran-2-yl)-9-sulfonyldecaneperoxoic acid
N	
	10,11-dihydroxy-9-sulfooctadecaneperoxoic acid
O	
	8,9-dihydroxy-10-sulfoctadecaneperoxoic acid

In some embodiments, the starting material for the preparation of the compounds of the present invention is a sulfonated fatty acid. Without wishing to be bound by any particular theory, it is thought that the sulfo-group is inert in an oxidative environment. Further, it is thought that the hydrophilicity of the sulfo-group is not as impacted by pH as other substituents. In some embodiments, the sulfonated percarboxylic acids of the present invention are formed from commercially available

sulfonated fatty acids. In other embodiments, the compounds of the present invention are formed from commercially available non-sulfonated fatty acids, which can be sulfonated. In some embodiments, the starting fatty acid will be sulfonated prior to conversion to a peroxy-carboxylic acid. In other embodiments, the starting fatty acid
5 will be sulfonated at the same time or after the formation of the peroxy-carboxylic acid. Sulfonated fatty acids suitable for use in forming compounds of the present invention include, but are not limited to, 11-sulfoundecanoic acid, 10,11-disulfoundecanoic acid, sulfonated oleic acid, sulfonated linoleic acid, sulfonated palmitoleic acid and sulfonated stearic acid.

10 Without wishing to be bound by any particular theory, it is thought that the peracid formed from certain commercially available sulfonated oleic acid starting materials includes a mixture of the compounds of the present invention. It is thought that this is due, in part, to the nature of the sulfonated oleic acid starting material. That is, it is thought that because the sulfonated oleic acid starting material is derived from
15 naturally occurring sources, it is not chemically pure, i.e., does not contain only one form of the sulfonated oleic acid. Thus, without wishing to be bound by any particular theory it is thought that sulfonated peroleic acid formed can include (hereinafter referred to as the "sulfonated peroleic acid product") a mixture of Compounds A, N, I, and O as the primary components. Without wishing to be bound by any particular
20 theory it is thought that in some embodiments, the sulfonated peroleic acid product includes about 20-25 wt% Compound A (10-Hydroxy-9-sulfooctadecaneperoxoic acid) about 20-25 wt% Compound N (10,11-dihydroxy-9-sulfooctadecaneperoxoic acid), about 20-25 wt% Compound I (9-Hydroxy-10-sulfooctadecaneperoxoic acid), and about 20-25 wt% Compound O (8,9-dihydroxy-10-sulfooctadecaneperoxoic acid). The
25 remainder of the product is thought to include about 5 to about 10 wt% of a mixture of these compounds.

The sulfoperoxyacids can be formed using a variety of reaction mechanisms. For example, in some embodiments, the peracids are formed by the direct acid catalyzed equilibrium action of hydrogen peroxide with the starting materials.

In some embodiments, the sulfonated carboxylic acids for use in forming the compounds of the present invention are not sulfonated at the α position. It has been found that having the sulfonate group at the α position of the fatty acid prohibits the oxidation and/or perhydrolysis of the carboxylic acid group to form the corresponding peroxycarboxylic acid. Without wishing to be bound by any particular theory, it is thought that the α -sulfo group makes the carboxylic acid group on the fatty acid electronically deficient, and thus oxidation and/or perhydrolysis and formation of the corresponding percarboxylic acid does not occur.

10 *Sulfonated Peroxycarboxylic Acid Compositions*

In some aspects, the present invention relates to compositions including a sulfonated peroxycarboxylic acid compound, or mixture thereof, of Formula I. The compositions of the present invention can be used as bleaching compositions for a variety of substrates and surfaces, e.g., textiles, hard surfaces. The compositions of the present invention can also be used as disinfectant or antimicrobial compositions. Further, compounds of the present invention can be used as coupling agents in compositions for various applications, e.g., food contact sanitizing, hard surface disinfection, textile disinfection. In some embodiments, compositions containing compounds of the present invention can be multipurpose. That is, the compositions of the present invention can, for example, act as both antimicrobials and bleaches, or as both coupling agents, and bleaching agents.

The compositions of the present invention also show enhanced stability compared to conventional peroxygen containing compositions. In some embodiments, the compositions of the present invention are stable for at least about 1 year at room temperature. In some embodiments, the compositions of the present invention are stable at about 100°F for at least 30 days. In other embodiments, the compositions of the present invention are stable at about 140°F for at least 30 days. For example, 11-sulfoundecanoic peroxyacid (Compound D) is stable as a powder system at about 140°F for at least 30 days.

The compositions of the present invention have no or low odor. For example, in some embodiments, compositions of the present invention have an odor less unpleasant than (e.g., as measured by an hedonic tone rating) than 5, 4, 3, 2, or 1 wt-% acetic acid in water. In other embodiments, the compositions of the present invention have no odor
 5 detectable by a user.

In some embodiments, the compositions of the present invention include a sulfonated peracid or mixture thereof of Formula I, and at least one additional ingredient. Additional ingredients suitable for use with the compositions of the present invention include, but are not limited to, oxidizing agents, carboxylic acids, surfactants,
 10 stabilizing agents (e.g., metal chelators), and mixtures thereof. The compounds and compositions of the invention can also be used in conjunction with conventional cleaning agents, e.g., alkaline detergents.

In some embodiments, the compositions of the present invention can be used as a sanitizing composition for articles cleaned using a clean in place (CIP) technique.
 15 Such compositions can include an oxidizing agent, a stabilizing agent, an acidulant and a surfactant or mixture thereof, in the following concentrations.

Table A – Concentrate CIP Sanitizer by Weight %

Oxidizing Agent	0.1 - 10	2-8	5-7
Stabilizing Agent	0.1-10	0.5-5	1-2
Acidulant	0-50	10-40	20-30
Surfactant	0-50	10-40	25-35

In other embodiments, the compositions of the present invention can be used as
 20 a textile disinfectant/sanitizer. Such compositions can include oxidizing agent, stabilizing agent and a carboxylic acid in the following concentrations.

Table B. – Concentrate Textile Disinfectant/Sanitizer by Weight %

Oxidizing Agent	10-75	25-60	30-50
Stabilizing Agent	0.1-10	0.5-5	2-4
Carboxylic Acid	1-40	10-30	20-25

Oxidizing agents

In some aspects, the compositions of the present invention include a compound of Formula I. In some embodiments, the compositions of the present invention further
5 include at least one oxidizing agent. In some embodiments, the compositions of the present invention are substantially free of an oxidizing agent. When present, the present composition can include any of a variety of oxidizing agents, for example, hydrogen peroxide. The oxidizing agent can be present at an amount effective to convert a sulfonated carboxylic acid to a sulfonated peroxy-carboxylic acid. In some
10 embodiments, the oxidizing agent can also have antimicrobial activity. In other embodiments, the oxidizing agent is present in an amount insufficient to exhibit antimicrobial activity.

In some embodiments, the compositions of the present invention include about 0.001 wt % oxidizing agent to about 99 wt% oxidizing agent. In other embodiments,
15 the compositions of the present invention include about 1 wt% to about 60 wt% oxidizing agent. In some embodiments, the compositions of the invention include about 50 wt% to about 80 wt% oxidizing agent. In other embodiments, the compositions of the invention include about 15 wt% to about 30 wt% oxidizing agent. In yet other
20 embodiments, the compositions of the present invention include about 25 wt% oxidizing agent. It is to be understood that all ranges and values between these ranges and values are encompassed by the present invention.

Examples of inorganic oxidizing agents include the following types of compounds or sources of these compounds, or alkali metal salts including these types of
25 compounds, or forming an adduct therewith: hydrogen peroxide, urea-hydrogen peroxide complexes or hydrogen peroxide donors of: group 1 (IA) oxidizing agents, for example lithium peroxide, sodium peroxide; group 2 (IIA) oxidizing agents, for example magnesium peroxide, calcium peroxide, strontium peroxide, barium peroxide; group 12 (IIB) oxidizing agents, for example zinc peroxide; group 13 (IIIA) oxidizing
30 agents, for example boron compounds, such as perborates, for example sodium perborate hexahydrate of the formula $\text{Na}_2[\text{B}_2(\text{O}_2)_2(\text{OH})_4] \cdot 6\text{H}_2\text{O}$ (also called sodium

perborate tetrahydrate); sodium peroxyborate tetrahydrate of the formula $\text{Na}_2\text{B}_2(\text{O}_2)_2[(\text{OH})_4] \cdot 4\text{H}_2\text{O}$ (also called sodium perborate trihydrate); sodium peroxyborate of the formula $\text{Na}_2[\text{B}_2(\text{O}_2)_2(\text{OH})_4]$ (also called sodium perborate monohydrate); group 14 (IVA) oxidizing agents, for example persilicates and peroxycarbonates, which are also called percarbonates, such as persilicates or peroxycarbonates of alkali metals; group 15 (VA) oxidizing agents, for example peroxyphosphoric acids and their salts, for example, perphosphates; group 16 (VIA) oxidizing agents, for example peroxyphosphoric acids and their salts, such as peroxydisulfuric acids, and their salts, such as persulfates, for example, sodium persulfate; and group VIIa oxidizing agents such as sodium periodate, potassium perchlorate. Other active inorganic oxygen compounds can include transition metal peroxides; and other such peroxygen compounds, and mixtures thereof.

In some embodiments, the compositions of the present invention employ one or more of the inorganic oxidizing agents listed above. Suitable inorganic oxidizing agents include ozone, hydrogen peroxide, hydrogen peroxide adduct, group IIIA oxidizing agent, or hydrogen peroxide donors of group VIA oxidizing agent, group VA oxidizing agent, group VIIA oxidizing agent, or mixtures thereof. Suitable examples of such inorganic oxidizing agents include percarbonate, perborate, persulfate, perphosphate, persilicate, or mixtures thereof.

Carboxylic and Percarboxylic Acids

In some embodiments, the compositions of the present invention include at least one sulfoperoxy-carboxylic acid of the present invention, and at least one carboxylic and/or percarboxylic acid. In some embodiments, the compositions of the present invention include at least two, at least three, or at least four or more carboxylic and/or percarboxylic acids.

In some embodiments, the carboxylic acid for use with the compositions of the present invention includes a C_1 to C_{22} carboxylic acid. In some embodiments, the carboxylic acid for use with the compositions of the present invention is a C_5 to C_{11}

carboxylic acid. In some embodiments, the carboxylic acid for use with the compositions of the present invention is a C₁ to C₄ carboxylic acid. Examples of suitable carboxylic acids include, but are not limited to, formic, acetic, propionic, butanoic, pentanoic, hexanoic, heptanoic, octanoic, nonanoic, decanoic, undecanoic, 5 dodecanoic, as well as their branched isomers, lactic, malic, ascorbic, citric, hydroxyacetic, neopentanoic, neoheptanoic, neodecanoic, oxalic, malonic, succinic, glutaric, adipic, pimelic subric acid, and mixtures thereof.

In some embodiments, the compositions of the present invention include about 0.1wt% to about 80 wt% of a carboxylic acid. In other embodiments, the compositions 10 of the present invention include about 1 wt% to about 60 wt% of a carboxylic acid. In yet other embodiments, the compositions of the present invention include about 20 wt%, about 30 wt%, or about 40 wt% of a carboxylic acid. In some embodiments, the compositions of the present invention include about 5 wt% to about 10 wt% of acetic acid. In other embodiments, the compositions of the present invention include about 5 15 wt% to about 10 wt% of octanoic acid. In other embodiments, the compositions of the present invention include a combination of octanoic acid and acetic acid.

In some embodiments, the compositions of the present invention include a compound of Formula I, and at least one peroxy-carboxylic acid. Peroxy-carboxylic acids useful in the compositions and methods of the present invention include 20 peroxyformic, peroxyacetic, peroxypropionic, peroxybutanoic, peroxy-pentanoic, peroxyhexanoic, peroxyheptanoic, peroxyoctanoic, peroxy-nonanoic, peroxydecanoic, peroxyundecanoic, peroxydodecanoic, or the peroxyacids of their branched chain isomers, peroxy-lactic, peroxy-maleic, peroxy-ascorbic, peroxyhydroxyacetic, peroxyoxalic, peroxy-malonic, peroxy-succinic, peroxyglutaric, peroxyadipic, 25 peroxy-pimelic and peroxy-subric acid and mixtures thereof. In some embodiments, the compositions of the invention utilize a combination of several different peroxy-carboxylic acids. For example, in some embodiments, the composition includes one or more C₁ to C₄ peroxy-carboxylic acids and one or more C₅ to C₁₁ peroxy-carboxylic acids. In some embodiments, the C₁ to C₄ peroxy-carboxylic acid is peroxyacetic acid and the C₅ to C₁₁ acid is peroxyoctanoic acid. 30

In some embodiments, the compositions of the present invention include peroxyacetic acid. Peroxyacetic (or peracetic) acid is a peroxycarboxylic acid having the formula: CH_3COOOH . Generally, peroxyacetic acid is a liquid having an acrid odor at higher concentrations and is freely soluble in water, alcohol, ether, and sulfuric acid.

5 Peroxyacetic acid can be prepared through any number of methods known to those of skill in the art including preparation from acetaldehyde and oxygen in the presence of cobalt acetate. A solution of peroxyacetic acid can be obtained by combining acetic acid with hydrogen peroxide. A 50% solution of peroxyacetic acid can be obtained by combining acetic anhydride, hydrogen peroxide and sulfuric acid.

10 In some embodiments, the compositions of the present invention include peroxyoctanoic acid, peroxynonanoic acid, or peroxyheptanoic acid. In some embodiments, the compositions include peroxyoctanoic acid. Peroxyoctanoic (or perooctanoic) acid is a peroxycarboxylic acid having the formula, for example, of n-peroxyoctanoic acid: $\text{CH}_3(\text{CH}_2)_6\text{COOOH}$. Peroxyoctanoic acid can be an acid with a straight chain alkyl moiety, an acid with a branched alkyl moiety, or a mixture thereof.

15 Peroxyoctanoic acid can be prepared through any number of methods known to those of skill in the art. A solution of peroxyoctanoic acid can be obtained by combining octanoic acid and hydrogen peroxide and a hydrotrope, solvent or carrier.

In some embodiments, the compositions of the present invention include about 20 0.1 wt% to about 90wt% of one or more peroxycarboxylic acids. In other embodiments, the compositions of the present invention include about 1 wt% to about 25 wt% of one or more peroxycarboxylic acids. In yet other embodiments, the compositions of the present invention include about 5 wt% to about 10 wt% of one or more peroxycarboxylic acids. In some embodiments, the compositions of the present invention include about 1 wt% to about 25 wt% of peroxyacetic acid. In other 25 embodiments, the compositions of the present invention include about 0.1 wt% to about 10 wt% of peroxyoctanoic acid. In still yet other embodiments, the compositions of the present invention include a mixture of about 5 wt% peroxyacetic acid, and about 1.5 wt% peroxyoctanoic acid.

30

Surfactants

In some embodiments, the compositions of the present invention include a surfactant. Surfactants suitable for use with the compositions of the present invention include, but are not limited to, nonionic surfactants, anionic surfactants, and zwitterionic surfactants. In some embodiments, the compositions of the present invention include about 10wt% to about 50wt% of a surfactant. In other embodiments the compositions of the present invention include about 15wt% to about 30% of a surfactant. In still yet other embodiments, the compositions of the present invention include about 25wt% of a surfactant. In some embodiments, the compositions of the present invention include about 100 ppm to about 1000 ppm of a surfactant.


Nonionic Surfactants

Suitable nonionic surfactants suitable for use with the compositions of the present invention include alkoxyated surfactants. Suitable alkoxyated surfactants include EO/PO copolymers, capped EO/PO copolymers, alcohol alkoxyates, capped alcohol alkoxyates, mixtures thereof, or the like. Suitable alkoxyated surfactants for use as solvents include EO/PO block copolymers, such as the Pluronic and reverse Pluronic surfactants; alcohol alkoxyates, such as Dehypon LS-54 (R-(EO)₅(PO)₄) and Dehypon LS-36 (R-(EO)₃(PO)₆); and capped alcohol alkoxyates, such as Plurafac LF221 and Tegoten EC11; mixtures thereof, or the like.

Semi-Polar Nonionic Surfactants

The semi-polar type of nonionic surface active agents are another class of nonionic surfactant useful in compositions of the present invention. Semi-polar nonionic surfactants include the amine oxides, phosphine oxides, sulfoxides and their alkoxyated derivatives.

Amine oxides are tertiary amine oxides corresponding to the general formula:



wherein the arrow is a conventional representation of a semi-polar bond; and, R^1 , R^2 , and R^3 may be aliphatic, aromatic, heterocyclic, alicyclic, or combinations thereof. Generally, for amine oxides of detergent interest, R^1 is an alkyl radical of from about 8 to about 24 carbon atoms; R^2 and R^3 are alkyl or hydroxyalkyl of 1-3 carbon atoms or a mixture thereof; R^2 and R^3 can be attached to each other, e.g. through an oxygen or nitrogen atom, to form a ring structure; R^4 is an alkylene or a hydroxyalkylene group containing 2 to 3 carbon atoms; and n ranges from 0 to about 20. An amine oxide can be generated from the corresponding amine and an oxidizing agent, such as hydrogen peroxide.

Useful water soluble amine oxide surfactants are selected from the octyl, decyl, dodecyl, isododecyl, coconut, or tallow alkyl di-(lower alkyl) amine oxides, specific examples of which are octyldimethylamine oxide, nonyldimethylamine oxide, dodecyldimethylamine oxide, undecyldimethylamine oxide, dodecyldimethylamine oxide, iso-dodecyldimethylamine oxide, tridecyldimethylamine oxide, tetradecyldimethylamine oxide, pentadecyldimethylamine oxide, hexadecyldimethylamine oxide, heptadecyldimethylamine oxide, octadecyldimethylamine oxide, dodecyldipropylamine oxide, tetradecyldipropylamine oxide, hexadecyldipropylamine oxide, tetradecyldibutylamine oxide, octadecyldibutylamine oxide, bis(2-hydroxyethyl)dodecylamine oxide, bis(2-hydroxyethyl)-3-dodecoxy-1-hydroxypropylamine oxide, dimethyl-(2-hydroxydodecyl)amine oxide, 3,6,9-trioctadecyldimethylamine oxide and 3-dodecoxy-2-hydroxypropyl-di-(2-hydroxyethyl)amine oxide.

25 *Anionic surfactants*


Anionic sulfate surfactants suitable for use in the present compositions include alkyl ether sulfates, alkyl sulfates, the linear and branched primary and secondary alkyl


sulfates, alkyl ethoxysulfates, fatty oleyl glycerol sulfates, alkyl phenol ethylene oxide ether sulfates, the C₅-C₁₇ acyl-N-(C₁-C₄ alkyl) and -N-(C₁-C₂ hydroxyalkyl) glucamine sulfates, and sulfates of alkylpolysaccharides such as the sulfates of alkylpolyglucoside, and the like. Also included are the alkyl sulfates, alkyl
 5 poly(ethyleneoxy) ether sulfates and aromatic poly(ethyleneoxy) sulfates such as the sulfates or condensation products of ethylene oxide and nonyl phenol (usually having 1 to 6 oxyethylene groups per molecule).


Anionic sulfonate surfactants suitable for use in the present compositions also include alkyl sulfonates, the linear and branched primary and secondary alkyl
 10 sulfonates, and the aromatic sulfonates with or without substituents.

Anionic carboxylate surfactants suitable for use in the present compositions include carboxylic acids (and salts), such as alkanolic acids (and alkanooates), ester carboxylic acids (e.g. alkyl succinates), ether carboxylic acids, and the like. Such carboxylates include alkyl ethoxy carboxylates, alkyl aryl ethoxy carboxylates, alkyl
 15 polyethoxy polycarboxylate surfactants and soaps (e.g. alkyl carboxyls). Secondary carboxylates useful in the present compositions include those which contain a carboxyl unit connected to a secondary carbon. The secondary carbon can be in a ring structure, e.g. as in p-octyl benzoic acid, or as in alkyl-substituted cyclohexyl carboxylates. The secondary carboxylate surfactants typically contain no ether linkages, no ester linkages
 20 and no hydroxyl groups. Further, they typically lack nitrogen atoms in the head-group (amphiphilic portion). Suitable secondary soap surfactants typically contain 11-13 total carbon atoms, although more carbon atoms (e.g., up to 16) can be present. Suitable carboxylates also include acylamino acids (and salts), such as acylgluamates, acyl peptides, sarcosinates (e.g. N-acyl sarcosinates), taurates (e.g. N-acyl taurates and fatty
 25 acid amides of methyl tauride), and the like.

Suitable anionic surfactants include alkyl or alkylaryl ethoxy carboxylates of the following formula:



in which R is a C₈ to C₂₂ alkyl group or , in which R¹ is a C₄-C₁₆ alkyl group; n is an integer of 1-20; m is an integer of 1-3; and X is a counter ion, such as hydrogen, sodium, potassium, lithium, ammonium, or an amine salt such as monoethanolamine, diethanolamine or triethanolamine. In some embodiments, n is an integer of 4 to 10 and m is 1. In some embodiments, R is a C₈-C₁₆ alkyl group. In some embodiments, R is a C₁₂-C₁₄ alkyl group, n is 4, and m is 1.

In other embodiments, R is  and R¹ is a C₆-C₁₂ alkyl group. In still yet other embodiments, R¹ is a C₉ alkyl group, n is 10 and m is 1.

Such alkyl and alkylaryl ethoxy carboxylates are commercially available. These ethoxy carboxylates are typically available as the acid forms, which can be readily converted to the anionic or salt form. Commercially available carboxylates include, Neodox 23-4, a C₁₂₋₁₃ alkyl polyethoxy (4) carboxylic acid (Shell Chemical), and Emcol CNP-110, a C₉ alkylaryl polyethoxy (10) carboxylic acid (Witco Chemical). Carboxylates are also available from Clariant, e.g. the product Sandopan[®] DTC, a C₁₃ alkyl polyethoxy (7) carboxylic acid.

Amphoteric Surfactants


Amphoteric, or ampholytic, surfactants contain both a basic and an acidic hydrophilic group and an organic hydrophobic group. These ionic entities may be any of anionic or cationic groups described herein for other types of surfactants. A basic nitrogen and an acidic carboxylate group are the typical functional groups employed as the basic and acidic hydrophilic groups. In a few surfactants, sulfonate, sulfate, phosphonate or phosphate provide the negative charge.

Amphoteric surfactants can be broadly described as derivatives of aliphatic secondary and tertiary amines, in which the aliphatic radical may be straight chain or branched and wherein one of the aliphatic substituents contains from about 8 to 18 carbon atoms and one contains an anionic water solubilizing group, e.g., carboxy, sulfo,

sulfato, phosphato, or phosphono. Amphoteric surfactants are subdivided into two major classes known to those of skill in the art and described in "Surfactant Encyclopedia" Cosmetics & Toiletries, Vol. 104 (2) 69-71 (1989). The first class includes acyl/dialkyl ethylenediamine derivatives (e.g. 2-alkyl hydroxyethyl imidazoline derivatives) and their salts. The second class includes N-alkylamino acids and their salts. Some amphoteric surfactants can be envisioned as fitting into both classes.

Amphoteric surfactants can be synthesized by methods known to those of skill in the art. For example, 2-alkyl hydroxyethyl imidazoline is synthesized by condensation and ring closure of a long chain carboxylic acid (or a derivative) with dialkyl ethylenediamine. Commercial amphoteric surfactants are derivatized by subsequent hydrolysis and ring-opening of the imidazoline ring by alkylation -- for example with chloroacetic acid or ethyl acetate. During alkylation, one or two carboxy-alkyl groups react to form a tertiary amine and an ether linkage with differing alkylating agents yielding different tertiary amines.

Long chain imidazole derivatives having application in the present invention generally have the general formula:



20

wherein R is an acyclic hydrophobic group containing from about 8 to 18 carbon atoms and M is a cation to neutralize the charge of the anion, generally sodium.

Commercially prominent imidazoline-derived ampherics that can be employed in the present compositions include for example: Cocoamphopropionate,

25 Cocoamphocarboxy-propionate, Cocoamphoglycinate, Cocoamphocarboxy-glycinate,

Cocoamphopropyl-sulfonate, and Cocoamphocarboxy-propionic acid.

Amphocarboxylic acids can be produced from fatty imidazolines in which the dicarboxylic acid functionality of the amphodicarboxylic acid is diacetic acid and/or dipropionic acid.

5 The carboxymethylated compounds (glycinates) described herein above frequently are called betaines. Betaines are a special class of amphoteric discussed herein below in the section entitled, Zwitterion Surfactants.

 Long chain N-alkylamino acids are readily prepared by reaction RNH_2 , in which $R=C_8-C_{18}$ straight or branched chain alkyl, fatty amines with halogenated carboxylic
10 acids. Alkylation of the primary amino groups of an amino acid leads to secondary and tertiary amines. Alkyl substituents may have additional amino groups that provide more than one reactive nitrogen center. Most commercial N-alkylamine acids are alkyl derivatives of beta-alanine or beta-N(2-carboxyethyl) alanine. Examples of commercial N-alkylamino acid ampholytes having application in this invention include alkyl beta-
15 amino dipropionates, $RN(C_2H_4COOM)_2$ and $RNHC_2H_4COOM$. In an embodiment, R can be an acyclic hydrophobic group containing from about 8 to about 18 carbon atoms, and M is a cation to neutralize the charge of the anion.

 Suitable amphoteric surfactants include those derived from coconut products such as coconut oil or coconut fatty acid. Additional suitable coconut derived
20 surfactants include as part of their structure an ethylenediamine moiety, an alkanolamide moiety, an amino acid moiety, e.g., glycine, or a combination thereof; and an aliphatic substituent of from about 8 to 18 (e.g., 12) carbon atoms. Such a surfactant can also be considered an alkyl amphodicarboxylic acid. These amphoteric surfactants can include chemical structures represented as: C_{12} -alkyl-C(O)-NH-CH₂-CH₂-N⁻(CH₂-
25 CH₂-CO₂Na)₂-CH₂-CH₂-OH or C_{12} -alkyl-C(O)-N(H)-CH₂-CH₂-N⁻(CH₂-CO₂Na)₂-CH₂-CH₂-OH. Disodium cocoampho dipropionate is one suitable amphoteric surfactant and is commercially available under the tradename Miranol™ FBS from Rhodia Inc., Cranbury, N.J. Another suitable coconut derived amphoteric surfactant with the chemical name disodium cocoampho diacetate is sold under the tradename Mirataine™
30 JCHA, also from Rhodia Inc., Cranbury, N.J.

A typical listing of amphoteric classes, and species of these surfactants, is given in U.S. Pat. No. 3,929,678 issued to Laughlin and Huring on Dec. 30, 1975. Further examples are given in "Surface Active Agents and Detergents" (Vol. I and II by Schwartz, Perry and Berch).


5

Zwitterionic Surfactants

Zwitterionic surfactants can be thought of as a subset of the amphoteric surfactants and can include an anionic charge. Zwitterionic surfactants can be broadly described as derivatives of secondary and tertiary amines, derivatives of heterocyclic secondary and tertiary amines, or derivatives of quaternary ammonium, quaternary phosphonium or tertiary sulfonium compounds. Typically, a zwitterionic surfactant includes a positive charged quaternary ammonium or, in some cases, a sulfonium or phosphonium ion; a negative charged carboxyl group; and an alkyl group. Zwitterionics generally contain cationic and anionic groups which ionize to a nearly equal degree in the isoelectric region of the molecule and which can develop strong "inner-salt" attraction between positive-negative charge centers. Examples of such zwitterionic synthetic surfactants include derivatives of aliphatic quaternary ammonium, phosphonium, and sulfonium compounds, in which the aliphatic radicals can be straight chain or branched, and wherein one of the aliphatic substituents contains from 8 to 18 carbon atoms and one contains an anionic water solubilizing group, e.g., carboxy, sulfonate, sulfate, phosphate, or phosphonate. Bctaine and sultaine surfactants are exemplary zwitterionic surfactants for use herein.

20

A general formula for these compounds is:




25 wherein R¹ contains an alkyl, alkenyl, or hydroxyalkyl radical of from 8 to 18 carbon atoms having from 0 to 10 ethylene oxide moieties and from 0 to 1 glyceryl moiety; Y is selected from the group consisting of nitrogen, phosphorus, and sulfur atoms; R² is an alkyl or monohydroxy alkyl group containing 1 to 3 carbon atoms; x is 1 when Y is a

sulfur atom and 2 when Y is a nitrogen or phosphorus atom, R³ is an alkylene or hydroxy alkylene or hydroxy alkylene of from 1 to 4 carbon atoms and Z is a radical selected from the group consisting of carboxylate, sulfonate, sulfate, phosphonate, and phosphate groups.

- 5 Examples of zwitterionic surfactants having the structures listed above include: 4-[N,N-di(2-hydroxyethyl)-N-octadecylammonio]-butane-1-carboxylate; 5-[S-3-hydroxypropyl-S-hexadecylsulfonio]-3-hydroxypentane-1-sulfate; 3-[P,P-diethyl-P-3,6,9-trioxatetracosanephosphonio]-2-hydroxypropane-1-phosphate; 3-[N,N-dipropyl-N-3-dodecoxy-2-hydroxypropyl-ammonio]-propane-1-phosphonate; 3-(N,N-dimethyl-N-hexadecylammonio)-propane-1-sulfonate; 3-(N,N-dimethyl-N-hexadecylammonio)-2-hydroxy-propane-1-sulfonate; 4-[N,N-di(2(2-hydroxyethyl)-N(2-hydroxydodecyl)ammonio)-butane-1-carboxylate; 3-[S-ethyl-S-(3-dodecoxy-2-hydroxypropyl)sulfonio]-propane-1-phosphate; 3-[P,P-dimethyl-P-dodecylphosphonio]-propane-1-phosphonate; and S[N,N-di(3-hydroxypropyl)-N-hexadecylammonio]-2-hydroxy-pentane-1-sulfate. The alkyl groups contained in said detergent surfactants can be straight or branched and saturated or unsaturated.

The zwitterionic surfactant suitable for use in the present compositions includes a betaine of the general structure:



- 20 These surfactant betaines typically do not exhibit strong cationic or anionic characters at pH extremes nor do they show reduced water solubility in their isoelectric range. Unlike "external" quaternary ammonium salts, betaines are compatible with anionics. Examples of suitable betaines include coconut acylamidopropyldimethyl betaine; hexadecyl dimethyl betaine; C₁₂₋₁₄ acylamidopropylbetaine; C₈₋₁₄ acylamidohexyldiethyl betaine; 4-C₁₄₋₁₆ acylmethylamidodiethylammonio-1-carboxybutane; C₁₆₋₁₈ acylamidodimethylbetaine; C₁₂₋₁₆ acylamidopentanedimethylbetaine; and C₁₂₋₁₆ acylmethylamidodimethylbetaine.

Sultaines useful in the present invention include those compounds having the formula $(R(R^1)_2 N^+ R^2 SO_3^-)$, in which R is a C₆-C₁₈ hydrocarbyl group, each R¹ is typically independently C₁-C₃ alkyl, e.g. methyl, and R² is a C₁-C₆ hydrocarbyl group, e.g. a C₁-C₃ alkylene or hydroxyalkylene group.

5 A typical listing of zwitterionic classes, and species of these surfactants, is given in U.S. Pat. No. 3,929,678 issued to Laughlin and Heuring on Dec. 30, 1975. Further examples are given in "Surface Active Agents and Detergents" (Vol. I and II by Schwartz, Perry and Berch).

In an embodiment, the compositions of the present invention include a betaine.
10 For example, the compositions can include cocoamidopropyl betaine.

Other Additional Ingredients

In some embodiments, the compositions of the present invention can include other additional ingredients. Additional ingredients suitable for use with the
15 compositions of the present invention include, but are not limited to, acidulants, stabilizing agents, e.g., chelating agents or sequestrants, buffers, detergents, wetting agents, defoaming agents, thickeners, foaming agents solidification agents, aesthetic enhancing agents (i.e., colorants, odorants, or perfumes) and other cleaning agents. These additional ingredients can be preformulated with the compositions of the
20 invention or added to the system before, after, or substantially simultaneously with the addition of the compositions of the present invention. Additionally, the compositions can be used in conjunction with one or more conventional cleaning agents, e.g., an alkaline detergent.

25 Acidulants

In some embodiments, the compositions of the present invention include an acidulant. The acidulant can act as a catalyst for conversion of carboxylic acid to peroxycarboxylic acid. The acidulant can be effective to form a concentrate
composition with pH of about 1 or less. The acidulant can be effective to form a use
30 composition with pH of about 5, about 5 or less, about 4, about 4 or less, about 3, about

3 or less, about 2, about 2 or less, or the like. In some embodiments, an acidulant can be used to lower the pH of an alkaline cleaning solution to a pH of about 10, about 10 or less, about 9, about 9 or less, about 8, about 8 or less, about 7, about 7 or less, about 6, or about 6 or less. In an embodiment, the acidulant includes an inorganic acid. Suitable
5 inorganic acids include, but are not limited to, sulfuric acid, sodium bisulfate, phosphoric acid, nitric acid, hydrochloric acid. In some embodiments, the acidulant includes an organic acid. Suitable organic acids include, but are not limited to, methane sulfonic acid, ethane sulfonic acid, propane sulfonic acid, butane sulfonic acid, xylene sulfonic acid, benzene sulfonic acid, formic acid, acetic acid, mono, di, or tri-
10 halocarboxylic acids, picolinic acid, dipicolinic acid, and mixtures thereof. In some embodiments, the compositions of the present invention are free or substantially free of a phosphorous based acid.

In some embodiments, acidulant selected can also function as a stabilizing agent. Thus, the compositions of the present invention can be substantially free of an
15 additional stabilizing agent.

In certain embodiments, the present composition includes about 0.5 to about 80 wt-% acidulant, about 1 to about 50 wt%, about 5 to about 30 wt-% acidulant, or about 7 to about 14 wt-% acidulant. It is to be understood that all values and ranges between these values and ranges are encompassed by the compositions of the present invention.
20

Stabilizing Agents

In some embodiments, the compositions of the present invention include one or more stabilizing agents. The stabilizing agents can be used, for example, to stabilize the peracid and hydrogen peroxide and prevent the premature oxidation of this constituent
25 within the composition of the invention.

In some embodiments, an acidic stabilizing agent can be used. Thus, in some embodiments, the compositions of the present invention can be substantially free of an additional acidulant.

Suitable stabilizing agents include, for example, chelating agents or
30 sequestrants. Suitable sequestrants include, but are not limited to, organic chelating

compounds that sequester metal ions in solution, particularly transition metal ions. Such sequestrants include organic amino- or hydroxy-polyphosphonic acid complexing agents (either in acid or soluble salt forms), carboxylic acids (e.g., polymeric polycarboxylate), hydroxycarboxylic acids, aminocarboxylic acids, or heterocyclic
5 carboxylic acids, e.g., pyridine-2,6-dicarboxylic acid (dipicolinic acid).

In some embodiments, the compositions of the present invention include dipicolinic acid as a stabilizing agent. Compositions including dipicolinic acid can be formulated to be free or substantially free of phosphorous. It has also been observed that the inclusion of dipicolinic acid in a composition of the present invention aids in
10 achieving the phase stability of the compositions, compared to other conventional stabilizing agents, e.g., 1-hydroxy ethylidene-1,1-diphosphonic acid ($\text{CH}_3\text{C}(\text{PO}_3\text{H}_2)_2\text{OH}$) (HEDP).

In other embodiments, the sequesterant can be or include phosphonic acid or phosphonate salt. Suitable phosphonic acids and phosphonate salts include HEDP;
15 ethylenediamine tetrakis methylenephosphonic acid (EDTMP); diethylenetriamine pentakis methylenephosphonic acid (DTPMP); cyclohexane-1,2-tetramethylene phosphonic acid; amino[tri(methylene phosphonic acid)]; (ethylene diamine[tetra methylene-phosphonic acid]); 2-phosphene butane-1,2,4-tricarboxylic acid; or salts thereof, such as the alkali metal salts, ammonium salts, or alkylol amine salts, such as
20 mono, di, or tetra-ethanolamine salts; picolinic, dipicolinic acid or mixtures thereof. In some embodiments, organic phosphonates, e.g., HEDP are included in the compositions of the present invention.

Commercially available food additive chelating agents include phosphonates sold under the trade name DEQUEST® including, for example, 1-hydroxyethylidene-
25 1,1-diphosphonic acid, available from Monsanto Industrial Chemicals Co., St. Louis, MO, as DEQUEST® 2010; amino(tri(methylenephosphonic acid)), ($\text{N}[\text{CH}_2\text{PO}_3\text{H}_2]_3$), available from Monsanto as DEQUEST® 2000; ethylenediamine[tetra(methylenephosphonic acid)] available from Monsanto as DEQUEST® 2041; and 2-phosphonobutane-1,2,4-tricarboxylic acid available from

Mobay Chemical Corporation, Inorganic Chemicals Division, Pittsburgh, PA, as Bayhibit AM.

The sequestrant can be or include aminocarboxylic acid type sequestrant. Suitable aminocarboxylic acid type sequestrants include the acids or alkali metal salts thereof, e.g., amino acetates and salts thereof. Suitable aminocarboxylates include N-hydroxyethylaminodiacetic acid; hydroxyethylenediaminetetraacetic acid, nitrilotriacetic acid (NTA); ethylenediaminetetraacetic acid (EDTA); N-hydroxyethyl-
5 ethylenediaminetriacetic acid (HEDTA); diethylenetriaminopentaacetic acid (DTPA); and
10 alanine-N,N-diacetic acid; and the like; and mixtures thereof.

The sequestrant can be or include a polycarboxylate. Suitable polycarboxylates include, for example, polyacrylic acid, maleic/olefin copolymer, acrylic/maleic copolymer, polymethacrylic acid, acrylic acid-methacrylic acid copolymers, hydrolyzed polyacrylamide, hydrolyzed polymethacrylamide, hydrolyzed polyamide-
15 methacrylamide copolymers, hydrolyzed polyacrylonitrile, hydrolyzed polymethacrylonitrile, hydrolyzed acrylonitrile-methacrylonitrile copolymers, polymaleic acid, polyfumaric acid, copolymers of acrylic and itaconic acid, phosphino polycarboxylate, acid or salt forms thereof, mixtures thereof, and the like.

In certain embodiments, the present composition includes about 0.01 to about 10
20 wt-% stabilizing agent, about 0.4 to about 4 wt-% stabilizing agent, about 0.6 to about 3 wt-% stabilizing agent, about 1 to about 2 wt-% stabilizing agent. It is to be understood that all values and ranges within these values and ranges are encompassed by the present invention.

25 Wetting or Defoaming Agents

Also useful in the compositions of the invention are wetting and defoaming agents. Wetting agents function to increase the surface contact or penetration activity of the antimicrobial composition of the invention. Wetting agents which can be used in the composition of the invention include any of those constituents known within the art
30 to raise the surface activity of the composition of the invention.

Generally, defoamers which can be used in accordance with the invention include silica and silicones; aliphatic acids or esters; alcohols; sulfates or sulfonates; amines or amides; halogenated compounds such as fluorochlorohydrocarbons; vegetable oils, waxes, mineral oils as well as their sulfonated or sulfated derivatives; 5 fatty acids and/or their soaps such as alkali, alkaline earth metal soaps; and phosphates and phosphate esters such as alkyl and alkaline diphosphates, and tributyl phosphates among others; and mixtures thereof.

In some embodiments, the compositions of the present invention can include antifoaming agents or defoamers which are of food grade quality given the application 10 of the method of the invention. To this end, one of the more effective antifoaming agents includes silicones. Silicones such as dimethyl silicone, glycol polysiloxane, methylphenol polysiloxane, trialkyl or tetraalkyl silanes, hydrophobic silica defoamers and mixtures thereof can all be used in defoaming applications. Commercial defoamers commonly available include silicones such as Ardefoam® from Armour Industrial 15 Chemical Company which is a silicone bound in an organic emulsion; Foam Kill® or Kresseo® available from Krusale Chemical Company which are silicone and non-silicone type defoamers as well as silicone esters; and Anti-Foam A® and DC-200 from Dow Corning Corporation which are both food grade type silicones among others. These defoamers can be present at a concentration range from about 0.01 wt-% to 20 wt-%, from about 0.01 wt-% to 5 wt-%, or from about 0.01 wt-% to about 1 wt-%.

Thickening or Gelling Agents

The compositions of the present invention can include any of a variety of known thickeners. Suitable thickeners include natural gums such as xanthan gum, guar gum, or 25 other gums from plant mucilage; polysaccharide based thickeners, such as alginates, starches, and cellulosic polymers (e.g., carboxymethyl cellulose); polyacrylates thickeners; and hydrocolloid thickeners, such as pectin. In an embodiment, the thickener does not leave contaminating residue on the surface of an object. For example, the thickeners or gelling agents can be compatible with food or other sensitive 30 products in contact areas. Generally, the concentration of thickener employed in the

present compositions or methods will be dictated by the desired viscosity within the final composition. However, as a general guideline, the viscosity of thickener within the present composition ranges from about 0.1 wt-% to about 5 wt-%, from about 0.1 wt-% to about 1.0 wt-%, or from about 0.1 wt-% to about 0.5 wt-%.

5

Solidification Agent

The present compositions can include a solidification agent, which can participate in maintaining the compositions in a solid form. In some embodiments, the solidification agent can form and/or maintain the composition as a solid. In other
10 embodiments, the solidification agent can solidify the composition without unacceptably detracting from the eventual release of the sulfonated peroxycarboxylic acid. The solidification agent can include, for example, an organic or inorganic solid compound having a neutral inert character or making a functional, stabilizing or
15 deterrent contribution to the present composition. Suitable solidification agents include solid polyethylene glycol (PEG), solid polypropylene glycol, solid EO/PO block copolymer, amide, urea (also known as carbamide), nonionic surfactant (which can be employed with a coupler), anionic surfactant, starch that has been made water-soluble (e.g., through an acid or alkaline treatment process), cellulose that has been made water-
20 soluble, inorganic agent, poly(maleic anhydride/methyl vinyl ether), polymethacrylic acid, other generally functional or inert materials with high melting points, mixtures thereof, and the like;

Suitable glycol solidification agents include a solid polyethylene glycol or a solid polypropylene glycol, which can, for example, have molecular weight of about 1,400 to about 30,000. In certain embodiments, the solidification agent includes or is
25 solid PEG, for example PEG 1500 up to PEG 20,000. In certain embodiments, the PEG includes PEG 1450, PEG 3350, PEG 4500, PEG 8000, PEG 20,000, and the like. Suitable solid polyethylene glycols are commercially available from Union Carbide under the tradename CARBOWAX.

Suitable amide solidification agents include stearic monoethanolamide, lauric
30 diethanolamide, stearic diethanolamide, stearic monoethanol amide, cocodithylenc

amide, an alkylamide, mixtures thereof, and the like. In an embodiment, the present composition can include glycol (e.g., PEG) and amide.

Suitable nonionic surfactant solidification agents include nonylphenol ethoxylate, linear alkyl alcohol ethoxylate, ethylene oxide/propylene oxide block copolymer, mixtures thereof, or the like. Suitable ethylene oxide/propylene oxide block copolymers include those sold under the Pluronic tradename (e.g., Pluronic 108 and Pluronic F68) and commercially available from BASF Corporation. In some embodiments, the nonionic surfactant can be selected to be solid at room temperature or the temperature at which the composition will be stored or used. In other embodiments, the nonionic surfactant can be selected to have reduced aqueous solubility in combination with the coupling agent. Suitable couplers that can be employed with the nonionic surfactant solidification agent include propylene glycol, polyethylene glycol, mixtures thereof, or the like.

Suitable anionic surfactant solidification agents include linear alkyl benzene sulfonate, alcohol sulfate, alcohol ether sulfate, alpha olefin sulfonate, mixtures thereof, and the like. In an embodiment, the anionic surfactant solidification agent is or includes linear alkyl benzene sulfonate. In an embodiment, the anionic surfactant can be selected to be solid at room temperature or the temperature at which the composition will be stored or used.

Suitable inorganic solidification agents include phosphate salt (e.g., alkali metal phosphate), sulfate salt (e.g., magnesium sulfate, sodium sulfate or sodium bisulfate), acetate salt (e.g., anhydrous sodium acetate), Borates (e.g., sodium borate), Silicates (e.g., the precipitated or fumed forms (e.g., Sipernat 50® available from Degussa), carbonate salt (e.g., calcium carbonate or carbonate hydrate), other known hydratable compounds, mixtures thereof, and the like. In an embodiment, the inorganic solidification agent can include organic phosphonate compound and carbonate salt, such as an E-Form composition.

In some embodiments, the compositions of the present invention can include any agent or combination of agents that provide a requisite degree of solidification and aqueous solubility can be included in the present compositions. In other embodiments,

increasing the concentration of the solidification agent in the present composition can tend to increase the hardness of the composition. In yet other embodiments, decreasing the concentration of solidification agent can tend to loosen or soften the concentrate composition.

5 In some embodiments, the solidification agent can include any organic or inorganic compound that imparts a solid character to and/or controls the soluble character of the present composition, for example, when placed in an aqueous environment. For example, a solidifying agent can provide controlled dispensing if it has greater aqueous solubility compared to other ingredients in the composition. Urea
10 can be one such solidification agent. By way of further example, for systems that can benefit from less aqueous solubility or a slower rate of dissolution, an organic nonionic or amide hardening agent may be appropriate.

 In some embodiments, the compositions of the present invention can include a solidification agent that provides for convenient processing or manufacture of the
15 present composition. For example, the solidification agent can be selected to form a composition that can harden to a solid form under ambient temperatures of about 30 to about 50 °C after mixing ceases and the mixture is dispensed from the mixing system, within about 1 minute to about 3 hours, or about 2 minutes to about 2 hours, or about 5 minutes to about 1 hour.

20 The compositions of the present invention can include solidification agent at any effective amount. The amount of solidification agent included in the present composition can vary according to the type of composition, the ingredients of the composition, the intended use of the composition, the quantity of dispensing solution applied to the solid composition over time during use, the temperature of the dispensing
25 solution, the hardness of the dispensing solution, the physical size of the solid composition, the concentration of the other ingredients, the concentration of the cleaning agent in the composition, and other like factors. Suitable amounts can include about 1 to about 99 wt-%, about 1.5 to about 85 wt-%, about 2 to about 80 wt-%, about 10 to about 45 wt-%, about 15% to about 40 wt-%, about 20% to about 30 wt-%, about

30% to about 70%, about 40% to about 60%, up to about 50 wt-%, about 40% to about 50%

Carrier

In some embodiments, the compositions of the present invention include a
5 carrier. The carrier provides a medium which dissolves, suspends, or carries the other
components of the composition. For example, the carrier can provide a medium for
solubilization, suspension, or production of a sulfonated peroxycarboxylic acid and for
forming an equilibrium mixture. The carrier can also function to deliver and wet the
composition of the invention on an object. To this end, the carrier can contain any
10 component or components that can facilitate these functions.

In some embodiments, the carrier includes primarily water which can promote
solubility and work as a medium for reaction and equilibrium. The carrier can include
or be primarily an organic solvent, such as simple alkyl alcohols, e.g., ethanol,
isopropanol, n-propanol, benzyl alcohol, and the like. Polyols are also useful carriers,
15 including glycerol, sorbitol, and the like.

Suitable carriers include glycol ethers. Suitable glycol ethers include diethylene
glycol n-butyl ether, diethylene glycol n-propyl ether, diethylene glycol ethyl ether,
diethylene glycol methyl ether, diethylene glycol t-butyl ether, dipropylene glycol n-
butyl ether, dipropylene glycol methyl ether, dipropylene glycol ethyl ether,
20 dipropylene glycol propyl ether, dipropylene glycol tert-butyl ether, ethylene glycol
butyl ether, ethylene glycol propyl ether, ethylene glycol ethyl ether, ethylene glycol
methyl ether, ethylene glycol methyl ether acetate, propylene glycol n-butyl ether,
propylene glycol ethyl ether, propylene glycol methyl ether, propylene glycol n-propyl
ether, tripropylene glycol methyl ether and tripropylene glycol n-butyl ether, ethylene
25 glycol phenyl ether (commercially available as DOWANOL EPH™ from Dow
Chemical Co.), propylene glycol phenyl ether (commercially available as DOWANOL
PPH™ from Dow Chemical Co.), and the like, or mixtures thereof. Additional suitable
commercially available glycol ethers (all of which are available from Union Carbide
Corp.) include Butoxyethyl PROPASOL™, Butyl CARBITOL™ acetate, Butyl

CARBITOL™, Butyl CELLOSOLVE™ acetate, Butyl CELLOSOLVE™, Butyl
DIPROPASOL™, Butyl PROPASOL™, CARBITOL™ PM-600, CARBITOL™ Low
Gravity, CELLOSOLVE™ acetate, CELLOSOLVE™, Ester EEP™, FILMER IBT™,
Hexyl CARBITOL™, Hexyl CELLOSOLVE™, Methyl CARBITOL™, Methyl
5 CELLOSOLVE™ acetate, Methyl CELLOSOLVE™, Methyl DIPROPASOL™, Methyl
PROPASOL™ acetate, Methyl PROPASOL™, Propyl CARBITOL™, Propyl
CELLOSOLVE™, Propyl DIPROPASOL™ and Propyl PROPASOL™.

In some embodiments, the carrier makes up a large portion of the composition of
the invention and may be the balance of the composition apart from the sulfonated
10 peroxy-carboxylic acid, oxidizing agent, additional ingredients, and the like. The carrier
concentration and type will depend upon the nature of the composition as a whole, the
environmental storage, and method of application including concentration of the
sulfonated peroxy-carboxylic acid, among other factors. Notably the carrier should be
chosen and used at a concentration which does not inhibit the efficacy of the sulfonated
15 peroxy-carboxylic acid in the composition of the invention for the intended use, e.g.,
bleaching, sanitizing, disinfecting.

In certain embodiments, the present composition includes about 5 to about 90
wt-% carrier, about 10 to about 80 wt-% carrier, about 20 to about 60 wt-% carrier, or
about 30 to about 40 wt-% carrier. It is to be understood that all values and ranges
20 between these values and ranges are encompassed by the present invention.

Use Compositions

The compositions of the present invention include concentrate compositions and
use compositions. For example, a concentrate composition can be diluted, for example
25 with water, to form a use composition. In an embodiment, a concentrate composition
can be diluted to a use solution before to application to an object. For reasons of
economics, the concentrate can be marketed and an end user can dilute the concentrate
with water or an aqueous diluent to a use solution.

The level of active components in the concentrate composition is dependent on the intended dilution factor and the desired activity of the sulfonated peroxycarboxylic acid compound. Generally, a dilution of about 1 fluid ounce to about 10 gallons of water to about 10 fluid ounces to about 1 gallon of water is used for aqueous compositions of the present invention. In some embodiments, higher use dilutions can be employed if elevated use temperature (greater than 25 °C) or extended exposure time (greater than 30 seconds) can be employed. In the typical use locus, the concentrate is diluted with a major proportion of water using commonly available tap or service water mixing the materials at a dilution ratio of about 3 to about 40 ounces of concentrate per 10 100 gallons of water.

In some embodiments, when used in a laundry application, the concentrated compositions can be diluted at a dilution ratio of about 0.1g/L to about 100g/L concentrate to diluent, about 0.5g/L to about 10.0g/L concentrate to diluent, about 1.0g/L to about 4.0g/L concentrate to diluent, or about 1.0 g/L to about 2.0 g/L 15 concentrate to diluent.

In other embodiments, a use composition can include about 0.01 to about 10 wt-% of a concentrate composition and about 90 to about 99.99 wt-% diluent; or about 0.1 to about 1 wt-% of a concentrate composition and about 99 to about 99.9 wt-% diluent.

Amounts of an ingredient in a use composition can be calculated from the 20 amounts listed above for concentrate compositions and these dilution factors. In some embodiments, for example when used in a laundry application, the concentrated compositions of the present invention are diluted such that the sulfopercarboxylic acid is present at from about 20 ppm to about 80 ppm. In other embodiments, the concentrated compositions of the present invention are diluted such that the 25 sulfopercarboxylic acid is present at about 20 ppm, about 40 ppm, about 60 ppm, about 80 ppm, about 500 ppm, about 1000 ppm, or about 10,000 to about 20,000 ppm. It is to be understood that all values and ranges between these values and ranges are encompassed by the present invention.

Methods Employing the Sulfoperoxydicarboxylic Acid Compounds and Compositions

In some aspects, the present invention includes methods of using the sulfoperoxydicarboxylic acid compounds and compositions of the present invention. In some embodiments, these methods employ the antimicrobial and/or bleaching activity
5 of the sulfoperoxydicarboxylic acid. For example, the invention includes a method for reducing a microbial population, a method for reducing the population of a microorganism on skin, a method for treating a disease of skin, a method for reducing an odor, and/or a method for bleaching. These methods can operate on an article, surface, in a body or stream of water or a gas, or the like, by contacting the article,
10 surface, body, or stream with a sulfoperoxydicarboxylic acid compound or composition of the invention. Contacting can include any of numerous methods for applying a compound or composition of the invention, such as spraying the compounds or compositions, immersing the article in the compounds or compositions, foam or gel treating the article with the compounds or composition, or a combination thereof.

15 In some aspects, a composition of the present invention includes an amount of sulfoperoxydicarboxylic acid of the present invention effective for killing one or more of the food-borne pathogenic bacteria associated with a food product, including, but not limited to, *Salmonella typhimurium*, *Salmonella javiana*, *Campylobacter jejuni*, *Listeria monocytogenes*, and *Escherichia coli* O157:H7, yeast, and mold. In some
20 embodiments, the compositions of the present invention include an amount of sulfoperoxydicarboxylic acid effective for killing one or more of the pathogenic bacteria associated with a health care surfaces and environments including, but not limited to, *Salmonella typhimurium*, *Staphylococcus aureus*, methicillin resistant *Staphylococcus aureus*, *Salmonella choleraesuis*, *Pseudomonas aeruginosa*, *Escherichia coli*,
25 mycobacteria, yeast, and mold. The compounds and compositions of the present invention have activity against a wide variety of microorganisms such as Gram positive (for example, *Listeria monocytogenes* or *Staphylococcus aureus*) and Gram negative (for example, *Escherichia coli* or *Pseudomonas aeruginosa*) bacteria, yeast, molds, bacterial spores, viruses, etc. The compounds and compositions of the present
30 invention, as described above, have activity against a wide variety of human pathogens.

The present compounds and compositions can kill a wide variety of microorganisms on a food processing surface, on the surface of a food product, in water used for washing or processing of food product, on a health care surface, or in a health care environment.

5 The compounds of the invention can be used for a variety of domestic or industrial applications, e.g., to reduce microbial or viral populations on a surface or object or in a body or stream of water. The compounds can be applied in a variety of areas including kitchens, bathrooms, factories, hospitals, dental offices and food plants, and can be applied to a variety of hard or soft surfaces having smooth, irregular or porous topography. Suitable hard surfaces include, for example, architectural surfaces
10 (e.g., floors, walls, windows, sinks, tables, counters and signs); eating utensils; hard-surface medical or surgical instruments and devices; and hard-surface packaging. Such hard surfaces can be made from a variety of materials including, for example, ceramic, metal, glass, wood or hard plastic. Suitable soft surfaces include, for example paper; filter media; hospital and surgical linens and garments; soft-surface medical or surgical
15 instruments and devices; and soft-surface packaging. Such soft surfaces can be made from a variety of materials including, for example, paper, fiber, woven or nonwoven fabric, soft plastics and elastomers. The compounds of the invention can also be applied to soft surfaces such as food and skin (e.g., a hand). The present compounds can be employed as a foaming or nonfoaming environmental sanitizer or disinfectant.

20 The compounds and compositions of the invention can be included in products such as sterilants, sanitizers, disinfectants, preservatives, deodorizers, antiseptics, fungicides, germicides, sporicides, virucides, detergents, bleaches, hard surface cleaners, hand soaps, waterless hand sanitizers, and pre- or post-surgical scrubs.

25 The compounds can also be used in veterinary products such as mammalian skin treatments or in products for sanitizing or disinfecting animal enclosures, pens, watering stations, and veterinary treatment areas such as inspection tables and operation rooms. The present compounds can be employed in an antimicrobial foot bath for livestock or people. The compounds of the present invention can also be employed as an antimicrobial teat dip.

In some aspects, the compounds of the present invention can be employed for reducing the population of pathogenic microorganisms, such as pathogens of humans, animals, and the like. The compounds exhibit activity against pathogens including fungi, molds, bacteria, spores, and viruses, for example, *S. aureus*, *E. coli*, *Streptococci*,
5 *Legionella*, *Pseudomonas aeruginosa*, mycobacteria, tuberculosis, phages, or the like. Such pathogens can cause a variety of diseases and disorders, including mastitis or other mammalian milking diseases, tuberculosis, and the like. The compounds of the present invention can reduce the population of microorganisms on skin or other external or mucosal surfaces of an animal. In addition, the present compounds can kill pathogenic
10 microorganisms that spread through transfer by water, air, or a surface substrate. The compounds need only be applied to the skin, other external or mucosal surfaces of an animal water, air, or surface.

In some embodiments, the compounds and compositions of the present invention can be used to reduce the population of prions on a surface. Prions are
15 proteinaceous infectious particles free of nucleic acid. Prions are known to cause several brain diseases including kuru, Creutzfeldt-Jakob disease, Gerstmann-Straussler-Scheinker disease, and fatal familial insomnia in humans; scrapie in sheep; bovine spongiform encephalopathy (Mad Cow Disease) in cattle; transmissible mink encephalopathy in mink; chronic wasting disease in deer and elk; and feline spongiform
20 encephalopathy in cats. These diseases lead to symptoms including dementia, ataxia, behavioral disturbances, dizziness, involuntary movement, and death. Prions can be transmitted by exposure to infected tissue and brain tissue, spinal cord tissue, pituitary tissue, and eye tissue in particular. In some embodiments, the compounds and compositions of the present invention can be used to reduce a population of prions
25 according to a method as described in US Patent No. 7470655, the entire contents of which are hereby incorporated by reference.

The antimicrobial compounds can also be used on foods and plant species to reduce surface microbial populations; used at manufacturing or processing sites handling such foods and plant species; or used to treat process waters around such sites.
30 For example, the compounds can be used on food transport lines (e.g., as belt sprays);

boot and hand-wash dip-pans; food storage facilities; anti-spoilage air circulation systems; refrigeration and cooler equipment; beverage chillers and warmers, blanchers, cutting boards, third sink areas, and meat chillers or scalding devices. The compounds of the invention can be used to treat produce transport waters such as those found in
5 flumes, pipe transports, cutters, slicers, blanchers, retort systems, washers, and the like. Particular foodstuffs that can be treated with compounds of the invention include eggs, meats, seeds, leaves, fruits and vegetables. Particular plant surfaces include both harvested and growing leaves, roots, seeds, skins or shells, stems, stalks, tubers, corms, fruit, and the like. The compounds may also be used to treat animal carcasses to reduce
10 both pathogenic and non-pathogenic microbial levels.

The antimicrobial compounds can also be used to treat waste water where both its antimicrobial function and its oxidant properties can be utilized. Aside from the microbial issues surrounding waste water, it is often rich in malodorous compounds of reduced sulfur, nitrogen or phosphorous. A strong oxidant such as the present invention
15 converts these compounds efficiently to their odor free derivatives e.g. the sulfates, phosphates and amine oxides. These same properties are very useful in the pulp and paper industry where the property of bleaching is also of great utility.

In some aspects, the compounds of the present invention can be employed for epoxidations. The polymer industry is a major consumer of peracids, especially
20 peroxyacetic acid but the typical equilibrium peroxyacetic acid also includes some strong acid residues which are problematic for the epoxide derivatives. A stable peracid isolate is therefore potentially of great utility in this industry.

In some aspects, the compounds and compositions of the present invention are useful in the cleaning or sanitizing of containers, processing facilities, or equipment in
25 the food service or food processing industries. The compounds and compositions have particular value for use on food packaging materials and equipment, and especially for cold or hot aseptic packaging. Examples of process facilities in which the compound of the invention can be employed include a milk line dairy, a continuous brewing system, food processing lines such as pumpable food systems and beverage lines, etc. Food
30 service wares can be disinfected with the compound of the invention. For example, the

compounds can also be used on or in ware wash machines, low temperature ware wash machines, dishware, bottle washers, bottle chillers, warmers, third sink washers, cutting areas (e.g., water knives, slicers, cutters and saws) and egg washers. Particular treatable surfaces include packaging such as cartons, bottles, films and resins; dish ware such as
5 glasses, plates, utensils, pots and pans; ware wash and low temperature ware wash machines; exposed food preparation area surfaces such as sinks, counters, tables, floors and walls; processing equipment such as tanks, vats, lines, pumps and hoses (e.g., dairy processing equipment for processing milk, cheese, ice cream and other dairy products); and transportation vehicles. Containers include glass bottles, PVC or polyolefin film
10 sacks, cans, polyester, PEN or PET bottles of various volumes (100 ml to 2 liter, etc.), one gallon milk containers, paper board juice or milk containers, etc.

The compounds and compositions can also be used on or in other industrial equipment and in other industrial process streams such as heaters, cooling towers, boilers, retort waters, rinse waters, aseptic packaging wash waters, and the like. The
15 compounds can be used to treat microbes and odors in recreational waters such as in pools, spas, recreational flumes and water slides, fountains, and the like.

A filter containing the compound can reduce the population of microorganisms in air and liquids. Such a filter can remove water and air-born pathogens such as
Legionella.

20 The present compounds can be employed for reducing the population of microbes, fruit flies, or other insect larva on a drain or other surface.

The compounds of the present invention can also be employed by dipping food processing equipment into the use solution, soaking the equipment for a time sufficient to sanitize the equipment, and wiping or draining excess solution off the equipment,
25 The compound may be further employed by spraying or wiping food processing surfaces with the use solution, keeping the surfaces wet for a time sufficient to sanitize the surfaces, and removing excess solution by wiping, draining vertically, vacuuming, etc.

The compounds of the present invention may also be used in a method of sanitizing hard surfaces such as institutional type equipment, utensils, dishes, health care equipment or tools, and other hard surfaces.

5 The antimicrobial compounds can be applied to microbes or to soiled or cleaned surfaces using a variety of methods. These methods can operate on an object, surface, in a body or stream of water or a gas, or the like, by contacting the object, surface, body, or stream with a compound of the invention. Contacting can include any of numerous methods for applying a compound, such as spraying the compound, immersing the object in the compound, foam or gel treating the object with the
10 compound, or a combination thereof.

A concentrate or use concentration of a compound of the present invention can be applied to or brought into contact with an object by any conventional method or apparatus for applying an antimicrobial or cleaning compound to an object. For example, the object can be wiped with, sprayed with, foamed on, and/or immersed in
15 the compound, or a use solution made from the compound. The compound can be sprayed, foamed, or wiped onto a surface; the compound can be caused to flow over the surface, or the surface can be dipped into the compound. Contacting can be manual or by machine. Food processing surfaces, food products, food processing or transport waters, and the like can be treated with liquid, foam, gel, aerosol, gas, wax, solid, or
20 powdered stabilized compounds according to the invention, or solutions containing these compounds.

Laundry Applications

In some aspects, the compounds can also be employed in sanitizing articles, e.g.,
25 textiles, which have become contaminated. The articles are contacted with the compounds of the invention at use temperatures in the range of about 4 °C to 80 °C, for a period of time effective to sanitize, disinfect, and/or sterilize the articles. In some embodiments, the compounds of the present invention can be used to bleach and/or sanitize articles at a temperature of about 30°C to about 50°C or about 40°C. For
30 example, in some embodiments, the compounds of the present invention can be injected

into the wash or rinse water of a laundry machine and contacted with contaminated fabric for a time sufficient to sanitize the fabric. In some embodiments, the contaminated fabric is contacted with the compounds and compositions of the present invention for about 5 to about 30 minutes. Excess solution can then be removed by
5 rinsing or centrifuging the fabric.

In some aspects, the compounds of the present invention can be used as a bleaching agent to whiten or lighten or remove stains from a substrate, e.g., hard surface, or fabric. The compounds of the present invention can be used to bleach or remove stains from any conventional textile, including but not limited to, cotton, poly-
10 cotton blends, wool, and polyesters. The compounds of the present invention are also textile tolerant, i.e., they will not substantially degrade the textile to which they are applied. The compounds of the present invention can be used to remove a variety of stains from a variety of sources including, but not limited to, lipstick, pigment/sebum, pigment/lanolin, soot, olive oil, mineral oil, motor oil, blood, make-up, red wine, tea,
15 ketchup, and combinations thereof.

In some embodiments, the compounds of the present invention can be used as a low odor, acidic bleaching agent. In some embodiments, the compounds of the present invention can be used as a low odor bleaching agent at a neutral pH, i.e., about 7. In some embodiments, the compounds of the present invention can be used at an alkaline
20 pH, e.g., about 8, 9, or 10. In still yet other embodiments, the compounds of the present invention can be used as an all in one sour, bleaching and sterilant product.

The compounds and compositions of the present invention can be used alone to treat the articles, e.g., textiles, or can be used in conjunction with conventional detergents suitable for the articles to be treated. The compounds and compositions of
25 the invention can be used with conventional detergents in a variety of ways, for example, the compounds and compositions of the invention can be formulated with a conventional detergent. In other embodiments, the compounds and compositions of the invention can be used to treat the article as a separate additive from a conventional detergent. When used as a separate additive, the compounds and compositions of the
30 present invention can contact the article to be treated at any time. For example, the

compounds and compositions of the invention can contact the article before, after, or substantially simultaneously as the articles are contacted with the selected detergent.

In some embodiments, when used as a bleaching and/or sanitizing/disinfecting agent for a laundry application, a compound or mixture of compounds of the present invention will be present in a composition at about 5 ppm to about 1000ppm. In other embodiments, when used as a bleaching and/or sanitizing/disinfecting agent for a laundry application, a compound or mixture of compounds of the present invention will be present in a composition at about 25ppm to about 100 ppm. In other embodiments, when used as a bleaching and/or sanitizing/disinfecting agent in a laundry application, a compound or mixture thereof of the present invention will be present at about 20, about 40, about 60, or about 80ppm. In still yet other embodiments, a compound or mixture of compounds of the present invention itself will be used as a bleaching agent, i.e., the compound or mixture of compounds will be present in a composition at about 100 wt%.

15 *Clean in Place*

Other hard surface cleaning applications for the compounds of the present invention include clean-in-place systems (CIP), clean-out-of-place systems (COP), washer-decontaminators, sterilizers, textile laundry machines, ultra and nano-filtration systems and indoor air filters. COP systems can include readily accessible systems including wash tanks, soaking vessels, mop buckets, holding tanks, scrub sinks, vehicle parts washers, non-continuous batch washers and systems, and the like. CIP systems include the internal components of tanks, lines, pumps and other process equipment used for processing typically liquid product streams such as beverages, milk, juices.

Generally, the actual cleaning of the in-place system or other surface (i.e., removal of unwanted offal therein) is accomplished with a different material such as a formulated detergent which is introduced with heated water. After this cleaning step, the instant composition would be applied or introduced into the system at a use solution concentration in unheated, ambient temperature water. CIP typically employ flow rates on the order of about 40 to about 600 liters per minute, temperatures from ambient up to about 70°C, and contact times of at least about 10 seconds, for example, about 30 to

about 120 seconds. The present composition can remain in solution in cold (e.g., 40°F/4°C.) water and heated (e.g., 140°F/60° C.) water. Although it is not normally necessary to heat the aqueous use solution of the present composition, under some circumstances heating may be desirable to further enhance its activity. These materials
5 are useful at any conceivable temperatures.

A method of sanitizing substantially fixed in-place process facilities includes the following steps. The use solution of the invention is introduced into the process facilities at a temperature in the range of about 4 °C to 60 °C. After introduction of the use solution, the solution is held in a container or circulated throughout the system for a
10 time sufficient to sanitize the process facilities (e.g., to kill undesirable microorganisms). After the surfaces have been sanitized by means of the present composition, the use solution is drained. Upon completion of the sanitizing step, the system optionally may be rinsed with other materials such as potable water. The composition can be circulated through the process facilities for 10 minutes or less.

15 The present method can include delivering the present composition via air delivery to the clean-in-place or other surfaces such as those inside pipes and tanks. This method of air delivery can reduce the volume of solution required.

Contacting a Food Product with the Sulfoperoxy-carboxylic Acid Compounds

20 In some aspects, the present invention provides methods for contacting a food product with a sulfoperoxy-carboxylic acid compounds or composition employing any method or apparatus suitable for applying such a compound or composition. For example, in some embodiments, the food product is contacted by a compound of the present invention with a spray of the compound, by immersion in the compound, by
25 foam or gel treating with the compound. Contact with a spray, a foam, a gel, or by immersion can be accomplished by a variety of methods known to those of skill in the art for applying antimicrobial agents to food. Contacting the food product can occur in any location in which the food product might be found, such as field, processing site or plant, vehicle, warehouse, store, restaurant, or home. These same methods can also be
30 adapted to apply the compounds of the present invention to other objects.

The present methods require a certain minimal contact time of the compound with food product for occurrence of significant antimicrobial effect. The contact time can vary with concentration of the use compound, method of applying the use compound, temperature of the use compound, amount of soil on the food product,
5 number of microorganisms on the food product, type of antimicrobial agent, or the like. The exposure time can be at least about 5 to about 15 seconds. In some embodiments, the exposure time is about 15 to about 30 seconds. In other embodiments, the exposure time is at least about 30 seconds.

In some embodiments, the method for washing a food product employs a
10 pressure spray including a compound of the present invention. During application of the spray solution on the food product, the surface of the food product can be moved with mechanical action, e.g., agitated, rubbed, brushed, etc. Agitation can be by physical scrubbing of the food product, through the action of the spray solution under pressure, through sonication, or by other methods. Agitation increases the efficacy of
15 the spray solution in killing micro-organisms, perhaps due to better exposure of the solution into the crevasses or small colonies containing the micro-organisms. The spray solution, before application, can also be heated to a temperature of about 15 to 20 °C, for example, about 20 to 60 °C to increase efficacy. The spray stabilized compound can be left on the food product for a sufficient amount of time to suitably reduce the
20 population of microorganisms, and then rinsed, drained, or evaporated off the food product.

Application of the material by spray can be accomplished using a manual spray wand application, an automatic spray of food product moving along a production line using multiple spray heads to ensure complete contact, or other spray apparatus. One
25 automatic spray application involves the use of a spray booth. The spray booth substantially confines the sprayed compound to within the booth. The production line moves the food product through the entryway into the spray booth in which the food product is sprayed on all its exterior surfaces with sprays within the booth. After a complete coverage of the material and drainage of the material from the food product
30 within the booth, the food product can then exit the booth. The spray booth can include

steam jets that can be used to apply the stabilized compounds of the invention. These steam jets can be used in combination with cooling water to ensure that the treatment reaching the food product surface is less than 65°C, e.g., less than 60°C. The temperature of the spray on the food product is important to ensure that the food product is not substantially altered (cooked) by the temperature of the spray. The spray pattern can be virtually any useful spray pattern.

Immersing a food product in a liquid stabilized compound of the present invention can be accomplished by any of a variety of methods known to those of skill in the art. For example, the food product can be placed into a tank or bath containing the stabilized compound. Alternatively, the food product can be transported or processed in a flume of the stabilized compound. The washing solution can be agitated to increase the efficacy of the solution and the speed at which the solution reduces micro-organisms accompanying the food product. Agitation can be obtained by conventional methods, including ultrasonics, aeration by bubbling air through the solution, by mechanical methods, such as strainers, paddles, brushes, pump driven liquid jets, or by combinations of these methods. The washing solution can be heated to increase the efficacy of the solution in killing micro-organisms. After the food product has been immersed for a time sufficient for the desired antimicrobial effect, the food product can be removed from the bath or flume and the stabilized compound can be rinsed, drained, or evaporated off the food product.

In other embodiments, a food product can be treated with a foaming version of the compound of the present invention. The foam can be prepared by mixing foaming surfactants with the washing solution at time of use. The foaming surfactants can be nonionic, anionic or cationic in nature. Examples of useful surfactant types include, but are not limited to the following: alcohol ethoxylates, alcohol ethoxylate carboxylate, amine oxides, alkyl sulfates, alkyl ether sulfate, sulfonates, including, for example, alkyl aryl sulfonates, quaternary ammonium compounds, alkyl sarcosines, betaines and alkyl amides. The foaming surfactant is typically mixed at time of use with the washing solution. Use solution levels of the foaming agents is from about 50 ppm to about 2.0 wt-%. At time of use, compressed air can be injected into the mixture, then applied to

the food product surface through a foam application device such as a tank foamer or an aspirated wall mounted foamer.

In some embodiments, a food product can be treated with a thickened or gelled version of a compound of the present invention. In the thickened or gelled state the washing solution remains in contact with the food product surface for longer periods of
5 time, thus increasing the antimicrobial efficacy. The thickened or gelled solution will also adhere to vertical surfaces. The compound or the washing solution can be thickened or gelled using existing technologies such as: xanthan gum, polymeric thickeners, cellulose thickeners, or the like. Rod micelle forming systems such as
10 amine oxides and anionic counter ions could also be used. The thickeners or gel forming agents can be used either in the concentrated product or mixing with the washing solution, at time of use. Typical use levels of thickeners or gel agents range from about 100 ppm to about 10 wt-%.

15 *Methods for Beverage, Food, and Pharmaceutical Processing*

The sulfoperoxy-carboxylic acid compounds of the present invention can be used in the manufacture of beverage, food, and pharmaceutical materials including fruit juice, dairy products, malt beverages, soybean-based products, yogurts, baby foods, bottled water products, teas, cough medicines, drugs, and soft drinks. The compounds
20 of the present invention can be used to sanitize, disinfect, act as a sporicide for, or sterilize bottles, pumps, lines, tanks and mixing equipment used in the manufacture of such beverages. Further, the sulfoperoxy-carboxylic acid antimicrobial compounds of the present invention can be used in aseptic, cold filling operations in which the interior of the food, beverage, or pharmaceutical container is sanitized or sterilized prior to
25 filling. In such operations, a container can be contacted with the sanitizing sulfoperoxy-carboxylic acid compound, typically using a spray, dipping, or filling device to intimately contact the inside of the container with the sulfoperoxy-carboxylic acid compound, for a sufficient period of time to reduce microorganism populations within the container. The container can then be emptied of the amount of sanitizer or sterilant
30 used. After emptying, the container can be rinsed with potable water or sterilized water

and again emptied. After rinsing, the container can be filled with the beverage, food, or pharmaceutical. The container can then be sealed, capped or closed and then packed for shipment for ultimate sale. The sealed container can be autoclaved or retorted for added microorganism kill.

5 In food, beverage, or pharmaceutical manufacturing, fungal microorganisms of the genus *Chaetomium* or *Arthrimum*, and spores or bacteria of the genus *Bacillus spp.* can be a significant problem in bottling processes, particularly in cold aseptic bottling processes. The sulfoperoxycarboxylic acid compounds of the present invention can be used for the purpose of controlling or substantially reducing (by more than a 5 log₁₀
10 reduction) the number of *Chaetomium* or *Arthrimum* or *Bacillus* microorganisms in beverage or food or pharmaceutical bottling lines using cold aseptic bottling techniques.

 In such techniques, metallic, aluminum or steel cans can be filled, glass bottles or containers can be filled, or plastic (PET or PBT or PEN) bottles, and the like can be filled using cold aseptic filling techniques. In such processes, the
15 sulfoperoxycarboxylic acid materials of the invention can be used to sanitize the interior of beverage containers prior to filling with the carbonated (or noncarbonated) beverage. Typical carbonated beverages in this application include, but are not limited to, cola beverages, fruit beverages, ginger ale beverages, root beer beverages, iced tea beverages which may be non-carbonated, and other common beverages considered soft drinks.
20 The sulfoperoxycarboxylic acid materials of the invention can be used to sanitize both the tanks, lines, pumps, and other equipment used for the manufacture and storage of the soft drink material and also used in the bottling or containers for the beverages. In an embodiment, the sulfoperoxycarboxylic acid sanitizing materials are useful for killing both bacterial and fungal microorganisms that can be present on the surfaces of
25 the production equipment and beverage containers.

 The sulfoperoxycarboxylic acid compounds of the present invention can effectively kill microorganisms (e.g., > 1 log₁₀ or up to about 5 log₁₀ reduction in 30 seconds) from a concentration level of at least about 50 ppm, for example, about 150, about 500 ppm or about 1000 ppm of a sulfoperoxycarboxylic acid compound. In an
30 embodiment, the sulfoperoxycarboxylic acid compound, excluding water, would be

present at a concentration of about 0.001 to about 1 wt-%, for example, about 0.01 to about 0.15 wt-%, or about 0.05 to about 0.1 wt-%.

All acid, salt, base and other ionic and non-ionic forms of the compounds described are included as compounds of the invention. For example, if a compound is shown as an acid herein, the salt forms of the compound are also included. Likewise, if a compound is shown as a salt, the acid and/or basic forms are also included.

Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, numerous equivalents to the specific procedures, embodiments, claims, and examples described herein. Such equivalents are considered to be within the scope of this invention and covered by the claims appended hereto. The contents of all references, patents, and patent applications cited throughout this application are hereby incorporated by reference. The invention is further illustrated by the following examples, which should not be construed as further limiting.

15

EXAMPLES

Some of the following Examples were performed using a sulfonated peroleic acid product. Without wishing to be bound by any particular theory, it is thought that the peracid formed from a commercially available sulfonated oleic acid starting material includes a mixture of the compounds of the present invention. It is thought that this is due, in part, to the nature of the sulfonated oleic acid starting material. That is, it is thought that because the sulfonated oleic acid starting material is derived from naturally occurring sources, it is not chemically pure, i.e., does not contain only one form of the sulfonated oleic acid. Thus, without wishing to be bound by any particular theory it is thought that sulfonated peroleic acid (hereinafter referred to as the "sulfonated peroleic acid product") used in these examples included a mixture of about 20-25 wt% Compound A (10-Hydroxy-9-sulfooctadecaneperoxoic acid) about 20-25 wt% Compound N (10,11-dihydroxy-9-sulfooctadecaneperoxoic acid), about 20-25 wt% Compound I (9-Hydroxy-10-sulfooctadecaneperoxoic acid), and about 20-25 wt% Compound O (8,9-dihydroxy-10-sulfooctadecaneperoxoic acid). The remainder of the

peracid composition is thought to include about 5 to about 10 wt% of a mixture of these compounds.

Example 1- Use of a Sulfoperoxycarboxylic acid as a Coupler under High Level

5 *Disinfection Application Conditions*

Peroxyoctanoic acid (POOA) stability experiments were performed under high level disinfection (HLD) conditions to evaluate the stability of a composition of the present invention including a sulfonated peroleic acid product, compared with known commercially available disinfectants.

- 10 Octave FS®, a peroxyoctanoic containing product, commercially available from Ecolab Inc. was tested against Formulas A, B, and C, and mixtures thereof. Formula A was a mixture of: 2.5 wt% Dequest 2010 (commercially available from thermPhos), peracid grade; 61wt% hydrogen peroxide (35%); 2.50 wt% sulfuric acid (98%); 6.0 wt% octanoic acid, 19 wt% Hostapur SAS (40%) (commercially available from
- 15 Clariant); and 9.00 wt% SXS-40 (commercially available from Stepan Company). Formula B was a mixture of about 20 wt% of the sulfonated peroleic acid product, about 10 % peroctanoic acid, about 15 wt% octanoic acid, and about 0.5 wt% hydrogen peroxide. Formula C was a mixture of about 25 wt% of the sulfonated peroleic acid product, and about 0.50 wt% hydrogen peroxide. Mixtures of Formulas A, B, and C
- 20 were also tested. The test solutions were diluted with DI water to make a solution with about 1000ppm POOA present at a pH of about 6.5. The table below shows the five solutions tested, and the amount of sulfonated peroleic acid product, POOA, and hydrogen peroxide available in ppm in each of the solutions as tested.

Table 2.

	Test solution composition				
	#1	#2	#3	#4	#5
Octave FS® (wt%)	10.00	0	0	0	0
Formula A (wt%)	0	4.2	0	0	0
Formula B (wt%)	0	0	0.88	0.55	0.33

Formula C (wt%)	0	0	0.22	0.55	0.77
Final weight with added DI water (g)	100	100	100	100	100
Sulfonated peroleic acid product (ppm)	0	0	2318	2459	2554
POOA (ppm)	1000	1000	800	500	300
H ₂ O ₂	8050	8928	55	55	55

The samples were stored at 40°C and the amount of POOA present was measured by high performance liquid chromatography at the selected times. The following table shows the results of the HPLC analysis of the samples at various times.

5 *Table 3.*

	Test solution				
	1	2	3	4	5
Time (hrs)	POOA (ppm)	POOA (ppm)	POOA (ppm)	POOA (ppm)	POOA (ppm)
0	490	870	700	470	290
6	310	730	590	400	250
24	0	120	350	240	150
48	0	10	240	160	100
72	0	0	180	130	80
9 days	0	0	20	0	0

10 These results are also graphically depicted in Figure 1. As can be seen from the table above, and Figure 1, the test solutions including a compound of the present invention, i.e., test solutions 3, 4, and 5, lost less POOA over the course of the first 24 hours compared to the other two test solutions. Even after 48 hours, a greater amount of POOA remained in the test solutions including a compound of the present invention,

than in the other solutions tested. For each of the test solutions including a compound of the present invention, it was shown that the loss of POOA in the solutions was not linear, and that the decomposition rate of POOA slowed down dramatically at higher ratios of the sulfonated peroleic acid product to POOA.

5 Another stability study was performed to evaluate the stability of a composition of the present invention at an elevated temperature, i.e., 100°F. A solution including about 2 wt% of the sulfonated peroleic acid product, and about 55 wt% H₂O₂, among other ingredients, was used. The amount of the sulfonated peroleic acid product and H₂O₂ was measured over the course of 48 days. The results are shown in Figure 2. As
10 can be seen in this figure, the peracid compound, the sulfonated peroleic acid product maintained its activity over the course of the trial, even at this accelerated temperature.

Yet another stability study was performed to evaluate the stability of peroxyoctanoic acid when contacted by a compound of the present invention, i.e., the sulfonated peroleic acid product, under ambient conditions. For this study, the pH was
15 constant at about 6 to about 6.5. Three different formulas were tested for this study: Formula D included about 5 grams of a mixture of the sulfonated peroleic acid product , peroxyoctanoic acid, hydrogen peroxide and sodium cumene sulfate, among other ingredients; Formula E included about 0.5g of a mixture of the sulfonated peroleic acid product , and peroxyoctanoic acid; and Formula F included Octave®, commercially
20 available from Ecolab Inc. The amount of active peroxyoctanoic acid available at various times over the course of 15 days was measured. The results are shown in the table below.

Table 4.

	Formula D	Formula E	Formula F
Time (days)	POOA (ppm)	POOA (ppm)	POOA (ppm)
0	590	640	570
1	550	590	500
4	470	480	360
6	420	400	240

8	410	360	160
11	360	270	70
14	310	230	30

These results are also graphically depicted in Figure 3. As can be seen in this table, and figure, the formulas including a compound of the present invention, i.e., Formulas D and E, retained a higher level of POOA over the course of 15 days. Thus, without wishing to be bound by any particular theory it is thought that the addition of a composition including compounds of the present invention acts to stabilize other percarboxylic acids present in the composition.

Example 2- Use of a Sulfoperoxy-carboxylic Acid as a Bleaching Agent

The use of a compound of the present invention as a bleaching agent was evaluated. The soil removal ability of the cleaning composition was determined by washing with artificially soiled fabric swatches. The soiled swatches were purchased from a manufacturer or distributor (e.g. Test Fabrics, Inc., West Pittston, Pa.). Soil types such as olive oil, sebum, makeup, wine are characteristic of natural soils found in laundry applications.

Soiled swatches were washed with the cleaning composition in a device such as a Terg-o-tometer (United States Testing Co., Hoboken, N.J.). The Terg-o-tometer is a laboratory washing device that consists of multiple pots that reside in a single temperature-controlled water bath, with overhead agitators under time and speed control. Wash test parameters include: wash temperature, wash duration, pH, mechanical agitation, dose of cleaning composition, water hardness, wash formula, and cloth/liquor ratio. After completing the appropriate exposure times the fabric samples were removed. The test chemistries were immediately flushed, and the swatches rinsed with cold synthetic 5 grain water until 5 cycles of fills and rinses were complete. The swatches were then laid flat and dried overnight on white polyester-cotton towels before reflectance readings were taken using a spectrophotometer, e.g., Hunter ColorQuest XE (reflectance) Spectrophotometer.

To determine the percent (%) soil removal (SR), e.g., bleaching ability, the reflectance of the fabric sample was measured on a spectrophotometer. The “L value” is a direct reading supplied by the spectrophotometer. L generally is indicative of broad visible spectrum reflectance, where a value of 100% would be absolute white. The % soil removal is calculated from the difference between the initial (before washing) lightness (L) value and the final L value (after washing):

$$SR = ((L_{final} - L_{initial}) / (96 - L_{initial})) \times 100\%$$

A bleach test was run comparing a composition including a sulfonated peroleic acid product with the following commercially available bleaching /cleaning compositions: Ozonit®, and Oxysan® both available from Ecolab Inc. Ozonit® represents a 4.5% peroxyacetic acid product while Oxysan® represents a 0.6% peroxyoctanoic acid product. Formula A was a composition including about 2 wt% of sulfonated peroleic acid product, about 5 wt% peroxyacetic acid and about 1.5 wt% of peroxyoctanoic acid. Formula A was used at a concentration of 1200ppm and further treated in two of the three cases with additional acetic acid to produce lowered pH test solutions. Ozonit® was used at a concentration of 2000ppm. Oxysan® was tested at concentrations of 1272 and 2545ppm. All of the wash solutions were further treated with Detergent MP® and TurboCharge II ®, both available from Ecolab Inc and used at 500 and 750ppm respectively. The bath/wash temperature was maintained at 100°F. Detergent MP® and TurboCharge II provide a common alkaline builder detergent base. The results from the bleaching test are shown in the table below.

Table 5.

Bleach Type	Stain Removal (%) from Cotton			Conc. of Bleach (mg/L)	pH
	Tea	Red Wine	Ketchup		
Ozonit®	29	59	27	2000	9.50
Oxysan®	21	66	19	1272	8.00
1X Oxysan®	33	69	27	2545	8.00
Formula A, pH 8.0	37	73	38	1000	8.00
Formula A, pH 8.5	38	72	41	1000	8.50
Formula A, pH 9.0	34	69	36	1000	9.00

As can be seen from this table, the compositions of Formula A achieved a higher percent stain removal than the commercially available solutions tested at all pH levels tested, especially in the cases of ketchup which represents a hydrophobic stain.

Formula A was also tested using a full scale Wash Wheel Bleach Test. The test was run with a commercial 35 lb side loading washing machine (*UniMac UX35PVXR*). Multipaneled pre-stained test sheets (Ecomon No.1 & Ecomon No.4 included 14 bleachable and 12 pigment/unbleachable stained panels) were added to the otherwise empty machine before initiating a 20 minute washing program (typically at 40° C). The chemistries were added in a 30 second staggered sequence via the overhead dispensing cups once the machine was filled with 48L of 5 grain synthetic soft water. The initial chemistry added was the alkaline detergent product (about 84g of Turbocmulsion, commercially available from Ecolab Inc.). The bleaching chemistry was then added ~30 seconds after the surfactant-caustic blend and a 20 minute wash cycle was begun. After

the wash cycle the machine was drained and 3 rinse cycles were executed. The sheets were retrieved and air dried at 70° F, overnight before measuring each swatch panel's reflectance with a Hunter ColorQuest XE (reflectance) Spectrophotometer (UV filter "TN"). The results are shown in the table below.

5 Table 6.

	I. Reflectance Values				³ Stain Removal, %	
	Initial stained swatch	³ Turboemulsion only	TE + ⁶ Formula A	TE + ⁴ Ozonit	TE + Formula A	TE + Ozonit
<i>Bleachable Stains</i>						
Tea on CO	80.64	80.67	91.62	88.94	71.48	54.01
Tea on PES/CO	80.43	79.24	91.17	88.28	68.96	50.40
Red Wine on CO	73.66	85.94	93.03	92.06	86.72	82.36
Red Wine on PES/CO, aged	73.82	82.98	91.71	90.67	80.67	75.97
Coffee on CO	78.92	90.72	93.10	92.70	83.04	80.70
Coffee on PES/CO	79.77	92.27	93.62	93.28	85.34	83.26
Black currant juice on CO	64.40	88.37	93.54	92.82	92.22	89.94
Black currant juice on PES/CO	63.57	85.02	93.30	92.07	91.68	87.89
Blood on CO IEC 456, aged	46.25	89.51	90.60	91.48	89.14	90.91
Blood on CO IEC 456, not aged	49.36	93.06	93.81	93.88	95.30	95.45
Blood / Milk / Ink on CO	45.26	61.00	51.10	51.89	11.51	13.06
Cocoa on CO IEC 456, not aged	75.22	83.76	83.47	83.27	39.72	38.74
Blood / Milk / Soot on CO	58.87	86.37	69.87	70.54	29.62	31.44

Egg / Soot on CO	62.87	76.36	76.09	75.81	39.89	39.05
average / 14	66.65	83.95	86.15	85.55	68.95	65.23
Unbleachable Stains						
Pigment / Lanolin on CO	71.98	80.90	78.63	80.55	27.70	35.68
Pigment / Lanolin on PES / CO	66.65	82.38	73.28	81.72	22.60	51.35
Pigment / Sebum on CO	73.19	87.70	84.02	86.76	47.49	59.48
Pigment / Sebum on PES / CO	70.64	87.97	77.82	86.74	28.33	63.49
Soot / Olive Oil on CO	47.93	69.90	62.45	64.87	30.21	35.23
Soot / Olive Oil on PES / CO	40.77	62.89	56.23	58.57	27.99	32.23
Soot / Mineral Oil on CO	59.76	72.35	68.93	71.80	25.30	33.21
Soot / Mineral Oil on PES / CO	55.62	80.15	73.89	78.78	45.25	57.36
Used Motor Oil on CO	65.91	73.06	70.99	71.77	16.89	19.47
Used Motor Oil on PES / CO	61.10	68.27	64.08	66.01	8.53	14.08
Makeup on CO	84.81	90.06	89.50	90.14	41.94	47.63
Makeup on PES / CO	85.16	92.57	91.91	92.14	62.24	64.42
average / 12	70.85	86.01	81.49	84.62	32.04	42.80

Notes: 3. Turboemulsion (TE) is a commercially available all-in-one emulsion of alkaline metal chelators emulsified with a surfactant blend made by Ecolab, Inc. and was used in this test at 1750ppm. 4. Ozonit is a Peracetic acid - Hydrogen peroxide bleach disinfectant used at a concentration of 2000ppm. Ozonit is a blend of Peracetic acid and Hydrogen peroxide made by Ecolab, Inc.. 5. The "Stain Removal, %" was calculated using the following formula: $SR = ((I_{final} - I_{initial}) / (96 - I_{initial})) \times 100\%$ CO: Cotton; PES/CO: Polyester-Cotton blend

As can be seen from this table, Formula A averages superior bleaching to Ozonit®. Although the superiority on these “bleachable” stains is only 3.7 points (5.4%), on those stains which better resist wash removal e.g. tea, the difference was as many as 17 points (24%) higher.

- 5 Another full scale wash testing was conducted using a wash wheel (full size side loading washing machine), but rather than individual soiled swatches this test utilized multipanled sheets combining 14 “bleachable” stained swatches (Ecomon 4) and a second sheet which combined 12 “unbleachable” pigment/hydrocarbon stained swatches (Ecomon 1). These panels are custom made for Ecolab by wfk Testgewebe
- 10 Gmbh of Bruggen, Germany. This extensive bleach test utilized a design experiment which varied concentrations sometimes simultaneously with temperatures etc. Following completion of the specified wash time, all Ecomon sheets were rinsed thoroughly, dried and their broad spectrum light reflectivities were measured, again with UV filtering to removal possible interference from optical brightener effects.
- 15 Unlike the tergotometer data, the % stain removal wasn’t calculated but was rather directly measured from the reflectance instrument (*Minolta CM-2610d Spektrophotometer*). A “Y” value representing broad spectrum reflectivity was reported. The higher the “Y” value, the whiter the material, and therefore, the greater the bleaching or stain removal.
- 20 In this test, Formula A was compared to Ozonit®, Ozonit Super® (a 15% peroxyacetic acid product available from Ecolab) and Oxysan® these were variously combined with the following commercially available alkaline-builder cleaning agents: Triplex Emulsion®, available from Ecolab Inc.; Turbo Usona®, available from Ecolab Inc.; Ozonit Super®, available from Ecolab Inc.; and Oxysan®, available from Ecolab
- 25 Inc. The results are shown in the tables below.

Table 7. Bleaching Results

Procedure	Tea on CO	Tea on PES/CO	Red Wine on CO aged	Red Wine on PES/CO aged	Coffee on CO	Coffee on PES/CO	Black Currant Juice on CO	Black Currant Juice on PES/CO	Ave.
1.5 ml/l ² Triplex Emulsion + 1ml/l Formula A Conditions: 15' 40°C	72.7	70.0	75.4	74.6	80.2	84.6	82.5	84.1	78.0
1.5 ml/l Triplex Emulsion + 2ml/l Formula A Conditions: 15' 40°C	80.6	79.6	82.5	80.5	83.5	86.0	85.5	86.1	83.0
1.5 ml/l Triplex Emulsion + 2.5ml/l Formula A Conditions: 20' 40°C	82.6	83.1	84.3	80.9	84.3	86.0	86.2	86.3	84.2
1.5 ml/l Triplex Emulsion + 1ml/l Ozonit Super Conditions: 10'70°C	78.5	79.0	82.2	82.1	84.8	86.2	86.7	86.5	83.3

4 ml/l ³ Turbo Usona + 2ml/l ⁴ Ozonii Performance Conditions: 20' 40°C	80.8	80.5	81.5	79.0	81.1	84.2	82.0	80.8	81.2
4 ml/l Turbo Usona + 4ml/l ⁵ Oxyvan Conditions: 20' 40°C	79.2	77.9	78.9	76.3	79.9	83.3	77.6	75.9	78.6
4 ml/l Turbo Usona + 2ml/l Formula A Conditions: 15' 40°C	82.2	81.7	82.1	80.5	82.6	85.2	82.6	82.6	82.4
LSD	1.8	3	1.9	2.4	1.1	0.8	1.7	1.8	1.9

Table 8. Bleaching results

	1.5 ml/l ² Triple x Emulsi on + 1 ml/l Formul a A Condit ions: 15' 40°C	1.5 ml/l <i>Triplex Emulsion</i> + 2ml/l Formula A Conditio ns: 15' 40°C	1.5 ml/l <i>Triplex Emulsion</i> + 2.5ml/l Formula A Conditio ns: 20' 40°C	1.5 ml/l <i>Triplex Emulsion</i> + 1ml/l <i>Ozonit Super</i> Conditio ns: 10'70°C	4 ml/l ³ Turbo Usona + 2ml/l ⁴ Ozonit <i>Performan ce</i> Conditio ns: 20' 40°C	4 ml/l <i>Turbo Usona +</i> 4ml/l ⁵ Oxysan Conditio ns: 20' 40°C	4 ml/l <i>Turbo Usona</i> + 2ml/l Formula A Conditio ns: 15' 40°C	LSD
Pig men t/ Lan olin on CO	54.3	55.6	56.8	67.3	57.5	56.6	54.8	6.1
Pig men t/ Lan olin on PES /CO	53.4	51.4	48.8	60.6	46.1	44.9	46.2	5.8
Pig men t/ Lan olin on PES /CO	68.3	59.7	59.6	67.5	60.1	58.0	60.9	6.7

Sebum on CO								
Pigment / Sebum on PES /CO	66.0	54.2	54.7	73.4	53.2	50.5	54.3	6.3
Soot / Olive Oil on CO	47.7	42.5	32.2	46.9	24.7	25.1	24.3	7.8
Soot / Olive Oil on PES /CO	33.8	28.5	24.2	38.4	15.7	14.4	13.0	9.6
Soot / Min. Oil on CO	36.9	34.3	36.0	34.0	33.4	30.6	30.6	4.5

Soot / Min. Oil on PES /CO	42.4	43.9	35.8	46.6	31.0	32.7	37.7	8.2
Used Motor Oil on CO	42.7	43.9	42.6	46.0	44.0	44.9	46.3	2.6
Used Motor Oil on PES /CO	37.7	34.7	33.7	36.4	32.2	33.5	33.8	1.6
Makeup on CO	75.3	74.1	75.7	84.1	73.3	72.4	73.5	4
Makeup on PES	79.8	77.9	76.8	86.6	75.7	74.0	76.9	3.8

/CO								
Lip-stick on CO	87.6	87.4	87.3	87.7	85.9	86.9	87.3	1.4
Lip-stick on PES /CO								3.04 7061 8
Average	53.2	50.1	48.1	57.3	45.6	44.8	46.0	5.6

Notes:

1. Y-value refers to a reflectance value calculated by the Minolta CM-2610d Spektrophotometer. It is very similar to the L-value calculated by the Hunter Lab's Spectrophotometers.
 2. Triplex Emulsion is a commercially available all-in-one emulsion of alkaline metal chelators emulsified with a surfactant blend by Ecolab, Inc. (Europe).
 3. Turbo Usona is a commercially available all-in-one emulsion of alkaline metal chelators emulsified with a surfactant blend made by Ecolab, Inc. (Europe).
 4. Ozonit Super is a Peracetic acid -Hydrogen peroxide bleach disinfectant, made by Ecolab, Inc. (Europe).
 5. Oxysan is a Peracetic acid -Hydrogen peroxide bleach disinfectant which also contains Peroxyoctanoic acid, and is made by Ecolab, Inc. (Europe).
- CO: Cotton
PES/CO: Polyester-Cotton blend

As can be seen from these results, overall the samples washed with compositions of the present invention, i.e., Formula A, achieved similar bleaching compared with commercially available bleaching agents.

5

Example 3- Use of a Sulfoperoxydicarboxylic Acid as a Bleaching Agent

A bleach test was run comparing a composition including a sulfoperoxydicarboxylic acid of the present invention, i.e., 11-sulfoundecaneperoxydicarboxylic acid (Compound D) with the following commercially available bleaching/cleaning

compositions: Tsunami 100®, available from Ecolab Inc.; Oxonia Active®, available from Ecolab Inc.; hydrogen peroxide (35%); and PAP-70®, available from Solvay. These chemistries were used as is except for pH adjustments to pH 8 using sodium bicarbonate, and pH 12 by the addition of sodium hydroxide, in 5 grain hardwater.

- 5 Fabric swatches soiled with tea, blood, or wine were used for this example. The soil swatches were washed using the same experimental procedure described above in Example 2. However, for this example, the soil swatches were washed for 10 minutes at 120°F. The pH of the wash solution for all samples was about 9. The percent soil removal (SR) was determined according to the method described above in Example 2.
- 10 The following table shows the results of this study.

Table 9.

Removal of Tea Stains						
Bleach Type	pH	Temp(F)	Wash Time (min)	%SR	Bleach mg/L use solution	Use Solution Available Oxygen (ppm)
Composition Including Compound D	9	120	10	37	1350	56
Tsunami 100®	9	120	10	34	770	56
Oxonia Active®	9	120	10	27	410	56
H ₂ O ₂ (35%)	9	120	10	24	340	56
PAP-70	9	120	10	63	1386	56
Water	9	120	10	11	0	56

(control)						
Removal of Blood Stains						
Bleach Type	pH	Temp(F)	Wash Time (min)	%SR	Bleach mg/L use solution	Use Solution Available Oxygen (ppm)
Composition Including Compound D	9	120	10	90	1350	56
Tsunami 100®	9	120	10	81	770	56
Oxonia Active®	9	120	10	80	410	56
H ₂ O ₂ (35%)	9	120	10	82	340	56
PAP-70	9	120	10	88	1386	56
Water (control)	9	120	10	36	0	0

Removal of Red Wine Stains						
Bleach Type	pH	Temp(F)	Wash Time (min)	%SR	Bleach use solution mg/L	Use Solution Available Oxygen (ppm)
Composition including Compound D	9	120	10	62	1350	56
Tsunami 100®	9	120	10	57	770	56
Oxonia Active®	9	120	10	41	410	56
H ₂ O ₂ (35%)	9	120	10	45	340	56
PAP-70	9	120	10	74	1386	56
Water (control)	9	120	10	36	0	56

As can be seen from this table, with respect to tea stains, the PAP-70® composition achieved the greatest soil removal. The composition containing a compound of the present invention achieved the next highest percent soil removal.

- 5 With respect to blood stains, the composition containing the sulfoperoxy-carboxylic acid of the present invention achieved the greatest soil removal. However, all concentrated oxidizers performed well in removing the blood stains. With respect to the red wine stains, the sulfoperoxy-carboxylic acid of the present invention performed well compared to the PAP-70®.

Example 4- Stability Studies

The stability of a sulfoperoxydicarboxylic acid of the present invention, i.e., 11-sulfoundecaneperoxydicarboxylic acid (Compound D), was compared to that of phthalimidoperoxyhexanoic acid (PAP). The stability data for the PAP sample were taken from U.S. Patent No. 5,994,284, assigned to Clariant GmbH. Samples of the compound of the present invention were stored for four (4) weeks at various temperatures. The loss of active oxygen was measured by titrimetry. The results are shown in the table below.

Table 10.

Compound	Storage Time (weeks)	Temperature (°C)	Loss of Active Oxygen (%)
Compound D	4	Room Temp.	0.78
Compound D	4	38	7.9
Compound D	4	50	15.7
PAP	4	25	1.4
PAP	4	40	2.0
PAP	4	50	12.0

10

As can be seen from this table, the compound of the present invention was more stable, i.e., lost less active oxygen, at room temperature, i.e., about 23°C, than the PAP at 25°C.

15 *Example 5- Bleaching Performance of Various Formulas of the Present Invention*

A test was run to compare the bleaching properties of compositions of the present invention with the following commercially available bleaching agents: Ozonit®, available from Ecolab Inc.; and PAP®, available from Clariant. The following compositions of the present invention were used: Formula A, which included about 25 wt% of the sulfonated peroleic acid product, about 70 wt% H₂O₂ (35%), and about 5 wt% HEDP 60; Formula B which included about 24 wt% of a mixture of the sulfonated peroleic acid product and peroxyoctanoic acid, about 72 wt% H₂O₂ (35%), and about 4

20

wt% HEDP 60; and Formula C which included about 20 wt% of a mixture of the sulfonated peroleic acid product and peroxyoctanoic acid, about 62 wt% H₂O₂ (35%), about 4 wt% HEDP 60, and about 13 wt% acetic acid. These formulas were compared with the commercially available bleaching agents at 40°C at a pH of between 7 to 8.

5 The Ozonit® was also tested at 60°C.

To measure the bleaching ability of the formulations, a bleaching test as described in Example 2 was performed. The results are shown in Figure 4. As can be seen in this figure, Formulas A, B and C had far superior bleaching ability compared to Ozonit® at 40°C. When the Ozonit® was used at 60°C, Formulas A, B, and C had very similar bleaching ability. Formula C also had similar bleaching performance compared to the PAP. Thus, Formulas A, B, and C showed equal, if not better, bleaching properties compared to known commercially available bleaching agents at 40°C.

Example 6- Antimicrobial Studies

15 (a) Bactericidal Efficacy

An experiment was performed to determine the bactericidal efficacy of a composition according to the present invention, with and without a surfactant, as compared to other commercially available products. Formula A included about 1190ppm of a sulfonated peroleic acid product, as well as peroxyoctanoic acid, and peracetic acid. The surfactant used for this example was Turboemulsion® (TE), commercially available from Ecolab Inc. The compositions were tested against *Clostridium difficile* ATCC 9689, MRSA ATCC 33592, *Enterococcus hirac* ATCC 10541, *Escheria coli* ATCC 11229, and *Pseudomonas aeruginosa* ATCC 15442, at 5 and 60 minute exposure times. The commercially available compositions, Ozonit®, and PAP were also tested. The following formulations were tested:

25 *Table 11.*

Test Formulation	Desired Concentration of Active Agent	Diluent	Test Use Solution (Volume of Test Substance/Total	pH

			Volume)	
Formula A with surfactant	1190ppm	Sterile MilliQ Water	0.194g Formula A + 170µL TE/100g	7.59
Formula A without Surfactant	1190ppm		0.194g Formula A /100g	8.53
PAP®	1820ppm		0.182g PAP + 1.5g TE/ 100g	8.50
Ozonit®	2000		0.200g Ozonit + 1.5g TE/100g	7.21

The test method followed was according to European Standard EN 13704: *Quantitative Suspension Test for the Evaluation of Sporicidal Activity of Chemical Disinfectants and Antiseptics Used in Food, Industrial, Domestic and Institutional Areas*. Generally, a test suspension of bacterial spores in a solution of interfering substance, simulating clean conditions, was added to a prepared sample of the test formulation diluted in hard water. The mixture was maintained at the specific temperature and time desired. At this contact time, an aliquot is taken; the sporicidal action in this portion was immediately neutralized or suppressed by a validated method.

5 The number of surviving bacterial spores in each sample was determined and the reduction in viable counts was calculated.

The disinfectant properties of each of the formulations at 5 minutes at 40°C is shown below in Table 12.

15 Table 12.

Test / System	Formula A with Surfactant	PAP	Ozonit	Formula A without Surfactant
---------------	---------------------------	-----	--------	------------------------------

MRSA	>6.66	>6.66	>6.66	>6.66
Enterococcus hirae ATCC 10541	>6.26	>6.26	>6.26	>6.26
Escherichia coli ATCC 11229	>6.74	>6.74	>6.74	>6.74
Pseudomonas aeruginosa ATCC 15442	>6.32	>6.32	>6.32	>6.32
Clostridium difficile ATCC 9689	>3.87	1.17	2.57	3.09

As can be seen from this table, the compositions of the present invention that were tested were as effective as a disinfectant as the commercially available formulations tested. Further, with respect to *Clostridium difficile*, the compositions of the present invention were more effective than the commercially available products tested.

(b) *Stability and sporicidal efficacy at 14 days*

A test was run to determine the stability and sporicidal efficacy of a composition of the present invention against spores. The composition tested included the sulfonated peroleic acid product, and an amount of peroxyoctanoic acid. The test method used was the European Standard EN 13704: *Quantitative Suspension Test for the Evaluation of Sporicidal Activity of Chemical Disinfectants and Antiseptics Used in Food, Industrial, Domestic and Institutional Areas*, described above. The table below shows the results of this study.

Table 13.

Sulfonated Peroleic Acid Product + POOA (14 Day Retention)	
DI Water	
pH 6.5	
B. subtilis	C. difficile
Clean Conditions	
20°C	
60 min	60 min
Log reduction 3.84	Log reduction 2.71

A composition including 30ppm peroxyoctanoic acid was also tested. The composition of peroxyoctanoic acid alone did not result in a reduction.

Figure 5 shows the stability impact that the compound of the present invention used, i.e., the sulfonated peroleic acid product, had on the amount of POOA over time during this study. As can be seen from this figure, the amount of POOA available over time was higher with the sample of POOA that was stabilized using a composition of the present invention, compared to a sample of POOA that was not stabilized using a composition of the present invention.

10

(c) Synergistic Effect of a Composition of the Present Invention with a Known Sanitizer

For this study, the ASME 1052-96: Standard Test Method for Efficacy of Antimicrobial Agents against Viruses in Suspension was used. A composition including 1000ppm peroxyacetic acid (POAA) was tested alone, and in combination with sulfonated peroleic acid product.

The POAA solution alone did not display complete inactivation of Poliovirus Type 1 after an exposure time of four minutes. The reductions in viral titer were ≤ 0.75 and $\leq 0.50 \log 10$. When the POAA solution was tested with 1000ppm of the sulfonated peroleic acid product, the solution displayed complete inactivation of

20

Poliovirus Type 1 after an exposure time of a few minutes, and was therefore efficacious against the virus. The reduction in viral titer was $\geq 5.75 \log_{10}$.

(d) Synergistic Effect of a Compound of the Present Invention with

5 Peroxyoctanoic Acid

For this study, the MS103: Quantitative Tuberculocidal Test was used. The sulfonated peroleic acid product was tested alone, and in combination with peroxyoctanoic acid at various concentrations against *Mycobacterium bovis* BCG. The compositions were tested at a pH of 6.5 at room temperature. The results are shown in
10 the table below.

Table 14.

Test Substance	Exposure Time	Log Reduction
1000ppm Sulfonated Peroleic Acid Product	2.5 min	4.46
	5 min	5.11
300ppm POOA	2.5 min	3.48
	5 min	<4.31
1000ppm Sulfonated Peroleic Acid Product and 300ppm POOA	2.5 min	>7.31
	5 min	>7.31
1000ppm Sulfonated Peroleic Acid Product and 150ppm POOA	2.5 min	≥ 7.31
	5 min	>7.31

As can be seen from this table, the samples treated with both a composition of
15 the present invention including the sulfonated peroleic acid product, and POOA had a higher log reduction of *Mycobacterium bovis* BCG than those samples treated with either the sulfonated peroleic acid product or POOA alone. Although it was found that

the samples treated with just the sulfonated peroleic acid product did have a higher log reduction of bacteria than the samples treated with just POOA.

(e) Use of a Compound of the Invention as a Hospital Disinfectant

- 5 For this test, the AOAC Official Method 955.15- Testing Disinfectant Against *Staphylococcus aureus* and the AOAC Official Method 964.02- Testing Disinfectants Against *Pseudomonas aeruginosa* were used. The composition used included the sulfonated peroleic acid product, and peroxyoctanoic acid (POOA), at various concentrations. The following chart summarizes the test procedure used, and the
- 10 results.

Table 15.

Test Substance	Desired Concentration	Diluent	Dilution (Volume of Test System / Total Volume)	Test pH
Sulfonated Peroleic Acid Product + POOA	1000ppm Sulfonated Peroleic Acid Product 300ppm POOA	400ppm Synthetic	2.910g Sulfonated Peroleic Acid Product + 0.2345g POOA / 1500g	6.5
	1000ppm Sulfonated Peroleic Acid Product 150ppm POOA	Hard Water	0.1852g Sulfonated Peroleic Acid Product + 0.4690g POOA / 1500g	6.5
Test system	Test Substance		# Negative Tubes / # Carriers Tested	

<i>Staphylococcus aureus</i> ATCC 6538	1000ppm Sulfonated Peroleic Acid Product + 300ppm POOA	60 / 60
<i>Staphylococcus aureus</i> ATCC 6538	1000ppm Sulfonated Peroleic Acid Product + 150ppm POOA	60 / 60
<i>Pseudomonas aeruginosa</i> ATCC 15442	1000ppm Sulfonated Peroleic Acid Product + 300ppm POOA	60 / 60
<i>Pseudomonas aeruginosa</i> ATCC 15442	1000ppm Sulfonated Peroleic Acid Product + 150ppm POOA	60 / 60

As can be seen from this table, the compositions tested were effective against each of the test systems.

5 *Example 7- Coupling Abilities of Compounds of the Present Invention*

The ability of a composition of the present invention including the sulfonated peroleic acid product to couple octanoic acid was compared to the coupling abilities of two known commercially available coupling agents, NAS and linear alkylbenzene sulphonate (LAS).

- 10 The results can be seen in Figure 6. As can be seen from this figure, one gram of the sulfonated peroleic acid product was able to couple twice as much octanoic acid compared to the other coupling agents tested.



Example 8 – Formation of Sulfonated Carboxylic Acids and Their Pericarboxylic Salts

- 15 A study was run to determine the effect of the position of the sulfonate group on the carboxylic acid in forming a peracid. Specifically, a study was run to determine whether having the sulfonate group at the α position prohibits the oxidation and/or perhydrolysis of the carboxylic acid group to form the corresponding peroxycarboxylic acid.

- 20 Commercially available sulfonated fatty acid salts (methyl esters) are predominantly α sulfonated, including, for example, Alpha-Step PC-48 (commercially



available from the Stepan Comp.), Alpha-Step MC-48 (MC-48)(commercially available from the Stepan Comp.), Alpha-Step BSS-45 (commercially available from the Stepan Comp.), and MES (commercially available from the Lion Corporation). Structurally, these compounds are sodium alphasulfo methyl C₁₂-C₁₈ esters and disodium alphasulfo

- 5 C₁₂-C₁₈ fatty acid salts. Their structures are shown below:



Sulfonated oleic acid is another commercially available sulfonated fatty acid. These compounds are mainly 8-sulfo-octadecenoic acid salts, with a minority of 9-sulfo-10-hydroxy-octadecanoic acid salts. They are not sulfonated at the α position.

- 10 The structures of these types of compounds are shown below:



α -sulfonated fatty acids were prepared by the hydrolysis of the mixture of α -sulfonated fatty acid methyl ester and the acid (MC-48). To a beaker containing 25g of

MC-48, 12g of 50% NaOH solution was added. The mixture was stirred at ambient temperatures for 3 hours. The mixture was then acidified by adding H₂SO₄ (50%) until the pH of the mixture reached about 0-1. The white solid precipitate was filtered, washed with cold water, and dried. The white solid powder yield was evaluated using
5 ¹³C NMR (DMSO-d₆). The methyl group of the methyl ester in the raw material was not observed, indicating complete hydrolysis.

In order to try and form the peracid using an acid catalyzed hydroxide reaction the following reaction was performed. 0.5g of the MC-48 derived fatty acid sulfonate, as prepared above, was weighed into a 50ml beaker. To this beaker, 30g of H₂O₂ (35%)
10 was added. then, 5g of H₂SO₄ (985) was slowly added, producing a clear solution. After sitting at 50°C for 24 hours, the solution was analyzed to determine the presence of a peracid.

To determine the presence of a peracid, a kinetic iodometric titration similar to the method disclosed in Sully and Williams ("The Analysis of Per-Acids and Hydrogen
15 Peroxide," *The Analyst*, 87:1037, p. 653 (Aug. 1962)) was used. This method has demonstrated a lower detection limit of about 0.3 ppm for POAA. Given the molecular weight ratio of POAA to the perspective percarboxylic acid of PC-48, the detection limit was estimated to be about 1.4ppm (3.93 X 10⁻⁶ M). No peracid formation was observed. This is equivalent to a percarboxylic acid formation constant (K_{eq}) less than
20 0.002, suggesting substantially no peracid was formed.

Alternatively, formation of the peracid was determined using ¹³C NMR (D₂O). Using this technique, no carbonyl resonance signal from the peracid was observed.

Other α-sulfonated fatty acid sources such as Alpha-Step PC-48 and Alpha-Step BSS-45 were also reacted with H₂O₂ in a similar manner, and in both cases, no
25 corresponding peracids were detected.

Non- α-sulfonated fatty acids were also tested to determine the likelihood of peracid formation. For the sulfonated oleic acid discussed above, the measured formation constant was 1.42. The sulfonated undecenoic acid was collected as a stable solid powder, so the formation constant was not measured. Although the formation
30 constant of the peracid of sulfonated oleic acid is significantly lower than that of the

most common commercialized peracid, peroxyacetic acid ($K_{eq} = 2.70$), it is still high enough to make practical yields.

Overall, without wishing to be bound by any particular theory, it is thought that the α -sulfo group prohibits the oxidation and/or perhydrolysis of the carboxylic acid group by H_2O_2 to the corresponding peracid. This may in part be due to its strong electron withdrawing effects.

Example 9 – Clean in Place Sanitizing Compositions

A study was run to determine the efficacy of compositions of the present invention as sanitizers used in a clean in place cleaning method. A composition including about 5.85 wt% of the sulfonated peroleic acid product, and about 11.6% hydrogen peroxide, about 1wt% of a chelating agent, about 12.75wt% of H_2SO_4 , about 13.6wt% NAS-FAL (sodium octane sulfonate), and about 1.5wt% of SXS (commercially available from the Stepan Company) was prepared. Synthetic hard water was used to dilute the test composition to the desired peracid concentration. The peracid was tested at concentrations of 1000ppm, 750ppm and 500ppm. The pH of the use solutions were as follows:

Concentration of Peracid in Use Solution	pH
500ppm peracid	1.65
750ppm	1.46
1000ppm	1.38

The use solutions were tested against *Staphylococcus aureus* ATCC 6538 and *Pseudomonas aeruginosa* ATCC 15442. The organic soil used was 5% Fetal Bovine Serum. The exposure time of the test was 5 minutes at a temperature of $20 \pm 1^\circ C$. A neutralizer screen was also performed as part of the testing to verify that the neutralizer adequately neutralized the product and was not detrimental to the tested organisms. The plates were incubated at $35^\circ C$ for 48 hours with the test systems prior to exposure to the peracids. The results are shown in the table below.

Table 16.

Staphylococcus aureus ATCC 6538

Test Substance	# Negative Tubes / # Carriers Tested
1000ppm Peracid Composition	60 / 60

5

Pseudomonas aeruginosa ATCC 15442

Test Substance	# Negative Tubes / # Carriers Tested
1000ppm Peracid Composition	60 / 60

Test Controls

10

Control	Test System	Results
Negative Carrier		1 negative of 1 tested
Positive Carrier	<i>Staphylococcus aureus</i> ATCC 6538	1 positive of 1 tested
Positive Carrier	<i>Pseudomonas aeruginosa</i> ATCC 15442	1 positive of 1 tested
Organic Soil		1 negative of 1 tested
Neutralization (1000ppm)	<i>Staphylococcus aureus</i> ATCC 6538	6 positive of 6 tested
Neutralization (1000ppm)	<i>Pseudomonas aeruginosa</i> ATCC 15442	6 positive of 6 tested
Culture Enumeration	<i>Staphylococcus aureus</i> ATCC 6538	9.0×10^8 CFU/mL
Culture Enumeration	<i>Pseudomonas aeruginosa</i> ATCC 15442	1.0×10^9 CFU/mL
Carrier Enumeration	<i>Staphylococcus aureus</i> ATCC 6538	1.0×10^6 CFU/mL 1.0×10^7 CFU/Carrier
Carrier Enumeration	<i>Pseudomonas aeruginosa</i> ATCC 15442	2.3×10^6 CFU/mL 2.3×10^7 CFU/Carrier

Staphylococcus aureus ATCC 6538

Test Substance	# Negative Tubes / # Carriers Tested
----------------	--------------------------------------

500ppm Peracid Composition	59 / 60
750ppm Peracid Composition	60 / 60

Pseudomonas aeruginosa ATCC 15442

Test Substance	# Negative Tubes / # Carriers Tested
500ppm Peracid Composition	58 / 60
750ppm Peracid Composition	60 / 60

5 Test Controls

Control	Test System	Results
Negative Carrier		1 negative of 1 tested
Positive Carrier	<i>Staphylococcus aureus</i> ATCC 6538	1 positive of 1 tested
Positive Carrier	<i>Pseudomonas aeruginosa</i> ATCC 15442	1 positive of 1 tested
Organic Soil		1 negative of 1 tested
Neutralization	<i>Staphylococcus aureus</i> ATCC 6538	3 positive of 3 tested
Neutralization	<i>Pseudomonas aeruginosa</i> ATCC 15442	3 positive of 3 tested
Culture Enumeration	<i>Staphylococcus aureus</i> ATCC 6538	1.0 x 10 ⁹ CFU/mL
Culture Enumeration	<i>Pseudomonas aeruginosa</i> ATCC 15442	1.0 x 10 ⁹ CFU/mL
Carrier Enumeration	<i>Staphylococcus aureus</i> ATCC 6538	7.2 x 10 ⁵ CFU/mL 7.2 x 10 ⁶ CFU/Carrier
Carrier Enumeration	<i>Pseudomonas aeruginosa</i> ATCC 15442	2.0 x 10 ⁶ CFU/mL 2.0 x 10 ⁷ CFU/Carrier

As can be seen from these results, the use solutions tested were effective disinfectants against both *Staphylococcus aureus*, and *Pseudomonas aeruginosa* at the concentrations tested.

Another study was run to determine the sanitizing efficacy of the test solution against *Staphylococcus aureus* ATCC 6538 and *Escherichia coli* ATCC 11229 after a 30 second exposure time. For this experiment the solutions were diluted to have a

concentration of 50ppm, 75ppm or 100ppm of the sulfonated peroleic acid product.
 The pH of the use solutions were as follows:

Concentration of Peracid in Use Solution	pH
50ppm peracid	2.70
75ppm	2.47
100ppm	2.30

The use solutions were tested against *Staphylococcus aureus* ATCC 6538 and
 5 *Escherichia coli* ATCC 11229. The exposure time was 30 seconds at a temperature of
 25 ± 1°C. A neutralizer screen was also performed as part of the testing to verify that
 the neutralizer adequately neutralized the product and was not detrimental to the tested
 organisms. The plates were incubated at 35°C for 48 hours with the test systems prior
 to exposure to the peracids. The results are shown in the table below.

10

Table 17.

Inoculum Numbers

Test System	CFU/mL	Log ₁₀ Growth	Average Log ₁₀ Growth
<i>Staphylococcus aureus</i> ATCC 6538	107 x 10 ⁶ , 109 x 10 ⁶	8.03, 8.04	8.04
<i>Escherichia coli</i> ATCC 11229	138 x 10 ⁶ , 151 x 10 ⁶	8.14, 8.18	8.16

15

***Staphylococcus aureus* ATCC 6538**

Test Substance	Survivors (CFU/mL)	Log ₁₀ Growth	Average Log ₁₀ Growth	Log Reduction
50ppm Peracid Composition	28 x 10 ¹ , 20 x 10 ¹	2.45, 2.30	2.38	5.66
75ppm Peracid	0 x 10 ¹ , 100 x 10 ¹	<1.00, 3.00	<2.00	>6.04

Composition				
100ppm Peracid Composition	$0 \times 10^1, 0 \times 10^1$	<1.00, <1.00	<1.00	>7.04

Escherichia coli ATCC 11229

Test Substance	Survivors (CFU/mL)	Log ₁₀ Growth	Average Log ₁₀ Growth	Log Reduction
50ppm Peracid Composition	$0 \times 10^1, 2 \times 10^1$	<1.00, 1.30	<1.15	>7.01
75ppm Peracid Composition	$0 \times 10^1, 0 \times 10^1$	<1.00, <1.00	<1.00	>7.16
100ppm Peracid Composition	$0 \times 10^1, 0 \times 10^1$	<1.00, <1.00	<1.00	>7.16

5

As can be seen from these results the use solutions tested were effective sanitizers against both *Staphylococcus aureus* and *Escherichia coli*. The test solution containing 100ppm of the sulfonated peroleic acid product was the most effective sanitizer.

10

Example 10 – Foam Properties of Selected Compositions of the Present Invention

A study was performed to determine the foam properties of selected compositions of the present invention, compared to compositions including commercially available surfactants. The following compositions were prepared:

- 15 Formula A included 50ppm of the sulfonated peroleic acid product at a pH of 2.48;
 Formula B included 50ppm of the sulfonated peroleic acid product at a pH6.75;
 Formula C included 64ppm of a commercially available sulfonated oleic acid (SOA)(Lankropol OPA (50%) available from Akzo Nobel) at a pH of 2.48; Formula D
 20 included 64ppm of a commercially available sulfonated oleic acid (Lankropol OPA (50%) available from Akzo Nobel) at a pH of 6.56; Formula E included 128ppm of a commercially available sulfonated oleic acid (Lankropol OPA (50%) available from

Akzo Nobel) at a pH of 2.48; Formula F included 128ppm of a commercially available sulfonated oleic acid (Lankropol OPA (50%) available from Akzo Nobel) at a pH of 7.20; and Formula G included 93ppm of sodium octane sulfonate (NAS) (commercially available from Ecolab) at a pH of 2.48. The foam heights were determined using the following method. First 3000 ml of each formula was prepared and gently poured into Glewwe cylinder. A ruler was attached to the side of the cylinder, and the solution was level with the bottom of the ruler. The pump was turned on. Foam height was estimated by reading the average level of foaming according to the ruler. Foam height readings were taken versus time with a stopwatch or timer. The pump was turned off and height of the foam was recorded at various times. The results are shown in the table below.

Table 18.

Sample	Pump On Time (sec)			Pump Off Time (sec)		
	30	60	300	30	60	300
	Foam Height (inches)	Foam Height (inches)	Foam Height (inches)	Foam Height (inches)	Foam Height (inches)	Foam Height (inches)
Formula A	2.5	3.8	5.5	3.5	2.0	0.5
Formula B	1.5	2.0	2.5	0.2	<0.1	NA
Formula C	4.0	6.2	9.2	8.7	8.5	5.5
Formula D	3.1	4.5	10	9.8	8.5	4.0
Formula E	2.6	4.5	8.5	8.2	8.0	5.0
Formula F	0.15	0.15	0.2	<0.1	<0.1	<0.1
Formula G	1.0	1.0	1.2	0.4	0.2	<0.1

As can be seen from these results, the formulas including compositions of the present invention, i.e., Formulas A and B had much lower foam heights than Formulas C and D which included the non-peracid form of the sulfonated material, i.e., sulfonated oleic acid. The reduced foam height of the compositions of the present invention is useful when using the compositions in applications where the production of foam is

detrimental to the application, for example, in a clean in place cleaning and/or sanitizing application.

Example 11 –Laundry Sanitizing Compositions

5 A study was run to determine the ability of a composition of the present invention to sanitize laundry. A composition containing the sulfonated peroleic acid product was tested against the commercially available cleaning compositions Ozonit®, commercially available from Ecolab Inc., and PAP-70®, available from Solvay. The compositions were tested against *Staphylococcus aureus* ATCC 6538 and *Pseudomonas*
 10 *aeruginosa* ATCC 15442 at 104°F for 6 minutes. The test method was as follows. Fabric samples that had been rinsed with boiling water containing 300 grams sodium carbonate and 1.5 grams of a non-ionic wetting agent (e.g., Triton X-100), followed by a cold water rinse until all visible traces of the wetting agent were removed, were obtained. The fabric samples were allowed to completely dry. The fabric samples were
 15 then autoclaved to sterilize them.

The test substances were then prepared, and the fabric samples were inoculated with the test substances. The inoculated swatches were then dried. The samples were then secured in a laundrometer and agitated in wash water. The wash water was removed from the chamber of the laundrometer, and the wash water and fabric samples
 20 are evaluated for the reduction of the tested microorganism population.

The results are shown in the table below.

Table 19.

Test/System	Composition including Sulfonated Peroleic Acid Product	PAP-70®	Ozonit®
Sanitizer Screen	>3.82	>3.82	NA
Disinfectant (Cloth Carrier	9 negative/9 total	9 negative/ 9 total	5 negative / 9 total

Screen)			
---------	--	--	--

As can be seen from these results, both the composition of the present invention tested showed a greater than 3 log reduction in both the wash water and fabric carriers against *P. aeruginosa* and on the fabric carriers against *S. aureus*.

The present invention also relates to novel compounds and the synthesis thereof. Accordingly, the following examples are presented to illustrate how some of those compounds may be prepared.


Synthesis of Selected Compounds of the Invention

Preparation of the sulfonated peroleic acid product.

417.8g of OA5-R (Intertrade Organic's, 40% active Sulfonated Oleic acid) was added to a 2-L beaker immersed in a large ice-bath, to which was subsequently added, 66.4g of Dequest 2010 (60% active Hydroxyethylenediphosphonic acid, Monsanto) and 535g of Hydrogen peroxide (46% active, Solvay-Interox) . The beaker was fitted with a magnetic stir bar and the solution was stirred aggressively while adding 940g of sulfuric acid (96% active, Mallinkrodt). The rate of the sulfuric acid addition was controlled to produce a 120° F exotherm in the reaction solution, and while this was occasionally exceeded by several degrees F, it wasn't allowed to exceed 125° F. Several minutes after completing the sulfuric acid addition, the ice bath was removed and the heterogenous solution was stirred for 72 hours allowing the temperature to equilibrate to ambient (70° F) conditions.

Several hours after discontinuing the stirring, the two phase reaction solution was added to a separatory funnel and the upper and lower phase were separated. 239.4g of upper phase were collected and the upper phase was further purified by centrifugation at 3000rpm for 10 minutes. The final upper phase yield was 206g (theoretical yield 174g) and titrated as 55% Peroxyacid based upon an assumed molecular weight of 380. In addition the upper phase contained 1.8% Hydrogen peroxide. A centrifuged lower phase sample titrated as 13% Peroxyacid (MW 380) and 8.8% hydrogen peroxide.

Synthesis of 11- Sulfoundecanoic Acid and 10, 11-Disulfoundecanoic Acid




11-Sulfoundecanoic acid: Deionized water (150 ml), isopropyl alcohol(200 ml) and 11-undecylenic acid (28.56 g, 0.155mol) were placed in a 1.0 liter flask equipped with stirrer, additional funnel, reflux condenser, thermometer and a gas inlet tube. To the additional funnel was added a premix which contained 15.2 g (0.08 mol) of sodium metabisulfite and 1.28 g of NaOH in 55 g of water. The whole device was purged with nitrogen gently. After heating to reflux (82 °C), a small portion of *t-butyl* perbenzoate (out of total amount of 0.5 g, 2.5 mmol) was added to the flask. Then the sodium metabisulfite/ NaOH premix was added continuously over a five hour period to the reaction solution through an addition funnel. The remaining *t-butyl* perbenzoate was also added in small portions during this time.

The solvent was then removed under reduced pressure using a rotavapour, and the residue washed with acetone, and dried, yielding 31.0 g of white solid. NMR analysis of the solid indicated no presence of the residual raw materials. The white solid obtained was dissolved in hot water (100 ml, 75 °C), and neutralized to pH 5.5 with NaOH. Then 2.0 g of 50% H₂O₂ was added to the solution. The solution was then allowed to cool down to room temperature, and the solid precipitated was filtered, washed with cold water, and dried, affording 21.0 g of white solid, characterized as pure 11-sulfoundecanoic acid. ¹³C NMR (D₂O): 180, 51, 34, 28-29 (multiple), 27.5, 24.5, 24 ppm. MS (ESI): 265.1 (M⁺ -H).

10, 11-Disulfoundecanoic acid: this compound was obtained as a byproduct from the 11-Sulfoundecanoic acid reaction as described above. The filtrate, after collecting 11-sulfoundecanoic acid through filtration, was concentrated to ~ 50 ml when precipitate start to form. The mixture was cooled down in the refrigerator, and the additional solid formed was filtered, washed with a small amount of ice water, and dried, yielding 5.0 g of white solid. ^{13}C NMR (D_2O): 184, 57, 51.5, 37.5, 28-29 (multiple), 27.5, 26, 24 ppm. MS (ESI): 345.0.


Synthesis of 11- Sulfoundecaneperoxoic acid (Compound D) and 10, 11- Disulfoundecaneperoxoic acid (Compound E)

11- Sulfoperoxyundecanoic Acid:



1.3 g of 11- sulfoundecanoic acid was dissolved in 2.5 g of 98% sulfuric acid. To this solution (the temperature of the solution did not exceed 60°C) 1.5 g of 50% H_2O_2 was added, and the resulting mixture was stirred at room temperature for 1.5 hr. At this point, a white solid precipitated from the solution. The mixture was reheated to 50°C with a water bath until the solution was clear. The solution was then stirred at room temperature for 0.5 hr, and cooled down in the freezer. Then 20 ml of ice water was added to the mixture, and the solid filtered, washed with ice water, and dried under vacuum, yielding 0.6 g of a white solid. ^{13}C NMR (D_2O): 176, 51.5, 30.5, 27.5-29 (multiple), 24.5, 24 ppm. MS (ESI): 281.5 ($\text{M}^+ - \text{H}$). Available oxygen (iodometric): 5.41 % (theoretical: 5.64%).


10, 11- Disulfoundecaneperoxoic acid (Compound E):



To 1.5 g of 10, 11- disulfoundecanoic acid was added 2.5 g of 96 % H_2SO_4 , and the mixture was stirred at room temperature. Then 1.0 g of 50 % H_2O_2 was added slowly (the temperature not exceeding 60°C) to the mixture, and after addition, the mixture was heated to 50°C with water bath, and the solution stirred for 2.0 hrs. The solution was then cooled down in the freezer, and 20 ml of ice water was added with stirring. The solid precipitated was filtered, washed with ice water, and dried under vacuum, affording 1.0 g of white solid. ^{13}C NMR (D_2O): 175.5, 57, 30.5, 27.5-29 (multiple), 24.5, 24 ppm. Available oxygen (iodometric): 4.10 % (theoretical: 4.41%).

10

Synthesis of 9/10-Sulfostearic Acid (Sulfonated Stearic Acid)



Deionized water (150 ml), isopropyl alcohol (200 ml) and oleic acid (43.78 g, 0.155mol) were placed in a 1.0 liter flask equipped with stirrer, additional funnel, reflux condenser, thermometer and a gas inlet tube. To the additional funnel was added a premix which contained 15.2 g (0.08 mol) of sodium metabisulfite ($\text{Na}_2\text{S}_2\text{O}_5$) and 1.28 g of NaOH in 55 g of water. The whole device was bubbled gently with nitrogen. After heating to reflux (82°C), a small portion of *t-butyl* perbenzoate (out of total amount of 0.5 g, 2.5 mmol) was added to the flask. Then the $\text{Na}_2\text{S}_2\text{O}_5$ / NaOH premix was added through the addition funnel continuously over the course of five hours. The remaining *t-butyl* perbenzoate was also added in portions during this time.

The solvent was then removed under reduced pressure using rotavapour. To the residue was added 100 ml of DI water, the pH of the solution was adjusted to 2.5 with H₂SO₄. The resulting mixture/solution was transferred to a separation funnel, and the top oily layer (non reacted oleic acid) was removed. The aqueous layer was extracted
5 with petroleum ether (2 x 50 ml), and after removal of the water, afforded 12.5 g of white waxy solid. ¹³C NMR (D₂O): 179, 60, 34.5, 32, 28.5-30 (multiple), 24.5, 22.5, 14 ppm. MS (ESI): 363.4 (M⁺ - H).


Preparation of 9/10-Sulfoperoxystearic Acid (in formulation)

10

To a 2.0 g mixture of 9 or 10-Sulfostearic acid was added 2.0 g of 50% H₂O₂. The mixture was stirred at room temperature until all the solid was dissolved. Then, 2.0 g of 75% H₃PO₄ was added, and the resulting solution was stirred at room temperature overnight. No attempt was made to isolate the pure 9 or 10-sulfoperoxystearic acid from
15 solution. ¹³C NMR (D₂O) of the solution showed a peracid peak (COOOH) at 174 ppm and the parent the carboxylic acid peak at 178 ppm. The iodometric titration (QATM-202) indicated 18.96% of sulfoperoxystearic acid.

CLAIMS

1. An aqueous, antimicrobial bleaching composition comprising:
 - a) a compound according to Formula I:



(Formula I)

wherein:

- R_1 is a sulfonate substituted C_1 alkyl group;
 - R_2 is a substituted or unsubstituted C_n alkyl group;
 - X is hydrogen, a cationic group, or an ester forming moiety;
 - n is 1 to 10;
 - or salts or esters thereof,
- b) a C_1 to C_4 carboxylic acid;
 - c) a C_5 to C_{11} carboxylic acid; and
 - d) an oxidizing agent wherein the pH of the composition is about 7 to about 14.
2. The composition of claim 1, wherein the C_1 to C_4 carboxylic acid is acetic acid.
 3. The composition of claim 1 or claim 2, wherein the C_5 to C_{11} carboxylic acid is octanoic acid.
 4. The composition of any one of claims 1 to 3, wherein the compound of Formula I includes at least one of 10-hydroxy-9-sulfooctadecaneperoxoic acid, 10,11-dihydroxy-9-sulfooctadecaneperoxoic acid, 9-hydroxy-10-sulfooctadecaneperoxoic acid, and 10-sulfo-8,9-dihydroxyoctadecaneperoxoic acid.
 5. The composition of any one of claims 1 to 4, wherein the oxidizing agent comprises hydrogen peroxide.
 6. The composition of any one of claims 1 to 5, further comprising a stabilizing agent.
 7. The composition of any one of claims 1 to 6, wherein the composition is substantially phosphorous free.

8. The composition of any one of claims 1 to 7, wherein the composition comprises:
- (a) about 1 wt% to about 5 wt% of a compound according to Formula I;
 - (b) about 1 wt% to about 10 wt% of the C₁ to C₄ carboxylic acid;
 - (c) about 1 wt% to about 10 wt% of the C₅ to C₁₁ carboxylic acid; and
 - (d) about 10 wt% to about 30 wt% of the oxidizing agent.
9. A method for treating laundry, the method comprising:
- (a) applying the composition according to any one of claims 1 to 8 to the laundry;
 - (b) draining the composition from the laundry.


Ecolab Inc.

Patent Attorneys for the Applicant/Nominated Person

SPRUSON & FERGUSON

2012201801 26 Feb 2014

Figure 1






Figure 2




Figure 3

Bleaching Performance of PSOA formulations




Figure 4




Figure 5




Figure 6