



(86) Date de dépôt PCT/PCT Filing Date: 2003/06/05
(87) Date publication PCT/PCT Publication Date: 2003/12/18
(85) Entrée phase nationale/National Entry: 2004/12/06
(86) N° demande PCT/PCT Application No.: US 2003/017592
(87) N° publication PCT/PCT Publication No.: 2003/104399
(30) Priorité/Priority: 2002/06/07 (60/386,793) US

(51) Cl.Int.⁷/Int.Cl.⁷ C07K 16/30, A61K 39/395, A61K 38/17,
C07K 14/705, C12Q 1/68, C12Q 1/02
(71) Demandeur/Applicant:
AVALON PHARMACEUTICALS, INC, US
(72) Inventeurs/Inventors:
RICK, JENNIFER A., US;
EBNER, REINHARD, US
(74) Agent: SMART & BIGGAR

(54) Titre : GENE LIE AU CANCER SERVANT DE CIBLE EN CHIMIOOTHERAPIE
(54) Title: CANCER-LINKED GENE AS TARGET FOR CHEMOTHERAPY

(57) **Abrégé/Abstract:**

Cancer-linked gene sequences, and derived amino acid sequences, are disclosed along with processes for assaying potential antitumor agents based on their modulation of the expression of these cancer-linked genes. Also disclosed are antibodies that react with the disclosed polypeptides and methods of using the antibodies to treat cancerous conditions, such as by using the antibody to target cancerous cells in vivo for purposes of delivering therapeutic agents thereto. Also described are methods of diagnosing using the gene sequences.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
18 December 2003 (18.12.2003)

PCT

(10) International Publication Number
WO 03/104399 A2

- (51) International Patent Classification⁷: C12N
- (21) International Application Number: PCT/US03/17592
- (22) International Filing Date: 5 June 2003 (05.06.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/386,793 7 June 2002 (07.06.2002) US
- (71) Applicant (for all designated States except US): **AVALON PHARMACEUTICALS, INC** [US/US]; 20358 Seneca Meadows Parkway, Germantown, MD 20876 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **RICK, Jennifer, A.** [US/US]; 13356 Rushing Water Way, Germantown, MD 20874 (US). **EBNER, Reinhard** [US/US]; 9906 Shelbourne Terrace, #316, Gaithersburg, MD 20878 (US).
- (74) Agents: **GRANT, Alan, J.** et al.; Carella, Byrne, Bain, Gilfillan, Cecchi, Stewart & Olstein, 6 Becker Farm Road, Roseland, NJ 07068 (US).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:**
— without international search report and to be republished upon receipt of that report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: CANCER-LINKED GENE AS TARGET FOR CHEMOTHERAPY

(57) Abstract: Cancer-linked gene sequences, and derived amino acid sequences, are disclosed along with processes for assaying potential antitumor agents based on their modulation of the expression of these cancer-linked genes. Also disclosed are antibodies that react with the disclosed polypeptides and methods of using the antibodies to treat cancerous conditions, such as by using the antibody to target cancerous cells *in vivo* for purposes of delivering therapeutic agents thereto. Also described are methods of diagnosing using the gene sequences.



WO 03/104399 A2

CANCER-LINKED GENE AS TARGET FOR CHEMOTHERAPY

5

This application claims priority of U.S. Provisional Patent Application 60/386,793, filed 7 June 2002, the disclosure of which is hereby incorporated
10 by reference in its entirety.

15

FIELD OF THE INVENTION

The present invention relates to methods of screening cancer-linked
20 genes and expression products for involvement in the cancer initiation and facilitation process as a means of cancer diagnosis as well as the use of such genes for screening potential anti-cancer agents, including small organic compounds and other molecules, and development of therapeutic agents.

25

BACKGROUND OF THE INVENTION

Cancer-linked genes are valuable in that they indicate genetic
30 differences between cancer cells and normal cells, such as where a gene is expressed in a cancer cell but not in a non-cancer cell, or where said gene is over-expressed or expressed at a higher level in a cancer as opposed to normal or non-cancer cell. In addition, the expression of such a gene in a normal cell but not in a cancer cell, especially of the same type of tissue, can
35 indicate important functions in the cancerous process. For example, screening assays for novel drugs are based on the response of model cell based systems *in vitro* to treatment with specific compounds. Such genes are also

useful in the diagnosis of cancer and the identification of a cell as cancerous. Gene activity is readily measured by measuring the rate of production of gene products, such as RNAs and polypeptides encoded by such genes. Where genes encode cell surface proteins, appearance of, or alterations in, such proteins, as cell surface markers, are an indication of neoplastic activity. Some such screens rely on specific genes, such as oncogenes (or gene mutations). In accordance with the present invention, a cancer-linked gene has been identified and its putative amino acid sequence worked out. Such gene is useful in the diagnosing of cancer, the screening of anticancer agents and the treatment of cancer using such agents, especially in that these genes encode polypeptides that can act as markers, such as cell surface markers, thereby providing ready targets for anti-tumor agents such as antibodies, preferably antibodies complexed to cytotoxic agents, including apoptotic agents.

15

BRIEF SUMMARY OF THE INVENTION

20 In accordance with the present invention, there is provided herein a cancer specific gene, linked especially to breast or endometrium cancer, or otherwise involved in the cancer initiating and facilitating process and the derived amino acid sequence thereof, including a number of different transcripts derived from said gene.

25

In one aspect, the present invention relates to a process for identifying an agent that modulates the activity of a cancer-related gene comprising:

(a) contacting a compound with a cell containing a gene that corresponds to a polynucleotide having a sequence selected from the group consisting of SEQ ID NO: 1, 2, 3 and 4 and under conditions promoting the expression of said gene; and

30

(b) detecting a difference in expression of said gene relative to when said compound is not present

thereby identifying an agent that modulates the activity of a cancer-related gene.

5

In various embodiments of such a process, the cell is a cancer cell and the difference in expression is a decrease in expression. Such polynucleotides may also include those that have sequences identical to SEQ ID NO: 1, 2, 3 and 4.

10

In another aspect, the present invention relates to a process for identifying an anti-neoplastic agent comprising contacting a cell exhibiting neoplastic activity with a compound first identified as a cancer related gene modulator using an assay process disclosed herein and detecting a decrease

15 in said neoplastic activity after said contacting compared to when said contacting does not occur. Such neoplastic activity may include accelerated cellular replication and/or metastasis, and the decrease in neoplastic activity preferably results from the death of the cell, or senescence, terminal differentiation or growth inhibition.

20

The present invention also relates to a process for identifying an anti-neoplastic agent comprising administering to an animal exhibiting a cancer condition an effective amount of an agent first identified according to a process of one of one of the assays disclosed according to the invention and

25 detecting a decrease in said cancerous condition.

The present invention further relates to a process for determining the cancerous status of a cell, comprising determining an increase in the level of expression in said cell of at least one gene that corresponds to a

30 polynucleotide having a sequence selected from the group consisting of SEQ ID NO: 1, 2, 3 and 4 wherein an elevated expression relative to a known non-

cancerous cell indicates a cancerous state or potentially cancerous state. Such elevated expression may be due to an increased copy number.

5 The present invention additionally relates to an isolated polypeptide, encoded by one of the polynucleotide transcripts disclosed herein, comprising an amino acid sequence homologous to an amino acid sequence selected from the group consisting of SEQ ID NO: 5 and 6 wherein any difference between said amino acid sequence and the sequence of SEQ ID NO: 5 and 6 is due solely to conservative amino acid substitutions and wherein said
10 isolated polypeptide comprises at least one immunogenic fragment. In a preferred embodiment, the present invention encompasses an isolated polypeptide comprising an amino acid sequence homologous to an amino acid sequence selected from the group consisting of SEQ ID NO: 5 and 6.

15 The present invention also relates to an antibody that reacts with a polypeptide as disclosed herein, preferably a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 5 and 6. Such an antibody may be polyclonal, monoclonal, recombinant or synthetic in origin.

20 In one such embodiment, said antibody is associated, either covalently or non-covalently, with a cytotoxic agent, for example, an apoptotic agent. Thus, the present invention relates to an immunoconjugate comprising an antibody of the invention and a cytotoxic
25 agent.

The present invention also relates to a process for treating cancer comprising contacting a cancerous cell with an agent having activity against an expression product encoded by a gene sequence selected from the group
30 consisting of SEQ ID NO: 1, 2, 3 and 4. In one such embodiment, the cancerous cell is contacted *in vivo*. In another such embodiment, said agent has affinity for said expression product. In a preferred embodiment, such

agent is an antibody disclosed herein, such as an antibody that is specific or selective for, or otherwise reacts with, a polypeptide of the invention. In a preferred embodiment, the expression product is a polypeptide incorporating an amino acid sequence selected from SEQ ID NO: 5 and 6.

5

The present invention further encompasses an immunogenic composition comprising a polypeptide disclosed herein, as well as compositions formed using antibodies specific for these polypeptides.

10 The present invention is also directed to uses of such compositions. Such uses include a method for treating cancer in an animal afflicted therewith comprising administering to said animal an amount of an immunogenic composition of one or more of the polypeptides disclosed herein where such amount is an amount sufficient to elicit the production of cytotoxic
15 T lymphocytes specific for a polypeptide of the invention, preferably a polypeptide incorporating a sequence of SEQ ID NO: 5 and 6. In a preferred embodiment, the animal to be so treated is a human patient.

20

DEFINITIONS

As used herein, the terms "portion," "segment," and "fragment," when used in relation to polypeptides, refer to a continuous sequence of residues,
25 such as amino acid residues, which sequence forms a subset of a larger sequence. For example, if a polypeptide were subjected to treatment with any of the common endopeptidases, such as trypsin or chymotrypsin, the oligopeptides resulting from such treatment would represent portions, segments or fragments of the starting polypeptide. When used in relation to a polynucleotides, such
30 terms refer to the products produced by treatment of said polynucleotides with any of the common endonucleases.

As used herein, the term "isolated" means that the material is removed from its original environment (e.g., the natural environment if it is naturally occurring). It could also be produced recombinantly and subsequently purified. For example, a naturally-occurring polynucleotide or polypeptide present in a living animal is not isolated, but the same polynucleotide or polypeptide, separated from some or all of the coexisting materials in the natural system, is isolated. Such polynucleotides, for example, those prepared recombinantly, could be part of a vector and/or such polynucleotides or polypeptides could be part of a composition, and still be isolated in that such vector or composition is not part of its natural environment. In one embodiment of the present invention, such isolated, or purified, polypeptide is useful in generating antibodies for practicing the invention, or where said antibody is attached to a cytotoxic or cytolytic agent, such as an apoptotic agent.

The term "percent identity" or "percent identical," when referring to a sequence, means that a sequence is compared to a claimed or described sequence after alignment of the sequence to be compared (the "Compared Sequence") with the described or claimed sequence (the "Reference Sequence"). The Percent Identity is then determined according to the following formula:

$$\text{Percent Identity} = 100 [1 - (C/R)]$$

wherein C is the number of differences between the Reference Sequence and the Compared Sequence over the length of alignment between the Reference Sequence and the Compared Sequence wherein (i) each base or amino acid in the Reference Sequence that does not have a corresponding aligned base or amino acid in the Compared Sequence and (ii) each gap in the Reference Sequence and (iii) each aligned base or amino acid in the Reference Sequence that is different from an aligned base or amino acid in the Compared Sequence, constitutes a difference; and R is the number of bases or amino acids in the Reference Sequence over the length of the alignment with the Compared

Sequence with any gap created in the Reference Sequence also being counted as a base or amino acid.

5 If an alignment exists between the Compared Sequence and the Reference Sequence for which the percent identity as calculated above is about equal to or greater than a specified minimum Percent Identity then the Compared Sequence has the specified minimum percent identity to the Reference Sequence even though alignments may exist in which the hereinabove calculated Percent Identity is less than the specified Percent
10 Identity.

As known in the art "similarity" between two polypeptides is determined by comparing the amino acid sequence and its conserved amino acid substitutes of one polypeptide to the sequence of a second polypeptide.
15

In accordance with the present invention, the term "DNA segment" or "DNA sequence" refers to a DNA polymer, in the form of a separate fragment or as a component of a larger DNA construct, which has been derived from DNA isolated at least once in substantially pure form, i.e., free of
20 contaminating endogenous materials and in a quantity or concentration enabling identification, manipulation, and recovery of the segment and its component nucleotide sequences by standard biochemical methods, for example, using a cloning vector. Such segments are provided in the form of an open reading frame uninterrupted by internal nontranslated sequences, or
25 introns, which are typically present in eukaryotic genes. Sequences of non-translated DNA may be present downstream from the open reading frame, where the same do not interfere with manipulation or expression of the coding regions.

30 The term "coding region" refers to that portion of a gene which either naturally or normally codes for the expression product of that gene in its natural genomic environment, i.e., the region coding *in vivo* for the native expression product of the gene. The coding region can be from a normal,

mutated or altered gene, or can even be from a DNA sequence, or gene, wholly synthesized in the laboratory using methods well known to those of skill in the art of DNA synthesis.

5 In accordance with the present invention, the term "nucleotide sequence" refers to a heteropolymer of deoxyribonucleotides. Generally, DNA segments encoding the proteins provided by this invention are assembled from cDNA fragments and short oligonucleotide linkers, or from a series of oligonucleotides, to provide a synthetic gene which is capable of being
10 expressed in a recombinant transcriptional unit comprising regulatory elements derived from a microbial, eukaryotic or viral operon.

 The term "expression product" means that polypeptide or protein that is the natural translation product of the gene and any nucleic acid sequence
15 coding equivalents resulting from genetic code degeneracy and thus coding for the same amino acid(s).

 The term "active fragment," when referring to a coding sequence, means a portion comprising less than the complete coding region whose expression
20 product retains essentially the same biological function or activity as the expression product of the complete coding region.

 The term "primer" means a short nucleic acid sequence that is paired with one strand of DNA and provides a free 3'-OH end at which a DNA
polymerase starts synthesis of a deoxyribonucleotide chain.

25

 The term "promoter" means a region of DNA involved in binding of RNA polymerase to initiate transcription. The term "enhancer" refers to a region of DNA that, when present and active, has the effect of increasing expression of a different DNA sequence that is being expressed, thereby increasing the
30 amount of expression product formed from said different DNA sequence.

The term "open reading frame (ORF)" means a series of triplets coding for amino acids without any termination codons and is a sequence (potentially) translatable into protein.

5 As used herein, reference to a "DNA sequence" includes both single stranded and double stranded DNA. Thus, the specific sequence, unless the context indicates otherwise, refers to the single strand DNA of such sequence, the duplex of such sequence with its complement (double stranded DNA) and the complement of such sequence.

10

As used herein, "corresponding genes" refers to genes that encode an RNA that is at least 90% identical, preferably at least 95% identical, most preferably at least 98% identical, and especially identical, to an RNA encoded by one of the nucleotide sequences disclosed herein (i.e., SEQ ID NO: 1, 2, 3 and 4). Such genes will also encode the same polypeptide sequence as any of the sequences disclosed herein, preferably SEQ ID NO: 1, 2, 3 and 4, but may include differences in such amino acid sequences where such differences are limited to conservative amino acid substitutions, such as where the same overall three dimensional structure, and thus the same antigenic character, is maintained. Thus, amino acid sequences may be within the scope of the present invention where they react with the same antibodies that react with polypeptides comprising the sequences of SEQ ID NO: 5 and 6. A "corresponding gene" includes splice variants thereof.

25 The genes identified by the present disclosure are considered "cancer-related" genes, as this term is used herein, and include genes expressed at higher levels (due, for example, to elevated rates of expression, elevated extent of expression or increased copy number) in cancer cells relative to expression of these genes in normal (i.e., non-cancerous) cells where said cancerous state or status of test cells or tissues has been determined by
30 methods known in the art, such as by reverse transcriptase polymerase chain reaction (RT-PCR) as described in the Examples herein. In specific

embodiments, this relates to the genes whose sequences correspond to the sequences of SEQ ID NO: 1, 2, 3 and 4.

As used herein, the term "conservative amino acid substitutions" are defined herein as exchanges within one of the following five groups:

I. Small aliphatic, nonpolar or slightly polar residues:

Ala, Ser, Thr, Pro, Gly;

II. Polar, negatively charged residues and their amides:

Asp, Asn, Glu, Gln;

10 III. Polar, positively charged residues:

His, Arg, Lys;

IV. Large, aliphatic, nonpolar residues:

Met Leu, Ile, Val, Cys

V. Large, aromatic residues:

15 Phe, Tyr, Trp

DETAILED SUMMARY OF THE INVENTION

20

The present invention relates to processes for utilizing a nucleotide sequence for a cancer-linked gene, polypeptides encoded by such sequences and antibodies reactive with such polypeptides in methods of treating and diagnosing cancer, preferably breast or endometrium cancer, and in carrying out screening assays for agents effective in reducing the activity of cancer-linked genes and thereby treating a cancerous condition.

25

The polypeptides disclosed herein incorporate various polynucleotide transcripts (SEQ ID NO: 1, 2, 3 and 4) and the derived amino acid sequence (SEQ ID NO: 5 and 6) from said transcripts are available as targets for
30 chemotherapeutic agents, especially anti-cancer agents, including antibodies specific for said polypeptides.

The cancer-related polynucleotide sequences disclosed herein correspond to gene sequences whose expression is indicative of the cancerous status of a given cell. Such sequences are substantially identical to
 5 SEQ ID NO: 1, 2, 3 and 4, which represent different transcripts identified from the GenBank EST database and which exhibit cancer-specific expression. The polynucleotides of the invention are those that correspond to a sequence of SEQ ID NO: 1, 2, 3 and 4. Such sequences have been searched within the GenBank database, especially the EST database, with the following results:

10

Type:	cell-surface tumor antigen therapeutic antibody target
Tissue:	breast and endometrium
AffyFragment-ID(s):	125026, 142673 hypothetical protein FLJ22418
Accession(s):	AA075632, AI799522
Unigene cluster-ID(s):	Hs.36563
Chromosomal location:	1

25

The nucleotides and polypeptides, as gene products, used in the processes of the present invention may comprise a recombinant polynucleotide or polypeptide, a natural polynucleotide or polypeptide, or a synthetic polynucleotide or polypeptide, or a recombinant polynucleotide or polypeptide.

30

Fragments of such polynucleotides and polypeptides as are disclosed herein may also be useful in practicing the processes of the present invention. For example, a fragment, derivative or analog of the polypeptide (SEQ ID NO: 5 and 6) may be (i) one in which one or more of the amino acid residues are
 35 substituted with a conserved or non-conserved amino acid residue (preferably a conserved amino acid residue) and such substituted amino acid residue may or

may not be one encoded by the genetic code, or (ii) one in which one or more of the amino acid residues includes a substituent group, or (iii) one in which the mature polypeptide is fused with another compound, such as a compound to increase the half-life of the polypeptide (for example, polyethylene glycol), or (iv)
5 one in which the additional amino acids are fused to the mature polypeptide, such as a leader or secretory sequence or a sequence which is employed for purification of the mature polypeptide (such as a histidine hexapeptide) or a proprotein sequence. Such fragments, derivatives and analogs are deemed to be within the scope of those skilled in the art from the teachings herein.

10

In another aspect, the present invention relates to an isolated polypeptide, including a purified polypeptide, comprising an amino acid sequence at least 90% identical to the amino acid sequence of SEQ ID NO: 5 and/or 6. In preferred embodiments, said isolated polypeptide comprises an
15 amino acid sequence having sequence identity of at least 95%, preferably at least about 98%, and especially is identical to, the sequence of SEQ ID NO: 5 and/or 6. The present invention also includes isolated active fragments of such polypeptides where said fragments retain the biological activity of the polypeptide or where such active fragments are useful as specific targets for
20 cancer treatment, prevention or diagnosis. Thus, the present invention relates to any polypeptides, or fragments thereof, with sufficient sequence homology to the sequences disclosed herein as to be useful in the production of antibodies that react with (i.e., are selective or specific for) the polypeptides of SEQ ID NO: 5 and 6 so as to be useful in targeting cells that exhibit such polypeptides, or
25 fragments, on their surfaces, thereby providing targets for such antibodies and therapeutic agents associated with such antibodies.

The polynucleotides and polypeptides useful in practicing the processes of the present invention may likewise be obtained in an isolated or purified form.
30 In addition, the polypeptide disclosed herein as being useful in practicing the processes of the invention are believed to be surface proteins present on cells, such as cancerous cells. Precisely how such cancer-linked proteins are used in

the processes of the invention may thus differ depending on the therapeutic approach used. For example, cell-surface proteins, such as receptors, are desirable targets for cytotoxic antibodies that can be generated against the polypeptides disclosed herein.

5

The sequence information disclosed herein, as derived from the GenBank submissions, can readily be utilized by those skilled in the art to prepare the corresponding full-length polypeptide by peptide synthesis. The same is true for either the polynucleotides or polypeptides disclosed herein for use in the methods of the invention.

10

The present invention relates to an isolated polypeptide, encoded by one of the polynucleotide transcripts disclosed herein, comprising an amino acid sequence homologous to an amino acid sequence selected from the group consisting of SEQ ID NO: 5 and 6, wherein any difference between amino acid sequence in the isolated polypeptide and the sequence of SEQ ID NO: 5 and 6 is due solely to conservative amino acid substitutions and wherein said isolated polypeptide comprises at least one immunogenic fragment. In a preferred embodiment, the present invention encompasses an isolated polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 5 and 6.

15

20

Methods of producing recombinant cells and vectors useful in preparing the polynucleotides and polypeptides disclosed herein are well known to those skilled in the molecular biology art. See, for example, Sambrook, et al., *Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbor, N.Y., (1989), Wu et al., *Methods in Gene Biotechnology* (CRC Press, New York, NY, 1997), and *Recombinant Gene Expression Protocols*, in *Methods in Molecular Biology*, Vol. 62, (Tuan, ed., Humana Press, Totowa, NJ, 1997), the disclosures of which are hereby incorporated by reference.

25

30

In one aspect, the present invention relates to a process for identifying an agent that modulates the activity of a cancer-related gene comprising:

(a) contacting a compound with a cell containing a gene that corresponds to a polynucleotide having a sequence selected from the group consisting of SEQ ID NO: 1, 2, 3 and 4 and under conditions promoting the expression of said gene; and

5 (b) detecting a difference in expression of said gene relative to when said compound is not present

thereby identifying an agent that modulates the activity of a cancer-related gene.

10 In specific embodiments of such process the cell is a cancer cell and the difference in expression is a decrease in expression. Such polynucleotides may also include those that have sequences identical to SEQ ID NO: 1, 2, 3 and 4.

15 In another aspect, the present invention relates to a process for identifying an anti-neoplastic agent comprising contacting a cell exhibiting neoplastic activity with a compound first identified as a cancer related gene modulator using an assay process disclosed herein and detecting a decrease
20 in said neoplastic activity after said contacting compared to when said contacting does not occur. Such neoplastic activity may include accelerated cellular replication and/or metastasis, and the decrease in neoplastic activity preferably results from the death of the cell.

The present invention also relates to a process for identifying an anti-
25 neoplastic agent comprising administering to an animal exhibiting a cancer condition an effective amount of an agent first identified according to a process of one of one of the assays disclosed according to the invention and detecting a decrease in said cancerous condition.

30 In specific embodiments of the present invention, the genes useful for the invention comprise genes that correspond to polynucleotides having a sequence selected from SEQ ID NO: 1, 2, 3 and 4, or may comprise the

sequence of any of the polynucleotides disclosed herein (where the latter are cDNA sequences).

In accordance with the present invention, such assays rely on methods
5 of determining the activity of the gene in question. Such assays are
advantageously based on model cellular systems using cancer cell lines,
primary cancer cells, or cancerous tissue samples that are maintained in
growth medium and treated with compounds at a single concentration or at a
range of concentrations. At specific times after treatment, cellular RNAs are
10 conveniently isolated from the treated cells or tissues, which RNAs are
indicative of expression of selected genes. The cellular RNA is then divided
and subjected to differential analysis that detects the presence and/or quantity
of specific RNA transcripts, which transcripts may then be amplified for
detection purposes using standard methodologies, such as, for example,
15 reverse transcriptase polymerase chain reaction (RT-PCR), etc. The presence
or absence, or concentration levels, of specific RNA transcripts are
determined from these measurements. The polynucleotide sequences
disclosed herein are readily used as probes for the detection of such RNA
transcripts and thus the measurement of gene activity and expression.

20

The polynucleotides of the invention can include fully operational genes
with attendant control or regulatory sequences or merely a polynucleotide
sequence encoding the corresponding polypeptide or an active fragment or
analog thereof.

25

Because expression of the polynucleotide sequences disclosed herein
are specific to the cancerous state, useful gene modulation is downward
modulation, so that, as a result of exposure to an antineoplastic agent
identified by the screening assays herein, the corresponding gene of the
30 cancerous cell is expressed at a lower level (or not expressed at all) when
exposed to the agent as compared to the expression when not exposed to the
agent. For example, the gene sequences disclosed herein (SEQ ID NO: 1, 2,

3 and 4) correspond to a gene expressed at a higher level in cells of breast or endometrium cancer than in normal breast or endometrium cells. Thus, where said chemical agent causes this gene of the tested cell to be expressed at a lower level than the same genes of the reference, this is indicative of downward modulation and indicates that the chemical agent to be tested has anti-neoplastic activity.

In carrying out the assays disclosed herein, relative antineoplastic activity may be ascertained by the extent to which a given chemical agent modulates the expression of genes present in a cancerous cell. Thus, a first chemical agent that modulates the expression of a gene associated with the cancerous state (i.e., a gene corresponding to one or more of the polynucleotide transcripts disclosed herein) to a larger degree than a second chemical agent tested by the assays of the invention is thereby deemed to have higher, or more desirable, or more advantageous, anti-neoplastic activity than said second chemical agent.

The gene expression to be measured is commonly assayed using RNA expression as an indicator. Thus, the greater the level of RNA (for example, messenger RNA or mRNA) detected the higher the level of expression of the corresponding gene. Thus, gene expression, either absolute or relative, is determined by the relative expression of the RNAs encoded by such genes.

RNA may be isolated from samples in a variety of ways, including lysis and denaturation with a phenolic solution containing a chaotropic agent (e.g., trizol) followed by isopropanol precipitation, ethanol wash, and resuspension in aqueous solution; or lysis and denaturation followed by isolation on solid support, such as a Qiagen resin and reconstitution in aqueous solution; or lysis and denaturation in non-phenolic, aqueous solutions followed by enzymatic conversion of RNA to DNA template copies.

30

Normally, prior to applying the processes of the invention, steady state RNA expression levels for the genes, and sets of genes, disclosed herein will

have been obtained. It is the steady state level of such expression that is affected by potential anti-neoplastic agents as determined herein. Such steady state levels of expression are easily determined by any methods that are sensitive, specific and accurate. Such methods include, but are in no way limited to, real time quantitative polymerase chain reaction (PCR), for example, using a Perkin-Elmer 7700 sequence detection system with gene specific primer probe combinations as designed using any of several commercially available software packages, such as Primer Express software., solid support based hybridization array technology using appropriate internal controls for quantitation, including filter, bead, or microchip based arrays, solid support based hybridization arrays using, for example, chemiluminescent, fluorescent, or electrochemical reaction based detection systems.

The gene expression indicative of a cancerous state need not be characteristic of every cell of a given tissue. Thus, the methods disclosed herein are useful for detecting the presence of a cancerous condition within a tissue where less than all cells exhibit the complete pattern. Thus, for example, a selected gene corresponding to the sequence of SEQ ID NO: 1, may be found, using appropriate probes, either DNA or RNA, to be present in as little as 60% of cells derived from a sample of tumorous, or malignant, tissue. In a highly preferred embodiment, such gene pattern is found to be present in at least 100% of cells drawn from a cancerous tissue and absent from at least 100% of a corresponding normal, non-cancerous, tissue sample.

Expression of a gene may be related to copy number, and changes in expression may be measured by determining copy number. Such change in gene copy number may be determined by determining a change in expression of messenger RNA encoded by a particular gene sequence, especially that of SEQ ID NO: 1, 2, 3 and 4. Also in accordance with the present invention, said gene may be a cancer initiating or facilitating gene. In carrying out the methods of the present invention, a cancer facilitating gene is a gene that, while not directly initiating tumor formation or growth, acts, such as through

the actions of its expression product, to direct, enhance, or otherwise facilitate the progress of the cancerous condition, including where such gene acts against genes, or gene expression products, that would otherwise have the effect of decreasing tumor formation and/or growth.

5

Although the expression of a gene corresponding to a sequence of SEQ ID NO: 1, 2, 3 and 4 may be indicative of a cancerous status for a given cell, the mere presence of such a gene may not alone be sufficient to achieve a malignant condition and thus the level of expression of such gene may also
10 be a significant factor in determining the attainment of a cancerous state. Thus, it becomes essential to also determine the level of expression of a gene as disclosed herein, including substantially similar sequences, as a separate means of diagnosing the presence of a cancerous status for a given cell, groups of cells, or tissues, either in culture or *in situ*.

15

The level of expression of the polypeptides disclosed herein is also a measure of gene expression, such as polypeptides having sequence identical, or similar to, any polypeptide encoded by a sequence of SEQ ID NO: 1, 2, 3 and 4, especially a polypeptide whose amino acid sequence is the sequence
20 of SEQ ID NO: 5 and 6.

In accordance with the foregoing, the present invention specifically contemplates a method for determining the cancerous status of a cell to be tested, comprising determining the level of expression in said cell of a gene
25 that includes one of the nucleotide sequences selected from the sequences of SEQ ID NO: 1, 2, 3 and 4, including sequences substantially identical to said sequences, or characteristic fragments thereof, or the complements of any of the foregoing and then comparing said expression to that of a cell known to be non-cancerous whereby the difference in said expression indicates that
30 said cell to be tested is cancerous.

In accordance with the invention, although gene expression for a gene that includes as a portion thereof one of the sequences of SEQ ID NO: 1, 2, 3 and 4, is preferably determined by use of a probe that is a fragment of such nucleotide sequence, it is to be understood that the probe may be formed
5 from a different portion of the gene. Expression of the gene may be determined by use of a nucleotide probe that hybridizes to messenger RNA (mRNA) transcribed from a portion of the gene other than the specific nucleotide sequence disclosed herein.

10 It should be noted that there are a variety of different contexts in which genes have been evaluated as being involved in the cancerous process. Thus, some genes may be oncogenes and encode proteins that are directly involved in the cancerous process and thereby promote the occurrence of cancer in an animal. In addition, other genes may serve to suppress the
15 cancerous state in a given cell or cell type and thereby work against a cancerous condition forming in an animal. Other genes may simply be involved either directly or indirectly in the cancerous process or condition and may serve in an ancillary capacity with respect to the cancerous state. All such types of genes are deemed with those to be determined in accordance
20 with the invention as disclosed herein. Thus, the gene determined by said process of the invention may be an oncogene, or the gene determined by said process may be a cancer facilitating gene, the latter including a gene that directly or indirectly affects the cancerous process, either in the promotion of a cancerous condition or in facilitating the progress of cancerous growth or
25 otherwise modulating the growth of cancer cells, either *in vivo* or *ex vivo*. In addition, the gene determined by said process may be a cancer suppressor gene, which gene works either directly or indirectly to suppress the initiation or progress of a cancerous condition. Such genes may work indirectly where their expression alters the activity of some other gene or gene expression
30 product that is itself directly involved in initiating or facilitating the progress of a cancerous condition. For example, a gene that encodes a polypeptide, either wild or mutant in type, which polypeptide acts to suppress of tumor

suppressor gene, or its expression product, will thereby act indirectly to promote tumor growth.

As noted previously, polynucleotides encoding the same proteins as
5 any of SEQ ID NO: 1, 2, 3 and 4, regardless of the percent identity of such
sequences, are also specifically contemplated by any of the methods of the
present invention that rely on any or all of said sequences, regardless of how
they are otherwise described or limited. Thus, any such sequences are
available for use in carrying out any of the methods disclosed according to the
10 invention. Such sequences also include any open reading frames, as defined
herein, present within the sequence of SEQ ID NO: 1, 2, 3 and 4.

Because a gene disclosed according to the invention "corresponds to"
a polynucleotide having a sequence of SEQ ID NO: 1, 2, 3 and 4, said gene
15 encodes an RNA (processed or unprocessed, including naturally occurring
splice variants and alleles) that is at least 90% identical, preferably at least
95% identical, most preferably at least 98% identical to, and especially
identical to, an RNA that would be encoded by, or be complementary to, such
as by hybridization with, a polynucleotide having the indicated sequence. In
20 addition, genes including sequences at least 90% identical to a sequence
selected from SEQ ID NO: 1, 2, 3 and 4, preferably at least about 95%
identical to such a sequence, more preferably at least about 98% identical to
such sequence and most preferably comprising such sequence are
specifically contemplated by all of the processes of the present invention.
25 Sequences encoding the same proteins as any of these sequences,
regardless of the percent identity of such sequences, are also specifically
contemplated by any of the methods of the present invention that rely on any
or all of said sequences, regardless of how they are otherwise described or
limited. The polynucleotide sequences of the invention also include any open
30 reading frames, as defined herein, present within any of the sequences of
SEQ ID NO: 1, 2, 3 and 4.

The sequences disclosed herein may be genomic in nature and thus represent the sequence of an actual gene, such as a human gene, or may be a cDNA sequence derived from a messenger RNA (mRNA) and thus represent contiguous exonic sequences derived from a corresponding genomic sequence, or they may be wholly synthetic in origin for purposes of practicing the processes of the invention. Because of the processing that may take place in transforming the initial RNA transcript into the final mRNA, the sequences disclosed herein may represent less than the full genomic sequence. They may also represent sequences derived from ribosomal and transfer RNAs. Consequently, the gene as present in the cell (and representing the genomic sequence) and the polynucleotide transcripts disclosed herein, including cDNA sequences, may be identical or may be such that the cDNAs contain less than the full genomic sequence. Such genes and cDNA sequences are still considered "corresponding sequences" (as defined elsewhere herein) because they both encode the same or related RNA sequences (i.e., related in the sense of being splice variants or RNAs at different stages of processing). Thus, by way of non-limiting example only, a gene that encodes an RNA transcript, which is then processed into a shorter mRNA, is deemed to encode both such RNAs and therefore encodes an RNA complementary to (using the usual Watson-Crick complementarity rules), or that would otherwise be encoded by, a cDNA (for example, a sequence as disclosed herein). Thus, the sequences disclosed herein correspond to genes contained in the cancerous cells (here, breast or endometrium cancer) and are used to determine gene activity or expression because they represent the same sequence or are complementary to RNAs encoded by the gene. Such a gene also includes different alleles and splice variants that may occur in the cells used in the methods of the invention, such as where recombinant cells are used to assay for anti-neoplastic agents and such cells have been engineered to express a polynucleotide as disclosed herein, including cells that have been engineered to express such polynucleotides at a higher level than is found in non-engineered cancerous cells or where such recombinant cells express such polynucleotides only after having been engineered to do

so. Such engineering includes genetic engineering, such as where one or more of the polynucleotides disclosed herein has been inserted into the genome of such cell or is present in a vector.

5 Such cells, especially mammalian cells, may also be engineered to express on their surfaces one or more of the polypeptides of the invention for testing with antibodies or other agents capable of masking such polypeptides and thereby removing the cancerous nature of the cell. Such engineering includes both genetic engineering, where the genetic complement of the cells
10 is engineered to express the polypeptide, as well as non-genetic engineering, whereby the cell has been physically manipulated to incorporate a polypeptide of the invention in its plasma membrane, such as by direct insertion using chemical and/or other agents to achieve this result.

15 In accordance with the foregoing, the present invention includes anti-cancer agents that are themselves either polypeptides, or small chemical entities, that affect the cancerous process, including initiation, suppression or facilitation of tumor growth, either *in vivo* or *ex vivo*. Said cancer modulating agent will have the effect of decreasing gene expression.

20 The present invention thus also relates to a method for treating cancer comprising contacting a cancerous cell with an agent having activity against an expression product encoded by a gene or polynucleotide sequence as disclosed herein, such as one having, or corresponding to, the nucleotide
25 sequence of SEQ ID NO: 1, 2, 3 and 4. The present invention also relates to a process for treating cancer comprising contacting a cancerous cell with an agent having activity against an expression product encoded by a gene or polynucleotide sequence corresponding to a sequence selected from the group consisting of SEQ ID NO: 1, 2, 3 and 4. In one such embodiment, the
30 cancerous cell is contacted *in vivo*. In another such embodiment, said agent has affinity for said expression product. In a preferred embodiment, such agent is an antibody disclosed herein, such as an antibody that is specific or

selective for, or otherwise reacts with, a polypeptide of the invention. In a preferred embodiment, the expression product is a polypeptide incorporating an amino acid sequence selected from SEQ ID NO: 5 and 6.

5 The present invention is also directed to such uses of the compositions of polypeptides and antibodies disclosed herein. Such uses include a process for treating cancer in an animal afflicted therewith comprising administering to said animal an amount of an immunogenic composition of one or more of the polypeptides disclosed herein where such amount is an amount sufficient to
10 elicit the production of cytotoxic T lymphocytes specific for a polypeptide of the invention, preferably a polypeptide incorporating a sequence of SEQ ID NO: 5 and 6. In a preferred embodiment, the animal to be so treated is a human patient.

15 The proteins encoded by the genes disclosed herein due to their expression, or elevated expression, in cancer cells, represent highly useful therapeutic targets for "targeted therapies" utilizing such affinity structures as, for example, antibodies coupled to some cytotoxic agent. In such methodology, it is advantageous that nothing need be known about the
20 endogenous ligands or binding partners for such cell surface molecules. Rather, an antibody or equivalent molecule that can specifically recognize the cell surface molecule (which could include an artificial peptide, a surrogate ligand, and the like) that is coupled to some agent that can induce cell death or a block in cell cycling offers therapeutic promise against these proteins.
25 Thus, such approaches include the use of so-called suicide "bullets" against intracellular proteins. For example, monoclonal antibodies may readily be produced by methods well known in the art, for example, the method of Kohler and Milstein (see: *Nature*, **256**:495 (1975)).

30 With the advent of methods of molecular biology and recombinant technology, it is now possible to produce antibody molecules by recombinant means and thereby generate gene sequences that code for specific amino

acid sequences found in the polypeptide structure of the antibodies. Such antibodies can be produced by either cloning the gene sequences encoding the polypeptide chains of said antibodies or by direct synthesis of said polypeptide chains, with *in vitro* assembly of the synthesized chains to form
5 active tetrameric (H₂L₂) structures with affinity for specific epitopes and antigenic determinants. This has permitted the ready production of antibodies having sequences characteristic of neutralizing antibodies from different species and sources.

Regardless of the source of the antibodies, or how they are
10 recombinantly constructed, or how they are synthesized, *in vitro* or *in vivo*, using transgenic animals, such as cows, goats and sheep, using large cell cultures of laboratory or commercial size, in bioreactors or by direct chemical synthesis employing no living organisms at any stage of the process, all antibodies have a similar overall 3 dimensional structure. This structure is
15 often given as H₂L₂ and refers to the fact that antibodies commonly comprise 2 light (L) amino acid chains and 2 heavy (H) amino acid chains. Both chains have regions capable of interacting with a structurally complementary antigenic target. The regions interacting with the target are referred to as "variable" or "V" regions and are characterized by differences in amino acid
20 sequence from antibodies of different antigenic specificity.

The variable regions of either H or L chains contains the amino acid sequences capable of specifically binding to antigenic targets. Within these sequences are smaller sequences dubbed "hypervariable" because of their
25 extreme variability between antibodies of differing specificity. Such hypervariable regions are also referred to as "complementarity determining regions" or "CDR" regions. These CDR regions account for the basic specificity of the antibody for a particular antigenic determinant structure.

30 The CDRs represent non-contiguous stretches of amino acids within the variable regions but, regardless of species, the positional locations of these critical amino acid sequences within the variable heavy and light chain

regions have been found to have similar locations within the amino acid sequences of the variable chains. The variable heavy and light chains of all antibodies each have 3 CDR regions, each non-contiguous with the others (termed L1, L2, L3, H1, H2, H3) for the respective light (L) and heavy (H) chains. The accepted CDR regions have been described by Kabat et al., *J. Biol. Chem.* **252**:6609-6616 (1977).

In all mammalian species, antibody polypeptides contain constant (i.e., highly conserved) and variable regions, and, within the latter, there are the CDRs and the so-called "framework regions" made up of amino acid sequences within the variable region of the heavy or light chain but outside the CDRs.

The antibodies disclosed according to the invention may also be wholly synthetic, wherein the polypeptide chains of the antibodies are synthesized and, possibly, optimized for binding to the polypeptides disclosed herein as being receptors. Such antibodies may be chimeric or humanized antibodies and may be fully tetrameric in structure, or may be dimeric and comprise only a single heavy and a single light chain. Such antibodies may also include fragments, such as Fab and F(ab₂)' fragments, capable of reacting with and binding to any of the polypeptides disclosed herein as being receptors.

In one aspect, the present invention relates to immunoglobulins, or antibodies, as described herein, that react with, especially where they are specific for, the polypeptides having amino acid sequences as disclosed herein, preferably those having an amino acid sequence of one of SEQ ID NO: 5 and 6. Such antibodies may commonly be in the form of a composition, especially a pharmaceutical composition. Such antibodies, by themselves, may have therapeutic value in that they are able to bind to, and thereby tie up, surface sites on cancerous cells. Where such sites have some type of function to perform (i.e., where they are surface enzymes, or channel structures, or structures that otherwise facilitate, actively or passively, the

transport of nutrients and other vital materials to the cell. Such nutrients serve to facilitate the growth and replication of the cell and molecules that bind to such sites and thereby interfere with such activities can prove to have a therapeutic effect in that the result of such binding is to remove sources of nutrients from such cells, thereby interfering with growth and replication. In like manner, such binding may serve to remove vital enzyme activities from the cell's functional repertoire, thereby also interfering with viability and/or the ability of the cell to multiply or metastasize. In addition, by binding to such surface sites, the antibodies may serve to prevent the cells from reacting to environmental agents, such as cytokines and the like, that may facilitate growth, replication and metastasis, thereby further reducing the cancerous status of such cell and ameliorating the cancerous condition in a patient, even without proving fatal to the cell or cells so affected.

The methods of the present invention also include processes wherein the cancer cell is contacted *in vivo* as well as *ex vivo* with an agent that comprises a portion, or is part of an overall molecular structure, having affinity for an expression product of a gene corresponding to a polynucleotide sequence as disclosed herein, preferably where the expression product is a cell surface structure, most preferably a polypeptide as disclosed herein, such as one that comprises an amino acid sequence of SEQ ID NO: 5 and 6. In one such embodiment, said portion having affinity for said expression product is an antibody, especially where said expression product is a polypeptide or oligopeptide or comprises an oligopeptide portion, or comprises a polypeptide.

In another aspect, the present invention also relates to an antibody that reacts with a polypeptide as disclosed herein, preferably a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 5 and 6. Such an antibody may be polyclonal, monoclonal, recombinant or synthetic in origin. In one such embodiment, said antibody is associated, either covalently or non-covalently, with a cytotoxic agent, for example, an apoptotic agent. It is thus contemplated that the antibody acts a targeted vector for guiding an associated therapeutic agent to a cancerous

cell, such as a cell expressing a polypeptide homologous to, if not identical to, a polypeptide as disclosed herein.

Where the cytotoxic agent is itself a polypeptide, said may be linked
5 directly to an antibody specific for a surface target on a cancer cell, such as
where the polypeptide represents an extension of the amino acid chain of the
antibody. In alternative embodiments, such molecules may be covalently
linked through a linker sequence of long or short duration, such as an amino
acid sequence of 5 to 10 residues in length. Where the cytotoxic agents is
10 some small organic molecule, such as a small organic compound, or some
type of apoptotic agent, this may be covalently bonded to the antibody
molecule or may be attached by some other type of non-covalent linkage,
including hydrophobic and electrostatic linkages. Methods for forming such
linkages, especially covalent linkages, are well known to those skilled in the
15 art.

The antibodies disclosed herein may also serve as targeting vectors for
much larger structures, such as liposomes. In one such embodiment, an
antibody is part of, or otherwise linked to, or associated with, a membranous
20 structure, preferably a liposome or possibly some type of cellular organelle,
which acts as a reservoir for a cytotoxic agent, such as ricin. The antibody
then acts to target said liposome to a cancerous tissue in an animal,
whereupon the liposome provides a source of cytotoxic agents for localized
treatment of a solid tumor or other type of neoplasm.

25 The present invention further encompasses an immunogenic
composition comprising a polypeptide disclosed herein, as well as
compositions formed using antibodies specific for these polypeptides.

30 Methods well known in the art for making formulations are found in, for
example, *Remington: The Science and Practice of Pharmacy*, (19th ed.) Ed.
A.R. Gennaro, 1995, Mack Publishing Company, Easton, PA. Formulations

for parenteral administration may, for example, contain excipients, sterile water, or saline, polyalkylene glycols such as polyethylene glycol, oils of vegetable origin, or hydrogenated naphthalenes. Biocompatible, biodegradable lactide polymer, lactide/glycolide copolymer, or polyoxyethylene-
5 polyoxypropylene copolymers may be used to control the release of the compounds. Other potentially useful parenteral delivery systems for agonists of the invention include ethylenevinyl acetate copolymer particles, osmotic pumps, implantable infusion systems, and liposomes. Formulations for inhalation may contain excipients, or example, lactose, or may be aqueous
10 solutions containing, for example, polyoxyethylene-9-lauryl ether, glycocholate and deoxycholate, or may be oily solutions for administration in the form of nasal drops, or as a gel. It should be noted that, where the therapeutic agent to be administered is an immunoconjugate, these sometimes contain chemical linkages that are somewhat labile in aqueous media and therefor
15 must be stored prior to administration in a more stable environment, such as in the form of a lyophilized powder.

Such an agent can be a single molecular structure, comprising both affinity portion and anti-cancer activity portions, wherein said portions are
20 derived from separate molecules, or molecular structures, possessing such activity when separated and wherein such agent has been formed by combining said portions into one larger molecular structure, such as where said portions are combined into the form of an adduct. Said anti-cancer and affinity portions may be joined covalently, such as in the form of a single
25 polypeptide, or polypeptide-like, structure or may be joined non-covalently, such as by hydrophobic or electrostatic interactions, such structures having been formed by means well known in the chemical arts. Alternatively, the anti-cancer and affinity portions may be formed from separate domains of a single molecule that exhibits, as part of the same chemical structure, more than one
30 activity wherein one of the activities is against cancer cells, or tumor formation or growth, and the other activity is affinity for an expression product produced by expression of genes related to the cancerous process or condition.

In one embodiment of the present invention, a chemical agent, such as a protein or other polypeptide, is joined to an agent, such as an antibody, having affinity for an expression product of a cancerous cell, such as a polypeptide or protein encoded by a gene related to the cancerous process, preferably a gene as disclosed herein according to the present invention, most preferably a polypeptide sequence disclosed herein. Thus, where the presence of said expression product is essential to tumor initiation and/or growth, binding of said agent to said expression product will have the effect of negating said tumor promoting activity. In one such embodiment, said agent is an apoptosis-inducing agent that induces cell suicide, thereby killing the cancer cell and halting tumor growth.

Other genes within the cancer cell that are regulated in a manner similar to that of the genes disclosed herein and thus change their expression in a coordinated way in response to chemical compounds represent genes that are located within a common metabolic, signaling, physiological, or functional pathway so that by analyzing and identifying such commonly regulated groups of genes (groups that include the gene, or similar sequences, disclosed according to the invention, one can (a) assign known genes and novel genes to specific pathways and (b) identify specific functions and functional roles for novel genes that are grouped into pathways with genes for which their functions are already characterized or described. For example, one might identify a group of 10 genes, at least one of which is the gene as disclosed herein, that change expression in a coordinated fashion and for which the function of one, such as the polypeptide encoded by the sequence disclosed herein, is known then the other genes are thereby implicated in a similar function or pathway and may thus play a role in the cancer-initiating or cancer-facilitating process. In the same way, if a gene were found in normal cells but not in cancer cells, or happens to be expressed at a higher level in normal as opposed to cancer cells, then a similar conclusion may be drawn as to its involvement in cancer, or other diseases. Therefore, the processes disclosed according to the present invention at once

provide a novel means of assigning function to genes, i.e. a novel method of functional genomics, and a means for identifying chemical compounds that have potential therapeutic effects on specific cellular pathways. Such chemical compounds may have therapeutic relevance to a variety of diseases
5 outside of cancer as well, in cases where such diseases are known or are demonstrated to involve the specific cellular pathway that is affected.

The polypeptides disclosed herein, preferably those of SEQ ID NO: 5 and 6, also find use as vaccines in that, where the polypeptide represents a
10 surface protein present on a cancer cell, such polypeptide may be administered to an animal, especially a human being, for purposes of activating cytotoxic T lymphocytes (CTLs) that will be specific for, and act to lyse, cancer cells in said animal. Where used as vaccines, such polypeptides are present in the form of a pharmaceutical composition. The present
15 invention may also employ polypeptides that have the same, or similar, immunogenic character as the polypeptides of SEQ ID NO: 5 and 6 and thereby elicit the same, or similar, immunogenic response after administration to an animal, such as an animal at risk of developing cancer, or afflicted therewith. Thus, the polypeptides disclosed according to the invention will
20 commonly find use as immunogenic compositions.

Expression of a gene corresponding to a polynucleotide disclosed herein, when in normal tissues, may indicate a predisposition towards development of breast or endometrium cancer. The encoded polypeptide
25 might then present a potentially useful cell surface target for therapeutic molecules such as cytolytic antibodies, or antibodies attached to cytotoxic, or cytolytic, agents. .

The present invention specifically contemplates use of antibodies
30 against the polypeptides encoded by the polynucleotides corresponding to the genes disclosed herein, whereby said antibodies are conjugates to one or more cytotoxic agents so that the antibodies serve to target the conjugated

immunotoxins to a region of cancerous activity, such as a solid tumor. For many known cytotoxic agents, lack of selectivity has presented a drawback to their use as therapeutic agents in the treatment of malignancies. For example, the class of two-chain toxins, consisting of a binding subunit (or B-chain) 5 linked to a toxic subunit (A-chain) are extremely cytotoxic. Thus, such agents as ricin, a protein isolated from castor beans, kills cells at very low concentrations (even less than 10^{-11} M) by inactivating ribosomes in said cells (see, for example, Lord et al., Ricin: structure, mode of action, and some current applications. *Faseb J*, 8: 201-208 (1994), and Blättler et al., Realizing 10 the full potential of immunotoxins. *Cancer Cells*, 1: 50-55 (1989)). While isolated A-chains of protein toxins that functionally resemble ricin A-chain are only weakly cytotoxic for intact cells (in the concentration range of 10^{-7} to 10^{-6} M), they are very potent cytotoxic agents inside the cells. Thus, a single molecule of the A-subunit of diphtheria toxin can kill a cell once inside (see: 15 Yamaizumi et al., One molecule of diphtheria toxin fragment A introduced into a cell can kill the cell. *Cell*, 15: 245-250, 1978).

The present invention solves this selectivity problem by using antibodies specific for antigens present on cancer cells to target the cytotoxins to said cells. In addition, use of antibodies decreases toxicity because the 20 antibodies are non-toxic until they reach the tumor and, because the cytotoxin is bound to the antibody, it is presented with less opportunity to cause damage to non-targeted tissues.

In addition, use of such antibodies alone can provide therapeutic 25 effects on the tumor through the antibody-dependent cellular cytotoxic response (ADCC) and complement-mediated cell lysis mechanisms.

A number of recombinant immunotoxins (for example, consisting of Fv regions of cancer specific antibodies fused to truncated bacterial toxins) are 30 well known (see, for example, Smyth et al., Specific targeting of chlorambucil to tumors with the use of monoclonal antibodies, *J. Natl. Cancer Inst.*, 76(3):503-510 (1986); Cho et al., Single-chain Fv/folate conjugates mediate

efficient lysis of folate-receptor-positive tumor cells, *Bioconjug. Chem.*, 8(3):338-346 (1997)). As noted in the literature, these may contain, for example, a truncated version of *Pseudomonas* exotoxin as a toxic moiety but the toxin is modified in such a manner that by itself it does not bind to normal human cells, but it retains all other functions of cytotoxicity. Here, recombinant antibody fragments target the modified toxin to cancer cells which are killed, such as by direct inhibition of protein synthesis, or by concomitant induction of apoptosis. Cells that are not recognized by the antibody fragment, because they do not carry the cancer antigen, are not affected. Good activity and specificity has been observed for many recombinant immunotoxins in *in vitro* assays using cultured cancer cells as well as in animal tumor models. Ongoing clinical trials provide examples where the promising pre-clinical data correlate with successful results in experimental cancer therapy. (see, for example, Brinkmann U., Recombinant antibody fragments and immunotoxin fusions for cancer therapy, *In Vivo* (2000) 14:21-27).

While the safety of employing immunoconjugates in humans has been established, *in vivo* therapeutic results have been less impressive. Because clinical use of mouse MAbs in humans is limited by the development of a foreign anti-globulin immune response by the human host, genetically engineered chimeric human-mouse MAbs have been developed by replacing the mouse Fc region with the human constant region. In other cases, the mouse antibodies have been "humanized" by replacing the framework regions of variable domains of rodent antibodies by their human equivalents. Such humanized and engineered antibodies can even be structurally arranged to have specificities and effector functions determined by design and which characteristics do not appear in nature. The development of bispecific antibodies, having different binding ends so that more than one antigenic site can be bound, have proven useful in targeting cancer cells. Thus, such antibody specificity has been improved by chemical coupling to various agents such as bacterial or plant toxins, radionuclides or cytotoxic drugs and other agents. (see, for example, Bodey, B. et al). Genetically engineered monoclonal antibodies for direct anti-neoplastic treatment and cancer cell

specific delivery of chemotherapeutic agents. *Curr Pharm Des* (2000) Feb;6(3):261-76). See also, Garnett, M. C., Targeted drug conjugates: principles and progress. *Adv. Drug Deliv. Rev.* (2001 Dec 17) **53**(2):171-216; Brinkmann et al., Recombinant immunotoxins for cancer therapy. *Expert Opin Biol Ther.* (2001) **1**(4):693-702.

Among the cytotoxic agents specifically contemplated for use as immunoconjugates according to the present invention are Calicheamicin, a highly toxic enediyne antibiotic isolated from *Micromonospora echinospora ssp. Calichensis*, and which binds to the minor groove of DNA to induce double strand breaks and cell death (see: Lee et al., Calicheamicins, a novel family of antitumor antibiotics. 1. Chemistry and partial structure of calicheamicin g₁. *J Am Chem Soc*, **109**: 3464-3466 (1987); Zein et al., Calicheamicin gamma 1I: an antitumor antibiotic that cleaves double-stranded DNA site specifically, *Science*, **240**: 1198-1201 (1988)). Useful derivatives of the calicheamicins include mylotarg and 138H11-Cam θ . Mylotarg is an immunoconjugate of a humanized anti-CD33 antibody (CD33 being found in leukemic cells of most patients with acute myeloid leukemia) and N-acetyl gamma colicheamicin dimethyl hydrazide, the latter of which is readily coupled to an antibody of the present invention (in place of the anti-CD33 but which can also be humanized by substitution of human framework regions into the antibody during production as described elsewhere herein) to form an immunoconjugate of the invention. (see: Hamann et al. Gemtuzumab Ozogamicin, A Potent and Selective Anti-CD33 Antibody-Calicheamicin Conjugate for Treatment of Acute Myeloid Leukemia, *Bioconjug. Chem.* **13**, 47-58 (2002)) For use with 138H11-Cam θ , 138H11 is an anti- γ -glutamyl transferase antibody coupled to theta calicheamicin through a disulfide linkage and found useful *in vitro* against cultured renal cell carcinoma cells. (see: Knoll et al., Targeted therapy of experimental renal cell carcinoma with a novel conjugate of monoclonal antibody 138H11 and calicheamicin θ_1^I , *Cancer Res*, **60**: 6089-6094 (2000) The same linkage may be utilized to link

this cytotoxic agent to an antibody of the present invention, thereby forming a targeting structure for breast or endometrium cancer cells.

Also useful in forming the immunoconjugates of the invention is DC1, a
5 disulfide-containing analog of adozelesin, that kills cells by binding to the
minor groove of DNA, followed by alkylation of adenine bases. Adozelesin is a
structural analog of CC-1065, an anti-tumor antibiotic isolated from microbial
fermentation of *Streptomyces zelensis*, and is about 1,000 fold more toxic to
10 cultured cell lines than other DNA interacting agents, such as cis-platin and
doxorubicin. This agent is readily linked to antibodies through the disulfide
bond of adozelesin. (see: Chari et al., Enhancement of the selectivity and
antitumor efficacy of a CC-1065 analogue through immunoconjugate
formation, *Cancer Res*, **55**: 4079-4084 (1995)).

Maytansine, a highly cytotoxic microtubular inhibitor isolated from the
15 shrub *Maytenus serrata* found to have little value in human clinical trials, is
much more effective in its derivatized form, denoted DM1, containing a
disulfide bond to facilitate linkage to antibodies, is up to 10-fold more cytotoxic
(see: Chari et al., Immunoconjugates containing novel maytansinoids:
promising anticancer drugs, *Cancer Res*, **52**: 127-131 (1992)). These same *in*
20 *vitro* studies showed that up to four DM1 molecules could be linked to a single
immunoglobulin without destroying the binding affinity. Such conjugates have
been used against breast cancer antigens, such as the *neu/HER2/erbB-2*
antigen. (see: Goldmacher et al., Immunogen, Inc., (2002) *in press*); also see
Liu, C. et al., Eradication of large colon tumor xenografts by targeted delivery
25 of maytansinoids, *Proc. Natl. Acad. Sci. USA*, **93**, 8618-8623 (1996)). For
example, Liu et al. (1996) describes formation of an immunoconjugate of the
maytansinoid cytotoxin DM1 and C242 antibody, a murine IgG1
immunoglobulin, available from Pharmacia and which has affinity for a mucin-
like glycoprotein variably expressed by human colorectal cancers. The latter
30 immunoconjugate was prepared according to Chari et al., *Cancer Res.*,
52:127-131 (1992) and was found to be highly cytotoxic against cultured colon
cancer cells as well as showing anti-tumor effects *in vivo* in mice bearing

subcutaneous COLO 205 human colon tumor xenografts using doses well below the maximum tolerated dose.

In addition, there are a variety of protein toxins (cytotoxic proteins), which include a number of different classes, such as those that inhibit protein synthesis: ribosome-inactivating proteins of plant origin, such as ricin, abrin, gelonin, and a number of others, and bacterial toxins such as pseudomonas exotoxin and diphtheria toxin.

Another useful class is the one including taxol, taxotere, and taxoids. Specific examples include paclitaxel (taxol), its analog docetaxel (taxotere), and derivatives thereof. The first two are clinical drugs used in treating a number of tumors while the taxoids act to induce cell death by inhibiting the de-polymerization of tubulin. Such agents are readily linked to antibodies through disulfide bonds without disadvantageous effects on binding specificity.

In one instance, a truncated *Pseudomonas* exotoxin was fused to an anti-CD22 variable fragment and used successfully to treat patients with chemotherapy-resistant hairy-cell leukemia. (see: Kreitman et al., Efficacy of the anti-CD22 recombinant immunotoxin BL22 in chemotherapy-resistant hairy-cell leukemia, *N Engl J Med*, **345**: 241-247 (2001)) Conversely, the cancer-linked peptides of the present invention offer the opportunity to prepare antibodies, recombinant or otherwise, against the appropriate antigens to target solid tumors, preferably those of malignancies of breast or endometrium tissue, using the same or similar cytotoxic conjugates. Thus, many of the previously used immunoconjugates have been formed using antibodies against general antigenic sites linked to cancers whereas the antibodies formed using the peptides disclosed herein are more specific and target the antibody-cytotoxic agent to a particular tissue or organ, thus further reducing toxicity and other undesirable side effects.

In addition, the immunoconjugates formed using the antibodies prepared against the cancer-linked antigens disclosed herein can be formed by any type of chemical coupling. Thus, the cytotoxic agent of choice, along with the immunoglobulin, can be coupled by any type of chemical linkage, 5 covalent or non-covalent, including electrostatic linkage, to form the immunoconjugates of the present invention.

When used as immunoconjugates, the antitumor agents of the present invention represent a class of pro-drugs that are relatively non-toxic when first 10 administered to an animal (due mostly to the stability of the immunoconjugate), such as a human patient, but which are targeted by the conjugated immunoglobulin to a cancer cell where they then exhibit good toxicity. The tumor-related, associated, or linked, antigens, preferably those presented herein, serve as targets for the antibodies (monoclonal, 15 recombinant, and the like) specific for said antigens. The end result is the release of active cytotoxic agent inside the cell after binding of the immunoglobulin portion of the immunoconjugate.

The cited references describe a number of useful procedures for the 20 chemical linkage of cytotoxic agents to immunoglobulins and the disclosures of all such references cited herein are hereby incorporated by reference in their entirety. For other reviews see Ghetie et al., Immunotoxins in the therapy of cancer: from bench to clinic, *Pharmacol Ther*, **63**: 209-234 (1994), Pietersz et al. The use of monoclonal antibody immunoconjugates in cancer therapy, 25 *Adv Exp Med Biol*, **353**:169-179 (1994), and Pietersz, G. A. The linkage of cytotoxic drugs to monoclonal antibodies for the treatment of cancer, *Bioconjug Chem*, **1**:89-95 (1990).

Thus, the present invention provides highly useful cancer-associated 30 antigens for generation of antibodies for linkage to a number of different cytotoxic agents which are already known to have some *in vitro* toxicity and possess chemical groups available for linkage to antibodies.

The present invention also relates to a process that comprises a method for producing a product, including the generation of test data, comprising identifying an agent according to one of the disclosed processes for identifying such an agent (i.e., the therapeutic agents identified according to the assay procedures disclosed herein) wherein said product is the data collected with respect to said agent as a result of said identification process, or assay, and wherein said data is sufficient to convey the chemical character and/or structure and/or properties of said agent. For example, the present invention specifically contemplates a situation whereby a user of an assay of the invention may use the assay to screen for compounds having the desired enzyme modulating activity and, having identified the compound, then conveys that information (i.e., information as to structure, dosage, etc) to another user who then utilizes the information to reproduce the agent and administer it for therapeutic or research purposes according to the invention. For example, the user of the assay (user 1) may screen a number of test compounds without knowing the structure or identity of the compounds (such as where a number of code numbers are used the first user is simply given samples labeled with said code numbers) and, after performing the screening process, using one or more assay processes of the present invention, then imparts to a second user (user 2), verbally or in writing or some equivalent fashion, sufficient information to identify the compounds having a particular modulating activity (for example, the code number with the corresponding results). This transmission of information from user 1 to user 2 is specifically contemplated by the present invention.

It should be cautioned that, in carrying out the procedures of the present invention as disclosed herein, whether to form immunoconjugates or screen for other antitumor agents using the genes and polypeptides disclosed herein, any reference to particular buffers, media, reagents, cells, culture conditions and the like are not intended to be limiting, but are to be read so as to include all related materials that one of ordinary skill in the art would

recognize as being of interest or value in the particular context in which that discussion is presented. For example, it is often possible to substitute one buffer system or culture medium for another and still achieve similar, if not identical, results. Those of skill in the art will have sufficient knowledge of such systems and methodologies so as to be able, without undue experimentation, to make such substitutions as will optimally serve their purposes in using the methods and procedures disclosed herein.

The present invention will now be further described by way of the following non-limiting example. In applying the disclosure of the example, it should be kept clearly in mind that other and different embodiments of the methods disclosed according to the present invention will no doubt suggest themselves to those of skill in the relevant art. The following example shows how a potential anti-neoplastic agent may be identified using one or more of the genes disclosed herein.

EXAMPLE

Determination of Gene Inhibitory Activity of an Anti-neoplastic Agent

SW480 cells are grown to a density of 10^5 cells/cm² in Leibovitz's L-15 medium supplemented with 2 mM L-glutamine (90%) and 10% fetal bovine serum. The cells are collected after treatment with 0.25% trypsin, 0.02% EDTA at 37°C for 2 to 5 minutes. The trypsinized cells are then diluted with 30 ml growth medium and plated at a density of 50,000 cells per well in a 96 well plate (100 µl/well). The following day, cells are treated with either compound buffer alone, or compound buffer containing a chemical agent to be tested, for 24 hours. The media is then removed, the cells lysed and the RNA recovered using the RNAeasy reagents and protocol obtained from Qiagen. RNA is quantitated and 10 ng of sample in 1 µl are added to 24 µl of Taqman reaction mix containing 1X PCR buffer, RNAsin, reverse transcriptase, nucleoside triphosphates, amplitaq gold, tween 20, glycerol, bovine serum albumin (BSA)

and specific PCR primers and probes for a reference gene (18S RNA) and a test gene (Gene X). Reverse transcription is then carried out at 48°C for 30 minutes. The sample is then applied to a Perlin Elmer 7700 sequence detector and heat denatured for 10 minutes at 95°C. Amplification is performed through 40 cycles using 15 seconds annealing at 60°C followed by a 60 second extension at 72°C and 30 second denaturation at 95°C. Data files are then captured and the data analyzed with the appropriate baseline windows and thresholds.

10 The quantitative difference between the target and reference gene is then calculated and a relative expression value determined for all of the samples used. In this way, the ability of a chemotherapeutic agent to effectively and selectively reduce the activity of a cancer-specific gene is readily ascertained. The overall expression of the cancer-specific gene, as modulated by one chemical agent relative to another, is also determined. Chemical agents having the most effect in reducing gene activity are thereby identified as the most anti-neoplastic.

20

References:

25 Walter A. Blättler and Ravi Chari: Drugs to enhance the therapeutic potency of anti-cancer antibodies: antibody-drug conjugates as tumor-activated prodrugs. In *Anticancer Agents - Frontiers in Cancer Chemotherapy* (Iwao Ojima, Gregory D. Vite, Karl-Heinz Altmann, Eds.), American Chemical Society, pp. 317-338 (2001).

30 Dan L. Longo, Patricia L. Duffey, John G. Gribben, Elaine S. Jaffe, Brendan D. Curti, Barry L. Gause, John E. Janik, Virginia M. Braman, Dixie Esseltine, Wyndham H. Wilson, Dwight Kaufman, Robert E. Wittes, Lee M. Nadler, and Walter J. Urba: Combination chemotherapy followed by an Immunotoxin (Anti-B4-blocked Ricin) in patients with indolent lymphoma: results of a Phase II study. *Cancer J.* 6, 146-150 (2000).

35 Walter A. Blättler and John M. Lambert: Preclinical immunotoxin development. In *Monoclonal Antibody-Based Therapy of Cancer* (M. Grossbard, Ed.), Marcel Dekker, Inc. NY, NY, pp. 1-22 (1998).

- Ravi V. J. Chari: Targeted delivery of chemotherapeutics: tumor-activated prodrug therapy. In *Advanced Drug Delivery Reviews*, Elsevier Science B.V., pp. 89-104 (1998).
- 5 David T. Scadden, David P. Schenkein, Zale Bernstein, Barry Luskey, John Doweiko, Anil Tulpule, and Alexandra M. Levine: Immunotoxin combined with chemotherapy for patients with AIDS-related Non-Hodgkin's Lymphoma. *Cancer* 83, 2580-2587 (1998).
- 10 Changnian Liu and Ravi VJ Chari: The development of antibody delivery systems to target cancer with highly potent maytansinoids. *Exp. Opi. Invest. Drugs* 6, 169-172 (1997).
- A. C. Goulet, Viktor S. Goldmacher, John M. Lambert, C. Baron, Dennis C. Roy and E. Kouassi: Conjugation of blocked ricin to an anti-CD19 monoclonal antibody increases antibody-induced cell calcium mobilization and CD19 internalization. *Blood* 90, 2364-2375 (1997).
- 15 Changnian Liu, John M. Lambert, Beverly A. Teicher, Walter A. Blättler, and Rosemary O'Connor: Cure of multidrug-resistant human B-cell lymphoma xenografts by combinations of anti-B4-blocked ricin and chemotherapeutic drugs. *Blood* 87, 3892-3898 (1996).
- 20 Rajeeva Singh, Lana Kats, Walter A. Blättler, and John M. Lambert: Formation of N-Substituted 2-Iminothiolanes when amino groups in proteins and peptides are modified by 2-Iminothiolane. *Anal. Biochem.* 236, 114-125 (1996).
- 25 Changnian Liu, B. Mitra Tadayoni, Lizabeth A. Bourret, Kristin M. Mattocks, Susan M. Derr, Wayne C. Widdison, Nancy L. Kedersha, Pamela D. Ariniello, Victor S. Goldmacher, John M. Lambert, Walter A. Blättler, and Ravi V.J. Chari: Eradication of large colon tumor xenografts by targeted delivery of maytansinoids. *Proc. Natl. Acad. Sci. USA* 93, 8618-8623 (1996).
- 30 Denis C. Roy, Sophie Ouellet, Christiane Le Houiller, Pamela D. Ariniello, Claude Perreault and John M. Lambert: Elimination of neuroblastoma and small-cell lung cancer cells with an anti-neural cell adhesion molecule immunotoxin. *J. Natl. Cancer Inst.* 88, 1136-1145 (1996).
- Walter A. Blättler, Ravi V.J. Chari and John M. Lambert: Immunoconjugates. In *Cancer Therapeutics: Experimental and Clinical Agents*. (B. Teicher, Ed.), Humana Press, Totowa, NJ, pp. 371-394 (1996).
- 35 Michael L Grossbard, John M. Lambert, Victor S. Goldmacher, Arnold S. Freedman, Jeanne Kinsella, Danny P. Ducello, Susan N. Rabinowe, Laura Elisea, Felice Carol, James A. Taylor, Walter A. Blättler, Carol L. Epstein, and Lee M. Nadler: Anti-B4-blocked Ricin: A phase I trial of 7 day continuous infusion in patients with B-cell neoplasms. *J. Clin. Oncol.* 11, 726-737 (1993).
- 40 Michael L. Grossbard, John G. Gribben, Arnold S. Freedman, John M. Lambert, Jeanne Kinsella, Susan N. Rabinowe, Laura Eliseo, James A. Taylor, Walter A. Blättler, Carol L. Epstein, and Lee M. Nadler: Adjuvant

immunotoxin therapy with anti-B4-blocked ricin following autologous bone marrow transplantation for patients with B-cell Non-Hodgkin's lymphoma. *Blood* 81, 2263-2271 (1993).

5 Sudhir A. Shah, Patricia M. Halloran, Cynthia A. Ferris, Beth A. Levine, Elizabeth A. Bourret, Victor S. Goldmacher, and Walter A. Blättler: Anti-B4-blocked Ricin immunotoxin shows therapeutic efficacy in four different SCID mouse tumor models. *Cancer Res.* 53, 1360-1367 (1993).

10 Ravi V.J. Chari, Bridget A. Martell, Jonathan L. Gross, Sherilyn B. Cook, Sudhir A. Shah, Walter A. Blättler, Sara J. McKenzie, and Victor S. Goldmacher: Immunoconjugates containing novel maytansinoids: promising anti-cancer drugs. *Cancer Res.* 52, 127-131 (1992).

John M. Lambert, Peter D. Senter, Annie Yau-Young, Walter A. Blättler, and Victor S. Goldmacher: Purified immunotoxins that are reactive with human lymphoid cells. *J. Biol. Chem.* 250, 12035-12041 (1985).

SEQUENCE LISTING

<110> Avalon Pharmaceuticals

<120> Cancer-Linked Gene as Target for Chemotherapy

<130> 689290-163

<140>

<141>

<150> US/60/386,793

<151> 2002-06-07

<160> 6

<170> PatentIn version 3.0

<210> 1

<211> 1669

<212> DNA

<213> Homo sapiens

<400> 1

tgtgagtcac	caaggaaggc	agcggcagct	ccactcagcc	agtacccaga	tacgctggga	60
accttcccca	gccatggctt	ccctggggca	gatcctcttc	tggagcataa	ttagcatcat	120
cattattctg	gctggagcaa	ttgcactcat	cattggcttt	ggtatttcag	aagtctctgt	180
ctggctttca	gcaatgaagg	gtttggttgt	agaagttcca	aggcttcctt	tagcattgat	240
ctttgcttcc	tgaactgcag	ggagacactc	catcacagtc	actactgtcg	cctcagctgg	300
gaacattggg	gaggatggaa	tcctgagctg	cacttttgaa	cctgacatca	aactttctga	360
tatcgtgata	caatggctga	aggaaggtgt	tttaggcttg	gtccatgagt	tcaaagaagg	420
caaagatgag	ctgtcggagc	aggatgaaat	gttcagaggc	cggacagcag	tgtttgctga	480
tcaagtgata	gttggcaatg	cctctttgcg	gctgaaaaac	gtgcaactca	cagatgctgg	540
cacctacaaa	tgttatatca	tcacttctaa	aggcaagggg	aatgctaacc	ttgagtataa	600
aactggagcc	ttcagcatgc	cggaagtga	tgtggactat	aatgccagct	cagagacctt	660
gcggtgtgag	gctccccgat	ggttcccca	gcccacagtg	gtctgggcat	cccaagttga	720
ccagggagcc	aacttctcgg	aagtctccaa	taccagcttt	gagctgaact	ctgagaatgt	780
gaccatgaag	gttgtgtctg	tgctctacaa	tgttacgatc	aacaacacat	actcctgtat	840
gattgaaaat	gacattgcca	aagcaacagg	ggatatcaaa	gtgacagaat	cggagatcaa	900
aaggcggagt	cacctacagc	tgctaaactc	aaaggcttct	ctgtgtgtct	cttctttctt	960
tgccatcagc	tgggcacttc	tgctctcag	cccttacctg	atgctaaaat	aatgtgcctc	1020
ggccacaaaa	aagcatgcaa	agtcattggt	acaacagggg	tctacagaac	tatttcacca	1080
ccagatatga	cctagtttta	tatttctggg	aggaaatgaa	ttcatatcta	gaagtctgga	1140
gtgagcaaac	aagagcaaga	aacaaaaaga	agccaaaagc	agaaggctcc	aatatgaaca	1200
agataaatct	atcttcaaag	acatattaga	agttgggaaa	ataattcatg	tgaactagac	1260
aagtgtgtta	agagtgataa	gtaaaatgca	cgtggagaca	agtgcattcc	cagatctcag	1320
ggacctcccc	ctgcctgtca	cctggggagt	gagaggacag	gatagtgcac	gttctttgtc	1380
tctgaattht	tagttatatg	tgctgtaatg	ttgctctgag	gaagcccctg	gaaagtctat	1440
cccaacatat	ccacatctta	tattccacaa	attaagctgt	agtatgtacc	ctaagacgct	1500
gctaattgac	tgccacttcg	caactcaggg	gcggtctgac	tttagtaatg	ggtcaaatga	1560
ttcactthtt	atgatgcttc	caaaggtgcc	ttggcttctc	ttcccaactg	acaaatgcca	1620
aagttgagaa	aaatgatcat	aatttttagca	taaacagagc	agtcggcga		1669

<210> 2

<211> 1004

<212> DNA

<213> Homo sapiens

<400> 2

cttcttttta	aacaaacaaa	tgcgggttta	tttctcagat	gatgttcac	cgtgaatggt	60
ccaggggaag	acctttcacc	ttgtctatat	ggcattatgt	catcacaagc	tctgaggctt	120
ctcctttcca	tcctgcgtgg	acagctaaga	cctcagtttt	caatagcatc	tagagcagtg	180
ggactcagct	gggggtgattt	cgccccccat	ctccggggga	atgtctgaag	acaattttgg	240
ttacctcaat	gagggagtgg	aggaggatac	agtgctacta	ccaactagtg	gataaaggcc	300
agggatgctg	ctcaacctcc	taccatgtac	aggacgtctc	cccattacaa	ctacccaatc	360
cgaagtgtca	actgtgtcag	gactaagaaa	ccctggtttt	gagtagaaaa	gggctggaa	420
agaggggagc	caacaaatct	gtctgcttcc	tcacattagt	cattggcaaa	taagcattct	480
gtctctttgg	ctgctgcctc	agcacagaga	gccagaactc	tatcgggcac	caggataaca	540
tctctcagtg	aacagagttg	acaaggccta	tgggaaatgc	ctgatgggat	tatcttcagc	600
ttgttgagct	tctaagtttc	tttcccttca	ttctaccctg	caagccaagt	tctgtaagag	660
aatgcctga	gttctagctc	aggttttctt	actctgaatt	tagatctcca	gacccttctt	720
ggccacaatt	caaattaag	caacaaacat	ataccttcca	tgaagcacac	acagactttt	780
gaaagcaagg	acaatgactg	cttgaattga	ggccttgagg	aatgaagctt	tgaaggaaaa	840
gaatactttg	tttccagccc	ccttcccaca	ctcttcatgt	gttaaccact	gccttcctgg	900
accttgagc	cacggtgact	gtattacatg	ttgttataga	aaactgattt	tagagttctg	960
atcgttcaag	agaatgatta	aatatacatt	tcctaaaaaa	atgt		1004

<210> 3

<211> 1808

<212> DNA

<213> Homo sapiens

<400> 3

tgtgagtcac	caaggaaggc	agcggcagct	ccactcagcc	agtaccaga	tacgctggga	60
accttcccca	gccatggctt	ccctggggca	gatcctcttc	tggagcataa	ttagcatcat	120
cattattctg	gctggagcaa	ttgcactcat	cattggcttt	ggtatttcag	ggagacactc	180
catcacagtc	actactgtcg	cctcagctgg	gaacattggg	gaggatggaa	tcctgagctg	240
cacttttgaa	cctgacatca	aactttctga	tatcgtgata	caatggctga	aggaagggtg	300
tttaggcttg	gtccatgagt	tcaaagaagg	caaagatgag	ctgtcggagc	aggatgaaat	360
gttcagaggc	cggacagcag	tgtttgctga	tcaagtgata	gttggcaatg	cctctttgcg	420
gctgaaaaac	gtgcaactca	cagatgctgg	cacctacaaa	tgttatatca	tcacttctaa	480
aggcaagggg	aatgctaacc	ttgagtataa	aactggagcc	ttcagcatgc	cggagtgaa	540
tgtggactat	aatgccagct	cagagacctt	gcggtgtgag	gctccccgat	ggttccccca	600
gccacagtg	gtctgggcat	cccaagttga	ccagggagcc	aacttctcgg	aagtctccaa	660
taccagcttt	gagctgaact	ctgagaatgt	gaccatgaag	gttgtgtctg	tgctctacaa	720
tgttacgatc	aacaacacat	actcctgtat	gattgaaaat	gacattgcca	aagcaacagg	780
ggatatcaaa	gtgacagaat	cggagatcaa	aaggcggagt	cacctacagc	tgctaaactc	840
aaaggcttct	ctgtgtgtct	cttctttctt	tgccatcagc	tgggcacttc	tgctctcag	900
cccttacctg	atgctaaaat	aatgtgcctc	ggccacaaaa	aagcatgcaa	agtcattggt	960
acaacagggg	tctacagaac	tatttcacca	ccagatatga	cctagtttta	tatttctggg	1020
aggaaatgaa	ttcatatcta	gaagtctgga	gtgagcaaac	aagagcaaga	aacaaaaaga	1080
agccaaaagc	agaaggctcc	aatatgaaca	agataaatct	atcttcaaag	acatattaga	1140
agttgggaaa	ataattcatg	tgaactagat	gtcaactgtg	tcaggactaa	gaaaccctgg	1200
ttttgagtag	aaaagggcct	ggaaagaggg	gagccaacaa	atctgtctgc	ttcctcacat	1260
tagtcattgg	caaataagca	ttctgtctct	ttggctgctg	cctcagcaca	gagagccaga	1320
actctatcgg	gcaccaggat	aacatctctc	agtgaacaga	gttgacaagg	cctatgggaa	1380
atgcctgatg	ggattatctt	cagcttggtg	agcttctaag	tttctttccc	ttcattctac	1440
cctgcaagcc	aagttctgta	agagaaatgc	ctgagttcta	gctcaggttt	tcttactctg	1500
aatttagatc	tccagaccct	tcttgccac	aattcaaatt	aaggcaacaa	acataacct	1560
tccatgaagc	acacacagac	ttttgaaagc	aaggacaatg	actgcttgaa	ttgaggcctt	1620
gaggaatgaa	gctttgaagg	aaaagaatac	tttgtttcca	gcccccttcc	cacactcttc	1680
atgtgttaac	cactgccttc	ctggaccttg	gagccacggg	gactgtatta	catgttgta	1740
tagaaaactg	atthtagagt	tctgatcggt	caagagaatg	attaaatata	catttcctaa	1800
aaaaatgt						1808

<210> 4

<211> 1898

<212> DNA

<213> Homo sapiens

<400> 4

```

tgtgagtcac caaggaaggc agcggcagct ccaactcagcc agtaccacaga tacgctggga      60
accttcccca gccatggctt ccctggggca gatcctcttc tggagcataa ttagcatcat      120
cattattctg gctggagcaa ttgcactcat cattggcttt ggtatttcag aagtctctgt      180
ctggctttca gcaatgaagg gtttggttgt agaagttcca aggcttccct tagcattgat      240
ctttgcttcc tgaactgcag ggagacactc catcacagtc actactgtcg cctcagctgg      300
gaacattggg gaggatggaa tcctgagctg cacttttgaa cctgacatca aactttctga      360
tatcgtgata caatggctga aggaagggtg tttaggcttg gtccatgagt tcaaagaagg      420
caaagatgag ctgtcggagc aggatgaaat gttcagaggc cggacagcag tgtttgctga      480
tcaagtgata gttggcaatg cctctttgcg gctgaaaaac gtgcaactca cagatgctgg      540
cacctacaaa tgttatatca tcacttctaa aggcaagggg aatgctaacc ttgagtataa      600
aactggagcc ttcagcatgc cggaagtgaa tgtggactat aatgccagct cagagacctt      660
gcggtgtgag gctccccgat ggttccccca gccacacagt gtctgggcat cccaagttga      720
ccagggagcc aacttctcgg aagtctccaa taccagcttt gagctgaact ctgagaatgt      780
gaccatgaag gttgtgtctg tgctctacaa tgttacgatc aacaacacat actcctgtat      840
gattgaaaat gacattgcca aagcaacagg ggatatcaaa gtgacagaat cggagatcaa      900
aaggcggagt cacctacagc tgctaaactc aaaggcttct ctgtgtgtct cttctttctt      960
tgccatcagc tgggcacttc tgctctcag cccttacctg atgctaaaat aatgtgcctc     1020
ggccacaaaa aagcatgcaa agtcattggt acaacagggg tctacagaac tatttcacca     1080
ccagatatga cctagtttta ttttctggg aggaaatgaa ttcatatcta gaagtctgga     1140
gtgagcaaac aagagcaaga acaaaaaaga agccaaaagc agaaggctcc aatatgaaca     1200
agataaatct atcttcaaag acatattaga agttgggaaa ataattcatg tgaactagat     1260
gtcaactgtg tcaggactaa gaaaccctgg ttttgagtag aaaagggcct ggaaagaggg     1320
gagccaacaa atctgtctgc ttcctcacat tagtcattgg caaataagca ttctgtctct     1380
ttggctgctg cctcagcaca gagagccaga actctatcgg gcaccaggat aacatctctc     1440
agtgaacaga gttgacaagg cctatgggaa atgcctgatg ggattatctt cagcttggtg     1500
agcttctaag tttctttccc ttcattctac cctgcaagcc aagttctgta agagaaatgc     1560
ctgagttcta gctcaggttt tcttactctg aatttagatc tccagaccct tcctggccac     1620
aattcaaatt aaggcaacaa acatatacct tccatgaagc acacacagac ttttgaaagc     1680
aaggacaatg actgcttgaa ttgaggcctt gaggaatgaa gctttgaagg aaaagaatac     1740
tttgtttcca gcccccttcc cacactcttc atgtgttaac cactgccttc ctggaccttg     1800
gagccacggt gactgtatta catggttgta tagaaaactg attttagagt tctgatcgtt     1860
caagagaatg attaaatata catttcctaa aaaaatgt      1898

```

<210> 5

<211> 336

<212> PRT

<213> Homo sapiens

<400> 5

```

Val Ser His Gln Gly Arg Gln Arg Gln Leu His Ser Ala Ser Thr Gln
1          5          10          15

Ile Arg Trp Glu Pro Ser Pro Ala Met Ala Ser Leu Gly Gln Ile Leu
20          25          30

Phe Trp Ser Ile Ile Ser Ile Ile Ile Ile Leu Ala Gly Ala Ile Ala
35          40          45

Leu Ile Ile Gly Phe Gly Ile Ser Glu Val Ser Val Trp Leu Ser Ala
50          55          60

Met Lys Gly Leu Val Val Glu Val Pro Arg Leu Pro Leu Ala Leu Ile
65          70          75          80

Phe Ala Ser Cys Thr Ala Gly Arg His Ser Ile Thr Val Thr Thr Val
85          90          95

```

Ala Ser Ala Gly Asn Ile Gly Glu Asp Gly Ile Leu Ser Cys Thr Phe
100 105 110

Glu Pro Asp Ile Lys Leu Ser Asp Ile Val Ile Gln Trp Leu Lys Glu
115 120 125

Gly Val Leu Gly Leu Val His Glu Phe Lys Glu Gly Lys Asp Glu Leu
130 135 140

Ser Glu Gln Asp Glu Met Phe Arg Gly Arg Thr Ala Val Phe Ala Asp
145 150 155 160

Gln Val Ile Val Gly Asn Ala Ser Leu Arg Leu Lys Asn Val Gln Leu
165 170 175

Thr Asp Ala Gly Thr Tyr Lys Cys Tyr Ile Ile Thr Ser Lys Gly Lys
180 185 190

Gly Asn Ala Asn Leu Glu Tyr Lys Thr Gly Ala Phe Ser Met Pro Glu
195 200 205

Val Asn Val Asp Tyr Asn Ala Ser Ser Glu Thr Leu Arg Cys Glu Ala
210 215 220

Pro Arg Trp Phe Pro Gln Pro Thr Val Val Trp Ala Ser Gln Val Asp
225 230 235 240

Gln Gly Ala Asn Phe Ser Glu Val Ser Asn Thr Ser Phe Glu Leu Asn
245 250 255

Ser Glu Asn Val Thr Met Lys Val Val Ser Val Leu Tyr Asn Val Thr
260 265 270

Ile Asn Asn Thr Tyr Ser Cys Met Ile Glu Asn Asp Ile Ala Lys Ala
275 280 285

Thr Gly Asp Ile Lys Val Thr Glu Ser Glu Ile Lys Arg Arg Ser His
290 295 300

Leu Gln Leu Leu Asn Ser Lys Ala Ser Leu Cys Val Ser Ser Phe Phe
305 310 315 320

Ala Ile Ser Trp Ala Leu Leu Pro Leu Ser Pro Tyr Leu Met Leu Lys
325 330 335

<210> 6
<211> 306
<212> PRT
<213> Homo sapiens

<400> 6
Val Ser His Gln Gly Arg Gln Arg Gln Leu His Ser Ala Ser Thr Gln
1 5 10 15

Ile Arg Trp Glu Pro Ser Pro Ala Met Ala Ser Leu Gly Gln Ile Leu
20 25 30

Phe Trp Ser Ile Ile Ser Ile Ile Ile Ile Leu Ala Gly Ala Ile Ala
35 40 45

Leu Ile Ile Gly Phe Gly Ile Ser Gly Arg His Ser Ile Thr Val Thr
 50 . 55 60
 Thr Val Ala Ser Ala Gly Asn Ile Gly Glu Asp Gly Ile Leu Ser Cys
 65 70 75 80
 Thr Phe Glu Pro Asp Ile Lys Leu Ser Asp Ile Val Ile Gln Trp Leu
 85 90 95
 Lys Glu Gly Val Leu Gly Leu Val His Glu Phe Lys Glu Gly Lys Asp
 100 105 110
 Glu Leu Ser Glu Gln Asp Glu Met Phe Arg Gly Arg Thr Ala Val Phe
 115 120 125
 Ala Asp Gln Val Ile Val Gly Asn Ala Ser Leu Arg Leu Lys Asn Val
 130 135 140
 Gln Leu Thr Asp Ala Gly Thr Tyr Lys Cys Tyr Ile Ile Thr Ser Lys
 145 150 155 160
 Gly Lys Gly Asn Ala Asn Leu Glu Tyr Lys Thr Gly Ala Phe Ser Met
 165 170 175
 Pro Glu Val Asn Val Asp Tyr Asn Ala Ser Ser Glu Thr Leu Arg Cys
 180 185 190
 Glu Ala Pro Arg Trp Phe Pro Gln Pro Thr Val Val Trp Ala Ser Gln
 195 200 205
 Val Asp Gln Gly Ala Asn Phe Ser Glu Val Ser Asn Thr Ser Phe Glu
 210 215 220
 Leu Asn Ser Glu Asn Val Thr Met Lys Val Val Ser Val Leu Tyr Asn
 225 230 235 240
 Val Thr Ile Asn Asn Thr Tyr Ser Cys Met Ile Glu Asn Asp Ile Ala
 245 250 255
 Lys Ala Thr Gly Asp Ile Lys Val Thr Glu Ser Glu Ile Lys Arg Arg
 260 265 270
 Ser His Leu Gln Leu Leu Asn Ser Lys Ala Ser Leu Cys Val Ser Ser
 275 280 285
 Phe Phe Ala Ile Ser Trp Ala Leu Leu Pro Leu Ser Pro Tyr Leu Met
 290 295 300
 Leu Lys
 305

WHAT IS CLAIMED IS:

1. A process for identifying an agent that modulates the activity of a cancer-related gene comprising:

5 (a) contacting a compound with a cell containing a gene that corresponds to a polynucleotide having a sequence selected from the group consisting of SEQ ID NO: 1, 2, 3 and 4 and under conditions promoting the expression of said gene; and

10 (b) detecting a difference in expression of said gene relative to when said compound is not present

thereby identifying an agent that modulates the activity of a cancer-related gene.

2 The process of claim 1 wherein said gene has a sequence selected
15 from the group consisting of SEQ ID NO: 1, 2, 3 and 4.

3. The process of claim 1 wherein the cell is a cancer cell and the difference in expression is a decrease in expression.

20 4. The process of claim 3 wherein said cancer cell is a breast or endometrium cancer cell.

5. A process for identifying an anti-neoplastic agent comprising
25 contacting a cell exhibiting neoplastic activity with a compound first identified as a cancer related gene modulator using the process of claim 1 and detecting a decrease in said neoplastic activity after said contacting compared to when said contacting does not occur.

30 6. The process of claim 5 wherein said neoplastic activity is accelerated cellular replication.

7. The process of claim 5 wherein said decrease in neoplastic activity results from the death of the cell.

5 8. A process for identifying an anti-neoplastic agent comprising administering to an animal exhibiting a cancer condition an effective amount of an agent first identified according to the process of claim 1 and detecting a decrease in said cancerous condition.

10 9. A process for determining the cancerous status of a cell, comprising determining an increase in the level of expression in said cell of a gene that corresponds to a polynucleotide having a sequence selected from the group consisting of SEQ ID NO: 1, 2, 3 and 4 wherein an elevated expression relative to a known non-cancerous cell indicates a cancerous state or potentially cancerous state.

15

10. The process of claim 9 wherein said elevated expression is due to an increased copy number.

20 11. An isolated polypeptide comprising an amino acid sequence homologous to an amino acid sequence selected from the group consisting of SEQ ID NO: 5 and 6 wherein any difference between said amino acid sequence and the sequence of SEQ ID NO: 5 and 6 is due solely to conservative amino acid substitutions and wherein said isolated polypeptide comprises at least one immunogenic fragment.

25

12. An isolated polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 5 and 6.

30 13. An antibody that reacts with a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 5 and 6.

14. The antibody of claim 13 wherein said antibody is a recombinant antibody.

5 15. The antibody of claim 13 wherein said antibody is a synthetic antibody.

16. The antibody of claim 13 wherein said antibody is a humanized antibody.

10 17. An immunoconjugate comprising the antibody of claim 13 and a cytotoxic agent.

15 18. The antibody of claim 17 wherein said cytotoxic agent is a member selected from the group consisting of a calicheamicin, a maytansinoid, an adozelesin, a cytotoxic protein, a taxol, a taxotere, a taxoid and DC1.

20 19. The immunoconjugate of claim 18 wherein said calicheamicin is calicheamicin γ_1^I , N-acetyl gamma calicheamicin dimethyl hydrazide or calicheamicin θ_1^I .

20. The immunoconjugate of claim 18 wherein said maytansinoid is DM1.

25 21. The immunoconjugate of claim 18 wherein said cytotoxic protein is ricin, abrin, gelonin, pseudomonas exotoxin or diphtheria toxin.

22. The immunoconjugate of claim 18 wherein said taxol is paclitaxel.

30 23. The immunoconjugate of claim 18 wherein said taxotere is docetaxel.

24. A process for treating cancer comprising contacting a cancerous cell *in vivo* with an agent having activity against an expression product encoded by a gene sequence selected from the group consisting of SEQ ID NO: 1, 2, 3 and 4.

5

25. The process of claim 24 wherein said agent is an antibody of claim 13.

26. The process of claim 24 wherein said agent is an immunoconjugate
10 of claim 17.

27. An immunogenic composition comprising a polypeptide of claim 11.

28. An immunogenic composition comprising a polypeptide of claim 12.
15

29. The process of claim 24 wherein said cancer is breast or endometrium cancer.

30. A process for treating cancer in an animal afflicted therewith
20 comprising administering to said animal an amount of an immunogenic composition of claim 27 sufficient to elicit the production of cytotoxic T lymphocytes specific for the polypeptide of claim 11.

31. A process for treating cancer in an animal afflicted therewith
25 comprising administering to said animal an amount of an immunogenic composition of claim 28 sufficient to elicit the production of cytotoxic T lymphocytes specific for the polypeptide of claim 12.

32. A process for treating a cancerous condition in an animal afflicted
30 therewith comprising administering to said animal a therapeutically effective amount of an agent first identified as having anti-neoplastic activity using the process of claim 8.

33. A process for protecting an animal against cancer comprising administering to an animal at risk of developing cancer a therapeutically effective amount of an agent first identified as having anti-neoplastic activity using the process of claim 8.

5

34. The process of claim 30 wherein said animal is a human being.

35. The process of claim 30 wherein said cancer is breast or endometrium cancer.

10

36. A method for producing test data with respect to the gene modulating activity of a compound comprising:

15 (a) contacting a compound with a cell containing a polynucleotide comprising a nucleotide sequence corresponding to a gene whose expression is increased in a cancerous cell over that in a non-cancerous cell and under conditions wherein said polynucleotide is being expressed,

(b) determining a change in expression of polynucleotides as a result of said contacting, and

20 (c) producing test data with respect to the gene modulating activity of said compound based on a decrease in the expression of the determined gene whose expression is otherwise increased in a cancerous cell over that in a non-cancerous cell indicating gene modulating activity.