

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
5 August 2010 (05.08.2010)

PCT

(10) International Publication Number
WO 2010/088387 A1

(51) International Patent Classification:

C12N 9/42 (2006.01) C12P 19/14 (2006.01)
C12N 15/82 (2006.01)

(21) International Application Number:

PCT/US2010/022405

(22) International Filing Date:

28 January 2010 (28.01.2010)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/147,923 28 January 2009 (28.01.2009) US

(71) Applicants (for all designated States except US):

NOVOZYMES, INC. [US/US]; 1445 Drew Avenue,
Davis, California 95618 (US). **NOVOZYMES A/S** [DK/
DK]; Krogshoejvej 36, DK-2880 Bagsvaerd (DK).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **SCHNORR, Kirk**
[US/DK]; Sollerodgardsvej 38, DK-2840 Holte (DK).
REY, Michael [US/US]; 605 Robin Place, Davis, Cali-
fornia 95616 (US).

(74) Agent: **STARNES, Robert**; Novozymes, Inc., 1445
Drew Avenue, Davis, California 95618 (US).

(81) Designated States (unless otherwise indicated, for every

kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ,

CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO,
DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP,
KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD,
ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI,
NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD,
SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR,
TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every

kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ,
TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,
MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, SM,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— as to applicant's entitlement to apply for and be granted
a patent (Rule 4.17(ii))

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments (Rule 48.2(h))
- with sequence listing part of description (Rule 5.2(a))



WO 2010/088387 A1

(54) Title: POLYPEPTIDES HAVING BETA-GLUCOSIDASE ACTIVITY AND POLYNUCLEOTIDES ENCODING SAME

(57) Abstract: The present invention relates to isolated polypeptides having beta-glucosidase activity and isolated polynucleotides encoding the polypeptides. The invention also relates to nucleic acid constructs, vectors, and host cells comprising the polynucleotides as well as methods of producing and using the polypeptides.

**POLYPEPTIDES HAVING BETA-GLUCOSIDASE ACTIVITY
AND POLYNUCLEOTIDES ENCODING SAME**

5 **Reference to a Sequence Listing**

This application contains a Sequence Listing filed electronically by EFS, which is incorporated herein by reference.

Reference to a Deposit of Biological Material

10 This application contains a reference to a deposit of biological material, which deposit is incorporated herein by reference.

Background of the Invention

15 **Field of the Invention**

The present invention relates to isolated polypeptides having beta-glucosidase activity and isolated polynucleotides encoding the polypeptides. The invention also relates to nucleic acid constructs, vectors, and host cells comprising the polynucleotides as well as methods of producing and using the polypeptides.

20

Description of the Related Art

25 Cellulose is a polymer of glucose linked by beta-1,4-bonds. Many microorganisms produce enzymes that hydrolyze beta-linked glucans. These enzymes include endoglucanases, cellobiohydrolases, and beta-glucosidases. Endoglucanases digest the cellulose polymer at random locations, opening it to attack by cellobiohydrolases. Cellobiohydrolases sequentially release molecules of cellobiose from the ends of the cellulose polymer. Cellobiose is a water-soluble beta-1,4-linked dimer of glucose. Beta-glucosidases hydrolyze cellobiose to glucose.

30 The conversion of lignocellulosic feedstocks into ethanol has the advantages of the ready availability of large amounts of feedstock, the desirability of avoiding burning or land filling the materials, and the cleanliness of the ethanol fuel. Wood, agricultural residues, herbaceous crops, and municipal solid wastes have been considered as feedstocks for ethanol production. These materials primarily consist of cellulose, hemicellulose, and lignin. Once the cellulose is converted to glucose, the glucose is easily fermented by yeast into
35 ethanol. Since glucose is readily fermented to ethanol by a variety of yeasts while cellobiose is not, any cellobiose remaining at the end of the hydrolysis represents a loss of yield of ethanol. More importantly, cellobiose is a potent inhibitor of endoglucanases and

cellobiohydrolases. The accumulation of cellobiose during hydrolysis is undesirable for ethanol production.

Cellobiose accumulation has been a major problem in enzymatic hydrolysis because cellulase-producing microorganisms may produce little beta-glucosidase. The low amount of beta-glucosidase results in a shortage of capacity to hydrolyze the cellobiose to glucose. Several approaches have been used to increase the amount of beta-glucosidase in cellulose conversion.

One approach is to produce beta-glucosidase using microorganisms that produce little cellulase, and add the beta-glucosidase exogenously to endoglucanase and cellobiohydrolase to enhance the hydrolysis.

A second approach is to carry out cellulose hydrolysis simultaneously with fermentation. This process is known as simultaneous saccharification and fermentation (SSF). In an SSF system, fermentation of the glucose removes it from solution.

A third approach to overcome the shortage of beta-glucosidase is to overexpress the beta-glucosidase in a host, thereby increasing the yield of beta-glucosidase.

It would be an advantage in the art to provide new beta-glucosidases for degrading cellulosic materials.

The present invention provides polypeptides having beta-glucosidase activity and polynucleotides encoding the polypeptides.

Summary of the Invention

The present invention relates to isolated polypeptides having beta-glucosidase activity selected from the group consisting of:

(a) a polypeptide comprising an amino acid sequence having at least 60% sequence identity to the mature polypeptide of SEQ ID NO: 2;

(b) a polypeptide encoded by a polynucleotide that hybridizes under at least medium stringency conditions with (i) the mature polypeptide coding sequence of SEQ ID NO: 1, (ii) the genomic DNA sequence comprising the mature polypeptide coding sequence of SEQ ID NO: 1, or (iii) a full-length complementary strand of (i) or (ii);

(c) a polypeptide encoded by a polynucleotide comprising a nucleotide sequence having at least 60% sequence identity to the mature polypeptide coding sequence of SEQ ID NO: 1; and

(d) a variant comprising a substitution, deletion, and/or insertion of one or more (several) amino acids of the mature polypeptide of SEQ ID NO: 2.

The present invention also relates to isolated polynucleotides encoding polypeptides having beta-glucosidase activity, selected from the group consisting of:

(a) a polynucleotide encoding a polypeptide comprising an amino acid sequence having at least 60% sequence identity to the mature polypeptide of SEQ ID NO: 2;

(b) a polynucleotide that hybridizes under at least medium stringency conditions with (i) the mature polypeptide coding sequence of SEQ ID NO: 1, (ii) the genomic DNA sequence comprising the mature polypeptide coding sequence of SEQ ID NO: 1, or (iii) a full-length complementary strand of (i) or (ii);

(c) a polynucleotide comprising a nucleotide sequence having at least 60% sequence identity to the mature polypeptide coding sequence of SEQ ID NO: 1; and

(d) a polynucleotide encoding a variant comprising a substitution, deletion, and/or insertion of one or more (several) amino acids of the mature polypeptide of SEQ ID NO: 2.

The present invention also relates to nucleic acid constructs, recombinant expression vectors, and recombinant host cells comprising the polynucleotides, and to methods of producing the polypeptides having beta-glucosidase activity.

The present invention also relates to methods of inhibiting the expression of a polypeptide having beta-glucosidase activity in a cell, comprising administering to the cell or expressing in the cell a double-stranded RNA (dsRNA) molecule, wherein the dsRNA comprises a subsequence of a polynucleotide of the present invention. The present invention also relates to such a double-stranded inhibitory RNA (dsRNA) molecule, wherein optionally the dsRNA is a siRNA or a miRNA molecule.

The present invention also relates to methods for degrading or converting a cellulosic material, comprising: treating the cellulosic material with an enzyme composition in the presence of a polypeptide having beta-glucosidase activity.

The present invention also relates to methods of producing a fermentation product, comprising: (a) saccharifying a cellulosic material with an enzyme composition in the presence of a polypeptide having beta-glucosidase activity; (b) fermenting the saccharified cellulosic material with one or more (several) fermenting microorganisms to produce the fermentation product; and (c) recovering the fermentation product from the fermentation.

The present invention also relates to methods of fermenting a cellulosic material, comprising: fermenting the cellulosic material with one or more (several) fermenting microorganisms, wherein the cellulosic material is saccharified with an enzyme composition in the presence of a polypeptide having beta-glucosidase activity.

The present invention also relates to plants comprising an isolated polynucleotide encoding a polypeptide having beta-glucosidase activity.

The present invention also relates to methods of producing a polypeptide having beta-glucosidase activity, comprising: (a) cultivating a transgenic plant or a plant cell comprising a polynucleotide encoding the polypeptide having beta-glucosidase activity under conditions conducive for production of the polypeptide; and (b) recovering the polypeptide.

The present invention also relates to an isolated polynucleotide encoding a signal peptide comprising or consisting of amino acids 1 to 19 of SEQ ID NO: 2; nucleic acid constructs, expression vectors, and recombinant host cells comprising the polynucleotide; and methods of producing a protein.

5

Brief Description of the Figures

Figures 1A and 1B show the cDNA sequence and the deduced amino acid sequence of a *Trichophaea saccata* CBS 804.70 beta-glucosidase gene (SEQ ID NOs: 1 and 2, respectively).

10

Figure 2 shows a restriction map of pAHYG-33.

Definitions

Beta-glucosidase activity: The term "beta-glucosidase" is defined herein as a beta-D-glucoside glucohydrolase (E.C. 3.2.1.21), which catalyzes the hydrolysis of terminal non-reducing beta-D-glucose residues with the release of beta-D-glucose. For purposes of the present invention, beta-glucosidase activity is determined according to the basic procedure described by Venturi *et al.*, 2002, *J. Basic Microbiol.* 42: 55-66, except different conditions were employed as described herein. One unit of beta-glucosidase activity is defined as 1.0 μmole of p-nitrophenol produced per minute at 50°C, pH 5 from 1 mM p-nitrophenyl-beta-D-glucopyranoside as substrate in 50 mM sodium acetate, 0.01% TWEEN® 20.

15

20

The polypeptides of the present invention have preferably at least 20%, more preferably at least 40%, more preferably at least 50%, more preferably at least 60%, more preferably at least 70%, more preferably at least 80%, even more preferably at least 90%, most preferably at least 95%, and even most preferably at least 100% of the beta-glucosidase activity of the mature polypeptide of SEQ ID NO: 2.

25

Family 3 or Family GH3 or CEL3: The term "Family 3" or "Family GH3" or "CEL3" is defined herein as a polypeptide falling into the glycoside hydrolase Family 3 according to Henrissat B., 1991, A classification of glycosyl hydrolases based on amino-acid sequence similarities, *Biochem. J.* 280: 309-316, and Henrissat and Bairoch, 1996, Updating the sequence-based classification of glycosyl hydrolases, *Biochem. J.* 316: 695-696.

30

Cellulolytic activity: The term "cellulolytic activity" is defined herein as a biological activity that hydrolyzes a cellulosic material. The two basic approaches for measuring cellulolytic activity include: (1) measuring the total cellulolytic activity, and (2) measuring the individual cellulolytic activities (endoglucanases, cellobiohydrolases, and beta-glucosidases) as reviewed in Zhang *et al.*, Outlook for cellulase improvement: Screening and selection

35

strategies, 2006, *Biotechnology Advances* 24: 452-481. Total cellulolytic activity is usually measured using insoluble substrates, including Whatman №1 filter paper, microcrystalline cellulose, bacterial cellulose, algal cellulose, cotton, pretreated lignocellulose, etc. The most common total cellulolytic activity assay is the filter paper assay using Whatman №1 filter paper as the substrate. The assay was established by the International Union of Pure and Applied Chemistry (IUPAC) (Ghose, 1987, Measurement of cellulase activities, *Pure Appl. Chem.* 59: 257-68).

For purposes of the present invention, cellulolytic activity is determined by measuring the increase in hydrolysis of a cellulosic material by cellulolytic enzyme(s) under the following conditions: 1-20 mg of cellulolytic protein/g of cellulose in PCS for 3-7 days at 50-65°C compared to a control hydrolysis without addition of cellulolytic protein. Typical conditions are 1 ml reactions, washed or unwashed PCS, 5% insoluble solids, 50 mM sodium acetate pH 5, 1 mM MnSO₄, 50-65°C, 72 hours, sugar analysis by AMINEX® HPX-87H column (Bio-Rad Laboratories, Inc., Hercules, CA, USA).

Endoglucanase: The term “endoglucanase” is defined herein as an endo-1,4-(1,3;1,4)-beta-D-glucan 4-glucanohydrolase (E.C. 3.2.1.4), which catalyses endohydrolysis of 1,4-beta-D-glycosidic linkages in cellulose, cellulose derivatives (such as carboxymethyl cellulose and hydroxyethyl cellulose), lichenin, beta-1,4 bonds in mixed beta-1,3 glucans such as cereal beta-D-glucans or xyloglucans, and other plant material containing cellulosic components. Endoglucanase activity can be determined based on a reduction in substrate viscosity or increase in reducing ends determined by a reducing sugar assay (Zhang *et al.*, 2006, *Biotechnology Advances* 24: 452-481). For purposes of the present invention, endoglucanase activity is determined using carboxymethyl cellulose (CMC) hydrolysis according to the procedure of Ghose, 1987, *Pure and Appl. Chem.* 59: 257-268.

Cellobiohydrolase: The term “cellobiohydrolase” is defined herein as a 1,4-beta-D-glucan cellobiohydrolase (E.C. 3.2.1.91), which catalyzes the hydrolysis of 1,4-beta-D-glucosidic linkages in cellulose, cellooligosaccharides, or any beta-1,4-linked glucose containing polymer, releasing cellobiose from the reducing or non-reducing ends of the chain (Teeri, 1997, Crystalline cellulose degradation: New insight into the function of cellobiohydrolases, *Trends in Biotechnology* 15: 160-167; Teeri *et al.*, 1998, *Trichoderma reesei* cellobiohydrolases: why so efficient on crystalline cellulose?, *Biochem. Soc. Trans.* 26: 173-178). For purposes of the present invention, cellobiohydrolase activity is determined using a fluorescent disaccharide derivative 4-methylumbelliferyl-β-D-lactoside according to the procedures described by van Tilbeurgh *et al.*, 1982, *FEBS Letters* 149: 152-156 and van Tilbeurgh and Claeysens, 1985, *FEBS Letters* 187: 283-288.

Cellulolytic enhancing activity: The term “cellulolytic enhancing activity” is defined herein as a biological activity that enhances the hydrolysis of a cellulosic material by

polypeptides having cellulolytic activity. For purposes of the present invention, cellulolytic enhancing activity is determined by measuring the increase in reducing sugars or the increase of the total of cellobiose and glucose from the hydrolysis of a cellulosic material by cellulolytic protein under the following conditions: 1-50 mg of total protein/g of cellulose in PCS, wherein total protein is comprised of 50-99.5% w/w cellulolytic protein and 0.5-50% w/w protein of cellulolytic enhancing activity for 1-7 day at 50-65°C compared to a control hydrolysis with equal total protein loading without cellulolytic enhancing activity (1-50 mg of cellulolytic protein/g of cellulose in PCS). In a preferred aspect, a mixture of CELLUCLAST® 1.5L (Novozymes A/S, Bagsværd, Denmark) in the presence of 3% of total protein weight *Aspergillus oryzae* beta-glucosidase (recombinantly produced in *Aspergillus oryzae* according to WO 02/095014) or 3% of total protein weight *Aspergillus fumigatus* beta-glucosidase (recombinantly produced in *Aspergillus oryzae* as described in WO 2002/095014) of cellulase protein loading is used as the source of the cellulolytic activity.

The polypeptides having cellulolytic enhancing activity have at least 20%, preferably at least 40%, more preferably at least 50%, more preferably at least 60%, more preferably at least 70%, more preferably at least 80%, even more preferably at least 90%, most preferably at least 95%, and even most preferably at least 100% of the cellulolytic enhancing activity of the mature polypeptide of a GH61 polypeptide.

The polypeptides having cellulolytic enhancing activity enhance the hydrolysis of a cellulosic material catalyzed by proteins having cellulolytic activity by reducing the amount of cellulolytic enzyme required to reach the same degree of hydrolysis preferably at least 1.01-fold, more preferably at least 1.05-fold, more preferably at least 1.10-fold, more preferably at least 1.25-fold, more preferably at least 1.5-fold, more preferably at least 2-fold, more preferably at least 3-fold, more preferably at least 4-fold, more preferably at least 5-fold, even more preferably at least 10-fold, and most preferably at least 20-fold.

Family 61 glycoside hydrolase: The term "Family 61 glycoside hydrolase" or "Family GH61" is defined herein as a polypeptide falling into the glycoside hydrolase Family 61 according to Henrissat B., 1991, A classification of glycosyl hydrolases based on amino-acid sequence similarities, *Biochem. J.* 280: 309-316, and Henrissat B., and Bairoch A., 1996, Updating the sequence-based classification of glycosyl hydrolases, *Biochem. J.* 316: 695-696. Presently, Henrissat lists the GH61 Family as unclassified indicating that properties such as mechanism, catalytic nucleophile/base, and catalytic proton donors are not known for polypeptides belonging to this family.

Xylan degrading activity: The terms "xylan degrading activity" or "xylanolytic activity" are defined herein as a biological activity that hydrolyzes xylan-containing material. The two basic approaches for measuring xylanolytic activity include: (1) measuring the total xylanolytic activity, and (2) measuring the individual xylanolytic activities (endoxylanases,

beta-xylosidases, arabinofuranosidases, alpha-glucuronidases, acetylxylan esterases, feruloyl esterases, and alpha-glucuronoyl esterases). Recent progress in assays of xylanolytic enzymes was summarized in several publications including Biely and Puchard, Recent progress in the assays of xylanolytic enzymes, 2006, *Journal of the Science of Food and Agriculture* 86(11): 1636-1647; Spanikova and Biely, 2006, Glucuronoyl esterase - Novel carbohydrate esterase produced by *Schizophyllum commune*, *FEBS Letters* 580(19): 4597-4601; Herrmann, Vrsanska, Jurickova, Hirsch, Biely, and Kubicek, 1997, The beta-D-xylosidase of *Trichoderma reesei* is a multifunctional beta-D-xylan xylohydrolase, *Biochemical Journal* 321: 375-381.

Total xylan degrading activity can be measured by determining the reducing sugars formed from various types of xylan, including oat spelt, beechwood, and larchwood xylans, or by photometric determination of dyed xylan fragments released from various covalently dyed xylans. The most common total xylanolytic activity assay is based on production of reducing sugars from polymeric 4-O-methyl glucuronoxylan as described in Bailey, Biely, Poutanen, 1992, Interlaboratory testing of methods for assay of xylanase activity, *Journal of Biotechnology* 23(3): 257-270.

For purposes of the present invention, xylan degrading activity is determined by measuring the increase in hydrolysis of birchwood xylan (Sigma Chemical Co., Inc., St. Louis, MO, USA) by xylan-degrading enzyme(s) under the following typical conditions: 1 ml reactions, 5 mg/ml substrate (total solids), 5 mg of xylanolytic protein/g of substrate, 50 mM sodium acetate pH 5, 50°C, 24 hours, sugar analysis using p-hydroxybenzoic acid hydrazide (PHBAH) assay as described by Lever, 1972, A new reaction for colorimetric determination of carbohydrates, *Anal. Biochem* 47: 273-279.

Xylanase activity: The term "xylanase activity" is defined herein as a 1,4-beta-D-xylan-xylohydrolase activity (E.C. 3.2.1.8) that catalyzes the endo-hydrolysis of 1,4-beta-D-xylosidic linkages in xylans. For purposes of the present invention, xylanase activity is determined using birchwood xylan as substrate. One unit of xylanase activity is defined as 1.0 μ mole of reducing sugar (measured in glucose equivalents as described by Lever, 1972, A new reaction for colorimetric determination of carbohydrates, *Anal. Biochem* 47: 273-279) produced per minute during the initial period of hydrolysis at 50°C, pH 5 from 2 g of birchwood xylan per liter as substrate in 50 mM sodium acetate containing 0.01% TWEEN® 20.

Beta-xylosidase activity: The term "beta-xylosidase activity" is defined herein as a beta-D-xyloside xylohydrolase (E.C. 3.2.1.37) that catalyzes the exo-hydrolysis of short beta (1→4)-xylooligosaccharides, to remove successive D-xylose residues from the non-reducing termini. For purposes of the present invention, one unit of beta-xylosidase activity is defined as 1.0 μ mole of p-nitrophenol produced per minute at 40°C, pH 5 from 1 mM p-nitrophenyl-

beta-D-xyloside as substrate in 100 mM sodium citrate containing 0.01% TWEEN® 20.

Acetylxylan esterase activity: The term “acetylxylan esterase activity” is defined herein as a carboxylesterase activity (EC 3.1.1.72) that catalyses the hydrolysis of acetyl groups from polymeric xylan, acetylated xylose, acetylated glucose, alpha-naphthyl acetate, and *p*-nitrophenyl acetate. For purposes of the present invention, acetylxylan esterase activity is determined using 0.5 mM *p*-nitrophenylacetate as substrate in 50 mM sodium acetate pH 5.0 containing 0.01% TWEEN™ 20. One unit of acetylxylan esterase activity is defined as the amount of enzyme capable of releasing 1 μmole of *p*-nitrophenolate anion per minute at pH 5, 25°C.

Feruloyl esterase activity: The term “feruloyl esterase activity” is defined herein as a 4-hydroxy-3-methoxycinnamoyl-sugar hydrolase activity (EC 3.1.1.73) that catalyzes the hydrolysis of the 4-hydroxy-3-methoxycinnamoyl (feruloyl) group from an esterified sugar, which is usually arabinose in "natural" substrates, to produce ferulate (4-hydroxy-3-methoxycinnamate). Feruloyl esterase is also known as ferulic acid esterase, hydroxycinnamoyl esterase, FAE-III, cinnamoyl ester hydrolase, FAEA, cinnAE, FAE-I, or FAE-II. For purposes of the present invention, feruloyl esterase activity is determined using 0.5 mM *p*-nitrophenylferulate as substrate in 50 mM sodium acetate pH 5.0. One unit of feruloyl esterase activity equals the amount of enzyme capable of releasing 1 μmole of *p*-nitrophenolate anion per minute at pH 5, 25°C.

Alpha-glucuronidase activity: The term “alpha-glucuronidase activity” is defined herein as an alpha-D-glucosiduronate glucuronohydrolase activity (EC 3.2.1.139) that catalyzes the hydrolysis of an alpha-D-glucuronoside to D-glucuronate and an alcohol. For purposes of the present invention, alpha-glucuronidase activity is determined according to de Vries, 1998, *J. Bacteriol.* 180: 243-249. One unit of alpha-glucuronidase activity equals the amount of enzyme capable of releasing 1 μmole of glucuronic or 4-O-methylglucuronic acid per minute at pH 5, 40°C.

Alpha-L-arabinofuranosidase activity: The term “alpha-L-arabinofuranosidase activity” is defined herein as an alpha-L-arabinofuranoside arabinofuranohydrolase activity (EC 3.2.1.55) that catalyzes the hydrolysis of terminal non-reducing alpha-L-arabinofuranoside residues in alpha-L-arabinosides. The enzyme activity acts on alpha-L-arabinofuranosides, alpha-L-arabinans containing (1,3)- and/or (1,5)-linkages, arabinoxylans, and arabinogalactans. Alpha-L-arabinofuranosidase is also known as arabinosidase, alpha-arabinosidase, alpha-L-arabinosidase, alpha-arabinofuranosidase, polysaccharide alpha-L-arabinofuranosidase, alpha-L-arabinofuranoside hydrolase, L-arabinosidase, or alpha-L-arabinanase. For purposes of the present invention, alpha-L-arabinofuranosidase activity is determined using 5 mg of medium viscosity wheat arabinoxylan (Megazyme International Ireland, Ltd., Bray, Co. Wicklow, Ireland) per ml of

100 mM sodium acetate pH 5 in a total volume of 200 µl for 30 minutes at 40°C followed by arabinose analysis by AMINEX® HPX-87H column chromatography (Bio-Rad Laboratories, Inc., Hercules, CA, USA).

Cellulosic material: The cellulosic material can be any material containing cellulose.

5 The predominant polysaccharide in the primary cell wall of biomass is cellulose, the second most abundant is hemicellulose, and the third is pectin. The secondary cell wall, produced after the cell has stopped growing, also contains polysaccharides and is strengthened by polymeric lignin covalently cross-linked to hemicellulose. Cellulose is a homopolymer of anhydrocellobiose and thus a linear beta-(1-4)-D-glucan, while hemicelluloses include a
10 variety of compounds, such as xylans, xyloglucans, arabinoxylans, and mannans in complex branched structures with a spectrum of substituents. Although generally polymorphous, cellulose is found in plant tissue primarily as an insoluble crystalline matrix of parallel glucan chains. Hemicelluloses usually hydrogen bond to cellulose, as well as to other hemicelluloses, which help stabilize the cell wall matrix.

15 Cellulose is generally found, for example, in the stems, leaves, hulls, husks, and cobs of plants or leaves, branches, and wood of trees. The cellulosic material can be, but is not limited to, herbaceous material, agricultural residue, forestry residue, municipal solid waste, waste paper, and pulp and paper mill residue (see, for example, Wiselogel *et al.*, 1995, in Handbook on Bioethanol (Charles E. Wyman, editor), pp.105-118, Taylor & Francis, Washington D.C.; Wyman, 1994, *Bioresource Technology* 50: 3-16; Lynd, 1990, *Applied Biochemistry and Biotechnology* 24/25: 695-719; Mosier *et al.*, 1999, Recent Progress in Bioconversion of Lignocellulosics, in *Advances in Biochemical Engineering/Biotechnology*, T. Scheper, managing editor, Volume 65, pp.23-40, Springer-Verlag, New York). It is
20 understood herein that the cellulose may be in the form of lignocellulose, a plant cell wall material containing lignin, cellulose, and hemicellulose in a mixed matrix. In a preferred aspect, the cellulosic material is lignocellulose.

In one aspect, the cellulosic material is herbaceous material. In another aspect, the cellulosic material is agricultural residue. In another aspect, the cellulosic material is forestry residue. In another aspect, the cellulosic material is municipal solid waste. In another
30 aspect, the cellulosic material is waste paper. In another aspect, the cellulosic material is pulp and paper mill residue.

In another aspect, the cellulosic material is corn stover. In another aspect, the cellulosic material is corn fiber. In another aspect, the cellulosic material is corn cob. In another aspect, the cellulosic material is orange peel. In another aspect, the cellulosic
35 material is rice straw. In another aspect, the cellulosic material is wheat straw. In another aspect, the cellulosic material is switch grass. In another aspect, the cellulosic material is miscanthus. In another aspect, the cellulosic material is bagasse.

In another aspect, the cellulosic material is microcrystalline cellulose. In another aspect, the cellulosic material is bacterial cellulose. In another aspect, the cellulosic material is algal cellulose. In another aspect, the cellulosic material is cotton linter. In another aspect, the cellulosic material is amorphous phosphoric-acid treated cellulose. In another aspect, the cellulosic material is filter paper.

The cellulosic material may be used as is or may be subjected to pretreatment, using conventional methods known in the art, as described herein. In a preferred aspect, the cellulosic material is pretreated.

Pretreated corn stover: The term "PCS" or "Pretreated Corn Stover" is defined herein as a cellulosic material derived from corn stover by treatment with heat and dilute sulfuric acid.

Isolated polypeptide: The term "isolated polypeptide" as used herein refers to a polypeptide that is isolated from a source. In a preferred aspect, the polypeptide is at least 1% pure, preferably at least 5% pure, more preferably at least 10% pure, more preferably at least 20% pure, more preferably at least 40% pure, more preferably at least 60% pure, even more preferably at least 80% pure, and most preferably at least 90% pure, as determined by SDS-PAGE.

Substantially pure polypeptide: The term "substantially pure polypeptide" denotes herein a polypeptide preparation that contains at most 10%, preferably at most 8%, more preferably at most 6%, more preferably at most 5%, more preferably at most 4%, more preferably at most 3%, even more preferably at most 2%, most preferably at most 1%, and even most preferably at most 0.5% by weight of other polypeptide material with which it is natively or recombinantly associated. It is, therefore, preferred that the substantially pure polypeptide is at least 92% pure, preferably at least 94% pure, more preferably at least 95% pure, more preferably at least 96% pure, more preferably at least 97% pure, more preferably at least 98% pure, even more preferably at least 99% pure, most preferably at least 99.5% pure, and even most preferably 100% pure by weight of the total polypeptide material present in the preparation. The polypeptides of the present invention are preferably in a substantially pure form, *i.e.*, that the polypeptide preparation is essentially free of other polypeptide material with which it is natively or recombinantly associated. This can be accomplished, for example, by preparing the polypeptide by well-known recombinant methods or by classical purification methods.

Mature polypeptide: The term "mature polypeptide" is defined herein as a polypeptide in its final form following translation and any post-translational modifications, such as N-terminal processing, C-terminal truncation, glycosylation, phosphorylation, etc. In one aspect, the mature polypeptide is amino acids 20 to 856 of SEQ ID NO: 2 based on the SignalP program (Nielsen *et al.*, 1997, *Protein Engineering* 10: 1-6) that predicts amino acids

1 to 19 of SEQ ID NO: 2 are a signal peptide.

Mature polypeptide coding sequence: The term “mature polypeptide coding sequence” is defined herein as a nucleotide sequence that encodes a mature polypeptide having beta-glucosidase activity. In one aspect, the mature polypeptide coding sequence is nucleotides 58 to 2568 of SEQ ID NO: 1 based on the SignalP program (Nielsen *et al.*, 1997, *supra*) that predicts nucleotides 1 to 57 of SEQ ID NO: 1 encode a signal peptide.

Sequence Identity: The relatedness between two amino acid sequences or between two nucleotide sequences is described by the parameter “sequence identity”.

For purposes of the present invention, the degree of sequence identity between two amino acid sequences is determined using the Needleman-Wunsch algorithm (Needleman and Wunsch, 1970, *J. Mol. Biol.* 48: 443-453) as implemented in the Needle program of the EMBOSS package (EMBOSS: The European Molecular Biology Open Software Suite, Rice *et al.*, 2000, *Trends in Genetics* 16: 276-277), preferably version 3.0.0 or later. The optional parameters used are gap open penalty of 10, gap extension penalty of 0.5, and the EBLOSUM62 (EMBOSS version of BLOSUM62) substitution matrix. The output of Needle labeled “longest identity” (obtained using the –nobrief option) is used as the percent identity and is calculated as follows:

$$(\text{Identical Residues} \times 100) / (\text{Length of Alignment} - \text{Total Number of Gaps in Alignment})$$

For purposes of the present invention, the degree of sequence identity between two deoxyribonucleotide sequences is determined using the Needleman-Wunsch algorithm (Needleman and Wunsch, 1970, *supra*) as implemented in the Needle program of the EMBOSS package (EMBOSS: The European Molecular Biology Open Software Suite, Rice *et al.*, 2000, *supra*), preferably version 3.0.0 or later. The optional parameters used are gap open penalty of 10, gap extension penalty of 0.5, and the EDNAFULL (EMBOSS version of NCBI NUC4.4) substitution matrix. The output of Needle labeled “longest identity” (obtained using the –nobrief option) is used as the percent identity and is calculated as follows:

$$(\text{Identical Deoxyribonucleotides} \times 100) / (\text{Length of Alignment} - \text{Total Number of Gaps in Alignment})$$

Homologous sequence: The term “homologous sequence” is defined herein as a predicted protein having an E value (or expectancy score) of less than 0.001 in a tfasty search (Pearson, W.R., 1999, in *Bioinformatics Methods and Protocols*, S. Misener and S. A. Krawetz, ed., pp. 185-219) with the *Trichophaea saccata* beta-glucosidase of SEQ ID NO: 2 or the mature polypeptide thereof.

Polypeptide fragment: The term “polypeptide fragment” is defined herein as a polypeptide having one or more (several) amino acids deleted from the amino and/or carboxyl terminus of the mature polypeptide of SEQ ID NO: 2; or a homologous sequence thereof; wherein the fragment has beta-glucosidase activity. In one aspect, a fragment

contains at least 740 amino acid residues, more preferably at least 770 amino acid residues, and most preferably at least 800 amino acid residues of SEQ ID NO: 2, of the mature polypeptide of SEQ ID NO: 2 or a homologous sequence thereof.

Subsequence: The term "subsequence" is defined herein as a nucleotide sequence having one or more (several) nucleotides deleted from the 5' and/or 3' end of the mature polypeptide coding sequence of SEQ ID NO: 1; or a homologous sequence thereof; wherein the subsequence encodes a polypeptide fragment having beta-glucosidase activity. In one aspect, a subsequence contains at least 2220 nucleotides, more preferably at least 2310 nucleotides, and most preferably at least 2400 nucleotides of the mature polypeptide coding sequence of SEQ ID NO: 1 or a homologous sequence thereof.

Allelic variant: The term "allelic variant" denotes herein any of two or more (several) alternative forms of a gene occupying the same chromosomal locus. Allelic variation arises naturally through mutation, and may result in polymorphism within populations. Gene mutations can be silent (no change in the encoded polypeptide) or may encode polypeptides having altered amino acid sequences. An allelic variant of a polypeptide is a polypeptide encoded by an allelic variant of a gene.

Isolated polynucleotide: The term "isolated polynucleotide" as used herein refers to a polynucleotide that is isolated from a source. In a preferred aspect, the polynucleotide is preferably at least 1% pure, more preferably at least 5% pure, more preferably at least 10% pure, more preferably at least 20% pure, more preferably at least 40% pure, more preferably at least 60% pure, even more preferably at least 80% pure, and most preferably at least 90% pure, as determined by agarose electrophoresis.

Substantially pure polynucleotide: The term "substantially pure polynucleotide" as used herein refers to a polynucleotide preparation free of other extraneous or unwanted nucleotides and in a form suitable for use within genetically engineered protein production systems. Thus, a substantially pure polynucleotide contains preferable at most 10%, more preferably at most 8%, more preferably at most 6%, more preferably at most 5%, more preferably at most 4%, more preferably at most 3%, even more preferably at most 2%, most preferably at most 1%, and even most preferably at most 0.5% by weight of other polynucleotide material with which it is natively or recombinantly associated. A substantially pure polynucleotide may, however, include naturally occurring 5' and 3' untranslated regions, such as promoters and terminators. It is preferred that the substantially pure polynucleotide is preferably at least 90% pure, more preferably at least 92% pure, more preferably at least 94% pure, more preferably at least 95% pure, more preferably at least 96% pure, more preferably at least 97% pure, even more preferably at least 98% pure, most preferably at least 99% pure, and even most preferably at least 99.5% pure by weight. The polynucleotides of the present invention are preferably in a substantially pure form, *i.e.*, that

the polynucleotide preparation is essentially free of other polynucleotide material with which it is natively or recombinantly associated. The polynucleotides may be of genomic, cDNA, RNA, semisynthetic, synthetic origin, or any combinations thereof.

Coding sequence: When used herein the term "coding sequence" means a nucleotide sequence, which directly specifies the amino acid sequence of its protein product. The boundaries of the coding sequence are generally determined by an open reading frame, which usually begins with the ATG start codon or alternative start codons such as GTG and TTG and ends with a stop codon such as TAA, TAG, and TGA. The coding sequence may be a DNA, cDNA, synthetic, or recombinant nucleotide sequence.

cDNA: The term "cDNA" is defined herein as a DNA molecule that can be prepared by reverse transcription from a mature, spliced, mRNA molecule obtained from a eukaryotic cell. cDNA lacks intron sequences that may be present in the corresponding genomic DNA. The initial, primary RNA transcript is a precursor to mRNA that is processed through a series of steps before appearing as mature spliced mRNA. These steps include the removal of intron sequences by a process called splicing. cDNA derived from mRNA lacks, therefore, any intron sequences.

Nucleic acid construct: The term "nucleic acid construct" as used herein refers to a nucleic acid molecule, either single- or double-stranded isolated from a naturally occurring gene, which is modified to contain segments of nucleic acids in a manner that would not otherwise exist in nature. The nucleic acid construct can be also synthetic. The term nucleic acid construct is synonymous with the term "expression cassette" when the nucleic acid construct contains the control sequences required for expression of a coding sequence of the present invention.

Control sequences: The term "control sequences" is defined herein to include all components necessary for the expression of a polynucleotide encoding a polypeptide of the present invention. Each control sequence may be native or foreign to the nucleotide sequence encoding the polypeptide or native or foreign to each other. Such control sequences include, but are not limited to, a leader, polyadenylation sequence, propeptide sequence, promoter, signal peptide sequence, and transcription terminator. At a minimum, the control sequences include a promoter, and transcriptional and translational stop signals. The control sequences may be provided with linkers for the purpose of introducing specific restriction sites facilitating ligation of the control sequences with the coding region of the nucleotide sequence encoding a polypeptide.

Operably linked: The term "operably linked" denotes herein a configuration in which a control sequence is placed at an appropriate position relative to the coding sequence of a polynucleotide sequence such that the control sequence directs the expression of the coding sequence of a polypeptide.

Expression: The term "expression" includes any step involved in the production of a polypeptide including, but not limited to, transcription, post-transcriptional modification, translation, post-translational modification, and secretion.

Expression vector: The term "expression vector" is defined herein as a linear or circular DNA molecule that comprises a polynucleotide encoding a polypeptide of the present invention and is operably linked to additional nucleotides that provide for its expression.

Host cell: The term "host cell", as used herein, includes any cell type that is susceptible to transformation, transfection, transduction, and the like with a nucleic acid construct or expression vector comprising a polynucleotide of the present invention.

Modification: The term "modification" means herein any chemical modification of the polypeptide comprising or consisting of the mature polypeptide of SEQ ID NO: 2; or a homologous sequence thereof; as well as genetic manipulation of the DNA encoding such a polypeptide. The modification can be a substitution, a deletion and/or an insertion of one or more (several) amino acids as well as replacements of one or more (several) amino acid side chains.

Artificial variant: When used herein, the term "artificial variant" means a polypeptide having beta-glucosidase activity produced by an organism expressing a modified polynucleotide sequence of the mature polypeptide coding sequence of SEQ ID NO: 1; or a homologous sequence thereof. The modified nucleotide sequence is obtained through human intervention by modification of the polynucleotide sequence disclosed in SEQ ID NO: 1; or a homologous sequence thereof.

Detailed Description of the Invention

Polypeptides Having Beta-Glucosidase Activity

In a first aspect, the present invention relates to isolated polypeptides comprising amino acid sequences having a degree of sequence identity to the mature polypeptide of SEQ ID NO: 2 of preferably at least 60%, more preferably at least 65%, more preferably at least 70%, more preferably at least 75%, more preferably at least 80%, more preferably at least 85%, even more preferably at least 90%, most preferably at least 95%, and even most preferably at least 96%, at least 97%, at least 98%, or at least 99%, which have beta-glucosidase activity (hereinafter "homologous polypeptides"). In a preferred aspect, the homologous polypeptides comprise amino acid sequences that differ by ten amino acids, preferably by five amino acids, more preferably by four amino acids, even more preferably by three amino acids, most preferably by two amino acids, and even most preferably by one amino acid from the mature polypeptide of SEQ ID NO: 2.

A polypeptide of the present invention preferably comprises the amino acid sequence of SEQ ID NO: 2 or an allelic variant thereof; or a fragment thereof having beta-glucosidase activity. In a preferred aspect, the polypeptide comprises the amino acid sequence of SEQ ID NO: 2. In another preferred aspect, the polypeptide comprises the mature polypeptide of SEQ ID NO: 2. In another preferred aspect, the polypeptide comprises amino acids 20 to 856 of SEQ ID NO: 2, or an allelic variant thereof; or a fragment thereof having beta-glucosidase activity. In another preferred aspect, the polypeptide comprises amino acids 20 to 856 of SEQ ID NO: 2. In another preferred aspect, the polypeptide consists of the amino acid sequence of SEQ ID NO: 2 or an allelic variant thereof; or a fragment thereof having beta-glucosidase activity. In another preferred aspect, the polypeptide consists of the mature polypeptide of SEQ ID NO: 2. In another preferred aspect, the polypeptide consists of amino acids 20 to 856 of SEQ ID NO: 2 or an allelic variant thereof; or a fragment thereof having beta-glucosidase activity. In another preferred aspect, the polypeptide consists of amino acids 20 to 856 of SEQ ID NO: 2.

In a second aspect, the present invention relates to isolated polypeptides having beta-glucosidase activity that are encoded by polynucleotides that hybridize under preferably very low stringency conditions, more preferably low stringency conditions, more preferably medium stringency conditions, more preferably medium-high stringency conditions, even more preferably high stringency conditions, and most preferably very high stringency conditions with (i) the mature polypeptide coding sequence of SEQ ID NO: 1, (ii) the genomic DNA sequence comprising the mature polypeptide coding sequence of SEQ ID NO: 1, or (iii) a full-length complementary strand of (i) or (ii) (J. Sambrook, E.F. Fritsch, and T. Maniatis, 1989, *Molecular Cloning, A Laboratory Manual*, 2d edition, Cold Spring Harbor, New York).

The nucleotide sequence of SEQ ID NO: 1; or a subsequence thereof; as well as the amino acid sequence of SEQ ID NO: 2; or a fragment thereof; may be used to design nucleic acid probes to identify and clone DNA encoding polypeptides having beta-glucosidase activity from strains of different genera or species according to methods well known in the art. In particular, such probes can be used for hybridization with the genomic or cDNA of the genus or species of interest, following standard Southern blotting procedures, in order to identify and isolate the corresponding gene therein. Such probes can be considerably shorter than the entire sequence, but should be at least 14, preferably at least 25, more preferably at least 35, and most preferably at least 70 nucleotides in length. It is, however, preferred that the nucleic acid probe is at least 100 nucleotides in length. For example, the nucleic acid probe may be at least 200 nucleotides, preferably at least 300 nucleotides, more preferably at least 400 nucleotides, or most preferably at least 500 nucleotides in length.

Even longer probes may be used, e.g., nucleic acid probes that are preferably at least 600 nucleotides, more preferably at least 700 nucleotides, even more preferably at least 800 nucleotides, or most preferably at least 900 nucleotides in length. Both DNA and RNA probes can be used. The probes are typically labeled for detecting the corresponding gene (for example, with ^{32}P , ^3H , ^{35}S , biotin, or avidin). Such probes are encompassed by the present invention.

A genomic DNA or cDNA library prepared from such other strains may, therefore, be screened for DNA that hybridizes with the probes described above and encodes a polypeptide having beta-glucosidase activity. Genomic or other DNA from such other strains may be separated by agarose or polyacrylamide gel electrophoresis, or other separation techniques. DNA from the libraries or the separated DNA may be transferred to and immobilized on nitrocellulose or other suitable carrier material. In order to identify a clone or DNA that is homologous with SEQ ID NO: 1, or a subsequence thereof, the carrier material is preferably used in a Southern blot.

For purposes of the present invention, hybridization indicates that the nucleotide sequence hybridizes to a labeled nucleic acid probe corresponding to the mature polypeptide coding sequence of SEQ ID NO: 1; the genomic DNA sequence comprising the mature polypeptide coding sequence of SEQ ID NO: 1; its full-length complementary strand; or a subsequence thereof; under very low to very high stringency conditions. Molecules to which the nucleic acid probe hybridizes under these conditions can be detected using, for example, X-ray film.

In a preferred aspect, the nucleic acid probe is the mature polypeptide coding sequence of SEQ ID NO: 1. In another preferred aspect, the nucleic acid probe is nucleotides 58 to 2568 of SEQ ID NO: 1. In another preferred aspect, the nucleic acid probe is a polynucleotide sequence that encodes the polypeptide of SEQ ID NO: 2, or a subsequence thereof. In another preferred aspect, the nucleic acid probe is SEQ ID NO: 1. In another preferred aspect, the nucleic acid probe is the polynucleotide sequence contained in plasmid pAHYG-33 which is contained in *E. coli* NRRL B-50214, wherein the polynucleotide sequence thereof encodes a polypeptide having beta-glucosidase activity. In another preferred aspect, the nucleic acid probe is the mature polypeptide coding region contained in plasmid pAHYG-33 which is contained in *E. coli* NRRL B-50214.

For long probes of at least 100 nucleotides in length, very low to very high stringency conditions are defined as prehybridization and hybridization at 42°C in 5X SSPE, 0.3% SDS, 200 µg/ml sheared and denatured salmon sperm DNA, and either 25% formamide for very low and low stringencies, 35% formamide for medium and medium-high stringencies, or 50% formamide for high and very high stringencies, following standard Southern blotting procedures for 12 to 24 hours optimally.

For long probes of at least 100 nucleotides in length, the carrier material is finally washed three times each for 15 minutes using 2X SSC, 0.2% SDS preferably at 45°C (very low stringency), more preferably at 50°C (low stringency), more preferably at 55°C (medium stringency), more preferably at 60°C (medium-high stringency), even more preferably at 65°C (high stringency), and most preferably at 70°C (very high stringency).

For short probes of about 15 nucleotides to about 70 nucleotides in length, stringency conditions are defined as prehybridization, hybridization, and washing post-hybridization at about 5°C to about 10°C below the calculated T_m using the calculation according to Bolton and McCarthy (1962, *Proceedings of the National Academy of Sciences USA* 48:1390) in 0.9 M NaCl, 0.09 M Tris-HCl pH 7.6, 6 mM EDTA, 0.5% NP-40, 1X Denhardt's solution, 1 mM sodium pyrophosphate, 1 mM sodium monobasic phosphate, 0.1 mM ATP, and 0.2 mg of yeast RNA per ml following standard Southern blotting procedures for 12 to 24 hours optimally.

For short probes of about 15 nucleotides to about 70 nucleotides in length, the carrier material is washed once in 6X SSC plus 0.1% SDS for 15 minutes and twice each for 15 minutes using 6X SSC at 5°C to 10°C below the calculated T_m .

In a third aspect, the present invention relates to isolated polypeptides having beta-glucosidase activity encoded by polynucleotides comprising or consisting of nucleotide sequences having a degree of sequence identity to the mature polypeptide coding sequence of SEQ ID NO: 1 of preferably at least 60%, more preferably at least 65%, more preferably at least 70%, more preferably at least 75%, more preferably at least 80%, more preferably at least 85%, even more preferably at least 90%, most preferably at least 95%, and even most preferably at least 96%, at least 97%, at least 98%, or at least 99%, which encode a polypeptide having beta-glucosidase activity. See polynucleotide section herein.

In a fourth aspect, the present invention relates to artificial variants comprising a substitution, deletion, and/or insertion of one or more (or several) amino acids of the mature polypeptide of SEQ ID NO: 2, or a homologous sequence thereof. Preferably, amino acid changes are of a minor nature, that is conservative amino acid substitutions or insertions that do not significantly affect the folding and/or activity of the protein; small deletions, typically of one to about 30 amino acids; small amino- or carboxyl-terminal extensions, such as an amino-terminal methionine residue; a small linker peptide of up to about 20-25 residues; or a small extension that facilitates purification by changing net charge or another function, such as a poly-histidine tract, an antigenic epitope or a binding domain.

Examples of conservative substitutions are within the group of basic amino acids (arginine, lysine and histidine), acidic amino acids (glutamic acid and aspartic acid), polar amino acids (glutamine and asparagine), hydrophobic amino acids (leucine, isoleucine and valine), aromatic amino acids (phenylalanine, tryptophan and tyrosine), and small amino

acids (glycine, alanine, serine, threonine and methionine). Amino acid substitutions that do not generally alter specific activity are known in the art and are described, for example, by H. Neurath and R.L. Hill, 1979, *In, The Proteins*, Academic Press, New York. The most commonly occurring exchanges are Ala/Ser, Val/Ile, Asp/Glu, Thr/Ser, Ala/Gly, Ala/Thr, Ser/Asn, Ala/Val, Ser/Gly, Tyr/Phe, Ala/Pro, Lys/Arg, Asp/Asn, Leu/Ile, Leu/Val, Ala/Glu, and Asp/Gly.

In addition to the 20 standard amino acids, non-standard amino acids (such as 4-hydroxyproline, 6-N-methyl lysine, 2-aminoisobutyric acid, isovaline, and alpha-methyl serine) may be substituted for amino acid residues of a wild-type polypeptide. A limited number of non-conservative amino acids, amino acids that are not encoded by the genetic code, and unnatural amino acids may be substituted for amino acid residues. "Unnatural amino acids" have been modified after protein synthesis, and/or have a chemical structure in their side chain(s) different from that of the standard amino acids. Unnatural amino acids can be chemically synthesized, and preferably, are commercially available, and include pipercolic acid, thiazolidine carboxylic acid, dehydroproline, 3- and 4-methylproline, and 3,3-dimethylproline.

Alternatively, the amino acid changes are of such a nature that the physico-chemical properties of the polypeptides are altered. For example, amino acid changes may improve the thermal stability of the polypeptide, alter the substrate specificity, change the pH optimum, and the like.

Essential amino acids in the parent polypeptide can be identified according to procedures known in the art, such as site-directed mutagenesis or alanine-scanning mutagenesis (Cunningham and Wells, 1989, *Science* 244: 1081-1085). In the latter technique, single alanine mutations are introduced at every residue in the molecule, and the resultant mutant molecules are tested for biological activity (*i.e.*, beta-glucosidase activity) to identify amino acid residues that are critical to the activity of the molecule. See also, Hilton *et al.*, 1996, *J. Biol. Chem.* 271: 4699-4708. The active site of the enzyme or other biological interaction can also be determined by physical analysis of structure, as determined by such techniques as nuclear magnetic resonance, crystallography, electron diffraction, or photoaffinity labeling, in conjunction with mutation of putative contact site amino acids. See, for example, de Vos *et al.*, 1992, *Science* 255: 306-312; Smith *et al.*, 1992, *J. Mol. Biol.* 224: 899-904; Wlodaver *et al.*, 1992, *FEBS Lett.* 309: 59-64. The identities of essential amino acids can also be inferred from analysis of identities with polypeptides that are related to a polypeptide according to the invention.

Single or multiple amino acid substitutions, deletions, and/or insertions can be made and tested using known methods of mutagenesis, recombination, and/or shuffling, followed by a relevant screening procedure, such as those disclosed by Reidhaar-Olson and Sauer,

1988, *Science* 241: 53-57; Bowie and Sauer, 1989, *Proc. Natl. Acad. Sci. USA* 86: 2152-2156; WO 95/17413; or WO 95/22625. Other methods that can be used include error-prone PCR, phage display (e.g., Lowman *et al.*, 1991, *Biochem.* 30: 10832-10837; U.S. Patent No. 5,223,409; WO 92/06204), and region-directed mutagenesis (Derbyshire *et al.*, 1986, *Gene* 5: 46: 145; Ner *et al.*, 1988, *DNA* 7: 127).

Mutagenesis/shuffling methods can be combined with high-throughput, automated screening methods to detect activity of cloned, mutagenized polypeptides expressed by host cells (Ness *et al.*, 1999, *Nature Biotechnology* 17: 893-896). Mutagenized DNA molecules that encode active polypeptides can be recovered from the host cells and rapidly sequenced using standard methods in the art. These methods allow the rapid determination of the importance of individual amino acid residues in a polypeptide of interest, and can be applied to polypeptides of unknown structure.

The total number of amino acid substitutions, deletions and/or insertions of the mature polypeptide of SEQ ID NO: 2 is 10, preferably 9, more preferably 8, more preferably 7, more preferably 6, more preferably 5, more preferably 4, even more preferably 3, most preferably 2, and even most preferably 1.

Polypeptides of the present invention also include fused polypeptides or cleavable fusion polypeptides in which another polypeptide is fused at the N-terminus or the C-terminus of the polypeptide or fragment thereof. A fused polypeptide is produced by fusing a nucleotide sequence (or a portion thereof) encoding another polypeptide to a nucleotide sequence (or a portion thereof) of the present invention. Techniques for producing fusion polypeptides are known in the art, and include ligating the coding sequences encoding the polypeptides so that they are in frame and that expression of the fused polypeptide is under control of the same promoter(s) and terminator.

A fusion polypeptide can further comprise a cleavage site. Upon secretion of the fusion protein, the site is cleaved releasing the polypeptide having beta-glucosidase activity from the fusion protein. Examples of cleavage sites include, but are not limited to, a Kex2 site that encodes the dipeptide Lys-Arg (Martin *et al.*, 2003, *J. Ind. Microbiol. Biotechnol.* 3: 568-576; Svetina *et al.*, 2000, *J. Biotechnol.* 76: 245-251; Rasmussen-Wilson *et al.*, 1997, *Appl. Environ. Microbiol.* 63: 3488-3493; Ward *et al.*, 1995, *Biotechnology* 13: 498-503; and Contreras *et al.*, 1991, *Biotechnology* 9: 378-381), an Ile-(Glu or Asp)-Gly-Arg site, which is cleaved by a Factor Xa protease after the arginine residue (Eaton *et al.*, 1986, *Biochem.* 25: 505-512); a Asp-Asp-Asp-Asp-Lys site, which is cleaved by an enterokinase after the lysine (Collins-Racie *et al.*, 1995, *Biotechnology* 13: 982-987); a His-Tyr-Glu site or His-Tyr-Asp site, which is cleaved by Genenase I (Carter *et al.*, 1989, *Proteins: Structure, Function, and Genetics* 6: 240-248); a Leu-Val-Pro-Arg-Gly-Ser site, which is cleaved by thrombin after the Arg (Stevens, 2003, *Drug Discovery World* 4: 35-48); a Glu-Asn-Leu-Tyr-Phe-Gln-Gly site,

which is cleaved by TEV protease after the Gln (Stevens, 2003, *supra*); and a Leu-Glu-Val-Leu-Phe-Gln-Gly-Pro site, which is cleaved by a genetically engineered form of human rhinovirus 3C protease after the Gln (Stevens, 2003, *supra*).

5 Sources of Polypeptides Having Beta-Glucosidase Activity

A polypeptide having beta-glucosidase activity of the present invention may be obtained from microorganisms of any genus. For purposes of the present invention, the term "obtained from" as used herein in connection with a given source shall mean that the polypeptide encoded by a nucleotide sequence is produced by the source or by a strain in
10 which the nucleotide sequence from the source has been inserted. In a preferred aspect, the polypeptide obtained from a given source is secreted extracellularly.

A polypeptide having beta-glucosidase activity of the present invention may be a bacterial polypeptide. For example, the polypeptide may be a gram positive bacterial polypeptide such as a *Bacillus*, *Streptococcus*, *Streptomyces*, *Staphylococcus*,
15 *Enterococcus*, *Lactobacillus*, *Lactococcus*, *Clostridium*, *Geobacillus*, or *Oceanobacillus* polypeptide having beta-glucosidase activity, or a Gram negative bacterial polypeptide such as an *E. coli*, *Pseudomonas*, *Salmonella*, *Campylobacter*, *Helicobacter*, *Flavobacterium*, *Fusobacterium*, *Ilyobacter*, *Neisseria*, or *Ureaplasma* polypeptide having beta-glucosidase activity.

In a preferred aspect, the polypeptide is a *Bacillus alkalophilus*, *Bacillus amyloliquefaciens*, *Bacillus brevis*, *Bacillus circulans*, *Bacillus clausii*, *Bacillus coagulans*, *Bacillus firmus*, *Bacillus lautus*, *Bacillus lentus*, *Bacillus licheniformis*, *Bacillus megaterium*, *Bacillus pumilus*, *Bacillus stearothermophilus*, *Bacillus subtilis*, or *Bacillus thuringiensis* polypeptide having beta-glucosidase activity.
20

In another preferred aspect, the polypeptide is a *Streptococcus equisimilis*, *Streptococcus pyogenes*, *Streptococcus uberis*, or *Streptococcus equi* subsp. *Zooepidemicus* polypeptide having beta-glucosidase activity.
25

In another preferred aspect, the polypeptide is a *Streptomyces achromogenes*, *Streptomyces avermitilis*, *Streptomyces coelicolor*, *Streptomyces griseus*, or *Streptomyces lividans* polypeptide having beta-glucosidase activity.
30

A polypeptide having beta-glucosidase activity of the present invention may also be a fungal polypeptide, and more preferably a yeast polypeptide such as a *Candida*, *Kluyveromyces*, *Pichia*, *Saccharomyces*, *Schizosaccharomyces*, or *Yarrowia* polypeptide having beta-glucosidase activity; or more preferably a filamentous fungal polypeptide such
35 as an *Acremonium*, *Agaricus*, *Alternaria*, *Aspergillus*, *Aureobasidium*, *Botryosphaeria*, *Ceriporiopsis*, *Chaetomidium*, *Chrysosporium*, *Claviceps*, *Cochliobolus*, *Coprinopsis*, *Coptotermes*, *Corynascus*, *Cryphonectria*, *Cryptococcus*, *Diplodia*, *Exidia*, *Filibasidium*,

Fusarium, *Gibberella*, *Holomastigotoides*, *Humicola*, *Irpex*, *Lentinula*, *Leptosphaeria*, *Magnaporthe*, *Melanocarpus*, *Meripilus*, *Mucor*, *Myceliophthora*, *Neocallimastix*, *Neurospora*, *Paecilomyces*, *Penicillium*, *Phanerochaete*, *Piromyces*, *Poitrasia*, *Pseudoplectania*, *Pseudotriconympha*, *Rhizomucor*, *Schizophyllum*, *Scytalidium*, *Talaromyces*,
5 *Thermoascus*, *Thielavia*, *Tolypocladium*, *Trichoderma*, *Trichophaea*, *Verticillium*, *Volvariella*,
or *Xylaria* polypeptide having beta-glucosidase activity.

In a preferred aspect, the polypeptide is a *Saccharomyces carlsbergensis*, *Saccharomyces cerevisiae*, *Saccharomyces diastaticus*, *Saccharomyces douglasii*, *Saccharomyces kluyveri*, *Saccharomyces norbensis*, or *Saccharomyces oviformis*
10 polypeptide having beta-glucosidase activity.

In another preferred aspect, the polypeptide is an *Acremonium cellulolyticus*, *Aspergillus aculeatus*, *Aspergillus awamori*, *Aspergillus fumigatus*, *Aspergillus foetidus*, *Aspergillus japonicus*, *Aspergillus nidulans*, *Aspergillus niger*, *Aspergillus oryzae*, *Chrysosporium keratinophilum*, *Chrysosporium lucknowense*, *Chrysosporium tropicum*,
15 *Chrysosporium merdarium*, *Chrysosporium inops*, *Chrysosporium pannicola*, *Chrysosporium queenslandicum*, *Chrysosporium zonatum*, *Fusarium bactridioides*, *Fusarium cerealis*, *Fusarium crookwellense*, *Fusarium culmorum*, *Fusarium graminearum*, *Fusarium graminum*, *Fusarium heterosporum*, *Fusarium negundi*, *Fusarium oxysporum*, *Fusarium reticulatum*, *Fusarium roseum*, *Fusarium sambucinum*, *Fusarium sarcochromum*, *Fusarium sporotrichioides*, *Fusarium sulphureum*, *Fusarium torulosum*, *Fusarium trichothecioides*,
20 *Fusarium venenatum*, *Humicola grisea*, *Humicola insolens*, *Humicola lanuginosa*, *Irpex lacteus*, *Mucor miehei*, *Myceliophthora thermophila*, *Neurospora crassa*, *Penicillium funiculosum*, *Penicillium purpurogenum*, *Phanerochaete chrysosporium*, *Thielavia achromatica*, *Thielavia albomyces*, *Thielavia albopilosa*, *Thielavia australeinsis*, *Thielavia fimeti*, *Thielavia microspora*, *Thielavia ovispora*, *Thielavia peruviana*, *Thielavia spededonium*, *Thielavia setosa*, *Thielavia subthermophila*, *Thielavia terrestris*, *Trichoderma harzianum*, *Trichoderma koningii*, *Trichoderma longibrachiatum*, *Trichoderma reesei*, or *Trichoderma viride* polypeptide having beta-glucosidase activity.

In another preferred aspect, the polypeptide is a *Trichophaea abundans*,
30 *Trichophaea Trichophaea contradicta*, *Trichophaea hemisphaerioides*, *Trichophaea minuta*,
or *Trichophaea saccata* polypeptide having beta-glucosidase activity.

In a more preferred aspect, the polypeptide is a *Trichophaea saccata* polypeptide having beta-glucosidase activity. In a most preferred aspect, the polypeptide is a *Trichophaea saccata* CBS 804.70 polypeptide having beta-glucosidase activity, e.g., the
35 polypeptide comprising the amino acid sequence of SEQ ID NO: 2.

It will be understood that for the aforementioned species the invention encompasses both the perfect and imperfect states, and other taxonomic equivalents, e.g., anamorphs,

regardless of the species name by which they are known. Those skilled in the art will readily recognize the identity of appropriate equivalents.

Strains of these species are readily accessible to the public in a number of culture collections, such as the American Type Culture Collection (ATCC), Deutsche Sammlung von
5 Mikroorganismen und Zellkulturen GmbH (DSMZ), Centraalbureau Voor Schimmelcultures (CBS), and Agricultural Research Service Patent Culture Collection, Northern Regional Research Center (NRRL).

Furthermore, such polypeptides may be identified and obtained from other sources including microorganisms isolated from nature (*e.g.*, soil, composts, water, etc.) using the
10 above-mentioned probes. Techniques for isolating microorganisms from natural habitats are well known in the art. The polynucleotide may then be obtained by similarly screening a genomic or cDNA library of such a microorganism. Once a polynucleotide encoding a polypeptide has been detected with the probe(s), the polynucleotide can be isolated or cloned by utilizing techniques that are well known to those of ordinary skill in the art (see,
15 *e.g.*, Sambrook *et al.*, 1989, *supra*).

Polynucleotides

The present invention also relates to isolated polynucleotides comprising or consisting of nucleotide sequences that encode polypeptides having beta-glucosidase
20 activity of the present invention.

In a preferred aspect, the nucleotide sequence comprises or consists of SEQ ID NO: 1. In another more preferred aspect, the nucleotide sequence comprises or consists of the sequence contained in plasmid pAHYG-33 which is contained in *E. coli* NRRL B-50214. In another preferred aspect, the nucleotide sequence comprises or consists of the mature
25 polypeptide coding sequence of SEQ ID NO: 1. In another preferred aspect, the nucleotide sequence comprises or consists of nucleotides 58 to 2568 of SEQ ID NO: 1. In another more preferred aspect, the nucleotide sequence comprises or consists of the mature polypeptide coding sequence contained in plasmid pAHYG-33 which is contained in *E. coli* NRRL B-50214. The present invention also encompasses nucleotide sequences that
30 encode polypeptides comprising or consisting of the amino acid sequence of SEQ ID NO: 2 or the mature polypeptide thereof, which differ from SEQ ID NO: 1 or the mature polypeptide coding sequence thereof by virtue of the degeneracy of the genetic code. The present invention also relates to subsequences of SEQ ID NO: 1 that encode fragments of SEQ ID NO: 2 having beta-glucosidase activity.

The present invention also relates to mutant polynucleotides comprising or consisting
35 of at least one mutation in the mature polypeptide coding sequence of SEQ ID NO: 1, in which the mutant nucleotide sequence encodes the mature polypeptide of SEQ ID NO: 2.

The techniques used to isolate or clone a polynucleotide encoding a polypeptide are known in the art and include isolation from genomic DNA, preparation from cDNA, or a combination thereof. The cloning of the polynucleotides of the present invention from such genomic DNA can be effected, e.g., by using the well known polymerase chain reaction (PCR) or antibody screening of expression libraries to detect cloned DNA fragments with shared structural features. See, e.g., Innis *et al.*, 1990, *PCR: A Guide to Methods and Application*, Academic Press, New York. Other nucleic acid amplification procedures such as ligase chain reaction (LCR), ligation activated transcription (LAT) and nucleotide sequence-based amplification (NASBA) may be used. The polynucleotides may be cloned from a strain of *Trichophaea*, or another or related organism and thus, for example, may be an allelic or species variant of the polypeptide encoding region of the nucleotide sequence.

The present invention also relates to isolated polynucleotides comprising or consisting of nucleotide sequences having a degree of sequence identity to the mature polypeptide coding sequence of SEQ ID NO: 1 of preferably at least 60%, more preferably at least 65%, more preferably at least 70%, more preferably at least 75%, more preferably at least 80%, more preferably at least 85%, even more preferably at least 90%, most preferably at least 95%, and even most preferably at least 96%, at least 97%, at least 98%, or at least 99%, which encode a polypeptide having beta-glucosidase activity.

Modification of a nucleotide sequence encoding a polypeptide of the present invention may be necessary for the synthesis of polypeptides substantially similar to the polypeptide. The term "substantially similar" to the polypeptide refers to non-naturally occurring forms of the polypeptide. These polypeptides may differ in some engineered way from the polypeptide isolated from its native source, e.g., artificial variants that differ in specific activity, thermostability, pH optimum, or the like. The variant sequence may be constructed on the basis of the nucleotide sequence presented as the mature polypeptide coding sequence of SEQ ID NO: 1, e.g., a subsequence thereof, and/or by introduction of nucleotide substitutions that do not give rise to another amino acid sequence of the polypeptide encoded by the nucleotide sequence, but which correspond to the codon usage of the host organism intended for production of the enzyme, or by introduction of nucleotide substitutions that may give rise to a different amino acid sequence. For a general description of nucleotide substitution, see, e.g., Ford *et al.*, 1991, *Protein Expression and Purification 2*: 95-107.

It will be apparent to those skilled in the art that such substitutions can be made outside the regions critical to the function of the molecule and still result in an active polypeptide. Amino acid residues essential to the activity of the polypeptide encoded by an isolated polynucleotide of the invention, and therefore preferably not subject to substitution, may be identified according to procedures known in the art, such as site-directed

mutagenesis or alanine-scanning mutagenesis (see, *e.g.*, Cunningham and Wells, 1989, *supra*). In the latter technique, mutations are introduced at every positively charged residue in the molecule, and the resultant mutant molecules are tested for beta-glucosidase activity to identify amino acid residues that are critical to the activity of the molecule. Sites of substrate-enzyme interaction can also be determined by analysis of the three-dimensional structure as determined by such techniques as nuclear magnetic resonance analysis, crystallography or photoaffinity labeling (see, *e.g.*, de Vos *et al.*, 1992, *supra*; Smith *et al.*, 1992, *supra*; Wlodaver *et al.*, 1992, *supra*).

The present invention also relates to isolated polynucleotides encoding polypeptides of the present invention, which hybridize under preferably very low stringency conditions, more preferably low stringency conditions, more preferably medium stringency conditions, more preferably medium-high stringency conditions, even more preferably high stringency conditions, and most preferably very high stringency conditions with (i) the mature polypeptide coding sequence of SEQ ID NO: 1, (ii) the genomic DNA sequence comprising the mature polypeptide coding sequence of SEQ ID NO: 1, or (iii) a full-length complementary strand of (i) or (ii); or allelic variants and subsequences thereof (Sambrook *et al.*, 1989, *supra*), as defined herein.

The present invention also relates to isolated polynucleotides obtained by (a) hybridizing a population of DNA under preferably very low stringency conditions, more preferably low stringency conditions, more preferably medium stringency conditions, more preferably medium-high stringency conditions, even more preferably high stringency conditions, and most preferably very high stringency conditions with (i) the mature polypeptide coding sequence of SEQ ID NO: 1, (ii) the genomic DNA sequence comprising the mature polypeptide coding sequence of SEQ ID NO: 1, or (iii) a full-length complementary strand of (i) or (ii); and (b) isolating the hybridizing polynucleotide, which encodes a polypeptide having beta-glucosidase activity.

Nucleic Acid Constructs

The present invention also relates to nucleic acid constructs comprising an isolated polynucleotide of the present invention operably linked to one or more (several) control sequences that direct the expression of the coding sequence in a suitable host cell under conditions compatible with the control sequences.

An isolated polynucleotide encoding a polypeptide of the present invention may be manipulated in a variety of ways to provide for expression of the polypeptide. Manipulation of the polynucleotide's sequence prior to its insertion into a vector may be desirable or necessary depending on the expression vector. The techniques for modifying polynucleotide sequences utilizing recombinant DNA methods are well known in the art.

The control sequence may be an appropriate promoter sequence, a nucleotide sequence that is recognized by a host cell for expression of a polynucleotide encoding a polypeptide of the present invention. The promoter sequence contains transcriptional control sequences that mediate the expression of the polypeptide. The promoter may be any nucleotide sequence that shows transcriptional activity in the host cell of choice including mutant, truncated, and hybrid promoters, and may be obtained from genes encoding extracellular or intracellular polypeptides either homologous or heterologous to the host cell.

Examples of suitable promoters for directing the transcription of the nucleic acid constructs of the present invention, especially in a bacterial host cell, are the promoters obtained from the *E. coli lac* operon, *Streptomyces coelicolor* agarase gene (*dagA*), *Bacillus subtilis* levansucrase gene (*sacB*), *Bacillus licheniformis* alpha-amylase gene (*amyL*), *Bacillus stearothermophilus* maltogenic amylase gene (*amyM*), *Bacillus amyloliquefaciens* alpha-amylase gene (*amyQ*), *Bacillus licheniformis* penicillinase gene (*penP*), *Bacillus subtilis xylA* and *xylB* genes, and prokaryotic beta-lactamase gene (Villa-Kamaroff *et al.*, 1978, *Proceedings of the National Academy of Sciences USA* 75: 3727-3731), as well as the *tac* promoter (DeBoer *et al.*, 1983, *Proceedings of the National Academy of Sciences USA* 80: 21-25). Further promoters are described in "Useful proteins from recombinant bacteria" in *Scientific American*, 1980, 242: 74-94; and in Sambrook *et al.*, 1989, *supra*.

Examples of suitable promoters for directing the transcription of the nucleic acid constructs of the present invention in a filamentous fungal host cell are promoters obtained from the genes for *Aspergillus oryzae* TAKA amylase, *Rhizomucor miehei* aspartic proteinase, *Aspergillus niger* neutral alpha-amylase, *Aspergillus niger* acid stable alpha-amylase, *Aspergillus niger* or *Aspergillus awamori* glucoamylase (*glaA*), *Rhizomucor miehei* lipase, *Aspergillus oryzae* alkaline protease, *Aspergillus oryzae* triose phosphate isomerase, *Aspergillus nidulans* acetamidase, *Fusarium venenatum* amyloglucosidase (WO 00/56900), *Fusarium venenatum* Daria (WO 00/56900), *Fusarium venenatum* Quinn (WO 00/56900), *Fusarium oxysporum* trypsin-like protease (WO 96/00787), *Trichoderma reesei* beta-glucosidase, *Trichoderma reesei* cellobiohydrolase I, *Trichoderma reesei* cellobiohydrolase II, *Trichoderma reesei* endoglucanase I, *Trichoderma reesei* endoglucanase II, *Trichoderma reesei* endoglucanase III, *Trichoderma reesei* endoglucanase IV, *Trichoderma reesei* endoglucanase V, *Trichoderma reesei* xylanase I, *Trichoderma reesei* xylanase II, *Trichoderma reesei* beta-xylosidase, as well as a NA2-tpi promoter (a modified promoter including a gene encoding a neutral alpha-amylase in *Aspergilli* in which the untranslated leader has been replaced by an untranslated leader from a gene encoding triose phosphate isomerase in *Aspergilli*; non-limiting examples include modified promoters including the gene encoding neutral alpha-amylase in *Aspergillus niger* in which the untranslated leader has been replaced by an untranslated leader from the gene encoding triose phosphate

isomerase in *Aspergillus nidulans* or *Aspergillus oryzae*); and mutant, truncated, and hybrid promoters thereof.

In a yeast host, useful promoters are obtained from the genes for *Saccharomyces cerevisiae* enolase (ENO-1), *Saccharomyces cerevisiae* galactokinase (GAL1),
5 *Saccharomyces cerevisiae* alcohol dehydrogenase/glyceraldehyde-3-phosphate dehydrogenase (ADH1, ADH2/GAP), *Saccharomyces cerevisiae* triose phosphate isomerase (TPI), *Saccharomyces cerevisiae* metallothionein (CUP1), and *Saccharomyces cerevisiae* 3-phosphoglycerate kinase. Other useful promoters for yeast host cells are described by Romanos *et al.*, 1992, *Yeast* 8: 423-488.

10 The control sequence may also be a suitable transcription terminator sequence, a sequence recognized by a host cell to terminate transcription. The terminator sequence is operably linked to the 3' terminus of the nucleotide sequence encoding the polypeptide. Any terminator that is functional in the host cell of choice may be used in the present invention.

Preferred terminators for filamentous fungal host cells are obtained from the genes
15 for *Aspergillus oryzae* TKA amylose, *Aspergillus niger* glucoamylase, *Aspergillus nidulans* anthranilate synthase, *Aspergillus niger* alpha-glucosidase, and *Fusarium oxysporum* trypsin-like protease.

Preferred terminators for yeast host cells are obtained from the genes for
20 *Saccharomyces cerevisiae* enolase, *Saccharomyces cerevisiae* cytochrome C (CYC1), and *Saccharomyces cerevisiae* glyceraldehyde-3-phosphate dehydrogenase. Other useful terminators for yeast host cells are described by Romanos *et al.*, 1992, *supra*.

The control sequence may also be a suitable leader sequence, when transcribed is a
25 nontranslated region of an mRNA that is important for translation by the host cell. The leader sequence is operably linked to the 5' terminus of the nucleotide sequence encoding the polypeptide. Any leader sequence that is functional in the host cell of choice may be used in the present invention.

Preferred leaders for filamentous fungal host cells are obtained from the genes for
Aspergillus oryzae TKA amylose and *Aspergillus nidulans* triose phosphate isomerase.

Suitable leaders for yeast host cells are obtained from the genes for *Saccharomyces*
30 *cerevisiae* enolase (ENO-1), *Saccharomyces cerevisiae* 3-phosphoglycerate kinase, *Saccharomyces cerevisiae* alpha-factor, and *Saccharomyces cerevisiae* alcohol dehydrogenase/glyceraldehyde-3-phosphate dehydrogenase (ADH2/GAP).

The control sequence may also be a polyadenylation sequence, a sequence
35 operably linked to the 3' terminus of the nucleotide sequence and, when transcribed, is recognized by the host cell as a signal to add polyadenosine residues to transcribed mRNA. Any polyadenylation sequence that is functional in the host cell of choice may be used in the present invention.

Preferred polyadenylation sequences for filamentous fungal host cells are obtained from the genes for *Aspergillus oryzae* TAKA amylase, *Aspergillus niger* glucoamylase, *Aspergillus nidulans* anthranilate synthase, *Fusarium oxysporum* trypsin-like protease, and *Aspergillus niger* alpha-glucosidase.

5 Useful polyadenylation sequences for yeast host cells are described by Guo and Sherman, 1995, *Molecular Cellular Biology* 15: 5983-5990.

The control sequence may also be a signal peptide coding sequence that encodes a signal peptide linked to the amino terminus of a polypeptide and directs the encoded polypeptide into the cell's secretory pathway. The 5' end of the coding sequence of the nucleotide sequence may inherently contain a signal peptide coding sequence naturally linked in translation reading frame with the segment of the coding sequence that encodes the secreted polypeptide. Alternatively, the 5' end of the coding sequence may contain a signal peptide coding sequence that is foreign to the coding sequence. The foreign signal peptide coding sequence may be required where the coding sequence does not naturally contain a signal peptide coding sequence. Alternatively, the foreign signal peptide coding sequence may simply replace the natural signal peptide coding sequence in order to enhance secretion of the polypeptide. However, any signal peptide coding sequence that directs the expressed polypeptide into the secretory pathway of a host cell of choice, *i.e.*, secreted into a culture medium, may be used in the present invention.

20 Effective signal peptide coding sequences for bacterial host cells are the signal peptide coding sequences obtained from the genes for *Bacillus* NCIB 11837 maltogenic amylase, *Bacillus stearothermophilus* alpha-amylase, *Bacillus licheniformis* subtilisin, *Bacillus licheniformis* beta-lactamase, *Bacillus stearothermophilus* neutral proteases (*nprT*, *nprS*, *nprM*), and *Bacillus subtilis* *prSA*. Further signal peptides are described by Simonen and Palva, 1993, *Microbiological Reviews* 57: 109-137.

30 Effective signal peptide coding sequences for filamentous fungal host cells are the signal peptide coding sequences obtained from the genes for *Aspergillus oryzae* TAKA amylase, *Aspergillus niger* neutral amylase, *Aspergillus niger* glucoamylase, *Rhizomucor miehei* aspartic proteinase, *Humicola insolens* cellulase, *Humicola insolens* endoglucanase V, and *Humicola lanuginosa* lipase.

Useful signal peptides for yeast host cells are obtained from the genes for *Saccharomyces cerevisiae* alpha-factor and *Saccharomyces cerevisiae* invertase. Other useful signal peptide coding sequences are described by Romanos *et al.*, 1992, *supra*.

35 In a preferred aspect, the signal peptide comprises or consists of amino acids 1 to 19 of SEQ ID NO: 2. In another preferred aspect, the signal peptide coding sequence comprises or consists of nucleotides 1 to 57 of SEQ ID NO: 1.

The control sequence may also be a propeptide coding sequence that encodes a

propeptide positioned at the amino terminus of a polypeptide. The resultant polypeptide is known as a proenzyme or propolypeptide (or a zymogen in some cases). A propeptide is generally inactive and can be converted to a mature active polypeptide by catalytic or autocatalytic cleavage of the propeptide from the propolypeptide. The propeptide coding sequence may be obtained from the genes for *Bacillus subtilis* alkaline protease (*aprE*), *Bacillus subtilis* neutral protease (*nprT*), *Saccharomyces cerevisiae* alpha-factor, *Rhizomucor miehei* aspartic proteinase, and *Myceliophthora thermophila* laccase (WO 95/33836).

Where both signal peptide and propeptide sequences are present at the amino terminus of a polypeptide, the propeptide sequence is positioned next to the amino terminus of a polypeptide and the signal peptide sequence is positioned next to the amino terminus of the propeptide sequence.

It may also be desirable to add regulatory sequences that allow the regulation of the expression of the polypeptide relative to the growth of the host cell. Examples of regulatory systems are those that cause the expression of the gene to be turned on or off in response to a chemical or physical stimulus, including the presence of a regulatory compound. Regulatory systems in prokaryotic systems include the *lac*, *tac*, and *trp* operator systems. In yeast, the ADH2 system or GAL1 system may be used. In filamentous fungi, the TAKA alpha-amylase promoter, *Aspergillus niger* glucoamylase promoter, and *Aspergillus oryzae* glucoamylase promoter may be used as regulatory sequences. Other examples of regulatory sequences are those that allow for gene amplification. In eukaryotic systems, these regulatory sequences include the dihydrofolate reductase gene that is amplified in the presence of methotrexate, and the metallothionein genes that are amplified with heavy metals. In these cases, the nucleotide sequence encoding the polypeptide would be operably linked with the regulatory sequence.

Expression Vectors

The present invention also relates to recombinant expression vectors comprising a polynucleotide of the present invention, a promoter, and transcriptional and translational stop signals. The various nucleic acids and control sequences described herein may be joined together to produce a recombinant expression vector that may include one or more (several) convenient restriction sites to allow for insertion or substitution of the nucleotide sequence encoding the polypeptide at such sites. Alternatively, a polynucleotide sequence of the present invention may be expressed by inserting the nucleotide sequence or a nucleic acid construct comprising the sequence into an appropriate vector for expression. In creating the expression vector, the coding sequence is located in the vector so that the coding sequence is operably linked with the appropriate control sequences for expression.

The recombinant expression vector may be any vector (e.g., a plasmid or virus) that can be conveniently subjected to recombinant DNA procedures and can bring about expression of the nucleotide sequence. The choice of the vector will typically depend on the compatibility of the vector with the host cell into which the vector is to be introduced. The
5 vectors may be linear or closed circular plasmids.

The vector may be an autonomously replicating vector, i.e., a vector that exists as an extrachromosomal entity, the replication of which is independent of chromosomal replication, e.g., a plasmid, an extrachromosomal element, a minichromosome, or an artificial chromosome. The vector may contain any means for assuring self-replication. Alternatively,
10 the vector may be one that, when introduced into the host cell, is integrated into the genome and replicated together with the chromosome(s) into which it has been integrated. Furthermore, a single vector or plasmid or two or more vectors or plasmids that together contain the total DNA to be introduced into the genome of the host cell, or a transposon, may be used.

The vectors of the present invention preferably contain one or more (several) selectable markers that permit easy selection of transformed, transfected, transduced, or the like cells. A selectable marker is a gene the product of which provides for biocide or viral resistance, resistance to heavy metals, prototrophy to auxotrophs, and the like.
15

Examples of bacterial selectable markers are the *dal* genes from *Bacillus subtilis* or
20 *Bacillus licheniformis*, or markers that confer antibiotic resistance such as ampicillin, kanamycin, chloramphenicol, or tetracycline resistance. Suitable markers for yeast host cells are ADE2, HIS3, LEU2, LYS2, MET3, TRP1, and URA3. Selectable markers for use in a filamentous fungal host cell include, but are not limited to, *amdS* (acetamidase), *argB* (ornithine carbamoyltransferase), *bar* (phosphinothricin acetyltransferase), *hph* (hygromycin phosphotransferase), *niaD* (nitrate reductase), *pyrG* (orotidine-5'-phosphate decarboxylase),
25 *sC* (sulfate adenylyltransferase), and *trpC* (anthranilate synthase), as well as equivalents thereof. Preferred for use in an *Aspergillus* cell are the *amdS* and *pyrG* genes of *Aspergillus nidulans* or *Aspergillus oryzae* and the *bar* gene of *Streptomyces hygroscopicus*.

The vectors of the present invention preferably contain an element(s) that permits
30 integration of the vector into the host cell's genome or autonomous replication of the vector in the cell independent of the genome.

For integration into the host cell genome, the vector may rely on the polynucleotide's sequence encoding the polypeptide or any other element of the vector for integration into the genome by homologous or non-homologous recombination. Alternatively, the vector may
35 contain additional nucleotide sequences for directing integration by homologous recombination into the genome of the host cell at a precise location(s) in the chromosome(s). To increase the likelihood of integration at a precise location, the integrational elements

should preferably contain a sufficient number of nucleic acids, such as 100 to 10,000 base pairs, preferably 400 to 10,000 base pairs, and most preferably 800 to 10,000 base pairs, which have a high degree of sequence identity to the corresponding target sequence to enhance the probability of homologous recombination. The integrational elements may be any sequence that is homologous with the target sequence in the genome of the host cell. Furthermore, the integrational elements may be non-encoding or encoding nucleotide sequences. On the other hand, the vector may be integrated into the genome of the host cell by non-homologous recombination.

For autonomous replication, the vector may further comprise an origin of replication enabling the vector to replicate autonomously in the host cell in question. The origin of replication may be any plasmid replicator mediating autonomous replication that functions in a cell. The term "origin of replication" or "plasmid replicator" is defined herein as a nucleotide sequence that enables a plasmid or vector to replicate *in vivo*.

Examples of bacterial origins of replication are the origins of replication of plasmids pBR322, pUC19, pACYC177, and pACYC184 permitting replication in *E. coli*, and pUB110, pE194, pTA1060, and pAM β 1 permitting replication in *Bacillus*.

Examples of origins of replication for use in a yeast host cell are the 2 micron origin of replication, ARS1, ARS4, the combination of ARS1 and CEN3, and the combination of ARS4 and CEN6.

Examples of origins of replication useful in a filamentous fungal cell are AMA1 and ANS1 (Gems *et al.*, 1991, *Gene* 98: 61-67; Cullen *et al.*, 1987, *Nucleic Acids Research* 15: 9163-9175; WO 00/24883). Isolation of the AMA1 gene and construction of plasmids or vectors comprising the gene can be accomplished according to the methods disclosed in WO 00/24883.

More than one copy of a polynucleotide of the present invention may be inserted into a host cell to increase production of the gene product. An increase in the copy number of the polynucleotide can be obtained by integrating at least one additional copy of the sequence into the host cell genome or by including an amplifiable selectable marker gene with the polynucleotide where cells containing amplified copies of the selectable marker gene, and thereby additional copies of the polynucleotide, can be selected for by cultivating the cells in the presence of the appropriate selectable agent.

The procedures used to ligate the elements described above to construct the recombinant expression vectors of the present invention are well known to one skilled in the art (see, *e.g.*, Sambrook *et al.*, 1989, *supra*).

Host Cells

The present invention also relates to recombinant host cells, comprising an isolated

polynucleotide of the present invention operably linked to one or more (several) control sequences that direct the production of a polypeptide having beta-glucosidase activity. A construct or vector comprising a polynucleotide of the present invention is introduced into a host cell so that the construct or vector is maintained as a chromosomal integrant or as a self-replicating extra-chromosomal vector as described earlier. The term "host cell" encompasses any progeny of a parent cell that is not identical to the parent cell due to mutations that occur during replication. The choice of a host cell will to a large extent depend upon the gene encoding the polypeptide and its source.

The host cell may be any cell useful in the recombinant production of a polypeptide of the present invention, e.g., a prokaryote or a eukaryote.

The prokaryotic host cell may be any Gram positive bacterium or a Gram negative bacterium. Gram positive bacteria include, but not limited to, *Bacillus*, *Streptococcus*, *Streptomyces*, *Staphylococcus*, *Enterococcus*, *Lactobacillus*, *Lactococcus*, *Clostridium*, *Geobacillus*, and *Oceanobacillus*. Gram negative bacteria include, but not limited to, *E. coli*, *Pseudomonas*, *Salmonella*, *Campylobacter*, *Helicobacter*, *Flavobacterium*, *Fusobacterium*, *Ilyobacter*, *Neisseria*, and *Ureaplasma*.

The bacterial host cell may be any *Bacillus* cell. *Bacillus* cells useful in the practice of the present invention include, but are not limited to, *Bacillus alkalophilus*, *Bacillus amyloliquefaciens*, *Bacillus brevis*, *Bacillus circulans*, *Bacillus clausii*, *Bacillus coagulans*, *Bacillus firmus*, *Bacillus lautus*, *Bacillus lentus*, *Bacillus licheniformis*, *Bacillus megaterium*, *Bacillus pumilus*, *Bacillus stearothermophilus*, *Bacillus subtilis*, and *Bacillus thuringiensis* cells.

In a preferred aspect, the bacterial host cell is a *Bacillus amyloliquefaciens* cell. In another preferred aspect, the bacterial host cell is a *Bacillus clausii* cell. In another preferred aspect, the bacterial host cell is a *Bacillus lentus* cell. In another preferred aspect, the bacterial host cell is a *Bacillus licheniformis* cell. In another preferred aspect, the bacterial host cell is a *Bacillus stearothermophilus* cell. In another preferred aspect, the bacterial host cell is a *Bacillus subtilis* cell.

The bacterial host cell may also be any *Streptococcus* cell. *Streptococcus* cells useful in the practice of the present invention include, but are not limited to, *Streptococcus equisimilis*, *Streptococcus pyogenes*, *Streptococcus uberis*, and *Streptococcus equi* subsp. *Zooepidemicus* cells.

In a preferred aspect, the bacterial host cell is a *Streptococcus equisimilis* cell. In another preferred aspect, the bacterial host cell is a *Streptococcus pyogenes* cell. In another preferred aspect, the bacterial host cell is a *Streptococcus uberis* cell. In another preferred aspect, the bacterial host cell is a *Streptococcus equi* subsp. *Zooepidemicus* cell.

The bacterial host cell may also be any *Streptomyces* cell. *Streptomyces* cells useful

in the practice of the present invention include, but are not limited to, *Streptomyces achromogenes*, *Streptomyces avermitilis*, *Streptomyces coelicolor*, *Streptomyces griseus*, and *Streptomyces lividans* cells.

In a preferred aspect, the bacterial host cell is a *Streptomyces achromogenes* cell. In another preferred aspect, the bacterial host cell is a *Streptomyces avermitilis* cell. In another preferred aspect, the bacterial host cell is a *Streptomyces coelicolor* cell. In another preferred aspect, the bacterial host cell is a *Streptomyces griseus* cell. In another preferred aspect, the bacterial host cell is a *Streptomyces lividans* cell.

The introduction of DNA into a *Bacillus* cell may, for instance, be effected by protoplast transformation (see, e.g., Chang and Cohen, 1979, *Molecular General Genetics* 168: 111-115), by using competent cells (see, e.g., Young and Spizizen, 1961, *Journal of Bacteriology* 81: 823-829, or Dubnau and Davidoff-Abelson, 1971, *Journal of Molecular Biology* 56: 209-221), by electroporation (see, e.g., Shigekawa and Dower, 1988, *Biotechniques* 6: 742-751), or by conjugation (see, e.g., Koehler and Thorne, 1987, *Journal of Bacteriology* 169: 5271-5278). The introduction of DNA into an *E coli* cell may, for instance, be effected by protoplast transformation (see, e.g., Hanahan, 1983, *J. Mol. Biol.* 166: 557-580) or electroporation (see, e.g., Dower *et al.*, 1988, *Nucleic Acids Res.* 16: 6127-6145). The introduction of DNA into a *Streptomyces* cell may, for instance, be effected by protoplast transformation and electroporation (see, e.g., Gong *et al.*, 2004, *Folia Microbiol. (Praha)* 49: 399-405), by conjugation (see, e.g., Mazodier *et al.*, 1989, *J. Bacteriol.* 171: 3583-3585), or by transduction (see, e.g., Burke *et al.*, 2001, *Proc. Natl. Acad. Sci. USA* 98: 6289-6294). The introduction of DNA into a *Pseudomonas* cell may, for instance, be effected by electroporation (see, e.g., Choi *et al.*, 2006, *J. Microbiol. Methods* 64: 391-397) or by conjugation (see, e.g., Pinedo and Smets, 2005, *Appl. Environ. Microbiol.* 71: 51-57). The introduction of DNA into a *Streptococcus* cell may, for instance, be effected by natural competence (see, e.g., Perry and Kuramitsu, 1981, *Infect. Immun.* 32: 1295-1297), by protoplast transformation (see, e.g., Catt and Jollick, 1991, *Microbios.* 68: 189-207, by electroporation (see, e.g., Buckley *et al.*, 1999, *Appl. Environ. Microbiol.* 65: 3800-3804) or by conjugation (see, e.g., Clewell, 1981, *Microbiol. Rev.* 45: 409-436). However, any method known in the art for introducing DNA into a host cell can be used.

The host cell may also be a eukaryote, such as a mammalian, insect, plant, or fungal cell.

In a preferred aspect, the host cell is a fungal cell. "Fungi" as used herein includes the phyla Ascomycota, Basidiomycota, Chytridiomycota, and Zygomycota (as defined by Hawksworth *et al.*, *In, Ainsworth and Bisby's Dictionary of The Fungi*, 8th edition, 1995, CAB International, University Press, Cambridge, UK) as well as the Oomycota (as cited in Hawksworth *et al.*, 1995, *supra*, page 171) and all mitosporic fungi (Hawksworth *et al.*, 1995,

supra).

In a more preferred aspect, the fungal host cell is a yeast cell. "Yeast" as used herein includes ascosporogenous yeast (Endomycetales), basidiosporogenous yeast, and yeast belonging to the Fungi Imperfecti (Blastomycetes). Since the classification of yeast may change in the future, for the purposes of this invention, yeast shall be defined as described in *Biology and Activities of Yeast* (Skinner, F.A., Passmore, S.M., and Davenport, R.R., eds, *Soc. App. Bacteriol. Symposium Series No. 9*, 1980).

In an even more preferred aspect, the yeast host cell is a *Candida*, *Hansenula*, *Kluyveromyces*, *Pichia*, *Saccharomyces*, *Schizosaccharomyces*, or *Yarrowia* cell.

In a most preferred aspect, the yeast host cell is a *Saccharomyces carlsbergensis* cell. In another most preferred aspect, the yeast host cell is a *Saccharomyces cerevisiae* cell. In another most preferred aspect, the yeast host cell is a *Saccharomyces diastaticus* cell. In another most preferred aspect, the yeast host cell is a *Saccharomyces douglasii* cell. In another most preferred aspect, the yeast host cell is a *Saccharomyces kluyveri* cell. In another most preferred aspect, the yeast host cell is a *Saccharomyces norbensis* cell. In another most preferred aspect, the yeast host cell is a *Saccharomyces oviformis* cell. In another most preferred aspect, the yeast host cell is a *Kluyveromyces lactis* cell. In another most preferred aspect, the yeast host cell is a *Yarrowia lipolytica* cell.

In another more preferred aspect, the fungal host cell is a filamentous fungal cell. "Filamentous fungi" include all filamentous forms of the subdivision Eumycota and Oomycota (as defined by Hawksworth *et al.*, 1995, *supra*). The filamentous fungi are generally characterized by a mycelial wall composed of chitin, cellulose, glucan, chitosan, mannan, and other complex polysaccharides. Vegetative growth is by hyphal elongation and carbon catabolism is obligately aerobic. In contrast, vegetative growth by yeasts such as *Saccharomyces cerevisiae* is by budding of a unicellular thallus and carbon catabolism may be fermentative.

In an even more preferred aspect, the filamentous fungal host cell is an *Acremonium*, *Aspergillus*, *Aureobasidium*, *Bjerkandera*, *Ceriporiopsis*, *Chrysosporium*, *Coprinus*, *Coriolus*, *Cryptococcus*, *Filibasidium*, *Fusarium*, *Humicola*, *Magnaporthe*, *Mucor*, *Myceliophthora*, *Neocallimastix*, *Neurospora*, *Paecilomyces*, *Penicillium*, *Phanerochaete*, *Phlebia*, *Piromyces*, *Pleurotus*, *Schizophyllum*, *Talaromyces*, *Thermoascus*, *Thielavia*, *Tolypocladium*, *Trametes*, or *Trichoderma* cell.

In a most preferred aspect, the filamentous fungal host cell is an *Aspergillus awamori*, *Aspergillus fumigatus*, *Aspergillus foetidus*, *Aspergillus japonicus*, *Aspergillus nidulans*, *Aspergillus niger* or *Aspergillus oryzae* cell. In another most preferred aspect, the filamentous fungal host cell is a *Fusarium bactridioides*, *Fusarium cerealis*, *Fusarium crookwellense*, *Fusarium culmorum*, *Fusarium graminearum*, *Fusarium graminum*, *Fusarium*

heterosporum, *Fusarium negundi*, *Fusarium oxysporum*, *Fusarium reticulatum*, *Fusarium roseum*, *Fusarium sambucinum*, *Fusarium sarcochroum*, *Fusarium sporotrichioides*, *Fusarium sulphureum*, *Fusarium torulosum*, *Fusarium trichothecioides*, or *Fusarium venenatum* cell. In another most preferred aspect, the filamentous fungal host cell is a
5 *Bjerkandera adusta*, *Ceriporiopsis aneirina*, *Ceriporiopsis aneirina*, *Ceriporiopsis caregiea*, *Ceriporiopsis gilvescens*, *Ceriporiopsis pannocinta*, *Ceriporiopsis rivulosa*, *Ceriporiopsis subrufa*, *Ceriporiopsis subvermispora*, *Chrysosporium keratinophilum*, *Chrysosporium lucknowense*, *Chrysosporium tropicum*, *Chrysosporium merdarium*, *Chrysosporium inops*, *Chrysosporium pannicola*, *Chrysosporium queenslandicum*, *Chrysosporium zonatum*,
10 *Coprinus cinereus*, *Coriolus hirsutus*, *Humicola insolens*, *Humicola lanuginosa*, *Mucor miehei*, *Myceliophthora thermophila*, *Neurospora crassa*, *Penicillium purpurogenum*, *Phanerochaete chrysosporium*, *Phlebia radiata*, *Pleurotus eryngii*, *Thielavia terrestris*, *Trametes villosa*, *Trametes versicolor*, *Trichoderma harzianum*, *Trichoderma koningii*, *Trichoderma longibrachiatum*, *Trichoderma reesei*, or *Trichoderma viride* cell.

15 In another most preferred aspect, the filamentous fungal host cell is an *Aspergillus niger* cell. In another most preferred aspect, the filamentous fungal host cell is an *Aspergillus oryzae* cell. In another most preferred aspect, the filamentous fungal host cell is a *Chrysosporium lucknowense* cell. In another most preferred aspect, the filamentous fungal host cell is a *Fusarium venenatum* cell. In another a most preferred spect, the
20 filamentous fungal host cell is a *Myceliophthora thermophila* cell. In another most preferred aspect, the filamentous fungal host cell is a *Trichoderma reesei* cell.

Fungal cells may be transformed by a process involving protoplast formation, transformation of the protoplasts, and regeneration of the cell wall in a manner known *per se*. Suitable procedures for transformation of *Aspergillus* and *Trichoderma* host cells are
25 described in EP 238 023 and Yelton *et al.*, 1984, *Proceedings of the National Academy of Sciences USA* 81: 1470-1474. Suitable methods for transforming *Fusarium* species are described by Malardier *et al.*, 1989, *Gene* 78: 147-156, and WO 96/00787. Yeast may be transformed using the procedures described by Becker and Guarente, *In* Abelson, J.N. and Simon, M.I., editors, *Guide to Yeast Genetics and Molecular Biology, Methods in*
30 *Enzymology*, Volume 194, pp 182-187, Academic Press, Inc., New York; Ito *et al.*, 1983, *Journal of Bacteriology* 153: 163; and Hinnen *et al.*, 1978, *Proceedings of the National Academy of Sciences USA* 75: 1920.

Methods of Production

35 The present invention also relates to methods of producing a polypeptide of the present invention, comprising: (a) cultivating a cell, which in its wild-type form produces the polypeptide, under conditions conducive for production of the polypeptide; and (b) recovering

the polypeptide. In a preferred aspect, the cell is of the genus *Trichophaea*. In a more preferred aspect, the cell is *Trichophaea saccata*. In a most preferred aspect, the cell is *Trichophaea saccata* CBS 804.70.

5 The present invention also relates to methods of producing a polypeptide of the present invention, comprising: (a) cultivating a recombinant host cell, as described herein, under conditions conducive for production of the polypeptide; and (b) recovering the polypeptide.

10 The present invention also relates to methods of producing a polypeptide of the present invention, comprising: (a) cultivating a recombinant host cell under conditions conducive for production of the polypeptide, wherein the host cell comprises a mutant nucleotide sequence having at least one mutation in the mature polypeptide coding sequence of SEQ ID NO: 1, wherein the mutant nucleotide sequence encodes a polypeptide that comprises or consists of the mature polypeptide of SEQ ID NO: 2; and (b) recovering the polypeptide.

15 In the production methods of the present invention, the cells are cultivated in a nutrient medium suitable for production of the polypeptide using methods well known in the art. For example, the cell may be cultivated by shake flask cultivation, and small-scale or large-scale fermentation (including continuous, batch, fed-batch, or solid state fermentations) in laboratory or industrial fermentors performed in a suitable medium and under conditions
20 allowing the polypeptide to be expressed and/or isolated. The cultivation takes place in a suitable nutrient medium comprising carbon and nitrogen sources and inorganic salts, using procedures known in the art. Suitable media are available from commercial suppliers or may be prepared according to published compositions (e.g., in catalogues of the American Type Culture Collection). If the polypeptide is secreted into the nutrient medium, the polypeptide
25 can be recovered directly from the medium. If the polypeptide is not secreted into the medium, it can be recovered from cell lysates.

The polypeptides may be detected using methods known in the art that are specific for the polypeptides. These detection methods may include use of specific antibodies, formation of an enzyme product, or disappearance of an enzyme substrate. For example,
30 an enzyme assay may be used to determine the activity of the polypeptide as described herein.

The resulting polypeptide may be recovered using methods known in the art. For example, the polypeptide may be recovered from the nutrient medium by conventional procedures including, but not limited to, centrifugation, filtration, extraction, spray-drying,
35 evaporation, or precipitation.

The polypeptides of the present invention may be purified by a variety of procedures known in the art including, but not limited to, chromatography (e.g., ion exchange, affinity,

hydrophobic, chromatofocusing, and size exclusion), electrophoretic procedures (e.g., preparative isoelectric focusing), differential solubility (e.g., ammonium sulfate precipitation), SDS-PAGE, or extraction (see, e.g., *Protein Purification*, J.-C. Janson and Lars Ryden, editors, VCH Publishers, New York, 1989) to obtain substantially pure polypeptides.

5

Plants

The present invention also relates to plants, e.g., a transgenic plant, plant part, or plant cell, comprising an isolated polynucleotide encoding a polypeptide having beta-glucosidase activity of the present invention so as to express and produce the polypeptide in recoverable quantities. The polypeptide may be recovered from the plant or plant part. Alternatively, the plant or plant part containing the recombinant polypeptide may be used as such for improving the quality of a food or feed, e.g., improving nutritional value, palatability, and rheological properties, or to destroy an antinutritive factor.

The transgenic plant can be dicotyledonous (a dicot) or monocotyledonous (a monocot). Examples of monocot plants are grasses, such as meadow grass (blue grass, Poa), forage grass such as Festuca, Lolium, temperate grass, such as Agrostis, and cereals, e.g., wheat, oats, rye, barley, rice, sorghum, and maize (corn).

Examples of dicot plants are tobacco, legumes, such as lupins, potato, sugar beet, pea, bean and soybean, and cruciferous plants (family Brassicaceae), such as cauliflower, rape seed, and the closely related model organism *Arabidopsis thaliana*.

Examples of plant parts are stem, callus, leaves, root, fruits, seeds, and tubers as well as the individual tissues comprising these parts, e.g., epidermis, mesophyll, parenchyme, vascular tissues, meristems. Specific plant cell compartments, such as chloroplasts, apoplasts, mitochondria, vacuoles, peroxisomes and cytoplasm are also considered to be a plant part. Furthermore, any plant cell, whatever the tissue origin, is considered to be a plant part. Likewise, plant parts such as specific tissues and cells isolated to facilitate the utilisation of the invention are also considered plant parts, e.g., embryos, endosperms, aleurone and seeds coats.

Also included within the scope of the present invention are the progeny of such plants, plant parts, and plant cells.

The transgenic plant or plant cell expressing a polypeptide of the present invention may be constructed in accordance with methods known in the art. In short, the plant or plant cell is constructed by incorporating one or more (several) expression constructs encoding a polypeptide of the present invention into the plant host genome or chloroplast genome and propagating the resulting modified plant or plant cell into a transgenic plant or plant cell.

The expression construct is conveniently a nucleic acid construct that comprises a polynucleotide encoding a polypeptide of the present invention operably linked with

appropriate regulatory sequences required for expression of the nucleotide sequence in the plant or plant part of choice. Furthermore, the expression construct may comprise a selectable marker useful for identifying host cells into which the expression construct has been integrated and DNA sequences necessary for introduction of the construct into the plant in question (the latter depends on the DNA introduction method to be used).

The choice of regulatory sequences, such as promoter and terminator sequences and optionally signal or transit sequences, is determined, for example, on the basis of when, where, and how the polypeptide is desired to be expressed. For instance, the expression of the gene encoding a polypeptide of the present invention may be constitutive or inducible, or may be developmental, stage or tissue specific, and the gene product may be targeted to a specific tissue or plant part such as seeds or leaves. Regulatory sequences are, for example, described by Tague *et al.*, 1988, *Plant Physiology* 86: 506.

For constitutive expression, the 35S-CaMV, the maize ubiquitin 1, and the rice actin 1 promoter may be used (Franck *et al.*, 1980, *Cell* 21: 285-294; Christensen *et al.*, 1992, *Plant Mol. Biol.* 18: 675-689; Zhang *et al.*, 1991, *Plant Cell* 3: 1155-1165). Organ-specific promoters may be, for example, a promoter from storage sink tissues such as seeds, potato tubers, and fruits (Edwards and Coruzzi, 1990, *Ann. Rev. Genet.* 24: 275-303), or from metabolic sink tissues such as meristems (Ito *et al.*, 1994, *Plant Mol. Biol.* 24: 863-878), a seed specific promoter such as the glutelin, prolamin, globulin, or albumin promoter from rice (Wu *et al.*, 1998, *Plant and Cell Physiology* 39: 885-889), a *Vicia faba* promoter from the legumin B4 and the unknown seed protein gene from *Vicia faba* (Conrad *et al.*, 1998, *Journal of Plant Physiology* 152: 708-711), a promoter from a seed oil body protein (Chen *et al.*, 1998, *Plant and Cell Physiology* 39: 935-941), the storage protein *napA* promoter from *Brassica napus*, or any other seed specific promoter known in the art, *e.g.*, as described in WO 91/14772. Furthermore, the promoter may be a leaf specific promoter such as the *rbcs* promoter from rice or tomato (Kyojuka *et al.*, 1993, *Plant Physiology* 102: 991-1000, the chlorella virus adenine methyltransferase gene promoter (Mitra and Higgins, 1994, *Plant Molecular Biology* 26: 85-93), or the *aldP* gene promoter from rice (Kagaya *et al.*, 1995, *Molecular and General Genetics* 248: 668-674), or a wound inducible promoter such as the potato *pin2* promoter (Xu *et al.*, 1993, *Plant Molecular Biology* 22: 573-588). Likewise, the promoter may be inducible by abiotic treatments such as temperature, drought, or alterations in salinity or induced by exogenously applied substances that activate the promoter, *e.g.*, ethanol, oestrogens, plant hormones such as ethylene, abscisic acid, and gibberellic acid, and heavy metals.

A promoter enhancer element may also be used to achieve higher expression of a polypeptide of the present invention in the plant. For instance, the promoter enhancer element may be an intron that is placed between the promoter and the nucleotide sequence

encoding a polypeptide of the present invention. For instance, Xu *et al.*, 1993, *supra*, disclose the use of the first intron of the rice actin 1 gene to enhance expression.

The selectable marker gene and any other parts of the expression construct may be chosen from those available in the art.

5 The nucleic acid construct is incorporated into the plant genome according to conventional techniques known in the art, including *Agrobacterium*-mediated transformation, virus-mediated transformation, microinjection, particle bombardment, biolistic transformation, and electroporation (Gasser *et al.*, 1990, *Science* 244: 1293; Potrykus, 1990, *Bio/Technology* 8: 535; Shimamoto *et al.*, 1989, *Nature* 338: 274).

10 Presently, *Agrobacterium tumefaciens*-mediated gene transfer is the method of choice for generating transgenic dicots (for a review, see Hooykas and Schilperoort, 1992, *Plant Molecular Biology* 19: 15-38) and can also be used for transforming monocots, although other transformation methods are often used for these plants. Presently, the method of choice for generating transgenic monocots is particle bombardment (microscopic gold or tungsten particles coated with the transforming DNA) of embryonic calli or developing embryos (Christou, 1992, *Plant Journal* 2: 275-281; Shimamoto, 1994, *Current Opinion Biotechnology* 5: 158-162; Vasil *et al.*, 1992, *Bio/Technology* 10: 667-674). An alternative method for transformation of monocots is based on protoplast transformation as described by Omirulleh *et al.*, 1993, *Plant Molecular Biology* 21: 415-428. Additional transformation methods for use in accordance with the present disclosure include those described in U.S. Patent Nos. 6,395,966 and 7,151,204 (both of which are herein incorporated by reference in their entirety).

20 Following transformation, the transformants having incorporated the expression construct are selected and regenerated into whole plants according to methods well-known in the art. Often the transformation procedure is designed for the selective elimination of selection genes either during regeneration or in the following generations by using, for example, co-transformation with two separate T-DNA constructs or site specific excision of the selection gene by a specific recombinase.

30 The present invention also relates to methods of producing a polypeptide of the present invention comprising: (a) cultivating a transgenic plant or a plant cell comprising a polynucleotide encoding the polypeptide having beta-glucosidase activity of the present invention under conditions conducive for production of the polypeptide; and (b) recovering the polypeptide.

35 In addition to direct transformation of a particular plant genotype with a construct prepared according to the present invention, transgenic plants may be made by crossing a plant having a construct of the present invention to a second plant lacking the construct. For example, a construct encoding a polypeptide having beta-glucosidase activity or a portion

thereof can be introduced into a particular plant variety by crossing, without the need for ever directly transforming a plant of that given variety. Therefore, the present invention not only encompasses a plant directly regenerated from cells which have been transformed in accordance with the present invention, but also the progeny of such plants. As used herein, 5 progeny may refer to the offspring of any generation of a parent plant prepared in accordance with the present invention. Such progeny may include a DNA construct prepared in accordance with the present invention, or a portion of a DNA construct prepared in accordance with the present invention. Crossing results in a transgene of the present invention being introduced into a plant line by cross pollinating a starting line with a donor 10 plant line that includes a transgene of the present invention. Non-limiting examples of such steps are further articulated in U.S. Patent No: 7,151,204.

It is envisioned that plants including a polypeptide having beta-glucosidase activity of the present invention include plants generated through a process of backcross conversion. For examples, plants of the present invention include plants referred to as a backcross 15 converted genotype, line, inbred, or hybrid.

Genetic markers may be used to assist in the introgression of one or more transgenes of the invention from one genetic background into another. Marker assisted selection offers advantages relative to conventional breeding in that it can be used to avoid errors caused by phenotypic variations. Further, genetic markers may provide data 20 regarding the relative degree of elite germplasm in the individual progeny of a particular cross. For example, when a plant with a desired trait which otherwise has a non-agronomically desirable genetic background is crossed to an elite parent, genetic markers may be used to select progeny which not only possess the trait of interest, but also have a relatively large proportion of the desired germplasm. In this way, the number of generations 25 required to introgress one or more traits into a particular genetic background is minimized.

Removal or Reduction of Beta-Glucosidase Activity

The present invention also relates to methods of producing a mutant of a parent cell, which comprises disrupting or deleting a polynucleotide, or a portion thereof, encoding a 30 polypeptide of the present invention, which results in the mutant cell producing less of the polypeptide than the parent cell when cultivated under the same conditions.

The mutant cell may be constructed by reducing or eliminating expression of a nucleotide sequence encoding a polypeptide of the present invention using methods well known in the art, for example, insertions, disruptions, replacements, or deletions. In a 35 preferred aspect, the nucleotide sequence is inactivated. The nucleotide sequence to be modified or inactivated may be, for example, the coding region or a part thereof essential for activity, or a regulatory element required for the expression of the coding region. An

example of such a regulatory or control sequence may be a promoter sequence or a functional part thereof, *i.e.*, a part that is sufficient for affecting expression of the nucleotide sequence. Other control sequences for possible modification include, but are not limited to, a leader, polyadenylation sequence, propeptide sequence, signal peptide sequence, transcription terminator, and transcriptional activator.

Modification or inactivation of the nucleotide sequence may be performed by subjecting the parent cell to mutagenesis and selecting for mutant cells in which expression of the nucleotide sequence has been reduced or eliminated. The mutagenesis, which may be specific or random, may be performed, for example, by use of a suitable physical or chemical mutagenizing agent, by use of a suitable oligonucleotide, or by subjecting the DNA sequence to PCR generated mutagenesis. Furthermore, the mutagenesis may be performed by use of any combination of these mutagenizing agents.

Examples of a physical or chemical mutagenizing agent suitable for the present purpose include ultraviolet (UV) irradiation, hydroxylamine, N-methyl-N'-nitro-N-nitrosoguanidine (MNNG), O-methyl hydroxylamine, nitrous acid, ethyl methane sulphonate (EMS), sodium bisulphite, formic acid, and nucleotide analogues.

When such agents are used, the mutagenesis is typically performed by incubating the parent cell to be mutagenized in the presence of the mutagenizing agent of choice under suitable conditions, and screening and/or selecting for mutant cells exhibiting reduced or no expression of the gene.

Modification or inactivation of the nucleotide sequence may be accomplished by introduction, substitution, or removal of one or more (several) nucleotides in the gene or a regulatory element required for the transcription or translation thereof. For example, nucleotides may be inserted or removed so as to result in the introduction of a stop codon, the removal of the start codon, or a change in the open reading frame. Such modification or inactivation may be accomplished by site-directed mutagenesis or PCR generated mutagenesis in accordance with methods known in the art. Although, in principle, the modification may be performed *in vivo*, *i.e.*, directly on the cell expressing the nucleotide sequence to be modified, it is preferred that the modification be performed *in vitro* as exemplified below.

An example of a convenient way to eliminate or reduce expression of a nucleotide sequence by a cell is based on techniques of gene replacement, gene deletion, or gene disruption. For example, in the gene disruption method, a nucleic acid sequence corresponding to the endogenous nucleotide sequence is mutagenized *in vitro* to produce a defective nucleic acid sequence that is then transformed into the parent cell to produce a defective gene. By homologous recombination, the defective nucleic acid sequence replaces the endogenous nucleotide sequence. It may be desirable that the defective

nucleotide sequence also encodes a marker that may be used for selection of transformants in which the nucleotide sequence has been modified or destroyed. In a particularly preferred aspect, the nucleotide sequence is disrupted with a selectable marker such as those described herein.

5 Alternatively, modification or inactivation of the nucleotide sequence may be performed by established anti-sense or RNAi techniques using a sequence complementary to the nucleotide sequence. More specifically, expression of the nucleotide sequence by a cell may be reduced or eliminated by introducing a sequence complementary to the nucleotide sequence of the gene that may be transcribed in the cell and is capable of
10 hybridizing to the mRNA produced in the cell. Under conditions allowing the complementary anti-sense nucleotide sequence to hybridize to the mRNA, the amount of protein translated is thus reduced or eliminated.

 The present invention further relates to a mutant cell of a parent cell that comprises a disruption or deletion of a nucleotide sequence encoding the polypeptide or a control
15 sequence thereof, which results in the mutant cell producing less of the polypeptide or no polypeptide compared to the parent cell.

 The polypeptide-deficient mutant cells so created are particularly useful as host cells for the expression of native and/or heterologous polypeptides. Therefore, the present invention further relates to methods of producing a native or heterologous polypeptide,
20 comprising: (a) cultivating the mutant cell under conditions conducive for production of the polypeptide; and (b) recovering the polypeptide. The term "heterologous polypeptides" is defined herein as polypeptides that are not native to the host cell, a native protein in which modifications have been made to alter the native sequence, or a native protein whose expression is quantitatively altered as a result of a manipulation of the host cell by
25 recombinant DNA techniques.

 In a further aspect, the present invention relates to a method of producing a protein product essentially free of beta-glucosidase activity by fermentation of a cell that produces both a polypeptide of the present invention as well as the protein product of interest by adding an effective amount of an agent capable of inhibiting beta-glucosidase activity to the
30 fermentation broth before, during, or after the fermentation has been completed, recovering the product of interest from the fermentation broth, and optionally subjecting the recovered product to further purification.

 In a further aspect, the present invention relates to a method of producing a protein product essentially free of beta-glucosidase activity by cultivating the cell under conditions
35 permitting the expression of the product, subjecting the resultant culture broth to a combined pH and temperature treatment so as to reduce the beta-glucosidase activity substantially, and recovering the product from the culture broth. Alternatively, the combined pH and

temperature treatment may be performed on an enzyme preparation recovered from the culture broth. The combined pH and temperature treatment may optionally be used in combination with a treatment with an beta-glucosidase inhibitor.

5 In accordance with this aspect of the invention, it is possible to remove at least 60%, preferably at least 75%, more preferably at least 85%, still more preferably at least 95%, and most preferably at least 99% of the beta-glucosidase activity. Complete removal of beta-glucosidase activity may be obtained by use of this method.

10 The combined pH and temperature treatment is preferably carried out at a pH in the range of 2-4 or 9-11 and a temperature in the range of at least 60-70°C for a sufficient period of time to attain the desired effect, where typically, 30 to 60 minutes is sufficient.

The methods used for cultivation and purification of the product of interest may be performed by methods known in the art.

15 The methods of the present invention for producing an essentially beta-glucosidase-free product is of particular interest in the production of eukaryotic polypeptides, in particular fungal proteins such as enzymes. The enzyme may be selected from, *e.g.*, an amylolytic enzyme, lipolytic enzyme, proteolytic enzyme, cellulolytic enzyme, oxidoreductase, or plant cell-wall degrading enzyme. Examples of such enzymes include an aminopeptidase, amylase, amyloglucosidase, carbohydrase, carboxypeptidase, catalase, cellobiohydrolase, cellulase, chitinase, cutinase, cyclodextrin glycosyltransferase, deoxyribonuclease, 20 endoglucanase, esterase, galactosidase, beta-galactosidase, glucoamylase, glucose oxidase, glucosidase, haloperoxidase, hemicellulase, invertase, isomerase, laccase, ligase, lipase, lyase, mannosidase, oxidase, pectinolytic enzyme, peroxidase, phytase, phenoxidase, polyphenoxidase, proteolytic enzyme, ribonuclease, transferase, transglutaminase, or xylanase. The beta-glucosidase-deficient cells may also be used to 25 express heterologous proteins of pharmaceutical interest such as hormones, growth factors, receptors, and the like.

30 It will be understood that the term "eukaryotic polypeptides" includes not only native polypeptides, but also those polypeptides, *e.g.*, enzymes, which have been modified by amino acid substitutions, deletions or additions, or other such modifications to enhance activity, thermostability, pH tolerance and the like.

In a further aspect, the present invention relates to a protein product essentially free from beta-glucosidase activity that is produced by a method of the present invention.

Methods of Inhibiting Expression of a Polypeptide Having Beta-Glucosidase Activity

35 The present invention also relates to methods of inhibiting the expression of a polypeptide having beta-glucosidase activity in a cell, comprising administering to the cell or

expressing in the cell a double-stranded RNA (dsRNA) molecule, wherein the dsRNA comprises a subsequence of a polynucleotide of the present invention. In a preferred aspect, the dsRNA is about 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25 or more duplex nucleotides in length.

5 The dsRNA is preferably a small interfering RNA (siRNA) or a micro RNA (miRNA). In a preferred aspect, the dsRNA is small interfering RNA (siRNAs) for inhibiting transcription. In another preferred aspect, the dsRNA is micro RNA (miRNAs) for inhibiting translation.

10 The present invention also relates to such double-stranded RNA (dsRNA) molecules, comprising a portion of the mature polypeptide coding sequence of SEQ ID NO: 1 for inhibiting expression of a polypeptide having beta-glucosidase activity in a cell. While the present invention is not limited by any particular mechanism of action, the dsRNA can enter a cell and cause the degradation of a single-stranded RNA (ssRNA) of similar or identical sequences, including endogenous mRNAs. When a cell is exposed to dsRNA, mRNA from the homologous gene is selectively degraded by a process called RNA interference (RNAi).

15 The dsRNAs of the present invention can be used in gene-silencing. In one aspect, the invention provides methods to selectively degrade RNA using the dsRNAs of the present invention. The process may be practiced *in vitro*, *ex vivo* or *in vivo*. In one aspect, the dsRNA molecules can be used to generate a loss-of-function mutation in a cell, an organ or an animal. Methods for making and using dsRNA molecules to selectively degrade RNA are well known in the art, see, for example, U.S. Patent No. 6,506,559; U.S. Patent No. 6,511,824; U.S. Patent No. 6,515,109; and U.S. Patent No. 6,489,127.

Compositions

25 The present invention also relates to compositions comprising a polypeptide of the present invention. Preferably, the compositions are enriched in such a polypeptide. The term "enriched" indicates that the beta-glucosidase activity of the composition has been increased, *e.g.*, with an enrichment factor of at least 1.1.

30 The composition may comprise a polypeptide of the present invention as the major enzymatic component, *e.g.*, a mono-component composition. Alternatively, the composition may comprise multiple enzymatic activities, such as an aminopeptidase, amylase, carbohydrase, carboxypeptidase, catalase, cellulase, chitinase, cutinase, cyclodextrin glycosyltransferase, deoxyribonuclease, esterase, alpha-galactosidase, beta-galactosidase, glucoamylase, alpha-glucosidase, beta-glucosidase, haloperoxidase, invertase, laccase, lipase, mannosidase, oxidase, pectinolytic enzyme, peptidoglutaminase, peroxidase, 35 phytase, polyphenoloxidase, proteolytic enzyme, ribonuclease, transglutaminase, or

xylanase. The additional enzyme(s) may be produced, for example, by a microorganism belonging to the genus *Aspergillus*, preferably *Aspergillus aculeatus*, *Aspergillus awamori*, *Aspergillus fumigatus*, *Aspergillus foetidus*, *Aspergillus japonicus*, *Aspergillus nidulans*, *Aspergillus niger*, or *Aspergillus oryzae*; *Fusarium*, preferably *Fusarium bactridioides*, *Fusarium cerealis*, *Fusarium crookwellense*, *Fusarium culmorum*, *Fusarium graminearum*, *Fusarium graminum*, *Fusarium heterosporum*, *Fusarium negundi*, *Fusarium oxysporum*, *Fusarium reticulatum*, *Fusarium roseum*, *Fusarium sambucinum*, *Fusarium sarcochroum*, *Fusarium sulphureum*, *Fusarium toruloseum*, *Fusarium trichothecioides*, or *Fusarium venenatum*; *Humicola*, preferably *Humicola insolens* or *Humicola lanuginosa*; or *Trichoderma*, preferably *Trichoderma harzianum*, *Trichoderma koningii*, *Trichoderma longibrachiatum*, *Trichoderma reesei*, or *Trichoderma viride*.

The polypeptide compositions may be prepared in accordance with methods known in the art and may be in the form of a liquid or a dry composition. For instance, the polypeptide composition may be in the form of a granulate or a microgranulate. The polypeptide to be included in the composition may be stabilized in accordance with methods known in the art.

Examples are given below of preferred uses of the polypeptide compositions of the invention. The dosage of the polypeptide composition of the invention and other conditions under which the composition is used may be determined on the basis of methods known in the art.

Processing of Cellulosic Material

The present invention also relates to methods for degrading or converting a cellulosic material, comprising: treating the cellulosic material with an enzyme composition in the presence of a polypeptide having beta-glucosidase activity of the present invention. In a preferred aspect, the method further comprises recovering the degraded or converted cellulosic material.

The present invention also relates to methods of producing a fermentation product, comprising: (a) saccharifying a cellulosic material with an enzyme composition in the presence of a polypeptide having beta-glucosidase activity of the present invention; (b) fermenting the saccharified cellulosic material with one or more (several) fermenting microorganisms to produce the fermentation product; and (c) recovering the fermentation product from the fermentation.

The present invention also relates to methods of fermenting a cellulosic material, comprising: fermenting the cellulosic material with one or more (several) fermenting microorganisms, wherein the cellulosic material is saccharified with an enzyme composition in the presence of a polypeptide having beta-glucosidase activity of the present invention. In

a preferred aspect, the fermenting of the cellulosic material produces a fermentation product. In another preferred aspect, the method further comprises recovering the fermentation product from the fermentation.

The methods of the present invention can be used to saccharify a cellulosic material to fermentable sugars and convert the fermentable sugars to many useful substances, e.g., fuel, potable ethanol, and/or fermentation products (e.g., acids, alcohols, ketones, gases, and the like). The production of a desired fermentation product from cellulosic material typically involves pretreatment, enzymatic hydrolysis (saccharification), and fermentation.

The processing of cellulosic material according to the present invention can be accomplished using processes conventional in the art. Moreover, the methods of the present invention can be implemented using any conventional biomass processing apparatus configured to operate in accordance with the invention.

Hydrolysis (saccharification) and fermentation, separate or simultaneous, include, but are not limited to, separate hydrolysis and fermentation (SHF); simultaneous saccharification and fermentation (SSF); simultaneous saccharification and cofermentation (SSCF); hybrid hydrolysis and fermentation (HHF); separate hydrolysis and co-fermentation (SHCF); hybrid hydrolysis and fermentation (HHCF); and direct microbial conversion (DMC). SHF uses separate process steps to first enzymatically hydrolyze cellulosic material to fermentable sugars, e.g., glucose, cellobiose, cellotriose, and pentose sugars, and then ferment the fermentable sugars to ethanol. In SSF, the enzymatic hydrolysis of cellulosic material and the fermentation of sugars to ethanol are combined in one step (Philippidis, G. P., 1996, Cellulose bioconversion technology, in *Handbook on Bioethanol: Production and Utilization*, Wyman, C. E., ed., Taylor & Francis, Washington, DC, 179-212). SSCF involves the cofermentation of multiple sugars (Sheehan, J., and Himmel, M., 1999, Enzymes, energy and the environment: A strategic perspective on the U.S. Department of Energy's research and development activities for bioethanol, *Biotechnol. Prog.* 15: 817-827). HHF involves a separate hydrolysis step, and in addition a simultaneous saccharification and hydrolysis step, which can be carried out in the same reactor. The steps in an HHF process can be carried out at different temperatures, i.e., high temperature enzymatic saccharification followed by SSF at a lower temperature that the fermentation strain can tolerate. DMC combines all three processes (enzyme production, hydrolysis, and fermentation) in one or more (several) steps where the same organism is used to produce the enzymes for conversion of the cellulosic material to fermentable sugars and to convert the fermentable sugars into a final product (Lynd, L. R., Weimer, P. J., van Zyl, W. H., and Pretorius, I. S., 2002, Microbial cellulose utilization: Fundamentals and biotechnology, *Microbiol. Mol. Biol. Reviews* 66: 506-577). It is understood herein that any method known in the art comprising pretreatment, enzymatic hydrolysis (saccharification), fermentation, or a combination thereof,

can be used in the practicing the methods of the present invention.

A conventional apparatus can include a fed-batch stirred reactor, a batch stirred reactor, a continuous flow stirred reactor with ultrafiltration, and/or a continuous plug-flow column reactor (Fernanda de Castilhos Corazza, Flávio Faria de Moraes, Gisella Maria Zanin and Ivo Neitzel, 2003, Optimal control in fed-batch reactor for the cellobiose hydrolysis, *Acta Scientiarum. Technology* 25: 33-38; Gusakov, A. V., and Sinitsyn, A. P., 1985, Kinetics of the enzymatic hydrolysis of cellulose: 1. A mathematical model for a batch reactor process, *Enz. Microb. Technol.* 7: 346-352), an attrition reactor (Ryu, S. K., and Lee, J. M., 1983, Bioconversion of waste cellulose by using an attrition bioreactor, *Biotechnol. Bioeng.* 25: 53-65), or a reactor with intensive stirring induced by an electromagnetic field (Gusakov, A. V., Sinitsyn, A. P., Davydkin, I. Y., Davydkin, V. Y., Protas, O. V., 1996, Enhancement of enzymatic cellulose hydrolysis using a novel type of bioreactor with intensive stirring induced by electromagnetic field, *Appl. Biochem. Biotechnol.* 56: 141-153). Additional reactor types include: fluidized bed, upflow blanket, immobilized, and extruder type reactors for hydrolysis and/or fermentation.

Pretreatment. In practicing the methods of the present invention, any pretreatment process known in the art can be used to disrupt plant cell wall components of cellulosic material (Chandra *et al.*, 2007, Substrate pretreatment: The key to effective enzymatic hydrolysis of lignocellulosics? *Adv. Biochem. Engin./Biotechnol.* 108: 67-93; Galbe and Zacchi, 2007, Pretreatment of lignocellulosic materials for efficient bioethanol production, *Adv. Biochem. Engin. / Biotechnol.* 108: 41-65; Hendriks and Zeeman, 2009, Pretreatments to enhance the digestibility of lignocellulosic biomass, *Bioresource Technol.* 100: 10-18; Mosier *et al.*, 2005, Features of promising technologies for pretreatment of lignocellulosic biomass, *Bioresource Technol.* 96: 673-686; Taherzadeh and Karimi, 2008, Pretreatment of lignocellulosic wastes to improve ethanol and biogas production: A review, *Int. J. of Mol. Sci.* 9: 1621-1651; Yang and Wyman, 2008, Pretreatment: the key to unlocking low-cost cellulosic ethanol, *Biofuels Bioproducts and Biorefining-Biofpr.* 2: 26-40).

The cellulosic material can also be subjected to particle size reduction, pre-soaking, wetting, washing, or conditioning prior to pretreatment using methods known in the art.

Conventional pretreatments include, but are not limited to, steam pretreatment (with or without explosion), dilute acid pretreatment, hot water pretreatment, alkaline pretreatment, lime pretreatment, wet oxidation, wet explosion, ammonia fiber explosion, organosolv pretreatment, and biological pretreatment. Additional pretreatments include ammonia percolation, ultrasound, electroporation, microwave, supercritical CO₂, supercritical H₂O, ozone, and gamma irradiation pretreatments.

The cellulosic material can be pretreated before hydrolysis and/or fermentation. Pretreatment is preferably performed prior to the hydrolysis. Alternatively, the pretreatment can

be carried out simultaneously with enzyme hydrolysis to release fermentable sugars, such as glucose, xylose, and/or cellobiose. In most cases the pretreatment step itself results in some conversion of biomass to fermentable sugars (even in absence of enzymes).

5 Steam Pretreatment. In steam pretreatment, cellulosic material is heated to disrupt the plant cell wall components, including lignin, hemicellulose, and cellulose to make the cellulose and other fractions, e.g., hemicellulose, accessible to enzymes. Cellulosic material is passed to or through a reaction vessel where steam is injected to increase the temperature to the required temperature and pressure and is retained therein for the desired reaction time. Steam pretreatment is preferably done at 140-230°C, more preferably 160-10 200°C, and most preferably 170-190°C, where the optimal temperature range depends on any addition of a chemical catalyst. Residence time for the steam pretreatment is preferably 1-15 minutes, more preferably 3-12 minutes, and most preferably 4-10 minutes, where the optimal residence time depends on temperature range and any addition of a chemical catalyst. Steam pretreatment allows for relatively high solids loadings, so that cellulosic material is generally only moist during the pretreatment. The steam pretreatment is often 15 combined with an explosive discharge of the material after the pretreatment, which is known as steam explosion, that is, rapid flashing to atmospheric pressure and turbulent flow of the material to increase the accessible surface area by fragmentation (Duff and Murray, 1996, *Bioresource Technology* 855: 1-33; Galbe and Zacchi, 2002, *Appl. Microbiol. Biotechnol.* 59: 20 618-628; U.S. Patent Application No. 20020164730). During steam pretreatment, hemicellulose acetyl groups are cleaved and the resulting acid autocatalyzes partial hydrolysis of the hemicellulose to monosaccharides and oligosaccharides. Lignin is removed to only a limited extent.

A catalyst such as H₂SO₄ or SO₂ (typically 0.3 to 3% w/w) is often added prior to 25 steam pretreatment, which decreases the time and temperature, increases the recovery, and improves enzymatic hydrolysis (Ballesteros *et al.*, 2006, *Appl. Biochem. Biotechnol.* 129-132: 496-508; Varga *et al.*, 2004, *Appl. Biochem. Biotechnol.* 113-116: 509-523; Sassner *et al.*, 2006, *Enzyme Microb. Technol.* 39: 756-762).

30 Chemical Pretreatment: The term "chemical treatment" refers to any chemical pretreatment that promotes the separation and/or release of cellulose, hemicellulose, and/or lignin. Examples of suitable chemical pretreatment processes include, for example, dilute acid pretreatment, lime pretreatment, wet oxidation, ammonia fiber/freeze explosion (AFEX), ammonia percolation (APR), and organosolv pretreatments.

In dilute acid pretreatment, cellulosic material is mixed with dilute acid, typically H₂SO₄, 35 and water to form a slurry, heated by steam to the desired temperature, and after a residence time flashed to atmospheric pressure. The dilute acid pretreatment can be performed with a number of reactor designs, e.g., plug-flow reactors, counter-current reactors,

or continuous counter-current shrinking bed reactors (Duff and Murray, 1996, *supra*; Schell *et al.*, 2004, *Bioresource Technol.* 91: 179-188; Lee *et al.*, 1999, *Adv. Biochem. Eng. Biotechnol.* 65: 93-115).

Several methods of pretreatment under alkaline conditions can also be used. These alkaline pretreatments include, but are not limited to, lime pretreatment, wet oxidation, ammonia percolation (APR), and ammonia fiber/freeze explosion (AFEX).

Lime pretreatment is performed with calcium carbonate, sodium hydroxide, or ammonia at low temperatures of 85-150°C and residence times from 1 hour to several days (Wyman *et al.*, 2005, *Bioresource Technol.* 96: 1959-1966; Mosier *et al.*, 2005, *Bioresource Technol.* 96: 673-686). WO 2006/110891, WO 2006/11899, WO 2006/11900, and WO 2006/110901 disclose pretreatment methods using ammonia.

Wet oxidation is a thermal pretreatment performed typically at 180-200°C for 5-15 minutes with addition of an oxidative agent such as hydrogen peroxide or over-pressure of oxygen (Schmidt and Thomsen, 1998, *Bioresource Technol.* 64: 139-151; Palonen *et al.*, 2004, *Appl. Biochem. Biotechnol.* 117: 1-17; Varga *et al.*, 2004, *Biotechnol. Bioeng.* 88: 567-574; Martin *et al.*, 2006, *J. Chem. Technol. Biotechnol.* 81: 1669-1677). The pretreatment is performed at preferably 1-40% dry matter, more preferably 2-30% dry matter, and most preferably 5-20% dry matter, and often the initial pH is increased by the addition of alkali such as sodium carbonate.

A modification of the wet oxidation pretreatment method, known as wet explosion (combination of wet oxidation and steam explosion), can handle dry matter up to 30%. In wet explosion, the oxidizing agent is introduced during pretreatment after a certain residence time. The pretreatment is then ended by flashing to atmospheric pressure (WO 2006/032282).

Ammonia fiber explosion (AFEX) involves treating cellulosic material with liquid or gaseous ammonia at moderate temperatures such as 90-100°C and high pressure such as 17-20 bar for 5-10 minutes, where the dry matter content can be as high as 60% (Gollapalli *et al.*, 2002, *Appl. Biochem. Biotechnol.* 98: 23-35; Chundawat *et al.*, 2007, *Biotechnol. Bioeng.* 96: 219-231; Alizadeh *et al.*, 2005, *Appl. Biochem. Biotechnol.* 121: 1133-1141; Teymouri *et al.*, 2005, *Bioresource Technol.* 96: 2014-2018). AFEX pretreatment results in the depolymerization of cellulose and partial hydrolysis of hemicellulose. Lignin-carbohydrate complexes are cleaved.

Organosolv pretreatment delignifies cellulosic material by extraction using aqueous ethanol (40-60% ethanol) at 160-200°C for 30-60 minutes (Pan *et al.*, 2005, *Biotechnol. Bioeng.* 90: 473-481; Pan *et al.*, 2006, *Biotechnol. Bioeng.* 94: 851-861; Kurabi *et al.*, 2005, *Appl. Biochem. Biotechnol.* 121: 219-230). Sulphuric acid is usually added as a catalyst. In organosolv pretreatment, the majority of hemicellulose is removed.

Other examples of suitable pretreatment methods are described by Schell *et al.*, 2003,

Appl. Biochem. and Biotechnol. Vol. 105-108, p. 69-85, and Mosier *et al.*, 2005, *Bioresource Technology* 96: 673-686, and U.S. Published Application 2002/0164730.

In one aspect, the chemical pretreatment is preferably carried out as an acid treatment, and more preferably as a continuous dilute and/or mild acid treatment. The acid is typically sulfuric acid, but other acids can also be used, such as acetic acid, citric acid, nitric acid, phosphoric acid, tartaric acid, succinic acid, hydrogen chloride, or mixtures thereof. Mild acid treatment is conducted in the pH range of preferably 1-5, more preferably 1-4, and most preferably 1-3. In one aspect, the acid concentration is in the range from preferably 0.01 to 20 wt % acid, more preferably 0.05 to 10 wt % acid, even more preferably 0.1 to 5 wt % acid, and most preferably 0.2 to 2.0 wt % acid. The acid is contacted with cellulosic material and held at a temperature in the range of preferably 160-220°C, and more preferably 165-195°C, for periods ranging from seconds to minutes to, *e.g.*, 1 second to 60 minutes.

In another aspect, pretreatment is carried out as an ammonia fiber explosion step (AFEX pretreatment step).

In another aspect, pretreatment takes place in an aqueous slurry. In preferred aspects, cellulosic material is present during pretreatment in amounts preferably between 10-80 wt%, more preferably between 20-70 wt%, and most preferably between 30-60 wt%, such as around 50 wt%. The pretreated cellulosic material can be unwashed or washed using any method known in the art, *e.g.*, washed with water.

Mechanical Pretreatment: The term "mechanical pretreatment" refers to various types of grinding or milling (*e.g.*, dry milling, wet milling, or vibratory ball milling).

Physical Pretreatment: The term "physical pretreatment" refers to any pretreatment that promotes the separation and/or release of cellulose, hemicellulose, and/or lignin from cellulosic material. For example, physical pretreatment can involve irradiation (*e.g.*, microwave irradiation), steaming/steam explosion, hydrothermolysis, and combinations thereof.

Physical pretreatment can involve high pressure and/or high temperature (steam explosion). In one aspect, high pressure means pressure in the range of preferably about 300 to about 600 psi, more preferably about 350 to about 550 psi, and most preferably about 400 to about 500 psi, such as around 450 psi. In another aspect, high temperature means temperatures in the range of about 100 to about 300°C, preferably about 140 to about 235°C. In a preferred aspect, mechanical pretreatment is performed in a batch-process, steam gun hydrolyzer system that uses high pressure and high temperature as defined above, *e.g.*, a Sunds Hydrolyzer available from Sunds Defibrator AB, Sweden.

Combined Physical and Chemical Pretreatment: Cellulosic material can be pretreated both physically and chemically. For instance, the pretreatment step can involve dilute or mild acid treatment and high temperature and/or pressure treatment. The physical and chemical pretreatments can be carried out sequentially or simultaneously, as desired. A mechanical

pretreatment can also be included.

Accordingly, in a preferred aspect, cellulosic material is subjected to mechanical, chemical, or physical pretreatment, or any combination thereof, to promote the separation and/or release of cellulose, hemicellulose, and/or lignin.

5 Biological Pretreatment: The term "biological pretreatment" refers to any biological pretreatment that promotes the separation and/or release of cellulose, hemicellulose, and/or lignin from cellulosic material. Biological pretreatment techniques can involve applying lignin-solubilizing microorganisms (see, for example, Hsu, T.-A., 1996, Pretreatment of biomass, in *Handbook on Bioethanol: Production and Utilization*, Wyman, C. E., ed., Taylor & Francis, Washington, DC, 179-212; Ghosh and Singh, 1993, Physicochemical and biological treatments for enzymatic/microbial conversion of cellulosic biomass, *Adv. Appl. Microbiol.* 39: 295-333; McMillan, J. D., 1994, Pretreating lignocellulosic biomass: a review, in *Enzymatic Conversion of Biomass for Fuels Production*, Himmel, M. E., Baker, J. O., and Overend, R. P., eds., ACS Symposium Series 566, American Chemical Society, 10 Washington, DC, chapter 15; Gong, C. S., Cao, N. J., Du, J., and Tsao, G. T., 1999, Ethanol production from renewable resources, in *Advances in Biochemical Engineering/Biotechnology*, Scheper, T., ed., Springer-Verlag Berlin Heidelberg, Germany, 65: 207-241; Olsson and Hahn-Hagerdal, 1996, Fermentation of lignocellulosic hydrolysates for ethanol production, *Enz. Microb. Tech.* 18: 312-331; and Vallander and Eriksson, 1990, Production of ethanol from lignocellulosic materials: State of the art, *Adv. Biochem. Eng./Biotechnol.* 42: 63-95).

Saccharification. In the hydrolysis step, also known as saccharification, the cellulosic material, e.g., pretreated, is hydrolyzed to break down cellulose and alternatively also hemicellulose to fermentable sugars, such as glucose, cellobiose, xylose, xylulose, 25 arabinose, mannose, galactose, and/or soluble oligosaccharides. The hydrolysis is performed enzymatically by an enzyme composition in the presence of a polypeptide having beta-glucosidase activity of the present invention. The composition can further comprise one or more (several) hemicellulolytic enzymes. The enzymes of the compositions can also be added sequentially.

30 Enzymatic hydrolysis is preferably carried out in a suitable aqueous environment under conditions that can be readily determined by one skilled in the art. In a preferred aspect, hydrolysis is performed under conditions suitable for the activity of the enzyme(s), i.e., optimal for the enzyme(s). The hydrolysis can be carried out as a fed batch or continuous process where the pretreated cellulosic material (substrate) is fed gradually to, for example, an enzyme 35 containing hydrolysis solution.

The saccharification is generally performed in stirred-tank reactors or fermentors under controlled pH, temperature, and mixing conditions. Suitable process time, temperature

and pH conditions can readily be determined by one skilled in the art. For example, the saccharification can last up to 200 hours, but is typically performed for preferably about 12 to about 96 hours, more preferably about 16 to about 72 hours, and most preferably about 24 to about 48 hours. The temperature is in the range of preferably about 25°C to about 70°C, more preferably about 30°C to about 65°C, and more preferably about 40°C to 60°C, in particular about 50°C. The pH is in the range of preferably about 3 to about 8, more preferably about 3.5 to about 7, and most preferably about 4 to about 6, in particular about pH 5. The dry solids content is in the range of preferably about 5 to about 50 wt %, more preferably about 10 to about 40 wt %, and most preferably about 20 to about 30 wt %.

The enzyme composition preferably comprises enzymes having cellulolytic activity and/or xylan degrading activity. In one aspect, the enzyme composition comprises one or more (several) cellulolytic enzymes. In another aspect, the enzyme composition comprises one or more (several) xylan degrading enzymes. In another aspect, the enzyme composition comprises one or more (several) cellulolytic enzymes and one or more (several) xylan degrading enzymes.

The one or more (several) cellulolytic enzymes are preferably selected from the group consisting of an endoglucanase, a cellobiohydrolase, and a beta-glucosidase. The one or more (several) xylan degrading enzymes are preferably selected from the group consisting of a xylanase, an acetyxylan esterase, a feruloyl esterase, an arabinofuranosidase, a xylosidase, and a glucuronidase.

In another aspect, the enzyme composition further or even further comprises a polypeptide having cellulolytic enhancing activity (see, for example, WO 2005/074647, WO 2005/074656, and WO 2007/089290). In another aspect, the enzyme composition may further or even further comprise one or more (several) additional enzyme activities to improve the degradation of the cellulose-containing material. Preferred additional enzymes are hemicellulases (e.g., alpha-D-glucuronidases, alpha-L-arabinofuranosidases, endo-mannanases, beta-mannosidases, alpha-galactosidases, endo-alpha-L-arabinanases, beta-galactosidases), carbohydrate-esterases (e.g., acetyl-xylan esterases, acetyl-mannan esterases, ferulic acid esterases, coumaric acid esterases, glucuronoyl esterases), pectinases, proteases, ligninolytic enzymes (e.g., laccases, manganese peroxidases, lignin peroxidases, H₂O₂-producing enzymes, oxidoreductases), expansins, swollenins, or mixtures thereof. In the methods of the present invention, the additional enzyme(s) can be added prior to or during fermentation, e.g., during saccharification or during or after propagation of the fermenting microorganism(s).

One or more (several) components of the enzyme composition may be wild-type proteins, recombinant proteins, or a combination of wild-type proteins and recombinant proteins. For example, one or more (several) components may be native proteins of a cell,

which is used as a host cell to express recombinantly one or more (several) other components of the enzyme composition. One or more (several) components of the enzyme composition may be produced as monocomponents, which are then combined to form the enzyme composition. The enzyme composition may be a combination of multicomponent and monocomponent protein preparations.

The enzymes used in the methods of the present invention may be in any form suitable for use in the processes described herein, such as, for example, a crude fermentation broth with or without cells removed, a cell lysate with or without cellular debris, a semi-purified or purified enzyme preparation, or a host cell as a source of the enzymes. The enzyme composition may be a dry powder or granulate, a non-dusting granulate, a liquid, a stabilized liquid, or a stabilized protected enzyme. Liquid enzyme preparations may, for instance, be stabilized by adding stabilizers such as a sugar, a sugar alcohol or another polyol, and/or lactic acid or another organic acid according to established processes.

The optimum amounts of the enzymes and polypeptides having beta-glucosidase activity depend on several factors including, but not limited to, the mixture of component cellulolytic enzymes, the cellulosic substrate, the concentration of cellulosic substrate, the pretreatment(s) of the cellulosic substrate, temperature, time, pH, and inclusion of fermenting organism (e.g., yeast for Simultaneous Saccharification and Fermentation).

In a preferred aspect, an effective amount of cellulolytic enzyme(s) to cellulosic material is about 0.5 to about 50 mg, preferably at about 0.5 to about 40 mg, more preferably at about 0.5 to about 25 mg, more preferably at about 0.75 to about 20 mg, more preferably at about 0.75 to about 15 mg, even more preferably at about 0.5 to about 10 mg, and most preferably at about 2.5 to about 10 mg per g of cellulosic material.

In another preferred aspect, an effective amount of polypeptide(s) having beta-glucosidase activity to cellulosic material is about 0.01 to about 50.0 mg, preferably about 0.01 to about 40 mg, more preferably about 0.01 to about 30 mg, more preferably about 0.01 to about 20 mg, more preferably about 0.01 to about 10 mg, more preferably about 0.01 to about 5 mg, more preferably at about 0.025 to about 1.5 mg, more preferably at about 0.05 to about 1.25 mg, more preferably at about 0.075 to about 1.25 mg, more preferably at about 0.1 to about 1.25 mg, even more preferably at about 0.15 to about 1.25 mg, and most preferably at about 0.25 to about 1.0 mg per g of cellulosic material.

In another preferred aspect, an effective amount of polypeptide(s) having beta-glucosidase activity to cellulolytic enzyme(s) is about 0.005 to about 1.0 g, preferably at about 0.01 to about 1.0 g, more preferably at about 0.15 to about 0.75 g, more preferably at about 0.15 to about 0.5 g, more preferably at about 0.1 to about 0.5 g, even more preferably at about 0.1 to about 0.5 g, and most preferably at about 0.05 to about 0.2 g per g of cellulolytic enzyme(s).

The enzymes can be derived or obtained from any suitable origin, including, bacterial, fungal, yeast, plant, or mammalian origin. The term "obtained" means herein that the enzyme may have been isolated from an organism that naturally produces the enzyme as a native enzyme. The term "obtained" also means herein that the enzyme may have been produced recombinantly in a host organism employing methods described herein, wherein the recombinantly produced enzyme is either native or foreign to the host organism or has a modified amino acid sequence, e.g., having one or more (several) amino acids that are deleted, inserted and/or substituted, i.e., a recombinantly produced enzyme that is a mutant and/or a fragment of a native amino acid sequence or an enzyme produced by nucleic acid shuffling processes known in the art. Encompassed within the meaning of a native enzyme are natural variants and within the meaning of a foreign enzyme are variants obtained recombinantly, such as by site-directed mutagenesis or shuffling.

A polypeptide having cellulolytic enzyme activity or xylan degrading activity may be a bacterial polypeptide. For example, the polypeptide may be a gram positive bacterial polypeptide such as a *Bacillus*, *Streptococcus*, *Streptomyces*, *Staphylococcus*, *Enterococcus*, *Lactobacillus*, *Lactococcus*, *Clostridium*, *Geobacillus*, or *Oceanobacillus* polypeptide having cellulolytic enzyme activity or xylan degrading activity, or a Gram negative bacterial polypeptide such as an *E. coli*, *Pseudomonas*, *Salmonella*, *Campylobacter*, *Helicobacter*, *Flavobacterium*, *Fusobacterium*, *Ilyobacter*, *Neisseria*, or *Ureaplasma* polypeptide having cellulolytic enzyme activity or xylan degrading activity.

In a preferred aspect, the polypeptide is a *Bacillus alkalophilus*, *Bacillus amyloliquefaciens*, *Bacillus brevis*, *Bacillus circulans*, *Bacillus clausii*, *Bacillus coagulans*, *Bacillus firmus*, *Bacillus lautus*, *Bacillus lentus*, *Bacillus licheniformis*, *Bacillus megaterium*, *Bacillus pumilus*, *Bacillus stearothermophilus*, *Bacillus subtilis*, or *Bacillus thuringiensis* polypeptide having cellulolytic enzyme activity or xylan degrading activity.

In another preferred aspect, the polypeptide is a *Streptococcus equisimilis*, *Streptococcus pyogenes*, *Streptococcus uberis*, or *Streptococcus equi* subsp. *Zooepidemicus* polypeptide having cellulolytic enzyme activity or xylan degrading activity.

In another preferred aspect, the polypeptide is a *Streptomyces achromogenes*, *Streptomyces avermitilis*, *Streptomyces coelicolor*, *Streptomyces griseus*, or *Streptomyces lividans* polypeptide having cellulolytic enzyme activity or xylan degrading activity.

The polypeptide having cellulolytic enzyme activity or xylan degrading activity may also be a fungal polypeptide, and more preferably a yeast polypeptide such as a *Candida*, *Kluyveromyces*, *Pichia*, *Saccharomyces*, *Schizosaccharomyces*, or *Yarrowia* polypeptide having cellulolytic enzyme activity or xylan degrading activity; or more preferably a filamentous fungal polypeptide such as an *Acremonium*, *Agaricus*, *Alternaria*, *Aspergillus*, *Aureobasidium*, *Botryosphaeria*, *Ceriporiopsis*, *Chaetomidium*, *Chrysosporium*, *Claviceps*,

Cochliobolus, *Coprinopsis*, *Coptotermes*, *Corynascus*, *Cryphonectria*, *Cryptococcus*, *Diplodia*, *Exidia*, *Filibasidium*, *Fusarium*, *Gibberella*, *Holomastigotoides*, *Humicola*, *Irpex*, *Lentinula*, *Leptosphaeria*, *Magnaporthe*, *Melanocarpus*, *Meripilus*, *Mucor*, *Myceliophthora*, *Neocallimastix*, *Neurospora*, *Paecilomyces*, *Penicillium*, *Phanerochaete*, *Piromyces*,
5 *Poitrasia*, *Pseudoplectania*, *Pseudotriconympha*, *Rhizomucor*, *Schizophyllum*, *Scytalidium*,
Talaromyces, *Thermoascus*, *Thielavia*, *Tolypocladium*, *Trichoderma*, *Trichophaea*,
Verticillium, *Volvariella*, or *Xylaria* polypeptide having cellulolytic enzyme activity or xylan
degrading activity.

In a preferred aspect, the polypeptide is a *Saccharomyces carlsbergensis*,
10 *Saccharomyces cerevisiae*, *Saccharomyces diastaticus*, *Saccharomyces douglasii*,
Saccharomyces kluyveri, *Saccharomyces norbensis*, or *Saccharomyces oviformis*
polypeptide having cellulolytic enzyme activity or xylan degrading activity.

In another preferred aspect, the polypeptide is an *Acremonium cellulolyticus*,
Aspergillus aculeatus, *Aspergillus awamori*, *Aspergillus fumigatus*, *Aspergillus foetidus*,
15 *Aspergillus japonicus*, *Aspergillus nidulans*, *Aspergillus niger*, *Aspergillus oryzae*,
Chrysosporium keratinophilum, *Chrysosporium lucknowense*, *Chrysosporium tropicum*,
Chrysosporium merdarium, *Chrysosporium inops*, *Chrysosporium pannicola*, *Chrysosporium*
queenslandicum, *Chrysosporium zonatum*, *Fusarium bactridioides*, *Fusarium cerealis*,
Fusarium crookwellense, *Fusarium culmorum*, *Fusarium graminearum*, *Fusarium graminum*,
20 *Fusarium heterosporum*, *Fusarium negundi*, *Fusarium oxysporum*, *Fusarium reticulatum*,
Fusarium roseum, *Fusarium sambucinum*, *Fusarium sarcochromum*, *Fusarium*
sporotrichioides, *Fusarium sulphureum*, *Fusarium torulosum*, *Fusarium trichothecioides*,
Fusarium venenatum, *Humicola grisea*, *Humicola insolens*, *Humicola lanuginosa*, *Irpex*
lacteus, *Mucor miehei*, *Myceliophthora thermophila*, *Neurospora crassa*, *Penicillium*
25 *funiculosum*, *Penicillium purpurogenum*, *Phanerochaete chrysosporium*, *Thielavia*
achromatica, *Thielavia albomyces*, *Thielavia albopilosa*, *Thielavia australeinsis*, *Thielavia*
fimeti, *Thielavia microspora*, *Thielavia ovispora*, *Thielavia peruviana*, *Thielavia*
spededonium, *Thielavia setosa*, *Thielavia subthermophila*, *Thielavia terrestris*, *Trichoderma*
harzianum, *Trichoderma koningii*, *Trichoderma longibrachiatum*, *Trichoderma reesei*,
30 *Trichoderma viride*, or *Trichophaea saccata* polypeptide having cellulolytic enzyme activity or
xylan degrading activity.

Chemically modified or protein engineered mutants of polypeptides having cellulolytic
enzyme activity or xylan degrading activity may also be used.

One or more (several) components of the enzyme composition may be a
35 recombinant component, *i.e.*, produced by cloning of a DNA sequence encoding the single
component and subsequent cell transformed with the DNA sequence and expressed in a
host (see, for example, WO 91/17243 and WO 91/17244). The host is preferably a

heterologous host (enzyme is foreign to host), but the host may under certain conditions also be a homologous host (enzyme is native to host). Monocomponent cellulolytic proteins may also be prepared by purifying such a protein from a fermentation broth.

5 Examples of commercial cellulolytic protein preparations suitable for use in the present invention include, for example, CELLIC™ Ctec (Novozymes A/S), CELLUCLAST™ (Novozymes A/S), NOVOZYM™ 188 (Novozymes A/S), CELLUZYME™ (Novozymes A/S), CEREFLO™ (Novozymes A/S), and ULTRAFLO™ (Novozymes A/S), ACCELERASE™ (Genencor Int.), LAMINEX™ (Genencor Int.), SPEZYME™ CP (Genencor Int.), ROHAMENT™ 7069 W (Röhm GmbH), FIBREZYME® LDI (Dyadic International, Inc.),
10 FIBREZYME® LBR (Dyadic International, Inc.), or VISCOSTAR® 150L (Dyadic International, Inc.). The cellulase enzymes are added in amounts effective from about 0.001 to about 5.0 wt % of solids, more preferably from about 0.025 to about 4.0 wt % of solids, and most preferably from about 0.005 to about 2.0 wt % of solids. The cellulase enzymes are added in amounts effective from about 0.001 to about 5.0 wt % of solids, more preferably
15 from about 0.025 to about 4.0 wt % of solids, and most preferably from about 0.005 to about 2.0 wt % of solids.

Examples of bacterial endoglucanases that can be used in the methods of the present invention, include, but are not limited to, an *Acidothermus cellulolyticus* endoglucanase (WO 91/05039; WO 93/15186; U.S. Patent No. 5,275,944; WO 96/02551; U.S. Patent No. 5,536,655, WO 00/70031, WO 05/093050); *Thermobifida fusca* endoglucanase III (WO 05/093050); and *Thermobifida fusca* endoglucanase V (WO 05/093050).

Examples of fungal endoglucanases that can be used in the methods of the present invention, include, but are not limited to, a *Trichoderma reesei* endoglucanase I (Penttila *et al.*, 1986, *Gene* 45: 253-263; GENBANK™ accession no. M15665); *Trichoderma reesei* endoglucanase II (Saloheimo, *et al.*, 1988, *Gene* 63:11-22; GENBANK™ accession no. M19373); *Trichoderma reesei* endoglucanase III (Okada *et al.*, 1988, *Appl. Environ. Microbiol.* 64: 555-563; GENBANK™ accession no. AB003694); *Trichoderma reesei* endoglucanase IV (Saloheimo *et al.*, 1997, *Eur. J. Biochem.* 249: 584-591; GENBANK™ accession no. Y111113); and *Trichoderma reesei* endoglucanase V (Saloheimo *et al.*, 1994, *Molecular Microbiology* 13: 219-228; GENBANK™ accession no. Z33381); *Aspergillus aculeatus* endoglucanase (Ooi *et al.*, 1990, *Nucleic Acids Research* 18: 5884); *Aspergillus kawachii* endoglucanase (Sakamoto *et al.*, 1995, *Current Genetics* 27: 435-439); *Erwinia carotovora* endoglucanase (Saarilahti *et al.*, 1990, *Gene* 90: 9-14); *Fusarium oxysporum* endoglucanase (GENBANK™ accession no. L29381); *Humicola grisea* var. *thermoidea* endoglucanase (GENBANK™ accession no. AB003107); *Melanocarpus albomyces*
25
30
35

endoglucanase (GENBANK™ accession no. MAL515703); *Neurospora crassa* endoglucanase (GENBANK™ accession no. XM_324477); *Humicola insolens* endoglucanase V (SEQ ID NO: 4); *Myceliophthora thermophila* CBS 117.65 endoglucanase (SEQ ID NO: 6); basidiomycete CBS 495.95 endoglucanase (SEQ ID NO: 8); basidiomycete CBS 494.95 endoglucanase (SEQ ID NO: 10); *Thielavia terrestris* NRRL 8126 CEL6B endoglucanase (SEQ ID NO: 12); *Thielavia terrestris* NRRL 8126 CEL6C endoglucanase (SEQ ID NO: 14); *Thielavia terrestris* NRRL 8126 CEL7C endoglucanase (SEQ ID NO: 16); *Thielavia terrestris* NRRL 8126 CEL7E endoglucanase (SEQ ID NO: 18); *Thielavia terrestris* NRRL 8126 CEL7F endoglucanase (SEQ ID NO: 20); *Cladorrhinum foecundissimum* ATCC 62373 CEL7A endoglucanase (SEQ ID NO: 22); and *Trichoderma reesei* strain No. VTT-D-80133 endoglucanase (SEQ ID NO: 24; GENBANK™ accession no. M15665). The endoglucanases of SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 14, SEQ ID NO: 16, SEQ ID NO: 18, SEQ ID NO: 20, SEQ ID NO: 22, and SEQ ID NO: 24 described above are encoded by the mature polypeptide coding sequence of SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21, and SEQ ID NO: 23, respectively.

Examples of cellobiohydrolases useful in the methods of the present invention include, but are not limited to, *Trichoderma reesei* cellobiohydrolase I (SEQ ID NO: 26); *Trichoderma reesei* cellobiohydrolase II (SEQ ID NO: 28); *Humicola insolens* cellobiohydrolase I (SEQ ID NO: 30), *Myceliophthora thermophila* cellobiohydrolase II (SEQ ID NO: 32 and SEQ ID NO: 34), *Thielavia terrestris* cellobiohydrolase II (CEL6A) (SEQ ID NO: 36), *Chaetomium thermophilum* cellobiohydrolase I (SEQ ID NO: 38), and *Chaetomium thermophilum* cellobiohydrolase II (SEQ ID NO: 40). The cellobiohydrolases of SEQ ID NO: 24, SEQ ID NO: 26, SEQ ID NO: 28, SEQ ID NO: 30, SEQ ID NO: 32, SEQ ID NO: 34, SEQ ID NO: 36, SEQ ID NO: 38, and SEQ ID NO: 40 described above are encoded by the mature polypeptide coding sequence of SEQ ID NO: 23, SEQ ID NO: 25, SEQ ID NO: 27, SEQ ID NO: 29, SEQ ID NO: 31, SEQ ID NO: 33, SEQ ID NO: 35, SEQ ID NO: 37, and SEQ ID NO: 39, respectively.

Examples of beta-glucosidases useful in the methods of the present invention include, but are not limited to, *Aspergillus oryzae* beta-glucosidase (SEQ ID NO: 42); *Aspergillus fumigatus* beta-glucosidase (SEQ ID NO: 44); *Penicillium brasilianum* IBT 20888 beta-glucosidase (SEQ ID NO: 46); *Aspergillus niger* beta-glucosidase (SEQ ID NO: 48); and *Aspergillus aculeatus* beta-glucosidase (SEQ ID NO: 50). The beta-glucosidases of SEQ ID NO: 40, SEQ ID NO: 42, SEQ ID NO: 44, SEQ ID NO: 46, SEQ ID NO: 48, and SEQ ID NO: 50 described above are encoded by the mature polypeptide coding sequence SEQ

ID NO: 39, SEQ ID NO: 41, SEQ ID NO: 43, SEQ ID NO: 45, SEQ ID NO: 47, and SEQ ID NO: 49, respectively.

The *Aspergillus oryzae* polypeptide having beta-glucosidase activity can be obtained according to WO 2002/095014. The *Aspergillus fumigatus* polypeptide having beta-glucosidase activity can be obtained according to WO 2005/047499. The *Penicillium brasilianum* polypeptide having beta-glucosidase activity can be obtained according to WO 2007/019442. The *Aspergillus niger* polypeptide having beta-glucosidase activity can be obtained according to Dan *et al.*, 2000, *J. Biol. Chem.* 275: 4973-4980. The *Aspergillus aculeatus* polypeptide having beta-glucosidase activity can be obtained according to Kawaguchi *et al.*, 1996, *Gene* 173: 287-288.

The beta-glucosidase may be a fusion protein. In one aspect, the beta-glucosidase is the *Aspergillus oryzae* beta-glucosidase variant BG fusion protein of SEQ ID NO: 52 or the *Aspergillus oryzae* beta-glucosidase fusion protein of SEQ ID NO: 54. In another aspect, the *Aspergillus oryzae* beta-glucosidase variant BG fusion protein is encoded by the polynucleotide of SEQ ID NO: 51 or the *Aspergillus oryzae* beta-glucosidase fusion protein is encoded by the polynucleotide of SEQ ID NO: 53.

Other endoglucanases, cellobiohydrolases, and beta-glucosidases are disclosed in numerous Glycosyl Hydrolase families using the classification according to Henrissat B., 1991, A classification of glycosyl hydrolases based on amino-acid sequence similarities, *Biochem. J.* 280: 309-316, and Henrissat B., and Bairoch A., 1996, Updating the sequence-based classification of glycosyl hydrolases, *Biochem. J.* 316: 695-696.

Other cellulolytic enzymes that may be used in the present invention are described in EP 495,257, EP 531,315, EP 531,372, WO 89/09259, WO 94/07998, WO 95/24471, WO 96/11262, WO 96/29397, WO 96/034108, WO 97/14804, WO 98/08940, WO 98/012307, WO 98/13465, WO 98/015619, WO 98/015633, WO 98/028411, WO 99/06574, WO 99/10481, WO 99/025846, WO 99/025847, WO 99/031255, WO 2000/009707, WO 2002/050245, WO 2002/0076792, WO 2002/101078, WO 2003/027306, WO 2003/052054, WO 2003/052055, WO 2003/052056, WO 2003/052057, WO 2003/052118, WO 2004/016760, WO 2004/043980, WO 2004/048592, WO 2005/001065, WO 2005/028636, WO 2005/093050, WO 2005/093073, WO 2006/074005, WO 2006/117432, WO 2007/071818, WO 2007/071820, WO 2008/008070, WO 2008/008793, U.S. Patent No. 4,435,307, U.S. Patent No. 5,457,046, U.S. Patent No. 5,648,263, U.S. Patent No. 5,686,593, U.S. Patent No. 5,691,178, U.S. Patent No. 5,763,254, and U.S. Patent No. 5,776,757.

In the methods of the present invention, any polypeptide having cellulolytic enhancing activity can be used.

In a first aspect, the polypeptide having cellulolytic enhancing activity comprises the

following motifs:

[ILMV]-P-X(4,5)-G-X-Y-[ILMV]-X-R-X-[EQ]-X(4)-[HNQ] and [FW]-[TF]-K-[AIV],

wherein X is any amino acid, X(4,5) is any amino acid at 4 or 5 contiguous positions, and X(4) is any amino acid at 4 contiguous positions.

5 The polypeptide comprising the above-noted motifs may further comprise:

H-X(1,2)-G-P-X(3)-[YW]-[AILMV],

[EQ]-X-Y-X(2)-C-X-[EHQN]-[FILV]-X-[ILV], or

H-X(1,2)-G-P-X(3)-[YW]-[AILMV] and [EQ]-X-Y-X(2)-C-X-[EHQN]-[FILV]-X-[ILV],

10 wherein X is any amino acid, X(1,2) is any amino acid at 1 position or 2 contiguous positions, X(3) is any amino acid at 3 contiguous positions, and X(2) is any amino acid at 2 contiguous positions. In the above motifs, the accepted IUPAC single letter amino acid abbreviation is employed.

In a preferred aspect, the polypeptide having cellulolytic enhancing activity further comprises H-X(1,2)-G-P-X(3)-[YW]-[AILMV]. In another preferred aspect, the isolated polypeptide having cellulolytic enhancing activity further comprises [EQ]-X-Y-X(2)-C-X-[EHQN]-[FILV]-X-[ILV]. In another preferred aspect, the polypeptide having cellulolytic enhancing activity further comprises H-X(1,2)-G-P-X(3)-[YW]-[AILMV] and [EQ]-X-Y-X(2)-C-X-[EHQN]-[FILV]-X-[ILV].

20 In a second aspect, the polypeptide having cellulolytic enhancing activity comprises the following motif:

[ILMV]-P-x(4,5)-G-x-Y-[ILMV]-x-R-x-[EQ]-x(3)-A-[HNQ],

wherein x is any amino acid, x(4,5) is any amino acid at 4 or 5 contiguous positions, and x(3) is any amino acid at 3 contiguous positions. In the above motif, the accepted IUPAC single letter amino acid abbreviation is employed.

25 Examples of isolated polypeptides having cellulolytic enhancing activity include *Thielavia terrestris* polypeptides having cellulolytic enhancing activity (the mature polypeptide of SEQ ID NO: 56, SEQ ID NO: 58, SEQ ID NO: 60, SEQ ID NO: 62, SEQ ID NO: 64, or SEQ ID NO: 66); *Thermoascus auranticus* (the mature polypeptide of SEQ ID NO: 68), or *Trichoderma reesei* (the mature polypeptide of SEQ ID NO: 70). The polypeptides having cellulolytic enhancing activity of SEQ ID NO: 56, SEQ ID NO: 58, SEQ ID NO: 60, SEQ ID NO: 62, SEQ ID NO: 64, SEQ ID NO: 66, SEQ ID NO: 68, and SEQ ID NO: 70 described above are encoded by the mature polypeptide coding sequence of SEQ ID NO: 55, SEQ ID NO: 57, SEQ ID NO: 59, SEQ ID NO: 61, SEQ ID NO: 63, SEQ ID NO: 65, SEQ ID NO: 67, and SEQ ID NO: 69, respectively. See, for example, WO 2005/074647, 35 WO 2005/074656, and WO 2007/089290.

Examples of commercial xylan degrading enzyme preparations suitable for use in the present invention include, for example, SHEARZYME™ (Novozymes A/S), CELLIC™ Htec

(Novozymes A/S), VISCOZYME® (Novozymes A/S), ULTRAFLO® (Novozymes A/S), PULPZYME® HC (Novozymes A/S), MULTIFECT® Xylanase (Genencor), ECOPULP® TX-200A (AB Enzymes), HSP 6000 Xylanase (DSM), DEPOL™ 333P (Biocatalysts Limit, Wales, UK), DEPOL™ 740L. (Biocatalysts Limit, Wales, UK), and DEPOL™ 762P (Biocatalysts Limit, Wales, UK).

Examples of xylanases useful in the methods of the present invention include, but are not limited to, *Aspergillus aculeatus* xylanase (GeneSeqP:AAR63790; WO 94/21785), *Aspergillus fumigatus* xylanases (WO 2006/078256), and *Thielavia terrestris* NRRL 8126 xylanases (WO 2009/079210).

Examples of beta-xylosidases useful in the methods of the present invention include, but are not limited to, *Trichoderma reesei* beta-xylosidase (UniProtKB/TrEMBL accession number Q92458), *Talaromyces emersonii* (SwissProt accession number Q8X212), and *Neurospora crassa* (SwissProt accession number Q7SOW4).

Examples of acetylxylan esterases useful in the methods of the present invention include, but are not limited to, *Hypocrea jecorina* acetylxylan esterase (WO 2005/001036), *Neurospora crassa* acetylxylan esterase (UniProt accession number q7s259), *Thielavia terrestris* NRRL 8126 acetylxylan esterase (WO 2009/042846), *Chaetomium globosum* acetylxylan esterase (Uniprot accession number Q2GWX4), *Chaetomium gracile* acetylxylan esterase (GeneSeqP accession number AAB82124), *Phaeosphaeria nodorum* acetylxylan esterase (Uniprot accession number Q0UHJ1), and *Humicola insolens* DSM 1800 acetylxylan esterase (WO 2009/073709).

Examples of ferulic acid esterases useful in the methods of the present invention include, but are not limited to, *Humicola insolens* DSM 1800 feruloyl esterase (WO 2009/076122), *Neurospora crassa* feruloyl esterase (UniProt accession number Q9HGR3), and *Neosartorya fischeri* feruloyl esterase (UniProt Accession number A1D9T4).

Examples of arabinofuranosidases useful in the methods of the present invention include, but are not limited to, *Humicola insolens* DSM 1800 arabinofuranosidase (WO 2009/073383) and *Aspergillus niger* arabinofuranosidase (GeneSeqP accession number AAR94170).

Examples of alpha-glucuronidases useful in the methods of the present invention include, but are not limited to, *Aspergillus clavatus* alpha-glucuronidase (UniProt accession number alcc12), *Trichoderma reesei* alpha-glucuronidase (Uniprot accession number Q99024), *Talaromyces emersonii* alpha-glucuronidase (UniProt accession number Q8X211), *Aspergillus niger* alpha-glucuronidase (Uniprot accession number Q96WX9), *Aspergillus terreus* alpha-glucuronidase (SwissProt accession number Q0CJP9), and *Aspergillus fumigatus* alpha-glucuronidase (SwissProt accession number Q4WW45).

The enzymes and proteins used in the methods of the present invention may be

produced by fermentation of the above-noted microbial strains on a nutrient medium containing suitable carbon and nitrogen sources and inorganic salts, using procedures known in the art (see, e.g., Bennett, J.W. and LaSure, L. (eds.), *More Gene Manipulations in Fungi*, Academic Press, CA, 1991). Suitable media are available from commercial suppliers or may be prepared according to published compositions (e.g., in catalogues of the American Type Culture Collection). Temperature ranges and other conditions suitable for growth and enzyme production are known in the art (see, e.g., Bailey, J.E., and Ollis, D.F., *Biochemical Engineering Fundamentals*, McGraw-Hill Book Company, NY, 1986).

The fermentation can be any method of cultivation of a cell resulting in the expression or isolation of an enzyme. Fermentation may, therefore, be understood as comprising shake flask cultivation, or small- or large-scale fermentation (including continuous, batch, fed-batch, or solid state fermentations) in laboratory or industrial fermentors performed in a suitable medium and under conditions allowing the enzyme to be expressed or isolated. The resulting enzymes produced by the methods described above may be recovered from the fermentation medium and purified by conventional procedures.

Fermentation. The fermentable sugars obtained from the pretreated and hydrolyzed cellulosic material can be fermented by one or more (several) fermenting microorganisms capable of fermenting the sugars directly or indirectly into a desired fermentation product. "Fermentation" or "fermentation process" refers to any fermentation process or any process comprising a fermentation step. Fermentation processes also include fermentation processes used in the consumable alcohol industry (e.g., beer and wine), dairy industry (e.g., fermented dairy products), leather industry, and tobacco industry. The fermentation conditions depend on the desired fermentation product and fermenting organism and can easily be determined by one skilled in the art.

In the fermentation step, sugars, released from cellulosic material as a result of the pretreatment and enzymatic hydrolysis steps, are fermented to a product, e.g., ethanol, by a fermenting organism, such as yeast. Hydrolysis (saccharification) and fermentation can be separate or simultaneous, as described herein.

Any suitable hydrolyzed cellulosic material can be used in the fermentation step in practicing the present invention. The material is generally selected based on the desired fermentation product, i.e., the substance to be obtained from the fermentation, and the process employed, as is well known in the art.

The term "fermentation medium" is understood herein to refer to a medium before the fermenting microorganism(s) is(are) added, such as, a medium resulting from a saccharification process, as well as a medium used in a simultaneous saccharification and fermentation process (SSF).

"Fermenting microorganism" refers to any microorganism, including bacterial and

5 fungal organisms, suitable for use in a desired fermentation process to produce a fermentation product. The fermenting organism can be C₆ and/or C₅ fermenting organisms, or a combination thereof. Both C₆ and C₅ fermenting organisms are well known in the art. Suitable fermenting microorganisms are able to ferment, *i.e.*, convert, sugars, such as glucose, xylose, xylulose, arabinose, maltose, mannose, galactose, or oligosaccharides, directly or indirectly into the desired fermentation product.

Examples of bacterial and fungal fermenting organisms producing ethanol are described by Lin *et al.*, 2006, *Appl. Microbiol. Biotechnol.* 69: 627-642.

10 Examples of fermenting microorganisms that can ferment C₆ sugars include bacterial and fungal organisms, such as yeast. Preferred yeast includes strains of the *Saccharomyces* spp., preferably *Saccharomyces cerevisiae*.

15 Examples of fermenting organisms that can ferment C₅ sugars include bacterial and fungal organisms, such as yeast. Preferred C₅ fermenting yeast include strains of *Pichia*, preferably *Pichia stipitis*, such as *Pichia stipitis* CBS 5773; strains of *Candida*, preferably *Candida boidinii*, *Candida brassicae*, *Candida sheatae*, *Candida diddensii*, *Candida pseudotropicalis*, or *Candida utilis*.

20 Other fermenting organisms include strains of *Zymomonas*, such as *Zymomonas mobilis*; *Hansenula*, such as *Hansenula anomala*; *Kluyveromyces*, such as *K. fragilis*; *Schizosaccharomyces*, such as *S. pombe*; and *E. coli*, especially *E. coli* strains that have been genetically modified to improve the yield of ethanol.

25 In a preferred aspect, the yeast is a *Saccharomyces* spp. In a more preferred aspect, the yeast is *Saccharomyces cerevisiae*. In another more preferred aspect, the yeast is *Saccharomyces distaticus*. In another more preferred aspect, the yeast is *Saccharomyces uvarum*. In another preferred aspect, the yeast is a *Kluyveromyces*. In another more preferred aspect, the yeast is *Kluyveromyces marxianus*. In another more preferred aspect, the yeast is *Kluyveromyces fragilis*. In another preferred aspect, the yeast is a *Candida*. In another more preferred aspect, the yeast is *Candida boidinii*. In another more preferred aspect, the yeast is *Candida brassicae*. In another more preferred aspect, the yeast is *Candida diddensii*. In another more preferred aspect, the yeast is *Candida pseudotropicalis*.
30 In another more preferred aspect, the yeast is *Candida utilis*. In another preferred aspect, the yeast is a *Clavispora*. In another more preferred aspect, the yeast is *Clavispora lusitaniae*. In another more preferred aspect, the yeast is *Clavispora opuntiae*. In another preferred aspect, the yeast is a *Pachysolen*. In another more preferred aspect, the yeast is *Pachysolen tannophilus*. In another preferred aspect, the yeast is a *Pichia*. In another more preferred aspect, the yeast is a *Pichia stipitis*. In another preferred aspect, the yeast is a *Bretannomyces*. In another more preferred aspect, the yeast is *Bretannomyces clausenii* (Philippidis, G. P., 1996, Cellulose bioconversion technology, in *Handbook on Bioethanol*:

Production and Utilization, Wyman, C. E., ed., Taylor & Francis, Washington, DC, 179-212).

Bacteria that can efficiently ferment hexose and pentose to ethanol include, for example, *Zymomonas mobilis* and *Clostridium thermocellum* (Philippidis, 1996, *supra*).

In a preferred aspect, the bacterium is a *Zymomonas*. In a more preferred aspect, the bacterium is *Zymomonas mobilis*. In another preferred aspect, the bacterium is a *Clostridium*. In another more preferred aspect, the bacterium is *Clostridium thermocellum*.

Commercially available yeast suitable for ethanol production includes, e.g., ETHANOL RED™ yeast (available from Fermentis/Lesaffre, USA), FALI™ (available from Fleischmann's Yeast, USA), SUPERSTART™ and THERMOSACC™ fresh yeast (available from Ethanol Technology, WI, USA), BIOFERM™ AFT and XR (available from NABC - North American Bioproducts Corporation, GA, USA), GERT STRAND™ (available from Gert Strand AB, Sweden), and FERMIOL™ (available from DSM Specialties).

In a preferred aspect, the fermenting microorganism has been genetically modified to provide the ability to ferment pentose sugars, such as xylose utilizing, arabinose utilizing, and xylose and arabinose co-utilizing microorganisms.

The cloning of heterologous genes into various fermenting microorganisms has led to the construction of organisms capable of converting hexoses and pentoses to ethanol (cofermentation) (Chen and Ho, 1993, Cloning and improving the expression of *Pichia stipitis* xylose reductase gene in *Saccharomyces cerevisiae*, *Appl. Biochem. Biotechnol.* 39-40: 135-147; Ho *et al.*, 1998, Genetically engineered *Saccharomyces* yeast capable of effectively cofermenting glucose and xylose, *Appl. Environ. Microbiol.* 64: 1852-1859; Kotter and Ciriacy, 1993, Xylose fermentation by *Saccharomyces cerevisiae*, *Appl. Microbiol. Biotechnol.* 38: 776-783; Walfridsson *et al.*, 1995, Xylose-metabolizing *Saccharomyces cerevisiae* strains overexpressing the TKL1 and TAL1 genes encoding the pentose phosphate pathway enzymes transketolase and transaldolase, *Appl. Environ. Microbiol.* 61: 4184-4190; Kuyper *et al.*, 2004, Minimal metabolic engineering of *Saccharomyces cerevisiae* for efficient anaerobic xylose fermentation: a proof of principle, *FEMS Yeast Research* 4: 655-664; Beall *et al.*, 1991, Parametric studies of ethanol production from xylose and other sugars by recombinant *Escherichia coli*, *Biotech. Bioeng.* 38: 296-303; Ingram *et al.*, 1998, Metabolic engineering of bacteria for ethanol production, *Biotechnol. Bioeng.* 58: 204-214; Zhang *et al.*, 1995, Metabolic engineering of a pentose metabolism pathway in ethanologenic *Zymomonas mobilis*, *Science* 267: 240-243; Deanda *et al.*, 1996, Development of an arabinose-fermenting *Zymomonas mobilis* strain by metabolic pathway engineering, *Appl. Environ. Microbiol.* 62: 4465-4470; WO 2003/062430, xylose isomerase).

In a preferred aspect, the genetically modified fermenting microorganism is *Saccharomyces cerevisiae*. In another preferred aspect, the genetically modified fermenting microorganism is *Zymomonas mobilis*. In another preferred aspect, the genetically modified

fermenting microorganism is *Escherichia coli*. In another preferred aspect, the genetically modified fermenting microorganism is *Klebsiella oxytoca*. In another preferred aspect, the genetically modified fermenting microorganism is *Kluyveromyces* sp.

5 It is well known in the art that the organisms described above can also be used to produce other substances, as described herein.

The fermenting microorganism is typically added to the degraded lignocellulose or hydrolysate and the fermentation is performed for about 8 to about 96 hours, such as about 24 to about 60 hours. The temperature is typically between about 26°C to about 60°C, in particular about 32°C or 50°C, and at about pH 3 to about pH 8, such as around pH 4-5, 6,
10 or 7.

In a preferred aspect, the yeast and/or another microorganism is applied to the degraded cellulosic material and the fermentation is performed for about 12 to about 96 hours, such as typically 24-60 hours. In a preferred aspect, the temperature is preferably between about 20°C to about 60°C, more preferably about 25°C to about 50°C, and most preferably about 32°C to about 50°C, in particular about 32°C or 50°C, and the pH is generally from about pH 3 to about pH 7, preferably around pH 4-7. However, some fermenting organisms, e.g., bacteria, have higher fermentation temperature optima. Yeast or another microorganism is preferably applied in amounts of approximately 10^5 to 10^{12} , preferably from approximately 10^7 to 10^{10} , especially approximately 2×10^8 viable cell count
15 per ml of fermentation broth. Further guidance in respect of using yeast for fermentation can be found in, e.g., "The Alcohol Textbook" (Editors K. Jacques, T.P. Lyons and D.R. Kelsall, Nottingham University Press, United Kingdom 1999), which is hereby incorporated by reference.

For ethanol production, following the fermentation the fermented slurry is distilled to
25 extract the ethanol. The ethanol obtained according to the methods of the invention can be used as, e.g., fuel ethanol, drinking ethanol, i.e., potable neutral spirits, or industrial ethanol.

A fermentation stimulator can be used in combination with any of the processes described herein to further improve the fermentation process, and in particular, the performance of the fermenting microorganism, such as, rate enhancement and ethanol yield.
30 A "fermentation stimulator" refers to stimulators for growth of the fermenting microorganisms, in particular, yeast. Preferred fermentation stimulators for growth include vitamins and minerals. Examples of vitamins include multivitamins, biotin, pantothenate, nicotinic acid, meso-inositol, thiamine, pyridoxine, para-aminobenzoic acid, folic acid, riboflavin, and Vitamins A, B, C, D, and E. See, for example, Alfenore *et al.*, Improving ethanol production and viability of *Saccharomyces cerevisiae* by a vitamin feeding strategy during fed-batch
35 process, Springer-Verlag (2002), which is hereby incorporated by reference. Examples of minerals include minerals and mineral salts that can supply nutrients comprising P, K, Mg, S,

Ca, Fe, Zn, Mn, and Cu.

Fermentation products: A fermentation product can be any substance derived from the fermentation. The fermentation product can be, without limitation, an alcohol (e.g., arabinitol, butanol, ethanol, glycerol, methanol, 1,3-propanediol, sorbitol, and xylitol); an organic acid (e.g., acetic acid, acetic acid, adipic acid, ascorbic acid, citric acid, 2,5-diketo-D-gluconic acid, formic acid, fumaric acid, glucaric acid, gluconic acid, glucuronic acid, glutaric acid, 3-hydroxypropionic acid, itaconic acid, lactic acid, malic acid, malonic acid, oxalic acid, oxaloacetic acid, propionic acid, succinic acid, and xylonic acid); a ketone (e.g., acetone); an amino acid (e.g., aspartic acid, glutamic acid, glycine, lysine, serine, and threonine); and a gas (e.g., methane, hydrogen (H₂), carbon dioxide (CO₂), and carbon monoxide (CO)). The fermentation product can also be protein as a high value product.

In a preferred aspect, the fermentation product is an alcohol. It will be understood that the term "alcohol" encompasses a substance that contains one or more hydroxyl moieties. In a more preferred aspect, the alcohol is arabinitol. In another more preferred aspect, the alcohol is butanol. In another more preferred aspect, the alcohol is ethanol. In another more preferred aspect, the alcohol is glycerol. In another more preferred aspect, the alcohol is methanol. In another more preferred aspect, the alcohol is 1,3-propanediol. In another more preferred aspect, the alcohol is sorbitol. In another more preferred aspect, the alcohol is xylitol. See, for example, Gong, C. S., Cao, N. J., Du, J., and Tsao, G. T., 1999, Ethanol production from renewable resources, in *Advances in Biochemical Engineering/Biotechnology*, Scheper, T., ed., Springer-Verlag Berlin Heidelberg, Germany, 65: 207-241; Silveira, M. M., and Jonas, R., 2002, The biotechnological production of sorbitol, *Appl. Microbiol. Biotechnol.* 59: 400-408; Nigam, P., and Singh, D., 1995, Processes for fermentative production of xylitol – a sugar substitute, *Process Biochemistry* 30 (2): 117-124; Ezeji, T. C., Qureshi, N. and Blaschek, H. P., 2003, Production of acetone, butanol and ethanol by *Clostridium beijerinckii* BA101 and *in situ* recovery by gas stripping, *World Journal of Microbiology and Biotechnology* 19 (6): 595-603.

In another preferred aspect, the fermentation product is an organic acid. In another more preferred aspect, the organic acid is acetic acid. In another more preferred aspect, the organic acid is acetic acid. In another more preferred aspect, the organic acid is adipic acid. In another more preferred aspect, the organic acid is ascorbic acid. In another more preferred aspect, the organic acid is citric acid. In another more preferred aspect, the organic acid is 2,5-diketo-D-gluconic acid. In another more preferred aspect, the organic acid is formic acid. In another more preferred aspect, the organic acid is fumaric acid. In another more preferred aspect, the organic acid is glucaric acid. In another more preferred aspect, the organic acid is gluconic acid. In another more preferred aspect, the organic acid is glucuronic acid. In another more preferred aspect, the organic acid is glutaric acid. In

another preferred aspect, the organic acid is 3-hydroxypropionic acid. In another more preferred aspect, the organic acid is itaconic acid. In another more preferred aspect, the organic acid is lactic acid. In another more preferred aspect, the organic acid is malic acid. In another more preferred aspect, the organic acid is malonic acid. In another more preferred aspect, the organic acid is oxalic acid. In another more preferred aspect, the organic acid is propionic acid. In another more preferred aspect, the organic acid is succinic acid. In another more preferred aspect, the organic acid is xylonic acid. See, for example, Chen, R., and Lee, Y. Y., 1997, Membrane-mediated extractive fermentation for lactic acid production from cellulosic biomass, *Appl. Biochem. Biotechnol.* 63-65: 435-448.

In another preferred aspect, the fermentation product is a ketone. It will be understood that the term "ketone" encompasses a substance that contains one or more ketone moieties. In another more preferred aspect, the ketone is acetone. See, for example, Qureshi and Blaschek, 2003, *supra*.

In another preferred aspect, the fermentation product is an amino acid. In another more preferred aspect, the organic acid is aspartic acid. In another more preferred aspect, the amino acid is glutamic acid. In another more preferred aspect, the amino acid is glycine. In another more preferred aspect, the amino acid is lysine. In another more preferred aspect, the amino acid is serine. In another more preferred aspect, the amino acid is threonine. See, for example, Richard, A., and Margaritis, A., 2004, Empirical modeling of batch fermentation kinetics for poly(glutamic acid) production and other microbial biopolymers, *Biotechnology and Bioengineering* 87 (4): 501-515.

In another preferred aspect, the fermentation product is a gas. In another more preferred aspect, the gas is methane. In another more preferred aspect, the gas is H₂. In another more preferred aspect, the gas is CO₂. In another more preferred aspect, the gas is CO. See, for example, Kataoka, N., A. Miya, and K. Kiriya, 1997, Studies on hydrogen production by continuous culture system of hydrogen-producing anaerobic bacteria, *Water Science and Technology* 36 (6-7): 41-47; and Gunaseelan V.N. in *Biomass and Bioenergy*, Vol. 13 (1-2), pp. 83-114, 1997, Anaerobic digestion of biomass for methane production: A review.

Recovery. The fermentation product(s) can be optionally recovered from the fermentation medium using any method known in the art including, but not limited to, chromatography, electrophoretic procedures, differential solubility, distillation, or extraction. For example, alcohol is separated from the fermented cellulosic material and purified by conventional methods of distillation. Ethanol with a purity of up to about 96 vol.% can be obtained, which can be used as, for example, fuel ethanol, drinking ethanol, *i.e.*, potable neutral spirits, or industrial ethanol.

Signal Peptide

The present invention also relates to an isolated polynucleotide encoding a signal peptide comprising or consisting of amino acids 1 to 19 of SEQ ID NO: 2.

5 The present invention also relates to nucleic acid constructs comprising a gene encoding a protein, wherein the gene is operably linked to one or both of the isolated polynucleotide encoding the signal peptide and the isolated polynucleotide encoding the propeptide, wherein the gene is foreign to the polynucleotides encoding the signal peptide and propeptide.

10 In a preferred aspect, the isolated polynucleotide encoding a signal peptide comprises or consists of nucleotides 1 to 57 of SEQ ID NO: 1.

The present invention also relates to recombinant expression vectors and recombinant host cells comprising such nucleic acid constructs.

15 The present invention also relates to methods of producing a protein, comprising: (a) cultivating a recombinant host cell comprising a gene encoding a protein operably linked to the such a polynucleotide encoding a signal peptide, wherein the gene is foreign to the polynucleotide under conditions conducive for production of the protein; and (b) recovering the protein.

20 The protein may be native or heterologous to a host cell. The term "protein" is not meant herein to refer to a specific length of the encoded product and, therefore, encompasses peptides, oligopeptides, and proteins. The term "protein" also encompasses two or more polypeptides combined to form the encoded product. The proteins also include hybrid polypeptides that comprise a combination of partial or complete polypeptide sequences obtained from at least two different proteins wherein one or more (several) may be heterologous or native to the host cell. Proteins further include naturally occurring allelic and engineered variations of the above mentioned proteins and hybrid proteins.

25 Preferably, the protein is a hormone or variant thereof, enzyme, receptor or portion thereof, antibody or portion thereof, or reporter. In a more preferred aspect, the protein is an oxidoreductase, transferase, hydrolase, lyase, isomerase, or ligase. In an even more preferred aspect, the protein is an aminopeptidase, amylase, carbohydrase, carboxypeptidase, catalase, cellulase, chitinase, cutinase, cyclodextrin glycosyltransferase, deoxyribonuclease, esterase, alpha-galactosidase, beta-galactosidase, glucoamylase, alpha-glucosidase, beta-glucosidase, invertase, laccase, another lipase, mannosidase, mutanase, oxidase, pectinolytic enzyme, peroxidase, phytase, polyphenoloxidase, 30 proteolytic enzyme, ribonuclease, transglutaminase or xylanase.

35 The gene may be obtained from any prokaryotic, eukaryotic, or other source.

The present invention is further described by the following examples that should not be construed as limiting the scope of the invention.

5

Examples

Chemicals used as buffers and substrates were commercial products of at least reagent grade.

10 Materials

Chemicals used as buffers and substrates were commercial products of at least reagent grade.

Strains

15 *Trichophaea saccata* strain CBS 804.70 was used as the source of a gene encoding a Family GH3A beta-glucosidase. *Aspergillus oryzae* JAL250 strain (WO 99/61651) was used for expression of the *Trichophaea saccata* CBS 804.70 beta-glucosidase.

Media

20 MEX-1 medium was composed of 20 g of soya bean meal, 15 g of wheat bran, 10 g of microcrystalline cellulose (AVICEL®; FMC, Philadelphia, PA, USA), 5 g of maltodextrin, 3 g of Bactopeptone, 0.2 g of pluronic, 1 g of olive oil, and deionized water to 1 liter.

PDA plates were composed of 39 grams of potato dextrose agar and deionized water to 1 liter.

25 LB medium was composed of 10 g of tryptone, 5 g of yeast extract, 5 g of sodium chloride, and deionized water to 1 liter.

LB plates were composed of LB medium and 15 g of Bacto agar per liter.

30 SOC medium was composed of 2% tryptone, 0.5% yeast extract, 10 mM NaCl, 2.5 mM KCl, 10 mM MgCl₂, and 10 mM MgSO₄, and deionized water to 1 liter; sterilized by autoclaving and then filter-sterilized glucose was added to 20 mM.

MDU2BP medium was composed of 45 g of maltose, 1 g of MgSO₄·7H₂O, 1 g of NaCl, 2 g of K₂HPO₄, 12 g of KH₂PO₄, 2 g of urea, 500 µl of AMG trace metals solution, and deionized water to 1 liter. The pH was adjusted to 5.0 and then filter sterilized using a 0.22 µm filtering unit.

35 AMG trace metals solution was composed of 14.3 g of ZnSO₄·7H₂O, 2.5 g of CuSO₄·5H₂O, 0.5 g of NiCl₂·6H₂O, 13.8 g of FeSO₄·7H₂O, 8.5 g of MnSO₄·7H₂O, 3 g of citric acid, and deionized water to 1 liter.

2X YT plates were composed per liter of 16 g of tryptone, 10 g of yeast extract, 5 g of NaCl, 15 g of bacto agar, and deionized water to 1 liter.

YPG medium was composed of 4 g of yeast extract, 1 g of K₂HPO₄, 0.5 g of MgSO₄, 15.0 g of glucose, and deionized water to 1 liter (pH 6.0).

5 YPM medium was composed of 10 g of yeast extract, 20 g of Bacto peptone, 20 g of maltose, and deionized water to 1 liter.

M410 medium was composed of 50 g of glucose, 50 g of maltose, 2 g of MgSO₄·7H₂O, 2 g of KH₂PO₄, 4 g of citric acid, 8 g of yeast extract, 2 g of urea, 0.5 g of CaCl₂, 0.5 ml of AMG trace metals solution, and deionized water to 1 liter (pH 6.0).

10 **Example 1: Preparation of *Trichophaea saccata* strain CBS 804.70 mycelia for cDNA library production**

Trichophaea saccata CBS 804.70 was inoculated onto a PDA plate and incubated for 7 days at 28°C. Several mycelia-PDA agar plugs were inoculated into 750 ml shake flasks containing 100 ml of MEX-1 medium. The flasks were incubated at 37°C for 9 days with shaking at 150 rpm. The fungal mycelia were harvested by filtration through MIRACLOTH® (Calbiochem, San Diego, CA, USA) before being frozen in liquid nitrogen. The mycelia were then pulverized into a powder by milling the frozen mycelia together with an equal volume of dry ice in a coffee grinder precooled with liquid nitrogen. The powdered mycelial material was kept at -80°C until use.

Example 2: *Trichophaea saccata* strain CBS 804.70 RNA isolation

Total RNA was prepared from the frozen, powdered mycelia of *Trichophaea saccata* CBS 804.70 by extraction with guanidium thiocyanate followed by ultracentrifugation through a 5.7 M CsCl cushion according to Chirgwin *et al.*, 1979, *Biochemistry* 18: 5294-5299. The polyA enriched RNA was isolated by oligo (dT)-cellulose affinity chromatography according to Aviv *et al.*, 1972, *Proc. Natl. Acad. Sci. USA* 69: 1408-1412.

Example 3: Construction of a *Trichophaea saccata* strain CBS 804.70 cDNA library

30 Double stranded cDNA was synthesized according to the general methods of Gubler and Hoffman, 1983, *Gene* 25: 263-269; Sambrook, J., Fritsch, E.F., and Maniatis, T. *Molecular cloning: A Laboratory Manual*, 2nd ed., 1989, Cold Spring Harbor Laboratory, Cold Spring Harbor, New York; and Kofod *et al.*, 1994, *J. Biol. Chem.* 269: 29182-29189, using a polyA-Not I primer (Promega Corp., Madison, Wisconsin, USA). After synthesis, the cDNA was treated with mung bean nuclease, blunt ended with T4 DNA polymerase, and ligated to a 50-fold molar excess of *Eco* RI adaptors (Invitrogen Corp., Carlsbad, CA, USA). The cDNA was cleaved with *Not* I and size fractionated by 0.8% agarose gel electrophoresis

using 44 mM Tris base, 44 mM boric acid, 0.5 mM EDTA (TBE) buffer. The fraction of cDNA of 700 bp and larger was excised from the gel and purified using a GFX® PCR DNA and Gel Band Purification Kit (GE Healthcare, United Kingdom) according to the manufacturer's instructions.

5 The prepared cDNA was then directionally cloned by ligation into *Eco* RI-Not I cleaved pMHas5 (WO 03/044049) using a Rapid Ligation Kit (Roche Diagnostics GmbH, Penzberg, Germany) according to the manufacturer's instructions. The ligation mixture was electroporated into *E. coli* DH10B cells (Invitrogen Corp., Carlsbad, CA, USA) using a GENE PULSER® and Pulse Controller (Bio-Rad Laboratories, Inc., Hercules, CA, USA) at 50 μ F,
10 25 mAmp, 1.8 kV with a 2 mm gap cuvette according to the manufacturer's procedure.

The electroporated cells were spread onto LB plates supplemented with 50 μ g of kanamycin per ml. A cDNA plasmid pool was prepared from approximately 30,000 total transformants of the original cDNA-pMHas5 vector ligation. Plasmid DNA was prepared directly from the pool of colonies using a QIAPREP® Spin Midi/Maxiprep Kit (QIAGEN
15 GmbH Corporation, Hilden, Germany). The cDNA library was designated SBL521-2.

Example 4: Construction of a SigA4 transposon containing the β -lactamase reporter gene

A transposon containing plasmid designated pSigA4 was constructed from the
20 pSigA2 transposon containing plasmid described in WO 01/77315 in order to create an improved version of the signal trapping transposon of pSigA2 with decreased selection background. The pSigA2 transposon contains a signal-less beta-lactamase construct encoded on the transposon itself. PCR was used to create a deletion of the intact beta-lactamase gene found on the plasmid backbone using a proofreading PROOFSTART® DNA
25 polymerase (QIAGEN GmbH Corporation, Hilden, Germany) and the following 5' phosphorylated primers (TAG Copenhagen, Denmark):

SigA2NotU-P:

5'-TCGCGATCCGTTTTTCGCATTTATCGTGAAACGCT-3' (SEQ ID NO: 71)

SigA2NotD-P:

30 5'-CCGCAAACGCTGGTGAAAGTAAAAGATGCTGAA-3' (SEQ ID NO: 72)

The amplification reaction was composed of 1 μ l of pSigA2 (10 ng/ μ l), 5 μ l of 10X PROOFSTART™ Buffer (QIAGEN GmbH Corporation, Hilden, Germany), 2.5 μ l of dNTP mix (20 mM), 0.5 μ l of SigA2NotU-P (10 mM), 0.5 μ l of SigA2NotD-P (10 mM), 10 μ l of Q solution (QIAGEN GmbH Corporation, Hilden, Germany), and 31.25 μ l of deionized water. A
35 DNA ENGINE™ Thermal Cycler (MJ Research Inc., Waltham, MA, USA) was used for the amplification programmed for 1 cycle at 95°C for 5 minutes; and 20 cycles each at 94°C for 30 seconds, 62°C for 30 seconds, and 72°C for 4 minutes.

A 3.9 kb PCR reaction product was isolated by 0.8% agarose gel electrophoresis using 40 mM Tris base-20 mM sodium acetate-1 mM disodium EDTA (TAE) buffer and 0.1 µg of ethidium bromide per ml. The DNA band was visualized with the aid of an Eagle Eye Imaging System (Stratagene, La Jolla, CA, USA) at 360 nm. The 3.9 kb DNA band was excised from the gel and purified using a GFX® PCR DNA and Gel Band Purification Kit according to the manufacturer's instructions.

The 3.9 kb fragment was self-ligated at 16°C overnight with 10 units of T4 DNA ligase (New England Biolabs, Inc., Beverly, MA, USA), 9 µl of the 3.9 kb PCR fragment, and 1 µl of 10X ligation buffer (New England Biolabs, Inc., Beverly, MA, USA). The ligation was heat inactivated for 10 minutes at 65°C and then digested with *Dpn* I at 37°C for 2 hours. After incubation, the digestion was purified using a GFX® PCR DNA and Gel Band Purification Kit.

The purified material was then transformed into *E. coli* TOP10 competent cells (Invitrogen Corp., Carlsbad, CA, USA) according to the manufacturer's instructions. The transformation mixture was plated onto LB plates supplemented with 25 µg of chloramphenicol per ml. Plasmid minipreps were prepared from several transformants and digested with *Bgl* II. One plasmid with the correct construction was chosen and designated pSigA4. Plasmid pSigA4 contains the *Bgl* II flanked transposon SigA2 (SEQ ID NO: 71) identical to that disclosed in WO 01/77315.

A 60 µl sample of plasmid pSigA4 DNA (0.3 µg/µl) was digested with *Bgl* II and separated by 0.8% agarose gel electrophoresis using TAE buffer. A SigA2 transposon DNA band of 2 kb was eluted with 200 µl of EB buffer (QIAGEN GmbH Corporation, Hilden, Germany) and purified using a GFX® PCR DNA and Gel Band Purification Kit according to the manufacturer's instructions and eluted in 200 µl of EB buffer. SigA2 was used for transposon assisted signal trapping.

Example 5: Transposon Assisted Signal Trapping of *Trichophaea saccata* CBS 804.70

A complete description of transposon assisted signal trapping can be found in WO 01/77315. A cDNA plasmid pool was prepared from 30,000 total transformants of the original cDNA-pMHas5 vector ligation. Plasmid DNA was prepared directly from a pool of colonies recovered from solid LB selective medium using a QIAPREP® Spin Midi/Maxiprep Kit. The plasmid pool was treated with transposon SigA2 and MuA transposase (Finnzymes OY, Espoo, Finland) according to the manufacturer's instructions.

For *in vitro* transposon tagging of the *Trichophaea saccata* CBS 804.70 cDNA library, 4 or 8 µl of SigA2 transposon containing approximately 2.6 µg of DNA were mixed with 1 µl of the plasmid DNA pool of the *Trichophaea saccata* CBS 804.70 cDNA library containing 2 µg of DNA, 2 µl of MuA transposase (0.22 µg/µl), and 5 µl of 5X buffer (Finnzymes OY,

Espoo, Finland) in a total volume of 50 μ l and incubated at 30°C for 3.5 hours followed by heat inactivation at 75°C for 10 minutes. The DNA was precipitated by addition of 5 μ l of 3 M sodium acetate pH 5 and 110 μ l of 96% ethanol and centrifuged for 30 minutes at 10,000 \times g. The pellet was washed in 70% ethanol, air dried at room temperature, and resuspended in 10 μ l of 10 mM Tris, pH 8, 1 mM EDTA (TE) buffer.

A 1.5 μ l volume of the transposon tagged plasmid pool was electroporated into 20 μ l of *E. coli* DH10B ultracompetent cells (Gibco-BRL, Gaithersburg MD, USA) according to the manufacturer's instructions using a GENE PULSER® and Pulse Controller (Bio-Rad Laboratories, Inc., Hercules, CA, USA) at 50 μ F, 25 mA, 1.8 kV with a 2 mm gap cuvette according to the manufacturer's procedure.

The electroporated cells were incubated in SOC medium with shaking at 250 rpm for 2 hours at 28°C before being plated on the following selective media: LB medium supplemented with 50 μ g of kanamycin per ml; LB medium supplemented with 50 μ g of kanamycin per ml and 15 μ g of chloramphenicol per ml; and/or LB medium supplemented with 50 μ g of kanamycin per ml, 15 μ g of chloramphenicol per ml, and 12.5 μ g of ampicillin per ml.

From dilution plating of the electroporation onto LB medium supplemented with kanamycin and chloramphenicol medium, it was determined that approximately 72,000 colonies were present containing a cDNA plasmid library with a SigA2 transposition per electroporation and that approximately 69 colonies were recovered under triple selection (LB, kanamycin, chloramphenicol, ampicillin). Further electroporation and plating experiments were performed until 445 total colonies were recovered under triple selection. The colonies were miniprepmed using a QIAPREP® 96 Turbo Miniprep Kit (QIAGEN GmbH Corporation, Hilden, Germany). Plasmids were sequenced with the transposon forward and reverse primers (primers A and B), shown below, according to the procedure disclosed in WO 2001/77315 (page 28)

Primer A:

5'-AGCGTTTGCGGCCGCGATCC-3' (SEQ ID NO: 73)

Primer B:

5'-TTATTCGGTCGAAAAGGATCC-3' (SEQ ID NO: 74)

Example 6: Sequence assembly and annotation

DNA sequences were obtained for the reactions using an ABI PRISM Automated DNA Sequencer Model 3700 (Applied Biosystems, Foster City, CA, USA). Primer A and primer B sequence reads for each plasmid were trimmed to remove vector and transposon sequence. This resulted in 225 assembled sequences, which were grouped into 148 contigs by using the program PhredPhrap (Ewing *et al.*, 1998, *Genome Research* 8: 175-185; Ewing

and Green, 1998, *Genome Research* 8: 186-194). All 148 contigs were subsequently compared to sequences available in standard public DNA and protein sequences databases (TrEMBL, SWALL, PDB, EnsemblPep, GeneSeqP) by using the program BLASTX 2.0a19MP-WashU [14-Jul-1998] [Build linux-x86 18:51:44 30-Jul-1998] (Gish *et al.*, 1993, *Nat. Genet.* 3: 266-72). The Family GH3A beta-glucosidase was identified directly by analysis of the BlastX results.

The open reading frame for the DNA contig was extracted from the assembly of 15 independent transposition events positioned randomly along the coding region. The nucleotide sequence (SEQ ID NO: 1) and deduced amino acid sequence (SEQ ID NO: 2) of the *Trichophaea saccata* beta-glucosidase are shown in Figures 1A and 1B. The cDNA fragment encodes a polypeptide of 856 amino acids. The %G+C content of the gene is 58.1%. Using the SignalP software program (Nielsen *et al.*, 1997, *Protein Engineering* 10: 1-6), a signal peptide of 19 residues was predicted. The predicted mature protein contains 837 amino acids with a molecular mass of 90.35 kDa.

A comparative alignment of beta-glucosidase Family 3 sequences was determined using the Needleman-Wunsch algorithm (Needleman and Wunsch, 1970, *J. Mol. Biol.* 48: 443-453) as implemented in the Needle program of EMBOSS with gap open penalty of 10, gap extension penalty of 0.5, and the EBLOSUM62 matrix. The alignment showed that the deduced amino acid sequence of the *Trichophaea saccata* Cel3A mature polypeptide shared 59.3% identity (excluding gaps) to the deduced amino acid sequence of a hypothetical predicted protein from *Sclerotinia sclerotiorum* (Uniprot accession number A7EI88).

Example 7: Cloning of the *Trichophaea saccata* Family GH3A gene into an *Aspergillus oryzae* expression vector

Two synthetic oligonucleotide primers shown below were designed to PCR amplify the *Trichophaea saccata gh3a* gene from cDNA. An IN-FUSION™ PCR Cloning Kit (Clontech Laboratories, Inc., Mountain View, CA, USA) was used to clone the fragment directly into the expression vector, pAILo2 (WO 2005/074647).

Forward primer:

5'-ACTGGATTTACCATGTTTGGCATCACCTCGACTGCG-3' (SEQ ID NO: 75)

Reverse primer:

5'-TCACCTCTAGTTAATTAATCAATAATGATCACTCGG-3' (SEQ ID NO: 76)

Bold letters represent coding sequence. The remaining sequence is homologous to the insertion sites of pAILo2.

Fifty picomoles of each of the primers above were used in a PCR reaction containing 1 µl (239 µg) of the SBL521-2 cDNA library, 1X EXPAND® High Fidelity buffer with MgCl₂ (Roche Diagnostics Corp., Indianapolis, IN, USA), 1 µl of a 10 mM blend of dATP, dTTP,

dGTP, and dCTP, and 2.6 units of EXPAND® High Fidelity enzyme mix (Roche, Indianapolis, IN, USA) in a final volume of 50 µl. An EPPENDORF® MASTERCYCLER® epgradient S (Eppendorf Scientific, Inc., Westbury, NY, USA) was used to amplify the fragment programmed for 1 cycle at 94°C for 2 minutes; 10 cycles each at 94°C for 15
5 seconds, 58°C for 30 seconds, and 72°C for 2 minutes; 20 cycles each at 94°C for 15 seconds, 58°C for 30 seconds, and 72°C for 120 seconds plus 5 seconds per successive cycle. The heat block was then held at 72°C for 7 minutes followed by a 4°C soak cycle.

The reaction products were isolated by 1.0% agarose gel electrophoresis using TAE buffer and an approximately 2.5 kb product band was excised from the gel and purified using
10 a MINELUTE® Gel Extraction Kit (QIAGEN Inc., Valencia, CA, USA) according to the manufacturer's instructions.

The purified PCR fragment was then cloned into pAILo2 using an IN-FUSION™ PCR Cloning Kit. The vector was digested with *Nco* I and *Pac* I. The digested fragment was purified by gel electrophoresis as above, excised from the gel, and purified using a
15 QIAQUICK® Gel Extraction Kit (QIAGEN Inc., Valencia, CA, USA). The gene fragment and the digested vector were combined together in a reaction resulting in the expression plasmid pAG39, in which transcription of the *gh3a* gene was under the control of the NA2-tpi promoter (a hybrid of the promoters from the genes for *Aspergillus niger* neutral alpha-amylase and *Aspergillus nidulans* triose phosphate isomerase). The recombination reaction
20 (20 µl) was composed of 1X IN-FUSION™ Buffer (Clontech, Mountain View, CA, USA), 1X BSA (Clontech, Mountain View, CA, USA), 1 µl of IN-FUSION™ enzyme (diluted 1:10) (Clontech, Mountain View, CA, USA), 249 ng of pAILo2 digested with *Nco* I and *Pac* I, and 82.5 ng of the *Trichophaea saccata gh3a* purified PCR product. The reaction was incubated at 37°C for 15 minutes followed by 15 minutes at 50°C. The reaction was diluted with 40 µl
25 of TE buffer and 2.5 µl of the diluted reaction was used to transform *E. coli* TOP10 competent cells (Invitrogen, Carlsbad, CA, USA).

Plasmid DNA was prepared using a BIOROBOT® 9600 (QIAGEN Inc., Valencia, CA, USA) and putative pAG39 clones were digested with *Bam* HI. The plasmid DNA from these clones was then sequenced to identify clones without PCR induced errors. Sequencing
30 reactions contained 1.5 µl of plasmid DNA, 4.5 µl of water, and 4 µl of sequencing master-mix containing 1 µl of 5X sequencing buffer (Millipore, Billerica, MA, USA), 1 µl of BIGDYE™ terminator (Applied Biosystems, Inc., Foster City, CA, USA), 1 µl of water and one of the following primers at 3.2 pmoles per reaction.

Primer pAILo2 5':

5'-TGTCCCTTGTCGATGCG-3' (SEQ ID NO: 77)

Primer pAILo2 3':

5'-CACATGACTTGGCTTCC-3' (SEQ ID NO: 78)

Primer Ts3AF1:

5'-GTCTACCTCTGGCCATTCCA-3' (SEQ ID NO: 79)

Primer Ts3AF2:

5'-TACCGCAATCCTCCAGAACA-3' (SEQ ID NO: 80)

Primer Tsac *gh3a* 1675-17000 Rev:

5'-ACGCTTGTCGGCCCATTTCTCCATGA-3' (SEQ ID NO: 81)

Primer Tsac *gh3a* 925-950 Rev:

5'-TGGGCCAGCAAGGGCACACGATCAT-3' (SEQ ID NO: 82)

10

Example 8: Expression of the *Trichophaea saccata* cDNA encoding a GH3A polypeptide having beta-glucosidase activity in *Aspergillus oryzae* JaL355

Aspergillus oryzae JaL355 protoplasts were prepared according to the method of Christensen *et al.*, 1988, *Bio/Technology* 6: 1419-1422. Five µg of pAG39 were used to transform *Aspergillus oryzae* JaL355. Twenty-four transformants were isolated to individual PDA plates.

Confluent PDA plates of 24 transformants were washed with 5 ml of 0.01% TWEEN® 20 and spores collected. Eight µl of each spore stock was added to 1 ml of YPG, YPM, and M410 media separately in 24 well plates and incubated at 34°C. Three days after incubation, 7.5 µl of supernatant from the cultures were analyzed using a CRITERION® stain-free, 8-16% gradient SDS-PAGE gel (BioRad Laboratories, Inc., Hercules, CA, USA) according to the manufacturer's instructions. SDS-PAGE profiles of the cultures showed that several transformants had a new major band of approximately 100 kDa and the best expression in M410 medium. After a total of five days of incubation, all M410 cultures were sampled and analyzed by SDS-PAGE as described above at which point the transformant exhibiting the best expression was selected.

A confluent PDA plate of the top transformant was washed with 5 ml of 0.01% TWEEN® 20 and inoculated into five 500 ml shake flasks containing 100 ml of M410 medium to generate broth for characterization of the enzyme. The flasks were harvested on day 5. Broths were filtered using a 0.22 µm stericup suction filter (Millipore, Bedford, MA, USA).

Example 9: Characterization of *Trichophaea saccata* GH3 beta-glucosidase

Activity on cellobiose and 4-nitrophenyl-beta-D-glucopyranoside: Cellobiose was dissolved in 50 mM sodium acetate-0.01% TWEEN® 20 pH 5 buffer to 2 g/L. Enzyme was diluted at various concentrations using the same buffer. To 50 µl of the enzyme dilutions

(0.04 - 5 µg/ml), 100 µl of the cellobiose solution (2 g/L buffer) was added. The reaction was incubated at 50°C for 30 minutes before the reaction was stopped with 50 µl of 0.5 M sodium hydroxide. Glucose produced was measured using the Glucose Oxidase reagent (Pointe Scientific, Inc., Canton, MI, USA) according to the manufacturer.

5 4-Nitrophenyl-beta-D-glucopyranoside was dissolved in DMSO to 100 mM. Just before assay, the 100 mM 4-nitrophenyl-beta-D-glucopyranoside stock solution was diluted 100X in 50 mM sodium acetate-0.01% TWEEN® 20 buffer (1 mM 4-nitrophenyl-beta-D-glucopyranoside). Enzyme was diluted at various concentrations using the same buffer. To 20 µl of each enzyme dilution, 100 µl of the 4-nitrophenyl-beta-D-glucopyranoside solution
10 were added. The reaction was incubated at 40°C for 20 minutes before the reaction was stopped with 50 µl of 1 M sodium carbonate pH 10 buffer. The liberated 4-nitrophenol was determined spectrophotometrically at 405 nm.

The results demonstrated that the *Trichophaea saccata* GH3 beta-glucosidase was active toward both substrates

15 Thermostability: The *Trichophaea saccata* GH3A beta-glucosidase was diluted in 50 mM sodium acetate-0.01% TWEEN® 20 pH 5 buffer to 1 g/L, and then incubated at 50°C for 72 hours, and 60°C for 3 hours or 24 hours. The same sample was stored at 4°C to serve as a control. After incubation, the residual activity of the samples was measured using 4-nitrophenyl-beta-D-glucopyranoside as substrate, following the assay described above, but
20 using one enzyme loading which gave less than 5% conversion in the assay. The activity of sample at 4°C was normalized to 100%, and the activities of samples at other incubation conditions were compared to the 4°C activity. The thermostability of *Trichophaea saccata* GH3A beta-glucosidase is shown below in Table I.

Table I: Thermostability of *Trichophaea saccata* GH3A beta-glucosidase

Incubation condition	Residual activity
4°C	100%
50°C, 72 hours	95%
60°C, 3 hours	23%
60°C, 24 hours	6%

25 pH profile: The pH activity profile of the *Trichophaea saccata* GH3A beta-glucosidase was determined using the same protocol above for determining activity, except the enzyme was incubated at five different pHs: pH 4, 5, 6, 7, 8, and one enzyme loading was used which gave less than 5% conversion of the substrate in the assay. Britton-Robinson buffer
30 was used as the buffer system. To prepare the Britton-Robinson buffer, a 100 mM stock solution was made composed of 0.1 mole boric acid, 0.1 mole acetic acid, and 0.1 mol phosphoric acid in 1 liter of deionized water. The 100 mM stock solution was then titrated to

a certain pH (4~8) using 5 M sodium hydroxide and then diluted to 40 mM. The 4-nitrophenyl-beta-D-glucopyranoside was prepared with the buffer, and the activity was measured at 50°C. The highest activity was normalized to be 100%, and activities at other pHs were compared to the highest activity and expressed in % activity. The pH activity profile of the *Trichophaea saccata* GH3 beta-glucosidase is shown in Table II.

Table II: pH Activity Profile

pH value	Activity
4.0	100%
5.0	94%
6.0	64%
7.0	26%
8.0	1%

Deposit of Biological Material

The following biological material has been deposited under the terms of the Budapest Treaty with the Agricultural Research Service Patent Culture Collection, Northern Regional Research Center, 1815 University Street, Peoria, IL, USA, and given the following accession number:

Deposit	Accession Number	Date of Deposit
<i>E. coli</i> pAHYG-33	NRRL B-50214	December 22, 2008

The strain has been deposited under conditions that assure that access to the culture will be available during the pendency of this patent application to one determined by foreign patent laws to be entitled thereto. The deposit represents a substantially pure culture of the deposited strain. The deposit is available as required by foreign patent laws in countries wherein counterparts of the subject application, or its progeny are filed. However, it should be understood that the availability of a deposit does not constitute a license to practice the subject invention in derogation of patent rights granted by governmental action.

The present invention is further described by the following numbered paragraphs:

[1] An isolated polypeptide having beta-glucosidase activity, selected from the group consisting of:

(a) a polypeptide comprising an amino acid sequence having at least 60% sequence identity to the mature polypeptide of SEQ ID NO: 2;

(b) a polypeptide encoded by a polynucleotide that hybridizes under at least medium stringency conditions with (i) the mature polypeptide coding sequence of SEQ ID NO: 1, (ii) the genomic DNA sequence comprising the mature polypeptide coding sequence

of SEQ ID NO: 1, or (iii) a full-length complementary strand of (i) or (ii);

(c) a polypeptide encoded by a polynucleotide comprising a nucleotide sequence having at least 60% sequence identity to the mature polypeptide coding sequence of SEQ ID NO: 1; and

5 (d) a variant comprising a substitution, deletion, and/or insertion of one or more (several) amino acids of the mature polypeptide of SEQ ID NO: 2.

[2] The polypeptide of paragraph 1, comprising an amino acid sequence having at least 60% sequence identity to the mature polypeptide of SEQ ID NO: 2.

10 [3] The polypeptide of paragraph 2, comprising an amino acid sequence having at least 65% sequence identity to the mature polypeptide of SEQ ID NO: 2.

[4] The polypeptide of paragraph 3, comprising an amino acid sequence having at least 70% sequence identity to the mature polypeptide of SEQ ID NO: 2.

[5] The polypeptide of paragraph 4, comprising an amino acid sequence having at least 75% sequence identity to the mature polypeptide of SEQ ID NO: 2.

15 [6] The polypeptide of paragraph 5, comprising an amino acid sequence having at least 80% sequence identity to the mature polypeptide of SEQ ID NO: 2.

[7] The polypeptide of paragraph 6, comprising an amino acid sequence having at least 85% sequence identity to the mature polypeptide of SEQ ID NO: 2.

20 [8] The polypeptide of paragraph 7, comprising an amino acid sequence having at least 90% sequence identity to the mature polypeptide of SEQ ID NO: 2.

[9] The polypeptide of paragraph 8, comprising an amino acid sequence having at least 95% sequence identity to the mature polypeptide of SEQ ID NO: 2.

[10] The polypeptide of paragraph 9, comprising an amino acid sequence having at least 97% sequence identity to the mature polypeptide of SEQ ID NO: 2.

25 [11] The polypeptide of paragraph 1, comprising or consisting of the amino acid sequence of SEQ ID NO: 2; or a fragment thereof having beta-glucosidase activity.

[12] The polypeptide of paragraph 11, comprising or consisting of the amino acid sequence of SEQ ID NO: 2.

30 [13] The polypeptide of paragraph 1, comprising or consisting of the mature polypeptide of SEQ ID NO: 2.

[14] The polypeptide of paragraph 1, which is encoded by a polynucleotide that hybridizes under at least medium stringency conditions with (i) the mature polypeptide coding sequence of SEQ ID NO: 1, (ii) the genomic DNA sequence comprising the mature polypeptide coding sequence of SEQ ID NO: 1, or (iii) a full-length complementary strand of (i) or (ii).

35 [15] The polypeptide of paragraph 14, which is encoded by a polynucleotide that hybridizes under at least medium-high stringency conditions with (i) the mature polypeptide

coding sequence of SEQ ID NO: 1, (ii) the genomic DNA sequence comprising the mature polypeptide coding sequence of SEQ ID NO: 1, or (iii) a full-length complementary strand of (i) or (ii).

5 [16] The polypeptide of paragraph 15, which is encoded by a polynucleotide that hybridizes under at least high stringency conditions with (i) the mature polypeptide coding sequence of SEQ ID NO: 1, (ii) the genomic DNA sequence comprising the mature polypeptide coding sequence of SEQ ID NO: 1, or (iii) a full-length complementary strand of (i) or (ii).

10 [17] The polypeptide of paragraph 16, which is encoded by a polynucleotide that hybridizes under at least very high stringency conditions with (i) the mature polypeptide coding sequence of SEQ ID NO: 1, (ii) the genomic DNA sequence comprising the mature polypeptide coding sequence of SEQ ID NO: 1, or (iii) a full-length complementary strand of (i) or (ii).

15 [18] The polypeptide of paragraph 17, which is encoded by a polynucleotide comprising a nucleotide sequence having at least 60% sequence identity to the mature polypeptide coding sequence of SEQ ID NO: 1.

[19] The polypeptide of paragraph 18, which is encoded by a polynucleotide comprising a nucleotide sequence having at least 65% sequence identity to the mature polypeptide coding sequence of SEQ ID NO: 1.

20 [20] The polypeptide of paragraph 19, which is encoded by a polynucleotide comprising a nucleotide sequence having at least 70% sequence identity to the mature polypeptide coding sequence of SEQ ID NO: 1.

25 [21] The polypeptide of paragraph 20, which is encoded by a polynucleotide comprising a nucleotide sequence having at least 75% sequence identity to the mature polypeptide coding sequence of SEQ ID NO: 1.

[22] The polypeptide of paragraph 21, which is encoded by a polynucleotide comprising a nucleotide sequence having at least 80% sequence identity to the mature polypeptide coding sequence of SEQ ID NO: 1.

30 [23] The polypeptide of paragraph 22, which is encoded by a polynucleotide comprising a nucleotide sequence having at least 85% sequence identity to the mature polypeptide coding sequence of SEQ ID NO: 1.

[24] The polypeptide of paragraph 23, which is encoded by a polynucleotide comprising a nucleotide sequence having at least 90% sequence identity to the mature polypeptide coding sequence of SEQ ID NO: 1.

35 [25] The polypeptide of paragraph 24, which is encoded by a polynucleotide comprising a nucleotide sequence having at least 95% sequence identity to the mature polypeptide coding sequence of SEQ ID NO: 1.

[26] The polypeptide of paragraph 25, which is encoded by a polynucleotide comprising a nucleotide sequence having at least 97% sequence identity to the mature polypeptide coding sequence of SEQ ID NO: 1.

5 [27] The polypeptide of paragraph 1, which is encoded by a polynucleotide comprising or consisting of the nucleotide sequence of SEQ ID NO: 1; or a subsequence thereof encoding a fragment having beta-glucosidase activity.

[28] The polypeptide of paragraph 27, which is encoded by a polynucleotide comprising or consisting of the nucleotide sequence of SEQ ID NO: 1.

10 [29] The polypeptide of paragraph 1, which is encoded by a polynucleotide comprising or consisting of the mature polypeptide coding sequence of SEQ ID NO: 1.

[30] The polypeptide of paragraph 1, wherein the polypeptide is a variant comprising a substitution, deletion, and/or insertion of one or more (several) amino acids of the mature polypeptide of SEQ ID NO: 2.

15 [31] The polypeptide of paragraph 1, which is encoded by the polynucleotide contained in plasmid pAHYG-33 which is contained in *E. coli* NRRL B-50214.

[32] The polypeptide of any of paragraphs 1-31, wherein the mature polypeptide is amino acids 20 to 856 of SEQ ID NO: 2.

[33] The polypeptide of any of paragraphs 1-32, wherein the mature polypeptide coding sequence is nucleotides 58 to 2568 of SEQ ID NO: 1.

20 [34] An isolated polynucleotide comprising a nucleotide sequence that encodes the polypeptide of any of paragraphs 1-33.

[35] The isolated polynucleotide of paragraph 34, comprising at least one mutation in the mature polypeptide coding sequence of SEQ ID NO: 1, in which the mutant nucleotide sequence encodes the mature polypeptide of SEQ ID NO: 2.

25 [36] A nucleic acid construct comprising the polynucleotide of paragraph 34 or 35 operably linked to one or more (several) control sequences that direct the production of the polypeptide in an expression host.

[37] A recombinant expression vector comprising the polynucleotide of paragraph 34 or 35.

30 [38] A recombinant host cell comprising the polynucleotide of paragraph 34 or 35 operably linked to one or more (several) control sequences that direct the production of a polypeptide having alpha-glucuronidase activity.

[39] A method of producing the polypeptide of any of paragraphs 1-33, comprising: (a) cultivating a cell, which in its wild-type form produces the polypeptide, under conditions conducive for production of the polypeptide; and (b) recovering the polypeptide.

35 [40] A method of producing the polypeptide of any of paragraphs 1-33, comprising: (a) cultivating a host cell comprising a nucleic acid construct comprising a nucleotide

sequence encoding the polypeptide under conditions conducive for production of the polypeptide; and (b) recovering the polypeptide.

[41] A method of producing a mutant of a parent cell, comprising disrupting or deleting a polynucleotide encoding the polypeptide, or a portion thereof, of any of paragraphs 1-33, which results in the mutant producing less of the polypeptide than the parent cell.

[42] A mutant cell produced by the method of paragraph 41.

[43] The mutant cell of paragraph 42, further comprising a gene encoding a native or heterologous protein.

[44] A method of producing a protein, comprising: (a) cultivating the mutant cell of paragraph 43 under conditions conducive for production of the protein; and (b) recovering the protein.

[45] The isolated polynucleotide of paragraph 34 or 35, obtained by (a) hybridizing a population of DNA under at least medium stringency conditions with (i) the mature polypeptide coding sequence of SEQ ID NO: 1, (ii) the genomic DNA sequence comprising the mature polypeptide coding sequence of SEQ ID NO: 1, or (iii) a full-length complementary strand of (i) or (ii); and (b) isolating the hybridizing polynucleotide, which encodes a polypeptide having beta-glucosidase activity.

[46] The isolated polynucleotide of paragraph 45, obtained by (a) hybridizing a population of DNA under at least medium-high stringency conditions with (i) the mature polypeptide coding sequence of SEQ ID NO: 1, (ii) the genomic DNA sequence comprising the mature polypeptide coding sequence of SEQ ID NO: 1, or (iii) a full-length complementary strand of (i) or (ii); and (b) isolating the hybridizing polynucleotide, which encodes a polypeptide having beta-glucosidase activity.

[47] The isolated polynucleotide of paragraph 46, obtained by (a) hybridizing a population of DNA under at least high stringency conditions with (i) the mature polypeptide coding sequence of SEQ ID NO: 1, (ii) the genomic DNA sequence comprising the mature polypeptide coding sequence of SEQ ID NO: 1, or (iii) a full-length complementary strand of (i) or (ii); and (b) isolating the hybridizing polynucleotide, which encodes a polypeptide having beta-glucosidase activity.

[48] The isolated polynucleotide of paragraph 47, obtained by (a) hybridizing a population of DNA under at least very high stringency conditions with (i) the mature polypeptide coding sequence of SEQ ID NO: 1, (ii) the genomic DNA sequence comprising the mature polypeptide coding sequence of SEQ ID NO: 1, or (iii) a full-length complementary strand of (i) or (ii); and (b) isolating the hybridizing polynucleotide, which encodes a polypeptide having beta-glucosidase activity.

[49] The isolated polynucleotide of any of paragraphs 45-48, wherein the mature

polypeptide coding sequence is nucleotides 58 to 2568 of SEQ ID NO: 1.

[50] A method of producing a polynucleotide comprising a mutant nucleotide sequence encoding a polypeptide having beta-glucosidase activity, comprising: (a) introducing at least one mutation into the mature polypeptide coding sequence of SEQ ID NO: 1, wherein the mutant nucleotide sequence encodes a polypeptide comprising or consisting of the mature polypeptide of SEQ ID NO: 2; and (b) recovering the polynucleotide comprising the mutant nucleotide sequence.

[51] A mutant polynucleotide produced by the method of paragraph 50.

[52] A method of producing a polypeptide, comprising: (a) cultivating a cell comprising the mutant polynucleotide of paragraph 51 encoding the polypeptide under conditions conducive for production of the polypeptide; and (b) recovering the polypeptide.

[53] A method of producing the polypeptide of any of paragraphs 1-33, comprising: (a) cultivating a transgenic plant or a plant cell comprising a polynucleotide encoding the polypeptide under conditions conducive for production of the polypeptide; and (b) recovering the polypeptide.

[54] A transgenic plant, plant part or plant cell transformed with a polynucleotide encoding the polypeptide of any of paragraphs 1-33.

[55] A double-stranded inhibitory RNA (dsRNA) molecule comprising a subsequence of the polynucleotide of paragraph 34 or 35, wherein optionally the dsRNA is a siRNA or a miRNA molecule.

[56] The double-stranded inhibitory RNA (dsRNA) molecule of paragraph 55, which is about 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25 or more duplex nucleotides in length.

[57] A method of inhibiting the expression of a polypeptide having beta-glucosidase activity in a cell, comprising administering to the cell or expressing in the cell a double-stranded RNA (dsRNA) molecule, wherein the dsRNA comprises a subsequence of the polynucleotide of paragraph 34 or 35.

[58] The method of paragraph 57, wherein the dsRNA is about 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25 or more duplex nucleotides in length.

[59] An isolated polynucleotide encoding a signal peptide comprising or consisting of amino acids 1 to 19 of SEQ ID NO: 2.

[60] A nucleic acid construct comprising a gene encoding a protein operably linked to the polynucleotide of paragraph 59, wherein the gene is foreign to the polynucleotide encoding the signal peptide.

[61] A recombinant expression vector comprising a gene encoding a protein operably linked to the polynucleotide of paragraph 59, wherein the gene is foreign to the polynucleotide encoding the signal peptide.

[62] A recombinant host cell comprising a gene encoding a protein operably linked to

the polynucleotide of paragraph 59, wherein the gene is foreign to the polynucleotide encoding the signal peptide.

[63] A method of producing a protein, comprising: (a) cultivating a recombinant host cell comprising a gene encoding a protein operably linked to the polynucleotide of paragraph 59, wherein the gene is foreign to the polynucleotide encoding the signal peptide, under conditions conducive for production of the protein; and (b) recovering the protein.

[64] A composition comprising the polypeptide of any of paragraphs 1-33.

[65] A method for degrading or converting a cellulosic material, comprising: treating the cellulosic material with an enzyme composition in the presence of the polypeptide having beta-glucosidase activity of any of paragraphs 1-33.

[66] The method of paragraph 65, wherein the cellulosic material is pretreated.

[67] The method of paragraph 65 or 66, wherein the enzyme composition comprises one or more cellulolytic enzymes selected from the group consisting of an endoglucanase, a cellobiohydrolase, and a beta-glucosidase.

[68] The method of any of paragraphs 65-67, wherein the enzyme composition further comprises a polypeptide having cellulolytic enhancing activity.

[69] The method of any of paragraphs 65-68, wherein the enzyme composition further comprises one or more enzymes selected from the group consisting of a xylanase, a hemicellulase, an esterase, a protease, a laccase, or a peroxidase.

[70] The method of any of paragraphs 65-69, further comprising recovering the degraded cellulosic material.

[71] The method of paragraph 70, wherein the degraded cellulosic material is a sugar.

[72] The method of paragraph 71, wherein the sugar is selected from the group consisting of glucose, xylose, mannose, galactose, and arabinose.

[73] A method for producing a fermentation product, comprising: (a) saccharifying a cellulosic material with an enzyme composition in the presence of the polypeptide having beta-glucosidase activity of any of paragraphs 1-33; (b) fermenting the saccharified cellulosic material with one or more fermenting microorganisms to produce the fermentation product; and (c) recovering the fermentation product from the fermentation.

[74] The method of paragraph 73, wherein the cellulosic material is pretreated.

[75] The method of paragraph 73 or 74, wherein the enzyme composition comprises one or more cellulolytic enzymes selected from the group consisting of an endoglucanase, a cellobiohydrolase, and a beta-glucosidase.

[76] The method of any of paragraphs 73-75, wherein the enzyme composition further comprises a polypeptide having cellulolytic enhancing activity.

[77] The method of any of paragraphs 73-76, wherein the enzyme composition

further comprises one or more enzymes selected from the group consisting of a xylanase, a hemicellulase, an esterase, a protease, a laccase, or a peroxidase.

[78] The method of any of paragraphs 73-77, wherein steps (a) and (b) are performed simultaneously in a simultaneous saccharification and fermentation.

5 [79] The method of any of paragraphs 73-78, wherein the fermentation product is an alcohol, an organic acid, a ketone, an amino acid, or a gas.

[80] A method of fermenting a cellulosic material, comprising: fermenting the cellulosic material with one or more fermenting microorganisms, wherein the cellulosic material is saccharified with an enzyme composition in the presence of a polypeptide having
10 beta-glucosidase activity of any of paragraphs 1-33.

[81] The method of paragraph 80, wherein the fermenting of the cellulosic material produces a fermentation product.

[82] The method of paragraph 81, further comprising recovering the fermentation product from the fermentation.

15 [83] The method of any of paragraphs 80-82, wherein the cellulosic material is pretreated before saccharification.

[84] The method of any of paragraphs 80-83, wherein the enzyme composition comprises one or more cellulolytic enzymes selected from the group consisting of an endoglucanase, a cellobiohydrolase, and a beta-glucosidase.

20 [85] The method of any of paragraphs 80-84, wherein the enzyme composition further comprises a polypeptide having cellulolytic enhancing activity.

[86] The method of any of paragraphs 80-85, wherein the enzyme composition further comprises one or more enzymes selected from the group consisting of a xylanase, a hemicellulase, an esterase, a protease, a laccase, or a peroxidase.

25 [88] The method of any of paragraphs 80-86, wherein the fermentation product is an alcohol, an organic acid, a ketone, an amino acid, or a gas.

The invention described and claimed herein is not to be limited in scope by the specific aspects herein disclosed, since these aspects are intended as illustrations of several
30 aspects of the invention. Any equivalent aspects are intended to be within the scope of this invention. Indeed, various modifications of the invention in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description. Such modifications are also intended to fall within the scope of the appended claims. In the case of conflict, the present disclosure including definitions will control.

35

Claims

What is claimed is:

- 5 1. An isolated polypeptide having beta-glucosidase activity, selected from the group consisting of:
- (a) a polypeptide comprising an amino acid sequence having at least 60% sequence identity to the mature polypeptide of SEQ ID NO: 2;
 - 10 (b) a polypeptide encoded by a polynucleotide that hybridizes under at least medium stringency conditions with (i) the mature polypeptide coding sequence of SEQ ID NO: 1, (ii) the genomic DNA sequence comprising the mature polypeptide coding sequence of SEQ ID NO: 1, or (iii) a full-length complementary strand of (i) or (ii);
 - 15 (c) a polypeptide encoded by a polynucleotide comprising a nucleotide sequence having at least 60% sequence identity to the mature polypeptide coding sequence of SEQ ID NO: 1; and
 - (d) a variant comprising a substitution, deletion, and/or insertion of one or more (several) amino acids of the mature polypeptide of SEQ ID NO: 2.
2. The polypeptide of claim 1, comprising or consisting of the amino acid sequence of 20 SEQ ID NO: 2; or a fragment thereof having beta-glucosidase activity.
3. The polypeptide of claim 1, which is encoded by the polynucleotide contained in plasmid pAHYG-33 which is contained in *E. coli* NRRL B-50214.
- 25 4. An isolated polynucleotide comprising a nucleotide sequence that encodes the polypeptide of any of claims 1-3.
- 30 5. A method of producing the polypeptide of any of claims 1-3, comprising: (a) cultivating a cell, which in its wild-type form produces the polypeptide, under conditions conducive for production of the polypeptide; and (b) recovering the polypeptide.
- 35 6. A method of producing the polypeptide of any of claims 1-3, comprising: (a) cultivating a host cell comprising a nucleic acid construct comprising a nucleotide sequence encoding the polypeptide under conditions conducive for production of the polypeptide; and (b) recovering the polypeptide.
7. A method of producing a mutant of a parent cell, comprising disrupting or deleting a

polynucleotide encoding the polypeptide, or a portion thereof, of any of claims 1-3, which results in the mutant producing less of the polypeptide than the parent cell.

5 8. A method of producing the polypeptide of any of claims 1-3, comprising: (a) cultivating a transgenic plant or a plant cell comprising a polynucleotide encoding the polypeptide under conditions conducive for production of the polypeptide; and (b) recovering the polypeptide.

10 9. A transgenic plant, plant part or plant cell transformed with a polynucleotide encoding the polypeptide of any of claims 1-3.

15 10. A double-stranded inhibitory RNA (dsRNA) molecule comprising a subsequence of the polynucleotide of claim 4, wherein optionally the dsRNA is a siRNA or a miRNA molecule.

20 11. A method of inhibiting the expression of a polypeptide having beta-glucosidase activity in a cell, comprising administering to the cell or expressing in the cell a double-stranded RNA (dsRNA) molecule, wherein the dsRNA comprises a subsequence of the polynucleotide of claim 4.

25 12. An isolated polynucleotide encoding a signal peptide comprising or consisting of amino acids 1 to 19 of SEQ ID NO: 2.

30 13. A method of producing a protein, comprising: (a) cultivating a recombinant host cell comprising a gene encoding a protein operably linked to the polynucleotide of claim 12, wherein the gene is foreign to the polynucleotide encoding the signal peptide, under conditions conducive for production of the protein; and (b) recovering the protein.

35 14. A composition comprising the polypeptide of any of claims 1-3.

15. A method for degrading or converting a cellulosic material, comprising: treating the cellulosic material with an enzyme composition in the presence of the polypeptide having beta-glucosidase activity of any of claims 1-3.

16. The method of claim 15, further comprising recovering the degraded cellulosic material.

17. A method for producing a fermentation product, comprising: (a) saccharifying a

cellulosic material with an enzyme composition in the presence of the polypeptide having beta-glucosidase activity of any of claims 1-3; (b) fermenting the saccharified cellulosic material with one or more fermenting microorganisms to produce the fermentation product; and (c) recovering the fermentation product from the fermentation.

5

18. A method of fermenting a cellulosic material, comprising: fermenting the cellulosic material with one or more fermenting microorganisms, wherein the cellulosic material is saccharified with an enzyme composition in the presence of a polypeptide having beta-glucosidase activity of any of claims 1-3.

10

19. The method of claim 18, wherein the fermenting of the cellulosic material produces a fermentation product.

15

20. The method of claim 19, further comprising recovering the fermentation product from the fermentation.

1/3

M F G I T S T A L L L A T A L L A S S T
 1 ATGTTTGGCATCACCTCGACTGCGCTGCTGCTTGCCACTGCGCTCCTTGCTTCTTCGACG
 P L E V R Q S G L S T A S P P W Y P A P
 61 CCCCTCGAAGTCCGTCAGTCCGGCTTGTCCACTGCCAGCCCGCCATGGTACCCTGCTCCC
 H G G T D R L W A Q S Y D K A A K L V S
 121 CATGGAGGAACCGATAGGCTGTGGGCTCAGTCCTATGACAAGGCAGCGAAGCTGGTGTGCG
 Q M T L A E K V N I T T G T G W S M G P
 181 CAGATGACGCTGGCCGAGAAGGTGAACATCACCACCGGTACCGGATGGTCCATGGGCCCA
 C V G N T G P V P R L G F P S L C L Q D
 241 TGCGTAGGCAACACTGGACCCGTTCCCGTCTCGGCTTCCCGTCTCTGTGTCTGCAGGAT
 G P L G V R F A D K I T S F P A G I T T
 301 GGTCTCTCGGTGTCCGCTTTGCCGACAAGATCACATCGTTCCCGCTGGAATCACCCT
 G G T W D K D L M R L R G E A I G A E N
 361 GGAGGTACTGGGACAAGGACCTTATGCGGCTTCGTGGTGAAGCAATCGGTGCGGAGAAC
 R G K G V H I T L G P A M G P L G K H P
 421 CGTGGAAAGGGAGTTACATCACTTTGGGTCCCGCTATGGGTCTCTGGGAAAGCACCCC
 A G G R N W E G F G S D P Y L Q G V A A
 481 GCTGGTGGAAAGAACTGGGAAGGTTTCGGATCCGATCCTTACCTCCAGGGCGTTGCTGCC
 Y E S I L G I Q S Q G V V A T A K H Y I
 541 TACGAGAGCATCCTTGGAAATCCAGTCTCAGGGTGTCTGCTACCGCGAAGCACTACATC
 G N E Q E H F R Q G P N A I S A N I D D
 601 GGTAACTGAGCAGGAACACTTCCGCCAGGGCCCAATGCCATCTCCGCGAACATTGACGAC
 R T L H E V Y L W P F Q D S V K A G V G
 661 CGAACTCTGCACGAAGTCTACCTCTGGCCATTCCAGGACAGTGTCAAGGCTGGCGTTGGC
 A V M C S Y N Q V N N S A A C Q N S W L
 721 GCGGTATGTGCTCCTACAACCAAGTCAACAACCTCCGCCGCTTGTGCAAGCACTTGGGTC
 Q N N V L K D Q L G F Q G F I M S D W L
 781 CAGAACAACGTGCTCAAGGACCAACTCGGTTTCCAGGGTTTCATCATGAGTGACTGGCTG
 A Q M S G V A S V L A G M D M T M P G D
 841 GCGCAGATGTCTGGTGTGCTTCTGTCTCGCTGGAATGGACATGACCATGCCTGGTGAT
 G L N W D D R V P L L G P R L T E A V L
 901 GGTTTGAACTGGGATGATCGTGTGCCCTTGTGGGCCACGTCTGACCGAGGCCGTGCTG
 N S S V P I D R L D D M V T R I V A T W
 961 AACTCTTCCGTCCTATCGACCGTCTGGATGACATGGTAACCCGTATCGTTGCCACTTGG
 Y Q M G G Q D K N H P E P N F S S W T K N
 1021 TACCAGATGGCCAGGACAAGAACCACCCGAGCCCAACTCAGCTCATGGACCAAGAAC
 T T G N I Y F G S G E G P E G I V N Q H
 1081 ACCACCGGCAACATTTACTTCCGGCTCTGGCGAGGGACCGGAAGGAATTGTCAACCAGCAC
 V D V M A D H P T A V A R K V A A D A I
 1141 GTCGACGTGATGGCCGACCACCCACCGCTGTGGCCCCAAAGGTTGCGGCCGATGCCATC
 A L L K N D K N T L P L K K P M K I G V
 1201 GCTCTGCTCAAGAATGACAAGAACCCTTCCGCTCAAGAAGCCGATGAAGATCGGTGTC
 Y G E D A A L A P G G A N V C A D R G C
 1261 TACGGTGAGGATGCTGCGCTTGTCTCCCGAGGTGCCAACGTCTGTGCCGACCGTGGCTGC
 N E G T L A V G W G S G S T D F E Y L V
 1321 AACGAGGGCACTTGTCCGTCGGATGGGGCAGTGGCTCTACCGACTTTGAGTACCTTGTT
 D P L S A I T R Q A Q S Y G G K V T A I
 1381 GATCCTCTCTCGCCATCACCAGGCAGGCCAGTCCTACGGTGGCAAGGTTACCGCAATC
 L Q N N A T D I I A A S A K K Q D V C L
 1441 CTCCAGAACAATGCGACCGACATCATTGCCGCTCGGCCAAGAAGCAGGACGCTGTGCTC
 V F V N A D A G E G Y I Q W E D V A G D
 1501 GTTTTCGTGAACGCCGATGCCGGTGAGGGTTACATTAGTGGGAAGACGTGGCGGGCGAC
 R P N L I T Q K G G D D L V L A V A N N
 1561 CGTCCCAACCTGATCAGCAGAAGGGCGGTGATGACCTCGTGTGCTGTTGCCAACAAC
 C D N T V V V V H S V G P I I M E K W A
 1621 TGCGACAATACTGTTGTCGTCGTCCACAGCGTCGGCCCCGATCATGAGAAATGGGCC
 D K R Q V R S I L W A H L P G M E S G N

Fig. 1A

2/3

1681 GACAAGCGTCAAGTCCGTTTCGATTCTCTGGGCACACCTTCCGGGCATGGAAAGCGGAAAC
A L V D V L W G S T N P S G K L P Y T I
1741 GCCCTGTGCGATGTTCTTTGGGGCTCCACCAACCCCTCCGGCAAGCTCCCTTACACCATC
G K S L A D Y G P A A G V L Y T E N A Q
1801 GGAAAGTCACTCGCCGACTATGGCCCTGCTGCTGGCGTTCTCTACACCGAGAACGCACAG
P P Q Q D F T D G V F I D Y R H F D A N
1861 CCTCCTCAGCAAGACTTCACCGACGGCGTCTTCATCGACTACCGCCACTTCGACGCCAAC
K I E P R Y E F G F G L S Y T T F S Y S
1921 AAGATTGAGCCCCGTTACGAGTTTGGCTTCGGATTGTCGTACACTACCTTCTCGTACAGC
N L V K V K K G A F T P L P A P R P K D
1981 AACCTCGTCAAGGTCAAGAAGGGAGCCTTCACTCCTCTCCCTGCGCCTAGGCCGAAGGAC
S A T P P K Y D S K L P K P E E C T F P
2041 AGTGCCACCCCTCCGAAGTACGACTCCAAGCTCCCCAAGCCTGAAGAGTGCACCTTCCCC
D G F K K I N R M I Y P Y L E S A D T V
2101 GATGGGTTCAAGAAGATCAACAGAATGATTTACCCCTACCTGGAGTCTGCGGACACCGTC
K V G P Y P Y P E G Y D T K Q T P S Q A
2161 AAGGTCGGCCCATACCCCTACCCCGAGGGCTACGACACCAAGCAAACCTCCCTCGCAAGCC
G G A Q G G N P A L W E V L A E V S V T
2221 GCGGGCGCTCAGGGTGGTAACCCTGCCCTCTGGGAGGTGCTGGCTGAGGTCTCGGTACA
V K N T G K V A G A E V A Q L Y L G F P
2281 GTGAAGAACACTGGAAAGGTGCGGGTTCGCGAGGTTCGCGCAGTTGTACCTTGGATTCCCC
Q N G P V P F P P K Q L R G F E K V F L
2341 CAGAATGGACCCGTTCCGTTCCCGCCCAAGCAGCTTAGAGGATTTGAGAAGGTCTTCTTG
Q P G E S K R V T F P L T R R D L S Y W
2401 CAGCCCGGTGAGAGTAAGCGCGTGACATCCCACTCACTAGGAGGGATCTTAGCTACTGG
D V T K Q N W V I P K G G F G V M V G T
2461 GATGTTACCAAGCAGAAGTGGGTCATCCCCAAGGGCGGATTTGGAGTTATGGTTGGAAC
S S R K I K A Q G Y I P S D H Y *
2521 TCGAGTAGGAAGATTAAGGCACAGGGTTATATCCCGAGTGATCATTATTGA

Fig. 1B

3/3




Fig. 2

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2010/022405

A. CLASSIFICATION OF SUBJECT MATTER

INV. C12N9/42 C12N15/82 C12P19/14

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C12N C12P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, Sequence Search, WPI Data, BIOSIS, EMBASE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2007/019442 A2 (NOVOZYMES INC [US]; NOVOZYMES AS [DK]; KROGH KRISTIAN [DK]; HARRIS PAU) 15 February 2007 (2007-02-15)	1,4-9, 14-20
Y	claims; sequence 1	2,3
X	WO 2008/148131 A1 (NOVOZYMES INC [US]; LOPEZ DE LEON ALFREDO [US]; DING HANSHU [US]; BROW) 4 December 2008 (2008-12-04)	1,4-11, 14-20
Y	page 2, line 22 - page 4, line 14; claims 18-20	2,3
X	WO 2008/151079 A2 (NOVOZYMES INC [US]; MERINO SANDRA [US]; MCFARLAND KEITH [US]; CHERRY J) 11 December 2008 (2008-12-11)	1,4-6, 14-20
Y	page 46, paragraph 1; claims; sequence 21	2,3,7-11
	-/--	

 Further documents are listed in the continuation of Box C. See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

6 April 2010

Date of mailing of the international search report

24/06/2010

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Huber, Angelika

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2010/022405

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X Y	US 2004/091469 A1 (FUKASAWA TOMOYUKI [JP] ET AL) 13 May 2004 (2004-05-13) claims; sequence 2	1,4-6, 14-20 2,3,7-11
X,P Y	----- WO 2009/085864 A2 (NOVOZYMES AS [DK]; HARRIS PAUL [US]; MAIYURAN SUCHINDRA [US]; BROWN KI) 9 July 2009 (2009-07-09) page 18, lines 18-21; claims; examples -----	1,4-11, 14-20 2,3

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US2010/022405

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.

2. As all searchable claims could be searched without effort justifying an additional fees, this Authority did not invite payment of additional fees.

3. As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

see additional sheet(s)

Remark on Protest

- The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. claims: 1-11, 14-20

Polypeptide having beta-glucosidase activity comprising an amino acid sequence having at least 60% sequence identity to the mature polypeptide of SEQ ID NO: 2; nucleic acid encoding said polypeptide; method of producing said polypeptide, method of producing a mutant of a parent cell, transgenic plant transformed with the polynucleotide, dsRNA molecule comprising a subsequence of the polynucleotide, method of inhibiting the expression of a polypeptide with said dsRNA molecule, method of degrading or converting a cellulosic material, method of producing a fermentation product

2. claims: 12, 13

Isolated polynucleotide encoding a signal peptide comprising amino acids 1-19 of SEQ ID NO: 2; method of producing a protein comprising cultivating a recombinant host cells comprising a heterologous gene linked to said signal peptide

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No PCT/US2010/022405

Patent document cited in search report	Publication date	Patent family member(s)	Publication date	
WO 2007019442	A2	15-02-2007	CA 2625933 A1	15-02-2007
			CN 101287751 A	15-10-2008
			EP 1928901 A2	11-06-2008
			JP 2009502209 T	29-01-2009
			US 2008201805 A1	21-08-2008
WO 2008148131	A1	04-12-2008	CA 2687609 A1	04-12-2008
			EP 2069489 A1	17-06-2009
			US 2009019608 A1	15-01-2009
WO 2008151079	A2	11-12-2008	AU 2008259937 A1	11-12-2008
			AU 2008259986 A1	11-12-2008
			CA 2689261 A1	11-12-2008
			CA 2689910 A1	11-12-2008
			EP 2064323 A1	03-06-2009
			EP 2069492 A2	17-06-2009
			KR 20100020977 A	23-02-2010
			US 2010129860 A1	27-05-2010
WO 2008151043 A1	11-12-2008			
US 2004091469	A1	13-05-2004	AU 9229201 A	08-04-2002
			WO 0226979 A1	04-04-2002
WO 2009085864	A2	09-07-2009	NONE	