



US 20200063147A1

(19) **United States**

(12) **Patent Application Publication**

Zhang et al.

(10) **Pub. No.: US 2020/0063147 A1**

(43) **Pub. Date: Feb. 27, 2020**

(54) **CRISPR-CAS COMPONENT SYSTEMS, METHODS AND COMPOSITIONS FOR SEQUENCE MANIPULATION**

plication No. 61/791,409, filed on Mar. 15, 2013, provisional application No. 61/835,931, filed on Jun. 17, 2013.

(71) Applicants: **The Broad Institute, Inc.**, Cambridge, MA (US); **Massachusetts Institute of Technology**, Cambridge, MA (US); **President and Fellows of Harvard College**, Cambridge, MA (US)

(72) Inventors: **Feng Zhang**, Cambridge, MA (US); **Le Cong**, Cambridge, MA (US); **David Benjamin Turitz Cox**, Cambridge, MA (US); **Patrick Hsu**, Cambridge, MA (US); **Shuailiang Lin**, Cambridge, MA (US); **Fei Ran**, Boston, MA (US); **Randall Jeffrey Platt**, Cambridge, MA (US); **Neville Espi Sanjana**, Cambridge, MA (US)

(21) Appl. No.: **16/532,442**

(22) Filed: **Aug. 5, 2019**

Related U.S. Application Data

(63) Continuation of application No. 15/230,161, filed on Aug. 5, 2016, which is a continuation of application No. 14/105,035, filed on Dec. 12, 2013, now abandoned.

(60) Provisional application No. 61/736,527, filed on Dec. 12, 2012, provisional application No. 61/748,427, filed on Jan. 2, 2013, provisional application No. 61/768,959, filed on Feb. 25, 2013, provisional ap-

Publication Classification

(51) **Int. Cl.**

C12N 15/74 (2006.01)
C12N 15/113 (2006.01)
C12N 9/22 (2006.01)
C12N 15/63 (2006.01)
C12N 15/85 (2006.01)
C12N 15/70 (2006.01)
C12N 15/90 (2006.01)

(52) **U.S. Cl.**

CPC *C12N 15/746* (2013.01); *C12N 15/113* (2013.01); *C12N 9/22* (2013.01); *C12N 15/63* (2013.01); *C12N 15/85* (2013.01); *C12N 15/70* (2013.01); *C12N 15/102* (2013.01); *C12N 15/8509* (2013.01); *C12N 15/907* (2013.01); *C12N 2310/3519* (2013.01); *C12N 2310/531* (2013.01); *C12N 2310/20* (2017.05); *C12N 15/74* (2013.01)

(57)

ABSTRACT

The invention provides for systems, methods, and compositions for manipulation of sequences and/or activities of target sequences. Provided are vectors and vector systems, some of which encode one or more components of a CRISPR complex, as well as methods for the design and use of such vectors. Also provided are methods of directing CRISPR complex formation in eukaryotic cells and methods for selecting specific cells by introducing precise mutations utilizing the CRISPR/Cas system.

Specification includes a Sequence Listing.




FIG. 1




FIG. 2A




FIG. 2B




FIG. 2D

FIG. 2E

Target locus 5' - . . . AGCTGGAGGAGGAGGGCCCTGAGTCCGAGCAGAAAGAAGGGCTCCCAC. . . -3'
 PAM
 |||||
 3' - . . . TCGACCTCCTCCCTCCGGACTCAGGGCTGGTCTTCTTCCCGAGGGTG. . . -5'
 |||||
 crRNA 5' - GAGUCCGAGCAGAAAGAAGUUUAGAGC. . . -3'




FIG. 2F

human EMX1 protospacer target (mutation in 5 of 43 sequenced clones = 11.6%)
 WT 5' - . . . CTGGAGGAGGAAAGGGCCCTGAGTCCGAGCAGAAAGAAGGGCTCCCACATC. . . -3'
 Δ1 CTGGAGGAGGAAAGGGCCCTGAGTCCGAGCAGAAAGAAGGGCTCCCACATC
 +1 CTGGAGGAGGAAAGGGCCCTGAGTCCGAGCAGAAAGAAGGGCTCCCACATC
 Δ3 CTGGAGGAGGAAAGGGCCCTGAGTCCGAGCAGAAAGAAGGGCTCCCACATC
 m1, Δ6 CTGGAGGAGGAAAGGGCCCTGAGCCCGAGCAGAAAG-----GGCTCCCACATC

FIG. 4A




FIG. 4B





FIG. 4C






FIG. 4D

FIG. 5A




FIG. 5B




FIG. 5C




FIG. 5D





FIG. 5E






FIG. 5F




FIG. 5G

Cas9	target species	gene	protospacer ID	protospacer sequence (5' to 3')	PAM	strand	cell line tested	% indel (pre-crRNA + tracerRNA)	% indel (chimeric RNA)
<i>S. pyogenes</i> SF370 type II			1	GGAGGGCTCGATCCGAGCAGAGAGAA	GGG	+	293FT	20 ± 1.8	6.7 ± 0.62
			2	CATTGGAGTGCATCGATGCTCTCCCAT	TGG	-	293FT	2.1 ± 0.31	N.D.
			3	GGACATCGATGTACCTCANTGACTAGCG	TGG	+	293FT	14 ± 1.1	N.D.
			4	CATCGATGCTCCGCTTCCGCTGCTTGG	TGG	-	293FT	11 ± 1.7	N.D.
			5	TTCGTGGCANTGCCACCCGCTTGATGTGA	TGG	-	293FT	4.3 ± 0.46	2.1 ± 0.51
			6	TGCTGGCANTGCCACCCGCTTGATGTGAT	GGG	-	293FT	4.0 ± 0.69	0.41 ± 0.25
			7	TCCAGCTTCTGCGTTTGTACTTTGTCCTC	TGG	-	293FT	1.5 ± 0.12	N.D.
			8	GGAGGGAGGGCCACAGATGAGAACTCAGG	AGG	-	293FT	7.8 ± 0.63	2.3 ± 1.2
CRISPR	<i>Homo sapiens</i>		9	AGAGGCCGAGATTGGTGTTCAGGGCAGAG	AGG	+	293FT	21 ± 2.6	6.5 ± 0.32
			10	ATCCAGAGGTGCGCAGAGGGGCGAGAT	TGG	+	293FT	N.D.	N.D.
			11	GGTGGCAGAGGGCCGAGATTGGGTGTC	AGG	+	293FT	N.D.	N.D.
<i>Mus musculus</i>			12	CAGCCACTGAGTGGCATTACTAATTCAT	AGG	-	Neuro2A	27 ± 4.3	4.1 ± 2.2
			13	AATGCATAGGATACCCACCCAGGTGCCAG	GGG	-	Neuro2A	4.8 ± 1.2	N.D.
			14	ACACACATGGGAAAGCCCTTGGGCAAGAA	AGG	+	Neuro2A	11.3 ± 1.3	N.D.
			15	GGAGGGGTAGTATACGAAACACAGAGA	GTAGCAT	-	293FT	14 ± 0.88	N.T.
LMO-9 CRISPR1	<i>Homo sapiens</i>	EMX1	16	AGATGTAGAGGATCAGCAAACTCAGCA	CTTAGAA	-	293FT	7.8 ± 0.77	N.T.

FIG. 6

FIG. 7A





FIG. 7B






FIG. 7C




FIG. 8

FIG. 9A




FIG. 9B




FIG. 10A




FIG. 10B




FIG. 10C




FIG. 10D




FIG. 11C

Chr	NGG		NNAGAAW	
	median	mean	median	mean
1	7	12.8	67	115.8
2	8	12.7	64	100.8
3	8	13.0	63	98.5
4	9	14.0	61	94.5
5	8	13.1	63	97.9
6	8	13.1	63	98.5
7	8	12.4	64	102.9
8	8	12.8	64	100.9
9	7	13.9	65	120.5
10	7	12.1	66	107.0
11	7	12.0	65	105.8
12	8	12.4	65	103.5
13	8	13.6	62	94.8
14	8	12.0	65	101.5
15	7	11.5	68	107.7
16	7	11.7	74	136.8
17	6	10.3	76	127.9
18	8	13.4	63	101.8
19	6	9.4	82	145.4
20	7	11.1	72	121.8
21	7	13.4	64	111.4
22	6	9.2	85	140.3
X	6	13.2	63	99.0
Y	8	29.2	62	223.7

FIG. 11A




FIG. 11B




FIG. 12A




FIG. 12B





FIG. 12C







FIG. 15A




FIG. 15B






FIG. 16

Primer name	Assay	Genomic Target	Primer sequence
Sp-EMX1-F	SURVEYOR assay, sequencing	<i>EMX1</i>	AAAACCACCCCTTCTCTCTGGC
Sp-EMX1-R	SURVEYOR assay, sequencing	<i>EMX1</i>	GGAGATTGGAGACACGGAGAG
Sp-PVALB-F	SURVEYOR assay, sequencing	<i>PVALB</i>	CTGGAAAGCCAATGCCTGAC
Sp-PVALB-R	SURVEYOR assay, sequencing	<i>PVALB</i>	GGCAGCAAACCTCCTTGTCCT
Sp-Th-F	SURVEYOR assay, sequencing	<i>Th</i>	GTGCTTTGCAGAGGCTTACC
Sp-Th-R	SURVEYOR assay, sequencing	<i>Th</i>	CCTGGAGCGCATGCAGTAGT
St-EMX1-F	SURVEYOR assay, sequencing	<i>EMX1</i>	ACCTTCTGTGTTTCCACCATTCC
St-EMX1-R	SURVEYOR assay, sequencing	<i>EMX1</i>	TTGGGGAGTGCACAGACTTC
Sp-EMX1- RFLP-F	RFLP, sequencing	<i>EMX1</i>	GGCTCCCTGGGTTCAAAGTA
Sp-EMX1- RFLP-R	RFLP, sequencing	<i>EMX1</i>	AGAGGGGTCTGGATGTCGTAA
Pb_EMX1_sp1	Northern Blot Probe	Not applicable	TAGCTCTAAAACCTTCTTCTTCTGCTCGGAC
Pb_tracrRNA	Northern Blot Probe	Not applicable	CTAGCCTTATTTTAACTTGCTATGCTGTTT

FIG. 17




FIG. 18A




FIG. 18B




FIG. 18C

FIG. 19B





FIG. 19A






FIG. 21

FIG. 23A




FIG. 23B

Editing Construct	cfu/ug of crR6M DNA	Kan ^r transformants	
		Ed. Un.	
None	2.6 ± 0.9		
R6(<i>srtA</i>)	14 ± 1.8		
R6 ^{370.1}	25 ± 2.6		

FIG. 24A




FIG. 24B

		2 nd PAM position			
		A	C	G	T
1 st PAM position	A	AAA 1.02	ACA 1.06	AGA 0.95	ATA 1.02
	C	CAB 1.03	CCA 1.03	CGA 0.97	CTA 1.00
	G	GAA 1.10	GCA 1.09	GGA 0.93	GTA 1.01
	T	TAA 1.02	TCA 1.11	TGA 0.91	TTA 1.11
1 st PAM position	A	AAC 1.07	ACC 1.02	AGC 1.00	ATC 1.09
	C	CAC 1.06	CCC 1.04	CGC 1.00	CTC 1.08
	G	GAC 1.05	GCC 1.02	GGC 1.02	GTC 1.22
	T	TAC 1.05	TCC 1.05	TGC 1.03	TTC 1.06
1 st PAM position	A	AAU 0.99	ACU 0.97	AGU 0.96	AU 0.96
	C	CAU 0.99	CCU 1.08	CGU 0.98	CU 1.03
	G	GAU 0.99	GCU 1.08	GU 0.98	GU 1.20
	T	TAU 0.95	TGU 0.91	TU 0.97	TU 1.06
1 st PAM position	A	AAT 1.01	ACT 1.09	AGT 1.04	ATT 1.03
	C	CAT 1.03	CCT 1.03	CGT 1.07	CTT 1.08
	G	GAT 1.04	GCT 1.08	GT 1.08	GT 1.12
	T	TAT 1.06	TCT 1.03	TGT 1.06	TTT 1.01

FIG. 24C




FIG. 25A




FIG. 25B




FIG. 25C




FIG. 25D





FIG. 25E



FIG. 25F

FIG. 26A




FIG. 26B




FIG. 26C




FIG. 26D




FIG. 27A




FIG. 27B

	PAM	Protospacer
<i>rpsL</i>	TCCTA	AAAAACCGAAGCTCCGGCCCTGGCGTAAAGTA
	P	K K P N S A L R K V
<i>rpsL</i> mut	TCCTA	AAAAACCGAAGCTCCGGCCCTGGCGTAAAGTA
	P	T K P N S A L R K V




FIG. 27C




FIG. 27D

FIG. 28A





FIG. 28B






FIG. 30A




FIG. 30B





FIG. 30C






FIG. 31




FIG. 32

FIG. 33A




FIG. 33B




FIG. 33C





FIG. 33D

FIG. 34A




FIG 34B




FIG. 34C





FIG. 34D

FIG. 35A




FIG. 35B




FIG. 35C




FIG. 36




FIG. 37

FIG. 38A





FIG. 38B






FIG. 40B




FIG. 40C




FIG. 40E




FIG. 40F

SpCas9 mutation positions

hSpCas9




FIG. 41A




FIG. 41B




FIG. 41C




FIG. 41D




FIG. 41E




FIG. 41F




FIG. 41G




FIG. 41H




FIG. 411




FIG. 41J




FIG. 41K




FIG. 41L




FIG. 41M




FIG. 42




FIG. 43A




FIG. 43B




FIG. 43C




FIG. 43D




FIG. 44A




FIG. 44B




FIG. 45




FIG. 46




FIG. 47

FIG. 48A





FIG. 48B



Conditional Cas9, Rosa26 targeting vector map




FIG. 49A

Constitutive Cas9, Rosa26 targeting vector map




FIG. 49B

ATCCAGCTGGTGCAGACCTACAACCAGCTGTTCGAGGAAAACCCCATCAACGCCAG
CGGCGTGGACGCCAAGGCCATCCTGTCTGCCAGACTGAGCAAGAGCAGACGGCTGG
AAAATCTGATCGCCCAGCTGCCCGGCGAGAAGAAGAATGGCCTGTTCCGgAACCTG
ATTGCCCTGAGCCTGGGCCTGACCCCCAACTTCAAGAGCAACTTCGACCTGGCCGAG
GATGCCAAACTGCAGCTGAGCAAGGACACCTACGACGACGACCTGGACAACCTGCT
GGCCCAGATCGGCGACCAGTACGCCGACCTGTTTCTGGCCGCCAAGAACCTGTCCG
ACGCCATCCTGCTGAGCGACATCCTGAGAGTGAACACCGAGATCACCAAGGCCCCC
CTGAGCGCCTCTATGATCAAGAGATACGACGAGCACCACCAGGACCTGACCCTGCT
GAAAGCTCTCGTGC GG CAGCAGCTGCCTGAGAAGTACAAAGAGATTTTCTTCGACC
AGAGCAAGAACGGCTACGCCGGCTACATTGACGGCGGAGCCAGCCAGGAAGAGTTC
TACAAGTTCATCAAGCCCATCCTGGAAAAGATGGACGGCACCGAGGAACCTGCTCGT
GAAGCTGAACAGAGAGGACCTGCTGCGGAAGCAGCGGACCTTCGACAACGGCAGC
ATCCCCCACCAGATCCACCTGGGAGAGCTGCACGCCATTCTGCGGCGGCAGGAAGA
TTTTTACCCATTCTGAAGGACAACCGGGAAAAGATCGAGAAGATCCTGACCTTCCG
CATCCCCTACTACGTGGGCCCCTCTGGCCAGGGGAAAACAGCAGATTTCGCTGGATGA
CCAGAAAGAGCGAGGAAACCATCACCCCTGGAACCTTCGAGGAAGTGGTGGACAA
GGGCGCTTCCGCCCAGAGCTTCATCGAGCGGATGACCAACTTCGATAAGAACCTGC
CCAACGAGAAGGTGCTGCCCAAGCACAGCCTGCTGTACGAGTACTTCACCGTGTAT
AACGAGCTGACCAAAGTGAAATACGTGACCGAGGGAATGAGAAAGCCCCGCCTTCCT
GAGCGGCGAGCAGAAAAAGGCCATCGTGGACCTGCTGTTCAAGACCAACCGGAAA
GTGACCGTGAAGCAGCTGAAAGAGGACTACTTCAAGAAAATCGAGTGCTTCGACTC
CGTGGAATCTCCGGCGTGGAAGATCGGTTCAACGCCTCCCTGGGCACATACCACG
ATCTGCTGAAAATTATCAAGGACAAGGACTTCCTGGACAATGAGGAAAACGAGGAC
ATTCTGGAAGATATCGTGCTGACCCTGACACTGTTTGAGGACAGAGAGATGATCGA
GGAACGGCTGAAAACCTATGCCACCTGTTTCGACGACAAAGTGATGAAGCAGCTGA
AGCGGCGGAGATACACCGGCTGGGGCAGGCTGAGCCGGAAGCTGATCAACGGCATC
CGGGACAAGCAGTCCGGCAAGACAATCCTGGATTTCTGAAGTCCGACGGCTTCGC
CAACAGAACTTCATGCAGCTGATCCACGACGACAGCCTGACCTTTAAAGAGGACA
TCCAGAAAGCCCAGGTGTCCGGCCAGGGCGATAGCCTGCACGAGCACATTGCCAAT
CTGGCCGGCAGCCCCGCCATTAAGAAGGGCATCCTGCAGACAGTGAAGGTGGTGA
CGAGCTCGTGAAAGTGATGGGCCGGCACAAGCCCAGAACATCGTGATCGAAATGG
CCAGAGAGAACCAGACCACCCAGAAGGGACAGAAGAACAGCCGCGAGAGAATGAA
GCGGATCGAAGAGGGCATCAAAGAGCTGGGCAGCCAGATCCTGAAAGAACACCCC
GTGGAAAACACCCAGCTGCAGAACGAGAAGCTGTACCTGTACTACCTGCAGAATGG
GCGGGATATGTACGTGGACCAGGAACTGGACATCAACCGGCTGTCCGACTACGAT

FIG. 50B

GTGGACCATATCGTGCCTCAGAGCTTTCTGAAGGACGACTCCATCGACAACAAGGTG
CTGACCAGAAGCGACAAGAACCAGGGGCAAGAGCGACAACGTGCCCTCCGAAGAGG
TCGTGAAGAAGATGAAGAATACTGGCGGCAGCTGCTGAACGCCAAGCTGATTACC
CAGAGAAAGTTCGACAATCTGACCAAGGCCGAGAGAGGGCGCCTGAGCGAACTGG
ATAAGGCCGGCTTCATCAAGAGACAGCTGGTGGAAACCCGGCAGATCACAAAGCAC
GTGGCACAGATCCTGGACTCCCGGATGAACACTAAGTACGACGAGAATGACAAGCT
GATCCGGGAAGTGAAAGTGATCACCTGAAGTCCAAGCTGGTGTCCGATTTCCGGA
AGGATTTCCAGTTTACAAAGTGCGCGAGATCAACAATACTACCACCACGCCACGAC
GCCTACCTGAACGCCGTCGTGGGAACCCGCCCTGATCAAAAAGTACCCTAAGCTGGA
AAGCGAGTTCGTGTACGGCGACTACAAGGTGTACGACGTGCGGAAGATGATCGCCA
AGAGCGAGCAGGAAATCGGCAAGGCTACCGCCAAGTACTTCTTCTACAGCAACATC
ATGAACTTTTTCAAGACCGAGATTACCCTGGCCAACGGCGAGATCCGGAAGCGGCC
TCTGATCGAGACAAACGGCGAAACCCGGGAGATCGTGTGGGATAAGGGCCGGGATT
TTGCCACCGTGCAGGAAAGTGCTGAGCATGCCCAAGTGAATATCGTGAAAAAGACC
GAGGTGCAGACAGGCGGCTTCAGCAAAGAGTCTATCCTGCCCAAGAGGAACAGCGA
TAAGCTGATCGCCAGAAAGAAGGACTGGGACCCTAAGAAGTACGGCGGCTTCGACA
GCCCCACCGTGGCCTATTCTGTGCTGGTGGTGGCCAAAGTGGAAAAGGGCAAGTCC
AAGAACTGAAGAGTGTGAAAGAGCTGCTGGGGATCACCATCATGGAAAGAAGCA
GCTTCGAGAAGAATCCCATCGACTTTCTGGAAGCCAAGGGCTACAAAGAAGTGAAA
AAGGACCTGATCATCAAGCTGCCTAAGTACTCCCTGTTTCGAGCTGGAAAACGGCCG
GAAGAGAATGCTGGCCTCTGCCGGCGAAGTGCAGAAGGGAAACGAAGTGGCCCTGC
CCTCCAAATATGTGAACTTCTGTACCTGGCCAGCCACTATGAGAAGCTGAAGGGCT
CCCCGAGGATAATGAGCAGAAACAGCTGTTTGTGGAACAGCACAAAGCACTACCTG
GACGAGATCATCGAGCAGATCAGCGAGTTCTCCAAGAGAGTGATCCTGGCCGACGC
TAATCTGGACAAAGTGCTGTCCGCCTACAACAAGCACCCGGGATAAGCCCATCAGAG
AGCAGGCCGAGAAATATCATCCACCTGTTTACCCTGACCAATCTGGGAGCCCCTGCCG
CCTTCAAGTACTTTGACACCACCATCGACCGGAAGAGGTACACCAGCACCAAGAG
GTGCTGGACGCCACCCTGATCCACCAGAGCATCACCGGCCTGTACGAGACACGGAT
CGACCTGTCTCAGCTGGGAGGCGACAAAAGGCCGGCGGCCACGAAAAAGGCCGGC
CAGGCAAAAAAGAAAAAG

P2A-EGFP

ggaagcggagccaacttctccctgttgaacaagcaggggatgtcgaagagaatccgggccaGTGAGCAAGGGCGA
GGAGCTGTTACCGGGGTGGTGGCCATCCTGGTTCGAGCTGGACGGCGACGTAAACG
GCCACAAGTTCAGCGTGTCCGGCGAGGGCGAGGGCGATGCCACCTACGGCAAGCTG
ACCCTGAAGTTCATCT

FIG. 50C

GCACCACCGGCAAGCTGCCCGTGCCCTGGCCCACCCCTCGTGACCACCCTGACCTACG
CGGTGCAGTGCTTCAGCCGCTACCCCGACCACATGAAGCAGCACGACTTCTTCAAGT
CCGCCATGCCCGAAGGCTACGTCCAGGAGCGCACCATCTTCTTCAAGGACGACGGC
AACTACAAGACCCGCGCCGAGGTGAAGTTGAGGGCGACACCCTGGTGAACCGCAT
CGAGCTGAAGGGCATCGACTTCAAGGAGGACGGCAACATCCTGGGGCACAAGCTGG
AGTACAAC TACAACAGCCACAACGTCTATATCATGGCCGACAAGCAGAAGAACGGC
ATCAAGGTGAACTTCAAGATCCGCCACAACATCGAGGACGGCAGCGTGCAGCTCGC
CGACCACTACCAGCAGAACACCCCCATCGGCGACGGCCCCGTGCTGCTGCCCGACA
ACCACTACCTGAGCACCCAGTCCGCCCTGAGCAAAGACCCCAACGAGAAGCGCGAT
CACATGGTCTGCTGGAGTTCGTGACCGCCGCCGGGATCACTCTCGGCATGGACGAG
CTGTACAAG

WPRE

Cgataatcaacctctggattacaaaattgtgaaagattgactggattcctaactatgttgccttttaacgctatgtggatacgtgctttaatgc
ctttgatcatgctattgcttcccgatggtttcatttctcctctgtataaatctggttgcctccttatgaggagtgtggcccgtgtcagg
caacgtggcgtgggtgcaactgtgtttctgacgcaacccccactgggtggggcattgcccaccacgtcagctccttccgggactttcgtt
tccccctccctattgcccaggggaactcatgcccctgcttcccgctgctggacaggggctcgggtgttgggcaactgacaattccgtg
gtgtgtcggggaaatcatgctccttccctgtgctgctgcctgtgttcccactgattctgcccgggacgctcctctgctacgtcccttggc
cctcaatccagcggacccttcccgcggtgctgctgcccgtctgcccctctccgctcttccgctccttccgctccttccgctcagacgagtgatc
cctttggcgcctcccccacatg

bGHpolyA

cgaactCGA Ctggtcctctagtgccagccatctgtgtttgccctccccgtgccttcttgaccctggaagggtgccactcccactgt
ccttccataataaatgaggaaattgcatcgcattgtctgagtaggttcattctattctgggggggtgggtggggcaggacagcaagggg
aggattgggaagacaatGgcagggcatg

loxP-SV40polyAx3-loxP

ataacttcgtataatgtatgctatcgaagtatttcgcatgaataaataaagccttcagatctgogactctagaggatctgcgactctagagg
atcataatcagccntaccacattttgtagaggtttactngctttaaaaaacctccacacctccccctgaacctgaaacataaaaatgaatgcaa
ttgtgtgttaactgtttattgcaactfataatggttacaaaataaagcaatgacatcacaatttcacaaataaagcatttttcaactgcattctagt
tgtggtttgccaactcatcaatgtatcttatcatgtctggatctgcgactctagaggatcataatcagccataccacattttgtagaggtttactt
gctttaaaaaacctccacacctccccctgaacctgaaacataaaaatgaatgcaattgtgtgttaactgtttattgcaactfataatggttaca
aataaagcaatagcacaacaaatttcacaaataaagcatttttcaactgcattctagtgtgtttgccaactcatcaatgtatcttatcatgct
ggatctgcgactctagaggatcataatcagccataccacattttgtagaggtttacttgcctttaaaaaacctccacacctccccctgaacctga
aacataaaaatgaatgcaattgtgtgttaactgtttattgcaactfataatggttacaata

FIG. 50D

aagcaatagcaccacaatttcacaataaagcatitititcaactgcattctagttggtttgccaactcacaatgtatcttatcatgtctggat
ccccatcaagctgafccggaaaccttaataacttcgtataatgtatgctatacgaagtat

Rosa26 short homology arm

caggccctccgagcgtggtggagccgittotgtgagacagccgggtacgagtcgtgacgctggaaggggcaagcgggtggtgggcagg
aatgagggtccgacctgacgcaaccggagggggagggagaagggagcggaaaagtcaccaccggagcggccatggtcggggggg
ggggggcagcggaggaGcgcttccggccgacgctcgtcgtgattggcttCttttctcccggctgtgtgaaaacacaaatggcgtgt
ttggttggcgtaaaggcgcctgtcagtaacggcagccggagtgccagccggcagccctgctctgcccactgggtggggggggag
gtaggtgggggtgagggcagctgGacgtgcgggcggcggcctctggcggggcgggggaggggagggagggcagcgaagga
gtcgcgcgcgagcggccgccaccctcccctccctcctgggggagctgittaccggccggcggcggcctcgtcgtctatggctc
ggggcccagaaaactggcccttgccttggctcgtgtcgtgcaagttgagtcaccctccggccagcggggggcggcagggagggcctc
ccaggttccggccctcccctggccccggccggcagagctggccggcggccctggcgaacgtggcaggaagcgcgctggggggc
ggggacgggcagtaggggtgagcggctggcgggggggggcaagcagcttccgactigagttgctcaagagggggcgtgctgagccag
acctccatcgccactccggggagtgagggggaaggagcggagggctcagttgggctgtttggagggcaggaagcacttgcctcccaagt
cgtctgagttgtatcagtaaggagctgcagtgaggaggggggagaaagggccacccttctccggaggggggagggggaggtttgc
aatacctttctgggagttcctctgctgctcctggcttctgaggacggcctgggctgggagaaatcccttccctctccctctgctgca
ctccagctcttctag

Rosa26 long homology arm

agatgggggggagctctcggggcaggctfaaaggctaacctgggtgtgtgggcgttgcctgcaggggaaatgaacaggtgtaaaatggag
ggacaagacttcccacagatttccgggtttgtcgggaagtittitaatagggcaataaaggaaaatgggagggataggtatcctgagggttt
atgcagcaaaactacagggtatttctgtgafccgccctggagfaffttccatcgaggtagatfaaagacatgctcaccggagtttatactct
cctgcttGAGATCCTTACTACAGTATGAAATTACAGTGTCTCGAGTTAGACTATGTAAGC
AGAATTTTAAATCA Ttttaagagccagtactcatalccatttctcccgtctctcgaagccttatcaaaaggta Ttttagaaca
ctcaatitagccccattttcatttatactggcttatccaacccttagacagagcattggcattttcccttctgacttagaagctgtagactca
tgaaccagacagattagttatatacaccacaatcagagcgttagctggggcctcaaacactgcagttctttataactccttagtacacttttg
ttgatcctttgccttgatccctaaatctcagtgctatcacctctcccgtcagggtggtgttccacatttgggccatttctcagctccaggagttitaca
acaatagatgtatgagaatccaaactaaagcttaactttccactcccataatgcctctctccttttctccattTATAAACTGAGCT
ATTAACCATTAATGGTTTCCAGGTGGATGTCTCCTCCCCCAATATTACCTGATGTATC
TTACATATTGCCAGGCTGATATTTTAAGACATTA AAAAGGTATATTTTATTATTGAGCC
ACATGGTATTGATTACTGCTTactaaaattttgcattgtacacatctgtaaaagggtggttcccttttggaaatgcaaaagttaggt
gtttgtgtctttctgacctaaaggtctgtgagctgtattttctatttaagcagtgctttctctggactggcctgactcattggcattctacaggtta
ttgctggctaaatgtgattttgccaaagctctcaggacctataatttgtcgtgactgttagccaacacaagtaaaatgaitaagcaacaaatgt
atttgtgaagcttggtttttaggttgtgtgtgtgtgctgtgctctataataactatccaggggctggagaggtggctcggagttcaagag
cacagactgctctccagaagtc

FIG. 50E

ctgagttcaattccagcaaccacatgggtggctcacaaccatctgtaatgggatctgatgccctctctgggtgtgtgaagaccacaagtgtat
tcacafaaataaataaaTCCTCCTTCTTCTTCTTTTTTTTTTTTTTTTAAAGAGAATACTGTCTCCAG
TAGAAiTTACTGAAGTAATGAAATACTTTGTGTTTGTTCGAATATGGTAGCCAATAAT
CAAATiACTCTTTaAGCACTGGAAATGTtACCAAGGAACTAaTTTTtATTTgAAGTGTaA
CTGTGGACAGAGGAGCCATAACTGCAGACTTGTGGGATACAGAAGACCAATGCAGAC
CITTAATGTCTTTTTCTCTTACACTAAGCAATAAAGAAAATAAAAATTGAACTTCTAGTA
TCCTATTTGTTiAAACTGCTAGCTTTACiTAACTTTTTGTGCTTCATCTATACAAAGCTG
AAAGCTAAGTCTGCAGCCATTACTAAACATGAAAGCAAGTAATGATAATTTTGGATT
TCAAAAATGTAGGGCCAGAGTTTAGCCAGCCAGTGGTGGTGGCTTGCCTTTATGCCiTT
AATCCCAGCACTCTGGAGGCAGAGACAGGCAGATCTCTGAGTTTGGAGCCAGCCTG
GICTACACATCAAGTTCTATCTAGGATAGCCAGGAATACACACAGAAACCCTGTTGG
GGAGGGGGGCTCTGAGATTTCAATAAAATTATAATTGAAGCATTCCCTAATGAGCCAC
TATGGATGTGGCTAAATCCGTCTACCTTTCTGATGAGATTTGGGTATTATTTTTTCTG
TCTCTGCTGTIGGTTGGGTCITTTGACACTGTGGGCTTTCTTtAAAGCCTCCTTCTGC
CATGTGGTCTCTTGTTTGCTACTAACTTCCCATGGCTTAAATGGCATGGCTTTTTGCC
TTCTAAGGGCAGCTGCTGAGATTTGCAGCCTGATTTCCAGGGTGGGGTTGGGAAATC
TTTCAAACACTAAAATTGTCCITTAATTTTTTTTTTAAAAAATGGGTTATATAATAAA
CCTCATAAAAATAGTTATGAGGAGTGAGGTGGACTAATATTAAaTGAGTCCCTCCCCT
ATAAAAGAGCTATTAAGGCTTTTTGTCTTATACTiAACTTTTTTTTTTAAATGTGGTATC
TTTAGAACCAAGGGTCTTAGAGTTTTAGTATACAGAAACTGTTGCATCGCTTAATCA
GATTTTCTAGTTTCAAATCCAGAGAAATCCAAATTCCTTACAGCCAAAGTCAAATTA
GAATTTCTGACTTTiAATGTTAaTTTGCtTACTGTGAATATaAAAATGATAGCTTTTCCT
GAGGCAGGGTCTCACTATGTATCTCTGCCTGATCTGCAACAAGATATGTAGACTAAA
GTTCTGCCTGCTTTTTGTCTCCTGAATACTAAGGTTAAAATGTAGTAATACTTTTTGGAA
CTTGCAGGTCAGATTCTTTTATAGGGGACACACTAAGGGAGCTTGGGTGATAGTTGG
TAAAtgtgttaagtgatgaaaactgaattattatcacgcaacctacttttaaaaaaaaaagccaggcctgttagagcatgctTaaggg
atccctaggacttgcgagcacacaAGAGTAGiTACTTGGCAGGCTCCTGGTGAGAGCATAATTTCAA
AAAACAAGGCAGACAACCAAGAACTACAGTiAAGGTTACCTGTCTTTaAACCATCT
GCATATACACAGGGATATTAATAATTTCCAAATAATATTTTCAATCAAGTTTTCCCCC
ATCAAATTTGGGACATGGATTTCTCCGGTGAATAGGCAGAGTTGGAAACTAAACAAA
TGTTGGTTTTGTGATTTGTGAAATTGTTTTCAAGTGATAGTTAAAGCCCATGAGATAC
AGAACAAAGCTGCTATTTTCGAGGTCTCTTGGTTiATACTCAGAAGCACTTCTTTGGGT
TTCCCTGCACTATCCTGATCATGTGCTAGGCCTACCTTAGGCTGATTGTTGTTCAAAT
aAACTTAAGTTTCTGTGAGGTGATGTCATATGATTTTCATATATCAAGGCCAAAACATG
TTATATATGTTAAACATTTGTACTTAATGTGAAAGTTAGGTCTTTGTGGGTT

FIG. 50F

TGATTTTtAAAtTTTCAAAAACCTGAGCTAAATAAGTCATTTTtACATGTCTTACATTTGGT
GgAATTGTATaATTGTGGTTTTGCAGGCAAGACTCTCTGACCTAGTAACCCTaCCTATA
GAGCACTTTGCTGGGTCACAAGTCTAGGAGTCAAGCATTTCACCTTGAAGTTGAGAC
GTTTTGTTAGTGTATACTAGTTtATATGTTGGAGGACATGTTTATCCAGAAGATATTC
AGGACTATTTTTGACTGGGCTAAGGAATTGATTCTGATTAGCACTGTTAGTGAGCAT
TGAGTGGCCTTTAGGCTTGAATTggagtcactgtatatctcaaataatgctggcctttttaaagccctgtctttatca
ccctgtttctacataaattttgtcaaagaaactgtttggaTCTCCTTTTGACAACAATAGCATGTTTTCAAG
CCATATTTTTTTICCTTTTTTTTTTTTTTTGGTTTTTCGAGACAGGGTTTCTCTGTAT
AGCCCTGGCTGTCTGGAACCTCACTTTGTAGACCAGGCTGGCCTCGAACTCAGAAAT
CCGCTGCCCTCTGCCCTCTGAGTGCCGGGATTAAGGGCGTGCACCACCACGCCTGGC
TAAGTTGGATATTTTGTtATATAACTATAACCAATACTAACTCCACTGGGTGGATTTT
TAATTCAGTCAGTAGTCTTAAGTGGTCTTTATTGGCCCTTcATTAAAATCTACTGTTT
ACTCTAACAGAGGCTGTTGGtACTAGTGGCACHtAAGCAACTTCCTACGGATATACTA
GCAGAtAAGGGTCAGGGATAGAACTAGTCTAGCGTTTTGTATACCTACCAGCTtA
TACTACCTTGTCTGATAGAAATATTTcAGGACATCTAGCTT

pPGK-Neo-pPGK-polyA

aatctaccgggtaggggagggcgcttttccaaggcagctctggagcatgoccttttagcagcccccgtggcacttggogctacacaagtgg
cctctggcctcgcacacattccacatccaccggtagggcgccaaccggctccgttttggggcccccfcggccaccttctactcctcccc
agtcaggaagttccccccgccccgagctcgcctcgtcaggacgtgacaaatggaagtagcacgtctcactagctcgtgcagatgga
cagcaccgctgagcaatggaagcgggtagggcctttggggcagcggccaatagcagctttgctcctcgtttctgggctcagaggctggg
aaggggtgggtccgggggcccggctcaggggcccggctcaggggcccgggcccggcggccgaaggtcctcgggagggcccggcattctgc
acgctcaaaaagcgcagctcgcggcgtgctcctctcctcctcctcctcgggccccttcgacctgcaatcggcgtagcgaagtctctattct
agaaagtataggaacttcgccaccatgggatcggccaitgaacaagatggatgacgcaggtctccggccttgggtggagaggctat
tcggctatgactgggcacaacagacaatcggctgctctgatccgcctgttccggctgacgcagggggcggccgggtcttttgtcaag
accgacctgctcgggtgccctgaatgaactgcaggacgaggcagcggcctatcgtggctggccacgacggggcgttcttgcgcagctgt
gctcgcaggtgtcactgaagcgggaaggactggctgctatggggcgaagtgcggggcaggatcctcgtcctcactcctgctcctgccc
gagaaagtatccatcagctgatgcaatgcggcggctgcatcactgactcggctacctgcccaitegccaccaagcgaacatcgc
tcgagcagcacgtactcggatggaagccggtcttctcgaatcaggatgafctggacgaagagcatcaggggctcggccagccgaactg
ttcggcaggtcaaggcgcgatgcccagcggcggatgatcctcgtgacccatggcgatgctgcttggcgaatcctatgtggaaaatg
ggcgttttctggatcactgactgtggccggctgggtgtggcggaccctatcaggacatagcgttggctaccctgatattgctgaagag
cttggcggcgaatgggctgaccgctctcgtgcttttaoggtatcgcgctcccgatcgcagcgcacgcctctctcctctgacgagt
tcttctgaggggatccgctgtaagctgcagaaatgatgatctatfaaacaataaagatgtccactaaaatggaagtittcctgctacacttgtt
aagaagggtgagaacagagtagctacatttgaatggaaggattggagctacgggggtgggggtgggggtgggattagataaatgcctgct
cttactgaaggctcttactattgcittatgataatgtttcatagtg

FIG. 50G

gatataaaittaaacaaagcaaaaccaaaitaagggccagctcattectcccactcatgatctatagatctatagatctctcgtgggatcattgt
ttttcttctgattccactttgtgggttaagtaactgtggtttccaaatgtgtcagtttcatagcctgaagaacgagatcagcagcctctgtccaca
tacacttcattctcagtaattgtttgccaagttcfaatfccatcagaaagc

pPGK-DTA

TACCGGGTAGGGGAGGCGCTTTTCCcAAGGCAGTCTGgAGCATGCGCtTTAGCAGCCC
CGCTgGGCACTTGGCGCTACACAAGTGGCCTCTGGCCTCGCACACATTCCACATCCA
CCGGTAGGGCCAAACCGGCTCCGTTCTTTGGTGGCCCCCTTCGCGCCACCTTCTACTCC
TCCCCTAGTCAGGAAGTTCCCCCCCCGCCCGCAGCTCGCGTCGTGcAGGACGTGACA
AATGGAAGTAGCACGTCTCACTAGTCTCGTgCAGATGGACAGCACCGCTGAGCAATG
GAAGCGGGTAGGCCCTTTGGGGCAGCGGCCAATAGCAGCTTTGCTCCTTCGCTTTCTG
GGCTCAGAGGCTGGGAAGGGGTGGGTCCGGGGGGCGGGCTCAGGGGGCGGGCTCAGG
GGCGGGGCGGGCGCCCGAAGGTCCCTCCGGAGGCCCGGCATTCTGCACGCTTCAAAA
GCGCACGTCTGCCGCGCTGTTCTCCTCTTCCTCATCTCCGGGCCTTTCGACCTGCAGG
TCCTCGCCATggatcctgatgatgttgttGattctctaaAtctttgtGatggaaaacttttcttctgaccacgggactaaacctggtt
atgtagattccattcaaaaaggatatacaaaaagcacaatctggfacacaaggaaatfatgacgatgattggaagggtttatagfaccgaca
taaatacagcctcgggatactctgtagataatgaaaaccgctctctggaaaagctggaggcgtggtcaaatgacgfatccaggactg
acgaagggtctcgcactaaaagtggaataatccgaaactaataagaaagagttaggttaagtctactgaaccgtgatggagcaagtcgga
acggaagagttaataaaaaggtcgggtgatggtgcttcgctgtagtgctcagccttcccttcgctgaggggagttctagcgttgaatalata
taactgggaacaggcgaagcgttaagcgtagaacttgagattaatttgaaacccgtggaaaacgtggccaagatgcatgatgatgagtata
tggctcaagcctgtgcaggaaatcgtgcaggcgatctctttgtgaaggaacccttctgtggtgtgacataattggacaactacctacag
agattaaagctcgaagtaataataaaattttaagtgataatgtgttaactactgattcgaattgtgtattitagattccaacctatggaact
gatgaatgggagcagtggtggaatgcagatccagagctcgtgatcagcctcactgtgcttctagttgccagccatctgtgtttgccct
ccccgtgcttctgacctggaagggtgcactcccactgtccttcttaataaaatgaggaaattgcatcgcattgtctgagtaggtgtcat
tctattctggggggtggggggggcaggacagcaagggggaggagtgggaagacaatagcagggcatg

FIG. 50H




FIG. 51




FIG. 52

Validation of Cas9 nuclease activity by Surveyor




FIG. 53

		Average	StDev
pCAG-loxp(pA)loxp-NLS-hSpCas9-NLS-2A-GFP	Clone 1	32.1	7.1
	Clone 2	27.3	3.5
	Clone 3	35.9	1.4
	Clone 4	39.0	4.7
pCAG-NLS-hSpCas9-NLS-2A-GFP	Clone 1	26.9	1.3
	Clone 2	33.1	2.7

FIG. 54




FIG. 55




FIG. 56




FIG. 57




FIG. 59




FIG. 60

FIG. 62A




FIG. 62B




FIG. 62C




FIG. 62D

sgRNAs	Approx. deletion size (bp)	Number of clones				Total
		+/+	Δ/+	Δ/Δ	Inversion	
3.1 + 4.1	282	12	10	1	0	23
3.2 + 4.1	237	20	16	2	0	38
3.1 + 4.2	425	4	14	5	0	23
3.2 + 4.2	259	5	11	4	0	21

FIG. 63A




FIG. 63B





FIG. 63C







FIG. 65



gRNA sequences for Chd8 targeting:

- Chd8.1 - agctgtttactggtcggct
- Chd8.2 - aatggatacacctggtcgaa
- Chd8.3 - caatggatacacctggtcga

FIG. 66




FIG. NLS1

FIG. 67




FIG. 68





FIG. 69B



NLS architecture optimization for SpCas9





FIG. 70



QQ plot for the MCGNN sequences

FIG. 71




FIG. 72

FIG. 73A




FIG. 73C




FIG. 73B



FIG. 74A




FIG. 74B




FIG. 74C





FIG. 74D




FIG. 74E

**CRISPR-CAS COMPONENT SYSTEMS,
METHODS AND COMPOSITIONS FOR
SEQUENCE MANIPULATION**

**RELATED APPLICATIONS AND
INCORPORATION BY REFERENCE**

[0001] This application is a continuation of U.S. application Ser. No. 15/230,161 filed Aug. 5, 2016, which is a continuation of U.S. application Ser. No. 14/105,035 filed Dec. 12, 2013, which claims priority to U.S. provisional patent applications 61/736,527, 61/748,427, 61/768,959, 61/791,409 and 61/835,931, all entitled SYSTEMS METHODS AND COMPOSITIONS FOR SEQUENCE MANIPULATION filed on Dec. 12, 2012, Jan. 2, 2013, Feb. 25, 2013, Mar. 15, 2013 and Jun. 17, 2013, respectively.

[0002] Reference is made to U.S. provisional patent applications 61/758,468; 61/769,046; 61/802,174; 61/806,375; 61/814,263; 61/819,803 and 61/828,130, each entitled ENGINEERING AND OPTIMIZATION OF SYSTEMS, METHODS AND COMPOSITIONS FOR SEQUENCE MANIPULATION, filed on Jan. 30, 2013; Feb. 25, 2013; Mar. 15, 2013; Mar. 28, 2013; Apr. 20, 2013; May 6, 2013 and May 28, 2013 respectively. Reference is also made to U.S. provisional patent applications 61/835,936, 61/836,127, 61/836,101, 61/836,080, 61/836,123 and 61/835,973 each filed Jun. 17, 2013. Reference is also made to U.S. provisional patent application 61/842,322 and U.S. patent application Ser. No. 14/054,414, each having Broad reference BI-2011/008A, entitled CRISPR-CAS SYSTEMS AND METHODS FOR ALTERING EXPRESSION OF GENE PRODUCTS filed on Jul. 2, 2013 and Oct. 15, 2013 respectively.

[0003] The foregoing applications, and all documents cited therein or during their prosecution (“appln cited documents”) and all documents cited or referenced in the appln cited documents, and all documents cited or referenced herein (“herein cited documents”), and all documents cited or referenced in herein cited documents, together with any manufacturer’s instructions, descriptions, product specifications, and product sheets for any products mentioned herein or in any document incorporated by reference herein, are hereby incorporated herein by reference, and may be employed in the practice of the invention. More specifically, all referenced documents are incorporated by reference to the same extent as if each individual document was specifically and individually indicated to be incorporated by reference.

**STATEMENT AS TO FEDERALLY SPONSORED
RESEARCH**

[0004] This invention was made with government support under the NIH Pioneer Award DP1MH100706, awarded by the National Institutes of Health. The government has certain rights in the invention.

SEQUENCE LISTING

[0005] The instant application contains a Sequence Listing which has been submitted electronically in ASCII format and is hereby incorporated by reference in its entirety. Said ASCII copy, created on Mar. 31, 2015, is named 44790.07.2003_SL.txt and is 308.802 bytes in size.

FIELD OF THE INVENTION

[0006] The present invention generally relates to systems, methods and compositions used for the control of gene expression involving sequence targeting, such as genome perturbation or gene-editing, that may use vector systems related to Clustered Regularly Interspaced Short Palindromic Repeats (CRISPR) and components thereof.

BACKGROUND OF THE INVENTION

[0007] Recent advances in genome sequencing techniques and analysis methods have significantly accelerated the ability to catalog and map genetic factors associated with a diverse range of biological functions and diseases. Precise genome targeting technologies are needed to enable systematic reverse engineering of causal genetic variations by allowing selective perturbation of individual genetic elements, as well as to advance synthetic biology, biotechnological, and medical applications. Although genome-editing techniques such as designer zinc fingers, transcription activator-like effectors (TALEs), or homing meganucleases are available for producing targeted genome perturbations, there remains a need for new genome engineering technologies that are affordable, easy to set up, scalable, and amenable to targeting multiple positions within the eukaryotic genome.

SUMMARY OF THE INVENTION

[0008] There exists a pressing need for alternative and robust systems and techniques for sequence targeting with a wide array of applications. This invention addresses this need and provides related advantages. The CRISPR/Cas or the CRISPR-Cas system (both terms are used interchangeably throughout this application) does not require the generation of customized proteins to target specific sequences but rather a single Cas enzyme can be programmed by a short RNA molecule to recognize a specific DNA target, in other words the Cas enzyme can be recruited to a specific DNA target using said short RNA molecule. Adding the CRISPR-Cas system to the repertoire of genome sequencing techniques and analysis methods may significantly simplify the methodology and accelerate the ability to catalog and map genetic factors associated with a diverse range of biological functions and diseases. To utilize the CRISPR-Cas system effectively for genome editing without deleterious effects, it is critical to understand aspects of engineering and optimization of these genome engineering tools, which are aspects of the claimed invention.

[0009] In one aspect, the invention provides a vector system comprising one or more vectors. In some embodiments, the system comprises: (a) a first regulatory element operably linked to a tracr mate sequence and one or more insertion sites for inserting one or more guide sequences upstream of the tracr mate sequence, wherein when expressed, the guide sequence directs sequence-specific binding of a CRISPR complex to a target sequence in a eukaryotic cell, wherein the CRISPR complex comprises a CRISPR enzyme complexed with (1) the guide sequence that is hybridized to the target sequence, and (2) the tracr mate sequence that is hybridized to the tracr sequence; and (b) a second regulatory element operably linked to an enzyme-coding sequence encoding said CRISPR enzyme comprising a nuclear localization sequence; wherein components (a) and (b) are located on the same or different vectors of the system. In some embodiments, component (a)

further comprises the tracr sequence downstream of the tracr mate sequence under the control of the first regulatory element. In some embodiments, component (a) further comprises two or more guide sequences operably linked to the first regulatory element, wherein when expressed, each of the two or more guide sequences direct sequence specific binding of a CRISPR complex to a different target sequence in a eukaryotic cell. In some embodiments, the system comprises the tracr sequence under the control of a third regulatory element, such as a polymerase III promoter. In some embodiments, the tracr sequence exhibits at least 50%, 60%, 70%, 80%, 90%, 95%, or 99% of sequence complementarity along the length of the tracr mate sequence when optimally aligned. Determining optimal alignment is within the purview of one of skill in the art. For example, there are publically and commercially available alignment algorithms and programs such as, but not limited to, ClustalW, Smith-Waterman in matlab, Bowtie, Geneious, Biopython and SeqMan. In some embodiments, the CRISPR complex comprises one or more nuclear localization sequences of sufficient strength to drive accumulation of said CRISPR complex in a detectable amount in the nucleus of a eukaryotic cell. Without wishing to be bound by theory, it is believed that a nuclear localization sequence is not necessary for CRISPR complex activity in eukaryotes, but that including such sequences enhances activity of the system, especially as to targeting nucleic acid molecules in the nucleus. In some embodiments, the CRISPR enzyme is a type II CRISPR system enzyme. In some embodiments, the CRISPR enzyme is a Cas9 enzyme. In some embodiments, the Cas9 enzyme is *S. pneumoniae*, *S. pyogenes*, or *S. thermophilus* Cas9, and may include mutated Cas9 derived from these organisms. The enzyme may be a Cas9 homolog or ortholog. In some embodiments, the CRISPR enzyme is codon-optimized for expression in a eukaryotic cell. In some embodiments, the CRISPR enzyme directs cleavage of one or two strands at the location of the target sequence. In some embodiments, the CRISPR enzyme lacks DNA strand cleavage activity. In some embodiments, the first regulatory element is a polymerase III promoter. In some embodiments, the second regulatory element is a polymerase II promoter. In some embodiments, the guide sequence is at least 15, 16, 17, 18, 19, 20, 25 nucleotides, or between 10-30, or between 15-25, or between 15-20 nucleotides in length. In general, and throughout this specification, the term “vector” refers to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. Vectors include, but are not limited to, nucleic acid molecules that are single-stranded, double-stranded, or partially double-stranded; nucleic acid molecules that comprise one or more free ends, no free ends (e.g. circular); nucleic acid molecules that comprise DNA, RNA, or both; and other varieties of polynucleotides known in the art. One type of vector is a “plasmid,” which refers to a circular double stranded DNA loop into which additional DNA segments can be inserted, such as by standard molecular cloning techniques. Another type of vector is a viral vector, wherein virally-derived DNA or RNA sequences are present in the vector for packaging into a virus (e.g. retroviruses, replication defective retroviruses, adenoviruses, replication defective adenoviruses, and adeno-associated viruses). Viral vectors also include polynucleotides carried by a virus for transfection into a host cell. Certain vectors are capable of autonomous replication in a host cell into which they are introduced (e.g. bacterial

vectors having a bacterial origin of replication and episomal mammalian vectors). Other vectors (e.g., non-episomal mammalian vectors) are integrated into the genome of a host cell upon introduction into the host cell, and thereby are replicated along with the host genome. Moreover, certain vectors are capable of directing the expression of genes to which they are operatively-linked. Such vectors are referred to herein as “expression vectors.” Common expression vectors of utility in recombinant DNA techniques are often in the form of plasmids.

[0010] Recombinant expression vectors can comprise a nucleic acid of the invention in a form suitable for expression of the nucleic acid in a host cell, which means that the recombinant expression vectors include one or more regulatory elements, which may be selected on the basis of the host cells to be used for expression, that is operatively-linked to the nucleic acid sequence to be expressed. Within a recombinant expression vector, “operably linked” is intended to mean that the nucleotide sequence of interest is linked to the regulatory element(s) in a manner that allows for expression of the nucleotide sequence (e.g. in an in vitro transcription/translation system or in a host cell when the vector is introduced into the host cell).

[0011] The term “regulatory element” is intended to include promoters, enhancers, internal ribosomal entry sites (IRES), and other expression control elements (e.g. transcription termination signals, such as polyadenylation signals and poly-U sequences). Such regulatory elements are described, for example, in Goeddel, GENE EXPRESSION TECHNOLOGY: METHODS IN ENZYMOLOGY 185, Academic Press, San Diego, Calif. (1990). Regulatory elements include those that direct constitutive expression of a nucleotide sequence in many types of host cell and those that direct expression of the nucleotide sequence only in certain host cells (e.g., tissue-specific regulatory sequences). A tissue-specific promoter may direct expression primarily in a desired tissue of interest, such as muscle, neuron, bone, skin, blood, specific organs (e.g. liver, pancreas), or particular cell types (e.g. lymphocytes). Regulatory elements may also direct expression in a temporal-dependent manner, such as in a cell-cycle dependent or developmental stage-dependent manner, which may or may not also be tissue or cell-type specific. In some embodiments, a vector comprises one or more pol III promoter (e.g. 1, 2, 3, 4, 5, or more pol III promoters), one or more pol II promoters (e.g. 1, 2, 3, 4, 5, or more pol II promoters), one or more pol I promoters (e.g. 1, 2, 3, 4, 5, or more pol I promoters), or combinations thereof. Examples of pol III promoters include, but are not limited to, U6 and H1 promoters. Examples of pol II promoters include, but are not limited to, the retroviral Rous sarcoma virus (RSV) LTR promoter (optionally with the RSV enhancer), the cytomegalovirus (CMV) promoter (optionally with the CMV enhancer) [see, e.g., Boshart et al, Cell, 41:521-530 (1985)], the SV40 promoter, the dihydrofolate reductase promoter, the β -actin promoter, the phosphoglycerol kinase (PGK) promoter, and the EF1 α promoter. Also encompassed by the term “regulatory element” are enhancer elements, such as WPRE; CMV enhancers; the R-U5' segment in LTR of HTLV-I (Mol. Cell. Biol., Vol. 8(1), p. 466-472, 1988); SV40 enhancer; and the intron sequence between exons 2 and 3 of rabbit β -globin (Proc. Natl. Acad. Sci. USA., Vol. 78(3), p. 1527-31, 1981). It will be appreciated by those skilled in the art that the design of the expression vector can depend on such factors as the

choice of the host cell to be transformed, the level of expression desired, etc. A vector can be introduced into host cells to thereby produce transcripts, proteins, or peptides, including fusion proteins or peptides, encoded by nucleic acids as described herein (e.g., clustered regularly interspersed short palindromic repeats (CRISPR) transcripts, proteins, enzymes, mutant forms thereof, fusion proteins thereof, etc.).

[0012] Advantageous vectors include lentiviruses and adeno-associated viruses, and types of such vectors can also be selected for targeting particular types of cells.

[0013] In one aspect, the invention provides a vector comprising a regulatory element operably linked to an enzyme-coding sequence encoding a CRISPR enzyme comprising one or more nuclear localization sequences. In some embodiments, said regulatory element drives transcription of the CRISPR enzyme in a eukaryotic cell such that said CRISPR enzyme accumulates in a detectable amount in the nucleus of the eukaryotic cell. In some embodiments, the regulatory element is a polymerase II promoter. In some embodiments, the CRISPR enzyme is a type II CRISPR system enzyme. In some embodiments, the CRISPR enzyme is a Cas9 enzyme. In some embodiments, the CRISPR enzyme is *S. pneumoniae*, *S. pyogenes* or *S. thermophilus* Cas9, and may include mutated Cas9 derived from these organisms. In some embodiments, the CRISPR enzyme is codon-optimized for expression in a eukaryotic cell. In some embodiments, the CRISPR enzyme directs cleavage of one or two strands at the location of the target sequence. In some embodiments, the CRISPR enzyme lacks DNA strand cleavage activity.

[0014] In one aspect, the invention provides a CRISPR enzyme comprising one or more nuclear localization sequences of sufficient strength to drive accumulation of said CRISPR enzyme in a detectable amount in the nucleus of a eukaryotic cell. In some embodiments, the CRISPR enzyme is a type II CRISPR system enzyme. In some embodiments, the CRISPR enzyme is a Cas9 enzyme. In some embodiments, the Cas9 enzyme is *S. pneumoniae*, *S. pyogenes* or *S. thermophilus* Cas9, and may include mutated Cas9 derived from these organisms. The enzyme may be a Cas9 homolog or ortholog. In some embodiments, the CRISPR enzyme lacks the ability to cleave one or more strands of a target sequence to which it binds.

[0015] In one aspect, the invention provides a eukaryotic host cell comprising (a) a first regulatory element operably linked to a tracr mate sequence and one or more insertion sites for inserting one or more guide sequences upstream of the tracr mate sequence, wherein when expressed, the guide sequence directs sequence-specific binding of a CRISPR complex to a target sequence in a eukaryotic cell, wherein the CRISPR complex comprises a CRISPR enzyme complexed with (1) the guide sequence that is hybridized to the target sequence, and (2) the tracr mate sequence that is hybridized to the tracr sequence; and/or (b) a second regulatory element operably linked to an enzyme-coding sequence encoding said CRISPR enzyme comprising a nuclear localization sequence. In some embodiments, the host cell comprises components (a) and (b). In some embodiments, component (a), component (b), or components (a) and (b) are stably integrated into a genome of the host eukaryotic cell. In some embodiments, component (a) further comprises the tracr sequence downstream of the tracr mate sequence under the control of the first regulatory

element. In some embodiments, component (a) further comprises two or more guide sequences operably linked to the first regulatory element, wherein when expressed, each of the two or more guide sequences direct sequence specific binding of a CRISPR complex to a different target sequence in a eukaryotic cell. In some embodiments, the eukaryotic host cell further comprises a third regulatory element, such as a polymerase III promoter, operably linked to said tracr sequence. In some embodiments, the tracr sequence exhibits at least 50%, 60%, 70%, 80%, 90%, 95%, or 99% of sequence complementarity along the length of the tracr mate sequence when optimally aligned. In some embodiments, the CRISPR enzyme comprises one or more nuclear localization sequences of sufficient strength to drive accumulation of said CRISPR enzyme in a detectable amount in the nucleus of a eukaryotic cell. In some embodiments, the CRISPR enzyme is a type II CRISPR system enzyme. In some embodiments, the CRISPR enzyme is a Cas9 enzyme. In some embodiments, the Cas9 enzyme is *S. pneumoniae*, *S. pyogenes* or *S. thermophilus* Cas9, and may include mutated Cas9 derived from these organisms. The enzyme may be a Cas9 homolog or ortholog. In some embodiments, the CRISPR enzyme is codon-optimized for expression in a eukaryotic cell. In some embodiments, the CRISPR enzyme directs cleavage of one or two strands at the location of the target sequence. In some embodiments, the CRISPR enzyme lacks DNA strand cleavage activity. In some embodiments, the first regulatory element is a polymerase III promoter. In some embodiments, the second regulatory element is a polymerase II promoter. In some embodiments, the guide sequence is at least 15, 16, 17, 18, 19, 20, 25 nucleotides, or between 10-30, or between 15-25, or between 15-20 nucleotides in length. In an aspect, the invention provides a non-human eukaryotic organism; preferably a multicellular eukaryotic organism, comprising a eukaryotic host cell according to any of the described embodiments. In other aspects, the invention provides a eukaryotic organism; preferably a multicellular eukaryotic organism, comprising a eukaryotic host cell according to any of the described embodiments. The organism in some embodiments of these aspects may be an animal; for example a mammal. Also, the organism may be an arthropod such as an insect. The organism also may be a plant. Further, the organism may be a fungus.

[0016] In one aspect, the invention provides a kit comprising one or more of the components described herein. In some embodiments, the kit comprises a vector system and instructions for using the kit. In some embodiments, the vector system comprises (a) a first regulatory element operably linked to a tracr mate sequence and one or more insertion sites for inserting one or more guide sequences upstream of the tracr mate sequence, wherein when expressed, the guide sequence directs sequence-specific binding of a CRISPR complex to a target sequence in a eukaryotic cell, wherein the CRISPR complex comprises a CRISPR enzyme complexed with (1) the guide sequence that is hybridized to the target sequence, and (2) the tracr mate sequence that is hybridized to the tracr sequence; and/or (b) a second regulatory element operably linked to an enzyme-coding sequence encoding said CRISPR enzyme comprising a nuclear localization sequence. In some embodiments, the kit comprises components (a) and (b) located on the same or different vectors of the system. In some embodiments, component (a) further comprises the

tracr sequence downstream of the tracr mate sequence under the control of the first regulatory element. In some embodiments, component (a) further comprises two or more guide sequences operably linked to the first regulatory element, wherein when expressed, each of the two or more guide sequences direct sequence specific binding of a CRISPR complex to a different target sequence in a eukaryotic cell. In some embodiments, the system further comprises a third regulatory element, such as a polymerase III promoter, operably linked to said tracr sequence. In some embodiments, the tracr sequence exhibits at least 50%, 60%, 70%, 80%, 90%, 95%, or 99% of sequence complementarity along the length of the tracr mate sequence when optimally aligned. In some embodiments, the CRISPR enzyme comprises one or more nuclear localization sequences of sufficient strength to drive accumulation of said CRISPR enzyme in a detectable amount in the nucleus of a eukaryotic cell. In some embodiments, the CRISPR enzyme is a type II CRISPR system enzyme. In some embodiments, the CRISPR enzyme is a Cas9 enzyme. In some embodiments, the Cas9 enzyme is *S. pneumoniae*, *S. pyogenes* or *S. thermophilus* Cas9, and may include mutated Cas9 derived from these organisms. The enzyme may be a Cas9 homolog or ortholog. In some embodiments, the CRISPR enzyme is codon-optimized for expression in a eukaryotic cell. In some embodiments, the CRISPR enzyme directs cleavage of one or two strands at the location of the target sequence. In some embodiments, the CRISPR enzyme lacks DNA strand cleavage activity. In some embodiments, the first regulatory element is a polymerase III promoter. In some embodiments, the second regulatory element is a polymerase II promoter. In some embodiments, the guide sequence is at least 15, 16, 17, 18, 19, 20, 25 nucleotides, or between 10-30, or between 15-25, or between 15-20 nucleotides in length.

[0017] In one aspect, the invention provides a method of modifying a target polynucleotide in a eukaryotic cell. In some embodiments, the method comprises allowing a CRISPR complex to bind to the target polynucleotide to effect cleavage of said target polynucleotide thereby modifying the target polynucleotide, wherein the CRISPR complex comprises a CRISPR enzyme complexed with a guide sequence hybridized to a target sequence within said target polynucleotide, wherein said guide sequence is linked to a tracr mate sequence which in turn hybridizes to a tracr sequence. In some embodiments, said cleavage comprises cleaving one or two strands at the location of the target sequence by said CRISPR enzyme. In some embodiments, said cleavage results in decreased transcription of a target gene. In some embodiments, the method further comprises repairing said cleaved target polynucleotide by homologous recombination with an exogenous template polynucleotide, wherein said repair results in a mutation comprising an insertion, deletion, or substitution of one or more nucleotides of said target polynucleotide. In some embodiments, said mutation results in one or more amino acid changes in a protein expressed from a gene comprising the target sequence. In some embodiments, the method further comprises delivering one or more vectors to said eukaryotic cell, wherein the one or more vectors drive expression of one or more of: the CRISPR enzyme, the guide sequence linked to the tracr mate sequence, and the tracr sequence. In some embodiments, said vectors are delivered to the eukaryotic cell in a subject. In some embodiments, said modifying takes place in said eukaryotic cell in a cell culture. In some

embodiments, the method further comprises isolating said eukaryotic cell from a subject prior to said modifying. In some embodiments, the method further comprises returning said eukaryotic cell and/or cells derived therefrom to said subject.

[0018] In one aspect, the invention provides a method of modifying expression of a polynucleotide in a eukaryotic cell. In some embodiments, the method comprises allowing a CRISPR complex to bind to the polynucleotide such that said binding results in increased or decreased expression of said polynucleotide; wherein the CRISPR complex comprises a CRISPR enzyme complexed with a guide sequence hybridized to a target sequence within said polynucleotide, wherein said guide sequence is linked to a tracr mate sequence which in turn hybridizes to a tracr sequence. In some embodiments, the method further comprises delivering one or more vectors to said eukaryotic cells, wherein the one or more vectors drive expression of one or more of: the CRISPR enzyme, the guide sequence linked to the tracr mate sequence, and the tracr sequence.

[0019] In one aspect, the invention provides a method of generating a model eukaryotic cell comprising a mutated disease gene. In some embodiments, a disease gene is any gene associated an increase in the risk of having or developing a disease. In some embodiments, the method comprises (a) introducing one or more vectors into a eukaryotic cell, wherein the one or more vectors drive expression of one or more of: a CRISPR enzyme, a guide sequence linked to a tracr mate sequence, and a tracr sequence; and (b) allowing a CRISPR complex to bind to a target polynucleotide to effect cleavage of the target polynucleotide within said disease gene, wherein the CRISPR complex comprises the CRISPR enzyme complexed with (1) the guide sequence that is hybridized to the target sequence within the target polynucleotide, and (2) the tracr mate sequence that is hybridized to the tracr sequence, thereby generating a model eukaryotic cell comprising a mutated disease gene. In some embodiments, said cleavage comprises cleaving one or two strands at the location of the target sequence by said CRISPR enzyme. In some embodiments, said cleavage results in decreased transcription of a target gene. In some embodiments, the method further comprises repairing said cleaved target polynucleotide by homologous recombination with an exogenous template polynucleotide, wherein said repair results in a mutation comprising an insertion, deletion, or substitution of one or more nucleotides of said target polynucleotide. In some embodiments, said mutation results in one or more amino acid changes in a protein expression from a gene comprising the target sequence.

[0020] In one aspect, the invention provides a method for developing a biologically active agent that modulates a cell signaling event associated with a disease gene. In some embodiments, a disease gene is any gene associated an increase in the risk of having or developing a disease. In some embodiments, the method comprises (a) contacting a test compound with a model cell of any one of the described embodiments; and (b) detecting a change in a readout that is indicative of a reduction or an augmentation of a cell signaling event associated with said mutation in said disease gene, thereby developing said biologically active agent that modulates said cell signaling event associated with said disease gene.

[0021] In one aspect, the invention provides a recombinant polynucleotide comprising a guide sequence upstream of a

tracr mate sequence, wherein the guide sequence when expressed directs sequence-specific binding of a CRISPR complex to a corresponding target sequence present in a eukaryotic cell. In some embodiments, the target sequence is a viral sequence present in a eukaryotic cell. In some embodiments, the target sequence is a proto-oncogene or an oncogene.

[0022] In one aspect the invention provides for a method of selecting one or more prokaryotic cell(s) by introducing one or more mutations in a gene in the one or more prokaryotic cell (s), the method comprising: introducing one or more vectors into the prokaryotic cell (s), wherein the one or more vectors drive expression of one or more of: a CRISPR enzyme, a guide sequence linked to a tracr mate sequence, a tracr sequence, and an editing template; wherein the editing template comprises the one or more mutations that abolish CRISPR enzyme cleavage; allowing homologous recombination of the editing template with the target polynucleotide in the cell(s) to be selected; allowing a CRISPR complex to bind to a target polynucleotide to effect cleavage of the target polynucleotide within said gene, wherein the CRISPR complex comprises the CRISPR enzyme complexed with (1) the guide sequence that is hybridized to the target sequence within the target polynucleotide, and (2) the tracr mate sequence that is hybridized to the tracr sequence, wherein binding of the CRISPR complex to the target polynucleotide induces cell death, thereby allowing one or more prokaryotic cell(s) in which one or more mutations have been introduced to be selected. In a preferred embodiment, the CRISPR enzyme is Cas9. In another aspect of the invention the cell to be selected may be a eukaryotic cell. Aspects of the invention allow for selection of specific cells without requiring a selection marker or a two-step process that may include a counter-selection system.

[0023] Accordingly, it is an object of the invention not to encompass within the invention any previously known product, process of making the product, or method of using the product such that Applicants reserve the right and hereby disclose a disclaimer of any previously known product, process, or method. It is further noted that the invention does not intend to encompass within the scope of the invention any product, process, or making of the product or method of using the product, which does not meet the written description and enablement requirements of the USPTO (35 U.S.C. § 112, first paragraph) or the EPO (Article 83 of the EPC), such that Applicants reserve the right and hereby disclose a disclaimer of any previously described product, process of making the product, or method of using the product.

[0024] It is noted that in this disclosure and particularly in the claims and/or paragraphs, terms such as “comprises”, “comprising”, “comprising” and the like can have the meaning attributed to it in U.S. Patent law; e.g., they can mean “includes”, “included”, “including”, and the like; and that terms such as “consisting essentially of” and “consists essentially of” have the meaning ascribed to them in U.S. Patent law, e.g., they allow for elements not explicitly recited, but exclude elements that are found in the prior art or that affect a basic or novel characteristic of the invention. These and other embodiments are disclosed or are obvious from and encompassed by, the following Detailed Description.

BRIEF DESCRIPTION OF THE DRAWINGS

[0025] The novel features of the invention are set forth with particularity in the appended claims. A better understanding of the features and advantages of the present invention will be obtained by reference to the following detailed description that sets forth illustrative embodiments, in which the principles of the invention are utilized, and the accompanying drawings of which:

[0026] FIG. 1 shows a schematic model of the CRISPR system. The Cas9 nuclease from *Streptococcus pyogenes* (yellow) is targeted to genomic DNA by a synthetic guide RNA (sgRNA) consisting of a 20-nt guide sequence (blue) and a scaffold (red). The guide sequence base-pairs with the DNA target (blue), directly upstream of a requisite 5'-NGG protospacer adjacent motif (PAM; magenta), and Cas9 mediates a double-stranded break (DSB) ~3 bp upstream of the PAM (red triangle).

[0027] FIGS. 2A-2F show an exemplary CRISPR system, a possible mechanism of action, an example adaptation for expression in eukaryotic cells, and results of tests assessing nuclear localization and CRISPR activity. FIG. 2C discloses SEQ ID NOS 279-280, respectively, in order of appearance. FIG. 2E discloses SEQ ID NOS 281-283, respectively, in order of appearance. FIG. 2F discloses SEQ ID NOS 284-288, respectively, in order of appearance.

[0028] FIG. 3A-3C shows an exemplary expression cassette for expression of CRISPR system elements in eukaryotic cells, predicted structures of example guide sequences, and CRISPR system activity as measured in eukaryotic and prokaryotic cells (SEQ ID NOS 289-298, respectively, in order of appearance).

[0029] FIGS. 4A-4D show results of an evaluation of SpCas9 specificity for an example target. FIG. 4A discloses SEQ ID NOS 299, 282 and 300-310, respectively, in order of appearance. FIG. 4C discloses SEQ ID NO: 299.

[0030] FIGS. 5A-5G show an exemplary vector system and results for its use in directing homologous recombination in eukaryotic cells. FIG. 5E discloses SEQ ID NO: 311. FIG. 5F discloses SEQ ID NOS 312-313, respectively, in order of appearance. FIG. 5G discloses SEQ ID NOS 314-318, respectively, in order of appearance.

[0031] FIG. 6 provides a table of protospacer sequences (SEQ ID NOS 33, 32, 31, 322-327, 35, 34 and 330-334, respectively, in order of appearance) and summarizes modification efficiency results for protospacer targets designed based on exemplary *S. pyogenes* and *S. thermophilus* CRISPR systems with corresponding PAMs against loci in human and mouse genomes. Cells were transfected with Cas9 and either pre-crRNA/tracrRNA or chimeric RNA, and analyzed 72 hours after transfection. Percent indels are calculated based on Surveyor assay results from indicated cell lines (N=3 for all protospacer targets, errors are S.E.M., N.D. indicates not detectable using the Surveyor assay, and N.T. indicates not tested in this study).

[0032] FIGS. 7A-7C show a comparison of different tracrRNA transcripts for Cas9-mediated gene targeting. FIG. 7A discloses SEQ ID NOS 335-336, respectively, in order of appearance.

[0033] FIG. 8 shows a schematic of a surveyor nuclease assay for detection of double strand break-induced micro-insertions and -deletions.

[0034] FIGS. 9A-9B show exemplary bicistronic expression vectors for expression of CRISPR system elements in eukaryotic cells. FIG. 9A discloses SEQ ID NOS 337-339,

respectively, in order of appearance. FIG. 9B discloses SEQ ID NOS 340-342, respectively, in order of appearance.

[0035] FIG. 10A-10D shows a bacterial plasmid transformation interference assay, expression cassettes and plasmids used therein, and transformation efficiencies of cells used therein. FIG. 10A discloses SEQ ID NOS 343-345, respectively, in order of appearance.

[0036] FIGS. 11A-11C show histograms of distances between adjacent *S. pyogenes* SF370 locus 1 PAM (NGG) (FIG. 10A) and *S. thermophilus* LMD9 locus 2 PAM (NNA-GAAW) (FIG. 10B) in the human genome; and distances for each PAM by chromosome (Chr) (FIG. 10C).

[0037] FIGS. 12A-12C show an exemplary CRISPR system, an example adaptation for expression in eukaryotic cells, and results of tests assessing CRISPR activity. FIG. 12B discloses SEQ ID NOS 346-347, respectively, in order of appearance. FIG. 12C discloses SEQ ID NO: 348.

[0038] FIGS. 13A-13C show exemplary manipulations of a CRISPR system for targeting of genomic loci in mammalian cells. FIG. 13A discloses SEQ ID NO: 349. FIG. 13B discloses SEQ ID NOS 350-352, respectively, in order of appearance.

[0039] FIGS. 14A-14B show the results of a Northern blot analysis of crRNA processing in mammalian cells. FIG. 14A discloses SEQ ID NO: 353.

[0040] FIG. 15A-15B shows an exemplary selection of protospacers in the human PVALB and mouse Th loci. FIG. 15A discloses SEQ ID NO: 354. FIG. 15B discloses SEQ ID NO: 355.

[0041] FIG. 16 shows example protospacer and corresponding PAM sequence targets of the *S. thermophilus* CRISPR system in the human EMX1 locus (SEQ ID NO: 348).

[0042] FIG. 17 provides a table of sequences for primers and probes (SEQ ID NOS 36-39 and 356-363, respectively, in order of appearance) used for Surveyor, RFLP, genomic sequencing, and Northern blot assays.

[0043] FIGS. 18A-18C show exemplary manipulation of a CRISPR system with chimeric RNAs and results of SURVEYOR assays for system activity in eukaryotic cells. FIG. 18A discloses SEQ ID NO: 364, respectively, in order of appearance.

[0044] FIGS. 19A-19B show a graphical representation of the results of SURVEYOR assays for CRISPR system activity in eukaryotic cells (SEQ ID NOS 365-443, respectively, in order of appearance).

[0045] FIG. 20 shows an exemplary visualization of some *S. pyogenes* Cas9 target sites in the human genome using the UCSC genome browser.

[0046] FIG. 21 shows predicted secondary structures for exemplary chimeric RNAs comprising a guide sequence, tracr mate sequence, and tracr sequence (SEQ ID NOS 444-463, respectively, in order of appearance).

[0047] FIG. 22 shows exemplary bicistronic expression vectors for expression of CRISPR system elements in eukaryotic cells (SEQ ID NOS 464 and 341-342, respectively, in order of appearance).

[0048] FIG. 23A-23B shows that Cas9 nuclease activity against endogenous targets may be exploited for genome editing. (a) Concept of genome editing using the CRISPR system. The CRISPR targeting construct directed cleavage of a chromosomal locus and was co-transformed with an editing template that recombined with the target to prevent cleavage. Kanamycin-resistant transformants that survived

CRISPR attack contained modifications introduced by the editing template. tracr, trans-activating CRISPR RNA; aphA-3, kanamycin resistance gene. (b) Transformation of crR6M DNA in R6^{8232.5} cells with no editing template, the R6 wild-type srtA or the R6370.1 editing templates. Recombination of either R6 srtA or R6^{370.1} prevented cleavage by Cas9. Transformation efficiency was calculated as colony forming units (cfu) per μg of crR6M DNA; the mean values with standard deviations from at least three independent experiments are shown. PCR analysis was performed on 8 clones in each transformation. "Un." indicates the unedited srtA locus of strain R6^{8232.5}; "Ed." shows the editing template. R6^{8232.5} and R6^{370.1} targets are distinguished by restriction with EaeI.

[0049] FIG. 24A-24C shows analysis of PAM and seed sequences that eliminate Cas9 cleavage. (a) PCR products with randomized PAM sequences or randomized seed sequences were transformed in crR6 cells (SEQ ID NOS 465-469, respectively, in order of appearance). These cells expressed Cas9 loaded with a crRNA that targeted a chromosomal region of R6^{8232.5} cells (highlighted in pink) that is absent from the R6 genome. More than 2×10^5 chloramphenicol-resistant transformants, carrying inactive PAM or seed sequences, were combined for amplification and deep sequencing of the target region. (b) Relative proportion of number of reads after transformation of the random PAM constructs in crR6 cells (compared to number of reads in R6 transformants). The relative abundance for each 3-nucleotide PAM sequence is shown. Severely underrepresented sequences (NGG) are shown in red; partially underrepresented one in orange (NAG) (c) Relative proportion of number of reads after transformation of the random seed sequence constructs in crR6 cells (compared to number of reads in R6 transformants). The relative abundance of each nucleotide for each position of the first 20 nucleotides of the protospacer sequence is shown (SEQ ID NO: 470). High abundance indicates lack of cleavage by Cas9, i.e. a CRISPR inactivating mutation. The grey line shows the level of the WT sequence. The dotted line represents the level above which a mutation significantly disrupts cleavage (See section "Analysis of deep sequencing data" in Example 5)

[0050] FIG. 25A-25F shows introduction of single and multiple mutations using the CRISPR system in *S. pneumoniae*. (a) Nucleotide and amino acid sequences of the wild-type and edited (green nucleotides; underlined amino acid residues) bgaA. The protospacer, PAM and restriction sites are shown (SEQ ID NOS 471-475 and 472, respectively, in order of appearance). (b) Transformation efficiency of cells transformed with targeting constructs in the presence of an editing template or control. (c) PCR analysis for 8 transformants of each editing experiment followed by digestion with BtgZI (R \rightarrow A) and TseI (NE \rightarrow AA). Deletion of bgaA was revealed as a smaller PCR product. (d) Miller assay to measure the β -galactosidase activity of WT and edited strains. (e) For a single-step, double deletion the targeting construct contained two spacers (in this case matching srtA and bgaA) and was co-transformed with two different editing templates (f) PCR analysis for 8 transformants to detect deletions in srtA and bgaA loci. 6/8 transformants contained deletions of both genes.

[0051] FIG. 26A-26D provides mechanisms underlying editing using the CRISPR system. (a) A stop codon was introduced in the erythromycin resistance gene ermAM to generate strain JEN53. The wild-type sequence can be

restored by targeting the stop codon with the CRISPR:ermAM(stop) construct, and using the ermAM wild-type sequence as an editing template. (b) Mutant and wild-type ermAM sequences (SEQ ID NOS 476-479, respectively, in order of appearance). (c) Fraction of erythromycin-resistant (erm^R) cfu calculated from total or kanamycin-resistant (kan^R) cfu. (d) Fraction of total cells that acquire both the CRISPR construct and the editing template. Co-transformation of the CRISPR targeting construct produced more transformants (t-test, $p=0.011$). In all cases the values show the mean \pm s.d. for three independent experiments.

[0052] FIG. 27A-27D illustrates genome editing with the CRISPR system in *E. coli*. (a) A kanamycin-resistant plasmid carrying the CRISPR array (pCRISPR) targeting the gene to edit may be transformed in the HME63 recombinering strain containing a chloramphenicol-resistant plasmid harboring cas9 and tracr (pCas9), together with an oligonucleotide specifying the mutation. (b) A K42T mutation conferring streptomycin resistance was introduced in the rpsL gene (SEQ ID NOS 480-483, respectively, in order of appearance) (c) Fraction of streptomycin-resistant ($strep^R$) cfu calculated from total or kanamycin-resistant (kan^R) cfu. (d) Fraction of total cells that acquire both the pCRISPR plasmid and the editing oligonucleotide. Co-transformation of the pCRISPR targeting plasmid produced more transformants (t-test, $p=0.004$). In all cases the values showed the mean \pm s.d. for three independent experiments.

[0053] FIG. 28A-28B illustrates the transformation of crR6 genomic DNA leads to editing of the targeted locus (a) The IS1167 element of *S. pneumoniae* R6 was replaced by the CRISPR01 locus of *S. pyogenes* SF370 to generate crR6 strain. This locus encodes for the Cas9 nuclease, a CRISPR array with six spacers, the tracrRNA that is required for crRNA biogenesis and Cas1, Cas2 and Csn2, proteins not necessary for targeting. Strain crR6M contains a minimal functional CRISPR system without cas1, cas2 and csn2. The aphA-3 gene encodes kanamycin resistance. Protospacers from the streptococcal bacteriophages ϕ 8232.5 and ϕ 370.1 were fused to a chloramphenicol resistance gene (cat) and integrated in the srtA gene of strain R6 to generate strains R68232.5 and R6370.1. (b) Left panel: Transformation of crR6 and crR6M genomic DNA in R6^{8232.5} and R6^{370.1}. As a control of cell competence a streptomycin resistant gene was also transformed. Right panel: PCR analysis of 8 R6^{8232.5} transformants with crR6 genomic DNA. Primers that amplify the srtA locus were used for PCR. 7/8 genotyped colonies replaced the R68232.5 srtA locus by the WT locus from the crR6 genomic DNA.

[0054] FIG. 29A-29F provides chromatograms of DNA sequences of edited cells obtained in this study. In all cases the wild-type and mutant protospacer and PAM sequences (or their reverse complement) are indicated. When relevant, the amino acid sequence encoded by the protospacer is provided. For each editing experiment, all strains for which PCR and restriction analysis corroborated the introduction of the desired modification were sequenced. A representative chromatogram is shown. (a) Chromatogram for the introduction of a PAM mutation into the R6^{8232.5} target (FIG. 23d) (SEQ ID NOS 484-485, respectively, in order of appearance). (b) Chromatograms for the introduction of the R>A and NE>AA mutations into β -galactosidase (bgaA) (FIG. 25c) (SEQ ID NOS 471-475 and 472, respectively, in order of appearance). (c) Chromatogram for the introduction of a 6664 bp deletion within bgaA ORF (FIGS. 25c and 25f).

The dotted line indicates the limits of the deletion (SEQ ID NOS 486-488, respectively, in order of appearance). (d) Chromatogram for the introduction of a 729 bp deletion within srtA ORF (FIG. 25f). The dotted line indicates the limits of the deletion (SEQ ID NOS 489-491, respectively, in order of appearance). (e) Chromatograms for the generation of a premature stop codon within ermAM (FIG. 33) (SEQ ID NOS 492-495, respectively, in order of appearance). (f) rpsL editing in *E. coli* (FIG. 27) (SEQ ID NOS 480-483, respectively, in order of appearance).

[0055] FIG. 30A-30C illustrates CRISPR immunity against random *S. pneumoniae* targets containing different PAMs. (a) Position of the 10 random targets on the *S. pneumoniae* R6 genome. The chosen targets have different PAMs and are on both strands. (b) Spacers corresponding to the targets were cloned in a minimal CRISPR array on plasmid pLZ12 and transformed into strain crR6Rc, which supplies the processing and targeting machinery in trans. (c) Transformation efficiency of the different plasmids in strain R6 and crR6Rc. No colonies were recovered for the transformation of pDB99-108 (T1-T10) in crR6Rc. The dashed line represents limit of detection of the assay.

[0056] FIG. 31 provides a general scheme for targeted genome editing. To facilitate targeted genome editing, crR6M was further engineered to contain tracrRNA, Cas9 and only one repeat of the CRISPR array followed by kanamycin resistance marker (aphA-3), generating strain crR6Rk. DNA from this strain is used as a template for PCR with primers designed to introduce a new spacer (green box designated with N). The left and right PCRs are assembled using the Gibson method to create the targeting construct. Both the targeting and editing constructs are then transformed into strain crR6Rc, which is a strain equivalent to crR6Rk but has the kanamycin resistance marker replaced by a chloramphenicol resistance marker (cat). About 90% of the kanamycin-resistant transformants contain the desired mutation.

[0057] FIG. 32 illustrates the distribution of distances between PAMs. NGG and CCN that are considered to be valid PAMs. Data is shown for the *S. pneumoniae* R6 genome as well as for a random sequence of the same length and with the same GC-content (39.7%). The dotted line represents the average distance (12) between PAMs in the R6 genome.

[0058] FIG. 33A-33D illustrates CRISPR-mediated editing of the ermAM locus using genomic DNA as targeting construct. To use genomic DNA as targeting construct it is necessary to avoid CRISPR autoimmunity, and therefore a spacer against a sequence not present in the chromosome must be used (in this case the ermAM erythromycin resistance gene). (a) Nucleotide and amino acid sequences of the wild-type and mutated (red letters) ermAM gene. The protospacer and PAM sequences are shown (SEQ ID NOS 492-495, respectively, in order of appearance). (b) A schematic for CRISPR-mediated editing of the ermAM locus using genomic DNA. A construct carrying an ermAM-targeting spacer (blue box) is made by PCR and Gibson assembly, and transformed into strain crR6Rc, generating strain JEN37. The genomic DNA of JEN37 was then used as a targeting construct, and was co-transformed with the editing template into JEN38, a strain in which the srtA gene was replaced by a wild-type copy of ermAM. Kanamycin-resistant transformants contain the edited genotype (JEN43). (c) Number of kanamycin-resistant cells obtained after co-

transformation of targeting and editing or control templates. In the presence of the control template 5.4×10^3 cfu/ml were obtained, and 4.3×10^5 cfu/ml when the editing template was used. This difference indicates an editing efficiency of about 99% [$(4.3 \times 10^5 - 5.4 \times 10^3) / 4.3 \times 10^5$]. (d) To check for the presence of edited cells seven kanamycin-resistant clones and JEN38 were streaked on agar plates with (erm+) or without (erm-) erythromycin. Only the positive control displayed resistance to erythromycin. The ermAM mut genotype of one of these transformants was also verified by DNA sequencing (FIG. 29e).

[0059] FIG. 34A-34D illustrates sequential introduction of mutations by CRISPR-mediated genome editing. (a) A schematic for sequential introduction of mutations by CRISPR-mediated genome editing. First, R6 is engineered to generate crR6Rk. crR6Rk is co-transformed with a *srtA*-targeting construct fused to cat for chloramphenicol selection of edited cells, along with an editing construct for a Δ *srtA* in-frame deletion. Strain crR6 Δ *srtA* is generated by selection on chloramphenicol. Subsequently, the Δ *srtA* strain is co-transformed with a *bgaA*-targeting construct fused to *aphA-3* for kanamycin selection of edited cells, and an editing construct containing a Δ *bgaA* in-frame deletion. Finally, the engineered CRISPR locus can be erased from the chromosome by first co-transforming R6 DNA containing the wild-type IS1167 locus and a plasmid carrying a *bgaA* protospacer (pDB97), and selection on spectinomycin. (b) PCR analysis for 8 chloramphenicol (Cam)-resistant transformants to detect the deletion in the *srtA* locus. (c) β -galactosidase activity as measured by Miller assay. In *S. pneumoniae*, this enzyme is anchored to the cell wall by sortase A. Deletion of the *srtA* gene results in the release of β -galactosidase into the supernatant. Δ *bgaA* mutants show no activity. (d) PCR analysis for 8 spectinomycin (Spec)-resistant transformants to detect the replacement of the CRISPR locus by wild-type IS1167.

[0060] FIG. 35A-35C illustrates the background mutation frequency of CRISPR in *S. pneumoniae*. (a) Transformation of the CRISPR: \emptyset or CRISPR:erm(stop) targeting constructs in JEN53, with or without the ermAM editing template. The difference in kan^R CFU between CRISPR: \emptyset and CRISPR:erm(stop) indicates that Cas9 cleavage kills non-edited cells. Mutants that escape CRISPR interference in the absence of editing template are observed at a frequency of 3×10^{-3} . (b) PCR analysis of the CRISPR locus of escapers shows that 7/8 have a spacer deletion. (c) Escaper #2 carries a point mutation in *cas9* (SEQ ID NOS 496-499, respectively, in order of appearance).

[0061] FIG. 36 illustrates that the essential elements of the *S. pyogenes* CRISPR locus 1 are reconstituted in *E. coli* using pCas9. The plasmid contained tracrRNA, Cas9, as well as a leader sequence driving the crRNA array. The pCRISPR plasmids contained the leader and the array only. Spacers may be inserted into the crRNA array between BsaI sites using annealed oligonucleotides (SEQ ID NOS 343, 500 and 127, respectively, in order of appearance). Oligonucleotide design is shown at bottom. pCas9 carried chloramphenicol resistance (CmR) and is based on the low-copy pACYC184 plasmid backbone. pCRISPR is based on the high-copy number pZE21 plasmid. Two plasmids were required because a pCRISPR plasmid containing a spacer targeting the *E. coli* chromosome may not be constructed using this organism as a cloning host if Cas9 is also present (it will kill the host).

[0062] FIG. 37 illustrates CRISPR-directed editing in *E. coli* MG1655. An oligonucleotide (W542) carrying a point mutation that both confers streptomycin resistance and abolishes CRISPR immunity, together with a plasmid targeting *rpsL* (pCRISPR:*rpsL*) or a control plasmid (pCRISPR: \emptyset) were co-transformed into wild-type *E. coli* strain MG1655 containing pCas9. Transformants were selected on media containing either streptomycin or kanamycin. Dashed line indicates limit of detection of the transformation assay.

[0063] FIG. 38A-38B illustrates the background mutation frequency of CRISPR in *E. coli* HME63. (a) Transformation of the pCRISPR: \emptyset or pCRISPR:*rpsL* plasmids into HME63 competent cells. Mutants that escape CRISPR interference were observed at a frequency of 2.6×10^{-4} . (b) Amplification of the CRISPR array of escapers showed that 8/8 have deleted the spacer.

[0064] FIGS. 39A-39D show a circular depiction of the phylogenetic analysis revealing five families of Cas9s, including three groups of large Cas9s (~1400 amino acids) and two of small Cas9s (~1100 amino acids).

[0065] FIGS. 40A-40F show the linear depiction of the phylogenetic analysis revealing five families of Cas9s, including three groups of large Cas9s (~1400 amino acids) and two of small Cas9s (~1100 amino acids).

[0066] FIG. 41A-41M shows sequences where the mutation points are located within the SpCas9 gene (SEQ ID NOS 501-502, respectively, in order of appearance).

[0067] FIG. 42 shows a schematic construct in which the transcriptional activation domain (VP64) is fused to Cas9 with two mutations in the catalytic domains (D10 and H840).

[0068] FIG. 43A-43D shows genome editing via homologous recombination. (a) Schematic of SpCas9 nickase, with D10A mutation in the RuvC I catalytic domain. (b) Schematic representing homologous recombination (HR) at the human EMXJ locus using either sense or antisense single stranded oligonucleotides as repair templates. Red arrow above indicates sgRNA cleavage site; PCR primers for genotyping (Tables J and K) are indicated as arrows in right panel. (c) Sequence of region modified by HR. d, SURVEYOR assay for wildtype (wt) and nickase (D10A) SpCas9-mediated indels at the EMXJ target 1 locus (n=3) (SEQ ID NOS 503-505, 503, 506 and 505, respectively, in order of appearance). Arrows indicate positions of expected fragment sizes.

[0069] FIGS. 44A-44B show single vector designs for SpCas9. FIG. 44A discloses SEQ ID NOS 320-321 and 328, respectively, in order of appearance. FIG. 44B discloses SEQ ID NO: 329.

[0070] FIG. 45 shows quantification of cleavage of NLS-Csn1 constructs NLS-Csn1, Csn1, Csn1-NLS, NLS-Csn1-NLS, NLS-Csn1-GFP-NLS and UnTFN.

[0071] FIG. 46 shows index frequency of NLS-Cas9, Cas9, Cas9-NLS and NLS-Cas9-NLS.

[0072] FIG. 47 shows a gel demonstrating that SpCas9 with nickase mutations (individually) do not induce double strand breaks.

[0073] FIG. 48A-48B shows a design of the oligo DNA used as Homologous Recombination (HR) template in this experiment and a comparison of HR efficiency induced by different combinations of Cas9 protein and HR template.

[0074] FIG. 49A shows the Conditional Cas9, Rosa26 targeting vector map.

[0075] FIG. 49B shows the Constitutive Cas9, Rosa26 targeting vector map.

[0076] FIG. 50A-50H show the sequences of each element present in the vector maps of FIGS. 49A-B (SEQ ID NOS 507-516, respectively, in order of appearance).

[0077] FIG. 51 shows a schematic of the important elements in the Constitutive and Conditional Cas9 constructs.

[0078] FIG. 52 shows the functional validation of the expression of Constitutive and Conditional Cas9 constructs.

[0079] FIG. 53 shows the validation of Cas9 nuclease activity by Surveyor.

[0080] FIG. 54 shows the quantification of Cas9 nuclease activity.

[0081] FIG. 55 shows construct design and homologous recombination (HR) strategy.

[0082] FIG. 56 shows the genomic PCR genotyping results for the constitutive (Right) and conditional (Left) constructs at two different gel exposure times (top row for 3 min and bottom row for 1 min).

[0083] FIG. 57 shows Cas9 activation in mESCs.

[0084] FIG. 58 shows a schematic of the strategy used to mediate gene knockout via NHEJ using a nickase version of Cas9 along with two guide RNAs.

[0085] FIG. 59 shows how DNA double-strand break (DSB) repair promotes gene editing. In the error-prone non-homologous end joining (NHEJ) pathway, the ends of a DSB are processed by endogenous DNA repair machineries and rejoined together, which can result in random insertion/deletion (indel) mutations at the site of junction. Indel mutations occurring within the coding region of a gene can result in frame-shift and a premature stop codon, leading to gene knockout. Alternatively, a repair template in the form of a plasmid or single-stranded oligodeoxynucleotides (ssODN) can be supplied to leverage the homology-directed repair (HDR) pathway, which allows high fidelity and precise editing.

[0086] FIG. 60 shows the timeline and overview of experiments. Steps for reagent design, construction, validation, and cell line expansion. Custom sgRNAs (light blue bars) for each target, as well as genotyping primers, are designed in silico via our online design tool (available at the website genome-engineering.org/tools). sgRNA expression vectors are then cloned into a plasmid containing Cas9 (PX330) and verified via DNA sequencing. Completed plasmids (pCRISPRs), and optional repair templates for facilitating homology directed repair, are then transfected into cells and assayed for ability to mediate targeted cleavage. Finally, transfected cells can be clonally expanded to derive isogenic cell lines with defined mutations.

[0087] FIG. 61A-61C shows Target selection and reagent preparation. (a) For *S. pyogenes* Cas9, 20-bp targets (highlighted in blue) must be followed by 5'-NGG, which can occur in either strand on genomic DNA. We recommend using the online tool described in this protocol in aiding target selection (www.genome-engineering.org/tools). (b) Schematic for co-transfection of Cas9 expression plasmid (PX165) and PCR-amplified U6-driven sgRNA expression cassette. Using a U6 promoter-containing PCR template and a fixed forward primer (U6 Fwd), sgRNA-encoding DNA can be appended onto the U6 reverse primer (U6 Rev) and synthesized as an extended DNA oligo (Ultrasmer oligos from IDT). Note the guide sequence (blue N's) in U6 Rev is the reverse complement of the 5'-NGG flanking target sequence (SEQ ID NOS 517 and 517-519, respectively, in

order of appearance). (c) Schematic for scarless cloning of the guide sequence oligos into a plasmid containing Cas9 and sgRNA scaffold (PX330). The guide oligos (blue N's) contain overhangs for ligation into the pair of BbsI sites on PX330, with the top and bottom strand orientations matching those of the genomic target (i.e. top oligo is the 20-bp sequence preceding 5'-NGG in genomic DNA). Digestion of PX330 with BbsI allows the replacement of the Type II restriction sites (blue outline) with direct insertion of annealed oligos. It is worth noting that an extra G was placed before the first base of the guide sequence. Applicants have found that an extra G in front of the guide sequence does not adversely affect targeting efficiency. In cases when the 20-nt guide sequence of choice does not begin with guanine, the extra guanine will ensure the sgRNA is efficiently transcribed by the U6 promoter, which prefers a guanine in the first base of the transcript (SEQ ID NOS 320-321 and 328, respectively, in order of appearance).

[0088] FIG. 62A-62D shows the anticipated results for multiplex NHEJ. (a) Schematic of the SURVEYOR assay used to determine indel percentage. First, genomic DNA from the heterogeneous population of Cas9-targeted cells is amplified by PCR. Amplicons are then reannealed slowly to generate heteroduplexes. The reannealed heteroduplexes are cleaved by SURVEYOR nuclease, whereas homoduplexes are left intact. Cas9-mediated cleavage efficiency (% indel) is calculated based on the fraction of cleaved DNA, as determined by integrated intensity of gel bands. (b) Two sgRNAs (orange and blue bars) are designed to target the human GRIN2B and DYRK1A loci. SURVEYOR gel shows modification at both loci in transfected cells. Colored arrows indicated expected fragment sizes for each locus. (c) A pair of sgRNAs (light blue and green bars) are designed to excise an exon (dark blue) in the human EMX1 locus. Target sequences and PAMs (red) are shown in respective colors, and sites of cleavage indicated by red triangle. Predicted junction is shown below. Individual clones isolated from cell populations transfected with sgRNA 3, 4, or both are assayed by PCR (OUT Fwd, OUT Rev), reflecting a deletion of ~270-bp. Representative clones with no modification (12/23), mono-allelic (10/23), and bi-allelic (1/23) modifications are shown. IN Fwd and IN Rev primers are used to screen for inversion events (FIG. 6d) (SEQ ID NOS 520-522, respectively, in order of appearance). (d) Quantification of clonal lines with EMX1 exon deletions. Two pairs of sgRNAs (3.1, 3.2 left-flanking sgRNAs; 4.1, 4.2, right flanking sgRNAs) are used to mediate deletions of variable sizes around one EMX1 exon. Transfected cells are clonally isolated and expanded for genotyping analysis for deletions and inversion events. Of the 105 clones are screened, 51 (49%) and 11 (10%) carrying heterozygous and homozygous deletions, respectively. Approximate deletion sizes are given since junctions may be variable.

[0089] FIG. 63A-63C shows the application of ssODNs and targeting vector to mediate HR with both wildtype and nickase mutant of Cas9 in HEK293FT and HUES9 cells with efficiencies ranging from 1.0-27%. FIG. 63B discloses SEQ ID NOS 503-505, 503, 506 and 505, respectively, in order of appearance.

[0090] FIG. 64 shows a schematic of a PCR-based method for rapid and efficient CRISPR targeting in mammalian cells. A plasmid containing the human RNA polymerase III promoter U6 is PCR-amplified using a U6-specific forward primer and a reverse primer carrying the reverse comple-

ment of part of the U6 promoter, the sgRNA(+85) scaffold with guide sequence, and 7 T nucleotides for transcriptional termination. The resulting PCR product is purified and co-delivered with a plasmid carrying Cas9 driven by the CBh promoter (SEQ ID NOS 517, 523, 518 and 524-525, respectively, in order of appearance).

[0091] FIG. 65 shows SURVEYOR Mutation Detection Kit from Transgenomics results for each gRNA and respective controls. A positive SURVEYOR result is one large band corresponding to the genomic PCR and two smaller bands that are the product of the SURVEYOR nuclease making a double-strand break at the site of a mutation. Each gRNA was validated in the mouse cell line, Neuro-N2a, by liposomal transient co-transfection with hSpCas9. 72 hours post-transfection genomic DNA was purified using Quick-Extract DNA from Epicentre. PCR was performed to amplify the locus of interest.

[0092] FIG. 66 shows Surveyor results for 38 live pups (lanes 1-38) 1 dead pup (lane 39) and 1 wild-type pup for comparison (lane 40). Pups 1-19 were injected with gRNA Chd8.2 and pups 20-38 were injected with gRNA Chd8.3. Of the 38 live pups, 13 were positive for a mutation. The one dead pup also had a mutation. There was no mutation detected in the wild-type sample. Genomic PCR sequencing was consistent with the SURVEYOR assay findings (SEQ ID NOS 526-528, respectively, in order of appearance).

[0093] FIG. 67 shows a design of different Cas9 NLS constructs. All Cas9 were the human-codon-optimized version of the Sp Cas9. NLS sequences are linked to the cas9 gene at either N-terminus or C-terminus. All Cas9 variants with different NLS designs were cloned into a backbone vector containing so it is driven by EF1a promoter. On the same vector there is a chimeric RNA targeting human EMX1 locus driven by U6 promoter, together forming a two-component system.

[0094] FIG. 68 shows the efficiency of genomic cleavage induced by Cas9 variants bearing different NLS designs. The percentage indicate the portion of human EMX1 genomic DNA that were cleaved by each construct. All experiments are from 3 biological replicates. n=3, error indicates S.E.M.

[0095] FIG. 69A shows a design of the CRISPR-TF (Transcription Factor) with transcriptional activation activity. The chimeric RNA is expressed by U6 promoter, while a human-codon-optimized, double-mutant version of the Cas9 protein (hSpCas9m), operably linked to triple NLS and a VP64 functional domain is expressed by a EF1a promoter. The double mutations, D10A and H840A, renders the cas9 protein unable to introduce any cleavage but maintained its capacity to bind to target DNA when guided by the chimeric RNA.

[0096] FIG. 69B shows transcriptional activation of the human SOX2 gene with CRISPR-TF system (Chimeric RNA and the Cas9-NLS-VP64 fusion protein). 293FT cells were transfected with plasmids bearing two components: (1) U6-driven different chimeric RNAs targeting 20-bp sequences within or around the human SOX2 genomic locus, and (2) EF 1a-driven hSpCas9m (double mutant)-NLS-VP64 fusion protein. 96 hours post transfection, 293FT cells were harvested and the level of activation is measured by the induction of mRNA expression using a qRT-PCR assay. All expression levels are normalized against the control group (grey bar), which represents results from cells transfected with the CRISPR-TF backbone plasmid without chimeric RNA. The qRT-PCR probes used for

detecting the SOX2 mRNA is Taqman Human Gene Expression Assay (Life Technologies). All experiments represents data from 3 biological replicates, n=3, error bars show s.e.m. **[0097]** FIG. 70 depicts NLS architecture optimization for SpCas9.

[0098] FIG. 71 shows a QQ plot for NGGNN sequences. **[0099]** FIG. 72 shows a histogram of the data density with fitted normal distribution (black line) and 0.99 quantile (dotted line).

[0100] FIG. 73A-73C shows RNA-guided repression of bgaA expression by dgRNA:cas9**. a. The Cas9 protein binds to the tracrRNA, and to the precursor CRISPR RNA which is processed by RNaseIII to form the crRNA. The crRNA directs binding of Cas9 to the bgaA promoter and represses transcription. b. The targets used to direct Cas9** to the bgaA promoter are represented (SEQ ID NO: 529). Putative -35, -10 as well as the bgaA start codon are in bold. c. Betagalactosidase activity as measure by Miller assay in the absence of targeting and for the four different targets.

[0101] FIG. 74A-74E shows characterization of Cas9** mediated repression. a. The gflimut2 gene and its promoter, including the -35 and -10 signals are represented together with the position of the different target sites used the study. b. Relative fluorescence upon targeting of the coding strand. c. Relative fluorescence upon targeting of the non-coding strand. d. Northern blot with probes B477 and B478 on RNA extracted from T5, T10, B10 or a control strain without a target. e. Effect of an increased number of mutations in the 5' end of the crRNA of B 1, T5 and B10.

[0102] The figures herein are for illustrative purposes only and are not necessarily drawn to scale.

DETAILED DESCRIPTION OF THE INVENTION

[0103] The terms “polynucleotide”, “nucleotide”, “nucleotide sequence”, “nucleic acid” and “oligonucleotide” are used interchangeably. They refer to a polymeric form of nucleotides of any length, either deoxyribonucleotides or ribonucleotides, or analogs thereof. Polynucleotides may have any three dimensional structure, and may perform any function, known or unknown. The following are non-limiting examples of polynucleotides: coding or non-coding regions of a gene or gene fragment, loci (locus) defined from linkage analysis, exons, introns, messenger RNA (mRNA), transfer RNA, ribosomal RNA, short interfering RNA (siRNA), short-hairpin RNA (shRNA), micro-RNA (miRNA), ribozymes, cDNA, recombinant polynucleotides, branched polynucleotides, plasmids, vectors, isolated DNA of any sequence, isolated RNA of any sequence, nucleic acid probes, and primers. A polynucleotide may comprise one or more modified nucleotides, such as methylated nucleotides and nucleotide analogs. If present, modifications to the nucleotide structure may be imparted before or after assembly of the polymer. The sequence of nucleotides may be interrupted by non-nucleotide components. A polynucleotide may be further modified after polymerization, such as by conjugation with a labeling component.

[0104] In aspects of the invention the terms “chimeric RNA”, “chimeric guide RNA”, “guide RNA”, “single guide RNA” and “synthetic guide RNA” are used interchangeably and refer to the polynucleotide sequence comprising the guide sequence, the tracr sequence and the tracr mate sequence. The term “guide sequence” refers to the about 20 bp sequence within the guide RNA that specifies the target

site and may be used interchangeably with the terms “guide” or “spacer”. The term “tracer sequence” may also be used interchangeably with the term “direct repeat(s)”.

[0105] As used herein the term “wild type” is a term of the art understood by skilled persons and means the typical form of an organism, strain, gene or characteristic as it occurs in nature as distinguished from mutant or variant forms.

[0106] As used herein the term “variant” should be taken to mean the exhibition of qualities that have a pattern that deviates from what occurs in nature.

[0107] The terms “non-naturally occurring” or “engineered” are used interchangeably and indicate the involvement of the hand of man. The terms, when referring to nucleic acid molecules or polypeptides mean that the nucleic acid molecule or the polypeptide is at least substantially free from at least one other component with which they are naturally associated in nature and as found in nature.

[0108] “Complementarity” refers to the ability of a nucleic acid to form hydrogen bond(s) with another nucleic acid sequence by either traditional Watson-Crick base pairing or other non-traditional types. A percent complementarity indicates the percentage of residues in a nucleic acid molecule which can form hydrogen bonds (e.g., Watson-Crick base pairing) with a second nucleic acid sequence (e.g., 5, 6, 7, 8, 9, 10 out of 10 being 50%, 60%, 70%, 80%, 90%, and 100% complementary). “Perfectly complementary” means that all the contiguous residues of a nucleic acid sequence will hydrogen bond with the same number of contiguous residues in a second nucleic acid sequence. “Substantially complementary” as used herein refers to a degree of complementarity that is at least 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%, 98%, 99%, or 100% over a region of 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 30, 35, 40, 45, 50, or more nucleotides, or refers to two nucleic acids that hybridize under stringent conditions.

[0109] As used herein, “stringent conditions” for hybridization refer to conditions under which a nucleic acid having complementarity to a target sequence predominantly hybridizes with the target sequence, and substantially does not hybridize to non-target sequences. Stringent conditions are generally sequence-dependent, and vary depending on a number of factors. In general, the longer the sequence, the higher the temperature at which the sequence specifically hybridizes to its target sequence. Non-limiting examples of stringent conditions are described in detail in Tijssen (1993), *Laboratory Techniques In Biochemistry And Molecular Biology-Hybridization With Nucleic Acid Probes Part I, Second Chapter “Overview of principles of hybridization and the strategy of nucleic acid probe assay”*, Elsevier, N.Y.

[0110] “Hybridization” refers to a reaction in which one or more polynucleotides react to form a complex that is stabilized via hydrogen bonding between the bases of the nucleotide residues. The hydrogen bonding may occur by Watson Crick base pairing, Hoogstein binding, or in any other sequence specific manner. The complex may comprise two strands forming a duplex structure, three or more strands forming a multi stranded complex, a single self hybridizing strand, or any combination of these. A hybridization reaction may constitute a step in a more extensive process, such as the initiation of PCR, or the cleavage of a polynucleotide by an enzyme. A sequence capable of hybridizing with a given sequence is referred to as the “complement” of the given sequence.

[0111] As used herein, “expression” refers to the process by which a polynucleotide is transcribed from a DNA template (such as into mRNA or other RNA transcript) and/or the process by which a transcribed mRNA is subsequently translated into peptides, polypeptides, or proteins. Transcripts and encoded polypeptides may be collectively referred to as “gene product.” If the polynucleotide is derived from genomic DNA, expression may include splicing of the mRNA in a eukaryotic cell.

[0112] The terms “polypeptide”, “peptide” and “protein” are used interchangeably herein to refer to polymers of amino acids of any length. The polymer may be linear or branched, it may comprise modified amino acids, and it may be interrupted by non amino acids. The terms also encompass an amino acid polymer that has been modified; for example, disulfide bond formation, glycosylation, lipidation, acetylation, phosphorylation, or any other manipulation, such as conjugation with a labeling component. As used herein the term “amino acid” includes natural and/or unnatural or synthetic amino acids, including glycine and both the D or L optical isomers, and amino acid analogs and peptidomimetics.

[0113] The terms “subject,” “individual,” and “patient” are used interchangeably herein to refer to a vertebrate, preferably a mammal, more preferably a human. Mammals include, but are not limited to, murines, simians, humans, farm animals, sport animals, and pets. Tissues, cells and their progeny of a biological entity obtained in vivo or cultured in vitro are also encompassed.

[0114] The terms “therapeutic agent”, “therapeutic capable agent” or “treatment agent” are used interchangeably and refer to a molecule or compound that confers some beneficial effect upon administration to a subject. The beneficial effect includes enablement of diagnostic determinations; amelioration of a disease, symptom, disorder, or pathological condition; reducing or preventing the onset of a disease, symptom, disorder or condition; and generally counteracting a disease, symptom, disorder or pathological condition.

[0115] As used herein, “treatment” or “treating,” or “palliating” or “ameliorating” are used interchangeably. These terms refer to an approach for obtaining beneficial or desired results including but not limited to a therapeutic benefit and/or a prophylactic benefit. By therapeutic benefit is meant any therapeutically relevant improvement in or effect on one or more diseases, conditions, or symptoms under treatment. For prophylactic benefit, the compositions may be administered to a subject at risk of developing a particular disease, condition, or symptom, or to a subject reporting one or more of the physiological symptoms of a disease, even though the disease, condition, or symptom may not have yet been manifested.

[0116] The term “effective amount” or “therapeutically effective amount” refers to the amount of an agent that is sufficient to effect beneficial or desired results. The therapeutically effective amount may vary depending upon one or more of: the subject and disease condition being treated, the weight and age of the subject, the severity of the disease condition, the manner of administration and the like, which can readily be determined by one of ordinary skill in the art. The term also applies to a dose that will provide an image for detection by any one of the imaging methods described herein. The specific dose may vary depending on one or more of: the particular agent chosen, the dosing regimen to

be followed, whether it is administered in combination with other compounds, timing of administration, the tissue to be imaged, and the physical delivery system in which it is carried.

[0117] The practice of the present invention employs, unless otherwise indicated, conventional techniques of immunology, biochemistry, chemistry, molecular biology, microbiology, cell biology, genomics and recombinant DNA, which are within the skill of the art. See Sambrook, Fritsch and Maniatis, *MOLECULAR CLONING: A LABORATORY MANUAL*, 2nd edition (1989); *CURRENT PROTOCOLS IN MOLECULAR BIOLOGY* (F. M. Ausubel, et al. eds., (1987)); the series *METHODS IN ENZYMOLOGY* (Academic Press, Inc.): *PCR 2: A PRACTICAL APPROACH* (M. J. MacPherson, B. D. Hames and G. R. Taylor eds. (1995)), Harlow and Lane, eds. (1988) *ANTIBODIES, A LABORATORY MANUAL*, and *ANIMAL CELL CULTURE* (R. I. Freshney, ed. (1987)).

[0118] Several aspects of the invention relate to vector systems comprising one or more vectors, or vectors as such. Vectors can be designed for expression of CRISPR transcripts (e.g. nucleic acid transcripts, proteins, or enzymes) in prokaryotic or eukaryotic cells. For example, CRISPR transcripts can be expressed in bacterial cells such as *Escherichia coli*, insect cells (using baculovirus expression vectors), yeast cells, or mammalian cells. Suitable host cells are discussed further in Goeddel, *GENE EXPRESSION TECHNOLOGY: METHODS IN ENZYMOLOGY 185*, Academic Press, San Diego, Calif. (1990). Alternatively, the recombinant expression vector can be transcribed and translated in vitro, for example using T7 promoter regulatory sequences and T7 polymerase.

[0119] Vectors may be introduced and propagated in a prokaryote. In some embodiments, a prokaryote is used to amplify copies of a vector to be introduced into a eukaryotic cell or as an intermediate vector in the production of a vector to be introduced into a eukaryotic cell (e.g. amplifying a plasmid as part of a viral vector packaging system). In some embodiments, a prokaryote is used to amplify copies of a vector and express one or more nucleic acids, such as to provide a source of one or more proteins for delivery to a host cell or host organism. Expression of proteins in prokaryotes is most often carried out in *Escherichia coli* with vectors containing constitutive or inducible promoters directing the expression of either fusion or non-fusion proteins. Fusion vectors add a number of amino acids to a protein encoded therein, such as to the amino terminus of the recombinant protein. Such fusion vectors may serve one or more purposes, such as: (i) to increase expression of recombinant protein; (ii) to increase the solubility of the recombinant protein; and (iii) to aid in the purification of the recombinant protein by acting as a ligand in affinity purification. Often, in fusion expression vectors, a proteolytic cleavage site is introduced at the junction of the fusion moiety and the recombinant protein to enable separation of the recombinant protein from the fusion moiety subsequent to purification of the fusion protein. Such enzymes, and their cognate recognition sequences, include Factor Xa, thrombin and enterokinase. Example fusion expression vectors include pGEX (Pharmacia Biotech Inc; Smith and Johnson, 1988. *Gene* 67: 31-40), pMAL (New England Biolabs, Beverly, Mass.) and pRIT5 (Pharmacia, Piscataway, N.J.)

that fuse glutathione S-transferase (GST), maltose E binding protein, or protein A, respectively, to the target recombinant protein.

[0120] Examples of suitable inducible non-fusion *E. coli* expression vectors include pTrc (Amrann et al., (1988) *Gene* 69:301-315) and pET 11d (Studier et al., *GENE EXPRESSION TECHNOLOGY: METHODS IN ENZYMOLOGY 185*, Academic Press, San Diego, Calif. (1990) 60-89).

[0121] In some embodiments, a vector is a yeast expression vector. Examples of vectors for expression in yeast *Saccharomyces cerevisiae* include pYepSec1 (Baldari, et al., 1987. *EMBO J.* 6: 229-234), pMFa (Kuijan and Herskowitz, 1982. *Cell* 30: 933-943), pJRY88 (Schultz et al., 1987. *Gene* 54: 113-123), pYES2 (Invitrogen Corporation, San Diego, Calif.), and picZ (Invitrogen Corp, San Diego, Calif.).

[0122] In some embodiments, a vector drives protein expression in insect cells using baculovirus expression vectors. Baculovirus vectors available for expression of proteins in cultured insect cells (e.g., SF9 cells) include the pAc series (Smith, et al., 1983. *Mol. Cell. Biol.* 3: 2156-2165) and the pVL series (Lucklow and Summers, 1989. *Virology* 170: 31-39).

[0123] In some embodiments, a vector is capable of driving expression of one or more sequences in mammalian cells using a mammalian expression vector. Examples of mammalian expression vectors include pCDM8 (Seed, 1987. *Nature* 329: 840) and pMT2PC (Kaufman, et al., 1987. *EMBO J.* 6: 187-195). When used in mammalian cells, the expression vector's control functions are typically provided by one or more regulatory elements. For example, commonly used promoters are derived from polyoma, adenovirus 2, cytomegalovirus, simian virus 40, and others disclosed herein and known in the art. For other suitable expression systems for both prokaryotic and eukaryotic cells see, e.g., Chapters 16 and 17 of Sambrook, et al., *MOLECULAR CLONING: A LABORATORY MANUAL*, 2nd ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989.

[0124] In some embodiments, the recombinant mammalian expression vector is capable of directing expression of the nucleic acid preferentially in a particular cell type (e.g., tissue-specific regulatory elements are used to express the nucleic acid). Tissue-specific regulatory elements are known in the art. Non-limiting examples of suitable tissue-specific promoters include the albumin promoter (liver-specific; Pinkert, et al., 1987. *Genes Dev.* 1: 268-277), lymphoid-specific promoters (Calame and Eaton, 1988. *Adv. Immunol.* 43: 235-275), in particular promoters of T cell receptors (Winoto and Baltimore, 1989. *EMBO J.* 8: 729-733) and immunoglobulins (Baneiji, et al., 1983. *Cell* 33: 729-740; Queen and Baltimore, 1983. *Cell* 33: 741-748), neuron-specific promoters (e.g., the neurofilament promoter; Byrne and Ruddle, 1989. *Proc. Natl. Acad. Sci. USA* 86: 5473-5477), pancreas-specific promoters (Edlund, et al., 1985. *Science* 230: 912-916), and mammary gland-specific promoters (e.g., milk whey promoter; U.S. Pat. No. 4,873,316 and European Application Publication No. 264,166). Developmentally-regulated promoters are also encompassed, e.g., the murine hox promoters (Kessel and Gruss, 1990. *Science* 249: 374-379) and the α -fetoprotein promoter (Campes and Tilghman, 1989. *Genes Dev.* 3: 537-546).

[0125] In some embodiments, a regulatory element is operably linked to one or more elements of a CRISPR system so as to drive expression of the one or more elements

of the CRISPR system. In general, CRISPRs (Clustered Regularly Interspaced Short Palindromic Repeats), also known as SPIDRs (Spacer Interspersed Direct Repeats), constitute a family of DNA loci that are usually specific to a particular bacterial species. The CRISPR locus comprises a distinct class of interspersed short sequence repeats (SSRs) that were recognized in *E. coli* (Ishino et al., *J. Bacteriol.*, 169:5429-5433 [1987]; and Nakata et al., *J. Bacteriol.*, 171:3553-3556 [1989]), and associated genes. Similar interspersed SSRs have been identified in *Haloferax mediterranei*, *Streptococcus pyogenes*, *Anabaena*, and *Mycobacterium tuberculosis* (See, Groenen et al., *Mol. Microbiol.*, 10:1057-1065 [1993]; Hoe et al., *Emerg. Infect. Dis.*, 5:254-263 [1999]; Masepohl et al., *Biochim. Biophys. Acta* 1307: 26-30 [1996]; and Mojica et al., *Mol. Microbiol.*, 17:85-93 [1995]). The CRISPR loci typically differ from other SSRs by the structure of the repeats, which have been termed short regularly spaced repeats (SRSRs) (Janssen et al., *OMICS J. Integ. Biol.*, 6:23-33 [2002]; and Mojica et al., *Mol. Microbiol.*, 36:244-246 [2000]). In general, the repeats are short elements that occur in clusters that are regularly spaced by unique intervening sequences with a substantially constant length (Mojica et al., [2000], supra). Although the repeat sequences are highly conserved between strains, the number of interspersed repeats and the sequences of the spacer regions typically differ from strain to strain (van Embden et al., *J. Bacteriol.*, 182:2393-2401 [2000]). CRISPR loci have been identified in more than 40 prokaryotes (See e.g., Jansen et al., *Mol. Microbiol.*, 43:1565-1575 [2002]; and Mojica et al., [2005]) including, but not limited to *Aeropyrum*, *Pyrobaculum*, *Sulfolobus*, *Archaeoglobus*, *Halocarcularia*, *Methanobacterium*, *Methanococcus*, *Methanosarcina*, *Methanopyrus*, *Pyrococcus*, *Picrophilus*, *Thermoplasma*, *Corynebacterium*, *Mycobacterium*, *Streptomyces*, *Aquifex*, *Porphyromonas*, *Chlorobium*, *Thermus*, *Bacillus*, *Listeria*, *Staphylococcus*, *Clostridium*, *Thermoanaerobacter*, *Mycoplasma*, *Fusobacterium*, *Azarcus*, *Chromobacterium*, *Neisseria*, *Nitrosomonas*, *Desulfovibrio*, *Geobacter*, *Mycococcus*, *Campylobacter*, *Wolinella*, *Acinetobacter*, *Envinia*, *Escherichia*, *Legionella*, *Methylococcus*, *Pasteurella*, *Photobacterium*, *Salmonella*, *Xanthomonas*, *Yersinia*, *Treponema*, and *Thermotoga*.

[0126] In general, “CRISPR system” refers collectively to transcripts and other elements involved in the expression of or directing the activity of CRISPR-associated (“Cas”) genes, including sequences encoding a Cas gene, a tracr (trans-activating CRISPR) sequence (e.g. tracrRNA or an active partial tracrRNA), a tracr-mate sequence (encompassing a “direct repeat” and a tracrRNA-processed partial direct repeat in the context of an endogenous CRISPR system), a guide sequence (also referred to as a “spacer” in the context of an endogenous CRISPR system), or other sequences and transcripts from a CRISPR locus. In some embodiments, one or more elements of a CRISPR system is derived from a type I, type II, or type III CRISPR system. In some embodiments, one or more elements of a CRISPR system is derived from a particular organism comprising an endogenous CRISPR system, such as *Streptococcus pyogenes*. In general, a CRISPR system is characterized by elements that promote the formation of a CRISPR complex at the site of a target sequence (also referred to as a protospacer in the context of an endogenous CRISPR system). In the context of formation of a CRISPR complex, “target sequence” refers to a sequence to which a guide sequence is designed to have

complementarity, where hybridization between a target sequence and a guide sequence promotes the formation of a CRISPR complex. Full complementarity is not necessarily required, provided there is sufficient complementarity to cause hybridization and promote formation of a CRISPR complex. A target sequence may comprise any polynucleotide, such as DNA or RNA polynucleotides. In some embodiments, a target sequence is located in the nucleus or cytoplasm of a cell. In some embodiments, the target sequence may be within an organelle of a eukaryotic cell, for example, mitochondrion or chloroplast. A sequence or template that may be used for recombination into the targeted locus comprising the target sequences is referred to as an “editing template” or “editing polynucleotide” or “editing sequence”. In aspects of the invention, an exogenous template polynucleotide may be referred to as an editing template. In an aspect of the invention the recombination is homologous recombination.

[0127] Typically, in the context of an endogenous CRISPR system, formation of a CRISPR complex (comprising a guide sequence hybridized to a target sequence and complexed with one or more Cas proteins) results in cleavage of one or both strands in or near (e.g. within 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 50, or more base pairs from) the target sequence. Without wishing to be bound by theory, the tracr sequence, which may comprise or consist of all or a portion of a wild-type tracr sequence (e.g. about or more than about 20, 26, 32, 45, 48, 54, 63, 67, 85, or more nucleotides of a wild-type tracr sequence), may also form part of a CRISPR complex, such as by hybridization along at least a portion of the tracr sequence to all or a portion of a tracr mate sequence that is operably linked to the guide sequence. In some embodiments, the tracr sequence has sufficient complementarity to a tracr mate sequence to hybridize and participate in formation of a CRISPR complex. As with the target sequence, it is believed that complete complementarity is not needed, provided there is sufficient to be functional. In some embodiments, the tracr sequence has at least 50%, 60%, 70%, 80%, 90%, 95% or 99% of sequence complementarity along the length of the tracr mate sequence when optimally aligned. In some embodiments, one or more vectors driving expression of one or more elements of a CRISPR system are introduced into a host cell such that expression of the elements of the CRISPR system direct formation of a CRISPR complex at one or more target sites. For example, a Cas enzyme, a guide sequence linked to a tracr-mate sequence, and a tracr sequence could each be operably linked to separate regulatory elements on separate vectors. Alternatively, two or more of the elements expressed from the same or different regulatory elements, may be combined in a single vector, with one or more additional vectors providing any components of the CRISPR system not included in the first vector. CRISPR system elements that are combined in a single vector may be arranged in any suitable orientation, such as one element located 5' with respect to (“upstream” of) or 3' with respect to (“downstream” of) a second element. The coding sequence of one element may be located on the same or opposite strand of the coding sequence of a second element, and oriented in the same or opposite direction. In some embodiments, a single promoter drives expression of a transcript encoding a CRISPR enzyme and one or more of the guide sequence, tracr mate sequence (optionally operably linked to the guide sequence), and a tracr sequence

embedded within one or more intron sequences (e.g. each in a different intron, two or more in at least one intron, or all in a single intron). In some embodiments, the CRISPR enzyme, guide sequence, tracr mate sequence, and tracr sequence are operably linked to and expressed from the same promoter.

[0128] In some embodiments, a vector comprises one or more insertion sites, such as a restriction endonuclease recognition sequence (also referred to as a “cloning site”). In some embodiments, one or more insertion sites (e.g. about or more than about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more insertion sites) are located upstream and/or downstream of one or more sequence elements of one or more vectors. In some embodiments, a vector comprises an insertion site upstream of a tracr mate sequence, and optionally downstream of a regulatory element operably linked to the tracr mate sequence, such that following insertion of a guide sequence into the insertion site and upon expression the guide sequence directs sequence-specific binding of a CRISPR complex to a target sequence in a eukaryotic cell. In some embodiments, a vector comprises two or more insertion sites, each insertion site being located between two tracr mate sequences so as to allow insertion of a guide sequence at each site. In such an arrangement, the two or more guide sequences may comprise two or more copies of a single guide sequence, two or more different guide sequences, or combinations of these. When multiple different guide sequences are used, a single expression construct may be used to target CRISPR activity to multiple different, corresponding target sequences within a cell. For example, a single vector may comprise about or more than about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, or more guide sequences. In some embodiments, about or more than about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more such guide-sequence-containing vectors may be provided, and optionally delivered to a cell.

[0129] In some embodiments, a vector comprises a regulatory element operably linked to an enzyme-coding sequence encoding a CRISPR enzyme, such as a Cas protein. Non-limiting examples of Cas proteins include Cas1, Cas1B, Cas2, Cas3, Cas4, Cas5, Cas6, Cas7, Cas8, Cas9 (also known as Csn1 and Csx12), Cas10, Csy1, Csy2, Csy3, Cse1, Cse2, Csc1, Csc2, Csa5, Csn2, Csm2, Csm3, Csm4, Csm5, Csm6, Cmr1, Cmr3, Cmr4, Cmr5, Cmr6, Csb1, Csb2, Csb3, Csx17, Csx14, Csx10, Csx16, CsaX, Csx3, Csx1, Csx15, Csf1, Csf2, Csf3, Csf4, homologs thereof, or modified versions thereof. These enzymes are known; for example, the amino acid sequence of *S. pyogenes* Cas9 protein may be found in the SwissProt database under accession number Q99ZW2. In some embodiments, the unmodified CRISPR enzyme has DNA cleavage activity, such as Cas9. In some embodiments the CRISPR enzyme is Cas9, and may be Cas9 from *S. pyogenes* or *S. pneumoniae*. In some embodiments, the CRISPR enzyme directs cleavage of one or both strands at the location of a target sequence, such as within the target sequence and/or within the complement of the target sequence. In some embodiments, the CRISPR enzyme directs cleavage of one or both strands within about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 50, 100, 200, 500, or more base pairs from the first or last nucleotide of a target sequence. In some embodiments, a vector encodes a CRISPR enzyme that is mutated to with respect to a corresponding wild-type enzyme such that the mutated CRISPR enzyme lacks the ability to cleave one or both strands of a target polynucleotide containing a target

sequence. For example, an aspartate-to-alanine substitution (D10A) in the RuvC I catalytic domain of Cas9 from *S. pyogenes* converts Cas9 from a nuclease that cleaves both strands to a nickase (cleaves a single strand). Other examples of mutations that render Cas9 a nickase include, without limitation, H840A, N854A, and N863A. In some embodiments, a Cas9 nickase may be used in combination with guide sequence(s), e.g., two guide sequences, which target respectively sense and antisense strands of the DNA target. This combination allows both strands to be nicked and used to induce NHEJ. Applicants have demonstrated (data not shown) the efficacy of two nickase targets (i.e., sgRNAs targeted at the same location but to different strands of DNA) in inducing mutagenic NHEJ. A single nickase (Cas9-D10A with a single sgRNA) is unable to induce NHEJ and create indels but Applicants have shown that double nickase (Cas9-D10A and two sgRNAs targeted to different strands at the same location) can do so in human embryonic stem cells (hESCs). The efficiency is about 50% of nuclease (i.e., regular Cas9 without D10 mutation) in hESCs.

[0130] As a further example, two or more catalytic domains of Cas9 (RuvC I, RuvC II, and RuvC III) may be mutated to produce a mutated Cas9 substantially lacking all DNA cleavage activity. In some embodiments, a D10A mutation is combined with one or more of H840A, N854A, or N863A mutations to produce a Cas9 enzyme substantially lacking all DNA cleavage activity. In some embodiments, a CRISPR enzyme is considered to substantially lack all DNA cleavage activity when the DNA cleavage activity of the mutated enzyme is less than about 25%, 10%, 5%, 1%, 0.1%, 0.01%, or lower with respect to its non-mutated form. Other mutations may be useful; where the Cas9 or other CRISPR enzyme is from a species other than *S. pyogenes*, mutations in corresponding amino acids may be made to achieve similar effects.

[0131] In some embodiments, an enzyme coding sequence encoding a CRISPR enzyme is codon optimized for expression in particular cells, such as eukaryotic cells. The eukaryotic cells may be those of or derived from a particular organism, such as a mammal, including but not limited to human, mouse, rat, rabbit, dog, or non-human primate. In general, codon optimization refers to a process of modifying a nucleic acid sequence for enhanced expression in the host cells of interest by replacing at least one codon (e.g. about or more than about 1, 2, 3, 4, 5, 10, 15, 20, 25, 50, or more codons) of the native sequence with codons that are more frequently or most frequently used in the genes of that host cell while maintaining the native amino acid sequence. Various species exhibit particular bias for certain codons of a particular amino acid. Codon bias (differences in codon usage between organisms) often correlates with the efficiency of translation of messenger RNA (mRNA), which is in turn believed to be dependent on, among other things, the properties of the codons being translated and the availability of particular transfer RNA (tRNA) molecules. The predominance of selected tRNAs in a cell is generally a reflection of the codons used most frequently in peptide synthesis. Accordingly, genes can be tailored for optimal gene expression in a given organism based on codon optimization. Codon usage tables are readily available, for example, at the “Codon Usage Database”, and these tables can be adapted in a number of ways. See Nakamura, Y., et al. “Codon usage tabulated from the international DNA sequence databases:

status for the year 2000” Nucl. Acids Res. 28:292 (2000). Computer algorithms for codon optimizing a particular sequence for expression in a particular host cell are also available, such as Gene Forge (Aptagen; Jacobus, Pa.), are also available. In some embodiments, one or more codons (e.g. 1, 2, 3, 4, 5, 10, 15, 20, 25, 50, or more, or all codons) in a sequence encoding a CRISPR enzyme correspond to the most frequently used codon for a particular amino acid.

[0132] In some embodiments, a vector encodes a CRISPR enzyme comprising one or more nuclear localization sequences (NLSs), such as about or more than about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more NLSs. In some embodiments, the CRISPR enzyme comprises about or more than about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more NLSs at or near the amino-terminus, about or more than about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more NLSs at or near the carboxy-terminus, or a combination of these (e.g. one or more NLS at the amino-terminus and one or more NLS at the carboxy terminus). When more than one NLS is present, each may be selected independently of the others, such that a single NLS may be present in more than one copy and/or in combination with one or more other NLSs present in one or more copies. In a preferred embodiment of the invention, the CRISPR enzyme comprises at most 6 NLSs. In some embodiments, an NLS is considered near the N- or C-terminus when the nearest amino acid of the NLS is within about 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 40, 50, or more amino acids along the polypeptide chain from the N- or C-terminus. Typically, an NLS consists of one or more short sequences of positively charged lysines or arginines exposed on the protein surface, but other types of NLS are known. Non-limiting examples of NLSs include an NLS sequence derived from: the NLS of the SV40 virus large T-antigen, having the amino acid sequence PKKKRKY (SEQ ID NO: 1); the NLS from nucleoplasmin (e.g. the nucleoplasmin bipartite NLS with the sequence KRPAATKKAGQAKKKK (SEQ ID NO: 2)); the c-myc NLS having the amino acid sequence PAAKRVKLD (SEQ ID NO: 3) or RQQRNELKRSP (SEQ ID NO: 4); the hRNPA1 M9 NLS having the sequence NQSSNFGPMKGGNFGGRSSGPYGGGGQYFAKPRN-QGGY (SEQ ID NO: 5); the sequence RMRIZFKNKGKD-TAELRRRRVEVSVELRKAKKDEQILKRRNV (SEQ ID NO: 6) of the IBB domain from importin-alpha; the sequences VSRKRPRP (SEQ ID NO: 7) and PPKKARED (SEQ ID NO: 8) of the myoma T protein; the sequence PPKKKKPL (SEQ ID NO: 9) of human p53; the sequence SALIKKKKKMAP (SEQ ID NO: 10) of mouse c-abl IV; the sequences DRLRR (SEQ ID NO: 11) and PKQKKRK (SEQ ID NO: 12) of the influenza virus NS1; the sequence RKLKKKIKKL (SEQ ID NO: 13) of the Hepatitis virus delta antigen; the sequence REKKKFLKRR (SEQ ID NO: 14) of the mouse Mx1 protein; the sequence KRK-GDEVGDVDEVAKKSKK (SEQ ID NO: 15) of the human poly(ADP-ribose) polymerase; and the sequence RKCLQAGMNLARKTKK (SEQ ID NO: 16) of the steroid hormone receptors (human) glucocorticoid.

[0133] In general, the one or more NLSs are of sufficient strength to drive accumulation of the CRISPR enzyme in a detectable amount in the nucleus of a eukaryotic cell. In general, strength of nuclear localization activity may derive from the number of NLSs in the CRISPR enzyme, the particular NLS(s) used, or a combination of these factors. Detection of accumulation in the nucleus may be performed by any suitable technique. For example, a detectable marker

may be fused to the CRISPR enzyme, such that location within a cell may be visualized, such as in combination with a means for detecting the location of the nucleus (e.g. a stain specific for the nucleus such as DAPI). Examples of detectable markers include fluorescent proteins (such as Green fluorescent proteins, or GFP; RFP; CFP), and epitope tags (HA tag, flag tag, SNAP tag). Cell nuclei may also be isolated from cells, the contents of which may then be analyzed by any suitable process for detecting protein, such as immunohistochemistry, Western blot, or enzyme activity assay. Accumulation in the nucleus may also be determined indirectly, such as by an assay for the effect of CRISPR complex formation (e.g. assay for DNA cleavage or mutation at the target sequence, or assay for altered gene expression activity affected by CRISPR complex formation and/or CRISPR enzyme activity), as compared to a control not exposed to the CRISPR enzyme or complex, or exposed to a CRISPR enzyme lacking the one or more NLSs.

[0134] In general, a guide sequence is any polynucleotide sequence having sufficient complementarity with a target polynucleotide sequence to hybridize with the target sequence and direct sequence-specific binding of a CRISPR complex to the target sequence. In some embodiments, the degree of complementarity between a guide sequence and its corresponding target sequence, when optimally aligned using a suitable alignment algorithm, is about or more than about 50%, 60%, 75%, 80%, 85%, 90%, 95%, 97.5%, 99%, or more. Optimal alignment may be determined with the use of any suitable algorithm for aligning sequences, non-limiting example of which include the Smith-Waterman algorithm, the Needleman-Wunsch algorithm, algorithms based on the Burrows-Wheeler Transform (e.g. the Burrows Wheeler Aligner), ClustalW, Clustal X, BLAT, Novoalign (Novocraft Technologies, ELAND (Illumina, San Diego, Calif.), SOAP (available at soap.genomics.org.cn), and Maq (available at maq.sourceforge.net). In some embodiments, a guide sequence is about or more than about 5, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 35, 40, 45, 50, 75, or more nucleotides in length. In some embodiments, a guide sequence is less than about 75, 50, 45, 40, 35, 30, 25, 20, 15, 12, or fewer nucleotides in length. The ability of a guide sequence to direct sequence-specific binding of a CRISPR complex to a target sequence may be assessed by any suitable assay. For example, the components of a CRISPR system sufficient to form a CRISPR complex, including the guide sequence to be tested, may be provided to a host cell having the corresponding target sequence, such as by transfection with vectors encoding the components of the CRISPR sequence, followed by an assessment of preferential cleavage within the target sequence, such as by Surveyor assay as described herein. Similarly, cleavage of a target polynucleotide sequence may be evaluated in a test tube by providing the target sequence, components of a CRISPR complex, including the guide sequence to be tested and a control guide sequence different from the test guide sequence, and comparing binding or rate of cleavage at the target sequence between the test and control guide sequence reactions. Other assays are possible, and will occur to those skilled in the art.

[0135] A guide sequence may be selected to target any target sequence. In some embodiments, the target sequence is a sequence within a genome of a cell. Exemplary target sequences include those that are unique in the target genome. For example, for the *S. pyogenes* Cas9, a unique

target sequence in a genome may include a Cas9 target site of the form MMMMMMMNNNNNNNNNNNNXGG (SEQ ID NO: 530) where NNNNNNNNNNNXGG (SEQ ID NO: 531) (N is A, G, T, or C; and X can be anything) has a single occurrence in the genome. A unique target sequence in a genome may include an *S. pyogenes* Cas9 target site of the form MMMMMMMNNNNNNNNNNNNXXG (SEQ ID NO: 532) where NNNNNNNNNNNXGG (SEQ ID NO: 533) (N is A, G, T, or C; and X can be anything) has a single occurrence in the genome. For the *S. thermophilus* CRISPR1 Cas9, a unique target sequence in a genome may include a Cas9 target site of the form MMMMMMM-MNNNNNNNNNNNNXXAGAAW (SEQ ID NO: 17) where NNNNNNNNNNNXXAGAAW (SEQ ID NO: 18) (N is A, G, T, or C; X can be anything; and W is A or T) has a single occurrence in the genome. A unique target sequence in a genome may include an *S. thermophilus* CRISPR1 Cas9 target site of the form MMMMMMM-MNNNNNNNNNNNNXXAGAAW (SEQ ID NO: 19) where NNNNNNNNNNNXXAGAAW (SEQ ID NO: 20) (N is A, G, T, or C; X can be anything; and W is A or T) has a single occurrence in the genome. For the *S. pyogenes* Cas9, a unique target sequence in a genome may include a Cas9 target site of the form MMMMMMM-MNNNNNNNNNNNNXXGGXG (SEQ ID NO: 534) where NNNNNNNNNNNXXGGXG (SEQ ID NO: 535) (N is A, G, T, or C; and X can be anything) has a single occurrence in the genome. A unique target sequence in a genome may include an *S. pyogenes* Cas9 target site of the form MMMMMMMNNNNNNNNNNNNXXGGXG (SEQ ID No.: 536) where NNNNNNNNNNNXXGGXG (SEQ ID NO: 537) (N is A, G, T, or C; and X can be anything) has a single occurrence in the genome. In each of these sequences "M" may be A, G, T, or C, and need not be considered in identifying a sequence as unique.

[0136] In some embodiments, a guide sequence is selected to reduce the degree of secondary structure within the guide sequence. Secondary structure may be determined by any suitable polynucleotide folding algorithm. Some programs are based on calculating the minimal Gibbs free energy. An example of one such algorithm is mFold, as described by Zuker and Stiegler (Nucleic Acids Res. 9 (1981), 133-148). Another example folding algorithm is the online webserver RNAfold, developed at Institute for Theoretical Chemistry at the University of Vienna, using the centroid structure prediction algorithm (see e.g. A. R. Gruber et al., 2008, *Cell* 106(1): 23-24; and P A Carr and G M Church, 2009, *Nature Biotechnology* 27(12): 1151-62). Further algorithms may be found in U.S. application Serial No. TBA (attorney docket 44790.11.2022; Broad Reference BI-2013/004A); incorporated herein by reference.

[0137] In general, a tracr mate sequence includes any sequence that has sufficient complementarity with a tracr sequence to promote one or more of: (1) excision of a guide sequence flanked by tracr mate sequences in a cell containing the corresponding tracr sequence; and (2) formation of a CRISPR complex at a target sequence, wherein the CRISPR complex comprises the tracr mate sequence hybridized to the tracr sequence. In general, degree of complementarity is with reference to the optimal alignment of the tracr mate sequence and tracr sequence, along the length of the shorter of the two sequences. Optimal alignment may be determined by any suitable alignment algorithm, and may further account for secondary structures, such as self-

complementarity within either the tracr sequence or tracr mate sequence. In some embodiments, the degree of complementarity between the tracr sequence and tracr mate sequence along the length of the shorter of the two when optimally aligned is about or more than about 25%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, 97.5%, 99%, or higher. Example illustrations of optimal alignment between a tracr sequence and a tracr mate sequence are provided in FIGS. 12B and 13B. In some embodiments, the tracr sequence is about or more than about 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 40, 50, or more nucleotides in length. In some embodiments, the tracr sequence and tracr mate sequence are contained within a single transcript, such that hybridization between the two produces a transcript having a secondary structure, such as a hairpin. Preferred loop forming sequences for use in hairpin structures are four nucleotides in length, and most preferably have the sequence GAAA. However, longer or shorter loop sequences may be used, as may alternative sequences. The sequences preferably include a nucleotide triplet (for example, AAA), and an additional nucleotide (for example C or G). Examples of loop forming sequences include CAAA and AAAG. In an embodiment of the invention, the transcript or transcribed polynucleotide sequence has at least two or more hairpins. In preferred embodiments, the transcript has two, three, four or five hairpins. In a further embodiment of the invention, the transcript has at most five hairpins. In some embodiments, the single transcript further includes a transcription termination sequence; preferably this is a polyT sequence, for example six T nucleotides. An example illustration of such a hairpin structure is provided in the lower portion of FIG. 13B, where the portion of the sequence 5' of the final "N" and upstream of the loop corresponds to the tracr mate sequence, and the portion of the sequence 3' of the loop corresponds to the tracr sequence. Further non-limiting examples of single polynucleotides comprising a guide sequence, a tracr mate sequence, and a tracr sequence are as follows (listed 5' to 3'), where "N" represents a base of a guide sequence, the first block of lower case letters represent the tracr mate sequence, and the second block of lower case letters represent the tracr sequence, and the final poly-T sequence represents the transcription terminator: (1) NNNNNNNNNNNNNNNNNNNNNgTTTTgtactctcaagatttGAAAtaatcttcgagaagctacaagataaggctt catgcgaaatcaacacctgtcattttatggcagggtgttttcgtatttaaTTTTTT (SEQ ID NO: 21); (2) NNNNNNNNNNNNNNNNNNNNNgTTTTgtactctcaGAAAtgcagaagctacaagataaggcttcctgcgaaatca acacctgtcattttatggcagggtgttttcgtatttaaTTTTTT (SEQ ID NO: 22); (3) NNNNNNNNNNNNNNNNNNNNNgTTTTgtactctcaGAAAtgcagaagctacaagataaggcttcctgcgaaatca acacctgtcattttatggcagggtgtTTTTTT (SEQ ID NO: 23); (4) NNNNNNNNNNNNNNNNNNNNNgTTTTtagagctaGAAAtagcaagttaaaataaggctagtcctgtatcaacttgaaa agtgccaccgagtcgtgcTTTTTT (SEQ ID NO: 24); (5) NNNNNNNNNNNNNNNNNNNNNgTTTTtagagctaGAAATAGcaagttaaaataaggctagtcctgtatcaacttgaa aaagt-gTTTTTTT (SEQ ID NO: 25); and (6) NNNNNNNNNNNNNNNNNNNNNgTTTTtagagctaGAAATAGcaagttaaaataaggctagtcctgtatcaTTTT TTT (SEQ ID NO: 26). In some embodiments, sequences (1) to (3) are used in combination with Cas9 from *S. thermophilus* CRISPR1. In some embodiments, sequences (4) to (6) are used in combination with Cas9 from *S. pyogenes*. In some

embodiments, the tracr sequence is a separate transcript from a transcript comprising the tracr mate sequence (such as illustrated in the top portion of FIG. 13B).

[0138] In some embodiments, a recombination template is also provided. A recombination template may be a component of another vector as described herein, contained in a separate vector, or provided as a separate polynucleotide. In some embodiments, a recombination template is designed to serve as a template in homologous recombination, such as within or near a target sequence nicked or cleaved by a CRISPR enzyme as a part of a CRISPR complex. A template polynucleotide may be of any suitable length, such as about or more than about 10, 15, 20, 25, 50, 75, 100, 150, 200, 500, 1000, or more nucleotides in length. In some embodiments, the template polynucleotide is complementary to a portion of a polynucleotide comprising the target sequence. When optimally aligned, a template polynucleotide might overlap with one or more nucleotides of a target sequences (e.g. about or more than about 1, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, 100 or more nucleotides). In some embodiments, when a template sequence and a polynucleotide comprising a target sequence are optimally aligned, the nearest nucleotide of the template polynucleotide is within about 1, 5, 10, 15, 20, 25, 50, 75, 100, 200, 300, 400, 500, 1000, 5000, 10000, or more nucleotides from the target sequence.

[0139] In some embodiments, the CRISPR enzyme is part of a fusion protein comprising one or more heterologous protein domains (e.g. about or more than about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more domains in addition to the CRISPR enzyme). A CRISPR enzyme fusion protein may comprise any additional protein sequence, and optionally a linker sequence between any two domains. Examples of protein domains that may be fused to a CRISPR enzyme include, without limitation, epitope tags, reporter gene sequences, and protein domains having one or more of the following activities: methylase activity, demethylase activity, transcription activation activity, transcription repression activity, transcription release factor activity, histone modification activity, RNA cleavage activity and nucleic acid binding activity. Non-limiting examples of epitope tags include histidine (His) tags, V5 tags, FLAG tags, influenza hemagglutinin (HA) tags, Myc tags, VSV-G tags, and thioredoxin (Trx) tags. Examples of reporter genes include, but are not limited to, glutathione-S-transferase (GST), horseradish peroxidase (HRP), chloramphenicol acetyltransferase (CAT) beta-galactosidase, beta-glucuronidase, luciferase, green fluorescent protein (GFP), HcRed, DsRed, cyan fluorescent protein (CFP), yellow fluorescent protein (YFP), and auto-fluorescent proteins including blue fluorescent protein (BFP). A CRISPR enzyme may be fused to a gene sequence encoding a protein or a fragment of a protein that bind DNA molecules or bind other cellular molecules, including but not limited to maltose binding protein (MBP), S-tag, Lex A DNA binding domain (DBD) fusions, GAL4 DNA binding domain fusions, and herpes simplex virus (HSV) BP16 protein fusions. Additional domains that may form part of a fusion protein comprising a CRISPR enzyme are described in US20110059502, incorporated herein by reference. In some embodiments, a tagged CRISPR enzyme is used to identify the location of a target sequence.

[0140] In some aspects, the invention provides methods comprising delivering one or more polynucleotides, such as or one or more vectors as described herein, one or more

transcripts thereof, and/or one or proteins transcribed therefrom, to a host cell. In some aspects, the invention further provides cells produced by such methods, and organisms (such as animals, plants, or fungi) comprising or produced from such cells. In some embodiments, a CRISPR enzyme in combination with (and optionally complexed with) a guide sequence is delivered to a cell. Conventional viral and non-viral based gene transfer methods can be used to introduce nucleic acids in mammalian cells or target tissues. Such methods can be used to administer nucleic acids encoding components of a CRISPR system to cells in culture, or in a host organism. Non-viral vector delivery systems include DNA plasmids, RNA (e.g. a transcript of a vector described herein), naked nucleic acid, and nucleic acid complexed with a delivery vehicle, such as a liposome. Viral vector delivery systems include DNA and RNA viruses, which have either episomal or integrated genomes after delivery to the cell. For a review of gene therapy procedures, see Anderson, *Science* 256:808-813 (1992); Nabel & Felgner, *TIBTECH* 11:211-217 (1993); Mitani & Caskey, *TIBTECH* 11:162-166 (1993); Dillon, *TIBTECH* 11:167-175 (1993); Miller, *Nature* 357:455-460 (1992); Van Brunt, *Biotechnology* 6(10):1149-1154 (1988); Vigne, *Restorative Neurology and Neuroscience* 8:35-36 (1995); Kremer & Perricaudet, *British Medical Bulletin* 51(1):31-44 (1995); Haddada et al., in *Current Topics in Microbiology and Immunology*, Doerfler and Bohm (eds) (1995); and Yu et al., *Gene Therapy* 1:13-26 (1994).

[0141] Methods of non-viral delivery of nucleic acids include lipofection, nucleofection, microinjection, biolistics, virosomes, liposomes, immunoliposomes, polycation or lipid:nucleic acid conjugates, naked DNA, artificial virions, and agent-enhanced uptake of DNA. Lipofection is described in e.g., U.S. Pat. Nos. 5,049,386, 4,946,787; and 4,897,355) and lipofection reagents are sold commercially (e.g., Transfectam™ and Lipofectin™). Cationic and neutral lipids that are suitable for efficient receptor-recognition lipofection of polynucleotides include those of Felgner, WO 91/17424; WO 91/16024. Delivery can be to cells (e.g. in vitro or ex vivo administration) or target tissues (e.g. in vivo administration).

[0142] The preparation of lipid:nucleic acid complexes, including targeted liposomes such as immunolipid complexes, is well known to one of skill in the art (see, e.g., *Crystal*, *Science* 270:404-410 (1995); Blaese et al., *Cancer Gene Ther.* 2:291-297 (1995); Behr et al., *Bioconjugate Chem.* 5:382-389 (1994); Remy et al., *Bioconjugate Chem.* 5:647-654 (1994); Gao et al., *Gene Therapy* 2:710-722 (1995); Ahmad et al., *Cancer Res.* 52:4817-4820 (1992); U.S. Pat. Nos. 4,186,183, 4,217,344, 4,235,871, 4,261,975, 4,485,054, 4,501,728, 4,774,085, 4,837,028, and 4,946,787).

[0143] The use of RNA or DNA viral based systems for the delivery of nucleic acids takes advantage of highly evolved processes for targeting a virus to specific cells in the body and trafficking the viral payload to the nucleus. Viral vectors can be administered directly to patients (in vivo) or they can be used to treat cells in vitro, and the modified cells may optionally be administered to patients (ex vivo). Conventional viral based systems could include retroviral, lentivirus, adenoviral, adeno-associated and herpes simplex virus vectors for gene transfer. Integration in the host genome is possible with the retrovirus, lentivirus, and adeno-associated virus gene transfer methods, often result-

ing in long term expression of the inserted transgene. Additionally, high transduction efficiencies have been observed in many different cell types and target tissues.

[0144] The tropism of a retrovirus can be altered by incorporating foreign envelope proteins, expanding the potential target population of target cells. Lentiviral vectors are retroviral vectors that are able to transduce or infect non-dividing cells and typically produce high viral titers. Selection of a retroviral gene transfer system would therefore depend on the target tissue. Retroviral vectors are comprised of cis-acting long terminal repeats with packaging capacity for up to 6-10 kb of foreign sequence. The minimum cis-acting LTRs are sufficient for replication and packaging of the vectors, which are then used to integrate the therapeutic gene into the target cell to provide permanent transgene expression. Widely used retroviral vectors include those based upon murine leukemia virus (MuLV), gibbon ape leukemia virus (GaLV), Simian Immuno deficiency virus (SIV), human immuno deficiency virus (HIV), and combinations thereof (see, e.g., Buchscher et al., *J. Virol.* 66:2731-2739 (1992); Johann et al., *J. Virol.* 66:1635-1640 (1992); Sommerfelt et al., *Virol.* 176:58-59 (1990); Wilson et al., *J. Virol.* 63:2374-2378 (1989); Miller et al., *J. Virol.* 65:2220-2224 (1991); PCT/US94/05700). In applications where transient expression is preferred, adenoviral based systems may be used. Adenoviral based vectors are capable of very high transduction efficiency in many cell types and do not require cell division. With such vectors, high titer and levels of expression have been obtained. This vector can be produced in large quantities in a relatively simple system. Adeno-associated virus ("AAV") vectors may also be used to transduce cells with target nucleic acids, e.g., in the *in vitro* production of nucleic acids and peptides, and for *in vivo* and *ex vivo* gene therapy procedures (see, e.g., West et al., *Virology* 160:38-47 (1987); U.S. Pat. No. 4,797,368; WO 93/24641; Kotin, *Human Gene Therapy* 5:793-801 (1994); Muzyczka, *J. Clin. Invest.* 94:1351 (1994). Construction of recombinant AAV vectors are described in a number of publications, including U.S. Pat. No. 5,173,414; Tratschin et al., *Mol. Cell. Biol.* 5:3251-3260 (1985); Tratschin, et al., *Mol. Cell. Biol.* 4:2072-2081 (1984); Hermonat & Muzyczka, *PNAS* 81:6466-6470 (1984); and Samulski et al., *J. Virol.* 63:03822-3828 (1989).

[0145] Packaging cells are typically used to form virus particles that are capable of infecting a host cell. Such cells include 293 cells, which package adenovirus, and ψ 2 cells or PA317 cells, which package retrovirus. Viral vectors used in gene therapy are usually generated by producing a cell line that packages a nucleic acid vector into a viral particle. The vectors typically contain the minimal viral sequences required for packaging and subsequent integration into a host, other viral sequences being replaced by an expression cassette for the polynucleotide(s) to be expressed. The missing viral functions are typically supplied in trans by the packaging cell line. For example, AAV vectors used in gene therapy typically only possess ITR sequences from the AAV genome which are required for packaging and integration into the host genome. Viral DNA is packaged in a cell line, which contains a helper plasmid encoding the other AAV genes, namely rep and cap, but lacking ITR sequences. The cell line may also be infected with adenovirus as a helper. The helper virus promotes replication of the AAV vector and expression of AAV genes from the helper plasmid. The helper plasmid is not packaged in significant amounts due to

a lack of ITR sequences. Contamination with adenovirus can be reduced by, e.g., heat treatment to which adenovirus is more sensitive than AAV. Additional methods for the delivery of nucleic acids to cells are known to those skilled in the art. See, for example, US20030087817, incorporated herein by reference.

[0146] In some embodiments, a host cell is transiently or non-transiently transfected with one or more vectors described herein. In some embodiments, a cell is transfected as it naturally occurs in a subject. In some embodiments, a cell that is transfected is taken from a subject. In some embodiments, the cell is derived from cells taken from a subject, such as a cell line. A wide variety of cell lines for tissue culture are known in the art. Examples of cell lines include, but are not limited to, C8161, CCRF-CEM, MOLT, mIMCD-3, NHDF, HeLa-S3, Huh1, Huh4, Huh7, HUVEC, HASMC, HEK293, HEK293T, MiaPaCell, Panc1, PC-3, TF1, CTLL-2, C1R, Rat6, CV1, RPTE, A10, T24, J82, A375, ARH-77, Calu1, SW480, SW620, SKOV3, SK-UT, CaCo2, P388D1, SEM-K2, WEHI-231, HB56, TIB55, Jurkat, J45.01, LRMB, Bcl-1, BC-3, IC21, DLD2, Raw264.7, NRK, NRK-52E, MRCS, MEF, Hep G2, HeLa B, HeLa T4, COS, COS-1, COS-6, COS-M6A, BS-C-1 monkey kidney epithelial, BALB/3T3 mouse embryo fibroblast, 3T3 Swiss, 3T3-L1, 132-d5 human fetal fibroblasts; 10.1 mouse fibroblasts, 293-T, 3T3, 721, 9L, A2780, A2780ADR, A2780cis, A172, A20, A253, A431, A-549, ALC, B16, B35, BCP-1 cells, BEAS-2B, bEnd.3, BHK-21, BR 293, BxPC3, C3H-10T1/2, C6/36, Cal-27, CHO, CHO-7, CHO-IR, CHO-K1, CHO-K2, CHO-T, CHO Dhfr $-/-$, COR-L23, COR-L23/CPR, COR-L23/5010, COR-L23/R23, COS-7, COV-434, CML T1, CMT, CT26, D17, DH82, DU145, DuCaP, EL4, EM2, EM3, EMT6/AR1, EMT6/AR10.0, FM3, H1299, H69, HB54, HB55, HCA2, HEK-293, HeLa, Hepa1c1c7, HL-60, HMEC, HT-29, Jurkat, JY cells, K562 cells, Ku812, KCL22, KG1, KYO1, LNCap, Ma-Mel 1-48, MC-38, MCF-7, MCF-10A, MDA-MB-231, MDA-MB-468, MDA-MB-435, MDCK II, MDCK II, MOR/0.2R, MONO-MAC 6, MTD-1A, MyEnd, NCI-H69/CPR, NCI-H69/LX10, NCI-H69/LX20, NCI-H69/LX4, NIH-3T3, NALM-1, NW-145, OPCN/OPCT cell lines, Peer, PNT-1A/PNT 2, RenCa, RIN-5F, RMA/RMAS, Saos-2 cells, Sf-9, SkBr3, T2, T-47D, T84, THP1 cell line, U373, U87, U937, VCaP, Vero cells, WM39, WT-49, X63, YAC-1, YAR, and transgenic varieties thereof. Cell lines are available from a variety of sources known to those with skill in the art (see, e.g., the American Type Culture Collection (ATCC) (Manassas, Va.)). In some embodiments, a cell transfected with one or more vectors described herein is used to establish a new cell line comprising one or more vector-derived sequences. In some embodiments, a cell transiently transfected with the components of a CRISPR system as described herein (such as by transient transfection of one or more vectors, or transfection with RNA), and modified through the activity of a CRISPR complex, is used to establish a new cell line comprising cells containing the modification but lacking any other exogenous sequence. In some embodiments, cells transiently or non-transiently transfected with one or more vectors described herein, or cell lines derived from such cells are used in assessing one or more test compounds.

[0147] In some embodiments, one or more vectors described herein are used to produce a non-human transgenic animal or transgenic plant. In some embodiments, the transgenic animal is a mammal, such as a mouse, rat, or

rabbit. In certain embodiments, the organism or subject is a plant. In certain embodiments, the organism or subject or plant is algae. Methods for producing transgenic plants and animals are known in the art, and generally begin with a method of cell transfection, such as described herein.

[0148] In one aspect, the invention provides for methods of modifying a target polynucleotide in a eukaryotic cell. In some embodiments, the method comprises allowing a CRISPR complex to bind to the target polynucleotide to effect cleavage of said target polynucleotide thereby modifying the target polynucleotide, wherein the CRISPR complex comprises a CRISPR enzyme complexed with a guide sequence hybridized to a target sequence within said target polynucleotide, wherein said guide sequence is linked to a tracr mate sequence which in turn hybridizes to a tracr sequence.

[0149] In one aspect, the invention provides a method of modifying expression of a polynucleotide in a eukaryotic cell. In some embodiments, the method comprises allowing a CRISPR complex to bind to the polynucleotide such that said binding results in increased or decreased expression of said polynucleotide; wherein the CRISPR complex comprises a CRISPR enzyme complexed with a guide sequence hybridized to a target sequence within said polynucleotide, wherein said guide sequence is linked to a tracr mate sequence which in turn hybridizes to a tracr sequence.

[0150] With recent advances in crop genomics, the ability to use CRISPR-Cas systems to perform efficient and cost effective gene editing and manipulation will allow the rapid selection and comparison of single and multiplexed genetic manipulations to transform such genomes for improved production and enhanced traits. In this regard reference is made to US patents and publications: U.S. Pat. No. 6,603,061—*Agrobacterium*-Mediated Plant Transformation Method; U.S. Pat. No. 7,868,149—Plant Genome Sequences and Uses Thereof and US 2009/0100536—Transgenic Plants with Enhanced Agronomic Traits, all the contents and disclosure of each of which are herein incorporated by reference in their entirety. In the practice of the invention, the contents and disclosure of Morrell et al “Crop genomics: advances and applications” Nat Rev Genet. 2011 Dec. 29; 13(2):85-96 are also herein incorporated by reference in their entirety. In an advantageous embodiment of the invention, the CRISPR/Cas9 system is used to engineer microalgae (Example 15). Accordingly, reference herein to animal cells may also apply, *mutatis mutandis*, to plant cells unless otherwise apparent.

[0151] In one aspect, the invention provides for methods of modifying a target polynucleotide in a eukaryotic cell, which may be *in vivo*, *ex vivo* or *in vitro*. In some embodiments, the method comprises sampling a cell or population of cells from a human or non-human animal or plant (including micro-algae), and modifying the cell or cells. Culturing may occur at any stage *ex vivo*. The cell or cells may even be re-introduced into the non-human animal or plant (including micro-algae).

[0152] In plants, pathogens are often host-specific. For example, *Fusarium oxysporum* f sp. *lycopersici* causes tomato wilt but attacks only tomato, and *F. oxysporum* f. *dianthii Puccinia graminis* f sp. *tritici* attacks only wheat. Plants have existing and induced defenses to resist most pathogens. Mutations and recombination events across plant generations lead to genetic variability that gives rise to susceptibility, especially as pathogens reproduce with more

frequency than plants. In plants there can be non-host resistance, e.g., the host and pathogen are incompatible. There can also be Horizontal Resistance, e.g., partial resistance against all races of a pathogen, typically controlled by many genes and Vertical Resistance, e.g., complete resistance to some races of a pathogen but not to other races, typically controlled by a few genes. In a Gene-for-Gene level, plants and pathogens evolve together, and the genetic changes in one balance changes in other. Accordingly, using Natural Variability, breeders combine most useful genes for Yield, Quality, Uniformity, Hardiness, Resistance. The sources of resistance genes include native or foreign Varieties, Heirloom Varieties, Wild Plant Relatives, and Induced Mutations, e.g., treating plant material with mutagenic agents. Using the present invention, plant breeders are provided with a new tool to induce mutations. Accordingly, one skilled in the art can analyze the genome of sources of resistance genes, and in Varieties having desired characteristics or traits employ the present invention to induce the rise of resistance genes, with more precision than previous mutagenic agents and hence accelerate and improve plant breeding programs.

[0153] In one aspect, the invention provides kits containing any one or more of the elements disclosed in the above methods and compositions. In some embodiments, the kit comprises a vector system and instructions for using the kit. In some embodiments, the vector system comprises (a) a first regulatory element operably linked to a tracr mate sequence and one or more insertion sites for inserting a guide sequence upstream of the tracr mate sequence, wherein when expressed, the guide sequence directs sequence-specific binding of a CRISPR complex to a target sequence in a eukaryotic cell, wherein the CRISPR complex comprises a CRISPR enzyme complexed with (1) the guide sequence that is hybridized to the target sequence, and (2) the tracr mate sequence that is hybridized to the tracr sequence; and/or (b) a second regulatory element operably linked to an enzyme-coding sequence encoding said CRISPR enzyme comprising a nuclear localization sequence. Elements may be provide individually or in combinations, and may be provided in any suitable container, such as a vial, a bottle, or a tube. In some embodiments, the kit includes instructions in one or more languages, for example in more than one language.

[0154] In some embodiments, a kit comprises one or more reagents for use in a process utilizing one or more of the elements described herein. Reagents may be provided in any suitable container. For example, a kit may provide one or more reaction or storage buffers. Reagents may be provided in a form that is usable in a particular assay, or in a form that requires addition of one or more other components before use (e.g. in concentrate or lyophilized form). A buffer can be any buffer, including but not limited to a sodium carbonate buffer, a sodium bicarbonate buffer, a borate buffer, a Tris buffer, a MOPS buffer, a HEPES buffer, and combinations thereof. In some embodiments, the buffer is alkaline. In some embodiments, the buffer has a pH from about 7 to about 10. In some embodiments, the kit comprises one or more oligonucleotides corresponding to a guide sequence for insertion into a vector so as to operably link the guide sequence and a regulatory element. In some embodiments, the kit comprises a homologous recombination template polynucleotide.

[0155] In one aspect, the invention provides methods for using one or more elements of a CRISPR system. The CRISPR complex of the invention provides an effective means for modifying a target polynucleotide. The CRISPR complex of the invention has a wide variety of utility including modifying (e.g., deleting, inserting, translocating, inactivating, activating) a target polynucleotide in a multiplicity of cell types. As such the CRISPR complex of the invention has a broad spectrum of applications in, e.g., gene therapy, drug screening, disease diagnosis, and prognosis. An exemplary CRISPR complex comprises a CRISPR enzyme complexed with a guide sequence hybridized to a target sequence within the target polynucleotide. The guide sequence is linked to a tracr mate sequence, which in turn hybridizes to a tracr sequence.

[0156] The target polynucleotide of a CRISPR complex can be any polynucleotide endogenous or exogenous to the eukaryotic cell. For example, the target polynucleotide can be a polynucleotide residing in the nucleus of the eukaryotic cell. The target polynucleotide can be a sequence coding a gene product (e.g., a protein) or a non-coding sequence (e.g., a regulatory polynucleotide or a junk DNA). Without wishing to be bound by theory, it is believed that the target sequence should be associated with a PAM (protospacer adjacent motif); that is, a short sequence recognized by the CRISPR complex. The precise sequence and length requirements for the PAM differ depending on the CRISPR enzyme used, but PAMs are typically 2-5 base pair sequences adjacent the protospacer (that is, the target sequence). Examples of PAM sequences are given in the examples section below, and the skilled person will be able to identify further PAM sequences for use with a given CRISPR enzyme.

[0157] The target polynucleotide of a CRISPR complex may include a number of disease-associated genes and polynucleotides as well as signaling biochemical pathway-associated genes and polynucleotides as listed in U.S. provisional patent applications 61/736,527 and 61/748,427, both entitled SYSTEMS METHODS AND COMPOSITIONS FOR SEQUENCE MANIPULATION filed on Dec. 12, 2012 and Jan. 2, 2013, respectively, the contents of all of which are herein incorporated by reference in their entirety.

[0158] Examples of target polynucleotides include a sequence associated with a signaling biochemical pathway, e.g., a signaling biochemical pathway-associated gene or polynucleotide. Examples of target polynucleotides include a disease associated gene or polynucleotide. A “disease-associated” gene or polynucleotide refers to any gene or polynucleotide which is yielding transcription or translation products at an abnormal level or in an abnormal form in cells derived from a disease-affected tissues compared with tissues or cells of a non disease control. It may be a gene that becomes expressed at an abnormally high level; it may be a gene that becomes expressed at an abnormally low level, where the altered expression correlates with the occurrence and/or progression of the disease. A disease-associated gene also refers to a gene possessing mutation(s) or genetic variation that is directly responsible or is in linkage disequilibrium with a gene(s) that is responsible for the etiology of a disease. The transcribed or translated products may be known or unknown, and may be at a normal or abnormal level.

[0159] Examples of disease-associated genes and polynucleotides are available from McKusick-Nathans Institute of Genetic Medicine, Johns Hopkins University (Baltimore, Md.) and National Center for Biotechnology Information, National Library of Medicine (Bethesda, Md.), available on the World Wide Web.

[0160] Examples of disease-associated genes and polynucleotides are listed in Tables A and B. Disease specific information is available from McKusick-Nathans Institute of Genetic Medicine, Johns Hopkins University (Baltimore, Md.) and National Center for Biotechnology Information, National Library of Medicine (Bethesda, Md.), available on the World Wide Web. Examples of signaling biochemical pathway-associated genes and polynucleotides are listed in Table C.

[0161] Mutations in these genes and pathways can result in production of improper proteins or proteins in improper amounts which affect function. Further examples of genes, diseases and proteins are hereby incorporated by reference from U.S. Provisional application 61/736,527 filed on Dec. 12, 2012 and 61/748,427 filed on Feb. 2, 2013. Such genes, proteins and pathways may be the target polynucleotide of a CRISPR complex.

TABLE A

DISEASE/DISORDERS	GENE(S)
Neoplasia	PTEN; ATM; ATR; EGFR; ERBB2; ERBB3; ERBB4; Notch1; Notch2; Notch3; Notch4; AKT; AKT2; AKT3; HIF; HIF1a; HIF3a; Met; HRG; Bel2; PPAR alpha; PPAR gamma; WT1 (Wilms Tumor); FGF Receptor Family members (5 members: 1, 2, 3, 4, 5); CDKN2a; APC; RB (retinoblastoma); MEN1; VHL; BRCA1; BRCA2; AR (Androgen Receptor); TSG101; IGF; IGF Receptor; Igfl (4 variants); Igf2 (3 variants); Igf 1 Receptor; Igf 2 Receptor; Bax; Bcl2; caspases family (9 members: 1, 2, 3, 4, 6, 7, 8, 9, 12); Kras; Apc
Age-related Macular Degeneration	Aber; Ccl2; Cc2; ep (ceruloplasmin); Timp3; cathepsinD; Vldlr; Ccr2
Schizophrenia	Neuregulin1 (Nrg1); Erb4 (receptor for Neuregulin); Complexin1 (Cplx1); Tph1 Tryptophan hydroxylase; Tph2 Tryptophan hydroxylase 2; Neurexin 1; GSK3; GSK3a; GSK3b
Disorders	5-HTT (Slc6a4); COMT; DRD (Drd1a); SLC6A3; DAOA; DTNBP1; Dao (Dao1)

TABLE A-continued

DISEASE/DISORDERS	GENE(S)
Trinucleotide Repeat Disorders	HTT (Huntington's Dx); SBMA/SMAX1/AR (Kennedy's Dx); FXN/X25 (Friedrich's Ataxia); ATX3 (Machado-Joseph's Dx); ATXN1 and ATXN2 (spinocerebellar ataxias); DMPK (myotonic dystrophy); Atrophin-1 and Atn1 (DRPLA Dx); CBP (Creb-BP - global instability); VLDLR (Alzheimer's); Atxn7; Atxn10
Fragile X Syndrome	FMR2; FXR1; FXR2; mGLUR5
Secretase Related Disorders	APH-1 (alpha and beta); Presenilin (Psen1); nicastrin (Ncstn); PEN-2
Others	Nos1; Parp1; Nat1; Nat2
Prion - related disorders	Prp
ALS	SOD1; ALS2; STEX; FUS; TARDBP; VEGF (VEGF-a; VEGF-b; VEGF-c)
Drug addiction	Prkce (alcohol); Drd2; Drd4; ABAT (alcohol); GRIA2; Grm5; Grin1; Htr1b; Grin2a; Drd3; Pdyn; Gria1 (alcohol)
Autism	Mecp2; BZRAP1; MDGA2; Sema5A; Neurexin 1; Fragile X (FMR2 (AFF2)); FXR1; FXR2; Mglur5
Alzheimer's Disease	E1; CHIP; UCH; UBB; Tau; LRP; PICALM; Clusterin; PS1; SORL1; CR1; Vldlr; Uba1; Uba3; CHIP28 (Aqp1, Aquaporin 1); Uchl1; Uchl3; APP
Inflammation	IL-10; IL-1 (IL-1a; IL-1b); IL-13; IL-17 (IL-17a (CTLA8); IL-17b; IL-17c; IL-17d; IL-17f); IL-23; Cx3cr1; ptpn22; TNFa; NOD2/CARD15 for IBD; IL-6; IL-12 (IL-12a; IL-12b); CTLA4; Cx3cl1
Parkinson's Disease	α-Synuclein; DJ-1; LRRK2; Parkin; PINK1

TABLE B

Blood and coagulation diseases and disorders	Anemia (CDAN1, CDA1, RPS19, DBA, PKLR, PK1, NT5C3, UMPH1, PSN1, RHAG, RH50A, NRAMP2, SPTB, ALAS2, ANH1, ASB, ABCB7, ABC7, ASAT); Bare lymphocyte syndrome (TAPBP, TPSN, TAP2, ABCB3, PSF2, RING11, MHC2TA, C2TA, RFX5, RFXAP, RFX5); Bleeding disorders (TBXA2R, P2RX1, P2X1); Factor H and factor H-like 1 (HF1, CFH, HUS); Factor V and factor VIII (MCFD2); Factor VII deficiency (F7); Factor X deficiency (F10); Factor XI deficiency (F11); Factor XII deficiency (F12, HAF); Factor XIIIa deficiency (F13A1, F13A); Factor XIIIb deficiency (F13B); Fanconi anemia (FANCA, FACA, FA1, FA, FAA, FAAP95, FAAP90, FLJ34064, FANCB, FANCC, FACC, BRCA2, FANCD1, FANCD2, FANCD, FACD, FAD, FANCE, FACE, FANCF, XRCC9, FANCG, BRIP1, BACH1, FANCI, PHF9, FANCL, FANCM, KIAA1596); Hemophagocytic lymphohistiocytosis disorders (PRF1, HPLH2, UNC13D, MUNC13-4, HPLH3, HLH3, FHL3); Hemophilia A (F8, F8C, HEMA); Hemophilia B (F9, HEMB), Hemorrhagic disorders (PI, ATT, F5); Leukocyte deficiencies and disorders (ITGB2, CD18, LCAMB, LAD, EIF2B1, EIF2BA, EIF2B2, EIF2B3, EIF2B5, LVWM, CACH, CLE, EIF2B4); Sickle cell anemia (HBB); Thalassemia (HBA2, HBB, HBD, LCRB, HBA1).
Cell dysregulation and oncology diseases and disorders	B-cell non-Hodgkin lymphoma (BCL7A, BCL7); Leukemia (TAL1, TCL5, SCL, TAL2, FLT3, NBS1, NBS, ZNFN1A1, IK1, LYF1, HOXD4, HOX4B, BCR, CML, PHL, ALL, ARNT, KRAS2, RASK2, GMPS, AF10, ARHGEF12, LARG, KIAA0382, CALM, CLTH, CEBPA, CEBP, CHIC2, BTL, FLT3, KIT, PBT, LPP, NPM1, NUP214, D9S46E, CAN, CAIN, RUNX1, CBFA2, AML1, WHSC1L1, NSD3, FLT3, AF1Q, NPM1, NUMA1, ZNF145, PLZF, PML, MYL, STAT5B, AF10, CALM, CLTH, ARL11, ARLTS1, P2RX7, P2X7, BCR, CML, PHL, ALL, GRAF, NF1, VRNF, WSS, NFNS, PTPN11, PTP2C, SHP2, NS1, BCL2, CCND1, PRAD1, BCL1, TCRA, GATA1, GF1, ERYF1, NFE1, ABL1, NQO1, DIA4, NMOR1, NUP214, D9S46E, CAN, CAIN).
Inflammation and immune related diseases and disorders	AIDS (KIR3DL1, NKAT3, NKB1, AMB11, KIR3DS1, IFNG, CXCL12, SDF1); Autoimmune lymphoproliferative syndrome (TNFRSF6, APT1, FAS, CD95, ALPS1A); Combined immunodeficiency, (IL2RG, SCIDX1, SCIDX, IMD4); HIV-1 (CCL5, SCYA5, D17S136E, TCP228), HIV susceptibility or infection (IL10, CSIF, CMKBR2, CCR2, CMKBR5, CCKCR5 (CCR5)); Immunodeficiencies (CD3E, CD3G, AICDA, AID, HIGM2, TNFRSF5, CD40, UNG, DGU, HIGM4, TNFRSF5, CD40LG, HIGM1, IGM, FOXP3, IPEX, AIID, XPID, PIDX, TNFRSF14B, TACD); Inflammation (IL-10, IL-1 (IL-1a, IL-1b), IL-13, IL-17 (IL-17a (CTLA8), IL-17b, IL-17c, IL-17d, IL-17f), IL-23, Cx3cr1, ptpn22, TNFa, NOD2/CARD15 for IBD, IL-6, IL-12 (IL-12a, IL-12b),

TABLE B-continued

Metabolic, liver, kidney and protein diseases and disorders	<p>CTLA4, Cx3c11); Severe combined immunodeficiencies (SCIDs)(JAK3, JAKL, DCLRE1C, ARTEMIS, SCIDA, RAG1, RAG2, ADA, PTPRC, CD45, LCA, IL7R, CD3D, T3D, IL2RG, SCIDX1, SCIDX, IMD4). Amyloid neuropathy (TTR, PALB); Amyloidosis (APOA1, APP, AAA, CVAP, AD1, GSN, FGA, LYZ, TTR, PALB); Cirrhosis (KRT18, KRT8, CIRH1A, NAIC, TEX292, KIAA1988); Cystic fibrosis (CFTR, ABCC7, CF, MRP7); Glycogen storage diseases (SLC2A2, GLUT2, G6PC, G6PT, G6PT1, GAA, LAMP2, LAMPB, AGL, GDE, GBE1, GYS2, PYGL, PFKM); Hepatic adenoma, 142330 (TCF1, HNF1A, MODY3), Hepatic failure, early onset, and neurologic disorder (SCOD1, SCO1), Hepatic lipase deficiency (LIPC), Hepatoblastoma, cancer and carcinomas (CTNNB1, PDGFRL, PDGRL, PRLTS, AXIN1, AXIN, CTNNB1, TP53, P53, LFS1, IGF2R, MPRI, MET, CASP8, MCH5; Medullary cystic kidney disease (UMOD, HNFJ, FJHN, MCKD2, ADMCKD2); Phenylketonuria (PAH, PKU1, QDPR, DHPR, PTS); Polycystic kidney and hepatic disease (FCYT, PKHD1, ARPKD, PKD1, PKD2, PKD4, PKDTS, PRKCSH, G19P1, PCLD, SEC63).</p>
Muscular/Skeletal diseases and disorders	<p>Becker muscular dystrophy (DMD, BMD, MYF6), Duchenne Muscular Dystrophy (DMD, BMD); Emery-Dreifuss muscular dystrophy (LMNA, LMN1, EMD2, FPLD, CMD1A, HGPS, LGMD1B, LMNA, LMN1, EMD2, FPLD, CMD1A); Facioscapulohumeral muscular dystrophy (FSHMD1A, FSHD1A); Muscular dystrophy (FKRP, MDC1C, LGMD2I, LAMA2, LAMM, LARGE, KIAA0609, MDC1D, FCMD, TTID, MYOT, CAPN3, CANP3, DYSF, LGMD2B, SGCG, LGMD2C, DMDA1, SCG3, SGCA, ADL, DAG2, LGMD2D, DMDA2, SGCB, LGMD2E, SGCD, SGD, LGMD2F, CMD1L, TCAP, LGMD2G, CMD1N, TRIM32, HT2A, LGMD2H, FKRP, MDC1C, LGMD2I, TTN, CMD1G, TMD, LGMD2J, POMT1, CAV3, LGMD1C, SEPN1, SELN, RSMD1, PLEC1, PLTN, EBS1); Osteopetrosis (LRP5, BMND1, LRP7, LR3, OPG, VBCH2, CLCN7, CLC7, OPTA2, OSTM1, GL, TCIRG1, TIRC7, OC116, OPTB1); Muscular atrophy (VAPB, VAPC, ALS8, SMN1, SMA1, SMA2, SMA3, SMA4, BSCL2, SPG17, GARS, SMAD1, CMT2D, HEXB, IGHMBP2, SMUBP2, CATF1, SMARD1).</p>
Neurological and neuronal diseases and disorders	<p>ALS (SOD1, ALS2, STEX, FUS, TARDBP, VEGF (VEGF-a, VEGF-b, VEGF-c); Alzheimer disease (APP, AAA, CVAP, AD1, APOE, AD2, PSEN2, AD4, STM2, APBB2, FE65L1, NOS3, PLAU, URK, ACE, DCP1, ACE1, MPO, PACIP1, PAXIP1L, PTP, A2M, BLMH, BMH, PSEN1, AD3); Autism (Mecp2, BZRAP1, MDGA2, Sema5A, Neurexin1, GLO1, MECP2, RTT, PPMX, MRX16, MRX79, NLGN3, NLGN4, KIAA1260, AUTSX2); Fragile X Syndrome (FMR2, FXR1, FXR2, mGLUR5); Huntington's disease and disease like disorders (HD, IT15, PRNP, PRIP, JPH3, JP3, HDL2, TBP, SCA17); Parkinson disease (NR4A2, NURR1, NOT, TINUR, SNCAIP, TBP, SCA17, SNCA, NACP, PARK1, PARK4, DJ1, PARK7, LRRK2, PARK8, PINK1, PARK6, UCHL1, PARK5, SNCA, NACP, PARK1, PARK4, PRKN, PARK2, PDJ, DBH, NDUFV2); Rett syndrome (MECP2, RTT, PPMX, MRX16, MRX79, CDKL5, STK9, MECP2, RTT, PPMX, MRX16, MRX79, x-Synuclein, DJ-1); Schizophrenia (Neuregulin1 (Nrg1), Erb4 (receptor for Neuregulin), Complexin1 (Cplx1), Tph1 Tryptophan hydroxylase, Tph2, Tryptophan hydroxylase 2, Neurexin 1, GSK3, GSK3a, GSK3b, 5-HTT (Slc6a4), COMT, DRD (Drd1a), SLC6A3, DAOA, DTNBP1, Dao (Dao1)); Secretase Related Disorders (APH-1 (alpha and beta), Presenilin (Psen1), nicastrin, (Ncstn), PEN-2, Nos1, Parp1, Nat1, Nat2); Trinucleotide Repeat Disorders (HTT (Huntington's Dx), SBMA/SMAX1/AR (Kennedy's Dx), FXN/X25 (Friedrich's Ataxia), ATX3 (Machado-Joseph's Dx), ATXN1 and ATXN2 (spinocerebellar ataxias), DMPK (myotonic dystrophy), Atrophin-1 and Atn1 (DRPLA Dx), CBP (Creb-BP-global instability), VLDLR (Alzheimer's), Atxn7, Atxn10).</p>
Ocular diseases and disorders	<p>Age-related macular degeneration (Aber, Ccl2, Cc2, cp (ceruloplasmin), Timp3, cathepsinD, Vldlr, Cer2); Cataract (CRYAA, CRYA1, CRYBB2, CRYB2, PITX3, BFSP2, CP49, CP47, CRYAA, CRYA1, PAX6, AN2, MGDA, CRYBA1, CRYB1, CRYGC, CRYG3, CCL, LIM2, MP19, CRYGD, CRYG4, BFSP2, CP49, CP47, HSF4, CTM, HSF4, CTM, MIP, AQP0, CRYAB, CRYA2, CTTP2, CRYBB1, CRYGD, CRYG4, CRYBB2, CRYB2, CRYGC, CRYG3, CCL, CRYAA, CRYA1, GJA8, CX50, CAE1, GJA3, CX46, CZP3, CAE3, CCM1, CAM, KRIT1); Corneal clouding and dystrophy (APOA1, TGFBI, CSD2, CDGG1, CSD, BIGH3, CDG2, TACSTD2, TROP2, M1S1, VSX1, RINX, PPCD, PPD, KTCN, COL8A2, FECD, PPCD2, PIP5K3, CFD); Cornea plana congenital (KERA, CNA2); Glaucoma (MYOC, TIGR, GLC1A, JOAG, GPOA, OPTN, GLC1E, FIP2, HYPL, NRP, CYP1B1, GLC3A, OPA1, NTG, NPG, CYP1B1, GLC3A); Leber congenital amaurosis (CRB1, RP12, CRX, CORD2, CRD, RPGRIP1, LCA6, CORD9, RPE65, RP20, AIPL1, LCA4, GUCY2D, GUC2D, LCA1, CORD6, RDH12, LCA3);</p>

TABLE B-continued

Macular dystrophy (ELOVL4, ADMD, STGD2, STGD3, RDS, RP7, PRPH2, PRPH, AVMD, AOFMD, VMD2).

TABLE C

CELLULAR FUNCTION	GENES
PI3K/AKT Signaling	PRKCE; ITGAM; ITGA5; IRAK1; PRKAA2; EIF2AK2; PTEN; EIF4E; PRKCZ; GRK6; MAPK1; TSC1; PLK1; AKT2; IKKBK; PIK3CA; CDK8; CDKN1B; NFKB2; BCL2; PIK3CB; PPP2R1A; MAPK8; BCL2L1; MAPK3; TSC2; ITGA1; KRAS; EIF4EBP1; RELA; PRKCD; NOS3; PRKAA1; MAPK9; CDK2; PPP2CA; PIM1; ITGB7; YWHAZ; ILK; TP53; RAF1; IKBK; RELB; DYRK1A; CDKN1A; ITGB1; MAP2K2; JAK1; AKT1; JAK2; PIK3R1; CHUK; PDPK1; PPP2R5C; CTNNB1; MAP2K1; NFKB1; PAK3; ITGB3; CCND1; GSK3A; FRAP1; SFN; ITGA2; TTK; CSNK1A1; BRAF; GSK3B; AKT3; FOXO1; SGK; HSP90AA1; RPS6KB1
ERK/MAPK Signaling	PRKCE; ITGAM; ITGA5; HSPB1; IRAK1; PRKAA2; EIF2AK2; RAC1; RAP1A; TLN1; EIF4E; ELK1; GRK6; MAPK1; RAC2; PLK1; AKT2; PIK3CA; CDK8; CREB1; PRKCI; PTK2; FOS; RPS6KA4; PIK3CB; PPP2R1A; PIK3C3; MAPK8; MAPK3; ITGA1; ETS1; KRAS; MYCN; EIF4EBP1; PPARG; PRKCD; PRKAA1; MAPK9; SRC; CDK2; PPP2CA; PIM1; PIK3C2A; ITGB7; YWHAZ; PPP1CC; KSR1; PXN; RAF1; FYN; DYRK1A; ITGB1; MAP2K2; PAK4; PIK3R1; STAT3; PPP2R5C; MAP2K1; PAK3; ITGB3; ESR1; ITGA2; MYC; TTK; CSNK1A1; CRKL; BRAF; ATF4; PRKCA; SRF; STAT1; SGK
Glucocorticoid Receptor Signaling	RAC1; TAF4B; EP300; SMAD2; TRAF6; PCAF; ELK1; MAPK1; SMAD3; AKT2; IKKBK; NCOR2; UBE2I; PIK3CA; CREB1; FOS; HSPA5; NFKB2; BCL2; MAP3K14; STAT5B; PIK3CB; PIK3C3; MAPK8; BCL2L1; MAPK3; TSC2D3; MAPK10; NRIP1; KRAS; MAPK13; RELA; STAT5A; MAPK9; NOS2A; PBX1; NR3C1; PIK3C2A; CDKN1C; TRAF2; SERPINE1; NCOA3; MAPK14; TNF; RAF1; IKBK; MAP3K7; CREBBP; CDKN1A; MAP2K2; JAK1; IL8; NCOA2; AKT1; JAK2; PIK3R1; CHUK; STAT3; MAP2K1; NFKB1; TGFBR1; ESR1; SMAD4; CEBPB; JUN; AR; AKT3; CCL2; MMP1; STAT1; IL6; HSP90AA1
Axonal Guidance Signaling	PRKCE; ITGAM; ROCK1; ITGA5; CXCR4; ADAM12; IGF1; RAC1; RAP1A; EIF4E; PRKCZ; NRP1; NTRK2; ARHGEF7; SMO; ROCK2; MAPK1; PGF; RAC2; PTPN11; GNAS; AKT2; PIK3CA; ERBB2; PRKCI; PTK2; CFL1; GNAQ; PIK3CB; CXCL12; PIK3C3; WNT11; PRKD1; GNB2L1; ABL1; MAPK3; ITGA1; KRAS; RHOA; PRKCD; PIK3C2A; ITGB7; GLI2; PXN; VASP; RAF1; FYN; ITGB1; MAP2K2; PAK4; ADAM17; AKT1; PIK3R1; GLI1; WNT5A; ADAM10; MAP2K1; PAK3; ITGB3; CDC42; VEGFA; ITGA2; EPHA8; CRKL; RND1; GSK3B; AKT3; PRKCA
Ephrin Receptor Signaling	PRKCE; ITGAM; ROCK1; ITGA5; CXCR4; IRAK1; PRKAA2; EIF2AK2; RAC1; RAP1A; GRK6; ROCK2; MAPK1; PGF; RAC2; PTPN11; GNAS; PLK1; AKT2; DOK1; CDK8; CREB1; PTK2; CFL1; GNAQ; MAP3K14; CXCL12; MAPK8; GNB2L1; ABL1; MAPK3; ITGA1; KRAS; RHOA; PRKCD; PRKAA1; MAPK9; SRC; CDK2; PIM1; ITGB7; PXN; RAF1; FYN; DYRK1A; ITGB1; MAP2K2; PAK4; AKT1; JAK2; STAT3; ADAM10; MAP2K1; PAK3; ITGB3; CDC42; VEGFA; ITGA2; EPHA8; TTK; CSNK1A1; CRKL; BRAF; PTPN13; ATF4; AKT3; SGK
Actin Cytoskeleton Signaling	ACTN4; PRKCE; ITGAM; ROCK1; ITGA5; IRAK1; PRKAA2; EIF2AK2; RAC1; INS; ARHGEF7; GRK6; ROCK2; MAPK1; RAC2; PLK1; AKT2; PIK3CA; CDK8; PTK2; CFL1; PIK3CB; MYH9; DIAPH1; PIK3C3; MAPK8; F2R; MAPK3; SLC9A1; ITGA1; KRAS; RHOA; PRKCD; PRKAA1; MAPK9; CDK2; PIM1; PIK3C2A; ITGB7; PPP1CC; PXN; VIL2; RAF1; GSN; DYRK1A; ITGB1;

TABLE C-continued

CELLULAR FUNCTION	GENES
Huntington's Disease Signaling	MAP2K2; PAK4; PIP5K1A; PIK3R1; MAP2K1; PAK3; ITGB3; CDC42; APC; ITGA2; TTK; CSNK1A1; CRKL; BRAF; VAV3; SGK PRKCE; IGF1; EP300; RCOR1; PRKCZ; HDAC4; TGM2; MAPK1; CAPNS1; AKT2; EGFR; NCOR2; SP1; CAPN2; PIK3CA; HDAC5; CREB1; PRKCI; HSPA5; REST; GNAQ; PIK3CB; PIK3C3; MAPK8; IGF1R; PRKD1; GNB2L1; BCL2L1; CAPN1; MAPK3; CASP8; HDAC2; HDAC7A; PRKCD; HDAC11; MAPK9; HDAC9; PIK3C2A; HDAC3; TP53; CASP9; CREBBP; AKT1; PIK3R1; PDPK1; CASP1; APAF1; FRAP1; CASP2; JUN; BAX; ATF4; AKT3; PRKCA; CLTC; SGK; HDAC6; CASP3
Apoptosis Signaling	PRKCE; ROCK1; BID; IRAK1; PRKAA2; EIF2AK2; BAK1; BIRC4; GRK6; MAPK1; CAPNS1; PLK1; AKT2; IKBKB; CAPN2; CDK8; FAS; NFKB2; BCL2; MAP3K14; MAPK8; BCL2L1; CAPN1; MAPK3; CASP8; KRAS; RELA; PRKCD; PRKAA1; MAPK9; CDK2; PIM1; TP53; TNF; RAF1; IKBKG; RELB; CASP9; DYRK1A; MAP2K2; CHUK; APAF1; MAP2K1; NFKB1; PAK3; LMNA; CASP2; BIRC2; TTK; CSNK1A1; BRAF; BAX; PRKCA; SGK; CASP3; BIRC3; PARP1
B Cell Receptor Signaling	RAC1; PTEN; LYN; ELK1; MAPK1; RAC2; PTPN11; AKT2; IKBKB; PIK3CA; CREB1; SYK; NFKB2; CAMK2A; MAP3K14; PIK3CB; PIK3C3; MAPK8; BCL2L1; ABL1; MAPK3; ETS1; KRAS; MAPK13; RELA; PTPN6; MAPK9; EGR1; PIK3C2A; BTK; MAPK14; RAF1; IKBKG; RELB; MAP3K7; MAP2K2; AKT1; PIK3R1; CHUK; MAP2K1; NFKB1; CDC42; GSK3A; FRAP1; BCL6; BCL10; JUN; GSK3B; ATF4; AKT3; VAV3; RPS6KB1
Leukocyte Extravasation Signaling	ACTN4; CD44; PRKCE; ITGAM; ROCK1; CXCR4; CYBA; RAC1; RAP1A; PRKCZ; ROCK2; RAC2; PTPN11; MMP14; PIK3CA; PRKCI; PTK2; PIK3CB; CXCL12; PIK3C3; MAPK8; PRKD1; ABL1; MAPK10; CYBB; MAPK13; RHOA; PRKCD; MAPK9; SRC; PIK3C2A; BTK; MAPK14; NOX1; PXN; VIL2; VASP; ITGB1; MAP2K2; CTNND1; PIK3R1; CTNNB1; CLDN1; CDC42; F11R; ITK; CRKL; VAV3; CTTN; PRKCA; MMP1; MMP9
Integrin Signaling	ACTN4; ITGAM; ROCK1; ITGA5; RAC1; PTEN; RAP1A; TLN1; ARHGEF7; MAPK1; RAC2; CAPNS1; AKT2; CAPN2; PIK3CA; PTK2; PIK3CB; PIK3C3; MAPK8; CAV1; CAPN1; ABL1; MAPK3; ITGA1; KRAS; RHOA; SRC; PIK3C2A; ITGB7; PPP1CC; ILK; PXN; VASP; RAF1; FYN; ITGB1; MAP2K2; PAK4; AKT1; PIK3R1; TNK2; MAP2K1; PAK3; ITGB3; CDC42; RND3; ITGA2; CRKL; BRAF; GSK3B; AKT3
Acute Phase Response Signaling	IRAK1; SOD2; MYD88; TRAF6; ELK1; MAPK1; PTPN11; AKT2; IKBKB; PIK3CA; FOS; NFKB2; MAP3K14; PIK3CB; MAPK8; RIPK1; MAPK3; IL6ST; KRAS; MAPK13; IL6R; RELA; SOCS1; MAPK9; FTL; NR3C1; TRAF2; SERPINE1; MAPK14; TNF; RAF1; PDK1; IKKBG; RELB; MAP3K7; MAP2K2; AKT1; JAK2; PIK3R1; CHUK; STAT3; MAP2K1; NFKB1; FRAP1; CEBPB; JUN; AKT3; IL1R1; IL6
PTEN Signaling	ITGAM; ITGA5; RAC1; PTEN; PRKCZ; BCL2L11; MAPK1; RAC2; AKT2; EGFR; IKBKB; CBL; PIK3CA; CDKN1B; PTK2; NFKB2; BCL2; PIK3CB; BCL2L1; MAPK3; ITGA1; KRAS; ITGB7; ILK; PDGFRB; INSR; RAF1; IKBKG; CASP9; CDKN1A; ITGB1; MAP2K2; AKT1; PIK3R1; CHUK; PDGFRA; PDPK1; MAP2K1; NFKB1; ITGB3; CDC42; CCND1; GSK3A; ITGA2; GSK3B; AKT3; FOXO1; CASP3; RPS6KB1
p53 Signaling	PTEN; EP300; BBC3; PCAF; FASN; BRCA1; GADD45A; BIRC5; AKT2; PIK3CA; CHEK1; TP53INP1; BCL2; PIK3CB; PIK3C3; MAPK8; THBS1; ATR; BCL2L1; E2F1; PMAIP1; CHEK2; TNFRSF10B; TP73; RB1; HDAC9; CDK2; PIK3C2A; MAPK14; TP53; LRDD; CDKN1A; HIPK2; AKT1; PIK3R1; RRM2B; APAF1; CTNNB1; SIRT1; CCND1; PRKDC; ATM; SFN; CDKN2A; JUN; SNAI2; GSK3B; BAX; AKT3
Aryl Hydrocarbon Receptor Signaling	HSPB1; EP300; FASN; TGM2; RXRA; MAPK1; NQO1; NCOR2; SP1; ARNT; CDKN1B; FOS; CHEK1; SMARCA4; NFKB2; MAPK8; ALDH1A1; ATR; E2F1;

TABLE C-continued

CELLULAR FUNCTION	GENES
	MAPK3; NRIP1; CHEK2; RELA; TP73; GSTP1; RB1; SRC; CDK2; AHR; NFE2L2; NCOA3; TP53; TNF; CDKN1A; NCOA2; APAF1; NFKB1; CCND1; ATM; ESR1; CDKN2A; MYC; JUN; ESR2; BAX; IL6; CYP1B1; HSP90AA1
Xenobiotic Metabolism Signaling	PRKCE; EP300; PRKCZ; RXRA; MAPK1; NQO1; NCOR2; PIK3CA; ARNT; PRKCI; NFKB2; CAMK2A; PIK3CB; PPP2R1A; PIK3C3; MAPK8; PRKD1; ALDH1A1; MAPK3; NRIP1; KRAS; MAPK13; PRKCD; GSTP1; MAPK9; NOS2A; ABCB1; AHR; PPP2CA; FTL; NFE2L2; PIK3C2A; PPARGC1A; MAPK14; TNF; RAF1; CREBBP; MAP2K2; PIK3R1; PPP2R5C; MAP2K1; NFKB1; KEAP1; PRKCA; EIF2AK3; IL6; CYP1B1; HSP90AA1
SAPK/JNK Signaling	PRKCE; IRAK1; PRKAA2; EIF2AK2; RAC1; ELK1; GRK6; MAPK1; GADD45A; RAC2; PLK1; AKT2; PIK3CA; FADD; CDK8; PIK3CB; PIK3C3; MAPK8; RIPK1; GNB2L1; IRS1; MAPK3; MAPK10; DAXX; KRAS; PRKCD; PRKAA1; MAPK9; CDK2; PIM1; PIK3C2A; TRAF2; TP53; LCK; MAP3K7; DYRK1A; MAP2K2; PIK3R1; MAP2K1; PAK3; CDC42; JUN; TTK; CSNK1A1; CRKL; BRAF; SGK
PPAr/RXR Signaling	PRKAA2; EP300; INS; SMAD2; TRAF6; PPARA; FASN; RXRA; MAPK1; SMAD3; GNAS; IKBKB; NCOR2; ABCA1; GNAQ; NFKB2; MAP3K14; STAT5B; MAPK8; IRS1; MAPK3; KRAS; RELA; PRKAA1; PPARGC1A; NCOA3; MAPK14; INSR; RAF1; IKBKG; RELB; MAP3K7; CREBBP; MAP2K2; JAK2; CHUK; MAP2K1; NFKB1; TGFBF1; SMAD4; JUN; IL1R1; PRKCA; IL6; HSP90AA1; ADIPOQ
NF-KB Signaling	IRAK1; EIF2AK2; EP300; INS; MYD88; PRKCZ; TRAF6; TBK1; AKT2; EGFR; IKBKB; PIK3CA; BTRC; NFKB2; MAP3K14; PIK3CB; PIK3C3; MAPK8; RIPK1; HDAC2; KRAS; RELA; PIK3C2A; TRAF2; TLR4; PDGFRB; TNF; INSR; LCK; IKBKG; RELB; MAP3K7; CREBBP; AKT1; PIK3R1; CHUK; PDGFRA; NFKB1; TLR2; BCL10; GSK3B; AKT3; TNFAIP3; IL1R1
Neuregulin Signaling	ERBB4; PRKCE; ITGAM; ITGA5; PTEN; PRKCZ; ELK1; MAPK1; PTPN11; AKT2; EGFR; ERBB2; PRKCI; CDKN1B; STAT5B; PRKD1; MAPK3; ITGA1; KRAS; PRKCD; STAT5A; SRC; ITGB7; RAF1; ITGB1; MAP2K2; ADAM17; AKT1; PIK3R1; PDPK1; MAP2K1; ITGB3; EREG; FRAP1; PSEN1; ITGA2; MYC; NRG1; CRKL; AKT3; PRKCA; HSP90AA1; RPS6KB1
Wnt & Beta catenin Signaling	CD44; EP300; LRP6; DVL3; CSNK1E; GJA1; SMO; AKT2; PIN1; CDH1; BTRC; GNAQ; MARK2; PPP2R1A; WNT11; SRC; DKK1; PPP2CA; SOX6; SFRP2; ILK; LEF1; SOX9; TP53; MAP3K7; CREBBP; TCF7L2; AKT1; PPP2R5C; WNT5A; LRP5; CTNNB1; TGFBF1; CCND1; GSK3A; DVL1; APC; CDKN2A; MYC; CSNK1A1; GSK3B; AKT3; SOX2
Insulin Receptor Signaling	PTEN; INS; EIF4E; PTPN1; PRKCZ; MAPK1; TSC1; PTPN11; AKT2; CBL; PIK3CA; PRKCI; PIK3CB; PIK3C3; MAPK8; IRS1; MAPK3; TSC2; KRAS; EIF4EBP1; SLC2A4; PIK3C2A; PPP1CC; INSR; RAF1; FYN; MAP2K2; JAK1; AKT1; JAK2; PIK3R1; PDPK1; MAP2K1; GSK3A; FRAP1; CRKL; GSK3B; AKT3; FOXO1; SGK; RPS6KB1
IL-6 Signaling	HSPB1; TRAF6; MAPKAPK2; ELK1; MAPK1; PTPN11; IKKBK; FOS; NFKB2; MAP3K14; MAPK8; MAPK3; MAPK10; IL6ST; KRAS; MAPK13; IL6R; RELA; SOCS1; MAPK9; ABCB1; TRAF2; MAPK14; TNF; RAF1; IKBKG; RELB; MAP3K7; MAP2K2; IL8; JAK2; CHUK; STAT3; MAP2K1; NFKB1; CEBPB; JUN; IL1R1; SRF; IL6
Hepatic Cholestasis	PRKCE; IRAK1; INS; MYD88; PRKCZ; TRAF6; PPARA; RXRA; IKBKB; PRKCI; NFKB2; MAP3K14; MAPK8; PRKD1; MAPK10; RELA; PRKCD; MAPK9; ABCB1; TRAF2; TLR4; TNF; INSR; IKBKG; RELB; MAP3K7; IL8; CHUK; NR1H2; TJP2; NFKB1; ESR1; SREBF1; FGFR4; JUN; IL1R1; PRKCA; IL6
IGF-1 Signaling	IGF1; PRKCZ; ELK1; MAPK1; PTPN11; NEDD4; AKT2; PIK3CA; PRKCI; PTK2; FOS; PIK3CB; PIK3C3; MAPK8; IGF1R; IRS1; MAPK3; IGFBP7; KRAS; PIK3C2A;

TABLE C-continued

CELLULAR FUNCTION	GENES
	YWHAZ; PXN; RAF1; CASP9; MAP2K2; AKT1; PIK3R1; PDPK1; MAP2K1; IGFBP2; SFN; JUN; CYR61; AKT3; FOXO1; SRF; CTGF; RPS6KB1
NRF2-mediated Oxidative Stress Response	PRKCE; EP300; SOD2; PRKCZ; MAPK1; SQSTM1; NQO1; PIK3CA; PRKCI; FOS; PIK3CB; PIK3C3; MAPK8; PRKD1; MAPK3; KRAS; PRKCD; GTP1; MAPK9; FTL; NFE2L2; PIK3C2A; MAPK14; RAF1; MAP3K7; CREBBP; MAP2K2; AKT1; PIK3R1; MAP2K1; PPIB; JUN; KEAP1; GSK3B; ATF4; PRKCA; EIF2AK3; HSP90AA1
Hepatic Fibrosis/Hepatic Stellate Cell Activation	EDN1; IGF1; KDR; FLT1; SMAD2; FGFR1; MET; PGF; SMAD3; EGFR; FAS; CSF1; NFKB2; BCL2; MYH9; IGF1R; IL6R; RELA; TLR4; PDGFRB; TNF; RELB; IL8; PDGFRA; NFKB1; TGFB1; SMAD4; VEGFA; BAX; IL1R1; CCL2; HGF; MMP1; STAT1; IL6; CTGF; MMP9
PPAR Signaling	EP300; INS; TRAF6; PPARA; RXRA; MAPK1; IKBKB; NCOR2; FOS; NFKB2; MAP3K14; STAT5B; MAPK3; NRIP1; KRAS; PPARG; RELA; STAT5A; TRAF2; PPARGC1A; PDGFRB; TNF; INSR; RAF1; IKBKG; RELB; MAP3K7; CREBBP; MAP2K2; CHUK; PDGFRA; MAP2K1; NFKB1; JUN; IL1R1; HSP90AA1
Fc Epsilon RI Signaling	PRKCE; RAC1; PRKCZ; LYN; MAPK1; RAC2; PTPN11; AKT2; PIK3CA; SYK; PRKCI; PIK3CB; PIK3C3; MAPK8; PRKD1; MAPK3; MAPK10; KRAS; MAPK13; PRKCD; MAPK9; PIK3C2A; BTK; MAPK14; TNF; RAF1; FYN; MAP2K2; AKT1; PIK3R1; PDPK1; MAP2K1; AKT3; VAV3; PRKCA
G-Protein Coupled Receptor Signaling	PRKCE; RAP1A; RGS16; MAPK1; GNAS; AKT2; IKBKB; PIK3CA; CREB1; GNAQ; NFKB2; CAMK2A; PIK3CB; PIK3C3; MAPK3; KRAS; RELA; SRC; PIK3C2A; RAF1; IKBKG; RELB; FYN; MAP2K2; AKT1; PIK3R1; CHUK; PDPK1; STAT3; MAP2K1; NFKB1; BRAF; ATF4; AKT3; PRKCA
Inositol Phosphate Metabolism	PRKCE; IRAK1; PRKAA2; EIF2AK2; PTEN; GRK6; MAPK1; PLK1; AKT2; PIK3CA; CDK8; PIK3CB; PIK3C3; MAPK8; MAPK3; PRKCD; PRKAA1; MAPK9; CDK2; PIM1; PIK3C2A; DYRK1A; MAP2K2; PIP5K1A; PIK3R1; MAP2K1; PAK3; ATM; TTK; CSNK1A1; BRAF; SGK
PDGF Signaling	EIF2AK2; ELK1; ABL2; MAPK1; PIK3CA; FOS; PIK3CB; PIK3C3; MAPK8; CAV1; ABL1; MAPK3; KRAS; SRC; PIK3C2A; PDGFRB; RAF1; MAP2K2; JAK1; JAK2; PIK3R1; PDGFRA; STAT3; SPHK1; MAP2K1; MYC; JUN; CRKL; PRKCA; SRF; STAT1; SPHK2
VEGF Signaling	ACTN4; ROCK1; KDR; FLT1; ROCK2; MAPK1; PGF; AKT2; PIK3CA; ARNT; PTK2; BCL2; PIK3CB; PIK3C3; BCL2L1; MAPK3; KRAS; HIF1A; NOS3; PIK3C2A; PXN; RAF1; MAP2K2; ELAVL1; AKT1; PIK3R1; MAP2K1; SFN; VEGFA; AKT3; FOXO1; PRKCA
Natural Killer Cell Signaling	PRKCE; RAC1; PRKCZ; MAPK1; RAC2; PTPN11; KIR2DL3; AKT2; PIK3CA; SYK; PRKCI; PIK3CB; PIK3C3; PRKD1; MAPK3; KRAS; PRKCD; PTPN6; PIK3C2A; LCK; RAF1; FYN; MAP2K2; PAK4; AKT1; PIK3R1; MAP2K1; PAK3; AKT3; VAV3; PRKCA
Cell Cycle: G1/S Checkpoint Regulation	HDAC4; SMAD3; SUV39H1; HDAC5; CDKN1B; BTRC; ATR; ABL1; E2F1; HDAC2; HDAC7A; RB1; HDAC11; HDAC9; CDK2; E2F2; HDAC3; TP53; CDKN1A; CCND1; E2F4; ATM; RBL2; SMAD4; CDKN2A; MYC; NRG1; GSK3B; RBL1; HDAC6
T Cell Receptor Signaling	RAC1; ELK1; MAPK1; IKBKB; CBL; PIK3CA; FOS; NFKB2; PIK3CB; PIK3C3; MAPK8; MAPK3; KRAS; RELA; PIK3C2A; BTK; LCK; RAF1; IKBKG; RELB; FYN; MAP2K2; PIK3R1; CHUK; MAP2K1; NFKB1; ITK; BCL10; JUN; VAV3
Death Receptor Signaling	CRADD; HSPB1; BID; BIRC4; TBK1; IKBKB; FADD; FAS; NFKB2; BCL2; MAP3K14; MAPK8; RIPK1; CASP8; DAXX; TNFRSF10B; RELA; TRAF2; TNF; IKBKG; RELB; CASP9; CHUK; APAF1; NFKB1; CASP2; BIRC2; CASP3; BIRC3
FGF Signaling	RAC1; FGFR1; MET; MAPKAPK2; MAPK1; PTPN11; AKT2; PIK3CA; CREB1; PIK3CB; PIK3C3; MAPK8; MAPK3; MAPK13; PTPN6; PIK3C2A; MAPK14; RAF1;

TABLE C-continued

CELLULAR FUNCTION	GENES
	AKT1; PIK3R1; STAT3; MAP2K1; FGFR4; CRKL; ATF4; AKT3; PRKCA; HGF
GM-CSF Signaling	LYN; ELK1; MAPK1; PTPN11; AKT2; PIK3CA; CAMK2A; STAT5B; PIK3CB; PIK3C3; GNB2L1; BCL2L1; MAPK3; ETS1; KRAS; RUNX1; PIM1; PIK3C2A; RAF1; MAP2K2; AKT1; JAK2; PIK3R1; STAT3; MAP2K1; CCND1; AKT3; STAT1
Amyotrophic Lateral Sclerosis Signaling	BID; IGF1; RAC1; BIRC4; PGF; CAPNS1; CAPN2; PIK3CA; BCL2; PIK3CB; PIK3C3; BCL2L1; CAPN1; PIK3C2A; TP53; CASP9; PIK3R1; RAB5A; CASP1; APAF1; VEGFA; BIRC2; BAX; AKT3; CASP3; BIRC3
JAK/Stat Signaling	PTPN1; MAPK1; PTPN11; AKT2; PIK3CA; STAT5B; PIK3CB; PIK3C3; MAPK3; KRAS; SOCS1; STAT5A; PTPN6; PIK3C2A; RAF1; CDKN1A; MAP2K2; JAK1; AKT1; JAK2; PIK3R1; STAT3; MAP2K1; FRAP1; AKT3; STAT1
Nicotinate and Nicotinamide Metabolism	PRKCE; IRAK1; PRKAA2; EIF2AK2; GRK6; MAPK1;
	PLK1; AKT2; CDK8; MAPK8; MAPK3; PRKCD; PRKAA1; PBEF1; MAPK9; CDK2; PIM1; DYRK1A; MAP2K2; MAP2K1; PAK3; NT5E; TTK; CSNK1A1; BRAF; SGK
Chemokine Signaling	CXCR4; ROCK2; MAPK1; PTK2; FOS; CFL1; GNAQ; CAMK2A; CXCL12; MAPK8; MAPK3; KRAS; MAPK13; RHOA; CCR3; SRC; PPP1CC; MAPK14; NOX1; RAF1; MAP2K2; MAP2K1; JUN; CCL2; PRKCA
IL-2 Signaling	ELK1; MAPK1; PTPN11; AKT2; PIK3CA; SYK; FOS; STAT5B; PIK3CB; PIK3C3; MAPK8; MAPK3; KRAS; SOCS1; STAT5A; PIK3C2A; LCK; RAF1; MAP2K2; JAK1; AKT1; PIK3R1; MAP2K1; JUN; AKT3
Synaptic Long Term Depression	PRKCE; IGF1; PRKCZ; PRDX6; LYN; MAPK1; GNAS; PRKCI; GNAQ; PPP2R1A; IGF1R; PRKD1; MAPK3; KRAS; GRN; PRKCD; NOS3; NOS2A; PPP2CA; YWHAZ; RAF1; MAP2K2; PPP2R5C; MAP2K1; PRKCA
Estrogen Receptor Signaling	TAF4B; EP300; CARM1; PCAF; MAPK1; NCOR2; SMARCA4; MAPK3; NRIP1; KRAS; SRC; NR3C1; HDAC3; PPARGC1A; RBM9; NCOA3; RAF1; CREBBP; MAP2K2; NCOA2; MAP2K1; PRKDC; ESR1; ESR2
Protein Ubiquitination Pathway	TRAF6; SMURF1; BIRC4; BRCA1; UCHL1; NEDD4; CBL; UBE2I; BTRC; HSPA5; USP7; USP10; FBXW7; USP9X; STUB1; USP22; B2M; BIRC2; PARK2; USP8; USP1; VHL; HSP90AA1; BIRC3
IL-10 Signaling	TRAF6; CCR1; ELK1; IKKBK; SP1; FOS; NFKB2; MAP3K14; MAPK8; MAPK13; RELA; MAPK14; TNF; IKKBK; RELB; MAP3K7; JAK1; CHUK; STAT3; NFKB1; JUN; IL1R1; IL6
VDR/RXR Activation	PRKCE; EP300; PRKCZ; RXRA; GADD45A; HES1; NCOR2; SP1; PRKCI; CDKN1B; PRKD1; PRKCD; RUNX2; KLF4; YY1; NCOA3; CDKN1A; NCOA2; SPP1; LRP5; CEBPB; FOXO1; PRKCA
TGF-beta Signaling	EP300; SMAD2; SMURF1; MAPK1; SMAD3; SMAD1; FOS; MAPK8; MAPK3; KRAS; MAPK9; RUNX2; SERPINE1; RAF1; MAP3K7; CREBBP; MAP2K2; MAP2K1; TGFBF1; SMAD4; JUN; SMAD5
Toll-like Receptor Signaling	IRAK1; EIF2AK2; MYD88; TRAF6; PPARA; ELK1;
	IKKBK; FOS; NFKB2; MAP3K14; MAPK8; MAPK13; RELA; TLR4; MAPK14; IKKBK; RELB; MAP3K7; CHUK; NFKB1; TLR2; JUN
p38 MAPK Signaling	HSPB1; IRAK1; TRAF6; MAPKAPK2; ELK1; FADD; FAS; CREB1; DDIT3; RPS6KA4; DAXX; MAPK13; TRAF2; MAPK14; TNF; MAP3K7; TGFBF1; MYC; ATF4; IL1R1; SRF; STAT1
Neurotrophin/TRK Signaling	NTRK2; MAPK1; PTPN11; PIK3CA; CREB1; FOS;
	PIK3CB; PIK3C3; MAPK8; MAPK3; KRAS; PIK3C2A; RAF1; MAP2K2; AKT1; PIK3R1; PDPK1; MAP2K1; CDC42; JUN; ATF4
FXR/RXR Activation	INS; PPARA; FASN; RXRA; AKT2; SDC1; MAPK8; APOB; MAPK10; PPARG; MTPP; MAPK9; PPARGC1A; TNF; CREBBP; AKT1; SREBF1; FGFR4; AKT3; FOXO1
Synaptic Long Term Potentiation	PRKCE; RAPIA; EP300; PRKCZ; MAPK1; CREB1; PRKCI; GNAQ; CAMK2A; PRKD1; MAPK3; KRAS; PRKCD; PPP1CC; RAF1; CREBBP; MAP2K2; MAP2K1; ATF4; PRKCA

TABLE C-continued

CELLULAR FUNCTION	GENES
Calcium Signaling	RAP1A; EP300; HDAC4; MAPK1; HDAC5; CREB1; CAMK2A; MYH9; MAPK3; HDAC2; HDAC7A; HDAC11; HDAC9; HDAC3; CREBBP; CALR; CAMKK2; ATF4; HDAC6
EGF Signaling	ELK1; MAPK1; EGFR; PIK3CA; FOS; PIK3CB; PIK3C3; MAPK8; MAPK3; PIK3C2A; RAF1; JAK1; PIK3R1; STAT3; MAP2K1; JUN; PRKCA; SRF; STAT1
Hypoxia Signaling in the Cardiovascular System	EDN1; PTEN; EP300; NQO1; UBE2I; CREB1; ARNT; HIF1A; SLC2A4; NOS3; TP53; LDHA; AKT1; ATM; VEGFA; JUN; ATF4; VHL; HSP90AA1
LPS/IL-1 Mediated Inhibition of RXR Function	IRAK1; MYD88; TRAF6; PPARA; RXRA; ABCA1; MAPK8; ALDH1A1; GSTP1; MAPK9; ABCB1; TRAF2; TLR4; TNF; MAP3K7; NR1H2; SREBF1; JUN; IL1R1
LXR/RXR Activation	FASN; RXRA; NCOR2; ABCA1; NFKB2; IRF3; RELA; NOS2A; TLR4; TNF; RELB; LDLR; NR1H2; NFKB1; SREBF1; IL1R1; CCL2; IL6; MMP9
Amyloid Processing	PRKCE; CSNK1E; MAPK1; CAPNS1; AKT2; CAPN2; CAPN1; MAPK3; MAPK13; MAPT; MAPK14; AKT1; PSEN1; CSNK1A1; GSK3B; AKT3; APP
IL-4 Signaling	AKT2; PIK3CA; PIK3CB; PIK3C3; IRS1; KRAS; SOCS1; PTPN6; NR3C1; PIK3C2A; JAK1; AKT1; JAK2; PIK3R1; FRAP1; AKT3; RPS6KB1
Cell Cycle: G2/M DNA Damage Checkpoint Regulation	EP300; PCAF; BRCA1; GADD45A; PLK1; BTRC; CHEK1; ATR; CHEK2; YWHAZ; TP53; CDKN1A; PRKDC; ATM; SFN; CDKN2A
Nitric Oxide Signaling in the Cardiovascular System	KDR; FLT1; PGF; AKT2; PIK3CA; PIK3CB; PIK3C3; CAV1; PRKCD; NOS3; PIK3C2A; AKT1; PIK3R1; VEGFA; AKT3; HSP90AA1
Purine Metabolism	NME2; SMARCA4; MYH9; RRM2; ADAR; EIF2AK4; PKM2; ENTPD1; RAD51; RRM2B; TJP2; RAD51C; NT5E; POLD1; NME1
cAMP-mediated Signaling	RAP1A; MAPK1; GNAS; CREB1; CAMK2A; MAPK3; SRC; RAF1; MAP2K2; STAT3; MAP2K1; BRAF; ATF4
Mitochondrial Dysfunction	SOD2; MAPK8; CASP8; MAPK10; MAPK9; CASP9; PARK7; PSEN1; PARK2; APP; CASP3
Notch Signaling	HES1; JAG1; NUMB; NOTCH4; ADAM17; NOTCH2; PSEN1; NOTCH3; NOTCH1; DLL4
Endoplasmic Reticulum Stress Pathway	HSPA5; MAPK8; XBP1; TRAF2; ATF6; CASP9; ATF4; EIF2AK3; CASP3
Pyrimidine Metabolism	NME2; AICDA; RRM2; EIF2AK4; ENTPD1; RRM2B; NT5E; POLD1; NME1
Parkinson's Signaling	UCHL1; MAPK8; MAPK13; MAPK14; CASP9; PARK7; PARK2; CASP3
Cardiac & Beta Adrenergic Signaling	GNAS; GNAQ; PPP2R1A; GNB2L1; PPP2CA; PPP1CC; PPP2R5C
Glycolysis/Gluconeogenesis	HK2; GCK; GPI; ALDH1A1; PKM2; LDHA; HK1
Interferon Signaling	IRF1; SOCS1; JAK1; JAK2; IFITM1; STAT1; IFIT3
Sonic Hedgehog Signaling	ARRB2; SMO; GLI2; DYRK1A; GLI1; GSK3B; DYRK1B
Glycerophospholipid Metabolism	PLD1; GRN; GPAM; YWHAZ; SPHK1; SPHK2
Phospholipid Degradation	PRDX6; PLD1; GRN; YWHAZ; SPHK1; SPHK2
Tryptophan Metabolism	SLAH2; PRMT5; NEDD4; ALDH1A1; CYP1B1; SLAH1
Lysine Degradation	SUV39H1; EHMT2; NSD1; SETD7; PPP2R5C
Nucleotide Excision Repair Pathway	ERCC5; ERCC4; XPA; XPC; ERCC1
Starch and Sucrose Metabolism	UCHL1; HK2; GCK; GPI; HK1
Aminosugars Metabolism	NQO1; HK2; GCK; HK1
Arachidonic Acid Metabolism	PRDX6; GRN; YWHAZ; CYP1B1
Circadian Rhythm Signaling	CSNK1E; CREB1; ATF4; NR1D1
Coagulation System	BDKRB1; F2R; SERPINE1; F3
Dopamine Receptor Signaling	PPP2R1A; PPP2CA; PPP1CC; PPP2R5C
Glutathione Metabolism	IDH2; GSTP1; ANPEP; IDH1

TABLE C-continued

CELLULAR FUNCTION	GENES
Glycerolipid Metabolism	ALDH1A1; GPAM; SPHK1; SPHK2
Linoleic Acid Metabolism	PRDX6; GRN; YWHAZ; CYP1B1
Methionine Metabolism	DNMT1; DNMT3B; AHCY; DNMT3A
Pyruvate Metabolism	GLO1; ALDH1A1; PKM2; LDHA
Arginine and Proline Metabolism	ALDH1A1; NOS3; NOS2A
Eicosanoid Signaling	PRDX6; GRN; YWHAZ
Fructose and Mannose Metabolism	HK2; GCK; HK1
Galactose Metabolism	HK2; GCK; HK1
Stilbene, Coumarine and Lignin Biosynthesis	PRDX6; PRDX1; TYR
Antigen Presentation Pathway	CALR; B2M
Biosynthesis of Steroids	NQO1; DHCR7
Butanoate Metabolism	ALDH1A1; NLGN1
Citrate Cycle	IDH2; IDH1
Fatty Acid Metabolism	ALDH1A1; CYP1B1
Glycerophospholipid Metabolism	PRDX6; CHKA
Histidine Metabolism	PRMT5; ALDH1A1
Inositol Metabolism	ERO1L; APEX1
Metabolism of Xenobiotics	GSTP1; CYP1B1
by Cytochrome p450	
Methane Metabolism	PRDX6; PRDX1
Phenylalanine Metabolism	PRDX6; PRDX1
Propanoate Metabolism	ALDH1A1; LDHA
Selenoamino Acid Metabolism	PRMT5; AHCY
Sphingolipid Metabolism	SPHK1; SPHK2
Aminophosphonate Metabolism	PRMT5
Androgen and Estrogen Metabolism	PRMT5
Ascorbate and Aldarate Metabolism	ALDH1A1
Bile Acid Biosynthesis	ALDH1A1
Cysteine Metabolism	LDHA
Fatty Acid Biosynthesis	FASN
Glutamate Receptor Signaling	GNB2L1
NRF2-mediated Oxidative Stress Response	PRDX1
Pentose Phosphate Pathway	GPI
Pentose and Glucuronate Interconversions	UCHL1
Retinol Metabolism	ALDH1A1
Riboflavin Metabolism	TYR
Tyrosine Metabolism	PRMT5, TYR
Ubiquinone Biosynthesis	PRMT5
Valine, Leucine and Isoleucine Degradation	ALDH1A1
Glycine, Serine and Threonine Metabolism	CHKA
Lysine Degradation	ALDH1A1
Pain/Taste	TRPM5; TRPA1
Pain	TRPM7; TRPC5; TRPC6; TRPC1; Cnr1; cnr2; Grk2; Trpa1; Pomc; Cgrp; Crf; Pka; Era; Nr2b; TRPM5; Prkaca; Prkacb; Prkar1a; Prkar2a
Mitochondrial Function	AIF; CytC; SMAC (Diablo); Aifm-1; Aifm-2
Developmental Neurology	BMP-4; Chordin (Chrd); Noggin (Nog); WNT (Wnt2; Wnt2b; Wnt3a; Wnt4; Wnt5a; Wnt6; Wnt7b; Wnt8b; Wnt9a; Wnt9b; Wnt10a; Wnt10b; Wnt16); beta-catenin; Dkk-1; Frizzled related proteins; Otx-2; Gbx2; FGF-8; Reelin; Dab1; unc-86 (Pou4f1 or Brn3a); Numb; Reln

[0162] Embodiments of the invention also relate to methods and compositions related to knocking out genes, amplifying genes and repairing particular mutations associated with DNA repeat instability and neurological disorders (Robert D. Wells, Tetsuo Ashizawa, *Genetic Instabilities and Neurological Diseases*, Second Edition, Academic Press, Oct. 13, 2011—Medical). Specific aspects of tandem repeat sequences have been found to be responsible for more than twenty human diseases (New insights into repeat instability: role of RNA-DNA hybrids. McIvor EI, Polak U, Napierala M. *RNA Biol.* 2010 September-October; 7(5):551-8). The CRISPR-Cas system may be harnessed to correct these defects of genomic instability.

[0163] A further aspect of the invention relates to utilizing the CRISPR-Cas system for correcting defects in the EMP2A and EMP2B genes that have been identified to be associated with Lafora disease. Lafora disease is an autosomal recessive condition which is characterized by progressive myoclonus epilepsy which may start as epileptic seizures in adolescence. A few cases of the disease may be caused by mutations in genes yet to be identified. The disease causes seizures, muscle spasms, difficulty walking, dementia, and eventually death. There is currently no therapy that has proven effective against disease progression. Other genetic abnormalities associated with epilepsy may also be targeted by the CRISPR-Cas system and the underlying genetics is further described in *Genetics of Epilepsy and Genetic Epilepsies*, edited by Giuliano Avanzini, Jeffrey L. Noebels, Mariani Foundation Paediatric Neurology;20; 2009).

[0164] In yet another aspect of the invention, the CRISPR-Cas system may be used to correct ocular defects that arise from several genetic mutations further described in *Genetic Diseases of the Eye*, Second Edition, edited by Elias I. Traboulsi, Oxford University Press, 2012.

[0165] Several further aspects of the invention relate to correcting defects associated with a wide range of genetic diseases which are further described on the website of the National Institutes of Health under the topic subsection Genetic Disorders (website at health.nih.gov/topic/GeneticDisorders). The genetic brain diseases may include but are not limited to Adrenoleukodystrophy, Agenesis of the Corpus Callosum, Aicardi Syndrome, Alpers' Disease, Alzheimer's Disease, Barth Syndrome, Batten Disease, CADASIL, Cerebellar Degeneration, Fabry's Disease, Gerstmann-Straussler-Scheinker Disease, Huntington's Disease and other Triplet Repeat Disorders, Leigh's Disease, Lesch-Nyhan Syndrome, Menkes Disease, Mitochondrial Myopathies and NINDS Colpocephaly. These diseases are further described on the website of the National Institutes of Health under the subsection Genetic Brain Disorders.

[0166] In some embodiments, the condition may be neoplasia. In some embodiments, where the condition is neoplasia, the genes to be targeted are any of those listed in Table A (in this case PTEN and so forth). In some embodiments, the condition may be Age-related Macular Degeneration. In some embodiments, the condition may be a Schizophrenic Disorder. In some embodiments, the condition may be a Trinucleotide Repeat Disorder. In some embodiments, the condition may be Fragile X Syndrome. In some embodiments, the condition may be a Secretase Related Disorder. In some embodiments, the condition may be a Prion-related disorder. In some embodiments, the condition may be ALS. In some embodiments, the condition

may be a drug addiction. In some embodiments, the condition may be Autism. In some embodiments, the condition may be Alzheimer's Disease. In some embodiments, the condition may be inflammation. In some embodiments, the condition may be Parkinson's Disease.

[0167] Examples of proteins associated with Parkinson's disease include but are not limited to α -synuclein, DJ-1, LRRK2, PINK1, Parkin, UCHL1, Synphilin-1, and NURR1.

[0168] Examples of addiction-related proteins may include ABAT for example.

[0169] Examples of inflammation-related proteins may include the monocyte chemoattractant protein-1 (MCP1) encoded by the *Ccr2* gene, the C-C chemokine receptor type 5 (CCRS) encoded by the *Ccr5* gene, the IgG receptor IIB (FCGR2b, also termed CD32) encoded by the *Fcgr2b* gene, or the Fc epsilon R1g (FCER1g) protein encoded by the *Fcer1g* gene, for example.

[0170] Examples of cardiovascular diseases associated proteins may include IL1B (interleukin 1, beta), XDH (xanthine dehydrogenase), TP53 (tumor protein p53), PTGIS (prostaglandin I2 (prostacyclin) synthase), MB (myoglobin), IL4 (interleukin 4), ANGPT1 (angiopoietin 1), ABCG8 (ATP-binding cassette, sub-family G (WHITE), member 8), or CTSK (cathepsin K), for example.

[0171] Examples of Alzheimer's disease associated proteins may include the very low density lipoprotein receptor protein (VLDLR) encoded by the *VLDLR* gene, the ubiquitin-like modifier activating enzyme 1 (UBA1) encoded by the *UBA1* gene, or the NEDD8-activating enzyme E1 catalytic subunit protein (UBE1C) encoded by the *UBA3* gene, for example.

[0172] Examples of proteins associated Autism Spectrum Disorder may include the benzodiazapine receptor (peripheral) associated protein 1 (BZRAP1) encoded by the *BZRAP1* gene, the AF4/FMR2 family member 2 protein (AFF2) encoded by the *AFF2* gene (also termed MFR2), the fragile X mental retardation autosomal homolog 1 protein (FXR1) encoded by the *FXR1* gene, or the fragile X mental retardation autosomal homolog 2 protein (FXR2) encoded by the *FXR2* gene, for example.

[0173] Examples of proteins associated Macular Degeneration may include the ATP-binding cassette, sub-family A (ABC1) member 4 protein (ABCA4) encoded by the *ABCR* gene, the apolipoprotein E protein (APOE) encoded by the *APOE* gene, or the chemokine (C-C motif) Ligand 2 protein (CCL2) encoded by the *CCL2* gene, for example.

[0174] Examples of proteins associated Schizophrenia may include NRG1, ErbB4, CPLX1, TPH1, TPH2, NRXN1, GSK3A, BDNF, DISC1, GSK3B, and combinations thereof.

[0175] Examples of proteins involved in tumor suppression may include ATM (ataxia telangiectasia mutated), ATR (ataxia telangiectasia and Rad3 related), EGFR (epidermal growth factor receptor), ERBB2 (v-erb-b2 erythroblastic leukemia viral oncogene homolog 2), ERBB3 (v-erb-b2 erythroblastic leukemia viral oncogene homolog 3), ERBB4 (v-erb-b2 erythroblastic leukemia viral oncogene homolog 4), Notch 1, Notch2, Notch 3, or Notch 4, for example.

[0176] Examples of proteins associated with a secretase disorder may include PSENEN (presenilin enhancer 2 homolog (*C. elegans*)), CTSB (cathepsin B), PSEN1 (presenilin 1), APP (amyloid beta (A4) precursor protein), APH1B (anterior pharynx defective 1 homolog B (*C. elegans*)), PSEN2 (presenilin 2 (Alzheimer disease 4)), or BACE1 (beta-site APP-cleaving enzyme 1), for example.

[0177] Examples of proteins associated with Amyotrophic Lateral Sclerosis may include SOD1 (superoxide dismutase 1), ALS2 (amyotrophic lateral sclerosis 2), FUS (fused in sarcoma), TARDBP (TAR DNA binding protein), VAGFA (vascular endothelial growth factor A), VAGFB (vascular endothelial growth factor B), and VAGFC (vascular endothelial growth factor C), and any combination thereof.

[0178] Examples of proteins associated with prion diseases may include SOD1 (superoxide dismutase 1), ALS2 (amyotrophic lateral sclerosis 2), FUS (fused in sarcoma), TARDBP (TAR DNA binding protein), VAGFA (vascular endothelial growth factor A), VAGFB (vascular endothelial growth factor B), and VAGFC (vascular endothelial growth factor C), and any combination thereof.

[0179] Examples of proteins related to neurodegenerative conditions in prion disorders may include A2M (Alpha-2-Macroglobulin), AATF (Apoptosis antagonizing transcription factor), ACPP (Acid phosphatase prostate), ACTA2 (Actin alpha 2 smooth muscle aorta), ADAM22 (ADAM metallopeptidase domain), ADORA3 (Adenosine A3 receptor), or ADRA1D (Alpha-1D adrenergic receptor for Alpha-1D adrenoceptor), for example.

[0180] Examples of proteins associated with Immunodeficiency may include A2M [alpha-2-macroglobulin]; AANAT [arylalkylamine N-acetyltransferase]; ABCA1 [ATP-binding cassette, sub-family A (ABC1), member 1]; ABCA2 [ATP-binding cassette, sub-family A (ABC1), member 2]; or ABCA3 [ATP-binding cassette, sub-family A (ABC1), member 3]; for example.

[0181] Examples of proteins associated with Trinucleotide Repeat Disorders include AR (androgen receptor), FMR1 (fragile X mental retardation 1), HTT (huntingtin), or DMPK (dystrophia myotonica-protein kinase), FXN (frataxin), ATXN2 (ataxin 2), for example.

[0182] Examples of proteins associated with Neurotransmission Disorders include SST (somatostatin), NOS1 (nitric oxide synthase 1 (neuronal)), ADRA2A (adrenergic, alpha-2A-, receptor), ADRA2C (adrenergic, alpha-2C-, receptor), TACR1 (tachykinin receptor 1), or HTR2c (5-hydroxytryptamine (serotonin) receptor 2C), for example.

[0183] Examples of neurodevelopmental-associated sequences include A2BP1 [ataxin 2-binding protein 1], AADAT [aminoadipate aminotransferase], AANAT [arylalkylamine N-acetyltransferase], ABAT [4-aminobutyrate aminotransferase], ABCA1 [ATP-binding cassette, sub-family A (ABC1), member 1], or ABCA13 [ATP-binding cassette, sub-family A (ABC1), member 13], for example.

[0184] Further examples of preferred conditions treatable with the present system include may be selected from: Aicardi-Goutières Syndrome; Alexander Disease; Allan-Herndon-Dudley Syndrome; POLG-Related Disorders; Alpha-Mannosidosis (Type II and III); Alström Syndrome; Angelman; Syndrome; Ataxia-Telangiectasia; Neuronal Ceroid-Lipofuscinoses; Beta-Thalassemia; Bilateral Optic Atrophy and (Infantile) Optic Atrophy Type 1; Retinoblastoma (bilateral); Canavan Disease; Cerebrooculofacioskeletal Syndrome 1 [COF S1]; Cerebrotendinous Xanthomatosis; Cornelia de Lange Syndrome; MAPT-Related Disorders; Genetic Prion Diseases; Dravet Syndrome; Early-Onset Familial Alzheimer Disease; Friedrich Ataxia [FRDA]; Frys Syndrome; Fucosidosis; Fukuyama Congenital Muscular Dystrophy; Galactosialidosis; Gaucher Disease; Organic Acidemias; Hemophagocytic Lymphohistiocytosis; Hutchinson-Gilford Progeria Syndrome; Mucopolysaccharidosis II; Infantile Free Sialic Acid Storage Disease; PLA2G6-Associated Neurodegeneration; Jervell and Lange-Nielsen Syndrome; Junctional Epidermolysis Bullosa; Huntington Disease; Krabbe Disease (Infantile); Mitochondrial DNA-Associated Leigh Syndrome and NARP; Lesch-Nyhan Syndrome; LIS1-Associated Lissencephaly; Lowe Syndrome; Maple Syrup Urine Disease; MECP2 Duplication Syndrome; ATP7A-Related Copper Transport Disorders; LAMA2-Related Muscular Dystrophy; Arylsulfatase A Deficiency; Mucopolysaccharidosis Types I, II or III; Peroxisome Biogenesis Disorders, Zellweger Syndrome Spectrum; Neurodegeneration with Brain Iron Accumulation Disorders; Acid Sphingomyelinase Deficiency; Niemann-Pick Disease Type C; Glycine Encephalopathy; ARX-Related Disorders; Urea Cycle Disorders; COL1A1/2-Related Osteogenesis Imperfecta; Mitochondrial DNA Deletion Syndromes; PLP1-Related Disorders; Perry Syndrome; Phelan-McDermid Syndrome; Glycogen Storage Disease Type II (Pompe Disease) (Infantile); MAPT-Related Disorders; MECP2-Related Disorders; Rhizomelic Chondrodysplasia Punctata Type 1; Roberts Syndrome; Sandhoff Disease; Schindler Disease—Type 1; Adenosine Deaminase Deficiency; Smith-Lemli-Opitz Syndrome; Spinal Muscular Atrophy; Infantile-Onset Spinocerebellar Ataxia; Hexosaminidase A Deficiency; Thanatophoric Dysplasia Type 1; Collagen Type VI-Related Disorders; Usher Syndrome Type I; Congenital Muscular Dystrophy; Wolf-Hirschhorn Syndrome; Lysosomal Acid Lipase Deficiency; and Xeroderma Pigmentosum.

lipidosis II; Infantile Free Sialic Acid Storage Disease; PLA2G6-Associated Neurodegeneration; Jervell and Lange-Nielsen Syndrome; Junctional Epidermolysis Bullosa; Huntington Disease; Krabbe Disease (Infantile); Mitochondrial DNA-Associated Leigh Syndrome and NARP; Lesch-Nyhan Syndrome; LIS1-Associated Lissencephaly; Lowe Syndrome; Maple Syrup Urine Disease; MECP2 Duplication Syndrome; ATP7A-Related Copper Transport Disorders; LAMA2-Related Muscular Dystrophy; Arylsulfatase A Deficiency; Mucopolysaccharidosis Types I, II or III; Peroxisome Biogenesis Disorders, Zellweger Syndrome Spectrum; Neurodegeneration with Brain Iron Accumulation Disorders; Acid Sphingomyelinase Deficiency; Niemann-Pick Disease Type C; Glycine Encephalopathy; ARX-Related Disorders; Urea Cycle Disorders; COL1A1/2-Related Osteogenesis Imperfecta; Mitochondrial DNA Deletion Syndromes; PLP1-Related Disorders; Perry Syndrome; Phelan-McDermid Syndrome; Glycogen Storage Disease Type II (Pompe Disease) (Infantile); MAPT-Related Disorders; MECP2-Related Disorders; Rhizomelic Chondrodysplasia Punctata Type 1; Roberts Syndrome; Sandhoff Disease; Schindler Disease—Type 1; Adenosine Deaminase Deficiency; Smith-Lemli-Opitz Syndrome; Spinal Muscular Atrophy; Infantile-Onset Spinocerebellar Ataxia; Hexosaminidase A Deficiency; Thanatophoric Dysplasia Type 1; Collagen Type VI-Related Disorders; Usher Syndrome Type I; Congenital Muscular Dystrophy; Wolf-Hirschhorn Syndrome; Lysosomal Acid Lipase Deficiency; and Xeroderma Pigmentosum.

[0185] As will be apparent, it is envisaged that the present system can be used to target any polynucleotide sequence of interest. Some examples of conditions or diseases that might be usefully treated using the present system are included in the Tables above and examples of genes currently associated with those conditions are also provided there. However, the genes exemplified are not exhaustive.

EXAMPLES

[0186] The following examples are given for the purpose of illustrating various embodiments of the invention and are not meant to limit the present invention in any fashion. The present examples, along with the methods described herein are presently representative of preferred embodiments, are exemplary, and are not intended as limitations on the scope of the invention. Changes therein and other uses which are encompassed within the spirit of the invention as defined by the scope of the claims will occur to those skilled in the art.

Example 1

CRISPR Complex Activity in the Nucleus of a Eukaryotic Cell

[0187] An example type II CRISPR system is the type II CRISPR locus from *Streptococcus pyogenes* SF370, which contains a cluster of four genes Cas9, Cas1, Cas2, and Csn1, as well as two non-coding RNA elements, tracrRNA and a characteristic array of repetitive sequences (direct repeats) interspaced by short stretches of non-repetitive sequences (spacers, about 30 bp each). In this system, targeted DNA double-strand break (DSB) is generated in four sequential steps (FIG. 2A). First, two non-coding RNAs, the pre-crRNA array and tracrRNA, are transcribed from the CRISPR locus. Second, tracrRNA hybridizes to the direct

repeats of pre-crRNA, which is then processed into mature crRNAs containing individual spacer sequences. Third, the mature crRNA:tracrRNA complex directs Cas9 to the DNA target consisting of the protospacer and the corresponding PAM via heteroduplex formation between the spacer region of the crRNA and the protospacer DNA. Finally, Cas9 mediates cleavage of target DNA upstream of PAM to create a DSB within the protospacer (FIG. 2A). This example describes an example process for adapting this RNA-programmable nuclease system to direct CRISPR complex activity in the nuclei of eukaryotic cells.

[0188] Cell Culture and Transfection

[0189] Human embryonic kidney (HEK) cell line HEK 293FT (Life Technologies) was maintained in Dulbecco's modified Eagle's Medium (DMEM) supplemented with 10% fetal bovine serum (HyClone), 2 mM GlutaMAX (Life Technologies), 100 U/mL penicillin, and 100 µg/mL streptomycin at 37° C. with 5% CO₂ incubation. Mouse neuro2A (N2A) cell line (ATCC) was maintained with DMEM supplemented with 5% fetal bovine serum (HyClone), 2 mM GlutaMAX (Life Technologies), 100 U/mL penicillin, and 100 µg/mL streptomycin at 37° C. with 5% CO₂.

[0190] HEK 293FT or N2A cells were seeded into 24-well plates (Corning) one day prior to transfection at a density of 200,000 cells per well. Cells were transfected using Lipofectamine 2000 (Life Technologies) following the manufacturer's recommended protocol. For each well of a 24-well plate a total of 800 ng of plasmids were used.

[0191] Surveyor Assay and Sequencing Analysis for Genome Modification

[0192] HEK 293FT or N2A cells were transfected with plasmid DNA as described above. After transfection, the cells were incubated at 37° C. for 72 hours before genomic DNA extraction. Genomic DNA was extracted using the QuickExtract DNA extraction kit (Epicentre) following the manufacturer's protocol. Briefly, cells were resuspended in QuickExtract solution and incubated at 65° C. for 15 minutes and 98° C. for 10 minutes. Extracted genomic DNA was immediately processed or stored at -20° C.

[0193] The genomic region surrounding a CRISPR target site for each gene was PCR amplified, and products were purified using QiaQuick Spin Column (Qiagen) following manufacturer's protocol. A total of 400 ng of the purified PCR products were mixed with 2 µl 10× Taq polymerase PCR buffer (Enzymatics) and ultrapure water to a final volume of 20 µl, and subjected to a re-annealing process to enable heteroduplex formation: 95° C. for 10 min, 95° C. to 85° C. ramping at -2° C./s, 85° C. to 25° C. at -0.25° C/s, and 25° C. hold for 1 minute. After re-annealing, products were treated with Surveyor nuclease and Surveyor enhancer S (Transgenomics) following the manufacturer's recommended protocol, and analyzed on 4-20% Novex TBE poly-acrylamide gels (Life Technologies). Gels were stained with SYBR Gold DNA stain (Life Technologies) for 30 minutes and imaged with a Gel Doc gel imaging system (Bio-rad). Quantification was based on relative band intensities, as a measure of the fraction of cleaved DNA. FIG. 8 provides a schematic illustration of this Surveyor assay.

[0194] Restriction fragment length polymorphism assay for detection of homologous recombination

[0195] HEK 293FT and N2A cells were transfected with plasmid DNA, and incubated at 37° C. for 72 hours before genomic DNA extraction as described above. The target genomic region was PCR amplified using primers outside

the homology arms of the homologous recombination (HR) template. PCR products were separated on a 1% agarose gel and extracted with MinElute GelExtraction Kit (Qiagen). Purified products were digested with HindIII (Fermentas) and analyzed on a 6% Novex TBE poly-acrylamide gel (Life Technologies).

[0196] RNA Secondary Structure Prediction and Analysis

[0197] RNA secondary structure prediction was performed using the online webserver RNAfold developed at Institute for Theoretical Chemistry at the University of Vienna, using the centroid structure prediction algorithm (see e.g. A. R. Gruber et al., 2008, Cell 106(1): 23-24; and P A Carr and G M Church, 2009, Nature Biotechnology 27(12): 1151-62).

[0198] Bacterial Plasmid Transformation Interference Assay

[0199] Elements of the *S. pyogenes* CRISPR locus 1 sufficient for CRISPR activity were reconstituted in *E. coli* using pCRISPR plasmid (schematically illustrated in FIG. 10A). pCRISPR contained tracrRNA, SpCas9, and a leader sequence driving the crRNA array. Spacers (also referred to as "guide sequences") were inserted into the crRNA array between BsaI sites using annealed oligonucleotides, as illustrated. Challenge plasmids used in the interference assay were constructed by inserting the protospacer (also referred to as a "target sequence") sequence along with an adjacent CRISPR motif sequence (PAM) into pUC19 (see FIG. 10B). The challenge plasmid contained ampicillin resistance. FIG. 10C provides a schematic representation of the interference assay. Chemically competent *E. coli* strains already carrying pCRISPR and the appropriate spacer were transformed with the challenge plasmid containing the corresponding protospacer-PAM sequence. pUC19 was used to assess the transformation efficiency of each pCRISPR-carrying competent strain. CRISPR activity resulted in cleavage of the pPSP plasmid carrying the protospacer, precluding ampicillin resistance otherwise conferred by pUC19 lacking the protospacer. FIG. 10D illustrates competence of each pCRISPR-carrying *E. coli* strain used in assays illustrated in FIG. 4C.

[0200] RNA Purification

[0201] HEK 293FT cells were maintained and transfected as stated above. Cells were harvested by trypsinization followed by washing in phosphate buffered saline (PBS). Total cell RNA was extracted with TRI reagent (Sigma) following manufacturer's protocol. Extracted total RNA was quantified using Naonodrop (Thermo Scientific) and normalized to same concentration.

[0202] Northern Blot Analysis of crRNA and tracrRNA Expression in Mammalian Cells

[0203] RNAs were mixed with equal volumes of 2× loading buffer (Ambion), heated to 95° C. for 5 min, chilled on ice for 1 min, and then loaded onto 8% denaturing polyacrylamide gels (SequaGel, National Diagnostics) after pre-running the gel for at least 30 minutes. The samples were electrophoresed for 1.5 hours at 40W limit. Afterwards, the RNA was transferred to Hybond N+ membrane (GE Healthcare) at 300 mA in a semi-dry transfer apparatus (Bio-rad) at room temperature for 1.5 hours. The RNA was crosslinked to the membrane using autocrosslink button on Stratagene UV Crosslinker the Stratalink (Stratagene). The membrane was pre-hybridized in ULTRAhyb-Oligo Hybridization Buffer (Ambion) for 30 min with rotation at 42° C., and probes were then added and hybridized overnight. Probes

were ordered from IDT and labeled with [γ - 32 P] ATP (Perkin Elmer) with T4 polynucleotide kinase (New England Biolabs). The membrane was washed once with pre-warmed (42° C.) 2xSSC, 0.5% SDS for 1 min followed by two 30 minute washes at 42° C. The membrane was exposed to a phosphor screen for one hour or overnight at room temperature and then scanned with a phosphorimager (Typhoon).

[0204] Bacterial CRISPR System Construction and Evaluation

[0205] CRISPR locus elements, including tracrRNA, Cas9, and leader were PCR amplified from *Streptococcus pyogenes* SF370 genomic DNA with flanking homology arms for Gibson Assembly. Two BsaI type IIS sites were introduced in between two direct repeats to facilitate easy insertion of spacers (FIG. 9). PCR products were cloned into EcoRV-digested pACYC184 downstream of the tet promoter using Gibson Assembly Master Mix (NEB). Other endogenous CRISPR system elements were omitted, with the exception of the last 50 bp of Csn2. Oligos (Integrated DNA Technology) encoding spacers with complimentary overhangs were cloned into the BsaI-digested vector pDCC000 (NEB) and then ligated with T7 ligase (Enzymatics) to generate pCRISPR plasmids. Challenge plasmids containing spacers with PAM sequences (also referred to herein as “CRISPR motif sequences”) were created by ligating hybridized oligos carrying compatible overhangs (Integrated DNA Technology) into BamHI-digested pUC19. Cloning for all constructs was performed in *E. coli* strain JM109 (Zymo Research).

[0206] pCRISPR-carrying cells were made competent using the Z-Competent *E. coli* Transformation Kit and Buffer Set (Zymo Research, T3001) according to manufacturer’s instructions. In the transformation assay, 50 μ L aliquots of competent cells carrying pCRISPR were thawed on ice and transformed with 1 ng of spacer plasmid or pUC19 on ice for 30 minutes, followed by 45 second heat shock at 42° C. and 2 minutes on ice. Subsequently, 250 μ L SOC (Invitrogen) was added followed by shaking incubation at 37° C. for 1 hr, and 100 μ L of the post-SOC outgrowth was plated onto double selection plates (12.5 μ g/ml chloramphenicol, 100 μ g/ml ampicillin). To obtain cfu/ng of DNA, total colony numbers were multiplied by 3.

[0207] To improve expression of CRISPR components in mammalian cells, two genes from the SF370 locus 1 of *Streptococcus pyogenes* (*S. pyogenes*) were codon-optimized, Cas9 (SpCas9) and RNase III (SpRNase III). To facilitate nuclear localization, a nuclear localization signal (NLS) was included at the amino (N)- or carboxyl (C)-termini of both SpCas9 and SpRNase III (FIG. 2B). To facilitate visualization of protein expression, a fluorescent protein marker was also included at the N- or C-termini of both proteins (FIG. 2B). A version of SpCas9 with an NLS attached to both N- and C-termini (2xNLS-SpCas9) was also generated. Constructs containing NLS-fused SpCas9 and SpRNase III were transfected into 293FT human embryonic kidney (HEK) cells, and the relative positioning of the NLS to SpCas9 and SpRNase III was found to affect their nuclear localization efficiency. Whereas the C-terminal NLS was sufficient to target SpRNase III to the nucleus, attachment of a single copy of these particular NLS’s to either the N- or C-terminus of SpCas9 was unable to achieve adequate nuclear localization in this system. In this example, the C-terminal NLS was that of nucleoplasmin (KRPAATKK-

AGQAKKKK (SEQ ID NO: 2)), and the C-terminal NLS was that of the SV40 large T-antigen (PKKKRKV (SEQ ID NO: 1)). Of the versions of SpCas9 tested, only 2xNLS-SpCas9 exhibited nuclear localization (FIG. 2B).

[0208] The tracrRNA from the CRISPR locus of *S. pyogenes* SF370 has two transcriptional start sites, giving rise to two transcripts of 89-nucleotides (nt) and 171nt that are subsequently processed into identical 75nt mature tracrRNAs. The shorter 89nt tracrRNA was selected for expression in mammalian cells (expression constructs illustrated in FIG. 7A, with functionality as determined by results of the Surveyor assay shown in FIG. 7B). Transcription start sites are marked as +1, and transcription terminator and the sequence probed by northern blot are also indicated. Expression of processed tracrRNA was also confirmed by Northern blot. FIG. 7C shows results of a Northern blot analysis of total RNA extracted from 293FT cells transfected with U6 expression constructs carrying long or short tracrRNA, as well as SpCas9 and DR-EMX1(1)-DR. Left and right panels are from 293FT cells transfected without or with SpRNase III, respectively. U6 indicate loading control blotted with a probe targeting human U6 snRNA. Transfection of the short tracrRNA expression construct led to abundant levels of the processed form of tracrRNA (~75bp). Very low amounts of long tracrRNA are detected on the Northern blot.

[0209] To promote precise transcriptional initiation, the RNA polymerase III-based U6 promoter was selected to drive the expression of tracrRNA (FIG. 2C). Similarly, a U6 promoter-based construct was developed to express a pre-crRNA array consisting of a single spacer flanked by two direct repeats (DRs, also encompassed by the term “tracr-mate sequences”; FIG. 2C). The initial spacer was designed to target a 33-base-pair (bp) target site (30-bp protospacer plus a 3-bp CRISPR motif (PAM) sequence satisfying the NGG recognition motif of Cas9) in the human EMX1 locus (FIG. 2C), a key gene in the development of the cerebral cortex.

[0210] To test whether heterologous expression of the CRISPR system (SpCas9, SpRNase III, tracrRNA, and pre-crRNA) in mammalian cells can achieve targeted cleavage of mammalian chromosomes, HEK 293FT cells were transfected with combinations of CRISPR components. Since DSBs in mammalian nuclei are partially repaired by the non-homologous end joining (NHEJ) pathway, which leads to the formation of indels, the Surveyor assay was used to detect potential cleavage activity at the target EMX1 locus (FIG. 8) (see e.g. Guschin et al., 2010, Methods Mol Biol 649: 247). Co-transfection of all four CRISPR components was able to induce up to 5.0% cleavage in the protospacer (see FIG. 2D). Co-transfection of all CRISPR components minus SpRNase III also induced up to 4.7% indel in the protospacer, suggesting that there may be endogenous mammalian RNases that are capable of assisting with crRNA maturation, such as for example the related Dicer and Drosha enzymes. Removing any of the remaining three components abolished the genome cleavage activity of the CRISPR system (FIG. 2D). Sanger sequencing of amplicons containing the target locus verified the cleavage activity: in 43 sequenced clones, 5 mutated alleles (11.6%) were found. Similar experiments using a variety of guide sequences produced indel percentages as high as 29% (see FIGS. 4-7, 12, and 13). These results define a three-component system for efficient CRISPR-mediated genome modification in mammalian cells. To optimize the cleavage efficiency,

Applicants also tested whether different isoforms of tracrRNA affected the cleavage efficiency and found that, in this example system, only the short (89-bp) transcript form was able to mediate cleavage of the human EMX1 genomic locus (FIG. 7B).

[0211] FIG. 14 provides an additional Northern blot analysis of crRNA processing in mammalian cells. FIG. 14A illustrates a schematic showing the expression vector for a single spacer flanked by two direct repeats (DR-EMX1(1)-DR). The 30 bp spacer targeting the human EMX1 locus protospacer 1 (see FIG. 6) and the direct repeat sequences are shown in the sequence beneath FIG. 14A. The line indicates the region whose reverse-complement sequence was used to generate Northern blot probes for EMX1(1) crRNA detection. FIG. 14B shows a Northern blot analysis of total RNA extracted from 293FT cells transfected with U6 expression constructs carrying DR-EMX1(1)-DR. Left and right panels are from 293FT cells transfected without or with SpRNase III respectively. DR-EMX1(1)-DR was processed into mature crRNAs only in the presence of SpCas9 and short tracrRNA and was not dependent on the presence of SpRNase III. The mature crRNA detected from transfected 293FT total RNA is ~33 bp and is shorter than the 39-42 bp mature crRNA from *S. pyogenes*. These results demonstrate that a CRISPR system can be transplanted into eukaryotic cells and reprogrammed to facilitate cleavage of endogenous mammalian target polynucleotides.

[0212] FIG. 2 illustrates the bacterial CRISPR system described in this example. FIG. 2A illustrates a schematic showing the CRISPR locus 1 from *Streptococcus pyogenes* SF370 and a proposed mechanism of CRISPR-mediated DNA cleavage by this system. Mature crRNA processed from the direct repeat-spacer array directs Cas9 to genomic targets consisting of complimentary protospacers and a protospacer-adjacent motif (PAM). Upon target-spacer base pairing, Cas9 mediates a double-strand break in the target DNA. FIG. 2B illustrates engineering of *S. pyogenes* Cas9 (SpCas9) and RNase III (SpRNase III) with nuclear localization signals (NLs) to enable import into the mammalian nucleus. FIG. 2C illustrates mammalian expression of SpCas9 and SpRNase III driven by the constitutive EF1 α promoter and tracrRNA and pre-crRNA array (DR-Spacer-DR) driven by the RNA Pol3 promoter U6 to promote precise transcription initiation and termination. A protospacer from the human EMX1 locus with a satisfactory PAM sequence is used as the spacer in the pre-crRNA array. FIG. 2D illustrates surveyor nuclease assay for SpCas9-mediated minor insertions and deletions. SpCas9 was expressed with and without SpRNase III, tracrRNA, and a pre-crRNA array carrying the EMX1-target spacer. FIG. 2E illustrates a schematic representation of base pairing between target locus and EMX1-targeting crRNA, as well as an example chromatogram showing a micro deletion adjacent to the SpCas9 cleavage site. FIG. 2F illustrates mutated alleles identified from sequencing analysis of 43 clonal amplicons showing a variety of micro insertions and deletions. Dashes indicate deleted bases, and non-aligned or mismatched bases indicate insertions or mutations. Scale bar=10 μ m.

[0213] To further simplify the three-component system, a chimeric crRNA-tracrRNA hybrid design was adapted, where a mature crRNA (comprising a guide sequence) is fused to a partial tracrRNA via a stem-loop to mimic the natural crRNA:tracrRNA duplex (FIG. 3A). To increase co-delivery efficiency, a bicistronic expression vector was

created to drive co-expression of a chimeric RNA and SpCas9 in transfected cells (FIGS. 3A and 8). In parallel, the bicistronic vectors were used to express a pre-crRNA (DR-guide sequence-DR) with SpCas9, to induce processing into crRNA with a separately expressed tracrRNA (compare FIG. 13B top and bottom). FIG. 9 provides schematic illustrations of bicistronic expression vectors for pre-crRNA array (FIG. 9A) or chimeric crRNA (represented by the short line downstream of the guide sequence insertion site and upstream of the EF1 α promoter in FIG. 9B) with hSpCas9, showing location of various elements and the point of guide sequence insertion. The expanded sequence around the location of the guide sequence insertion site in FIG. 9B also shows a partial DR sequence (GTTTTAGAGCTA (SEQ ID NO: 27)) and a partial tracrRNA sequence (TAGCAAGT-TAAAATAAGGCTAGTCCGTTTTT (SEQ ID NO: 28)). Guide sequences can be inserted between BbsI sites using annealed oligonucleotides. Sequence design for the oligonucleotides are shown below the schematic illustrations in FIG. 9, with appropriate ligation adapters indicated. WPRE represents the Woodchuck hepatitis virus post-transcriptional regulatory element. The efficiency of chimeric RNA-mediated cleavage was tested by targeting the same EMX1 locus described above. Using both Surveyor assay and Sanger sequencing of amplicons, Applicants confirmed that the chimeric RNA design facilitates cleavage of human EMX1 locus with approximately a 4.7% modification rate (FIG. 4).

[0214] Generalizability of CRISPR-mediated cleavage in eukaryotic cells was tested by targeting additional genomic loci in both human and mouse cells by designing chimeric RNA targeting multiple sites in the human EMX1 and PVALB, as well as the mouse Th loci. FIG. 15 illustrates the selection of some additional targeted protospacers in human PVALB (FIG. 15A) and mouse Th (FIG. 15B) loci. Schematics of the gene loci and the location of three protospacers within the last exon of each are provided. The underlined sequences include 30 bp of protospacer sequence and 3 bp at the 3' end corresponding to the PAM sequences. Protospacers on the sense and anti-sense strands are indicated above and below the DNA sequences, respectively. A modification rate of 6.3% and 0.75% was achieved for the human PVALB and mouse Th loci respectively, demonstrating the broad applicability of the CRISPR system in modifying different loci across multiple organisms (FIGS. 3B and 6). While cleavage was only detected with one out of three spacers for each locus using the chimeric constructs, all target sequences were cleaved with efficiency of indel production reaching 27% when using the co-expressed pre-crRNA arrangement (FIG. 6).

[0215] FIG. 13 provides a further illustration that SpCas9 can be reprogrammed to target multiple genomic loci in mammalian cells. FIG. 13A provides a schematic of the human EMX1 locus showing the location of five protospacers, indicated by the underlined sequences. FIG. 13B provides a schematic of the pre-crRNA/tracrRNA complex showing hybridization between the direct repeat region of the pre-crRNA and tracrRNA (top), and a schematic of a chimeric RNA design comprising a 20 bp guide sequence, and tracr mate and tracr sequences consisting of partial direct repeat and tracrRNA sequences hybridized in a hairpin structure (bottom). Results of a Surveyor assay comparing the efficacy of Cas9-mediated cleavage at five protospacers in the human EMX1 locus is illustrated in FIG. 13C.

Each protospacer is targeted using either processed pre-crRNA/tracrRNA complex (crRNA) or chimeric RNA (chiRNA).

[0216] Since the secondary structure of RNA can be crucial for intermolecular interactions, a structure prediction algorithm based on minimum free energy and Boltzmann-weighted structure ensemble was used to compare the putative secondary structure of all guide sequences used in our genome targeting experiment (FIG. 3B) (see e.g. Gruber et al., 2008, *Nucleic Acids Research*, 36: W70). Analysis revealed that in most cases, the effective guide sequences in the chimeric crRNA context were substantially free of secondary structure motifs, whereas the ineffective guide sequences were more likely to form internal secondary structures that could prevent base pairing with the target protospacer DNA. It is thus possible that variability in the spacer secondary structure might impact the efficiency of CRISPR-mediated interference when using a chimeric crRNA.

[0217] FIG. 3 illustrates example expression vectors. FIG. 3A provides a schematic of a bi-cistronic vector for driving the expression of a synthetic crRNA-tracrRNA chimera (chimeric RNA) as well as SpCas9. The chimeric guide RNA contains a 20-bp guide sequence corresponding to the protospacer in the genomic target site. FIG. 3B provides a schematic showing guide sequences targeting the human EMX1, PVALB, and mouse Th loci, as well as their predicted secondary structures. The modification efficiency at each target site is indicated below the RNA secondary structure drawing (EMX1, n=216 amplicon sequencing reads; PVALB, n=224 reads; Th, n=265 reads). The folding algorithm produced an output with each base colored according to its probability of assuming the predicted secondary structure, as indicated by a rainbow scale that is reproduced in FIG. 3B in gray scale. Further vector designs for SpCas9 are shown in FIG. 4A, which illustrates single expression vectors incorporating a U6 promoter linked to an insertion site for a guide oligo, and a Cbh promoter linked to SpCas9 coding sequence. The vector shown in FIG. 4B includes a tracrRNA coding sequence linked to an H1 promoter.

[0218] To test whether spacers containing secondary structures are able to function in prokaryotic cells where CRISPRs naturally operate, transformation interference of protospacer-bearing plasmids were tested in an *E. coli* strain heterologously expressing the *S. pyogenes* SF370 CRISPR locus 1 (FIG. 10). The CRISPR locus was cloned into a low-copy *E. coli* expression vector and the crRNA array was replaced with a single spacer flanked by a pair of DRs (pCRISPR). *E. coli* strains harboring different pCRISPR plasmids were transformed with challenge plasmids containing the corresponding protospacer and PAM sequences (FIG. 10C). In the bacterial assay, all spacers facilitated efficient CRISPR interference (FIG. 4C). These results suggest that there may be additional factors affecting the efficiency of CRISPR activity in mammalian cells.

[0219] To investigate the specificity of CRISPR-mediated cleavage, the effect of single-nucleotide mutations in the guide sequence on protospacer cleavage in the mammalian genome was analyzed using a series of EMX1-targeting chimeric crRNAs with single point mutations (FIG. 4A). FIG. 4B illustrates results of a Surveyor nuclease assay comparing the cleavage efficiency of Cas9 when paired with different mutant chimeric RNAs. Single-base mismatch up

to 12-bp 5' of the PAM substantially abrogated genomic cleavage by SpCas9, whereas spacers with mutations at farther upstream positions retained activity against the original protospacer target (FIG. 4B). In addition to the PAM, SpCas9 has single-base specificity within the last 12-bp of the spacer. Furthermore, CRISPR is able to mediate genomic cleavage as efficiently as a pair of TALE nucleases (TALEN) targeting the same EMX1 protospacer. FIG. 4C provides a schematic showing the design of TALENs targeting EMX1, and FIG. 4D shows a Surveyor gel comparing the efficiency of TALEN and Cas9 (n=3).

[0220] Having established a set of components for achieving CRISPR-mediated gene editing in mammalian cells through the error-prone NHEJ mechanism, the ability of CRISPR to stimulate homologous recombination (HR), a high fidelity gene repair pathway for making precise edits in the genome, was tested. The wild type SpCas9 is able to mediate site-specific DSBs, which can be repaired through both NHEJ and HR. In addition, an aspartate-to-alanine substitution (D10A) in the RuvC I catalytic domain of SpCas9 was engineered to convert the nuclease into a nickase (SpCas9n; illustrated in FIG. 5A) (see e.g. Sapranaukas et al., 2011, *Nucleic Acids Research*, 39: 9275; Gasiunas et al., 2012, *Proc. Natl. Acad. Sci. USA*, 109: E2579), such that nicked genomic DNA undergoes the high-fidelity homology-directed repair (HDR). Surveyor assay confirmed that SpCas9n does not generate indels at the EMX1 protospacer target. As illustrated in FIG. 5B, co-expression of EMX1-targeting chimeric crRNA with SpCas9 produced indels in the target site, whereas co-expression with SpCas9n did not (n=3). Moreover, sequencing of 327 amplicons did not detect any indels induced by SpCas9n. The same locus was selected to test CRISPR-mediated HR by co-transfecting HEK 293FT cells with the chimeric RNA targeting EMX1, hSpCas9 or hSpCas9n, as well as a HR template to introduce a pair of restriction sites (HindIII and NheI) near the protospacer. FIG. 5C provides a schematic illustration of the HR strategy, with relative locations of recombination points and primer annealing sequences (arrows). SpCas9 and SpCas9n indeed catalyzed integration of the HR template into the EMX1 locus. PCR amplification of the target region followed by restriction digest with HindIII revealed cleavage products corresponding to expected fragment sizes (arrows in restriction fragment length polymorphism gel analysis shown in FIG. 5D), with SpCas9 and SpCas9n mediating similar levels of HR efficiencies. Applicants further verified HR using Sanger sequencing of genomic amplicons (FIG. 5E). These results demonstrate the utility of CRISPR for facilitating targeted gene insertion in the mammalian genome. Given the 14-bp (12-bp from the spacer and 2-bp from the PAM) target specificity of the wild type SpCas9, the availability of a nickase can significantly reduce the likelihood of off-target modifications, since single strand breaks are not substrates for the error-prone NHEJ pathway.

[0221] Expression constructs mimicking the natural architecture of CRISPR loci with arrayed spacers (FIG. 2A) were constructed to test the possibility of multiplexed sequence targeting. Using a single CRISPR array encoding a pair of EMX1- and PVALB-targeting spacers, efficient cleavage at both loci was detected (FIG. 4F, showing both a schematic design of the crRNA array and a Surveyor blot showing efficient mediation of cleavage). Targeted deletion of larger genomic regions through concurrent DSBs using spacers

against two targets within EMX1 spaced by 119 bp was also tested, and a 1.6% deletion efficacy (3 out of 182 amplicons; FIG. 4G) was detected. This demonstrates that the CRISPR system can mediate multiplexed editing within a single genome.

Example 2

CRISPR System Modifications and Alternatives

[0222] The ability to use RNA to program sequence-specific DNA cleavage defines a new class of genome engineering tools for a variety of research and industrial applications. Several aspects of the CRISPR system can be further improved to increase the efficiency and versatility of CRISPR targeting. Optimal Cas9 activity may depend on the availability of free Mg^{2+} at levels higher than that present in the mammalian nucleus (see e.g. Jinek et al., 2012, *Science*, 337:816), and the preference for an NGG motif immediately downstream of the protospacer restricts the ability to target on average every 12-bp in the human genome (FIG. 11, evaluating both plus and minus strands of human chromosomal sequences). Some of these constraints can be overcome by exploring the diversity of CRISPR loci across the microbial metagenome (see e.g. Makarova et al., 2011, *Nat Rev Microbiol*, 9:467). Other CRISPR loci may be transplanted into the mammalian cellular milieu by a process similar to that described in Example 1. For example, FIG. 12 illustrates adaptation of the Type II CRISPR system from CRISPR 1 of *Streptococcus thermophilus* LMD-9 for heterologous expression in mammalian cells to achieve CRISPR-mediated genome editing. FIG. 12A provides a Schematic illustration of CRISPR 1 from *S. thermophilus* LMD-9. FIG. 12B illustrates the design of an expression system for the *S. thermophilus* CRISPR system. Human codon-optimized hStCas9 is expressed using a constitutive EF1 α promoter. Mature versions of tracrRNA and crRNA are expressed using the U6 promoter to promote precise transcription initiation. Sequences from the mature crRNA and tracrRNA are illustrated. A single base indicated by the lower case “a” in the crRNA sequence is used to remove the polyU sequence, which serves as a RNA polIII transcriptional terminator. FIG. 12C provides a schematic showing guide sequences targeting the human EMX1 locus as well as their predicted secondary structures. The modification efficiency at each target site is indicated below the RNA secondary structures. The algorithm generating the structures colors each base according to its probability of assuming the predicted secondary structure, which is indicated by a rainbow scale reproduced in FIG. 12C in gray scale. FIG. 12D shows the results of hStCas9-mediated cleavage in the target locus using the Surveyor assay. RNA guide spacers 1 and 2 induced 14% and 6.4%, respectively. Statistical analysis of cleavage activity across biological replica at these two protospacer sites is also provided in FIG. 6. FIG. 16 provides a schematic of additional protospacer and corresponding PAM sequence targets of the *S. thermophilus* CRISPR system in the human EMX1 locus. Two protospacer sequences are highlighted and their corresponding PAM sequences satisfying NNAGAAW motif are indicated by underlining 3' with respect to the corresponding highlighted sequence. Both protospacers target the anti-sense strand.

Example 3

Sample Target Sequence Selection Algorithm

[0223] A software program is designed to identify candidate CRISPR target sequences on both strands of an input DNA sequence based on desired guide sequence length and a CRISPR motif sequence (PAM) for a specified CRISPR enzyme. For example, target sites for Cas9 from *S. pyogenes*, with PAM sequences NGG, may be identified by searching for 5'-N_x-NGG-3' both on the input sequence and on the reverse-complement of the input. Likewise, target sites for Cas9 of *S. thermophilus* CRISPR1, with PAM sequence NNAGAAW, may be identified by searching for 5'-N_x-NNAGAAW-3' (SEQ ID NO: 29) both on the input sequence and on the reverse-complement of the input. Likewise, target sites for Cas9 of *S. thermophilus* CRISPR3, with PAM sequence NGGNG, may be identified by searching for 5'-N_x-NGGNG-3' both on the input sequence and on the reverse-complement of the input. The value “x” in N_x may be fixed by the program or specified by the user, such as 20.

[0224] Since multiple occurrences in the genome of the DNA target site may lead to nonspecific genome editing, after identifying all potential sites, the program filters out sequences based on the number of times they appear in the relevant reference genome. For those CRISPR enzymes for which sequence specificity is determined by a ‘seed’ sequence, such as the 11-12 bp 5' from the PAM sequence, including the PAM sequence itself, the filtering step may be based on the seed sequence. Thus, to avoid editing at additional genomic loci, results are filtered based on the number of occurrences of the seed:PAM sequence in the relevant genome. The user may be allowed to choose the length of the seed sequence. The user may also be allowed to specify the number of occurrences of the seed:PAM sequence in a genome for purposes of passing the filter. The default is to screen for unique sequences. Filtration level is altered by changing both the length of the seed sequence and the number of occurrences of the sequence in the genome. The program may in addition or alternatively provide the sequence of a guide sequence complementary to the reported target sequence(s) by providing the reverse complement of the identified target sequence(s).

[0225] Further details of methods and algorithms to optimize sequence selection can be found in U.S. application Ser. No. 61/836,080 (attorney docket 44790.11.2022); incorporated herein by reference.

Example 4

Evaluation of Multiple Chimeric crRNA-tracrRNA Hybrids

[0226] This example describes results obtained for chimeric RNAs (chiRNAs; comprising a guide sequence, a tracr mate sequence, and a tracr sequence in a single transcript) having tracr sequences that incorporate different lengths of wild-type tracrRNA sequence. FIG. 18a illustrates a schematic of a bicistronic expression vector for chimeric RNA and Cas9. Cas9 is driven by the CBh promoter and the chimeric RNA is driven by a U6 promoter. The chimeric guide RNA consists of a 20 bp guide sequence (Ns) joined to the tracr sequence (running from the first “U” of the lower strand to the end of the transcript), which is truncated at

various positions as indicated. The guide and tracr sequences are separated by the tracr-mate sequence GUUUUA-GAGCUA (SEQ ID NO: 30) followed by the loop sequence GAAA. Results of SURVEYOR assays for Cas9-mediated indels at the human EMX1 and PVALB loci are illustrated in FIGS. 18b and 18c, respectively. Arrows indicate the expected SURVEYOR fragments. ChiRNAs are indicated by their "+n" designation, and crRNA refers to a hybrid RNA where guide and tracr sequences are expressed as separate transcripts. Quantification of these results, performed in triplicate, are illustrated by histogram in FIGS. 19a and 19b, corresponding to FIGS. 18b and 18c, respectively ("N.D." indicates no indels detected). Protospacer IDs and their corresponding genomic target, protospacer sequence, PAM sequence, and strand location are provided in Table D. Guide sequences were designed to be complementary to the entire protospacer sequence in the case of separate transcripts in the hybrid system, or only to the underlined portion in the case of chimeric RNAs.

TABLE D

protospacer ID	genomic target	protospacer sequence (5' to 3')	PAM	strand
1	EMX1	<u>GGACATCGATGTCACCTCCA</u> <u>ATGACTAGGG</u> (SEQ ID NO: 31)	TGG	+
2	EMX1	<u>CATTGGAGGTGACATCGATG</u> <u>TCCTCCCAT</u> (SEQ ID NO: 32)	TGG	-
3	EMX1	<u>GGAAGGGCTGAGTCCGAGC</u> <u>AGAGAGAGAA</u> (SEQ ID NO: 33)	GGG	+
4	PVALB	<u>GGTGGCGAGAGGGGCCGAGA</u> <u>TTGGGTGTTC</u> (SEQ ID NO: 34)	AGG	+
5	PVALB	<u>ATGCAGGAGGGTGGCGAGAG</u> <u>GGGCCGAGAT</u> (SEQ ID NO: 35)	TGG	+

[0227] Cell Culture and Transfection

[0228] Human embryonic kidney (HEK) cell line 293FT (Life Technologies) was maintained in Dulbecco's modified Eagle's Medium (DMEM) supplemented with 10% fetal bovine serum (HyClone), 2 mM GlutaMAX (Life Technologies), 100 U/mL penicillin, and 100 m/mL streptomycin at 37° C. with 5% CO₂ incubation. 293FT cells were seeded onto 24-well plates (Corning) 24 hours prior to transfection at a density of 150,000 cells per well. Cells were transfected using Lipofectamine 2000 (Life Technologies) following the manufacturer's recommended protocol. For each well of a 24-well plate, a total of 500 ng plasmid was used.

[0229] SURVEYOR Assay for Genome Modification

[0230] 293FT cells were transfected with plasmid DNA as described above. Cells were incubated at 37° C. for 72 hours post-transfection prior to genomic DNA extraction. Genomic DNA was extracted using the QuickExtract DNA Extraction Solution (Epicentre) following the manufacturer's protocol. Briefly, pelleted cells were resuspended in QuickExtract solution and incubated at 65° C. for 15 minutes and 98° C. for 10 minutes. The genomic region flanking the CRISPR target site for each gene was PCR amplified (primers listed in Table E), and products were purified using

QiaQuick Spin Column (Qiagen) following the manufacturer's protocol. 400 ng total of the purified PCR products were mixed with 2 µl 10× Taq DNA Polymerase PCR buffer (Enzymatics) and ultrapure water to a final volume of 20 µl, and subjected to a re-annealing process to enable heteroduplex formation: 95° C. for 10 min, 95° C. to 85° C. ramping at -2° C./s, 85° C. to 25° C. at -0.25° C./s, and 25° C. hold for 1 minute. After re-annealing, products were treated with SURVEYOR nuclease and SURVEYOR enhancer S (Transgenomics) following the manufacturer's recommended protocol, and analyzed on 4-20% Novex TBE poly-acrylamide gels (Life Technologies). Gels were stained with SYBR Gold DNA stain (Life Technologies) for 30 minutes and imaged with a Gel Doc gel imaging system (Bio-rad). Quantification was based on relative band intensities.

TABLE E

primer name	genomic target	primer sequence (5' to 3')
Sp-EMX1-F	EMX1	AAAACCACCTTCTCTCTGGC (SEQ ID NO: 36)
Sp-EMX1-R	EMX1	GGAGATTGGAGACCGGAGAG (SEQ ID NO: 37)
Sp-PVALB-F	PVALB	CTGGAAAGCCAATGCCTGAC (SEQ ID NO: 38)
Sp-PVALB-R	PVALB	GGCAGCAACTCCTTGTCTCT (SEQ ID NO: 39)

[0231] Computational Identification of Unique CRISPR Target Sites

[0232] To identify unique target sites for the *S. pyogenes* SF370 Cas9 (SpCas9) enzyme in the human, mouse, rat, zebrafish, fruit fly, and *C. elegans* genome, we developed a software package to scan both strands of a DNA sequence and identify all possible SpCas9 target sites. For this example, each SpCas9 target site was operationally defined as a 20 bp sequence followed by an NGG protospacer adjacent motif (PAM) sequence, and we identified all sequences satisfying this 5'-N₂₀-NGG-3' definition on all chromosomes. To prevent non-specific genome editing, after identifying all potential sites, all target sites were filtered based on the number of times they appear in the relevant reference genome. To take advantage of sequence specificity of Cas9 activity conferred by a 'seed' sequence, which can be, for example, approximately 11-12 bp sequence 5' from the PAM sequence, 5'-NNNNNNNNNN-NGG-3' sequences were selected to be unique in the relevant genome. All genomic sequences were downloaded from the UCSC Genome Browser (Human genome hg19, Mouse genome mm9, Rat genome rn5, Zebrafish genome danRer7, *D. melanogaster* genome dm4 and *C. elegans* genome ce10). The full search results are available to browse using UCSC Genome Browser information. An example visualization of some target sites in the human genome is provided in FIG. 21.

[0233] Initially, three sites within the EMX1 locus in human HEK 293FT cells were targeted. Genome modification efficiency of each chiRNA was assessed using the SURVEYOR nuclease assay, which detects mutations resulting from DNA double-strand breaks (DSBs) and their subsequent repair by the non-homologous end joining (NHEJ) DNA damage repair pathway. Constructs designated

chiRNA(+n) indicate that up to the +n nucleotide of wild-type tracrRNA is included in the chimeric RNA construct, with values of 48, 54, 67, and 85 used for n. Chimeric RNAs containing longer fragments of wild-type tracrRNA (chiRNA(+67) and chiRNA(+85)) mediated DNA cleavage at all three EMX1 target sites, with chiRNA(+85) in particular demonstrating significantly higher levels of DNA cleavage than the corresponding crRNA/tracrRNA hybrids that expressed guide and tracr sequences in separate transcripts (FIGS. 18b and 19a). Two sites in the PVALB locus that yielded no detectable cleavage using the hybrid system (guide sequence and tracr sequence expressed as separate transcripts) were also targeted using chiRNAs. chiRNA(+67) and chiRNA(+85) were able to mediate significant cleavage at the two PVALB protospacers (FIGS. 18c and 19b).

[0234] For all five targets in the EMX1 and PVALB loci, a consistent increase in genome modification efficiency with increasing tracr sequence length was observed. Without wishing to be bound by any theory, the secondary structure formed by the 3' end of the tracrRNA may play a role in enhancing the rate of CRISPR complex formation. An illustration of predicted secondary structures for each of the chimeric RNAs used in this example is provided in FIG. 21. The secondary structure was predicted using RNAfold using minimum free energy and partition function algorithm. Pseudocolor for each based (reproduced in grayscale) indicates the probability of pairing. Because chiRNAs with

longer tracr sequences were able to cleave targets that were not cleaved by native CRISPR crRNA/tracrRNA hybrids, it is possible that chimeric RNA may be loaded onto Cas9 more efficiently than its native hybrid counterpart. To facilitate the application of Cas9 for site-specific genome editing in eukaryotic cells and organisms, all predicted unique target sites for the *S. pyogenes* Cas9 were computationally identified in the human, mouse, rat, zebra fish, *C. elegans*, and *D. melanogaster* genomes. Chimeric RNAs can be designed for Cas9 enzymes from other microbes to expand the target space of CRISPR RNA-programmable nucleases.

[0235] FIG. 22 illustrates an exemplary bicistronic expression vector for expression of chimeric RNA including up to the +85 nucleotide of wild-type tracr RNA sequence, and SpCas9 with nuclear localization sequences. SpCas9 is expressed from a CBh promoter and terminated with the bGH polyA signal (bGH pA). The expanded sequence illustrated immediately below the schematic corresponds to the region surrounding the guide sequence insertion site, and includes, from 5' to 3', 3'-portion of the U6 promoter (first shaded region), BbsI cleavage sites (arrows), partial direct repeat (tracr mate sequence GTTTTACAGCTA (SEQ ID NO: 27), underlined), loop sequence GAAA, and +85 tracr sequence (underlined sequence following loop sequence). An exemplary guide sequence insert is illustrated below the guide sequence insertion site, with nucleotides of the guide sequence for a selected target represented by an "N".

[0236] Sequences described in the above examples are as follows (polynucleotide sequences are 5' to 3'):

```

U6-short tracrRNA (Streptococcus pyogenes SF370): (SEQ ID NO: 40)
GAGGGCCTATTTCCCATGATTCTTCATATTTGCATATACGATACAAGGCT
GTTAGAGAGATAATTGGAATTAATTTGACTGTAAACACAAAGATATTAGTACAAAA
TACGTGACGTAGAAAAGTAATAATTTCTGGGTAGTTTGCAGTTTTAAAATTATGTTTT
AAAATGGACTATCATATGCTTACCGTAACTTGAAAGTATTTTCGATTTCTGGCTTTAT
ATATCTTGTGGAAAGGACGAAACACCGGAACCATTCAAACAGCATAGCAAGTTA
AAATAAGGCTAGTCCGTATCAACTTGAAAAAGTGGCACCGAGTCGGTCTTTT
TTT (bold = tracrRNA sequence;
underline = terminator sequence)

U6-long tracrRNA (Streptococcus pyogenes SF370): (SEQ ID NO: 41)
GAGGGCCTATTTCCCATGATTCTTCATATTTGCATATACGATACAAGGCT
GTTAGAGAGATAATTGGAATTAATTTGACTGTAAACACAAAGATATTAGTACAAAA
TACGTGACGTAGAAAAGTAATAATTTCTGGGTAGTTTGCAGTTTTAAAATTATGTTTT
AAAATGGACTATCATATGCTTACCGTAACTTGAAAGTATTTTCGATTTCTGGCTTTAT
ATATCTTGTGGAAAGGACGAAACACCGGTAGTATTAAGTATTGTTTTATGGCTGATA
AATTTCTTTGAATTTCTCCTTGATTATTTGTTATAAAAGTTATAAAATAATCTTGTTG
GAACCATTCAAAACAGCATAGCAAGTTAAAATAAGGCTAGTCCGTATCAACTTGA
AAAAGTGGCACCGAGTCGGTCTTTTTTT

U6-DR-BbsI backbone-DR (Streptococcus pyogenes SF370): (SEQ ID NO: 42)
GAGGGCCTATTTCCCATGATTCTTCATATTTGCATATACGATACAAGGCT
GTTAGAGAGATAATTGGAATTAATTTGACTGTAAACACAAAGATATTAGTACAAAA
TACGTGACGTAGAAAAGTAATAATTTCTGGGTAGTTTGCAGTTTTAAAATTATGTTTT

```

- continued

AAAATGGACTATCATATGCTTACCGTAACTTGAAAGTATTCGATTTCTGGCTTTAT
ATATCTTGTGGAAAGGACGAAACACCGGGTTTTAGAGCTATGCTGTTTTGAATGGTC
CCAAAACGGGTCTTCGAGAAGACGTTTTAGAGCTATGCTGTTTTGAATGGTCCCAA
AC

U6-chimeric RNA-BbsI backbone (*Streptococcus pyogenes* SF370)
(SEQ ID NO: 43)

GAGGGCCTATTTCCATGATTCCTTCATATTTGCATATACGATACAAGGCT
GTTAGAGAGATAATGGAATTAATTTGACTGTAAACACAAGATATTAGTACAAAA
TACGTGACGTAGAAAGTAATAATTTCTGGGTAGTTGCAGTTTTAAAATTATGTTTT
AAAATGGACTATCATATGCTTACCGTAACTTGAAAGTATTCGATTTCTGGCTTTAT
ATATCTTGTGGAAAGGACGAAACACCGGGTCTTCGAGAAGACCTGTTTTAGAGCTA
GAAATAGCAAGTAAAATAAGGCTAGTCCG

NLS-SpCas9-EGFP:
(SEQ ID NO: 44)

MDYKDHDGDYKDHDIDYKDDDDKMAPKKRKGVIHGVPAAADKKYSIGLDI
GTNSVGWAVITDEYKVPSKPKVLGNTDRHSIKKNLIGALLFDSGETAEATRLKRTARR
RYTRRNRI CYLQEIFSNEMAKVDDSPFHRLEESFLVEEDKKHERHPFGNIVDEVAYHE
KYPTIYHLRKKLVDSTDKADLRLIYLALAHMIKFRGHFLIEGDLNPDNSDVKLFIQLVQ
TYNQLFEBENPINASGVDAKAIL SARLSKSRLENLIAQLPGEKKNLFGNLIALSLGLTPN
FKSNFDLAEDAKLQLSKD TYDDLDNLLAQIGDQYADLFLAAKNLSDAILLSDILRVNT
EITKAPLSASMIKRYDEHHQDLTLKALVRRQQLPEKYKEIFPDQSKNGYAGYIDGGASQ
EEFYKFIKPILEKMDGTEELLVKLNREDLLRQRTFDNGSIPHQIHLGELHAILRRQEDFY
PFLKDNREKIEKILTFRIPYVYVGPLARGNSRFWMTRKSEETITPWNFEVVDKGSASQ
FIERMTNFDKNLPNEKVLPHSLLYEYFTVYNELTKVKYVTEGMRKPAFLS GEQKKAIV
DLLFKTNRKVTVKQLKEDYFKKIECFDSVEISGVEDRFNASLGT YHDL LKIKDKDFLDN
EENEDI LEDIVLTLTLFEDREMI EERLKYAHLFDDKVMKQLKRRRYTGWGRLSRKLIN
GIRDKQSGKTI LDFLKS DGFANRNFQLIHDDSLTFKEDIQKAQVSGQDLSLHEHIANLA
GSPAIIKKGILQTVKVVDELVKVMGRHKPENIVIEMARENQTTQKGQKNSRERMKRIEEG
IKELGSQILKEHPVENTQLQNEKLYLYLQNGRDMYVDQELDINRLSDYDVDHIVPQSF
LKDDSIDNKVLRSDKNRGS DNVPS EEVVKMKNYWRQLLNAKLITQRKFDNLTKAE
RGGLELKDAGFIKRLVETRQITKHVAQILD SRMNTKYDENDKLIREVKVITLKS KLVS
DFRKDFQFYKVIENNYHHAHDAYLNAVVG TALI KKYPKLESEFVYGDYKVYDVRKMI
AKSEQEIGKATAKYFFYSNIMNFFKTEITLANGEIRKRPLIETNGETGEIVWDKGRDFATV
RKVLSMPQVNI VKKTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKGFDSP TVAYS
VLVVAKEKGSKKLKSVKELLGITIMERS SFEKNPIDFLEAKGYKEVKKDLIIKLPKYS
LFELENGRKRMLASAGELQKGNELALPSKYVNFY LASHYEKLGSPEDNEQKQLFVE
QHKHYLDEIEIQISEFSKRVI LADANLDKVL SAYNKHRDKPIREQAENI IHLFTLTNLGAP
AAFKYFDTTIDRKYRSTKEVLDATLIHQSI TGLYETRIDLSQLGGDAAAVSKGEELFTG
VVPILVELDGDVNGHKFSVSGEGEDATY GKLTLKFICTTGKLPVPWPTLVTTLT YGVQ
CFSRYPDHMKQHDFPKSAMPEGYVQERTIFFKDDGNYKTRAEVKFEGDTLVNRIELKGI

- continued

DFKEDGNI LGHKLEYNYN SHNVYIMADKQKNGIKVNFKIRHNI EDGSVQLADHYQQNT

PIGDGPVLLPDNHYLSTQSALS KDPNEKRDHMLLEFVTAAGITLGMDELYK

SpCas9 - EGFP - NLS :

(SEQ ID NO: 45)

MDKKYS IGLDIGTNSVGWAVI TDEYKVP SKKPKVLGNTDRHSI KKNLIGALLF

DSGETAEATRLKRTARRRYTRRKNR ICYLQEIFSNEMAKVDDSPFHRLEESFLVEEDKK

HERHPIFGNIVDEVAYHEKYPTIYHLRKKLVSTDKADLR LIYLALAHMIKFRGHFLIEG

DLNPDNSDVKLFIQLVQTYNQLFEENPINASGVDAKAILSARLSKSRRENLI AQLPGE

KKNLFGNLI ALSLGLTPNFKSNFDLAEDAKLQLSKDTYDDDLNLLAQIGDQYADLFL

AAKNLSDAILLSDILRVNTEI TKAPLSASMI KRYDEHHQDLTLLKALVRQQLP EKYKEIFF

DQSKNGYAGYIDGGASQEEFYKFIKPILEKMDGTEELLVKLNREDLLRKQRTFDNGSIPH

QIHLGELHAILRRQEDFYFPLKDNREKIEKILTPRI PYVGPLARGNSRFAMWTRKSEETI

TPWNFEVVVDKGSASQSPIERMTNFDKNLPNEKVL PKHSLLEYEFTVYNELTKVKYVTE

GMRKPAFLSGEQKKAIVDLLFKTNRKVTVKQLKEDYFKKIECFDSVEISGVEDRFNASL

GTYHDLKIIKDKDFLDNEENEDI EDIVLTLTLFEDREMI EERLKYAHLFDDKVMKQL

KRRRYTGWRLSRKLINGIRDKQSGKTI LDFLKS DGFANRNFQILHDDSLTFKEDIQKA

QVSGQGDSLHEHIANLAGSPA I KKGILQTVKVVDELVKVMGRHKPENIV IEMARENQTT

QKQKNSRERMKRI EEGIKELGSI LKEHPVENTQLQNEKLYLYLQNGRDMYVDQEL

DINRLSDYDVDHIVPQSFLKDDSIDNKVLRSDKNR GSDNVPSEEVKMKNYWRQL

LNAKLI TQRKFDNLTKAERGGSEL DKA GF I KRQLVETRQITKHVAQILDSRMNTKYDE

NDKLIREVKVITLKS KLVSDPRKDFQFYKREINNYHHAHDAYLNAVGTALIKKYPKL

ESEFVYGDYKVVDRKMI AKSEQEIGKATAKYFFYSNIMNFFKTEITLANGEIRKRPLIET

NGETGEIVWDKGRDFATVRKVL SMPQVNI VKKTEVQTGGFSKESILPKRNSDKLIARKK

DWDPKKYGGFDSPTVAYSVLVAVKVEK GSKK LKSVKELLGITIMERS SFEKNPIDFLE

AKGYKEVKKDLI IKLPKYSLFELENGRKRMLASAGELQKGNELALPSKYVNFY LASHY

EKLKGS PEDNEBQKQLFEQHKHYLDEI IEQISEFSKRVLADANLDKVL SAYNKH RDKPI

REQAENI IHLFTLNLGAPAAF KYFDTTIDRKRYTSTKEVL DATLIHQ SITGLYETRIDL SQ

LGGDAAAVSKGEELFTGVVPI LVELDGDVNGHKFSVSGEGEDATYGLTLKFICTTGK

LPVPWPTLVTTLYGVQCF SRYPDHMKQHDFPKSAMPEGYVQERTIFFKDDGNYKTRA

EVKFEGDTLVNRIELKGI DFKEDGNILGHKLEYNYN SHNVYIMADKQKNGIKVNFKIRH

NIEDGSVQLADHYQQNTPIGDGPVLLPDNHYLSTQSALS KDPNEKRDHMLLEFVTAAG

ITLGMDELYKKRPAATKAGQAKKKK

NLS - SpCas9 - EGFP - NLS :

(SEQ ID NO: 46)

MDYKDHDGDYKDHDIDYKDDDDKMAPKKR KRVGIHGVPAAADKKYS IGLDI

GTNSVGWAVI TDEYKVP SKKPKVLGNTDRHSI KKNLIGALLFDSGETAEATRLKRTARR

RYTRRKNR ICYLQEIFSNEMAKVDDSPFHRLEESFLVEEDKKHERHPIFGNIVDEVAYHE

KYPTIYHLRKKLVSTDKADLR LIYLALAHMIKFRGHFLIEGDLNPDNSDVKLFIQLVQ

TYNQLFEENPINASGVDAKAILSARLSKSRRENLI AQLPGEKKNLFGNLI ALSLGLTPN

FKSNFDLAEDAKLQLSKDTYDDDLNLLAQIGDQYADLFLAAKNLSDAILLSDILRVNT

EITKAPLSASMI KRYDEHHQDLTLLKALVRQQLP EKYKEIFFDQSKNGYAGYIDGGASQ

- continued

EEFYKFIKPILEKMDGTEELLVKNREDLLRKQRTFDNGSIPHQIHLGELHAILRRQEDFY
PFLKDNREKIEKILTFRIPYYVGPLARGNSRFAMWTRKSEETITPWNFEVVDKGASAQS
FIERMTNFDKNLNEKVLPHKSLLYEYFTVYNELTKVKYVTEGMRKPAFLSGEQKKAIV
DLLFKTNRKVTVKQLKEDYFKKIECFDSVEISGVEDRFNASLGTYHDLKIKDKDFLDN
EENEDILEDIVLTLTLFEDREMIERLKTYAHLFDDKVMKQLKRRRYTGWGRLSRKLIN
GIRDKQSGKTIILDFLKSDFANRNFQLIHDDSLTFKEDIQKAQVSGQGDSLHEHIANLA
GSPAIIKKGILQTVKVVDELVKVMGRHKPENIVIAMARENQTTQKGQKNSRERMKRIEEG
IKELGSQILKEHPVENTQLQNEKLYLYLQNGRDMYVDQELDINRLSDYDVDHIVPQSF
LKDDSIDNKVLRSDKNRGSNDVPSSEVVKMKNYWRQLLNAKLITQRKFDNLTKAE
RGGLESELDKAGFIKQRLVETRQITKHVAQILDSRMNTKYDENDKLIREVKVITLKSCLVS
DFRKDFQFYKVRINNYHHAHDAYLNAVVGITALIKKYPKLESEFVYGDYKVDVRKMI
AKSEQEIGKATAKYFFYSNIMNPFKTEITLANGEIRKRPLIETNGETGEIVWDKGRDFATV
RKVLSMPQVNIIVKTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFDSPTVAYS
VLVVAKVEKGSKKLKSVKELLGITIMERSSEKNPIDFLEAKGYKEVKKDLIKLPKYS
LFELENGRKRMLASAGELQKGNELALPSKYVNFYLYLASHYEKLGKSPEDNEQKQLFVE
QHKHYLDEIIEQISEFSKRVI LADANLDKVL SAYNKHDKPIREQAENI IHLFTLTNLGAP
AAFKYFDTTIDRKRVTSTKEVLDATLIHQSI TGLYETRIDLSQLGGDAAAVSKGEELFTG
VVPILVELDGDVNGHKFSVSGEGEDATYGLTLKFICTTGKLPVPWPTLVTTLYGVQ
CFSRYPDHMKQHDFKFSAMPEGYVQERTIFFKDDGNYKTRAEVKFEGDTLVNRIELKGI
DFKEDGNI LGHKLEYNYNSHNVYIMADKQKNGIKVNFKIRHNI EDGSVQLADHYQONT
PIGDGPVLLPDNHVLSQSAKSKDPNEKRDHMLLEFVTAAGITLGMDELYKKRPAATK
KAGQAKKKK

NLS - SpCas9 - NLS : (SEQ ID NO: 47)

MDYKDHDGDYKDHDIDYKDDDDKMAPKKRKGIVHGVPAAADKKYSIGLDI
GTNSVGWAVI TDEYKVP SKKPKVLGNTDRHSIKKNLIGALLFDSGETAEATRLKRTARR
RYTRRKNRICYLQEIFSNEMAKVDDSPFHRLEESFLVEEDKKHERHPIFGNIVDEVAYHE
KYPTIYHLRKKLVDSTDKADLRLIYLALAHMIKFRGHFLIEGDLNPDNSDVKLFIQLVQ
TYNQLFEENPINASGVDAKAIL SARLSKSRRLLENLIAQLPGEKKNLFGNLIALSLGLTPN
FKSNFDLAEDAKLQLSKD TYDDDLNLLAQIGDQYADLFLAAKNLSDAILLSDILRVNT
EITKAPLSASMIKRDEHHQDLTLLKALVRQQLPEKYKEIFFDQSKNGYAGYIDGGASQ
EEFYKFIKPILEKMDGTEELLVKNREDLLRKQRTFDNGSIPHQIHLGELHAILRRQEDFY
PFLKDNREKIEKILTFRIPYYVGPLARGNSRFAMWTRKSEETITPWNFEVVDKGASAQS
FIERMTNFDKNLNEKVLPHKSLLYEYFTVYNELTKVKYVTEGMRKPAFLSGEQKKAIV
DLLFKTNRKVTVKQLKEDYFKKIECFDSVEISGVEDRFNASLGTYHDLKIKDKDFLDN
EENEDILEDIVLTLTLFEDREMIERLKTYAHLFDDKVMKQLKRRRYTGWGRLSRKLIN
GIRDKQSGKTIILDFLKSDFANRNFQLIHDDSLTFKEDIQKAQVSGQGDSLHEHIANLA
GSPAIIKKGILQTVKVVDELVKVMGRHKPENIVIAMARENQTTQKGQKNSRERMKRIEEG
IKELGSQILKEHPVENTQLQNEKLYLYLQNGRDMYVDQELDINRLSDYDVDHIVPQSF
LKDDSIDNKVLRSDKNRGSNDVPSSEVVKMKNYWRQLLNAKLITQRKFDNLTKAE

- continued

RGGLESELDKAGFIKRQLVETRQITKHVAQIILDSRMNTKYDENDKLIREVKVITLKSCLVS
 DFRKDFQFYKVINNYHHADAYLNAVVGITALIKKYPKLESEFVYGDYKVDVRKMI
 AKSEQEIGKATAKYFFYSNIMNFFKTEITLANGEIRKRPLIETNGETGEIVWDKGRDFATV
 RKLVSMPQVNVIVKTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFDSPTVAYS
 VLVVAKVEKKGSKLKSVKELGITIMERSSEKNPIDFLEAKGYKEVKKDLIIKLPKYS
 LFELENGRKRMLASAGELQKGNELALPSKYVNFYLAHYEKLKGSPEDEQKQLFVE
 QHKHYLDEIIEQISEFSKRVILADANLDKVL SAYNKHRDKPIREQAENI IHLFTLTNLGAP
 AAFKYFDTTIDRKRYSSTKEVLDATLIHQSI TGLYETRIDLSQLGGDKRPAATKKAGQAK
 KKK

NLS-mCherry-SpRNase3:

(SEQ ID NO: 48)

MFLFLSLTSFLSSSRTLVSKEEDNMAIIKEFMRFKVHMEGVSNGHEFEIEGE
 GEGRPYEGTQTAKLKVTKGGPLPFAWDILSPQFMYGSKAYVKHPADIPDYLKLSFPEGF
 KWERMNFEDGGVVTVDSSLDGGEFIYKVKLRGTNFPDGPVMQKKTMGWEASSE
 RMYPEDGALKGEIKQRLKLDGGHYDAEVKTTYKAKKPVQLPGAYNVNIKLDITSHNE
 DYTIVEQYERAEGRHSTGGMDELYKGSKQLEELLSTSPDIQFNDLTLETAFTHTSYANE
 HRLLNVSHERLEFLGDAVLQLIISEYLFAYPKKTEGDMSKLRSMIVREESLAGFSRFC
 SFDAYIKLGKGEKSGRRRDTILGDLFEAPLGAALLDKGIDAVRRFLKQVMIPQVEKG
 NFERVKDYKTCLQEFLQTKGDVAIDYQVISEKGAHAKQFEVSI VVNGAVLSKGLGKSK
 KLAEQDAAKNALAQLSEV

SpRNase3-mCherry-NLS:

(SEQ ID NO: 49)

MKQLEELLSTSPDIQFNDLTLETAFTHTSYANEHRLLNVSHERLEFLGDAV
 LQLIISEYLFAYPKKTEGDMSKLRSMIVREESLAGFSRFC SFDAYIKLGKGEKSGRR
 RDTILGDLFEAPLGAALLDKGIDAVRRFLKQVMIPQVEKGNFERVKDYKTCLQEFLQTK
 GDVAIDYQVISEKGAHAKQFEVSI VVNGAVLSKGLGKSKLAEQDAAKNALAQLSEV
 GSVSKGEEDNMAIIKEFMRFKVHMEGVSNGHEFEIEGEGEGRPYEGTQTAKLKVTKGGP
 LPFAWDILSPQFMYGSKAYVKHPADIPDYLKLSFPEGFKWERMNFEDGGVVTVDSS
 SLQDGEFIYKVKLRGTNFPDGPVMQKKTMGWEASSERMYPEDGALKGEIKQRLKLD
 GGHYDAEVKTTYKAKKPVQLPGAYNVNIKLDITSHNEDYTIVEQYERAEGRHSTGGMD
 ELYKKRPAATKKAGQAKKKK

NLS-SpCas9n-NLS (the D10A nickase mutation is lowercase):

(SEQ ID NO: 50)

M DYKDHGDYKDHDIDYKDDDDKMAPKKRKGIVHGVPAAADKKYSIGLaI
 GTNSVGWAVITDEYKVPKSKFKVLGNTDRHSIKKNLIGALLFDSGETAEATRLKRTARR
 RYTRRKNRICYLQEIFSNEMAKVDDSFPHRLEESFLVEEDKKHERHPIFGNIVDEVAYHE
 KYPTIYHLRKKLVDSTDKADLRLIYLALAHMIKFRGHFLIEGDLNPDNSDVKLFIQLVQ
 TYNQLFEENPINASGVDAKAIL SARLSKSRLENLIAQLPGEKKNGLFGNLIALSLGLTPN
 FKSNDLAEDAKLQLSKD TYDDDLNLLAQIGDQYADLFLAAKNLSDAILLSDILRVNT
 EITKAPLSASMIKRYDEHHQDLTLLKALVRQQLPEKYKEIPFDQSKNGYAGYIDGGASQ
 EEFYKFIKPILEKMDGTEELLVKNREDLLRKQRTFDNGSIPHQIHLGELHAILRRQEDFY
 PFLKDNREKIEKILTFRIPYVVGPLARGNSRFAWMTRKSEETITPWNPEEVVDKGAASQ

- continued

FIERMTNFDKNLPEKVLPHKSLLYEYFTVYNELTKVKYVTEGMRKPAFLSGEQKKAIV
 DLLPKTNRKVTVKQLKEDYFKKIECFDSVEISGVEDRFNASLGTYHDLKIKDKDFLDN
 EENEDILEDIVLTLTLFEDREMIERLKYAHLFDDKVMKQLKRRRYTGWGRLSRKLIN
 GIRDKQSGKITLDFLKSDFANRNFMLIHDDSLTFKEDIQKAQVSGQGDSLHEHIANLA
 GSPAIIKKGILQTVKVVDELVKVMGRHKPENIVIEMARENQTTQKGQKNSRERMKRIEEG
 IKELGSQILKEHPVENTQLQNEKLYLYLQNGRDMYVDQELDINRLSDYDVIDHIVPQSF
 LKDDSIDNKVLTRSDKNRGSNDVPSSEVVKMKNYWRQLLNAKLITQRKFDNLTKAE
 RGGELSELDKAGFIKRLVETRQITKHVAQILD SRMNTKYDENDKLIREVKVITLKS KLVS
 DFRKDFQFYKREINNYHHAHDAYLNAVVGITALIKKYPKLESEFVYGDYKVDVRKMI
 AKSEQEIGKATAKYFFYSNIMNFPKTEITLANGEIRKRPLIETNGETGEIVWDKGRDFATV
 RKVLSMPQVNIIVKKEVQTGGFSKESILPKRNSDKLIARKKDWDPKKGFGFDSPTVAYS
 VLVVAKVEKKGSKLKSVKELLGITIMERSSEKPNIDFLEAKGYKEVKKDLIIKLPKYS
 LFELENGRKRMLASAGELQKGNELALPSKYVNFYLYLASHYEKLGSPEDNEQKQLFVE
 QHKHYLDEIIEQISEFSKRVI LADANLDKVL SAYNKHRDKPIREQAENI IHLFTLTNLGAP
 AAPKYFDTTIDRKYRSTKEVLDATLIHQSIITGLYETRIDLSQLGGDKRPAATKKAGQAK
 KKK

hEMX1-HR Template-HindII-NheI:

(SEQ ID NO: 51)

GAATGCTGCCCTCAGACCCGCTTCCCTCCCTGTCTGTCTGTCACAGGAGA
 ATGAGGTCTACTGGTGGATTTCGGACTACCCCTGAGGAGCTGGCACCTGAGGGACA
 AGGCCCCCACCTGCCCAGCTCCAGCCTCTGATGAGGGGTGGGAGAGACTACATG
 AGGTTGCTAAGAAAGCCTCCCTGAAGGAGACCACACAGTGTGTGAGGTTGGAGTC
 TCTAGCAGCGGTTCTGTGCCCCAGGGATAGTCTGGCTGTCCAGGCCTGCTCTTG
 ATATAAACACCACCCTCCTAGTTATGAAACCATGCCATCTCTGCTCTCTGTATGGAA
 AAGAGCATGGGGCTGGCCCGTGGGGTGGTGTCCACTTTAGGCCCTGTGGGAGATCA
 TGGGAACCCACGAGTGGGTATAGGCTCTCTCATTACTACTCACATCCACTCTGT
 GAAGAAGCGATTATGATCTCTCCTCTAGAACTCGTAGAGTCCCATGTCTGCCGGCT
 TCCAGAGCCTGCACCTCCACCTTGGCTTGGCTTTGCTGGGGCTAGAGGAGCTAGG
 ATGCACAGCAGCTCTGTGACCCTTTGTTTGAGAGGAACAGGAAAACCCCTTCTCT
 CTGGCCCACTGTGTCTCTCTCTGCCCCGTCATCCCTTCTGTGAATGTTAGACCCAT
 GGGAGCAGCTGGTCAGAGGGGACCCCGCCTGGGGCCCTAACCTATGTAGCCTC
 AGTCTTCCCATCAGGCTCTCAGCTCAGCCTGAGTGTGAGGCCCCAGTGGCTGCTCT
 GGGGGCTCCTGAGTTTCTCATCTGTGCCCTCCCTCCCTGGCCAGGTGAAGGTGT
 GGTTCAGAACCGGAGGACAAAGTACAACCGCAGAAAGCTGGAGGAGGAAGGGCC
 TGAGTCCGAGCAGAAGAAGGGCTCCCATCACATCAACCGGTGGCGCATTGCCA
 CGAAGCAGGCCAATGGGAGGACATCGATGTCACCTCCAATGACaagctgctagcGGTGG
 GCAACCACAAACCCACGAGGGCAGAGTGTCTGCTTGTCTGCTGGCCAGGCCCTGCGT
 GGGCCCAAGCTGGACTCTGGCCACTCCCTGGCCAGGCTTTGGGGAGGCTGGAGTC
 ATGGCCCCACAGGGCTTGAAGCCCGGGCCGCCATTGACAGAGGGACAAGCAATGG
 GCTGGCTGAGGCTGGGACCACTTGGCCTTCTCTCGGAGAGCCTGCCTGCCTGGGC

- continued

GGGCCCCCGCCACCCGAGCCTCCAGCTGCTCTCCGTGTCTCCAATCTCCCTTTTG
 TTTTGATGCATTTCTGTTTAAATTTATTTTCCAGGCACCACTGTAGTTTAGTGATCCCC
 AGTGTCCCCCTTCCCTATGGGAATAATAAAAGTCTCTCTCTTAATGACACGGGCATC
 CAGCTCCAGCCCCAGAGCCTGGGGTGGTAGATTCCGGCTCTGAGGGCCAGTGGGGG
 CTGGTAGAGCAAACGCGTT CAGGGCCTGGGAGCCTGGGGTGGGGTACTGGTGGAGG
 GGGTCAAGGGTAATTCATTAACCTCTCTTTTGTGGGGACCCCTGGTCTCTACCTC
 CAGCTCCACAGCAGGAGAAACAGGCTAGACATAGGAAGGGCCATCCTGTATCTTG
 AGGGAGGACAGGCCAGGTCTTTCTTAACGTATTGAGAGGTGGGAATCAGGCCAG
 GTAGTTCAATGGGAGAGGAGAGTGCTTCCCTCTGCCTAGAGACTCTGGTGGCTTCT
 CCAGTTGAGGAGAAACCAGAGGAAAGGGGAGGATTGGGGTCTGGGGGAGGGAACA
 CCATTCACAAAGGCTGACGGTTCAGTCCGAAGTCGTGGGCCACCCAGGATGCTCA
 CCTGTCTTGGAGAACCCTGGGCAGGTTGAGACTGCAGAGACAGGGCTTAAGGCT
 GAGCCTGCAACCAGTCCCCAGTCACTCAGGGCCTCCTCAGCCAAAGAAAGAGCAAC
 GTGCCAGGGCCCGCTGAGCTCTTGTGTTACCTG

NLS - St Csn1 - NLS :

(SEQ ID NO: 52)

MKRPAATKKAGQAKKKKSDLVLGLDIGISVGVGILNKVTGEI IHKNSRIFPA
 AQAENNLVRRTRNRQRRLLARRKKHRRVRLNRLFEEGLITDFTKISINLNPYQLRVKGL
 TDELSNEELFIALKNMVKHRGISYLLDDASDDGNSVGDYAQIVKENS KQLETKTPGQIQ
 ERYQTYGQLRGDFTVEKDGKHLINVPPTSAYRSEALRILQTQQEFNPQITDEFINRYL
 EILTGKRKYHGPNEKSRDYGRYRSTGETLDNIFGILIGKCTFPYDFRAAKASYTAQ
 EFNLLNDLNNLTVPTETKKLSKEQKNQI INYVKNEKAMGPAKLPKYIAKLLSCDVADIK
 GYRIDKSGKAEIHTFEAYRKMKLTLETLDIEQMDRETLDKLAYVLTNTEREGIQEALHE
 FADGFSFSQKQVDELVQFRKANSSIFGKGWHNFSVKLMMELIPELYETSEEQMTILTRLG
 KQKTTSSSNKTKYIDEKLLTEEIYNPVVAKSVRQAIKIVNAAIKEYGDFDNIVIEMARETN
 EDDEKKAIQKIQKANKDEKDAAMLKAANQYNGKAE LPHSVFPHGKQLATKIRLWHQQ
 GERCLYTGKTI S IHDLINNSNQFEVDHILPLSITFDDSLANKVLVYATANQEKQRTPYQ
 ALDSMDDAWSFRELKAFVRESKTL SNKKKEYLLTEEDI SKFDVRKKFIERNLVDTRYAS
 RVVLNALQEHFRAHKIDTKVSVVRGQFTSQLRRHWGIEKTRDTHHHAVDALIIAASSQ
 LNLWKKQKNTLVSYSEDQLLDIETGELISSDEYKESVFKAPYQHPVDTLKSKEFEDSILF
 SYQVDSKFNRKISDATIYATRQAKVGDKADETYVLGKIKDIYTDGYDAFMKIYKGD
 KSKFLMYRHPQTFEKVIEPILENYPNKQINEKQKEVPCNPFLKYKEEHGYIRKYSKKG
 GPEIKSLKYYDSKLGNHIDITPKDSMNKVVLSVSPWRADVFNKTTGKYEILGLKYAD
 LQFEKGTGTYKISQEKYNDIKKKEGVDSDSEFKFTLYKNDLLLVKDETETKEQQLFRFLSR
 TMPKQKHVELKPYDKQKFEGGEALIKVLGNVANSQCKKGLGKSNI S IYKVRTDVLG
 NQHI I KNEGDKPKLDFKRPAATKKAGQAKKKK

U6-St_tracrRNA (7-97) :

(SEQ ID NO: 53)

GAGGGCCTATTCCCATGATTCCTTCATATTGCATATACGATACAAGGCT
 GTTAGAGAGATAATTGGAATTAATTTGACTGTAACACAAGATATTAGTACAAAA
 TACGTGACGTAGAAAGTAATAATTTCTGGGTAGTTGCAGTTTTAAATATGTTTT

- continued

AAAATGGACTATCATATGCTTACCGTAACTTGAAAGTATTCGATTTCTGGCTTTAT
 ATATCTTGTGGAAAGGACGAAACACCGTTACTTAAATCTGCAGAAGCTACAAAGA
 TAAGGCTTCATGCCGAAATCAACACCCGTGCTATTTATGGCAGGGTGTTCGTTATT
 TAA

U6-DR-spacer-DR (*S. pyogenes* SF370)

(SEQ ID NO: 54)

gagggcctatttcccatgattccttcatatttgcatatcagatacaaggctgtagagag
 ataattggaattaatttgactgtaaacacaaagatattagtacaaaatcgtgacgtaga
 aagtaataatttcttgggtagtttgcagttttaaattatgttttaaaggactatcat
 atgcttacgtaactgaaagtatttcgatttcttggctttatatcttgtggaaggac
 gaaacaccgggttttagagctatgctgttttgaatgggtcccaaacNNNNNNNNNNNNNNN

NNNNNNNNNNNNNNgttttagagctatgctgttttgaatgggtcccaaacT

TTTTTT (lowercase underline = direct repeat; N = guide
 sequence; bold = terminator)

Chimeric RNA containing +48 tracr RNA (*S. pyogenes* SF370)

(SEQ ID NO: 55)

gagggcctatttcccatgattccttcatatttgcatatcagatacaaggctgtagagaga
 taattggaattaatttgactgtaaacacaaagatattagtacaaaatcgtgacgtagaaa
 gtaataatttcttgggtagtttgcagttttaaattatgttttaaaggactatcatatgc
 ttaccgtaactgaaagtatttcgatttcttggctttatatcttgtggaaggacgaaa
 caccNNNNNNNNNNNNNNNNgttttagagctagaaatagcaagttaaaataaggcta

gtccg**TTTTTT** (N = guide

sequence; first underline = tracr mate sequence; second
 underline = tracr sequence; bold = terminator)

Chimeric RNA containing +54 tracr RNA (*S. pyogenes* SF370)

(SEQ ID NO: 56)

gagggcctatttcccatgattccttcatatttgcatatcagatacaaggctgtagagaga
 taattggaattaatttgactgtaaacacaaagatattagtacaaaatcgtgacgtagaaag
 taataatttcttgggtagtttgcagttttaaattatgttttaaaggactatcatatgc
 ttaccgtaactgaaagtatttcgatttcttggctttatatcttgtggaaggacgaaac
 accNNNNNNNNNNNNNNNNgttttagagctagaaatagcaagttaaaataaggctagt

ccgtta**TTTTTT** (N = guide

sequence; first underline = tracr mate sequence; second
 underline = tracr sequence; bold = terminator)

Chimeric RNA containing +67 tracr RNA (*S. pyogenes* SF370)

(SEQ ID NO: 57)

gagggcctatttcccatgattccttcatatttgcatatcagatacaaggctgtagagagat
 aattggaattaatttgactgtaaacacaaagatattagtacaaaatcgtgacgtagaaagtaat
 aatttcttgggtagtttgcagttttaaattatgttttaaaggactatcatatgcttacgtaa
 cttgaaagtatttcgatttcttggctttatatcttgtggaaggacgaaacaccNNNNNNNNNN

- continued

NNNNNNNNNNqtttttagagctagaatagcaagttaaaataaaggctagtcggttatcaacttgaa

aaagtgTTTTTTT

(N = guide sequence; first underline = tracr mate sequence; second underline = tracr sequence; bold = terminator)

Chimeric RNA containing +85 tracr RNA (*S. pyogenes* SF370) (SEQ ID NO: 58)

gagggcctatttcccatgattccttcatatttgcataacgatacaaggctgtagagagataat
tggaattaatttgactgtaaacacaaagatattagtaaaaaacgtgacgtagaaagtaataatt
tcttgggtagtttgcagttttaaattatgttttaaaatggactatcatatgcttaccgtaacttg
aaagtatttcgatttcttggctttatatacttgtggaaggacgaaacaccNNNNNNNNNNNNNN

NNNNNNqtttttagagctagaatagcaagttaaaataaaggctagtcggttatcaacttgaaaaagtg

gcaccgagtcggtgcTTTTTTT

(N = guide sequence; first underline = tracr mate sequence; second underline = tracr sequence; bold = terminator)
CBh-NLS-SpCas9-NLS

(SEQ ID NO: 59)

CGTTACATAACTTACGGTAAATGGCCCCCTGGCTGACCGCCCAACGACC
CCCGCCCATGACGTCAATAATGACGTATGTTCCCATAGTAACGCCAATAGGGACTT
TCCATTGACGTCAATGGGTGGAGTATTTACGGTAACTGCCCACTGGCAGTACATC
AAGTGTATCATATGCCAAGTACGCCCTTATGACGTCAATGACGGTAAATGGCCCCG
CCTGGCATTATGCCAGTACATGACCTTATGGGACTTTCTACTTGGCAGTACATCTA
CGTATTAGTCATCGTATTACCATGGTCGAGGTGAGCCCCAGTTCTGCTTCACTCTC
CCCATCTCCCCCTCCCCACCCCAATTTGTATTTATTTATTTTTAATTATTTTG
TGCAGCGATGGGGCGGGGGGGGGGGGGGGCGCGCCAGGCGGGCGGGGGCGG
GGCGAGGGCGGGCGGGCGAGCGGAGAGGTGCGCGGCAGCCAATCAGAGCG
GCGCGCTCCGAAAGTTTCTTTTATGGCGAGGCGCGCGGCGGCGCCCTATAAA
AAGCGAAGCGCGGGCGGGGAGTTCGCTGCGACGCTGCCTTCGCCCCGTGCCCC
GCTCCGCCCGCCTCGCGCCCGCCCGGCTCTGACTGACCGGTTACTCCAC
AGGTGAGCGGGCGGACGCGCTTCTCCTCCGGCTGTAATTAGCTGAGCAAGAGG
TAAGGGTTAAGGGATGGTTGGTTGGTGGGTATTAATGTTAATTACCTGGAGCAC
CTGCCTGAAATCACTTTTTTTCAGGTTGaccggtgccaccATGGACTATAAGGACCACGA
CGGAGACTACAAGGATCATGATATTGATTACAAAGACGATGACGATAAGATGGCCC
CAAAGAAGAAGCGGAAGGTCCGATCCACGGAGTCCAGCAGCCGACAGAAGTA
CAGCATCGGCCTGGACATCGGCACCAACTCTGTGGGCTGGGCGGTGATCACCGACG
AGTACAAGGTGCCAGCAAGAAATCAAGGTGCTGGCAACACCGACCGGCACAGC
ATCAAGAAGAACCTGATCGGAGCCCTGCTGTTTCGACAGCGGCGAAGACGCGGAGC
CACCCGGCTGAAGAGAACCAGCAGAAAGATACACCAGACGGAAGAACCAGGATC
TGCTATCTGCAAGAGATCTTCAGCAACGAGATGGCCAAGGTGGACGACAGCTTCTTC
CACAGACTGGAAGAGTCTTCTGTTGGAAGAGGATAAGAAGCACGAGCGGCACCC
CATCTTCGGCAACATCGTGGACGAGGTGGCCTACCACGAGAAGTACCCACCATCT
ACCACCTGAGAAAGAACTGGTGGACAGCACCGCAAGGCCGACCTGCGGCTGATC
TATCTGGCCCTGGCCACATGATCAAGTTCGGGGCCACTTCTGATCGAGGGCGAC
CTGAACCCCGACAACAGCGAGCTGGACAGCTGTTTCATCCAGCTGGTGCAGACCTA

- continued

CAACCAGCTGTTTCGAGGAAAACCCCATCAACGCCAGCGGCGTGGACGCCAAGGCCA
TCCTGTCTGCCAGACTGAGCAAGAGCAGACGGCTGGAAAATCTGATCGCCCAGCTG
CCCGGCGAGAAGAAGAATGGCCTGTTCGGCAACCTGATTGCCCTGAGCCTGGGCCT
GACCCCCAACTTCAAGAGCAACTTCGACCTGGCCGAGGATGCCAACTGCAGCTGA
GCAAGGACACCTACGACGACGACCTGGACAACCTGCTGGCCAGATCGGCGACCAG
TACGCCGACCTGTTTCTGGCCGCCAAGAACCTGTCCGACGCCATCCTGCTGAGCGAC
ATCCTGAGAGTGAACACCGAGATCACC AAGGCCCCCTGAGCGCCTCTATGATCAA
GAGATACGACGAGCACACCAGGACCTGACCTGTGAAAGCTCTCGTGGCGCAGC
AGCTGCCTGAGAAGTACAAAGAGATTTTCTTCGACCAGAGCAAGAACGGCTACGCC
GGCTACATTGACGGCGGAGCCAGCCAGGAAGAGTCTACAAGTTCATCAAGCCCAT
CCTGGAAAAGATGGACGGCACCGAGGAAGTCTCGTGAAGCTGAACAGAGAGGAC
CTGCTGCGGAAGCAGCGGACCTTCGACAACGGCAGCATCCCCACCAGATCCACCT
GGGAGAGCTGCACGCCATTCTGCGGCGCAGGAAGATTTTACCATTCTGAAGG
ACAACCGGAAAAGATCGAGAAGATCCTGACCTTCGCATCCCCACTACGTGGGC
CCTCTGGCCAGGGGAACAGCAGATTCGCTGGATGACCAGAAAGAGCGAGGAAA
CCATCACCCCCTGAACTTCGAGGAAGTGGTGGACAAGGGCGCTTCGCCCCAGAGC
TTATCGAGCGGATGACCAACTTCGATAAGAACC TGCCAACGAGAAGGTGCTGCC
CAAGCACAGCCTGCTGTACGAGTACTTCACCGTGATAACGAGCTGACCAAAGTGA
AATACGTGACCAGGGAATGAGAAAAGCCCGCCTTCTGAGCGGCAGCAGAAAAA
GGCCATCGTGGACCTGCTGTTCAAGACCAACCGAAAGTGACCGTGAAGCAGCTGA
AAGAGGACTACTTCAAGAAAATCGAGTGCTTCGACTCCGTGGAAATCTCCGGCGTG
GAAGATCGGTTCAACGCCTCCCTGGGCACATACCAGATCTGCTGAAAATTATCAAG
GACAAGGACTTCCTGGCAATGAGGAAAACGAGGACATTCGGAAGATATCGTGCT
GACCCTGACACTGTTTGAGGACAGAGAGATGATCGAGGAACGGCTGAAAACCTATG
CCCACCTGTTTCGACGACAAAGTGATGAAGCAGCTGAAGCGGCGAGATACACCGGC
TGGGGCAGGCTGAGCCGGAAGCTGATCAACGGCATCCGGGACAAGCAGTCCGGCAA
GACAATCCTGGATTTCCCTGAAGTCCGACGGCTTCGCCAACAGAACTTCATGCAGCT
GATCCACGACGACAGCCTGACCTTTAAAGAGGACATCCAGAAAGCCAGGTGTCCG
GCCAGGGCGATAGCCTGCACGAGCACATTGCCAATCTGGCCGGCAGCCCCGCCATT
AAGAAGGGCATCCTGCAGACAGTGAAGGTGGTGGACGAGCTCGTGAAGTGTGGG
CCGGCACAAAGCCCGAAGACATCGTGATCGAAATGGCCAGAGAGAACCAGACCACCC
AGAAGGGACAGAAGAACAGCCCGGAGAGAATGAAGCGGATCGAAGAGGGCATCAA
AGAGCTGGGCAGCCAGATCCTGAAAGAACCCCCGTGAAAACACCCAGCTGCAGA

- continued

ACGAGAAGCTGTACCTGTACTACCTGCAGAATGGGCGGGATATGTACGTGGACCAG
GAACTGGACATCAACCCGGCTGTCCGACTACGATGTGGACCATATCGTGCCTCAGAG
CTTTCTGAAGGACGACTCCATCGACAACAAGGTGCTGACCAGAAGCGACAAGAACC
GGGGCAAGAGCGACAACGTGCCCTCCGAAGAGGTCGTGAAGAAGATGAAGAATA
CTGGCGGCAGCTGCTGAACGCCAAGCTGATTACCCAGAGAAAAGTTCGACAATCTGA
CCAAGGCCGAGAGAGGCGCCCTGAGCGAACTGGATAAGGCCGGCTTCATCAAGAG
ACAGCTGGTGGAAACCCGGCAGATCACAAAGCACGTGGCACAGATCCTGGACTCCC
GGATGAACACTAAGTACGACGAGAAATGACAAGCTGATCCGGGAAGTGAAAGTGATC
ACCTGAAGTCCAAGCTGGTGTCCGATTTCCGGAAGGATTTCCAGTTTACAAAGTG
CGCGAGATCAACAACCTACCACCACGCCACGACGCTACCTGAACGCCGTCTGGG
AACCGCCCTGATCAAAAAGTACCCTAAGCTGGAAAGCGAGTTCGTGTACGGCGACT
ACAAGGTGTACGACGTGCGGAAGATGATCGCCAAGAGCGAGCAGGAAATCGGCAA
GGCTACCGCCAAGTACTTCTTCTACAGCAACATCATGAACTTTTCAAGACCGAGAT
TACCTGGCCAAACGGCGAGATCCGGAAGCGGCCTCTGATCGAGACAAACGGCGAAA
CCGGGGAGATCGTGTGGGATAAGGGCCGGGATTTTGCCACCGTGCGGAAAGTGCTG
AGCATGCCCAAGTGAATATCGTGAAAAGACCGAGGTGCAGACAGGCGGCTTCAG
CAAAGAGTCTATCTGCCCAAGAGGAACAGCGATAAGCTGATCGCCAGAAAGAAGG
ACTGGGACCCTAAGAAGTACGGCGGCTTCGACAGCCACCGTGGCTATTCTGTGC
TGGTGGTGGCCAAAGTGGAAAAGGGCAAGTCCAAGAACTGAAGAGTGTGAAAGA
GCTGCTGGGGATCACCATCATGGAAGAAGCAGCTTCGAGAAGAATCCCATCGACT
TTCTGGAAGCCAAAGGCTACAAGAAGTGAAAAGGACCTGATCATCAAGCTGCCT
AAGTACTCCCTGTTTCGAGCTGAAAACGGCCGGAAGAGAATGCTGGCCTCTGCCG
CGAATGCAGAAAGGAAACGAACTGGCCCTGCCCTCCAAATATGTGAACTTCTGT
ACCTGGCCAGCCACTATGAGAAGCTGAAGGGCTCCCCGAGGATAATGAGCAGAAA
CAGCTGTTTGTGGAACAGCACAAAGCACTACCTGGACGAGATCATCGAGCAGATCAG
CGAGTTCTCCAAAGAGAGTGATCCTGGCCGACGCTAATCTGGACAAAGTGCTGTCCGC
CTACAACAAGCACCCGGGATAAGCCCATCAGAGAGCAGGCCGAGAATATCATCCACC
TGTTTACCCTGACCAATCTGGGAGCCCTGCCGCTTCAAGTACTTTGACACCACCA
TCGACCGGAAGAGGTACACCAGCACCAAGAGGTGCTGGACGCCACCTGATCCAC
CAGAGCATCACCGCCTGTACGAGACACGGATCGACCTGTCTCAGCTGGGAGCGA
CTTTCTTTTCTTAGCTTGACCAGCTTTCTTAGTAGCAGCAGGACGCTTTAA
 (underline = NLS-hSpCas9-NLS)

[0237] Example chimeric RNA for *S. thermophilus* LMD-9 CRISPR1 Cas9 (with PAM of NNAGAAW)

[0238] Example chimeric RNA for *S. thermophilus* LMD-9 CRISPR1 Cas9 (with PAM of NNAGAAW)

(SEQ ID NO: 21)
 NNNNNNNNNNNNNNNNNNNgtttttgtactctcaagatttaGAAAtaaa
tcttgcagaagctacaaaqataaaggcttcatgccgaaatcaacaccctgt
cattttatggcagggtgttttcgttatttaaTTTTT
 (N = guide sequence; first underline = tracr
 mate sequence; second underline = tracr sequence;
 bold = terminator)

(SEQ ID NO: 22)
 NNNNNNNNNNNNNNNNNNNgtttttgtactctcaGAAAtgcagaagcta
caaaqataaaggcttcatgccgaaatcaacaccctgtcattttatggcagg
gtgttttcgttatttaaTTTTT
 (N = guide sequence; first underline = tracr
 mate sequence; second underline = tracr sequence;
 bold = terminator)

[0239] Example chimeric RNA for *S. thermophilus* LMD-9 CRISPR1 Cas9 (with PAM of NNAGAAW)

(SEQ ID NO: 23)
NNNNNNNNNNNNNNNNNNNNgtttttgtactctcaGAAAtgcagaagcta
caaaqataaggcttcatgccgaaatcaacaccctgtcattttatggcagg
gtgtTTTTTT
(N = guide sequence; first underline = tracr
mate sequence; second underline = tracr sequence;
bold = terminator)

[0240] Example chimeric RNA for *S. thermophilus* LMD-9 CRISPR1 Cas9 (with PAM of NNAGAAW)

(SEQ ID NO: 60)
NNNNNNNNNNNNNNNNNNNNgttattgtactctcaagatttaGAAAtaaa
tcttgcagaagctacaaaagataaggcttcatgccgaaatcaacaccctgt
cattttatggcagggtgttttcggtatttaaTTTTTT
(N = guide sequence; first underline = tracr
mate sequence; second underline = tracr sequence;
bold = terminator)

[0241] Example chimeric RNA for *S. thermophilus* LMD-9 CRISPR1 Cas9 (with PAM of NNAGAAW)

(SEQ ID NO: 61)
NNNNNNNNNNNNNNNNNNNNgttattgtactctcaGAAAtgcagaagcta
caaaqataaggcttcatgccgaaatcaacaccctgtcattttatggcagg
gtgttttcggtatttaaTTTTTT
(N = guide sequence; first underline = tracr
mate sequence; second underline = tracr sequence;
bold = terminator)

[0242] Example chimeric RNA for *S. thermophilus* LMD-9 CRISPR1 Cas9 (with PAM of NNAGAAW)

(SEQ ID NO: 62)
NNNNNNNNNNNNNNNNNNNNgttattgtactctcaGAAAtgcagaagcta
caaaqataaggcttcatgccgaaatcaacaccctgtcattttatggcagg
gtgtTTTTTT
(N = guide sequence; first underline = tracr
mate sequence; second underline = tracr sequence;
bold = terminator)

[0243] Example chimeric RNA for *S. thermophilus* LMD-9 CRISPR1 Cas9 (with PAM of NNAGAAW)

(SEQ ID NO: 63)
NNNNNNNNNNNNNNNNNNNNgttattgtactctcaagatttaGAAAtaaa
tcttgcagaagctacaaatgataaggcttcatgccgaaatcaacaccctgt
cattttatggcagggtgttttcggtatttaaTTTTTT
(N = guide sequence; first underline = tracr
mate sequence; second underline = tracr sequence;
bold = terminator)

[0244] Example chimeric RNA for *S. thermophilus* LMD-9 CRISPR1 Cas9 (with PAM of NNAGAAW)

(SEQ ID NO: 64)
NNNNNNNNNNNNNNNNNNNNgttattgtactctcaGAAAtgcagaagcta
caatgataaggcttcatgccgaaatcaacaccctgtcattttatggcagg
gtgttttcggtatttaaTTTTTT
(N = guide sequence; first underline = tracr
mate sequence; second underline = tracr sequence;
bold = terminator)

[0245] Example chimeric RNA for *S. thermophilus* LMD-9 CRISPR1 Cas9 (with PAM of NNAGAAW)

(SEQ ID NO: 65)
NNNNNNNNNNNNNNNNNNNNgttattgtactctcaGAAAtgcagaagcta
caatgataaggcttcatgccgaaatcaacaccctgtcattttatggcagg
gtgtTTTTTT
(N = guide sequence; first underline = tracr
mate sequence; second underline = tracr sequence;
bold = terminator)

[0246] Example chimeric RNA for *S. thermophilus* LMD-9 CRISPR3 Cas9 (with PAM of NGGNG)

(SEQ ID NO: 66)
NNNNNNNNNNNNNNNNNNNNgttttagagctgtgAAAcacagcgagtta
aaataaggcttagtccgtactcaactgaaaagggtggcaccgattccggtg
tTTTTTT
(N = guide sequence; first underline = tracr
mate sequence; second underline = tracr sequence;
bold = terminator)

[0247] Codon-optimized version of Cas9 from *S. thermo-philus* LMD-9 CRISPR3 locus (with an NLS at both 5' and 3' ends)

(SEQ ID NO: 67)
ATGAAAAGCCCGCGGCCACGAAAAGCCCGCCAGGCAAAAAGAAAA
GACCAAGCCCTACAGCATCGGCCTGGACATCGGCACCAATAGCTGGGCT
GGGCCGTGACCACCGACAACCTACAAGGTGCCAGCAAGAAAATGAAGGTG
CTGGGCAACACCTCCAAGAAGTACATCAAGAAAAACCTGCTGGGCGTGCT
GCTGTTTCGACAGCGGCATTACAGCCGAGGGCAGACGGCTGAAGAGAACC
CCAGACGGCGGTACACCCGCGGAGAAAACAGAATCCTGTATCTGCAAGAG
ATCTTCAGCACCGAGATGGCTACCCTGGACGACGCTTCTTCCAGCGGCT
GGACGACAGCTTCCTGGTGCCCGACGACAAGCGGACAGCAAGTACCCCA
TCTTCGGCAACCTGGTGAAGAGAAGGCCTACCACGACGAGTCCCCACC
ATCTACCACCTGAGAAAAGTACCTGGCCGACAGCAACAAGAAGGCCGACCT
GAGACTGGTGTATCTGGCCCTGGCCACATGATCAAGTACCGGGGCCACT
TCCTGATCGAGGGCGAGTTCAACAGCAAGAACAACGACATCCAGAAGAAC
TTCCAGGACTTCTTGACACCTACACGCCATCTTCGAGAGCGACCTGTC
CCTGAAAACAGCAAGCAGCTGGAAGAGATCGTGAAGGACAAGATCAGCA

-continued

AGCTGGAAAAGAAGGACCGCATCTCTGAAGCTGTTCCCGGCGAGAAGAAC
 AGCGGAATCTTACGCGAGTTTCTGAAGCTGATCGTGGCAACCAGGCCGA
 CTTCAGAAAAGTCTTCAACCTGGACGAGAAAGCCAGCCTGCACCTCAGCA
 AAGAGAGCTACGACGAGGACCTGGAAACCTGCTGGGATATATCGGGAC
 GACTACAGCGAGCTGTTCTGAAGGCCAAGAAGCTGTACGACGCTATCTT
 GCTGAGCGGCTTCTGACCGTGACCGACAACGAGACAGAGGCCCACTGA
 GCAGCGCCATGATTAAGCGGTACAACGAGCACAAAGAGGATCTGGCTCTG
 CTGAAAGAGTACATCCGGAACATCAGCCTGAAAACTACAATGAGGTGTT
 CAAGGACGACACCAAGAACGGCTACGCCGGCTACATCGACGCAAGACCA
 ACCAGGAAGATTTCTATGTGTACTGAAGAAGCTGCTGGCCGAGTTCGAG
 GGGGCCGACTACTTTCTGGAAAAATCGACCGCGAGGATTTCTGCGGAA
 GCAGCGGACCTTCGACAACGGCAGCATCCCCACAGATCCATCTGCAGG
 AAATGCGGGCCATCTGGACAAGCAGGCCAAGTTCTACCCATTCCTGGCC
 AAGAACAAGAGCGGATCGAGAAGATCTGACCTTCCGCATCCCTTACTA
 CGTGGCCCCCTGGCCAGAGGCAACAGCGATTTTGCCTGGTCCATCCGGA
 AGCGCAATGAGAAGATCACCCCTGGAACCTCGAGGACGTGATCGACAAA
 GAGTCCAGCGCCGAGGCCCTTCAACCGGATGACCAGCTTCGACCTGTA
 CCTGCCGAGGAAAAGGTGCTGCCAAGCACAGCCTGCTGTACGAGACAT
 TCAATGTGTATAACGAGCTGACCAAAGTGGGTTTATCGCCGAGTCTATG
 CGGGACTACAGTTCCTGGACTCCAAGCAGAAAAAGGACATCGTGGCGCT
 GTACTTCAAGGACAAGCGGAAAGTACCGGATAAGGACATCATCGAGTACC
 TGCACGCCATCTACGGCTACGATGGCATCGAGCTGAAGGGCATCGAGAAG
 CAGTTCAACTCCAGCCTGAGCACATACCACGACCTGCTGAACATTATCAA
 CGACAAGAATTTCTGGACGACTCCAGCAACGAGGCCATCATCGAAGAGA
 TCATCCACACCTTGACCATCTTTGAGGACCGCGAGATGATCAAGCAGCGG
 CTGAGCAAGTTCGAGAACATCTTCGACAAGAGCGTGTGAAAAGCTGAG
 CAGACGGCACTACACCGCTGGGGCAAGCTGAGCGCCAAGCTGATCAACG
 GCATCCGGGACGAGAAGTCCGGCAACACAATCTGGACTACCTGATCGAC
 GACGGCATCAGCAACCGGAACCTCATGCAGCTGATCCAGCAGCAGCCCT
 GAGCTTCAAGAAGAAGATCCAGAAGGCCAGATCATCGGGGACGAGGACA
 AGGGCAACATCAAAGAAGTCTGTAAGTCCCTGCCCGGACGCCGCCATC
 AAGAAGGGAATCTGCAGAGCATCAAGATCGTGGACGAGCTCGTGAAAGT
 GATGGCGGCGAGAAAGCCGAGAGCATCGTGGTGGAAATGGCTAGAGAGA
 ACCAGTACCAATCAGGGCAAGAGCAACAGCCAGCAGAGACTGAAGAGA
 CTGGAAGAGTCCCTGAAAGAGCTGGGCAGCAAGATTCTGAAAGAGAATAT
 CCCTGCCAAGCTGTCCAAGATCGACAACAACGCCCTGCAGAACGACCGGC
 TGTACCTGTACTACCTGCAGAAATGGCAAGGACATGTATACAGGCGACGAC
 CTGGATATCGACCGCTGAGCAACTACGACATCGACCATATTATCCCCCA
 GGCCTTCTGAAAGACAACAGCATTGACAACAAGTGTGGTGTCTCTCCG

-continued

CCAGCAACCGCGGCAAGTCCGATGATGTGCCAGCCTGGAAGTCTGTGAAA
 AAGAGAAAGACCTTCTGGTATCAGCTGCTGAAAAGCAAGCTGATTAGCCA
 GAGGAAGTTCGACACCTGACCAAGGCCGAGAGAGGCCCTGAGCCCTG
 AAGATAAGGCCGGCTTCATCCAGAGACAGCTGGTGGAAACCGGCAGATC
 ACCAAGCAGTGGCCAGACTGCTGGATGAGAAGTTTAAACAACAAGAAGGA
 CGAGAACAACCGGGCGTGGGACCGTGAAGATCATCACCTGAAAGTCCA
 CCTGGTGTCCAGTTCGGAAAGGACTTCGAGCTGTATAAAGTGCAGGAG
 ATCAATGACTTTCACACGCCACGACGCTTACCTGAATGCCGTGGTGGC
 TTCCGCCCTGCTGAAGAAGTACCCTAAGCTGGAACCCGAGTTCGTGTACG
 GCGACTACCCCAAGTACAACCTCTTCAGAGAGCGGAAGTCCGCCACCGAG
 AAGGTGACTTCTACTCCAACATCATGAATATCTTTAAGAAGTCCATCTC
 CCTGGCCGATGGCAGAGTGTGAGCGGCCCTGATCGAAGTGAACGAAG
 AGACAGGCGAGAGCGTGTGGAACAAGAAAGCGACCTGGCCACCGTGGCG
 CGGGTGTGAGTTATCTCAAGTGAATGCTGTAAGAAGGTGGAAGAACA
 GAACCACGGCCTGGATCGGGCAAGCCAAGGGCTGTTCAACGCCAACCC
 TGTCAGCAAGCCTAAGCCCAACTCCAACGAGAATCTCGTGGGGCCAAA
 GAGTACCTGGACCTAAGAAGTACGGCGGATACGCCGGCATCTCCAATAG
 CTTACCGTGTCTGTAAGGGCACAAATCGAGAAGGGCCTAAGAAAAAGA
 TCACAAACGTGCTGGAATTCAGGGGATCTCTATCCTGGACCGGATCAAC
 TACCGGAAGGATAAGCTGAACCTTCTGCTGGAAGAAAGGCTACAAGGACAT
 TGAGCTGATTATCGAGCTGCCCTAAGTACTCCCTGTTTCAAGTGAAGCAG
 GCTCCAGACGGATGCTGGCCTCCATCTGTCCACCAACAACAAGCGGGGC
 GAGATCCACAAGGGAAACCAGATCTTCTGAGCCAGAAATTTGTGAAACT
 GCTGTACCACGCCAAGCGGATCTCCAACACCATCAATGAGAACCACCGGA
 AATACGTGGAAGAACACAAGAAAGAGTTTGAAGAACTGTTCTACTACATC
 CTGGAGTTCAACGAGAACTATGTGGGACCAAGAAGAAGCGCAACTGCT
 GAATCCGCTTCCAGAGCTGGCAGAACCACAGCATCGACGAGCTGTGCA
 GCTCCTTATCGGCCCTACCGGACGAGCGGAAGGGACTGTTTGAAGCTG
 ACCTCCAGAGGCTTCCGCGGACTTTGAGTTCCTGGGAGTGAAGATCCC
 CCGGTACAGAGACTACACCCCTCTAGTCTGCTGAAGGACGCCACCTGTA
 TCCACCAGAGCGTGAACCGCTGTACGAAACCGGATCGACCTGGCTAAG
 CTGGGCGAGGAAAGCGTCTGCTGCTACTAAGAAAGCTGGTCAAGCTAA
 GAAAAAGAAATAA

Example 5

RNA-Guided Editing of Bacterial Genomes using CRISPR-Cas Systems

[0248] Applicants used the CRISPR-associated endonuclease Cas9 to introduce precise mutations in the genomes of *Streptococcus pneumoniae* and *Escherichia coli*. The approach relied on Cas9-directed cleavage at the targeted site to kill unmutated cells and circumvented the need for

selectable markers or counter-selection systems. Cas9 specificity was reprogrammed by changing the sequence of short CRISPR RNA (crRNA) to make single- and multi-nucleotide changes carried on editing templates. Simultaneous use of two crRNAs enabled multiplex mutagenesis. In *S. pneumoniae*, nearly 100% of cells that survived Cas9 cleavage contained the desired mutation, and 65% when used in combination with recombineering in *E. coli*. Applicants exhaustively analyzed Cas9 target requirements to define the range of targetable sequences and showed strategies for editing sites that do not meet these requirements, suggesting the versatility of this technique for bacterial genome engineering.

[0249] The understanding of gene function depends on the possibility of altering DNA sequences within the cell in a controlled fashion. Site-specific mutagenesis in eukaryotes is achieved by the use of sequence-specific nucleases that promote homologous recombination of a template DNA containing the mutation of interest. Zinc finger nucleases (ZFNs), transcription activator-like effector nucleases (TALENs) and homing meganucleases can be programmed to cleave genomes in specific locations, but these approaches require engineering of new enzymes for each target sequence. In prokaryotic organisms, mutagenesis methods either introduce a selection marker in the edited locus or require a two-step process that includes a counter-selection system. More recently, phage recombination proteins have been used for recombineering, a technique that promotes homologous recombination of linear DNA or oligonucleotides. However, because there is no selection of mutations, recombineering efficiency can be relatively low (0.1-10% for point mutations down to 10^{-5} - 10^{-6} for larger modifications), in many cases requiring the screening of a large number of colonies. Therefore new technologies that are affordable, easy to use and efficient are still in need for the genetic engineering of both eukaryotic and prokaryotic organisms.

[0250] Recent work on the CRISPR (clustered, regularly interspaced, short palindromic repeats) adaptive immune system of prokaryotes has led to the identification of nucleases whose sequence specificity is programmed by small RNAs. CRISPR loci are composed of a series of repeats separated by 'spacer' sequences that match the genomes of bacteriophages and other mobile genetic elements. The repeat-spacer array is transcribed as a long precursor and processed within repeat sequences to generate small crRNA that specify the target sequences (also known as protospacers) cleaved by CRISPR systems. Essential for cleavage is the presence of a sequence motif immediately downstream of the target region, known as the protospacer-adjacent motif (PAM). CRISPR-associated (cas) genes usually flank the repeat-spacer array and encode the enzymatic machinery responsible for crRNA biogenesis and targeting. Cas9 is a dsDNA endonuclease that uses a crRNA guide to specify the site of cleavage. Loading of the crRNA guide onto Cas9 occurs during the processing of the crRNA precursor and requires a small RNA antisense to the precursor, the tracrRNA, and RNase III. In contrast to genome editing with ZFNs or TALENs, changing Cas9 target specificity does not require protein engineering but only the design of the short crRNA guide.

[0251] Applicants recently showed in *S. pneumoniae* that the introduction of a CRISPR system targeting a chromosomal locus leads to the killing of the transformed cells. It

was observed that occasional survivors contained mutations in the target region, suggesting that Cas9 dsDNA endonuclease activity against endogenous targets could be used for genome editing. Applicants showed that marker-less mutations can be introduced through the transformation of a template DNA fragment that will recombine in the genome and eliminate Cas9 target recognition. Directing the specificity of Cas9 with several different crRNAs allows for the introduction of multiple mutations at the same time. Applicants also characterized in detail the sequence requirements for Cas9 targeting and show that the approach can be combined with recombineering for genome editing in *E. coli*.

[0252] RESULTS: Genome Editing by Cas9 Cleavage of a Chromosomal Target

[0253] *S. pneumoniae* strain crR6 contains a Cas9-based CRISPR system that cleaves a target sequence present in the bacteriophage ϕ 8232.5. This target was integrated into the *srtA* chromosomal locus of a second strain R6^{8232.5}. An altered target sequence containing a mutation in the PAM region was integrated into the *srtA* locus of a third strain R6^{370.1}, rendering this strain 'immune' to CRISPR cleavage (FIG. 28a). Applicants transformed R6^{8232.5} and R6^{370.1} cells with genomic DNA from crR6 cells, expecting that successful transformation of R6^{8232.5} cells should lead to cleavage of the target locus and cell death. Contrary to this expectation, Applicants isolated R6^{8232.5} transformants, albeit with approximately 10-fold less efficiency than R6^{370.1} transformants (FIG. 28b). Genetic analysis of eight R6^{8232.5} transformants (FIG. 28) revealed that the great majority are the product of a double recombination event that eliminates the toxicity of Cas9 targeting by replacing the ϕ 8232.5 target with the crR6 genome's wild-type *srtA* locus, which does not contain the protospacer required for Cas9 recognition. These results were proof that the concurrent introduction of a CRISPR system targeting a genomic locus (the targeting construct) together with a template for recombination into the targeted locus (the editing template) led to targeted genome editing (FIG. 23a).

[0254] To create a simplified system for genome editing, Applicants modified the CRISPR locus in strain crR6 by deleting *cas1*, *cas2* and *csn2*, genes which have been shown to be dispensable for CRISPR targeting, yielding strain crR6M (FIG. 28a). This strain retained the same properties of crR6 (FIG. 28b). To increase the efficiency of Cas9-based editing and demonstrate that a template DNA of choice can be used to control the mutation introduced, Applicants co-transformed R6^{8232.5} cells with PCR products of the wild-type *srtA* gene or the mutant R6^{370.1} target, either of which should be resistant to cleavage by Cas9. This resulted in a 5- to 10-fold increase of the frequency of transformation compared with genomic crR6 DNA alone (FIG. 23b). The efficiency of editing was also substantially increased, with 8/8 transformants tested containing a wild-type *srtA* copy and 7/8 containing the PAM mutation present in the R6^{370.1} target (FIG. 23b and FIG. 29a). Taken together, these results showed the potential of genome editing assisted by Cas9.

[0255] Analysis of Cas9 target requirements: To introduce specific changes in the genome, one must use an editing template carrying mutations that abolish Cas9-mediated cleavage, thereby preventing cell death. This is easy to achieve when the deletion of the target or its replacement by another sequence (gene insertion) is sought. When the goal is to produce gene fusions or to generate single-nucleotide

mutations, the abolishment of Cas9 nuclease activity will only be possible by introducing mutations in the editing template that alter either the PAM or the protospacer sequences. To determine the constraints of CRISPR-mediated editing, Applicants performed an exhaustive analysis of PAM and protospacer mutations that abrogate CRISPR targeting.

[0256] Previous studies proposed that *S. pyogenes* Cas9 requires an NGG PAM immediately downstream of the protospacer. However, because only a very limited number of PAM-inactivating mutations have been described so far, Applicants conducted a systematic analysis to find all 5-nucleotide sequences following the protospacer that eliminate CRISPR cleavage. Applicants used randomized oligonucleotides to generate all possible 1,024 PAM sequences in a heterogeneous PCR product that was transformed into crR6 or R6 cells. Constructs carrying functional PAMs were expected to be recognized and destroyed by Cas9 in crR6 but not R6 cells (FIG. 24a). More than 2×10^5 colonies were pooled together to extract DNA for use as template for the co-amplification of all targets. PCR products were deep sequenced and found to contain all 1,024 sequences, with coverage ranging from 5 to 42,472 reads (See section "Analysis of deep sequencing data"). The functionality of each PAM was estimated by the relative proportion of its reads in the crR6 sample over the R6 sample. Analysis of the first three bases of the PAM, averaging over the two last bases, clearly showed that the NGG pattern was under-represented in crR6 transformants (FIG. 24b). Furthermore, the next two bases had no detectable effect on the NGG PAM (See section "Analysis of deep sequencing data"), demonstrating that the NNGGN sequence was sufficient to license Cas9 activity. Partial targeting was observed for NAG PAM sequences (FIG. 24b). Also the NNGGN pattern partially inactivated CRISPR targeting (Table G), indicating that the NGG motif can still be recognized by Cas9 with reduced efficiency when shifted by 1 bp. These data shed light onto the molecular mechanism of Cas9 target recognition, and they revealed that NGG (or CCN on the complementary strand) sequences are sufficient for Cas9 targeting and that NGG to NAG or NNGGN mutations in the editing template should be avoided. Owing to the high frequency of these tri-nucleotide sequences (once every 8 bp), this means that almost any position of the genome can be edited. Indeed, Applicants tested ten randomly chosen targets carrying various PAMs and all were found to be functional (FIG. 30).

[0257] Another way to disrupt Cas9-mediated cleavage is to introduce mutations in the protospacer region of the editing template. It is known that point mutations within the 'seed sequence' (the 8 to 10 protospacer nucleotides immediately adjacent to the PAM) can abolish cleavage by CRISPR nucleases. However, the exact length of this region is not known, and it is unclear whether mutations to any nucleotide in the seed can disrupt Cas9 target recognition. Applicants followed the same deep sequencing approach described above to randomize the entire protospacer sequence involved in base pair contacts with the crRNA and to determine all sequences that disrupt targeting. Each position of the 20 matching nucleotides (14) in the *spc1* target present in R6^{8232.5} cells (FIG. 23a) was randomized and transformed into crR6 and R6 cells (FIG. 24a). Consistent with the presence of a seed sequence, only mutations in the 12 nucleotides immediately upstream of the PAM abrogated cleavage by Cas9 (FIG. 24c). However, different

mutations displayed markedly different effects. The distal (from the PAM) positions of the seed (12 to 7) tolerated most mutations and only one particular base substitution abrogated targeting. In contrast, mutations to any nucleotide in the proximal positions (6 to 1, except 3) eliminated Cas9 activity, although at different levels for each particular substitution. At position 3, only two substitutions affected CRISPR activity and with different strength. Applicants concluded that, although seed sequence mutations can prevent CRISPR targeting, there are restrictions regarding the nucleotide changes that can be made in each position of the seed. Moreover, these restrictions can most likely vary for different spacer sequences. Therefore Applicants believe that mutations in the PAM sequence, if possible, should be the preferred editing strategy. Alternatively, multiple mutations in the seed sequence may be introduced to prevent Cas9 nuclease activity.

[0258] Cas9-mediated genome editing in *S. pneumoniae*: To develop a rapid and efficient method for targeted genome editing, Applicants engineered strain crR6Rk, a strain in which spacers can be easily introduced by PCR (FIG. 33). Applicants decided to edit the β -galactosidase (*bgaA*) gene of *S. pneumoniae*, whose activity can be easily measured. Applicants introduced alanine substitutions of amino acids in the active site of this enzyme: R481A (R→A) and N563A,E564A (NE→AA) mutations. To illustrate different editing strategies, Applicants designed mutations of both the PAM sequence and the protospacer seed. In both cases the same targeting construct with a crRNA complementary to a region of the β -galactosidase gene that is adjacent to a TGG PAM sequence (CCA in the complementary strand, FIG. 26) was used. The R→A editing template created a three-nucleotide mismatch on the protospacer seed sequence (CGT to GCA, also introducing a BtgZI restriction site). In the NE→AA editing template Applicants simultaneously introduced a synonymous mutation that created an inactive PAM (TGG to TTG) along with mutations that are 218 nt downstream of the protospacer region (AAT GAA to GCT GCA, also generating a TseI restriction site). This last editing strategy demonstrated the possibility of using a remote PAM to make mutations in places where a proper target may be hard to choose. For example, although the *S. pneumoniae* R6 genome, which has a 39.7% GC content, contains on average one PAM motif every 12 bp, some PAM motifs are separated by up to 194 bp (FIG. 33). In addition Applicants designed a *AbgaA* in-frame deletion of 6,664 bp. In all three cases, co-transformation of the targeting and editing templates produced 10-times more kanamycin-resistant cells than co-transformation with a control editing template containing wild-type *bgaA* sequences (FIG. 25b). Applicants genotyped 24 transformants (8 for each editing experiment) and found that all but one incorporated the desired change (FIG. 25c). DNA sequencing also confirmed not only the presence of the introduced mutations but also the absence of secondary mutations in the target region (FIG. 29b,c). Finally, Applicants measured β -galactosidase activity to confirm that all edited cells displayed the expected phenotype (FIG. 25d).

[0259] Cas9-mediated editing can also be used to generate multiple mutations for the study of biological pathways. Applicants decided to illustrate this for the sortase-dependent pathway that anchors surface proteins to the envelope of Gram-positive bacteria. Applicants introduced a sortase deletion by co-transformation of a chloramphenicol-resis-

tant targeting construct and a *AsrtA* editing template (FIG. 33*a,b*), followed by a Δ *bgaA* deletion using a kanamycin-resistant targeting construct that replaced the previous one. In *S. pneumoniae*, β -galactosidase is covalently linked to the cell wall by sortase. Therefore, deletion of *srtA* results in the release of the surface protein into the supernatant, whereas the double deletion has no detectable β -galactosidase activity (FIG. 34*c*). Such a sequential selection can be iterated as many times as required to generate multiple mutations.

[0260] These two mutations may also be introduced at the same time. Applicants designed a targeting construct containing two spacers, one matching *srtA* and the other matching *bgaA*, and co-transformed it with both editing templates at the same time (FIG. 25*e*). Genetic analysis of transformants showed that editing occurred in 6/8 cases (FIG. 25*f*). Notably, the remaining two clones each contained either a Δ *srtA* or a Δ *bgaA* deletion, suggesting the possibility of performing combinatorial mutagenesis using Cas9. Finally, to eliminate the CRISPR sequences, Applicants introduced a plasmid containing the *bgaA* target and a spectinomycin resistance gene along with genomic DNA from the wild-type strain R6. Spectinomycin-resistant transformants that retain the plasmid eliminated the CRISPR sequences (FIG. 34*a,d*).

[0261] Mechanism and efficiency of editing: To understand the mechanisms underlying genome editing with Cas9, Applicants designed an experiment in which the editing efficiency was measured independently of Cas9 cleavage. Applicants integrated the *ermAM* erythromycin resistance gene in the *srtA* locus, and introduced a premature stop codon using Cas9-mediated editing (FIG. 33). The resulting strain (JEN53) contains an *ermAM*(stop) allele and is sensitive to erythromycin. This strain may be used to assess the efficiency at which the *ermAM* gene is repaired by measuring the fraction of cells that restore antibiotic resistance with or without the use of Cas9 cleavage. JEN53 was transformed with an editing template that restores the wild-type allele, together with either a kanamycin-resistant CRISPR construct targeting the *ermAM*(stop) allele (CRISPR:*ermAM*(stop)) or a control construct without a spacer (CRISPR: \emptyset) (FIG. 26*a,b*). In the absence of kanamycin selection, the fraction of edited colonies was on the order of 10^{-2} (erythromycin-resistant cfu/total cfu) (FIG. 26*c*), representing the baseline frequency of recombination without Cas9-mediated selection against unedited cells. However, if kanamycin selection was applied and the control CRISPR construct was co-transformed, the fraction of edited colonies increased to about 10^{-1} (kanamycin- and erythromycin-resistant cfu/kanamycin-resistant cfu) (FIG. 26*c*). This result shows that selection for the recombination of the CRISPR locus co-selected for recombination in the *ermAM* locus independently of Cas9 cleavage of the genome, suggesting that a subpopulation of cells is more prone to transformation and/or recombination. Transformation of the CRISPR:*ermAM*(stop) construct followed by kanamycin selection resulted in an increase of the fraction of erythromycin-resistant, edited cells to 99% (FIG. 26*c*). To determine if this increase is caused by the killing of non-edited cells, Applicants compared the kanamycin-resistant colony forming units (cfu) obtained after co-transformation of JEN53 cells with the CRISPR:*ermAM*(stop) or CRISPR: \emptyset constructs.

[0262] Applicants counted 5.3 times less kanamycin-resistant colonies after transformation of the *ermAM*(stop) construct ($2.5 \times 10^4 / 4.7 \times 10^3$, FIG. 35*a*), a result that suggests that indeed targeting of a chromosomal locus by Cas9 leads

to the killing of non-edited cells. Finally, because the introduction of dsDNA breaks in the bacterial chromosome is known to trigger repair mechanisms that increase the rate of recombination of the damaged DNA, Applicants investigated whether cleavage by Cas9 induces recombination of the editing template. Applicants counted 2.2 times more colonies after co-transformation with the CRISPR:*erm*(stop) construct than with the CRISPR: \emptyset construct (FIG. 26*d*), indicating that there was a modest induction of recombination. Taken together, these results showed that co-selection of transformable cells, induction of recombination by Cas9-mediated cleavage and selection against non-edited cells, each contributed to the high efficiency of genome editing in *S. pneumoniae*.

[0263] As cleavage of the genome by Cas9 should kill non-edited cells, one would not expect to recover any cells that received the kanamycin resistance-containing Cas9 cassette but not the editing template. However, in the absence of the editing template Applicants recovered many kanamycin-resistant colonies after transformation of the CRISPR:*ermAM*(stop) construct (FIG. 35*a*). These cells that ‘escape’ CRISPR-induced death produced a background that determined a limit of the method. This background frequency may be calculated as the ratio of CRISPR:*ermAM*(stop)/CRISPR: \emptyset cfu, 2.6×10^{-3} ($7.1 \times 10^1 / 2.7 \times 10^4$) in this experiment, meaning that if the recombination frequency of the editing template is less than this value, CRISPR selection may not efficiently recover the desired mutants above the background. To understand the origin of these cells, Applicants genotyped 8 background colonies and found that 7 contained deletions of the targeting spacer (FIG. 35*b*) and one harbored a presumably inactivating mutation in Cas9 (FIG. 35*c*).

[0264] Genome editing with Cas9 in *E. coli*: The activation of Cas9 targeting through the chromosomal integration of a CRISPR-Cas system is only possible in organisms that are highly recombinogenic. To develop a more general method that is applicable to other microbes, Applicants decided to perform genome editing in *E. coli* using a plasmid-based CRISPR-Cas system. Two plasmids were constructed: a pCas9 plasmid carrying the *tracrRNA*, Cas9 and a chloramphenicol resistance cassette (FIG. 36), and a pCRISPR kanamycin-resistant plasmid carrying the array of CRISPR spacers. To measure the efficiency of editing independently of CRISPR selection, Applicants sought to introduce an A to C transversion in the *rpsL* gene that confers streptomycin resistance. Applicants constructed a pCRISPR:*rpsL* plasmid harboring a spacer that would guide Cas9 cleavage of the wild-type, but not the mutant *rpsL* allele (FIG. 27*b*). The pCas9 plasmid was first introduced into *E. coli* MG1655 and the resulting strain was co-transformed with the pCRISPR:*rpsL* plasmid and W542, an editing oligonucleotide containing the A to C mutation. streptomycin-resistant colonies after transformation of the pCRISPR:*rpsL* plasmid were only recovered, suggesting that Cas9 cleavage induces recombination of the oligonucleotide (FIG. 37). However, the number of streptomycin-resistant colonies was two orders of magnitude lower than the number of kanamycin-resistant colonies, which are presumably cells that escape cleavage by Cas9. Therefore, in these conditions, cleavage by Cas9 facilitated the introduction of the mutation, but with an efficiency that was not enough to select the mutant cells above the background of ‘escapers’.

[0265] To improve the efficiency of genome editing in *E. coli*, Applicants applied their CRISPR system with recombineering, using Cas9-induced cell death to select for the desired mutations. The pCas9 plasmid was introduced into the recombineering strain HME63 (31), which contains the Gam, Exo and Beta functions of the λ -red phage. The resulting strain was co-transformed with the pCRISPR:rpsL plasmid (or a pCRISPR: \emptyset control) and the W542 oligonucleotide (FIG. 27a). The recombineering efficiency was 5.3×10^{-5} , calculated as the fraction of total cells that became streptomycin-resistant when the control plasmid was used (FIG. 27c). In contrast, transformation with the pCRISPR:rpsL plasmid increased the percentage of mutant cells to $65 \pm 14\%$ (FIGS. 27c and 29f). Applicants observed that the number of cfu was reduced by about three orders of magnitude after transformation of the pCRISPR:rpsL plasmid than the control plasmid ($4.8 \times 10^5 / 5.3 \times 10^2$, FIG. 38a), suggesting that selection results from CRISPR-induced death of non-edited cells. To measure the rate at which Cas9 cleavage was inactivated, an important parameter of Applicants' method, Applicants transformed cells with either pCRISPR:rpsL or the control plasmid without the W542 editing oligonucleotide (FIG. 38a). This background of CRISPR 'escapers', measured as the ratio of pCRISPR:rpsL/pCRISPR: \emptyset cfu, was 2.5×10^{-4} ($1.2 \times 10^2 / 4.8 \times 10^5$). Genotyping eight of these escapers revealed that in all cases there was a deletion of the targeting spacer (FIG. 38b). This background was higher than the recombineering efficiency of the rpsL mutation, 5.3×10^{-5} , which suggested that to obtain 65% of edited cells, Cas9 cleavage must induce oligonucleotide recombination. To confirm this, Applicants compared the number of kanamycin- and streptomycin-resistant cfu after transformation of pCRISPR:rpsL or pCRISPR: \emptyset (FIG. 27d). As in the case for *S. pneumoniae*, Applicants observed a modest induction of recombination, about 6.7 fold ($2.0 \times 10^{-4} / 3.0 \times 10^{-5}$). Taken together, these results indicated that the CRISPR system provided a method for selecting mutations introduced by recombineering.

[0266] Applicants showed that CRISPR-Cas systems may be used for targeted genome editing in bacteria by the co-introduction of a targeting construct that killed wild-type cells and an editing template that both eliminated CRISPR cleavage and introduced the desired mutations. Different types of mutations (insertions, deletions or scar-less single-nucleotide substitutions) may be generated. Multiple mutations may be introduced at the same time. The specificity and versatility of editing using the CRISPR system relied on several unique properties of the Cas9 endonuclease: (i) its target specificity may be programmed with a small RNA, without the need for enzyme engineering, (ii) target specificity was very high, determined by a 20 bp RNA-DNA interaction with low probability of non-target recognition, (iii) almost any sequence may be targeted, the only requirement being the presence of an adjacent NGG sequence, (iv) almost any mutation in the NGG sequence, as well as mutations in the seed sequence of the protospacer, eliminates targeting.

[0267] Applicants showed that genome engineering using the CRISPR system worked not only in highly recombinogenic bacteria such as *S. pneumoniae*, but also in *E. coli*. Results in *E. coli* suggested that the method may be applicable to other microorganisms for which plasmids may be introduced. In *E. coli*, the approach complements recombineering of mutagenic oligonucleotides. To use this method-

ology in microbes where recombineering is not a possible, the host homologous recombination machinery may be used by providing the editing template on a plasmid. In addition, because accumulated evidence indicates that CRISPR-mediated cleavage of the chromosome leads to cell death in many bacteria and archaea, it is possible to envision the use of endogenous CRISPR-Cas systems for editing purposes.

[0268] In both *S. pneumoniae* and *E. coli*, Applicants observed that although editing was facilitated by a co-selection of transformable cells and a small induction of recombination at the target site by Cas9 cleavage, the mechanism that contributed the most to editing was the selection against non-edited cells. Therefore the major limitation of the method was the presence of a background of cells that escape CRISPR-induced cell death and lack the desired mutation. Applicants showed that these 'escapers' arose primarily through the deletion of the targeting spacer, presumably after the recombination of the repeat sequences that flank the targeting spacer. Future improvements may focus on the engineering of flanking sequences that can still support the biogenesis of functional crRNAs but that are sufficiently different from one another to eliminate recombination. Alternatively, the direct transformation of chimeric crRNAs may be explored. In the particular case of *E. coli*, the construction of the CRISPR-Cas system was not possible if this organism was also used as a cloning host. Applicants solved this issue by placing Cas9 and the tracrRNA on a different plasmid than the CRISPR array. The engineering of an inducible system may also circumvent this limitation.

[0269] Although new DNA synthesis technologies provide the ability to cost-effectively create any sequence with a high throughput, it remains a challenge to integrate synthetic DNA in living cells to create functional genomes. Recently, the co-selection MAGE strategy was shown to improve the mutation efficiency of recombineering by selecting a subpopulation of cells that has an increased probability to achieve recombination at or around a given locus. In this method, the introduction of selectable mutations is used to increase the chances of generating nearby non-selectable mutations. As opposed to the indirect selection provided by this strategy, the use of the CRISPR system makes it possible to directly select for the desired mutation and to recover it with a high efficiency. These technologies add to the toolbox of genetic engineers, and together with DNA synthesis, they may substantially advance both the ability to decipher gene function and to manipulate organisms for biotechnological purposes. Two other studies also relate to CRISPR-assisted engineering of mammalian genomes. It is expected that these crRNA-directed genome editing technologies may be broadly useful in the basic and medical sciences.

[0270] Strains and culture conditions. *S. pneumoniae* strain R6 was provided by Dr. Alexander Tomasz. Strain crR6 was generated in a previous study. Liquid cultures of *S. pneumoniae* were grown in THYE medium (30g/l Todd-Hewitt agar, 5 g/l yeast extract). Cells were plated on tryptic soy agar (TSA) supplemented with 5% defibrinated sheep blood. When appropriate, antibiotics were added as followings: kanamycin (400 μ g/ml), chloramphenicol (5 μ g/ml), erythromycin (1 μ g/ml) streptomycin (100 μ g/ml) or spectinomycin (100 μ g/ml). Measurements of β -galactosidase activity were made using the Miller assay as previously described.

[0271] *E. coli* strains MG1655 and HME63 (derived from MG1655, A(argF-lac) U169 λ c1857 Δ cro-bioA galK tyr 145 UAG mutS \rightarrow amp) (31) were provided by Jeff Roberts and Donald Court, respectively. Liquid cultures of *E. coli* were grown in LB medium (Difco). When appropriate, antibiotics were added as followings: chloramphenicol (25 μ g/ml), kanamycin (25 μ g/ml) and streptomycin (50 μ g/ml).

[0272] *S. pneumoniae* transformation. Competent cells were prepared as described previously (23). For all genome editing transformations, cells were gently thawed on ice and resuspended in 10 volumes of M2 medium supplemented with 100 ng/ml of competence-stimulating peptide CSP1 (40), and followed by addition of editing constructs (editing constructs were added to cells at a final concentration between 0.7 ng/ μ l to 2.5 μ g/ μ l). Cells were incubated 20 min at 37° C. before the addition of 2 μ l of targeting constructs and then incubated 40 min at 37° C. Serial dilutions of cells were plated on the appropriate medium to determine the colony forming units (cfu) count.

[0273] *E. coli* Lambda-red recombineering. Strain HME63 was used for all recombineering experiments. Recombineering cells were prepared and handled according to a previously published protocol (6). Briefly, a 2 ml overnight culture (LB medium) inoculated from a single colony obtained from a plate was grown at 30° C. The overnight culture was diluted 100-fold and grown at 30° C. with shaking (200 rpm) until the OD600 is from 0.4-0.5 (approximately 3 hrs). For Lambda-red induction, the culture was transferred to a 42° C. water bath to shake at 200 rpm for 15 min. Immediately after induction, the culture was swirled in an ice-water slurry and chilled on ice for 5-10 min. Cells were then washed and aliquoted according to the protocol. For electro-transformation, 50 μ l of cells were mixed with 1 mM of salt-free oligos (IDT) or 100-150 ng of plasmid DNA (prepared by QIAprep Spin Miniprep Kit, Qiagen). Cells were electroporated using 1 mm Gene Pulser cuvette (Bio-rad) at 1.8 kV and were immediately resuspended in 1 ml of room temperature LB medium. Cells were recovered at 30° C. for 1-2 hrs before being plated on LB agar with appropriate antibiotic resistance and incubated at 32° C. overnight.

[0274] Preparation of *S. pneumoniae* genomic DNA. For transformation purposes, *S. pneumoniae* genomic DNA was extracted using the Wizard Genomic DNA Purification Kit, following instructions provided by the manufacturer (Promega). For genotyping purposes, 700 μ l of overnight *S. pneumoniae* cultures were pelleted, resuspended in 60 μ l of lysozyme solution (2 mg/ml) and incubated 30 min at 37° C. The genomic DNA was extracted using QIAprep Spin Miniprep Kit (Qiagen).

[0275] Strain construction. All primers used in this study are provided in Table G. To generate *S. pneumoniae* crR6M, an intermediate strain, LAM226, was made. In this strain the *aphA-3* gene (providing kanamycin resistance) adjacent to the CRISPR array of *S. pneumoniae* crR6 strain was replaced by a *cat* gene (providing chloramphenicol resistance). Briefly, crR6 genomic DNA was amplified using primers L448/L444 and L447/L481, respectively. The *cat* gene was amplified from plasmid pC194 using primers L445/L446. Each PCR product was gel-purified and all three were fused by SOEing PCR with primers L448/L481. The resulting PCR product was transformed into competent *S. pneumoniae* crR6 cells and chloramphenicol-resistant transformants were selected. To generate *S. pneumoniae* crR6M,

S. pneumoniae crR6 genomic DNA was amplified by PCR using primers L409/L488 and L448/L481, respectively. Each PCR product was gel-purified and they were fused by SOEing PCR with primers L409/L481. The resulting PCR product was transformed into competent *S. pneumoniae* LAM226 cells and kanamycin-resistant transformants were selected.

[0276] To generate *S. pneumoniae* crR6Rc, *S. pneumoniae* crR6M genomic DNA was amplified by PCR using primers L430/W286, and *S. pneumoniae* LAM226 genomic DNA was amplified by PCR using primers W288/L481. Each PCR product was gel-purified and they were fused by SOEing PCR with primers L430/L481. The resulting PCR product was transformed into competent *S. pneumoniae* crR6M cells and chloramphenicol-resistant transformants were selected.

[0277] To generate *S. pneumoniae* crR6Rk, *S. pneumoniae* crR6M genomic DNA was amplified by PCR using primers L430/W286 and W287/L481, respectively. Each PCR product was gel-purified and they were fused by SOEing PCR with primers L430/L481. The resulting PCR product was transformed into competent *S. pneumoniae* crR6Rc cells and kanamycin-resistant transformants were selected.

[0278] To generate JEN37, *S. pneumoniae* crR6Rk genomic DNA was amplified by PCR using primers L430/W356 and W357/L481, respectively. Each PCR product was gel-purified and they were fused by SOEing PCR with primers L430/L481. The resulting PCR product was transformed into competent *S. pneumoniae* crR6Rc cells and kanamycin-resistant transformants were selected.

[0279] To generate JEN38, R6 genomic DNA was amplified using primers L422/L461 and L459/L426, respectively. The *ermAM* gene (specifying erythromycin resistance) was amplified from plasmid pFW15⁴⁻³ using primers L457/L458. Each PCR product was gel-purified and all three were fused by SOEing PCR with primers L422/L426. The resulting PCR product was transformed into competent *S. pneumoniae* crR6Rc cells and erythromycin-resistant transformants were selected.

[0280] *S. pneumoniae* JEN53 was generated in two steps. First JEN43 was constructed as illustrated in FIG. 33. JEN53 was generated by transforming genomic DNA of JEN25 into competent JEN43 cells and selecting on both chloramphenicol and erythromycin.

[0281] To generate *S. pneumoniae* JEN62, *S. pneumoniae* crR6Rk genomic DNA was amplified by PCR using primers W256/W365 and W366/L403, respectively. Each PCR product was purified and ligated by Gibson assembly. The assembly product was transformed into competent *S. pneumoniae* crR6Rc cells and kanamycin-resistant transformants were selected.

[0282] Plasmid construction. pDB97 was constructed through phosphorylation and annealing of oligonucleotides B296/B297, followed by ligation in pLZ12spec digested by EcoRI/BamHI. Applicants fully sequenced pLZ12spec and deposited its sequence in genebank (accession: KC112384).

[0283] pDB98 was obtained after cloning the CRISPR leader sequence was cloned together with a repeat-spacer-repeat unit into pLZ12spec. This was achieved through amplification of crR6Rc DNA with primers B298/B320 and B299/B321, followed by SOEing PCR of both products and cloning in pLZ12spec with restriction sites BamHI/EcoRI. In this way the spacer sequence in pDB98 was engineered to contain two BsaI restriction sites in opposite directions that allow for the scar-less cloning of new spacers.

[0284] pDB99 to pDB108 were constructed by annealing of oligonucleotides B300/B301 (pDB99), B302/B303 (pDB100), B304/B305 (pDB101), B306/B307 (pDB102), B308/B309 (pDB103), B310/B311 (pDB104), B312/B313 (pDB105), B314/B315 (pDB106), B315/B317 (pDB107), B318/B319 (pDB108), followed by ligation in pDB98 cut by BsaI.

[0285] The pCas9 plasmid was constructed as follow. Essential CRISPR elements were amplified from *Streptococcus pyogenes* SF370 genomic DNA with flanking homology arms for Gibson Assembly. The tracrRNA and Cas9 were amplified with oligos HC008 and HC010. The leader and CRISPR sequences were amplified HC011/HC014 and HC015/HC009, so that two BsaI type IIS sites were introduced in between two direct repeats to facilitate easy insertion of spacers.

[0286] pCRISPR was constructed by subcloning the pCas9 CRISPR array in pZE21-MCS1 through amplification with oligos B298+B299 and restriction with EcoRI and BamHI. The rpsL targeting spacer was cloned by annealing of oligos B352+B353 and cloning in the BsaI cut pCRISPR giving pCRISPR:rpsL.

[0287] Generation of targeting and editing constructs. Targeting constructs used for genome editing were made by Gibson assembly of Left PCRs and Right PCRs (Table G). Editing constructs were made by SOEing PCR fusing PCR products A (PCR A), PCR products B (PCR B) and PCR products C (PCR C) when applicable (Table G). The CRISPR:Ø and CRISPR:ermAM(stop) targeting constructs were generated by PCR amplification of JEN62 and crR6 genomic DNA respectively, with oligos L409 and L481.

[0288] Generation of targets with randomized PAM or protospacer sequences. The 5 nucleotides following the spacer 1 target were randomized through amplification of R6^{8232.5} genomic DNA with primers W377/L426. This PCR product was then assembled with the cat gene and the srtA upstream region that were amplified from the same template with primers L422/W376. 80 ng of the assembled DNA was used to transform strains R6 and crR6. Samples for the randomized targets were prepared using the following primers: B280-B290/L426 to randomize bases 1-10 of the target and B269-B278/L426 to randomize bases 10-20. Primers L422/B268 and L422/B279 were used to amplify the cat gene and srtA upstream region to be assembled with the first and last 10 PCR products respectively. The assembled constructs were pooled together and 30 ng was transformed in R6 and crR6. After transformation, cells were plated on chloramphenicol selection. For each sample more than 2x10⁵ cells were pooled together in 1 ml of THYE and genomic DNA was extracted with the Promega Wizard kit. Primers B250/B251 were used to amplify the target region. PCR products were tagged and run on one Illumina MiSeq paired-end lane using 300 cycles.

[0289] Analysis of Deep Sequencing Data

[0290] Randomized PAM: For the randomized PAM experiment 3,429,406 reads were obtained for crR6 and 3,253,998 for R6. It is expected that only half of them will correspond to the PAM-target while the other half will sequence the other end of the PCR product. 1,623,008 of the crR6 reads and 1,537,131 of the R6 reads carried an error-free target sequence. The occurrence of each possible PAM among these reads is shown in supplementary file. To estimate the functionality of a PAM, its relative proportion in the crR6 sample over the R6 sample was computed and is denoted r_{ijklm} where l,j,k,l,m are one of the 4 possible bases. The following statistical model was constructed:

$$\log(r_{ijklm}) = \mu + b_2i + b_3j + b_4k + b_2b_3i_j + b_3b_4j_k + \epsilon_{ijklm},$$

[0291] where ϵ is the residual error, b_2 is the effect of the 2nd base of the PAM, b_3 of the third, b_4 of the fourth, b_2b_3 is the interaction between the second and third bases, b_3b_4 between the third and fourth bases. An analysis of variance was performed:

Anova table					
	Df	Sum Sq	Mean Sq	F value	Pr (>F)
b3	3	151.693	50.564	601.8450	<2.2e-16***
b2	3	90.521	30.174	359.1454	<2.2e-16***
b4	3	1.881	0.627	7.4623	6.070e-05***
b3:b2	9	228.940	25.438	302.7738	<2.2e-16***
b3:b4	9	3.010	0.334	3.9809	5.227e-05***
Residuals	996	83.680	0.084		

[0292] When added to this model, b_1 or b_5 do not appear to be significant and other interactions than the ones included can also be discarded. The model choice was made through successive comparisons of more or less complete models using the anova method in R. Tukey's honest significance test was used to determine if pairwise differences between effects are significant.

[0293] NGGNN patterns are significantly different from all other patterns and carry the strongest effect (see table below).

[0294] In order to show that positions 1, 4 or 5 do not affect the NGGNN pattern Applicants looked at these sequences only. Their effect appears to be normally distributed (see QQ plot in FIG. 71), and model comparisons using the anova method in R shows that the null model is the best one, i.e. there is no significant role of b_1 , b_4 and b_5 .

[0295] Model Comparison using the Anova Method in R for the NGGNN Sequences

Model 1: ratio.log~1 Model 2: ratio.log~b1 + b4 + b5						
	Res. Df	RSS	Df	Sum of Sq	F	Pr (>F)
1	63	14.579				
2	54	11.295	9	3.2836	1.7443	0.1013

[0296] Partial Interference of NAGNN and NNGGN Patterns

[0297] NAGNN patterns are significantly different from all other patterns but carry a much smaller effect than NNGGN (see Tukey's honest significance test below).

[0298] Finally, NTGGN and NCGGN patterns are similar and show significantly more CRISPR interference than NTGHN and NCGHN patterns (where H is A,T or C), as shown by a bonferroni adjusted pairwise student-test.

[0299] Pairwise Comparisons of the Effect of b_4 on NYGNN Sequences Using t Tests with Pooled SD

Data: b4			
	A	C	G
C	1.00	—	—
G	9.2e-05	2.4e-06	—
T	0.31	1.00	1.2e-08

[0300] Taken together, these results allow concluding that NNGGN patterns in general produce either a complete interference in the case of NGGGN, or a partial interference in the case of NAGGN, NTGGN or NCGGN.

[0301] Tukey multiple comparisons of means: 95% family-wise confidence level

	diff	lwr	upr	p adj
	Sb2:b3			
G:G-A:A	-2.76475	-2.94075	-2.58875	<1E-07
G:G-C:A	-2.79911	-2.97511	-2.62311	<1E-07
G:G-T:A	-2.7809	-2.9569	-2.6049	<1E-07
G:G-A:C	-2.81643	-2.99244	-2.64043	<1E-07
G:G-C:C	-2.77903	-2.95504	-2.60303	<1E-07
G:G-G:C	-2.64867	-2.82468	-2.47267	<1E-07
G:G-T:C	-2.79718	-2.97319	-2.62118	<1E-07
G:G-A:G	-2.67068	-2.84668	-2.49468	<1E-07
G:G-C:G	-2.73525	-2.91125	-2.55925	<1E-07
G:G-T:G	-2.7976	-2.62159	-2.9736	<1E-07
G:G-A:T	-2.76727	-2.59127	-2.94328	<1E-07
G:G-C:T	-2.84114	-2.66513	-3.01714	<1E-07
G:G-G:T	-2.76409	-2.58809	-2.94009	<1E-07
G:G-T:T	-2.76781	-2.59181	-2.94381	<1E-07
G:G-G:A	-2.13964	-2.31565	-1.96364	<1E-07
G:A-A:A	-0.62511	-0.80111	-0.4491	<1E-07
G:A-C:A	-0.65947	-0.83547	-0.48346	<1E-07
G:A-T:A	-0.64126	-0.46525	-0.81726	<1E-07
G:A-A:C	-0.67679	-0.50078	-0.85279	<1E-07
G:A-C:C	-0.63939	-0.46339	-0.81539	<1E-07
G:A-G:C	-0.50903	-0.33303	-0.68503	<1E-07
G:A-T:C	-0.65754	-0.48154	-0.83354	<1E-07
G:A-A:G	-0.53104	-0.35503	-0.70704	<1E-07
G:A-C:G	-0.59561	-0.4196	-0.77161	<1E-07

-continued

	diff	lwr	upr	p adj
G:A-T:G	-0.65795	-0.48195	-0.83396	<1E-07
G:A-A:T	-0.62763	-0.45163	-0.80363	<1E-07
G:A-C:T	-0.70149	-0.52549	-0.8775	<1E-07
G:A-G:T	-0.62445	-0.44844	-0.80045	<1E-07
G:A-T:T	-0.62817	-0.45216	-0.80417	<1E-07
		Sb3:b4		
G:G-G:A	-0.33532	-0.51133	-0.15932	<1E-07
G:G-G:C	-0.18118	-0.35719	-0.00518	0.036087
G:G-G:T	-0.31626	-0.14026	-0.49226	<1E-07

[0302] Randomized Target

[0303] For the randomized target experiment **540,726** reads were obtained for crR6 and **753,570** for R6. As before, only half of the reads are expected to sequence the interesting end of the PCR product. After filtering for reads that carry a target that is error-free or with a single point mutation, **217,656** and **353,141** reads remained for crR6 and R6 respectively. The relative proportion of each mutant in the crR6 sample over the R6 sample was computed (FIG. 24c). All mutations outside of the seed sequence (13-20 bases away from the PAM) show full interference. Those sequences were used as a reference to determine if other mutations inside the seed sequence can be said to significantly disrupt interference. A normal distribution was fitted to these sequences using the fitdistr function of the MASS R package. The 0.99 quantile of the fitted distribution is shown as a dotted line in FIG. 24c. FIG. 72 shows a histogram of the data density with fitted normal distribution (black line) and 0.99 quantile (dotted line).

TABLE F

Relative abundance of PAM sequences in the crR6/R6 samples averaged over bases 1 and 5.										
3rd position										
		A	C	G	T					
2nd position	A	AAA	1.04	ACA	1.12	AGA	0.73	ATA	1.10	A 4th position
		AAC	1.07	ACC	1.04	AGC	0.54	ATC	0.97	C
		AAG	1.00	ACG	1.09	AGG	0.61	ATG	1.07	G
		AAT	0.98	ACT	1.02	AGT	0.65	ATT	1.01	T
	C	CAA	1.05	CCA	1.05	CGA	0.99	CTA	1.07	A
		CAC	1.04	CCC	1.02	CGC	1.08	CTC	1.04	C
		CAG	1.08	CCG	1.08	CGG	0.81	CTG	1.05	G
		CAT	1.13	CCT	1.05	CGT	1.07	CTT	1.08	T
	G	GAA	0.97	GCA	1.05	GGA	0.08	GTA	0.99	A
		GAC	0.92	GCC	1.00	GGC	0.05	GTC	1.15	C
		GAG	0.96	GCG	0.98	GGG	0.07	GTG	0.98	G
		GAT	0.98	GCT	0.99	GGT	0.06	GTT	1.05	T
	T	TAA	1.08	TCA	1.16	TGA	1.05	TTA	1.14	A
		TAC	1.00	TCC	1.08	TGC	1.08	TTC	1.05	C
		TAG	1.02	TCG	1.11	TGG	0.77	TTG	1.01	G
		TAT	1.01	TCT	1.12	TGT	1.21	TTT	1.02	T

TABLE G

Primers used in this study (SEQ ID NOS 68-183, respectively, in order of appearance).	
Primer	Sequence 5'-3'
B217	TCCTAGCAGGATTTCTGATATTACTGTCACGTTTTAGAGCTATGCTGTTTTGA
B218	GTGACAGTAATATCAGAAATCCTGCTAGGAGTTTTGGGACCATTCAACAGC

TABLE G-continued

Primers used in this study (SEQ ID NOS 68-183, respectively, in order of appearance).	
Primer	Sequence 5'-3'
B229	GGGTTTCAAGTCTTTGTAGCAAGAG
B230	GCCAATGAACGGGAACCCCTTGGTC
B250	NNNNGACGAGGCAATGGCTGAAATC
B251	NNNNTTATTTGGCTCATATTTGCTG
B255	CTTTACACCAATCGCTGCAACAGAC
B256	CAAAATTTCTAGTCTTCTTTGCCTTCCCCATAAACCCCTCCTTA
B257	AGGGTTTTATGGGAAAGGCAAGAAGACTAGAAATTTGATACC
B258	CTTACGGTGCATAAAGTCAATTTCC
B269	TGGCTCGATTTTCAGCCATTGC
B270	CTTTGACGAGGCAATGGCTGAAATCGAGCCAANAAGCGCAAG
B271	CTTTGACGAGGCAATGGCTGAAATCGAGCCAANAAGCGCAAG
B272	CTTTGACGAGGCAATGGCTGAAATCGAGCCAANAAGCGCAAG
B273	CTTTGACGAGGCAATGGCTGAAATCGAGCCAANAAGCGCAAG
B274	CTTTGACGAGGCAATGGCTGAAATCGAGCCAANAAGCGCAAG
B275	CTTTGACGAGGCAATGGCTGAAATCGAGCCAANAAGCGCAAG
B276	CTTTGACGAGGCAATGGCTGAAATCGAGCCAANAAGCGCAAG
B277	CTTTGACGAGGCAATGGCTGAAATCGAGCCAANAAGCGCAAG
B278	CTTTGACGAGGCAATGGCTGAAATCGAGCCAANAAGCGCAAG
B279	GCGCTTTTTTGGCTCGATTTTCAG
B280	CAATGGCTGAAATCGAGCCAANAAGCGCAANAAGAAATC
B281	CAATGGCTGAAATCGAGCCAANAAGCGCAANAAGAAATC
B282	CAATGGCTGAAATCGAGCCAANAAGCGCAANAAGAAATC
B283	CAATGGCTGAAATCGAGCCAANAAGCGCAANAAGAAATC
B284	CAATGGCTGAAATCGAGCCAANAAGCGCAANAAGAAATC
B285	CAATGGCTGAAATCGAGCCAANAAGCGCAANAAGAAATC
B286	CAATGGCTGAAATCGAGCCAANAAGCGCAANAAGAAATC
B287	CAATGGCTGAAATCGAGCCAANAAGCGCAANAAGAAATC
B288	CAATGGCTGAAATCGAGCCAANAAGCGCAANAAGAAATC
B289	CAATGGCTGAAATCGAGCCAANAAGCGCAANAAGAAATC
B290	CAATGGCTGAAATCGAGCCAANAAGCGCAANAAGAAATC
B296	gatacTCCATCCGTACAACCACAACCTGg
B297	aattcCAGGGTTGTGGTTGTACGGATGGAg
B298	CATGGATCCTATTTCTTAATAACTAAAAATATGG
B299	CATGAATTCAACTCAACAAGTCTCAGTGTGCTG
B300	AAACATTTTTTCTCCATTTAGGAAAAGGATGCTG
B301	AAAACAGCATCCTTTTTCTAAATGGAGAAAAAT

TABLE G-continued

Primers used in this study (SEQ ID NOS 68-183, respectively, in order of appearance).	
Primer	Sequence 5'-3'
B302	AAACCTTAAATCAGTCACAAATAGCAGCAAATG
B303	AAAACAATTTTGCTGCTATTTGTGACTGATTTAAG
B304	AAACTTTTCATCATACGACCAATCTGCTTTATTTG
B305	AAAACAATAAAGCAGATTGGTCGTATGATGAAAA
B306	AAACTCGTCCAGAAGTTATCGTAAAAGAAATCGAG
B307	AAAACCTGATTTCTTTTACGATAACTTCTGGACGA
B308	AAACAATCTCTCCAAGGTTTCTTAAAAATCTCTG
B309	AAAACAGAGATTTTAAAGGAAACCTTGGAGAGATT
B310	AAACGCCATCGTCAGGAAGAAGCTATGCTTGAGTG
B311	AAAACACTCAAGCATAGCTTCTCTCTGACGATGGC
B312	AAACATCTCTATACTTATTGAAATTTCTTTGTATG
B313	AAAACATACAAGAAATTTCAATAAGTATAGAGAT
B314	AAACTAGCTGTGATAGTCCGCAAACCAGCCTTCG
B315	AAAACGAAGGCTGGTTTTGCGGACTATCACAGCTA
B316	AAACATCGGAAGTTCGAGCAAGTAATTATCTTTTG
B317	AAAACAAAAGATAATTACTTGTCTCGACCTCCGAT
B318	AAACAAGATGGTATCGCAAATAAGTGACAATAAG
B319	AAAACCTATTGTCACTTACTTTGCGATACCATCTT
B320	GAGACCTTTGAGCTTCCGAGACTGGTCTCAGTTTTGGACCATTCAAACAG
B321	TGAGACCAGTCTCGGAAGCTCAAAGTCTCGTTTTAGAGCTATGCTGTTTTG
B352	aaacTACTTTACGCAGCGCGGAGTTCGGTTTTTTg
B353	aaaacAAAAACCGAACTCCGCGCTGCGTAAAGTA
HC008_SP	ATGCCGGTACTGCCGGGCCTCTTCCGGGATTACGAAATCATCCTG
HC009_SP	GTGACTGGCGATGCTGTCCGAATGGACGATCACACTACTCTTCTT
HC010_SP	TTAAGAAATAATCTTCATCTAAAATATACTTCAGTCACCTCCTAGCTGAC
HC011_SP	ATTGATTTGAGTCAGCTAGGAGGTGACTGAAGTATATTTTAGATGAAG
HC014_SP	GAGACCTTTGAGCTTCCGAGACTGGTCTCAGTTTTGGACCATTCAAACAGCATAGCTCTAAAACCTCGTAGACTA TTTTTGTCT
HC015_SP	GAGACCAGTCTCGGAAGCTCAAAGTCTCGTTTTAGAGCTATGCTGTTTTGAATGGTCCCAAACTTCCAGCACCTG AGACTTG
L403	AGTCATCCCAGCAACAAATGG
L409	CGTGGTAAATCGGATAACGTTCCAAGTGAAG
L422	Tgctcttcttcacaaacaaggg
L426	AAGCCAAAGTTTGGCACCACC
L430	GTAGCTTATTTCAGTCTTAGTGG
L444	CGTTTGTGAACTAATGGGTGCAAATACGAATCTTCTCCTGACG
L445	CGTCAGGAGAAGATTCGTAATTTGCACCCATTAGTTCACAAACG

TABLE G-continued

Primers used in this study (SEQ ID NOS 68-183, respectively, in order of appearance).	
Primer	Sequence 5'-3'
L446	GATATTATGGAGCCTATTTTGTGGGTTTTTAGGCATAAACTATATG
L447	CATATAGTTTTATGCCTAAAAACCcACAAAATAGGCTCCATAATATC
L448	ATTATTTCTTAATAACTAAAAATATGG
L457	CGTgtacaattgctagecgtacggc
L458	GCACCGGTGATCACTAGTCCTAGG
L459	cctaggactagtgatcaccggtGCAAATATGAGCCAATAAATATAT
L461	GCCGTACGCTAGCAATTGTACACGTTTGTGAACTAATGGGTGC
L481	TTCAAATTTTCCATTTGATTCTCC
L488	CCATATTTTTAGTTATTAAGAAATAATACCAGCCATCAGTCACCTCC
W256	AGACGATTCAATAGACAATAAGG
W286	GTTTTGGGACCATTCAAACAGCATAGCTCTAAAACCTCGTAGAC
W287	GCTATGCTGTTTTGAATGGTCCCAAACcattatthtaacacacagaggtg
W288	GCTATGCTGTTTTGAATGGTCCCAAACGCACCCATTAGTTCAACAAACG
W326	AATTCTTTTCTTCATCATCGGTC
W327	AAGAAAGAATGAAGATTGTTTCATG
W341	GGTACTAATCAAATAGTGAGGAGG
W354	GTTTTTCAAATCTGCGTTGCG
W355	AAAAATTGAAAAATGGTGGAAACAC
W356	ATTCGTAACCGGTATCGGTTTCTTTTAAAGTTTTGGGACCATTCAAACAGC
W357	TTTAAAGAAACCGATACCGTTTACGAAATGTTTTAGAGCTATGCTGTTTTGA
W365	AAACGGTATCGGTTTCTTTTAAATCAATTGTTTTGGGACCATTCAAACAGC
W366	AATTGAATTTAAAGAAACCGATACCGTTTGTGTTTTAGAGCTATGCTGTTTTGA
W370	GTTCTTAAACCAAACGGTATCGGTTTCTTTTAAATTC
W371	GAAACCGATACCGTTTTGGTTTAAAGGAACAGGTAAGGGCATTTAAC
W376	CGATTTCAGCCATTGCCTCGTC
W377	GCCTTTGACGAGGCAATGGCTGAAATCGNNNNAAAAAGCGCAAGAAGAAATCAAC
W391	TCCGTACAACCCACAACCTGCTAGTGAGCGTTTTGGGACCATTCAAACAGC
W392	GCTCACTAGCAGGGTTGTGGTTGTACGGAGTTTTAGAGCTATGCTGTTTTGA
W393	TTGTTGCCACTCTTCTTCTTTC
W397	CAGGGTTGTGGTTGTTGCGATGGAGTTAACTCCCATCTCC
W398	GGGAGTTAACTCCATCGCAACAACCCACAACCTGCTAGTG
W403	GTGGTATCTATCGTGATGTGAC
W404	TTACCGAAACGGAATTTATCTGC
W405	AAAGCTAGAGTTCCGCAATTGG

TABLE G-continued

Primers used in this study (SEQ ID NOS 68-183, respectively, in order of appearance).	
Primer	Sequence 5'-3'
W431	GTGGGTTGTACGGATTGAGTTAACTCCCATCTCCTTC
W432	GATGGGAGTTAACTCAATCCGTACAACCCACAACCCCTG
W433	GCTTCACCTATTGCAGCACCATTGACCACATGAAGATAG
W434	GTGGTCAATTGGTCTGCAATAGGTGAAGCTAATGGTGATG
W463	CTGATTTGTATTAATTTTGAGACATTATGCTTCACCTTC
W464	GCATAATGTCTCAAATAATACAATCAGTGAAATCATG
W465	GTTTTGGGACCATTCAAACAGCATAGCTCTAAAACGTGACAGTAATATCAG
W466	GTTTTAGAGCTATGCTGTTTTGAATGGTCCAAAACGCTCACTAGCAGGGTTG
W542	ATACTTTACGCAGCGCGGAGTTCGGTTTTgTAGGAGTGGTAGTATATACACGAGTACAT

TABLE H

Design of targeting and editing constructs used in this study (SEQ ID NOS 184, 184, 184, 185, and 186, respectively, in order of appearance).								
Targeting Constructs								
Edition	Template DNA	Left PCR	Right PCR	Spacer sequence	PAM			
bgaA R > A	crR6Rk	W256/W391	W392/L403	GCTCACTAGCAGGGTTGTGGTTGTACGGA	TGG			
bgaA NE > AA	crR6Rk	W256/W391	W392/L403	GCTCACTAGCAGGGTTGTGGTTGTACGGA	TGG			
AbgaA	crR6Rk	W256/W391	W392/L403	GCTCACTAGCAGGGTTGTGGTTGTACGGA	TGG			
AsrtA	crR6Rc	W256/B218	B217/L403	TCCTAGCAGGATTTCTGATATTACTGTAC	TGG			
ermB Stop	crR6Rk	W256/W356	W357/L403	TTTAAAAGAAACCGATACCGTTTACGAAAT	TGG			
AsrtA AbgaA	JEN51 (for Left PCR) and JEN52 (for Right PCR)	W256/W465	W466/W403	same as the ones used for AsrtA and AbgaA				
Editing Constructs								
Edition	Template DNA	PCR A	PCR B	PCR C	SOEing PCR	Name of resulting strains	Primers used to verify edited genotype	
bgaA R > A	R6	W403/W397	W398/W404	N/A	W403/W404	JEN56	W403/W404	
bgaA NE > AA	R6	W403/W431	W432/W433	W434/W404	W403/W404	JEN60	W403/W404	
AbgaA	R6	B255/B256	B257/B258	N/A	B255/B258	JEN52	W393/W405	
AsrtA	R6	B230/W463	W464/B229	N/A	B230/B229	JEN51	W422/W426	
ermB Stop	JEN38	L422/W370	W371/L426	N/A	L422/L426	JEN43	L457/L458	
AsrtA AbgaA	same as the ones used for AsrtA and AbgaA					JEN64	same as the ones used for AsrtA and AbgaA	

Example 6

Optimization of the Guide RNA for *Streptococcus pyogenes* Cas9 (Referred to as SpCas9).

[0304] Applicants mutated the tracrRNA and direct repeat sequences, or mutated the chimeric guide RNA to enhance the RNAs in cells.

[0305] The optimization is based on the observation that there were stretches of thymines (Ts) in the tracrRNA and guide RNA, which might lead to early transcription termination by the pol 3 promoter. Therefore Applicants generated the following optimized sequences. Optimized tracrRNA and corresponding optimized direct repeat are presented in pairs.

[0306] Optimized tracrRNA 1 (mutation underlined):

(SEQ ID NO: 187)
GGAACCATTCAACAGCATAGCAAGTTAAATAAGGCTAGTCCGTTATC
AACTTGAAAAAGTGGCACCAGTTCGGTGCTTTTT

[0307] Optimized direct repeat 1 (mutation underlined):

(SEQ ID NO: 188)
GTTaTAGAGCTATGCTGTTaTGAATGGTCCCAAAC

[0308] Optimized tracrRNA 2 (mutation underlined):

(SEQ ID NO: 189)
GGAACCATTCAACAGCATAGCAAGTTAAATAAGGCTAGTCCGTTATC
AACTTGAAAAAGTGGCACCAGTTCGGTGCTTTTT

[0309] Optimized direct repeat 2 (mutation underlined):

(SEQ ID NO: 190)
GTaTTAGAGCTATGCTGTaTTGAATGGTCCCAAAC

[0310] Applicants also optimized the chimeric guideRNA for optimal activity in eukaryotic cells.

[0311] Original guide RNA:

(SEQ ID NO: 191)
NNNNNNNNNNNNNNNNNNNNNGTTTTAGAGCTAGAAATAGCAAGTTAAAT
AAGGCTAGTCCGTTATCAACTTGAAAAAGTGGCACCAGTTCGGTGCTTTTT
TTT

[0312] Optimized chimeric guide RNA sequence 1:

(SEQ ID NO: 192)
NNNNNNNNNNNNNNNNNNNNNGTATTAGAGCTAGAAATAGCAAGTTAAAT
AAGGCTAGTCCGTTATCAACTTGAAAAAGTGGCACCAGTTCGGTGCTTTTT
TTT

[0313] Optimized chimeric guide RNA sequence 2:

(SEQ ID NO: 193)
NNNNNNNNNNNNNNNNNNNNNGTTTTAGAGCTATGCTGTTTTGGAACAAA
ACAGCATAGCAAGTTAAATAAGGCTAGTCCGTTATCAACTTGAAAAAGT
GGCACCAGTTCGGTGCTTTTTTTT

[0314] Optimized chimeric guide RNA sequence 3:

(SEQ ID NO: 194)
NNNNNNNNNNNNNNNNNNNNNGTATTAGAGCTATGCTGTATTGGAACAAT
ACAGCATAGCAAGTTAAATAAGGCTAGTCCGTTATCAACTTGAAAAAGT
GGCACCAGTTCGGTGCTTTTTTTT

[0315] Applicants showed that optimized chimeric guide RNA works better as indicated in FIG. 3. The experiment was conducted by co-transfecting 293FT cells with Cas9 and a U6-guide RNA DNA cassette to express one of the four RNA forms shown above. The target of the guide RNA is the

same target site in the human Emx1 locus: "GTCACCTC-CAATGACTAGGG (SEQ ID NO: 195)"

Example 7

Optimization of *Streptococcus Thermophiles* LMD-9 CRISPR1 Cas9 (Referred to as St1Cas9).

[0316] Applicants designed guide chimeric RNAs as shown in FIG. 4.

[0317] The St1Cas9 guide RNAs can undergo the same type of optimization as for SpCas9 guide RNAs, by breaking the stretches of poly thymines (Ts)

Example 8

Cas9 Diversity and Mutations

[0318] The CRISPR-Cas system is an adaptive immune mechanism against invading exogenous DNA employed by diverse species across bacteria and archaea. The type II CRISPR-Cas9 system consists of a set of genes encoding proteins responsible for the "acquisition" of foreign DNA into the CRISPR locus, as well as a set of genes encoding the "execution" of the DNA cleavage mechanism; these include the DNA nuclease (Cas9), a non-coding transactivating cr-RNA (tracrRNA), and an array of foreign DNA-derived spacers flanked by direct repeats (crRNAs). Upon maturation by Cas9, the tracrRNA and crRNA duplex guide the Cas9 nuclease to a target DNA sequence specified by the spacer guide sequences, and mediates double-stranded breaks in the DNA near a short sequence motif in the target DNA that is required for cleavage and specific to each CRISPR-Cas system. The type II CRISPR-Cas systems are found throughout the bacterial kingdom and highly diverse in in Cas9 protein sequence and size, tracrRNA and crRNA direct repeat sequence, genome organization of these elements, and the motif requirement for target cleavage. One species may have multiple distinct CRISPR-Cas systems.

[0319] Applicants evaluated 207 putative Cas9s from bacterial species identified based on sequence homology to known Cas9s and structures orthologous to known subdomains, including the HNH endonuclease domain and the RuvC endonuclease domains [information from the Eugene Koonin and Kira Makarova]. Phylogenetic analysis based on the protein sequence conservation of this set revealed five families of Cas9s, including three groups of large Cas9s (~1400 amino acids) and two of small Cas9s (~1100 amino acids) (FIGS. 39 and 40A-F).

[0320] In this example, Applicants show that the following mutations can convert SpCas9 into a nicking enzyme: D10A, E762A, H840A, N854A, N863A, D986A.

[0321] Applicants provide sequences showing where the mutation points are located within the SpCas9 gene (FIG. 41). Applicants also show that the nickases are still able to mediate homologous recombination (Assay indicated in FIG. 2). Furthermore, Applicants show that SpCas9 with these mutations (individually) do not induce double strand break (FIG. 47).

Example 9

Supplement to DNA Targeting Specificity of the RNA-Guided Cas9 Nuclease

[0322] Cell culture and Transfection

[0323] Human embryonic kidney (HEK) cell line 293FT (Life Technologies) was maintained in Dulbecco's modified Eagle's Medium (DMEM) supplemented with 10% fetal

bovine serum (HyClone), 2 mM GlutaMAX (Life Technologies), 100 U/mL penicillin, and 100 mg/mL streptomycin at 37° C. with 5% CO₂ incubation.

[0324] 293FT cells were seeded either onto 6-well plates, 24-well plates, or 96-well plates (Corning) 24 hours prior to transfection. Cells were transfected using Lipofectamine 2000 (Life Technologies) at 80-90% confluence following the manufacturer's recommended protocol. For each well of a 6-well plate, a total of 1 µg of Cas9+sgRNA plasmid was used. For each well of a 24-well plate, a total of 500 ng Cas9+sgRNA plasmid was used unless otherwise indicated. For each well of a 96-well plate, 65 ng of Cas9 plasmid was used at a 1:1 molar ratio to the U6-sgRNA PCR product.

[0325] Human embryonic stem cell line HUES9 (Harvard Stem Cell Institute core) was maintained in feeder-free conditions on GelTrex (Life Technologies) in mTesR medium (Stemcell Technologies) supplemented with 100 µg/ml Normocin (InvivoGen). HUES9 cells were transfected with Amaxa P3 Primary Cell 4-D Nucleofector Kit (Lonza) following the manufacturer's protocol.

[0326] SURVEYOR Nuclease Assay for Genome Modification

[0327] 293FT cells were transfected with plasmid DNA as described above. Cells were incubated at 37° C. for 72 hours post-transfection prior to genomic DNA extraction. Genomic DNA was extracted using the QuickExtract DNA Extraction Solution (Epicentre) following the manufacturer's protocol. Briefly, pelleted cells were resuspended in QuickExtract solution and incubated at 65° C. for 15 minutes and 98° C. for 10 minutes.

[0328] The genomic region flanking the CRISPR target site for each gene was PCR amplified (primers listed in Tables J and K), and products were purified using QiaQuick Spin Column (Qiagen) following the manufacturer's protocol. 400ng total of the purified PCR products were mixed with 2 µl 10× Taq DNA Polymerase PCR buffer (Enzymatics) and ultrapure water to a final volume of 20 µl, and subjected to a re-annealing process to enable heteroduplex formation: 95° C. for 10 min, 95° C. to 85° C. ramping at -2° C/s, 85° C. to 25° C. at -0.25° C/s, and 25° C. hold for 1 minute. After re-annealing, products were treated with SURVEYOR nuclease and SURVEYOR enhancer S (Transgenomics) following the manufacturer's recommended protocol, and analyzed on 4-20% Novex TBE poly-acrylamide gels (Life Technologies). Gels were stained with SYBR Gold DNA stain (Life Technologies) for 30 minutes and imaged with a Gel Doc gel imaging system (Bio-rad). Quantification was based on relative band intensities.

[0329] Northern Blot Analysis of tracrRNA Expression in Human Cells

[0330] Northern blots were performed as previously described. Briefly, RNAs were heated to 95° C. for 5 min before loading on 8% denaturing polyacrylamide gels (SequaGel, National Diagnostics). Afterwards, RNA was transferred to a pre-hybridized Hybond N+ membrane (GE Healthcare) and crosslinked with Stratagene UV Crosslinker (Stratagene). Probes were labeled with [γ -32P] ATP (Perkin Elmer) with T4 polynucleotide kinase (New England Biolabs). After washing, membrane was exposed to phosphor screen for one hour and scanned with phosphorimager (Typhoon).

[0331] Bisulfite Sequencing to Assess DNA Methylation Status

[0332] HEK 293FT cells were transfected with Cas9 as described above. Genomic DNA was isolated with the DNeasy Blood & Tissue Kit (Qiagen) and bisulfite converted with EZ DNA Methylation-Lightning Kit (Zymo

Research). Bisulfite PCR was conducted using KAPA2G Robust HotStart DNA Polymerase (KAPA Biosystems) with primers designed using the Bisulfite Primer Seeker (Zymo Research, Tables J and K). Resulting PCR amplicons were gel-purified, digested with EcoRI and HindIII, and ligated into a pUC19 backbone prior to transformation. Individual clones were then Sanger sequenced to assess DNA methylation status.

[0333] In Vitro Transcription and Cleavage Assay

[0334] HEK 293FT cells were transfected with Cas9 as described above. Whole cell lysates were then prepared with a lysis buffer (20 mM HEPES, 100 mM KCl, 5 mM MgCl₂, 1 mM DTT, 5% glycerol, 0.1% Triton X-100) supplemented with Protease Inhibitor Cocktail (Roche). T7-driven sgRNA was in vitro transcribed using custom oligos (Example 10) and HiScribe T7 In Vitro Transcription Kit (NEB), following the manufacturer's recommended protocol. To prepare methylated target sites, pUC19 plasmid was methylated by M.SssI and then linearized by NheI. The in vitro cleavage assay was performed as follows: for a 20 µL cleavage reaction, 10 µL of cell lysate with incubated with 2 µL cleavage buffer (100 mM HEPES, 500 mM KCl, 25 mM MgCl₂, 5 mM DTT, 25% glycerol), the in vitro transcribed RNA, and 300 ng pUC19 plasmid DNA.

[0335] Deep Sequencing to Assess Targeting Specificity

[0336] HEK 293FT cells plated in 96-well plates were transfected with Cas9 plasmid DNA and single guide RNA (sgRNA) PCR cassette 72 hours prior to genomic DNA extraction (FIG. 72). The genomic region flanking the CRISPR target site for each gene was amplified (FIG. 74, FIG. 80, (Example 10) by a fusion PCR method to attach the Illumina P5 adapters as well as unique sample-specific barcodes to the target amplicons (schematic described in FIG. 73). PCR products were purified using EconoSpin 96-well Filter Plates (Epoch Life Sciences) following the manufacturer's recommended protocol.

[0337] Barcoded and purified DNA samples were quantified by Quant-iT PicoGreen dsDNA Assay Kit or Qubit 2.0 Fluorometer (Life Technologies) and pooled in an equimolar ratio. Sequencing libraries were then deep sequenced with the Illumina MiSeq Personal Sequencer (Life Technologies).

[0338] Sequencing Data Analysis and Indel Detection

[0339] MiSeq reads were filtered by requiring an average Phred quality (Q score) of at least 23, as well as perfect sequence matches to barcodes and amplicon forward primers. Reads from on- and off-target loci were analyzed by first performing Smith-Waterman alignments against amplicon sequences that included 50 nucleotides upstream and downstream of the target site (a total of 120 bp). Alignments, meanwhile, were analyzed for indels from 5 nucleotides upstream to 5 nucleotides downstream of the target site (a total of 30 bp). Analyzed target regions were discarded if part of their alignment fell outside the MiSeq read itself, or if matched base-pairs comprised less than 85% of their total length.

[0340] Negative controls for each sample provided a gauge for the inclusion or exclusion of indels as putative cutting events. For each sample, an indel was counted only if its quality score exceeded $\mu - \sigma$, where μ was the mean quality-score of the negative control corresponding to that sample and σ was the standard deviation of same. This yielded whole target-region indel rates for both negative controls and their corresponding samples. Using the negative control's per-target-region-per-read error rate, q , the sample's observed indel count n , and its read-count R , a maximum-likelihood estimate for the fraction of reads hav-

ing target-regions with true-indels, p , was derived by applying a binomial error model, as follows.

[0341] Letting the (unknown) number of reads in a sample having target regions incorrectly counted as having at least 1 indel be E , we can write (without making any assumptions about the number of true indels)

$$Prob(E | p) = \binom{R(1-p)}{E} q^E (1-q)^{R(1-p)-E}$$

[0342] since $R(1-p)$ is the number of reads having target-regions with no true indels. Meanwhile, because the number of reads observed to have indels is n , $n-E+Rp$, in other words the number of reads having target-regions with errors but no true indels plus the number of reads whose target-regions correctly have indels. We can then re-write the above

$$Prob(E | p) = Prob(n = E + Rp | p) = \binom{R(1-p)}{n-Rp} q^{n-Rp} (1-q)^{R-n}$$

[0343] Taking all values of the frequency of target-regions with true-indels p to be equally probable a priori, $Prob(n|p) \propto Prob(p|n)$. The maximum-likelihood estimate (MLE) for the frequency of target regions with true-indels was therefore set as the value of P that maximized $Prob(n|p)$. This was evaluated numerically.

[0344] In order to place error bounds on the true-indel read frequencies in the sequencing libraries themselves, Wilson score intervals (2) were calculated for each sample, given the MLE-estimate for true-indel target-regions, R_p , and the number of reads R . Explicitly, the lower bound l and upper bound u were calculated as

$$l = \left(Rp + \frac{z^2}{2} - z\sqrt{Rp(1-p) + \frac{z^2}{4}} \right) / (R + z^2)$$

$$u = \left(Rp + \frac{z^2}{2} + z\sqrt{Rp(1-p) + \frac{z^2}{4}} \right) / (R + z^2)$$

[0345] where z , the standard score for the confidence required in normal distribution of variance 1, was set to 1.96, meaning a confidence of 95%. The maximum upper bounds and minimum lower bounds for each biological replicate are listed in FIGS. 80-83.

[0346] qRT-PCR Analysis of Relative Cas9 and sgRNA Expression

[0347] 293FT cells plated in 24-well plates were transfected as described above. 72 hours post-transfection, total RNA was harvested with miRNeasy Micro Kit (Qiagen). Reverse-strand synthesis for sgRNAs was performed with qScript Flex cDNA kit (VWR) and custom first-strand synthesis primers (Tables J and K). qPCR analysis was performed with Fast SYBR Green Master Mix (Life Technologies) and custom primers (Tables J and K), using GAPDH as an endogenous control. Relative quantification was calculated by the AACT method.

TABLE I

Target site genomic				
ID	target	Target site sequence (5' to 3')	PAM	strand
1	EMX1	GTCACCTCCAATGACTAGGG (SEQ ID NO: 319)	TGG	+
2	EMX1	GACATCGATGTCCTCCCAT (SEQ ID NO: 196)	TGG	-
3	EMX1	GAGTCCGAGCAGAAGAAGAA (SEQ ID NO: 197)	GGG	+
6	EMX1	GCGCCACCGTTGATGTGAT (SEQ ID NO: 198)	GGG	-
10	EMX1	GGGGCACAGATGAGAACTC (SEQ ID NO: 199)	AGG	-
11	EMX1	GTACAAACGGCAGAAGCTGG (SEQ ID NO: 200)	AGG	+
12	EMX1	GGCAGAAGCTGGAGGAGGAA (SEQ ID NO: 201)	GGG	+
13	EMX1	GGAGCCCTTCTTCTTGCT (SEQ ID NO: 202)	CGG	-
14	EMX1	GGGCAACCACAAACCACGA (SEQ ID NO: 203)	GGG	+
15	EMX1	GCTCCCATCACATCAACCGG (SEQ ID NO: 204)	TGG	+

Target site sequences. Tested target sites for *S. pyogenes* type II CRISPR system with the requisite PAM. Cells were transfected with Cas9 and either crRNA-tracrRNA or chimeric sgRNA for each target.

TABLE I-continued

Target site sequences. Tested target sites for *S. pyogenes* type II CRISPR system with the requisite PAM. Cells were transfected with Cas9 and either crRNA-tracrRNA or chimeric sgRNA for each target.

Target site ID	genomic target	Target site sequence (5' to 3')	PAM strand
16	EMX1	GTGGCGCATTGCCACGAAGC (SEQ ID NO: 205)	AGG +
17	EMX1	GGCAGAGTGCTGCTTGCTGC (SEQ ID NO: 206)	TGG +
18	EMX1	GCCCCTGCGTGGGCCAAGC (SEQ ID NO: 207)	TGG +
19	EMX1	GAGTGGCCAGAGTCCAGCTT (SEQ ID NO: 208)	GGG -
20	EMX1	GGCCTCCCCAAAGCCTGGCC (SEQ ID NO: 209)	AGG -
4	PVALB	GGGGCCGAGATTGGGTGTTT (SEQ ID NO: 210)	AGG +
5	PVALB	GTGGCGAGAGGGGCCGAGAT (SEQ ID NO: 211)	TGG +
1	SERPINB5	GAGTGCCCGGAGGCGGGC (SEQ ID NO: 212)	GGG +
2	SERPINB5	GGAGTGCCCGGAGGCGGGG (SEQ ID NO: 213)	CGG +
3	SERPINB5	GGAGAGGAGTGCCCGGAGG (SEQ ID NO: 214)	CGG +

TABLE J

Primer sequences		
SURVEYOR assay		
primer name	genomic target	primer sequence (5' to 3')
Sp-EMX1-F1	EMX1	AAAACCACCCTTCTCTCTGGC (SEQ ID NO: 36)
Sp-EMX1-R1	EMX1	GGAGATTGGAGACACGGAGAG (SEQ ID NO: 37)
Sp-EMX1-F2	EMX1	CCATCCCCTTCTGTGAATGT (SEQ ID NO: 215)
Sp-EMX1-R2	EMX1	GGAGATTGGAGACACGGAGA (SEQ ID NO: 216)
Sp-PVALB-F	PVALB	CTGGAAAAGCCAATGCCTGAC (SEQ ID NO: 38)
Sp-PVALB-R	PVALB	GGCAGCAAACCTTGTCTCT (SEQ ID NO: 39)
qRT-PCR for Cas9 and sgRNA expression		
primer name	primer sequence (5' to 3')	
sgRNA reverse-strand synthesis	AAGCACCGACTCGGTGCCAC (SEQ ID NO: 217)	

TABLE J-continued

Primer sequences	
EMX1.1 sgRNA qPCR F	TCACCTCCAATGACTAGGGG (SEQ ID NO: 218)
EMX1.1 sgRNA qPCR R	CAAGTTGATAACGGACTAGCCT (SEQ ID NO: 219)
EMX1.3 sgRNA qPCR F	AGTCCGAGCAGAAGAAGAAGTTT (SEQ ID NO: 220)
EMX1.3 sgRNA qPCR R	TTTCAAGTTGATAACGGACTAGCCT (SEQ ID NO: 221)
Cas9 qPCR F	AAACAGCAGATTCCGCTGGA (SEQ ID NO: 222)
Cas9 qPCR R	TCATCCGCTCGATGAAGCTC (SEQ ID NO: 223)
GAPDH qPCR F	TCCAAAATCAAGTGGGGCGA (SEQ ID NO: 224)
GAPDH qPCR R	TGATGACCCTTTTGGCTCCC (SEQ ID NO: 225)
Bisulfite PCR and sequencing	
primer name	primer sequence (5' to 3')
Bisulfite PCR F (SERPINB5 locus)	GAGGAATCTTTTTTGTTYGAATATGTTGGAG (SEQ ID NO: 226)
Bisulfite PCR R (SERPINB5 locus)	GAGAAGCTTAAATAAAAAACRACAATACTCAACC CAACAACC (SEQ ID NO: 227)
pUC19 sequencing	CAGGAAACAGCTATGAC (SEQ ID NO: 228)

TABLE K

Sequences for primers to test sgRNA architecture. Primers hybridize to the reverse strand of the U6 promoter unless otherwise indicated. The U6 priming site is in italics, the guide sequence is indicated as a stretch of Ns, the direct repeat sequence is highlighted in bold, and the tracrRNA sequence underlined. The secondary structure of each sgRNA architecture is shown in FIG. 43.

primer name	primer sequence (5' to 3')
U6-Forward	<i>GCCTCTAGAGGTACCTGAGGGCCTATTTCCCATGATTCC</i> (SEQ ID NO: 229)
I: sgRNA(DR + 12, tracrRNA + 85)	<u>ACCTCTAGAAAAAAGCACCGACTCGGTGCCACTTTTTCAAGT</u> <u>TGATAACGGACTAGCCTTATTTAACTTGCTATTCTAGCTCT</u> AAAACNNNNNNNNNNNNNNNNNNNNNNNGGTGTTTCGTCCTTCC ACAAG (SEQ ID NO: 230)
II: sgRNA(DR + 12, tracrRNA + 85) mut2	<u>ACCTCTAGAAAAAAGCACCGACTCGGTGCCACTTTTTCAAGT</u> <u>TGATAACGGACTAGCCTTATATTAAGTTGCTATTCTAGCTCT</u> AATACNNNNNNNNNNNNNNNNNNNNNNNGGTGTTTCGTCCTTTCCA CAAG (SEQ ID NO: 231)
III: sgRNA(DR + 22, tracrRNA + 85)	<u>ACCTCTAGAAAAAAGCACCGACTCGGTGCCACTTTTTCAAGT</u> <u>TGATAACGGACTAGCCTTATTTAACTTGCTATGCTGTTTGTG</u> TCCAAAACAGCATAGCTCTAAAACNNNNNNNNNNNNNNNNNN NNNNGGTGTTCGTCCTTTCCACAAG (SEQ ID NO: 232)
IV: sgRNA(DR + 22, tracrRNA + 85) mut4	<u>ACCTCTAGAAAAAAGCACCGACTCGGTGCCACTTTTTCAAGT</u> <u>TGATAACGGACTAGCCTTATATTAAGTTGCTATGCTGTATTGT</u> TTCCAATACAGCATAGCTCTAATACNNNNNNNNNNNNNNNNNN NNNNGGTGTTCGTCCTTTCCACAAG (SEQ ID NO: 233)

TABLE L

Target sites with alternate PAMs for testing PAM specificity of Cas9. All target sites for PAM specificity testing are found within the human EMX1 locus.

Target site sequence (5' to 3')	PAM
AGGCCCCAGTGGCTGCTCT (SEQ ID NO: 234)	NAA
ACATCAACCGGTGGCGCAT (SEQ ID NO: 235)	NAT
AAGGTGTGGTTCCAGAACC (SEQ ID NO: 236)	NAC
CCATCACATCAACCGTGG (SEQ ID NO: 237)	NAG
AAACGGCAGAAGCTGGAGG (SEQ ID NO: 238)	NTA
GGCAGAAGCTGGAGGAGGA (SEQ ID NO: 239)	NTT
GGTGTGGTTCCAGAACCGG (SEQ ID NO: 240)	NTC
AACCGGAGGACAAAGTACA (SEQ ID NO: 241)	NTG
TTCCAGAACCGGAGGACAA (SEQ ID NO: 242)	NCA
GTGTGGTTCCAGAACCGGA (SEQ ID NO: 243)	NCT
TCCAGAACCGGAGGACAAA (SEQ ID NO: 244)	NCC
CAGAAGCTGGAGGAGGAAG (SEQ ID NO: 245)	NCG
CATCAACCGGTGGCGCATT (SEQ ID NO: 246)	NGA
GCAGAAGCTGGAGGAGGAA (SEQ ID NO: 247)	NGT
CCTCCCTCCCTGGCCAGG (SEQ ID NO: 248)	NGC
TCATCTGTGCCCTCCCTC (SEQ ID NO: 249)	NAA

TABLE L-continued

Target sites with alternate PAMs for testing PAM specificity of Cas9. All target sites for PAM specificity testing are found within the human EMX1 locus.

Target site sequence (5' to 3')	PAM
GGGAGGACATCGATGTCAC (SEQ ID NO: 250)	NAT
CAAACGGCAGAAGCTGGAG (SEQ ID NO: 251)	NAC
GGGTGGGCAACCACAAACC (SEQ ID NO: 252)	NAG
GGTGGGCAACCACAAACC (SEQ ID NO: 253)	NTA
GGCTCCCATCACATCAACC (SEQ ID NO: 254)	NTT
GAAGGGCCTGAGTCCGAGC (SEQ ID NO: 255)	NTC
CAACCGGTGGCGCATTGCC (SEQ ID NO: 256)	NTG
AGGAGGAAGGCCTGAGTC (SEQ ID NO: 257)	NCA
AGCTGGAGGAGGAAGGGCC (SEQ ID NO: 258)	NCT
GCATTGCCACGAAGCAGGC (SEQ ID NO: 259)	NCC
ATTGCCACGAAGCAGGCCA (SEQ ID NO: 260)	NCG
AGAACCGGAGGACAAAGTA (SEQ ID NO: 261)	NGA
TCAACCGGTGGCGCATTGC (SEQ ID NO: 262)	NGT
GAAGCTGGAGGAGGAAGGG (SEQ ID NO: 263)	NGC

-continued

ttaccgtaacttgaaagtatttcgatttcttggctttatatatcttgtggaaggacgaaacaccNNNNNNNNNNNNNNNNNNNN
NNggttttagagctagaaa**tagcaag**tt**aaaa**ta**aggctagtc**cg**ttatca**acttg**aaaa**agt**ggcaccgagtc**cg**gtgc**TTTTTT

T
(guide sequence is in bold Ns and the tracrRNA fragment is in bold)

>CBh-NLS-SpCas9-NLS

(SEQ ID NO: 59)

CGTTACATAACTTACGGTAAATGGCCCCCTGGCTGACCGCCCAACGACC
CCCGCCCATTTGACGTCAATAATGACGTATGTTCCCATAGTAACGCCAATAGGGACTT
TCCATTGACGTCAATGGGTGGAGTATTTACGGTAAACTGCCCACTGGCAGTACATC
AAGTGTATCATATGCCAAGTACGCCCCCTATTGACGTCAATGACGGTAAATGGCCCG
CCTGGCATTATGCCAGTACATGACCTTATGGGACTTCTACTTGGCAGTACATCTA
CGTATTAGTCATCGTATTACCATGGTCGAGGTGAGCCCCAGTTCTGCTTCACTCTC
CCCATCTCCCCCCCCTCCCCACCCCAATTTTGTATTTATTTATTTTAAATTATTTTGT
TGCAGCGATGGGGCGGGGGGGGGGGGGCGCGCCAGGGCGGGCGGGGGCGG
GGCGAGGGGGCGGGGGCGAGGCGGAGAGGTGCGGGCGGCAATCAGAGCG
GCGCGCTCCGAAAGTTTCTTTTATGGCGAGGCGGCGGGCGGCCCTATAAA
AAGCGAAGCGCGGGGGGGAGTCGCTGCGACGCTGCCTTCGCCCCGTGCCCC
GCTCCGCCGCCCTCGCGCCGCCCGCCCCGGCTCTGACTGACCGGTTACTCCAC
AGGTGAGCGGGCGGACGGCCCTTCTCCTCCGGGTGTAATTAGCTGAGCAAGAGG
TAAGGGTTAAGGGATGGTTGGTTGGTGGGTATTAATGTTAATTACCTGGAGCAC
CTGCCTGAAATCACTTTTTTTCAGGTGGaccggtgccacc**ATGGACTATAAGGACCAG**
ACGGAGACTACAAGGATCATGATATTGATTACAAAGACGATGACGATAAGATG
GCCCCAAAGAAGAAAGCGAAGGTCGGTATCCACGGAGTCCCAGCAGCCGACAA
GAAGTACAGCATCGGCTGGACATCGGCACCAACTCTGTGGGCTGGGCGGTGA
TCACCGACGAGTACAAGGTGCCAGCAAGAAATCAAGGTGCTGGGCAACACC
GACCGGCACAGCATCAAGAAGAACCTGATCGGAGCCCTGCTGTTGACAGCGG
CGAAACAGCCGAGGCCACCCGGCTGAAAGAGAACCGCCAGAAGAAGATACACCA
GACGGAAGAACC GGATCTGCTATCTGCAAGAGATCTTCAGCAACGAGATGGCC
AAGTGGAAGCAGCTTCTTCCACAGACTGGAAGAGTCTTCTGTTGGAAGA
GGATAAGAAGCAGAGCGGCAACCCATCTTCGGCAACATCGTGGACGAGGTGG
CCTACCACGAGAAGTACCCACCATCTACCACCTGAGAAAGAACTGGTGGAC
AGCACCGACAAGGCCGACCTGCGGCTGATCTATCTGGCCCTGGCCACATGAT
CAAGTTCGGGGCCACTTCTGATCGAGGGCGACCTGAACCCGCAACAGCG
ACGTGGACAAGCTGTTTCCAGCTGGTGCAGACCTACAACCAGCTGTTTCGAG
GAAAACCCCATCAACGCCAGCGGCTGGACGCCAAGGCCATCCTGTCTGCCAG
ACTGAGCAAGAGCAGACGGCTGAAAAATCTGATCGCCAGCTGCCCGCGAGA
AGAAGAATGGCCTGTTTCGGCAACCTGATGGCCCTGAGCCTGGGCTGACCCCC
AACTTCAAGAGCAACTTCGACCTGGCCGAGGATGCCAACTGCAGCTGAGCAA
GGACACCTACGACGACACCTGGACAACTGCTGGCCAGATCGGGGACCAAT
ACGCCGACCTGTTTCTGGCCCAAGAACCTGTCGACGCCATCCTGCTGAGC

-continued

GACATCCTGAGAGTGAACACCGAGATCACCAAGGCCCCCTGAGCGCCTCTAT
GATCAAGAGATACGACGAGCACCCACGACCTGACCTGTGAAAGCTCTCG
TGCGGCAGCAGCTGCCTGAGAAGTACAAAGAGATTTCTTCGACCAGAGCAAG
AACGGCTACGCCGGCTACATTGACGGCGGAGCCAGCCAGGAAGAGTTCTACAA
GTTTCATCAAGCCCATCCTGAAAAGATGGACGGCACCGAGGAACTGCTCGTGA
AGCTGAACAGAGAGGACCTGCTGCGGAAGCAGCGGACCTTCGACAACGGCAG
CATCCCCCACCAGATCCACCTGGGAGAGCTGCACGCCATTCTGCGGGCGCAGG
AAGATTTTACCCATTCTGAAGGACAACCGGAAAAGATCGAGAAGATCCTG
ACCTTCGCATCCCCTACTACGTGGGCCCTCTGGCCAGGGGAAAACAGCAGATT
CGCCTGGATGACCAGAAAGAGCGAGGAAACCATCACCCCTGGAACTTCGAGG
AAGTGGTGGACAAGGGCGCTTCGCCCCAGAGCTTCATCGAGCGGATGACCAAC
TTCGATAAGAACCTGCCAACGAGAAGGTGCTGCCAACGACAGCCTGCTGTA
CGAGTACTTCACCGTGTATAACGAGCTGACCAAAGTAAAATACGTGACCGAGG
GAATGAGAAAGCCCGCTTCTGAGCGCGAGCAGAAAAGGCCATCGTGGAC
CTGCTGTTCAAGACCAACCGGAAAGTGACCGTGAAGCAGCTGAAAAGGACTA
CTTCAAGAAAATCGAGTCTTCCGACTCCGTGAAAATCTCCGGCGTGGAAAGATC
GGTTCAACGCCTCCCTGGGCACATACCACGATCTGCTGAAAATATCAAGGAC
AAGGACTTCTGGACAAAGAGGAAAACGAGGACATTCTGGAAGATATCGTGCT
GACCTGACACTGTTGAGGACAGAGAGATGATCGAGGAACGGCTGAAAACCT
ATGCCACCTGTTTCGACGACAAAAGTGATGAAGCAGCTGAAGCGGCGGAGATAC
ACCGGCTGGGGCAGGCTGAGCCGGAAGCTGATCAACGGCATCCGGGACAAGC
AGTCCGGCAAGACAATCCTGGATTTCTGAAGTCCGACGGCTTCGCCAACAGA
AACTTCATGCAGCTGATCCACGACGACAGCCTGACCTTTAAGAGGACATCCA
GAAAGCCAGGTGTCCGGCCAGGGCGATAGCCTGCAAGGACATTTGCCAATC
TGGCCGGCAGCCCCGCCATTAAGAAAGGGCATCCTGCAGACAGTGAAGTGGTG
GACGAGCTCGTGAAAAGTGTGGCCGGCACAAAGCCGAGAACATCGTGATCGA
AATGGCCAGAGAGAACCAGACCAACCGAAGGGACAGAAAGAACAGCCGCGAG
AGAAAGAAAGCGGATCGAAGAGGGCATCAAAGAGCTGGGCAGCCAGATCCTGAA
AGAACAACCCGTGGAAAACACCCAGCTGCAGAACGAGAAGCTGTACTGTACT
ACCTGCAGAAATGGGCGGATATGTACGTGGACCAGGAACTGGACATCAAACGG
CTGTCCGACTACGATGTGGACCATATCGTGCCTCAGAGCTTTCTGAAGGACGA
CTCCATCGACAAACAAGGTGCTGACCAGAAGCGACAAAGAACCGGGGCAAGAGCG
ACAACGTGCCCTCCGAAAGAGTGTGAAAGAGATGAAGAACTACTGGCGGCGAG
CTGCTGAAACGCCAAGCTGATTACCCAGAGAAAAGTTCGACAATCTGACCAAGGC
CGAGAGAGGGCCCTGAGCGAACTGGATAAGGCCGGCTTCATCAAGAGACAG
CTGGTGGAAAACCGGCAGATCAAAAAGCACGTGGCACAGATCCTGGACTCCCG
GATGAACACTAAGTACGACGAGAAATGACAAGCTGATCCGGGAAGTGAAGTGA
TCACCCCTGAAGTCCAAGCTGGTGTCCGATTTCCGGAAGGATTTCCAGTTTTACA
AAGTGCAGGAGATCAACAACTACCAACCGCCACGACGCCTACCTGAAACGCC

-continued

GTCGTGGGAACCGCCCTGATCAAAAAGTACCCCTAAGCTGGAAAGCGAGTTCGT
 GTACGGCGACTACAAGGTGTACGACGTGCGGAAGATGATCGCCAAGAGCGAGC
 AGGAAATCGGCAAGGCTACCGCCAAGTACTTCTTCTACAGCAACATCATGAACT
 TTTTCAAGACCGAGATTACCCTGGCCAACGGCGAGATCCGGAAGCGGCCTCTG
 ATCGAGACAAACGGCGAAAACGGGGAGATCGTGTGGGATAAGGGCCGGGATT
 TTGCCACCGTGCAGAAAAGTGTGAGCATGCCCAAGTGAATATCGTGAAAAAG
 ACCGAGGTGCAGACAGGCGGCTTCAGCAAAGAGTCTATCCTGCCCAAGAGGAA
 CAGCGATAAGCTGATCGCCAGAAAAGAGACTGGGACCCTAAGAAGTACGGCG
 GCTTCGACAGCCCCACCGTGCCTATTCTGTGCTGGTGGTGGCCAAAAGTGGAA
 AAGGGCAAGTCCAAGAACTGAAGAGTGTGAAAGAGCTGCTGGGGATCACCAT
 CATGGAAAAGAGCAGCTTCGAGAAGAATCCCATCGACTTCTGGAAGCCAAGG
 GCTACAAAAGAAAGTAAAAAGGACCTGATCATCAAGCTGCCTAAGTACTCCCTG
 TTCGAGCTGGAAAACGGCCGGAAGAGAATGCTGGCCTCTGCCGGCGAACTGCA
 GAAGGAAACGAACTGGCCCTGCCCTCCAATATGTGAACCTCCTGTACCTGG
 CCAGCCACTATGAGAAGCTGAAGGGCTCCCCGAGGATAATGAGCAGAAAACAG
 CTGTTTGTGGAAACAGCACAAGCACTACCTGGACGAGATCATCGAGCAGATCAG
 CGAGTTCTCCAAGAGAGTGATCCTGGCCGACGCTAATCTGGACAAAAGTCTGT
 CCGCCTACAACAAGCACCGGGATAAGCCCATCAGAGAGCAGGCCGAGAATATC
 ATCCACCTGTTTACCCTGACCAATCTGGGAGCCCCTGCCGCTTCAAGTACTTT
 GACACCACCATCGACCGAAGAGGTACACCAGCACCAAAAGAGGTGCTGGACGC
 CACCCTGATCCACCAGGCATCACCGGCCTGTACGAGACACGGATCGACCTGT
 CTCAGCTGGGAGGCGACTTCTTTTCTTAGCTTGACCAGCTTCTTAGTAGCA

GCAGGACGCTTFAA

(NLS-hSpCas9-NLS is highlighted in bold)

>Sequencing amplicon for EMX1 guides 1.1, 1.14, 1.17

(SEQ ID NO: 264)

CCAATGGGGAGGACATCGATGTCACCTCCAATGACTAGGGTGGGCAACC
 ACAAAACCACGAGGGCAGAGTGCTGCTTGCTGCTGGCCAGGCCCTGCGTGGGCC
 AAGCTGGACTCTGGCCAC

>Sequencing amplicon for EMX1 guides 1.2, 1.16

(SEQ ID NO: 265)

CGAGCAGAAGAAGAAGGGCTCCCATCACATCAACCGGTGGCGCATTGCC
 ACGAAGCAGGCCAATGGGGAGGACATCGATGTCACCTCCAATGACTAGGGTGGGCA
 ACCACAAACCCACGAG

>Sequencing amplicon for EMX1 guides 1.3, 1.13, 1.15

(SEQ ID NO: 266)

GGAGGACAAAGTACAAACGGCAGAAGCTGGAGGAGGAAGGGCCTGAGTC
 CGAGCAGAAGAAGAAGGGCTCCCATCACATCAACCGGTGGCGCATTGCCACGAAGC
 AGCCCAATGGGAGGACATCGAT

>Sequencing amplicon for EMX1 guides 1.6

(SEQ ID NO: 267)

AGAAGCTGGAGGAGGAAGGGCTGAGTCCGAGCAGAAGAAGAAGGGCTC
 CCATCACATCAACCGGTGGCGCATTGCCACGAAGCAGGCCAATGGGGAGGACATCG
 ATGTCACCTCCAATGACTAGGGTGG

-continued

>Sequencing amplicon for EMX1 guides 1.10 (SEQ ID NO: 268)
 CCTCAGTCTTCCCATCAGGCTCTCAGCTCAGCCTGAGTGTGAGGCCCCAG
 TGGCTGCTCTGGGGCCCTCCTGAGTTTCTCATCTGTGCCCTCCCTCCCTGGCCAGG
 TGAAGGTGTGGTTCCA

>Sequencing amplicon for EMX1 guides 1.11, 1.12 (SEQ ID NO: 269)
 TCATCTGTGCCCTCCCTCCCTGGCCAGGTGAAGGTGTGGTTCCAGAACC
 GGAGGACAAAGTACAAACGGCAGAAGCTGGAGGAGGAAGGGCCTGAGTCCGAGCA
 GAAGAAGAAGGGCTCCCATCACA

>Sequencing amplicon for EMX1 guides 1.18, 1.19 (SEQ ID NO: 270)
 CTCCAATGACTAGGGTGGGCAACCACAAACCCACGAGGGCAGAGTGCTG
 CTTGCTGCTGGCCAGGCCCTGCGTGGGCCCAAGCTGGACTCTGGCCACTCCCTGGC
 CAGGCTTTGGGAGGCCCTGGAGT

>Sequencing amplicon for EMX1 guides 1.20 (SEQ ID NO: 271)
 CTGCTTGCTGCTGGCCAGGCCCTGCGTGGGCCCAAGCTGGACTCTGGCC
 ACTCCCTGGCCAGGCTTTGGGGAGGCCCTGGAGTCATGGCCCCACAGGGCTTGAAGC
 CCGGGGCCGCCATTGACAGAG

>T7 promoter F primer for annealing with target strand (SEQ ID NO: 272)
 GAAATTAATACGACTCACTATAGGG

>oligo containing pUC19 target site 1 for methylation (T7 reverse) (SEQ ID NO: 273)
 AAAAAAGCACCGACTCGGTGCCACTTTTTCAAGTTGATAACGGACTAGCC
 TTATTTTAACTTGCTATTTCTAGCTCTAAAACAACGACGAGCGTGACACCACCTAT
 AGTGAGTCGTATTAATTTT

>oligo containing pUC19 target site 2 for methylation (T7 reverse) (SEQ ID NO: 274)
 AAAAAAGCACCGACTCGGTGCCACTTTTTCAAGTTGATAACGGACTAGCC
 TTATTTTAACTTGCTATTTCTAGCTCTAAAACGCAACAATTAATAGACTGGACCTATA
 GTGAGTCGTATTAATTTT

Example 11

Oligo-Mediated Cas9-Induced Homologous
Recombination

[0349] The oligo homologous recombination test is a comparison of efficiency across different Cas9 variants and different HR template (oligo vs. plasmid).

[0350] 293FT cells were used. SpCas9=Wildtype Cas9 and SpCas9n=nickase Cas9 (D10A). The chimeric RNA target is the same EMX1 Protospacer Target 1 as in Examples 5, 9 and 10 and oligos synthesized by IDT using PAGE purification.

[0351] FIG. 44 depicts a design of the oligo DNA used as Homologous Recombination (HR) template in this experiment. Long oligos contain 100 bp homology to the EMX1 locus and a HindIII restriction site. 293FT cells were co-transfected with: first, a plasmid containing a chimeric RNA targeting human EMX1 locus and wild-type cas9 protein, and second, the oligo DNA as HR template. Samples are from 293FT cells collected 96 hours post transfection with Lipofectamine 2000. All products were amplified with an

EMX1 HR Primer, gel purified, followed by digestion with HindIII to detect the efficiency of integration of HR template into the human genome.

[0352] FIGS. 45 and 46 depict a comparison of HR efficiency induced by different combination of Cas9 protein and HR template. The Cas9 construct used were either wild-type Cas9 or the nickase version of Cas9 (Cas9n). The HR template used were: antisense oligo DNA (Antisense-Oligo in above figure), or sense oligo DNA (Sense-Oligo in above figure), or plasmid HR template (HR template in above figure). The sense/anti-sense definition is that the actively-transcribed strand with sequence corresponding to the transcribed mRNA is defined as the sense strand of genome. HR Efficiency is shown as percentage of HindIII digestion band as against all genomic PCR amplified product (bottom numbers).

Example 12

Autistic Mouse

[0353] Recent large-scale sequencing initiatives have produced a large number of genes associated with disease.

Discovering the genes is only the beginning in understanding what the gene does and how it leads to a diseased phenotype. Current technologies and approaches to study candidate genes are slow and laborious. The gold standards, gene targeting and genetic knockouts, require a significant investment in time and resources, both monetary and in terms of research personnel. Applicants set out to utilize the hSpCas9 nuclease to target many genes and do so with higher efficiency and lower turnaround compared to any other technology. Because of the high efficiency of hSpCas9 Applicants can do RNA injection into mouse zygotes and immediately get genome-modified animals without the need to do any preliminary gene targeting in mESCs.

[0354] Chromodomain helicase DNA binding protein 8 (CHD8) is a pivotal gene involved in early vertebrate development and morphogenesis. Mice lacking CHD8 die during embryonic development. Mutations in the CHD8 gene have been associated with autism spectrum disorder in humans. This association was made in three different papers published simultaneously in Nature. The same three studies identified a plethora of genes associated with autism spectrum disorder. Applicants' aim was to create knockout mice for the four genes that were found in all papers, Chd8, Katnal2, Kctd13, and Scn2a. In addition, Applicants chose two other genes associated with autism spectrum disorder, schizophrenia, and ADHD, GIT1, CACNA1C, and CACNB2. And finally, as a positive control Applicants decide to target MeCP2.

[0355] For each gene Applicants designed three gRNAs that would likely knockout the gene. A knockout would occur after the hSpCas9 nuclease makes a double strand break and the error prone DNA repair pathway, non-homologous end joining, corrects the break, creating a mutation. The most likely result is a frameshift mutation that would knockout the gene. The targeting strategy involved finding proto-spacers in the exons of the gene that had a PAM sequence, NGG, and was unique in the genome. Preference was given to proto-spacers in the first exon, which would be most deleterious to the gene.

[0356] Each gRNA was validated in the mouse cell line, Neuro-N2a, by liposomal transient co-transfection with hSp-Cas9. 72 hours post-transfection genomic DNA was purified using QuickExtract DNA from Epicentre. PCR was performed to amplify the locus of interest. Subsequently the SURVEYOR Mutation Detection Kit from Transgenomics was followed. The SURVEYOR results for each gRNA and respective controls are shown in FIG. A1. A positive SURVEYOR result is one large band corresponding to the genomic PCR and two smaller bands that are the product of the SURVEYOR nuclease making a double-strand break at the site of a mutation. The average cutting efficiency of each gRNA was also determined for each gRNA. The gRNA that was chosen for injection was the highest efficiency gRNA that was the most unique within the genome.

[0357] RNA (hSpCas9+gRNA RNA) was injected into the pronucleus of a zygote and later transplanted into a foster mother. Mothers were allowed to go full term and pups were sampled by tail snip 10 days postnatal. DNA was extracted and used as a template for PCR, which was then processed by SURVEYOR. Additionally, PCR products were sent for sequencing. Animals that were detected as being positive in either the SURVEYOR assay or PCR sequencing would have their genomic PCR products cloned into a pUC19 vector and sequenced to determine putative mutations from each allele.

[0358] So far, mice pups from the Chd8 targeting experiment have been fully processed up to the point of allele

sequencing. The Surveyor results for 38 live pups (lanes 1-38) 1 dead pup (lane 39) and 1 wild-type pup for comparison (lane 40) are shown in FIG. A2. Pups 1-19 were injected with gRNA Chd8.2 and pups 20-38 were injected with gRNA Chd8.3. Of the 38 live pups, 13 were positive for a mutation. The one dead pup also had a mutation. There was no mutation detected in the wild-type sample. Genomic PCR sequencing was consistent with the SURVEYOR assay findings.

Example 13

CRISPR/Cas-Mediated Transcriptional Modulation

[0359] FIG. 67 depicts a design of the CRISPR-TF (Transcription Factor) with transcriptional activation activity. The chimeric RNA is expressed by U6 promoter, while a human-codon-optimized, double-mutant version of the Cas9 protein (hSpCas9m), operably linked to triple NLS and a VP64 functional domain is expressed by a EF1a promoter. The double mutations, D10A and H840A, renders the cas9 protein unable to introduce any cleavage but maintained its capacity to bind to target DNA when guided by the chimeric RNA.

[0360] FIG. 68 depicts transcriptional activation of the human SOX2 gene with CRISPR-TF system (Chimeric RNA and the Cas9-NLS-VP64 fusion protein). 293FT cells were transfected with plasmids bearing two components: (1) U6-driven different chimeric RNAs targeting 20-bp sequences within or around the human SOX2 genomic locus, and (2) EF1a-driven hSpCas9m (double mutant)-NLS-VP64 fusion protein. 96 hours post transfection, 293FT cells were harvested and the level of activation is measured by the induction of mRNA expression using a qRT-PCR assay. All expression levels are normalized against the control group (grey bar), which represents results from cells transfected with the CRISPR-TF backbone plasmid without chimeric RNA. The qRT-PCR probes used for detecting the SOX2 mRNA is Taqman Human Gene Expression Assay (Life Technologies). All experiments represents data from 3 biological replicates, n=3, error bars show s.e.m.

Example 14

NLS: Cas9 NLS

[0361] 293FT cells were transfected with plasmid containing two components: (1) EF1a promoter driving the expression of Cas9 (wild-type human-codon-optimized Sp Cas9) with different NLS designs (2) U6 promoter driving the same chimeric RNA targeting human EMX1 locus.

[0362] Cells were collected at 72h time point post transfection, and then extracted with 50 μ l of the QuickExtract genomic DNA extraction solution following manufacturer's protocol. Target EMX1 genomic DNA were PCR amplified and then Gel-purify with 1% agarose gel. Genomic PCR product were re-anneal and subjected to the Surveyor assay following manufacturer's protocol. The genomic cleavage efficiency of different constructs were measured using SDS-PAGE on a 4-12% TBE-PAGE gel (Life Technologies), analyzed and quantified with ImageLab (Bio-rad) software, all following manufacturer's protocol.

[0363] FIG. 69 depicts a design of different Cas9 NLS constructs. All Cas9 were the human-codon-optimized version of the Sp Cas9. NLS sequences are linked to the cas9 gene at either N-terminus or C-terminus. All Cas9 variants with different NLS designs were cloned into a backbone vector containing so it is driven by EF1a promoter. On the same vector there is a chimeric RNA targeting human EMX1 locus driven by U6 promoter, together forming a two-component system.

TABLE M

Cas9 NLS Design Test Results. Quantification of genomic cleavage of different cas9-nls constructs by surveyor assay.

Percentage Genome Cleavage as measured by Surveyor assay	Biological Replicate 1 (%)	Biological Replicate 2 (%)	Biological Replicate 3 (%)	Average (%)	Error (S.E.M., standard error of the mean)
Cas9 (No NLS)	2.50	3.30	2.73	2.84	0.24
Cas9 with N-term NLS	7.61	6.29	5.46	6.45	0.63
Cas9 with C-term NLS	5.75	4.86	4.70	5.10	0.33
Cas9 with Double (N-term and C-term) NLS	9.08	9.85	7.78	8.90	0.60

[0364] FIG. 70 depicts the efficiency of genomic cleavage induced by Cas9 variants bearing different NLS designs. The percentage indicate the portion of human EMX1 genomic DNA that were cleaved by each construct. All experiments are from 3 biological replicates. n=3, error indicates S.E.M.

Example 15

Engineering of Microalgae using Cas9

[0365] Methods of Delivering Cas9

[0366] Method 1: Applicants deliver Cas9 and guide RNA using a vector that expresses Cas9 under the control of a constitutive promoter such as Hsp70A-Rbc S2 or Beta2-tubulin.

[0367] Method 2: Applicants deliver Cas9 and T7 polymerase using vectors that expresses Cas9 and T7 polymerase under the control of a constitutive promoter such as Hsp70A-Rbc S2 or Beta2-tubulin. Guide RNA will be delivered using a vector containing T7 promoter driving the guide RNA.

[0368] Method 3: Applicants deliver Cas9 mRNA and in vitro transcribed guide RNA to algae cells. RNA can be in vitro transcribed. Cas9 mRNA will consist of the coding region for Cas9 as well as 3'UTR from Cop1 to ensure stabilization of the Cas9 mRNA.

[0369] For Homologous recombination, Applicants provide an additional homology directed repair template.

[0370] Sequence for a cassette driving the expression of Cas9 under the control of beta-2 tubulin promoter, followed by the 3' UTR of Cop 1.

(SEQ ID NO: 275)

TCTTTCTTGGCTATGACACTTCCAGCAAAAGGTAGGGCGGGCTGCGAGA
CGGCTTCCCGCGCTGCATGCAACACCGATGATGCTTCCACCCCGAAG
CTCCTTCCGGGCTGCATGGGCGCTCCGATGCCGCTCCAGGGCGAGCGCTG
TTTAAATAGCCAGGCCCGGATTGCAAAGACATTATAGCGAGTACCAA
GCCATATTCAAACCTTAGATCACTACCACTTCTACACAGGCCACTCGAG
CTTGTGATCGCACTCCGCTAAGGGGGCGCTTCTCTTTCGTTTCAGTC
ACAACCCGCAACATGTACCATACGATGTTCCAGATTACGCTTCGCCGA
AGAAAAGCGCAAGGTGCAAGCGTCCGACAAGAAGTACAGCATCGGCCTG

- continued

GACATCGGCACCAACTCTGTGGGCTGGGCGGTGATCACCACGAGTACAA
GGTGCCAGCAAGAAATTCAGGTGCTGGGCAACACCGACCGGCACAGCA
TCAAGAAGAACCTGATCGGAGCCCTGCTGTTGACAGCGGGAAACAGCC
GAGGCCACCCGGCTGAAGAGAACCGCCAGAAGAGATACACCAGACGGAA
GAACCGGATCTGCTATCTGCAAGAGATCTTCAGCAACGAGATGGCCAAGG
TGGACGACAGCTTCTCCACAGACTGGAAGAGTCTTCTGTTGGAAGAG
GATAAGAAGCACGAGCGGCACCCCATCTTCGGCAACATCGTGGACGAGGT
GGCCTACCACGAGAAGTACCCACCATCTACCACCTGAGAAAAGAACTGG
TGGACAGCACCGACAAGCCGACCTGCGGCTGATCTATCTGGCCCTGGCC
CACATGATCAAGTTCGGGGCCACTTCTGATCGAGGGCGACCTGAACCC
CGACAACAGCGACGTGGACAAGCTGTTTCATCCAGCTGGTGCAGACTACA
ACCAGCTGTTTCGAGGAAAACCCCATCAACGCCAGCGGCGTGGACGCCAAG
GCCATCTGTCTGCCAGACTGAGCAAGAGCAGACCGCTGGAAAATCTGAT
CGCCAGCTGCCCGCGAGAAGAAGAAATGGCTGTTTCGGCAACCTGATTG
CCCTGAGCCTGGGCTGACCCCAACTTCAAGAGCAACTTCGACCTGGCC
GAGGATGCCAAACTGCAGCTGAGCAAGGACACCTACGACGACGACCTGGA
CAACCTGCTGGCCAGATCGGCGACAGTACGCCACCTGTTTCTGGCCG
CCAAGAACCTGTCCGACGCCATCTGCTGAGCGACATCCTGAGAGTGAAC
ACCGAGATCACAAGGCCCCCTGAGCGCCTTATGATCAAGAGATACGA
CGAGACCACCAGGACCTGACCTGCTGAAAGCTCTCGTGGCGCAGCAGC
TGCTTGAGAAGTACAAGAGATTTTCTTTCGACCAGAGCAAGAACGGCTAC
GCCGGCTACATTGACGGCGGAGCCAGCCAGGAAGAGTTCTACAAGTTTCA
CAAGCCCATCTGGAAAAGATGGACGGCACCGAGGAACCTGCTCGTGAAGC
TGAAACAGAGAGGACCTGCTGCGGAAGCAGCGGACCTTCGACAACGGCAGC
ATCCCCCACCAGATCCACCTGGGAGAGCTGCACGCCATTCTGCGGGCGCA
GGAAGATTTTACCATTCTGAAGGACAACCGGAAAAGATCGAGAAGA
TCTTGACCTTCCGCATCCCTACTACGTGGGCCCTCTGCCCAGGGGAAAC
AGCAGATTCGCCTGGATGACCAGAAGAGCGAGGAACCATCACCCCTG

-continued

GAACCTCGAGGAAGTGGTGGACAAGGGCGCTTCCGCCAGAGCTTCATCG
 AGCGGATGACCAACTTCGATAAAGAACTGCCCAACGAGAAGGTGCTGCC
 AAGCACAGCCTGCTGTACGAGTACTTACCCTGTATAACGAGCTGACCAA
 AGTGAATACGTGACCGAGGAATGAGAAAGCCCGCTTCTGAGCGGG
 AGCAGAAAAGGCCATCGTGGACCTGCTGTTCAGACCAACCGAAAGTG
 ACCGTGAAGCAGCTGAAAGAGGACTACTTCAAGAAAATCGAGTGTTCGA
 CTCCGTGGAAATCTCCGGCGTGAAGATCGGTTCAACGCCTCCCTGGGCA
 CATAACCAGATCTGTGAAAATATCAAGGACAAGGACTTCTGGACAAT
 GAGGAAAACGAGGACATCTGGAAGATATCGTGTGACCTGACACTGTT
 TGAGGACAGAGAGATGATCGAGGAACGGCTGAAAACCTATGCCACCTGT
 TCGACGACAAAGTGATGAAGCAGCTGAAGCGCGGAGATACACCGCTGG
 GGCAGGCTGAGCCGGAAGCTGATCAACGGCATCCGGGACAAGCAGTCCGG
 CAAGACAATCTGGATTCTGAAGTCCGACGGCTTCCGCAACAGAACT
 TCATCGAGCTGATCCACGACGACAGCTGACCTTTAAGAGGACATCCAG
 AAAGCCAGGTGTCCGGCCAGGGCAGATAGCTGCACGACGACATTGCCAA
 TCTGGCCGGCAGCCCGCCATTAAGAGGGCATCTGCAGACAGTGAAGG
 TGGTGGACGAGCTCGTGAAGTGATGGGCCGACAAAGCCGAGAACATC
 GTGATCGAAAATGGCCAGAGAGAACAGACCACCCAGAAGGGCAGAAAGAA
 CAGCCGCGAGAGAAATGAAGCGGATCGAAGAGGGCATCAAAGAGCTGGGCA
 GCCAGATCTGAAAGAACACCCCGTGGAAAACACCCAGCTGCAGAACGAG
 AAGCTGTACTGTACTACCTGAGAAATGGCGGGATATGTACGTGGACCA
 GGAACCTGGACATCAACCGGCTGTCCGACTACGATGTGGACCATATCGTGC
 CTCAGAGCTTTCTGAAGGACGACTCCATCGACAACAAGTGTGACCAGA
 AGCGACAAGAACCAGGGCAAGAGCGACAACGTGCCCTCCGAAGAGGTGCT
 GAAGAAGATGAAGAACTACTGGCGGAGCTGCTGAACGCCAAGCTGATTA
 CCCAGAGAAAGTTCGACAATCTGACCAAGGCCGAGAGAGGGCCCTGAGC
 GAACTGGATAAAGCCGGCTTTCATCAAGAGACAGCTGGTGGAAAACCGCA
 GATCACAAGCAGCTGGCACAGATCTGGACTCCCGGATGAACACTAAGT
 ACGACGAGAATGACAAGCTGATCCGGGAAGTGAAGTGTACCCCTGAAG
 TCCAAGCTGGTGTCCGATTTCGGAAGGATTTCAGTTTTACAAAGTGC
 CGAGATCAACAATACCACCACGCCACGACGCTTACCTGAACGCGCTCG
 TGGGAACCGCCTGATCAAAAAGTACCCTAAGCTGGAAGCGAGTTCGTG
 TACGCGACTACAAGGTGTACGACGTGCGGAAGATGATCGCAAGAGCGA
 GCAGAAATCGCAAGGCTACCGCAAGTACTTCTTCTACAGCAACATCA
 TGAACTTTTTCAAGACCAGATTACCCTGGCCAACGGCGAGATCCGGAAG
 CGGCTCTGATCGAGACAACGGCGAAAACCGGGGAGATCGTGTGGGATAA
 GGGCCGGGATTTTGGCCACCGTGGGAAAGTGTGAGCATGCCCAAGTGA
 ATATCGTGAAGAAGACCAGGTGCAGACAGGCGGCTTCAGCAAGAGTCT
 ATCTGCCCAAGAGGAACAGCGATAAGCTGATCGCCAGAAAAGAGGACTG

-continued

GGACCCCTAAGAAGTACGGCGGCTTCGACAGCCCCACCCTGGCCTATTCTG
 TGCTGGTGGTGGCCAAAGTGGAAAAGGGCAAGTCCAAGAACTGAAGAGT
 GTGAAAGAGCTGCTGGGGATCACCATCATGGAAGAAGCAGCTTCGAGAA
 GAATCCCATCGACTTCTGGAAGCCAAGGGCTACAAGAAGTGAAAAAGG
 ACCTGATCATCAAGCTGCCTAAGTACTCCCTGTTTCGAGCTGAAAACGGC
 CGGAAGAGAATGCTGGCCTCTGCCGGCAACTGCAGAAGGGAAACGAACT
 GGCCCTGCCCTCCAAATATGTGAACTTCTGTACTGGCCAGCCACTATG
 AGAAGCTGAAGGGCTCCCCGAGGATAATGAGCAGAAAACAGCTGTTTGTG
 GAACAGCACAAAGCACTACCTGGACGAGATCATCGAGCAGATCAGCGAGTT
 CTCCAAGAGAGTGATCTGGCCGACGCTAATCTGGACAAAAGTGTGTCCG
 CCTACAACAAGCACCGGGATAAGCCCATCAGAGAGCAGGCCGAGAAATATC
 ATCCACCTGTTTACCCTGACCAATCTGGGAGCCCTGCCGCTTCAAGTA
 CTTTGACACCACCATCGACCGAAGAGGTACACCAGCACCAAAGAGGTGC
 TGGACGCCACCTGATCCACCAGAGCATCACCGGCTGTACGAGACACGG
 ATCGACCTGTCTCAGCTGGGAGGCGACAGCCCAAGAAGAAGAGAAAGGT
 GGAGGCCAGCTAAGGATCCGGCAAGACTGGCCCGCTTGGCAACGCAACA
 GTGAGCCCTCCCTAGTGTGTTGGGGATGTGACTATGTATTCGTGTGTT
 GGCCAACGGGTCAACCCGAACAGATTGATACCCGCTTGGCATTTCCTGT
 CAGAATGTAACGTGAGTTGATGGTACT

[0371] Sequence for a cassette driving the expression of T7 polymerase under the control of beta-2 tubulin promoter, followed by the 3' UTR of Cop1:

(SEQ ID NO: 276)

TCTTTCTTGCGCTATGACACTTCCAGCAAAAAGGTAGGGCGGGCTGCGAGA
 CGGCTTCCCGGCGCTGCATGCAACACCCGATGATGCTTCGACCCCCGAAG
 CTCTTTCGGGGCTGCATGGCGCTCCGATGCCGCTCCAGGGCGAGCGCTG
 TTTAAATAGCCAGGCCCCGATTGCAAAGACATTATAGCGAGCTACCAAA
 GCCATATTCAAACACCTAGATCACTACCATTCTACACAGGCCACTCGAG
 CTTGTGATCGCACTCCGCTAAGGGGGCGCTTCTCTTCTGTTTCAGTC
 ACAACCCGCAAAc atg cct aagaagaagaggaagggttaacacgattaaaca
 tcgctaagaacgacttctctgacatcgaactggctgctatccggttcaac
 actctggctgaccattacgggtgagcgttagctcgcaaacagttggccct
 tgagcatgagtcctacgagatgggtgaagcagcctccgcaagatggttg
 agcgtcaactaaagctggtaggttgcgataaacgctgcccaagcct
 ctcatcactaccctactccctaaagatgattgcacgcatcaacgactgggt
 tgaggaagtgaagctaaagcgggcaagcggccgacagcctccagttcc
 tgcaagaaatcaagcgggaagccgtagcgtacatcaccattaaagaccact
 ctggcttgcttaaccagtgctgacaatacaaccgttcaggctgtagcaag
 cgcaatcggctgggcatgaggacagggctcgttcggctgctatccggtg
 acctgaaagctaaagcacttcaagaaaaagcttgaggaacaactcaacaag

-continued

cgcgtagggcagctctacaagaaagcatttatgcaagtgtcgaggctga
 catgctctctaagggctactcgggtggcgaggcgtggtcttcggtgcata
 aggaagactctattcatgtaggagtacgctgcacgagatgctcattgag
 tcaaccggaatggttagcttacaccgccaaaatgctggcgtagtaggtca
 agactctgagactatcgaaactcgcacctgaatacgcgtgaggctatcgcaa
 cccgtgcagggtgcgctggtggcatctctccgatgttccaaacctgcgta
 gttcctcctaagcctggactggcattactggtggtggtcattgggctaa
 cggtegtgctcctctggcgctggtgcgtaactcacagtaagaaagcactga
 tgcgctacgaagacgctttacatgcctgagggtgtacaaagcattaacatt
 gcgcaaaacaccgcatggaatacaacaagaaagtcctagcggctgcgcaa
 cgtaatcaccaagtggaagcattgtccggctcgaggacatccctgcgattg
 agcgtgaagaactcccgatgaaaccggaagacatcgacatgaatcctgag
 gctctcacgcgctggaacgtgctgccgctgctgtgtaccgcaaggacaa
 ggctcgcaagtcctgcgctatcagccttgagttcatgcttgagcaagcca
 ataagttgctaaccataagggcattcgggtcccttacaacatggactgg
 cgcggtcgtggttacgctgtgtcaatgttcaaccgcaaggtaacgatat
 gaccaaaaggactgcttacgctggcgaaggtaaaccaatcggtaaggaaag
 gttactactggctgaaataccacggtgcaaacctgtgcgggtgctgacaag
 gttccggtccctgagcgcataaagttcattgaggaataaccacgagaacat
 catggccttgcgtaagctccactgggaaacacttggtgggctgagcaag
 attctccgtctgcttccctgcttctgctttgagtagcgtgggtgacag
 caccacggcctgagctataaactgctccctccgctggcggttgacgggtc
 ttgctctggcatccagcactctccgcgatgctccgagatgaggtagggtg
 gtcgcgcggttaacttgcttccctagtgaaaccggtcaggacatctacggg
 attgttgctaagaaagtcacagagattctacaagcagacgcaatcaatgg
 gaccgataacgaagtagttaccgtagccgatgagaacactggtgaaatct
 ctgagaaagtcaagctgggcaactaaggcactggctggtcaatggctggt
 tacgggtgtactcgcagtgtagtaagcgttcagtcacgctgctgctta
 cgggtccaaagagtccggtccgtcaacaagtgctggaagataccattc
 agccagctattgatccggcaagggtctgatgttactcagccgaatcag
 gctgctggatacatggctaagctgatttgggaatctgtgagcgtgacgg
 ggtagctcgggtgaaagcaatgaactggcttaagctcgtgctgtaagctgc
 tggctgctgaggtcaagataagaagactggagagattcttcgcaagcgt
 tgcgctgtgcatgggtaactcctgatggttccctgtgtggcaggaata
 caagaagcctattcagacgcgcttgaacctgatgttccctcggtcagttcc
 gcttacagcctaccattaacaccaacaagatagcgagattgatgcacac
 aaacaggagtctggatcgctcctaactttgtacacagccaagcggtag
 ccacctcgtgaagactgtagtggtggcaccagagaagtacggaatcgaat
 cttttgcactgattcacgactcctcggtagcatccggctgacgctgog

-continued

aacctgttcaaagcagtgccgcaaacatggttgacacatatgagtcttg
 tgatgtagtggctgatttctacgaccagttcgctgaccagttgcacgag
 ctcaattggacaaaaatgccagcacttccggctaaaggtaacctgacacct
 cgtgacatcttagagtcggacttccgcttcgctgaagGATCCGGCAAGAC
 TGGCCCCGCTTGGCAACGCAACAGTGAGCCCCCTCCCTAGTGTGTTGGGG
 ATGTGACTATGTATTCTGTGTTGGCCAACGGGTCAACCCGACAGATTG
 ATACCCGCCTTGGCATTTCCTGTCAGAATGTAACGTCAGTTGATGGTACT

[0372] Sequence of guide RNA driven by the T7 promoter (T7 promoter, Ns represent targeting sequence):

(SEQ ID NO: 277)

gaaatTAATACGACTCACTATANNNNNNNNNNNNNNNNNNNNNNgttttaga
 gctaGAAAtagcaagttaaaataaggctagtcgcttatcaacttgaaaaa
 gtggcaaccgagtcggtgcttttttt

[0373] Gene Delivery:

[0374] Chlamydomonas reinhardtii strain CC-124 and CC-125 from the Chlamydomonas Resource Center will be used for electroporation. Electroporation protocol follows standard recommended protocol from the GeneArt Chlamydomonas Engineering kit.

[0375] Also, Applicants generate a line of Chlamydomonas reinhardtii that expresses Cas9 constitutively. This can be done by using pChlamy1 (linearized using PvuI) and selecting for hygromycin resistant colonies. Sequence for pChlamy1 containing Cas9 is below. In this way to achieve gene knockout one simply needs to deliver RNA for the guideRNA. For homologous recombination Applicants deliver guideRNA as well as a linearized homologous recombination template.

pChlamy1-Cas9:

(SEQ ID NO: 278)

TGGCGTATTTACACCGCATCAGGTGGCACTTTTTCGGGAAATGTGCGCG
 GAACCCCTATTGTTTATTTTCTAAATACATTCAAATATGTATCCGCTC
 ATGAGATTATCAAAAAGGATCTTACCTAGATCCTTTTAAATTAATAATG
 AAGTTTTAAATCAATCTAAAGTATATATGAGTAAACTTGGTCTGACAGTT
 ACCAATGCTTAATCAGTGAGGCACCTATCTCAGCGATCTGTCTATTTTCT
 TCATCCATAGTTGCCTGACTCCCCGTCGTGTAGATAACTACGATACGGGA
 GGGCTTACCATCTGGCCCCAGTGTGCAATGATACCGCGAGACCCACGCT
 CACCGGCTCCAGATTTATCAGCAATAAACAGCCAGCCGGAAGGGCCGAG
 CGCAGAAGTGGTCTGCAACTTTATCCGCCCTCCATCCAGTCTATTAATTG
 TTGCCGGGAAGCTAGAGTAAGTAGTTCGCGAGTTAATAGTTTGCACAACG
 TTGTTGCCATTGCTACAGGCATCGTGGTGTACGCTCGTCTTTGGTATG
 GCTTCATTGAGTCCGGTCCCAACGATCAAGGCGAGTTACATGATCCCC
 CATGTTGTGCAAAAAGCGGTTAGCTCCTTCGGTCTCCGATCGTTGTCA
 GAAGTAAGTTGGCCGAGTGTATCACTCATGTTTATGGCAGCACTGCAT

-continued

AATTCTCTTACTGTATGCCATCCGTAAGATGCTTTTCTGTGACTGGTGA
GTACTCAACCAAGTCATTCTGAGAATAGTGTATGCGCGCACCAGTTGCT
CTTGCCCGCGCTCAATACGGGATAATACCGGCCACATAGCAGAACTTTA
AAAGTGCTCATCATTGAAAAAGCTTCTCGGGGCGAAAACTCTCAAGGAT
CTTACCGCTGTTGAGATCCAGTTTCGATGTAACCCACTCGTGCACCCA
GATCTTCAGCATCTTTACTTTTACCCAGCGTTTCTGGGTGAGCAAAAACA
GGAAGGCAAAATGCCGCAAAAAGGGAATAAGGGCGACACGGAATGTTG
AATACTCATACTCTTCTTTTCAATATATTGAAGCATTATCAGGGTT
ATTGTCTCATGACCAAAATCCCTTAACGTGAGTTTTCTGTTCCACTGAGCG
TCAGACCCCGTAGAAAAGATCAAAGGATCTTCTTGAGATCTTTTTTTCT
GCGCGTAATCTGCTGCTTGCAACAAAAAACCCGCTACCAGCGGTGG
TTTGTGTCGCGGATCAAGAGCTACCAACTCTTTTTCGAAGTA
TTCAGCAGAGCGCAGATACCAATACTGTTCTTCTAGTGTAGCCGTAGTT
AGGCCACCACTTCAAGAACTCTGTAGCACCCTACATACCTCGCTCTGC
TAATCCTGTTACAGTGCTGTTGCGCAGTGCGATAAGTCTGTTCTTACC
GGGTGACTCAAGACGATAGTTACCGGATAAGGCGCAGCGGTGGGGCTG
AACGGGGGTTCTGTGCACACAGCCAGCTTGGAGCGAACGACCTACACCG
AACTGAGATACCTACAGCGTATGAGAAAGCGCCACGCTTCCCGAA
GGGAGAAAGCGGACAGGTATCCGTAAGCGGCGGGTGGAAACAGGAGA
GCGCACGAGGGAGCTTCCAGGGGAAACGCTGATCTTTATAGTCTCTG
TCGGGTTTCGCCACCTCTGACTTGAGCGTCGATTTTGTGATGCTCGTCA
GGGGGGCGGAGCTATGGAAAAACCCAGCAACCGCGCTTTTTACGGTT
CCTGGCCTTTTGTGCTGCTTTTGTCTCACATGTTCTTTCTGCGTTATCCC
CTGATTCTGTGGATAACCGTATTACCGCCTTTGAGTGAAGCTGATACCGCT
CGCCGACCCGAACGACCGAGCGCAGCGAGTCTGAGCGAGGAAGCGGT
CGCTGAGGCTTGACATGATTTGGTGCATGTTTGTATGAAGTACAGGAC
TGATTTGGCGGCTATGAGGGCGGGGAAGCTCTGGAAGGGCGCGATGG
GGCGCGGGGCTCCAGAAGGCCATACGGCCGCTGGCGGCACCCATCC
GGTATAAAGCCCGCAGCCCGAACGGTACCTCCACTTTCAGCGACAAA
CGAGCACTTATACATACCGACTATTCTGCCCTATACATAACCACTCAG
CTAGCTTAAGATCCCATCAAGCTTGCATGCCGGGCGGCCAGAAGGAGCG
CAGCCAAACAGGATGATGTTTGTATGGGTATTTGAGCACTTGCAACCT
TATCCGGAAGCCCTGGCCCAAAAGGCTAGGCGCAATGCAAGCAGTT
CGCATGCAGCCCTGGAGCGGTGCCCTCCTGATAAACCGGCCAGGGGGC
TATGTTCTTACTTTTTTACAAGAGAAGTCACTCAACATCTTAAATGGC
CAGGTGAGTCGACGAGCAAGCCCGCGGATCAGGCGCGTCTTGCAGAT
TTGACTTGCAACGCCCGCATTTGTGTCGACGAAGGCTTTTGGCTCCTCTGT
CGCTGCTCAAGCAGCATCTAACCTGCGTGCCTGCTTCCATTTGCAGGA
GATTCGAGGTACCATGTACCCATACGATGTTCCAGATTACGCTTCGCCGA

-continued

AGAAAAAGCGCAAGGTCGAAGCGTCCGACAAGAGTACAGCATCGGCCCTG
GACATCGGCACCAACTCTGTGGGCTGGGCGGTGATACCCGACGAGTACAA
GGTGCCAGCAAGAAATTCGAAGTCTGGGCAACACCGACCGGCACAGCA
TCAAGAAGAACCTGATCGGAGCCCTGCTGTTTCGACAGCGGCGAAACAGCC
GAGGCCACCCGGCTGAAGAGAACCAGCAGAGAAGATACCCAGACGGAA
GAACCGGATCTGCTATCTGCAAGAGATCTTCAGCAACGAGATGGCCAAGG
TGGACGACAGCTTCTCCACAGACTGGAAGAGTCTTCTGTTGGTGAAGAG
GATAAGAAGCAGCAGCGGCACCCATCTTCGGCAACATCGTGGACGAGGT
GGCTTACCAGGAGTACCCACCATCTACCACTGAGAAAGAACTGG
TGGACAGCACCGACAAGGCGACCTGCGGCTGATCTATCTGGCCCTGGCC
CACATGATCAAGTTCGCGGCGCACTTCTGATCGAGGGCGACCTGAACCC
CGACAAACAGCGACGTGGACAAGCTGTTTCATCCAGTGGTGCAGACTACA
ACCAGCTGTTTCGAGGAAAAACCCATCAACGCCAGCGGCTGGACGCCAAG
GCCATCTGCTGTCGAGACTGAGCAAGAGCAGACGGCTGGAAATCTGAT
CGCCAGCTGCCCCGCGAGAAGAAGTGGCTGTTTCGGCAACCTGATTG
CCCTGAGCCTGGGCTGACCCCACTTCAAGAGCAACTTCGACCTGGCC
GAGGATGCCAAACTGCAGCTGAGCAAGGACACCTACGACGACGACCTGGA
CAACTGCTGGCCAGATCGGCGACAGTACCGCACCTGTTTCTGGCCG
CCAAGAACCTGTCGACGCCATCTGCTGAGCGACATCTGAGAGTGAAC
ACCGAGATCACCAAGGCCCTGAGCGCTCTATGATCAAGAGATACGA
CGAGCACACCAGGACCTGACCTGCTGAAAGCTCTGTCGCGGCGAGCAGC
TGCTGAGAAGTACAAGAGATTTTCTTCGACCAGAGCAAGAACGGCTAC
GCCGCTACATGACGGCGGAGCCAGCCAGGAAGGTTCTACAAGTTTCT
CAAGCCATCTGGAAGATGGACGGCACCGGAGAACTGCTGCTGAAGC
TGAAACAGAGAGGACCTGCTGCGGAAGCAGCGGACCTTCGACAACGGCAGC
ATCCCCCACCAGATCCACTGGGAGAGCTGCACGCCATTCTGCGGCGCA
GGAAGATTTTACCATTCTGAAGGACAACCGGAAAAAGATCGAGAAGA
TCTGACCTTCCGCATCCCTACTACGTGGGCCCTTGGCCAGGGGAAAC
AGCAGATTCGCTGGATGACAGAAAGAGCAGGAAACCATCACCCCTG
GAACCTCGAGGAAGTGGTGGACAAGGGCGCTTCCGCCAGAGCTTCATCG
AGCGGATGACCAACTTCGATAAAGAACCTGCCAACGAGAAGGTGCTGCC
AAGCACAGCTGCTGTACGAGTACTTCCCGTGTATAACGAGCTGACCAA
AGTGAATAACGTGACCGAGGGAATGAGAAAGCCGCTTCTGAGCGGCG
AGCAGAAAAAGGCCATCGTGGACCTGCTGTTCAAGACCAACCGAAAGTG
ACCGTGAAGCAGCTGAAAGAGGACTACTTCAAGAAAAATCGAGTCTTCGA
CTCCGTGGAATCTCCGGCGTGAAGATCGGTTCAACCGCTCCCTGGGCA
CATAACCAGATCTGCTGAAAATTATCAAGGACAAGGACTTCTGGACAAT
GAGGAAAAAGGACATCTGGAAGATATCGTGTGACCCCTGACACTGTT
TGAGGACAGAGAGATGATCGAGGAACGGCTGAAAACCTATGCCACCTGT

-continued

TCGACGACAAAGTGATGAAGCAGCTGAAGCGGCGGAGATACACCGGCTGG
GGCAGGCTGAGCCGGAAGCTGATCAACGGCATCCGGGACAAGCAGTCCGG
CAAGACAATCCTGGATTTCTGAAGTCCGACGGCTTCGCCAACAGAAACT
TCATGCAGCTGATCCACGACGACAGCTTGACCTTTAAAGAGGACATCCAG
AAAGCCAGGTGTCCGGCCAGGGCGATAGCTGCACGACGACATTGCCAA
TCTGGCCGGCAGCCCCGCCATTAAGAAGGGCATCCTGCAGACAGTGAAGG
TGGTGGACGAGCTCGTAAAGTGATGGGCCGGCACAAGCCCCGAGAACATC
GTGATCGAAATGGCCAGAGAGAACCAGACCACCCAGAAGGGCAGAAAGAA
CAGCCGCGAGAGAATGAAGCGGATCGAAGAGGGCATCAAAGAGCTGGGCA
GCCAGATCCTGAAAGAACACCCCGTGGAAAACACCCAGCTGCAGAACGAG
AAGCTGTACTGTACTACCTGCAGAATGGGCGGGATATGTACTGTGACCA
GGAACTGGACATCAACCGGCTGTCCGACTACGATGTGGACCATATCGTGC
CTCAGAGCTTTCTGAAGGACGACTCCATCGACAACAAGGTGCTGACCAGA
AGCGACAAGAACCGGGGCAAGAGCGACAACTGCCCTCCGAAGAGGTGCT
GAAGAAGATGAAGAATACTGGCGGAGCTGCTGAACGCCAAGCTGATTA
CCCAGAGAAAGTTCGACAATCTGACCAAGGCCGAGAGAGGCGGCTGAGC
GAATGGATAAGGCCGGCTTCATCAAGAGACAGCTGGTGGAAACCCGGCA
GATCACAAGCACGCTGGCACAGATCCTGGACTCCCGGATGAACACTAAGT
ACGACGAGAATGACAAGCTGATCCGGGAAGTGAAGTGATCACCTGAAG
TCCAAGCTGGTGTCCGATTTCCGGAAGGATTTCCAGTTTACAAAAGTGGC
CGAGATCAACAATACTACCACACGCCACGACGCTACCTGAACGCGCTCG
TGGGAACCGCCCTGATCAAAAAGTACCCTAAGCTGGAAAGCGAGTTCTGTG
TACGGCGACTACAAGGTGTACGACGTGCGGAAGATGATCGCCAAGAGCGA
GCAGGAAATCGGCAAGGCTACCGCAAGTACTTCTTCTACAGCAACATCA
TGAACTTTTCAAGACCGAGATTACCTGGCCAACGGCGAGATCCGGAAG
CGGCTCTGATCGAGACAAACGGCGAAACCGGGGAGATCGTGTGGGATAA
GGGCCGGGATTTTGCCACCGTCCGGAAGTGTGAGCATGCCCAAGTGA
ATATCGTGAAGAGACCGAGGTGACGACAGGCGGCTTCAGCAAAGAGTCT
ATCCTGCCCAAGAGGAACAGCGATAAGCTGATCGCCAGAAGAAGGACTG
GGACCTAAGAAGTACGGCGGCTTCGACAGCCCCACCGTGGCCTATTCTG
TGCTGGTGGTGGCCAAAGTGGAAAAGGGCAAGTCCAAGAACTGAAGAGT
GTGAAAGAGCTGCTGGGATCACCATCATGGAAGAAGCAGCTTCGAGAA
GAATCCATCGACTTTCTGGAAGCCAAGGGCTACAAAGAAGTGAAGAAAGG
ACCTGATCATCAAGCTGCCTAAGTACTCCCTGTTTCGAGCTGGAAAACGGC
CGGAAGAGAATGCTGGCCTCTGCCGGCGAACTGCAGAAGGGAACGAACT
GGCCTGCCCTCCAAATATGTGAACCTCCTGTACTTGGCCAGCCACTATG
AGAAGCTGAAGGGCTCCCCGAGGATAATGAGCAGAAACAGCTGTTTGTG
GAACAGCACAAGCACTACCTGGACGAGATCATCGAGCAGATCAGCGAGTT
CTCCAAGAGAGTGATCCTGGCCGACGCTAATCTGGACAAGTGCTGTCCG

-continued

CCTACAACAAGCACC CGGATAAGCCCATCAGAGAGCAGGCCGAGAATATC
ATCCACCTGTTTACCCTGACCAATCTGGGAGCCCTGCCGCCCTTCAAGTA
CTTTGACACCACCATCGACCGGAAGAGGTACACCAGCACCAAAGAGGTGC
TGGACGCCACCCTGATCCACCAGAGCATCACCGGCTGTACGAGACACGG
ATCGACCTGTCTCAGCTGGGAGGCGACAGCCCCAAGAAGAAGAGAAAGGT
GGAGGCCAGCTAACATATGATTCGAATGTCTTTCTTGGCTATGACACTT
CCAGCAAAAGGTAGGGCGGGCTGCAGAGCGGCTTCCCGGCGCTGCATGCA
ACACCGATGATGCTTCGACCCCCGAAGCTCCTTCGGGGCTGCATGGGCG
CTCCGATGCCGCTCCAGGGCGAGCGCTGTTTAAATAGCCAGGCCCCCGAT
TGCAAGACATTATAGCGAGCTACCAAAGCCATATTCAAACACCTAGATC
ACTACCCTTCTACACAGGCCACTCGAGCTGTGATCGCACTCCGCTAAG
GGGGCGCCTCTTCTCTCTCGTTTCAGTACAACC CGCAAAACATGACACAA
GAATCCCTGTACTTCTCGACCGTATTGATTCGGATGATTCTACGCGAG
CCTGCGGAACGACCAGGAATTCGGGAGGTGAGTGCAGAGCAAGCCCGG
CGGATCAGGCAGCGTGTTCGAGATTGACTTGC AACCCCGCATTGTGT
CGACGAAGGCTTTTGGCTCCTCTGTGCTGTCTCAAGCAGCATCTAACCC
TGCGTGC CGTTCATTTGCAGCCGCTGGCCCGCGAGCCCTGGAGGAG
CTCGGGCTGCCCGGTGCCCGGTGCTGCGGGTGC CCGCGAGAGCACCAA
CCCCGTACTGGTCCGCGAGCCCGGCCGGTGATCAAGCTGTTCCGGCGAGC
ACTGGTGC CGTCCGGAGAGCCTCGCGTCCGAGTCCGAGGCGTACGCGGTC
CTGGCGGACGCCCCGGTGC CGGTGCCCGCCTCCTCGCGCGCGGAGCT
GCGGCCCGGCACCGGAGCCTGGCCGTGGCCCTACCTGGTGATGAGCCGGA
TGACCGGCACCACCTGGCGGTCCGCGATGAGACGGCACGACCAGCCGAAAC
GCGCTGCTCGCCCTGGCCCGGAACTCGGCCGGGTGCTCGGCCGGCTGCA
CAGGGTGC CGCTGACCGGAAACCGTGCTCACCCCCATTCGAGGTCT
TCCCGGAACTGCTGCGGAAACCGCGCGGGGACCGCTCGAGGACACC CGC
GGGTGGGCTACCTCTCGCCCCGGTGTGGACCGCTGGAGGACTGGCT
GCCGGACGTGGACACGCTGCTGGCCGGCCGCGAAACCCCGTTCGTCCACG
GCGACCTGCACGGGACCAACATCTTCGTGGACCTGGCCGCGACCAGGTC
ACCGGGATCGTGC ACTTACCAGCTCTATGCGGAGACTCCCGCTACAG
CCTGGTGCAACTGCATCTCAACGCTTCCGGGGGACCCGCGAGATCCTGG
CCGCGTGTCTGACGGGGCGCAGTGAAGCGGACCGAGGACTTCGCCCCG
GAACTGCTCGCCTTACCTTCTGCACGACTTCGAGGTGTTTCGAGGAGAC
CCCCTGGATCTCTCCGGCTTACCAGATCCGGAGGAACTGGCGCAGTTCC
TCTGGGGCCCGCGGACACCGCCCCCGCGCTGATAAGGATCCGGCAAG
ACTGGCCCCGCTGGCAACGCAACAGTGAAGCCCCCTCCTAGTGTGTTGG
GGATGTGACTATGTATTCTGTGTGTTGGCCAACGGGTCAACCCGAACAGAT
TGATACCCGCTTGGCATTCTCTGTGAGAATGTAACGTCAGTTGATGGTA
CT

[0376] For all modified *Chlamydomonas reinhardtii* cells, Applicants used PCR, SURVEYOR nuclease assay, and DNA sequencing to verify successful modification.

Example 16

Use of Cas9 as a Transcriptional Repressor in Bacteria

[0377] The ability to artificially control transcription is essential both to the study of gene function and to the construction of synthetic gene networks with desired properties. Applicants describe here the use of the RNA-guided Cas9 protein as a programmable transcriptional repressor.

[0378] Applicants have previously demonstrated how the Cas9 protein of *Streptococcus pyogenes* SF370 can be used to direct genome editing in *Streptococcus pneumoniae*. In this study Applicants engineered the crR6Rk strain containing a minimal CRISPR system, consisting of cas9, the tracrRNA and a repeat. The D10A-H840 mutations were introduced into cas9 in this strain, giving strain crR6Rk**. Four spacers targeting different positions of the bgaA β -galactosidase gene promoter were cloned in the CRISPR array carried by the previously described pDB98 plasmid. Applicants observed a X to Y fold reduction in β -galactosidase activity depending on the targeted position, demonstrating the potential of Cas9 as a programmable repressor (FIG. 73).

[0379] To achieve Cas9** repression in *Escherichia coli* a green fluorescence protein (GFP) reporter plasmid (pDB127) was constructed to express the gfpmut2 gene from a constitutive promoter. The promoter was designed to carry several NPP PAMs on both strands, to measure the effect of Cas9** binding at various positions. Applicants introduced the D10A-H840 mutations into pCas9, a plasmid described carrying the tracrRNA, cas9 and a minimal CRISPR array designed for the easy cloning of new spacers. Twenty-two different spacers were designed to target different regions of the gfpmut2 promoter and open reading frame. An approximately 20-fold reduction of fluorescence of was observed upon targeting regions overlapping or adjacent to the -35 and -10 promoter elements and to the Shine-Dalgarno sequence. Targets on both strands showed similar repression levels. These results suggest that the binding of Cas9** to any position of the promoter region prevents transcription initiation, presumably through steric inhibition of RNAP binding.

[0380] To determine whether Cas9** could prevent transcription elongation, Applicants directed it to the reading frame of gfpmut2. A reduction in fluorescence was observed both when the coding and non-coding strands where targeted, suggesting that Cas9 binding is actually strong enough to represent an obstacle to the running RNAP. However, while a 40% reduction in expression was observed when the coding strand was the target, a 20-fold reduction was observed for the non-coding strand (FIG. 21b, compare T9, T10 and T11 to B9, B10 and B11). To directly determine the effects of Cas9** binding on transcription, Applicants extracted RNA from strains carrying either the T5, T10, B10 or a control construct that does not target pDB 127 and subjected it to Northern blot analysis using either a probe binding before (B477) or after (B510) the B10 and T10 target sites. Consistent with Applicants' fluorescence methods, no gfpmut2 transcription was detected when Cas9** was directed to the promoter region (T5 target) and a transcription was observed after the targeting of the T10

region. Interestingly, a smaller transcript was observed with the B477 probe. This band corresponds to the expected size of a transcript that would be interrupted by Cas9**, and is a direct indication of a transcriptional termination caused by dgRNA:Cas9** binding to the coding strand. Surprisingly, Applicants detected no transcript when the non-coding strand was targeted (B10). Since Cas9** binding to the B10 region is unlikely to interfere with transcription initiation, this result suggests that the mRNA was degraded. DgRNA:Cas9 was shown to bind ssRNA in vitro. Applicants speculate that binding may trigger degradation of the mRNA by host nucleases. Indeed, ribosome stalling can induce cleavage on the translated mRNA in *E. coli*.

[0381] Some applications require a precise tuning gene expression rather than its complete repression. Applicants sought to achieve intermediate repression levels through the introduction of mismatches that will weaken the crRNA/target interactions. Applicants created a series of spacers based on the B1, T5 and B10 constructs with increasing numbers of mutations in the 5' end of the crRNA. Up to 8 mutations in B1 and T5 did not affect the repression level, and a progressive increased in fluorescence was observed for additional mutations.

[0382] The observed repression with only an 8 nt match between the crRNA and its target raises the question of off-targeting effects of the use of Cas9** as a transcriptional regulator. Since a good PAM (NGG) is also required for Cas9 binding, the number of nucleotides to match to obtain some level of repression is 10. A 10 nt match occurs randomly once every ~1 Mbp, and such sites are thus likely to be found even in small bacterial genomes. However, to effectively repress transcription, such site needs to be in the promoter region of gene, which makes off-targeting much less likely. Applicants also showed that gene expression can be affected if the non-coding strand of a gene is targeted. For this to happen, a random target would have to be in the right orientation, but such events relatively more likely to happen. As a matter of fact, during the course of this study Applicants were unable to construct one of the designed spacer on pCas9**. Applicants later found this spacer showed a 12 bp match next to a good PAM in the essential murC gene. Such off-targeting could easily be avoided by a systematic blast of the designed spacers.

[0383] Aspects of the invention are further described in the following numbered paragraphs:

[0384] 1. A vector system comprising one or more vectors, wherein the system comprises

[0385] a. a first regulatory element operably linked to a traer mate sequence and one or more insertion sites for inserting a guide sequence upstream of the traer mate sequence, wherein when expressed, the guide sequence directs sequence-specific binding of a CRISPR complex to a target sequence in a eukaryotic cell, wherein the CRISPR complex comprises a CRISPR enzyme complexed with (1) the guide sequence that is hybridized to the target sequence, and (2) the traer mate sequence that is hybridized to the traer sequence; and

[0386] b. a second regulatory element operably linked to an enzyme-coding sequence encoding said CRISPR enzyme comprising a nuclear localization sequence;

[0387] wherein components (a) and (b) are located on the same or different vectors of the system.

[0388] 2. The vector system of paragraph 1, wherein component (a) further comprises the traer sequence downstream of the traer mate sequence under the control of the first regulatory element.

[0389] 3. The vector system of paragraph 1, wherein component (a) further comprises two or more guide sequences operably linked to the first regulatory element, wherein when expressed, each of the two or more guide sequences direct sequence specific binding of a CRISPR complex to a different target sequence in a eukaryotic cell.

[0390] 4. The vector system of paragraph 1, wherein the system comprises the traer sequence under the control of a third regulatory element.

[0391] 5. The vector system of paragraph 1, wherein the traer sequence exhibits at least 50% of sequence complementarity along the length of the traer mate sequence when optimally aligned.

[0392] 6. The vector system of paragraph 1, wherein the CRISPR enzyme comprises one or more nuclear localization sequences of sufficient strength to drive accumulation of said CRISPR enzyme in a detectable amount in the nucleus of a eukaryotic cell.

[0393] 7. The vector system of paragraph 1, wherein the CRISPR enzyme is a type II CRISPR system enzyme.

[0394] 8. The vector system of paragraph 1, wherein the CRISPR enzyme is a Cas9 enzyme.

[0395] 9. The vector system of paragraph 1, wherein the CRISPR enzyme is codon-optimized for expression in a eukaryotic cell.

[0396] 10. The vector system of paragraph 1, wherein the CRISPR enzyme directs cleavage of one or two strands at the location of the target sequence.

[0397] 11. The vector system of paragraph 1, wherein the CRISPR enzyme lacks DNA strand cleavage activity.

[0398] 12. The vector system of paragraph 1, wherein the first regulatory element is a polymerase III promoter.

[0399] 13. The vector system of paragraph 1, wherein the second regulatory element is a polymerase II promoter.

[0400] 14. The vector system of paragraph 4, wherein the third regulatory element is a polymerase III promoter.

[0401] 15. The vector system of paragraph 1, wherein the guide sequence is at least 15 nucleotides in length.

[0402] 16. The vector system of paragraph 1, wherein fewer than 50% of the nucleotides of the guide sequence participate in self-complementary base-pairing when optimally folded.

[0403] 17. A vector comprising a regulatory element operably linked to an enzyme-coding sequence encoding a CRISPR enzyme comprising one or more nuclear localization sequences, wherein said regulatory element drives transcription of the CRISPR enzyme in a eukaryotic cell such that said CRISPR enzyme accumulates in a detectable amount in the nucleus of the eukaryotic cell.

[0404] 18. The vector of paragraph 17, wherein said regulatory element is a polymerase II promoter.

[0405] 19. The vector of paragraph 17, wherein said CRISPR enzyme is a type IICRISPR system enzyme.

[0406] 20. The vector of paragraph 17, wherein said CRISPR enzyme is a Cas9 enzyme.

[0407] 21. The vector of paragraph 17, wherein said CRISPR enzyme lacks the ability to cleave one or more strands of a target sequence to which it binds.

[0408] 22. A CRISPR enzyme comprising one or more nuclear localization sequences of sufficient strength to drive

accumulation of said CRISPR enzyme in a detectable amount in the nucleus of a eukaryotic cell.

[0409] 23. The CRISPR enzyme of paragraph 22, wherein said CRISPR enzyme is a type IICRISPR system enzyme.

[0410] 24. The CRISPR enzyme of paragraph 22, wherein said CRISPR enzyme is a Cas9 enzyme.

[0411] 25. The CRISPR enzyme of paragraph 22, wherein said CRISPR enzyme lacks the ability to cleave one or more strands of a target sequence to which it binds.

[0412] 26. A eukaryotic host cell comprising:

[0413] a. a first regulatory element operably linked to a traer mate sequence and one or more insertion sites for inserting a guide sequence upstream of the traer mate sequence, wherein when expressed, the guide sequence directs sequence-specific binding of a CRISPR complex to a target sequence in a eukaryotic cell, wherein the CRISPR complex comprises a CRISPR enzyme complexed with (1) the guide sequence that is hybridized to the target sequence, and (2) the traer mate sequence that is hybridized to the traer sequence; and/or

[0414] b. a second regulatory element operably linked to an enzyme-coding sequence encoding said CRISPR enzyme comprising a nuclear localization sequence.

[0415] 27. The eukaryotic host cell of paragraph 26, wherein said host cell comprises components (a) and (b).

[0416] 28. The eukaryotic host cell of paragraph 26, wherein component (a), component (b), or components (a) and (b) are stably integrated into a genome of the host eukaryotic cell.

[0417] 29. The eukaryotic host cell of paragraph 26, wherein component (a) further comprises the traer sequence downstream of the traer mate sequence under the control of the first regulatory element.

[0418] 30. The eukaryotic host cell of paragraph 26, wherein component (a) further comprises two or more guide sequences operably linked to the first regulatory element, wherein when expressed, each of the two or more guide sequences direct sequence specific binding of a CRISPR complex to a different target sequence in a eukaryotic cell.

[0419] 31. The eukaryotic host cell of paragraph 26, further comprising a third regulatory element operably linked to said traer sequence.

[0420] 32. The eukaryotic host cell of paragraph 26, wherein the traer sequence exhibits at least 50% of sequence complementarity along the length of the traer mate sequence when optimally aligned.

[0421] 33. The eukaryotic host cell of paragraph 26, wherein the CRISPR enzyme comprises one or more nuclear localization sequences of sufficient strength to drive accumulation of said CRISPR enzyme in a detectable amount in the nucleus of a eukaryotic cell.

[0422] 34. The eukaryotic host cell of paragraph 26, wherein the CRISPR enzyme is a type II CRISPR system enzyme.

[0423] 35. The eukaryotic host cell of paragraph 26, wherein the CRISPR enzyme is a Cas9 enzyme.

[0424] 36. The eukaryotic host cell of paragraph 26, wherein the CRISPR enzyme is codon-optimized for expression in a eukaryotic cell.

[0425] 37. The eukaryotic host cell of paragraph 26, wherein the CRISPR enzyme directs cleavage of one or two strands at the location of the target sequence.

- [0426] 38. The eukaryotic host cell of paragraph 26, wherein the CRISPR enzyme lacks DNA strand cleavage activity.
- [0427] 39. The eukaryotic host cell of paragraph 26, wherein the first regulatory element is a polymerase III promoter.
- [0428] 40. The eukaryotic host cell of paragraph 26, wherein the second regulatory element is a polymerase II promoter.
- [0429] 41. The eukaryotic host cell of paragraph 31, wherein the third regulatory element is a polymerase III promoter.
- [0430] 42. The eukaryotic host cell of paragraph 26, wherein the guide sequence is at least 15 nucleotides in length.
- [0431] 43. The eukaryotic host cell of paragraph 26, wherein fewer than 50% of the nucleotides of the guide sequence participate in self-complementary base-pairing when optimally folded.
- [0432] 44. A non-human animal comprising a eukaryotic host cell of any one of paragraphs 26-43.
- [0433] 45. A kit comprising a vector system and instructions for using said kit, the vector system comprising:
- [0434] a. a first regulatory element operably linked to a traer mate sequence and one or more insertion sites for inserting a guide sequence upstream of the traer mate sequence, wherein when expressed, the guide sequence directs sequence-specific binding of a CRISPR complex to a target sequence in a eukaryotic cell, wherein the CRISPR complex comprises a CRISPR enzyme complexed with (1) the guide sequence that is hybridized to the target sequence, and (2) the traer mate sequence that is hybridized to the traer sequence; and/or
- [0435] b. a second regulatory element operably linked to an enzyme-coding sequence encoding said CRISPR enzyme comprising a nuclear localization sequence.
- [0436] 46. The kit of paragraph 45, wherein said kit comprises components (a) and (b) located on the same or different vectors of the system.
- [0437] 47. The kit of paragraph 45, wherein component (a) further comprises the traer sequence downstream of the traer mate sequence under the control of the first regulatory element.
- [0438] 48. The kit of paragraph 45, wherein component (a) further comprises two or more guide sequences operably linked to the first regulatory element, wherein when expressed, each of the two or more guide sequences direct sequence specific binding of a CRISPR complex to a different target sequence in a eukaryotic cell.
- [0439] 49. The kit of paragraph 45, wherein the system comprises the traer sequence under the control of a third regulatory element.
- [0440] 50. The kit of paragraph 45, wherein the traer sequence exhibits at least 50% of sequence complementarity along the length of the traer mate sequence when optimally aligned.
- [0441] 51. The kit of paragraph 45, wherein the CRISPR enzyme comprises one or more nuclear localization sequences of sufficient strength to drive accumulation of said CRISPR enzyme in a detectable amount in the nucleus of a eukaryotic cell.
- [0442] 52. The kit of paragraph 45, wherein the CRISPR enzyme is a type II CRISPR system enzyme.
- [0443] 53. The kit of paragraph 45, wherein the CRISPR enzyme is a Cas9 enzyme.
- [0444] 54. The kit of paragraph 45, wherein the CRISPR enzyme is codon-optimized for expression in a eukaryotic cell.
- [0445] 55. The kit of paragraph 45, wherein the CRISPR enzyme directs cleavage of one or two strands at the location of the target sequence.
- [0446] 56. The kit of paragraph 45, wherein the CRISPR enzyme lacks DNA strand cleavage activity.
- [0447] 57. The kit of paragraph 45, wherein the first regulatory element is a polymerase III promoter.
- [0448] 58. The kit of paragraph 45, wherein the second regulatory element is a polymerase II promoter.
- [0449] 59. The kit of paragraph 49, wherein the third regulatory element is a polymerase III promoter.
- [0450] 60. The kit of paragraph 45, wherein the guide sequence is at least 15 nucleotides in length.
- [0451] 61. The kit of paragraph 45, wherein fewer than 50% of the nucleotides of the guide sequence participate in self-complementary base-pairing when optimally folded.
- [0452] 62. A computer system for selecting a candidate target sequence within a nucleic acid sequence in a eukaryotic cell for targeting by a CRISPR complex, the system comprising:
- [0453] a. a memory unit configured to receive and/or store said nucleic acid sequence; and
- [0454] b. one or more processors alone or in combination programmed to (i) locate a CRISPR motif sequence within said nucleic acid sequence, and (ii) select a sequence adjacent to said located CRISPR motif sequence as the candidate target sequence to which the CRISPR complex binds.
- [0455] 63. The computer system of paragraph 62, wherein said locating step comprises identifying a CRISPR motif sequence located less than about 500 nucleotides away from said target sequence.
- [0456] 64. The computer system of paragraph 62, wherein said candidate target sequence is at least 10 nucleotides in length.
- [0457] 65. The computer system of paragraph 62, wherein the nucleotide at the 3' end of the candidate target sequence is located no more than about 10 nucleotides upstream of the CRISPR motif sequence.
- [0458] 66. The computer system of paragraph 62, wherein the nucleic acid sequence in the eukaryotic cell is endogenous to the eukaryotic genome.
- [0459] 67. The computer system of clam 62, wherein the nucleic acid sequence in the eukaryotic cell is exogenous to the eukaryotic genome.
- [0460] 68. A computer-readable medium comprising codes that, upon execution by one or more processors, implements a method of selecting a candidate target sequence within a nucleic acid sequence in a eukaryotic cell for targeting by a CRISPR complex, said method comprising: (a) locating a CRISPR motif sequence within said nucleic acid sequence, and (b) selecting a sequence adjacent to said located CRISPR motif sequence as the candidate target sequence to which the CRISPR complex binds.
- [0461] 69. The computer-readable medium of paragraph 68, wherein said locating comprises locating a CRISPR motif sequence that is less than about 500 nucleotides away from said target sequence.

[0462] 70. The computer-readable of paragraph 68, wherein said candidate target sequence is at least 10 nucleotides in length.

[0463] 71. The computer-readable of paragraph 68, wherein the nucleotide at the 3' end of the candidate target sequence is located no more than about 10 nucleotides upstream of the CRISPR motif sequence.

[0464] 72. The computer-readable of paragraph 68, wherein the nucleic acid sequence in the eukaryotic cell is endogenous the eukaryotic genome.

[0465] 73. The computer-readable of paragraph 68, wherein the nucleic acid sequence in the eukaryotic cell is exogenous to the eukaryotic genome.

[0466] 74. A method of modifying a target polynucleotide in a eukaryotic cell, the method comprising allowing a CRISPR complex to bind to the target polynucleotide to effect cleavage of said target polynucleotide thereby modifying the target polynucleotide, wherein the CRISPR complex comprises a CRISPR enzyme complexed with a guide sequence hybridized to a target sequence within said target polynucleotide, wherein said guide sequence is linked to a traer mate sequence which in tum hybridizes to a traer sequence.

[0467] 75. The method of paragraph 74, wherein said cleavage comprises cleaving one or two strands at the location of the target sequence by said CRISPR enzyme.

[0468] 76. The method of paragraph 74, wherein said cleavage results in decreased transcription of a target gene.

[0469] 77. The method of paragraph 74, further comprising repairing said cleaved target polynucleotide by homologous recombination with an exogenous template polynucleotide, wherein said repair results in a mutation comprising an insertion, deletion, or substitution of one or more nucleotides of said target polynucleotide.

[0470] 78. The method of paragraph 77, wherein said mutation results in one or more amino acid changes in a protein expressed from a gene comprising the target sequence.

[0471] 79. The method of paragraph 74, further comprising delivering one or more vectors to said eukaryotic cell, wherein the one or more vectors drive expression of one or more of: the CRISPR enzyme, the guide sequence linked to the traer mate sequence, and the traer sequence.

[0472] 80. The method of paragraph 79, wherein said vectors are delivered to the eukaryotic cell in a subject.

[0473] 81. The method of paragraph 74, wherein said modifying takes place in said eukaryotic cell in a cell culture.

[0474] 82. The method of paragraph 74, further comprising isolating said eukaryotic cell from a subject prior to said modifying.

[0475] 83. The method of paragraph 82, further comprising returning said eukaryotic cell and/or cells derived therefrom to said subject.

[0476] 84. A method of modifying expression of a polynucleotide in a eukaryotic cell, the method compnsmg: allowing a CRISPR complex to bind to the polynucleotide such that said binding results in increased or decreased expression of said polynucleotide; wherein the CRISPR complex comprises a CRISPR enzyme complexed with a guide sequence hybridized to a target sequence within said polynucleotide, wherein said guide sequence is linked to a traer mate sequence which in tum hybridizes to a traer sequence.

[0477] 85. The method of paragraph 74, further comprising delivering one or more vectors to said eukaryotic cells, wherein the one or more vectors drive expression of one or more of: the CRISPR enzyme, the guide sequence linked to the traer mate sequence, and the traer sequence.

[0478] 86. A method of generating a model eukaryotic cell comprising a mutated disease gene, the method comprising: [0479] a. introducing one or more vectors into a eukaryotic cell, wherein the one or more vectors drive expression of one or more of: a CRISPR enzyme, a guide sequence linked to a traer mate sequence, and a traer sequence; and [0480] b. allowing a CRISPR complex to bind to a target polynucleotide to effect cleavage of the target polynucleotide within said disease gene, wherein the CRISPR complex comprises the CRISPR enzyme complexed with (1) the guide sequence that is hybridized to the target sequence within the target polynucleotide, and (2) the traer mate sequence that is hybridized to the traer sequence, thereby generating a model eukaryotic cell comprising a mutated disease gene.

[0481] 87. The method of paragraph 86, wherein said cleavage comprises cleaving one or two strands at the location of the target sequence by said CRISPR enzyme.

[0482] 88. The method of paragraph 86, wherein said cleavage results in decreased transcription of a target gene.

[0483] 89. The method of paragraph 86, further comprising repairing said cleaved target polynucleotide by homologous recombination with an exogenous template polynucleotide, wherein said repair results in a mutation comprising an insertion, deletion, or substitution of one or more nucleotides of said target polynucleotide.

[0484] 90. The method of paragraph 89, wherein said mutation results in one or more amino acid changes in a protein expressed from a gene comprising the target sequence.

[0485] 91. A method of developing a biologically active agent that modulates a cell signaling event associated with a disease gene, comprising:

[0486] a. contacting a test compound with a model cell of any one of paragraphs 86-90; and

[0487] b. detecting a change in a readout that is indicative of a reduction or an augmentation of a cell signaling event associated with said mutation in said disease gene, thereby developing said biologically active agent that modulates said cell signaling event associated with said disease gene.

[0488] 92. A recombinant polynucleotide comprising a guide sequence upstream of a traer mate sequence, wherein the guide sequence when expressed directs sequence-specific binding of a CRISPR complex to a corresponding target sequence present in a eukaryotic cell.

[0489] 93. The recombinant polynucleotide of paragraph 89, wherein the target sequence is a viral sequence present in a eukaryotic cell.

[0490] 94. The recombinant polynucleotide of paragraph 89, wherein the target sequence is a proto-oncogene or an oncogene.

[0491] While preferred embodiments of the present invention have been shown and described herein, it will be obvious to those skilled in the art that such embodiments are provided by way of example only. Numerous variations, changes, and substitutions will now occur to those skilled in the art without departing from the invention. It should be understood that various alternatives to the embodiments of the invention described herein may be employed in practic-

ing the invention. It is intended that the following claims define the scope of the invention and that methods and structures within the scope of these claims and their equivalents be covered thereby.

REFERENCES

- [0492] 1. Urnov, F. D., Rebar, E. J., Holmes, M. C., Zhang, H. S. & Gregory, P. D. Genome editing with engineered zinc finger nucleases. *Nat. Rev. Genet.* 11, 636-646 (2010).
- [0493] 2. Bogdanove, A. J. & Voytas, D. F. TAL effectors: customizable proteins for DNA targeting. *Science* 333, 1843-1846 (2011).
- [0494] 3. Stoddard, B. L. Homing endonuclease structure and function. *Q. Rev. Biophys.* 38, 49-95 (2005).
- [0495] 4. Bae, T. & Schneewind, O. Allelic replacement in *Staphylococcus aureus* with inducible counter-selection. *Plasmid* 55, 58-63 (2006).
- [0496] 5. Sung, C. K., Li, H., Claverys, J. P. & Morrison, D. A. An rpsL cassette, janus, for gene replacement through negative selection in *Streptococcus pneumoniae*. *Appl. Environ. Microbiol.* 67, 5190-5196 (2001).
- [0497] 6. Sharan, S. K., Thomason, L. C., Kuznetsov, S. G. & Court, D. L. Recombineering: a homologous recombination-based method of genetic engineering. *Nat. Protoc.* 4, 206-223 (2009).
- [0498] 7. Jinek, M. et al. A programmable dual-RNA-guided DNA endonuclease in adaptive bacterial immunity. *Science* 337, 816-821 (2012).
- [0499] 8. Deveau, H., Garneau, J. E. & Moineau, S. CRISPR/Cas system and its role in phage-bacteria interactions. *Annu. Rev. Microbiol.* 64, 475-493 (2010).
- [0500] 9. Horvath, P. & Barrangou, R. CRISPR/Cas, the immune system of bacteria and archaea. *Science* 327, 167-170 (2010).
- [0501] 10. Terns, M. P. & Terns, R. M. CRISPR-based adaptive immune systems. *Curr. Opin. Microbiol.* 14, 321-327 (2011).
- [0502] 11. van der Oost, J., Jore, M. M., Westra, E. R., Lundgren, M. & Brouns, S. J. CRISPR-based adaptive and heritable immunity in prokaryotes. *Trends. Biochem. Sci.* 34, 401-407 (2009).
- [0503] 12. Brouns, S. J. et al. Small CRISPR RNAs guide antiviral defense in prokaryotes. *Science* 321, 960-964 (2008).
- [0504] 13. Carte, J., Wang, R., Li, H., Terns, R. M. & Terns, M. P. Cas6 is an endoribonuclease that generates guide RNAs for invader defense in prokaryotes. *Genes Dev.* 22, 3489-3496 (2008).
- [0505] 14. Deltcheva, E. et al. CRISPR RNA maturation by trans-encoded small RNA and host factor RNase III. *Nature* 471, 602-607 (2011).
- [0506] 15. Hatoum-Aslan, A., Maniv, I. & Marraffini, L. A. Mature clustered, regularly interspaced, short palindromic repeats RNA (crRNA) length is measured by a ruler mechanism anchored at the precursor processing site. *Proc. Natl. Acad. Sci. U.S.A.* 108, 21218-21222 (2011).
- [0507] 16. Haurwitz, R. E., Jinek, M., Wiedenheft, B., Zhou, K. & Doudna, J. A. Sequence- and structure-specific RNA processing by a CRISPR endonuclease. *Science* 329, 1355-1358 (2010).
- [0508] 17. Deveau, H. et al. Phage response to CRISPR-encoded resistance in *Streptococcus thermophilus*. *J. Bacteriol.* 190, 1390-1400 (2008).
- [0509] 18. Gasiunas, G., Barrangou, R., Horvath, P. & Siksnys, V. Cas9-crRNA ribonucleoprotein complex mediates specific DNA cleavage for adaptive immunity in bacteria. *Proc. Natl. Acad. Sci. U.S.A.* (2012).
- [0510] 19. Makarova, K. S., Aravind, L., Wolf, Y. I. & Koonin, E. V. Unification of Cas protein families and a simple scenario for the origin and evolution of CRISPR-Cas systems. *Biol. Direct.* 6, 38 (2011).
- [0511] 20. Barrangou, R. RNA-mediated programmable DNA cleavage. *Nat. Biotechnol.* 30, 836-838 (2012).
- [0512] 21. Brouns, S. J. Molecular biology. A Swiss army knife of immunity. *Science* 337, 808-809 (2012).
- [0513] 22. Carroll, D. A CRISPR Approach to Gene Targeting. *Mol. Ther.* 20, 1658-1660 (2012).
- [0514] 23. Bikard, D., Hatoum-Aslan, A., Mucida, D. & Marraffini, L. A. CRISPR interference can prevent natural transformation and virulence acquisition during in vivo bacterial infection. *Cell Host Microbe* 12, 177-186 (2012).
- [0515] 24. Sapranaukas, R. et al. The *Streptococcus thermophilus* CRISPR/Cas system provides immunity in *Escherichia coli*. *Nucleic Acids Res.* (2011).
- [0516] 25. Semenova, E. et al. Interference by clustered regularly interspaced short palindromic repeat (CRISPR) RNA is governed by a seed sequence. *Proc. Natl. Acad. Sci. U.S.A.* (2011).
- [0517] 26. Wiedenheft, B. et al. RNA-guided complex from a bacterial immune system enhances target recognition through seed sequence interactions. *Proc. Natl. Acad. Sci. U.S.A.* (2011).
- [0518] 27. Zahner, D. & Hakenbeck, R. The *Streptococcus pneumoniae* beta-galactosidase is a surface protein. *J. Bacteriol.* 182, 5919-5921 (2000).
- [0519] 28. Marraffini, L. A., Dedent, A. C. & Schneewind, O. Sortases and the art of anchoring proteins to the envelopes of gram-positive bacteria. *Microbiol. Mol. Biol. Rev.* 70, 192-221 (2006).
- [0520] 29. Motamedi, M. R., Szigety, S. K. & Rosenberg, S. M. Double-strand-break repair recombination in *Escherichia coli*: physical evidence for a DNA replication mechanism in vivo. *Genes Dev.* 13, 2889-2903 (1999).
- [0521] 30. Hosaka, T. et al. The novel mutation K87E in ribosomal protein S12 enhances protein synthesis activity during the late growth phase in *Escherichia coli*. *Mol. Genet. Genomics* 271, 317-324 (2004).
- [0522] 31. Costantino, N. & Court, D. L. Enhanced levels of lambda Red-mediated recombinants in mismatch repair mutants. *Proc. Natl. Acad. Sci. U.S.A.* 100, 15748-15753 (2003).
- [0523] 32. Edgar, R. & Qimron, U. The *Escherichia coli* CRISPR system protects from lambda lysogenization, lysogens, and prophage induction. *J. Bacteriol.* 192, 6291-6294 (2010).
- [0524] 33. Marraffini, L. A. & Sontheimer, E. J. Self versus non-self discrimination during CRISPR RNA-directed immunity. *Nature* 463, 568-571 (2010).
- [0525] 34. Fischer, S. et al. An archaeal immune system can detect multiple Protospacer Adjacent Motifs (PAMs) to target invader DNA. *J. Biol. Chem.* 287, 33351-33363 (2012).

- [0526] 35. Gudbergdottir, S. et al. Dynamic properties of the *Sulfolobus* CRISPR/Cas and CRISPR/Cmr systems when challenged with vector-borne viral and plasmid genes and protospacers. *Mol. Microbiol.* 79, 35-49 (2011).
- [0527] 36. Wang, H. H. et al. Genome-scale promoter engineering by coselection MAGE. *Nat Methods* 9, 591-593 (2012).
- [0528] 37. Cong, L. et al. Multiplex Genome Engineering Using CRISPR/Cas Systems. *Science* In press (2013).
- [0529] 38. Mali, P. et al. RNA-Guided Human Genome Engineering via Cas9. *Science* In press (2013).
- [0530] 39. Hoskins, J. et al. Genome of the bacterium *Streptococcus pneumoniae* strain R6. *J. Bacteriol.* 183, 5709-5717 (2001).
- [0531] 40. Havarstein, L. S., Coomaraswamy, G. & Morrison, D. A. An unmodified heptadecapeptide pheromone induces competence for genetic transformation in *Streptococcus pneumoniae*. *Proc. Natl. Acad. Sci. U.S.A.* 92, 11140-11144 (1995).
- [0532] 41. Horinouchi, S. & Weisblum, B. Nucleotide sequence and functional map of pC194, a plasmid that specifies inducible chloramphenicol resistance. *J. Bacteriol.* 150, 815-825 (1982).
- [0533] 42. Horton, R. M. In Vitro Recombination and Mutagenesis of DNA:SOEing Together Tailor-Made Genes. *Methods Mol. Biol.* 15, 251-261 (1993).
- [0534] 43. Podbielski, A., Spellerberg, B., Woischnik, M., Pohl, B. & Luttmann, R. Novel series of plasmid vectors for gene inactivation and expression analysis in group A streptococci (GAS). *Gene* 177, 137-147 (1996).
- [0535] 44. Husmann, L. K., Scott, J. R., Lindahl, G. & Stenberg, L. Expression of the Arp protein, a member of the M protein family, is not sufficient to inhibit phagocytosis of *Streptococcus pyogenes*. *Infection and immunity* 63, 345-348 (1995).
- [0536] 45. Gibson, D. G. et al. Enzymatic assembly of DNA molecules up to several hundred kilobases. *Nat Methods* 6, 343-345 (2009).

 SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 537

<210> SEQ ID NO 1
 <211> LENGTH: 7
 <212> TYPE: PRT
 <213> ORGANISM: Simian virus 40

<400> SEQUENCE: 1

Pro Lys Lys Lys Arg Lys Val
 1 5

<210> SEQ ID NO 2
 <211> LENGTH: 16
 <212> TYPE: PRT
 <213> ORGANISM: Unknown
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Unknown:
 Nucleoplasmin bipartite NLS sequence

<400> SEQUENCE: 2

Lys Arg Pro Ala Ala Thr Lys Lys Ala Gly Gln Ala Lys Lys Lys Lys
 1 5 10 15

<210> SEQ ID NO 3
 <211> LENGTH: 9
 <212> TYPE: PRT
 <213> ORGANISM: Unknown
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Unknown:
 C-myc NLS sequence

<400> SEQUENCE: 3

Pro Ala Ala Lys Arg Val Lys Leu Asp
 1 5

<210> SEQ ID NO 4
 <211> LENGTH: 11
 <212> TYPE: PRT
 <213> ORGANISM: Unknown
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Unknown:
 C-myc NLS sequence

<400> SEQUENCE: 4

-continued

Arg Gln Arg Arg Asn Glu Leu Lys Arg Ser Pro
1 5 10

<210> SEQ ID NO 5
<211> LENGTH: 38
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 5

Asn Gln Ser Ser Asn Phe Gly Pro Met Lys Gly Gly Asn Phe Gly Gly
1 5 10 15

Arg Ser Ser Gly Pro Tyr Gly Gly Gly Gly Gln Tyr Phe Ala Lys Pro
20 25 30

Arg Asn Gln Gly Gly Tyr
35

<210> SEQ ID NO 6
<211> LENGTH: 42
<212> TYPE: PRT
<213> ORGANISM: Unknown
<220> FEATURE:
<223> OTHER INFORMATION: Description of Unknown:
IBB domain from importin-alpha sequence

<400> SEQUENCE: 6

Arg Met Arg Ile Glx Phe Lys Asn Lys Gly Lys Asp Thr Ala Glu Leu
1 5 10 15

Arg Arg Arg Arg Val Glu Val Ser Val Glu Leu Arg Lys Ala Lys Lys
20 25 30

Asp Glu Gln Ile Leu Lys Arg Arg Asn Val
35 40

<210> SEQ ID NO 7
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Unknown
<220> FEATURE:
<223> OTHER INFORMATION: Description of Unknown:
Myoma T protein sequence

<400> SEQUENCE: 7

Val Ser Arg Lys Arg Pro Arg Pro
1 5

<210> SEQ ID NO 8
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Unknown
<220> FEATURE:
<223> OTHER INFORMATION: Description of Unknown:
Myoma T protein sequence

<400> SEQUENCE: 8

Pro Pro Lys Lys Ala Arg Glu Asp
1 5

<210> SEQ ID NO 9
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 9

-continued

Pro Gln Pro Lys Lys Lys Pro Leu
1 5

<210> SEQ ID NO 10
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 10

Ser Ala Leu Ile Lys Lys Lys Lys Lys Met Ala Pro
1 5 10

<210> SEQ ID NO 11
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: Influenza virus

<400> SEQUENCE: 11

Asp Arg Leu Arg Arg
1 5

<210> SEQ ID NO 12
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Influenza virus

<400> SEQUENCE: 12

Pro Lys Gln Lys Lys Arg Lys
1 5

<210> SEQ ID NO 13
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Hepatitis delta virus

<400> SEQUENCE: 13

Arg Lys Leu Lys Lys Lys Ile Lys Lys Leu
1 5 10

<210> SEQ ID NO 14
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 14

Arg Glu Lys Lys Lys Phe Leu Lys Arg Arg
1 5 10

<210> SEQ ID NO 15
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 15

Lys Arg Lys Gly Asp Glu Val Asp Gly Val Asp Glu Val Ala Lys Lys
1 5 10 15

Lys Ser Lys Lys
20

<210> SEQ ID NO 16
<211> LENGTH: 17

-continued

```

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      oligonucleotide
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (1)..(11)
<223> OTHER INFORMATION: a, c, t or g
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (12)..(13)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 20

nnnnnnnnnn nnnagaaw                               18

<210> SEQ ID NO 21
<211> LENGTH: 137
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      polynucleotide
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (1)..(20)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 21

nnnnnnnnnn nnnnnnnnnn gttttgtac tctcaagatt tagaaataaa tcttgcagaa    60
gctacaaaga taaggcttca tgccgaaatc aacaccctgt cattttatgg caggggtgtt    120
tcggtatttta attttttt                                     137

<210> SEQ ID NO 22
<211> LENGTH: 123
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      polynucleotide
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (1)..(20)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 22

nnnnnnnnnn nnnnnnnnnn gttttgtac tctcagaaat gcagaagcta caagataag    60
gcttcatgcc gaaatcaaca cctgtcatt ttatggcagg gtgttttcgt tatttaattt    120
ttt                                                         123

<210> SEQ ID NO 23
<211> LENGTH: 110
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      polynucleotide
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (1)..(20)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 23

nnnnnnnnnn nnnnnnnnnn gttttgtac tctcagaaat gcagaagcta caagataag    60

```

-continued

gcttcacgcc gaaatcaaca cctgtcatt ttatggcagg gtgttttttt 110

<210> SEQ ID NO 24
 <211> LENGTH: 102
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide
 <220> FEATURE:
 <221> NAME/KEY: modified_base
 <222> LOCATION: (1)..(20)
 <223> OTHER INFORMATION: a, c, t, g, unknown or other
 <400> SEQUENCE: 24

nnnnnnnnnn nnnnnnnnnn gttttagagc tagaaatagc aagttaaaat aaggctagtc 60

cgttatcaac ttgaaaaagt ggcaccgagt cggtgctttt tt 102

<210> SEQ ID NO 25
 <211> LENGTH: 88
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide
 <220> FEATURE:
 <221> NAME/KEY: modified_base
 <222> LOCATION: (1)..(20)
 <223> OTHER INFORMATION: a, c, t, g, unknown or other
 <400> SEQUENCE: 25

nnnnnnnnnn nnnnnnnnnn gttttagagc tagaaatagc aagttaaaat aaggctagtc 60

cgttatcaac ttgaaaaagt gttttttt 88

<210> SEQ ID NO 26
 <211> LENGTH: 76
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide
 <220> FEATURE:
 <221> NAME/KEY: modified_base
 <222> LOCATION: (1)..(20)
 <223> OTHER INFORMATION: a, c, t, g, unknown or other
 <400> SEQUENCE: 26

nnnnnnnnnn nnnnnnnnnn gttttagagc tagaaatagc aagttaaaat aaggctagtc 60

cgttatcatt tttttt 76

<210> SEQ ID NO 27
 <211> LENGTH: 12
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide
 <400> SEQUENCE: 27

gttttagagc ta 12

<210> SEQ ID NO 28
 <211> LENGTH: 31

-continued

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 28

tagcaagtta aaataaggct agtccgtttt t 31

<210> SEQ ID NO 29
<211> LENGTH: 27
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (1)..(22)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 29

nnnnnnnnnn nnnnnnnnnn nnagaaw 27

<210> SEQ ID NO 30
<211> LENGTH: 12
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 30

guuuuagagc ua 12

<210> SEQ ID NO 31
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 31

ggacatcgat gtcacctcca atgactaggg tgg 33

<210> SEQ ID NO 32
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 32

cattggaggt gacatcgatg tctctcccat tgg 33

<210> SEQ ID NO 33
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 33

ggaagggcct gagtccgagc agaagaagaa ggg 33

<210> SEQ ID NO 34
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

-continued

<400> SEQUENCE: 34

ggtggcgaga ggggccgaga ttgggtgttc agg 33

<210> SEQ ID NO 35

<211> LENGTH: 33

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 35

atgcaggagg gtggcgagag gggccgagat tgg 33

<210> SEQ ID NO 36

<211> LENGTH: 21

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 36

aaaaccaccc ttctctctgg c 21

<210> SEQ ID NO 37

<211> LENGTH: 21

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 37

ggagattgga gacacggaga g 21

<210> SEQ ID NO 38

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 38

ctggaagcc aatgcctgac 20

<210> SEQ ID NO 39

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 39

ggcagcaaac tccttgtct 20

<210> SEQ ID NO 40

<211> LENGTH: 335

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 40

-continued

```

gagggcctat ttcccatgat tccttcatat ttgcatatac gatacaaggc tgtagagag    60
ataattggaa ttaatttgac tgtaaacaca aagatattag tacaaaatac gtgacgtaga    120
aagtaataat ttcttgggta gtttgcagtt ttaaaattat gttttaaaat ggactatcat    180
atgcttaccg taacttgaaa gtatttcgat ttcttggcctt tatatatctt gtggaagga    240
cgaaacaccc gaaccattca aaacagcata gcaagttaaa ataaggctag tccggtatca    300
acttgaaaaa gtggcaccga gtcggtgctt ttttt                                335

```

```

<210> SEQ ID NO 41
<211> LENGTH: 423
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        polynucleotide

```

```

<400> SEQUENCE: 41

```

```

gagggcctat ttcccatgat tccttcatat ttgcatatac gatacaaggc tgtagagag    60
ataattggaa ttaatttgac tgtaaacaca aagatattag tacaaaatac gtgacgtaga    120
aagtaataat ttcttgggta gtttgcagtt ttaaaattat gttttaaaat ggactatcat    180
atgcttaccg taacttgaaa gtatttcgat ttcttggcctt tatatatctt gtggaagga    240
cgaaacaccc gtagtattaa gtattgtttt atggctgata aatttctttg aatttctcct    300
tgattatttg ttataaaagt tataaaataa tcttgttggga accattcaaa acagcatagc    360
aagttaaaat aaggctagtc cgttatcaac ttgaaaaagt ggcaccgagt cggtgctttt    420
ttt                                                                423

```

```

<210> SEQ ID NO 42
<211> LENGTH: 339
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        polynucleotide

```

```

<400> SEQUENCE: 42

```

```

gagggcctat ttcccatgat tccttcatat ttgcatatac gatacaaggc tgtagagag    60
ataattggaa ttaatttgac tgtaaacaca aagatattag tacaaaatac gtgacgtaga    120
aagtaataat ttcttgggta gtttgcagtt ttaaaattat gttttaaaat ggactatcat    180
atgcttaccg taacttgaaa gtatttcgat ttcttggcctt tatatatctt gtggaagga    240
cgaaacaccc ggttttagag ctatgctgtt ttgaatggtc ccaaaacggg tcttcgagaa    300
gacgttttag agctatgctg ttttgaatgg tcccaaac                                339

```

```

<210> SEQ ID NO 43
<211> LENGTH: 309
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        polynucleotide

```

```

<400> SEQUENCE: 43

```

```

gagggcctat ttcccatgat tccttcatat ttgcatatac gatacaaggc tgtagagag    60

```


-continued

```

ataattggaa ttaatttgac tgtaaacaca aagatattag tacaaaatac gtagcgtaga 120
aagtaataat ttcttgggta gtttgcagtt ttaaaattat gttttaaata ggactatcat 180
atgcttacgc taacttgaaa gtatttcgat ttcttggcctt tatatatcctt gtggaaagga 240
cgaaacacgc ggtcttcgag aagacctgtt ttagagctag aaatagcaag ttaaaataag 300
gctagtccg 309

```

```

<210> SEQ ID NO 44
<211> LENGTH: 1648
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        polypeptide

```

```

<400> SEQUENCE: 44

```

```

Met Asp Tyr Lys Asp His Asp Gly Asp Tyr Lys Asp His Asp Ile Asp
1          5          10         15
Tyr Lys Asp Asp Asp Lys Met Ala Pro Lys Lys Lys Arg Lys Val
20         25         30
Gly Ile His Gly Val Pro Ala Ala Asp Lys Lys Tyr Ser Ile Gly Leu
35         40         45
Asp Ile Gly Thr Asn Ser Val Gly Trp Ala Val Ile Thr Asp Glu Tyr
50         55         60
Lys Val Pro Ser Lys Lys Phe Lys Val Leu Gly Asn Thr Asp Arg His
65         70         75         80
Ser Ile Lys Lys Asn Leu Ile Gly Ala Leu Leu Phe Asp Ser Gly Glu
85         90         95
Thr Ala Glu Ala Thr Arg Leu Lys Arg Thr Ala Arg Arg Arg Tyr Thr
100        105        110
Arg Arg Lys Asn Arg Ile Cys Tyr Leu Gln Glu Ile Phe Ser Asn Glu
115        120        125
Met Ala Lys Val Asp Asp Ser Phe Phe His Arg Leu Glu Glu Ser Phe
130        135        140
Leu Val Glu Glu Asp Lys Lys His Glu Arg His Pro Ile Phe Gly Asn
145        150        155        160
Ile Val Asp Glu Val Ala Tyr His Glu Lys Tyr Pro Thr Ile Tyr His
165        170        175
Leu Arg Lys Lys Leu Val Asp Ser Thr Asp Lys Ala Asp Leu Arg Leu
180        185        190
Ile Tyr Leu Ala Leu Ala His Met Ile Lys Phe Arg Gly His Phe Leu
195        200        205
Ile Glu Gly Asp Leu Asn Pro Asp Asn Ser Asp Val Asp Lys Leu Phe
210        215        220
Ile Gln Leu Val Gln Thr Tyr Asn Gln Leu Phe Glu Glu Asn Pro Ile
225        230        235        240
Asn Ala Ser Gly Val Asp Ala Lys Ala Ile Leu Ser Ala Arg Leu Ser
245        250        255
Lys Ser Arg Arg Leu Glu Asn Leu Ile Ala Gln Leu Pro Gly Glu Lys
260        265        270
Lys Asn Gly Leu Phe Gly Asn Leu Ile Ala Leu Ser Leu Gly Leu Thr
275        280        285
Pro Asn Phe Lys Ser Asn Phe Asp Leu Ala Glu Asp Ala Lys Leu Gln

```


-continued

Leu Ile Asn Gly Ile Arg Asp Lys Gln Ser Gly Lys Thr Ile Leu Asp
 705 710 715 720
 Phe Leu Lys Ser Asp Gly Phe Ala Asn Arg Asn Phe Met Gln Leu Ile
 725 730 735
 His Asp Asp Ser Leu Thr Phe Lys Glu Asp Ile Gln Lys Ala Gln Val
 740 745 750
 Ser Gly Gln Gly Asp Ser Leu His Glu His Ile Ala Asn Leu Ala Gly
 755 760 765
 Ser Pro Ala Ile Lys Lys Gly Ile Leu Gln Thr Val Lys Val Val Asp
 770 775 780
 Glu Leu Val Lys Val Met Gly Arg His Lys Pro Glu Asn Ile Val Ile
 785 790 795 800
 Glu Met Ala Arg Glu Asn Gln Thr Thr Gln Lys Gly Gln Lys Asn Ser
 805 810 815
 Arg Glu Arg Met Lys Arg Ile Glu Glu Gly Ile Lys Glu Leu Gly Ser
 820 825 830
 Gln Ile Leu Lys Glu His Pro Val Glu Asn Thr Gln Leu Gln Asn Glu
 835 840 845
 Lys Leu Tyr Leu Tyr Tyr Leu Gln Asn Gly Arg Asp Met Tyr Val Asp
 850 855 860
 Gln Glu Leu Asp Ile Asn Arg Leu Ser Asp Tyr Asp Val Asp His Ile
 865 870 875 880
 Val Pro Gln Ser Phe Leu Lys Asp Asp Ser Ile Asp Asn Lys Val Leu
 885 890 895
 Thr Arg Ser Asp Lys Asn Arg Gly Lys Ser Asp Asn Val Pro Ser Glu
 900 905 910
 Glu Val Val Lys Lys Met Lys Asn Tyr Trp Arg Gln Leu Leu Asn Ala
 915 920 925
 Lys Leu Ile Thr Gln Arg Lys Phe Asp Asn Leu Thr Lys Ala Glu Arg
 930 935 940
 Gly Gly Leu Ser Glu Leu Asp Lys Ala Gly Phe Ile Lys Arg Gln Leu
 945 950 955 960
 Val Glu Thr Arg Gln Ile Thr Lys His Val Ala Gln Ile Leu Asp Ser
 965 970 975
 Arg Met Asn Thr Lys Tyr Asp Glu Asn Asp Lys Leu Ile Arg Glu Val
 980 985 990
 Lys Val Ile Thr Leu Lys Ser Lys Leu Val Ser Asp Phe Arg Lys Asp
 995 1000 1005
 Phe Gln Phe Tyr Lys Val Arg Glu Ile Asn Asn Tyr His His Ala
 1010 1015 1020
 His Asp Ala Tyr Leu Asn Ala Val Val Gly Thr Ala Leu Ile Lys
 1025 1030 1035
 Lys Tyr Pro Lys Leu Glu Ser Glu Phe Val Tyr Gly Asp Tyr Lys
 1040 1045 1050
 Val Tyr Asp Val Arg Lys Met Ile Ala Lys Ser Glu Gln Glu Ile
 1055 1060 1065
 Gly Lys Ala Thr Ala Lys Tyr Phe Phe Tyr Ser Asn Ile Met Asn
 1070 1075 1080
 Phe Phe Lys Thr Glu Ile Thr Leu Ala Asn Gly Glu Ile Arg Lys
 1085 1090 1095

-continued

Arg	Pro	Leu	Ile	Glu	Thr	Asn	Gly	Glu	Thr	Gly	Glu	Ile	Val	Trp
1100						1105					1110			
Asp	Lys	Gly	Arg	Asp	Phe	Ala	Thr	Val	Arg	Lys	Val	Leu	Ser	Met
1115						1120					1125			
Pro	Gln	Val	Asn	Ile	Val	Lys	Lys	Thr	Glu	Val	Gln	Thr	Gly	Gly
1130						1135					1140			
Phe	Ser	Lys	Glu	Ser	Ile	Leu	Pro	Lys	Arg	Asn	Ser	Asp	Lys	Leu
1145						1150					1155			
Ile	Ala	Arg	Lys	Lys	Asp	Trp	Asp	Pro	Lys	Lys	Tyr	Gly	Gly	Phe
1160						1165					1170			
Asp	Ser	Pro	Thr	Val	Ala	Tyr	Ser	Val	Leu	Val	Val	Ala	Lys	Val
1175						1180					1185			
Glu	Lys	Gly	Lys	Ser	Lys	Lys	Leu	Lys	Ser	Val	Lys	Glu	Leu	Leu
1190						1195					1200			
Gly	Ile	Thr	Ile	Met	Glu	Arg	Ser	Ser	Phe	Glu	Lys	Asn	Pro	Ile
1205						1210					1215			
Asp	Phe	Leu	Glu	Ala	Lys	Gly	Tyr	Lys	Glu	Val	Lys	Lys	Asp	Leu
1220						1225					1230			
Ile	Ile	Lys	Leu	Pro	Lys	Tyr	Ser	Leu	Phe	Glu	Leu	Glu	Asn	Gly
1235						1240					1245			
Arg	Lys	Arg	Met	Leu	Ala	Ser	Ala	Gly	Glu	Leu	Gln	Lys	Gly	Asn
1250						1255					1260			
Glu	Leu	Ala	Leu	Pro	Ser	Lys	Tyr	Val	Asn	Phe	Leu	Tyr	Leu	Ala
1265						1270					1275			
Ser	His	Tyr	Glu	Lys	Leu	Lys	Gly	Ser	Pro	Glu	Asp	Asn	Glu	Gln
1280						1285					1290			
Lys	Gln	Leu	Phe	Val	Glu	Gln	His	Lys	His	Tyr	Leu	Asp	Glu	Ile
1295						1300					1305			
Ile	Glu	Gln	Ile	Ser	Glu	Phe	Ser	Lys	Arg	Val	Ile	Leu	Ala	Asp
1310						1315					1320			
Ala	Asn	Leu	Asp	Lys	Val	Leu	Ser	Ala	Tyr	Asn	Lys	His	Arg	Asp
1325						1330					1335			
Lys	Pro	Ile	Arg	Glu	Gln	Ala	Glu	Asn	Ile	Ile	His	Leu	Phe	Thr
1340						1345					1350			
Leu	Thr	Asn	Leu	Gly	Ala	Pro	Ala	Ala	Phe	Lys	Tyr	Phe	Asp	Thr
1355						1360					1365			
Thr	Ile	Asp	Arg	Lys	Arg	Tyr	Thr	Ser	Thr	Lys	Glu	Val	Leu	Asp
1370						1375					1380			
Ala	Thr	Leu	Ile	His	Gln	Ser	Ile	Thr	Gly	Leu	Tyr	Glu	Thr	Arg
1385						1390					1395			
Ile	Asp	Leu	Ser	Gln	Leu	Gly	Gly	Asp	Ala	Ala	Ala	Val	Ser	Lys
1400						1405					1410			
Gly	Glu	Glu	Leu	Phe	Thr	Gly	Val	Val	Pro	Ile	Leu	Val	Glu	Leu
1415						1420					1425			
Asp	Gly	Asp	Val	Asn	Gly	His	Lys	Phe	Ser	Val	Ser	Gly	Glu	Gly
1430						1435					1440			
Glu	Gly	Asp	Ala	Thr	Tyr	Gly	Lys	Leu	Thr	Leu	Lys	Phe	Ile	Cys
1445						1450					1455			
Thr	Thr	Gly	Lys	Leu	Pro	Val	Pro	Trp	Pro	Thr	Leu	Val	Thr	Thr
1460						1465					1470			
Leu	Thr	Tyr	Gly	Val	Gln	Cys	Phe	Ser	Arg	Tyr	Pro	Asp	His	Met

-continued

1475	1480	1485
Lys Gln His Asp Phe Phe Lys Ser Ala Met Pro Glu Gly Tyr Val		
1490	1495	1500
Gln Glu Arg Thr Ile Phe Phe Lys Asp Asp Gly Asn Tyr Lys Thr		
1505	1510	1515
Arg Ala Glu Val Lys Phe Glu Gly Asp Thr Leu Val Asn Arg Ile		
1520	1525	1530
Glu Leu Lys Gly Ile Asp Phe Lys Glu Asp Gly Asn Ile Leu Gly		
1535	1540	1545
His Lys Leu Glu Tyr Asn Tyr Asn Ser His Asn Val Tyr Ile Met		
1550	1555	1560
Ala Asp Lys Gln Lys Asn Gly Ile Lys Val Asn Phe Lys Ile Arg		
1565	1570	1575
His Asn Ile Glu Asp Gly Ser Val Gln Leu Ala Asp His Tyr Gln		
1580	1585	1590
Gln Asn Thr Pro Ile Gly Asp Gly Pro Val Leu Leu Pro Asp Asn		
1595	1600	1605
His Tyr Leu Ser Thr Gln Ser Ala Leu Ser Lys Asp Pro Asn Glu		
1610	1615	1620
Lys Arg Asp His Met Val Leu Leu Glu Phe Val Thr Ala Ala Gly		
1625	1630	1635
Ile Thr Leu Gly Met Asp Glu Leu Tyr Lys		
1640	1645	

<210> SEQ ID NO 45

<211> LENGTH: 1625

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polypeptide

<400> SEQUENCE: 45

Met Asp Lys Lys Tyr Ser Ile Gly Leu Asp Ile Gly Thr Asn Ser Val			
1	5	10	15
Gly Trp Ala Val Ile Thr Asp Glu Tyr Lys Val Pro Ser Lys Lys Phe			
	20	25	30
Lys Val Leu Gly Asn Thr Asp Arg His Ser Ile Lys Lys Asn Leu Ile			
	35	40	45
Gly Ala Leu Leu Phe Asp Ser Gly Glu Thr Ala Glu Ala Thr Arg Leu			
	50	55	60
Lys Arg Thr Ala Arg Arg Arg Tyr Thr Arg Arg Lys Asn Arg Ile Cys			
65	70	75	80
Tyr Leu Gln Glu Ile Phe Ser Asn Glu Met Ala Lys Val Asp Asp Ser			
	85	90	95
Phe Phe His Arg Leu Glu Glu Ser Phe Leu Val Glu Glu Asp Lys Lys			
	100	105	110
His Glu Arg His Pro Ile Phe Gly Asn Ile Val Asp Glu Val Ala Tyr			
	115	120	125
His Glu Lys Tyr Pro Thr Ile Tyr His Leu Arg Lys Lys Leu Val Asp			
	130	135	140
Ser Thr Asp Lys Ala Asp Leu Arg Leu Ile Tyr Leu Ala Leu Ala His			
145	150	155	160

-continued

Met Ile Lys Phe Arg Gly His Phe Leu Ile Glu Gly Asp Leu Asn Pro
165 170 175

Asp Asn Ser Asp Val Asp Lys Leu Phe Ile Gln Leu Val Gln Thr Tyr
180 185 190

Asn Gln Leu Phe Glu Glu Asn Pro Ile Asn Ala Ser Gly Val Asp Ala
195 200 205

Lys Ala Ile Leu Ser Ala Arg Leu Ser Lys Ser Arg Arg Leu Glu Asn
210 215 220

Leu Ile Ala Gln Leu Pro Gly Glu Lys Lys Asn Gly Leu Phe Gly Asn
225 230 235 240

Leu Ile Ala Leu Ser Leu Gly Leu Thr Pro Asn Phe Lys Ser Asn Phe
245 250 255

Asp Leu Ala Glu Asp Ala Lys Leu Gln Leu Ser Lys Asp Thr Tyr Asp
260 265 270

Asp Asp Leu Asp Asn Leu Leu Ala Gln Ile Gly Asp Gln Tyr Ala Asp
275 280 285

Leu Phe Leu Ala Ala Lys Asn Leu Ser Asp Ala Ile Leu Leu Ser Asp
290 295 300

Ile Leu Arg Val Asn Thr Glu Ile Thr Lys Ala Pro Leu Ser Ala Ser
305 310 315 320

Met Ile Lys Arg Tyr Asp Glu His His Gln Asp Leu Thr Leu Leu Lys
325 330 335

Ala Leu Val Arg Gln Gln Leu Pro Glu Lys Tyr Lys Glu Ile Phe Phe
340 345 350

Asp Gln Ser Lys Asn Gly Tyr Ala Gly Tyr Ile Asp Gly Gly Ala Ser
355 360 365

Gln Glu Glu Phe Tyr Lys Phe Ile Lys Pro Ile Leu Glu Lys Met Asp
370 375 380

Gly Thr Glu Glu Leu Leu Val Lys Leu Asn Arg Glu Asp Leu Leu Arg
385 390 395 400

Lys Gln Arg Thr Phe Asp Asn Gly Ser Ile Pro His Gln Ile His Leu
405 410 415

Gly Glu Leu His Ala Ile Leu Arg Arg Gln Glu Asp Phe Tyr Pro Phe
420 425 430

Leu Lys Asp Asn Arg Glu Lys Ile Glu Lys Ile Leu Thr Phe Arg Ile
435 440 445

Pro Tyr Tyr Val Gly Pro Leu Ala Arg Gly Asn Ser Arg Phe Ala Trp
450 455 460

Met Thr Arg Lys Ser Glu Glu Thr Ile Thr Pro Trp Asn Phe Glu Glu
465 470 475 480

Val Val Asp Lys Gly Ala Ser Ala Gln Ser Phe Ile Glu Arg Met Thr
485 490 495

Asn Phe Asp Lys Asn Leu Pro Asn Glu Lys Val Leu Pro Lys His Ser
500 505 510

Leu Leu Tyr Glu Tyr Phe Thr Val Tyr Asn Glu Leu Thr Lys Val Lys
515 520 525

Tyr Val Thr Glu Gly Met Arg Lys Pro Ala Phe Leu Ser Gly Glu Gln
530 535 540

Lys Lys Ala Ile Val Asp Leu Leu Phe Lys Thr Asn Arg Lys Val Thr
545 550 555 560

Val Lys Gln Leu Lys Glu Asp Tyr Phe Lys Lys Ile Glu Cys Phe Asp

-continued

565				570				575							
Ser	Val	Glu	Ile	Ser	Gly	Val	Glu	Asp	Arg	Phe	Asn	Ala	Ser	Leu	Gly
		580						585					590		
Thr	Tyr	His	Asp	Leu	Leu	Lys	Ile	Ile	Lys	Asp	Lys	Asp	Phe	Leu	Asp
		595					600					605			
Asn	Glu	Glu	Asn	Glu	Asp	Ile	Leu	Glu	Asp	Ile	Val	Leu	Thr	Leu	Thr
	610					615					620				
Leu	Phe	Glu	Asp	Arg	Glu	Met	Ile	Glu	Glu	Arg	Leu	Lys	Thr	Tyr	Ala
	625				630					635					640
His	Leu	Phe	Asp	Asp	Lys	Val	Met	Lys	Gln	Leu	Lys	Arg	Arg	Arg	Tyr
			645						650					655	
Thr	Gly	Trp	Gly	Arg	Leu	Ser	Arg	Lys	Leu	Ile	Asn	Gly	Ile	Arg	Asp
			660						665				670		
Lys	Gln	Ser	Gly	Lys	Thr	Ile	Leu	Asp	Phe	Leu	Lys	Ser	Asp	Gly	Phe
		675					680					685			
Ala	Asn	Arg	Asn	Phe	Met	Gln	Leu	Ile	His	Asp	Asp	Ser	Leu	Thr	Phe
	690					695					700				
Lys	Glu	Asp	Ile	Gln	Lys	Ala	Gln	Val	Ser	Gly	Gln	Gly	Asp	Ser	Leu
	705				710					715					720
His	Glu	His	Ile	Ala	Asn	Leu	Ala	Gly	Ser	Pro	Ala	Ile	Lys	Lys	Gly
			725						730					735	
Ile	Leu	Gln	Thr	Val	Lys	Val	Val	Asp	Glu	Leu	Val	Lys	Val	Met	Gly
			740					745					750		
Arg	His	Lys	Pro	Glu	Asn	Ile	Val	Ile	Glu	Met	Ala	Arg	Glu	Asn	Gln
		755					760					765			
Thr	Thr	Gln	Lys	Gly	Gln	Lys	Asn	Ser	Arg	Glu	Arg	Met	Lys	Arg	Ile
	770					775					780				
Glu	Glu	Gly	Ile	Lys	Glu	Leu	Gly	Ser	Gln	Ile	Leu	Lys	Glu	His	Pro
	785				790					795					800
Val	Glu	Asn	Thr	Gln	Leu	Gln	Asn	Glu	Lys	Leu	Tyr	Leu	Tyr	Tyr	Leu
			805						810					815	
Gln	Asn	Gly	Arg	Asp	Met	Tyr	Val	Asp	Gln	Glu	Leu	Asp	Ile	Asn	Arg
			820					825					830		
Leu	Ser	Asp	Tyr	Asp	Val	Asp	His	Ile	Val	Pro	Gln	Ser	Phe	Leu	Lys
		835					840					845			
Asp	Asp	Ser	Ile	Asp	Asn	Lys	Val	Leu	Thr	Arg	Ser	Asp	Lys	Asn	Arg
	850					855					860				
Gly	Lys	Ser	Asp	Asn	Val	Pro	Ser	Glu	Glu	Val	Val	Lys	Lys	Met	Lys
	865				870					875				880	
Asn	Tyr	Trp	Arg	Gln	Leu	Leu	Asn	Ala	Lys	Leu	Ile	Thr	Gln	Arg	Lys
			885						890					895	
Phe	Asp	Asn	Leu	Thr	Lys	Ala	Glu	Arg	Gly	Gly	Leu	Ser	Glu	Leu	Asp
			900					905					910		
Lys	Ala	Gly	Phe	Ile	Lys	Arg	Gln	Leu	Val	Glu	Thr	Arg	Gln	Ile	Thr
		915					920					925			
Lys	His	Val	Ala	Gln	Ile	Leu	Asp	Ser	Arg	Met	Asn	Thr	Lys	Tyr	Asp
	930					935					940				
Glu	Asn	Asp	Lys	Leu	Ile	Arg	Glu	Val	Lys	Val	Ile	Thr	Leu	Lys	Ser
	945				950					955					960
Lys	Leu	Val	Ser	Asp	Phe	Arg	Lys	Asp	Phe	Gln	Phe	Tyr	Lys	Val	Arg
			965						970					975	

-continued

Glu Ile Asn Asn Tyr His His Ala His Asp Ala Tyr Leu Asn Ala Val
 980 985 990

Val Gly Thr Ala Leu Ile Lys Lys Tyr Pro Lys Leu Glu Ser Glu Phe
 995 1000 1005

Val Tyr Gly Asp Tyr Lys Val Tyr Asp Val Arg Lys Met Ile Ala
 1010 1015 1020

Lys Ser Glu Gln Glu Ile Gly Lys Ala Thr Ala Lys Tyr Phe Phe
 1025 1030 1035

Tyr Ser Asn Ile Met Asn Phe Phe Lys Thr Glu Ile Thr Leu Ala
 1040 1045 1050

Asn Gly Glu Ile Arg Lys Arg Pro Leu Ile Glu Thr Asn Gly Glu
 1055 1060 1065

Thr Gly Glu Ile Val Trp Asp Lys Gly Arg Asp Phe Ala Thr Val
 1070 1075 1080

Arg Lys Val Leu Ser Met Pro Gln Val Asn Ile Val Lys Lys Thr
 1085 1090 1095

Glu Val Gln Thr Gly Gly Phe Ser Lys Glu Ser Ile Leu Pro Lys
 1100 1105 1110

Arg Asn Ser Asp Lys Leu Ile Ala Arg Lys Lys Asp Trp Asp Pro
 1115 1120 1125

Lys Lys Tyr Gly Gly Phe Asp Ser Pro Thr Val Ala Tyr Ser Val
 1130 1135 1140

Leu Val Val Ala Lys Val Glu Lys Gly Lys Ser Lys Lys Leu Lys
 1145 1150 1155

Ser Val Lys Glu Leu Leu Gly Ile Thr Ile Met Glu Arg Ser Ser
 1160 1165 1170

Phe Glu Lys Asn Pro Ile Asp Phe Leu Glu Ala Lys Gly Tyr Lys
 1175 1180 1185

Glu Val Lys Lys Asp Leu Ile Ile Lys Leu Pro Lys Tyr Ser Leu
 1190 1195 1200

Phe Glu Leu Glu Asn Gly Arg Lys Arg Met Leu Ala Ser Ala Gly
 1205 1210 1215

Glu Leu Gln Lys Gly Asn Glu Leu Ala Leu Pro Ser Lys Tyr Val
 1220 1225 1230

Asn Phe Leu Tyr Leu Ala Ser His Tyr Glu Lys Leu Lys Gly Ser
 1235 1240 1245

Pro Glu Asp Asn Glu Gln Lys Gln Leu Phe Val Glu Gln His Lys
 1250 1255 1260

His Tyr Leu Asp Glu Ile Ile Glu Gln Ile Ser Glu Phe Ser Lys
 1265 1270 1275

Arg Val Ile Leu Ala Asp Ala Asn Leu Asp Lys Val Leu Ser Ala
 1280 1285 1290

Tyr Asn Lys His Arg Asp Lys Pro Ile Arg Glu Gln Ala Glu Asn
 1295 1300 1305

Ile Ile His Leu Phe Thr Leu Thr Asn Leu Gly Ala Pro Ala Ala
 1310 1315 1320

Phe Lys Tyr Phe Asp Thr Thr Ile Asp Arg Lys Arg Tyr Thr Ser
 1325 1330 1335

Thr Lys Glu Val Leu Asp Ala Thr Leu Ile His Gln Ser Ile Thr
 1340 1345 1350

-continued

Gly Leu Tyr Glu Thr Arg Ile Asp Leu Ser Gln Leu Gly Gly Asp
 1355 1360 1365

Ala Ala Ala Val Ser Lys Gly Glu Glu Leu Phe Thr Gly Val Val
 1370 1375 1380

Pro Ile Leu Val Glu Leu Asp Gly Asp Val Asn Gly His Lys Phe
 1385 1390 1395

Ser Val Ser Gly Glu Gly Glu Gly Asp Ala Thr Tyr Gly Lys Leu
 1400 1405 1410

Thr Leu Lys Phe Ile Cys Thr Thr Gly Lys Leu Pro Val Pro Trp
 1415 1420 1425

Pro Thr Leu Val Thr Thr Leu Thr Tyr Gly Val Gln Cys Phe Ser
 1430 1435 1440

Arg Tyr Pro Asp His Met Lys Gln His Asp Phe Phe Lys Ser Ala
 1445 1450 1455

Met Pro Glu Gly Tyr Val Gln Glu Arg Thr Ile Phe Phe Lys Asp
 1460 1465 1470

Asp Gly Asn Tyr Lys Thr Arg Ala Glu Val Lys Phe Glu Gly Asp
 1475 1480 1485

Thr Leu Val Asn Arg Ile Glu Leu Lys Gly Ile Asp Phe Lys Glu
 1490 1495 1500

Asp Gly Asn Ile Leu Gly His Lys Leu Glu Tyr Asn Tyr Asn Ser
 1505 1510 1515

His Asn Val Tyr Ile Met Ala Asp Lys Gln Lys Asn Gly Ile Lys
 1520 1525 1530

Val Asn Phe Lys Ile Arg His Asn Ile Glu Asp Gly Ser Val Gln
 1535 1540 1545

Leu Ala Asp His Tyr Gln Gln Asn Thr Pro Ile Gly Asp Gly Pro
 1550 1555 1560

Val Leu Leu Pro Asp Asn His Tyr Leu Ser Thr Gln Ser Ala Leu
 1565 1570 1575

Ser Lys Asp Pro Asn Glu Lys Arg Asp His Met Val Leu Leu Glu
 1580 1585 1590

Phe Val Thr Ala Ala Gly Ile Thr Leu Gly Met Asp Glu Leu Tyr
 1595 1600 1605

Lys Lys Arg Pro Ala Ala Thr Lys Lys Ala Gly Gln Ala Lys Lys
 1610 1615 1620

Lys Lys
 1625

<210> SEQ ID NO 46
 <211> LENGTH: 1664
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 polypeptide

<400> SEQUENCE: 46

Met Asp Tyr Lys Asp His Asp Gly Asp Tyr Lys Asp His Asp Ile Asp
 1 5 10 15

Tyr Lys Asp Asp Asp Asp Lys Met Ala Pro Lys Lys Lys Arg Lys Val
 20 25 30

Gly Ile His Gly Val Pro Ala Ala Asp Lys Lys Tyr Ser Ile Gly Leu
 35 40 45

-continued

Asp Ile Gly Thr Asn Ser Val Gly Trp Ala Val Ile Thr Asp Glu Tyr
 50 55 60
 Lys Val Pro Ser Lys Lys Phe Lys Val Leu Gly Asn Thr Asp Arg His
 65 70 75 80
 Ser Ile Lys Lys Asn Leu Ile Gly Ala Leu Leu Phe Asp Ser Gly Glu
 85 90 95
 Thr Ala Glu Ala Thr Arg Leu Lys Arg Thr Ala Arg Arg Arg Tyr Thr
 100 105 110
 Arg Arg Lys Asn Arg Ile Cys Tyr Leu Gln Glu Ile Phe Ser Asn Glu
 115 120 125
 Met Ala Lys Val Asp Asp Ser Phe Phe His Arg Leu Glu Glu Ser Phe
 130 135 140
 Leu Val Glu Glu Asp Lys Lys His Glu Arg His Pro Ile Phe Gly Asn
 145 150 155 160
 Ile Val Asp Glu Val Ala Tyr His Glu Lys Tyr Pro Thr Ile Tyr His
 165 170 175
 Leu Arg Lys Lys Leu Val Asp Ser Thr Asp Lys Ala Asp Leu Arg Leu
 180 185 190
 Ile Tyr Leu Ala Leu Ala His Met Ile Lys Phe Arg Gly His Phe Leu
 195 200 205
 Ile Glu Gly Asp Leu Asn Pro Asp Asn Ser Asp Val Asp Lys Leu Phe
 210 215 220
 Ile Gln Leu Val Gln Thr Tyr Asn Gln Leu Phe Glu Glu Asn Pro Ile
 225 230 235 240
 Asn Ala Ser Gly Val Asp Ala Lys Ala Ile Leu Ser Ala Arg Leu Ser
 245 250 255
 Lys Ser Arg Arg Leu Glu Asn Leu Ile Ala Gln Leu Pro Gly Glu Lys
 260 265 270
 Lys Asn Gly Leu Phe Gly Asn Leu Ile Ala Leu Ser Leu Gly Leu Thr
 275 280 285
 Pro Asn Phe Lys Ser Asn Phe Asp Leu Ala Glu Asp Ala Lys Leu Gln
 290 295 300
 Leu Ser Lys Asp Thr Tyr Asp Asp Asp Leu Asp Asn Leu Leu Ala Gln
 305 310 315 320
 Ile Gly Asp Gln Tyr Ala Asp Leu Phe Leu Ala Ala Lys Asn Leu Ser
 325 330 335
 Asp Ala Ile Leu Leu Ser Asp Ile Leu Arg Val Asn Thr Glu Ile Thr
 340 345 350
 Lys Ala Pro Leu Ser Ala Ser Met Ile Lys Arg Tyr Asp Glu His His
 355 360 365
 Gln Asp Leu Thr Leu Leu Lys Ala Leu Val Arg Gln Gln Leu Pro Glu
 370 375 380
 Lys Tyr Lys Glu Ile Phe Phe Asp Gln Ser Lys Asn Gly Tyr Ala Gly
 385 390 395 400
 Tyr Ile Asp Gly Gly Ala Ser Gln Glu Glu Phe Tyr Lys Phe Ile Lys
 405 410 415
 Pro Ile Leu Glu Lys Met Asp Gly Thr Glu Glu Leu Leu Val Lys Leu
 420 425 430
 Asn Arg Glu Asp Leu Leu Arg Lys Gln Arg Thr Phe Asp Asn Gly Ser
 435 440 445

-continued

Ile	Pro	His	Gln	Ile	His	Leu	Gly	Glu	Leu	His	Ala	Ile	Leu	Arg	Arg
450						455					460				
Gln	Glu	Asp	Phe	Tyr	Pro	Phe	Leu	Lys	Asp	Asn	Arg	Glu	Lys	Ile	Glu
465					470					475					480
Lys	Ile	Leu	Thr	Phe	Arg	Ile	Pro	Tyr	Tyr	Val	Gly	Pro	Leu	Ala	Arg
				485					490						495
Gly	Asn	Ser	Arg	Phe	Ala	Trp	Met	Thr	Arg	Lys	Ser	Glu	Glu	Thr	Ile
			500					505					510		
Thr	Pro	Trp	Asn	Phe	Glu	Glu	Val	Val	Asp	Lys	Gly	Ala	Ser	Ala	Gln
		515					520					525			
Ser	Phe	Ile	Glu	Arg	Met	Thr	Asn	Phe	Asp	Lys	Asn	Leu	Pro	Asn	Glu
530						535					540				
Lys	Val	Leu	Pro	Lys	His	Ser	Leu	Leu	Tyr	Glu	Tyr	Phe	Thr	Val	Tyr
545					550					555					560
Asn	Glu	Leu	Thr	Lys	Val	Lys	Tyr	Val	Thr	Glu	Gly	Met	Arg	Lys	Pro
				565					570					575	
Ala	Phe	Leu	Ser	Gly	Glu	Gln	Lys	Lys	Ala	Ile	Val	Asp	Leu	Leu	Phe
			580					585					590		
Lys	Thr	Asn	Arg	Lys	Val	Thr	Val	Lys	Gln	Leu	Lys	Glu	Asp	Tyr	Phe
		595					600					605			
Lys	Lys	Ile	Glu	Cys	Phe	Asp	Ser	Val	Glu	Ile	Ser	Gly	Val	Glu	Asp
610						615					620				
Arg	Phe	Asn	Ala	Ser	Leu	Gly	Thr	Tyr	His	Asp	Leu	Leu	Lys	Ile	Ile
625					630					635					640
Lys	Asp	Lys	Asp	Phe	Leu	Asp	Asn	Glu	Glu	Asn	Glu	Asp	Ile	Leu	Glu
				645					650					655	
Asp	Ile	Val	Leu	Thr	Leu	Thr	Leu	Phe	Glu	Asp	Arg	Glu	Met	Ile	Glu
			660					665					670		
Glu	Arg	Leu	Lys	Thr	Tyr	Ala	His	Leu	Phe	Asp	Asp	Lys	Val	Met	Lys
		675					680					685			
Gln	Leu	Lys	Arg	Arg	Arg	Tyr	Thr	Gly	Trp	Gly	Arg	Leu	Ser	Arg	Lys
690						695					700				
Leu	Ile	Asn	Gly	Ile	Arg	Asp	Lys	Gln	Ser	Gly	Lys	Thr	Ile	Leu	Asp
705					710					715					720
Phe	Leu	Lys	Ser	Asp	Gly	Phe	Ala	Asn	Arg	Asn	Phe	Met	Gln	Leu	Ile
				725					730					735	
His	Asp	Asp	Ser	Leu	Thr	Phe	Lys	Glu	Asp	Ile	Gln	Lys	Ala	Gln	Val
			740					745					750		
Ser	Gly	Gln	Gly	Asp	Ser	Leu	His	Glu	His	Ile	Ala	Asn	Leu	Ala	Gly
		755					760					765			
Ser	Pro	Ala	Ile	Lys	Lys	Gly	Ile	Leu	Gln	Thr	Val	Lys	Val	Val	Asp
	770					775					780				
Glu	Leu	Val	Lys	Val	Met	Gly	Arg	His	Lys	Pro	Glu	Asn	Ile	Val	Ile
785					790					795					800
Glu	Met	Ala	Arg	Glu	Asn	Gln	Thr	Thr	Gln	Lys	Gly	Gln	Lys	Asn	Ser
				805					810					815	
Arg	Glu	Arg	Met	Lys	Arg	Ile	Glu	Glu	Gly	Ile	Lys	Glu	Leu	Gly	Ser
			820					825					830		
Gln	Ile	Leu	Lys	Glu	His	Pro	Val	Glu	Asn	Thr	Gln	Leu	Gln	Asn	Glu
		835					840					845			
Lys	Leu	Tyr	Leu	Tyr	Tyr	Leu	Gln	Asn	Gly	Arg	Asp	Met	Tyr	Val	Asp

-continued

850			855			860									
Gln	Glu	Leu	Asp	Ile	Asn	Arg	Leu	Ser	Asp	Tyr	Asp	Val	Asp	His	Ile
865					870					875					880
Val	Pro	Gln	Ser	Phe	Leu	Lys	Asp	Asp	Ser	Ile	Asp	Asn	Lys	Val	Leu
				885					890						895
Thr	Arg	Ser	Asp	Lys	Asn	Arg	Gly	Lys	Ser	Asp	Asn	Val	Pro	Ser	Glu
			900					905						910	
Glu	Val	Val	Lys	Lys	Met	Lys	Asn	Tyr	Trp	Arg	Gln	Leu	Leu	Asn	Ala
		915					920							925	
Lys	Leu	Ile	Thr	Gln	Arg	Lys	Phe	Asp	Asn	Leu	Thr	Lys	Ala	Glu	Arg
930					935						940				
Gly	Gly	Leu	Ser	Glu	Leu	Asp	Lys	Ala	Gly	Phe	Ile	Lys	Arg	Gln	Leu
945					950					955					960
Val	Glu	Thr	Arg	Gln	Ile	Thr	Lys	His	Val	Ala	Gln	Ile	Leu	Asp	Ser
				965					970						975
Arg	Met	Asn	Thr	Lys	Tyr	Asp	Glu	Asn	Asp	Lys	Leu	Ile	Arg	Glu	Val
			980					985						990	
Lys	Val	Ile	Thr	Leu	Lys	Ser	Lys	Leu	Val	Ser	Asp	Phe	Arg	Lys	Asp
		995					1000							1005	
Phe	Gln	Phe	Tyr	Lys	Val	Arg	Glu	Ile	Asn	Asn	Tyr	His	His	Ala	
	1010					1015						1020			
His	Asp	Ala	Tyr	Leu	Asn	Ala	Val	Val	Gly	Thr	Ala	Leu	Ile	Lys	
	1025					1030						1035			
Lys	Tyr	Pro	Lys	Leu	Glu	Ser	Glu	Phe	Val	Tyr	Gly	Asp	Tyr	Lys	
	1040					1045						1050			
Val	Tyr	Asp	Val	Arg	Lys	Met	Ile	Ala	Lys	Ser	Glu	Gln	Glu	Ile	
	1055					1060						1065			
Gly	Lys	Ala	Thr	Ala	Lys	Tyr	Phe	Phe	Tyr	Ser	Asn	Ile	Met	Asn	
	1070					1075						1080			
Phe	Phe	Lys	Thr	Glu	Ile	Thr	Leu	Ala	Asn	Gly	Glu	Ile	Arg	Lys	
	1085					1090						1095			
Arg	Pro	Leu	Ile	Glu	Thr	Asn	Gly	Glu	Thr	Gly	Glu	Ile	Val	Trp	
	1100					1105						1110			
Asp	Lys	Gly	Arg	Asp	Phe	Ala	Thr	Val	Arg	Lys	Val	Leu	Ser	Met	
	1115					1120						1125			
Pro	Gln	Val	Asn	Ile	Val	Lys	Lys	Thr	Glu	Val	Gln	Thr	Gly	Gly	
	1130					1135						1140			
Phe	Ser	Lys	Glu	Ser	Ile	Leu	Pro	Lys	Arg	Asn	Ser	Asp	Lys	Leu	
	1145					1150						1155			
Ile	Ala	Arg	Lys	Lys	Asp	Trp	Asp	Pro	Lys	Lys	Tyr	Gly	Gly	Phe	
	1160					1165						1170			
Asp	Ser	Pro	Thr	Val	Ala	Tyr	Ser	Val	Leu	Val	Val	Ala	Lys	Val	
	1175					1180						1185			
Glu	Lys	Gly	Lys	Ser	Lys	Lys	Leu	Lys	Ser	Val	Lys	Glu	Leu	Leu	
	1190					1195						1200			
Gly	Ile	Thr	Ile	Met	Glu	Arg	Ser	Ser	Phe	Glu	Lys	Asn	Pro	Ile	
	1205					1210						1215			
Asp	Phe	Leu	Glu	Ala	Lys	Gly	Tyr	Lys	Glu	Val	Lys	Lys	Asp	Leu	
	1220					1225						1230			
Ile	Ile	Lys	Leu	Pro	Lys	Tyr	Ser	Leu	Phe	Glu	Leu	Glu	Asn	Gly	
	1235					1240						1245			

-continued

Arg Lys	Arg Met	Leu Ala	Ser	Ala Gly	Glu Leu	Gln	Lys Gly	Asn		
1250			1255			1260				
Glu Leu	Ala Leu	Pro Ser	Lys	Tyr Val	Asn Phe	Leu	Tyr Leu	Ala		
1265			1270			1275				
Ser His	Tyr Glu	Lys Leu	Lys	Gly Ser	Pro Glu	Asp	Asn Glu	Gln		
1280			1285			1290				
Lys Gln	Leu Phe	Val Glu	Gln	His Lys	His Tyr	Leu	Asp Glu	Ile		
1295			1300			1305				
Ile Glu	Gln Ile	Ser Glu	Phe	Ser Lys	Arg Val	Ile	Leu Ala	Asp		
1310			1315			1320				
Ala Asn	Leu Asp	Lys Val	Leu	Ser Ala	Tyr Asn	Lys	His Arg	Asp		
1325			1330			1335				
Lys Pro	Ile Arg	Glu Gln	Ala	Glu Asn	Ile Ile	His	Leu Phe	Thr		
1340			1345			1350				
Leu Thr	Asn Leu	Gly Ala	Pro	Ala Ala	Phe Lys	Tyr	Phe Asp	Thr		
1355			1360			1365				
Thr Ile	Asp Arg	Lys Arg	Tyr	Thr Ser	Thr Lys	Glu	Val Leu	Asp		
1370			1375			1380				
Ala Thr	Leu Ile	His Gln	Ser	Ile Thr	Gly Leu	Tyr	Glu Thr	Arg		
1385			1390			1395				
Ile Asp	Leu Ser	Gln Leu	Gly	Gly Asp	Ala Ala	Ala	Val Ser	Lys		
1400			1405			1410				
Gly Glu	Glu Leu	Phe Thr	Gly	Val Val	Pro Ile	Leu	Val Glu	Leu		
1415			1420			1425				
Asp Gly	Asp Val	Asn Gly	His	Lys Phe	Ser Val	Ser	Gly Glu	Gly		
1430			1435			1440				
Glu Gly	Asp Ala	Thr Tyr	Gly	Lys Leu	Thr Leu	Lys	Phe Ile	Cys		
1445			1450			1455				
Thr Thr	Gly Lys	Leu Pro	Val	Pro Trp	Pro Thr	Leu	Val Thr	Thr		
1460			1465			1470				
Leu Thr	Tyr Gly	Val Gln	Cys	Phe Ser	Arg Tyr	Pro	Asp His	Met		
1475			1480			1485				
Lys Gln	His Asp	Phe Phe	Lys	Ser Ala	Met Pro	Glu	Gly Tyr	Val		
1490			1495			1500				
Gln Glu	Arg Thr	Ile Phe	Phe	Lys Asp	Asp Gly	Asn	Tyr Lys	Thr		
1505			1510			1515				
Arg Ala	Glu Val	Lys Phe	Glu	Gly Asp	Thr Leu	Val	Asn Arg	Ile		
1520			1525			1530				
Glu Leu	Lys Gly	Ile Asp	Phe	Lys Glu	Asp Gly	Asn	Ile Leu	Gly		
1535			1540			1545				
His Lys	Leu Glu	Tyr Asn	Tyr	Asn Ser	His Asn	Val	Tyr Ile	Met		
1550			1555			1560				
Ala Asp	Lys Gln	Lys Asn	Gly	Ile Lys	Val Asn	Phe	Lys Ile	Arg		
1565			1570			1575				
His Asn	Ile Glu	Asp Gly	Ser	Val Gln	Leu Ala	Asp	His Tyr	Gln		
1580			1585			1590				
Gln Asn	Thr Pro	Ile Gly	Asp	Gly Pro	Val Leu	Leu	Pro Asp	Asn		
1595			1600			1605				
His Tyr	Leu Ser	Thr Gln	Ser	Ala Leu	Ser Lys	Asp	Pro Asn	Glu		
1610			1615			1620				

-continued

Lys Arg Asp His Met Val Leu Leu Glu Phe Val Thr Ala Ala Gly
1625 1630 1635

Ile Thr Leu Gly Met Asp Glu Leu Tyr Lys Lys Arg Pro Ala Ala
1640 1645 1650

Thr Lys Lys Ala Gly Gln Ala Lys Lys Lys Lys
1655 1660

<210> SEQ ID NO 47

<211> LENGTH: 1423

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polypeptide

<400> SEQUENCE: 47

Met Asp Tyr Lys Asp His Asp Gly Asp Tyr Lys Asp His Asp Ile Asp
1 5 10 15

Tyr Lys Asp Asp Asp Lys Met Ala Pro Lys Lys Lys Arg Lys Val
20 25 30

Gly Ile His Gly Val Pro Ala Ala Asp Lys Lys Tyr Ser Ile Gly Leu
35 40 45

Asp Ile Gly Thr Asn Ser Val Gly Trp Ala Val Ile Thr Asp Glu Tyr
50 55 60

Lys Val Pro Ser Lys Lys Phe Lys Val Leu Gly Asn Thr Asp Arg His
65 70 75 80

Ser Ile Lys Lys Asn Leu Ile Gly Ala Leu Leu Phe Asp Ser Gly Glu
85 90 95

Thr Ala Glu Ala Thr Arg Leu Lys Arg Thr Ala Arg Arg Arg Tyr Thr
100 105 110

Arg Arg Lys Asn Arg Ile Cys Tyr Leu Gln Glu Ile Phe Ser Asn Glu
115 120 125

Met Ala Lys Val Asp Asp Ser Phe Phe His Arg Leu Glu Glu Ser Phe
130 135 140

Leu Val Glu Glu Asp Lys Lys His Glu Arg His Pro Ile Phe Gly Asn
145 150 155 160

Ile Val Asp Glu Val Ala Tyr His Glu Lys Tyr Pro Thr Ile Tyr His
165 170 175

Leu Arg Lys Lys Leu Val Asp Ser Thr Asp Lys Ala Asp Leu Arg Leu
180 185 190

Ile Tyr Leu Ala Leu Ala His Met Ile Lys Phe Arg Gly His Phe Leu
195 200 205

Ile Glu Gly Asp Leu Asn Pro Asp Asn Ser Asp Val Asp Lys Leu Phe
210 215 220

Ile Gln Leu Val Gln Thr Tyr Asn Gln Leu Phe Glu Glu Asn Pro Ile
225 230 235 240

Asn Ala Ser Gly Val Asp Ala Lys Ala Ile Leu Ser Ala Arg Leu Ser
245 250 255

Lys Ser Arg Arg Leu Glu Asn Leu Ile Ala Gln Leu Pro Gly Glu Lys
260 265 270

Lys Asn Gly Leu Phe Gly Asn Leu Ile Ala Leu Ser Leu Gly Leu Thr
275 280 285

Pro Asn Phe Lys Ser Asn Phe Asp Leu Ala Glu Asp Ala Lys Leu Gln
290 295 300

-continued

Leu Ser Lys Asp Thr Tyr Asp Asp Asp Leu Asp Asn Leu Leu Ala Gln
 305 310 315 320
 Ile Gly Asp Gln Tyr Ala Asp Leu Phe Leu Ala Ala Lys Asn Leu Ser
 325 330 335
 Asp Ala Ile Leu Leu Ser Asp Ile Leu Arg Val Asn Thr Glu Ile Thr
 340 345 350
 Lys Ala Pro Leu Ser Ala Ser Met Ile Lys Arg Tyr Asp Glu His His
 355 360 365
 Gln Asp Leu Thr Leu Leu Lys Ala Leu Val Arg Gln Gln Leu Pro Glu
 370 375 380
 Lys Tyr Lys Glu Ile Phe Phe Asp Gln Ser Lys Asn Gly Tyr Ala Gly
 385 390 395 400
 Tyr Ile Asp Gly Gly Ala Ser Gln Glu Glu Phe Tyr Lys Phe Ile Lys
 405 410 415
 Pro Ile Leu Glu Lys Met Asp Gly Thr Glu Glu Leu Leu Val Lys Leu
 420 425 430
 Asn Arg Glu Asp Leu Leu Arg Lys Gln Arg Thr Phe Asp Asn Gly Ser
 435 440 445
 Ile Pro His Gln Ile His Leu Gly Glu Leu His Ala Ile Leu Arg Arg
 450 455 460
 Gln Glu Asp Phe Tyr Pro Phe Leu Lys Asp Asn Arg Glu Lys Ile Glu
 465 470 475 480
 Lys Ile Leu Thr Phe Arg Ile Pro Tyr Tyr Val Gly Pro Leu Ala Arg
 485 490 495
 Gly Asn Ser Arg Phe Ala Trp Met Thr Arg Lys Ser Glu Glu Thr Ile
 500 505 510
 Thr Pro Trp Asn Phe Glu Glu Val Val Asp Lys Gly Ala Ser Ala Gln
 515 520 525
 Ser Phe Ile Glu Arg Met Thr Asn Phe Asp Lys Asn Leu Pro Asn Glu
 530 535 540
 Lys Val Leu Pro Lys His Ser Leu Leu Tyr Glu Tyr Phe Thr Val Tyr
 545 550 555 560
 Asn Glu Leu Thr Lys Val Lys Tyr Val Thr Glu Gly Met Arg Lys Pro
 565 570 575
 Ala Phe Leu Ser Gly Glu Gln Lys Lys Ala Ile Val Asp Leu Leu Phe
 580 585 590
 Lys Thr Asn Arg Lys Val Thr Val Lys Gln Leu Lys Glu Asp Tyr Phe
 595 600 605
 Lys Lys Ile Glu Cys Phe Asp Ser Val Glu Ile Ser Gly Val Glu Asp
 610 615 620
 Arg Phe Asn Ala Ser Leu Gly Thr Tyr His Asp Leu Leu Lys Ile Ile
 625 630 635 640
 Lys Asp Lys Asp Phe Leu Asp Asn Glu Glu Asn Glu Asp Ile Leu Glu
 645 650 655
 Asp Ile Val Leu Thr Leu Thr Leu Phe Glu Asp Arg Glu Met Ile Glu
 660 665 670
 Glu Arg Leu Lys Thr Tyr Ala His Leu Phe Asp Asp Lys Val Met Lys
 675 680 685
 Gln Leu Lys Arg Arg Arg Tyr Thr Gly Trp Gly Arg Leu Ser Arg Lys
 690 695 700

-continued

Leu Ile Asn Gly Ile Arg Asp Lys Gln Ser Gly Lys Thr Ile Leu Asp
 705 710 715 720
 Phe Leu Lys Ser Asp Gly Phe Ala Asn Arg Asn Phe Met Gln Leu Ile
 725 730 735
 His Asp Asp Ser Leu Thr Phe Lys Glu Asp Ile Gln Lys Ala Gln Val
 740 745 750
 Ser Gly Gln Gly Asp Ser Leu His Glu His Ile Ala Asn Leu Ala Gly
 755 760 765
 Ser Pro Ala Ile Lys Lys Gly Ile Leu Gln Thr Val Lys Val Val Asp
 770 775 780
 Glu Leu Val Lys Val Met Gly Arg His Lys Pro Glu Asn Ile Val Ile
 785 790 795 800
 Glu Met Ala Arg Glu Asn Gln Thr Thr Gln Lys Gly Gln Lys Asn Ser
 805 810 815
 Arg Glu Arg Met Lys Arg Ile Glu Glu Gly Ile Lys Glu Leu Gly Ser
 820 825 830
 Gln Ile Leu Lys Glu His Pro Val Glu Asn Thr Gln Leu Gln Asn Glu
 835 840 845
 Lys Leu Tyr Leu Tyr Tyr Leu Gln Asn Gly Arg Asp Met Tyr Val Asp
 850 855 860
 Gln Glu Leu Asp Ile Asn Arg Leu Ser Asp Tyr Asp Val Asp His Ile
 865 870 875 880
 Val Pro Gln Ser Phe Leu Lys Asp Asp Ser Ile Asp Asn Lys Val Leu
 885 890 895
 Thr Arg Ser Asp Lys Asn Arg Gly Lys Ser Asp Asn Val Pro Ser Glu
 900 905 910
 Glu Val Val Lys Lys Met Lys Asn Tyr Trp Arg Gln Leu Leu Asn Ala
 915 920 925
 Lys Leu Ile Thr Gln Arg Lys Phe Asp Asn Leu Thr Lys Ala Glu Arg
 930 935 940
 Gly Gly Leu Ser Glu Leu Asp Lys Ala Gly Phe Ile Lys Arg Gln Leu
 945 950 955 960
 Val Glu Thr Arg Gln Ile Thr Lys His Val Ala Gln Ile Leu Asp Ser
 965 970 975
 Arg Met Asn Thr Lys Tyr Asp Glu Asn Asp Lys Leu Ile Arg Glu Val
 980 985 990
 Lys Val Ile Thr Leu Lys Ser Lys Leu Val Ser Asp Phe Arg Lys Asp
 995 1000 1005
 Phe Gln Phe Tyr Lys Val Arg Glu Ile Asn Asn Tyr His His Ala
 1010 1015 1020
 His Asp Ala Tyr Leu Asn Ala Val Val Gly Thr Ala Leu Ile Lys
 1025 1030 1035
 Lys Tyr Pro Lys Leu Glu Ser Glu Phe Val Tyr Gly Asp Tyr Lys
 1040 1045 1050
 Val Tyr Asp Val Arg Lys Met Ile Ala Lys Ser Glu Gln Glu Ile
 1055 1060 1065
 Gly Lys Ala Thr Ala Lys Tyr Phe Phe Tyr Ser Asn Ile Met Asn
 1070 1075 1080
 Phe Phe Lys Thr Glu Ile Thr Leu Ala Asn Gly Glu Ile Arg Lys
 1085 1090 1095
 Arg Pro Leu Ile Glu Thr Asn Gly Glu Thr Gly Glu Ile Val Trp

-continued

1100	1105	1110
Asp Lys Gly Arg Asp Phe Ala Thr Val Arg Lys Val Leu Ser Met		
1115	1120	1125
Pro Gln Val Asn Ile Val Lys Lys Thr Glu Val Gln Thr Gly Gly		
1130	1135	1140
Phe Ser Lys Glu Ser Ile Leu Pro Lys Arg Asn Ser Asp Lys Leu		
1145	1150	1155
Ile Ala Arg Lys Lys Asp Trp Asp Pro Lys Lys Tyr Gly Gly Phe		
1160	1165	1170
Asp Ser Pro Thr Val Ala Tyr Ser Val Leu Val Val Ala Lys Val		
1175	1180	1185
Glu Lys Gly Lys Ser Lys Lys Leu Lys Ser Val Lys Glu Leu Leu		
1190	1195	1200
Gly Ile Thr Ile Met Glu Arg Ser Ser Phe Glu Lys Asn Pro Ile		
1205	1210	1215
Asp Phe Leu Glu Ala Lys Gly Tyr Lys Glu Val Lys Lys Asp Leu		
1220	1225	1230
Ile Ile Lys Leu Pro Lys Tyr Ser Leu Phe Glu Leu Glu Asn Gly		
1235	1240	1245
Arg Lys Arg Met Leu Ala Ser Ala Gly Glu Leu Gln Lys Gly Asn		
1250	1255	1260
Glu Leu Ala Leu Pro Ser Lys Tyr Val Asn Phe Leu Tyr Leu Ala		
1265	1270	1275
Ser His Tyr Glu Lys Leu Lys Gly Ser Pro Glu Asp Asn Glu Gln		
1280	1285	1290
Lys Gln Leu Phe Val Glu Gln His Lys His Tyr Leu Asp Glu Ile		
1295	1300	1305
Ile Glu Gln Ile Ser Glu Phe Ser Lys Arg Val Ile Leu Ala Asp		
1310	1315	1320
Ala Asn Leu Asp Lys Val Leu Ser Ala Tyr Asn Lys His Arg Asp		
1325	1330	1335
Lys Pro Ile Arg Glu Gln Ala Glu Asn Ile Ile His Leu Phe Thr		
1340	1345	1350
Leu Thr Asn Leu Gly Ala Pro Ala Ala Phe Lys Tyr Phe Asp Thr		
1355	1360	1365
Thr Ile Asp Arg Lys Arg Tyr Thr Ser Thr Lys Glu Val Leu Asp		
1370	1375	1380
Ala Thr Leu Ile His Gln Ser Ile Thr Gly Leu Tyr Glu Thr Arg		
1385	1390	1395
Ile Asp Leu Ser Gln Leu Gly Gly Asp Lys Arg Pro Ala Ala Thr		
1400	1405	1410
Lys Lys Ala Gly Gln Ala Lys Lys Lys Lys		
1415	1420	

<210> SEQ ID NO 48

<211> LENGTH: 483

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polypeptide

<400> SEQUENCE: 48

-continued

Met	Phe	Leu	Phe	Leu	Ser	Leu	Thr	Ser	Phe	Leu	Ser	Ser	Ser	Arg	Thr
1				5					10					15	
Leu	Val	Ser	Lys	Gly	Glu	Glu	Asp	Asn	Met	Ala	Ile	Ile	Lys	Glu	Phe
			20					25					30		
Met	Arg	Phe	Lys	Val	His	Met	Glu	Gly	Ser	Val	Asn	Gly	His	Glu	Phe
		35					40					45			
Glu	Ile	Glu	Gly	Glu	Gly	Glu	Gly	Arg	Pro	Tyr	Glu	Gly	Thr	Gln	Thr
	50					55					60				
Ala	Lys	Leu	Lys	Val	Thr	Lys	Gly	Gly	Pro	Leu	Pro	Phe	Ala	Trp	Asp
65					70					75					80
Ile	Leu	Ser	Pro	Gln	Phe	Met	Tyr	Gly	Ser	Lys	Ala	Tyr	Val	Lys	His
				85					90					95	
Pro	Ala	Asp	Ile	Pro	Asp	Tyr	Leu	Lys	Leu	Ser	Phe	Pro	Glu	Gly	Phe
			100					105					110		
Lys	Trp	Glu	Arg	Val	Met	Asn	Phe	Glu	Asp	Gly	Gly	Val	Val	Thr	Val
		115					120					125			
Thr	Gln	Asp	Ser	Ser	Leu	Gln	Asp	Gly	Glu	Phe	Ile	Tyr	Lys	Val	Lys
	130					135					140				
Leu	Arg	Gly	Thr	Asn	Phe	Pro	Ser	Asp	Gly	Pro	Val	Met	Gln	Lys	Lys
145					150					155					160
Thr	Met	Gly	Trp	Glu	Ala	Ser	Ser	Glu	Arg	Met	Tyr	Pro	Glu	Asp	Gly
				165					170					175	
Ala	Leu	Lys	Gly	Glu	Ile	Lys	Gln	Arg	Leu	Lys	Leu	Lys	Asp	Gly	Gly
			180					185					190		
His	Tyr	Asp	Ala	Glu	Val	Lys	Thr	Thr	Tyr	Lys	Ala	Lys	Lys	Pro	Val
		195					200					205			
Gln	Leu	Pro	Gly	Ala	Tyr	Asn	Val	Asn	Ile	Lys	Leu	Asp	Ile	Thr	Ser
	210					215					220				
His	Asn	Glu	Asp	Tyr	Thr	Ile	Val	Glu	Gln	Tyr	Glu	Arg	Ala	Glu	Gly
225					230					235					240
Arg	His	Ser	Thr	Gly	Gly	Met	Asp	Glu	Leu	Tyr	Lys	Gly	Ser	Lys	Gln
				245					250					255	
Leu	Glu	Glu	Leu	Leu	Ser	Thr	Ser	Phe	Asp	Ile	Gln	Phe	Asn	Asp	Leu
			260					265					270		
Thr	Leu	Leu	Glu	Thr	Ala	Phe	Thr	His	Thr	Ser	Tyr	Ala	Asn	Glu	His
		275					280					285			
Arg	Leu	Leu	Asn	Val	Ser	His	Asn	Glu	Arg	Leu	Glu	Phe	Leu	Gly	Asp
	290					295					300				
Ala	Val	Leu	Gln	Leu	Ile	Ile	Ser	Glu	Tyr	Leu	Phe	Ala	Lys	Tyr	Pro
305					310					315					320
Lys	Lys	Thr	Glu	Gly	Asp	Met	Ser	Lys	Leu	Arg	Ser	Met	Ile	Val	Arg
				325					330					335	
Glu	Glu	Ser	Leu	Ala	Gly	Phe	Ser	Arg	Phe	Cys	Ser	Phe	Asp	Ala	Tyr
			340					345					350		
Ile	Lys	Leu	Gly	Lys	Gly	Glu	Glu	Lys	Ser	Gly	Gly	Arg	Arg	Arg	Asp
		355					360					365			
Thr	Ile	Leu	Gly	Asp	Leu	Phe	Glu	Ala	Phe	Leu	Gly	Ala	Leu	Leu	Leu
	370				375						380				
Asp	Lys	Gly	Ile	Asp	Ala	Val	Arg	Arg	Phe	Leu	Lys	Gln	Val	Met	Ile
385					390					395					400
Pro	Gln	Val	Glu	Lys	Gly	Asn	Phe	Glu	Arg	Val	Lys	Asp	Tyr	Lys	Thr

-continued

260	265	270
Pro Tyr Glu Gly Thr Gln Thr Ala Lys Leu Lys Val Thr Lys Gly Gly		
275	280	285
Pro Leu Pro Phe Ala Trp Asp Ile Leu Ser Pro Gln Phe Met Tyr Gly		
290	295	300
Ser Lys Ala Tyr Val Lys His Pro Ala Asp Ile Pro Asp Tyr Leu Lys		
305	310	315
Leu Ser Phe Pro Glu Gly Phe Lys Trp Glu Arg Val Met Asn Phe Glu		
325	330	335
Asp Gly Gly Val Val Thr Val Thr Gln Asp Ser Ser Leu Gln Asp Gly		
340	345	350
Glu Phe Ile Tyr Lys Val Lys Leu Arg Gly Thr Asn Phe Pro Ser Asp		
355	360	365
Gly Pro Val Met Gln Lys Lys Thr Met Gly Trp Glu Ala Ser Ser Glu		
370	375	380
Arg Met Tyr Pro Glu Asp Gly Ala Leu Lys Gly Glu Ile Lys Gln Arg		
385	390	395
Leu Lys Leu Lys Asp Gly Gly His Tyr Asp Ala Glu Val Lys Thr Thr		
405	410	415
Tyr Lys Ala Lys Lys Pro Val Gln Leu Pro Gly Ala Tyr Asn Val Asn		
420	425	430
Ile Lys Leu Asp Ile Thr Ser His Asn Glu Asp Tyr Thr Ile Val Glu		
435	440	445
Gln Tyr Glu Arg Ala Glu Gly Arg His Ser Thr Gly Gly Met Asp Glu		
450	455	460
Leu Tyr Lys Lys Arg Pro Ala Ala Thr Lys Lys Ala Gly Gln Ala Lys		
465	470	475
480		

Lys Lys Lys

<210> SEQ ID NO 50
 <211> LENGTH: 1423
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polypeptide

<400> SEQUENCE: 50

Met Asp Tyr Lys Asp His Asp Gly Asp Tyr Lys Asp His Asp Ile Asp
1 5 10 15
Tyr Lys Asp Asp Asp Lys Met Ala Pro Lys Lys Lys Arg Lys Val
20 25 30
Gly Ile His Gly Val Pro Ala Ala Asp Lys Lys Tyr Ser Ile Gly Leu
35 40 45
Ala Ile Gly Thr Asn Ser Val Gly Trp Ala Val Ile Thr Asp Glu Tyr
50 55 60
Lys Val Pro Ser Lys Lys Phe Lys Val Leu Gly Asn Thr Asp Arg His
65 70 75 80
Ser Ile Lys Lys Asn Leu Ile Gly Ala Leu Leu Phe Asp Ser Gly Glu
85 90 95
Thr Ala Glu Ala Thr Arg Leu Lys Arg Thr Ala Arg Arg Arg Tyr Thr
100 105 110
Arg Arg Lys Asn Arg Ile Cys Tyr Leu Gln Glu Ile Phe Ser Asn Glu

-continued

115		120				125									
Met	Ala	Lys	Val	Asp	Asp	Ser	Phe	Phe	His	Arg	Leu	Glu	Glu	Ser	Phe
	130					135					140				
Leu	Val	Glu	Glu	Asp	Lys	Lys	His	Glu	Arg	His	Pro	Ile	Phe	Gly	Asn
	145				150					155					160
Ile	Val	Asp	Glu	Val	Ala	Tyr	His	Glu	Lys	Tyr	Pro	Thr	Ile	Tyr	His
				165					170					175	
Leu	Arg	Lys	Lys	Leu	Val	Asp	Ser	Thr	Asp	Lys	Ala	Asp	Leu	Arg	Leu
			180					185					190		
Ile	Tyr	Leu	Ala	Leu	Ala	His	Met	Ile	Lys	Phe	Arg	Gly	His	Phe	Leu
		195					200					205			
Ile	Glu	Gly	Asp	Leu	Asn	Pro	Asp	Asn	Ser	Asp	Val	Asp	Lys	Leu	Phe
	210					215					220				
Ile	Gln	Leu	Val	Gln	Thr	Tyr	Asn	Gln	Leu	Phe	Glu	Glu	Asn	Pro	Ile
	225				230					235					240
Asn	Ala	Ser	Gly	Val	Asp	Ala	Lys	Ala	Ile	Leu	Ser	Ala	Arg	Leu	Ser
				245					250					255	
Lys	Ser	Arg	Arg	Leu	Glu	Asn	Leu	Ile	Ala	Gln	Leu	Pro	Gly	Glu	Lys
			260					265					270		
Lys	Asn	Gly	Leu	Phe	Gly	Asn	Leu	Ile	Ala	Leu	Ser	Leu	Gly	Leu	Thr
		275					280					285			
Pro	Asn	Phe	Lys	Ser	Asn	Phe	Asp	Leu	Ala	Glu	Asp	Ala	Lys	Leu	Gln
	290					295					300				
Leu	Ser	Lys	Asp	Thr	Tyr	Asp	Asp	Asp	Leu	Asp	Asn	Leu	Leu	Ala	Gln
	305				310					315					320
Ile	Gly	Asp	Gln	Tyr	Ala	Asp	Leu	Phe	Leu	Ala	Ala	Lys	Asn	Leu	Ser
				325					330					335	
Asp	Ala	Ile	Leu	Leu	Ser	Asp	Ile	Leu	Arg	Val	Asn	Thr	Glu	Ile	Thr
		340						345					350		
Lys	Ala	Pro	Leu	Ser	Ala	Ser	Met	Ile	Lys	Arg	Tyr	Asp	Glu	His	His
		355					360					365			
Gln	Asp	Leu	Thr	Leu	Leu	Lys	Ala	Leu	Val	Arg	Gln	Gln	Leu	Pro	Glu
	370					375					380				
Lys	Tyr	Lys	Glu	Ile	Phe	Phe	Asp	Gln	Ser	Lys	Asn	Gly	Tyr	Ala	Gly
	385				390					395					400
Tyr	Ile	Asp	Gly	Gly	Ala	Ser	Gln	Glu	Glu	Phe	Tyr	Lys	Phe	Ile	Lys
				405					410					415	
Pro	Ile	Leu	Glu	Lys	Met	Asp	Gly	Thr	Glu	Glu	Leu	Leu	Val	Lys	Leu
		420						425					430		
Asn	Arg	Glu	Asp	Leu	Leu	Arg	Lys	Gln	Arg	Thr	Phe	Asp	Asn	Gly	Ser
		435					440					445			
Ile	Pro	His	Gln	Ile	His	Leu	Gly	Glu	Leu	His	Ala	Ile	Leu	Arg	Arg
	450					455					460				
Gln	Glu	Asp	Phe	Tyr	Pro	Phe	Leu	Lys	Asp	Asn	Arg	Glu	Lys	Ile	Glu
	465				470					475					480
Lys	Ile	Leu	Thr	Phe	Arg	Ile	Pro	Tyr	Tyr	Val	Gly	Pro	Leu	Ala	Arg
				485					490					495	
Gly	Asn	Ser	Arg	Phe	Ala	Trp	Met	Thr	Arg	Lys	Ser	Glu	Glu	Thr	Ile
			500					505					510		
Thr	Pro	Trp	Asn	Phe	Glu	Glu	Val	Val	Asp	Lys	Gly	Ala	Ser	Ala	Gln
		515					520					525			

-continued

Ser Phe Ile Glu Arg Met Thr Asn Phe Asp Lys Asn Leu Pro Asn Glu
 530 535 540
 Lys Val Leu Pro Lys His Ser Leu Leu Tyr Glu Tyr Phe Thr Val Tyr
 545 550 555 560
 Asn Glu Leu Thr Lys Val Lys Tyr Val Thr Glu Gly Met Arg Lys Pro
 565 570 575
 Ala Phe Leu Ser Gly Glu Gln Lys Lys Ala Ile Val Asp Leu Leu Phe
 580 585 590
 Lys Thr Asn Arg Lys Val Thr Val Lys Gln Leu Lys Glu Asp Tyr Phe
 595 600 605
 Lys Lys Ile Glu Cys Phe Asp Ser Val Glu Ile Ser Gly Val Glu Asp
 610 615 620
 Arg Phe Asn Ala Ser Leu Gly Thr Tyr His Asp Leu Leu Lys Ile Ile
 625 630 635 640
 Lys Asp Lys Asp Phe Leu Asp Asn Glu Glu Asn Glu Asp Ile Leu Glu
 645 650 655
 Asp Ile Val Leu Thr Leu Thr Leu Phe Glu Asp Arg Glu Met Ile Glu
 660 665 670
 Glu Arg Leu Lys Thr Tyr Ala His Leu Phe Asp Asp Lys Val Met Lys
 675 680 685
 Gln Leu Lys Arg Arg Arg Tyr Thr Gly Trp Gly Arg Leu Ser Arg Lys
 690 695 700
 Leu Ile Asn Gly Ile Arg Asp Lys Gln Ser Gly Lys Thr Ile Leu Asp
 705 710 715 720
 Phe Leu Lys Ser Asp Gly Phe Ala Asn Arg Asn Phe Met Gln Leu Ile
 725 730 735
 His Asp Asp Ser Leu Thr Phe Lys Glu Asp Ile Gln Lys Ala Gln Val
 740 745 750
 Ser Gly Gln Gly Asp Ser Leu His Glu His Ile Ala Asn Leu Ala Gly
 755 760 765
 Ser Pro Ala Ile Lys Lys Gly Ile Leu Gln Thr Val Lys Val Val Asp
 770 775 780
 Glu Leu Val Lys Val Met Gly Arg His Lys Pro Glu Asn Ile Val Ile
 785 790 795 800
 Glu Met Ala Arg Glu Asn Gln Thr Thr Gln Lys Gly Gln Lys Asn Ser
 805 810 815
 Arg Glu Arg Met Lys Arg Ile Glu Glu Gly Ile Lys Glu Leu Gly Ser
 820 825 830
 Gln Ile Leu Lys Glu His Pro Val Glu Asn Thr Gln Leu Gln Asn Glu
 835 840 845
 Lys Leu Tyr Leu Tyr Tyr Leu Gln Asn Gly Arg Asp Met Tyr Val Asp
 850 855 860
 Gln Glu Leu Asp Ile Asn Arg Leu Ser Asp Tyr Asp Val Asp His Ile
 865 870 875 880
 Val Pro Gln Ser Phe Leu Lys Asp Asp Ser Ile Asp Asn Lys Val Leu
 885 890 895
 Thr Arg Ser Asp Lys Asn Arg Gly Lys Ser Asp Asn Val Pro Ser Glu
 900 905 910
 Glu Val Val Lys Lys Met Lys Asn Tyr Trp Arg Gln Leu Leu Asn Ala
 915 920 925

-continued

Lys Leu Ile Thr Gln Arg Lys Phe Asp Asn Leu Thr Lys Ala Glu Arg
 930 935 940

Gly Gly Leu Ser Glu Leu Asp Lys Ala Gly Phe Ile Lys Arg Gln Leu
 945 950 955 960

Val Glu Thr Arg Gln Ile Thr Lys His Val Ala Gln Ile Leu Asp Ser
 965 970 975

Arg Met Asn Thr Lys Tyr Asp Glu Asn Asp Lys Leu Ile Arg Glu Val
 980 985 990

Lys Val Ile Thr Leu Lys Ser Lys Leu Val Ser Asp Phe Arg Lys Asp
 995 1000 1005

Phe Gln Phe Tyr Lys Val Arg Glu Ile Asn Asn Tyr His His Ala
 1010 1015 1020

His Asp Ala Tyr Leu Asn Ala Val Val Gly Thr Ala Leu Ile Lys
 1025 1030 1035

Lys Tyr Pro Lys Leu Glu Ser Glu Phe Val Tyr Gly Asp Tyr Lys
 1040 1045 1050

Val Tyr Asp Val Arg Lys Met Ile Ala Lys Ser Glu Gln Glu Ile
 1055 1060 1065

Gly Lys Ala Thr Ala Lys Tyr Phe Phe Tyr Ser Asn Ile Met Asn
 1070 1075 1080

Phe Phe Lys Thr Glu Ile Thr Leu Ala Asn Gly Glu Ile Arg Lys
 1085 1090 1095

Arg Pro Leu Ile Glu Thr Asn Gly Glu Thr Gly Glu Ile Val Trp
 1100 1105 1110

Asp Lys Gly Arg Asp Phe Ala Thr Val Arg Lys Val Leu Ser Met
 1115 1120 1125

Pro Gln Val Asn Ile Val Lys Lys Thr Glu Val Gln Thr Gly Gly
 1130 1135 1140

Phe Ser Lys Glu Ser Ile Leu Pro Lys Arg Asn Ser Asp Lys Leu
 1145 1150 1155

Ile Ala Arg Lys Lys Asp Trp Asp Pro Lys Lys Tyr Gly Gly Phe
 1160 1165 1170

Asp Ser Pro Thr Val Ala Tyr Ser Val Leu Val Val Ala Lys Val
 1175 1180 1185

Glu Lys Gly Lys Ser Lys Lys Leu Lys Ser Val Lys Glu Leu Leu
 1190 1195 1200

Gly Ile Thr Ile Met Glu Arg Ser Ser Phe Glu Lys Asn Pro Ile
 1205 1210 1215

Asp Phe Leu Glu Ala Lys Gly Tyr Lys Glu Val Lys Lys Asp Leu
 1220 1225 1230

Ile Ile Lys Leu Pro Lys Tyr Ser Leu Phe Glu Leu Glu Asn Gly
 1235 1240 1245

Arg Lys Arg Met Leu Ala Ser Ala Gly Glu Leu Gln Lys Gly Asn
 1250 1255 1260

Glu Leu Ala Leu Pro Ser Lys Tyr Val Asn Phe Leu Tyr Leu Ala
 1265 1270 1275

Ser His Tyr Glu Lys Leu Lys Gly Ser Pro Glu Asp Asn Glu Gln
 1280 1285 1290

Lys Gln Leu Phe Val Glu Gln His Lys His Tyr Leu Asp Glu Ile
 1295 1300 1305

Ile Glu Gln Ile Ser Glu Phe Ser Lys Arg Val Ile Leu Ala Asp

-continued

1310	1315	1320
Ala Asn Leu Asp Lys Val Leu Ser Ala Tyr Asn Lys His Arg Asp		
1325	1330	1335
Lys Pro Ile Arg Glu Gln Ala Glu Asn Ile Ile His Leu Phe Thr		
1340	1345	1350
Leu Thr Asn Leu Gly Ala Pro Ala Ala Phe Lys Tyr Phe Asp Thr		
1355	1360	1365
Thr Ile Asp Arg Lys Arg Tyr Thr Ser Thr Lys Glu Val Leu Asp		
1370	1375	1380
Ala Thr Leu Ile His Gln Ser Ile Thr Gly Leu Tyr Glu Thr Arg		
1385	1390	1395
Ile Asp Leu Ser Gln Leu Gly Gly Asp Lys Arg Pro Ala Ala Thr		
1400	1405	1410
Lys Lys Ala Gly Gln Ala Lys Lys Lys Lys		
1415	1420	

<210> SEQ ID NO 51
 <211> LENGTH: 2012
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 51

```

gaatgctgcc ctcagaccgc cttcctcct gtccttgtct gtccaaggag aatgaggctc 60
cactggtgga tttcggacta ccctgaggag ctggcacctg agggacaagg cccccacct 120
gccagctcc agcctctgat gaggggtggg agagagctac atgaggttgc taagaaagcc 180
tccccgaag gagaccacac agtgtgtgag gttggagtct ctagcagcgg gttctgtgcc 240
cccagggata gtctggctgt ccaggcactg ctcttgatat aaacaccacc tcctagtatt 300
gaaaccatgc ccattctgcc tctctgtatg gaaaagagca tggggctggc ccgtgggggtg 360
gtgtccactt taggcctgt gggagatcat gggaaaccac gcagtgggctc ataggctctc 420
tcatttacta ctcacatcca ctctgtgaag aagcgattat gatctctcct ctagaaactc 480
gtagagtccc atgtctgccg gcttccagag cctgcactcc tccacctgg cttggctttg 540
ctggggctag aggagctagg atgcacagca gctctgtgac cctttgtttg agaggaacag 600
gaaaaccacc cttctctctg gccactgtg tcctcttctc gccctgceat ccccttctgt 660
gaatgttaga cccatgggag cagctgggtca gaggggaccc cggcctgggg ccctaaacc 720
tatgtagcct cagtcttccc atcaggctct cagctcagcc tgagtgttga ggccccagtg 780
gctgctctgg gggcctcctg agtttctcat ctgtgccct ccctccctgg cccaggtgaa 840
ggtgtggttc cagaaccgga ggacaaagta caaacggcag aagctggagg aggaagggcc 900
tgagtccgag cagaagaaga agggctocca tcacatcaac cgggtggcgca ttgccacgaa 960
gcaggccaat ggggaggaca tcgatgtcac ctccaatgac aagcttgcta gcggtgggca 1020
accacaaacc cacgagggca gagtgtctct tgetgtggc caggcccctg cgtgggcccc 1080
agctggactc tggccactcc ctggccagge tttggggagg cctggagtca tggccccaca 1140
gggcttgaag cccggggccg ccattgacag agggacaagc aatgggctgg ctgaggcctg 1200
ggaccacttg gccttctcct cggagagcct gcctgcctgg gggggcccgc ccgccaccgc 1260
    
```


-continued

```

agcctcccag ctgctctccg tgtctccaat ctccttttg tttgatgca tttctgtttt 1320
aatttatttt ccaggcacca ctgtagttta gtgatcccca gtgtcccct tccctatggg 1380
aataataaaa gtctctctct taatgacacg ggcattccagc tccagcccca gagcctgggg 1440
tggtagattc cggctctgag ggccagtggg ggctggtaga gcaaacgcgt tcagggctctg 1500
ggagcctggg gtggggact ggtggagggg gtcaagggta attcattaac tcctctcttt 1560
tgttggggga ccctggtctc tacctccagc tccacagcag gagaaacagg ctagacatag 1620
ggaagggcca tcctgtatct tgagggagga caggcccagg tctttcttaa cgtattgaga 1680
ggtgggaatc agggccaggt agttcaatgg gagagggaga gtgcttcctt ctgcctagag 1740
actctggtgg cttctccagt tgaggagaaa ccagaggaaa ggggaggatt ggggtctggg 1800
ggagggaaaca ccattcacia aggctgacgg ttccagtcog aagtcgtggg cccaccagga 1860
tgctcacctg tccttgagaga accgctgggc aggttgagac tgcagagaca gggcttaagg 1920
ctgagcctgc aaccagtccc cagtgactca gggcctctc agcccaagaa agagcaactg 1980
gccagggccc gctgagctct tgtgttcacc tg 2012
    
```

```

<210> SEQ ID NO 52
<211> LENGTH: 1153
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        polypeptide
    
```

<400> SEQUENCE: 52

```

Met Lys Arg Pro Ala Ala Thr Lys Lys Ala Gly Gln Ala Lys Lys Lys
 1          5          10          15
Lys Ser Asp Leu Val Leu Gly Leu Asp Ile Gly Ile Gly Ser Val Gly
 20          25          30
Val Gly Ile Leu Asn Lys Val Thr Gly Glu Ile Ile His Lys Asn Ser
 35          40          45
Arg Ile Phe Pro Ala Ala Gln Ala Glu Asn Asn Leu Val Arg Arg Thr
 50          55          60
Asn Arg Gln Gly Arg Arg Leu Ala Arg Arg Lys Lys His Arg Arg Val
 65          70          75          80
Arg Leu Asn Arg Leu Phe Glu Glu Ser Gly Leu Ile Thr Asp Phe Thr
 85          90          95
Lys Ile Ser Ile Asn Leu Asn Pro Tyr Gln Leu Arg Val Lys Gly Leu
 100         105         110
Thr Asp Glu Leu Ser Asn Glu Glu Leu Phe Ile Ala Leu Lys Asn Met
 115         120         125
Val Lys His Arg Gly Ile Ser Tyr Leu Asp Asp Ala Ser Asp Asp Gly
 130         135         140
Asn Ser Ser Val Gly Asp Tyr Ala Gln Ile Val Lys Glu Asn Ser Lys
 145         150         155         160
Gln Leu Glu Thr Lys Thr Pro Gly Gln Ile Gln Leu Glu Arg Tyr Gln
 165         170         175
Thr Tyr Gly Gln Leu Arg Gly Asp Phe Thr Val Glu Lys Asp Gly Lys
 180         185         190
Lys His Arg Leu Ile Asn Val Phe Pro Thr Ser Ala Tyr Arg Ser Glu
 195         200         205
    
```

-continued

Ala Leu Arg Ile Leu Gln Thr Gln Gln Glu Phe Asn Pro Gln Ile Thr
210 215 220

Asp Glu Phe Ile Asn Arg Tyr Leu Glu Ile Leu Thr Gly Lys Arg Lys
225 230 235 240

Tyr Tyr His Gly Pro Gly Asn Glu Lys Ser Arg Thr Asp Tyr Gly Arg
245 250 255

Tyr Arg Thr Ser Gly Glu Thr Leu Asp Asn Ile Phe Gly Ile Leu Ile
260 265 270

Gly Lys Cys Thr Phe Tyr Pro Asp Glu Phe Arg Ala Ala Lys Ala Ser
275 280 285

Tyr Thr Ala Gln Glu Phe Asn Leu Leu Asn Asp Leu Asn Asn Leu Thr
290 295 300

Val Pro Thr Glu Thr Lys Lys Leu Ser Lys Glu Gln Lys Asn Gln Ile
305 310 315 320

Ile Asn Tyr Val Lys Asn Glu Lys Ala Met Gly Pro Ala Lys Leu Phe
325 330 335

Lys Tyr Ile Ala Lys Leu Leu Ser Cys Asp Val Ala Asp Ile Lys Gly
340 345 350

Tyr Arg Ile Asp Lys Ser Gly Lys Ala Glu Ile His Thr Phe Glu Ala
355 360 365

Tyr Arg Lys Met Lys Thr Leu Glu Thr Leu Asp Ile Glu Gln Met Asp
370 375 380

Arg Glu Thr Leu Asp Lys Leu Ala Tyr Val Leu Thr Leu Asn Thr Glu
385 390 395 400

Arg Glu Gly Ile Gln Glu Ala Leu Glu His Glu Phe Ala Asp Gly Ser
405 410 415

Phe Ser Gln Lys Gln Val Asp Glu Leu Val Gln Phe Arg Lys Ala Asn
420 425 430

Ser Ser Ile Phe Gly Lys Gly Trp His Asn Phe Ser Val Lys Leu Met
435 440 445

Met Glu Leu Ile Pro Glu Leu Tyr Glu Thr Ser Glu Glu Gln Met Thr
450 455 460

Ile Leu Thr Arg Leu Gly Lys Gln Lys Thr Thr Ser Ser Ser Asn Lys
465 470 475 480

Thr Lys Tyr Ile Asp Glu Lys Leu Leu Thr Glu Glu Ile Tyr Asn Pro
485 490 495

Val Val Ala Lys Ser Val Arg Gln Ala Ile Lys Ile Val Asn Ala Ala
500 505 510

Ile Lys Glu Tyr Gly Asp Phe Asp Asn Ile Val Ile Glu Met Ala Arg
515 520 525

Glu Thr Asn Glu Asp Asp Glu Lys Lys Ala Ile Gln Lys Ile Gln Lys
530 535 540

Ala Asn Lys Asp Glu Lys Asp Ala Ala Met Leu Lys Ala Ala Asn Gln
545 550 555 560

Tyr Asn Gly Lys Ala Glu Leu Pro His Ser Val Phe His Gly His Lys
565 570 575

Gln Leu Ala Thr Lys Ile Arg Leu Trp His Gln Gln Gly Glu Arg Cys
580 585 590

Leu Tyr Thr Gly Lys Thr Ile Ser Ile His Asp Leu Ile Asn Asn Ser
595 600 605

Asn Gln Phe Glu Val Asp His Ile Leu Pro Leu Ser Ile Thr Phe Asp

-continued

610			615			620									
Asp	Ser	Leu	Ala	Asn	Lys	Val	Leu	Val	Tyr	Ala	Thr	Ala	Asn	Gln	Glu
625					630					635					640
Lys	Gly	Gln	Arg	Thr	Pro	Tyr	Gln	Ala	Leu	Asp	Ser	Met	Asp	Asp	Ala
				645						650					655
Trp	Ser	Phe	Arg	Glu	Leu	Lys	Ala	Phe	Val	Arg	Glu	Ser	Lys	Thr	Leu
			660							665					670
Ser	Asn	Lys	Lys	Lys	Glu	Tyr	Leu	Leu	Thr	Glu	Glu	Asp	Ile	Ser	Lys
										680					685
Phe	Asp	Val	Arg	Lys	Lys	Phe	Ile	Glu	Arg	Asn	Leu	Val	Asp	Thr	Arg
										695					700
Tyr	Ala	Ser	Arg	Val	Val	Leu	Asn	Ala	Leu	Gln	Glu	His	Phe	Arg	Ala
										710					720
His	Lys	Ile	Asp	Thr	Lys	Val	Ser	Val	Val	Arg	Gly	Gln	Phe	Thr	Ser
										725					735
Gln	Leu	Arg	Arg	His	Trp	Gly	Ile	Glu	Lys	Thr	Arg	Asp	Thr	Tyr	His
										740					750
His	His	Ala	Val	Asp	Ala	Leu	Ile	Ile	Ala	Ala	Ser	Ser	Gln	Leu	Asn
										755					765
Leu	Trp	Lys	Lys	Gln	Lys	Asn	Thr	Leu	Val	Ser	Tyr	Ser	Glu	Asp	Gln
										770					780
Leu	Leu	Asp	Ile	Glu	Thr	Gly	Glu	Leu	Ile	Ser	Asp	Asp	Glu	Tyr	Lys
										785					800
Glu	Ser	Val	Phe	Lys	Ala	Pro	Tyr	Gln	His	Phe	Val	Asp	Thr	Leu	Lys
										805					815
Ser	Lys	Glu	Phe	Glu	Asp	Ser	Ile	Leu	Phe	Ser	Tyr	Gln	Val	Asp	Ser
										820					830
Lys	Phe	Asn	Arg	Lys	Ile	Ser	Asp	Ala	Thr	Ile	Tyr	Ala	Thr	Arg	Gln
										835					845
Ala	Lys	Val	Gly	Lys	Asp	Lys	Ala	Asp	Glu	Thr	Tyr	Val	Leu	Gly	Lys
										850					860
Ile	Lys	Asp	Ile	Tyr	Thr	Gln	Asp	Gly	Tyr	Asp	Ala	Phe	Met	Lys	Ile
										865					880
Tyr	Lys	Lys	Asp	Lys	Ser	Lys	Phe	Leu	Met	Tyr	Arg	His	Asp	Pro	Gln
										885					895
Thr	Phe	Glu	Lys	Val	Ile	Glu	Pro	Ile	Leu	Glu	Asn	Tyr	Pro	Asn	Lys
										900					910
Gln	Ile	Asn	Glu	Lys	Gly	Lys	Glu	Val	Pro	Cys	Asn	Pro	Phe	Leu	Lys
										915					925
Tyr	Lys	Glu	Glu	His	Gly	Tyr	Ile	Arg	Lys	Tyr	Ser	Lys	Lys	Gly	Asn
										930					940
Gly	Pro	Glu	Ile	Lys	Ser	Leu	Lys	Tyr	Tyr	Asp	Ser	Lys	Leu	Gly	Asn
										945					960
His	Ile	Asp	Ile	Thr	Pro	Lys	Asp	Ser	Asn	Asn	Lys	Val	Val	Leu	Gln
										965					975
Ser	Val	Ser	Pro	Trp	Arg	Ala	Asp	Val	Tyr	Phe	Asn	Lys	Thr	Thr	Gly
										980					990
Lys	Tyr	Glu	Ile	Leu	Gly	Leu	Lys	Tyr	Ala	Asp	Leu	Gln	Phe	Glu	Lys
										995					1005
Gly	Thr	Gly	Thr	Tyr	Lys	Ile	Ser	Gln	Glu	Lys	Tyr	Asn	Asp	Ile	
										1010					1020

-continued

Lys	Lys	Lys	Glu	Gly	Val	Asp	Ser	Asp	Ser	Glu	Phe	Lys	Phe	Thr
1025						1030					1035			
Leu	Tyr	Lys	Asn	Asp	Leu	Leu	Leu	Val	Lys	Asp	Thr	Glu	Thr	Lys
1040						1045					1050			
Glu	Gln	Gln	Leu	Phe	Arg	Phe	Leu	Ser	Arg	Thr	Met	Pro	Lys	Gln
1055						1060					1065			
Lys	His	Tyr	Val	Glu	Leu	Lys	Pro	Tyr	Asp	Lys	Gln	Lys	Phe	Glu
1070						1075					1080			
Gly	Gly	Glu	Ala	Leu	Ile	Lys	Val	Leu	Gly	Asn	Val	Ala	Asn	Ser
1085						1090					1095			
Gly	Gln	Cys	Lys	Lys	Gly	Leu	Gly	Lys	Ser	Asn	Ile	Ser	Ile	Tyr
1100						1105					1110			
Lys	Val	Arg	Thr	Asp	Val	Leu	Gly	Asn	Gln	His	Ile	Ile	Lys	Asn
1115						1120					1125			
Glu	Gly	Asp	Lys	Pro	Lys	Leu	Asp	Phe	Lys	Arg	Pro	Ala	Ala	Thr
1130						1135					1140			
Lys	Lys	Ala	Gly	Gln	Ala	Lys	Lys	Lys	Lys					
1145						1150								

<210> SEQ ID NO 53
 <211> LENGTH: 340
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 53

```

gagggcctat ttcccatgat tccttcatat ttgcatatac gatacaaggc tgtagagag      60
ataattggaa ttaatttgac tgtaaacaca aagatattag tacaaaatac gtgacgtaga      120
aagtaataat ttcttgggta gtttgcagtt ttaaaattat gttttaaata ggactatcat      180
atgcttacgc taacttgaaa gtatttogat ttcttggcctt tatatatctt gtggaagga      240
cgaaacaccg ttacttaaat ctgcagaag ctacaaagat aaggcttcat gccgaaatca      300
acaccctgtc attttatggc aggggtgttt cgttatntaa      340
    
```

<210> SEQ ID NO 54
 <211> LENGTH: 360
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide
 <220> FEATURE:
 <221> NAME/KEY: modified_base
 <222> LOCATION: (288)..(317)
 <223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 54

```

gagggcctat ttcccatgat tccttcatat ttgcatatac gatacaaggc tgtagagag      60
ataattggaa ttaatttgac tgtaaacaca aagatattag tacaaaatac gtgacgtaga      120
aagtaataat ttcttgggta gtttgcagtt ttaaaattat gttttaaata ggactatcat      180
atgcttacgc taacttgaaa gtatttogat ttcttggcctt tatatatctt gtggaagga      240
cgaaacaccg gggttttagag ctatgctgtt ttgaatggtc ccaaaaacnnn mnnnnnnnnn      300
    
```

-continued

```
nnnnnnnnnn nnnnnnngtt ttagagctat gctgttttga atggtcccaa aacttttttt 360
```

```
<210> SEQ ID NO 55
<211> LENGTH: 318
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      polynucleotide
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (250)..(269)
<223> OTHER INFORMATION: a, c, t, g, unknown or other
```

```
<400> SEQUENCE: 55
gagggcctat ttcccatgat tccttcatat ttgcatatac gatacaaggc tgtagagag 60
ataattggaa ttaatttgac tgtaaacaca aagatattag tacaaaatac gtgacgtaga 120
aagtaataat ttcttgggta gtttgcagtt ttaaaattat gttttaaata ggactatcat 180
atgcttacgc taacttgaaa gtatttcgat ttcttggcctt tatatatcctt gtggaaggaa 240
cgaaacaccc nnnnnnnnnn nnnnnnnnng ttttagagct agaaatagca agttaaata 300
aggctagtcc gttttttt 318
```

```
<210> SEQ ID NO 56
<211> LENGTH: 325
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      polynucleotide
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (250)..(269)
<223> OTHER INFORMATION: a, c, t, g, unknown or other
```

```
<400> SEQUENCE: 56
gagggcctat ttcccatgat tccttcatat ttgcatatac gatacaaggc tgtagagag 60
ataattggaa ttaatttgac tgtaaacaca aagatattag tacaaaatac gtgacgtaga 120
aagtaataat ttcttgggta gtttgcagtt ttaaaattat gttttaaata ggactatcat 180
atgcttacgc taacttgaaa gtatttcgat ttcttggcctt tatatatcctt gtggaaggaa 240
cgaaacaccc nnnnnnnnnn nnnnnnnnng ttttagagct agaaatagca agttaaata 300
aggctagtcc gttatcattt ttttt 325
```

```
<210> SEQ ID NO 57
<211> LENGTH: 337
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      polynucleotide
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (250)..(269)
<223> OTHER INFORMATION: a, c, t, g, unknown or other
```

```
<400> SEQUENCE: 57
gagggcctat ttcccatgat tccttcatat ttgcatatac gatacaaggc tgtagagag 60
ataattggaa ttaatttgac tgtaaacaca aagatattag tacaaaatac gtgacgtaga 120
aagtaataat ttcttgggta gtttgcagtt ttaaaattat gttttaaata ggactatcat 180
```

-continued

```

atgcttacgc taacttgaaa gtatttcgat ttcttggtt tataatcctt gtggaaagga 240
cgaaacaccc nnnnnnnnnn nnnnnnnnng ttttagagct agaaatagca agttaaata 300
aggctagtcc gttatcaact tgaaaaagtg tttttttt 337

```

```

<210> SEQ ID NO 58
<211> LENGTH: 352
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      polynucleotide
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (250)..(269)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

```

```

<400> SEQUENCE: 58
gagggcctat ttcccatgat tcttcacat ttgcatatac gatacaaggc tgtagagag 60
ataattggaa ttaatttgac tgtaaacaca aagatattag taaaaatac gtgacgtaga 120
aagtaataat ttcttggtgta gtttgagtt ttaaaattat gttttaaata ggactatcat 180
atgcttacgc taacttgaaa gtatttcgat ttcttggtt tataatcctt gtggaaagga 240
cgaaacaccc nnnnnnnnnn nnnnnnnnng ttttagagct agaaatagca agttaaata 300
aggctagtcc gttatcaact tgaaaaagtg gcaccgagtc ggtgcttttt tt 352

```

```

<210> SEQ ID NO 59
<211> LENGTH: 5101
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      polynucleotide

```

```

<400> SEQUENCE: 59
cgttacataa cttacggtaa atggcccgcg tggtgaccg cccaacgacc cccgcccatt 60
gacgtcaata atgacgtatg ttcccatagt aacccaata gggactttcc attgacgtca 120
atgggtggag tatttacggt aaactgccca ctggcagta catcaagtgt atcatatgcc 180
aagtacgccc cctattgacg tcaatgacgg taaatggccc gcctggcatt atgcccagta 240
catgacctta tgggactttc ctacttgcca gtacatctac gtattagta tcgctattac 300
catggtcgag gtgagcccca cgttctgctt cactctcccc atctcccccc cctccccacc 360
cccaattttg tatttattta ttttttaatt attttggtgca gcgatggggg cggggggggg 420
ggggggggcg cgcgccagggc gggcgggggc gggcgagggg cggggcgggg cgaggcggag 480
agggtcggcg gcagccaatc agagcggcgc gctccgaaag tttcctttta tggcgaggcg 540
gcggcggcgg cggccctata aaaagcgaag cgcgcggcgg gcgggagtcg ctgacgacgt 600
gccttcgccc cgtgccccgc tccgcgcgcg cctcgcgcgc cccgccccgg ctctgactga 660
ccgcgttact cccacaggtg agcgggcccg acggcccttc tcctccgggc tgtaattagc 720
tgagcaagag gtaagggttt aagggatggt tggttggtgg ggtattaatg ttaattacc 780
tggagcacct gcctgaaate acttttttcc aggttggtacc ggtgccacca tggactataa 840
ggaccacgac ggagactaca aggatcatga tattgattac aaagacgatg acgataagat 900
ggccccaag aagaagcggg aggtcgggat ccacggagtc ccagcagccc acaagaagta 960

```

-continued

cagcatcggc ctggacatcg gcaccaactc tgtgggctgg gccgtgatca ccgacgagta 1020
caaggtgccc agcaagaaat tcaaggtgct gggcaacacc gaccggcaca gcatcaagaa 1080
gaacctgatc ggagccctgc tgttcgacag cggcgaaaca gccgaggcca cccggctgaa 1140
gagaaccgcc agaagaagat acaccagacg gaagaaccgg atctgctatc tgcaagagat 1200
cttcagcaac gagatggcca agtgggacga cagcttcttc cacagactgg aagagtctt 1260
cctggtggaa gaggataaga agcaccagacg gcacccatc ttcggcaaca tcgtggacga 1320
ggtggcctac caccgagaagt accccaccat ctaccacctg agaaagaaac tggtgacag 1380
caccgacaag gccgacctgc ggctgatcta tctggccctg gccacatga tcaagttccg 1440
gggccacttc ctgatcagag ggcacctgaa ccccgacaac agcgacctgg acaagctgtt 1500
catccagctg gtgcagacct acaaccagct gttcagggaa aaccccatca acccagcgg 1560
cgtggacgcc aaggccatcc tgtctgccag actgagcaag agcagacggc tggaaaatct 1620
gatcggcccag ctgcccggcg agaagaagaa tggcctgttc ggcaacctga ttgccctgag 1680
cctgggctcg acccccactc tcaagagcaa cttegacctg gccgaggatg ccaaactgca 1740
gctgagcaag gacacctacg acgacgacct ggacaacctg ctggcccaga tcggcgacca 1800
gtacgcccag ctgtttctgg ccgccaagaa cctgtccgac gccatcctgc tgagcgacat 1860
cctgagagtg aacaccgaga tcaccaaggc ccccctgagc gcctctatga tcaagagata 1920
cgacgagcac caccaggacc tgacctgct gaaagctctc gtgcggcagc agctgcctga 1980
gaagtacaaa gagatcttct tcgaccagag caagaacggc tacgcccggc acattgacgg 2040
cggagccagc caggaagagt tctacaagtt catcaagccc atcctggaaa agatggacgg 2100
caccgaggaa ctgctcgtga agctgaacag agaggacctg ctgcggaagc agcggacctt 2160
cgacaacggc agcatcccc accagatcca cctgggagag ctgcacgcca ttctgcccgg 2220
gcaggaagat ttttaccatc tctgaagga caaccgggaa aagatcgaga agatcctgac 2280
cttccgcatc ccctactacg tggccctctt ggccagggga aacagcagat tcgctggat 2340
gaccgaaaag agcagggaaa ccatacccc ctggaacttc gaggaagtgg tgacaagggg 2400
cgcttccgcc cagagcttca tcgagcggat gaccaacttc gataagaacc tgcccacga 2460
gaagtgctg cccaagcaca gcctgctgta cgagtacttc accgtgtata acgagctgac 2520
caaagtgaat tacgtgaccg agggaaatgag aaagcccgcc ttctgagcg gcgagcagaa 2580
aaaggccatc gtggacctgc tgttcaagac caaccggaaa gtgaccgtga agcagctgaa 2640
agaggactac ttcaagaaaa tcgagtgctt cgactccgtg gaaatctccg gcgtggaaga 2700
tcggttcaac gcctccctgg gcacatacca cgatctgctg aaaattatca aggacaagga 2760
cttctggac aatgaggaaa acgaggacat tctggaagat atcgtgctga cctgacact 2820
gtttgaggac agagagatga tcgaggaacg gctgaaaacc tatgcccacc tgttcgacga 2880
caaagtgatg aagcagctga agcggcggag atacaccggc tggggcagggc tgagccggaa 2940
gctgatcaac ggcatccggg acaagcagtc cggcaagaca atcctggatt tcctgaagtc 3000
cgacggcttc gccaacagaa acttcatgca gctgatccac gacgacagcc tgacctttaa 3060
agaggacatc cagaagccc agtggtccgg ccagggcgat agcctgcacg agcaccattgc 3120
caatctggcc ggcagccccg ccattaagaa gggcatcctg cagacagtga agtggtgga 3180
cgagctcgtg aaagtgatgg gccggcacia gcccgagaac atcgtgatcg aaatggccag 3240

-continued

```

agagaaccag accaccaga agggacagaa gaacagccgc gagagaatga agcggatcga 3300
agagggcatc aaagagctgg gcagccagat cctgaaagaa caccocgtgg aaaacacca 3360
gctgcagaac gagaagctgt acctgtacta cctgcagaat gggcgggata tgtacttgga 3420
ccaggaactg gacatcaacc ggtgtccga ctacgatgtg gaccatatcg tgcctcagag 3480
ctttctgaag gacgactcca tcgacaacaa ggtgctgacc agaagcgaca agaaccgggg 3540
caagagcgac aacgtgcocct ccgaagaggt cgtgaagaag atgaagaact actggcggca 3600
gctgctgaac gccaaactga ttaccagag aaagtctgac aatctgacca aggccgagag 3660
aggcggcctg agcgaactgg ataagcccg cttcatcaag agacagctgg tggaaacccg 3720
gcagatcaca aagcacgtgg cacagatcct ggactcccg atgaacacta agtacgcga 3780
gaatgacaag ctgatccggg aagtgaagt gatcaccctg aagtccaagc tgggtgccga 3840
tttccggaag gatttccagt ttacaaaagt gcgcgagatc aacaactacc accacgcca 3900
cgacgcctac ctgaacccg tcgtgggaac cgccctgatc aaaaagtacc ctaagctgga 3960
aagcgagttc gtgtacggcg actacaaggt gtacgacgtg cggaagatga tcgccaagag 4020
cgagcaggaa atcggcaagg ctaccgcaa gtacttcttc tacagcaaca tcatgaactt 4080
tttcaagacc gagattacc tggccaacgg cgagatccgg aagcggcctc tgatcgagac 4140
aaacggcgaa accggggaga tcgtgtggga taagggccgg gattttgcca ccgtgcgaa 4200
agtgtgagc atgccccaa gtaatatcgt gaaaaagacc gaggtgcaga caggcggctt 4260
cagcaaagag tctatcctgc ccaagaggaa cagcgataag ctgatcgcca gaaagaagga 4320
ctgggaccct aagaagtacg gcggttcga cagcccccacc gtggcctatt ctgtgctggt 4380
ggtggccaaa gtggaaaagg gcaagtcca gaaactgaag agtgtgaaag agctgctggg 4440
gatcaccatc atgaaaagaa gcagcttcga gaagaatccc atcgactttc tggaagccaa 4500
gggtacaaa gaagtgaaaa aggacctgat catcaagctg cctaagtact ccctgttcga 4560
gctggaaaac ggccggaaga gaatgctggc ctctgccggc gaactgcaga agggaaacga 4620
actggccctg ccctccaaat atgtgaactt cctgtacctg gccagccact atgagaagct 4680
gaagggctcc cccgaggata atgagcagaa acagctgttt gtggaacagc acaagcacta 4740
cctggacgag atcatcgagc agatcagcga gttctccaag agagtgatcc tggccgacgc 4800
taatctggac aaagtgtgt cccgctacaa caagcaccgg gataagccca tcagagagca 4860
ggccgagaat atcatccacc tgtttaccct gaccaatctg ggagcccctg ccgcttcaa 4920
gtactttgac accaccatcg accggaagag gtacaccagc accaaagagg tgctggacgc 4980
caccctgatc caccagagca tcaccggcct gtacgagaca cggatcgacc tgtctcagct 5040
gggagggcag tttcttttct ttagcttgac cagctttctt agtagcagca ggacgcttta 5100
a 5101

```

```

<210> SEQ ID NO 60
<211> LENGTH: 137
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      polynucleotide
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (1)..(20)

```

-continued

<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 60

nnnnnnnnnn nnnnnnnnnn gttattgtac tctcaagatt tagaaataaa tcttgcagaa 60

gctacaaaga taaggcttca tgccgaaatc aacaccctgt cattttatgg caggggtgtt 120

tcggtattta atttttt 137

<210> SEQ ID NO 61

<211> LENGTH: 123

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<220> FEATURE:

<221> NAME/KEY: modified_base

<222> LOCATION: (1)..(20)

<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 61

nnnnnnnnnn nnnnnnnnnn gttattgtac tctcagaaat gcagaagcta caaagataag 60

gcttcatgcc gaaatcaaca cctgtcatt ttatggcagg gtgttttctg tatttaattt 120

ttt 123

<210> SEQ ID NO 62

<211> LENGTH: 110

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<220> FEATURE:

<221> NAME/KEY: modified_base

<222> LOCATION: (1)..(20)

<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 62

nnnnnnnnnn nnnnnnnnnn gttattgtac tctcagaaat gcagaagcta caaagataag 60

gcttcatgcc gaaatcaaca cctgtcatt ttatggcagg gtgttttttt 110

<210> SEQ ID NO 63

<211> LENGTH: 137

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<220> FEATURE:

<221> NAME/KEY: modified_base

<222> LOCATION: (1)..(20)

<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 63

nnnnnnnnnn nnnnnnnnnn gttattgtac tctcaagatt tagaaataaa tcttgcagaa 60

gctacaatga taaggcttca tgccgaaatc aacaccctgt cattttatgg caggggtgtt 120

tcggtattta atttttt 137

<210> SEQ ID NO 64

<211> LENGTH: 123

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

-continued

```

<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      polynucleotide
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (1)..(20)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 64

nnnnnnnnnn nnnnnnnnnn gttattgtac tctcagaaat gcagaagcta caatgataag      60
gcttcatgcc gaaatcaaca cctctgcatt ttatggcagg gtgttttctg tatttaattt    120
ttt                                                                    123

<210> SEQ ID NO 65
<211> LENGTH: 110
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      polynucleotide
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (1)..(20)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 65

nnnnnnnnnn nnnnnnnnnn gttattgtac tctcagaaat gcagaagcta caatgataag      60
gcttcatgcc gaaatcaaca cctctgcatt ttatggcagg gtgttttttt                110

<210> SEQ ID NO 66
<211> LENGTH: 107
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      polynucleotide
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (1)..(20)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 66

nnnnnnnnnn nnnnnnnnnn gttttagagc tgtggaaaca cagcgagtta aaataaggct      60
tagtccgtac tcaactttaa aaggtggcac cgattcggty ttttttt                107

<210> SEQ ID NO 67
<211> LENGTH: 4263
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      polynucleotide

<400> SEQUENCE: 67

atgaaaaggc cggcggccac gaaaaaggcc ggcaggcaa aaaagaaaa gaccaagccc      60
tacagcatcg gcctggacat cggcaccaat agcgtgggct gggcctgac caccgacaac    120
tacaaggctg ccagcaagaa aatgaaggty ctgggcaaca cctccaagaa gtacatcaag    180
aaaaacctgc tggcgtgct gctgttcgac agcggcatta cagccgagg cagacggctg    240
aagagaaccg ccagacggcg gtacaccggy cggagaaaca gaatcctgta tctgcaagag    300
atcttcagca ccgagatggc taccctggac gacgccttct tccagcggct ggacgacagc    360

```

-continued

ttcctggtgc	ccgacgacaa	gcgggacagc	aagtacccca	tcttcggcaa	cctggtggaa	420
gagaaggcct	accacgaoga	gttccccacc	atctaccacc	tgagaaagta	cctggccgac	480
agcaccaaga	aggccgacct	gagactggtg	tatctggccc	tggccacat	gatcaagtac	540
cggggccact	tcctgatoga	gggaggttc	aacagcaaga	acaacgacat	ccagaagaac	600
ttccaggact	tcctggacac	ctacaacgcc	atcttcgaga	gcgacctgtc	cctggaaaaac	660
agcaagcagc	tggaagagat	cgtgaaggac	aagatcagca	agctggaaaa	gaaggaccgc	720
atcctgaagc	tgttccccgg	cgagaagaac	agcggaatct	tcagcgagtt	tctgaagctg	780
atcgtgggca	accaggccga	cttcagaaag	tgcttcaacc	tggacgagaa	agccagcctg	840
cacttcagca	aagagagcta	cgacgaggac	ctggaacccc	tgctgggata	tatcggcgac	900
gactacagcg	acgtgttctc	gaaggccaag	aagctgtacg	acgtatcctc	gctgagcggc	960
ttcctgacog	tgaccgacaa	cgagacagag	gccccactga	gcagcgccat	gattaagcgg	1020
tacaacgagc	acaagagga	tctggctctg	ctgaaagagt	acatccggaa	catcagcctg	1080
aaaacctaca	atgaggtggt	caaggacgac	accaagaacg	gctacgccgg	ctacatcgac	1140
ggcaagacca	accaggaaga	tttctatgtg	tacctgaaga	agctgctggc	cgagttcgag	1200
ggggccgact	actttctgga	aaaaatcgac	cgcgaggatt	tcctgcgga	gcagcggacc	1260
ttcgacaacg	gcagcatccc	ctaccagatc	catctgcagg	aaatgcgggc	catcctggac	1320
aagcaggcca	agttctaccc	attcctggcc	aagaacaaag	agcggatcga	gaagatcctg	1380
accttcgca	tcccttacta	cgtgggcccc	ctggccagag	gcaacagcga	ttttgctgg	1440
tccatccgga	agcgaatga	gaagatcacc	ccttggaaact	tcgaggacgt	gatcgacaaa	1500
gagtccagcg	ccgaggcctt	catcaaccgg	atgaccagct	tcgacctgta	cctgcccagag	1560
gaaaaggtgc	tgcccaagca	cagcctgctg	tacgagacat	tcaatgtgta	taacgagctg	1620
accaaagtgc	ggtttatcgc	cgagtctatg	cgggactacc	agttcctgga	ctccaagcag	1680
aaaaaggaca	tcgtgctggc	gtacttcaag	gacaagcggg	aagtgaccga	taaggacatc	1740
atcgagtacc	tgcacgccat	ctacggctac	gatggcatcg	agctgaaagg	catcgagaag	1800
cagttcaact	ccagcctgag	cacataccac	gacctgctga	acattatcaa	cgacaaagaa	1860
tttctggacg	actccagcaa	cgaggccatc	atcgaagaga	tcatccacac	cctgaccatc	1920
tttgaggacc	gcgagatgat	caagcagcgg	ctgagcaagt	tcgagaacat	cttcgacaag	1980
agcgtgctga	aaaagctgag	cagacggcac	tacaccggct	ggggcaagct	gagcgccaag	2040
ctgatcaacg	gcattccggg	cgagaagtcc	ggcaacacaa	tcttgacta	cctgatcgac	2100
gacggcatca	gcaaccggaa	cttcatgcag	ctgatccacg	acgacgccct	gagcttcaag	2160
aagaagatcc	agaaggccca	gatcatcggg	gacgaggaca	agggcaacat	caaagaagtc	2220
gtgaagtccc	tgcccggcag	ccccgccatc	aagaagggaa	tctgcagag	catcaagatc	2280
gtggacgagc	tcgtgaaagt	gatggcgggc	agaaagcccg	agagcatcgt	ggtggaatg	2340
gctagagaga	accagtacac	caatcagggc	aagagcaaca	gccagcagag	actgaagaga	2400
ctggaaaagt	ccctgaaaga	gctgggcagc	aagattctga	aagagaatat	ccctgccaag	2460
ctgtccaaga	tcgacaacaa	cgccctgcag	aacgaccggc	tgtaacctga	ctacctgcag	2520
aatggcaagg	acatgtatac	aggcagcagc	ctggatatcg	accgctgag	caactacgac	2580
atcgaccata	ttatccccca	ggccttctctg	aaagacaaca	gcattgacaa	caaagtgctg	2640

-continued

```

gtgtcctccg ccagcaaccg cggcaagtcc gatgatgtgc ccagcctgga agtcgtgaaa 2700
aagagaaaga ccttctggta tcagctgctg aaaagcaagc tgattagcca gaggaagttc 2760
gacaacctga ccaaggccga gagaggcggc ctgagccctg aagataaggc cggcttcac 2820
cagagacagc tgggtgaaac ccggcagatc accaagcacg tggccagact gctggatgag 2880
aagtttaaca acaagaagga cgagaacaac cgggccgtgc ggaccgtgaa gatcatcacc 2940
ctgaagtcca ccctgggtgc ccagttccgg aaggacttcg agctgataa agtgcgcgag 3000
atcaatgact ttcaccacgc ccacgacgcc tacctgaatg ccgtggtggc ttcgcctctg 3060
ctgaagaagt accctaagct ggaaccgag ttcgtgtacg gcgactaccc caagtacaac 3120
tccttcagag agcgggaagtc cgccaccgag aaggtgtact tctactccaa catcatgaat 3180
atccttaaga agtccatctc cctggccgat ggcagagtga tcgagcggcc cctgatcgaa 3240
gtgaacgaag agacaggcga gagcgtgtgg aacaaagaaa gcgacctggc caccgtgagg 3300
cgggtgctga gttatcctca agtgaatgtc gtgaagaagg tggaagaaca gaaccacggc 3360
ctggatcggg gcaagcccaa gggcctgttc aacgccaaac tgtccagcaa gcctaagccc 3420
aactccaacg agaatctcgt gggggccaaa gagtaoctgg accctaagaa gtacggcgga 3480
tacgccggca tctccaatag cttcacctg ctcgtgaagg gcacaatcga gaaggcgct 3540
aagaaaaaga tcacaaaagt gctggaattt caggggatct ctatcctgga ccggatcaac 3600
taccggaagg ataagctgaa ctttctgctg gaaaaaggct acaaggacat tgagctgatt 3660
atcgagctgc ctaagtactc cctgttcgaa ctgagcagc gctccagacg gatgctggcc 3720
tccatcctgt ccaccaacaa caagcggggc gagatocaca agggaaacca gatcttctctg 3780
agccagaaat ttgtgaaact gctgtaccac gccaaagcga tctccaacac catcaatgag 3840
aaccaccgga aatacgtgga aaaccacaag aaagagtgtg aggaactggt ctactacatc 3900
ctggagttca acgagaacta tgtgggagcc aagaagaacg gcaaaactgct gaactccgcc 3960
ttccagagct ggcagaacca cagcatcgac gagctgtgca gctccttcat cggccctacc 4020
ggcagcagc ggaagggact gtttgagctg acctccagag gctctgccgc cgactttgag 4080
ttcctgggag tgaagatccc ccggtacaga gactaccccc cctctagtct gctgaaggac 4140
gccaccctga tccaccagag cgtgaccggc ctgtacgaaa cccggatcga cctggctaag 4200
ctggcgagg gaaagcgtcc tgctgctact aagaaagctg gtcaagctaa gaaaaagaaa 4260
taa 4263

```

```

<210> SEQ ID NO 68
<211> LENGTH: 53
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

```

```

<400> SEQUENCE: 68

```

```

tcctagcagg atttctgata ttactgtcac gttttagagc tatgctgttt tga 53

```

```

<210> SEQ ID NO 69
<211> LENGTH: 53
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

```

-continued

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 69

gtgacagtaa tatcagaat cctgctagga gttttgggac cattcaaac agc 53

<210> SEQ ID NO 70
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 70

gggtttcaag tctttgtagc aagag 25

<210> SEQ ID NO 71
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 71

gccaatgaac gggaaacctt ggtc 24

<210> SEQ ID NO 72
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (1)..(4)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 72

nnnngacgag gcaatggctg aaatc 25

<210> SEQ ID NO 73
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (1)..(4)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 73

nnnnttattt ggctcatatt tgctg 25

<210> SEQ ID NO 74
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

-continued

<400> SEQUENCE: 74

ctttacacca atcgctgcaa cagac 25

<210> SEQ ID NO 75

<211> LENGTH: 45

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 75

caaaatttct agtcttcttt gcctttcccc ataaaaccct cctta 45

<210> SEQ ID NO 76

<211> LENGTH: 45

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 76

agggttttat ggggaaaggc aaagaagact agaaattttg atacc 45

<210> SEQ ID NO 77

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 77

cttacggtgc ataaagtcaa ttcc 25

<210> SEQ ID NO 78

<211> LENGTH: 21

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 78

tggtcgcatt tcagccattg c 21

<210> SEQ ID NO 79

<211> LENGTH: 43

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<220> FEATURE:

<221> NAME/KEY: modified_base

<222> LOCATION: (33)..(33)

<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 79

ctttgacgag gcaatggctg aaatcgagcc aanaaagcgc aag 43

<210> SEQ ID NO 80

<211> LENGTH: 43

-continued

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (34)..(34)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 80

ctttgacgag gcaatggctg aaatcgagcc aaanaagcgc aag 43

<210> SEQ ID NO 81
<211> LENGTH: 43
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (35)..(35)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 81

ctttgacgag gcaatggctg aaatcgagcc aaaaagcgc aag 43

<210> SEQ ID NO 82
<211> LENGTH: 43
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (36)..(36)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 82

ctttgacgag gcaatggctg aaatcgagcc aaaaangcgc aag 43

<210> SEQ ID NO 83
<211> LENGTH: 43
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (37)..(37)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 83

ctttgacgag gcaatggctg aaatcgagcc aaaaaancgc aag 43

<210> SEQ ID NO 84
<211> LENGTH: 43
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (38)..(38)

-continued

<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 84

ctttgacgag gcaatggctg aaatcgagcc aaaaaagngc aag 43

<210> SEQ ID NO 85
<211> LENGTH: 46
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (39)..(39)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 85

ctttgacgag gcaatggctg aaatcgagcc aaaaaagcnc aagaag 46

<210> SEQ ID NO 86
<211> LENGTH: 46
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (40)..(40)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 86

ctttgacgag gcaatggctg aaatcgagcc aaaaaagcgn aagaag 46

<210> SEQ ID NO 87
<211> LENGTH: 46
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (41)..(41)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 87

ctttgacgag gcaatggctg aaatcgagcc aaaaaagcgc nagaag 46

<210> SEQ ID NO 88
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 88

gcgctttttt ggctcgattt cag 23

<210> SEQ ID NO 89
<211> LENGTH: 40
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (31)..(31)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 89

caatggctga aatcgagcca aaaaagcgca ngaagaaatc 40

<210> SEQ ID NO 90
<211> LENGTH: 40
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (32)..(32)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 90

caatggctga aatcgagcca aaaaagcgca anaagaaatc 40

<210> SEQ ID NO 91
<211> LENGTH: 40
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (33)..(33)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 91

caatggctga aatcgagcca aaaaagcgca agnagaaatc 40

<210> SEQ ID NO 92
<211> LENGTH: 40
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (34)..(34)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 92

caatggctga aatcgagcca aaaaagcgca agangaaatc 40

<210> SEQ ID NO 93
<211> LENGTH: 40
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (35)..(35)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 93

-continued

caatggctga aatcgagcca aaaaagcgca agaanaaatc 40

<210> SEQ ID NO 94
<211> LENGTH: 44
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (36)..(36)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 94

caatggctga aatcgagcca aaaaagcgca agaagnaatc aacc 44

<210> SEQ ID NO 95
<211> LENGTH: 44
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (37)..(37)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 95

caatggctga aatcgagcca aaaaagcgca agaaganatc aacc 44

<210> SEQ ID NO 96
<211> LENGTH: 44
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (38)..(38)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 96

caatggctga aatcgagcca aaaaagcgca agaagaantc aacc 44

<210> SEQ ID NO 97
<211> LENGTH: 44
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (39)..(39)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 97

caatggctga aatcgagcca aaaaagcgca agaagaaanc aacc 44

<210> SEQ ID NO 98
<211> LENGTH: 47
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (40)..(40)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 98

caatggctga aatcgagcca aaaaagcgca agaagaaatn aaccgc 47

<210> SEQ ID NO 99
<211> LENGTH: 47
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (41)..(41)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 99

caatggctga aatcgagcca aaaaagcgca agaagaaatc naccgc 47

<210> SEQ ID NO 100
<211> LENGTH: 31
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 100

gatcctccat cgtacaacc cacaacctg g 31

<210> SEQ ID NO 101
<211> LENGTH: 31
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 101

aattccaggg ttgtgggtg tacggatgga g 31

<210> SEQ ID NO 102
<211> LENGTH: 34
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 102

catggatcct atttcttaat aactaaaaat atgg 34

<210> SEQ ID NO 103
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

-continued

<400> SEQUENCE: 103

catgaattca actcaacaag tctcagtgtg ctg 33

<210> SEQ ID NO 104

<211> LENGTH: 35

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 104

aaacattttt tctccattta ggaaaaagga tgctg 35

<210> SEQ ID NO 105

<211> LENGTH: 35

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 105

aaaacagcat cctttttcct aaatggagaa aaaat 35

<210> SEQ ID NO 106

<211> LENGTH: 35

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 106

aaaccttaaa tcagtcacaa atagcagcaa aattg 35

<210> SEQ ID NO 107

<211> LENGTH: 35

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 107

aaaacaattt tgetgctatt tgtgactgat ttaag 35

<210> SEQ ID NO 108

<211> LENGTH: 35

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 108

aaacttttca tcatacgacc aatctgcttt atttg 35

<210> SEQ ID NO 109

<211> LENGTH: 35

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

-continued

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 109

aaaacaaata aagcagattg gtcgtatgat gaaaa 35

<210> SEQ ID NO 110
<211> LENGTH: 35
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 110

aaactcgtcc agaagttatc gtaaaagaaa tcgag 35

<210> SEQ ID NO 111
<211> LENGTH: 35
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 111

aaaactcgat ttcttttacg ataacttctg gacga 35

<210> SEQ ID NO 112
<211> LENGTH: 35
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 112

aaacaatctc tccaagggtt ccttaaaaat ctctg 35

<210> SEQ ID NO 113
<211> LENGTH: 35
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 113

aaaacagaga tttttaagga aaccttgag agatt 35

<210> SEQ ID NO 114
<211> LENGTH: 35
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 114

aaacgccatc gtcaggaaga agctatgctt gagtg 35

<210> SEQ ID NO 115
<211> LENGTH: 35
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 115

aaaacactca agcatagctt cttcctgacg atggc 35

<210> SEQ ID NO 116
<211> LENGTH: 35
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 116

aaacatctct atacttattg aaatttcttt gtatg 35

<210> SEQ ID NO 117
<211> LENGTH: 35
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 117

aaaacataca aagaaatttc aataagtata gagat 35

<210> SEQ ID NO 118
<211> LENGTH: 35
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 118

aaactagctg tgatagtccg caaaaccagc cttcg 35

<210> SEQ ID NO 119
<211> LENGTH: 35
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 119

aaaacgaagg ctggttttgc ggactatcac agcta 35

<210> SEQ ID NO 120
<211> LENGTH: 35
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 120

aaacatcgga aggtcgagca agtaattatc ttttg 35

<210> SEQ ID NO 121

-continued

<211> LENGTH: 35
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 121

aaaacaaaag ataattactt gctcgacctt ccgat 35

<210> SEQ ID NO 122
<211> LENGTH: 35
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 122

aaacaagatg gtatcgcaaa gtaagtgaca ataag 35

<210> SEQ ID NO 123
<211> LENGTH: 35
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 123

aaaacttatt gtcacttact ttgcgatacc atctt 35

<210> SEQ ID NO 124
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 124

gagacctttg agcttccgag actggtctca gttttgggac cattcaaac ag 52

<210> SEQ ID NO 125
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 125

tgagaccagt ctcggaagct caaaggtctc gttttagagc tatgctgttt tg 52

<210> SEQ ID NO 126
<211> LENGTH: 35
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 126

aaactacttt acgcagcgcg gagttcgggt ttttg 35

-continued

<210> SEQ ID NO 127
<211> LENGTH: 35
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 127

aaaacaaaa accgaactcc gcgctgcgta aagta 35

<210> SEQ ID NO 128
<211> LENGTH: 45
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 128

atgccggtac tgccgggcct cttgcgggat tacgaaatca tcctg 45

<210> SEQ ID NO 129
<211> LENGTH: 45
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 129

gtgactggcg atgctgtogg aatggacgat cacactactc ttctt 45

<210> SEQ ID NO 130
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 130

ttaagaaata atcttcatct aaaatatact tcagtcacct cctagctgac 50

<210> SEQ ID NO 131
<211> LENGTH: 48
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 131

attgatttga gtcagctagg aggtgactga agtatatfff agatgaag 48

<210> SEQ ID NO 132
<211> LENGTH: 85
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 132

-continued

gagacctttg agcttccgag actggtctca gttttgggac cattcaaac agcatagctc 60

taaaacctcg tagactatctt ttgtc 85

<210> SEQ ID NO 133
<211> LENGTH: 84
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 133

gagaccagtc tcggaagctc aaaggtctcg ttttagagct atgctgtttt gaatgtccc 60

aaaacttcag cacactgaga cttg 84

<210> SEQ ID NO 134
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 134

agtcaccca gcaacaaatg g 21

<210> SEQ ID NO 135
<211> LENGTH: 31
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 135

cgtagtaaat cggataacgt tccaagtga g 31

<210> SEQ ID NO 136
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 136

tgctcttctt cacaaacaag gg 22

<210> SEQ ID NO 137
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 137

aagccaaagt ttggcaccac c 21

<210> SEQ ID NO 138
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 138

gtagcttatt cagtcctagt gg 22

<210> SEQ ID NO 139
<211> LENGTH: 45
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 139

cgtttggtga actaatgggt gcaaattacg aatcttctcc tgacg 45

<210> SEQ ID NO 140
<211> LENGTH: 45
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 140

cgtcaggaga agattcgtaa tttgcacca ttagttcaac aaacg 45

<210> SEQ ID NO 141
<211> LENGTH: 48
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 141

gatattatgg agcctatddd tgtgggtddd taggcataaa actatatg 48

<210> SEQ ID NO 142
<211> LENGTH: 48
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 142

catatagttt tatgcctaaa aaccacaaa aataggctcc ataatatc 48

<210> SEQ ID NO 143
<211> LENGTH: 27
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 143

attatttttt aataactaaa aatatgg 27

<210> SEQ ID NO 144
<211> LENGTH: 24

-continued

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 144

cgtgtacaat tgctagcgta cggc 24

<210> SEQ ID NO 145
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 145

gcaccggtga tcactagtcc tagg 24

<210> SEQ ID NO 146
<211> LENGTH: 47
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 146

cctaggacta gtgatcaccg gtgcaaatat gagcceaata aatatat 47

<210> SEQ ID NO 147
<211> LENGTH: 44
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 147

gccgtacgct agcaattgta cacgtttggt gaactaatgg gtgc 44

<210> SEQ ID NO 148
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 148

ttcaaatttt cccatttgat tctcc 25

<210> SEQ ID NO 149
<211> LENGTH: 47
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 149

ccatattttt agttattaag aaataatacc agccatcagt cacctcc 47

-continued

<210> SEQ ID NO 150
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 150

agacgattca atagacaata agg 23

<210> SEQ ID NO 151
<211> LENGTH: 45
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 151

gttttgggac cattcaaaac agcatagctc taaaacctcg tagac 45

<210> SEQ ID NO 152
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 152

gctatgctgt tttgaatggt cccaaaacca ttattttaac acacgaggtg 50

<210> SEQ ID NO 153
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 153

gctatgctgt tttgaatggt cccaaaacgc acccattagt tcaacaaacg 50

<210> SEQ ID NO 154
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 154

aattcttttc ttcacatcatcg gtc 23

<210> SEQ ID NO 155
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 155

aagaagaat gaagattggt catg 24

-continued

<210> SEQ ID NO 156
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 156

ggtactaatc aaaatagtga ggagg 25

<210> SEQ ID NO 157
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 157

gtttttcaaa atctgcggtt gcg 23

<210> SEQ ID NO 158
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 158

aaaaattgaa aaaatggtgg aaacac 26

<210> SEQ ID NO 159
<211> LENGTH: 53
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 159

atctcgtaaa cggtatcggg ttcttttaaa gttttgggac cattcaaac agc 53

<210> SEQ ID NO 160
<211> LENGTH: 53
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 160

tttaaaagaa accgataccg tttaacgaaat gtttttagagc tatgctgttt tga 53

<210> SEQ ID NO 161
<211> LENGTH: 53
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 161

-continued

aaacgggtatc ggtttctttt aaattcaatt gttttgggac cattcaaac agc 53

<210> SEQ ID NO 162
<211> LENGTH: 53
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 162

aattgaattt aaaagaaacc gataccgttt gtttagagc tatgctgttt tga 53

<210> SEQ ID NO 163
<211> LENGTH: 39
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 163

gttccttaaa ccaaacggt atcggtttct ttaaattc 39

<210> SEQ ID NO 164
<211> LENGTH: 47
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 164

gaaaccgata cgttttgggt ttaaggaaca ggtaaagggc atttaac 47

<210> SEQ ID NO 165
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 165

cgatttcagc cattgcctcg tc 22

<210> SEQ ID NO 166
<211> LENGTH: 56
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (29)..(33)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 166

gcctttgacg aggcaatggc tgaatcggnn mnnaaaagc gcaagaagaa atcaac 56

<210> SEQ ID NO 167
<211> LENGTH: 53
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 167

tccgtacaac ccacaaccct gctagtgagc gttttgggac cattcaaac agc 53

<210> SEQ ID NO 168
<211> LENGTH: 53
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 168

gctcactagc agggttgtgg gttgtacgga gttttagagc tatgctgttt tga 53

<210> SEQ ID NO 169
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 169

ttggtgccac tcttccttct ttc 23

<210> SEQ ID NO 170
<211> LENGTH: 41
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 170

cagggttggtg ggttggtgag atggagtaa ctcccatctc c 41

<210> SEQ ID NO 171
<211> LENGTH: 41
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 171

gggagttaac tccatcgcaa caaccacaaa cctgctagt g 41

<210> SEQ ID NO 172
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 172

gtggtatcta tcgtgatgtg ac 22

<210> SEQ ID NO 173

-continued

<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 173

ttaccgaaac ggaatttadc tgc 23

<210> SEQ ID NO 174
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 174

aaagctagag ttccgcaatt gg 22

<210> SEQ ID NO 175
<211> LENGTH: 37
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 175

gtgggttgta cggattgagt taactcccat ctccttc 37

<210> SEQ ID NO 176
<211> LENGTH: 38
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 176

gatgggagtt aactcaatcc gtacaacca caaccctg 38

<210> SEQ ID NO 177
<211> LENGTH: 40
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 177

gcttcaccta ttgcagcacc aattgaccac atgaagatag 40

<210> SEQ ID NO 178
<211> LENGTH: 41
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 178

gtggtaatt ggtgctgcaa taggtgaagc taatggtgat g 41

-continued

<210> SEQ ID NO 179
<211> LENGTH: 39
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 179

ctgatttgta ttaattttga gacattatgc ttcaccttc 39

<210> SEQ ID NO 180
<211> LENGTH: 40
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 180

gcataatgtc tcaaaattaa tacaaatcag tgaaatcatg 40

<210> SEQ ID NO 181
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 181

gttttgggac cattcaaaac agcatagctc taaaacgtga cagtaatatc ag 52

<210> SEQ ID NO 182
<211> LENGTH: 53
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 182

gttttagagc tatgctgttt tgaatggctc caaaacgctc actagcaggg ttg 53

<210> SEQ ID NO 183
<211> LENGTH: 59
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 183

atactttaag cagcgcggag ttcggttttg taggagtggg agtatatata cgagtacat 59

<210> SEQ ID NO 184
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide

<400> SEQUENCE: 184

-continued

gctcactagc agggttgtgg gttgtacgga tgg 33

<210> SEQ ID NO 185
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 185

tcctagcagg atttctgata ttactgtcac tgg 33

<210> SEQ ID NO 186
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 186

tttaaaagaa accgataccg tttacgaaat tgg 33

<210> SEQ ID NO 187
<211> LENGTH: 84
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 187

ggaaccattc ataacagcat agcaagttat aataaggcta gtccggtatc aacttgaaaa 60

agtggcaccg agtcggtgct tttt 84

<210> SEQ ID NO 188
<211> LENGTH: 36
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 188

gttatagagc tatgctgtta tgaatggtcc caaaac 36

<210> SEQ ID NO 189
<211> LENGTH: 84
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 189

ggaaccattc aatacagcat agcaagttaa tataaggcta gtccggtatc aacttgaaaa 60

agtggcaccg agtcggtgct tttt 84

<210> SEQ ID NO 190
<211> LENGTH: 36
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 190

gtattagagc tatgctgat tgaatggtcc caaaac 36

<210> SEQ ID NO 191
<211> LENGTH: 103
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
polynucleotide
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (1)..(20)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 191

nnnnnnnnnn nnnnnnnnnn gttttagagc tagaaatagc aagttaaaat aaggctagtc 60

cgttatcaac ttgaaaaagt ggcaccgagt cggtgctttt ttt 103

<210> SEQ ID NO 192
<211> LENGTH: 103
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
polynucleotide
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (1)..(20)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 192

nnnnnnnnnn nnnnnnnnnn gtattagagc tagaaatagc aagttaatat aaggctagtc 60

cgttatcaac ttgaaaaagt ggcaccgagt cggtgctttt ttt 103

<210> SEQ ID NO 193
<211> LENGTH: 123
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
polynucleotide
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (1)..(20)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 193

nnnnnnnnnn nnnnnnnnnn gttttagagc tatgctggtt tggaacaaa acagcatagc 60

aagttaaaat aaggctagtc cgttatcaac ttgaaaaagt ggcaccgagt cggtgctttt 120

ttt 123

<210> SEQ ID NO 194
<211> LENGTH: 123
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
polynucleotide
<220> FEATURE:

-continued

<221> NAME/KEY: modified_base
<222> LOCATION: (1)..(20)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 194

nnnnnnnnnn nnnnnnnnnn gtattagagc tatgctgtat tggaacaat acagcatagc 60
aagttaatat aaggctagtc cgttatcaac ttgaaaaagt ggcaccgagt cggtgctttt 120
ttt 123

<210> SEQ ID NO 195
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 195

gtcacctcca atgactaggg 20

<210> SEQ ID NO 196
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 196

gacatcgatg tcctcccat tgg 23

<210> SEQ ID NO 197
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 197

gagtccgagc agaagaagaa ggg 23

<210> SEQ ID NO 198
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 198

gcgccaccgg ttgatgtgat ggg 23

<210> SEQ ID NO 199
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 199

ggggcacaga tgagaaactc agg 23

<210> SEQ ID NO 200
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 200

gtacaaacgg cagaagctgg agg 23

<210> SEQ ID NO 201
<211> LENGTH: 23
<212> TYPE: DNA

-continued

<213> ORGANISM: Homo sapiens
<400> SEQUENCE: 201
ggcagaagct ggaggaggaa ggg 23

<210> SEQ ID NO 202
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<400> SEQUENCE: 202
ggagcccttc ttcttctgct cgg 23

<210> SEQ ID NO 203
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<400> SEQUENCE: 203
gggcaaccac aaaccacga ggg 23

<210> SEQ ID NO 204
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<400> SEQUENCE: 204
gctcccatca catcaaccgg tgg 23

<210> SEQ ID NO 205
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<400> SEQUENCE: 205
gtggcgcatt gccacgaagc agg 23

<210> SEQ ID NO 206
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<400> SEQUENCE: 206
ggcagagtgc tgcttgctgc tgg 23

<210> SEQ ID NO 207
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<400> SEQUENCE: 207
gccctgcgt gggccaagc tgg 23

<210> SEQ ID NO 208
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<400> SEQUENCE: 208
gagtggccag agtccagctt ggg 23

-continued

<210> SEQ ID NO 209
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 209
ggcctcccca aagcctggcc agg 23

<210> SEQ ID NO 210
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 210
ggggccgaga ttgggtgttc agg 23

<210> SEQ ID NO 211
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 211
gtggcgagag gggccgagat tgg 23

<210> SEQ ID NO 212
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 212
gagtgcgcc gaggcggggc ggg 23

<210> SEQ ID NO 213
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 213
ggagtccgc cgagccgggg cgg 23

<210> SEQ ID NO 214
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 214
ggagaggagt gccgccgagg cgg 23

<210> SEQ ID NO 215
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 215
ccateccett ctgtgaatg 20

<210> SEQ ID NO 216
<211> LENGTH: 20

-continued

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 216
ggagattgga gacacggaga 20

<210> SEQ ID NO 217
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 217
aagcaccgac tcggtgccac 20

<210> SEQ ID NO 218
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 218
tcacctccaa tgactagggg 20

<210> SEQ ID NO 219
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 219
caagttgata acggactagc ct 22

<210> SEQ ID NO 220
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 220
agtccgagca gaagaagaag ttt 23

<210> SEQ ID NO 221
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 221
tttcaagttg ataacggact agcct 25

-continued

<210> SEQ ID NO 222
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 222

aaacagcaga ttcgcctgga 20

<210> SEQ ID NO 223
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 223

tcatacgcctc gatgaagctc 20

<210> SEQ ID NO 224
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 224

tccaaaatca agtggggcga 20

<210> SEQ ID NO 225
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 225

tgatgacctt tttggctccc 20

<210> SEQ ID NO 226
<211> LENGTH: 45
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 226

gaggaattct tttttgtty gaatatgttg gaggtttttt ggaag 45

<210> SEQ ID NO 227
<211> LENGTH: 42
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 227

gagaagctta aataaaaaac racaatactc aaccaacaa cc 42

-continued

<210> SEQ ID NO 228
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 228

caggaaacag ctatgac 17

<210> SEQ ID NO 229
<211> LENGTH: 39
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 229

gcctctagag gtacctgagg gcctatttcc catgattcc 39

<210> SEQ ID NO 230
<211> LENGTH: 133
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (92)..(111)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 230

acctctagaa aaaaagcacc gactcgggtgc cactttttca agttgataac ggactagcct 60
tattttaact tgctatttct agctctaaaa cnnnnnnnnn nnnnnnnnn nggtgtttcg 120
tcctttccac aag 133

<210> SEQ ID NO 231
<211> LENGTH: 133
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (92)..(111)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 231

acctctagaa aaaaagcacc gactcgggtgc cactttttca agttgataac ggactagcct 60
tatattaact tgctatttct agctctaata cnnnnnnnnn nnnnnnnnn nggtgtttcg 120
tcctttccac aag 133

<210> SEQ ID NO 232
<211> LENGTH: 153
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic

-continued

```

    primer
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (112)..(131)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 232
acctctagaa aaaagcacc gactcgggtgc cactttttca agttgataac ggactagcct    60
tattttaact tgctatgctg tttgtttcc aaaacagcat agctctaaaa cnnnnnnnnn    120
nnnnnnnnnn nggtgtttcg tcctttccac aag                                153

<210> SEQ ID NO 233
<211> LENGTH: 153
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
    primer
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (112)..(131)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 233
acctctagaa aaaagcacc gactcgggtgc cactttttca agttgataac ggactagcct    60
tatattaact tgctatgctg tattgtttcc aatacagcat agctctaaata cnnnnnnnnn    120
nnnnnnnnnn nggtgtttcg tcctttccac aag                                153

<210> SEQ ID NO 234
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (20)..(20)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 234
aggccccagt ggctgctctn aa                                          22

<210> SEQ ID NO 235
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (20)..(20)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 235
acatcaaccg gtggcgcacn at                                          22

<210> SEQ ID NO 236
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (20)..(20)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 236
aaggtgtggt tccagaaccn ac                                          22

```

-continued

<210> SEQ ID NO 237
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (20)..(20)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 237

ccatcacatc aaccggtggn ag 22

<210> SEQ ID NO 238
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (20)..(20)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 238

aaacggcaga agctggaggn ta 22

<210> SEQ ID NO 239
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (20)..(20)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 239

ggcagaagct ggaggaggn tt 22

<210> SEQ ID NO 240
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (20)..(20)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 240

ggtgtggttc cagaaccggn tc 22

<210> SEQ ID NO 241
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (20)..(20)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 241

aaccggagga caaagtacan tg 22

<210> SEQ ID NO 242
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

-continued

<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (20)..(20)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 242

ttccagaacc ggaggacaan ca 22

<210> SEQ ID NO 243
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (20)..(20)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 243

gtgtggttcc agaaccggn ct 22

<210> SEQ ID NO 244
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (20)..(20)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 244

tccagaaccg gaggacaaan cc 22

<210> SEQ ID NO 245
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (20)..(20)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 245

cagaagctgg aggaggaagn cg 22

<210> SEQ ID NO 246
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (20)..(20)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 246

catcaaccgg tggcgcattn ga 22

<210> SEQ ID NO 247
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (20)..(20)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 247

-continued

gcagaagctg gaggaggaan gt 22

<210> SEQ ID NO 248
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (20)..(20)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 248

cctccctccc tggcccaggn gc 22

<210> SEQ ID NO 249
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (20)..(20)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 249

tcatctgtgc ccctccctcn aa 22

<210> SEQ ID NO 250
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (20)..(20)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 250

gggaggacat cgatgtcaen at 22

<210> SEQ ID NO 251
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (20)..(20)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 251

caaacggcag aagctggagn ac 22

<210> SEQ ID NO 252
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (20)..(20)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 252

gggtgggcaa ccacaaacn ag 22

<210> SEQ ID NO 253
<211> LENGTH: 22

-continued

<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (20)..(20)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 253

ggtgggcaac cacaaaccn ta 22

<210> SEQ ID NO 254
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (20)..(20)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 254

ggctcccatc acatcaaccn tt 22

<210> SEQ ID NO 255
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (20)..(20)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 255

gaagggcctg agtccgagcn tc 22

<210> SEQ ID NO 256
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (20)..(20)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 256

caaccggtgg cgcatcgccn tg 22

<210> SEQ ID NO 257
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (20)..(20)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 257

aggaggaagg gcctgagtcn ca 22

<210> SEQ ID NO 258
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (20)..(20)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

-continued

<400> SEQUENCE: 258

agctggagga ggaagggccn ct 22

<210> SEQ ID NO 259

<211> LENGTH: 22

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: modified_base

<222> LOCATION: (20)..(20)

<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 259

gcattgccac gaagcaggcn cc 22

<210> SEQ ID NO 260

<211> LENGTH: 22

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: modified_base

<222> LOCATION: (20)..(20)

<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 260

attgccacga agcaggccan cg 22

<210> SEQ ID NO 261

<211> LENGTH: 22

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: modified_base

<222> LOCATION: (20)..(20)

<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 261

agaaccggag gacaaagtan ga 22

<210> SEQ ID NO 262

<211> LENGTH: 22

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: modified_base

<222> LOCATION: (20)..(20)

<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 262

tcaaccggtg ggcattgcn gt 22

<210> SEQ ID NO 263

<211> LENGTH: 22

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: modified_base

<222> LOCATION: (20)..(20)

<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 263

gaagctggag gaggaagggn gc 22

-continued

<210> SEQ ID NO 264
 <211> LENGTH: 123
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 264
 ccaatgggga ggacatcgat gtcacctcca atgactaggg tgggcaacca caaacccacg 60
 agggcagagt gctgcttgct gctggccagg cccctgctg ggcccaagct ggactctggc 120
 cac 123

<210> SEQ ID NO 265
 <211> LENGTH: 121
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 265
 cgagcagaag aagaagggt cccatcacat caaccggtgg cgcattgcca cgaagcaggc 60
 caatggggag gacatcgatg tcacctcaa tgactagggg gggcaaccac aaaccacga 120
 g 121

<210> SEQ ID NO 266
 <211> LENGTH: 128
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 266
 ggaggacaaa gtacaaacgg cagaagctgg aggaggaagg gcctgagtcc gagcagaaga 60
 agaagggctc ccatcacatc aaccggtggc gcattgccac gaagcaggcc aatggggagg 120
 acatcgat 128

<210> SEQ ID NO 267
 <211> LENGTH: 130
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 267
 agaagctgga ggaggaaggg cctgagtccg agcagaagaa gaagggctcc catcacatca 60
 accggtggcg cattgccacg aagcaggcca atggggagga catcgatgtc acctccaatg 120
 actaggggtgg 130

<210> SEQ ID NO 268
 <211> LENGTH: 125
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

-continued

<400> SEQUENCE: 268

ctcagcttt cccatcaggc tctcagctca gctgagtggt tgaggcccca gtggctgctc 60
 tgggggctc ctgagtttct catctgtgcc cctccctccc tggcccagggt gaaggtgtgg 120
 ttcca 125

<210> SEQ ID NO 269

<211> LENGTH: 129

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 269

tcatctgtgc cctccctccc ctggcccagg tgaaggtgtg gttccagaac cggaggacaa 60
 agtacaaaacg gcagaagctg gaggaggaag ggcctgagtc cgagcagaag aagaagggt 120
 cccatcaca 129

<210> SEQ ID NO 270

<211> LENGTH: 129

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 270

ctccaatgac tagggtgggc aaccacaaac ccacgagggc agagtgtgctc ttgctgctgg 60
 ccaggcccct gcgtgggccc aagctggact ctggcactc cctggccagg ctttggggag 120
 gcttgaggt 129

<210> SEQ ID NO 271

<211> LENGTH: 127

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 271

ctgcttgtg ctggccaggc ccttgcgtgg gcccaagctg gactctggcc actccctggc 60
 caggctttgg ggaggcctgg agtcatggcc ccacagggct tgaagcccgg ggccgccatt 120
 gacagag 127

<210> SEQ ID NO 272

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide

<400> SEQUENCE: 272

gaaattaata cgactcacta taggg 25

<210> SEQ ID NO 273

<211> LENGTH: 126

<212> TYPE: DNA

-continued

```

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      polynucleotide

<400> SEQUENCE: 273

aaaaaagcac cgactcggcg ccactttttc aagttgataa cggactagcc ttattttaac      60
ttgctatttc tagctctaaa acaacgacga gcgtgacacc accctatagt gagtcgtatt      120
aatttc                                             126

<210> SEQ ID NO 274
<211> LENGTH: 126
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      polynucleotide

<400> SEQUENCE: 274

aaaaaagcac cgactcggcg ccactttttc aagttgataa cggactagcc ttattttaac      60
ttgctatttc tagctctaaa acgcaacaat taatagactg gacctatagt gagtcgtatt      120
aatttc                                             126

<210> SEQ ID NO 275
<211> LENGTH: 4677
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      polynucleotide

<400> SEQUENCE: 275

tctttcttgc gctatgacac ttccagcaaa aggtagggcg ggctgcgaga cggcttcccg      60
gcgctgcatg caacaccgat gatgcttcca cccccgaag ctcttcggg gctgcatggg      120
cgctccgatg ccgctccagg gcgagcgtg tttaaatagc caggcccccg attgcaaaga      180
cattatagcg agctacaaa gccatattca aacacctaga tcaactaccac ttctacacag      240
gccactcgag cttgtgatcg cactccgcta agggggcgcc tcttctctt cgtttcagtc      300
acaacccgca aacatgtacc catacgatgt tccagattac gcttcgccga agaaaaagcg      360
caaggtcgaa gcgtccgaca agaagtacag catcggcctg gacatcggca ccaactctgt      420
gggctgggcc gtgatcaccg acgagtacaa ggtgcccgag aagaaattca aggtgctggg      480
caacaccgac cggcacagca tcaagaagaa cctgatcggg gccctgctgt tcgacagcgg      540
cgaaacagcc gaggcccccc ggctgaagag aaccgccaga agaagatata ccagacggaa      600
gaaccggatc tgctatctgc aagagatctt cagcaacgag atggccaagg tggacgacag      660
cttcttccac agactggaag agtccttctt ggtggaagag gataagaagc acgagcggca      720
ccccatcttc ggcaacatcg tggacgaggt ggcctaccac gagaagtacc ccaccatcta      780
ccacctgaga aagaaactgg tggacagcac cgacaaggcc gacctcggc tgatctatct      840
ggcctgggcc cacatgatca agttccgggg ccacttctct atcgagggcg acctgaacct      900
cgacaacagc gacgtggaca agctgttcat ccagctggtg cagacctaca accagctggt      960
cgaggaaaac cccatcaacg ccagcggcgt ggacgccaag gccatcctgt ctgccagact      1020
gagcaagagc agacggctgg aaaatctgat cgcccagctg cccggcgaga agaagaatgg      1080

```

-continued

cctgttcggc	aacctgattg	cctgagcct	gggectgacc	cccaacttca	agagcaactt	1140
cgacctggcc	gaggatgcca	aactgcagct	gagcaaggac	acctacgacg	acgacctgga	1200
caacctgctg	gccagatcg	gcgaccagta	cgccgacctg	tttctggccg	ccaagaacct	1260
gtccgacgcc	atcctgctga	gcgacatcct	gagagtgaac	accgagatca	ccaaggcccc	1320
cctgagcgcc	tctatgatca	agagatacga	cgagcaccac	caggacctga	cctgctgaa	1380
agctctcgtg	cggcagcagc	tgctgagaa	gtacaaagag	atcttcttcg	accagagcaa	1440
gaacggctac	gccggctaca	ttgacggcgg	agccagccag	gaagagtctt	acaagttcat	1500
caagcccatc	ctggaaaaga	tggacggcac	cgaggaactg	ctcgtgaagc	tgaacagaga	1560
ggacctgctg	cggaaagcagc	ggaccttcga	caacggcagc	atccccacc	agatccacct	1620
gggagagctg	cacgccattc	tgccggcgca	ggaagatctt	taccatttcc	tgaaggacaa	1680
ccgggaaaag	atcgagaaga	tctgacctt	ccgcatcccc	tactacgtgg	gccctctggc	1740
caggggaaac	agcagattcg	cctggatgac	cagaaagagc	gaggaaacca	tcacccctg	1800
gaacttcgag	gaagtgtg	acaagggcgc	ttccgccag	agcttcacg	agcggatgac	1860
caacttcgat	aagaacctgc	ccaacgagaa	ggtgctgccc	aagcacagcc	tgctgtacga	1920
gtacttcacc	gtgtataacg	agctgaccaa	agtgaatac	gtgaccgagg	gaatgagaaa	1980
gcccccttc	ctgagcggcg	agcagaaaaa	ggccatcgtg	gacctgctgt	tcaagaccaa	2040
ccgaaaagtg	accgtgaagc	agctgaaaga	ggactacttc	aagaaaatcg	agtgcttcga	2100
ctccgtggaa	atctccggcg	tggaagatcg	gttcaacgcc	tcctgggca	cataccacga	2160
tctgctgaaa	attatcaagg	acaaggactt	cctggacaat	gaggaaaacg	aggacattct	2220
ggaagatc	gtgctgacc	tgacctgtt	tgaggacaga	gagatgatcg	aggaacggct	2280
gaaaacctat	gccccctgt	tcgacgacaa	agtgatgaag	cagctgaagc	ggcggagata	2340
caccggctgg	ggcaggctga	gcccgaagct	gatcaacggc	atccgggaca	agcagtcgg	2400
caagacaatc	ctggatttcc	tgaagtccga	cggcttcgcc	aacagaaact	tcatgcagct	2460
gatccacgac	gacagcctga	cctttaaaga	ggacatccag	aaagcccagg	tgtccggcca	2520
gggcgatagc	ctgacagcgc	acattgccaa	tctggccggc	agccccgcca	ttaagaaggg	2580
catcctgcag	acagtgaagg	tggcggacga	gctcgtgaaa	gtgatgggcc	ggcacaagcc	2640
cgagaacatc	gtgatcga	tggccagaga	gaaccagacc	accagaagg	gacagaagaa	2700
cagccgcgag	agaatgaagc	ggatcgaaga	gggcatcaaa	gagctgggca	gccagatcct	2760
gaaagaacac	cccgtgga	acccccagct	gcagaacgag	aagctgtacc	tgtactacct	2820
gcagaatggg	cgggatatgt	acgtggacca	ggaactggac	atcaaccggc	tgtccgacta	2880
cgatgtggac	catatcgtgc	ctcagagctt	tctgaaggac	gactccatcg	acaacaaggt	2940
gctgaccaga	agcgacaaga	accggggcaa	gagcgacaac	gtgccctccg	aagaggtcgt	3000
gaagaagatg	aagaactact	ggcggcagct	gctgaacgcc	aagctgatta	cccagagaaa	3060
gttcgacaat	ctgaccaag	ccgagagagg	cggcctgagc	gaactggata	aggccggctt	3120
catcaagaga	cagctgtg	aaacccggca	gatcacaag	cacgtggcac	agatcctgga	3180
ctcccgatg	aacactaagt	acgacgagaa	tgacaagctg	atccgggaag	tgaagtgat	3240
cacctgaag	tccaagctgg	tgtccgattt	ccggaaggat	ttccagttt	acaaagtgcg	3300
cgagatcaac	aactaccacc	acgcccacga	cgcctacctg	aacgocgtcg	tgggaaccgc	3360

-continued

```

cctgatcaaa aagtacccta agctggaaag cgagttcgtg tacggcgact acaaggtgta 3420
cgacgtgctg aagatgatcg ccaagagcga gcagaaaatc ggcaaggcta ccgccaagta 3480
cttcttttac agcaacatca tgaacttttt caagaccgag attaccctgg ccaacggcga 3540
gatccggaag cggcctctga tcgagacaaa cggcgaaaacc ggggagatcg tgtgggataa 3600
gggccgggat tttgccaccg tgcggaaagt gctgagcatg cccaagtga atatcgtgaa 3660
aaagaccgag gtgcagacag gcggcttcag caaagagtct atcctgcca agaggaacag 3720
cgataagctg atgccagaa agaaggactg ggaccctaag aagtacggcg gcttcgacag 3780
ccccaccgtg gcctattctg tctgtgtggt ggccaaagtg gaaaaggga agtccaagaa 3840
actgaagagt gtgaagagc tctgtgggat caccatcatg gaaagaagca gcttcgagaa 3900
gaatcccatc gactttctgg aagccaaggg ctacaaagaa gtgaaaagg acctgatcat 3960
caagctgcct aagtactccc tgttcgagct ggaaaacggc cggaagagaa tgctggctc 4020
tgccggcgaa ctgcagaag gaaacgaact gcccctgccc tccaaatag tgaacttcct 4080
gtacctggcc agccactatg agaagctgaa gggctcccc gaggataatg agcagaaaca 4140
gctgtttgtg gaacagcaca agcactacct ggacgagatc atcgagcaga tcagcgagtt 4200
ctccaagaga gtgatcctgg ccgacgctaa tctggacaaa gtgctgtccg cctacaacaa 4260
gcaccgggat aagcccatca gagagcaggc cgagaatata atccacctgt ttacctgac 4320
caatctggga gccctgccc ccttcaagta ctttgacacc accatcgacc ggaagaggt 4380
caccagcacc aaagaggctg tggacgccac cctgatccac cagagcatca ccggcctgta 4440
cgagacacgg atcgacctgt ctacagctgg aggcgacagc cccaagaaga agagaaagg 4500
ggaggccagc taaggatccg gcaagactgg ccccgcttgg caacgcaaca gtgagcccct 4560
ccctagtgtg tttgggatg tgactatgta ttcgtgtgtt ggccaacggg tcaaccggaa 4620
cagattgata ccgccttgg catttctgt cagaatgtaa cgtcagtga tggctact 4677

```

<210> SEQ ID NO 276

<211> LENGTH: 3150

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 276

```

tctttcttgc gctatgacac ttccagcaaa aggtaggcgg ggctgcgaga cggcttccc 60
gcgctgcatg caacaccgat gatgcttoga cccccgaag ctcttcggg gctgcatgg 120
cgctccgatg ccgctccagg gcgagcgtg ttaaatagc caggccccg attgcaaaga 180
cattatagcg agtaccacaa gccatattca aacacctaga tcaactaccac ttctacacag 240
gccactcgag cttgtgatcg cactccgcta agggggcgcc tcttctctt cgtttcagtc 300
acaaccgcga aacatgccta agaagaagag gaaggttaac acgattaaca tcgctaagaa 360
cgacttctct gacatcgaac tggtgctat cccgttcaac actctggctg accattacgg 420
tgagcgttta gctcgcgaac agttggccct tgagcatgag tcttacgaga tgggtgaagc 480
acgcttccgc aagatgtttg agcgtcaact taaagctggt gaggttcggg ataacgctgc 540
cgccaagcct ctcatcacta ccctactccc taagatgatt gcacgcatca acgactggtt 600

```

-continued

tgaggaagtg aaagctaagc gcggaagcg cccgacagcc ttccagttcc tgcaagaaat	660
caagccggaa gccgtagcgt acatcaccat taagaccact ctggcttgcc taaccagtgc	720
tgacaataca accgttcagg ctgtagcaag cgcaatcggg cgggccattg aggacgaggc	780
tcgcttcggg cgtatccgtg accttgaagc taagcacttc aagaaaaacg ttgaggaaca	840
actcaacaag cgcgtagggc acgtctacaa gaaagcattt atgcaagtgg tcgaggetga	900
catgctctct aagggtctac tcgggtggcg ggcgtggtct tcgtggcata aggaagactc	960
tattcatgta ggagtacgct gcatcgagat gctcattgag tcaaccggaa tggttagctt	1020
acaccgcaa aatgctggcg tagtaggtca agactctgag actatcgaac tcgcacctga	1080
atacgctgag gctatcgcaa cccgtgcagg tgcgctggct ggcatctctc cgatgttcca	1140
accttgcgta gttcctccta agccgtggac tggcattact ggtggtggct attgggctaa	1200
cggctcgtcg cctctggcgc tgggtcgtac tcacagtaag aaagcactga tgcgctacga	1260
agacgtttac atgctcgagg tgtacaaagc gattaacatt gcgcaaaaca ccgcatggaa	1320
aatcaacaag aaagtcttag cggctgcgca cgtaatcacc aagtggaagc attgtccggg	1380
cgaggacatc cctcggattg agcgtgaaga actcccgatg aaaccggaag acatcgacat	1440
gaatcctgag gctctcaccg cgtggaacg tgcctgctgct gctgtgtacc gcaaggacaa	1500
ggctcgcaag tctcgcgcta tcagccttga gttcatgctt gagcaagcca ataagtttgc	1560
taaccataag gccatctggg tcaccttaca catggactgg cgcggctcgtg tttacgctgt	1620
gtcaatgttc aacccgcaag gtaacgatat gaccaaagga ctgcttacgc tggcgaaagg	1680
taaaccaatc ggtaaggaag gttactactg gctgaaaatc cacggtgcaa actgtgcggg	1740
tgctgacaag gttccgttcc ctgagcgcct caagtccatt gaggaaaacc acgagaacat	1800
catggcttgc gctaagtctc cactggagaa cacttgggtg gctgagcaag attctccggt	1860
ctgcttcctt gcgttctgct ttgagtacgc tggggtacag caccacggcc tgagctataa	1920
ctgctccctt ccgctggcgt ttgacgggtc ttgctctggc atccagcact tctcccgat	1980
gctccgagat gaggtagggt gtcgctgggt taacttgcct cctagtgaac ccgttcagga	2040
catctacggg attgttgcta agaaagtcaa cgagattcta caagcagacg caatcaatgg	2100
gaccgataac gaagtgttca ccgtgaccga tgagaacctt ggtgaaatct ctgagaaagt	2160
caagctgggc actaaggcac tggctgggtc atggctggct tacgggtgta ctgcgagtgt	2220
gactaagcgt tcagtcatga cgtctggctt cgggtccaaa gagttcggct tccgtcaaca	2280
agtgtcggaa gataaccatt agccagctat tgattccggc aagggtctga tgttactca	2340
gccgaatcag gctgctggat acatggctaa gctgatttgg gaatctgtga gcgtgacggg	2400
ggtagctgcg gttgaagcaa tgaactggct taagtctgct gctaagctgc tggctgctga	2460
ggtcaaatgat aagaagactg gagagattct tcgcaagcgt tgcgctgtgc attgggtaac	2520
tcctgatggg ttcctctgtg ggcaggaata caagaagcct attcagacgc gcttgaacct	2580
gatgttcctc ggtcagttcc gcttacagcc taccattaac accaacaag atagcgagat	2640
tgatgcacac aaacaggagt ctggatcgc tcctaacttt gtacacagcc aagacggtag	2700
ccacctctgt aagactgtag tgtgggcaca cgagaagtac ggaatcgaat cttttgcaact	2760
gattcacgac tccttcggta cgattccggc tgacgctgcg aacctgttca aagcagtgcg	2820
cgaaactatg gttgacacat atgagtcttg tgatgtaactg gctgatttct acgaccagtt	2880

-continued

```

cgctgaccag ttgcacgagt ctcaattgga caaaatgcc a gcacttccgg ctaaaggtaa 2940
cttgaacctc cgtgacatct tagagtcgga cttcgcgttc gcgtaaggat cgggcaagac 3000
tggccccgct tggcaacgca acagtgagcc cctccctagt gtgtttgggg atgtgactat 3060
gtattcgtgt gttggccaac gggtaaccc gaacagattg ataccgcct tggcatttcc 3120
tgtcagaatg taacgtcagt tgatgggtact 3150

```

```

<210> SEQ ID NO 277
<211> LENGTH: 125
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      polynucleotide
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (23)..(42)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

```

```

<400> SEQUENCE: 277
gaaattaata cgactcacta tannnnnnnn nnnnnnnnnn nngttttaga gctagaaata 60
gcaagttaaa ataaggctag tccgttatca acttgaaaaa gtggcaccga gtcggtgctt 120
ttttt 125

```

```

<210> SEQ ID NO 278
<211> LENGTH: 8452
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      polynucleotide

```

```

<400> SEQUENCE: 278
tgcggtatth cacaccgat caggtggcac ttttcgggga aatgtgcgcg gaaccctat 60
tgtttatth ttctaaatac attcaaatat gtatccgctc atgagattat caaaaaggat 120
cttcacctag atccttttaa attaaaaatg aagttttaaa tcaatctaaa gtatatatga 180
gtaaaacttg tctgacagtt accaatgctt aatcagtgag gcacctatct cagcgatctg 240
tctatttcgt tcctccatag ttgcctgact ccccgctcgt tagataacta cgatacggga 300
gggcttacca tctggcccca gtgctgcaat gataccgcga gaccacgct caccggctcc 360
agatttatca gcaataaacc agccagccgg aagggccgag cgcagaagtg gtcctgcaac 420
tttatccgcc tccatccagt ctattaattg ttgcccggaa gctagagtaa gtagttcgcc 480
agttaaatag ttgcgcaacg ttgttgccat tgctacaggc atcgtggtgt cacgctcgtc 540
gtttggtatg gcttcattca gctccgggtc ccaacgatca aggcgagtta catgatcccc 600
catgttgctc aaaaaagcgg ttagctcctt cggctcctcg atcgttgta gaagtaagt 660
ggccgcagtg ttatcactca tggttatgac agcactgcat aattctctta ctgtcatgcc 720
atccgtaaga tgctttctg tgactggtga gtactcaacc aagtcattct gagaatagtg 780
tatgcggcga ccgagttgct cttgccgggc gtaataacgg gataataccg cgccacatag 840
cagaacttta aaagtgtca tcattggaaa acgttcttcg gggcgaaaac tctcaaggat 900
cttaccgctg ttgagatcca gttcgatgta acccactcgt gcaccaact gatcttcagc 960
atcttttact ttcaccagcg tttctgggtg agcaaaaaa ggaaggcaaa atgccgcaaa 1020

```

-continued

aaaggaata	agggcgacac	ggaatggtg	aatactcata	ctctccttt	ttcaatatta	1080
ttgaagcatt	tatcaggggt	attgtctcat	gacaaaaatc	ccttaacgtg	agttttcggt	1140
ccactgagcg	tcagaccocg	tagaaaagat	caaaggatct	tcttgagatc	ctttttttct	1200
gcgcgtaatc	tgctgcttgc	aaacaaaaa	accacogcta	ccagcgggtg	ttgtttgccc	1260
ggatcaagag	ctaccaactc	ttttccgaa	ggttaactggc	ttcagcagag	cgagataacc	1320
aaatactggt	cttctagtgt	agccgtagtt	aggccaccac	ttcaagaact	ctgtagcacc	1380
gcctacatac	ctcgccttgc	taatcctggt	accagtggct	gttgccagtg	gagataagtc	1440
gtgtcttacc	gggttggtgact	caagacgata	gttaccggat	aaggcgcagc	ggtcgggctg	1500
aacgggggggt	tcgtgcacac	agcccagctt	ggagcgaacg	acctacaccg	aactgagata	1560
cctacagcgt	gagctatgag	aaagcggcac	gcttccgaa	gggagaaagg	cggacaggta	1620
tccggtaagc	ggcagggctg	gaacaggaga	gcgcacgagg	gagcttcag	gggaaacgc	1680
ctggtatctt	tatagtctctg	tcgggttctg	ccacctctga	cttgagcgtc	gatttttctg	1740
atgctcgtca	ggggggcgga	gcctatgaa	aaacgccagc	aacgcggcct	tttacgggt	1800
cctggccttt	tgctggcctt	ttgctcacat	gttctttcct	gcgttatccc	ctgattctgt	1860
ggataaacctg	attaccgctt	ttgagtgagc	tgataaccgt	cgccgcagcc	gaacgaccga	1920
gcgagcagag	tcagtgagcg	aggaagcgggt	cgctgaggct	tgacatgatt	ggtgcgtatg	1980
tttgtatgaa	gctacaggac	tgatttgccg	ggctatgagg	gcgggggaag	ctctggaagg	2040
gcccgcgatg	ggcgcggcgc	gtccagaagg	cgccatacgg	cccgcctggc	gcaccatcc	2100
ggtataaaag	cccgcgaccc	cgaacgggtg	cctccacttt	cagcgacaaa	cgagcactta	2160
tacatacgcg	actattctgc	cgctatacat	aaccactcag	ctagcttaag	atcccatcaa	2220
gcttgcatgc	cgggcgcgcc	agaaggagcg	cagccaaaac	aggatgatgt	ttgatggggt	2280
atgtgagcac	ttgcaaccct	tatccggaag	ccccctggcc	cacaaaggct	aggcgccaat	2340
gcaagcagtt	cgcatgcagc	ccttgagcgg	gtgcctcct	gataaacggg	ccagggggcc	2400
tatgttcttt	acttttttac	aagagaagtc	actcaacatc	ttaaaatggc	caggtgagtc	2460
gacgagcaag	cccggcggat	caggcagcgt	gcttgagat	ttgacttgca	acgcccgcct	2520
tgtgtcagc	aagccttttg	gctcctctgt	cgctgtctca	agcagcatct	aaccctgcgt	2580
cgccgtttcc	atgtgcagga	gattcgaggt	acctgtacc	catacagatg	tccagattac	2640
gcttcgccga	agaaaaagcg	caaggtcgaa	gcgtccgaca	agaagtacag	catcgccctg	2700
gacatcggca	ccaactctgt	gggctgggccc	gtgatcaccg	acgagtacaa	ggtgccccagc	2760
aagaaattca	aggtgctggg	caacaccgac	cggcacagca	tcaagaagaa	cctgatcgga	2820
gccctgctgt	tcgacagcgg	cgaaacagcc	gagggcacc	ggctgaagag	aaccgcccaga	2880
agaagataca	ccagacggaa	gaaccggatc	tgctatctgc	aagagatctt	cagcaacgag	2940
atggccaagg	tggacgacag	cttcttcocac	agactggaag	agtccttcc	ggtggaagag	3000
gataagaagc	acgagcggca	ccccatcttc	ggcaacatcg	tggacgaggt	ggcctaccac	3060
gagaagtacc	ccaccatcta	ccacctgaga	aagaaactgg	tggacagcac	cgacaaggcc	3120
gacctgcggc	tgatctatct	ggccctggcc	cacatgatca	agttccgggg	ccacttctctg	3180
atcgagggcg	acctgaaccc	cgacaacagc	gacgtggaca	agctgttcat	ccagctgggtg	3240
cagacctaca	accagctggt	cgaggaaaac	cccatcaacg	ccagcggcgt	ggacgccaaag	3300

-continued

gccatcctgt	ctgccagact	gagcaagagc	agacggctgg	aaaatctgat	cgcccagctg	3360
cccggcgaga	agaagaatgg	cctgttcggc	aacctgattg	ccctgagcct	gggcctgacc	3420
cccaacttca	agagcaactt	cgacctggcc	gaggatgcca	aactgcagct	gagcaaggac	3480
acctacgaag	acgacctgga	caacctgctg	gcccagatcg	gcgaccagta	cgccgacctg	3540
tttctggcgg	ccaagaacct	gtccgacgcc	atcctgctga	gcgacatcct	gagagtgaac	3600
accgagatca	ccaaggcccc	cctgagcgcc	tctatgatca	agagatacga	cgagcaccac	3660
caggacctga	ccctgctgaa	agctctcgtg	cggcagcagc	tgctgagaa	gtacaaagag	3720
atcttctctg	accagagcaa	gaacggctac	gccggctaca	ttgacggcgg	agccagccag	3780
gaagagttct	acaagttcat	caagcccatc	ctggaaaaga	tggaacggac	cgaggaactg	3840
ctcgtgaagc	tgaacagaga	ggacctgctg	cggaagcagc	ggaccttcga	caacggcagc	3900
atccccacc	agatccacct	gggagagctg	cacgccattc	tgcggcggca	ggaagatctt	3960
taccatttcc	tgaaggacaa	ccgggaaaag	atcgagaaga	tcctgacctt	ccgcatcccc	4020
tactacgtgg	gccctctggc	caggggaaac	agcagattcg	cctggatgac	cagaaagagc	4080
gaggaacca	tcacccccctg	gaacttcgag	gaagtggctg	acaagggcgc	ttccgcccag	4140
agcttcatcg	agcggatgac	caacttcgat	aagaacctgc	ccaacgagaa	ggtgctgccc	4200
aagcacagcc	tgctgtacga	gtacttcacc	gtgtataacg	agctgaccaa	agtgaaatac	4260
gtgaccgagg	gaatgagaaa	gcccgccttc	ctgagcggcg	agcagaaaaa	ggccatcgtg	4320
gacctgctgt	tcaagaccaa	ccgaaagtg	accgtgaagc	agctgaaaga	ggactacttc	4380
aagaaaatcg	agtgcttcga	ctccgtgaa	atctccggcg	tggaagatcg	gttcaacgcc	4440
tccctgggca	cataccacga	tctgctgaaa	attatcaagg	acaaggactt	cctggacaat	4500
gaggaaaaag	aggacattct	ggaagatc	gtgctgacct	tgacactgtt	tgaggacaga	4560
gagatgatcg	aggaacggct	gaaaacctat	gcccacctgt	tcgacgacaa	agtgatgaag	4620
cagctgaagc	ggcggagata	caccggctgg	ggcaggctga	gccggaagct	gatcaacggc	4680
atccgggaca	agcagtcggg	caagacaatc	ctggatttcc	tgaagtccga	cggcttcgcc	4740
aacagaaact	tcatgcagct	gatccacgac	gacagcctga	cctttaaaga	ggacatccag	4800
aaagcccagg	tgtccggcca	ggcgatagc	ctgcacgagc	acattgccaa	tctggccggc	4860
agccccgcca	ttaagaaggg	catcctgcag	acagtgaagg	tggtggacga	gctcgtgaaa	4920
gtgatgggcc	ggcacaagcc	cgagaacatc	gtgatcga	tgccagaga	gaaccagacc	4980
accagaagg	gacagaagaa	cagcccgag	agaatgaagc	ggatcgaaga	gggcatcaaa	5040
gagctgggca	gccagatcct	gaaagaacac	cccgtggaaa	acaccagct	gcagaacgag	5100
aagctgtacc	tgtactacct	gcagaatggg	cgggatatgt	acgtggacca	ggaactggac	5160
atcaaccggc	tgtccgacta	cgatgtggac	catatcgtgc	ctcagagctt	tctgaaggac	5220
gactccatcg	acaacaaggt	gctgaccaga	agcgacaaga	accggggcaa	gagcgacaac	5280
gtgcctccg	aagaggtcgt	gaagaagatg	aagaactact	ggcggcagct	gctgaacgcc	5340
aagctgatta	cccagagaaa	gttcgacaat	ctgaccaagg	ccgagagagg	cggcctgagc	5400
gaactggata	aggccggctt	catcaagaga	cagctggtgg	aaaccggca	gatcacaag	5460
cacgtggcac	agatcctgga	ctcccggatg	aacactaagt	acgacgagaa	tgacaagctg	5520
atccgggaag	tgaagtgat	caccctgaag	tccaagctgg	tgtccgattt	ccggaaggat	5580

-continued

ttccagtttt	acaaagtgcg	cgagatcaac	aactaccacc	acgcccacga	cgctacctg	5640
aacgccgtcg	tgggaaccgc	cctgatcaaa	aagtacccta	agctggaaag	cgagttcgtg	5700
tacggcgact	acaaggtgta	cgacgtgcgg	aagatgatcg	ccaagagcga	gcaggaaatc	5760
ggcaaggcta	ccgccaagta	cttctttctac	agcaacatca	tgaacttttt	caagaccgag	5820
attaccctgg	ccaacggcga	gatccggaag	cggcctctga	tcgagacaaa	cggcgaaacc	5880
ggggagatcg	tgtgggataa	gggccgggat	tttgccacgg	tgcggaaagt	gctgagcatg	5940
ccccaagtga	atatcgtgaa	aaagaccgag	gtgcagacag	gcggttcag	caaagagtct	6000
atcctgcccc	agaggaacag	cgataagctg	atcgccagaa	agaaggactg	ggaccctaag	6060
aagtacggcg	gcttcgacag	ccccaccgtg	gcctattctg	tgctgggtgg	ggccaaagtg	6120
gaaaagggca	agtccaagaa	actgaagagt	gtgaaagagc	tgctggggat	caccatcatg	6180
gaaagaagca	gcttcgagaa	gaatcccatc	gactttctgg	aagccaaggg	ctacaaagaa	6240
gtgaaaaagg	acctgatcat	caagctgcct	aagtactccc	tgctcgagct	ggaaaacggc	6300
cggaagagaa	tgctggcctc	tgccggcgaa	ctgcagaagg	gaaacgaact	ggcctgccc	6360
tccaaatgat	tgaacttctc	gtacctggcc	agccactatg	agaagctgaa	gggctcccc	6420
gaggataatg	agcagaaaca	gctgttgtg	gaacagcaca	agcactacct	ggacgagatc	6480
atcgagcaga	tcagcgagtt	ctccaagaga	gtgatcctgg	ccgacgctaa	tctggacaaa	6540
gtgctgtccg	cctacaacaa	gcaccgggat	aagcccatca	gagagcaggc	cgagaatatc	6600
atccacctgt	ttaccctgac	caatctggga	gcccctgccc	ccttcaagta	ctttgacacc	6660
accatcgacc	ggaagaggta	caccagcacc	aaagagggtc	tggacgccac	cctgatccac	6720
cagagcatca	ccggcctgta	cgagacacgg	atcgacctgt	ctcagctggg	aggcgacagc	6780
cccaagaaga	agagaaaagt	ggaggccagc	taacatatga	ttcgaatgtc	tttcttgccg	6840
tatgacactt	ccagcaaaag	gtagggcggg	ctgcgagacg	gcttcccggc	gctgcatgca	6900
acaccgatga	tgcttcgacc	ccccgaagct	ccttcggggc	tgcatggggc	ctccgatgcc	6960
gctccagggc	gagcgtggtt	taaatagcca	ggccccgat	tgcaaaagaca	ttatagcgag	7020
ctaccaaagc	catattcaaa	cacctagatc	actaccactt	ctacacaggc	cactcgagct	7080
tgtgatcgca	ctccgctaag	ggggcgccctc	ttcctcttcg	tttcagtcac	aaaccgcaaa	7140
catgacacaa	gaatcctctg	tactttctga	ccgtattgat	tcggatgatt	cctacgcgag	7200
cctgcggaac	gaccaggaat	tctgggaggt	gagtcgacga	gcaagcccgg	cgatcaggc	7260
agcgtgcttg	cagatttgac	ttgcaacgcc	cgcattgtgt	cgacgaaggc	ttttggctcc	7320
tctgtcgtg	tctcaagcag	catctaacc	tgcgtcggcg	tttccatttg	cagccgctgg	7380
cccgccgagc	cctggaggag	ctcgggctgc	cggtgccgcc	ggtgctcggg	gtgcccggcg	7440
agagcaccaa	ccccgtactg	gtcggcgagc	ccggcccggg	gatcaagctg	ttcggcgagc	7500
actggtgcgg	tccggagagc	ctcgcgtcgg	agtcggaggc	gtacgcggtc	ctggcgagc	7560
ccccggtgcc	ggtgccccgc	ctcctcggcc	gcgggcggct	gcgccccggc	accggagcct	7620
ggcctgggcc	ctacctgggtg	atgagccgga	tgaccggcac	cacctggcgg	tccgcatggg	7680
acggcacgac	cgaccggaac	gcgctgctcg	ccctggcccg	cgaactcggc	cggtgctcg	7740
gcccgtgca	cagggtgccg	ctgaccggga	acaccgtgct	cacccccat	tccgaggtct	7800
tcccggaaact	gctgcgggaa	cgccgcggcg	cgaccgtcga	ggaccaccgc	gggtggggct	7860

-continued

```

acctctcgcc cggctgctg gaccgcctgg aggactggct gccggacgtg gacacgctgc 7920
tggccggcgc cgaaccccg ttcgtccacg gcgacctgca cgggaccaac atcttcgtgg 7980
acctggccgc gaccgaggtc accgggatcg tcgacttcac cgacgtctat gcgggagact 8040
cccgtacag cctggtgcaa ctgcatctca acgccttccg gggcgaccgc gagatcctgg 8100
ccgcgctgct cgacggggcg cagtggaagc ggaccgagga cttegccgc gaactgctcg 8160
ccttcacett cctgcacgac ttcgaggtgt tcgaggagac cccgctggat ctctccggt 8220
tcaccgatcc ggaggaactg gcgcagttcc tctgggggcc gccggacacc gccccggcg 8280
cctgataagg atccggcaag actggccccg cttggcaacg caacagtgag cccctccta 8340
gtgtgtttgg ggatgtgact atgtattcgt gtgttgcca acgggtcaac ccgaacagat 8400
tgataccgc cttggcattt cctgtcagaa tgtaacgtca gttgatggta ct 8452

```

```

<210> SEQ ID NO 279
<211> LENGTH: 102
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        polynucleotide

```

```

<400> SEQUENCE: 279
gttttagagc tatgctgttt tgaatggtcc caaacggaa gggcctgagt ccgagcagaa 60
gaagaagttt tagagctatg ctgttttgaa tggccccaaa ac 102

```

```

<210> SEQ ID NO 280
<211> LENGTH: 100
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 280
cggaggacaa agtataaacg gcagaagctg gaggaggaag ggcctgagtc cgagcagaag 60
aagaagggtc cccatcacat caaccggtgg cgcattgcca 100

```

```

<210> SEQ ID NO 281
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 281
agctggagga ggaagggcct gactccgagc agaagaagaa gggctcccac 50

```

```

<210> SEQ ID NO 282
<211> LENGTH: 30
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        oligonucleotide

```

```

<400> SEQUENCE: 282
gaguccgagc agaagaagaa guuuuagagc 30

```

```

<210> SEQ ID NO 283
<211> LENGTH: 49
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

```

-continued

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide

<400> SEQUENCE: 283

agctggagga ggaagggcct gagtccgagc agaagagaag ggctcccat 49

<210> SEQ ID NO 284
 <211> LENGTH: 53
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 284

ctggaggagg aagggcctga gtccgagcag aagaagaagg gctcccatca cat 53

<210> SEQ ID NO 285
 <211> LENGTH: 52
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 285

ctggaggagg aagggcctga gtccgagcag aagagaagg ctcccatcac at 52

<210> SEQ ID NO 286
 <211> LENGTH: 54
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 286

ctggaggagg aagggcctga gtccgagcag aagaaagaag ggctcccatc acat 54

<210> SEQ ID NO 287
 <211> LENGTH: 50
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 287

ctggaggagg aagggcctga gtccgagcag aagaagggt cccatcacat 50

<210> SEQ ID NO 288
 <211> LENGTH: 47
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 288

ctggaggagg aagggcctga gcccgagcag aagggtccc atcacat 47

<210> SEQ ID NO 289
 <211> LENGTH: 66
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence

<220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide

<220> FEATURE:
 <223> OTHER INFORMATION: Description of Combined DNA/RNA Molecule: Synthetic oligonucleotide

<220> FEATURE:
 <221> NAME/KEY: modified_base
 <222> LOCATION: (1)..(20)
 <223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 289

nnnnnnnnnn nnnnnnnnnn guuuuagagc uagaaauagc aaguuaaaau aaggctagtc 60

-continued

cguuuu 66

<210> SEQ ID NO 290
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 290

gaguccgagc agaagaagaa 20

<210> SEQ ID NO 291
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 291

gacaucgaug uccuccccau 20

<210> SEQ ID NO 292
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 292

gucaccucca augacuaggg 20

<210> SEQ ID NO 293
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 293

auuggguguu cagggcagag 20

<210> SEQ ID NO 294
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 294

guggcgagag gggccgagau 20

<210> SEQ ID NO 295
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

-continued

<400> SEQUENCE: 295

ggggccgaga uggguguuc 20

<210> SEQ ID NO 296

<211> LENGTH: 20

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 296

gugccauag cuaaugcau 20

<210> SEQ ID NO 297

<211> LENGTH: 20

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 297

guaccacca caggugccag 20

<210> SEQ ID NO 298

<211> LENGTH: 20

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 298

gaaagccucu gggccaggaa 20

<210> SEQ ID NO 299

<211> LENGTH: 48

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 299

ctggaggagg aagggcctga gtccgagcag aagaagaagg gctcccat 48

<210> SEQ ID NO 300

<211> LENGTH: 20

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 300

gaguccgagc agaagaagau 20

<210> SEQ ID NO 301

<211> LENGTH: 20

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

-continued

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 301

gaguccgagc agaagaagua 20

<210> SEQ ID NO 302
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 302

gaguccgagc agaagaacaa 20

<210> SEQ ID NO 303
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 303

gaguccgagc agaagaugaa 20

<210> SEQ ID NO 304
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 304

gaguccgagc agaaguagaa 20

<210> SEQ ID NO 305
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 305

gaguccgagc agaugaagaa 20

<210> SEQ ID NO 306
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 306

gaguccgagc acaagaagaa 20

<210> SEQ ID NO 307
<211> LENGTH: 20
<212> TYPE: RNA

-continued

<213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 oligonucleotide

 <400> SEQUENCE: 307

 gaguccgagg agaagaagaa 20

 <210> SEQ ID NO 308
 <211> LENGTH: 20
 <212> TYPE: RNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 oligonucleotide

 <400> SEQUENCE: 308

 gaguccgugc agaagaagaa 20

 <210> SEQ ID NO 309
 <211> LENGTH: 20
 <212> TYPE: RNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 oligonucleotide

 <400> SEQUENCE: 309

 gagucggagc agaagaagaa 20

 <210> SEQ ID NO 310
 <211> LENGTH: 20
 <212> TYPE: RNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 oligonucleotide

 <400> SEQUENCE: 310

 gagaccgagc agaagaagaa 20

 <210> SEQ ID NO 311
 <211> LENGTH: 24
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 oligonucleotide

 <400> SEQUENCE: 311

 aatgacaagc ttgctagcgg tggg 24

 <210> SEQ ID NO 312
 <211> LENGTH: 39
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 oligonucleotide

 <400> SEQUENCE: 312

 aaaacggaag ggcctgagtc cgagcagaag aagaagttt 39

 <210> SEQ ID NO 313

-continued

<211> LENGTH: 39
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 313

aaacaggggc cgagattggg tggtcagggc agaggtttt 39

<210> SEQ ID NO 314
<211> LENGTH: 38
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 314

aaaacggaag ggctgagtc cgagcagaag aagaagtt 38

<210> SEQ ID NO 315
<211> LENGTH: 40
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 315

aacggaggga ggggcacaga tgagaaactc agggtttag 40

<210> SEQ ID NO 316
<211> LENGTH: 38
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 316

agcccttctt cttctgctcg gactcaggcc cttcctcc 38

<210> SEQ ID NO 317
<211> LENGTH: 40
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 317

cagggaggga ggggcacaga tgagaaactc aggaggcccc 40

<210> SEQ ID NO 318
<211> LENGTH: 80
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 318

ggcaatgcgc caccggttga tgtgatggga gcccttctag gaggccccca gagcagccac 60

tggggcctca acactcagggc 80

-continued

<210> SEQ ID NO 319
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 319

gtcacctcca atgactaggg tgg 23

<210> SEQ ID NO 320
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (6)..(25)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 320

caccgnnnnn nnnnnnnnnn nnnnn 25

<210> SEQ ID NO 321
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (5)..(24)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 321

aaacnnnnnn nnnnnnnnnn nnnnc 25

<210> SEQ ID NO 322
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 322

gatc gatgtc ctccccattg gctgcttcg tgg 33

<210> SEQ ID NO 323
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 323

ttcgtggcaa tgcgccaccg gttgatgtga tgg 33

<210> SEQ ID NO 324
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 324

tcgtggcaat gcgccaccg ttgatgtgat ggg 33

<210> SEQ ID NO 325
<211> LENGTH: 33

-continued

<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 325
tccagcttct gccgtttgta ctttgctctc cgg 33

<210> SEQ ID NO 326
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 326
ggagggaggg gcacagatga gaaactcagg agg 33

<210> SEQ ID NO 327
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 327
aggggccgag attgggtgtt cagggcagag agg 33

<210> SEQ ID NO 328
<211> LENGTH: 54
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 328
aacaccgggt cttcgagaag acctgtttta gagctagaaa tagcaagtta aaat 54

<210> SEQ ID NO 329
<211> LENGTH: 54
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 329
caaaacgggt cttcgagaag acgttttaga gctatgctgt tttgaatggt ccca 54

<210> SEQ ID NO 330
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 330
caagcactga gtgccattag ctaaattgcat agg 33

<210> SEQ ID NO 331
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 331
aatgcatagg gtaccacca caggtgccag ggg 33

<210> SEQ ID NO 332
<211> LENGTH: 33

-continued

<212> TYPE: DNA
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 332

acacacatgg gaaagcctct gggccaggaa agg 33

<210> SEQ ID NO 333
 <211> LENGTH: 37
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 333

ggaggaggta gtatacagaa acacagagaa gtagaat 37

<210> SEQ ID NO 334
 <211> LENGTH: 37
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 334

agaatgtaga ggagtcacag aaactcagca ctagaaa 37

<210> SEQ ID NO 335
 <211> LENGTH: 98
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 oligonucleotide

<400> SEQUENCE: 335

ggacgaaaca ccggaacat tcaaaacagc atagcaagtt aaaataaggc tagtccgta 60

tcaacttgaa aaagtggcac cgagtcggtg cttttttt 98

<210> SEQ ID NO 336
 <211> LENGTH: 186
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 polynucleotide

<400> SEQUENCE: 336

ggacgaaaca ccggtagtat taagtattgt tttatggctg ataaatttct ttgaatttct 60

ccttgattat ttgttataaa agttataaaa taatcttggt ggaaccattc aaaacagcat 120

agcaagttaa aataaggcta gtccgttatc aacttgaaaa agtggcaccg agtcggtgct 180

tttttt 186

<210> SEQ ID NO 337
 <211> LENGTH: 95
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 oligonucleotide

<400> SEQUENCE: 337

gggttttaga gctatgctgt tttgaatggt cccaaaacgg gtcttcgaga agacgtttta 60

gagctatgct gttttgaatg gtcccaaac ttttt 95

-continued

<210> SEQ ID NO 338
<211> LENGTH: 36
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (5)..(34)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 338

aaacnnnnnn nnnnnnnnnn nnnnnnnnnn nnnngt 36

<210> SEQ ID NO 339
<211> LENGTH: 36
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (7)..(36)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 339

taaaacnnnn nnnnnnnnnn nnnnnnnnnn nnnnnn 36

<210> SEQ ID NO 340
<211> LENGTH: 84
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 340

gtggaaagga cgaaacacccg ggtcttcgag aagacctgtt ttagagctag aaatagcaag 60

ttaaataag gctagtcggt tttt 84

<210> SEQ ID NO 341
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (6)..(24)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 341

caccgnnnnn nnnnnnnnnn nnnn 24

<210> SEQ ID NO 342
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide
<220> FEATURE:
<221> NAME/KEY: modified_base

-continued

<222> LOCATION: (5)..(23)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 342

aaacnnnnnn nnnnnnnnnn nnnnc 24

<210> SEQ ID NO 343
<211> LENGTH: 88
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 343

gttttagagc tatgctgttt tgaatggtcc caaaactgag accaaaggtc tcgttttaga 60
gctatgctgt tttgaatggt cccaaaac 88

<210> SEQ ID NO 344
<211> LENGTH: 35
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 344

aaacggaagg gcctgagtcc gacgagaaga agaag 35

<210> SEQ ID NO 345
<211> LENGTH: 35
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 345

aaaacttctt cttctgctcg gactcaggcc ctcc 35

<210> SEQ ID NO 346
<211> LENGTH: 46
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (1)..(19)
<223> OTHER INFORMATION: a, c, u, g, unknown or other

<400> SEQUENCE: 346

nnnnnnnnnn nnnnnnnnng uuauguacu cucaagauuu auuuuu 46

<210> SEQ ID NO 347
<211> LENGTH: 91
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 347

-continued

 guuacuuuaaa ucuugcagaa gcuacaaaga uaaggcuuca ugccgaaauc aacaccucgu 60

cauuuuauagg caggguuuuu ucguuuuuu a 91

<210> SEQ ID NO 348

<211> LENGTH: 70

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 348

ttttctagtgt ctgagtttct gtgactctctac tacattctac ttctctgtgt ttctgtatac 60

tacctcctcc 70

<210> SEQ ID NO 349

<211> LENGTH: 122

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 349

ggaggaaggg cctgagtccg agcagaagaa gaagggctcc catcacatca accggtggcg 60

cattgccacg aagcaggcca atggggagga catcgatgtc acctccaatg actaggggtg 120

gc 122

<210> SEQ ID NO 350

<211> LENGTH: 48

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide

<220> FEATURE:

<221> NAME/KEY: modified_base

<222> LOCATION: (3)..(32)

<223> OTHER INFORMATION: a, c, u, g, unknown or other

<400> SEQUENCE: 350

acnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnguuiuaga gcuaugcu 48

<210> SEQ ID NO 351

<211> LENGTH: 67

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide

<220> FEATURE:

<223> OTHER INFORMATION: Description of Combined DNA/RNA Molecule: Synthetic oligonucleotide

<400> SEQUENCE: 351

agcauagcaa guuuaaaaua ggctaguccg uuaucaacu gaaaagugg caccgagucg 60

gugcuuu 67

<210> SEQ ID NO 352

<211> LENGTH: 62

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide

<220> FEATURE:

<221> NAME/KEY: modified_base

<222> LOCATION: (1)..(20)

-continued

<223> OTHER INFORMATION: a, c, u, g, unknown or other

<400> SEQUENCE: 352

nnnnnnnnnn nnnnnnnnnn guuuuagagc uagaaaauagc aaguuaaaau aaggcuaguc 60

cg 62

<210> SEQ ID NO 353
 <211> LENGTH: 73
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 oligonucleotide

<400> SEQUENCE: 353

tgaatggtcc caaaacggaa gggcctgagt ccgagcagaa gaagaagttt tagagctatg 60

ctgttttgaa tgg 73

<210> SEQ ID NO 354
 <211> LENGTH: 69
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 354

ctggtcttcc acctctctgc cctgaacacc caatctcgcc ccctctcgcc acctctctgc 60

atttctggt 69

<210> SEQ ID NO 355
 <211> LENGTH: 138
 <212> TYPE: DNA
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 355

acccaagcac tgagtgccat tagctaaatg catagggtac caccacacagg tgccaggggc 60

ctttcccaaa gttccagcc ccttctccaa cctttcctgg ccagagggt ttcccatgtg 120

tgtggctgga ccctttga 138

<210> SEQ ID NO 356
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 primer

<400> SEQUENCE: 356

gtgctttgca gaggcctacc 20

<210> SEQ ID NO 357
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 primer

<400> SEQUENCE: 357

cctggagcgc atgcagtagt 20

-continued

<210> SEQ ID NO 358
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 358

accttctgtg ttccaccat tc 22

<210> SEQ ID NO 359
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 359

ttggggagtg cacagacttc 20

<210> SEQ ID NO 360
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 360

ggctccctgg gttcaaagta 20

<210> SEQ ID NO 361
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 361

agaggggtct ggatgtcgta a 21

<210> SEQ ID NO 362
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 362

tagctctaaa acttcttctt ctgctcgac 30

<210> SEQ ID NO 363
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 363

ctagccttat ttaaacttgc tatgctgttt 30

-continued

<210> SEQ ID NO 364
 <211> LENGTH: 99
 <212> TYPE: RNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 oligonucleotide
 <220> FEATURE:
 <221> NAME/KEY: modified_base
 <222> LOCATION: (1)..(20)
 <223> OTHER INFORMATION: a, c, u, g, unknown or other

 <400> SEQUENCE: 364

 nnnnnnnnnn nnnnnnnnnn guuuuagagc uagaaaagc aaguuaaaau aaggcuaguc 60
 cguaaucaac ugaaaaaagu ggcaccgagu cggugcuuu 99

 <210> SEQ ID NO 365
 <211> LENGTH: 12
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

 <400> SEQUENCE: 365

 tagcgggtaa gc 12

 <210> SEQ ID NO 366
 <211> LENGTH: 12
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

 <400> SEQUENCE: 366

 tcggtgacat gt 12

 <210> SEQ ID NO 367
 <211> LENGTH: 12
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

 <400> SEQUENCE: 367

 actccccgta gg 12

 <210> SEQ ID NO 368
 <211> LENGTH: 12
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

 <400> SEQUENCE: 368

 actgcgtggt aa 12

 <210> SEQ ID NO 369
 <211> LENGTH: 12
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

 <400> SEQUENCE: 369

 acgtcgctg at 12

 <210> SEQ ID NO 370
 <211> LENGTH: 12
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

-continued

<400> SEQUENCE: 370
taggtcgacc ag 12

<210> SEQ ID NO 371
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 371
ggcgtaatg at 12

<210> SEQ ID NO 372
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 372
tgtegcgatg ta 12

<210> SEQ ID NO 373
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 373
atgaaaacgc at 12

<210> SEQ ID NO 374
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 374
gccgaattcc tc 12

<210> SEQ ID NO 375
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 375
gcatggtacg ga 12

<210> SEQ ID NO 376
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 376
eggtactctt ac 12

<210> SEQ ID NO 377
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 377
gcctgtgccg ta 12

<210> SEQ ID NO 378

-continued

<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 378

tacggtaagt cg 12

<210> SEQ ID NO 379
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 379

cacgaaatta cc 12

<210> SEQ ID NO 380
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 380

aaccaagata cg 12

<210> SEQ ID NO 381
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 381

gagtcgatac gc 12

<210> SEQ ID NO 382
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 382

gtctcacgat cg 12

<210> SEQ ID NO 383
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 383

tcgtcgggtg ca 12

<210> SEQ ID NO 384
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 384

actccgtagt ga 12

<210> SEQ ID NO 385
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 385

-continued

caggacgtcc gt 12

<210> SEQ ID NO 386
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 386

tcgtatccct ac 12

<210> SEQ ID NO 387
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 387

tttcaaggcc gg 12

<210> SEQ ID NO 388
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 388

cgccggtgga at 12

<210> SEQ ID NO 389
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 389

gaaccctcc ta 12

<210> SEQ ID NO 390
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 390

gattcatcag cg 12

<210> SEQ ID NO 391
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 391

acaccgtct tc 12

<210> SEQ ID NO 392
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 392

atcgtgccct aa 12

<210> SEQ ID NO 393
<211> LENGTH: 12
<212> TYPE: DNA

-continued

<213> ORGANISM: Homo sapiens
<400> SEQUENCE: 393
g c g t c a a t g t t c 12

<210> SEQ ID NO 394
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<400> SEQUENCE: 394
c t c c g t a t c t c g 12

<210> SEQ ID NO 395
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<400> SEQUENCE: 395
c c g a t t c c t t c g 12

<210> SEQ ID NO 396
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<400> SEQUENCE: 396
t g c g c c t c c a g t 12

<210> SEQ ID NO 397
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<400> SEQUENCE: 397
t a a c g t c g g a g c 12

<210> SEQ ID NO 398
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<400> SEQUENCE: 398
a a g g t c g c c c a t 12

<210> SEQ ID NO 399
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<400> SEQUENCE: 399
g t c g g g g a c t a t 12

<210> SEQ ID NO 400
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<400> SEQUENCE: 400
t t c g a g c g a t t t 12

-continued

<210> SEQ ID NO 401
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 401

tgagtcgtcg ag 12

<210> SEQ ID NO 402
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 402

tttacgcaga gg 12

<210> SEQ ID NO 403
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 403

aggaagtatc gc 12

<210> SEQ ID NO 404
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 404

actcgatacc at 12

<210> SEQ ID NO 405
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 405

cgctacatag ca 12

<210> SEQ ID NO 406
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 406

ttcataaccg gc 12

<210> SEQ ID NO 407
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 407

ccaaacggtt aa 12

<210> SEQ ID NO 408
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

-continued

<400> SEQUENCE: 408
cgattccttc gt 12

<210> SEQ ID NO 409
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 409
cgtcatgaat aa 12

<210> SEQ ID NO 410
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 410
agtggcgatg ac 12

<210> SEQ ID NO 411
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 411
cccctacggc ac 12

<210> SEQ ID NO 412
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 412
gccaaaccgc ac 12

<210> SEQ ID NO 413
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 413
tgggacaccg gt 12

<210> SEQ ID NO 414
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 414
ttgactgcgg cg 12

<210> SEQ ID NO 415
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 415
actatgcgta gg 12

<210> SEQ ID NO 416

-continued

<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 416

tcacccaaag cg 12

<210> SEQ ID NO 417
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 417

gcaggacgtc cg 12

<210> SEQ ID NO 418
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 418

acaccgaaaa cg 12

<210> SEQ ID NO 419
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 419

cggtgtattg ag 12

<210> SEQ ID NO 420
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 420

cacgaggtat gc 12

<210> SEQ ID NO 421
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 421

taaagcgacc cg 12

<210> SEQ ID NO 422
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 422

cttagtcggc ca 12

<210> SEQ ID NO 423
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 423

-continued

cgaaaacgtg gc 12

<210> SEQ ID NO 424
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 424

cgtgccctga ac 12

<210> SEQ ID NO 425
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 425

tttaccatcg aa 12

<210> SEQ ID NO 426
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 426

cgtagccatg tt 12

<210> SEQ ID NO 427
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 427

cccaaacggt ta 12

<210> SEQ ID NO 428
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 428

gcggtatcag aa 12

<210> SEQ ID NO 429
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 429

tcgatggtaa ac 12

<210> SEQ ID NO 430
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 430

cgactttttg ca 12

<210> SEQ ID NO 431
<211> LENGTH: 12
<212> TYPE: DNA

-continued

<213> ORGANISM: Homo sapiens
<400> SEQUENCE: 431
tcgacgactc ac 12

<210> SEQ ID NO 432
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<400> SEQUENCE: 432
acgcgtcaga ta 12

<210> SEQ ID NO 433
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<400> SEQUENCE: 433
cgtacggcac ag 12

<210> SEQ ID NO 434
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<400> SEQUENCE: 434
ctatgccgtg ca 12

<210> SEQ ID NO 435
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<400> SEQUENCE: 435
cgcgtcagat at 12

<210> SEQ ID NO 436
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<400> SEQUENCE: 436
aagatcggta gc 12

<210> SEQ ID NO 437
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<400> SEQUENCE: 437
cttcgcaagg ag 12

<210> SEQ ID NO 438
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<400> SEQUENCE: 438
gtcgtggact ac 12

-continued

<210> SEQ ID NO 439
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 439

ggtcgcatc aa 12

<210> SEQ ID NO 440
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 440

gttaacagcg tg 12

<210> SEQ ID NO 441
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 441

tagctaaccg tt 12

<210> SEQ ID NO 442
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 442

agtaaaggcg ct 12

<210> SEQ ID NO 443
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 443

ggtaatttcg tg 12

<210> SEQ ID NO 444
<211> LENGTH: 69
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 444

gucaccucca augacuaggg guuuuagagc uagaaaagc aaguuaaaau aaggcuaguc 60

cguuuuuuu 69

<210> SEQ ID NO 445
<211> LENGTH: 69
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 445

-continued

```
gacaucgaug uccuccccau guuuuagagc uagaaauagc aaguuaaaau aaggcuaguc 60
cguuuuuuu 69
```

```
<210> SEQ ID NO 446
<211> LENGTH: 69
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
    oligonucleotide
```

```
<400> SEQUENCE: 446
```

```
gaguccgagc agaagaagaa guuuuagagc uagaaauagc aaguuaaaau aaggcuaguc 60
cguuuuuuu 69
```

```
<210> SEQ ID NO 447
<211> LENGTH: 69
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
    oligonucleotide
```

```
<400> SEQUENCE: 447
```

```
ggggccgaga ugggguguuc guuuuagagc uagaaauagc aaguuaaaau aaggcuaguc 60
cguuuuuuu 69
```

```
<210> SEQ ID NO 448
<211> LENGTH: 69
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
    oligonucleotide
```

```
<400> SEQUENCE: 448
```

```
guggcgagag gggccgagau guuuuagagc uagaaauagc aaguuaaaau aaggcuaguc 60
cguuuuuuu 69
```

```
<210> SEQ ID NO 449
<211> LENGTH: 76
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
    oligonucleotide
```

```
<400> SEQUENCE: 449
```

```
gucaccucca augacuaggg guuuuagagc uagaaauagc aaguuaaaau aaggcuaguc 60
cguuaucauu uuuuuu 76
```

```
<210> SEQ ID NO 450
<211> LENGTH: 76
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
    oligonucleotide
```

```
<400> SEQUENCE: 450
```

```
gacaucgaug uccuccccau guuuuagagc uagaaauagc aaguuaaaau aaggcuaguc 60
```

-continued

cguaucuu uuuuuu 76

<210> SEQ ID NO 451
<211> LENGTH: 76
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 451

gaguccgagc agaagaagaa guuuuagagc uagaaaagc aaguuaaaau aaggcuagc 60

cguaucuu uuuuuu 76

<210> SEQ ID NO 452
<211> LENGTH: 76
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 452

ggggccgaga ugggguguuc guuuuagagc uagaaaagc aaguuaaaau aaggcuagc 60

cguaucuu uuuuuu 76

<210> SEQ ID NO 453
<211> LENGTH: 76
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 453

guggcgagag gggccgagau guuuuagagc uagaaaagc aaguuaaaau aaggcuagc 60

cguaucuu uuuuuu 76

<210> SEQ ID NO 454
<211> LENGTH: 88
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 454

gucaccucca augacuaggg guuuuagagc uagaaaagc aaguuaaaau aaggcuagc 60

cguaucaac ugaaaaagu guuuuuuu 88

<210> SEQ ID NO 455
<211> LENGTH: 88
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 455

gacaucgaug uccuccccau guuuuagagc uagaaaagc aaguuaaaau aaggcuagc 60

-continued

cguaaucaac uugaaaaagu guuuuuuu 88

<210> SEQ ID NO 456
 <211> LENGTH: 88
 <212> TYPE: RNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 oligonucleotide

<400> SEQUENCE: 456

gaguccgagc agaagaagaa guuuuagagc uagaaaauagc aaguuaaaa aaggcuaguc 60

cguaaucaac uugaaaaagu guuuuuuu 88

<210> SEQ ID NO 457
 <211> LENGTH: 88
 <212> TYPE: RNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 oligonucleotide

<400> SEQUENCE: 457

ggggccgaga ugggguguuc guuuuagagc uagaaaauagc aaguuaaaa aaggcuaguc 60

cguaaucaac uugaaaaagu guuuuuuu 88

<210> SEQ ID NO 458
 <211> LENGTH: 88
 <212> TYPE: RNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 oligonucleotide

<400> SEQUENCE: 458

guggccgagag gggccgagau guuuuagagc uagaaaauagc aaguuaaaa aaggcuaguc 60

cguaaucaac uugaaaaagu guuuuuuu 88

<210> SEQ ID NO 459
 <211> LENGTH: 103
 <212> TYPE: RNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 oligonucleotide

<400> SEQUENCE: 459

gucaccucca augacuaggg guuuuagagc uagaaaauagc aaguuaaaa aaggcuaguc 60

cguaaucaac uugaaaaagu ggcaccgagu cggugcuuuu uuu 103

<210> SEQ ID NO 460
 <211> LENGTH: 103
 <212> TYPE: RNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 oligonucleotide

<400> SEQUENCE: 460

gacaucgaug uccuccccau guuuuagagc uagaaaauagc aaguuaaaa aaggcuaguc 60

cguaaucaac uugaaaaagu ggcaccgagu cggugcuuuu uuu 103

-continued

<210> SEQ ID NO 461
<211> LENGTH: 103
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 461

gaguccgagc agaagaagaa guuuuagagc uagaaaagc aaguuaaaau aaggcuaguc 60
cguaaucaac uugaaaaagu ggcaccgagu cggugcuuuu uuu 103

<210> SEQ ID NO 462
<211> LENGTH: 103
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 462

ggggccgaga uuggguguuc guuuuagagc uagaaaagc aaguuaaaau aaggcuaguc 60
cguaaucaac uugaaaaagu ggcaccgagu cggugcuuuu uuu 103

<210> SEQ ID NO 463
<211> LENGTH: 103
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 463

guggcgagag gggccgagau guuuuagagc uagaaaagc aaguuaaaau aaggcuaguc 60
cguaaucaac uugaaaaagu ggcaccgagu cggugcuuuu uuu 103

<210> SEQ ID NO 464
<211> LENGTH: 120
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
polynucleotide

<400> SEQUENCE: 464

gtgaaagga cgaaacaccg ggtcttcgag aagacctgtt ttagagctag aaatagcaag 60
ttaaataag gctagtccgt tatcaacttg aaaaagtggc accgagtcgg tgcttttttt 120

<210> SEQ ID NO 465
<211> LENGTH: 40
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 465

tcggtgcgct ggttgatttc ttcttgcgct tttttggctt 40

<210> SEQ ID NO 466

-continued

<211> LENGTH: 26
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 466

gauuucuucu ugcgcuuuuu guuuua 26

<210> SEQ ID NO 467
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (22)..(26)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 467

tgatttcttc ttgcgctttt tnnnnn 26

<210> SEQ ID NO 468
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (21)..(21)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 468

tgatttcttc ttgcgctttt ntggct 26

<210> SEQ ID NO 469
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (2)..(2)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 469

tnatttcttc ttgcgctttt ttggct 26

<210> SEQ ID NO 470
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 470

gatttcttct tgcgcttttt tgg 23

-continued

<210> SEQ ID NO 471
<211> LENGTH: 34
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(33)

<400> SEQUENCE: 471

tcc atc cgt aca acc cac aac cct gct agt gag c 34
Ser Ile Arg Thr Thr His Asn Pro Ala Ser Glu
1 5 10

<210> SEQ ID NO 472
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
peptide

<400> SEQUENCE: 472

Ser Ile Arg Thr Thr His Asn Pro Ala Ser Glu
1 5 10

<210> SEQ ID NO 473
<211> LENGTH: 34
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(33)

<400> SEQUENCE: 473

tcc atc gca aca acc cac aac cct gct agt gag c 34
Ser Ile Ala Thr Thr His Asn Pro Ala Ser Glu
1 5 10

<210> SEQ ID NO 474
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
peptide

<400> SEQUENCE: 474

Ser Ile Ala Thr Thr His Asn Pro Ala Ser Glu
1 5 10

<210> SEQ ID NO 475
<211> LENGTH: 34
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(33)

<400> SEQUENCE: 475

-continued

```
tca atc cgt aca acc cac aac cct gct agt gag c          34
Ser Ile Arg Thr Thr His Asn Pro Ala Ser Glu
1           5           10
```

```
<210> SEQ ID NO 476
<211> LENGTH: 42
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(36)
```

```
<400> SEQUENCE: 476
```

```
caa ttg aat tta aaa gaa acc gat acc gtt ttg gtt taagga      42
Gln Leu Asn Leu Lys Glu Thr Asp Thr Val Leu Val
1           5           10
```

```
<210> SEQ ID NO 477
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
```

```
<400> SEQUENCE: 477
```

```
Gln Leu Asn Leu Lys Glu Thr Asp Thr Val Leu Val
1           5           10
```

```
<210> SEQ ID NO 478
<211> LENGTH: 42
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(42)
```

```
<400> SEQUENCE: 478
```

```
caa ttg aat tta aaa gaa acc gat acc gtt tac gaa att gga      42
Gln Leu Asn Leu Lys Glu Thr Asp Thr Val Tyr Glu Ile Gly
1           5           10
```

```
<210> SEQ ID NO 479
<211> LENGTH: 14
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
```

```
<400> SEQUENCE: 479
```

```
Gln Leu Asn Leu Lys Glu Thr Asp Thr Val Tyr Glu Ile Gly
1           5           10
```

```
<210> SEQ ID NO 480
<211> LENGTH: 34
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (2)..(34)
```

```
<400> SEQUENCE: 480
```

```
t cct aaa aaa ccg aac tcc gcg ctg cgt aaa gta          34
Pro Lys Lys Pro Asn Ser Ala Leu Arg Lys Val
1           5           10
```

```
<210> SEQ ID NO 481
<211> LENGTH: 11
<212> TYPE: PRT
```

-continued

 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 481

 Pro Lys Lys Pro Asn Ser Ala Leu Arg Lys Val
 1 5 10

<210> SEQ ID NO 482

<211> LENGTH: 34

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: CDS

<222> LOCATION: (2)..(34)

<400> SEQUENCE: 482

 t cct aca aaa ccg aac tcc gcg ctg cgt aaa gta 34
 Pro Thr Lys Pro Asn Ser Ala Leu Arg Lys Val
 1 5 10

<210> SEQ ID NO 483

<211> LENGTH: 11

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 483

 Pro Thr Lys Pro Asn Ser Ala Leu Arg Lys Val
 1 5 10

<210> SEQ ID NO 484

<211> LENGTH: 33

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 484

tgcgctgggtt gatttcttct tgcgcttttt tgg 33

<210> SEQ ID NO 485

<211> LENGTH: 33

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 485

tacgctgggtt gatttcttct tgcgcttttt ttg 33

<210> SEQ ID NO 486

<211> LENGTH: 27

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 486

ggaggggtttt atggggaag gccattg 27

<210> SEQ ID NO 487

<211> LENGTH: 29

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 487

gtaaaaaaga agactagaaa ttttgatac 29

<210> SEQ ID NO 488

<211> LENGTH: 46

-continued

<212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 488

ggagggtttt atggggaaag gcaaagaaga ctagaaattt tgatac 46

<210> SEQ ID NO 489
 <211> LENGTH: 27
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 489

aggtgaagca taatgtctca aaaaata 27

<210> SEQ ID NO 490
 <211> LENGTH: 29
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 490

atatttattaa tacaatcag tgaatcat 29

<210> SEQ ID NO 491
 <211> LENGTH: 46
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 491

aggtgaagca taatgtctca aaattaatac aatcagtga aatcat 46

<210> SEQ ID NO 492
 <211> LENGTH: 36
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1)..(36)

<400> SEQUENCE: 492

aat tta aaa gaa acc gat acc gtt tac gaa att gga 36
 Asn Leu Lys Glu Thr Asp Thr Val Tyr Glu Ile Gly
 1 5 10

<210> SEQ ID NO 493
 <211> LENGTH: 12
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 493

Asn Leu Lys Glu Thr Asp Thr Val Tyr Glu Ile Gly
 1 5 10

<210> SEQ ID NO 494
 <211> LENGTH: 36
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1)..(30)

<400> SEQUENCE: 494

aat tta aaa gaa acc gat acc gtt ttg gtt taagga 36
 Asn Leu Lys Glu Thr Asp Thr Val Leu Val

-continued

 1 5 10

<210> SEQ ID NO 495
 <211> LENGTH: 10
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 495

Asn Leu Lys Glu Thr Asp Thr Val Leu Val
 1 5 10

<210> SEQ ID NO 496
 <211> LENGTH: 36
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1)..(36)

<400> SEQUENCE: 496

tgg gat cca aaa aaa tat ggt ggt ttt gat agt cca 36
 Trp Asp Pro Lys Lys Tyr Gly Gly Phe Asp Ser Pro
 1 5 10

<210> SEQ ID NO 497
 <211> LENGTH: 12
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 497

Trp Asp Pro Lys Lys Tyr Gly Gly Phe Asp Ser Pro
 1 5 10

<210> SEQ ID NO 498
 <211> LENGTH: 36
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 oligonucleotide
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1)..(36)

<400> SEQUENCE: 498

tgg gat cca aaa aaa tat tgt ggt ttt gat agt cca 36
 Trp Asp Pro Lys Lys Tyr Cys Gly Phe Asp Ser Pro
 1 5 10

<210> SEQ ID NO 499
 <211> LENGTH: 12
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 peptide

<400> SEQUENCE: 499

Trp Asp Pro Lys Lys Tyr Cys Gly Phe Asp Ser Pro
 1 5 10

<210> SEQ ID NO 500
 <211> LENGTH: 35
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 500

aaactacttt acgcagcgcg gagttcgggtt ttttg 35

<210> SEQ ID NO 501

<211> LENGTH: 4104

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: CDS

<222> LOCATION: (1)..(4104)

<400> SEQUENCE: 501

atg gac aag aag tac agc atc ggc ctg gac atc ggc acc aac tct gtg 48
Met Asp Lys Lys Tyr Ser Ile Gly Leu Asp Ile Gly Thr Asn Ser Val
1 5 10 15ggc tgg gcc gtg atc acc gac gag tac aag gtg ccc agc aag aaa ttc 96
Gly Trp Ala Val Ile Thr Asp Glu Tyr Lys Val Pro Ser Lys Lys Phe
20 25 30aag gtg ctg ggc aac acc gac cgg cac agc atc aag aag aac ctg atc 144
Lys Val Leu Gly Asn Thr Asp Arg His Ser Ile Lys Lys Asn Leu Ile
35 40 45gga gcc ctg ctg ttc gac agc ggc gaa aca gcc gag gcc acc cgg ctg 192
Gly Ala Leu Leu Phe Asp Ser Gly Glu Thr Ala Glu Ala Thr Arg Leu
50 55 60aag aga acc gcc aga aga aga tac acc aga cgg aag aac cgg atc tgc 240
Lys Arg Thr Ala Arg Arg Arg Tyr Thr Arg Arg Lys Asn Arg Ile Cys
65 70 75 80tat ctg caa gag atc ttc agc aac gag atg gcc aag gtg gac gac agc 288
Tyr Leu Gln Glu Ile Phe Ser Asn Glu Met Ala Lys Val Asp Asp Ser
85 90 95ttc ttc cac aga ctg gaa gag tcc ttc ctg gtg gaa gag gat aag aag 336
Phe Phe His Arg Leu Glu Glu Ser Phe Leu Val Glu Glu Asp Lys Lys
100 105 110cac gag cgg cac ccc atc ttc ggc aac atc gtg gac gag gtg gcc tac 384
His Glu Arg His Pro Ile Phe Gly Asn Ile Val Asp Glu Val Ala Tyr
115 120 125cac gag aag tac ccc acc atc tac cac ctg aga aag aaa ctg gtg gac 432
His Glu Lys Tyr Pro Thr Ile Tyr His Leu Arg Lys Lys Leu Val Asp
130 135 140agc acc gac aag gcc gac ctg cgg ctg atc tat ctg gcc ctg gcc cac 480
Ser Thr Asp Lys Ala Asp Leu Arg Leu Ile Tyr Leu Ala Leu Ala His
145 150 155 160atg atc aag ttc cgg ggc cac ttc ctg atc gag ggc gac ctg aac ccc 528
Met Ile Lys Phe Arg Gly His Phe Leu Ile Glu Gly Asp Leu Asn Pro
165 170 175gac aac agc gac gtg gac aag ctg ttc atc cag ctg gtg cag acc tac 576
Asp Asn Ser Asp Val Asp Lys Leu Phe Ile Gln Leu Val Gln Thr Tyr
180 185 190aac cag ctg ttc gag gaa aac ccc atc aac gcc agc ggc gtg gac gcc 624
Asn Gln Leu Phe Glu Glu Asn Pro Ile Asn Ala Ser Gly Val Asp Ala
195 200 205aag gcc atc ctg tct gcc aga ctg agc aag agc aga cgg ctg gaa aat 672
Lys Ala Ile Leu Ser Ala Arg Leu Ser Lys Ser Arg Arg Leu Glu Asn
210 215 220

ctg atc gcc cag ctg ccc ggc gag aag aag aat ggc ctg ttc ggc aac 720

-continued

Leu Ile Ala Gln Leu Pro Gly Glu Lys Lys Asn Gly Leu Phe Gly Asn 225 230 235 240	
ctg att gcc ctg agc ctg ggc ctg acc ccc aac ttc aag agc aac ttc Leu Ile Ala Leu Ser Leu Gly Leu Thr Pro Asn Phe Lys Ser Asn Phe 245 250 255	768
gac ctg gcc gag gat gcc aaa ctg cag ctg agc aag gac acc tac gac Asp Leu Ala Glu Asp Ala Lys Leu Gln Leu Ser Lys Asp Thr Tyr Asp 260 265 270	816
gac gac ctg gac aac ctg ctg gcc cag atc ggc gac cag tac gcc gac Asp Asp Leu Asp Asn Leu Leu Ala Gln Ile Gly Asp Gln Tyr Ala Asp 275 280 285	864
ctg ttt ctg gcc gcc aag aac ctg tcc gac gcc atc ctg ctg agc gac Leu Phe Leu Ala Ala Lys Asn Leu Ser Asp Ala Ile Leu Leu Ser Asp 290 295 300	912
atc ctg aga gtg aac acc gag atc acc aag gcc ccc ctg agc gcc tct Ile Leu Arg Val Asn Thr Glu Ile Thr Lys Ala Pro Leu Ser Ala Ser 305 310 315 320	960
atg atc aag aga tac gac gag cac cac cag gac ctg acc ctg ctg aaa Met Ile Lys Arg Tyr Asp Glu His His Gln Asp Leu Thr Leu Leu Lys 325 330 335	1008
gct ctc gtg cgg cag cag ctg cct gag aag tac aaa gag att ttc ttc Ala Leu Val Arg Gln Gln Leu Pro Glu Lys Tyr Lys Glu Ile Phe Phe 340 345 350	1056
gac cag agc aag aac ggc tac gcc ggc tac att gac ggc gga gcc agc Asp Gln Ser Lys Asn Gly Tyr Ala Gly Tyr Ile Asp Gly Gly Ala Ser 355 360 365	1104
cag gaa gag ttc tac aag ttc atc aag ccc atc ctg gaa aag atg gac Gln Glu Glu Phe Tyr Lys Phe Ile Lys Pro Ile Leu Glu Lys Met Asp 370 375 380	1152
ggc acc gag gaa ctg ctc gtg aag ctg aac aga gag gac ctg ctg cgg Gly Thr Glu Glu Leu Leu Val Lys Leu Asn Arg Glu Asp Leu Leu Arg 385 390 395 400	1200
aag cag cgg acc ttc gac aac ggc agc atc ccc cac cag atc cac ctg Lys Gln Arg Thr Phe Asp Asn Gly Ser Ile Pro His Gln Ile His Leu 405 410 415	1248
gga gag ctg cac gcc att ctg cgg cgg cag gaa gat ttt tac cca ttc Gly Glu Leu His Ala Ile Leu Arg Arg Gln Glu Asp Phe Tyr Pro Phe 420 425 430	1296
ctg aag gac aac cgg gaa aag atc gag aag atc ctg acc ttc cgc atc Leu Lys Asp Asn Arg Glu Lys Ile Glu Lys Ile Leu Thr Phe Arg Ile 435 440 445	1344
ccc tac tac gtg ggc cct ctg gcc agg gga aac agc aga ttc gcc tgg Pro Tyr Tyr Val Gly Pro Leu Ala Arg Gly Asn Ser Arg Phe Ala Trp 450 455 460	1392
atg acc aga aag agc gag gaa acc atc acc ccc tgg aac ttc gag gaa Met Thr Arg Lys Ser Glu Glu Thr Ile Thr Pro Trp Asn Phe Glu Glu 465 470 475 480	1440
gtg gtg gac aag ggc gct tcc gcc cag agc ttc atc gag cgg atg acc Val Val Asp Lys Gly Ala Ser Ala Gln Ser Phe Ile Glu Arg Met Thr 485 490 495	1488
aac ttc gat aag aac ctg ccc aac gag aag gtg ctg ccc aag cac agc Asn Phe Asp Lys Asn Leu Pro Asn Glu Lys Val Leu Pro Lys His Ser 500 505 510	1536
ctg ctg tac gag tac ttc acc gtg tat aac gag ctg acc aaa gtg aaa Leu Leu Tyr Glu Tyr Phe Thr Val Tyr Asn Glu Leu Thr Lys Val Lys 515 520 525	1584
tac gtg acc gag gga atg aga aag ccc gcc ttc ctg agc ggc gag cag 1632	

-continued

Tyr	Val	Thr	Glu	Gly	Met	Arg	Lys	Pro	Ala	Phe	Leu	Ser	Gly	Glu	Gln	
530						535					540					
aaa	aag	gcc	atc	gtg	gac	ctg	ctg	ttc	aag	acc	aac	cgg	aaa	gtg	acc	1680
Lys	Lys	Ala	Ile	Val	Asp	Leu	Leu	Phe	Lys	Thr	Asn	Arg	Lys	Val	Thr	
545					550					555					560	
gtg	aag	cag	ctg	aaa	gag	gac	tac	ttc	aag	aaa	atc	gag	tgc	ttc	gac	1728
Val	Lys	Gln	Leu	Lys	Glu	Asp	Tyr	Phe	Lys	Lys	Ile	Glu	Cys	Phe	Asp	
				565					570					575		
tcc	gtg	gaa	atc	tcc	ggc	gtg	gaa	gat	cgg	ttc	aac	gcc	tcc	ctg	ggc	1776
Ser	Val	Glu	Ile	Ser	Gly	Val	Glu	Asp	Arg	Phe	Asn	Ala	Ser	Leu	Gly	
				580				585					590			
aca	tac	cac	gat	ctg	ctg	aaa	att	atc	aag	gac	aag	gac	ttc	ctg	gac	1824
Thr	Tyr	His	Asp	Leu	Leu	Lys	Ile	Ile	Lys	Asp	Lys	Asp	Phe	Leu	Asp	
		595					600					605				
aat	gag	gaa	aac	gag	gac	att	ctg	gaa	gat	atc	gtg	ctg	acc	ctg	aca	1872
Asn	Glu	Glu	Asn	Glu	Asp	Ile	Leu	Glu	Asp	Ile	Val	Leu	Thr	Leu	Thr	
610					615						620					
ctg	ttt	gag	gac	aga	gag	atg	atc	gag	gaa	cgg	ctg	aaa	acc	tat	gcc	1920
Leu	Phe	Glu	Asp	Arg	Glu	Met	Ile	Glu	Glu	Arg	Leu	Lys	Thr	Tyr	Ala	
625					630					635					640	
cac	ctg	ttc	gac	gac	aaa	gtg	atg	aag	cag	ctg	aag	cgg	cgg	aga	tac	1968
His	Leu	Phe	Asp	Asp	Lys	Val	Met	Lys	Gln	Leu	Lys	Arg	Arg	Arg	Tyr	
				645					650					655		
acc	ggc	tgg	ggc	agg	ctg	agc	cgg	aag	ctg	atc	aac	ggc	atc	cgg	gac	2016
Thr	Gly	Trp	Gly	Arg	Leu	Ser	Arg	Lys	Leu	Ile	Asn	Gly	Ile	Arg	Asp	
			660					665					670			
aag	cag	tcc	ggc	aag	aca	atc	ctg	gat	ttc	ctg	aag	tcc	gac	ggc	ttc	2064
Lys	Gln	Ser	Gly	Lys	Thr	Ile	Leu	Asp	Phe	Leu	Lys	Ser	Asp	Gly	Phe	
			675				680					685				
gcc	aac	aga	aac	ttc	atg	cag	ctg	atc	cac	gac	gac	agc	ctg	acc	ttt	2112
Ala	Asn	Arg	Asn	Phe	Met	Gln	Leu	Ile	His	Asp	Asp	Ser	Leu	Thr	Phe	
690						695					700					
aaa	gag	gac	atc	cag	aaa	gcc	cag	gtg	tcc	ggc	cag	ggc	gat	agc	ctg	2160
Lys	Glu	Asp	Ile	Gln	Lys	Ala	Gln	Val	Ser	Gly	Gln	Gly	Asp	Ser	Leu	
705					710					715					720	
cac	gag	cac	att	gcc	aat	ctg	gcc	ggc	agc	ccc	gcc	att	aag	aag	ggc	2208
His	Glu	His	Ile	Ala	Asn	Leu	Ala	Gly	Ser	Pro	Ala	Ile	Lys	Lys	Gly	
				725				730						735		
atc	ctg	cag	aca	gtg	aag	gtg	gtg	gac	gag	ctc	gtg	aaa	gtg	atg	ggc	2256
Ile	Leu	Gln	Thr	Val	Lys	Val	Val	Asp	Glu	Leu	Val	Lys	Val	Met	Gly	
			740					745					750			
cgg	cac	aag	ccc	gag	aac	atc	gtg	atc	gcc	atg	gcc	aga	gag	aac	cag	2304
Arg	His	Lys	Pro	Glu	Asn	Ile	Val	Ile	Ala	Met	Ala	Arg	Glu	Asn	Gln	
		755				760						765				
acc	acc	cag	aag	gga	cag	aag	aac	agc	cgc	gag	aga	atg	aag	cgg	atc	2352
Thr	Thr	Gln	Lys	Gly	Gln	Lys	Asn	Ser	Arg	Glu	Arg	Met	Lys	Arg	Ile	
770						775					780					
gaa	gag	ggc	atc	aaa	gag	ctg	ggc	agc	cag	atc	ctg	aaa	gaa	cac	ccc	2400
Glu	Glu	Gly	Ile	Lys	Glu	Leu	Gly	Ser	Gln	Ile	Leu	Lys	Glu	His	Pro	
785					790					795					800	
gtg	gaa	aac	acc	cag	ctg	cag	aac	gag	aag	ctg	tac	ctg	tac	tac	ctg	2448
Val	Glu	Asn	Thr	Gln	Leu	Gln	Asn	Glu	Lys	Leu	Tyr	Leu	Tyr	Tyr	Leu	
				805					810					815		
cag	aat	ggg	cgg	gat	atg	tac	gtg	gac	cag	gaa	ctg	gac	atc	aac	cgg	2496
Gln	Asn	Gly	Arg	Asp	Met	Tyr	Val	Asp	Gln	Glu	Leu	Asp	Ile	Asn	Arg	
			820					825					830			
ctg	tcc	gac	tac	gat	gtg	gac	gcc	atc	gtg	cct	cag	agc	ttt	ctg	aag	2544

-continued

Leu	Ser	Asp	Tyr	Asp	Val	Asp	Ala	Ile	Val	Pro	Gln	Ser	Phe	Leu	Lys		
		835					840					845					
gac	gac	tcc	atc	gac	gcc	aag	gtg	ctg	acc	aga	agc	gac	aag	gcc	cgg	2592	
Asp	Asp	Ser	Ile	Asp	Ala	Lys	Val	Leu	Thr	Arg	Ser	Asp	Lys	Ala	Arg		
	850					855					860						
ggc	aag	agc	gac	aac	gtg	ccc	tcc	gaa	gag	gtc	gtg	aag	aag	atg	aag	2640	
Gly	Lys	Ser	Asp	Asn	Val	Pro	Ser	Glu	Glu	Val	Val	Lys	Lys	Met	Lys		
	865				870					875					880		
aac	tac	tgg	cgg	cag	ctg	ctg	aac	gcc	aag	ctg	att	acc	cag	aga	aag	2688	
Asn	Tyr	Trp	Arg	Gln	Leu	Leu	Asn	Ala	Lys	Leu	Ile	Thr	Gln	Arg	Lys		
			885							890					895		
ttc	gac	aat	ctg	acc	aag	gcc	gag	aga	ggc	ggc	ctg	agc	gaa	ctg	gat	2736	
Phe	Asp	Asn	Leu	Thr	Lys	Ala	Glu	Arg	Gly	Gly	Leu	Ser	Glu	Leu	Asp		
		900						905					910				
aag	gcc	ggc	ttc	atc	aag	aga	cag	ctg	gtg	gaa	acc	cgg	cag	atc	aca	2784	
Lys	Ala	Gly	Phe	Ile	Lys	Arg	Gln	Leu	Val	Glu	Thr	Arg	Gln	Ile	Thr		
	915					920						925					
aag	cac	gtg	gca	cag	atc	ctg	gac	tcc	cgg	atg	aac	act	aag	tac	gac	2832	
Lys	His	Val	Ala	Gln	Ile	Leu	Asp	Ser	Arg	Met	Asn	Thr	Lys	Tyr	Asp		
	930					935					940						
gag	aat	gac	aag	ctg	atc	cgg	gaa	gtg	aaa	gtg	atc	acc	ctg	aag	tcc	2880	
Glu	Asn	Asp	Lys	Leu	Ile	Arg	Glu	Val	Lys	Val	Ile	Thr	Leu	Lys	Ser		
	945				950					955					960		
aag	ctg	gtg	tcc	gat	ttc	cgg	aag	gat	ttc	cag	ttt	tac	aaa	gtg	cgc	2928	
Lys	Leu	Val	Ser	Asp	Phe	Arg	Lys	Asp	Phe	Gln	Phe	Tyr	Lys	Val	Arg		
		965						970					975				
gag	atc	aac	aac	tac	cac	cac	gcc	cac	gcc	gcc	tac	ctg	aac	gcc	gtc	2976	
Glu	Ile	Asn	Asn	Tyr	His	His	Ala	His	Ala	Ala	Tyr	Leu	Asn	Ala	Val		
		980					985						990				
gtg	gga	acc	gcc	ctg	atc	aaa	aag	tac	cct	aag	ctg	gaa	agc	gag	ttc	3024	
Val	Gly	Thr	Ala	Leu	Ile	Lys	Lys	Tyr	Pro	Lys	Leu	Glu	Ser	Glu	Phe		
		995					1000						1005				
gtg	tac	ggc	gac	tac	aag	gtg	tac	gac	gtg	cgg	aag	atg	atc	gcc		3069	
Val	Tyr	Gly	Asp	Tyr	Lys	Val	Tyr	Asp	Val	Arg	Lys	Met	Ile	Ala			
	1010					1015						1020					
aag	agc	gag	cag	gaa	atc	ggc	aag	gct	acc	gcc	aag	tac	ttc	ttc		3114	
Lys	Ser	Glu	Gln	Glu	Ile	Gly	Lys	Ala	Thr	Ala	Lys	Tyr	Phe	Phe			
	1025					1030					1035						
tac	agc	aac	atc	atg	aac	ttt	ttc	aag	acc	gag	att	acc	ctg	gcc		3159	
Tyr	Ser	Asn	Ile	Met	Asn	Phe	Phe	Lys	Thr	Glu	Ile	Thr	Leu	Ala			
	1040					1045						1050					
aac	ggc	gag	atc	cgg	aag	cgg	cct	ctg	atc	gag	aca	aac	ggc	gaa		3204	
Asn	Gly	Glu	Ile	Arg	Lys	Arg	Pro	Leu	Ile	Glu	Thr	Asn	Gly	Glu			
	1055					1060						1065					
acc	ggg	gag	atc	gtg	tgg	gat	aag	ggc	cgg	gat	ttt	gcc	acc	gtg		3249	
Thr	Gly	Glu	Ile	Val	Trp	Asp	Lys	Gly	Arg	Asp	Phe	Ala	Thr	Val			
	1070					1075					1080						
cgg	aaa	gtg	ctg	agc	atg	ccc	caa	gtg	aat	atc	gtg	aaa	aag	acc		3294	
Arg	Lys	Val	Leu	Ser	Met	Pro	Gln	Val	Asn	Ile	Val	Lys	Lys	Thr			
	1085					1090						1095					
gag	gtg	cag	aca	ggc	ggc	ttc	agc	aaa	gag	tct	atc	ctg	ccc	aag		3339	
Glu	Val	Gln	Thr	Gly	Gly	Phe	Ser	Lys	Glu	Ser	Ile	Leu	Pro	Lys			
	1100					1105						1110					
agg	aac	agc	gat	aag	ctg	atc	gcc	aga	aag	aag	gac	tgg	gac	cct		3384	
Arg	Asn	Ser	Asp	Lys	Leu	Ile	Ala	Arg	Lys	Lys	Asp	Trp	Asp	Pro			
	1115					1120					1125						
aag	aag	tac	ggc	ggc	ttc	gac	agc	ccc	acc	gtg	gcc	tat	tct	gtg		3429	

-continued

Lys	Lys	Tyr	Gly	Gly	Phe	Asp	Ser	Pro	Thr	Val	Ala	Tyr	Ser	Val	
	1130					1135					1140				
ctg	gtg	gtg	gcc	aaa	gtg	gaa	aag	ggc	aag	tcc	aag	aaa	ctg	aag	3474
Leu	Val	Val	Ala	Lys	Val	Glu	Lys	Gly	Lys	Ser	Lys	Lys	Leu	Lys	
	1145					1150					1155				
agt	gtg	aaa	gag	ctg	ctg	ggg	atc	acc	atc	atg	gaa	aga	agc	agc	3519
Ser	Val	Lys	Glu	Leu	Leu	Gly	Ile	Thr	Ile	Met	Glu	Arg	Ser	Ser	
	1160					1165					1170				
ttc	gag	aag	aat	ccc	atc	gac	ttt	ctg	gaa	gcc	aag	ggc	tac	aaa	3564
Phe	Glu	Lys	Asn	Pro	Ile	Asp	Phe	Leu	Glu	Ala	Lys	Gly	Tyr	Lys	
	1175					1180					1185				
gaa	gtg	aaa	aag	gac	ctg	atc	atc	aag	ctg	cct	aag	tac	tcc	ctg	3609
Glu	Val	Lys	Lys	Asp	Leu	Ile	Ile	Lys	Leu	Pro	Lys	Tyr	Ser	Leu	
	1190					1195					1200				
ttc	gag	ctg	gaa	aac	ggc	cgg	aag	aga	atg	ctg	gcc	tct	gcc	ggc	3654
Phe	Glu	Leu	Glu	Asn	Gly	Arg	Lys	Arg	Met	Leu	Ala	Ser	Ala	Gly	
	1205					1210					1215				
gaa	ctg	cag	aag	gga	aac	gaa	ctg	gcc	ctg	ccc	tcc	aaa	tat	gtg	3699
Glu	Leu	Gln	Lys	Gly	Asn	Glu	Leu	Ala	Leu	Pro	Ser	Lys	Tyr	Val	
	1220					1225					1230				
aac	ttc	ctg	tac	ctg	gcc	agc	cac	tat	gag	aag	ctg	aag	ggc	tcc	3744
Asn	Phe	Leu	Tyr	Leu	Ala	Ser	His	Tyr	Glu	Lys	Leu	Lys	Gly	Ser	
	1235					1240					1245				
ccc	gag	gat	aat	gag	cag	aaa	cag	ctg	ttt	gtg	gaa	cag	cac	aag	3789
Pro	Glu	Asp	Asn	Glu	Gln	Lys	Gln	Leu	Phe	Val	Glu	Gln	His	Lys	
	1250					1255					1260				
cac	tac	ctg	gac	gag	atc	atc	gag	cag	atc	agc	gag	ttc	tcc	aag	3834
His	Tyr	Leu	Asp	Glu	Ile	Ile	Glu	Gln	Ile	Ser	Glu	Phe	Ser	Lys	
	1265					1270					1275				
aga	gtg	atc	ctg	gcc	gac	gct	aat	ctg	gac	aaa	gtg	ctg	tcc	gcc	3879
Arg	Val	Ile	Leu	Ala	Asp	Ala	Asn	Leu	Asp	Lys	Val	Leu	Ser	Ala	
	1280					1285					1290				
tac	aac	aag	cac	cgg	gat	aag	ccc	atc	aga	gag	cag	gcc	gag	aat	3924
Tyr	Asn	Lys	His	Arg	Asp	Lys	Pro	Ile	Arg	Glu	Gln	Ala	Glu	Asn	
	1295					1300					1305				
atc	atc	cac	ctg	ttt	acc	ctg	acc	aat	ctg	gga	gcc	cct	gcc	gcc	3969
Ile	Ile	His	Leu	Phe	Thr	Leu	Thr	Asn	Leu	Gly	Ala	Pro	Ala	Ala	
	1310					1315					1320				
ttc	aag	tac	ttt	gac	acc	acc	atc	gac	cgg	aag	agg	tac	acc	agc	4014
Phe	Lys	Tyr	Phe	Asp	Thr	Thr	Ile	Asp	Arg	Lys	Arg	Tyr	Thr	Ser	
	1325					1330					1335				
acc	aaa	gag	gtg	ctg	gac	gcc	acc	ctg	atc	cac	cag	agc	atc	acc	4059
Thr	Lys	Glu	Val	Leu	Asp	Ala	Thr	Leu	Ile	His	Gln	Ser	Ile	Thr	
	1340					1345					1350				
ggc	ctg	tac	gag	aca	cgg	atc	gac	ctg	tct	cag	ctg	gga	ggc	gac	4104
Gly	Leu	Tyr	Glu	Thr	Arg	Ile	Asp	Leu	Ser	Gln	Leu	Gly	Gly	Asp	
	1355					1360					1365				

<210> SEQ ID NO 502

<211> LENGTH: 1368

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 502

Met Asp Lys Lys Tyr Ser Ile Gly Leu Asp Ile Gly Thr Asn Ser Val
 1 5 10 15

Gly Trp Ala Val Ile Thr Asp Glu Tyr Lys Val Pro Ser Lys Lys Phe
 20 25 30

-continued

Lys Val Leu Gly Asn Thr Asp Arg His Ser Ile Lys Lys Asn Leu Ile
 35 40 45
 Gly Ala Leu Leu Phe Asp Ser Gly Glu Thr Ala Glu Ala Thr Arg Leu
 50 55 60
 Lys Arg Thr Ala Arg Arg Arg Tyr Thr Arg Arg Lys Asn Arg Ile Cys
 65 70 75 80
 Tyr Leu Gln Glu Ile Phe Ser Asn Glu Met Ala Lys Val Asp Asp Ser
 85 90 95
 Phe Phe His Arg Leu Glu Glu Ser Phe Leu Val Glu Glu Asp Lys Lys
 100 105 110
 His Glu Arg His Pro Ile Phe Gly Asn Ile Val Asp Glu Val Ala Tyr
 115 120 125
 His Glu Lys Tyr Pro Thr Ile Tyr His Leu Arg Lys Lys Leu Val Asp
 130 135 140
 Ser Thr Asp Lys Ala Asp Leu Arg Leu Ile Tyr Leu Ala Leu Ala His
 145 150 155 160
 Met Ile Lys Phe Arg Gly His Phe Leu Ile Glu Gly Asp Leu Asn Pro
 165 170 175
 Asp Asn Ser Asp Val Asp Lys Leu Phe Ile Gln Leu Val Gln Thr Tyr
 180 185 190
 Asn Gln Leu Phe Glu Glu Asn Pro Ile Asn Ala Ser Gly Val Asp Ala
 195 200 205
 Lys Ala Ile Leu Ser Ala Arg Leu Ser Lys Ser Arg Arg Leu Glu Asn
 210 215 220
 Leu Ile Ala Gln Leu Pro Gly Glu Lys Lys Asn Gly Leu Phe Gly Asn
 225 230 235 240
 Leu Ile Ala Leu Ser Leu Gly Leu Thr Pro Asn Phe Lys Ser Asn Phe
 245 250 255
 Asp Leu Ala Glu Asp Ala Lys Leu Gln Leu Ser Lys Asp Thr Tyr Asp
 260 265 270
 Asp Asp Leu Asp Asn Leu Leu Ala Gln Ile Gly Asp Gln Tyr Ala Asp
 275 280 285
 Leu Phe Leu Ala Ala Lys Asn Leu Ser Asp Ala Ile Leu Leu Ser Asp
 290 295 300
 Ile Leu Arg Val Asn Thr Glu Ile Thr Lys Ala Pro Leu Ser Ala Ser
 305 310 315 320
 Met Ile Lys Arg Tyr Asp Glu His His Gln Asp Leu Thr Leu Leu Lys
 325 330 335
 Ala Leu Val Arg Gln Gln Leu Pro Glu Lys Tyr Lys Glu Ile Phe Phe
 340 345 350
 Asp Gln Ser Lys Asn Gly Tyr Ala Gly Tyr Ile Asp Gly Gly Ala Ser
 355 360 365
 Gln Glu Glu Phe Tyr Lys Phe Ile Lys Pro Ile Leu Glu Lys Met Asp
 370 375 380
 Gly Thr Glu Glu Leu Leu Val Lys Leu Asn Arg Glu Asp Leu Leu Arg
 385 390 395 400
 Lys Gln Arg Thr Phe Asp Asn Gly Ser Ile Pro His Gln Ile His Leu
 405 410 415
 Gly Glu Leu His Ala Ile Leu Arg Arg Gln Glu Asp Phe Tyr Pro Phe
 420 425 430

-continued

Leu Lys Asp Asn Arg Glu Lys Ile Glu Lys Ile Leu Thr Phe Arg Ile
 435 440 445

Pro Tyr Tyr Val Gly Pro Leu Ala Arg Gly Asn Ser Arg Phe Ala Trp
 450 455 460

Met Thr Arg Lys Ser Glu Glu Thr Ile Thr Pro Trp Asn Phe Glu Glu
 465 470 475 480

Val Val Asp Lys Gly Ala Ser Ala Gln Ser Phe Ile Glu Arg Met Thr
 485 490 495

Asn Phe Asp Lys Asn Leu Pro Asn Glu Lys Val Leu Pro Lys His Ser
 500 505 510

Leu Leu Tyr Glu Tyr Phe Thr Val Tyr Asn Glu Leu Thr Lys Val Lys
 515 520 525

Tyr Val Thr Glu Gly Met Arg Lys Pro Ala Phe Leu Ser Gly Glu Gln
 530 535 540

Lys Lys Ala Ile Val Asp Leu Leu Phe Lys Thr Asn Arg Lys Val Thr
 545 550 555 560

Val Lys Gln Leu Lys Glu Asp Tyr Phe Lys Lys Ile Glu Cys Phe Asp
 565 570 575

Ser Val Glu Ile Ser Gly Val Glu Asp Arg Phe Asn Ala Ser Leu Gly
 580 585 590

Thr Tyr His Asp Leu Leu Lys Ile Ile Lys Asp Lys Asp Phe Leu Asp
 595 600 605

Asn Glu Glu Asn Glu Asp Ile Leu Glu Asp Ile Val Leu Thr Leu Thr
 610 615 620

Leu Phe Glu Asp Arg Glu Met Ile Glu Glu Arg Leu Lys Thr Tyr Ala
 625 630 635 640

His Leu Phe Asp Asp Lys Val Met Lys Gln Leu Lys Arg Arg Arg Tyr
 645 650 655

Thr Gly Trp Gly Arg Leu Ser Arg Lys Leu Ile Asn Gly Ile Arg Asp
 660 665 670

Lys Gln Ser Gly Lys Thr Ile Leu Asp Phe Leu Lys Ser Asp Gly Phe
 675 680 685

Ala Asn Arg Asn Phe Met Gln Leu Ile His Asp Asp Ser Leu Thr Phe
 690 695 700

Lys Glu Asp Ile Gln Lys Ala Gln Val Ser Gly Gln Gly Asp Ser Leu
 705 710 715 720

His Glu His Ile Ala Asn Leu Ala Gly Ser Pro Ala Ile Lys Lys Gly
 725 730 735

Ile Leu Gln Thr Val Lys Val Val Asp Glu Leu Val Lys Val Met Gly
 740 745 750

Arg His Lys Pro Glu Asn Ile Val Ile Ala Met Ala Arg Glu Asn Gln
 755 760 765

Thr Thr Gln Lys Gly Gln Lys Asn Ser Arg Glu Arg Met Lys Arg Ile
 770 775 780

Glu Glu Gly Ile Lys Glu Leu Gly Ser Gln Ile Leu Lys Glu His Pro
 785 790 795 800

Val Glu Asn Thr Gln Leu Gln Asn Glu Lys Leu Tyr Leu Tyr Tyr Leu
 805 810 815

Gln Asn Gly Arg Asp Met Tyr Val Asp Gln Glu Leu Asp Ile Asn Arg
 820 825 830

Leu Ser Asp Tyr Asp Val Asp Ala Ile Val Pro Gln Ser Phe Leu Lys

-continued

835			840			845									
Asp	Asp	Ser	Ile	Asp	Ala	Lys	Val	Leu	Thr	Arg	Ser	Asp	Lys	Ala	Arg
850					855						860				
Gly	Lys	Ser	Asp	Asn	Val	Pro	Ser	Glu	Glu	Val	Val	Lys	Lys	Met	Lys
865				870						875					880
Asn	Tyr	Trp	Arg	Gln	Leu	Leu	Asn	Ala	Lys	Leu	Ile	Thr	Gln	Arg	Lys
			885							890					895
Phe	Asp	Asn	Leu	Thr	Lys	Ala	Glu	Arg	Gly	Gly	Leu	Ser	Glu	Leu	Asp
			900					905						910	
Lys	Ala	Gly	Phe	Ile	Lys	Arg	Gln	Leu	Val	Glu	Thr	Arg	Gln	Ile	Thr
		915					920						925		
Lys	His	Val	Ala	Gln	Ile	Leu	Asp	Ser	Arg	Met	Asn	Thr	Lys	Tyr	Asp
930						935					940				
Glu	Asn	Asp	Lys	Leu	Ile	Arg	Glu	Val	Lys	Val	Ile	Thr	Leu	Lys	Ser
945				950						955					960
Lys	Leu	Val	Ser	Asp	Phe	Arg	Lys	Asp	Phe	Gln	Phe	Tyr	Lys	Val	Arg
			965					970							975
Glu	Ile	Asn	Asn	Tyr	His	His	Ala	His	Ala	Ala	Tyr	Leu	Asn	Ala	Val
		980						985							990
Val	Gly	Thr	Ala	Leu	Ile	Lys	Lys	Tyr	Pro	Lys	Leu	Glu	Ser	Glu	Phe
		995					1000						1005		
Val	Tyr	Gly	Asp	Tyr	Lys	Val	Tyr	Asp	Val	Arg	Lys	Met	Ile	Ala	
1010					1015						1020				
Lys	Ser	Glu	Gln	Glu	Ile	Gly	Lys	Ala	Thr	Ala	Lys	Tyr	Phe	Phe	
1025						1030						1035			
Tyr	Ser	Asn	Ile	Met	Asn	Phe	Phe	Lys	Thr	Glu	Ile	Thr	Leu	Ala	
1040					1045							1050			
Asn	Gly	Glu	Ile	Arg	Lys	Arg	Pro	Leu	Ile	Glu	Thr	Asn	Gly	Glu	
1055					1060							1065			
Thr	Gly	Glu	Ile	Val	Trp	Asp	Lys	Gly	Arg	Asp	Phe	Ala	Thr	Val	
1070						1075						1080			
Arg	Lys	Val	Leu	Ser	Met	Pro	Gln	Val	Asn	Ile	Val	Lys	Lys	Thr	
1085					1090							1095			
Glu	Val	Gln	Thr	Gly	Gly	Phe	Ser	Lys	Glu	Ser	Ile	Leu	Pro	Lys	
1100					1105							1110			
Arg	Asn	Ser	Asp	Lys	Leu	Ile	Ala	Arg	Lys	Lys	Asp	Trp	Asp	Pro	
1115					1120							1125			
Lys	Lys	Tyr	Gly	Gly	Phe	Asp	Ser	Pro	Thr	Val	Ala	Tyr	Ser	Val	
1130					1135							1140			
Leu	Val	Val	Ala	Lys	Val	Glu	Lys	Gly	Lys	Ser	Lys	Lys	Leu	Lys	
1145					1150							1155			
Ser	Val	Lys	Glu	Leu	Leu	Gly	Ile	Thr	Ile	Met	Glu	Arg	Ser	Ser	
1160					1165							1170			
Phe	Glu	Lys	Asn	Pro	Ile	Asp	Phe	Leu	Glu	Ala	Lys	Gly	Tyr	Lys	
1175					1180							1185			
Glu	Val	Lys	Lys	Asp	Leu	Ile	Ile	Lys	Leu	Pro	Lys	Tyr	Ser	Leu	
1190					1195							1200			
Phe	Glu	Leu	Glu	Asn	Gly	Arg	Lys	Arg	Met	Leu	Ala	Ser	Ala	Gly	
1205					1210							1215			
Glu	Leu	Gln	Lys	Gly	Asn	Glu	Leu	Ala	Leu	Pro	Ser	Lys	Tyr	Val	
1220					1225							1230			

-continued

Asn Phe Leu Tyr Leu Ala Ser His Tyr Glu Lys Leu Lys Gly Ser 1235 1240 1245	
Pro Glu Asp Asn Glu Gln Lys Gln Leu Phe Val Glu Gln His Lys 1250 1255 1260	
His Tyr Leu Asp Glu Ile Ile Glu Gln Ile Ser Glu Phe Ser Lys 1265 1270 1275	
Arg Val Ile Leu Ala Asp Ala Asn Leu Asp Lys Val Leu Ser Ala 1280 1285 1290	
Tyr Asn Lys His Arg Asp Lys Pro Ile Arg Glu Gln Ala Glu Asn 1295 1300 1305	
Ile Ile His Leu Phe Thr Leu Thr Asn Leu Gly Ala Pro Ala Ala 1310 1315 1320	
Phe Lys Tyr Phe Asp Thr Thr Ile Asp Arg Lys Arg Tyr Thr Ser 1325 1330 1335	
Thr Lys Glu Val Leu Asp Ala Thr Leu Ile His Gln Ser Ile Thr 1340 1345 1350	
Gly Leu Tyr Glu Thr Arg Ile Asp Leu Ser Gln Leu Gly Gly Asp 1355 1360 1365	
<p><210> SEQ ID NO 503 <211> LENGTH: 15 <212> TYPE: DNA <213> ORGANISM: Homo sapiens</p>	
<400> SEQUENCE: 503	
cagaagaaga agggc	15
<p><210> SEQ ID NO 504 <211> LENGTH: 51 <212> TYPE: DNA <213> ORGANISM: Homo sapiens</p>	
<400> SEQUENCE: 504	
ccaatgggga ggacatcgat gtcacctcca atgactaggg tgggtgggcaa c	51
<p><210> SEQ ID NO 505 <211> LENGTH: 15 <212> TYPE: DNA <213> ORGANISM: Homo sapiens</p>	
<400> SEQUENCE: 505	
ctctggccac tccct	15
<p><210> SEQ ID NO 506 <211> LENGTH: 52 <212> TYPE: DNA <213> ORGANISM: Homo sapiens</p>	
<400> SEQUENCE: 506	
acatcgatgt cacctccaat gacaagcttg ctageggtgg gcaaccacaa ac	52
<p><210> SEQ ID NO 507 <211> LENGTH: 1733 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide</p>	

-continued

<400> SEQUENCE: 507

```

ccgtttaaac aattctgcag gaatctagtt attaatagta atcaattacg gggtcattag    60
ttcatagccc atatatggag ttccgcggtta cataacttac ggtaaatggc ccgcctggct    120
gaccgcccac cgacccccgc ccattgacgt caataatgac gtatgttccc atagtaacgc    180
caatagggac tttccattga cgtcaatggg tggagtattt acggtaaaact gcccaacttg    240
cagtacatca agtgtatcat atgccaagta cgccccctat tgacgtcaat gacggtaaat    300
ggcccgcctg gcattatgcc cagtacatga ccttatggga ctttctact tggcagtaca    360
tctacgtatt agtcatcgct attaccatgg tcgagggtgag ccccacgttc tgcttactc    420
tccccatctc cccccctccc ccacccccaa ttttgtattt atttattttt taattatttt    480
gtgcagcgat gggggcgggg gggggggggg ggcgcgcgcc aggcggggcg gggcggggcg    540
aggggcgggg cggggcgagg cggagagggt ggcgcgcgac caatcagagc ggcgcgctcc    600
gaaagtttcc ttttatggcg aggcgcgggc ggcgcgcgcc ctataaaaag cgaagcgcgc    660
ggcggggcga agtgcgtgcg cgtgccttc gccccgtgcc ccgctccgcc gccgcctcgc    720
gccgccccgc cgggctctga ctgaccgctt tactcccaca ggtgagcggg cgggacggcc    780
cttctcctcc gggctgtaat tagcgttgg ttaaatgacg gcttgtttct tttctgtggc    840
tgcgtaaaag ccttgagggg ctccgggagg gccctttgtg cgggggggagc ggctcggggg    900
gtgcgtgctg gtgtgtgtgc gtggggagcg ccgcgtgcgg ctccgcgctg cccggcggct    960
gtgagcgtcg cgggcgcggc ggcgggcttt gtgcgctccg cagtgtgcgc gaggggagcg    1020
cggccggggg cgggtccccc cgggtcgggg ggggctgcga ggggaacaaa ggctgcgtgc    1080
ggggtgtgtg cgtggggggg tgagcagggg gtgtgggcgc gtcggtcggg ctgcaacccc    1140
ccctgcaccc cctcccccca gttgctgagc acggccccgc ttcgggtgcg gggctccgta    1200
cggggcgtgg cgcggggctc gccgtgccgg gcgggggggtg gcggcaggtg ggggtgccgg    1260
gcggggcggg gccgcctcgg gccggggagg gctcggggga ggggcgcggc ggcccccgga    1320
gcgcggcgcg ctgtcgaggc gcggcgagcc gcagccattg ccttttatgg taatcgtgcg    1380
agagggcgca gggacttctc ttgtcccaaa tctgtgcgga gccgaaatct gggagggccc    1440
gccgcacccc ctctagcggg cgcggggcga agcgggtcgg ccggcgagg aaggaaatgg    1500
gcggggaggg ccttcgtgcg tcgcgcgcc gccgtccctc tctccctc cagcctcggg    1560
gctgtccgcg gggggacggc tgccctcggg ggggacgggg cagggcgggg ttcggcttct    1620
ggcgtgtgac cggcgctctc agacccctcg ctaaccatgt tcatgccttc ttcttttcc    1680
tacagctcct gggcaacgtg ctggttattg tgctgtctca tcattttggc aaa          1733

```

<210> SEQ ID NO 508

<211> LENGTH: 4269

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 508

```

atggactata aggaccacga cggagactac aaggatcatg atattgatta caaagacgat    60
gacgataaga tggccccaaa gaagaagcgg aaggtcggta tccacggagt cccagcagcc    120

```

-continued

gacaagaagt acagcatcgg cctggacatc ggcaccaact ctgtgggctg ggccgtgate	180
accgacgagt acaagggtgcc cagcaagaaa ttcaagggtc tgggcaacac cgaccggcac	240
agcatcaaga agaacctgat cggagccctg ctgttcgaca gcggcgaaac agccgaggcc	300
acccggctga agagaaccgc cagaagaaga tacaccagac ggaagaaccg gatctgctat	360
ctgcaagaga tcttcagcaa cgagatggcc aagggtggacg acagcttctt ccacagactg	420
gaagagtctt tcctgggtga agaggataag aagcacgagc ggcaccccat cttcggaac	480
atcgtggaag aggtggccta ccacgagaag taccaccaca tctaccacct gagaaagaaa	540
ctggtggaca gcaccgacaa ggccgacctg cggctgatct atctggccct ggcccacatg	600
atcaagttcc ggggccactt cctgatcgag ggcgacctga accccgacaa cagcgcgctg	660
gacaagctgt tcattccagct ggtgcagacc tacaaccagc tggtcgagga aaaccccatc	720
aacgccagcg gcgtggagcg caaggccatc ctgtctgcca gactgagcaa gagcagacgg	780
ctggaaaatc tgatcgccca gctgcccggc gagaagaaga atggcctgtt cggaaaacctg	840
attgccctga gcctgggctt gacccccaac ttcaagagca acttcgacct ggccgaggat	900
gccaaaactgc agctgagcaa ggacacctac gacgacgacc tggacaacct gctggcccag	960
atcggcgacc agtacgcoga cctgtttctg gccgccaaga acctgtccga cgccatcctg	1020
ctgagcgaca tcctgagagt gaaccgcgag atcaccaggg cccccctgag cgctctatg	1080
atcaagagat acgacgagca ccaccaggac ctgacctgc tgaagctct cgtgcggcag	1140
cagctgcctg agaagtacaa agagattttc ttcgaccaga gcaagaacgg ctacgccggc	1200
tacattgacg gcggagccag ccaggaagag ttctacaagt tcatcaagcc catcctgga	1260
aagatggacg gcaccgagga actgctctgt aagctgaaca gagaggacct gctgcggaag	1320
cagcggacct tcgacaacgg cagcatcccc caccagatcc acctgggaga gctgcacgcc	1380
attctgcggc ggcaggaaga ttttaacca ttctgaagg acaaccggga aaagatcgag	1440
aagatcctga ccttcgcat ccctactac gtgggccctc tggccagggg aaacagcaga	1500
ttcgctgga tgaccagaaa gagcggaggaa accatcccc cctggaactt cgaggaagtg	1560
gtggacaagg gcgcttcgcg ccagagcttc atcgagcgga tgaccaactt cgataagaac	1620
ctgcccacg agaagggtct gcccaagcac agcctgctgt acgagtactt caccgtgat	1680
aacgagctga ccaaaagtga atacgtgacc gagggaatga gaaagcccgc cttcctgagc	1740
ggcagcaga aaaaggccat cgtggacctg ctgttcaaga ccaaccgga agtgacctg	1800
aagcagctga aagaggacta cttcaagaaa atcgagtgtc tcgactccgt ggaatctcc	1860
ggcgtggaag atcggttcaa cgctccctg ggcacatacc acgatctgct gaaaattatc	1920
aaggacaagg acttcctgga caatgaggaa aacgaggaca ttctggaaga tatcgtgctg	1980
accctgacac tgtttgagga cagagagatg atcgaggaac ggctgaaac ctatgccac	2040
ctgttcgacg acaaaagtgat gaagcagctg aagcggcgga gataaccgg ctggggcag	2100
ctgagccgga agctgatcaa cggcatccgg gacaagcagt ccggcaagac aatcctggat	2160
ttcctgaagt ccgacggctt cgccaacaga aacttcatgc agctgatcca cgacgacagc	2220
ctgaccttta aagaggacat ccagaaagcc caggtgtccg gccagggcga tagcctgcac	2280
gagcacattg ccaatctggc cggcagcccc gccattaaga agggcatcct gcagacagtg	2340
aagggtggtg acgagctcgt gaaagtgatg ggccggcaca agcccagaa catcgtgatc	2400

-continued

```

gaaatggcca gagagaacca gaccacccag aaggacaga agaacagccg cgagagaatg 2460
aagcggatcg aagagggcat caaagagctg ggcagccaga tcctgaaaga acaccccgctg 2520
gaaaacaccc agctgcagaa cgagaagctg tacctgtact acctgcagaa tgggcgggat 2580
atgtacgtgg accaggaact ggacatcaac cggctgtccg actacgatgt ggaccatatac 2640
gtgcctcaga gctttctgaa ggacgactcc atcgacaaca aggtgctgac cagaagcgac 2700
aagaaccggg gcaagagcga caacgtgcc tccgaagagg tcgtgaaaga gatgaagaac 2760
tactggcggc agctgctgaa cgccaagctg attacccaga gaaagttcga caatctgacc 2820
aaggccgaga gagggcgct gagcgaactg gataaggccg gcttcatcaa gagacagctg 2880
gtggaacccc ggcagatcac aaagcagctg gcacagatcc tggactcccc gatgaacact 2940
aagtacgacg agaatgacaa gctgatccgg gaagtgaaag tgatcacctt gaagtccaag 3000
ctggtgtccg atttccgtaa ggatttccag ttttacaag tgcgcgagat caacaactac 3060
caccacgccc acgacgccta cctgaacgcc gtcgtgggaa ccgcccctgat caaaaagtac 3120
cctaagctgg aaagcgagtt cgtgtacggc gactacaagg tgtacgacgt gcggaagatg 3180
atcgccaaga gcgagcagga aatcggaag gctaccgcca agtacttctt ctacagcaac 3240
atcatgaaact ttttcaagac cgagattacc ctggccaacg gcgagatccg gaagcggcct 3300
ctgatcgaga caaacggcga aaccggggag atcgtgtggg ataagggccg ggattttgcc 3360
accgtgcgga aagtgtctgag catgccccaa gtgaatatcg tgaaaaagac cgaggtgcag 3420
acagggcgct tcagcaaaga gtctatctcg cccaagagga acagcgataa gctgatcgcc 3480
agaaaagaag actgggaccc taagaagtac ggcggcttcg acagccccac cgtggcctat 3540
tctgtctggtg tgggtggccaa agtggaaaag ggcaagtcca agaaactgaa gagtgtgaaa 3600
gagctgctgg ggatcaccat catggaaga agcagcttcg agaagaatcc catcgacttt 3660
ctggaagcca agggctacaa agaagtgaaa aaggacctga tcatcaagct gcctaagtac 3720
tcccctgttcg agctggaaaa cggccggaag agaagtctgg cctctgccgg cgaactgcag 3780
aagggaaaag aactggccct gccctccaaa tatgtgaact tcctgtacct ggccagccac 3840
tatgagaagc tgaagggctc ccccaggat aatgagcaga aacagctgtt tgtggaacag 3900
cacaagcact acctggacga gatcatcgag cagatcagcg agttctccaa gagagtgatc 3960
ctggccgacg ctaatctgga caaagtctg tccgcctaca acaagcaccg ggataagccc 4020
atcagagagc agggcgagaa tatcatccac ctgtttaccc tgaccaatct gggagcccct 4080
gccgccttca agtactttga caccaccatc gaccggaaga ggtacaccag caccaaagag 4140
gtgctggacg ccaccctgat ccaccagac atcaccggcc tgtacgagac acggatcgac 4200
ctgtctcagc tgggagcgca caaaaggccg gcggccacga aaaaggccgg ccaggcaaaa 4260
aagaaaaag 4269

```

<210> SEQ ID NO 509

<211> LENGTH: 780

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 509

```

ggaagcggag ccactaactt ctcctgttg aaacaagcag gggatgtcga agagaatccc 60

```

-continued

```

gggccagtga gcaagggcga ggagctgttc accggggtgg tgcccatcct ggtcgagctg 120
gacggcgaag taaacggcca caagttcage gtgtccggcg agggcgaggg cgatgccacc 180
tacggcaagc tgaccctgaa gttcatctgc accaccggca agctgcccgt gccctggccc 240
accctcgtga ccaccctgac ctacggcgtg cagtgttca gccgctaccc cgaccacatg 300
aagcagcaag acttcttcaa gtccgccatg cccgaaggct acgtccagga gcgcaccatc 360
ttcttcaagg acgacggcaa ctacaagacc cgcgccgagg tgaagttcga gggcgacacc 420
ctggtgaacc gcacgcagct gaagggcacc gacttcaagg aggacggcaa catcctgggg 480
cacaagctgg agtacaacta caacagccac aacgtctata tcatggccga caagcagaag 540
aacggcatca aggtgaaact caagatccgc cacaacatcg aggacggcag cgtgcagctc 600
gccgaccact accagcagaa cccccccatc ggcgacggcc ccgtgctgct gcccgacaac 660
cactacctga gcaccagtc cgccctgagc aaagacccca acgagaagcg cgatcacatg 720
gtcctgctgg agttcgtgac cgcgcgctgg atcactctcg gcatggacga gctgtacaag 780

```

```

<210> SEQ ID NO 510
<211> LENGTH: 597
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        polynucleotide

```

```

<400> SEQUENCE: 510
cgataatcaa cctctggatt acaaaatttg tgaagattg actggtattc ttaactatgt 60
tgctcctttt acgctatgtg gatacgtgct ttaaatgcct ttgtatcatg ctattgcttc 120
ccgatatggt ttcattttct cctccttgta taaatcctgg ttgctgtctc tttatgagga 180
gttgtggccc gttgtcaggc aacgtggcgt ggtgtgcact gtgtttgctg acgcaacccc 240
cactggttgg ggcattgcca ccacctgtca gctcctttcc gggactttcg ctttcccctc 300
ccctattgcc acggcggaac tcategcgcg ctgccttgcc cgctgctgga caggggctcg 360
gctgttgggc actgacaatt ccgtggtgtt gtcggggaaa tcatgctcct tccttggtct 420
gctgcctctg gttgccacct ggattctgag cgggacgtcc ttctgctacg tccttcggc 480
cctcaatcca gcggaccttc cttcccggcg cctgctgccc gctctgcggc ctcttcggcg 540
tcttcgctt cgccctcaga cgagtcggat ctccccttgg gccgcctccc cgcatcg 597

```

```

<210> SEQ ID NO 511
<211> LENGTH: 210
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        polynucleotide

```

```

<400> SEQUENCE: 511
cgacctgac tgtgccttct agttgccagc catctgttgt ttgcccctcc cccgtgcctt 60
ccttgacctt ggaaggtgcc actcccactg tcctttccta ataaatgag gaaattgcat 120
cgcattgtct gagtagtgt cattctatc tggggggtgg ggtggggcag gacagcaag 180
gggaggattg ggaagacaat ggcaggcatg 210

```

-continued

```

<210> SEQ ID NO 512
<211> LENGTH: 906
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      polynucleotide
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (109)..(109)
<223> OTHER INFORMATION: a, c, t, g, unknown or other
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (135)..(135)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 512

ataacttcgt ataatgtatg ctatacgaag ttattcgcgga tgaataaatg aaagcttgca    60
gatctgcgac tctagaggat ctgcgactct agaggatcat aatcagcncnt accacatttt    120
gtagagggtt tactngcttt aaaaaacctc ccacacctcc ccctgaaact gaaacataaa    180
atgaatgcaa ttgttgttgt taacttggtt attgcagctt ataatggta caaataaagc    240
aatagcatca caaatctcac aaataaagca tttttttcac tgcattctag ttgtggtttg    300
tccaaactca tcaatgtatc ttatcatgtc tggatctgog actctagagg atcataatca    360
gccataccac attttagagag gttttacttg ctttaaaaaa cctcccacac ctccccctga    420
acctgaaaca taaaatgaat gcaattgttg ttgtaactt gtttattgca gcttataatg    480
gttacaataa aagcaatagc atcacaattc tcacaaataa agcatttttt tcaactgcatt    540
ctagttgtgg tttgtccaaa ctcatcaatg tatcttatca tgtctggatc tgcgactcta    600
gaggatcata atcagccata ccacatttgt agaggtttta cttgctttaa aaaacctccc    660
acacctcccc ctgaacctga aacataaaat gaatgcaatt gttgttgta acttgtttat    720
tgcagcttat aatggttaca aataaagcaa tagcatcaca aatttcacaa ataaagcatt    780
tttttcaact cattctagtt gtggtttgtc caaactcacc aatgtatcct atcatgtctg    840
gatccccatc aagctgatcc ggaaccctta atataacttc gtataatgta tgctatacga    900
agttat                                           906

```

```

<210> SEQ ID NO 513
<211> LENGTH: 1079
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      polynucleotide

<400> SEQUENCE: 513

caggccctcc gagcgtggtg gagccgttct gtgagacagc cgggtacgag tctgacgct    60
ggaaggggca agcgggtggt gggcaggaat gcggtccgcc ctgcagcaac cggaggggga    120
gggagaaggg agcggaaaag tctccaccgg acgcggccat ggctcggggg ggggggggca    180
gcggaggagc gcttccggcc gacgtctcgt cgctgattgg cttcttttcc tcccgccgtg    240
tgtgaaaaca caaatggcgt gttttggttg gcgtaaggcg cctgtcagtt aacggcagcc    300
ggagtgcgca gcccccgcca gctctgctct gcccaactggg tggggcgggg ggtaggtggg    360
gtgaggcgag ctggacgtgc gggcgcggtc ggcctctggc ggggcggggg aggggagggg    420
gggtcagcga aagtagctcg cgcgcgagcg gccgccacc ctccccttcc tctgggggag    480

```

-continued

tcgttttacc	cgccgcccgc	cgggcctcgt	cgctcgattg	gctctcgggg	cccagaaaaac	540
tggcccttgc	cattggctcg	tgctcgtgca	agttgagtc	atccgcccgc	cagcgggggc	600
ggcgaggagg	cgctcccagg	ttccggcct	cccctcggcc	ccgcgcccga	gagtctggcc	660
gcgcgcccct	gcgcaacgtg	gcaggaagcg	cgcgctgggg	gcggggacgg	gcagtagggc	720
tgagcggctg	cggggcgggt	gcaagcacgt	ttccgacttg	agttgcctca	agagggcgct	780
gctgagccag	acctccatcg	cgcactccgg	ggagtggagg	gaaggagcga	gggctcagtt	840
gggctgtttt	ggaggcagga	agcacttgct	ctcccaaagt	cgctctgagt	tgttatcagt	900
aagggagctg	cagtgagta	ggcggggaga	aggccgcacc	cttctccgga	ggggggaggg	960
gagtgttgca	atacctttct	gggagttctc	tgctgcctcc	tggtttctga	ggaccgccct	1020
gggctcggga	gaatcccttc	cccctcttcc	ctcgtgatct	gcaactccag	tctttctag	1079

<210> SEQ ID NO 514

<211> LENGTH: 4336

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 514

agatgggccc	gagctctctg	ggcaggctta	aaggctaacc	tggtgtgtgg	gcgttgcct	60
gcaggggaat	tgaacagggt	taaaattgga	gggacaagac	ttcccacaga	ttttcggtt	120
tgctcgggaag	tttttaata	ggggcaaata	aggaaaatgg	gaggataggt	agtcactctg	180
ggttttatgc	agcaaaaacta	cagggttatta	ttgcttctga	tccgcctcgg	agtattttcc	240
atcgaggtag	attaagaca	tgtcaccocg	agttttatac	tctcctgctt	gagatcctta	300
ctacagtatg	aaattacagt	gtcgcgagtt	agactatgta	agcagaat	taatcatttt	360
taaagagccc	agtacttcoat	atccatttct	cccgcctcct	ctgcagcctt	atcaaaagg	420
attttagaac	actcatttta	gccccatttt	catttattat	actggcttat	ccaaccctta	480
gacagagcat	tggcattttc	cctttcctga	tcttagaagt	ctgatgactc	atgaaaccag	540
acagattagt	tacatacacc	acaatcogag	gctgtagctg	gggcctcaac	actgcagttc	600
ttttataact	ccttagtaca	ctttttgttg	atcctttgcc	ttgatcctta	attttcagtg	660
tctatcacct	ctcccgtcag	gtgggtgtcc	acatttgggc	ctattctcag	tccagggagt	720
tttacaacaa	tagatgtatt	gagaatccaa	cctaaagctt	aactttccac	tcccataaat	780
gcctctctcc	ttttctccca	tttataaact	gagctattaa	ccattaatgg	tttccaggtg	840
gatgtctcct	cccccaatat	tacctgatgt	atcttacata	ttgccaggct	gatattttaa	900
gacattaaaa	ggatattttc	attattgagc	cacatgggat	tgattactgc	ttactaaaat	960
tttgtcattg	tacacatctg	taaaagggtg	ttccttttgg	aatgcaaaagt	tcaggtgttt	1020
gttgtctttc	ctgacctaa	gtcttctgag	cttgtatttt	ttctatttaa	gcagtgcttt	1080
ctcttggaact	ggcttgactc	atggcattct	acacgttatt	gctggctctaa	atgtgatttt	1140
gccaagcttc	ttcaggacct	ataattttgc	ttgacttgta	gccaacacaca	agtaaaatga	1200
ttaagcaaca	aatgtatttg	tgaagcttgg	tttttaggtt	gttggtgtgt	gtgtgcttgt	1260
gctctataat	aatactatcc	aggggctgga	gaggtggctc	ggagtccaag	agcacagact	1320

-continued

gctcttccag aagtcctgag tccaattccc agcaaccaca tgggtggetca caaccatctg	1380
taatgggatac tgatgccttc ttctgggtg tctgaagacc acaagtgtat tcacattaaa	1440
taaataaate ctccttcttc ttcttttttt ttttttttaa gagaatactg tctccagtag	1500
aatttactga agtaatgaaa tactttgtgt ttgttccaat atggtagcca ataatacaat	1560
tactctttaa gcactggaaa tgttaccacg gaactaattt ttatttgaag tgtaactgtg	1620
gacagaggag ccataactgc agacttgtgg gatacagaag accaatgcag actttaatgt	1680
cttttctctt acactaagca ataaagaaat aaaaattgaa cttctagtat cctatttgtt	1740
taactgcta gctttactta acttttgtgc ttcactata caaagctgaa agctaagtct	1800
gcagccatta ctaaacatga aagcaagtaa tgataatttt ggatttcaaa aatgtagggc	1860
cagagttag ccagccagtg gtgggtgctg cctttatgcc ttaataccca gcactctgga	1920
ggcagagaca ggcagatctc tgagtttgag cccagcctgg tctacacatc aagttctatc	1980
taggatagcc aggaatacac acagaaaccc tgttggggag gggggctctg agatttcata	2040
aaattataat tgaagcattc cctaatgagc cactatggat gtggctaaat cctgtctact	2100
ttctgatgag atttgggtat tattttttct gtctctgctg ttggttgggt cttttgacac	2160
tgtgggcttt ctttaaagcc tccttctgc catgtggtct cttgtttgct actaaecttc	2220
catggcctaa atggcatggc tttttgctt ctaagggcag ctgctgagat ttgcagcctg	2280
atctccaggg tggggttggg aaactcttca aacactaaaa ttgtccttta attttttttt	2340
taaaaaatgg gttatataat aaacctcata aaatagttat gaggagttag gtggactaat	2400
attaatgag tccctccct ataaaagagc tattaaggct tttgtctta tacttaactt	2460
tttttttaa tgtggatctc ttagaaccaa gggctctaga gtttttagtat acagaaactg	2520
ttgcatcgtc taatcagatt ttctagttc aaatccagag aatccaaatt cttcacagcc	2580
aaagtcaaat taagaatttc tgacttttaa tgttaatttg cttactgtga atataaaat	2640
gatagctttt cctgaggcag ggtctcacta tgtatctctg cctgatctgc aacaagatat	2700
gtagactaaa gttctgctg cttttgtctc ctgaatacta aggttaaaat gtagtaatac	2760
ttttggaact tgcaggtcag attcttttat aggggacaca ctaagggagc ttgggtgata	2820
gttggtaaat gtgtttaagt gatgaaaact tgaattatta tcaccgcaac ctacttttta	2880
aaaaaaaaag ccaggcctgt tagagcatgc ttaagggatc cctaggactt gctgagcaca	2940
caagagtagt tacttggcag gctcctggtg agagcatatt tcaaaaaaca aggcagacaa	3000
ccaagaaact acagttaagg ttacctgtct ttaaaccatc tgcataatac cagggatatt	3060
aaaatattcc aaataatatt tcattcaagt tttccccat caaattggga catggatttc	3120
tccggtgaat aggcagagtt gaaaactaaa caaatgttg ttttgtgatt tgtgaaattg	3180
ttttcaagtg atagttaaag cccatgagat acagaacaaa gctgctattt cgaggctctc	3240
tggtttatac tcagaagcac ttctttgggt ttcctgcac tctctgac atgtgctagg	3300
cctaccttag gctgattgtt gttcaataa acttaagttt cctgtcaggt gatgtcatat	3360
gatttcatat atcaaggcaa aacatgttat atatgttaa catttgtact taatgtgaaa	3420
gttaggtctt tgtgggtttg atttttaatt ttcaaacct gagctaaata agtcattttt	3480
acatgtctta catttgggtg aattgtataa ttgtgggttg caggcaagac tctctgacct	3540
agtaacccta cctatagagc actttgctgg gtcacaagtc taggagtcaa gcatttcacc	3600

-continued

ttgaagttga gacgttttgt tagtgatac tagtttatat gttggaggac atgtttatcc	3660
agaagatatt caggactatt tttgactggg ctaaggaatt gattctgatt agcactgtta	3720
gtgagcattg agtggccttt aggcctgaat tggagtcact tgatatctc aaataatgct	3780
ggcctttttt aaaagccctt gttctttatc accctgtttt ctacataatt tttgttcaaa	3840
gaaatacttg tttggatctc cttttgacaa caatagcatg ttttcaagcc atattttttt	3900
tccttttttt tttttttttt ggttttttga gacagggttt ctctgtatag ccttggtgt	3960
cctggaactc actttgtaga ccaggctggc ctggaactca gaaatccgcc tgcctctgcc	4020
tcctgagtgc cgggattaaa ggcgtgcacc accacgcctg gctaagttgg atattttgtt	4080
atataactat aaccaatact aactccactg ggtggatttt taattcagtc agtagtotta	4140
agtgtctttt attggccctt cattaaaatc tactgttcac tctaacagag gctggtgta	4200
ctagtggcac ttaagcaact tcctacggat atactagcag attaagggtc agggatagaa	4260
actagtctag cgttttgtat acctaccagc tttatactac cttgttctga tagaaatatt	4320
tcaggacatc tagctt	4336

<210> SEQ ID NO 515

<211> LENGTH: 1846

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 515

aattctacgg ggtaggggag ggccttttcc caaggcagtc tggagcatgc gcttttagcag	60
ccccgctggg cacttggggc tacacaagtg gcctctggcc tcgcacacat tccacatcca	120
ccggtaggcg ccaaccggct cegtcttttg gtggcccctt cgcgccacct tctactcctc	180
ccctagtcag gaagtcccc ccgcgccgcg agctcgcgtc gtgcaggacg tgacaaatgg	240
aagtagcacg tctcactagt ctctgcaga tggacagcac cgtgagcaa tggaaagcggg	300
taggcctttg gggcagcggc caatagcagc tttgtctcct cgctttctgg gctcagaggc	360
tgggaagggg tgggtccggg ggcgggctca ggggcgggct caggggcggg gcgggcgccc	420
gaagtcctc cggaggcccg gcattctgca cgcttcaaaa gcgcacgtct gccgcctgt	480
tctctcttc ctcatctccg ggccttttga cctgcaatcg ccgctagcga agttcctatt	540
ctctagaaag tataggaact tcgccaccat gggatcggcc attgaacaag atggattgca	600
cgcaggttct ccggccgctt gggtgagag gctattcggc tatgactggg cacaacagac	660
aateggctgc tctgatccg cegtgtccg gctgtcagcg caggggcgcc cggttctttt	720
tgtcaagacc gacctgtccg gtgccctgaa tgaactgcag gacgaggcag cgcggctatc	780
gtggctggcc acgacggggc ttccttgcgc agctgtgctc gacgttgta ctgaagcggg	840
aagggactgg ctgctattgg gcgaagtgcc ggggcaggat ctctgtcat ctcaccttgc	900
tcctgccgag aaagtatcca tcattgctga tgcaatgcgg cggctgcata cgcttgatcc	960
ggctacctgc ccattcgacc accaagcga acatcgcac gacgagcac gtactcggat	1020
ggaagccggg cttgtcgatc aggatgatct ggacgaagag catcaggggc tcgcccagc	1080
cgaactgttc gccaggctca agcgcgcgat gcccgacggc gatgatctcg tcgtgacca	1140
tggcgatgoc tgcttgccga atatcatggt ggaaaatggc cgcttttctg gattcatcga	1200

-continued

```

ctgtggccgg ctgggtgtgg cggaccgcta tcaggacata gcgttggcta cccgtgatat 1260
tgctgaagag cttggcggcg aatgggctga ccgcttctc gtgctttacg gtatcgccgc 1320
tcccgattcg cagcgcacgc ccttctatcg ccttcttgac gagttcttct gaggggatcc 1380
gctgtaagtc tgcagaaatt gatgatctat taaacaataa agatgtccac taaaatggaa 1440
gtttttctg tcatactttg ttaagaaggg tgagaacaga gtacctacat tttgaatgga 1500
aggattggag ctacgggggt gggggtgggg tgggattaga taaatgcctg ctctttactg 1560
aaggctcttt actattgctt tatgataatg tttcatagtt ggatatcata atttaacaa 1620
gcaaaccxaa attaagggcc agctcattcc tcccactcat gatctataga tctatagatc 1680
tctcgtggga tcattgtttt tctcttgatt cccactttgt ggttctaagt actgtggttt 1740
ccaaatgtgt cagtttcata gctgaagaa cgagatcagc agcctctgtt ccacatacac 1800
ttcattctca gtattgtttt gccaaattct aattccatca gaaagc 1846

```

<210> SEQ ID NO 516

<211> LENGTH: 1519

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 516

```

taccgggtag gggaggcgt tttcccaagg cagtctggag catgcgcttt agcagccccg 60
ctgggcactt ggcgctacac aagtggctc tggcctcgca cacattccac atccaccggt 120
aggcgccaac cggctccggt ctttggtggc cccttcgccc caccttctac tcctccccta 180
gtcaggaagt tccccccgc cccgcagtc gcgtcgtgca ggacgtgaca aatggaagta 240
gcacgtctca ctagtctcgt gcagatggac agcacccgtg agcaatggaa gcgggtaggc 300
ctttggggca gcgccaata gcagctttgc tccttcgctt tctgggctca gaggctggga 360
aggggtgggt ccggggcggt gctcaggggc gggctcaggg gcggggcggg cgcccgaagg 420
tcctccggag gcccggcatt ctgcacgctt caaaagcgca cgtctgcccg gctgttctcc 480
tcttctctat ctccgggcct ttgcacctgc aggtctctgc catggatcct gatgatgttg 540
ttgattcttc taaatctttt gtgatggaaa acttttcttc gtaccacggg actaaaactg 600
gttatgtaga ttccattcaa aaaggtatac aaaagccaaa atctggtaca caaggaaatt 660
atgacgatga ttgaaaaggg ttttatagta ccgacaataa atacgacgct gcgggatact 720
ctgtagataa tgaaaaccgc ctctctggaa aagctggagg cgtgggtcaaa gtgacgtatc 780
caggactgac gaaggttctc gcactaaaag tggataatgc cgaaactatt aagaaagagt 840
taggtttaag tctcactgaa ccgttgatgg agcaagtcgg aacggaagag tttatcaaaa 900
ggttcgtgta tgggtcttcg cgtgtagtgc tcagccttcc cttecgctgag gggagttcta 960
gcgttgaata tattaataac tgggaacagg cgaaagcgtt aagcgtagaa cttgagatta 1020
atthtgaaac ccgtgaaaaa cgtggccaag atgcatgata tgagtatatg gctcaagcct 1080
gtgcaggaaa tcgtgtcagg cgatctcttt gtgaaggaac cttacttctg tgggtgacaa 1140
taattggaca aactacctac agagatttaa agctctaagg taaatataaa atthttaaagt 1200
gtataatgtg ttaaactact gattctaatt gtttgtgtat tttagattcc aacctatgga 1260

```

-continued

```

actgatgaat gggagcagtg gtggaatgca gatcctagag ctcgctgac agcctcgact 1320
gtgccttcta gttgccagcc atctgttgtt tgcccctccc ccgtgccttc cttgaccctg 1380
gaagtgcca ctcccactgt cctttcctaa taaaatgagg aaattgcac gcattgtctg 1440
agtaggtgc attctattct ggggggtggg gtggggcagg acagcaagg ggaggattgg 1500
gaagacaata gcaggcatg 1519

```

```

<210> SEQ ID NO 517
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        primer

```

```

<400> SEQUENCE: 517

```

```

gagggcctat ttcccatgat tcc 23

```

```

<210> SEQ ID NO 518
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        oligonucleotide

```

```

<400> SEQUENCE: 518

```

```

cttgtgaaa ggacgaaaca cc 22

```

```

<210> SEQ ID NO 519
<211> LENGTH: 45
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        oligonucleotide
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (4)..(23)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

```

```

<400> SEQUENCE: 519

```

```

aacnnnnnnn nnnnnnnnnn nnnngtgttt cgtcctttcc acaag 45

```

```

<210> SEQ ID NO 520
<211> LENGTH: 28
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        oligonucleotide

```

```

<400> SEQUENCE: 520

```

```

cctgagtgtt gagccccag tggtgtct 28

```

```

<210> SEQ ID NO 521
<211> LENGTH: 37
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        oligonucleotide

```

```

<400> SEQUENCE: 521

```


-continued

acgagggcag agtgctgctt gctgctggcc aggeccc 37

<210> SEQ ID NO 522
 <211> LENGTH: 68
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 oligonucleotide

<400> SEQUENCE: 522

catcaggctc tcagctcagc ctgagtgttg aggcctgct ggccaggccc ctgctgggc 60

ccaagctg 68

<210> SEQ ID NO 523
 <211> LENGTH: 27
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 oligonucleotide

<400> SEQUENCE: 523

gagggcctat ttcccatgat tccttca 27

<210> SEQ ID NO 524
 <211> LENGTH: 125
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 polynucleotide
 <220> FEATURE:
 <221> NAME/KEY: modified_base
 <222> LOCATION: (84)..(103)
 <223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 524

aaaaaaagca ccgactcggg gccacttttt caagttgata acggactagc cttattttaa 60

cttgctatct ctgactctaa aacnnnnnnn nnnnnnnnnn nnnnggtgtt cgtcctttcc 120

acaag 125

<210> SEQ ID NO 525
 <211> LENGTH: 111
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 polynucleotide
 <220> FEATURE:
 <221> NAME/KEY: modified_base
 <222> LOCATION: (9)..(28)
 <223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 525

gaaacaccnn nnnnnnnnnn nnnnnnnngt tttagagcta gaaatagcaa gttaaaataa 60

ggctagtcg ttatcaactt gaaaaagtgg caccgagtcg gtgctttttt t 111

<210> SEQ ID NO 526
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 oligonucleotide

<400> SEQUENCE: 526

agctgtttta ctggtcggct 20

<210> SEQ ID NO 527
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 oligonucleotide

<400> SEQUENCE: 527

aatggataca cctggtcgaa 20

<210> SEQ ID NO 528
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 oligonucleotide

<400> SEQUENCE: 528

caatggatag acctggtcga 20

<210> SEQ ID NO 529
 <211> LENGTH: 68
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 529

accatgtata ccacttgggc ttggcagta gctaactgca ctaaataaa tataaggagg 60

gttttatg 68

<210> SEQ ID NO 530
 <211> LENGTH: 23
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 oligonucleotide

<220> FEATURE:
 <221> NAME/KEY: modified_base
 <222> LOCATION: (1)..(20)
 <223> OTHER INFORMATION: a, c, t or g

<220> FEATURE:
 <221> NAME/KEY: modified_base
 <222> LOCATION: (21)..(21)
 <223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 530

nnnnnnnnnn nnnnnnnnnn ngg 23

<210> SEQ ID NO 531
 <211> LENGTH: 15
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 oligonucleotide

<220> FEATURE:

-continued

<221> NAME/KEY: modified_base
<222> LOCATION: (1)..(12)
<223> OTHER INFORMATION: a, c, t or g
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (13)..(13)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 531

nnnnnnnnnn nnnngg 15

<210> SEQ ID NO 532
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (1)..(20)
<223> OTHER INFORMATION: a, c, t or g
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (21)..(21)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 532

nnnnnnnnnn nnnnnnnnnn ngg 23

<210> SEQ ID NO 533
<211> LENGTH: 14
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (1)..(11)
<223> OTHER INFORMATION: a, c, t or g
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (12)..(12)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 533

nnnnnnnnnn nngg 14

<210> SEQ ID NO 534
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (1)..(20)
<223> OTHER INFORMATION: a, c, t or g
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (21)..(21)
<223> OTHER INFORMATION: a, c, t, g, unknown or other
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (24)..(24)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 534

-continued

nnnnnnnnnn nnnnnnnnn nggng 25

<210> SEQ ID NO 535
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (1)..(12)
<223> OTHER INFORMATION: a, c, t or g
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (13)..(13)
<223> OTHER INFORMATION: a, c, t, g, unknown or other
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (16)..(16)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 535

nnnnnnnnnn nnnngng 17

<210> SEQ ID NO 536
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (1)..(20)
<223> OTHER INFORMATION: a, c, t or g
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (21)..(21)
<223> OTHER INFORMATION: a, c, t, g, unknown or other
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (24)..(24)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 536

nnnnnnnnnn nnnnnnnnn nggng 25

<210> SEQ ID NO 537
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (1)..(11)
<223> OTHER INFORMATION: a, c, t or g
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (12)..(12)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

-continued

<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (15)..(15)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 537

nnnnnnnnn nnggng

16

What is claimed is:

1. A method comprising:

introducing into, or expressing in, a eukaryotic cell having a DNA molecule,

(I) a Cas9 protein or a nucleotide sequence encoding the Cas9 protein, and

(II) RNA or a nucleotide sequence encoding the RNA, the RNA comprising:

(a) a first RNA comprising a first ribonucleotide sequence and a second ribonucleotide sequence, and

(b) a second RNA,

wherein (II) (a) and (II) (b) are fused to one another or are covalently linked to one another with intervening nucleotides; and

wherein, in the eukaryotic cell, the second RNA forms an RNA duplex with the second ribonucleotide sequence, and the first ribonucleotide sequence directs the Cas9 protein to a target sequence of the DNA molecule, and the DNA molecule is cleaved or edited or at least one product of the DNA molecule is altered.

* * * * *