



- (51) **International Patent Classification:**
C05F 11/00 (2006.01) *C05F 15/00* (2006.01)
- (21) **International Application Number:**
PCT/US2013/076858
- (22) **International Filing Date:**
20 December 2013 (20.12.2013)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**
61/739,869 20 December 2012 (20.12.2012) US
61/834,544 13 June 2013 (13.06.2013) US
- (71) **Applicant:** RHODIA OPERATIONS [FR/FR]; 40 rue de la Haie Coq, F-93306 Aubervilliers (FR).
- (72) **Inventors; and**
- (71) **Applicants :** ORTIZ-SUAREZ, Marivi [US/US]; 37 Steeplechase Blvd., Burlington, NJ 08016 (US). RUCH, Thomas [US/US]; 6124 Main Street, Voorhees, NJ 08043 (US). SHANMUGA, Krish [US/US]; 115 Hampshire Dr., Plainsboro, NJ 08536 (US).
- (74) **Agent:** McVEIGH, Kevin, E.; Solvay USA Inc., 8 Cedar Brook Drive, CN 7500, Cranbury, NJ 08512 (US).
- (81) **Designated States** (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM,

AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

- (84) **Designated States** (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- *with international search report (Art. 21(3))*
- *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))*



WO 2014/100561 A1

(54) **Title:** LIQUID DICYANDIAMIDE AND/OR ALKYL THIOPHOSPHORIC TRIAMIDE COMPOSITIONS AND THEIR USE IN AGRICULTURAL APPLICATIONS

(57) **Abstract:** An inhibitor composition contains dicyandiamide as a nitrification inhibitor, alkyl thiophosphoric triamide as a urease inhibitor, or a combination thereof, dissolved in a liquid medium comprising an organic solvent selected from, among others, one or more polar aprotic solvents, amine solvents, heterocyclic alcohol solvents, and mixtures thereof, is useful in making fertilizer compositions and in a method of fertilizing target plants.

LIQUID DICYANDIAMIDE AND/OR ALKYL THIOPHOSPHORIC TRIAMIDE COMPOSITIONS AND THEIR USE IN AGRICULTURAL APPLICATIONS

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the benefit of US Provisional Patent Application No. 61/739,869 filed December 20, 2012, incorporated herein by reference in its entirety, and claims the benefit of US Provisional Patent Application No. 61/834,544 filed June 13, 2013, incorporated herein by reference in its entirety.

FIELD OF THE INVENTION

[0002] This invention relates to liquid compositions comprising dicyandiamide and/or an alkyl thiophosphoric triamide, methods for incorporating dicyandiamide into agricultural fertilizer compositions, agricultural fertilizer compositions comprising dicyandiamide and/or an alkyl thiophosphoric triamide, and the use of such compositions.

BACKGROUND OF THE INVENTION

[0003] In the agrochemical industry, farmers use various fertilizers to impart macronutrients to plants either by application to the soil or application to plant leaves. Nitrogen, phosphorus, potassium, calcium, magnesium, and sulfur are macronutrients that must be supplied to the plants and soil manually by farmers. In many crops, the amount of nitrogen supplied is critical to the overall quality and growth of the crop. Nitrogen is typically supplied in the form of nitrogenous, i.e., nitrogen precursor-containing, fertilizer compounds, such as urea, ammonium nitrate, or ammonium phosphate fertilizer compounds. Due to the high water solubility of these salts, however, applied nitrogen values may be lost due to run-off and leaching of the nitrogenous fertilizer compounds. Once applied, the nitrogenous fertilizer compounds are typically degraded, for example, by microorganisms present in the soil, to

nitrogenous species such as NH_4^+ , NO_2^- , NO_3^- , and ammonia gas, that may be even more readily lost through evaporation, run-off, and leaching than the fertilizer compounds themselves. If degradation of the fertilizer compounds occurs at a rate that is faster than the nitrogenous degradation products can be used by the plants, then the nitrogen values in the degradation products are at increased risk of being lost.

[0004] Nitrification and/or urease inhibitors are of potential use in delaying degradation of fertilizer compounds and thereby reducing losses of nitrogenous degradation products that would otherwise occurred in the absence of the inhibitors. The use of nitrification and/or urease inhibitors in combination with nitrogenous fertilizer compounds tends to increase the amount of time the nitrogen source remains in the soil and available for absorption by the plants, which tends to increase the effectiveness of the fertilizer and positively impact crop yield and quality.

[0005] Aqueous end use fertilizer solutions are typically prepared in the field by diluting commercially available concentrated fertilizer compositions with water. Commonly used concentrated fertilizer compositions include concentrated ammonium nitrate compositions, such as, for example, UAN 18, UAN 28, UAN 30 and UAN 32.

[0006] Dicyandiamide is potentially useful as a nitrification inhibitor in such aqueous end use fertilizer compositions, but has very low solubility (about 41 grams per liter ("g/l")) in water and so is difficult to incorporate into the aqueous end use fertilizer compositions, particularly under field conditions

SUMMARY OF THE INVENTION

[0007] Urease inhibitors can be used with a fertilizer (i.e., incorporated into a urea-containing fertilizer, e.g., urea and urea ammonium nitrate (UAN)) to slow the conversion of ammonium to ammonia gas and thus slow the loss of ammonia to volatilization, thus making ammonium available to plants in the soil for longer periods of

time. Nitrification inhibitors can be used with a fertilizer (i.e., incorporated into a urea-containing fertilizer, e.g., urea and urea ammonium nitrate (UAN)) to slow the process of ammonium conversion to nitrate, and subsequently the loss of nitrate to leaching, thus making ammonium available to plants in the soil for longer periods of time. Ammonium is one of the main forms of nitrogen that can be utilized by plants. Increasing the amount of time that the nitrogen is available to the plant increases the effectiveness of the fertilizer which positively impacts crop yield and quality.

[0008] Fertilizers, in one embodiment, are common water soluble inorganic fertilizers that provide nutrients such as phosphorus-based, nitrogen-based, potassium-based or sulphur-based fertilizers. Examples of such fertilizers include: for nitrogen as the nutrient: nitrates and or ammonium salts such as ammonium nitrate, including in combination with urea e.g. as Uram type materials, calcium ammonium nitrate, ammonium sulphate nitrate, ammonium phosphates, particularly mono-ammonium phosphate, di-ammonium phosphate and ammonium polyphosphate, ammonium sulphate, and the less commonly used calcium nitrate, sodium nitrate, potassium nitrate and ammonium chloride. It is understood that a fertilizer composition can comprise one or a combination of the fertilizers described herein.

[0009] A typical urease inhibitor, NBPT (N-(n-butyl)-thiophosphoric triamide), however, faces drawbacks in its use as NBPT is extremely difficult to handle. NBPT is a sticky, waxy, heat and water sensitive material, which cannot be used in its solid form, as it is used at low concentrations making it difficult to evenly distribute on urea prills (i.e., large granules) and in soil. In order to evenly distribute the NBPT onto the urea, the NBPT should be dispersed into a carrier prior to being sprayed onto the urea. Thus, the use of a solvent system containing the NBPT is desirable as, in its liquid form, the solvent system is capable of distributing the NBPT into granular urea (e.g., urea prills) and into liquid fertilizers containing urea. By introducing the NBPT to liquid fertilizers containing urea (for example, urea-ammonium nitrate solutions or UAN) in a solvent system, the NBPT is capable of being better dispersed in the liquid fertilizer.

[0010] Dicyandiamide is useful as a nitrification inhibitor in aqueous agricultural applications, e.g., end use fertilizer compositions, but similar to urease inhibitors face similar drawbacks. Nitrification inhibitors, such as dicyandiamide, generally have very low solubility (about 41 grams per liter (“g/l”)) in water and so it is difficult to incorporate into the aqueous end use fertilizer compositions, particularly under field conditions. As nitrification inhibitors, such as dicyandiamide, have a generally low solubility, they are used at low concentrations in water making it difficult to evenly distribute on urea-containing prills (i.e., large granules) and in soil. In order to evenly distribute the dicyandiamide onto the urea-containing prills or granules, dicyandiamide should be dispersed into a solvent carrier prior to being sprayed onto the urea. Thus, the use of a solvent system containing dicyandiamide (herein, also termed “DCD”) is desirable as, in its liquid form, the solvent system is capable of distributing the dicyandiamide onto urea granules or prills, urea ammonium nitrate granules or prills or, otherwise, urea-containing granules or prills, and into liquid fertilizers containing urea or urea ammonium nitrate. By introducing the dicyandiamide to liquid fertilizers containing urea (for example, urea-ammonium nitrate solutions or UAN) in a solvent system, the dicyandiamide is capable of being better dispersed in the liquid fertilizer.

[0011] In one embodiment, concentrated fertilizer compositions include concentrated ammonium nitrate compositions, such as, for example, UAN 18, UAN 28, UAN 30 and UAN 32.

[0012] Thus, it is desirable to have a solvent system containing alkyl thiophosphoric triamide, and in particular, (N-(n-butyl)-thiophosphoric triamide), that has a favorable toxicological and/or ecological profile and desirable characteristics in terms of low volatility, biodegradability or ready biodegradability (i.e., readily biodegradable), low toxicity or low hazard level. It is desirable to have a solvent system containing dicyandiamide, that has a favorable toxicological and/or ecological profile and desirable characteristics in terms of low volatility, biodegradability or ready biodegradability (i.e., readily biodegradable), low toxicity or low hazard level. It is also desirable to have a

solvent system containing a combination of dicyandiamide and an alkyl thiophosphoric triamide, in particular, (N-(n-butyl)-thiophosphoric triamide), that has a favorable toxicological and/or ecological profile and desirable characteristics in terms of low volatility, biodegradability or ready biodegradability (i.e., readily biodegradable), low toxicity or low hazard level.

[0013] The present invention described herein will become apparent from the following detailed description and examples, which comprises in one aspect, a composition for use in agricultural applications comprising at least one of a dicyandiamide or an alkyl thiophosphoric triamide, which is dissolved in a liquid medium comprising an organic solvent selected from polar aprotic solvents, amine solvents, heterocyclic alcohol solvents, and mixtures thereof.

[0014] In another aspect, described herein are methods of making a solid or concentrated liquid fertilizer compositions comprising treating one or more nitrogenous fertilizer compounds with an inhibitor composition that comprises at least one of a dicyandiamide or an alkyl thiophosphoric triamide. The inhibitor composition, in one embodiment, is dissolved in a liquid medium comprising an organic solvent selected from polar aprotic solvents, amine solvents, heterocyclic alcohol solvents, and mixtures thereof. The term treating, in one embodiment, includes spray applying the inhibitor composition with the one or more nitrogenous fertilizer compounds. The term treating, in one embodiment, includes spray contacting the inhibitor composition with the one or more nitrogenous fertilizer compounds.

[0015] In yet another aspect, described herein are concentrated liquid fertilizer compositions comprising, based on weight of the composition: (a) up to about 99 wt%, by weight of composition, of one or more nitrogenous fertilizer compounds, (b) at least one of a dicyandiamide or an alkyl thiophosphoric triamide, (c) an organic solvent selected from polar aprotic solvents, amine solvents, heterocyclic alcohol solvents, and mixtures thereof, and (d) water.


[0016] In yet another aspect, described herein are solid or substantially solid fertilizer

compositions comprising: (a) solid particles of one or more nitrogenous fertilizer compounds, and (b) an inhibitor composition comprising at least one of a dicyandiamide or an alkyl thiophosphoric triamide supported on at least a portion of the solid particles.

[0017] In another aspect, described herein are methods of making an aqueous end use fertilizer composition comprising incorporating: (a) one or more nitrogenous fertilizer compounds, and (b) an inhibitor composition that comprises at least one of a dicyandiamide or an alkyl thiophosphoric triamide, the inhibitor composition dissolved in a liquid medium comprising an organic solvent selected from polar aprotic solvents, amine solvents, heterocyclic alcohol solvents, and mixtures thereof, in an aqueous medium.

[0018] In another aspect, described herein are aqueous end use fertilizer composition comprising, based on weight of the composition: (a) up to about 99 wt%, based on the weight of the composition, of one or more nitrogenous fertilizer compounds, (b) at least one of a dicyandiamide or an alkyl thiophosphoric triamide, (c) an organic solvent selected from polar aprotic solvents, amine solvents, heterocyclic alcohol solvents, and mixtures thereof, and (d), optionally, water.


[0019] In another aspect, described herein are methods for fertilizing target plants, comprising applying an aqueous end use fertilizer composition that comprises: (a) one or more nitrogenous fertilizer compounds, (b) at least one of a dicyandiamide or an alkyl thiophosphoric triamide, (c) an organic solvent selected from dimethyl sulfoxide, dimethyl formamide, the dimethyl ester of succinic acid, dimethyl ester of ethyl succinic acid, the dimethyl ester of glutaric acid, the dimethyl ester of methyl glutaric acid, and the dimethyl ester of adipic acid, diethylene triamine, or monoethanolamine, methyl-5-(dimethylamino)-2-methyl-oxopentanoate, dimethylaminoethanol, triethanol amine, a heterocyclic alcohol according to structure (III):



(III),


[0020] and mixtures thereof, and, optionally, (d) water, to the target plants or to an environment for the target plants. It is understood that the term heterocyclic alcohol includes dioxolane compounds.

[0021] In one embodiment, the alkyl thiophosphoric triamide is N-(n-butyl)-thiophosphoric triamide. In another embodiment, the liquid medium further comprises a solvent selected from the group consisting of: (a) at least one dioxolane compound of formula (III):



(III),

[0022] wherein R_6 and R_7 individually comprises a hydrogen, an alkyl group, an alkenyl group, or a phenyl group, wherein n is an integer of from 1 to 10; b) at least one dibasic ester; c) at least one compound of formula (IIa):



(IIa),

[0023] wherein R_3 comprises a C_1 - C_{36} alkyl group; wherein R_4 and R_5 individually comprise a C_1 - C_{36} alkyl group, wherein R_4 and R_5 can optionally together form a ring; and wherein A is a linear or a branched divalent C_2 - C_6 alkyl group; d) at least one alkyldimethylamide; e) at least one alkyl lactate; f) ethyl levulinate; g) at least one

alkoxyalcohol, ether alcohol, amine alcohol, amino alcohol or alcohol; h) at least one glycerine or glycerine derivative; i) at least one alkylene carbonate; j) dimethylsulfoxide; and k) any combination thereof. In one embodiment, the organic solvent is dimethylsulfoxide.

[0024] In another aspect, the present invention is directed to a nitrification inhibitor composition comprising dicyandiamide dissolved in a liquid medium that comprises an organic solvent selected from polar aprotic solvents, dibasic esters, amines, amino alcohols, heterocyclic alcohols, and mixtures thereof.

[0025] In yet another aspect, the present invention is directed to a method of making a solid or concentrated liquid fertilizer composition comprising treating one or more nitrogenous fertilizer compounds with a nitrification inhibitor composition that comprises dicyandiamide dissolved in a liquid medium that comprises an organic solvent selected from polar aprotic solvents, amine solvents, heterocyclic alcohol solvents, and mixtures thereof.

[0026] In a further aspect, the present invention is directed to a concentrated liquid fertilizer composition comprising, based on 100 parts by weight of the composition:

- (a) up to about 99 parts by weight of one or more nitrogenous fertilizer compounds,
- (b) dicyandiamide,
- (c) an organic solvent selected from polar aprotic solvents, amine solvents, heterocyclic alcohol solvents, and mixtures thereof, and
- (d) water.

[0027] In another aspect, the present invention is directed to a concentrated solid fertilizer composition comprising:

- (a) solid particles of one or more nitrogenous fertilizer compounds, and
- (b) dicyandiamide supported on at least a portion of the solid particles.

[0028] In yet another aspect, the present invention is directed to a method of making an aqueous end use fertilizer composition comprising incorporating:


- (a) one or more nitrogenous fertilizer compounds, and
- (b) a nitrification inhibitor composition that comprises dicyandiamide dissolved in a liquid medium that comprises an organic solvent selected from polar aprotic solvents, amine solvents, heterocyclic alcohol solvents, and mixtures thereof, in an aqueous medium.

[0029] In another aspect, the present invention is directed to an aqueous end use fertilizer composition comprising, based on 100 parts by weight of the composition:

- (a) up to about 99 parts by weight of one or more nitrogenous fertilizer compounds,
- (b) dicyandiamide,
- (c) an organic solvent selected from polar aprotic solvents, amine solvents, heterocyclic alcohol solvents, and mixtures thereof, and
- (d) water.

[0030] In a further aspect, the present invention is directed to a method for fertilizing target plants, comprising applying an aqueous end use fertilizer composition that comprises:

- (a) one or more nitrogenous fertilizer compounds
- (b) dicyandiamide,
- (c) an organic solvent selected from dimethyl sulfoxide, dimethyl formamide, the dimethyl ester of succinic acid, dimethyl ester of ethyl succinic acid, the dimethyl ester of glutaric acid, the dimethyl ester of methyl glutaric acid, and the dimethyl ester of adipic acid, diethylene triamine, or monoethanolamine, methyl-5-(dimethylamino)-2-methyl-oxopentanoate, dimethylaminoethanol, triethanol amine, a heterocyclic alcohol according to structure (III):



(III),

and mixtures thereof, and

(d) water,

to the target plants or to an environment for the target plants.

DETAILED DESCRIPTION


[0031] As used herein, the term "alkyl" means a saturated straight chain, branched chain, or cyclic hydrocarbon radical, including but not limited to, methyl, ethyl, n-propyl, iso-propyl, n-butyl, sec-butyl, t-butyl, pentyl, n-hexyl, and cyclohexyl.

[0032] As used herein, the term "aryl" means a monovalent unsaturated hydrocarbon radical containing one or more six-membered carbon rings in which the unsaturation may be represented by three conjugated double bonds, which may be substituted one or more of carbons of the ring with hydroxy, alkyl, alkenyl, halo, haloalkyl, or amino, including but not limited to, phenoxy, phenyl, methylphenyl, dimethylphenyl, trimethylphenyl, chlorophenyl, trichloromethylphenyl, aminophenyl, and tristyrylphenyl.

[0033] As used herein, the term "alkylene" means a divalent saturated straight or branched chain hydrocarbon radical, such as for example, methylene, dimethylene, trimethylene.

[0034] As used herein, the terminology "(C_r-C_s)" in reference to an organic group, wherein r and s are each integers, indicates that the group may contain from r carbon atoms to s carbon atoms per group.

[0035] Dicyandiamide is a known compound according to formula (I):



[0036] Dicyandiamide, also known as “2-cyanoguanidine”, is typically made by treating cyanamide with base and is commercially available.

[0037] In one embodiment, the compositions according to the present invention comprise a urease inhibitor, such as N-(n-butyl)-thiophosphoric triamide (“NBPT”) or ammonium thiosulfate, a nitrification inhibitor, or a combination of both a urease inhibitor and a nitrification inhibitor.

[0038] In one embodiment, alkyl thiophosphoric triamide is N-(n-butyl)-thiophosphoric triamide. The at least one of alkyl thiophosphoric triamide or dicyandiamide or combination thereof can be present in the composition at a lower range of 5% by weight of the composition. In another embodiment, at least one of alkyl thiophosphoric triamide and/or dicyandiamide can be present in the composition at a lower range of 6%, or 8%, or 10% or 12 % or 14%, by weight of the composition. In another embodiment, the at least one of alkyl thiophosphoric triamide or dicyandiamide can be present in the composition in an amount between about 7% by weight of the composition to about 13% by weight of the composition. . In another embodiment, the at least one of alkyl thiophosphoric triamide or dicyandiamide can be present in the composition in an amount between about 8% by weight of the composition to about 12% by weight of the composition.

[0039] The at least one of alkyl thiophosphoric triamide or dicyandiamide can be present in the composition in an amount between about 0.5 % by weight of the composition and about 50 % by weight of the composition or, in another embodiment, can be present in the composition in an amount between about 1 % by weight of the composition and about 40 % by weight of the composition, and, in another embodiment, can be present in the composition in an amount between about 0.5 % by weight of the composition and

about 20 % by weight of the composition. In one particular embodiment, the at least one of alkyl thiophosphoric triamide or dicyandiamide is present in the composition in an amount between about 1 % by weight of the composition and about 30 % by weight of the composition. The at least one of alkyl thiophosphoric triamide or dicyandiamide means that alkyl thiophosphoric triamide can be solely present, dicyandiamide can be solely present, or a combination of alkyl thiophosphoric triamide and dicyandiamide is present.

[0040] Compounds suitable as the organic solvent component of the composition and methods of the present invention are those polar aprotic solvents, amine solvents, heterocyclic alcohol solvents, and mixtures thereof that are liquid at temperatures of from 20 to 50°C, and that do not react with dicyandiamide under anticipated manufacturing, storage, and use conditions. In one embodiment, the solvent comprises a polar aprotic solvent, amine solvent, heterocyclic alcohol solvent, or mixtures thereof in which dicyandiamide has a solubility at 25 °C of greater than or equal to 50 grams per liter (g/l), more typically greater than or equal to 200 g/l.

[0041] Suitable polar aprotic organic solvents include, for example, dichloromethane, dimethyl acetamide, dimethyl formamide, dimethyl sulfoxide, ethyl acetate, hexamethylphosphoramide, dimethyl sulfone, sulfolane, 1,3-dimethyl-2-imidazolidinone, 1,3-dimethyl-3,4,5,6-tetrahydro-2(1H)-pyrimidone, methyl acetate, ethyl lactate, methylpyrrolidone, tetrahydrofuran, propylene carbonate, and dibasic ester solvents.

[0042] Suitable dibasic ester solvents include, for example, dialkyl esters of dicarboxylic acids, more typically, the di(C₁-C₁₂)alkyl esters of saturated linear or branched (C₂-C₈)aliphatic carboxylic acids or a mixture thereof. In one embodiment, the dibasic ester component comprises one or more compounds according to structure (II):



wherein:

A is a divalent linear or branched (C₂-C₈)aliphatic group, and

R¹, R², and R³ are each independently (C₁-C₁₂)alkyl, (C₁-C₁₂)aryl, (C₁-C₁₂)alkaryl or (C₁-C₁₂)arylalkyl, and R² and R³ may each optionally be substituted with one or more hydroxyl groups.

[0043] In one embodiment, the dibasic ester solvent component of the compositions and methods of the present invention comprises one or more dimethyl esters of saturated linear or branched (C₄-C₆)aliphatic carboxylic acids, such the dimethyl ester of succinic acid, dimethyl ester of ethyl succinic acid, the dimethyl ester of glutaric acid, the dimethyl ester of methyl glutaric acid, and the dimethyl ester of adipic acid, and mixtures thereof. In one embodiment, the dibasic ester component comprises the dimethyl ester of succinic acid, the dimethyl ester of glutaric acid, and optionally, the dimethyl ester of adipic acid, In another embodiment, the dibasic ester component comprises the dimethyl ester of ethyl succinic acid, the dimethyl ester of methyl glutaric acid, and optionally, the dimethyl ester of adipic acid.

[0044] In one embodiment, the dibasic ester solvent component of the compositions and methods of the present invention comprises one or more dialkyl esters of saturated linear or branched (C₄-C₆)aliphatic carboxylic acids, such the dialkyl ester of succinic acid, dialkyl ester of ethylsuccinic acid, the dialkyl ester of glutaric acid, the dialkyl ester of methylglutaric acid, and the dialkyl ester of adipic acid, and mixtures thereof. In one embodiment, the dibasic ester component comprises the dialkyl ester of succinic acid, the dialkyl ester of glutaric acid, and optionally, the dimethyl ester of adipic acid, In another embodiment, the dibasic ester component comprises the dialkyl dimethyl ester of ethylsuccinic acid, the dialkyl ester of methylglutaric acid, and optionally, the dialkyl ester of adipic acid. Each alkyl group in the dialkyl group, one embodiment, individually comprise a C₁-C₈ alkyl. In another embodiment, each alkyl group in the dialkyl group, one embodiment, individually comprise a C₁-C₄ alkyl. In another embodiment, each alkyl group in the dialkyl group, one embodiment, individually comprise a C₁-C₆ alkyl.

[0045] Suitable amine solvents include primary amines, including monoalkylamines,

such as propylamine, secondary amines, including dialkyl amines and diaryl amines, such as dimethylamine and diphenylamine, and tertiary amines, such as diethylene triamine and methyl-5-(dimethylamino)-2-methyl-oxopentanoate.


[0046] In one embodiment, the amine solvent component of the compositions and methods of the present invention is selected from aliphatic or aromatic primary, secondary, or tertiary amines that do not comprise any functional group other than one or more amino groups.

[0047] In one embodiment, the amine solvent component of the compositions and methods of the present invention is selected from aliphatic or aromatic primary, secondary, or tertiary amines may optionally further comprise one or more additional functional groups, such as hydroxyalkyl groups, hydroxyl groups, carbonyl groups, or alkyl ester groups, other than one or more amino groups.

[0048] In one embodiment, the organic solvent component of the compositions and methods of the present invention comprises an amino alcohol. Compounds suitable as the amino alcohol solvent component of the compositions and methods of the present invention are those compounds that comprise at least one primary, secondary, or tertiary amino moiety per molecule and at least one hydroxyalkyl moiety per molecule, more typically In one embodiment, the amino alcohol is a linear, branched, or cyclic, saturated or unsaturated hydrocarbon that is substituted on at least one carbon atom with an amino group and on at least one other carbon atom with hydroxyalkyl or hydroxyl group, such as monoethanolamine, ethylaminoethanol, dimethylaminoethanol, isopropylaminoethanol, diethanolamine, triethanolamine, methylaminoethanol, aminopropanol, methylaminopropanol, dimethylaminopropanol, aminobutanol, dimethylaminobutanol, aminobutanediol, trihydroxymethylaminoethane, diethylaminopropanediol, 1-amino-cyclopentane methanol, and aminobenzyl alcohol, or a heterocyclic ring that comprises at least one nitrogen atom as a ring member and/or is substituted on at least one carbon atom with an amino group and that is substituted on at least one other carbon atom with a hydroxyalkyl or hydroxyl group, such as


methylaminomethyl-1,3-dioxolane.

[0049] Suitable heterocyclic alcohol solvents include, for example, 5- or 6-membered heterocyclic rings that include 1 or 2 oxygen atoms as ring member, that are substituted on at least one carbon atom of the ring with a (C₁-C₆)hydroxyalkyl group, and that may optionally be substituted on one or more carbon atoms of the ring with one or more (C₁-C₄)alkyl groups. It is understood that the term heterocyclic alcohol includes dioxolane compounds. In one embodiment, the heterocyclic alcohol component of the present invention comprises a one or more compounds selected from heterocyclic alcohols according to structures (III), (IV), (V), (VI), and (VII):




(III)


wherein $n = 1$ or 2 ,




(IV)



(V)



(VI)



(VII)

[0050] In one embodiment, the organic solvent component of the composition and methods of the present invention comprises one or more polar aprotic solvents, one or more dibasic ester compounds according to structure (II), one or more amino alcohols, one or more tertiary amines, one or more heterocyclic alcohols according to structure (III), or a mixture thereof.


[0051] In one embodiment, the organic solvent component of the composition and methods of the present invention comprises dimethyl sulfoxide, dimethyl formamide, the dimethyl ester of succinic acid, dimethyl ester of ethyl succinic acid, the dimethyl ester of glutaric acid, the dimethyl ester of methyl glutaric acid, and the dimethyl ester of adipic acid, diethylene triamine, or monoethanolamine, methyl-5-(dimethylamino)-2-methyl-oxopentanoate, dimethylaminoethanol, triethanol amine, a heterocyclic alcohol according to structure (III), or a mixture thereof.

[0052] In one embodiment, the organic solvent component of the composition and methods of the present invention comprises dimethyl sulfoxide, dimethyl formamide, diethylene triamine, monoethanolamine, or a mixture thereof.

[0053] In one embodiment, the organic solvent component of the composition and methods of the present invention comprises dimethyl sulfoxide.

[0054]

[0055] In one embodiment, a compound utilized as the solvent or as a component in the solvent blend is a compound of general formula (II):



[0056] According to one embodiment, the expression "compound" denotes any compound corresponding to the general formula (II). In other embodiments, the term "compound" also refers to mixtures of several molecules corresponding to general formula (II). It may therefore be a molecule of formula (II) or a mixture of several molecules of formula (II), wherein both fall under the definition of the term "compound" when referring to formula (II).

[0057] The R₃, R₄ and R₅ groups can be, in some embodiments, identical or, in other embodiment, different. In one embodiment, may be groups chosen from C₁-C₂₀ alkyl, aryl, alkaryl or arylalkyl groups or the phenyl group. In another embodiment, may be

groups chosen from C₁-C₁₂ alkyl, aryl, alkaryl or arylalkyl groups or the phenyl group. Mention is made especially of Rhodiasolv® PolarClean (Manufactured by Rhodia Inc. of Cranbury, NJ). The R₄ and R₅ groups may optionally be substituted. In one particular embodiment, the groups are substituted with hydroxyl groups.

[0058] In one embodiment, R₃ group is chosen from methyl, ethyl, propyl, isopropyl, n-butyl, isobutyl, n-pentyl, isopentyl, isoamyl, n-hexyl, cyclohexyl, 2-ethylbutyl, n-octyl, isooctyl, 2-ethylhexyl, tridecyl groups.

[0059] R₄ and R₅ groups, which are identical or different, in one embodiment, may especially be chosen from methyl, ethyl, propyl (n-propyl), isopropyl, n-butyl, isobutyl, n-pentyl, amyl, isoamyl, hexyl, cyclohexyl or hydroxyethyl groups. The R₄ and R₅ groups may also be such that they form, together with the nitrogen atom, a morpholine, piperazine or piperidine group. According to some embodiments, R₄ and R₅ are each methyl, or R₄ and R₅ are each ethyl, or R₄ and R₅ are each hydroxyethyl.

[0060] According to one embodiment, if A comprises a linear group of formula --CH₂--CH₂-- and/or of formula --CH₂--CH₂--CH₂--CH₂-- and/or of formula --(CH₂)₈-- then it is a mixture of A groups. According to one particular embodiment, if A is linear, then it is a mixture of A groups, for example a mixture of two or three --CH₂--CH₂-- (ethylene); --CH₂--CH₂--CH₂-- (n-propylene); and --CH₂--CH₂--CH₂--CH₂-- (n-butylene) groups (or isomers thereof).

[0061] According to a first particular embodiment of the invention, the A group is a divalent linear alkyl group chosen from the groups of the following formulae: --CH₂--CH₂-- (ethylene); --CH₂--CH₂--CH₂-- (n-propylene); --CH₂--CH₂--CH₂--CH₂-- (n-butylene), and mixtures thereof.

[0062] In one embodiment, the inhibitor composition of the present invention comprises, based on 100 parts by weight ("pbw") of the composition:

[0063] from about 4 to about 45 pbw, more typically from about 10 to about 42 pbw, and even more typically from about 20 to about 40 pbw dicyandiamide, and

[0064] from about 55 to about 96 pbw, more typically from about 58 to about 90 pbw, and even more typically from about 60 to about 80 pbw of the organic solvent.

[0065] In one embodiment, the inhibitor composition of the present invention comprises one or more urease inhibitors, such as, for example, NBPT or ammonium thiosulfate.

[0066] The nitrogenous fertilizer compound is treated with the inhibitor composition by contacting the nitrogenous fertilizer composition with the inhibitor composition described herein (e.g., nitrification inhibitor or urease inhibitor or a combination of both). The nitrogenous fertilizer composition may be in solid or liquid form.

[0067] Suitable nitrogenous fertilizers are those containing a nitrogenous compound such as urea, nitrate salts, ammonium salt, or a mixture thereof, such as ammonium nitrate, ammonium sulfate, ammonium thiosulfate, ammonium polysulfide, ammonium phosphates, ammonium chloride, ammonium bicarbonate, anhydrous ammonia, calcium nitrate, nitrate soda, calcium cyanamide. In one embodiment, the nitrogenous fertilizer comprises ammonium nitrate. Suitable ammonium nitrate-containing fertilizers include, for example, UAN 18, UAN 28, and UAN 30.

[0068] In one embodiment, the nitrogenous fertilizer composition is in solid particulate form, and the contacting of the nitrogenous fertilizer composition with the inhibitor composition is conducted by, for example, spraying the composition of the present invention on the particles of solid fertilizer composition.

[0069] In one embodiment, the concentrated fertilizer composition of the present invention is a solid nitrification-inhibited fertilizer composition that comprises, based on 100 pbw of the composition:

from about 60 pbw to about 99.999, more typically from about 70 pbw to about 99.999, and even more typically from about 80 pbw to about 99.999 solid particles of one or more nitrogenous fertilizer compounds, and

from about 0.001 to about 40 pbw, more typically from about 0.001 to about 30

pbw, and even more typically from about 0.001 to about 20 pbw, dicyandiamide.

[0070] In one embodiment, the solid nitrification-inhibited fertilizer composition of the present invention further comprises one or more urease inhibitors, more typically NBPT.

[0071] In one embodiment, the end use fertilizer composition of the present invention is made by combining the inhibitor composition of the present invention with a solid nitrogenous fertilizer to form a solid nitrification-inhibited fertilizer composition and subsequently dissolving the solid nitrification-inhibited fertilizer composition in an aqueous medium, typically water, in a ratio of up to about 500 pbw, more typically from 100 to 500 pbw and even more typically from about 100 to about 300 pbw, of the aqueous medium per 1 pbw of the solid nitrification-inhibited fertilizer composition.

[0072] In one embodiment, the fertilizer compound is in liquid form and the contacting of the fertilizer composition with the inhibitor composition is conducted by mixing the inhibitor composition with the liquid fertilizer composition.

[0073] In one embodiment, the concentrated fertilizer composition of the present invention is a concentrated liquid nitrification-inhibited fertilizer composition that comprises, based on 100 pbw of the composition:

from about 20 to about 99.989 pbw, more typically from about 30 to about 99.985 pbw, and even more typically from about 40 to about 99.98 pbw of one or more nitrogenous fertilizer compounds,

from about 0.001 to 40 pbw, more typically from about 0.005 to 30 pbw, and even more typically from about 0.01 to 20 pbw dicyandiamide, and

from about 0.01 to 60 pbw, more typically from about 0.01 to about 40 pbw, and even more typically from about 0.01 to about 30 pbw of the organic solvent.

[0074] In one embodiment, the concentrated liquid nitrification-inhibited fertilizer composition of the present invention further comprises one or more urease inhibitors, more typically NBPT.

[0075] In one embodiment, the end use fertilizer composition of the present invention is made by combining the inhibitor composition of the present invention with a concentrated nitrogenous fertilizer to form a concentrated liquid nitrification-inhibited fertilizer composition and subsequently diluting the concentrated liquid nitrification-inhibited fertilizer composition with an aqueous medium, typically water in a ratio of up to about 500 pbw, more typically from about 10 to about 500 pbw and even more typically from about 100 to about 300 pbw, of the aqueous medium per 1 pbw concentrated liquid nitrogenous fertilizer composition.

[0076] In one embodiment, the end use fertilizer composition of the present invention is made by combining the inhibitor composition of the present invention, a solid or concentrated liquid nitrogenous fertilizer, and an aqueous medium.

[0077] In one embodiment, the end use fertilizer composition of the present invention is an aqueous liquid composition that comprises water, one or more nitrogenous fertilizer compounds, and dicyandiamide, typically in an amount of from 2×10^{-6} pbw to about 4 pbw dicyandiamide per 100 pbw of the end use fertilizer composition.

[0078] In one embodiment, the end use fertilizer composition of the present invention comprises water and based on 100 parts by weight of the composition:

from about 0.04 to about 10 pbw, more typically from about 0.06 to about 10 pbw, and even more typically from about 0.08 pbw to about 10 pbw of one or more nitrogenous fertilizer compounds,

from about 2×10^{-6} to about 4 pbw, more typically from about 1×10^{-5} to about 3 pbw, and even more typically from about 2×10^{-4} to about 2 pbw dicyandiamide, and

from about 2×10^{-4} to about 6 pbw, more typically from about 2×10^{-4} to about 4 pbw, and even more typically from about 2×10^{-4} to about 3 pbw of the organic solvent.

[0079] In one embodiment, the end use fertilizer composition of the present invention comprises one or more urease inhibitors, more typically NBPT, alone or in combination

with the nitrification inhibitor.

[0080] In one embodiment, the end use fertilizer composition of the present invention comprises from about 0.001 to about 5 pbw, more typically from about 0.01 to about 2 pbw dicyandiamide per 100 pbw of the one or more nitrogenous fertilizer compounds.

[0081] In one embodiment, the end use fertilizer composition is applied to target plants or to an environment for the target plants, i.e., to ground on or within which the target plants are growing or to be grown, at a rate of from about 0.01 pounds to about 5 pounds of the fertilizer composition, more typically from about 0.05 pounds to about 2 pounds of the fertilizer composition, per 100 square feet of ground.

[0082] In one embodiment, the end use fertilizer composition is applied to target plants or to an environment for the target plants at a rate effective to provide a dosage of nitrogenous fertilizer compound of from about 0.01 pounds to about 5 pounds of fertilizer compound, more typically from about 0.05 pounds to 2 pounds of fertilizer compound, per 100 square feet of ground.

[0083] In one embodiment, the end use fertilizer composition is applied to target plants or to an environment for the target plants at a rate effective to provide a dosage of dicyandiamide of from about 0.01 pounds to 5 pounds of dicyandiamide, more typically from about 0.05 pounds to 2 pounds of dicyandiamide, per 1000 square feet of ground.

[0084] The composition of the present invention provides improved ease of handling of dicyandiamide, improved solubility characteristics, low toxicity of the organic solvents; good storage characteristics, and excellent miscibility with aqueous compositions, such as aqueous nitrogenous fertilizer formulations.

Example 1

[0085] The composition of Example 1 was made by dissolving 30 pbw dicyandiamide (“DCD”) in 70 pbw dimethyl sulfoxide (“DMSO”).

[0086] Samples of the 30 DCD/70 DMSO solution were analyzed by NMR spectroscopy immediately after formation and at 3 weeks after formation and were in each case found to contain no evidence of any reaction having occurred between the DCD and the DMSO.

Samples 30 DCD/70 DMSO solution were stored at -16°C, 4°C, RT, 45 and 54°C and subjected to freeze-thaw cycling (cycling between 25 °C and -12 °C, with a 12 hour dwell at each temperature extreme and an 8 hour ramp between the temperature extremes) and the stability of the solution was evaluated after 1 week, after 2 weeks, and after 1 month of storage under each of the storage conditions and was found to be stable under each of the storage conditions.

[0087] Samples of 5 wt%, 10 wt%, and 20 wt% of the 30 DCD/70 DMSO solution in UAN 28, 32, and 34 were made and found to be stable at room temperature for over 2 months.

Examples 2, 3, and 4

[0088] The composition of Example 2 was made by dissolving 21 pbw DCD in 79 pbw monoethanol amine.

[0089] The composition of Example 3 was made by dissolving 12 pbw DCD in a mixture of 62 pbw of a mixture of methyl-5-(dimethylamino)-2-methyl-oxopentanoate,, 9 pbw propylene carbonate, and 17 pbw water

[0090] The composition of Example 4 was made by dissolving 30 pbw DCD and 15 pbw N-(n-butyl)-thiophosphoric triamide in 55 pbw dimethyl sulfoxide.

[0091] Each of the compositions of Examples 2, 3, and 4 was analyzed by NMR spectroscopy immediately after formation as were in each case found to be free of evidence of any reaction having occurred between the DCD and the respective organic solvent system. Each of the compositions of Examples 2, 3, and 4 was found to be stable after storage at room temperature for at least one month.


Example 5

[0092] NBPT (50 wt%, by weight of mixture) in DMSO (50 wt%) samples were put through stability testing at -16°C, 4°C, 25°C, 45°C, 54°C and subjected to freeze-thaw cycling (cycling between 25 °C and -12 °C, with a 12 hour dwell at each temperature extreme and an 8 hour ramp between the temperature extremes). The stability of the sample was evaluated after two weeks. All the samples were observed to remained in 1 phase, although the samples became darker at higher temperatures.

CLAIMS

What is claimed is:


1. A composition for use in agricultural applications comprising at least one of a dicyandiamide or an alkyl thiophosphoric triamide, dissolved in a liquid medium comprising an organic solvent comprising a polar aprotic solvent, an amine solvent, a heterocyclic alcohol solvent or a mixture thereof.
2. The composition of claim 1 wherein the alkyl thiophosphoric triamide is N-(n-butyl)-thiophosphoric triamide.
3. The composition of claim 1 wherein the organic solvent comprises:
 - (a) at least one dioxolane compound of formula (Ia):



wherein R_6 and R_7 individually comprises a hydrogen, an alkyl group, an alkenyl group, or a phenyl group, wherein n is an integer of from 1 to 10

b) at least one dibasic ester;

c) at least one compound of formula (IIa):



(IIa),

wherein R_3 comprises a C_1 - C_{36} alkyl group; wherein R_4 and R_5 individually comprise a C_1 - C_{36} alkyl group, wherein R_4 and R_5 can optionally together form a ring; and wherein A is a linear or a branched divalent C_2 - C_6 alkyl group;

d) at least one alkyldimethylamide;

e) at least one alkyl lactate;

f) ethyl levulinate;

g) at least one alkoxyalcohol, ether alcohol, amine alcohol, amino alcohol or alcohol;

h) at least one glycerine or glycerine derivative;

i) at least one alkylene carbonate;

j) dimethylsulfoxide; or

k) any combination thereof.

4. The composition of claim 1 wherein the organic solvent is dimethyl sulfoxide.

5. The composition of claim 1 wherein the polar aprotic solvent is selected from the group consisting of dichloromethane, dimethyl acetamide, dimethyl formamide, dimethyl sulfoxide, ethyl acetate, hexamethylphosphoramide, dimethyl sulfone, sulfolane, 1,3-dimethyl-2-imidazolidinone, 1,3-dimethyl-3,4,5,6-tetrahydro-2(1H)-pyrimidone, methyl


acetate, ethyl lactate, methylpyrrolidone, tetrahydrofuran, propylene carbonate, and dibasic ester solvents.

6. A method of making a solid or concentrated liquid fertilizer composition comprising treating one or more nitrogenous fertilizer compounds with an inhibitor composition that comprises at least one of a dicyandiamide or an alkyl thiophosphoric triamide, the inhibitor composition dissolved in a liquid medium comprising an organic solvent selected from polar aprotic solvents, amine solvents, heterocyclic alcohol solvents, and mixtures thereof.

7. The method of claim 6 wherein the alkyl thiophosphoric triamide is N-(n-butyl)-thiophosphoric triamide.

8. The method of claim 6 wherein the solvent comprises:

(a) at least one dioxolane compound of formula (Ia):



wherein R_6 and R_7 individually comprises a hydrogen, an alkyl group, an alkenyl group, or a phenyl group, wherein n is an integer of from 1 to 10

b) at least one dibasic ester;

c) at least one compound of formula (IIa):



(IIa),

wherein R_3 comprises a C_1 - C_{36} alkyl group; wherein R_4 and R_5 individually comprise a C_1 - C_{36} alkyl group, wherein R_4 and R_5 can optionally together form a ring; and wherein A is a linear or a branched divalent C_2 - C_6 alkyl group;

d) at least one alkyldimethylamide;

e) at least one alkyl lactate;

f) ethyl levulinate;

g) at least one alkoxyalcohol, ether alcohol, amine alcohol, amino alcohol or alcohol;

h) at least one glycerine or glycerine derivative;

i) at least one alkylene carbonate;

j) dimethylsulfoxide; or

k) any combination thereof.

9. The method of claim 6 wherein the organic solvent is dimethyl sulfoxide.

10. The method of claim 6 wherein the polar aprotic solvent is selected from the group consisting of dichloromethane, dimethyl acetamide, dimethyl formamide, dimethyl sulfoxide, ethyl acetate, hexamethylphosphoramide, dimethyl sulfone, sulfolane, 1,3-dimethyl-2-imidazolidinone, 1,3-dimethyl-3,4,5,6-tetrahydro-2(1H)-pyrimidone, methyl


acetate, ethyl lactate, methylpyrrolidone, tetrahydrofuran, propylene carbonate, and dibasic ester solvents.

11. A concentrated liquid fertilizer composition comprising, based on 100 parts by weight of the composition:

- (a) up to about 99 parts by weight of one or more nitrogenous fertilizer compounds,
- (b) at least one of a dicyandiamide or an alkyl thiophosphoric triamide,
- (c) an organic solvent selected from polar aprotic solvents, amine solvents, heterocyclic alcohol solvents, and mixtures thereof, and
- (d) water.

12. The liquid fertilizer composition of claim 11 wherein the organic solvent comprises:

(a) at least one dioxolane compound of formula (Ia):



wherein R_6 and R_7 individually comprises a hydrogen, an alkyl group, an alkenyl group, or a phenyl group, wherein n is an integer of from 1 to 10

b) at least one dibasic ester;

c) at least one compound of formula (IIa):



(IIa),

wherein R_3 comprises a C_1 - C_{36} alkyl group; wherein R_4 and R_5 individually comprise a C_1 - C_{36} alkyl group, wherein R_4 and R_5 can optionally together form a ring; and wherein A is a linear or a branched divalent C_2 - C_6 alkyl group;

d) at least one alkyldimethylamide;

e) at least one alkyl lactate;

f) ethyl levulinate;

g) at least one alkoxyalcohol, ether alcohol, amine alcohol, amino alcohol or alcohol;

h) at least one glycerine or glycerine derivative;

i) at least one alkylene carbonate;

j) dimethylsulfoxide; or

k) any combination thereof.

12. A solid fertilizer composition comprising:

(a) solid particles of one or more nitrogenous fertilizer compounds, and

(b) an inhibitor composition comprising at least one of a dicyandiamide or an alkyl thiophosphoric triamide supported on at least a portion of the solid particles.


13. A method of making an aqueous end use fertilizer composition comprising incorporating:

(a) one or more nitrogenous fertilizer compounds, and

- (b) an inhibitor composition that comprises at least one of a dicyandiamide or an alkyl thiophosphoric triamide, the inhibitor composition dissolved in a liquid medium comprising an organic solvent selected from polar aprotic solvents, amine solvents, heterocyclic alcohol solvents, and mixtures thereof, in an aqueous medium.

14. The method of claim 13 wherein the organic solvent comprises:

- (a) at least one dioxolane compound of formula (Ia):



wherein R_6 and R_7 individually comprises a hydrogen, an alkyl group, an alkenyl group, or a phenyl group, wherein n is an integer of from 1 to 10

- b) at least one dibasic ester;

- c) at least one compound of formula (IIa):



(IIa),

wherein R_3 comprises a C_1 - C_{36} alkyl group; wherein R_4 and R_5 individually comprise a C_1 - C_{36} alkyl group, wherein R_4 and R_5 can optionally together form a ring; and wherein A is a linear or a branched divalent C_2 - C_6 alkyl group;

- d) at least one alkyldimethylamide;
- e) at least one alkyl lactate;
- f) ethyl levulinate;
- g) at least one alkoxyalcohol, ether alcohol, amine alcohol, amino alcohol or alcohol;
- h) at least one glycerine or glycerine derivative;
- i) at least one alkylene carbonate;
- j) dimethylsulfoxide; or
- k) any combination thereof.


15. An aqueous end use fertilizer composition comprising, based on 100 parts by weight of the composition:

- (a) up to about 99 parts by weight of one or more nitrogenous fertilizer compounds,
- (b) at least one of a dicyandiamide or an alkyl thiophosphoric triamide,
- (c) an organic solvent selected from polar aprotic solvents, amine solvents, heterocyclic alcohol solvents, and mixtures thereof, and
- (d) water.

16. A method for fertilizing target plants, comprising applying an aqueous end use fertilizer composition that comprises:

- (a) one or more nitrogenous fertilizer compounds
- (b) at least one of a dicyandiamide or an alkyl thiophosphoric triamide,
- (c) an organic solvent selected from dimethyl sulfoxide, dimethyl formamide, the

dimethyl ester of succinic acid, dimethyl ester of ethyl succinic acid, the dimethyl ester of glutaric acid, the dimethyl ester of methyl glutaric acid, and the dimethyl ester of adipic acid, diethylene triamine, or monoethanolamine, methyl-5-(dimethylamino)-2-methyl-oxopentanoate, dimethylaminoethanol, triethanol amine, a heterocyclic alcohol according to structure (III):



and mixtures thereof, and

(d) water,

to the target plants or to an environment for the target plants.

A. CLASSIFICATION OF SUBJECT MATTER**C05F 11/00(2006.01)i, C05F 15/00(2006.01)i**

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C05F 11/00; C07F 9/22; C05C 9/00; C05F 15/00

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Korean utility models and applications for utility models

Japanese utility models and applications for utility models

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

eKOMPASS(KIPO internal) & Keywords: thiophosphoric triamide, NBPT, fertilizer, agriculture, solvent

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X A	WO 2010-096266 A1 (WEYERHAEUSER NR COMPANY) 26 August 2010 See abstract and claims 1-26.	1-3,6-8,11-16 4,5,9,10
X A	WO 2010-072184 A2 (AGRA GROUP, A. S.) 1 July 2010 See abstract: page 9, lines 4-15; and claims 1, 12.	1-3,6-8,11-16 4,5,9,10
A	EP 2032589 B1 (AGRA GROUP, A. S.) 28 September 2011 See abstract and claims 1, 12-15.	1-16
A	US 5698003 A (OMILINSKY, B. A. et al.) 16 December 1997 See abstract and claims 1-4, 8, 9.	1-16

 Further documents are listed in the continuation of Box C. See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

26 May 2014 (26.05.2014)

Date of mailing of the international search report

26 May 2014 (26.05.2014)

Name and mailing address of the ISA/KR

International Application Division
Korean Intellectual Property Office
189 Cheongsu-ro, Seo-gu, Daejeon Metropolitan City, 302-701,
Republic of Korea

Facsimile No. +82-42-472-7140

Authorized officer

HEO, Joo Hyung

Telephone No. +82-42-481-8150



INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/US2013/076858

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2010-096266 A1	26/08/2010	AR 075577 A1	20/04/2011
		AU 2011-216282 A1	08/09/2011
		AU 216282 B2	23/08/2012
		EP 2398752 A1	28/12/2011
		MX 2011008693 A	18/11/2011
		PE 07152012 A1	12/07/2012
		US 2010-0206031 A1	19/08/2010
		US 2011-0259068 A1	27/10/2011
		US 8163058 B2	24/04/2012
		UY 32453 A	30/09/2010
WO 2010-072184 A2	01/07/2010	AU 2010-329568 A1	01/07/2010
		CA 2745584 A1	01/07/2010
		CN 102264748 A	30/11/2011
		CZ 20080842 A3	30/06/2010
		EA 201100886 A1	28/02/2012
		EP 2376506 A2	19/10/2011
		NZ 593120 A	31/08/2012
		US 2011-0233474 A1	29/09/2011
		WO 2010-072184 A3	19/08/2010
		EP 2032589 B1	28/09/2011
AU 2008-264228 A1	03/01/2008		
AU 264228 B2	01/03/2012		
BR PI0713788 A2	30/10/2012		
CN 101479282 A	08/07/2009		
CN 101479282 B	18/01/2012		
CZ 301509 B6	31/03/2010		
DK 2032589 T3	09/01/2012		
EA 014854 B1	28/02/2011		
EA 200802353 A1	28/04/2009		
EP 2032589 A1	11/03/2009		
ES 2373991 T3	10/02/2012		
NZ 573326 A	30/09/2010		
PL 2032589 T3	29/02/2012		
PT 2032589 E	11/01/2012		
WO 2008-000196 A1	03/01/2008		
US 05698003 A	16/12/1997	AU 1997-77597 A	14/07/1997
		AU 708702 B2	12/08/1999
		BR 9612034 A	28/03/2000
		CA 2240284 A1	26/06/1997
		CA 2240284 C	18/01/2005
		CN 1163442 C	25/08/2004
		CN 1204310 A	06/01/1999
		DE 69637957 D1	06/08/2009
		EP 0869933 A1	14/10/1998
		EP 0869933 A4	20/10/2004
		EP 0869933 B1	24/06/2009

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/US2013/076858

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
		ES 2329201 T3	23/11/2009
		IN 186192 A1	07/07/2001
		JP 03282149 B2	13/05/2002
		JP 2000-502037 A	22/02/2000
		NZ 324974 A	30/08/1999
		WO 97-22568 A1	26/06/1997