



(86) Date de dépôt PCT/PCT Filing Date: 2001/09/17
 (87) Date publication PCT/PCT Publication Date: 2002/03/28
 (85) Entrée phase nationale/National Entry: 2003/03/18
 (86) N° demande PCT/PCT Application No.: EP 2001/010704
 (87) N° publication PCT/PCT Publication No.: 2002/024666
 (30) Priorité/Priority: 2000/09/20 (09/666,117) US

(51) Cl.Int.⁷/Int.Cl.⁷ C07D 239/94, A61K 31/517, A61P 9/10,
C07D 409/10, C07D 409/14, C07D 405/12, C07D 403/12


(71) Demandeur/Applicant:
MERCK PATENT GESELLSCHAFT MIT
BESCHRANKTER HAFTUNG, DE

(72) Inventeurs/Inventors:
MEDERSKI, WERNER, DE;
DEVANT, RALF, DE;
BARNICKEL, GERHARD, DE;
BERNOTAT-DANIELOWSKI, SABINE, DE;
VICKERS, JAMES, DE;
CEZANNE, BERTRAM, DE;

...

(74) Agent: FETHERSTONHAUGH & CO.

(54) Titre : 4-AMINO-QUINAZOLINES
 (54) Title: 4-AMINO-QUINAZOLINES



(57) Abrégé/Abstract:

Quinazolines of the formula I, in which R, R¹, R², R³, R⁴ and Y have the meaning indicated in Patent Claim 1, and their salts or solvates as glycoprotein IblX antagonists.

(72) **Inventeurs(suite)/Inventors(continued)**: DHANOA, DALJIT, US; ZHAO, BAO-PING, US; RINKER, JAMES, US; PLAYER, MARK R., US; JAEGER, EDWARD, US; SOLL, RICHARD, US

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
28 March 2002 (28.03.2002)

PCT

(10) International Publication Number
WO 02/024666 A3(51) International Patent Classification⁷: C07D 239/94,
409/10, 403/12, 405/12, A61K 31/517, A61P 9/10, C07D
409/141115 Fern Avenue, Kenhurst, PA 19607 (US). **PLAYER, Mark, R.** [US/US]; 5022 Swan Drive, Phoenixville, PA 19460 (US). **JAEGER, Edward** [US/US]; 210 Kelso Circle, Trappe, PA 19426 (US). **SOLL, Richard** [US/US]; 324 Glenn Avenue, Lawrenceville, NJ 08648 (US).

(21) International Application Number: PCT/EP01/10704

(22) International Filing Date:
17 September 2001 (17.09.2001)(74) Common Representative: **MERCK PATENT GMBH**;
Frankfurter Strasse 250, 64293 Darmstadt (DE).

(25) Filing Language: English

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(26) Publication Language: English

(30) Priority Data:
09/666,117 20 September 2000 (20.09.2000) US(71) Applicant (*for all designated States except US*): **MERCK PATENT GMBH** [DE/DE]; Frankfurter Strasse 250, 64293 Darmstadt (DE).


(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **MEDERSKI, Werner** [DE/DE]; Am Ohlenberg 29, 64390 Erzhausen (DE). **DEVANT, Ralf** [DE/DE]; Frankfurter Landstrasse 135, 64291 Darmstadt (DE). **BARNICKEL, Gerhard** [DE/DE]; Emilstrasse 27, 64298 Darmstadt (DE). **BERNOTAT-DANIELOWSKI, Sabine** [DE/DE]; Leibigstrasse 5, 61231 Bad Nauheim (DE). **VICKERS, James** [GB/DE]; Mörrikestrasse 6, 65719 Hofheim/Ts. (DE). **CEZANNE, Bertram** [DE/DE]; Goethestrasse 47, 64546 Mörfelden-Walldorf (DE). **DHANO, Daljit** [US/US]; 2037 Skiles Blvd, West Chester, PA 19382 (US). **ZHAO, Bao-Ping** [CN/US]; 3606 Hunters Glen Drive, Plainsboro, NJ 08536 (US). **RINKER, James** [US/US];(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).**Published:**

— with international search report

(88) Date of publication of the international search report:
26 September 2002*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*


(54) Title: 4-AMINO-QUINAZOLINES

(57) Abstract: Quinazolines of the formula I, in which R, R¹, R², R³, R⁴ and Y have the meaning indicated in Patent Claim 1, and their salts or solvates as glycoprotein IbIX antagonists.

4-Amino-quinazolines

The invention relates to substituted 4-amino-quinazolines of the formula I

5



in which


10

R and R¹ are independently of each other H, A, OH, OA, Hal, N(R⁵)₂, NO₂, CN, C(O)R², CON(R⁵)₂, COOR⁵, allyl, CH=CH-COOR⁵, CH=CHCON(R⁵)₂, SO₂A or phenyl, which is unsubstituted or mono-, di- or trisubstituted by A,

15

R² and R³ are independently of each other H, A, cycloalkyl, -Het³, -(CH₂)_o-OR⁵, -(CH₂)_o-OR⁶, -(CH₂)_o-Het¹, -(CH₂)_o-NR⁵-Het¹, -(CHA)_p-(CH₂)_o-N(R⁵)₂, -(CH₂)_p-(CHA)_p-(CH₂)_m-Ar, -(CH₂)_o-Z-(CH₂)_q-N(R⁵)₂,

20




25

provided that R² and R³ together are not H, or NR²R³ together form a saturated monocyclic heterocyclic radical having 5 to 6 ring members, where 1 or 2 N atoms are present and the heterocyclic radical can be mono- or disubstituted by OH, Ar, OAr or arylalkyl,

30

R⁴ is Ar or Het¹,
 R⁵ is H or A,
 R⁶ is benzo[1,3]dioxol-5-yl,
 Q is O or S,

- Y is (CH=CH)_n,
- Z is phenylene, cyclohexylene, -NR⁵-, O, -CH(OH)-, -CA₂- or
- 
- 5 A is unbranched or branched alkyl having 1 to 6 carbon atoms,
- Ar is phenyl, naphthyl or biphenyl, which is unsubstituted or mono-, di- or trisubstituted by A, OH, OA, cycloalkyloxy, O-(CH₂)_p-Ph, CF₃, OCF₃, Hal, CN, CHO, COA, COOR⁵, N(R⁵)₂, NR⁵-COA, NO₂, SO₂N(R⁵)₂, mor, SO₂-mor, 5-methyl-3-oxo-2,4-dihydropyrazol-2-yl, naphthyl or Het²,
- 10 Het¹ is a saturated, partially or completely unsaturated mono- or bicyclic heterocyclic radical having 5 to 10 ring members, where 1 or 2 N and/or 1 or 2 S or O atoms can be present and the heterocyclic radical can be mono- or disubstituted by A, Hal, OH, OA, CF₃, OCF₃, N(R⁵)₂, carbonyl oxygen, COOR⁵, Het², benzyl or phenyl which is unsubstituted or mono-, di- or trisubstituted by A, OH, OA, CF₃, OCF₃, Hal, CN, COOR⁵, N(R⁵)₂, NO₂ or SO₂N(R⁵)₂,
- 15 Het² is a unsaturated mono- or bicyclic heterocyclic radical having 5 to 10 ring members, where 1 or 2 N and/or 1 or 2 S or O atoms can be present and the heterocyclic radical can be mono- or disubstituted by A, Hal, OH, OA, CF₃, OCF₃, N(R⁵)₂ or COOR⁵,
- 20 Het³ is a partially or completely unsaturated mono- or bicyclic heterocyclic radical having 5 to 10 ring members, where 1 or 2 N atoms are present and the heterocyclic radical can be mono- or disubstituted by A, Hal, OH, OA, CF₃, OCF₃, N(R⁵)₂, SO₂A or COOR⁵ provided that the heterocyclic radical is not bonded via an N atom,
- 25 Hal is F, Cl, Br or I,
- 30 mor is morpholin-4-yl,

Ph is phenyl,
n is 1 or 2,
m is 0, 1, 2, 3, 4, 5 or 6,
o is 1, 2, 3, 4, 5, 6 or 7,
5 p is 0, 1, 2, 3 or 4,
q is 1, 2, 3 or 4,
and their pharmaceutically tolerable salts and solvates as glycoprotein IbIX antagonists.

10 Similar 4-amino substituted quinazolines are disclosed in WO 99/09986, Mastafanova, LI et al, Khim.-Farm.Zh. 1982, 16, 938-42 or DE 2135172.

The invention is based on the object of finding novel glycoprotein IbIX inhibitors which can be used for the production of medicaments.

15 It has been found that the compounds of the formula I according to claims 1 to 5 and their salts or solvates act especially as GPIbIX inhibitors, in particular inhibiting the interaction of this receptor with the ligand von Willebrand factor (vWF). This action can be demonstrated, for
20 example, by a method which is described by S. Meyer et al. in J. Biol. Chem. 1993, 268, 20555-20562. The property as GPIbIX alpha-thrombin receptor (N.J. Greco, Biochemistry 1996, 35, 915-921) can also be blocked by the compounds mentioned.

25 The significance of GPIbIX as an adhesion receptor on platelets, which mediates the primary interaction of platelets with an arteriosclerotically modified vascular wall via binding to the vWF expressed there, has been described by many authors (e.g. Z.M. Ruggeri in Thromb. Hemost. 1997, 78, 611-616). The activation of another platelet adhesion receptor,
30 GPIIbIIIa, following the GPIbIX-vWF interaction, leads to platelet aggregation and thus to thrombotic vascular occlusion.

A GPIbIX antagonist can thus prevent the start of thrombus formation and thus also release of active substances from the platelets which, for example, promote thrombus growth and have an additional trophic action on the vascular wall. This has been shown with inhibitory peptides or antibodies in various experimental models (e.g. H Yamamoto et al.,
5 Thromb. Hemost. 1998, 79, 202-210).

In the case of higher shear forces, the blocking action of GPIbIX inhibitors exerts its maximum effect, as described by J.J. Sixma et al. in
10 Arteriosclerosis, Thrombosis, and Vascular Biology 1996, 16, 64-71.
According to the flow chamber method used there, the compounds of the formula I can be characterized as GPIbIX inhibitors in whole blood.

The inhibition of thrombus formation of the GPIbIX inhibitors can be
15 measured by a modified Born method (Nature 1962, 4832, 927-929) using botrocetin or ristocetin as an aggregation stimulant.

The compounds of the formula I according to the invention can therefore be employed as pharmaceutical active compounds in human and
20 veterinary medicine. They act as adhesion receptor antagonists, in particular as glycoprotein IbIX antagonists, and are suitable for the prophylaxis and/or therapy of thrombotic disorders and sequelae deriving therefrom. The preferentially best action is to be expected in the case of thrombotic disorders in the arterial vascular system, but GPIbIX inhibitors
25 also have an effect in the case of thrombotic disorders in the venous vascular bed. The disorders are acute coronary syndromes, angina pectoris, myocardial infarct, peripheral circulatory disorders, stroke, transient ischaemic attacks, arteriosclerosis, reocclusion/restenosis after angioplasty/stent implantation. The compounds can furthermore be
30 employed as anti-adhesive substances where the body comes into contact with foreign surfaces such as implants, catheters or cardiac pacemakers.


Comparison medication introduced onto the market which may be mentioned are aspirin and GPIIb/IIIa antagonists.

The invention relates furthermore to novel compounds of the formula I and their salts or solvates, especially of compounds relating to group Ia to Ic, and to a process for the preparation of these novel compounds and their salts or solvates, characterized in that

a) a compound of the formula I according to claims 1 to 4 is liberated from one of its functional derivatives by treating with a solvolysing or hydrogenolysing agent,


or

b) in stage 1) a compound of the formula II




in which

R and R¹ have the meaning as given in Claims 1 to 4, is reacted with a compound of the formula III




in which R⁴ has the meaning indicated in Claims 1 to 4 and s is 0 or 1, to give a compound of formula IV



in which R, R¹ and R⁴ have the meaning indicated in Claims 1 to 4 and s is 0 or 1,

in stage 2) a compound of formula IV as indicated above is reacted with a chlorinating agent to give a compound of formula V


5



in which R, R¹ and R⁴ have the meaning indicated in Claims 1 to 4 and s is 0 or 1,

10

and in stage 3) a compound of formula V as indicated above is reacted with a compound of formula VI




15

in which R² and R³ or NR²R³ have the meaning indicated in Claims 1 to 4,
or

c) in stage 1) a compound of the formula II

20




in which

R and R¹ have the meaning as given in Claims 1 to 4,

25

is reacted with a chlorinating agent to give a compound of formula VII




30

in which

R and R¹ have the meaning as given in Claims 1 to 4,


in stage 2) a compound of formula VII as indicated above is reacted with a compound of formula VI



5

in which R^2 and R^3 or NR^2R^3 have the meaning indicated in Claims 1 to 4 to give a compound of formula VIII


10



in which R , R^1 , R^2 , R^3 and NR^2R^3 have the meaning indicated in Claims 1 to 4

15

and in stage 3) a compound of formula VIII as indicated above is reacted with a compound of formula III



20

in which R^4 has the meaning indicated in Claims 1 to 4 and s is 0 or 1 or

d) a radical R , R^1 , R^2 , R^3 and/or R^4 is converted into another radical R , R^1 , R^2 , R^3 and/or R^4 by, for example

25

- reducing a nitro group, sulfonyl group or sulfoxyl group,
- etherifying an OH group or subjecting an OA group to ether cleavage,
- alkylating a primary or secondary amino group,
- partially or completely hydrolysing a CN group,
- cleaving an ester group or esterifying a carboxylic acid radical,
- reacting an aryl bromide, aryl iodide, heteroaryl bromide or heteroaryliodide to give the corresponding coupling products by means of a Suzuki coupling with boronic acids,

30

- reacting a iodoquinazoline or bromoquinazoline to give the corresponding coupling products by means of a Stille coupling with allyltributyltin,
- 5 - reacting a iodoquinazoline or bromoquinazoline to give the corresponding coupling products by means of a Heck coupling with acrylates,
- or carrying out a nucleophilic or electrophilic substitution,

and/or

a base or acid of the formula I is converted into one of its salts or solvates.

10

The compounds of the formula I can have a chiral center and therefore occur in a number of stereoisomeric forms. All these forms (e.g. R and S forms) and their mixtures (e.g. the RS forms) are included in the formula I.

15

The compounds according to the invention also include so-called prodrug derivatives, i.e. compounds of the formula I modified with, for example, alkyl or acyl groups, sugars or oligopeptides and which are rapidly cleaved in the body to give the active compounds according to the invention.

20

Furthermore, free amino groups as substituents of compounds of the formula I can be provided with appropriate conventional protective groups. Solvates of the compounds of the formula I are understood as meaning adducts of inert solvent molecules to the compounds of the formula I which are formed on account of their mutual power of attraction. Solvates are, for

25

The abbreviations used have the following meanings:

- Ac acetyl,
- Bu n-butyl,
- 30 DBU 1,8-diazabicyclo[5.4.0]undec-7-ene,
- DMA dimethylacetamide,
- DMF dimethylformamide,

| | | |
|---|------|---------------------------------------|
| | dppf | 1,1'-bis(diphenylphosphino)ferrocene, |
| | Et | ethyl, |
| | Me | methyl, |
| | Ph | phenyl, |
| 5 | TEA | triethylamine, |
| | TFA | trifluoroacetic acid. |

In the above formulae, A is alkyl and has 1 to 6, preferably 1, 2, 3 or 4 C atoms. Alkyl is preferably methyl, furthermore ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl or tert-butyl, additionally also pentyl, 1-, 2- or 3-methylbutyl, 1,1-, 1,2- or 2,2-dimethylpropyl, 1-ethylpropyl, hexyl, 1-, 2-, 3- or 4-methylpentyl, 1,1-, 1,2-, 1,3-, 2,2-, 2,3- or 3,3-dimethylbutyl, 1- or 2-ethylbutyl, 1-ethyl-1-methylpropyl, 1-ethyl-2-methylpropyl, 1,1,2- or 1,2,2-trimethylpropyl.

15 A is preferentially methyl, ethyl, propyl, isopropyl, butyl or pentyl.

Ar is phenyl, naphthyl or biphenyl, which is unsubstituted or mono-, di- or trisubstituted by A, OH, OA, cycloalkyloxy, O-(CH₂)_p-Ph, CF₃, OCF₃, Hal, CN, CHO, COA, COOR⁵, N(R⁵)₂, NR⁵-COA, NO₂, SO₂N(R⁵)₂, mor, SO₂-mor, 5-methyl-3-oxo-2,4-dihydropyrazol-2-yl, naphthyl or Het².

Ar is preferentially phenyl, preferably - as indicated - mono- di- or trisubstituted phenyl, specifically preferentially phenyl, 2-, 3- or 4-methylphenyl, 2-, 3- or 4-ethylphenyl, 2-, 3- or 4-propylphenyl, 2-, 3- or 4-isopropylphenyl, 2-, 3- or 4-butylphenyl, 2-, 3- or 4-tert-butylphenyl, 2-, 3- or 4-aminophenyl, 2-, 3- or 4-N,N-dimethylaminophenyl, 2-, 3- or 4-sulfamoylphenyl, 2-, 3- or 4-nitrophenyl, 2-, 3- or 4-hydroxyphenyl, 2-, 3- or 4-methoxyphenyl, 2-, 3- or 4-ethoxyphenyl, 2-, 3- or 4-pentoxyphenyl, 2-, 3- or 4-phenoxyphenyl, 2-, 3- or 4-phenylmethoxyphenyl, 2-, 3- or 4-trifluoromethylphenyl, 2-, 3- or 4-trifluoromethoxyphenyl, 2-, 3- or 4-cyclopentyloxyphenyl, 2-, 3- or 4-carboxyphenyl, 2-, 3- or 4-(N,N-diethyl)sulfamoylphenyl, 2-, 3- or 4-cyanophenyl, 2-, 3- or 4-fluorophenyl,

2-, 3- or 4-chlorophenyl, 2-, 3- or 4-bromophenyl, 2,3-, 2,4-, 2,5-, 2,6-, 3,4-
 or 3,5-dichlorophenyl, 2,3-, 2,4-, 2,5-, 2,6-, 3,4- or 3,5-dibromophenyl, 2,3-,
 2,4-, 2,5-, 2,6-, 3,4- or 3,5-dimethoxyphenyl, 2,3-, 2,4-, 2,5-, 2,6-, 3,4- or
 3,5-di(trifluoromethyl)phenyl, 2,3-, 2,4-, 2,5-, 2,6-, 3,4- or 3,5-
 5 di(phenylmethoxy)phenyl, 2-chloro-6-methylphenyl, 2-chloro-4-
 fluorophenyl, 3-bromo-6-fluorophenyl, 3,4,5-trimethoxyphenyl, 4-
 (morpholin-4-yl)phenyl, 4-(morpholin-4-yl-sulfonyl)phenyl, 4-(5-methyl-3-
 oxo-2,4-dihydropyrazol-2-yl)phenyl, 4-(4,6-dimethoxy-pyrimidin-2-yl)phenyl,
 3-(4,6-dimethoxy-pyrimidin-2-yl)phenyl, 4-(pyrid-3-yl)phenyl, 3-(pyrid-3-
 10 yl)phenyl, 4-(thiophen-2-yl)phenyl, 3-(thiophen-2-yl)phenyl, 4-
 (benzo[c]thiophen-2-yl)phenyl or 4-(naphthalen-1-yl)phenyl.

Furthermore, however, Ar is also preferentially unsubstituted naphthyl or
 biphenyl - as indicated - or alternatively mono-, di- or trisubstituted
 15 biphenyl, specifically preferentially biphenyl-4-yl or biphenyl-3-yl,
 2'-methylbiphenyl-4-yl, 3'-methylbiphenyl-4-yl, 4'-methylbiphenyl-4-yl,
 2'-methylbiphenyl-3-yl, 3'-methylbiphenyl-3-yl, 4'-methylbiphenyl-3-yl,
 2-methylbiphenyl-4-yl, 3-methylbiphenyl-4-yl, 2-methylbiphenyl-3-yl,
 4-methylbiphenyl-3-yl, 2'-tert-butylbiphenyl-4-yl, 3'-tert-butylbiphenyl-4-yl,
 20 4'-tert-butylbiphenyl-4-yl, 2'-tert-butylbiphenyl-3-yl, 3'-tert-butylbiphenyl-3-yl,
 4'-tert-butylbiphenyl-3-yl, 2-tert-butylbiphenyl-4-yl, 3-tert-butylbiphenyl-4-yl,
 2-tertbutylbiphenyl-3-yl, 4-tert-butylbiphenyl-3-yl, 2'-isopropylbiphenyl-4-yl,
 3'-isopropylbiphenyl-4-yl, 4'-isopropylbiphenyl-4-yl, 2'-isopropylbiphenyl-3-
 yl, 3'-isopropylbiphenyl-3-yl, 4'-isopropylbiphenyl-3-yl, 2-isopropylbiphenyl-
 25 4-yl, 3-isopropylbiphenyl-4-yl, 2-isopropylbiphenyl), 4-isopropylbiphenyl-3-
 yl, 2'-fluorobiphenyl-4-yl, 3'-fluorobiphenyl-4-yl, 4'-fluorobiphenyl-4-yl,
 2'-fluorobiphenyl-3-yl, 3'-fluorobiphenyl-3-yl, 4'-fluorobiphenyl-3-yl,
 2-fluorobiphenyl-4-yl, 3-fluorobiphenyl-4-yl, 2-fluorobiphenyl-3-yl, 4-fluoro-
 biphenyl-3-yl, 2'-chlorobiphenyl-4-yl, 3'-chlorobiphenyl-4-yl, 4'-chloro-
 30 biphenyl-4-yl, 2'-chlorobiphenyl-3-yl, 3'-chlorobiphenyl-3-yl,
 4'-chlorobiphenyl-3-yl, 2-chlorobiphenyl-4-yl, 3-chlorobiphenyl-4-yl,

2-chlorobiphenyl-3-yl, 4-chlorobiphenyl-3-yl, 2'-methoxybiphenyl-4-yl,
3'-methoxybiphenyl-4-yl, 4'-methoxybiphenyl-4-yl, 2'-methoxybiphenyl-3-yl,
3'-methoxybiphenyl-3-yl, 4'-methoxybiphenyl-3-yl, 2-methoxybiphenyl-4-yl,
3-methoxybiphenyl-4-yl, 2-methoxybiphenyl-3-yl, 4-methoxybiphenyl-3-yl,
5 2'-nitrobiphenyl-4-yl, 3'-nitrobiphenyl-4-yl, 4'-nitrobiphenyl-4-yl, 2'-nitro-
biphenyl-3-yl, 3'-nitrobiphenyl-3-yl, 4'-nitrobiphenyl-3-yl, 2-nitrobiphenyl-4-yl,
3-nitrobiphenyl-4-yl, 2-nitrobiphenyl-3-yl, 4-nitrobiphenyl-3-yl,
2'-trifluoromethylbiphenyl-4-yl, 3'-trifluoromethylbiphenyl-4-yl, 4'-tri-
fluoromethyl-biphenyl-4-yl, 2'-trifluoromethylbiphenyl-3-yl,
10 3'-trifluoromethylbiphenyl-3-yl, 4'-trifluoromethylbiphenyl-3-yl,
2-trifluoromethylbiphenyl-4-yl, 3-trifluoromethylbiphenyl-4-yl,
2-trifluoromethylbiphenyl-3-yl, 4-trifluoromethylbiphenyl-3-yl,
2'-trifluoromethoxybiphenyl-4-yl, 3'-trifluoromethoxybiphenyl-4-yl, 4'-tri-
fluoromethoxybiphenyl-4-yl, 2'-trifluoromethoxybiphenyl-3-yl,
15 3'-trifluoromethoxybiphenyl-3-yl, 4'-trifluoromethoxybiphenyl-3-yl,
2-trifluoromethoxybiphenyl-4-yl, 3-trifluoromethoxybiphenyl-4-yl,
2-trifluoromethoxybiphenyl-3-yl, 4-trifluoromethoxybiphenyl-3-yl, 3'-
acetyl-biphenyl-4-yl, 3'-acetylamino-biphenyl-4-yl, 3'-amino-biphenyl-4-yl,
furthermore preferentially disubstituted biphenyls, such as 2'-methyl-3'-
20 nitrobiphenyl-4-yl, 2'-methyl-4'-nitrobiphenyl-4-yl, 2'-methyl-5'-nitrobiphenyl-
4-yl, 2'-methyl-6'-nitrobiphenyl-4-yl, 3'-methyl-2'-nitrobiphenyl-4-yl, 3'-
methyl-4'-nitrobiphenyl-4-yl, 3'-methyl-5'-nitrobiphenyl-4-yl, 3'-methyl-6'-
nitrobiphenyl-4-yl, 4'-methyl-2'-nitrobiphenyl-4-yl, 4'-methyl-3'-nitrobiphenyl-
4-yl, 2'-methyl-3'-nitrobiphenyl-3-yl, 2'-methyl-4'-nitrobiphenyl-3-yl,
25 2'-methyl-5'-nitrobiphenyl-3-yl, 2'-methyl-6'-nitrobiphenyl-3-yl, 3'-methyl-2'-
nitrobiphenyl-3-yl, 3'-methyl-4'-nitrobiphenyl-3-yl, 3'-methyl-5'-nitrobiphenyl-
3-yl, 3'-methyl-6'-nitrobiphenyl-3-yl, 4'-methyl-2'-nitrobiphenyl-3-yl,
4'-methyl-3'-nitrobiphenyl-3-yl, 2'-methoxy-2-methylbiphenyl-4-yl,
3'-methoxy-2-methylbiphenyl-4-yl, 4'-methoxy-2-methylbiphenyl-4-yl,
30 4'-methoxy-3-nitrobiphenyl-4-yl, 2'-chloro-3'-fluorobiphenyl-4-yl, 2'-chloro-4'-
fluorobiphenyl-4-yl, 2'-chloro-5'-fluorobiphenyl-4-yl, 2'-chloro-6'-
fluorobiphenyl-4-yl, 3'-chloro-2'-fluorobiphenyl-4-yl, 3'-chloro-4'-fluoro-

biphenyl-4-yl, 3'-chloro-5'-fluorobiphenyl-4-yl, 3'-chloro-6'-fluorobiphenyl-4-yl, 4'-chloro-2'-fluorobiphenyl-4-yl, 4'-chloro-3'-fluorobiphenyl-4-yl, 2'-chloro-3'-fluorobiphenyl-3-yl, 2'-chloro-4'-fluorobiphenyl-3-yl, 2'-chloro-5'-fluorobiphenyl-3-yl, 2'-chloro-6'-fluorobiphenyl-3-yl, 3'-chloro-2'-fluorobiphenyl-3-yl, 3'-chloro-4'-fluorobiphenyl-3-yl, 3'-chloro-5'-fluorobiphenyl-3-yl, 3'-chloro-6'-fluorobiphenyl-3-yl, 4'-chloro-2'-fluorobiphenyl-3-yl, 4'-chloro-3'-fluorobiphenyl-3-yl, (2,3'-diethyl)biphenyl-4-yl, (3,3'-diethyl)biphenyl-4-yl, (2,2'-diethyl)biphenyl-4-yl, (2,4'-diethyl)biphenyl-4-yl, (2',3'-dimethoxy)biphenyl-4-yl, (2',4'-dimethoxy)biphenyl-4-yl, (2',5'-dimethoxy)biphenyl-4-yl, (2',6'-dimethoxy)biphenyl-4-yl, (3',4'-dimethoxy)biphenyl-4-yl, (3',5'-dimethoxy)biphenyl-4-yl, (2',3'-dimethoxy)biphenyl-3-yl, (2',4'-dimethoxy)biphenyl-3-yl, (2',5'-dimethoxy)biphenyl-3-yl, (2',6'-dimethoxy)biphenyl-3-yl, (3',4'-dimethoxy)biphenyl-3-yl, (3',5'-dimethoxy)biphenyl-3-yl, (3',5'-dichloro)biphenyl-4-yl, (3',5'-dichloro)biphenyl-3-yl, (2',4'-dichloro)biphenyl-4-yl, (3',4',5'-trimethoxy)biphenyl-4-yl, (2',3'-di(trifluoromethyl))biphenyl-4-yl, (2',4'-di(trifluoromethyl))biphenyl-4-yl, (2',5'-di(trifluoromethyl))biphenyl-4-yl, (2',6'-di(trifluoromethyl))biphenyl-4-yl, (3',4'-di(trifluoromethyl))biphenyl-4-yl, (3',5'-di(trifluoromethyl))biphenyl-4-yl, (2',3'-di(trifluoromethyl))biphenyl-3-yl, (2',4'-di(trifluoromethyl))biphenyl-3-yl, (2',5'-di(trifluoromethyl))biphenyl-3-yl, (2',6'-di(trifluoromethyl))biphenyl-3-yl, (3',4'-di(trifluoromethyl))biphenyl-3-yl, (3',5'-di(trifluoromethyl))biphenyl-3-yl, (2,2'-dimethyl)biphenyl-4-yl, (2,3'-dimethyl)biphenyl-4-yl, (2,4'-dimethyl)biphenyl-4-yl, (2,2'-dimethyl)biphenyl-3-yl, (2,3'-dimethyl)biphenyl-3-yl or (2,4'-dimethyl)biphenyl-3-yl.

Phenyl, 2-, 3- or 4-fluorophenyl, 3- or 4-chlorophenyl, 4-bromophenyl, 2,4- or 3,4-dichlorophenyl, 2,3-dimethoxyphenyl, 4-aminophenyl, 4-dimethylaminophenyl, 2-, 3- or 4-methylphenyl, 2-, 3- or 4-methoxyphenyl, 4-propylphenyl, 4-isopropylphenyl, 4-butylphenyl, 2-, 3- or 4-nitrophenyl, 2-, 3- or 4-pentoxyphenyl, 2-, 3- or 4-phenoxyphenyl, 2-, or 4-phenylmethoxyphenyl, 2-, 3- or 4-trifluoromethylphenyl, 2-, 3- or 4-

trifluoromethoxyphenyl, 2- or 4-cyclopentyloxyphenyl, 3- or 4-carboxyphenyl, 2-, 3- or 4-(N,N-diethyl)sulfamoylphenyl, 3,4-di(phenylmethoxy)phenyl, 2-chloro-6-methylphenyl, 2-chloro-4-fluorophenyl, 5-bromo-2-fluorophenyl, 3,4,5-trimethoxyphenyl, 4-
5 (morpholin-4-yl)phenyl, 4-(morpholin-4-yl-sulfonyl)phenyl, 4-(5-methyl-3-oxo-2,4-dihydropyrazol-2-yl)phenyl, 4-(4,6-dimethoxy-pyrimidin-2-yl)phenyl, 3-(4,6-dimethoxy-pyrimidin-2-yl)phenyl, 4-(pyrid-3-yl)phenyl, 3-(pyrid-3-yl)phenyl, 4-(thiophen-2-yl)phenyl, 3-(thiophen-2-yl)phenyl, 4-(benzo[c]thiophen-2-yl)phenyl, 4-(naphthalen-1-yl)phenyl, naphthyl,
10 biphenyl-4-yl, 2'-fluorobiphenyl-4-yl, 4'-fluorobiphenyl-4-yl, 4'-fluorobiphenyl-3-yl, 4'-chlorobiphenyl-4-yl, 4'-chlorobiphenyl-3-yl, 4'-methoxybiphenyl-4-yl, 4'-methoxybiphenyl-3-yl, 3'-nitrobiphenyl-4-yl, 3'-acetylbiphenyl-4-yl, 3'-acetylamino-biphenyl-4-yl, 3'-aminobiphenyl-4-yl, (2,3'-diethyl)biphenyl-4-yl, (3',5'-dichloro)biphenyl-3-yl, (2',4'-dichloro)biphenyl-4-yl, (3',4',5'-
15 trimethoxy)biphenyl-4-yl, (3',5'-di(trifluoromethyl))biphenyl-4-yl is particularly preferred for Ar.

Arylalkyl is preferentially benzyl.

O-(CH₂)_p-Ph is phenylalkyloxy, in which p can be 0, 1, 2, 3 or 4. Benzyloxy
20 or phenyloxy is particularly preferred.

Cycloalkyl preferably has 3-7 C atoms and is preferably cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl, and further also cyclopentylmethyl, cyclopentylethyl or cyclohexylmethyl; cyclopentyl, cyclohexylmethyl or
25 cyclohexyl are particularly preferred.

Hal is preferably F, Cl, Br or I.

Het¹ is a saturated, partially or completely unsaturated mono- or bicyclic
30 heterocyclic radical having 5 to 10 ring members, where 1 or 2 N and/or 1 or 2 S or O atoms can be present and the heterocyclic radical can be mono- or disubstituted by A, Hal, OH, OA, CF₃, OCF₃, N(R⁵)₂, carbonyl

oxygen, COOR⁵, Het², benzyl or phenyl which is unsubstituted or mono-, di- or trisubstituted by A, OH, OA, CF₃, OCF₃, Hal, CN, COOR⁵, N(R⁵)₂, NO₂, SO₂N(R⁵)₂.

- 5 Het¹ is preferably unsubstituted 2- or 3-furyl, 2- or 3-thiophenyl, 1-, 2- or 3-pyrrolyl, 1-, 2-, 4- or 5-imidazolyl, 1-, 3-, 4- or 5-pyrazolyl, 2-, 4- or 5-oxazolyl, 3-, 4- or 5-isoxazolyl, 2-, 4- or 5-thiazolyl, 3-, 4- or 5-isothiazolyl, 2-, 3- or 4-pyridyl, 2-, 4-, 5- or 6-pyrimidinyl, furthermore preferably 1,2,3-triazol-1-, -4- or -5-yl, 1,2,4-triazol-1-, -4- or -5-yl,
 10 1- or 5-tetrazolyl, 1,2,3-oxadiazol-4- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,2,4-thiadiazol-3- or -5-yl, 1,2,3-thiadiazol-4- or -5-yl, 2-, 3-, 4-, 5- or 6-2H-thiopyranyl, 2-, 3- or 4-4H-thiopyranyl, 3- or 4-pyridazinyl, pyrazinyl, 2-, 3-, 4-, 5-, 6- or 7-benzofuryl, 2-, 3-, 4-, 5-, 6- or 7-benzothiophenyl, 1-, 2-, 3-, 4-, 5-, 6-
 15 or 7-1H-indolyl, 1-, 2-, 4- or 5-benzimidazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzopyrazolyl, 2-, 4-, 5-, 6- or 7-benzoxazolyl, 3-, 4-, 5-, 6- or 7-benzisoxazolyl, 2-, 4-, 5-, 6- or 7-benzothiazolyl, 2-, 4-, 5-, 6- or 7-benzisothiazolyl, 4-, 5-, 6- or 7-benz-2,1,3-oxadiazolyl, 1-, 2-, 3-, 4-, 5-, 6-, 7- or 8-quinolinyl, 1-, 3-, 4-, 5-, 6-, 7- or 8-isoquinolinyl, 1-,
 20 2-, 3-, 4- or 9-carbazolyl, 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8- or 9-acridinyl, 3-, 4-, 5-, 6-, 7- or 8-cinnolinyl, 2-, 4-, 5-, 6-, 7- or 8-quinazolinyl. The heterocyclic radicals can also be partially or completely hydrogenated. Het can thus also be 2,3-dihydro-2-, -3-, -4- or -5-furyl, 2,5-dihydro-2-, -3-, -4- or -5-furyl, tetrahydro-2- or -3-furyl, 1,3-dioxolan-4-yl,
 25 tetrahydro-2- or -3-thiophenyl, 2,3-dihydro-1-, -2-, -3-, -4- or -5-pyrrolyl, 2,5-dihydro-1-, -2-, -3-, -4- or -5-pyrrolyl, 1-, 2- or 3-pyrrolidinyl, tetrahydro-1-, -2- or -3-pyrrolyl, tetrahydro-1-, -2- or 4-imidazolyl, 2,3-dihydro-1-, -2-, -3-, -4-, -5-, -6-, -7-1H-indolyl, 2,3-dihydro-1-, -2-, -3-, -4- or -5-pyrazolyl, tetrahydro-1-, -3-
 30 or -4-pyrazolyl, 1,4-dihydro-1-, -2-, -3- or -4-pyridyl, 1,2,3,4-tetrahydro-1-, -2-, -3-, -4-, -5- or -6-pyridyl, 1,2,3,6-tetrahydro-1-, -2-, -3-, -4-, -5- or -6-pyridyl, 1-, 2-, 3- or

4-piperidinyl, 1-, 2-, 3- or 4-azepanyl, 2-, 3- or 4-morpholinyl, tetrahydro-2-, -3- or -4-pyranyl, 1,4-dioxanyl, 1,3-dioxan-2-, -4- or -5-yl, hexahydro-1-, -3- or -4-pyridazinyl, hexahydro-1-, -2-, -4- or -5-pyrimidinyl, 1-, 2- or 3-piperazinyl, 1,2,3,4-tetrahydro-1-, -2-, -3-,
 5 -4-, -5-, -6-, -7- or -8-quinolinyl, 1,2,3,4-tetrahydro-1-, -2-, -3-, -4-, -5-, -6-, -7- or -8-isoquinolinyl which can be substituted as indicated above or particularly substituted by A, OA, carbonyl oxygen, NO₂, Het² or phenyl which is substituted by Hal, CN or OA.

10 Thiophen-2-yl, tetrahydro-furan-2-yl, 1-methyl-octahydro-indol-3-yl, benzo[1,3]dioxol-5-yl, piperazin-1-yl, 4-methyl-piperazin-1-yl, piperidin-1-yl, piperidin-4-yl, 4-benzyl-piperidin-1-yl, 2-methyl-piperidin-1-yl, 1-ethyl-pyrrolidin-2-yl, 1-methyl-pyrrolidin-2-yl, 2-oxo-pyrrolidin-1-yl, pyridin-2-yl, pyridin-4-yl, 5-nitro-pyridin-2-yl, imidazol-1-yl, morpholin-4-yl, 5-methoxy-
 15 1H-indol-2-yl, 5-(3-chlorophenyl)-furan-2-yl, 5-(4-fluorophenyl)-thiophen-2-yl, 5-(2-methoxyphenyl)-thiophen-2-yl, 5-(2-cyanophenyl)-thiophen-2-yl, 5-(2,5-dimethoxyphenyl)-thiophen-2-yl, 2-[2,2']bithiophenyl-5-yl, 5-(pyridin-4-yl)-thiophen-2-yl, 5-(1H-indol-5-yl)-thiophen-2-yl, 5-quinolin-8-yl-thiophen-2-yl or 5-(benzo[b]thiophen-2-yl)-thiophen-2-yl is particularly preferred for
 20 Het¹.

Het² is a unsaturated mono- or bicyclic heterocyclic radical having 5 to 10 ring members, where 1 or 2 N and/or 1 or 2 S or O atoms can be present and the heterocyclic radical can be mono- or disubstituted by A,
 25 Hal, OH, OA, CF₃, OCF₃, N(R⁵)₂ or COOR⁵.

Thiophen-2-yl, pyridin-3-yl, pyridin-2-yl, pyridin-4-yl, indol-5-yl, quinolin-8-yl, 4,6-dimethoxy-pyrimidin-2-yl or benzo[b]thiophen-2-yl is particularly preferred for Het².

30

Het³ is a partially or completely unsaturated mono- or bicyclic heterocyclic radical having 5 to 10 ring members, where 1 or 2 N atoms are present and

the heterocyclic radical can be mono- or disubstituted by A, Hal, OH, OA, CF₃, OCF₃, N(R⁵)₂, SO₂A or COOR⁵ provided that the heterocyclic radical is not bonded via an N atom.

5 Quinolin-5-yl and 1-methanesulfonyl-2,3-dihydro-1H-indol-5-yl is particularly preferred for Het³.


(CH₂)_o-Het¹ is preferentially thiophen-2-yl-ethyl, tetrahydro-furan-2-yl-methyl, 1-methyl-octahydro-indol-3-yl-methyl, 1-methyl-octahydro-indol-3-yl-ethyl benzo[1,3]dioxol-5-yl-methyl, benzo[1,3]dioxol-5-yl-ethyl, piperazin-10 1-yl-ethyl, 4-methyl-piperazin-1-yl-propyl, piperidin-1-yl-ethyl, piperidin-4-yl-methyl, 1-methyl-piperidin-3-yl-ethyl, 4-benzyl-piperidin-1-yl-ethyl, 2-methyl-piperidin-1-yl-propyl, 1-ethyl-pyrrolidin-2-yl-methyl, 1-methyl-pyrrolidin-2-yl-methyl, 2-oxo-pyrrolidin-1-yl-propyl, pyridin-2-yl-ethyl, pyridin-4-yl-methyl, 15 pyridin-4-yl-ethyl, imidazol-1-yl-propyl, morpholin-4-yl-propyl or morpholin-4-yl-ethyl.

Piperidin-4-yl-methyl, 2-oxo-pyrrolidin-1-yl-propyl, pyridin-4-yl-methyl, imidazol-1-yl-propyl or morpholin-4-yl-propyl is particularly preferred for (CH₂)_o-Het¹.

20


(CH₂)_o-OR⁵ is preferentially (CH₂)₂-OCH₃, (CH₂)₃-OCH₃ or (CH₂)₃-O(iPr).
(CH₂)_o-OR⁶ is preferentially


25




(CH₂)_o-Z-(CH₂)_q-N(R⁵)₂ is preferentially

30







or $\text{---(CH}_2\text{)}_3\text{---O---(CH}_2\text{)}_3\text{---NH}_2$.





is particularly preferred for $(\text{CH}_2)_o\text{---Z---(CH}_2\text{)}_q\text{---N(R}^5\text{)}_2$.



$(\text{CH}_2)_p\text{---(CHA)}_p\text{---(CH}_2\text{)}_m\text{---Ar}$ is preferentially phenyl,











is particularly preferred for $(\text{CH}_2)_p\text{-(CHA)}_p\text{-(CH}_2)_m\text{-Ar}$.

15 R and R¹ are independently of each other H, A, OH, OA, Hal, N(R⁵)₂, NO₂, CN, C(O)R², CON(R⁵)₂, COOR⁵, allyl, CH=CH-COOR⁵, CH=CHCON(R⁵)₂, SO₂A or phenyl, which is unsubstituted or mono-, di- or trisubstituted by A, where A and Hal have a preferred meaning indicated beforehand and R⁵ have a preferred meaning indicated in the following.


R is preferentially H.

20 R¹ is preferentially H, Hal, allyl, CH=CH-COOR⁵, CH=CHCON(R⁵)₂ or phenyl, which is unsubstituted or monosubstituted by A. H, Cl, Br, I, CH=CH-COOEt, 4-methylphenyl, allyl or CH=CH-CONMe₂ is particularly preferred for R¹.

25 The preferred position of R¹ is the 6- or 7-position of the quinazoline ring system.

30 R² and R³ are independently of each other H, A, cycloalkyl, -Het³, -(CH₂)_o-OR⁵, -(CH₂)_o-OR⁶, -(CH₂)_o-Het¹, -(CH₂)_o-NR⁵-Het¹, -(CHA)_p-(CH₂)_o-N(R⁵)₂, -(CH₂)_p-(CHA)_p-(CH₂)_m-Ar, -(CH₂)_o-Z-(CH₂)_q-N(R⁵)₂,

5




provided that R^2 and R^3 together are not H, where A, Ar, cycloalkyl, Het¹ or Het³ have a preferred meaning indicated beforehand and R^5 , R^6 , Q and Z have a preferred meaning indicated in the following.

10

R^2 is preferentially H or A.

R^3 is preferentially A, cycloalkyl, -Het³, $-(CH_2)_o-OR^5$, $-(CH_2)_o-OR^6$, $-(CH_2)_o-Het^1$, $-(CH_2)_o-NR^5-Het^1$, $-(CHA)_p-(CH_2)_o-N(R^5)_2$, $-(CH_2)_p-(CHA)_p-(CH_2)_m-Ar$, $-(CH_2)_o-Z-(CH_2)_q-N(R^5)_2$,

15




20


Furthermore NR^2R^3 together form a saturated monocyclic heterocyclic radical having 5 to 6 ring members, where 1 or 2 N atoms are present and the heterocyclic radical can be mono- or disubstituted by OH, Ar, OAr or arylalkyl, where Ar or arylalkyl have a preferred meaning indicated beforehand.

25

Preferred saturated monocyclic heterocyclic radicals can be piperidine or piperazine.

30





5

are particularly preferred for NR^2R^3 .

R^4 is Ar or Het¹, where Ar or Het¹ have a preferred meaning indicated beforehand.

10


R^5 is H or A, where A has a preferred meaning indicated beforehand.

Q is O or S, preferentially O.

Y is $(\text{CH}=\text{CH})_n$, where n can be 1 or 2.

15

Z is phenylene, cyclohexylene, $-\text{NR}^5-$, O, $-\text{CH}(\text{OH})-$, $-\text{CA}_2-$ or



, where R^5 and A have a preferred meaning indicated

20

beforehand. Phenylene and/or cyclohexylene are particularly bonded in 1,4- or 1,3-position.

m is 0, 1, 2, 3, 4, 5 or 6, preferentially 0, 1 or 2.

o is 1, 2, 3, 4, 5, 6 or 7, preferentially 1, 2, 3 or 7.

p is 0, 1, 2, 3 or 4, preferentially 0, 1 or 2.

25

q is 1, 2, 3 or 4, preferentially 1, 2 or 3.

30

Some preferred groups of compounds of formula I which are novel can be expressed by the following groups or subformulae Ia to Ic, which correspond to the formula I and in which the radicals not designated in greater detail have the meaning indicated in formula I according to Claim 1, but

I. for the group Ia

in which in Ia-1

5 R and R¹ are independently of each other H, A, OH, OA, Hal, N(R⁵)₂, NO₂, CN, C(O)R², CON(R⁵)₂, COOR⁵, allyl, CH=CH-COOR⁵, CH=CHCON(R⁵)₂, SO₂A or phenyl, which is unsubstituted or mono-, di- or trisubstituted by A,

R² is H,


R³ is -(CH₂)_o-Z-(CH₂)_q-N(R⁵)₂,

R⁴ is Ar,

10 R⁵ is H or A,

Y is (CH=CH)_n,

Z is phenylene, cyclohexylene, -NR⁵-, O, -CH(OH)-, -CA₂- or



15

A is unbranched or branched alkyl having 1 to 6 carbon atoms,

Ar is phenyl or naphthyl, which is unsubstituted or mono-, di- or trisubstituted by A, OH, OA, CF₃, OCF₃, Hal, CN, CHO, COA, COOR⁵, N(R⁵)₂, NO₂ or SO₂N(R⁵)₂,

20 Hal is F, Cl, Br or I,

n is 1 or 2,

o is 1, 2, 3, 4, 5, 6 or 7 and

q is 1, 2, 3 or 4;

25 in Ia-2

R and R¹ are independently of each other H or Hal,

R² is H,


R³ is -(CH₂)_o-Z-(CH₂)_q-N(R⁵)₂,

R⁴ is Ar,

30 R⁵ is H or A,

Y is (CH=CH)_n,


Z is phenylene, cyclohexylene, -NR⁵-, O, -CH(OH)-, -CA₂- or



- 5 A is unbranched or branched alkyl having 1 to 6 carbon atoms,
 Ar is phenyl or naphthyl, which is unsubstituted or mono-, di- or
 trisubstituted by A, OH, OA, Hal, COOR⁵, N(R⁵)₂ or NO₂,
 Hal is F, Cl, Br or I,
 n is 1 or 2,
 o is 1, 2 or 3 and
 10 q is 1, 2 or 3;

in Ia-3

- R and R¹ are independently of each other H or Hal,
 R² is H,
 15 R³ is -(CH₂)_o-Z-(CH₂)_q-N(R⁵)₂,




-(CH₂)_o-Z-(CH₂)_q-N(R⁵)₂ is

- 20 R⁴ is Ar,
 R⁵ is H or A,
 Y is (CH=CH)_n,
 A is unbranched or branched alkyl having 1 to 6 carbon atoms,
 Ar is phenyl or naphthyl, which is unsubstituted or mono-, di- or
 trisubstituted by A, OH, OA, Hal, COOR⁵, N(R⁵)₂ or NO₂,
 25 Hal is F, Cl, Br or I and
 n is 1 or 2;

II. for the group Ib

in which in Ib-1

- 30 R and R¹ are independently of each other H, A, OH, OA, Hal, N(R⁵)₂,
 NO₂, CN, C(O)R², CON(R⁵)₂, COOR⁵, allyl, CH=CH-COOR⁵,

- CH=CHCON(R⁵)₂, SO₂A or phenyl, which is unsubstituted or mono-, di- or trisubstituted by A,
- R² and R³ are independently of each other H, A, cycloalkyl, -Het³,
 -(CH₂)_o-OR⁵, -(CH₂)_o-OR⁶, -(CH₂)_o-Het¹, -(CH₂)_o-NR⁵-Het¹,
 5 -(CHA)_p-(CH₂)_o-N(R⁵)₂, -(CH₂)_p-(CHA)_p-(CH₂)_m-Ar or
 -(CH₂)_o-Z-(CH₂)_q-N(R⁵)₂,
 provided that R² and R³ together are not H,
- R⁴ is Ar,
 R⁵ is H or A,
 10 R⁶ is benzo[1,3]dioxol-5-yl,
 Y is (CH=CH)_n,
 Z is phenylene, cyclohexylene, -NR⁵-, O, -CH(OH)-, -CA₂- or
- 
- 15 A is unbranched or branched alkyl having 1 to 6 carbon atoms,
 Ar is phenyl, which is mono-, di- or trisubstituted by O-(CH₂)_p-Ph,
 naphthyl or Het², or biphenyl, which is unsubstituted or mono-,
 di- or trisubstituted by A, OH, OA, CF₃, OCF₃, Hal, CN, CHO,
 20 COA, COOR⁵, N(R⁵)₂, NR⁵-COA, NO₂, SO₂N(R⁵)₂, naphthyl or
 Het²,
- Het¹ is a saturated, partially or completely unsaturated mono- or
 bicyclic heterocyclic radical having 5 to 10 ring members,
 where 1 or 2 N and/or 1 or 2 S or O atoms can be present and
 25 the heterocyclic radical can be mono- or disubstituted by A,
 Hal, OH, OA, CF₃, OCF₃, N(R⁵)₂, carbonyl oxygen, COOR⁵,
 Het², benzyl or phenyl which is unsubstituted or mono-, di- or
 trisubstituted by A, OH, OA, CF₃, OCF₃, Hal, CN, COOR⁵,
 N(R⁵)₂, NO₂ or SO₂N(R⁵)₂,
- 30 Het² is a unsaturated mono- or bicyclic heterocyclic radical having
 5 to 10 ring members, where 1 or 2 N and/or 1 or 2 S or
 O atoms can be present and the heterocyclic radical can be

- mono- or disubstituted by A, Hal, OH, OA, CF₃, OCF₃, N(R⁵)₂ or COOR⁵,
- 5 Het³ is a partially or completely unsaturated mono- or bicyclic heterocyclic radical having 5 to 10 ring members, where 1 or 2 N atoms are present and the heterocyclic radical can be mono- or disubstituted by A, Hal, OH, OA, CF₃, OCF₃, N(R⁵)₂, SO₂A or COOR⁵ provided that the heterocyclic radical is not bonded via an N atom,
- 10 Hal is F, Cl, Br or I,
- Ph is phenyl,
- n is 1 or 2,
- m is 0, 1, 2, 3, 4, 5 or 6,
- o is 1, 2, 3, 4, 5, 6 or 7,
- p is 0, 1, 2, 3 or 4 and
- 15 q is 1, 2, 3 or 4;

in which in Ib-2

- R and R¹ are independently of each other H, Hal, allyl, CH=CH-COOR⁵, CH=CHCON(R⁵)₂ or phenyl, which is unsubstituted or mono-,
- 20 R² and R³ are independently of each other H, cycloalkyl, -(CH₂)_o-Het¹, -(CHA)_p-(CH₂)_o-N(R⁵)₂, -(CH₂)_p-(CHA)_p-(CH₂)_m-Ar or -(CH₂)_o-Z-(CH₂)_q-N(R⁵)₂, provided that R² and R³ together are not H,
- 25 R⁴ is Ar,
- R⁵ is H or A,
- Y is (CH=CH)_n,
- Z is phenylene, cyclohexylene, -NR⁵-, O, -CH(OH)-, -CA₂- or





- A is unbranched or branched alkyl having 1 to 6 carbon atoms,

- Ar is phenyl, which is mono-, di- or trisubstituted by O-(CH₂)_p-Ph, naphthyl or Het², or biphenyl, which is unsubstituted or mono-, di- or trisubstituted by A, OH, OA, CF₃, OCF₃, Hal, CN, CHO, COA, COOR⁵, N(R⁵)₂, NR⁵-COA, NO₂, SO₂N(R⁵)₂, naphthyl or Het²,
- 5 Het¹ is a saturated, partially or completely unsaturated mono- or bicyclic heterocyclic radical having 5 to 10 ring members, where 1 or 2 N and/or 1 or 2 S or O atoms can be present and the heterocyclic radical can be mono- or disubstituted by A, Hal, OH, OA, CF₃, OCF₃, N(R⁵)₂, carbonyl oxygen, COOR⁵, Het², benzyl or phenyl which is unsubstituted or mono-, di- or trisubstituted by A, OH, OA, CF₃, OCF₃, Hal, CN, COOR⁵, N(R⁵)₂, NO₂ or SO₂N(R⁵)₂,
- 10 Het² is a unsaturated mono- or bicyclic heterocyclic radical having 5 to 10 ring members, where 1 or 2 N and/or 1 or 2 S or O atoms can be present and the heterocyclic radical can be mono- or disubstituted by A, Hal, OH, OA, CF₃, OCF₃, N(R⁵)₂ or COOR⁵,
- 15 Hal is F, Cl, Br or I,
- 20 Ph is phenyl,
- n is 1 or 2,
- m is 0, 1 or 2,
- o is 1, 2, 3 or 7,
- p is 0 or 1 and
- 25 q is 1, 2 or 3;

in which in Ib-3

- R and R¹ are independently of each other H, Hal, allyl, CH=CH-COOR⁵, CH=CHCON(R⁵)₂ or phenyl, which is unsubstituted or mono-, di- or trisubstituted by A,
- 30

- R^2 and R^3 are independently of each other H, cycloalkyl, $-(CH_2)_o-Het^1$,
 $-(CHA)_p-(CH_2)_o-N(R^5)_2$, $-(CH_2)_p-(CHA)_p-(CH_2)_m-Ar$ or
 $-(CH_2)_o-Z-(CH_2)_q-N(R^5)_2$,
 provided that R^2 and R^3 together are not H,
- 5 R^4 is Ar,
 R^5 is H or A,
 Y is $(CH=CH)_n$,
 Z is phenylene, cyclohexylene, $-NR^5-$, O, $-CH(OH)-$, $-CA_2-$ or
- 10 
- A is unbranched or branched alkyl having 1 to 6 carbon atoms,
 Ar is phenyl, which is mono-, di- or trisubstituted by $O-(CH_2)_p-Ph$,
 naphthyl or Het^2 , or biphenyl, which is unsubstituted or mono-,
 15 di- or trisubstituted by A, OH, OA, CF_3 , Hal, COA, $N(R^5)_2$,
 NO_2 , NR^5-COA or Het^2 ,
- Het^1 is thiophen-2-yl, tetrahydro-furan-2-yl, 1-methyl-octahydro-
 indol-3-yl, benzo[1,3]dioxol-5-yl, piperazin-1-yl, 4-methyl-
 piperazin-1-yl, piperidin-1-yl, piperidin-4-yl, 4-benzyl-piperidin-
 1-yl, 2-methyl-piperidin-1-yl, 1-ethyl-pyrrolidin-2-yl, 1-methyl-
 20 pyrrolidin-2-yl, 2-oxo-pyrrolidin-1-yl, pyridin-2-yl, pyridin-4-yl,
 5-nitro-pyridin-2-yl, imidazol-1-yl, morpholin-4-yl, 5-methoxy-
 1H-indol-2-yl, 5-(3-chlorophenyl)-furan-2-yl, 5-(4-
 fluorophenyl)-thiophen-2-yl, 5-(2-methoxyphenyl)-thiophen-2-
 25 yl, 5-(2-cyanophenyl)-thiophen-2-yl, 5-(2,5-dimethoxyphenyl)-
 thiophen-2-yl, 2-[2,2']bithiophenyl-5-yl, 2-(5-pyridin-4-yl-
 thiophen-2-yl), 5-(1H-indol-5-yl)-thiophen-2-yl, 5-quinolin-8-yl-
 thiophen-2-yl or 5-(benzo[b]thiophen-2-yl)-thiophen-2-yl,
- Het^2 is thiophen-2-yl, pyridin-3-yl, pyridin-2-yl, pyridin-4-yl, indol-5-
 30 yl, quinolin-8-yl, 4,6-dimethoxy-pyrimidin-2-yl or
 benzo[b]thiophen-2-yl,
- Hal is F, Cl, Br or I,

- Ph is phenyl,
 n is 1 or 2,
 m is 0, 1 or 2,
 o is 1, 2, 3 or 7,
 5 p is 0 or 1 and
 q is 1, 2 or 3;
- in Ib-4
 R and R¹ are independently of each other H, Hal, allyl, CH=CH-COOR⁵,
 10 CH=CHCON(R⁵)₂ or 4-methylphenyl,
 R² and R³ are independently of each other H, cyclohexylmethyl,
 -(CH₂)_o-Het¹, -(CHA)_p-(CH₂)_o-N(R⁵)₂, -(CH₂)_p-(CHA)_p-(CH₂)_m-
 Ar or -(CH₂)_o-Z-(CH₂)_q-N(R⁵)₂,
 provided that R² and R³ together are not H,
 15 R⁴ is Ar,
 R⁵ is H or A,
 Y is (CH=CH)_n,
 Z is phenylene, cyclohexylene, -NR⁵-, O, -CH(OH)-, -CA₂- or
 20 ,
 A is unbranched or branched alkyl having 1 to 6 carbon atoms,
 Ar is phenyl, which is mono-, di- or trisubstituted by O-(CH₂)_p-Ph,
 naphthyl or Het², or biphenyl, which is unsubstituted or mono-,
 25 di- or trisubstituted by A, OH, OA, CF₃, Hal, COA, N(R⁵)₂,
 NO₂, NR⁵-COA or Het²,
 Het¹ is 4-methyl-piperazin-1-yl, imidazol-1-yl or morpholin-4-yl,
 Het² is thiophen-2-yl, pyridin-3-yl or benzo[b]thiophen-2-yl,
 Hal is F, Cl, Br or I,
 30 Ph is phenyl,
 n is 1 or 2,
 m is 0, 1 or 2,


o is 1, 2, 3 or 7,
 p is 0 or 1 and
 q is 1, 2 or 3;

5 III. for group Ic

in Ic-1

R and R¹ are independently of each other H, A, OH, OA, Hal, N(R⁵)₂,
 NO₂, CN, C(O)R², CON(R⁵)₂, COOR⁵, allyl, CH=CH-COOR⁵,
 CH=CHCON(R⁵)₂, SO₂A or phenyl, which is unsubstituted or
 10 mono-, di- or trisubstituted by A,

R² and R³ are independently of each other H, A, cycloalkyl, -Het³,
 -(CH₂)_o-OR⁵, -(CH₂)_o-OR⁶, -(CH₂)_o-Het¹, -(CH₂)_o-NR⁵-Het¹,
 -(CHA)_p-(CH₂)_o-N(R⁵)₂, -(CH₂)_p-(CHA)_p-(CH₂)_m-Ar,
 15 -(CH₂)_o-Z-(CH₂)_q-N(R⁵)₂,



provided that R² and R³ together are not H,
 or NR²R³ together form a saturated monocyclic heterocyclic radical having
 5 to 6 ring members, where 1 or 2 N atoms are present and
 the heterocyclic radical can be mono- or disubstituted by OH,
 25 Ar, OAr or arylalkyl,

R⁴ is Het¹,


R⁵ is H or A,

R⁶ is benzo[1,3]dioxol-5-yl,


Q is O or S,

30 Y is (CH=CH)_n,

Z is phenylene, cyclohexylene, -NR⁵-, O, -CH(OH)-, -CA₂- or



- 5 A is unbranched or branched alkyl having 1 to 6 carbon atoms,
Ar is phenyl, naphthyl or biphenyl, which is unsubstituted or mono-, di- or trisubstituted by A, OH, OA, cycloalkyloxy, O-(CH₂)_p-Ph, CF₃, OCF₃, Hal, CN, CHO, COA, COOR⁵, N(R⁵)₂, NR⁵-COA, NO₂, SO₂N(R⁵)₂, mor, SO₂-mor, 5-methyl-3-oxo-2,4-dihydropyrazol-2-yl, naphthyl or Het²,
- 10 Het¹ is a saturated, partially or completely unsaturated mono- or bicyclic heterocyclic radical having 5 to 10 ring members, where 1 or 2 N and/or 1 or 2 S or O atoms can be present and the heterocyclic radical can be mono- or disubstituted by A, Hal, OH, OA, CF₃, OCF₃, N(R⁵)₂, carbonyl oxygen, COOR⁵, Het², benzyl or phenyl which is unsubstituted or mono-, di- or trisubstituted by A, OH, OA, CF₃, OCF₃, Hal, CN, COOR⁵, N(R⁵)₂, NO₂ or SO₂N(R⁵)₂,
- 15 Het² is a unsaturated mono- or bicyclic heterocyclic radical having 5 to 10 ring members, where 1 or 2 N and/or 1 or 2 S or O atoms can be present and the heterocyclic radical can be mono- or disubstituted by A, Hal, OH, OA, CF₃, OCF₃, N(R⁵)₂ or COOR⁵,
- 20 Het³ is a partially or completely unsaturated mono- or bicyclic heterocyclic radical having 5 to 10 ring members, where 1 or 2 N atoms are present and the heterocyclic radical can be mono- or disubstituted by A, Hal, OH, OA, CF₃, OCF₃, N(R⁵)₂, SO₂A or COOR⁵ provided that the heterocyclic radical is not bonded via an N atom,
- 25 Hal is F, Cl, Br or I,
30 mor is morpholin-4-yl,
Ph is phenyl,
n is 1 or 2,

- m is 0, 1, 2, 3, 4, 5 or 6,
 o is 1, 2, 3, 4, 5, 6 or 7,
 p is 0, 1, 2, 3 or 4 and
 q is 1, 2, 3 or 4;
- 5
- in Ic-2
- R and R¹ are independently of each other Hal, allyl, CH=CH-COOR⁵, CH=CHCON(R⁵)₂ or phenyl, which is unsubstituted or mono-, di- or trisubstituted by A,
- 10 R² and R³ are independently of each other H, cycloalkyl, -(CH₂)_o-Het¹, -(CHA)_p-(CH₂)_o-N(R⁵)₂ or -(CH₂)_o-Z-(CH₂)_q-N(R⁵)₂, provided that R² and R³ together are not H,
- R⁴ is Het¹,
- R⁵ is H or A,
- 15 Y is (CH=CH)_n,
- Z is phenylene, cyclohexylene, -NR⁵-, O, -CH(OH)-, -CA₂- or
- 
- 20 A is unbranched or branched alkyl having 1 to 6 carbon atoms,
- Het¹ is a saturated, partially or completely unsaturated mono- or bicyclic heterocyclic radical having 5 to 10 ring members, where 1 or 2 N and/or 1 or 2 S or O atoms can be present and the heterocyclic radical can be mono- or disubstituted by A, Hal, OH, OA, CF₃, OCF₃, N(R⁵)₂, carbonyl oxygen, COOR⁵,
- 25 Het², benzyl or phenyl which is unsubstituted or mono-, di- or trisubstituted by A, OH, OA, CF₃, OCF₃, Hal, CN, COOR⁵, N(R⁵)₂, NO₂ or SO₂N(R⁵)₂,
- Het² is a unsaturated mono- or bicyclic heterocyclic radical having
- 30 5 to 10 ring members, where 1 or 2 N and/or 1 or 2 S or O atoms can be present and the heterocyclic radical can be

mono- or disubstituted by A, Hal, OH, OA, CF₃, OCF₃, N(R⁵)₂ or COOR⁵,

Hal is F, Cl, Br or I,

n is 1 or 2,

5 o is 1, 2, 3 or 7,

p is 0, 1, 2 or 3 and

q is 1, 2 or 3;

in Ic-3

10 R and R¹ are independently of each other H or Hal,


R² and R³ are independently of each other H, cycloalkyl, -(CH₂)_o-Het¹, -(CHA)_p-(CH₂)_o-N(R⁵)₂ or -(CH₂)_o-Z-(CH₂)_q-N(R⁵)₂, provided that R² and R³ together are not H,

R⁴ is Het¹,

15 R⁵ is H or A,

Y is (CH=CH)_n,

Z is phenylene, cyclohexylene, -NR⁵-, O, -CH(OH)-, -CA₂- or




20


A is unbranched or branched alkyl having 1 to 6 carbon atoms,

Het¹ is thiophen-2-yl, tetrahydro-furan-2-yl, 1-methyl-octahydro-indol-3-yl, benzo[1,3]dioxol-5-yl, piperazin-1-yl, 4-methyl-piperazin-1-yl, piperidin-1-yl, piperidin-4-yl, 4-benzyl-piperidin-1-yl, 2-methyl-piperidin-1-yl, 1-ethyl-pyrrolidin-2-yl, 1-methyl-pyrrolidin-2-yl, 2-oxo-pyrrolidin-1-yl, pyridin-2-yl, pyridin-4-yl, 5-nitro-pyridin-2-yl, imidazol-1-yl, morpholin-4-yl, 5-methoxy-1H-indol-2-yl, 5-(3-chlorophenyl)-furan-2-yl, 5-(4-fluorophenyl)-thiophen-2-yl, 5-(2-methoxyphenyl)-thiophen-2-yl, 5-(2-cyanophenyl)-thiophen-2-yl, 5-(2,5-dimethoxyphenyl)-thiophen-2-yl, 2-[2,2']bithiophenyl-5-yl, 2-(5-pyridin-4-yl-

25

30

- thiophen-2-yl, 5-(1H-indol-5-yl)-thiophen-2-yl, 5-quinolin-8-yl-thiophen-2-yl or 5-(benzo[b]thiophen-2-yl)-thiophen-2-yl,
- Hal is F, Cl, Br or I,
- n is 1 or 2,
- 5 o is 1, 2, 3 or 7,
- p is 0, 1, 2 or 3 and
- q is 1, 2 or 3;
- in Ic-4
- 10 R and R¹ are independently of each other H or Hal,
R² and R³ are independently of each other H, cycloalkyl, -(CH₂)_o-Het¹,
-(CHA)_p-(CH₂)_o-N(R⁵)₂ or -(CH₂)_o-Z-(CH₂)_q-N(R⁵)₂, provided
that R² and R³ together are not H,
- R⁴ is Het¹,
- 15 Het¹ in R⁴ is 2-[2,2']bithiophenyl-5-yl or 5-(3-chlorophenyl)-furan-2-yl,
- R⁵ is H or A,
- Y is (CH=CH)_n,
- Z is phenylene, cyclohexylene, -NR⁵-, O, -CH(OH)-, -CA₂- or
- 20 
- A is unbranched or branched alkyl having 1 to 6 carbon atoms,
Het¹ in -(CH₂)_o-Het¹ is thiophen-2-yl, tetrahydro-furan-2-yl, 1-methyl-
octahydro-indol-3-yl, benzo[1,3]dioxol-5-yl, piperazin-1-yl, 4-
25 methyl-piperazin-1-yl, piperidin-1-yl, piperidin-4-yl, 4-benzyl-
piperidin-1-yl, 2-methyl-piperidin-1-yl, 1-ethyl-pyrrolidin-2-yl, 1-
methyl-pyrrolidin-2-yl, 2-oxo-pyrrolidin-1-yl, pyridin-2-yl,
pyridin-4-yl, 5-nitro-pyridin-2-yl, imidazol-1-yl, morpholin-4-yl,
5-methoxy-1H-indol-2-yl, 5-(3-chlorophenyl)-furan-2-yl, 5-(4-
30 fluorophenyl)-thiophen-2-yl, 5-(2-methoxyphenyl)-thiophen-2-
yl, 5-(2-cyanophenyl)-thiophen-2-yl, 5-(2,5-dimethoxyphenyl)-
thiophen-2-yl, 2-[2,2']bithiophenyl-5-yl, 2-(5-pyridin-4-yl-

- thiophen-2-yl, 5-(1H-indol-5-yl)-thiophen-2-yl, 5-quinolin-8-yl-thiophen-2-yl or 5-(benzo[b]thiophen-2-yl)-thiophen-2-yl,
- Hal is F, Cl, Br or I,
- n is 1 or 2,
- 5 o is 1, 2, 3 or 7,
- p is 0, 1, 2 or 3 and
- q is 1, 2 or 3;
- in Ic-5
- 10 R and R¹ are independently of each other H or Hal,
R² and R³ are independently of each other H, cycloalkyl, -(CH₂)_o-Het¹,
-(CHA)_p-(CH₂)_o-N(R⁵)₂ or -(CH₂)_o-Z-(CH₂)_q-N(R⁵)₂, provided
that R² and R³ together are not H,
- R⁴ is Het¹,
- 15 Het¹ in R⁴ is 2-[2,2']bithiophenyl-5-yl,
- R⁵ is H or A,
- Y is (CH=CH)_n,
- Z is phenylene, cyclohexylene, -NR⁵-, O, -CH(OH)-, -CA₂- or
- 20 
- A is unbranched or branched alkyl having 1 to 6 carbon atoms,
Het¹ in -(CH₂)_o-Het¹ is piperidin-4-yl or pyridin-4-yl,
- Hal is F, Cl, Br or I,
- 25 n is 1,
- o is 1, 2, 3 or 7,
- p is 0, 1, 2 or 3 and
- q is 1, 2 or 3.

30 The invention relates additionally to novel substituted 4-amino-quinazolines of the formula I according to group Ia and their pharmaceutically tolerable salts and solvates.

The invention relates additionally to novel substituted 4-amino-quinazolines of the formula I according to group Ib and their pharmaceutically tolerable salts and solvates.

5

The invention relates additionally to novel substituted 4-amino-quinazolines of the formula I according to group Ic and their pharmaceutically tolerable salts and solvates.

10

The invention relates further to novel substituted 4-amino-quinazolines of the formula I according to groups Ia-Ic and their pharmaceutically tolerable salts and solvates as a medicament.

15

The invention relates to novel substituted 4-amino-quinazolines of the formula I according to groups Ia-Ic and their pharmaceutically tolerable salts and solvates as a glycoprotein IbIX antagonist.

The invention relates further to novel special compounds of formula I selected from the group

20

a) (7-chloro-2-styryl-quinazolin-4-yl)-(3-imidazol-1-yl-propyl)-amine,
b) N¹-(7-chloro-2-styryl-quinazolin-4-yl)-N,N-diethyl-ethane-1,2-diamine,
c) N¹-(7-chloro-2-styryl-quinazolin-4-yl)-N,N-diethyl-propane-1,3-diamine,

25

d) (7-chloro-2-styryl-quinazolin-4-yl)-(3-morpholin-4-yl-propyl)-amine,
e) 1-[3-(7-chloro-2-styryl-quinazolin-4-ylamino)-propyl]-pyrrolidin-2-one,
f) [2-(4-amino-phenyl)-ethyl]-(7-chloro-2-styryl-quinazolin-4-yl)-amine,
g) N⁴-{2-[2-(4-bromo-phenyl)-vinyl]-7-chloro-quinazolin-4-yl}-N¹,N¹-diethyl-pentane-1,4-diamine and

30

h) N⁴-[7-chloro-2-(4-phenyl-buta-1,3-dienyl)-quinazolin-4-yl]-N¹,N¹-diethyl-pentane-1,4-diamine and their pharmaceutically tolerable salts and solvates.

The invention relates further to the novel substituted 4-amino-quinazolines a) to h) of the formula I and their pharmaceutically tolerable salts and solvates as a medicament.

5 The invention relates to the novel substituted 4-amino-quinazolines a) to h) of the formula I and their pharmaceutically tolerable salts and solvates as a glycoprotein IbIX antagonist.

10 The compounds of the formula I and also the starting substances for their preparation are otherwise prepared by methods known per se, such as are described in the literature (e.g. in the standard works such as Houben-Weyl, Methoden der organischen Chemie [Methods of Organic Chemistry], Georg-Thieme-Verlag, Stuttgart), namely under reaction conditions which are known and suitable for the reactions mentioned. In this case, use can
15 also be made of variants which are known per se, but not mentioned here in greater detail.

The starting substances, if desired, can also be formed in situ such that they are not isolated from the reaction mixture, but immediately reacted further to give the compounds of the formula I.

20

The compounds of the formula I according to claims 1 to 4 can be obtained by liberating them from their functional derivatives by solvolysis, in particular hydrolysis or by hydrogenolysis.

25 Preferred starting substances for the solvolysis or hydrogenolysis are those which otherwise correspond to the formula I according to claims 1 to 4, but instead of one or more free amino and/or hydroxyl groups contain corresponding protected amino and/or hydroxyl groups, in particular those which instead of an H-N- group carry an R'-N'- group, in which R' is an
30 amino protective group and/or those which instead of the H atom of a hydroxyl group carry a hydroxyl protective group, e.g. those which

correspond to the formula I, but instead of a group -COOH carry a group -COOR", in which R" is a hydroxyl protective group.

5 A number of - identical or different - protected amino and/or hydroxyl groups can also be present in the molecule of the starting substance. If the protective groups present are different from one another, in many cases they can be removed selectively (lit.: T.W. Greene, P.G.M. Wuts, *Protective Groups in Organic Chemistry*, 2nd ed., Wiley, New York 1991 or P.J. Kocienski, *Protecting Groups*, 1st ed., Georg Thieme Verlag, Stuttgart -
10 New-York, 1994).

The expression "amino protective group" is generally known and relates to groups which are suitable for protecting (for blocking) an amino group against chemical reactions, but which are easily removable after the
15 desired chemical reaction has been carried out at other positions in the molecule. Typical groups of this type are, in particular, unsubstituted or substituted acyl, aryl, aralkoxymethyl or aralkyl groups. Since the amino protective groups are removed after the desired reaction (or reaction
20 sequence), their nature and size is otherwise not critical; however, those having 1-20, in particular 1-8, C atoms are preferred. The expression "acyl group" is to be interpreted in the widest sense in connection with the present process. It includes acyl groups derived from aliphatic, araliphatic, aromatic or heterocyclic carboxylic acids or sulfonic acids and, in particular, alkoxy-carbonyl groups, aryloxy-carbonyl groups and especially
25 aralkoxy-carbonyl groups. Examples of acyl groups of this type are alkanoyl such as acetyl, propionyl, butyryl; aralkanoyl such as phenylacetyl; aroyl such as benzoyl or toluyl; aryloxyalkanoyl such as POA; alkoxy-carbonyl such as methoxycarbonyl, ethoxycarbonyl, 2,2,2-trichloroethoxycarbonyl, BOC, 2-iodoethoxycarbonyl; aralkyloxy-carbonyl such as CBZ
30 ("carbobenzoxy"), 4-methoxybenzyloxy-carbonyl (MOZ), 4-Nitro-benzyloxy-carbonyl oder 9-fluorenylmethoxycarbonyl (Fmoc); 2-(phenylsulfonyl)ethoxycarbonyl; trimethylsilylethoxycarbonyl (Teoc) or

arylsulfonyl such as 4-methoxy-2,3,6-trimethylphenyl-sulfonyl (Mtr). Preferred amino protective groups are BOC, furthermore CBZ, Fmoc, benzyl and acetyl; particularly preferred Fmoc.

- 5 The expression "hydroxyl protective group" is also generally known and relates to groups which are suitable for protecting a hydroxyl group against chemical reactions, but which are easily removable after the desired chemical reaction has been carried out at other positions in the molecule.
- 10 Typical groups of this type are the above mentioned unsubstituted or substituted aryl, aralkyl, aroyl or acyl groups, furthermore also alkyl groups, alkyl-, aryl- or aralkylsilyl groups or O,O- or O,S-acetals. The nature and size of the hydroxyl protective groups is not critical, since they are removed again after the desired chemical reaction or reaction sequence; groups
- 15 having 1-20, in particular 1-10 C atoms, are preferred. Examples of hydroxyl protective groups are, inter alia, benzyl, 4-methoxybenzyl or 2,4-dimethoxybenzyl, aroyl groups such as benzoyl or p-nitrobenzoyl, acyl groups such as acetyl or pivaloyl, p-toluolsulfonyl, alkyl groups such as methyl or tert-butyl, but also allyl, alkylsilyl groups such as trimethylsilyl
- 20 (TMS), triisopropylsilyl (TIPS), tert-butyldimethylsilyl (TBS) or triethylsilyl, trimethylsilylethyl, aralkylsilyl groups such as tert-butyldiphenylsilyl (TBDPS), cyclic acetals such as isopropylidene-, cyclopentylidene-, cyclohexylidene-, benzylidene-, p-methoxybenzylidene- or o,p-dimethoxybenzylideneacetal, acyclic acetals such as tetrahydropyranyl
- 25 (Thp), methoxymethyl (MOM), methoxyethoxymethyl (MEM), benzyloxymethyl (BOM) or methylthiomethyl (MTM). Acetyl, benzyl, tert-butyl or TBS being particularly preferred.

30 The liberation of the compounds of the formula I from their functional derivatives depending on the protective group used is known in the present literature such as T.W. Greene, P.G.M. Wuts, *Protective Groups in Organic Chemistry*, 2nd ed., Wiley, New York 1991, P.J. Kocienski, *Protecting*

Groups, 1st ed., Georg Thieme Verlag, Stuttgart - New-York, 1994. In this case, use can also be made of variants which are known per se, but not mentioned here in greater detail.

5 The groups BOC and O-tert-butyl can preferably be removed, for example, using TFA in dichloromethane or using approximately 3 to 5N HCl in dioxane at 15-30°C, the Fmoc group using an approximately 5 to 50% solution of dimethylamine, diethylamine or piperidine in DMF at 15-30°C.

10 Preferred starting substances for the solvolysis or hydrogenolysis includes also those which otherwise correspond to the formula I, but are attached to a solid phase. The liberation of the compounds of the formula I from the solid phase is known in the present literature such as Novabiochem - The Combinatorial Chemistry Catalog, March 99 and cited literature.

15 The solid phase with a carbonate moiety as terminal functional group can preferably be removed, for example, using TFA (50%) in dichloromethane.

20 The quinazolines of formula I according to claims 1 to 4 can also preferably be prepared, using either solution or solid-phase techniques.

The term solid phase indicates a resin for solid-phase chemistry, especially for combinatorial chemistry, i.e. by robot- and computer-assisted syntheses, and subjected to mass screening as indicated in US 5,463,564; 25 M. A. Gallop et al., J. Med. Chem. 1994, 37, 1233-1251 and 1385-1401 and M.J. Sofia, Drug Discovery Today 1996, 1, 27-34). The polymeric material of the solid phase is generally chosen from the group consisting of cross-linked polystyrene, cross-linked polyacrylamide or other resins, natural polymers or silicagels.

30 The group of cross-linked polystyrene, cross-linked polyacrylamide or other resins includes e.g. polyacrylamide, polymethacrylamide,

polyhydroxyethylmethacrylate, polyamide, polystyrene, (meth)acrylate copolymers, for instance from (meth)acrylic acid, esters of (meth)acrylic acid and/or 2-methylene-succinic acid, but-2-enoic acid or maleic acid, polyurethanes or other copolymers.

5

Suitable terminal functional groups or linkers on the surface of the resin have to be chosen to attach the compounds to the resin. There exists a variety of commercially available resins, e.g. in Novabiochem - The Combinatorial Chemistry Catalog, March 99. Examples for suitable resins are carbonate resins with a modified carbonate group as terminal functional group like p-nitrophenylcarbonate resin, halogenated resins like Merrifield resin (chloromethylpolystyrene) or carboxy resins like carboxy polystyrene resin or NovaSyn[®] TG Carboxy Resin. p-Nitrophenylcarbonate resin is particularly preferred. These and other types of resins well known in the art can be used in the subject invention.

10

15

The quinazolines of formula I according to claims 1 to 4 can therefore preferably be prepared by combining and reacting a 2-methyl-3H-quinazolin-4-one of formula II with an aldehyde of formula III, chlorinating the given formula IV and reacting the given formula V with an amine of formula VI.

20

The quinazolines of formula I according to claims 1 to 4 can furthermore be prepared by chlorinating a 2-methyl-3H-quinazolin-4-one of formula II, reacting the given formula VII with the amine of formula VI and reacting the given formula VIII with an aldehyde of formula III.

25

As a rule, the starting compounds of the formulae II, III and VI are known or commercially available.

The unknown compounds, however, can be prepared by methods known per se.

30

The 2-methyl-3H-quinazolin-4-ones of formula II in which R and R¹ have a meaning indicated in claims 1 to 4 can be prepared by reacting a substituted anthranilic acid with acetic anhydride and reacting the given 2-methyl-benzoxazin-4-one with ammonium acetate.

5

The aldehydes of formula III, as a rule, are also commercially available. Furthermore, syntheses for the preparation of aldehydes of formula III, such as, for example, the oxidation of an alcohol, can be used.

10

The amines of formula VI in which R² or R³ have a meaning indicated in claims 1 to 4, as a rule, are also commercially available and can be attached to the suitable resin or to a compound of formula V or VII by coupling procedures well known in the art and as described in the ensuing Examples. Furthermore, syntheses for the preparation of amines of formula

15

III, such as, for example, the Gabriel synthesis, can be used.

For the preparation of compounds of the formula I in which R⁴ is unsubstituted or substituted biphenyl, arylsubstituted furanyl or 5-[2,2']bithiophenyl, an appropriate compound of the formula I in which R⁴ is

20

phenyl chloride, phenyl bromide, phenyl iodide, furanyl chloride, furanyl bromide, furanyl iodide, thiophenyl chloride, thiophenyl bromide or thiophenyl iodide can be reacted with the appropriate boronic acid

25

derivatives in a Suzuki type coupling reaction. This reaction is expediently carried out under Palladium catalysis with different phosphines as coordination ligands, e.g. Pd(P(Ph)₃)₂, Pd(II)Cl₂dppf, PdOAc₂ + P(R*)₃ (R* = phenyl, cyclohexyl, tert-butyl) etc. in the presence of a base such as potassium carbonate, cesium carbonate, DBU, NaOH, in an inert solvent or solvent mixture, e.g. DMF or 1,4-dioxane at temperatures between 0° and 150°, preferably between 60° and 120°. Depending on the conditions used,

30

the reaction time is between a few minutes and a number of days. The boronic acid derivatives can be prepared by conventional methods or are commercially available. The reactions can be carried out in analogy to the

methods indicated in Suzuki et al., J. Am. Chem. Soc. 1989, 111, 314ff., Suzuki et al., Chem. Rev. 1995, 95, 2457ff and G.C. Fu et al. Angew. Chem 1998, 110, 3586.

5 The Suzuki type coupling reaction can be furthermore used to convert radicals R and R¹ into other radicals R and R¹, for e.g. to convert a halogen substituted quinazolines to a quinazoline substituted by substituted or unsubstituted phenyl.

10 For the preparation of compounds of the formula I in which R or R¹ is allyl, an appropriate compound of the formula I in which R⁴ is quinazoline chloride, quinazoline bromide or quinazoline iodide can be reacted with allyltributyltin in a Stille type coupling reaction. This reaction is expediently carried out under Palladium catalysis with different phosphines as coordination ligands, e.g. Pd(P(Ph)₃)₂, Pd(II)Cl₂dppf, PdOAc₂ + P(R*)₃ (R* = phenyl, cyclohexyl, tert-butyl) etc. in an inert solvent or solvent mixture, 15 e.g. DMF or 1,4-dioxane at temperatures between 0° and 150°, preferably between 60° and 120°. Depending on the conditions used, the reaction time is between a few minutes and a number of days.

20 For the preparation of compounds of the formula I in which R or R¹ is CH=CH-COOR⁵ or CH=CH-CON(R⁵)₂, an appropriate compound of the formula I in which R⁴ is quinazoline chloride, quinazoline bromide or quinazoline iodide can be reacted with substituted acrylate in a Heck type coupling reaction. This reaction is expediently carried out under Palladium 25 catalysis with different phosphines as coordination ligands, e.g. Pd(P(Ph)₃)₂, Pd(II)Cl₂dppf, PdOAc₂ + P(R*)₃ (R* = phenyl, cyclohexyl, tert-butyl) etc. in the presence of a base such as triethyl amine or a catalyst tetrabutylammonium iodide, in an inert solvent or solvent mixture, e.g. DMF or 1,4-dioxane at temperatures between 0° and 150°, preferably between 30 60° and 120°. Depending on the conditions used, the reaction time is between a few minutes and a number of days.

A base of the formula I can be converted into the associated acid addition salt using an acid, for example by reaction of equivalent amounts of the base and of the acid in an inert solvent such as ethanol and subsequent evaporation. Acids which give physiologically acceptable salts are particularly suitable for this reaction. Thus inorganic acids can be used, e.g. sulfuric acid, nitric acid, hydrohalic acids such as hydrochloric acid or hydrobromic acid, phosphoric acids such as orthophosphoric acid, sulfamic acid, furthermore organic acids, in particular aliphatic, alicyclic, araliphatic, aromatic or heterocyclic mono- or polybasic carboxylic, sulfonic or sulfuric acids, e.g. formic acid, acetic acid, propionic acid, pivalic acid, diethylacetic acid, malonic acid, succinic acid, pimelic acid, fumaric acid, maleic acid, lactic acid, tartaric acid, malic acid, citric acid, gluconic acid, ascorbic acid, nicotinic acid, isonicotinic acid, methane- or ethanesulfonic acid, p-toluenesulfonic acid, naphthalenemono- and disulfonic acids or laurylsulfuric acid. Salts with physiologically unacceptable acids, e.g. picrates, can be used for the isolation and/or purification of the compounds of the formula I.

On the other hand, compounds of the formula I with bases (e.g. sodium or potassium hydroxide or carbonate) can be converted into the corresponding metal salts, in particular alkali metal or alkaline earth metal salts, or into the corresponding ammonium salts.

The invention furthermore relates to pharmaceutical preparations comprising at least one compound of the formula I according to Claims 1 to 5 and/or one of its physiologically acceptable salts, which are prepared, in particular, in a non-chemical way. In this case, the compounds of the formula I according to the invention can be brought into a suitable dose form together with at least one solid, liquid and/or semi-liquid excipient or auxiliary and, if appropriate, in combination with one or more other active compounds.

These preparations can be used as medicaments in human or veterinary medicine. Possible excipients are organic or inorganic substances which are suitable for enteral (e.g. oral) or parenteral administration or topical application and do not react with the novel compounds, for example water, vegetable oils, benzyl alcohols, alkylene glycols, polyethylene glycols, glyceryl triacetate, gelatin, carbohydrates such as lactose or starch, magnesium stearate, talc and petroleum jelly. Tablets, pills, coated tablets, capsules, powders, granules, syrups, juices or drops are used, in particular, for oral administration, suppositories are used for rectal administration, solutions, preferably oily or aqueous solutions, furthermore suspensions, emulsions or implants, are used for parenteral administration, and ointments, creams or powders are used for topical application. The novel compounds can also be lyophilized and the lyophilizates obtained used, for example, for the production of injection preparations. The preparations indicated can be sterilized and/or can contain auxiliaries such as lubricants, preservatives, stabilizers and/or wetting agents, emulsifiers, salts for affecting the osmotic pressure, buffer substances, colorants, flavorings and/or one or more other active compounds, e.g. one or more vitamins.

The compounds of the formula I and their physiologically acceptable salts according to claims 1 to 5 act as adhesion receptor antagonists, in particular glycoprotein IbIX antagonists, and can be employed for the prophylaxis and/or therapy of thrombotic disorders and sequelae deriving therefrom. The disorders are acute coronary syndromes, angina pectoris, myocardial infarct, peripheral circulatory disorders, stroke, transient ischaemic attacks, arteriosclerosis and reocclusion/restenosis after angioplasty/stent implantation.

In this case, the substances according to the invention are as a rule administered in the dose of the glycoprotein IIbIIIa antagonist ReoPro® of preferably between approximately 1 and 500 mg, in particular between 5

and 100 mg, per dose unit. The daily dose is preferably between approximately 0.02 and 10 mg/kg of body weight. The specific dose for each patient depends, however, on all sorts of factors, for example on the efficacy of the specific compound employed, on the age, body weight,
5 general state of health and sex, on the diet, on the time and route of administration, and on the excretion rate, pharmaceutical combination and severity of the particular disorder to which the therapy applies. Oral administration is preferred.

10 Above and below, all temperatures are indicated in °C. In the following examples, "customary working-up" for solution reactions means: if necessary, water is added, if necessary, depending on the constitution of the final product, the mixture is adjusted to pHs between 2 and 10 and
15 extracted with ethyl acetate or dichloromethane, the organic phase is separated off, dried over sodium sulfate and evaporated, and the residue is purified by chromatography on silica gel and/or by crystallization.

"Customary working-up" for solid-phase reactions means: the crude reaction is filtered and washed with DMF twice, then successively with
20 methanol and methylene chloride three times, and finally once with methyl tert-butyl ether. The resin is then dried in vacuo.

Mass spectrometry (MS) apparatuses OMIT and Finnigan LCQ. (M+H)⁺ values or M⁺ values are determined.

25

EXAMPLES

Example 1:

(3-Aminomethyl-cyclohexylmethyl)-[2-(2-naphthalen-1-yl-vinyl)-quinazolin-4-yl]-amine
30

1. Anthranilic acid (0,29 mole) is added to 170 ml acetic anhydride. The solution is heated to 140° for 3 h. After cooling to room temperature (rt), the white solid is collected by filtration and washed with diethyl ether. Air drying give 2-methylbenzoxazin-4-one.
5
2. 2-Methylbenzoxazin-4-one (0,24 mole) and ammonium acetate (0,30 mole) is given in 50 ml of N,N-dimethylacetamide and then heated to 160° under a nitrogen blanket for 2 h. After cooling to rt, the white solid is collected by filtration and washed with diethyl ether. Air drying give 2-methyl-quinazolin-4-one.
10
3. 2-Methylquinazolin-4-one (50 mmol) and naphthalene-1-carbaldehyde (54 mmol) are suspended in 80 ml acetic acid. The mixture is heated to 100° for 24 h. After cooling to rt, the product crystallize out. After filtration, washing with ethyl acetate and air drying, 2-(2-naphthalen-1-yl-vinyl)-3H-quinazolin-4-one is given.
15
4. 2-(2-Naphthalen-1-yl-vinyl)-3H-quinazolin-4-one (11 mmol), phosphorus oxychloride (20 ml) and N,N-diethylaniline (1,0 ml) is added to a round-bottomed flask. The mixture is heated at 110° for 12 h. After cooling to rt, the reaction is quenched with ice-water and the crude product is collected by suction filtration. The crude solid is dissolved in ethyl acetate. Customary working up afforded 4-chloro-2-(2-naphthalen-1-yl-vinyl)-3,4-dihydro-quinazoline.
20
5. 4-Chloro-2-(2-naphthalen-1-yl-vinyl)-3,4-dihydro-quinazoline (0,08 mmol), C-(3-aminomethyl-cyclohexyl)-methylamine (0,24 mmol) and 2 ml ethyl alcohol are placed in an 8 ml glass vial sealed with a teflon-lined screw cap. The mixture is heated at 80° for 3 hrs. After cooling to rt, the ethyl alcohol is removed. Ethyl acetate (3ml) and water (3ml) are added to the vial. After agitation, the water was removed. The procedure is repeated with water and aq. NaCl (sat.). Purification with silica gel afforded (3-
25
30

aminomethyl-cyclohexylmethyl)-[2-(2-naphthalen-1-yl-vinyl)-quinazolin-4-yl]-
amine;

MS calc.: 422.6 ; found: 423.5.

5 Example 2:

Analogously to example 1, 2-methylquinazolin-4-one
is reacted

with 4-pentyloxy-benzaldehyde, chlorinated and reacted with C-(3-
aminomethyl-cyclohexyl)-methylamine to obtain

10 (3-aminomethyl-cyclohexylmethyl)-{2-[2-(4-pentyloxy-phenyl)-vinyl]-
quinazolin-4-yl}-amine;


MS calc.: 458.6 ; found: 459.5;

with 3-phenyl-propenal, chlorinated and reacted with C-(3-aminomethyl-
15 cyclohexyl)-methylamine to obtain

(3-aminomethyl-cyclohexylmethyl)-[2-(4-phenyl-buta-1,3-dienyl)-
quinazolin-4-yl]-amine;

MS calc.: 398.6 ; found: 399.4;

20



25

with 3-fluoro-benzaldehyde, chlorinated and reacted with C-(3-
aminomethyl-cyclohexyl)-methylamine to obtain

(3-aminomethyl-cyclohexylmethyl)-{2-[2-(3-fluoro-phenyl)-vinyl]-
quinazolin-4-yl}-amine;

30 MS calc.: 390.5 ; found: 391.4;

with 2-fluoro-benzaldehyde, chlorinated and reacted with C-(3-aminomethyl-cyclohexyl)-methylamine to obtain

(3-aminomethyl-cyclohexylmethyl)-{2-[2-(2-fluoro-phenyl)-vinyl]-quinazolin-4-yl}-amine;

5 MS calc.: 390.5 ; found: 391.3;

with 4-chloro-benzaldehyde, chlorinated and reacted with C-(3-aminomethyl-cyclohexyl)-methylamine to obtain

(3-aminomethyl-cyclohexylmethyl)-{2-[2-(4-chloro-phenyl)-vinyl]-quinazolin-4-yl}-amine;

10 MS calc.: 407.0 ; found: 410.3;

with 3-chloro-benzaldehyde, chlorinated and reacted with C-(3-aminomethyl-cyclohexyl)-methylamine to obtain

(3-aminomethyl-cyclohexylmethyl)-{2-[2-(3-chloro-phenyl)-vinyl]-quinazolin-4-yl}-amine;

15 MS calc.: 407.0 ; found: 407.3;

with 4-amino-benzaldehyde, chlorinated and reacted with C-(3-aminomethyl-cyclohexyl)-methylamine to obtain

(3-aminomethyl-cyclohexylmethyl)-{2-[2-(4-amino-phenyl)-vinyl]-quinazolin-4-yl}-amine;

20 MS calc.: 387.5 ; found: 388.4;

with 4-methoxy-benzaldehyde, chlorinated and reacted with C-(3-aminomethyl-cyclohexyl)-methylamine to obtain

(3-aminomethyl-cyclohexylmethyl)-{2-[2-(4-methoxy-phenyl)-vinyl]-quinazolin-4-yl}-amine;

25 MS calc.: 402.5 ; found: 403.3;

with 3-methoxy-benzaldehyde, chlorinated and reacted with C-(3-aminomethyl-cyclohexyl)-methylamine to obtain

30

(3-aminomethyl-cyclohexylmethyl)-{2-[2-(3-methoxy-phenyl)-vinyl]-quinazolin-4-yl}-amine;

MS calc.: 402.5 ; found: 403.3;

5 with 4-methyl-benzaldehyde, chlorinated and reacted with C-(3-aminomethyl-cyclohexyl)-methylamine to obtain

(3-aminomethyl-cyclohexylmethyl)-{2-[2-(4-methyl-phenyl)-vinyl]-quinazolin-4-yl}-amine;

MS calc.: 386.5 ; found: 387.3;

10

with 3-methyl-benzaldehyde, chlorinated and reacted with C-(3-aminomethyl-cyclohexyl)-methylamine to obtain

(3-aminomethyl-cyclohexylmethyl)-{2-[2-(3-methyl-phenyl)-vinyl]-quinazolin-4-yl}-amine;

15 MS calc.: 386.5 ; found: 387.3;

with 2-methyl-benzaldehyde, chlorinated and reacted with C-(3-aminomethyl-cyclohexyl)-methylamine to obtain

(3-aminomethyl-cyclohexylmethyl)-{2-[2-(2-methyl-phenyl)-vinyl]-quinazolin-4-yl}-amine;

20 MS calc.: 386.5 ; found: 387.3;

with 2-nitro-benzaldehyde, chlorinated and reacted with C-(3-aminomethyl-cyclohexyl)-methylamine to obtain

25 (3-aminomethyl-cyclohexylmethyl)-{2-[2-(2-nitro-phenyl)-vinyl]-quinazolin-4-yl}-amine;

MS calc.: 417.5 ; found: 418.3;

with 3-nitro-benzaldehyde, chlorinated and reacted with C-(3-aminomethyl-cyclohexyl)-methylamine to obtain

30 (3-aminomethyl-cyclohexylmethyl)-{2-[2-(3-nitro-phenyl)-vinyl]-quinazolin-4-yl}-amine;

MS calc.: 417.5 ; found: 418.3;

with 4-nitro-benzaldehyde, chlorinated and reacted with C-(3-aminomethyl-cyclohexyl)-methylamine to obtain

5 (3-aminomethyl-cyclohexylmethyl)-{2-[2-(4-nitro-phenyl)-vinyl]-quinazolin-4-yl}-amine;

MS calc.: 417.5 ; found: 418.3;

with 3,4,5-trimethoxy-benzaldehyde, chlorinated and reacted with C-(3-aminomethyl-cyclohexyl)-methylamine to obtain

10 (3-aminomethyl-cyclohexylmethyl)-{2-[2-(3,4,5-trimethoxy-phenyl)-vinyl]-quinazolin-4-yl}-amine;

with 4-carboxy-benzaldehyde, chlorinated and reacted with C-(3-aminomethyl-cyclohexyl)-methylamine to obtain

15 (3-aminomethyl-cyclohexylmethyl)-{2-[2-(4-carboxy-phenyl)-vinyl]-quinazolin-4-yl}-amine;

MS calc.: 416.5 ; found: 417.3 and

with 3-carboxy-benzaldehyde, chlorinated and reacted with C-(3-aminomethyl-cyclohexyl)-methylamine to obtain

20 (3-aminomethyl-cyclohexylmethyl)-{2-[2-(3-carboxy-phenyl)-vinyl]-quinazolin-4-yl}-amine;

MS calc.: 416.5 ; found: 417.3.

25

Analogously to example 1, 7-chloro-2-methylquinazolin-4-one is reacted with 4-bromo-benzaldehyde, chlorinated and reacted

with C-(3-aminomethyl-cyclohexyl)-methylamine to obtain

30 (3-aminomethyl-cyclohexylmethyl)-{2-[2-(4-bromo-phenyl)-vinyl]-7-chloro-quinazolin-4-yl}-amine;

MS calc.: 485.9 ; found: 487.2.

Example 3:

5 Analogously to example 1, 7-chloro-2-methylquinazolin-4-one
is reacted with 3-phenoxy-benzaldehyde, chlorinated and reacted

with N¹,N¹-diethyl-pentane-1,4-diamine to obtain

N⁴-{7-chloro-2-[2-(3-phenoxy-phenyl)-vinyl]-quinazolin-4-yl}-N¹,N¹-
diethyl-pentane-1,4-diamine;

10

with C-(3-aminomethyl-cyclohexyl)-methylamine to obtain

(3-aminomethyl-cyclohexylmethyl)-{7-chloro-2-[2-(3-phenoxy-phenyl)-
vinyl]-quinazolin-4-yl}-amine;

MS calc.: 499.1 ; found: 499.5;

15

with 3-aminomethyl-benzylamine to obtain

(3-aminomethyl-benzyl)-{7-chloro-2-[2-(3-phenoxy-phenyl)-vinyl]-
quinazolin-4-yl}-amine;

MS calc.: 493.0 ; found: 493.4 and

20

with heptane-1,7-diamine

N¹-{7-chloro-2-[2-(3-phenoxy-phenyl)-vinyl]-quinazolin-4-yl}-heptane-1,7-
diamine;

MS calc.: 487.0 ; found: 487.5.

25

Example 4:

Analogously to example 1, 2-methylquinazolin-4-one
is reacted with 4-benzyloxy-benzaldehyde, chlorinated and reacted

30 with C-(3-aminomethyl-cyclohexyl)-methylamine to obtain

(3-aminomethyl-cyclohexylmethyl)-{2-[2-(4-benzyloxy-phenyl)-vinyl]-
quinazolin-4-yl}-amine;

MS calc.: 478.6 ; found: 479.4;

with 3-(3-amino-propoxy)-propylamine to obtain

5 [3-(3-amino-propoxy)-propyl]-{2-[2-(4-benzyloxy-phenyl)-vinyl]-
quinazolin-4-yl}-amine;

MS calc.: 468.6 ; found: 469.3;

with 2,2-dimethyl-propane-1,3-diamine to obtain


10 N¹-{2-[2-(4-benzyloxy-phenyl)-vinyl]-quinazolin-4-yl}-2,2-dimethyl-
propane-1,3-diamine;

MS calc.: 438.6 ; found: 439.3;

with N¹-(2-diethylamino-ethyl)-ethane-1,2-diamine to obtain

15 N-{2-[2-(4-benzyloxy-phenyl)-vinyl]-quinazolin-4-yl}-N'-(2-diethylamino-
ethyl)-ethane-1,2-diamine;

MS calc.: 495.7 ; found: 496.3;



with heptane-1,7-diamine to obtain

25 N¹-{2-[2-(4-benzyloxy-phenyl)-vinyl]-quinazolin-4-yl}-heptane-1,7-
diamine;

MS calc.: 466.6 ; found: 467.3;

with 4-(2-amino-ethyl)-phenylamine to obtain


30 [2-(4-amino-phenyl)-ethyl]-{2-[2-(4-benzyloxy-phenyl)-vinyl]-quinazolin-4-
yl}-amine;

MS calc.: 472.6 ; found: 473.2;

with 3-morpholin-4-yl-propylamine to obtain

{2-[2-(4-benzyloxy-phenyl)-vinyl]-quinazolin-4-yl}-(3-morpholin-4-yl-propyl)-amine;

5 MS calc.: 480.6 ; found: 481.3;



with N¹,N¹-diethyl-propane-1,3-diamine to obtain

15 N¹-{2-[2-(4-benzyloxy-phenyl)-vinyl]-quinazolin-4-yl}-N,N-diethyl-propane-1,3-diamine;

MS calc.: 466.6 ; found: 467.3;

with N¹,N¹-diethyl-ethane-1,2-diamine to obtain


20 N¹-{2-[2-(4-benzyloxy-phenyl)-vinyl]-quinazolin-4-yl}-N,N-diethyl-ethane-1,2-diamine;

MS calc.: 452.6 ; found: 453.2;

with 3-imidazol-1-yl-propylamine to obtain

25 {2-[2-(4-benzyloxy-phenyl)-vinyl]-quinazolin-4-yl}-(3-imidazol-1-yl-propyl)-amine;

MS calc.: 461.6 ; found: 462.2;



Analogously to example 1, 7-chloro-2-methylquinazolin-4-one is reacted with 4-benzyloxy-benzaldehyde, chlorinated and reacted

5 with N¹,N¹-diethyl-pentane-1,4-diamine to obtain

N⁴-{2-[2-(4-benzyloxy-phenyl)-vinyl]-7-chloro-quinazolin-4-yl}- N¹,N¹-diethyl-pentane-1,4-diamine;

with C-(3-aminomethyl-cyclohexyl)-methylamine to obtain

10 (3-aminomethyl-cyclohexylmethyl)-{2-[2-(4-benzyloxy-phenyl)-vinyl]-7-chloro-quinazolin-4-yl}-amine;

MS calc.: 513.1 ; found: 513.3.

Analogously to example 1, 6-iodo-2-methylquinazolin-4-one

15 is reacted with 4-benzyloxy-benzaldehyde, chlorinated and reacted

with N¹,N¹-diethyl-pentane-1,4-diamine to obtain

N⁴-{2-[2-(4-benzyloxy-phenyl)-vinyl]-6-iodo-quinazolin-4-yl}- N¹,N¹-diethyl-pentane-1,4-diamine;

20 MS calc.: 620.6 ; found: 621.1;

with C-(3-aminomethyl-cyclohexyl)-methylamine to obtain

(3-aminomethyl-cyclohexylmethyl)-{2-[2-(4-benzyloxy-phenyl)-vinyl]-6-iodo-quinazolin-4-yl}-amine;

25 MS calc.: 604.5 ; found: 605.2.

Analogously to example 1, 6-bromo-2-methylquinazolin-4-one

is reacted with 4-benzyloxy-benzaldehyde, chlorinated and reacted

30 with N¹,N¹-diethyl-pentane-1,4-diamine to obtain

N⁴-{2-[2-(4-benzyloxy-phenyl)-vinyl]-6-bromo-quinazolin-4-yl}- N¹,N¹-diethyl-pentane-1,4-diamine;

MS calc.: 573.6 ; found: 575.1;

with C-(3-aminomethyl-cyclohexyl)-methylamine to obtain

(3-aminomethyl-cyclohexylmethyl)-{2-[2-(4-benzyloxy-phenyl)-vinyl]-6-
5 bromo-quinazolin-4-yl}-amine;

MS calc.: 557.5 ; found: 559.2.


Example 5:

3-{4-[(3-Aminomethyl-cyclohexylmethyl)-amino]-2-[2-(4-benzyloxy-phenyl)-
10 vinyl]-quinazolin-6-yl}-acrylic acid ethyl ester

1. *Synthesis of resin-bound (3-aminomethyl-cyclohexylmethyl)-carbamate
(1)*

p-Nitrophenyl carbonate Wang resin (2,7 mmol, 0,54 mmol/g), 1,3-
15 cyclohexane-bis(dimethylamine) (2,18 g, 15 mmol) and 75 ml DMF
(dimethylformamide) are added to a sealed fritted polypropylene tube. The
mixture is agitated for 72 h. After evacuation of the solvent, the resin is
customary worked up for solid phase reactions.

20



25


2. Analogously to example 1, 6-iodo-2-methylquinazolin-4-one
is reacted with 4-benzyloxy-benzaldehyde and chlorinated to obtain
2-[2-(4-benzyloxy-phenyl)-vinyl]-4-chloro-6-iodo-quinazoline.

30

3. The resin-bound carbamate (1) (4,8 g, 0,54 mmol/g), 2-[2-(4-benzyloxy-
phenyl)-vinyl]-4-chloro-6-iodo-quinazoline (6,6 mmol), triethylamine (1 ml)
and 50 ml DMF are placed in a fritted polypropylene tube. The mixture is

stirred at 80° for 60 hrs. After cooling to rt, the resin is customary worked up for solid phase reactions. Resin bound carbamate (2) is obtained.

5



10

15

20

4. The solid supported 6-iodoquinazoline (2) (0,054 mmol, 0,54 mmol/g), ethyl acrylate (50 mg, 0,5 mmol), Pd(PPh₃)₄ (20 mg), Bu₄NI (tetra-butylammonium-iodide, 32 mg, 0,08 mmol) and 2 ml DMF are placed in a fritted polypropylene tube. The mixture is agitated at 80° for 24 h. After cooling to rt, the solvent is evacuated and the resin is customary worked up for solid phase reactions. The solid supported 6-ethoxyacrylquinazoline and 2 ml of a mixture of H₂O, TFA (trifluoroacetic acid) and dichloromethane (1:49:50) is placed in a fritted polypropylene tube. The contents are shaken for 2 h at rt. The suspension is filtered and the resin is washed with dichloromethane (1 ml) and methanol (1 ml) respectively. Evaporation of the combined filtrates give

3-{4-[(3-aminomethyl-cyclohexylmethyl)-amino]-2-[2-(4-benzyloxy-phenyl)-vinyl]-quinazolin-6-yl}-acrylic acid ethyl ester;

MS calc.: 576.7 ; found: 577.3.

25

Analogously to example 5.4, solid supported 6-iodoquinazoline (2) is reacted with N,N-dimethyl-acrylamide to obtain 3-{4-[(3-aminomethyl-cyclohexylmethyl)-amino]-2-[2-(4-benzyloxy-phenyl)-vinyl]-quinazolin-6-yl}-N,N-dimethyl-acrylamide;

30

MS calc.: 575.8 ; found: 576.4.

Example 6:

5 Solid supported 6-iodoquinazoline (2) [synthesized according to example 5] (0,054 mmol, 0,54 mmol/g), allyltributyltin (140 mg, 0,5 mmol), Pd(PPh₃)₄ (20 mg), and 2 ml DMF are placed in a fritted polypropylene tube. The mixture is agitated at 80° for 24 h. After cooling to rt, the mixture is
5 customary worked up for solid phase reactions. The solid supported 6-allylquinazoline and 2 ml of a mixture of H₂O, TFA and dichloromethane (1:49:50) is placed in a fritted polypropylene tube. The contents are shaken for 2 h at rt. The suspension is filtered and the resin is washed with dichloromethane (1 ml) and methanol (1 ml) respectively. Evaporation of
10 the combined filtrates give
{6-allyl-2-[2-(4-benzyloxy-phenyl)-vinyl]-quinazolin-4-yl}-(3-aminomethyl-cyclohexylmethyl)-amine;
MS calc.: 518.7 ; found: 519.3.

15 Example 7:

Solid supported 6-iodoquinazoline (2) [synthesized according to example 5] (0,054 mmol, 0,54 mmol/g), 4-methylphenylboronic acid (0,5 mmol), Pd(PPh₃)₄ (20 mg), and 2 ml DMF are placed in a fritted polypropylene tube. The mixture is agitated at 80° for 24 h. After cooling to rt, the mixture
20 is customary worked up for solid phase reactions. The solid supported 6-(4-methylphenyl)quinazoline and 2 ml of a mixture of H₂O, TFA and dichloromethane (1:49:50) are placed in a fritted polypropylene tube. The contents are shaken for 2 h at rt. The suspension is filtered and the resin is washed with dichloromethane (1 ml) and methanol (1 ml) respectively.
25 Evaporation of the combined filtrates give
(3-aminomethyl-cyclohexylmethyl)-{2-[2-(4-benzyloxy-phenyl)-vinyl]-6-4-tolyl-quinazolin-4-yl}-amine;
MS calc.: 568.8 ; found: 569.4.

30 Example 8:

Analogously to example 1, 7-chloro-2-methylquinazolin-4-one is reacted with 3,4-bis-benzyloxy-benzaldehyde, chlorinated and reacted

with C-(3-aminomethyl-cyclohexyl)-methylamine to obtain

(3-aminomethyl-cyclohexylmethyl)-{2-[2-(3,4-bis-benzyloxy-phenyl)-vinyl]-7-chloro-quinazolin-4-yl}-amine;

5

with N¹,N¹-diethyl-pentane-1,4-diamine to obtain

N⁴-{2-[2-(3,4-bis-benzyloxy-phenyl)-vinyl]-7-chloro-quinazolin-4-yl}-N¹,N¹-diethyl-pentane-1,4-diamine;

10

with N¹,N¹-diethyl-propane-1,3-diamine to obtain

N¹-{2-[2-(3,4-bis-benzyloxy-phenyl)-vinyl]-7-chloro-quinazolin-4-yl}-N,N-diethyl-propane-1,3-diamine;


MS calc.: 466.6 ; found: 467.3;

15

with 3-(4-methyl-piperazin-1-yl)-propylamine to obtain

{2-[2-(3,4-bis-benzyloxy-phenyl)-vinyl]-7-chloro-quinazolin-4-yl}-[3-(4-methyl-piperazin-1-yl)-propyl]-amine;

20



25

with 2,2-dimethyl-propane-1,3-diamine to obtain

N¹-{2-[2-(3,4-bis-benzyloxy-phenyl)-vinyl]-7-chloro-quinazolin-4-yl}-2,2-dimethyl-propane-1,3-diamine;

30

with 3-aminomethyl-benzylamine to obtain

(3-aminomethyl-benzyl)-{2-[2-(3,4-bis-benzyloxy-phenyl)-vinyl]-7-chloro-quinazolin-4-yl}-amine;

with heptane-1,7-diamine to obtain

5 N¹-{2-[2-(3,4-bis-benzyloxy-phenyl)-vinyl]-7-chloro-quinazolin-4-yl}-heptane-1,7-diamine;


with N¹-(3-amino-propyl)-N¹-methyl-propane-1,3-diamine to obtain

10 N¹-(3-{2-[2-(3,4-bis-benzyloxy-phenyl)-vinyl]-7-chloro-quinazolin-4-ylamino}-propyl)-N¹-methyl-propane-1,3-diamine;

with 3-[4-(3-amino-propyl)-piperazin-1-yl]-propylamine to obtain

{3-[4-(3-amino-propyl)-piperazin-1-yl]-propyl}-{2-[2-(3,4-bis-benzyloxy-phenyl)-vinyl]-7-chloro-quinazolin-4-yl}-amine;

15



20

with C-cyclohexyl-methylamine to obtain

25 {2-[2-(3,4-bis-benzyloxy-phenyl)-vinyl]-7-chloro-quinazolin-4-yl}-cyclohexylmethyl-amine;

with 3-(3-amino-propoxy)-propylamine to obtain

[3-(3-amino-propoxy)-propyl]-{2-[2-(3,4-bis-benzyloxy-phenyl)-vinyl]-7-chloro-quinazolin-4-yl}-amine;

30

with 3-morpholin-4-yl-propylamine to obtain

{2-[2-(3,4-bis-benzyloxy-phenyl)-vinyl]-7-chloro-quinazolin-4-yl}-(3-morpholin-4-yl-propyl)-amine.

Example 9:

- 5 Analogously to example 1, 7-chloro-2-methylquinazolin-4-one is reacted with benzaldehyde, chlorinated and reacted with 3-imidazol-1-yl-propylamine to obtain (7-chloro-2-styryl-quinazolin-4-yl)-(3-imidazol-1-yl-propyl)-amine;
- 10 with N¹,N¹-diethyl-ethane-1,2-diamine to obtain N'-(7-chloro-2-styryl-quinazolin-4-yl)-N,N-diethyl-ethane-1,2-diamine;
- with N¹,N¹-diethyl-propane-1,3-diamine to obtain
- 15 N'-(7-chloro-2-styryl-quinazolin-4-yl)-N,N-diethyl-propane-1,3-diamine;
- with 3-morpholin-4-yl-propylamine to obtain (7-chloro-2-styryl-quinazolin-4-yl)-(3-morpholin-4-yl-propyl)-amine;
- 20 with 1-(3-amino-propyl)-pyrrolidin-2-one to obtain 1-[3-(7-chloro-2-styryl-quinazolin-4-yl-amino)-propyl]-pyrrolidin-2-one;
- with 4-(2-amino-ethyl)-phenylamine to obtain [2-(4-amino-phenyl)-ethyl]-(7-chloro-2-styryl-quinazolin-4-yl)-amine.
- 25 Analogously to example 1, 7-chloro-2-methylquinazolin-4-one is reacted with 4-bromo-benzaldehyde, chlorinated and reacted with N¹,N¹-diethyl-pentane-1,4-diamine to obtain
- 30 N⁴-{2-[2-(4-bromo-phenyl)-vinyl]-7-chloro-quinazolin-4-yl}- N¹,N¹-diethyl-pentane-1,4-diamine;
- MS calc.: 501.9 ; found: 501.9.

Analogously to example 1, 7-chloro-2-methylquinazolin-4-one is reacted with 3-phenyl-propenal, chlorinated and reacted

5 with N¹,N¹-diethyl-pentane-1,4-diamine to obtain

N⁴-[7-chloro-2-(4-phenyl-buta-1,3-dienyl)-quinazolin-4-yl]-N¹,N¹-diethyl-pentane-1,4-diamine;


MS calc.: 414.6 ; found: 415.3.

10 Example 10:

1. Analogously to example 1, 2-methylquinazolin-4-one is reacted with 4-bromo-benzaldehyde and chlorinated to obtain 2-[2-(4-bromo-phenyl)-vinyl]-4-chloro-quinazoline.

15 2. The resin-bound carbamate (1) (4,8 g, 0,54 mmol/g) [synthesized analogously to example 5.1], 2-[2-(4-bromo-phenyl)-vinyl]-4-chloro-quinazoline (6,6 mmol), triethylamine (1 ml) and 50 ml DMF are placed in a fritted polypropylene tube. The mixture is stirred at 80° for 60 hrs. After cooling to rt, the resin is customary worked up for solid phase reactions.

20 Resin bound carbamate (3) is obtained.




30 3. The solid supported 2-bromostyrylquinazoline (3) (0,054 mmol, 0,54 mmol/g), phenylboronic acid (0,5 mmol), Pd(PPh₃)₄ (20 mg), triethylamine (20 ml) and 2 ml DMF are placed in a fritted polypropylene tube. The mixture is agitated at 80° for 24 h. After cooling to rt, the mixture is

customary worked up for solid phase reactions. The solid-supported 2-(2-biphenyl-4-yl-vinyl)-quinazoline and 2 ml of a mixture of H₂O, TFA and dichloromethane (1:49:50) is placed in a fritted polypropylene tube. The contents are shaken for 2 h at rt. The suspension is filtered and the resin is washed with dichloromethane (1 ml) and methanol (1 ml) respectively. Evaporation of the combined filtrates give (3-aminomethyl-cyclohexylmethyl)-[2-(2-biphenyl-4-yl-vinyl)-quinazolin-4-yl]-amine;

MS calc.: 448.6 ; found: 449.4.

10

Analogously to example 10, 2-methylquinazolin-4-one is reacted with 4-bromo-benzaldehyde, chlorinated, reacted with resin bound carbamate (4) [synthesized analogously to example 5.1]



and phenylboronic acid to obtain

N⁴-[2-(2-biphenyl-4-yl-vinyl)-7-chloro-quinazolin-4-yl]-N¹,N¹-diethyl-pentane-1,4-diamine;

20 MS calc.: 464.7 ; found: 465.2.

Example 11:


Analogously to example 10, 7-chloro-2-methylquinazolin-4-one is reacted with 4-bromo-benzaldehyde, chlorinated, reacted with resin bound carbamate (1) and phenylboronic acid to obtain (3-aminomethyl-cyclohexylmethyl)-[2-(2-biphenyl-4-yl-vinyl)-7-chloro-quinazolin-4-yl]-amine;

MS calc.: 483.1 ; found: 483.3.

30 Analogously to example 10, 7-chloro-2-methylquinazolin-4-one is reacted with 4-bromo-benzaldehyde, chlorinated, reacted

with resin bound carbamate (5) [synthesized analogously to example 5.1]

5




and phenylboronic acid to obtain

N^1 -[2-(2-biphenyl-4-yl-vinyl)-7-chloro-quinazolin-4-yl]-heptane-1,7-diamine;

MS calc.: 471.0 ; found: 471.4;

10

with resin bound carbamate (6) [synthesized analogously to example 5.1]



15


and phenylboronic acid to obtain

[3-(3-amino-propoxy)-propyl]-[2-(2-biphenyl-4-yl-vinyl)-7-chloro-quinazolin-4-yl]-amine;

MS calc.: 473.0 ; found: 473.3;

20

with resin bound carbamate (7) [synthesized analogously to example 5.1]



and phenylboronic acid to obtain


25

1-amino-3-[2-(2-biphenyl-4-yl-vinyl)-7-chloro-quinazolin-4-ylamino]-propan-2-ol;

MS calc.: 430.9 ; found: 431.2;

30


with resin bound carbamate (8) [synthesized analogously to example 5.1]



and phenylboronic acid to obtain

N^1 -[2-(2-biphenyl-4-yl-vinyl)-7-chloro-quinazolin-4-yl]-2,2-dimethyl-propane-1,3-diamine;

5 with resin bound carbamate (9) [synthesized analogously to example 5.1]




and phenylboronic acid to obtain

10 {3-[4-(3-amino-propyl)-piperazin-1-yl]-propyl}-[2-(2-biphenyl-4-yl-vinyl)-7-chloro-quinazolin-4-yl]-amine;

MS calc.: 541.1 ; found: 541.3;

with resin bound carbamate (10) [synthesized analogously to example 5.1]



and phenylboronic acid to obtain

20 N^1 -{3-[2-(2-biphenyl-4-yl-vinyl)-7-chloro-quinazolin-4-ylamino]-propyl}- N^1 -methyl-propane-1,3-diamine;

MS calc.: 486.1 ; found: 486.2.

Example 12:

25 Analogously to example 10, 6-iodo-2-methylquinazolin-4-one is reacted with 4-bromo-benzaldehyde, chlorinated, reacted with resin bound carbamate (1) and phenylboronic acid to obtain

(3-aminomethyl-cyclohexylmethyl)-[2-(2-biphenyl-4-yl-vinyl)-6-iodo-quinazolin-4-yl]-amine;

MS calc.: 574.5 ; found: 575.2.

30

Analogously to example 10, 6-iodo-2-methylquinazolin-4-one

is reacted with 4-bromo-benzaldehyde, chlorinated, reacted

with resin bound carbamate (5) and phenylboronic acid to obtain

N^1 -[2-(2-biphenyl-4-yl-vinyl)-6-iodo-quinazolin-4-yl]-heptane-1,7-diamine;

5 MS calc.: 562.5 ; found: 563.3;

with resin bound carbamate (6) and phenylboronic acid to obtain

[3-(3-amino-propoxy)-propyl]-[2-(2-biphenyl-4-yl-vinyl)-6-iodo-quinazolin-4-yl]-amine;

10 MS calc.: 564.5 ; found: 565.2;

with resin bound carbamate (7) and phenylboronic acid to obtain

1-amino-3-[2-(2-biphenyl-4-yl-vinyl)-6-iodo-quinazolin-4-ylamino]-propan-2-ol;

15 MS calc.: 522.4 ; found: 523.2;

with resin bound carbamate (8) and phenylboronic acid to obtain

N^1 -[2-(2-biphenyl-4-yl-vinyl)-6-iodo-quinazolin-4-yl]-2,2-dimethyl-propane-1,3-diamine;

20 MS calc.: 534.4 ; found: 535.2;

with resin bound carbamate (9) and phenylboronic acid to obtain

{3-[4-(3-amino-propyl)-piperazin-1-yl]-propyl}-[2-(2-biphenyl-4-yl-vinyl)-6-iodo-quinazolin-4-yl]-amine;

25 MS calc.: 632.6 ; found: 633.2;

with resin bound carbamate (10) and phenylboronic acid to obtain

N^1 -{3-[2-(2-biphenyl-4-yl-vinyl)-6-iodo-quinazolin-4-ylamino]-propyl}- N^1 -methyl-propane-1,3-diamine;

30 MS calc.: 577.5 ; found: 578.1.

Example 13:

Analogously to example 10, 7-chloro-2-methylquinazolin-4-one is reacted with 4-bromo-benzaldehyde, chlorinated, reacted with resin bound carbamate (1) and

5 2-methylphenylboronic acid to obtain

(3-aminomethyl-cyclohexylmethyl)-{7-chloro-2-[2-(2'-methyl-biphenyl-4-yl)-vinyl]-quinazolin-4-yl}-amine;

MS calc.: 497.1 ; found: 497.4;

10 2,4-dichlorophenylboronic acid

(3-aminomethyl-cyclohexylmethyl)-{7-chloro-2-[2-(2',4'-dichloro-biphenyl-4-yl)-vinyl]-quinazolin-4-yl}-amine;

MS calc.: 551.9 ; found: 551.3;

15 4-fluorophenylboronic acid

(3-aminomethyl-cyclohexylmethyl)-{7-chloro-2-[2-(4'-fluoro-biphenyl-4-yl)-vinyl]-quinazolin-4-yl}-amine;


MS calc.: 501.1 ; found: 501.4;

20 naphthylboronic acid

(3-aminomethyl-cyclohexylmethyl)-{7-chloro-2-[2-(4-naphthalen-1-yl-phenyl)-vinyl]-quinazolin-4-yl}-amine;

MS calc.: 533.1 ; found: 533.4;

25



30

4-methoxyphenylboronic acid


- (3-aminomethyl-cyclohexylmethyl)-{7-chloro-2-[2-(4'-methoxy-biphenyl-4-yl)-vinyl]-quinazolin-4-yl}-amine;
MS calc.: 513.1 ; found: 513.4;
- 5 3,4,5-trimethoxyphenylboronic acid
(3-aminomethyl-cyclohexylmethyl)-{7-chloro-2-[2-(3',4',5'-trimethoxy-biphenyl-4-yl)-vinyl]-quinazolin-4-yl}-amine;
MS calc.: 573.1 ; found: 573.4;
- 10 3-acetylaminophenylboronic acid
N-[4'-(2-{4-[(3-aminomethyl-cyclohexylmethyl)-amino]-7-chloro-quinazolin-2-yl}-vinyl)-biphenyl-3-yl]-acetamide;
MS calc.: 540.1 ; found: 540.4;
- 15 3-acetylphenylboronic acid
1-[4'-(2-{4-[(3-aminomethyl-cyclohexylmethyl)-amino]-7-chloro-quinazolin-2-yl}-vinyl)-biphenyl-3-yl]-ethanone;
MS calc.: 525.1 ; found: 525.4;
- 20 benzo[b]-thiophen-2-ylboronic acid
(3-aminomethyl-cyclohexylmethyl)-{2-[2-(4-benzo[b]thiophen-2-yl-phenyl)-vinyl]-7-chloro-quinazolin-4-yl}-amine;
- 3,5-bis-trifluoromethylphenylboronic acid
- 25 (3-aminomethyl-cyclohexylmethyl)-{2-[2-(3',5'-bis-trifluoromethyl-biphenyl-4-yl)-vinyl]-7-chloro-quinazolin-4-yl}-amine;
- 3-nitrophenylboronic acid
(3-aminomethyl-cyclohexylmethyl)-{7-chloro-2-[2-(3'-nitro-biphenyl-4-yl)-vinyl]-quinazolin-4-yl}-amine;
MS calc.: 528.1 ; found: 528.3;
- 30

thiophenylboronic acid

(3-aminomethyl-cyclohexylmethyl)-{7-chloro-2-[2-(4-thiophen-2-yl-phenyl)-vinyl]-quinazolin-4-yl}-amine;

MS calc.: 489.1 ; found: 489.4;

5



10

3-aminophenylboronic acid

(3-aminomethyl-cyclohexylmethyl)-{7-chloro-2-[2-(3'-amino-biphenyl-4-yl)-vinyl]-quinazolin-4-yl}-amine;

15

MS calc.: 498.1 ; found: 498.4;

3-isopropylphenylboronic acid

(3-aminomethyl-cyclohexylmethyl)-{7-chloro-2-[2-(3'-isopropyl-biphenyl-4-yl)-vinyl]-quinazolin-4-yl}-amine;

20

MS calc.: 525.1 ; found: 525.4;

pyridin-3-ylboronic acid

(3-aminomethyl-cyclohexylmethyl)-{7-chloro-2-[2-(4-pyridin-3-yl-phenyl)-vinyl]-quinazolin-4-yl}-amine;

25

MS calc.: 484.0 ; found: 484.4.

Example 14:

Analogously to example 10, 7-chloro-2-methylquinazolin-4-one is reacted with 5-bromo-furan-2-carbaldehyde, chlorinated, reacted with resin bound carbamate (1) and

30

3-chlorophenylboronic acid to obtain

(3-aminomethyl-cyclohexylmethyl)-(7-chloro-2-{2-[5-(3-chloro-phenyl)-furan-2-yl]-vinyl}-quinazolin-4-yl)-amine;

MS calc.: 507.5 ; found: 507.8.

5


Example 15:

Analogously to example 10, 7-chloro-2-methylquinazolin-4-one is reacted with 5-bromo-thiophene-2-carbaldehyde, chlorinated, reacted with resin bound carbamate (1) and thiophen-2-ylboronic acid to obtain

10

(3-aminomethyl-cyclohexylmethyl)-[2-(2-[2,2']bithiophenyl-5-yl-vinyl)-7-chloro-quinazolin-4-yl]-amine

15



20

Analogously to example 10, 7-chloro-2-methylquinazolin-4-one is reacted with 5-bromo-thiophene-2-carbaldehyde, chlorinated, reacted with resin bound carbamate (4) and thiophenylboronic acid to obtain

25


N⁴-[2-(2-[2,2']bithiophenyl-5-yl-vinyl)-7-chloro-quinazolin-4-yl]-N¹,N¹-diethyl-pentane-1,4-diamine;

MS calc.: 511.2 ; found: 511.1;

with resin bound carbamate (11)

30

- 72 -




and thiophenylboronic acid to obtain

5 N'-[2-(2-[2,2']bithiophenyl-5-yl-vinyl)-7-chloro-quinazolin-4-yl]-N,N-diethyl-ethane-1,2-diamine;

MS calc.: 469.1 ; found: 469.1;

with resin bound carbamate (12)




and thiophenylboronic acid to obtain

15 N'-[2-(2-[2,2']bithiophenyl-5-yl-vinyl)-7-chloro-quinazolin-4-yl]-N,N-diethyl-propane-1,3-diamine;

MS calc.: 483.1 ; found: 483.1;

with resin bound carbamate (13)




and thiophenylboronic acid to obtain

25 N¹-[2-(2-[2,2']bithiophenyl-5-yl-vinyl)-7-chloro-quinazolin-4-yl]-propane-1,3-diamine;

MS calc.: 427.0 ; found: 427.4;

with resin bound carbamate (14)



and thiophenylboronic acid to obtain

(3-aminomethyl-benzyl)-[2-(2-[2,2']bithiophenyl-5-yl-vinyl)-7-chloro-quinazolin-4-yl]-amine;

MS calc.: 489.1 ; found: 489.1;

5 with resin bound carbamate (5) and thiophenylboronic acid to obtain
N¹-[2-(2-[2,2']bithiophenyl-5-yl-vinyl)-7-chloro-quinazolin-4-yl]-heptane-1,7-diamine;

MS calc.: 483.1 ; found: 483.2;


10 with resin bound carbamate (10) and thiophenylboronic acid to obtain
N¹-{3-[2-(2-[2,2']bithiophenyl-5-yl-vinyl)-7-chloro-quinazolin-4-ylamino]-propyl}-N¹-methyl-propane-1,3-diamine;

MS calc.: 498.1 ; found: 498.1;

15 with resin bound carbamate (9) and thiophenylboronic acid to obtain
{3-[4-(3-amino-propyl)-piperazin-1-yl]-propyl}-[2-(2-[2,2']bithiophenyl-5-yl-vinyl)-7-chloro-quinazolin-4-yl]-amine;

MS calc.: 553.2 ; found: 553.2;

20 with resin bound carbamate (15)



and thiophenylboronic acid to obtain

25 [2-(2-[2,2']bithiophenyl-5-yl-vinyl)-7-chloro-quinazolin-4-yl]-piperidin-4-ylmethyl-amine;


MS calc.: 467.1 ; found: 467.2;

with resin bound carbamate (6) and thiophenylboronic acid to obtain

30 [3-(3-amino-propoxy)-propyl]-[2-(2-[2,2']bithiophenyl-5-yl-vinyl)-7-chloro-quinazolin-4-yl]-amine;

MS calc.: 485.1 ; found: 485.2;

with resin bound carbamate (16)



and thiophenylboronic acid to obtain

[2-(2-[2,2']bithiophenyl-5-yl-vinyl)-7-chloro-quinazolin-4-yl]-pyridin-4-ylmethyl-amine;

MS calc.: 461.0 ; found: 461.2.

10

Analogously to example 10, 6-iodo-2-methylquinazolin-4-one is reacted with 5-bromo-thiophene-2-carbaldehyde, chlorinated, reacted

15

with resin bound carbamate (1) and thiophen-2-ylboronic acid to obtain

(3-aminomethyl-cyclohexylmethyl)-[2-(2-[2,2']bithiophenyl-5-yl-vinyl)-6-iodo-quinazolin-4-yl]-amine.

20

with resin bound carbamate (12) and thiophen-2-ylboronic acid to obtain

N'-[2-(2-[2,2']bithiophenyl-5-yl-vinyl)-6-iodo-quinazolin-4-yl]-N,N-diethylpropane-1,3-diamine;

MS calc.: 574.5 ; found: 575.2.

25

The following examples relate to pharmaceutical preparations:

Example A: Injection vials

30

A solution of 100 g of an active compound of the formula I and 5 g of disodium hydrogenphosphate is adjusted to pH 6.5 in 3 l of double-distilled water using 2N hydrochloric acid, sterile-filtered, dispensed into injection vials, lyophilized under sterile conditions and aseptically sealed. Each injection vial contains 5 mg of active compound.

Example B: Suppositories

A mixture of 20 g of an active compound of the formula I is melted with 100 g of soya lecithin and 1400 g of cocoa butter, poured into moulds and allowed to cool. Each suppository contains 20 mg of active compound.

5 **Example C: Solution**

A solution is prepared from 1 g of an active compound of the formula I, 9.38 g of $\text{NaH}_2\text{PO}_4 \cdot 2\text{H}_2\text{O}$, 28.48 g of $\text{Na}_2\text{HPO}_4 \cdot 12\text{H}_2\text{O}$ and 0.1 g of benzalkonium chloride in 940 ml of double-distilled water. The mixture is adjusted to pH 6.8, made up to 1 l and sterilized by irradiation. This solution
10 can be used in the form of eye drops.

Example D: Ointment

500 mg of an active compound of the formula I is mixed with 99.5 g of petroleum jelly under aseptic conditions.

15

Example E: Tablets

A mixture of 1 kg of active compound of the formula I, 4 kg of lactose, 1.2 kg of potato starch, 0.2 g of talc and 0.1 kg of magnesium stearate is compressed in a customary manner to give tablets such that each tablet
20 contains 10 mg of active compound.

Example F: Coated tablets

Analogously to Example E, tablets are pressed which are then coated with a coating of sucrose, potato starch, talc, tragacanth and colorant in a
25 customary manner.

Example G: Capsules

2 kg of active compound of the formula I are dispensed into hard gelatin capsules in a customary manner such that each capsule contains 20 mg of
30 the active compound.

Example H: Ampules

A solution of 1 kg of active compound of the formula I in 60 ml of double-distilled water is sterile-filtered, dispensed into ampoules, lyophilized under sterile conditions and aseptically sealed. Each ampoule contains 10 mg of active compound.

5

10

15

20


25

30

What is claimed is:

1. Compounds of the formula I

5



in which


10

R and R¹ are independently of each other H, A, OH, OA, Hal, N(R⁵)₂, NO₂, CN, C(O)R², CON(R⁵)₂, COOR⁵, allyl, CH=CH-COOR⁵, CH=CHCON(R⁵)₂, SO₂A or phenyl, which is unsubstituted or mono-, di- or trisubstituted by A,

15

R² and R³ are independently of each other H, A, cycloalkyl, -Het³, -(CH₂)_o-OR⁵, -(CH₂)_o-OR⁶, -(CH₂)_o-Het¹, -(CH₂)_o-NR⁵-Het¹, -(CHA)_p-(CH₂)_o-N(R⁵)₂, -(CH₂)_p-(CHA)_p-(CH₂)_m-Ar, -(CH₂)_o-Z-(CH₂)_q-N(R⁵)₂,

20




25

provided that R² and R³ together are not H, or NR²R³ together form a saturated monocyclic heterocyclic radical having 5 to 6 ring members, where 1 or 2 N atoms are present and the heterocyclic radical can be mono- or disubstituted by OH, Ar, OAr or arylalkyl,

30


R⁴ is Ar or Het¹,
 R⁵ is H or A,
 R⁶ is benzo[1,3]dioxol-5-yl,
 Q is O or S,

- Y is (CH=CH)_n,
- Z is phenylene, cyclohexylene, -NR⁵-, O, -CH(OH)-, -CA₂- or
- 
- 5
- A is unbranched or branched alkyl having 1 to 6 carbon atoms,
- Ar is phenyl, naphthyl or biphenyl, which is unsubstituted or mono-, di- or trisubstituted by A, OH, OA, cycloalkyloxy, O-(CH₂)_p-Ph, CF₃, OCF₃, Hal, CN, CHO, COA, COOR⁵, N(R⁵)₂, NR⁵-COA, NO₂, SO₂N(R⁵)₂, mor, SO₂-mor, 5-methyl-3-oxo-2,4-dihydropyrazol-2-yl, naphthyl or Het²,
- 10
- Het¹ is a saturated, partially or completely unsaturated mono- or bicyclic heterocyclic radical having 5 to 10 ring members, where 1 or 2 N and/or 1 or 2 S or O atoms can be present and the heterocyclic radical can be mono- or disubstituted by A, Hal, OH, OA, CF₃, OCF₃, N(R⁵)₂, carbonyl oxygen, COOR⁵, Het², benzyl or phenyl which is unsubstituted or mono-, di- or trisubstituted by A, OH, OA, CF₃, OCF₃, Hal, CN, COOR⁵, N(R⁵)₂, NO₂ or SO₂N(R⁵)₂,
- 15
- 20 Het² is a unsaturated mono- or bicyclic heterocyclic radical having 5 to 10 ring members, where 1 or 2 N and/or 1 or 2 S or O atoms can be present and the heterocyclic radical can be mono- or disubstituted by A, Hal, OH, OA, CF₃, OCF₃, N(R⁵)₂ or COOR⁵,
- 25 Het³ is a partially or completely unsaturated mono- or bicyclic heterocyclic radical having 5 to 10 ring members, where 1 or 2 N atoms are present and the heterocyclic radical can be mono- or disubstituted by A, Hal, OH, OA, CF₃, OCF₃, N(R⁵)₂, SO₂A or COOR⁵ provided that the heterocyclic radical is not bonded via an N atom,
- 30
- Hal is F, Cl, Br or I,
- mor is morpholin-4-yl,

Ph is phenyl,
 n is 1 or 2,
 m is 0, 1, 2, 3, 4, 5 or 6,
 o is 1, 2, 3, 4, 5, 6 or 7,
 5 p is 0, 1, 2, 3 or 4,
 q is 1, 2, 3 or 4,

and their pharmaceutically tolerable salts and solvates solvates as glycoprotein IbIX antagonists.

10 2. Compounds of the formula I



15

in which

R and R¹ are independently of each other H, A, OH, OA, Hal, N(R⁵)₂, NO₂, CN, C(O)R², CON(R⁵)₂, COOR⁵, allyl, CH=CH-COOR⁵, CH=CHCON(R⁵)₂, SO₂A or phenyl, which is unsubstituted or mono-, di- or trisubstituted by A,

20

R² is H,

R³ is -(CH₂)_o-Z-(CH₂)_q-N(R⁵)₂,

R⁴ is Ar,

R⁵ is H or A,

25

Y is (CH=CH)_n,

Z is phenylene, cyclohexylene, -NR⁵-, O, -CH(OH)-, -CA₂- or



30

A is unbranched or branched alkyl having 1 to 6 carbon atoms,

Ar is phenyl or naphthyl, which is unsubstituted or mono-, di- or trisubstituted by A, OH, OA, CF₃, OCF₃, Hal, CN, CHO, COA, COOR⁵, N(R⁵)₂, NO₂ or SO₂N(R⁵)₂,

Hal is F, Cl, Br or I,


5 n is 1 or 2,

o is 1, 2, 3, 4, 5, 6 or 7,

q is 1, 2, 3 or 4,

and their pharmaceutically tolerable salts and solvates.

10 3. Compounds of the formula I



15

in which

R and R¹ are independently of each other H, A, OH, OA, Hal, N(R⁵)₂, NO₂, CN, C(O)R², CON(R⁵)₂, COOR⁵, allyl, CH=CH-COOR⁵, CH=CHCON(R⁵)₂, SO₂A or phenyl, which is unsubstituted or mono-, di- or trisubstituted by A,

20

R² and R³ are independently of each other H, A, cycloalkyl, -Het³, -(CH₂)_o-OR⁵, -(CH₂)_o-OR⁶, -(CH₂)_o-Het¹, -(CH₂)_o-NR⁵-Het¹, -(CHA)_p-(CH₂)_o-N(R⁵)₂, -(CH₂)_p-(CHA)_p-(CH₂)_m-Ar or -(CH₂)_o-Z-(CH₂)_q-N(R⁵)₂,

25

provided that R² and R³ together are not H,

R⁴ is Ar,


R⁵ is H or A,

R⁶ is benzo[1,3]dioxol-5-yl,

Y is (CH=CH)_n,

30

Z is phenylene, cyclohexylene, -NR⁵-, O, -CH(OH)-, -CA₂- or



- 5 A is unbranched or branched alkyl having 1 to 6 carbon atoms,
Ar is phenyl, which is mono-, di- or trisubstituted by O-(CH₂)_p-Ph, naphthyl or Het², or biphenyl, which is unsubstituted or mono-, di- or trisubstituted by A, OH, OA, CF₃, OCF₃, Hal, CN, CHO, COA, COOR⁵, N(R⁵)₂, NR⁵-COA, NO₂, SO₂N(R⁵)₂, naphthyl or Het²,
- 10 Het¹ is a saturated, partially or completely unsaturated mono- or bicyclic heterocyclic radical having 5 to 10 ring members, where 1 or 2 N and/or 1 or 2 S or O atoms can be present and the heterocyclic radical can be mono- or disubstituted by A, Hal, OH, OA, CF₃, OCF₃, N(R⁵)₂, carbonyl oxygen, COOR⁵, Het², benzyl or phenyl which is unsubstituted or mono-, di- or trisubstituted by A, OH, OA, CF₃, OCF₃, Hal, CN, COOR⁵, N(R⁵)₂, NO₂ or SO₂N(R⁵)₂,
- 15 Het² is a unsaturated mono- or bicyclic heterocyclic radical having 5 to 10 ring members, where 1 or 2 N and/or 1 or 2 S or O atoms can be present and the heterocyclic radical can be mono- or disubstituted by A, Hal, OH, OA, CF₃, OCF₃, N(R⁵)₂ or COOR⁵,
- 20 Het³ is a partially or completely unsaturated mono- or bicyclic heterocyclic radical having 5 to 10 ring members, where 1 or 2 N atoms are present and the heterocyclic radical can be mono- or disubstituted by A, Hal, OH, OA, CF₃, OCF₃, N(R⁵)₂, SO₂A or COOR⁵ provided that the heterocyclic radical is not bonded via an N atom,
- 25 Hal is F, Cl, Br or I,
Ph is phenyl,
30 n is 1 or 2,
m is 0, 1, 2, 3, 4, 5 or 6,

o is 1, 2, 3, 4, 5, 6 or 7,

p is 0, 1, 2, 3 or 4,


q is 1, 2, 3 or 4,

and their pharmaceutically tolerable salts and solvates.

5

4. Compounds of the formula I

10




in which

R and R¹ are independently of each other H, A, OH, OA, Hal, N(R⁵)₂, NO₂, CN, C(O)R², CON(R⁵)₂, COOR⁵, allyl, CH=CH-COOR⁵, CH=CHCON(R⁵)₂, SO₂A or phenyl, which is unsubstituted or mono-, di- or trisubstituted by A,

15

R² and R³ are independently of each other H, A, cycloalkyl, -Het³, -(CH₂)_o-OR⁵, -(CH₂)_o-OR⁶, -(CH₂)_o-Het¹, -(CH₂)_o-NR⁵-Het¹, -(CHA)_p-(CH₂)_o-N(R⁵)₂, -(CH₂)_p-(CHA)_p-(CH₂)_m-Ar, -(CH₂)_o-Z-(CH₂)_q-N(R⁵)₂,

20




25

provided that R² and R³ together are not H, or NR²R³ together form a saturated monocyclic heterocyclic radical having 5 to 6 ring members, where 1 or 2 N atoms are present and the heterocyclic radical can be mono- or disubstituted by OH, Ar, OAr or arylalkyl,

30

R⁴ is Het¹,

- R^5 is H or A,
 R^6 is benzo[1,3]dioxol-5-yl,
 Q is O or S,
 Y is $(CH=CH)_n$,
 5 Z is phenylene, cyclohexylene, $-NR^5-$, O, $-CH(OH)-$, $-CA_2-$ or


 A is unbranched or branched alkyl having 1 to 6 carbon atoms,
 10 Ar is phenyl, naphthyl or biphenyl, which is unsubstituted or
 mono-, di- or trisubstituted by A, OH, OA, cycloalkyloxy,
 $O-(CH_2)_p-Ph$, CF_3 , OCF_3 , Hal, CN, CHO, COA, $COOR^5$,
 $N(R^5)_2$, NR^5-COA , NO_2 , $SO_2N(R^5)_2$, mor, SO_2-mor , 5-methyl-
 3-oxo-2,4-dihydropyrazol-2-yl, naphthyl or Het^2 ,
 15 Het^1 is a saturated, partially or completely unsaturated mono- or
 bicyclic heterocyclic radical having 5 to 10 ring members,
 where 1 or 2 N and/or 1 or 2 S or O atoms can be present and
 the heterocyclic radical can be mono- or disubstituted by A,
 Hal, OH, OA, CF_3 , OCF_3 , $N(R^5)_2$, carbonyl oxygen, $COOR^5$,
 20 Het^2 , benzyl or phenyl which is unsubstituted or mono-, di- or
 trisubstituted by A, OH, OA, CF_3 , OCF_3 , Hal, CN, $COOR^5$,
 $N(R^5)_2$, NO_2 or $SO_2N(R^5)_2$,
 Het^2 is a unsaturated mono- or bicyclic heterocyclic radical having
 5 to 10 ring members, where 1 or 2 N and/or 1 or 2 S or
 25 O atoms can be present and the heterocyclic radical can be
 mono- or disubstituted by A, Hal, OH, OA, CF_3 , OCF_3 , $N(R^5)_2$
 or $COOR^5$,
 Het^3 is a partially or completely unsaturated mono- or bicyclic
 heterocyclic radical having 5 to 10 ring members, where 1 or 2
 30 N atoms are present and the heterocyclic radical can be
 mono- or disubstituted by A, Hal, OH, OA, CF_3 , OCF_3 , $N(R^5)_2$,

SO₂A or COOR⁵ provided that the heterocyclic radical is not bonded via an N atom,

Hal is F, Cl, Br or I,

mor is morpholin-4-yl,

5 Ph is phenyl,

n is 1 or 2,

m is 0, 1, 2, 3, 4, 5 or 6,

o is 1, 2, 3, 4, 5, 6 or 7,

p is 0, 1, 2, 3 or 4,

10 q is 1, 2, 3 or 4,

and their pharmaceutically tolerable salts and solvates.

5. A compound selected from the group:

a) (7-chloro-2-styryl-quinazolin-4-yl)-(3-imidazol-1-yl-propyl)-amine,

15 b) N'-(7-chloro-2-styryl-quinazolin-4-yl)-N,N-diethyl-ethane-1,2-diamine,

c) N'-(7-chloro-2-styryl-quinazolin-4-yl)-N,N-diethyl-propane-1,3-diamine,

d) (7-chloro-2-styryl-quinazolin-4-yl)-(3-morpholin-4-yl-propyl)-amine,

e) 1-[3-(7-chloro-2-styryl-quinazolin-4-yl-amino)-propyl]-pyrrolidin-2-one,

f) [2-(4-amino-phenyl)-ethyl]-(7-chloro-2-styryl-quinazolin-4-yl)-amine,

20 g) N⁴-{2-[2-(4-bromo-phenyl)-vinyl]-7-chloro-quinazolin-4-yl}-N¹,N¹-diethyl-pentane-1,4-diamine and

h) N⁴-[7-chloro-2-(4-phenyl-buta-1,3-dienyl)-quinazolin-4-yl]-N¹,N¹-diethyl-pentane-1,4-diamine

and their pharmaceutically tolerable salts and solvates.

25

6. Compounds of the formula I according to Claim 4


a) N'-[2-(2-[2,2']bithiophenyl-5-yl-vinyl)-6-iodo-quinazolin-4-yl]-N,N-diethyl-propane-1,3-diamine,

30 b) (3-aminomethyl-cyclohexylmethyl)-[2-(2-[2,2']bithiophenyl-5-yl-vinyl)-7-chloro-quinazolin-4-yl]-amine

and their physiologically acceptable salts and solvates.

7. Process for the preparation of novel compounds of the formula I

5



in which

R, R¹, R², R³, R⁴ and Y have the meaning indicated in Claims 1 to 4 and their pharmaceutically tolerable salts and solvates, characterized in that


10

a) a compound of the formula I according to claims 1 to 4 is liberated from one of its functional derivatives by treating with a solvolysing or hydrogenolysing agent,

or

15

b) in stage 1) a compound of the formula II




20

in which


R and R¹ have the meaning as given in Claims 1 to 4, is reacted with a compound of the formula III

25




in which R⁴ has the meaning indicated in Claims 1 to 4 and s is 0 or 1, to give a compound of formula IV

30




in which R, R¹ and R⁴ have the meaning indicated in Claims 1 to 4
and s is 0 or 1,

10 in stage 2) a compound of formula IV as indicated above is reacted with
a chlorinating agent to give a compound of formula V



in which R, R¹ and R⁴ have the meaning indicated in Claims 1 to 4 and s
is 0 or 1


and in stage 3) a compound of formula V as indicated above is reacted
with a compound of formula VI



in which R² and R³ or NR²R³ have the meaning indicated in Claims 1 to
4,

or


25 c) in stage 1) a compound of the formula II



in which

R and R¹ have the meaning as given in Claims 1 to 4,

is reacted with a chlorinating agent to give a compound of formula VII




in which

R and R¹ have the meaning as given in Claims 1 to 4,


in stage 2) a compound of formula VII as indicated above is reacted with a compound of formula VI

10



in which R² and R³ or NR²R³ have the meaning indicated in Claims 1 to 4 to give a compound of formula VIII

15




20

in which R, R¹, R², R³ and NR²R³ have the meaning indicated in Claims 1 to 4

and in stage 3) a compound of formula VIII as indicated above is reacted with a compound of formula III

25



in which R⁴ has the meaning indicated in Claims 1 to 4 and s is 0 or 1 or

d) a radical R, R¹, R², R³ and/or R⁴ is converted into another radical R, R¹, R², R³ and/or R⁴ by, for example

30

- reducing a nitro group, sulfonyl group or sulfoxyl group,

- etherifying an OH group or subjecting an OA group to ether cleavage,
 - alkylating a primary or secondary amino group,
 - partially or completely hydrolysing a CN group,
 - 5 - cleaving an ester group or esterifying a carboxylic acid radical,
 - reacting an aryl bromide, aryl iodide, heteroaryl bromide or heteroaryliodide to give the corresponding coupling products by means of a Suzuki coupling with boronic acids,
 - reacting a iodoquinazoline or bromoquinazoline to give the
 - 10 corresponding coupling products by means of a Stille coupling with allyltributyltin,
 - reacting a iodoquinazoline or bromoquinazoline to give the corresponding coupling products by means of a Heck coupling with acrylates,
 - 15 - or carrying out a nucleophilic or electrophilic substitution, and/or
a base or acid of the formula I is converted into one of its salts or solvates.
- 20 8. Compounds of the formula I according to Claims 2 to 5 and their physiologically acceptable salts or solvates as pharmaceutical active compounds.
- 25 9. Compounds of the formula I according to Claim 8 and their physiologically acceptable salts or solvates as glycoprotein IbIX antagonists.
- 30 10. Compounds of the formula I according to Claims 1 and 8 and their physiologically acceptable salts or solvates as glycoprotein IbIX antagonists for the control of thrombotic disorders and sequelae deriving therefrom.

11. Pharmaceutical preparation characterized in that it contains at least one compound of the formula I according to Claim 10 and/or one of its physiologically acceptable salts or solvates.
- 5 12. Use of compounds of the formula I according to Claims 1 to 5 and/or their physiologically acceptable salts or solvates for the production of a pharmaceutical preparation for the control of thrombotic disorders and sequelae deriving therefrom or for use as anti-adhesive substances.
- 10 13. Use of compounds of the formula I according to Claims 1 to 5 and/or their physiologically acceptable salts or solvates for the production of a pharmaceutical preparation for the treatment of illnesses, such as for the prophylaxis and/or therapy of thrombotic disorders, as well as sequelae such as, for example, myocardial infarct, arteriosclerosis, angina pectoris, acute coronary syndromes, peripheral circulatory disorders, 15 stroke, transient ischaemic attacks, reocclusion/restenosis after angioplasty/stent implantations or as anti-adhesive substances for implants, catheters or heart pacemakers.
- 20
- 25
- 30

